



(43) International Publication Date  
24 October 2013 (24.10.2013)

(10) International Publication Number  
**WO 2013/157002 A1**

- (51) International Patent Classification:  
A61K 38/24 (2006.01)
- (21) International Application Number:  
PCT/IL2013/050332
- (22) International Filing Date:  
17 April 2013 (17.04.2013)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
61/635,483 19 April 2012 (19.04.2012) US
- (71) Applicant: PROLOR BIOTECH INC. [IL/IL]; 7 Golda Meir Street, 2nd floor, 74140 Nes Ziona (IL).
- (72) Inventors: FIMA, Udi Eyal; P.O.B 238, 85330 Dvira (IL). HERSHKOVITZ, Oren; 10/17 Rupin Street, 75256 Rishon Lezion (IL).
- (74) Agents: PEARL COHEN ZEDEK LATZER et al.; P.O Box 12704, 4673339 Herzlia (IL).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: LONG-ACTING OXYNTOMODULIN VARIANTS AND METHODS OF PRODUCING SAME




Figure 5A

(57) Abstract: This invention is directed to a chorionic gonadotrophin carboxy terminal peptide (CTP)modified dual GLP-1/Glucagon receptor agonist, and methods of producing and using the same. In one embodiment, the present invention provides a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist.

**LONG-ACTING OXYNTOMODULIN VARIANTS AND METHODS OF  
PRODUCING SAME**

**FIELD OF INVENTION**

5 [0001] This invention is directed to a chorionic gonadotrophin carboxy terminal peptide (CTP)- modified dual GLP-1/Glucagon receptor agonist, and methods of producing and using the same.

**BACKGROUND OF THE INVENTION**

10 [0002] Oxyntomodulin (OXM) is a 37aa peptide secreted by intestinal L cells into the bloodstream upon food ingestion and induces satiety in the brain. OXM is also a dual GLP-1/Glucagon receptor agonist. It is involved in food intake regulation and was shown to inhibit food intake and reduce body weight in both rodents and humans. However, OXM has a relatively very short half-life and repeated daily administrations of supraphysiological doses  
15 are required in order to achieve pharmacological effect in humans. Hence, there is a necessity for extending the serum half-life of OXM to make it latter more amenable for prophylactic and therapeutic purposes in animal and human subjects.

[0003] The present invention makes use of peptide-based technology for extending serum half-life of proteins and peptides, in particular, OXM. This technology is based on a natural  
20 peptide, the C-terminal peptide (CTP) of the beta chain of hCG.

**SUMMARY OF THE INVENTION**

[0004] In one embodiment, the present invention provides a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin  
25 carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist.

[0005] In another embodiment, the present invention provides a method of producing a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or  
30 carboxy terminus of said agonist, the method comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist.

[0006] In one embodiment, the present invention provides a method of extending the biological half life of a dual GLP-1/Glucagon receptor agonist, comprising the step of

attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of said agonist, thereby improving the biological half life of said agonist.

[0007] In another embodiment, the present invention provides a method of improving the area under the curve (AUC) of a dual GLP-1/Glucagon receptor agonist, comprising the step of  
5 attaching at least one chorionic gonadotrophin carboxy terminal peptides to a carboxy terminus of said agonist, thereby improving the area under the curve (AUC) of said agonist.

[0008] In one embodiment, the present invention provides a method of reducing the dosing frequency of a GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of said  
10 agonist, thereby reducing the dosing frequency of said agonist.

[0009] In another embodiment, the present invention provides a use of a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of said agonist for inducing glucose tolerance in a subject.

[00010] In one embodiment, the present invention provides a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist for preventing undesired weight gain by a subject. In another embodiment, the risk of gaining weight gain is due to a psychological condition, or due to a  
20 genetic predisposition to gain weight by the subject. In another embodiment, the psychological condition is depression, anxiety or post-traumatic stress disorder (PTSD).

[00011] In another embodiment, the present invention provides a use of a composition comprising CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino  
25 terminus or carboxy terminus of said agonist for preventing, reducing or suppressing food intake in a subject.

[00012] In one embodiment, the present invention provides a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus  
30 of said agonist for treating obesity in a subject.

[00013] In another embodiment, the present invention provides a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating type II diabetes in a subject.

[00014] In one embodiment, the present invention provides a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating a metabolic disorder in a subject.

5 [00015] Other features and advantages of the present invention will become apparent from the following detailed description examples and figures. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art  
10 from this detailed description.

### BRIEF DESCRIPTION OF THE DRAWINGS

[00016] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present disclosure, the inventions of which can be  
15 better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[00017] **Figure 1** shows PAGE and Western Blot (WB) analysis of purified OXM-CTP variants: OXM-CTP-CTP-CTP, OXM-CTP 4X, OXM-CTP 5X. A. Coomassie staining of OXM samples and OXM-CTP variants (10 and 2 µg protein/lane). B. WB analysis of OXM-  
20 CTP variants using anti OXM.

[00018] **Figure 2** shows PAGE analysis of samples from purification process of OXM-CTP variants: CTP-OXM-CTP, CTP-OXM-CTP-CTP, OXM-CTP-CTP and CTP-CTP-OXM.

[00019] **Figure 3** shows PK profiles of three sequential experiments of OXM peptide and OXM-CTP variants in SD-1 rats.

25 [00020] **Figure 4** shows the results of glucose tolerance test of OXM peptide and OXM-CTP variants as measured in C57BL/6 mice.

[00021] **Figure 5** shows the results of acute food intake of OXM peptide and OXM-CTP variants as measured in C57BL/6 mice. A. The percentage of accumulated food intake in mice receiving treatment compared to vehicle. B. Percentage of food intake for various time  
30 intervals over the study.

**DETAILED DESCRIPTION OF THE INVENTION**

[00022] In one embodiment, provided herein is a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist.

5 [00023] In another embodiment, the agonist is a protein, a polypeptide, or a peptide. In another embodiment, the peptide is an oxyntomodulin.

[00024] In another embodiment, the CTP-modified polypeptide comprises a peptide that comprises fewer than 50 amino acids and at least one chorionic gonadotrophin carboxy terminal peptide, attached to an amino or a carboxy terminus of the peptide. In another  
10 embodiment, the peptide is oxyntomodulin (OXM).

[00025] Oxyntomodulin peptide is useful for the treatment of metabolic disorders such as diabetes and obesity. However, due to the short half-life of the peptide and its low stability in-vivo, repeated daily administrations of supraphysiological doses are required in order to achieve pharmacological effect in humans. As demonstrated hereinbelow (see Examples), all  
15 OXM-CTP variants of the invention demonstrated superior pharmacokinetic profile in rats as compared to native OXM, with a substantial increase in the exposure and elongated half-life. Surprisingly, the OXM-CTP-CTP variant demonstrated superior PK parameters as compared to the CTP-OXM-CTP-CTP variant which is fused to 3 copies of CTP, one in the N-terminus and two in tandem in the C-terminus.

20 [00026] Fusion of 2 and 3 CTPs to the C-terminus of OXM resulted in similar fold of increase compared to the native peptide half life (21.6 and 21 respectively), following SC administration (see Example 3 herein). Therefore, it was expected that fusion of four and five CTP to the C-terminus of OXM would not significantly elongate the half life above 20-fold. However, a surprisingly marked increase in OXM half life (i.e. 50-fold) was demonstrated for  
25 the OXM-4CTP variants and the OXM-5CTP variant with 30-fold increase in the exposure as reflected by the area under the curve (AUC) parameter (see Example 3 herein).

[00027] Hence, in one embodiment, provided herein is a CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the  
30 oxyntomodulin peptide.

[00028] In one embodiment, a CTP-modified OXM as described herein comprises a full length OXM or an active fragment thereof connected via a peptide bond on its amino or carboxy terminus to at least one CTP unit with no CTPs on its amino terminus. In another embodiment, a CTP-modified OXM as described herein comprises a full length OXM or an

active fragment thereof connected via a peptide bond on its carboxy terminus to at least one CTP unit with no CTPs on its amino terminus. In another embodiment, a CTP-modified OXM as described herein comprises a full length OXM or an active fragment thereof connected via a peptide bond on its amino terminus to at least one CTP unit with no CTPs on its carboxy terminus. In another embodiment, the present invention provides a nucleic acid molecule encoding an engineered OXM as described herein above, which in one embodiment, comprises at least one CTP attached to its carboxy terminus or its amino terminus.

[00029] In another embodiment, the CTP sequence comprises: DPRFQDSSSSKAPPPSLPSPSRLPGPSDTPIL (SEQ ID NO: 1). In another embodiment, the CTP sequence comprises: SSSSKAPPPSLPSPSRLPGPSDTPILPQ (SEQ ID NO: 2). In another embodiment, the CTP sequence comprises an amino acid sequence selected from the sequences set forth in SEQ ID NO: 1 and SEQ ID NO: 2.

[00030] In one embodiment, the carboxy terminal peptide (CTP) peptide of the present invention comprises the amino acid sequence from amino acid 112 to position 145 of human chorionic gonadotrophin. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 118 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence also commences from any position between positions 112-118 and terminates at position 145 of human chorionic gonadotrophin. In some embodiments, the CTP sequence peptide is 28, 29, 30, 31, 32, 33 or 34 amino acids long and commences at position 112, 113, 114, 115, 116, 117 or 118 of the CTP amino acid sequence.

[00031] Thus, in one embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 112 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 113 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 114 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 115 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 116 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2. In another embodiment, the CTP sequence of the present

invention comprises the amino acid sequence from amino acid 117 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 2.

[00032] In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 112 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 113 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 114 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 115 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 116 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 117 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1. In another embodiment, the CTP sequence of the present invention comprises the amino acid sequence from amino acid 118 to position 145 of human chorionic gonadotropin, as set forth in SEQ ID NO: 1.

[00033] In one embodiment, the truncated CTP comprises SSSSKAPPPSLP (SEQ ID NO: 3).

In another embodiment, the truncated CTP comprises the first 10 amino acids of SEQ ID NO: 3. In another embodiment, the truncated CTP comprises the first 11 amino acids of SEQ ID NO: 3.

[00034] In one embodiment, the truncated CTP comprises the first 14 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 13 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 12 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 11 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 10 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 9 amino acids of SEQ ID NO: 2. In one embodiment, the truncated CTP comprises the first 8 amino acids of SEQ ID NO: 2 or SEQ ID NO: 3. In one embodiment, the truncated CTP comprises the first 7 amino acids of SEQ ID NO: 2 or SEQ ID NO: 3. In one embodiment, the truncated CTP comprises the first 6 amino acids of SEQ ID NO: 2 or SEQ ID NO: 3. In one embodiment, the truncated CTP comprises the first 5 amino acids of SEQ ID NO: 2 or SEQ ID NO: 3.

[00035] In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 1-5 conservative amino acid substitutions as described in U.S. Pat. No. 5,712,122, which is incorporated herein by reference in its entirety. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 1 conservative amino acid substitution. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 2 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 3 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 4 conservative amino acid substitutions. In another embodiment, the CTP peptide is a variant of chorionic gonadotrophin CTP which differs from the native CTP by 5 conservative amino acid substitutions.

[00036] In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 70% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 80% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 85% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 90% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 95% homologous to the native CTP amino acid sequence or a peptide thereof. In another embodiment, the CTP peptide amino acid sequence of the present invention is at least 98% homologous to the native CTP amino acid sequence or a peptide thereof.

[00037] In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 70% homologous to the native human CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 80% homologous to the native human CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 85% homologous to the native human CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 90% homologous to the native CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention

is at least 95% homologous to the native CTP DNA sequence or a peptide thereof. In another embodiment, the polynucleotide encoding the CTP peptide of the present invention is at least 98% homologous to the native CTP DNA sequence or a peptide thereof.

[00038] In one embodiment, at least one of the chorionic gonadotrophin CTP amino acid sequences is truncated. In another embodiment, both of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, two of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, three of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, four of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, five of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, two or more of the chorionic gonadotrophin CTP amino acid sequences are truncated. In another embodiment, all of the chorionic gonadotrophin CTP amino acid sequences are truncated.

[00039] In one embodiment, at least one CTP is attached to the agonist polypeptide via a linker. In another embodiment, at least one CTP is attached to the agonist polypeptide via a linker. In another embodiment, the linker is a peptide bond.

[00040] In one embodiment, at least one of the chorionic gonadotrophin CTP amino acid sequences is glycosylated. In another embodiment, both of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, two of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, three of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, four of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, five of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, two or more of the chorionic gonadotrophin CTP amino acid sequences are glycosylated. In another embodiment, all of the chorionic gonadotrophin CTP amino acid sequences are glycosylated.

[00041] In one embodiment, the CTP sequence of the present invention comprises at least one glycosylation site. In one embodiment, the CTP sequence of the present invention comprises two glycosylation sites. In one embodiment, the CTP sequence of the present invention comprises three glycosylation sites. In one embodiment, the CTP sequence of the present invention comprises four glycosylation sites. In one embodiment, one or more of the chorionic gonadotrophin CTP amino acid sequences is fully glycosylated. In another embodiment, one or more of the chorionic gonadotrophin CTP amino acid sequences is partially glycosylated. In one embodiment, partially glycosylated indicates that one of the

CTP glycosylation sites is glycosylated. In another embodiment, two of the CTP glycosylation sites are glycosylated. In another embodiment, three of the CTP glycosylation sites are glycosylated.

[00042] In some embodiments, the CTP sequence modification is advantageous in permitting the usage of lower dosages. In some embodiments, the CTP sequences modification is advantageous in permitting fewer dosages. In some embodiments, the CTP sequences modification is advantageous in permitting a safe, long-acting effect.

[00043] In some embodiments, "polypeptide", "engineered oxyntomodulin", or "protein" as used herein encompasses native polypeptides (either degradation products, synthetically synthesized polypeptides or recombinant polypeptides) and peptidomimetics (typically, synthetically synthesized polypeptides), as well as peptoids and semipeptoids which are polypeptide analogs, which have, in some embodiments, modifications rendering the polypeptides comprising a oxyntomodulin even more stable while in a human or animal subject or more capable of penetrating into cells.

[00044] In some embodiments, modifications include, but are limited to C terminus modification, polypeptide bond modification, including, but not limited to, CH<sub>2</sub>-NH, CH<sub>2</sub>-S, CH<sub>2</sub>-S=O, O=C-NH, CH<sub>2</sub>-O, CH<sub>2</sub>-CH<sub>2</sub>, S=C-NH, CH=CH or CF=CH, backbone modifications, and residue modification. Methods for preparing peptidomimetic compounds are well known in the art and are specified, for example, in Quantitative Drug Design, C.A. Ramsden Gd., Chapter 17.2, F. Choplin Pergamon Press (1992), which is incorporated by reference as if fully set forth herein. Further details in this respect are provided hereinunder.

[00045] In some embodiments, polypeptide bonds (-CO-NH-) within the polypeptide are substituted. In some embodiments, the polypeptide bonds are substituted by N-methylated bonds (-N(CH<sub>3</sub>)-CO-). In some embodiments, the polypeptide bonds are substituted by ester bonds (-C(R)H-C-O-O-C(R)-N-). In some embodiments, the polypeptide bonds are substituted by ketomethylen bonds (-CO-CH<sub>2</sub>-). In some embodiments, the polypeptide bonds are substituted by  $\alpha$ -aza bonds (-NH-N(R)-CO-), wherein R is any alkyl, e.g., methyl, carba bonds (-CH<sub>2</sub>-NH-). In some embodiments, the polypeptide bonds are substituted by hydroxyethylene bonds (-CH(OH)-CH<sub>2</sub>-). In some embodiments, the polypeptide bonds are substituted by thioamide bonds (-CS-NH-). In some embodiments, the polypeptide bonds are substituted by olefinic double bonds (-CH=CH-). In some embodiments, the polypeptide bonds are substituted by retro amide bonds (-NH-CO-). In some embodiments, the polypeptide bonds are substituted by polypeptide derivatives (-N(R)-CH<sub>2</sub>-CO-), wherein R is the "normal" side chain, naturally presented on the carbon atom. In some embodiments, these

modifications occur at any of the bonds along the polypeptide chain and in one embodiment at several (2-3 bonds) at the same time.

[00046] In one embodiment, natural aromatic amino acids of the polypeptide such as Trp, Tyr and Phe, are substituted for synthetic non-natural acid such as Phenylglycine, TIC, naphthylelanine (Nol), ring-methylated derivatives of Phe, halogenated derivatives of Phe or o-methyl-Tyr. In some embodiments, the polypeptides of the present invention include one or more modified amino acid or one or more non-amino acid monomers (e.g. fatty acid, complex carbohydrates etc).

[00047] In one embodiment, "amino acid" or "amino acid sequence" is understood to include the 20 naturally occurring amino acid; those amino acid often modified post-translationally in vivo, including, for example, hydroxyproline, phosphoserine and phosphothreonine; and other unusual amino acid including, but not limited to, 2-amino adipic acid, hydroxylysine, isodesmosine, nor-valine, nor-leucine and ornithine. In one embodiment, "amino acid" includes both D- and L-amino acids.

[00048] In one embodiment, the polypeptides of the present invention are utilized in therapeutics which requires the polypeptides comprising an oxyntomodulin to be in a soluble form. In some embodiments, the polypeptides of the present invention include one or more non-natural or natural polar amino acid, including but not limited to serine and threonine which are capable of increasing polypeptide solubility due to their hydroxyl-containing side chain.

[00049] In another embodiment, the engineered oxyntomodulin peptides of the present invention are biochemically synthesized such as by using standard solid phase techniques. In some embodiments, these biochemical methods include exclusive solid phase synthesis, partial solid phase synthesis, fragment condensation, or classical solution synthesis.

[00050] In one embodiment, recombinant protein techniques are used to generate the engineered oxyntomodulin peptides of the present invention. In some embodiments, recombinant protein techniques are used for the generation of relatively long polypeptides (e.g., longer than 18-25 amino acids). In some embodiments, recombinant protein techniques are used for the generation of large amounts of the engineered oxyntomodulin peptides of the present invention. In some embodiments, recombinant techniques are described by Bitter et al., (1987) *Methods in Enzymol.* 153:516-544, Studier et al. (1990) *Methods in Enzymol.* 185:60-89, Brisson et al. (1984) *Nature* 310:511-514, Takamatsu et al. (1987) *EMBO J.* 6:307-311, Coruzzi et al. (1984) *EMBO J.* 3:1671-1680 and Brogli et al., (1984) *Science* 224:838-843, Gurley et al. (1986) *Mol. Cell. Biol.* 6:559-565 and Weissbach & Weissbach,

1988, Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp 421-463, which are incorporated herein by reference in their entirety.

[00051] In another embodiment, the CTP-modified polypeptide comprises a peptide that comprises fewer than 50 amino acids and at least one chorionic gonadotrophin carboxy terminal peptide, attached to an amino or a carboxy terminus of the peptide. In one  
5 embodiment, the CTP-modified polypeptide provided herein comprises a peptide that comprises fewer than 40 amino acids and at least one chorionic gonadotrophin carboxy terminal peptide, attached to an amino or a carboxy terminus of the peptide. In another embodiment, the CTP-modified polypeptide provided herein comprises a peptide that  
10 comprises fewer than 30, 20, or 10 amino acids. In one embodiment, the peptide comprising fewer than 50 amino acids is a GLP-1/Glucagon receptor agonist. In another embodiment, the peptide comprising fewer than 50 amino acids is OXM.

[00052] In another embodiment, OXM comprises the following amino acid (AA) sequence: HSQGTFTSDYSKYLDSSRAQDFVQWLMNTKRNRNNIA (SEQ ID NO: 30). In another  
15 embodiment, OXM consists of the amino acid sequence of SEQ ID NO: 30. In another embodiment, OXM comprises or consists of the amino acid sequence depicted in CAS No. 62340-29-8.

[00053] In another embodiment, OXM is human OXM or any mammal OXM. In another embodiment, OXM is also referred to as glucagon-37 or bioactive enteroglucagon. In another  
20 embodiment, OXM is a dual GLP-1/Glucagon receptor agonist. In another embodiment, the term OXM includes a biologically active fragment of OXM. In another embodiment, biologically active OXM extends from amino acid 30 to amino acid 37 of SEQ ID NO: 30. In another embodiment, biologically active OXM extends from amino acid 19 to amino acid 37 of SEQ ID NO: 30. In another embodiment, OXM of the invention corresponds to an  
25 octapeptide from which the two C-terminal amino acids are deleted. In another embodiment, OXM of the invention corresponds to any fragment of SEQ ID NO: 30 which retains OXM activity as described herein.

[00054] In one embodiment, two chorionic gonadotrophin carboxy terminal peptides are attached to oxyntomodulin, one CTP on the carboxy terminus and one CTP on the amino  
30 terminus of the oxyntomodulin peptide. In another embodiment, two chorionic gonadotrophin carboxy terminal peptides are attached to oxyntomodulin on the carboxy terminus of the oxyntomodulin peptide. In another embodiment, two chorionic gonadotrophin carboxy terminal peptides are attached to oxyntomodulin, both on the amino terminus of the

oxyntomodulin peptide. In another embodiment, three chorionic gonadotrophin carboxy terminal peptides are attached to oxyntomodulin, one CTP on the amino terminus and two CTPs on the carboxy terminus of the oxyntomodulin peptide. In another embodiment, three chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of the oxyntomodulin peptide. In another embodiment, four chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of the oxyntomodulin peptide. In another embodiment, five chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of the oxyntomodulin peptide. In another embodiment, 1-10 CTP are attached to the amino or carboxy terminus of oxyntomodulin. In another embodiment, 1-10 CTPs are attached to the amino terminus of oxyntomodulin. In another embodiment, 1-10 CTPs are attached to the carboxy terminus of oxyntomodulin.

[00055] In another embodiment, provided herein is a method of producing a CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, the method comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide.

[00056] In one embodiment, provided herein is a method of extending the biological half life of a peptide comprising fewer than 50 amino acids, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the agonist, thereby improving the biological half life of the agonist.

[00057] In one embodiment, provided herein is a method of extending the biological half life of a dual GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the agonist, thereby improving the biological half life of the agonist.

[00058] In another embodiment, provided herein is a method of extending the biological half life of oxyntomodulin, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the oxyntomodulin, thereby improving the biological half life of oxyntomodulin.

[00059] In one embodiment, provided herein is a method of improving the area under the curve (AUC) of a peptide comprising fewer than 50 amino acids, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to a carboxy terminus of the agonist, thereby improving the area under the curve (AUC) of the agonist.

[00060] In one embodiment, provided herein is a method of improving the area under the curve (AUC) of a dual GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to a carboxy terminus of the agonist, thereby improving the area under the curve (AUC) of the agonist.

5 [00061] In another embodiment, provided herein is a method of improving the area under the curve (AUC) of oxyntomodulin, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to a carboxy terminus of the oxyntomodulin, thereby improving the area under the curve (AUC) of oxyntomodulin.

[00062] In one embodiment, the term oxyntomodulin further includes a homologue of a  
10 known oxyntomodulin. In one embodiment, the homologue is a functional homologue. In another embodiment, the term "functional" refers to the ability a homologue, peptide or protein provided herein has to suppress appetite. The term also refers to the ability a homologue, peptide or protein provided herein has to extend another protein's or peptide's biological half-life. In another embodiment, the biological half-life ( $T_{1/2}$ ) of a protein, peptide  
15 or homologue provided herein refers to the time it takes for half of the amount of the protein, peptide or homologue to be degraded or to not be present in a biological medium in a subject. In another embodiment, the biological medium is serum, cerebrospinal fluid, tissue, mucosa, and the like.

[00063] In one embodiment, homology according to the present invention also encompasses  
20 deletion, insertion, or substitution variants, including an amino acid substitution, thereof and biologically active polypeptide fragments thereof. In one embodiment, the variant provided herein comprises conservative substitutions, or deletions, insertions, or substitutions that do not significantly alter the three dimensional structure of the oxyntomodulin. In another embodiment, the deletion, insertion, or substitution does not alter the function of interest of  
25 the oxyntomodulin, which in one embodiment, is binding to a particular binding partner.

[00064] In one embodiment, oxyntomodulin binds to a receptor and mediates appetite suppression. In another embodiment, the receptor is a dual GLP-1/Glucagon receptor. In another embodiment, the receptor is a GLP-1 receptor. In another embodiment, the receptor is a glucagon receptor. In yet another embodiment, the receptor is any receptor known in the art  
30 to bind to oxyntomodulin.

[00065] In another embodiment, the invention includes a homologue of an oxyntomodulin. In one embodiment, the invention provides an OXM homologue, which in one embodiment, refers to a peptide homologue of the peptide of SEQ ID NO: 30.

[00066] In another embodiment, the invention includes a homologue of an oxyntomodulin having an appetite suppressing activity. In another embodiment, the invention includes a homologue of an oxyntomodulin having functional binding. In another embodiment, the invention includes homologues of an oxyntomodulin as described herein having an appetite suppression and activity. In another embodiment, the invention includes homologues of an oxyntomodulin as described herein having functional binding. In another embodiment, the invention also encompasses homologues e.g., polypeptides which are at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 87%, at least 89%, at least 91%, at least 93%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% homologous to an oxyntomodulin as determined using BlastP software of the National Center of Biotechnology Information (NCBI) using default parameters.

[00067] In one embodiment, provided herein is a pharmaceutical composition comprising the CTP-modified polypeptide provided herein.

[00068] It is to be understood that the compositions and methods of the present invention comprising the elements or steps as described herein may, in another embodiment, consist of those elements or steps, or in another embodiment, consist essentially of those elements or steps. In some embodiments, the term “comprise” refers to the inclusion of the indicated active agent, such as the CTP-modified GLP-1/Glucagon receptor agonist, as well as inclusion of other active agents, and pharmaceutically or physiologically acceptable carriers, excipients, emollients, stabilizers, etc., as are known in the pharmaceutical industry. In some embodiments, the term “consisting essentially of” refers to a composition, whose only active ingredient is the indicated active ingredient, however, other compounds may be included which are for stabilizing, preserving, etc. the formulation, but are not involved directly in the therapeutic effect of the indicated active ingredient. In some embodiments, the term “consisting essentially of” may refer to components which facilitate the release of the active ingredient. In some embodiments, the term “consisting” refers to a composition, which contains the active ingredient and a pharmaceutically acceptable carrier or excipient.

[00069] It is to be understood that the compositions and methods of the present invention comprising the elements or steps as described herein may, in another embodiment, consist of those elements or steps, or in another embodiment, consist essentially of those elements or steps. In some embodiments, the term “comprise” refers to the inclusion of the indicated active agent, such as the CTP-modified oxyntomodulin, as well as inclusion of other active agents, and pharmaceutically or physiologically acceptable carriers, excipients, emollients, stabilizers, etc., as are known in the pharmaceutical industry. In some embodiments, the term

“consisting essentially of” refers to a composition, whose only active ingredient is the indicated active ingredient, however, other compounds may be included which are for stabilizing, preserving, etc. the formulation, but are not involved directly in the therapeutic effect of the indicated active ingredient. In some embodiments, the term “consisting essentially of” may refer to components which facilitate the release of the active ingredient. In some embodiments, the term “consisting” refers to a composition, which contains the active ingredient and a pharmaceutically acceptable carrier or excipient.

[00070] In one embodiment, the present invention provides a polypeptide comprising a GLP-1/Glucagon receptor agonist and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to three CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and two to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00071] In one embodiment, the present invention provides a polypeptide comprising an oxyntomodulin and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to three CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to four CTPs attached to the carboxy terminus of the

oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and two to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00072] In one embodiment, the present invention provides a polypeptide comprising a GLP-1/Glucagon receptor agonist and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and three to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00073] In one embodiment, the present invention provides a polypeptide comprising an oxyntomodulin and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a

polypeptide comprising an oxyntomodulin and three to four CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and three to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00074] In one embodiment, the present invention provides a polypeptide comprising a GLP-1/Glucagon receptor agonist and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and four to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00075] In one embodiment, the present invention provides a polypeptide comprising an oxyntomodulin and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to five CTPs attached to the carboxy

terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and four to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00076] In one embodiment, the present invention provides a polypeptide comprising a GLP-1/Glucagon receptor agonist and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and five to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and five to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and five to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and five to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide comprising a GLP-1/Glucagon receptor agonist and five to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00077] In one embodiment, the present invention provides a polypeptide comprising an oxyntomodulin and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and five to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and five to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and five to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an

oxyntomodulin and five to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide comprising an oxyntomodulin and five to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00078] In one embodiment, the present invention provides a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to three CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and two to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00079] In one embodiment, the present invention provides a polypeptide consisting of an oxyntomodulin and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to three CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to four CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and

two to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to eight CTPs  
5 attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and two to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

10 [00080] In one embodiment, the present invention provides a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon  
15 receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment,  
20 provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide  
25 consisting of a GLP-1/Glucagon receptor agonist and three to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and three to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

30 [00081] In one embodiment, the present invention provides a polypeptide consisting of an oxyntomodulin and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to four CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to five CTPs attached to the carboxy terminus of

the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and three to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00082] In one embodiment, the present invention provides a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and four to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00083] In one embodiment, the present invention provides a polypeptide consisting of an oxyntomodulin and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

oxyntomodulin and four to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and four to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00084] In one embodiment, the present invention provides a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting of a GLP-1/Glucagon receptor agonist and five to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00085] In one embodiment, the present invention provides a polypeptide consisting of an oxyntomodulin and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and five to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and five to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and five to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and five to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting of an oxyntomodulin and five to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00086] In one embodiment, the present invention provides a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to three CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and two to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00087] In one embodiment, the present invention provides a polypeptide consisting essentially of an oxyntomodulin and two gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to three CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to four CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to seven CTPs

attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and two to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00088] In one embodiment, the present invention provides a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to four CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and three to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00089] In one embodiment, the present invention provides a polypeptide consisting essentially of an oxyntomodulin and three gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and three to four CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and three to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided

herein is a polypeptide consisting essentially of an oxyntomodulin and three to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and three to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided  
5 herein is a polypeptide consisting essentially of an oxyntomodulin and three to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and three to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided  
10 herein is a polypeptide consisting essentially of an oxyntomodulin and three to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00090] In one embodiment, the present invention provides a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-  
15 1/Glucagon receptor agonist and four to five CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and four to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist  
20 and four to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and four to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and four to nine  
25 CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and four to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00091] In one embodiment, the present invention provides a polypeptide consisting  
30 essentially of an oxyntomodulin and four gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to five CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to six CTPs

attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to nine CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and four to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00092] In one embodiment, the present invention provides a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five to six CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five to seven CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five to eight CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five to nine CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist. In another embodiment, provided herein is a polypeptide consisting essentially of a GLP-1/Glucagon receptor agonist and five to ten CTPs attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist.

[00093] In one embodiment, the present invention provides a polypeptide consisting essentially of an oxyntomodulin and five gonadotrophin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and five to six CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and five to seven CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and five to eight CTPs attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and five to nine CTPs

attached to the carboxy terminus of the oxyntomodulin. In another embodiment, provided herein is a polypeptide consisting essentially of an oxyntomodulin and five to ten CTPs attached to the carboxy terminus of the oxyntomodulin.

[00094] In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a GLP-1/Glucagon receptor agonist having no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a GLP-1/Glucagon receptor agonist lacking a CTP on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a GLP-1/Glucagon receptor agonist having at least one CTP on its carboxy terminus and no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of a GLP-1/Glucagon receptor agonist having the number of CTPs on its carboxy terminus as described herein and no CTPs on its amino terminus.

[00095] In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of an oxyntomodulin having no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of an oxyntomodulin lacking a CTP on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of an oxyntomodulin having at least one CTP on its carboxy terminus and no CTPs on its amino terminus. In another embodiment, provided herein is a polypeptide comprising, consisting essentially of, or consisting of an oxyntomodulin having the number of CTPs on its carboxy terminus as described herein and no CTPs on its amino terminus.

[00096] In one embodiment, the amino acid sequence of the CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29, further provided herein. In another embodiment, the amino acid sequence of the CTP-modified polypeptide is selected from the group consisting of: SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29, further provided herein.

[00097] In another embodiment, the present invention provides a polynucleotide encoding a polypeptide as described hereinabove. In another embodiment, the polynucleotide comprises SEQ ID NO: 8, 14, 20, 22, 24, 26 or 28, further provided herein. In another embodiment, the polynucleotide is selected from the group consisting of SEQ ID NO: 98, 14, 20, 22, 24, 26 or 28, further provided herein.

[00098] In one embodiment, provided herein is an expression vector comprising the polynucleotide provided herein. In another embodiment, provided herein is a cell comprising

the expression vector. In another embodiment, provided herein is a composition comprising the expression vector.

[00099] In another embodiment, the invention provides a cell comprising the expression vector as described herein. In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide comprising or consisting of a GLP-1/Glucagon receptor agonist and at least one gonadotropin carboxy terminal peptides (CTPs) attached to the amino or carboxy terminus of the agonist. In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of a GLP-1/Glucagon receptor agonist and at least one gonadotropin carboxy terminal peptides (CTPs) attached to the amino or carboxy terminus of the agonist.

[000100] In another embodiment, the invention provides a composition comprising the expression vector as described herein. In another embodiment, the present invention provides a composition comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide comprising or consisting of a GLP-1/Glucagon receptor agonist and at least one gonadotropin carboxy terminal peptides (CTPs) attached to the amino or carboxy terminus of the agonist. In another embodiment, the present invention provides a cell comprising an expression vector comprising a polynucleotide encoding a CTP-modified polypeptide consisting of the agonist and at least one gonadotropin carboxy terminal peptides (CTPs) attached to the amino or carboxy terminus of the agonist. In another embodiment, the agonist is a polypeptide, or a peptide. In another embodiment, the peptide is oxyntomodulin.

[000101] In another embodiment, the invention provides a composition comprising the cell as described herein. In another embodiment, the cell is a eukaryotic cell. In another embodiment, the cell is a prokaryotic cell.

[000102] In one embodiment, the present invention provides a GLP-1/Glucagon receptor agonist. In another embodiment, the present invention provides a recombinant oxyntomodulin as described hereinabove. In one embodiment, the present invention provides an engineered oxyntomodulin as described hereinabove. In one embodiment, the engineered oxyntomodulin as described hereinabove is referred to as a CTP-modified oxyntomodulin.

[000103] In one embodiment, the CTPs that are attached to the carboxy terminus of the oxyntomodulin are attached in tandem to the carboxy terminus. In one embodiment, the CTPs that are attached to the amino terminus of the oxyntomodulin are attached in tandem to the amino terminus.

[000104] In one embodiment, an engineered GLP-1/Glucagon receptor agonist as described herein has equivalent or improved biological activity compared to the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, an engineered GLP-1/Glucagon receptor agonist as provided herein has equivalent or improved pharmacological measurements compared to the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, an engineered GLP-1/Glucagon receptor agonist as provided herein has equivalent or improved pharmacokinetics compared to the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, an engineered GLP-1/Glucagon receptor agonist as provided herein has equivalent or improved pharmacodynamics compared to the non-CTP-modified GLP-1/Glucagon receptor agonist.

[000105] In another embodiment, the terms “CTP peptide,” “carboxy terminal peptide” and “CTP sequence” are used interchangeably herein. In another embodiment, the carboxy terminal peptide is a full-length CTP. Each possibility represents a separate embodiment of the invention.

[000106] In another embodiment, a signal peptide is attached to the amino terminus of the CTP, as described in US 7,553,940, which is incorporated by reference herein in its entirety.

[000107] In other embodiments, the term engineered oxyntomodulin refers to the amino acid sequence of a matured oxyntomodulin. In other embodiments, the term engineered oxyntomodulin refers to the amino acid sequence of the oxyntomodulin including its signal sequence or signal peptide.

[000108] In another embodiment, “signal sequence” and “signal peptide” are used interchangeably herein. In another embodiment, “sequence” when in reference to a polynucleotide molecule can refer to a coding portion. Each possibility represents a separate embodiment of the present invention.

[000109] In another embodiment, an engineered GLP-1/Glucagon receptor agonist comprising at least one CTP as described herein has enhanced in vivo biological activity compared the same GLP-1/Glucagon receptor agonist without at least one CTP. In one embodiment, the enhanced biological activity stems from the longer half-life of the engineered GLP-1/Glucagon receptor agonist while maintaining at least some biological activity. In another embodiment, the enhanced biological activity stems from enhanced biological activity resulting from the CTP modification. In another embodiment, the enhanced biological activity stems from both a longer half life and from enhanced functionality of the CTP-modified GLP-1/Glucagon receptor agonist.

[000110] In one embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist provides enhanced protection against degradation of a GLP-1/Glucagon receptor agonist. In another embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist provides enhanced protection against clearance. In another embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist provides prolonged clearance time. In another embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist enhances its C<sub>max</sub>. In another embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist enhances its T<sub>max</sub>. In another embodiment, the at least one CTP sequence at the carboxy terminal end of the GLP-1/Glucagon receptor agonist prolongs its T<sub>1/2</sub>.

[000111] In another embodiment, a conjugated GLP-1/Glucagon receptor agonist of this invention is used in the same manner as an unmodified conjugated GLP-1/Glucagon receptor agonist. In another embodiment, a conjugated oxyntomodulin of this invention is used in the same manner as an unmodified conjugated oxyntomodulin. In another embodiment, a conjugated oxyntomodulin of this invention has an increased circulating half-life and plasma residence time, decreased clearance, and increased clinical activity in vivo. In another embodiment, due to the improved properties of the conjugated oxyntomodulin as described herein, this conjugate is administered less frequently than the unmodified form of the same oxyntomodulin.

[000112] In another embodiment, decreased frequency of administration results in improved treatment strategy, which in one embodiment, leads to improved patient compliance leading to improved treatment outcomes, as well as improved patient quality of life. In another embodiment, compared to conventional conjugates of oxyntomodulins, it has been found that conjugates having the molecular weight and linker structure of the conjugates as described herein have an improved potency, improved stability, elevated AUC levels, and enhanced circulating half-life.

[000113] In another embodiment, the present invention further provides a pharmaceutical composition comprising a CTP-modified polypeptide consisting of a GLP-1/Glucagon receptor agonist and three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin.

[000114] In another embodiment, the present invention further provides a pharmaceutical composition comprising a CTP-modified polypeptide consisting of an oxyntomodulin and

three gonadotropin carboxy terminal peptides (CTPs) attached to the carboxy terminus of the oxyntomodulin.

[000115] In another embodiment, provided herein is a composition comprising a conjugated GLP-1/Glucagon receptor agonist as provided herein. In another embodiment, provided  
5 herein is a pharmaceutical composition comprising the conjugated GLP-1/Glucagon receptor agonist as provided herein. In another embodiment, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of the conjugated GLP-1/Glucagon receptor agonist as provided herein. In one embodiment, a therapeutically effective amount of a conjugated GLP-1/Glucagon receptor agonist is determined according  
10 to factors such as the specific condition being treated, the condition of the patient being treated, as well as the other ingredients in the composition.

[000116] In another embodiment, provided herein is a composition comprising a conjugated oxyntomodulin as provided herein. In another embodiment, provided herein is a pharmaceutical composition comprising the conjugated oxyntomodulin as provided herein. In  
15 another embodiment, provided herein is a pharmaceutical composition comprising a therapeutically effective amount of the conjugated oxyntomodulin as provided herein. In one embodiment, a therapeutically effective amount of a conjugated oxyntomodulin is determined according to factors such as the specific condition being treated, the condition of the patient being treated, as well as the other ingredients in the composition.

[000117] In one embodiment, a CTP-modified GLP-1/Glucagon receptor agonist of the  
20 present invention has therapeutic uses. In another embodiment, a CTP-modified GLP-1/Glucagon receptor agonist of the present invention has prophylactic uses.

[000118] In another embodiment, a CTP-modified oxyntomodulin of the present invention has therapeutic uses. In another embodiment, a CTP-modified oxyntomodulin of the present  
25 invention has prophylactic uses.

[000119] In one embodiment, provided herein is a method of reducing the dosing frequency of GLP-1/Glucagon receptor agonist in a subject, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the GLP-1/Glucagon receptor agonist, thereby reducing the dosing frequency of GLP-1/Glucagon  
30 receptor agonist. In another embodiment, the CTP sequence modification is advantageous in permitting the usage of lower dosages. In some embodiments, the CTP sequences modification is advantageous in permitting fewer dosages. In some embodiments, the CTP sequences modification is advantageous in permitting a safe, long-acting effect.

[000120] In another embodiment, provided herein is a method of reducing the dosing frequency of oxyntomodulin in a subject, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the oxyntomodulin, thereby reducing the dosing frequency of oxyntomodulin. In another embodiment, the CTP sequence modification is advantageous in permitting the usage of lower dosages. In some embodiments, the CTP sequences modification is advantageous in permitting fewer dosages. In some embodiments, the CTP sequences modification is advantageous in permitting a safe, long-acting effect.

[000121] In one embodiment, provided herein is a method of reducing cholesterol in a subject, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the oxyntomodulin, thereby reducing cholesterol in a subject.

[000122] In another embodiment, provided herein is a method of reducing glycerol in a subject, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the oxyntomodulin, thereby reducing glycerol in a subject.

[000123] In one embodiment, the present invention provides a long-acting OXM. In one embodiment, a long-acting OXM of the invention maintains the biological activity of unmodified OXM. In another embodiment, the long-acting OXM of the invention comprises OXM biological activity. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reducing digestive secretions. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reducing and delaying gastric emptying. In another embodiment, the biological activity of a long-acting OXM of the invention comprises the inhibition of the fed motility pattern in the small intestine. In another embodiment, the biological activity of a long-acting OXM of the invention comprises the inhibition of acid secretion stimulated by pentagastrin. In another embodiment, the biological activity of a long-acting OXM of the invention comprises an increase of gastric somatostatin release. In another embodiment, the biological activity of a long-acting OXM of the invention comprises potentiating the effects of peptide YY. In another embodiment, the biological activity of a long-acting OXM of the invention comprises the inhibition of ghrelin release. In another embodiment, the biological activity of a long-acting OXM of the invention comprises the stimulation of aminopyrine accumulation and cAMP production. In another embodiment, the biological activity of a long-acting OXM of the invention comprises binding the GLP-1 receptor. In another embodiment, the biological activity of a long-acting OXM of the

invention comprises stimulating H<sup>+</sup> production by activating adenylate cyclase. In another embodiment, the biological activity of a long-acting OXM of the invention comprises inhibiting histamine-stimulated gastric acid secretion. In another embodiment, the biological activity of a long-acting OXM of the invention comprises inhibiting food intake. In another embodiment, the biological activity of a long-acting OXM of the invention comprises stimulating insulin release. In another embodiment, the biological activity of a long-acting OXM of the invention comprises inhibiting exocrine pancreatic secretion.

[000124] In another embodiment, the biological activity of a long-acting OXM of the invention comprises inhibiting pancreatic secretion through a vagal neural indirect mechanism. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reducing hydromineral transport through the small intestine. In another embodiment, the biological activity of a long-acting OXM of the invention comprises stimulating glucose uptake. In another embodiment, the biological activity of a long-acting OXM of the invention comprises controlling/stimulating somatostatin secretion. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reduction in both food intake and body weight gain. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reduction in adiposity. In another embodiment, the biological activity of a long-acting OXM of the invention comprises appetite suppression. In another embodiment, the biological activity of a long-acting OXM of the invention comprises induction of anorexia. In another embodiment, the biological activity of a long-acting OXM of the invention comprises reducing body weight in overweight and obese subjects. In another embodiment, the biological activity of a long-acting OXM of the invention comprises inducing changes in the levels of the adipose hormones leptin and adiponectin. In another embodiment, the biological activity of a long-acting OXM of the invention comprises increasing energy expenditure in addition to decreasing energy intake in overweight and obese subjects.

[000125] Thus, in one embodiment, the present invention provides a method of effecting any of the above-mentioned biological activities of OXM by administering a CTP-modified OXM of the present invention. In another embodiment, the present invention further provides a method for reducing food intake, reducing body weight, or both in a subject, comprising the step of administering to the subject a CTP-modified oxyntomodulin peptide. In another embodiment, the subject has diabetes. In another embodiment, the subject is overweight. In another embodiment, the subject is obese.

[000126] In one embodiment, the terms “reducing, reduction, lowering, etc.” when used in relation to the methods provided herein refer to 100% reduction from a previously measured or determined level or from a normal level. In another embodiment, the reduction is by 89-99% from a previously determined level. In another embodiment, the reduction is by 79-88% from a previously determined level. In another embodiment, the reduction is by 69-78% from a previously determined level. In another embodiment, the reduction is by 59-68% from a previously determined level. In another embodiment, the reduction is by 49-58% from a previously determined level. In another embodiment, the reduction is by 39-48% from a previously determined level. In another embodiment, the reduction is by 29-38% from a previously determined level. In another embodiment, the reduction is by 19-28% from a previously determined level. In another embodiment, the reduction is by 9-18% from a previously determined level. In another embodiment, the reduction is by 5-8% from a previously determined level. In another embodiment, the reduction is by 1-4% from a previously determined level.

[000127] The *in-vivo* biological activity of OXM-CTP variants was assessed in two animal models, IPGTT in mice, which evaluate the ability of OXM to induce glucose tolerance following glucose administration, and food intake inhibition in lean rats, which assess the ability of OXM to inhibit animal’s food consumption. It was demonstrated that OXM-CTP variants: CTP-OXM-CTP-CTP, OXM-CTP-CTP, OXM-CTP-CTP-CTP, OXM-CTP-CTP-CTP-CTP and OXM-CTP-CTP-CTP-CTP-CTP induced glucose tolerance as reflected by reduction of 20-30% of blood glucose AUC compared to vehicle group (see Example 4, herein). These results indicate that the biological activity of OXM-CTP variants was not inhibited *in-vivo* due to potential steric interference of CTP fusion to the binding of OXM to its receptor. The marked reduction in glucose induced by these variants correlated with their improved PK profiles. Therefore, in one embodiment, provided herein is a method of inducing glucose tolerance in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of the oxyntomodulin peptide, thereby inducing glucose tolerance in a subject.

[000128] In another embodiment, provided herein is a method of inducing glucose tolerance in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP)

attached to the amino terminus or caboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby inducing glucose tolerance in a subject.

[000129] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of said agonist for inducing glucose tolerance in a subject. In one embodiment, said dual GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000130] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of said agonist in the preparation of a medicament for inducing glucose tolerance in a subject. In one embodiment, said dual GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000131] In one embodiment, provided herein is a method of increasing insulin sensitivity in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby increasing insulin sensitivity in a subject.

[000132] In one embodiment, provided herein is a method of reducing insulin resistance in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby reducing insulin resistance in a subject.

[000133] In one embodiment, provided herein is a method of increasing insulin sensitivity and reducing insulin resistance in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby increasing insulin sensitivity and reducing insulin resistance in a subject.

[000134] The ability of OXM-CTP variants OXM-3CTP, OXM-4CTP and OXM-5CTP to inhibit food intake was assessed. Native OXM inhibited food intake only in the first hour following administration, however, all variants manifested substantial prolonged improved

inhibition in food intake compared to OXM. The cumulative food inhibition of OXM-5CTP was surprisingly sustained for at least 141 hrs emphasizing the elongated half life of this variant (see Example 5, herein).

[000135] Hence, in one embodiment, provided herein is a method of inducing food intake inhibition in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, thereby inducing food intake inhibition in the subject.

[000136] In another embodiment, provided herein is a method of inducing food intake inhibition in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby inducing food intake inhibition in the subject.

[000137] In another embodiment, provided herein is a method of preventing, reducing or suppressing food intake in a subject, the method comprising the step of administering to the subject an effective dose of a composition comprising CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby preventing reducing or suppressing food intake by a subject. In another embodiment, preventing, reducing or suppressing food intake by a subject reduces the chances of the subject developing undesired weight gain. In another embodiment, preventing reducing or suppressing food intake in a subject reduces the chances of the subject developing obesity.

[000138] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist for preventing undesired weight gain by a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000139] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said

agonist in the preparation of a medicament for preventing undesired weight gain by a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000140] In another embodiment, provided herein is a method of preventing, reducing or suppressing food intake in a subject, the method comprising the step of administering to the  
5 subject an effective dose of a composition comprising CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, thereby preventing reducing or suppressing food intake by a subject. In another  
10 embodiment, preventing, reducing or suppressing food intake by a subject reduces the chances of the subject developing undesired weight gain. In another embodiment, preventing reducing or suppressing food intake in a subject reduces the chances of the subject developing obesity.

[000141] In another embodiment, provided herein is a method of preventing undesired weight gain by a subject, the method comprising the step of administering to the subject an effective  
15 dose of a composition comprising CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby preventing weight gain in a subject. In another embodiment, the weight gain leads to or results in obesity of the subject. In another embodiment, the risk of gaining weight  
20 gain is due to a psychological condition, or due to a genetic predisposition to gain weight by the subject. In another embodiment, the psychological condition is depression, anxiety or post-traumatic stress disorder (PTSD).

[000142] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin  
25 carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist for preventing undesired weight gain by a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000143] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin  
30 carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist in the preparation of a medicament for preventing undesired weight gain by a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000144] In another embodiment, provided herein is a method of preventing undesired weight gain by a subject, the method comprising the step of administering to the subject an effective

dose of a composition comprising CTP-modified polypeptide comprising an oxyntomodulin (OXM) peptide and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, thereby preventing weight gain in a subject. In another embodiment, the weight gain leads to or results in obesity of the subject. In another embodiment, the risk of gaining weight gain is due to a psychological condition, or due to a genetic predisposition to gain weight by the subject. In another embodiment, the psychological condition is depression, anxiety or post-traumatic stress disorder (PTSD).

[000145] In another embodiment, provided herein is a method of treating obesity in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist peptide and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby treating obesity in the subject. In another embodiment, the subject is genetically predisposed to being obese. In another embodiment, the method of treating obesity results in a reduction of body weight in a subject. In another embodiment, the reduction in body weight is due to body fat reduction.

[000146] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating obesity in a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000147] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist in the preparation of a medicament for treating obesity in a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000148] In another embodiment, provided herein is a method of treating obesity in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising an oxyntomodulin peptide and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, thereby treating obesity in the subject. In another embodiment, the subject is genetically predisposed to being obese.

[000149] In one embodiment, provided herein is a method of treating type II diabetes in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist peptide and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby treating type II diabetes in the subject.

[000150] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating type II diabetes in a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000151] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist in the preparation of a medicament for treating type II diabetes in a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000152] In one embodiment, provided herein is a method of treating type II diabetes in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising an oxyntomodulin peptide and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the oxyntomodulin peptide, thereby treating type II diabetes in the subject.

[000153] In another embodiment, provided herein is a method of treating a metabolic disorder in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of the GLP-1/Glucagon receptor agonist peptide, thereby treating a metabolic disorder in the subject.

[000154] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating a metabolic disorder in a subject. In one embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000155] In another embodiment, provided herein is a use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist in the preparation of a medicament for treating a metabolic disorder in a subject. In one  
5 embodiment, said GLP-1/Glucagon receptor agonist is oxyntomodulin.

[000156] In another embodiment, provided herein is a method of treating a metabolic disorder in a subject, the method comprising administering to the subject an effective dose of a composition comprising a CTP-modified polypeptide comprising an oxyntomodulin peptide and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino  
10 terminus or carboxy terminus of the oxyntomodulin peptide, thereby treating a metabolic disorder in the subject.

[000157] In another embodiment, the metabolic disorder is diabetic ketoacidosis, or diabetes mellitus or any glucose-related metabolic disorder known in the art. In another embodiment, the metabolic disorder results from a lack of insulin and an overabundance of glucose in a  
15 subject.

[000158] In another embodiment, the engineered GLP-1/Glucagon receptor agonist variants of the present invention are synthesized using a polynucleotide molecule encoding a polypeptide of the present invention. In another embodiment, the polynucleotide molecule encoding the engineered GLP-1/Glucagon receptor agonist of the present invention is ligated into an  
20 expression vector, comprising a transcriptional control of a cis-regulatory sequence (e.g., promoter sequence). In another embodiment, the cis-regulatory sequence is suitable for directing constitutive expression of an engineered GLP-1/Glucagon receptor agonist of the present invention. In another embodiment, the cis-regulatory sequence is suitable for directing tissue-specific expression of the engineered GLP-1/Glucagon receptor agonist peptides of the  
25 present invention. In another embodiment, the cis-regulatory sequence is suitable for directing inducible expression of the engineered GLP-1/Glucagon receptor agonist variants of the present invention.

[000159] In one embodiment, tissue-specific promoters suitable for use with the present invention include sequences which are functional in one or more specific cell populations.  
30 Examples include, but are not limited to, promoters such as albumin that is liver-specific [Pinkert et al., (1987) Genes Dev. 1:268-277], lymphoid-specific promoters [Calame et al., (1988) Adv. Immunol. 43:235-275]; in particular promoters of T-cell receptors [Winoto et al., (1989) EMBO J. 8:729-733] and immunoglobulins; [Banerji et al. (1983) Cell 33:729-740], neuron-specific promoters such as the neurofilament promoter [Byrne et al. (1989) Proc. Natl.

Acad. Sci. USA 86:5473-5477], pancreas-specific promoters [Edlunch et al. (1985) Science 230:912-916] or mammary gland-specific promoters such as the milk whey promoter (U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Inducible promoters suitable for use with the present invention include, for example, the tetracycline-inducible promoter (Srouf, M.A., et al., 2003. Thromb. Haemost. 90: 398-405).

[000160] In one embodiment, the phrase "a polynucleotide molecule" refers to a single or double stranded nucleic acid sequence which is isolated and provided in the form of an RNA sequence, a complementary polynucleotide sequence (cDNA), a genomic polynucleotide sequence and/or a composite polynucleotide sequences (e.g., a combination of the above).

[000161] In one embodiment, a "complementary polynucleotide sequence" refers to a sequence, which results from reverse transcription of messenger RNA using a reverse transcriptase or any other RNA-dependent DNA polymerase. In one embodiment, the sequence can be subsequently amplified in vivo or in vitro using a DNA polymerase.

[000162] In one embodiment, a "genomic polynucleotide sequence" refers to a sequence derived (isolated) from a chromosome and thus it represents a contiguous portion of a chromosome.

[000163] In one embodiment, a "composite polynucleotide sequence" refers to a sequence, which is at least partially complementary and at least partially genomic. In one embodiment, a composite sequence can include some exonal sequences required to encode the polypeptide of the present invention, as well as some intronic sequences interposing therebetween. In one embodiment, the intronic sequences can be of any source, including of other genes, and typically includes conserved splicing signal sequences. In one embodiment, intronic sequences include cis-acting expression regulatory elements.

[000164] In one embodiment, following expression and secretion, the signal peptides are cleaved from the precursor engineered oxyntomodulin peptides resulting in the mature engineered oxyntomodulin peptides.

[000165] In one embodiments, polynucleotides of the present invention are prepared using PCR techniques, or any other method or procedure known to one skilled in the art. In some embodiments, the procedure involves the ligation of two different DNA sequences (See, for example, "Current Protocols in Molecular Biology", eds. Ausubel et al., John Wiley & Sons, 1992).

[000166] In one embodiment, polynucleotides of the present invention which encode the engineered GLP-1/Glucagon receptor agonist provided herein are inserted into expression vectors (i.e., a nucleic acid construct) to enable expression of the recombinant polypeptide. In

one embodiment, the expression vector of the present invention includes additional sequences which render this vector suitable for replication and integration in prokaryotes. In one embodiment, the expression vector of the present invention includes additional sequences which render this vector suitable for replication and integration in eukaryotes. In another embodiment, the expression vector of the present invention includes a shuttle vector which renders this vector suitable for replication and integration in both prokaryotes and eukaryotes. In some embodiments, cloning vectors comprise transcription and translation initiation sequences (e.g., promoters, enhances) and transcription and translation terminators (e.g., polyadenylation signals).

[000167] In one embodiment, a variety of prokaryotic or eukaryotic cells can be used as host-expression systems to express the CTP-modified GLP-1/Glucagon receptor agonist provided herein. In another embodiment, these include, but are not limited to, microorganisms, such as bacteria transformed with a recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vector containing the polypeptide coding sequence; yeast transformed with recombinant yeast expression vectors containing the polypeptide coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors, such as Ti plasmid, containing the polypeptide coding sequence.

[000168] In one embodiment, non-bacterial expression systems are used (e.g. mammalian expression systems such as CHO cells) to express the GLP-1/Glucagon receptor agonist provided herein. In another embodiment, the GLP-1/Glucagon receptor agonist is oxyntomodulin. In another embodiment, the oxyntomodulin is modified by the addition of at least one CTP, as further provided herein. In another embodiment, the expression vector used to express polynucleotides of the present invention in mammalian cells is pCI-dhfr vector. Construction of the pCI-dhfr vector is described, according to one embodiment, in Example's Materials and Methods, below.

[000169] In one embodiment, in bacterial systems of the present invention, a number of expression vectors can be advantageously selected depending upon the use intended for the polypeptide expressed. In one embodiment, large quantities of polypeptide are desired. In one embodiment, vectors that direct the expression of high levels of the protein product, possibly as a fusion with a hydrophobic signal sequence, which directs the expressed product into the periplasm of the bacteria or the culture medium where the protein product is readily purified are desired. In one embodiment, certain fusion proteins are engineered with a specific cleavage site to aid in recovery of the polypeptide. In one embodiment, vectors adaptable to

such manipulation include, but are not limited to, the pET series of *E. coli* expression vectors [Studier et al., Methods in Enzymol. 185:60-89 (1990)].

[000170] In one embodiment, yeast expression systems are used. In one embodiment, a number of vectors containing constitutive or inducible promoters can be used in yeast as disclosed in U.S. Pat. Application. No: 5,932,447, which is incorporated by reference herein in its entirety. In another embodiment, vectors which promote integration of foreign DNA sequences into the yeast chromosome are used.

[000171] In one embodiment, the expression vector of the present invention can further include additional polynucleotide sequences that allow, for example, the translation of several proteins from a single mRNA such as an internal ribosome entry site (IRES) and sequences for genomic integration of the promoter-chimeric polypeptide.

[000172] In one embodiment, mammalian expression vectors include, but are not limited to, pcDNA3, pcDNA3.1(+/-), pGL3, pZeoSV2(+/-), pSecTag2, pDisplay, pEF/myc/cyto, pCMV/myc/cyto, pCR3.1, pSinRep5, DH26S, DHBB, pNMT1, pNMT41, pNMT81, which are available from Invitrogen, pCI which is available from Promega, pMbac, pPbac, pBK-RSV and pBK-CMV which are available from Strategene, pTRES which is available from Clontech, and their derivatives.

[000173] In one embodiment, expression vectors containing regulatory elements from eukaryotic viruses such as retroviruses are used in the present invention. SV40 vectors include pSVT7 and pMT2. In some embodiments, vectors derived from bovine papilloma virus include pBV-1MTHA, and vectors derived from Epstein Bar virus include pHEBO, and p2O5. Other exemplary vectors include pMSG, pAV009/A+, pMTO10/A+, pMAMneo-5, baculovirus pDSVE, and any other vector allowing expression of proteins under the direction of the SV-40 early promoter, SV-40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in eukaryotic cells.

[000174] In one embodiment, recombinant viral vectors are useful for *in vivo* expression of the oxyntomodulin peptides of the present invention since they offer advantages such as lateral infection and targeting specificity. In one embodiment, lateral infection is inherent in the life cycle of, for example, a retrovirus and is the process by which a single infected cell produces many progeny virions that bud off and infect neighboring cells. In one embodiment, the result is that a large area becomes rapidly infected, most of which was not initially infected by the original viral particles. In one embodiment, viral vectors are produced that are

unable to spread laterally. In one embodiment, this characteristic can be useful if the desired purpose is to introduce a specified gene into only a localized number of targeted cells.

[000175] In one embodiment, various methods can be used to introduce the expression vector of the present invention into cells. Such methods are generally described in Sambrook et al.,  
5 Molecular Cloning: A Laboratory Manual, Cold Springs Harbor Laboratory, New York (1989, 1992), in Ausubel et al., Current Protocols in Molecular Biology, John Wiley and Sons, Baltimore, Md. (1989), Chang et al., Somatic Gene Therapy, CRC Press, Ann Arbor, Mich. (1995), Vega et al., Gene Targeting, CRC Press, Ann Arbor Mich. (1995), Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworths, Boston Mass. (1988) and  
10 Gilboa et al. [Biotechniques 4 (6): 504-512, 1986] and include, for example, stable or transient transfection, lipofection, electroporation and infection with recombinant viral vectors. In addition, see U.S. Pat. Nos. 5,464,764 and 5,487,992, incorporated herein by reference, for positive-negative selection methods.

[000176] In one embodiment, introduction of nucleic acid by viral infection offers several  
15 advantages over other methods such as lipofection and electroporation, since higher transfection efficiency can be obtained due to the infectious nature of viruses.

[000177] It will be appreciated that the engineered GLP-1/Glucagon receptor agonist or oxyntomodulin peptides of the present invention can also be expressed from a nucleic acid construct administered to the individual employing any suitable mode of administration (e.g.,  
20 subcutaneous administration, oral administration, intra-nasal administration, intra-venal administration, or in vivo gene therapy). In one embodiment, the nucleic acid construct is introduced into a suitable cell via an appropriate gene delivery vehicle/method (transfection, transduction, homologous recombination, etc.) and an expression system as needed and then the modified cells are expanded in culture and returned to the individual (i.e., ex vivo gene  
25 therapy).

[000178] In one embodiment, plant expression vectors are used. In one embodiment, the expression of a polypeptide coding sequence is driven by a number of promoters. In some embodiments, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV [Brisson et al., Nature 310:511-514 (1984)], or the coat protein promoter to TMV [Takamatsu  
30 et al., EMBO J. 6:307-311 (1987)] are used. In another embodiment, plant promoters are used such as, for example, the small subunit of RUBISCO [Coruzzi et al., EMBO J. 3:1671-1680 (1984); and Brogli et al., Science 224:838-843 (1984)] or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B [Gurley et al., Mol. Cell. Biol. 6:559-565 (1986)]. In one embodiment, constructs are introduced into plant cells using Ti plasmid, Ri plasmid, plant

viral vectors, direct DNA transformation, microinjection, electroporation and other techniques well known to the skilled artisan. See, for example, Weissbach & Weissbach [Methods for Plant Molecular Biology, Academic Press, NY, Section VIII, pp 421-463 (1988)]. Other expression systems such as insects and mammalian host cell systems, which are well known  
5 in the art, can also be used by the present invention.

[000179] It will be appreciated that other than containing the necessary elements for the transcription and translation of the inserted coding sequence (encoding the polypeptide), the expression construct of the present invention can also include sequences engineered to optimize stability, production, purification, yield or activity of the expressed polypeptide.

10 [000180] In one embodiment, transformed cells are cultured under effective conditions, which allow for the expression of high amounts of recombinant engineered oxyntomodulin peptides. In another embodiment, effective culture conditions include, but are not limited to, effective media, bioreactor, temperature, pH and oxygen conditions that permit protein production. In one embodiment, an effective medium refers to any medium in which a cell is cultured to  
15 produce the recombinant polypeptide of the present invention. In some embodiments, a medium typically includes an aqueous solution having assimilable carbon, nitrogen and phosphate sources, and appropriate salts, minerals, metals and other nutrients, such as vitamins. Cells of the present invention can be cultured in conventional fermentation bioreactors, shake flasks, test tubes, microtiter dishes and petri plates. In another embodiment,  
20 culturing is carried out at a temperature, pH and oxygen content appropriate for a recombinant cell. In another embodiment, the determination of culturing conditions are within the expertise of one of ordinary skill in the art.

[000181] In one embodiment, depending on the vector and host system used for production, the resultant engineered CTP-modified GLP-1/Glucagon receptor agonist or, in another  
25 embodiment, the resultant CTP-modified oxyntomodulin peptides of the present invention are expressed within a recombinant cell for glycosylation of the CTP to take place, are secreted into the fermentation medium, or are retained on the outer surface of a mammalian cell.

[000182] In one embodiment, following a predetermined time in culture, recovery of the recombinant engineered GLP-1/Glucagon receptor agonist or oxyntomodulin is effected.

30 [000183] In one embodiment, the phrase "recovering the recombinant engineered oxyntomodulin" refers to collecting the whole fermentation medium containing the polypeptide and need not imply additional steps of separation or purification.

[000184] In one embodiment, engineered oxyntomodulin or variants thereof provided herein are purified using a variety of standard protein purification techniques, such as, but not

limited to, affinity chromatography, ion exchange chromatography, filtration, electrophoresis, hydrophobic interaction chromatography, gel filtration chromatography, reverse phase chromatography, concanavalin A chromatography, chromatofocusing and differential solubilization.

5 [000185] To facilitate recovery, the expressed coding sequence can be engineered to encode the engineered GLP-1/Glucagon receptor agonist or oxyntomodulin of the present invention and fused cleavable moiety. Further, a fusion protein can be designed so that the polypeptide can be readily isolated by affinity chromatography; e.g., by immobilization on a column specific for the cleavable moiety. A cleavage site is engineered between the engineered  
10 oxyntomodulin and the cleavable moiety and the polypeptide can be released from the chromatographic column by treatment with an appropriate enzyme or agent that specifically cleaves the fusion protein at this site [e.g., see Booth et al., Immunol. Lett. 19:65-70 (1988); and Gardella et al., J. Biol. Chem. 265:15854-15859 (1990)].

[000186] In one embodiment, the engineered oxyntomodulin provided herein is retrieved in  
15 "substantially pure" form.

[000187] In one embodiment, the phrase "substantially pure" refers to a purity that allows for the effective use of the protein in the applications described herein.

[000188] The engineered GLP-1/Glucagon receptor agonist or oxyntomodulin provided herein can also be synthesized using in vitro expression systems. In another embodiment, *in vitro*  
20 synthesis methods are well known in the art and the components of the system are commercially available.

[000189] In one embodiment, the recombinant engineered oxyntomodulin peptides are synthesized and purified; their therapeutic efficacy can be assayed either *in vivo* or *in vitro*. The binding activities of the recombinant engineered oxyntomodulin peptides of the present  
25 invention can be ascertained using various assays as known to one of skill in the art.

[000190] In another embodiment, the GLP-1/Glucagon receptor agonist or engineered oxyntomodulin of the present invention can be provided to the individual per se. In one embodiment, the engineered GLP-1/Glucagon receptor agonist or oxyntomodulin of the present invention can be provided to the individual as part of a pharmaceutical composition  
30 where it is mixed with a pharmaceutically acceptable carrier.

[000191] In another embodiment, a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

[000192] In another embodiment, "active ingredient" refers to the polypeptide sequence of interest, which is accountable for the biological effect.

[000193] In another embodiment, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier" which are interchangeably used refer to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound. An adjuvant is included under these phrases. In one embodiment, one of the ingredients included in the pharmaceutically acceptable carrier can be for example polyethylene glycol (PEG), a biocompatible polymer with a wide range of solubility in both organic and aqueous media (Mutter et al. (1979)).

[000194] In another embodiment, "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient. In one embodiment, excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

[000195] Techniques for formulation and administration of drugs are found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition, which is incorporated herein by reference.

[000196] Various embodiments of dosage ranges are contemplated by this invention. The dosage of the CTP-modified GLP-1/Glucagon receptor agonist of the present invention, in one embodiment, is in the range of 0.005-100 mg/day. In another embodiment, the dosage is in the range of 0.005-5 mg/day. In another embodiment, the dosage is in the range of 0.01-50 mg/day. In another embodiment, the dosage is in the range of 0.1-20 mg/day. In another embodiment, the dosage is in the range of 0.1-10 mg/day. In another embodiment, the dosage is in the range of 0.01-5 mg/day. In another embodiment, the dosage is in the range of 0.001-0.01 mg/day. In another embodiment, the dosage is in the range of 0.001-0.1 mg/day. In another embodiment, the dosage is in the range of 0.1-5 mg/day. In another embodiment, the dosage is in the range of 0.5-50 mg/day. In another embodiment, the dosage is in the range of 0.2-15mg/day. In another embodiment, the dosage is in the range of 0.8-65 mg/day. In another embodiment, the dosage is in the range of 1-50 mg/day. In another embodiment, the dosage is in the range of 5-10 mg/day. In another embodiment, the dosage is in the range of 8-15 mg/day. In another embodiment, the dosage is in a range of 10-20mg/day. In another embodiment, the dosage is in the range of 20-40 mg/day. In another embodiment, the dosage is in a range of 60-120 mg/day. In another embodiment, the dosage is in the range of 12-40 mg/day. In another embodiment, the dosage is in the range of 40-60 mg/day. In another

embodiment, the dosage is in a range of 50-100mg/day. In another embodiment, the dosage is in a range of 1-60 mg/day. In another embodiment, the dosage is in the range of 15-25 mg/day. In another embodiment, the dosage is in the range of 5-10 mg/day. In another embodiment, the dosage is in the range of 55-65 mg/day.

5 [000197] In another embodiment, the dosage is in a range of 50-500 mg/day. In another embodiment, the dosage is in a range of 50-150 mg/day. In another embodiment, the dosage is in a range of 100-200 mg/day. In another embodiment, the dosage is in a range of 150-250 mg/day. In another embodiment, the dosage is in a range of 200-300 mg/day. In another embodiment, the dosage is in a range of 250-400 mg/day. In another embodiment, the dosage is in a range of 300-500 mg/day. In another embodiment, the dosage is in a range of 350-500 mg/day.

[000198] In one embodiment, the dosage is 20 mg/day. In one embodiment, the dosage is 30 mg/day. In one embodiment, the dosage is 40 mg/day. In one embodiment, the dosage is 50 mg/day. In one embodiment, the dosage is 0.01 mg/day. In another embodiment, the dosage is 0.1 mg/day. In another embodiment, the dosage is 1 mg/day. In another embodiment, the dosage is 0.530 mg/day. In another embodiment, the dosage is 0.05 mg/day. In another embodiment, the dosage is 50 mg/day. In another embodiment, the dosage is 10 mg/day. In another embodiment, the dosage is 20-70 mg/day. In another embodiment, the dosage is 5 mg/day.

20 [000199] In one embodiment, the dosage of the CTP-modified GLP-1/Glucagon receptor agonist is 1-5 mg/day. In one embodiment, the dosage of the CTP-modified oxyntomodulin is 1-3 mg/day. In another embodiment, the dosage of the CTP-modified oxyntomodulin is 2 mg/day.

[000200] In another embodiment, the dosage is 1-90 mg/day. In another embodiment, the dosage is 1-90 mg/2 days. In another embodiment, the dosage is 1-90 mg/3 days. In another embodiment, the dosage is 1-90 mg/4 days. In another embodiment, the dosage is 1-90 mg/5 days. In another embodiment, the dosage is 1-90 mg/6 days. In another embodiment, the dosage is 1-90 mg/week. In another embodiment, the dosage is 1-90 mg/9 days. In another embodiment, the dosage is 1-90 mg/11 days. In another embodiment, the dosage is 1-90 mg/14 days.

30 [000201] In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist dosage is 10-50 mg/day. In another embodiment, the dosage is 10-50 mg/2 days. In another embodiment, the dosage is 10-50 mg/3 days. In another embodiment, the dosage is 10-50 mg/4 days. In another embodiment, the dosage is 10-50 micrograms mg/5 days. In another

embodiment, the dosage is 10-50 mg/6 days. In another embodiment, the dosage is 10-50 mg/week. In another embodiment, the dosage is 10-50 mg/9 days. In another embodiment, the dosage is 10-50 mg/11 days. In another embodiment, the dosage is 10-50 mg/14 days.

[000202] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is formulated in an intranasal dosage form. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is formulated in an injectable dosage form. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.0001 mg to 0.6 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.001 mg to 0.005 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.005 mg to 0.01 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.01 mg to 0.3 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose in a dose ranging from 0.2 mg to 0.6 mg. In another embodiment, the GLP-1/Glucagon receptor agonist is free of CTPs on its amino terminus.

[000203] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is formulated in an intranasal dosage form. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is formulated in an injectable dosage form. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.0001 mg to 0.6 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.001 mg to 0.005 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.005 mg to 0.01 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.01 mg to 0.3 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose in a dose ranging from 0.2 mg to 0.6 mg. In another embodiment, the GLP-1/Glucagon receptor agonist is free of CTPs on its amino terminus.

[000204] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 1-100 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-80 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 20-60 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-50 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 40-80 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-30 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 30-60 micrograms.

[000205] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 1-100 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-80 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 20-60 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-50 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 40-80 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 10-30 micrograms. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 30-60 micrograms.

[000206] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.2 mg to 2 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 2 mg to 6 mg. In

another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 4 mg to 10 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 5 mg and 15 mg.

5 [000207] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 0.2 mg to 2 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 2 mg to 6 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 4 mg to 10 mg. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject in a dose ranging from 5 mg and 15 mg.

[000208] In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 65% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 55% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 45% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 10% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 25% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 35% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 75% of the amount of the agonist than that administered using the non-CTP-modified GLP-1/Glucagon receptor agonist. In another embodiment, the dosage of CTP-modified GLP-1/Glucagon receptor agonist is such that it contains 100% of the amount of the agonist than that administered using the non-CTP-modified agonist. However, even if the dosage contains the same amount of agonist as non-CTP-modified GLP-

1/Glucagon receptor agonist, it is still advantageous to subjects in that it will be administered less frequently because of its increased half-life compared to normal agonist.

[000209] In another embodiment, a therapeutically effective amount of a conjugated GLP-1/Glucagon receptor agonist is between 50-500 IU per kg body weight administered once a day. In another embodiment, a therapeutically effective amount of a conjugated GLP-1/Glucagon receptor agonist is 150-250 IU per kg body weight, administered once a day. In another embodiment, a pharmaceutical composition comprising a conjugated GLP-1/Glucagon receptor agonist is formulated at a strength effective for administration by various means to a human patient.

[000210] In one embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 20-30 IU/dL in a subject. In another embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 25-50 IU/dL in a subject. In another embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 50-100 IU/dL in a subject. In another embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 100-200 IU/dL in a subject. In another embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 10-50 IU/dL in a subject. In another embodiment, the CTP-modified polypeptide comprising GLP-1/Glucagon receptor agonist is administered in an amount effective to bring circulating GLP-1/Glucagon receptor agonist activity to 20-100 IU/dL in a subject.

[000211] In one embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject on a weekly basis. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject twice a week. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject on a fortnightly (once every two weeks) basis. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject twice a month. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject once a month. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is

administered to a subject on a daily basis. In another embodiment, the CTP-modified GLP-1/Glucagon receptor agonist is administered to a subject every two days.

[000212] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every three days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every four days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every five days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every six days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every 7-14 days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every 10-20 days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every 5-15 days. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is administered to a subject once every 15-30 days.

[000213] In another embodiment, the methods of the invention include increasing the compliance in the use of GLP-1/Glucagon receptor agonist therapy, comprising providing to a subject in need thereof, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the carboxy terminus of the GLP-1/Glucagon receptor agonist, thereby increasing compliance in the use of GLP-1/Glucagon receptor agonist therapy.

[000214] In another embodiment, the methods of the invention include increasing the compliance of patients afflicted with chronic illnesses that are in need of a GLP-1/Glucagon receptor agonist therapy. In another embodiment, the methods of the invention enable reduction in the dosing frequency of a GLP-1/Glucagon receptor agonist by modifying the GLP-1/Glucagon receptor agonist with CTPs as described hereinabove.

[000215] In another embodiment, the term compliance comprises adherence. In another embodiment, the methods of the invention include increasing the compliance of patients in need of a GLP-1/Glucagon receptor agonist therapy by reducing the frequency of administration of the GLP-1/Glucagon receptor agonist. In another embodiment, reduction in the frequency of administration of the GLP-1/Glucagon receptor agonist is achieved due to the CTP modifications which render the CTP-modified GLP-1/Glucagon receptor agonist

more stable. In another embodiment, reduction in the frequency of administration of the GLP-1/Glucagon receptor agonist is achieved as a result of increasing  $T_{1/2}$  of the GLP-1/Glucagon receptor agonist. In another embodiment, reduction in the frequency of administration of the GLP-1/Glucagon receptor agonist is achieved as a result of increasing the clearance time or  
5 reducing the clearance rate of the GLP-1/Glucagon receptor agonist.

[000216] In another embodiment, the present invention provides a method of reducing the clearance rate of GLP-1/Glucagon receptor agonist, comprising the step of at least one chorionic gonadotrophin carboxy terminal peptides (CTPs) to the amino or carboxy terminus of GLP-1/Glucagon receptor agonist, thereby reducing the clearance rate of GLP-1/Glucagon  
10 receptor agonist.

[000217] In another embodiment, reduction in the frequency of administration of the GLP-1/Glucagon receptor agonist is achieved as a result of increasing the AUC measure of the GLP-1/Glucagon receptor agonist.

[000218] Oral administration, in one embodiment, comprises a unit dosage form comprising  
15 tablets, capsules, lozenges, chewable tablets, suspensions, emulsions and the like. Such unit dosage forms comprise a safe and effective amount of the desired GLP-1/Glucagon receptor agonist of the invention, each of which is in one embodiment, from about 0.7 or 3.5 mg to about 280 mg/70 kg, or in another embodiment, about 0.5 or 10 mg to about 210 mg/70 kg. The pharmaceutically-acceptable carriers suitable for the preparation of unit dosage forms for  
20 peroral administration are well-known in the art. In another embodiments, tablets typically comprise conventional pharmaceutically-compatible adjuvants as inert diluents, such as calcium carbonate, sodium carbonate, mannitol, lactose and cellulose; binders such as starch, gelatin and sucrose; disintegrants such as starch, alginic acid and croscarmellose; lubricants such as magnesium stearate, stearic acid and talc. In one embodiment, glidants such as silicon  
25 dioxide can be used to improve flow characteristics of the powder-mixture. In one embodiment, coloring agents, such as the FD&C dyes, can be added for appearance. Sweeteners and flavoring agents, such as aspartame, saccharin, menthol, peppermint, and fruit flavors, are useful adjuvants for chewable tablets. Capsules typically comprise one or more solid diluents disclosed above. In another embodiment, the selection of carrier components  
30 depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention, and can be readily made by a person skilled in the art.

[000219] In one embodiment, the oral dosage form comprises predefined release profile. In one embodiment, the oral dosage form of the present invention comprises an extended release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form of

the present invention comprises a slow release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form of the present invention comprises an immediate release tablets, capsules, lozenges or chewable tablets. In one embodiment, the oral dosage form is formulated according to the desired release profile of the pharmaceutical active ingredient as known to one skilled in the art.

[000220] Peroral compositions, in some embodiments, comprise liquid solutions, emulsions, suspensions, and the like. In some embodiments, pharmaceutically-acceptable carriers suitable for preparation of such compositions are well known in the art. In some embodiments, liquid oral compositions comprise from about 0.001% to about 0.933% of the desired compound or compounds, or in another embodiment, from about 0.01% to about 10%.

[000221] In one embodiment, compositions for use in the methods of this invention comprise solutions or emulsions, which in some embodiments are aqueous solutions or emulsions comprising a safe and effective amount of the compounds of the present invention and optionally, other compounds, intended for topical intranasal administration. In another embodiment, the compositions comprise from about 0.001% to about 10.0% w/v of a subject compound, more preferably from about 0.1% to about 2.0, which is used for systemic delivery of the compounds by the intranasal route.

[000222] In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is injected into the muscle (intramuscular injection). In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is injected below the skin (subcutaneous injection). In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is injected into the muscle. In another embodiment, a polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one CTP unit is injected into the skin. In another embodiment, a GLP-1/Glucagon receptor agonist as described herein is administered via systemic administration. In another embodiment, a GLP-1/Glucagon receptor agonist as described herein is administered by intravenous injection. In another embodiment, administration can be parenteral, pulmonary, oral, topical, intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, transnasal, intraocular, ophthalmic, epidural, buccal, rectal, transmucosal, intestinal or parenteral delivery, including intramedullary injections as well as intrathecal or direct intraventricular administration.

[000223] In another embodiment, the preparation is administered in a local rather than systemic manner, for example, via injection of the preparation directly into a specific region of a patient's body.

[000224] In one embodiment, the route of administration may be enteral. In another embodiment, the route may be conjunctival, transdermal, intradermal, intra-arterial, vaginal, rectal, intratumoral, paracancerous, transmucosal, intramuscular, intravascular, intraventricular, intracranial, intra-nasal, sublingual, or a combination thereof.

5 [000225] In another embodiment, the pharmaceutical compositions are administered by intravenous, intra-arterial, or intramuscular injection of a liquid preparation. In some embodiments, liquid formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In one embodiment, the pharmaceutical compositions are administered intravenously, and are thus formulated in a form suitable for intravenous administration. In  
10 another embodiment, the pharmaceutical compositions are administered intra-arterially, and are thus formulated in a form suitable for intra-arterial administration. In another embodiment, the pharmaceutical compositions are administered intramuscularly, and are thus formulated in a form suitable for intramuscular administration.

[000226] Further, and in another embodiment, the pharmaceutical compositions are  
15 administered topically to body surfaces, and are thus formulated in a form suitable for topical administration. Suitable topical formulations include gels, ointments, creams, lotions, drops and the like. For topical administration, the compounds of the present invention are combined with an additional appropriate therapeutic agent or agents, prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a  
20 pharmaceutical carrier.

[000227] In one embodiment, pharmaceutical compositions of the present invention are manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

25 [000228] In one embodiment, pharmaceutical compositions for use in accordance with the present invention is formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which, can be used pharmaceutically. In one embodiment, formulation is dependent upon the route of administration chosen.

30 [000229] In one embodiment, injectables of the invention are formulated in aqueous solutions. In one embodiment, injectables of the invention are formulated in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer. In some embodiments, for transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[000230] In one embodiment, the preparations described herein are formulated for parenteral administration, e.g., by bolus injection or continuous infusion. In some embodiments, formulations for injection are presented in unit dosage form, e.g., in ampoules or in multidose containers with optionally, an added preservative. In some embodiments, compositions are  
5 suspensions, solutions or emulsions in oily or aqueous vehicles, and contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[000231] The compositions also comprise, in some embodiments, preservatives, such as benzalkonium chloride and thimerosal and the like; chelating agents, such as edetate sodium and others; buffers such as phosphate, citrate and acetate; tonicity agents such as sodium  
10 chloride, potassium chloride, glycerin, mannitol and others; antioxidants such as ascorbic acid, acetylcystine, sodium metabisulfite and others; aromatic agents; viscosity adjusters, such as polymers, including cellulose and derivatives thereof; and polyvinyl alcohol and acid and bases to adjust the pH of these aqueous compositions as needed. The compositions also comprise, in some embodiments, local anesthetics or other active ingredients. The  
15 compositions can be used as sprays, mists, drops, and the like.

[000232] In one embodiment, pharmaceutical compositions for parenteral administration include aqueous solutions of the active preparation in water-soluble form. Additionally, suspensions of the active ingredients can be prepared as appropriate oil or water based injection suspensions. Suitable lipophilic solvents or vehicles include, in some embodiments,  
20 fatty oils such as sesame oil, or synthetic fatty acid esters such as ethyl oleate, triglycerides or liposomes. Aqueous injection suspensions contain, in some embodiments, substances, which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol or dextran. In another embodiment, the suspension also contains suitable stabilizers or agents which increase the solubility of the active ingredients to allow for the preparation of highly  
25 concentrated solutions.

[000233] In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez- Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; J. E. Diederichs and al.,  
30 Pharm./nd. 56 (1994) 267- 275).

[000234] In another embodiment, the pharmaceutical composition delivered in a controlled release system is formulated for intravenous infusion, implantable osmotic pump, transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump is used (see Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery

88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989). In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, 5 vol. 2, pp. 115-138 (1984). Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

[000235] In another embodiment, the active ingredient is in powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water based solution, before use. Compositions are formulated, in some embodiments, for atomization and inhalation administration. In 10 another embodiment, compositions are contained in a container with attached atomizing means.

[000236] In one embodiment, the preparation of the present invention is formulated in rectal compositions such as suppositories or retention enemas, using, e.g., conventional suppository bases such as cocoa butter or other glycerides.

15 [000237] In one embodiment, pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose. In another embodiment, a therapeutically effective amount means an amount of active ingredients effective to prevent, alleviate or ameliorate symptoms of disease or prolong the survival of the subject being treated.

20 [000238] In one embodiment, determination of a therapeutically effective amount is well within the capability of those skilled in the art.

[000239] Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose and its derivatives, such as sodium carboxymethyl 25 cellulose, ethyl cellulose, and methyl cellulose; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the Tween™ brand emulsifiers; wetting agents, such sodium lauryl 30 sulfate; coloring agents; flavoring agents; tableting agents, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions. The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the compound is basically determined by the way the compound is to be administered. If the subject compound is to be injected, in one embodiment, the pharmaceutically-acceptable carrier is sterile,

physiological saline, with a blood-compatible suspending agent, the pH of which has been adjusted to about 7.4.

[000240] In addition, the compositions further comprise binders (e.g. acacia, cornstarch, gelatin, carbomer, ethyl cellulose, guar gum, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, povidone), disintegrating agents (e.g. cornstarch, potato starch, alginic acid, silicon dioxide, croscarmellose sodium, crospovidone, guar gum, sodium starch glycolate), buffers (e.g., Tris-HCl, acetate, phosphate) of various pH and ionic strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), protease inhibitors, surfactants (e.g. sodium lauryl sulfate), permeation enhancers, solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite, butylated hydroxyanisole), stabilizers (e.g. hydroxypropyl cellulose, hydroxypropylmethyl cellulose), viscosity increasing agents (e.g. carbomer, colloidal silicon dioxide, ethyl cellulose, guar gum), sweeteners (e.g. aspartame, citric acid), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), lubricants (e.g. stearic acid, magnesium stearate, polyethylene glycol, sodium lauryl sulfate), flow-aids (e.g. colloidal silicon dioxide), plasticizers (e.g. diethyl phthalate, triethyl citrate), emulsifiers (e.g. carbomer, hydroxypropyl cellulose, sodium lauryl sulfate), polymer coatings (e.g., poloxamers or poloxamines), coating and film forming agents (e.g. ethyl cellulose, acrylates, polymethacrylates) and/or adjuvants.

[000241] Typical components of carriers for syrups, elixirs, emulsions and suspensions include ethanol, glycerol, propylene glycol, polyethylene glycol, liquid sucrose, sorbitol and water. For a suspension, typical suspending agents include methyl cellulose, sodium carboxymethyl cellulose, cellulose (e.g. Avicel™, RC-591), tragacanth and sodium alginate; typical wetting agents include lecithin and polyethylene oxide sorbitan (e.g. polysorbate 80).

Typical preservatives include methyl paraben and sodium benzoate. In another embodiment, peroral liquid compositions also contain one or more components such as sweeteners, flavoring agents and colorants disclosed above.

[000242] The compositions also include incorporation of the active material into or onto particulate preparations of polymeric compounds such as polylactic acid, polglycolic acid, hydrogels, etc, or onto liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts.) Such compositions will influence the physical state, solubility, stability, rate of *in vivo* release, and rate of *in vivo* clearance.

[000243] Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound coupled to antibodies directed

against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors.

[000244] In one embodiment, compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or polyproline. In another embodiment, the modified compounds exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds. In one embodiment, modifications also increase the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the immunogenicity and reactivity of the compound. In another embodiment, the desired in vivo biological activity is achieved by the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

[000245] Preparation of effective amount or dose can be estimated initially from in vitro assays. In one embodiment, a dose can be formulated in animal models and such information can be used to more accurately determine useful doses in humans.

[000246] In one embodiment, toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures in vitro, in cell cultures or experimental animals. In one embodiment, the data obtained from these in vitro and cell culture assays and animal studies can be used in formulating a range of dosage for use in human. In one embodiment, the dosages vary depending upon the dosage form employed and the route of administration utilized. In one embodiment, the exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. [See e.g., Fingl, et al., (1975) "The Pharmacological Basis of Therapeutics", Ch. 1 p.1].

[000247] In one embodiment, depending on the severity and responsiveness of the condition to be treated, dosing can be of a single or a plurality of administrations, with course of treatment lasting from several days to several weeks or until cure is effected or diminution of the disease state is achieved.

[000248] In one embodiment, the amount of a composition to be administered will be dependent on the subject being treated, the severity of the affliction, the manner of administration, the judgment of the prescribing physician, etc.

[000249] In one embodiment, compositions including the preparation of the present invention formulated in a compatible pharmaceutical carrier are also prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

[000250] In another embodiment, a GLP-1/Glucagon receptor agonist or a variant (CTP-modified) form thereof, as described herein, is lyophilized (i.e., freeze-dried) preparation in combination with complex organic excipients and stabilizers such as nonionic surface active agents (i.e., surfactants), various sugars, organic polyols and/or human serum albumin. In another embodiment, a pharmaceutical composition comprises a lyophilized GLP-1/Glucagon receptor agonist as described in sterile water for injection. In another embodiment, a pharmaceutical composition comprises a lyophilized GLP-1/Glucagon receptor agonist as described in sterile PBS for injection. In another embodiment, a pharmaceutical composition comprises a lyophilized GLP-1/Glucagon receptor agonist as described in sterile 0.9% NaCl for injection.

[000251] In another embodiment, the pharmaceutical composition comprises a CTP-modified GLP-1/Glucagon receptor agonist as described herein and complex carriers such as human serum albumin, polyols, sugars, and anionic surface active stabilizing agents. In another embodiment, the pharmaceutical composition comprises a CTP-modified GLP-1/Glucagon receptor agonist as described herein and lactobionic acid and an acetate/glycine buffer. In another embodiment, the pharmaceutical composition comprises a CTP-modified GLP-1/Glucagon receptor agonist as described herein and amino acids, such as arginine or glutamate that increase the solubility of interferon compositions in water. In another embodiment, the pharmaceutical composition comprises a lyophilized CTP-modified GLP-1/Glucagon receptor agonist as described herein and glycine or human serum albumin (HSA), a buffer (e.g. acetate) and an isotonic agent (e.g. NaCl). In another embodiment, the pharmaceutical composition comprises a lyophilized CTP-modified GLP-1/Glucagon receptor agonist as described herein and phosphate buffer, glycine and HSA.

[000252] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein is stabilized when placed in buffered solutions having a pH between about 4 and 7.2. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist is in a buffered solution having a pH between about 4 and 8.5. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist is in a buffered solution having a pH between about 6 and 7. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist is

in a buffered solution having a pH of about 6.5. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein is stabilized with an amino acid as a stabilizing agent and in some cases a salt (if the amino acid does not contain a charged side chain).

5 [000253] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein is a liquid composition comprising a stabilizing agent at between about 0.3% and 5% by weight which is an amino acid.

[000254] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein provides dosing accuracy and product safety. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein provides a biologically active, stable liquid formulation for use in injectable applications. In another embodiment, the pharmaceutical composition comprises a non-lyophilized CTP-modified GLP-1/Glucagon receptor agonist as described herein.

[000255] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist provided herein provides a liquid formulation permitting storage for a long period of time in a liquid state facilitating storage and shipping prior to administration.

20 [000256] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist provided herein comprises solid lipids as matrix material. In another embodiment, the injectable pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises solid lipids as matrix material. In another embodiment, the production of lipid microparticles by spray congealing was described by Speiser (Speiser and al., Pharm. Res. 8 (1991) 47-54) followed by lipid nanopellets for peroral administration (Speiser EP 0167825 (1990)). In another embodiment, lipids, which are used, are well tolerated by the body (e. g. glycerides composed of fatty acids which are present in the emulsions for parenteral nutrition).

[000257] In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises polymeric microparticles. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises nanoparticles. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises liposomes. In another

embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises lipid emulsion. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises microspheres. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises lipid nanoparticles. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises lipid nanoparticles comprising amphiphilic lipids. In another embodiment, the pharmaceutical composition comprising a CTP-modified GLP-1/Glucagon receptor agonist as described herein comprises lipid nanoparticles comprising a drug, a lipid matrix and a surfactant. In another embodiment, the lipid matrix has a monoglyceride content which is at least 50% w/w.

[000258] In one embodiment, compositions of the present invention are presented in a pack or dispenser device, such as an FDA approved kit, which contain one or more unit dosage forms containing the active ingredient. In one embodiment, the pack, for example, comprise metal or plastic foil, such as a blister pack. In one embodiment, the pack or dispenser device is accompanied by instructions for administration. In one embodiment, the pack or dispenser is accommodated by a notice associated with the container in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions or human or veterinary administration. Such notice, in one embodiment, is labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert.

[000259] In one embodiment, it will be appreciated that the GLP-1/Glucagon receptor agonist of the present invention can be provided to the individual with additional active agents to achieve an improved therapeutic effect as compared to treatment with each agent by itself. In another embodiment, measures (e.g., dosing and selection of the complementary agent) are taken to avoid adverse side effects which are associated with combination therapies.

[000260] In one embodiment, the term "about", means in quantitative terms plus or minus 5%, or in another embodiment plus or minus 10%, or in another embodiment plus or minus 15%, or in another embodiment plus or minus 20%.

[000261] The term "subject" refers in one embodiment to a mammal including a human in need of therapy for, or susceptible to, a condition or its sequelae. The subject may include dogs, cats, pigs, cows, sheep, goats, horses, rats, and mice and humans. The term "subject" does not exclude an individual that is normal in all respects.

[000262] It is to be understood that in one embodiment, methods of treatment and prevention of the CTP-modified polypeptide as described herein may be substituted by uses of the CTP-modified polypeptide as described herein for accomplishing the methods for treatment and prevention as described herein and/or uses of the CTP-modified polypeptide as described herein in the preparation of a medicament for accomplishing the methods for treatment and prevention as described herein.

[000263] Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

## EXAMPLES

### Materials and Methods:

#### *Plasmid Construction*

[000264] Seven OXM plasmids were constructed in eukaryotic expression vector (pCI-dhfr) based on the OXM amino acid sequence (accession # NP\_002045) and the CTP amino acid sequence (accession # NP\_149032).

[000265] A schematic presentation of these plasmids is shown in Table 1. A detailed description of plasmids construction follows.

[000266] **Table 1:** Schematic description of plasmids

|   | Plasmid variant description | Nucleic Acid Sequence |
|---|-----------------------------|-----------------------|
| 1 | OXM-CTP-CTP                 | SEQ ID NO: 20         |
| 2 | CTP-OXM-CTP                 | SEQ ID NO: 8          |
| 3 | CTP-CTP-OXM                 | SEQ ID NO: 22         |
| 4 | CTP-OXM-CTP-CTP             | SEQ ID NO: 14         |
| 5 | OXM-CTP-CTP-CTP             | SEQ ID NO: 24         |
| 6 | OXM-CTP-CTP-CTP-CTP         | SEQ ID NO: 26         |
| 7 | OXM-CTP-CTP-CTP-CTP-CTP     | SEQ ID NO: 28         |

[000267] A nucleic acid sequence of the OXM variant CTP-OXM-CTP (0.5C CTP-OXM-CTP, GenArt, GA # 0804377) was synthesized after a codon usage optimization for DG44

expression system. An XbaI–NotI fragment containing 0.5C CTP-OXM-CTP sequence was isolated. The fragment was inserted into the eukaryotic expression vector pCI-dhfr to yield the 601-0-p142-1 clone.

### **Construction of CTP-OXM-CTP**

5 [000268] The following primers were used in order to synthesize 601-6-p149-1 (CTP-OXM-CTP).

[000269] **Primer 25** 5' CTCTAGAGGACATGGCCAC 3' (SEQ ID NO: 4).

[000270] **Primer 85R** 5' CTGGCTGTGCTGGGGCAGAATGGGTGT 3' (SEQ ID NO: 5).

10 [000271] **Primer 86** 5' CCCCAGCACAGCCAGGG 3' (SEQ ID NO: 6).

[000272] **Primer 74R** 5' GCGGCCGCATCCAGACCT 3' (SEQ ID NO: 7).

[000273] Three PCR reactions were performed. The first reaction was conducted with primer 25 and primer 85R and plasmid DNA of 402-3-p81-4 (CTP-hGH-CTP-CTP) as a template; as a result of the PCR amplification, a ~181 bp product was formed. The second reaction was  
 15 conducted with primer 86 and primer 74R and plasmid DNA of 601-0-p142-1 (0.5C CTP-OXM-CTP) as a template; as a result of the PCR amplification, a ~224 bp product was formed. The last reaction was conducted with primers 25 and 74R and a mixture of the products of the previous two reactions as a template; as a result of the PCR amplification, a ~391 bp product was formed and ligated into TA cloning vector (Invitrogen, catalog K2000-  
 20 01). An XbaI–NotI fragment containing CTP-OXM-CTP sequence was isolated. The fragment was inserted into our eukaryotic expression vector pCI-dhfr to yield 601-6-p149-1 clone.

[000274] The nucleic acid sequence encoding CTP-OXM-CTP is as follows:

tctagaggacatggccaccggcagcaggaccagcctgctgctggccttcggcctgctgtgcctgccatggctgcaggagggcagcg  
 25 ccagctcttcttetaaggctccacccccatctctgccagccccagcagactgccgggccccagcgacacaccattctgccccagca  
 cagccagggcaccttcaccagcgactacagcaagtacctggacagcagaagggcccaggacttcgtccagtggctgatgaacacca  
 agaggaacaggaacaacatcgcttccttagctccaaggccctccaccctctctgcctagcccctctcggctgctggcccatccga  
 cacaccaatcctgccacagtgatgaaggtctggatgcggccgc (SEQ ID NO: 8).

[000275] The amino acid sequence encoding CTP-OXM-CTP is as follows:

MATGSRTSLLLAFLGLCLPWLQEGSASSSSKAPPPSLPSPSRLPGPSDTPILPQHSQGTFTSDYSKYLDSRRAQDFVQWLMNTRNRNNIASSSSKAPPPSLPSPSRLPGPSDTPILPQ (SEQ ID NO: 9).

### **Construction of CTP-OXM-CTP-CTP**

5 [000276] The following primers were used in order to synthesize 601-3-p158-2 (CTP-OXM-CTP-CTP):

[000277] **Primer 25** 5' CTCTAGAGGACATGGCCAC 3' (contains the restriction site of XbaI ) (SEQ ID NO: 10).

10 [000278] **Primer 87<sup>R</sup>** 5' GCTGGAGCTAGCGATGTTGTTCTGTTCC 3' (contains the 3' end of OXM and the 5' end of CTP) (SEQ ID NO: 11).

[000279] **Primer 88** 5' ACATCGCTAGCTCCAGCAGCAAGGCC 3' (contains the 3' end of OXM and the 5' end of CTP) (SEQ ID NO: 12).

[000280] **Primer 74<sup>R</sup>** 5' GCGGCCGCATCCAGACCT 3' (contains the restriction site of NotI) (SEQ ID NO: 13).

15 [000281] Three PCR reactions were performed. The first reaction was conducted with primer 25 and primer 87R and plasmid DNA of 601-6-p149-1 (CTP-OXM-CTP-CTP) as a template; as a result of the PCR amplification, a ~ 290 bp product was formed. The second reaction was conducted with primer 88 and primer 74R and plasmid DNA of 402-3-p81-4 (CTP-hGH-CTP-CTP) as a template; as a result of the PCR amplification, a ~200 bp product was formed.

20 The last reaction was conducted with primers 25 and 74R and a mixture of the products of the previous two reactions as a template; as a result of the PCR amplification, a ~ 450 bp product was formed and ligated into TA cloning vector (Invitrogen, catalog K2000-01). XbaI –NotI fragment containing CTP-OXM-CTP-CTP sequence was isolated. The fragment was inserted in to our eukaryotic expression vector pCI-dhfr to yield 601-3-p158-2 clone.

25 [000282] The nucleic acid sequence encoding CTP-OXM-CTP-CTP is as follows:

tctagaggacatggccaccggcagcaggaccagcctgctgctggccttcggcctgctgtgcctgccatggctgcaggagggcagcgcagctcttcttctaaggctccaccccatctctgccagccccagcagactgccgggccccagcgacacaccattctgccccagcacgccagggcaccttcaccagcgactacagcaagtacctggacagcagaagggccccaggacttcgtccagtggctgatgaacaccaagaggaacaggaacaacatcgctagctccagcagcaaggccccctccccgagcctgccctccccagcaggctgcctgggcccctcgcacacaccaatcctgccacagagcagctcctctaaggccccctcctccatccctgccatccccctcccggctgcctggcccctctgacaccctatcctgcctcagtgatgaaggtctggatgcggccgc (SEQ ID NO: 14).

30

[000283] The amino acid sequence encoding CTP-OXM-CTP-CTP is as follows:

MATGSRTSLLLAFLGLCLPWLQEGSASSSSKAPPPSLPSPSRLPGPSDTPILPQHSQGTF  
TSDYSKYLDSRRAQDFVQWLMNTRNRNNIASSSSKAPPPSLPSPSRLPGPSDTPILPQ  
SSSSKAPPPSLPSPSRLPGPSDTPILPQ (SEQ ID NO: 15).

#### 5 Construction of OXM-CTP-CTP

[000284] The following primers were used in order to synthesize 601-2-p160-2 (OXM-CTP-CTP):

[000285] **Primer 75** 5' CTCCCAGTTCAATTACAGCT 3' (contains sequence of pCI-dhfr before XbaI restriction site) (SEQ ID NO: 16).

10 [000286] **Primer 89<sup>R</sup>** 5' GCTGTGAGCGCTGCCCTCCTGCAG 3' (contains 5' end of OXM) (SEQ ID NO: 17).

[000287] **Primer 90** 5' GCGCTCACAGCCAGGGCACCTTC 3' (contains the 5' end of OXM) (SEQ ID NO: 18).

15 [000288] **Primer 74<sup>R</sup>** 5' GCGGCCGCATCCAGACCT 3' (contains the restriction site of NotI) (SEQ ID NO: 19).

[000289] Three PCR reactions were performed. The first reaction was conducted with primer 75 and primer 89R and plasmid DNA of 601-0-p142-1 (0.5C CTP-OXM-CTP) as a template; as a result of the PCR amplification, a ~175 bp product was formed. The second reaction was conducted with primer 90 and primer 74R and plasmid DNA of 601-3-p158-2 (CTP-OXM-CTP-CTP) as a template; as a result of the PCR amplification, a ~ 200 bp product was formed. The last reaction was conducted with primers 25 and 74R and a mixture of the products of the previous two reactions as a template; as a result of the PCR amplification, a ~ 391 bp product was formed and ligated into TA cloning vector (Invitrogen, catalog K2000-01). XbaI –NotI fragment containing OXM-CTP-CTP sequence was isolated. The fragment was inserted into our eukaryotic expression vector pCI-dhfr to yield the 601-2-p160-2 clone.

[000290] The nucleic acid sequence encoding OXM-CTP-CTP is as follows:

tctagaggacatggccacagggagcaggaccagcctgctgctggctttcggcctgctgtgtctgccatggctgcaggagggcagcgc  
tcacagccagggcaccttcaccagcgactacagcaagtacctggacagcagaaggggccaggacttcgtccagtggctgatgaaca  
ccaagaggaacaggaacaacatcgctagctccagcagcaaggccctccccgagcctgccctcccaagcaggctgcctgggcc  
30 ctccgacacaccaatcctgccacagagcagctcctctaaggccctcctccatccctgccatccccctcccgctgctggccctctg  
acaccctatcctgcctcagtgatgaaggctctggatgcggccgc (SEQ ID NO: 20)

[000291] The amino acid sequence encoding OXM-CTP-CTP is as follows:

LEDMATGSRTSLLLAFGLLCLPWLQEGSAHSQGTFTSDYSKYLDSRRAQDFVQWLM  
NTRNRNNIASSSSKAPPPSLPSPSRLPGPSDTPILPQSSSSKAPPPSLPSPSRLPGPSDTPILPQ (SEQ ID NO: 21)

#### 5 **Construction of CTP- CTP-OXM**

[000292] A nucleic acid sequence of OXM variant (CTPx2-OXM, GenArt, GA # 1067101) was synthesized after a codon usage optimization for DG44 expression system. The fragment was inserted in to Excellgene eukaryotic expression vector to yield p162 clone.

[000293] The nucleic acid sequence encoding CTPx2-OXM is as follows:

10 ctatagggcgaattgaaggaaggccgtcaaggccgatgagctctctagaggacatggccaccggcagcaggaccagcctgctgct  
ggccttcggcctgctgtgcctgccatggctgcaggaggcgcagcgcagctccagcagcaaggccctccccgagcctgccctccc  
caagcaggctgcctgggccctccgacacaccaatcctgccacagagcagctcctctaaggccctcctccatccctgccatccccctc  
ccggctgcctggcccccttgacaccctatcctgcctcagcacagccagggcaccttcaccagcgactacagcaagtacctggacag  
cagaagggccaggacttcgtccagtggctgatgaacaccaagaggaacaggaacaacatcgcttgatgaaggtctggatgcggcc  
15 gcggtaccctgggcctcatgggccttccttcaactgcccgtttccag (SEQ ID NO: 22).

[000294] The amino acid sequence encoding CTPx2-OXM is as follows:

M A T G S R T S L L L A F G L L C L P W L Q E G S A S S S S K A P  
P P S L P S P S R L P G P S D T P I L P Q S S S S K A P P P S L P S P  
S R L P G P S D T P I L P Q H S Q G T F T S D Y S K Y L D S R R A  
20 Q D F V Q W L M N T K R N R N N I A \* \* R S G C G R G T L G L M  
G L P F T A R F P (SEQ ID NO: 23).

#### **Construction of OXM-CTPx3:**

[000295] A nucleic acid sequence of OXM variant (OXM-CTPx3, GenArt, GA # 1017864) was synthesized after a codon usage optimization for DG44 expression system. XbaI –NotI  
25 fragment containing OXM-CTPx3 sequence was isolated. The fragment was inserted in to our eukaryotic expression vector pCI-dhfr to yield pCI-dhfr-OXM-ctpx3-p216-4 clone.

[000296] The nucleic acid sequence encoding OXM-CTPx3 is as follows:

tctagactcgagcgatcgccatggccaccggctctaggacctccctgctgctggccttcggcctgctgtgcctgccctggctgcagga  
aggcagcgctcactcccagggcaccttcacctccgactactccaagtacctggactctcggagagcccaggacttcgtgcagtggctg  
30 atgaacaccaagcggaaccggaacaatatgcctcctcaagctccaaggcacctccaccttcctgcttagcccttcagactccctg

ggcccagtgacaccctatctgctcagtcagtcagcaaggccccaccctagcctgccttctccttctcggtgctggcccc  
agcgatactccaattctgccccagtcctccagcagtaaggctccccctccatcttgccatccccagcagactgccaggcccttctgat  
acaccatcctcccacagtgatgaggatccgcggccgc (SEQ ID NO: 24).

[000297] The amino acid sequence encoding OXM-CTPx3 is as follows:

5 M A T G S R T S L L L A F G L L C L P W L Q E G S A H S Q G T F T  
S D Y S K Y L D S R R A Q D F V Q W L M N T K R N R N N I A S S  
S S K A P P P S L P S P S R L P G P S D T P I L P Q S S S S K A P P P  
S L P S P S R L P G P S D T P I L P Q S S S S K A P P P S L P S P S R  
L P G P S D T P I L P Q \* \* (SEQ ID NO: 25).

#### 10 Construction of OXM-CTPx4:

[000298] A nucleic acid sequence of OXM variant (OXM-CTPx4, GenArt, GA # 1115769) was synthesized after a codon usage optimization for DG44 expression system. XbaI –NotI fragment containing OXM-CTPx4 sequence was isolated. The fragment was inserted in to the eukaryotic expression vector pCI-dhfr to yield pCI-dhfr-OXM-ctpx4 -p254-3 clone.

15 [000299] The nucleic acid sequence encoding OXM-CTPx4 is as follows:

tctagactcgagcgatcgccatggctaccggctccagaacctctctgctgctggccttcggcctgctgtgtctgccttggctgcaagagg  
gcagcgctcattcccagggcaccttcacctccgactactccaagtacctggactctgcagagcccaggacttcgtgcagtggctgatg  
aacaccaagcggaaaccggaacaatatcgctcctccagctccaaggccccctcctccatcttgccatccccagtagactgcctgggc  
cctctgacaccctatctgctcagtcagtcctctaaggccccaccaccttcctgctagcccttcaagactgccaggccctagcg  
20 atacaccaattctgccccagtcctccagcagcaaggctccccacctagcctgccttctccatcaaggctgcctggcccatccgatacc  
ccaattttgcctcagagcagctctagcaaggcacctccccccagctctgcctctccaagcagactccctggcccttcagacactccatt  
ctgccacagtgatgaggatccgcggccgc (SEQ ID NO: 26)

[000300] The amino acid sequence encoding OXM-CTPx4 is as follows:

M A T G S R T S L L L A F G L L C L P W L Q E G S A H S Q G  
25 T F T S D Y S K Y L D S R R A Q D F V Q W L M N T K R N R  
N N I A S S S S K A P P P S L P S P S R L P G P S D T P I L P Q  
S S S S K A P P P S L P S P S R L P G P S D T P I L P Q S S S S  
K A P P P S L P S P S R L P G P S D T P I L P Q S S S S K A P P  
P S L P S P S R L P G P S D T P I L P Q \* \* (SEQ ID NO: 27).

#### 30 Construction of OXM-CTPx5:

[000301] A nucleic acid sequence of OXM variant (OXM-CTPx5, GenArt, GA # 1115770) was synthesized after a codon usage optimization for DG44 expression system. XbaI –NotI fragment containing OXM-CTPx5 sequence was isolated. The fragment was inserted in to the eukaryotic expression vector pCI-dhfr to yield pCI-dhfr-OXM-ctpx5 -p255-1 clone.

5 [000302] The nucleic acid sequence encoding OXM-CTPx5 is as follows:

ctctagactcgagcgatcgccatggctaccggctccagaacctctctgctgctggcctcggcctgctgtgtctgccttggtgcaagag  
ggcagcgctcattcccagggcaccttcacctccgactactccaagtacctggactctcgagagcacaggacttcgtgcagtggctgat  
gaacaccaagcgggaaccggaacaatatcgctcctccagctccaagggccctcctccatctctgcatccccagtagactgcctggg  
ccctctgacacccctatctgctcagtcagctcctctaaggtccaccaccttcctgcctagccctcaagactgccaggccctagc  
10 gatacaccaattctgccccagtcctccagcagaaggctccccacctagcctgccttctccatcaagggtgcctggcccatccgatac  
cccaattttgctcagagcagctctagcaaggcacctccccagctctgccttctccaagcagactccctggcccttcagacactccaat  
cctcccacagtcctctagctctaaagtcacctccagcctgccagccctagtagactccccggaccttctgatacccccatcttgcc  
ccagtgatgaggatccgcggccgc (SEQ ID NO: 28).

15 [000303] The amino acid sequence encoding OXM-CTPx5 is as follows:

M A T G S R T S L L L A F G L L C L P W L Q E G S A H S Q G T F T  
S D Y S K Y L D S R R A Q D F V Q W L M N T K R N R N N I A S S  
S S K A P P P S L P S P S R L P G P S D T P I L P Q S S S S K A P P P  
S L P S P S R L P G P S D T P I L P Q S S S S K A P P P S L P S P S R  
20 L P G P S D T P I L P Q S S S S K A P P P S L P S P S R L P G P S D  
T P I L P Q S S S S K A P P P S L P S P S R L P G P S D T P I L P Q \*  
\* (SEQ ID NO: 29).

### **Expression, purification and characterization of OXM-CTP variants**

25 [000304] All seven variants of OXM-CTPs were transiently expressed and produced in XLG's CHO-Express cell line (Excellgene Company, Switzerland). Concentration levels of OXM-CTPs harvests were determined using an OXM ELISA kit (Bachem Cat# S-1393.0001). The harvest was purified using a DEAE column, followed by a Jacalin column as a second purification step. The final fractions were dialyzed against 10mM buffer citrate, 147 mM  
30 NaCl, pH 6. The concentration of purified variants was determined by absorbance at 280 nm using extinction coefficient of 1.9=1mg/ml. 1.9 is the extinction coefficient that theoretically was calculated for the OXM peptide. Since CTP peptide does not absorb at 280 nm, the same

extinction coefficient was applied for OXM-CTP variants. OXM native peptide was chemically synthesized (Almac Company, Ireland), and its peptide content was determined by amino acid analysis.

**Pharmacokinetic (PK) profile of OXM-CTP variants**

- 5 [000305] The pharmacokinetic profiles of the OXM peptide and OXM-CTP variants were assessed as follows. Male Sprague-Dawley (SD)-1 rats were administered subcutaneously (SC) or intravenously (IV) with a single dose of native OXM (n=6, 230µg/kg), of variants CTP-OXM-CTP, CTP-OXM-CTP-CTP, OXM-CTP-CTP and CTP-CTP-OXM (n=6, 230µg/kg) or variants # OXM-CTP-CTP-CTP, OXM-CTP-CTP-CTP-CTP, OXM-CTP-CTP-CTP-CTP-CTP (n=6, 153 µg/kg). Cohorts of 3 animals per group were bled at alternating time points. OXM serum concentration was analyzed using commercial ELISA kit (Bachem, Cat# S-1393.0001). The study design is summarized in Table 2. The study was divided into 3 sequential experiments, Experiment #1 (Groups 1-4), Experiment #2 (Groups 5-9) and Experiment #3 (Groups 10-13).

- 15 [000306] **Table 2:** Summary of the PK study design.

*IPGTT study in C57BL/6 mice*

| Gro<br>up # | Test<br>Article<br>Variant<br># | No. of<br>animals<br>/ group/<br>timepoi<br>nt | Dose<br>Route | Gend<br>er | Dose<br>Level<br>of<br>OXM<br>(µg/rat<br>) | Dose<br>Level<br>of<br>OXM<br>(µg/kg) | Time-Points<br>(hours post-dose)                                             |
|-------------|---------------------------------|------------------------------------------------|---------------|------------|--------------------------------------------|---------------------------------------|------------------------------------------------------------------------------|
| 1           | OXM                             | 6/3                                            | IV            | Male       | 34                                         | 230                                   | 0 (Pre-dose), 5min, 15min,<br>25min, 35min, 45min, 60min,<br>75min, 1.5h, 2h |
| 2           | CTP-<br>OXM-<br>CTP             | 6/3                                            | IV            | Male       | 34                                         | 230                                   | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 3h, 4h , 6h, 8h , 24h,<br>36h, 48h    |
| 3           | CTP-<br>OXM-<br>CTP-<br>CTP     | 6/3                                            | IV            | Male       | 34                                         | 230                                   | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 3h, 4h , 6h, 8h, 24h,<br>36h, 48h     |

|    |                             |     |    |      |    |     |                                                                             |
|----|-----------------------------|-----|----|------|----|-----|-----------------------------------------------------------------------------|
| 4  | CTP-<br>CTP-<br>OXM         | 6/3 | IV | Male | 34 | 230 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 3h, 4h , 6h, 8h , 24h,<br>36h, 48h   |
| 5  | OXM                         | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 5min, 15min,<br>25min, 35min, 45min, 60min<br>75min, 1.5h, 2h |
| 6  | CTP-<br>OXM-<br>CTP         | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h,4h , 6h, 8h ,24h, 36h,<br>2d, 3d      |
| 7  | CTP-<br>OXM-<br>CTP-<br>CTP | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h,4h , 6h, 8h , 24h, 36h,<br>2d, 3d     |
| 8  | CTP-<br>CTP-<br>OXM         | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h ,4h , 6h, 8h , 24h, 36h,<br>2d,3d     |
| 9  | OXM-<br>CTP-<br>CTP         | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 4h , 6h, 8h, 24h, 36h,<br>2d, 3d     |
| 10 | OXM                         | 6/3 | SC | Male | 34 | 230 | 0 (Pre-dose), 5min, 15min,<br>25min, 35min, 45min, 60min<br>75min, 1.5h, 2h |
| 11 | OXM-<br>CTPX3               | 6/3 | SC | Male | 23 | 153 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 4h , 6h, 8h, 24h, 36h,<br>2d, 3d, 4d |
| 12 | OXM-<br>CTPX4               | 6/3 | SC | Male | 23 | 153 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h ,4h , 6h, 8h, 24h, 36h,<br>2d, 3d, 4d |
| 13 | OXM-<br>CTPX5               | 6/3 | SC | Male | 23 | 153 | 0 (Pre-dose), 10min, 30min,<br>1h, 2h, 4h , 6h, 8h, 24h, 36h,<br>2d, 3d, 4d |

[000307] Overnight-fasted mice were measured for glucose (pre-dose glucose). Immediately after, they were IP injected with one of the test articles. Fifteen or 120 min post injection, glucose levels were measured (zero time for glucose) followed by glucose administration via IP injection (1.5 g/kg). Additional glucose measurement at 10, 20, 30, 60, 90, 120 and 180 min were performed. Blood glucose level were measured by tail vein sampling using a handheld glucometer.

#### **Food intake study in C57BL/6 mice**

[000308] At least one week before injection, mice were weighed and were transferred to a mini cage for acclimatization, (one mouse per cage). During the acclimatization period, they were handled daily and received two injections of vehicle to minimize stress during the study period. The day before the experiment, the mice were fasted. Seventeen hours after fasting, at the early light phase (900–1000h), the mice were weighed again (prior to the IP injection) followed by a single IP injection of 1700 nmol/kg (10 µl/1g mice) of OXM peptide or OXM-CTP variants OXM-CTPX3, OXM-CTPX4 and OXM-CTPX5. After injection, the mice were returned to their home cages (1 mouse per cage), and provided with a pre-weighed amount of chow. Food intake were measured 0, 1, 2, 4, 6, 8, 21, 32 and 44, 55, 68, 80, 93 and 141h post injection by weighing the chow. At the end of experiment, rats were weighed again.

### **RESULTS**

#### **EXAMPLE 1: CONSTRUCTION OF CTP-MODIFIED OXM**

[000309] By genetic engineering, CTP peptide cDNA was fused to human OXM cDNA, generating seven different OXM-CTP variant as detailed in Table 1. The nucleotide sequences of the plasmids were verified, and the plasmids were transiently transfected into XLG's CHO-Express cell line (Excellgene Company, Switzerland). The OXM-CTP variants were secreted into the growth medium, harvests were collected and OXM-CTPs levels were determined.

#### **EXAMPLE 2: EXPRESSION, PURIFICATION AND CHARACTERIZATION OF OXM-CTP VARIANTS**

[000310] The production media (harvests) were measured for secretion level of OXM-CTP variants. The secretion levels are summarized in Table 3. The secretion levels obtained were high considering the standard transient transfection expression levels of recombinant peptides.

[000311] **Table 3:** Summary of OXM-CTPs secretion level.

| Variant                       | CTP-OXM-CTP | CTP-OXM-CTP-CTP | OXM-CTP-CTP | CTP-CTP-OXM | OXM-CTPX4 | OXM-CTPX4 | OXM-CTPX5 |
|-------------------------------|-------------|-----------------|-------------|-------------|-----------|-----------|-----------|
| Harvest concentration (µg/ml) | 31.1        | 15.3            | 3.3         | 51.9        | 13.9      | 15.3      | 14.4      |

5

[000312] The harvests were purified according to the method described in *Materials and Methods*. The final samples of variants OXM-CTPX3, OXM-CTPX4 and OXM-CTPX5 were analyzed by SDS-PAGE (Coomassie staining (Figure 1A) and anti-OXM Western Blot Analysis (Figure 1B)). For variants CTP-OXM-CTP, CTP-OXM-CTP-CTP, OXM-CTP-CTP and CTP-CTP-OXM, samples from the purification process were analyzed by both

10 Coomassie staining and Western Blot analysis (Figure 2). As expected, OXM-CTP variants showed differences in size correlating to the number of CTP cassettes fused to the peptide. The relatively high molecular weights are an indication that there may be high occupancy of the O-glycan chains on the potential serine sites on the CTP peptide. The OXM peptide did

15 not react with the anti-OXM antibody, probably due to the technical difficulty in transferring a small sized peptide. The Coomassie staining shows that the OXM-CTP variants were highly purified and contained mainly the high form.

### **EXAMPLE 3: PHARMACOKINETIC (PK) PROFILE OF OXM-CTP VARIANTS**

20 [000313] The pharmacokinetic profile of OXM peptide compared to OXM-CTP variants was analyzed in male SD-1 rats. Animals were administered with a single IV (Experiment 1, Figure 3A) or single SC (Experiments 2&3, Figures 3B-3C) injection of native OXM or OXM-CTP variants CTP-OXM-CTP, CTP-OXM-CTP-CTP, OXM-CTP-CTP, CTP-CTP-OXM (230µg/kg peptide) or variants OXM-CTPX3, OXM-CTPX4 and OXM-CTPX5

25 (153µg/kg peptide). Serum concentration of OXM or OXM-CTP variants at the indicated time points was analyzed using commercial ELISA. The PK profile is shown in Figure 3, and the conventional noncompartmental PK parameters are summarized in Table 4. Addition of

CTP peptides to OXM resulted in prolonging the half-life of native OXM, from 0.22hr to 2.7 - 10 h for various OXM-CTP variants (Table 4, SC administration; Experiments 2 & 3).

[000314] Two copies of CTP were added to OXM peptide in three different ways to produce variants CTP-OXM-CTP, OXM-CTP-CTP and CTP-CTP-OXM. The most significant extension of serum half-life was received when the CTPs were added in tandem to the C-terminus of OXM; variant OXM-CTP-CTP with  $T_{1/2}$  of 4.76h. Fusion of three CTPs to the OXM C-terminus did not results in an elongated half-life as compared to fusion of two CTPs. Surprisingly, addition of four and five copies of CTPs to OXM c-terminus elongated the  $T_{1/2}$  up to 10h (Table 4).

[000315] Exposure as reflected by the AUC parameter, was most increased by ~17-fold for variant OXM-CTP-CTP (Experiment 2) and ~30-fold for variants OXM-CTPX4 and OXM-CTPX5 (Experiment 3). These results indicate that there is a superior prolonging effect after the addition of four or five copies of CTP to the OXM peptide.

[000316] The bioavailability as calculated from Experiments 1 & 2 was increased for CTP-OXM-CTP-CTP, but no significant improvement was found for CTP-OXM-CTP and CTP-CTP-OXM (Table 4).

[000317] **Table 4:** Conventional noncompartmental PK parameters of OXM peptide and OXM-CTP variants as were determined in three different experiments.

|                  |                 | $T_{1/2}$<br>hr | $T_{1/2B}$<br>hr | $C_{max}$<br>ng/ml | $T_{max}$<br>hr | $AUC(0-t)$<br>ng-hr/ml | $AUC_{\infty}$<br>ng-hr/ml | CL<br>ml/hr/kg | Vd<br>ml/kg | MRT<br>hr | Bioavail<br>ability<br>% |
|------------------|-----------------|-----------------|------------------|--------------------|-----------------|------------------------|----------------------------|----------------|-------------|-----------|--------------------------|
| Exp.<br>1<br>IV. | OXM             | 0.29            | NA               | NA                 | NA              | 211.7                  | 213.9                      | 993.0          | 417.0       | 0.34      |                          |
|                  | CTP-OXM-CTP     | 0.25            | 1.57             | NA                 | NA              | 592.7                  | 598.0                      | 355.3          | 806.9       | 0.81      |                          |
|                  | CTP-OXM-CTP-CTP | 0.28            | 1.61             | NA                 | NA              | 968.2                  | 977.4                      | 217.4          | 506.4       | 0.80      |                          |
|                  | CTP-CTP-OXM     | 0.27            | 1.20             | NA                 | NA              | 488.6                  | 490.4                      | 433.3          | 748.1       | 0.86      |                          |
| Exp.<br>2<br>SC. | OXM             | 0.22            |                  | 118.40             | 0.08            | 42.52                  | 42.84                      | 4960.39        | 1564.40     | 0.37      | 20.09                    |
|                  | CTP-OXM-CTP     | 3.28            |                  | 31.11              | 1.00            | 144.87                 | 170.82                     | 1243.97        | 5887.75     | 4.73      | 24.44                    |
|                  | CTP-OXM-CTP-CTP | 4.39            |                  | 30.18              | 6.00            | 313.98                 | 320.86                     | 662.28         | 4139.25     | 6.62      | 32.43                    |
|                  | OXM-CTP-CTP     | 4.76            |                  | 50.90              | 6.00            | 697.00                 | 702.10                     | 302.70         | 2078.90     | 8.20      | NA                       |
|                  | CTP-CTP-OXM     | 2.74            |                  | 27.77              | 1.00            | 114.15                 | 129.85                     | 1636.55        | 6459.76     | 4.10      | 23.36                    |

|                   |            |       |  |      |      |       |       |       |        |  |  |
|-------------------|------------|-------|--|------|------|-------|-------|-------|--------|--|--|
| <b>Exp. 3 SC.</b> | <b>OXM</b> | 0.18  |  | 36.4 | 0.08 | 13.1  | 13.3  | 16999 | 4375   |  |  |
|                   | OXM-CTPX3  | 3.78  |  | 25.1 | 2.0  | 308.9 | 312   | 512.8 | 2797   |  |  |
|                   | OXM-CTPX4  | 10.15 |  | 21.1 | 2.0  | 400.4 | 431.9 | 355.4 | 5379.6 |  |  |
|                   | OXM-CTPX5  | 9.63  |  | 21.2 | 8    | 431   | 442   | 346.2 | 4812   |  |  |

[000318] A comparison between the various groups showing the fold increase of T1/2 and AUC $\infty$  demonstrated that OXM-CTPx4 and OX-CTPx5 were superior in these parameters compared to other variants (Table 5).

[000319] **Table 5:** Fold of increase of OXM-CTP variants's PK parameters in compared to OXM peptide as calculated from Experiments 1-3.

|               | Variant                 | T1/2<br>Sample/OXM<br>peptide | AUC $\infty$<br>Sample/OXM peptide |
|---------------|-------------------------|-------------------------------|------------------------------------|
| Exp. 1<br>IV. | CTP-OXM-CTP             | NA                            | 2.8                                |
|               | CTP-OXM-CTP-CTP         | NA                            | 5.6 <sub>10</sub>                  |
|               | CTP-CTP-OXM             | NA                            | 2.29                               |
| Exp. 2<br>SC. | CTP-OXM-CTP             | 14.9                          | 3.98                               |
|               | CTP-OXM-CTP-CTP         | 19.9                          | 7.48                               |
|               | OXM-CTP-CTP             | 21.6                          | 16.3                               |
|               | CTP-CTP-OXM             | 12.4                          | 3.03                               |
| Exp. 3<br>SC. | OXM-CTP-CTP-CTP         | 21                            | 23.45                              |
|               | OXM-CTP-CTP-CTP-CTP     | 56.4                          | 32.478 <sub>15</sub>               |
|               | OXM-CTP-CTP-CTP-CTP-CTP | 53.5                          | 33.23                              |

#### **EXAMPLE 4: CTP-MODIFIED OXM INDUCES GLUCOSE TOLERANCE**

[000320] An IPGTT study was carried out in C57BL/6 mice demonstrated that OXM enhanced glucose clearance via stimulation of insulin secretion. The IP glucose tolerance test (IPGTT) evaluates the glucose lowering effect of OXM. In order to evaluate the *in vivo* activity of OXM or OXM-CTP variants, the IPGTT model was applied. Overnight fasted C57BL/6 mice were injected IP with OXM peptide or OXM-CTP variants followed by IP injection of glucose (1.5g/kg) and measurement of blood glucose levels from the tail vein by glucometer (Figure 4). OXM-CTP variants were assessed in two sequential experiments; Experiment 1 for variants CTP-OXM-CTP, CTP-OXM-CTP-CTP, OXM-CTP-CTP, CTP-CTP-OXM (Figures 4A and 4C) and Experiment 2 for variants OXM-CTP-CTP-CTP, OXM-

CTPX4 and OXM-CTPX5 (Figures 4B and 4D). OXM (100 nmol/kg), or OXM-CTP variants (100 nmol/kg) were administered IP 15 min or 2 hrs (OXM peptide and variant OXM-CTPX4) prior to glucose IP injection, and the induction of glucose tolerance was compared to vehicle (buffer) group. A marked effect was measured for OXM-CTP variants CTP-OXM-CTP-CTP, OXM-CTP-CTP, OXM-CTPX3, OXM-CTPX4 and OXM-CTPX5 as reflected by reduction of 20-30% of blood glucose AUC compared to vehicle group as compared to 100nmol/kg of OXM peptide which had low impact on the calculated AUC (Figure 4, reduction of 1%, Experiment 1 and 6.7%, Experiment 2). Surprisingly, CTP-OXM-CTP resulted in increased glucose levels while CTP-CTP-OXM had a minor impact on glucose tolerance. OXM-CTPX4 induced glucose tolerance activity even when administered 120 min prior to glucose while OXM peptide activity was no longer apparent. This result is aligned with the improved pharmacokinetics profile of this variant.

#### **EXAMPLE 5: CTP-MODIFIED OXM REDUCE FOOD INTAKE**

[000321] The pharmacological activity of OXM and OXM-CTP variants was further evaluated in C57BL/6 mice following single SC drug administration. In this study, the acute effect on food intake was measured. Male C57BL/6 mice (n=4 per group) were administered with a single SC administration of OXM peptide (2000nmol/kg) or OXM-CTP (1700nmol/kg) variants: OXM-CTPX3, OXM-CTPX4 or OXM-CTPX5. Food intake was measured daily for 6 days. OXM and all measured OXM-CTP variants resulted in a significant reduction in food intake, as was measured 1h and 2h post injection, 34-51% (compared to vehicle) food intake for OXM and OXM-CTP variant-treated mice, as shown in Figure 6. The OXM peptide effect was abolished 4h following injection, while all other OXM-CTP variants were effective for at least 24h. The most significant effect was found for variant OXM-CTPX5, which reduced food intake for 5 days compared to vehicle. The percentage of food intake compared to vehicle, for various time intervals over the experiment is presented in Figure 6B. The effect of native OXM disappeared after one hour, OXM-4/5 CTP maintained the inhibition for 2 hours and OXM-5CTP for about 80 hours (3 days). Of note, the superiority of variant OXM-CTPx5 in acute food intake study is in correlation to its improved PK profile. Surprisingly, although variant OXM-CTPx4 demonstrated an improved PK profile with  $T_{1/2}$  of about 10h, its effect did not last as long as OXM-CTPx5 or OXM-CTPx3.

[000322] Having described preferred embodiments of the invention with reference to the accompanying drawings, it is to be understood that the invention is not limited to the precise

embodiments, and that various changes and modifications may be effected therein by those skilled in the art without departing from the scope or spirit of the invention as defined in the appended claims.

**What is claimed is:**

1. A CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist.
2. The CTP-modified polypeptide of claim 1, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
3. The CTP-modified polypeptide of any one of claims 1-2, wherein the amino acid sequence of at least one CTP is selected from the group consisting of SEQ ID NO: 1 and SEQ ID NO: 2.
4. The CTP-modified polypeptide of any one of claims 1-3, wherein at least one CTP is glycosylated.
5. The CTP-modified polypeptide of any one of claims 1-4, wherein at least one CTP is truncated.
6. The CTP-modified polypeptide of any one of claims 1-5, wherein at least one CTP is attached to said agonist polypeptide via a linker.
7. The CTP-modified polypeptide of claim 6, wherein said linker is a peptide bond.
8. The CTP-modified polypeptide of any one of claims 1-7, wherein five chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
9. The CTP-modified polypeptide of any one of claims 1-7, wherein four chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
10. The CTP-modified polypeptide of any one of claims 1-7, wherein three chorionic gonadotrophin carboxy terminal peptides are attached to said agonist, one CTP on the amino terminus and two CTP on the carboxy terminus of said agonist.
11. The CTP-modified polypeptide of any one of claims 1-7, wherein three chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
12. The CTP-modified polypeptide of any one of claims 1-7, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist, one CTP on the carboxy terminus and one CTP on the amino terminus of said agonist.
13. The CTP-modified polypeptide of any one of claims 1-7, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist on the carboxy terminus of said agonist.

14. The CTP-modified polypeptide of any one of claims 1-7, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist on the amino terminus of said agonist.
15. The CTP-modified polypeptide of any one of claims 1-14, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
16. A pharmaceutical composition comprising the CTP-modified polypeptide of any one of claims 1-15.
17. A polynucleotide encoding the CTP-modified polypeptide of any one of claims 1-15.
18. The polynucleotide of claim 17, wherein the sequence of said polynucleotide comprises SEQ ID NO: 8, 14, 20, 22, 24, 26 or 28.
19. An expression vector comprising the polynucleotide of any one of claims 17-18.
20. A cell comprising the expression vector of claim 19.
21. A composition comprising the expression vector of claim 19.
22. A method of producing a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist, the method comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptide to the amino terminus or carboxy terminus of said agonist.
23. The CTP-modified polypeptide of claim 22, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
24. The method of any one of claims 22-23, wherein five chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
25. The method of any one of claims 22-23, wherein four chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
26. The method of any one of claims 22-23, wherein three chorionic gonadotrophin carboxy terminal peptides are attached to said agonist, one CTP on the amino terminus and two CTP on the carboxy terminus of said agonist.
27. The method of any one of claims 22-23, wherein three chorionic gonadotrophin carboxy terminal peptides are attached to the carboxy terminus of said agonist.
28. The method of any one of claims 22-23, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist, one CTP on the carboxy terminus and one CTP on the amino terminus of said agonist.

29. The method of any one of claims 22-23, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist on the carboxy terminus of said agonist.
30. The method of any one of claims 22-23, wherein two chorionic gonadotrophin carboxy terminal peptides are attached to said agonist on the amino terminus of said agonist.
31. The method of any one of claims 22-30, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
32. The method of any one of claims 22-31, wherein the amino acid sequence of at least one CTP is selected from the group consisting of SEQ ID NO: 1 and SEQ ID NO:2.
33. The method of any one of claims 22-32, wherein at least one CTP is glycosylated.
34. The method of any one of claims 22-33, wherein at least one CTP is truncated.
35. The method of any one of claims 22-34, wherein at least one CTP is attached to said agonist via a linker.
36. The method of claim 35, wherein said linker is a peptide bond.
37. A method of extending the biological half life of a dual GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of said agonist, thereby improving the biological half life of said agonist.
38. The CTP-modified polypeptide of claim 37, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
39. The method of any one of claims 37-38, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
40. The method of any one of claims 37-39, wherein at least one CTP is glycosylated.
41. The method of any one of claims 37-40, wherein at least one CTP is truncated.
42. The method of any one of claims 37-41, wherein at least one CTP is attached to said agonist via a linker.
43. The method of claim 42, wherein said linker is a peptide bond.
44. The method of any one of claims 37-43, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
45. A method of improving the area under the curve (AUC) of a dual GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to a carboxy terminus of said agonist, thereby improving the area under the curve (AUC) of said agonist.

46. The CTP-modified polypeptide of claim 45, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
47. The method of any one of claims 45-46, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
48. The method of any one of claims 45-47, wherein at least one CTP is glycosylated.
49. The method of any one of claims 45-48, wherein at least one CTP is truncated.
50. The method of any one of claims 45-49, wherein at least one CTP is attached to said agonist via a linker.
51. The method of claim 50, wherein said linker is a peptide bond.
52. The method of any one of claims 45-51, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
53. A method of reducing the dosing frequency of a dual GLP-1/Glucagon receptor agonist, comprising the step of attaching at least one chorionic gonadotrophin carboxy terminal peptides to an amino or a carboxy terminus of the GLP-1/Glucagon receptor agonist, thereby reducing the dosing frequency of GLP-1/Glucagon receptor agonist.
54. The method of claim 53, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
55. The method of any one of claims 53-54, wherein the amino acid sequence of at least one CTP is selected from the group consisting of SEQ ID NO: 1 and SEQ ID NO: 2.
56. The method of any one of claims 53-55, wherein at least one CTP is glycosylated.
57. The method of any one of claims 53-56, wherein at least one CTP is truncated.
58. The method of any one of claims 53-57, wherein at least one CTP is attached to the GLP-1/Glucagon receptor agonist via a linker.
59. The method of claim 58, wherein said linker is a peptide bond.
60. The method of any one of claims 53-59, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
61. Use of a CTP-modified polypeptide comprising a dual GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or caboxy terminus of said agonist for inducing glucose tolerance in a subject.
62. The use of claim 61, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
63. The use of any one of claims 61-62, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
64. The use of any one of claims 61-63, wherein at least one CTP is glycosylated.

65. The use of any one of claims 61-64, wherein at least one CTP is truncated.
66. The use of any one of claims 61-65, wherein at least one CTP is attached to said agonist polypeptide via a linker.
67. The use of claim 66, wherein said linker is a peptide bond.
68. The use of any one of claims 61-67, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
69. Use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist for preventing undesired weight gain by a subject.
70. The use of claim 69, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
71. The use of any one of claims 69-70, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
72. The use of any one of claims 69-71, wherein at least one CTP is glycosylated.
73. The use of any one of claims 69-72, wherein at least one CTP is truncated.
74. The use of any one of claims 69-73, wherein at least one CTP is attached to said agonist via a linker.
75. The use of claim 74, wherein said linker is a peptide bond.
76. The use of any one of claims 69-75, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
77. The use of any one of claims 69-76, wherein said weight gain results in obesity in the subject.
78. The use of any one of claims 69-77, wherein said weight gain is due to a psychological condition, or due to a genetic predisposition to weight gain by said subject.
79. The use of claim 78, wherein said psychological condition is depression, anxiety or post-traumatic stress disorder (PTSD).
80. Use of a composition comprising CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide (CTP) attached to the amino terminus or carboxy terminus of said agonist for preventing, reducing or suppressing food intake in a subject.
81. The use of claim 80, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.

82. The use of any one of claims 80-81, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
83. The use of any one of claims 80-82, wherein at least one CTP is glycosylated.
84. The use of any one of claims 80-83, wherein at least one CTP is truncated.
85. The use of any one of claims 80-84, wherein at least one CTP is attached to said agonist via a linker.
86. The use of claim 85, wherein said linker is a peptide bond.
87. The use of any one of claims 80-86, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
88. The use of any one of claims 80-87, wherein preventing, reducing or suppressing food intake by said subject reduces the chances of said subject developing undesired weight gain.
89. The use of any one of claims 80-87, wherein preventing, reducing or suppressing food intake by said subject reduces the chances of the subject developing obesity.
90. Use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating obesity in a subject.
91. The use of claim 90, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
92. The use of any one of claims 90-91, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
93. The use of any one of claims 90-92, wherein at least one CTP is glycosylated.
94. The use of any one of claims 90-93, wherein at least one CTP is truncated.
95. The use of any one of claims 90-94, wherein at least one CTP is attached to said agonist via a linker.
96. The use of claim 95, wherein said linker is a peptide bond.
97. The use of any one of claims 90-96, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
98. The use of any one of claims 90-97, wherein said subject is genetically predisposed to being obese.
99. The use of any one of claims 90-98, wherein said method of treating obesity results in a reduction of body weight in said subject.
100. The use of any one of claims 90-99, wherein said method of treating obesity results in body fat reduction in said subject.

101. Use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating type II diabetes in a subject.
102. The use of claim 101, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
103. The use of any one of claims 101-102, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
104. The use of any one of claims 101-103, wherein at least one CTP is glycosylated.
105. The use of any one of claims 101-104, wherein at least one CTP is truncated.
106. The use of any one of claims 101-105, wherein at least one CTP is attached to said agonist via a linker.
107. The use of claim 106, wherein said linker is a peptide bond.
108. The use of any one of claims 101-107, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.
109. The use of any one of claims 101-108, wherein said subject is genetically predisposed to being obese.
110. Use of a CTP-modified polypeptide comprising a GLP-1/Glucagon receptor agonist and at least one chorionic gonadotrophin carboxy terminal peptide attached to the amino terminus or carboxy terminus of said agonist for treating a metabolic disorder in a subject.
111. The use of claim 110, wherein said GLP-1/Glucagon receptor agonist is oxyntomodulin.
112. The use of any one of claims 110-111, wherein the amino acid sequence of at least one CTP is selected from the group consisting of: SEQ ID NO: 1 and SEQ ID NO: 2.
113. The use of any one of claims 110-112, wherein at least one CTP is glycosylated.
114. The use of any one of claims 110-113, wherein at least one CTP is truncated.
115. The use of any one of claims 110-114, wherein at least one CTP is attached to said agonist via a linker.
116. The use of claim 115, wherein said linker is a peptide bond.
117. The use of any one of claims 110-116, wherein the amino acid sequence of said CTP-modified polypeptide comprises SEQ ID NO: 9, 15, 21, 23, 25, 27, or 29.

118. The use of any one of claims 110-117, wherein said metabolic disorder is diabetic ketoacidosis, diabetes mellitus, or a combination thereof.
119. The use of any one of claims 110-118, wherein said metabolic disorder results from a lack of insulin and an overabundance of glucose in said subject.




Figure 1A




Figure 1B











Figure 2



Continue Figure 2




FIGURE 3A




FIGURE 3B




FIGURE 3C




Figure 4A




Figure 4B




Figure 4D




Figure 4C




Figure 5A




Figure 5B

## INTERNATIONAL SEARCH REPORT

International application no.

PCT/IL2013/050332

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 38/24 (2013.01)

USPC - 424/198.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A61K 38/24; C07K 14/59, 19/00 (2013.01)

USPC - 424/192.1, 198.1; 435/69.7; 530/397, 398; 930/110

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

CPC - A61K 38/00; C07K 14/59, 16/26, 2319/00; Y10S 930/11 (2013.01)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Patbase, Google Patent, PubMed

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category*       | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No.                                                                                |
|-----------------|------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| X<br>-----<br>Y | US 2012/0015437 A1 (FARES et al) 19 January 2012 (19.01.2012) entire document      | 1, 3, 22, 24-30, 37, 39,<br>45, 47, 53, 55, 61, 63, 69,<br>71, 80, 82, 90, 92, 101,<br>103, 110, 112 |
| Y               | US 2011/0034374 A1 (BLOOM et al) 10 February 2011 (10.02.2011) entire document     | 2, 23, 38, 46, 54, 62, 70,<br>81, 91, 102, 111                                                       |
| A               | US 5,792,460 A (BOIME) 11 August 1998 (11.08.1998) entire document                 | 2, 23, 38, 46, 54, 62, 70,<br>81, 91, 102, 111                                                       |
|                 |                                                                                    | 1-3, 22-30, 37-39, 45-47<br>53-55, 61-63, 69-71,<br>80-82, 90-92, 101-103,<br>110-112                |

☐ Further documents are listed in the continuation of Box C.


## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

08 August 2013

Date of mailing of the international search report

26 AUG 2013

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents  
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Blaine R. Copenheaver

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

# INTERNATIONAL SEARCH REPORT

international application no.

PCT/IL2013/050332

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-21, 31-36, 40-44, 48-52, 56-60, 64-68, 72-79, 83-89, 93-100, 104-109, 113-119  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

**Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)**

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing filed or furnished:

a. (means)

☐

on paper

☒

in electronic form

b. (time)

☒

in the international application as filed

☐

together with the international application in electronic form

☐

subsequently to this Authority for the purposes of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

Specifically, SEQ ID NOs: 1 and 2 were searched.

## 摘要

本發明涉及絨毛膜促性腺激素羧基末端肽(CTP)修飾的雙重 GLP-1/胰高血糖素受體激動劑及其生產和使用方法。在一個實施例中，本發明提供了 CTP 修飾的多肽，所述 CTP 修飾的多肽包含雙重 GLP-1/胰高血糖素受體激動劑和附著至所述激動劑的氨基末端或羧基末端的至少一個絨毛膜促性腺激素羧基末端肽(CTP)。