

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2012243714 B2**

**(54) Title**  
**Herbicidal composition comprising flazasulfuron and an inhibitor of protoporphyrinogen oxidase**

**(51) International Patent Classification(s)**

<b>A01P 13/00</b> (2006.01)	<b>A01N 43/76</b> (2006.01)
<b>A01N 43/54</b> (2006.01)	<b>A01N 43/82</b> (2006.01)
<b>A01N 43/56</b> (2006.01)	<b>A01N 43/84</b> (2006.01)
<b>A01N 43/58</b> (2006.01)	<b>A01N 43/90</b> (2006.01)
<b>A01N 43/653</b> (2006.01)	<b>A01N 47/36</b> (2006.01)

**(21) Application No:** **2012243714**      **(22) Date of Filing:** **2012.04.06**

**(87) WIPO No:** **WO12/141276**

**(30) Priority Data**

<b>(31) Number</b>	<b>(32) Date</b>	<b>(33) Country</b>
<b>2011-087546</b>	<b>2011.04.11</b>	<b>JP</b>

**(43) Publication Date:** **2012.10.18**

**(44) Accepted Journal Date:** **2015.01.15**

**(71) Applicant(s)**

**Ishihara Sangyo Kaisha, Ltd.**

**(72) Inventor(s)**

**Yamada, Ryu;Okamoto, Hiroyuki;Terada, Takashi**

**(74) Agent / Attorney**

**Davies Collison Cave, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000**

**(56) Related Art**

**D2 : WO 2007/105377 A2 (ISHIHARA SANGYO KAISHA LTD) 20 September 2007**

**D3 : DE 19933702 A1 (NOVARTIS AG) 11 November 1999**

**D1 : US 2002/0004457 A1 (NEVILL ET AL) 10 January 2002**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(10) International Publication Number

WO 2012/141276 A1

(43) International Publication Date  
18 October 2012 (18.10.2012)

(51) International Patent Classification:  
*A01P 13/00* (2006.01) *A01N 43/653* (2006.01)  
*A01N 47/36* (2006.01) *A01N 43/76* (2006.01)  
*A01N 43/54* (2006.01) *A01N 43/82* (2006.01)  
*A01N 43/56* (2006.01) *A01N 43/84* (2006.01)  
*A01N 43/58* (2006.01) *A01N 43/90* (2006.01)

(21) International Application Number:  
PCT/JP2012/060090

(22) International Filing Date:  
6 April 2012 (06.04.2012)

(25) Filing Language:  
English

(26) Publication Language:  
English

(30) Priority Data:  
2011-087546 11 April 2011 (11.04.2011) JP

(71) Applicant (for all designated States except US): **ISHIHARA SANGYO KAISHA, LTD.** [JP/JP]; 3-15, Edobori 1-chome, Nishi-ku, Osaka-shi, Osaka, 5500002 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **YAMADA, Ryu** [JP/JP]; c/o Ishihara Sangyo Kaisha, Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga, 5250025 (JP). **OKAMOTO, Hiroyuki** [JP/JP]; c/o Ishihara Sangyo Kaisha, Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga, 5250025 (JP). **TERADA, Takashi** [JP/JP]; c/o Ishihara Sangyo Kaisha, Ltd. Chuo Kenkyusho, 3-1, Nishi-shibukawa 2-chome, Kusatsu-shi, Shiga, 5250025 (JP).

(74) Agents: **SENMYO, Kenji** et al.; 4th Floor, SIA Kanda Square, 17, Kanda-konyacho, Chiyoda-ku, Tokyo, 1010035 (JP).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BII, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SF, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2012/141276 A1

(54) Title: HERBICIDAL COMPOSITION COMPRISING FLAZASULFURON AND AN INHIBITOR OF PROTOPORPHYRINOGEN OXIDASE

(57) Abstract: Many herbicidal compositions have been developed and are presently used. However, weeds to be controlled are various in types and their emergence extends over a long period. Accordingly, it is desired to develop a herbicidal composition which has a broad herbicidal spectrum, a high activity and a long-lasting effect. The present invention provides a herbicidal composition comprising (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide compound, a pyrimidindione compound, an oxadiazole compound, an oxazolidinedione compound, a thiadiazole compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts. According to the present invention, a herbicidal composition which has a broad herbicidal spectrum, a high activity and a long-lasting effect can be provided.

## DESCRIPTION

### TITLE OF INVENTION: HERBICIDAL COMPOSITION COMPRISING FLAZASULFURON AND AN INHIBITOR OF PROTOPORPHYRINOGEN OXIDASE

#### TECHNICAL FIELD

The present invention relates to a herbicidal composition comprising flazasulfuron or its salt and a protoporphyrinogen oxidase inhibitor.

#### BACKGROUND ART

Various herbicidal compositions have been studied to control undesired plants (hereinafter sometimes referred to simply as "weeds") in agricultural fields and non-agricultural fields.

For example, Patent Document 1 discloses microgranules comprising a compound which inhibits protoporphyrinogen oxidase when absorbed from the stem and leaves, a photosynthesis inhibiting herbicide and an acetolactate synthase inhibiting herbicide, which are to be directly applied to plants to be controlled. Patent Documents 2 and 3 also disclose various herbicidal compositions, and as one example, a combination of a protoporphyrinogen oxidase inhibitor and an acetolactate synthase inhibiting herbicide is disclosed.

However, a herbicidal composition comprising flazasulfuron or its salt and a protoporphyrinogen oxidase inhibitor is not disclosed in any of Patent Documents 1 to 3.

#### PRIOR ART DOCUMENTS

##### PATENT DOCUMENTS

Patent Document 1: JP-A-2005-68121

Patent Document 2: WO2003/024221

Patent Document 3: WO00/27203

#### DISCLOSURE OF INVENTION

Many herbicidal compositions have been developed and are presently used, but as weeds to be controlled are many in types and their emergence extends over a long period, it is desired to develop a herbicidal composition having a broader herbicidal spectrum, a high activity and a long-lasting effect.

It is possible to provide a herbicidal composition having a broader herbicidal spectrum, a high activity and a long-lasting effect, by use of flazasulfuron or its salt and a specific protoporphyrinogen oxidase inhibitor in combination.

That is, the present invention provides a herbicidal composition comprising (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide compound, a pyrimidindione compound, an oxadiazole compound, an oxazolidinedione compound, a thiadiazole compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts. Further, the present invention provides a method for controlling undesired plants or inhibiting their growth, which comprises applying a herbicidally effective amount of the above herbicidal composition.

The present invention further provides a method for controlling undesired plants or inhibiting their growth, which comprises applying herbicidally effective amounts of (A) and (B), to the undesired plants or to a place where they grow.

## 5 ADVANTAGEOUS EFFECTS OF INVENTION

The herbicidal composition of the present invention comprising flazasulfuron or its salt and a specific protoporphyrinogen oxidase inhibitor as active ingredients is capable of controlling a wide range of undesired plants in cropland or non-cropland, and it surprisingly presents a synergistic herbicidal effect i.e. a herbicidal effect higher than the mere addition of the respective herbicidal effects of the active ingredients, and it can be applied at a low dose as compared with a case where the respective active ingredients are applied individually. Such a herbicidal composition of the present invention has an enlarged herbicidal spectrum, and further its herbicidal effect will last over a long period of time.

15 When the herbicidal activity in a case where two active ingredients are combined, is larger than the simple sum of the respective herbicidal activities of the two active ingredients (the expected activity), it is called a synergistic effect. The activity expected by the combination of two active ingredients can be calculated as follows (Colby S.R., "Weed", vol. 15, p. 20-22, 1967).

$$20 E = \alpha + \beta - (\alpha \times \beta \div 100)$$

where  $\alpha$ : growth inhibition rate when treated with  $x$  (g/ha) of herbicide X,

$\beta$ : growth inhibition rate when treated with  $y$  (g/ha) of herbicide Y,

$E$ : growth inhibition rate expected when treated with  $x$  (g/ha) of herbicide X and  $y$  (g/ha) of herbicide Y.

25 That is, when the actual growth inhibition rate (measured value) is larger than the growth inhibition rate by the above calculation (calculated value), the activity by the combination can be regarded as showing a synergistic effect. The herbicidal composition of the present invention shows a synergistic effect when calculated by the above formula.

## 30 DESCRIPTION OF EMBODIMENTS

The herbicidal composition of the present invention comprises, as active ingredients, (A) flazasulfuron or its salt (hereinafter sometimes referred to as "compound A") and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide compound, a pyrimidindione compound, an oxadiazole compound, an oxazolidinedione compound, a thiadiazole compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts (hereinafter they will sometimes be referred to as "compound B").

40 In the compound A, flazasulfuron (common name) is 1-(4,6-dimethoxypyrimidin-2-yl)-3-(3-trifluoromethyl-2-pyridylsulfonyl)urea.

The compound B will be described in detail below. The compound B is represented by common names.

45 The phenylpyrazole compound may, for example, be pyraflufen-ethyl or fluazolate. Among them, pyraflufen-ethyl is preferred.

The triazolinone compound may, for example, be azafenidin, bencarbazone, carfentrazone-ethyl or sulfentrazone. Among them, azafenidin, carfentrazone-ethyl or sulfentrazone is preferred, and carfentrazone-ethyl or sulfentrazone is more preferred.

The N-phenylphthalimide compound may, for example, be cinidon-ethyl, flumiclorac-pentyl or flumioxazin. Among them, flumioxazin or flumiclorac-pentyl is preferred, and flumioxazin is more preferred.

The pyrimidindione compound may, for example, be benzfendizone, butafenacil, saflufenacil or ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy)pyridin-2-yloxy]acetate (test code: SYN-523). Among them, butafenacil or saflufenacil is preferred.

The oxadiazole compound may, for example, be oxadiargyl or oxadiazon. Among them, oxadiargyl is preferred.

10 The oxazolidinedione compound may, for example, be pentoxazone.

The thiadiazole compound may, for example, be fluthiacet-methyl or thidiazimin. Among them, fluthiacet-methyl is preferred.

15 Other compounds included in the compound B may, for example, be pyraclonil, profluazol and flufenpyr-ethyl. Among them, pyraclonil or flufenpyr-ethyl is preferred.

The compound B is preferably the phenylpyrazole compound, the triazolinone compound, the N-phenylphthalimide compound, the pyrimidindione compound or the oxadiazole compound, more preferably the phenylpyrazole compound, the triazolinone compound or the N-phenylphthalimide compound, capable of achieving a high herbicidal effect when combined with the compound A.

20 More specifically, preferred is pyraflufen-ethyl, carfentrazone-ethyl, sulfentrazone, flumioxazin, saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy)pyridin-2-yloxy]acetate (test code: SYN-523), oxadiargyl, fluthiacet-methyl, flufenpyr-ethyl, flumiclorac-pentyl, azafenidin, butafenacil, pentoxazone or pyraclonil, and more preferred is pyraflufen-ethyl, carfentrazone-ethyl, sulfentrazone, flumioxazin, saflufenacil, oxadiargyl, fluthiacet-methyl, flufenpyr-ethyl, butafenacil, pentoxazone or pyraclonil.

25 The salt included in the compound A or the compound B may be any salt so long as it is agriculturally acceptable. Examples thereof include alkali metal salts such as a sodium salt and a potassium salt; alkaline earth metal salts such as a magnesium salt and a calcium salt; ammonium salts such as a monomethylammonium salt, a dimethylammonium salt and a triethylammonium salt; inorganic acid salts such as a hydrochloride, a perchlorate, a sulfate and a nitrate, and organic acid salts such as an acetate and a methanesulfonate.

30 The mixing ratio of the compound A to the compound B cannot generally be defined, as it may vary depending upon various conditions such as the type of the formulation, the weather conditions, and the type and the growth stage of the undesired plants to be controlled, but it is preferably a mixing ratio to achieve the herbicidally effective amounts (synergistic herbicidally effective amount) with which the synergistic herbicidal effect is obtained, and for example, by the weight ratio, it is preferably from 100:1 to 1:100, more preferably from 50:1 to 1:64, particularly preferably from 20:1 to 1:32.

35 When pyraflufen-ethyl is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 20:1 to 1:8, more preferably from 20:1 to 1:3.2.

40 When carfentrazone-ethyl is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 40:1 to 1:40, more preferably from 20:1 to 1:18, particularly preferably from 5:1 to 1:18.

45 When sulfentrazone is used as the compound B, the mixing ratio of the

compound A to the compound B is, for example, by the weight ratio, preferably from 10:1 to 1:50, more preferably from 4:1 to 1:25, particularly preferably from 2:1 to 1:20.

When flumioxazin is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 100:1 to 1:75, more preferably from 50:1 to 1:24, particularly preferably from 20:1 to 1:20.

When saflufenacil is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 20:1 to 1:10, more preferably from 10:1 to 1:8.

When oxadiargyl is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 2:1 to 1:100, more preferably from 1:1 to 1:32.

When fluthiacet-methyl is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 20:1 to 1:10, more preferably from 10:1 to 1:2.

When flufenpyr-ethyl is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 20:1 to 1:10, more preferably from 10:1 to 1:2.

When butafenacil is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 20:1 to 1:25, more preferably from 5:1 to 1:8.

When pentoxazone is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 2:1 to 1:50, more preferably from 0.5:1 to 1:10.

When pyraclonil is used as the compound B, the mixing ratio of the compound A to the compound B is, for example, by the weight ratio, preferably from 2:1 to 1:50, more preferably from 0.5:1 to 1:10.

The doses of the compound A and the compound B cannot generally be defined, as they may vary depending upon various conditions such as the mixing ratio of the compound A to the compound B, the type of the formulation, the weather conditions, and the type and the growth stage of the undesired plants to be controlled. However, they are preferably doses to achieve the herbicidally effective amounts (synergistic herbicidally effective amount) with which the synergistic herbicidal effect is obtained, and for example, the dose of the compound A is preferably from 0.5 to 120 g/ha, more preferably from 1 to 110 g/ha, particularly preferably from 1 to 100 g/ha, and the dose of the compound B is preferably from 0.5 to 1,000 g/ha, more preferably from 1 to 900 g/ha, particularly preferably from 2 to 800 g/ha.

With respect to the doses of the compounds A and B when pyraflufen-ethyl is used as the compound B, for example, the dose of the compound A is preferably from 0.5 to 120 g/ha, more preferably from 1 to 110 g/ha, particularly preferably from 1 to 100 g/ha, and the dose of the compound B is preferably from 1 to 100 g/ha, more preferably from 1 to 90 g/ha, particularly preferably from 2 to 80 g/ha.

With respect to the doses of the compounds A and B when carfentrazone-ethyl is used as the compound B, for example, the dose of the compound A is preferably from 0.5 to 120 g/ha, more preferably from 0.5 to 110 g/ha, particularly preferably from 1 to 100 g/ha, and the dose of the compound B is preferably from 2.5 to 400 g/ha, more preferably from 10 to 250 g/ha, particularly preferably from 10 to 50 g/ha.

With respect to the doses of the compounds A and B when sulfentrazone is used as the compound B, for example, the dose of the compound A is preferably from 10 to

100 g/ha, more preferably from 20 to 100 g/ha, particularly preferably from 25 to 100 g/ha, and the dose of the compound B is preferably from 10 to 500 g/ha, more preferably from 25 to 500 g/ha.

With respect to the doses of the compounds A and B when flumioxazin is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 25 to 100 g/ha, particularly preferably from 25 to 50 g/ha, and the dose of the compound B is preferably from 1 to 750 g/ha, more preferably from 2 to 600 g/ha, particularly preferably from 2.5 to 500 g/ha.

With respect to the doses of the compounds A and B when saflufenacil is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 12.5 to 100 g/ha, and the dose of the compound B is preferably from 5 to 100 g/ha.

With respect to the doses of the compounds A and B when oxadiargyl is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 12.5 to 100 g/ha, and the dose of the compound B is preferably from 50 to 1,000 g/ha, more preferably from 150 to 800 g/ha.

With respect to the doses of the compounds A and B when fluthiacet-methyl is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 25 to 50 g/ha, and the dose of the compound B is preferably from 5 to 100 g/ha, more preferably from 5 to 50 g/ha.

With respect to the doses of the compounds A and B when flufenpyr-ethyl is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 25 to 50 g/ha, and the dose of the compound B is preferably from 5 to 100 g/ha, more preferably from 5 to 50 g/ha.

With respect to the doses of the compounds A and B when butafenacil is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 12.5 to 50 g/ha, and the dose of the compound B is preferably from 5 to 250 g/ha, more preferably from 10 to 100 g/ha.

With respect to the doses of the compounds A and B when pentozaone is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 25 to 50 g/ha, and the dose of the compound B is preferably from 50 to 500 g/ha, more preferably from 100 to 250 g/ha.

With respect to the doses of the compounds A and B when pyraclonil is used as the compound B, for example, the dose of the compound A is preferably from 10 to 100 g/ha, more preferably from 25 to 50 g/ha, and the dose of the compound B is preferably from 50 to 500 g/ha, more preferably from 100 to 250 g/ha.

The herbicidal composition of the present invention may be applied to undesired plants or may be applied to a place where they grow. Further, it may be applied at any time either before or after the emergence of the undesired plants. Further, the herbicidal composition of the present invention may take various application forms such as soil application, foliar application, irrigation application, and submerged application, and it can be applied to agricultural fields such as upland fields, orchards and paddy fields, and non-cropland such as ridges of fields, fallow fields, play grounds, golf courses, vacant lands, forests, factory sites, railway sides and roadsides.

The herbicidal composition of the present invention can control a broad range of undesired plants such as annual weeds and perennial weeds. The undesired plants to be controlled by the herbicidal composition of the present invention may, for example, be cyperaceae such as green kyllinga (Kyllinga brevifolia Rottb. var.

leolepis), or sedge (*Cyperus* spp.) [the sedge may, for example, be purple nutsedge (*Cyperus rotundus* L.), smallflower umbrella sedge (*Cyperus difformis* L.), yellow nutsedge (*Cyperus esculentus* L.) or amur cyperus (*Cyperus microtia* Steud.)]; *gramineae* such as barnyardgrass (*Echinochloa crus-galli* L., *Echinochloa oryzicola* vasing.), japanese millet (*Echinochloa utilis* Ohwi et Yabuno), crabgrass (*Digitaria* spp.) [the crabgrass may, for example, be summergrass (*Digitaria ciliaris* (Retz.) Koel), large crabgrass (*Digitaria sanguinalis* L.), violet crabgrass (*Digitaria violascens* Link) or *Digitaria horizontalis* Willd.], green foxtail (*Setaria viridis* L.), giant foxtail (*Setaria faberii* Herrm.), goosegrass (*Eleusine indica* L.), johnsongrass (*Sorghum halepense* (L.) Pers.), bermudagrass (*Cynodon dactylon* (L.) Pers.), wild oat (*Avena fatua* L.), annual bluegrass (*Poa annua* L.), panic grass (*Panicum* spp.) [the panic grass may, for example, be guinea grass (*Panicum maximum* Jacq.), or fall panicum (*Panicum dichotomiflorum* (L.) Michx.)], signal grass (*Brachiaria* spp.) [the signal grass may, for example, be plantain signal grass (*Brachiaria plantaginea* (LINK) Hitchc.), palisade signal grass (*Brachiaria decumbens* Stapf), or mauritius signal grass (*Brachiaria mutica* (Forssk.) Stapf)], paspalum (*Paspalum* spp.), itchgrass (*Rottboellia cochinchinensis* (LOUR.) W.D.CLAYTON); southern sandbur (*Cenchrus echinatus* L.), or shattercane (*Sorghum bicolor* (L.) Moench.); *scrophulariaceae* such as persian speedwell (*Veronica persica* Poir.), or corn speedwell (*Veronica arvensis* L.); *compositae* such as beggar ticks (*Bidens* spp.) [the beggar ticks may, for example, be hairy beggarticks (*Bidens pilosa* L.), devils berggarticks (*Bidens frondosa* L.), *Bidens biternata* (Lour.) Merr. et Sherff ], or beggarticks (*Bidens subalternans* DC.)], hairy fleabane (*Conyza bonariensis* (L.) Cronq.), horseweed (*Erigeron canadensis* L.), dandelion (*Taraxacum officinale* Weber), or common cocklebur (*Xanthium strumarium* L.); *leguminosae* such as rattlepod or rattlebox (*Crotalaria* spp.) [the rattlepod or rattlebox may, for example, be sunn-hemp (*Crotalaria juncea* L.)], poison bean (*Sesbania* spp.) [the poison bean may, for example, be rostrate sesbania (*Sesbania rostrata* Bremek. & Oberm.) or sesbania pea (*Sesbania cannabina* (Retz.) Pers.)], white clover (*Trifolium repens* L.); *caryophyllaceae* such as sticky chickweed (*Cerastium glomeratum* Thuill.), or common chickweed (*Stellaria media* L.); *euphorbiaceae* such as garden spurge (*Euphorbia hirta* L.), threeseeded copperleaf (*Acalypha australis* L.), or fireplant (*Euphorbia heterophylla* L.); *plantaginaceae* such as asiatic plantain (*Plantago asiatica* L.); *oxalidaceae* such as creeping woodsorrel (*Oxalis corniculata* L.); *apiaceae* such as lawn pennywort (*Hydrocotyle sibthorpioides* Lam.); *violaceae* such as violet (*Viola mandshurica* W. Becker); *iridaceae* such as blue-eyedgrass (*Sisyrinchium rosulatum* Bicknell); *geraniaceae* such as carolina geranium (*Geranium carolinianum* L.); *labiatae* such as purple deadnettle (*Lamium purpureum* L.), or henbit (*Lamium amplexicaule* L.); *malvaceae* such as velvetleaf (*Abutilon theophrasti* MEDIC.), or prickly sida (*Sida spinosa* L.); *convolvulaceae* such as ivy-leaved morningglory (*Ipomoea hederacea* (L.) Jacq.), common morningglory (*Ipomoea purpurea* ROTH), cypressvine morningglory (*Ipomoea quamoclit* L.), *Ipomoea grandifolia* (DAMMERMANN) O'DONNELL, hairy merremia (*Merremia aegyptia* (L.) URBAN), or field Bindweed (*Convolvulus arvensis* L.); *chenopodiaceae* such as common lambsquarters (*Chenopodium album* L.); *portulacaceae* such as common purslane (*Portulaca oleracea* L.); *amaranthaceae* such as pigweed (*Amaranthus* spp.) [the pigweed may, for example, be prostrate pigweed (*Amaranthus blitoides* S. Wats.), livid amaranth (*Amaranthus lividus* L.), purple amaranth (*Amaranthus blitum* L.), smooth pigweed (*Amaranthus hybridus* L.), *Amaranthus patulus* Bertol., powell amaranth (*Amaranthus powellii* S.Wats.), slender

amaranth (Amaranthus viridis L.), palmer amaranth (Amaranthus palmeri S.Wats.), redroot pigweed (Amaranthus retroflexus L.), tall waterhemp (Amaranthus tuberculatus (Moq.) Sauer.), common waterhemp (Amaranthus tamariscinus Nutt.), thorny amaranth (Amaranthus spinosus L.), ataco (Amaranthus quitensis Kunth.), or Amaranthus rudis Sauer]; solanaceae such as black nightshade (Solanum nigrum L.); polygonaceae such as spotted knotweed (Polygonum lapathifolium L.), or green smartweed (Polygonum scabrum MOENCH); cruciferae such as flexuous bittercress (Cardamine flexuosa WITH.); cucurbitaceae such as burcucumber (Sicyos angulatus L.); or commelinaceae such as common dayflower (Commelina communis L.); rosaceae such as mock strawberry (Duchesnea chrysanthia (Zoll. et Mor. ) Miq.); molluginacea such as carpetweed (Mollugo verticillata L.); or rubiaceae such as false cleavers (Galium spurium var. echinospermon (Wallr.) Hayek) or stickywilly (Galium aparine L.).

The herbicidal composition of the present invention can control even weeds against which the compound A has no satisfactory controlling effects depending upon various conditions such as the weather conditions, and the growth stage of the weeds. For example, the compound A has no satisfactory controlling effects against some weeds included in solanaceae, scrophulariaceae and gramineae in some cases depending upon various conditions such as the weather conditions and the growth stage of the weeds, however, the herbicidal composition of the present invention comprising the combination of the compounds A and B has excellent effect to control such weeds or to inhibit their growth.

Further, the herbicidal composition of the present invention can control perennial grass weeds such as quackgrass (Agropyron repens (L.) P. Beauv.), Cholorado bluestem (Agropyron tsukushense (Honda) Ohwi var. transiens (Hack.) Ohwi), redtop (Agrostis alba L.), orchardgrass (Dactylis glomerata L.), perennial ryegrass (Lolium perenne L.), eulaliagrass (Miscanthus sinensis Anderss.), knotgrass (Paspalum distichum L.), bahiagrass (Paspalum notatum Flugge), johnsongrass (Sorghum halepense L.), bermuda grass (Cynodon dactylon (L.) Pers.), dallisgrass (Paspalum dilatatum Poir.), cogongrass (Imperata cylindrica (L.) Beauv.), japanese paspalum (Paspalum thunbergii Kunth) which are problematic as strong weeds in agricultural fields such as orchards and non-cropland such as golf courses, railway sides and roadsides. Further, the herbicidal composition of the present invention has a high herbicidal activity also against weeds in late leaf stage, such as weeds in 5-leaf stage to heading stage, and such is particularly remarkable for grass weeds. The herbicidal composition of the present invention has favorable herbicidal effects against grass weeds and broad leaf weeds either by foliar application or soil application.

Further, as one of cultivation manners for crop plants, different crop plants may be cultivated in the same field by differentiating timing for their cultivation. For example, in the same field where corn was cultivated last year, sugarcane may be cultivated this year, and in such a case, the previous crop plant such as the corn may be an object to be controlled as an undesired plant. Further, along with spread of genetically modified crop plants or increase of cultivation fields, there may be a case where at the time of repeated cultivation, rotational cropping or change in cropping, the previous crop plant grown as weeds (volunteer crop plant) becomes an object to be controlled as an undesired plant. Even in such a situation, the herbicidal composition of the present invention is capable of controlling the undesired plant to be controlled and thus is very useful in such a practical application.

Further, in practical application in which the rapid herbicidal efficacy and regrowth

of the undesired plants after the herbicidal composition is applied are problematic, the herbicidal composition of the present invention is useful in view of the rapid herbicidal efficacy and a high effect of suppressing regrowth of the undesired plants.

The herbicidal composition of the present invention may further contain other herbicidal compounds in addition to the above-described active ingredients so long as such will meet the object of the present invention, and there may be a case where it is thereby possible to improve e.g. the range of undesired plants to be controlled, the timing for application of the herbicidal composition, the herbicidal activities, etc. to more desirable directions. Such other herbicidal compounds include, for example, the following compounds (common names including ones under application for approval by ISO, or test codes, here, "under application for approval by ISO" means common names before approval by ISO (International Organization for Standardization)), and one or more of them may suitably be selected for use. Even when not specifically mentioned here, in a case where such compounds have salts, alkyl esters, hydrates, different crystal forms, various structural isomers, etc., they are, of course, all included.

Further, in consideration of the application site of the herbicidal composition or the type or growth state of the undesired plants, the herbicidal composition of the present invention may be mixed with or may be used in combination with fungicides, antibiotics, plant hormones, insecticides, fertilizers, phytotoxicity-reducing agents, etc., whereby more excellent effects and activities may sometimes be obtained.

(1) Those which are believed to exhibit herbicidal effects by disturbing hormone activities of plants, such as a phenoxy type such as 2,4-D, 2,4-D-butyl, 2,4-D-butyl, 2,4-D-dimethylammonium, 2,4-D-diolamine, 2,4-D-ethyl, 2,4-D-2-ethylhexyl, 2,4-D-isobutyl, 2,4-D-isooctyl, 2,4-D-isopropyl, 2,4-D-isopropylammonium, 2,4-D-sodium, 2,4-D-isopropanolammonium, 2,4-D-trolamine, 2,4-DB, 2,4-DB-butyl, 2,4-DB-dimethylammonium, 2,4-DB-isooctyl, 2,4-DB-potassium, 2,4-DB-sodium, dichlorprop, dichlorprop-butyl, dichlorprop-dimethylammonium, dichlorprop-isooctyl, dichlorprop-potassium, dichlorprop-P, dichlorprop-P-dimethylammonium, dichlorprop-P-potassium, dichlorprop-P-sodium, MCPA, MCPA-butyl, MCPA-dimethylammonium, MCPA-2-ethylhexyl, MCPA-potassium, MCPA-sodium, MCPA-thioethyl, MCPB, MCPB-ethyl, MCPB-sodium, mecoprop, mecoprop-butyl, mecoprop-sodium, mecoprop-P, mecoprop-P-butyl, mecoprop-P-dimethylammonium, mecoprop-P-2-ethylhexyl, mecoprop-P-potassium, naproanilide or clomeprin; an aromatic carboxylic acid type such as 2,3,6-TBA, dicamba, dicamba-butyl, dicamba-diglycolamine, dicamba-dimethylammonium, dicamba-diolamine, dicamba-isopropylammonium, dicamba-potassium, dicamba-sodium, dichlobenil, picloram, picloram-dimethylammonium, picloram-isooctyl, picloram-potassium, picloram-triisopropanolammonium, picloram-triisopropylammonium, picloram-trolamine, triclopyr, triclopyr-butyl, triclopyr-triethylammonium, clopyralid, clopyralid-olamine, clopyralid-potassium, clopyralid-triisopropanolammonium or aminopyralid; and others such as naptalam, naptalam-sodium, benazolin, benazolin-ethyl, quinclorac, quinmerac, diflufenzopyr, diflufenzopyr-sodium, fluroxypyr, fluroxypyr-2-butoxy-1-methylethyl, fluroxypyr-meptyl, chlorflurenol, chlorflurenol-methyl, aminocyclopyrachlor, aminocyclopyrachlor-methyl or aminocyclopyrachlor-potassium.

(2) Those which are believed to exhibit herbicidal effects by inhibiting photosynthesis of plants, such as a urea type such as chlorotoluron, diuron, fluometuron, linuron, isoproturon, metobenzuron, tebuthiuron, dimefuron, isouron, karbutilate, methabenzthiazuron, metoxuron, monolinuron, neburon, siduron,

terbumeton, trietazine or metobromuron; a triazine type such as simazine, atrazine, atratone, simetryn, prometryn, dimethametryn, hexazinone, metribuzin, terbutylazine, cyanazine, ametryn, cybutryne, triaziflam, indaziflam, terbutryn, propazine, metamitron or prometon; a uracil type such as bromacil, bromacyl-lithium, lenacil or terbacil; an anilide type such as propanil or cypromid; a carbamate type such as swep, desmedipham or phenmedipham; a hydroxybenzonitrile type such as bromoxynil, bromoxynil-octanoate, bromoxynil-heptanoate, ioxynil, ioxynil-octanoate, ioxynil-potassium or ioxynil-sodium; and others such as pyridate, bentazone, bentazone-sodium, amicarbazone, methazole or pentanochlor.

(3) Quaternary ammonium salt type such as paraquat or diquat, which is believed to be converted to free radicals by itself to form active oxygen in the plant body and shows rapid herbicidal efficacy.

(4) Those which are believed to exhibit herbicidal effects by inhibiting chlorophyll biosynthesis of plants and abnormally accumulating a photosensitizing peroxide substance in the plant body, such as a diphenylether type such as nitrofen, chlomethoxyfen, bifenoxyfen, acifluorfen, acifluorfen-sodium, fomesafen, fomesafen-sodium, oxyfluorfen, lactofen, aclonifen, ethoxyfen-ethyl (HC-252), fluoroglycofen-ethyl or fluoroglycofen; a cyclic imide type such as chlorphthalim; and others such as isopropazole or flupoxam.

(5) Those which are believed to exhibit herbicidal effects characterized by bleaching activities by inhibiting chromogenesis of plants such as carotenoids, such as a pyridazinone type such as norflurazon, chloridazon or metflurazon; a pyrazole type such as pyrazolynate, pyrazoxyfen, benzofenap, topramezone or pyrasulfotole; and others such as amitrole, fluridone, flurtamone, diflufenican, methoxyphenone, clomazone, sulcotrione, mesotrione, tembotrione, tefuryltrione (AVH-301), bicyclopyprone, isoxaflutole, difenzoquat, difenzoquat-metilsulfate, isoxachlortole, benzobicyclon, picolinafen, beflubutamid, a compound (SW-065, H-965) disclosed in the claim of WO2003/016286, a compound (KIH-3653, KUH-110) disclosed in the claim of WO2009/016841, a compound disclosed in the claim of WO2005/118530, a compound disclosed in the claim of WO2008/065907, or a compound disclosed in the claim of WO2009/142318.

(6) Those which exhibit strong herbicidal effects specifically to gramineous plants, such as an aryloxyphenoxypropionic acid type such as diclofop-methyl, diclofop, pyriphenop-sodium, fluazifop-butyl, fluazifop, fluazifop-P, fluazifop-P-butyl, haloxyfop-methyl, haloxyfop, haloxyfop-etyl, haloxyfop-P, haloxyfop-P-methyl, quizalofop-ethyl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, cyhalofop-butyl, fenoxaprop-ethyl, fenoxaprop-P, fenoxaprop-P-ethyl, metamifop-propyl, metamifop, clodinafop-propargyl, clodinafop or propaquizafop; a cyclohexanedione type such as aloxydim-sodium, aloxydim, clethodim, sethoxydim, tralkoxydim, butroxydim, tepraloxydim, profoxydim or cycloxydim; and others such as flamprop-M-methyl, flamprop-M or flamprop-M-isopropyl.

(7) Those which are believed to exhibit herbicidal effects by inhibiting an amino acid biosynthesis of plants, such as a sulfonylurea type such as chlormuron-ethyl, chlorimuron, sulfometuron-methyl, sulfometuron, primisulfuron-methyl, primisulfuron, bensulfuron-methyl, bensulfuron, chlorsulfuron, metsulfuron-methyl, metsulfuron, cinosulfuron, pyrazosulfuron-ethyl, pyrazosulfuron, azimsulfuron, rimsulfuron, nicosulfuron, imazosulfuron, cyclosulfamuron, prosulfuron, flupysulfuron-methyl-sodium, flupysulfuron, triflusulfuron-methyl, triflusulfuron, halosulfuron-methyl,

halosulfuron, thifensulfuron-methyl, thifensulfuron, ethoxysulfuron, oxasulfuron, ethametsulfuron, ethametsulfuron-methyl, iodosulfuron, iodosulfuron-methyl-sodium, sulfosulfuron, triasulfuron, tribenuron-methyl, tribenuron, tritosulfuron, foramsulfuron, trifloxsulfuron, trifloxsulfuron-sodium, mesosulfuron-methyl, mesosulfuron, 5 orthosulfamuron, flucetosulfuron, amidosulfuron, propyrisulfuron (TH-547), metazosulfuron, iofensulfuron, or a compound disclosed in the claim of EP0645386; a triazolopyrimidinesulfonamide type such as flumetsulam, metosulam, diclosulam, cloransulam-methyl, florasulam, penoxsulam or pyroxsulam; an imidazolinone type such as imazapyr, imazapyr-isopropylammonium, imazethapyr, imazethapyr- 10 ammonium, imazaquin, imazaquin-ammonium, imazamox, imazamox-ammonium, imazamethabenz, imazamethabenz-methyl or imazapic; a pyrimidinylsalicylic acid type such as pyrithiobac-sodium, bispyribac-sodium, pyriminobac-methyl, pyribenzoxim, pyriftalid or pyrimisulfan; a sulfonylaminocarbonyltriazolinone type such as flucarbazone, flucarbazone-sodium, propoxycarbazone-sodium, propoxycarbazone or thiencarbazone; and others such as glyphosate, glyphosate-sodium, glyphosate- 15 potassium, glyphosate-ammonium, glyphosate-diammonium, glyphosate-isopropylammonium, glyphosate-trimesium, glyphosate-sesquisodium, glufosinate, glufosinate-ammonium, glufosinate-P, glufosinate-P-ammonium, glufosinate-P-sodium, bilanafos, bilanafos-sodium, cinmethylin or triafamone.

20 (8) Those which are believed to exhibit herbicidal effects by inhibiting cell mitoses of plants, such as a dinitroaniline type such as trifluralin, oryzalin, nitrinal, pendimethalin, ethalfluralin, benfluralin, prodiamine, butralin or dinitramine; an amide type such as bensulide, napropamide, propyzamide or pronamide; an organic phosphorus type such as amiprofos-methyl, butamifos, anilofos or piperophos; a 25 phenyl carbamate type such as prophan, chlorprophan, barban or carbetamide; a cumylamine type such as daimuron, cumyluron, bromobutide or methyldymron; and others such as asulam, asulam-sodium, dithiopyr, thiazopyr, chlorthal-dimethyl, chlorthal or diphenamid.

30 (9) Those which are believed to exhibit herbicidal effects by inhibiting protein biosynthesis or lipid biosynthesis of plants, such as a chloroacetamide type such as alachlor, metazachlor, butachlor, pretilachlor, metolachlor, S-metolachlor, thenylchlor, pethoxamid, acetochlor, propachlor, dimethenamid, dimethenamid-P, propisochlor or dimethachlor; a thiocarbamate type such as molinate, dimepiperate, pyributicarb, EPTC, butylate, vernolate, pebulate, cycloate, prosulfocarb, esprocarb, thiobencarb, 35 diallate, tri-allate or orbencarb; and others such as etobenzanid, mefenacet, flufenacet, tridiphane, cafenstrole, fentrazamide, oxaziclofone, indanofan, benfuresate, pyroxasulfone, fenoxasulfone, dalapon, dalapon-sodium, TCA-sodium or trichloroacetic acid.

40 (10) MSMA, DSMA, CMA, endothall, endothall-dipotassium, endothall-sodium, endothall-mono(N,N-dimethylalkylammonium), ethofumesate, sodium chlorate, pelargonic acid (nonanoic acid), fosamine, fosamine-ammonium, pinoxaden, ipfencarbazone (HOK-201), aclolein, ammonium sulfamate, borax, chloroacetic acid, sodium chloroacetate, cyanamide, methylarsonic acid, dimethylarsinic acid, sodium dimethylarsinate, dinoterb, dinoterb-ammonium, dinoterb-diolamine, dinoterb-acetate, 45 DNO, ferrous sulfate, flupropanate, flupropanate-sodium, isoxaben, mefluidide, mefluidide-diolamine, metam, metam-ammonium, metam-potassium, metam-sodium, methyl isothiocyanate, pentachlorophenol, sodium pentachlorophenoxyde, pentachlorophenol laurate, quinoclamine, sulfuric acid, urea sulfate, methiozolin (MRC-

01), etc.

(11) Those which are believed to exhibit herbicidal effects by being parasitic on plants, such as Xanthomonas campestris, Epicoccosirus nematosorus, Epicoccosirus nematosperus, Exserohilum monoseras or Drechsrela monoceras.

5 The herbicidal composition of the present invention may be prepared by mixing the compounds A and B, as active ingredients, with various agricultural additives in accordance with conventional formulation methods for agricultural chemicals, and applied in various formulations such as dusts, granules, water dispersible granules, wettable powders, tablets, pills, capsules (including a formulation packaged by a water 10 soluble film), water-based suspensions, oil-based suspensions, microemulsions, suspoemulsions, water soluble powders, emulsifiable concentrates, soluble 15 concentrates or pastes. It may be formed into any formulation which is commonly used in this field, so long as the object of the present invention is thereby met.

15 The type of the formulation of the herbicidal composition of the present invention is preferably a liquid formulation which can be applied as it is or a formulation to be applied after diluted with water, since by the formulation such that a solid formulation such as microgranules is applied as it is without being diluted with water, application to a wide area at once is difficult, and the application will take long. More specifically, 20 water dispersible granules, wettable powders, water-based suspensions, oil-based suspensions, emulsifiable concentrates, soluble concentrates and the like are preferred.

At the time of the formulation, the compounds A and B may be mixed together for the formulation, or they may be separately formulated.

25 The additives to be used for the formulation include, for example, a solid carrier such as kaolinite, sericite, diatomaceous earth, slaked lime, calcium carbonate, talc, white carbon, kaoline, bentonite, clay, sodium carbonate, sodium bicarbonate, mirabilite, zeolite or starch; a solvent such as water, toluene, xylene, solvent naphtha, dioxane, dimethylsulfoxide, N,N-dimethylformamide, dimethylacetamide, N-methyl-2-pyrrolidone or an alcohol; an anionic surfactant such as a salt of fatty acid, a benzoate, 30 a polycarboxylate, a salt of alkylsulfuric acid ester, an alkyl sulfate, an alkylaryl sulfate, an alkyl diglycol ether sulfate, a salt of alcohol sulfuric acid ester, an alkyl sulfonate, an alkylaryl sulfonate, an aryl sulfonate, a lignin sulfonate, an alkylidiphenylether disulfonate, a polystyrene sulfonate, a salt of alkylphosphoric acid ester, an alkylaryl 35 phosphate, a styrylaryl phosphate, a salt of polyoxyethylene alkyl ether sulfuric acid ester, a polyoxyethylene alkylaryl ether sulfate, a salt of polyoxyethylene alkylaryl ether sulfuric acid ester, a polyoxyethylene alkyl ether phosphate, a salt of polyoxyethylene alkylaryl phosphoric acid ester, a salt of polyoxyethylene aryl ether phosphoric acid ester, a naphthalene sulfonic acid condensed with formaldehyde or a salt of alkylnaphthalene sulfonic acid condensed with formaldehyde; a nonionic surfactant 40 such as a sorbitan fatty acid ester, a glycerin fatty acid ester, a fatty acid polyglyceride, a fatty acid alcohol polyglycol ether, acetylene glycol, acetylene alcohol, an oxyalkylene block polymer, a polyoxyethylene alkyl ether, a polyoxyethylene alkylaryl ether, a polyoxyethylene styrylaryl ether, a polyoxyethylene glycol alkyl ether, 45 polyethylene glycol, a polyoxyethylene fatty acid ester, a polyoxyethylene sorbitan fatty acid ester, a polyoxyethylene glycerin fatty acid ester, a polyoxyethylene hydrogenated castor oil or a polyoxypropylene fatty acid ester; and a vegetable oil or mineral oil such as olive oil, kapok oil, castor oil, palm oil, camellia oil, coconut oil, sesame oil, corn oil, rice bran oil, peanut oil, cottonseed oil; soybean oil, rapeseed oil, linseed oil, tung oil or

liquid paraffins; transesterified vegetable oil such as methylated rapeseed oil or ethylated rapeseed oil. These additives may suitably be selected for use alone or in combination as a mixture of two or more of them, so long as the object of the present invention is met. Further, additives other than the above-mentioned may be suitably selected for use among those known in this field. For example, various additives commonly used, such as a filler, a thickener, an anti-settling agent, an anti-freezing agent, a dispersion stabilizer, a safener, an anti-mold agent, a bubble agent, a disintegrator and a binder, may be used. The mix ratio by weight of the active ingredient to such various additives in the herbicidal composition of the present invention may be from 0.001:99.999 to 95:5, preferably from 0.005:99.995 to 90:10.

As a method of applying the herbicidal composition of the present invention, a proper method can be employed among various methods depending upon various conditions such as the application site, the type of the formulation, and the type and the growth stage of the undesired plants to be controlled, and for example, the following methods may be mentioned.

1. The compound A and the compound B are formulated together, and the formulation is applied as it is.
2. The compound A and the compound B are formulated together, the formulation is diluted to a predetermined concentration with e.g. water, and as the case requires, a spreader (such as a surfactant, a vegetable oil or a mineral oil) is added for application.
- 20 3. The compound A and the compound B are separately formulated and applied as they are.
4. The compound A and the compound B are separately formulated, and they are diluted to a predetermined concentration with e.g. water, and as the case requires, a spreader (such as a surfactant, a vegetable oil or a mineral oil) is added for application.
- 25 5. The compound A and the compound B are separately formulated, and the formulations are mixed when diluted to a predetermined concentration with e.g. water, and as the case requires, a spreader (such as a surfactant, a vegetable oil or a mineral oil) is added for application.

30 Now, preferred embodiments of the present invention will be described below.

However, the present invention is by no means restricted thereto.

(1) A herbicidal composition comprising (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide

35 compound, a pyrimidindione compound, an oxadiazole compound, an oxazolidinedione compound, a thiadiazole compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts.

(2) The compound according to the above (1), wherein (B) is at least one member selected from the group consisting of a phenylpyrazole compound, a triazolinone

40 compound, a N-phenylphthalimide compound, a pyrimidindione compound, an oxadiazole compound and their salts.

(3) The compound according to the above (2), wherein (B) is at least one member selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide compound, a pyrimidindione compound and their salts.

(4) The compound according to the above (3), wherein (B) is at least one member selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, a N-phenylphthalimide compound and their salts.

(5) The composition according to the above (1), wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl, fluazolate, azafenidin, bencarbazone, carfentrazone-ethyl, sulfentrazone, cinidon-ethyl, flumiclorac-pentyl, flumioxazin, benzfendizone, butafenacil, saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy)pyridin-2-yloxy]acetate (test code: SYN-523), oxadiargyl, oxadiazon, pentozacone, fluthiacet-methyl, thidiazimin, pyraclonil, profluazol, flufenpyr-ethyl and their salts.

5 (6) The composition according to the above (1), wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl, carfentrazone-ethyl, sulfentrazone, flumioxazin, saflufenacil, oxadiargyl, fluthiacet-methyl, flufenpyr-ethyl, butafenacil, pentozacone, pyraclonil and their salts.

10 (7) The composition according to the above (1), wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl and carfentrazone-ethyl.

(8) The composition according to the above (1), wherein (B) is pyraflufen-ethyl.

15 (9) The composition according to the above (1), wherein (B) is carfentrazone-ethyl.

(10) The composition according to any one of the above (1) to (9), which contains synergistic herbicidally effective amounts of (A) and (B).

(11) The composition according to any one of the above (1) to (10), wherein the mixing ratio of (A) to (B) is from 100:1 to 1:100 by the weight ratio.

20 (12) A method for controlling undesired plants or inhibiting their growth, which comprises applying a herbicidally effective amount of a herbicidal composition comprising (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, an oxadiazole compound, an oxazolidinedione compound, a 25 N-phenylphthalimide compound, a thiadiazole compound, a pyrimidindione compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts, to the undesired plants or to a place where they grow.

(13) A method for controlling undesired plants or inhibiting their growth, which comprises applying herbicidally effective amounts of (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of a phenylpyrazole compound, a triazolinone compound, an oxadiazole compound, an oxazolidinedione compound, a N-phenylphthalimide compound, a thiadiazole compound, a pyrimidindione compound, pyraclonil, profluazol, flufenpyr-ethyl and their salts, to the undesired plants or to a place where they grow.

30 (14) The method according to the above (12) or (13), wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl and carfentrazone-ethyl.

(15) The method according to the above (12) or (13), wherein (B) is pyraflufen-ethyl.

(16) The method according to the above (12) or (13), wherein (B) is carfentrazone-ethyl.

40 (17) The method according to the above (12) or (13), wherein synergistic herbicidally effective amounts of (A) and (B) are applied.

(18) The method according to the above (12) or (13), wherein (A) is applied in an amount of from 0.5 to 120 g/ha, and (B) is applied in an amount of from 0.5 to 1,000 g/ha.

45 (19) The method according to the above (12) or (13), wherein (A) is applied in an amount of from 10 to 100 g/ha, and (B) is applied in an amount of from 1 to 1,000 g/ha.

(20) The method according to the above (15), wherein (A) is applied in an amount of from 10 to 100 g/ha, and (B) is applied in an amount of from 5 to 80 g/ha.

(21) The method according to the above (16), wherein (A) is applied in an amount of from 10 to 100 g/ha, and (B) is applied in an amount of from 2.5 to 400 g/ha.

In a further embodiment, the present invention provides a method for controlling undesired plants or inhibiting their growth, which comprises

(I) applying a herbicidally effective amount of a herbicidal composition comprising

- (A) flazasulfuron or its salt and
- (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of pyraflufen-ethyl, fluazolate, azafenidin, bencarbazone, carfentrazone-ethyl, sulfentrazone, cinidon-ethyl, flumiclorac-pentyl, flumioxazin, benzfendizone, butafenacil, saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy]pyridin-2-ylloxy]acetate (test code: SYN-523), oxadiargyl, oxadazon, pentozacone, fluthiacet-methyl, thidiazimin, pyraclonil, profluazol and flufenpyr-ethyl and their salts, or which comprises

(II) applying herbicidally effective amounts of components (A) and (B), to the undesired plants or to a place where they grow, wherein the undesired plants is persian speedwell, black nightshade, sticky chickweek, common lambsquarters, wild oat, rostrate sesbania, sunn-hemp, velvetleaf, bermudagrass, japanese millet, corn, ivy-leaved morningglory or shattercane, and wherein said composition presenting a synergistic herbicidal effect, or wherein the combination of components (A) and (B) presenting a synergistic herbicidal effect.

In a further embodiment, the present invention provides A herbicidal composition comprising (A) flazasulfuron or its salt and (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy]pyridin-2-ylloxy]acetate, bencarbazone, butafenacil, pyranclonil, profluazol, flufenpyr-ethyl and their salts.

In a further embodiment, the present invention provides A herbicidal composition consisting of (A) flazasulfuron or its salt and (B) flumiclorac-pentyl or its salt, and additives to be used for the formulation.

## EXAMPLES

The present invention will be described in further detail with reference to Examples. However, it should be understood that the present invention is by no means restricted thereto.

### TEST EXAMPLE 1

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of persian speedwell (Veronica persica Poir.) were sown. When persian speedwell reached from 7 to 8 leaf-stage, water dispersible granules containing flazasulfuron as an active ingredient (tradename: SHIBAGEN DF, manufactured by Ishihara Sangyo Kaisha, Ltd.) and SC agent containing pyraflufen-ethyl as an active ingredient (tradename: ECOPART FLOWABLE, manufactured by NIHON NOYAKU CO., LTD.) in predetermined amounts were diluted with water (corresponding to 1,000 L/ha)

containing 0.05 vol% of an agricultural adjuvant (tradename: KUSARINO, manufactured by NIHON NOYAKU CO., LTD.) and applied for foliar treatment by a small sprayer.

On the 21st day after the treatment, the state of growth of persian speedwell was visually observed and evaluated in accordance with the following evaluation standard. The growth inhibition rate (%) (measured value) and the growth inhibition rate (%) (calculated value) calculated in accordance with the Colby's formula are shown in Table 1.

Growth inhibition rate (%) = 0 (equivalent to the non-treated area) to 100 (complete kill)

TABLE 1

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of persian speedwell	
		Measured value	Calculated value
Flazasulfuron	25	10	-
Pyraflufen-ethyl	10	43	-
Flazasulfuron + Pyraflufen-ethyl	25+10	63	49

#### TEST EXAMPLE 2

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of black nightshade (Solanum nigrum L.) were sown. When black nightshade reached from 3.2 to 3.5 leaf-stage, SHIBAGEN DF (tradename) and ECOPART FLOWABLE (tradename) in predetermined amounts were diluted with water (corresponding to 1,000 L/ha) containing 0.05 vol% of KUSARINO (tradename) and applied for foliar treatment by a small sprayer.

On the 21st day after the treatment, the state of growth of black nightshade was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 2.

TABLE 2

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of black nightshade	
		Measured value	Calculated value
Flazasulfuron	50	74	-
Pyraflufen-ethyl	2.5	88	-
Flazasulfuron + Pyraflufen-ethyl	50+2.5	99	97

As shown in Table 2, in order to completely suppress growth of black nightshade with flazasulfuron alone, a dose of 50 g/ha or more is required. On the other hand, although not shown in the above Table, the growth inhibition rate of black nightshade was 100% (calculated value: 96%) when flazasulfuron (12.5 g/ha) and pyraflufen-ethyl (5 g/ha) were used in combination, and accordingly it was found that the total dose can be reduced to 17.5 g/ha by using the herbicidal composition of the present invention.

## TEST EXAMPLE 3

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of sticky chickweed (*Cerastium glomeratum* Thuill.) were sown. When sticky chickweed reached from 3.3 to 4.0 leaf-stage, SHIBAGEN DF (tradename) and ECOPART FLOWABLE (tradename) in predetermined amounts were diluted with water (corresponding to 1,000 L/ha) and applied for foliar treatment by a small sprayer.

On the 21st day after the treatment, the state of growth of sticky chickweed was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 3.

TABLE 3

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of sticky chickweed	
		Measured value	Calculated value
Flazasulfuron	6.3	92	-
Pyraflufen-ethyl	10	58	-
Flazasulfuron + Pyraflufen-ethyl	6.3+10	99	97

## TEST EXAMPLE 4

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of common lambsquarters (*Chenopodium album* L.) were sown. When common lambsquarters reached from 6 to 7 leaf-stage, SHIBAGEN DF (tradename) and water dispersible granules containing carfentrazone-ethyl as an active ingredient (tradename: TASK DF, manufactured by Ishihara Sangyo Kaisha, Ltd.) in predetermined amounts were diluted with water (corresponding to 1,000 L/ha) and applied for foliar treatment by a small sprayer.

On the 21st day after the treatment, the state of growth of common lambsquarters was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 4.

TABLE 4

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of common lambsquarters	
		Measured value	Calculated value
Flazasulfuron	1.6	3	-
Carfentrazone-ethyl	27.4	92	-
Flazasulfuron + Carfentrazone-ethyl	1.6+27.4	96	92

## TEST EXAMPLE 5

Upland field soil was put into a 1/300,000 ha pot, and seeds of wild oat (*Avena fatua* L.) were sown. One day later, SHIBAGEN DF (tradename), water dispersible granules containing saflufenacil as an active ingredient (tradename: Treevix, manufactured by BASF) and a wettable powder containing oxadiargyl (manufactured by SIGMA-ALDRICH) as an active ingredient prepared in accordance with a conventional preparation method, in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 13th day after the treatment, the state of growth of wild oat was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 5.

TABLE 5

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of wild oat	
		Measured value	Calculated value
Flazasulfuron	25	40	-
	50	70	-
Saflufenacil	25	15	-
	50	5	-
Oxadiargyl	50	5	-
	800	20	-
Flazasulfuron + Saflufenacil	25+25	70	49
	50+50	80	72
Flazasulfuron + Oxadiargyl	50+50	80	72
	25+800	78	52

## TEST EXAMPLE 6

Upland field soil was put into a 1/300,000 ha pot, and seeds of rostrate sesbania (*Sesbania rostrata* Bremek. & Oberm.) were sown. One day later, SHIBAGEN DF (tradename), water dispersible granules containing flumioxazin as an active ingredient (tradename: Chateau, manufactured by Valent) and a wettable powder containing oxadiargyl (manufactured by SIGMA-ALDRICH) as an active ingredient prepared in accordance with a conventional preparation method, in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of rostrate sesbania was

visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 6.

TABLE 6

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of rostrate sesbania	
		Measured value	Calculated value
Flazasulfuron	12.5	25	-
Flumioxazin	250	70	-
Oxadiargyl	400	20	-
Flazasulfuron + Flumioxazin	12.5+250	95	78
Flazasulfuron + Oxadiargyl	12.5+400	65	40

5

## TEST EXAMPLE 7

Upland field soil was put into a 1/300,000 ha pot, and seeds of sunn-hemp (*Crotalaria juncea* L.) were sown. One day later, SHIBAGEN DF (tradename), ECOPART FLOWABLE (tradename) and a wettable powder containing oxadiargyl (manufactured by SIGMA-ALDRICH) as an active ingredient prepared in accordance with a conventional preparation method, in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of sunn-hemp was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 7.

TABLE 7

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of sunn-hemp	
		Measured value	Calculated value
Flazasulfuron	50	55	-
Pyraflufen-ethyl	10	20	-
	50	45	-
Oxadiargyl	400	40	-
Flazasulfuron + Pyraflufen-ethyl	50+10	70	64
	50+50	85	75
Flazasulfuron + Oxadiargyl	50+400	78	73

## TEST EXAMPLE 8

Upland field soil was put into a 1/300,000 ha pot, and seeds of velvetleaf (*Abutilon theophrasti* Medic.) were sown. One day later, SHIBAGEN DF (tradename), water dispersible granules containing sulfentrazone as an active ingredient (tradename: Authority, manufactured by FMC Corporation) and Treevix (tradename) in

20

predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of velvetleaf was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 8.

TABLE 8

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of velvetleaf	
		Measured value	Calculated value
Flazasulfuron	50	88	-
	100	88	-
Sulfentrazone	25	20	-
Saflufenacil	5	0	-
	10	65	-
Flazasulfuron + Sulfentrazone	50+25	100	90
Flazasulfuron + Saflufenacil	50+5	100	88
	100+10	100	96

TEST EXAMPLE 9

Upland field soil was put into a 1/300,000 ha pot, and seeds of black nightshade (Solanum nigrum L.) were sown. One day later, SHIBAGEN DF (tradename) and Chateau (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of black nightshade was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 9.

TABLE 9

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of black nightshade	
		Measured value	Calculated value
Flazasulfuron	50	85	-
Flumioxazin	2.5	0	-
Flazasulfuron + Flumioxazin	50+2.5	90	85

TEST EXAMPLE 10

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of bermudagrass (Cynodon dactylon (L.) Pers.) were sown. When bermudagrass reached from 2.2 to 2.5 leaf-stage, SHIBAGEN DF (tradename), TASK DF (tradename), Authority (tradename) and Chateau (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) containing 0.2 vol% of KUSARINO (tradename) and applied for foliar treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of bermudagrass was

visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 10.

TABLE 10

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of bermudagrass	
		Measured value	Calculated value
Flazasulfuron	50	75	-
	100	75	-
Carfentrazone-ethyl	20	0	-
	50	0	-
Sulfentrazone	50	30	-
Flumioxazin	5	20	-
Flazasulfuron + Carfentrazone-ethyl	50+50	85	75
	100+20	88	75
Flazasulfuron + Sulfentrazone	100+50	90	83
Flazasulfuron + Flumioxazin	100+5	88	80

5

## TEST EXAMPLE 11

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of bermudagrass (*Cynodon dactylon* (L.) Pers.) were sown. When bermudagrass reached from 3.8 to 4.3 leaf-stage, SHIBAGEN DF (tradename) and ECOPART FLOWABLE (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) containing 0.2 vol% of KUSARINO (tradename) and applied for foliar treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of bermudagrass was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 11.

TABLE 11

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of bermudagrass	
		Measured value	Calculated value
Flazasulfuron	12.5	15	-
Pyraflufen-ethyl	40	0	-
Flazasulfuron + Pyraflufen-ethyl	12.5+40	60	15

## TEST EXAMPLE 12

Upland field soil was put into a 1/1,000,000 ha pot, and seeds of persian speedwell (*Veronica persica* Poir.) were sown. When persian speedwell reached

from 4.0 to 4.3 leaf-stage, SHIBAGEN DF (tradename), ECOPART FLOWABLE (tradename) and TASK DF (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) containing 0.2 vol% of KUSARINO (tradename) and applied for foliar treatment by a small sprayer.

5 On the 28th day after the treatment, the state of growth of persian speedwell was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 12.

TABLE 12

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of persian speedwell	
		Measured value	Calculated value
Flazasulfuron	25	0	-
Pyraflufen-ethyl	80	80	-
Carfentrazone-ethyl	25	65	-
Flazasulfuron + Pyraflufen-ethyl	25+80	100	80
Flazasulfuron + Carfentrazone-ethyl	25+25	85	65

10

## TEST EXAMPLE 13

Upland field soil was put into a 1/300,000 ha pot, and seeds of japanese millet (Echinochloa esculenta (A.Braun) H.Scholz.) were sown. One day later, SHIBAGEN DF (tradename), a SC agent containing pyraclonil as an active ingredient (tradename: PYRACLON FLOWABLE, manufactured by Kyoyu Agri Co., Ltd.) and a SC agent containing pentozaone as an active ingredient (tradename: WECHSER FLOWABLE, manufactured by MITSUI CHEMICALS AGRO, INC.) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

20

On the 14th day after the treatment, the state of growth of japanese millet was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 13.

TABLE 13

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of japanese millet	
		Measured value	Calculated value
Flazasulfuron	50	80	-
Pyraclonil	100	30	-
Pentozaone	100	0	-
Flazasulfuron + Pyraclonil	50+100	95	86
Flazasulfuron + Pentozaone	50+100	90	80

25

## TEST EXAMPLE 14

Upland field soil was put into a 1/300,000 ha pot, and seeds of rostrate sesbania

(*Sesbania rostrata* Bremek. & Oberm.) were sown. One day later, SHIBAGEN DF (tradename), Chateau (tradename), a wettable powder containing flufenpyr-ethyl (manufactured by Wako Pure Chemical Industries, Ltd.) as an active ingredient prepared in accordance with a conventional preparation method, and water dispersible granules containing fluthiacet-methyl as an active ingredient (tradename: Cadet, manufactured by FMC Corporation) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 14th day after the treatment, the state of growth of rostrate sesbania was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 14.

TABLE 14

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of rostrate sesbania	
		Measured value	Calculated value
Flazasulfuron	50	50	-
Flumioxazin	200	75	-
Flufenpyr-ethyl	5	0	-
Fluthiacet-methyl	5	0	-
Flazaſulfuron + Flumioxazin	50+200	99	88
Flazasulfuron + Flufenpyr-ethyl	50+5	70	50
Flazasulfuron + Fluthiacet-methyl	50+5	100	50

## TEST EXAMPLE 15

Upland field soil was put into a 1/300,000 ha pot, and seeds of sunn-hemp (*Crotalaria juncea* L.) were sown. One day later, SHIBAGEN DF (tradename), Authority (tradename), Chateau (tradename), a wettable powder containing flufenpyr-ethyl (manufactured by Wako Pure Chemical Industries, Ltd.) as an active ingredient prepared in accordance with a conventional preparation method, and Cadet (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 28th day after the treatment, the state of growth of sunn-hemp was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 15.

TABLE 15

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of sunn-hemp	
		Measured value	Calculated value
Flazasulfuron	25	0	-
Sulfentrazone	500	30	-
Flumioxazin	100	0	-
Flufenpyr-ethyl	50	0	-
Fluthiacet-methyl	50	0	-
Flazasulfuron + Sulfentrazone	25+500	100	30
Flazasulfuron + Flumioxazin	25+100	70	0
Flazasulfuron + Flufenpyr-ethyl	25+50	90	0
Flazasulfuron + Fluthiacet-methyl	25+50	80	0

## TEST EXAMPLE 16

5 Upland field soil was put into a 1/300,000 ha pot, and seeds of wild oat (*Avena* *fatua* L.) were sown. One day later, SHIBAGEN DF (tradename), Chateau (tradename), Treevix (tradename), PYRACLON FLOWABLE (tradename) and WECHSER FLOWABLE (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

10 On the 28th day after the treatment, the state of growth of wild oat was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 16.

TABLE 16

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of wild oat	
		Measured value	Calculated value
Flazasulfuron	12.5	20	-
	25	75	-
Flumioxazin	250	90	-
Saflufenacil	50	0	-
Pyraclonil	250	40	-
Pentoxazone	250	0	-
Flazasulfuron + Flumioxazin	12.5+250	100	92
Flazasulfuron + Saflufenacil	12.5+50	70	20
Flazasulfuron + Pyraclonil	25+250	90	85
Flazasulfuron + Pentoxazone	25+250	80	75

## TEST EXAMPLE 17

Upland field soil was put into a 1/300,000 ha pot, and seeds of wild oat (*Avena fatua* L.) were sown. One day later, SHIBAGEN DF (tradename) and Chateau (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 14th day after the treatment, the state of growth of wild oat was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 17.

TABLE 17

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of wild oat	
		Measured value	Calculated value
Flazasulfuron	25	30	-
Flumioxazin	500	75	-
Flazasulfuron + Flumioxazin	25+500	90	83

## TEST EXAMPLE 18

Upland field soil was put into a 1/300,000 ha pot, and seeds of corn (*Zea mays* L.) were sown. One day later, SHIBAGEN DF (tradename) and a wettable powder containing butafenacil (synthesized by Ishihara Sangyo Kaisha, Ltd.) as an active ingredient prepared in accordance with a conventional preparation method, in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

On the 14th day after the treatment, the state of growth of corn was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test

Example 1 is shown in Table 18.

TABLE 18

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of corn	
		Measured value	Calculated value
Flazasulfuron	12.5	70	-
Butafenacil	100	20	-
Flazasulfuron + Butafenacil	12.5+100	83	76

## 5 TEST EXAMPLE 19

Upland field soil was put into a 1/300,000 ha pot, and seeds of ivy-leaved morningglory (Ipomoea hederacea Jacq.) were sown. One day later, SHIBAGEN DF (tradename) and a wettable powder containing butafenacil (synthesized by Ishihara Sangyo Kaisha, Ltd.) as an active ingredient prepared in accordance with a conventional preparation method, in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

10 On the 14th day after the treatment, the state of growth of ivy-leaved morningglory was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 19.

15 TABLE 19

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of ivy-leaved morningglory	
		Measured value	Calculated value
Flazasulfuron	50	60	-
Butafenacil	10	0	-
Flazasulfuron + Butafenacil	50+10	85	60

## TEST EXAMPLE 20

Upland field soil was put into a 1/300,000 ha pot, and seeds of shattercane (Sorghum bicolor (L.) Moench) were sown. One day later, SHIBAGEN DF (tradename) and Treevix (tradename) in predetermined amounts were diluted with water (corresponding to 300 L/ha) and applied for soil treatment by a small sprayer.

20 On the 14th day after the treatment, the state of growth of shattercane was visually observed, and the growth inhibition rate (%) obtained in the same manner as in Test Example 1 is shown in Table 20.

TABLE 20

Active ingredient	Dose (g/ha)	Growth inhibition rate (%) of shattercane	
		Measured value	Calculated value
Flazasulfuron	12.5	0	-
Saflufenacil	100	20	-
Flazasulfuron + Saflufenacil	12.5+100	98	20

## INDUSTRIAL APPLICABILITY

According to the present invention, it is possible to provide a herbicidal composition which has a broad herbicidal spectrum and also has a high activity and a long-lasting effect.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgement or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or groups of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

2012243714 25 Nov 2014

## THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for controlling undesired plants or inhibiting their growth, which comprises
  - (I) applying a herbicidally effective amount of a herbicidal composition comprising
    - (A) flazasulfuron or its salt and
    - (B) at least one protoporphyrinogen oxidase inhibitor selected from the group consisting of pyraflufen-ethyl, fluazolate, azafenidin, bencarbazone, carfentrazone-ethyl, sulfentrazone, cinidon-ethyl, flumiclorac-pentyl, flumioxazin, benzfendizone, butafenacil, saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy)pyridin-2-yloxy]acetate (test code: SYN-523), oxadiargyl, oxadiaxon, pentozacone, fluthiacet-methyl, thidiazimin, pyraclonil, profluazol and flufenpyr-ethyl and their salts, or which comprises
      - (II) applying herbicidally effective amounts of components (A) and (B), to the undesired plants or to a place where they grow, wherein the undesired plants is persian speedwell, black nightshade, sticky chickweed, common lambsquarters, wild oat, rostrate sesbania, sunn-hemp, velvetleaf, bermudagrass, japanese millet, corn, ivy-leaved morningglory or shattercane, and wherein said composition presenting a synergistic herbicidal effect, or wherein the combination of components (A) and (B) presenting a synergistic herbicidal effect, wherein (A) is applied in an amount of from 1 to 100 g/ha, and (B) is applied in an amount of from 2 to 800 g/ha.
2. The method according to Claim 1, wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl, carfentrazone-ethyl, sulfentrazone, flumioxazin, saflufenacil, ethyl [3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl)phenoxy)pyridin-2-yloxy]acetate, oxadiargyl, fluthiacet-methyl, flufenpyr-ethyl, flumiclorac-pentyl, azafenidin, butafenacil, pentozacone, pyraclonil and their salts.
3. The method according to Claim 1, wherein (B) is at least one member selected from the group consisting of pyraflufen-ethyl, carfentrazone-ethyl, sulfentrazone, flumioxazin, saflufenacil, oxadiargyl, fluthiacet-methyl, flufenpyr-ethyl, butafenacil, pentozacone, pyraclonil and their salts.