




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00, 31/135, 31/13		A2	(11) International Publication Number: WO 98/22100 (43) International Publication Date: 28 May 1998 (28.05.98)
(21) International Application Number: PCT/EP97/06408	(22) International Filing Date: 17 November 1997 (17.11.97)	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(30) Priority Data: 9624038.7 19 November 1996 (19.11.96) GB			
(71) Applicant (for all designated States except US): NOVARTIS AG [CH/CH]; Schwarzwaldallee 215, CH-4058 Basel (CH).			
(72) Inventors; and (75) Inventors/Applicants (for US only): COTTENS, Sylvain [CH/CH]; In den Reben 12, CH-4108 Witterswil (CH). HOF, Robert, Paul [CH/CH]; Im Huebacher 12, CH-4460 Gelterkinden (CH). WENGER, Roland [CH/CH]; Grenzacherweg 45, CH-4125 Riehen (CH).		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(74) Agent: ROTH, Bernhard, M.; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).			

(54) Title: NEW USE FOR 1,3-PROPANEDIOL DERIVATIVES

(57) Abstract

Use of a compound of formula (I) wherein R₁ is an optionally substituted straight- or branched carbon chain having 12 to 22 carbon atoms which may be optionally interrupted by an optionally substituted phenylene, and each of R₂, R₃, R₄ and R₅, independently, is H or lower alkyl, in free form or in pharmaceutically acceptable salt form, in the prevention or treatment of chronic rejection in a recipient of organ or tissue allo- or xenotransplant, or of acute rejection in a xenograft transplant recipient.



FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

New Use For 1,3-Propanediol Derivatives

The present invention relates to a new use for a compound group comprising 2-amino-1,3-propanediol derivatives.

Compounds for use according to the invention are compounds of formula I



wherein

R_1 is an optionally substituted straight- or branched carbon chain having 12 to 22 carbon atoms which may be optionally interrupted by an optionally substituted phenylene, and
each of R_2 , R_3 , R_4 and R_5 , independently, is H or lower alkyl, in free form or in pharmaceutically acceptable salt form.

When the carbon chain as R_1 is substituted, it is preferably substituted by halogen, nitro, amino, hydroxy or carboxy. When the carbon chain is interrupted by an optionally substituted phenylene, the carbon chain is preferably unsubstituted. When the phenylene moiety is substituted, it is preferably substituted by halogen, nitro, amino, methoxy, hydroxy or carboxy.

Such compounds are disclosed in EP-A1-627,406 the relevant disclosure of which, in particular with respect to the compounds, is incorporated herein by reference.

Preferred compounds of formula I are those wherein R_1 is a straight or branched, preferably straight, chain alkyl having 13 to 20 carbon atoms, optionally substituted by nitro, halogen,

amino, hydroxy or carboxy, and, more preferably those wherein R₁ is phenylalkyl substituted by a straight or branched C₆₋₁₄ alkyl chain optionally substituted by halogen and the alkyl moiety is a C₁₋₆ alkyl optionally substituted by hydroxy. More preferably, R₁ is phenyl-C₁₋₆ alkyl substituted on the phenyl by a straight or branched, preferably straight, C₆₋₁₄ alkyl chain. The C₆₋₁₄ alkyl chain may be in ortho, meta or para, preferably in para.

Preferably each of R₂ to R₅ is H.

Examples of the pharmaceutically acceptable salts of the compounds of the formula (I) include salts with inorganic acids, such as hydrochloride, hydrobromide and sulfate, salts with organic acids, such as acetate, fumarate, maleate, benzoate, citrate, malate, methanesulfonate and benzenesulfonate salts, and when a carboxy group is present, salts with metals such as sodium, potassium, calcium and aluminium, salts with amines, such as triethylamine and salts with dibasic amino acids, such as lysine. The compounds and salts of the present invention encompass hydrate and solvate forms.

When the compounds of formula I have one or more asymmetric centers in the molecule, the present invention is to be understood as embracing the various optical isomers, as well as racemates, diastereoisomers and mixtures thereof are embraced.

Particularly preferred compounds of formula I are 2-amino-2-tetradecyl-1,3-propanediol and especially 2-amino-2-[2-(4-octylphenyl)ethyl]-1,3-propanediol, (hereinafter Compound A) e.g. in hydrochloride form.

Compounds of formula I have, on the basis of observed activity, e.g. as described in EP-A1-627,406 been found to be useful e.g. as immunosuppressant, e.g. in the treatment of acute allograft rejection.

Organ transplants of liver, kidney, lung and heart are now regularly performed as treatment for endstage organ disease. Allograft as well as xenograft transplants have been performed. However, because of problems with long-term chronic rejection, organ transplantation is not yet a permanent solution to irreversible organ disease. Chronic rejection, which manifests as progressive and irreversible graft dysfunction, is the leading cause of organ transplant loss, in some cases already after the first postoperative

year. The clinical problem of chronic rejection is clear from transplantation survival times; about half of kidney allografts are lost within 5 years after transplantation, and a similar value is observed in patients with a heart allograft.

Chronic rejection is considered as a multifactorial process in which not only the immune reaction towards the graft but also the response of the blood vessel wall in the grafted organ to injury ("response-to-injury" reaction) plays a role. The variant of chronic rejection with the worst prognosis is an arteriosclerosis-like alteration, also called transplant vasculopathy graft vessel disease, graft atherosclerosis, transplant coronary disease, etc. This vascular lesion is characterized by migration and proliferation of smooth muscle cells under influence of growth factors, that are amongst others synthesized by endothelium. It appears to progress also through repetitive endothelial injury induced amongst others by host antibody or antigen-antibody complexes, through intimal proliferation and thickening, smooth muscle cell hypertrophy repair, and finally to gradual luminal obliteration. Also so-called non-immunological factors like hypertension, hyperlipidemia, hypercholesterolemia etc. play a role.

Chronic rejection appears to be inexorable and uncontrollable because there is no known effective treatment or prevention modality. Thus, there continues to exist a need for a treatment effective in preventing, controlling or reversing manifestations of chronic graft vessel diseases.

In accordance with the present invention, it has now surprisingly been found that compounds of formula I in free form or in pharmaceutically acceptable salt form inhibit graft vessel disease and are particularly indicated to prevent or treat chronic rejection in a transplanted organ.

Furthermore, it has also been found that compounds of formula I in free form or in pharmaceutically acceptable salt form suppress xenograft rejection.

In accordance with the particular findings of the present invention, there is provided:

- 1.1. A method of preventing or treating manifestations of chronic rejection, e.g. to avoid, reduce or restrict chronic rejection, in a recipient of organ or tissue allo- or xeno-transplant, e.g. heart, lung, combined heart-lung, liver, kidney or pancreatic

transplant, comprising the step of administering to said recipient a therapeutically effective amount of a compound of formula I in free form or in pharmaceutically acceptable salt form;

- 1.2. A method of preventing or treating graft vessel diseases, e.g. transplant vasculopathy, arteriosclerosis or atherosclerosis, in a recipient of organ or tissue allo- or xeno-transplant, e.g. heart, lung, combined heart-lung, liver, kidney or pancreatic transplants, comprising the step of administering to said recipient a therapeutically effective amount of a compound of formula I in free form or in pharmaceutically acceptable salt form;

In a series of further specific or alternative embodiments, the present invention also provides:

2. A method of preventing or controlling acute rejection in a xenograft transplant recipient, e.g. a patient receiving a heart, lung, combined heart-lung, kidney, liver, bone marrow, pancreatic bowel, skin or corneal xenotransplant, comprising administering to said recipient a therapeutically effective amount of a compound of formula I in free form or in pharmaceutically acceptable salt form.

As alternative to the above the present invention also provides:

3. A compound of formula I in free form or in pharmaceutically acceptable salt form for use in any method as defined under 1 or 2 above; or
4. A compound of formula I in free form or in pharmaceutically acceptable salt form for use in the preparation of a pharmaceutical composition for use in any method as defined under 1 or 2 above; or
5. A pharmaceutical composition for use in any method as defined under 1 or 2 above comprising a compound of formula I in free form or in pharmaceutically acceptable salt form together with one or more pharmaceutically acceptable diluents or carriers therefor.

Utility of the compounds of formula I in free form or in pharmaceutically acceptable salt form in chronic rejection, as well as utility in treating diseases and conditions as hereinabove specified, may be demonstrated in animal tests for example in accordance with the methods hereinafter described, as well as in clinic where e.g. the transplanted organ or tissue may be submitted to regular biopsy controls and in the case of heart transplant additionally to ultrasound scanning.

A. Prevention of Graft Vessel Disease

Experimental animals:

Inbred rat strains DA (RT1^a, donors) and Lewis (RT1¹, recipients of allografts), weighing between 200 and 350 grams are used. The animals are allowed unrestricted access to food and water before and after the operation.

Carotid artery transplantation:

The rats are anaesthetised with isofluorane (Abbott), (4-5 % for induction, 1.5-2 % for maintenance) and 300 µg atropin sulphate is injected subcutaneously following the induction. The left carotid artery is dissected free. The artery is clamped proximally and distally and a segment of about 7-10 mm is removed. The gap is bridged by an allograft which had also been subjected to 45 min cold ischemia. Ethilon 10/o sutures are used. Finally the skin is closed with 4/0 sutures. If needed, an Alzet osmotic minipump (Alza Corp. Palo Alto, Calif.) is then implanted subcutaneously over the back (alternatively the animals are treated orally).

The rats are subject to one of the following treatments: A compound of formula I alone at the doses of 0.1 to 10 mg/kg, or in combination with Cyclosporin A at the dose of 0.03, 0.3, or 1 mg·kg⁻¹·day⁻¹ is administered for 8 weeks either by using Alzet osmotic minipumps implanted subcutaneously or alternatively by oral administration.

At 8 weeks the rats are sacrificed, the carotid arteries are perfused for 1 min with 0.1 M phosphate buffered saline solution (PBS, pH 7.4) and then for 15 min with 2.5 % glutaraldehyde in phosphate buffer (pH 7.4). The carotid arteries are then excised and stained in Giemsa solution for histological evaluation.

Morphometric analysis includes the measurement of the thickness of the media and intima. A qualitative analysis of the morphological changes includes a scoring on an

0-3 scale for adventitial infiltration of mononuclear cells and necrosis (vacuolar degeneration, hypertrophy of cells), the number of smooth muscle cells (SMC) nuclei in the media (0-10, <100, >100 and >>100 nuclei for scores, 0, 1, 2 and 3 respectively, SMC necrosis (vacuolar degeneration and hypertrophy of SMC) and the intimal infiltration of mononuclear cells (13).

In both experiments, the compounds of formula I, particularly Compound A in hydrochloride form, significantly inhibit graft infiltration and neointima formation.

B. In vivo heart xenotransplantation (hamster-to-rat)

The hamster-into-rat xenograft combination is a so-called difficult concordant combination. Rats do not have natural anti-hamster antibody in sufficient amounts to yield immediate hyperacute rejection as observed in concordant combinations; however, rejection in untreated recipients occurs within 3-4 days, by antibodies in combination with complement. This is visualized in histology by destruction of blood vessels, exsudation and extravasation of erythrocytes, and influx by polymorphonuclear granulocytes; often there are signs of hemorrhage and thrombosis. Once this rejection has been overcome by effective inhibition of antibody synthesis or complement inactivation, a cellular rejection can emerge later on. This is visualized in histology by influx of mononuclear cells, including lymphocytes, lymphoblastoid cells, and macrophages, and destruction of the myocyte parenchyma. The inhibition of cellular rejection requires more immuno-suppression than that of allografts. Congenitally athymic (rnu/rnu) rats lack a competent (thymus-dependent) cellular immune system and generally are unable to reject allografts. Such animals do reject a hamster xenograft within 3-4 days in a similar fashion as euthymic rats, indicative that (at least part of) anti-hamster antibody synthesis in rats occurs following a thymus-independent B-cell response. Such recipients are useful in hamster xenografting to evaluate rejection by thymus-independent antibody-mediated rejection.

The heart of a Syrian hamster is heterotopically transplanted in the abdomen of a male Lewis (RT1') rat with anastomoses between the donor and recipient's aorta and the donor right pulmonary artery to the recipient's inferior vena cava. The graft is monitored daily by palpation of the abdomen. Rejection is concluded in case of cessation of heart beat. Animals are weighed weekly. In the present series of

experiments, the endpoint is set to 28 days. Animals are subjected to autopsy; apart from the graft, weight and histology is assessed for thymus, spleen, liver, seminal vesicles and testes. Blood is taken and processed to serum for the determination of cytolytic anti-hamster erythrocyte antibody and hemolytic complement activity.

Compounds are dissolved in water and administered daily orally in a volume of 2 ml/kg body weight. Administration of 5-30 mg/kg/day, of a compound of Formula I, e.g. Compound A in hydrochloride form, results in prolonged graft survival, in both athymic and euthymic recipients.

Daily dosages required in practicing the method of the present invention will vary depending upon, for example, the compound of formula I employed, the host, the mode of administration, the severity of the condition to be treated, and the optionally concomitantly used immunosuppressive drug e.g. CysA. A preferred daily dosage range is about from 0.03 to 2.5 mg/kg per day, particularly 0.1 to 2.5 mg/kg per day, e.g. 0.5 to 2.5 mg/kg per day as a single dose or in divided doses. Suitable daily dosages for patients are on the order of from e.g. 1 to 100 mg p.o. Suitable unit dosage forms for oral administration comprise from ca. 1 to 50 mg, usually 5 to 30 mg active ingredient, e.g. Compound A, e.g. in hydrochloride form, together with one or more pharmaceutically acceptable diluents or carriers therefor. As an alternative, the compound of formula I in free form or in pharmaceutically acceptable salt form may also be administered twice or three times a week, e.g. at a dosage as indicated above.

The compounds of formula I may be administered by any conventional route, in particular enterally, e.g. orally, for example in the form of solutions for drinking, tablets or capsules or parenterally, for example in the form of injectable solutions or suspensions. Pharmaceutical compositions comprising the compounds of formula I may be manufactured in conventional manner, e.g. as described in EP-A1-627,406.

The compounds of formula I may be administered as the sole active ingredient or together with other drugs in immunomodulating regimens or other anti-inflammatory agents. For example, the compounds of formula I may be used in combination with cyclosporins, rapamycins or ascomycins, or their immunosuppressive analogs, e.g. cyclosporin A, cyclosporin G, FK-506, rapamycin, 40-O-(2-hydroxy)ethyl-rapamycin, etc.; corticosteroids;

cyclophosphamide; azathioprene; methotrexate; brequinar; leflunomide; mizoribine; mycophenolic acid; mycophenolate mofetil; 15-deoxyspergualine; immuno-suppressive monoclonal antibodies, e.g., monoclonal antibodies to leukocyte receptors, e.g., to MHC, CD2, CD3, CD4, CD7, CD25, CD28, B7, CD40, CD45, or CD58 or to their ligands; or other immunomodulatory compounds, e.g. CTLA4-Ig.

Where the compounds of formula I are administered in conjunction with other immuno-suppressive / immunomodulatory therapy, e.g. for preventing or treating chronic rejection as hereinabove specified, dosages of the co-administered immuno-suppressant or immuno-modulatory compound will of course vary depending on the type of co-drug employed, e.g. whether it is a steroid or a cyclosporin, on the specific drug employed, on the condition being treated, and so forth. In accordance with the foregoing the present invention provides in a yet further aspect:

6. A method as defined above comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a compound of formula I and a second drug substance, said second drug substance being an immuno-suppressant or immunomodulatory drug, e.g. as set forth above.
7. A kit or package for use in any method as defined under 1 or 2 above, comprising a compound of formula I, in free form or in pharmaceutically acceptable salt form, with at least one pharmaceutical composition comprising an immuno-suppressant or immunomodulatory drug. The kit or package may comprise instructions for its administration.

Formulation Example: soft capsules

Compound of formula I,


e.g. Compound A	30 mg
Polyethylene glycol 300	300 mg
Polysorbate 80	20 mg
Total	350 mg

- 9 -

Compounds of formula I in free form or in pharmaceutically acceptable salt form are well tolerated at dosages required for use in accordance with the present invention. For example, the acute LD₅₀ is > 10 mg/kg p.o. in rats and monkeys.

CLAIMS

1. Use of a compound of formula I




wherein

R_1 is an optionally substituted straight- or branched carbon chain having 12 to 22 carbon atoms which may be optionally interrupted by an optionally substituted phenylene, and each of R_2 , R_3 , R_4 and R_5 , independently, is H or lower alkyl, in free form or in pharmaceutically acceptable salt form,

in the preparation of a pharmaceutical composition for preventing or treating manifestations of chronic rejection in a recipient of organ or tissue allo- or xenotransplant.

2. Use of a compound of formula I




wherein

R_1 is an optionally substituted straight- or branched carbon chain having 12 to 22 carbon atoms which may be optionally interrupted by an optionally substituted phenylene, and each of R_2 , R_3 , R_4 and R_5 , independently, is H or lower alkyl, in free form or in pharmaceutically acceptable salt form,

- 11 -

in the preparation of a pharmaceutical composition for preventing or treating graft vessel diseases in a recipient of organ or tissue allo- or xenotransplant.

3. Use of a compound of formula I



wherein

R_1 is an optionally substituted straight- or branched carbon chain having 12 to 22 carbon atoms which may be optionally interrupted by an optionally substituted phenylene, and each of R_2 , R_3 , R_4 and R_5 , independently, is H or lower alkyl, in free form or in pharmaceutically acceptable salt form,

in the preparation of a pharmaceutical composition for preventing or controlling acute rejection in a xenograft transplant recipient.

4. Use of a compound of formula I as defined in claim 1, in free form or in a pharmaceutically acceptable salt form, for preventing or treating manifestations of chronic rejection in a recipient of organ or tissue allo- or xenotransplant.

5. Use of a compound of formula I as defined in claim 1, in free form or in a pharmaceutically acceptable salt form, for preventing or controlling acute rejection in a xenograft transplant recipient.

6. Use according to any one of the preceding claims, wherein the compound of formula I is 2-amino-2-[2-(4-octylphenyl)ethyl]-1,3-propanediol, in free form or in pharmaceutically acceptable salt form.

7. A pharmaceutical composition for preventing or treating manifestations of chronic rejection in a recipient of organ or tissue allo- or xenotransplant, or for preventing or controlling acute rejection in a xenograft transplant recipient, comprising a compound of

formula I as defined in claim 1, in free form or in a pharmaceutically acceptable salt form, together with one or more pharmaceutically acceptable diluents or carriers therefor.

8. A kit or package for preventing or treating manifestations of chronic rejection in a recipient of organ or tissue allo- or xenotransplant, or for preventing or controlling acute rejection in a xenograft transplant recipient, comprising a compound of formula I, in free form or in a pharmaceutically acceptable salt form, with a pharmaceutical composition comprising an immunosuppressant or immunomodulatory drug.
9. A composition according to claim 7 or a kit or package according to claim 8, wherein the compound of formula I is 2-amino-2-[2-(4-octylphenyl)ethyl]-1,3-propanediol, in free form or in pharmaceutically acceptable salt form.
10. A method of preventing or treating manifestations of chronic rejection in a recipient of organ or tissue allo- or xenotransplant, or of preventing or controlling acute rejection in a xenograft transplant recipient, comprising the step of administering to said recipient a therapeutically effective amount of a compound of formula I as defined in claim 1 in free form or in pharmaceutically acceptable salt form.