



(22) Date de dépôt/Filing Date: 2012/05/02
(41) Mise à la disp. pub./Open to Public Insp.: 2012/11/08
(45) Date de délivrance/Issue Date: 2021/05/18
(62) Demande originale/Original Application: 2 834 900
(30) Priorités/Priorities: 2011/05/02 (US61/481,522);
2011/10/06 (US61/544,054)

(51) Cl.Int./Int.Cl. *A61K 39/395* (2006.01),
A61P 1/00 (2006.01), *C07K 16/28* (2006.01),
C07K 16/46 (2006.01), *C12N 15/13* (2006.01)

(72) Inventeurs/Inventors:
DILUZIO, WILLOW, US;
NGUYEN, PHUONG M., US;
VARGA, CSANAD M., US;
PALANIAPPAN, VAITHIANATHAN, US;
BROWN, JASON, US;
FOX, IRVING H., US;
...

(73) Propriétaire/Owner:
MILLENNIUM PHARMACEUTICALS, INC., US

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : PROCEDES DE TRAITEMENT DES AFFECTIONS INTESTINALES INFLAMMATOIRES

(54) Title: METHODS OF TREATING INFLAMMATORY BOWEL DISEASE

(57) Abrégé/Abstract:

Antibody formulations are described comprising a mixture of an anti- $\alpha 4\beta 7$ antibody, an antioxidant or chelator, and at least one free amino acid. The disclosed formulations may have improved stability, reduced aggregate formation, or both. The present invention further provides a safe dosing regimen of these antibody formulations that is easy to follow, and which results in a therapeutically effective amount of the anti- $\alpha 4\beta 7$ antibody in vivo.

(72) Inventeurs(suite)/Inventors(continued): SCHOLZ, CATHERINE, US; ROSARIO, MARIA, US; JENKINS, HELEN, GB

ABSTRACT

Antibody formulations are described comprising a mixture of an anti- $\alpha 4\beta 7$ antibody, an antioxidant or chelator, and at least one free amino acid. The disclosed formulations may have improved stability, reduced aggregate formation, or both. The present invention further provides a safe dosing regimen of these antibody formulations that is easy to follow, and which results in a therapeutically effective amount of the anti- $\alpha 4\beta 7$ antibody in vivo.

METHODS OF TREATING INFLAMMATORY BOWEL DISEASE

This application is a divisional application of Canadian Patent Application number 2,834,900 filed May 2, 2012.

BACKGROUND OF THE INVENTION

- 5 Advances in biotechnology have made it possible to produce a variety of proteins for pharmaceutical applications using recombinant DNA techniques. Because proteins are larger and more complex than traditional organic and inorganic drugs (i.e., possessing multiple functional groups in addition to complex three-dimensional structures), the formulation of such proteins poses special problems.
- 10 For a protein to remain biologically active, a formulation must preserve the conformational integrity of at least a core sequence of the protein's amino acids, while at the same time protecting the protein's multiple functional groups from degradation. Proteins may suffer from a lack of stability, and monoclonal and polyclonal antibodies in particular may be relatively unstable (See e.g., Wang, et al.,
- 15 *J. Pharm Sci.* 96:1-26 (2007)). A large number of formulation options are available, and not one approach or system is suitable for all proteins. Several factors to be considered have been reported (See e.g., Wang et al.).

Numerous characteristics may affect a protein's stability. In fact, even in the case of purified antibodies, the antibody structures may be heterogenous, which further complicates the formulation of such systems. Moreover, the excipients included in antibody formulations preferably minimize any potential immune response.

5 In the case of antibodies, preservation of the conformational integrity is even more important. Degradation pathways for proteins can involve chemical instability (i.e., any process which involves modification of the protein by bond formation or cleavage resulting in a new chemical entity) or physical instability (i.e., changes in the higher order structure of the protein). Chemical instability is manifested in, for
10 example, deamidation, isomerization, hydrolysis, oxidation, fragmentation, glycan beta elimination or disulfide exchange. Physical instability can result from denaturation, aggregation, precipitation or adsorption, for example. The four most common protein degradation pathways are protein fragmentation, aggregation, deamidation, and oxidation. Consequences of chemical or physical instability of
15 therapeutic protein include a lowering of the effective administered dose, decreased safety of the therapy due to, for example irritation or immunological reactivity, and more frequent manufacturing due to short shelf life.

 Several publications have disclosed generally various methods of treating inflammatory bowel diseases, and provided dosing schemes for administration of
20 agents designed to treat inflammatory bowel disease. For example, WO 96/24673 discloses mucosal vascular addressins and treatment of diseases associated with leukocyte recruitment to the gastrointestinal tract as a result of leukocyte binding to cells expressing MAdCAM. U.S. 2005/0095238 describes methods of treating a disease associated with leukocyte infiltration of mucosal tissue and administration to
25 a human an effective amount of a human or humanized immunoglobulin or antigen binding fragment having binding specificity for $\alpha 4\beta 7$ integrin. U.S. 2005/0095238 further describes various doses (e.g. 0.15, about 0.5, about 1.0, about 1.5 or about 2.0 mg immunoglobulin or fragment per kg body weight) and various intervals between doses (7, 14, 21, 28, or 30 days). However, the aforementioned patents
30 and publications do not disclose specific formulations of the anti- $\alpha 4\beta 7$ antibody or

the specific doses and dose regimens described and claimed herein. Importantly, the aforementioned patents do not disclose formulations, doses, and dose regimens that provide for the methods of treatment (supported by clinical trial data) described and claimed herein.

5 The antibody formulations of the present invention may be useful for inhibiting leukocyte binding to cells expressing MAdCAM and therefore aid in treatment of inflammatory bowel diseases in patients. There is, accordingly, an urgent need to discover suitable dosages and dosing schedules of these compounds, and to develop formulations, preferably subcutaneous formulations, which give rise
10 to steady, therapeutically effective blood levels of the antibody formulations over an extended period of time in a stable and convenient form.

SUMMARY OF THE INVENTION

 The invention relates to the identification of an antioxidant or chelator, and at least one amino acid, as useful excipients for formulating anti- $\alpha 4\beta 7$ antibody
15 formulations whose instability makes them susceptible to deamidation, oxidation, isomerization and/or aggregation. The formulation improves stability, reduces aggregate formation and retards degradation of the antibody therein.

 Thus, in a first aspect, the invention relates to a stable liquid pharmaceutical formulation comprising a mixture of an anti- $\alpha 4\beta 7$ antibody, an antioxidant or
20 chelator and at least one free amino acid.

 In some embodiments, the stable liquid pharmaceutical formulation has less than about 1.0% aggregate formation after 12 months at room temperature. The stable liquid pharmaceutical formulation can have less than about 0.2% aggregate formation after 12 months at room temperature.

25 In some embodiments, the antioxidant or chelator is citrate. In some embodiments the chelator is EDTA.

 In some embodiments, the free amino acid of the formulation is histidine, alanine, arginine, glycine, glutamic acid, or any combination thereof. The formulation can comprise between about 50 mM to about 175 mM of free amino
30 acid. The formulation can comprise between about 100 mM and about 175 mM of

free amino acid. The ratio of free amino acid to antibody molar ratio can be at least 250:1.

The formulation can also contain a surfactant. The surfactant can be polysorbate 20, polysorbate 80, a poloxamer, or any combination thereof.

5 In some embodiments, the molar ratio of the antioxidant to the surfactant is about 3:1 to about 156:1.

The formulation can have a pH between about 6.3 and about 7.0. The pH of the formulation can be between about 6.5 and about 6.8. The formulation can have a pH between about 6.1 and about 7.0, or between about 6.2 and 6.8.

10 In some embodiments, the stable liquid pharmaceutical formulation contains at least about 60 mg/ml to about 160 mg/ml anti- $\alpha 4\beta 7$ antibody. The formulation can contain at least about 160 mg/ml anti- $\alpha 4\beta 7$ antibody. The formulation can contain about 150 to about 180 mg/ml antibody or about 165 mg/ml antibody.

In another aspect, the invention relates to a stable liquid pharmaceutical
15 formulation comprising at least about 60 mg/ml to about 160 mg/ml anti- $\alpha 4\beta 7$ antibody, a buffering agent and at least about 10 mM citrate. The buffering agent can be a histidine buffer.

In another aspect, the invention relates to a stable liquid pharmaceutical formulation comprising at least about 60 mg/ml to about 180 mg/ml anti- $\alpha 4\beta 7$
20 antibody, a buffering agent and at least about 5 mM citrate. The buffering agent can be a histidine buffer.

In another aspect, the invention relates to a stable liquid pharmaceutical formulation comprising at least about 160 mg/ml anti- $\alpha 4\beta 7$ antibody and at least about 10 mM citrate. The formulation can further contain polysorbate 80.

25 In another aspect, the invention relates to a stable liquid pharmaceutical formulation comprising about 160 mg/ml anti- $\alpha 4\beta 7$ antibody and at least about 5 mM citrate. The formulation can further contain polysorbate 80.

In another aspect, the invention relates to a stable liquid pharmaceutical formulation comprising a mixture of anti- $\alpha 4\beta 7$ antibody, citrate, histidine, arginine
30 and polysorbate 80. The formulation can be present in a container, such as a vial, cartridge, syringe or autoinjector.

The anti- $\alpha 4\beta 7$ antibody in the stable liquid pharmaceutical formulation of the invention can be vedolizumab. The formulation of the invention can be for subcutaneous, intravenous, or intramuscular administration.

In some aspects, the formulation can minimize immunogenicity of the anti-
5 $\alpha 4\beta 7$ antibody.

In another aspect, the invention relates to a method of treating inflammatory bowel disease, comprising administering to a patient in need thereof the stable liquid pharmaceutical formulation described herein. The administering can be subcutaneous administering. The administering can be self-administering.

10 In yet another aspect, the invention relates to an article of manufacture, comprising a container, a stable liquid pharmaceutical formulation described herein, and instructions for its use.

In one aspect, the invention relates to a method for treating a human patient suffering from inflammatory bowel disease, wherein the method comprises the step
15 of administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to the following dosing regimen: (a) initial doses, e.g., in an induction phase treatment
20 regimen, of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection every other day for six doses; (b) followed at week six by a seventh and subsequent doses, e.g., in a maintenance phase treatment regimen, of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection every two weeks or every four weeks as needed;
25 wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and
30 three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below: light chain: CDR1

SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1
SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In one aspect, the invention relates to a method for treating a human patient suffering from inflammatory bowel disease, wherein the method comprises the step
5 of administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized
10 immunoglobulin or antigen-binding fragment thereof is administered to the patient according to the following dosing regimen comprising an induction phase of intravenous doses and a maintenance phase of subcutaneous doses: (a) an initial intravenous dose of 300 mg of the humanized immunoglobulin or antigen-binding fragment thereof as an intravenous infusion; (b) followed by a second intravenous
15 subsequent dose of 300 mg of the humanized immunoglobulin or antigen-binding fragment thereof as an intravenous infusion at about two weeks after the initial dose; (c) followed beginning at week six by a third and subsequent doses of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection every week, every two weeks, every three weeks or every four weeks as
20 needed; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain
25 variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below:
light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11;
heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In another aspect, the invention relates to a dosing regimen for the
30 therapeutic treatment of inflammatory bowel disease, wherein the dosing regimen comprises the step of: administering to a patient suffering from inflammatory bowel

disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein

5 the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to a subcutaneous or intramuscular dosing regimen which maintains a mean steady state trough serum concentration of the immunoglobulin or antigen-binding fragment thereof of about 9 to about 13 $\mu\text{g/mL}$; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory

10 bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain

15 variable region of the amino acid sequence set forth below: light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In another aspect, the invention relates to a dosing regimen for the therapeutic treatment of inflammatory bowel disease, wherein the dosing regimen

20 comprises the step of: administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein

25 the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to a subcutaneous or intramuscular dosing regimen which maintains a mean steady state trough serum concentrations of the humanized immunoglobulin or antigen-binding fragment thereof of about 35 to about 40 $\mu\text{g/mL}$; wherein the dosing regimen induces a clinical response and clinical

30 remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for

the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below:

- 5 light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11;
heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

- In another aspect, the invention relates to a method of treating a human patient suffering from inflammatory bowel disease, wherein the method comprises the step of: administering to a patient suffering from inflammatory bowel disease, a
- 10 humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient
- 15 according to the following dosing regimen: (a) a plurality of induction phase doses of the humanized immunoglobulin or antigen-binding fragment thereof sufficient to achieve a mean trough serum concentration of about 20 to about 30 $\mu\text{g/mL}$ of the humanized immunoglobulin or antigen-binding fragment thereof by about six weeks of initial dosing; (b) followed by a plurality of maintenance phase doses of the
- 20 humanized immunoglobulin or antigen-binding fragment thereof as needed to maintain a mean steady state trough serum concentration of about 9 to about 13 $\mu\text{g/mL}$ or about 35 to 40 $\mu\text{g/mL}$ of the immunoglobulin or antigen-binding fragment thereof; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the
- 25 humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below:
- 30 light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11;
heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In some aspects, the formulation, method of treatment, dose and/or dose regimen ensure minimal likelihood that a patient will develop antibodies reactive to the anti- $\alpha 4\beta 7$ antibody.

5 The patient may have had a lack of an adequate response with, loss of response to, or was intolerant to treatment with at least one of an immunomodulator, a tumor necrosis factor-alpha (TNF- α) antagonist or combinations thereof.

The inflammatory bowel disease can be Crohn's disease or ulcerative colitis. The inflammatory bowel disease can be moderate to severely active ulcerative colitis.

10 The dosing regimen can result in mucosal healing in patients suffering from moderate to severely active ulcerative colitis.

The patient may have previously received treatment with at least one corticosteroid for the inflammatory bowel disease. The patient may concurrently receive treatment with at least one corticosteroid for the inflammatory bowel
15 disease. The dosing regimen can result in a reduction, elimination or reduction and elimination of corticosteroid use by the patient.

In some aspects, the humanized immunoglobulin or antigen-binding fragment thereof is administered in a final dosage form at a concentration of between about 1.0 mg/ml to about 1.4 mg/ml. The humanized immunoglobulin or
20 antigen-binding fragment thereof can be administered in a final dosage form of about 1.2 mg/ml.

In some aspects, the humanized immunoglobulin or antigen-binding fragment is administered in a final dosage form having an amount of anti- $\alpha 4\beta 7$ antibody between about 70 to about 250 mg, between about 90 to about 200 mg,
25 between about 150 to about 180 mg, or at least 160 mg.

In some aspects, the dosing regimen does not alter the ratio of CD4 to CD8 in cerebrospinal fluid of patients receiving said treatment.

The patient can be a person 65 years of age or older and does not require any adjustment of the dosing regimen.

In some aspects the method of treatment with the anti- $\alpha 4\beta 7$ antibody formulation, the dose, or the dose regimen can minimize immunogenicity of the anti- $\alpha 4\beta 7$ antibody.

In one aspect, there is provided a stable liquid pharmaceutical formulation
5 comprising at least 60 mg/mL of an anti- $\alpha 4\beta 7$ antibody, an antioxidant or chelator,
and at least one free amino acid, wherein the formulation is in liquid form, wherein
the anti- $\alpha 4\beta 7$ antibody comprises a light chain variable region comprising a CDR1
comprising SEQ ID NO: 11, a CDR2 comprising SEQ ID NO: 12, and a CDR3
comprising SEQ ID NO: 13, and comprises a heavy chain variable region
10 comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9,
and a CDR3 comprising SEQ ID NO: 10.

In one aspect, there is provided a stable liquid pharmaceutical formulation
comprising at least 60 mg/mL of an anti- $\alpha 4\beta 7$ antibody, citrate, and at least one free
amino acid, wherein the anti- $\alpha 4\beta 7$ antibody comprises a light chain variable region
15 comprising a complementarity determining region1 (CDR1) comprising SEQ ID
NO:11, a CDR2 comprising SEQ ID NO:12, and a CDR3 comprising SEQ ID
NO:13, and comprises a heavy chain variable region comprising a CDR1 comprising
SEQ ID NO:8, a CDR2 comprising SEQ ID NO:9, and a CDR3 comprising SEQ ID
NO:10.

20 In one aspect, there is provided a stable liquid pharmaceutical formulation
comprising at least 140 mg/ml of an anti- $\alpha 4\beta 7$ antibody, a buffering agent, a free
amino acid, and at least 5 mM citrate, wherein the anti- $\alpha 4\beta 7$ antibody comprises a
light chain variable region comprising a CDR1 comprising SEQ ID NO: 11, a CDR2
comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and
25 comprises a heavy chain variable region comprising a CDR1 comprising SEQ ID
NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10.

In one aspect, there is provided a stable liquid pharmaceutical formulation
comprising at least 60 mg/mL of an anti- $\alpha 4\beta 7$ antibody, citrate, histidine, arginine,
and polysorbate 80, wherein the anti- $\alpha 4\beta 7$ antibody comprises a light chain variable
30 region comprising a CDR1 comprising SEQ ID NO: 11, a CDR2 comprising SEQ
ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprises a heavy chain

variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10.

In one aspect, there is provided a prefilled syringe comprising a stable liquid formulation comprising 60 to 190 mg/ml of an anti- $\alpha 4\beta 7$ antibody comprising a
5 light chain variable region comprising a CDR1 comprising SEQ ID NO: 11, a CDR2 comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprising a heavy chain variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10, wherein the formulation has a pH of 5.5 to 7.5, and a surfactant concentration of
10 0.01% to 0.15%, and wherein the prefilled syringe comprises 0.2 to 0.8 mg silicone oil.

In one aspect, there is provided a stable liquid pharmaceutical formulation comprising 150mg/ml to 180 mg/ml of an anti- $\alpha 4\beta 7$ antibody, citrate, at least one free amino acid, a buffering agent, and a surfactant, wherein the anti- $\alpha 4\beta 7$ antibody
15 is an IgG1 isotype comprising a light chain variable region comprising a CDR1 comprising SEQ ID NO: 11, a CDR2 comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprising a heavy chain variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10, and wherein the formulation has a pH of 6
20 to 7, and wherein the formulation has greater than or equal to 96% monomeric anti- $\alpha 4\beta 7$ antibody as determined by size exclusion chromatography (SEC) after storage at 40°C for 4 weeks.

In one aspect, there is provided a stable liquid pharmaceutical formulation comprising an anti- $\alpha 4\beta 7$ antibody at a concentration of 150 mg/ml to 180 mg/ml, citrate, histidine, arginine, and polysorbate 80, wherein the anti- $\alpha 4\beta 7$ antibody
25 comprises a light chain variable region comprising a CDR1 comprising SEQ ID NO: 11, a CDR2 comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprises a heavy chain variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO:
30 10, and wherein the formulation has greater than or equal to 96% monomeric anti-

α 4 β 7 antibody as determined by size exclusion chromatography (SEC) after storage at 40°C for 4 weeks.

In one aspect, there is provided a stable liquid pharmaceutical formulation comprising 140 mg/ml to 170 mg/ml of an anti- α 4 β 7 antibody, 20 mM to 30 mM
5 citrate, 10 mM to 75 mM histidine, 0.01% to 0.5% (w/v) polysorbate 80, and 50 mM to 150 mM arginine, wherein the anti- α 4 β 7 antibody is an IgG1 isotype comprising a light chain variable region comprising a CDR1 comprising SEQ ID NO:11, a CDR2 comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprising a heavy chain variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2
10 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10, and wherein the formulation has greater than or equal to 96% monomeric anti- α 4 β 7 antibody as determined by size exclusion chromatography (SEC) after storage at 40°C for 4 weeks.

In one aspect, there is provided a prefilled syringe or an autoinjector comprising a 108 mg dose of an anti- α 4 β 7 antibody, wherein the antibody comprises
15 a light chain variable region comprising a CDR1 comprising SEQ ID NO:11, a CDR2 comprising SEQ ID NO: 12, and a CDR3 comprising SEQ ID NO: 13, and comprises a heavy chain variable region comprising a CDR1 comprising SEQ ID NO:8, a CDR2 comprising SEQ ID NO: 9, and a CDR3 comprising SEQ ID NO: 10.

BRIEF DESCRIPTION OF THE DRAWINGS

20 FIG. 1 is an illustration of a nucleotide sequence (SEQ ID NO:1) encoding the heavy chain of a humanized anti- α 4 β 7 immunoglobulin, and the deduced amino acid sequence of the heavy chain (SEQ ID NO:2). The nucleotide sequence contains cloning sites (lower case), Kozak sequence (upper case, nucleotides 18-23 of SEQ ID NO:1) and leader sequence (lower case, nucleotides 24-86 of SEQ ID NO:1) at
25 the 5' end of the heavy chain. The open reading frame of the nucleotide sequence is nucleotides 24-1433 of SEQ ID NO:1.

FIG. 2 is an illustration of a nucleotide sequence (SEQ ID NO:3) encoding the light chain of a humanized immunoglobulin referred to herein as vedolizumab, and the deduced amino acid sequence (SEQ ID NO: 4) of the light chain. The
30 nucleotide sequence contains cloning sites (lower case), Kozak sequence (upper case, nucleotides 18-23 of SEQ ID NO:3) and leader sequence (lower case,

nucleotides 24-80 of SEQ ID NO:3) at the 5' end of the heavy chain. The open reading frame of the nucleotide sequence is nucleotides 24-737 of SEQ ID NO:3.

FIG. 3 is an alignment of the amino acid sequences of (A) the mature humanized light chain (amino acids 20-238 of SEQ ID NO:4) of the humanized immunoglobulin referred to herein as vedolizumab and (B) the mature humanized light chain of the humanized immunoglobulin referred to herein as LDP-02 (SEQ ID NO:5). (Regarding LDP-02, see, WO 98/06248 and Feagan et al., *N. Eng. J. Med.* 352:2499-2507 (2005)). Feagan et al. describe a clinical study of LDP-02, but in the article they refer to LDP-02 as MLN02.) The alignment illustrates that the amino acid sequences of the light chains of vedolizumab and LDP-02 differ at positions 114 and 115 of the mature light chains.

FIG. 4 is an alignment of amino acid sequences of (A) a generic human kappa light chain constant region (SEQ ID NO:6) and (B) a generic murine kappa light chain constant region (SEQ ID NO:7). The amino acid residues Thr and Val (which are present at positions 114 and 115 of the mature vedolizumab light chain (amino acids 133 and 134 of SEQ ID NO:4)) are present in the constant region of the human kappa light chain, whereas the amino acid residues Ala and Asp (which are present at positions 114 and 115 of the mature LDP-02 light chain (SEQ ID NO:5)) are present in the constant region of the mouse kappa light chain.

FIG. 5 is a map of vector pLKTOK38D (also referred to as pTOK38MLN02-TV), which encodes the humanized heavy chain and the humanized light chain of MLN02, and is suitable for producing vedolizumab in CHO cells. (See, U.S. Patent Application Publication No. 2004/0033561 A1 which discloses pLKTOK38. pLKTOK38D is a variant of pLKTOK38 in which the restriction sites indicated on the map flank the sequence encoding the light chain variable region.)

FIG. 6 shows the SEC aggregates slope of formation (% per day) as a result of changes to protein concentration, pH and surfactant:protein molar ratio. At a pH range of 6.0 to 6.5, the formation of aggregates was similar for formulation with the polysorbate 80:protein molar ratio range of 0.7 to 1.5.

FIG. 7 is a graph showing that at polysorbate 80:protein molar ratios greater than 1.5, the aggregate formation rate increases as pH increases.

FIG. 8 is a graph showing the effect of excipients on the formation of aggregates. 25mM citrate, 5 mM citrate, 5 mM EDTA, 25 mM cysteine, or 5 mM cysteine was added to formulations. All three excipients reduced the formation of aggregates.

5 FIG. 9 is a set of graphs that shows reduction in aggregate formation with the presence of 25 mM citrate in the formulation, and a correlation between increased protein concentration and increased rate of aggregate formation.

FIG. 10 is a graph showing the results of the CEX species degradation at 40°C. The data shows the influence of pH change on CEX degradation.

10 FIG. 11 is a graph showing the effect of temperature on the pH of formulations. The pH of formulations containing histidine decrease with temperature, whereas the pH of citrate formulations is not affected by temperature.

FIG. 12 is a graph showing the percentage of CEX major isoform over a period of twelve months. Formulations having a pH of 6.0-6.2 showed about 1-2%
15 fewer major isoform than formulations having a pH of 6.3-6.4.

FIG. 13 shows a set of graphs that demonstrate that viscosity is affected mainly by protein concentration and pH. Sucrose, histidine and arginine additions are shown to have a minor affect on the viscosity of the formulation.

FIG. 14 shows the amino acid sequences of (A) the mature human
20 GM607'CL antibody kappa light chain variable region and (B) the the human 21/28'CL heavy chain variable region.

FIG. 15 shows components of a protein product in a pre-filled syringe.

FIGS. 16A-B show the effect of (A) protein concentration and (B) viscosity on the injection force of various syringes tested.

25 FIG. 17 (A) shows the initial glide force as a function of protein concentration and the needle size. FIG. 17 (B) shows the initial glide force for each syringe manufacturer and needle size.

FIG. 18 shows the absorption profile of vedolizumab. The graph shows that concentrations of the intramuscular and subcutaneous doses generally overlap.

30 There are no apparent gross differences in the absorption profiles of these routes of administration.

DETAILED DESCRIPTION OF THE INVENTION

The invention relates to a pharmaceutical formulation comprising anti- $\alpha 4\beta 7$ antibodies. The pharmaceutical formulation may be a mixture comprising an antioxidant or chelator (e.g., citrate), anti- $\alpha 4\beta 7$ antibody and a free amino acid. The pharmaceutical formulation may be in a solid or liquid form.

Definitions

The term "pharmaceutical formulation" refers to a preparation that contains an anti- $\alpha 4\beta 7$ antibody in such form as to permit the biological activity of the antibody to be effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

A "stable" formulation is one in which the antibody therein substantially retains its physical stability and/or its chemical stability and/or its biological activity upon storage. In one aspect, the formulation substantially retains its physical and chemical stability, as well as its biological activity upon storage. The storage period is generally selected based on the intended shelf-life of the formulation. Various analytical techniques for measuring protein stability are available in the art and are reviewed, for example, in *Peptide and Protein Drug Delivery*, 247-301, Vincent Lee Ed., Marcel Dekker, Inc., New York, N.Y., Pubs. (1991) and Jones, A. *Adv. Drug Delivery Rev.* 10: 29-90 (1993). Stability can be measured at a selected temperature for a selected time period. For example, the liquid formulation is stable at about 40 °C for at least about 3 days, 5 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks or 6 weeks. In another aspect, the lyophilized formulation is stable at about 40 °C for at least about 2-4 weeks, at least about 3 months, at least about 6 months, at least about 9 months, at least about 12 months, or at least about 18 months. The liquid and/or lyophilized formulation in another aspect is stable at about 5 °C and/or 25 °C for at least about 1 month, at least about 3 months, at least about 6 months, at least about 9 months, at least about 12 months, at least about 18 months, at least about 24 months, at least about 30 months, or at least about 36 months; and/or stable at about -20 °C and/or -70 °C for at least about 1 month, at least about 3 months, at least about 6 months, at least about 9 months, at least about 12 months, at least about 18 months,

at least about 24 months, at least about 30 months, at least about 36 months, at least about 42 months, or at least about 48 months. Furthermore, the liquid formulation may, in some embodiments, be stable following freezing (to, e.g., -80 °C) and thawing, for example following 1, 2 or 3 cycles of freezing and thawing.

- 5 The stability of a liquid formulation can be evaluated qualitatively and/or quantitatively in a variety of different ways, including evaluation of dimer, multimer and/or aggregate formation (for example using size exclusion chromatography (SEC), matrix-assisted laser desorption-ionization time-of-flight mass spectrometry (MALDI-TOF MS), analytical ultracentrifugation, light scattering (photon
10 correlation spectroscopy, dynamic light scattering (DLS), static light scattering, multi-angle laser light scattering (MALLS)), flow-based microscopic imaging, electronic impedance (coulter) counting, light obscuration or other liquid particle counting system, by measuring turbidity, and/or by visual inspection); by assessing charge heterogeneity using cation exchange chromatography (CEX), isoelectric
15 focusing (IEF), e.g. capillary technique (cIEF), or capillary zone electrophoresis; amino-terminal or carboxy-terminal sequence analysis; mass spectrometric analysis; SDS-PAGE or SEC analysis to compare fragmented, intact and multimeric (i.e., dimeric, trimeric, etc.) antibody; peptide map (for example tryptic or LYS-C)
analysis; evaluating biological activity or antigen binding function of the antibody;
20 and the like. Stability of a solid-state formulation can also be evaluated qualitatively and/or quantitatively in a variety of different ways, including direct tests, such as identifying crystal structure by X-Ray Powder Diffraction (XRPD); evaluating antibody structure in the solid state using Fourier Transform Infrared Spectroscopy (FTIR); and measuring thermal transitions in the lyophilized solid (melting, glass
25 transition, etc.) using Differential Scanning Calorimetry (DSC) and indirect tests such as measuring moisture content by Karl Fisher test, e.g., to extrapolate the likelihood of chemical instability through hydrolysis. Instability may involve any one or more of: aggregation (e.g., non-covalent soluble aggregation, covalent
soluble aggregation (e.g., disulfide bond rearrangement/scrambling), insoluble
30 aggregation), deamidation (e.g. Asn deamidation), oxidation (e.g. Met oxidation), isomerization (e.g. Asp isomeriation), clipping/hydrolysis/fragmentation (e.g. hinge

region fragmentation), succinimide formation, N-terminal extension, C-terminal processing, glycosylation differences, and the like.

A “deamidated” monoclonal antibody is one in which one or more asparagine or glutamine residue thereof has been derivatized, e.g. to an aspartic acid
5 or an iso-aspartic acid.

An antibody which is “susceptible to deamidation” is one comprising one or more residue which has been found to be prone to deamidate.

An antibody which is “susceptible to oxidation” is an antibody comprising one or more residue which has been found to be prone to oxidation.

10 An antibody which is “susceptible to aggregation” is one which has been found to aggregate with other antibody molecule(s), especially upon freezing, heating, drying, reconstituting and/or agitation.

An antibody which is “susceptible to fragmentation” is one which has been found to be cleaved into two or more fragments, for example at a hinge region
15 thereof.

By “reducing deamidation, oxidation, aggregation, or fragmentation” is intended to mean preventing or decreasing (e.g., to 80%, 60%, 50%, 40%, 30%, 20% or 10% of) the amount of deamidation, aggregation, or fragmentation relative to the monoclonal antibody formulated at a different pH or in a different buffer.

20 An “aggregate”, “SEC aggregate”, or “soluble aggregate” is more than one and less than or equal to ten antibody proteins and/or fragments associated together through covalent, ionic, or hydrophobic interactions to form a larger protein body.

An “insoluble aggregate” or “particle” is greater than ten antibody proteins and/or fragments associated together through covalent, ionic, or hydrophobic
25 interactions to form a larger protein body.

As used herein, “biological activity” of a monoclonal antibody refers to the ability of the antibody to bind to antigen and result in a measurable biological response which can be measured *in vitro* or *in vivo*. Such activity may be antagonistic or agonistic.

30 The cell surface molecule, “ $\alpha 4\beta 7$ integrin,” or “ $\alpha 4\beta 7$,” is a heterodimer of an $\alpha 4$ chain (CD49D, ITGA4) and a $\beta 7$ chain (ITGB7). Each chain can form a

heterodimer with an alternative integrin chain, to form for example $\alpha_4\beta_1$ or $\alpha_E\beta_7$. Human α_4 and β_7 genes (GenBank (National Center for Biotechnology Information, Bethesda, MD) RefSeq Accession numbers NM_000885 and NM_000889, respectively) are expressed by B and T lymphocytes, particularly memory CD4+ lymphocytes. Typical of many integrins, $\alpha_4\beta_7$ can exist in either a resting or activated state. Ligands for $\alpha_4\beta_7$ include vascular cell adhesion molecule (VCAM), fibronectin and mucosal addressin (MAdCAM, e.g., MAdCAM-1).

As used herein, a human immunoglobulin or antigen-binding fragment thereof that has "binding specificity for the $\alpha_4\beta_7$ complex" binds to $\alpha_4\beta_7$, but not to $\alpha_4\beta_1$ or $\alpha_E\beta_7$.

As used herein, an "isotonic" formulation has substantially the same osmotic pressure as human blood. Isotonic formulations will generally have an osmotic pressure from about 250 to 350 mOsm. Isotonicity can be measured using a vapor pressure or ice-freezing type osmometer, for example.

As used herein, "buffering agent" refers to a buffer that resists changes in pH by the action of its acid-base conjugate components. The buffering agent may be present in a liquid or solid formulation of the invention. In some embodiments, the buffering agent of this invention adjusts the pH of the formulation to about 5.0 to about 7.5, to about pH 5.5 to about 7.5, to about pH 6.0 to about 7.0, or to a pH of about 6.3 to about 6.5. In one aspect, examples of buffering agents that alone or in combination, will control the pH in the 5.0 to 7.5 range include acetate, succinate, gluconate, histidine, citrate, phosphate, maleate, cacodylate, 2-[N-morpholino]ethanesulfonic acid (MES), bis(2-hydroxyethyl)iminotris[hydroxymethyl]methane (Bis-Tris), N-[2-acetamido]-2-iminodiacetic acid (ADA), glycylglycine and other organic acid buffers. In another aspect, the buffering agent herein is histidine or citrate.

A "histidine buffer" is a buffer comprising histidine ions. Examples of histidine buffers include histidine chloride, histidine acetate, histidine phosphate, histidine sulfate solutions. The histidine buffer or histidine-HCl buffer has a pH between about pH 5.5 to about 7.0, between about pH 6.1 to about 6.9, or about pH 6.5.

A "citrate buffer" is a buffer comprising citrate ions. Examples of citrate buffers include sodium citrate, ammonium citrate, calcium citrate, and potassium citrate solutions. The citrate buffer has a pH of about 3.0 to 6.2, about pH 5.5 to 6.5, about pH 6.1 to about 6.5, about pH 6.1, about pH 6.2, or about pH 6.5.

5 A "saccharide" herein is a compound that has a general formula $(CH_2O)_n$ and derivatives thereof, including monosaccharides, disaccharides, trisaccharides, polysaccharides, sugar alcohols, reducing sugars, nonreducing sugars, and the like. Examples of saccharides herein include glucose, sucrose, trehalose, lactose, fructose, maltose, dextran, erythritol, glycerol, arabitol, sylitol, sorbitol, mannitol, mellibiose,
10 melezitose, raffinose, mannotriose, stachyose, maltose, lactulose, maltulose, glucitol, maltitol, lactitol, iso-maltulose, and the like. A saccharide can be a lyoprotectant. In one aspect, a saccharide herein is a nonreducing disaccharide, such as sucrose.

 Herein, a "surfactant" refers to an agent that lowers surface tension of a liquid. In one aspect, the surfactant is a nonionic surfactant. Examples of
15 surfactants herein include polysorbate (polyoxyethylene sorbitan monolaurate, for example, polysorbate 20 and polysorbate 80); TRITON (t-Octylphenoxy polyethoxy ethanol, nonionic detergent, Union Carbide subsidiary of Dow Chemical Co., Midland MI); sodium dodecyl sulfate (SDS); sodium laurel sulfate; sodium octyl glycoside; lauryl-, myristyl-, linoleyl-, or stearyl-sulfobetaine;
20 lauryl-, myristyl-, linoleyl- or stearyl-sarcosine; linoleyl-, myristyl-, or cetyl-betaine; lauroamidopropyl-, cocamidopropyl-, linoleamidopropyl-, myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-betaine (e.g. lauroamidopropyl); myristamidopropyl-, palmidopropyl-, or isostearamidopropyl-dimethylamine; sodium methyl cocoyl-, or disodium methyl oleyl-taurate; sorbitan monopalmitate;
25 and the MONAQUAT series (Mona Industries, Inc., Paterson, N.J.); polyethyl glycol (PEG), polypropylene glycol (PPG), and copolymers of poloxyethylene and poloxypropylene glycol (e.g. Pluronics/Poloxamer, PF68 etc); etc. In another aspect, the surfactant herein is polysorbate 80.

 The term "chelator" refers to an agent that binds to an atom through more
30 than one bond. In one aspect, examples of chelators herein include citrate, ethylenediaminetetraacetic acid, ethyleneglycoltetraacetic acid (EGTA),

dimercaprol, diethylenetriaminepentaacetic acid, and N,N-bis(carboxymethyl)glycine. In another aspect, the chelator is citrate or EDTA.

The term “antioxidant” refers to an agent that inhibits the oxidation of other molecules. Examples of antioxidants herein include citrate, lipoic acid, uric acid, glutathione, tocopherol, carotene, lycopene, cysteine, phosphonate compounds, e.g., etidronic acid, desferoxamine and malate.

The term “antibody” herein is used in the broadest sense and specifically covers full length monoclonal antibodies, immunoglobulins, polyclonal antibodies, multispecific antibodies (e.g. bispecific antibodies) formed from at least two full length antibodies, e.g., each to a different antigen or epitope, and individual antigen binding fragments, including dAbs, scFv, Fab, F(ab)₂, Fab', including human, humanized and antibodies from non-human species and recombinant antigen binding forms such as monobodies and diabodies.

Molar amounts and ratios of anti- $\alpha 4\beta 7$ antibody to other excipients described herein are calculated on the assumption of an approximate molecular weight of about 150,000 daltons for the antibody. The actual antibody molecular weight may differ from 150,000 daltons, depending on amino acid composition or post-translational modification, e.g., as dependent on the cell line used to express the antibody. Actual antibody molecular weight can be +/- 5% of 150,000 daltons.

The term “human antibody” includes an antibody that possesses a sequence that is derived from a human germ-line immunoglobulin sequence, such as an antibody derived from transgenic mice having human immunoglobulin genes (e.g., XENOMOUSE genetically engineered mice (Abgenix, Fremont, CA), HUMAB-MOUSE®, KIRIN TC MOUSE™ transchromosome mice, KMMOUSE® (MEDAREX, Princeton, NJ)), human phage display libraries, human myeloma cells, or human B cells.

The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variants that may arise during production of the monoclonal antibody, such variants generally being present in minor amounts. In

contrast to polyclonal antibody preparations that typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.*, *Nature*, 256:495 (1975), or may be made by recombinant DNA methods (see, e.g., U.S. Pat. No. 4,816,567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.*, *Nature*, 352:624-628 (1991) and Marks *et al.*, *J. Mol. Biol.*, 222:581-597 (1991), for example.

The monoclonal antibodies herein specifically include “chimeric” antibodies in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Pat. No. 4,816,567; and Morrison *et al.*, *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). Chimeric antibodies of interest herein include “primatized” antibodies comprising variable domain antigen binding sequences derived from a non-human primate (e.g. Old World Monkey, Ape etc) and human constant region sequences.

“Antigen binding fragments” of the humanized immunoglobulin prepared in the formulation of the invention comprise at least the variable regions of the heavy and/or light chains of an anti- $\alpha 4\beta 7$ antibody. For example, an antigen binding fragment of vedolizumab comprises amino acid residues 20-131 of the humanized light chain sequence of SEQ ID NO:4. Examples of such antigen binding fragments include Fab fragments, Fab' fragments, scFv and F(ab')₂ fragments of a humanized immunoglobulin known in the art. Antigen binding fragments of the humanized

immunoglobulin of the invention can be produced by enzymatic cleavage or by recombinant techniques. For instance, papain or pepsin cleavage can be used to generate Fab or F(ab')₂ fragments, respectively. Antibodies can also be produced in a variety of truncated forms using antibody genes in which one or more stop codons
5 have been introduced upstream of the natural stop site. For example, a recombinant construct encoding the heavy chain of an F(ab')₂ fragment can be designed to include DNA sequences encoding the CH₁ domain and hinge region of the heavy chain. In one aspect, antigen binding fragments inhibit binding of $\alpha 4\beta 7$ integrin to one or more of its ligands (e.g. the mucosal addressin MAdCAM (e.g., MAdCAM-
10 1), fibronectin).

Papain digestion of antibodies produces two identical antigen binding fragments, called "Fab" fragments, each with a single antigen binding site, and a residual "Fc" fragment, whose name reflects its ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen binding sites and is still
15 capable of cross-linking antigen.

"Fv" is an antibody fragment which consists of a dimer of one heavy chain variable domain and one light chain variable domain in non-covalent association.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH₁) of the heavy chain. Fab' fragments differ from Fab
20 fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH₁ domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear at least one free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines
25 between them. Other chemical couplings of antibody fragments are also known.

"Single-chain Fv" or "scFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. In one aspect, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the scFv to form the desired
30 structure for antigen binding. For a review of scFv see Pluckthun in *The*

Pharmacology of Monoclonal Antibodies, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term “diabodies” refers to small antibody fragments with two antigen binding sites, which fragments comprise a variable heavy domain (V_H) connected to
5 a variable light domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger *et al.*, *Proc. Natl. Acad. Sci. USA*,
10 90:6444-6448 (1993).

A “full length antibody” is one which comprises an antigen binding variable region as well as a light chain constant domain (C_L) and heavy chain constant domains, C_{H1}, C_{H2} and C_{H3}. The constant domains may be native sequence constant domains (e.g. human native sequence constant domains) or amino acid sequence
15 variants thereof. In one aspect, the full length antibody has one or more effector functions.

An “amino acid sequence variant” antibody herein is an antibody with an amino acid sequence which differs from a main species antibody. Ordinarily, amino acid sequence variants will possess at least about 70%, at least about 80%, at least
20 about 85%, at least about 90%, or at least about 95% homology with the main species antibody. The amino acid sequence variants possess substitutions, deletions, and/or additions at certain positions within or adjacent to the amino acid sequence of the main species antibody, but retain antigen binding activity. Variations in sequence of the constant regions of the antibody will have less effect on the antigen
25 binding activity than variations in the variable regions. In the variable regions, amino acid sequence variants will be at least about 90% homologous, at least about 95% homologous, at least about 97% homologous, at least about 98% homologous, or at least about 99% homologous with the main species antibody.

“Homology” is defined as the percentage of residues in the amino acid
30 sequence variant that are identical after aligning the sequences and introducing gaps,

if necessary, to achieve the maximum percent homology. Methods and computer programs for the alignment are well known in the art.

A “therapeutic monoclonal antibody” is an antibody used for therapy of a human subject. Therapeutic monoclonal antibodies disclosed herein include anti-
5 $\alpha 4\beta 7$ antibodies.

A “glycosylation variant” antibody herein is an antibody with one or more carbohydrate moieties attached thereto which differ from one or more carbohydrate moieties attached to a main species antibody. Examples of glycosylation variants herein include antibody with a G1 or G2 oligosaccharide structure, instead of a G0
10 oligosaccharide structure, attached to an Fc region thereof, antibody with one or two carbohydrate moieties attached to one or two light chains thereof, antibody with no carbohydrate attached to one or two heavy chains of the antibody, etc, and combinations of glycosylation alterations.

Antibody “effector functions” refer to those biological activities attributable
15 to the Fc region (a native sequence Fc region or amino acid sequence variant Fc region) of an antibody. Examples of antibody effector functions include C1q binding; complement dependent cytotoxicity; Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor; BCR), and the like.

Depending on the amino acid sequence of the constant domain of their heavy
20 chains, full length antibodies can be assigned to different “classes”. There are five major classes of full length antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into “subclasses” (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2. The heavy-chain constant domains that correspond to the
25 different classes of antibodies are called α , δ , ϵ , γ , and μ , respectively. The subunit structures and three-dimensional configurations of different classes of immunoglobulins are well known.

The “light chains” of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the
30 amino acid sequences of their constant domains.

“Antibody-dependent cell-mediated cytotoxicity” and “ADCC” refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express FcγRIII only, whereas monocytes express FcγRI, FcγRII and FcγRIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in U.S. Pat. Nos. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be assessed *in vivo*, e.g., in an animal model such as that disclosed in Clynes *et al. PNAS (USA)* 95:652-656 (1998).

The terms “Fc receptor” or “FcR” are used to describe a receptor that binds to the Fc region of an antibody. In one aspect, the FcR is a native sequence human FcR. In another aspect, the FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors of the FcγRI, FcγRII, and FcγRIII subclasses, including allelic variants and alternatively spliced forms of these receptors. FcγRII receptors include FcγRIIA (an “activating receptor”) and FcγRIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor FcγRIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor FcγRIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain. (See, review in M. Daeron, *Annu. Rev. Immunol.* 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991); Capel *et al., Immunomethods* 4:25-34 (1994); and de Haas *et al., J. Lab. Clin. Med.* 126:33-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of

maternal IgGs to the fetus (Guyer *et al.*, *J. Immunol.* 117:587 (1976) and Kim *et al.*, *J. Immunol.* 24:249 (1994)).

The term “hypervariable region” when used herein refers to the amino acid residues of an antibody which are responsible for antigen binding. The
5 hypervariable region generally comprises amino acid residues from a “complementarity determining region” or “CDR” (e.g. residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and 31-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National
10 Institutes of Health, Bethesda, Md. (1991)) and/or those residues from a “hypervariable loop” (e.g. residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55 (H2) and 96-101 (H3) in the heavy chain variable domain; Chothia and Lesk *J. Mol. Biol.* 196:901-917 (1987)). “Framework Region” or “FR” residues are those variable domain residues other than
15 the hypervariable region residues as herein defined. The hypervariable region or the CDRs thereof can be transferred from one antibody chain to another or to another protein to confer antigen binding specificity to the resulting (composite) antibody or binding protein.

“Humanized” forms of non-human (e.g., rodent) antibodies are chimeric
20 antibodies that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a hypervariable region of the recipient are replaced by residues from a hypervariable region of a non-human species (donor antibody) such as mouse, rat, rabbit or nonhuman primate having the desired specificity,
25 affinity, and capacity. In some instances, framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, humanized antibodies may comprise residues that are not found in the recipient antibody or in the donor antibody. These modifications are made to further refine antibody performance. In general, the humanized antibody will comprise
30 substantially all of at least one, and typically two, variable domains, in which all or substantially all of the hypervariable loops correspond to those of a non-human

immunoglobulin and all or substantially all of the FRs are those of a human immunoglobulin sequence. The humanized antibody optionally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones *et al.*, *Nature* 321:522-525 (1986); Riechmann *et al.*, *Nature* 332:323-329 (1988); and Presta, *Curr. Op. Struct. Biol.* 2:593-596 (1992).

An "affinity matured" antibody is one with one or more alterations in one or more hypervariable regions thereof which result an improvement in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). In one aspect, affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are produced by procedures known in the art. Marks *et al. Bio/Technology* 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas *et al. Proc Nat. Acad. Sci, USA* 91:3809-3813 (1994); Schier *et al. Gene* 169:147-155 (1995); Yelton *et al. J. Immunol.* 155:1994-2004 (1995); Jackson *et al., J. Immunol.* 154(7):3310-9 (1995); and Hawkins *et al., J. Mol. Biol.* 226:889-896 (1992).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. In certain embodiments, the antibody will be purified (1) to greater than 95% by weight of protein as determined by the Lowry method, and alternatively, more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures. Those in need of treatment include those already with the disease as well as those in which the disease or its recurrence is to be prevented.

Hence, the patient to be treated herein may have been diagnosed as having the disease or may be predisposed or susceptible to the disease. The terms “patient” and “subject” are used interchangeably herein.

The antibody which is formulated is substantially pure and desirably
5 substantially homogeneous (i.e. free from contaminating proteins etc).
“Substantially pure” antibody means a composition comprising at least about 90% antibody by weight, based on total weight of the protein, alternatively, at least about 95% or 97% by weight. “Substantially homogeneous” antibody means a
composition comprising protein wherein at least about 99% by weight of protein is
10 specific antibody, e.g., anti- $\alpha 4\beta 7$ antibody, based on total weight of the protein.

“Clinical remission” as used herein with reference to ulcerative colitis subjects refers to a complete Mayo score of 2 or less points and no individual subscore greater than 1 point. Crohn’s disease “clinical remission” refers to a CDAI score of 150 points or less.

15 A “clinical response” as used herein with reference to ulcerative colitis subjects refers to a reduction in complete Mayo score of 3 or greater points and 30% from baseline, (or a partial Mayo score of 2 or greater points and 25% or greater from baseline, if the complete Mayo score was not performed at the visit) with an accompanying decrease in rectal bleeding subscore of 1 or greater points or absolute
20 rectal bleeding score of 1 or less point. A “clinical response” as used herein with reference to Crohn’s disease subjects refers to a 70 point or greater decrease in CDAI score from baseline (week 0).

“Mucosal healing” as used herein with reference to ulcerative colitis subjects refers to an endoscopic subscore of 1 point or less.

25 As used herein, “treatment failure” refers to disease worsening, a need for rescue medications or surgical intervention for treatment of ulcerative colitis or Crohn’s disease. A rescue medication is any new medication or any increase in dose of a baseline medication required to treat new or unresolved ulcerative colitis or Crohn’s disease symptoms (other than antidiarrheals for control of chronic diarrhea).

30

Formulations

As described herein, it has been discovered that anti- $\alpha 4\beta 7$ antibodies are more stable when formulated with an antioxidant or chelator. In addition, as described herein, anti- $\alpha 4\beta 7$ antibodies may be formulated to reduce aggregate formation (e.g., the amount of polysorbate 80 in the formulation may be reduced). For example, formulations that comprise citrate or EDTA and anti- $\alpha 4\beta 7$ antibodies decrease the rate of antibody aggregate formation during storage. Formulations may also be stored without oxygen to reduce aggregate formation. In one embodiment, the formulation has an antibody aggregate formation of less than about 2.5% at 25 °C after 12 months. In one embodiment, the formulation has an antibody aggregate formation of less than about 2.0% at 25 °C after 12 months. In one embodiment, the formulation has an antibody aggregate formation of less than about 1.6% at 25 °C after 12 months. In one embodiment, the formulation has an antibody aggregate formation of less than about 1.3% at 25 °C after 12 months. In one embodiment, the formulation has an antibody aggregate formation of less than about 1.0% at 25 °C after 12 months. In another embodiment, the formulation has an antibody aggregate formation of less than about 0.5% at 5°C after 12 months. In another embodiment, the formulation has an antibody aggregate formation of less than about 0.3% at 5°C after 12 months.

The present invention provides, in a first aspect, a stable anti- $\alpha 4\beta 7$ antibody formulation. The formulation comprises an anti- $\alpha 4\beta 7$ antibody and an antioxidant or chelator. The formulation also comprises a buffering agent that may be one or more free amino acids. The formulation may optionally further comprise a surfactant. The antibody in the formulation may be a full length antibody or an antigen binding fragment thereof, such as a Fab, Fv, scFv, Fab' or F(ab')₂ fragment.

Aggregate formation can be reduced by removing oxygen from the formulation. Alternatively, the formulation can contain an antioxidant or chelator. In one aspect, exemplary antioxidants and chelators that can be included in the formulation include lipoic acid, uric acid, glutathione, tocopherol, carotene, lycopene, cysteine, ethylenediaminetetraacetic acid (EDTA), ethyleneglycoltetraacetic acid (EGTA), dimercaprol, diethylenetriaminepentaacetic

acid, and N,N-bis(carboxymethyl)glycine, phosphonate compounds, e.g., etidronic acid, desferoxamine, malate and citrate. Some antioxidants and chelators can decrease the rate of aggregate formation during storage of the formulation. In another aspect, the chelator and/or antioxidant is citrate or EDTA. Exemplary
5 chelator concentrations for liquid formulations are in the range of from about greater than 0 mM to about 60 mM, about 5 mM to about 50 mM, about 5 mM to about 15 mM, about 10 mM to about 25 mM, and about 20 to about 30 mM. In another aspect, the chelator concentration is from about 0 mM to about 30 mM. In one embodiment, the chelator and/or antioxidant is citrate, and the citrate concentration
10 is from about 0 mM to about 15 mM, about 0 mM to about 10 mM, or about 0 mM to about 5 mM.

The formulation can contain any desired one free amino acid, which can be in the L-form, the D-form or any desired mixture of these forms. In one aspect, free amino acids that can be included in the formulation include, for example, histidine,
15 alanine, arginine, glycine, glutamic acid, serine, lysine, tryptophan, valine, cysteine and combinations thereof. Some amino acids can stabilize the proteins against degradation during manufacturing, drying, lyophilization and/or storage, e.g., through hydrogen bonds, salt bridges, antioxidant properties, or hydrophobic interactions or by exclusion from the protein surface. Amino acids can act as
20 tonicity modifiers or can act to decrease viscosity of the formulation. In another aspect, free amino acids, such as histidine and arginine, can act as lyoprotectants, and do not crystallize when lyophilized as components of the formulation. Free amino acids, such as glutamic acid and histidine, alone or in combination, can act as buffering agents in aqueous solution in the pH range of 5 to 7.5. In still yet another
25 aspect, the formulation contains histidine, arginine, or a combination of histidine and arginine. In still yet another aspect, free amino acid concentrations for liquid formulations are in the range from about 9 mM to about 0.5 M, for example, from about 10 mM to about 90 mM, about 10 mM to about 75 mM, about 10 mM to about 40 mM, about 25 mM to about 50 mM, about 15 mM to about 300 mM, about 20
30 mM to about 200 mM, about 25 mM to about 150 mM, about 50 mM to about 75

mM, about 50 mM to about 120 mM, about 50 to about 150 mM, or about 50 mM or about 125 mM.

- The formulation can optionally further contain at least one surfactant, e.g., to control soluble and insoluble aggregate formation. In one aspect, the surfactant is a non-ionic surfactant. In another aspect, the surfactant is an ionic surfactant. Exemplary surfactants that can be included in the formulation include, for example, polysorbate 20, polysorbate 80, a poloxamer (Pluronic®) and combinations thereof. When present, the surfactant is generally included in an amount which reduces formation of insoluble aggregates of antibody, e.g., during bottling, freezing, drying, lyophilization and/or reconstitution, in the presence of silicone, filling vials, prefilled syringes, and/or cartridges. The surfactant concentration is generally from about 0.0001% to about 1.0%, from about 0.01% to about 0.5%, for example, about 0.05%, 0.1%, 0.15%, 0.20%, 0.3%, 0.4%, or 0.5% (w/v). Higher concentrations of surfactant, e.g., polysorbate 80 can lead to more SEC aggregate formation.
- Reducing the concentration of polysorbate 80 can reduce SEC aggregate formation upon storage. In one aspect, the surfactant:antibody molar ratio is from about 0.7:1 to about 2.0:1. In another aspect, the surfactant:antibody molar ratio is 1.5:1.

- An embodiment of an anti- $\alpha 4\beta 7$ antibody formulation contains a high concentration of anti- $\alpha 4\beta 7$ antibody. For example, in one embodiment, the liquid formulations can comprise at least about 60 mg/ml, at least about 70 mg/ml, at least about 80 mg/ml, at least about 90 mg/ml, at least about 100 mg/ml, at least about 110 mg/ml, at least about 120 mg/ml, at least about 130 mg/ml, at least about 140 mg/ml, at least about 150 mg/ml, at least about 160 mg/ml, at least about 170 mg/ml, at least about 180 mg/ml, at least about 190 mg/ml, at least about 200 mg/ml, at least about 250 mg/ml, at least about 300 mg/ml, from about 60 mg/ml to about 190 mg/ml, from about 60 mg/ml to about 170 mg/ml anti- $\alpha 4\beta 7$ antibody, from about 150 mg/ml to about 180 mg/ml, or about 160 mg/ml or about 165 mg/ml anti- $\alpha 4\beta 7$ antibody. Alternatively, in another aspect, the liquid formulations can comprise at least about 154 mg/ml, at least about 176 mg/ml.

- The formulation can be a liquid or a solid. Liquid formulations are aqueous solutions or suspensions, prepared in a suitable aqueous solvent, such as water or an

aqueous/organic mixture, such as water alcohol mixtures. Liquid formulations have a pH between about 5.5 and about 7.5, between about 6.0 and 7.3, between about 6.0 and about 7.0, between about 6.0 and 6.5, between about 6.0 and 6.3, between about 6.3 and 7.1, or between about 6.4 and 7.0, or between 6.3 and 6.8, such as about 6.0,
5 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, or 6.9. Liquid formulations can be kept at room temperature, refrigerated (e.g., 2-8°C), or frozen (e.g., -20 °C or -70 °C) for storage.

A solid formulation can be prepared in any suitable way and can be in the form of a cake or powder, for example, with the addition of a lyoprotectant. In one aspect, the solid formulation is prepared by drying a liquid formulation as described
10 herein, for example by lyophilization or spray drying. When the formulation is a solid formulation, the formulation can have a moisture content of no more than about 5%, no more than about 4.5%, no more than about 4%, no more than about 3.5%, no more than about 3%, no more than about 2.5%, no more than about 2%, no more than about 1.5%, no more than about 1%, or is substantially anhydrous. A
15 solid formulation can be dissolved, i.e. reconstituted, in a suitable medium or solvent to become liquid suitable for administration. Suitable solvents for reconstituting the solid formulation include water, isotonic saline, buffer, e.g., phosphate-buffered saline, Ringer's (lactated or dextrose) solution, minimal essential medium, alcohol/aqueous solutions, dextrose solution, etc. The amount of solvent can result
20 in a therapeutic protein concentration higher, the same, or lower than the concentration prior to drying. In another aspect, the reconstituted anti- $\alpha 4\beta 7$ antibody concentration is the same concentration as in the pre-drying liquid formulation.

The formulation may be sterile, and this can be achieved according to the
25 procedures known to the skilled person for generating sterile pharmaceutical formulations suitable for administration to human subjects, prior to, or following, preparation of the formulation. The formulation can be sterilized as a liquid, e.g., before drying and/or after reconstitution by filtration through small pores, through aseptic processing or by exposure to ultraviolet radiation. Filter pore sizes can be
30 0.1 μm or 0.2 μm to filter microorganisms or 10 to 20 nm to filter virus particles. Alternatively, or additionally, the dried formulation can be sterilized, e.g., by

exposure to gamma radiation. In one aspect, the anti- $\alpha 4\beta 7$ antibody liquid formulation is sterilized by filtration before drying.

In one aspect, the formulation is stable upon storage. Various stability assays are available to the skilled practitioner for confirming the stability of the formulation. For example, the antibody in the liquid formulation may be stable upon storage at about 25°C for at least about 4 weeks, at least about 2 months, at least about 3 months, or at least about 6 months, or at least about 9 months, or at least about 12 months; at about 2-8°C at least about 3 months, at least about 1 year, at least about 2 years, at least about 3 years or longer. Alternatively or in addition, the antibody in the formulation may be stable upon storage at about 15°C for at least about 4 weeks, at least about 3 months, at least about 6 months, at least about 9 months, at least about 1 year, or longer. Alternatively or in addition, the antibody in the formulation may be stable upon storage at about -20°C or -70°C for at least about 4 weeks; at least about 3 months, at least about 6 months, at least about 9 months, at least about 1 year, at least about 2 years, at least about 3 years, at least about 4 years or longer.

Stability can be tested by evaluating physical stability, chemical stability, and/or biological activity of the antibody in the formulation around the time of formulation as well as following storage at the noted temperatures. Physical and/or chemical stability of a liquid formulation or a reconstituted dry powder can be evaluated qualitatively and/or quantitatively in a variety of different ways (see, e.g., *Analytical Techniques for Biopharmaceutical Development*, Rodriguez-Diaz *et al.* eds. Informa Healthcare (2005)), including evaluation of soluble and insoluble aggregate formation (for example using size exclusion chromatography, analytical ultracentrifugation, MALDI-TOF MS, light scattering (dynamic (DLS) or MALLS), flow-based microscopic imaging, or other liquid particle counting system, by measuring turbidity, by density gradient centrifugation and/or by visual inspection); by assessing charge heterogeneity using cation exchange chromatography (see also Vlasak and Ionescu, *Curr. Pharm. Biotechnol.* 9:468-481 (2008) and Harris *et al. J. Chromatogr. B Biomed. Sci. Appl.* 752:233-245 (2001)), isoelectric focusing or capillary zone electrophoresis; amino-terminal or carboxy terminal sequence

analysis; mass spectrometric analysis; SDS-PAGE analysis to compare fragmented, intact and multimeric (i.e., dimeric, trimeric, etc.) antibody; peptide map (for example tryptic or LYS-and the like). Instability may result in aggregation, deamidation (e.g., Asn deamidation), oxidation (e.g., Met oxidation), isomerization (e.g., Asp isomerization), denaturation, clipping/hydrolysis/fragmentation (e.g., hinge region fragmentation), succinimide formation, unpaired cysteine(s), N-terminal extension, C-terminal processing, glycosylation differences, etc. Biological activity or antigen binding function, e.g., binding of the anti- $\alpha 4\beta 7$ antibody to MAdCAM (e.g., MAdCAM-1) or inhibition of the binding of a cell expressing $\alpha 4\beta 7$ integrin to MAdCAM (e.g., MAdCAM-1), e.g., immobilized MAdCAM (e.g., MAdCAM-1), can be evaluated using various techniques available to the skilled practitioner (see e.g., Soler *et al.*, *J. Pharmacol. Exper. Ther.* 330:864-875 (2009)). Measurement of the moisture content of a dry formulation can indicate how likely a formulation will undergo chemical or physical degradation, with higher moisture leading to more degradation.

A stable formulation can contribute to a low immunogenicity of an anti- $\alpha 4\beta 7$ antibody. An immunogenic anti- $\alpha 4\beta 7$ antibody can lead to a human-anti-human antibody (HAHA) response in human subjects or patients. Patients who develop a HAHA response to an anti- $\alpha 4\beta 7$ antibody can have adverse events (e.g., site infusion reaction) upon treatment or can eliminate anti- $\alpha 4\beta 7$ antibody quickly, resulting in a lower dose than planned by treatment. A report (Feagen et al. (2005) *N. Engl. J. Med.* 352:2499-2507) of early study of an anti- $\alpha 4\beta 7$ antibody treatment indicated that human antihuman antibodies developed by week 8 in 44% of treated patients. The antibody in this study was stored as a liquid and did not contain any polysorbate.

In some embodiments, the formulation can increase the proportion of HAHA negative patients to at least 40%, at least 50%, at least 60%, at least 70%, at least 80% or at least 90% of patients compared to the HAHA results of a less stable formulation.

In some embodiments, an anti- $\alpha 4\beta 7$ antibody formulation has $\geq 50\%$ major charged isoform, $\geq 55\%$ major charged isoform, or 65 to 70% major charged

isoform. In other aspects, a stable anti- $\alpha 4\beta 7$ antibody formulation has $\leq 45\%$ acidic charged isoforms, $\leq 40\%$ acidic charged isoforms, $\leq 30\%$ acidic charged isoforms or 22 to 28% acidic isoforms. In still other aspects, a stable anti- $\alpha 4\beta 7$ antibody formulation has $\leq 25\%$ basic isoforms, $\leq 20\%$ basic isoforms, $\leq 15\%$ basic isoforms, about 5% basic isoforms or about 10% basic isoforms. In one aspect, a stable anti- $\alpha 4\beta 7$ antibody formulation has $\geq 55\%$ major isoform, $\leq 30\%$ acidic isoforms and/or $\leq 20\%$ basic isoforms, e.g., as determined by CEX. In another aspect, a stable anti- $\alpha 4\beta 7$ antibody formulation has $\geq 50\%$ major isoform, $\leq 45\%$ acidic isoforms and/or $< 10\%$ basic isoforms, e.g., as determined by cIEF.

10 In some aspects, an anti- $\alpha 4\beta 7$ antibody dry, solid formulation has $\leq 10\%$ moisture content, $\leq 5\%$ moisture content or $< 2.5\%$ moisture content. The time required for reconstitution is ≤ 60 minutes, ≤ 50 minutes or ≤ 40 minutes or ≤ 30 minutes or ≤ 20 minutes.

Monomeric content and/or aggregate content (e.g., as dimers, trimers, tetramers, pentamers, oligomers and higher-order aggregates), i.e., in the liquid formulation, or in the reconstituted formulation, can be measured by SEC, analytical ultracentrifugation, light scattering (DLS or MALLS), MALDI-TOF MS or nanoscale measurement, such as nanoparticle tracking analysis NTA, NanoSight Ltd, Wiltshire, UK). Resolution, characterization and quantification of aggregate can be achieved in a number of ways, including increasing the length of the SEC column separation, e.g., by a longer column or by serial attachment of a second or more SEC column(s) in line with the initial analytical SEC column, supplementing SEC quantification of monomers with light scattering, or by using NTA.

25 In one embodiment, an anti- $\alpha 4\beta 7$ antibody formulation has $\geq 90\%$ monomeric antibody, $\geq 95\%$ monomeric antibody, or 97 to 99% monomeric antibody. In another embodiment, the majority of the material in an anti- $\alpha 4\beta 7$ antibody formulation has an average radius of ≤ 20 nm, ≤ 15 nm, ≤ 10 nm, or about 5 to about 7 nm. In one aspect, an anti- $\alpha 4\beta 7$ antibody formulation has $\geq 80\%$ amount heavy plus light chain by protein analysis. In one aspect, there is $\geq 90\%$ heavy plus light chain. In another aspect, an anti- $\alpha 4\beta 7$ antibody formulation has \leq

10% aggregate, $\leq 5\%$ aggregate, $\leq 2.5\%$ aggregate, $\leq 1.5\%$ aggregate, $\leq 1.0\%$ aggregate or $\leq 0.5\%$ aggregate. In another aspect, stable anti- $\alpha 4\beta 7$ antibody formulation has $\geq 96\%$ monomer and/or $\leq 2.5\%$ aggregate. In yet another aspect, a stable anti- $\alpha 4\beta 7$ antibody formulation has about 99% monomer and/or about $< 1\%$ aggregate.

Particle sizes, greater than 1 to 2 micron, e.g., of aggregates or undissolved excipient, i.e., in the liquid formulation, or in the reconstituted formulation can be measured by light obscuration (e.g., liquid particle counting system (HIAC) by Hach Ultra Analytics (Grants Pass, OR)), microscopy, coulter counter, or digital (e.g., flow-based) microscopic imaging based system such as microfluidics imaging (MFI) by Brightwell (Ottawa, CA) or FLOWCAM® Image particle analyzer by Fluid Imaging Technologies (Yarmouth, ME). In one aspect, particle size in an anti- $\alpha 4\beta 7$ antibody preparation is about 30 μm , about 25 μm , about 10 μm , about 5 μm , about 2 μm or 1 μm or less. The amount of particles should be minimized in antibody formulations. In one aspect, an amount of particles in an anti- $\alpha 4\beta 7$ antibody formulation is < 6000 particles $\geq 10 \mu\text{m}$ diameter and/or < 600 particles $\geq 25 \mu\text{m}$ diameter in one dose (U.S. Pharmacopoeia Chp. 788, light obscuration counting method; half those amounts by microscopic quantification method). In another aspect, an amount of particles in a dose of an anti- $\alpha 4\beta 7$ antibody formulation is about 1000 particles $\geq 10 \mu\text{m}$ and about 0-100 particles $\geq 25 \mu\text{m}$ (MFI method). In yet another aspect, an amount of particles per milliliter, e.g., by MFI measurement, in a dose of an anti- $\alpha 4\beta 7$ antibody formulation is about 500 to about 2000 of 2-10 μm particles per ml, about 50 to about 350 of $\geq 10 \mu\text{m}$ particles per ml and about 0 to about 50 of $\geq 25 \mu\text{m}$ particles per ml. In yet another aspect, an amount of particles in a dose of an anti- $\alpha 4\beta 7$ antibody formulation is about 500 to about 100,000, about 1000 to about 5000 or about 1500 to about 3000 of 2-10 μm particles per ml.

The viscosity of an anti- $\alpha 4\beta 7$ antibody formulation can be controlled for subcutaneous or intramuscular administration. The viscosity can be affected by protein concentration and pH. For example, as the protein concentration increases, the viscosity can increase. An increase in pH can decrease the viscosity of the anti-

$\alpha 4\beta 7$ antibody formulation. In some protein formulations, sodium chloride is added to reduce the viscosity of the formulation. Additional components that can affect viscosity of an anti- $\alpha 4\beta 7$ antibody formulation are amino acids such as histidine and arginine.

5 An anti- $\alpha 4\beta 7$ antibody formulation can be isotonic (e.g., 250-350 mOsm) or hypertonic (e.g., greater than 350 mOsm, greater than 450 mOsm, greater than 550 mOsm or greater than 650 mOsm), e.g., for subcutaneous or intramuscular administration. In one aspect, the anti- $\alpha 4\beta 7$ antibody formulation is not hypotonic, e.g., less than 250 mOsm. In another aspect, the anti- $\alpha 4\beta 7$ antibody formulation is
10 about 350 to about 400 mOsm, about 400 to about 450 mOsm or about 350 to about 450 mOsm.

 Instability leading to denaturation can be assessed by differential scanning calorimetry (DSC). Antibodies have two melting temperatures (T_m) in DSC, e.g., T_{m1} and T_{m2} . Certain excipients can affect the stability of the native anti- $\alpha 4\beta 7$
15 antibody. A finding of a higher melting temperature when comparing formulations by DSC can indicate a more stable anti- $\alpha 4\beta 7$ antibody formulation with the higher T_m . For example, at pH 5.7, the T_m of an anti- $\alpha 4\beta 7$ antibody formulation is lower, and thus less stable than at pH 6.5. In one aspect, T_{m1} of an anti- $\alpha 4\beta 7$ antibody formulation is $>60^\circ\text{C}$. In another aspect, the T_{m1} of an anti- $\alpha 4\beta 7$ antibody
20 formulation is about 65°C to about 70°C or about 69°C . In one aspect, T_{m2} of an anti- $\alpha 4\beta 7$ antibody formulation is $>80^\circ\text{C}$. In another aspect, the T_{m2} of an anti- $\alpha 4\beta 7$ antibody formulation is about 82°C to about 88°C or about 86°C .

 In one embodiment, an anti- $\alpha 4\beta 7$ antibody formulation has a binding affinity or EC_{50} value of about 60% to about 140% of the reference standard anti- $\alpha 4\beta 7$
25 antibody. In one aspect, an anti- $\alpha 4\beta 7$ antibody in a formulation described herein binds to $\alpha 4\beta 7$, e.g., on a cell (WO98/06248 or U.S. Patent No. 7,147,851), at a value of about 80% to about 120% of the reference standard. In another embodiment, an anti- $\alpha 4\beta 7$ antibody formulation has the ability to inhibit at least 50%, or at least 60% of the binding of a cell expressing $\alpha 4\beta 7$ integrin to MAdCAM (e.g., MAdCAM-1),
30 e.g., a MAdCAM-Ig chimera (see U.S. Patent Application Publication No. 20070122404, also for reference standard examples).

As noted above, freezing of the formulation is specifically contemplated herein. Hence, the formulation can be tested for stability upon freezing and thawing. Accordingly, the antibody in a liquid formulation may be stable upon freezing and thawing the formulation, for example the antibody can be stable after one, two,
5 three, four, five or more freeze/thaw cycles.

In some embodiments, the pharmaceutical formulation is a liquid formulation comprising at least about 60 mg/ml to about 170 mg/ml anti- $\alpha 4\beta 7$ antibody, a buffering agent (e.g., histidine), and at least about 5 mM citrate. In other
10 mg/ml to about 170 mg/ml anti- $\alpha 4\beta 7$ antibody, a buffering agent (e.g., citrate), amino acid (e.g., arginine) and surfactant (e.g., polysorbate 80).

In another embodiment, the formulation comprises at least about 140 mg/ml or about 150 mg/ml to about 170 mg/ml, for example, about 160 mg/ml of an anti- $\alpha 4\beta 7$ antibody, a buffering agent (e.g., histidine), at least about 5 mM citrate and a
15 free amino acid (e.g., arginine).

In yet another embodiment, the formulation comprises at least about 160 mg/ml of an anti- $\alpha 4\beta 7$ antibody, a buffering agent (e.g., histidine), at least about 5 mM citrate, 0.2% polysorbate 80, and a free amino acid (e.g., arginine). In an
20 embodiment, the buffer concentration in the formulation is about 15 to about 75 mM, about 25 to about 65 mM, or about 50 mM. The free amino acid concentration in the formulation is about 50 to about 250 mM, about 75 to about 200 mM, about 100 to about 150 mM or about 125 mM; the polysorbate 80 concentration in the formulation is about 0.05% to 0.4%, about 0.1% to 0.4%, about 0.1% to 0.3%, about 0.1% to 0.25%, about 0.1% to 0.2%, or about 0.2%.

25 In some embodiments, the formulation is a solid formulation (e.g., a lyophilized formulation), comprising a mixture of an anti- $\alpha 4\beta 7$ antibody, citrate, histidine, arginine, polysorbate 80, and a lyoprotectant or a saccharide, such as a non-reducing sugar. Saccharide can be included in the liquid formulation to reach concentrations from 0% to 20%, or about 6% to about 10%.

30 In one embodiment, the formulation is lyophilized and stored as a single dose in one container, e.g., vial, syringe, cartridge, and/or autoinjector. The

container can be stored at about 2-8°C or 25 °C until it is administered to a subject in need thereof. The vial may for example be a 5, 10 or 20 cc vial (for example for a 160 mg/ml dose). The vial may contain at least about 20 mg, at least about 50 mg, at least about 70 mg, at least about 80 mg, at least about 100 mg, at least about 120
5 mg, at least about 155 mg, at least about 180 mg, at least about 200 mg, at least about 240 mg, at least about 300 mg, at least about 360 mg, at least about 400 mg, at least about 540 mg, or at least about 900 mg of anti- α 4 β 7 antibody. In one aspect, the container contains about 165 mg of anti- α 4 β 7 antibody.

In another embodiment, the formulation is liquid and stored as a single dose
10 in one or two vials, cartridges, syringes, or autoinjectors. The vial, cartridge, syringe, or autoinjector can be stored at about 2-8°C until its contents, e.g., an anti- α 4 β 7 antibody, are administered to a subject in need thereof. The vial may, for example, be a 5, 10 or 20 cc vial (for example for a 160 mg/ml dose). The vial may contain at least about 20 mg, at least about 50 mg, at least about 70 mg, at least
15 about 80 mg, at least about 100 mg, at least about 120 mg, at least about 155 mg, at least about 180 mg, at least about 200 mg, at least about 240 mg, at least about 300 mg, at least about 360 mg, at least about 400 mg, at least about 540 mg, or at least about 900 mg of anti- α 4 β 7 antibody. In one aspect, the vial contains about 165 mg of anti- α 4 β 7 antibody. The syringe or cartridge may be a 1 mL or 2 mL container
20 (for example for a 160 mg/mL dose) or more than 2 ml, e.g., for a higher dose (at least 320 mg or 400 mg or higher). The syringe or cartridge may contain at least about 20 mg, at least about 50 mg, at least about 70 mg, at least about 80 mg, at least about 100 mg, at least about 120 mg, at least about 155 mg, at least about 180 mg, at least about 200 mg, at least about 240 mg, at least about 300 mg, at least about 360
25 mg, at least about 400 mg, or at least about 500 mg of anti- α 4 β 7 antibody.

One or more other pharmaceutically acceptable carriers, excipients or stabilizers such as those described in *Remington: The Science and Practice of Pharmacy*, 21st Edition, Hendrickson, R. Ed. (2005) may be included in the formulation provided that they do not adversely affect the desired characteristics of
30 the formulation. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed and include; additional

buffering agents; co-solvents; antioxidants including citrate and cysteine; chelating agents such as EDTA; metal complexes (e.g., Zn-protein complexes); biodegradable polymers such as polyesters; preservatives; container wall lubricants, e.g., silicone, mineral oil, glycerin, or TRIBOGLIDE® (Tribo Film Research, Inc.)

- 5 perfluoropolyether derivative, for injection ease and/or salt-forming counterions such as sodium.

$\alpha 4\beta 7$ Antibodies

- 10 Anti- $\alpha 4\beta 7$ antibodies suitable for use in the formulations include antibodies from any desired source, such as fully human antibodies, murine antibodies, rabbit antibodies and the like, and any desired engineered antibodies, such as chimeric antibodies, humanized antibodies, and the like. Antigen-binding fragments of any of these types of antibodies, such as Fab, Fv, scFv, Fab' and F(ab')₂ fragments, are also suitable for use in the formulations.

- 15 The anti- $\alpha 4\beta 7$ antibody can bind to an epitope on the $\alpha 4$ chain (e.g., humanized MAb 21.6 (Bendig *et al.*, U.S. Pat. No. 5,840,299)), on the $\beta 7$ chain (e.g., FIB504 or a humanized derivative (e.g., Fong *et al.*, U.S. Pat. No. 7,528,236)), or to a combinatorial epitope formed by the association of the $\alpha 4$ chain with the $\beta 7$ chain. In one aspect, the antibody binds a combinatorial epitope on the $\alpha 4\beta 7$ complex, but does not bind an epitope on the $\alpha 4$ chain or the $\beta 7$ chain unless the chains are in association with each other. The association of $\alpha 4$ integrin with $\beta 7$ integrin can create a combinatorial epitope for example, by bringing into proximity residues present on both chains which together comprise the epitope or by conformationally exposing on one chain, e.g., the $\alpha 4$ integrin chain or the $\beta 7$ integrin chain, an epitopic binding site that is inaccessible to antibody binding in the absence of the proper integrin partner or in the absence of integrin activation. In another aspect, the anti- $\alpha 4\beta 7$ antibody binds both the $\alpha 4$ integrin chain and the $\beta 7$ integrin chain, and thus, is specific for the $\alpha 4\beta 7$ integrin complex. Such antibodies can bind $\alpha 4\beta 7$ but not bind $\alpha 4\beta 1$, and/or not bind $\alpha_E\beta 7$, for example. In another aspect, the anti- $\alpha 4\beta 7$ antibody binds to the same or substantially the same epitope as the Act-1 antibody (Lazarovits, A. I. *et al.*, *J. Immunol.*, 133(4): 1857-1862 (1984),
- 20
25
30

Schweighoffer *et al.*, *J. Immunol.*, 151(2): 717-729, 1993; Bednarczyk *et al.*, *J. Biol. Chem.*, 269(11): 8348-8354, 1994). Murine ACT-1 Hybridoma cell line, which produces the murine Act-1 monoclonal antibody, was deposited under the provisions of the Budapest Treaty on Aug. 22, 2001, on behalf of Millennium Pharmaceuticals, Inc., 40 Landsdowne Street, Cambridge, Mass. 02139, U.S.A., at the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110-2209, U.S.A., under Accession No. PTA-3663. In another aspect, the anti- $\alpha 4\beta 7$ antibody is a human antibody or an $\alpha 4\beta 7$ binding protein using the CDRs provided in U.S. Patent Application Publication No. 2010/0254975.

10 In one aspect, the anti- $\alpha 4\beta 7$ antibody inhibits binding of $\alpha 4\beta 7$ to one or more of its ligands (e.g. the mucosal addressin, e.g., MAdCAM (e.g., MAdCAM-1), fibronectin, and/or vascular addressin (VCAM)). Primate MAdCAMs (e.g., MAdCAM-1) are described in the PCT publication WO 96/24673. In another aspect, the anti- $\alpha 4\beta 7$ antibody inhibits binding of $\alpha 4\beta 7$ to MAdCAM (e.g.,
15 MAdCAM-1) and/or fibronectin without inhibiting the binding of VCAM.

 In one aspect, the anti- $\alpha 4\beta 7$ antibodies for use in the formulations are humanized versions of the mouse Act-1 antibody. Suitable methods for preparing humanized antibodies are well-known in the art. Generally, the humanized anti-
20 $\alpha 4\beta 7$ antibody will contain a heavy chain that contains the 3 heavy chain complementarity determining regions (CDRs, CDR1, SEQ ID NO:8, CDR2, SEQ ID NO:9 and CDR3, SEQ ID NO:10) of the mouse Act-1 antibody and suitable human heavy chain framework regions; and also contain a light chain that contains the 3 light chain CDRs (CDR1, SEQ ID NO:11, CDR2, SEQ ID NO:12 and CDR3,
25 SEQ ID NO:13) of the mouse Act-1 antibody and suitable human light chain framework regions. The humanized Act-1 antibody can contain any suitable human framework regions, including consensus framework regions, with or without amino acid substitutions. For example, one or more of the framework amino acids can be replaced with another amino acid, such as the amino acid at the corresponding
30 position in the mouse Act-1 antibody. The human constant region or portion thereof, if present, can be derived from the κ or λ light chains, and/or the γ (e.g., $\gamma 1$,

$\gamma 2$, $\gamma 3$, $\gamma 4$), μ , α (e.g., $\alpha 1$, $\alpha 2$), δ or ϵ heavy chains of human antibodies, including allelic variants. A particular constant region (e.g., IgG1), variant or portions thereof can be selected in order to tailor effector function. For example, a mutated constant region (variant) can be incorporated into a fusion protein to minimize binding to Fc
5 receptors and/or ability to fix complement (see e.g., Winter *et al.*, GB 2,209,757 B; Morrison *et al.*, WO 89/07142; Morgan *et al.*, WO 94/29351, Dec. 22, 1994). Humanized versions of Act-1 antibody were described in PCT publications nos. WO98/06248 and WO07/61679.

10 In another aspect, the anti- $\alpha 4\beta 7$ humanized antibodies for use in the formulation comprise a heavy chain variable region comprising amino acids 20 to 140 of SEQ ID NO:2, and a light chain variable region comprising amino acids 20 to 131 of SEQ ID NO:4 or amino acids 21 to 132 of SEQ ID NO:5. If desired, a suitable human constant region(s) can be present. For example, the humanized anti-
15 $\alpha 4\beta 7$ antibody can comprise a heavy chain that comprises amino acids 20 to 470 of SEQ ID NO:2 and a light chain comprising amino acids 21 to 239 of SEQ ID NO:5. In another example, the humanized anti- $\alpha 4\beta 7$ antibody can comprise a heavy chain that comprises amino acids 20 to 470 of SEQ ID NO:2 and a light chain comprising amino acids 20 to 238 of SEQ ID NO:4. Figure 4 shows an alignment which
20 compares the generic light chains of human antibodies with murine antibodies. The alignment illustrates that the humanized light chain of vedolizumab (e.g., Chemical Abstract Service (CAS, American Chemical Society) Registry number 943609-66-3), with two mouse residues switched for human residues, is more human than the light chain of LDP-02 (Figure 3). In addition, LDP-02 has the somewhat
25 hydrophobic, flexible alanine 114 and a hydrophilic site (Aspartate 115) that is replaced in vedolizumab with the slightly hydrophilic hydroxyl-containing threonine 114 and hydrophobic, potentially inward facing valine 115 residue.

Further substitutions to the antibody sequence can be, for example, mutations to the heavy and light chain framework regions, such as a mutation of isoleucine to
30 valine on residue 2 of SEQ ID NO:14; a mutation of methionine to valine on residue 4 of SEQ ID NO:14; a mutation of alanine to glycine on residue 24 of SEQ ID

NO:15; a mutation of arginine to lysine at residue 38 of SEQ ID NO:15; a mutation of alanine to arginine at residue 40 of SEQ ID NO:15; a mutation of methionine to isoleucine on residue 48 of SEQ ID NO:15; a mutation of isoleucine to leucine on residue 69 of SEQ ID NO:15; a mutation of arginine to valine on residue 71 of SEQ ID NO:15; a mutation of threonine to isoleucine on residue 73 of SEQ ID NO:15; or any combination thereof; and replacement of the heavy chain CDRs with the CDRs (CDR1, SEQ ID NO:8, CDR2, SEQ ID NO:9 and CDR3, SEQ ID NO:10) of the mouse Act-1 antibody; and replacement of the light chain CDRs with the light chain CDRs (CDR1, SEQ ID NO:11, CDR2, SEQ ID NO:12 and CDR3, SEQ ID NO:13) of the mouse Act-1 antibody.

In some embodiments, the anti- $\alpha 4\beta 7$ humanized antibodies for use in the formulation comprise a heavy chain variable region that has about 95%, 96%, 97%, 98%, or 99% sequence identity to amino acids 20 to 140 of SEQ ID NO:2, and a light chain variable region that has about 95%, 96%, 97%, 98%, or 99% sequence identity to amino acids 20 to 131 of SEQ ID NO:4 or amino acids 21 to 132 of SEQ ID NO:5. Amino acid sequence identity can be determined using a suitable sequence alignment algorithm, such as the Lasergene system (DNASTAR, Inc., Madison, Wis.), using the default parameters. In an embodiment, the anti- $\alpha 4\beta 7$ antibody for use in the formulation is vedolizumab (CAS, American Chemical Society, Registry number 943609-66-3).

Other $\alpha 4\beta 7$ antibodies may also be used in the formulations and dosing regimes described herein. For example, the $\alpha 4\beta 7$ antibodies described in US 2010/0254975 (Amgen, Inc.), are suitable for use in the formulations and methods of treating inflammatory bowel disease in an individual.

The anti- $\alpha 4\beta 7$ antibody can be produced by expression of nucleic acid sequences encoding each chain in living cells, e.g., cells in culture. A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide

coding sequences, express an anti- $\alpha 4\beta 7$ antibody in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*)
5 transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression
10 vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3, NS0 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). For example,
15 mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus, is an effective expression system for antibodies (Foecking *et al.*, *Gene* 45:101 (1986); Cockett *et al.*, *Bio/Technology* 8:2 (1990)).

In bacterial systems, a number of expression vectors may be advantageously
20 selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli*
25 expression vector pUR278 (Ruther *et al.*, *EMBO J.* 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, *Nucleic Acids Res.* 13:3101-3109 (1985); Van Heeke & Schuster, *J. Biol. Chem.* 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to
30 express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from

lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

5 In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin
10 promoter).

 In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This
15 chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts (e.g., see Logan & Shenk, *Proc. Natl. Acad. Sci. USA* 81:355-359 (1984)). Specific initiation signals may also be required for
20 efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The
25 efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner *et al.*, *Methods in Enzymol.* 153:51-544 (1987)).

 In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific
30 fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein.

Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products.

Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end,

5 eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to Chinese hamster ovary (CHO), NS0, HeLa, VERY, baby hamster kidney (BHK), monkey kidney (COS), MDCK, 293, 3T3, WI38, human hepatocellular carcinoma cells (e.g., Hep
10 G2), breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

The glycosylation machinery of different cell types can produce antibodies with different glycosylation composition than in another cell type, or no
15 glycosylation, as with bacterial cells. In one aspect, cell types for production of the anti- $\alpha\beta 7$ antibody are mammalian cells, such as NS0 or CHO cells. In one aspect, the mammalian cells can comprise the deletion of an enzyme involved in cell metabolism and the exogenous gene of interest can be operably linked to a replacement enzyme, e.g., in a construct or vector for introduction into the cells, e.g.,
20 by transformation or transfection. The construct or vector with the exogenous gene confers to the cells which host the construct or vector a selection advantage to encourage production of the polypeptide encoded by the exogenous gene. In one embodiment, CHO cells are DG44 cells (Chasin and Urlaub (1980) *PNAS USA* 77:4216), comprising the deletion or inactivation of the dihydrofolate reductase
25 gene. In another embodiment, CHO cells are CHO K1 cells comprising the deletion or inactivation of the glutamine synthase gene (see, e.g., U.S. Patent Nos. 5,122,464 or 5,827,739).

Solid Formulations

Solid formulations of the invention are generally prepared by drying a liquid
30 formulation. Any suitable method of drying can be used, such as lyophilization or

spray drying. In one aspect, a lyoprotectant is added to the formulation prior to lyophilization. Lyophilization involves freezing a liquid formulation, usually in the container that will be used to store, ship and distribute the formulation (e.g., a vial, syringe (e.g., a single- or dual-chamber syringe), or cartridge (e.g., a single- or dual-chamber cartridge) (See, e.g., Gatlin and Nail in *Protein Purification Process Engineering*, ed. Roger G. Harrison, Marcel Dekker Inc., 317-367 (1994).) Once the formulation is frozen, the atmospheric pressure is reduced and the temperature is adjusted to allow removal of the frozen solvent e.g., through sublimation. This step of the lyophilization process is sometimes referred to as primary drying. If desired, the temperature can then be raised to remove any solvent that is still bound to the dry formulation by evaporation. This step of the lyophilization process is sometimes referred to as secondary drying. When the formulation has reached the desired degree of dryness, the drying process is concluded and the containers are sealed. The final solid formulation is sometimes referred to as a "lyophilized formulation" or a "cake." The lyophilization process can be performed using any suitable equipment. Suitable lyophilization equipment is available from a number of commercial sources (e.g., SP Scientific, Stone Ridge, NY).

A variety of suitable apparatuses can be used to dry liquid formulations to produce a solid (e.g., lyophilized) formulation. Generally, lyophilized formulations are prepared by those of skill in the art using a sealed chamber that contains shelves, on which vials of the liquid formulation to be dried are placed. The temperature of the shelves, as well as cooling and heating rate can be controlled, as can the pressure inside the chamber. It will be understood that various process parameters discussed herein refer to processes performed using this type of apparatus. Persons of ordinary skill can easily adapt the parameters described herein to other types of drying apparatuses if desired.

Suitable temperatures and the amount of vacuum for primary and secondary drying can be readily determined by a person of ordinary skill. In general, the formulation is frozen at a temperature of about -30°C or less, such as -40°C or -50°C. The rate of cooling can affect the amount and size of ice crystals in the matrix. Primary drying is generally conducted at a temperature that is about 10°C,

about 20°C, about 30°C, about 40°C or about 50°C warmer than the freezing temperature. In one aspect, the primary drying conditions can be set to maintain the anti- $\alpha 4\beta 7$ antibody below the glass transition temperature or collapse temperature of the formulation. Above the collapse temperature, the amorphous frozen matrix can flow (collapse), with a result that the protein molecules may not be surrounded by a rigid, solid matrix, and the protein molecules may not be stable in the collapsed matrix. Also, the formulation can be difficult to fully dry if collapse occurs. The resulting higher amounts of moisture in the formulation can lead to higher rates of protein degradation and a decrease in the amount of time that the lyophilized product can be stored before its quality diminishes to unacceptable levels. In one aspect, the shelf temperature and chamber pressure are selected to maintain the product temperature below the collapse temperature during primary drying. The glass transition temperature of a frozen formulation can be measured by methods known in the art, e.g., by differential scanning calorimetry (DSC). The collapse temperature can be measured by methods known in the art, e.g. freeze-drying microscopy, optical coherence tomography. The drying step can remove at least 50%, at least 60%, at least 70% or more of the solvent. In one aspect, the primary drying step removes more than 80% of the solvent from the anti- $\alpha 4\beta 7$ antibody formulation.

Vial size can be selected based on the surface area which will be exposed to the shelf and to the vacuum during lyophilization. Drying time is directly proportional to cake height, thus the vial size may be chosen based upon what is determined to be a reasonable cake height. A vial with a large diameter relative to volume can provide a high amount of contact with the shelf for efficient heat transfer during the lyophilization cycle. A dilute antibody solution in a high volume of liquid will require more time for drying. A balance in vial size versus formulation volume needs to be struck, because larger vials can be more expensive to store and ship and have a larger headspace to formulation ratio, and may expose a high proportion of the formulation to the degradative effects of moisture during long term storage. For a 165 mg dose, the vial size of the anti- $\alpha 4\beta 7$ antibody formulation

can be 3 mL, 5 ml or 10 ml. In one aspect, the vial size is 5 ml for a 160 mg/ml solution.

The principles for choosing a cartridge or syringe size for lyophilization are similar to that of the vial. The depth of the cake height will also increase the drying
5 time as the height increases. The diameter and size of the syringe or cartridge must be balanced out with the final formulation volume. Larger diameters can increase the rate of moisture uptake in the lyophilized cake, thus increasing the degradative effects of moisture during storage. For a 165 mg dose, the anti- $\alpha 4\beta 7$ antibody formulation volume can be 1 ml or 2 mL. In one aspect, the syringe or cartridge size
10 is greater than 1 mL for a 160 mg/mL solution.

After lyophilization, the vial, syringe, or cartridge can be sealed, e.g., stoppered, under a vacuum. Alternatively, a gas, e.g., dry air or nitrogen, can be allowed into the container prior to sealing. Where oxidation is a concern, the gas allowed into the lyophilization chamber can comprise a gas which retards or
15 prevents oxidation of the lyophilized product. In one aspect the gases are non-oxygenated gases, e.g., nitrogen, or an inert gas, e.g., helium, neon, argon, krypton or xenon. In another aspect, the gas is nitrogen or argon.

Treatment With the Antibody Formulation

20 In one aspect, the invention provides a method of treating a disease or disorder in a subject comprising administering to a subject the anti- $\alpha 4\beta 7$ antibody formulation described herein in an amount effective to treat the disease or disorder, e.g., in humans. The human subject may be an adult (e.g., 18 years or older), an adolescent, or a child. The human subject may be a person 65 years or older. In
25 contrast to alternative therapeutic dosing regimens, a human subject 65 years or older does not require any modification of the dosing regimen described herein, and may be administered the conventional anti- $\alpha 4\beta 7$ antibody formulation described herein.

The subject may have had a lack of an adequate response with, loss of
30 response to, or was intolerant to treatment with an immunomodulator, a TNF- α antagonist, or combinations thereof. The patient may have previously received

treatment with at least one corticosteroid (e.g., prednisone) for the inflammatory bowel disease. An inadequate response to corticosteroids refers to signs and symptoms of persistently active disease despite a history of at least one 4-week induction regimen that included a dose equivalent to prednisone 30 mg daily orally for 2 weeks or intravenously for 1 week. A loss of response to corticosteroids refers to two failed attempts to taper corticosteroids to below a dose equivalent to prednisone 10 mg daily orally. Intolerance of corticosteroids includes a history of Cushing's syndrome, osteopenia/osteoporosis, hyperglycemia, insomnia and/or infection.

10 An immunomodulator may be, for example, oral azathioprine, 6-mercaptopurine, or methotrexate. An inadequate response to an immunomodulator refers to signs and symptoms of persistently active disease despite a history of at least one 8 week regimen of oral azathioprine (≥ 1.5 mg/kg), 6-mercaptopurine (≥ 0.75 mg/kg), or methotrexate (≥ 12.5 mg/week). Intolerance of an
15 immunomodulator includes, but is not limited to, nausea/vomiting, abdominal pain, pancreatitis, LFT abnormalities, lymphopenia, TPMT genetic mutation and/or infection.

 A TNF- α antagonist is, for example, an agent that inhibits the biological activity of TNF- α , and preferably binds TNF- α , such as a monoclonal antibody,
20 e.g., REMICADE (infliximab), HUMIRA (adalimumab), CIMZIA (certolizumab pegol), SIMPONI (golimumab) or a circulating receptor fusion protein such as ENBREL (etanercept). An inadequate response to a TNF- α antagonist refers to signs and symptoms of persistently active disease despite a history of at least one 4 week induction regimen of infliximab 5 mg/kg IV, 2 doses at least 2 weeks apart;
25 one 80 mg subcutaneous dose of adalimumab, followed by one 40 mg dose at least two weeks apart; or 400 mg subcutaneously of certolizumab pegol, 2 doses at least 2 weeks apart. A loss of response to a TNF- α antagonist refers to recurrence of symptoms during maintenance dosing following prior clinical benefit. Intolerance of a TNF- α antagonist includes, but is not limited to infusion related reaction,
30 demyelination, congestive heart failure, and/or infection.

A loss of maintenance of remission, as used herein for ulcerative colitis subjects, refers to an increase in Mayo score of at least 3 points and a Modified Baron Score of at least 2.

In another aspect, the present invention provides anti- $\alpha 4\beta 7$ antibody
5 formulations which (1) can bind $\alpha 4\beta 7$ integrin *in vitro* and/or *in vivo*; and (2) can modulate an activity or function of an $\alpha 4\beta 7$ integrin, such as (a) binding function (e.g., the ability of $\alpha 4\beta 7$ integrin to bind to MAdCAM (e.g., MAdCAM-1), fibronectin and/or VCAM-1) and/or (b) leukocyte infiltration function, including recruitment and/or accumulation of leukocytes in tissues (e.g., the ability to inhibit
10 lymphocyte migration to intestinal mucosal tissue). In one embodiment, an antibody in the formulation can bind an $\alpha 4\beta 7$ integrin, and can inhibit binding of the $\alpha 4\beta 7$ integrin to one or more of its ligands (e.g., MAdCAM (e.g., MAdCAM-1), VCAM-1, fibronectin), thereby inhibiting leukocyte infiltration of tissues (including recruitment and/or accumulation of leukocytes in tissues). In another embodiment,
15 an antibody in the formulation can bind an $\alpha 4\beta 7$ integrin, and can selectively inhibit binding of the $\alpha 4\beta 7$ integrin to one or more of its ligands (e.g., MAdCAM (e.g., MAdCAM-1), VCAM-1, fibronectin), thereby inhibiting leukocyte infiltration of tissues (including recruitment and/or accumulation of leukocytes in tissues). Such anti- $\alpha 4\beta 7$ antibody formulations can inhibit cellular adhesion of cells bearing an
20 $\alpha 4\beta 7$ integrin to vascular endothelial cells in mucosal tissues, including gut-associated tissues, lymphoid organs or leukocytes (especially lymphocytes such as T or B cells) *in vitro* and/or *in vivo*. In yet another embodiment, the anti- $\alpha 4\beta 7$ antibody formulation of the present invention can inhibit the interaction of $\alpha 4\beta 7$ with MAdCAM (e.g., MAdCAM-1) and/or fibronectin. In still yet another
25 embodiment, the anti- $\alpha 4\beta 7$ antibody formulation of the present invention can inhibit the interaction of $\alpha 4\beta 7$ with MAdCAM (e.g., MAdCAM-1) and/or fibronectin selectively, e.g., without inhibiting the interaction of $\alpha 4\beta 7$ with VCAM.

The anti- $\alpha 4\beta 7$ antibody formulations of the present invention can be used to modulate (e.g., inhibit (reduce or prevent)) binding function and/or leukocyte (e.g.,
30 lymphocyte, monocyte) infiltration function of $\alpha 4\beta 7$ integrin. For example, humanized immunoglobulins which inhibit the binding of $\alpha 4\beta 7$ integrin to a ligand

(i.e., one or more ligands) can be administered according to the method in the treatment of diseases associated with leukocyte (e.g., lymphocyte, monocyte) infiltration of tissues (including recruitment and/or accumulation of leukocytes in tissues), particularly of tissues which express the molecule MAdCAM (e.g.,
5 MAdCAM-1).

An effective amount of an anti- $\alpha 4\beta 7$ antibody formulation of the present invention (i.e., one or more) is administered to an individual (e.g., a mammal, such as a human or other primate) in order to treat such a disease. For example, inflammatory diseases, including diseases which are associated with leukocyte
10 infiltration of the gastrointestinal tract (including gut-associated endothelium), other mucosal tissues, or tissues expressing the molecule MAdCAM (e.g., MAdCAM-1) (e.g., gut-associated tissues, such as venules of the lamina propria of the small and large intestine; and mammary gland (e.g., lactating mammary gland)), can be treated according to the present method. Similarly, an individual having a disease
15 associated with leukocyte infiltration of tissues as a result of binding of leukocytes to cells (e.g., endothelial cells) expressing MAdCAM (e.g., MAdCAM-1) can be treated according to the present invention.

In one embodiment, diseases which can be treated accordingly include inflammatory bowel disease (IBD), such as ulcerative colitis, Crohn's disease, ileitis,
20 Celiac disease, nontropical Sprue, enteropathy associated with seronegative arthropathies, microscopic or collagenous colitis, eosinophilic gastroenteritis, or pouchitis resulting after proctocolectomy, and ileoanal anastomosis. In some embodiments, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. The ulcerative colitis may be moderate to severely active ulcerative colitis.
25 Treatment may result in mucosal healing in patients suffering from moderate to severely active ulcerative colitis. Treatment may also result in a reduction, elimination, or reduction and elimination of corticosteroid use by the patient.

Pancreatitis and insulin-dependent diabetes mellitus are other diseases which can be treated using the formulations of the invention. It has been reported that
30 MAdCAM (e.g., MAdCAM-1) is expressed by some vessels in the exocrine pancreas from NOD (nonobese diabetic) mice, as well as from BALB/c and SJL

mice. Expression of MAdCAM (e.g., MAdCAM-1) was reportedly induced on endothelium in inflamed islets of the pancreas of the NOD mouse, and MAdCAM (e.g., MAdCAM-1) was the predominant addressin expressed by NOD islet endothelium at early stages of insulinitis (Hanninen, A., *et al.*, *J. Clin. Invest.*, 92: 2509-2515 (1993)). Treatment of NOD mice with either anti-MAdCAM or anti- β 7 antibodies prevented the development of diabetes (Yang *et al.*, *Diabetes*, 46:1542-1547 (1997)). Further, accumulation of lymphocytes expressing α 4 β 7 within islets was observed, and MAdCAM-1 was implicated in the binding of lymphoma cells via α 4 β 7 to vessels from inflamed islets (Hanninen, A., *et al.*, *J. Clin. Invest.*, 92: 2509-2515 (1993)) or to the gastrointestinal tract in mantle cell lymphoma (Geissmann *et al.*, *Am. J. Pathol.*, 153:1701-1705 (1998)).

Examples of inflammatory diseases associated with mucosal tissues which can be treated using a formulation of the invention include cholecystitis, cholangitis (Adams and Eksteen *Nature Reviews* 6:244-251 (2006) Grant *et al.*, *Hepatology* 33:1065-1072 (2001)), e.g., primary sclerosing cholangitis, Behcet's disease, e.g., of the intestine, or pericholangitis (bile duct and surrounding tissue of the liver), and graft versus host disease (e.g., in the gastrointestinal tract (e.g., after a bone marrow transplant) (Petrovic *et al.* *Blood* 103:1542-1547 (2004)). As seen in Crohn's disease, inflammation often extends beyond the mucosal surface, accordingly chronic inflammatory diseases, such as sarcoidosis, chronic gastritis, e.g., autoimmune gastritis (Katakai *et al.*, *Int. Immunol.*, 14:167-175 (2002)) and other idiopathic conditions can be amenable to treatment.

The invention also relates to a method of inhibiting leukocyte infiltration of mucosal tissue. The invention also relates to a method for treating cancer (e.g., an α 4 β 7 positive tumor, such as a lymphoma). Other examples of inflammatory diseases associated with mucosal tissues which can be treated using a formulation of the invention include mastitis (mammary gland) and irritable bowel syndrome.

Diseases or pathogens whose etiologies exploit the interaction of MAdCAM (e.g., MAdCAM-1) with α 4 β 7 can be treated with an anti- α 4 β 7 antibody in a formulation described herein. Examples of such diseases include immunodeficiency

disorders, such as caused by human immunodeficiency virus (See, e.g., WO2008140602).

A formulation of the invention is administered in an effective amount which inhibits binding of $\alpha 4\beta 7$ integrin to a ligand thereof. For therapy, an effective
5 amount will be sufficient to achieve the desired therapeutic (including prophylactic) effect (such as an amount sufficient to reduce or prevent $\alpha 4\beta 7$ integrin-mediated binding and/or signaling, thereby inhibiting leukocyte adhesion and infiltration and/or associated cellular responses). An effective amount of an anti- $\alpha 4\beta 7$ antibody, e.g., an effective titer sufficient to maintain saturation, e.g., neutralization, of $\alpha 4\beta 7$
10 integrin, can induce clinical response or remission in inflammatory bowel disease. An effective amount of an anti- $\alpha 4\beta 7$ antibody can lead to mucosal healing in ulcerative colitis or Crohn's disease. A formulation of the invention can be administered in a unit dose or multiple doses. The dosage can be determined by methods known in the art and can be dependent, for example, upon the individual's
15 age, sensitivity, tolerance and overall well-being. Examples of modes of administration include topical routes such as nasal or inhalational or transdermal administration, enteral routes, such as through a feeding tube or suppository, and parenteral routes, such as intravenous, intramuscular, subcutaneous, intraarterial, intraperitoneal, or intravitreal administration. Suitable dosages for antibodies can be
20 from about 0.1 mg/kg body weight to about 10.0 mg/kg body weight per treatment, for example about 2 mg/kg to about 7 mg/kg, about 3 mg/kg to about 6 mg/kg, or about 3.5 to about 5 mg/kg. In particular embodiments, the dose administered is about 0.3 mg/kg, about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about
25 9 mg/kg, or about 10 mg/kg. The total dose may be about 22 mg, about 50 mg, about 72 mg, about 125 mg, about 165 mg, or about 432 mg. The total dose may be at least 77 mg, at least 125 mg or at least 356 mg. In one embodiment, the total dose is 165 mg. In another embodiment, the total dose is 108 mg. In another embodiment, the total dose is 216 mg.

30 Modeling and simulations (BERKELEY MADONNA™ software, University of California) using pharmacokinetic (PK) data from studies of

availability of anti- $\alpha 4\beta 7$ antibody over time after administration can assess potential dosing regimens for subcutaneous or intramuscular administration. PK data can be evaluated for induction and for maintenance regimens. Another modeling approach is population pharmacokinetic/pharmacodynamic analysis (NONMEM® nonlinear
5 mixed effects modeling tool, ICON plc, Dublin, Ireland). Both exposure levels and trough levels can be analyzed.

Typically, after target, e.g., $\alpha 4\beta 7$ integrin, saturation is reached, the antibody concentration in the blood has a linear relationship to the dose administered. An anti- $\alpha 4\beta 7$ antibody administered by the subcutaneous or intramuscular route has
10 about 60% to about 90% of the bioavailability of an anti- $\alpha 4\beta 7$ antibody administered by an intravenous route. In an example of this relationship, if an IV dose is assumed to have a 100% bioavailability and a subcutaneous dose is found to have a 69.5% bioavailability, then a 300 mg intravenous dose can be matched with a 432 mg dose by subcutaneous administration. Accordingly, a 150 mg intravenous
15 dose can be matched by a 216 mg subcutaneous dose at 69.5% relative bioavailability. Similarly, if a subcutaneous dose is found to have a 75% availability and an intramuscular dose is found to have an 80% bioavailability, then to match a 300 mg intravenous dose, the subcutaneous dose can be 400 mg and the intramuscular dose can be 375 mg. Tables 40-43 in the examples illustrate these
20 relationships and provide useful doses and dosing regimens of an anti- $\alpha 4\beta 7$ antibody.

In some aspects, the dosing regimen has two phases, an induction phase and a maintenance phase. In the induction phase, the antibody or antigen-binding fragment thereof is administered in a way that quickly provides an effective amount
25 of the antibody or antigen binding fragment thereof suitable for certain purposes, such as inducing immune tolerance to the antibody or antigen-binding fragment thereof or for inducing a clinical response and ameliorating inflammatory bowel disease symptoms. A patient can be administered an induction phase treatment when first being treated by an anti- $\alpha 4\beta 7$ antibody, when being treated after a long
30 absence from therapy, e.g., more than three months, more than four months, more than six months, more than nine months, more than one year, more than eighteen

months or more than two years since anti- $\alpha 4\beta 7$ antibody therapy or during maintenance phase of anti- $\alpha 4\beta 7$ antibody therapy if there has been a return of inflammatory bowel disease symptoms, e.g., a relapse from remission of disease. In some embodiments, the induction phase regimen results in a higher mean trough
5 serum concentration, e.g., the concentration just before the next dose, than the mean steady state trough serum concentration maintained during the maintenance regimen.

In the maintenance phase, the antibody or antigen-binding fragment thereof is administered in a way that continues the response achieved by induction therapy with a stable level of antibody or antigen-binding fragment thereof. A maintenance
10 regimen can prevent return of symptoms or relapse of inflammatory bowel disease. A maintenance regimen can provide convenience to the patient, e.g., be a simple dosing regimen or require infrequent trips for treatment. In some embodiments, the maintenance regimen can include administration of the anti- $\alpha 4\beta 7$ antibody or antigen-binding fragment thereof, e.g., in a formulation described herein, by a
15 strategy selected from the group consisting of low dose, infrequent administration, self-administration and a combination any of the foregoing.

In one embodiment, e.g., during an induction phase of therapy, the dosing regimen provides an effective amount of an anti- $\alpha 4\beta 7$ antibody or antigen-binding fragment in a formulation described herein for inducing remission of an
20 inflammatory bowel disease in a human patient. In some embodiments, the effective amount of the anti- $\alpha 4\beta 7$ antibody is sufficient to achieve about 5 $\mu\text{g/ml}$ to about 60 $\mu\text{g/ml}$, about 15 $\mu\text{g/ml}$ to about 45 $\mu\text{g/ml}$, about 20 $\mu\text{g/ml}$ to about 30 $\mu\text{g/ml}$, or about 25 $\mu\text{g/ml}$ to about 35 $\mu\text{g/ml}$ mean trough serum concentration of the anti- $\alpha 4\beta 7$ antibody by the end of the induction phase. The duration of induction phase can be
25 about four weeks, about five weeks, about six weeks, about seven weeks, or about eight weeks of treatment. In some embodiments, the induction regimen can utilize a strategy selected from the group consisting of high dose, frequent administration, and a combination of high dose and frequent administration of the anti- $\alpha 4\beta 7$ antibody or antigen-binding fragment thereof, e.g., in a formulation described
30 herein. Induction dosing can be once, or a plurality of more than one dose, e.g., at least two doses. During induction phase, a dose can be administered once per day,

every other day, twice per week, once per week, once every ten days, once every two weeks or once every three weeks. In some embodiments, the induction doses are administered within the first two weeks of therapy with the anti- $\alpha 4\beta 7$ antibody. In one embodiment, induction dosing can be once at initiation of treatment (day 0) and once at about two weeks after initiation of treatment. In another embodiment, the induction phase duration is six weeks. In another embodiment, the induction phase duration is six weeks and a plurality of induction doses are administered during the first two weeks.

In some embodiments, e.g., when initiating treatment of a patient with severe inflammatory bowel disease (e.g., in patients who have failed anti-TNF α therapy), the induction phase needs to have a longer duration than for patients with mild or moderate disease. In some embodiments, the induction phase for a patient with a severe disease can have a duration of at least 6 weeks, at least 8 weeks, at least 10 weeks, at least 12 weeks or at least 14 weeks. In one embodiment, an induction dosing regimen for a patient with a severe disease can include a dose at week 0 (initiation of treatment), a dose at week 2 and a dose at week 6. In another embodiment, an induction dosing regimen for a patient with a severe disease can comprise a dose at week 0 (initiation of treatment), a dose at week 2, a dose at week 6 and a dose at week 10.

In one embodiment, e.g., during a maintenance phase of therapy, the dosing regimen maintains a mean steady state trough serum concentration, e.g., the plateau concentration just before the next dose, of about 5 to about 25 $\mu\text{g/mL}$, about 7 to about 20 $\mu\text{g/mL}$, about 5 to about 10 $\mu\text{g/mL}$, about 10 to about 20 $\mu\text{g/mL}$, about 15 to about 25 $\mu\text{g/mL}$ or about 9 to about 13 $\mu\text{g/mL}$ of anti- $\alpha 4\beta 7$ antibody. In another embodiment, the dosing regimen e.g., during a maintenance phase of therapy, maintains a mean steady state trough serum concentration of about 20 to about 30 $\mu\text{g/mL}$, about 20 to about 55 $\mu\text{g/mL}$, about 30 to about 45 $\mu\text{g/mL}$, about 45 to about 55 $\mu\text{g/mL}$ or about 35 to about 40 $\mu\text{g/mL}$ of anti- $\alpha 4\beta 7$ antibody. In another embodiment, the dosing regimen e.g., during a maintenance phase of therapy, maintains a long term mean serum concentration, e.g., exposure (e.g., area under the curve - concentration-time) of about 15 to about 40 $\mu\text{g/mL}$, about 10 to about 50

µg/mL, about 18 to about 26 µg/mL, or about 22 to about 33 µg/mL of anti-α4β7 antibody. In yet another embodiment, the dosing regimen e.g., during a maintenance therapy, maintains a long term mean serum concentration, e.g., exposure (e.g., area under the curve - concentration-time) of about 35 to about 90 µg/mL, about 45 to
5 about 75 µg/mL, about 52 to about 60 µg/mL or about 50 to about 65 µg/mL of anti-α4β7 antibody.

The final dosage form can comprise the entire dose in about 0.5 ml, in about 1 ml, in about 1.5 ml in about 2 ml, in about 2.5 ml, in about 3 ml of the antibody formulation.

10 The final dosage form for intravenous administration may be at a concentration of between about 1.0 mg/ml to about 1.4 mg/ml, about 1.0 mg/ml to about 1.3 mg/ml, about 1.0 mg/ml to about 1.2 mg/ml, about 1.0 to about 1.1 mg/ml, about 1.1 mg/ml to about 1.4 mg/ml, about 1.1 mg/ml to about 1.3 mg/ml, about 1.1 mg/ml to about 1.2 mg/ml, about 1.2 mg/ml to about 1.4 mg/ml, about 1.2 mg/ml to
15 about 1.3 mg/ml, or about 1.3 mg/ml to about 1.4 mg/ml. The final dosage form may be at a concentration of about 0.6 mg/ml, 0.8 mg/ml, 1.0 mg/ml, 1.1 mg/ml, about 1.2 mg/ml, about 1.3 mg/ml, about 1.4 mg/ml, about 1.5 mg/ml, about 1.6 mg/ml, about 1.8 mg/ml or about 2.0 mg/ml.

The dose can be administered once per week, once every 2 weeks, once
20 every 3 weeks, once every 4 weeks, once every 6 weeks, once every 8 weeks or once every 10 weeks. A higher or more frequent dose, e.g., every other day, once per week, once every 2 weeks, once every 3 weeks or once every 4 weeks can be useful for inducing remission of active disease or for treating a new patient, e.g., for inducing tolerance to the anti- α4β7 antibody. A dose once every 2 weeks, once
25 every 3 weeks, once every 4 weeks, once every 5 weeks, once every 6 weeks, once every 8 weeks or once every 10 weeks, can be useful for preventative therapy, e.g., to maintain remission of a patient with chronic disease. In one aspect, the treatment regimen is treatment at day 0, about week 2, about week 6 and every 1 or 2 weeks thereafter. In another aspect, the induction treatment regimen is treatment every
30 other day for a total of 6 treatments.

The dosing regimen can be optimized to induce a clinical response and clinical remission in the inflammatory bowel disease of the patient. In some embodiments, the dosing regimen does not alter the ratio of CD4 to CD8 in cerebrospinal fluid of patients receiving treatment.

5 In some aspects, a durable clinical remission, for example, a clinical remission which is sustained through at least two, at least three, at least four visits with a caretaking physician within a six month or one year period after beginning treatment, may be achieved with an optimized dosing regimen.

10 In some aspects, a durable clinical response, for example, a clinical response which is sustained for at least 6 months, at least 9 months, at least a year, after the start of treatment, may be achieved with an optimized dosing regimen.

 The formulation may be administered subcutaneously in single or multiple injections. For example, the volume of a single injection may range from about 0.5 ml to about 3 ml. In an embodiment, the volume of a single injection may be about 15 0.6 ml to about 1.1 ml or about 1 ml to about 3 ml. In one aspect, the volume of a single injection is about 1 ml. The gauge of the needle used to administer the formulation subcutaneously may be about 25, about 26, about 27, about 28, about 29 or about 30G.

 The formulation may be administered intramuscularly in single or multiple 20 injections. For example, the volume of a single injection may range from about 0.5 ml to about 5 ml. In an embodiment, the volume of a single injection may be about 2 ml to about 5 ml, about 0.6 ml to about 1.1 ml or about 1 ml to about 3 ml. In one aspect, the volume of a single injection is about 1 ml, about 2 ml, about 3 ml, about 4 ml, or about 5 ml. The needle used to administer the formulation intramuscularly 25 may be about 5/8", about 7/8", about 1", about 1.25", about 1.5", about 2", or about 3". The gauge of the needle may be between 20-22G for intramuscular administration.

 In one aspect, the invention relates to a method for treating a human patient suffering from inflammatory bowel disease, wherein the method comprises the step 30 of administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding

specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to the following dosing regimen: (a) initial doses of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection
5 every other day for six doses; (b) followed by a seventh and subsequent doses of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection every two weeks or every four weeks as needed; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized
10 immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below: light chain: CDR1
15 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In one aspect, the invention relates to a method for treating a human patient suffering from inflammatory bowel disease, wherein the method comprises the step of administering to a patient suffering from inflammatory bowel disease, a
20 humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient
25 according to the following dosing regimen: (a) an initial intravenous dose of 300 mg of the humanized immunoglobulin or antigen-binding fragment thereof as an intravenous infusion; (b) followed by a second intravenous subsequent dose of 300 mg of the humanized immunoglobulin or antigen-binding fragment thereof as an intravenous infusion at about two weeks after the initial dose; (c) followed
30 beginning at week six by a third and subsequent doses of 165 mg of the humanized immunoglobulin or antigen-binding fragment thereof as a subcutaneous injection

every week, every two weeks or every three weeks as needed; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below: light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In another aspect, the invention relates to a dosing regimen for the therapeutic treatment of inflammatory bowel disease, wherein the dosing regimen comprises the step of: administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to a subcutaneous or intramuscular dosing regimen which maintains a mean steady state serum trough concentration of about 9 to about 13 $\mu\text{g/mL}$ of the antibody or antigen-binding fragment thereof; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below: light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1 SEQ ID NO:12, CDR2 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In another aspect, the invention relates to a dosing regimen for the therapeutic treatment of inflammatory bowel disease, wherein the dosing regimen comprises the step of: administering to a patient suffering from inflammatory bowel disease, a humanized immunoglobulin or antigen-binding fragment thereof having
5 binding specificity for human $\alpha 4\beta 7$ integrin, wherein the humanized immunoglobulin or antigen-binding fragment comprises an antigen-binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized immunoglobulin or antigen-binding fragment thereof is administered to the patient according to a subcutaneous or intramuscular dosing regimen which
10 maintains a steady state serum trough concentrations of about 35 to about 40 $\mu\text{g/mL}$ of the antibody or antigen-binding fragment thereof; wherein the dosing regimen induces a clinical response and clinical remission in the inflammatory bowel disease of the patient; and further wherein the humanized immunoglobulin or antigen-binding fragment has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-
15 binding region comprises three complementarity determining regions (CDR1, CDR2, and CDR3) of a light chain variable region and three complementarity determining regions (CDR1, CDR2, and CDR3) of a heavy chain variable region of the amino acid sequence set forth below: light chain: CDR1 SEQ ID NO:9, CDR2 SEQ ID NO:10, CDR3 SEQ ID NO:11; heavy chain: CDR1 SEQ ID NO:12, CDR2
20 SEQ ID NO:13, CDR3 SEQ ID NO:14.

In some embodiments, the method of treatment, dose or dosing regimen reduces the likelihood that a patient will develop a HAHA response to the anti- $\alpha 4\beta 7$ antibody. The development of HAHA, e.g., as measured by antibodies reactive to the anti- $\alpha 4\beta 7$ antibody, can increase the clearance of the anti- $\alpha 4\beta 7$ antibody, e.g.,
25 reduce the serum concentration of the anti- $\alpha 4\beta 7$ antibody, e.g., lowering the number of anti- $\alpha 4\beta 7$ antibody bound to $\alpha 4\beta 7$ integrin, thus making the treatment less effective. In some embodiments, to prevent HAHA, the patient can be treated with an induction regimen followed by a maintenance regimen. In some embodiments, there is no break between the induction regimen and the maintenance regimen. In
30 some embodiments, the induction regimen comprises administering a plurality of doses of anti- $\alpha 4\beta 7$ antibody to the patient. To prevent HAHA, the patient can be

treated with a high initial dose, e.g., at least 1.5 mg/kg, at least 2 mg/kg, at least 2.5 mg/kg, at least 3 mg/kg, at least 5 mg/kg, at least 8 mg/kg, at least 10 mg/kg or about 2 to about 6 mg/kg, or frequent initial administrations, e.g., about once per week, about once every two weeks or about once every three weeks, of the standard
5 dose when beginning therapy with an anti- $\alpha 4\beta 7$ antibody. In some embodiments, the method of treatment maintains at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90% or at least 95% of patients as HAHA-negative. In other embodiments, the method of treatment maintains patients as HAHA-negative for at least 6 weeks, at least 10 weeks at least 15 weeks, at least six
10 months, at least 1 year, at least 2 years, or for the duration of therapy. In some embodiments, the patients, or at least 30%, at least 40%, at least 50% or at least 60% of patients who develop HAHA maintain a low titer, e.g., ≤ 125 , of anti- $\alpha 4\beta 7$ antibody. In an embodiment, the method of treatment maintains at least 70% of patients as HAHA-negative for at least 12 weeks after beginning therapy with an
15 anti- $\alpha 4\beta 7$ antibody.

The formulation may be administered to an individual (e.g., a human) alone or in conjunction with another agent. A formulation of the invention can be administered before, along with or subsequent to administration of the additional agent. In one embodiment, more than one formulation which inhibits the binding of
20 $\alpha 4\beta 7$ integrin to its ligands is administered. In such an embodiment, an agent, e.g., a monoclonal antibody, such as an anti-MAdCAM or an anti-VCAM-1 monoclonal antibody can be administered. In another embodiment, the additional agent inhibits the binding of leukocytes to an endothelial ligand in a pathway different from the $\alpha 4\beta 7$ pathway. Such an agent can inhibit the binding, e.g. of chemokine (C-C motif)
25 receptor 9 (CCR9)-expressing lymphocytes to thymus expressed chemokine (TECK or CCL25) or an agent which prevents the binding of LFA-1 to intercellular adhesion molecule (ICAM). For example, an anti-TECK or anti-CCR9 antibody or a small molecule CCR9 inhibitor, such as inhibitors disclosed in PCT publication WO03/099773 or WO04/046092, or anti-ICAM-1 antibody or an oligonucleotide
30 which prevents expression of ICAM, is administered in addition to a formulation of the present invention. In yet another embodiment, an additional active ingredient

(e.g., an anti-inflammatory compound, such as sulfasalazine, azathioprine, 6-mercaptapurine, 5-aminosalicylic acid containing anti-inflammatories, another non-steroidal anti-inflammatory compound, a steroidal anti-inflammatory compound, or antibiotics commonly administered for control of IBD (e.g. ciprofloxacin, metronidazole), or another biologic agent (e.g. TNF alpha antagonists) can be administered in conjunction with a formulation of the present invention.

In an embodiment, the dose of the co-administered medication can be decreased over time during the period of treatment by the formulation comprising the anti- $\alpha 4\beta 7$ antibody. For example, a patient being treated with a steroid (e.g. prednisone, prednisolone) at the beginning, or prior to, treating with the anti- $\alpha 4\beta 7$ antibody formulation would undergo a regimen of decreasing doses of steroid beginning as early as 6 weeks of treatment with the anti- $\alpha 4\beta 7$ antibody formulation. The steroid dose will be reduced by about 25% within 4-8 weeks of initiating tapering, by 50 % at about 8-12 weeks and 75% at about 12-16 weeks of tapering during treatment with the anti- $\alpha 4\beta 7$ antibody formulation. In one aspect, by about 16-24 weeks of treatment with the anti- $\alpha 4\beta 7$ antibody formulation, the steroid dose can be eliminated. In another example, a patient being treated with an anti-inflammatory compound, such as 6-mercaptapurine at the beginning, or prior to, treating with the anti- $\alpha 4\beta 7$ antibody formulation would undergo a regimen of decreasing doses of anti-inflammatory compound similar to the tapering regimen for steroid dosing as noted above.

In one embodiment, the method comprises subcutaneously administering or intramuscularly administering an effective amount of a formulation of the invention to a patient. In another embodiment, the formulation can be prepared for self-administration.

If the formulation is in a solid, e.g., dry state, the process of administration can comprise a step of converting the formulation to a liquid state. In one aspect, a dry formulation can be reconstituted, e.g., by a liquid as described above, for use in injection, e.g. intravenous, intramuscular or subcutaneous injection. In another aspect, a solid or dry formulation can be administered topically, e.g., in a patch, cream, aerosol or suppository.

The invention also relates to a method for treating a disease associated with leukocyte infiltration of tissues expressing the molecule MAdCAM (e.g., MAdCAM-1). The method comprises administering to a patient in need thereof an effective amount of an anti- $\alpha 4\beta 7$ antibody formulation of the invention. In an embodiment, the disease is graft versus host disease. In some embodiments, the disease is a disease associated with leukocyte infiltration of tissues as a result of binding of leukocytes expressing $\alpha 4\beta 7$ integrin to gut-associated endothelium expressing the molecule MAdCAM (e.g., MAdCAM-1). In other embodiments, the disease is gastritis (e.g., eosinophilic gastritis or autoimmune gastritis), pancreatitis, or insulin-dependent diabetes mellitus. In yet other embodiments, the disease is cholecystitis, cholangitis, or pericholangitis.

The invention also relates to a method for treating inflammatory bowel disease in a patient. In one embodiment, the method comprises subcutaneously administering to the patient an effective amount of an anti- $\alpha 4\beta 7$ antibody formulation of the invention. In some embodiments, the inflammatory bowel disease is ulcerative colitis or Crohn's disease. In other embodiments, the inflammatory bowel disease is Celiac disease, enteropathy associated with seronegative arthropathies, microscopic or collagenous colitis, gastroenteritis (e.g., eosinophilic gastroenteritis), or pouchitis.

In some embodiments, treatment with an anti- $\alpha 4\beta 7$ antibody does not alter the ratio of CD4:CD8 lymphocytes. CD4:CD8 ratios can be measured in blood, lymph node aspirate, and cerebro-spinal fluid (CSF). The CSF CD4+:CD8+ lymphocyte ratios in healthy individuals are typically greater than or equal to about 1. (Svenningsson et al., J. Neuroimmunol. 1995;63:39-46; Svenningsson et al., Ann Neurol. 1993; 34:155-161). An immunomodulator can alter the CD4:CD8 ratio to less than 1.

Articles of Manufacture

In another aspect, the invention is an article of manufacture which contains the pharmaceutical formulation of the present invention and provides instructions for its use. The article of manufacture comprises a container. Suitable containers

include, for example, bottles, vials (e.g., dual chamber vials, a vial of liquid formulation with or without a needle, a vial of solid formulation with or without a vial of reconstitution liquid with or without a needle), syringes (such as dual chamber syringes, preloaded syringes, an auto-injector), cartridges, and test tubes.

5 The container may be formed from a variety of materials such as glass, metal or plastic. The container holds the formulation and a label on, or associated with, the container may indicate directions for use. In another embodiment, the formulation can be prepared for self-administration and/or contain instructions for self-administration. In one aspect, the container holding the formulation may be a
10 single-use vial. In another aspect, the container holding the formulation may be a multi-use vial, which allows for repeat administration (e.g., from 2-6 administrations) of the formulation, e.g., using more than one portion of a reconstituted formulation. The article of manufacture may further include other materials desirable from a commercial and user standpoint, including other buffers,
15 diluents, filters, needles, syringes and package inserts with instructions for use as noted in the previous section.

In one embodiment, the article of manufacture is a syringe with a needle. The gauge of the needle may be 25 G, 26 G, 27 G, 29 G, 30 G. A thin wall needle, e.g., 19 G or 23 G, or greater, can facilitate injection of a high viscosity formulation.
20 In one aspect, the needle gauge is 27 G or greater. Needle length can be suitable for subcutaneous administration, and can be about 1/2 inch, about 5/8 inch or 1 inch long. In some embodiments, the syringe is a pre-filled syringe.

Pre-filled Syringe Product Development

25 In some aspects, there are several product attributes that are desired for a protein product (e.g., an anti- $\alpha\beta 7$ antibody) in a pre-filled syringe (PFS) (e.g., for use in administration of a formulation for subcutaneous or intramuscular delivery). It is helpful to balance some of the attributes to mitigate competing effects. For example, when a low injection volume is desired, a high protein concentration for
30 the formulation may be preferred. However, in the case of a high protein concentration, there can be higher rates of impurity formation (e.g., aggregated

impurities that leach into formulation from syringe) and higher manual forces needed to operate the syringe. A small needle size used for patient comfort at the injection site, may require high forces to operate the syringe. An understanding of how product stability and performance is affected by both formulation and syringe parameters such as protein concentration, pH, and needle inner diameter aids in the development of a protein product (e.g., an anti- $\alpha 4\beta 7$ antibody in a pre-filled syringe).

In one aspect, a method of developing a protein product (e.g., an anti- $\alpha 4\beta 7$ antibody) for use in a pre-filled syringe comprises varying syringe parameters and formulation parameters together, e.g., in a coordinate fashion or simultaneously. This can lead to a better understanding of the range of protein stability and product performance that can be expected from a protein product in a pre-filled syringe than if each aspect is varied separately or in series.

The development of a pre-filled syringe product (e.g., an anti- $\alpha 4\beta 7$ antibody) relates to understanding that at some point, there is a liquid formulation in contact with several components of the pre-filled syringe (Figure 15). For example, the formulation can be in contact with a syringe barrel, which can be constructed of glass (e.g., type I borosilicate glass) or plastic (e.g., cyclic olefin polymer (COP), cyclic olefin copolymer (COC), polypropylene or polytetrafluoroethylene). The formulation can be in contact with the syringe, plunger and/or tip cap, which can be elastomeric (e.g., of the same or different materials (e.g., plastic, such as polyethylene, polystyrene or polypropylene or elastic, such as rubber (natural, synthetic, butyl) or silicone)). The formulation may be in contact with a lubricant that is added to an inner surface of the barrel for ease of plunger movement. The lubricant can be, for example, silicone oil, mineral oil or glycerin. In the embodiment of a staked needle syringe, there can be a metal alloy needle (e.g., stainless steel needle and adhesive used to glue the needle in place). A consideration for a protein product in a pre-filled syringe is that the liquid protein solution is in direct contact with one or more of these syringe components throughout the shelf life of the product. Both the formulation and syringe components can have an impact on the stability of the product.

Formulation parameters that can affect pre-filled syringe product stability include protein concentration, pH, buffer type, buffer concentration, ionic strength, stabilizer type, and stabilizer concentration. Examples of stabilizers for protein formulations include, for example, ionic salts, polysaccharides, amino acids, antioxidants, chelators, and surfactants as described in earlier sections.

Syringe components that can affect pre-filled syringe product stability include, for example, lubricant, composition of plunger and tip cap, and impurities. The amount of lubricant (e.g., silicone oil on the syringe barrel) may affect product stability. The composition of the plunger and tip cap, which can affect oxygen permeability of these components and introduce leachables from these components into the protein product (e.g., the anti- $\alpha 4\beta 7$ antibody formulation) may also affect product stability. Another syringe parameter that can affect product stability includes the type and/or amount of impurities (e.g., heavy metal (e.g., tungsten)) that can leach into the product formulation (e.g., from in the barrel (e.g., glass barrel) and/or needle (e.g., stainless steel needle)). (See also Ludwig et al. *J.Pharm. Sci.* 99:1721-1733 (2010); Nashed-Samuel et al., *American Pharmaceutical Review* Jan/Feb:74-80 (2011); Badkar et al. *AAPS PharmSciTech* 12:564-572)

A pre-filled syringe can be injected manually or used with an auto-injector device. Functional testing of the pre-filled syringe includes measuring the break-loose force, the force required to begin movement of the plunger, and the gliding force, the force needed to inject the contents of the syringe at a constant rate. The mechanical performance of the pre-filled syringe can be dependent on several formulation and syringe parameters such as the viscosity of the formulation and the amount of lubricant (e.g., silicone oil) in the syringe.

Several attributes of protein products in pre-filled syringes and formulation or syringe factors that can impact those product attributes are shown in Table 1. Many product attributes can be a complex function of several formulation and syringe parameters. For example, syringe gliding force is a function of formulation viscosity, although viscosity can be dependent on several formulation factors, such as protein concentration, stabilizer concentrations, and pH.

Table 1: Product Attributes for Protein Products in Pre-filled Syringes and the Potential Formulation and Syringe Parameters that may Impact These Attributes

Product Attribute	Protein Formulation Parameters that may Impact Product Attribute	Syringe Parameters that may Impact Product Attribute
Osmolality	Stabilizer concentrations, pH, protein concentration	None
Viscosity	Stabilizer concentrations, pH, protein concentration	None
Syringe Break Loose and Gliding Force	Viscosity, protein concentration, surfactant concentration	Injection speed, needle length, needle ID, syringe barrel ID, silicone oil amount, plunger formulation and shape
Rate of Protein Deamidation	Stabilizer concentrations, pH, protein concentration	None
Rate of Protein Oxidation	Stabilizer/antioxidant concentrations, pH, protein concentration, surfactant concentration, dissolved oxygen	Plunger and tip cap formulation (oxygen permeability), heavy metal impurity levels, size of air bubble
Rate of Soluble Aggregate Formation	Stabilizer concentrations, pH, protein concentration, surfactant concentration, dissolved oxygen in solution	Silicone oil amount, heavy metal impurity levels, size of air bubble
Rate of Sub-visible and Visible Proteinaceous Particulate Formation	Stabilizer concentrations, pH, protein concentration, surfactant concentration	Silicone oil amount, Heavy metal impurity levels, , size of air bubble, syringe internal surface area

In one aspect, a surfactant, such as polysorbate 20 or polysorbate 80 can be
 5 added to protein formulations in pre-filled syringes (e.g., to prevent protein
 molecules from adsorbing and denaturing at the liquid/air and/or liquid/lubricant
 (e.g., silicone oil) interfaces). Surface adsorption and denaturation of protein
 molecules can be one mechanism for the nucleation of sub-visible and visible
 proteinaceous particles. Addition of a surfactant to a pre-filled syringe, therefore,
 10 can reduce the formation of sub-visible and visible particles in pre-filled syringe
 products. In one embodiment, a small amount of surfactant can emulsify lubricant
 (e.g., silicone oil droplets in the solution and thereby reduce the formation of
 subvisible and visible lubricant (e.g., silicone oil droplets)) (Ludwig et al., *supra*).

In another embodiment, the amount of surfactant in a formulation is minimized, due to potential harmful effects of high amounts of surfactants on protein formulations. Peroxide impurities present in polysorbates can lead to increased protein oxidation (Wang and Wang *J. Pharm. Sci.* 91:2252-2264 (2002)). High amounts of surfactant can emulsify a significant amount of silicone oil from the walls of the syringe and lead to an increase in the functional gliding force over the shelf life. Product development studies should be designed to examine the effect of varying surfactant levels on both product stability and syringe performance.

The complex interactions between the formulation and the syringe parameters in protein/PFS systems are amenable to examination of these systems using a Quality by Design (QbD) or Design of Experiments (DOE) approach. Studies can be designed that simultaneously vary formulation and syringe parameters to gain a better understanding of these complex systems. This results in a comprehensive approach to the development of pre-filled syringe products. Table 2 shows an example of the input parameters and levels that may go into a design of experiment for a pre-filled syringe product and an example of the analytical testing to be employed. Depending on the type of experimental design that is used for QbD study, the number of experiments could vary from 9 for a screening design to 81 for a full-factorial design (all possible combinations). The higher the number of experiments, the higher the number of interactions between product parameters that can be resolved. Software designed for this analysis, for example, JMP® statistical discovery software (Cary, NC), can be helpful for QbD studies. This analysis results in a quantitative understanding of how formulation and syringe parameters interact to impact product attributes.

Table 2: Example of an Experimental Design for a Liquid Protein Product in a Pre-filled Syringe

Formulation Input Parameters	Levels	Analytical Testing
Protein Concentration	50, 100, 150 mg/mL	• Protein stability by size exclusion

pH	5.5, 6.5, 7.5	chromatography and ion exchange chromatography • Break loose and gliding force over time using force testing • Formation of sub-visible proteinaceous particles and silicone oil droplets over time using microflow imaging or /coulter counter
Surfactant Concentration	0.01, 0.08, 0.15 %	
Amount of Silicone Oil in Syringe	0.2, 0.5, 0.8 mg/syringe	

An example of a predictive model that can be obtained from the example experiment shown in Table 2 is given below, where C_n are numerical constants.

5

Soluble Aggregate Formation over Time = C₀ + C₁[Protein Concentration] + C₂[Protein Concentration]² + C₃[pH] + C₄[Surfactant Concentration] + C₅[Lubricant Amount]

10

Numerous syringe parameters can affect product stability and performance, therefore an embodiment includes characterization of how the allowable tolerances in syringe parameters affect product stability and performance. The amount of lubricant (e.g., silicone oil) on the syringe barrel may vary 50-100% from syringe to syringe. This variation in the amount may affect several product characteristics as shown in Table 1. The inner diameter of the syringe barrel can vary from syringe to syringe which affects injection forces. For staked-needle syringes, the needle inner diameter may vary from lot to lot or from manufacturer to manufacturer, which will affect injection forces. By using a QbD approach to examine how syringe parameters affect performance, predictive models can be obtained that can be used to estimate how the allowable tolerances in syringe parameters may affect product performance. Predictive models that are obtained using a QbD approach can be used to select formulation and syringe parameters that meet desired product attributes and to predict product stability and performance.

20

The pre-filled syringe may contain an addition of silicone emulsion or tungsten to the protein formulation. Exemplary amounts of silicone that may be present in the pre-filled syringe range from about 0.3 mg to about 0.8 mg. In one aspect, the amount of silicone that may be present in the pre-filled syringe is about

25

0.3 mg, about 0.4 mg, about 0.5 mg, about 0.6 mg, about 0.7 mg, or about 0.8 mg. In another aspect, the viscosity of the formulation will range from 2 to 60 cP, resulting in injection forces of 5N to 80N at a speed of 200 mm/min. In still yet another aspect, the viscosity of the formulation will range from 4 to 27 cP resulting in injection forces of 10 N to 40 N at a speed of 200 mm/min.

The invention will be more fully understood by reference to the following examples. They should not, however, be construed as limiting the scope of the invention.

10 PROTOCOL FOR MAKING FORMULATION

A solution of anti- $\alpha\beta 7$ antibody is diafiltered in a tangential flow filtration system to reach a specified concentration in citrate, histidine, arginine buffer, then pooled and mixed with a solution of polysorbate 80 in citrate, histidine, arginine
15 buffer. The solution is stored at -70 °C in either 2L or 5L bottles. The solution is then thawed and filtered twice through a 0.2 μ m filter. Approximately 1.0 mL is filled into a sterilized syringe and closed with a sterilized plunger (stopper). The formulation is stored and the final drug product is shipped in syringes at 2-8 °C.

20 **EXAMPLES**

EXAMPLE 1

FORMULATION MANUFACTURING

Factors

25

Excipient Concentrations

The formation of aggregates in the antibody formulation was tested. An SEC aggregates model was developed from experimental data that examined protein concentration, pH, and surfactant:protein molar ratio. At a pH range from 6.0 to 6.5,
30 the formation of aggregates was similar with the polysorbate 80 to protein molar

ratio range from 0.7 to 1.5. (Fig 6) Generally, at PS80:Protein ratios greater than 1.5, the aggregates formation rate increases with increasing pH. (Fig 7)

An experiment examining the formation of SEC aggregates in the presence of air was performed. Eleven different formulations of varying composition were put into borosilicate vials and capped with elastomeric stoppers with an air headspace. An identical set of formulations were created, and the air headspace was displaced with argon. These samples were placed on stability at 40°C for two weeks. All the samples with the air headspace resulted in large amounts of aggregates at the end of the experiment in comparison to the same formulation with the argon headspace.

Table 3

Sample	Protein Conc. (mg/ml)	Sucrose (%)	Histidine (mM)	Arginine (mM)	PS 80 (%)	pH	Aggregates Air Samples (%)	Aggregates Argon Samples (%)
1	60	2	25	75	0.05	6.2	0.64	0.48
2	60	4	25	75	0.05	7	0.62	0.42
3	160	4	50	75	0.14	6.2	0.92	0.73
4	160	2	50	75	0.14	7	1.16	0.74
5	60	2	50	125	0.05	7	1.28	0.33
6	60	4	50	125	0.05	6.6	0.48	0.36
7	160	4	25	125	0.14	7	1.04	0.70
8	160	2	25	125	0.14	6.2	1.06	0.75
9	160	3	25	75	0.14	6.6	1.09	0.78
10	110	3	50	125	0.10	6.2	0.65	0.47
11	110	2	25	75	0.10	6.6	0.90	0.62

Based on these experiments, SEC aggregates were hypothesized to form by oxidation or by disulfide bond formation. The addition of antioxidants and/or chelators was explored. A formulation containing 40 mM Histidine, 90 mM Arginine, and 160 mg/mL protein with a polysorbate 80 to protein molar ratio of 1.5 at pH 6.6 was made. To the formulation, 25 mM citrate, 5 mM citrate, 5 mM EDTA, 25 mM cysteine, or 5 mM cysteine was added. All 3 additional excipients reduced the formation of aggregates (Fig 8). The addition of the antioxidants and/or chelators were ranked in order of performance as citrate>EDTA>cysteine. Either

or 25 mM citrate reduced the formation of SEC aggregates as compared to the control formulation.

5 An experiment was performed to determine the effects of pH, protein concentration, citrate concentration, histidine concentration and the polysorbate 80 to protein molar ratio. The pH was varied from 6.0 to 6.3, the protein concentration was varied from 60 to 160 mg/mL, the citrate concentration was varied from 0 to 25 mM, the histidine concentration was varied from 25 to 50 mM, and the polysorbate 80 to protein molar ratio was varied from 0.7 to 1.5. Formulations were filled in 1 ml long, 27G1/2" syringes (0.55 +/- 0.2 mg silicone). All formulations contained
10 approximately 125 mM arginine.

Stability was tested at 40°C for two weeks, using CEX and SEC. The results (Fig. 9 and Table 4) show a reduction in aggregate formation with the presence of 25 mM citrate in the formulation, while increasing the protein concentration increased the rate of aggregate formation. The amount of monomer shows opposite trends to
15 the aggregate formation at 25 °C and 40 °C, while at 5 °C the amount of monomer is essentially unchanged for up to 24 months (Table 5).

Another set of formulations explored the rate of formation of SEC aggregates in the presence of 40-63 mM citrate but with no histidine at 40 °C, 25 °C, 5 °C. The rate of aggregate formation in these formulations was slightly higher than
20 formulations with histidine at 40 °C. However, at 5 °C, the rate of formation of aggregates in the formulations with citrate and no histidine were comparable to the formulations containing citrate and histidine (Table 6). Also at 5 °C, the amount of monomer is essentially unchanged for up to 24 months (Table 7).

Table 4

Formulation #	Protein Conc. (mg/mL)	pH	Histidine Conc. (mM)	Citrate Conc. (mM)	Arginine Conc. (mM)	PS80: Protein Molar Ratio	Initial Amount of Aggregates (%)	Change in Aggregates After 12 Months at 5 °C	Change in Aggregates After 24 Months at 5 °C	Change in Aggregates After 12 Months at 25 °C	Change in Aggregates After 1 Month at 40 °C
1	62	6.4	50	25	125	0.7	0.4	0.1	0.1	0.7	0.2
2	60	6.4	50	0	125	1.5	0.4	0.5	1.1	1.5	0.4
3	157	6.4	50	25	125	1.5	0.4	0.2	0.3	1.3	0.5
4	161	6.3	50	0	125	0.7	0.4	0.6	0.7	2.5	0.8
5	60	6.2	50	25	125	1.5	0.4	0.2	0.2	0.5	0.2
6	110	6.0	50	0	125	0.7	0.4	0.4	0.6	1.7	0.7
7	162	6.2	50	25	125	0.7	0.4	0.3	0.3	1.1	0.5
8	160	6.0	50	0	125	1.5	0.4	0.4	0.6	2.2	0.9
9	169	6.3	25	25	125	0.7	0.5	--	0.3	--	0.6
10	158	6.3	25	25	123	1.0	0.5	--	--	--	0.6

Table 5

Formulation #	Protein Conc. (mg/mL)	pH	Histidine Conc. (mM)	Citrate Conc. (mM)	Arginine Conc. (mM)	PS80: Protein Molar Ratio	Initial Amount of Monomer (%)	Change in Monomer After 12 Months at 5 °C	Change in Monomer After 24 Months at 5 °C	Change in Monomer After 12 Months at 25 °C	Change in Monomer After 1 Month at 40 °C
1	62	6.4	50	25	125	0.7	98.3	0.4	0.1	-2.5	-1.3
2	60	6.4	50	0	125	1.5	98.3	-0.2	-1.0	-3.9	-2.0
3	157	6.4	50	25	125	1.5	98.2	0.2	0.0	-3.1	-1.6
4	161	6.3	50	0	125	0.7	98.2	0.0	-0.4	-4.5	-2.1
5	60	6.2	50	25	125	1.5	98.3	0.3	0.2	-2.2	-1.4
6	110	6.0	50	0	125	0.7	98.3	0.1	-0.3	-3.6	-2.1
7	162	6.2	50	25	125	0.7	98.3	0.2	-0.1	-2.8	-1.7
8	160	6.0	50	0	125	1.5	98.3	-0.1	-0.4	-4.2	-2.3
9	169	6.3	25	25	125	0.7	98.2	--	-0.1	--	-1.8
10	158	6.3	25	25	123	1.0	98.1	--	--	--	-1.7

Table 6

Formulation #	Protein Conc. (mg/mL)	pH	Histidine Conc. (mM)	Citrate Conc. (mM)	Arginine Conc. (mM)	PS80: Protein Molar Ratio	Initial Amount of Aggregates (%)	Change in Aggregates After 12 Months at 5 °C	Change in Aggregates After 24 Months at 5 °C	Change in Aggregates After 12 Months at 25 °C	Change in Aggregates After 1 Month at 40 °C
11	160	6.3	0	40	125	0.7	0.5	--	0.4	--	0.9
12	165	6.3	0	40	125	1.5	0.6	0.3	--	--	0.9
13	62	6.2	0	40	125	1.5	0.5	--	--	1.7	0.4
14	170	6.1	0	40	125	1.5	0.5	--	--	--	0.9
15	165	6.5	0	63	125	1.5	0.5	--	--	--	1.0
16	160	6.3	0	40	125	1.0	0.6	0.3	--	--	0.9

Table 7

Formulation #	Protein Conc. (mg/mL)	pH	Histidine Conc. (mM)	Citrate Conc. (mM)	Arginine Conc. (mM)	PS80: Protein Molar Ratio	Initial Amount of Monomer (%)	Change in Monomer After 12 Months at 5 °C	Change in Monomer After 24 Months at 5 °C	Change in Monomer After 12 Months at 25 °C	Change in Monomer After 1 Month at 40 °C
11	160	6.3	0	40	125	0.7	98.3	--	-0.3	--	-2.2
12	165	6.3	0	40	125	1.5	98.2	0.1	--	-3.4	-2.1
13	62	6.2	0	40	125	1.5	98.1	--	--	--	-1.4
14	170	6.1	0	40	125	1.5	98.1	--	--	--	-2.0
15	165	6.5	0	63	125	1.5	98.2	--	--	--	-2.3
16	160	6.3	0	40	125	1.0	98.2	0.1	--	--	-2.0

5 pH

Several pH experiments were done to determine the effects of pH on CEX degradation at 5 °C. The vedolizumab antibody formulation comprised 160 mg/ml of anti- $\alpha 4\beta 7$ antibody, 125 mM arginine, 50 mM histidine, and 25 mM citrate.

Several different pH levels, 6.3, 6.5, 6.7 and 6.9 were tested for stability at 40°C, 25°C, and 5°C.

The CEX models at 40°C show (FIG. 10) that pH influences CEX degradation the most. The pH of formulations containing histidine decrease with

increasing temperature, however the pH of citrate formulations was shown to not be affected by temperature (FIG. 11). The histidine/citrate formulation was determined to have good stability at a pH of 6.8 at 40 °C after 1 week, 6.3-6.5 at 25 °C after 6 months and 6.3-6.5 at 5 °C after 6 months. Based on additional studies, the stability of the formulations were similar at 25°C and 5°C for the pH range of 6.2 to 6.9 (Tables 8 and 9).

Table 8

Protein Concentration (mg/mL)	pH	Histidine Concentration (mM)	Citrate Concentration (mM)	Arginine Concentration (mM)	PS80:Protein Molar Ratio	Initial Amount of Acidic Species (%)	Initial Amount of Basic Species (%)	Initial Amount of Major Isoform (%)	Difference in % Relative Area Over Time at 25 °C					
									Change in CE X Acidic After 6 Months	Change in CE X Acidic After 12 Months	Change in CE X Basic After 6 Months	Change in CE X Basic After 12 Months	Change in CE X Major After 6 Months	Change in CE X Major After 12 Months
157	6.4	50	25	125	1.5	23.9	6.8	69.3	8.5	17.4	7.1	3.4	-15.6	-20.8
162	6.2	50	25	125	0.7	24.0	6.9	69.1	4.4	12.8	10.8	8.6	-15.2	-21.4
158	6.3	50	25	125	1.5	24.8	5.5	69.7	7.2	--	4.9	--	-14.4	--
160	6.4	42	25	125	1.5	24.9	5.5	69.7	9.5	--	1.0	--	-14.4	--
147	6.7	45	25	125	2.1	24.9	4.7	70.4	14.8	--	3.6	--	-16.5	--
147	6.9	45	25	125	2.2	25.0	4.9	70.1	14.5	--	3.2	--	-17.9	--
153	6.7	46	25	125	1.5	24.8	5.5	69.7	14.7	--	0.3	--	-17.3	--
154	6.9	46	25	125	1.5	24.9	5.3	69.8	19.7	--	0.5	--	-20.2	--
170	6.5	50	25	125	1.0	25.7	4.6	69.7	10.8	--	4.4	--	-15.2	--
170	6.5	50	25	125	1.5	25.7	4.6	69.7	11.1	--	5.2	--	-16.4	--
160	6.5	50	25	125	1.5	26.3	7.0	66.7	11.8	--	-2.6	--	-11.9	--

10

15

Table 9

Protein Conc. (mg/mL)	pH	Histidine Conc. (mM)	Citrate Conc. (mM)	Arginine Conc. (mM)	PS80:Protein Molar Ratio	Difference in % Relative Area Over Time at 5 °C					
						Change in CEX Acidic After 6 Months	Change in CEX Acidic After 24 Months	Change in CEX Basic After 6 Months	Change in CEX Basic After 24 Months	Change in CEX Major After 6 Months	Change in CEX Major After 24 Months
157	6.4	50	25	125	1.5	0.2	0.6	2.3	-0.9	-2.5	0.3
162	6.2	50	25	125	0.7	-0.2	-0.7	4.3	2.2	-4.1	-1.5
158	6.3	50	25	125	1.5	0.0	--	1.9	--	-1.9	--
160	6.4	42	25	125	1.5	0.1	--	1.7	--	-1.8	--
147	6.7	45	25	125	2.1	1.7	--	1.7	--	-3.4	--
147	6.9	45	25	125	2.2	1.6	--	1.1	--	-2.7	--
153	6.7	46	25	125	1.5	1.7	--	0.4	--	-2.1	--
154	6.9	46	25	125	1.5	2.1	--	0.4	--	-2.4	--
170	6.5	50	25	125	1.0	0.9	--	1.6	--	-2.5	--
170	6.5	50	25	125	1.5	0.8	--	1.6	--	-2.5	--
160	6.5	50	25	125	1.5	11.8	--	-2.6	--	-11.9	--

EXAMPLE 2

STABILITY

Four different anti- $\alpha 4\beta 7$ antibody formulations were tested for stability over the course of twelve months. Formulations having a pH of 6.0-6.2 showed approximately 1-2% less major species than formulations having a pH of 6.3-6.4 (FIG. 12). Formulations having a pH of 6.3-6.4 showed less than 1% change in basic or major species at 5 °C.

Ten different anti- $\alpha 4\beta 7$ antibody formulations were tested for stability by SEC over the course of twelve months (Table 10). The formulations with 60 mg/mL protein concentration and containing 25 mM citrate had a change in aggregates of 0.1-0.2% after 1 year, while formulations containing 160 mg/mL protein and 25 mM citrate had an increase of aggregates from 0.2-0.3% over 1 year. There was an increase of 0.4-0.6% aggregates for formulations containing 60, 110, or 160 mg/mL protein with no citrate.

Table 10

Formulation #	Protein Concentration (mg/mL)	pH	Histidine Concentration (mM)	Citrate Concentration (mM)	Arginine Concentration (mM)	PS80 Molar Ratio	Change in % Aggregates at 5 °C After 1 Year
1	62	6.41	50	25	125	0.7	0.11
2	60	6.35	50	0	125	1.5	0.50
3	157	6.44	50	25	125	1.5	0.23
4	161	6.3	50	0	125	0.7	0.56
5	60	6.19	50	25	125	1.5	0.16
6	110	6.03	50	0	125	0.7	0.39
7	162	6.19	50	25	125	0.7	0.26
8	160	6	50	0	125	1.5	0.44
9	165	6.28	0	40	125	1.5	0.30
10	160	6.3	0	40	125	1.0	0.33

EXAMPLE 3

VISCOSITY

The injection force needed to administer the pharmaceutical formulation is related to the viscosity of the formulation. Formulations with varying pH and varying concentrations of protein, arginine, histidine, citrate, sucrose, and polysorbate 80 were made. The viscosity of these formulations was tested. A statistical model of the Ln (viscosity) was developed. The model showed that the viscosity is affected mainly by protein concentration and pH (Fig. 13). Sucrose, histidine and arginine also can have a minor effect on viscosity. In some protein formulations, sodium chloride is added to reduce the viscosity of the formulation. It is known, however, that the effect of sodium chloride on viscosity is protein and formulation dependent.

Sodium chloride was added to a formulation containing 140 mg/ml vedolizumab, 125 mM arginine, 25 mM histidine, 25 mM citrate, and polysorbate 80 at a 1.5 polysorbate 80 to protein molar ratio, and a pH of 6.4. The NaCl did not have any effect on the viscosity of the formulation.

The effects of the viscosity on the injection force of various syringes tested are shown in Figures 16A and 16B.

EXAMPLE 4

METHODS

Cation Exchange Chromatography (CEX)

A phosphate/sodium chloride gradient on a weak cation exchange column is used in a high performance liquid chromatography system to separate charged species in anti- $\alpha 4\beta 7$ antibody formulations and determine the charge composition of the antibody species. Acidic Isoforms elute before the Major Isoform and Basic Isoforms elute after the Major Isoform.

Stability data for a vedolizumab formulation generated using a CEX assay indicated that the % Major Isoform was above 55.0%.

Capillary Isoelectric Focusing (cIEF)

cIEF is performed using an iCE280 whole column detection cIEF system (Convergent Biosciences, Toronto, Ontario). Choice of ampholyte can be as recommended by the manufacturer or can be a combination of commercially available ampholytes. A useful combination is a mixture of 3-10 and 5-8 PHARMALYTE™ (GE Healthcare, Piscataway, NJ).

Stability data for a vedolizumab formulation generated using a cIEF assay indicated that the % Major Isoform was about 53%, the % Acidic Species was about 42% and the % Basic Species was about 5%.

Size Exclusion Chromatography (SEC)

SEC is performed using an analytical SEC column (Tosoh Bioscience, LLC, King of Prussia, PA). The mobile phase is a phosphate-buffered saline solution and the absorbance is monitored at 280 nm.

Stability data for a vedolizumab formulation generated using an SEC assay indicated that the % Monomer was 99.0%, the % Aggregates was <0.5% and the % Low Molecular Weight substances was <1.0%.

SDS-PAGE Assay

SDS-PAGE is performed using an Invitrogen (Carlsbad, CA) Tris-Glycine gel, 4-20% for reducing condition and 4-12% for non-reducing condition. The reconstituted antibody formulation sample is diluted in liquid formulation buffer then diluted one to two with Tris-Glycine SDS Sample Buffer (2X, Invitrogen) either with 10% 2-mercaptoethanol (reducing sample buffer) or without 2-mercaptoethanol (non-reducing sample buffer). Samples are briefly heated and loaded in comparison with a molecular weight marker (Invitrogen). The gels are stained with colloidal coomassie blue (Invitrogen) according to the manufacturer's instruction. Protein bands are analyzed by densitometry to identify the % heavy and light chain for reduced gels and % IgG for non-reduced gels.

Binding Efficacy

HuT78 cells (human T cell lymphoma cells, American Type Culture Collection, Manassas, VA) suspended in 1% BSA in PBS, 0.01% sodium azide are contacted with serial dilutions of primary test antibody. After incubation on ice, the cells are washed and treated with fluorescently labeled secondary antibody. After a further wash, the cells are fixed and suspended in FACS reagent for analysis by flow cytometry (Becton Dickinson Franklin Lakes, NJ); also see U.S. Patent No. 7,147,851.

10 Moisture by Karl Fischer

The formulation is titrated with methanol for a coulometric Karl Fischer moisture determination.

EXAMPLE 5

15 Effects of Silicone From Syringe Products Pre-filled with Anti- α 4 β 7 Antibody Formulation

A subcutaneous formulation consisting of 60-160 mg/mL of anti- α 4 β 7 protein in a buffer containing L-Histidine, L-Arginine Hydrochloride, Citrate and Polysorbate 80 is used to study the effects of silicone on the stability of the protein formulations and container/closure attributes. The study is performed with a 0.5 mL fill.

Parameters including the protein concentration, the polysorbate 80 to protein molar ratio, and the amount of silicone that is sprayed onto the syringe barrels are explored. The range of each of the input parameters is shown in Table 11.

30

Table 11. Input Parameter Ranges

Parameters	Low	High
Protein Concentration (mg/mL)	100	160
Polysorbate 80:Protein Ratio	0	2
Silicone Amount (mg)	0.4	0.8

A design of experiment is used to determine the set of formulations to study. A reasonable number of formulations range from 6 to 8 formulations. An example of the formulations that are tested is shown in Table 12.

Table 12

Run	Protein Conc. (mg/mL)	PS80:Protein Ratio	PS80 Concentration (%)	Silicone Level (mg)
1	100	1	0.087	0.8
2	100	2	0.174	0.8
3	160	0	0	0.8
4	160	2	0.279	0.4
5	100	0	0	0.4
6	160	2	0.279	0.8
7	100	1	0.087	0.4
8	160	0	0	0.4
9	100	0	0	0
10	100	2	0.174	0
11	160	0	0	0
12	160	2	0.279	0

Some controls may be added to the set of formulations and tested at a few select time points.

These formulations are placed on stability at several different temperatures (e.g., 5°C , 25°C/60%RH, 40°C/75% RH) and pulled at various time points (e.g., 0 week, 1 week, 2 weeks, 4 weeks, 8 weeks, 12 weeks, 6 months, and 12 months) for testing. Controls are tested at 0 weeks, 12 weeks, 6 months and 12 months.

The tests that are performed at each stability time pull include SEC, CEX, Instron, MFI, and Silicone Quantification. 1 syringe is tested for Instron, with the

expelled material being used for SEC, CEX, injection force measurements and microflow imaging (MFI), and silicone quantification.

EXAMPLE 6

5 Analysis of Pre-filled Syringe Components filled with Anti- α 4 β 7 Antibody Formulation

This study explored how various syringe manufacturers, plunger (stopper) elastomeric materials, and the amount of PS80 in the formulation affected the mechanical properties of the system and the stability of the formulation.

10 A design of experiment was created exploring 3 different syringe manufacturers, 2 different plunger (stopper) material types, and 2 different PS80 to protein molar ratios. The rest of the formulation was kept constant at 170 mg/mL of protein, 125 mM arginine, 50 mM histidine, 25 mM citrate, and a pH of 6.5. The needle size on these pre-filled syringes was 27G ½' or 29G ½" thin wall. The
15 experiments performed are detailed in Table 15.

The experimental design inputs for the active portion of the experiment are shown below in Table 13, while the constants are shown in Table 14. The experimental design was created utilizing the inputs shown in Table 13.

20 The list of experiments is shown in Table 10.

Table 13. DOE variables and levels with active formulation

Variable	Values		
PS80: Protein Molar Ratio	1.0	1.5	
Syringe Manufacturer	A	B	C
Plunger (Stopper) Type	4432	4023 Coated	

25

Table 14. Constants for active formulation

Constant	Value
Protein Concentration (mg/mL)	170
Arginine Concentration (mM)	125
Histidine Concentration (mM)	50
Citrate Concentration (mM)	25
pH	6.5

Table 15. Experimental details

Run #	Syringe Type	Plunger (Stopper)	PS80
1	C	4432	1
2	B	4432	1
3	A	4023	1
4	C	4023	1
5	B	4023	1
6	A	4023	1.5
7	C	4023	1.5
8	B	4023	1.5

5

A concentrated formulation anti- $\alpha 4\beta 7$ formulation is spiked with polysorbate 80 and diluted down to 170 mg/mL. The composition of the starting formulation is shown below in Table 16.

10

Table 16. Starting Formulation buffer details

Protein (mg/ml)	Total His (mM)	Total Citrate (mM)	Arg (mM)	pH
183	50	25	125	6.48

For the dilution of the material to the desired formulation composition, stock solutions of PS80 in 25 mM Citrate, 50 mM Histidine, 125 mM Arginine, pH 6.48 are made.

15

Table 17. Stock solution details

Excipient	Concentration
PS 80 (%)	5

The dilution scheme for the formulations is detailed in Table 18.

5 Table 18. Dilution detail

Starting Formulation (uL)	PS80 in His/Arg/Citrate Buffer (uL)	50 mM Histidine, 125 mM Arginine, 25 mM Citrate pH 6.48 Buffer	Total Volume (uL)
27868.9	890.8	1240.3	30000.0
18579.2	890.8	530.0	20000.0

Compounding is performed based on the dilution scheme, and the starting formulation should be weighed, while the other stock solutions can be pipetted volumetrically. Formulations are filtered. 0.5 mL of formulation is aliquotted into as many 1 mL Long syringes as possible. The syringes are stoppered by the stoppering machine with a 2-4 mm bubble. For each time point, there is one syringe stored needle down and one syringe stored sideways. The extra syringes are stored needle down.

15 The syringes are tested at 5, 25 and 40 °C on week 2 and at one month. Analytical testing (appearance, Instron, pH, osmolality, density, viscosity, SEC, CEX, and Brightwell) is performed initially and then again at 2 weeks at 25 and 40 °C and at 4 weeks at 25 °C.

EXAMPLE 7

Analysis of Subcutaneous Container Closures Used in 27 G Thin Wall Needle Syringes Pre-filled with Anti- $\alpha 4\beta 7$ Antibody Formulation

5 This study explores how various syringe models with a 27G thin wall needle and various plunger (stopper) manufacturers and models affect the mechanical properties of the system and the stability of the formulation over time.

This study explores how the stability of the anti- $\alpha 4\beta 7$ subcutaneous liquid formulation in a prefilled syringe and the mechanical properties of the syringe are
10 affected by the syringe manufacturer and the plunger (stopper) model for syringes with a 27GTW needle. The data generated from this study may determine the container/closure components for the liquid subcutaneous anti- $\alpha 4\beta 7$ formulation.

The experimental design inputs are shown below in Table 19, while the constants are shown in Table 20. The experimental design was created utilizing the
15 inputs shown in Table 19.

The list of experiments to be performed is shown in Table 21.

Table 19. DOE variables and levels with active formulation

20

Variable	Values			
Syringe Manufacturer	A	B	C	
Plunger (Stopper) Type	4432	4023 Coated	D	E

Table 20. Constants for active formulation

Constant	Value
Protein Concentration (mg/mL)	160
Arginine Concentration (mM)	125
Histidine Concentration (mM)	50
Citrate Concentration (mM)	25
PS80 (%)	0.2
pH	6.5

25

Table 21. Experimental details

Run #	Syringe	Plunger (Stopper)
1	B	D
2	B	4432
3	A	4432
4	B	4023 Coated
5	A	D
6	C	4023 Coated
7	A	4023 Coated
8	C	D
9	C	4432
10	C	E

A concentrated anti- α 4 β 7 formulation is spiked with polysorbate 80 and diluted down to 160 mg/mL. The composition of the starting formulation is shown below in Table 22.

Table 22. Starting Formulation buffer details

Protein (mg/ml)	Total His (mM)	Total Citrate (mM)	Arg (mM)	pH
180	50	25	125	6.3

For the dilution of the material to the desired formulation composition, stock solutions of PS80 in 25 mM Citrate, 50 mM Histidine and 125 mM Arginine, pH 6.3 are made.

Table 23. Stock solution details

Excipient	Concentration
PS 80 (%) in His/ Arg/ Citrate buffer pH 6.3	1.68

The dilution scheme for the formulations is detailed in Table 24.

Table 24. Dilution detail

Starting Formulation in His Arg Citrate buffer (mL)	PS80 in His/Arg/Citrate Buffer (mL)	Total Volume (mL)
78	10 (1.68%)	88

5 Compounding is performed based on the dilution scheme, and the starting formulation should be weighed, while the other stock solutions can be pipetted volumetrically. Formulations are filtered. 0.5 mL of formulation is aliquotted into as many 1 mL Long syringes as possible. The syringes are stoppered by the stoppering machine with a 2-4 mm bubble. For each time point, there is one syringe
10 stored needle down (horizontal position).

 The syringes are tested at 5 °C, 25 °C/60% RH, and 40 °C/75% RH at 1 month, 3 months, 6 months, 9 months (optional), 12 months, 18 months and 24 months.

 The liquid formulations are analytically tested (concentration, osmolality,
15 pH, Instron, MFI, SEC, and/or CEX) at 1, 3, 6, 9, 12, 18, 24 month (5°C); 1, 3, 6, 9, 12, 18, month (25°C); 1, 3, 6, 9, 12, month (40°C); and 1, 3, month (40°C).

EXAMPLE 8

20 Analysis of Subcutaneous Anti- α 4 β 7 Antibody Formulation in Plastic Prefilled Syringes

 This study is initiated to research the use of plastic syringes as the container/closure system for an anti- α 4 β 7 antibody subcutaneous formulation. The stability of a representative anti- α 4 β 7 antibody subcutaneous formulation in candidate plastic prefilled syringes is studied. The data generated from this study
25 helps to judge the applicability of using a plastic syringe for a liquid subcutaneous anti- α 4 β 7 antibody formulation.

Stability test samples are prepared as shown below. Stability tests are conducted under the storage conditions of 40 °C/75% RH, 25 °C/60% RH, and 5 °C.

Two types of plastic syringes and one glass syringe (Control) in Table 25 are tested with a liquid subcutaneous anti- $\alpha 4\beta 7$ antibody formulation shown in Table

26. Table 27 shows the details of each set of samples to be tested in the experiment.

Table 25. Plastic syringes

		Sample #1 Plastic syringe 1	Sample #2 Plastic syringe 2	Sample #3 Glass syringe (Control)
Syringe	Vendor	F	B	A
	Components	Syringe: polymer Needle: 27G(TW) Rigid needle shield	Syringe: polymer Needle: 26G(RW) Luer lock tip cap	Syringe: Glass Needle: 27G(TW) Rigid needle shield
	Silicon coating	Free	Not free	Not free
Plunger	Vendor	F	F	←
	Product description	1mL material A	1mL material B	←
	Silicon coating	No	Yes	←

Table 26. Anti- $\alpha 4\beta 7$ antibody subcutaneous formulation (pH 6.5)

Component	Composition
Anti- $\alpha 4\beta 7$ antibody	160 mg/mL
Arginine	125 mM
Histidine	50 mM
Citrate	25 mM
PS80 (Protein Molar Ratio)	1.5 (0.2w/v%)

10

Table 27. Sample details

Sample #	Plastic syringe vendor	Protein (MW:150000)		Arg (MW:174.20) (mM)	His (MW:155.15) (mM)	Citrate (MW:210.14) (mM)	PS 80 (MW:1309.68)		pH
		(mg/ml)	(mM)				(w/v%)	Protein Molar Ratio	
1	F	160	1.067	125	50	25	0.21	1.5	6.5
2	B	160	1.067	125	50	25	0.21	1.5	6.5
3 (Cont.)	A	160	1.067	125	50	25	0.21	1.5	6.5

Previously prepared liquid subcutaneous anti- $\alpha 4\beta 7$ formulation are used for this investigation. Formulations are filtered. Sampling the filtered solution for the

quality test as “before filling” sample (Appearance, MFI, DLS). 0.5 mL of formulation are aliquotted into 1 mL plastic syringes. The syringes are stoppered by the vacuum stoppering machine. The syringes are stored needle down.

An initial check is performed to measure pH, osmolality, density, viscosity, and protein concentration. Analytical testing (appearance, SEC(Aggregates, Monomer, LMW), CEX(Acidic, Main, Basic), glide force, MFI, DLS, and/or weight) is performed after 1 week, at 40 °C, 2 weeks, 40 °C, 1 month, 5, 25 and 40 °C, 3 months, 5 and 25 °C, 6 months at 5 and 25 °C, 9 months at 5 and 25 °C, and 12 months at 5 and 25 °C.

Samples are taken at 1 month, 3 months, 6 months, 9 months and 12 months at 5 °C and 25 °C. Samples are taken at 1 week, 2 weeks and 1 month at 40 °C.

Example 9:

The samples were analyzed for appearance, injection force, SEC, CEX, and micro-flow imaging at 5 °C and 25 °C at various time points that may have included 0, 1, 3, 6, and 12 months. The stability of the formulation as measured by SEC and CEX were similar to what was discussed in Examples 1 and 2. For injection force testing, the glide force were measured (Table 28). A statistical model determined that the only significant factor affecting the glide force was the syringe manufacturer, where A had higher glide forces than B, which was greater than C (Figure 17). The changes in glide force of the syringes over 12 months at 5 °C and 6 months at 25 °C were less than 10 N, but mostly less than 5 N.

Table 28

Run #	Syringe Manufacturer	Needle Size	Plunger (Stopper) Type	PS80:Protein Molar Ratio	Initial Glide Force (N)
1	C	27G	D	1	19.2
2	B	27G	D	1	22.9
3	A	29GTW	E	1	25.0
4	C	27G	E	1	18.5

5	B	27G	E	1	22.7
6	A	29GTW	E	1.5	28.8
7	C	27G	E	1.5	18.7
8	B	27G	E	1.5	23.7

EXAMPLE 10

Analysis of Pre-filled Syringe Components Used in 27 G Thin Wall Needle Syringes filled with Anti- α 4 β 7 Antibody Formulation

5 This study explored how various syringe manufacturers with a 27G thin wall needle and various plunger (stopper) manufacturers and elastomeric materials affected the mechanical properties of the pre-filled syringe system and the stability of the formulation over time.

10 Three different syringe manufacturers and 4 different plunger (stopper) models were tested with 27G ½” thin wall needles and a formulation containing 160 mg/mL protein, 125 mM arginine, 50 mM histidine, 25 mM citrate, 0.2% PS80, at a pH of 6.5. All of the samples created and tested are shown in Table 29.

Table 29. Experimental details

15

Run #	Syringe	Plunger (Stopper)
1	B	F
2	B	D
3	A	D
4	B	E
5	A	F
6	C	E
7	A	E
8	C	F
9	C	D
10	C	G

20 The samples were analyzed for appearance, injection force, SEC, CEX, and micro-flow imaging at 5 °C, 25 °C, and 40 °C at various time points that may have included 0, 1, 3, 6, and 12 months. The stability of the formulation as measured by SEC and CEX were similar to what was discussed in Examples 1 and 2. For

injection force testing, the breakloose and glide force were measured. The results at the initial time point are shown in Table 30.

5 **Table 30**

Run #	Syringe Manufacturer	Plunger (Stopper) Type	Initial Glide Force (N)	Initial Breakloose Force (N)	Breakloose Force at 12 Months at 5 °C (N)	Breakloose Force at 12 Months at 25 °C (N)	Breakloose Force at 12 Months at 40 °C (N)
1	B	F	12.0	4.0	3.8	12.9	28.7
2	B	D	11.9	3.9	4.6	12.4	36.0
3	A	D	7.0	4.0	6.5	5.1	6.4
4	B	E	13.9	4.5	4.7	5.8	17.2
5	A	F	5.7	4.1	3.0	17.5	23.9
6	C	E	6.7	4.1	5.0	5.8	11.4
7	A	E	7.9	7.6	4.3	10.4	6.1
8	C	F	6.3	4.2	4.1	15.0	33.3
9	C	D	5.9	4.8	3.9	4.4	10.0
10	C	G	7.2	4.6	6.1	9.8	13.0

A statistical model showed that syringe manufacturers A and C were similar and had lower glide forces than manufacturer B, while plunger (stopper) E have slightly higher glide force than the other plunger (stoppers).

10 Generally, the initial breakloose forces were similar between all the samples that were tested.

Over 12 months at 5 °C, 25 °C, and 40 °C, the glide forces did not significantly change. However, the breakloose force for syringes with plunger (stopper) F increased by 12 months at 25 °C and 40 °C.

15

EXAMPLE 11

Analysis of Anti- α 4 β 7 Antibody Formulation in Prefilled Syringes

This study determines how varying levels of protein concentration, polysorbate 80 concentration, citrate concentration, and pH affects anti- $\alpha 4\beta 7$ antibody formulations in a prefilled syringe format.

Part of the experimental design is created in JMP with a fraction factorial of two levels of protein concentration (60 to 160 mg/mL), pH (6.0 to 6.3), polysorbate 80:protein molar ratio (0.723 to 1.5), and citrate concentration (0 to 25 mM). These formulations have a constant value of Histidine concentration (50 mM) and Arginine (125 mM) (Formulations 1-8). Variations of these formulations with 25 mM Histidine are added (Formulations 9-10).

An additional set of formulations are developed to explore formulations with no histidine present and only citrate acting as the buffer (Formulations 11-16). The levels of the inputs for all formulations being explored are shown in Table 31. The constants used for all formulations are shown in Table 32.

Table 31. DOE variables and levels

Variable	Nominal Values	
	Low	High
Protein Concentration (mg/mL)	60	160
pH	6.0	6.3
PS80: Protein Molar Ratio	0.723	1.5
Citrate Concentration (mM)	0	40
Histidine Concentration (mM)	0	50

Table 32. Constants

Constant	Value
Arginine Concentration (mM)	125

Table 33 lists the formulations to be tested.

Table 33. Formulation details

Formulation #	Protein (mg/ml)	Protein (mM)	His (mM)	Arg (mM)	PS 80 %	pH	PS80:Protein Molar Ratio	Antioxidant	Antioxidant Concentration (mM)
1	60	0.400	50	125	0.038	6.3	0.723	Citric Acid	25
2	60	0.400	50	125	0.079	6.3	1.5	Citric Acid	0
3	157	1.047	50	125	0.206	6.3	1.5	Citric Acid	25
4	160	1.067	50	125	0.101	6.3	0.723	Citric Acid	0
5	60	0.400	50	125	0.079	6.0	1.5	Citric Acid	25
6	110	0.733	50	125	0.069	6.0	0.723	Citric Acid	0
7	160	1.067	50	125	0.101	6.0	0.723	Citric Acid	25
8	160	1.067	50	125	0.210	6.0	1.5	Citric Acid	0
9*	160	1.067	25	125	0.101	6.0	0.723	Citric Acid	25
10*	160	1.067	25	125	0.140	6.0	1	Citric Acid	25
11	160	1.067	0	125	0.101	6.3	0.723	Citric Acid	40
12	160	1.067	0	125	0.210	6.3	1.5	Citric Acid	40
13	60	0.400	0	125	0.079	6.3	1.5	Citric Acid	40
14*	160	1.067	0	125	0.210	6.1	1.5	Citric Acid	40
15*	160	1.067	0	125	0.210	6.6	1.5	Citric Acid	40
16*	160	1.067	0	125	0.140	6.3	1	Citric Acid	40

Each formulation is generated from a starting stock formulation containing anti- $\alpha 4\beta 7$ antibody and diluted down with various excipient stock solutions. In order to achieve reasonable dilution volumes, the anti- $\alpha 4\beta 7$ antibody stock solution used is shown in Table 34. Two different TFF operations are performed to achieve the formulations TFF 1 and 2. A portion of TFF 1 is used in a dialysis to achieve the formulation labeled “Dialysis”.

Table 34. Starting Formulation buffer details

Starting Formulation	Protein (mg/ml)	Total His (mM)	Total Citrate (mM)	Arg (mM)	pH
TFF 1	192.1	50	0	125	6.1
TFF 2	206.1	0	40	125	6.3
Dialysis	169.65	25	25	125	6.0

For the dilution of the material to the desired formulation composition, stock solutions of each excipient in water are made at the concentrations specified by Table 35.

Table 35. Stock solution details

Excipient	Concentration
Histidine (mM)	220
Arginine Hydrochloride (mM)	625
PS 80 (%)	2.5
Histidine Hydrochloride (mM)	600
Citric Acid (mM) (pH 6.3)	1500
Citric Acid (mM) (pH 6.0)	1500
Citrate (mM)	600
Sodium Citrate (mM)	800

The dilution scheme for the formulations is detailed in Table 36 and 37.

5

Table 36. Dilution detail

	Starting Formulation (uL)	Starting Formulation (mg)	His (uL)	His*HCl (uL)	Arg (uL)	Citrate Solution (uL)	PS80 (uL)	WFI (uL)
1	4685.06	4961.01	1612.2	268.4	2063.0	250	227.3	5894.0
2	4685.06	4961.01	1612.2	268.4	2063.0	0	471.6	5899.7
3	12259.2	12981.31	724.1	0.0	548.2	250.0	1234.0	0.0
4	12493.49	13229.36	696.7	0.0	501.3	0	606.2	702.4
5	4685.06	4961.01	1002.8	491.9	2063.0	250	471.6	6035.7
6	8589.28	9095.18	545.0	334.4	1282.1	0	416.7	3832.4
7	12493.49	13229.36	87.2	176.9	501.3	250	606.2	884.9
8	12493.49	13229.36	87.2	176.9	501.3	0	1257.6	483.5
9	12260.8	12941.30	38.2	16.8	147.8	12.3	525.3	0.0
10	12260.8	12941.30	38.2	16.8	147.8	12.3	726.6	0.0

Table 37. Dilution details

	Starting Formulation (uL)	Starting Formulation (mg)	Citrate (uL)	NaCitrate (uL)	Arg (uL)	PS80 (uL)	WFI (uL)
11	9315.87	9944.69	8.3	128.0	536.8	484.9	1526.1
12	9315.87	9944.69	8.3	128.0	536.8	1006.1	1005.0
13	3493.45	3729.26	30.5	402.5	1701.3	377.3	5995.0

14	5822.42	6215.43	22.0	67.4	335.5	628.8	623.9
15	5822.42	6215.43	0.0	300.0	335.5	628.8	413.3
16	5822.42	6215.43	5.2	80.0	335.5	419.2	837.7

Compounding is performed based on the dilution scheme, and the starting formulation is weighed, while the other stock solutions are pipetted volumetrically. Formulations are filtered. 0.5 mL of formulation is aliquotted into as many 1 mL Long syringes as possible. The syringes are stoppered by the stoppering machine. The syringes are stored needle down.

Liquid formulations are tested analytically (appearance, pH, osmolality, density, DLS, SEC, CEX, and/or Brightwell) initially, and at 1 week, 40 °C, 2 weeks, 40 °C, 1 month 25 and 40 °C, 2 months, 5 and 25 °C, 3 months, 5 and 25 °C, 6 months, 5 and 25 °C, 9 months, 5 and 25 °C, and 12 months, 5 and 25 °C.

Specific formulation pulls according to Table 38 are also performed.

Table 38: Specific formulation pulls

Formulation	Temperature	1 Week	2 Week	1 Month	2 Month	3 Month	6 Month	9 Month	12 Month	Extras
1	5	--	--	--	X	X	X	X	X	1
2	5	--	--	--	X	X	X	X	X	1
3	5	--	--	--	X	X	X	X	X	1
4	5	--	--	--	X	X	X	X	X	1
5	5	--	--	--	X	X	X	X	X	1
6	5	--	--	--	X	X	X	X	X	1
7	5	--	--	--	X	X	X	X	X	0
8	5	--	--	--	X	X	X	X	X	1
9	5	--	--	--	--	--	--	--	--	3
10	5	--	--	--	X	X	X	--	X	0
11	5	--	--	--	--	--	--	--	--	5
12	5	--	--	--	X	--	--	--	--	4
13	5	--	--	--	--	--	--	--	--	5
14	5	--	--	--	X	--	--	--	--	1
15	5	--	--	--	--	--	--	--	--	2
16	5	--	--	--	X	--	--	--	--	1
1	25	--	--	X	X	X	X	X	X	1
2	25	--	--	X	X	X	X	X	X	1
3	25	--	--	X	X	X	X	X	X	1
4	25	--	--	X	X	X	X	X	X	1
5	25	--	--	X	X	X	X	X	X	1
6	25	--	--	X	X	X	X	X	X	1
7	25	--	--	X	X	X	X	X	X	1

8	25	--	--	X	X	X	X	X	X	1
9	25	--	--	X	--	--	--	--	--	3
10	25	--	--	X	X	--	X	--	--	0
11	25	--	--	X	--	--	--	--	--	5
12	25	--	--	X	X	--	--	--	--	4
13	25	--	--	X	--	--	--	--	--	6
14	25	--	--	X	X	--	--	--	--	1
15	25	--	--	X	--	--	--	--	--	2
16	25	--	--	X	X	--	--	--	--	1
1	40	X	X	X	--	--	--	--	--	1
2	40	X	X	X	--	--	--	--	--	1
3	40	X	X	X	--	--	--	--	--	1
4	40	X	X	X	--	--	--	--	--	1
5	40	X	X	X	--	--	--	--	--	1
6	40	X	X	X	--	--	--	--	--	1
7	40	X	X	X	--	--	--	--	--	1
8	40	X	X	X	--	--	--	--	--	1
9	40	X	X	X	--	--	--	--	--	0
10	40	X	X	X	--	--	--	--	--	0
11	40	X	X	X	--	--	--	--	--	0
12	40	X	X	X	--	--	--	--	--	0
13	40	X	X	X	--	--	--	--	--	0
14	40	X	X	X	--	--	--	--	--	0
15	40	X	X	X	--	--	--	--	--	0
16	40	X	X	X	--	--	--	--	--	0

EXAMPLE 12

A formulation containing 160 mg/mL protein, 50 mM histidine, 25 mM citrate, 125 mM arginine at pH 6.5 was tested for stability in either a glass syringe or two different COP plastic syringes. At 5 °C and 25 °C after 12 months, the amount of aggregates and monomer were comparable between the plastic and glass syringes.

Table 39

Formulation #	PS80:Protein Molar Ratio	Syringe Material	Change in SEC Aggregates After 12 Months at 5 °C (%)	Change in SEC Aggregates After 12 Months at 25 °C (%)	Amount of Monomer After 12 Months at 5 °C (%)	Amount of Monomer After 12 Months at 25 °C (%)
1	1.5	COP Manufacturer	0.2	1.0	98.3	96.8

		1				
2	1.5	COP Manufacturer 2	0.2	1.6	98.3	96.9
3	1.5	Glass	0.2	1.4	98.4	96.8
4	1	Glass	0.2	1.6	98.3	96.8

Example 13: Bioavailability of Vedolizumab Administered by Subcutaneous and Intramuscular Injection

A phase I study of the bioavailability of vedolizumab administered by subcutaneous and intramuscular injection to healthy male subjects was completed. A total of 42 healthy males were enrolled in the study. The subjects were divided into three groups (subcutaneous, intramuscular, and intravenous administration) of 14 subjects each. The subjects were administered 180 mg of vedolizumab on one day. The dose was reconstituted from a lyophilized formulation of 60 mg/ml antibody in 50 mM histidine, 125 mM arginine, 0.06% polysorbate 80, 10% sucrose, at pH 6.3. For the intramuscular and subcutaneous subjects, the dose was divided into two injections of 1.5 ml each. Blood was sampled to determine the plasma vedolizumab concentration and the bioavailability of vedolizumab in each set of subjects was determined.

No serious adverse events or significant infections, clinically significant abnormalities, positive subjective/objective RAMP checklists, nor clinically significant ECG findings were reported.

PK/PD modeling and simulation was completed to determine the dose and regimens of extra-vascular doses that result in similar exposures as the intravenous doses in order to maintain this desired serum concentrations at trough levels.

The absorption profile (FIG. 18) showed that concentrations of the intramuscular and subcutaneous doses generally overlap. There are no apparent gross differences in the absorption profiles of these routes of administration. The absolute bioavailability of vedolizumab following SC injection was approximately 75% and following IM injection was approximately 80%.

Example 14. Modeling Subcutaneous dose Regimens

PK/PD modeling and simulation was completed to determine the dose and regimens of extra-vascular doses that result in similar exposures as the intravenous doses in order to maintain certain serum concentrations at trough levels.

5

A final combined dataset (IV, SC and IM data) showed two compartment linear models parameterized in terms of clearance (CL) and central volume of distribution (V2), peripheral volume of distribution (V3), an extra-vascular route dependent absorption rate constant (KA) and the relative bioavailability (compared to intravenous administration) of the extra-vascular doses (F). IIV terms were included on CL, V2 and V3 with body weight as the only covariate influencing CL and V3 through an allometric effect.

Model acceptability and predictability was demonstrated through bootstrap parameter estimates, visual predictive checks and goodness of fit plot. Analysis of the model identified body weight as a predictor for the PK of vedolizumab with the variability in the PK attributed to between subject and within subject components.

Once the model was demonstrated to be adequate for simulation, simulations were performed in order to assess the effect of route of administration (IV, IM or SC), and assess the effect of frequency of dosing (weekly, every 2 weeks, every 4 weeks, and every 8 weeks) on the steady state trough concentrations. Based on these values and the relative bioavailability of vedolizumab following IM and SC administration (F=69.5%), doses were selected to achieve similar trough concentrations as the IV doses.

25

Simulations modeled doses and regimens to match intravenous induction and maintenance regimens. The targets were both exposure (area under serum drug concentration-time curve (AUC)) and trough drug concentration. Tables 40-43 provide results of simulations.

30

Table 40. Induction regimen matching an IV AUC during weeks 0-6

Route	Dose	Frequency
IV	300 mg	Week 0 & 2
SC	485 mg	Week 0 & 2
SC	160 mg	Every other day (6 doses)
SC	>160 mg	Weekly (6 doses)

Table 41. Induction regimen matching an IV trough concentration, weeks 0-6

Route	Dose	Frequency
IV	300 mg	Week 0 & 2
SC	>160 mg	Week 0 & 2
SC	100 mg	Every week (6 doses)
SC	160 mg	Every other day (for 2 weeks)

Table 42. Maintenance regimen matching a 300 mg IV dose every 4 weeks

Frequency	Route	Dose matching 4wk IV steady state trough concentration	Dose matching 4wk IV AUC
Once every 4 weeks	IV	300	300
	IM	432	432
	SC	432	432
Once every 2 weeks	IV	115	150
	IM	165	216
	SC	165	216
Every week	IV	50	75
	IM	72	108
	SC	72	108

5

Table 43. Maintenance regimen matching a 300 mg IV dose every 8 weeks

Frequency	Route	Dose matching	Dose matching
-----------	-------	---------------	---------------

		8wk IV steady state trough concentration	8wk IV AUC
Once every 8 weeks	IV	300	300
	IM	432	432
	SC	432	432
Once every 4 weeks	IV	90	150
	IM	125	216
	SC	125	216
Once every 2 weeks	IV	35	75
	IM	50	108
	SC	50	108
Every week	IV	15	37.5
	IM	22	54
	SC	22	54

Example 15: Phase 2a Multiple Dose Study

A Phase 2a multiple dose study can assess the safety, tolerability and steady state PK of vedolizumab following multiple doses of vedolizumab by the subcutaneous administration route and to assess the relative bioavailability of the subcutaneous regimen compared with the intravenous regimen. The development of HAMA and neutralizing HAMA and the effect on PD of multiple doses of vedolizumab following subcutaneous administration can be assessed.

Ulcerative colitis patients having a partial Mayo score of 1-12 and Crohn's disease patients having a CDAI greater than 150 can be included in the study. Cohorts can receive an induction regimen of vedolizumab (300 mg) administered IV at weeks 0 and 2, followed by a maintenance regimen of either

- Vedolizumab (300 mg) administered IV every 4 weeks at weeks 6-22
- Vedolizumab (300 mg) administered IV every 8 weeks at weeks 6-22
- Vedolizumab (108 mg) administered SC every week at weeks 6-22
- Vedolizumab (108 mg) administered SC every 2 weeks at weeks 6-22

Vedolizumab (165 mg) administered SC every 3 weeks at weeks 6-22.

Samples can be collected before dosing on day 1, and then again on day 1
(12 hours), 2, 3, 5, 8, 15, 29, 43, 127, 127 (12 hours), 128, 129, 131, 134, 141, and
5 155 to assess PK and PD.

Example 16: Long-Term Clinical Experience with Vedolizumab for the Treatment of IBD

A phase 2 open-label safety extension study was completed to assess the
10 long-term pharmacokinetics (PK), pharmacodynamics (PD), safety, and efficacy of
vedolizumab. Patients were aged 18 to 75 years old, and had either previously
participated in an earlier PK/PD/safety study in ulcerative colitis patients or had IBD
symptoms for at least 2 months confirmed endoscopically and/or histopathologically
and/or radiologically within 36 months of screening.

15 All patients received an intravenous dosing regimen of either 2 mg/kg or 6
mg/kg of vedolizumab (5 mg/mL antibody, 20 mM citrate/citric acid, 125 mM
sodium chloride, 0.05% polysorbate 80, pH 6.0 (stored long term -70°C and up to 3
mo -20°C)) on days 1, 15 and 43, followed by a dose every 8 weeks for up to a total
of 78 weeks. Patients were either treatment-naïve ulcerative colitis or Crohn's
20 disease patients, or ulcerative colitis patients that had participated in an earlier
clinical trial.

Efficacy/quality of life (QoL); partial Mayo score (PMS), Crohn's disease
activity index (CDAI), and Inflammatory Bowel Disease Questionnaire (IBDQ)
were used to assess the results of the study.

25 PK Results

Mean pre-infusion vedolizumab concentrations were dose proportional, and
remained steady and detectable throughout the study.

30

PD Results

Receptors (%ACT-1 + [CD4+CD45RO HIGH] and % MADCAM+ [CD4+CD45RO HIGH] were almost fully inhibited throughout the study period at all dose levels.

5

Partial Mayo Score

Baseline mean PMS was higher for treatment-naïve ulcerative colitis patients (5.4) than for ulcerative colitis rollover patients (2.3). By day 43, mean PMS showed a pronounced decrease for both rollover and treatment-naïve ulcerative colitis patients. By day 155, mean scores of the two groups were similar. Mean PMS continued to decrease through day 267, and leveled off thereafter.

10

Crohn's Disease Activity Index

CD patients' mean CDAI decreased from 294.6 at baseline to 237.7 at Day 43, and continued to decrease through day 155 (156.1).

15

IBDQ

Ulcerative colitis rollover patients had the highest mean IBDQ scores at baseline. By day 43, mean IBDQ scores had increased in all three disease groups. Mean IBDQ scores continued to increase over time in all 3 disease groups, reaching a maximum at day 155 for Crohn's Disease patients, and at day 491 for treatment-naïve ulcerative colitis patients and ulcerative colitis rollover patients.

20

C- reactive protein

Both ulcerative colitis rollover and Crohn's disease patients showed decreased mean CRP levels through day 155 and then leveled off. Treatment-naïve ulcerative colitis patients had a lower mean CRP level at baseline than ulcerative colitis rollover patients (2.28 v. 7.09). Mean CRP levels of the treatment-naïve ulcerative colitis patients remained relatively constant at all time points assessed.

25

30

Other Safety Results

No systematic opportunistic infections (including PML) were reported during the study. One patient tested positive for JC viremia at a single time point, though was negative for JCV at all other time points. Three of 72 patients (4%) had
5 positive HAHA results (two of these were transiently positive). The study showed no evidence of liver toxicity, lymphocytosis, or lymphopenia, or any other drug-associated laboratory changes.

Conclusions

10 Vedolizumab administered at 2.0 or 6.0 mg/kg once every 8 weeks for up to 78 weeks achieved target receptor saturations, was associated with durable mean decreases in disease activity and improved IBDQ scores, was generally safe and well tolerated, and demonstrated acceptable immunogenicity.

15 Example 17: Induction and Maintenance of Response and Remission in Patients with Moderately to Severely Active Ulcerative Colitis

A single trial comprising two randomized, double blind, multi-center studies designed to evaluate induction and maintenance of response and remission in
20 patients with moderately to severely active ulcerative colitis. Demographic and baseline disease characteristics were comparable across all treatment groups.

The induction study, using intravenous administration, compared placebo against vedolizumab, at a 300 mg dose reconstituted from a lyophilized formulation of 60 mg/ml antibody in 50 mM histidine, 125 mM arginine, 0.06% polysorbate 80,
25 10% sucrose, at pH 6.3, with an endpoint at 6 weeks after 2 doses of vedolizumab.

The maintenance study, using the same formulation and route of administration as the induction study, compared placebo against vedolizumab dosed every four weeks, and placebo against vedolizumab dosed every eight weeks. The endpoint of this study was at 52 weeks, analyzing the induction responder
30 population.

Blood samples were collected to measure concentrations of vedolizumab during the study. The mean serum concentration of vedolizumab at the end of the induction phase was 20 to 30 µg/mL. The mean vedolizumab trough serum concentrations at steady state after 30 min IV infusion of 300mg dose administration were between 9 to 13 µg/mL for the q8wks regimen (8 week regimen) and between 35 to 40 µg/mL for the q4wks regimen (4 week regimen). At the end of infusion, the vedolizumab median plasma concentrations were between 98 and 101 µg/mL for the q8wks regimen and around 129 and 137 µg/mL for the q4 wks regimen.

Summaries of the responses of the induction and maintenance studies are provided in Tables 44-47. A significantly greater proportion of vedolizumab-treated patients achieved clinical response, remission, and mucosal healing at 6 weeks, compared with placebo (Table 44). 39% of the induction phase intent-to-treat population had prior anti-TNFα failure. Clinical response and remission rates were higher in vedolizumab than placebo patients among both those with prior anti-TNF failure and those with no prior anti-TNF exposure. In preliminary analyses through week 6, rates of adverse events (AEs), serious AEs, and adverse events leading to study discontinuation were higher in the placebo group than vedolizumab group. A significantly greater proportion of vedolizumab patients than placebo patients achieved clinical remission, mucosal healing, and corticosteroid-free remission at 52 wks and durable response and remission (Table 45). 32% of the maintenance study population had prior anti-TNFα failure. Clinical remission and durable clinical response rates were greater with vedolizumab than placebo in both TNF failure and TNF naïve patients. In the safety population (N=895) for wks 0–52, rates of adverse events (AEs), serious AEs, and serious infections were similar between vedolizumab and placebo groups. No increase in rates of opportunistic or enteric infections was observed in the vedolizumab group.

Table 44: Induction Study Results—Primary and Key Secondary Endpoints

Efficacy Endpoints	Placebo	Vedolizumab	Difference/RR	P value

Clinical Response (%)	25.5%	47.1%	21.7%/1.8	<0.0001
Clinical Remission (%)	5.4%	16.9%	11.5%/3.1	0.0010
Mucosal Healing (%)	24.8%	40.9	16.1%/1.6	0.0013

Table 45: Maintenance Study Results—Primary and Key Secondary Endpoints

Efficacy Endpoint	Placebo N=126	VDZ Q8 N=122	VDZ Q4 N=125	Difference/RR Q8 vs. Pb Q4 vs. Pb	P value
Clinical Remission (%)	15.9	41.8	44.8	26.1/2.7 29.1/2.8	<0.0001 <0.0001
Durable Response (%)	23.8	56.6	52.0	32.8/2.4 28.5/2.2	<0.0001 <0.0001
Mucosal Healing (%)	19.8	51.6	56.0	32.0/2.6 36.3/2.8	<0.0001 <0.0001
Durable Remission (%)	8.7	20.5	24.0	11.8/2.4 15.3/2.8	0.0090 0.0011
Corticosteroid-free Remission (%)	13.9 n=72	31.4 n=70	45.2 N=73	17.6/2.3 31.4/3.3	0.0133 <0.0001

5 Table 46: Induction Study: Clinical Response and Remission at 6 Weeks in Patients with Prior Anti-TNF- α Antagonist Failure and Without Anti-TNF Exposure, ITT Population

Patients with Prior Anti-TNF- α Antagonist Failure (39%)				
Endpoint	Placebo N=63	Vedolizumab N=82	Difference	95% CI
Clinical	20.6	39.0	18.4	3.9, 32.9

Response (%)				
Clinical	3.2	9.8	6.6	-9.8, 22.8
Remission (%)				
Patients Without Anti-TNF- α Antagonist Exposure (55%)				
	Placebo N=76	Vedolizumab N=130	Difference	95% CI
Clinical	26.3	53.1	26.8	13.7, 39.9
Response (%)				
Clinical	6.6	23.1	16.5	2.4, 30.2
Remission (%)				

Table 47: Clinical Remission and Durable Clinical Response at 52 Weeks:
Patients with Prior Anti-TNF- α Antagonist Failure or Without Anti-TNF- α
Antagonist Exposure ITT Population

Patients with Prior Anti-TNF- α Antagonist Failure (32%)					
Endpoint	Placebo N=38	VDZ Q8Wks N=43	VDZ Q4Wks N=40	Difference Q8wks vs Placebo Q4 wks vs. Placebo	95% CI
Clinical remission (%)	5.3	37.2	35.0	31.9 29.7	10.3, 51.4 7.4, 49.4
Durable Clinical Response (%)	15.8	46.5	42.5	30.7 26.7	11.8, 49.6 7.5, 45.9
Patients without Anti-TNF- α Antagonist Exposure (60%)					
	Placebo	VDZ	VDZ	Difference	95% CI

	N=79	Q8wks N=72	Q4wks N=73	Q8wks vs. Placebo Q4wks vs. Placebo	
Clinical Remission (%)	19.0	45.8	47.9	26.8 29.0	12.4, 41.2 14.6, 43.3
Durable Clinical Response (%)	26.6	65.3	56.2	38.7 29.6	24.0, 53.4 14.6, 44.6

Example 18: Induction and Maintenance of Response and Remission in Patients with Moderately to Severely Active Crohn's Disease

A single trial comprising two randomized, double blind, multi-center studies designed to evaluate induction and maintenance of response and remission in patients with moderately to severely active Crohn's Disease. Demographic and baseline disease characteristics were comparable across all treatment groups.

The induction study, using intravenous administration, compared placebo against vedolizumab, at a 300 mg dose reconstituted from a lyophilized formulation of 60 mg/ml antibody in 50 mM histidine, 125 mM arginine, 0.06% polysorbate 80, 10% sucrose, at pH 6.3, with an endpoint at 6 weeks after 2 doses of vedolizumab.

The maintenance study, using the same formulation and route of administration as the induction study, compared placebo against vedolizumab dosed every four weeks, and placebo against vedolizumab dosed every eight weeks. The endpoint of this study was at 52 weeks, analyzing the induction responder population.

Surprisingly, this study showed that Q4 and Q8 week groups yielded very similar results. Summaries of the responses of the induction and maintenance studies are provided in Tables 48-51. A significantly greater proportion of

vedolizumab-treated patients achieved clinical remission and enhanced response, compared with placebo (Table 48). Clinical remission and enhanced response rates were higher in vedolizumab than placebo patients among both those with prior anti-TNF failure and those with no prior anti-TNF exposure. Rates of adverse events (AEs), serious AEs, and serious infections were similar between vedolizumab and placebo groups. No increase in rates of opportunistic or enteric infections was observed in the vedolizumab group.

Table 48: Induction Study Results—Primary and Secondary Endpoints

Endpoints	Placebo N=148	Vedolizumab N=220	Adjusted Difference/RR	P value
Clinical Remission (%)	6.8%	14.5%	7.8%/2.1	0.0206
Enhanced Response (%)	25.7%	31.4%	5.7%/1.2	0.2322
Mean CRP Change (µg/mL)	-3.6 N=147	-2.9 N=220		0.9288

Table 49: Maintenance Study Results—Primary and Key Secondary Endpoints

Efficacy Endpoint	Placebo N=153	VDZ Q8 N=154	VDZ Q4 N=154	Adj. Difference/RR Q8 vs. Pb Q4 vs. Pb	P value
Clinical Remission (%)	21.6	39.0	36.4	17.4/1.8 14.7/1.7	0.0007 0.0042
Enhanced Response (%)	30.1	43.5	45.5	13.4/1.4 15.3/1.5	0.0132 0.0053
Corticosteroid-free	15.9	31.7	28.8	15.9/2.0	0.0154

Remission (%)	N=82	N=82	N=80	12.9/1.8	0.0450
Durable Remission (%)	14.4	21.4	16.2	7.2/1.5	0.1036
				2.0/1.1	0.6413

Table 50: Clinical Remission and Enhanced Response at 6 Weeks in Patients with Prior Anti-TNF- α Antagonist Failure and Without Anti-TNF Exposure, ITT Population

Patients with Prior Anti-TNF- α Antagonist Failure (48%)				
Endpoint	Placebo N=70	Vedolizumab N=105	Difference	95% CI
Clinical Remission (%)	4.3	10.5	6.2	(-9.1, 21.3)
Enhanced Response (%)	22.9	23.8	1.0	(-11.8, 13.7)
Patients Without Anti-TNF- α Antagonist Exposure (50%)				
	Placebo N=76	Vedolizumab N=130	Difference	95% CI
Clinical Remission (%)	9.2	17.4	8.2	(-1.4, 17.9)
Enhanced Response (%)	30.3	42.2	11.9	(-1.9, 25.8)

5

Table 51: Clinical Remission and Enhanced Response at 52 Weeks: Patients with Prior Anti-TNF- α Antagonist Failure or Without Anti-TNF- α Antagonist Exposure ITT Population

Patients with Prior Anti-TNF- α Antagonist Failure (51%)					
Endpoint	Placebo N=78	VDZ Q8Wks N=82	VDZ Q4Wks N=77	Difference Q8wks vs Placebo Q4 wks vs.	95% CI

				Placebo	
Clinical remission (%)	12.8	28.0	27.3	15.2 14.5	(3.0, 27.5) (2.0, 26.9)
Enhanced Response (%)	20.5	29.3	37.7	8.8 17.1	(-4.6, 22.1) (3.1, 31.2)
Patients without Anti-TNF- α Antagonist Exposure (45%)					
	Placebo N=71	VDZ Q8wks N=66	VDZ Q4wks N=71	Difference Q8wks vs. Placebo Q4wks vs. Placebo	95% CI
Clinical Remission (%)	26.8	51.1	46.5	24.8 19.7	(8.9, 40.6) (4.2, 35.2)
Enhanced Response (%)	38.0	60.6	53.5	22.6 15.5	(6.3, 38.9) (-0.7, 31.7)

Example 19: Induction of Response and Remission in Patients with Moderate to Severely Active Crohn's Disease

A randomized, double blind, placebo controlled multi-center study was completed to evaluate the induction effect of vedolizumab at 300 mg doses (reconstituted from a formulation of 60 mg/ml antibody in 50 mM histidine, 125 mM arginine, 0.06% polysorbate 80, 10% sucrose, at pH6.3 which was lyophilized), in TNF α antagonist failure patients at week 6 (after 2 doses--0 and 2 weeks) and at

week 10 (after 3 doses). The study consisted of 416 patients, 75% of whom were TNF α antagonist failures, and 25% of whom were TNF α naïve. Demographics and concomitant IBD medication were balanced across treatment groups. Baseline disease characteristics were also balanced across treatment groups, except for

5 baseline disease activity.

The primary endpoint designated for the study was week 6 remission (%) in anti-TNF- α antagonist failure population. The key secondary endpoints that were evaluated (sequential testing procedure) were: week 6 remission (%) in overall population, week 10 remission (%) in anti-TNF- α antagonist failure and overall

10 population (using Hochberg procedure), week 6 and 10 sustained remission (%) in anti-TNF- α antagonist failure and overall population (using Hochberg procedure), and week 6 enhanced response (%) in anti-TNF- α antagonist failure population.

Table 52: Baseline CDAI:

	<u>Placebo</u>	<u>Vedolizumab</u>	<u>p-value</u>
<u>TNF ITT: Mean</u> <u>(Std Dev)</u>	<u>306.1 (55.43)</u>	<u>316.1 (52.63)</u>	<u>0.0945</u>
<u>Overall ITT: Mean</u> <u>(Std Dev)</u>	<u>301.3 (54.97)</u>	<u>313.9 (53.17)</u>	<u>0.0153</u>

15

Table 53: Induction Study Results: Primary and Key Secondary Endpoints

Endpoints	TNF ITT (N=315)				Overall ITT (N=416)			
	PLA N=157	VDZ V=158	Diff (RR)	P- value	PLA N=207	VDZ N=209	Diff (RR)	P-value
Primary Wk6 Remission	12.1 %	15.2 %	3.0 % (1.2)	0.4332				
1st Secondary Wk6					12.1 %	19.1 %	6.9 % (1.6)	0.0478

Remission								
2nd Secondary Wk10 Remission	12.1 %	26.6 %	14.4 % (2.2)	0.0012	13 %	28.7 %	15.5 % (2.2)	<0.0001
Sustained Remission (both Wk 6&10)	8.3 %	12.0 %	3.7 % (1.4)	0.2755	8.2 %	15.3 %	7% (1.9)	0.0249
Enhanced Response (CDAI100)	22.3 %	39.2 %	16.9% (1.8)	0.0011				

Table 54: Results in Anti-TNF- α Antagonist Naïve Patients (n=101, 24% of overall)

	Placebo %	Vedolizumab %	Difference %	95% CI
Remission Week 6	12	31.4	19.1	(3.3, 35.0)
Remission Week 10	16	35.3	19.2	(2.4, 35.8)

5 Table 55: Study Results: Clinical Remission at Weeks 6 and 10, Key Subgroup—Previous Tx Failures, ITT Overall

Subgroup	Variable	Placebo	VDZ	Diff	95% CI
Any prior anti-TNF failure (75% of ITT)	N	156	155		
	Wk6 Rem (%)	12.8	14.8	2	(-5.7, 9.7)
	Wk10 Rem (%)	12.8	26.5	13.6	(4.9, 22.3)
Prior immunomodulator failure but not	N	45	44		
	Wk 6 Rem (%)	11.1	31.8	20.7	(-0.5, 39.7)

anti-TNF failure (21% ITT)	Wk10 Rem (%)	15.6	31.8	16.3	(-1.1, 33.6)
Prior corticosteroid failure only (3% ITT)	N	5	9		
	Wk6 Rem (%)	0	33.3	33.3	(-23.9, 75.7)
	Wk10 Rem (%)	0	44.4	44.4	(-13.4, 85.3)

The study showed that TNF- α antagonist failure patients required 3 doses for induction of remission. Remission rates in TNF- α antagonist failure patients increased between week 6 and week 10, but only for the vedolizumab group (not placebo). Remission rates for TNF- α antagonist naïve patients did not increase substantially between week 6 and 10. Of the TNF- α antagonist failure population with a high degree of disease severity, 43% never responded to a TNF- α antagonist, and 45% lost response.

Example 20: Stability

Various different anti- $\alpha 4\beta 7$ antibody formulations were tested for stability over the course of 6 to 24 months at 5°C (Tables 6 and 7). Formulations having a pH of 6.0-6.2 showed approximately less than 4% major species degradation after 6 months and at 24 months.

Various different anti- $\alpha 4\beta 7$ antibody formulations were tested for stability by SEC for up to 24 months (Tables 4 and 5). The formulations with 60 mg/mL protein concentration and containing 25 mM citrate had a change in aggregates of 0.1-0.2% after 2 years, while formulations containing 160 mg/mL protein and 25 mM citrate had an increase of aggregates of approximately 0.3% over 2 years. There was an increase of 0.6-1.1% aggregates for formulations containing 60, 110, or 160 mg/mL protein with no citrate. For the formulations tested containing citrate, but no histidine, after 12 months and 24 months, there was approximately 0.3-0.4% growth of aggregates.

Example 21: Determination of the effect of vedolizumab on the CD4:CD8 ratio

Healthy subjects ages 18–45 were treated with a single 450 mg dose of vedolizumab reconstituted from a lyophilized formulation of 10% sucrose and diluted into an infusion system of 0.9% saline. Cerebrospinal fluid (CSF) was collected by lumbar puncture before (baseline) and 5 weeks after the single 450-mg dose of vedolizumab. Each subject served as his/her own control.

A 5-week time point was selected based on a previous study that showed patients with MS treated with natalizumab demonstrated effects on CSF CD4+:CD8+ lymphocyte ratio and reduction in number of brain lesions after only one dose (Stuve et al. *Arch Neurol*.2006;63:1383-1387; Stuve et al. *Ann Neurol*. 2006;59:743-747. Miller et al. *N Engl J Med*. 2003;348(1):15-23); and also because at 5 weeks, a 450-mg dose of vedolizumab is sufficient to saturate the target and provides serum concentrations that exceed estimated steady-state trough levels associated with the phase 3 dose regimen of 300 mg every 4 weeks.

Approximately 15 mL CSF was obtained from each subject for immunophenotyping. CSF samples were included for analyses if they met the following criteria: ≤ 10 RBCs/ μ L per sample (to minimize peripheral blood contamination); negative CSF culture result; adequate T-lymphocyte numbers in each flow cytometry sample; and no detection of serum antibodies to vedolizumab.

Week 5 median (34.80 μ g/mL) and individual subject serum vedolizumab concentrations (range 24.9–47.9 μ g/mL) were higher than projected steady-state trough concentration (~ 24 μ g/mL) for the phase 3 dose regimen. A high degree ($>90\%$) of $\alpha 4\beta 7$ receptor saturation was observed at week 5 as measured by MAdCAM-1-Fc, indicating vedolizumab's saturation of its target at the time of endpoint assessment.

Vedolizumab was not detected in any CSF sample (detection limit = 0.125 μ g/mL).

30

Effect on CD4+ and CD8+ T Lymphocyte Numbers and Ratio

Vedolizumab did not significantly reduce CD4+:CD8+ ratio (Table 56). None of the subjects had a postdose CD4+:CD8+ ratio <1 ($p < 0.0001$ (1-sided t-test)). Vedolizumab did not significantly reduce the number of CD4+ or CD8+ T lymphocytes in CSF. In addition, there were no significant changes in CSF % CD4+ and % CD8+ T lymphocytes (Table 57). Also, no significant changes in peripheral blood WBC, CD4+ and CD8+ memory T lymphocytes (Table 58) were observed.

Table 56: Effect of Treatment on CSF CD4+:CD8+ Ratio (Evaluable Population, n=13)

	Baseline	Week 5	CD4+:CD8+ Ratio Difference†
CD4+:CD8+ ratio	3.59 (0.273)	3.60 (0.265)*	0.01 (0.197)
Mean (SE) Range	1.53-5.67	1.42-5.15	
90% 2-sided CI for ratio	3.00-4.19	3.132, 4.077	
90% 2-sided CI for difference			-0.337, 0.363

CI=confidence interval

* $p < 0.0001$ (one sided one sample t-test for $H_0: \mu < 1$ vs $H_1: \mu \geq 1$).

†Difference is defined as week 5 ratio minus baseline ratio

Table 57: Treatment Effect on CSF CD4+ and CD8+ Lymphocyte Count (Evaluable Population, n=13)

	Baseline	Week 5
CD4+ as % of Lymphocytes, mean (SD)	75.160 (7.3831)	74.215 (6.3732)
CD8+ as % of	22.272 (5.4320)	22.007 (6.1624)

Lymphocytes, mean (SD)		
------------------------	--	--

Table 58: Peripheral Blood Memory T Lymphocytes (RO+) Counts
(Evaluable Population, n=13)

	Baseline	Week 5
	Mean (SD)	Mean (SD)
CD4+CD45RO+	27.85 (4.98)	27.06 (5.02)
CD8+CD45RO+(%)	11.24 (3.40)	10.78 (2.98)

5 Summary

Vedolizumab did not affect CSF CD4+ and CD8+ cell counts or CD4+:CD8+ ratio in healthy volunteers after a single 450 mg dose. None of the subjects had a reduction in the post-dose CSF CD4+:CD8+ ratio to less than 1. Vedolizumab was not detected in CSF. In addition, there was no change observed in the total WBCs or memory T lymphocyte CD4+ and CD8+ subsets in peripheral blood. Saturation of the target ($\alpha 4\beta 7$) in blood occurred in all subjects at the time of endpoint assessment. The CSF CD4+ and CD8+ lymphocyte levels and ratio were similar to those previously reported in the literature.

10 These results are consistent with vedolizumab's lack of effect on both
15 physiologic CNS immune surveillance and pathologic CNS inflammation of monkeys.

20 While this invention has been particularly shown and described with
references to preferred embodiments thereof, it will be understood by those skilled
in the art that various changes in form and details may be made therein without
departing from the scope of the invention encompassed by the appended claims.

25

Table 59: Sequences

SEQ ID NO:	Sequence Shown	Description
1	FIG. 1	DNA encoding heavy chain of humanized anti- $\alpha 4\beta 7$ immunoglobulin
2	FIG. 1	Amino acid sequence of heavy chain of humanized anti- $\alpha 4\beta 7$ immunoglobulin
3	FIG. 2	DNA encoding the light chain of humanized anti- $\alpha 4\beta 7$ immunoglobulin
4	FIG. 2	Amino acid sequence of light chain of humanized anti- $\alpha 4\beta 7$ immunoglobulin
5	FIG. 3	Mature humanized light chain of LDP-02
6	FIG. 4	Generic human kappa light chain constant region
7	FIG. 4	Generic murine kappa light chain constant region
8	Referenced on page 31 SYWMH	CDR1 of heavy chain mouse ACT-1 antibody

9	Referenced on page 31 EIDPSESNTNYNQKFKG	CDR2 of heavy chain mouse ACT-1 antibody
10	Referenced on page 31 GGYDGWDY AidY	CDR3 of heavy chain mouse ACT-1 antibody
11	Referenced on page 31 RSSQSLAKSYGNTYLS	CDR1 of light chain mouse ACT-1 antibody
12	Referenced on page 31 GISNRFS	CDR2 of light chain mouse ACT-1 antibody
13	Referenced on page 31 LQGTHQPYT	CDR3 of light chain mouse ACT-1 antibody
14	FIG. 7	human GM607 CL antibody kappa light chain variable region
15	FIG. 7	Human 21/28 CL antibody heavy chain variable region

CLAIMS:

1. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating a human patient having inflammatory bowel disease, wherein the medicament is for administration to the patient according to the following dosing regimen:

(a) initial doses of 165 mg of the humanized anti- $\alpha 4\beta 7$ antibody as a subcutaneous injection every other day for six doses;

(b) followed by a seventh and subsequent doses of 165 mg of the humanized anti- $\alpha 4\beta 7$ antibody as a subcutaneous injection every two weeks;

wherein the dosing regimen is for inducing a clinical response, clinical remission, or both, in the inflammatory bowel disease of the patient;

further wherein the humanized anti- $\alpha 4\beta 7$ antibody comprises an antigen binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized anti- $\alpha 4\beta 7$ antibody has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises the CDRs:

Light chain: CDR1 SEQ ID NO:11,
CDR2 SEQ ID NO:12, and
CDR3 SEQ ID NO:13;

Heavy chain: CDR1 SEQ ID NO:8,
CDR2 SEQ ID NO:9, and
CDR3 SEQ ID NO:10.

2. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating a human patient having inflammatory bowel disease, wherein the medicament is for administration to the patient according to the following dosing regimen:

(a) initial doses of 165 mg of the humanized anti- $\alpha 4\beta 7$ antibody as a subcutaneous injection every other day for six doses;

(b) followed by a seventh and subsequent doses of 165 mg of the humanized anti- $\alpha 4\beta 7$ antibody as subcutaneous injections every four weeks;

wherein the dosing regimen is for inducing a clinical response, clinical remission, or both, in the inflammatory bowel disease of the patient;

further wherein the humanized anti- $\alpha 4\beta 7$ antibody comprises an antigen binding region of nonhuman origin and at least a portion of an antibody of human origin, wherein the humanized anti- $\alpha 4\beta 7$ antibody has binding specificity for the $\alpha 4\beta 7$ complex, wherein the antigen-binding region comprises the CDRs:

Light chain: CDR1 SEQ ID NO:11,
CDR2 SEQ ID NO:12, and
CDR3 SEQ ID NO:13;

Heavy chain: CDR1 SEQ ID NO:8,
CDR2 SEQ ID NO:9, and
CDR3 SEQ ID NO:10.

3. The use of claim 1 or claim 2, wherein the patient having inflammatory bowel disease had a lack of an adequate response with, loss of response to, or was intolerant to treatment with at least one of an immunomodulator, a tumor necrosis factor-alpha antagonist or combinations thereof.

4. The use of any one of claims 1 to 3, wherein the inflammatory bowel disease is pouchitis, Crohn's disease or ulcerative colitis.

5. The use of claim 4, wherein the inflammatory bowel disease is ulcerative colitis.

6. The use of claim 5, wherein the inflammatory bowel disease is moderate to severely active ulcerative colitis.

7. The use of claim 6, wherein the dosing regimen results in mucosal healing in patients suffering from moderate to severely active ulcerative colitis.
8. The use of any one of claims 1 to 7, wherein the dosing regimen results in a reduction, elimination or reduction and elimination of corticosteroid use by the patient.
9. The use of any one of claims 1 to 7, wherein the patient previously received treatment with at least one corticosteroid for the inflammatory bowel disease.
10. The use of any one of claims 1 to 9, wherein the dosing regimen does not alter the ratio of CD4 to CD8 in cerebrospinal fluid of patients receiving the treatment.
11. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient subcutaneously every two weeks at a dose of 108 mg of the antibody, and wherein the humanized anti- $\alpha 4\beta 7$ antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.
12. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient according to the following dosing regimen:

a first intravenous dose of 300 mg of the antibody, a second intravenous dose of 300 mg of the antibody for administration to the human patient two weeks after the first intravenous dose, a third intravenous dose of 300 mg of the antibody for administration to the human patient six weeks after the first intravenous dose, and

fourth and subsequent doses of 108 mg of the antibody for subcutaneous administration to the human patient beginning ten weeks after the first dose, and every two weeks thereafter,

wherein the antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

13. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient according to the following dosing regimen:

a first intravenous dose of 300 mg of the antibody, a second intravenous dose of 300 mg of the antibody for administration to the human patient two weeks after the first intravenous dose, a third intravenous dose of 300 mg of the antibody for administration to the human patient six weeks after the first intravenous dose, and

fourth and subsequent doses of 108 mg of the antibody for subcutaneous administration to the human patient beginning fourteen weeks after the first intravenous dose, and every two weeks thereafter,

wherein the antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

14. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient according to the following dosing regimen:

a first subcutaneous dose of 432 mg of the antibody,

a second subcutaneous dose of 432 mg of the antibody for administration to the human patient two weeks after the first subcutaneous dose, and

third and subsequent subcutaneous doses of 108 mg of the antibody beginning six weeks after the first subcutaneous dose, and every two weeks thereafter,

wherein the antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

15. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient according to the following dosing regimen:

a first subcutaneous dose of 216 mg of the antibody,

a second subcutaneous dose of 216 mg of the antibody for administration to the human patient two weeks after the first subcutaneous dose, and

third and subsequent subcutaneous doses of 108 mg of the antibody beginning six weeks after the first subcutaneous dose, and every two weeks thereafter,

wherein the antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

16. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the medicament is for administration to the human patient according to the following dosing regimen:

a subcutaneous dose of 160 mg of the antibody every other day for two weeks, followed by a subcutaneous dose of 108 mg of the antibody beginning six weeks after the first subcutaneous dose, and every two weeks thereafter,

wherein the antibody comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

17. Use of a humanized anti- $\alpha 4\beta 7$ antibody in the manufacture of a medicament for treating inflammatory bowel disease in a human patient, wherein the antibody is for administration according to a dosing regimen comprising a first intravenous dose of 300 mg of the antibody, a second intravenous dose of 300 mg of the antibody for administration to the human patient two weeks after the first intravenous dose, and third and subsequent doses of 108 mg of the antibody for subcutaneous administration to the human patient beginning six weeks after the first intravenous dose and every two weeks thereafter, wherein the antibody comprises a heavy chain variable region comprising a CDR1 as set forth in SEQ ID NO:8, a CDR2 as set forth in SEQ ID NO:9, and a CDR3 as set forth in SEQ ID NO:10, and comprises a light chain variable region comprising a CDR1 as set forth in SEQ ID NO:11, a CDR2 as set forth in SEQ ID NO:12, and a CDR3 as set forth in SEQ ID NO:13.

18. The use according to any one of claims 11-17, wherein the inflammatory bowel disease is Crohn's disease, ulcerative colitis, pouchitis, or celiac disease.

19. The use according to claim 18, wherein the ulcerative colitis is moderate to severely active ulcerative colitis.

20. The use according to claim 18 or 19, wherein the use results in mucosal healing in the human patient having ulcerative colitis.

21. The use according to claim 18, wherein the Crohn's disease is moderately to severely active Crohn's disease.
22. The use according to any one of claims 11-21, where the human patient previously received treatment with at least one corticosteroid for the inflammatory bowel disease.
23. The use according to any one of claims 11-22, wherein the human patient having inflammatory bowel disease had a lack of an adequate response to, loss of response to, or was intolerant to treatment with at least one of an immunomodulator, a corticosteroid, or a tumor necrosis factor-alpha antagonist.
24. The use according to any one of claims 11-23, wherein the use results in a reduction, elimination or reduction and elimination of corticosteroid use in the human patient having inflammatory bowel disease.
25. The use according to any one of claims 1-24, wherein the human patient having inflammatory bowel disease had an inadequate response to, loss of response to, or was intolerant to either conventional therapy or infliximab.
26. The use according to any one of claims 11-22, wherein the human patient having inflammatory bowel disease had an inadequate response to, loss of response to, or was intolerant to an immunomodulator or a tumor necrosis factor-alpha antagonist.
27. The use according to any one of claims 11-22, wherein the human patient having inflammatory bowel disease had an inadequate response to, intolerance to, or demonstrated dependence on corticosteroids.
28. The use according to any one of claims 1-27, wherein the antibody comprises a heavy chain variable region comprising amino acids 20-140 of SEQ ID NO: 2 and a light chain variable region comprising amino acids 20-131 of SEQ ID NO:4.
29. The use according to any one of claims 1-27, wherein the antibody is an IgG1 isotype.

30. The use according to any one of claims 1-27, wherein the antibody comprises a heavy chain comprising amino acids 20 to 470 of SEQ ID NO:2, and a light chain comprising amino acids 20 to 238 of SEQ ID NO:4.
31. The use according to any one of claims 1-27, wherein the antibody is vedolizumab.
32. The use according to any one of claims 1-31, wherein the medicament for subcutaneous administration is for self-administration.
33. The use according to any one of claims 1-32, wherein the medicament for subcutaneous administration is in a vial.
34. The use according to any one of claims 1-32, wherein the medicament for subcutaneous administration is in a cartridge.
35. The use according to any one of claims 1-32, wherein the medicament for subcutaneous administration is in a syringe.
36. The use according to claim 35, wherein the syringe comprises a 25 gauge, a 26 gauge, a 27 gauge, a 28 gauge, a 29 gauge or a 30 gauge needle.
37. The use according to any one of claims 1-32, wherein the medicament for subcutaneous administration is in an autoinjector.
38. The use according to any one of claims 1-37, wherein the medicament for subcutaneous administration is in a volume of 1 mL.
39. The use according to any one of claims 1-37, wherein the medicament for subcutaneous administration is in a volume of 2 mL.
40. The use according to any one of claims 1-37, wherein the medicament for subcutaneous administration is in a volume of 1.5 mL.

41. The use according to any one of claims 1-40, wherein the medicament for intravenous administration is in a vial.

FIG. 1A

New LDP02 Heavy DNA -- contains cloning sites (lower case),Kozak sequence (upper case) and Leader (lower case)

```

gaattctcgatcgcgatCTCACCatgggatggagctgtatcatcctcttcttggtagcaacagctacaggtgtccactcccag
gtgCAATTGGTGCAGTCTGGGGCTGAGGTTAAGAAAGCCTGGGGCTTCAGTGAAG
GGTGCTCTGCAAGGGTCTGGGTACACCTTCAACCAGCTACTGGATGCATTGGG
TGAGGCAGGCGCCTGGCCAAACGTCTAGAGTGGATCGGAGAGATTGATCCTTC
TGAGAGTAATACTAATACTAATAATAATAATTCAAGGACGCGTCACATTGACT
GTAGACATTTCGCTAGCACAGCCTACATGGAGCTCTCCAGCCTGAGATCTG
AGGACACTGCGGTCTACTATTGTGCAAGAGGGGGTTACGACGGATGGGACTA
TGCTATTGACTACTGGGTCAAGGCACCTGGTCAACCGTCAGCTCAGCCTCCA
CCAAGGGCCCATCGGTCTTCCCCCTGGCACCTCTCCAAAGAGCACCTCTGGG
GGCACAGCGGCCCTGGGTGCTGCTGTCAAGGACTACTTCCCCGAACCGGTGA
CGGTGTCGTGGAACCTCAGGCGCCCTGACCAAGCGCGTGCACACCTTCCCCGGC
TGTCTACAGTCTCAGGACTCTACTCTCCCTCAGCAGCGTGGTGACCGTGCCCT
CCAGCAGCTTGGCACCCAGACCTACATCTGCAACGTGAATCACAAAGCCCCAG
CAACACCAAGGTGGACAAAGAAAGTTGAGCCCAATCTTGTGACAAAACTCAC
ACATGCCCAACCGTGCCCAAGCACCTGAACCTCGCGGGGCAACCGTCAGTCTTCC
TCTTCCCCCAAAACCCAAAGGACACCTCATGATCTCCCGGACCCCTGAGGTC
ACATGCGTGGTGTGACGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAACT
GGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGG
AGCAGTACAACAGCACGTACCGTGTGGTCAAGCGTCTCACCGTCTCTGCACCA
GGACTGGCTGAATGGC

```

AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGA
 AAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACACACAGGTGTACACCCCT
 GCCCCATCCCGGATGAGCTGACCAAGAACAGGTGAGCTGAGCTGACCTGCCTG
 GTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGAGAGCAATGGGC
 AGCCGGAGAACTACAAGACCAGCCTCCCGTGGTGGACTCCGACGGCTC
 CTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGG
 AACGTCTTCTCATGCTCCGTGATGATGAGGCTCTGCACAACCACTACACGCA
 GAAGAGCCTCTCCCTGTCTCCGGGTAAAtaatctagagca

New LDP02 Heavy Protein (space between VHL, VH and Human IgG1-FcRmut)
 MGWSCILFLVATATGVHS
 QVQLVQSGAEVKKPGASVKVSCKGSGYTFTSYWMHWVRQAPGQRLEWIGEIDP
SESNTNYNQKFGRVTLTVDISASTAYMELSSLRSEDTAVYYCARGGYDGDWDY
AIDYWGQGTLVTVSS
 ASTKGPSVFPPLAPSSKSTSGGTAALGCLVKDYFPEPTVSWNSGALTSGVHTFPA
 VLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCP
 PCPAPELAGAPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV
 EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
 SKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN
 YKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVMHEALHNHYTQKSLSLS
 PGK

FIG. 1B

FIG. 2

New LDP02 Light DNA – contains cloning sites (lower case), Kozak sequence (upper case) and Leader (lower case)

gaattctcgatcgatCTCACCatggatggagctgtatcatctcttcttgtagcaacagctacaggtgtccactccgat
GTAGTGATGACTCAAAAGTCCACTCTCCCTGCCCTGTCACCCCCTGGAGAACCCAGC
TTCTATCTCTTGCAGGTCTAGTCAGAGTCTTGCAAAGAGTTATGGGAACACCT
ATTTGTCTTGGTACCTGCAGAACCTGGCCAGTCTCCACAGCTCCTCATCTAT
GGGATTTCCACACAGATTTTCTGGGTGCCAGACAGGTTCAAGTGGCAGTGGTT
CAGGGACAGATTTCACTCACTCAAGATCTCGCGAGTAGAGGCTGAGGACGTGGG
AGTGATTACTGCTTACAAGGTACACATCAGCCGTACACGTTTCGGACAGGGG
ACCAAGGTGGAGATCAAGCGTACGGTGGCTGCCACCATCTGTCTTCATCTTCCC
GCCATCTGATGAGCAGTTGAAATCTGGAACCTGCCTCTGTGTGTGCCTGCTGA
ATAACTTCTATCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGATAACGCCCT
CCAAATCGGGTAACTCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAG
CACCTACAGCCTCAGCAGCACCTGACCTGAGCAAGCAGACTACGAGAAA
CACAAAGTCTACGCCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCCGTCA
CAAAGAGCTTCAACAGGGGAGAGTGThagttagagcagc

New LDP02 Light Protein (space between VKL, VK and Human C Kappa)

MGWSCILFLVATATGVHS
DVVMTQSP^{LP}SLPVT^{PG}EPASISCRSSQSLAKSYGNTYLSWY^{LQ}KPGQSPQLLIYGI
SNRFSGVVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCLQGTHQPYTFGGTKVEI
K
RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQE
SVTEQDSKDS^{TY}SLSS^{TL}TL^{SK}ADY^EKKH^KVYACEVTHQGLSSPVT^KSFNRGEC

PAIRWISE newMLN02-no sig

4/18

Pairwise Hum kappa-const
MI:Mur kappa-const.txt

```

A 1 rtvaapsvfifppsdeqlksgtasvvc1lnnfypreakvqwkvdnalqsg 50
  | | | | | | | | | | | | | | | | | | | | | | | | | | | |
  | | | . | | | | | | | | | | | | | | | | | | | | | | | |
B 1 radaaptvsifppsseq1tsggasvvcflnnfypkdinvkwidgserqn 50
  .
51 nsqesvteqdskdstyslsst1t1skadyekhkvyacevthqglsspvtk 100
  | | : | | | | | | | | | | | | | | | | | | | | | | | |
51 gvlnswtdqdskdstysmsst1t1tkdeyerhnsytceathktstspivk 100

101 sfnrgec 107
    | | | | | |
101 sfnrnec 107

```




FIG. 5

7/18




FIG. 6





FIG. 7

FIG. 8



Prediction Profiler




FIG. 9

10/18

FIG. 10




FIG. 11

25 mM Citrate, 50 mM Histidine. 125 mM Arginine




FIG. 12




FIG. 13




FIG. 14A

**GM607'Cl antibody kappa light
chain variable region**

SEQ ID NO: 14

```

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
 1           5           10           15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Leu His Ser
          20           25           30
Asn Gly Tyr Asn Tyr Leu Asp Trp Tyr Leu Gln Lys Pro Gly Gln Ser
          35           40           45
Pro Gln Leu Leu Ile Tyr Leu Gly Ser Asn Arg Ala Ser Gly Val Pro
          50           55           60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65           70           75           80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Met Gln Ala
          85           90           95
Leu Gln Thr Pro Gln Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
          100          105          110

```

**21/28'CL antibody heavy
chain variable region**

SEQ ID NO: 15

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	1	5	10	15
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	20	25	30	
Ala	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Arg	Leu	Glu	Trp	Met	35	40	45	
Gly	Trp	Ile	Asn	Ala	Gly	Asn	Gly	Asn	Thr	Lys	Tyr	Ser	Gln	Lys	Phe	50	55	60	
Gln	Gly	Arg	Val	Thr	Ile	Thr	Arg	Asp	Thr	Ser	Ala	Ser	Thr	Ala	Tyr	65	70	75	80
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	85	90	95	
Ala	Arg	Gly	Gly	Tyr	Tyr	Gly	Ser	Gly	Ser	Asn	Tyr	Trp	Gly	Gln	Gly	100	105	110	
Thr	Leu	Val	Thr	Val	Ser	Ser										115			

FIG. 14B

FIG. 15

Components of a Protein Product in a Pre-filled Syringe




FIG. 16





FIG. 17A

Prediction Profiler: Initial Glide Force




FIG. 17B

Prediction Profiler: Initial Glide Force




FIG. 18

