

(19) AUSTRALIAN PATENT OFFICE

- (54) Title
Method of treating corneal transplant rejection by using VEGF antagonists
- (51)⁶ International Patent Classification(s)
C07K 16/24 (2006.01) 31/7088
A61K 31/00 (2006.01) 20060101ALI2005100
A61K 31/7088 ^{8BMEP} **A61K**
(2006.01) 45/06
A61K 45/06 (2006.01) 20060101ALI2005100
C07K 14/71 (2006.01) ^{8BMEP} **C07K**
C07K 19/00 (2006.01) 14/71
C07K 16/24 20060101ALI2005100
20060101AFI2005100 ^{8BMEP} **C07K**
^{8BMEP} **A61K** 19/00
31/00 20060101ALI2005100
20060101ALI2005100 ^{8BMEP}
^{8BMEP} **A61K** PCT/US2004/012540
- (21) Application No: 2004242586 (22) Application Date: 2004 .04 .23
- (87) WIPO No: W004/106378
- (30) Priority Data
- | | | |
|-------------|--------------|--------------|
| (31) Number | (32) Date | (33) Country |
| 60/473,734 | 2003 .05 .28 | US |
| 60/492,865 | 2003 .08 .06 | US |
- (43) Publication Date : 2004 .12 .09
- (71) Applicant(s)
The Schepens Eye Research Institute, Inc., Regeneron Pharmaceuticals, Inc.
- (72) Inventor(s)
Wiegand, Stanley, Cursiefen, Claus, Cao, Jingtai
- (74) Agent/Attorney
Phillips Ormonde Fitzpatrick, 367 Collins Street, Melbourne, VIC, 3000
- (56) Related Art
AU 2003215337 (ALLERGAN, INC.) 09 September 2003 (equivalent to WO 2003/072029)
WO 2000/075319 A1 (REGENERON PHARMACEUTICALS, INC.) 14 December 2000
WO 1999/013909 A1 (TOAGOSEI CO., LTD) 25 March 1999
YATOH S. et al. "Effect of a topically applied neutralizing antibody against vascular endothelial growth factor on corneal allograft rejection of rat." Transplantation, 15 December 1998. vol. 66, no. 11, pages 1519-1524.
LAI, C-M, "Inhibition of corneal neovascularization by recombinant adenovirus mediated anti-sense VEGF RNA." Experimental Eye Research, December 2002. vol. 75, no. 6, pages 625-634.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 December 2004 (09.12.2004)

PCT

(10) International Publication Number
WO 2004/106378 A3

- (51) International Patent Classification⁷: **C07K 16/24**,
A61K 31/7088, C07K 19/00, 14/71, A61P 37/06
- (21) International Application Number:
PCT/US2004/012540
- (22) International Filing Date: 23 April 2004 (23.04.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/473,734 28 May 2003 (28.05.2003) US
60/492,865 6 August 2003 (06.08.2003) US
- (71) Applicants (for all designated States except US): **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US). **THE SCHEPENS EYE RESEARCH INSTITUTE** [US/US]; 20 Staniford Street, Boston, MA 02114 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **WIEGAND, Stanley** [US/US]; 15 Fox Run Road, Croton on Hudson, NY 10520 (US). **CAO, Jingtai** [CN/US]; 308 N. Greeley Avenue, Chappaqua, NY 10514 (US). **CURSIEFFEN, Claus** [DE/DE]; Nordliche Stadtmauerstr. 14, 91054 Erlangen (DE).
- (74) Agent: **VALETA, Gregg**, Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AF, AG, AI, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CI, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- (88) Date of publication of the international search report:
19 May 2005
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2004/106378 A3

(54) Title: METHOD OF TREATING CORNEAL TRANSPLANT REJECTION BY USING VEGF ANTAGONISTS

(57) Abstract: Methods of preventing, reducing, or treating corneal transplant rejection to improve transplant survival in a subject in need thereof comprising administering an agent capable of blocking or inhibiting vascular endothelial growth factor (VEGF) are provided. The methods are useful for inhibiting or preventing corneal transplant rejection in a human subject who is the recipient of a transplanted cornea.

METHOD OF TREATING CORNEAL TRANSPLANT REJECTION**BACKGROUND****Field of the Invention**

[0001] The field of the invention is related to methods of using VEGF antagonists to reduce, prevent, or treat corneal transplant rejection, thus improving long-term transplant survival.

Description of Related Art

[0002] It has previously been reported that topical application of an anti-VEGF neutralizing antibody suppresses acute allograft rejection in a rat corneal transplant model (Yatoh et al. (1998) Transplantation 66(11):1519-24). As the leading cause of human corneal transplant failure is transplant rejection, there is a need for a therapeutic for use in preventing corneal transplant rejection in humans who receive a corneal transplant.

BRIEF SUMMARY OF THE INVENTION

[0003] The invention is based in part on the finding that administration of an agent capable of blocking or inhibiting vascular endothelial growth factor (VEGF) prevents corneal transplant rejection. The experiments, described below, conducted in an animal model of corneal transplantation show that long-term transplant survival is promoted by blocking VEGF-mediated activity.

[0004] In a first aspect, the invention features a method of improving transplant survival in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that transplant survival is improved.

[0004a] In another aspect, the invention features a use of an first agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity in the preparation of a medicament for treating or preventing corneal transplant rejection in a mammalian subject, wherein the agent is VEGFR1R2-FcΔC1(a).

[0005] In specific embodiments, the agent capable of blocking, inhibiting, or ameliorating VEGF-mediated activity is a VEGF antagonist. The VEGF antagonist may be a polypeptide, an antibody, a small molecule, or a nucleic acid. More specifically, the VEGF antagonist includes a VEGF trap selected from the group consisting of acetylated Flt-1(1-3)-Fc, Flt-1(1-3_{R→N})-Fc, Flt-1(1-3_{ΔB})-Fc, Flt-1(2-3_{ΔB})-Fc, Flt-1(2-3)-Fc, Flt-1D2-VEGFR3D3-FcΔC1(a), Flt-1D2-Flk-1D3-FcΔC1(a), and VEGFR1R2-FcΔC1(a). In a specific and preferred embodiment, the VEGF trap is

VEGFR1R2-FcΔC1(a) (also termed VEGF trap_{R1R2}) having the nucleotide sequence set forth in SEQ ID NO: 1 and the amino acid sequence set forth in SEQ ID NO: 2. The invention encompasses the use of a VEGF trap that is at least 90%, 95%, 98%, or at least 99% homologous with the nucleotide sequence set forth in SEQ ID NO: 1 and/or the amino acid sequence set forth in SEQ ID NO:2.

[0006] In other embodiments, the agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity is a nucleic acid-based antagonist capable of interfering with the expression of VEGF. A specific example of this embodiment is one in which the nucleic acid-based antagonist is an aptamer, an siRNA, or an antisense molecule.

[0007] Administration of the agent may be by any method known in the art, including subcutaneous, intramuscular, intradermal, intraperitoneal, intravenous, intranasal, oral, or topical routes of administration. Preferable, administration to the subject in need of the agent is topical administration to the eye or subconjunctival administration. Administration may occur prior to or following corneal transplantation, preferably following surgery. Administration may also include a second agent, such as an immunosuppressive agent.

[0008] The subject to be treated is preferably a human subject who has or will receive a corneal transplant.

[0009] In a related second aspect, the invention features the use of an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity in the preparation of a medicament for improving transplant survival in a mammalian subject.

[0010] In a third aspect, the invention features a method of preventing corneal transplant rejection in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that corneal transplant rejection is prevented.

[0011] In a related fourth aspect, the invention features the use of an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity in the preparation of a medicament for the treatment of corneal transplant rejection in a mammalian subject.

[0012] In a fifth aspect, the invention features a method of reducing the incidence of corneal transplant rejection in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that the incidence of corneal transplant rejection is reduced, wherein the agent is VEGFR1R2-FcΔC1(a).

[0012a] In a further aspect, the invention features a method of treating corneal transplant rejection in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that corneal transplant rejection is treated, wherein the agent is VEGFR1R2-FcΔC1(a).

[0013] In a related sixth aspect, the invention features the use of an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity in the preparation of a medicament for reducing the incidence of corneal transplant rejection in a mammalian subject receiving a corneal transplant.

[0013a] In a further aspect, the invention features a pharmaceutical composition when used for prevention or treatment of corneal transplant rejection, comprising a vascular endothelial growth factor (VEGF) antagonist, and a pharmaceutically acceptable carrier, wherein the antagonist is VEGFR1R2-FcΔC1(a).

[0014] In a seventh aspect, the invention features a pharmaceutical composition comprising a VEGF antagonist, for example the VEGF trap VEGFR1R2-FcΔC1(a), in a pharmaceutically acceptable carrier. Such pharmaceutical compositions may be liquid, gel, ointment, salve, slow release formulations or other formulations suitable for ophthalmic administration.

[0015] In an eighth aspect, the invention features an article of manufacture comprising packaging materials and a pharmaceutical agent contained within the packaging materials, wherein the pharmaceutical agent comprises at least one VEGF-specific fusion protein of the invention, and the packaging material comprises a label or package insert which indicates that the VEGF-specific fusion protein can be used for the treatment or prevention of corneal transplant rejection.

[0015b] In a further aspect, the invention features an article of manufacturing when used for prevention or treatment of corneal transplant rejection comprising:

- (a) packaging material; and
- (b) a pharmaceutical agent contained within the packaging material;

wherein the pharmaceutical agent comprises at least one VEGF-specific fusion protein of the invention and wherein the packaging material comprises a label or package insert which indicates that the VEGF-specific fusion protein can be used to treat or prevent corneal transplant rejection in a mammalian subject, wherein the agent is VEGFR1R2-FcΔC1(a).

[0016] Other aspects will become apparent from a review of the ensuing detailed description.

DETAILED DESCRIPTION

[0017] Before the present methods are described, it is to be understood that this invention is not limited to particular methods, and experimental conditions described, as such methods and conditions may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0018] As used in this specification and the appended claims, the singular forms "a", "an", and "the" include plural references unless the context clearly dictates otherwise. Thus for example, a reference to "a method" includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure and so forth.

[0019] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference in their entirety.

General Description

[0020] Experiments were undertaken to evaluate occurrence and time course of hem- and lymphangiogenesis after normal-risk corneal transplantation and to test whether pharmacologic strategies inhibiting both processes improve long-term graft survival. As described in the experimental section below, normal-risk allogeneic (C57BL/6 to BALB/c) and syngeneic (BALB/c to BALB/c) corneal transplantations were performed and occurrence and time course

of hem- and lymphangiogenesis after keratoplasty was observed using double immunofluorescence of corneal flatmounts (with CD31 as panendothelial and LYVE-1 as lymphatic vascular endothelial specific marker). A molecular trap designed to eliminate VEGF-A ("VEGF Trap_{R1R2}"; 12.5 mg/kg) was tested for its ability to inhibit both processes after keratoplasty and to promote long-term graft survival (intraperitoneal injections on the day of surgery and 3, 7, and 14 days later). The results show that no blood or lymph vessels were detectable immediately after normal-risk transplantation in either donor or host cornea, but hem- and lymphangiogenesis were clearly visible at day 3 after transplantation. Both vessel types reached donor tissue at one week after allo- and similarly after syngeneic grafting. Early postoperative trapping of VEGF-A significantly reduced both hem- and lymphangiogenesis and significantly improved long-term graft survival (78% versus 40%; $p < 0.05$). There is concurrent, VEGF-A-dependent hem- and lymphangiogenesis after normal-risk keratoplasty within the preoperatively avascular recipient bed. Inhibition of hem- and lymphangiogenesis (which mediate the efferent and afferent arms of an immune response) after normal-risk corneal transplantation improves long-term graft survival, establishing that early postoperative hem- and lymphangiogenesis are risk factors for graft rejection even in low-risk eyes.

Definitions

[0021] By the term "therapeutically effective dose" is meant a dose that produces the desired effect for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, for example, Lloyd (1999) *The Art, Science and Technology of Pharmaceutical Compounding*).

[0022] By the term "blocker", "inhibitor", or "antagonist" is meant a substance that retards or prevents a chemical or physiological reaction or response. Common blockers or inhibitors include but are not limited to antisense molecules, antibodies, antagonists and their derivatives. More specifically, an example of a VEGF blocker or inhibitor is a VEGF receptor-based antagonist including, for example, an anti-VEGF antibody, or a VEGF trap such as VEGFR1R2-FcΔC1(a) (SEQ ID NOs:1-2). For a complete description of VEGF-receptor based antagonists including VEGFR1R2-FcΔC1(a), see PCT publication WO/00/75319, the contents of which is incorporated in its entirety herein by reference.

[0023] A "small molecule" is defined herein to have a molecular weight below about 500 Daltons, and may include chemical as well as peptide molecules.

VEGF Antagonists

[0024] In one aspect of the invention, VEGF-mediated activity is blocked or inhibited by the use

of VEGF receptor-based blockers of VEGF-mediated activity. A non-limiting example of a VEGF receptor-based blocker includes, but is not limited to, VEGFR1R2-FcAC1(a). Other suitable receptor-based blockers include acetylated Flt-1(1-3)-Fc, Flt-1(1-3_{R→N})-Fc, Flt-1(1-3_{ΔB})-Fc, Flt-1(2-3_{ΔB})-Fc, Flt-1(2-3)-Fc, Flt-1D2-VEGFR3D3-FcAC1(a), Flt-1D2-Flk-1D3-FcAC1(a). For a complete description of these and other VEGF-receptor-based blockers, including pegylated receptor-based blockers, see PCT Publication No. WO/00/75319, the contents of which is incorporated in its entirety herein by reference.

[0025] In addition to the VEGF receptor-based blockers described in PCT Publication No. WO/00/75319, variants and derivatives of such VEGF receptor-based blockers are also contemplated by the invention. The sequence of the variants or derivatives may differ by a change which is one or more additions, insertions, deletions and/or substitutions of one or more nucleotides of the sequence set forth in SEQ ID NO:1. Changes to a nucleotide sequence may result in an amino acid change at the protein level, or not, as determined by the genetic code. Thus, nucleic acid according to the present invention may include a sequence different from the sequence shown in SEQ ID NO:1, yet encode a polypeptide with the same amino acid sequence as SEQ ID NO: 2. On the other hand, the encoded polypeptide may comprise an amino acid sequence which differs by one or more amino acid residues from the amino acid sequence shown in SEQ ID NO:2. Nucleic acid encoding a polypeptide which is an amino acid sequence variant or derivative of the sequence shown in SEQ ID NO:2 is further provided by the present invention. Nucleic acid encoding such a polypeptide may show at the nucleotide sequence and/or encoded amino acid level greater than about 90%, 95%, 98%, or 99% homology with the coding sequence shown in SEQ ID NO:1 and/or the amino acid sequence shown in SEQ ID NO:2. For amino acid "homology", this may be understood to be similarity (according to the established principles of amino acid similarity, e.g. as determined using the algorithm GAP (Genetics Computer Group, Madison, Wis.)) or identity. GAP uses the Needleman and Wunsch algorithm to align two complete sequences that maximizes the number of matches and minimizes the number of gaps. Generally, the default parameters are used, with a gap creation penalty=12 and gap extension penalty=4.

[0026] Individual components of the VEGF-specific fusion proteins of the invention may be constructed by molecular biological methods known to the art with the instructions provided by the instant specification. These components are selected from a first cellular receptor protein, such as, for example, VEGFR1; a second cellular receptor protein, such as, for example, VEGFR2; a multimerizing component, such as an Fc.

[0027] Specific embodiments of the VEGF-specific fusion proteins useful in the methods of the invention comprise a multimerizing component which allows the fusion proteins to associate,

e.g., as multimers, preferably dimers. Preferably, the multimerizing component comprises an immunoglobulin derived domain. Suitable multimerizing components are sequences encoding an immunoglobulin heavy chain hinge region (Takahashi et al. 1982 Cell 29:671-679); immunoglobulin gene sequences, and portions thereof.

[0028] The nucleic acid constructs encoding the fusion proteins useful in the methods of the invention are inserted into an expression vector by methods known to the art, wherein the nucleic acid molecule is operatively linked to an expression control sequence. Host-vector systems for the production of proteins comprising an expression vector introduced into a host cell suitable for expression of the protein are known in the art. The suitable host cell may be a bacterial cell such as *E. coli*, a yeast cell, such as *Pichia pastoris*, an insect cell, such as *Spodoptera frugiperda*, or a mammalian cell, such as a COS, CHO, 293, BHK or NS0 cell.

Antisense Nucleic Acids

[0029] In one aspect of the invention, VEGF-mediated activity is blocked or inhibited by the use of VEGF antisense nucleic acids. The present invention provides the therapeutic or prophylactic use of nucleic acids comprising at least six nucleotides that are antisense to a gene or cDNA encoding VEGF or a portion thereof. As used herein, a VEGF "antisense" nucleic acid refers to a nucleic acid capable of hybridizing by virtue of some sequence complementarity to a portion of an RNA (preferably mRNA) encoding VEGF. The antisense nucleic acid may be complementary to a coding and/or noncoding region of an mRNA encoding VEGF. Such antisense nucleic acids have utility as compounds that prevent VEGF expression, and can be used in the treatment or prevention of corneal transplant rejection. The antisense nucleic acids of the invention are double-stranded or single-stranded oligonucleotides, RNA or DNA or a modification or derivative thereof, and can be directly administered to a cell or produced intracellularly by transcription of exogenous, introduced sequences.

[0028] The VEGF antisense nucleic acids are of at least six nucleotides and are preferably oligonucleotides ranging from 6 to about 50 oligonucleotides. In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 15 nucleotides, at least 100 nucleotides, or at least 200 nucleotides. The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof and can be single-stranded or double-stranded. In addition, the antisense molecules may be polymers that are nucleic acid mimics, such as PNA, morpholino oligos, and LNA. Other types of antisense molecules include short double-stranded RNAs, known as siRNAs, and short hairpin RNAs, and long dsRNA (>50 bp but usually ≥ 500 bp).

Short interfering RNAs

[0029] In another embodiment, VEGF-mediated activity is blocked by blocking VEGF expression. One method for inhibiting VEGF expression is the use of short interfering RNA (siRNA) through RNA interference (RNAi) or post-transcriptional gene silencing (PTGS) (see, for example, Ketting et al. (2001) *Genes Develop.* 15:2654-2659). siRNA molecules can target homologous mRNA molecules for destruction by cleaving the mRNA molecule within the region spanned by the siRNA molecule. Accordingly, siRNAs capable of targeting and cleaving homologous VEGF mRNA are useful for treating, reducing or preventing corneal transplant rejection.

Inhibitory Ribozymes

[0030] In aspect of the invention, corneal transplant rejection may be treated or prevented in a subject suffering from such disease by decreasing the level of VEGF activity by using ribozyme molecules designed to catalytically cleave gene mRNA transcripts encoding VEGF, preventing translation of target gene mRNA and, therefore, expression of the gene product.

[0031] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see, e.g., U.S. Patent No. 5,093,246. While ribozymes that cleave mRNA at site-specific recognition sequences can be used to destroy mRNAs encoding VEGF, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA has the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art. The ribozymes of the present invention also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one that occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA). The Cech-type ribozymes have an eight base pair active site that hybridizes to a target RNA sequence where after cleavage of the target RNA takes place. The invention encompasses those Cech-type ribozymes that target eight base-pair active site sequences that are present in the gene encoding VEGF.

Generation of Antibodies to VEGF Proteins

[0032] In another aspect of the invention, the invention may be practiced with an anti-VEGF

antibody or antibody fragment capable of binding and blocking VEGF activity. Anti-VEGF antibodies are disclosed, for example, in US Patent No. 6,121,230, herein specifically incorporated by reference. The term "antibody" as used herein refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant regions, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD, and IgE, respectively. Within each IgG class, there are different isotypes (eg. IgG₁, IgG₂, etc.). Typically, the antigen-binding region of an antibody will be the most critical in determining specificity and affinity of binding.

[0033] Antibodies exist as intact immunoglobulins, or as a number of well-characterized fragments produced by digestion with various peptidases. For example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce F(ab)₂, a dimer of Fab which itself is a light chain joined to V_H-C_H1 by a disulfide bond. The F(ab)₂ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the F(ab)₂ dimer into an Fab' monomer. The Fab' monomer is essentially Fab with part of the hinge region. While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized *de novo* either chemically or by using recombinant DNA methodology. Thus, the terms antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies, or those synthesized *de novo* using recombinant DNA methodologies (e.g., single chain Fv)(scFv) or those identified using phase display libraries (see, for example, McCafferty et al. (1990) Nature 348:552-554).

[0034] Methods for preparing antibodies are known to the art. See, for example, Kohler & Milstein (1975) Nature 256:495-497; Harlow & Lane (1988) Antibodies: a Laboratory Manual, Cold Spring Harbor Lab., Cold Spring Harbor, NY). The genes encoding the heavy and light chains of an antibody of interest can be cloned from a cell, e.g., the genes encoding a monoclonal antibody can be cloned from a hybridoma and used to produce a recombinant monoclonal antibody. Gene libraries encoding heavy and light chains of monoclonal antibodies can also be made from hybridoma or plasma cells. Random combinations of the heavy and light chain gene products generate a large pool of antibodies with different antigenic specificity. Techniques for the production of single chain antibodies or recombinant antibodies (US 4,946,778; US 4,816,567) can be adapted to produce antibodies used in the fusion proteins and methods of the instant invention. Also, transgenic mice, or other organisms such as other mammals, may be

used to express human or humanized antibodies. Alternatively, phage display technology can be used to identify antibodies and heteromeric Fab fragments that specifically bind to selected antigens.

Antibody Screening and Selection

[0035] Screening and selection of preferred antibodies can be conducted by a variety of methods known to the art. Initial screening for the presence of monoclonal antibodies specific to a target antigen may be conducted through the use of ELISA-based methods, for example. A secondary screen is preferably conducted to identify and select a desired monoclonal antibody for use in construction of the multi-specific fusion proteins of the invention. Secondary screening may be conducted with any suitable method known to the art. One preferred method, termed "Biosensor Modification-Assisted Profiling" ("BiaMAP") is described in co-pending USSN 60/423,017 filed 01 Nov 2002, herein specifically incorporated by reference in its entirety. BiaMAP allows rapid identification of hybridoma clones producing monoclonal antibodies with desired characteristics. More specifically, monoclonal antibodies are sorted into distinct epitope-related groups based on evaluation of antibody:antigen interactions.

Treatment Population

[0036] A suitable subject for treatment by the method of the invention is a human who has received or will receive a corneal transplant. Corneal transplantation is the oldest, most successful and most commonly performed tissue transplantation, with nearly 40,000 transplantations a year alone in the US. When corneal grafts are placed into an avascular recipient bed (so-called normal-risk keratoplasty), 2-year graft survival rates approach 90% under cover of topical steroids, even without HLA-matching. This very successful outcome is attributed to corneal immune privilege, i.e. the phenomenon of suppressed corneal inflammation induced by an array of endogenous mechanisms downregulating alloimmune and inflammatory responses in the cornea and its bed. These mechanisms include the lack of both afferent lymphatic and efferent blood vessels in the normal-risk recipient cornea, lack of MHC II⁺ antigen presenting cells (APCs), FASL-expression on corneal epithelium and endothelium, and the anterior chamber associated immune privilege (ACAID) directed at graft antigens etc. (Streilein et al. (1999) Transplant Proc. 31:1472-1475).

[0037] In contrast, survival rates of cornea grafts placed into vascularized, not immune-privileged recipient beds (so called high-risk keratoplasty) drop significantly to below 50% (even with local and systemic immune suppression). Pre-existing corneal stromal blood vessels have been identified as strong risk factors for immune rejection after corneal transplantation, both in

the clinical setting as well as in the well-defined mouse model of corneal transplantation (Sano et al. (1995) Invest. Ophthalmol. Vis. Sci. 36:2176-85). Recently, in addition to blood vessels, biomicroscopically undetectable lymphatic vessels have been found in association with blood vessels in vascularized high-risk human corneas (Cursiefen et al. (2003) Cornea. 22:273-81) and it is likely that corneal lymphatic vessels enable effective access of donor and host APCs and antigenic material to regional lymph nodes where accelerated sensitisation to graft antigens occurs (Liu et al. (2002) J. Exp. Med. 195:259-68) even in the normal-risk setting (with a preoperatively avascular recipient bed), where mild corneal hemangiogenesis develops after keratoplasty. Outgrowth of new blood vessels from the limbal arcade towards the graft can be observed within the first postoperative year in about 50% of patients undergoing normal-risk keratoplasty, and in 10% of patients these new blood vessels even reach the interface or invade donor tissue (Cursiefen et al. (2001) Graefes Arch. clin. Exp. Ophthalmol. 39:514-21) at corneal suture sites, and then proceed centrally.

Methods of Administration

[0038] The invention provides methods of treatment comprising administering to a subject an effective amount of an agent of the invention. In a preferred aspect, the agent is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, e.g., such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

[0039] Various delivery systems are known and can be used to administer an active agent of the invention, e.g., delivery systems suitable for topical administration, preferably topical administration directly to the eye, or subconjunctival administration, as well as other delivery systems such as those that utilize encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, 1987, J. Biol. Chem. 262:4429-4432), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction are preferably topical or subconjunctival, but may be enteral or parenteral including but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, and oral routes. The active agents may be administered by any convenient route, for example by absorption through epithelial (e.g. topical administration to the eye) or mucocutaneous linings (e.g., oral mucosa, intestinal mucosa, etc.) or infusion or bolus injection, and may be administered together with other biologically active agents. Administration can be systemic or local. Administration can be acute or chronic (e.g. daily, weekly, monthly, etc.) or in combination or alteration with other agents. Pulmonary

administration can also be employed, *e.g.*, by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[0040] In another embodiment, the active agent can be delivered in a vesicle, in particular a liposome (see Langer (1990) *Science* 249:1527-1533). In yet another embodiment, the active agent can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer (1990) *supra*). In another embodiment, polymeric materials can be used (see Howard et al. (1989) *J. Neurosurg.* 71:105). In another embodiment where the active agent of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, *e.g.*, by use of a retroviral vector (see, for example, U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (*e.g.*, a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see *e.g.*, Joliet et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:1864-1868), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

[0041] In a specific embodiment, it may be desirable to administer the pharmaceutical compositions of the invention locally to the area in need of treatment; this may be achieved, for example, and not by way of limitation, by topical administration, subconjunctival administration, local infusion during surgery, *e.g.*, by injection, by means of a catheter, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, fibers, or commercial skin substitutes.

Cellular Transfection and Gene Therapy

[0042] The present invention encompasses the use of nucleic acids encoding the VEGF-specific fusion proteins of the invention for transfection of cells *in vitro* and *in vivo*. These nucleic acids can be inserted into any of a number of well-known vectors for transfection of target cells and organisms. The nucleic acids are transfected into cells *ex vivo* and *in vivo*, through the interaction of the vector and the target cell. Reintroduction of transfected cells may be accomplished by any method known to the art, including re-implantation of encapsulated cells. The compositions are administered (*e.g.*, by injection into a muscle) to a subject in an amount sufficient to elicit a therapeutic response. An amount adequate to accomplish this is defined as "a therapeutically effective dose or amount."

[0043] In another aspect, the invention provides a method of treating or preventing corneal transplant rejection in a human comprising transfecting a cell with a nucleic acid encoding a

VEGF-specific fusion protein of the invention, wherein the nucleic acid comprises an inducible promoter operably linked to the nucleic acid encoding the VEGF-specific fusion protein. For gene therapy procedures in the treatment or prevention of human disease, see for example, Van Brunt (1998) *Biotechnology* 6:1149-1154.

Pharmaceutical Compositions

[0044] Pharmaceutical compositions useful in the practice of the method of the invention include a therapeutically effective amount of an active agent, and a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly, in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin.

[0045] In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous, subcutaneous, or intramuscular administration to human beings. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lidocaine to ease pain at the site of the injection. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0046] The active agents of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with free amino groups such as those

derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with free carboxyl groups such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[0047] The amount of the active agent of the invention that will be effective in the treatment or prevention of corneal transplant rejection can be determined by standard clinical techniques based on the present description. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the condition, and should be decided according to the judgment of the practitioner and each subject's circumstances. However, suitable dosage ranges for intravenous administration are generally about 50-5000 micrograms of active compound per kilogram body weight. Suitable dosage ranges for intranasal administration are generally about 0.01 pg/kg body weight to 1 mg/kg body weight. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0048] For systemic administration, a therapeutically effective dose can be estimated initially from *in vitro* assays. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the IC_{50} as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Initial dosages can also be estimated from *in vivo* data, e.g., animal models, using techniques that are well known in the art. One having ordinary skill in the art could readily optimize administration to humans based on animal data.

[0049] Dosage amount and interval may be adjusted individually to provide plasma levels of the compounds that are sufficient to maintain therapeutic effect. One having skill in the art will be able to optimize therapeutically effective local dosages without undue experimentation.

[0050] The amount of compound administered will, of course, be dependent on the subject being treated, on the subject's weight, the severity of the affliction, the manner of administration, and the judgment of the prescribing physician. The therapy may be repeated intermittently while symptoms are detectable or even when they are not detectable. The therapy may be provided alone or in combination with other drugs.

Combination Therapies

[0051] In numerous embodiments, the VEGF blockers of the present invention may be administered in combination with one or more additional compounds or therapies or medical procedures. For example, suitable therapeutic agents for use in combination, either alternating or simultaneously, with the VEGF blockers may include topically administered immunosuppressive

agents such as corticosteroids, dexamethasone, cyclosporin A, or anti-metabolic agents or systemically administered immunosuppressive agents such as corticosteroids, dexamethasone, cyclosporin A, FK506, or anti-metabolic agents, as well as other agents effective to treat, reduce, or prevent corneal transplant rejection (see Barker, NH, *et al.*, (2000) Clin Exp Ophthalmol 28:357-360). Other suitable therapeutic agents for use in combination, either alternating or simultaneously, with the VEGF blockers of the subject invention may include blockers that can block other VEGF family members such as VEGF-C and VEGF-D.

Kits

[0052] The invention also provides an article of manufacturing comprising packaging material and a pharmaceutical agent contained within the packaging material, wherein the pharmaceutical agent comprises at least one VEGF-specific fusion protein of the invention and wherein the packaging material comprises a label or package insert which indicates that the VEGF-specific fusion protein can be used for treating corneal transplant rejection.

[0053] Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

[0054] The following example is put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the methods and compositions of the invention, and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1: Inhibition of corneal lymphangiogenesis and angiogenesis after low-risk keratoplasty using VEGFR1R2-FcΔC1(a).

[0055] **Mice and anesthesia.** Six to 8 weeks old male C57BL/6 mice were used as donors and same-aged male BALB/c mice (Taconic, Germantown, NY) as recipients in the mouse model of normal-risk keratoplasty (Sonoda *et al.* (1992) Transplantation 54:694-704). For syngeneic transplantations, 6-8 weeks old male BALB/c mice were used both as donors as well as recipients. For the dose response studies, 8 weeks old male C57BL/6 mice were used. All animals were treated in accordance with the ARVO Statement for the Use of Animals in

Ophthalmic and Vision Research. Mice were anesthetized using a mixture of ketamine and xylazine (120 mg/kg body weight and 20 mg/kg body weight respectively).

[0056] Dose response of VEGF Trap_{R1R2}. Five different doses of VEGF-Trap_{R1R2} (SEQ ID NO:2) were tested in mice that received three interrupted intrastromal sutures (10-0 nylon, 50- μ m-diameter, Sharpoint, Surgical Specialties Corporation, Reading, PA). Gentamicine and ophthalmic ointment were applied immediately after surgery. Following surgery (day 0), mice received a single subcutaneous injection of VEGF Trap_{R1R2} (25 mg/kg, 12.5 mg/kg, 6.25 mg/kg, 2.5 mg/kg or 0.5 mg) or human Fc (12.5 mg/kg; control). Corneas were harvested on day 9 after suture placement, following an intravenous administration of an endothelial-specific fluorescein-conjugated lectin (*Lycopersicon esculentum*, Vector Laboratories, Burlingame, CA). The isolated corneas were flat-mounted on glass slides, and images of lectin-labeled vessels were captured using a Spot RT Digital camera (Diagnostic Instrument, Inc. Sterling Heights, MI) attached to a Nikon Microphot-FXA microscope (Nikon Inc. Garden City, NY). Scion Image 1.62c (Scion Corporation, Frederick, MD) was used to quantify the extent of corneal neovascularization.

[0057] Corneal transplantation in mice. Orthotopic corneal allografting in the mouse model of normal-risk keratoplasty was performed as described previously (Sonoda et al. (1992) *supra*). Donor corneas were excised by trephination using a 2.0 mm bore and cut with a curved vannas scissor. Until grafting, corneal tissue was placed in chilled phosphate-buffered saline. Recipients were anesthetized and the graft bed was prepared by trephining a 1.5 mm site in the central cornea of the right eye and discarding the excised cornea. The donor cornea was immediately applied to the bed and secured in place with 8 interrupted sutures (11-0 nylon, 70 μ m diameter needles, Arosurgical, Newport Beach, CA). Antibiotic ointment (Oxymycin, Pharmafair, Hauppauge, NY) was placed on the corneal surface and the eyelids sutured with 8-0 suture (Sharpoint, Reading, PA). Recipients of grafts in which bleeding developed in the immediate postoperative period were discarded from further evaluation. All grafted eyes were examined after 72 hours, and grafts with technical difficulties (hyphema, cataract, infection, loss of anterior chamber) were excluded from further consideration. Tarsorrhaphy and corneal sutures were removed after 7 days and grafts were then examined at least twice a week until week 8 post transplantation by slit-lamp microscopy and scored for opacity. The survival experiment was performed twice and comprised 10 and 12 mice per experiment in both groups, respectively. Clinical scores of corneal grafts for opacity were as follows: 0= clear; +1=minimal, superficial (nonstromal) opacity; pupil margin and iris vessels readily visible through the cornea; +2=minimal, deep (stroma) opacity; pupil margins and iris vessels visible; +3= moderate stromal opacity; only pupil margin visible; +4= intense stromal opacity; only a portion of pupil margin

visible; +5= maximum stromal opacity; anterior chamber not visible. Grafts with opacity scores of +2 or greater after 2 weeks were considered to have been rejected. Syngeneic transplantations were performed and evaluated in a similar manner.

[0058] Immunohistochemistry and morphometry of angiogenesis and lymphangiogenesis in the cornea. Briefly, corneal flat mounts were rinsed in PBS, fixed in acetone, rinsed in PBS, blocked in 2% bovine serum albumin, stained with FITC-conjugated CD31/PECAM-1 overnight (Santa Cruz Biotechnology, Santa Cruz, CA; 1:100), washed, blocked, stained with LYVE-1 (1:500; a lymphatic endothelium specific hyaluronic acid receptor (Cursiefen et al. (2002) Invest. Ophthalmol. Vis. Sci. 43:2127-35) washed, blocked, and stained with Cy3 (1:100; Jackson ImmunoResearch Laboratories, West Grove, PA) and analyzed using a Zeiss Axiophot microscope. Digital pictures of the flat mounts were taken using Spot Image Analysis system. Then the area covered by CD31⁺⁺⁺/LYVE-1⁻ blood vessels and CD31⁺/LYVE-1⁺⁺⁺ lymph vessels was measured morphometrically on these flat-mounts using NIH Image software. The total corneal area was outlined using the innermost vessel of the limbal arcade as the border. The total area of blood versus lymphatic neovascularization was then normalized to the total corneal area and the percentage of the cornea covered by each vessel type calculated.

[0059] Neutralization of VEGF-A using VEGF Trap_{R1R2}. The VEGF trap_{R1R2} (Regeneron Pharmaceuticals Inc, Tarrytown, NY (Holash et al. (2002) Proc. Natl. Acad. Sci. USA 99:11393-8, herein specifically incorporated by reference in its entirety) was used in the transplant survival experiment at a concentration of 12.5 mg/kg intraperitoneally (i.p.) at time of surgery (CHO hVEGFR1 [Ig domain 2] R2 [Ig domain 3]-Fc), and 3, 7, and 14 days after surgery. Human Fc-fragment given i.p. at same concentration and times was used in the control mice (sCHO h Fc).

[0060] Statistical analysis. Statistical significance was analyzed by Mann-Whitney's test. Differences were considered significant at $P < 0.05$. Each experiment was performed at least twice with similar results. Graphs were drawn using Graph Pad Prism, Version 3.02.

[0061] Results. Dose response of angiogenesis inhibition by VEGF Trap_{R1R2}. VEGF-Trap_{R1R2} at doses of either 25 mg/kg or 12.5 mg/kg completely inhibited suture-induced inflammatory corneal neovascularization. In contrast, doses of 6.25mg/kg and 2.5mg/kg produced ~50% and ~20% inhibition of corneal neovascularization, respectively, while the lowest dose tested, 0.5 mg/kg, had a negligible effect (<5% inhibition). Therefore, for subsequent experiments a dose of 12.5 mg/kg VEGF Trap_{R1R2} was chosen.

[0062] Rapid and parallel onset of hemangiogenesis and lymphangiogenesis after normal-risk allogeneic corneal transplantation. To determine whether the mild and temporary hemangiogenesis occurring after normal-risk keratoplasty is accompanied by lymphatic vessel outgrowth from the limbus into the normally alymphatic cornea, we studied the time course of

25.8±4.4%; day 7: 48.3±12.8%; $p<0.0001$). This was also true for the lymphovascularized area comparing Trap- (9.5±9.4%) and Fc-treated mice on day 3 (21.5±9.3%; $p<0.0001$). At day 7, the lymphovascularized area was smaller, but not significantly different in the Trap-group (28.7±20.3%) compared to the Fc-group (51.5±23.8%; $p=0.06$). In contrast to results obtained in corneal injury models neither hem- or lymphangiogenesis were completely inhibited by the VEGF Trap_{R1R2} following corneal transplantation. However, the number of lymphatic vessels reaching the graft-host interface (10.6±0.6 versus 1.3±1.5 vessels) and the number of hours where the interface was filled with draining lymphatic vessels were much larger in the Fc-treated compared to the Trap-treated group (3±2 versus 0.2±0.3 hours; not significant due to small sample size) at day 7. This might indicate that lymphovascularized area per se is less decisive for host sensitisation than the contact area with donor tissue.

[0065] Partial inhibition of early postoperative hem- and lymphangiogenesis by trapping VEGF-A after normal-risk surgery improves long-term graft survival.

Since hem- and lymphangiogenesis occurring *after* normal-risk keratoplasty peaked around day 7, and regressed thereafter, and since both vascular processes could be significantly inhibited by early postoperative neutralization of VEGF-A, we determined whether inhibition of postkeratoplasty hem- and lymphangiogenesis during this interval improves graft survival. The long-term survival of C57BL/6 grafts placed into avascular BALB/c recipient beds was compared between mice receiving an i.p. injection of 12.5 mg/kg VEGF Trap_{R1R2}, or Fc-fragment alone, at surgery and 3, 7, and 14 days later. Trapping of VEGF-A postoperatively caused a significantly improved long-term graft survival at 8 weeks (78%), compared to grafts in eyes of Fc-treated controls (40%; $p=0.044$; $n=22$ in both groups).

[0066] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof.

[0066a] The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

[0066b] Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

comparing Trap- ($9.5 \pm 9.4\%$) and Fc-treated mice on day 3 ($21.5 \pm 9.3\%$; $p < 0.0001$). At day 7, the lymphovascularized area was smaller, but not significantly different in the Trap-group ($28.7 \pm 20.3\%$) compared to the Fc-group ($51.5 \pm 23.8\%$; $p = 0.06$). In contrast to results obtained in corneal injury models neither hem- or lymphangiogenesis were completely inhibited by the VEGF Trap_{R1R2} following corneal transplantation. However, the number of lymphatic vessels reaching the graft-host interface (10.6 ± 0.6 versus 1.3 ± 1.5 vessels) and the number of hours where the interface was filled with draining lymphatic vessels were much larger in the Fc-treated compared to the Trap-treated group (3 ± 2 versus 0.2 ± 0.3 hours; not significant due to small sample size) at day 7. This might indicate that lymphovascularized area per se is less decisive for host sensitisation than the contact area with donor tissue.

[0065] Partial inhibition of early postoperative hem- and lymphangiogenesis by trapping VEGF-A after normal-risk surgery improves long-term graft survival.

Since hem- and lymphangiogenesis occurring *after* normal-risk keratoplasty peaked around day 7, and regressed thereafter, and since both vascular processes could be significantly inhibited by early postoperative neutralization of VEGF-A, we determined whether inhibition of postkeratoplasty hem- and lymphangiogenesis during this interval improves graft survival. The long-term survival of C57BL/6 grafts placed into avascular BALB/c recipient beds was compared between mice receiving an i.p. injection of 12.5 mg/kg VEGF Trap_{R1R2}, or Fc-fragment alone, at surgery and 3, 7, and 14 days later. Trapping of VEGF-A postoperatively caused a significantly improved long-term graft survival at 8 weeks (78%), compared to grafts in eyes of Fc-treated controls (40%; $p = 0.044$; $n = 22$ in both groups).

[0066] The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof.

EDITORIAL NOTE

APPLICATION NUMBER – 2004242586

The next page number is 18. The claims also start from page 18.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. Use of an first agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity in the preparation of a medicament for treating or preventing corneal transplant rejection in a mammalian subject, wherein the agent is VEGFR1R2-FcΔC1(a).
2. The use of claim 1, wherein administration is subcutaneous, intramuscular, intradermal, intraperitoneal, intravenous, intranasal, oral, subconjunctival, or topical. Administration may also include a second agent, such as an immunosuppressive agent.
3. The use of claim 1, further comprising administering a second agent.
4. The use of claim 3, wherein the second agent is an immunosuppressive agent.
5. The use of claim 1, wherein the mammalian subject is a human.
6. The use of claim 5, wherein the human subject has received a corneal transplant.
7. A method of reducing the incidence of corneal transplant rejection in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that the incidence of corneal transplant rejection is reduced, wherein the agent is VEGFR1R2-FcΔC1(a).
8. A method of treating corneal transplant rejection in a subject in need thereof, comprising administering to the subject an agent capable of blocking, inhibiting, or ameliorating vascular endothelial growth factor (VEGF)-mediated activity, such that corneal transplant rejection is treated, wherein the agent is VEGFR1R2-FcΔC1(a).
9. A pharmaceutical composition when used for prevention or treatment of corneal transplant rejection, comprising a vascular endothelial growth factor (VEGF) antagonist, and a pharmaceutically acceptable carrier, wherein the antagonist is VEGFR1R2-FcΔC1(a).

10. The pharmaceutical composition of claim 9, in the form of a liquid, gel, ointment, salve, or ophthalmic solution.

11. An article of manufacturing when used for prevention or treatment of corneal transplant rejection comprising:

(a) packaging material; and

(b) a pharmaceutical agent contained within the packaging material;

wherein the pharmaceutical agent comprises at least one VEGF-specific fusion protein of the invention and wherein the packaging material comprises a label or package insert which indicates that the VEGF-specific fusion protein can be used to treat or prevent corneal transplant rejection in a mammalian subject, wherein the agent is VEGFR1R2-FcΔC1(a).

12. A use of claim 1, substantially as hereinbefore described with reference to the Examples.

13. The method of claims 7 and 8, substantially as hereinbefore described with reference to the Examples.

14. A pharmaceutical composition according to claim 9, substantially as hereinbefore described with reference to the Examples.

15. An article of manufacturing of claim 11, substantially as hereinbefore described with reference to the Examples.

SEQUENCE LISTING

<110> Regeneron Pharmaceuticals, Inc.
The Schepens Eye Research Institute

<120> Method of Treating Corneal Transplant Rejection

<130> REG 713B-WO

<140> To be Assigned

<141> 2004-04-23

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 1377
<212> DNA
<213> homo sapiens

<400> 1
atgggcagct actgggacac cggggtcctg ctgtgcgcgc tgcctcagctg tctgcttctc 60
acaggatcta gttccggaag tgataccggt agacctttcg tagagatgta cagtgaatac 120
cccgaaatta tacacatgac tgaaggaag gagctcgtca ttccctgccg gggtacgtca 180
cctaacaatca ctgttacttt aaaaaagttt ccacttgaca ctttgatccc tgaatggaaa 240
cgcataatct gggacagtag aaagggcttc atcatatcaa atgcaacgta caaagaaata 300
gggcttctga cctgtgaagc aacagtcaat gggaatttgt ataagacaaa ctatctcaca 360
catcgacaaa ccaatacaat catagatgtg gttctgagtc cgtctcatgy aattgaacta 420
tctgttggag aaaagcttgt cttaaattgt acagcaagaa ctgaactaaa tgtggggatt 480
gacttcaact gggaataccc ttcttcgaag catcagcata agaaacttgt aaaccgagac 540
ctaaaaaccc agtctgggag tgagatgaag aaatttttga gcaccttaac tatagatggg 600
gtaaccgcga gtgaccaaag attgtacacc tgtgcagcat ccagtgggct gatgaccaag 660
aagaacagca ctttgttcag ggtccatgaa aaggacaaaa ctccacatgy cccaccgtgc 720
ccagcacctg aactcctggg gggaccgtca gtcttctctt tcccccaaaa acccaaggac 780
accctcatga tctcccgac ccctgaggtc acatgcgtgy tgggtggacgt gagccacgaa 840
gaccttgagg tcaagttcaa ctggtacgtg gacggcgtgy aggtgcataa tgccaagaca 900
aagccgcggg aggagcagla caacagcacg tacogtgytg tcagcgtcct caccgtcctg 960
caccaggact ggctgaatgg caaggagtac aagtgcagg tctccaacaa agccctccca 1020
gcccccatcg agaaaacat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac 1080
accctgcccc catcccgga tgagctgacc aagaaccagc tcagcctgac ctgcctggtc 1140
aaaggtctct atcccagcga catcgccgtg gagtgggaga gcaatgggca gccggagaa 1200
aactacaaga ccacgcctcc cgtgtggac tccgacggct ccttcttctc ctacagcaag 1260
ctcaccgtgg acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat 1320
gaggctctgc acaaccacta cagcagaag agcctctccc tgtctccggg taaatga 1377

<210> 2
<211> 458
<212> PRT
<213> homo sapiens

<400> 2
Met Val Ser Tyr Trp Asp Thr Gly Val Leu Leu Cys Ala Leu Leu Ser
1 5 10 15
Cys Leu Leu Leu Thr Gly Ser Ser Ser Gly Ser Asp Thr Gly Arg Pro

```

      20      25      30
Phe Val Glu Met Tyr Ser Glu Ile Pro Glu Ile Ile His Met Thr Glu
      35      40      45
Gly Arg Glu Leu Val Ile Pro Cys Arg Val Thr Ser Pro Asn Ile Thr
      50      55      60
Val Thr Leu Lys Lys Phe Pro Leu Asp Thr Leu Ile Pro Asp Gly Lys
      65      70      75      80
Arg Ile Ile Trp Asp Ser Arg Lys Gly Phe Ile Ile Ser Asn Ala Thr
      85      90      95
Tyr Lys Glu Ile Gly Leu Leu Thr Cys Glu Ala Thr Val Asn Gly His
      100      105      110
Leu Tyr Lys Thr Asn Tyr Leu Thr His Arg Gln Thr Asn Thr Ile Ile
      115      120      125
Asp Val Val Leu Ser Pro Ser His Gly Ile Glu Leu Ser Val Gly Glu
      130      135      140
Lys Leu Val Leu Asn Cys Thr Ala Arg Thr Glu Leu Asn Val Gly Ile
      145      150      155      160
Asp Phe Asn Trp Glu Tyr Pro Ser Ser Lys His Gln His Lys Lys Leu
      165      170      175
Val Asn Arg Asp Leu Lys Thr Gln Ser Gly Ser Glu Met Lys Lys Phe
      180      185      190
Leu Ser Thr Leu Thr Ile Asp Gly Val Thr Arg Ser Asp Gln Gly Leu
      195      200      205
Tyr Thr Cys Ala Ala Ser Ser Gly Leu Met Thr Lys Lys Asn Ser Thr
      210      215      220
Phe Val Arg Val His Glu Lys Asp Lys Thr His Thr Cys Pro Pro Cys
      225      230      235      240
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      245      250      255
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      260      265      270
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
      275      280      285
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      290      295      300
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
      305      310      315      320
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      325      330      335
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      340      345      350
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu
      355      360      365
Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      370      375      380
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      385      390      395      400
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      405      410      415
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      420      425      430
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      435      440      445
Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
      450      455

```