





(86) Date de dépôt PCT/PCT Filing Date: 2006/08/16
(87) Date publication PCT/PCT Publication Date: 2007/02/22
(45) Date de délivrance/Issue Date: 2010/03/30
(85) Entrée phase nationale/National Entry: 2007/03/16
(86) N° demande PCT/PCT Application No.: EP 2006/065380
(87) N° publication PCT/PCT Publication No.: 2007/020277
(30) Priorité/Priority: 2005/08/17 (EP05107582.8)

(51) Cl.Int./Int.Cl. *C07D 475/08* (2006.01),
A61K 31/505 (2006.01), *A61K 31/519* (2006.01),
A61P 35/00 (2006.01), *C07D 475/04* (2006.01)
(72) Inventeur/Inventor:
STOICESCU, DAN, CY
(73) Propriétaire/Owner:
STOICESCU, DAN, CY
(74) Agent: SIM & MCBURNEY

(54) Titre : NOUVEAUX INHIBITEURS D'ENZYMES DEPENDANT DE L'ACIDE FOLIQUE
(54) Title: NOVEL INHIBITORS OF FOLIC ACID-DEPENDENT ENZYMES



(57) **Abrégé/Abstract:**

The invention relates to compounds of the formula I, or pharmaceutically acceptable salts thereof: wherein: Z = O or S; n = 1 - 3; R³ = -CO₂R⁸, -C(O)SR⁸, -C(O)NHR⁸, -C(S)OR⁸, -C(S)SR⁸, -C(S)NHR⁸, -C(NH)SR⁸ or -C(NH)NHR⁸, wherein R⁸ is -H or alkyl; R⁴ = -H, -CH₂R⁵ or -CH₂CH₂R⁵, wherein R⁵ independently has one of the meanings of R³; B = -NR²-, -CH₂NR²-, -CH₂CH₂NR²-, -CH₂CHR⁷- or -CH₂O-, wherein R² is H or a C₁₋₃ alkyl, alkenyl or alkynyl group, and R⁷ is H or a C₁₋₃ alkyl or alkoxy group; A = wherein R¹ = -NH₂ or -OH, and C and D are each, independently, a 5- or 6- membered, substituted or unsubstituted, aromatic or non-aromatic ring which may also contain one or more heteroatoms, and C is connected to group B in any available position.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2007 (22.02.2007)

PCT

(10) International Publication Number
WO 2007/020277 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/EP2006/065380

(22) International Filing Date: 16 August 2006 (16.08.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05107582.8 17 August 2005 (17.08.2005) EP

(71) Applicant and

(72) Inventor: STOICESCU, Dan [RO/CY]; Gr. Afxenthiou
5, Agios Tychonas, Limassol, 4521 (CY).

(74) Agent: KILLIN, Stephen; 20 Little Britain, London
EC1A 7DH (GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP,
KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT,
LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ,
NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU,
SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT,
TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.



(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL INHIBITORS OF FOLIC ACID-DEPENDENT ENZYMES



(57) Abstract: The invention relates to compounds of the formula I, or pharmaceutically acceptable salts thereof: wherein: Z = O or S; n = 1 - 3; R³ = -CO₂R⁸, -C(O)SR⁸, -C(O)NHR⁸, -C(S)OR⁸, -C(S)SR⁸, -C(S)NHR⁸, -C(NH)SR⁸ or -C(NH)NHR⁸, wherein R⁸ is -H or alkyl; R⁴ = .H, -CH₂R⁵ or -CH₂CH₂R⁵, wherein R⁵ independently has one of the meanings of R³; B = -NR²-, -CH₂NR²-, -CH₂CH₂NR²-, -CH₂CHR⁷- or -CH₂O-, wherein R² is H or a C₁₋₃ alkyl, alkenyl or alkynyl group, and R⁷ is H or a C₁₋₃ alkyl or alkoxy group; A = wherein R¹ = -NH₂ or -OH, and C and D are each, independently, a 5- or 6- membered, substituted or unsubstituted, aromatic or non-aromatic ring which may also contain one or more heteroatoms, and C is connected to group B in any available position.



WO 2007/020277 A2

Novel inhibitors of folic acid-dependent enzymes


Description

The present invention relates to a novel class of compounds. In particular, the
5 present invention relates to novel compounds that inhibit enzymes whose natural
substrates are folic acid or folic acid derivatives (folates), and that may be used in
the treatment of diseases such as cancer.


Cancer cells replicate more rapidly than most other cells and so have a greater
10 demand for nucleotides, the precursors of deoxyribonucleic acid (DNA) and
ribonucleic acid (RNA). Since all cells do not maintain a residual store of
nucleotides (except for adenosine triphosphate, ATP), they must be synthesized
continually during DNA and RNA synthesis. Accordingly, the replication of cancer
cells tends to be more sensitive than that of healthy cells to inhibition of nucleotide
15 biosynthesis, and for this reason interest is increasing in chemotherapeutic agents
capable of effecting such inhibition.

Nucleotides may be synthesised biologically *via* de novo pathways from fundamental
metabolic precursors, or *via* salvage pathways whereby the products of nucleic acid
20 degradation, free bases and nucleosides, are recycled. The de novo pathways are of
primary interest with regard to the search for new chemotherapeutic agents.

The nucleotides 2'-deoxyadenosine-5'-monophosphate (dAMP) and
2'-deoxyguanosine-5'-monophosphate (dGMP) are both derived de novo from
25 inosine monophosphate (IMP), which in turn is derived from 5-phosphoribosyl-1-
pyrophosphate (PPRP). Two enzymes involved in the biosynthetic pathway
between PPRP and IMP are GAR transformylase and AICAR transformylase. GAR
transformylase converts glycinamide ribonucleotide (GAR) to formylglycinamide
ribonucleotide (FGAR) using N¹⁰-formyltetrahydrofolate, whereas AICAR
30 transformylase uses the same compound to convert 5-aminoimidazole-4-
carboxamide ribonucleotide (AICAR) to N-formylaminoimidazole-4-carboxamide
ribonucleotide (FAICAR), as shown below.




- 5 The nucleotide 2'-deoxythymidine-5'-monophosphate (dTMP), on the other hand, is produced by de novo synthesis from 2'-deoxyuridine-5'-monophosphate (dUMP), a conversion catalysed by the enzyme thymidylate synthase. During the conversion, N⁵,N¹⁰-methylene-tetrahydrofolate is reduced to 7,8-dihydrofolate; the former is regenerated *via* tetrahydrofolate using the enzymes dihydrofolate reductase (DHFR)
- 10 and serine hydroxymethyl-transferase. These processes are illustrated below.



Interference with these mechanisms has been exploited in the treatment of cancer. For example, US 2,512,572 discloses a number of substituted pteridines including the potent chemotherapeutic agent methotrexate, which belongs to the class of “folate antagonists”, and inhibits DNA synthesis by competitively antagonising dihydrofolate reductase, binding with about 100 times higher affinity than its natural substrate, thereby preventing the regeneration of tetrahydrofolate which is essential for the synthesis of dTMP. This leads to so-called “thymine-less death” in cancer cells. Methotrexate also inhibits GAR transformylase, AICAR transformylase and thymidylate synthase, albeit to a lesser degree. The structures of methotrexate and other related anti-folates are shown below.

Compound	Structure
Methotrexate	
Aminopterin	
Pemetrexed	
Lometrexol	


US 4,684,653 discloses compounds of the formula:



5


wherein R^1 is OH or NH_2 , and R^3 is H, Me or Et, and their corresponding 5,6,7,8-tetrahydro derivatives. These compounds are disclosed to have an effect on one or more enzymes that utilize folic acid and its metabolic derivatives as a substrate.

US 5,344,932 discloses glutamic acid derivatives of the formula:



- 5 wherein R^5 is H or NH_2 , R^4 is H or OMe, and R^2 is H or a pharmaceutically acceptable cation, and discloses that they have an inhibitory effect on one or more enzymes which utilise folic acid and its metabolic derivatives as a substrate.


US 4,077,957 discloses a method of synthesising various pteridine compounds,
10 including:



Although such compounds have proved useful in developing new therapeutic
15 strategies for treating cancer, there are still a number of problems associated with their use, including low efficacy, intrinsic and acquired resistance to such drugs in some patients, toxicity and adverse side effects. Consequently, there remains a need for alternative compounds that can be used in treating cancer and may address one or more of the aforementioned problems.

20

Accordingly, in a first aspect of the invention, there is provided a compound of the formula I, or a pharmaceutically acceptable salt thereof:



wherein:

Z = O or S;

n = 1 – 3;

5 $R^3 = -CO_2R^8, -C(O)SR^8, -C(O)NHR^8, -C(S)OR^8, -C(S)SR^8, -C(S)NHR^8,$
 $-C(NH)SR^8$ or $-C(NH)NHR^8,$

wherein R^8 is -H or alkyl;

$R^4 = -H, -CH_2R^5$ or $-CH_2CH_2R^5,$


wherein R^5 independently has one of the meanings of R^3 ;

10 $B = -NR^2-, -CH_2NR^2-, -CH_2CH_2NR^2-, -CH_2CHR^7-$ or $-CH_2O-,$

wherein R^2 is H or a C_{1-3} alkyl, alkenyl or alkynyl group, and

R^7 is H or a C_{1-3} alkyl or alkoxy group;

A =



15

wherein $R^1 = -NH_2$ or $-OH,$

C and D are each, independently, a 5- or 6-membered, substituted or unsubstituted, aromatic or non-aromatic ring which may also contain one or more heteroatoms,

20

and C is connected to group B in any available position.

In a second aspect of the invention, there is provided a compound according to the invention in its first aspect, for use in therapy.


In a third aspect of the invention, there is provided a pharmaceutical composition comprising a compound according to the invention in its first or second aspects.

In a fourth aspect of the invention, there is provided the use of a compound
 5 according to the invention in its first or second aspects for the manufacture of a medicament for use in the treatment of a condition responsive to inhibition of an enzyme dependent upon folic acid or a folic acid derivative.

In a fifth aspect of the invention, there is provided the use of a compound
 10 according to the invention in its first or second aspects for the manufacture of a medicament for use in the treatment of cancer.


In a sixth aspect of the invention, there is provided a method of preparing a compound according to the invention in its first or second aspects, which comprises
 15 the step of:

(a) reacting a compound of the formula II



20

wherein A is as previously defined, m is 0, 1 or 2 and X is a leaving group, with a compound of the formula III




25 wherein Z, n, R², R³ and R⁴ are as previously defined;

(b) reacting a compound of the formula IV



wherein A is as previously defined and X is a leaving group, with a compound of the formula V




5

wherein Z, n, R³ and R⁴ are as previously defined; or

(c) converting one of the following compounds VI or VII into a corresponding organometallic reagent


10




wherein A, Z, n, R³, R⁴, and R⁷ are as previously defined, and Y is, in each case
independently, a halide, and reacting said reagent with the other of compounds VI
or VII.

In a seventh aspect of the invention, there is provided a method of preparing a
compound according to the invention in its first or second aspects having the
formula I-A or a pharmaceutically acceptable salt thereof


20



wherein n , R^1 , R^3 and R^4 are as previously defined, which comprises reacting a compound of the formula II-A




5 wherein X is Cl, Br or I, with a compound of the formula III-A




In an eighth aspect of the invention, there is provided a compound of the formula III, V or VII as previously defined, except that R^4 is $-\text{CH}_2\text{R}^5$ or $-\text{CH}_2\text{CH}_2\text{R}^5$.

10 In accordance with an aspect of the present invention, there is provided a compound of the formula I-A1, or a pharmaceutically acceptable salt thereof:




15 In accordance with another aspect of the present invention, there is provided a compound of the formula I-A2, or a pharmaceutically acceptable salt thereof:

9a




In accordance with another aspect of the present invention, there is provided a compound of the formula I-A3, or a pharmaceutically acceptable salt thereof:




5

In accordance with another aspect of the present invention, there is provided a compound of the formula I-A4, or a pharmaceutically acceptable salt thereof:




10 In accordance with another aspect of the present invention, there is provided a compound of the formula I-A5, or a pharmaceutically acceptable salt thereof:



15 In accordance with another aspect of the present invention, there is provided a compound of the formula I-A6, or a pharmaceutically acceptable salt thereof:

9b



Preferred embodiments of the invention in any of its various aspects are as described below or as defined in the sub-claims.

5

In all of the various aspects of the invention, the carbon marked C^* may be asymmetric (when R^4 is not H) and in this event it will be appreciated that compounds of the formula I may exist in racemic form, or may be separated into their (+) or (-) enantiomers by conventional methods. In addition, other chiral centres may be present in some compounds giving rise to one or more further pairs

10

of enantiomers. For example, a second chiral centre exists in those compounds wherein $B = -CH_2CHR^7-$ wherein R^7 is a C_{1-3} alkyl or alkoxy group. All such racemic or enantiomeric forms are intended to lie within the scope of the present invention. Furthermore, it will be understood that the compounds of formula I may
5 exist in one or more tautomeric forms, and each of these forms are also intended to lie within the scope of the present invention.


As described in more detail hereinafter, the compounds of formula I are structural analogs of folic acid and have been found to possess activity as inhibitors of those
10 enzymes that are dependent upon folic acid or folic acid derivatives (folates), such as dihydrofolate reductase (DHFR), at levels comparable to that of methotrexate *in vitro*. The compounds of formula I have also been shown to be active in inhibiting tumor growth in animal models *in vivo*. It is expected that the latter activity may be due to the compounds' ability to act as competitive antagonists of DHFR, although
15 details of the mechanism are not presented here. The compounds of formula I may be used in treating cancer, as well as conditions that are responsive to inhibition of an enzyme dependent upon folic acid or a folic acid derivative.

In preferred compounds of the formula I, one or more of the following conditions
20 are satisfied:


- Z is O;
- n is 1;
- R^3 is $-CO_2R^8$ and R^4 is $-CH_2CH_2CO_2R^8$;
- R^8 is -H, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or tertiary
25 butyl, preferably -H, -Me or -Et, preferably -H;
- B is $-CH_2NR^2-$, $-CH_2CHR^7-$ or $-CH_2O-$, preferably $-CH_2NR^2-$;
- R^2 is -H, -Me, -Et or $-CH_2-C\equiv CH$, preferably H;
- R^7 is -H, -Me, -Et or -OMe, preferably H.

30 In addition, preferred compounds of the formula III, V or VII display one or more of the preferred designations of Z, n, R^2 , R^3 , R^4 and/or R^7 set out above.

In group A in formulae I, II, IV or VI, D is preferably a 5-membered heteroaromatic ring. Preferably, A is




- 5 In group A, C may be one of the following groups (points of attachment to the adjacent ring and to group B are shown):




- 10 wherein X is CH or N, and either: Y is C and R⁶ is H, Me, Et or HCO; or Y is N and R⁶ is a lone pair of electrons. In preferred embodiments, X and Y are both N and R⁶ is a lone pair of electrons.

- 15 Especially preferred A groups are those of the following structures, which closely mimic those found in naturally occurring pteridines and other heterocyclic bases:





Of particular interest are the following two A groups:




5




Especially preferred are compounds of the formula I having the above two A groups, wherein B is $-\text{CH}_2\text{NR}^2-$, R^2 is $-\text{H}$, $-\text{Me}$, $-\text{Et}$ or $-\text{CH}_2\text{C}\equiv\text{CH}$, Z is O, n is 1, and R^3 is $-\text{CO}_2\text{R}^8$, preferably any hydrolysable ester group. Individual examples of

10



15





5

10

15

20

The compounds of the formula I can be prepared by the methods of the invention from readily available and inexpensive starting materials. In the case where B is -NR²-, -CH₂NR²- or -CH₂CH₂NR²-, for example, the compounds of formula I can be prepared by coupling a compound of the formula II with a compound of the formula III. The leaving group X will generally be a halogen such as chlorine, bromine or iodine, especially bromine or iodine. This reaction is preferably performed in a dipolar aprotic solvent such as dimethyl formamide (DMF) or dimethylacetamide (DMAc). A basic catalyst such as potassium fluoride may be used, which affords a higher yield than tertiary amines or sodium bicarbonate. Where necessary, sensitive groups may be protected prior to the reaction using suitable protecting groups known in the art, and later deprotected *via* standard methods. For example, when R³ is H and R⁴ is CH₂CH₂CO₂H, these acid groups may be protected for instance as methyl ester groups, with subsequent deprotection by known methods such as alkaline hydrolysis with sodium hydroxide in ethanol and precipitation by addition of acid, such as glacial acetic acid. Accordingly, it will be understood that the method of the invention encompasses the reaction of

compounds of the formulae II and III wherein either or both of these compounds are in a protected form.


10 In the case where B is $-\text{CH}_2\text{O}-$, the compounds of formula I may be prepared, for example, by coupling a compound of the formula IV with a compound of the formula V by a Williamson ether type reaction. In this reaction, the compound of formula V is generally converted into its aroxide ion form prior to reaction with the compound of formula IV, using a base such as NaH, for instance. X may be any suitable leaving group, in particular a halide.

10

15 In the case where B is $-\text{CH}_2\text{CHR}^7-$, the compound of formula I may be prepared, for example, by coupling a compound of the formula VI with a compound of the formula VII by any known carbon-carbon bond forming reaction, especially those involving the use or formation of organometallic reagents such as Grignard reagents and lithium or copper-lithium compounds. For instance, a compound of the formula VII may be converted into its corresponding Grignard reagent or lithium cuprate reagent and reacted with a compound of the formula VI. Alternatively, a compound of the formula VI may be converted into its corresponding Grignard reagent or lithium cuprate reagent and reacted with a compound of the formula VII.


20 Once again, suitable protecting groups for any reactive substituent groups will be well-known to those skilled in the art.

The intermediates II to VII may be prepared by conventional methods. By way of illustration, compounds of the formula III, V or VII may be prepared by reacting a
25 compound of the formula



with a compound of the formula

30



wherein B' is -NHR², -OH or -CHYR⁷ and X is a leaving group, in the presence of a base. This may be followed by removal of the cyano group by hydrolysis and
 5 decarboxylation, with the use of suitable protecting groups where necessary.

Compounds of formula I have an inhibitory effect on one or more enzymes which utilize folic acid, and in particular metabolic derivatives of folic acid as a substrate. These enzymes include GAR transformylase, AICAR transformylase, dihydrofolate
 10 reductase and thymidylate synthase. The compounds appear to be particularly active as inhibitors of dihydrofolate reductase. They can be used, alone or in combination, to treat neoplasms which in the past have been treated with methotrexate, including choriocarcinoma, leukaemia, adenocarcinoma of the female breast, epidermid cancers of the head and neck, squamous or small-cell lung cancer,
 15 and various lymphosarcomas. Although not wishing to be limited by theory, it is believed that the modified ketomethylenic or thioketomethylenic side chain of the compounds of formula I leads to a lower renal toxicity compared with methotrexate. Inactivation due to hydrolysis is minimal due to the lower lability of the ketomethylenic or thioketomethylenic group, allowing a longer half-life. In
 20 addition, compounds of the formula I exhibit improved physico-chemical characteristics compared with those of the prior art.

As discussed above, primary interest in the compounds of the invention relates to their use in the treatment of cancer, since cancerous cells replicate more rapidly
 25 than healthy cells and so have a greater demand for nucleotides. The therapeutic utility of the compounds of the invention extends further than this, however, since all fast-growing cells have a similar high demand for nucleotides. For example, methotrexate has been used in the treatment of psoriasis and mycosis fungoides, and in inducing miscarriage in patients with ectopic pregnancy. Methotrexate has

also been used in the treatment of rheumatoid arthritis, although the mechanism of action in this instance is not fully understood.

The compounds may be administered either orally or preferably parenterally, alone
5 or in combination with other anti-neoplastic agents or with other therapeutic agents, such as steroids, to a mammal, preferably a human, suffering from neoplasm and in need of treatment. Parenteral routes of administration include intramuscular, intrathecal, intravenous or intra-arterial.

10 In order that the invention may be more fully understood, it will now be described by way of example only with reference to the accompanying drawing, wherein

Figure 1 is a schematic representation of the total synthesis of two compounds according to the present invention (compounds 3 and 4), as described in detail
15 below in Examples 1 and 2.

The following examples are intended to demonstrate the invention but are not intended to limit the invention in any manner.

20 **Example 1**

Synthesis of compound 3

Source of starting materials

3-chloropropanoyl chloride and ethyl cyanoacetate were obtained commercially from Sigma-Aldrich Company Ltd., The Old Brickyard, New Road, Gillingham,
25 Dorset SP8 4XT, United Kingdom, or synthesised by standard methods. α -bromo-p-nitro-acetophenone 7 was obtained by bromination of p-nitroacetophenone with bromine in tetrahydrofuran (THF). 2,4-diamino-6-bromomethylpteridine 2 was obtained by standard methods (see, for example, US 4,077,957 and US 4,224,446).

30 **Step A:**

Synthesis of compound 6

3-chloropropanoyl chloride was esterified with ethanol in the presence of pyridine or triethylacetate to produce ethyl 3-chloropropionate 5. The latter was condensed

with ethyl cyanoacetate according to the procedure of L. Ruzicka *et. al.*, *Helv. Chim. Acta* 17, 183 – 200 (1934), CA 28:2584, or Koelsch, C.F., *J. Am. Chem.* 65, 2458-9 (1943), to form diethyl α -cyanoglutarate **6**. ^1H -NMR confirmed the expected structure. GC: 97 % purity.

5

Step B:**Synthesis of compound 8**

175 g (0,71 mmol) α -bromo-p-nitro-acetophenone **7** was added in portions at 0 – 5 °C to a suspension of 175 g (0.82 mmol) diethyl α -cyanoglutarate **6** and 175 g (3 mmol) KF in 500 ml DMF. The reaction was monitored by thin layer chromatography (TLC). After 4 hours, the reaction mixture was suspended in 2 l of water containing 0.1 % acetic acid at pH 5. After decanting the water, the gummy precipitate was washed with water (2 x 750 ml) then triturated with 300 ml methanol. When crystallisation was complete, the precipitate was filtered and washed successively with an excess of methanol and ether, affording 210 g of compound **8**, a yellow solid with m.p. 92.1 °C. (Yield 68 %) After chromatography on silica gel (50:50:5 benzene-cyclohexane-ethanol) the product had m.p. 99.7 °C. TLC on silica gel plates (5:1:3:10:0.1 benzene-ethanol-cyclohexane-petroleum ether-AcOH) showed a single spot with R_f (retention factor) 0.38. HPLC: 97 % purity.

20

Step C:**Synthesis of compound 9**

30 g (0.08 mmol) compound **8** was dissolved in 400 ml methanol and hydrogenated in a hydrogenation flask at room temperature in the presence of 6 g 20 % Pd/C catalyst. The theoretical volume of hydrogen (c. 6200 ml; 0.28 mmol) was absorbed in 1 hour (TLC control). The platinum catalyst was filtered and the methanol was evaporated. The crude product obtained solidified on drying *in vacuo*, resulting in 27.6 g compound **9**, a yellow solid (yield 99 %) which was used without further purification in the conversion to crude compound **10** described below. The purity was acceptable by TLC analysis. TLC (4:1 chloroform-methanol) showed a single spot, R_f 0.5 (characteristic reaction with 4-dimethylamino benzaldehyde). The HCl salt was isolated after reflux in HCl. LC-MS and ^1H -NMR confirmed the expected structure; HPLC: 99 % purity.

30

Step D:**Synthesis of compound 10**

A solution of 52.2 g (0.15 mmol) intermediate **9** in 1000 ml methanol was prepared.
5 188 ml 6N NaOH was added dropwise at room temperature for 1 hour and the solution was allowed to stand for 12 hours. The reaction mixture was then diluted with 300 ml water and concentrated under high vacuum. 700 ml 37 % HCl was added to the residue and the resulting mixture was heated to reflux for 4 hours. The mixture obtained was diluted with 1.5 l methanol and the NaCl precipitate was
10 removed by filtration. The filtrate was used in step E. A small amount of diacid **10** was isolated before dilution, by filtering the suspension and washing the precipitate successively with an excess of water, acetone and ether. TLC (4:1 chloroform-methanol) showed a single spot, R_f 0.26.

Step E:**Synthesis of compound 1**

The methanolic solution of the dicarboxylic acid **10** obtained in step D was cooled at 0 – 5 °C and 100 ml thionyl chloride was added dropwise. The reaction mixture was stirred under reflux for 3 hours, then cooled to room temperature and the
20 solvent was evaporated off. The precipitate obtained was filtered and washed with ether, resulting in 27 g compound **1** (yield 63 %), a solid with m.p. 115 – 116 °C. After recrystallisation from tetrahydrofuran, 17.5 g white crystals of **1** were obtained having m.p. 116 – 117 °C. TLC (4:1 chloroform-methanol) showed a single spot, R_f 0.73. UV spectra: 234, 319 nm (MeOH). ¹H NMR spectra: 2.0 (2H, m, CH₂CH₂COOCH₃), 2.5 (2H, t, CH₂CH₂COOCH₃), 3.1 (2H, m, COCH₂), 3.5 (1H, m, COCH₂CH), 3.75 (6H, s, COOCH₃), 7.6 – 8.0 (4H, m, CH arom.). HPLC: 99 %
25 purity.

Step F:**Synthesis of N-[4-[[2,4-diamino-6-pteridiny]methyl]amino]benzoyl] pseudoglutamic ester (compound 3)**

30 A mixture of 7 g (27.4 mmol) 2,4-diamino-6-bromomethylpteridine **2** and 7 g (23.4 mmol) dimethyl N-[4-methyl-amino]benzoyl]pseudogluamate **1** in 70 ml N,N-

dimethylacetamide was stirred for 30 minutes at 70 °C then allowed to stand at room temperature overnight protected from light, then heated again for 10 minutes to 100 °C. The reaction was controlled by TLC. After cooling, the reaction mixture was poured into water acidified with AcOH at pH 4 (1000 ml). The dark-yellow precipitate that formed was filtered and washed three times with water and allowed to air dry. 2.6 g orange-yellow product **3** was obtained, m.p. 200 – 210 °C. The filtrate was treated with 10 % NaHCO₃ and the precipitate formed was separated in the same manner, resulting in a second fraction of dimethyl ester **3** (2 g). Total yield: 36 %. TLC (4:1 chloroform-methanol) showed a single spot, R_f 0.48. UV Spectra: 210, 242, 332 (0.1 N HCl); 238, 335 (MeOH).

Example 2

Synthesis of N-[4-[[2,4-diamino-6-pteridinyl)methyl]amino]benzoyl]pseudoglutamic acid (compound 4)

1 g (2.1 mmol) dimethyl ester **3** was added in portions to a solution of 10 ml 2N NaOH and 25 ml ethanol, and the mixture was stirred at room temperature for 4 hours. The precipitate formed was filtered and dissolved in distilled water. The alkaline solution was treated with charcoal, filtered and the pH was adjusted to 4.5 with 10 % AcOH. The precipitate was filtered and washed with water at pH 4.5, then with acetone, resulting in 0.8 g compound **4** (yield 85 %). The product, a brown solid, was purified by preparative HPTLC (High Performance Thin Layer Chromatography). After elution with 50:50:5 CH₃CN-H₂O-NH₄OH, the diacid was extracted from silica gel with 100 ml NaOH solution at pH 8. The water was removed by freeze-drying. TLC (7:2:1 CH₃CN-H₂O-NH₄OH) showed a single spot, R_f 0.80. Mass spectrum: m/z 120 (M⁺, 100%). IR spectra (KBr): 1651 (C=O), 1594 (C=C), 1563, 1403 (C=O acid), 1176 (C-O), 823 (CH). UV spectra: 242, 332 nm (0.1 N HCl); 232, 259, 325 nm (0.1 N NaOH); 229, 262, 318 (MeOH). ¹H NMR: 1.6 (3H, m, CH₃-CH₂), 2.2 (2H, t, CH₂CH₂COOH), 2.9 (2H, m, COCH₂), 4.6 (2H, s, CH₂NH), 6.8 – 7.8 (4H, m, CH arom.), 9 (1H, s, 7-CH). HPLC: 97 % purity.

Other compounds of formula I can be produced by adapting the procedures set out above in an appropriate manner. For example, intermediate **1** and analogous compounds can be converted to their corresponding N-methyl derivatives by

reaction with formaldehyde and sodium cyanoborohydride. Intermediate 6 can also be produced by reacting ethyl cyanoacetate with ethyl acrylate according to standard procedures.

5 **Example 3**

***In vitro* inhibition of DHFR**

The ability of compounds of formula I to inhibit dihydrofolate reductase (DHFR) *in vitro* was measured using a standard DHFR enzyme inhibition assay. DHFR enzyme was purified from rat livers or the commercially available DHFR was used, this
 10 being produced by recombinant expression in *E. coli*. Assays of enzyme activity were performed at 37 °C by monitoring changes in UV absorbance at 340 nm of a solution containing 50 mM *N*-tris(hydroxymethyl)methyl-2-aminoethanesulfonic acid (pH 7.0), 1 mM EDTA, 75 µM 2-mercaptoethanol, 0.1 % bovine serum albumin, 20 µM dihydrofolate, and 100 µM NADPH. Reactions were initiated by
 15 adding dihydrofolate. Each titration of the inhibitor was performed twice, and mean DHFR activity was plotted against inhibitor concentration to obtain IC₅₀ values. The ratio of IC_{50(Compound)}/ IC_{50(MTX)} was designated relative IC₅₀ and the results of one representative experiment (out of 5 experiments) for compounds 438 and 497 relative to methotrexate (MTX) are shown in Table 1. The results obtained
 20 overall suggest that most compounds of formula I tested possessed an *in vitro* potency similar to that of MTX.

Table 1.

Compound	Chemical formula	Relative IC ₅₀
MTX		1
438	Compound 4, R ³ = H	12.5
497	Compound 4	0.75

25

N.B. Compound 438 has the same structure as compound 4 in Figure 1 except that R³ (-CH₂CH₂CO₂H) is replaced by H.

Example 4

***In vitro* cytotoxicity**

The cytotoxicity of compounds of formula I against a number of tumor cell lines (CCRF-CEM, HepG2, HeLa, KB, L1210, A549 and COLO205) was assayed by measuring cell viability at different time points following drug addition up to 72 hours, and compared with the corresponding cytotoxicity of methotrexate and pemetrexed (obtained as Alimta®). The compounds of formula I demonstrated potent inhibitory effects against growth of all the cell lines tested, with the strongest inhibitory effects being seen against the L1210 cell line. Compared with methotrexate and pemetrexed, the compounds of formula I exhibited similar or stronger inhibitory effects against all the cell lines, and in some cases the onset of cytotoxicity was more rapid. Further, compounds of formula I wherein B = -CH₂NH- were found to be particularly cytotoxic.

Cytotoxicity was not reversed by addition of purines, such as hypoxanthine, or by addition of aminoimidazole carboxamide (up to very high concentration), but was reversed by addition of leucovorin, indicating cytotoxicity was due to antagonism of a folate-related mechanism. Consistent with a proposed mechanism of action in which DHFR is the main target, the addition of thymidine reversed cytotoxicity induced by the compounds of formula I only at high concentrations. These effects indicate specific inhibition of de novo purine synthesis and a less significant inhibition of the thymidylate cycle, however more pronounced than in the case of methotrexate. The compounds of formula I also inhibited glycineamide ribonucleotide transformylase in a comparable concentration range to methotrexate.

Example 5

Tumor inhibition in animal models *in vivo*

The ability of compounds of formula I to inhibit tumor growth in mice was tested as follows. 5 x 10⁶ L 1210 cells were injected subcutaneously in the axillary region of DBA/2 mice (groups of 8 mice/treatment). Following intraperitoneal administration of saline solution only or saline solution containing a compound of formula I the length and width of the control tumor (receiving only saline) was measured at the indicated time and compared to those of animals receiving test

compound to calculate the percentage of inhibition. Intraperitoneal administration of a saline solution containing the test compound, daily for 6 days, led to 60 % tumor-free long term survivors (tumor weight zero). Median survival times for saline-treated control animals and animals receiving Compound 4 (0.5 mg/kg) were 6.7 and 15.6 days, respectively. Oral administration of the compound required a higher dosage of the inhibitor and led to a less-marked, but still significant reduction of the tumor weight and 25 % long-term survivors as compared with the saline-treated control group.

Compounds of formula I were also active *in vivo* against the carcinoma W256 (TGI = 28 %), most likely due to the higher solubility and thus passive transport into the tumor compared with other anti-folate agents. Median survival times for saline-treated control animals and animals receiving Compound 4 (1 mg/kg) intraperitoneally in the Walker-256 rat tumor model were 22.5 and > 46.3 days, respectively.


The tumor evolution was measured following different treatment regimes and results are summarized in Table 2.

Table 2. Effect of various concentration of Compound 4 in a rat tumor model

Group	Treatment	No. animals surviving at day 30 days post-transplant	Tumor volume (cm ³) $\pm$ SE
1	Saline Control	2/10	37.3 $\pm$ 0.3
2	0.5 mg/kg – 3 doses every 4 days	9/10	13.4 $\pm$ 3.07
3	0.25 mg/kg – 3 doses every 4 days	8/10	23 $\pm$ 3.5
4	0.1 mg/kg – 5 doses every day	10/10	14.5 $\pm$ 3.8
5	1 mg/kg – one dose	8/10	17.1 $\pm$ 3.3

Claims

1. A compound of the formula I, or a pharmaceutically acceptable salt thereof:



wherein:

Z = O or S;

n = 1 - 3;

$R^3 = -CO_2R^8, -C(O)SR^8, -C(O)NHR^8, -C(S)OR^8, -C(S)SR^8, -C(S)NHR^8,$
 $-C(NH)SR^8$ or $-C(NH)NHR^8,$

wherein R^8 is -H or alkyl;

$R^4 = -H, -CH_2R^5$ or $-CH_2CH_2R^5,$


wherein R^5 independently has one of the meanings of R^3 ;

$B = -NR^2-, -CH_2NR^2-, -CH_2CH_2NR^2-, -CH_2CHR^7-$ or $-CH_2O-$,

wherein R^2 is H or a C_{1-3} alkyl, alkenyl or alkynyl group, and

R^7 is H or a C_{1-3} alkyl or alkoxy group;


A =





wherein $R^1 = -NH_2$ or $-OH,$

C and D are each, independently, a 5- or 6-membered, substituted or unsubstituted, aromatic or non-aromatic ring which may also contain one or more heteroatoms, and C is connected to group B in any available position.


2. A compound as claimed in claim 1, wherein Z is O.
3. A compound as claimed in claim 1 or 2, wherein n is 1.
4. A compound as claimed in any one of claims 1 to 3, wherein R^8 is - H, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl or tertiary butyl.
5. A compound as claimed in any one of claims 1 to 4, wherein R^3 is - CO_2R^8 and R^4 is H or - $CH_2CH_2CO_2R^8$.
6. A compound as claimed in claim 5, wherein R^8 is H, methyl or ethyl.
7. A compound as claimed in any one of claims 1 to 6, wherein B is - CH_2NR^2 -, - CH_2CHR^7 - or - CH_2O -.
8. A compound as claimed in any one of claims 1 to 7, wherein R^2 is - H, -Me, - Et or - $CH_2C \equiv CH$.
9. A compound as claimed in any one of claims 1 to 8, wherein R^7 is - H, -Me, - Et or -OMe.
10. A compound as claimed in any one of claims 1 to 9, wherein D is a 5-membered aromatic ring.
11. A compound as claimed in claim 10, wherein A is



12. A compound as claimed in claim 10, wherein A is of the formula A-i or A-ii:



13. A compound as claimed in any one of claims 1 to 12, wherein C is:





wherein X is CH or N, and either:

Y is C and R⁶ is H, Me, Et or HCO; or

Y is N and R⁶ is a lone pair of electrons.

14. A compound as claimed in claim 13, wherein A is of the formula A-i-1 or A-ii-1:



15. A compound as claimed in claim 14, wherein B is $-\text{CH}_2\text{NR}^2-$; R^2 is -H, -Me, -Et or $-\text{CH}_2\text{C}\equiv\text{CH}$; Z is O; n is 1; and R^3 is $-\text{CO}_2\text{R}^8$.


16. A compound as claimed in claim 13, wherein B is $-\text{CH}_2\text{NR}^2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CHCH}_3-$ or $-\text{CH}_2\text{O}-$; R^2 is -H, -Me, -Et or $-\text{CH}_2\text{C}\equiv\text{CH}$; Z is O; n is 1; R^3 is $-\text{CO}_2\text{R}^8$; R^4 is -H or $-\text{CH}_2\text{CH}_2\text{CO}_2\text{R}^8$; R^8 is independently -H, -Me or -Et; if A is A-i and Y is C, R^6 is -H; and if A is A-ii, R^1 is -OH.

17. A compound as claimed in any one of claims 1 to 13 and 16, wherein R^4 is $-\text{CH}_2\text{CH}_2\text{CO}_2\text{R}^8$.


18. A compound as claimed in any one of claims 1 to 13, 16, and 17 wherein B is $-\text{CH}_2\text{NR}^2-$.

19. A method of preparing a compound as claimed in claim 1, which comprises the step of:

(a) reacting a compound of the formula II



wherein A is as defined in claim 1, m is 0, 1 or 2 and X is a leaving group, with a compound of the formula III




wherein Z, n, R^2 , R^3 and R^4 are as defined in claim 1;

(b) reacting a compound of the formula IV




wherein A is as defined in claim 1 and X is a leaving group, with a compound of the formula V




wherein Z, n, R³ and R⁴ are as defined in claim 1; or

(c) converting one of the following compounds VI or VII into a corresponding organometallic reagent




wherein A, Z, n, R³, R⁴, and R⁷ are as defined in claim 1, and Y is, in each case independently, a halide, and reacting said reagent with the other of compounds VI or VII.


20. A method of preparing a compound as claimed in claim 1 having the formula I-A or a pharmaceutically acceptable salt thereof:



wherein n, R¹, R³ and R⁴ are as defined in claim 1, which comprises reacting a compound of the formula II-A




wherein X is Cl, Br or I, with a compound of the formula III-A




21. A compound of the formula V or VII, as defined in claim 19, wherein R^4 is $-CH_2R^5$ or $-CH_2CH_2R^5$.
22. A compound of the formula III, as defined in claim 19, wherein R^4 is $-CH_2R^5$ or $-CH_2CH_2R^5$, and R^2 is a C_{1-3} alkyl, alkenyl or alkenyl group.
23. A compound of the formula III, as defined in claim 19 or 22, wherein R^4 is $-CH_2R^5$ or $-CH_2CH_2R^5$, and Z is S.
24. A compound of the formula III, as defined in any one of claims 19, 22, and 23, wherein R^4 is $-CH_2R^5$.
25. A compound of the formula III, as defined in any one of claims 19 and 22 to 24, wherein R^3 is $-C(O)SR^8$, $-C(O)NHR^8$, $-C(S)OR^8$, $-C(S)SR^8$, $-C(S)NHR^8$, $-C(NH)SR^8$ or $-C(NH)NHR^8$.
26. A compound of the formula III, as defined in any one of claims 19 and 22 to 25, wherein R^5 is $-C(O)SR^8$, $-C(O)NHR^8$, $-C(S)OR^8$, $-C(S)SR^8$, $-C(S)NHR^8$, $-C(NH)SR^8$ or $-C(NH)NHR^8$.


27. A compound of the formula I-A1, or a pharmaceutically acceptable salt thereof:




28. A compound of the formula I-A2, or a pharmaceutically acceptable salt thereof:




29. A compound of the formula I-A3, or a pharmaceutically acceptable salt thereof:




30. A compound of the formula I-A4, or a pharmaceutically acceptable salt thereof:



31. A compound of the formula I-A5, or a pharmaceutically acceptable salt thereof:



32. A compound of the formula I-A6, or a pharmaceutically acceptable salt thereof:



33. A compound as claimed in any one of claims 1 to 18 and 23 to 32, for use in therapy.
34. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 18 and 23 to 33 and a pharmaceutically acceptable excipient.
35. Use of a compound as claimed in any one of claims 1 to 18 and 23 to 32, or the composition of claim 34, for the manufacture of a medicament for use in the treatment of a condition responsive to inhibition of an enzyme dependent upon folic acid or a folic acid derivative.
36. Use of a compound as claimed in any one of claims 1 to 18 and 23 to 32, or the composition of claim 34, for the manufacture of a medicament for use in the treatment of cancer.

37. Use of a compound as claimed in any one of claims 1 to 18 and 23 to 32, or the composition of claim 34, for the treatment of a condition responsive to inhibition of an enzyme dependent upon folic acid or a folic acid derivative.

38. Use of a compound as claimed in any one of claims 1 to 18 and 23 to 32, or the composition of claim 34, for the treatment of cancer.

Figure 1

