



- (51) **International Patent Classification:**
C12N 15/29 (2006.01)
- (21) **International Application Number:**
PCT/US2013/071340
- (22) **International Filing Date:**
21 November 2013 (21.11.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/729,233 8 January 2013 (08.01.2013) US
- (71) **Applicant:** ACADEMIA SINICA; No. 128, Sec. 2, Academia Road, Nankang, Taipei, 115 (TW).
- (72) **Inventor; and**
- (71) **Applicant :** HO, Tuan-Hua, David [US/—]; 1512 Baxter Lane Court, Chesterfield, MO 63017 (US).
- (72) **Inventors:** YU, Su-May; 4th F, No. 18, Alley 3, Lane 61, Sec. 2, Academia Road, Nankang, Taipei, 115 (TW). CHEN, Yi-Shih; No.8-10, Kezhuang, Yuanli Township, Miaoli County, 358 (TW). LO, Shuen-Fang; No. 24, Alley 2, Lane 2, Gongye Road, Dali City, Taichung County, 412 (TW).
- (74) **Agent:** KING, Anthony; WPAT, P.C., 1100 Quail Street, Suite 202, Newport Beach, CA 92660 (US).
- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))


[Continued on next page]

(54) **Title:** METHODS FOR ENHANCING ROOT GROWTH OF PLANTS(A) **Fig. 1****ABRC321**

gggtaccGCAACGCGTGTCTCCTACGTGGCGGCGcgagATTGCCACCGGctctaga

CE3 A2 CE1

(B)



(C)



(57) **Abstract:** The present invention relates to a method for enhancing root growth of a plant by introducing a polynucleotide encoding late embryogenesis abundant protein, group 3 (LEA3) into the plant. Plant root architecture is essential for its functions in water and nutrient uptake, anchorage and interactions with microbes in the soil.



— with sequence listing part of description (Rule 5.2(a))

TITLE OF THE INVENTION

METHODS FOR ENHANCING ROOT GROWTH OF PLANTS

TECHNOLOGY FIELD

5 **[0001]** The present invention relates to a method for enhancing root growth of plants. More particularly, the invention relates to a method for enhancing root growth of plants by introducing a polynucleotide encoding group 3 late embryogenesis abundant protein (LEA3) into the plant.

10 BACKGROUND OF THE INVENTION

[0002] Plant root architecture is essential for its functions in water and nutrient uptake, anchorage and interactions with microbes in the soil.

[0003] Studies in Arabidopsis have greatly advanced our knowledge on mechanisms controlling root development (Potters G *et al.*, (2007) *Trends Plant Sci* 12(3):98-105; Peret B, *et al.* (2009) *Trends Plant Sci* 14(7):399-408; Lavenus J, *et al.* (2013) *Trends Plant Sci* 18(8):450-458); however, similar studies in cereals are relatively scarce (Coudert Y *et al.*, (2010) *Trends Plant Sci* 15(4):219-226). Unlike Arabidopsis which has a primary root that iteratively branches to generate several orders of lateral roots, the cereals have several types of branched roots including shoot-born crown roots and root-borne lateral roots. Approximately 675 quantitative trait loci (QTLs) control 29 root parameters in rice (Courtois B, *et al.* (2009) *Rice* 2(2-3):115-128). Several homologous genes that play similar roles regulating root formation between Arabidopsis and cereals have been identified; however, distinct hormonal and developmental pathways are also found to be involved in root formation in cereals (Orman-Ligeza B, *et al.* (2013) *Trends Plant Sci* 18(8):459-467). In Arabidopsis, six classical hormones control primary root growth by targeting cells in distinct tissues (Ubeda-Tomas S *et al.*, (2012) *Trends Plant Sci* 17(6):326-331). Among these hormones, auxin is shown to act as a common integrator to many endogenous and environmental signals regulating lateral root development in both dicots and monocots (Ubeda-Tomas S *et al.*, *supra*; Laurie S *et al.*, *J Exp Bot* 54(383):739-747). In Arabidopsis, the universal stress hormone abscisic acid (ABA) stimulates main root elongation in response to drought and osmotic stresses; however, ABA and auxin signals act antagonistically during lateral root initiation, with ABA as a repressing while auxin a promoting agent (De Smet I *et al.*, (2006) *Trends Plant Sci*

11(9):434-439).

[0004] LEA proteins are a set of proteins highly accumulated at the onset of seed desiccation and in response to water deficit in plant vegetative tissues (Dure L (1981) *Biochemistry* 20:4162-4168; Dure L (1992) *Control of Plant Gene Expression*. CRC Press, Boca Raton, FL, pp. 325-335). LEA proteins have been classified into six groups based on conservation in amino acid sequence domains and expression patterns (Dure L, 3rd (1993) *Plant J* 3(3):363-369; Wise MJ (2003) *BMC Bioinformatics* 4:52). HVA1 is a group 3 LEA (LEA3) protein specifically expressed in barley aleurone layers and embryos during late seed development undergoing desiccation (Jefferson RA (1987) *Plant Mol Biol Rep* 5:387-405). HVA1 contains an 11-amino acid consensus motif which is repeated 9 times, forming an α -helical dimer suitable for accommodating positively and negatively charged ions, thus has been proposed to function as an ion sequester (Liang Y & Harris JM (2005) *Am J Bot* 92(10):1675-1683). Functions of HVA1 in protection against environmental stresses have been reported (Hong B *et al.*, (1992) *Plant Mol Biol* 18:663-674; Sutton F *et al.*, *Plant Physiol* 99(1):338-340). However, no one reports HVA1's function on promoting root growth of plants.

BRIEF SUMMARY OF THE INVENTION

[0005] In the present invention, it is unexpectedly found that a transgenic plant transformed with group 3 late embryogenesis abundant protein (LEA3) gene exhibits surprisingly improved root growth as compared with a control (wild type) plant without transformation with LEA3 gene. Compared with control plants, the transgenic plants of the invention exhibit significant increase in elongation, number, and density of primary and branch root system and initiation of lateral root. Especially, the transgenic plants exhibit improved root growth in the presence of stress hormone abscisic acid (ABA) or under nutrient deficient conditions.

[0006] Therefore, the present invention provides a method for enhancing root growth of a plant, comprising

- (i) introducing a polynucleotide encoding group 3 late embryogenesis abundant protein (LEA3) into a plant cell to obtain a transformed plant cell, wherein the polynucleotide is operably linked to an expression control sequence;
- (ii) producing a transformed plant from said transformed plant; and
- (iii) selecting a transformed plant exhibiting improved root growth compared to a control plant which is not introduced with the polynucleotide.

[0007] In some certain embodiments, the expression control sequence comprising a promoter sequence, which is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, preferably SEQ ID NO: 3.

[0008] In some certain embodiments, the LEA3 protein is the one originated from
5 barely or rice, for example, having the amino acid sequence of SEQ ID NO: 4 or 5.

[0009] In some certain embodiments, the transgenic plant is monocotyledon, including but not limited to rice, barley, wheat, rye, oat, corn, bamboo, sugar cane, onion, leek and ginger.

[0010] In some certain embodiments, the transgenic plant is dicotyledon, including
10 but not limited to Arabidopsis, eggplant, soybean, mung bean, kidney bean, pea, tobacco, lettuce, spinach, sweet potato, carrot, melon, cucumber and pumpkin.

[0011] To perform the method of the invention, it is preferably to select the transgenic plants with improved root growth, in the presence of stress hormone abscisic acid (ABA) or under nutrient deficient conditions.

[0012] The details of one or more embodiments of the invention are set forth in the
15 description below. Other features or advantages of the present invention will be apparent from the following detailed description of several embodiments, and also from the appending claims.

20 BRIEF DESCRIPTION OF THE DRAWINGS

[0013] The foregoing summary, as well as the following detailed description of the invention, will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, there are shown in the drawings embodiments which are presently preferred. It should be understood,
25 however, that the invention is not limited to the precise arrangements and instrumentalities shown.

[0014] In the drawings:

[0015] Fig. 1 shows (A) the nucleotide sequence of *ABRC321* containing CE3 and A2 from HVA1 ABRC3 and CE1 from HVA22 ABRC1; (B) constructs contain 0-3
30 tandem repeats of *ABRC321* for GUS expression (the lengths of DNA fragments constituting the expression cassette being: *ABRC321* (including ABRE, linker and enzyme restriction sites), 56 bp; *Amy64* minimal promoter (mini P), 99 bp; Intron1-exon2-intron2 (In1-Ex2-In2) of *HVA22*, 240 bp; *GUS* cDNA, 2 kb; *HVA22* terminator (3'), 120 bp; and (C) constructs containing 3 tandem repeats of *ABRC321*

for rHVA1 expression (lengths of *HVA1* cDNA and *Nos* 3' being 630 and 300 bp, respectively, SEQ ID NO: 6).

[0016] Fig. 2 shows that *ABRC321* is an ABA-inducible synthetic promoter with low-background but high ABA-inducible activity in leaves and roots in transgenic rice.

- 5 (A) Three copies of *ABRC321* confer high ABA inducibility in a rice half-seed transient expression assay. Rice half seeds were transfected with plasmids shown in Fig. 1 by particle bombardment, and incubated with (+) or without (-) 10 μ M ABA for 16 h, and GUS activity was analyzed. (B) Three copies of *ABRC321* confer high ABA inducibility in leaves and roots in transgenic rice. Constructs shown in Fig. 1
10 were introduced into the rice genome. Leaves and roots of 10-day-old T2 seedlings of four transgenic lines were collected and incubated in water with (+) or without (-) 10 μ M ABA for 16 h, and GUS activity was analyzed. Numbers above bars indicate fold induction of GUS activity by ABA. Error bars represent SD (n = 3).

- [0017]** Fig. 3 shows that *3xABRC321* has low-background but high
15 ABA-inducibility in leaves (A) and roots (B) in transgenic rice. (B1) and (B2): without ABA treatment; (B3) and (B4): with ABA treatment. Abbreviation for leaf tissues: Bc: bulliform cell; Bs: bundle sheath; Cc: companion cell; Ec: epidermal cell; Mc: mesophyll cell; Ms: mestome sheath; Sc: sclerenchyma; Se: sieve element; Ve: vessel element. (C1) longitudinal sections of roots; (C2) a LRP indicated in the red
20 box in (C1); (C3): two newly initiated LRP from a crown root; (C4) cross section of crown roots; (C5): crown root tip indicated in the green box in (C1); (C6): a LRP from a primary lateral root of a crown root; (C7): a LRP from a primary lateral root. Asterisk indicates GUS is also expressed in cortex and exodermis cells nearby LRP. Abbreviation for root tissues: C: cortex; Crc: columella root cap; En: endodermis; Ep:
25 epidermis; Ex: exodermis; Lrc: lateral root cap; LRP: lateral root primorida; Mx: metaxylem vessel; P, pericycle; Pc: procambium; QC: quiescent center; Sc: sclerenchyma.

- [0018]** Fig. 4 shows that amino acid sequence of barley HVA1 (SEQ ID NO: 3) is homologous to that of the rice LEA3 (SEQ ID NO: 4). Amino acid alignment of
30 barley HVA1 and rice LEA3 using Vector NTI 9.0 software. The light gray and dark gray highlights indicate identical and similar amino acid sequences, respectively. The underline indicates the tandem 11-mer conserved amino acid repeats.

[0019] Fig. 5 shows that rHVA1 expression regulated by *3xABRC321* has similar tissue-specificity but at higher-level than the endogenous HVA1 in transgenic rice in

response to ABA. (A) Cross sections of root elongation zone. (B) Longitudinal sections of root elongation zone. (C) Longitudinal sections of root tip. F: image of fluorescence field. C: composite images of fluorescence and transmission fields. P: pericycle. Scale bar represents 50 μ m.

- 5 [0020] Fig. 6 shows *3xABRC321:HVA1* induces lateral root growth in transgenic rice in the presence of ABA. Three-day-old seedlings of Tainung 67 (T) transgenic lines were transferred to half-strength MS medium without sugar but containing ABA or sorbitol, and root morphology was examined. (A) Morphology of various branch roots. (B) Treatments with 0.1-2.0 μ M ABA for 14 days, wherein the growth of
10 secondary lateral roots (arrowheads) was observed in transgenic seedlings treated with 2.0 μ M ABA.

[0021] Fig. 7 shows *3xABRC321:HVA1* induces root growth in transgenic rice at ABA concentration of 0.1 μ M. Left panel: crown and adventitious roots. Right panel: seminal root.

- 15 [0022] Fig. 8 shows that expression of *3xABRC321:HVA1* induces root growth in transgenic lines under nutrient deficient conditions. Four-day-old seedlings of Tainung 67 (T) and Kitaake (K) transgenic lines were germinated on MS medium for four days and transfer to full strength Yoshida solution containing full nutrients, nutrients diluted 10,000-fold (nutrient starvation), full nutrients minus nitrogen (N),
20 and full nutrients minus phosphorus (P) for 21 days. (A) Root morphology. (B) Lateral root density. (C) Total root number.

- [0023] Fig. 9 shows that rHVA1 expression controlled by *3xABRC321* induces root growth in transgenic lines under nutrient deficiency. Three-day-old seedlings of Tainung 67 (T) or Kitaake (K) wild type (WT) and transgenic lines were transferred to
25 Yoshida solutions containing (A) full nutrients, (B) nutrients diluted 10,000-fold (nutrient starvation), (C) full nutrients minus nitrogen (N), and (D) full nutrients minus phosphorus (P) for 18 days. Left panel: crown and adventitious roots. Right panel: seminal root. Significance levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

- [0024] Fig. 10 shows that *3xABRC321:HVA1* promote lateral root initiation and
30 growth in transgenic rice. Three-day-old seedlings of Tainung 67 (T) transgenic lines were transferred to half-strength MS ($\frac{1}{2}$) medium without sugar but containing 0.2 μ M ABA for 11 days. Roots were treated with ethanol and the number of lateral root primordia (LRP) and lateral roots (LR) in three different zones were counted

under microscope. Zone 1 (upper part) and Zone 2 (medium part) cover maturation zone and Zone 3 (bottom part) covers elongation and division zones of the root system. (A) Wild type plants. (B) The transgenic plants. Error bars represent SD ($n = 12$). Significance levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

5 [0025] Fig. 11 shows that auxin transporters are up-regulated in *3xABRC321:HVA1* transgenic rice. Four-day-old seedlings of Tainung 67 (T) transgenic lines were germinated on MS medium for four days and transfer to full strength Yoshida solution with or without 0.2 μM ABA for 14 days. Total RNA was extracted from roots and subjected to quantitative real-time RT-PCR.

10 [0026] Fig. 12 shows that induction of root growth by *3xABRC321:HVA1* in the presence ABA requires auxin. Three-day-old seedlings of Tainung 67 (T) transgenic lines that containing seminal roots only were transferred to half-strength MS ($1/2$) medium without sugar but containing ABA and/or NPA for 11 days, and root morphology was examined. (A) Medium only, (B) 0.2 μM ABA only, (C) 1 μM NPA only, (D) 0.2 μM ABA plus 1 μM NPA. The dots indicate starting points of root growth after transferring to $1/2$ MS medium containing ABA and/or NPA.

[0027] Fig. 13 shows that rHVA1 expression regulated by *3xABRC321* enhances WUE and grain yield in transgenic lines grown in non-irrigated field. (A) Three-day-old seedlings of Tainung 67 (T) and Kitaake (K) transgenic lines were
20 treated with two cycles of 250 mM sorbitol (3 days) and water (5 days) to induce root growth. Seedlings were then grown in Yoshida solution and transferred to same amount of fresh solution every two days. Total water consumption was measured up to 19 days, and whole-plant dry weight was measured. Panel 1, dry weight of 10 plants at the end of experiment; Panel 2, total water use of 10 plants; Panel 3, WUE
25 determined by dividing total plant dry weight with total water use. (B) Kitaake transgenic lines were grown in irrigated and non-irrigated fields during August to December in 2011. Grain yield was determined after harvest. Error bars represent SD ($n = 10$). Significance levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

30 DETAILED DESCRIPTION OF THE INVENTION

[0028] Unless defined otherwise, all technical and scientific terms used herein have the same meanings as is commonly understood by one of skill in the art to which this invention belongs.

[0029] As used herein, the articles “a” and “an” refer to one or more than one (*i.e.*,

at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0030] The term “polynucleotide” or “nucleic acid” refers to a polymer composed of nucleotide units. Polynucleotides include naturally occurring nucleic acids, such as deoxyribonucleic acid (“DNA”) and ribonucleic acid (“RNA”) as well as nucleic acid analogs including those which have non-naturally occurring nucleotides. Polynucleotides can be synthesized, for example, using an automated DNA synthesizer. The term “nucleic acid” typically refers to large polynucleotides. It will be understood that when a nucleotide sequence is represented by a DNA sequence (i.e., A, T, G, C), this also includes an RNA sequence (i.e., A, U, G, C) in which “U” replaces “T.” The term “cDNA” refers to a DNA that is complementary or identical to an mRNA, in either single stranded or double stranded form.

[0031] The term “complementary” refers to the topological compatibility or matching together of interacting surfaces of two polynucleotides. Thus, the two molecules can be described as complementary, and furthermore the contact surface characteristics are complementary to each other. A first polynucleotide is complementary to a second polynucleotide if the nucleotide sequence of the first polynucleotide is identical to the nucleotide sequence of the polynucleotide binding partner of the second polynucleotide. Thus, the polynucleotide whose sequence 5'-TATAC-3' is complementary to a polynucleotide whose sequence is 5'-GTATA-3'.”

[0032] The term “encoding” refers to the inherent property of specific sequences of nucleotides in a polynucleotide (e.g., a gene, a cDNA, or an mRNA) to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Therefore, a gene encodes a protein if transcription and translation of mRNA produced by that gene produces the protein in a cell or other biological system. It is understood by a skilled person that numerous different polynucleotides and nucleic acids can encode the same polypeptide as a result of the degeneracy of the genetic code. It is also understood that skilled persons may, using routine techniques, make nucleotide substitutions that do not affect the polypeptide sequence encoded by the polynucleotides described there to reflect the codon usage of any particular host organism in which the polypeptides are to be expressed. Therefore, unless otherwise specified, a “nucleotide sequence encoding an amino acid sequence” includes all

nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. Nucleotide sequences that encode proteins and RNA may include introns.

[0033] The term “recombinant nucleic acid” refers to a polynucleotide or nucleic acid having sequences that are not naturally joined together. A recombinant nucleic acid may be present in the form of a vector. “Vectors” may contain a given nucleotide sequence of interest and a regulatory sequence. Vectors may be used for expressing the given nucleotide sequence or maintaining the given nucleotide sequence for replicating it, manipulating it or transferring it between different locations (e.g., between different organisms). Vectors can be introduced into a suitable host cell for the above mentioned purposes.

[0034] As used herein, the term “operably linked” may mean that a polynucleotide is linked to an expression control sequence in such a manner to enable expression of the polynucleotide when a proper molecule (such as a transcriptional factor) is bound to the expression control sequence.

[0035] As used herein, the term “expression control sequence” or “regulatory sequence” means a DNA sequence that regulates the expression of the operably linked nucleic acid sequence in a certain host cell.

[0036] Examples of vectors include, but are not limited to, plasmids, cosmids, phages, YACs or PACs. Typically, in vectors, the given nucleotide sequence is operatively linked to the regulatory sequence such that when the vectors are introduced into a host cell, the given nucleotide sequence can be expressed in the host cell under the control of the regulatory sequence. The regulatory sequence may comprises, for example and without limitation, a promoter sequence (e.g., the cytomegalovirus (CMV) promoter, simian virus 40 (SV40) early promoter, T7 promoter, and alcohol oxidase gene (*AOXI*) promoter), a start codon, a replication origin, enhancers, an operator sequence, a secretion signal sequence (e.g., α -mating factor signal) and other control sequence (e.g., Shine-Dalgarno sequences and termination sequences). Preferably, vectors may further contain a marker sequence (e.g., an antibiotic resistant marker sequence) for the subsequent screening procedure. For purpose of protein production, in vectors, the given nucleotide sequence of interest may be connected to another nucleotide sequence other than the above-mentioned regulatory sequence such that a fused polypeptide is produced and

beneficial to the subsequent purification procedure. Said fused polypeptide includes, but is not limited to, a His-tag fused polypeptide and a GST fused polypeptide.

[0037] Where the expression vector is constructed for a plant cell, several suitable promoters known in the art may be used, including but not limited to the Figwort mosaic virus 35S promoter, the cauliflower mosaic virus (CaMV) 35S promoter, the commelina yellow mottle virus promoter, the rice cytosolic triosephosphate isomerase (TPI) promoter, the rice actin 1 (Act1) gene promoter, the ubiquitin (Ubi) promoter, the rice amylase gene promoter, the adenine phosphoribosyltransferase (APRT) promoter of Arabidopsis, the mannopine synthase and octopine synthase promoters.

10 [0038] In certain embodiments, a promoter sequence as used in the invention is a synthetic promoter, having SEQ ID NO: 1 (1xABRC321), SEQ ID NO: 2 (1xABRC321) or SEQ ID NO: 3 (1xABRC321), preferably SEQ ID NO: 3 (1xABRC321).

[0039] To prepare a transgenic plant, it is preferably that the expression vector as
15 used herein carries one or more selection markers for selection of the transformed plants, for example, genes conferring the resistance to antibiotics such as hygromycin, ampicillin, gentamycin, chloramphenicol, streptomycin, kanamycin, neomycin, geneticin and tetracycline, URA3 gene, genes conferring the resistance to any other toxic compound such as certain metal ions or herbicide, such as glufosinate or
20 bialaphos.

[0040] As used herein, the term “transgenic plant” or “transgenic line” refers to a plant that contains a recombinant nucleotide sequence. The transgenic plant can be grown from a recombinant cell.

[0041] A variety of procedures that can be used to engineer a stable transgenic
25 plant are available in this art. In one embodiment of the present invention, the transgenic plant is produced by transforming a tissue of a plant, such as a protoplast or leaf-disc of the plant, with a recombinant *Agrobacterium* cell comprising a polynucleotide encoding a desired protein (e.g. LEA3) and generating a whole plant from the transformed plant tissue. In another embodiment, a polynucleotide
30 encoding a desired protein can be introduced into a plant via gene gun technology, particularly if transformation with a recombinant *Agrobacterium* cell is not efficient in the plant.

[0042] The term “polypeptide” refers to a polymer composed of amino acid residues linked via peptide bonds. The term “protein” typically refers to relatively

large polypeptides. The term “peptide” typically refers to relatively short polypeptides.

[0043] As used herein, LEA3 protein is group 3 LEA proteins that are a set of proteins highly accumulated at the onset of seed desiccation and in response to water deficit in plant vegetative tissues. HVA1 is a LEA3 protein in barley, which has been identified to have an 11-amino acid consensus motif which is repeated 9 times, forming an α -helical dimer suitable for accommodating positively and negatively charged ions. In certain embodiments, the LEA3 protein as used herein is the one originated from barley (HVA1) or rice (OsLEA3), particularly have the amino acid sequence of SEQ ID NO: 4 or 5.

[0044] It is understandable that the LEA3 protein as used herein also include its biological equivalent, which means that there is a limited number of changes or modifications that may be made within a certain portion of the molecule irrelevant to the activity or function of the protein and still result in a molecule with an acceptable level of equivalent biological activity. Biologically equivalent polypeptides are thus defined herein as those polypeptides in which certain amino acid residues may be substituted. Polypeptides with different substitutions may be made and used in accordance with the invention. Modifications and changes may be made in the structure of such polypeptides and still obtain a molecule having similar or desirable characteristics. For example, certain amino acids may be substituted for other amino acids in the peptide/polypeptide structure without appreciable loss of activity. Amino acid substitutions are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. For example, arginine (Arg), lysine (Lys), and histidine (His) are all positively charged residues; and alanine (Ala), glycine (Gly) and serine (Ser) are all in a similar size. Therefore, based upon these considerations, arginine (Arg), lysine (Lys) and histidine (His); and alanine (Ala), glycine (Gly) and serine (Ser) may be defined as biologically functional equivalents. One can readily design and prepare recombinant genes for microbial expression of polypeptides having equivalent amino acid residues.

[0045] In some embodiments, the biological equivalent of LEA3 protein comprises the amino acid sequence having at least 50%, 60%, or 65%, or 70%, or 75%, or 80%, or 85%, or 90%, or 95% identity with the amino acid sequence of SEQ ID NO: 4 or 5,

and possess the conserved structure, i.e. an 11-amino acid consensus motif, which is repeated 9 or more times (such as 10 to 25 repeats, 10 to 20 repeats, or 10 to 15 repeats), forming an α -helical dimer suitable for accommodating positively and negatively charged ions.

5 **[0046]** To determine the percent identity of two amino acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid sequence for optimal alignment with a second amino acid sequence). In calculating percent identity, typically exact matches are counted. The determination of percent homology or identity between two sequences
10 can be accomplished using a mathematical algorithm known in the art, such as BLAST and Gapped BLAST programs, the NBLAST and XBLAST programs, or the ALIGN program.

[0047] As used herein, the term “improved root growth” refers to the growth of root system of a transgenic plant is improved or enhanced when compared with a
15 control plant that is not introduced with the same gene. Such improvement, for example, includes one or more characteristics of roots, e.g. increase in total number, length, density, dry weight and initiation, of primary root or secondary root, or crown root, lateral root, adventitious root, or seminal root.

[0048] The present invention is based on the unexpected finding that
20 transformation of LEA3 gene into a plant surprisingly promotes the root growth in the plant when compared with a control plant that is not introduced with the gene.

[0049] Therefore, the present invention provides a method for enhancing root growth of a plant, comprising

(i) introducing a polynucleotide encoding group 3 late embryogenesis abundant
25 (LEA3) protein into a plant cell to obtain a transformed plant cell, wherein the polynucleotide is operably linked to an expression control sequence;

(ii) producing a transformed plant from said transformed plant; and

(iii) selecting a transformed plant exhibiting improved root growth compared to a control plant which is not introduced with the polynucleotide.

30 **[0050]** According to the invention, the LEA3 protein can be the one originated from monocotyledon or dicotyledon.

[0051] Examples of monocotyledon includes but not limited to rice, barley, wheat, rye, oat, corn, bamboo, sugar cane, onion, leek and ginger.

[0052] Examples of monocotyledon includes but not limited to Arabidopsis, eggplant, soybean, mung bean, kidney bean, pea, tobacco, lettuce, spinach, sweet potato, carrot, melon, cucumber and pumpkin.

[0053] In particular examples, the LEA3 protein is the one from barley and rice,
5 more particularly having the amino acid sequence of SEQ ID NO: 4 or 5.

[0054] According to the present invention, the transgenic plants transformed with the LEA3 gene surprisingly exhibit improved root growth, even without ABA treatment, and have more significant improvement of root growth under ABA induction.

10 [0055] Therefore, in one certain embodiment, it is preferably to select the transgenic plants improved root growth, in the presence of ABA. Particularly, ABA is applied at a concentration from 0.1 to 5 μ M, more particularly, from 0.1 to 2 μ M, even more particularly from 0.1 to 0.5 μ M.

[0056] It is also found that the transgenic plants transformed with the LEA3 gene
15 exhibit improved root growth under nutrient deficient conditions.

[0057] As known in the art, normal nutrients of plant include major nutrients required in largest amount in plants, including nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg) and sulfur (S), and minor nutrients required in relatively smaller amount, including iron (Fe), copper (Cu), zinc (Zn),
20 boron (B), molybdenum (Mo), manganese (Mn) and chlorine (Cl). Additional mineral elements that are good for plants but not necessarily essential include sodium (Na), cobalt (Co), vanadium (Va), nickel (Ni), selenium (Se), aluminum (Al) and silicon (Si). In one embodiment, for hydroponic culture of rice seedlings, a commercial product, Yoshida's culture solution can be used, for example see From
25 IRRI (1976) Laboratory Manual for Physiological Studies of Rice. As used herein, nutrient deficient conditions for plants refers to a condition lacking one or more chemical elements for plant growth, such as lacking one or more of those as set forth above, particularly the major nutrients, e.g. nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg) and sulfur (S), more particularly nitrogen (N),
30 phosphorus (P), potassium (K), calcium (Ca), magnesium (Mg), and even more particularly nitrogen (N), phosphorus (P) and potassium (K), and/or the minor nutrients, e.g. iron (Fe), copper (Cu), zinc (Zn), boron (B), molybdenum (Mo), manganese (Mn) and chlorine (Cl), and/or additional mineral elements, e.g. sodium

(Na), cobalt (Co), vanadium (Va), nickel (Ni), selenium (Se), aluminum (Al) and silicon (Si). In some embodiments, nutrient deficient conditions also mean that the concentration of the chemical elements as set forth above is overly low, for example, less than 80%, 70%, 60%, 50%, 40%, 30%, 20%, or 10% of regular amount of total nutrients or each nutrient element as required for plant growth.

[0058] The present invention is further illustrated by the following examples, which are provided for the purpose of demonstration rather than limitation. Those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

[0059] 1. Materials and Methods

[0060] 1.1 Plant Materials

[0061] Rice cultivars *Oryza sativa* L. cv Tainung 67 and *Oryza sativa* L. cv Kitaake were used for all experiments. Plasmid was introduced into *Agrobacterium tumefaciens* strain EHA101 and rice transformation was performed as described (Chen P-W, *et al.* (2002) *J Biol Chem* 277:13641-13649). Homozygous transgenic lines were used in all experiments. For observation of root growth, transgenic rice seeds were germinated on the surface of half-strength MS medium without sugar but with or without ABA. For hydroponic culture, Yoshida solution was used. Seedlings were normally grown in 28 °C incubator with 12 hr daily light.

[0062] 1.2 Plasmids

[0063] Plasmid pAHC18 contains the luciferase (*Luc*) cDNA fused between the *Ubi* promoter and the *Nos* terminator (Bruce WB et al., (1989) *Proc Natl Acad Sci U S A* 86(24):9692-9696). Plasmid MP64 contains the barley *Amy64* minimal promoter (-60 relative to the transcription start site) and its 5' untranslated region (+57 relative to the transcription start site), *HVA22* intron1-exon2-intron2, the *GUS* coding region, and the *HVA22* 3'untranslated region (Shen Q & Ho TH (1995) *Plant Cell* 7(3):295-307). Plasmid QS115 contains a copy of *HVA22* ABRC1 fused upstream of the *Amy64* minimal promoter in plasmid MP64 (Shen Q & Ho TH, *supra*).

[0064] 1.3 Plasmid Construction

[0065] Two 56-bp complementary oligonucleotides, containing the CE3 and A2 elements from the *HVA1* promoter and the CE1 element from the *HVA22* promoter (Shen Q, Zhang P, & Ho TH (1996) *Plant Cell* 8(7):1107-1119) and restriction sites

KpnI, *XhoI* and *XbaI* were synthesized, annealed, and designated as *ABRC321* (Fig. 1A). *ABRC321* was self-ligated in two or three copies in correct orientations. For expression of GUS under the control of *ABRC321*, 1-3 copies *ABRC321* was inserted into the *XbaI* site in MP64, so that the *ABRC321* was fused upstream of the barley
 5 *Amy64* minimal promoter, generating constructs *1xABRC321-GUS*, *2x-ABRC321-GUS* and *3xABRC321-GUS* (Fig. 1(B)). For expression of *HVA1* under the control of *3xABRC321*, *GUS* cDNA in construct *3xABRC321-GUS* was replaced with *HVA1* cDNA, generating construct *3xABRC321-HVA1* (Fig. 1(C)) (SEQ ID NO: 6).

10 [0066] 1.4 GUS Activity Staining

[0067] Sections of leaf and root from 10-day-old seedlings were cut with Microslicers DTK-1000 (TED PELLA, Inc.), incubated in water containing or lacking 10 μ M ABA at 28 °C for 24 h, and subjected to histochemical staining with a buffer (0.1 M NaPO₄, pH 7.0, 10 mM EDTA, 0.1 % Triton X-100, 0.5 mM potassium
 15 ferriocyanide, pH 7.0 and 1 mM X-glucuronide) at 37 °C as described (Jefferson RA (1987) *Plant Mol Biol Rep* 5:387-405). After GUS staining, leave samples were incubated in 70 % ethanol at 65 °C for 1 h to remove chlorophyll.

[0068] 1.5 Immuno-histological Fluorescence Staining

[0069] Tissue localization of *HVA1* was examined by modification of a described
 20 method (Long SP et al., (2006) *Plant Cell Environ* 29(3):315-330). Rice roots were fixed with 2% paraformaldehyde (w/v) in 0.1 M NaPO₄ buffer, pH 7.0, and then embedded in 5% agar. Sections were sliced to 30- μ m thickness using Microslicers DTK-1000 and incubated in PBS buffer containing 0.3% (v/v) Triton X-100 (PBS-T). The nonspecific reaction was blocked with 5% (w/v) bovine serum albumin in PBS-T.
 25 Samples were then incubated with purified rabbit anti-barley *HVA1* polyclonal antibodies and subsequently with secondary antibodies (Alexa Fluor 555 goat anti-rabbit IgG; Molecular Probes). Samples were examined with a laser scanning confocal microscope. (LSM510 META; Zeiss).

[0070] 1.6 Measurement of Roots

30 [0071] Rice branch roots could be classified into four types: seminal root, also called radical, the first root grows during germination; crown roots, roots emerge from the node of coleoptile; adventitious roots, roots differentiate from the nodes of main stem and tillers, lateral roots, roots branch from above three types of roots and can bear

additional large or small lateral roots until the fifth order of branching. The number of each type of root was measured by simple counting. Root length was measured, number of roots was counted, and lateral root density was determined by dividing lateral root number with root length. Error bars represent SD ($n = 12$). Significance levels: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

[0072] 1.7 Field test

[0073] To evaluate grain yield and biomass production in the field, 25-day-old seedlings were transplanted to field with 25 x 25 cm of space between each plant. Irrigated field was flooded with 1–5 cm of water (soil water content 37%, v/v) until the end of active tillering stage (30–40 days after transplanting), then water was drained (soil water content 27%, v/v) for 10–15 days at late tillering stage, and flooded again with 3–10 cm of water until the milky stage. Soil in the non-irrigated field was kept moist (soil water content 20~25 %, v/v) by intermittent irrigation during the entire planting period. Soil water content was measured using a Theta probe and meter (models ML2x and HH1, Delta-T devices, Cambridge, UK) (Ji K, *et al.* (2012) *J Plant Physiol* 169(4):336-344). Seeds were harvested, dried and yield determined.

[0074] 2. Results

[0075] 2.1 ABRC321 has low-background but high ABA-inducibility in leaves and roots in transgenic rice

[0076] In this study, we used a tissue specific promoter, ABRC321, to express recombinant HVA1 (rHVA1) in plants. Fig. 1(A) shows the sequence of ABRC321. *ABRC321* was generated by fusion of coupling element CE3 and ABRE2 (A2) from the *HVA1* ABRC3 and coupling element CE1 from the *HVA22* ABRC1 (Shen Q, Zhang P, & Ho TH (1996) *Plant Cell* 8(7):1107-1119). The reporter gene *GUS* was fused downstream of 1-3 tandem repeats of *ABRC321* and the *Amy64* minimal promoter. See Fig.1 (B). Without ABA treatment, *GUS* activity was relatively low and only slightly increased with the copy number of *ABRC321*; in contrast, with ABA, *GUS* activity controlled by *1xABRC321*, *2xABRC321* and *3xABRC321* was induced by 19-, 20-, and 33-fold, respectively (Fig. 2 (A)).

[0077] All *ABRC321*:*GUS* constructs were also used for rice stable transformation. As shown in Fig. 2(B), four representative transgenic lines for each construct exhibited positional effects of transgene insertion site on absolute promoter activities. Among them, the *GUS* activity in leaves and roots controlled by *3xABRC321* was most highly induced by ABA among all constructs, and its absolute level and fold

induction were generally higher in leaves than in roots in response to ABA. These studies show that the promoter activity of *3xABRC321* in vegetative tissues have relatively low background but is highly inducible by ABA.

[0078] According to the staining results, in leaves, *GUS* controlled by *3xABRC321* was weakly expressed in phloems without ABA. (Fig. 3, B1 and B2), but strongly expressed in phloems, particularly in the companion cells and mesophyll cells after ABA treatment (Fig. 3, B3 and B4). In roots, *GUS* was barely detectable without ABA, but was detected at high levels mainly in lateral root primordia (LRP), pericycle and endodermis of developing roots (Fig. 3, C1-C4), and in stele, procambium and quiescent center of root tip after ABA treatment (Fig. 3, C1 and C5). *GUS* was also detected in the cortex and exodermis nearby the developing LRP derived from the crown and primary lateral roots treated with ABA (Fig. 3, C1-C4, C6 and C7).

[0079] 2.2 HVA1 expression regulated by *3xABRC321* has similar tissue-specificity and at higher-level than the endogenous HVA1 in transgenic rice in response to ABA.

[0080] The highly ABA-inducible *3xABRC321* was used to control the expression of HVA1 in transgenic rice. The expression pattern of recombinant HVA1 (rHVA1) and the rice HVA1 homolog (*Lea3*) in wild type (WT) and transgenic rice roots was examined by immunocytochemistry assays using the anti-barley HVA1 antibodies, likely due to high homology of amino acid sequences between the rice *Lea3* and barley HVA1s (see Fig. 4). The accumulation of *Lea3* in WT and rHVA1 in transgenic rice was barely detectable in roots without ABA treatment, but was significantly increased in pericycle and endodermis and slightly increased in cortex and exodermis of roots treated with ABA (Fig. 5A). The *Lea3* and rHVA1 were also detected in lateral root primordia (LRP) (Fig. 5B) and in stele, procambium and quiescent center in root tips (Fig. 5C), and levels were all increased by ABA. The accumulation of rHVA1 was significantly higher than the *Lea3* prior to and after ABA treatment.

[0081] 2.3 *3xABRC321:HVA1* induces root growth in different conditions.

[0082] The rice line, Tainung 67, seedlings were grown in agar medium containing various concentrations of ABA. Rice contains seminal, crown, adventitious and lateral roots (Fig. 6(A)) that were longer and lateral root density was greater in three transgenic lines even without ABA treatment (Fig. 6(B), panel 1), and were all significantly increased at ABA concentrations of 0.1 μ M (see the quantitative results, Fig. 7). However, lateral roots became shorter and thicker and root density

decreased with ABA concentrations higher than 0.5 μ M (Fig.6 (B), panels 3-5). Importantly, all root systems were longer and lateral root density was greater in transgenic lines than wild type (WT) at all ABA concentrations. Secondary lateral roots were observed only in transgenic lines with ABA concentration of 2 μ M (Fig. 6(B), panel 5). Similar results were also observed in roots of transgenic rice cultivar Kitaake (data not shown).

[0083] Root growth of cultivars Tainung 67 and Kitaake transgenic lines were also examined under nutrient deficient conditions. Fig. 8 (A) shows the root morphology. Total root number (Fig. 8 (C)) and root length, lateral root number, and lateral root density (Fig. 8(B) and Fig. 9 (A)-(D)) of different root systems were not significantly different when seedlings were grown with full nutrients, but were generally greater under nutrient starvation or full nutrients without nitrogen or phosphate.

[0084] 2.4 3xABRC321:HVA1 promotes root elongation and lateral root initiation in transgenic rice involves an auxin-dependent pathway

[0085] To understand whether HVA1 promotes lateral root initiation and/or elongation, root systems of seedlings treated with ABA for 11 days were examined. As shown in Fig. 10 (A)-(B), the number of lateral root primordium (LRP) was reduced but lateral roots increased in zone 1 (upper part) and zone 2 (middle part), that both cover the maturation zones, in transgenic lines as compared with WT, indicating that the majority of lateral roots initiated in transgenic lines were elongated in transgenic lines. The number of LRP and lateral root elongated were both greater in zone 3 (bottom part), which covers elongation and division zones, in transgenic lines than in WT. These results indicate that HVA1 promotes lateral root initiation and the primary root.

[0086] To determine whether the promotion of root growth by HVA1 in transgenic rice involves the auxin (IAA)-dependent pathway, the expression of various genes involved in auxin transport and signaling were determined. PIN proteins play a key role in auxin efflux from cells and have been shown to be involved in lateral root formation in cereals (Orman-Ligeza B, *et al.* (2013) Post-embryonic root organogenesis in cereals: branching out from model plants. *Trends Plant Sci* 18(8):459-467). The rice PIN family is encoded by 12 genes, with expression of each being spatially and temporally regulated (Wang JR, *et al.* (2009) *Mol Plant* 2(4):823-831). Real-time quantitative RT-PCR analysis revealed that expression of

several PIN genes was up-regulated by ABA as well as by overexpression of HVA1 in transgenic rice (Fig. 11).

[0087] To further demonstrate that auxin is involved for the ABA-induced root growth, 3-day-old rice seedlings with seminal roots only were transferred to medium containing ABA, with or without the polar auxin transport inhibitor N-(1-naphthyl) phtalamic acid (NPA) (Reed RC et al., (1998) *Plant Physiol* 118(4):1369-1378) for 11 days. In medium without any treatment, newly grown crown and lateral roots were more abundant in Tainung 67 transgenic lines than in WT (Figure 12(A)). In the presence of ABA, newly grown seminal and crown roots were significantly longer and lateral roots were more abundant in transgenic lines than in WT (Fig. 12 (B)). In the presence of NPA, growth of all types of roots was inhibited (Fig. 12(C)). In the presence of both ABA and NPA, growth of crown roots and lateral roots were also inhibited except seminal roots continued to grow (Fig. 12(D)).

[0088] **2.5 3xABRC321:HVA1 enhances water use efficiency (WUE) and grain yield in transgenic rice grown in non-irrigated field**

[0089] The WUE of transgenic Tainung 67 and Kitaake expressing rHVA1 was determined. Although transgenic lines consumed higher amounts of water (Fig. 13(A), panel 1), they produced even more biomass compared to the WT (Fig. 13(A), panel 2), thus had higher WUE than the WT (Fig. 13(A), panel 3). The performance of transgenic Kitaake grown in the field was evaluated in the fall of 2011. Rice plants were grown in two separated irrigated and non-irrigated fields, but both fields were subject to natural rainfall occasionally. Grain yield in transgenic lines was lower than WT in irrigated field, but was generally higher than WT in non-irrigated field, and surprisingly, the grain yield in transgenic lines was generally higher in non-irrigated field than in irrigated field (Fig. 13(B)). The result indicates that transgenic plants adapted the non-irrigated but naturally rain-fed field better than WT.

[0090]

[0091] **III. Conclusions**

[0092] In this study, we found that in transgenic rice with HVA1 (the LEA3 protein in barley), the accumulation of the LEA3 protein is highly inducible by ABA, accumulated in root apical meristem and lateral root primordia (LRP), and the expansion of primary and branch root systems is significantly enhanced by ABA (especially at low concentration) and also by nutrient deficiency. We also found that the root elongation and lateral root initiation in the LEA3 protein transgenic plants

involves an auxin-dependent pathway. Furthermore, the water use efficiency, biomass production, and grain yield in non-irrigated field is also increased. Our study demonstrates a successful application of transformation of plants with LEA3 gene, especially expressed by a highly active and root-specific promoter, *3xABRC321*,
5 in promoting root growth of plants, leading to advantages of at least better water use efficiency, biomass production, and grain yield.

CLAIMS

What is claimed is:

1. A method for enhancing root growth of a plant, comprising
 - (i) introducing a polynucleotide encoding group 3 late embryogenesis abundant (LEA3) protein into a plant cell to obtain a transformed plant cell, wherein the polynucleotide is operably linked to an expression control sequence,
 - (ii) producing a transformed plant from said transformed plant; and
 - (iii) selecting a transformed plant exhibiting improved root growth compared to a control plant which is not introduced with the polynucleotide.
2. The method of claim 1, wherein the expression control sequence comprising a promoter sequence.
3. The method of claim 2, wherein the promoter sequence is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3.
4. The method of claim 1, wherein the LEA3 protein is originated from monocotyledon or dicotyledon.
5. The method of claim 4, wherein the monocotyledon is selected from the group consisting of rice, barley, wheat, rye, oat, corn, bamboo, sugar cane, onion, leek and ginger.
6. The method of claim 4, wherein the monocotyledon is selected from the group consisting of rice and barley.
7. The method of claim 1, the LEA3 protein has the amino acid sequence of SEQ ID NO: 4 or 5.
8. The method of claim 4, wherein the dicotyledon is selected from the group consisting of Arabidopsis, eggplant, soybean, mung bean, kidney bean, pea, tobacco, lettuce, spinach, sweet potato, carrot, melon, cucumber and pumpkin.
9. The method of claim 1, wherein the transformed plant exhibits improved root growth compared to the control plant, in the presence of abscisic acid (ABA).
10. The method of claim 9, wherein the ABA is at a concentration from 0.1 to 5 μ M.
11. The method of claim 9, wherein the ABA is at a concentration from 0.1 to 2 μ M.
12. The method of claim 9, wherein the ABA is at a concentration of from 0.1 to

0.5 μ M.

13. The method of claim 1, wherein the transformed plant exhibits improved root growth compared to the control plant, under nutrient deficient conditions.

14. The method of claim 1, wherein the transgenic plant is monocotyledon or
5 dicotyledon.

15. The method of claim 14, wherein the monocotyledon is selected from the group consisting of rice, barley, wheat, rye, oat, corn, bamboo, sugar cane, onion, leek and ginger.

16. The method of claim 14, wherein the monocotyledon is rice.

10 17. The method of claim 14, wherein the dicotyledon is selected from the group consisting of Arabidopsis, eggplant, soybean, mung bean, kidney bean, pea, tobacco, lettuce, spinach, sweet potato, carrot, melon, cucumber and pumpkin.

15

1/20


(A)

ABRC321

ggtaccGCAACGCGTGTCTCCCTACGTGGCGGCtcgagATTGCCACCGGtctaga

CE3
A2
CE1

(B)



(C)




Fig. 1

2/20





Fig. 2

3/20

(A)



(B)




Fig. 3

4/20

Section 1									
	(1)	1	10	20	30	40	50	60	65
LEA3	(1)	MASHQDQASVRAGETRAHYEEKLGGVMDAEDKA---SEAFDASERAGSHAGGGGTFEATKD							
HVA1	(1)	MASHQDQASVRAGETRAHYEEKLGGVMDAEDKA---SEAFDASERAGSHAGGGGTFEATKD							
Consensus	(1)	MAS Q Q SY RGETKA TEEK GOMMSASF KA H KA E A N DT EA K							
Section 2									
	(66)	66	80	90	100	110	120	130	
LEA3	(62)	TAGAARDKASITAGARDCHTSSTSCARDKAAESKDDQIGGEGEKTEGAKINAAET-----							
HVA1	(66)	TAGAARDKASITAGARDCHTSSTSCARDKAAESKDDQIGGEGEKTEGAKINAAET-----							
Consensus	(66)	KA AEDK A TAGAARDCHT TAGAARDFAA KDQIG LGKTE ARDAET							
Section 3									
	(131)	131	140	150	160	170	180	195	
LEA3	(118)	--AGARDKQKTFSTAGYTFDEAAGKDKTSSVLQDASEQVNSTVVSARDAVNTEGHTEDAGTDS							
HVA1	(131)	--AGARDKQKTFSTAGYTFDEAAGKDKTSSVLQDASEQVNSTVVSARDAVNTEGHTEDAGTDS							
Consensus	(131)	AARDK STAGYTFDEAAGKDKTSSVLQQA E V VVSARDAV TEGH D T							
Section 4									
	(196)	196	215						
LEA3	(181)	GANKEDSATAAATETIAREN							
HVA1	(194)	TKEDATGATVDETTITRHH							
Consensus	(196)	T AT T TT R H							

Fig. 4

5/20




Fig. 5

6/20

(A)




Fig. 6 (Cont')

7/20

(B)





Fig. 6

8/20

(A)



(B)




Fig. 7

9/20

(A)




Fig. 8 (Cont')

10/20

(B)




Fig. 8 (Cont')

11/20

(C)




Fig. 8

12/20

(A)




Fig. 9 (Cont')

13/20

(B)




Fig. 9 (Cont')

14/20

(C)




Fig. 9 (Cont')

15/20

(D)




Fig. 9

16/20




Fig. 10

17/20




Fig. 11

18/20




Fig. 12

19/20




Fig. 13 (Cont')

20/20


B

Fig. 13

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2013/071340

A. CLASSIFICATION OF SUBJECT MATTER IPC(8) - C12N 15/29 (2014.01) USPC - 800/290 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC(8) - A01H 5/00, 5/06, 5/10; C07K 14/415; C12N 5/04, 15/29, 15/82 (2014.01) USPC - 435/410, 413, 418, 468; 536/23.6, 24.1; 800/290, 320.2 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched CPC - C07K 14/415; C12N 15/8227; 15/8261, 15/8293 (2013.01) Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Patbase, Google Patents, Google, Pubmed		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/031581 A2 (SANZ MOLINERO) 22 March 2007 (22.03.2007) entire document	1, 2, 4-6, 8, 9, 14-17
Y		3, 7, 10-13
Y	US 2009/0133154 A1 (YU et al) 21 May 2009 (21.05.2009) entire document	3, 7, 13
Y	MULKEY et al. 'The kinetics of abscisic acid action on root growth and gravitropism.' Planta. 01 March 1983 (01.03.2983), Vol. 157 (2). Pgs. 150-157. entire document	10-12
P,Y	TSENG et al. 'Abscisic Acid- and Stress-Induced Highly Proline-Rich Glycoproteins Regulate Root Growth in Rice.' Plant Physiology. 25 July 2013 (25.07.2013), Vol. 163. Pgs. 118-134. entire document	1-17
A	WO 2012/021494 A1 (STICKLEN et al) 16 February 2012 (16.02.2012) entire document	1-17
☐ Further documents are listed in the continuation of Box C. ☐		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 28 January 2014		Date of mailing of the international search report 20 FEB 2014
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-3201		Authorized officer: Blaine R. Copenheaver PCT Helpdesk: 571-272-4300 PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2013/071340

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing filed or furnished:

a. (means)

☐

on paper

☒

in electronic form

b. (time)

☒

in the international application as filed

☐

together with the international application in electronic form

☐

subsequently to this Authority for the purposes of search

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

Specifically SEQ ID NO's: 1, 2, 3, 4, and 5 were searched.