



(86) Date de dépôt PCT/PCT Filing Date: 2007/03/15
(87) Date publication PCT/PCT Publication Date: 2007/10/25
(45) Date de délivrance/Issue Date: 2011/08/23
(85) Entrée phase nationale/National Entry: 2008/06/16
(86) N° demande PCT/PCT Application No.: JP 2007/055931
(87) N° publication PCT/PCT Publication No.: 2007/119496
(30) Priorité/Priority: 2006/03/17 (JP2006-075307)

(51) Cl.Int./Int.Cl. *C07D 417/04* (2006.01),
A61K 31/4439 (2006.01), *A61P 11/00* (2006.01)

(72) Inventeurs/Inventors:
AOKI, SATOSHI, JP;
NAKAYA, KENJI, JP;
SOTA, MASAHIRO, JP;
ISHIGAMI, MASASHI, JP

(73) Propriétaire/Owner:
OTSUKA PHARMACEUTICAL CO., LTD., JP

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : POLYMORPHES DE TETOMILAST
(54) Title: TETOMILAST POLYMORPHS

(57) **Abrégé/Abstract:**

The present invention provides a tetomilast crystal that is industrially easily produced in a large volume, (1) a tetomilast hydrate crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 2; (2) an anhydrous tetomilast type A crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 4; (3) an anhydrous tetomilast type C crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 8; (4) a tetomilast acetonitrile solvate crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 10; and (5) a mixture consisting of the above anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal. These crystals are stable towards heat and moisture, and are also excellent in terms of the disintegration property and dissolution property of tablets. Accordingly, these crystals are preferably used as pharmaceutical compositions.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2007 (25.10.2007)

PCT

(10) International Publication Number
WO 2007/119496 A1

(51) International Patent Classification:

C07D 417/04 (2006.01) **A61P 11/00** (2006.01)
A61K 31/4439 (2006.01)

(74) Agents: ASAMURA, Kiyoshi et al.; Room 331, New Ohtemachi Bldg., 2-1, Ohtemachi 2-chome, Chiyoda-ku, Tokyo, 1000004 (JP).

(21) International Application Number:

PCT/JP2007/055931

(22) International Filing Date: 15 March 2007 (15.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

2006-075307 17 March 2006 (17.03.2006) JP

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): OTSUKA PHARMACEUTICAL CO., LTD. [JP/JP]; 9, Kanda-Tsukasacho 2-chome, Chiyoda-ku, Tokyo, 1018535 (JP).

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): AOKI, Satoshi [JP/JP]; c/o Otsuka Pharmaceutical Co., Ltd., 224-18, Ebisuno, Hiraishi, Kawauchi-cho, Tokushima-shi, Tokushima, 7710182 (JP). NAKAYA, Kenji [JP/JP]; c/o Otsuka Pharmaceutical Co., Ltd., 224-18, Ebisuno, Hiraishi, Kawauchi-cho, Tokushima-shi, Tokushima, 7710182 (JP). SOTA, Masahiro [JP/JP]; c/o Otsuka Pharmaceutical Co., Ltd., 224-18, Ebisuno, Hiraishi, Kawauchi-cho, Tokushima-shi, Tokushima, 7710182 (JP). ISHIGAMI, Masashi [JP/JP]; c/o Otsuka Pharmaceutical Co., Ltd., 5006-5, Aza Higashiyama, Oaza Omagari, Yoshinogari-cho, Kanzaki-gun, Saga, 8420197 (JP).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: TETOMILAST POLYMORPHS

(57) Abstract: The present invention provides a tetomilast crystal that is industrially easily produced in a large volume. (1) a tetomilast hydrate crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 2; (2) an anhydrous tetomilast type A crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 4; (3) an anhydrous tetomilast type C crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 8; (4) a tetomilast acetonitrile solvate crystal having a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 10; and (5) a mixture consisting of the above anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal. These crystals are stable towards heat and moisture, and are also excellent in terms of the disintegration property and dissolution property of tablets. Accordingly, these crystals are preferably used as pharmaceutical compositions.

WO 2007/119496 A1

DESCRIPTION

TETOMILAST POLYMORPHS

TECHNICAL FIELD

[0001]

The present invention relates to a novel tetomilast crystal.

5 BACKGROUND ART

[0002]

2-(3,4-diethoxyphenyl)-4-(2-carboxy-6-pyridyl)thiazole (or 6-[2-(3,4-diethoxyphenyl)thiazol-4-yl]pyridin-2-carboxylic acid) is a known compound.

10 This compound is named as tetomilast. This tetomilast has active oxygen inhibitory action, cytokine generation inhibitory action, adhesion inhibitory action, etc., and it is useful for the treatment of ulcerative colitis, Crohn's disease, asthma, and the
15 like (JP-A-5-51318 (paragraph [0015], Example 371) and JP-A-10-152437 (paragraphs [0024] and [0029])). In addition, tetomilast is also useful as a therapeutic agent for chronic obstructive pulmonary disease (JP-A-2003-104890). Moreover, such a tetomilast crystal is
20 stable towards heat and moisture, and is excellent in terms of the disintegration property and dissolution property of tablets.

[0003]

According to Example 371 of JP-A-5-51318, an anhydrous tetomilast crystal (hereinafter referred to as an "anhydrous tetomilast type B crystal") is produced by allowing 3,4-diethoxythiobenzamide to react with 2-(2-chloroacetyl)-6-pyridine carboxylic acid and then recrystallizing the obtained tetomilast crude product from ethanol, for example.

[0004]

Furthermore, according to Journal of Medicinal Chemistry, 1995, 38, pp. 353-358, such an anhydrous tetomilast type B crystal is produced by allowing 3,4-diethoxythiobenzamide to react with 2-(2-bromoacetyl)-6-pyridine carboxylic acid, then hydrolyzing the obtained methyl 6-[2-(3,4-diethoxyphenyl)thiazol-4-yl]pyridin-2-carboxylic acid methyl ester, and then recrystallizing the obtained tetomilast crude product from ethyl acetate.

[0005]

However, since an anhydrous tetomilast type B crystal has the property of causing clogging, if such an anhydrous tetomilast type B crystal is produced by the conventional recrystallization method, an operating efficiency significantly deteriorates during filtration. Thus, it is difficult to produce anhydrous tetomilast type B crystals industrially in a large volume. Accordingly, it is strongly desired that a novel tetomilast crystal, which is advantageous in terms of industrial mass production, be developed.

DISCLOSURE OF THE INVENTION

[0006]

It is a main object of the present invention to provide a novel tetomilast crystal that is
5 industrially produced in a large volume.

[0007]

The present inventors have conducted intensive studies directed towards achieving the aforementioned object. As a result, the inventors have
10 found that a novel tetomilast crystal, which can be produced industrially in a large volume, can be obtained by stirring anhydrous tetomilast type B crystals in an aqueous solvent, and that the above tetomilast crystal is further recrystallized using a
15 suitable solvent, or is further heated while it is suspended in a specific solvent, so as to obtain a tetomilast crystal having various physical properties. The present invention has been completed based on these findings.

20 [0008]

That is to say, the present invention relates to the following tetomilast crystal and pharmaceutical composition:

1. A tetomilast hydrate crystal having a powder
25 X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 2.
2. An anhydrous tetomilast type A crystal having

a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 4.

3. An anhydrous tetomilast type C crystal having
5 a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 8.

4. A tetomilast acetonitrile solvate crystal having a powder X-ray diffraction spectrum that is
10 substantially the same as the powder X-ray diffraction spectrum shown in Fig. 10.

5. A mixture of the anhydrous tetomilast type A crystal described in item 2 and an anhydrous tetomilast type B crystal.

15 6. A pharmaceutical composition comprising at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast type A crystal, an anhydrous tetomilast type C crystal, and a tetomilast
20 acetonitrile solvate crystal.

7. The pharmaceutical composition according to item 6, which is an agent for preventing and/or treating gastrointestinal ulcer, an agent for preventing and/or treating cardiac ischemic disease, an
25 agent for preventing and/or treating cerebrovascular disease, a liver and kidney function improver used for disorders caused by transplantation, microcirculation failure, etc., or an agent for preventing and/or

treating Behcet's disease, cutaneous vasculitis, ulcerative colitis, malignant rheumatism, arthritis, arteriosclerosis or diabetes.

8. The pharmaceutical composition according to
5 item 6, which is an agent for preventing and/or treating chronic rheumatoid arthritis, endotoxin shock, ARDS, thermal burn, asthma, chronic heart failure, myocardial infarction, viral myocarditis, or an agent for preventing and/or treating ischemic reperfusion
10 abnormality, transition from SIRS (systemic inflammatory response syndrome) to organ failure, multiple organ failure, inflammatory bowel disease, autoimmune disease, metastasis, immunological rejection occurring during transplantation, monoclonal B cell
15 abnormality, polyclonal B cell abnormality, atrial myxoma, Castleman's syndrome, primary glomerulonephritis, mesangial proliferative nephritis, cancer cachexia, Lennert's lymphoma, psoriasis, atopic dermatitis, Kaposi's sarcoma developed due to AIDS,
20 postmenopausal osteoporosis, septicemia, inflammatory disease or chronic obstructive pulmonary disease.

9. The pharmaceutical composition according to item 8, wherein the inflammatory intestinal disease is ulcerative colitis or Crohn's disease.

25 10. The pharmaceutical composition according to item 8, which is an agent for preventing and/or treating chronic obstructive pulmonary disease.

11. A process for the preparation of a tetomilast

hydrate, which is characterized in that it comprises stirring an anhydrous tetomilast type B crystal in an aqueous solvent.

12. A process for the preparation of an anhydrous
5 tetomilast type A crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving an anhydrous tetomilast type B crystal in a solvent.

13. The process according to item 12, wherein the
10 solvent is ethanol, acetone, or acetone-water (wherein the acetone content is 40% or more).

14. A process for the preparation of an anhydrous
tetomilast type A crystal, which is characterized in that it comprises recrystallization from a solution
15 formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast type C crystal, and a tetomilast acetonitrile solvate crystal.

20 15. The process according to item 14, wherein the solvent is a mixed solvent consisting of water, and at least one organic solvent selected from the group consisting of methanol, ethanol, acetone, and tetrahydrofuran.

25 16. A process for the preparation of an anhydrous tetomilast type C crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving an anhydrous tetomilast type B

crystal in a solvent.

17. The process according to item 16, wherein the solvent is methanol or ethanol.

18. A process for the preparation of an anhydrous
5 tetomilast type C crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous
10 tetomilast type A crystal, and a tetomilast acetonitrile solvate crystal.

19. The production method according to item 18, wherein the solvent is methanol or ethanol.

20. A process for the preparation of a tetomilast
15 acetonitrile solvate crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving an anhydrous tetomilast type B crystal in acetonitrile.

21. A process for the preparation of an anhydrous
20 tetomilast acetonitrile solvate crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an
25 anhydrous tetomilast type A crystal, and an anhydrous tetomilast type C crystal.

22. A process for the preparation of a mixture consisting of an anhydrous tetomilast type A crystal

and an anhydrous tetomilast type B crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving the anhydrous tetomilast type B crystal in a solvent.

5 23. The process according to item 22, wherein the solvent is acetone-water (wherein the acetone content is 40% to 95% by volume).

24. A process for the preparation of a mixture consisting of an anhydrous tetomilast type A crystal
10 and an anhydrous tetomilast type B crystal, which is characterized in that it comprises recrystallization from a solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an
15 anhydrous tetomilast type A crystal, a tetomilast acetonitrile solvate crystal, and an anhydrous tetomilast type C crystal.

25. The process according to item 24, wherein the solvent is acetone-water (wherein the acetone content
20 is 40% to 95% by volume).

[0009]

In the present invention, the term "novel tetomilast crystals" is used to mean a tetomilast hydrate crystal, an anhydrous tetomilast type A
25 crystal, an anhydrous tetomilast type C crystal, a tetomilast acetonitrile solvate crystal, and a mixture of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal.

[0010]

In addition, in the present invention, the term "tetomilast crystal" is used to simply mean a generic name for such novel tetomilast crystals and an
5 anhydrous tetomilast type B crystal that is known as a tetomilast crystal.

BRIEF DESCRIPTION OF THE DRAWINGS

[0223]

Fig. 1 is a view showing the
10 thermogravimetry/differential thermal analysis of the tetomilast hydrate crystal obtained in Example 5 (1);

Fig. 2 is a view showing the powder X-ray diffraction of the tetomilast hydrate crystal obtained in Example 5 (1);

15 Fig. 3 is a view showing the thermogravimetry/differential thermal analysis of the anhydrous tetomilast type A crystal obtained in Example 1 (1);

Fig. 4 is a view showing the powder X-ray
20 diffraction of the anhydrous tetomilast type A crystal obtained in Example 1 (1);

Fig. 5 is a view showing the thermogravimetry/differential thermal analysis of the anhydrous tetomilast type B crystal obtained in
25 Reference example 1 (2);

Fig. 6 is a view showing the powder X-ray diffraction of the anhydrous tetomilast type B crystal

obtained in Reference example 1 (2);

Fig. 7 is a view showing the thermogravimetry/differential thermal analysis of the anhydrous tetomilast type C crystal obtained in Example 2;

Fig. 8 is a view showing the powder X-ray diffraction of the anhydrous tetomilast type C crystal obtained in Example 2;

Fig. 9 is a view showing the thermogravimetry/differential thermal analysis of the tetomilast acetonitrile solvate crystal obtained in Example 3;

Fig. 10 is a view showing the powder X-ray diffraction of the tetomilast acetonitrile solvate crystal obtained in Example 3;

Fig. 11 is a view showing the thermogravimetry/differential thermal analysis of the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal obtained in Example 4(1);

Fig. 12 is a view showing the powder X-ray diffraction of the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal obtained in Example 4(1);

Fig. 13 is a view showing the thermogravimetry/differential thermal analysis of the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal

obtained in Example 4(2); and

Fig. 14 is a view showing the powder X-ray diffraction of the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast
5 type B crystal obtained in Example 4 (2).

BEST MODE FOR CARRYING OUT THE INVENTION

[0011]

Tetomilast hydrate crystal

The tetomilast hydrate crystal of the present
10 invention includes 0.5 to 3 hydrate crystals. Among the tetomilast hydrate crystals of the present invention, a monohydrate crystal has the physicochemical properties described in (1) to (3) below:

15 [0012]

(1) The monohydrate crystal has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal analysis (temperature-rising rate/min) endothermic curve shown
20 in Fig. 1. Specifically, such a monohydrate crystal is characterized in that it has an endothermic peak around 189°C and a wide peak around 102°C.

[0013]

(2) The monohydrate crystal has a powder X-
25 ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 2. Specifically, it has characteristic peaks at $2\theta =$

10.6°, 12.9°, 21.1°, 22.3°, and 25.0°.

[0014]

(3) The monohydrate crystal has significant infrared absorption bands at 3516, 3433, 1742, 1709,
5 1587, 1472, 1267, 1143, 1040, 758, and 716 cm⁻¹ in the IR (KBr) spectrum.

[0015]

Method for producing tetomilast hydrate crystal

The tetomilast hydrate crystal of the present
10 invention can be obtained by stirring in an aqueous solvent the known anhydrous tetomilast type B crystal or an anhydrous tetomilast type B crystal obtained by the method as described below.

[0016]

15 Such an aqueous solvent is a mixed solvent obtained by mixing an organic solvent such as methanol, ethanol, isopropanol, acetone, or ethylmethylketone, with water. Examples of such a mixed solvent may include methanol-water (a methanol content of 10% to
20 80% by volume), ethanol-water (an ethanol content of 10% to 70% by volume), isopropanol-water (an isopropanol content of 10% to 60% by volume), acetone-water (an acetone content of 10% to 80% by volume), and ethylmethylketone-water (an ethylmethylketone content
25 of 10% to 80% by volume). Of these, acetone-water (an acetone content of 10% to 60% by volume) and ethylmethylketone-water (an ethylmethylketone content of 10% to 60% by volume) are particularly preferable.

Acetone-water (an acetone content of 35% to 55% by volume) is more preferable.

[0017]

The amount of an aqueous solvent used is not limited. It is used in an amount of at least 10 ml, and preferably between 10 and 50 ml, with respect to 1 g of an anhydrous tetomilast type B crystal.

[0018]

The stirring temperature is not particularly limited. It is preferably between approximately 10°C and 35°C, and more preferably between approximately 20°C and 30°C. The stirring time is preferably between approximately 5 minutes and 3 hours, and more preferably between approximately 30 and 90 minutes.

[0019]

In addition, when a tetomilast hydrate crystal is produced from an anhydrous tetomilast type B crystal, it is preferable to allow the tetomilast hydrate crystal produced by the aforementioned method separately to exist as a seed crystal in a suspension.

[0020]

The time of adding a seed crystal is not particularly limited. It is preferable to add such a seed crystal before or during stirring.

[0021]

The obtained tetomilast hydrate crystal can be separated by isolation operations such as filtration, concentration, or extraction. Moreover,

after separation, the separated crystal may be subjected to a drying treatment by a known method. Furthermore, the crystal can be purified by a known purification operation.

5 [0022]

The thus obtained tetomilast hydrate crystal has a purity of 95% or more, and it can be crushed using an ordinary crusher (for example, an atomizer). Thus, a tetomilast crushed product having a mean
10 particle size between 10 and 50 μm and a 90% cumulative particle size of 80 μm or smaller, which is suitable for formulation, can be obtained.

[0023]

Anhydrous tetomilast type B crystal

15 An anhydrous tetomilast type B crystal has the physicochemical properties described in (4) to (6) below:

[0024]

(4) The anhydrous tetomilast type B crystal
20 has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal analysis (temperature-rising rate/min) endothermic curve shown in Fig. 5. Specifically, such an anhydrous tetomilast type B crystal is characterized in that it has
25 endothermic peaks around 177°C and around 188°C.

[0025]

(5) The anhydrous tetomilast type B crystal has a powder X-ray diffraction spectrum that is

substantially the same as the powder X-ray diffraction spectrum shown in Fig. 6. Specifically, it has characteristic peaks at $2\theta = 4.1^\circ, 8.1^\circ, 11.9^\circ, 16.1^\circ,$ and 24.2° .

5 [0026]

(6) The anhydrous tetomilast type B crystal has significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 762, and 706 cm^{-1} in the IR (KBr) spectrum.

10 [0027]

The anhydrous tetomilast type B crystal can be produced by the method described in JP-A-5-51318 or Journal of Medicinal Chemistry, 1995, 38, pp. 353-358.

[0028]

15 Moreover, the anhydrous tetomilast type B crystal may be obtained by completely dissolving a novel tetomilast crystal in a solvent while stirring under heating to reflux, and then allowing the obtained solution to stand to cool. Herein, such a novel
20 tetomilast crystal can be used singly or as a mixture consisting of two or more types.

[0029]

Examples of a solvent may include isopropanol, ethyl acetate, and a mixed solvent
25 thereof.

[0030]

The amount of a solvent used is not particularly limited, as long as it is able to

completely dissolve a novel tetomilast crystal while stirring under heating to reflux. In the case of isopropanol, the amount used is preferably between 70 and 600 ml with respect to 1 g of a novel tetomilast
5 crystal. In the case of ethyl acetate, it is preferably between 30 and 300 ml with respect to 1 g of a novel tetomilast crystal. When a mixed solvent consisting of isopropanol and ethyl acetate is used, isopropanol is mixed with ethyl acetate at any given
10 ratio, and the amount of such a mixed solvent used may be adjusted so as to completely dissolve a novel tetomilast crystal while stirring under heating to reflux.

[0031]

15 The obtained solution is cooled to approximately 30°C over about 5 minutes to 1 hour, or is naturally cooled, so as to obtain an anhydrous tetomilast type B crystal. Further, after the suspension has been stood to cool, it may be cooled at
20 a temperature of 10°C or lower, and preferably at a temperature between approximately 0°C and 10°C. By this operation, an anhydrous tetomilast type B crystal is obtained at a higher yield.

[0032]

25 Furthermore, such an anhydrous tetomilast type B crystal is also obtained by allowing a basic compound to act on a novel tetomilast crystal so as to form salts, dissolving the salts in a suitable aqueous

solvent, and then adding suitable acid to the obtained solution.

[0033]

Examples of a basic compound may include
5 carbonate, alkali metal hydroxide, and alkali earth
metal hydroxide. Of these, alkali metal hydroxide is
particularly preferable. Examples of carbonate may
include sodium carbonate, potassium carbonate, sodium
bicarbonate, and potassium bicarbonate. Examples of
10 alkali metal hydroxide may include sodium hydroxide and
potassium hydroxide. Examples of alkali earth metal
hydroxide may include calcium hydroxide, barium
hydroxide, and magnesium hydroxide. These compounds
are used singly or as a mixture of two or more types.
15 Of these, potassium hydroxide and sodium hydroxide are
particularly preferable.

[0034]

The additive amount of a basic compound is
not particularly limited. It is generally 1 equivalent
20 or greater, and preferably between approximately 1 and
1.5 equivalent, with respect to 1 equivalent of a novel
tetomilast crystal to be treated.

[0035]

Examples of the above described acid may
25 include inorganic acids such as hydrochloric acid,
sulfuric acid, and hydrobromic acid.

[0036]

The additive amount of acid is generally 1

equivalent (neutralization amount) or greater, and preferably between approximately 1 and 1.5 equivalent, with respect to the above described basic compound used.

5 [0037]

Examples of an aqueous solvent used herein may include those as described above in the method for producing a tetomilast hydrate crystal. Among others, acetone-water (an acetone content of 30% to 70% by
10 volume) is particularly preferable.

[0038]

The amount of an aqueous solvent used is not limited. It is 5 ml or greater, preferably between 5 and 300 ml, and more preferably between 30 and 70 ml,
15 with respect to 1 g of a novel tetomilast crystal.

[0039]

The temperature of a solution is preferably between 10°C and 30°C when acid is added.

[0040]

20 A solution is converted to a suspension by addition of acid. Thus, such a suspension is cooled generally at 10°C or lower, and preferably at a temperature between 0°C and 10°C, so as to efficiently extract an anhydrous tetomilast type B crystal.

25 [0041]

In addition, these methods are applied to anhydrous tetomilast type B crystals, so as to obtain anhydrous tetomilast type B crystals with higher

purity.

[0042]

Anhydrous tetomilast type A crystal

An anhydrous tetomilast type A crystal has
5 the physicochemical properties described in (7) to (9)
below:

[0043]

(7) The anhydrous tetomilast type A crystal
has an endothermic curve that is substantially the same
10 as the thermogravimetry/differential thermal analysis
(temperature-rising rate/min) endothermic curve shown
in Fig. 3. Specifically, such an anhydrous tetomilast
type A crystal is characterized in that it has an
endothermic peak around 188°C.

15 [0044]

(8) The anhydrous tetomilast type A crystal
has a powder X-ray diffraction spectrum that is
substantially the same as the powder X-ray diffraction
spectrum shown in Fig. 4. Specifically, it has
20 characteristic peaks at $2\theta = 10.5^\circ, 13.1^\circ, 18.4^\circ, 21.9^\circ,$
and 25.8° .

[0045]

(9) The anhydrous tetomilast type A crystal
has significant infrared absorption bands at 3306,
25 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758,
and 704 cm^{-1} in the IR (KBr) spectrum.

[0046]

Method for producing anhydrous tetomilast type A

crystal

The anhydrous tetomilast type A crystal can be obtained by recrystallization from a solution that is obtained by dissolving a known anhydrous tetomilast
5 type B crystal or an anhydrous tetomilast type B crystal obtained by the aforementioned method in a suitable solvent.

[0047]

Examples of the aforementioned solvent that
10 can be used herein may include ethanol, acetone, and acetone-water (an acetone content of 40% or more by volume). Of these, acetone-water (an acetone content of 40% or more by volume) is particularly preferable.

[0048]

15 The amount of a solvent used is not particularly limited, as long as it is able to completely dissolve an anhydrous tetomilast type B crystal while stirring under heating to reflux. In the case of ethanol, the amount used is preferably between
20 70 and 400 ml with respect to 1 g of an anhydrous tetomilast type B crystal. In the case of acetone, it is preferably between 30 and 120 ml with respect to 1 g of the above anhydrous tetomilast type B crystal. In the case of acetone-water (an acetone content of 40% to
25 80% by volume), it is preferably between 30 and 500 ml with respect to 1 g of the above anhydrous tetomilast type B crystal.

[0049]

Such an anhydrous tetomilast type B crystal is preferably dissolved in a solvent while stirring under heating to reflux. At that time, the heating temperature is not particularly limited. It is
5 generally between approximately 40°C and 85°C, and preferably between approximately 55°C and 80°C.

[0050]

After such dissolution, the temperature of the obtained solution is decreased, so that the
10 anhydrous tetomilast type A crystal of the present invention can be crystallized.

[0051]

The temperature-decreasing rate is not particularly limited. When ethanol is used as a
15 solvent, for example, the temperature-decreasing rate is preferably 0.8°C/min or less. In addition, when acetone-water (an acetone content of 40% or more by volume) is used as a solvent, the temperature-decreasing rate is preferably 0.4°C/min or less. By
20 setting the temperature-decreasing rate within such a range, the anhydrous tetomilast type A crystal of the present invention can be obtained more efficiently.

[0052]

When acetone-water (an acetone content of 40%
25 or more by volume) is used as a solvent, the aforementioned solution is retained at 40°C to 50°C for 60 minutes or longer, and it is then cooled, so that an anhydrous tetomilast type A crystal can be

crystallized. The temperature-decreasing rate during the aforementioned cooling does not particularly affect crystallization of the anhydrous tetomilast type A crystal.

5 [0053]

In addition, during such a decrease in the temperature, while the temperature is retained in several temperature ranges such as 40°C to 50°C, 30°C to 40°C, 15°C to 25°C, or 0°C to 10°C, the solution is
10 stirred for approximately 30 minutes to 5 hours, so that the temperature may be decreased in a stepwise manner. In this stepwise temperature-decreasing method, at a temperature between 40°C and 50°C, an anhydrous tetomilast type A crystal, which has been
15 produced separately, may be added as a seed crystal.
[0054]

Moreover, such an anhydrous tetomilast type A crystal is produced by using a novel tetomilast crystal (excluding an anhydrous tetomilast type A crystal)
20 instead of the known anhydrous tetomilast type B crystal, and recrystallizing from a solution formed by dissolving the above novel tetomilast crystal in a suitable solvent.

Specifically, an anhydrous tetomilast type A
25 crystal is produced by recrystallization from a solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an

anhydrous tetomilast type C crystal, and a tetomilast acetonitrile solvate crystal.

Herein, a novel tetomilast crystal is used singly or as a mixture consisting of two or more types.

5 [0055]

Such an anhydrous tetomilast type A crystal can also be produced by suspending the known anhydrous tetomilast type B crystal or a novel tetomilast crystal (excluding an anhydrous tetomilast type A crystal) in
10 an aqueous solvent (a water content of 90% or less by volume), and then stirring the suspension.

[0056]

Examples of an aqueous solvent that can be used herein may include mixed solvents formed by mixing
15 organic solvents having high compatibility with water, such as methanol, ethanol, acetone, or tetrahydrofuran, with water.

Specifically, the aqueous solvent that can be used as a mixed solvent is consisting of water, and at
20 least one organic solvent selected from the group consisting of methanol, ethanol, acetone, and tetrahydrofuran. In particular, acetone-water (an acetone content of 30% to 60% by volume) is preferable as such an aqueous solvent.

25 [0057]

The temperature of a suspension is not particularly limited during stirring. It is generally between 0°C and 65°C, and preferably between 10°C and

60°C.

[0058]

The stirring time is generally between 10 minutes and 48 hours, and preferably between 10 minutes and 3 hours.

[0059]

In addition, these methods are applied to anhydrous tetomilast type A crystals, so as to obtain anhydrous tetomilast type A crystals with higher purity.

[0060]

The obtained anhydrous tetomilast type A crystal can be separated by isolation operations such as filtration, concentration, or extraction. Moreover, after separation, the separated crystal may be subjected to a drying treatment by a known method. Furthermore, the crystal can be purified by a known purification operation.

[0061]

The thus obtained anhydrous tetomilast type A crystal has a purity of 95% or more, and it can be crushed using an ordinary crusher (for example, an atomizer). Thus, a tetomilast crushed product having a mean particle size between 10 and 50 μm and a 90% cumulative particle size of 80 μm or smaller, which is suitable for formulation, can be obtained.

[0062]

Anhydrous tetomilast type C crystal

An anhydrous tetomilast type C crystal has the physicochemical properties described in (10) to (12) below:

[0063]

5 (10) The anhydrous tetomilast type C crystal has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal analysis (temperature-rising rate/min) endothermic curve shown in Fig. 7. Specifically, such an anhydrous tetomilast
10 type C crystal is characterized in that it has endothermic peaks around 184°C and around 189°C.

[0064]

(11) The anhydrous tetomilast type C crystal has a powder X-ray diffraction spectrum that is
15 substantially the same as the powder X-ray diffraction spectrum shown in Fig. 8. Specifically, it has characteristic peaks at $2\theta = 4.2^\circ, 8.2^\circ, 12.0^\circ, 16.4^\circ, 24.7^\circ,$ and 25.9° .

[0065]

20 (12) The anhydrous tetomilast type C crystal has significant infrared absorption bands at 3300, 3088, 1744, 1593, 1476, 1346, 1267, 1132, 1045, 754, and 704 cm^{-1} in the IR (KBr) spectrum.

[0066]

25 Method for producing anhydrous tetomilast type C crystal

The anhydrous tetomilast type C crystal can be produced by recrystallization from a solution that

is obtained by dissolving the known anhydrous tetomilast type B crystal or an anhydrous tetomilast type B crystal obtained by the aforementioned method in a suitable solvent.

5 [0067]

Examples of a solvent that can be used herein may include methanol and ethanol. Of these, methanol is particularly preferable.

[0068]

10 The amount of a solvent used is not particularly limited, as long as it is able to completely dissolve the aforementioned anhydrous tetomilast type B crystal while stirring under heating to reflux. The amount of a solvent used is preferably
15 between 70 and 200 ml, and more preferably between 80 and 120 ml, with respect to 1 g of the aforementioned anhydrous tetomilast type B crystal.

[0069]

After such dissolution, the temperature of
20 the obtained solution is decreased to a temperature between 10°C and 30°C, so that the anhydrous tetomilast type C crystal of the present invention can be crystallized. The temperature-decreasing rate is not particularly limited, when the solvent is methanol. It
25 may be approximately 0.4°C to 0.6°C/min. When ethanol is used as a solvent, the obtained solution may rapidly be cooled at a temperature-decreasing rate of 5°C/min or more, and preferably of 10°C/min or more. By setting

the temperature-decreasing rate within such a range, the anhydrous tetomilast type C crystal of the present invention can be obtained more efficiently.

[0070]

5 In addition, such an anhydrous tetomilast type C crystal is produced by recrystallization from a solution obtained by dissolving in a suitable solvent a novel tetomilast crystal (excluding the anhydrous tetomilast type C crystal) that is used instead of the
10 known anhydrous tetomilast type B crystal.

Specifically, the known anhydrous tetomilast type B crystal is first added to methanol, and it is then dissolved therein while stirring under heating to reflux. The obtained solution is stood to cool to
15 approximately 30°C for about 40 minutes to 1 hour. Thereafter, the suspension obtained by the aforementioned cooling is cooled at 10°C or lower, and preferably at a temperature between approximately 0°C and 10°C, for 30 minutes to 3 hours, so that the
20 anhydrous tetomilast type C crystal of the present invention can be obtained in the form of a crystal.

[0071]

Specifically, an anhydrous tetomilast type C crystal is produced by recrystallization from a
25 solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast type A crystal, and a tetomilast

acetonitrile solvate crystal. Herein, a novel tetomilast crystal is used singly or as a mixture consisting of two or more types. The solvent and recrystallization conditions applied herein are
5 equivalent to those in the case of using the aforementioned known anhydrous tetomilast type B crystal as a starting material.

[0072]

Moreover, the aforementioned method is
10 applied to anhydrous tetomilast type C crystals, so as to obtain anhydrous tetomilast type C crystals with higher purity.

[0073]

The obtained anhydrous tetomilast type C
15 crystal can be separated by isolation operations such as filtration, concentration, or extraction. Moreover, after separation, the separated crystal may be subjected to a drying treatment by a known method. Furthermore, the crystal can be purified by a known
20 purification operation.

[0074]

The thus obtained anhydrous tetomilast type C crystal has a purity of 95% or more, and it can be crushed using an ordinary crusher (for example, an
25 atomizer). Thus, a tetomilast crushed product having a mean particle size between 10 and 50 μm and a 90% cumulative particle size of 80 μm or smaller, which is suitable for formulation, can be obtained.

[0075]

Tetomilast acetonitrile solvate crystal

A tetomilast acetonitrile solvate crystal has the physicochemical properties described in (13) to
5 (15) below:

[0076]

(13) The tetomilast acetonitrile solvate crystal has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal
10 analysis (temperature-rising rate/min) endothermic curve shown in Fig. 9. Specifically, such a tetomilast acetonitrile solvate crystal is characterized in that it has endothermic peaks around 91°C, around 176°C, and around 189°C.

15 [0077]

(14) The tetomilast acetonitrile solvate crystal has a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 10. Specifically, it has
20 characteristic peaks at $2\theta = 3.6^\circ, 7.1^\circ, 10.6^\circ, 14.2^\circ,$ and 24.8° .

[0078]

(15) The tetomilast acetonitrile solvate crystal has significant infrared absorption bands at
25 3300, 3090, 2249 (nitrile group), 1744, 1593, 1476, 1346, 1269, 1132, 1045, 752, and 704 cm^{-1} in the IR (KBr) spectrum.

[0079]

Method for producing tetomilast acetonitrile solvate crystal

The tetomilast acetonitrile solvate crystal can be produced by recrystallization from a solution that is obtained by dissolving in acetonitrile the known anhydrous tetomilast type B crystal or an anhydrous tetomilast type B crystal obtained by the aforementioned method.

[0080]

The amount of acetonitrile used is not particularly limited, as long as it is able to completely dissolve the above anhydrous tetomilast type B crystal while stirring under heating to reflux. The amount of acetonitrile used is preferably between 70 and 150 ml, and more preferably between 70 and 100 ml, with respect to 1 g of the above anhydrous tetomilast type B crystal.

[0081]

The above anhydrous tetomilast type B crystal may be dissolved in acetonitrile while stirring under heating to reflux, for example. After such dissolution, the temperature of the solution in which the anhydrous tetomilast type B crystal has been dissolved is decreased, so that the tetomilast acetonitrile solvate crystal of the present invention can be crystallized. The temperature-decreasing rate is not particularly limited, and it may be between approximately 0.1°C/min and 1.5°C/min. The tetomilast

acetonitrile solvate crystal of the present invention is not particularly affected by such a temperature-decreasing rate, and it can preferably be obtained.

[0082]

5 Specifically, the known anhydrous tetomilast type B crystal is added to acetonitrile, and it is then dissolved therein while stirring under heating to reflux. The obtained solution is stood to cool to approximately 30°C for about 30 minutes to 8 hours.

10 Thereafter, the suspension obtained by the aforementioned cooling is cooled at 10°C or lower, and preferably at a temperature between approximately 0°C and 10°C, for 30 minutes to 3 hours, so that the tetomilast acetonitrile solvate crystal of the present

15 invention can be obtained in the form of a crystal.

[0083]

 In addition, such a tetomilast acetonitrile solvate crystal is produced by recrystallization from a solution obtained by dissolving in acetonitrile a novel

20 tetomilast crystal (excluding the tetomilast acetonitrile solvate crystal) that is used instead of the known anhydrous tetomilast type B crystal.

 Specifically, an tetomilast acetonitrile solvate crystal is produced by recrystallization from a

25 solution formed by dissolving in acetonitrile at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast type A, and an anhydrous

tetomilast type C crystal. Herein, a novel tetomilast crystal is used singly or as a mixture consisting of two or more types. The solvent and recrystallization conditions applied herein are equivalent to those in the case of using the aforementioned known anhydrous tetomilast type B crystal as a starting material.

[0084]

Moreover, the aforementioned method is applied to tetomilast acetonitrile solvate crystals, so as to obtain tetomilast acetonitrile solvate crystals with higher purity.

[0085]

The obtained tetomilast acetonitrile solvate crystal can be separated by isolation operations such as filtration, concentration, or extraction. Moreover, after separation, the separated crystal may be subjected to a drying treatment by a known method. Furthermore, the crystal can be purified by a known purification operation.

[0086]

The thus obtained tetomilast acetonitrile solvate crystal has a purity of 95% or more, and it can be crushed using an ordinary crusher (for example, an atomizer). Thus, a tetomilast crushed product having a mean particle size between 10 and 50 μm and a 90% cumulative particle size of 80 μm or smaller, which is suitable for formulation, can be obtained.

[0087]

Mixture consisting of anhydrous tetomilast type A crystal and anhydrous tetomilast type B crystal

As a mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal, mixtures with various types of ratios are generated depending on production conditions. As an example, a mixture having the physicochemical properties described in (16) to (18) below can be generated:

10 [0088]

(16) The level of an endothermic peak depends on the mixing ratio of type A crystals and type B crystals. Fig. 11 shows the endothermic peak of a sample having a mixing ratio of A:B = 40:60. The mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal analysis (temperature-rising rate/min) endothermic curve shown in Fig. 11. Specifically, such a mixture is characterized in that it has endothermic peaks around 175°C and around 189°C.

[0089]

(17) The powder X-ray diffraction spectrum of the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal is the sum of the powder X-ray diffraction spectrum of a pure anhydrous tetomilast type A crystal

and the powder X-ray diffraction spectrum of a pure anhydrous tetomilast type B crystal. The strength of a peak derived from each crystal form is affected by the mixing ratio between the anhydrous tetomilast type A
5 crystal and the anhydrous tetomilast type B crystal. Fig. 12 shows the powder X-ray diffraction spectrum of a sample having a mixing ratio of the anhydrous tetomilast type A crystal: the anhydrous tetomilast type B crystal = 40:60.

10 [0090]

(18) The sample having a mixing ratio of the anhydrous tetomilast type A crystal: the anhydrous tetomilast type B crystal = 40:60 has significant infrared absorption bands at 3298, 3088, 1744, 1593,
15 1474, 1348, 1269, 1132, 1045, 760, and 704 cm^{-1} in the IR (KBr) spectrum. In addition, a deviation of $\pm 5\text{cm}^{-1}$ may be generated in an absorption peak due to a difference in the mixing ratio between the anhydrous tetomilast type A crystal and the anhydrous tetomilast
20 type B crystal.

[0091]

Moreover, a sample having a mixing ratio of A:B = 10:90 has the physicochemical properties described in (19) to (21) below:

25 [0092]

(19) The sample has an endothermic curve that is substantially the same as the thermogravimetry/differential thermal analysis

(temperature-rising rate/min) endothermic curve shown in Fig. 13. Specifically, such a sample is characterized in that it has endothermic peaks around 176°C and around 189°C.

5 [0093]

(20) The sample has a powder X-ray diffraction spectrum that is substantially the same as the powder X-ray diffraction spectrum shown in Fig. 14. Specifically, it has characteristic peaks at $2\theta = 4.1^\circ$,
10 11.9° , 16.1° , 17.2° , 19.3° , 24.2° , 25.1° , 25.9° , and 27.3° .

[0094]

(21) The sample has significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348,
15 1269, 1132, 1045, 756, and 706 cm^{-1} in the IR (KBr) spectrum.

[0095]

Method for producing mixture consisting of anhydrous tetomilast type A crystal and anhydrous tetomilast type
20 B crystal

A mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal can be produced by recrystallization from a solution that is obtained by dissolving in a
25 suitable solvent the known anhydrous tetomilast type B crystal or an anhydrous tetomilast type B crystal obtained by the aforementioned method.

[0096]

The type of the aforementioned solvent is not particularly limited. Acetone-water (an acetone content of 40% to 95% by volume) is preferable.

[0097]

5 The amount of the aforementioned solvent used is not particularly limited, as long as it is able to completely dissolve the above tetomilast crystal while stirring under heating to reflux. The amount of the solvent used is preferably between 30 and 160 ml, and
10 more preferably between 30 and 50 ml, with respect to 1 g of the above anhydrous tetomilast type B crystal.

[0098]

 The above tetomilast crystal may be dissolved in the aforementioned solvent while stirring under
15 heating to reflux, for example. After such dissolution, the temperature of the solution in which the anhydrous tetomilast type B crystal has been dissolved is decreased, so that the mixture consisting of an anhydrous tetomilast type A crystal and an
20 anhydrous tetomilast type B crystal of the present invention can be obtained. The temperature-decreasing rate may be between approximately 0.4°C/min and 1.9°C/min. In particular, the temperature-decreasing rate is adjusted, so as to control the mixing ratio of
25 the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal of the present invention.

[0099]

Specifically, the anhydrous tetomilast type B crystal is first added to acetone-water (an acetone content of 40% to 95% by volume), and it is then dissolved therein while stirring under heating to reflux (approximately 60°C). The obtained solution is stood to cool to approximately 30°C for about 15 minutes to 1 hour. Thereafter, the suspension obtained by the aforementioned cooling is cooled at 10°C or lower, and preferably at a temperature between approximately 0°C and 10°C, for 30 minutes to 3 hours, so that the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal of the present invention can be obtained in the form of a crystal. In a case where the solution obtained after stirring under heating to reflux is quenched (for example, the solution is cooled to 10°C or lower, and preferably to a temperature between approximately 0°C and 10°C, for about 5 minutes to 1 hour), a mixture having a ratio of the anhydrous tetomilast type A crystal: the anhydrous tetomilast type B crystal = approximately 10:90 (weight ratio) can be obtained.

[0100]

The ratio of the mixture consisting of the anhydrous tetomilast type A crystal and the anhydrous tetomilast type B crystal is not particularly limited.

[0101]

In addition, the aforementioned mixture can also be produced using a novel tetomilast crystal

(excluding the mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal) instead of the known anhydrous tetomilast type B crystal. Specifically, the

5 aforementioned mixture is produced by recrystallization from a solution formed by dissolving in a solvent at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast type A crystal, an anhydrous

10 tetomilast type C crystal, and a tetomilast acetonitrile solvate crystal.

The solvent and recrystallization conditions applied herein are equivalent to those in the case of using the aforementioned known anhydrous tetomilast type B

15 crystal as a starting material.

[0102]

The obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal can be separated by isolation

20 operations such as filtration, concentration, or extraction. Moreover, after separation, the separated crystal may be subjected to a drying treatment by a known method. Furthermore, the crystal can be purified by a known purification operation.

25 [0103]

The thus obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal has a purity (the content

ratio of the anhydrous tetomilast type A crystal and the anhydrous tetomilast type B crystal to the total content) of 95% or more, and it can be crushed using an ordinary crusher (for example, an atomizer). Thus, a
5 tetomilast crushed product having a mean particle size between 10 and 50 μm and a 90% cumulative particle size of 80 μm or smaller, which is suitable for formulation, can be obtained.

[0104]

10 Pharmaceutical composition

The pharmaceutical composition of the present invention comprises at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, an anhydrous tetomilast
15 type A crystal, an anhydrous tetomilast type C crystal, and a tetomilast acetonitrile solvate crystal.

[0105]

The pharmaceutical composition of the present invention further comprises an anhydrous tetomilast
20 type B crystal. An example of such a pharmaceutical composition comprises a mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal.

[0106]

25 The novel tetomilast crystal of the present invention has activity of suppressing the release of active oxygen from neutrophils, or of eliminating reactive oxygen species. Thus, the above novel

tetomilast crystal has action to prevent generation of peroxide lipid in a living body, or to decrease such generation. Accordingly, the novel tetomilast crystal of the present invention is useful as an agent for preventing and/or treating various types of disorders or diseases that are caused by the aforementioned excessive generation of reactive oxygen species, accumulation of peroxide lipid in a living body, or a defect in defense mechanism towards these phenomena.

More specifically, an agent comprising the pharmaceutical composition of the present invention is useful, in a pharmaceutical field, as an agent for protecting various types of tissue cells from ischemia and disorders associated with revascularization, such as an agent for preventing and/or treating gastrointestinal ulcer including stress ulcer; an agent for preventing and/or treating cardiac ischemic diseases such as myocardial infarct or arrhythmia; an agent for preventing and/or treating cerebrovascular diseases such as cerebral hemorrhage, cerebral infarct, or transient ischemic attack; a liver and kidney function improver used for disorders caused by transplantation, microcirculation failure, etc.; or an agent for suppressing various types of cell injury that seem to be caused by active oxygen that is abnormally generated due to causes other than ischemia, such as an agent for preventing and/or treating Behcet's disease, cutaneous vasculitis, ulcerative colitis, malignant

rheumatism, arthritis, arteriosclerosis or diabetes.

[0107]

Moreover, the novel tetomilast crystal of the present invention is effective for various types of diseases associated with abnormal generation of cytokines, and particularly, abnormal generation of TNF- α , IL- β , IL-6, IFN- γ , etc., or various types of diseases associated with an acceleration state of adhesive action. In particular, the novel tetomilast crystal of the present invention can preferably be used: as an agent for preventing and/or treating various diseases such as chronic rheumatoid arthritis, endotoxin shock, ARDS caused by accidental ingestion of gastric juice, toxic gas, or septicemia, thermal burn or asthma, or myocardial infarction that is a myocardial ischemic state, viral myocarditis such as the acute stage of viral myocarditis, chronic heart failure such as ischemic myocardiopathy, spontaneous dilated cardiomyopathy, etc.; and as an agent for preventing and/or treating ischemic reperfusion abnormality occurring during coronary-artery bypass surgery (CABG) or during the use of artificial heart and lung, transition from SIRS (systemic inflammatory response syndrome) to organ failure (grave acute pancreatitis, DIC, etc.), or multiple organ failure caused by grave acute pancreatitis, liver failure occurring after hepatectomy for liver cancer, etc.; , inflammatory bowel diseases such as Crohn's disease,

ulcerative colitis, etc.; a series of autoimmune diseases such as hypergammaglobulinemia, systemic erythematoses (SLE) or multiple sclerosis, metastasis, immunological rejection occurring during

5 transplantation, monoclonal B cell abnormality (myeloma, etc.), polyclonal B cell abnormality, atrial myxoma, Castleman's syndrome, primary glomerulonephritis, mesangial proliferative nephritis, cancer cachexia, Lennert's lymphoma, psoriasis, atopic

10 dermatitis, Kaposi's sarcoma developed due to AIDS, postmenopausal osteoporosis, diabetes, septicemia, arteriosclerosis, or inflammatory diseases such as angitis or hepatitis or an agent for preventing and/or treating chronic obstructive pulmonary disease.

15 [0108]

Specifically, the novel tetomilast crystal of the present invention has action to improve a decrease in lung functions, such as airflow obstruction, and it exhibits an extremely high therapeutic effect upon

20 chronic obstructive pulmonary disease.

[0109]

The tetomilast crystals of the present invention can be used with at least one of the members selected from the group consisting of:

25 1. leukotriene biosynthesis inhibitors (5-lipoxygenase inhibitors and a 5-lipoxygenase activating protein (FLAP) antagonists);

2. receptor antagonists for leukotrienes

LTB₄, LTC₄, LTD₄ or LTE₄;

3. PDE4 inhibitors including inhibitors of isoform PDE4D;

4. 5-lipoxygenase inhibitors and a 5-lipoxygenase activating protein (FLAP) antagonists;

5. dual inhibitors of 5-lipoxygenase and antagonists of platelet activating factor (PAF);

6. leukotriene antagonists (LTRAs) including those for LTB₄, LTC₄, LTD₄ and LTE₄;

7. Antihistaminic H₁ receptor antagonists;

8. H₂ receptor antagonists;

9. α ₁ and α ₂ adrenoceptor agonist vasoconstrictor sympathomimetic agents administered orally or topically for decongestant use;

10. α ₁ and α ₂ adrenoceptor agonists in combination with inhibitors of 5-lipoxygenase;

11. anticholinergic agents;

12. β ₁- to β ₄- adrenoceptor agonists;

13. methylxanthines;

14. sodium cromoglicate;

15. muscarinic receptor (M₁, M₂ and M₃) antagonists;

16. NSAIDs (Non-steroid anti-inflammatory drugs) including COX-1 inhibitors, COX-2 selective inhibitors, and nitric monoxide;

17. insulin like growth factor-1 (IGF-1) and its mimetics;

18. ciclesonide,

19. inhaled glucocorticoids (of which side effects are reduced);
20. tryptase inhibitors;
21. platelet activating factor antagonists;
- 5 22. monoclonal antibodies active against endogenous inflammatory entities;
23. IPL576;
24. antitumor necrosis factor (TNF- α) agents;
25. DMARD (including leflunomide);
- 10 26. TCR peptides;
27. interleukin converting enzyme (ICE) inhibitors;
28. IMPDH inhibitors;
29. adhesion molecule inhibitors including a
- 15 VLA-4 antagonists;
30. cathepsins;
31. MAP kinase inhibitors;
32. glucose-6-phosphate dehydrogenase inhibitors;
- 20 33. kinin-B1 or kinin-B2 receptor antagonists;
34. gold in the form of an aurothio group in combination with hydrophilic groups;
35. immunosuppressive agents;
- 25 36. antigout agents;
37. xanthine oxidase inhibitors;
38. uricosuric agents;
39. antitumor agents;

45

- 40 growth hormone secretagogues;
- 41. MMP(matrix metalloproteases) inhibitors;
- 42. TGF- β (transforming growth factor);
- 43. PDGF(
5 platelet-derived growth factor);
- 44. fibroblast growth factor (for example,
basic fibroblast growth factor: b-FGF);
- 45. granulocyte-macrophage colony stimulating
factor (GM-CSF);
- 10 46. capsaicin cream;
- 47. tachykinin NK1 and NK3 receptor
antagonists;
- 48. elastase inhibitors;
- 49. PDE3 inhibitors;
- 15 50. H4 receptor antagonist or a inverse
agonists;
- 51. antioxidant agents;
- 52. radical scavenging agents;
- 53. combinations of β 2 adrenoceptor agonists
20 and glucocorticoids;
- 54. agents increasing protein level of
hypoxia-inducible factor-1 α (HIF-1 α);
- 55. antioxidant proteins upregulated by HIF-
1 α ;
- 25 56. vascular endothelial growth factor (VEGF)
secretagogus; and
- 57. VEGF receptor agonists.

[0110]

The novel tetomilast crystal of the present invention is generally used in the form of a common pharmaceutical preparation. Such a pharmaceutical preparation is prepared using commonly used diluents or
5 excipients such as a filler, an extender, a binder, a wetting agent, a disintegrator, a surfactant, or a lubricant. For such a pharmaceutical preparation, various types of forms can be selected depending on therapeutic purpose. Typical forms of such a
10 pharmaceutical preparation may include a tablet, a pill, a powder, solution, suspension, emulsion, a granule, a capsule, a suppository, and an injection (solution, suspension, etc.). When the present tetomilast crystal is molded into the form of a tablet,
15 various types of carriers, which have previously been known in the present field, can widely be used. Examples of such a carrier that can be used herein may include: excipients such as lactose, saccharose, sodium chloride, grape sugar, urea, starch, calcium
20 carbonate, kaoline, crystalline cellulose, or silica; binders such as water, ethanol, propanol, simple syrup, grape sugar in water, starch solution, gelatin solution, carboxymethyl cellulose, shellac, methylcellulose, potassium phosphate, or polyvinyl
25 pyrrolidone; disintegrators such as dry starch, sodium alginate, agar powder, laminaran powder, sodium bicarbonate, calcium carbonate, polyoxyethylene sorbitan fatty acid esters, sodium lauryl sulfate,

monoglyceride stearate, starch, or lactose;
disintegration inhibitors such as saccharose, stearin,
cacao butter, or hydrogenated oil; absorbefaciants such
as quaternary ammonium base or sodium lauryl sulfate;
5 humectant such as glycerin or starch; adsorbents such
as starch, lactose, kaoline, bentonite, or colloidal
silica; and lubricants such as purified talc, stearate,
boric acid powder, or polyethylene glycol. Such a
tablet may be further processed into a tablet coated
10 with an ordinary tablet coat, such as a sugarcoated
tablet, a gelatin-coated tablet, an enteric-coated
tablet, a film-coated tablet, a double-coated tablet,
and a multiple layer tablet, as necessary. When the
present tetomilast crystal is molded into the form of a
15 pill, carriers, which have previously been known in the
present field, can widely be used. Examples of such a
carrier that can be used herein may include:
excipients such as grape sugar, lactose, starch, cacao
butter, hydrogenated vegetable oil, kaoline, or talc;
20 binders such as gum Arabic, powdered tragacanth,
gelatin, or ethanol; and disintegrators such as
laminaran or agar. When the present tetomilast crystal
is molded into the form of a suppository, carriers,
which have previously been known in the present field,
25 can widely be used. Examples of such a carrier may
include polyethylene glycol, cacao butter, higher
alcohol, higher alcohol esters, gelatin, and
semisynthetic glyceride. As a capsule, an active

ingredient compound is generally mixed with various types of carriers as described above according to a common method, and the obtained mixture is then filled into a hard gelatin capsule, a soft capsule, etc. When
5 the present tetomilast crystal is molded into an injection, it is preferable that a liquid agent, an emulsion, and a suspension be sterilized and be isotonic to blood. When the present crystal is molded into such a form, all types of diluents, which have
10 commonly been used in the present field, can be used. Examples of such diluents that can be used herein may include water, ethyl alcohol, macrogol, propylene glycol, ethoxylated isostearyl alcohol, polyoxylated isostearyl alcohol, and polyoxyethylene sorbitan fatty
15 acid esters. In this case, it is possible that common salt, grape sugar, or glycerin, be mixed into a pharmaceutical preparation in an amount sufficient to prepare an isotonic solution. It is also possible that a common solubilizer, buffer, soothing agent, or the
20 like, be added to the pharmaceutical preparation. Further, it is also possible that a coloring agent, a preservative, an aromatic, a flavor, a sweetener, or other pharmaceuticals, be mixed into the pharmaceutical preparation, as necessary.

25 [0111]

The amount of an active ingredient compound contained in the above pharmaceutical preparation is not particularly limited, and it is appropriately

selected from a wide range. In general, approximately 1% to 70% by weight of such an active ingredient compound may be contained in the preparation.

[0112]

5 The administration method of the above pharmaceutical preparation is not particularly limited, and the pharmaceutical preparation is administered by a method that depends on various types of pharmaceutical forms, the age, sex, and other conditions of a patient,
10 the degree of disease, etc. For example, in the case of a tablet, a pill, solution, suspension, emulsion, a granule, and a capsule, these preparations are orally administered. In the case of injection, it is intravenously administered singly or as a mixture with
15 an ordinary replacement fluid such as grape sugar or amino acid. Further, as necessary, such injection is administered as a single use, intramuscularly, intradermally, subcutaneously, or intraperitoneally. A suppository is administered intrarectally.

20 The dosage of the above pharmaceutical preparation is appropriately selected depending on usage, the age, sex, and other conditions of a patient, the degree of disease, etc. In general, the amount of an active ingredient compound may be determined to be
25 approximately 0.2 to 200 mg per kg of body weight per day.

[0113]

With regard to the tetomilast hydrate

crystal, anhydrous tetomilast type A crystal, anhydrous
tetomilast type C crystal, tetomilast acetonitrile
solvate crystal, and a mixture of the above anhydrous
tetomilast type A crystal and an anhydrous tetomilast
5 type B crystal of the present invention, the crystal
forms thereof are easily controlled by
recrystallization, and these crystals are excellent in
terms of filterability. Thus, these crystals are
suitable for industrial mass production.

10 [0114]

In addition, these tetomilast crystals have
performance that is equivalent to or greater than that
of an anhydrous tetomilast type B crystal, in terms of
stability towards heat and moisture, and the
15 disintegration property and dissolution property of
tablets. Accordingly, these tetomilast crystals can
preferably be used as pharmaceutical compositions.

EXAMPLES

[0115]

20 The present invention will be described more
in detail in the following reference examples,
examples, and formulation example.

[0116]

Analytical method

25 (1) Thermogravimetry/differential thermal analysis

Thermogravimetry/differential thermal
analysis was carried out using TA60WS controller and

DTG-60A thermogravimetry/differential thermal analysis simultaneous measuring instrument manufactured by Shimadzu Corporation. Specifically, using the aforementioned apparatuses, 5 to 10 mg of a sample was
5 heated from 20°C (room temperature) to 250°C at a temperature-rising rate of 5°C/min in a dry nitrogen atmosphere. As a reference substance, α -alumina was used.

[0117]

10 (2) Powder X-ray diffraction

The powder X-ray diffraction spectrum was measured in an angle of diffraction that ranged from 3° to 40°, according to the general test method of the Pharmacopoeia of Japan, using RAD-2B diffractometer
15 (radiation source: $\text{CuK}\alpha$) manufactured by Rigaku Denki. During the measurement, voltage/current was set at 35 kV/20 mA, and the scan speed was set at 5°/min.

[0118]

The mixing ratio of a mixture consisting of
20 an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal was obtained by comparing the powder X-ray diffraction spectrum of the above mixture, with the powder X-ray diffraction spectrum of a mixture formed by mixing a pure anhydrous tetomilast type A
25 crystal and a pure anhydrous tetomilast type B crystal at various ratios.

[0119]

(3) Infrared spectroscopic analysis

The IR spectrum was measured by the KBr method.

[0120]

(4) ^1H -NMR measurement

5 The ^1H -NMR spectrum was measured in DMSO- d_6 , using TMS as a reference.

[0121]

(5) Purity measurement

The purity was measured using high
10 performance liquid chromatography (HPLC). The measurement conditions were as follows.

Sample: 0.03 g of a sample was dissolved in 80 ml of acetonitrile, and 20 ml of water was then added to the solution, so as to prepare a sample
15 solution. The measurement was carried out using 10 μl of the sample solution.

Detector: Ultraviolet photometric detector (UV 254 nm)

Column: Wakosil 5C18 HG

20 Mobile phase: Acetonitrile/10 mM Na_2SO_4 aqueous solution/phosphoric acid (500:500:1)

[0122]

(6) Particle size measurement

For the measurement of the particle size, 0.1
25 g of particles to be measured was suspended in 20 ml of n-hexane solution that contained 0.2 g of 0.1 w/v % polyoxyethylene (10) octylphenyl ether, and ultrasonication was then performed on the suspension.

Thereafter, the measurement was carried out using a particle size distribution measurement equipment (Microtrac HRA; manufactured by Microtrac).

[0123]

5 (7) Moisture value measurement

The moisture contained in the sample was measured by the Karl Fischer method.

[0124]

(8) Melting point measurement (corrected)

10 For the measurement of the melting point, the sample was heated using a heater (product name: LK6000PM; manufactured by Japan High Tech Co., Ltd.) under the condition of a temperature-rising rate of 5°C/min, and the melting state was then observed using a
15 VH-7000C microscope manufactured by Keyence Corporation.

[0125]

Reference example 1

Production of anhydrous tetomilast type B crystal

20 An anhydrous tetomilast type B crystal was produced by the methods described in (1) to (3) below.

[0126]

(1) An anhydrous tetomilast type B crystal was obtained by the method described in Journal of
25 Medicinal Chemistry, 1995, 38, pp. 353-358. That is to say, an anhydrous tetomilast type B crystal was obtained by the following method.

[0127]

First, methyl 6-[2-(3,4-diethoxyphenyl)thiazol-4-yl]pyridin-2-carboxylate (49 g; 127 mmol) and 10% sodium hydroxide (100 ml) were added to ethanol (1.4 L), and the obtained mixture was then stirred for 4 hours under heating to reflux. A majority of solvent was eliminated from the obtained solution, and water and ethyl acetate were then added to the residue for separation. The water layer obtained as a result of the separation was changed to acidic by addition of 10% hydrochloric acid, followed by extraction with ethyl acetate. Thereafter, the extract was quickly washed with a saturated sodium chloride aqueous solution, and was then dried over an excessive amount of magnesium sulfate. The obtained mixture was recrystallized from ethyl acetate, so as to obtain an anhydrous tetomilast type B crystal.

[0128]

A part of the obtained anhydrous tetomilast type B crystal melted around 175°C, and it was changed to a needle crystal. Thereafter, the above crystal completely melted (decomposed) at a temperature between 187°C and 190°C.

[0129]

The obtained anhydrous tetomilast type B crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peaks as those shown in Fig. 5 were observed.

[0130]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 6 was observed.

[0131]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that the obtained anhydrous tetomilast type B crystal had significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 762, and 706 cm^{-1} in the IR (KBr) spectrum thereof.

[0132]

(2) 5 g of an anhydrous tetomilast type A crystal obtained by the method of Example 1 as described later was dissolved in 400 ml of isopropanol while stirring under heating to reflux. The obtained solution was cooled to approximately 30°C over about 1 hour, and it was then further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so as to obtain 4.6 g of an anhydrous tetomilast type B crystal in the form of a white needle crystal (yield: 92%).

[0133]

A part of the obtained anhydrous tetomilast type B crystal melted around 175°C, and it was changed

to a needle crystal. Thereafter, the above crystal completely melted (decomposed) at a temperature between 187°C and 190°C.

[0134]

5 The obtained anhydrous tetomilast type B crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that as shown in Fig. 5, endothermic peaks were observed around 177°C and 188°C.

10 [0135]

 The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that as shown in Fig. 6, the above crystal had characteristic peaks at 2θ
15 = 4.1°, 8.1°, 11.9°, 16.1°, and 24.2°.

[0136]

 The IR(KBr) spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that the obtained anhydrous
20 tetomilast type B crystal had significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 762, and 706 cm^{-1} in the IR (KBr) spectrum thereof.

[0137]

25 (3) 10 g of an anhydrous tetomilast type A crystal obtained by the method of Example 1 as described later was dissolved in 400 ml of ethyl acetate while stirring under heating to reflux. The

obtained solution was cooled to approximately 30°C over about 1 hour, and it was then further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so as to obtain 9.3 g of an anhydrous tetomilast type B crystal in the form of a white needle crystal (yield: 93%).

[0138]

A part of the obtained anhydrous tetomilast type B crystal melted around 175°C, and it was changed to a needle crystal. Thereafter, the above crystal completely melted (decomposed) at a temperature between 187°C and 190°C.

[0139]

The obtained anhydrous tetomilast type B crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peaks as those shown in Fig. 5 were observed.

[0140]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 6 was observed.

[0141]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type B crystal was measured. As a result, it was found that the obtained anhydrous

tetomilast type B crystal had significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 762, and 706 cm^{-1} in the IR (KBr) spectrum thereof.

5 [0142]

Example 1

Production of anhydrous tetomilast type A crystal

An anhydrous tetomilast type A crystal was produced by the methods described in (1) to (7) below.

10 [0143]

(1) 5 g of the anhydrous tetomilast type B crystal obtained in Reference example 1 (3) was dissolved in a solution consisting of 140 ml of acetone and 35 ml of water, while stirring under heating to reflux. The obtained solution was cooled to 40°C over about 100 minutes (at this point, an anhydrous tetomilast type A crystal was precipitated), and it was then further cooled to 10°C or lower, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 60°C for 18 hours, so as to obtain 4.0 g of an anhydrous tetomilast type A crystal in the form of a white columnar crystal (yield: 80%).

[0144]

25 The obtained anhydrous tetomilast type A crystal melted (decomposed) at a temperature between 187°C and 189°C.

[0145]

The obtained anhydrous tetomilast type A crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that as shown in Fig. 3, an endothermic peak was observed
5 around 188°C.

[0146]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that as shown in
10 Fig. 4, the above crystal had characteristic peaks at 2θ = 10.5°, 13.1°, 18.4°, 21.9°, and 25.8°.

[0147]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a
15 result, it was found that the obtained anhydrous tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

20 [0148]

(2) 5 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) as described later was dissolved in 400 ml of ethanol while stirring under heating to reflux. The obtained solution was cooled to
25 approximately 30°C over about 1 hour, and it was then further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3

hours, so as to obtain 4.3 g of an anhydrous tetomilast type A crystal in the form of a white columnar crystal (yield: 86%).

The melting point of the obtained anhydrous tetomilast type A crystal was between 188°C and 190°C (decomposition).

[0149]

The obtained anhydrous tetomilast type A crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was observed.

[0150]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 4 was observed.

[0151]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0152]

(3) 10 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) as described later

was dissolved in 400 ml of acetone while stirring under heating to reflux. The obtained solution was cooled to approximately 30°C over about 1 hour, and it was then further cooled at 10°C or lower for 1 hour, so as to
5 obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so as to obtain 8.3 g of an anhydrous tetomilast type A crystal in the form of a white columnar crystal (yield: 83%).

10 [0153]

The obtained anhydrous tetomilast type A crystal melted (decomposed) at a temperature between 188°C and 190°C.

[0154]

15 The obtained anhydrous tetomilast type A crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was observed.

20 [0155]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 4 was observed.

25 [0156]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous

tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

5 [0157]

(4) 10 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) as described later was dissolved in a solution consisting of 320 ml of acetone and 80 ml of water, while stirring under
10 heating to reflux. The obtained solution was cooled, in a stepwise manner, to 30°C over about 3 hours, it was then cooled to 20°C over 1 hour, and it was then further cooled to 10°C over 0.5 hours, so as to obtain the precipitated crystal by filtration. The above
15 precipitated crystal was dried at 50°C for 3 hours, so as to obtain 8.3 g of an anhydrous tetomilast type A crystal in the form of a white columnar crystal (yield: 83%).

[0158]

20 The obtained anhydrous tetomilast type A crystal melted (decomposed) at a temperature between 187°C and 189°C.

[0159]

The obtained anhydrous tetomilast type A
25 crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was observed.

[0160]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same
5 spectrum as that shown in Fig. 4 was observed.

[0161]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous
10 tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0162]

15 (5) 5 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) as described later was dissolved in a solution consisting of 450 ml of acetone and 300 ml of water, while stirring under heating to reflux. The obtained solution was cooled to
20 approximately 30°C over about 1 hour, and it was then further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3
25 hours, so as to obtain 4.2 g of an anhydrous tetomilast type A crystal in the form of a white columnar crystal (yield: 84%).

[0163]

The obtained anhydrous tetomilast type A

crystal melted (decomposed) at a temperature between 188°C and 190°C.

[0164]

The obtained anhydrous tetomilast type A
5 crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was observed.

[0165]

10 The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 4 was observed.

[0166]

15 The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348,
20 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0167]

(6) 41.4 g of ethyl 3-oxo-3-(6-methoxycarbonyl-2-pyridyl)propionate was dissolved in a
25 solution consisting of 42 ml of water and 414 ml of ethyl acetate, and the obtained solution was then cooled to a temperature between 5°C and 10°C. Thereafter, a solution obtained by dissolving 35.6 g of

sulfuryl chloride in 83 ml of ethyl acetate was added dropwise to the above cooled solution over approximately 30 minutes, while stirring. Thereafter, the obtained mixture was stirred at a temperature between 10°C and 20°C for 1 hour. Subsequently, the reaction solution was heated up to approximately 90°C, while the solvent was distilled away from the reaction solution. The reaction solution was continuously heated at a temperature between approximately 90°C and 100°C for 2 hours while stirring. Thereafter, the obtained mixed suspension (containing crystals) was cooled to approximately 10°C, and it was then stirred for 1 hour, followed by filtration, so as to obtain 27.99 g of 2-(2-chloroacetyl)-6-pyridine carboxylic acid in the form of yellow-brown crystals (melting point: 184°C to 189°C; purity: 98% to 99%).

[0168]

20 g of 2-(2-chloroacetyl)-6-pyridine carboxylic acid and 22.6 g of 3,4-diethoxythiobenzamide were dissolved in a solution consisting of 100 ml of water and 200 ml of dimethoxyethane. The obtained solution was heated to reflux for 2 hours while stirring, and the reaction solution was then cooled to 5°C or lower, so as to obtain a yellow-brown precipitate by filtration.

[0169]

Subsequently, the above precipitated crystal was dissolved in a solution formed by dissolving 6.18 g

of potassium hydroxide in 372 ml of water. The obtained solution was extracted with ethyl acetate twice (186 ml × 2). Thereafter, 1 g of activated carbon was added to the separated water layer, and the
5 obtained solution was then stirred at approximately 30°C for 30 minutes. Thereafter, the activated carbon was removed by filtration, and 372 ml of acetone and 11.2 g of concentrated hydrochloric acid were then added to the obtained filtrate, so as to obtain a suspension (a
10 mixture consisting of a tetomilast hydrate crystal and an anhydrous tetomilast type B crystal). In order to transfer the mixture consisting of a tetomilast hydrate crystal and an anhydrous tetomilast type B crystal to an anhydrous tetomilast type A crystal, the above
15 suspension was heated at 60°C for 30 minutes, and it was then cooled to room temperature, so as to obtain a crystal by filtration, thereby obtaining an anhydrous tetomilast type A crude crystal (34.82 g; wet state).
[0170]

20 8.67 g of the crude crystal was dissolved in a solution consisting of 213 ml of acetone and 53 ml of water by heating at 60°C without drying it, followed by filtration during a hot state. Subsequently, the obtained filtrate was heated again, and dissolution of
25 the crystal was then confirmed. Thereafter, the filtrate was cooled to 50°C. 79 mg of an anhydrous tetomilast type A crystal was added as a seed crystal to the cooled filtrate, and the obtained mixture was

then stirred at a temperature between 42°C and 50°C (internal temperature) for 2 hours. Thereafter, the obtained solution was cooled to 20°C over approximately 20 minutes, and it was then stirred at a temperature between 19°C and 25°C for 2 hours. Thereafter, the solution was cooled to 5°C over 35 minutes, and it was then stirred at a temperature between 4°C and 5°C for 2 hours, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 80°C overnight, so as to obtain 7.25 g of an anhydrous tetomilast type A crystal (whose yield was 78.4% when 6-chloroacetyl-2-pyridine carboxylic acid was used as a reference). The HPLC purity of the obtained anhydrous tetomilast type A crystal was 99.9%.

[0171]

A part of the obtained anhydrous tetomilast type A crystal melted around 175°C, and it was changed to a needle crystal. Thereafter, the above crystal completely melted (decomposed) at a temperature between 187°C and 190°C.

[0172]

The obtained anhydrous tetomilast type A crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was observed.

[0173]

The powder X-ray diffraction spectrum of the

obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same spectrum as that shown in Fig. 4 was observed.

[0174]

5 The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348,
10 1271, 1132, 1045, 758, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0175]

 The obtained anhydrous tetomilast type A crystal was crushed with an atomizer, so as to obtain
15 powders having a mean particle size of 30.4 μm and a 90% cumulative particle size of 57 μm .

[0176]

(7) 32.36 g of the tetomilast hydrate crude crystal obtained in Example 5 (2) as described later
20 was dissolved in a solution consisting of 197 ml of purified water and 793 ml of acetone by heating at approximately 60°C, followed by filtration during a hot state. Subsequently, the obtained filtrate was heated again, and dissolution of the crystal was then
25 confirmed. Thereafter, the filtrate was cooled to 45°C. 290 mg of an anhydrous tetomilast type A crystal was added to the cooled filtrate, and the obtained mixture was then stirred at 45°C for 2 hours. Thereafter, the

obtained solution was cooled to 20°C over about 1 hour, and it was then stirred at a temperature between 20°C and 24°C for 2 hours. Thereafter, the solution was cooled to 5°C over about 2 hours, and it was then
5 stirred at a temperature between -1°C and 5°C for 2 hours, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 80°C for 4 hours, so as to obtain 24.11 g of an anhydrous tetomilast type A crystal.

10 [0177]

The obtained anhydrous tetomilast type A crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that the same endothermic peak as that shown in Fig. 3 was
15 observed.

[0178]

The powder X-ray diffraction spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the same
20 spectrum as that shown in Fig. 4 was observed.

[0179]

The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous
25 tetomilast type A crystal had significant infrared absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758, and 704 cm⁻¹ in the IR (KBr) spectrum thereof.

[0180]

Example 2

Production of anhydrous tetomilast type C crystal

5 g of the anhydrous tetomilast type A
5 crystal obtained in Example 1 (6) was dissolved in 500
ml of methanol, while stirring under heating to reflux.
The obtained solution was cooled to approximately 30°C
over about 1 hour, and it was then further cooled at
10°C or lower for 1 hour, so as to obtain the
10 precipitated crystal by filtration. The above
precipitated crystal was dried at 50°C for 3 hours, so
as to obtain 3.8 g of an anhydrous tetomilast type C
crystal in the form of a white platy crystal (yield:
76%).

15 [0181]

A needle crystal was generated from the
crystal surface of the obtained anhydrous tetomilast
type C crystal at 184°C, and the above crystal melted
(decomposed) at a temperature between 187°C and 190°C.

20 [0182]

The obtained anhydrous tetomilast type C
crystal was subjected to thermogravimetry/differential
thermal analysis. As a result, it was found that as
shown in Fig. 7, endothermic peaks were observed around
25 184°C and around 189°C.

[0183]

The powder X-ray diffraction spectrum of the
obtained anhydrous tetomilast type C crystal was

measured. As a result, it was found that as shown in Fig. 8, the above crystal had characteristic peaks at $2\theta = 4.2^\circ, 8.2^\circ, 12.0^\circ, 16.4^\circ, 24.7^\circ,$ and 25.9° .

[0184]

5 The IR(KBr) spectrum of the obtained anhydrous tetomilast type A crystal was measured. As a result, it was found that the obtained anhydrous tetomilast type A crystal had significant infrared absorption bands at 3300, 3088, 1744, 1593, 1476, 1346,
10 1267, 1132, 1045, 754, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0185]

Example 3

Production of tetomilast acetonitrile solvate crystal

15 5 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) was dissolved in 400 ml of acetonitrile, while stirring under heating to reflux. The obtained solution was cooled to approximately 30°C over about 1 hour, and it was then
20 further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so as to obtain 5.1 g of a tetomilast acetonitrile solvate crystal in the form of a while
25 platy crystal (yield: quantitative).

[0186]

The obtained tetomilast acetonitrile solvate crystal became clouded at 90°C , and the above crystal

melted (decomposed) at a temperature between 187°C and 190°C.

[0187]

The obtained tetomilast acetonitrile solvate
5 crystal was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that as shown in Fig. 9, endothermic peaks were observed around 91°C, around 176°C, and around 189°C.

[0188]

10 The powder X-ray diffraction spectrum of the obtained tetomilast acetonitrile solvate crystal was measured. As a result, it was found that as shown in Fig. 10, the above crystal had characteristic peaks at $2\theta = 3.6^\circ$, 7.1° , 10.6° , 14.2° , and 24.8° .

15 [0189]

The IR(KBr) spectrum of the obtained tetomilast acetonitrile solvate crystal was measured. As a result, it was found that the obtained tetomilast acetonitrile solvate crystal had significant infrared
20 absorption bands at 3300, 3090, 2249 (nitrile group), 1744, 1593, 1476, 1346, 1269, 1132, 1045, 752, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0190]

The NMR (DMSO- d_6) spectrum of the obtained
25 tetomilast acetonitrile solvate crystal was measured. As a result, it was found that a methyl group peak of acetonitrile was observed at $\delta 2.1$ ppm.

[0191]

Example 4

Production of mixture consisting of anhydrous tetomilast type A crystal and anhydrous tetomilast type B crystal

5 A mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal was produced by the method described in (1) or (2) below:

[0192]

10 (1) 10 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) was dissolved in a solution consisting of 320 ml of acetone and 80 ml of water, while stirring under heating to reflux. The obtained solution was cooled to approximately 30°C over
15 about 1 hour, and it was then cooled to 10°C over about 10 minutes. Thereafter, the solution was further cooled at 10°C or lower for 1 hour, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so
20 as to obtain 8.5 g of a mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 40:60) in the form of a white columnar crystal (yield: 85%).

[0193]

25 Only a small part of the obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal melted at 178°C, and it was crystallized in the form of a needle

crystal. Thereafter, the mixture melted (decomposed) at a temperature between 188°C and 190°C.

[0194]

The obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 40:60) was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that as shown in Fig. 11, endothermic peaks were observed around 175°C and around 189°C.

[0195]

In addition, the powder X-ray diffraction spectrum of the obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 40:60) was measured. As a result, it was found that as shown in Fig. 12, the above mixture had characteristic peaks at $2\theta = 4.2^\circ$, 11.9° , 13.2° , 16.2° , 17.3° , 24.3° , 25.3° , 25.9° , and 27.5° .

[0196]

Moreover, the IR(KBr) spectrum of the obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 40:60) was measured. As a result, it was found that the obtained mixture had significant infrared absorption bands at 3298, 3088, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 760, and 704 cm^{-1} in the IR (KBr) spectrum thereof.

[0197]

(2) 10 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) was dissolved in a solution consisting of 320 ml of acetone and 80 ml of water, while stirring under heating to reflux. The obtained solution was quenched to 10°C over 30 minutes, so as to obtain the precipitated crystal by filtration. The above precipitated crystal was dried at 50°C for 3 hours, so as to obtain 7.8 g of a mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 10:90) in the form of white powders (yield: 78%). A part of the obtained mixture melted at 176°C, and it was crystallized in the form of a needle crystal. Thereafter, the mixture melted (decomposed) at a temperature between 187°C and 190°C.

[0198]

The obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 10:90) was subjected to thermogravimetry/differential thermal analysis. As a result, it was found that as shown in Fig. 13, endothermic peaks were observed around 176°C and around 189°C.

[0199]

The powder X-ray diffraction spectrum of the obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B

crystal (A:B = 10:90) was measured. As a result, it was found that as shown in Fig. 14, the above mixture had characteristic peaks at $2\theta = 4.1^\circ, 11.9^\circ, 16.1^\circ, 17.2^\circ, 19.3^\circ, 24.2^\circ, 25.1^\circ, 25.9^\circ,$ and 27.3° .

5 [0200]

The IR(KBr) spectrum of the obtained mixture consisting of an anhydrous tetomilast type A crystal and an anhydrous tetomilast type B crystal (A:B = 10:90) was measured. As a result, it was found that
10 the obtained mixture had significant infrared absorption bands at 3298, 3090, 1744, 1593, 1474, 1348, 1269, 1132, 1045, 756, and 706 cm^{-1} in the IR (KBr) spectrum thereof.

[0201]

15 Example 5

Production of tetomilast monohydrate crystal

A tetomilast monohydrate crystal was produced by the methods described in (1) to (3) below:

[0202]

20 (1) 18.7 g of 2-(2-chloroacetyl)-6-pyridine carboxylic acid and 21.1 g of 3,4-diethoxythiobenzamide were dissolved in a solution consisting of 94 ml of water and 187 ml of dimethoxyethane, and the obtained mixture was then stirred at approximately 80°C (reflux)
25 for 2 hours. The obtained solution was cooled to 5°C , and it was then stirred for 1 hour, so as to obtain a yellow-brown precipitate by filtration. The above precipitate was dissolved in a solution formed by

dissolving 5.78 g of potassium hydroxide in 348 ml of water. The obtained mixture solution was washed with 174 ml of ethyl acetate twice. Thereafter, a solution formed by suspending 0.9 g of activated carbon in 1.9
5 ml of water was added to the separated water layer, and the obtained mixture was then stirred at a temperature between 30°C and 31°C for 30 minutes. Thereafter, the activated carbon was removed by filtration, and 348 ml of acetone was then added to the obtained filtrate.
10 Subsequently, 10.4 g of concentrated hydrochloric acid was added thereto while stirring, and the obtained mixture was then stirred for 1 hour. Thereafter, the precipitated crystal was collected by filtration. This crystal was suspended in 348 ml of water, and the
15 obtained mixture was then stirred at a temperature between 27°C and 30°C for 30 minutes. Thereafter, the obtained crystal was collected by filtration. The obtained crystal was washed with a solution consisting of 35 ml of acetone and 35 ml of water, so as to obtain
20 29.53 g of a tetomilast monohydrate crystal.

[0203]

The obtained tetomilast monohydrate crystal was subjected to thermogravimetry/differential thermal analysis. As a result, as shown in Fig. 1, an
25 endothermic peak was observed around 189°C, and further, a wide peak was observed around 102°C.

[0204]

The powder X-ray diffraction spectrum of the

obtained tetomilast monohydrate crystal was measured. As a result, as shown in Fig. 2, it was found that the above crystal had characteristic peaks at $2\theta = 10.6^\circ$, 12.9° , 21.1° , 22.3° , and 25.0° .

5 [0205]

The IR(KBr) spectrum of the obtained tetomilast monohydrate crystal was measured. As a result, it was found that the obtained crystal had significant infrared absorption bands at 3516, 3433, 10 1742, 1709, 1587, 1472, 1267, 1143, 1040, 758, and 716 cm^{-1} in the IR (KBr) spectrum thereof.

[0206]

(2) 50 g of the anhydrous tetomilast type A crystal obtained in Example 1 (6) was dissolved in a 15 solution formed by dissolving 8.33 g of potassium hydroxide in 500 ml of water. This solution was filtrated, and 500 ml of acetone was then added to the obtained filtrate. Thereafter, 13 ml (1.1 eq) of concentrated hydrochloric acid was added thereto while 20 stirring (The product precipitated at that time was an anhydrous tetomilast type B crystal). The obtained solution was stirred at room temperature for approximately 10 minutes. After completion of the stirring, 2.5 g of the tetomilast monohydrate crystal 25 obtained in the aforementioned Example 5 (1) was added as a seed crystal to the resultant, and the obtained mixture was continuously stirred for 2 hours (During such stirring, transformation progressed via the

solvent. If the stirring time was short, a mixture consisting of a tetomilast hydrate crystal and an anhydrous tetomilast type B crystal was obtained.) The precipitated crystal was collected by filtration, and
5 the obtained crystal was then suspended in 400 ml of water. The obtained mixture was stirred at a temperature between 20°C and 30°C for 30 minutes. Thereafter, the crystal was collected by filtration, and it was then washed with 80 ml of acetone-water (an
10 acetone content of 50% by volume). The resultant was dried under reduced pressure all night, so as to obtain 51.5 g of a tetomilast monohydrate crystal.

[0207]

The obtained tetomilast monohydrate crystal
15 was subjected to thermogravimetry/differential thermal analysis. As a result, the same endothermic peak as that shown in Fig. 1 was obtained.

[0208]

The powder X-ray diffraction spectrum of the
20 obtained tetomilast monohydrate crystal was measured. As a result, the same spectrum as that shown in Fig. 2 was observed.

[0209]

The IR(KBr) spectrum of the obtained
25 tetomilast monohydrate crystal was measured. As a result, it was found that the obtained crystal had significant infrared absorption bands at 3516, 3433, 1742, 1709, 1587, 1472, 1267, 1143, 1040, 758, and 716

cm⁻¹ in the IR (KBr) spectrum thereof.

[0210]

(3) 2.37 g of the anhydrous tetomilast type B crystal obtained in Reference example 1(3) was
5 suspended in a solution consisting of 50 ml of acetone and 50 ml of water, and the obtained mixture was then stirred for approximately 5 minutes. Thereafter, a tetomilast hydrate crystal was added as a seed crystal to the above mixture, and the thus obtained mixture was
10 further stirred at 30°C for 1 hour. The crystal was collected by filtration, and it was then dried at 60°C all night, so as to obtain 2.34 g of a tetomilast hydrate crystal.

[0211]

15 The moisture value of the obtained tetomilast hydrate crystal was 4.68%. This value was almost the same as the theoretical moisture value (4.64%) of a tetomilast monohydrate crystal.

[0212]

20 The above monohydrate became clouded around 100°C, and melted (decomposed) at a temperature between 188°C and 189°C.

[0213]

The obtained tetomilast monohydrate crystal
25 was subjected to thermogravimetry/differential thermal analysis. As a result, the same endothermic peak as that shown in Fig. 1 was obtained.

[0214]

The powder X-ray diffraction spectrum of the obtained tetomilast monohydrate crystal was measured. As a result, the same spectrum as that shown in Fig. 2 was observed.

5 [0215]

The IR(KBr) spectrum of the obtained tetomilast monohydrate crystal was measured. As a result, it was found that the obtained crystal had significant infrared absorption bands at 3516, 3433, 10 1742, 1709, 1587, 1472, 1267, 1143, 1040, 758, and 716 cm^{-1} in the IR (KBr) spectrum thereof.

[0216]

Example 6

Thermal transformation from anhydrous tetomilast type B 15 crystal to anhydrous tetomilast type A crystal

Thermal transformation from an anhydrous tetomilast type B crystal to an anhydrous tetomilast type A crystal was confirmed by the means described in (1) to (3) below:

20 [0217]

(1) The anhydrous tetomilast type B crystal obtained in Reference example 1 (3) was left at a temperature between 20°C and 30°C for approximately 1 year, and thereafter, the crystal form thereof was 25 examined by powder X-ray diffraction. As a result, it was confirmed that the above crystal maintained the form as an anhydrous tetomilast type B crystal.

[0218]

(2) 5 g of the anhydrous tetomilast type B crystal obtained in Reference example 1 (3) was suspended in a solution consisting of 40 ml of acetone and 10 ml of water, and the obtained mixture was then stirred at 20°C. At that time, sampling was carried out at intervals of 15 minutes, 30 minutes, 60 minutes, and 120 minutes, and each of the obtained samples was subjected to powder X-ray diffraction to examine the crystal form. As a result, it could be confirmed that each of the samples collected after stirring for 15 minutes, 30 minutes, and 60 minutes, maintained the form as an anhydrous tetomilast type B crystal.

On the other hand, it could be confirmed that the sample collected after stirring for 120 minutes was a mixture consisting of an anhydrous tetomilast type B crystal and an anhydrous tetomilast type A crystal (A:B = 70:30).

[0219]

(3) 5 g of the anhydrous tetomilast type B crystal obtained in Reference example 1 (3) was suspended in a solution consisting of 40 ml of acetone and 10 ml of water, and the obtained mixture was then stirred at 40°C. At that time, sampling was carried out at intervals of 15 minutes, 30 minutes, 60 minutes, and 120 minutes, and each of the obtained samples was subjected to powder X-ray diffraction to examine the crystal form. As a result, it could be confirmed that the sample collected after stirring for 15 minutes was

a mixture consisting of an anhydrous tetomilast type B crystal and an anhydrous tetomilast type A crystal (A:B = 50:50).

[0220]

5 On the other hand, it could be confirmed that each of the samples that had been stirred for 30 minutes, 60 minutes, and 120 minutes, was changed to an anhydrous tetomilast type A crystal.

[0221]

10 When tetomilast crystals with different crystal forms are obtained depending on the type of a solvent used, the obtained crystal forms do not depend on the crystal forms of tetomilast crystals used as raw materials. Thus, an example of using an anhydrous
15 tetomilast type A crystal (purity of 99.9%) was given herein. When a novel tetomilast crystal other than the anhydrous tetomilast type A crystal was used as well, the same results were obtained.

[0222]

20 Formulation example

A table comprising 5 mg of an anhydrous tetomilast type C crystal, 132 mg of starch, 18 mg of magnesium stearate and 45 mg of lactose, in a single tablet, was produced by a common method.

25711-857

CLAIMS:

1. An anhydrous tetomilast type A crystal having a powder X-ray diffraction spectrum that is the same as the powder X-ray diffraction spectrum shown in Fig. 4.
- 5 2. An anhydrous tetomilast type A crystal having a powder X-ray diffraction spectrum with peaks at 10.5, 13.1, 18.4, 21.9 and 25.8 degrees 2θ .
3. An anhydrous tetomilast type A crystal as defined in claim 1 or 2, having an infrared KBr spectrum with absorption bands at 3306, 3084, 1746, 1593, 1474, 1348, 1271, 1132, 1045, 758 and 704 cm^{-1} .
- 10 4. An anhydrous tetomilast type A crystal as defined in claim 1 or 2, having a thermogravimetry/differential thermal analysis spectrum with an endothermic peak at around 188°C .
5. An anhydrous tetomilast type A crystal as defined in claim 1 or 2, having a thermogravimetry/differential thermal analysis spectrum the same as the
15 thermogravimetry/differential thermal analysis spectrum shown in Fig. 3.
6. A mixture consisting of an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5, and an anhydrous tetomilast type B crystal wherein said anhydrous tetomilast type B crystal has a powder X-ray diffraction spectrum having characteristic peaks at $2\theta=4.1^{\circ}$, 8.1° , 11.9° , 16.1° and 24.2° .
- 20 7. A pharmaceutical composition comprising an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5 and a pharmaceutically acceptable diluent.
8. A pharmaceutical composition comprising an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5 and an anhydrous tetomilast type B crystal and a pharmaceutically acceptable diluent wherein said anhydrous tetomilast type B

25711-857

85

crystal has a powder X-ray diffraction spectrum having characteristic peaks at $2\theta=4.1^\circ$, 8.1° , 11.9° , 16.1° and 24.2° .

9. The pharmaceutical composition according to claim 7 or 8, which is for use in the prevention and/or treatment of: gastrointestinal ulcer, cardiac ischemic
5 disease, cerebrovascular disease, Behcet's disease, cutaneous vasculitis, ulcerative colitis, malignant rheumatism, arthritis, arteriosclerosis or diabetes; or for use in improving liver or kidney function in respect of a disorder caused by transplantation or microcirculation failure.

10. The pharmaceutical composition according to claim 7 or 8, which is for
10 use in the prevention and/or treatment of: chronic rheumatoid arthritis, endotoxin shock, ARDS, thermal burn, asthma, chronic heart failure, myocardial infarction, viral myocarditis, ischemic reperfusion abnormality, transition from SIRS (systemic inflammatory response syndrome) to organ failure, multiple organ failure, inflammatory bowel disease, autoimmune disease, metastasis, immunological rejection occurring
15 during transplantation, monoclonal B cell abnormality, polyclonal B cell abnormality, atrial myxoma, Castleman's syndrome, primary glomerulonephritis, mesangial proliferative nephritis, cancer cachexia, Lennert's lymphoma, psoriasis, atopic dermatitis, Kaposi's sarcoma developed due to AIDS, postmenopausal osteoporosis, septicemia, inflammatory disease or chronic obstructive pulmonary disease.

20 11. The pharmaceutical composition according to claim 10, wherein the inflammatory bowel disease is ulcerative colitis.

12. The pharmaceutical composition according to claim 10, wherein the inflammatory bowel disease is Crohn's disease.

13. The pharmaceutical composition according to claim 10, which is for use
25 in the prevention and/or treatment of chronic obstructive pulmonary disease.

25711-857

14. A process for the preparation of an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5, wherein the process comprises recrystallisation from a solution formed by dissolving an anhydrous tetomilast type B crystal in ethanol or acetone, wherein said anhydrous tetomilast type B crystal has a powder X-ray
5 diffraction spectrum having characteristic peaks at $2\theta=4.1^\circ$, 8.1° , 11.9° , 16.1° and 24.2° .

15. A process for the preparation of an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5, wherein the process comprises dissolving an anhydrous tetomilast type B crystal in a mixture of acetone and water wherein the acetone content is 40% or more by volume, followed by reacting the obtained solution
10 at 40 to 50°C for 60 minutes or longer, and then cooling the solution in order to obtain the anhydrous tetomilast type A crystal, wherein said anhydrous tetomilast type B crystal has a powder X-ray diffraction spectrum having characteristic peaks at $2\theta=4.1^\circ$, 8.1° , 11.9° , 16.1° and 24.2° .

16. A process for the preparation of an anhydrous tetomilast type A crystal
15 as defined in claim 1, 2, 3, 4 or 5, wherein the process comprises stirring from a suspension formed by suspending at least one type of tetomilast crystal selected from the group consisting of a tetomilast hydrate crystal, anhydrous tetomilast type B crystal, an anhydrous tetomilast type C crystal, and a tetomilast acetonitrile solvate crystal in a mixed solvent consisting of water, and at least one organic solvent
20 selected from the group consisting of methanol, ethanol, acetone, and tetrahydrofuran, wherein the temperature of suspension is between 0 and 65°C and stirring time in between 10 minutes and 48 hours, wherein said tetomilast hydrate crystal has a powder X-ray diffraction spectrum having characteristic peaks at $2\theta=10.6^\circ$, 12.9° , 21.1° , 22.3° and 25.0° wherein said anhydrous tetomilast type B
25 crystal has a powder x-ray diffraction spectrum having characteristic peaks at $2\theta=4.1^\circ$, 8.1° , 11.9° , 16.1° and 24.2° wherein said anhydrous tetomilast type C crystal has a powder X-ray diffraction spectrum having characteristic peaks at $2\theta=4.2^\circ$, 8.2° , 12.0° , 16.4° , 24.7° and 25.9° wherein said tetomilast acetonitrile solvate crystal has a

25711-857

87

powder X-ray diffraction spectrum having characteristic peaks at $2\theta=3.6^\circ$, 7.1° , 10.9° , 14.2° and 24.8° .

17. Use of an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4 or 5 for the prevention and/or treatment of: gastrointestinal ulcer, cardiac

5 ischemic disease or cerebrovascular disease, Behcet's disease, cutaneous vasculitis, ulcerative colitis, malignant rheumatism, arthritis, arteriosclerosis or diabetes; or for improving liver or kidney function in respect of a disorder caused by transplantation or microcirculation failure.

18. Use of an anhydrous tetomilast type A crystal as defined in claim 1, 2, 3, 4

10 or 5 for the prevention and/or treatment of: chronic rheumatoid arthritis, endotoxin shock, ARDS, thermal burn, asthma, chronic heart failure, myocardial infarction, viral myocarditis, ischemic reperfusion abnormality, transition from SIRS (systemic inflammatory response syndrome) to organ failure, multiple organ failure, inflammatory bowel disease, autoimmune disease, metastasis, immunological rejection occurring
15 during transplantation, monoclonal B cell abnormality, polyclonal B cell abnormality, atrial myxoma, Castleman's syndrome, primary glomerulonephritis, mesangial proliferative nephritis, cancer cachexia, Lennert's lymphoma, psoriasis, atopic dermatitis, Kaposi's sarcoma developed due to AIDS, postmenopausal osteoporosis, septicemia, inflammatory disease or chronic obstructive pulmonary disease.


20 19. Use according to claim 18, wherein the inflammatory bowel disease is ulcerative colitis.

20. Use according to claim 18, wherein the inflammatory bowel disease is Crohn's disease.

21. Use according to claim 18, which is for the prevention and/or treatment
25 of chronic obstructive pulmonary disease.


1/14

FIG.1




2/14

FIG.2



3/14


FIG.3



4/14

FIG.4

PEAK WIDTH THRESHOLD : 0.20
PEAK STRENGTH THRESHOLD : 300.000



5/14

FIG.5





FIG.6

PEAK WIDTH THRESHOLD : 0.20
PEAK STRENGTH THRESHOLD : 300.000



7/14

FIG.7





FIG.8

PEAK WIDTH THRESHOLD : 0.20
PEAK STRENGTH THRESHOLD : 300.000



9/14

FIG.9





FIG.10

PEAK WIDTH THRESHOLD : 0.20
PEAK STRENGTH THRESHOLD : 300.000



11/14

FIG.11





FIG.12



13/14

FIG.13




FIG.14

PEAK WIDTH THRESHOLD : 0.30
PEAK STRENGTH THRESHOLD : 300.000

