



(51) International Patent Classification:

C07D 487/04 (2006.01) *C12Q 1/66* (2006.01)
A61K 49/00 (2006.01) *G01N 21/76* (2006.01)

(21) International Application Number:

PCT/US2017/017761

(22) International Filing Date:

14 February 2017 (14.02.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:


62/295,363 15 February 2016 (15.02.2016) US

(71) Applicant: **PROMEGA CORPORATION** [US/US];
2800 Woods Hollow Road, Madison, WI 53711-5399
(US).(72) Inventors: **SHAKHMIN, Anton**; 1534 Brighton Avenue,
Apt. 2, Grover Beach, CA 93433 (US). **KIRKLAND,
Thomas**; 13350 San Antonio Road, Atascadero, CA 93422
(US). **WALKER, Joel**; 1445 Prefumo Canyon Road, Unit
3, San Luis Obispo, CA 93405 (US). **MACHLEIDT,
Thomas**; 6401 Landfall Drive, Madison, WI 53505 (US).
HALL, Mary; 504 Skyview Drive, Waunakee, WI 53597
(US). **WOOD, Keith, V.**; 8380 Swan Road, Mount Horeb,
WI 53572 (US).(74) Agent: **HOFMEISTER, Todd, S.**; Michael Best &
Friedrich LLP, 100 East Wisconsin Avenue, Suite 3300,
Milwaukee, WI 53202-4108 (US).(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
ZA, ZM, ZW.(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COELENTERAZINE ANALOGUES



(I),

(57) Abstract: Described are coelenterazine analogues of formula (I), methods for making the analogues, kits comprising the analogues, and methods of using the compounds for the detection of luminescence in luciferase-based assays.

COELENTERAZINE ANALOGUES

CROSS-REFERNECE TO RELATED APPLICATIONS

[0001] This application claims priority to U.S. Provisional Patent Application No. 62/295,363, filed on February 15, 2016, the entire contents of which are incorporated herein by reference.

TECHNICAL FIELD

[0002] The present disclosure relates to coelenterazine analogues, methods for making coelenterazine analogues, and methods of using coelenterazine analogues in luciferase-based assays.

BACKGROUND


[0002] Bioluminescent assays are used extensively in the investigation of cellular physiology, especially processes associated with gene expression. In particular, luciferase reporter enzymes are quite valuable tools in this field, and, to date, there has been intense protein engineering to obtain small and environmentally insensitive luciferases that may be useful in bioluminescent assays. There exist a number of efficient luciferase reporters enabling whole-cell biosensor measurements, drug discovery through high-throughput screening, and in vivo imaging, which also permits the study of protein-protein interactions in living cells, apoptosis, and cell viability. Luciferases that use coelenterazine and coelenterazine analogues as substrates are among the most widely used systems due to their brightness and acceptance in whole cell applications.

SUMMARY OF THE INVENTION

[0003] Many known coelenterazine analogues have deficiencies, which limit their effectiveness as luciferase substrates and usefulness in luciferase-based luminescence assays. These deficiencies include cell toxicity, light sensitivity, thermodynamic instability, low aqueous solubility, and low cell permeability. Accordingly, there exists a need for

coelenterazine analogues with improved properties and methods for synthesizing the analogues.

[0004] In one aspect, disclosed are compounds of formula (I),



(I),

or tautomers, or pharmaceutically acceptable salts thereof, wherein R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and q is 0-2; wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently substituted or unsubstituted.

[0005] Also disclosed are methods of making the compounds, kits comprising the compounds, and methods of using the compounds as luciferase substrates in luciferase-based luminescence assays.

BRIEF DESCRIPTION OF THE DRAWINGS

[0006] FIG. 1 is a graph depicting cellular bioluminescent activity of exemplary compounds in HEK293 cells in comparison to coelenterazine.

[0007] FIG. 2 is a graph depicting the rank ordering of compounds of formula (I) by signal strength relative to coelenterazine.

[0008] FIG. 3 and FIG. 4 are graphs depicting results of a cell viability assay employing exemplary compounds in HeLa cells.

[0009] FIG. 5 and FIG. 6 are graphs depicting results of a cell viability assay employing exemplary compounds in HEK293 cells.

DETAILED DESCRIPTION

[0010] Disclosed herein are coelenterazine analogues. The coelenterazine analogues can be compounds of formula (I) and useful substrates of proteins that utilize coelenterazine to produce luminescence, including, but not limited to, luciferases and photoproteins found in various marine organisms such as cnidarians (e.g., *Renilla* luciferase), jellyfish (e.g., aequorin from the *Aequorea* jellyfish) and decapods luciferases (e.g., luciferase complex of *Oplophorus gracilirostris*). In comparison to coelenterazine, compounds of formula (I) may have at least one of improved water solubility, improved stability, improved cell permeability, increased biocompatibility with cells, reduced autoluminescence, and reduced toxicity.

[0011] Also disclosed herein are methods of making coelenterazine analogues [compounds of formula (I)]. Disclosed are three robust and versatile approaches towards the preparation of coelenterazine analogues. The methods allow for the preparation of analogues that could not have been prepared using existing synthetic methods. The described methodology enables access to a variety of substituents at the R¹ position and can be performed under mild conditions utilizing a wide variety of readily available starting materials. The disclosed synthetic methodology unexpectedly provides a variety of new applications and advancements in bioluminescence technology based on coelenterazine analogues.

1. Definitions

[0012] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. In case of conflict, the present document, including definitions, will control. Preferred methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting.

[0013] The terms “comprise(s),” “include(s),” “having,” “has,” “can,” “contain(s),” and variants thereof, as used herein, are intended to be open-ended transitional phrases, terms, or words that do not preclude the possibility of additional acts or structures. The singular forms “a,” “an” and “the” include plural references unless the context clearly dictates otherwise.

The present disclosure also contemplates other embodiments “comprising,” “consisting of” and “consisting essentially of,” the embodiments or elements presented herein, whether explicitly set forth or not.

[0014] The modifier “about” used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context (for example, it includes at least the degree of error associated with the measurement of the particular quantity). The modifier “about” should also be considered as disclosing the range defined by the absolute values of the two endpoints. For example, the expression “from about 2 to about 4” also discloses the range “from 2 to 4.” The term “about” may refer to plus or minus 10% of the indicated number. For example, “about 10%” may indicate a range of 9% to 11%, and “about 1” may mean from 0.9-1.1. Other meanings of “about” may be apparent from the context, such as rounding off, so, for example “about 1” may also mean from 0.5 to 1.4.

[0015] Definitions of specific functional groups and chemical terms are described in more detail below. For purposes of this disclosure, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in *Organic Chemistry*, Thomas Sorrell, University Science Books, Sausalito, 1999; Smith and March *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd Edition, Cambridge University Press, Cambridge, 1987; the entire contents of each of which are incorporated herein by reference.

[0016] The term “alkoxy” as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy and tert-butoxy.

[0017] The term “alkyl” as used herein, means a straight or branched, saturated hydrocarbon chain containing from 1 to 10 carbon atoms. The term “lower alkyl” or “C₁-C₆-alkyl” means a straight or branched chain hydrocarbon containing from 1 to 6 carbon atoms. The term “C₁-C₃-alkyl” means a straight or branched chain hydrocarbon containing from 1 to 3 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl, isopentyl,

neopentyl, *n*-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, *n*-heptyl, *n*-octyl, *n*-nonyl, and *n*-decyl.

[0018] The term “alkenyl” as used herein, means a hydrocarbon chain containing from 2 to 10 carbon atoms with at least one carbon-carbon double bond. The alkenyl group may be substituted or unsubstituted. For example, the alkenyl group may be substituted with an aryl group, such as a phenyl.

[0019] The term “alkynyl” as used herein, means a hydrocarbon chain containing from 2 to 10 carbon atoms with at least one carbon-carbon triple bond. The alkynyl group may be substituted or unsubstituted. For example, the alkynyl group may be substituted with an aryl group, such as a phenyl.

[0020] The term “alkoxyalkyl” as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

[0021] The term “alkylene”, as used herein, refers to a divalent group derived from a straight or branched chain hydrocarbon of 1 to 10 carbon atoms, for example, of 2 to 5 carbon atoms. Representative examples of alkylene include, but are not limited to, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, and -CH₂CH₂CH₂CH₂CH₂-.

[0022] The term “aryl” as used herein, refers to a phenyl group, or bicyclic aryl or tricyclic aryl fused ring systems. Bicyclic fused ring systems are exemplified by a phenyl group appended to the parent molecular moiety and fused to a phenyl group. Tricyclic fused ring systems are exemplified by a phenyl group appended to the parent molecular moiety and fused to two other phenyl groups. Representative examples of bicyclic aryls include, but are not limited to, naphthyl. Representative examples of tricyclic aryls include, but are not limited to, anthracenyl. The monocyclic, bicyclic, and tricyclic aryls are connected to the parent molecular moiety through any carbon atom contained within the rings, and can be unsubstituted or substituted.

[0023] The term “cycloalkyl” as used herein, refers to a carbocyclic ring system containing three to ten carbon atoms, zero heteroatoms and zero double bonds. Representative examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl and cyclodecyl.

[0024] The term “cycloalkenyl” as used herein, means a non-aromatic monocyclic or multicyclic ring system containing at least one carbon-carbon double bond and preferably having from 5-10 carbon atoms per ring. Exemplary monocyclic cycloalkenyl rings include cyclopentenyl, cyclohexenyl or cycloheptenyl.

[0025] The term “fluoroalkyl” as used herein, means an alkyl group, as defined herein, in which one, two, three, four, five, six, seven or eight hydrogen atoms are replaced by fluorine. Representative examples of fluoroalkyl include, but are not limited to, 2-fluoroethyl, 2,2,2-trifluoroethyl, trifluoromethyl, difluoromethyl, pentafluoroethyl, and trifluoropropyl such as 3,3,3-trifluoropropyl.

[0026] The term “alkoxyfluoroalkyl” as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through a fluoroalkyl group, as defined herein.

[0027] The term “fluoroalkoxy” as used herein, means at least one fluoroalkyl group, as defined herein, is appended to the parent molecular moiety through an oxygen atom. Representative examples of fluoroalkoxy include, but are not limited to, difluoromethoxy, trifluoromethoxy and 2,2,2-trifluoroethoxy.

[0028] The term “halogen” or “halo” as used herein, means Cl, Br, I, or F.

[0029] The term “haloalkyl” as used herein, means an alkyl group, as defined herein, in which one, two, three, four, five, six, seven or eight hydrogen atoms are replaced by a halogen.

[0030] The term “haloalkoxy” as used herein, means at least one haloalkyl group, as defined herein, is appended to the parent molecular moiety through an oxygen atom.

[0031] The term “heteroalkyl” as used herein, means an alkyl group, as defined herein, in which one or more of the carbon atoms has been replaced by a heteroatom selected from S, Si, O, P and N. The heteroatom may be oxidized. Representative examples of heteroalkyls include, but are not limited to, alkyl ethers, secondary and tertiary alkyl amines, amides, and alkyl sulfides.

[0032] The term “heteroaryl” as used herein, refers to an aromatic monocyclic ring or an aromatic bicyclic ring system or an aromatic tricyclic ring system. The aromatic monocyclic rings are five or six membered rings containing at least one heteroatom independently selected from the group consisting of N, O and S (e.g. 1, 2, 3, or 4 heteroatoms independently selected from O, S, and N). The five membered aromatic monocyclic rings have two double bonds and the six membered aromatic monocyclic rings have three double bonds. The bicyclic heteroaryl groups are exemplified by a monocyclic heteroaryl ring appended to the parent molecular moiety and fused to a monocyclic cycloalkyl group, as defined herein, a monocyclic aryl group, as defined herein, a monocyclic heteroaryl group, as defined herein, or a monocyclic heterocycle, as defined herein. The tricyclic heteroaryl

groups are exemplified by a monocyclic heteroaryl ring appended to the parent molecular moiety and fused to two of a monocyclic cycloalkyl group, as defined herein, a monocyclic aryl group, as defined herein, a monocyclic heteroaryl group, as defined herein, or a monocyclic heterocycle, as defined herein. Representative examples of monocyclic heteroaryl include, but are not limited to, pyridinyl (including pyridin-2-yl, pyridin-3-yl, pyridin-4-yl), pyrimidinyl, pyrazinyl, thienyl, furyl, thiazolyl, thiadiazolyl, isoxazolyl, pyrazolyl, and 2-oxo-1,2-dihydropyridinyl. Representative examples of bicyclic heteroaryl include, but are not limited to, chromenyl, benzothienyl, benzodioxolyl, benzotriazolyl, quinolinyl, thienopyrrolyl, thienothienyl, imidazothiazolyl, benzothiazolyl, benzofuranyl, indolyl, quinolinyl, imidazopyridine, benzooxadiazolyl, and benzopyrazolyl. Representative examples of tricyclic heteroaryl include, but are not limited to, dibenzofuranyl and dibenzothienyl. The monocyclic, bicyclic, and tricyclic heteroaryls are connected to the parent molecular moiety through any carbon atom or any nitrogen atom contained within the rings, and can be unsubstituted or substituted.

[0033] The term “heterocycle” or “heterocyclic” as used herein, means a monocyclic heterocycle, a bicyclic heterocycle, or a tricyclic heterocycle. The monocyclic heterocycle is a three-, four-, five-, six-, seven-, or eight-membered ring containing at least one heteroatom independently selected from the group consisting of O, N, and S. The three- or four-membered ring contains zero or one double bond, and one heteroatom selected from the group consisting of O, N, and S. The five-membered ring contains zero or one double bond and one, two or three heteroatoms selected from the group consisting of O, N and S. The six-membered ring contains zero, one or two double bonds and one, two, or three heteroatoms selected from the group consisting of O, N, and S. The seven- and eight-membered rings contains zero, one, two, or three double bonds and one, two, or three heteroatoms selected from the group consisting of O, N, and S. Representative examples of monocyclic heterocycles include, but are not limited to, azetidiny, azepanyl, aziridiny, diazepanyl, 1,3-dioxanyl, 1,3-dioxolanyl, 1,3-dithiolanyl, 1,3-dithianyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dione, imidazolinyl, imidazolidinyl, isothiazolinyl, isothiazolidinyl, isoxazolinyl, isoxazolidinyl, morpholinyl, oxadiazolinyl, oxadiazolidinyl, oxazolinyl, oxazolidinyl, oxetanyl, piperazinyl, piperidinyl, pyranyl, pyrazolinyl, pyrazolidinyl, pyrrolinyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydrothienyl, thiadiazolinyl, thiadiazolidinyl, 1,2-thiazinanyl, 1,3-thiazinanyl, thiazolinyl, thiazolidinyl, thiomorpholinyl, 1,1-dioxidothiomorpholinyl (thiomorpholine sulfone), thiopyranyl, and

trithianyl. The bicyclic heterocycle is a monocyclic heterocycle fused to a phenyl group, or a monocyclic heterocycle fused to a monocyclic cycloalkyl, or a monocyclic heterocycle fused to a monocyclic cycloalkenyl, or a monocyclic heterocycle fused to a monocyclic heterocycle, or a spiro heterocycle group, or a bridged monocyclic heterocycle ring system in which two non-adjacent atoms of the ring are linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, or an alkenylene bridge of two, three, or four carbon atoms. Representative examples of bicyclic heterocycles include, but are not limited to, benzopyranyl, benzothiopyranyl, chromanyl, 2,3-dihydrobenzofuranyl, 2,3-dihydrobenzothienyl, 2,3-dihydroisoquinoline, 2-azaspiro[3.3]heptan-2-yl, azabicyclo[2.2.1]heptyl (including 2-azabicyclo[2.2.1]hept-2-yl), 2,3-dihydro-1*H*-indolyl, isoindolyl, octahydrocyclopenta[*c*]pyrrolyl, octahydropyrrolopyridinyl, and tetrahydroisoquinolyl. Tricyclic heterocycles are exemplified by a bicyclic heterocycle fused to a phenyl group, or a bicyclic heterocycle fused to a monocyclic cycloalkyl, or a bicyclic heterocycle fused to a monocyclic cycloalkenyl, or a bicyclic heterocycle fused to a monocyclic heterocycle, or a bicyclic heterocycle in which two non-adjacent atoms of the bicyclic ring are linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, or an alkenylene bridge of two, three, or four carbon atoms. Examples of tricyclic heterocycles include, but are not limited to, octahydro-2,5-epoxypentalene, hexahydro-2*H*-2,5-methanocyclopenta[*b*]furan, hexahydro-1*H*-1,4-methanocyclopenta[*c*]furan, aza-adamantane (1-azatricyclo[3.3.1.1^{3,7}]decane), and oxa-adamantane (2-oxatricyclo[3.3.1.1^{3,7}]decane). The monocyclic, bicyclic, and tricyclic heterocycles are connected to the parent molecular moiety through any carbon atom or any nitrogen atom contained within the rings, and can be unsubstituted or substituted.

[0034] The term “hydroxyl” as used herein, means an –OH group.

[0035] In some instances, the number of carbon atoms in a hydrocarbyl substituent (e.g., alkyl or cycloalkyl) is indicated by the prefix “C_x-C_y–”, wherein x is the minimum and y is the maximum number of carbon atoms in the substituent. Thus, for example, “C₁-C₃-alkyl” refers to an alkyl substituent containing from 1 to 3 carbon atoms.

[0036] The term “substituents” refers to a group “substituted” on an aryl, heteroaryl, phenyl or pyridinyl group at any atom of that group. Any atom can be substituted.

[0037] The term “substituted” refers to a group that may be further substituted with one or more non-hydrogen substituent groups. Substituent groups include, but are not limited to, halogen, =O, =S, cyano, nitro, fluoroalkyl, alkoxyfluoroalkyl, fluoroalkoxy, alkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, heteroalkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl,


heterocycle, cycloalkylalkyl, heteroarylalkyl, arylalkyl, hydroxy, hydroxyalkyl, alkoxy, alkoxyalkyl, alkylene, aryloxy, phenoxy, benzyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, -COOH, ketone, amide, carbamate, and acyl.

[0038] For compounds described herein, groups and substituents thereof may be selected in accordance with permitted valence of the atoms and the substituents, such that the selections and substitutions result in a stable compound, e.g., which does not spontaneously undergo transformation such as by rearrangement, cyclization, elimination, etc.

[0039] For the recitation of numeric ranges herein, each intervening number there between with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the number 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, and 7.0 are explicitly contemplated.

2. Compounds

[0040] Disclosed are compounds of formula (I):



(I),


or tautomers, or pharmaceutically acceptable salts thereof, wherein R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and q is 0-2; wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently substituted or unsubstituted.

[0041] In certain embodiments, R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and q is 0-2; wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently

substituted or unsubstituted with 1, 2, 3, 4, 5, 6, or 7 functional groups independently selected from the group consisting halogen, =O, =S, cyano, carbamate, nitro, fluoroalkyl, alkoxyfluoroalkyl, fluoroalkoxy, alkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, heteroalkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, heterocycloalkyl, cycloalkylalkyl, heteroarylalkyl, arylalkyl, hydroxy, hydroxyalkyl, alkoxy, allyloxy, alkoxyalkyl, alkylene, aryloxy, phenoxy, benzyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, -COOH, ketone, amide, carbamate, silyl, substituted silyloxy, t-butyltrimethylsilyloxy, alkylsulfanyl, sulfanyl, thiotriazolyl, and acyl.




[0042] In certain embodiments, R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and q is 0-2; wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently substituted or unsubstituted;

provided that if q is 0, then

R^1 is not C_1 - C_5 alkyl or , wherein R^a is NH_2 , halogen, OH,

$NHC(O)C_1$ - C_7 -alkyl, or CO_2C_1 - C_7 -alkyl;

provided that if q is 1, then

R^1 is not C_1 - C_4 alkyl, , , or , wherein R^c is H, OH,

$OC(O)C_1$ - C_7 -alkyl, $OCH_2OC(O)C_1$ - C_7 -alkyl, X is S, O, NH, NCH_3 , or NCH_2CH_3 , and Z is CH or N; and


provided that if q is 2, then

R^1 is not C_1 - C_3 alkyl; and wherein the compound of formula (I) is not 8-benzyl-2-((1-methyl-1H-imidazol-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.

[0043] In certain embodiments, R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and q is 0-2; wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently substituted or unsubstituted with 1, 2, 3, 4, 5, 6, or 7 functional groups independently selected




from the group consisting of halogen, =O, =S, cyano, carbamate, nitro, fluoroalkyl, alkoxyfluoroalkyl, fluoroalkoxy, alkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, heteroalkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, heterocycloalkyl, cycloalkylalkyl, heteroarylalkyl, arylalkyl, hydroxy, hydroxyalkyl, alkoxy, allyloxy, alkoxyalkyl, alkylene, aryloxy, phenoxy, benzyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, -COOH, ketone, amide, carbamate, silyl, substituted silyloxy, t-butyl dimethylsilyloxy, alkylsulfanyl, sulfanyl, thiotriazolyl, and acyl;

provided that if q is 0, then

R^1 is not C_1 - C_5 alkyl or , wherein R^a is NH_2 , halogen, OH,

$NHC(O)C_1$ - C_7 -alkyl, or CO_2C_1 - C_7 -alkyl;

provided that if q is 1, then

R^1 is not C_1 - C_4 alkyl, , , or , wherein R^c is H, OH,


$OC(O)C_1$ - C_7 -alkyl, $OCH_2OC(O)C_1$ - C_7 -alkyl, X is S, O, NH, NCH_3 , or NCH_2CH_3 , and Z is CH or N; and

provided that if q is 2, then

R^1 is not C_1 - C_3 alkyl; and wherein the compound of formula (I) is not 8-benzyl-2-((1-methyl-1H-imidazol-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.


[0044] In certain embodiments, R^1 is phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C_{10} - C_{12} alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl or naphthalenyl; and q is 0-2; wherein said phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C_{10} - C_{12} alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl and naphthalenyl, at each occurrence, are independently substituted or unsubstituted;

provided that if q is 0, then

R^1 is not C_1 - C_5 alkyl or , wherein R^a is NH_2 , halogen, OH,

$NHC(O)C_1$ - C_7 -alkyl, or CO_2C_1 - C_7 -alkyl;

provided that if q is 1, then


R^1 is not C_1 - C_4 alkyl, , , or , wherein R^c is H, OH, $OC(O)C_1$ - C_7 -alkyl, $OCH_2OC(O)C_1$ - C_7 -alkyl, X is S, O, NH, NCH_3 or NCH_2CH_3 , and Z is CH or N; and

provided that if q is 2, then


R^1 is not C_1 - C_3 alkyl; and wherein the compound of formula (I) is not 8-benzyl-2-((1-methyl-1H-imidazol-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.

[0045] In certain embodiments, R^1 is phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C_{10} - C_{12} alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl or naphthalenyl; and q is 0-2; wherein said phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C_{10} - C_{12} alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl and naphthalenyl, at each occurrence, are independently substituted or unsubstituted with 1, 2, 3, 4, 5, 6, or 7 functional groups independently selected from the group consisting of halogen, =O, =S, cyano, carbamate, nitro, fluoroalkyl, alkoxyfluoroalkyl, fluoroalkoxy, alkyl, alkenyl, alkynyl, haloalkyl, haloalkoxy, heteroalkyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, heterocycloalkyl, cycloalkylalkyl, heteroarylalkyl, arylalkyl, hydroxy, hydroxyalkyl, alkoxy, allyloxy, alkoxyalkyl, alkylene, aryloxy, phenoxy, benzyloxy, amino, alkylamino, acylamino, aminoalkyl, arylamino, sulfonylamino, sulfinylamino, sulfonyl, alkylsulfonyl, arylsulfonyl, aminosulfonyl, sulfinyl, -COOH, ketone, amide, carbamate, silyl, substituted silyloxy, t-butyl dimethylsilyloxy, alkylsulfanyl, sulfanyl, thiotriazolyl, and acyl;

provided that if q is 0, then

R^1 is not C_1 - C_5 alkyl or , wherein R^a is NH_2 , halogen, OH, $NHC(O)C_1$ - C_7 -alkyl, or CO_2C_1 - C_7 -alkyl;

provided that if q is 1, then

R^1 is not C_1 - C_4 alkyl, , , or , wherein R^c is H, OH, $OC(O)C_1$ - C_7 -alkyl, $OCH_2OC(O)C_1$ - C_7 -alkyl, X is S, O, NH, NCH_3 , or NCH_2CH_3 , and Z is CH or N; and

provided that if q is 2, then

R^1 is not C_1 - C_3 alkyl; and wherein the compound of formula (I) is not 8-benzyl-2-((1-methyl-1H-imidazol-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.

[0046] Representative compounds of formula (I) include, but are not limited to:

8-benzyl-2-(4-fluorobenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(2,4-difluorobenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(4-fluoro-3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(3-hydroxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(4-fluoro-3-hydroxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(3-fluoro-4-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(4-fluoro-2-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(4-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 tert-butyl (3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)carbamate;
 8-benzyl-2-((5-methoxythiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 2-(3-aminobenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(4-(tert-butyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(naphthalen-2-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 2-([1,1'-biphenyl]-4-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 2-((2H-chromen-3-yl)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-((1,3-diphenyl-1H-pyrazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-(dibenzo[b,d]furan-4-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 8-benzyl-2-cinnamyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 2-((3-acetylbenzo[d]thiazol-2(3H)-ylidene)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

tert-butyl 4-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)carbamate;

2-(4-aminobenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(4-(allyloxy)-3-methoxybenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(4-hydroxy-3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-((4-methyl-4H-1,2,4-triazol-3-yl)thio)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((5-(pyridin-2-yl)thiophen-2-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one

2-(benzo[b]thiophen-3-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-dodecyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(benzo[b]thiophen-2-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyridin-4-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-cyclohexylthiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-([2,2'-bithiophen]-5-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-isobutylthiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-(1-methyl-5-(trifluoromethyl)-1H-pyrazol-3-yl)thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((5-(trifluoromethyl)furan-2-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one;

4-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzoic acid;

8-benzyl-2-((5-(methoxymethyl)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((perfluorophenyl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3,5-bis(trifluoromethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-chloro-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-methylfuran-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzoic acid;

8-benzyl-2-((5-(morpholinomethyl)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(3-(aminomethyl)benzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

5-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)-1,3-dimethylpyrimidine-2,4(1H,3H)-dione;

8-benzyl-2-(imidazo[2,1-b]thiazol-6-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

N-(3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)methanesulfonamide;

8-benzyl-2-((3,5-dimethylisoxazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyridin-3-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

diethyl (8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)phosphonate;

8-benzyl-2-(4-methoxy-3-(methoxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(dimethylamino)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-(methoxymethyl)thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(methoxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(hydroxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-methoxypyrimidin-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((3,4-dimethylthieno[2,3-b]thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyrazin-2-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((2-(propylthio)pyrimidin-5-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((4-methyl-4H-thieno[3,2-b]pyrrol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-((1,2,3-thiadiazol-5-yl)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3,4-dimethoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(methylthio)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(benzo[d][1,3]dioxol-5-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-((dimethylamino)methyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-ethoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-methylpyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-(dimethylamino)pyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(2,4,6-trifluorobenzyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-dimethoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-chloro-4-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-fluoro-2-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-fluoro-6-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-difluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-bromopyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-chloropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-dichloropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-styryl-1,7-dihydroimidazo[1,2-a]pyrazin-3(2H)-one;

8-benzyl-2-(3-(2-methoxyethoxy)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzonitrile;

and pharmaceutically acceptable salts thereof.

[0047] Compound names are assigned by using Struct=Name naming algorithm as part of CHEMDRAW® ULTRA v. 12.0.

[0048] The compounds may exist as stereoisomers wherein asymmetric or chiral centers are present. The stereoisomers are “*R*” or “*S*” depending on the configuration of substituents around the chiral carbon atom. The terms “*R*” and “*S*” used herein are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, in Pure Appl. Chem., 1976, 45: 13-30. The disclosure contemplates various stereoisomers and mixtures thereof, and these are specifically included within the scope of this invention. Stereoisomers include enantiomers and diastereomers and mixtures of enantiomers or diastereomers. Individual stereoisomers of the compounds may be prepared synthetically from commercially available starting materials, which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by methods of resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography, and optional liberation of the optically pure product from the auxiliary as described in Furniss, Hannaford, Smith, and Tatchell, “Vogel’s Textbook of Practical Organic Chemistry”, 5th edition (1989), Longman Scientific & Technical, Essex CM20 2JE, England, or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns, or (3) fractional recrystallization methods.

[0049] It should be understood that the compounds may possess tautomeric forms, as well as geometric isomers, and that these also constitute an aspect of the invention.

[0050] The present disclosure also includes isotopically-labeled compounds, which are identical to those recited in formula (I), but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes suitable for inclusion in the compounds of the invention are hydrogen, carbon, nitrogen, oxygen, phosphorus, sulfur, fluorine, and chlorine, such as, but not limited to, ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively. Substitution with heavier isotopes such as deuterium, i.e., ^2H , can afford certain therapeutic advantages resulting from greater metabolic stability, for example increased in vivo half-life or reduced dosage requirements and, hence, may be preferred in some circumstances. The compound may incorporate positron-emitting isotopes for medical imaging and positron-emitting tomography (PET) studies for determining the distribution of receptors. Suitable positron-emitting isotopes that can be incorporated in compounds of formula (I) are ^{11}C , ^{13}N , ^{15}O , and ^{18}F . Isotopically-labeled compounds of formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples using appropriate isotopically-labeled reagent in place of non-isotopically-labeled reagent.

A. Properties of the compounds of formula (I)

[0051] The compounds of formula (I) may be substrates of luciferases to produce luminescence. The compounds may have improved water solubility, improved stability, improved cell permeability, increased biocompatibility with cells, reduced autoluminescence, and reduced toxicity.

[0052] “Luminescence” refers to the light output of a luciferase under appropriate conditions, e.g., in the presence of a suitable substrate such as a coelenterazine analogue. The light output may be measured as an instantaneous or near-instantaneous measure of light output (which is sometimes referred to as “T=0” luminescence or “flash”) at the start of the luminescence reaction, which may be initiated upon addition of the coelenterazine substrate. The luminescence reaction in various embodiments is carried out in a solution. In other embodiments, the luminescence reaction is carried out on a solid support. The solution may contain a lysate, for example from the cells in a prokaryotic or eukaryotic expression system. In other embodiments, expression occurs in a cell-free system, or the luciferase protein is secreted into an extracellular medium, such that, in the latter case, it is not necessary to

produce a lysate. In some embodiments, the reaction is started by injecting appropriate materials, e.g., coelenterazine analogue, buffer, etc., into a reaction chamber (e.g., a well of a multiwell plate such as a 96-well plate) containing the luminescent protein. In still other embodiments, the luciferase and/or coelenterazine analogues (e.g., compounds of formula (I)) are introduced into a host and measurements of luminescence are made on the host or a portion thereof, which can include a whole organism or cells, tissues, explants, or extracts thereof. The reaction chamber may be situated in a reading device which can measure the light output, e.g., using a luminometer or photomultiplier. The light output or luminescence may also be measured over time, for example in the same reaction chamber for a period of seconds, minutes, hours, etc. The light output or luminescence may be reported as the average over time, the half-life of decay of signal, the sum of the signal over a period of time, or the peak output. Luminescence may be measured in Relative Light Units (RLUs).

[0053] Compounds of formula (I) can have an RLU of greater than or equal to 1, greater than or equal to 2, greater than or equal to 3, greater than or equal to 4, greater than or equal to 5, greater than or equal to 10, greater than or equal to 20, greater than or equal to 30, greater than or equal to 40, greater than or equal to 50, or greater than or equal to 100, relative to coelenterazine or a known coelenterazine analogue.

[0054] “Biocompatibility” refers to the tolerance of a cell (e.g., prokaryotic or eukaryotic) to a coelenterazine analogue (e.g., compounds of formula (I)). Biocompatibility of a coelenterazine analogue is related to the stress it causes on the host cell.

[0055] Enhanced biocompatibility of the coelenterazine analogues (e.g., compounds of formula (I)), may be determined by measuring cell viability and/or growth rate of cells. For example, enhanced biocompatibility of the coelenterazine analogues may be determined by measuring cell viability in the absence of luciferase expression of cells exposed to the coelenterazine analogues compared to native or known coelenterazines to determine how compatible and/or toxic the coelenterazine analogues are to the cells.

[0056] In particular, enhanced biocompatibility may be determined using cell viability analysis (e.g., using the CELLTITER-GLO® Luminescent Cell Viability assay), an apoptosis assay (e.g., using the CASPASE-GLO® technology), or another method known in the art. The effect of the compounds of formula (I) on cell viability or apoptosis may be compared to the effect of native or known coelenterazine analogues on cell viability or apoptosis.


[0057] Enhanced biocompatibility may also be determined by measuring the effect of the coelenterazine analogues (e.g., compounds of formula (I)) on cell growth or gene expression.

For example, enhanced biocompatibility of the compounds of formula (I) may be determined by measuring the cell number after a period of time or by determining the expression of stress response genes in a sample of cells that are exposed to compounds of formula (I) compared to cells exposed to a native or known coelenterazine or no coelenterazine. The effect of the compounds of formula (I) on cell growth or gene expression may be compared to a native or known coelenterazine.

[0058] Basic addition salts may be prepared during the final isolation and purification of the disclosed compounds by reaction of a carboxyl group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation such as lithium, sodium, potassium, calcium, magnesium, or aluminum, or an organic primary, secondary, or tertiary amine. Quaternary amine salts can be prepared, such as those derived from methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, 1-phenamine and N,N'-dibenzylethylenediamine, ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine, and the like.

B. Synthesis of compounds of formula (I)

[0059] Current methods that produce coelenterazine analogues employ harsh reaction conditions. The limitations of these synthetic protocols have slowed the investigation of new analogues. In addition, incompatibility of the conditions of these methods with a variety of functional groups and the inability to synthesize ketoaldehyde/ketoacid precursors have led to limited structural alterations at the C-2 position of coelenterazine.



Coelenterazine

[0060] Herein, three robust and versatile methods that accomplish the preparation of compounds of formula (I) are described. These methods are useful in preparing analogues that could not be made using existing synthetic methods. The described methodologies enable


access to a variety of substituents at the C-2 position and can be performed under relatively mild conditions, utilizing a wide variety of readily available aldehydes as starting materials.

[0061] Compounds of formula (I) may be prepared by synthetic processes or by metabolic processes. Preparation of the compounds by metabolic processes includes those occurring in the human or animal body (*in vivo*) or processes occurring *in vitro*.

[0062] Compounds of formula (I), wherein the groups R^1 , and q have the meanings as set forth in the Summary of the Invention section unless otherwise noted, can be synthesized as shown in Schemes 1-3.


[0063] Abbreviations which have been used in the descriptions of the Schemes that follow are: Ac_2O for acetic anhydride; CDI for carbonyldiimidazole; MeOH for methanol; TMG for 1,1,3,3-tetramethylguanidine; and TFA for trifluoroacetic acid.

Scheme 1. Synthesis of intermediates **A-C**




[0064] As shown in Scheme 1, intermediates **A-C** can be prepared from aminopyrazine **i**. Treatment of **i** with diazocarbonyls **ii-iv**, in the presence of $Rh_2(OAc)_4$, can result in formation of aminopyrazine acetophosphonates **A-C**. Intermediates **A-C** may be stable at room temperature and provide starting materials for varied analogues.

Scheme 2. Synthesis of the compound of formula (I)



[0065] Scheme 2 illustrates the conversion of intermediates **A** and **B** to the compound of formula (I), wherein n is 1. Intermediates **A** and **B** can be treated with 1,1,3,3-tetramethylguanidine and undergo Horner-Wadsworth-Emmons olefination with aldehyde **v**, wherein R^1 is as defined in the Summary of Invention, to yield intermediate **vi**. Intermediate **vi** can be reduced to give the compound of formula (I).

Scheme 3. Synthesis of the compound of formula (I)



[0066] Scheme 3 illustrates the preparation of the compound of formula (I) by two routes from intermediate **C**. Intermediate **C** can be treated with 1,1,3,3-tetramethylguanidine and undergo Horner-Wadsworth-Emmons olefination with aldehyde **v**, wherein R^1 is as defined in the Summary of Invention, to give intermediate **vii**. Intermediate **vii** can be hydrogenated to provide intermediate **viii**, which can then be converted to the compound of formula (I) by treatment with TFA and subsequent cyclization promoted by the addition of carbonyldiimidazole. Alternatively, intermediate **vii** can be treated with TFA, resulting in hydrolysis of the t-butyl ester and formation of intermediate **ix**. Addition of acetic anhydride followed by sodium borohydride can then provide the compound of formula (I).

[0067] Optimum reaction conditions and reaction times for each individual step can vary depending on the particular reactants employed and substituents present in the reactants used. Specific procedures are provided in the Examples section. Reactions can be worked up in the

conventional manner, e.g. by eliminating the solvent from the residue and further purified according to methodologies generally known in the art such as, but not limited to, crystallization, distillation, extraction, trituration, and chromatography. Unless otherwise described, the starting materials and reagents are either commercially available or can be prepared by one skilled in the art from commercially available materials using methods described in the chemical literature. Starting materials, if not commercially available, can be prepared by procedures selected from standard organic chemical techniques, techniques that are analogous to the synthesis of known, structurally similar compounds, or techniques that are analogous to the above described schemes or the procedures described in the synthetic examples section.

[0068] Routine experimentations, including appropriate manipulation of the reaction conditions, reagents and sequence of the synthetic route, protection of any chemical functionality that cannot be compatible with the reaction conditions, and deprotection at a suitable point in the reaction sequence of the method are included in the scope of the invention. Suitable protecting groups and the methods for protecting and deprotecting different substituents using such suitable protecting groups are well known to those skilled in the art; examples of which can be found in PGM Wuts and TW Greene, in Greene's book titled *Protective Groups in Organic Synthesis* (4th ed.), John Wiley & Sons, NY (2006), which is incorporated herein by reference in its entirety. Synthesis of the compounds of the invention can be accomplished by methods analogous to those described in the synthetic schemes described hereinabove and in specific examples.

[0069] When an optically active form of a disclosed compound is required, it can be obtained by carrying out one of the procedures described herein using an optically active starting material (prepared, for example, by asymmetric induction of a suitable reaction step) or by resolution of a mixture of the stereoisomers of the compound or intermediates using a standard procedure (such as chromatographic separation, recrystallization or enzymatic resolution).

[0070] Similarly, when a pure geometric isomer of a compound is required, it can be obtained by carrying out one of the above procedures using a pure geometric isomer as a starting material or by resolution of a mixture of the geometric isomers of the compound or intermediates using a standard procedure such as chromatographic separation.

[0071] It can be appreciated that the synthetic schemes and specific examples as described are illustrative and are not to be read as limiting the scope of the invention as it is defined in

the appended claims. All alternatives, modifications, and equivalents of the synthetic methods and specific examples are included within the scope of the claims.

3. Methods of Use and Kits

[0072] The compounds and proteins of the disclosure may be used in any way that luciferase substrates, e.g., coelenterazine analogues, have been used. For example, they may be used in a bioluminogenic method which employs an analogue of coelenterazine to detect one or more molecules in a sample, e.g., an enzyme, a cofactor for an enzymatic reaction, an enzyme substrate, an enzyme inhibitor, an enzyme activator, or OH radicals, or one or more conditions, e.g., redox conditions. The sample may include an animal (e.g., a vertebrate), a plant, a fungus, physiological fluid (e.g., blood, plasma, urine, mucous secretions), a cell, a cell lysate, a cell supernatant, or a purified fraction of a cell (e.g., a subcellular fraction). The presence, amount, spectral distribution, emission kinetics, or specific activity of such a molecule may be detected or quantified. The molecule may be detected or quantified in solution, including multiphasic solutions (e.g., emulsions or suspensions), or on solid supports (e.g., particles, capillaries, or assay vessels).

[0073] In certain embodiments, the compounds of formula (I) may be used to quantify coelenterazine. In some embodiments, a coelenterazine (e.g., a native or known coelenterazine or a compound of formula (I)) can be used as a probe of a specific biochemical activity, e.g., apoptosis or drug metabolism. In some embodiments, the coelenterazine concentration is coupled to a specific enzyme activity by a “pro-coelenterazine” or “pro-substrate” that can be acted on by the specific enzyme of interest. In some embodiments, the pro-coelenterazine is a molecule that cannot support luminescence directly when combined with a luciferase, but can be converted into coelenterazine through catalytic processing by a specific enzyme of interest. In some embodiments, the approach can be used for enzymes such as those used in drug metabolism, e.g., cytochrome P450 enzymes, monoamine oxidase, and glutathione S-transferase; and apoptosis, e.g., caspases. For example, coelenterazine (e.g., a native or known coelenterazine, or a compound of formula (I)) can be modified to contain a cleavable group, such as 6'-O-methyl. In some embodiments, when incubated with a specific cytochrome P450 enzyme, the 6'-O-methyl is cleaved, and the pro-coelenterazine converted to coelenterazine, which can be detected with a luciferase. In some embodiments, the pro-coelenterazine can be combined with other components necessary to support luminescence, e.g., luminescent protein such as a luciferase, to provide a single reagent and a

homogeneous assay. For example, when the reagent is added to a sample, luminescence is generated as pro-coelenterazine is converted to coelenterazine. In various embodiments, similar assays can be developed for other enzymes, small molecules, or other cellular processes that can be linked to the generation of coelenterazines from pro-coelenterazines.


[0074] In certain embodiments, the compounds of formula (I) can be used for detecting luminescence in live cells. In some embodiments, a luciferase can be expressed in cells (as a reporter or otherwise), and the cells treated with a coelenterazine (e.g., a compound of formula (I)), which will permeate cells in culture, react with the luciferase and generate luminescence. In addition to being cell permeant, the compounds of formula (I) show comparable biocompatibility to native coelenterazine in terms of cell viability. In some embodiments, the compounds of formula (I) containing chemical modifications known to increase the stability of native coelenterazine in media can be synthesized and used for more robust, live cell luciferase-based reporter assays. In still other embodiments, a sample (including cells, tissues, animals, etc.) containing a luciferase and a compound of formula (I) may be assayed using various microscopy and imaging techniques. In still other embodiments, a secretable luciferase is expressed in cells as part of a live-cell reporter system.

[0075] In certain embodiments, the compounds of formula (I) disclosed herein may be provided as part of a kit. In some embodiments, the kit may include one or more luciferases (in the form of a polypeptide, a polynucleotide, or both) and a coelenterazine, along with suitable reagents and instructions to enable a user to perform assays such as those disclosed herein. The coelenterazine may be any of the native, known, or compounds of formula (I) disclosed herein. The kit may also include one or more buffers, such as those disclosed herein.

4. Examples

Example 1. Synthesis of Common Intermediates (A, B, C)

Scheme 4. Synthesis of Intermediates A-C



[0076] General Procedure for the synthesis of A-C: In a 5 mL round bottom flask were placed aminopyrazine (**1**, 500 mg, 1.9 mmol), diazo compound (**2-4**, 3.83 mmol, 2 eq.), $\text{Rh}_2(\text{OAc})_4$ (84.57 mg, 10 mol %) and 3 mL of chlorobenzene. The reaction mixture was heated at 100°C for 24 hours. The progress of the reaction was monitored by LCMS. After 24 hours, the reaction reached 100% conversion. The mixture was adsorbed on celite and purified on silica column using 40% EtOAc in heptane as eluent. The desired product was isolated pure as a brown solid with a 78-90% yield. The required diazophosphonoacetates were synthesized according to Wang et al. *Synlett*, **2007**, 11, 1683-1686.

[0077] Methyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-

(diethoxyphosphoryl)acetate (A): Yield 90%; ^1H NMR (300 MHz, CDCl_3) δ = 8.40 (s, 1H), 7.95-7.91 (m, 2H), 7.49 – 7.40 (m, 2H), 7.40 – 7.19 (m, 6H), 5.33-5.19 (m, 2H), 4.25 (s, 2H), 4.18 – 3.85 (m, 4H), 3.73 (s, 3H), 1.25 (t, $J=6$, 3H), 1.20 (t, $J=6$, 3H); ^{13}C NMR (75 MHz, cdcl_3) δ = 168.34, 168.32, 149.76, 149.64, 142.24, 141.72, 137.14, 136.66, 136.40, 128.85, 128.77, 128.76, 128.06, 126.96, 125.78, 63.72, 63.65, 63.63, 63.56, 53.55, 52.96, 51.61, 40.84, 16.33, 16.29, 16.25, 16.21. ESI-MS (m/z) [$\text{M}+\text{H}$] ($\text{C}_{24}\text{H}_{29}\text{N}_3\text{O}_5\text{P}$) observed 470.

[0078] Benzyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-

(diethoxyphosphoryl)acetate (B): Yield 78%; ^1H NMR (300 MHz, CDCl_3) δ = 8.35 (s, 1H), 7.94 (d, $J=7.2$, 2H), 7.50-7.42 (m, 2H), 7.40 – 7.18 (m, 1H), 5.43 – 5.06 (m, 4H), 4.25 (s, 2H), 4.17 – 3.76 (m, 4H), 1.19 (t, $J=7.1$, 3H), 1.16 (t, $J=7.0$, 3H); ^{13}C NMR (75 MHz, cdcl_3) δ = 167.70, 167.68, 149.82, 149.70, 142.27, 141.73, 137.18, 136.63, 136.39, 135.20, 128.85, 128.79, 128.77, 128.43, 128.30, 128.29, 128.07, 126.94, 125.80, 67.57, 63.72, 63.64, 63.60, 63.51, 53.96, 52.04, 40.84, 16.29, 16.24, 16.21, 16.16; ESI-MS (m/z) [$\text{M}+\text{H}$] ($\text{C}_{30}\text{H}_{32}\text{N}_3\text{O}_5\text{P}$) observed 546.


[0079] tert-Butyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-

(diethoxyphosphoryl)acetate (C): Yield 83%; ^1H NMR (300 MHz, CDCl_3) δ = 8.41 (s, 1H), 7.95 (d, $J=5.2$, 2H), 7.52 – 7.16 (m, 8H), 5.30 – 5.06 (m, 2H), 4.31 – 4.17 (m, 2H), 4.15 –

3.83 (m, 4H), 1.44 (s, 9H), 1.25 (t, $J=9$, 3H), 1.19 (t, $J=9$ 3H); ^{13}C NMR (75 MHz, cdcl_3) δ = 166.54, 166.52, 149.89, 149.79, 141.94, 141.60, 137.25, 136.58, 136.49, 128.80, 128.75, 127.98, 126.84, 125.73, 82.86, 63.46, 63.38, 63.29, 63.20, 54.43, 52.50, 40.75, 27.84, 16.34, 16.26; ESI-MS (m/z) [$\text{M}+\text{H}$] ($\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_5\text{P}$) observed 512.

Example 2. Method I for the synthesis of compounds of formula (I)

Scheme 5. Method I




[0080] The development of methodology for synthesizing compounds of formula (I) was initiated by investigating reaction conditions for the HWE olefination shown in the scheme above. A variety of bases were screened in a HWE olefination reaction between *p*-fluorobenzaldehyde and compound **A** in THF. These initial experiments revealed that when the pK_a of the base employed is less than 11 (K_2CO_3 or DIPEA), the reaction did not proceed. However, stronger nitrogenous or alkali metal bases with $\text{pK}_\text{a} \geq 12$ produced complex mixtures of decomposition products. Surprisingly, employment of 1,1,3,3-tetramethylguanidine (TMG) in THF ($\text{pK}_\text{a} \sim 13.6$) gave rise to the olefination product in low yield (see entry 2 of Table 1).

[0081] Most aldehydes, when employed in the HWE reaction with compound **A**, did not exclusively provide the unsaturated ester **x**. Instead, **x** tended to cyclize to generate **vi** (see Scheme 5). However, these compounds were generally unstable, especially in non-protic

solvents and at elevated pH. Further experiments with the solvent/base combination revealed that a wide range of protic solvents stabilize the cyclization product **vi**, with MeOH being preferred (see scheme below and Table 1), although pretreatment of **A** with TMG in methanol led to the cyclized diethylphosponate [diethyl (8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)phosphonate, compound **1**].

[0082] In a model system employing p-fluorobenzaldehyde, several nitrogenous bases led to the desired cyclized product **E** in the reaction between **A** and p-fluorobenzaldehyde. In general, there was no advantage to using bases other than TMG in MeOH (Table 1).



entry	Solvent	Base	time	SM conversion%	D (%)*	E (%)*
1	Dichloromethane	TMG	4h	95	30.9	15
2	THF	TMG	4h	100	10	37
3	Toluene	TMG	4h	100	34	16
4	Acetonitrile	TMG	4h	100	9	0
5	DMF	TMG	4h	100	5	0
6	Pyridine	TMG	4h	100	12	0
7	Dioxane	TMG	4h	79	43	31
8	Methanol	TMG	4h	100	0	94
9	Methanol	DBU	4h	100	0	79
10	Methanol	<i>t</i> -Bu-TMG	4h	100	0	97
11	Methanol	TBD	4h	100	0	77
12	Methanol	MeTBD	4h	100	0	56
13	Ethanol	TMG	4h	100	0	70
14	Ethylene glycol	TMG	4h	100	11	78
15	Butanol	TMG	4h	100	17	28
16	isopropanol	TMG	4h	100	5	24














Acronyms of bases and solvents used in the screening of the reaction conditions: TMG – 1,1,3,3-Tetramethylguanidine; DBU – 1,8-Diazabicycloundec-7-ene; *t*-BuTMG – 2-*tert*-Butyl-1,1,3,3-tetramethylguanidine; TBD – 1,5,7-Triazabicyclo[4.4.0]dec-5-ene; MeTBD – 7-Methyl-1,5,7-triazabicyclo[4.4.0]dec-5-ene;















*Yields of the products **D** and **E** were obtained through integration of the LCMS traces of the corresponding reaction mixtures. Conditions: room temperature, 10 mg of **A**, solvent (4 mL), TMG (3 eq.), 3 eq. 4-fluorobenzaldehyde















[0083] It was surmised that in the reaction between **A** and *p*-fluorobenzaldehyde, the product ratio of **D** and **E** was determined by an equilibrium established in the TMG-methanol solution, and the instability of **E** under these conditions led to a complete decay of the reaction components within 24 hours. Individually isolated **D** and **E** re-exposed to the reaction conditions provided similar ratios of **D** and **E**. Furthermore, when the reaction was carried out in *n*-butanol, similar ratios of **D** and **E** were generated, but the *n*-Bu-ester ($R^5 = n\text{-Bu}$) also formed. Therefore, for the widest scope of substrates, methanol as solvent provided reliable access to **E**. Typically, synthesis of **E** in methanol required less than 4 h and proceeded in good yield if the reaction was quenched before significant decomposition of **E** was observed.














[0084] **General procedure for Method I:** In a 20 mL vial was placed methyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-(diethoxyphosphoryl)acetate (**A**) (100mg, 0.21 mmol, 1 eq.), aldehyde (0.23 mmol, 1.1 eq.), and 12 mL of methanol. To that solution, 1,1,3,3-tetramethylguanidine (74 mg, 0.64 mmol, 3 eq.) was added. The reaction mixture was stirred at room temperature until it reached maximum conversion (2-6 hours). The progress of the reaction was monitored by LCMS. The mixture was poured into water, extracted with ethyl acetate, and dried over MgSO_4 . The drying agent was filtered off, and the solvent was concentrated under reduced pressure. The residue was subjected to flash chromatography on silica gel using dichloromethane as eluent. The corresponding dehydrocoelenterazine with the general structure **vi** was isolated as a red solid and used in the next step without further purification. Dehydrocoelenterazine **vi** was dissolved in 25 mL of dichloromethane and 10 mL of methanol and cooled to 0°C. To this solution, NaBH_4 (24.2 mg, 0.64 mmol, 3 eq.) was added, and the reaction mixture stirred at 0°C for 30 minutes. The reaction mixture was quenched with the 50 mL of 0.1 M HCl , extracted with dichloromethane, and dried over MgSO_4 . The drying agent was filtered off, the solvent was concentrated under reduced pressure, and the residue was purified on silica gel using dichloromethane/methanol as eluent. The target coelenterazine analogue was isolated pure as a yellow solid and dried on high vacuum.







[0085] The following compounds were made using common intermediate **A** and the general procedure of Method I above. Yields were calculated for the 2 step process starting from intermediate **A**.

compound	R ¹	Yield (%)	MS [M+H]
2		63	410
3		85	428
4		59	442
5		60	448
6		45	448
7		72	448
8		37	422
9		18	407
10		20	436
11		32	447
12		36	478
13		56	411
14		47	534

15		39	424
16		45	454
17		37	422
18		21	440
19		9	408
20		15	426
21		32	440
22		36	440
23		20	507
24		31	428
25		87	407
26		37	468
27		40	446
28		26	482


29		7	418
30		43	507
31		48	478
32		63	438
33		7	495
34		13	475
35		18	470
36		4	393
37		5	480
38		6	480
39		11	454
40		7	546
41		11	450
42		31	426
43		10	482

44		10	528
45		5	498
46		13	396
47		43	436
48		29	421
49		8	485
50		21	466
51		29	435
52		27	442
53		27	436
54		27	422
55		7	482
56		36	468

57		25	452
58		37	438
59		24	436
60		19	449
61		45	436
62		30	466

Example 3. Modification of Method I for the synthesis of compounds of formula (I)









Scheme 6. Modified Method I










[0086] Further investigation of Method I showed that the use of intermediate **B** was advantageous in cases where the equilibrium between the cyclized and uncyclized olefin intermediate is slow. After the olefination reaction was complete, **vi** quickly formed, and the uncyclized intermediate was not detected in the reaction mixture. However, **vi** did undergo ring opening by methoxide attack, affording **x**. To avoid formation of **x**, the reaction was halted after 100% conversion of **B** (10-20 minutes). Reduction of **vi** afforded the compound of formula (I) smoothly, though. This two-step process can be completed within 10-20 minutes. Moreover, yields of challenging substrates can be increased.

[0087] Throughout these studies it was observed that electron withdrawing substituents at the R¹ position tend to favor the formation of closed form **vi**. This was taken advantage of and modified Method I was used to prepare compounds of formula (I) containing pyridinyl groups at the R¹ position.

[0088] The following compounds were made in an analogous fashion using common intermediate **B** and the general procedure of Method I in Example 2. Yields were calculated for the 2 step process starting from intermediate **B**.

compound	R ¹	Yield (%)	MS [M+H]
63	 Mixture of regioisomers	33	441
64		13.7	393
65		27.9	429
66		35.9	411
67		29.2	411
68		16	411
69		28	427
70		25.4	471


71		72.1	423
72		34.8	423
73		42	453
74		32.2	457
75		25.9	461
76		35.1	407
77		27.5	436

Example 4. Methods II and III for the synthesis of compounds of formula (I)

[0089] Method I is a convenient and fast method with short reaction times that enables access to a variety of compounds of formula (I). However, many compounds of formula (I) containing electron deficient aromatic rings or reactive heterocyclic rings at the R¹ position could not be made by Method I. In some cases, olefination and subsequent cyclization were slow, and significant decomposition of the products was observed. On the other hand, electron rich aldehydes tended to have low conversion rates in the HWE olefination reaction which led to the formation of significant amounts of **1**, an unproductive intermediate (see Example 2).

[0090] To circumvent these challenges, two alternate routes were developed. Utilization of **C** prevented cyclization to the imidazolinone ring during the HWE reaction, providing only unsaturated ester **vii**. Subsequent treatment with TFA generated unsaturated acid **ix**, which was converted to the compound of formula (I) through a two-step cyclization and reduction sequence (Method II). In this method, **vii** was typically isolated in excellent yield. Following the hydrolysis approach, Method II avoids exposing sensitive intermediates to high pH conditions. Thus, in cases when the product **ix** is stable to TFA treatment, Method II afforded compounds of formula (I) with higher yields than Method I, although the process was lengthier.

Scheme 7. Method II




[0091] General procedure for Method II: In a 20 mL vial was placed *tert*-butyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-(diethoxyphosphoryl)acetate (**C**) (200 mg, 0.39 mmol, 1 eq.), aldehyde (0.43 mmol, 1.1 eq.), and 15 mL of methanol. To that solution, 1,1,3,3-tetramethylguanidine (135 mg, 1.2 mmol, 3 eq.) was added, and the reaction mixture was stirred at room temperature for 2 hours. The progress of the reaction was monitored by LCMS. The reaction mixture was poured into water extracted with ethyl acetate and dried over MgSO_4 . The drying agent was filtered off, the solvent was concentrated under reduced pressure, and the product with the general structure **vii** was used in the next step without further purification.

[0092] To a solution of **vii** in dichloromethane (5 mL), trifluoroacetic acid (2 mL) was added. The reaction mixture stirred at room temperature for 4 hours. The progress of the reaction was monitored by LCMS. After the reaction was complete, all volatiles were removed under reduced pressure, and the residue was dried under high vacuum. The resulting unsaturated acid was dissolved in THF (10 mL), and to this solution, acetic anhydride (398 mg, 3.9 mmol, 10 eq.), triethylamine (395 mg, 3.9 mmol, 10 eq.) and DMAP (4.8 mg, 0.039 mmol, 0.1 eq.) were added. The reaction mixture stirred at room temperature for 30 min and poured in the water, then extracted with dichloromethane, and dried over MgSO_4 . The drying agent was filtered off, and the solvent was concentrated under reduced pressure. The residue was subjected to a flash chromatography on silica gel using dichloromethane as eluent. The corresponding dehydrocoelenterazine with the general structure **vi** was isolated as red solid and used in the next step without further purification.

[0093] Dehydrocoelenterazine **vi** was dissolved in 25 mL of dichloromethane and 10 mL of methanol and cooled to 0°C. To this solution, NaBH₄ (44.26 mg, 1.2 mmol, 3 eq.) was added, and the reaction mixture stirred at 0°C for 30 minutes. The reaction mixture was quenched with 100 mL of 0.1 M HCl, extracted with dichloromethane, and dried over MgSO₄. The drying agent was filtered off, the solvent was concentrated under reduced pressure, and the residue was purified on silica gel using dichloromethane/methanol as eluent. The target coelenterazine analogue of formula (I) was isolated pure as a yellow solid and dried on high vacuum.

[0094] Method III is a second alternative that bypasses all potentially unstable synthetic intermediates. Hydrogenation of unsaturated ester **vii**, which was achieved in the presence of Wilkinson's catalyst under elevated pressure of hydrogen, yielded intermediate **viii**, which was typically easy to isolate and purify. TFA treatment of **viii**, followed by intramolecular cyclization of the corresponding acid using carbonyl diimidazole (CDI) afforded the compound of formula (I). Method III employs mild reaction conditions and avoids oxygen and moisture sensitive reaction intermediates; it is suitable both for scale-up and preparation of coelenterazine analogue containing sensitive functional groups. However, high pressure and prolonged reaction times are required.

Scheme 8. Method III




[0095] **General procedure for Method III:** In a 20 mL vial was placed *tert*-butyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-2-(diethoxyphosphoryl)acetate (**C**) (200 mg, 0.39 mmol, 1 eq.), aldehyde (0.43 mmol, 1.1 eq.), and 15 mL of methanol. To that solution, 1,1,3,3-

tetramethylguanidine (135 mg, 1.2 mmol, 3 eq.) was added, and the reaction mixture was stirred at room temperature for 1-2 hours. The progress of the reaction was monitored by LCMS. After the reaction was complete, the mixture was poured into water, extracted with ethyl acetate, and dried over MgSO₄. The drying agent was filtered off, and the solvent was concentrated under reduced pressure. The residue was purified using flash chromatography on silica gel with heptane/ethyl acetate as eluent and target compound **vii** was isolated as colorless oil.


[0096] In a parr-shaker reactor flask was placed *tert*-butyl 2-(3-benzyl-5-phenylpyrazin-2-ylamino)-acrylate derivative **vii** (0.3 mmol, 1 eq.), Rh(PPh₃)₃Cl (0.03 mmol, 0.1 eq.), and 50 mL of ethanol. The reactor was charged with H₂ (50 psi) and shaken for 20 hours at room temperature. The reaction was monitored by LCMS. Then, all volatiles were removed under reduced pressure, and the residue was subjected to flash chromatography on silica gel using heptane/ethylacetate as eluent. The reduced aminopirazine derivative **viii** was isolated pure as colorless oil.

[0097] Compound **viii** (0.25 mmol, 1 eq.) was dissolved in dichloromethane (5 mL), and to this solution, trifluoroacetic acid (2 mL) was added. The reaction mixture stirred at room temperature for 12 hours. The progress of the reaction was monitored by LCMS. After the reaction was complete, all volatiles were removed under reduced pressure, and the residue was co-evaporated 3 times with 10 mL of toluene to remove all TFA and provide compound **xi**. After drying under high vacuum, compound **xi** was dissolved in dichloromethane (15 mL), and to this solution was added carbodiimidazole (CDI) (122 mg, 0.75 mmol, 3 eq.). The reaction mixture stirred at room temperature for 20 minutes and then poured into 0.01 M HCl (50 mL), extracted with dichloromethane, and dried over MgSO₄. The drying agent was filtered off, and the solvent was concentrated under reduced pressure. The residue was subjected to a flash chromatography on silica gel using dichloromethane/methanol as eluent. The corresponding coelenterazine of formula (I) was isolated pure as yellow solid.


[0098] The following exemplary intermediates were isolated and characterized in the employment of Method III:

***tert*-Butyl (E)-2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-3-(3-cyanophenyl)acrylate**


^1H NMR (300 MHz, CDCl_3) δ = 10.03 (s, 1H), 8.31 (s, 1H), 8.15 (td, J =0.6, 1.6, 1H), 8.12 – 8.08 (m, 1H), 7.96 – 7.87 (m, 2H), 7.68 (t, J =7.7, 1H), 7.51 – 7.35 (m, 4H), 7.33 – 7.23 (m, 3H), 7.21 – 7.07 (m, 1H), 6.98 (s, 1H), 6.34 (s, 1H), 4.30 (s, 2H), 1.46 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3) δ = 189.91, 164.29, 148.16, 143.74, 142.89, 137.18, 136.85, 136.79, 136.76, 136.62, 136.13, 133.27, 133.12, 132.29, 132.23, 131.08, 130.23, 130.09, 129.20, 129.12, 128.83, 128.54, 128.49, 127.41, 126.02, 121.99, 118.39, 117.54, 113.68, 112.55, 104.99, 82.43, 41.15, 27.89; ESI-MS (m/z) [$M+H$] ($\text{C}_{31}\text{H}_{29}\text{N}_4\text{O}_2$) observed 489.

***tert*-Butyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-3-(3-cyanophenyl)propanoate**

^1H NMR (300 MHz, CDCl_3) δ = 8.40 (s, 1H), 7.98 – 7.92 (m, 2H), 7.50 – 7.43 (m, 3H), 7.38 – 7.23 (m, 7H), 7.22 – 7.18 (m, 1H), 7.11 (dt, J =1.5, 7.8, 1H), 5.02 (d, J =6.8, 1H), 4.93 – 4.79 (m, 1H), 4.15 (q, J =15.3, 3H), 3.10 (ddd, J =5.8, 13.9, 34.0, 2H), 1.38 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3) δ = 170.81, 150.01, 141.21, 141.14, 138.34, 137.35, 136.79, 136.45, 133.91, 132.94, 130.36, 128.93, 128.91, 128.77, 128.58, 127.88, 127.03, 125.66, 118.72, 112.15, 82.50, 55.09, 40.94, 37.46, 27.96; ESI-MS (m/z) [$M+H$] ($\text{C}_{31}\text{H}_{31}\text{N}_4\text{O}_2$) observed 491.

***tert*-Butyl (E)-2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-3-(1,2,3-thiadiazol-5-yl)acrylate**

^1H NMR (300 MHz, CDCl_3) δ = 9.45 (s, 1H), 8.51 (s, 1H), 8.27 (s, 1H), 8.03 – 7.96 (m, 2H), 7.52 – 7.45 (m, 4H), 7.43 – 7.31 (m, 3H), 7.30 – 7.22 (m, 1H), 6.70 (s, 1H), 4.43 (s, 2H), 1.48 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3) δ = 164.35, 159.03, 147.85, 144.44, 143.61, 136.90, 136.87, 136.42, 135.80, 133.05, 129.07, 128.85, 128.74, 128.54, 126.81, 126.09, 104.18, 82.10, 40.34, 27.87; ESI-MS (m/z) [$\text{M}+\text{H}$] ($\text{C}_{26}\text{H}_{26}\text{N}_5\text{O}_2\text{S}$) observed 472









***tert*-Butyl 2-((3-benzyl-5-phenylpyrazin-2-yl)amino)-3-(1,2,3-thiadiazol-5-yl)propanoate**

^1H NMR (300 MHz, CDCl_3) δ = 8.42 (s, 1H), 7.99-7.93 (m, 2H), 7.51 – 7.42 (m, 3H), 7.41 – 7.23 (m, 6H), 5.41 (d, $J=6.6$, 1H), 5.00 (dt, $J=5.0$, 6.5, 1H), 4.19 (dd, $J=15.2$, 58.4, 2H), 3.82 – 3.62 (m, 2H), 1.36 (s, 9H); ^{13}C NMR (75 MHz, CDCl_3) δ = 170.52, 158.41, 149.87, 141.26, 141.12, 137.38, 136.91, 136.87, 132.95, 129.02, 128.81, 128.78, 127.88, 127.01, 125.65, 82.60, 53.92, 40.98, 30.06, 27.83; ESI-MS (m/z) [$\text{M}+\text{H}$] ($\text{C}_{26}\text{H}_{28}\text{N}_5\text{O}_2\text{S}$) observed 474.

[0099] The following compounds were made by Methods II and III described above.

Yields were calculated for the 4 step process starting from intermediate C.

Compound	R ¹	Method	Yield (%)	MS [$\text{M}+\text{H}$]
78		II	78%	446

79		III	30%	417
80		II	32%	438
81		II	25%	481
82		III	17%	400
83 ^a		II	6	491
84		II	12	394
85		II	7	451
86 ^a	 trans	II	10	404

a:

compounds were made by Method II, but excluding the final reduction step with NaBH₄

[00100] Thus, having established multiple synthetic methodologies, the scope and utility of these methods were demonstrated by synthesizing compounds of formula (I) with a wide variety of substituents at the R¹ position. As shown above, aromatic groups with electron-withdrawing substituents, aromatic groups with electron-donating substituents, sterically demanding groups, and polar heterocyclic motifs were all amenable to synthesis using one of these new methods.

Example 5. Luminescent Properties

[00101] **Luminescence Assay Procedure:** Each compound to be screened was dissolved in DMSO (5mM) and then further diluted to 100uM in NANO-GLO® Luciferase Assay Buffer. Each diluted substrate was then combined in equal volumes with purified NanoLuc® Luciferase diluted into CO₂ independent media+10% FBS. Initial light output for each substrate was measured in a GloMax®-Multi+ luminometer three minutes after substrate addition and then at five minute intervals as a means to determine signal half-life.

[00102] A subset of the synthesized coelenterazine analogues (compounds of formula (I)) were evaluated for their suitability as luciferase substrates. NANOLUC® luciferase was employed for the screening because it is a small (19 kDa), stable, and particularly bright enzyme. Table 2 demonstrates that relative light unit (RLUs) and half-life data (ratios in comparison to native coelenterazine) indicated that moderately electron-donating and electron-withdrawing functionalities were well tolerated. A variety of pyridinyl groups were also tolerated with a significant number demonstrating superior performance to coelenterazine. Overall, a significant number of analogues were superior to coelenterazine in generating luminescence.

Table 2.

Compound	I	Half-life	Compound	I	Half-life
coelenterazine	1	1	66	1.625	0.98
1	--	--	67	4.583	1.14
2	40	0.16	68	12.916	0.93
3	38.75	0.23	69	7.5	0.35
4	9.166	0.72	70	5	0.19
5	3.875	0.43	71	27.083	0.14
6	13.75	0.33	72	1.208	0.88
7	4.166	0.52	73	6.666	0.93
8	23.333	0.38	74	3	1.09
9	1.041	0.93	75	0.958	0.57
10	0.142	1.02	76	1.666	1.14
11	0.3	1.04	77	1.75	0.57
12	0.192	0.57	78	8.75	0.43
13	0.583	2.64	79	13.87	0.52
14	0.042	2.15	80	2.166	1.62
15	0.583	0.88	81	0.246	1.19
16	0.0042	0.005	82	5.416	1.19
64	2.75	3.03			
65	10.833	0.62			

[00103] The compounds of formula (I) were also evaluated for their ability to be luciferase substrates in an additional assay. In this assay, the analogues were compared to a known coelenterazine analogue, 8-benzyl-2-(furan-2-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one (**C-1**), which is known to be superior to coelenterazine as a luciferase substrate. Table 3 demonstrates that a significant number of the analogues were well tolerated and performed similarly to or better than 8-benzyl-2-(furan-2-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.

Table 3

Compound	RLU (@100uM)	half-life (@100uM)	Km
C-1	1	1	1
1	0.00048	0.4	3
2	0.96	0.32	2.68
3	0.93	0.45	2.47
4	0.22	1.4	4.2
5	0.093	0.83	6.4
6	0.33	0.646	3.7
7	0.1	1	3
8	0.56	0.73	1.83
9	0.025	1.8	5.57
10	0.0034	1.96	7.4
11	0.0072	2	3.3
12	0.0046	1.1	3.4
13	0.014	5.1	4.96
14	0.001		14
15	0.014	1.7	4.3
16	0.0001	0.01	3.7
17	0.83	1.2	2.28
18	0.4	1.28	4.05
19	0.1	3.34	3.63
20	0.29 (75)	1.87 (75)	4.95
21	0.32	0.81	2.26
22	0.3 (75)	1.79 (75)	2.32
23	0.001	5	4.61
24	0.68	0.55	1.98
25	0.15	3.22	4.29
26	0.046	1.09	4.5
27	0.1	1.19	2.2
28	0.01		2.47
29	0.13		14.3
30	0.001	3.5	4.62
31	0.054	1.56	4.44
32	0.052	1.35	6.5
33	0.017	1.9	8.3
34	0.02	1.2	1.89
35	0.0005	4.92	7.5
36	0.0017	N/A	0.97
37	0.05	0.45	1.25
38	0.054	0.6	1.25
39	0.29	0.92	6.8
40	0.0042	0.71	6.2
41	0.82	0.6	2
42	0.52	1.3	4.5
43	0.052	0.56	3.2
44	0.019	1.1	8.2

45	0.0012	2.4	7.3
46			
47	0.013	2.9	4
48	0.0013	4.4	11.7
49	0.002	4.4	4.8
50	0.023	2.2	4
51	0.49	1.2	3.2
52	0.45	1	2.2
53	0.47	1.6	2.1
54	0.04	2.8	3.8
55	0.0063	1.4	4.4
56	0.0098	0.9	6.1
57	0.063	1.7	4.5
58	0.47	1.2	3.2
59	0.72	0.53	1.7
60	0.00038	4.1	6.9
61	0.49	1.7	3.4
62	0.057	3.36	3.97
63	0.28	0.5	1.6
64	0.066	5.85	1.82
65	0.26	1.2	7.1
66	0.039	1.9	2.8
67	0.11	2.2	6.7
68	0.31	1.8	11
69	0.18	0.68	3.6
70	0.12	0.37	3.1
71	0.65	0.28	1.9
72	0.029	1.7	3
73	0.16	1.8	2.7
74	0.072	2.1	3.4
75	0.023	1.1	6.5
76	0.04	2.2	6.3
77	0.042	1.1	5.8
78	0.21	0.83	3.8
79	0.33		4.06
80	0.052	3.12	7.1
81	0.0059	2.3	6.7
82	0.13	2.3	10
83	0.02		11.1
84	0.003	3.1	8
85	0.038	0.5	NA
86	0.0002	NA	NA

Example 6. Cell Permeability and Bioluminescent Activity

[00104] **Cell Culture:** HeLa and HEK293 cells were maintained in DMEM containing 0.3 mg/ml glutamine, 100 IU/ml penicillin, 100 µg/ml streptomycin, and 10% fetal calf serum at

37°C in 5% CO₂. Dulbecco's modified eagle medium (DMEM), Opti-MEM, Penicillin/Streptomycin, and Trypsin-EDTA were purchased from Life Technologies (Carlsbad). Fetal calf serum (FBS) was purchased from HyClone (GE Healthcare). Microtiter plates were purchased from Corning.

[00105] Cell based luciferase assay: HEK293 cells stably expressing NANOLUC® luciferase under the control of a CMV promotor were plated in 100 µl growth medium (DMEM supplemented with 10% FBS) into wells of white, TC-treated, 96-well plates at a density of 10000 cells per well and incubated for 24h. The growth medium was then replaced with 100 µl OptiMEM containing 12.5 µM of the indicated substrate. The luminescent signal was analyzed immediately following substrate addition using a GLOMAX® Discover multimode detection plate reader (Promega).

[00106] Cell permeability and bioluminescent activity of certain compounds (**64-77**) were determined in comparison to coelenterazine using HEK293 cells stably expressing NANOLUC® luciferase (FIG. 1). A majority of the compounds tested exhibited a luminescent signal equal or stronger than with coelenterazine, and at least six showed a substantial increase (> 5-fold) over coelenterazine. Furthermore, rank ordering compounds by signal strength relative to coelenterazine showed a strong correlation between the biochemical and cell-based luciferase assay (FIG. 2). Together, these results imply that the tested analogues were freely permeable across biological membranes and show good bioluminescent activity within living cells.

Example 7. Cell Viability

[00107] Cell viability assay: HEK293 or HeLa cells were plated in 100 µl growth medium (DMEM supplemented with 10% FBS) into wells of white, TC-treated, 96-well plates at a density of 10000 cells per well and incubated for 24h. The growth medium was then replaced with 100 µl Opti-MEM medium, which contained a serial dilution of the indicated compound. Changes in cell viability were then measured after incubation for 24h using the CELLTITER® Green cell viability assay (Promega) according to manufacturer instructions. All luminescent measurements were performed on a GLOMAX® Discover multimode plate reader (Promega).

[00108] Evaluation of compound-induced cellular toxicity in HeLa (FIG. 3 and 4) and Hek293 (FIG. 5 and 6) cells showed no fundamental difference in the substrate toxicity pattern throughout the two different cell lines. Interestingly, fluorinated compounds **65-68** generally demonstrated reduced toxicity in both cell lines when compared to coelenterazine.

In addition compound **76** also exhibited notably reduced toxicity in both cell lines, which might enable the use of higher substrate concentrations in cell based assays.


[00109] It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents.

[00110] Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, compositions, formulations, or methods of use of the invention, may be made without departing from the spirit and scope thereof.

CLAIMS

What is claimed is:

1. A compound of formula (I)



(I),


or a tautomer, or a pharmaceutically acceptable salt thereof, wherein

R^1 is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate; and

q is 0-2;


wherein said alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, and phosphonate, at each occurrence, are independently substituted or unsubstituted;

provided that if q is 0, then



R^1 is not C_1 - C_5 alkyl or $NHC(O)C_1$ - C_7 -alkyl, or CO_2C_1 - C_7 -alkyl;

provided that if q is 1, then



R^1 is not C_1 - C_4 alkyl, $OC(O)C_1$ - C_7 -alkyl, $OCH_2OC(O)C_1$ - C_7 -alkyl, X is S, O, NH, NCH_3 , or NCH_2CH_3 , and Z is CH or N; and

provided that if q is 2, then

R^1 is not C_1 - C_3 alkyl; and wherein the compound of formula (I) is not 8-benzyl-2-((1-methyl-1H-imidazol-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one.

2. The compound of claim 1, wherein

R¹ is phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C₁₀-C₁₂ alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl or naphthalenyl.

3. The compound of claim 1, selected from the group consisting of:
 - 8-benzyl-2-(4-fluorobenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(2,4-difluorobenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(4-fluoro-3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(3-hydroxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(4-fluoro-3-hydroxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(3-fluoro-4-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(4-fluoro-2-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(4-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - tert-butyl (3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)carbamate;
 - 8-benzyl-2-((5-methoxythiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 2-(3-aminobenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(4-(tert-butyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(naphthalen-2-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 2-([1,1'-biphenyl]-4-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 2-((2H-chromen-3-yl)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-((1,3-diphenyl-1H-pyrazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-(dibenzo[b,d]furan-4-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 8-benzyl-2-cinnamyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;
 - 2-((3-acetylbenzo[d]thiazol-2(3H)-ylidene)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

tert-butyl 4-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)carbamate;

2-(4-aminobenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(4-(allyloxy)-3-methoxybenzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(4-hydroxy-3-methoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-((4-methyl-4H-1,2,4-triazol-3-yl)thio)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((5-(pyridin-2-yl)thiophen-2-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one

2-(benzo[b]thiophen-3-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-dodecyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(benzo[b]thiophen-2-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyridin-4-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-cyclohexylthiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-([2,2'-bithiophen]-5-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-isobutylthiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-(1-methyl-5-(trifluoromethyl)-1H-pyrazol-3-yl)thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((5-(trifluoromethyl)furan-2-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one;

4-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzoic acid;

8-benzyl-2-((5-(methoxymethyl)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((perfluorophenyl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((1-methyl-3-(thiophen-2-yl)-1H-pyrazol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3,5-bis(trifluoromethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-chloro-1-methyl-3-(trifluoromethyl)-1H-pyrazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-methylfuran-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzoic acid;

8-benzyl-2-((5-(morpholinomethyl)furan-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(3-(aminomethyl)benzyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

5-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)-1,3-dimethylpyrimidine-2,4(1H,3H)-dione;

8-benzyl-2-(imidazo[2,1-b]thiazol-6-ylmethyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

N-(3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)phenyl)methanesulfonamide;

8-benzyl-2-((3,5-dimethylisoxazol-4-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyridin-3-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

diethyl (8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)phosphonate;

8-benzyl-2-(4-methoxy-3-(methoxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(dimethylamino)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-(methoxymethyl)thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(methoxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(hydroxymethyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-methoxypyrimidin-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((3,4-dimethylthieno[2,3-b]thiophen-2-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(pyrazin-2-ylmethyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-((2-(propylthio)pyrimidin-5-yl)methyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((4-methyl-4H-thieno[3,2-b]pyrrol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-((1,2,3-thiadiazol-5-yl)methyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3,4-dimethoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-(methylthio)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

2-(benzo[d][1,3]dioxol-5-ylmethyl)-8-benzyl-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-((dimethylamino)methyl)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-(3-ethoxybenzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-methylpyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-(dimethylamino)pyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-(2,4,6-trifluorobenzyl)imidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-dimethoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-chloro-4-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-fluoro-2-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-fluoro-6-methoxypyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((5-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-difluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-bromopyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((6-chloropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2-fluoropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-2-((2,6-dichloropyridin-3-yl)methyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

8-benzyl-6-phenyl-2-styryl-1,7-dihydroimidazo[1,2-a]pyrazin-3(2H)-one;

8-benzyl-2-(3-(2-methoxyethoxy)benzyl)-6-phenylimidazo[1,2-a]pyrazin-3(7H)-one;

and


3-((8-benzyl-3-oxo-6-phenyl-3,7-dihydroimidazo[1,2-a]pyrazin-2-yl)methyl)benzonitrile;

or a pharmaceutically acceptable salt thereof.

4. A kit comprising a compound of claim 1.
5. The kit of claim 4, further comprising a luciferase.
6. The kit of claim 4, further comprising a buffer reagent.
7. A method for detecting luminescence in a sample, the method comprising contacting a sample with a compound of claim 1;
contacting the sample with a coelenterazine-utilizing luciferase, if it is not present in the sample; and
detecting luminescence.
8. The method of claim 7, wherein the sample contains live cells.
9. The method of claim 7, wherein the sample contains a coelenterazine-utilizing luciferase.
10. A method for detecting luminescence in a transgenic animal comprising

administering a compound of claim 1 to a transgenic animal; and
 detecting luminescence;
 wherein the transgenic animal expresses a coelenterazine-utilizing luciferase.

11. A method of preparing a compound of formula (I),




(I),

wherein R¹ is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate;
 and

q is 0-2;

the method comprising:

converting compound (vi),




(vi),

to a compound of formula (I).

12. The method of claim 11, wherein the converting of compound (vi) to a compound of formula (I) comprises reduction with a hydride reducing agent.


13. The method of claim 2, wherein the hydride reducing agent is sodium borohydride.

14. The method of claim 11, further comprising:
 reacting compound (A),




with a base and $R^1-C(O)H$ in the presence of a solvent;
to form compound (vi).

15. The method of claim 14, wherein the base is 1,1,3,3-tetramethylguanidine.
16. The method of claim 14, wherein the solvent is methanol.
17. The method of claim 11, further comprising:
reacting compound (B),




with a base and $R^1-C(O)H$ in the presence of a solvent;
to form compound (vi).

18. The method of claim 17, wherein the base is 1,1,3,3-tetramethylguanidine.
19. The method of claim 17, wherein the solvent is methanol.
20. The method of claim 11, further comprising:
reacting compound (ix),




with acetic anhydride and a base;
to form compound (vi).

21. The method of claim 20, further comprising:
reacting compound (vii),




with trifluoroacetic acid to form compound (ix).

22. The method of claim 21, further comprising:
reacting compound (C),



with a base and $R^1-C(O)H$ in the presence of a solvent;
to form compound (vii).

23. The method of claim 22, wherein the base is 1,1,3,3-tetramethylguanidine.
24. The method of claim 22, wherein the solvent is methanol.
25. A method of preparing a compound of formula (I),




(I),

wherein R¹ is alkyl, alkenyl, alkynyl, aryl, bicyclic aryl, tricyclic aryl, heteroaryl, bicyclic heteroaryl, tricyclic heteroaryl, heterocycle, cycloalkyl, heteroarylcarbonyl, or phosphonate;
and

q is 0-2;

the method comprising:


converting compound (xi),



to a compound of formula (I).


26. The method of claim 25, wherein the converting of compound (xi) to a compound of formula (I) comprises reaction of compound (xi) with carbonyldiimidazole.

27. The method of claim 25, further comprising:
reacting compound (viii),




with trifluoroacetic acid to form compound (xi).

28. The method of claim 27, further comprising:
hydrogenating compound (vii),



in the presence of a metal catalyst to form compound (viii).

29. The method of claim 28, wherein the metal catalyst comprises rhodium.
30. The method of claim 28, further comprising:
reacting compound (C),



with a base and $R^1-C(O)H$ in the presence of a solvent;
to form compound (vii).

31. The method of claim 30, wherein the base is 1,1,3,3-tetramethylguanidine.
32. The method of claim 30, wherein the solvent is methanol.
33. The method of claim 11 or claim 25, wherein
 R^1 is phenyl, pyridinyl, benzodioxolyl, benzotriazolyl, thiazolyl, thiadiazolyl, thienopyrrolyl, pyrimidinyl, pyrazinyl, thienothienyl, thienyl, diethylphosphonate, isoxazolyl, imidazothiazolyl, 1,3-dimethylpyrimidine-2,4(1H,3H)-dionyl, furanyl, pyrazolyl, benzothienyl, C_{10} - C_{12} alkyl, benzothiazolyl, cinnamyl, dibenzofuranyl, chromenyl or naphthalenyl.

1/6




FIG. 1

2/6




FIG. 2

3/6




FIG. 3




FIG. 4




FIG. 5




FIG. 6

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/017761

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D487/04 A61K49/00 C12Q1/66 G01N21/76
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2012/117667 A1 (KLAUBERT DIETER H [US] ET AL) 10 May 2012 (2012-05-10) Reference example 4, [0272]-[0273]; Scheme on p.33; example 8; claims 1,8,10,11,14,17	1-33
A	US 2015/212078 A1 (ZHOU WENHUI [US] ET AL) 30 July 2015 (2015-07-30) claims 1,17,20	1-33
A	US 2014/302539 A1 (INOUE SATOSHI [JP] ET AL) 9 October 2014 (2014-10-09) claims 1,5,9,13	1-33
	----- -/-	



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

13 March 2017

Date of mailing of the international search report

23/03/2017

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Schuemacher, Anne

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/017761

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	ANTON SHAKHMIN ET AL: "Three Efficient Methods for Preparation of Coelenterazine Analogues", CHEMISTRY - A EUROPEAN JOURNAL, vol. 22, no. 30, 15 June 2016 (2016-06-15) , pages 10369-10375, XP055353558, ISSN: 0947-6539, DOI: 10.1002/chem.201601111 the whole document -----	1-33

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2017/017761

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-33

compounds of formula (I) according to claim 1, kit comprising such compounds and method for detecting luminescence using such compounds .

1.1. claims: 11-33

process for the preparation of a compound of formula (I)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/017761

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2012117667	A1	10-05-2012	
		AU 2011323418 A1	30-05-2013
		AU 2011323419 A1	30-05-2013
		AU 2016213854 A1	01-09-2016
		BR 112013010487 A2	22-11-2016
		CA 2815227 A1	10-05-2012
		CA 2817102 A1	10-05-2012
		CN 103370319 A	23-10-2013
		CN 103443121 A	11-12-2013
		CN 106148362 A	23-11-2016
		DK 2635582 T3	14-03-2016
		EP 2635582 A1	11-09-2013
		EP 2635595 A2	11-09-2013
		ES 2567410 T3	22-04-2016
		JP 5889910 B2	22-03-2016
		JP 2014500011 A	09-01-2014
		JP 2014503485 A	13-02-2014
		JP 2016144454 A	12-08-2016
		JP 2016145219 A	12-08-2016
		KR 20130084312 A	24-07-2013
		KR 20130099997 A	06-09-2013
		SG 189496 A1	31-05-2013
		SG 190104 A1	28-06-2013
		SG 10201607774R A	29-11-2016
		SG 10201608550S A	29-12-2016
		US 2012117667 A1	10-05-2012
		US 2012174242 A1	05-07-2012
		US 2014223590 A1	07-08-2014
		US 2014227759 A1	14-08-2014
		US 2015064731 A1	05-03-2015
		US 2016002703 A1	07-01-2016
		WO 2012061529 A1	10-05-2012
		WO 2012061530 A2	10-05-2012

US 2015212078	A1	30-07-2015	
		EP 3099691 A1	07-12-2016
		US 2015212078 A1	30-07-2015
		WO 2015116806 A1	06-08-2015

US 2014302539	A1	09-10-2014	
		GB 2516136 A	14-01-2015
		GB 2534288 A	20-07-2016
		JP 6040846 B2	07-12-2016
		JP 2014201565 A	27-10-2014
		US 2014302539 A1	09-10-2014
