




- (51) **International Patent Classification:**
A23L 1/227 (2006.01) C07C 233/49 (2006.01)
C07C 233/47 (2006.01)
- (21) **International Application Number:**
PCT/US2013/034363
- (22) **International Filing Date:**
28 March 2013 (28.03.2013)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/617,796 30 March 2012 (30.03.2012) US
- (71) **Applicant:** GIVAUDAN S.A. [CH/CH]; Chemin De La Parfumerie 5, CH-1214 Vernier (CH).
- (72) **Inventors:** **SHI, Feng**; 6873 Charleston Park Drive, Mason, OH 45040 (US). **RENES, Harry**; Poseidonsingel 58A, 1363TR Almere (NL). **VAN OMMEREN, Esther**; Eaglelaan 173, NL-8241 As Lelystad (NL). **VORSTER, Santi**; Eaglelaan 173, NL-8241 As Lelystad (NL). **WANG, Yili**; 3823 Blossom Court, Mason, OH 45040 (US). **DE KLERK, Adri**; Meeuwenlaan 11, NL-4921 Vk Made (NL).
- (74) **Agent:** **PARFOMAK, Andrew, N.**; NORRIS, MCLAUGHLIN & MARCUS P.A., 875 Third Ave., 8th Floor, New York, NY 10022 (US).

- (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** N-ACYL AMINO ACID DERIVATIVES AS FOOD FLAVOURING COMPOUNDS



(57) **Abstract:** A flavour composition comprising a compound of formula (I) represented by the formula (I) and edible salts thereof, wherein R₁ is an alkyl residue containing 6 to 20 carbon atoms, or an alkene residue containing from 9 to 25 carbon atoms with 1 to 6 double bonds, R₁ together with the carbonyl group to which it is attached is a residue of a carboxylic acid, R₃ is hydrogen or methyl, and R₄ is methyl, ethyl or iso-propyl.

N-ACYL AMINO ACID DERIVATIVES AS FOOD FLAVOURING COMPOUNDS

The invention is concerned with certain carboxylic acid-amino acid conjugates, flavour compositions containing said conjugates, and their use in edible compositions.

5


Many food flavour ingredients are known in the art that exert their own very pronounced flavour characteristics to a food product. Such ingredients may be of great value in niche areas for particular types of food categories, but can be incongruous, or even offensive, when employed in other food categories.

10

There is a need to provide broad spectrum flavour ingredients that act to complement or accentuate the taste and mouthfeel characteristics of the edible compositions into which they are incorporated.

15 The applicant has now found a group of compounds that may be employed flavour ingredients that complement or accentuate the taste and mouthfeel characteristics of edible compositions into which they are incorporated across a wide category of flavour applications.

20 The present invention provides in a first aspect compounds of formula (I) represented by the formula



25 and their edible salts, and their use in edible compositions wherein

R₁ is an alkyl residue containing 6 to 20 carbon atoms, or an alkene residue containing from 9 to 25 carbon atoms with 1 to 6 double bonds, R₁ together with the carbonyl group to which it is attached is a residue of a carboxylic acid, and

- 5 R₃ is hydrogen or methyl, and
R₄ is methyl, ethyl or iso-propyl.

Edible salts include those typically employed in the food and beverage industry and include chlorides, sulphates, phosphates, gluconates, sodium, citrates, carbonates, acetates and
10 lactates.

Particular compounds are those in which R₃ is hydrogen and R₄ is iso-propyl; R₃ is methyl and R₄ is methyl; and R₃ is methyl and R₄ is ethyl. The skilled person will appreciate that the amino acid residue in which R₃ is hydrogen and R₄ is iso-propyl is the residue of Leucine
15 (Leu); whereas the amino acid residue in which R₃ is methyl and R₄ is methyl is the residue of Valine (Val); and the amino acid residue in which R₃ is methyl and R₄ is ethyl is the residue of iso-Leucine (Ile).

The compounds in which R₃ is hydrogen and R₄ is iso-propyl; R₃ is methyl and R₄ is methyl;
20 and R₃ is methyl and R₄ is ethyl, as well as their use in edible compositions, are all embodiments of the present invention.

Henceforth, when referring to the amino acids or their residues, use may be made of the three letter abbreviations used above in parentheses.
25

The carboxylic acids can likewise be represented by abbreviations. Henceforth, the carboxylic acid residues may be referred to by the abbreviation C_n, wherein "n" represents the number of carbon atoms in the residue. For example, the residue of an 18 carbon acid may be abbreviated as C₁₈. Still further, if the 18 carbon acid is saturated, e.g. stearic acid. It
30 may be abbreviated as C₁₈:0 (because it contains zero double bonds), whereas an 18 carbon acid having one double bond- e.g. oleic acid - may be abbreviated as C₁₈:1. Still

further, if the C18 acid has a single double bond in the cis configuration, then it can be abbreviated as C18:1c. Similarly, if the double bond was in the trans configuration, then the abbreviation becomes C18:1t.

- 5 In a particular embodiment of the present invention the carboxylic acid residue is a residue of a fatty acid.

The fatty acid residue may be the residue of a C8 to C22 fatty acid. The fatty acid may be mammalian or non-mammalian. A mammalian fatty acid is a natural or synthetic fatty acid
10 that is identical in structure to one naturally produced in a mammal, including, but not limited to, myristic acid, palmitic acid, stearic acid, oleic acid, linoleic acid, linolenic acid, eicosatrienoic acid, arachidonic acid, eicosapentenoic acid, and docosatetraenoic acid. A non-mammalian fatty acid is a natural or synthetic fatty acid not normally produced by a mammal, including, but not limited to, pentadecanoic acid; heptadecanoic acid;
15 nonadecanoic acid; heneicosanoic acid; 9-trans-tetradecenoic acid,; 10-trans-pentadecenoic acid, ; 9-trans-hexadecenoic acid, ; 10-trans-heptadecenoic acid,; 10-trans-heptadecenoic acid, ; 7-trans-nonadecenoic acid, ; 10,13-nonadecadienoic acid, ; 11-trans-eicosenoic acid, ; and 12-transhenicosenoic acid, .

- 20 The fatty acid residues may be saturated or unsaturated. If they are unsaturated, it is preferred that they have 1, 2 or 3 double bonds, which may in cis- or trans- configuration. More particularly, the preferred fatty acid residues are C16 to C18, and may be saturated or unsaturated.

- 25 The skilled person will appreciate, however, that natural sources of these fatty acids, for example almond oil, avocado oil, castor oil, coconut oil, corn oil, cottonseed oil, olive oil, peanut oil, rice bran oil, safflower oil, sesame oil, soybean oil, sunflower oil, palm oil and canola oil, each consist of a complex mixture of fatty acids. For example, safflower oil is predominately a source of the C18:2 linoleic acid, nevertheless it may contain other fatty
30 acids, such as linolenic acid (C18:3) and palmitic acid (C16:0), amongst others. Accordingly, reference herein to a compound containing a particular fatty acid residue, for example a

residue of C18 fatty acid, may be a reference to a pure, or substantially pure C18 fatty acid residue, or it may relate to a mixture of fatty acid residues with the predominant residue being a C18 residue. Preferred fatty acid residues are C16 to C18.

5 The compounds of formula (I) can also be represented using these abbreviations. For example, the compound of formula (I) consisting of a residue of a C18 carboxylic acid and a residue of the amino acid Leucine can be represented by the abbreviation C18-Leu. For simplicity the compounds of formula (I) henceforth may be represented in this abbreviated form.

10

As is evident from the above formula (I), the amino nitrogen atom on the amino acid residue is bound to a carbonyl carbon atom of the carboxylic acid residue to form an amide linkage. Thus, the abbreviated form C18-Leu represents the compound of formula (I) in which the residue of Leucine is connected via its nitrogen atom to the carbonyl carbon atom
15 of a C18 carboxylic acid.

These compounds are particularly useful to enhance authentic fruit profiles, They may also find use in fruit flavoured milk, yoghurt and ice creams.

20 The compounds include C8-Leu, C9- Leu, C10- Leu, C12- Leu, C14- Leu, C16- Leu, C18- Leu, C20- Leu and C22- Leu.

The compounds include C8-Leu, C9- Leu, C10- Leu, C12- Leu, C14- Leu, C16- Leu, C18- Leu, C20- Leu and C22- Leu, wherein the carboxylic acid residue is saturated.

25

The compounds include C8-Leu, C9- Leu, C10- Leu, C12- Leu, C14- Leu, C16- Leu, C18- Leu, C20- Leu and C22- Leu, wherein the carboxylic acid residue is unsaturated and contains 1, 2 or 3 double bonds. The double bonds may be in cis-configuration, trans-configuration or a mixture of cis- and trans-configuration.

30

In a particular embodiment the compounds do not include C18:2-Leu.

Particular compounds bearing the Leu residue include N-palmitenoyl-L-leucine, N-palmitoyl-L-leucine, N-linolenoyl-L-leucine, N-linoleoyl-L-leucine and N-oleoyl-L-leucine.

5 The compounds include C8-Ile, C9- Ile, C10- Ile, C12- Ile, C14- Ile, C16- Ile, C18- Ile, C20- Ile and C22- Ile.

The compounds include C8-Ile, C9- Ile, C10- Ile, C12- Ile, C14- Ile, C16- Ile, C18- Ile, C20- Ile and C22- Ile, wherein the carboxylic acid residue is saturated.

10

The compounds include C8-Ile, C9- Ile, C10- Ile, C12- Ile, C14- Ile, C16- Ile, C18- Ile, C20- Ile and C22- Ile, wherein the carboxylic acid residue is unsaturated and contains 1, 2 or 3 double bonds. The double bonds may be in cis-configuration, trans-configuration or a mixture of cis- and trans-configuration.

15

A particularly preferred compound bearing the Ile residue is N-oleoyl-Ile.

In a particular embodiment C18:1-Ile and C23:1-Ile are not included in a compound of formula (I).

20

The compounds include C8-Val, C9- Val, C10- Val, C12- Val, C14- Val, C16- Val, C18- Val, C20- Val and C22- Val.

25 The compounds include C8-Val, C9- Val, C10- Val, C12- Val, C14- Val, C16- Val, C18- Val, C20- Val and C22- Val, wherein the carboxylic acid residue is saturated.

30 The compounds include C8-Val, C9- Val, C10- Val, C12- Val, C14- Val, C16- Val, C18- Val, C20- Val and C22- Val, wherein the carboxylic acid residue is unsaturated and contains 1, 2 or 3 double bonds. The double bonds may be in cis-configuration, trans-configuration or a mixture of cis- and trans-configuration.

Particularly preferred compounds bearing the Val residue include N-palmitenoyl-L-valine, N-palmitoyl-L-valine, N-linolenoyl-L-valine, N-linoleoyl-L-valine and N-oleoyl-L-valine.

5 The compounds of formula (I) can be formed by known methods using commercially available starting materials, reagents and solvents, and a detailed discussion is not warranted here. In an embodiment of the present invention, the conjugates can be formed by the reaction of an amino acid with a fatty acid halide, e.g. a chloride under basic conditions in aqueous conditions such as a water/THF solvent system. Yield and reaction
10 times may be improved by applying heat to the reaction mixture. In an alternative embodiment, a fatty acid can be reacted with an amino acid in dioxane in the presence of DCC (dicyclohexylcarbodiimide) and 1-hydroxypyrrolidine-2,5-dione.

In yet another embodiment, an amino acid alkyl ester may be reacted with a fatty acid
15 chloride under basic conditions in an aqueous-based solvent, such as a water/THF solvent system. Thereafter, the ester can be hydrolysed carefully without affecting the amide bond in basic methanol water solution

In yet another embodiment, a fatty acid and an amino acid alkyl ester can be reacted in
20 dioxane in the presence of DCC (dicyclohexylcarbodiimide) and 1-hydroxypyrrolidine-2,5-dione. The ester can be hydrolysed carefully without affecting the amide bond in dilute basic methanol water solution

In yet another embodiment, a (mixed) anhydride of a fatty acid is reacted with an amino
25 acid in dioxane.

In yet another embodiment, a fatty acid alkyl ester can be reacted with an amino acid in dioxane

In still another embodiment, an amino acid alkyl ester is reacted with a triglyceride, optionally in the presence of a co-solvent. The amino acid ester thus formed is then hydrolysed according to a method described above.

5 In yet another embodiment, an amino acid is reacted with a triglyceride, optionally in the presence of a co-solvent.

In yet another embodiment, an amino acid is reacted with a triglyceride in the presence of a lipase, esterase, peptidase, amidase or acylase, optionally in the presence of a cosolvent
10 and/or water.

In yet another embodiment a fatty acid alkyl ester is reacted with an amino acid in the presence of a lipase or an acylase, optionally in the presence of a co-solvent and/or water.

15 The compounds of formula (I) impart remarkable organoleptic properties to edible compositions to which they are added. In particular, they impart highly intense, authentic and harmonious flavour, and a roundness and fullness to edible compositions containing them.

20 More particularly, the compounds can be incorporated into an edible product to impart a remarkable mouthfeel, body and enhanced fat perception; or an enhanced umami or salt taste; or a cooling and richness. They are particularly useful in applications low in fat, salt and umami. They are also useful in fat-free formulations such as beverages and oral care applications. They also find use in dairy applications and in vanilla, cocoa and chocolate

25 This finding was all the more surprising considering that when applicant tasted the compounds in dilute aqueous solution, they were either tasteless or they exhibited a disappointing, faintly fatty taste profile. As such, they appeared to be quite unsuitable for use in flavour applications. Only their combination with flavour co-ingredients and the
30 judicious selection of their usage levels was it possible to discover the remarkable organoleptic properties of these compounds. Their effect on edible compositions is quite

unusual in that rather than exerting a characteristic flavour profile to a foodstuff or a beverage, they actually complement, lift or accentuate the essential or authentic flavour and mouth feel characteristics of the foods or beverages in which they are incorporated. Accordingly, the compounds of the present invention find utility in a broad spectrum of applications in the food and beverage industry, as well as in health and wellness.

Accordingly, the invention provides in another of its aspects, a method of conferring flavour and/or mouthfeel to, or improving taste and/or mouthfeel of an edible composition, which method comprises adding to said composition a compound of formula (I) defined herein.

The remarkable organoleptic effects are observed when the compounds of formula (I) are incorporated into an edible composition containing one or more flavour co-ingredients.

The flavour co-ingredients may be sugars, fats, salt (e.g. sodium chloride), MSG, calcium ions, phosphate ions, organic acids, proteins, purines and mixtures thereof.

In a particular embodiment, sugars are present in amounts of 0.001 % to 90 %, more particularly 0.001 % to 50 %, still more particularly 0.001 % to 20 % based on the total weight of an edible composition.

In a particular embodiment, fats are present in amounts of 0.001 % to 100 %, more particularly 0.001 % to 80 %, more particularly 0.001 % to 30 %, still more particularly 0.001 % to 5 % based on the total weight of an edible composition.

In a particular embodiment, salt (e.g. sodium chloride) is present in amounts of 0.001 % to 20 %, more particularly 0.001 % to 5 % based on the total weight of an edible composition.

In a particular embodiment, MSG is present in amounts of 0.001 % to 2 % based on the total weight of an edible composition.

In a particular embodiment, calcium is present in amounts of 0.001% to 50% more particularly 0.001 % to 20 %, still more particularly 0.001 % to 1 % based on the total weight of an edible composition.

5 In a particular embodiment, organic acids are present in amounts of 0.001 % to 10 %, more particularly 0.001 % to 7 % based on the total weight of an edible composition.

Types of organic acids include citric, malic, tartaric, fumaric, lactic, acetic, succinic. Types of edible compositions containing organic acids include beverages, such as carbonated soft
10 drink beverages, still beverages, Juices, powdered soft drinks, liquid concentrates, alcoholic beverages and functional beverages.

In a particular embodiment, phosphorus is present in an amount up to 0.5 % by weight of an edible composition. Typically phosphorus is present as a phosphate or phosphoric acid.
15

In a particular embodiment, purines are present in an amount up to 0.5 % by weight of an edible composition. The term "purines" include ribonucleotides such as IMP and GMP.

Despite their interesting organoleptic properties, nevertheless, applicant found that
20 formulating the compounds of formula (I) was not a trivial matter. The discovered potency of the compounds suggested that they could be employed at very low levels in flavour applications, and so for ease of handling, mixing and processing with other ingredients, although it is possible to use the compounds in neat form, it is desirable to extend or add
25 volume to the physical form of the compounds by incorporating them into a suitable vehicle, for example a diluent, such as a solvent. However, the compounds are solids or viscous oils at ambient temperatures, and have very limited solubility in water. Applicant found that an at least about 0.01 % stock solution, more particularly about 0.01 – 1 % stock solution of a compound of formula (I) achieved a balance regarding acceptable solvent
30 levels for ease of handling and mixing, and the desire to limit the amount of solvent that would have to be removed from the stock solution when further processing of the compounds in flavour compositions and edible products for reasons of palatability,

efficiency, cost and the like. Applicant found that suitable solvents for the stock solution include ethanol, triacetine, glycerol and miglyol.

In order to aid in the process of solubilization and produce a stock solution and minimize
5 the amount of solvent, it is preferred to use compounds of the formula (I) formed from a mixture of carboxylic acids, rather than a pure carboxylic acid.

Accordingly, the invention provides in another of its aspects an at least about 0.01 % stock solution, more particularly about 0.01 – 1 % stock solution of a compound of formula (I).

10

The stock solution may contain other materials such as carrier materials and/or adjuvants more fully described below. In a particular embodiment, the stock solution contains an anti-oxidant selected from the group consisting of vitamin C, vitamin E, rosemary extract, antrancine, butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT). Anti-oxidants are preferably employed to prevent, or significantly reduce, generation of volatile
15 off notes as a result of degradation of the compounds of formula (I). Anti-oxidants are particularly preferred when the compounds of formula (I) bear a residue of an unsaturated fatty acid. Anti-oxidants are particularly preferred if the fatty acid residue contains more than 1 double bond. Determination of an effective amount of anti-oxidant is within the
20 purview of the skilled person, however amounts in the range of about 10 ppm to 1000 ppm based on the weight of the stock solution may be present.

In preparing the flavour compositions of the present invention, the compounds of formula (I) may be employed in any physical form. They may be used in neat form, in the form of a
25 stock solution described above; they may be used in the form of an emulsion; or they may be used in a powder form. If the compounds of formula (I) are presented in the form of a powder, the powder form can be produced by a dispersive evaporation process, such as a spray drying process as is more fully described below. The powder form may be prepared by subjecting a liquid formulation containing a compound of formula (I) to a dispersive
30 evaporation process. The liquid formulation may comprise a solution, suspension or emulsion comprising the compound of formula (I). In particular, the liquid formulation may

take the form of the stock solution described hereinabove. The liquid formulation may contain other ingredients such as a carrier material and/or an adjuvant as described more fully below.

5 A powder comprising a compound of formula (I) forms another aspect of the present invention.

The compounds of formula (I) may be incorporated into an edible composition alone, or in the form of a flavour composition comprising one or more flavour co-ingredients.

10

A flavour composition comprising a compound according to the formula (I) forms another aspect of the present invention.

15 In an embodiment of the present invention, the flavour formulation comprises a compound of formula (I) and at least one flavour co-ingredient.

In a particular embodiment of the present invention the flavour composition comprises:

- 20
- i) a compound according to formula (I);
 - ii) at least one flavour co-ingredient;
 - iii) optionally a carrier material; and
 - iv) optionally at least one adjuvant.

25 By the term "flavour co-ingredient" is an ingredient that is able to contribute or impart or modify in a positive or pleasant way the taste of an edible composition.

25

All manner of flavour co-ingredients may be employed in a composition according to the present invention, including, but not limited to natural flavours, artificial flavours, spices, seasonings, and the like. Flavour co-ingredients include synthetic flavour oils and flavouring aromatics and/or oils, oleoresins, essences, distillates, and extracts derived from plants, leaves, flowers, fruits, and so forth, and combinations comprising at least one of the foregoing.

30

Flavour oils include spearmint oil, cinnamon oil, oil of wintergreen (methyl salicylate), peppermint oil, Japanese mint oil, clove oil, bay oil, anise oil, eucalyptus oil, thyme oil, cedar leaf oil, oil of nutmeg, allspice, oil of sage, mace, oil of bitter almonds, and cassia oil; useful
5 flavouring agents include artificial, natural and synthetic fruit flavours such as vanilla, and citrus oils including lemon, orange, lime, grapefruit, yuzu, sudachi, and fruit essences including apple, pear, peach, grape, blueberry, strawberry, raspberry, cherry, plum, prune, raisin, cola, guarana, neroli, pineapple, apricot, banana, melon, apricot, ume, cherry, raspberry, blackberry, tropical fruit, mango, mangosteen, pomegranate, papaya and the like.

10

Additional exemplary flavours imparted by a flavouring agent include a milk flavour, a butter flavour, a cheese flavour, a cream flavour, and a yogurt flavour; a vanilla flavour; tea or coffee flavours, such as a green tea flavour, an oolong tea flavour, a tea flavour, a cocoa flavour, a chocolate flavour, and a coffee flavour; mint flavours, such as a peppermint
15 flavour, a spearmint flavour, and a Japanese mint flavour; spicy flavours, such as an asafetida flavour, an ajowan flavour, an anise flavour, an angelica flavour, a fennel flavour, an allspice flavour, a cinnamon flavour, a chamomile flavour, a mustard flavour, a cardamom flavour, a caraway flavour, a cumin flavour, a clove flavour, a pepper flavour, a coriander flavour, a saffron flavour, a savoury flavour, a Zanthoxyli Fructus flavour, a
20 perilla flavour, a juniper berry flavour, a ginger flavour, a star anise flavour, a horseradish flavour, a thyme flavour, a tarragon flavour, a dill flavour, a capsicum flavour, a nutmeg flavour, a basil flavour, a marjoram flavour, a rosemary flavour, a bayleaf flavour, and a wasabi (Japanese horseradish) flavour; a nut flavour such as an almond flavour, a hazelnut flavour, a macadamia nut flavour, a peanut flavour, a pecan flavour, a pistachio flavour, and
25 a walnut flavour; alcoholic flavours, such as a wine flavour, a whisky flavour, a brandy flavour, a rum flavour, a gin flavour, and a liqueur flavour; floral flavours; and vegetable flavours, such as an onion flavour, a garlic flavour, a cabbage flavour, a carrot flavour, a celery flavour, mushroom flavour, and a tomato flavour.

30 In some embodiments, said flavour co-ingredients include aldehydes and esters such as cinnamyl acetate, cinnamaldehyde, citral diethylacetal, dihydrocarvyl acetate, eugenyl 49

formate, p-methylamisol, and so forth can be used. Further examples of aldehyde flavourings include acetaldehyde (apple), benzaldehyde (cherry, almond), anisic aldehyde (licorice, anise), cinnamic aldehyde (cinnamon), citral, i.e., alpha-citral (lemon, lime), neral, i.e., beta-citral (lemon, lime), decanal (orange, lemon), ethyl vanillin (vanilla, cream),
5 heliotrope, i.e., piperonal (vanilla, cream), vanillin (vanilla, cream), alpha-amyl cinnamaldehyde (spicy fruity flavours), butyraldehyde (butter, cheese), valeraldehyde (butter, cheese), citronellal (modifies, many types), decanal (citrus fruits), aldehyde C-8 (citrus fruits), aldehyde C-9 (citrus fruits), aldehyde C-12 (citrus fruits), 2-ethyl butyraldehyde (berry fruits), hexenal, i.e., trans-2 (berry fruits), tolyl aldehyde (cherry,
10 almond), veratraldehyde (vanilla), 2,6-dimethyl-5-heptenal, i.e., melonal (melon), 2,6-dimethyloctanal (green fruit), and 2-dodecenal (citrus, mandarin), and the like.

Further examples of other flavour co-ingredients can be found in "Chemicals Used in Food Processing", publication 1274, pages 63-258, by the National Academy of Sciences.

15

Flavour co-ingredients can also include salt tastants, umami tastants, and savoury flavour compounds. Non limiting examples include: NaCl, KCl, MSG, guanosine monophosphate (GMP), inosin monophosphate (IMP), ribonucleotides such as disodium inosinate, disodium guanylate, N-(2-hydroxyethyl)-lactamide, N-lactoyl -GMP, N-lactoyl tyramine, gamma amino
20 butyric acid, allyl cysteine, 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridine-2-yl)propan-1-one, arginine, potassium chloride, ammonium chloride, succinic acid, N-(2-methoxy-4-methyl benzyl)-N'-(2-(pyridin-2-yl)ethyl) oxalamide, N-(heptan-4-yl)benzo(D)(1,3)dioxole-5-carboxamide, N-(2,4-dimethoxybenzyl)-N'-(2-(pyridin-2-yl)ethyl) oxalamide, N-(2-methoxy-4-methyl benzyl)-N'-2-(5-methyl pyridin-2-yl)ethyl) oxalamide, cyclopropyl-
25 E,Z-2,6-nonadienamide.

In particular embodiments of the present invention, the flavour co-ingredient is selected from the compounds and compositions disclosed in WO2005102701, WO2006009425, WO2005096843, WO2006046853 and WO2005096844, all of which references are herein
30 incorporated by reference in their entirety.

Flavour co-ingredients may include known salt tastants, umami tastants, and savoury flavour compounds. Non limiting examples include: NaCl, KCl, MSG, guanosine monophosphate (GMP), inosin monophosphate (IMP), ribonucleotides such as disodium inosinate, disodium guanylate, N-(2-hydroxyethyl)-lactamide, N-lactoyl -GMP, N-lactoyl tyramine, gamma amino butyric acid, allyl cysteine, 1-(2-hydroxy-4-methoxyphenyl)-3-(pyridine-2-yl)propan-1-one, arginine, potassium chloride, ammonium chloride, succinic acid, N-(2-methoxy-4-methyl benzyl)-N'-(2-(pyridin-2-yl)ethyl) oxalamide, N-(heptan-4-yl)benzo(D)(1,3)dioxole-5-carboxamide, N-(2,4-dimethoxybenzyl)-N'-(2-(pyridin-2-yl)ethyl) oxalamide, N-(2-methoxy-4-methyl benzyl)-N'-2-(2-(5-methyl pyridin-2-yl)ethyl) oxalamide, cyclopropyl-E,Z-2,6-nonadienamide.

The carrier material may be employed in compositions according to the invention to encapsulate or to entrap in a matrix the other components of the composition. The role of the carrier material may be merely that of a processing aid or a bulking agent, or it might be employed to shield or protect the other components from the effects of moisture or oxygen or any other aggressive media. The carrier material might also act as a means of controlling the release of flavour from edible compositions.

Carrier materials may include mono, di- or trisaccharides, natural or modified starches, hydrocolloids, cellulose derivatives, polyvinyl acetates, polyvinylalcohols, proteins or pectins. Example of particular carrier materials include sucrose, glucose, lactose, levulose, fructose, maltose, ribose, dextrose, isomalt, sorbitol, mannitol, xylitol, lactitol, maltitol, pentitol, arabinose, pentose, xylose, galactose, maltodextrin, dextrin, chemically modified starch, hydrogenated starch hydrolysate, succinylated or hydrolysed starch, agar, carrageenan, gum arabic, gum accacia, tragacanth, alginates, methyl cellulose, carboxymethyl cellulose, hydroxyethyl cellulose, hydroxypropylmethyl cellulose, derivatives and mixtures thereof. Of course, the skilled addressee will appreciate that the cited materials are hereby given by way of example and are not to be interpreted as limiting the invention.

30

By "flavour adjuvant" is meant an ingredient capable of imparting additional added benefit to compositions of the present invention such as a colour, light resistance, chemical stability and the like. Suitable adjuvants include solvents (including water, alcohol, ethanol, triacetone, oils, fats, vegetable oil and miglyol), binders, diluents, disintegrating agents, 5 lubricants, colouring agents, preservatives, antioxidants, emulsifiers, stabilisers, anti-caking agents, and the like. In a particular embodiment, the flavour composition comprises an anti-oxidant. Said anti-oxidants may include vitamin C, vitamin E, rosemary extract, antrancine, butylated hydroxyanisole (BHA) and butylated hydroxytoluene (BHT).

10 Examples of such carriers or adjuvants for flavour compositions may be found in for example, "Perfume and Flavour Materials of Natural Origin", S. Arctander, Ed., Elizabeth, N.J., 1960; in "Perfume and Flavour Chemicals", S. Arctander, Ed., Vol. I & II, Allured Publishing Corporation, Carol Stream, USA, 1994; in "Flavourings", E. Ziegler and H. Ziegler (ed.), Wiley-VCH Weinheim, 1998, and "CTFA Cosmetic Ingredient Handbook", J.M. Nikitakis 15 (ed.), 1st ed., The Cosmetic, Toiletry and Fragrance Association, Inc., Washington, 1988.

Other suitable and desirable ingredients of flavour compositions are described in standard texts, such as "Handbook of Industrial Chemical Additives", ed. M. and I. Ash, 2nd Ed., (Synapse 2000).

20 Flavour compositions according to the present invention may be provided in any suitable physical form. For example, they may be in the form of oils, emulsions or dispersions in a hydrous liquid or organic liquid suitable for use in edible compositions, or solid form, such as powders.

25 If the flavour compositions are to be provided in the form of a powdered composition, they may be prepared by dispersive evaporation techniques generally known in the art, such as spray drying.

30 Accordingly, in another aspect of the present invention there is provided a method of forming a powder composition, comprising the steps of providing a liquid composition

comprising a compound of the formula (I) and one or more optional ingredients selected from at least one flavour co-ingredient, a carrier material and an adjuvant, and dispersively evaporating said liquid composition to form a powder composition.

- 5 In this manner, a compound of formula (I) or a flavour composition comprising said compound may be presented in a powder form.

The liquid composition used in the preparation of a powder may be in the form of a solution, emulsion, dispersion or slurry. The liquid may contain water, and/or an organic
10 liquid, such as ethanol, glycerol, triacetine, miglyol (MCT) that is acceptable for use in edible compositions.

Powder compositions according to the present invention may be prepared according to methods and apparatus known in the art for producing powders on an industrial scale. A
15 particularly suitable method is spray drying. Spray drying techniques and apparatus are well known in the art and need no detailed discussion herein. The spray drying techniques, apparatus and methods described in US2005/0031769 and US2013/0022728, as well as those techniques, apparatus and methods described in those documents are suitable for producing powder compositions of the present invention and are herein incorporated by
20 reference in their entirety.

The manner in which the compounds of formula (I) are incorporated into powder flavour compositions of the invention is not critical. For example, the compounds may be added to the liquid composition described above and be subjected to a dispersive evaporation
25 process along with all the other components of the flavour composition. Alternatively, compounds may be added to the flavour composition after it has been formed as a powder.

Many of the flavour co-ingredients described herein above are volatile and/or may be sensitive to oxidative degradation, particularly when subjected to elevated temperature,
30 and under humid conditions. Accordingly, particular problems can arise when subjecting flavour co-ingredients described above to dispersive evaporation processes such as spray

drying. A non-exhaustive list of ingredients that can be particularly susceptible include, those ingredients containing artificial, natural or synthetic fruit flavours such as vanilla, chocolate, coffee, cocoa and citrus oil, including lemon, orange, grape, lime and grapefruit, and fruit essences including apple, pear, peach, strawberry, raspberry, cherry, plum,
5 pineapple, apricot and the like. The volatile components of these flavour co-ingredients may include, but are not limited to, acetaldehyde, dimethyl sulfide, ethyl acetate, ethyl propionate, methyl butyrate, and ethyl butyrate. Flavour co-ingredients containing volatile aldehydes or esters include, e.g., cinnamyl acetate, cinnamaldehyde, citral, diethylacetal, dihydrocarvyl acetate, eugenyl formate, and p-methylanisole. Further examples of volatile
10 compounds that may be present as co-ingredients include acetaldehyde (apple); benzaldehyde (cherry, almond); cinnamic aldehyde (cinnamon); citral, i.e., alpha citral (lemon, lime); neral, i.e., beta citral (lemon, lime); decanal (orange, lemon); ethyl vanillin (vanilla, cream); heliotropine, i.e., piperonal (vanilla, cream); vanillin (vanilla, cream); alpha-amyl cinnamaldehyde (spicy fruity flavors); butyraldehyde (butter, cheese);
15 valeraldehyde (butter, cheese); citronellal (modifies, many types); decanal (citrus fruits); aldehyde C-8 (citrus fruits); aldehyde C-9 (citrus fruits); aldehyde C-12 (citrus fruits); 2-ethyl butyraldehyde (berry fruits); hexenal, i.e., trans-2 (berry fruits); tolyl aldehyde (cherry, almond); veratraldehyde (vanilla); 2,6-dimethyl-5-heptenal, i.e., melonal (melon); 2-6-dimethyloctanal (green fruit); and 2-dodecenal (citrus, mandarin); cherry; or grape and
20 mixtures thereof.

Applicant surprisingly found that the inclusion of a compound of formula (I) in a powder flavour composition, it was possible to maintain the flavour quality of flavour composition converted into powdered form by a dispersive evaporation process.

25 Accordingly, the invention provides in another of its aspects a method of maintaining flavour quality of a powder flavour composition comprising the step of including in said powder flavour composition a compound of formula (I)

30 As stated hereinabove, compounds of formula (I) or flavour compositions containing such compounds can be incorporated into edible compositions, and an edible composition

containing such a compound or flavour composition forms another aspect of the present invention.

The term "edible composition" refers to products for consumption by a subject, typically via
5 the oral cavity (although consumption may occur via non-oral means such as inhalation),
for at least one of the purposes of enjoyment, nourishment, or health and wellness benefits.
Edible compositions may be present in any form including, but not limited to, liquids, solids,
semi-solids, tablets, capsules, lozenges, strips, powders, gels, gums, pastes, slurries, syrups,
aerosols and sprays. The term also refers to, for example, dietary and nutritional
10 supplements. Edible compositions include compositions that are placed within the oral
cavity for a period of time before being discarded but not swallowed. It may be placed in the
mouth before being consumed, or it may be held in the mouth for a period of time before
being discarded. An edible composition as herein above defined includes compositions
whose taste is modified in the manner described herein by the addition of compounds of
15 formula (I) or whose taste is so modified by processing such that it is enriched in a
compound of formula (I).

In a particular embodiment the term "edible compositions" refers to products for
consumption by a subject, typically via the oral cavity (although consumption may occur via
20 non-oral means such as inhalation), for one of the purposes of enjoyment or nourishment.

In a more particular embodiment the term "edible compositions" refers to products for
consumption by a subject, typically via the oral cavity (although consumption may occur via
non-oral means such as inhalation), for the purpose of enjoyment. Still more particularly,
25 the term refers to foodstuffs and beverages.

In a particular embodiment, the term "edible composition" does not relate to
pharmaceutical compositions.

30 In another embodiment, the term "edible composition" does not relate to nutritional
supplements.

Broadly, the edible composition includes, but is not limited to foodstuffs of all kinds, confectionery products, baked products, sweet products, savoury products, fermented products, dairy products, beverages and oral care products.

5

Exemplary foodstuffs include, but are not limited to, chilled snacks, sweet and savoury snacks, fruit snacks, chips/crisps, extruded snacks, tortilla/corn chips, popcorn, pretzels, nuts, other sweet and savoury snacks, snack bars, granola bars, breakfast bars, energy bars, fruit bars, other snack bars, meal replacement products, slimming products, convalescence
10 drinks, ready meals, canned ready meals, frozen ready meals, dried ready meals, chilled ready meals, dinner mixes, frozen pizza, chilled pizza, soup, canned soup, dehydrated soup, instant soup, chilled soup, uht soup, frozen soup, pasta, canned pasta, dried pasta, chilled/fresh pasta, noodles, plain noodles, instant noodles, cups/bowl instant noodles, pouch instant noodles, chilled noodles, snack noodles, dried food, dessert mixes, sauces,
15 dressings and condiments, herbs and spices, spreads, jams and preserves, honey, chocolate spreads, nut-based spreads, and yeast-based spreads.

Exemplary confectionery products include, but are not limited to, chewing gum (which includes sugarized gum, sugar-free gum, functional gum and bubble gum), centerfill
20 confections, chocolate and other chocolate confectionery, medicated confectionery, lozenges, tablets, pastilles, mints, standard mints, power mints, chewy candies, hard candies, boiled candies, breath and other oral care films or strips, candy canes, lollipops, gummies, jellies, fudge, caramel, hard and soft panned goods, toffee, taffy, liquorice, gelatin
25 candies, gum drops, jelly beans, nougats, fondants, combinations of one or more of the above, and edible flavour compositions incorporating one or more of the above.

Exemplary baked products include, but are not limited to, alfajores, bread, packaged/industrial bread, unpackaged/artisanal bread, pastries, cakes, packaged/industrial cakes, unpackaged/artisanal cakes, cookies, chocolate coated biscuits,
30 sandwich biscuits, filled biscuits, savoury biscuits and crackers, bread substitutes,

Exemplary sweet products include, but are not limited to, breakfast cereals, ready-to-eat ("rte") cereals, family breakfast cereals, flakes, muesli, other ready to eat cereals, children's breakfast cereals, hot cereals,

- 5 Exemplary savoury products include, but are not limited to, salty snacks (potato chips, crisps, nuts, tortilla-tostada, pretzels, cheese snacks, corn snacks, potato-snacks, ready-to-eat popcorn, microwaveable popcorn, pork rinds, nuts, crackers, cracker snacks, breakfast cereals, meats, aspic, cured meats (ham, bacon), luncheon/breakfast meats (hotdogs, cold cuts, sausage), tomato products, margarine, peanut butter, soup (clear, canned, cream,
- 10 instant, UHT),canned vegetables, pasta sauces.

- Exemplary dairy products include, but are not limited to, cheese, cheese sauces, cheese-based products, ice cream, impulse ice cream, single portion dairy ice cream, single portion water ice cream, multi-pack dairy ice cream, multi-pack water ice cream, take-home ice
- 15 cream, take-home dairy ice cream, ice cream desserts, bulk ice cream, take-home water ice cream, frozen yoghurt, artisanal ice cream, dairy products, milk, fresh/pasteurized milk, full fat fresh/pasteurized milk, semi skimmed fresh/pasteurized milk, long-life/uht milk, full fat long life/uht milk, semi skimmed long life/uht milk, fat-free long life/uht milk, goat milk, condensed/evaporated milk, plain condensed/evaporated milk, flavoured, functional and
- 20 other condensed milk, flavoured milk drinks, dairy only flavoured milk drinks, flavoured milk drinks with fruit juice, soy milk, sour milk drinks, fermented dairy drinks, coffee whiteners, powder milk, flavoured powder milk drinks, cream, yoghurt, plain/natural yoghurt, flavoured yoghurt, fruited yoghurt, probiotic yoghurt, drinking yoghurt, regular drinking yoghurt, probiotic drinking yoghurt, chilled and shelf-stable desserts, dairy-based
- 25 desserts, soy-based desserts.

- Exemplary beverages include, but are not limited to, flavoured water, soft drinks, fruit drinks, coffee-based drinks, tea-based drinks, juice-based drinks (includes fruit and vegetable), milk-based drinks, gel drinks, carbonated or non-carbonated drinks, powdered
- 30 drinks, alcoholic or non-alcoholic drinks.

Exemplary fermented foods include, but are not limited to, Cheese and cheese products, meat and meat products, soy and soy products, fish and fish products, grain and grain products, fruit and fruit products.

- 5 In a particular embodiment the consumable product is selected from the group consisting of soy sauce, cheese, soup, hot and cold sauces, fruits, vegetables, ketchups, tea, coffee, snacks such as potato chips or extruded snacks.

10 The compounds of formula (I), when added to a flavour composition and/or an edible composition act on the composition to complement its flavour and/or mouthfeel to render it more delicious and authentic. The effects may be temporal or related to intensity, for example the compounds may act by enhancing, strengthening, softening, sharpening a flavour, or making more salivating. The compounds of formula (I) may also affect the temporal profile of a flavour, that is, they may affect the initial impact of a flavour, the body
15 of a flavour, or its lingering effect.

The compounds of formula (I) may modify any aspect of the temporal profile of taste or flavour of an edible composition. In particular, the compounds improve mouthfeel and impart more creamy and fatty sensations.

20 Compounds of formula (I) or flavour compositions containing same may be added to edible compositions in widely carrying amounts. The amount will depend on the nature of the edible composition to be flavoured, and on the desired effect, as well as on the nature of the ingredients present in said flavour composition. In order to obtain the remarkable
25 beneficial effects attributed to the presence of the compounds of formula (I), the flavour composition should be employed in amounts such that the compounds of formula (I) are present in amounts of 1 part per billion to 10 parts per million based on the total weight of the edible composition. Whereas amounts higher than this can be employed, the beneficial effects are considerably less apparent and undesirable off-notes can become increasingly
30 apparent.

Interesting organoleptic effects, e.g. salt, alcohol or coolant boosting effects, in edible compositions containing salt or alcohol or coolant compounds can be achieved when compounds of the formula (I) are employed at levels of 1 to 100 ppb.

- 5 Interesting organoleptic effects, for example umami boosting effects, in edible compositions containing umami tastants can be achieved when compounds of the formula (I) are employed at levels of 100 to 250 ppb.

- 10 Interesting organoleptic effects, in particular mouthfeel boosting effects, in edible compositions can be achieved when compounds of the formula (I) are employed at levels of 250 to 500 ppb.

- 15 Interesting organoleptic effects, e.g. fat boosting effects, in edible compositions containing fats can be achieved when compounds of the formula (I) are employed at levels of 500 to 1000 ppb.

- 20 It is particularly advantageous to incorporate compounds of formula (I) into edible compositions that are formed under conditions of high temperature, such as baking, frying or which are processed by heat treatments such as pasteurization or under UHT conditions. Under high preparation or processing temperatures, volatile flavour ingredients may be lost or degraded with the result that flavour intensity can be reduced and the essential and authentic flavour characteristics can be diminished. Such edible products include dairy products, snack foods, baked products, powdered soft drinks and similar dry mixes, and the like, fats and condiments, mayonnaise, dressings, soups and bouillons, and beverages.

- 30 A particularly preferred class of edible composition according to the present invention are powdered soft drinks and similar dry mix applications. Dry mix applications are known in the art and included products in powder form that are intended to be reconstituted before consumption. They include powdered soups, powdered cake mixes, powdered chocolate drinks, instant coffees, seasonings and fonds, and the like.

Dry powders formed by dispersive evaporation processes, such as spray drying, represent a very convenient vehicle to deliver flavour oil quality flavours to edible compositions.

5 Unfortunately, flavour oils, and in particular citrus flavour oils can be particularly sensitive to dispersive evaporation processes, especially processes carried out at high temperature. Flavour oils tend to evaporate or degrade to form products having unfavourable off-notes. Powdered flavour compositions, particularly those containing citrus oils, can be of poor quality and exhibit relatively short self-life, as a result.

10

Surprisingly, the incorporation of compounds of formula (I) or flavour compositions containing same into powder compositions, results in powder compositions that exhibit the impact and authenticity of the flavour oils used in their preparation, essentially maintaining flavour oil quality in a powdered flavour formulations.

15

Accordingly, the invention provides in another aspect a powder flavour composition comprising a compound according to formula (I) and at least one additional flavour co-ingredient.

20 In another aspect of the invention there is provided a powder soft drink composition or other dry mix composition comprising a compound according to formula (I).

In yet another aspect of the present invention there is provided a powder soft drink composition or other dry mix composition comprising a powder flavour composition
25 comprising a compound of formula (I).

In yet another aspect of the present invention there is provided a method of forming a powder flavour composition comprising the step of incorporating into said composition a compound according to formula (I).

30

In a particular embodiment of the compound of formula (I) may be added to the formed powder flavour composition, or it may be added to flavour composition before forming the powder.

5 Another particularly preferred class of edible composition according to the present invention are snack foods. Snack foods are a category of product well known to the skilled person in the food industry. These products are described above and include, without limitation, pretzels, corn chips, potato chips, puffed products, extruded products, tortilla chips and the like. Still more particularly, the invention is concerned with low fat snack food
10 compositions. Low fat snack food compositions contain less than 30 % by weight fat, more particularly between 5 to 25 % by weight of fat.

A problem with reducing fat in a snack food composition is the loss in taste and texture. Fats play an important role in the way that dough behaves during processing and greatly
15 affect the quality, flavor and texture of ready-to-eat products. As the fat content in snack products is reduced or replaced with other ingredients (e.g., non-digestible fat, protein, fiber, gums), adverse organoleptic effects (e.g., mouth coating, drying, lack of crispness and lack of flavour) are increased. The adverse organoleptic effects result in products having reduced palatability.

20 Considerable efforts have been expended in devising flavour compositions to overcome the problems associated with low fat snack food products. Flavours may be applied to a snack food as topical coatings in the form of dry powders and/or as liquids (e.g., oil-based, water-based). Another approach has been to add flavour to the dough.

25 Despite these various approaches which have been taken to improve consumer appeal and palatability of snack foods, and particularly low fat snack foods, there is still a need for improved low-fat snack foods having coatings applied thereto with the visual appeal, flavor, and texture of full-fat snack foods.

30

Compounds according to formula (I) or flavour compositions containing same can be incorporated into snack foods to impart an impactful flavour and a mouthfeel with a remarkable roundness and fullness. Furthermore, the taste and mouthfeel effects can be achieved even in low fat snack foods.

5

Accordingly, the invention provides in another of its aspects a snack food comprising a flavour composition as hereinabove described. In a particular embodiment of the invention the snack food has a fat content of about 40 % or less by weight based on the total weight of the snack food, more particularly about 30 % or less, still more particularly 25 % or less, more particularly still about 10 % or less, still more particularly about 5 % or less, still more particularly about 3 % or less.

10

Examples of snack foods are described above and include products processed by oven baking, extrusion or frying, and which are made from potato and/or corn and/or various grains such as rice or wheat.

15

Another particularly preferred class of edible composition according to the present invention is alcoholic beverages.

Applicant surprisingly found that compounds according to formula (I) incorporated into an alcoholic beverage had the effect of increasing the alcohol impact of the beverage.

20

Accordingly, the invention provides in another of its aspects an alcoholic beverage comprising a compound according to formula (I).

25

In yet another aspect of the invention there is provided a method of producing a heightened alcoholic impression in an alcoholic beverage by incorporating into said beverage a compound according to formula (I).

Compounds of formula (I) may be incorporated into said alcoholic beverage in amounts of 1 ppb to 1 ppm.

30

Another preferred class of edible compositions are products ingested in the form of tablets, capsules, powders, multiparticulates and the like, which may include pharmaceuticals and nutraceuticals.

5

Certain groups of people have problems swallowing tablets or capsules, powders, multiparticulates and the like. This problem can be particularly pronounced in certain consumer groups, such as children and the very old or infirm. Applicant surprisingly found that compounds according to the formula (I) when taken into the oral cavity produce a pronounced salivating effect. Incorporating the compounds into these dosage forms, particularly as part of a coating around said dosage forms can ease the swallowing process for consumers, in particular children and the old or infirm.

10

Accordingly, the invention provides in another of its aspects an orally administrable dosage form, in particular in the form of tablets capsules, powders or multiparticulates comprising a compound according to the formula (I).

15

Another preferred class of edible composition is baked goods. Compounds of the formula (I) may be incorporated topically or in-dough. Incorporated at levels of 1 ppb to 1 ppm, the compounds of formula (I) render baked products less dry and more succulent.

20

Other preferred class of edible compositions are caloric or non-caloric beverages containing carbohydrate sweeteners, such as sucrose, high fructose corn syrup, fructose and glucose, or high intensity, non-nutritive sweeteners such as aspartame, acesulfame K, sucralose, cyclamate, sodium saccharin, neotame, rebaudioside A, and/or other stevia-based sweeteners; as well as other optional ingredients such as juices, organic acids such as citric acid, alcohol and functional ingredients.

25

Incorporated at levels of 1 ppb to 10 ppm, compounds of formula (I) impart to said beverages containing sweeteners at levels of less than 1 % and up to about 20 %, an upfront sweetness and mouthfeel that is reminiscent of sugar.

30

Other preferred edible compositions are savoury compositions, in particular those that are soy-based or fish-based.

- 5 Incorporated at levels of 1 ppb to 10 ppm, in a soy-based composition (such as soy sauce) or a fish-based composition (such as fish sauce) containing 5 to 40 % salt, the compositions are found to exhibit strong umami tastes that are long-lasting and rich.

Another preferred edible composition is a clouded beverage composition.

10

Certain beverages such as juices have relatively higher turbidity and thus have an opaque appearance. Often, it is desired that the beverage have a relatively high turbidity. This might be desirable to provide a more natural appearance to beverages with low juice content, or it might be for reasons related to masking sedimentation or "ringing" (where flavour or colour oils rise to the surface of a container during storage). Clouded beverages are usually formed by means of a clouding agent. Clouding agents are usually supplied in the form of emulsions, or the clouding agent may be part of a powdered beverage that upon reconstitution will formed an emulsion providing a permanent cloud to the beverage.

15

- 20 Compounds of the formula (I), in addition to their remarkable organoleptic properties, can lend stability to clouding agents and to beverage compositions containing same.

Accordingly, the invention provides in another of its aspects a composition comprising a beverage clouding composition and a compound of formula (I).

25

In a particular embodiment of the invention, a flavour composition as herein defined may be provided in the form of an emulsion. This emulsion composition may be particularly useful in clouded beverage applications, in particular, in which it is intended to employ a clouding agent.

30

In yet another aspect of the invention there is provided a clouded beverage composition comprising a clouding agent and a compound of the formula (I).

5 Other preferred edible compositions are those compositions that are formed by a process of ripening.

In food processing, it frequently occurs that a food needs to remain for a prolonged period of time and under well-defined conditions to obtain the food with the requisite and recognised quality. A commonly used term for this process is ripening. Ripening is well
10 known in the processing of certain types of cheese, meat, soy-sauce and wine, as well as beer sausage, sauerkraut, tempeh and tofu. There are also specific steps that are carried out for specific reasons (such as water-removal, or off-note removal) that have beneficial effects on the food products. Examples of this are the conching of chocolate and the drying
15 are induced by chemical conversions, enzymatically catalysed conversions or fermentative transformations. All of these conversions are slow and therefore expensive; they are also not fully predictable or controllable.

The compounds of formula (I), having regard to their remarkable property of adding to the
20 authentic taste characteristics of the edible compositions in which they are incorporated, may be added to an edible product during its ripening process in order to reduce storage time without adversely influencing the taste quality of the ripened product.


Accordingly, in another aspect of the invention there is provided a method of ripening a
25 product selected from the group consisting of cheese, meat, soy-sauce and wine, beer, sausage, sauerkraut, tempeh and tofu, comprising the step of ripening the product in the presence of a compound according to the formula (I).

In another aspect of the invention there is provided a method of conching chocolate, said
30 method comprising the step of adding to the chocolate a compound according to the formula (I), or a flavour composition containing same.

There now follows a series of non-limiting examples that serve to illustrate the invention.

Synthesis Examples

5 1. General structure



$R^1 = R^2 = R^3 = H$ or CH_3

10

R^4 is a straight chain hydrocarbon group having 9 to 21 carbon atoms and containing zero to three double bonds

or

15 R^4 is a [(1*E*)-2,6-dimethylhepta-1,5-dien-1-yl]-group.


Several compounds belonging to that general formula have been synthesised according to one of the 2 procedure described below.

20

2. Synthesis

2.1 Example for synthesis route A: (DCC method)


Synthesis of (Z)-2-(hexadec-9-enamido)-4-methylpentanoic acid



In a 250 mL round-bottomed flask was mixed Palmitoleic acid (1 g, 3.93 mmol) with 1-
 5 hydroxypyrrolidine-2,5-dione (0.498 g, 4.32 mmol) in dioxane (50 ml) to give a colorless
 solution. The solution was cooled to 10°C and DCC (0.892 g, 4.32 mmol) was added while
 stirring. Stirring was continued for three hours at room temperature.
 The formed solids were filtered (dicyclohexylurea) and the filtrate was added to a solution
 of Leucine (0.85 g, 6.48 mmol) in a 2% solution of sodiumbicarbonate (0.363 g, 4.32 mmol)
 10 in water.
 The reaction mixture was stirred for 4 hours at 50°C.

Dioxane was evaporated and the aqueous residue was further diluted with water, acidified
 with a diluted hydrochloric acid solution and extracted with ethylacetate.
 15 Organic layers were combined, washed with brine, dried and evaporated to yield 1.3 g of a
 white solid.
 Product was purified by flash column chromatography, eluent DCM/methanol.
 1 g of 85-90% pure product could be obtained.

20



^1H NMR (600 MHz, CHLOROFORM-*d*) δ ppm 0.85 - 0.90 (m, 3 H, H-C(16)) 0.91 - 0.98 (m, 6
 H, H-C(22, 23)) 1.19 - 1.40 (m, 14 H, H-C(4, 5, 6, 7, 12, 13, 14)) 1.49 - 1.75 (m, 7 H, H-C(3, 15,

20, 21)) 2.01 (q, $J=6.07$ Hz, 4 H, H-C(8, 11)) 2.24 (t, $J=7.73$ Hz, 2 H, H-C(2)) 4.54 - 4.59 (m, 1 H, H-C(19)) 5.24 - 5.43 (m, 2 H, H-C(9, 10)) 6.14 (d, $J=8.25$ Hz, 1 H, H-N(19))

^{13}C NMR (150 MHz, CHLOROFORM- d) δ ppm 14.11 (C16)) 21.9 (C(15)) 22.66 (C(22)) 22.86 (C(23)) 24.91 (C21)) 25.63 (C(3)) 27.18 (C(11)) 27.23 (C(8)) 28.99 (C(6)) 29.16 (C(4)) 29.20 (C(5)) 29.25 (C(13)) 29.71 (C(7)) 29.73 (C(12)) 31.79 (C(14)) 36.51 (C(2)) 41.32 (C(20)) 50.87 (C(19)) 129.73 (C(9)) 130.00 (C(10)) 173.95 (C(1)) 176.38 (C(18))

10 **2.2 Route B (DCC method with protection group)**

Step 1:

To a solution of an O-methylated amino acid (16.51 mmol) in DCM (100 ml) was added triethylamine (1.519 g, 15.01 mmol) at minus 15°C. A fatty acid (.01 mmol) was added while stirring. A solution of DCC (15.01 mmol) in 10 mL of DCM was added dropwise at 0°C. The reaction mixture was stirred at 0°C for 1 hour and stirring was continued at room temperature for 3 hours. The dicyclohexylurea was removed by filtration from the reaction mixture. Filtrate was washed with a saturated sodium bicarbonate solution, diluted hydrochloric acid solution and water. Organic layer was separated, dried and evaporated to yield 3 g of an oil. This oil was purified by flash column chromatography, eluent DCM/methanol The intermediate ester compound could be isolated in a purity of 95%.

Step 2:

The O-methylated N-acyl-amino-acid (4.91 mmol) was dissolved in a mixture of Ethanol (8.00 ml) and water (8 ml). To this mixture was added a 32% solution of sodium hydroxide (2.453 g, 19.63 mmol) and mixture was stirred at room temperature for three hours. Mixture stand over for 14 hours. After 14 hours the mixture was acidified with a concentrated hydrochloric acid solution (1.612 ml, 19.63 mmol), diluted with water and extracted with mtbe. Organic layer was separated, dried and evaporated. 1.3 g of a half solid yellow residue was obtained. NMR confirmed the structure of the title compound, purity 95%

30

2.3 Route C (acid chloride)

An amino acid (20 mmol) was dissolved in a solution of sodiumhydroxide (54.5 mmol) in water (40 ml).




Tetrahydrofuran (60 ml) was added. Fatty acid chloride (18.18 mmol) was added dropwise at room temperature. Stirring was continued for 2 hours. Mixture was diluted with water, acidified with a 37% solution of hydrochloric acid (2.99 ml, 36.4 mmol) and extracted with ethylacetate.






Organic layers were combined, dried and evaporated.






The residue contains about 20% free fatty acid according NMR. The solids were stirred with heptane for 30 minutes, filtered and dried. This resulted in 2.4 g of the title compound as a creamy colored solid. (purity 95%).

2.4 All synthesized compounds

15 Table 1: List synthesized compounds


Structure	Amino acid	Carboxylic acid	Structure	Route
1	Leucine	C-8:0		C
2	Leucine	C10:2		B
3	Leucine	C16:0		C

4	Leucine	C16:1		A
5	Leucine	C18:0		C
6	Leucine	C18:1		C
7	Leucine	C18:2		B
8	Leucine	C22:1		A

9	Isoleucine	C18:1		C
10	Valine	C16:0		C
11	Valine	C18:0		C
12	Valine	C18:1		C
13	Valine	C18:2		A

3 NMR data (examples)

3.1 Structure 6 Leu-C18:1




^1H NMR (600 MHz, CHLOROFORM-*d*) δ ppm 0.77 - 0.84 (m, 3 H, H-C(18)) 0.85 - 0.93 (m, 6 H, H-C(24, 25)) 1.14 - 1.29 (m, 20 H, H-C(4, 5, 6, 7, 12, 13, 14, 15, 16, 17)) 1.48 - 1.59 (m, 3 H, H-C(3, 22)) 1.60 - 1.69 (m, 2 H, C-H(22, 23)) 1.90 - 1.99 (m, 4 H, H-C(8, 11)) 2.17 (t, $J=7.39$ Hz, 2 H, H-C(2)) 4.55 (td, $J=8.51, 4.64$ Hz, 1 H, H-C(21)) 5.15 - 5.35 (m, 2 H, H-C(9,10)) 5.95 (d, $J=7.56$ Hz, 1 H, H-N(19))

^{13}C NMR (150 MHz, CHLOROFORM-*d*) δ ppm 13.68 (C(18)) 21.43 (C(17)) 22.24 (C(25)) 22.40 (C(24)) 24.45 (C(23)) 25.14 (C(3)) 26.74 (C(11)) 26.78 (C(8)) 28.71 (C(6)) 28.73 (C(4)) 28.78 (C(5)) 28.88 (C(13, 15)) 29.09 (C(14)) 29.26 (C(7)) 31.46 (C(16)) 36.04 (C(2)) 40.70 (C(22)) 50.41 (C(21)) 129.28 (C(9, 10)) 173.64 (C(1)) 176.11 (C(20))

15

20

25 Structure 10 Val-C16:0



¹H NMR (600 MHz, CHLOROFORM-*d*) δ ppm 0.88 (t, *J*=7.05 Hz, 3 H, H-C(16)) 0.95 (d, *J*=6.87 Hz, 3 H, H-C(21)) 0.98 (d, *J*=6.87 Hz, 3 H, H-C(20)) 1.19 - 1.37 (m, 24 H, H-C(3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14)) 1.59 - 1.71 (m, 2 H, H-C(3)) 2.20 - 2.32 (m, 3 H, H-C(3)) 4.59 (dd, *J*=8.59, 4.81 Hz, 1 H, H-C(18)) 6.19 (d, *J*=8.59 Hz, 1 H, H-N(18))

5

¹³C NMR (150 MHz, CHLOROFORM-*d*) δ ppm 14.13 C(16) 17.70 C(20) 19.02 C(21) 22.71 C(15) 25.78 C(3) 29.25 C(6) 29.35 C(9) 29.38 (C(13) 29.52 C(5)) 29.64 C(4) 29.68 (C(7, 10)) 29.72 C(8, 11, 12) 31.00 (C(19)) 31.94 (C(14)) 36.69 (C(2)) 57.08 C(18) 174.23 (C(1)) 175.49 C(22))

10

Application Examples

Testing Leucine, IsoLeucine and Valine derivatives

15

Samples were evaluated by expert tasters. Tasters were asked to describe the samples focusing on authentic taste, mouthfeel, fullness, salivation, salty-ness, umami, sweetness, juiciness, richness, long lastingness and fattiness.

20 Comparison in a mango juice drink

Base is water, 8% sucrose, 0.1% citric acid, 1% clear mango juice, proprietary mango flavour @0.05%

Base: fruity mango

25 Base plus C16:0-leu at 2 ppm very fruity, ripe mango taste, strawberry note

Base plus C18:1-leu at 2 ppm very long lasting, ripe, fruity, juicy

Base plus C18:0-leu at 2 ppm fruity strawberry like, ripe, long lasting

Base: fruity mango

30 Base plus C18:2-val at 1 ppm ripe, very juicy nice, full body, long lasting

Test in a strawberry drink

Base is 7% sucrose, 0.1% citric acid, proprietary strawberry flavour @ 0.015%

35 Base: sweet, fruity, estery strawberry

Base plus C18:0-leu at 0.5 ppm fruity authentic strawberry, stronger strawberry, ripe, long lasting, juicy

Test in a strawberry yoghurt

- 5 Base: Yoghurt 1.5% fat yoghurt, 8% sucrose, proprietary strawberry flavour @ 0,015%

Base: fruity, strawberry

Base plus C18:0-leu at 2 ppm fruity authentic strawberry , stronger strawberry, ripe, long lasting, juicy, creamy yoghurt.

10

Test in peach flavoured drink:

Base is water, 8% sugar, 0.1% citric acid, proprietary peach flavour at 0.05%

Base: fruity peach

Base plus C18:1-val at 1 ppm tropical, fatty, skin, juicy

- 15 Base plus C16:0-val at 1ppm very juicy

Test in beef bouillon:


Base is Maggi beef bouillon 1 tablet in 500 ml of hot water

- 20 Base: salty, umami, powdery

Base plus C18:2- val at 1 ppm more umami, salivating, salty

Claims:

1. A compound of formula (I) represented by the formula



5

and edible salts thereof,

wherein

R₁ is an alkyl residue containing 6 to 20 carbon atoms, or an alkene residue containing from 9 to 25 carbon atoms with 1 to 6 double bonds, R₁ together with the carbonyl group to which it is attached is a residue of a carboxylic acid,

10

R₃ is hydrogen or methyl, and

R₄ is methyl, ethyl or iso-propyl.

2. A compound according to claim 1 in which R₃ is hydrogen when R₄ is iso-propyl; R₃ is methyl when R₄ is methyl; and R₃ is methyl when R₄ is ethyl.

15

3. A compound according to claim 1 or claim 2 selected from the group consisting of N-palmitenoyl-L-leucine, N-palmitoyl-L-leucine, N-linolenoyl-L-leucine, N-linoleoyl-L-leucine; N-oleoyl-L-leucine; N-oleoyl-Iso leucine; N-palmitenoyl-L-valine, N-palmitoyl-L-valine, N-linolenoyl-L-valine, N-linoleoyl-L-valine and N-oleoyl-L-valine.

20

4. A stock solution comprising 0.01 to 1% by weight of a compound according to formula (I).

5. A stock solution according to claim 4 wherein the solvent is selected from the group consisting of ethanol, triacetine, glycerol and miglyol or mixtures thereof.

25

6. The use of a compound according to any of the preceding claims in an edible composition.
7. A flavour composition comprising a compound according to any of the claims 1 to 3.
- 5 8. A flavour composition according to claim 6 comprising at least one flavour co-ingredient.
9. A flavour composition according to claim 7 or 8 comprising an ingredient selected from the group consisting of sugars, fats, salt, MSG, calcium ions, phosphate ions, organic acids, proteins, purines and mixtures thereof.
- 10 10. A flavour composition according to any of the claims 7 to 9 comprising a carrier material and an adjuvant.
- 15 11. A flavour composition according to claim 10 wherein the adjuvant is an anti-oxidant.
12. A flavour composition according to any of the claims 7 through 11 in the form of an emulsion.
- 20 13. A flavour composition according to any of the claims 7 through 11 in the form of a powder.
14. A flavour composition according to claim 13 wherein the powder is formed by spray drying.
15. A edible composition comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 25 16. An edible composition according to claim 15 in the form of a powdered soft drink or a dry mix composition comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 30

17. A snack food according to claim 15 comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14 incorporated in the snack food or applied topically to the snack food as a powder.
- 5
18. An alcoholic beverage according to claim 15 comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 10
19. An orally administrable tablet, capsule, powder or multiparticulate according to claim 15 comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 15
20. A caloric or non-caloric beverage according to claim 15 containing carbohydrate sweeteners, selected from sucrose, high fructose corn syrup, fructose and glucose, or high intensity, non-nutritive sweeteners selected from aspartame, acesulfame K, sucralose, cyclamate, sodium saccharin, neotame, rebaudioside A, and/or other stevia-based sweeteners, comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 20
21. A soy-based edible composition according to claim 15 comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 25
22. A clouded beverage composition according to claim 15 comprising a compound of formula (I) according to any of the claims 1 through 3 or a flavour composition according to any of the claims 7 through 14.
- 30
23. An edible product according to claim 15, formed by a process of ripening selected from the group consisting of cheese, meat, soy-sauce and wine, as well as beer

sausage, sauerkraut, chocolate tempeh and tofu, comprising the step of ripening the product in the presence of a compound of formula (I) according to any of the claims 1 through 3.

- 5 24. A method of ripening a product selected from the group consisting of cheese, meat, soy-sauce and wine, beer, sausage, sauerkraut, chocolate tempeh and tofu, comprising the step of ripening the product in the presence of compound of formula (I) according to any of the claims 1 through 3.
- 10 25. A method of conching chocolate in the presence of a compound according to the formula (I) as defined in any of the claims 1 through 3.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/034363

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A23L1/227 C07C233/47 C07C233/49
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 A23L C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 442 090 A (KAKEYA NOBUHARU [JP] ET AL) 10 April 1984 (1984-04-10) column 3, line 65 - column 4, line 17 -----	1-5,7,9,10
X	US 4 016 287 A (EBERHARDT HANS ET AL) 5 April 1977 (1977-04-05) column 3, line 65 - column 4, line 17 -----	1,2,4,5,7,9,10
X	US 3 947 589 A (MISATO TOMOMASA ET AL) 30 March 1976 (1976-03-30) column 5, line 55 - column 6, line 10 -----	1,2,4,5,7
X	US 2007/282002 A1 (MAEZONO KATSUMI [JP] ET AL) 6 December 2007 (2007-12-06) page 4 - page 5 ----- -/--	1,2,4-7,9,10,15,19

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 5 July 2013	Date of mailing of the international search report 18/07/2013
--	--

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Panday, Narendra
--	--

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2013/034363

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01/59067 A2 (YANG ZHENHUA [US]) 16 August 2001 (2001-08-16) claims 6, 26 -----	1,2,4-7, 9,10,15, 19
X	EP 0 891 764 A1 (KASHIMA OIL [JP]) 20 January 1999 (1999-01-20) claim 1 -----	1,2,4,5, 7,9,10, 12
X	FR 2 878 439 A1 (SEPPIC SA [FR]) 2 June 2006 (2006-06-02) claim 9 -----	1-5,7,9, 10,12
X	GB 1 426 545 A (KAO CORP) 3 March 1976 (1976-03-03) claims 6, 7 -----	1-4,7,9, 10
X	WO 2009/141294 A1 (NESTEC SA [CH]; RIERA CELINE [CH]; MENOZZI CANDICE [CH]; ROBERT FABIEN) 26 November 2009 (2009-11-26) cited in the application claim 1 figure 2 -----	1-25
Y		1-25
X	EP 2 140 770 A1 (NESTEC SA [CH]) 6 January 2010 (2010-01-06) claim 1 -----	1-25
Y		1-25
Y	DATABASE WPI Week 199626 Thomson Scientific, London, GB; AN 1996-253848 XP002699514, & JP H08 103242 A (HASEGAWA CO LTD) 23 April 1996 (1996-04-23) abstract -----	1-25
T	WO 2013/010991 A1 (GIVAUDAN SA [CH]; YANG XIAOGEN [US]; LUBIAN ELISABETTA [CH]; RENES HAR) 24 January 2013 (2013-01-24) claim 1 examples 3, 5, 7 -----	
T	EP 2 597 082 A1 (SYMRISE AG [DE]) 29 May 2013 (2013-05-29) claim 1 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2013/034363

Patent document cited in search report	Publication date	Publication date	Patent family member(s)	Publication date
US 4442090	A	10-04-1984	NONE	

US 4016287	A	05-04-1977	NONE	

US 3947589	A	30-03-1976	NONE	

US 2007282002	A1	06-12-2007	JP 2008105945	A 08-05-2008
			US 2007282002	A1 06-12-2007
			US 2010063151	A1 11-03-2010
			WO 2006082978	A1 10-08-2006

WO 0159067	A2	16-08-2001	AU 778366	B2 02-12-2004
			AU 3442901	A 20-08-2001
			CA 2398923	A1 16-08-2001
			EP 1259470	A2 27-11-2002
			JP 2003523975	A 12-08-2003
			US 2004034242	A1 19-02-2004
			WO 0159067	A2 16-08-2001

EP 0891764	A1	20-01-1999	DE 69821685	D1 25-03-2004
			DE 69821685	T2 09-12-2004
			EP 0891764	A1 20-01-1999

FR 2878439	A1	02-06-2006	NONE	

GB 1426545	A	03-03-1976	DE 2443646	A1 27-03-1975
			GB 1426545	A 03-03-1976
			JP S5053545	A 12-05-1975
			JP S5247022	B2 29-11-1977

WO 2009141294	A1	26-11-2009	CA 2725155	A1 26-11-2009
			CN 102036571	A 27-04-2011
			EP 2296492	A1 23-03-2011
			JP 2011520457	A 21-07-2011
			US 2011129583	A1 02-06-2011
			WO 2009141294	A1 26-11-2009

EP 2140770	A1	06-01-2010	NONE	

JP H08103242	A	23-04-1996	NONE	

WO 2013010991	A1	24-01-2013	NONE	

EP 2597082	A1	29-05-2013	NONE	
