Office de la Propriété Intellectuelle du Canada
Un organisme d'Industrie Canada

Canadian Intellectual Property Office
An agency of Industry Canada

CA 2494236 A1 2004/02/19

(21) 2 494 236

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2003/08/06
(87) Date publication PCT/PCT Publication Date: 2004/02/19
(85) Entrée phase nationale/National Entry: 2005/02/22
(86) N° demande PCT/PCT Application No.: US 2003/024551
(87) N° publication PCT/PCT Publication No.: 2004/015071
(30) Priorité/Priority: 2002/08/07 (60/401,739) US

(51) Cl.Int.^{7}/Int.Cl.^{7} C12Q 1/00, A61K 49/00, C12Q 1/68, G01N 33/53, G01N 33/00

(71) Demandeur/Applicant:
EXELIXIS, INC., US

(72) Inventeurs/Inventors:
FRANCIS-LANG, HELEN, US;
FRIEDMAN, LORI, US;
KIDD, THOMAS, US;
ROCHE, SIOBHAN, IE;
ZHANG, HAIGUANG, US

(74) Agent: BENNETT JONES LLP

(54) Titre : GENES CSNK1GS UTILISES EN TANT QUE MODIFICATEURS DE LA VOIE p21 ET PROCEDE D'UTILISATION ASSOCIE

(54) Title: CSNK1GS AS MODIFIERS OF THE p21 PATHWAY AND METHODS OF USE

(57) Abrégé/Abstract:
Human CSNK1G genes are identified as modulators of the p21 pathway, and thus are therapeutic targets for disorders associated with defective p21 function. Methods for identifying modulators of p21, comprising screening for agents that modulate the activity of CSNK1G are provided.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
19 February 2004 (19.02.2004)

PCT

(10) International Publication Number
WO 2004/015071 A3

(51) International Patent Classification[7]: C12Q 1/00, 1/68, G01N 33/53, 33/00, A61K 49/00

(21) International Application Number: PCT/US2003/024551

(22) International Filing Date: 6 August 2003 (06.08.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/401,739 7 August 2002 (07.08.2002) US

(71) Applicant *(for all designated States except US)*: **EXELIXIS, INC.** [US/US]; P.O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **FRANCIS-LANG, Helen** [GB/US]; 1782 Pacific Avenue, Apt. 2, San Francisco, CA 94109 (US). **FRIEDMAN, Lori** [US/US]; 113 Arundel Road, San Carlos, CA 94070 (US). **KIDD, Thomas** [GB/US]; 643 Lake Street, San Fransisco, CA 94118 (US). **ROCHE, Siobhan** [IE/IE]; 30 Moatfield Park, Coolock, Dublin 5 (IE). **ZHANG, HaiGuang** [US/US]; 4833 El Grande Place, El Sobrante, CA 94803 (US).

(74) Agents: **SHAYESTEH, Laleh** et al.; Exelixis, Inc., P. O. Box 511, 170 Harbor Way, South San Francisco, CA 94083-0511 (US).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report*

(88) Date of publication of the international search report:
12 August 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CSNK1GS AS MODIFIERS OF THE p21 PATHWAY AND METHODS OF USE

(57) Abstract: Human CSNK1G genes are identified as modulators of the p21 pathway, and thus are therapeutic targets for disorders associated with defective p21 function. Methods for identifying modulators of p21, comprising screening for agents that modulate the activity of CSNK1G are provided.

WO 2004/015071 A3

CSNK1GS AS MODIFIERS OF THE p21 PATHWAY AND METHODS OF USE

REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. provisional patent application US 60/401,739 filed 8/7/2002. The content of the prior application is hereby incorporated in its entirety.

BACKGROUND OF THE INVENTION

The p21/CDKN1/WAF1/CIP1 protein(El-Deiry, W. S.; et al. Cell 75: 817-825, 1993; Harper, J. W.; et al. Cell 75: 805-816, 1993; Huppi, Ket al. Oncogene 9: 3017-3020, 1994) is a cell cycle control protein that inhibits cyclin-kinase activity, is tightly regulated at the transcriptional level by p53, and mediates p53 suppression of tumor cell growth. Along with p53, p21 appears to be essential for maintaining the G2 checkpoint in human cells (Bunz, F.; Dutriaux, A.; et al. Science 282:1497-1501, 1998). Sequences of P21 are well-conserved throughout evolution, and have been identified in species as diverse as human (Genbank Identifier 13643057), Drosophila melanogaster (GI# 1684911), Caenorhabditis elegans (GI#4966283), and yeast (GI#2656016).

Casein kinase I is the most abundant serine/threonine kinase in eukaryotic cell extracts, and preferentially phosphorylate acidic substrates using ATP as a phosphate donor. Multiple isoforms of the enzyme exist. The gamma-1 isoform (CSNK1G1) is involved in growth and morphogenesis of eukaryotic cells (Kusuda, J., et al (2000) Cytogenet Cell Genet 90:298-302), the gamma-2 (CSNK1G2) and gamma-3 (CSNK1G3) isoforms may play a role in signal transduction (Kitabayashi, A. N., et al. (1997) Genomics 46, 133-7; Kusuda, J., et al (1998) Cytogenet Cell Genet 83:101-3).

The ability to manipulate the genomes of model organisms such as *Drosophila* provides a powerful means to analyze biochemical processes that, due to significant evolutionary conservation, have direct relevance to more complex vertebrate organisms. Due to a high level of gene and pathway conservation, the strong similarity of cellular processes, and the functional conservation of genes between these model organisms and mammals, identification of the involvement of novel genes in particular pathways and their functions in such model organisms can directly contribute to the understanding of the correlative pathways and methods of modulating them in mammals (see, for example, Mechler BM et al., 1985 EMBO J 4:1551-1557; Gateff E. 1982 Adv. Cancer Res. 37: 33-74; Watson KL., et al., 1994 J Cell Sci. 18: 19-33; Miklos GL, and Rubin GM. 1996 Cell

86:521-529; Wassarman DA, et al., 1995 Curr Opin Gen Dev 5: 44-50; and Booth DR. 1999 Cancer Metastasis Rev. 18: 261-284). For example, a genetic screen can be carried out in an invertebrate model organism having underexpression (e.g. knockout) or overexpression of a gene (referred to as a "genetic entry point") that yields a visible phenotype. Additional genes are mutated in a random or targeted manner. When a gene mutation changes the original phenotype caused by the mutation in the genetic entry point, the gene is identified as a "modifier" involved in the same or overlapping pathway as the genetic entry point. When the genetic entry point is an ortholog of a human gene implicated in a disease pathway, such as p21, modifier genes can be identified that may be attractive candidate targets for novel therapeutics.

All references cited herein, including patents, patent applications, publications, and sequence information in referenced Genbank identifier numbers, are incorporated herein in their entireties.

SUMMARY OF THE INVENTION

We have discovered genes that modify the p21 pathway in *Drosophila*, and identified their human orthologs, hereinafter referred to as casein kinase I gamma isoforms (CSNK1G). The invention provides methods for utilizing these p21 modifier genes and polypeptides to identify CSNK1G-modulating agents that are candidate therapeutic agents that can be used in the treatment of disorders associated with defective or impaired p21 function and/or CSNK1G function. Preferred CSNK1G-modulating agents specifically bind to CSNK1G polypeptides and restore p21 function. Other preferred CSNK1G-modulating agents are nucleic acid modulators such as antisense oligomers and RNAi that repress CSNK1G gene expression or product activity by, for example, binding to and inhibiting the respective nucleic acid (i.e. DNA or mRNA).

CSNK1G modulating agents may be evaluated by any convenient *in vitro* or *in vivo* assay for molecular interaction with a CSNK1G polypeptide or nucleic acid. In one embodiment, candidate CSNK1G modulating agents are tested with an assay system comprising a CSNK1G polypeptide or nucleic acid. Agents that produce a change in the activity of the assay system relative to controls are identified as candidate p21 modulating agents. The assay system may be cell-based or cell-free. CSNK1G-modulating agents include CSNK1G related proteins (e.g. dominant negative mutants, and biotherapeutics); CSNK1G -specific antibodies; CSNK1G -specific antisense oligomers and other nucleic acid modulators; and chemical agents that specifically bind to or interact with CSNK1G or

compete with CSNK1G binding partner (e.g. by binding to a CSNK1G binding partner). In one specific embodiment, a small molecule modulator is identified using a kinase assay. In specific embodiments, the screening assay system is selected from a binding assay, an apoptosis assay, a cell proliferation assay, an angiogenesis assay, and a hypoxic induction assay.

In another embodiment, candidate p21 pathway modulating agents are further tested using a second assay system that detects changes in the p21 pathway, such as angiogenic, apoptotic, or cell proliferation changes produced by the originally identified candidate agent or an agent derived from the original agent. The second assay system may use cultured cells or non-human animals. In specific embodiments, the secondary assay system uses non-human animals, including animals predetermined to have a disease or disorder implicating the p21 pathway, such as an angiogenic, apoptotic, or cell proliferation disorder (e.g. cancer).

The invention further provides methods for modulating the CSNK1G function and/or the p21 pathway in a mammalian cell by contacting the mammalian cell with an agent that specifically binds a CSNK1G polypeptide or nucleic acid. The agent may be a small molecule modulator, a nucleic acid modulator, or an antibody and may be administered to a mammalian animal predetermined to have a pathology associated the p21 pathway.

DETAILED DESCRIPTION OF THE INVENTION

Genetic screens were designed to identify modifiers of the p21 pathway in *Drosophila*, where a dominant loss of function screen was carried out to identify genes that interact with the cyclin dependent kinase inhibitor, p21 (Bourne HR, et al., Nature (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95). Expression of the p21 gene in the eye causes deterioration of normal eye morphology. Modifiers of the eye phenotype were identified as members of the p21 pathway. The GISH (CG6963) gene was identified as a modifier of the p21 pathway. Accordingly, vertebrate orthologs of these modifiers, and preferably the human orthologs, CSNK1G genes (i.e., nucleic acids and polypeptides) are attractive drug targets for the treatment of pathologies associated with a defective p21 signaling pathway, such as cancer.

In vitro and in vivo methods of assessing CSNK1G function are provided herein. Modulation of the CSNK1G or their respective binding partners is useful for understanding the association of the p21 pathway and its members in normal and disease

conditions and for developing diagnostics and therapeutic modalities for p21 related pathologies. CSNK1G-modulating agents that act by inhibiting or enhancing CSNK1G expression, directly or indirectly, for example, by affecting a CSNK1G function such as enzymatic (e.g., catalytic) or binding activity, can be identified using methods provided herein. CSNK1G modulating agents are useful in diagnosis, therapy and pharmaceutical development.

Nucleic acids and polypeptides of the invention

Sequences related to CSNK1G nucleic acids and polypeptides that can be used in the invention are disclosed in Genbank (referenced by Genbank identifier (GI) number) as GI#s 11545750 (SEQ ID NO:1), 16878051 (SEQ ID NO:2), 10439939 (SEQ ID NO:3), 21733126 (SEQ ID NO:4), 27375099 (SEQ ID NO:5), 21314777 (SEQ ID NO:6), 18088089 (SEQ ID NO:7), 2199528 (SEQ ID NO:8), 9956036 (SEQ ID NO:9), 4758079 (SEQ ID NO:10), 4590041 (SEQ ID NO:11), and 28837292 (SEQ ID NO:12) for nucleic acid, and GI#s 11545751 (SEQ ID NO:13), 5579454 (SEQ ID NO:14), 21314778 (SEQ ID NO:15), and 4758080 (SEQ ID NO:16) for polypeptides.

The term "CSNK1G polypeptide" refers to a full-length CSNK1G protein or a functionally active fragment or derivative thereof. A "functionally active" CSNK1G fragment or derivative exhibits one or more functional activities associated with a full-length, wild-type CSNK1G protein, such as antigenic or immunogenic activity, enzymatic activity, ability to bind natural cellular substrates, etc. The functional activity of CSNK1G proteins, derivatives and fragments can be assayed by various methods known to one skilled in the art (Current Protocols in Protein Science (1998) Coligan *et al.*, eds., John Wiley & Sons, Inc., Somerset, New Jersey) and as further discussed below. In one embodiment, a functionally active CSNK1G polypeptide is a CSNK1G derivative capable of rescuing defective endogenous CSNK1G activity, such as in cell based or animal assays; the rescuing derivative may be from the same or a different species. For purposes herein, functionally active fragments also include those fragments that comprise one or more structural domains of a CSNK1G, such as a kinase domain or a binding domain. Protein domains can be identified using the PFAM program (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2). For example, the kinase domain (PFAM 00069) of CSNK1G from GI#s 11545751, 21314778, and 4758080 (SEQ ID NOs:13, 15, and 16, respectively) is located respectively at approximately amino acid residues 44-312, 46-312, and 43-308. Methods for obtaining CSNK1G polypeptides are also further described below. In some

embodiments, preferred fragments are functionally active, domain-containing fragments comprising at least 25 contiguous amino acids, preferably at least 50, more preferably 75, and most preferably at least 100 contiguous amino acids of any one of SEQ ID NOs:13-16 (a CSNK1G). In further preferred embodiments, the fragment comprises the entire kinase (functionally active) domain.

The term "CSNK1G nucleic acid" refers to a DNA or RNA molecule that encodes a CSNK1G polypeptide. Preferably, the CSNK1G polypeptide or nucleic acid or fragment thereof is from a human, but can also be an ortholog, or derivative thereof with at least 70% sequence identity, preferably at least 80%, more preferably 85%, still more preferably 90%, and most preferably at least 95% sequence identity with human CSNK1G. Methods of identifying orthlogs are known in the art. Normally, orthologs in different species retain the same function, due to presence of one or more protein motifs and/or 3-dimensional structures. Orthologs are generally identified by sequence homology analysis, such as BLAST analysis, usually using protein bait sequences. Sequences are assigned as a potential ortholog if the best hit sequence from the forward BLAST result retrieves the original query sequence in the reverse BLAST (Huynen MA and Bork P, Proc Natl Acad Sci (1998) 95:5849-5856; Huynen MA *et al.*, Genome Research (2000) 10:1204-1210). Programs for multiple sequence alignment, such as CLUSTAL (Thompson JD et al, 1994, Nucleic Acids Res 22:4673-4680) may be used to highlight conserved regions and/or residues of orthologous proteins and to generate phylogenetic trees. In a phylogenetic tree representing multiple homologous sequences from diverse species (e.g., retrieved through BLAST analysis), orthologous sequences from two species generally appear closest on the tree with respect to all other sequences from these two species. Structural threading or other analysis of protein folding (e.g., using software by ProCeryon, Biosciences, Salzburg, Austria) may also identify potential orthologs. In evolution, when a gene duplication event follows speciation, a single gene in one species, such as *Drosophila*, may correspond to multiple genes (paralogs) in another, such as human. As used herein, the term "orthologs" encompasses paralogs. As used herein, "percent (%) sequence identity" with respect to a subject sequence, or a specified portion of a subject sequence, is defined as the percentage of nucleotides or amino acids in the candidate derivative sequence identical with the nucleotides or amino acids in the subject sequence (or specified portion thereof), after aligning the sequences and introducing gaps, if necessary to achieve the maximum percent sequence identity, as generated by the program WU-BLAST-2.0a19 (Altschul *et al.*, J. Mol. Biol. (1997) 215:403-410) with all

the search parameters set to default values. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched. A % identity value is determined by the number of matching identical nucleotides or amino acids divided by the sequence length for which the percent identity is being reported. "Percent (%) amino acid sequence similarity" is determined by doing the same calculation as for determining % amino acid sequence identity, but including conservative amino acid substitutions in addition to identical amino acids in the computation.

A conservative amino acid substitution is one in which an amino acid is substituted for another amino acid having similar properties such that the folding or activity of the protein is not significantly affected. Aromatic amino acids that can be substituted for each other are phenylalanine, tryptophan, and tyrosine; interchangeable hydrophobic amino acids are leucine, isoleucine, methionine, and valine; interchangeable polar amino acids are glutamine and asparagine; interchangeable basic amino acids are arginine, lysine and histidine; interchangeable acidic amino acids are aspartic acid and glutamic acid; and interchangeable small amino acids are alanine, serine, threonine, cysteine and glycine.

Alternatively, an alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman (Smith and Waterman, 1981, Advances in Applied Mathematics 2:482-489; database: European Bioinformatics Institute; Smith and Waterman, 1981, J. of Molec.Biol., 147:195-197; Nicholas et al., 1998, "A Tutorial on Searching Sequence Databases and Sequence Scoring Methods" (www.psc.edu) and references cited therein.; W.R. Pearson, 1991, Genomics 11:635-650). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff (Dayhoff: Atlas of Protein Sequences and Structure, M. O. Dayhoff ed., 5 suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA), and normalized by Gribskov (Gribskov 1986 Nucl. Acids Res. 14(6):6745-6763). The Smith-Waterman algorithm may be employed where default parameters are used for scoring (for example, gap open penalty of 12, gap extension penalty of two). From the data generated, the "Match" value reflects "sequence identity."

Derivative nucleic acid molecules of the subject nucleic acid molecules include sequences that hybridize to the nucleic acid sequence of any of SEQ ID NOs:1-12. The stringency of hybridization can be controlled by temperature, ionic strength, pH, and the presence of denaturing agents such as formamide during hybridization and washing.

Conditions routinely used are set out in readily available procedure texts (*e.g.*, Current Protocol in Molecular Biology, Vol. 1, Chap. 2.10, John Wiley & Sons, Publishers (1994); Sambrook *et al.*, Molecular Cloning, Cold Spring Harbor (1989)). In some embodiments, a nucleic acid molecule of the invention is capable of hybridizing to a nucleic acid molecule containing the nucleotide sequence of any one of SEQ ID NOs:1-12 under high stringency hybridization conditions that are: prehybridization of filters containing nucleic acid for 8 hours to overnight at 65° C in a solution comprising 6X single strength citrate (SSC) (1X SSC is 0.15 M NaCl, 0.015 M Na citrate; pH 7.0), 5X Denhardt's solution, 0.05% sodium pyrophosphate and 100 μg/ml herring sperm DNA; hybridization for 18-20 hours at 65° C in a solution containing 6X SSC, 1X Denhardt's solution, 100 μg/ml yeast tRNA and 0.05% sodium pyrophosphate; and washing of filters at 65° C for 1h in a solution containing 0.1X SSC and 0.1% SDS (sodium dodecyl sulfate).

In other embodiments, moderately stringent hybridization conditions are used that are: pretreatment of filters containing nucleic acid for 6 h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.1% PVP, 0.1% Ficoll, 1% BSA, and 500 μg/ml denatured salmon sperm DNA; hybridization for 18-20h at 40° C in a solution containing 35% formamide, 5X SSC, 50 mM Tris-HCl (pH7.5), 5mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 μg/ml salmon sperm DNA, and 10% (wt/vol) dextran sulfate; followed by washing twice for 1 hour at 55° C in a solution containing 2X SSC and 0.1% SDS.

Alternatively, low stringency conditions can be used that are: incubation for 8 hours to overnight at 37° C in a solution comprising 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 μg/ml denatured sheared salmon sperm DNA; hybridization in the same buffer for 18 to 20 hours; and washing of filters in 1 x SSC at about 37° C for 1 hour.

Isolation, Production, Expression, and Mis-expression of CSNK1G Nucleic Acids and Polypeptides

CSNK1G nucleic acids and polypeptides are useful for identifying and testing agents that modulate CSNK1G function and for other applications related to the involvement of CSNK1G in the p21 pathway. CSNK1G nucleic acids and derivatives and orthologs thereof may be obtained using any available method. For instance, techniques for isolating cDNA or genomic DNA sequences of interest by screening DNA libraries or by using polymerase chain reaction (PCR) are well known in the art. In general, the

particular use for the protein will dictate the particulars of expression, production, and purification methods. For instance, production of proteins for use in screening for modulating agents may require methods that preserve specific biological activities of these proteins, whereas production of proteins for antibody generation may require structural integrity of particular epitopes. Expression of proteins to be purified for screening or antibody production may require the addition of specific tags (*e.g.*, generation of fusion proteins). Overexpression of a CSNK1G protein for assays used to assess CSNK1G function, such as involvement in cell cycle regulation or hypoxic response, may require expression in eukaryotic cell lines capable of these cellular activities. Techniques for the expression, production, and purification of proteins are well known in the art; any suitable means therefore may be used (e.g., Higgins SJ and Hames BD (eds.) Protein Expression: A Practical Approach, Oxford University Press Inc., New York 1999; Stanbury PF et al., Principles of Fermentation Technology, 2nd edition, Elsevier Science, New York, 1995; Doonan S (ed.) Protein Purification Protocols, Humana Press, New Jersey, 1996; Coligan JE et al, Current Protocols in Protein Science (eds.), 1999, John Wiley & Sons, New York). In particular embodiments, recombinant CSNK1G is expressed in a cell line known to have defective p21 function, such as HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). The recombinant cells are used in cell-based screening assay systems of the invention, as described further below.

The nucleotide sequence encoding a CSNK1G polypeptide can be inserted into any appropriate expression vector. The necessary transcriptional and translational signals, including promoter/enhancer element, can derive from the native CSNK1G gene and/or its flanking regions or can be heterologous. A variety of host-vector expression systems may be utilized, such as mammalian cell systems infected with virus (*e.g.* vaccinia virus, adenovirus, *etc.*); insect cell systems infected with virus (*e.g.* baculovirus); microorganisms such as yeast containing yeast vectors, or bacteria transformed with bacteriophage, plasmid, or cosmid DNA. An isolated host cell strain that modulates the expression of, modifies, and/or specifically processes the gene product may be used.

To detect expression of the CSNK1G gene product, the expression vector can comprise a promoter operably linked to a CSNK1G gene nucleic acid, one or more origins of replication, and, one or more selectable markers (*e.g.* thymidine kinase activity, resistance to antibiotics, *etc.*). Alternatively, recombinant expression vectors can be identified by assaying for the expression of the CSNK1G gene product based on the

physical or functional properties of the CSNK1G protein in *in vitro* assay systems (*e.g.* immunoassays).

The CSNK1G protein, fragment, or derivative may be optionally expressed as a fusion, or chimeric protein product (i.e. it is joined via a peptide bond to a heterologous protein sequence of a different protein), for example to facilitate purification or detection. A chimeric product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other using standard methods and expressing the chimeric product. A chimeric product may also be made by protein synthetic techniques, *e.g.* by use of a peptide synthesizer (Hunkapiller *et al.*, Nature (1984) 310:105-111).

Once a recombinant cell that expresses the CSNK1G gene sequence is identified, the gene product can be isolated and purified using standard methods (*e.g.* ion exchange, affinity, and gel exclusion chromatography; centrifugation; differential solubility; electrophoresis). Alternatively, native CSNK1G proteins can be purified from natural sources, by standard methods (*e.g.* immunoaffinity purification). Once a protein is obtained, it may be quantified and its activity measured by appropriate methods, such as immunoassay, bioassay, or other measurements of physical properties, such as crystallography.

The methods of this invention may also use cells that have been engineered for altered expression (mis-expression) of CSNK1G or other genes associated with the p21 pathway. As used herein, mis-expression encompasses ectopic expression, over-expression, under-expression, and non-expression (*e.g.* by gene knock-out or blocking expression that would otherwise normally occur).

Genetically modified animals

Animal models that have been genetically modified to alter CSNK1G expression may be used in *in vivo* assays to test for activity of a candidate p21 modulating agent, or to further assess the role of CSNK1G in a p21 pathway process such as apoptosis or cell proliferation. Preferably, the altered CSNK1G expression results in a detectable phenotype, such as decreased or increased levels of cell proliferation, angiogenesis, or apoptosis compared to control animals having normal CSNK1G expression. The genetically modified animal may additionally have altered p21 expression (e.g. p21 knockout). Preferred genetically modified animals are mammals such as primates, rodents (preferably mice or rats), among others. Preferred non-mammalian species include

zebrafish, *C. elegans*, and *Drosophila.* Preferred genetically modified animals are transgenic animals having a heterologous nucleic acid sequence present as an extrachromosomal element in a portion of its cells, i.e. mosaic animals (see, for example, techniques described by Jakobovits, 1994, Curr. Biol. 4:761-763.) or stably integrated into its germ line DNA (i.e., in the genomic sequence of most or all of its cells). Heterologous nucleic acid is introduced into the germ line of such transgenic animals by genetic manipulation of, for example, embryos or embryonic stem cells of the host animal.

Methods of making transgenic animals are well-known in the art (for transgenic mice see Brinster et al., Proc. Nat. Acad. Sci. USA 82: 4438-4442 (1985), U.S. Pat. Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Pat. No. 4,873,191 by Wagner et al., and Hogan, B., Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1986); for particle bombardment see U.S. Pat. No., 4,945,050, by Sandford *et al.*; for transgenic *Drosophila* see Rubin and Spradling, Science (1982) 218:348-53 and U.S. Pat. No. 4,670,388; for transgenic insects see Berghammer A.J. *et al.*, A Universal Marker for Transgenic Insects (1999) Nature 402:370-371; for transgenic Zebrafish see Lin S., Transgenic Zebrafish, Methods Mol Biol. (2000);136:375-3830); for microinjection procedures for fish, amphibian eggs and birds see Houdebine and Chourrout, Experientia (1991) 47:897-905; for transgenic rats see Hammer *et al.*, Cell (1990) 63:1099-1112; and for culturing of embryonic stem (ES) cells and the subsequent production of transgenic animals by the introduction of DNA into ES cells using methods such as electroporation, calcium phosphate/DNA precipitation and direct injection see, e.g., Teratocarcinomas and Embryonic Stem Cells, A Practical Approach, E. J. Robertson, ed., IRL Press (1987)). Clones of the nonhuman transgenic animals can be produced according to available methods (see Wilmut, I. *et al.* (1997) Nature 385:810-813; and PCT International Publication Nos. WO 97/07668 and WO 97/07669).

In one embodiment, the transgenic animal is a "knock-out" animal having a heterozygous or homozygous alteration in the sequence of an endogenous CSNK1G gene that results in a decrease of CSNK1G function, preferably such that CSNK1G expression is undetectable or insignificant. Knock-out animals are typically generated by homologous recombination with a vector comprising a transgene having at least a portion of the gene to be knocked out. Typically a deletion, addition or substitution has been introduced into the transgene to functionally disrupt it. The transgene can be a human gene (e.g., from a human genomic clone) but more preferably is an ortholog of the human gene derived from the transgenic host species. For example, a mouse CSNK1G gene is

used to construct a homologous recombination vector suitable for altering an endogenous CSNK1G gene in the mouse genome. Detailed methodologies for homologous recombination in mice are available (see Capecchi, Science (1989) 244:1288-1292; Joyner *et al.*, Nature (1989) 338:153-156). Procedures for the production of non-rodent transgenic mammals and other animals are also available (Houdebine and Chourrout, *supra*; Pursel *et al.*, Science (1989) 244:1281-1288; Simms *et al.*, Bio/Technology (1988) 6:179-183). In a preferred embodiment, knock-out animals, such as mice harboring a knockout of a specific gene, may be used to produce antibodies against the human counterpart of the gene that has been knocked out (Claesson MH et al., (1994) Scan J Immunol 40:257-264; Declerck PJ et al., (1995) J Biol Chem. 270:8397-400).

In another embodiment, the transgenic animal is a "knock-in" animal having an alteration in its genome that results in altered expression (e.g., increased (including ectopic) or decreased expression) of the CSNK1G gene, e.g., by introduction of additional copies of CSNK1G, or by operatively inserting a regulatory sequence that provides for altered expression of an endogenous copy of the CSNK1G gene. Such regulatory sequences include inducible, tissue-specific, and constitutive promoters and enhancer elements. The knock-in can be homozygous or heterozygous.

Transgenic nonhuman animals can also be produced that contain selected systems allowing for regulated expression of the transgene. One example of such a system that may be produced is the cre/loxP recombinase system of bacteriophage P1 (Lakso *et al.*, PNAS (1992) 89:6232-6236; U.S. Pat. No. 4,959,317). If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase. Another example of a recombinase system is the FLP recombinase system of Saccharomyces cerevisiae (O'Gorman et al. (1991) Science 251:1351-1355; U.S. Pat. No. 5,654,182). In a preferred embodiment, both Cre-LoxP and Flp-Frt are used in the same system to regulate expression of the transgene, and for sequential deletion of vector sequences in the same cell (Sun X et al (2000) Nat Genet 25:83-6).

The genetically modified animals can be used in genetic studies to further elucidate the p21 pathway, as animal models of disease and disorders implicating defective p21 function, and for *in vivo* testing of candidate therapeutic agents, such as those identified in

screens described below. The candidate therapeutic agents are administered to a genetically modified animal having altered CSNK1G function and phenotypic changes are compared with appropriate control animals such as genetically modified animals that receive placebo treatment, and/or animals with unaltered CSNK1G expression that receive candidate therapeutic agent.

In addition to the above-described genetically modified animals having altered CSNK1G function, animal models having defective p21 function (and otherwise normal CSNK1G function), can be used in the methods of the present invention. For example, a p21 knockout mouse can be used to assess, *in vivo*, the activity of a candidate p21 modulating agent identified in one of the *in vitro* assays described below. p21 knockout mice are described in the literature (Umanoff H, et al., Proc Natl Acad Sci U S A 1995 Feb 28;92(5):1709-13). Preferably, the candidate p21 modulating agent when administered to a model system with cells defective in p21 function, produces a detectable phenotypic change in the model system indicating that the p21 function is restored, i.e., the cells exhibit normal cell cycle progression.

Modulating Agents

The invention provides methods to identify agents that interact with and/or modulate the function of CSNK1G and/or the p21 pathway. Modulating agents identified by the methods are also part of the invention. Such agents are useful in a variety of diagnostic and therapeutic applications associated with the p21 pathway, as well as in further analysis of the CSNK1G protein and its contribution to the p21 pathway. Accordingly, the invention also provides methods for modulating the p21 pathway comprising the step of specifically modulating CSNK1G activity by administering a CSNK1G-interacting or -modulating agent.

As used herein, an "CSNK1G-modulating agent" is any agent that modulates CSNK1G function, for example, an agent that interacts with CSNK1G to inhibit or enhance CSNK1G activity or otherwise affect normal CSNK1G function. CSNK1G function can be affected at any level, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In a preferred embodiment, the CSNK1G - modulating agent specifically modulates the function of the CSNK1G. The phrases "specific modulating agent", "specifically modulates", etc., are used herein to refer to modulating agents that directly bind to the CSNK1G polypeptide or nucleic acid, and preferably inhibit, enhance, or otherwise alter, the function of the CSNK1G. These

phrases also encompass modulating agents that alter the interaction of the CSNK1G with a binding partner, substrate, or cofactor (e.g. by binding to a binding partner of a CSNK1G, or to a protein/binding partner complex, and altering CSNK1G function). In a further preferred embodiment, the CSNK1G- modulating agent is a modulator of the p21 pathway (e.g. it restores and/or upregulates p21 function) and thus is also a p21-modulating agent.

Preferred CSNK1G-modulating agents include small molecule compounds; CSNK1G-interacting proteins, including antibodies and other biotherapeutics; and nucleic acid modulators such as antisense and RNA inhibitors. The modulating agents may be formulated in pharmaceutical compositions, for example, as compositions that may comprise other active ingredients, as in combination therapy, and/or suitable carriers or excipients. Techniques for formulation and administration of the compounds may be found in "Remington's Pharmaceutical Sciences" Mack Publishing Co., Easton, PA, 19th edition.

Small molecule modulators

Small molecules are often preferred to modulate function of proteins with enzymatic function, and/or containing protein interaction domains. Chemical agents, referred to in the art as "small molecule" compounds are typically organic, non-peptide molecules, having a molecular weight less than 10,000, preferably less than 5,000, more preferably less than 1,000, and most preferably less than 500 daltons. This class of modulators includes chemically synthesized molecules, for instance, compounds from combinatorial chemical libraries. Synthetic compounds may be rationally designed or identified based on known or inferred properties of the CSNK1G protein or may be identified by screening compound libraries. Alternative appropriate modulators of this class are natural products, particularly secondary metabolites from organisms such as plants or fungi, which can also be identified by screening compound libraries for CSNK1G–modulating activity. Methods for generating and obtaining compounds are well known in the art (Schreiber SL, Science (2000) 151: 1964-1969; Radmann J and Gunther J, Science (2000) 151:1947-1948).

Small molecule modulators identified from screening assays, as described below, can be used as lead compounds from which candidate clinical compounds may be designed, optimized, and synthesized. Such clinical compounds may have utility in treating pathologies associated with the p21 pathway. The activity of candidate small molecule modulating agents may be improved several-fold through iterative secondary

functional validation, as further described below, structure determination, and candidate modulator modification and testing. Additionally, candidate clinical compounds are generated with specific regard to clinical and pharmacological properties. For example, the reagents may be derivatized and re-screened using *in vitro* and *in vivo* assays to optimize activity and minimize toxicity for pharmaceutical development.

Protein Modulators

Specific CSNK1G-interacting proteins are useful in a variety of diagnostic and therapeutic applications related to the p21 pathway and related disorders, as well as in validation assays for other CSNK1G-modulating agents. In a preferred embodiment, CSNK1G-interacting proteins affect normal CSNK1G function, including transcription, protein expression, protein localization, and cellular or extra-cellular activity. In another embodiment, CSNK1G-interacting proteins are useful in detecting and providing information about the function of CSNK1G proteins, as is relevant to p21 related disorders, such as cancer (e.g., for diagnostic means).

A CSNK1G-interacting protein may be endogenous, i.e. one that naturally interacts genetically or biochemically with a CSNK1G, such as a member of the CSNK1G pathway that modulates CSNK1G expression, localization, and/or activity. CSNK1G-modulators include dominant negative forms of CSNK1G-interacting proteins and of CSNK1G proteins themselves. Yeast two-hybrid and variant screens offer preferred methods for identifying endogenous CSNK1G-interacting proteins (Finley, R. L. et al. (1996) in DNA Cloning-Expression Systems: A Practical Approach, eds. Glover D. & Hames B. D (Oxford University Press, Oxford, England), pp. 169-203; Fashema SF et al., Gene (2000) 250:1-14; Drees BL Curr Opin Chem Biol (1999) 3:64-70; Vidal M and Legrain P Nucleic Acids Res (1999) 27:919-29; and U.S. Pat. No. 5,928,868). Mass spectrometry is an alternative preferred method for the elucidation of protein complexes (reviewed in, e.g., Pandley A and Mann M, Nature (2000) 405:837-846; Yates JR 3rd, Trends Genet (2000) 16:5-8).

An CSNK1G-interacting protein may be an exogenous protein, such as a CSNK1G-specific antibody or a T-cell antigen receptor (see, e.g., Harlow and Lane (1988) Antibodies, A Laboratory Manual, Cold Spring Harbor Laboratory; Harlow and Lane (1999) Using antibodies: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press). CSNK1G antibodies are further discussed below.

In preferred embodiments, a CSNK1G-interacting protein specifically binds a CSNK1G protein. In alternative preferred embodiments, a CSNK1G-modulating agent binds a CSNK1G substrate, binding partner, or cofactor.

Antibodies

In another embodiment, the protein modulator is a CSNK1G specific antibody agonist or antagonist. The antibodies have therapeutic and diagnostic utilities, and can be used in screening assays to identify CSNK1G modulators. The antibodies can also be used in dissecting the portions of the CSNK1G pathway responsible for various cellular responses and in the general processing and maturation of the CSNK1G.

Antibodies that specifically bind CSNK1G polypeptides can be generated using known methods. Preferably the antibody is specific to a mammalian ortholog of CSNK1G polypeptide, and more preferably, to human CSNK1G. Antibodies may be polyclonal, monoclonal (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab').sub.2 fragments, fragments produced by a FAb expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above. Epitopes of CSNK1G which are particularly antigenic can be selected, for example, by routine screening of CSNK1G polypeptides for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein (Hopp and Wood (1981), Proc. Nati. Acad. Sci. U.S.A. 78:3824-28; Hopp and Wood, (1983) Mol. Immunol. 20:483-89; Sutcliffe et al., (1983) Science 219:660-66) to the amino acid sequence shown in any of SEQ ID NOs:13-16. Monoclonal antibodies with affinities of 10^8 M^{-1} preferably 10^9 M^{-1} to 10^{10} M^{-1}, or stronger can be made by standard procedures as described (Harlow and Lane, *supra*; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed) Academic Press, New York; and U.S. Pat. Nos. 4,381,292; 4,451,570; and 4,618,577). Antibodies may be generated against crude cell extracts of CSNK1G or substantially purified fragments thereof. If CSNK1G fragments are used, they preferably comprise at least 10, and more preferably, at least 20 contiguous amino acids of a CSNK1G protein. In a particular embodiment, CSNK1G-specific antigens and/or immunogens are coupled to carrier proteins that stimulate the immune response. For example, the subject polypeptides are covalently coupled to the keyhole limpet hemocyanin (KLH) carrier, and the conjugate is emulsified in Freund's complete adjuvant, which enhances the immune response. An appropriate immune system such as a laboratory rabbit or mouse is immunized according to conventional protocols.

The presence of CSNK1G-specific antibodies is assayed by an appropriate assay such as a solid phase enzyme-linked immunosorbant assay (ELISA) using immobilized corresponding CSNK1G polypeptides. Other assays, such as radioimmunoassays or fluorescent assays might also be used.

Chimeric antibodies specific to CSNK1G polypeptides can be made that contain different portions from different animal species. For instance, a human immunoglobulin constant region may be linked to a variable region of a murine mAb, such that the antibody derives its biological activity from the human antibody, and its binding specificity from the murine fragment. Chimeric antibodies are produced by splicing together genes that encode the appropriate regions from each species (Morrison et al., Proc. Natl. Acad. Sci. (1984) 81:6851-6855; Neuberger et al., Nature (1984) 312:604-608; Takeda et al., Nature (1985) 31:452-454). Humanized antibodies, which are a form of chimeric antibodies, can be generated by grafting complementary-determining regions (CDRs) (Carlos, T. M., J. M. Harlan. 1994. Blood 84:2068-2101) of mouse antibodies into a background of human framework regions and constant regions by recombinant DNA technology (Riechmann LM, et al., 1988 Nature 323: 323-327). Humanized antibodies contain ~10% murine sequences and ~90% human sequences, and thus further reduce or eliminate immunogenicity, while retaining the antibody specificities (Co MS, and Queen C. 1991 Nature 351: 501-501; Morrison SL. 1992 Ann. Rev. Immun. 10:239-265). Humanized antibodies and methods of their production are well-known in the art (U.S. Pat. Nos. 5,530,101, 5,585,089, 5,693,762, and 6,180,370).

CSNK1G-specific single chain antibodies which are recombinant, single chain polypeptides formed by linking the heavy and light chain fragments of the Fv regions via an amino acid bridge, can be produced by methods known in the art (U.S. Pat. No. 4,946,778; Bird, Science (1988) 242:423-426; Huston et al., Proc. Natl. Acad. Sci. USA (1988) 85:5879-5883; and Ward et al., Nature (1989) 334:544-546).

Other suitable techniques for antibody production involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors (Huse et al., Science (1989) 246:1275-1281). As used herein, T-cell antigen receptors are included within the scope of antibody modulators (Harlow and Lane, 1988, *supra*).

The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, antibodies will be labeled by joining, either covalently or non-covalently, a substance that provides for a detectable signal, or that is toxic to cells

that express the targeted protein (Menard S, et al., Int J. Biol Markers (1989) 4:131-134). A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, fluorescent emitting lanthanide metals, chemiluminescent moieties, bioluminescent moieties, magnetic particles, and the like (U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241). Also, recombinant immunoglobulins may be produced (U.S. Pat. No. 4,816,567). Antibodies to cytoplasmic polypeptides may be delivered and reach their targets by conjugation with membrane-penetrating toxin proteins (U.S. Pat. No. 6,086,900).

When used therapeutically in a patient, the antibodies of the subject invention are typically administered parenterally, when possible at the target site, or intravenously. The therapeutically effective dose and dosage regimen is determined by clinical studies. Typically, the amount of antibody administered is in the range of about 0.1 mg/kg –to about 10 mg/kg of patient weight. For parenteral administration, the antibodies are formulated in a unit dosage injectable form (e.g., solution, suspension, emulsion) in association with a pharmaceutically acceptable vehicle. Such vehicles are inherently nontoxic and non-therapeutic. Examples are water, saline, Ringer's solution, dextrose solution, and 5% human serum albumin. Nonaqueous vehicles such as fixed oils, ethyl oleate, or liposome carriers may also be used. The vehicle may contain minor amounts of additives, such as buffers and preservatives, which enhance isotonicity and chemical stability or otherwise enhance therapeutic potential. The antibodies' concentrations in such vehicles are typically in the range of about 1 mg/ml to about10 mg/ml. Immunotherapeutic methods are further described in the literature (US Pat. No. 5,859,206; WO0073469).

Nucleic Acid Modulators

Other preferred CSNK1G-modulating agents comprise nucleic acid molecules, such as antisense oligomers or double stranded RNA (dsRNA), which generally inhibit CSNK1G activity. Preferred nucleic acid modulators interfere with the function of the CSNK1G nucleic acid such as DNA replication, transcription, translocation of the CSNK1G RNA to the site of protein translation, translation of protein from the CSNK1G RNA, splicing of the CSNK1G RNA to yield one or more mRNA species, or catalytic activity which may be engaged in or facilitated by the CSNK1G RNA.

In one embodiment, the antisense oligomer is an oligonucleotide that is sufficiently complementary to a CSNK1G mRNA to bind to and prevent translation, preferably by binding to the 5´ untranslated region. CSNK1G-specific antisense oligonucleotides, preferably range from at least 6 to about 200 nucleotides. In some embodiments the oligonucleotide is preferably at least 10, 15, or 20 nucleotides in length. In other embodiments, the oligonucleotide is preferably less than 50, 40, or 30 nucleotides in length. The oligonucleotide can be DNA or RNA or a chimeric mixture or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone. The oligonucleotide may include other appending groups such as peptides, agents that facilitate transport across the cell membrane, hybridization-triggered cleavage agents, and intercalating agents.

In another embodiment, the antisense oligomer is a phosphothioate morpholino oligomer (PMO). PMOs are assembled from four different morpholino subunits, each of which contain one of four genetic bases (A, C, G, or T) linked to a six-membered morpholine ring. Polymers of these subunits are joined by non-ionic phosphodiamidate intersubunit linkages. Details of how to make and use PMOs and other antisense oligomers are well known in the art (e.g. see WO99/18193; Probst JC, Antisense Oligodeoxynucleotide and Ribozyme Design, Methods. (2000) 22(3):271-281; Summerton J, and Weller D. 1997 Antisense Nucleic Acid Drug Dev. :7:187-95; US Pat. No. 5,235,033; and US Pat No. 5,378,841).

Alternative preferred CSNK1G nucleic acid modulators are double-stranded RNA species mediating RNA interference (RNAi). RNAi is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. Methods relating to the use of RNAi to silence genes in *C. elegans*, *Drosophila*, plants, and humans are known in the art (Fire A, et al., 1998 Nature 391:806-811; Fire, A. Trends Genet. 15, 358-363 (1999); Sharp, P. A. RNA interference 2001. Genes Dev. 15, 485-490 (2001); Hammond, S. M., et al., Nature Rev. Genet. 2, 110-1119 (2001); Tuschl, T. Chem. Biochem. 2, 239-245 (2001); Hamilton, A. et al., Science 286, 950-952 (1999); Hammond, S. M., et al., Nature 404, 293-296 (2000); Zamore, P. D., et al., Cell 101, 25-33 (2000); Bernstein, E., et al., Nature 409, 363-366 (2001); Elbashir, S. M., et al., Genes Dev. 15, 188-200 (2001); WO0129058; WO9932619; Elbashir SM, et al., 2001 Nature 411:494-498).

Nucleic acid modulators are commonly used as research reagents, diagnostics, and therapeutics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used to elucidate the function of particular genes (see, for example, U.S. Pat. No. 6,165,790). Nucleic acid modulators are also used, for example, to distinguish between functions of various members of a biological pathway. For example, antisense oligomers have been employed as therapeutic moieties in the treatment of disease states in animals and man and have been demonstrated in numerous clinical trials to be safe and effective (Milligan JF, *et al*, Current Concepts in Antisense Drug Design, J Med Chem. (1993) 36:1923-1937; Tonkinson JL *et al.*, Antisense Oligodeoxynucleotides as Clinical Therapeutic Agents, Cancer Invest. (1996) 14:54-65). Accordingly, in one aspect of the invention, a CSNK1G-specific nucleic acid modulator is used in an assay to further elucidate the role of the CSNK1G in the p21 pathway, and/or its relationship to other members of the pathway. In another aspect of the invention, a CSNK1G-specific antisense oligomer is used as a therapeutic agent for treatment of p21-related disease states.

Assay Systems

The invention provides assay systems and screening methods for identifying specific modulators of CSNK1G activity. As used herein, an "assay system" encompasses all the components required for performing and analyzing results of an assay that detects and/or measures a particular event. In general, primary assays are used to identify or confirm a modulator's specific biochemical or molecular effect with respect to the CSNK1G nucleic acid or protein. In general, secondary assays further assess the activity of a CSNK1G modulating agent identified by a primary assay and may confirm that the modulating agent affects CSNK1G in a manner relevant to the p21 pathway. In some cases, CSNK1G modulators will be directly tested in a secondary assay.

In a preferred embodiment, the screening method comprises contacting a suitable assay system comprising a CSNK1G polypeptide or nucleic acid with a candidate agent under conditions whereby, but for the presence of the agent, the system provides a reference activity (e.g. kinase activity), which is based on the particular molecular event the screening method detects. A statistically significant difference between the agent-biased activity and the reference activity indicates that the candidate agent modulates CSNK1G activity, and hence the p21 pathway. The CSNK1G polypeptide or nucleic acid used in the assay may comprise any of the nucleic acids or polypeptides described above.

Primary Assays

The type of modulator tested generally determines the type of primary assay.

Primary assays for small molecule modulators

For small molecule modulators, screening assays are used to identify candidate modulators. Screening assays may be cell-based or may use a cell-free system that recreates or retains the relevant biochemical reaction of the target protein (reviewed in Sittampalam GS *et al.*, Curr Opin Chem Biol (1997) 1:384-91 and accompanying references). As used herein the term "cell-based" refers to assays using live cells, dead cells, or a particular cellular fraction, such as a membrane, endoplasmic reticulum, or mitochondrial fraction. The term "cell free" encompasses assays using substantially purified protein (either endogenous or recombinantly produced), partially purified or crude cellular extracts. Screening assays may detect a variety of molecular events, including protein-DNA interactions, protein-protein interactions (*e.g.*, receptor-ligand binding), transcriptional activity (*e.g.*, using a reporter gene), enzymatic activity (*e.g.*, via a property of the substrate), activity of second messengers, immunogenicty and changes in cellular morphology or other cellular characteristics. Appropriate screening assays may use a wide range of detection methods including fluorescent, radioactive, colorimetric, spectrophotometric, and amperometric methods, to provide a read-out for the particular molecular event detected.

Cell-based screening assays usually require systems for recombinant expression of CSNK1G and any auxiliary proteins demanded by the particular assay. Appropriate methods for generating recombinant proteins produce sufficient quantities of proteins that retain their relevant biological activities and are of sufficient purity to optimize activity and assure assay reproducibility. Yeast two-hybrid and variant screens, and mass spectrometry provide preferred methods for determining protein-protein interactions and elucidation of protein complexes. In certain applications, when CSNK1G-interacting proteins are used in screens to identify small molecule modulators, the binding specificity of the interacting protein to the CSNK1G protein may be assayed by various known methods such as substrate processing (e.g. ability of the candidate CSNK1G-specific binding agents to function as negative effectors in CSNK1G-expressing cells), binding equilibrium constants (usually at least about 10^7 M^{-1}, preferably at least about 10^8 M^{-1}, more preferably at least about 10^9 M^{-1}), and immunogenicity (e.g. ability to elicit CSNK1G specific antibody in a heterologous host such as a mouse, rat, goat or rabbit).

For enzymes and receptors, binding may be assayed by, respectively, substrate and ligand processing.

The screening assay may measure a candidate agent's ability to specifically bind to or modulate activity of a CSNK1G polypeptide, a fusion protein thereof, or to cells or membranes bearing the polypeptide or fusion protein. The CSNK1G polypeptide can be full length or a fragment thereof that retains functional CSNK1G activity. The CSNK1G polypeptide may be fused to another polypeptide, such as a peptide tag for detection or anchoring, or to another tag. The CSNK1G polypeptide is preferably human CSNK1G, or is an ortholog or derivative thereof as described above. In a preferred embodiment, the screening assay detects candidate agent-based modulation of CSNK1G interaction with a binding target, such as an endogenous or exogenous protein or other substrate that has CSNK1G –specific binding activity, and can be used to assess normal CSNK1G gene function.

Suitable assay formats that may be adapted to screen for CSNK1G modulators are known in the art. Preferred screening assays are high throughput or ultra high throughput and thus provide automated, cost-effective means of screening compound libraries for lead compounds (Fernandes PB, Curr Opin Chem Biol (1998) 2:597-603; Sundberg SA, Curr Opin Biotechnol 2000, 11:47-53). In one preferred embodiment, screening assays uses fluorescence technologies, including fluorescence polarization, time-resolved fluorescence, and fluorescence resonance energy transfer. These systems offer means to monitor protein-protein or DNA-protein interactions in which the intensity of the signal emitted from dye-labeled molecules depends upon their interactions with partner molecules (*e.g.,* Selvin PR, Nat Struct Biol (2000) 7:730-4; Fernandes PB, *supra*; Hertzberg RP and Pope AJ, Curr Opin Chem Biol (2000) 4:445-451).

A variety of suitable assay systems may be used to identify candidate CSNK1G and p21 pathway modulators (e.g. U.S. Pat. No. 6,165,992 (kinase assays); U.S. Pat. Nos. 5,550,019 and 6,133,437 (apoptosis assays); and U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434 (angiogenesis assays), among others). Specific preferred assays are described in more detail below.

Kinase assays. In some preferred embodiments the screening assay detects the ability of the test agent to modulate the kinase activity of a CSNK1G polypeptide. In further embodiments, a cell-free kinase assay system is used to identify a candidate p21 modulating agent, and a secondary, cell-based assay, such as an apoptosis or hypoxic

induction assay (described below), may be used to further characterize the candidate p21 modulating agent. Many different assays for kinases have been reported in the literature and are well known to those skilled in the art (e.g. U.S. Pat. No. 6,165,992; Zhu et al., Nature Genetics (2000) 26:283-289; and WO0073469). Radioassays, which monitor the transfer of a gamma phosphate are frequently used. For instance, a scintillation assay for p56 (lck) kinase activity monitors the transfer of the gamma phosphate from gamma -^{33}P ATP to a biotinylated peptide substrate; the substrate is captured on a streptavidin coated bead that transmits the signal (Beveridge M *et al.*, J Biomol Screen (2000) 5:205-212). This assay uses the scintillation proximity assay (SPA), in which only radio-ligand bound to receptors tethered to the surface of an SPA bead are detected by the scintillant immobilized within it, allowing binding to be measured without separation of bound from free ligand.

Other assays for protein kinase activity may use antibodies that specifically recognize phosphorylated substrates. For instance, the kinase receptor activation (KIRA) assay measures receptor tyrosine kinase activity by ligand stimulating the intact receptor in cultured cells, then capturing solubilized receptor with specific antibodies and quantifying phosphorylation via phosphotyrosine ELISA (Sadick MD, Dev Biol Stand (1999) 97:121-133).

Another example of antibody based assays for protein kinase activity is TRF (time-resolved fluorometry). This method utilizes europium chelate-labeled anti-phosphotyrosine antibodies to detect phosphate transfer to a polymeric substrate coated onto microtiter plate wells. The amount of phosphorylation is then detected using time-resolved, dissociation-enhanced fluorescence (Braunwalder AF, et al., Anal Biochem 1996 Jul 1;238(2):159-64).

Apoptosis assays. Assays for apoptosis may be performed by terminal deoxynucleotidyl transferase-mediated digoxigenin-11-dUTP nick end labeling (TUNEL) assay. The TUNEL assay is used to measure nuclear DNA fragmentation characteristic of apoptosis (Lazebnik *et al.*, 1994, Nature 371, 346), by following the incorporation of fluorescein-dUTP (Yonehara *et al.*, 1989, J. Exp. Med. 169, 1747). Apoptosis may further be assayed by acridine orange staining of tissue culture cells (Lucas, R., et al., 1998, Blood 15:4730-41). Other cell-based apoptosis assays include the caspase-3/7 assay and the cell death nucleosome ELISA assay. The caspase 3/7 assay is based on the activation of the caspase cleavage activity as part of a cascade of events that occur during programmed cell

death in many apoptotic pathways. In the caspase 3/7 assay (commercially available Apo-ONE™ Homogeneous Caspase-3/7 assay from Promega, cat# 67790), lysis buffer and caspase substrate are mixed and added to cells. The caspase substrate becomes fluorescent when cleaved by active caspase 3/7. The nucleosome ELISA assay is a general cell death assay known to those skilled in the art, and available commercially (Roche, Cat# 1774425). This assay is a quantitative sandwich-enzyme-immunoassay which uses monoclonal antibodies directed against DNA and histones respectively, thus specifically determining amount of mono- and oligonucleosomes in the cytoplasmic fraction of cell lysates. Mono and oligonucleosomes are enriched in the cytoplasm during apoptosis due to the fact that DNA fragmentation occurs several hours before the plasma membrane breaks down, allowing for accumalation in the cytoplasm. Nucleosomes are not present in the cytoplasmic fraction of cells that are not undergoing apoptosis. An apoptosis assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the apoptosis assay system and changes in induction of apoptosis relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, an apoptosis assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using a cell-free assay system. An apoptosis assay may also be used to test whether CSNK1G function plays a direct role in apoptosis. For example, an apoptosis assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in apoptotic response compared to wild type cells suggests that the CSNK1G plays a direct role in the apoptotic response. Apoptosis assays are described further in US Pat. No. 6,133,437.

Cell proliferation and cell cycle assays. Cell proliferation may be assayed via bromodeoxyuridine (BRDU) incorporation. This assay identifies a cell population undergoing DNA synthesis by incorporation of BRDU into newly-synthesized DNA. Newly-synthesized DNA may then be detected using an anti-BRDU antibody (Hoshino *et al.*, 1986, Int. J. Cancer 38, 369; Campana *et al.*, 1988, J. Immunol. Meth. 107, 79), or by other means.

Cell proliferation is also assayed via phospho-histone H3 staining, which identifies a cell population undergoing mitosis by phosphorylation of histone H3. Phosphorylation of histone H3 at serine 10 is detected using an antibody specfic to the phosphorylated form

of the serine 10 residue of histone H3. (Chadlee,D.N. 1995, J. Biol. Chem 270:20098-105). Cell Proliferation may also be examined using [^{3}H]-thymidine incorporation (Chen, J., 1996, Oncogene 13:1395-403; Jeoung, J., 1995, J. Biol. Chem. 270:18367-73). This assay allows for quantitative characterization of S-phase DNA syntheses. In this assay, cells synthesizing DNA will incorporate [^{3}H]-thymidine into newly synthesized DNA. Incorporation can then be measured by standard techniques such as by counting of radioisotope in a scintillation counter (e.g., Beckman LS 3800 Liquid Scintillation Counter). Another proliferation assay uses the dye Alamar Blue (available from Biosource International), which fluoresces when reduced in living cells and provides an indirect measurement of cell number (Voytik-Harbin SL et al., 1998, In Vitro Cell Dev Biol Anim 34:239-46). Yet another proliferation assay, the MTS assay, is based on in vitro cytotoxicity assessment of industrial chemicals, and uses the soluble tetrazolium salt, MTS. MTS assays are commercially available, for example, the Promega CellTiter 96® AQueous Non-Radioactive Cell Proliferation Assay (Cat.# G5421).

Cell proliferation may also be assayed by colony formation in soft agar (Sambrook et al., Molecular Cloning, Cold Spring Harbor (1989)). For example, cells transformed with CSNK1G are seeded in soft agar plates, and colonies are measured and counted after two weeks incubation.

Cell proliferation may also be assayed by measuring ATP levels as indicator of metabolically active cells. Such assays are commercially available, for example Cell Titer-Glo™, which is a luminescent homogeneous assay available from Promega.

Involvement of a gene in the cell cycle may be assayed by flow cytometry (Gray JW et al. (1986) Int J Radiat Biol Relat Stud Phys Chem Med 49:237-55). Cells transfected with a CSNK1G may be stained with propidium iodide and evaluated in a flow cytometer (available from Becton Dickinson), which indicates accumulation of cells in different stages of the cell cycle.

Accordingly, a cell proliferation or cell cycle assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the assay system and changes in cell proliferation or cell cycle relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the cell proliferation or cell cycle assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system such as a cell-free assay system. A cell proliferation assay may also be used to test

whether CSNK1G function plays a direct role in cell proliferation or cell cycle. For example, a cell proliferation or cell cycle assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in proliferation or cell cycle compared to wild type cells suggests that the CSNK1G plays a direct role in cell proliferation or cell cycle.

Angiogenesis. Angiogenesis may be assayed using various human endothelial cell systems, such as umbilical vein, coronary artery, or dermal cells. Suitable assays include Alamar Blue based assays (available from Biosource International) to measure proliferation; migration assays using fluorescent molecules, such as the use of Becton Dickinson Falcon HTS FluoroBlock cell culture inserts to measure migration of cells through membranes in presence or absence of angiogenesis enhancer or suppressors; and tubule formation assays based on the formation of tubular structures by endothelial cells on Matrigel® (Becton Dickinson). Accordingly, an angiogenesis assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the angiogenesis assay system and changes in angiogenesis relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the angiogenesis assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system. An angiogenesis assay may also be used to test whether CSNK1G function plays a direct role in cell proliferation. For example, an angiogenesis assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in angiogenesis compared to wild type cells suggests that the CSNK1G plays a direct role in angiogenesis. U.S. Pat. Nos. 5,976,782, 6,225,118 and 6,444,434, among others, describe various angiogenesis assays.

Hypoxic induction. The alpha subunit of the transcription factor, hypoxia inducible factor-1 (HIF-1), is upregulated in tumor cells following exposure to hypoxia in vitro. Under hypoxic conditions, HIF-1 stimulates the expression of genes known to be important in tumour cell survival, such as those encoding glyolytic enzymes and VEGF. Induction of such genes by hypoxic conditions may be assayed by growing cells transfected with CSNK1G in hypoxic conditions (such as with 0.1% O2, 5% CO2, and balance N2, generated in a Napco 7001 incubator (Precision Scientific)) and normoxic

conditions, followed by assessment of gene activity or expression by Taqman®. For example, a hypoxic induction assay system may comprise a cell that expresses a CSNK1G, and that optionally has defective p21 function (e.g. p21 is over-expressed or under-expressed relative to wild-type cells). A test agent can be added to the hypoxic induction assay system and changes in hypoxic response relative to controls where no test agent is added, identify candidate p21 modulating agents. In some embodiments of the invention, the hypoxic induction assay may be used as a secondary assay to test a candidate p21 modulating agents that is initially identified using another assay system. A hypoxic induction assay may also be used to test whether CSNK1G function plays a direct role in the hypoxic response. For example, a hypoxic induction assay may be performed on cells that over- or under-express CSNK1G relative to wild type cells. Differences in hypoxic response compared to wild type cells suggests that the CSNK1G plays a direct role in hypoxic induction.

Cell adhesion. Cell adhesion assays measure adhesion of cells to purified adhesion proteins, or adhesion of cells to each other, in presence or absence of candidate modulating agents. Cell-protein adhesion assays measure the ability of agents to modulate the adhesion of cells to purified proteins. For example, recombinant proteins are produced, diluted to 2.5g/mL in PBS, and used to coat the wells of a microtiter plate. The wells used for negative control are not coated. Coated wells are then washed, blocked with 1% BSA, and washed again. Compounds are diluted to 2× final test concentration and added to the blocked, coated wells. Cells are then added to the wells, and the unbound cells are washed off. Retained cells are labeled directly on the plate by adding a membrane-permeable fluorescent dye, such as calcein-AM, and the signal is quantified in a fluorescent microplate reader.

Cell-cell adhesion assays measure the ability of agents to modulate binding of cell adhesion proteins with their native ligands. These assays use cells that naturally or recombinantly express the adhesion protein of choice. In an exemplary assay, cells expressing the cell adhesion protein are plated in wells of a multiwell plate. Cells expressing the ligand are labeled with a membrane-permeable fluorescent dye, such as BCECF , and allowed to adhere to the monolayers in the presence of candidate agents. Unbound cells are washed off, and bound cells are detected using a fluorescence plate reader.

High-throughput cell adhesion assays have also been described. In one such assay, small molecule ligands and peptides are bound to the surface of microscope slides using a microarray spotter, intact cells are then contacted with the slides, and unbound cells are washed off. In this assay, not only the binding specificity of the peptides and modulators against cell lines are determined, but also the functional cell signaling of attached cells using immunofluorescence techniques in situ on the microchip is measured (Falsey JR et al., Bioconjug Chem. 2001 May-Jun;12(3):346-53).

Tubulogenesis. Tubulogenesis assays monitor the ability of cultured cells, generally endothelial cells, to form tubular structures on a matrix substrate, which generally simulates the environment of the extracellular matrix. Exemplary substrates include Matrigel™ (Becton Dickinson), an extract of basement membrane proteins containing laminin, collagen IV, and heparin sulfate proteoglycan, which is liquid at 4° C and forms a solid gel at 37° C. Other suitable matrices comprise extracellular components such as collagen, fibronectin, and/or fibrin. Cells are stimulated with a pro-angiogenic stimulant, and their ability to form tubules is detected by imaging. Tubules can generally be detected after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Tube formation assays are well known in the art (e.g., Jones MK et al., 1999, Nature Medicine 5:1418-1423). These assays have traditionally involved stimulation with serum or with the growth factors FGF or VEGF. Serum represents an undefined source of growth factors. In a preferred embodiment, the assay is performed with cells cultured in serum free medium, in order to control which process or pathway a candidate agent modulates. Moreover, we have found that different target genes respond differently to stimulation with different pro-angiogenic agents, including inflammatory angiogenic factors such as TNF-alpa. Thus, in a further preferred embodiment, a tubulogenesis assay system comprises testing a CSNK1G's response to a variety of factors, such as FGF, VEGF, phorbol myristate acetate (PMA), TNF-alpha, ephrin, etc.

Cell Migration. An invasion/migration assay (also called a migration assay) tests the ability of cells to overcome a physical barrier and to migrate towards pro-angiogenic signals. Migration assays are known in the art (e.g., Paik JH et al., 2001, J Biol Chem 276:11830-11837). In a typical experimental set-up, cultured endothelial cells are seeded onto a matrix-coated porous lamina, with pore sizes generally smaller than typical cell size. The matrix generally simulates the environment of the extracellular matrix, as

described above. The lamina is typically a membrane, such as the transwell polycarbonate membrane (Corning Costar Corporation, Cambridge, MA), and is generally part of an upper chamber that is in fluid contact with a lower chamber containing pro-angiogenic stimuli. Migration is generally assayed after an overnight incubation with stimuli, but longer or shorter time frames may also be used. Migration is assessed as the number of cells that crossed the lamina, and may be detected by staining cells with hemotoxylin solution (VWR Scientific, South San Francisco, CA), or by any other method for determining cell number. In another exemplary set up, cells are fluorescently labeled and migration is detected using fluorescent readings, for instance using the Falcon HTS FluoroBlok (Becton Dickinson). While some migration is observed in the absence of stimulus, migration is greatly increased in response to pro-angiogenic factors. As described above, a preferred assay system for migration/invasion assays comprises testing a CSNK1G's response to a variety of pro-angiogenic factors, including tumor angiogenic and inflammatory angiogenic agents, and culturing the cells in serum free medium.

Sprouting assay. A sprouting assay is a three-dimensional *in vitro* angiogenesis assay that uses a cell-number defined spheroid aggregation of endothelial cells ("spheroid"), embedded in a collagen gel-based matrix. The spheroid can serve as a starting point for the sprouting of capillary-like structures by invasion into the extracellular matrix (termed "cell sprouting") and the subsequent formation of complex anastomosing networks (Korff and Augustin, 1999, J Cell Sci 112:3249-58). In an exemplary experimental set-up, spheroids are prepared by pipetting 400 human umbilical vein endothelial cells into individual wells of a nonadhesive 96-well plates to allow overnight spheroidal aggregation (Korff and Augustin: J Cell Biol 143: 1341-52, 1998). Spheroids are harvested and seeded in 900μl of methocel-collagen solution and pipetted into individual wells of a 24 well plate to allow collagen gel polymerization. Test agents are added after 30 min by pipetting 100 μl of 10-fold concentrated working dilution of the test substances on top of the gel. Plates are incubated at 37°C for 24h. Dishes are fixed at the end of the experimental incubation period by addition of paraformaldehyde. Sprouting intensity of endothelial cells can be quantitated by an automated image analysis system to determine the cumulative sprout length per spheroid.

Primary assays for antibody modulators

For antibody modulators, appropriate primary assays test is a binding assay that tests the antibody's affinity to and specificity for the CSNK1G protein. Methods for testing antibody affinity and specificity are well known in the art (Harlow and Lane, 1988, 1999, *supra*). The enzyme-linked immunosorbant assay (ELISA) is a preferred method for detecting CSNK1G-specific antibodies; others include FACS assays, radioimmunoassays, and fluorescent assays.

In some cases, screening assays described for small molecule modulators may also be used to test antibody modulators.

Primary assays for nucleic acid modulators

For nucleic acid modulators, primary assays may test the ability of the nucleic acid modulator to inhibit or enhance CSNK1G gene expression, preferably mRNA expression. In general, expression analysis comprises comparing CSNK1G expression in like populations of cells (*e.g.*, two pools of cells that endogenously or recombinantly express CSNK1G) in the presence and absence of the nucleic acid modulator. Methods for analyzing mRNA and protein expression are well known in the art. For instance, Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR (*e.g.*, using the TaqMan®, PE Applied Biosystems), or microarray analysis may be used to confirm that CSNK1G mRNA expression is reduced in cells treated with the nucleic acid modulator (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, *eds.*, John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm DH and Guiseppi-Elie, A Curr Opin Biotechnol 2001, 12:41-47). Protein expression may also be monitored. Proteins are most commonly detected with specific antibodies or antisera directed against either the CSNK1G protein or specific peptides. A variety of means including Western blotting, ELISA, or in situ detection, are available (Harlow E and Lane D, 1988 and 1999, *supra*).

In some cases, screening assays described for small molecule modulators, particularly in assay systems that involve CSNK1G mRNA expression, may also be used to test nucleic acid modulators.

Secondary Assays

Secondary assays may be used to further assess the activity of CSNK1G-modulating agent identified by any of the above methods to confirm that the modulating

agent affects CSNK1G in a manner relevant to the p21 pathway. As used herein, CSNK1G-modulating agents encompass candidate clinical compounds or other agents derived from previously identified modulating agent. Secondary assays can also be used to test the activity of a modulating agent on a particular genetic or biochemical pathway or to test the specificity of the modulating agent's interaction with CSNK1G.

Secondary assays generally compare like populations of cells or animals (*e.g.*, two pools of cells or animals that endogenously or recombinantly express CSNK1G) in the presence and absence of the candidate modulator. In general, such assays test whether treatment of cells or animals with a candidate CSNK1G–modulating agent results in changes in the p21 pathway in comparison to untreated (or mock- or placebo-treated) cells or animals. Certain assays use "sensitized genetic backgrounds", which, as used herein, describe cells or animals engineered for altered expression of genes in the p21 or interacting pathways.

Cell-based assays

Cell based assays may use a variety of mammalian cell lines known to have defective p21 function such as HCT116 colon cancer cells, among others, available from American Type Culture Collection (ATCC), Manassas, VA). Cell based assays may detect endogenous p21 pathway activity or may rely on recombinant expression of p21 pathway components. Any of the aforementioned assays may be used in this cell-based format. Candidate modulators are typically added to the cell media but may also be injected into cells or delivered by any other efficacious means.

Animal Assays

A variety of non-human animal models of normal or defective p21 pathway may be used to test candidate CSNK1G modulators. Models for defective p21 pathway typically use genetically modified animals that have been engineered to mis-express (*e.g.*, over-express or lack expression in) genes involved in the p21 pathway. Assays generally require systemic delivery of the candidate modulators, such as by oral administration, injection, etc.

In a preferred embodiment, p21 pathway activity is assessed by monitoring neovascularization and angiogenesis. Animal models with defective and normal p21 are used to test the candidate modulator's affect on CSNK1G in Matrigel® assays. Matrigel® is an extract of basement membrane proteins, and is composed primarily of laminin,

collagen IV, and heparin sulfate proteoglycan. It is provided as a sterile liquid at 4° C, but rapidly forms a solid gel at 37° C. Liquid Matrigel® is mixed with various angiogenic agents, such as bFGF and VEGF, or with human tumor cells which over-express the CSNK1G. The mixture is then injected subcutaneously(SC) into female athymic nude mice (Taconic, Germantown, NY) to support an intense vascular response. Mice with Matrigel® pellets may be dosed via oral (PO), intraperitoneal (IP), or intravenous (IV) routes with the candidate modulator. Mice are euthanized 5 - 12 days post-injection, and the Matrigel® pellet is harvested for hemoglobin analysis (Sigma plasma hemoglobin kit). Hemoglobin content of the gel is found to correlate the degree of neovascularization in the gel.

In another preferred embodiment, the effect of the candidate modulator on CSNK1G is assessed via tumorigenicity assays. Tumor xenograft assays are known in the art (see, e.g., Ogawa K et al., 2000, Oncogene 19:6043-6052). Xenografts are typically implanted SC into female athymic mice, 6-7 week old, as single cell suspensions either from a pre-existing tumor or from *in vitro* culture. The tumors which express the CSNK1G endogenously are injected in the flank, 1 x 10^5 to 1 x 10^7 cells per mouse in a volume of 100 µL using a 27gauge needle. Mice are then ear tagged and tumors are measured twice weekly. Candidate modulator treatment is initiated on the day the mean tumor weight reaches 100 mg. Candidate modulator is delivered IV, SC, IP, or PO by bolus administration. Depending upon the pharmacokinetics of each unique candidate modulator, dosing can be performed multiple times per day. The tumor weight is assessed by measuring perpendicular diameters with a caliper and calculated by multiplying the measurements of diameters in two dimensions. At the end of the experiment, the excised tumors maybe utilized for biomarker identification or further analyses. For immunohistochemistry staining, xenograft tumors are fixed in 4% paraformaldehyde, 0.1M phosphate, pH 7.2, for 6 hours at 4°C, immersed in 30% sucrose in PBS, and rapidly frozen in isopentane cooled with liquid nitrogen.

In another preferred embodiment, tumorogenicity is monitored using a hollow fiber assay, which is described in U.S. Pat No. US 5,698,413. Briefly, the method comprises implanting into a laboratory animal a biocompatible, semi-permeable encapsulation device containing target cells, treating the laboratory animal with a candidate modulating agent, and evaluating the target cells for reaction to the candidate modulator. Implanted cells are generally human cells from a pre-existing tumor or a tumor cell line. After an appropriate period of time, generally around six days, the implanted samples are harvested for

evaluation of the candidate modulator. Tumorogenicity and modulator efficacy may be evaluated by assaying the quantity of viable cells present in the macrocapsule, which can be determined by tests known in the art, for example, MTT dye conversion assay, neutral red dye uptake, trypan blue staining, viable cell counts, the number of colonies formed in soft agar, the capacity of the cells to recover and replicate in vitro, etc.

In another preferred embodiment, a tumorogenicity assay use a transgenic animal, usually a mouse, carrying a dominant oncogene or tumor suppressor gene knockout under the control of tissue specific regulatory sequences; these assays are generally referred to as transgenic tumor assays. In a preferred application, tumor development in the transgenic model is well characterized or is controlled. In an exemplary model, the "RIP1-Tag2" transgene, comprising the SV40 large T-antigen oncogene under control of the insulin gene regulatory regions is expressed in pancreatic beta cells and results in islet cell carcinomas (Hanahan D, 1985, Nature 315:115-122; Parangi S et al, 1996, Proc Natl Acad Sci USA 93: 2002-2007; Bergers G et al, 1999, Science 284:808-812). An "angiogenic switch," occurs at approximately five weeks, as normally quiescent capillaries in a subset of hyperproliferative islets become angiogenic. The RIP1-TAG2 mice die by age 14 weeks. Candidate modulators may be administered at a variety of stages, including just prior to the angiogenic switch (e.g., for a model of tumor prevention), during the growth of small tumors (e.g., for a model of intervention), or during the growth of large and/or invasive tumors (e.g., for a model of regression). Tumorogenicity and modulator efficacy can be evaluating life-span extension and/or tumor characteristics, including number of tumors, tumor size, tumor morphology, vessel density, apoptotic index, etc.

Diagnostic and therapeutic uses

Specific CSNK1G-modulating agents are useful in a variety of diagnostic and therapeutic applications where disease or disease prognosis is related to defects in the p21 pathway, such as angiogenic, apoptotic, or cell proliferation disorders. Accordingly, the invention also provides methods for modulating the p21 pathway in a cell, preferably a cell pre-determined to have defective or impaired p21 function (e.g. due to overexpression, underexpression, or misexpression of p21, or due to gene mutations), comprising the step of administering an agent to the cell that specifically modulates CSNK1G activity. Preferably, the modulating agent produces a detectable phenotypic change in the cell indicating that the p21 function is restored. The phrase "function is restored", and equivalents, as used herein, means that the desired phenotype is achieved, or is brought

closer to normal compared to untreated cells. For example, with restored p21 function, cell proliferation and/or progression through cell cycle may normalize, or be brought closer to normal relative to untreated cells. The invention also provides methods for treating disorders or disease associated with impaired p21 function by administering a therapeutically effective amount of a CSNK1G -modulating agent that modulates the p21 pathway. The invention further provides methods for modulating CSNK1G function in a cell, preferably a cell pre-determined to have defective or impaired CSNK1G function, by administering a CSNK1G -modulating agent. Additionally, the invention provides a method for treating disorders or disease associated with impaired CSNK1G function by administering a therapeutically effective amount of a CSNK1G -modulating agent.

The discovery that CSNK1G is implicated in p21 pathway provides for a variety of methods that can be employed for the diagnostic and prognostic evaluation of diseases and disorders involving defects in the p21 pathway and for the identification of subjects having a predisposition to such diseases and disorders.

Various expression analysis methods can be used to diagnose whether CSNK1G expression occurs in a particular sample, including Northern blotting, slot blotting, ribonuclease protection, quantitative RT-PCR, and microarray analysis. (*e.g.*, Current Protocols in Molecular Biology (1994) Ausubel FM *et al.*, *eds.*, John Wiley & Sons, Inc., chapter 4; Freeman WM *et al.*, Biotechniques (1999) 26:112-125; Kallioniemi OP, Ann Med 2001, 33:142-147; Blohm and Guiseppi-Elie, Curr Opin Biotechnol 2001, 12:41-47). Tissues having a disease or disorder implicating defective p21 signaling that express a CSNK1G, are identified as amenable to treatment with a CSNK1G modulating agent. In a preferred application, the p21 defective tissue overexpresses a CSNK1G relative to normal tissue. For example, a Northern blot analysis of mRNA from tumor and normal cell lines, or from tumor and matching normal tissue samples from the same patient, using full or partial CSNK1G cDNA sequences as probes, can determine whether particular tumors express or overexpress CSNK1G. Alternatively, the TaqMan® is used for quantitative RT-PCR analysis of CSNK1G expression in cell lines, normal tissues and tumor samples (PE Applied Biosystems).

Various other diagnostic methods may be performed, for example, utilizing reagents such as the CSNK1G oligonucleotides, and antibodies directed against a CSNK1G, as described above for: (1) the detection of the presence of CSNK1G gene mutations, or the detection of either over- or under-expression of CSNK1G mRNA relative to the non-disorder state; (2) the detection of either an over- or an under-

abundance of CSNK1G gene product relative to the non-disorder state; and (3) the detection of perturbations or abnormalities in the signal transduction pathway mediated by CSNK1G.

Thus, in a specific embodiment, the invention is drawn to a method for diagnosing a disease or disorder in a patient that is associated with alterations in CSNK1G expression, the method comprising: a) obtaining a biological sample from the patient; b) contacting the sample with a probe for CSNK1G expression; c) comparing results from step (b) with a control; and d) determining whether step (c) indicates a likelihood of the disease or disorder. Preferably, the disease is cancer, most preferably a cancer as shown in TABLE 1. The probe may be either DNA or protein, including an antibody.

EXAMPLES

The following experimental section and examples are offered by way of illustration and not by way of limitation.

I. Drosophila p21 screen

A dominant loss of function screen was carried out in *Drosophila* to identify genes that interact with the cyclin dependent kinase inhibitor, p21 (Bourne HR, et al., Nature (1990) 348(6297):125-132; Marshall CJ, Trends Genet (1991) 7(3):91-95). Expression of the p21 gene from GMR-p21 transgene (Hay, B. A., et al. (1994) Development120:2121-2129) in the eye causes deterioration of normal eye morphology, resulting in reduced, rough eyes. Flies carrying this transgene were maintained as a stock (P 1025 F, genotype: y w; P{p21-pExp-gl-w[+]Hsp70(3'UTR)-5}). Females of this stock were crossed to a collection of males carrying piggyBac insertions (Fraser M *et al.*, Virology (1985) 145:356-361). Resulting progeny carrying both the transgene and transposons were scored for the effect of the transposon on the eye phenotype, i.e. whether the transposon enhanced or suppressed (or had no effect) the eye phenotype. All data was recorded and all modifiers were retested with a repeat of the original cross, and the retests were scored at least twice. Modifiers of the eye phenotype were identified as members of the p21 pathway. GISH (CG6963) was an enhancer of the eye phenotype. Orthologs of the modifiers are referred to herein as CSNK1G.

BLAST analysis (Altschul et al., *supra*) was employed to identify orthologs of Drosophila modifiers.

Various domains, signals, and functional subunits in proteins were analyzed using the PSORT (Nakai K., and Horton P., Trends Biochem Sci, 1999, 24:34-6; Kenta Nakai, Protein sorting signals and prediction of subcellular localization, Adv. Protein Chem. 54, 277-344 (2000)), PFAM (Bateman A., et al., Nucleic Acids Res, 1999, 27:260-2), SMART (Ponting CP, et al., SMART: identification and annotation of domains from signaling and extracellular protein sequences. Nucleic Acids Res. 1999 Jan 1;27(1):229-32), TM-HMM (Erik L.L. Sonnhammer, Gunnar von Heijne, and Anders Krogh: A hidden Markov model for predicting transmembrane helices in protein sequences. In Proc. of Sixth Int. Conf. on Intelligent Systems for Molecular Biology, p 175-182 Ed J. Glasgow, T. Littlejohn, F. Major, R. Lathrop, D. Sankoff, and C. Sensen Menlo Park, CA: AAAI Press, 1998), and clust (Remm M, and Sonnhammer E. Classification of transmembrane protein families in the Caenorhabditis elegans genome and identification of human orthologs. Genome Res. 2000 Nov;10(11):1679-89) programs. For example, the kinase domain (PFAM 00069) of CSNK1G from GI#s 11545751, 21314778, and 4758080 (SEQ ID NOs:13, 15, and 16, respectively) is located respectively at approximately amino acid residues 44-312, 46-312, and 43-308.

II. <u>High-Throughput In Vitro Fluorescence Polarization Assay</u>

Fluorescently-labeled CSNK1G peptide/substrate are added to each well of a 96-well microtiter plate, along with a test agent in a test buffer (10 mM HEPES, 10 mM NaCl, 6 mM magnesium chloride, pH 7.6). Changes in fluorescence polarization, determined by using a Fluorolite FPM-2 Fluorescence Polarization Microtiter System (Dynatech Laboratories, Inc), relative to control values indicates the test compound is a candidate modifier of CSNK1G activity.

III. <u>High-Throughput In Vitro Binding Assay.</u>

^{33}P-labeled CSNK1G peptide is added in an assay buffer (100 mM KCl, 20 mM HEPES pH 7.6, 1 mM $MgCl_2$, 1% glycerol, 0.5% NP-40, 50 mM beta-mercaptoethanol, 1 mg/ml BSA, cocktail of protease inhibitors) along with a test agent to the wells of a Neutralite-avidin coated assay plate and incubated at 25°C for 1 hour. Biotinylated substrate is then added to each well and incubated for 1 hour. Reactions are stopped by washing with PBS, and counted in a scintillation counter. Test agents that cause a difference in activity relative to control without test agent are identified as candidate p21 modulating agents.

IV. Immunoprecipitations and Immunoblotting

For coprecipitation of transfected proteins, 3×10^6 appropriate recombinant cells containing the CSNK1G proteins are plated on 10-cm dishes and transfected on the following day with expression constructs. The total amount of DNA is kept constant in each transfection by adding empty vector. After 24 h, cells are collected, washed once with phosphate-buffered saline and lysed for 20 min on ice in 1 ml of lysis buffer containing 50 mM Hepes, pH 7.9, 250 mM NaCl, 20 mM -glycerophosphate, 1 mM sodium orthovanadate, 5 mM p-nitrophenyl phosphate, 2 mM dithiothreitol, protease inhibitors (complete, Roche Molecular Biochemicals), and 1% Nonidet P-40. Cellular debris is removed by centrifugation twice at 15,000 × g for 15 min. The cell lysate is incubated with 25 μl of M2 beads (Sigma) for 2 h at 4 °C with gentle rocking.

After extensive washing with lysis buffer, proteins bound to the beads are solubilized by boiling in SDS sample buffer, fractionated by SDS-polyacrylamide gel electrophoresis, transferred to polyvinylidene difluoride membrane and blotted with the indicated antibodies. The reactive bands are visualized with horseradish peroxidase coupled to the appropriate secondary antibodies and the enhanced chemiluminescence (ECL) Western blotting detection system (Amersham Pharmacia Biotech).

V. Kinase assay

A purified or partially purified CSNK1G is diluted in a suitable reaction buffer, e.g., 50 mM Hepes, pH 7.5, containing magnesium chloride or manganese chloride (1-20 mM) and a peptide or polypeptide substrate, such as myelin basic protein or casein (1-10 μg/ml). The final concentration of the kinase is 1-20 nM. The enzyme reaction is conducted in microtiter plates to facilitate optimization of reaction conditions by increasing assay throughput. A 96-well microtiter plate is employed using a final volume 30-100 μl. The reaction is initiated by the addition of ^{33}P-gamma-ATP (0.5 μCi/ml) and incubated for 0.5 to 3 hours at room temperature. Negative controls are provided by the addition of EDTA, which chelates the divalent cation ($Mg2^+$ or Mn^{2+}) required for enzymatic activity. Following the incubation, the enzyme reaction is quenched using EDTA. Samples of the reaction are transferred to a 96-well glass fiber filter plate (MultiScreen, Millipore). The filters are subsequently washed with phosphate-buffered saline, dilute phosphoric acid (0.5%) or other suitable medium to remove excess radiolabeled ATP. Scintillation cocktail is added to the filter plate and the incorporated radioactivity is quantitated by scintillation counting (Wallac/Perkin Elmer). Activity is

defined by the amount of radioactivity detected following subtraction of the negative control reaction value (EDTA quench).

VI. Expression analysis

All cell lines used in the following experiments are NCI (National Cancer Institute) lines, and are available from ATCC (American Type Culture Collection, Manassas, VA 20110-2209). Normal and tumor tissues were obtained from Impath, UC Davis, Clontech, Stratagene, Ardais, Genome Collaborative, and Ambion.

TaqMan analysis was used to assess expression levels of the disclosed genes in various samples.

RNA was extracted from each tissue sample using Qiagen (Valencia, CA) RNeasy kits, following manufacturer's protocols, to a final concentration of 50ng/µl. Single stranded cDNA was then synthesized by reverse transcribing the RNA samples using random hexamers and 500ng of total RNA per reaction, following protocol 4304965 of Applied Biosystems (Foster City, CA).

Primers for expression analysis using TaqMan assay (Applied Biosystems, Foster City, CA) were prepared according to the TaqMan protocols, and the following criteria: a) primer pairs were designed to span introns to eliminate genomic contamination, and b) each primer pair produced only one product. Expression analysis was performed using a 7900HT instrument.

Taqman reactions were carried out following manufacturer's protocols, in 25 µl total volume for 96-well plates and 10 µl total volume for 384-well plates, using 300nM primer and 250 nM probe, and approximately 25ng of cDNA. The standard curve for result analysis was prepared using a universal pool of human cDNA samples, which is a mixture of cDNAs from a wide variety of tissues so that the chance that a target will be present in appreciable amounts is good. The raw data were normalized using 18S rRNA (universally expressed in all tissues and cells).

For each expression analysis, tumor tissue samples were compared with matched normal tissues from the same patient. A gene was considered overexpressed in a tumor when the level of expression of the gene was 2 fold or higher in the tumor compared with its matched normal sample. In cases where normal tissue was not available, a universal pool of cDNA samples was used instead. In these cases, a gene was considered overexpressed in a tumor sample when the difference of expression levels between a tumor sample and the average of all normal samples from the same tissue type was greater

than 2 times the standard deviation of all normal samples (i.e., Tumor – average(all normal samples) > 2 x STDEV(all normal samples)).

Results are shown in Table 1. Number of pairs of tumor samples and matched normal tissue from the same patient are shown for each tumor type. Percentage of the samples with at least two-fold overexpression for each tumor type is provided. A modulator identified by an assay described herein can be further validated for therapeutic effect by administration to a tumor in which the gene is overexpressed. A decrease in tumor growth confirms therapeutic utility of the modulator. Prior to treating a patient with the modulator, the likelihood that the patient will respond to treatment can be diagnosed by obtaining a tumor sample from the patient, and assaying for expression of the gene targeted by the modulator. The expression data for the gene(s) can also be used as a diagnostic marker for disease progression. The assay can be performed by expression analysis as described above, by antibody directed to the gene target, or by any other available detection method.

Table 1

SEQ ID NO	1	8	11
Breast	7%	17%	7%
# of Pairs	30	30	30
Colon	3%	12%	8%
# of Pairs	36	33	36
Kidney	0%	5%	0%
# of Pairs	20	20	20
Lung	3%	9%	0%
# of Pairs	35	32	35
Ovary	11%	26%	0%
# of Pairs	19	19	19
Prostate	7%	7%	7%
# of Pairs	15	14	15
Uterus	5%	5%	5%
# of Pairs	19	19	19

VII. CSNK1G functional assays

RNAi experiments were carried out to knock down expression of CSNK1Gs in various cell lines using small interfering RNAs (siRNA, Elbashir et al, *supra*).

Effect of CSNK1G RNAi on cell proliferation. BrdU and Cell Titer-Glo™ assays, as described above, were employed to study the effects of decreased CSNK1G expression on cell proliferation. The results of these experiments indicated that RNAi of CSNK1G of SEQ ID NOs:1, 8, and 11 decreases proliferation in LX1 small cell lung cancer cells, and in 231T breast cancer cells. MTS cell proliferation assay, as described above, was also employed to study the effects of decreased CSNK1G expression on cell proliferation. The results of this experiment indicated that RNAi of CSNK1G of SEQ ID NOs:1, 8, and 11 decreased proliferation in the above cell lines in addition to A549lung cancer cells.

Effect of CSNK1G RNAi on apoptosis. Nucleosome ELISA apoptosis assay, as described above, was employed to study the effects of decreased CSNK1G expression on apoptosis. Results indicated that RNAi of SEQ ID NO:1 caused increased apoptosis in A549 cells; RNAi of SEQ ID NO:8 did not have an affect on apoptosis; and RNAi of SEQ ID NO:11 caused increased apoptosis in LX1 cells.

SEQUENCE LISTING

<110> EXELIXIS, INC.

<120> CSNK1GS AS MODIFIERS OF THE P21 PATHWAY AND METHODS OF USE

<130> EX03-057C-PC

<150> US 60/401,739
<151> 2002-08-07

<160> 16

<170> PatentIn version 3.2

<210> 1
<211> 1950
<212> DNA
<213> Homo sapiens

<400> 1
aatatggtgc taaaagaact ccttaccttt ctctgactac aatttatttg gacatacttt 60

tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag actctgtcct 120

gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca actaaaccca 180

tggcacaaag gagtgcacac tgctctcgac catctggctc ctcatcgtcc tctggggttc 240

ttatggtggg acccaacttc agggttggca agaagatagg atgtgggaac ttcggagagc 300

tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg gaaccaataa 360

aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt ggcagtgcag 420

gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat gccatggtgc 480

tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga acatttactt 540

tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac gtgcactcaa 600

agaacctcat ttaccgagat gtcaagccag agaacttcct gattggtcga caaggcaata 660

agaaagagca tgttatacac attatagact ttggactggc caaggaatac attgaccccg 720

aaaccaaaaa acacatacct tatagggaac acaaaagttt aactggaact gcaagatata 780

tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg gaagccctag 840

gccatatgtt catgtatttc cttcgaggca gcctcccctg gcaaggactc aaggctgaca 900

cattaaaaga gagatatcaa aaaattggtg acaccaaaag gaatactccc attgaagctc 960

tctgtgagaa ctttccagag gagatggcaa cctaccttcg atatgtcagg cgactggact 1020

tctttgaaaa acctgattat gagtatttac ggaccctctt cacagacctc tttgaaaaga 1080

aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct actccagtag 1140

ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac acacataggg 1200

atcggccatc acaacagcag cctcttcgaa atcagagcct taggactgtt acagctgagc 1260

attatgatgt taacaactca gccatctggc acaggggaag aggcacctaa acagattctt 1320

ctggacaaaa gtcacatatg agcaagatca gaatagtagc agatgtaaaa agcagaatgt 1380

atcatcagag cgccgaggag agtgggaaat tcagcccagc cggcagacca atacctcata 1440

cctaacgtct cacttggctg cagaccgcca tgggggatca gtgcaggtgg ttagctcaac 1500

caatggagag ctgaatgttg atgatcccac gggagcccac tccaatgcac caatcacagc 1560

tcatgccgag gtggaggtag tggaggaagc taagtgctgc tgtttcttta agaggaaaag 1620

gaagaagact gctcagcgcc acaagtgacc agtgcctccc aggagtcctc aggccctggg 1680

gactctgact caattgtacc tgcagctcct gccatttctc attggaaggg actcctcttt 1740

gggggagggt ggatatccaa actaaaaaga agaaaacaga tgcccccaga aggggccagt 1800

gcgggcagcc agggcctagt gggtcattgg ccatctccgc ctgcctaagg ctctgagcag 1860

gtcccagagc tgctgttcct ccactgcttg cccatagggc tgcctggttg actctccttc 1920

ccattgttta cagtgaaggt gtcattcaca 1950

<210> 2
<211> 2066
<212> DNA
<213> Homo sapiens

<400> 2
ggcacgaggc tgctctcgac catctggctc ctcatcgtcc tctggggttc ttatggtggg 60

acccaacttc agggttggca agaagatagg atgtgggaac ttcggagagc tcagattagg 120

taaaaatctc tacaccaatg aatatgtagc aatcaaactg gaaccaataa aatcacgtgc 180

tccacagctt catttagagt acagatttta taaacagctt ggcagtgcag gtgaaggtct 240

cccacaggtg tattactttg gaccatgtgg gaaatataat gccatggtgc tggagctcct 300

tggccctagc ttggaggact tgtttgacct ctgtgaccga acatttactt tgaagacggt 360

gttaatgata gccatccagc tgctttctcg aatggaatac gtgcactcaa agaacctcat 420

ttaccgagat gtcaagccag agaacttcct gattggtcga caaggcaata agaaagagca 480

tgttatacac attatagact ttggactggc caaggaatac attgaccccg aaaccaaaaa 540

acacatacct tatagggaac acaaaagttt aactggaact gcaagatata tgtctatcaa 600

cacgcatctt ggcaaagagc aaagccggag agatgatttg gaagccctag gccatatgtt 660

catgtatttc cttcgaggca gcctcccctg gcaaggactc aaggctgaca cattaaaaga 720

gagatatcaa aaaattggtg acaccaaaag gaatactccc attgaagctc tctgtgagaa 780

ctttccagag gagatggcaa cctaccttcg atatgtcagg cgactggact tctttgaaaa 840

acctgattat gagtatttac ggaccctctt cacagacctc tttgaaaaga aaggctacac 900

ctttgactat gcctatgatt gggttgggag acctattcct actccagtag ggtcagttca 960

cgtagattct ggtgcatctg caataactcg agaaagccac acacataggg atcggccatc 1020

acaacagcag cctcttcgaa atcaggtggt tagctcaacc aatggagagc tgaatgttga 1080

tgatcccacg ggagcccact ccaatgcacc aatcacagct catgccgagg tggaggtagt 1140

ggaggaagct aagtgctgct gtttctttaa gaggaaaagg aagaagactg ctcagcgcca 1200

caagtgacca gtgcctccca ggagtcctca ggccctgggg actctgactc aattgtacct 1260

ggaggtggtg ggatttgtga ttggaaggga ctcctctttg ggggagggtg gatatccaaa 1320

gcagctcctg ccatttctca ttggaaggga ctcctctttg ggggagggtg gatatccaaa 1320

ccaaaaagaa gaaaacagat gcccccagaa ggggccagtg cgggcagcca gggcctagtg 1380

ggtcattggc catctccgcc tgcctaaggc tctgagcagg tcccagagct gctgttcctc 1440

cactgcttgc ccatagggct gcctggttga ctctccttcc cattgtttac agtgaaggtg 1500

tcattcacaa aaactcaagg actgctattc tccttcttcc ccttagttta ctcctggttt 1560

ttaccccacc ctcaaccctc tccagcataa aacctagtga gctaaaggct ttgtctgcag 1620

aaggagatca agaggctggg ggtaaggcca agaaggtagg aggaaaatgg cagacctggg 1680

ctggagaaga accttctccg tatcccaggt gtgcctggca gtatggtttc ctcttcctct 1740

gtgcctgtgc agcattcatc ccagctggcc ttggggttca ggttccttct tccctccctc 1800

ctgtgaagtt acactgtagg acacaagctg tgagcaatct gcagtctact gtccctgtgt 1860

gttggcgttc ttagcttttt tgacaaactc ttttctccag gtagtaggac aatgaaaatt 1920

gttctaagca aaggaaagaa aactgacttt gttgcacttt tagtttttt aaaaaaaaca 1980

aaaacaaaaa catggcagat gcaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2040

aaaaaaaaaa aaaaaaaaaa aaaaaa 2066

<210> 3
<211> 3552
<212> DNA
<213> Homo sapiens

<400> 3
ggagacggtg ttaatgatag ccatccagct gctttctcga atggaatacg tgcactcaaa 60

gaacctcatt taccgagatg tcaagccaga gaacttcctg attggtcgac gaggcaataa 120

gaaagagcat gttatacaca ttatagactt tggactggcc aaggaataca ttgaccccga 180

aaccaaaaaa cacatacctt atagggaaca caaaagttta actggaactg caagatatat 240

gtctatcaac acgcatcttg gcaaagagca aagccggaga gatgatttgg aagccctagg 300

ccatatgttc atgtatttcc ttcgaggcag cctcccctgg caaggactca aggctgacac 360

attaaaagag agatatcaaa aaattggtga caccaaaagg aatactccca ttgaagctct 420

ctgtgagaac tttccagagg agatggcaac ctaccttcga tatgtcaggc gactggactt 480

ctttgaaaaa cctgattatg agtatttacg gaccctcttc acagacctct ttgaaaagaa 540

aggctacacc tttgactatg cctatgattg ggttgggaga cctattccta ctccagtagg 600

gtcagttcac gtagattctg gtgcatctgc aataactcga gaaagccaca cacataggga 660

tcggccatca caacagcagc ctcttcgaaa tcaggtggtt agctcaacca atggagagct 720

gaatgttgat gatcccacgg gagcccactc caatgcacca atcacagctc atgccgaggt 780

ggaggtagtg gaggaagcta agtgctgctg tttctttaag aggaaaagga agaagactgc 840

tcagcgccac aagtgaccag tgcctcccag gagtcctcag gccctgggga ctctgactca 900

attgtacctg cagctcctgc catttctcat tggaagggac tcctctttgg gggagggtgg 960

atatccaaac caaaaagaag aaaacagatg cccccagaag gggccagtgc gggcagccag 1020

ggcctagtgg gtcattggcc atctccgcct gcctaaggct ctgagcaggt cccagagctg 1080

ctgttcctcc actgcttgcc catagggctg cctggttgac tctccttccc attgtttaca 1140

gtgaaggtgt cattcacaaa aactcaagga ctgctattct ccttcttccc cttagtttac 1200

tcctggtttt taccccaccc tcaaccctct ccagcataaa acctagtgag ctaaaggctt 1260

tgtctgcaga aggagatcaa gaggctgggg gtaaggccaa gaaggtagga ggaaaatggc 1320

agacctgggc tggagaagaa ccttctccgt atcccaggtg tgcctggcag tatggtttcc 1380

tcttcctctg tgcctgtgca gcattcatcc cggctggcct tggggttcag gttccttctt 1440

ccctccctcc tgtgaagtta cactgtagga cacaagctgt gagcaatctg cagtctactg 1500

tccccgtgtg taggcgttct tagcttttt gacaaactct tttctccagg tagtaggaca 1560

atgaaaattg ttctaagcaa aggaaagaaa actgactttg ttgcactttt agtttttta 1620

aaaaaaacaa aaacaaaaac atggcagatg catattgtgt ctggttatat tgggggtttt 1680

acttttacct gttttgaggg ggatggggcc ggccaagcca ttcagagaga acatgggtcc 1740

agaggacatt ctcagtggaa agagtttgat ctgcagcacc cagaagagaa gccaaactcg 1800

gtgtcattct gagtgaacac tcaggttggc aagaaaacat acttgaattt tcattcatct 1860

tctcagcagc tgaagaatgt ccctgccaga gcatcttgac ctaatcagct tacagtttga 1920

aaacctagct ctccagaaca tgagatgagc cagccgagcc agactgtgac caggaaacag 1980

ctcatcccag agaaggagat gcttaacaaa aaaaaattga aattgtttcc catgctgcca 2040

gggacttcca actagatagc catgtgacgt cctggtgact tgagggaaaa attagtgatg 2100

aaacagccac caccatattg ccattagtga aaaaaaagag gacagtgaac ctgccttcca 2160

cctgccagag ggacctcagg gtgtggcatt atagggccag gaaaagaaaa tcggtgtatc 2220

ctatctgccc caatagctga gctgtagcat ttgggctggc ctgccttatc agaaaccaag 2280

cttatgaaga tcttctccca gcaggtccat agcagtaggc ttaggatgca gtatatgggg 2340

ccgcatttaa aaggagggaa agattgtttg gtgctggaac attccaggga aaaggagact 2400

ggaatgaaag gtctgaaatt atcttctcaa ttggactcct tccagaaagg tggccgtgcc 2460

tctaagcatg tttttcccag tatgccctag gcctcccccc atggtgtttt catatgaggt 2520

acaactgtga aggatctggt tcctcattca ctgtttgaca agtctttcat gtgtggagtt 2580

actcttctca tgcccaattt tcatttgagt ttagtggctt aaccaaacaa tgactcctca 2640

ttccagcggt gacagaagag aaagggtcat ttacatcagg aaagaggtct tgtatctggg 2700

agtagagagc taaccatgga gcacagtggc tggtgggtga cttagtctga tggtttgtgg 2760

accatagaag tcttcacctc tggtttgagg tgcagggctg tcttttgtac tggagggtgt 2820

ggggatattt tctgatagtt gccatttctt gaaaaattcc cttgatgtac cttacacaga 2880

gcagaaataa cattaacatg gatcagaggt actgggcttc atctgttcca ttggaccttg 2940

gctagggaat atcatttcac tggcatcaaa cctgcttagc ttatgaaaag atggtaatat 3000

gtcatttcta taaatgtttc tatatatgaa acataaagtg gcagggagat acaatatcac 3060

acccccttccc cacaaggact gtgaatattg ggatttatgt ccttgccatt acctagtggt 3120

tacagcccta tcactaaaat ttacatcgtt tctcagttgg gatttgggca ttgctaactt 3180

actgtataga aagtttaact tttcctcacc cctgtataga aaatgccttg cctctcaaga 3240

gagggcagag gggggggccag gtgcagtggc tcacgcctgt aatcccagca gtttgggagg 3300

ccaaggcaag tggatcatgt gaggtcaaga gttcgagacc agcctggcca acatggtgaa 3360

accccgtctc tacaaaaaat acaaaaatta gctgggcatg gtggcatgct cccgtagtcc 3420

cagctactcg ggaggctgag gcaggagaat cacttgagcc tgggaggcag aagttgcagt 3480

gagccgagat cgcaccactg cactccagcc tgggcaacag agtgagactc tgtctaaaaa 3540

aaaaaaaaaa aa 3552

<210> 4
<211> 4222
<212> DNA
<213> Homo sapiens

<400> 4
gcctcatttg cgccttgcag cactgctgga ccaggttaca agatgttcac ctaagattga 60

gacctagtga ctacatttcc tacgggaaca aataaatggt ttttcatctc ccggagatac 120

attacaaaca aatatggtgc taaaagaact ccttaccttt ctctgactac aatttatttg 180

gacatacttt tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag 240

actctgtcct gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca 300

actaaaccca tggcacaaag gagtgcacac tgctctcgac catctggctc ctcatcgtcc 360

tctggggttc ttatggtggg acccaacttc agggttggca agaagatagg atgtgggaac 420
ttcggagagc tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg 480
gaaccaataa aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt 540
ggcagtgcag gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat 600
gccatggtgc tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga 660
acatttactt tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac 720
gtgcactcaa agaacctcat ttaccgagat gtcaagccag agaacttcct gattggtcga 780
caaggcaata agaaagagca tgttatacac attatagact ttggactggc caaggaatac 840
attgaccccg aaaccaaaaa acacatacct tatagggaac acaaaagttt aactggaact 900
gcaagatata tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg 960
gaagccctag gccatatgtt catgtatttc cttcgaggca gcctcccctg gcaaggactc 1020
aaggctgaca cattaaaaga gagatatcaa aaaattggtg acaccaaaag gaatactccc 1080
attgaagctc tctgtgagaa ctttccagag gagatggcaa cctaccttcg atatgtcagg 1140
cgactggact tctttgaaaa acctgattat gagtatttac ggaccctctt cacagacctc 1200
tttgaaaaga aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct 1260
actccagtag ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac 1320
acacataggg atcggccatc acaacagcag cctcttcgaa atcaggtggt tagctcaacc 1380
aatggagagc tgaatgttga tgatcccacg ggagcccact ccaatgcacc aatcacagct 1440
catgccgagg tggaggtagt ggaggaagct aagtgctgct gtttctttaa gaggaaaagg 1500
aagaagactg ctcagcgcca caagtgacca gtgcctccca ggagtcctca ggccctgggg 1560
actctgactc aattgtacct gcagctcctg ccatttctca ttggaaggga ctcctctttg 1620
ggggagggtg gatatccaaa ccaaaaagaa gaaaacagat gcccccagaa ggggccagtg 1680
cgggcagcca gggcctagtg ggtcattggc catctccgcc tgcctaaggc tctgagcagg 1740
tcccagagct gctgttcctc cactgcttgc ccatagggct gcctggttga ctctccttcc 1800
cattgtttac agtgaaggtg tcattcacaa aaactcaagg actgctattc tccttcttcc 1860
ccttagttta ctcctggttt ttaccccacc ctcaaccctc tccagcataa aacctagtga 1920
gctaaaggct ttgtctgcag aaggagatca agaggctggg ggtaaggcca agaaggtagg 1980
aggaaaatgg cagacctggg ctggagaaga accttctccg tatcccaggt gtgcctggca 2040
gtatggtttc ctcttcctct gtgcctgtgc agcattcatc ccagctggcc ttggggttca 2100
ggttccttct tccctccctc ctgtgaagtt acactgtagg acacaagctg tgagcaatct 2160
gcagtctact gtccctgtgt gttggcgttc ttagcttttt tgacaaactc ttttctccag 2220

gtagtaggac aatgaaaatt gttctaagca aaggaaagaa aactgacttt gttgcacttt 2280
tagttttttt aaaaaaaaca aaaacaaaaa catggcagat gcatattgtg tctggttata 2340
ttgggggttt tacttttacc tgttttgagg gggatggggc cggccaagcc attcagagag 2400
aacatgggtc cagaggacat tctcagtgga aagagtttga tctgcagcac ccagaagaga 2460
agccaaactc ggtgtcattc tgagtgaaca ctcaggttgg caagaaaaca tacttgaatt 2520
ttcattcatc ttctcagcag ctgaagaatg tccctaccag agcatcttga cctaatcagc 2580
ttacagtttg aaaacctagc tctccagaac atgagatgag ccagccgagc cagactgtga 2640
ccaggaaaca gctcatccca gagaaggaga tgcttaacaa aaaaaaattg aaattgtttc 2700
ccatgctgcc agggacttcc aactagatag ccatgtgacg tcctggtgac ttgggggaaa 2760
aattagtgat gaaacagcca ccaccatatt gccattagtg gaaaaaaaga ggacagtgaa 2820
cctgccttcc acctgccaga gggacctcag ggtgtggcat tatagggcca ggaaaagaaa 2880
atcggtgtat cctatctgcc ccaatagctg agctgtagca tttgggctgg cctgccttat 2940
cagaaaccaa gcttatgaag atcttctccc agcaggtcca tagcagtagg cttaggatgc 3000
agtatatggg gccgcattta aaaggaggga aagattgttt ggtgctggaa cattccaggg 3060
aaaaggagac tggaatgaaa ggtctgaaat tatcttctca attggactcc ttccagaaag 3120
gtggccgtgc ctctaagcat gtttttccca gtatgcccta ggcctccccc catggtgttt 3180
tcatatgagg tactactgtg aaggatctgg ttcctcattc actgtttgac aagtctttca 3240
tgtgtggagt tactcttctc atgcccaatt ttcatttgag tttagtggct taaccaaaca 3300
atgactcctc attccagcgg tgacagaaga gaaagggtca tttacatcag gaaagaggtc 3360
ttgtatctgg gagtagagag ctaaccatgg agcacagtgg ctggtgggtg acttagtctg 3420
atggtttgtg gaccatagaa gtcttcacct ctggtttgag gtgcagggct gtcttttgta 3480
ctggagggtg tggggatatt ttctgatagt tgccatttct tgaaaaattc ccttgatgta 3540
ccttacacag agcagaaata acattaacat ggatcagagg tactgggctt catctgttcc 3600
attggacctt ggctagggaa tatcatttca ctggcatcaa acctgcttag cttatgaaaa 3660
gatggtaata tgtcatttct ataaatgttt ctatatatga aacataaagt ggcagggaga 3720
tacaatatca cacccttcc ccacaaggac tgtgaatatt gggatttatg tccttgccat 3780
tacctagtgg ttacagccct atcactaaaa tttacatcgt ttctcagttg ggatttgggc 3840
attgctaact tactgtatag aaagtttaac ttttcctcac ccctgtatag aaaatgcctt 3900
gcctctcaag agagggcaga gggggggcca ggtgcagtgg ctcacgcctg taatcccagc 3960
agtttgggag gccaaggcaa gtggatcatg tgaggtcaag agttcgagac cagcctggcc 4020
aacatggtga aaccccgtct ctacaaaaaa tacaaaaatt agctgggcat ggtggcatgc 4080

tcccgtagtc ccagctactc gggaggctga ggcaggagaa tcacttgagc ctgggaggca 4140

gaagttgcag tgagccgaga tcgcaccact gcactccagc ctgggcaaca gagtgagact 4200

ctgtctaaaa aaaaaaaaaa aa 4222

<210> 5
<211> 1754
<212> DNA
<213> Homo sapiens

<400> 5
gacctagtga ctacatttcc tacgggaaca ataaatggt ttttcatctc ccggagatac 60

attacaaaca aatatggtgc taaaagaact ccgtaccttt ctctgactac aatttatttg 120

gacatacttt tgtattgaag agaggtatac atactgaagc tacttgctgt actataggag 180

actctgtcct gtaggatcat ggaccatcct agtagggaaa aggatgaaag acaacggaca 240

actaaaccca tggcacaaag gagtgcacac tgctctcgac catctggctc ctcatcgtcc 300

tctggggttc ttatggtggg acccaacttc agggttggca agaagatagg atgtgggaac 360

ttcggagagc tcagattagg taaaaatctc tacaccaatg aatatgtagc aatcaaactg 420

gaaccaataa aatcacgtgc tccacagctt catttagagt acagatttta taaacagctt 480

ggcagtgcag gtgaaggtct cccacaggtg tattactttg gaccatgtgg gaaatataat 540

gccatggtgc tggagctcct tggccctagc ttggaggact tgtttgacct ctgtgaccga 600

acatttactt tgaagacggt gttaatgata gccatccagc tgctttctcg aatggaatac 660

gtgcactcaa agaacctcat ttaccgagat gtcaagccag agaacttcct gattggtcga 720

caaggcaata agaaagagca tgttatacac attatagact ttggactggc caaggaatac 780

attgaccccg aaaccaaaaa acacatacct tatagggaac acaaaagttt aactggaact 840

gcaagatata tgtctatcaa cacgcatctt ggcaaagagc aaagccggag agatgatttg 900

gaagccctag gccatatgtt catgtatttc cttcgaggca gcctcccctg gcaaggactc 960

aaggctgaca cattaaaaga gagatatcaa aaaattggtg acaccaaaag gaatactccc 1020

attgaagctc tctgtgagaa ctttccagag gagatggcaa cctaccttcg atatgtcagg 1080

cgactggact tctttgaaaa acctgattat gagtatttac ggacctctt cacagacctc 1140

tttgaaaaga aaggctacac ctttgactat gcctatgatt gggttgggag acctattcct 1200

actccagtag ggtcagttca cgtagattct ggtgcatctg caataactcg agaaagccac 1260

acacataggg atcggccatc acaacagcag cctcttcgaa atcaggtggt tagctcaacc 1320

aatggagagc tgaatgttga tgatcccacg ggagcccact ccaatgcacc aatcacagct 1380

catgccgagg tgggggtagt ggaggaagct aaatgtataa tgtttcataa accctggcac 1440

tggtcacaaa gctcagctgt gaaaatgaaa tttgtagtat ttttaaacat gaatgtcaat 1500

ttcaagtgta tttgaaatgg ttcctccagg agagatttgt gcaccattag gaaaatcttc 1560

tctgcagagg aagtagcctt ctttggagaa aatggaaaat gggttctgat atgtgatctc 1620

agagtagccc atttcctagg gcaccatgga aaacacaaat gtgatcttta agtatacctc 1680

ttccccagtt tggggaggaa aggactcagt ttgcaccctt tttgtatgta aaataaaatg 1740

tcttaccttt cttg 1754

<210> 6
<211> 2446
<212> DNA
<213> Homo sapiens

<400> 6
gggatttgca cggcagcaga gtcaccgtgg agaggccagg gtatcacaaa cttatggatt 60

ttgacaagaa aggagggaaa ggggagacgg aggagggccg gagaatgtcc aaggccggcg 120

ggggccggag cagccacggc atccggagct cggggaccag ctcgggggtc ctgatggtgg 180

gccccaactt ccgcgtcggc aagaagatcg gctgcggcaa cttcggggag ctccgcctag 240

gaaagaatct ctatacaaat gaatacgtgg ctatcaaatt ggagccgatc aagtcccggg 300

ccccgcagct gcacctggag taccggttct acaagcagct cagcgccaca gagggcgtcc 360

ctcaggtcta ctacttcggt ccgtgcggga attacaacgc catggtgctg gagctgctgg 420

ggcccagcct ggaggacctg ttcgacctgt gcgaccggac cttcacgctc aagacggtgc 480

tgatgatcgc catccagctg atcacgcgca tggagtatgt gcacaccaag agcctaatct 540

accgggacgt gaagcccgag aacttcctgg tgggccgccc ggggaccaag cggcagcatg 600

ccatccacat catcgacttc gggctggcca aggagtacat cgaccccgag accaagaagc 660

acatcccgta ccgcgagcac aagagcctga cgggcacggc gcgctacatg agcatcaaca 720

cgcacctggg caaggagcag agccgccgcg acgacctgga ggcgctgggc cacatgttca 780

tgtacttcct gcgcggcagc ctcccctggc aggggctcaa ggccgacacg ctcaaggagc 840

ggtaccagaa gatcggggac accaaacgcg ccacgcccat cgaggtgctc tgcgagaact 900

tcccagagga gatggccacg tacctgcgct atgtgcggcg cctggacttc ttcgagaagc 960

ccgactatga ctacctgcgg aagctcttca ccgacctctt cgaccgcagt ggcttcgtgt 1020

tcgactatga gtacgactgg gccgggaagc ccctgccgac ccccatcggc accgtccaca 1080

ccgacctgcc ctcccagcct cagctccggg acaaaaccca gccgcacagc aaaaaccagg 1140

cgttgaactc caccaacggg gagctgaatg cggacgaccc cacggccggc cactccaacg 1200

ccccgatcac agcgcctgca gaggtggagg tggccgatga aaccaaatgc tgctgtttct 1260

tcaagaggag aaagagaaaa tcgctgcagc gacacaagtg accctgggcg cgtgcagccc 1320

cctgaatctt ctccgtgcag ccccttgggg cgcgaccttg tgcgaggccc tcggggccca 1380

cccacagcgg cccagggcca gaccctggct ggaagccaga acgcagactg caggggccgc 1440

gcctggctca ggcggcccca cccccgggac gtggggtcac ttccttcatg taagactttg 1500

gccgaaattt ctacacctgt gtctagtcct cccctccaag agcattaact atttaaaaca 1560

aggaaaagag gaaaaaaaaa acagaggccc gccctacccc actcctgccc ctccgtttct 1620

ttgctgaagt gagtagtgtg atcctggagg ccccccggcc tggccccgcc ccgccagccg 1680

cccccgttag cgtcataaag tccagcttgt ctccctcgat ccaaaggccg ttttctcgag 1740

gggagggcag gcccggcctg gaggggtgct gtggagctgt cttgcccagg ccctcctggg 1800

aggggacag gcattgttgc caggggtgag gccgtgcccc aggcctcccc gaaaccaaag 1860

gggaaggcag gggtggggcc gtggctgaag ccggctcccc aaccaaaatg ctgcaccaaa 1920

gctcgggcgc cgcgggcacg gctgctgcag tctcttccca gcctggccct ggcaaggggc 1980

gggtgggcgc tgccaggcgg gtgcttctcg acgcacttgc tcccggaggc tgcgccccgg 2040

cgcctggaac ccgaggtggg aggaccggtt ggtgtcaccc tgctcggccc tcagccctgc 2100

cgcgtggggc gcgtgggcac ggagcttcct gcctctgctc cgacacccgg caagcagccg 2160

gagacaaaac gccttaaagc ccccggccca gccctgcagg tatattgcag gggcctgggg 2220

gcggccctgg actggcgggc ggttccccag tggggtgccc tggaggctgc cgggcagagt 2280

ggagcagctt ggggccgtgc ccagggcggt ggctgtgagt ctagtttttg ctttaccaag 2340

tgtacagaaa tggcatttac gtttctctga tgctcccttg aagccataga atttaggggc 2400

ttttttaaaa aaataaaaga aaaatgaaac caaaaaaaaa aaaaaa 2446

<210> 7
<211> 2446
<212> DNA
<213> Homo sapiens

<400> 7
gggatttgca cggcagcaga gtcaccgtgg agaggccagg gtatcacaaa cttatggatt 60

ttgacaagaa aggagggaaa ggggagacgg aggagggccg gagaatgtcc aaggccggcg 120

ggggccggag cagccacggc atccggagct cggggaccag ctcgggggtc ctgatggtgg 180

gccccaactt ccgcgtcggc aagaagatcg gctgcggcaa cttcggggag ctccgcctag 240

gaaagaatct ctatacaaat gaatacgtgg ctatcaaatt ggagccgatc aagtcccggg 300

ccccgcagct gcacctggag taccggttct acaagcagct cagcgccaca gagggcgtcc 360

ctcaggtcta ctacttcggt ccgtgcggga attacaacgc catggtgctg gagctgctgg 420

ggcccagcct ggaggacctg ttcgacctgt gcgaccggac cttcacgctc aagacggtgc 480

tgatgatcgc catccagctg atcacgcgca tggagtatgt gcacaccaag agcctaatct 540

accgggacgt gaagcccgag aacttcctgg tgggccgccc ggggaccaag cggcagcatg 600

ccatccacat catcgacttc gggctggcca aggagtacat cgaccccgag accaagaagc 660

acatcccgta ccgcgagcac aagagcctga cgggcacggc gcgctacatg agcatcaaca 720

cgcacctggg caaggagcag agccgccgcg acgacctgga ggcgctgggc cacatgttca 780

tgtacttcct gcgcggcagc ctcccctggc aggggctcaa ggccgacacg ctcaaggagc 840

ggtaccagaa gatcggggac accaaacgcg ccacgcccat cgaggtgctc tgcgagaact 900

tcccagagga gatggccacg tacctgcgct atgtgcggcg cctggacttc ttcgagaagc 960

ccgactatga ctacctgcgg aagctcttca ccgacctctt cgaccgcagt ggcttcgtgt 1020

tcgactatga gtacgactgg gccgggaagc ccctgccgac ccccatcggc accgtccaca 1080

ccgacctgcc ctcccagcct cagctccggg acaaaaccca gccgcacagc aaaaaccagg 1140

cgttgaactc caccaacggg gagctgaatg cggacgaccc cacggccggc cactccaacg 1200

ccccgatcac agcgcctgca gaggtggagg tggccgatga aaccaaatgc tgctgtttct 1260

tcaagaggag aaagagaaaa tcgctgcagc gacacaagtg accctgggcg cgtgcagccc 1320

cctgaatctt ctccgtgcag ccccttgggg cgcgaccttg tgcgaggccc tcggggccca 1380

cccacagcgg cccagggcca gaccctggct ggaagccaga acgcagactg caggggccgc 1440

gcctggctca ggcggcccca cccccgggac gtggggtcac ttccttcatg taagactttg 1500

gccgaaattt ctacacctgt gtctagtcct cccctccaag agcattaact atttaaaaca 1560

aggaaaagag gaaaaaaaaa acagaggccc gccctacccc actcctgccc ctccgtttct 1620

ttgctgaagt gagtagtgtg atcctggagg ccccccggcc tggccccgcc ccgccagccg 1680

cccccgttag cgtcataaag tccagcttgt ctccctcgat ccaaaggccg ttttctcgag 1740

gggagggcag gcccggcctg gaggggtgct gtggagctgt cttgcccagg ccctcctggg 1800

aggggacag gcattgttgc caggggtgag gccgtgcccc aggcctcccc gaaaccaaag 1860

gggaaggcag gggtggggcc gtggctgaag ccggctcccc aaccaaaatg ctgcaccaaa 1920

gctcgggcgc cgcgggcacg gctgctgcag tctcttccca gcctggccct ggcaaggggc 1980

gggtgggcgc tgccaggcgg gtgcttctcg acgcacttgc tcccggaggc tgcgccccgg 2040

cgcctggaac ccgaggtggg aggaccggtt ggtgtcaccc tgctcggccc tcagccctgc 2100

cgcgtggggc gcgtgggcac ggagcttcct gcctctgctc cgacacccgg caagcagccg 2160

gagacaaaac gccttaaagc ccccggccca gccctgcagg tatattgcag gggcctgggg 2220

gcggccctgg actggcgggc ggttccccag tggggtgccc tggaggctgc cgggcagagt 2280

ggagcagctt ggggccgtgc ccagggcggt ggctgtgagt ctagtttttg ctttaccaag 2340

tgtacagaaa tggcatttac gtttctctga tgctcccttg aagccataga atttaggggc 2400

ttttttaaaa aaataaaaga aaaatgaaac caaaaaaaaa aaaaaa 2446

<210> 8
<211> 2773
<212> DNA
<213> Homo sapiens

<400> 8
cggcacgagc agcagaatgt ctcctgcccc cgagagcgac cccgaggcca ctgagaagag 60

cagcgcggcc tggccggccc gaacgcctgc gtctcagtag ctgggagcca cgggcccacg 120

cccgcccacc ggccgcagtg atgttctagc cacagaggag ccaagacctc aggtttccag 180

agacttggga tttgcacggc agcagagtca ccgtggagag gccagggtat cacaaactta 240

tggattttga caagaaagga gggaaagggg agacggagga gggccggaga atgtccaagg 300

ccggcggggg ccggagcagc cacggcatcc ggagctcggg gaccagctcg ggggtcctga 360

tggtgggccc caacttccgc gtcggcaaga agatcggctg cggcaacttc ggggagctcc 420

gcctagtgca gagtagaaga gccgggaagg accccatcac catcagatgc atcagagctt 480

caaaggcccg cccatcaaga cgctgccaag agcatgaggc cagccaggct ggggaggacg 540

ttggcgagca tccgcctgcc cgtcagggag ccacagctgg atgatacatc gacagacagc 600

agagggacgg tgtagcacac aggcgagggc ttgtcctcct cagcctgggg gagcgcagct 660

cagtgcttgg tggtcagcac gtgactgaca ctttgccgga gaaggggacg gctgcccgca 720

ggaagagcta cacagccgtg ttcgccggaa ctccgctctg gagagtcaga actggagacg 780

ggagggcgtg cctcaagggc agatgggcaa accaaacccc atcctccctg tgagccgcgg 840

cacagcacca ggcgggcaca cgcagcacga cacggcgagg agcccaccgc aggggcagcc 900

ttttgggcgg ggaaggggggg atgaatccac tgtgaatccc acctccgaag ggcagcgcag 960

gaaggacacg caaatgctgg actgaacctg cagctgattc agggggcagt gtcctgagct 1020

ctaaaacttc aaacctgggg cagtgtcctg agctctaaaa cttcgaaatc tgaccaggca 1080

cagtggctca cgcctgtaat cccaacactt tgggaggctg atgtgggcgg atcacctgac 1140

gtcaggagtt cgagaccagc ctggacaaca tggagaaagc ccatctctag taaaaataca 1200

aaaattagcc aggttgatgg tgcctgcctg taattccagc tactcaggag gctgaggcag 1260

gagaatcgct tgaacccagg aggcagaggt tccagtgagc cgtgatcgcg ccagtgcact 1320

ccagcctggg caacagaacg agactctgct ccaaaaagaa aaaaactttg aaatccctct 1380

aaaaataaaa tggaccggtg gatggtagct gtgaggatgg accgtgtggc aggaaagaat 1440

ctctatacaa atgaatacgt ggctatcaaa ttggagccga tcaagtcccg ggccccgcag 1500

ctgcacctgg agtaccggtt ctacaagcag ctcagcgcca cagagggcgt ccctcaggtc 1560

tactacttcg gtccgtgcgg gaagtacaac gccatggtgc tggagctgct ggggcccagc 1620

ctggaggacc tgttcgacct gtgcgaccgg accttcacgc tcaagacggt gctgatgatc 1680

gccatccagc tgatcacgcg catggagtat gtgcacacca agagcctaat ctaccgggac 1740

gtgaagcccg agaacttcct ggtgggccgc ccggggacca agcggcagca tgccatccac 1800

atcatcgact tcgggctggc caaggagtac atcgaccccg agaccaagaa gcacatcccg 1860

taccgcgagc acaagagcct gacgggcacg gcgcgctaca tgagcatcaa cacgcacctg 1920

ggcaaggagc agagccgccg cgacgacctg gaggcgctgg gccacatgtt catgtacttc 1980

ctgcgcggca gcctcccctg gcaggggctc aaggccgaca cgctcaagga gcggtaccag 2040

aagatcgggg acaccaaacg cgccacgccc atcgaggtgc tctgcgagaa cttcccagag 2100

gagatggcca cgtacctgcg ctatgtgcgg cgcctggact tcttcgagaa gcccgactat 2160

gactacctgc ggaagctctt caccgacctc ttcgaccgca gtggcttcgt gttcgactat 2220

gagtacgact gggccgggaa gcccctgccg acccccatcg gcaccgtcca caccgacctg 2280

ccctcccagc ctcagctccg ggacaaaacc cagccgcaca gcaaaaacca ggcgttgaac 2340

tccaccaacg gggagctgaa tgcggacgac cccacggccg gccactccaa cgccccgatc 2400

acagcgcctg cagaggtgga ggtggccgat gaaaccaaat gctgctgttt cttcaagagg 2460

agaaagagaa aatcgctgca gcgacacaag tgaccctggg cgcgtgcagc cccctgaatc 2520

ttctccgtgc agccccttgg ggcgcgacct tgtgcgaggc cctcggggcc cacccacagc 2580

ggcccagggc cagaccctgg ctggaagcca gaacgcagac tgcaggggcc gcgcctggct 2640

caggcggccc caccccgggg acgtggggtc acttccttca tgtaagactt tggccgaaat 2700

ttctacacct gtgtctagtc ctcccctcca agagcattaa ctatttaaaa caaggaaaaa 2760

aaaaaaaaaa aaa 2773

<210> 9
<211> 1615
<212> DNA
<213> Homo sapiens

<400> 9
agtacatcga ccccgagacc aagaagcaca tcccgtaccg cgagcacaag agcctgacgg 60

gcacggcgcg ctacatgagc atcaacacgc acctgggcaa ggagcagagc cgccgcgacg 120

acctggaggc gctgggccac atgttcatgt acttcctgcg cggcagcctc ccctggcagg 180

ggctcaaggc cgacacgctc aaggagcggt accagaagat cggggacacc aaacgcgcca 240

cgcccatcga ggtgctctgc gagaacttcc cagaggagat ggccacgtac ctgcgctatg 300

tgcggcgcct ggacttcttc gagaagcccg actatgacta cctgcggaag ctcttcaccg 360

acctcttcga ccgcagtggc ttcgtgttcg actatgagta cgactgggcc gggaagcccc 420

tgccgacccc catcggcacc gtccacaccg acctgccctc ccagcctcag ctccgggaca 480

aaacccagcc ccctgaatct tctccgtgca gcccccttggg gcgcgacctt gtgcgaggcc 540

ctcggggccc acccacagcg gcccagggcc agaccctggc tggaagccag aacgcagact 600

gcaggggccg cgcctggctc aggcggcccc acccccggga cgtggggtca cttccttcat 660

gtaagacttt ggccgaaatt tctacacctg tgtctagtcc tcccctccaa gagcattaac 720

tatttaaaac aaggaaaaga ggaaaaaaaa aacagaggcc cgccctaccc cactcctgcc 780

cctccgtttc tttgctgaag tgagtagtgt gatcctggag gcccccggc ctggccccgc 840

cccgccagcc gcccccgtta gcgtcataaa gtccagcttg tctccctcga tccaaaggcc 900

gttttctcga ggggagggca ggcccggcct ggaggggtgc tgtggagctg tcttgcccag 960

gccctcctgg gaggggggaca ggcattgttg ccaggggtga ggccgtgccc caggcctccc 1020

cgaaaccaaa ggggaaggca ggggtggggc cgtggctgaa gccggctccc caaccaaaat 1080

gctgcaccaa agctcgggcg ccgcgggcac ggctgctgca gtctcttccc agcctggccc 1140

tggcaagggg cgggtgggcg ctgccaggcg ggtgcttctc gacgcacttg ctcccggagg 1200

ctgcgccccg gcgcctggaa cccgaggtgg gaggaccggt tggtgtcacc ctgctcggcc 1260

ctcagccctg ccgcgtgggg cgcgtgggca cggagcttcc tgcctctctg ctccgacacc 1320

cggcaagcag ccggagacaa aacgccttaa agcccccggc ccagccctgc aggtatattg 1380

caggggcctg ggggcggccc tggactggcg ggcggttccc cagtggggtg ccctggaggc 1440

tgccgggcag agtggagcag cttggggccg tgcccagggc ggtggctgtg agtctagttt 1500

ttgctttacc aagtgtacag aaatggcatt tacgtttctc tgatgctccc ttgaagccat 1560

agaatttagg ggctttttta aaaaaaataa aagaaaaatg aaaccaaaaa aaaaa 1615

<210> 10
<211> 2421
<212> DNA
<213> Homo sapiens

<400> 10
gaattcaaag tggagtaccg caaacttgat atggaaaata aaaagaaaga caaggacaaa 60

tcagatgata gaatggcacg acctagtggt cgatcgggac acaacactcg aggaactggg 120

tcttcatcgt ctggagtttt aatggttgga cctaacttta gagttggaaa aaaaattgga 180

tgtggcaatt ttggagaatt acgattaggg aaaaatttat acacaaatga atatgtggca 240

attaagttgg agcccatgaa atcaagagca ccacagctac atttggaata cagattctat 300

aagcagttag gatctggaga tggtatacct caagtttact atttcggccc ttgtggtaaa 360

tacaatgcta tggtgctgga actgctggga cctagtttgg aagacttgtt tgacttgtgt 420

gacagaacat tttctcttaa aacagttctc atgatagcta tacaactgat ttctcgcatg 480

gaatatgtcc attcaaagaa cttgatatac agagatgtaa aacctgagaa cttcttaata 540

ggacgaccaa gaaacaaaac ccagcaagtt attcacatta tagattttgg tttggcaaag 600

gaatatattg atccggagac aaagaaacac ataccataca gagaacacaa gagccttaca 660

ggaacagcta gatatatgag cataaacaca catttaggaa aagaacaaag tagaagagac 720

gatttagaag ctttaggtca tatgttcatg tattttctga gaggcagtct tccttggcaa 780

ggcttaaagg ctgacacatt aaaggagagg tatcagaaaa ttggagatac aaaacgggct 840

acaccaatag aagtgttatg tgaaaatttt ccagaaatgg caacatatct tcgttatgta 900

agaaggctag atttttttga aaaaccagac tatgaatact taagaaagct ttttactgac 960

ttgtttgatc gaaaaggata tatgtttgat tatgaatatg actggattgg taaacagttg 1020

cctactccag tgggtgcagt tcagcaagat cctgctctgt catcaaacag agaagcacat 1080

caacacagag ataagatgca acaatccaaa aaccagtcgg cagaccacag ggcagcttgg 1140

gactcccagc aggcaaatcc ccaccatttg agagctcacc ttgcagcaga cagacatggt 1200

ggctcggtac aggttgtaag ttctacaaat ggagagttaa acacagatga ccccaccgca 1260

ggacgttcaa atgcacccat cacagcccct actgaagtag aagtgatgga tgaaaccaag 1320

tgctgctgct ttttcaaacg aaggaaaagg aaaaccatac agcgccacaa atgactctgg 1380

acacagacag atcctgggga gttacttaca tgttcatctg ctgtcttgtg attaaaatca 1440

tctctgtagt gaccacgtat attttcaagg actcactctt agaaacaaaa atgtcatact 1500

atcatacttc attttgtggt tgtcttacat tctttttctt ttttttttc tctaatttaa 1560

cctttatgga agctttaaag ttttgtcaaa acatgagtgc tttgcccatc agtgaatgga 1620

atggaccaat gaggtggtat caatgaatat agttccatag aacattttcc agaagttctt 1680

ctgttgtaga aagcagtaca gtatcttaag tgtcaaccag ttatatacct aatctggttt 1740

tttataactt ctgtaagagc ataatcaaac aggaattttc ttttctcagt ggataataca 1800

acagagaaaa cagagttgcc caaatattta aaagaagtta ttccttgaga agttcatatt 1860

ttgtgacatc tgcattgatt tcagtattac tgatggtact gttattcata agtcatatta 1920

acattctctc cgtgaaatca tggtacagtc actgcccaga ggtactgagg aaaagcaata 1980

tgggttcggc agatggtggt ggtaaaatga atcttaagga gtgtggtaaa tatgtgctcc 2040

gcttttgttg catcactatg tgaagtactg tgttgcagaa gtggcaaaag cgcttatttt 2100

taaaaatgca aaatatttgt acaatgtaac tttatgcttc caaataataa tgtatgttag 2160

acagcaagaa atgaatactt taaaaagtga tatatgttgg agttataaag aaatacacta 2220

aggagaggta gtaaatgtga accttgttgc agtgtataag gtggaagcct aaagaaatct 2280

caccgaaact tactgctgaa tgattacatt ctcccttaag cagaaaactt tggatgtgcc 2340

atgcaatggt gtctgtgtaa ttattttgct ctttgattaa aaaaaagacc cccagcaata 2400

aaaagtgggt cactctatgc c 2421

<210> 11
<211> 2453
<212> DNA
<213> Homo sapiens

<400> 11
gaattcaaag tggagtaccg caaacttgat atggaaaata aaaagaaaga caaggacaaa 60

tcagatgata gaatggcacg acctagtggt cgatcgggac acaacactcg aggaactggg 120

tcttcatcgt ctggagtttt aatggttgga cctaacttta gagttggaaa aaaaattgga 180

tgtggcaatt ttggagaatt acgattaggg aaaaatttat acacaaatga atatgtggca 240

attaagttgg agcccatgaa atcaagagca ccacagctac atttggaata cagattctat 300

aagcagttag gatctggaga tggtatacct caagtttact atttcggccc ttgtggtaaa 360

tacaatgcta tggtgctgga actgctggga cctagtttgg aagacttgtt tgacttgtgt 420

gacagaacat tttctcttaa aacagttctc atgatagcta tacaactgat ttctcgcatg 480

gaatatgtcc attcaaagaa cttgatatac agagatgtaa aacctgagaa cttcttaata 540

ggacgaccaa gaaacaaaac ccagcaagtt attcacatta tagattttgg tttggcaaag 600

gaatatattg atccggagac aaagaaacac ataccataca gagaacacaa gagccttaca 660

ggaacagcta gatatatgag cataaacaca catttaggaa aagaacaaag tagaagagac 720

gatttagaag ctttaggtca tatgttcatg tattttctga gaggcagtct tccttggcaa 780

ggcttaaagg ctgacacatt aaaggagagg tatcagaaaa ttggagatac aaaacgggct 840

acaccaatag aagtgttatg tgaaaatttt ccagaaatgg caacatatct tcgttatgta 900

agaaggctag attttttga aaaaccagac tatgaatact taagaaagct ttttactgac 960

ttgtttgatc gaaaaggata tatgtttgat tatgaatatg actggattgg taaacagttg 1020

cctactccag tgggtgcagt tcagcaagat cctgctctgt catcaaacag agaagcacat 1080

caacacagag ataagatgca acaatccaaa aaccagtcgg cagaccacag ggcagcttgg 1140

gactcccagc aggcaaatcc ccaccatttg agagctcacc ttgcagcaga cagacatggt 1200

ggctcggtac aggttgtaag ttctacaaat ggagagttaa acacagatga ccccaccgca 1260

ggacgttcaa atgcacccat cacagcccct actgaagtag aagtgatgga tgaaaccaac 1320

tgccagaaag tgttgaacat gtggtgctgc tgctttttca aacgaaggaa aaggaaaacc 1380

atacagcgcc acaaatgact ctggacacag acagatcctg gggagttact tacatgttca 1440

tctgctgtct tgtgattaaa atcatctctg tagtgaccac gtatattttc aaggactcac 1500

tcttagaaac aaaaatgtca tactatcata cttcattttg tggttgtctt acattctttt 1560

tctttttttt tttctctaat ttaaccttta tggaagcttt aaagttttgt caaaacatga 1620

gtgctttgcc catcagtgaa tggaatggac caatgaggtg gtatcaatga atatagttcc 1680

atagaacatt ttccagaagt tcttctgttg tagaaagcag tacagtatct taagtgtcaa 1740

ccagttatat acctaatctg gtttttttata acttctgtaa gagcataatc aaacaggaat 1800

tttcttttct cagtggataa tacaacagag aaaacagagt tgcccaaata tttaaaagaa 1860

gttattcctt gagaagttca tattttgtga catctgcatt gatttcagta ttactgatgg 1920

tactgttatt cataagtcat attaacattc tctccgtgaa atcatggtac agtcactgcc 1980

cagaggtact gaggaaaagc aatatgggtt cggcagatgg tggtggtaaa atgaatctta 2040

aggagtgtgg taaatatgtg ctccgctttt gttgcatcac tatgtgaagt actgtgttgc 2100

agaagtggca aaagcgctta tttttaaaaa tgcaaaatat ttgtacaatg taactttatg 2160

cttccaaata ataatgtatg ttagacagca agaaatgaat actttaaaaa gtgatatatg 2220

ttggagttat aaagaaatac actaaggaga ggtagtaaat gtgaaccttg ttgcagtgta 2280

taaggtggaa gcctaaagaa atctcaccga aacttactgc tgaatgatta cattctccct 2340

taagcagaaa actttggatg tgccatgcaa tggtgtctgt gtaattattt tgctctttga 2400

ttaaaaaaaa gacccccagc aataaaaagt gggtcactct atgccctctg tgc 2453

<210> 12
<211> 2717
<212> DNA
<213> Homo sapiens

<400> 12
gctttcctcc tccggccgcc ggcgggtgtg atgtgccgcc gccgctgccc ccgccggcgc 60

tgacggactg ggggcgccgc ccgcgcccgg gaccgacccc tctgctcgcg gccgcgcctt 120

tgagctctct atcaatatca gctcacatca ttgaaaagat aattttgaag acatgttttg 180

ctgaaaagac actaagaaaa attttacgaa tgggatgaac atgctccagt taattgacta 240

cctactgcaa tttgaatgtt aacattaccc atctggtaca gttacctagt gatgtaccta 300

ttttcacaat accctgtttc agtgtgcttg tcttgattaa agaattcaaa gtggagtacc 360

gcaaacttga tatggaaaat aaaaagaaag acaaggacaa atcagatgat agaatggcac 420

gacctagtgg tcgatcggga cacaacactc gaggaactgg gtcttcatcg tctggagttt 480

taatggttgg acctaacttt agagttggaa aaaaaattgg atgtggcaat tttggagaat 540

tacgattagg gaaaaattta tacacaaatg aatatgtggc aattaagttg gagcccatga 600

aatcaagagc accacagcta catttggaat acagattcta tgagcagtta ggatctggag 660

atggtatacc tcaagtttac tatttcggcc cttgtggtaa atacaatgct atggtgctgg 720

aactgctggg acctagtttg gaagacttgt ttgacttgtg tgacagaaca ttttctctta 780

aaacagttct catgatagct atacaactga tttctcgcat ggaatatgtc cattcaaaga 840

acttgatata cagagatgta aaacctgaga acttcttaat aggacgacca ggaaacaaaa 900

cccagcaagt tattcacatt atagattttg gtttggcaaa ggaatatatt gatccggaga 960

caaagaaaca cataccatac agagaacaca agagccttac aggaacagct agatatatga 1020

gcataaacac acatttagga aaagaacaaa gtagaagaga cgatttagaa gctttaggtc 1080

atatgttcat gtattttctg agaggcagtc ttccttggca aggcttaaag gctgacacat 1140

taaaggagag gtatcagaaa attggagata caaaacgggc tacaccaata gaagtgttat 1200

gtgaaaattt tccagaaatg gcaacatatc ttcgttatgt aagaaggcta gatttttttg 1260

aaaaaccaga ctatgactac ttaagaaagc tttttactga cttgtttgat cgaaaaggat 1320

atatgtttga ttatgaatat gactggattg gtaaacagtt gcctactcca gtgggtgcag 1380

ttcagcaaga tcctgctctg tcatcaaaca gagaagcaca tcaacacaga gataagatgc 1440

aacaatccaa aaaccaggtt gtaagttcta caaatggaga gttaaacaca gatgacccca 1500

ccgcaggacg ttcaaatgca cccatcacag cccctactga agtagaagtg atggatgaaa 1560

ccaactgcca gaaagtgttg aacatgtggt gctgctgttt tttcaaacga aggaaaagga 1620

aaaccataca gcgccacaaa tgactctgga cacagacaga tcctggggag ttacttacat 1680

gttcatctgc tgtcttgtga ttaaaatcat ctctgtagtg accacgtata ttttcaagga 1740

ctcactctta gaaacaaaaa tgtcatactt tcatacttca ttttgtggtt gtcttacatt 1800

ctttttcttt ttttttttct ctctaattta acatttatgg aagctttaaa gttttgtcaa 1860

aacatgagtg ctttgcccat cagtgaatgg aatggaccaa tgaggtggta tcaatgaata 1920

tagttccata gaacattttc cagaagttct tctgttgtag aaagcagtac agtatcttaa 1980

gtgtcaacca gttatatacc taatctggtt ttttataact tctgtaagag cataatcaaa 2040

caggaatttt cttttctcag tggataatac aacagagaaa acagagttgc ccaaatattt 2100

aaaagaagtt attccttgag aagttcatat tttgtgacat ctgcattgat ttcagtatta 2160

ctgatggtac tgttattcat aagtcatatt aacattctct ccgtgaaatc atggtacagt 2220

cactgcccag aggtactgag gaaaaagcaa tatgggttcg gcagatggtg gtggtaaaat 2280

gaatcttaag gagtgtggta aatatgtgct ccgcttttgt tgcatcacta tgtgaagtac 2340

tgtgttgcag aagtggcaaa agcgcttatt tttaaaaatg caaaatattt gtacaatgta 2400

actttatgct tccaaataat aatgtatgtt agacagcaag aaatgaatac tttaaaaagt 2460

gatgtatgtt ggagttataa agaaatacac taaggagagg tagtaaatgt gaaccttgtt 2520

gcagtgtata aggtggaagc ctaaagaaat ctcaccaaaa cttactgctg aatgattaca 2580

ttctccctta agcagaaaac tttggatgtg ccatgcaatg gtgtctgtgt aattattttg 2640

ctctttgatt aaaaaaaaga cccccagcaa taaaaagtgg gtcactctat gccctctgtg 2700

caaaaaaaaa aaaaaaa 2717

<210> 13
<211> 393
<212> PRT
<213> Homo sapiens

<400> 13

Met Asp His Pro Ser Arg Glu Lys Asp Glu Arg Gln Arg Thr Thr Lys
1 5 10 15

Pro Met Ala Gln Arg Ser Ala His Cys Ser Arg Pro Ser Gly Ser Ser
20 25 30

Ser Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val Gly Lys
35 40 45

Lys Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys Asn Leu
50 55 60

Tyr Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Ile Lys Ser Arg
65 70 75 80

Ala Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu Gly Ser
85 90 95

Ala Gly Glu Gly Leu Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly Lys
100 105 110

Tyr Asn Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp Leu
115 120 125

Phe Asp Leu Cys Asp Arg Thr Phe Thr Leu Lys Thr Val Leu Met Ile
130 135 140

Ala Ile Gln Leu Leu Ser Arg Met Glu Tyr Val His Ser Lys Asn Leu
145 150 155 160

Ile Tyr Arg Asp Val Lys Pro Glu Asn Phe Leu Ile Gly Arg Gln Gly
165 170 175

Asn Lys Lys Glu His Val Ile His Ile Ile Asp Phe Gly Leu Ala Lys
180 185 190

Glu Tyr Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu His
195 200 205

Lys Ser Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His Leu
210 215 220

Gly Lys Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His Met
225 230 235 240

Phe Met Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys Ala
245 250 255

Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg Asn
260 265 270

Thr Pro Ile Glu Ala Leu Cys Glu Asn Phe Pro Glu Glu Met Ala Thr
275 280 285

Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp Tyr
290 295 300

Glu Tyr Leu Arg Thr Leu Phe Thr Asp Leu Phe Glu Lys Lys Gly Tyr
305 310 315 320

Thr Phe Asp Tyr Ala Tyr Asp Trp Val Gly Arg Pro Ile Pro Thr Pro
325 330 335

Val Gly Ser Val His Val Asp Ser Gly Ala Ser Ala Ile Thr Arg Glu
340 345 350

Ser His Thr His Arg Asp Arg Pro Ser Gln Gln Gln Pro Leu Arg Asn
355 360 365

Gln Ser Leu Arg Thr Val Thr Ala Glu His Tyr Asp Val Asn Asn Ser
370 375 380

Ala Ile Trp His Arg Gly Arg Gly Thr
385 390

<210> 14
<211> 415
<212> PRT
<213> Homo sapiens

<400> 14

Met Asp Phe Asp Lys Lys Gly Gly Lys Gly Glu Thr Glu Glu Gly Arg

```
1               5                   10                  15

Arg Met Ser Lys Ala Gly Gly Gly Arg Ser Ser His Gly Ile Arg Ser
            20                  25                  30

Ser Gly Thr Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val
        35                  40                  45

Gly Lys Lys Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys
    50                  55                  60

Asn Leu Tyr Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Ile Lys
65                  70                  75                  80

Ser Arg Ala Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu
                85                  90                  95

Ser Ala Thr Glu Gly Val Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly
            100                 105                 110

Lys Tyr Asn Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp
        115                 120                 125

Leu Phe Asp Leu Cys Asp Arg Thr Phe Thr Leu Lys Thr Val Leu Met
    130                 135                 140

Ile Ala Ile Gln Leu Ile Thr Arg Met Glu Tyr Val His Thr Lys Ser
145                 150                 155                 160

Leu Ile Tyr Arg Asp Val Lys Pro Glu Asn Phe Leu Val Gly Arg Pro
                165                 170                 175

Gly Thr Lys Arg Gln His Ala Ile His Ile Ile Asp Phe Gly Leu Ala
            180                 185                 190

Lys Glu Tyr Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu
        195                 200                 205

His Lys Ser Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His
    210                 215                 220

Leu Gly Lys Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His
225                 230                 235                 240

Met Phe Met Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys
                245                 250                 255
```

Ala Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg
260 265 270

Ala Thr Pro Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Glu Met Ala
275 280 285

Thr Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp
290 295 300

Tyr Asp Tyr Leu Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Ser Gly
305 310 315 320

Phe Val Phe Asp Tyr Glu Tyr Asp Trp Ala Gly Lys Pro Leu Pro Thr
325 330 335

Pro Ile Gly Thr Val His Thr Asp Leu Pro Ser Gln Pro Gln Leu Arg
340 345 350

Asp Lys Thr Gln Pro His Ser Lys Asn Gln Ala Leu Asn Ser Thr Asn
355 360 365

Gly Glu Leu Asn Ala Asp Asp Pro Thr Ala Gly His Ser Asn Ala Pro
370 375 380

Ile Thr Ala Pro Ala Glu Val Glu Val Ala Asp Glu Thr Lys Cys Cys
385 390 395 400

Cys Phe Phe Lys Arg Arg Lys Arg Lys Ser Leu Gln Arg His Lys
405 410 415

<210> 15
<211> 415
<212> PRT
<213> Homo sapiens

<400> 15

Met Asp Phe Asp Lys Lys Gly Gly Lys Gly Glu Thr Glu Glu Gly Arg
1 5 10 15

Arg Met Ser Lys Ala Gly Gly Gly Arg Ser Ser His Gly Ile Arg Ser
20 25 30

Ser Gly Thr Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val
35 40 45

Gly Lys Lys Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys
50 55 60

Asn Leu Tyr Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Ile Lys
65 70 75 80

Ser Arg Ala Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu
85 90 95

Ser Ala Thr Glu Gly Val Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly
100 105 110

Asn Tyr Asn Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp
115 120 125

Leu Phe Asp Leu Cys Asp Arg Thr Phe Thr Leu Lys Thr Val Leu Met
130 135 140

Ile Ala Ile Gln Leu Ile Thr Arg Met Glu Tyr Val His Thr Lys Ser
145 150 155 160

Leu Ile Tyr Arg Asp Val Lys Pro Glu Asn Phe Leu Val Gly Arg Pro
165 170 175

Gly Thr Lys Arg Gln His Ala Ile His Ile Ile Asp Phe Gly Leu Ala
180 185 190

Lys Glu Tyr Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu
195 200 205

His Lys Ser Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His
210 215 220

Leu Gly Lys Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His
225 230 235 240

Met Phe Met Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys
245 250 255

Ala Asp Thr Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg
260 265 270

Ala Thr Pro Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Glu Met Ala
275 280 285

Thr Tyr Leu Arg Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp
290 295 300

Tyr Asp Tyr Leu Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Ser Gly
305 310 315 320

Phe Val Phe Asp Tyr Glu Tyr Asp Trp Ala Gly Lys Pro Leu Pro Thr
325 330 335

Pro Ile Gly Thr Val His Thr Asp Leu Pro Ser Gln Pro Gln Leu Arg
340 345 350

Asp Lys Thr Gln Pro His Ser Lys Asn Gln Ala Leu Asn Ser Thr Asn
355 360 365

Gly Glu Leu Asn Ala Asp Asp Pro Thr Ala Gly His Ser Asn Ala Pro
370 375 380

Ile Thr Ala Pro Ala Glu Val Glu Val Ala Asp Glu Thr Lys Cys Cys
385 390 395 400

Cys Phe Phe Lys Arg Arg Lys Arg Lys Ser Leu Gln Arg His Lys
405 410 415

<210> 16
<211> 447
<212> PRT
<213> Homo sapiens

<400> 16

Met Glu Asn Lys Lys Lys Asp Lys Asp Lys Ser Asp Asp Arg Met Ala
1 5 10 15

Arg Pro Ser Gly Arg Ser Gly His Asn Thr Arg Gly Thr Gly Ser Ser
20 25 30

Ser Ser Gly Val Leu Met Val Gly Pro Asn Phe Arg Val Gly Lys Lys
35 40 45

Ile Gly Cys Gly Asn Phe Gly Glu Leu Arg Leu Gly Lys Asn Leu Tyr
50 55 60

Thr Asn Glu Tyr Val Ala Ile Lys Leu Glu Pro Met Lys Ser Arg Ala
65 70 75 80

Pro Gln Leu His Leu Glu Tyr Arg Phe Tyr Lys Gln Leu Gly Ser Gly
85 90 95

Asp Gly Ile Pro Gln Val Tyr Tyr Phe Gly Pro Cys Gly Lys Tyr Asn
100 105 110

Ala Met Val Leu Glu Leu Leu Gly Pro Ser Leu Glu Asp Leu Phe Asp
115 120 125

Leu Cys Asp Arg Thr Phe Ser Leu Lys Thr Val Leu Met Ile Ala Ile
130 135 140

Gln Leu Ile Ser Arg Met Glu Tyr Val His Ser Lys Asn Leu Ile Tyr
145 150 155 160

Arg Asp Val Lys Pro Glu Asn Phe Leu Ile Gly Arg Pro Arg Asn Lys
165 170 175

Thr Gln Gln Val Ile His Ile Ile Asp Phe Gly Leu Ala Lys Glu Tyr
180 185 190

Ile Asp Pro Glu Thr Lys Lys His Ile Pro Tyr Arg Glu His Lys Ser
195 200 205

Leu Thr Gly Thr Ala Arg Tyr Met Ser Ile Asn Thr His Leu Gly Lys
210 215 220

Glu Gln Ser Arg Arg Asp Asp Leu Glu Ala Leu Gly His Met Phe Met
225 230 235 240

Tyr Phe Leu Arg Gly Ser Leu Pro Trp Gln Gly Leu Lys Ala Asp Thr
245 250 255

Leu Lys Glu Arg Tyr Gln Lys Ile Gly Asp Thr Lys Arg Ala Thr Pro
260 265 270

Ile Glu Val Leu Cys Glu Asn Phe Pro Glu Met Ala Thr Tyr Leu Arg
275 280 285

Tyr Val Arg Arg Leu Asp Phe Phe Glu Lys Pro Asp Tyr Glu Tyr Leu
290 295 300

Arg Lys Leu Phe Thr Asp Leu Phe Asp Arg Lys Gly Tyr Met Phe Asp
305 310 315 320

Tyr Glu Tyr Asp Trp Ile Gly Lys Gln Leu Pro Thr Pro Val Gly Ala
325 330 335

Val Gln Gln Asp Pro Ala Leu Ser Ser Asn Arg Glu Ala His Gln His
340 345 350

Arg Asp Lys Met Gln Gln Ser Lys Asn Gln Ser Ala Asp His Arg Ala

355 360 365

Ala Trp Asp Ser Gln Gln Ala Asn Pro His His Leu Arg Ala His Leu
370 375 380

Ala Ala Asp Arg His Gly Gly Ser Val Gln Val Val Ser Ser Thr Asn
385 390 395 400

Gly Glu Leu Asn Thr Asp Asp Pro Thr Ala Gly Arg Ser Asn Ala Pro
405 410 415

Ile Thr Ala Pro Thr Glu Val Glu Val Met Asp Glu Thr Lys Cys Cys
420 425 430

Cys Phe Phe Lys Arg Arg Lys Arg Lys Thr Ile Gln Arg His Lys
435 440 445

WHAT IS CLAIMED IS:

1. A method of identifying a candidate p21 pathway modulating agent, said method comprising the steps of:

(a) providing an assay system comprising a CSNK1G polypeptide or nucleic acid;

(b) contacting the assay system with a test agent under conditions whereby, but for the presence of the test agent, the system provides a reference activity; and

(c) detecting a test agent-biased activity of the assay system, wherein a difference between the test agent-biased activity and the reference activity identifies the test agent as a candidate p21 pathway modulating agent.

2. The method of Claim 1 wherein the assay system comprises cultured cells that express the CSNK1G polypeptide.

3. The method of Claim 2 wherein the cultured cells additionally have defective p21 function.

4. The method of Claim 1 wherein the assay system includes a screening assay comprising a CSNK1G polypeptide, and the candidate test agent is a small molecule modulator.

5. The method of Claim 4 wherein the assay is a kinase assay.

6. The method of Claim 1 wherein the assay system is selected from the group consisting of an apoptosis assay system, a cell proliferation assay system, an angiogenesis assay system, and a hypoxic induction assay system.

7. The method of Claim 1 wherein the assay system includes a binding assay comprising a CSNK1G polypeptide and the candidate test agent is an antibody.

8. The method of Claim 1 wherein the assay system includes an expression assay comprising a CSNK1G nucleic acid and the candidate test agent is a nucleic acid modulator.

9. The method of claim 8 wherein the nucleic acid modulator is an antisense oligomer.

10. The method of Claim 8 wherein the nucleic acid modulator is a PMO.

11. The method of Claim 1 additionally comprising:

(d) administering the candidate p21 pathway modulating agent identified in (c) to a model system comprising cells defective in p21 function and, detecting a phenotypic change in the model system that indicates that the p21 function is restored.

12. The method of Claim 11 wherein the model system is a mouse model with defective p21 function.

13. A method for modulating a p21 pathway of a cell comprising contacting a cell defective in p21 function with a candidate modulator that specifically binds to a CSNK1G polypeptide, whereby p21 function is restored.

14. The method of Claim 13 wherein the candidate modulator is administered to a vertebrate animal predetermined to have a disease or disorder resulting from a defect in p21 function.

15. The method of Claim 13 wherein the candidate modulator is selected from the group consisting of an antibody and a small molecule.

16. The method of Claim 1, comprising the additional steps of:

(e) providing a secondary assay system comprising cultured cells or a non-human animal expressing CSNK1G ,

(f) contacting the secondary assay system with the test agent of (b) or an agent derived therefrom under conditions whereby, but for the presence of the test agent or agent derived therefrom, the system provides a reference activity; and

(g) detecting an agent-biased activity of the second assay system,

wherein a difference between the agent-biased activity and the reference activity of the second assay system confirms the test agent or agent derived therefrom as a candidate p21 pathway modulating agent,

and wherein the second assay detects an agent-biased change in the p21 pathway.

17. The method of Claim 16 wherein the secondary assay system comprises cultured cells.

18. The method of Claim 16 wherein the secondary assay system comprises a non-human animal.

19. The method of Claim 18 wherein the non-human animal mis-expresses a p21 pathway gene.

20. A method of modulating p21 pathway in a mammalian cell comprising contacting the cell with an agent that specifically binds a CSNK1G polypeptide or nucleic acid.

21. The method of Claim 20 wherein the agent is administered to a mammalian animal predetermined to have a pathology associated with the p21 pathway.

22. The method of Claim 20 wherein the agent is a small molecule modulator, a nucleic acid modulator, or an antibody.

23. A method for diagnosing a disease in a patient comprising:

(a) obtaining a biological sample from the patient;

(b) contacting the sample with a probe for CSNK1G expression;

(c) comparing results from step (b) with a control;

(d) determining whether step (c) indicates a likelihood of disease.

24. The method of Claim 23 wherein said disease is cancer.

25. The method according to Claim 24, wherein said cancer is a cancer as shown in Table 1 as having >25% expression level.