

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017386417 B2**

(54) Title
1H-pyrrole-2,5-dione compounds and methods of using them to induce self-renewal of stem/progenitor supporting cells

(51) International Patent Classification(s)
C07D 487/04 (2006.01) **A61P 1/04** (2006.01)
A61K 31/437 (2006.01) **A61P 43/00** (2006.01)
A61K 31/5517 (2006.01) **C07D 519/00** (2006.01)

(21) Application No: **2017386417** (22) Date of Filing: **2017.12.21**

(87) WIPO No: **WO18/125746**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/484,282	2017.04.11	US
62/441,060	2016.12.30	US

(43) Publication Date: **2018.07.05**

(44) Accepted Journal Date: **2022.07.14**

(71) Applicant(s)
Frequency Therapeutics, Inc.

(72) Inventor(s)
Loose, Christopher;Tait, Bradley;Manchanda, Rajesh;McLean, Will

(74) Agent / Attorney
Pizzeys Patent and Trade Mark Attorneys Pty Ltd, GPO Box 1374, Brisbane, QLD, 4001, AU

(56) Related Art
WO 2016/037016 A1
WO 2009/006043 A2

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
05 July 2018 (05.07.2018)



(10) International Publication Number
WO 2018/125746 A1

(51) International Patent Classification:

C07D 487/04 (2006.01) *C07D 519/00* (2006.01)
A61K 31/437 (2006.01) *A61P 1/04* (2006.01)
A61K 31/5517 (2006.01) *A61P 43/00* (2006.01)

Published:

— with international search report (Art. 21(3))

(21) International Application Number:

PCT/US2017/067885

(22) International Filing Date:

21 December 2017 (21.12.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/441,060 30 December 2016 (30.12.2016) US
62/484,282 11 April 2017 (11.04.2017) US

(71) Applicant: **FREQUENCY THERAPEUTICS, INC.**
[US/US]; 19 Presidential Way, Woburn, Massachusetts
01801 (US).

(72) Inventors: **LOOSE, Christopher**; 6 Wendell St., Win-
chester, Massachusetts 01890 (US). **TAIT, Bradley**; 22
Bartlett St., Malden, Massachusetts 02148 (US). **MAN-
CHANDA, Rajesh**; 30 Conant St., Acton, Massachusetts
01720 (US). **MCLEAN, Will**; 155 Bayard Ave., North
Haven, Connecticut 06473 (US).

(74) Agent: **FARMER, J. Dean**; Cooley LLP, 1299 Pennsylv-
ania Avenue, Suite 700, Washington, District of Columbia
20004 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,
KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(54) Title: 1H-PYRROLE-2,5-DIONE COMPOUNDS AND METHODS OF USING THEM TO INDUCE SELF-RENEWAL OF STEM/PROGENITOR SUPPORTING CELLS

(57) Abstract: The present invention relates to 1H-pyrrole-2,5-dione compounds and methods of using them to induce self-renewal of stem/progenitor supporting cells, including inducing the stem/progenitor cells to proliferate while maintaining, in the daughter cells, the capacity to differentiate into tissue cells.



WO 2018/125746 A1

1H-PYRROLE-2,5-DIONE COMPOUNDS AND METHODS OF USING THEM TO INDUCE SELF-RENEWAL OF STEM/PROGENITOR SUPPORTING CELLS

RELATED APPLICATION

[0001] This application claims priority to U.S. Application Nos. 62/484,282, filed April 11, 2017, and 62/441,060, filed December 30, 2016, the entire contents of each of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention relates to 1H-pyrrole-2,5-dione compounds and methods of using them to induce self-renewal of stem/progenitor supporting cells, including inducing the stem/progenitor cells to proliferate while maintaining, in the daughter cells, the capacity to differentiate into tissue cells.

BACKGROUND OF THE INVENTION

[0003] Stem cells exhibit an extraordinary ability to generate multiple cell types in the body. Besides embryonic stem cells, tissue specific stem cells serve a critical role during development as well as in homeostasis and injury repair in the adult. Stem cells renew themselves through proliferation as well as generate tissue specific cell types through differentiation. The characteristics of different stem cells vary from tissue to tissue, and are determined by their intrinsic genetic and epigenetic status. However, the balance between self-renewal and differentiation of different stem cells are all stringently controlled. Uncontrolled self-renewal may lead to overgrowth of stem cells and possibly tumor formation, while uncontrolled differentiation may exhaust the stem cell pool, leading to an impaired ability to sustain tissue homeostasis. Thus, stem cells continuously sense their environment and appropriately respond with proliferation, differentiation or apoptosis. It would be desirable to drive regeneration by controlling the timing and extent of stem cell proliferation and differentiation. Controlling the proliferation with small molecules that are cleared over time would allow for control of the timing and extent of stem cell proliferation and differentiation. Remarkably, tissue stem cells from different tissues share a limited number of signaling pathways for the regulation of their

self-renewal and differentiation, albeit in a very context dependent manner. Some of these pathways are the Wnt and GSK3 pathways.

[0004] *Lgr5* is expressed across a diverse range of tissues and has been identified as a biomarker of adult stem cells in a variety of tissues such as the gut epithelia (Barker et al. 2007), kidney, hair follicle, and stomach (Barker et al, 2010; Haegebarth & Clevers, 2009). For example, it was first published in 2011, that mammalian inner ear hair cells are derived from *LGR5*⁺ cells (Chai et al, 2011, Shi et al. 2012). *Lgr5* is a known component of the Wnt/beta-catenin pathway, which has been shown to play major roles in differentiation, proliferation, and inducing stem cell characteristics (Barker et al. 2007).


[0005] Permanent damage to the hair cells of the inner ear results in sensorineural hearing loss, leading to communication difficulties in a large percentage of the population. Hair cells are the receptor cells that transduce the acoustic stimulus. Regeneration of damaged hair cells would provide an avenue for the treatment of a condition that currently has no therapies other than prosthetic devices. Although hair cells do not regenerate in the mammalian cochlea, new hair cells in lower vertebrates are generated from epithelial cells, called supporting cells, that surround hair cells.

[0006] Prior work has focused on transdifferentiation of supporting cells into hair cells through activation or forced expression of genes that lead to hair cell formation, with a particular focus on mechanisms to enhance expression of *Atoh1* (Birmingham et al., 1999; Zheng and Gao, 2000; Izumikawa et al., 2005; Mizutari et al., 2013). Interestingly, cells transduced with *Atoh1* vectors have been shown to acquire vestibular phenotypes (Kawamoto et al., 2003; Huang et al., 2009; Yang et al., 2012, 2013), and lack complete development. As mentioned, upregulating *Atoh1* via gene insertion has been shown to create non-cochlear cell types that behave in a manner that is not found within the native cochlea. In addition, these methods increase hair cell numbers but decrease supporting cell numbers. Since supporting cells are known to have specialized roles (Ramirez-Camancho 2006, Dale and Jagger 2010), loss of these cells could create problems in proper cochlear function.

[0007] Thus, there remains a long-felt need for new compounds that can preserve/promote the function of existing cells after injury.

SUMMARY OF THE INVENTION

[0008] The present disclosure provides compounds of Formula (I),



and pharmaceutically acceptable salts and tautomers thereof, wherein:

Q^1 is CH or N;

Q^2 is C or N;


Q^3 is C or N;

wherein at least one of Q^1 , Q^2 , and Q^3 is N;

R^1 is selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NHC(O) R^{1a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{1a} is C_1 - C_4 alkyl;

R^2 is selected from the group consisting of halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NH(C_1 - C_4 alkyl), -N(C_1 - C_4 alkyl)₂, -NHC(O) R^{2a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{2a} is C_1 - C_4 alkyl;

R^3 is selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NHC(O) R^{3a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{3a} is C_1 - C_4 alkyl;



-Z-W-X-Y- is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$, $-C(R^Z)_2-C(R^W)_2-CH(R^X)-C(R^Y)_2-$,
or $-C(R^W)_2-CH(R^X)-C(R^Y)_2-$;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^Z groups together form C_3 - C_6 cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^W groups together form C_3 - C_6 cycloalkyl or oxo;

or R^Z and R^W together with the carbons to which they are attached form a C_3 - C_6 cycloalkyl;

R^X is selected from the group consisting of $-COR^{X1}$, $-SO_2R^{X1}$, heteroaryl, and $-(C_1$ - C_4 alkylene)- $(C_3$ - C_8 cycloalkyl), and wherein the $-(C_1$ - C_4 alkylene)- $(C_3$ - C_8 cycloalkyl) is optionally substituted with one to four halo on the C_1 - C_4 alkylene;



wherein R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, $-[C(R^{X1a})_2]_p-CN$, $-CF_3$, C_1 - C_4 alkyl, $-(CH_2)_p-OH$, $-[C(R^{X1a})_2]_p-OH$, $-[C(R^{X1a})_2]_p-O-C_1$ - C_4 alkyl, $-NHCOC_1$ - C_4 alkyl, $-CONHC_1$ - C_4 alkyl, $-COH$, $-CO_2H$, $-[C(R^{X1a})_2]_p-COO-C_1$ - C_4 alkyl, $-(CH_2)_p-NH_2$, $-[C(R^{X1a})_2]_p-NH_2$, $-[C(R^{X1a})_2]_p-NH-C_1$ - C_4 alkyl, $-[C(R^{X1a})_2]_p-N-(C_1$ - C_4 alkyl) $_2$; wherein p is 0, 1, 2, or 3; wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, $-CF_3$, and C_1 - C_4 alkyl, or both R^{X1a} groups together form C_3 - C_6 cycloalkyl;

or R^{X1} is $N(R^{X2})_2$ wherein R^{X2} is independently selected from hydrogen, alkyl, substituted alkyl, wherein the alkyl substitution can be halo, heterocycles and substituted heterocycles; each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^Y groups together form C_3 - C_6 cycloalkyl or oxo; and




m is 0, 1, or 2.


[0009] In some embodiments, the compounds of Formula (I) have one or more of the following features:

a) provided that the compound is not




or

b) provided that when Ar is , and  is  or



[0010] The present disclosure provides compounds of Formula (Ia):



and pharmaceutically acceptable salts and tautomers thereof, wherein:

Q^1 is CH or N;

Q^2 is C or N;


Q^3 is C or N;


wherein at least one of Q^1 , Q^2 , and Q^3 is N;

R^1 is selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NHC(O) R^{3a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{1a} is C_1 - C_4 alkyl;

R^2 is selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NH(C_1 - C_4 alkyl), -N(C_1 - C_4 alkyl)₂, -NHC(O) R^{2a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{2a} is C_1 - C_4 alkyl;

R^3 is selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 alkenyl, C_1 - C_4 alkynyl, -CN, -OH, -O- C_1 - C_4 alkyl, -NH₂, -NHC(O) R^{3a} , and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{3a} is C_1 - C_4 alkyl;





; wherein Ar is optionally substituted with deuterium, halo, alkyl, alkoxy, and CN;

Q^7 is selected from S, O, CH_2 , and NR^{Q7} ; wherein R^{Q7} is hydrogen or optionally substituted C_1 - C_4 alkyl;

-Z-W-X-Y- is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$, $-C(R^Z)_2-C(R^W)_2-CH(R^X)-C(R^Y)_2-$, or $-C(R^W)_2-CH(R^X)-C(R^Y)_2-$;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^Z groups together form C_3 - C_6 cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^W groups together form C_3 - C_6 cycloalkyl or oxo;

or R^Z and R^W together with the carbons to which they are attached form a C_3 - C_6 cycloalkyl;

R^X is selected from the group consisting of hydrogen, R^{X1} , $-COR^{X1}$, $-SO_2R^{X1}$, $-(C_1-C_4$ alkylene)- R^{X1} , and wherein the $-(C_1-C_4$ alkylene)- R^{X1} is optionally substituted with one to four halo on the C_1 - C_4 alkylene;


wherein R^{X1} is C_3 - C_8 cycloalkyl, heteroaryl, or heterocyclyl, wherein the heterocyclyl is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, $-[C(R^{X1a})_2]_p-CN$, $-CF_3$, C_1 - C_4 alkyl, $-(CH_2)_p-OH$, $-[C(R^{X1a})_2]_p-OH$, $-[C(R^{X1a})_2]_p-O-C_1-C_4$ alkyl, $-NHCOC_1-C_4$ alkyl, $CONHC_1-C_4$ alkyl, COH , $-CO_2H$, $-[C(R^{X1a})_2]_p-COO-C_1-C_4$ alkyl, $-(CH_2)_p-NH_2$, $-[C(R^{X1a})_2]_p-NH_2$, $-[C(R^{X1a})_2]_p-NH-C_1-C_4$ alkyl, $-[C(R^{X1a})_2]_p-N-(C_1-C_4$ alkyl) $_2$; wherein p is 0, 1, 2, or 3; wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, $-CF_3$, and C_1 - C_4 alkyl, or both R^{X1a} groups together form C_3 - C_6 cycloalkyl;

or R^{X1} is $N(R^{X2})_2$ wherein R^{X2} is independently selected from hydrogen, alkyl, substituted alkyl, wherein the alkyl substitution can be halo, heterocycles and substituted heterocycles;

each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^Y groups together form C_3 - C_6 cycloalkyl or oxo; and

m is 0, 1, or 2.

[0011] The present disclosure provides compounds of Formula (Ib):



and pharmaceutically acceptable salts and tautomers thereof, wherein:

Q^1 is CH or N;

Q^2 is C or N;


Q^3 is C or N;

wherein at least one of Q^1 , Q^2 , and Q^3 is N; and provided that when Q^1 is CH and Q^3 is C, Q^2 is not N;

R^1 is selected from the group consisting of hydrogen, halo, C₁-C₄alkyl, C₁-C₄alkenyl, C₁-C₄alkynyl, -CN, -OH, -O-C₁-C₄alkyl, -NH₂, -NHC(O)R^{3a}, and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{1a} is C₁-C₄alkyl;

R^2 is selected from the group consisting of hydrogen, halo, C₁-C₄alkyl, C₁-C₄alkenyl, C₁-C₄alkynyl, -CN, -OH, -O-C₁-C₄alkyl, -NH₂, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHC(O)R^{2a}, and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{2a} is C₁-C₄alkyl;

R^3 is selected from the group consisting of hydrogen, halo, C₁-C₄alkyl, C₁-C₄alkenyl, C₁-C₄alkynyl, -CN, -OH, -O-C₁-C₄alkyl, -NH₂, -NHC(O)R^{3a}, and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{3a} is C₁-C₄alkyl;



each Q^6 is independently selected from CR^{Q6} and N; wherein R^{Q6} is hydrogen, halo, -CN, lower alkyl, or substituted alkyl;

Q^7 is selected from S, O, CH_2 , and NR^{Q7} ; wherein R^{Q7} is hydrogen or optionally substituted C_1 - C_4 alkyl;

-Z-W-X-Y- is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$, $-C(R^Z)_2-C(R^W)_2-CH(R^X)-C(R^Y)_2-$, or $-C(R^W)_2-CH(R^X)-C(R^Y)_2-$;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^Z groups together form C_3 - C_6 cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C_1 - C_4 alkyl, or both R^W groups together form C_3 - C_6 cycloalkyl or oxo;

or R^Z and R^W together with the carbons to which they are attached form a C_3 - C_6 cycloalkyl;

R^X is selected from the group consisting of hydrogen, R^{X1} , $-COR^{X1}$, $-SO_2R^{X1}$, $-(C_1-C_4alkylene)-R^{X1}$, and wherein the $-(C_1-C_4alkylene)-R^{X1}$ is optionally substituted with one to four halo on the C_1 - C_4 alkylene;

wherein R^{X1} is C_3 - C_8 cycloalkyl, heteroaryl, or heterocyclyl, wherein the heterocyclyl is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, $-[C(R^{X1a})_2]_p-CN$, $-CF_3$, C_1 - C_4 alkyl, $-(CH_2)_p-OH$, $-[C(R^{X1a})_2]_p-OH$,

-[C(R^{X1a})₂]_p-O-C₁-C₄alkyl, -NHCOC₁-C₄alkyl, CONHC₁-C₄alkyl, COH, -CO₂H, -[C(R^{X1a})₂]_p-COO-C₁-C₄alkyl, -(CH₂)_p-NH₂, -[C(R^{X1a})₂]_p-NH₂, -[C(R^{X1a})₂]_p-NH-C₁-C₄alkyl, -[C(R^{X1a})₂]_p-N-(C₁-C₄alkyl)₂; wherein p is 0, 1, 2, or 3; wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, -CF₃, and C₁-C₄alkyl, or both R^{X1a} groups together form C₃-C₆cycloalkyl;

or R^{X1} is N(R^{X2})₂ wherein R^{X2} is independently selected from hydrogen, alkyl, substituted alkyl, wherein the alkyl substitution can be halo, heterocycles and substituted heterocycles;

each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Y groups together form C₃-C₆cycloalkyl or oxo; and

m is 0, 1, or 2.

[0012] In one aspect the present disclosure provides a method for proliferation of stem cells comprising administering to a cell population an effective amount of a composition provided herein. In some embodiments, proliferation occurs in the absence of a notch activator or an HDAC inhibitor.

[0013] Among the various aspects of the present disclosure, therefore, may be noted a method for activating the Wnt pathway in a cell population to increase the capacity of the population for self-renewal, *i.e.*, the capacity for repeated generation of daughter cells with equivalent proliferation and 'cell fate specification' potential, and differentiation, *i.e.*, the capacity for generation of daughter cells specified for differentiation. In one embodiment, the cell population is a cochlear supporting cell population. Preferably, the Wnt pathway is activated upstream of the c-myc gene in members of the population and without any genetic modification of the population. Instead, the Wnt pathway is preferably activated by small molecules that transiently induce such activity. Additionally, the supporting cell population preferably includes supporting cells that are LGR5⁺ and endogenous to the Organ of Corti.

[0014] A further aspect of the present disclosure is a method for inducing the self-renewal of stem/progenitor supporting cells comprised by a cochlear cell population. That is, the stem/progenitor supporting cells are induced to proliferate (*i.e.*, divide and form daughter cells) while maintaining, in the daughter cells, the capacity to differentiate into hair cells. In contrast, if the stem/progenitor supporting cells were merely induced to proliferate (without maintaining multi-potency), the daughter cells would lack the capacity to divide into hair cells. Further, merely enforcing differentiation of a pre-existing stem/progenitor cell population has the

potential to exhaust the stem cell pool. Proliferation is preferably activated by small molecules that transiently induce such activity. Additionally, in certain embodiments the supporting cell population preferably includes supporting cells that are LGR5+ and endogenous to the Organ of Corti.

[0015] In a first aspect methods of using 1H-pyrrole-2,5-dione compounds for inducing the self-renewal of stem/progenitor supporting cells is provided. In some embodiments, 1H-pyrrole-2,5-dione compounds are compounds of Formula (I).

[0016] In certain embodiments, therefore, the present disclosure provides methods to induce self-renewal of a population of supporting cells by activating pathways and mechanisms that are known to be involved in inducing stem cell properties, such as those used to create “induced pluripotent stem cells”. Preferably, the pathways are activated with small molecules. For example, a compound when applied *in vitro* to a supporting cell population induces the population to proliferate to a high degree and in high purity in a Stem Cell Proliferation Assay, and also allows the population to differentiate into a high purity population of a tissue cell in a Stem Cell Differentiation Assay. In one such embodiment, the compound induces and maintains stem cell properties by proliferating to produce stem cells that can divide for many generations and maintain the ability to have a high proportion of the resulting cells differentiate into tissue cells. Further, the proliferating stem cells express stem cell markers which may include one or more of Lgr5, Sox2, Opem1, Phex, lin28, Lgr6, cyclin D1, Msx1, Myb, Kit, Gdnf3, Zic3, Dppa3, Dppa4, Dppa5, Nanog, Esrrb, Rex1, Dnmt3a, Dnmt3b, Dnmt3l, Utf1, Tcf1, Oct4, Klf4, Pax6, Six2, Zic1, Zic2, Otx2, Bmi1, CDX2, STAT3, Smad1, Smad2, smad2/3, smad4, smad5, and smad7.

[0017] In certain embodiments, the disclosure provides a method for expanding a population of cochlear cells in a cochlear tissue comprising a parent population of cells. In this embodiment, the method comprises contacting the cochlear tissue with a stem cell proliferator to form an expanded population of cells in the cochlear tissue, wherein

the stem cell proliferator is capable of (i) forming a proliferation assay final cell population from a proliferation assay initial cell population over a proliferation assay time period in a stem cell proliferation assay and (ii) forming a differentiation assay final cell population from a differentiation assay initial cell population over a differentiation assay time period in a stem cell differentiation assay wherein:

(a) the proliferation assay initial cell population has (i) a proliferation assay initial number of total cells, (ii) a proliferation assay initial number of Lgr5⁺ cells, (iii) a proliferation assay initial number of hair cells, (iv) a proliferation assay initial Lgr5⁺ cell fraction that equals the ratio of the proliferation assay initial number of Lgr5⁺ cells to the proliferation assay initial number of total cells, and (v) a proliferation assay initial hair cell fraction that equals the ratio of the proliferation assay initial number of hair cells to the proliferation assay initial number of total cells;

(b) the proliferation assay final cell population has (i) a proliferation assay final number of total cells, (ii) a proliferation assay final number of Lgr5⁺ cells, (iii) a proliferation assay final number of hair cells, (iv) a proliferation assay final Lgr5⁺ cell fraction that equals the ratio of the proliferation assay final number of Lgr5⁺ cells to the proliferation assay final number of total cells and (v) a proliferation assay final hair cell fraction that equals the ratio of the proliferation assay final number of hair cells to the proliferation assay final number of total cells;

(c) the differentiation assay initial cell population has (i) a differentiation assay initial number of total cells, (ii) a differentiation assay initial number of Lgr5⁺ cells, (iii) a differentiation assay initial number of hair cells, (iv) a differentiation assay initial Lgr5⁺ cell fraction that equals the ratio of the differentiation assay initial number of Lgr5⁺ cells to the differentiation assay initial number of total cells, and (v) a differentiation assay initial hair cell fraction that equals the ratio of the differentiation assay initial number of hair cells to the differentiation assay initial number of total cells;

(d) the differentiation assay final cell population has (i) a differentiation assay final number of total cells, (ii) a differentiation assay final number of Lgr5⁺ cells, (iii) a differentiation assay final number of hair cells, (iv) a differentiation assay final Lgr5⁺ cell fraction that equals the ratio of the differentiation assay final number of Lgr5⁺ cells to the differentiation assay final number of total cells, and (v) a differentiation assay final hair cell fraction that equals the ratio of the differentiation assay final number of hair cells to the differentiation assay final number of total cells;

(e) the proliferation assay final number of Lgr5⁺ cells exceeds the proliferation assay initial number of Lgr5⁺ cells by a factor of at least 10; and

(f) the differentiation assay final number of hair cells is a non-zero number.

The assay described above does not include applying a notch activator or an HDAC inhibitor.

[0018] In certain embodiments, the disclosure provides a method for increasing the cell density of supporting cells in a population of cochlear cells. The method comprises activating pathways and mechanisms that induce stem cell properties in the supporting cells, proliferating the activated supporting cells (while maintaining the multi-potent character of the supporting cells in the newly formed daughter cells) and thereafter allowing (or even inducing) the expanded population to differentiate into hair cells to form an expanded cochlear cell population wherein the cell density of hair cells in the expanded cochlear cell population exceeds the cell density of hair cells in the original (non-expanded) cochlear cell population. In some embodiments, such proliferation occurs in the absence of a notch activator or an HDAC inhibitor. In some embodiments, the supporting cell population is an *in vitro* supporting cell population. In other embodiments, the supporting cell population is an *in vivo* supporting cell population. Additionally, the proliferation stage is preferably controlled to substantially maintain the native organization of the cochlear structure. The proliferation is induced by the compound described herein that transiently induces such activity rather than by induction of c-myc and without any genetic modification of the population. In some embodiments, such proliferation occurs in the absence of a notch activator or an HDAC inhibitor. Additionally, in certain embodiments the supporting cell population preferably includes supporting cells that are LGR5⁺ and endogenous to the Organ of Corti.

[0019] In certain embodiments, the disclosure provides a method for increasing the cell density of Lgr5⁺ supporting cells in a population of cochlear cells. The method comprises activating pathways and mechanisms that induce or maintain stem cell properties in the Lgr5⁺ supporting cells, proliferating the activated Lgr5⁺ supporting cells (while maintaining such stem cell properties) and thereafter allowing (or even inducing) the expanded population to differentiate into hair cells to form an expanded cochlear cell population wherein the cell density of hair cells in the expanded cochlear cell population exceeds the cell density of hair cells in the original (non-expanded) cochlear cell population. In some embodiments for increasing the cell density of Lgr5⁺ supporting cells in a population of cochlear cells, such increasing of the cell density occurs in the absence of a notch activator or an HDAC inhibitor. In some embodiments, the Lgr5⁺ supporting cell population is an *in vitro* Lgr5⁺ stem cell population. In other

embodiments, the Lgr5⁺ supporting cell population is an *in vivo* supporting cell population. Additionally, in certain embodiments the proliferation stage is preferably controlled to substantially maintain the native organization of the cochlear structure.

[0020] In certain embodiments, the disclosure provides a method for increasing the cell density of hair cells in an initial population of cochlear cells, the initial population (which may be an *in vivo* or an *in vitro* population) comprises hair cells, Lgr5⁻ supporting cells, and Lgr5⁺ supporting cells. In some embodiments for increasing the cell density of hair cells in an initial population of cochlear cells, such increasing of the cell density occurs in the absence of a notch activator or an HDAC inhibitor. The method comprises administering to the initial population a compound described herein.

[0021] In certain embodiments, the method produces stem cells in a Stem Cell Proliferation Assay that express stem cells markers Lgr5⁺. In certain embodiments, if a mixed population of Lgr5⁺ and non-Lgr5⁺ stems are placed in a Stem Cell Proliferation Assay, the method increases the fraction of cells in the population that are Lgr5⁺. In some embodiments, such production of stem cells in a Stem Cell Proliferation Assay occurs in the absence of a notch activator or an HDAC inhibitor.

[0022] Expanding supporting cell populations to a degree that destroys the native organization of the cochlear structure could inhibit cochlear function. Driving proliferation of existing supporting cells with a small molecule signal may allow for a more controlled regeneration of hair cells than using gene delivery, which is incapable of targeting a specific cell type and permanently alters a cell's genetic information. An approximately normal cochlear structure is desired with rows of hair cells that have supporting cells between them, and hair cells do not contact other hair cells. Further, it would be desirable to avoid using genetic modification to drive proliferation to create large cell aggregations in the cochlea that disrupt the organ's anatomy.

[0023] In certain embodiments, the disclosure provides a method for increasing the cell density of hair cells in an initial population of cochlear cells comprising hair cells and supporting cells. The method comprises selectively expanding the number of supporting cells in the initial population to form an intermediate cochlear cell population wherein the ratio of the number of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of the number of supporting cells to hair cells in the initial cochlear cell population. The method

further comprises generating hair cells in the intermediate cochlear cell population to form an expanded cochlear cell population wherein the ratio of the number of hair cells to supporting cells in the expanded cochlear cell population exceeds the ratio of the number of hair cells to supporting cells in the intermediate cochlear cell population. In some embodiments, the method does not comprise the use of a notch activator or an HDAC inhibitor.

[0024] In certain embodiments, the disclosure provides a method for increasing the number of Lgr5⁺ supporting cells or increasing the Lgr5⁺ activity in an initial population of cochlear cells, wherein the initial population comprises supporting cells and hair cells. For example, in one such method an intermediate population is formed in which the number of Lgr5⁺ supporting cells is expanded relative to the initial population. Alternatively, in one such method an intermediate population is formed in which the Lgr5⁺ activity of the supporting cells relative to the initial population is increased. Alternatively, a method where the number of Lgr5⁺ cells is increased relative to the initial cell population by activating Lgr5⁺ expression in cell types that normally lack or have very low levels of Lgr5⁺. In some embodiments, these alternative methods do not comprise the use of a notch activator or an HDAC inhibitor. By way of further example, an intermediate population is formed in which the number of Lgr5⁺ supporting cells is expanded and the Lgr5 activity is increased relative to the initial cochlear cell population. Thereafter, hair cells in the intermediate cochlear cell population may be generated to form an expanded cochlear cell population wherein the ratio of hair cells to supporting cells in the expanded cochlear cell population exceeds the ratio of the number of hair cells to supporting cells in the intermediate cochlear cell population.

[0025] In each of the afore mentioned embodiments of the present disclosure, stemness is induced by activating Wnt or inhibiting GSK3 activity. In some embodiments, inducing stemness does not comprise the use of a notch activator or an HDAC inhibitor.

[0026] In certain embodiments, the disclosure provides methods for preventing and treating auditory dysfunction. For example, in certain embodiments, the disclosure provides methods for preventing or treating auditory impairments in a subject comprising administering to said subject an effective amount of a compound provided herein.

[0027] In certain embodiments, the present disclosure also relates to ex-vivo uses of cells described herein. For example, approaches described herein can be used for high through screens and for discovery purposes. For example, certain embodiments of the present disclosure

are useful for identifying agents that proliferate hair cell progenitors and/or increase numbers of hair cells, and also agents that protect supporting cells and/or hair cells (*e.g.* to support their survival), and also for identifying agents that are toxic or not toxic to supporting cells or differentiated progeny including hair cells.

[0028] In certain embodiments, the disclosure provides for methods for inhibiting the loss or death of the cells of the auditory system in a subject comprising administering to said subject an effective amount of the compound described herein or derivative thereof or pharmaceutically acceptable salt thereof and an acceptable carrier or excipient, thereby inhibiting loss or death of the cells of the auditory system in the subject. In some embodiments, the method does not comprise the use of a notch activator or an HDAC inhibitor.

[0029] In certain embodiments, the disclosure provides methods for maintaining or promoting the growth of cells of the auditory system in a subject comprising administering to said subject the compound described herein or derivative thereof or pharmaceutically acceptable salt thereof in an effective amount so as to augment or initiate endogenous repair, thereby maintaining or promoting the growth of cells of the auditory system in the subject.

[0030] Also described herein is a method for expanding a population of cochlear cells in a cochlear tissue comprising a parent population of cells, the parent population including supporting cells and a number of Lgr5⁺ cells, the method comprising contacting the cochlear tissue with a stem cell proliferator to form an expanded population of cells in the cochlear tissue, wherein the stem cell proliferator is capable (i) in a stem cell proliferation assay of increasing the number of Lgr5⁺ cells in a stem cell proliferation assay cell population by a factor of at least 10 and (ii) in a stem cell differentiation assay of forming hair cells from a cell population comprising Lgr5⁺ cells. In some embodiments for expanding a population of cochlear cells, the method does not comprise the use of a notch activator or an HDAC inhibitor.

[0031] Also described herein is a method for expanding a population of cochlear cells in a cochlear tissue comprising a parent population of cells, the parent population including supporting cells, the method comprising contacting the cochlear tissue with a stem cell proliferator to form an expanded population of cells in the cochlear tissue. The stem cell proliferator can be capable of (i) forming a proliferation assay final cell population from a proliferation assay initial cell population over a proliferation assay time period in a stem cell proliferation assay and (ii) forming a differentiation assay final cell population from a

differentiation assay initial cell population over a differentiation assay time period in a stem cell differentiation assay wherein: (a) the proliferation assay initial cell population has (i) a proliferation assay initial number of total cells, (ii) a proliferation assay initial number of Lgr5⁺ cells, (iii) a proliferation assay initial number of hair cells, (iv) a proliferation assay initial Lgr5⁺ cell fraction that equals the ratio of the proliferation assay initial number of Lgr5⁺ cells to the proliferation assay initial number of total cells, and (v) a proliferation assay initial hair cell fraction that equals the ratio of the proliferation assay initial number of hair cells to the proliferation assay initial number of total cells; (b) the proliferation assay final cell population has (i) a proliferation assay final number of total cells, (ii) a proliferation assay final number of Lgr5⁺ cells, (iii) a proliferation assay final number of hair cells, (iv) a proliferation assay final Lgr5⁺ cell fraction that equals the ratio of the proliferation assay final number of Lgr5⁺ cells to the proliferation assay final number of total cells and (v) a proliferation assay final hair cell fraction that equals the ratio of the proliferation assay final number of hair cells to the proliferation assay final number of total cells; (c) the differentiation assay initial cell population has (i) a differentiation assay initial number of total cells, (ii) a differentiation assay initial number of Lgr5⁺ cells, (iii) a differentiation assay initial number of hair cells, (iv) a differentiation assay initial Lgr5⁺ cell fraction that equals the ratio of the differentiation assay initial number of Lgr5⁺ cells to the differentiation assay initial number of total cells, and (v) a differentiation assay initial hair cell fraction that equals the ratio of the differentiation assay initial number of hair cells to the differentiation assay initial number of total cells; (d) the differentiation assay final cell population has (i) a differentiation assay final number of total cells, (ii) a differentiation assay final number of Lgr5⁺ cells, (iii) a differentiation assay final number of hair cells, (iv) a differentiation assay final Lgr5⁺ cell fraction that equals the ratio of the differentiation assay final number of Lgr5⁺ cells to the differentiation assay final number of total cells, and (v) a differentiation assay final hair cell fraction that equals the ratio of the differentiation assay final number of hair cells to the differentiation assay final number of total cells; (e) the proliferation assay final number of Lgr5⁺ cells exceeds the proliferation assay initial number of Lgr5⁺ cells by a factor of at least 10; and (f) the differentiation assay final number of hair cells is a non-zero number. In some embodiments of the assay described above, the assay does not comprise the use of a notch activator or an HDAC inhibitor.

[0032] The proliferation assay final number of Lgr5⁺ cells can be greater than the proliferation assay initial number of Lgr5⁺ cells by a factor of at least 50, or by a factor of at least 100. The expanded population of cells in the cochlear tissue can include a greater number of hair cells than does the parent population. The proliferation assay final Lgr5⁺ cell fraction can be greater than the differentiation assay initial Lgr5⁺ cell fraction by at least a factor of 2. The differentiation assay final hair cell fraction can be greater than the proliferation assay initial hair cell fraction by at least a factor of 2. The proliferation assay final hair cell fraction can be at least 25% less than the proliferation assay initial hair cell fraction. The proliferation assay final Lgr5⁺ cell fraction can be at least 10% greater than proliferation assay initial Lgr5⁺ cell fraction. One or more morphological characteristics of the cochlear tissue can be maintained. Native morphology can be maintained. The stem cell proliferator can be dispersed in a biocompatible matrix, which can be a biocompatible gel or foam. The cochlear tissue can be an in vivo cochlear tissue or an ex vivo cochlear tissue. The method can produce a population of Lgr5⁺ cells that are in s-phase. The cochlear tissue can be in a subject, and contacting the cochlear tissue with the compound can be achieved by administering the compound trans-tympanically to the subject. Contacting the cochlear tissue with the compound can result in improved auditory functioning of the subject.

[0033] Also described herein is a method of treating a subject who has, or is at risk of developing, hearing loss. The method can include trans-tympanically administering to a cochlear tissue of the subject compound provided herein.

[0034] Also described herein is a method of generating Myo7a⁺ cochlear cells. The method can include contacting Lgr5⁺ cochlear cells with a compound provided herein, thereby generating an expanded population of Lgr5⁺ cells, thereby generating Myo7a⁺ cochlear cells.

[0035] Other objects and features will be in part apparent and in part pointed out hereinafter.

Definitions

[0036] In this application, the use of “or” means “and/or” unless stated otherwise. As used in this application, the term “comprise” and variations of the term, such as “comprising” and “comprises,” are not intended to exclude other additives, components, integers or steps. As used in this application, the terms “about” and “approximately” are used as equivalents. Any numerals used in this application with or without about/approximately are meant to cover any normal fluctuations appreciated by one of ordinary skill in the relevant art. In certain

embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0037] “Administration” refers to introducing a substance into a subject. In some embodiments, administration is auricular, intraauricular, intracochlear, intravestibular, or transtympanically, *e.g.*, by injection. In some embodiments, administration is directly to the inner ear, *e.g.* injection through the round or oval, otic capsule, or vestibular canals. In some embodiments, administration is directly into the inner ear via a cochlear implant delivery system. In some embodiments, the substance is injected transtympanically to the middle ear. In certain embodiments “causing to be administered” refers to administration of a second component after a first component has already been administered (*e.g.*, at a different time and/or by a different actor).

[0038] An “antibody” refers to an immunoglobulin polypeptide, or fragment thereof, having immunogen binding ability.

[0039] As used herein, an “agonist” is an agent that causes an increase in the expression or activity of a target gene, protein, or a pathway, respectively. Therefore, an agonist can bind to and activate its cognate receptor in some fashion, which directly or indirectly brings about this physiological effect on the target gene or protein. An agonist can also increase the activity of a pathway through modulating the activity of pathway components, for example, through inhibiting the activity of negative regulators of a pathway. Therefore, a “Wnt agonist” can be defined as an agent that increases the activity of Wnt pathway, which can be measured by increased TCF/LEF-mediated transcription in a cell. Therefore, a “Wnt agonist” can be a true Wnt agonist that bind and activate a Frizzled receptor family member, including any and all of the Wnt family proteins, an inhibitor of intracellular beta-catenin degradation, and activators of TCF/LEF.

[0040] An “antagonist” refers to an agent that binds to a receptor, and which in turn decreases or eliminates binding by other molecules.

[0041] “Anti-sense” refers to a nucleic acid sequence, regardless of length, that is complementary to the coding strand or mRNA of a nucleic acid sequence. Antisense RNA can

be introduced to an individual cell, tissue or organoid. An anti-sense nucleic acid can contain a modified backbone, for example, phosphorothioate, phosphorodithioate, or other modified backbones known in the art, or may contain non-natural internucleoside linkages.

[0042] As referred to herein, a “complementary nucleic acid sequence” is a nucleic acid sequence capable of hybridizing with another nucleic acid sequence comprised of complementary nucleotide base pairs. By “hybridize” is meant pair to form a double-stranded molecule between complementary nucleotide bases (e.g., adenine (A) forms a base pair with thymine (T), as does guanine (G) with cytosine (C) in DNA) under suitable conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507).

[0043] “Auricular administration” refers to a method of using a catheter or wick device to administer a composition across the tympanic membrane to the inner ear of the subject. To facilitate insertion of the wick or catheter, the tympanic membrane may be pierced using a suitably sized syringe or pipette. The devices could also be inserted using any other methods known to those of skill in the art, e.g., surgical implantation of the device. In particular embodiments, the wick or catheter device may be a stand-alone device, meaning that it is inserted into the ear of the subject and then the composition is controllably released to the inner ear. In other particular embodiments, the wick or catheter device may be attached or coupled to a pump or other device that allows for the administration of additional compositions. The pump may be automatically programmed to deliver dosage units or may be controlled by the subject or medical professional.

[0044] “Biocompatible Matrix” as used herein is a polymeric carrier that is acceptable for administration to humans for the release of therapeutic agents. A Biocompatible Matrix may be a biocompatible gel or foam.

[0045] “Cell Aggregate” as used herein shall mean a body cells in the Organ of Corti that have proliferated to form a cluster of a given cell type that is greater than 40 microns in diameter and/or produced a morphology in which greater than 3 cell layers reside perpendicular to the basilar membrane. A “Cell Aggregate” can also refer a process in which cell division creates a body of cells that cause one or more cell types to breach the reticular lamina, or the boundary between endolymph and perilymph

[0046] “Cell Density” as used herein in connection with a specific cell type is the mean number of that cell type per area in a Representative Microscopy Sample. The cell types may include but are not limited to Lgr5⁺ cells, hair cells, or supporting cells. The Cell Density may be assessed with a given cell type in a given organ or tissue, including but not limited to the cochlea or Organ of Corti. For instance, the Lgr5⁺ Cell Density in the Organ of Corti is the Cell Density of Lgr5⁺ cells as measured across the Organ of Corti. Typically, supporting cells and Lgr5⁺ cells will be enumerated by taking cross sections of the Organ of Corti. Typically, hair cells will be enumerated by looking down at the surface of the Organ of Corti, though cross sections may be used in some instances, as described in a Representative Microscopy Sample. Typically, Cell Density of Lgr5⁺ cells will be measured by analyzing whole mount preparations of the Organ of Corti and counting the number of Lgr5 cells across a given distance along the surface of the epithelia, as described in a Representative Microscopy Sample. Hair cells may be identified by their morphological features such as bundles or hair cell specific stains (*e.g.*, Myosin VIIa, Prestin, vGlut3, Pou4f3, Espin, conjugated-Phalloidin, PMCA2, Ribeye, Atoh1, etc). Lgr5⁺ cells may be identified by specific stains or antibodies (*e.g.* Lgr5-GFP transgenic reporter, anti-Lgr5 antibody, *etc.*)

[0047] “Cochlear Concentration” as used herein will be the concentration of a given agent as measured through sampling cochlear fluid. Unless otherwise noted, the sample should contain a substantial enough portion of the cochlear fluid so that it is approximately representative of the average concentration of the agent in the cochlea. For example, samples may be drawn from a vestibular canal, and a series of fluid samples drawn in series such that individual samples are comprised of cochlear fluid in specified portions of the cochlea

[0048] “Complementary nucleic acid sequence” refers to a nucleic acid sequence capable of hybridizing with another nucleic acid sequence comprised of complementary nucleotide base pairs.

[0049] “Cross-Sectional Cell Density” as used herein in connection with a specific cell type is the mean number of that cell type per area of cross section through a tissue in a Representative Microscopy Sample. Cross sections of the Organ of Corti can also be used to determine the number of cells in a given plane. Typically, hair cells Cross-sectional Cell Density will be measured by analyzing whole mount preparations of the Organ of Corti and counting the number of hair cells across a given distance in cross sections taken along a portion of the epithelia, as

described in a Representative Microscopy Sample. Typically, Cross-sectional Cell Density of Lgr5⁺ cells will be measured by analyzing whole mount preparations of the Organ of Corti and counting the number of Lgr5⁺ cells across a given distance in cross sections taken along a portion of the epithelia, as described in a Representative Microscopy Sample. Hair cells may be identified by their morphological features such as bundles or hair cell specific stains (suitable stains include *e.g.*, Myosin VIIa, Prestin, vGlut3, Pou4f3, conjugated-Phalloidin, PMCA2, Atoh1, *etc.*). Lgr5⁺ cells may be identified by specific stains or antibodies (suitable stains and antibodies include fluorescence *in situ* hybridization of Lgr5 mRNA, Lgr5-GFP transgenic reporter system, anti-Lgr5 antibodies, *etc.*).

[0050] “Decreasing” refers to decreasing by at least 5%, for example, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 99 or 100%, for example, as compared to the level of reference.

[0051] “Decreases” also means decreases by at least 1-fold, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 1000-fold or more, for example, as compared to the level of a reference.

[0052] “Differentiation Period” as used herein is the duration of time in which there is an Effective Stemness Driver Concentration without an Effective Differentiation Inhibition Concentration.

[0053] “Effective Concentration” may be the Effective Stemness Driver Concentration for a Stemness Driver or the Effective Diffusion Inhibition Concentration for a Diffusion Inhibitor.

[0054] “Effective Differentiation Inhibition Concentration” is the minimum concentration of a Differentiation Inhibitor that does not allow more than a 50% increase in the fraction of the total population of cells that are hair cells at the end of the Stem Cell Proliferation Assay compared to the start of the Stem Cell Proliferation Assay. In measuring the Effective Differentiation Inhibition Concentration, a Hair Cell stain for cells may be used with flow cytometry to quantify hair cells for a mouse strain that is not an Atoh1-GFP mouse. Alternatively, an Atoh1-GFP mouse strain may be used.

[0055] “Effective Release Rate” (mass/time) as used herein is the Effective Concentration (mass/volume) * 30 uL / 1 hour.

[0056] “Effective Stemness Driver Concentration” is the minimum concentration of a Stemness Driver that induces at least 1.5-fold increase in number of LGR5⁺ cells in a Stem Cell

Proliferation Assay compared to the number of Lgr5+ cells in a Stem Cell Proliferation Assay performed without the Stemness Driver and with all other components present at the same concentrations.

[0057] “Eliminate” means to decrease to a level that is undetectable.

[0058] “Engraft” or “engraftment” refers to the process of stem or progenitor cell incorporation into a tissue of interest *in vivo* through contact with existing cells of the tissue. “Epithelial progenitor cell” refers to a multipotent cell which has the potential to become restricted to cell lineages resulting in epithelial cells.

[0059] “Epithelial stem cell” refers to a multipotent cell which has the potential to become committed to multiple cell lineages, including cell lineages resulting in epithelial cells.

[0060] “Fragment” refers to a portion of a polypeptide or nucleic acid molecule. This portion contains, preferably, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the entire length of the reference nucleic acid molecule or polypeptide. A fragment may contain 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or amino acids.

[0061] “GSK3 inhibitor” is a composition that inhibits the activity of GSK3, GSK-3alpha, and/or GSK-3beta.

[0062] “GSK3beta,” “GSK3 β ,” and “GSK3B” as used interchangeably herein are acronyms for glycogen synthase kinase 3 beta,

[0063] “GSK3beta inhibitor” is a composition that inhibits the activity of GSK3beta.

[0064] “Hybridize” refers to pairing to form a double-stranded molecule between complementary nucleotide bases (e.g., adenine (A) forms a base pair with thymine (T), as does guanine (G) with cytosine (C) in DNA) under suitable conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507).

[0065] An “inhibitor” refers to an agent that causes a decrease in the expression or activity of a target gene or protein, respectively. An “antagonist” can be an inhibitor, but is more specifically an agent that binds to a receptor, and which in turn decreases or eliminates binding by other molecules.

[0066] As used herein, an “inhibitory nucleic acid” is a double-stranded RNA, RNA interference, miRNA, siRNA, shRNA, or antisense RNA, or a portion thereof, or a mimetic

thereof, that when administered to a mammalian cell results in a decrease in the expression of a target gene. Typically, a nucleic acid inhibitor comprises at least a portion of a target nucleic acid molecule, or an ortholog thereof, or comprises at least a portion of the complementary strand of a target nucleic acid molecule. Typically, expression of a target gene is reduced by 10%, 25%, 50%, 75%, or even 90-100%.

[0067] “In Vitro Lgr5 activity” refers to the level of expression or activity of Lgr5 in an in vitro population of cells. It may be measured, for example, in cells derived from a Lgr5-GFP expressing mouse such as a B6.129P2-Lgr5^{tm1(cre/ERT2)C1e/J} mouse (also known as Lgr5-EGFP-IRES-creERT2 or Lgr5-GFP mouse, Jackson Lab Stock No: 008875) by dissociating cells to single cells, staining with propidium iodide (PI), and analyzing the cells using a flow cytometer for Lgr5-GFP expression. Inner ear epithelial cells from wild-type (non-Lgr5-GFP) mice that passing the same culturing and analyzing procedures can be used as a negative control. Typically, two population of cells are shown in the bivariate plot with GFP/FITC as one variable, which include both GFP positive and GFP negative populations. Lgr5-positive cells are identified by gating GFP positive cell population. The percentage of Lgr5-positive cells are measured by gating GFP positive cell population against both GFP negative population and the negative control. The number of Lgr5-positive cells is calculated by multiplying the total number of cells by the percentage of Lgr5-positive cells. For cells derived from non-Lgr5-GFP mice, Lgr5 activity can be measured using an anti-Lgr5 antibody or quantitative-PCR on the Lgr5 gene.

[0068] “*In Vivo* Lgr5 activity” as used herein is the level of expression or activity of Lgr5 in a subject. It may be measured, for example, by removing an animal’s inner ear and measuring Lgr5 protein or Lgr5 mRNA. Lgr5 protein production can be measured using an anti-Lgr5 antibody to measure fluorescence intensity as determined by imaging cochlear samples, where fluorescence intensity is used as a measure of Lgr5 presence. Western blots can be used with an anti-Lgr5 antibody, where cells can be harvested from the treated organ to determine increases in Lgr5 protein. Quantitative-PCR or RNA in situ hybridization can be used to measure relative changes in Lgr5 mRNA production, where cells can be harvested from the inner ear to determine changes in Lgr5 mRNA. Alternatively, Lgr5 expression can be measured using an Lgr5 promoter driven GFP reporter transgenic system, where the presence or intensity GFP fluoresce can be directly detected using flow cytometry, imaging, or indirectly using an anti-GFP antibody.

[0069] “Increases” also means increases by at least 1-fold, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 500, 1000-fold or more, for example, as compared to the level of a as compared to the level of a reference standard.

[0070] “Increasing” refers to increasing by at least 5%, for example, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 99, 100% or more, for example, as compared to the level of a reference.

[0071] “Intraauricular administration” refers to administration of a composition to the middle or inner ear of a subject by directly injecting the composition.

[0072] “Intracochlear” administration refers to direct injection of a composition across the tympanic membrane and across the round or oval membrane into the cochlea.

[0073] “Intravestibular” administration refers to direct injection of a composition across the tympanic membrane and across the round or oval membrane into the vestibular organs.

[0074] “Isolated” refers to a material that is free to varying degrees from components which normally accompany it as found in its native state. “Isolate” denotes a degree of separation from original source or surroundings.

[0075] “Lgr5” is an acronym for the Leucine-rich repeat-containing G-protein coupled receptor 5, also known as G-protein coupled receptor 49 (GPR49) or G-protein coupled receptor 67 (GPR67). It is a protein that in humans is encoded by the Lgr5 gene.

[0076] “Lgr5 activity” is defined as the level of activity of Lgr5 in a population of cells. In an *in vitro* cell population, Lgr5 activity may be measured in an *in vitro* Lgr5 Activity assay. In an *in vivo* cell population, Lgr5 activity may be measured in an *in vivo* Lgr5 Activity assay.

[0077] “Lgr5⁺ cell” or “Lgr5-positive cell” as used herein is a cell that expresses Lgr5. “Lgr5⁻ cell” as used herein is a cell that is not Lgr5⁺.

[0078] “Lineage Tracing” as used herein is using a mouse line that enables fate tracing of any cell that expresses a target gene at the time of reporter induction. This can include hair cell or supporting cells genes (Sox2, Lgr5, Myosin VIIa, Pou4f3, etc). For example, lineage tracing may use an Lgr5-EGFP-IRES-creERT2 mouse crossed with a reporter mouse, which upon induction, allows one to trace the fate of cells that expressed Lgr5 at the time of induction. By further example, Lgr5 cells can be isolated into single cells and cultured in a Stem Cell Proliferation Assay to generate colonies, then subsequently differentiated in a Differentiation Assay and analyzed for cell fate by staining for hair cell and/or supporting cell proteins and determining the

reporter colocalization with either hair cell or supporting cell staining to determine the Lgr5 cells' fate. In addition, lineage tracing can be performed in cochlear explants to track supporting cell or hair cell fate within the intact organ after treatment. For example, Lgr5 cell fate can be determined by isolating the cochlea from a Lgr5-EGFP-IRES-creERT2 mouse crossed with a reporter mouse, and inducing the reporter in Lgr5 cells before or during treatment. The organ can then be analyzed for cell fate by staining for hair cell and/or supporting cell proteins and determining the reporter colocalization with either hair cell or supporting cell staining to determine the Lgr5 cells' fate. In addition, lineage tracing can be performed *in vivo* track supporting cell or hair cell fate within the intact organ after treatment. For example, Lgr5 cell fate can be determined inducing a reporter in an Lgr5-EGFP-IRES-creERT2 mouse crossed with a reporter mouse, treating the animal, then isolating the cochlea. The organ can then be analyzed for cell fate by staining for hair cell and/or supporting cell proteins and determining the reporter colocalization with either hair cell or supporting cell staining to determine the Lgr5 cells' fate. Lineage tracing may be performed using alternative reporters of interest as is standard in the art.

[0079] "Mammal" refers to any mammal including but not limited to human, mouse, rat, sheep, monkey, goat, rabbit, hamster, horse, cow or pig.

[0080] "Mean Release Time" as used herein is the time in which one-half of an agent is released into phosphate buffered saline from a carrier in a Release Assay.

[0081] "Native Morphology" as used herein is means that tissue organization largely reflects the organization in a healthy tissue.

[0082] "Non-human mammal", as used herein, refers to any mammal that is not a human.

[0083] As used in relevant context herein, the term "number" of cells can be 0, 1, or more cells.

[0084] "Organ of Corti" as used herein refers to the sensory cells (inner and outer hair cells) of the hearing organ located in the cochlea.

[0085] "Organoid" or "epithelial organoid" refers to a cell cluster or aggregate that resembles an organ, or part of an organ, and possesses cell types relevant to that particular organ.

[0086] "Population" of cells refers to any number of cells greater than 1, but is preferably at least 1×10^3 cells, at least 1×10^4 cells, at least at least 1×10^5 cells, at least 1×10^6 cells, at least 1×10^7 cells, at least 1×10^8 cells, at least 1×10^9 cells, or at least 1×10^{10} cells.

[0087] “Progenitor cell” as used herein refers to a cell that, like a stem cell, has the tendency to differentiate into a specific type of cell, but is already more specific than a stem cell and is pushed to differentiate into its “target” cell.

[0088] “Reference” means a standard or control condition (e.g., untreated with a test agent or combination of test agents).

[0089] “Release Assay” as used herein is a test in which the rate of release of an agent from a Biocompatible Matrix through dialysis membrane to a saline environment. An exemplary Release Assay may be performed by placing 30 microliters of a composition in 1 ml Phosphate Buffered Saline inside saline dialysis bag with a suitable cutoff, and placing the dialysis bag within 10 ml of Phosphate Buffered Saline at 37 °C. The dialysis membrane size may be chosen based on agent size in order to allow the agent being assessed to exit the membrane. For small molecule release, a 3.5-5 kDa cutoff may be used. The Release Rate for a composition may change over time and may be measured in 1 hour increments.

[0090] “Representative Microscopy Sample” as used herein describes a sufficient number of fields of view within a cell culture system, a portion of extracted tissue, or an entire extracted organ that the average feature size or number being measured can reasonably be said to represent the average feature size or number if all relevant fields were measured. For example, in order to assess the hair cell counts at a frequency range on the Organ of Corti, ImageJ software (NIH) can be used to measure the total length of cochlear whole mounts and the length of individual counted segments. The total number of inner hair cells, outer hair cells, and supporting cells can be counted in the entire or fraction of any of the four cochlear segments of 1200–1400 μm (apical, mid-apical, mid-basal, and basal) at least 3 fields of view at 100μm field size would be reasonably considered a Representative Microscopy Sample. A Representative Microscopy sample can include measurements within a field of view, which can be measured as cells per a given distance. A Representative Microscopy sample can be used to assess morphology, such as cell-cell contacts, cochlear architecture, and cellular components (e.g., bundles, synapses).

[0091] “Rosette Patterning” is a characteristic cell arrangement in the cochlea in which <5% hair cells are adjacent to other hair cells.

[0092] The term “sample” refers to a volume or mass obtained, provided, and/or subjected to analysis. In some embodiments, a sample is or comprises a tissue sample, cell sample, a fluid sample, and the like. In some embodiments, a sample is taken from (or is) a subject (e.g., a

human or animal subject). In some embodiments, a tissue sample is or comprises brain, hair (including roots), buccal swabs, blood, saliva, semen, muscle, or from any internal organs, or cancer, precancerous, or tumor cells associated with any one of these. A fluid may be, but is not limited to, urine, blood, ascites, pleural fluid, spinal fluid, and the like. A body tissue can include, but is not limited to, brain, skin, muscle, endometrial, uterine, and cervical tissue or cancer, precancerous, or tumor cells associated with any one of these. In an embodiment, a body tissue is brain tissue or a brain tumor or cancer. Those of ordinary skill in the art will appreciate that, in some embodiments, a “sample” is a “primary sample” in that it is obtained from a source (e.g., a subject); in some embodiments, a “sample” is the result of processing of a primary sample, for example to remove certain potentially contaminating components and/or to isolate or purify certain components of interest.

[0093] “Self-renewal” refers to the process by which a stem cell divides to generate one (asymmetric division) or two (symmetric division) daughter cells with development potentials that are indistinguishable from those of the mother cell. Self-renewal involves both proliferation and the maintenance of an undifferentiated state.

[0094] “siRNA” refers to a double stranded RNA. Optimally, an siRNA is 18, 19, 20, 21, 22, 23 or 24 nucleotides in length and has a 2 base overhang at its 3' end. These dsRNAs can be introduced to an individual cell or culture system. Such siRNAs are used to downregulate mRNA levels or promoter activity.

[0095] “Stem cell” refers to a multipotent cell having the capacity to self-renew and to differentiate into multiple cell lineages.

[0096] “Stem Cell Differentiation Assay” as used herein is an assay to determine the differentiation capacity of stem cells. In an exemplary Stem Cell Differentiation Assay, the number of cells for an initial cell population is harvested from a Atoh1-GFP mouse between the age of 3 to 7 days, by isolating the Organ of Corti sensory epithelium, dissociating the epithelium into single cells, and passing the cells through a 40um cell strainer. Approximately 5000 cells are entrapped in 40 µl of culture substrate (for example: Matrigel (Corning, Growth Factor Reduced)) and placed at the center of wells in a 24-well plate with 500 µl of an appropriate culture media, growth factors and agent being tested. Appropriate culture media and growth factors include Advanced DMEM/F12 with media Supplements (1X N2, 1X B27, 2 mM Glutamax, 10 mM HEPES, 1 mM N-acetylcysteine, and 100 U/ml Penicillin/100 µg/ml

Streptomycin) and growth factors (50 ng/ml EGF, 50 ng/ml bFGF, and 50 ng/ml IGF-1) as well as the agent(s) being assessed are added into each well. Cells are cultured for 10 days in a standard cell culture incubator at 37 °C and 5% CO₂, with media change every 2 days. These cells are then cultured by removing the Stem Cell Proliferation Assay agents and replacing with Basal culture media and molecules to drive differentiation. An appropriate Basal culture media is Advanced DMEM/F12 supplemented with 1X N2, 1X B27, 2 mM Glutamax, 10 mM HEPES, 1 mM N-acetylcysteine, and 100 U/ml Penicillin/100 µg/ml Streptomycin and appropriate molecules to drive differentiation are 3 µM CHIR99021 and 5 µM DAPT for 10 days, with media change every 2 days. The number of hair cells in a population may be measured by using flow cytometry for GFP. Hair cell differentiation level can further be assessed using qPCR to measure hair cell marker (*e.g.*, *Myo7a*) expression level normalized using suitable and unregulated references or housekeeping genes (*e.g.*, *Hprt*). Hair cell differentiation level can also be assessed by immunostaining for hair cell markers (*eg.* *Myosin7a*, *vGlut3*, *Espin*, *PMCA*s, *Ribeye*, conjugated-phalloidin, *Atoh1*, *Pou4f3*, etc). Hair cell differentiation level can also be assessed by Western Blot for *Myosin7a*, *vGlut3*, *Espin*, *PMCA*s, *Prestin*, *Ribeye*, *Atoh1*, *Pou4f3*.

[0097] “Stem Cell Assay” as used herein is an assay in which a cell or a cell population are tested for a series of criteria to determine whether the cell or cell population are stem cells or enriched in stem cells or stem cell markers. In a stem cell assay, the cell/cell population are tested for stem cell characteristics such as expression of Stem Cell Markers, and *further optionally* are tested for stem cell function, including the capacity of self-renewal and differentiation.

[0098] “Stem Cell Proliferator” as used herein is a compound that induces an increase in a population of cells which have the capacity for self-renewal and differentiation.

[0099] “Stem Cell Proliferation Assay” as used herein is an assay to determine the capacity for agent(s) to induce the creation of stem cells from a starting cell population. In an exemplary Stem Cell Proliferation Assay, the number of cells for an initial cell population is harvested from a *Lgr5-GFP* mouse such as a B6.129P2-*Lgr5*^{tm1}(*cre/ERT2*)*Cle/J* mouse (also known as *Lgr5-EGFP-IRES-creERT2* or *Lgr5-GFP* mouse, Jackson Lab Stock No: 008875) between the age of 3 to 7 days, by isolating the Organ of Corti sensory epithelium and dissociating the epithelium into single cells. Approximately 5000 cells are entrapped in 40 µl of culture substrate (for

example: Matrigel (Corning, Growth Factor Reduced)) and placed at the center of wells in a 24-well plate with 500 μ l of an appropriate culture media, growth factors and agent being tested. Appropriate culture media and growth factors include Advanced DMEM/F12 with media Supplements (1X N2, 1X B27, 2 mM Glutamax, 10 mM HEPES, 1 mM N-acetylcysteine, and 100 U/ml Penicillin/100 μ g/ml Streptomycin) and growth factors (50 ng/ml EGF, 50 ng/ml bFGF, and 50 ng/ml IGF-1) as well as the agent(s) being assessed are added into each well. Cells are cultured for 10 days in a standard cell culture incubator at 37 °C and 5% CO₂, with media change every 2 days. The number of Lgr5⁺ cells is quantified by counting the number of cells identified as Lgr5⁺ in an *In Vitro* Lgr5 activity assay. The fraction of cells that are Lgr5⁺ is quantified by dividing the number of cells identified as Lgr5⁺ in a cell population by the total number of cells present in the cell population. The average Lgr5⁺ activity of a population is quantified by measuring the average mRNA expression level of Lgr5 of the population normalized using suitable and unregulated references or housekeeping genes (*e.g.*, Hprt). The number of hair cells in a population may be measured by staining with hair cell marker (*e.g.*, MyosinVIIa), or using an endogenous reporter of hair cell genes (*eg.* Pou4f3-GFP, Atoh1-nGFP) and analyzing using flow cytometry. The fraction of cells that are hair cells is quantified by dividing the number of cells identified as hair cells in a cell population by the total number of cells present in the cell population. Lgr5 activity can be measured by qPCR.

[00100] “Stem Cell Markers” as used herein can be defined as gene products (*e.g.* protein, RNA, etc) that specifically expressed in stem cells. One type of stem cell marker is gene products that are directly and specifically support the maintenance of stem cell identity. Examples include Lgr5 and Sox2. Additional stem cell markers can be identified using assays that were described in the literatures. To determine whether a gene is required for maintenance of stem cell identity, gain-of-function and loss-of-function studies can be used. In gain-of-function studies, over expression of specific gene product (the stem cell marker) would help maintain the stem cell identity. While in loss-of-function studies, removal of the stem cell marker would cause loss of the stem cell identity or induced the differentiation of stem cells. Another type of stem cell marker is gene that only expressed in stem cells but does not necessary to have specific function to maintain the identity of stem cells. This type of markers can be identified by comparing the gene expression signature of sorted stem cells and non-stem cells by assays such as micro-array and qPCR. This type of stem cell marker can be found in the

literature. (e.g. Liu Q. et al., *Int J Biochem Cell Biol.* 2015 Mar;60:99-111. <http://www.ncbi.nlm.nih.gov/pubmed/25582750>). Potential stem cell markers include Ccdc121, Gdf10, Opcm1, Phex, etc. The expression of stem cell markers such as Lgr5 or Sox2 in a given cell or cell population can be measured using assays such as qPCR, immunohistochemistry, western blot, and RNA hybridization. The expression of stem cell markers can also be measured using transgenic cells express reporters which can indicate the expression of the given stem cell markers, e.g. Lgr5-GFP or Sox2-GFP. Flow cytometry analysis can then be used to measure the activity of reporter expression. Fluorescence microscopy can also be used to directly visualize the expression of reporters. The expression of stem cell markers may further be determined using microarray analysis for global gene expression profile analysis. The gene expression profile of a given cell population or purified cell population can be compared with the gene expression profile of the stem cell to determine similarity between the 2 cell populations. Stem cell function can be measured by colony forming assay or sphere forming assay, self-renewal assay and differentiation assay. In colony (or sphere) forming assay, when cultured in appropriate culture media, the stem cell should be able to form colonies, on cell culture surface (e.g. cell culture dish) or embedded in cell culture substrate (e.g. Matrigel) or be able to form spheres when cultured in suspension. In colony/sphere forming assay, single stem cells are seeded at low cell density in appropriate culture media and allowed to proliferate for a given period of time (7-10 days). Colony formed are then counted and scored for stem cell marker expression as an indicator of stemness of the original cell. Optionally, the colonies that formed are then picked and passaged to test its self-renewal and differentiation potential. In self-renewal assay, when cultured in appropriate culture media, the cells should maintain stem cell marker (e.g. Lgr5) expression over at least one (e.g. 1, 2, 3, 4, 5, 10, 20, etc) cell divisions. In a Stem Cell Differentiation Assay, when cultured in appropriate differentiation media, the cells should be able to generate hair cell which can be identified by hair cell marker expression measured by qPCR, immunostaining, western blot, RNA hybridization or flow cytometry.

[0100] “Stemness Driver” as used herein is a composition that induces proliferation of LGR5⁺ cells, upregulates Lgr5 in cells, or maintains Lgr5 expression in cells, while maintaining the potential for self-renewal and the potential to differentiate into hair cells. Generally, stemness drivers upregulate at least one biomarker of post-natal stem cells. Stemness Drivers include but are not limited to Wnt agonists and GSK3 inhibitors.

[0101] “Subject” includes humans and mammals (*e.g.*, mice, rats, pigs, cats, dogs, and horses). In many embodiments, subjects are mammals, particularly primates, especially humans. In some embodiments, subjects are livestock such as cattle, sheep, goats, cows, swine, and the like; poultry such as chickens, ducks, geese, turkeys, and the like; and domesticated animals particularly pets such as dogs and cats. In some embodiments (*e.g.*, particularly in research contexts) subject mammals will be, for example, rodents (*e.g.*, mice, rats, hamsters), rabbits, primates, or swine such as inbred pigs and the like.

[0102] “Supporting Cell” as used herein in connection with a cochlear epithelium comprises epithelial cells within the organ of Corti that are not hair cells. This includes inner pillar cells, outer pillar cells, inner phalangeal cells, Deiter cells, Hensen cells, Boettcher cells, and/or Claudius cells.

[0103] “Synergy” or “synergistic effect” is an effect which is greater than the sum of each of the effects taken separately; a greater than additive effect.

[0104] “TGF Beta inhibitor” as used herein is a composition that reduces activity of TGF Beta.

[0105] “Tissue” is an ensemble of similar cells from the same origin that together carry out a specific function including, for example, tissue of cochlear, such as the Organ of Corti.

[0106] “Transtympanic” administration refers to direct injection of a composition across the tympanic membrane into the middle ear.

[0107] “Treating” as used herein in connection with a cell population means delivering a substance to the population to effect an outcome. In the case of *in vitro* populations, the substance may be directly (or even indirectly) delivered to the population. In the case of *in vivo* populations, the substance may be delivered by administration to the host subject.

[0108] “Wnt activation” as used herein is an activation of the Wnt signaling pathway.

[0109] The term “alkyl” as used herein refers to a straight or branched saturated hydrocarbon. For example, an alkyl group can have 1 to 8 carbon atoms (*i.e.*, (C₁-C₈)alkyl) or 1 to 6 carbon atoms (*i.e.*, (C₁-C₆ alkyl) or 1 to 4 carbon atoms.

[0110] The term “alkenyl” as used herein refers to a linear or branched hydrocarbon radical which includes one or more double bonds and can include divalent radicals, having from 2 to about 15 carbon atoms. Examples of alkenyl groups include but are not limited to, ethenyl, propenyl, butenyl, and higher homologs and isomers.

[0111] The term “alkynyl” as used herein refers to a linear or branched hydrocarbon radical which includes one or more triple bonds and can include divalent radicals, having from 2 to about 15 carbon atoms. Examples of alkynyl groups include but are not limited to, ethynyl, propynyl, butynyl, and higher homologs and isomers.

[0112] The term “halo” or “halogen” as used herein refers to fluoro, chloro, bromo and iodo.

[0113] The term “aryl” as used herein refers to a single all carbon aromatic ring or a multiple condensed all carbon ring system wherein at least one of the rings is aromatic. For example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 12 carbon atoms. Aryl includes a phenyl radical. Aryl also includes multiple condensed ring systems (e.g., ring systems comprising 2, 3 or 4 rings) having about 9 to 20 carbon atoms in which at least one ring is aromatic and wherein the other rings may be aromatic or not aromatic (i.e., carbocycle). Such multiple condensed ring systems may be optionally substituted with one or more (e.g., 1, 2 or 3) oxo groups on any carbocycle portion of the multiple condensed ring system. The rings of the multiple condensed ring system can be connected to each other via fused, spiro and bridged bonds when allowed by valency requirements. It is to be understood that the point of attachment of a multiple condensed ring system, as defined above, can be at any position of the ring system including an aromatic or a carbocycle portion of the ring.

[0114] The term “heteroaryl” as used herein refers to a single aromatic ring that has at least one atom other than carbon in the ring, wherein the atom is selected from the group consisting of oxygen, nitrogen and sulfur; the term also includes multiple condensed ring systems that have at least one such aromatic ring, which multiple condensed ring systems are further described below. Thus, the term includes single aromatic rings of from about 1 to 6 carbon atoms and about 1-4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur in the rings. The sulfur and nitrogen atoms may also be present in an oxidized form provided the ring is aromatic. The term also includes multiple condensed ring systems (e.g., ring systems comprising 2, 3 or 4 rings) wherein a heteroaryl group, as defined above, can be condensed with one or more rings selected from heteroaryls (to form for example a naphthyridinyl such as 1,8-naphthyridinyl), heterocycles, (to form for example a 1, 2, 3, 4-tetrahydronaphthyridinyl such as 1, 2, 3, 4-tetrahydro-1,8-naphthyridinyl), carbocycles (to form for example 5,6,7,8-tetrahydroquinolyl) and aryls (to form for example indazolyl) to form the multiple condensed ring system. Thus, a heteroaryl (a single aromatic ring or multiple condensed ring system) has about 1-20 carbon

atoms and about 1-6 heteroatoms within the heteroaryl ring. Such multiple condensed ring systems may be optionally substituted with one or more (e.g., 1, 2, 3 or 4) oxo groups on the carbocycle or heterocycle portions of the condensed ring. The rings of the multiple condensed ring system can be connected to each other via fused, spiro and bridged bonds when allowed by valency requirements. It is to be understood that the individual rings of the multiple condensed ring system may be connected in any order relative to one another. It is also to be understood that the point of attachment of a multiple condensed ring system (as defined above for a heteroaryl) can be at any position of the multiple condensed ring system including a heteroaryl, heterocycle, aryl or carbocycle portion of the multiple condensed ring system and at any suitable atom of the multiple condensed ring system including a carbon atom and heteroatom (e.g., a nitrogen).

[0115] The term “cycloalkyl” as used herein refers to a saturated or partially saturated ring structure having about 3 to about 8 ring members that has only carbon atoms as ring atoms and can include divalent radicals. Examples of cycloalkyl groups include but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexene, cyclopentenyl, cyclohexenyl.

[0116] The terms “heterocyclyl” or “heterocyclic” refer to monocyclic or polycyclic 3 to 24-membered rings containing carbon and heteroatoms selected from oxygen, phosphorous, nitrogen, or sulfur and wherein there are no delocalized π electrons (aromaticity) shared among the ring carbon or heteroatoms. Examples of heterocyclyl include, but are not limited to, oxetanyl, azetadiny, tetrahydrofuranyl, pyrrolidinyl, oxazoliny, oxazolidiny, thiazoliny, thiazolidiny, pyranyl, thiopyranyl, tetrahydropyranyl, dioxaliny, piperidiny, morpholiny, thiomorpholiny, thiomorpholiny S-oxide, thiomorpholiny S-dioxide, piperaziny, azepiny, oxepiny, diazepiny, tropany, and homotropany. Examples of heterocyclyl also include, but are not limited to, fused rings, bridged rings (e.g., 2,5-diazabicyclo[2,2,1]heptane), and spirocyclic rings, (e.g., 2,8-diazaspiro[4,5]decane).

[0117] The use of “or” means “and/or” unless stated otherwise. As used in this application, the term “comprise” and variations of the term, such as “comprising” and “comprises,” are not intended to exclude other additives, components, integers or steps. As used in this application, the terms “about” and “approximately” are used as equivalents. Any numerals used in this application with or without about/approximately are meant to cover any normal fluctuations appreciated by one of ordinary skill in the relevant art. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%,

17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

[0118] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0119] As used herein "pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, surfactant, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals. Exemplary pharmaceutically acceptable carriers include, but are not limited to, to sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; tragacanth; malt; gelatin; talc; cocoa butter, waxes, animal and vegetable fats, paraffins, silicones, bentonites, silicic acid, zinc oxide; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; and any other compatible substances employed in pharmaceutical formulations.

[0120] "Pharmaceutically acceptable salt" includes both acid and base addition salts.

[0121] "Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic

acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pantoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, / toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

[0122] "Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. For example, inorganic salts include, but are not limited to, ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Example organic bases used in certain embodiments include isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

[0123] Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening,

flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

[0124] Examples of pharmaceutically-acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.


[0125] Compounds or compositions described herein can be formulated in any manner suitable for a desired delivery route, e.g., transtympanic injection, transtympanic wicks and catheters, and injectable depots. Typically, formulations include all physiologically acceptable compositions including derivatives or prodrugs, solvates, stereoisomers, racemates, or tautomers thereof with any physiologically acceptable carriers, diluents, and/or excipients.

DETAILED DESCRIPTION OF THE INVENTION

[0126] A description of example embodiments of the invention follows.

Compounds



[0127] The present disclosure provides compounds of Formula (I),






and pharmaceutically acceptable salts and tautomers thereof, wherein Q¹, Q², Q³, R¹, R², R³, Ar, -Z-W-X-Y- and m are as defined above for Formula (I).


[0128] In some embodiments, the compounds of Formula (I) have one or more of the following features:

- provided that the compound is not




or

b) provided that when Ar is , and  is  or



[0129] In certain embodiments, the present disclosure provides a compound of Formula (I) that is not disclosed in WO 2003/076442 (PCT/US03/05050), which is incorporated herein by reference.

[0130] In certain embodiments of Formula (I), R^X is $-COR^{X1}$ or $-SO_2R^{X1}$.



[0132] In certain embodiments, R^{X1} is heterocyclyl optionally substituted with one to twelve substituents that is halo. In certain embodiments, R^{X1} is heterocyclyl which is deuterated. In certain embodiments, the heterocyclyl is a monocyclic or bicyclic ring. In some embodiments, the heterocyclyl is a fused, bridged, or spirocyclic ring. In certain embodiments, the heterocyclyl contains one to three nitrogens (i.e., 1, 2, or 3 nitrogens) and/or one to three oxygens (i.e., 1, 2, or 3 oxygens). In certain embodiments, the heterocyclyl contains one nitrogen and/or one oxygen. In certain embodiments, the heterocyclyl contains one nitrogen. In certain embodiments, the heterocyclyl contains two nitrogens. In certain embodiments, the heterocyclyl contains one nitrogen and one oxygen.

[0133] In some embodiments, R^{X1} is piperidine, 2,8-diazaspiro[4,5]decane, 2,5-diazabicyclo[2,2,1]heptane, or 8-oxa-3-azabicyclo[3.2.1]octane, each of which is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, C_1 - C_4 alkyl, $-[C(R^{X1a})_2]_p-OH$, $-(CH_2)_p-NMe_2$, $-(CH_2)_p-NHMe$, $-(CH_2)_p-NH_2$; wherein p is 0, 1, 2, or 3. In some embodiments, R^{X1} is piperidine, optionally substituted with one to six halo substituents. In some embodiments, R^{X1} is piperidine optionally substituted with $-[C(R^{X1a})_2]_p-OH$, $-(CH_2)_p-NMe_2$.


[0134] In certain embodiments of Formula (I), the heterocyclyl is optionally substituted with C_1 - C_4 alkyl, $-(CH_2)_p-OH$, or $-(CH_2)_p-NH_2$; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is

heterocyclyl substituted with C₁-C₄alkyl. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)_p-OH; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is heterocyclyl substituted with -CH₂-OH. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)_p-NH₂; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is heterocyclyl substituted with -CH₂-NH₂.

[0135] In certain embodiments of Formula (I), R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with -[C(R^{X1a})₂]_p-CN. In certain embodiments, R^{X1} is heterocyclyl substituted with -[C(R^{X1a})₂]_p-OH, -[C(R^{X1a})₂]_p-O-C₁-C₄alkyl, -NHCOC₁-C₄alkyl, -CONHC₁-C₄alkyl, COH, -CO₂H, -[C(R^{X1a})₂]_p-COO-C₁-C₄alkyl, -[C(R^{X1a})₂]_p-NH₂, -[C(R^{X1a})₂]_p-NH-C₁-C₄alkyl, or -[C(R^{X1a})₂]_p-N-(C₁-C₄alkyl)₂. In certain embodiments, R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with -CONHC₁-C₄alkyl, -COH, -CO₂H, or -[C(R^{X1a})₂]_p-COO-C₁-C₄alkyl.


[0136] In certain embodiments of Formula (I), each R^{X1a} is independently selected from the group consisting of hydrogen and halo. In certain embodiments, both R^{X1a} groups together form C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

[0137] In certain embodiments of Formula (I), R^X is heteroaryl. In certain embodiments, the heteroaryl is monocyclic or bicyclic. In certain embodiments, the heteroaryl contains one to three nitrogens (i.e., 1, 2, or 3 nitrogens) and/ or one to three oxygens (i.e., 1, 2, or 3 oxygens). In certain embodiments, the heteroaryl contains one nitrogen and/ or one oxygen. In certain embodiments, the heteroaryl contains one nitrogen. In certain embodiments, the heteroaryl contains two nitrogens. In certain embodiments, the heteroaryl contains one nitrogen and one




oxygen. In certain embodiments, R^X is

[0138] In certain embodiments of Formula (I), R^X is -(C₁-C₄alkylene)-(C₃-C₈cycloalkyl). In certain embodiments, the-(C₁-C₄alkylene)-(C₃-C₈cycloalkyl) is substituted with one to two halo on the C₁-C₄alkylene. In certain embodiments, the C₃-C₈cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, R^X is -(C₁-C₄alkylene)-(C₃-C₈cycloalkyl), wherein the -(C₁-C₄alkylene)-(C₃-C₈cycloalkyl) is optionally substituted with one or two halo on the C₁-C₄alkylene and wherein C₃-C₈cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl, or




cyclohexyl. In certain embodiments, R^X is

[0139] The present disclosure provides a compound having the Formula (Ia),




and pharmaceutically acceptable salts and tautomers thereof, wherein Q^1 , Q^2 , Q^3 , R^1 , R^2 , R^3 , Ar, -Z-W-X-Y- and m are as defined above for Formula (Ia).

[0140] The present disclosure provides a compound having the Formula (Ib),




and pharmaceutically acceptable salts and tautomers thereof, wherein Q^1 , Q^2 , Q^3 , R^1 , R^2 , R^3 , Ar, -Z-W-X-Y- and m are as defined above for Formula (Ib).

[0141] In certain embodiments, Q^1 is CH; Q^2 is N; and Q^3 is C. In certain embodiments, Q^1 is N; Q^2 is C; and Q^3 is N. In certain embodiments, Q^1 is CH; Q^2 is C; and Q^3 is N. In certain embodiments, Q^1 is N; Q^2 is N; and Q^3 is C.



[0142] In certain embodiments, the [structure] is selected from the group consisting of





[0143] In certain embodiments, R¹ is hydrogen or halo. In certain embodiments, R¹ is C₁-C₄alkyl, wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH. In certain embodiments, R¹ is C₁-C₄alkynyl, -CN, -OH, or -S(O)₂NH₂. In certain embodiments, R¹ is -NH₂ or -NHC(O)R^{1a}, wherein R^{1a} is C₁-C₄alkyl. In certain embodiments, R¹ is C₁-C₄alkenyl. In certain embodiments, R¹ is -O-C₁-C₄alkyl.








[0144] In certain embodiments, R² is hydrogen or halo. In certain embodiments, R² is C₁-C₄alkyl, wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH. In certain embodiments, R² is C₁-C₄alkynyl, -CN, -OH, or -S(O)₂NH₂. In certain embodiments, R² is -NH₂ or -NHC(O)R^{2a}, wherein R^{2a} is C₁-C₄alkyl. In certain embodiments, R² is -S(O)₂NH₂.

[0145] In certain embodiments, R² is C₁-C₄alkenyl. In certain embodiments, R² is -O-C₁-C₄alkyl. In certain embodiments, R² is -NH₂, -NH(C₁-C₄alkyl), or -N(C₁-C₄alkyl)₂.

[0146] In certain embodiments, R² is selected from the group consisting of halo, C₁-C₄alkyl, C₁-C₄alkenyl, C₁-C₄alkynyl, -CN, -OH, -O-C₁-C₄alkyl, -NH₂, -NH(C₁-C₄alkyl), -N(C₁-C₄alkyl)₂, -NHC(O)R^{2a}, and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{2a} is C₁-C₄alkyl. In certain embodiments, R² is selected from the group consisting of halo, C₁-C₄alkyl, C₁-C₄alkynyl, -CN, -OH, -NH₂, -NHC(O)R^{2a}, and -S(O)₂NH₂; wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH; and wherein R^{2a} is C₁-C₄alkyl. In certain embodiments, R² is not hydrogen.

[0147] In certain embodiments, R³ is hydrogen or halo. In certain embodiments, R³ is C₁-C₄alkyl, wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and -OH. In certain embodiments, R³ is C₁-C₄alkynyl, -CN, -OH, or -S(O)₂NH₂. In certain embodiments, R³ is -NH₂ or -NHC(O)R^{3a}, wherein R^{3a} is C₁-C₄alkyl. In certain embodiments, R³ is C₁-C₄alkenyl. In certain embodiments, R³ is -O-C₁-C₄alkyl.



[0150] In certain embodiments of Formula (Ib), Ar is . In certain embodiments of Formula (Ib), Ar is . In certain embodiments of Formula (Ib), Ar is  or , wherein Q⁷ is selected from S, O, CH₂, and NR^{Q7}; wherein R^{Q7} is hydrogen or optionally substituted C₁-C₄alkyl. In certain embodiments of Formula (Ib), Ar is , , wherein Q⁷ is selected from S, O, CH₂, and NR^{Q7}; wherein R^{Q7} is hydrogen or optionally substituted C₁-C₄alkyl. In certain embodiments of Formula (Ib), Ar is , wherein each Q⁶ is independently selected from CR^{Q6} and N; wherein R^{Q6} is hydrogen, halo, -CN, lower alkyl, or substituted alkyl.

[0151] In certain embodiments, -Z-W-X-Y- is -C(R^Z)₂-C(R^W)₂-N(R^X)-C(R^Y)₂-. In certain embodiments, -Z-W-X-Y- is -C(R^Z)₂-C(R^W)₂-CH(R^X)-C(R^Y)₂-. In certain embodiments, -Z-W-X-Y- is -C(R^W)₂-CH(R^X)-C(R^Y)₂-.

[0152] In certain embodiments, each R^Z is independently selected from the group consisting of hydrogen and halo. In certain embodiments, both R^Z groups together form C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, both R^Z groups together form oxo. In certain embodiments, R^Z and R^W together with the carbons to which they are attached form a C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

[0153] In certain embodiments, each R^W is independently selected from the group consisting of hydrogen and halo. In certain embodiments, both R^W groups together form C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, both R^W groups together form oxo. In certain embodiments, R^Z and R^W together with the carbons to


which they are attached form a C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

[0154] In certain embodiments, each R^Y is independently selected from the group consisting of hydrogen and halo. In certain embodiments, both R^Y groups together form C₃-C₆cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, both R^Y groups together form oxo.

[0155] In certain embodiments of Formula (Ia) and (Ib), R^X is H. In certain embodiments of Formula (Ia) and (Ib), R^X is R^{X1}, which is C₃-C₈cycloalkyl, heteroaryl, or heterocyclyl, wherein the heterocyclyl is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, C₁-C₄alkyl, -(CH₂)_p-OH, -[C(R^{X1a})₂]_p-OH, -[C(R^{X1a})₂]_p-O-C₁-C₄alkyl, -NHCOC₁-C₄alkyl, -CONHC₁-C₄alkyl, -(CH₂)_p-NH₂, -[C(R^{X1a})₂]_p-NH₂, -[C(R^{X1a})₂]_p-NH-C₁-C₄alkyl, -[C(R^{X1a})₂]_p-N-(C₁-C₄alkyl)₂; wherein p is 0, 1, 2, or 3; wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^{X1a} groups together form C₃-C₆cycloalkyl.


[0156] In certain embodiments of Formula (Ia) and (Ib), R^X is -COR^{X1} or -SO₂R^{X1}.

[0157] In certain embodiments of Formula (Ia) and (Ib), R^X is selected from



[0158] In certain embodiments of Formula (Ia) and (Ib), R^X is -(C₁-C₄alkylene)-R^{X1}, wherein the -(C₁-C₄alkylene)-R^{X1} is optionally substituted with one to four halo on the C₁-C₄alkylene. In certain embodiments, the -(C₁-C₄alkylene)-R^{X1} is substituted with one to four halo on the C₁-

C₄alkylene. In certain embodiments, the -(C₁-C₄alkylene)-R^{X1} is substituted with one or two halo on the C₁-C₄alkylene. In certain embodiments, R^X is -(C₁-C₄alkylene)-R^{X1}, wherein the -(C₁-C₄alkylene)-R^{X1} is optionally substituted with one or two halo on the C₁-C₄alkylene and wherein R^{X1} is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. In certain embodiments, R^X



[0159] In certain embodiments of Formula (Ia) and (Ib), R^{X1} is C₃-C₈cycloalkyl. In certain embodiments, R^{X1} is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.


[0160] In certain embodiments of Formula (Ia) and (Ib), R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with one to twelve substituents that is halo. In certain embodiments, R^{X1} is heterocyclyl which is deuterated. In certain embodiments, the heterocyclyl is monocyclic or bicyclic. In certain embodiments, the heterocyclyl contains one to three nitrogens (i.e., 1, 2, or 3 nitrogens) and/ or one to three oxygens (i.e., 1, 2, or 3 oxygens). In certain embodiments, the heterocyclyl contains one nitrogen and/ or one oxygen. In certain embodiments, the heterocyclyl contains one nitrogen. In certain embodiments, the heterocyclyl contains two nitrogens. In certain embodiments, the heterocyclyl contains one nitrogen and one oxygen.

[0161] In certain embodiments of Formula (Ia) and (Ib), R^{X1} is heterocyclyl optionally substituted with C₁-C₄alkyl, -(CH₂)_p-OH, or -(CH₂)_p-NH₂; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is heterocyclyl substituted with C₁-C₄alkyl. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)_p-OH; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)-OH. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)_p-NH₂; wherein p is 1, 2, or 3. In certain embodiments, R^{X1} is heterocyclyl substituted with -(CH₂)-NH₂.

[0162] In certain embodiments of Formula (Ia) and (Ib), R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with -[C(R^{X1a})₂]_p-CN. In certain embodiments, R^{X1} is heterocyclyl substituted with -[C(R^{X1a})₂]_p-OH, -[C(R^{X1a})₂]_p-O-C₁-C₄alkyl, -NHCOC₁-C₄alkyl, -[C(R^{X1a})₂]_p-NH₂, -[C(R^{X1a})₂]_p-NH-C₁-C₄alkyl, or -[C(R^{X1a})₂]_p-N-(C₁-C₄alkyl)₂. In certain embodiments, R^{X1} is heterocyclyl, wherein the heterocyclyl is optionally substituted with -CONHC₁-C₄alkyl, -COH, -CO₂H, or -[C(R^{X1a})₂]_p-COO-C₁-C₄alkyl.

[0163] In certain embodiments of Formula (Ia) and (Ib), each R^{X1a} is independently selected from the group consisting of hydrogen and halo. In certain embodiments, both R^{X1a} groups together form C_3 - C_6 cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.


[0164] In certain embodiments of Formula (Ia) and (Ib), R^{X1} is heteroaryl. In certain embodiments, the heteroaryl is monocyclic or bicyclic. In certain embodiments, the heteroaryl contains one to three nitrogens (i.e., 1, 2, or 3 nitrogens) and/ or one to three oxygens (i.e., 1, 2, or 3 oxygens). In certain embodiments, the heteroaryl contains one nitrogen and/ or one oxygen. In certain embodiments, the heteroaryl contains one nitrogen. In certain embodiments, the heteroaryl contains two nitrogens. In certain embodiments, the heteroaryl contains one nitrogen





and one oxygen. In certain embodiments, R^{X1} is


[0165] In certain embodiments of Formula (Ia) and (Ib), R^X is $-\text{CON}(\text{R}^{X2})_2$. In certain embodiments, R^X is $-\text{CON}(\text{R}^{X2})_2$, wherein R^{X2} is hydrogen or methyl. In certain embodiments, R^X is $-\text{CONH}_2$. In certain embodiments, R^X is $-\text{CON}(\text{R}^{X2})_2$, wherein R^{X2} is C_1 - C_4 alkyl. In certain embodiments, R^X is $-\text{CON}(\text{R}^{X2})_2$, wherein R^{X2} is methyl.

[0166] In certain embodiments, m is 0. In certain embodiments, m is 1. In certain embodiments, m is 2.




[0167] In one variation of the compounds herein, Ar is  and Q^1 is CH; Q^2 is N; Q^3 is C; Q^4 is C; and Q^5 is C.





[0168] In one variation of the compounds herein, Ar is  and Q^1 is CH; Q^2 is N; Q^3 is C; Q^4 is C; and Q^5 is C.

[0169] The present disclosure provides compounds herein of Formula (I):




having one, two, three, or more of the following features:





- Ar is  ;
- Q^1 is CH; Q^2 is N; and Q^3 is C;
- R^2 is hydrogen or halo;
- $-Z-W-X-Y-$ is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$;
- R^X is $-COR^{X1}$.

[0170] The present disclosure provides compounds herein of Formula (I):




having one, two, three, or more of the following features:





- Ar is  ;
- Q^1 is CH; Q^2 is N; and Q^3 is C;
- R^2 is C_1-C_4 alkyl, wherein the alkyl is optionally substituted with one to 3 substituents independently selected from the group consisting of halo and $-OH$;
- $-Z-W-X-Y-$ is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$;
- R^X is $-COR^{X1}$.

[0171] The present disclosure provides compounds herein of Formula (I):












having one, two, three, or more of the following features:















- Ar is  ;
- Q^1 is CH; Q^2 is N; and Q^3 is C;
- R^2 is C_1 - C_4 alkynyl, -CN, -OH, -S(O)₂NH₂, -NH₂ or -NHC(O)R^{2a};
- Z-W-X-Y- is -C(R^Z)₂-C(R^W)₂-N(R^X)-C(R^Y)₂-;
- R^X is -COR^{X1}.






[0172] Nonlimiting examples of compounds of the invention are presented below.







Compound I-1	
Compound I-2	
Compound I-3	







<p>Compound I-4</p>	
<p>Compound I-5</p>	
<p>Compound I-6</p>	
<p>Compound I-7</p>	
<p>Compound I-8</p>	
<p>Compound I-9</p>	







Compound I-10	
Compound I-11	
Compound I-12	
Compound I-13	
Compound I-14	
Compound I-15	





Compound I-16	 <p>Chemical structure of Compound I-16: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a cyano group (-C≡N) at the 2-position and a carbonyl group (-C(=O)NH-) at the 4-position. The benzimidazole ring is further substituted with a piperidine ring at the 5-position, which is in turn substituted with a carbonyl group (-C(=O)-) and a difluoromethyl group (-CF₂-).</p>
Compound I-17	 <p>Chemical structure of Compound I-17: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a carbonyl group (-C(=O)NH-) at the 4-position and a trifluoromethyl group (-CF₃) at the 5-position. The benzimidazole ring is further substituted with a piperidine ring at the 2-position, which is in turn substituted with a carbonyl group (-C(=O)-) and a difluoromethyl group (-CF₂-).</p>
Compound I-18	 <p>Chemical structure of Compound I-18: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a carbonyl group (-C(=O)NH-) at the 4-position and a trifluoromethyl group (-CF₃) at the 5-position. The benzimidazole ring is further substituted with a bicyclic piperidine ring at the 2-position, which is in turn substituted with a carbonyl group (-C(=O)-).</p>
Compound I-19	 <p>Chemical structure of Compound I-19: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a carbonyl group (-C(=O)NH-) at the 4-position and a fluorine atom (-F) at the 5-position. The benzimidazole ring is further substituted with a piperidine ring at the 2-position, which is in turn substituted with a carbonyl group (-C(=O)-) and a primary amine group (-CH₂NH₂).</p>
Compound I-20	 <p>Chemical structure of Compound I-20: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a carbonyl group (-C(=O)NH-) at the 4-position and a trifluoromethyl group (-CF₃) at the 5-position. The benzimidazole ring is further substituted with a piperidine ring at the 2-position, which is in turn substituted with a carbonyl group (-C(=O)-) and a hydroxymethyl group (-CH₂OH).</p>
Compound I-21	 <p>Chemical structure of Compound I-21: A complex molecule featuring a central benzimidazole core. The benzimidazole ring is substituted with a carbonyl group (-C(=O)NH-) at the 4-position and a cyano group (-C≡N) at the 5-position. The benzimidazole ring is further substituted with a piperidine ring at the 2-position, which is in turn substituted with a carbonyl group (-C(=O)-) and a hydroxymethyl group (-CH₂OH).</p>

Compound I-22	 <p>Chemical structure of Compound I-22: A complex molecule featuring a central benzimidazole core. One nitrogen of the benzimidazole is substituted with a 2,2,3,3,4,4-hexafluoropiperidine ring. The other nitrogen is substituted with a 2,6-difluorophenyl ring. The 2-position of the benzimidazole is substituted with a 2,4-dioxo-1,2,3,4-tetrahydro-1H-quinazolin-5-yl group.</p>
Compound I-23	 <p>Chemical structure of Compound I-23: A complex molecule featuring a central benzimidazole core. One nitrogen of the benzimidazole is substituted with a 2,2,3,3,4,4-hexafluoropiperidine ring. The other nitrogen is substituted with a 2,6-difluorophenyl ring. The 2-position of the benzimidazole is substituted with a 2,4-dioxo-1,2,3,4-tetrahydro-1H-quinazolin-5-yl group.</p>
Compound I-24	 <p>Chemical structure of Compound I-24: A complex molecule featuring a central benzimidazole core. One nitrogen of the benzimidazole is substituted with a 2,2,3,3,4,4-hexafluoro-1,3-dioxepane ring. The other nitrogen is substituted with a 2,6-difluorophenyl ring. The 2-position of the benzimidazole is substituted with a 2,4-dioxo-1,2,3,4-tetrahydro-1H-quinazolin-5-yl group.</p>
Compound I-25	 <p>Chemical structure of Compound I-25: A complex molecule featuring a central benzimidazole core. One nitrogen of the benzimidazole is substituted with a 2,2,3,3,4,4-hexafluoro-1,3-dioxepane ring. The other nitrogen is substituted with a 2,6-difluorophenyl ring. The 2-position of the benzimidazole is substituted with a 2,4-dioxo-1,2,3,4-tetrahydro-1H-quinazolin-5-yl group.</p>
Compound I-26	 <p>Chemical structure of Compound I-26: A complex molecule featuring a central benzimidazole core. One nitrogen of the benzimidazole is substituted with a 2,2,3,3,4,4-hexafluoro-1,3-dioxepane ring. The other nitrogen is substituted with a 2,6-difluorophenyl ring. The 2-position of the benzimidazole is substituted with a 2,4-dioxo-1,2,3,4-tetrahydro-1H-quinazolin-5-yl group.</p>


<p>Compound I-27</p>	
<p>Compound I-28</p>	
<p>Compound I-29</p>	
<p>Compound I-30</p>	
<p>Compound I-31</p>	
<p>Compound I-32</p>	

Compound I-33	 <p>Chemical structure of Compound I-33: A central benzimidazole ring is substituted at the 2-position with a 1H-imidazole-4,5-dione ring. The 5-position of the benzimidazole is substituted with a 7-fluoro-1,2,3,4-tetrahydroquinoline ring. The 4-position of the tetrahydroquinoline is substituted with a piperidine ring via a carbonyl group. The piperidine ring is further substituted with a dimethylamino group (-N(CH₃)₂).</p>
Compound I-34	 <p>Chemical structure of Compound I-34: Similar to Compound I-33, but the piperidine ring is substituted with an amino group (-NH₂) instead of a dimethylamino group.</p>
Compound I-35	 <p>Chemical structure of Compound I-35: Similar to Compound I-33, but the piperidine ring is substituted with a methylamino group (-NHCH₃) instead of a dimethylamino group.</p>
Compound I-36	 <p>Chemical structure of Compound I-36: Similar to Compound I-33, but the piperidine ring is substituted with a dimethylamino group (-N(CH₃)₂) at the 4-position instead of the 1-position.</p>
Compound I-37	 <p>Chemical structure of Compound I-37: Similar to Compound I-33, but the piperidine ring is substituted with a piperidine ring via a carbonyl group, forming a piperidine-piperidine linkage.</p>
Compound I-38	 <p>Chemical structure of Compound I-38: Similar to Compound I-33, but the piperidine ring is substituted with a piperidine ring via a carbonyl group, forming a piperidine-piperidine linkage, and the piperidine ring is substituted with a methylamino group (-NHCH₃).</p>

<p>Compound I-39</p>	
<p>Compound I-40</p>	
<p>Compound I-41</p>	
<p>Compound I-42</p>	
<p>Compound I-43</p>	
<p>Compound I-44</p>	


Compound I-45	
Compound I-46	
Compound I-47	
Compound I-48	

[0100] The present disclosure provides a compound having Formula (IIa):




and pharmaceutically acceptable salts and tautomers thereof.

[0101] In some embodiments, R^{X1} is selected from the group consisting of:




[0106] The present disclosure provides a compound having Formula (IIc):





and pharmaceutically acceptable salts and tautomers thereof.


[0107] In some embodiments, R^{X1} is selected from the group consisting of:




[0108] In some embodiments, R^{X1} is selected from the group consisting of:



[0109] In some embodiments, R^{X1} is selected from the group consisting of:




[0110] The present disclosure provides a compound having Formula (IIe):




and pharmaceutically acceptable salts and tautomers thereof.


[0111] In some embodiments, R^{X1} is selected from the group consisting of:




[0112] In some embodiments, R^{X1} is selected from the group consisting of:



[0113] In some embodiments, R^{X1} is selected from the group consisting of:





[0114] Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structure except for the replacement of a hydrogen atom by deuterium or tritium, or the replacement of a carbon atom by ¹³C or ¹⁴C, or the replacement of a nitrogen atom by ¹⁵N, or the replacement of an oxygen atom with ¹⁷O or ¹⁸O are within the scope of the present disclosure. Such isotopically labeled compounds are useful as research or diagnostic tools. In certain embodiments, deuteration can be used to slow metabolism and thus potentially improve the compound half-life. Any or all hydrogens in the compound can be replaced with deuterium.

Methods of Synthesizing the Disclosed Compounds

[0115] The compounds of the present invention may be made by a variety of methods, including standard chemistry. Suitable synthetic routes are depicted in the schemes given below.

[0116] The compounds of any of the formulae described herein may be prepared by methods known in the art of organic synthesis as set forth in part by the following synthetic schemes and examples. In the schemes described below, it is well understood that protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles or chemistry. Protecting groups are manipulated according to standard methods of organic synthesis (T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", Third edition, Wiley, New York 1999). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection processes, as well as the reaction conditions and order of their execution, shall be consistent with the preparation of compounds of the present disclosure.

[0117] Those skilled in the art will recognize if a stereocenter exists in any of the compounds of the present disclosure. Accordingly, the present invention includes both possible stereoisomers (unless specified in the synthesis) and includes not only racemic compounds but


the individual enantiomers and/or diastereomers as well. When a compound is desired as a single enantiomer or diastereomer, it may be obtained by stereospecific synthesis or by resolution of the final product or any convenient intermediate. Resolution of the final product, an intermediate, or a starting material may be affected by any suitable method known in the art. See, for example, "Stereochemistry of Organic Compounds" by E. L. Eliel, S. H. Wilen, and L. N. Mander (Wiley-Interscience, 1994).

Methods of Making the Compounds


[0118] The compounds described herein may be made from commercially available starting materials or synthesized using known organic, inorganic, and/or enzymatic processes.

[0119] The compounds of the present invention can be prepared in a number of ways well known to those skilled in the art of organic synthesis. By way of example, compounds of the disclosure can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. These methods include but are not limited to those methods described below.

[0120] A representative synthesis for subject compounds is shown in Scheme 1.

Scheme 1. General synthesis of an illustrative compound of Formula (I)

[0121] In Scheme 1, Core 2 is an embodiment wherein Q^1 is CH, Q^2 is N, and Q^3 is C; R^2 is



[0122] Compound 1 and the allylic aldehyde are commercially available starting materials. Alternatively, compound 1 and the allylic aldehyde can be synthesized via a variety of different synthetic routes using commercially available starting materials and/or starting materials prepared by conventional synthetic methods.

[0123] With continued reference to Scheme 1, compound 1 and an allylic aldehyde are reacted to form compound 2 in a condensation reaction in a suitable solvent such as acetonitrile at a temperature, for example, from about 40⁰C to 100⁰C. Compound 2 is reacted with ammonia to form compound 3 in a suitable solvent such as methanol at a temperature, for example, in the range from 0⁰C to room temperature. Compound 3 can be used in a coupling reaction to be discussed below.

[0124] With continued reference to Scheme 1, compound 5 may be prepared from an alkylation of compound 4 with an alkyl halide in the presence of a base, such as an alkali metal hydride, such as sodium hydride. The reaction can be run in a suitable solvent, such as dimethylformamide (DMF) at a temperature, for example, in the range from 0⁰C to room temperature.

[0125] Compound 6 may be prepared from the reduction of compound 5. Suitable reduction reagents include borane pyridine complex. The reaction can be run in a suitable solvent, such as acetic acid at a temperature, for example, in the range from 0⁰C to room temperature.

[0126] Compound 7 may be prepared from reaction of compound 6. The reaction is carried with formaldehyde in the presence of acid such as sulfuric acid and acetic acid.

[0127] Compound 8 may be prepared from protection of the amino group of compound 7. Suitable reagents include BOC anhydride. The reaction can be run in a suitable solvent, such as tetrahydrofuran (THF) at a temperature, for example, in the range from 0⁰C to room temperature.

[0128] Core 1 may be prepared from the dehydrogenation of compound 8. Suitable reagents include DDQ (2,3-dichloro-5,6-dicyano-1,4-benzoquinone). The reaction can be run in a suitable solvent, such as tetrahydrofuran (THF) at a temperature, for example, in the range from 0⁰C to room temperature.

[0129] Compound 10 may be prepared from core 1 by deprotection of the amino group and then the subsequent reaction with acyl halide. The deprotection of the amino group can be performed under acidic conditions, if the protecting group is BOC. Then reaction with an acyl halide can result in compound 10. The reaction can be run in a suitable solvent, such as dimethylformamide (DMF) at a temperature, for example, in the range from 0⁰C to room temperature.

[0130] Compound 11 may be prepared from compound 10 by an acylation reaction, such as a Friedel Crafts acylation reaction. In this reaction, an acyl halide is reacted with compound 10 in

a suitable solvent, such as methylene chloride at a temperature, for example, in the range from 30⁰C to 100⁰C. The product is then reacted an alcohol and base to form an ester, as in compound 11.

[0131] Compound 11 and Compound 3 are reacted to form the 1H-pyrrole-2,5-dione compound. The reaction is carried out in an inert organic solvent such as dimethylformamide, tetrahydrofuran, and the like and in the presence of a base such as potassium tert-butoxide.

Methods of Using the Compounds

[0132] The present disclosure relates to methods to activate the Wnt pathway or inhibiting GSK3-beta activity. Although there are hundreds of purported GSK3 inhibitors in the patent and non patent literature, not all GSK3 inhibitors when administered in the absence of other therapeutic agents would be sufficient nor potent enough to promote activation of stem cell proliferation.

[0133] In another aspect the present disclosure relates to methods to prevent, reduce or treat the incidence and/or severity of disorders or diseases associated with absence or lack of certain tissue cells. In one aspect the present disclosure relates to methods to prevent, reduce or treat the incidence and/or severity of inner ear disorders and hearing impairments involving inner ear tissue, particularly inner ear hair cells, their progenitors, and optionally, the stria vascularis, and associated auditory nerves. Of particular interest are those conditions that lead to permanent hearing loss where reduced number of hair cells may be responsible and/or decreased hair cell function. Also of interest are those arising as an unwanted side-effect of ototoxic therapeutic drugs including cisplatin and its analogs, aminoglycoside antibiotics, salicylate and its analogs, or loop diuretics. In certain embodiments, the present disclosure relates to inducing, promoting, or enhancing the growth, proliferation or regeneration of inner ear tissue, particularly inner ear supporting cells and hair cells.

[0134] Among other things, the methods presented here are useful for the preparation of pharmaceutical formulations for the prophylaxis and/or treatment of acute and chronic ear disease and hearing loss, dizziness and balance problems especially of sudden hearing loss, acoustic trauma, hearing loss due to chronic noise exposure, presbycusis, trauma during implantation of the inner ear prosthesis (insertion trauma), dizziness due to diseases of the inner ear area, dizziness related and/or as a symptom of Meniere's disease, vertigo related and/or as a

symptom of Meniere's disease, tinnitus, and hearing loss due to antibiotics and cytostatics and other drugs.

[0135] When cochlea supporting cell populations are treated with the compound, whether the population is *in vivo* or *in vitro*, the treated supporting cells exhibit stem-like behavior in that the treated supporting cells have the capacity to proliferate and differentiate and, more specifically, differentiate into cochlear hair cells. Preferably, the compound induces and maintains the supporting cells to produce daughter stem cells that can divide for many generations and maintain the ability to have a high proportion of the resulting cells differentiate into hair cells. In certain embodiments, the proliferating stem cells express stem cell markers which may include *Lgr5*, *Sox2*, *Opem1*, *Phex*, *lin28*, *Lgr6*, *cyclin D1*, *Msx1*, *Myb*, *Kit*, *Gdnf3*, *Zic3*, *Dppa3*, *Dppa4*, *Dppa5*, *Nanog*, *Esrrb*, *Rex1*, *Dnmt3a*, *Dnmt3b*, *Dnmt3l*, *Utf1*, *Tcl1*, *Oct4*, *Klf4*, *Pax6*, *Six2*, *Zic1*, *Zic2*, *Otx2*, *Bmi1*, *CDX2*, *STAT3*, *Smad1*, *Smad2*, *smad2/3*, *smad4*, *smad5*, and/or *smad7*.

[0136] In some embodiments, the method of the present disclosure may be used to maintain, or even transiently increase stemness (*i.e.*, self-renewal) of a pre-existing supporting cell population prior to significant hair cell formation. In some embodiments, the pre-existing supporting cell population comprises inner pillar cells, outer pillar cells, inner phalangeal cells, Deiter cells, Hensen cells, Boettcher cells, and/or Claudius cells. Morphological analyses with immunostaining (including cell counts) and lineage tracing across a Representative Microscopy Samples may be used to confirm expansion of one or more of these cell-types. In some embodiments, the pre-existing supporting cells comprise *Lgr5*⁺ cells. Morphological analyses with immunostaining (including cell counts) and qPCR and RNA hybridization may be used to confirm *Lgr5* upregulation amongst the cell population.

[0137] Advantageously, the methods of the present disclosure achieve these goals without the use of genetic manipulation. Germ-line manipulation used in many academic studies is not a therapeutically desirable approach to treating hearing loss. In general, the therapy preferably involves the administration of a small molecule, peptide, antibody, or other non-nucleic acid molecule or nucleic acid delivery vector unaccompanied by gene therapy. In certain embodiments, the therapy involves the administration of a small organic molecule. Preferably, hearing protection or restoration is achieved through the use of a (non-genetic) therapeutic that is injected in the middle ear and diffuses into the cochlea.

[0138] The cochlea relies heavily on all present cell types, and the organization of these cells is important to their function. As supporting cells play an important role in neurotransmitter cycling and cochlear mechanics. Thus, maintaining a rosette patterning within the organ of Corti may be important for function. Cochlear mechanics of the basilar membrane activate hair cell transduction. Due to the high sensitivity of cochlear mechanics, it is also desirable to avoid masses of cells. In all, maintaining proper distribution and relation of hair cells and supporting cells along the basilar membrane, even after proliferation, is likely a desired feature for hearing as supporting cell function and proper mechanics is necessary for normal hearing.

[0139] In one embodiment of the present disclosure, the cell density of hair cells in a cochlear cell population is expanded in a manner that maintains, or even establishes, the rosette pattern characteristic of cochlear epithelia.

[0140] In accordance with one aspect of the present disclosure, the cell density of hair cells may be increased in a population of cochlear cells comprising both hair cells and supporting cells. The cochlear cell population may be an *in vivo* population (*i.e.*, comprised by the cochlear epithelium of a subject) or the cochlear cell population may be an *in vitro* (*ex vivo*) population. If the population is an *in vitro* population, the increase in cell density may be determined by reference to a Representative Microscopy Sample of the population taken prior and subsequent to any treatment. If the population is an *in vivo* population, the increase in cell density may be determined indirectly by determining an effect upon the hearing of the subject with an increase in hair cell density correlating to an improvement in hearing.

[0141] In one embodiment, supporting cells placed in a Stem Cell Proliferation Assay in the absence of neuronal cells form ribbon synapses.

[0142] In a native cochlea, patterning of hair cells and supporting cells occurs in a manner parallel to the basilar membrane. In one embodiment of the present disclosure, the proliferation of supporting cells in a cochlear cell population is expanded in a manner that the basilar membrane characteristic of cochlear epithelia.

[0143] In one embodiment, the number of supporting cells in an initial cochlear cell population is selectively expanded by treating the initial cochlear cell population with a composition provided herein to form an intermediate cochlear cell population and wherein the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population. The expanded

cochlear cell population may be, for example, an *in vivo* population, an *in vitro* population or even an *in vitro* explant. In one such embodiment, the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population. For example, in one such embodiment the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population by a factor of 1.1. By way of further example, in one such embodiment the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population by a factor of 1.5. By way of further example, in one such embodiment the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population by a factor of 2. By way of further example, in one such embodiment the ratio of supporting cells to hair cells in the intermediate cochlear cell population exceeds the ratio of supporting cells to hair cells in the initial cochlear cell population by a factor of 3. In each of the foregoing embodiments, the capacity of a composition of the present disclosure to expand a cochlear cell population as described in this paragraph may be determined by means of a Stem Cell Proliferation Assay.

[0144] In one embodiment, the number of stem cells in a cochlear cell population is expanded to form an intermediate cochlear cell population by treating a cochlear cell population with a composition provided herein wherein the cell density of stem cells in the intermediate cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population. The treated cochlear cell population may be, for example, an *in vivo* population, an *in vitro* population or even an *in vitro* explant. In one such embodiment, the cell density of stem cells in the treated cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population by a factor of at least 1.1. For example, in one such embodiment the cell density of stem cells in the treated cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population by a factor of at least 1.25. For example, in one such embodiment the cell density of stem cells in the treated cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population by a factor of at least 1.5. By way of further example, in one such embodiment the cell density of stem cells in the treated cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population by a factor of at least 2. By way of further example, in one such embodiment the cell density of stem cells

in the treated cochlear cell population exceeds the cell density of stem cells in the initial cochlear cell population by a factor of at least 3. *In vitro* cochlear cell populations may expand significantly more than *in vivo* populations; for example, in certain embodiments the cell density of stem cells in an expanded *in vitro* population of stem cells may be at least 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2,000 or even 3,000 times greater than the cell density of the stem cells in the initial cochlear cell population. In each of the foregoing embodiments, the capacity of a composition of the present disclosure to expand a cochlear cell population as described in this paragraph may be determined by means of a Stem Cell Proliferation Assay.

[0145] In accordance with one aspect of the present disclosure, a cochlea supporting cell population is treated with a composition provided herein to increase the Lgr5 activity of the population. For example, in one embodiment the composition provided herein has the capacity to increase and maintain the Lgr5 activity of an *in vitro* population of cochlea supporting cells by factor of at least 1.2. By way of further example, in one such embodiment the compound has the capacity to increase the Lgr5 activity of an *in vitro* population of cochlea supporting cells by factor of 1.5. By way of further example, in one such embodiment the compound has the capacity to increase the Lgr5 activity of an *in vitro* population of cochlea supporting cells by factor of 2, 3, 5, 10, 100, 500, 1,000, 2,000 or even 3,000. Increases in Lgr5 activity may also be observed for *in vivo* populations but the observed increase may be somewhat more modest. For example, in one embodiment the compound has the capacity to increase the Lgr5 activity of an *in vivo* population of cochlea supporting cells by at least 5%. By way of further example, in one such embodiment the compound has the capacity to increase the Lgr5 activity of an *in vivo* population of cochlea supporting cells by at least 10%. By way of further example, in one such embodiment the compound has the capacity to increase the Lgr5 activity of an *in vivo* population of cochlea supporting cells by at least 20%. By way of further example, in one such embodiment the compound has the capacity to increase the Lgr5 activity of an *in vivo* population of cochlea supporting cells by at least 30%. In each of the foregoing embodiments, the capacity of the compound for such an increase in Lgr5 activity may be demonstrated, for example, in an *In Vitro* Lgr5⁺ Activity Assay and in an *in vivo* population may be demonstrated, for example, in an *In Vivo* Lgr5⁺ Activity Assay, as measured by isolating the organ and performing morphological

analyses using immunostaining, endogenous fluorescent protein expression of Lgr5 (eg. Lgr5, Sox2), and qPCR for Lgr5.

[0146] In addition to increasing the Lgr5 activity of the population, the number of Lgr5⁺ supporting cells in a cochlea cell population may be increased by treating a cochlea cell population containing Lgr5⁺ supporting cells (whether *in vivo* or *in vitro*) with a composition provided herein. In general, the cell density of the stem/progenitor supporting cells may expand relative to the initial cell population via one or more of several mechanisms. For example, in one such embodiment, newly generated Lgr5⁺ supporting cells may be generated that have increased stem cell propensity (*i.e.*, greater capacity to differentiate into hair cell). By way of further example, in one such embodiment no daughter Lgr5⁺ cells are generated by cell division, but pre-existing Lgr5⁺ supporting cells are induced to differentiate into hair cells. By way of further example, in one such embodiment no daughter cells are generated by cell division, but Lgr5⁻ supporting cells are activated to a greater level of Lgr5 activity and the activated supporting cells are then able to differentiate into hair cells. Regardless of the mechanism, in one embodiment the compound of the present disclosure has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vitro* isolated cell population of cochlea supporting cells by factor of at least 5. By way of further example, in one such embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vitro* population of cochlea supporting cells by factor of at least 10. By way of further example, in one such embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vitro* population of cochlea supporting cells by factor of at least 100, at least 500, at least 1,000 or even at least 2,000. Increases in the cell density of Lgr5⁺ supporting cells may also be observed for *in vivo* populations but the observed increase may be somewhat more modest. For example, in one embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vivo* population of cochlea supporting cells by at least 5%. By way of further example, in one such embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vivo* population of cochlea supporting cells by at least 10%. By way of further example, in one such embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vivo* population of cochlea supporting cells by at least 20%. By way of further example, in one such embodiment the compound has the capacity to increase the cell density of Lgr5⁺ supporting cells in an *in vivo* population of cochlea supporting

cells by at least 30%. The capacity of the compound for such an increase in Lgr5⁺ supporting cells in an *in vitro* population may be demonstrated, for example, in a Stem Cell Proliferation Assay or in an appropriate *in vivo* assay. In one embodiment, a compound of the present disclosure has the capacity to increase the number of Lgr5⁺ cells in the cochlea by inducing expression of Lgr5 in cells with absent or low detection levels of the protein, while maintaining Native Morphology. In one embodiment, a compound of the present disclosure has the capacity to increase the number of Lgr5⁺ cells in the cochlea by inducing expression of Lgr5 in cells with absent or low detection levels of the protein, while maintaining Native Morphology and without producing Cell Aggregates.

[0147] In addition to increasing the cell density of Lgr5⁺ supporting cells, in one embodiment the method of the present disclosure has the capacity to increase the ratio of Lgr5⁺ cells to hair cells in a cochlear cell population. In one embodiment, the number of Lgr5⁺ supporting cells in an initial cochlear cell population is selectively expanded by treating the initial cochlear cell population with a compound of the present disclosure to form an expanded cell population and wherein the number of Lgr5⁺ supporting cells in the expanded cochlear cell population at least equals the number of hair cells. The expanded cochlear cell population may be, for example, an *in vivo* population, an *in vitro* population or even an *in vitro* explant. In one such embodiment, the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 1:1. For example, in one such embodiment the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 1.5:1. By way of further example, in one such embodiment the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 2:1. By way of further example, in one such embodiment the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 3:1. By way of further example, in one such embodiment the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 4:1. By way of further example, in one such embodiment the ratio of Lgr5⁺ supporting cells to hair cells in the expanded cochlear cell population is at least 5:1. In each of the foregoing embodiments, the capacity of the compound of the present disclosure to expand a cochlear cell population as described in this paragraph may be determined by means of a Stem Cell Proliferation Assay.

[0148] In certain embodiments, the method increases the fraction of the Lgr5⁺ cells to total cells on the sensory epithelium by at least 10%, 20%, 50%, 100%, 250%, 500%, 1,000% or 5000%.

[0149] In certain embodiments, the method increases the Lgr5⁺ cells until they become at least 10, 20, 30, 50, 70, or 85 % of the cells on the sensory epithelium, e.g. the Organ of Corti.

[0150] In general, excessive proliferation of supporting cells in the cochlea is preferably avoided. In one embodiment, the method of the present disclosure has the capacity to expand a cochlear cell population without creating a protrusion of new cells beyond the native surface of the cochlea, e.g. a Cell Aggregate. In some embodiments, 30 days after placing a composition provided herein on the round or oval membrane, the cochlear tissue has Native Morphology. In some embodiments, 30 days after placing the compound on the round or oval membrane, the cochlear tissue has Native Morphology and lacks Cell Aggregates. In some embodiments, 30 days after placing the compound on the round or oval membrane, the cochlear tissue has Native Morphology and at least 10, 20, 30, 50, 75, 90, 95, 98, or even at least 99% of the Lgr5⁺ cells in the Organ of Corti are not part of Cell Aggregates.

[0151] In addition to expanding supporting cell populations, generally, and Lgr5⁺ supporting cells, specifically, as described above, the method of the present disclosure has the capacity to maintain, in the daughter cells, the capacity to differentiate into hair cells. In *in vivo* populations, the maintenance of this capacity may be indirectly observed by an improvement in a subject's hearing. In *in vitro* populations, the maintenance of this capacity may be directly observed by an increase in the number of hair cells relative to a starting population or indirectly by measuring LGR5 activity, SOX2 activity or one or more of the other stem cell markers identified elsewhere herein.

[0152] In one embodiment, the capacity of the method to increase the stemness of a population of cochlear supporting cells, in general, or a population of Lgr5⁺ supporting cells, in particular, may be correlated with an increase of Lgr5 activity of an *in vitro* population of isolated Lgr5⁺ cells as determined by an Lgr5 Activity Assay. As previously noted, in one such embodiment, the compound has the capacity to increase the Lgr5 activity of stem cells in the intermediate cell population by a factor of 5 on average relative to the Lgr5 activity of the cells in the initial cell population. By way of further example, in one such embodiment the method has the capacity to increase the Lgr5 activity of the stem cells genes in the intermediate cell population by a factor

of 10 relative to the Lgr5 activity of the cells in the initial cell population. By way of further example, in one such embodiment the method has the capacity to increase the Lgr5 activity of the stem cells in the intermediate cell population by a factor of 100 relative to the Lgr5 activity of the cells in the initial cell population. By way of further example, in one such embodiment the method has the capacity to increase the Lgr5 activity of the stem cells in the intermediate cell population by a factor of 1000 relative to the Lgr5 activity of the cells in the initial cell population. In each of the foregoing embodiments, the increase in the activity of stem cells in the cell population may be determined *in vitro* by immunostaining or endogenous fluorescent protein expression for target genes and analysis of their relative intensities via imaging analysis or flow cytometry, or using qPCR for target stem cell genes. The identity of the resulting stem cell population may optionally be further determined by stem cell assays including stem cell marker expression assay, colony forming assay, self-renewal assay and differentiation assay as defined in Stem cell assay.

[0153] In some embodiments, the method applied to an adult mammal produces a population of adult mammalian Lgr5⁺ cells that are in S-phase.

[0154] In one embodiment, after applying the composition provided herein to the round or oval of a mouse, the *in vivo* Lgr5⁺ Activity of a cell population in the Organ of Corti increases 1.3x, 1.5x, up to 20x over baseline for a population that has not been exposed to the compound. In some embodiments, applying the compound to the round or oval of a mouse increases the average *In vivo* Lgr5⁺ Activity for cells in the Organ of Corti is increased 1.3x, 1.5x, up to 20x over baseline for a population that has not been exposed to the compound.

[0155] In certain embodiments, the method increases the Lgr5⁺ cells until they become at least 10%, 7.5%, 10%, up to 100% of the supporting cell population by number.

[0156] In certain embodiments, the compound has the capacity to increase the percentage of Lgr5⁺ cell in a cochlea by 5%, 10%, 25%, 50%, or 80%.

[0157] In certain embodiments, the stem cell population is of an *in vivo* subject, and the method is a treatment for hearing loss and/or vestibular dysfunction (*e.g.*, wherein the generation of inner ear hair cells from the expanded population of stem cells results in partial or full recovery of hearing loss and/or improved vestibular function). In certain embodiments, the stem cell population is of an *in vivo* subject, and the method further comprises delivering a drug to the subject (*e.g.*, for treatment of a disease and/or disorder unrelated to hearing loss and/or vestibular

dysfunction) at a higher concentration than a known safe maximum dosage of the drug for the subject (*e.g.*, the known safe maximum dosage if delivered in the absence of the generation of inner ear hair cells resulting from the method) (*e.g.*, due to a reduction or elimination of a dose-limiting ototoxicity of the drug).

[0158] In certain embodiments, the method further comprises performing high throughput screening using the generated inner ear hair cells. In certain embodiments, the method comprises using the generated inner ear hair cells to screen molecules for toxicity against inner ear hair cells. In certain embodiments, the method comprises using the generated inner ear hair cells to screen molecules for ability to improve survival of inner ear hair cells (*e.g.*, inner ear hair cells exposed to said molecules).

[0159] In another aspect, the disclosure is directed to a method of producing an expanded population of stem cells, the method comprising: administering or causing to be administered to a stem cell population (*e.g.*, of an *in vitro*, *ex vivo*, or *in vivo* sample/subject) a composition provided herein.

[0160] In certain embodiments, the administering step is carried out by performing one or more injections into the ear (*e.g.*, transtympanically into the middle ear and/or inner ear).

[0161] In certain embodiments, the administering step comprises administering the GSK3-beta inhibitor and/or Wnt agonist in a sustained manner.

[0162] In certain embodiments, the stem cells are inner ear stem cells and/or supporting cells.

[0163] In certain embodiments, the method further comprises performing high throughput screening using the generated expanded population of stem cells. In certain embodiments, the method further comprises using the generated stem cells to screen molecules for toxicity against stem cells and/or their progeny. In certain embodiments, the method comprises using the generated stem cells to screen molecules for ability to improve survival of stem cells and/or their progeny.

[0164] In another aspect, the disclosure is directed to a method of treating a subject who has, or is at risk of developing, hearing loss and/or vestibular dysfunction, the method comprising: identifying a subject who has experienced, or is at risk for developing, hearing loss and/or vestibular dysfunction, administering or causing to be administered a composition provided herein.

[0165] In certain embodiments, the stem cell population comprises Lgr5⁺ cells. In certain embodiments, the stem cell population comprises post-natal cells. In certain embodiments, the stem cell population comprises epithelial stem cells. In certain embodiments, stem cells include progenitor cells.

[0166] In certain embodiments, the step of administering is carried out by performing one or more injections into the ear (*e.g.*, transtympanically into the middle ear and/or inner ear).

[0167] In another aspect, the disclosure is directed to a method of generating inner ear hair cells, the method comprising: proliferating stem cells in an initial stem cell population (*e.g.*, of an *in vitro*, *ex vivo*, or *in vivo* sample/subject), resulting in an expanded population of stem cells (*e.g.*, such that the expanded population is a factor of at least 1.25, 1.5, 1.75, 2, 3, 5, 10, or 20 greater than the initial stem cell population); and facilitating generation of inner ear hair cells from the expanded population of stem cells.

[0168] In another aspect, the disclosure is directed to a method of generating inner ear hair cells, the method comprising administering a composition provided herein (*e.g.*, in a pharmaceutically acceptable form (*e.g.*, salt)) to a cell population in an inner ear of a subject, thereby facilitating generation of inner ear hair cells.

[0169] In another aspect, the disclosure is directed to a method of generating inner ear hair cells, the method comprising: proliferating post-natal LGR5⁺ cells in an initial population (*e.g.*, of an *in vitro*, *ex vivo*, or *in vivo* sample/subject), resulting in an expanded population of LGR5⁺ cells (*e.g.*, such that the expanded population is a factor of at least 1.25, 1.5, 1.75, 2, 3, 5, 10, or 20 greater than the initial stem cell population), said expanded population of LGR5⁺ cells resulting in generation of inner ear hair cells. In certain embodiments, stem cells include progenitor cells.

[0170] In another aspect, the disclosure is directed to a method of treating a disease or disorder, the method comprising: proliferating post-natal Lgr5⁺ epithelial cells in an initial population of a subject (*in vivo*), resulting in an expanded population of Lgr5⁺ epithelial cells (*e.g.*, such that the expanded population is a factor of at least 1.25, 1.5, 1.75, 2, 3, 5, 10, or 20 greater than the initial post-natal Lgr5⁺ epithelial cell population).

[0171] In some embodiments, Lgr5⁺ cells are differentiated into hair cells.

Hair Cell Regrowth

[0172] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of inner ear hair cells, the method comprising: administering a compound of present disclosure to expand the stem cell population of cochlear tissue.

[0173] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of inner ear hair cells, the method comprising: administering a composition comprising a compound of present disclosure and HDAC inhibitor to expand the stem cell population of cochlear tissue.

[0174] In certain embodiments, the present disclosure is directed to a method to regenerate hearing in mammals.

[0175] In certain embodiments, the stem cell population is of an in vivo subject, and the method is a treatment for hearing loss and/or vestibular dysfunction.

[0176] In certain embodiments, the present disclosure is directed to a method of generating inner ear hair cells using of a compound of the present disclosure to proliferate LGR5+ cells in an initial population in vivo, resulting in an expanded population of LGR5+ cells (e.g., such that the expanded population is at least 2 times, 3 times, 5 times, 10 times, or 20 times greater than the initial stem cell population), resulting in generation of inner ear hair cells.

[0177] In certain embodiments, the present disclosure is directed to a method of generating inner ear hair cells using of a composition comprising a compound of present disclosure and HDAC inhibitor to proliferate LGR5+ cells in an initial population in vivo, resulting in an expanded population of LGR5+ cells (e.g., such that the expanded population is at least 2 times, 3 times, 5 times, 10 times, or 20 times greater than the initial stem cell population), resulting in generation of inner ear hair cells.

Intestinal Regeneration

[0178] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of intestinal cells, the method comprising: administering a compound of the present disclosure to expand the stem cell population of intestinal epithelia.

[0179] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of intestinal cells, the method comprising: administering a composition comprising a compound of present disclosure and HDAC inhibitor to expand the stem cell population of intestinal epithelia.

[0180] In certain embodiments, the present disclosure is directed to a method to regenerate intestinal epithelia in mammals.

[0181] In certain embodiments, the stem cell population is of an in vivo subject. In certain embodiments, the method is a treatment for promoting the repair of damaged mucosa related to diseases such as chemotherapy-induced gastrointestinal mucositis, Graph Versus Host Disease, gastric ulcer, Crohns, or ulcerative colitis.

Intestinal Lgr5+ Proliferation

[0182] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of intestinal cells, the method comprising: administering a compound of the present disclosure to expand the Lgr5+ cell population of intestinal epithelia.

[0183] In certain embodiments, the present disclosure is directed to a method of facilitating the generation of intestinal cells, the method comprising: administering a composition comprising a compound of present disclosure and HDAC inhibitor to expand the Lgr5+ cell population of intestinal epithelia.

[0184] In certain embodiments, the present disclosure is directed to a method to regenerate Lgr5+ cell population intestinal cells in mammals.

[0185] In certain embodiments, the Lgr5+ cell population is in an in vivo subject. In certain embodiments, the method is a treatment for promoting the repair of damaged mucosa related to diseases such as chemotherapy-induced gastrointestinal mucositis, Graph Versus Host Disease, gastric ulcer, Crohns, or ulcerative colitis.

[0186] In certain embodiments, the present disclosure is directed to a method of treating a disease or disorder, the method comprising proliferating Lgr5+ epithelial cells in vivo, resulting in an expanded population of Lgr5+ epithelial cells (e.g., such that the expanded population is at least 2 times, 3 times, 5 times, 10 times, or 20 times greater than the initial post-natal Lgr5+ epithelial cell population).

Expansion of a population of vestibular cells

[0187] In certain embodiments, the pharmaceutical formulations containing can expand a population of vestibular cells in a vestibular tissue comprising contacting the vestibular tissue. In certain embodiments, the pharmaceutical formulations are capable in a stem cell proliferation assay of increasing the number of supporting cells in a stem cell proliferation assay cell population by a factor of at least 10 or at least 50. In certain embodiments, the pharmaceutical

formulations are capable in a stem cell differentiation assay of forming hair cells from a cell population comprising vestibular supporting cells.

[0188] In certain embodiments, the vestibular tissue maintains Native Morphology. In certain embodiments, the vestibular tissue is in a subject. In certain embodiments, the contacting the vestibular tissue with the composition is achieved by administering the composition trans-tympanically to the subject. In certain embodiments, the contacting the vestibular tissue with the composition results in improved vestibular functioning of the subject.

[0189] In certain embodiments, the present disclosure is directed to a method of treating a subject who has, or is at risk of developing, a disease associated with absence or lack of certain tissue cells, the method comprising administering or causing to be administered to said subject a compound of the present disclosure.

[0190] In certain embodiments, the compound is dispersed in a biocompatible matrix. In certain embodiments, the biocompatible matrix is a biocompatible gel or foam. In certain embodiments, the compound is administered trans-tympanically to a vestibular tissue of the subject.

[0191] In certain embodiments, the present disclosure provides a method for expanding a population of vestibular cells in a vestibular tissue comprising contacting the vestibular tissue with (i) a compound of the present disclosure, and (ii) a TGF- β Inhibitor to form an expanded population of cells in the vestibular tissue.

Generation of Dermal Papilla Cells

[0192] In certain embodiments, the present disclosure is directed to a method of facilitating generation of Dermal Papilla Cells, the method comprising: administering a compound of the present disclosure, alone or in combination with a BMP inhibitor, to expand the population of Dermal Papilla Cells. In certain embodiments, the compounds can regenerate hair in a mammal. In certain embodiments, the Dermal Papilla Cells population is of an in vivo subject. In certain embodiments, the Dermal Papilla Cells population is of an in vivo subject for the treatment for alopecia. In certain embodiments, the present disclosure provides a method of generating Dermal Papilla Cells using of a compound of the present disclosure, alone or in combination with BMP inhibitor to proliferate Dermal Papilla Cells in an initial population in vivo, resulting in an expanded population of Dermal Papilla Cells.

Administration

[0193] The membrane of the round or oval is the biological barrier to the inner ear space and represents the major obstacle for the local treatment of hearing impairment. The administered drug must overcome this membrane to reach the inner ear space. The drug can operatively (*e.g.*, injection through the tympanic membrane) be placed locally to the round or oval membrane and can then penetrate through the round or oval membrane. Substances that penetrate the round or oval typically distribute in the perilymph and thus reach the hair cells and supporting cells.

[0194] In certain embodiments, pharmaceutical formulations are adapted to administer the drug locally to the round or oval membrane. The pharmaceutical formulations may also contain a membrane penetration enhancer, which supports the passage of the agents mentioned herein through the round or oval membrane. Accordingly, liquid, gel or foam formulations may be used. It is also possible to apply the active ingredient orally or to employ a combination of delivery approaches.

[0195] Intratympanic (IT) delivery of drugs to the ear is increasingly used for both clinical and research purposes. Some groups have applied drugs in a sustained manner using microcatheters and microwicks, while the majority have applied them as single or as repeated IT injections (up to 8 injections over periods of up to 2 weeks).

[0196] Intratympanically applied drugs are thought to enter the fluids of the inner ear primarily by crossing the round or oval (RW) membrane. Calculations show that a major factor controlling both the amount of drug entering the ear and the distribution of drug along the length of the ear is the duration the drug remains in the middle ear space. Single, 'one-shot' applications or applications of aqueous solutions for few hours' duration result in steep drug gradients for the applied substance along the length of the cochlea and rapidly declining concentration in the basal turn of the cochlea as the drug subsequently becomes distributed throughout the ear.

[0197] Other injection approaches include by osmotic pump, or, by combination with implanted biomaterial, and more preferably, by injection or infusion. Biomaterials that can aid in controlling release kinetics and distribution of drug include hydrogel materials, degradable materials. One class of materials that is most preferably used includes in situ gelling materials. Other materials include collagen or other natural materials including fibrin, gelatin, and decellularized tissues. Gelfoam may also be suitable.

[0198] Delivery may also be enhanced via alternate means including but not limited to agents added to the delivered composition such as penetration enhancers, or could be through devices via ultrasound, electroporation, or high speed jet.

[0199] Methods described herein can also be used for inner ear cell types that may be produced using a variety of methods know to those skilled in the art including those cell types described in PCT Application No. WO2012103012 A1.

[0200] With regard to human and veterinary treatment, the amount of a particular agent(s) that is administered may be dependent on a variety of factors, including the disorder being treated and the severity of the disorder; activity of the specific agent(s) employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific agent(s) employed; the duration of the treatment; drugs used in combination or coincidental with the specific agent(s) employed; the judgment of the prescribing physician or veterinarian; and like factors known in the medical and veterinary arts.

[0201] The agents described herein may be administered in a therapeutically effective amount to a subject in need of treatment. Administration of compounds described herein can be via any of suitable route of administration, particularly by intratympanically. Other routes include ingestion, or alternatively parenterally, for example intravenously, intra-arterially, intraperitoneally, intrathecally, intraventricularly, intraurethrally, intrasternally, intracranially, intramuscularly, intranasally, subcutaneously, sublingually, transdermally, or by inhalation or insufflations, or topical by ear instillation for absorption through the skin of the ear canal and membranes of the eardrum. Such administration may be as a single or multiple oral dose, defined number of ear drops, or a bolus injection, multiple injections, or as a short- or long-duration infusion. Implantable devices (*e.g.*, implantable infusion pumps) may also be employed for the periodic parenteral delivery over time of equivalent or varying dosages of the particular formulation. For such parenteral administration, the compounds are preferably formulated as a sterile solution in water or another suitable solvent or mixture of solvents. The solution may contain other substances such as salts, sugars (particularly glucose or mannitol), to make the solution isotonic with blood, buffering agents such as acetic, citric, and/or phosphoric acids and their sodium salts, and preservatives.

[0202] Compounds described herein can be administered by a number of methods sufficient to deliver the compound to the inner ear. Delivering a compound to the inner ear includes

administering the compound to the middle ear, such that the compound may diffuse across the round or oval to the inner ear and administering a compound to the inner ear by direct injection through the round or oval membrane. Such methods include, but are not limited to auricular administration, by transtympanic wicks or catheters, or parenteral administration, for example, by intraauricular, transtympanic, or intracochlear injection.

[0203] In particular embodiments, the compounds, compositions and formulations of the disclosure are locally administered, meaning that they are not administered systemically.

[0204] In one embodiment, a syringe and needle apparatus is used to administer compounds or compositions to a subject using auricular administration. A suitably sized needle is used to pierce the tympanic membrane and a wick or catheter comprising the composition is inserted through the pierced tympanic membrane and into the middle ear of the subject. The device may be inserted such that it is in contact with the round or oval or immediately adjacent to the round or oval. Exemplary devices used for auricular administration include, but are not limited to, transtympanic wicks, transtympanic catheters, round or oval microcatheters (small catheters that deliver medicine to the round or oval), and Silverstein Microwicks™ (small tube with a "wick" through the tube to the round or oval, allowing regulation by subject or medical professional).

[0205] In another embodiment, a syringe and needle apparatus is used to administer compounds or compositions to a subject using transtympanic injection, injection behind the tympanic membrane into the middle and/or inner ear. The formulation may be administered directly onto the round or oval membrane via transtympanic injection or may be administered directly to the cochlea via intracochlear injection or directly to the vestibular organs via intravestibular injection.

[0206] In some embodiments, the delivery device is an apparatus designed for administration of compounds or compositions to the middle and/or inner ear. By way of example only: GYRUS Medical GmbH offers micro-otoscopes for visualization of and drug delivery to the round or oval niche; Arenberg has described a medical treatment device to deliver fluids to inner ear structures in U.S. Pat. Nos. 5,421,818; 5,474,529; and 5,476,446, each of which is incorporated by reference herein for such disclosure. U.S. patent application Ser. No. 08/874,208, which is incorporated herein by reference for such disclosure, describes a surgical method for implanting a fluid transfer conduit to deliver compositions to the inner ear. U.S. Patent Application Publication 2007/0167918, which is incorporated herein by reference for such disclosure, further

describes a combined otic aspirator and medication dispenser for trans-tympanic fluid sampling and medicament application.

[0207] In some embodiments, composition provided herein is administered to a subject in need thereof once. In some embodiments, composition provided herein is administered to a subject in need thereof more than once. In some embodiments, a first administration of composition provided herein is followed by a second, third, fourth, or fifth administration of composition provided herein.

[0208] The number of times a compound is administered to an subject in need thereof depends on the discretion of a medical professional, the disorder, the severity of the disorder, and the subject's response to the formulation. In some embodiments, the compound disclosed herein is administered once to a subject in need thereof with a mild acute condition. In some embodiments, the compound disclosed herein is administered more than once to a subject in need thereof with a moderate or severe acute condition. In the case wherein the subject's condition does not improve, upon the doctor's discretion the compound may be administered chronically, that is, for an extended period of time, including throughout the duration of the subject's life in order to ameliorate or otherwise control or limit the symptoms of the subject's disease or condition.

[0209] In the case wherein the subject's status does improve, upon the doctor's discretion the compound may administered continuously; alternatively, the dose of drug being administered may be temporarily reduced or temporarily suspended for a certain length of time (*i.e.*, a "drug holiday"). The length of the drug holiday varies between 2 days and 1 year, including by way of example only, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 10 days, 12 days, 15 days, 20 days, 28 days, 35 days, 50 days, 70 days, 100 days, 120 days, 150 days, 180 days, 200 days, 250 days, 280 days, 300 days, 320 days, 350 days, and 365 days. The dose reduction during a drug holiday may be from 10%- 100%, including by way of example only 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, and 100%.

[0210] Once the subject's hearing and/or balance has improved, a maintenance dose can be administered, if necessary. Subsequently, the dosage or the frequency of administration, or both, is optionally reduced, as a function of the symptoms, to a level at which the improved disease, disorder or condition is retained. In certain embodiments, subjects require intermittent treatment on a long-term basis upon any recurrence of symptoms.

[0211] In certain embodiments, the pharmaceutical formulations may also contain an additional agent selected from a Notch activator, HDAC inhibitor, a BMP4 antagonist, Noggin (Inhibits BMP4), Sox2, Vitamin D (calcitriol), Vitamin B (nicotinamide), Vitamin A, Vitamin C (pVc), Lgr4, p38/MAPK inhibition, ROCK inhibition, and/or Alk4/7 inhibition. In certain embodiments, the pharmaceutical formulations may also contain an epidermal growth factor (EGF), fibroblast growth factor (FGF), insulin-like growth factor (IGF), or a combination thereof.

Compositions with HDAC

[0212] In certain embodiments, the pharmaceutical formulations may also contain HDAC. In certain embodiments, the pharmaceutical formulations containing HDAC can enhance the formation of Lgr5+ cells, control differentiation, control stemness, and replication or restore hearing and intestinal regeneration.

[0213] In certain embodiments, the HDAC inhibitor is Valproic acid or a prodrug, ester, salt form, or amide thereof.

[0214] In certain embodiments, the HDAC inhibitor is a carboxylic acid containing compound. In certain embodiments, the carboxylic acid containing compound is C₆-C₂₀ carboxylic acid, wherein the carboxylic acid comprises alkyl, alkenyl, or alkynyl.

[0215] In certain embodiments, the carboxylic acid containing compound is a substituted or unsubstituted C₅-C₂₀ straight, branched, or cyclic chain alkyl-CO₂H, substituted or unsubstituted C₅-C₂₀ straight, branched, or cyclic chain alkenyl-CO₂H and substituted or unsubstituted C₅-C₂₀ straight, branched, or cyclic chain alkynyl-CO₂H. In certain embodiments, the carboxylic acid containing compound is a substituted C₅-C₂₀ straight or branched chain alkyl-CO₂H.


[0216] In certain embodiments, the carboxylic acid containing compound is a substituted C₅-C₂₀ straight or branched chain alkyl-CO₂H, wherein the substituent is -NH₂. In certain embodiments, the carboxylic acid containing compound is an amino substituted 2-propylpentanoic acid. In certain embodiments, the amino substituted 2-propylpentanoic acid is selected from the group consisting of 5-amino-2-propylpentanoic acid, 4-amino-2-propylpentanoic acid, 3-amino-2-propylpentanoic acid, and 2-amino-2-propylpentanoic acid.

[0217] In certain embodiments, the carboxylic acid containing compound is an unsubstituted C₅-C₂₀ straight or branched chain alkyl-CO₂H. In certain embodiments, the carboxylic acid containing compound is an unsubstituted C₆-C₉ branched straight chain alkyl-CO₂H. In certain

embodiments, the carboxylic acid containing compound is an unsubstituted C₈-C₉ branched straight chain alkyl-CO₂H. In certain embodiments, the carboxylic acid containing compound is an unsubstituted C₈ branched straight chain alkyl-CO₂H.

[0218] In certain embodiments, the carboxylic acid containing compound is Valproic acid.

[0219] In certain embodiments, the carboxylic acid containing compound is in the form of a prodrug of an unsubstituted C₈ branched straight chain alkyl-CO₂H wherein the prodrug is in the form of an amide or ester. In certain embodiments, the amide of unsubstituted C₈ branched straight chain alkyl-CO₂H is the condensation product with an amino acid. In certain embodiments, the amide of Valproic acid is selected from the group consisting of



[0220] In some embodiments, the HDAC inhibitor is any one of the inhibitors listed in Table 1.

Table 1 - HDAC Inhibitors

Column A	Column B	CAS Number
Class	Agent	
Aliphatic Acid	Valproic Acid	99-66-1
Aliphatic Acid	Phenyl butyrate	1821-12-1
Aliphatic Acid	Butyrate	107-92-6
Aliphatic Acid	2-(prop-2-yn-1-yl)octanoic acid	96017-59-3
Aliphatic Acid	(S)-2-(prop-2-yn-1-yl)octanoic acid	185463-37-0
Aliphatic Acid	(R)-2-(prop-2-yn-1-yl)octanoic acid	185463-38-1
Aliphatic Acid	2-(prop-2-yn-1-yl)heptanoic acid	176638-49-6
Aliphatic Acid	(S)-2-(prop-2-yn-1-yl) heptanoic acid	185463-37-0
Aliphatic Acid	(R)-2-(prop-2-yn-1-yl) heptanoic acid	185463-38-1
Aliphatic Acid	2-fluoro-2-propyl Pentanoic acid	197779-85-4
Aliphatic Acid Ester	AN-9	122110-53-6
Amine	932718-22-4	932718-22-4
Benzamide	Entinostat (MS-275)	209783-80-2
Benzamide	Mocetinostat (MGCD0103)	726169-73-9

Column A	Column B	CAS Number
Class	Agent	
Benzamide	Tacedinaline	112522-64-2
Benzamide	BML-210	537034-17-6
Benzamide	NKL 22	537034-15-4
Benzamide	RGFP109	1215493-56-3
Benzamide	RGFP136	1215493-97-2
Benzamide	RGFP966	1357389-11-7
Benzamide	4SC-202	1186222-89-8
Benzamide	HDAC Inhibitor IV	537034-15-4
Benzamide	Chidamide	743438-44-0
Benzamide	TC-H 106, HDAC Inhibitor VII	937039-45-7
Cyclic peptide	Romidepsin	128517-07-7
Cyclic peptide	Trapoxin A	133155-89-2
Cyclic peptide	HC Toxin	83209-65-8
Cyclic peptide	Apicidin	183506-66-3
Cyclic Peptide	Thailandepsin A	1269219-30-8
Cyclic peptide	Dihydrochlamydocin	52574-64-8
Epoxide	(-)-Depudecin	139508-73-9
Epoxide	Parthenolide	20554-84-1
Hydroxamate	Trichostatin A (TSA)	
Hydroxamate	Trichostatin A (TSA)	58880-19-6
Hydroxamate	SAHA (Zolinza, vorinostat)	149647-78-9
Hydroxamate	4-iodo-SAHA	1219807-87-0
Hydroxamate	SBHA	38937-66-5
Hydroxamate	CBHA	174664-65-4
Hydroxamate	LAQ-824	591207-53-3
Hydroxamate	PDX-101 (belinostat)	866323-14-0
Hydroxamate	LBH-589 (panobinostat)	404950-80-7
Hydroxamate	ITF2357 (Givinostat)	497833-27-9
Hydroxamate	PCI-34051	950762-95-5
Hydroxamate	PCI-24781 (Abexinostat)	783355-60-2
Hydroxamate	Tubastatin A	1252003-15-8
Hydroxamate	CUDC-101	1012054-59-9
Hydroxamate	Oxamflatin	151720-43-3
Hydroxamate	ITF2357	497833-27-9
Hydroxamate	Bufexamac	2438-72-4
Hydroxamate	APHA Compound 8	676599-90-9

Column A	Column B	CAS Number
Class	Agent	
Hydroxamate	HDAC Inhibitor XXIV	854779-95-6
Hydroxamate	Tubacin	537049-40-4
Hydroxamate	Butyrylhydroxamic acid	4312-91-8
Hydroxamate	MC 1568	852475-26-4
Hydroxamate	SB939 (Pracinostat)	929016-96-6
Hydroxamate	4SC-201 (Resminostat)	864814-88-0
Hydroxamate	Tefinostat (CHR-2845)	914382-60-8
Hydroxamate	CHR-3996	1256448-47-1
Hydroxamate	NSC 57457	6953-61-3
Hydroxamate	CG200745	936221-33-9
Hydroxamate	ACY1215	1316214-52-4
Hydroxamate	Nexturastat A	1403783-31-2
Hydroxamate	Droxinostat	99873-43-5
Hydroxamate	Scriptaid	287383-59-9
Hydroxamate	BRD9757	1423058-85-8
Hydroxamate	HPOB	1429651-50-2
Hydroxamate	CAY10603	1045792-66-2
Hydroxamate	HDAC6 Inhibitor III	1450618-49-1
Hydroxamate	M 344	251456-60-7
Hydroxamate	4-(dimethylamino)-N-[6-(hydroxyamino)-6-oxohexyl]-benzamide	193551-00-7
Hydroxamate	(S)-HDAC-42	935881-37-1
Hydroxamate	HNHA	926908-04-5
Hydroxamate	Pyroxamide	382180-17-8
Hydroxamate	HDAC Inhibitor VI	926908-04-5
Hydroxamate	HDAC Inhibitor II	174664-65-4
Hydroxamate	LMK235	1418033-25-6
Hydroxamate	HDAC-IN-1	1239610-44-6
Hydroxamate	VAHA	106132-78-9
Ketone - CF3	Compound 6e	946500-31-8
Ketone - CF3	Compound 6H	946500-39-6
Ketone - CF3	Compound 27	946499-86-1
Ketone	Compound 43	891259-76-0
Ketone - α -ketoamides	436150-82-2	436150-82-2
Polyketide	Ratjadone A	163564-92-9
Silylalcohol	1587636-32-5	1587636-32-5

Column A	Column B	CAS Number
Class	Agent	
Sulphonyl Urea	960130-17-0	960130-17-0
Sulphonamide	1587636-33-6	1587636-33-6
Sulphonamide	329967-25-1	329967-25-1
Thiol	1428536-05-3	1428536-05-3
Thiol	908860-21-9	908860-21-9
Thiol	828920-13-4	828920-13-4
Thiol	1368806-68-1	1368806-68-1
Thiol	827036-76-0	827036-76-0
Thioester	TCS HDAC6 20b	956154-63-5
Thioester	PTACH	848354-66-5
Thioester	KD 5170	940943-37-3
Thioester	HDAC Inhibitor XXII	848354-66-5
Thioketone	SIRT1/2 Inhibitor VII	143034-06-4
Tropones	46189-88-2	46189-88-2
Tropones	1411673-95-4	1411673-95-4
Non classical	TMP269	1314890-29-3
Non classical	Tasquinimod	254964-60-8

[0029] Classes of HDAC inhibitors for use in various embodiments of the compositions and methods disclosed herein include but are not limited to those listed in Column A of Table 1. Specific HDAC inhibitors for use in various embodiments of the compositions and methods disclosed herein include but are not limited to those listed in Column B of Table 1. All agents listed in Table 1 column B are understood to include derivatives or pharmaceutically acceptable salts thereof. All classes listed in Table 1 column A are understood to include both agents comprising that class and derivatives or pharmaceutically acceptable salts thereof.

[0221] In certain embodiments, the amount of the carboxylic acid containing compound is between least 2 wt % (weight carboxylic acid containing compound/weight pharmaceutical composition) and 20 wt %. In certain embodiments, the composition comprises at least 4 wt % carboxylic acid. In certain embodiments, the composition comprises at least 8 wt % carboxylic

acid. In certain embodiments, the composition comprises at least 12 wt % carboxylic acid. In certain embodiments, the composition comprises at least 16 wt % carboxylic acid. In certain embodiments, the composition comprises at least 20 wt % carboxylic acid.

Compositions with BMP inhibitor

[0222] In certain embodiments, the pharmaceutical formulations may also contain BMP inhibitor. Examples of BMP inhibitors are shown herein. Other examples are described in WO2014138088A1 and WO2016054406A1, which are incorporated herein by reference in their entireties.

BMP Inhibitor II	1206711-16-1
dorsomorphin	866405-64-3
ML347	1062368-49-3
LDN-193189	1062368-24-4

Compositions with TGF-beta inhibitor

[0223] In certain embodiments, the pharmaceutical formulations may also contain TGF-beta inhibitor. In certain embodiments, the pharmaceutical formulations containing TGF-beta inhibitor can expand a population of vestibular cells in a vestibular tissue comprising contacting the vestibular tissue. In certain embodiments, the pharmaceutical formulations containing TGF-beta inhibitor are capable in a stem cell proliferation assay of increasing the number of supporting cells in a stem cell proliferation assay cell population by a factor of at least 10 or at least 50. In certain embodiments, the pharmaceutical formulations containing TGF-beta inhibitor are capable in a stem cell differentiation assay of forming hair cells from a cell population comprising vestibular supporting cells.

[0224] In one embodiment, the TGF-beta inhibitor is selected from 616452 (Repsox), Galunisertib (LY2157299), EW-719, IN-1130, EW-7203, EW-7195, SM16, R 268712, GW788388, and PF-03671148.

[0225] Exemplary TGF-β Inhibitors appear in Table 2. TGF-beta type I receptor inhibitors include but are not limited to 2-(3-(6-Methylpyridin-2-yl)-1H-pyrazol-4-yl)-1,5 naphthyridine, [3-(Pyridin-2-yl)-4-(4-quinoyl)]-1H-pyrazole, and 3-(6-Methylpyridin-2-yl)-4-(4-quinoly)-1-phenylthiocarbamoyl-1H-pyrazole, which can be purchased from Calbiochem (San Diego, Calif.). Other small molecule inhibitors include, but are not limited to, SB-431542 (see e.g., Halder et al., 2005; Neoplasia 7(5):509-521), SM16 (see e.g., Fu, K et al., 2008; Arteriosclerosis,

Thrombosis and Vascular Biology 28(4):665), and SB-505124 (see e.g., Dacosta Byfield, S., et al., 2004; Molecular Pharmacology 65:744-52), among others.

Table 2. TGF- β Inhibitors

Class	Agent	CAS Number	Alternative Name
Tgf-beta-R1 inhibitor	LY-364947	396129-53-6	616451, TGF- β RI Kinase Inhibitor I, [3-(Pyridin-2-yl)-4-(4-quinonyl)]-1H-pyrazole, ALK5 Inhibitor I, LY-364947, HTS-466284
Tgf-beta-R1 inhibitor	Repsox	446859-33-2	616452, TGF- β RI Kinase Inhibitor II, 2-(3-(6-Methylpyridin-2-yl)-1H-pyrazol-4-yl)-1,5-naphthyridine
Tgf-beta-R1 inhibitor	SB-505124	356559-13-2	616453, TGF- β RI Kinase Inhibitor III, CAS 356559-13-2 2-(5-Benzo[1,3]dioxol-4-yl-2-tert-butyl-1H-imidazol-4-yl)-6-methylpyridine, HCl, ALK5 Inhibitor III,
Tgf-beta-R1 inhibitor	A-83-01	909910-43-6	616454, TGF- β RI Kinase Inhibitor IV - 3-(6-Methylpyridin-2-yl)-4-(4-quinolyl)-1-phenylthiocarbamoyl-1H-pyrazole, A-83-01, ALK5 Inhibitor IV
Tgf-beta-R1 inhibitor	SD-208	627536-09-8	616456, TGF- β RI Kinase Inhibitor V, 2-(5-Chloro-2-fluorophenyl)pteridin-4-yl)pyridin-4-yl amine, SD-208, ALK5 Inhibitor V
Tgf-beta-R1 inhibitor	SB-431542	301836-41-9	616461, TGF- β RI Kinase Inhibitor VI, 4-[4-(3,4-Methylenedioxyphenyl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzamide, Dihydrate, 4-[4-(1,3-Benzodioxol-5-yl)-5-(2-pyridyl)-1H-imidazol-2-yl]benzamide, Dihydrate
Tgf-beta-R1 inhibitor	TGF- β RI Kinase Inhibitor VII	666729-57-3	616458, TGF- β RI Kinase Inhibitor VII, 1-(2-((6,7-Dimethoxy-4-quinolyl)oxy)-(4,5-dimethylphenyl)-1-ethanone, ALK5 Inhibitor VII
Tgf-beta-R1 inhibitor	SB-525334	356559-20-1	616459, TGF- β RI Kinase Inhibitor VIII - SB-525334, 6-(2-tert-Butyl-5-(6-methyl-pyridin-2-yl)-1H-imidazol-4-yl)-quinoxaline, ALK5 Inhibitor VIII
Tgf-beta-R1 inhibitor	TGF- β RI Kinase Inhibitor IX	1117684-36-2	616463, TGF- β RI Kinase Inhibitor IX, 4-((4-((2,6-Dimethylpyridin-3-yl)oxy)pyridin-2-yl)amino)benzenesulfonamide, ALK5 Inhibitor IX
Tgf-beta-R1 inhibitor	GW788388	452342-67-5	4-(4-(3-(pyridin-2-yl)-1H-pyrazol-4-yl)pyridin-2-yl)-N-(tetrahydro-2H-pyran-4-yl)benzamide

Class	Agent	CAS Number	Alternative Name
Tgf-beta-R1 inhibitor	LY2109761	700874-71-1	7-(2-morpholinoethoxy)-4-(2-(pyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)quinoline
Tgf-beta-R1 inhibitor	Galunisertib (LY2157299)	700874-72-2	4-(2-(6-methylpyridin-2-yl)-5,6-dihydro-4H-pyrrolo[1,2-b]pyrazol-3-yl)quinoline-6-carboxamide
Tgf-beta-R1 inhibitor	EW-7197	1352608-82-2	N-(2-fluorophenyl)-5-(6-methyl-2-pyridinyl)-4-[1,2,4]triazolo[1,5-a]pyridin-6-yl-1H-imidazole-2-methanamine
Tgfb production inhibitor	Pirfenidone	53179-13-8	5-methyl-1-phenyl-2(1H)-Pyridinone,
Tgf-beta-R1 inhibitor	K02288	1431985-92-0	3-[(6-Amino-5-(3,4,5-trimethoxyphenyl)-3-pyridinyl]phenol
Tgf-beta-R1 inhibitor	D 4476	301836-43-1	4-[4-(2,3-Dihydro-1,4-benzodioxin-6-yl)-5-(2-pyridinyl)-1H-imidazol-2-yl]benzamide
Tgf-beta-R1 inhibitor	R 268712	879487-87-3	4-[2-Fluoro-5-[3-(6-methyl-2-pyridinyl)-1H-pyrazol-4-yl]phenyl]-1H-pyrazole-1-ethanol
Other	ITD 1	1099644-42-4	4-[1,1'-Biphenyl]-4-yl-1,4,5,6,7,8-hexahydro-2,7,7-trimethyl-5-oxo-3-quinolinecarboxylic acid ethyl ester
Smad3 inhibitor	SIS3	1009104-85-1	1,2,3,4-Tetrahydro-6,7-dimethoxy-2-[(2E)-3-(1-phenyl-1H-pyrrolo[2,3-b]pyridin-3-yl)-1-oxo-2-propenyl]-isoquinoline hydrochloride
Tgf-beta-R1 inhibitor	A77-01	909910-42-5	4-[5-(6-methylpyridin-2-yl)-1H-pyrazol-4-yl]quinoline
Tgf-beta-R1 inhibitor	SM16	614749-78-9	4-(5-(benzo[d][1,3]dioxol-5-yl)-4-(6-methylpyridin-2-yl)-1H-imidazol-2-yl)bicyclo[2.2.2]octane-1-carboxamide
Tgf-beta-R1 inhibitor	LY-550410	737791-20-7	5,6-dihydro-2-(2-pyridinyl)-4H-pyrrolo[1,2-b]pyrazol-3-yl]- Quinoline
Tgf-beta-R1 inhibitor	LY-580276	476475-07-7	3-(4-fluorophenyl)-5,6-dihydro-2-(6-methyl-2-pyridinyl)-4H-Pyrrolo[1,2-b]pyrazole
Tgf-beta-R1 inhibitor	EW-7203	1383123-98-5	3-[[[4-(6-methyl-2-pyridinyl)-5-[1,2,4]triazolo[1,5-a]pyridin-6-yl-2-thiazolyl]amino]methyl]- Benzonitrile,
Tgf-beta-R1 inhibitor	EW-7195	1352609-28-9	3-[[[5-(6-methyl-2-pyridinyl)-4-[1,2,4]triazolo[1,5-a]pyridin-6-yl-1H-imidazol-2-yl]methyl]amino]- Benzonitrile
Tgf-beta-R1 inhibitor	GW6604	452342-37-9	Pyridine, 2-phenyl-4-[3-(2-pyridinyl)-1H-pyrazol-4-yl]-
Tgf-beta-R1 inhibitor	Cmpd 3d	733806-89-8	4-Quinazolinamine, 2-(6-methyl-2-pyridinyl)-N-4-pyridinyl-

Class	Agent	CAS Number	Alternative Name
Tgf-beta-R1 inhibitor	LY-566578	607738-00-1	Pyridine, 2-[4-(4-fluorophenyl)-1H-pyrazol-3-yl]-6-methyl-
Tgf-beta-R1 inhibitor	Cmpd 5	607738-02-3	Phenol, 4-[3-(6-methyl-2-pyridinyl)-1H-pyrazol-4-yl]
Tgf-beta-R1 inhibitor	Cmpd 3	676331-30-9	Quinoline, 7-ethoxy-4-[3-(2-pyridinyl)-1H-pyrazol-4-yl]-
Tgf-beta-R1 inhibitor	Cmpd 8b	705263-50-9	1H-Benzimidazole, 6-[5,6-dihydro-2-(2-pyridinyl)-4H-pyrrolo[1,2-b]pyrazol-3-yl]-
Tgf-beta-R1 inhibitor	Cmpd 4b	1308760-90-8	N-(3-cyanophenyl)-3-(6-methyl-2-pyridinyl)-4-(6-quinolinyl)-1H-Pyrazole-1-acetamide
Tgf-beta-R1 inhibitor	Cmpd 21b	1607465-38-2?	1H-Pyrazole-1-acetamide, N-(3-cyanophenyl)-3-(6-methyl-2-pyridinyl)-4-[1,2,4]triazolo[1,5-a]pyridin-6-yl
Tgf-beta-R1 inhibitor	PF-03671148	1378524-25-4	3-methyl-6-[1-(6-methyl-2-pyridinyl)-1H-pyrazol-5-yl]-4(3H)-Quinazolinone,
Tgf-beta-R1 inhibitor	SB-203580	152121-47-6	Pyridine, 4-[4-(4-fluorophenyl)-2-[4-(methylsulfinyl)phenyl]-1H-imidazol-5-yl]-
Tgf-beta-R1 inhibitor	SB-202190	152121-30-7	4-[4-(4-Fluorophenyl)-5-(4-pyridinyl)-1H-imidazol-2-yl]phenol
Tgf-beta-R1 inhibitor	IN-1130	868612-83-3	3-[[5-(6-methyl-2-pyridinyl)-4-(6-quinoxaliny)-1H-imidazol-2-yl]methyl]-Benzamide,
Tgf-beta-R1 inhibitor	IN-1233	1093952-95-4	3-[[5-(6-methyl-2-pyridinyl)-4-(6-quinolinyl)-1H-imidazol-2-yl]methyl]-Benzamide,
Tgf-beta-R1 inhibitor	Cmpd 16i	864375-44-0	[[4-(6-benzothiazolyl)-5-(4-methyl-2-thiazolyl)-1H-imidazol-2-yl]methyl]-2-methylpropyl ester Carbamic acid
Tgf-beta-R1 inhibitor	LDN-214117	1627503-67-6	1-[4-[6-methyl-5-(3,4,5-trimethoxyphenyl)-3-pyridinyl]phenyl]-Piperazine
Tgf-beta-R1 inhibitor	LDN-193189	1627503-67-6	CAS 1062368-24-4, 4-[6-[4-(1-piperazinyl)phenyl]pyrazolo[1,5-a]pyrimidin-3-yl]-Quinoline
Tgf-beta-R1 inhibitor	Cmpd 12b	1415663-82-9	2-N-[(3-fluorophenyl)methyl]-4-(6-methyl-2-pyridinyl)-5-[1,2,4]triazolo[1,5-a]pyridin-6-yl Thiazolamine
Tgf-beta-R1 inhibitor	Cmpd 6d	1630024-29-1	5-[[2-cyclopropyl-6-(4-fluorophenyl)imidazo[2,1-b]-1,3,4-thiadiazol-5-yl]methylene]-4-oxo-2-thioxo-3-Thiazolidineacetic acid
Tgf-beta-R1 inhibitor	SD-093		Structure unknown
Tgf-beta-R1 inhibitor	Ki-26894		Structure unknown
Tgf-beta-R1 inhibitor	NPC-30345		Structure unknown
Tgf-beta-R1 inhibitor	SX-007		Structure unknown

Class	Agent	CAS Number	Alternative Name
Tgf-beta-R1 inhibitor	SKI-2162		Structure unknown
Other	Asiaticoside	16830-15-2	
Tgf-beta antibody	ID11		
Tgf-beta antibody	2G7		
Tgf-beta antibody	GC-1008		Fresolimumab
Tgf-beta antibody	CAT-152		Lerdelimumab
Tgf-beta antibody	CAT-192		Metelimumab
TGF-beta Receptor antibody	PF-03446962		
Tgf-beta antibody	SR-2F		
Tgf-beta antibody	2G7		
Tgf-beta antibody	LY2382770		
Tgf-beta antibody	IMC-TR1		
Tgf-beta antibody	STX-100		
TGF-beta antagonist	TGF-PRII:Fc		
Recombinant protein	betaglycan/TGF-PRIII		
Oligonucleotide inhibitor	AP12009		Trabedersen, antisense molecule
Oligonucleotide inhibitor	AP11014		
Oligonucleotide inhibitor	AP15012		
Is this TGF b inhibitor/YES	LY-573636	519055-62-0	N-[(5-bromo-2-thienyl)sulfonyl]-2,4-dichloro-Benzamide
	pyrrole-imidazole polyamide		Gene silencing
	U.S. Patent No. 7,087,626		Pyrrole derivatives as pharmaceutical agents
	U.S. Patent No. 6,476,031		Quinazoline derivatives as medicaments
	U.S. Patent No. 7,723,486, and EP 0945464		Antibodies to TGF- β
Peptide	Tryptopeptin A	1644153-72-9	
Peptide	Trx-xFoxHlb		Smad-interacting peptide aptamers
Peptide	Trx-Lefl		
Peptide	Distertide (p144)		
Peptide	p17		
Peptide	LSKL		

Class	Agent	CAS Number	Alternative Name
dihydropyrrlipyrazole-based scaffold	See US Patent US 8298825 B1		
imidazole-based scaffold	See US Patent US 8298825 B1		
pyrazolopyridine-based scaffold	See US Patent US 8298825 B1		
pyrazole-based scaffold			See US Patent US 8298825 B1
imidazopyridine-based scaffold	See US Patent US 8298825 B1		
triazole-based scaffold			See US Patent US 8298825 B1
pyridopyrimidine-based scaffold	See US Patent US 8298825 B1		
pyrrolopyrazole-based scaffold	See US Patent US 8298825 B1		
isothiazole-based scaffold	See US Patent US 8298825 B1		
oxazole-based scaffold	See US Patent US 8298825 B1		

Compositions with Poloxamers

[0226] In certain embodiments, the present disclosure provides a pharmaceutical composition comprising: a) a compound of the present disclosure and b) a poloxamer.

[0227] In certain embodiments, the pH of the pharmaceutical composition is between about 5 and about 9. In certain embodiments, the pH of the pharmaceutical composition is about 5, 6, 7, 8, or 9.

[0228] In certain embodiments, the solubility of the compound in the presence of the poloxamer is about 3-fold higher than the solubility of the compound at the same pH in the absence of poloxamer. In certain embodiments, the solubility of the compound in the presence of the poloxamer is about 2-, 3-, 4- or 5-fold higher than the solubility of the compound at the same pH in the absence of poloxamer.

[0229] In certain embodiments, the pharmaceutical formulations may also contain a poloxamer. Poloxamers are nonionic triblock copolymers composed of a central hydrophobic chain of polyoxypropylene (poly(propylene oxide)) flanked by two hydrophilic chains of polyoxyethylene (poly(ethylene oxide)). Poloxamers are often considered “functional excipients” because they are essential components and play an important role in a formulation.

[0230] In some embodiments, the poloxamer comprises at least one of Poloxamer 124, Poloxamer 188, Poloxamer 237, Poloxamer 338 or Poloxamer 407. In some embodiments, the poloxamer comprises mixtures of two or more of Poloxamer 124, Poloxamer 188, Poloxamer 237, Poloxamer 338 or Poloxamer 407. In some embodiments, the mixture of two or more poloxamers comprise Poloxamer 407 and Poloxamer 124. In another embodiment the Poloxamer comprises at least one of Poloxamer 188 and Poloxamer 407 or mixtures thereof. In some embodiments, the poloxamer is Poloxamer 407.

[0231] In some embodiments, the poloxamer is in a concentration between about 5 wt% and about 25 wt% relative to the composition. In some embodiments, the poloxamer is in a concentration between about 10 wt% and about 23 wt% relative to the composition. In some embodiments, the poloxamer is in a concentration between about 15 wt% and about 20 wt% relative to the composition. In some embodiments, the poloxamer is in a concentration is approximately 17 wt% relative to the composition. In some embodiments, the poloxamer is in a concentration is approximately 21 wt% relative to the composition.

[0232] In some embodiments, the poloxamer can be in a concentration between 21wt % and 40 wt % relative to the composition. In another embodiment the poloxamer is in a concentration between 21wt % and 30 wt % relative to the composition. In another embodiment the poloxamer is in a concentration between 23 wt % and 29 wt % relative to the composition. In another embodiment the poloxamer is in a concentration between 23 wt % and 27 wt % relative to the composition. In another embodiment the poloxamer is in a concentration of 25 wt % relative to the composition.

[0233] In some embodiments, the gelation temperature of the pharmaceutical composition is greater than about 10°C. In some embodiments, the gelation temperature of the pharmaceutical composition is between about 11°C and about 32°C. In some embodiments, the gelation temperature of the pharmaceutical composition is between about 15°C and about 30 °C. In some embodiments, the gelation temperature of the pharmaceutical composition is between about 20°C and about 28°C. In some embodiments, the gelation temperature of the pharmaceutical composition is between about 24°C and about 26°C.

[0234] In some embodiments, the gelation temperature of the pharmaceutical composition is about 15°C. In some embodiments, the gelation temperature of the pharmaceutical composition is about 20°C. In some embodiments, the gelation temperature of the pharmaceutical

composition is about 24°C. In some embodiments, the gelation temperature of the pharmaceutical composition is about 26°C. In some embodiments, the gelation temperature of the pharmaceutical composition is about 28°C. In some embodiments, the gelation temperature of the pharmaceutical composition is about 30°C. In some embodiments, the gelation temperature of the pharmaceutical composition is about 32°C.

[0235] In certain embodiments, the present disclosure provides a pharmaceutical composition having a gelation temperature greater than 10°C, the composition comprising: a) a pharmaceutically active compound of the present disclosure; b) a poloxamer at greater than or equal to 21 wt % of the pharmaceutical composition; and c) a HDAC inhibitor; wherein the pharmaceutical composition has a gelation temperature of greater than 10°C.

[0236] In certain embodiments, the present disclosure provides a pharmaceutical composition having a gelation temperature greater than 10°C, the composition comprising: a) a pharmaceutically active compound of the present disclosure; b) a poloxamer at greater than or equal to 21 wt % of the pharmaceutical composition; and c) a carboxylic acid containing compound; wherein the pharmaceutical composition has a gelation temperature of greater than 10°C.

EXAMPLES


General experimental methods

[0237] ¹H NMR spectra were recorded on Bruker Avance III 400 MHz and Bruker Fourier 300 MHz and TMS was used as an internal standard.


[0238] LCMS was taken on a quadrupole Mass Spectrometer on Agilent LC/MSD 1200 Series (Column: C18 (50 × 4.6 mm, 5 μm) operating in ES (+) or (-) ionization mode; T = 30 °C

Synthetic Schemes


Synthetic Scheme 1




Synthetic Scheme 2




Synthetic Scheme 3




Synthetic Scheme 4




Synthetic Scheme 5




Synthetic Scheme 6




Synthetic Scheme 7




Synthetic Scheme 8




Synthetic Scheme 9




Synthetic Scheme 10




Synthetic Scheme 11




Synthetic Scheme 12




Synthetic Scheme 13




Synthetic Scheme 14




Synthetic Scheme 15




Experimental Procedures:**Synthesis of intermediate 2.**

[0239] To a solution of intermediate 1 (20 g, 213 mmol) in MeCN (540 ml) was ethyl (E)-4-oxo-butenoate (28.6 g, 223 mmol). The reaction mixture was heated to 80 °C and stirred for 6 hrs. The reaction mixture was concentrated under reduced pressure, the residue was purified by flash column chromatography (eluted with Dichloromethane/MeOH from 1:0 to 200:1) to give the crude intermediate 2 (25 g) as brown solid.

Synthesis of intermediate 3.

[0240] To a solution of crude intermediate 2 (25 g) in MeOH (100 ml) was added NH_3/MeOH (6 M, 100 ml). The reaction mixture was stirred at room temperature overnight. The mixture was poured into EtOAc (500 ml), and then filtered. The filter cake was dried in vacuo to give intermediate 3 (13 g, 35% for two steps) as a brown solid.


[0241] $^1\text{H NMR}$ ($\text{DMSO}-d_6$, 400 MHz): δ (ppm) 8.30 (d, $J=6.8\text{Hz}$, 1H), 7.60 (s br, 1H), 7.54 (d, $J=5.2\text{Hz}$, 1H), 7.41 (s, 1H), 7.19-7.23 (m, 1H), 7.06 (s br, 1H), 6.89-6.93 (m, 1H), 3.80 (s, 2H).

Synthesis of intermediate 5.

[0242] To a solution of intermediate **4** (100 g, 0.51 mol) in DMF (1000 ml) was added NaH (60%, 61 g, 1.53 mol) at 0 °C. The mixture was stirred at room temperature for 20 mins. 2-chloroethylamine hydrochloride (89.2 g, 0.77 mol) was added to the mixture in portions at 0 °C. The mixture was stirred at room temperature for 2 hrs. TLC (Petroleum Ether/EtOAc=5/1) showed the reaction was complete. The mixture was poured into ice-water and extracted with EtOAc (600 ml × 3). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum to give intermediate **5** (110 g, 90 %) as yellow oil.

[0243] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.74-7.75 (d, 1H, J=1.2 Hz), 7.21-7.29 (m, 2H), 7.13-7.14 (d, 1H, J=3.2Hz), 6.44-6.45 (d, 1H, J=2.8Hz), 4.14-4.17 (t, 2H, J=6 Hz), 3.08-3.11 (t, 2H, J=6 Hz).


Synthesis of intermediate **6**.



[0244] To a solution of intermediate **5** (150 g, 0.63 mol) in AcOH (720 ml) was added Borane pyridine complex (9.3 M, 135.5 ml, 1.26 mol) at room temperature under N₂. The mixture was stirred at room temperature for overnight. Then the mixture was adjusted pH = 9-10 with aqueous NaOH, extracted with EtOAc (800 ml × 3). The combined organic phases were concentrated in vacuum to give crude compound. Water (720 ml) was added to the crude compound, followed by the slow addition of concentrated HCl (240 ml). The mixture was stirred at room temperature for 30 mins, adjusted pH = 10-11 with aqueous NaOH, extracted with EtOAc (800 ml × 3), concentrated to give crude compound. To a solution of the crude compound in methyl tertiary butyl ether (500 ml) was added AcOH (28 ml) at room temperature. The mixture was stirred at room temperature for 30 mins, then filtered, the filter cake was washed with methyl tertiary butyl ether, dried to give intermediate **6** (120 g, 63.5 %) as white solid.


[0245] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 7.14 (s, 1H), 7.09-7.12 (d, 1H, J=8.4 Hz), 6.43-6.45 (d, 1H, J=8.4 Hz), 3.33-3.37 (t, 2H, J=8.4 Hz), 3.06-3.09 (t, 2H, J=6.6 Hz), 2.87-2.92 (t, 2H, J=8.4 Hz), 2.76-2.79 (t, 2H, J=6.6 Hz).

Synthesis of intermediate **7**.



[0246] To a solution of H_2SO_4 (12.6 ml) in AcOH (80 ml) and HCHO (37% aqueous, 660 ml) was added intermediate **6** (100 g, 0.33 mol) in portions at room temperature. The mixture was stirred at 50 °C for 20 mins. Then the mixture was adjusted pH = 9-10 with aqueous NaOH, extracted with EtOAc (800 ml \times 3), concentrated to give crude intermediate **7** (100 g) as yellow solid, which was used directly for the next step without purification.

Synthesis of intermediate **8**.



[0247] A mixture of intermediate **7** (100 g, crude) and aqueous K_2CO_3 (300 ml, 1 M) in THF (700 ml) was added $(\text{Boc})_2\text{O}$ (94.4 g) at room temperature. The mixture was stirred at room temperature for overnight. TLC (dichloromethane/MeOH=10/1) showed the reaction was complete. Then H_2O was added, extracted with EtOAc (500 ml \times 3). The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 50:1 to 5:1) to give intermediate **8** (75 g, 64.6% for two steps) as yellow solid.


[0248] ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) 7.11 (bs, 1H), 6.99 (bs, 1H), 4.30-4.37 (m, 2H), 3.68 (m, 2H), 3.36-3.40 (m, 2H), 2.96-3.01 (m, 4H), 1.41 (s, 9H).

Synthesis of core **1**.

The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 20:1 to 1:1) to give intermediate **10** (7.5 g, 72.8%) as yellow solid.

[0253] $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ (ppm) 7.86 (d, $J=1.6\text{Hz}$, 1H), 7.26 (s, 1H), 7.02 (t, $J=1.6\text{Hz}$, 1H), 6.47 (d, $J=3.2\text{Hz}$, 1H), 4.63 (s, 2H), 4.22-4.20 (t, $J=4.8\text{Hz}$, 2H), 3.98-4.00 (t, $J=4.8\text{Hz}$, 2H), 3.18-3.19 (m, 4H), 1.58-1.40 (m, 6H).


Synthesis of intermediate 11.



[0254] To a solution of intermediate **10** (10 g, 27.6 mmol) in dichloromethane (100 ml) was added $(\text{COCl})_2$ (8.8 g, 69 mmol) at 0°C under N_2 . The mixture was stirred at 40°C for 1 hr. TLC (petroleum ether/EtOAc=1/1) showed the reaction was complete. Then a solution of NaOMe (3.7 g, 69 mmol) in MeOH (10 ml) was added at -60°C under N_2 . The mixture was stirred at room temperature for 1 hr. Water was added, extracted with dichloromethane (100 ml \times 3). The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 5:1 to 1:2) to give intermediate **11** (6 g, 48.5%) as yellow solid.

[0255] $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ (ppm) 8.70 (d, $J=1.6\text{Hz}$, 1H), 8.29 (s, 1H), 7.41 (s, 1H), 4.67 (s, 2H), 4.41-4.39 (m, 2H), 4.01-3.98 (m, 2H), 3.95 (s, 3H), 3.15-3.01 (m, 4H), 1.60-1.40 (m, 6H).

Synthesis of core 2.




[0256] A solution of intermediate **11** (10 g, 22.3 mmol) and intermediate **3** (3.9 g, 22.3 mmol) in DMF (180 ml) was added a solution of t-BuOK (6.4 g, 19.0 mmol) in THF (100 ml) at 0-10 °C. The mixture was stirred at 0-10 °C for 15 mins. TLC (dichloromethane/MeOH=15/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc/THF from 10:5:1 to 1:1:1) to give **core 2** (6.5 g, 50.7%) as orange solid.

[0257] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 11.25 (s, 1H), 8.02 (s, 1H), 7.91 (s, 1H), 7.65-7.68 (d, J = 6.8Hz, 1H), 7.59-7.61 (d, J = 6.8Hz, 1H), 7.19-7.23 (t, J = 7.6 Hz, 1H), 7.04 (s, 1H), 6.55-6.58 (t, J = 6.4 Hz, 1H), 6.08 (s, 1H), 4.63 (s, 2H), 4.50-4.62 (m, 2H), 3.82-3.86 (m, 2H), 2.94-3.06 (m, 4H), 1.40-1.60 (m, 6H).

[0258] LC/MS M+1=573.1


Synthesis of intermediate **12**.



[0259] To a solution of **core 1** (10 g, 28.5 mmol) in DMF (200 ml) was added CuI (5.4 g, 28.5 mmol) and methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (19.2 g, 100 mmol) at room temperature under N₂. The mixture is stirred at 80 °C for 2.5 hrs. TLC (petroleum ether/EtOAc=5/1) showed that the reaction was complete. The mixture was cooled to room temperature, and filtered. The filtrate was added water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 20:1 to 5:1) to give intermediate **12** (5.5 g, 56.7%) as yellow solid.

[0260] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.81 (s, 1H), 7.13-7.28 (m, 2H), 6.62 (s, 1H), 4.92-4.84 (m, 2H), 4.22 (s, 2H), 3.98-3.97 (m, 2H), 1.40 (s, 9H).


Synthesis of intermediate **13**.



[0261] To a solution of intermediate **12** (5.0 g, 14.7 mmol) in dichloromethane (75 ml) was added (COCl)₂ (4.6 g, 36.7 mmol) under N₂. The mixture was stirred at 40 °C for 1 hr. TLC (petroleum ether/EtOAc=1/1) showed the reaction was complete. Then a solution of NaOMe (1.98 g, 36.7 mmol) in MeOH (10 ml) was added at -60 °C under N₂. The mixture was stirred at room temperature for 1 hr. Water was added, extracted with dichloromethane (100 ml ×3). The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 10:1 to 5:1) to give intermediate **13** (3.8g, 60.7%) as a yellow solid.

[0262] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 8.64 (s, 1H), 8.44 (s, 1H), 7.40-7.31 (m, 1H), 4.98-4.88 (m, 2H), 4.51-4.50 (m, 2H), 4.04-4.01 (m, 2H), 3.96 (s, 3H), 1.40 (s, 9H).

Synthesis of intermediate **14**.




[0263] To a solution of intermediate **13** (5.5 g, 12.9 mmol) and intermediate **3** (2.25 g, 12.9 mmol) in DMF (110 ml) was added a solution of tBuOK (3.6 g, 32.2 mmol) in THF (10 ml) at 0-10 °C. The mixture was stirred at 0-10 °C for 15 min. TLC (petroleum ether/EtOAc=1/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography

(eluted with dichloromethane/MeOH from 100:1 to 30:1) to give intermediate **14** (3.5 g, 58.3%) as orange solid.

[0264] $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ (ppm) 8.16 (s, 1H), 8.02 (s, 1H), 7.96 (s, 1H), 7.35 (s, 1H), 7.27 (s, 1H), 7.12-7.08 (m, 2H), 6.40-6.38 (m, 2H), 4.89-4.77 (m, 2H), 4.50 (s, 2H), 4.04 (s, 2H), 1.48-1.37 (m, 9H).

Synthesis of compound I-1.




[0265] To a solution of intermediate **14** (5 g, 9.1 mmol) in dichloromethane (50 ml) was added HCl/dioxane (50 ml, 7M) at room temperature. The mixture was stirred at room temperature for 1 hr. TLC (dichloromethane/MeOH=15/1) showed the reaction was complete. The solvent was concentrated in vacuum to give white solid.

[0266] To a solution of the white solid and 1-Piperidinecarbonyl chloride (1.8 g, 12.3 mmol) in DMF (40 ml) was added Et_3N (2.49 g, 24.6 mmol) at room temperature. The mixture was stirred at room temperature for 1 hr. TLC (dichloromethane/MeOH =10/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/THF from 20:1 to 10:1) to give **compound I-1** (3.0 g, 70%) as red solid.

[0267] $^1\text{H NMR}$ ($\text{DMSO}-d_6$, 400 MHz): δ (ppm) 11.30 (s, 1H), 8.17 (s, 1H), 7.93 (s, 1H), 7.64-7.60 (m, 2H), 7.19 (s, 1H), 6.56 (m, 1H), 6.31 (s, 1H), 4.77-4.69 (m, 2H), 4.68-4.60 (m, 2H), 3.95-3.85 (m, 2H), 3.10-2.90 (m, 4H), 1.55-1.35 (m, 6H).

[0268] LC/MS M+1 563.1

Synthesis of compound I-2.




[0269] A solution of **core 2** (5 g, 8.7 mmol) in N-Methyl-2-Pyrrolidone (50 ml) was added CuCN (2.5 g, 28 mol) at room temperature. The mixture was stirred at 150 °C for 6 hr under N₂. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/MeOH from 1:0 to 50:1) to give **compound I-2** (2.2 g, 49%) as orange solid.

[0270] ¹H NMR (DMSO-*d*₆): δ (ppm) 11.30 (s, 1H), 8.07 (s, 1H), 7.97 (s, 1H), 7.66-7.68 (d, 1H, J = 9.2 Hz), 7.54-7.56 (d, 1H, J = 6.8 Hz), 7.29 (s, 1H), 7.19-7.23 (t, J = 8 Hz, 1H), 6.53-6.57 (t, 1H, J = 6.8 Hz), 6.44 (s, 1H), 4.68 (s, 4H), 4.60-4.63 (m, 4H), 3.84-3.88 (m, 2H), 2.96-3.05 (m, 4H), 1.43-1.47 (m, 6H).


[0271] LC/MS M+1 520.1

Synthesis of intermediate 15.



[0272] To a solution of **core 1** (50 g, 0.14 mol) in 1,4-dioxane (500 ml) was added NaI (42.7 g, 0.28 mol) and CuI (2.7g) and 2-Dimethylaminoethylamine (2.5g). The mixture was stirred at 140 °C overnight under N₂. The mixture was filtered, the filter cake was washed with dichloromethane. The combined organic phases were concentrated in vacuum. The residue was washed with methyl tertiary butyl ether to give intermediate **15** (50 g, 88.0%) as yellow solid.


Synthesis of intermediate 16.



[0273] To a solution of intermediate **15** (10 g, 25.1 mmol) in Et₃N (120 ml) and THF (60 ml) at room temperature, then Pd(PPh₃)₂Cl₂ (1.2 g, 2.4 mmol) and CuI (1.2 g, 2.4 mmol) was added under N₂. Then ethynyltrimethylsilane (4.75 g, 48.4 mmol) was added drop wise. The mixture was stirred at 60 °C overnight. The mixture was concentrated in vacuum to remove Et₃N. Then the mixture was poured into ice-water, extracted with CH₂Cl₂ (300 ml × 2). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 100:1 to 50:1) to give intermediate **16** (6 g, 64.8 %) as yellow solid.

[0274] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.63-7.67 (m, 1H), 7.04-7.41 (m, 2H), 6.46-6.49 (m, 1H), 4.77-4.84 (m, 2H), 4.23-4.29 (m, 2H), 3.90-3.97 (m, 2H), 1.43 (s, 9H), 0.26 (s, 9H).


Synthesis of intermediate 17.



[0275] A solution of intermediate **16** (10 g, 27 mmol) in dichloromethane (110 ml) was added (COCl)₂ (8.5 g, 67.4 mmol) at 0 °C under N₂. The mixture was stirred at 40 °C for 1 hr. TLC (petroleum ether/EtOAc=5/1) showed the reaction was complete. Then a solution of NaOMe (3.64 g, 67.4 mmol) in MeOH (10 ml) was added at -60 °C under N₂. The mixture was stirred at room temperature for 1 hr. Water was added, extracted with dichloromethane (100 ml × 3). The


combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 100:1 to 5:1) to give intermediate **17** (7 g, 56.7%) as yellow solid.

Synthesis of intermediate **18**.



[0276] To a solution of intermediate **17** (8 g, 17.6 mmol) and intermediate **3** (3.1 g, 17.6 mmol) in DMF (80 ml) was added a solution of tBuOK (4.9 g, 44 mmol) in THF (30 ml) at 0-10 °C. The mixture was stirred at 0-10 °C for 15 min. TLC (petroleum ether/EtOAc=1/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/MeOH from 100:1 to 50:1) to give intermediate **18** (5 g, 49%) as orange solid.

Synthesis of intermediate **19**.




[0277] To a solution of intermediate **18** (5 g, 8.6 mmol) in dichloromethane (50 ml) was added HCl/dioxane (50 ml, 7M) at room temperature. The mixture was stirred at room temperature for

1 hr. TLC (dichloromethane/MeOH=15/1) showed the reaction was complete. The solvent was concentrated in vacuum to give white solid.

[0278] To a solution of the white solid and 1-Piperidinecarbonyl chloride (1.4 g, 9.5mmol) in DMF (50 ml) was added Et₃N (2.6 g, 25.8 mmol) at room temperature. The mixture was stirred at room temperature for 1 hr. TLC (dichloromethane/MeOH =10/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum to give the crude intermediate **19** (5.2 g), which was used directly for the next step without purification.

Synthesis of compound I-3.




[0279] To a solution of the crude intermediate **19** (5.2 g) in DMF (100ml) was added K₂CO₃ (2 g). The mixture was heated to 50 °C and stirred for 1 hr. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/MeOH from 100:1 to 30:1) to give **compound I-3** (3 g, 18.6% for two steps) as red solid.

[0280] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.26 (s, 1H), 8.02 (s, 1H), 7.92 (s, 1H), 7.65-7.59 (m, 2H), 7.20 (t, 1H, *J* = 8Hz), 6.97 (s, 1H), 6.58-6.55 (t, *J* = 6.8Hz, 1H), 6.18 (s, 1H), 4.63 (s, 2H), 4.50-4.60 (m, 2H), 3.80-3.90 (m, 2H), 3.75 (s, 1H), 2.95-3.05 (m, 4H), 1.46-1.35 (m, 6H).

[0281] LC/MS M+1 519.2

Synthesis of compound I-10




[0282] A solution of **core 2** (5 g, 8.8 mmol) in dioxane (50 ml) was added acetamide (3.1 g, 53.3 mmol), CuI (1.1 g, 5.8 mmol), K_3PO_3 (5.5 g, 26.4 mmol) at RT under N_2 . The mixture was stirred at room temperature for 20 mins. N^1,N^2 -dimethylethane-1,2-diamine (1.56 g, 17.8 mmol) was added, and then the mixture was stirred at 115 °C for 5 hrs under N_2 . TLC (dichloromethane/MeOH=15/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc/THF (3/1, 100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/MeOH from 200:1 to 50:1) to give **compound I-10** (3 g, 62%) as red solid.

[0283] 1H NMR (DMSO- d_6 , 400 MHz): δ (ppm) 11.21 (s, 1H), 9.43 (s, 1H), 7.86 (s, 1H), 7.78 (s, 1H), 7.58-7.64 (m, 2H), 7.17-7.21 (t, 1H, $J = 8.0$ Hz), 7.00 (s, 1H), 6.77 (s, 1H), 6.56-6.58 (t, 1H, $J = 7.2$ Hz), 4.60 (s, 2H), 4.40-4.50 (m, 2H), 3.80-3.88 (m, 2H), 2.88-3.12 (m, 4H), 1.85 (s, 3H), 1.15-1.20 (m, 6H).

[0284] LC/MS $M+1$ 552.2

Synthesis of compound I-4.




[0285] A solution of compound I-10 (3 g, 5.4 mmol) in EtOH (10 ml) was added HCl (28 ml, 6 N) at room temperature. The mixture was stirred at 80 °C for 3 hrs. TLC (dichloromethane/MeOH =15/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with dichloromethane (100 ml ×2), then the water phase was adjusted PH =9-10 with aq. Na₂CO₃, extracted with dichloromethane (100 ml ×6), concentrated in vacuum. The residue was washed with methyl tertiary butyl ether, filtered. The filter cake was dried in vacuo to give **compound I-4** (2.1 g, 75 %) as red solid.

[0286] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.19 (s, 1H), 7.86 (s, 1H), 7.84 (s, 1H), 7.62 (bs, 2H), 7.18-7.22 (t, 1H, *J* = 7.2 Hz), 6.57-6.61 (t, 1H, *J* = 6.8 Hz), 6.24 (s, 1H), 5.27 (s, 1H), 4.50 (s, 2H), 4.36-4.39 (m, 2H), 4.14-4.17 (m, 2H), 3.75-3.78 (m, 2H), 2.88-3.05 (m, 4H), 1.44-1.48 (m, 6H).

[0287] MS/LC M+1 511.1


Synthesis of intermediate 22.



[0288] To a solution of intermediate **21** (100 g, 0.59 mol) in dry toluene (890 ml) was added n-BuOH (131.6 g, 1.78 mol) and TsOH (10 g) at room temperature. The mixture was stirred at 120 °C for overnight, and removed the water using a Dean-stark apparatus. TLC (petroleum ether/EtOAc=5/1) showed the reaction was completed. The mixture was concentrated in vacuum to give crude intermediate **22**. The crude intermediate **22** was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 100:1 to 20:1) to give intermediate **22** (120 g, 67.8%) as a yellow oil.

[0289] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.89-7.92 (dd, 1H, *J*=4.8Hz, 8.8 Hz), 7.51-7.54 (dd, 1H, *J*=2.8Hz, 9.6Hz), 7.09-7.14 (m, 1H), 6.04 (s, 1H), 3.50-3.56 (m, 4H), 1.55-1.62 (m, 4H), 1.33-1.42 (m, 4H), 0.83-0.93 (m, 6H).


Synthesis of intermediate 23.



[0290] To a solution of intermediate **22** (50 g, 0.17 mol) in dry THF (1500 ml) was added vinylmagnesium bromide solution (1 M, 668.8 ml, 668.8 mmol) dropwise at -40°C . The mixture was stirred at -40°C for 1 hr. TLC (petroleum ether/EtOAc = 5/1) showed the reaction was completed. Then the mixture was poured into aq. NH_4Cl , extracted with EtOAc (300 ml \times 3), the organic phases were concentrated to give crude compound. The crude compound was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 100:1 to 20:1) to give the compound (24 g) as yellow oil. To a solution of the compound (24 g) in THF (100 ml) was added HCl (0.5 N, 80 ml) drop wise at room temperature. The mixture was stirred at room temperature for 1 hr. TLC (petroleum ether/EtOAc = 5/1) showed the reaction was completed. The mixture was adjusted PH=10 with aq. NaOH, extracted with EtOAc (300 ml \times 3), concentrated in vacuum to give intermediate **23** (16 g, 58.8 %) as yellow solid.


[0291] $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ (ppm) 10.06 (s, 1H), 7.62 (dd, 1H, $J=2\text{Hz}$, 9.2Hz), 7.38-7.41 (m, 2H), 6.60(t, 1H, $J=2.4\text{Hz}$).

Synthesis of intermediate 24.



[0292] To a solution of intermediate **23** (140 g, 0.86 mol) in MeOH (2100 ml) was added 2-aminoethanol (78 g, 1.3 mol) and Pd/C (14 g) at room temperature under N_2 . The mixture was stirred at room temperature for 2 hrs under N_2 . Then the mixture was stirred at room temperature for overnight under H_2 . The mixture was filtered, concentrated to give intermediate **24** (200 g, crude) as a yellow oil, which was used directly for the next step without purification.


Synthesis of intermediate 25.



[0293] A mixture of intermediate **24** (90 g, crude) and K_2CO_3 (467 ml, 1 M) in THF (1300 ml) was added Boc_2O (141 g) at room temperature. The mixture was stirred at room temperature for overnight. TLC (dichloromethane/MeOH=10/1) showed the reaction was completed. Then H_2O was added, extracted with EtOAc (500 ml \times 3). The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 10:1 to 1:1) to give intermediate **25** (66.6 g, 56% for two steps) as yellow oil.

[0294] $^1\text{H NMR}$ (CDCl_3 , 400 MHz): δ (ppm) 10.17 (s, 1H), 7.23-7.26 (m, 2H), 6.78-6.82 (dd, 1H, $J = 9.2$ Hz, $J = 2$ Hz), 6.48-6.50 (t, 1H, $J = 2.4$ Hz), 4.67 (s, 2H), 3.66-3.71 (m, 2H), 3.25-3.5 (m, 2H), 1.40 (s, 9H).


Synthesis of intermediate 26.



[0295] To a solution of intermediate **25** (50 g, 0.26 mol) in THF (1000 ml) was added Et_3N (79 g, 0.79 mol) and Ms_2O (55 g, 0.32 mol) at 0°C under N_2 . The mixture was stirred at 0°C for 2 hrs. TLC (petroleum ether/EtOAc=3/1) showed the reaction was completed. Then it was poured into ice-water and extracted with EtOAc (400 ml \times 2). The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated in vacuum to give intermediate **26** (50 g, 79%) as yellow oil.

[0296] ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) 10.08 (bs, 1H), 7.24-7.29 (m, 2H), 6.81-6.84 (m, 1H), 6.49-6.50 (d, 1H, $J=2.4\text{Hz}$), 4.67 (s, 2H), 4.28-4.31 (m, 2H), 3.48-3.52 (m, 2H), 2.79 (s, 3H), 1.51 (s, 9H).


Synthesis of intermediate 27.



[0297] A solution of intermediate **26** (65 g, 0.19 mol) in DMF (722 ml) was added NaH (60%, 11.5 g, 0.29 mol) at 0 °C. The mixture was stirred at 0 °C for 1 hr under N_2 . TLC (petroleum ether/EtOAc=3/1) showed the reaction was completed. Then the mixture was poured into ice-water and extracted with EtOAc (500 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 50:1 to 10:1) to give intermediate **27** (26 g, 53.2%) as yellow solid.

[0298] ^1H NMR (CDCl_3 , 400 MHz): δ (ppm) 7.14-7.16 (d, 1H, $J=8\text{Hz}$), 7.07 (s, 1H), 6.72-6.83 (m, 1H), 6.48-6.49 (d, 1H, $J=2.8\text{Hz}$), 4.84 4.76 (s, 2H), 4.24-4.25 (m, 2H), 3.94 (m, 2H), 1.45-1.48 (m, 9H).

Synthesis of intermediate 28.




[0299] A solution of intermediate **27** (27.5 g, 95.0 mmol) in dichloromethane (900 ml) was added $(\text{COCl})_2$ (18 g, 142 mmol) at 0 °C under N_2 . The mixture was stirred at 0 °C for 2 hrs. TLC (petroleum ether/EtOAc=1/1) showed the reaction was completed. Then a solution of NaOMe (13.4 g, 247 mmol) in MeOH (40.8 ml) was added at -60 °C under N_2 . The mixture was stirred at

room temperature for 1 hr. Water was added, extracted with dichloromethane (200 ml \times 3). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 20:1 to 1:1) to give intermediate **28** (20 g, 56%) as white solid.

[0300] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 8.37 (s, 1H), 8.02-8.04 (d, 1H, J=8.4Hz), 6.84-6.91 (m, 1H), 4.80-4.90 (m, 2H), 4.44 (bs, 2H), 3.91-3.98 (m, 5H), 1.41-1.46 (m, 9H).


Synthesis of intermediate **29**.



[0301] A solution of intermediate **28** (10 g, 26.5 mmol) and intermediate **3** (4.6 g, 26.5 mmol) in DMF (120 ml) was added A solution of tBuOK (7.4 g, 66.2 mmol) in THF (100 ml) at 0-10 °C. The mixture was stirred at 0-10 °C for 15 min. TLC (petroleum ether/EtOAc =1/1) showed the reaction was completed. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 20:1 to 1:1) to give intermediate **29** (7 g, 52.5%) as red solid.

[0302] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.07 (s, 1H), 8.02-8.01 (d, 1H, J=4Hz), 8.96 (s, 1H), 7.61-7.63 (t, 1H, J=8.8Hz), 7.45 (s, 1H), 7.17 (s, 1H), 6.57-6.64 (m, 1H), 6.47-6.49 (d, 1H, J=8Hz), 5.77-5.80 (d, 1H J=8.4Hz), 4.73 4.78 (2H), 4.52 (bs, 2H), 3.96 (bs, 2H), 1.25-1.44 (m, 9H).


Synthesis of core **3**.



[0303] A solution of intermediate **29** (5 g, 9.9 mmol) in dichloromethane (50 ml) was added HCl/dioxane (50 ml, 7M) at room temperature. The mixture was stirred at room temperature for 1 hr. TLC (dichloromethane/MeOH=15/1) showed the reaction was completed. The solvent was concentrated in vacuum. The residue was washed with methyl tertiary butyl ether, filtered. The filter cake was dried in vacuo to give **core 3** (4 g, 91.9%) as orange solid.


[0304] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.59 (s, 1H), 10.46 (bs, 2H), 8.49 (s, 1H), 8.18 (s, 1H), 8.08-8.12 (m, 2H, J=14Hz), 7.88-7.92 (t, 1H, J=8.0Hz), 7.25-7.29 (t, 1H, J=6.8Hz), 7.04-7.07 (m, 1H), 6.22-6.25 (m, 1H), 4.76 (bs, 2H), 4.62 (bs, 2H), 3.68 (bs, 2H).

Synthesis of intermediate **31**.



[0305] A solution of intermediate **30** (20 g, 94 mmol) in dioxane (100 ml) was added HCl/dioxane (100 ml, 7M) at room temperature. The mixture was stirred at room temperature for 2 hrs. TLC (petroleum ether/EtOAc=3/1) showed the reaction was completed. The mixture was poured onto methyl tertiary butyl ether (300 ml), and filtered. The filter cake was dried in vacuo to give intermediate **31** hydrochloride (8 g, 57%) as white solid.


Synthesis of intermediate **32**.



[0306] A solution of intermediate **31** hydrochloride (7 g, 47 mmol) in dichloromethane (180 ml) was added Et₃N (14.2 g, 141 mmol) at room temperature. The mixture was stirred at room

temperature for 10 min. Then a solution of Triphosgene (5.6 g, 19 mmol) in dichloromethane (20 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH=10/1) showed the reaction was completed. Then the mixture was washed with aq. NaHCO₃, water, brine, dried over Na₂SO₄ and concentrated in vacuum. The crude product was distilled in vacuum to afford intermediate **32** (2.5 g, 30.5%) as colorless oil.


Synthesis of compound I-5.



[0307] A solution of **core 3** (5 g, 11.45 mmol) in DMF (70 ml) was added Et₃N (3.5 g, 34.35 mmol) at room temperature. The mixture was stirred at room temperature for 10 min. Then a solution of intermediate **32** (3.6 g, 20.5 mmol) in DMF (5 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH =15/1) showed the reaction was completed. The mixture was poured into ice-water and extracted with methyl tertiary butyl ether to remove impurities, then extracted with EtOAc (100 ml×5). The combined EtOAc phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum to give the crude **Compound I-5** (4.5 g, crude) as an orange solid.

[0308] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.3 (s, 1H), 9.58 (s, 1H), 8.05 (s, 1H), 7.92 (s, 1H), 7.65 (d, 1H, *J*=8.8Hz), 7.60 (d, 1H, *J*=6.8Hz), 7.21 (t, 1H, *J*=8Hz), 6.78 (d, 1H, *J*=9.6Hz), 6.58 (t, 1H, *J*=6.8Hz), 5.62 (d, 1H, *J*=2Hz), 4.64 (s, 2H), 4.55 (s, 2H), 3.85 (s, 2H), 3.21-3.37 (m, 2H), 2.73-2.76 (m, 2H), 1.75-1.76 (m, 2H), 1.33-1.47 (m, 3H).

Synthesis of compound I-6.




[0309] To solution of crude intermediate **33** (**Compound I-5**) (4.5 g) in THF (100 ml) was added NaBH₄ (0.16 g, 4.2 mmol) in portions below 5 °C. After addition, the reaction mixture was stirred below 5 °C for 0.5 hr. TLC (dichloromethane/MeOH =15/1) showed the reaction was completed. Then the mixture was poured into water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with dichloromethane/MeOH from 200:1 to 30:1) to give **compound I-6** (3 g, 48% for two steps) as orange solid.

[0310] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.24 (s, 1H), 8.06 (s, 1H), 7.92 (s, 1H), 7.64-7.66 (d, 1H, J=8.8Hz), 7.59-7.61 (d, 1H, J=6.4Hz), 7.18-7.22 (t, 1H J=8Hz), 6.78-6.80 (d, 1H, J=9.6Hz), 6.56-6.59 (t, 1H J=6.8Hz), 5.62-5.65 (dd, 1H J=2Hz J=10Hz), 4.63 (s, 2H), 4.55 (s, 2H), 4.45-4.47 (t, 2H, J=5.2Hz), 3.85 (s, 2H), 3.40-3.43 (d, 2H J=12.4Hz), 3.23-3.26 (t, 2H, J=4.8Hz), 2.58-2.65 (t, 2H, J=12Hz), 1.55-1.58 (d, 2H J=12.8Hz), 1.47-1.48 (d, 1H J=6.4Hz), 1.05-1.13 (m, 2H).

[0311] LC/MS M+1 543.1

Synthesis of intermediate 35.




[0312] A solution of intermediate **34** hydrochloride (8.6 g, 55.0 mmol) in dichloromethane (240 ml) was added Et₃N (16.7g, 165.0mmol) at room temperature. The mixture was stirred at room temperature for 10 min. Then a solution of Triphosgene (6.5g 22.0 mmol) in

dichloromethane (20 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH =10/1) showed the reaction was completed. Then the mixture was washed with aq. NaHCO₃, water, brine, dried over Na₂SO₄ and concentrated in vacuum. The crude product was distilled in vacuum to afford intermediate **35** (4.2 g, 42%) as colorless oil.

[0313] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 3.80-3.85 (bs, 2H), 3.71-3.75 (bs, 2H), 2.01-2.11 (m, 4H).

Synthesis of compound I-7.




[0314] A solution of **core 3** (3 g, 6.8 mmol) in DMF (40 ml) was added Et₃N (2.1 g, 20.6 mmol) at room temperature. The mixture was stirred at room temperature for 10 min. Then a solution of intermediate **35** (1.4 g 7.5 mmol) in DMF (5 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH =10/1) showed the reaction was completed. Then the mixture was poured into ice-water and extracted with methyl tertiary butyl ether to remove impurities, filtered. The filter cake was washed with water (100 ml × 3), dissolved by dichloromethane (200 ml), washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give **Compound I-7** (2.2 g, 58%) as orange solid.

[0315] ¹H NMR (DMSO-*d*₆, 400 MHz): δ (ppm) 11.24 (s, 1H), 8.06 (s, 1H), 7.91 (s, 1H), 7.62-7.66 (m, 2H), 7.19-7.23 (t, 1H, *J*=8Hz), 6.84-6.86 (d, 1H, *J*=8.0Hz), 6.58-6.61 (t, 1H, *J*=6.4Hz), 5.63-5.66 (d, 1H, *J*=8.0Hz), 4.70 (s, 2H), 4.58 (bs, 2H), 3.90 (bs, 2H), 3.15 (bs, 4H), 1.93-1.96 (m, 4H).

[0316] LC/MS M+1 549.1


Synthesis of intermediate 37.



[0317] A solution of intermediate **36** hydrochloride (15 g, 100 mmol) in dichloromethane (430 ml) was added Et₃N (30.6g, 300 mmol) at room temperature. The mixture was stirred at room temperature for 10 min. Then a solution of Triphosgene (11.9 g 40 mmol) in dichloromethane (20 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH =10/1) showed the reaction was completed. Then the mixture was washed with aq. NaHCO₃, water, brine, dried over Na₂SO₄ and concentrated in vacuum. The crude product was distilled in vacuum to afford intermediate **37** (9.2 g, 52%) as colorless oil.

[0318] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 4.40 (s, 2H), 3.90-3.94 (d, 2H, J=13.2Hz), 3.41-3.44 (d, 1H, J=13.2Hz), 3.23-3.36 (d, 1H, J=12.8Hz), 1.90-2.04 (m, 2H), 1.80-1.86 (m, 2H).

Synthesis of compound I-8.




[0319] A solution of **core 3** (3 g, 6.8 mmol) in DMF (40 ml) was added Et₃N (2.1 g, 20.6 mmol) at room temperature. The mixture was stirred at room temperature for 10 min. Then a solution of intermediate **37** (1.3g 7.5 mmol) in DMF (5 ml) was added to the mixture at 0 °C ~10 °C. The mixture was stirred at room temperature for 2 hrs. TLC (dichloromethane/MeOH =10/1) showed the reaction was completed. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml ×4). The combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum. The residue was purified by flash column

chromatography (eluted with dichloromethane/MeOH from 200:1 to 50:1) to give **compound I-8** (2.3 g, 62%) as orange solid.

[0320] $^1\text{H NMR}$ (DMSO- d_6 , 400 MHz): δ (ppm) 11.24 (s, 1H), 8.05 (s, 1H), 7.92 (s, 1H), 7.60-7.66 (m, 2H), 7.20-7.24 (t, 1H, $J=7.6\text{Hz}$), 6.80-6.82 (d, 1H, $J=9.2\text{Hz}$), 6.57-6.59 (t, 1H, $J=6.8\text{Hz}$), 5.64-5.67 (d, 1H, $J=10\text{Hz}$), 4.60 (s, 2H), 4.54 (bs, 2H), 4.19 (s, 2H), 3.84 (bs, 2H), 3.20-3.24 (d, 2H, $J=12.8\text{Hz}$), 3.01-3.04 (d, 2H, $J=12\text{Hz}$), 1.74 (s, 4H).


[0321] LC/MS M+1 541.1

Synthesis of intermediate 39.



[0322] To a solution of intermediate **38** (20 g, 113 mmol) in MeOH (100 ml) was added HCl/MeOH (4 M, 100 ml), then stirred at room temperature overnight. The mixture was concentrated in vacuum. To the residue was added water (500 ml), then extracted with EtOAc (200 ml \times 4). The combined organic phases were washed with brine, dried over Na_2SO_4 and concentrated in vacuum to give the crude intermediate **39** (21 g) as brown solid, which was used directly for the next step without purification.


Synthesis of intermediate 40.



[0323] To a solution of crude intermediate **39** (21 g) in MeOH (100 ml) was added NH_3/MeOH (6 M, 100 ml). The reaction mixture was stirred at room temperature overnight. The mixture was poured into EtOAc (500 ml), and then filtered. The filter cake was dried in vacuo to give intermediate **40** (8 g, 40% for two steps) as off-white solid.

[0324] ^1H NMR (DMSO- d_6 , 400 MHz): δ (ppm) 7.86 (d, $J=8.0\text{Hz}$, 1H), 7.80 (s br, 1H), 7.72 (d, $J=8.4\text{Hz}$, 1H), 7.62-7.66 (m, 1H), 7.37-7.41 (m, 1H), 7.22 (s br, 1H), 3.88 (s, 2H).


Synthesis of intermediate 41.



[0325] To a solution of intermediate **28** (10 g, 26.5 mmol) and intermediate **40** (4.6 g, 26.5 mmol) in DMF (120 ml) was added a solution of tBuOK (7.4 g, 66.2 mmol) in THF (100 ml) at 0-10 °C. The mixture was stirred at 0-10 °C for 15 min. TLC (petroleum ether/EtOAc=1/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc (100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was purified by flash column chromatography (eluted with petroleum ether/EtOAc from 10:5 to 1:2) to give intermediate **41** (5 g, 37.5%) as red solid.

[0326] ^1H NMR (DMSO- d_6 , 400 MHz): δ (ppm) 11.07 (s, 1H), 8.02-8.01 (d, 1H, $J=4\text{Hz}$), 8.96 (s, 1H), 7.61-7.63 (t, 1H, $J=8.8\text{Hz}$), 7.45 (s, 1H), 7.17 (s, 1H), 6.57-6.64 (m, 1H), 6.47-6.49 (d, 1H, $J=8\text{Hz}$), 5.77-5.80 (d, 1H, $J=8.4\text{Hz}$), 4.73-4.78 (d, 2H), 4.52 (bs, 2H), 3.96 (bs, 2H), 1.25-1.44 (m, 9H).

Synthesis of intermediate 42.




[0327] A solution of intermediate **41** (5 g, 9.9 mmol) in dichloromethane (50 ml) was added HCl/dioxane (50 ml, 7M) at room temperature. The mixture was stirred at room temperature for

1 hr. TLC (dichloromethane/MeOH=15/1) showed the reaction was complete. The solvent was concentrated to give intermediate **42** (4 g, 92%) as orange solid.

[0328] $^1\text{H NMR}$ (DMSO- d_6 , 400 MHz): δ (ppm) 11.59 (s, 1H), 10.46 (s, 1H), 8.49 (s, 1H), 8.17 (s, 1H), 8.08-8.12 (m, 2H, $J=14\text{Hz}$), 7.88-7.92 (t, 1H, $J=8.4\text{Hz}$), 7.25-7.29 (t, 1H, $J=13.6\text{Hz}$), 7.04-7.07 (t, 1H, $J=10\text{ Hz}$), 6.22-6.25 (m, 1H), 4.76 (s, 2H), 4.62 (s, 2H), 4.45 (s, 2H), 3.68 (s, 2H).

Synthesis of compound I-9.



[0329] To A solution of intermediate **42** (2 g, 4.9 mmol) and 1-Piperidinecarbonyl chloride (1.1 g, 7.4 mmol) in DMF (20 ml) was added Et_3N (1.5 g, 14.9 mmol) at room temperature. The mixture was stirred at room temperature for 30 min. TLC (dichloromethane/MeOH =15/1) showed the reaction was complete. The mixture was poured into ice-water and extracted with EtOAc (100 ml \times 4). The combined organic phases were washed with water, brine, dried over Na_2SO_4 and concentrated in vacuum. The residue was washed with methyl tertiary butyl ether, filtered. The filter cake was dried in vacuum to give **compound I-9** (1.1 g, 43.1%) as an orange solid.

[0330] $^1\text{H NMR}$ (DMSO- d_6 , 400 MHz): δ (ppm) 11.48 (s, 1H), 8.23 (s, 1H), 7.83-7.86 (d, 1H, $J=8.8\text{ Hz}$), 7.68-7.76 (m, 1H), 7.35-7.39 (t, 1H, $J=7.6\text{ Hz}$), 6.89-6.92 (m, 1H), 6.04-6.07 (m, 1H), 4.66 (s, 2H), 4.57-4.65 (bs, 2H), 3.83-3.90 (m, 2H), 3.01-3.05 (m, 4H), 1.46-1.49 (m, 6H).

[0331] LC/MS $M+514.1$

Synthesis of Compound I-12.

[0332] **Compound I-12** can be synthesized in a similar manner to **Compound I-7** using 3,3-difluoropiperidine as the starting material.


[0333] $^1\text{H NMR}$ (DMSO- d_6 , 400 MHz): δ (ppm) 11.25 (s, 1H), 8.06 (s, 1H), 7.93 (s, 1H), 7.58-7.66 (m, 2H), 7.18-7.23 (t, 1H, $J=8\text{Hz}$), 6.81-6.79 (d, 1H, $J=8.0\text{Hz}$), 6.58-6.61 (t, 1H,

$J=6.4\text{Hz}$), 5.65-5.67 (d, 1H, $J=8.0\text{Hz}$), 4.66 (s, 2H), 4.56 (bs, 2H), 3.88 (bs, 2H), 3.31-3.64 (bs, 2H), 3.07 (bs, 2H), 1.96-2.01 (bs, 2H), 1.69 (bs, 2H).

[0334] LC/MS $M+1$ 549.1

Synthesis of Compound I-13.


[0335] Compound I-13 as shown in Synthetic Scheme 9.



[0336] The synthesis of Intermediate 44 in a similar manner to Intermediate 47, yield 43 %.


[0337] ^1H NMR (400 MHz, CDCl_3) δ 1.47 (9H, s), 1.91-1.94 (2H, m), 3.37-3.56 (4H, m), 4.40-4.70 (1H, m), 4.71-4.74 (1H, m)

Synthesis of Intermediate 45



[0338] The synthesis of Intermediate 45 in a similar manner to Intermediate 48, crude.

Synthesis of Compound I-13



[0339] The synthesis of Compound I-13 is similar to Compound I-19, yield 20.7 % for two steps.

[0340] ^1H NMR (400 MHz, DMSO- d_6) δ 1.45-1.48 (1H, d, $J = 9.2$ Hz), 1.54-1.56 (1H, d, $J = 9.2$ Hz), 2.71-2.73 (1H, d, $J = 9.2$ Hz), 2.88-2.90 (1H, d, $J = 8.4$ Hz), 2.97-2.99 (1H, d, $J = 9.6$ Hz), 3.35-3.37 (1H, d, $J = 8.0$ Hz), 3.49 (1H, s), 3.75-3.76 (1H, m), 3.90-3.92 (1H, m), 4.04 (1H, s), 4.46-4.50 (1H, m), 4.58-4.74 (3H, m), 5.60-5.62 (1H, d, $J = 8.4$ Hz), 6.60-6.63 (1H, m), 6.78-6.80 (1H, d, $J = 9.6$ Hz), 7.20-7.24 (1H, m), 7.63-7.66 (2H, m), 7.90 (1H, s), 8.06 (1H, s).

[0341] MS Calcd: 525.5; MS Found: 526.2 $[\text{M}+1]^+$.

Synthesis of Compound I-14.

[0342] Compound I-14 can be synthesized in a similar manner to Compound I-8.

[0343] LC/MS $\text{M}+1$ 548.2

Synthesis of Compound I-15.

[0344] Compound I-15 can be synthesized in a similar manner to Compound I-7.

[0345] LC/MS $\text{M}+1$ 556.2

Synthesis of Compound I-16.

[0346] Compound I-16 can be synthesized in a similar manner to Compound I-7.

[0347] LC/MS $\text{M}+1$ 556.2

Synthesis of Compound I-17.

[0348] Compound I-17 can be synthesized in a similar manner to Compound I-7.


[0349] LC/MS $\text{M}+1$ 599.2

Synthesis of Compound I-18.

[0350] Compound I-18 can be synthesized in a similar manner to Compound I-8.

[0351] LC/MS $\text{M}+1$ 591.2

Synthesis of Compound I-19.



46


47

[0352] To a solution, Intermediate 46 (1.0 g, 4.67 mmol) in Dichloromethane (25 ml) was added Et_3N (1.41 g, 14.0 mmol). The suspension was stirred and treated dropwise with Triphosgene (0.55 g, 1.87 mmol) in DCM (5 ml) at 0~10 °C. After addition was completed, the suspension was stirred for 2 hrs at room temperature. The reaction mixture poured into iced water (20 ml), extracted with DCM, and the organic layer was washed with NaHCO_3 (aq), brine,

dried over Na_2SO_4 , filtered and concentrated in vacuum. The residue was purified by silica gel column chromatography to give Intermediate **47** (0.6 g, 46.5 %) as colorless oil.


[0353] $^1\text{H NMR}$ (400 MHz, CDCl_3) δ 1.20-1.27 (2H, m), 1.41 (9H, s), 1.72-1.79 (3H, m), 2.84-2.90 (1H, m), 3.03-3.09 (3H, m), 4.32-4.35 (2H, m), 4.66 (1H, br s).

Synthesis of Compound I-19



[0354] To a solution of **Core 3** (1.0 g, 2.29 mmol) in DMF (13 ml) was added Et_3N (0.70 g, 6.87 mmol). The suspension was stirred and treated dropwise with Intermediate **47** (0.70 g, 2.52 mmol) in DMF (2 ml) at 0~10 °C. After addition was completed, the suspension was stirred for 2 hrs at room temperature. The reaction mixture was poured into iced water (60 ml), filtered and concentrated in vacuo to give Intermediate **48** (1.0 g, crude) as red solid.

Synthesis of Compound I-19




[0355] To a solution of Intermediate **48** (1.0 g, crude) in DCM (30 ml) was added HCl/dioxane (10 ml, 8 mol/L) at room temperature, and the mixture was stirred at room temperature for 2 hrs. The reaction mixture was filtered, and the filter cake was dissolved in water, adjusted to pH=8~9 with Na_2CO_3 (aq), and filtered again, washed with water and concentrated in vacuo to give **Compound I-19** (130 mg, 10.5 %, two steps) as red solid.

[0356] $^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ 1.03-1.11 (2H, m), 1.13-1.62 (3H, m), 2.44-2.82 (4H, m), 3.41-3.44 (2H, m), 3.85 (2H, m), 4.54 (2H, m), 4.64 (2H, m), 5.64-5.66 (1H, d, $J = 9.6$ Hz),

6.56-6.58 (1H, m), 6.78-6.81 (1H, m), 7.19-7.23 (1H, m), 7.60-7.66 (2H, m), 7.92 (1H, s), 8.06 (1H, s).


[0357] MS Calcd: 541.6; MS Found: 542.2([M+1]⁺).

Synthesis of Compound I-20.



[0358] To a solution of Intermediate **49** (15 g, 130.2 mmol) in DCM (1.5 L) was added Et₃N (19.8 g, 195.3 mmol) and DMAP (0.8 g, 6.5 mmol). The suspension was stirred and treated dropwise with tert-butylchlorodiphenylsilane (53.7 g, 195.3 mmol) at 0~10 °C. After addition was completed, the suspension was stirred for 5 hrs at room temperature. The reaction mixture poured into iced water (500 ml), extracted with DCM (300 ml×2), and the organic layer was washed with brine (300 ml×2), dried over Na₂SO₄, filtered and concentrated in vacuum to give Intermediate **50** (30 g, crude) as yellow oil.


Synthesis of Intermediate 51



[0359] To a solution of Intermediate **50** (30 g, crude, 84.8 mmol) in DCM (900 ml) was added Et₃N (25.8 g, 254.5 mmol). The suspension was stirred and treated dropwise with Triphosgene (10.1 g, 33.9 mmol) in DCM (50 ml) at 0~10 °C. After addition was completed, the suspension was stirred for 2 hrs at room temperature. The reaction mixture poured into iced water (300 ml), extracted with DCM, and the organic layer was washed with NaHCO₃ (aq), brine, dried over Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by silica gel column chromatography to give Intermediate **51** (3.1 g, 8.8 %) as colorless oil.


[0360] ^1H NMR (400 MHz, CDCl_3) δ 1.05 (9H, s), 1.22-1.29 (2H, m), 1.78-1.83 (3H, m), 2.84-2.90 (1H, m), 3.01-3.07 (1H, m), 3.50-3.51 (2H, d, $J = 5.6$ Hz), 4.30-4.34 (2H, m), 7.36-7.45 (6H, m), 7.62-7.64 (4H, m).

Synthesis of Intermediate 52



[0361] To a solution of **CF₃ Core** (the synthesis is described in **Scheme 2**) (1.6 g, 3.28 mmol) in DMF (15 ml) was added Et_3N (1.0 g, 9.84 mmol). The suspension was stirred and treated dropwise with Intermediate **51** (1.5 g, 3.60 mmol) in DMF (5 ml) at 0–10 °C. After addition was completed, the suspension was stirred for 2 hrs at room temperature. The reaction mixture poured into iced water (80 ml), filtered and the filter cake was washed with water and MTBE concentrated *in vacuo* to give Intermediate **52** (0.98 g, crude) as red solid.

Synthesis of Compound I-20:




[0362] To a solution of Intermediate **52** (0.98 g, crude) in THF (20 ml) was added TBAF (0.56 g, 1.77 mmol) in THF (10 ml) at room temperature, and the mixture was stirred at room temperature for 2 hrs at room temperature. The reaction mixture was poured into iced water (40 ml), filtered, and the filter cake was concentrated *in vacuo*, purified by silica gel column chromatography to give to **Compound I-20** (160 mg, 22.8 %) as orange solid.

[0363] ^1H NMR (400 MHz, DMSO- d_6) δ 1.05-1.13 (2H, m), 1.48-1.58 (3H, m), 2.54-2.67 (2H, m), 3.23-3.26 (2H, m), 3.40-3.44 (2H, m), 3.87-3.90 (2H, m), 4.44-4.47 (1H, m), 4.62-4.69 (2H, m), 4.72 (2H, s), 6.31 (1H, s), 6.53-6.56 (1H, m), 7.14-7.20 (2H, m), 7.60-7.64 (2H, m), 7.92 (1H, s), 8.16 (1H, s), 11.29 (1H, s).


[0364] MS Calcd: 592.6; MS Found: 593.2([M+1] $^+$).

Synthesis of Compound I-21.



[0365] The synthesis of Intermediate **53** in a similar manner to Intermediate **52**. The synthesis of the CN Core can be prepared from Core 1 by displacing the Br using conditions as described in Scheme 3 followed by deblocking as using conditions as described for Core 1 to intermediate 10.

Synthesis of Compound I-21



[0366] The synthesis of **Compound I-21** is in a similar manner as the final step in the synthesis of **Compound I-20**, yield 28.6 %.

[0367] ^1H NMR (400 MHz, DMSO- d_6) δ 1.08-1.13 (2H, m), 1.47-1.56 (1H, m), 1.56-1.59 (2H, m), 2.58-2.65 (2H, m), 3.23-3.26 (2H, m), 3.40-3.43 (2H, m), 3.86 (2H, s), 4.45-4.61 (1H, m), 4.61 (2H, s), 4.68 (2H, s), 6.44 (1H, s), 6.53-6.57 (1H, m), 7.19-7.24 (1H, m), 7.28 (1H, s), 7.54-7.56 (1H, d, $J = 6.8$ Hz), 7.67-7.69 (2H, d, $J = 9.2$ Hz), 7.98 (1H, s), 8.08 (1H, s), 11.32 (1H, s).


[0368] MS Calcd: 549.6; MS Found: 550.2 [M+1] $^+$.

[0369] **Synthesis of Compounds I-22** MS Calcd: 620.14; MS Found: 621.1 $[M+1]^+$, **23, 24** MS Calcd: 586.14; MS Found: 587.1 $[M+1]^+$, **26, 27** MS Calcd: 584.17; MS Found: 565.1 $[M+1]^+$, can be synthesized in a similar manner to **Compound I-7**.

Synthesis of Compound I-25.


[0370] **Compound I-25** can be synthesized as shown in Synthetic Scheme 11.

Synthesis of intermediate 55




[0371] To a solution of compound **54** (2.3 g, 13.2 mmol) in DCM (30 ml) was added Et₃N (2.9 g, 29.1 mmol) and Boc₂O (3.5 g, 15.9 mmol). Then the mixture was stirred at R.T. for 2 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. The mixture was poured into ice water, extracted with DCM, the combined organic phase was washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give intermediate **55** (2.3 g, 57.8%) as white solid.

Synthesis of intermediate 56




[0372] To a solution of intermediate **55** (2.3 g, 9.7 mmol) in DMF (30 ml) was added NaH (0.5 g, 13.6 mmol) at 0~10 °C. Then BnBr (2.5 g, 14.6 mmol) was added to the mixture, after stirred at R.T. for 2 hrs. TLC (PE/EtOAc=3/1) showed the reaction was complete. The reaction was poured into ice-water, extracted with EtOAc. The combined organic phase was washed with water brine, dried over Na₂SO₄ and concentrated in vacuum to give a crude, which was purified by silica gel chromatography to give intermediate **56** (2.8 g, 87.5%) as white solid.

Synthesis of intermediate 57



[0373] To a solution of intermediate **56** (2.8 g, 8.6 mmol) in dioxane (10 ml) was added HCl/dioxane (7M, 30 ml). The mixture was stirred at RT for 2 hrs. TLC (PE/EtOAc=3/1) showed the reaction was complete. Then the mixture was concentrated in vacuum to give intermediate **57** (1.9 g, crude) as yellow oil.


Synthesis of intermediate 58



[0374] To a solution of intermediate **57** (1.9g, crude) in DCM (50 ml) was added Et₃N (2.1 g, 21.1 mmol) at 0~10 °C, then Triphosgene (0.8 g, 2.8 mmol) in DCM (10 ml) was added dropwise. The mixture was stirred at room temperature for 2 hrs. TLC (PE/EtOAc=5/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with DCM, the combined organic phases were washed with NaHCO₃ (aq), water, brine, dried over Na₂SO₄ and concentrated in vacuum to give a crude, which was purified by silica gel chromatography to give intermediate **58** (0.6 g, 24.2% for two steps) as white solid.

[0375] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.24~7.36 (m, 5H), 4.66~5.27 (m, 2H), 4.16~4.55 (m, 2H), 3.60~3.65 (m, 1H), 3.23~3.30 (m, 1H), 3.17~3.21 (m, 1H), 2.24~2.42 (m, 1H), 1.94~2.03 (m, 1H).


Synthesis of intermediate 59



[0376] To a solution of **core 3** (1.1 g, 2.6 mmol) in DMF (10 ml) was added Et₃N (0.6 g, 6.0 mmol), then intermediate **58** (0.6 g, 2.0 mmol) in DMF (3 ml) was added dropwise at 0~10 °C, the mixture was stirred at room temperature for 2 hr. TLC (DCM/MeOH=10/1) showed the reaction was complete. Then the mixture was poured into ice-water and MTBE, filtered and the filtered cake was washed with MTBE, concentrated in vacuum to give intermediate **59** (1.0 g, 76.3%) as yellow solid.

[0377] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 11.24 (s, 1H), 8.06 (s, 1H), 7.89 (s, 1H), 7.63~7.66 (m, 2H), 7.26~7.32 (m, 3H), 7.13~7.22 (m, 3H), 6.79~6.83 (m, 1H), 6.56 (t, 1H, *J*=6.4Hz), 5.64~5.67 (m, 1H), 4.40~4.72 (m, 6H), 3.87~3.89 (m, 1H), 3.70~3.79 (m, 2H), 3.38~3.39 (m, 1H), 3.33 (s, 2H), 3.17~3.16 (m, 1H), 2.06~2.14 (m, 1H), 1.86~1.90 (m, 1H).

Synthesis of Compound I-26



[0378] To a solution of intermediate **59** (1.0 g, 1.5 mmol) in EtOH (5 ml) was added HCl (12M, 10 ml). The mixture was heated to reflux for 4 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. Then the mixture was cooled, adjusted to pH=7~8 with Na₂CO₃ (aq), filtered and the filtered cake was washed with MTBE, dried in vacuum to give **Compound I-25** (0.4 g, 46.5 %) as orange solid.


[0379] ^1H NMR (DMSO- d_6 , 400 MHz): δ (ppm) 11.24 (s, 1H), 8.06 (s, 1H), 7.92 (s, 1H), 7.63~7.67 (m, 2H), 7.23 (t, 1H, $J=7.6\text{Hz}$), 6.82 (dd, 1H, $J=2.0\text{Hz}$, 9.6Hz), 6.62 (t, 1H, $J=6.4\text{Hz}$), 5.75~5.76 (m, 1H), 5.65 (dd, 1H, $J=2.4\text{Hz}$, 10.0Hz), 4.65~4.74 (m, 2H), 4.49~4.62 (m, 2H), 3.83~3.95 (m, 2H), 3.67~3.69 (m, 1H), 3.16~3.25 (m, 2H), 3.05~3.10 (m, 2H), 2.07~2.14 (m, 1H), 1.86~1.88 (m, 1H).

[0380] MS Calcd: 564.17; MS Found: 565.1 $[\text{M}+1]^+$,

Synthesis of Compound I-28.

[0381] **Compound I-28** MS Calcd: 513.19; MS Found: 514.2 $[\text{M}+1]^+$ can be synthesized as shown in Synthetic Scheme 12. Intermediate 60 is converted to **Intermediate 61** which is carried to **Compound I-28** using conditions as described for **Intermediate 2** to **Intermediate 3** and **Intermediate 28** to Core 3 and Core 3 to **Compound I-1**

Synthesis of intermediate 61.



To a solution of **60** (10 g, 91.7 mmol) in POCl_3 (200 ml) was added diethyl malonate (73.4 g, 458.3 mmol). The reaction mixture was heated to reflux and stirred for 3 hrs. After cooling, the reaction mixture was poured into ice-water, NaHCO_3 was added until $\text{pH}=7\sim 8$, extracted with EtOAc. The organic phase was washed with brine, dried over Na_2SO_4 , filtered, concentrated under reduced pressure. The residue was purified by flash column chromatography to give **61** (1.8 g, 9.5%) as white solid.


^1H NMR (DMSO- d_6): δ (ppm) 8.43 (d, 1H, $J=6.8\text{Hz}$), 7.75 (d, 1H, $J=9.2\text{Hz}$), 7.37-7.41 (t, 1H), 6.99-7.02 (t, 1H), 4.41 (s, 2H), 4.10-4.16 (q, 2H), 1.17-1.21 (t, 3H).

Synthesis of **Compound I-29** and **Compound 30** can be synthesized in a similar manner as the undeuterated material by utilizing the appropriate deuterated starting materials.

Synthesis of Compound I-31.

[0382] **Compound I-31** can be synthesized as shown in Synthetic Scheme 13.


Synthesis of intermediate 65



[0383] To a solution of compound **65** (5.0 g, 23.5 mmol) in THF (50 ml) was added TMSCF₃ (4.0 g, 28.2 mmol) and TBAF (28.2 ml, 28.2 mmol) at 0 °C. Then the mixture was warmed and stirred at 30 °C for 3 hrs. TLC (PE/EtOAc=1/1) showed the reaction was complete. The mixture was poured into ice-water, extracted with EtOAc (40 ml ×3). The combined organic phase were washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give intermediate **66** (6 g, crude) as yellow oil.


[0384] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 4.10~4.16 (m, 2H), 3.76~3.77 (m, 1H), 2.70~2.71 (m, 2H), 2.21 (d, 1H, *J*=6.0Hz), 1.85~1.89 (m, 2H), 1.56~1.64 (m, 1H), 1.53~1.54 (m, 3H), 1.46 (s, 9H).

Synthesis of intermediate 67




[0385] To a solution of intermediate **66** (5 g, 17.7 mmol) in DMF (50 ml) was added NaH (1.0g, 24.7 mmol) at 0~10 °C, then BnBr (4.5 g, 26.5 mmol) was added dropwise. The mixture was stirred at room temperature for 2 hrs. TLC (PE/EtOAc=2/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with EtOAc, the combined organic phases were washed with water, brine, dried over Na₂SO₄ and concentrated in vacuum to give intermediate **67** (7.5 g, crude) as yellow oil.

Synthesis of intermediate 68



[0386] To a solution of intermediate **67** (7.5 g, crude) in dioxane (10 ml) was added HCl/dioxane (7M, 50ml), then the mixture was stirred at room temperature for 3 hrs. TLC (PE/EtOAc=2/1) showed the reaction was complete. Then the mixture was concentrated in vacuum to give intermediate **68** (6.6 g, crude) as yellow oil.


Synthesis of intermediate 69



[0387] To a solution of intermediate **68** (6.6 g, crude) in DCM (130 ml) was added Et₃N (6.5 g, 63.9 mmol) at 0~10 °C, then Triphosgene (2.5 g, 8.5 mmol) in DCM (20 ml) was added dropwise. The mixture was stirred at room temperature for 2 hr. TLC (PE/EtOAc=5/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with DCM, the combined organic phases were washed with NaHCO₃ (aq), water, brine, dried over Na₂SO₄ and concentrated in vacuum to give a crude, which was purified by silica gel chromatography to give intermediate **69** (2.0 g, 33.8% for three steps) as white solid.

[0388] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.32~7.41 (m, 5H), 4.86 (d, 1H, *J*=11.2Hz), 4.52 (d, 1H, *J*=11.2Hz), 4.33 (d, 2H, *J*=12.4Hz), 3.55~3.57 (m, 1H), 3.00~3.03 (m, 1H), 2.81~2.84 (m, 1H), 1.89~1.97 (m, 2H), 1.61~1.64 (m, 1H), 1.45~1.53 (m, 2H).


Synthesis of intermediate 70



[0389] To a solution of **core 3** (2.3 g, 5.4 mmol) in DMF (40 ml) was added Et₃N (1.6 g, 16.2 mmol), then intermediate **69** (2.0 g, 6 mmol) in DMF (10 ml) was added dropwise at 0~10 °C, the mixture was stirred at room temperature for 2 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. Then the mixture was poured into ice-water, filtered and the filtered cake was washed with MTBE, concentrated in vacuum to give intermediate **70** (0.7 g, 17.9%) as red solid.

[0390] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 11.25 (s, 1H), 8.06 (s, 1H), 7.92 (s, 1H), 7.59~7.65 (m, 2H), 7.32~7.37 (m, 5H), 7.15~7.20 (m, 1H), 6.73 (dd, 1H, *J*=2.0Hz, 9.6Hz), 6.54~6.56 (m, 1H), 5.63 (dd, 1H, *J*=2.4Hz, 10.0Hz), 4.74 (d, 1H, *J*=11.2Hz), 4.54~4.64 (m, 5H), 4.01 (t, 1H, *J*=4.8Hz), 3.84~3.86 (m, 2H), 3.41~3.46 (m, 2H), 2.63~2.65 (m, 2H), 1.75~1.90 (m, 1H), 1.53~1.62 (m, 2H), 1.37~1.40 (m, 2H).


Synthesis of Compound I-31



[0391] To a solution of intermediate **70** (0.7g, 1.0 mmol) in EtOH (20 ml) was added HCl (12M, 10 ml). The mixture was heated to reflux for 5 hrs. TLC (D/M=10/1) showed the reaction

[0399] To a solution of compound **71** (5.0 g, 23.3 mmol) in THF (300 ml) was added 2,4-Dimethoxybenzaldehyde (3.9g, 23.3 mmol), then the mixture was stirred at R.T. for 2 hrs. NaBHAc₃ (7.4 g, 35 mmol) was added slowly at 0 °C. Then the mixture was stirred at R.T. for 3 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. The mixture was poured into ice-water, separated, the organic phase was washed with brine, dried over Na₂SO₄ and concentrated in vacuum to give intermediate **72** (9 g, crude) as yellow oil.


Synthesis of intermediate **73**



[0400] To a solution of intermediate **72** (9.0 g, crude) in DCM (300 ml) was added Et₃N (7.5 g, 74.1 mmol) at 0~10 °C, then Triphosgene (2.9 g, 9.9 mmol) in DCM (30 ml) was added dropwise. The mixture was stirred at room temperature for 2 hrs. TLC (PE/EtOAc=3/1) showed the reaction was complete. Then the mixture was poured into ice-water and extracted with DCM, the combined organic phases were washed with NaHCO₃ (aq), water, brine, dried over Na₂SO₄ and concentrated in vacuum to give a crude, which was purified by silica gel chromatography to give intermediate **73** (3.0 g, 30% for two steps) as yellow oil.

[0401] ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 7.07~7.19 (m, 1H), 6.45~6.49 (m, 2H), 4.65 (s, 1H), 4.54 (s, 1H), 4.09~4.14 (m, 2H), 3.81 (s, 3H), 3.80 (s, 3H), 3.29 (d, 1H, *J*=6.8Hz), 3.14 (s, 1H), 2.64~2.67 (m, 2H), 1.90~1.91 (m, 1H), 1.60~1.66 (m, 2H), 1.46 (s, 9H), 1.12~1.18 (m, 2H).


Synthesis of intermediate **74**



[0402] To a solution of **core 3** (2.8 g, 6.4 mmol) in DMF (30 ml) was added Et₃N (1.9 g, 19.2 mmol), then intermediate **73** (3.0 g, 7.0 mmol) in DMF (5 ml) was added dropwise at 0~10 °C, the mixture was stirred at room temperature for 2 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. Then the mixture was poured into ice-water and MTBE, filtered and the filtered cake was washed with MTBE, concentrated in vacuum to give intermediate **74** (1.2 g, 24%) as yellow solid.

[0403] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 11.23 (s, 1H), 8.09 (s, 1H), 7.91 (s, 1H), 7.57~7.64 (m, 2H), 7.08 (t, 1H, *J*=8.0Hz), 6.98 (d, 1H, *J*=8.4Hz), 6.71 (dd, 1H, *J*=1.6Hz, 9.2Hz), 6.49 (d, 1H, *J*=2.0Hz), 6.42~6.45 (m, 2H), 5.59~5.62 (m, 1H), 4.65 (s, 2H), 4.53 (s, 2H), 4.14 (s, 2H), 3.82~3.88 (m, 4H), 3.72 (s, 3H), 3.62 (s, 3H), 2.75 (d, 2H, *J*=6.8Hz), 2.50~2.51 (m, 2H), 1.58 (s, 1H), 1.42~1.45 (m, 2H), 1.34 (s, 9H), 0.71~0.75 (m, 2H).

Synthesis of Compound I-37



[0404] To a solution of intermediate **74** (1.2 g, 1.5 mmol) in DCM (30 ml) was added HCl/dioxane (7 M, 10 ml). The mixture was stirred at R.T. for 2 hrs. TLC (DCM/MeOH=10/1) showed the reaction was complete. Then the mixture was poured into ice-water, filtered and the filtered cake was dissolved with water adjusted to pH=7~8 with Na₂CO₃ (aq), filtered again and the filtered cake was washed with MTBE, dried in vacuum to give **Compound I-37** (0.27 g, 32.9%) as red solid.

[0405] ¹H NMR (DMSO-d₆, 400 MHz): δ (ppm) 8.07 (s, 1H), 7.90 (s, 1H), 7.64 (d, 2H, *J*=8.4Hz), 7.19~7.23 (m, 1H), 6.76~6.78 (m, 1H), 6.58~6.62 (t, 1H), 6.50 (s, 1H), 5.66 (dd, 1H, *J*=2.0Hz, 10.0 Hz), 4.78 (s, 2H), 4.53 (s, 2H), 3.90 (bm, 2H), 2.84 (s, 4H), 2.30~2.36 (m, 2H), 1.41 (bd, 3H, *J*=10.8Hz), 0.84~0.92 (m, 2H).

[0406] MS Calcd: 541.22; MS Found: 542.2 [M+1]⁺

Synthesis of Compound I-38.

[0407] **Compound I-38** MS Calcd: 555.24; MS Found: 556.2 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-19**.

Synthesis of Compound I-39.

[0408] **Compound I-39** MS Calcd: 569.26; MS Found: 570.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-8**.

Synthesis of Compound I-40.

[0409] **Compound I-40** MS Calcd: 539.21; MS Found: 540.2 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-8**.

Synthesis of Compound I-41.

[0410] **Compound I-41** MS Calcd: 581.26; MS Found: 582.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-8**.

Synthesis of Compound I-42.

[0411] **Compound I-42** MS Calcd: 581.26; MS Found: 582.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-8**.

Synthesis of Compound I-43.

[0412] **Compound I-43** MS Calcd: 636.14; MS Found: 637.1 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-18**.

Synthesis of Compound I-44.

[0413] **Compound I-44** MS Calcd: 614.17; MS Found: 615.1 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-18**.

Synthesis of Compound I-45.

[0414] **Compound I-45** MS Calcd: 562.24; MS Found: 563.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-36**.

Synthesis of Compound I-46.

[0415] **Compound I-46** MS Calcd: 576.26; MS Found: 577.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-39**.


Synthesis of Compound I-47.

[0416] **Compound I-47** MS Calcd: 588.26; MS Found: 589.3 [M+1]⁺ can be synthesized in a similar manner as the synthesis of **Compound I-42**.

Synthesis of Compound I-48.


[0417] **Compound I-48** MS Calcd: 530.19; MS Found: 531.2 [M+1]⁺ can be synthesized as shown in Synthetic Scheme 15 to produce **Intermediate 79**. **Intermediate 79** is carried to final product as shown in Scheme 6 (**Intermediate 23** to **core 3**) and Scheme 8 (**Intermediate 42** to **Intermediate I-9**).

[0418] **Synthesis of Intermediate 76**




To a solution of **Intermediate 75** (20 g, 98.5 mmol) in dry THF (130 ml) was added $\text{BH}_3 \cdot \text{Me}_2\text{S}$ (10 M, 49 ml, 492.6 mmol) at R.T. under N_2 . The mixture was stirred at RT for 2 hrs and then heated to 50 °C for 1hr. TLC (DCM/MeOH=5/1) showed the reaction was complete. The mixture was cooled to R.T. and MeOH (50 ml) was added at 0 °C slowly, the mixture solution was concentrated to afford the crude **Intermediate 76** (20 g, >100%) as a yellow solid.

[0419] ¹H NMR (DMSO-*d*₆): δ (ppm) 8.32 (dd, 1H, *J*=7.2Hz, 10.4Hz), 7.80 (dd, 1H, *J*=8.0Hz, 11.6Hz), 5.77-5.79 (t, 1H), 4.82 (d, 2 H, *J*=3.6Hz).

Synthetic of Intermediate 77.**Synthesis of Compound I-46.**


To a solution of **Intermediate 76** (20 g, crude) in THF (300 ml) was added MnO₂ (42.8 g, 492.5 mmol) at R.T. under N₂. The mixture was stirred at 80 °C for 5 hrs. TLC (PE/EtOAc=5/1) showed the reaction was complete. Then the mixture was filtered, concentrated to give **Intermediate 77** (20 g, >100%) as yellow oil. ¹H NMR (DMSO-d₆): δ (ppm) 10.17 (s, 1H), 8.45-8.50 (dd, 1H, *J*=6.8Hz, 10.0Hz), 7.99-8.04 (dd, 1H, *J*=8.0Hz, 10.0Hz).

[0420] Synthetic of Intermediate 78.

[0421] To a solution of **Intermediate 77** (20 g, crude) in dry toluene (180 ml) was added *n*-BuOH (26 g) and TsOH (2 g) at R.T. The mixture reaction was stirred at 120 °C for 3 hrs. And remove the water using a Dean-stark apparatus. TLC (PE/EtOAc=5/1) showed the reaction was complete. The mixture was concentrated in vacuum to give crude compound. The crude compound was purified by chromatography on silica gel (PE/EtOAc =1:0~200:1) to afford the **Intermediate 78** (15 g, 48% for three steps) as yellow oil.

¹H NMR (DMSO-d₆): δ (ppm) 8.22-8.27 (dd, 1H, *J*=7.2Hz, 10.0Hz), 7.63-7.68 (dd, 1H, *J*=8.4Hz, 11.2Hz), 5.88 (s, 1H), 3.47-3.59 (m, 4H), 1.47-1.54 (m, 4H), 1.27-1.40 (m, 4H), 0.83-0.89 (m, 6H).

[0422] Synthetic of Intermediate 79.



[0423] To a solution of Intermediate **78** (15 g, 47.3 mmol) in THF (450 ml) was added vinylmagnesium bromide solution (1 M, 189 ml, 189.2 mmol) dropwise at $-40\text{ }^{\circ}\text{C}$ under N_2 . The mixture was stirred at $-40\text{ }^{\circ}\text{C}$ for 1 hr. TLC (PE/EtOAc =5/1) showed the reaction was complete, aq. NH_4Cl was added at $-40\text{ }^{\circ}\text{C} \sim 0\text{ }^{\circ}\text{C}$, extracted with EtOAc, concentrated to give crude compound. The crude compound was purified by chromatography on silica gel to afford the compound (10 g) as yellow oil. To a solution of the compound (10 g) in EtOAc (100 ml) was added HCl (0.5 N, 100 ml) dropwise. The mixture was stirred at R.T. for 1hr. TLC (PE/EtOAc =5/1) showed the reaction was complete. The mixture was adjusted to pH=8~9 with aq. NaOH, extracted with EtOAc, the combined organic phase was washed with brine, dried with Na_2SO_4 , filtered, concentrated in vacuum to give Intermediate **79** (6 g, 70.5%) as yellow solid.

^1H NMR (DMSO- d_6): δ (ppm) 11.99 (s, 1H), 10.08 (s, 1H), 7.89-7.94 (dd, 1H, $J=7.6\text{Hz}$, 10.8Hz), 7.54 (d, 1H, $J=3.2\text{Hz}$), 6.72 (d, 1H, $J=3.2\text{Hz}$)

Biological Examples

Assay:

[0424] Enzymatic activities using recombinant human GSK3 using an in vitro enzymatic assay for inhibition GSK3 α and GSK3 β .

[0425] Enzymes and Substrates:

Assay	Enzyme Used (ng) / Reaction	Substrate/ATP
GSK3 α	13	0.1 mg/ml GSKtide/10 uM ATP
GSK3 β	13	0.1 mg/ml GSKtide/10 uM ATP

[0426] Assay Conditions:

[0427] The assay was performed using Kinase-Glo Max luminescence kinase assay kit (Promega). It measures kinase activity by quantitating the amount of ATP remaining in solution following a kinase reaction. The luminescent signal from the assay is correlated with the amount of ATP present and is inversely correlated with the amount of kinase activity. The compounds

were diluted in 10% DMSO and 5µl of the dilution was added to a 50µl reaction so that the final concentration of DMSO is 1% in all of reactions. The enzymatic reactions were conducted at 30 °C for 40 minutes. The 50µl reaction mixture contains 40 mM Tris, pH 7.4, 10 mM MgCl₂, 0.1 mg/ml BSA, 2 mM DTT, 0.1 mg/ml GSKtide substrate, 10 uM ATP and GSK3. After the enzymatic reaction, 50 µl of Kinase-Glo Max Luminescence kinase assay solution (Promega) was added to each reaction and incubate the plate for 15 minutes at room temperature. Luminescence signal was measured using a BioTek *Synergy 2* microplate reader.

[0428] Data Analysis:

[0429] Kinase activity assays were performed in duplicate at each concentration. The luminescence data were analyzed using the computer software, Graphpad Prism. The difference between luminescence intensities in the absence of Kinase (Lut) and in the presence of Kinase (Luc) was defined as 100 % activity (Lut – Luc). Using luminescence signal (Lu) in the presence of the compound, % activity was calculated as: % activity = {(Lut - Lu)/(Lut - Luc)}×100%, where Lu= the luminescence intensity in the presence of the compound (all percent activities below zero were shown zero in the table).

[0430] The values of % activity versus a series of compound concentrations were then plotted using non-linear regression analysis of Sigmoidal dose-response curve generated with the equation $Y=B+(T-B)/1+10((\text{LogEC50}-X)\times\text{Hill Slope})$, where Y=percent activity, B=minimum percent activity, T=maximum percent activity, X= logarithm of compound and Hill Slope=slope factor or Hill coefficient. The IC₅₀ value was determined by the concentration causing a half-maximal percent activity.

[0431] Table of activity for compounds vs GSK-alpha and GSK-beta:

[0432] The table herein shows activity of certain compounds. For potency, “+” indicates greater than about 1 µM; “++” indicates about 100 nM to 1 µM; “+++” indicates about 10 nM to 100 nM, and “++++” indicates about 1 nM to 10 nM.

Compound	Potency Alpha	Potency Beta
Compound I-1	++++	++++
Compound I-2	++++	++++
Compound I-3	++++	++++
Compound I-4	++++	++++

Compound I-6	++++	++++
Compound I-7	++++	++++
Compound I-8	++++	++++
Compound I-9	+++	++++
Compound I-10	++++	++++
Compound I-12	++++	++++
Compound I-13	++++	++++
Compound I-14	++++	++++
Compound I-15	++++	++++
Compound I-16	++++	++++
Compound I-17	++++	++++
Compound I-18	++++	++++
Compound I-19	++++	++++
Compound I-20	++++	++++
Compound I-21	++++	++++
Compound I-22	++++	++++
Compound I-24	++++	++++
Compound I-25	++++	++++
Compound I-27	++++	++++
Compound I-28	++++	++++
Compound I-31	++++	++++
Compound I-32	++++	++++
Compound I-33	++++	++++
Compound I-34	++++	++++
Compound I-35	++++	++++
Compound I-36	++++	++++
Compound I-37	++++	++++
Compound I-38	++++	++++
Compound I-39	++++	++++

Compound I-40	++++	++++
Compound I-41	++++	++++
Compound I-42	++++	++++
Compound I-43	++++	++++
Compound I-44	++++	++++
Compound I-45	++++	++++
Compound I-46	++++	++++
Compound I-47	++++	++++
Compound I-48	++++	++++

Assay: Mouse Strains

[0433] Lgr5-EGFP-IRES-Cre-ER mice (Barker et al., 2007)

(<http://jaxmice.jax.org/strain/008875.html>) are used to analyze the effects of small molecules on cochlear stem cell expansion.

[0434] Isolation of stem cells from the inner ear: All animal studies are conducted under an approved institutional protocol according to National Institutes of Health guidelines. For experiments with neonatal mice (postnatal days 1-3), the cochleae are dissected in HBSS and the organ of Corti are separated from the stria vascularis and the modiolus. The organs of Corti are then treated with Cell Recovery Solution (Corning) for 1 h to separate cochlear epithelium from the underlying mesenchyme. Epithelia are then collected and treated with TrypLE (Life Technologies) for 15-20 minutes at 37° C. Single cells obtained by mechanical trituration are filtered (40 µm) and suspended in Matrigel (Corning) for 3D culture.

Expansion of Lgr5-Positive Cells

[0435] Cells are cultured in a 1:1 mixture of DMEM and F12, supplemented with Glutamax (GIBCO), N2, B27 (Invitrogen), EGF (50 ng/ml; Chemicon), bFGF (50 ng/ml; Chemicon), IGF1 (50 ng/ml; Chemicon) and the composition provided herein. Media are changed every other day.

[0436] Differentiation of Lgr5-Positive Progenitor Cells Stem cell colonies are differentiated in a 1:1 mixture of DMEM and F12, supplemented with Glutamax (GIBCO), N2, B27 (Invitrogen), with addition of specific drugs or after removal of growth factors without drug addition. Small molecules are added to the culture to test their effect on differentiation.

Analysis

[0437] Lgr5-positive cells are quantified after 10 days (D10) in culture in multiple conditions. Cell colonies are dissociated into single cells using TrypLE (Gibco). The cells are then stained with propidium iodide (PI) and are analyzed using a flow cytometer for Lgr5-GFP expression. The number of GFP-positive cells and the percentage of GFP-positive cells are quantified.

[0438] Atoh1-nGFP-positive cells are quantified at day 0 (D0) and day 10 (D10) of differentiation treatment to determine the number of hair cells that have differentiated. Cell colonies are incubated in Cell Recovery Solution to release the colonies from Matrigel and dissociated into single cells using TrypLE. The total number and percentage of GFP-positive cells are quantified using a flow cytometer for multiple culture conditions. ANOVA is used to compare means across conditions, and the two-tailed Student's T-test is used to compare each condition to the treatment with the highest yield.


[0439] The table herein shows activity of certain compounds. For potency, “+” indicates greater than about 1 μ M; “++” indicates about 100 nM to 1 μ M; and “+++” indicates about 1 nM to 100 nM. For % of cells, “+” indicates about 0-5%; “++” indicates about 6-10%; and “+++” indicates greater than about 11%.

Compound	Potency	% of Cells
Compound I-1	+++	+++
Compound I-2	+++	+++
Compound I-3	+++	+++
Compound I-4	++	+++
Compound I-6	+++	+++
Compound I-7	+++	++
Compound I-8	+++	+++
Compound I-9	+	+++
Compound I-10	+	+
Compound I-12	+++	+++
Compound I-13	+	+++
Compound I-14	+	+
Compound I-15	+	++
Compound I-16	++	+++

Compound I-17	+	+
Compound I-19	+	+++
Compound I-20	+	+++
Compound I-21	+	+++
Compound I-22	+++	++
Compound I-24	+++	++
Compound I-27	+++	+++
Compound I-31	+++	+++
Compound I-32	+	+++
Compound I-33	++	+++
Compound I-34	+	++
Compound I-35	+	+
Compound I-37	+	+
Compound I-38	+	+
Compound I-40	+	+
Compound I-41	+	+++
Compound I-41	+	+++
Compound I-43	++	+++
Compound I-44	++	+++
Compound I-48	+++	++

CLAIMS

1. A compound of Formula (I):



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

Q¹ is CH;


Q² is N;

Q³ is C;

R¹ is selected from the group consisting of hydrogen and halo;

R² is halo;

R³ is selected from the group consisting of hydrogen and halo;



-Z-W-X-Y- is -C(R^Z)₂-C(R^W)₂-N(R^X)-C(R^Y)₂-;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Z groups together form C₃-C₆cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^W groups together form C₃-C₆cycloalkyl or oxo;

R^X is -COR^{X1};

wherein R^{X1} is either:

- (i) 2,8-diazaspiro[4,5]decane, 2,5-diazabicyclo[2,2,1]heptane, or 8-oxa-3-azabicyclo[3.2.1]octane, each of which is optionally substituted with one to twelve substituents independently selected from the group consisting of

deuterium, halo, C₁-C₄alkyl, -[C(RX^{1a})₂]_p-OH, -(CH₂)_p-NMe₂, -(CH₂)_p-NHMe, -(CH₂)_p-NH₂; wherein p is 0, 1, 2, or 3; or


- (ii) piperidine, 2,8-diazaspiro[4,5]decane, 2,5-diazabicyclo[2,2,1]heptane, or 8-oxa-3-azabicyclo[3.2.1]octane, each of which is substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, C₁-C₄alkyl, -[C(RX^{1a})₂]_p-OH, -(CH₂)_p-NMe₂, -(CH₂)_p-NHMe, -(CH₂)_p-NH₂; wherein p is 0, 1, 2, or 3;

wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, -CF₃, and C₁-C₄alkyl, or both R^{X1a} groups together form C₃-C₆cycloalkyl;

each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Y groups together form C₃-C₆cycloalkyl or oxo; and

m is 0.

2. A compound of Formula (I):



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

Q¹ is CH;


Q² is N;

Q³ is C;

R¹ is halo;

R² is halo;

R³ is selected from the group consisting of hydrogen and halo;



-Z-W-X-Y- is -C(R^Z)₂-C(R^W)₂-N(R^X)-C(R^Y)₂-;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Z groups together form C₃-C₆cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^W groups together form C₃-C₆cycloalkyl or oxo;

R^X is -COR^{X1};


wherein R^{X1} is piperidine, 2,8-diazaspiro[4,5]decane, 2,5-diazabicyclo[2,2,1]heptane, or 8-oxa-3-azabicyclo[3.2.1]octane, each of which is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, C₁-C₄alkyl, -[C(RX^{1a})₂]_p-OH, -(CH₂)_p-NMe₂, -(CH₂)_p-NHMe, -(CH₂)_p-NH₂; wherein p is 0, 1, 2, or 3;

wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, -CF₃, and C₁-C₄alkyl, or both R^{X1a} groups together form C₃-C₆cycloalkyl;

each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Y groups together form C₃-C₆cycloalkyl or oxo; and

m is 0.

3. A compound of Formula (I):



or a pharmaceutically acceptable salt or tautomer thereof, wherein:

Q¹ is CH;


Q² is N;

Q³ is C;

R¹ is selected from the group consisting of hydrogen and halo;

R² is selected from the group consisting of -CF₃, -CN, -C≡CH, and -NH₂;

R³ is selected from the group consisting of hydrogen and halo;



Ar is selected from the group consisting of

-Z-W-X-Y- is $-C(R^Z)_2-C(R^W)_2-N(R^X)-C(R^Y)_2-$;

each R^Z is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Z groups together form C₃-C₆cycloalkyl or oxo;

each R^W is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^W groups together form C₃-C₆cycloalkyl or oxo;

or R^Z and R^W together with the carbons to which they are attached form a C₃-C₆cycloalkyl;

R^X is $-COR^{X1}$;

wherein R^{X1} is piperidine, 2,8-diazaspiro[4,5]decane, 2,5-diazabicyclo[2,2,1]heptane, or 8-oxa-3-azabicyclo[3.2.1]octane, each of which is optionally substituted with one to twelve substituents independently selected from the group consisting of deuterium, halo, C₁-C₄alkyl, $-[C(RX^{1a})_2]_p-OH$, $-(CH_2)_p-NMe_2$, $-(CH_2)_p-NHMe$, $-(CH_2)_p-NH_2$; wherein p is 0, 1, 2, or 3;

wherein each R^{X1a} is independently selected from the group consisting of hydrogen, deuterium, halo, $-CF_3$, and C₁-C₄alkyl, or both R^{X1a} groups together form C₃-C₆cycloalkyl;

each R^Y is independently selected from the group consisting of hydrogen, deuterium, halo, and C₁-C₄alkyl, or both R^Y groups together form C₃-C₆cycloalkyl or oxo; and

m is 0.

4. The compound of any one of claim 1-3, wherein each R^Z is independently selected from the group consisting of hydrogen and halo.







5. The compound of any one of claim 1-3, wherein both R^Z groups together form C₃-C₆cycloalkyl.

6. The compound of any one of claims 1-3, wherein R^Z and R^W together with the carbons to which they are attached form a C₃-C₆cycloalkyl.

7. The compound of any one of claims 1-3, wherein each R^W is independently selected from the group consisting of hydrogen and halo.

8. The compound of any one of claims 1-3, wherein both R^W groups together form C₃-C₆cycloalkyl.
9. The compound of any one of claims 1-8, wherein each R^Y is independently selected from the group consisting of hydrogen and halo.
10. The compound of any one of claims 1-8, wherein both R^Y groups together form C₃-C₆cycloalkyl.
11. A compound selected from the group consisting of






Compound I-1	 ;
Compound I-2	 ;
Compound I-3	 ;
Compound I-4	 ;






Compound I-5	
Compound I-6	
Compound I-7	
Compound I-8	
Compound I-9	
Compound I-10	






Compound I-17	 ;
Compound I-18	 ;
Compound I-19	 ;
Compound I-20	 ;
Compound I-21	 ;
Compound I-22	 ;

Compound I-23	 ;
Compound I-24	 ;
Compound I-25	 ;
Compound I-26	 ;
Compound I-27	 ;

Compound I-28	<chem>C1CCN(C1)C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;
Compound I-29	<chem>C1CCN(C1)C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;
Compound I-30	<chem>C1CCN(C1)C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;
Compound I-31	<chem>CC1(C(F)(F)F)CCN1C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;
Compound I-32	<chem>CN(C)CC1CCN(C1)C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;
Compound I-33	<chem>CN(C)CC1CCN(C1)C(=O)N2CCN(C2)CC3=CN4C(=O)NC(=O)C4=C3C5=CC=NC6=C5N=CN=C6</chem> ;

Compound I-34	 <p>Chemical structure of Compound I-34: A central benzimidazole ring is substituted at the 2-position with a benzimidazole ring and at the 5-position with a 1H-imidazole-2,4-dione ring. The 4-position of the benzimidazole is substituted with a 7-fluoro-1,2,3,4-tetrahydroquinoline ring. The nitrogen at the 2-position of the tetrahydroquinoline is connected via a methylene group to a nitrogen atom, which is further connected to a piperidine ring substituted with an amino group (-NH₂).</p>
Compound I-35	 <p>Chemical structure of Compound I-35: Similar to Compound I-34, but the piperidine ring is substituted with a methylamino group (-NHCH₃).</p>
Compound I-36	 <p>Chemical structure of Compound I-36: Similar to Compound I-35, but the piperidine ring is substituted with a dimethylamino group (-N(CH₃)₂).</p>
Compound I-37	 <p>Chemical structure of Compound I-37: Similar to Compound I-34, but the piperidine ring is substituted with a secondary amine group (-NH-), which is further connected to a methylene group and another piperidine ring.</p>
Compound I-38	 <p>Chemical structure of Compound I-38: Similar to Compound I-37, but the piperidine ring is substituted with a tertiary amine group (-N-), which is further connected to a methylene group and another piperidine ring.</p>

Compound I-39	
Compound I-40	
Compound I-41	
Compound I-42	
Compound I-43	

Compound I-44	
Compound I-45	
Compound I-46	
Compound I-47	
Compound I-48	

or a pharmaceutically acceptable salt thereof.

12. A pharmaceutical composition comprising a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent, or excipient.

13. The pharmaceutical composition of claim 12, further comprising an HDAC inhibitor, a TGF beta inhibitor, or a BMP inhibitor.
14. The pharmaceutical composition of claim 12 or 13, further comprising a poloxamer.
15. A method of expanding a population of cochlear cells in a cochlear tissue, the method comprising contacting the cochlear tissue with a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof or a pharmaceutical composition of any one of claims 12-14.
16. A method of facilitating the generation of tissue cells, the method comprising administering to a stem cell population a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof or a pharmaceutical composition of any one of claims 12-14.
17. A method of treating a subject who has, or is at risk of developing, a disease associated with absence or lack of cochlear cells or inner ear hair cells, the method comprising administering to a stem cell population a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof or a pharmaceutical composition of any one of claims 12-14.
18. A method of treating a subject who has, or is at risk of developing, hearing loss, the method comprising administering to the subject a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof or a pharmaceutical composition of any one of claims 12-14, wherein the hearing loss is sensorineural hearing loss, sudden hearing loss, or tinnitus.
19. The method of claim 18, wherein the compound is administered transtympanically to a cochlear tissue of the subject.
20. A method comprising administering a compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof, alone or in combination with an HDAC inhibitor to a cell

population of Lgr5+ cells, wherein the administration results in an expanded population of Lgr5+ cells.

21. A method comprising administering a compound of any one of claims 1-11 or a pharmaceutically acceptable salt thereof, alone or in combination with an HDAC inhibitor, to a population of stem cells in an intestinal epithelium, wherein the administration results in an expanded population of stem cells in the intestinal epithelium.

22. A method comprising administering a compound of any one of claims 1-11 or a pharmaceutically acceptable salt thereof, alone or in combination with an HDAC inhibitor, to a population of Lgr5+ cells in an intestinal epithelium, wherein the administration results in an expanded population of Lgr5+ cells in the intestinal epithelium.

23. A method of proliferating Lgr5+ epithelial cells in vivo, the method comprising administering to a cell population a compound of any one of claims 1-11 or a pharmaceutically acceptable salt thereof.

24. A method for expanding a population of vestibular cells in a vestibular tissue, the method comprising contacting the vestibular tissue with (i) a compound of any one of claims 1-11 or a pharmaceutically acceptable salt thereof, and (ii) a TGF- β inhibitor, wherein the contacting results in an expanded population of vestibular cells in the vestibular tissue.

25. A compound of any one of claims 1-11, or a pharmaceutically acceptable salt, for use in treating a subject who has, or is at risk of developing, a disease associated with absence or lack of cochlear cells or inner ear hair cells.

26. A compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof, for use in treating a subject who has, or is at risk of developing, hearing loss, wherein the hearing loss is sensorineural hearing loss, sudden hearing loss, or tinnitus.

27. Use of a compound of any one of claims 1-11, or a pharmaceutically acceptable salt, in the manufacture of a medicament for treating a subject who has, or is at risk of developing, a disease associated with absence or lack of cochlear cells or inner ear hair cells.

28. Use of a compound of any one of claims 1-11, or a pharmaceutically acceptable salt, in the manufacture of a medicament for treating a subject who has, or is at risk of developing, hearing loss, wherein the hearing loss is sensorineural hearing loss, sudden hearing loss, or tinnitus.