




(86) Date de dépôt PCT/PCT Filing Date: 2004/08/30	(51) Cl.Int./Int.Cl. <i>A61K 38/00</i> (2006.01), <i>C07K 14/00</i> (2006.01)
(87) Date publication PCT/PCT Publication Date: 2005/03/10	
(45) Date de délivrance/Issue Date: 2019/09/10	(72) Inventeurs/Inventors: LADNER, ROBERT CHARLES, US; LEY, ARTHUR C., US; SATO, AARON K., US; STOCHL, MARK, US
(85) Entrée phase nationale/National Entry: 2006/02/24	
(86) N° demande PCT/PCT Application No.: US 2004/028257	(73) Propriétaire/Owner: DYAX CORP., US
(87) N° publication PCT/PCT Publication No.: 2005/021557	
(30) Priorités/Priorities: 2003/08/29 (US60/498,845); 2004/08/04 (US60/598,967)	(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : INHIBITEURS DE PROTEASE POLY-PEGYLEE
(54) Title: POLY-PEGYLATED PROTEASE INHIBITORS



(57) **Abrégé/Abstract:**
Disclosed are compounds that comprise: (i) a Kunitz domain polypeptide that comprises a Kunitz domain that binds to and inhibits a protease; and (ii) a plurality of polyethylene glycol moieties attached to the Kunitz domain polypeptide. Each accessible primary amine of the Kunitz domain polypeptide can be attached to one of the moieties. Also disclosed are related methods.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
10 March 2005 (10.03.2005)


PCT

(10) International Publication Number
WO 2005/021557 A2

- (51) International Patent Classification⁷: **C07F** (US). **STOCHL, Mark** [US/US]; 121 Achilles Way, N. Attleboro, MA 02763 (US).
- (21) International Application Number: PCT/US2004/028257 (74) **Agent: MYERS, Louis**; Fish & Richardson P.C., 225 Franklin Street, Boston, MA 02110-2804 (US).
- (22) International Filing Date: 30 August 2004 (30.08.2004) (81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English (84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (26) Publication Language: English
- (30) Priority Data:
60/498,845 29 August 2003 (29.08.2003) US
60/598,967 4 August 2004 (04.08.2004) US
- (71) Applicant (for all designated States except US): **DYAX CORP.** [US/US]; 300 Technology Square, 8th Floor, Cambridge, MA 02139 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **LADNER, Robert, Charles** [US/US]; 3827 Green Valley Road, Ijamsville, MD 21754 (US). **LEY, Arthur, C.** [US/US]; 122 Adrena Road, Newton, MA 02465 (US). **SATO, Aaron, K.** [US/US]; 11 Pembroke Street, #2, Somerville, MA 02145

[Continued on next page]

(54) Title: POLY-PEGYLATED PROTEASE INHIBITORS



5 10 15 20 25 30 35 40 45 50 55
EACNLPIVRGPCIAFFPRWAFDAVKGKCVLFPYGGCQGNNGKIFYSEKECREYCGVP

(57) **Abstract:** Disclosed are compounds that comprise: (i) a Kunitz domain polypeptide that comprises a Kunitz domain that binds to and inhibits a protease; and (ii) a plurality of polyethylene glycol moieties attached to the Kunitz domain polypeptide. Each accessible primary amine of the Kunitz domain polypeptide can be attached to one of the moieties. Also disclosed are related methods.

WO 2005/021557 A2

WO 2005/021557 A2



Published:

— *without international search report and to be republished
upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

POLY-PEGYLATED PROTEASE INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Application Serial No. 60/498,845, filed on August 29, 2003 and to 60/598,967, filed on August 4, 2004.

5

BACKGROUND

The invention relates to modified protease inhibitors.

SUMMARY

In one aspect, the invention features a compound that include: a) a polypeptide including a Kunitz domain that specifically binds and/or inhibits a protease; and b) a
10 plurality of non-protein moieties that are physically associated with the polypeptide and increases the molecular weight of the compound. The term “poly-PEGylated Kunitz domain” refers herein to the afore-mentioned compound. Typically, the non-protein moieties of a poly-PEGylated Kunitz domain include polyethylene glycol.

The compound (i.e., the polypeptide plus the plurality of non-protein moieties)
15 has a molecular weight of greater than 12, 14 or 16 kDa. In one embodiment, each non-protein moiety has an average molecular weight of between 3 and 20, 3 and 12 kDa, 3 and 10 kDa, 3 and 8 kDa, 4 and 6 kDa, e.g., about 4, 5, 6, 7, or 8 kDa.

The protease that is bound and/or inhibited can be, for example, an elastase (e.g., human neutrophil elastase (hNE)), a plasmin, a kallikrein, or other protease, e.g.,
20 a protease described herein. For example, the protease can be a serine protease.

In one embodiment, non-protein moieties are attached to each available primary amine on the Kunitz domain, e.g., the N-terminal primary amine and any solvent-accessible primary amines, e.g., accessible primary amines of lysine side chains in the Kunitz domain. For example, all possible primary amines are conjugated to one of the
25 non-protein moieties. The Kunitz domain may have at least one, two, three, or four lysines. For example, the Kunitz domain may have only one, two, three, four, or five lysines. In one embodiment, the polypeptide has an N-terminal primary amine. In another embodiment, the polypeptide does not include an N-terminal primary amine (e.g., the polypeptide can be chemically modified, e.g., with a non-polymeric


compound, at its N-terminal primary amine so that the polypeptide does not include a primary amine at that position).

A non-protein moiety can be attached at 2 or more of the primary amines in the polypeptide. For example, all lysines or all lysines that have a solvent accessible
5 primary amine are attached to a non-protein moiety. Preferably, the Kunitz domain does not include a lysine within one of its binding loops, e.g., about residues corresponding to amino acids 11-21 of BPTI and 31-42 of BPTI. Lysines within such binding loops can be replaced, e.g., with arginine residues. For example, the polypeptide is attached to at least three of molecules of the polymer. Each lysine of the
10 polypeptide, or one, two, three or more of the lysines can be attached to a molecule of the polymer. Unless otherwise stated, when it is said that a primary amine, e.g., that of a particular lysine or at the N terminus, is modified or has a non-protein moiety attached thereto, it is understood that the specified primary amine position on every molecule in a preparation may not be so modified. The preparations need not be
15 perfectly homogeneous to be within the invention. Homogeneity is desirable in some embodiments but it need not be absolute. In preferred embodiments, at least 60, 70, 80, 90, 95, 97, 98, 99, or 100% of a primary amine which is designated as modified will have a non-protein moiety attached thereto. Other embodiments however, include preparations that contains a mixture of species in which most of the molecules, e.g., at
20 least 60, 70, 80, 90, 95, 97, 98, 99, or 100% are PEGylated at two or more sites but the sites (and in some cases the number of sites modified) on molecules in the preparation will vary. E.g., some molecules will have lysines A, B, and D modified while other molecules will have the amino terminus and lysines A, B, C, and D modified.

In one embodiment, the non-protein moiety includes a hydrophilic polymer,
25 e.g., a substantially homogeneous polymer. The polymer can be branched or unbranched. For example, the moiety of polymer has a molecular weight (e.g., an average molecular weight of the moieties added to the compound) that is less than 20, 18, 15, 12, 10, 8, 7, or 6 or at least 1.5, 2, 2.5, 3, 5, 6, 10 kDa, e.g., about 5 kDa. In one embodiment, the sum of the molecular weight of the PEG moieties on the compound is
30 at least 15, 20, 25, 30, or 35, and/or less than 60, 50, 40, 35, 30, 25, or 23 kDa.

In one embodiment, the polymer is a polyalkylene oxide. For example, at least 20, 30, 50, 70, 80, 90, or 95% of the copolymer blocks of the polymer are ethylene glycol. In one embodiment, the polymer is polyethylene glycol.

In one embodiment, the compound has the following structure:



wherein P is the polypeptide,

each of R' and R'' is, independently, H, or C₁-C₁₂ alkyl ;

5 X⁰ is O, N-R¹, S, or absent, wherein R¹ is H, C₁-C₁₂ alkyl or aryl,

X¹ is O, N-R², S, wherein R² is H, alkyl or aryl,

X² is O, N-R³, S, or absent, wherein R³ is H, alkyl or aryl,

each n is between 1 and 5, e.g., 2,

a is at least 4,

10 m is between 0 and 5, and

R^t is H, C₁-C₁₂ alkyl or aryl.

R' and R'' can be H. In one embodiment, R' or R'' is independently, H, or C₁-C₄, C₁-C₆, or C₁-C₁₀ alkyl.

In one embodiment, the compound has the following structure:



wherein P is the polypeptide,

a is at least 4,

m is between 0 and 5,

X² is O, N-R¹, S, or absent, wherein R¹ is H, alkyl or aryl,

20 X⁰ is O, N-R², S, or absent, wherein R² is H, alkyl or aryl, and

R^t is H, C₁-C₁₂ alkyl or aryl. For example, X² is O, and R^t is CH₃. (The use of mPEG is preferred.)

In one embodiment, the Kunitz domain polypeptide is less than 14, 8, or 7 kDa in molecular weight. In one embodiment, the Kunitz domain polypeptide includes only one Kunitz domain. Generally, the compound includes only one Kunitz domain, but in
25 some embodiments, may include more than one.

In one embodiment, the Kunitz domain includes the amino acid sequence of DX-890, DX-88, or DX-1000 or an amino acid sequence that differs by at least one, but fewer than six, five, four, three, or two amino acid differences (e.g., substitutions,
30 insertions, or deletions) from the amino acid sequence of DX-890, DX-88, or DX-1000. Typically, the Kunitz domain does not naturally occur in humans. The Kunitz domain may include an amino acid sequence that differs by fewer than ten, seven, or four amino acids from a human Kunitz domain.

In one embodiment, the K_i of the compound is within a factor of 0.5 to 1.5, 0.8 to 1.2, 0.3 to 3.0, 0.1 to 10.0, or 0.02 to 50.0 of the K_i of the unmodified polypeptide for elastase. For example, the K_i for hNE can be less than 100, 50, 18, 12, 10, or 9 pM.

In one embodiment, the compound has a circulatory half life of the longest-lived component ("longest phase circulatory half life") in a rabbit or mouse model that is at least 1.5, 2, 4, or 8 fold greater than a substantially identical compound that does not include the polymer. The compound can have a longest phase circulatory half life in a rabbit or mouse model that has an amplitude at least 1.5, 2, 2.5, or 4 fold greater than a substantially identical compound that does not include the non-protein moiety.

10 The compound can have an alpha-phase circulatory half life in a rabbit or mouse model that has an amplitude at least 20, 30, 40, or 50% smaller than a substantially identical compound that does not include the non-protein moiety. For example, the compound has a longest phase circulatory half life with an amplitude of at least 40, 45, 46, 50, 53, 54, 60, or 65%. In one embodiment, the compound has a beta phase circulatory half

15 life in a mouse or rabbit model of at least 2, 3, 4, 5, 6, or 7 hours. In one embodiment, the compound has a longest phase circulatory half life in a 70 kg human of at least 6 hours, 12 hours, 24 hours, 2 days, 5 days, 7 days, or 10 days.

In one embodiment, the compound has a longest phase circulatory half life in a rabbit model of at least 4200 minutes, 4700 minutes, or 4980 minutes (or about 83 hours). In

20 one embodiment, the compound has a longest phase circulatory half-life that is longer than a similarly sized molecule with the same Kunitz domain, but only a single PEG moiety (i.e., a mono-PEGylated version of the same Kunitz domain). The longest phase half-life can be at least 5, 10, 20, 30, or 50% longer. In one embodiment, in a mouse, the longest phase circulatory half life has an amplitude of greater than 50, 55,

25 60, or 65%. The longest phase half life can be, e.g., greater than 550, 600, 700, 750, 900, 1000, 1100 minutes.

In one embodiment, the compound has increased solubility (e.g., 1.5, 2, 4, or 8 fold greater) in an aqueous solution having a pH between 5 and 8 and an ionic strength less than the ionic strength of 0.5 M NaCl than the polypeptide that does not include the

30 non-protein moiety.

In one embodiment, the polyethylene glycol is attached by coupling monomethoxy-PEG propionaldehyde or monomethoxy-PEG succinimidyl propionic acid to the polypeptide. The compound can be formed by coupling of mPEG ($\text{CH}_3\text{-(O-}$

CH₂-CH₂)_n-) at a pH that enables attachment to accessible amino groups on lysine side chains and to the N-terminal amino group, e.g., a pH 6.8 to 8.8, e.g., between 7.4 and 8.8.

In another aspect, the invention features a compound that includes (1) a
5 polypeptide including the amino acid sequence of DX-890, DX-88, or DX-1000 or an amino acid sequence that differs by at least one, but fewer than six, five, four, three, or two amino acid differences (e.g., substitutions, insertions, or deletions) from the amino acid sequence of DX-890, DX-88, or DX-1000, and (2) a plurality of polyethylene glycol moieties. Each polyethylene glycol moiety can be less than 20, 19, 18, 15, 12,
10 11, 10, 9, 8, 7, or 6 kDa in molecular weight and is attached. Each polyethylene moiety can be attached to the polypeptide by a single covalent bond.

In one embodiment, a molecule of polyethylene glycol is attached to each lysine side chain of the polypeptide, e.g., where the polypeptide includes more than one lysine, e.g., two, three or four lysines. For example, the polypeptide is identical to the
15 amino acid sequence of DX-890 and a molecule of polyethylene glycol is attached to each of the four lysine side chains of DX-890, and optionally also to the N-terminus. In another example, the polypeptide is identical to the amino acid sequence of DX-88 or DX-1000 and a molecule of polyethylene glycol is attached to each of the three lysine side chains of DX-88 or DX-1000, and, optionally, also to the N-terminus. In
20 one embodiment, the molecules of polyethylene glycol are between 4 and 12 kDa in molecular weight. In one embodiment, the polyethylene glycol is attached to the N-terminus and to each accessible lysine side chain.

In one embodiment, the amino acid sequence differs by at least one amino acid from the amino acid sequence of DX-890. The amino acid sequence is identical to the
25 amino acid sequence of DX-890 at one or more positions (e.g., at least two, three, five, seven, ten, twelve, thirteen, fourteen, or all) corresponding to positions 5, 13, 14, 16, 17, 18, 19, 30, 31, 32, 34, 38, 39, 51, and 55 according to the BPTI numbering.

In another aspect, the invention features a preparation that comprises Kunitz domain polypeptides that specifically bind and inhibit a protease. At least 40, 50, 70,
30 80, 85, 90, 92, 95, 97, 98, 99, or 99.5% of the Kunitz domain polypeptides in the preparation (i) bind and inhibit the protease, and (ii) have a polyethylene glycol moiety attached at a first common site and a polyethylene glycol moiety attached at a second common site. Typically, the average molecular weight of each attached polyethylene

glycol moiety is less than 12, 10, or 8 kDa. In one embodiment, the designated population of Kunitz domain polypeptides further have a polyethylene glycol moiety attached at the third common site and a polyethylene glycol moiety attached at the fourth common site. For example, the designated population of Kunitz domain polypeptides have a polyethylene glycol moiety attached to each accessible primary amine and/or an N-terminal primary amine.

In one embodiment, each of the Kunitz domain polypeptides in the preparation binds and inhibits the protease. For example, Kunitz domain polypeptides that are not members of the designated population also bind and inhibit a protease, e.g., the same or a different protease.

The Kunitz domain polypeptides of the population can include, for example, other features described herein.

The invention also features a preparation that includes a compound described herein, e.g., above. For example, the compound is present at a concentration of at least 0.1, 1, 2, or 5 mg of polypeptide per milliliter, e.g., in a solution between pH 6-8. In one embodiment, the compound produces a major peak by size exclusion chromatography that includes at least 70% the compound relative to the injectate. In one embodiment, the molecular weight of 95% of the species of the compound are within 5, 4, 3, 2, or 1 kDa of the average molecular weight of the compound.

In another aspect, the invention features a pharmaceutical preparation that includes (1) a compound described herein, and (2) a pharmaceutically acceptable carrier. In one embodiment, at least 60, 70, 80, 85, 90, 95, 97, 98, 99, or 100% of the compounds in the preparation have an identical distribution of PEG molecules attached thereto. The use of chemical reaction in which all available primary amines (e.g., all solvent accessible primary amines, or all primary amines) are modified can be used to provide a preparation in which the compounds have an identical distribution of PEG molecules. Of course, some variation will be present in the molecule weight of the moieties attached to different primary amines on a given molecule or among molecules since there is variation about an average molecular weight for the PEG reagent used in the chemical reaction. The preparation can also be made using a process that provides a greater than 25, 30, 40, 50, 60, 70, 75, 80, or 85% yield for input protein.

In one embodiment, the preparation is aqueous and the compound is present at a concentration of at least 0.1 mg of polypeptide per milliliter. In one embodiment,

injection of the preparation into a mouse results in less than 50, 30, 25, 15, or 10% of the compound is an SEC peak with higher mobility than the preparation after 12 hours.

The preparation can be suitable for pulmonary delivery or for gastrointestinal delivery (e.g., ingestion, rectal, etc.).

5 In another aspect, the invention features a pharmaceutical preparation that includes (1) a compound described herein, and (2) a pharmaceutically acceptable carrier. In one embodiment, at least 60, 70, 80, 85, 90, 95, 97, 98, 99, or 100% of the polypeptides in the preparation have at least 2, 3 or 4 primary amines modified with a non-protein moiety. The preparation can contain a mixture of species in which most of
10 the molecules, e.g., at least 60, 70, 80, 90, or 95 % are PEGylated at least 2 (or 3 or 4) sites but the sites (and in some cases the number of sites modified) on molecules in the preparation will vary. E.g., some molecules will have lysines A, B, and D modified while other molecules will have the amino terminus and lysines A, B, C, and D modified. In some embodiments, the preparation can include a small number of
15 compounds that are inactive (e.g., less than 5, 2, 1, or 0.1%), but generally, most of the compounds (e.g., at least 50%, 90, 95, 98, 99, 99.5, or 99.9%) in the preparation are active, e.g., can inhibit a protease.

In some aspects of the invention, the non-protein moiety attached to different sites will be the same, in terms of identity or size. In other aspects, a first non-protein
20 moiety is attached at a first primary amine, and a second non-protein moiety which is different, e.g., by type or size, is attached to a different primary amine. E.g., it may be desirable to attach a PEG of a first size to the primary amine of the N terminus but to attach a PEG of a different size to a lysine position.

The invention also features a medical device that includes a dispenser and a
25 compartment that includes a pharmaceutical preparation described herein. For example, the dispenser is configured to generate an inhalable form of the pharmaceutical preparation. The invention also features an implantable medical device that includes a dispenser and a compartment that includes a pharmaceutical preparation described herein wherein the dispenser is configured to delivery the pharmaceutical
30 preparation into the circulatory system of a subject. The invention also features a suppository that includes a pharmaceutical preparation described herein.

In another aspect, the invention features a preparation that includes a poly-pegylated Kunitz domain. The preparation can be substantially (e.g., at least 70, 75, 80,

85, 90, 95, or 100%) monodisperse. For example, the poly-PEGylated compound is present at a concentration of at least 0.05, 0.1, 0.2, 0.5, 0.8, 1.0, 1.5, 2.0, or 2.5 milligrams of polypeptide per milliliter or between 0.05 and 10 milligrams of polypeptide per milliliter. In one embodiment, the preparation is dry. For example, the preparation includes particles or is in the form of a powder.

In another aspect, the invention features a compound that includes a polypeptide including the amino acid sequence of DX-890, DX-88, or DX-1000 or other Kunitz domain sequence described herein in which at least one lysine is substituted with a non-lysine amino acid, e.g., arginine. The compound is useful for reducing the number of lysines to which PEG is coupled, e.g., without a substantial change in activity, to produce a substantially homogenous conjugate. In one embodiment, the amino acid sequence (e.g., of DX-890) has three lysine substitutions and a single remaining lysine. In another embodiment, the amino acid has one or two lysine substitutions. In one embodiment, the compound further includes a non-protein moiety, e.g., a hydrophilic polymer described herein. The polymer can be coupled to the remaining lysine residues, e.g., single remaining lysine (e.g., the first, second, third, or fourth lysine).

In another aspect, the invention features a preparation (e.g., an aqueous preparation) that includes: a compound that includes a Kunitz domain conjugated to a plurality of moieties of a hydrophilic and substantially homogeneous polymer. For example, the concentration of Kunitz domain component alone is greater than 2 mg per ml, the pH of the preparation is greater than 3, and the ionic strength of the preparation is less than the ionic strength of 0.5 M NaCl. In one embodiment, the Kunitz domain includes the amino acid sequence of DX-890, DX-88, or DX-1000 or an amino acid sequence that differs by at least one, but fewer than six, five, four, three, or two amino acid differences (e.g., substitutions, insertions, or deletions) from the amino acid sequence of DX-890. The invention also provides a sealed container that includes the preparation. The container can be opaque to light. The container can include printed information on an external region of the container.

In another aspect, the invention features a method that includes: providing a polypeptide that includes a Kunitz domain; contacting the polypeptide to a hydrophilic polymer (e.g., a polyalkylene oxide) that includes a single reactive group that can form a covalent bond with the polypeptide under conditions suitable for bond formation at a

plurality of available sites (e.g., a plurality of primary amines, e.g., all available primary amines), thereby providing a modified protease inhibitor.

In one embodiment, the hydrophilic polymer is mono-activated. For example, the hydrophilic polymer is alkoxy-terminated. In one embodiment, the polymer
5 includes a succinimidyl group.

In one embodiment, the polymer is a polyethylene glycol, e.g., monomethoxy-polyethylene glycol. For example, the polymer is mPEG propionaldehyde or mPEG succinimidyl propionic acid.

In one embodiment, the conditions are between pH 6.5 and 9.0, e.g., between
10 7.5 and 8.5. In one embodiment, the hydrophilic polymer is covalently attached to the N-terminus of the polypeptide. In another embodiment, the hydrophilic polymer is covalently attached to a lysine side chain of the polypeptide.

The method can further include separating polypeptides that have a single attached polymer from other products and reactants. The method can further include
15 chromatographically separating products of the contacting, e.g., using ion exchange chromatography or size exclusion chromatography.

The invention also features a modified Kunitz domain prepared by a method described herein, e.g., the above method.

In another aspect, the invention features a method of treating a disorder
20 characterized by excessive or undesired activity of a protease. The method includes: administering to a subject having the disorder or suspected of having the disorder to pharmaceutical composition comprising a compound or preparation described herein. The compound or preparation includes a Kunitz domain polypeptide that inhibits the protease. For example, a preparation has at least a certain percentage of molecules of
25 the Kunitz domain polypeptide in which a hydrophilic polymer is attached to a first common site and a second common site. For example, at least a certain percentage of molecules of the Kunitz domain polypeptide further include the hydrophilic polymer attached to a third, fourth, and optionally a fifth common site.

In one embodiment, the protease is elastase. For example, the Kunitz domain
30 polypeptide comprises the amino acid sequence of DX-890 or a sequence that differs by at least one, but fewer than six amino acid alterations from DX-890. Exemplary disorders that can be treated using a Kunitz domain that inhibits elastase (e.g., human neutrophil elastase) include cystic fibrosis, COPD, and an inflammatory disorder.

In one embodiment, the protease is a kallikrein. For example, the Kunitz domain polypeptide comprises the amino acid sequence of DX-88 or a sequence that differs by at least one, but fewer than six amino acid alterations from DX-88.

Exemplary disorders that can be treated using a Kunitz domain that inhibits a kallikrein
5 include disorders of coagulation, fibrinolysis, hypotensions, inflammation, hemophilia, post-operative bleeding, peri-operative bleeding, and hereditary angioedema.

In one embodiment, the protease is plasmin and the Kunitz domain polypeptide comprises the amino acid sequence of DX-1000 or a sequence that differs by at least one, but fewer than six amino acid alterations from DX-1000. Exemplary disorders
10 that can be treated using a Kunitz domain that inhibits plasmin include fibrinolysis or fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis.

In another aspect, the invention features a method of treating or preventing a pulmonary disorder. The method includes administering a compound described herein
15 to a subject, e.g., in an amount effective to ameliorate at least one symptom of the disorder. For example, the compound includes a) a polypeptide including a Kunitz domain that specifically binds and inhibits an elastase (e.g., human neutrophil elastase (hNE)); and b) a non-protein moiety that is physically associated with the polypeptide and increases the molecular weight of the compound. For example, the compound
20 includes (1) a polypeptide including the amino acid sequence of DX-890 or an amino acid sequence that differs by at least one, but fewer than six, five, four, three, or two amino acid differences (e.g., substitutions, insertions, or deletions) from the amino acid sequence of DX-890, and (2) polyethylene glycol wherein the sum of the polyethylene glycol moieties is at least 15, 18, 20, 25, 27, or 30 kDa in molecular weight.

25 In one embodiment, the compound is administered no more than once every 12, 24, 36, or 72 hours. In another embodiment, the compound is administered no more than once every four, seven, ten, twelve, or fourteen days. The compound can be administered once or at multiple times (e.g., regularly).

In one embodiment, the administering includes pulmonary delivery. For
30 example, the administering includes actuation of an inhaler and/or nebulization. In one embodiment, the administering includes delivery of the composition directly or indirectly into the circulatory system. For example, the administering includes injection or intravenous delivery.

In one embodiment, the subject has cystic fibrosis or a genetic defect in the cystic fibrosis gene. In another embodiment, the subject has chronic obstructive pulmonary disease.

The symptom can be lung tissue integrity or an index of tissue destruction.

5 In another aspect, the invention features a method of treating or preventing a inflammatory disorder. The method includes: administering a compound described herein to a subject, e.g., in an amount effective to ameliorate at least one symptom of the disorder. For example, the compound includes a) a polypeptide including a Kunitz domain that specifically binds and inhibits an elastase (e.g., human neutrophil elastase
10 (hNE)); and b) a plurality of non-protein moieties that are physically associated with the polypeptide and increase the molecular weight of the compound. For example, the compound includes (1) a polypeptide including the amino acid sequence of DX-890 or an amino acid sequence that differs by at least one, but fewer than six, five, four, three, or two amino acid differences (e.g., substitutions, insertions, or deletions) from the
15 amino acid sequence of DX-890, and (2) a plurality of polyethylene glycol moieties, e.g., wherein each polyethylene glycol moiety is less than 20, 18, 16, 12, 10, 9, 8, or 7 kDa in molecular weight.

In one embodiment, the disorder is an inflammatory bowel disorder, e.g., Crohn's disease or ulcerative colitis. In one embodiment, the compound is delivered by
20 a suppository.

In one embodiment, the compound is administered no more than once every 12, 24, 36, or 72 hours. In another embodiment, the compound is administered no more than once every four, seven, ten, twelve, or fourteen days. The compound can be administered once or at multiple times (e.g., regularly).

25 In another aspect, the invention features a method of treating or preventing a disorder characterized at least in part by inappropriate elastase activity or neutrophil activity. The method includes administering a compound described herein to a subject, e.g., in an amount effective to ameliorate at least one symptom of the disorder or to alter elastase or neutrophil activity, e.g., to reduce elastase-mediated proteolysis. For
30 example, the disorder is rheumatoid arthritis.

In one embodiment, the compound is administered no more than once every 12, 24, 36, or 72 hours. In another embodiment, the compound is administered no more

than once every four, seven, ten, twelve, or fourteen days. The compound can be administered once or at multiple times (e.g., regularly).

Many of the examples provided herein describe methods and compositions that relate to Kunitz domains and a particular protease target – elastase. However, these methods and compositions can be modified to provide corresponding methods and compositions that relate to other targets, e.g., other proteases or other proteins, e.g., protease other than Kunitz domains, particularly proteins that include one or more lysine residues. For example, the lysines may be positioned at a site where their modification does not interfere with function. Similarly the described methods and compositions can be modified to corresponding methods and compositions that relate to polypeptides that do not include a Kunitz domain or that include a Kunitz domain and other types of domains.

As used herein, “binding affinity” refers to the apparent association constant or K_a . The K_a is the reciprocal of the dissociation constant (K_d). A ligand may, for example, have a binding affinity of at least 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , or 10^{12} M^{-1} for a particular target molecule. Higher affinity binding of a ligand to a first target relative to a second target can be indicated by a higher K_a (or a smaller numerical value K_d) for binding the first target than the K_a (or numerical value K_d) for binding the second target. In such cases the ligand has specificity for the first target relative to the second target. K_a measurements for binding to hNE are typically made under the following conditions: 50 mM HEPES, pH 7.5, 150 mM NaCl, and 0.1% Triton X-100 at 30°C using 100 pM of the hNE.

Binding affinity can be determined by a variety of methods including equilibrium dialysis, equilibrium binding, gel filtration, ELISA, surface plasmon resonance, or spectroscopy (e.g., using a fluorescence assay). These techniques can be used to measure the concentration of bound and free ligand as a function of ligand (or target) concentration. The concentration of bound ligand ($[\text{Bound}]$) is related to the concentration of free ligand ($[\text{Free}]$) and the concentration of binding sites for the ligand on the target where (N) is the number of binding sites per target molecule by the following equation:

$$[\text{Bound}] = N \cdot [\text{Free}] / ((1/K_a) + [\text{Free}])$$

It is not always necessary to make an exact determination of K_a , though, since sometimes it is sufficient to obtain a quantitative measurement of affinity, e.g.,

determined using a method such as ELISA or FACS analysis, is proportional to K_a , and thus can be used for comparisons, such as determining whether a higher affinity is, e.g., 2 fold higher.

An "isolated composition" refers to a composition that is removed from at least 5 90% of at least one component of a natural sample from which the isolated composition can be obtained. Compositions produced artificially or naturally can be "compositions of at least" a certain degree of purity if the species or population of species of interests is at least 5, 10, 25, 50, 75, 80, 90, 95, 98, or 99% pure on a weight-weight basis.

An "epitope" refers to the site on a target compound that is bound by a ligand, 10 e.g., a polypeptide ligand such as a Kunitz domain, small peptide, or antibody. In the case where the target compound is a protein, for example, an epitope may refer to the amino acids that are bound by the ligand. Such amino acids may be contiguous or non-contiguous with respect to the underlying polypeptide backbone. Overlapping epitopes include at least one common amino acid residue.

15 As used herein, the term "substantially identical" (or "substantially homologous") is used herein to refer to a first amino acid or nucleotide sequence that contains a sufficient number of identical or equivalent (e.g., with a similar side chain, e.g., conserved amino acid substitutions) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or 20 nucleotide sequences have similar activities. In the case of Kunitz domains, the second domain has the same specificity and, for example, has at least 0.5, 5, or 50% of the binding affinity of the first domain. A sufficient degree of identity may be about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or higher.

Sequences similar or homologous (e.g., at least about 85% sequence identity) to 25 the sequences disclosed herein are also part of this application. In some embodiment, the sequence identity can be about 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or higher. Alternatively, substantial identity exists when the nucleic acid segments will hybridize under selective hybridization conditions (e.g., highly stringent hybridization conditions), to the complement of the strand. The nucleic acids may be 30 present in whole cells, in a cell lysate, or in a partially purified or substantially pure form.

Calculations of "homology" or "sequence identity" between two sequences (the terms are used interchangeably herein) are performed as follows. The sequences are

aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position (as used herein amino acid or nucleic acid "identity" is equivalent to amino acid or nucleic acid "homology"). The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch ((1970) *J. Mol. Biol.* 48:444-453) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package, using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used if the practitioner is uncertain about what parameters should be applied to determine if a molecule is within a sequence identity or homology limitation of the invention) are a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

As used herein, the term "homologous" is synonymous with "similarity" and means that a sequence of interest differs from a reference sequence by the presence of one or more amino acid substitutions (although modest amino acid insertions or

deletions) may also be present. Presently preferred means of calculating degrees of homology or similarity to a reference sequence are through the use of BLAST algorithms (available from the National Center of Biotechnology Information (NCBI), National Institutes of Health, Bethesda MD), in each case, using the algorithm default or recommended parameters for determining significance of calculated sequence relatedness. The percent identity between two amino acid or nucleotide sequences can also be determined using the algorithm of E. Meyers and W. Miller ((1989) *CABIOS*, 4:11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

As used herein, the term "hybridizes under low stringency, medium stringency, high stringency, or very high stringency conditions" describes conditions for hybridization and washing. Guidance for performing hybridization reactions can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Aqueous and nonaqueous methods are described in that reference and either can be used. Specific hybridization conditions referred to herein are as follows: 1) low stringency hybridization conditions in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by two washes in 0.2X SSC, 0.1% SDS at least at 50°C (the temperature of the washes can be increased to 55°C for low stringency conditions); 2) medium stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C; 3) high stringency hybridization conditions in 6X SSC at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C; and preferably 4) very high stringency hybridization conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C. Very high stringency conditions (4) are the preferred conditions and the ones that should be used unless otherwise specified. Accordingly, nucleic acids that hybridize with appropriate stringency to nucleic acids that encode a polypeptide described herein are provided as are polypeptides that are encoded by such nucleic acids. Such polypeptides can be similarly modified as described herein.

It is understood that a polypeptide described herein (e.g., a polypeptide that includes a Kunitz domain) may have mutations relative to a particular polypeptide described herein (e.g., a conservative or non-essential amino acid substitutions), which

do not have a substantial effect on the polypeptide functions. Whether or not a particular substitution will be tolerated, i.e., will not adversely affect desired biological properties, such as binding activity can be determined as described in Bowie, et al. (1990) *Science* 247:1306-1310. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). It is possible for many framework and CDR amino acid residues to include one or more conservative substitutions.

15 A “non-essential” amino acid residue is a residue that can be altered from the wild-type sequence of the binding agent, e.g., the antibody, without abolishing or more preferably, without substantially altering a biological activity, whereas an “essential” amino acid residue results in such a change.

The terms “polypeptide” or “peptide” (which may be used interchangeably) refer to a polymer of three or more amino acids linked by a peptide bond, e.g., between 3 and 30, 12 and 60, or 30 and 300, or over 300 amino acids in length. The polypeptide may include one or more unnatural amino acids. Typically, the polypeptide includes only natural amino acids. A “protein” can include one or more polypeptide chains. Accordingly, the term “protein” encompasses polypeptides. A protein or polypeptide can also include one or more modifications, e.g., a glycosylation, amidation, phosphorylation, and so forth. The term “small peptide” can be used to describe a polypeptide that is between 3 and 30 amino acids in length, e.g., between 8 and 24 amino acids in length.

30 The term “alkyl” refers to a hydrocarbon chain that may be a straight chain or branched chain, containing the indicated number of carbon atoms. For example, C₁-C₁₂ alkyl indicates that the group may have from 1 to 12 (inclusive) carbon atoms in it.

The term “aryl” refers to an aromatic monocyclic, bicyclic, or tricyclic hydrocarbon ring system, wherein any ring atom capable of substitution can be

50860-189

substituted by a substituent. Examples of aryl moieties include, but are not limited to, phenyl, naphthyl, and anthracenyl.

In one aspect, the invention provides a compound comprising: (i) a polypeptide that comprises a Kunitz domain that binds to and inhibits a protease, wherein the Kunitz domain is selected from the group consisting of: a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-88 amino acid sequence set forth in SEQ ID NO:24; and (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25; and (ii) a plurality of polyethylene glycol moieties, wherein the average molecular weight of each of the polyethylene glycol moieties is less than 12 kDa, and wherein each of at least four primary amines of the polypeptide is attached to one of the moieties of the plurality.

In another aspect, the invention provides a preparation that comprises (i) polypeptides that specifically bind and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of: a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-88 amino acid sequence set forth in SEQ ID NO:24; and (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25,

50860-189

wherein the at least 80% of the polypeptides comprise a polyethylene glycol moiety attached to each of at least four primary amines of the polypeptides, and wherein the average molecular weight of each of the attached polyethylene glycol moieties is less than 12 kDa, and (ii) a pharmaceutically acceptable carrier.

5 In another aspect, the invention provides a preparation that comprises (i) polypeptides that specifically bind and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of: a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but not
10 more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-88 amino acid sequence set forth in SEQ ID NO:24; and (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID
15 NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25, and wherein the at least 80% of the polypeptides comprise a polyethylene glycol moiety attached to each of at least five primary amines of the polypeptides and wherein the average molecular weight of each of the attached polyethylene glycol moieties is less
20 than 12 kDa, and (ii) a pharmaceutically acceptable carrier.

 In another aspect, the invention provides a method of providing a pegylated Kunitz domain, the method comprising: providing a polypeptide that comprises a Kunitz domain that has at least one lysine in the framework region of the Kunitz domain, wherein the Kunitz domain is selected from the group consisting of:
25 a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from

50860-189

the DX-88 amino acid sequence set forth in SEQ ID NO:24; and (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25; and contacting the

5 polypeptide with activated polyethylene glycol, of average molecular weight less than 12 kDa, under conditions in which a polyethylene glycol moiety is attached to each of at least four primary amines of the polypeptide, and wherein at least one polyethylene glycol moiety is attached to the N-terminal primary amine.

In another aspect, the invention provides a method of providing a

10 PEGylated Kunitz domain, the method comprising: providing a polypeptide that comprises a Kunitz domain that has at least two primary amine groups in the framework region of the Kunitz domain wherein the Kunitz domain is selected from the group consisting of: (a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but

15 not more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-88 amino acid sequence set forth in SEQ ID

20 SEQ ID NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25; and contacting the polypeptide with activated polyethylene glycol, of average molecular weight less than 12 kDa, under conditions in which a polypethylene glycol moiety is attached to each of at least four primary amines of the polypeptide.

25 In another aspect, the invention provides a composition for treating a disorder selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), an inflammatory disorder, hemophilia, post-operative bleeding, peri-operative bleeding, hereditary angioedema, fibrinolysis or fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative

50860-189

bleeding, peri-operative bleeding, and inappropriate androgenesis, in a subject having or suspected of having the disorder, wherein the composition comprises the preparation as described above, wherein the polypeptide of the preparation inhibits a protease.

- 5 In another aspect, the invention provides a preparation comprising: (i) polypeptides that specifically bind to and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of: a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO:23 or an amino acid sequence that differs by at least one, but not
10 more than five amino acids from the DX-890 amino acid sequence set forth in SEQ ID NO:23; (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO:24 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-88 amino acid sequence set forth in SEQ ID NO:24; and (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID
15 NO:25 or an amino acid sequence that differs by at least one, but not more than five amino acids from the DX-1000 amino acid sequence set forth in SEQ ID NO:25, a wherein the at least 80% of the polypeptides comprise a polyethylene glycol moiety attached to each lysine and an N-terminal primary amine of the polypeptide, and wherein the average molecular weight of each of the moieties is less than 12 kDa,
20 and (ii) a pharmaceutically acceptable carrier.

- In another aspect, the invention provides a composition for treating a disorder selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), an inflammatory disorder, hemophilia, post-operative bleeding, peri-operative bleeding, hereditary angioedema, fibrinolysis or
25 fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis, in a subject having the disorder or suspected of having the disorder, wherein the composition comprises the preparation as described above, and wherein the polypeptide of the preparation inhibits a protease.

50860-189

In another aspect, the invention provides use of a composition for treating a disorder selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), an inflammatory disorder, hemophilia, post-operative bleeding, peri-operative bleeding, hereditary angioedema, fibrinolysis or
5 fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis, in a subject having the disorder or suspected of having the disorder, wherein the composition comprises the preparation as described above, and wherein the polypeptide of the preparation inhibits a protease.

10 In another aspect, the invention provides use of a composition for the manufacture of a medicament for treating a disorder selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), an inflammatory disorder, hemophilia, post-operative bleeding, peri-operative bleeding, hereditary angioedema, fibrinolysis or fibrinogenolysis, excessive bleeding associated
15 with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis, in a subject having the disorder or suspected of having the disorder, wherein the composition comprises the preparation as described above, and wherein the polypeptide of the preparation inhibits a protease.

The details of one or more embodiments of the invention are set forth in
20 the accompanying drawings and the description below.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts the structure of DX-890 and the position of its four lysine residues.

FIG. 2 depicts the structure of DX-88 and the position of its three lysine
25 residues.

FIG. 3 depicts the structure of DX-1000 and the position of its three lysine residues.

50860-189

FIG. 4 is an exemplary PEGylation scheme. The reaction pH can be run at a pH of between 7.8 and 8.5.

FIG. 5 shows results of an exemplary MALDI analysis.

FIG. 6 shows results of exemplary GP-HPLC runs.

5 FIG. 7 shows exemplary results of SDS-PAGE analysis.

FIG. 8a shows exemplary results of clearance studies in mice and 8b shows clearance studies in rabbits. The data were plotted using a double 4-parameter exponential decay. Fig 8c shows an allometric extrapolation to humans. Extrapolated values for long half life clearance phase in a 70 Kg human were as
10 follows: DX-890, 8.4 hours; 5-PEG5-DX-890, 330 hours, or about 14 days; DX-1000, 1.7 hours; 4-PEG5-DX-1000, 210 hours, or about 9 days.

FIG. 9 shows exemplary results of DX-88 poly-PEGylation at various ratios by SDS-PAGE analysis.

DETAILED DESCRIPTION

The invention provides, in part, compounds that bind to and inhibit a protease (e.g., an elastase, e.g., a neutrophil elastase). The compounds include (i) a polypeptide that includes a Kunitz domain and (ii) a plurality of moieties (such as polymer
5 moieties) that increases the molecular weight of the compounds relative to the polypeptide alone. The addition of the moieties to the compound can increase the *in vivo* circulating half life of the compound. In some embodiments, the compounds can inhibit neutrophil elastase with high affinity and selectivity.

Polymers

10 A variety of moieties can be used to increase the molecular weight of a polypeptide that includes a Kunitz domain or other protease inhibitor. In one embodiment, the moiety is a polymer, e.g., a water soluble and/or substantially non-antigenic polymer such as a homopolymer or a non-biological polymer. Substantially non-antigenic polymers include, e.g., polyalkylene oxides or polyethylene oxides. The
15 moiety may improve stabilization and/or retention of the Kunitz domain in circulation, e.g., in blood, serum, lymph, or other tissues, e.g., by at least 1.5, 2, 5, 10, 50, 75, or 100 fold. A plurality of moieties are attached to a Kunitz domain. For example, the polypeptide is attached to at least three moieties of the polymer. Each lysine of the polypeptide can be attached to a moiety of the polymer.

20 Suitable polymers can vary substantially by weight. For example, it is possible to use polymers having average molecular weights ranging from about 200 Daltons to about 40 kDa, e.g., 1-20 kDa, 4-12 kDa or 3-8 kDa, e.g., about 4, 5, 6, or 7 kDa. In one embodiment, the average molecular weight of individual moieties of the polymer that are associated with the compound are less than 20, 18, 17, 15, 12, 10, 8, or 7 kDa. The
25 final molecular weight can also depend upon the desired effective size of the conjugate, the nature (e.g. structure, such as linear or branched) of the polymer, and the degree of derivatization.

A non-limiting list of exemplary polymers include polyalkylene oxide homopolymers such as polyethylene glycol (PEG) or polypropylene glycols,
30 polyoxyethylenated polyols, copolymers thereof and block copolymers thereof, provided that the water solubility of the block copolymers is maintained. The polymer can be a hydrophilic polyvinyl polymers, e.g. polyvinylalcohol and

polyvinylpyrrolidone. Additional useful polymers include polyoxyalkylenes such as polyoxyethylene, polyoxypropylene, and block copolymers of polyoxyethylene and polyoxypropylene (Pluronics); polylactic acid; polyglycolic acid; polymethacrylates; carbomers; branched or unbranched polysaccharides which comprise the saccharide
5 monomers D-mannose, D- and L-galactose, fucose, fructose, D-xylose, L-arabinose, D-glucuronic acid, sialic acid, D-galacturonic acid, D-mannuronic acid (e.g. polymannuronic acid, or alginic acid), D-glucosamine, D-galactosamine, D-glucose and neuraminic acid including homopolysaccharides and heteropolysaccharides such as lactose, cellulose, amylopectin, starch, hydroxyethyl starch, amylose, dextrane sulfate,
10 dextran, dextrans, glycogen, or the polysaccharide subunit of acid mucopolysaccharides, e.g. hyaluronic acid; polymers of sugar alcohols such as polysorbitol and polymannitol; heparin or heparon. In some embodiments, the polymer includes a variety of different copolymer blocks.

The polypeptide that includes a Kunitz domain can be physically associated
15 with the polymer in a variety of ways. Typically, the polypeptide is covalently linked to the polymer at a plurality of sites. For example, the polypeptide is conjugated to the polymer at a plurality of primary amines, e.g., all accessible primary amines or all primary amines. Other compounds can also be attached to the same polymer, e.g., a cytotoxin, a label, or another targeting agent, e.g., another ligand that binds to the same
20 target as the Kunitz domain or a ligand that binds to another target, e.g., a an unrelated ligand. Other compounds may also be attached to the polypeptide.


In one embodiment, the polymer is water soluble prior to conjugation to the polypeptide (although need not be). Generally, after conjugation to the polypeptide, the product is water soluble, e.g., exhibits a water solubility of at least about 0.01 mg/ml,
25 and more preferably at least about 0.1 mg/ml, and still more preferably at least about 1 mg/ml. In addition, the polymer should not be highly immunogenic in the conjugate form, nor should it possess viscosity that is incompatible with intravenous infusion or injection if the conjugate is intended to be administered by such routes.

In one embodiment, the polymer contains only a single group which is reactive.
30 This helps to avoid conjugation of one polymer to multiple protein molecules. Mono-activated, alkoxy-terminated polyalkylene oxides (PAO's), e.g., monomethoxy-terminated polyethylene glycols (mPEG's); C₁₋₄ alkyl-terminated polymers; and bis-

activated polyethylene oxides (glycols) can be used for conjugation to the polypeptide. See, e.g., U.S. 5,951,974.

In its most common form, poly(ethylene glycol), PEG, is a linear or branched polyether terminated with hydroxyl groups. Linear PEG can have the following

5 general structure:



PEG can be synthesized by anionic ring opening polymerization of ethylene oxide initiated by nucleophilic attack of a hydroxide ion on the epoxide ring. Particularly useful for polypeptide modification is monomethoxy PEG, mPEG, having the general

10 structure:



For further descriptions, see, e.g., Roberts et al. (2002) *Advanced Drug Delivery Reviews* 54:459-476. In one embodiment, the polymer units used for conjugation are mono-disperse or otherwise highly homogenous, e.g., present in a preparation in which

15 95% or all molecules are with 7, 5, 4, 3, 2, or 1 kDa of one another. In another embodiment, the polymer units are poly-disperse.

It is possible to select reaction conditions that reduce cross-linking between polymer units or conjugation to multiple polypeptides and to purify the reaction products through gel filtration or ion exchange chromatography to recover substantially

20 homogenous derivatives, e.g., derivatives that include only a single Kunitz domain polypeptide. In other embodiments, the polymer contains two or more reactive groups for the purpose of linking multiple polypeptides (e.g., multiple units of the Kunitz domain polypeptide) to the polymer. Again, gel filtration or ion exchange chromatography can be used to recover the desired derivative in substantially

25 homogeneous form.

The polypeptide that includes a Kunitz domain is generally attached to a plurality of PEG molecules. For example, to form a compound that is larger than 20 or 30 kDa, a Kunitz domain (about 7 kDa) can be attached to at least three 8 kDa molecules of PEG. Other combinations are possible, e.g., at least two, four, or five

30 molecules of PEG. The molecular weight of the PEG molecules can be selected so that the final molecular weight of the compound is equal to or larger than a desired molecular weight (e.g., between 17-35, or 20-25, or 27-33 kDa). The plurality of PEG molecules can be attached to any region of the Kunitz domain, preferably at least 5, 10,

or 15 Angstroms from a region that interacts with a target, or at least 2, 3, or 4 residues from an amino acid that interacts with a target. The PEG molecules can be attached, e.g., to lysine residues or a combination of lysine residues and the N-terminus.

A covalent bond can be used to attach a polypeptide (e.g., a polypeptide that includes a Kunitz domain) to a polymer, for example, conjugation to the N-terminal amino group and epsilon amino groups found on lysine residues, as well as other amino, imino, carboxyl, sulfhydryl, hydroxyl or other hydrophilic groups. The polymer may be covalently bonded directly to the polypeptide without the use of a multifunctional (ordinarily bifunctional) crosslinking agent. Covalent binding to amino groups can be accomplished by known chemistries based upon cyanuric chloride, carbonyl diimidazole, aldehyde reactive groups (PEG alkoxide plus diethyl acetyl of bromoacetaldehyde; PEG plus DMSO and acetic anhydride, or PEG chloride plus the phenoxide of 4-hydroxybenzaldehyde, activated succinimidyl esters, activated dithiocarbonate PEG, 2,4,5-trichlorophenylchloroformate or P-nitrophenylchloroformate activated PEG.) Carboxyl groups can be derivatized by coupling PEG-primary amine using carbodiimide. Sulfhydryl groups can be derivatized by coupling to maleimido-substituted PEG (see, e.g., WO 97/10847) or PEG-maleimide (e.g., commercially available from Shearwater Polymers, Inc., Huntsville, Ala.). Alternatively, free amino groups on the polypeptide (e.g. epsilon amino groups on lysine residues) can be thiolated with 2-imino-thiolane (Traut's reagent) and then coupled to maleimide-containing derivatives of PEG, e.g., as described in Pedley et al., Br. J. Cancer, 70: 1126-1130 (1994).

Functionalized PEG polymers that can be attached to a polypeptide that includes Kunitz domain include polymers that are commercially available, e.g., from Shearwater Polymers, Inc. (Huntsville, Ala.). Such PEG derivatives include, e.g., amino-PEG, PEG amino acid esters, PEG-hydrazide, PEG-thiol, PEG-succinate, carboxymethylated PEG, PEG-propionic acid, PEG amino acids, PEG succinimidyl succinate, PEG succinimidyl propionate, succinimidyl ester of carboxymethylated PEG, succinimidyl carbonate of PEG, succinimidyl esters of amino acid PEGs, PEG-oxycarbonylimidazole, PEG-nitrophenyl carbonate, PEG tresylate, PEG-glycidyl ether, PEG-aldehyde, PEG vinylsulfone, PEG-maleimide, PEG-orthopyridyl-disulfide, heterofunctional PEGs, PEG vinyl derivatives, PEG silanes, and PEG phospholides. The reaction conditions for coupling these PEG derivatives may vary depending on the

polypeptide, the desired degree of PEGylation, and the PEG derivative utilized. Some factors involved in the choice of PEG derivatives include: the desired point of attachment (such as lysine or cysteine R-groups), hydrolytic stability and reactivity of the derivatives, stability, toxicity and antigenicity of the linkage, suitability for analysis, etc.

The conjugates of a polypeptide that includes a Kunitz domain and a polymer can be separated from the unreacted starting materials using chromatographic methods, e.g., by gel filtration or ion exchange chromatography, e.g., HPLC. Heterologous species of the conjugates are purified from one another in the same fashion. Resolution of different species (e.g. containing one or two PEG residues) is also possible due to the difference in the ionic properties of the unreacted amino acids. See, e.g., WO 96/34015.

Kunitz Domains

As used herein, a "Kunitz domain" is a polypeptide domain having at least 51 amino acids and containing at least two, and preferably three, disulfides. The domain is folded such that the first and sixth cysteines, the second and fourth, and the third and fifth cysteines form disulfide bonds (e.g., in a Kunitz domain having 58 amino acids, cysteines can be present at positions corresponding to amino acids 5, 14, 30, 38, 51, and 55, according to the numbering of the BPTI sequence provided below, and disulfides can form between the cysteines at position 5 and 55, 14 and 38, and 30 and 51), or, if two disulfides are present, they can form between a corresponding subset of cysteines thereof. The spacing between respective cysteines can be within 7, 5, 4, 3 or 2 amino acids of the following spacing between positions corresponding to: 5 to 55, 14 to 38, and 30 to 51, according to the numbering of the BPTI sequence provided below. The BPTI sequence can be used as a reference to refer to specific positions in any generic Kunitz domain. Comparison of a Kunitz domain of interest to BPTI can be performed by identifying the best fit alignment in which the number of aligned cysteines is maximized.

The 3D structure (at high resolution) of the Kunitz domain of BPTI is known. One of the X-ray structures is deposited in the Brookhaven Protein Data Bank as "6PTI". The 3D structure of some BPTI homologues (Eigenbrot *et al.*, (1990) Protein Engineering, 3(7):591-598; Hynes *et al.*, (1990) Biochemistry, 29:10018-10022) are

known. At least seventy Kunitz domain sequences are known. Known human homologues include three Kunitz domains of LACI (Wun *et al.*, (1988) J. Biol. Chem. 263(13):6001-6004; Girard *et al.*, (1989) Nature, 338:518-20; Novotny *et al.*, (1989) J. Biol. Chem., 264(31):18832-18837) two Kunitz domains of Inter- α -Trypsin Inhibitor, APP-I (Kido *et al.*, (1988) J. Biol. Chem., 263(34):18104-18107), a Kunitz domain from collagen, and three Kunitz domains of TFPI-2 (Sprecher *et al.*, (1994) PNAS USA, 91:3353-3357). LACI is a human serum phosphoglycoprotein with a molecular weight of 39 kDa (amino acid sequence in Table 1) containing three Kunitz domains.

Table 1: Exemplary Natural Kunitz Domains

LACI: (SEQ ID NO. 1)	1 MIYTMKKVHA LWASVCLLLN LAPAPLNAds eedeehtiit dtelpplklm 51 HSFCFAFKADD GPCKAIMKRF FFNIFTRQCE EFIYGGCEGN QNRFESLEEC 101 KKMCTRDnan riiktttlqqe <u>kpdfCf</u> leed <u>pgiCrgyitr</u> yfynnqtkqC 151 erfkyggClg nmnnfetlee CkniCedgpn gfqvdnygtq lnavnnsltP 201 qstkvpSlfe fhgpswCltp adrglCrane nrfyynsvig kCrpfkysgC 251 ggnennftsk qeClraCkkg fiqriskggl iktkrkrkkq rvkiayeeif 301 vknm
	The signal sequence (1-28) is uppercase and underscored LACI-K1 is uppercase LACI-K2 is underscored LACI-K3 is bold
BPTI (SEQ ID NO: 2)	1 2 3 4 5 1234567890123456789012345678901234567890123456789012345678 RPDF <u>C</u> LEPPYTGP <u>C</u> KARIIRYFYNAKAGLC <u>Q</u> TFVYGG <u>C</u> RAKRNNFKSAED <u>C</u> MRT <u>C</u> GGA

10

The Kunitz domains above are referred to as LACI-K1 (residues 50 to 107), LACI-K2 (residues 121 to 178), and LACI-K3 (213 to 270). The cDNA sequence of LACI is reported in Wun *et al.* (J. Biol. Chem., 1988, 263(13):6001-6004). Girard *et al.* (Nature, 1989, 338:518-20) reports mutational studies in which the P1 residues of each of the three Kunitz domains were altered. LACI-K1 inhibits Factor VIIa (F.VIIa) when F.VIIa is complexed to tissue factor and LACI-K2 inhibits Factor Xa.

Proteins containing exemplary Kunitz domains include the following, with SWISS-PROT Accession Numbers in parentheses:

20 A4_HUMAN (P05067), A4_MACFA (P53601), A4_MACMU (P29216),
A4_MOUSE (P12023), A4_RAT (P08592), A4_SAISC (Q95241),
AMBP_PLEPL (P36992), APP2_HUMAN (Q06481), APP2_RAT (P15943),
AXP1_ANTAF (P81547), AXP2_ANTAF (P81548), BPT1_BOVIN (P00974),
BPT2_BOVIN (P04815), CA17_HUMAN (Q02388), CA36_CHICK (P15989),
25 CA36_HUMAN (P12111), CRPT_BOOMI (P81162), ELAC_MACEU (O62845),
ELAC_TRIVU (Q29143), EPPI_HUMAN (O95925), EPPI_MOUSE (Q9DA01),

HTIB_MANSE (P26227), IBP_CARCR (P00993), IBPC_BOVIN (P00976),
 IBPI_TACTR (P16044), IBPS_BOVIN (P00975), ICS3_BOMMO (P07481),
 IMAP_DROFU (P11424), IP52_ANESU (P10280), ISC1_BOMMO (P10831),
 ISC2_BOMMO (P10832), ISH1_STOHE (P31713), ISH2_STOHE (P81129),
 5 ISIK_HELPO (P00994), ISP2_GALME (P81906), IVB1_BUNFA (P25660),
 IVB1_BUNMU (P00987), IVB1_VIPAA (P00991), IVB2_BUNMU (P00989),
 IVB2_DABRU (P00990), IVB2_HEMHA (P00985), IVB2_NAJNI (P00986),
 IVB3_VIPAA (P00992), IVBB_DENPO (P00983), IVBC_NAJNA (P19859),
 IVBC_OPHHA (P82966), IVBE_DENPO (P00984), IVBI_DENAN (P00980),
 10 IVBI_DENPO (P00979), IVBK_DENAN (P00982), IVBK_DENPO (P00981),
 IVBT_ERIMA (P24541), IVBT_NAJNA (P20229), MCPI_MELCP (P82968),
 SBPI_SARBU (P26228), SPT3_HUMAN (P49223), TKD1_BOVIN (Q28201),
 TKD1_SHEEP (Q29428), TXCA_DENAN (P81658), UPTI_PIG (Q29100),
 AMBP_BOVIN (P00978), AMBP_HUMAN (P02760), AMBP_MERUN (Q62577),
 15 AMBP_MESAU (Q60559), AMBP_MOUSE (Q07456), AMBP_PIG (P04366),
 AMBP_RAT (Q64240), IATR_HORSE (P04365), IATR_SHEEP (P13371),
 SPT1_HUMAN (O43278), SPT1_MOUSE (Q9R097), SPT2_HUMAN (O43291),
 SPT2_MOUSE (Q9WU03), TFP2_HUMAN (P48307), TFP2_MOUSE (O35536),
 TFPI_HUMAN (P10646), TFPI_MACMU (Q28864), TFPI_MOUSE (O54819),
 20 TFPI_RABIT (P19761), TFPI_RAT (Q02445), YN81_CAEEL (Q03610)

Table 2: The amino-acid sequences of 19 human Kunitz domains. Amino-acid sequences of 19 Human Kunitz Domains Binding loops are underscored.

25

Collagen A1 VII (SEQ ID NO:3)

SDDPCSLPLDEGSCTAYTLRWYHRAVTEACHPFVYGGCGGNANRFGTREACERRCPPR

30 TFPI2-K1 (SEQ ID NO:4)

NAEICLLPLDYGPCRALLRYYYDRYTQSCRQFLYGGCEGNANNFYTWACDDACWRI

AppI (SEQ ID NO:5)

35

VREVCSEQAETGPCRAMISRWFYFDVTEGKCAPFFYGGCGGNRNNFDTEEYCMVCGSA

Hep GF AI T2, K2 (SEQ ID NO:6)

40 YEEYCTANAVTGPCRASFPWFYFDVERNSCNNFIYGGCRGNKNSYRSEEACMLRCFRQ

ITI, K1 (SEQ ID NO:7)

KEDSCQLGYSAGPCMGMTSRYFYNGTSMACETFQYGGCMGNGNMFVTEKECLQTCRTV

45

Chrome20 (SEQ ID NO:8)

FQEPCLPVRHGNHCNHEAQRWHFDFKNYRCTPFKYRGCEGNANNFLNEDACRTACMLI
R

50

Hep GF AI T1, K1 (SEQ ID NO:9)

- TEDYCLASNKVGRCRGSFPRWYYDPTEQICKSFVYGGCLGNKNNYLREEECILACRGV
- Hep GF AI T1, K2 (SEQ ID NO:10)
- 5 DKGHCVDLPDTGLCKESIPRWYYPFSEHCARFTYGGCYGNKNNFEEEQQCLESCRGI
- TFPI2-K3 (SEQ ID NO:11)
- 10 IPSFCYSPKDEGLCSANVTRYFNPRTCDAFTYTGCGGNDNNFVSREDCKRACAKA
- ITI, K2 (SEQ ID NO:12)
- AACNLPIVRGPCRAFIQLWAFDAVKGKCVLFPYGGCQGNGNKFYSEKECREYCGVP
- 15 Hep GF AI T2, K1 (SEQ ID NO:13)
- IHDFCLVSKVVGRCRASMPRWYNVTDGSCQLFVYGGCDGNSNNYLTKEECLKKCATV
- 20 App2 (SEQ ID NO:14)
- VKAVCSQEAMTGPCRAVMPRWYFDLSKGKCVRFIYGGCGGNRNNFESEDYCMAVCKAM
- TFPI1 K2=LACI-D2 (SEQ ID NO:15)
- 25 KPDFCFLEEDPGICRGYITRYFYNNQTKQCERFKYGGCLGNMNNFETLEECKNICEDG
- TFPI2-K2 (SEQ ID NO:16)
- 30 VPKVCRLQVSVDDQCEGSTEKYFFNLSSMTCEKFFSGGCHRNRIENRFPDEATCMGFC
- APK
- HKI B9 (SEQ ID NO:17)
- 35 LPNVCAFPMEKGPCQTYMTRWFFNFETGECELFAYGGCGGNSNNFLRKEKCEKFCKFT
- TFPI1 K1=LACI-D1 (SEQ ID NO:18)
- MHSFCAFKADDGPCKAIMKRFFFNIFTRQCEEFIYGGCEGNQNRFESLEECKKMCTRD
- 40 TFPI1 K3=LACI-D3 (SEQ ID NO:19)
- GPSWCLTPADRGGLCRANENRFYNSVIGKCRPFKYSGCGGNENNFTSKQECLRACKKG
- 45 Collagen A3 (SEQ ID NO:20)
- ETDICKLPKDEGTCRDFILKWYYDPNTKSCARFWYGGCGGNENKFGSQKECEKVCAPV
- CAB37635 (SEQ ID NO:21)

KQDVCEMPKETGPCLAYFLHWYDKKDNTCSMFVYGGCQGNNNFQSKANCLNTCKNK

End Table 2.

5

A variety of methods can be used to identify a Kunitz domain from a sequence database. For example, a known amino acid sequence of a Kunitz domain, a consensus sequence, or a motif (e.g., the ProSite Motif) can be searched against the GenBank sequence databases (National Center for Biotechnology Information, National Institutes of Health, Bethesda MD), e.g., using BLAST; against Pfam database of HMMs (Hidden Markov Models) (e.g., using default parameters for Pfam searching; against the SMART database; or against the ProDom database. For example, the Pfam Accession Number PF00014 of Pfam Release 9 provides numerous Kunitz domains and an HMM for identify Kunitz domains. A description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3):405-420 and a detailed description of HMMs can be found, for example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159; Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314. The SMART database (Simple Modular Architecture Research Tool, EMBL, Heidelberg, DE) of HMMs as described in Schultz *et al.* (1998), *Proc. Natl. Acad. Sci. USA* 95:5857 and Schultz *et al.* (2000) *Nucl. Acids Res* 28:231. The SMART database contains domains identified by profiling with the hidden Markov models of the HMMer2 search program (R. Durbin *et al.* (1998) *Biological sequence analysis: probabilistic models of proteins and nucleic acids*. Cambridge University Press). The database also is annotated and monitored. The ProDom protein domain database consists of an automatic compilation of homologous domains (Corpet *et al.* (1999), *Nucl. Acids Res.* 27:263-267). Current versions of ProDom are built using recursive PSI-BLAST searches (Altschul *et al.* (1997) *Nucleic Acids Res.* 25:3389-3402; Gouzy *et al.* (1999) *Computers and Chemistry* 23:333-340.) of the SWISS-PROT 38 and TREMBL protein databases. The database automatically generates a consensus sequence for each domain. Prosite lists the Kunitz domain as a motif and identifies proteins that include a Kunitz domain. See, e.g., Falquet *et al.* *Nucleic Acids Res.* 30:235-238(2002).

Useful Kunitz domains for selecting protease inhibitors can include Kunitz domains that have a framework region with a particular number of lysine residues. In one implementation, frameworks with four lysine residues are useful and can be modified, e.g., by attachment of PEG moieties of average molecular weight between 3-
5 8 kDa, e.g., about 5 kDa. For example, the ITI framework has four lysines. In another implementation, frameworks with three lysines are useful and can be modified e.g., by attachment of PEG moieties of average molecular weight between 4-10 kDa, e.g., about 5 kDa or 7 kDa. LACI is one such framework. Frameworks can also be altered to include fewer or additional lysines, for example, to reduce the number of lysines that
10 are within five, four, or three residues of a binding loop, or to introduce a sufficient number of lysines that the protein can be modified with small PEG moieties (e.g., between 3-8 kDa PEG moieties) to increase the size of the protein and stability of the protein in vivo.

Kunitz domains interact with target protease using, primarily, amino acids in
15 two loop regions ("binding loops"). The first loop region is between about residues corresponding to amino acids 11-21 of BPTI. The second loop region is between about residues corresponding to amino acids 31-42 of BPTI. An exemplary library of Kunitz domains varies one or more amino acid positions in the first and/or second loop regions. Particularly useful positions to vary include: positions 13, 16, 17, 18, 19, 31,
20 32, 34, and 39 with respect to the sequence of BPTI. At least some of these positions are expected to be in close contact with the target protease.

The "framework region" of a Kunitz domain is defined as those residues that are a part of the Kunitz domain, but specifically excluding residues in the first and second binding loops regions, i.e., about residues corresponding to amino acids 11-21
25 of BPTI and 31-42 of BPTI.

Conversely, residues that are not at these particular positions or which are not in the loop regions may tolerate a wider range of amino acid substitution (e.g., conservative and/or non-conservative substitutions) than these amino acid positions.

Elastase-Inhibiting Kunitz domains

30 One exemplary polypeptide that binds to and inhibits human neutrophil elastase (hNE) is DX-890 (also known as "EPI-hNE4"). DX-890 is a highly specific and potent

($K_i = 4 \times 10^{-12}$ M) inhibitor of human neutrophil elastase (hNE). DX-890 includes the following amino acid sequence:

Glu Ala Cys Asn Leu Pro Ile Val Arg Gly Pro Cys Ile Ala
Phe
5 Phe Pro Arg Trp Ala Phe Asp Ala Val Lys Gly Lys Cys Val
Leu
Phe Pro Tyr Gly Gly Cys Gln Gly Asn Gly Asn Lys Phe Tyr
Ser
Glu Lys Glu Cys Arg Glu Tyr Cys Gly Val Pro

10 DX-890 is derived from the second Kunitz-type domain of inter- α -inhibitor protein (ITI-D2) and can be produced by fermentation in *Pichia pastoris*. It includes 56 amino acids, with a predicted MW of 6,237 Daltons. DX-890 is resistant to oxidative and proteolytic inactivation.

In vitro, *ex vivo* and *in vivo* pharmacological studies have demonstrated hNE
15 inhibitory capacity and the protective effect of DX-890 against lesions induced by hNE of sputum from cystic fibrosis children (see ref. Delacourt et al. 2002). Acute and subchronic 4-week studies of aerosolised DX-890 in cynomolgus monkeys showed no signs of clinical or biological toxicity, nor of histopathological lesions induced by the administration of DX-890.

20 In clinical studies using healthy human volunteers, DX-890 was found to be safe for administration by inhalation at 8 increasing doses (up to 120 minutes of DX-890 in saline resulting in an inhaled mass of about 72 mg).

Some of the consequences of elastase activity include: cleavage of complement receptors and C3bi; cleavage of immunoglobulins; degradation of elastin (and
25 consequently plugging of airways, structural damage, bronchiectasis); secretion of macromolecules; increased interleukin-8; increase in PMN (and consequently release of oxygen, hydrogen peroxide, leukotriene B4 and interleukin-8); and persistence of bacteria. Inhibitors of elastase can be used to reduce one or more of these activities.

DX-890 can be used as an anti-inflammatory drug targeted against neutrophil
30 mediated inflammation, e.g., in pulmonary CF lesions. Exemplary pulmonary indications include Cystic Fibrosis (CF), Acute Respiratory Distress Syndrome (ARDS), and Chronic Obstructive Pulmonary Disease (COPD). In CF patients, the balance between proteinases and their inhibitors may become severely disturbed. Activated polymorphonuclear leukocytes (PMN) produce human neutrophil elastase
35 (hNE) and other proteases. hNE is considered to be a key cause of lung tissue damage

associated with cystic fibrosis. Inhibition of hNE is therefore a logical avenue for treatment of CF lung disease since it attacks the original source of damage rather than ameliorating symptoms and consequences of the damage.

It is possible, for example, to deliver DX-890 to the lung by nebulization.

5 DX-890 activity was detected in broncho-alveolar lavages of volunteer inhaling nebulized DX-890. 12 healthy volunteers received during 14 days a single daily dose of DX-890, by nebulization lasting 5 or 20 minutes, corresponding to estimated inhaled mass of 3.75 or 15 mg respectively. Tolerability was excellent; no significant adverse event was reported. No clinical or biological abnormalities were observed.

10 With respect to pulmonary indications, DX-890 can be used to treat, for example, Cystic Fibrosis (CF), Acute Respiratory Distress Syndrome (ARDS) and Chronic Obstructive Pulmonary Disease (COPD).

There are also known correlations between the structure of DX-890 and its ability to bind to hNE. See, e.g., U.S. 5,663,143. U.S. 5,663,143 also describes other
15 Kunitz domains that inhibit elastase. These and related domains (e.g., domains at least 70, 75, 80, 85, 90, or 95% identical) can also be used.

Exemplary Kunitz domains that inhibit plasma kallikrein are described, for example, in U.S. 6,057,287.

Exemplary Kunitz domains that inhibit plasmin are described, for example, in
20 U.S. 6,103,499.

Table 3: Exemplary Amino Acids for hNE inhibitors

Some preferred Amino acids in hNE-inhibiting Kunitz domains Position Allowed amino acids at amino acid positions corresponding to respective positions in BPTI	
5	C
10	YSVN
11	TARQP
12	G
13	PAV
14	C
15	IV
16	AG
17	FILVM
18	F
19	PSQKR
20	R
21	YWF
30	C
31	QEV
32	TLP
33	F
34	VQP
35	Y
36	G
37	G
38	C
39	MQ
40	GA
41	N
42	G
43	N
45	F
51	C
55	C

“Protection against acute lung injury by intravenous or intratracheal pretreatment with EPI-HNE4, a new potent neutrophil elastase inhibitor.” Delacourt C, Herigault S, Delclaux C, et al. Am J Respir Cell Mol Biol 2002;26:290-7 and Grimbart et al. (2003) “Characteristics of EPI-hNE4 aerosol: a new elastase inhibitor for treatment of cystic fibrosis” J Aerosol Med. 16(2):121-9.

Identifying Kunitz Domains and Other Protease Inhibitors

A variety of methods can be used to identify a protein that binds to and/or inhibits a protease. These methods can be used to identify natural and non-naturally occurring Kunitz domains that can be used as components of the compounds described
5 herein.

For example, a Kunitz domain can be identified from a library of proteins in which each of a plurality of library members includes a varied Kunitz domain. A variety of amino acids can be varied in the domain. See, e.g., U.S. 5,223,409; U.S. 5,663,143, and U.S. 6,333,402. Kunitz domains can varied, e.g., using DNA
10 mutagenesis, DNA shuffling, chemical synthesis of oligonucleotides (e.g., using codons as subunits), and cloning of natural genes. See, e.g., U.S. 5,223,409 and U.S. 2003-0129659.

The library can be an expression library that is used to produce proteins. The proteins can be arrayed, e.g., using a protein array. US 5,143,854; De Wildt *et al.*
15 (2000) *Nat. Biotechnol.* 18:989-994; Lueking *et al.* (1999) *Anal. Biochem.* 270:103-111; Ge (2000) *Nucleic Acids Res.* 28, e3, I-VII; MacBeath and Schreiber (2000) *Science* 289:1760-1763; WO 0/98534, WO01/83827, WO02/12893, WO 00/63701, WO 01/40803 and WO 99/51773.

The proteins can also be displayed on a replicable genetic package, e.g., in the
20 form of a phage library such as a phage display, yeast display library, ribosome display, or nucleic acid-protein fusion library. See, e.g., U.S. 5,223,409; Smith (1985) *Science* 228:1315-1317; WO 92/18619; WO 91/17271; WO 92/20791; WO 92/15679; WO 93/01288; WO 92/01047; WO 92/09690; WO 90/02809; de Haard *et al.* (1999) *J. Biol. Chem* 274:18218-30; Hoogenboom *et al.* (1998) *Immunotechnology* 4:1-20;
25 Hoogenboom *et al.* (2000) *Immunol Today* 2:371-8; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrard *et al.* (1991) *Bio/Technology* 9:1373-1377;
30 Rebar *et al.* (1996) *Methods Enzymol.* 267:129-49; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133-4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982 for examples of phage display and other methods. See, e.g., Boder and Wittrup (1997) *Nat. Biotechnol.* 15:553-557 and WO 03/029456 for examples of yeast cell display and other methods.

See, e.g., Mattheakis *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:9022 and Hanes *et al.* (2000) *Nat Biotechnol.* 18:1287-92; Hanes *et al.* (2000) *Methods Enzymol.* 328:404-30. and Schaffitzel *et al.* (1999) *J Immunol Methods.* 231(1-2):119-35 for examples of ribosome display and other methods. See, e.g., Roberts and Szostak (1997) *Proc. Natl.*
5 *Acad. Sci. USA* 94:12297-12302, and U.S. Patent No. 6,207,446 for examples of nucleic acid-protein fusions. Such libraries can be screened in a high throughput format. See, e.g., U.S. 2003-0129659.

Libraries of Kunitz domains can be generated by varying one or more binding site loop amino acid residues using a Kunitz domain described herein, e.g., a Kunitz
10 domain having a framework described herein, e.g., a modified or naturally occurring framework region. In one embodiment, the residues that are varied are varied among a plurality of amino acids. The plurality is chosen such that lysine is unavailable.

Screening Display Libraries

This section describes exemplary methods of screening a display library to
15 identify a polypeptide that interacts with an elastase. These methods can be modified to identify other polypeptides that interact with other targets, e.g., other proteases or other proteins. The methods can also be modified and used in combination with other types of libraries, e.g., an expression library or a protein array, and so forth.

In an exemplary display library screen, a phage library is contacted with and
20 allowed to bind to the target elastase protein (e.g., an active or an inactivated form (e.g., mutant or chemically inactivated protein) or a fragment thereof). To facilitate separation of binders and non-binders in the screening process, it is often convenient to immobilize the elastase on a solid support, although it is also possible to first permit binding to elastase in solution and then segregate binders from non-binders by coupling
25 the target compound to a support. By way of illustration, when incubated in the presence of the elastase, phage displaying a polypeptide that interacts with elastase form a complex with the elastase immobilized on a solid support whereas non-binding phage remain in solution and may be washed away with buffer. Bound phage may then be liberated from the elastase by a number of means, such as changing the buffer to a
30 relatively high acidic or basic pH (e.g., pH 2 or pH 10), changing the ionic strength of the buffer, adding denaturants, adding a competitor, adding a host cell which can be infected, or other known means.

For example, to identify elastase-binding peptides, elastase can be adsorbed to a solid surface, such as the plastic surface of wells in a multi-well assay plate.

Subsequently, an aliquot of a phage display library is added to a well under appropriate conditions that maintain the structure of the immobilized elastase and the phage, such as pH 6-7. Phage in the libraries that display polypeptides that bind the immobilized elastase are bound to the elastase and are retained in the well. Non-binding phage can be removed. It is also possible to include a blocking agent or competing ligand during the binding of the phage library to the immobilized elastase.

Phage bound to the immobilized elastase may then be eluted by washing with a buffer solution having a relatively strong acid pH (e.g., pH 2) or an alkaline pH (e.g., pH 8-9). The solutions of recovered phage that are eluted from the elastase are then neutralized and may, if desired, be pooled as an enriched mixed library population of phage displaying elastase binding peptides. Alternatively the eluted phage from each library may be kept separate as a library-specific enriched population of elastase binders. Enriched populations of phage displaying elastase binding peptides may then be grown up by standard methods for further rounds of screening and/or for analysis of peptide displayed on the phage and/or for sequencing the DNA encoding the displayed binding peptide.

One of many possible alternative screening protocols uses elastase target molecules that are biotinylated and that can be captured by binding to streptavidin, for example, coated on particles.

Recovered phage may then be amplified by infection of bacterial cells, and the screening process may be repeated with the new pool of phage that is now depleted in non-elastase binders and enriched in elastase binders. The recovery of even a few binding phage may be sufficient to carry the process to completion. After a few rounds of selection, the gene sequences encoding the binding moieties derived from selected phage clones in the binding pool are determined by conventional methods, revealing the peptide sequence that imparts binding affinity of the phage to the target. An increase in the number of phage recovered after each round of selection and the recovery of closely related sequences indicate that the screening is converging on sequences of the library having a desired characteristic.

After a set of binding polypeptides is identified, the sequence information may be used to design other, secondary libraries. For example, the secondary libraries can

explore a smaller segment of sequence space in more detail than the initial library. In some embodiments, the secondary library includes proteins that are biased for members having additional desired properties, e.g., sequences that have a high percentage identity to a human protein.

5 Display technology can also be used to obtain polypeptides that are specific to particular epitopes of a target. This can be done, for example, by using competing non-target molecules that lack the particular epitope or are mutated within the epitope, e.g., with alanine. Such non-target molecules can be used in a negative selection procedure as described below, as competing molecules when binding a display library to the
10 target, or as a pre-elution agent, e.g., to capture in a wash solution dissociating display library

Iterative Selection. In one preferred embodiment, display library technology is used in an iterative mode. A first display library is used to identify one or more proteins that interacts with a target. These identified proteins are then varied using a
15 mutagenesis method to form a second display library. Higher affinity proteins are then selected from the second library, e.g., by using higher stringency or more competitive binding and washing conditions.

In some implementations, the mutagenesis is targeted to regions known or likely to be at the binding interface. Some exemplary mutagenesis techniques include: error-prone PCR (Leung *et al.* (1989) *Technique* 1:11-15), recombination, DNA shuffling
20 using random cleavage (Stemmer (1994) *Nature* 389-391; termed “nucleic acid shuffling”), RACHITT™ (Coco *et al.* (2001) *Nature Biotech.* 19:354), site-directed mutagenesis (Zoller *et al.* (1987) *Nucl Acids Res* 10:6487-6504), cassette mutagenesis (Reidhaar-Olson (1991) *Methods Enzymol.* 208:564-586) and incorporation of
25 degenerate oligonucleotides (Griffiths *et al.* (1994) *EMBO J* 13:3245). For Kunitz domains, many positions near the binding interface are known. Such positions include, for example, positions 13, 16, 17, 18, 19, 31, 32, 34, and 39 with respect to the sequence of BPTI. (according to the BPTI numbering in U.S. 6,333,402). Such positions can be held constant and other positions can be varied or these positions
30 themselves may be varied.

In one example of iterative selection, the methods described herein are used to first identify proteins from a display library that bind an elastase with at least a minimal binding specificity for a target or a minimal activity, e.g., an equilibrium dissociation

constant for binding of greater than 1 nM, 10 nM, or 100 nM. The nucleic acid sequences encoding the initial identified proteins are used as a template nucleic acid for the introduction of variations, e.g., to identify a second protein ligand that has enhanced properties (e.g., binding affinity, kinetics, or stability) relative to the initial protein

5 ligand.

Off-Rate Selection. Since a slow dissociation rate can be predictive of high affinity, particularly with respect to interactions between proteins and their targets, the methods described herein can be used to isolate proteins with a desired kinetic dissociation rate (i.e. reduced) for a binding interaction to a target.

10 To select for slow dissociating proteins from a display library, the library is contacted to an immobilized target, e.g., immobilized elastase. The immobilized target is then washed with a first solution that removes non-specifically or weakly bound biomolecules. Then the immobilized target is eluted with a second solution that includes a saturation amount of free target, i.e., replicates of the target that are not
15 attached to the particle. The free target binds to biomolecules that dissociate from the target. Rebinding is effectively prevented by the saturating amount of free target relative to the much lower concentration of immobilized target.

The second solution can have solution conditions that are substantially physiological or that are stringent. Typically, the solution conditions of the second
20 solution are identical to the solution conditions of the first solution. Fractions of the second solution are collected in temporal order to distinguish early from late fractions. Later fractions include biomolecules that dissociate at a slower rate from the target than biomolecules in the early fractions.

Further, it is also possible to recover display library members that remain bound
25 to the target even after extended incubation. These can either be dissociated using chaotropic conditions or can be amplified while attached to the target. For example, phage bound to the target can be contacted to bacterial cells.

Selecting or Screening for Specificity. The display library screening methods described herein can include a selection or screening process that discards display
30 library members that bind to a non-target molecule, e.g., a protease other than elastase, such as trypsin. In one embodiment, the non-target molecule is elastase that has been activated by treatment with an irreversibly bound inhibitor, e.g., a covalent inhibitor.

In one implementation, a so-called “negative selection” step or “depletion” is used to discriminate between the target and a related, but distinct or an unrelated non-target molecules. The display library or a pool thereof is contacted to the non-target molecule. Members of the sample that do not bind the non-target are collected and
5 used in subsequent selections for binding to the target molecule or even for subsequent negative selections. The negative selection step can be prior to or after selecting library members that bind to the target molecule.

In another implementation, a screening step is used. After display library members are isolated for binding to the target molecule, each isolated library member is
10 tested for its ability to bind to a non-target molecule (e.g., a non-target listed above). For example, a high-throughput ELISA screen can be used to obtain this data. The ELISA screen can also be used to obtain quantitative data for binding of each library member to the target. The non-target and target binding data are compared (e.g., using a computer and software) to identify library members that specifically bind to the
15 target.

Modifying and Varying Polypeptides

It is also possible to vary a protein described herein to obtain useful variant protein that has similar or improved or altered properties. Typically, a number of variants are possible. A variant can be prepared and then tested, e.g., using a binding
20 assay described herein (such as fluorescence anisotropy).

One type of variant is a truncation of a ligand described herein or isolated by a method described herein. In this example, the variant is prepared by removing one or more amino acid residues of the ligand from the N or C terminus. In some cases, a series of such variants is prepared and tested. Information from testing the series is
25 used to determine a region of the ligand that is essential for binding the elastase protein. A series of internal deletions or insertions can be similarly constructed and tested. For Kunitz domains, it can be possible to remove, e.g., between one and five residues or one and three residues that are N-terminal to C₅, the first cysteine, and between one and five residues or one and three residues that are C-terminal to C₅₅, the final cysteine,
30 wherein each of the cysteines corresponds to a respectively numbered cysteine in BPTI.

Another type of variant is a substitution. In one example, the ligand is subjected to alanine scanning to identify residues that contribute to binding activity. In another

example, a library of substitutions at one or more positions is constructed. The library may be unbiased or, particularly if multiple positions are varied, biased towards an original residue. In some cases, the substitutions are all conservative substitutions.

Another type of variant includes one or more non-naturally occurring amino acids. Such variant ligands can be produced by chemical synthesis or modification. One or more positions can be substituted with a non-naturally occurring amino acid. In some cases, the substituted amino acid may be chemically related to the original naturally occurring residue (e.g., aliphatic, charged, basic, acidic, aromatic, hydrophilic) or an isostere of the original residue.

It may also be possible to include non-peptide linkages and other chemical modifications. For example, part or all of the ligand may be synthesized as a peptidomimetic, e.g., a peptoid (see, e.g., Simon *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:9367-71 and Horwell (1995) *Trends Biotechnol.* 13:132-4). See also other modifications discussed below.

Characterization of Binding Interactions

The binding properties of a protein (e.g., a polypeptide that includes a Kunitz domain) can be readily assessed using various assay formats. For example, the binding property of a protein can be measured in solution by fluorescence anisotropy, which provides a convenient and accurate method of determining a dissociation constant (K_D) or association constant (K_a) of the protein for a particular target. In one such procedure, the protein to be evaluated is labeled with fluorescein. The fluorescein-labeled protein is mixed in wells of a multi-well assay plate with various concentrations of the particular target (e.g., elastase). Fluorescence anisotropy measurements are carried out using a fluorescence polarization plate reader.

ELISA. The binding interactions can also be analyzed using an ELISA assay. For example, the protein to be evaluated is contacted to a microtitre plate whose bottom surface has been coated with the target, e.g., a limiting amount of the target. The molecule is contacted to the plate. The plate is washed with buffer to remove non-specifically bound molecules. Then the amount of the protein bound to the plate is determined by probing the plate with an antibody that recognizes the protein. For example, the protein can include an epitope tag. The antibody can be linked to an enzyme such as alkaline phosphatase, which produces a colorimetric product when

appropriate substrates are provided. In the case where a display library member includes the protein to be tested, the antibody can recognize a region that is constant among all display library members, e.g., for a phage display library member, a major phage coat protein.

5 **Homogeneous Assays.** A binding interaction between a protein and a particular target can be analyzed using a homogenous assay, i.e., after all components of the assay are added, additional fluid manipulations are not required. For example, fluorescence energy transfer (FET) can be used as a homogenous assay (see, for example, Lakowicz *et al.*, U.S. Patent No. 5,631,169; Stavrianopoulos, *et al.*, U.S. Patent No. 4,868,103). A fluorophore label on the first molecule (e.g., the molecule identified in the fraction) is selected such that its emitted fluorescent energy can be absorbed by a fluorescent label on a second molecule (e.g., the target) if the second molecule is in proximity to the first molecule. The fluorescent label on the second molecule fluoresces when it absorbs the transferred energy. Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, the spatial relationship between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the 'acceptor' molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (e.g., using a fluorimeter). By titrating the amount of the first or second binding molecule, a binding curve can be generated to estimate the equilibrium binding constant.

Surface Plasmon Resonance (SPR). A binding interaction between a protein and a particular target can be analyzed using SPR. For example, after sequencing of a display library member present in a sample, and optionally verified, e.g., by ELISA, the displayed protein can be produced in quantity and assayed for binding the target using SPR. SPR or real-time Biomolecular Interaction Analysis (BIA) detects biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore). Changes in the mass at the binding surface (indicative of a binding event) of the BIA chip result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)). The changes in the refractivity generate a detectable signal, which are measured as an indication of real-time reactions between biological molecules. Methods for using SPR are described, for example, in

U.S. Patent No. 5,641,640; Raether (1988) *Surface Plasmons* Springer Verlag;
Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345; Szabo *et al.*
(1995) *Curr. Opin. Struct. Biol.* 5:699-705.

Information from SPR can be used to provide an accurate and quantitative
5 measure of the equilibrium dissociation constant (K_d), and kinetic parameters, including
 k_{on} and k_{off} , for the binding of a biomolecule to a target. Such data can be used to
compare different biomolecules. For example, proteins selected from a display library
can be compared to identify individuals that have high affinity for the target or that
have a slow k_{off} . This information can also be used to develop structure-activity
10 relationship (SAR) if the biomolecules are related. For example, if the proteins are all
mutated variants of a single parental antibody or a set of known parental antibodies,
variant amino acids at given positions can be identified that correlate with particular
binding parameters, e.g., high affinity and slow k_{off} .

Additional methods for measuring binding affinities include fluorescence
15 polarization (FP) (see, e.g., U.S. Patent No. 5,800,989), nuclear magnetic resonance
(NMR), and binding titrations (e.g., using fluorescence energy transfer).

Other solution measures for studying binding properties include fluorescence
resonance energy transfer (FRET) and NMR.

Characterization of Elastase Inhibition

20 With respect to embodiments in which the compound includes a polypeptide
that has a Kunitz domain specific for elastase, it may be useful to characterize the
ability of the polypeptide to inhibit elastase.

Kunitz domains can be screened for binding to elastase and for inhibition of
elastase proteolytic activity. Kunitz domains can be selected for their potency and
25 selectivity of inhibition of elastase. In one example, elastase and its substrate are
combined under assay conditions permitting reaction of the protease with its substrate.
The assay is performed in the absence of the Kunitz domain, and in the presence of
increasing concentrations of the Kunitz domain. The concentration of test compound at
which 50% of the elastase activity is inhibited by the test compound is the IC_{50} value
30 (Inhibitory Concentration) or EC_{50} (Effective Concentration) value for that compound.
Within a series or group of Kunitz domain, those having lower IC_{50} or EC_{50} values are
considered more potent inhibitors of the elastase than those compounds having higher

IC₅₀ or EC₅₀ values. Preferred compounds according to this aspect have an IC₅₀ value of 100 nM or less as measured in an *in vitro* assay for inhibition of elastase activity.

Kunitz domain can also be evaluated for selectivity toward elastase. A test compound is assayed for its potency toward a panel of serine proteases and other enzymes and an IC₅₀ value is determined for each peptide. A Kunitz domain that demonstrates a low IC₅₀ value for the elastase enzyme, and a higher IC₅₀ value for other enzymes within the test panel (e. g., trypsin, plasmin, kallikrein), is considered to be selective toward elastase. Generally, a compound is deemed selective if its IC₅₀ value is at least one order of magnitude less than the next smallest IC₅₀ value measured in the panel of enzymes.

Specific methods for evaluating inhibition of elastase are described in the Example below.

It is also possible to evaluate Kunitz domain activity *in vivo* or in samples (e.g., pulmonary lavages) of subjects to which a compound described herein has been administered.

Protease Targets

Proteases are involved in a wide variety of biological processes, including inflammation and tissue injury. Serine proteases produced by inflammatory cells, including neutrophils, are implicated in various disorders, such as pulmonary emphysema. Neutrophil elastase is a serine protease produced by polymorphonuclear leukocytes with activity against extracellular matrix components and pathogens. Pulmonary emphysema is characterized by alveolar destruction leading to a major impairment in lung function.

A deficiency of a serine protease inhibitor, α 1-protease inhibitor (API, or α 1-PI, formerly known as α -1 antitrypsin) is a risk factor for the development of pulmonary emphysema (Laurell, C. B. and Eriksson, S. (1963) *Scand. J. Clin. Lab. Invest.* 15:132–140; Brantly, M. L., et al. (1988) *Am. Rev. Respir. Dis.* 138:327–336). API deficiency may lead to uncontrolled activity of neutrophil elastase and contribute to the destruction of lung tissue in pulmonary emphysema. Likewise, API inactivation and chronic inflammation can lead to excess neutrophil elastase activity and pathologic destruction of pulmonary tissue.

Human neutrophil elastase consists of approximately 218 amino acid residues,

contains 2 asparagine-linked carbohydrate side chains, and is joined together by 2 disulfide bonds (Sinha, S., et al. *Proc. Nat. Acad. Sci.* 84: 2228-2232, 1987). It is normally synthesized in the developing neutrophil as a proenzyme but stored in the primary granules in its active form, ready with full enzymatic activity when released
5 from the granules, normally at sites of inflammation (Gullberg U, et al. *Eur J Haematol.* 1997;58:137-153; Borregaard N, Cowland JB. *Blood.* 1997;89:3503-3521).

Other exemplary protease targets include: plasmin, kallikrein, Factor VIIa, Factor XIa, thrombin, urokinase, and Factor IIa. Classes of relevant proteases include: proteases associated with blood coagulation, proteases associated with complement,
10 proteases that digest extracellular matrix components, proteases that digest basement membranes, and proteases associated with endothelial cells. For example, the protease is a serine protease.

Protein Production

15 *Recombinant production of polypeptides.* Standard recombinant nucleic acid methods can be used to express a polypeptide component of a compound described herein (e.g., a polypeptide that includes a Kunitz domain). Generally, a nucleic acid sequence encoding the polypeptide is cloned into a nucleic acid expression vector. If the polypeptide is sufficiently small, e.g., the protein is a peptide of less than 50 amino
20 acids, the protein can be synthesized using automated organic synthetic methods.

The expression vector for expressing the polypeptide can include a segment encoding the polypeptide and regulatory sequences, for example, a promoter, operably linked to the coding segment. Suitable vectors and promoters are known to those of skill in the art and are commercially available for generating the recombinant constructs
25 of the present invention. See, for example, the techniques described in Sambrook & Russell, *Molecular Cloning: A Laboratory Manual*, 3rd Edition, Cold Spring Harbor Laboratory, N.Y. (2001) and Ausubel *et al.*, *Current Protocols in Molecular Biology* (Greene Publishing Associates and Wiley Interscience, N.Y. (1989).

Scopes (1994) *Protein Purification: Principles and Practice*, New
30 York:Springer-Verlag and other texts provide a number of general methods for purifying recombinant (and non-recombinant) proteins.

Synthetic production of peptides. The polypeptide component of a compound can also be produced by synthetic means. See, e.g., Merrifield (1963) J. Am. Chem.Soc., 85 : 2149. For example, the molecular weight of synthetic peptides or peptide mimetics can be from about 250 to about 8,0000 Daltons. A peptide can be
5 modified, e.g., by attachment to a moiety that increases the effective molecular weight of the peptide. If the peptide is oligomerized, dimerized and/or derivatized, e.g., with a hydrophilic polymer (e. g., to increase the affinity and/or activity of the peptides), its molecular weights can be greater and can range anywhere from about 500 to about 50,000 Daltons.

10 Pharmaceutical Compositions

Also featured is a composition, e.g., a pharmaceutically acceptable composition, that includes a poly-PEGylated Kunitz domain. In one embodiment, the Kunitz domain binds to a protease such as elastase, plasmin, or kallikrein. As used herein, “pharmaceutical compositions” encompass compounds (e.g., labeled compounds) for
15 diagnostic (e.g., in vivo imaging) use as well as compounds for therapeutic or prophylactic use.

As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. In one
20 embodiment, the carrier is other than water. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the
25 compound.

A “pharmaceutically acceptable salt” refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects (see e.g., Berge, S.M., *et al.* (1977) *J. Pharm. Sci.* 66:1-19). Examples of such salts include acid addition salts and base addition salts. Acid
30 addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous and the like, as well as from nontoxic organic acids such as aliphatic mono- and

dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-
5 dibenzylethylenediamine, N-methylglucamine, chlorprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

The compositions of this invention may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (e.g., injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders,
10 liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for administration of humans with antibodies. The preferred mode of administration is parenteral (e.g., intravenous, subcutaneous, intraperitoneal, intramuscular). In a
15 preferred embodiment, the compound is administered by intravenous infusion or injection. In another preferred embodiment, the compound is administered by intramuscular or subcutaneous injection.

The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration,
20 usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

Pharmaceutical compositions typically must be sterile and stable under the
25 conditions of manufacture and storage. A pharmaceutical composition can also be tested to insure it meets regulatory and industry standards for administration. For example, endotoxin levels in the preparation can be tested using the Limulus amebocyte lysate assay (e.g., using the kit from Bio Whittaker lot # 7L3790, sensitivity 0.125 EU/mL) according to the USP 24/NF 19 methods. Sterility of pharmaceutical
30 compositions can be determined using thioglycollate medium according to the USP 24/NF 19 methods. For example, the preparation is used to inoculate the thioglycollate medium and incubated at 35°C for 14 or more days. The medium is inspected periodically to detect growth of a microorganism.

The composition can be formulated as a solution, microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients
5 enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and
10 freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by
15 including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

The poly-PEGylated Kunitz domains described herein can be administered by a variety of methods known in the art. For many applications, the route/mode of administration is intravenous injection or infusion. For example, for therapeutic
20 applications, the compound can be administered by intravenous infusion at a rate of less than 30, 20, 10, 5, or 1 mg/min to reach a dose of about 1 to 100 mg/m² or 7 to 25 mg/m². The route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release
25 formulation, including implants, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known. *See, e.g., Sustained and Controlled Release Drug Delivery Systems*, J.R.
30 Robinson, ed., Marcel Dekker, Inc., New York, 1978. Pharmaceutical formulation is a well-established art, and is further described in Gennaro (ed.), Remington: The Science and Practice of Pharmacy, 20th ed., Lippincott, Williams & Wilkins (2000) (ISBN: 0683306472); Ansel et al., Pharmaceutical Dosage Forms and Drug Delivery Systems,

7th Ed., Lippincott Williams & Wilkins Publishers (1999) (ISBN: 0683305727); and Kibbe (ed.), Handbook of Pharmaceutical Excipients American Pharmaceutical Association, 3rd ed. (2000) (ISBN: 091733096X).

In certain embodiments, the composition may be orally administered, for example, with an inert diluent or an assimilable edible carrier. The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

Pharmaceutical compositions can be administered with medical devices known in the art. For example, in a preferred embodiment, a pharmaceutical composition of the invention can be administered with a needleless hypodermic injection device, such as the devices disclosed in U.S. Patent Nos. 5,399,163, 5,383,851, 5,312,335, 5,064,413, 4,941,880, 4,790,824, or 4,596,556. Examples of well-known implants and modules useful in the present invention include: U.S. Patent No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Patent No. 4,486,194, which discloses a therapeutic device for administering medicants through the skin; U.S. Patent No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Patent No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Patent No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and U.S. Patent No. 4,475,196, which discloses an osmotic drug delivery system. Of course, many other such implants, delivery systems, and modules are also known.

In certain embodiments, the compound can be formulated to ensure proper distribution *in vivo*. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds of the invention cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g., U.S. Patents 4,522,811; 5,374,548; and 5,399,331. The liposomes may comprise one or more moieties that are selectively

transported into specific cells or organs, thus enhance targeted drug delivery (see, e.g., V.V. Ranade (1989) *J. Clin. Pharmacol.* 29:685).

Also within the scope of the invention are kits comprising poly-PEGylated Kunitz domain and instructions for use, e.g., treatment, prophylactic, or diagnostic use.

5 In one embodiment, the kit includes (a) the compound, e.g., a composition that includes the compound, and, optionally, (b) informational material. The informational material can be descriptive, instructional, marketing or other material that relates to the methods described herein and/or the use of the compound for the methods described herein. For example, in the case of a Kunitz domain that inhibits elastase activity, the informational
10 material describes methods for administering the compound to reduce elastase activity or to treat or prevent a pulmonary disorder (e.g., CF or COPD), an inflammatory disorder (e.g., IBD), or a disorder characterized by excessive elastase activity.

In one embodiment, the informational material can include instructions to administer the compound in a suitable manner, e.g., in a suitable dose, dosage form, or
15 mode of administration (e.g., a dose, dosage form, or mode of administration described herein). In another embodiment, the informational material can include instructions for identifying a suitable subject, e.g., a human, e.g., a human having, or at risk for a disorder characterized by excessive elastase activity. The informational material can include information about production of the compound, molecular weight of the
20 compound, concentration, date of expiration, batch or production site information, and so forth. The informational material of the kits is not limited in its form. In many cases, the informational material, e.g., instructions, is provided in printed matter, e.g., a printed text, drawing, and/or photograph, e.g., a label or printed sheet. However, the informational material can also be provided in other formats, such as Braille, computer
25 readable material, video recording, or audio recording. In another embodiment, the informational material of the kit is a link or contact information, e.g., a physical address, email address, hyperlink, website, or telephone number, where a user of the kit can obtain substantive information about the compound and/or its use in the methods described herein. Of course, the informational material can also be provided in any
30 combination of formats.

In addition to the compound, the composition of the kit can include other ingredients, such as a solvent or buffer, a stabilizer or a preservative, and/or a second agent for treating a condition or disorder described herein, e.g. a pulmonary (e.g., CF or

COPD) or inflammatory (e.g., IBD or RA) disorder. Alternatively, the other ingredients can be included in the kit, but in different compositions or containers than the compound. In such embodiments, the kit can include instructions for admixing the compound and the other ingredients, or for using the compound together with the other
5 ingredients.

The compound can be provided in any form, e.g., liquid, dried or lyophilized form. It is preferred that the compound be substantially pure and/or sterile. When the compound is provided in a liquid solution, the liquid solution preferably is an aqueous solution, with a sterile aqueous solution being preferred. When the compound is
10 provided as a dried form, reconstitution generally is by the addition of a suitable solvent. The solvent, e.g., sterile water or buffer, can optionally be provided in the kit.

The kit can include one or more containers for the composition containing the compound. In some embodiments, the kit contains separate containers, dividers or compartments for the composition and informational material. For example, the
15 composition can be contained in a bottle, vial, or syringe, and the informational material can be contained in a plastic sleeve or packet. In other embodiments, the separate elements of the kit are contained within a single, undivided container. For example, the composition is contained in a bottle, vial or syringe that has attached thereto the informational material in the form of a label. In some embodiments, the kit
20 includes a plurality (e.g., a pack) of individual containers, each containing one or more unit dosage forms (e.g., a dosage form described herein) of the compound. For example, the kit includes a plurality of syringes, ampules, foil packets, or blister packs, each containing a single unit dose of the compound. The containers of the kits can be air tight, waterproof (e.g., impermeable to changes in moisture or evaporation), and/or
25 light-tight.

In one embodiment wherein the compound contains a polypeptide that binds to an elastase, the instructions for diagnostic applications include the use of the compound to detect elastase, *in vitro*, e.g., in a sample, e.g., a biopsy or cells from a patient having a pulmonary disorder, or *in vivo*. In another embodiment, the instructions for
30 therapeutic applications include suggested dosages and/or modes of administration in a patient with a pulmonary disorder. The kit can further contain a least one additional reagent, such as a diagnostic or therapeutic agent, e.g., a diagnostic or therapeutic agent as described herein, and/or one or more additional agents to treat the pulmonary

disorder (e.g., another elastase inhibitor), formulated as appropriate, in one or more separate pharmaceutical preparations.

Treatments

A poly-PEGylated Kunitz domain has therapeutic and prophylactic utilities.

5 In one embodiment, poly-PEGylated Kunitz domain inhibits an elastase, e.g., a neutrophil elastase. The compound can be administered to a subject to treat, prevent, and/or diagnose a variety of disorders, such as diseases characterized by unwanted or aberrant elastase activity. For example, the disease or disorder can be characterized by enhanced elastolytic activity of neutrophils. The disease or disorder may result from an
10 increased neutrophil burden on a tissue, e.g., an epithelial tissue such as the epithelial surface of the lung. For example, the polypeptide that inhibits elastase can be used to treat or prevent pulmonary diseases such as cystic fibrosis (CF) or chronic obstructive pulmonary disorder (COPD), e.g., emphysema. The compound can also be administered to cells, tissues, or organs in culture, e.g. *in vitro* or *ex vivo*.

15 Poly-PEGylated Kunitz domains that inhibit other proteases can also be used to treat or prevent disorders associated with the activity of such other respective proteases.

As used herein, the term “treat” or “treatment” is defined as the application or administration of poly-PEGylated Kunitz domain, alone or in combination with, a second agent to a subject, e.g., a patient, or application or administration of the agent to
20 an isolated tissue or cell, e.g., cell line, from a subject, e.g., a patient, who has a disorder (e.g., a disorder as described herein), a symptom of a disorder or a predisposition toward a disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disorder, the symptoms of the disorder or the predisposition toward the disorder. Treating a cell refers to the inhibition, ablation,
25 killing of a cell *in vitro* or *in vivo*, or otherwise reducing capacity of a cell, e.g., an aberrant cell, to mediate a disorder, e.g., a disorder as described herein (e.g., a pulmonary disorder). In one embodiment, “treating a cell” refers to a reduction in the activity and/or proliferation of a cell, e.g., a leukocyte or neutrophil. Such reduction does not necessarily indicate a total elimination of the cell, but a reduction, e.g., a
30 statistically significant reduction, in the activity or the number of the cell.

As used herein, an amount of a poly-PEGylated Kunitz domain effective to treat a disorder, or a “therapeutically effective amount” refers to an amount of the compound

which is effective, upon single or multiple dose administration to a subject, in treating a subject, e.g., curing, alleviating, relieving or improving at least one symptom of a disorder in a subject to a degree beyond that expected in the absence of such treatment. For example, the disorder can be a pulmonary disorder, e.g., a pulmonary disorder
5 described herein.

A “locally effective amount” refers to the amount (e.g., concentration) of the compound which is effective at detectably modulating activity of a target protein (e.g., elastase) in a tissue, e.g., in a region of the lung exposed to elastase, or a elastase-producing cell, such as a neutrophil. Evidence of modulation can include, e.g.,
10 increased amount of substrate, e.g., reduced proteolysis of the extracellular matrix.

As used herein, an amount of poly-PEGylated Kunitz domain effective to prevent a disorder, or a “a prophylactically effective amount” of the compound refers to an amount of an elastase-binding compound, e.g., a polypeptide-polymer compound described herein, which is effective, upon single- or multiple-dose administration to the
15 subject, in preventing or delaying the occurrence of the onset or recurrence of a disorder, e.g., a pulmonary disorder.

The terms “induce,” “inhibit,” “potentiate,” “elevate,” “increase,” “decrease” or the like, e.g., which denote quantitative differences between two states, refer to a difference, e.g., a statistically significant difference (e.g., $P < 0.05$, 0.02, or 0.005),
20 between the two states.

Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is
25 especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical
30 carrier.

An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of a compound described herein is 0.1-20 mg/kg, more preferably 1-10 mg/kg. The compound can be administered by intravenous infusion at a rate of less

than 20, 10, 5, or 1 mg/min to reach a dose of about 1 to 50 mg/m² or about 5 to 20 mg/m². It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need
5 and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are only exemplary.

A pharmaceutical composition may include a “therapeutically effective amount” or a “prophylactically effective amount” of a compound described herein, e.g., a
10 compound that includes a polypeptide that binds and inhibits a protease (e.g., elastase). A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the composition may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the compound to elicit a
15 desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the composition is outweighed by the therapeutically beneficial effects. A “therapeutically effective dosage” preferably inhibits a measurable parameter, e.g., an increase in pulmonary function, relative to untreated subjects. The ability of a compound to inhibit a measurable parameter can be
20 evaluated in an animal model system predictive of efficacy in a human disorder. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit, such inhibition *in vitro* by assays known to the skilled practitioner, e.g., an assay described herein.

A “prophylactically effective amount” refers to an amount effective, at dosages
25 and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount may be less than the therapeutically effective amount.

As used herein, the term “subject” is intended to include human and non-human
30 animals. The term “non-human animals” of the invention includes all vertebrates, e.g., non-mammals (such as chickens, amphibians, reptiles) and mammals, such as non-human primates, sheep, dog, cow, pig, etc.

In one embodiment, the subject is a human subject. Alternatively, the subject can be a non-human mammal expressing a human neutrophil elastase or an endogenous non-human neutrophil elastase protein or an elastase-like antigen to which an elastase-binding compound cross-reacts. A compound of the invention can be administered to a human subject for therapeutic purposes (discussed further below). Moreover, an elastase-binding compound can be administered to a non-human mammal expressing the elastase-like antigen to which the compound binds (e.g., a primate, pig or mouse) for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal models may be useful for evaluating the therapeutic efficacy of the compound (e.g., testing of dosages and time courses of administration).

The subject method can be used on cells in culture, e.g. *in vitro* or *ex vivo*. The method can be performed on cells present in a subject, as part of an *in vivo* (e.g., therapeutic or prophylactic) protocol. For *in vivo* embodiments, the contacting step is effected in a subject and includes administering the elastase-binding compound to the subject under conditions effective to permit both binding of the compound to a target (e.g., an elastase) in the subject.

The compounds which inhibit elastase can reduce elastase-mediated degradation and its sequelae, such as persistent infection and inflammation, leading to destruction of tissue (e.g., destruction of airway epithelium).

Methods of administering compounds are described in "Pharmaceutical Compositions". Suitable dosages of the compounds used will depend on the age and weight of the subject and the particular drug used. The compounds can be used as competitive agents to inhibit, reduce an undesirable interaction, e.g., between a natural or pathological agent and the elastase, e.g., between the extracellular matrix and elastase.

In one embodiment, the compounds are used to kill or ablate cells that express elastase *in vivo*. The compounds can be used by themselves or conjugated to an agent, e.g., a cytotoxic drug, radioisotope. This method includes: administering the compound alone or attached to a cytotoxic drug, to a subject requiring such treatment.

The terms "cytotoxic agent" and "cytostatic agent" refer to agents that have the property of inhibiting the growth or proliferation (e.g., a cytostatic agent), or inducing the killing of cells.

Poly-PEGylated Kunitz domain may also be used to deliver a variety of drugs including therapeutic drugs, a compound emitting radiation, molecules of plants, fungal, or bacterial origin, biological proteins, and mixtures thereof. For example, the Kunitz domain can be used to target the payload to a region of a subject which includes
5 a protease that specifically interacts with the Kunitz domain.

Enzymatically active toxins and fragments thereof are exemplified by diphtheria toxin A fragment, nonbinding active fragments of diphtheria toxin, exotoxin A (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, α -sacrin, certain *Aleurites fordii* proteins, certain Dianthin proteins, *Phytolacca americana*
10 proteins (PAP, PAPII and PAP-S), *Morodica charantia* inhibitor, curcin, crotin, *Saponaria officinalis* inhibitor, gelonin, mitogillin, restrictocin, phenomycin, and enomycin. Procedures for preparing enzymatically active polypeptides of the immunotoxins are described in WO 84/03508 and WO 85/03508. Examples of cytotoxic moieties that can be conjugated to the antibodies include adriamycin,
15 chlorambucil, daunomycin, methotrexate, neocarzinostatin, and platinum.

In the case of polypeptide toxins, recombinant nucleic acid techniques can be used to construct a nucleic acid that encodes the polypeptide including a Kunitz domain and the cytotoxin (or a polypeptide component thereof) as translational fusions. The recombinant nucleic acid is then expressed, e.g., in cells and the encoded fusion
20 polypeptide isolated. Then the fusion protein is physically associated with a moiety that increases the molecular weight of the compound, e.g., to stabilize half-life in vivo, and then attached to a moiety (e.g., a polymer).

Procedures for conjugating proteins with the cytotoxic agents have been previously described. For conjugating chlorambucil with proteins, see, e.g., Flechner
25 (1973) *European Journal of Cancer*, 9:741-745; Ghose et al. (1972) *British Medical Journal*, 3:495-499; and Szekerke, et al. (1972) *Neoplasma*, 19:211-215. For conjugating daunomycin and adriamycin to proteins, see, e.g., Hurwitz, E. et al. (1975) *Cancer Research*, 35:1175-1181 and Arnon et al. (1982) *Cancer Surveys*, 1:429-449. For preparing protein-ricin conjugates, see, e.g., U.S. 4,414,148 and by Osawa, T., et al.
30 (1982) *Cancer Surveys*, 1:373-388 and the references cited therein. Coupling procedures as also described in EP 226 419.

Also encompassed by the present invention is a method of killing or ablating which involves using the compound for prophylaxis. For example, these materials can be used to prevent or delay development or progression of a lung disease.

Use of the therapeutic methods of the present invention to treat lung diseases has a number of benefits. Since the polypeptide portion of the compound specifically recognizes elastase, other tissue is spared and high levels of the agent are delivered directly to the site where therapy is required. Treatment in accordance with the present invention can be effectively monitored with clinical parameters. Alternatively, these parameters can be used to indicate when such treatment should be employed.

10 Pulmonary Disorders and Methods and Formulations

hNE inhibitor polypeptides that are physically associated with a moiety (e.g., a polymer) can be used to treat pulmonary disorders such as emphysema, cystic fibrosis, COPD, bronchitis, pulmonary hypertension, acute respiratory distress syndrome, interstitial lung disease, asthma, smoke intoxication, bronchopulmonary dysplasia, pneumonia, thermal injury, and lung transplant rejection.

Cystic Fibrosis. Cystic fibrosis (CF) is a genetic disease affecting approximately 30,000 children and adults in the United States. A defect in the CF gene causes the body to produce an abnormally thick, sticky mucus that clogs the lungs and leads to life-threatening lung infections. A diagnostic for the genetic disorder includes a sweat test which can include measuring chloride concentration in sweat collected on gauze or filter paper, measuring sodium concentration in sweat collected on gauze or filter paper, and pilocarpine delivery and current density in sweat collection. The gene that causes CF has been identified and a number of mutations in the gene are known.

In one embodiment, a hNE inhibitor polypeptide that is physically associated with a moiety (e.g., a polymer) is used to ameliorate at least one symptom of CF, e.g., to reduce pulmonary lesions in the lungs of a CF patient.

This compound can also be used to ameliorate at least one symptom of a chronic obstructive pulmonary disease (COPD). Emphysema, along with chronic bronchitis, is part of chronic obstructive pulmonary disease (COPD). It is a serious lung disease and is progressive, usually occurring in elderly patients. COPD causes over-inflation of structures in the lungs known as alveoli or air sacs. The walls of the alveoli break down resulting in a decrease in the respiratory ability of the lungs.

Patients with this disease may first experience shortness of breath and cough. One clinical index for evaluating COPD is the destructive index which measures a measure of alveolar septal damage and emphysema, and has been proposed as a sensitive index of lung destruction that closely reflects functional abnormalities, especially loss of elastic recoil. See, e.g., Am Rev Respir Dis 1991 Jul;144(1):156-9. The compound can be used to reduce the destructive index in a patient, e.g., a statistically significant amount, e.g., at least 10, 20, 30, or 40% or at least to within 50, 40, 30, or 20% of normal of a corresponding age and gender-matched individual.

In one aspect, the invention provides a composition that poly-PEGylated Kunitz domain that is an hNE inhibitor for treatment of a pulmonary disorder (e.g., cystic fibrosis, COPD). The composition can be formulated for inhalation or other mode of pulmonary delivery. Accordingly, the compounds described herein can be administered by inhalation to pulmonary tissue. The term "pulmonary tissue" as used herein refers to any tissue of the respiratory tract and includes both the upper and lower respiratory tract, except where otherwise indicated. A hNE inhibitor polypeptide that is physically associated with a moiety (e.g., a polymer) can be administered in combination with one or more of the existing modalities for treating pulmonary diseases.

In one example the compound is formulated for a nebulizer. In one embodiment, the compound can be stored in a lyophilized form (e.g., at room temperature) and reconstituted in solution prior to inhalation. In another embodiment, the compound is stored at an acidic pH (e.g., a pH less than 5, 4, or 3) and then combined with a neutralizing buffer having a basic pH prior to inhalation.

It is also possible to formulate the compound for inhalation using a medical device, e.g., an inhaler. See, e.g., U.S. 6,102,035 (a powder inhaler) and 6,012,454 (a dry powder inhaler). The inhaler can include separate compartments for the active compound at an acidic pH and the neutralizing buffer and a mechanism for combining the compound with a neutralizing buffer immediately prior to atomization. In one embodiment, the inhaler is a metered dose inhaler.

The three common systems used to deliver drugs locally to the pulmonary air passages include dry powder inhalers (DPIs), metered dose inhalers (MDIs) and nebulizers. MDIs, the most popular method of inhalation administration, may be used to deliver medicaments in a solubilized form or as a dispersion. Typically MDIs

comprise a Freon or other relatively high vapor pressure propellant that forces aerosolized medication into the respiratory tract upon activation of the device. Unlike MDIs, DPIs generally rely entirely on the inspiratory efforts of the patient to introduce a medicament in a dry powder form to the lungs. Nebulizers form a medicament aerosol to be inhaled by imparting energy to a liquid solution. Direct pulmonary delivery of drugs during liquid ventilation or pulmonary lavage using a fluorochemical medium has also been explored. These and other methods can be used to deliver a hNE inhibitor polypeptide that is physically associated with a moiety (e.g., a polymer).

For example, for administration by inhalation, poly-PEGylated Kunitz domain that inhibits hNE are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant or a nebulizer. The compound may be in the form of a dry particle or as a liquid. Particles that include the compound can be prepared, e.g., by spray drying, by drying an aqueous solution of the poly-PEGylated Kunitz domain that inhibits hNE with a charge neutralizing agent and then creating particles from the dried powder or by drying an aqueous solution in an organic modifier and then creating particles from the dried powder.

The compound may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, diethyltetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol, the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges for use in an inhaler or insufflator may be formulated containing a powder mix of the poly-PEGylated Kunitz domain that inhibits hNE and a suitable powder base such as lactose or starch, if the particle is a formulated particle. In addition to the formulated or unformulated compound, other materials such as 100% DPPC or other surfactants can be mixed with the poly-PEGylated Kunitz domain that inhibit hNE to promote the delivery and dispersion of formulated or unformulated compound. Methods of preparing dry particles are described, for example, in PCT Publication WO 02/32406.

The poly-PEGylated Kunitz domain that inhibits hNE, e.g., as dry aerosol particles, when administered can be rapidly absorbed and can produce a rapid local or systemic therapeutic result. Administration can be tailored to provide detectable activity within 2 minutes, 5 minutes, 1 hour, or 3 hours of administration. In some

embodiments, the peak activity can be achieved even more quickly, e.g., within one half hour or even within ten minutes. Alternatively, a poly-PEGylated Kunitz domain that inhibits hNE can be formulated for longer biological half-life can be used as an alternative to other modes of administration, e.g., such that the compound enters
5 circulation from the lung and is distributed to other organs or to a particular target organ.

In one embodiment, poly-PEGylated Kunitz domain that inhibits hNE is delivered in an amount such that at least 5% of the mass of the polypeptide is delivered to the lower respiratory tract or the deep lung. Deep lung has an extremely rich
10 capillary network. The respiratory membrane separating capillary lumen from the alveolar air space is very thin ($\leq 6 \mu\text{m}$) and extremely permeable. In addition, the liquid layer lining the alveolar surface is rich in lung surfactants. In other embodiments, at least 2%, 3%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, or 80% of the composition of a poly-PEGylated Kunitz domain that inhibits hNE is delivered to the lower
15 respiratory tract or to the deep lung. Delivery to either or both of these tissues results in efficient absorption of the compound and high bioavailability. In one embodiment, the compound is provided in a metered dose using, e.g., an inhaler or nebulizer. For example, the compound is delivered in a dosage unit form of at least about 0.02, 0.1, 0.5, 1, 1.5, 2, 5, 10, 20, 40, or 50 mg/puff or more.

20 The percent bioavailability can be calculated as follows: the percent bioavailability = $(\text{AUC}_{\text{non-invasive}} / \text{AUC}_{\text{i.v. or s.c.}}) \times (\text{dose}_{\text{i.v. or s.c.}} / \text{dose}_{\text{non-invasive}}) \times 100$.

Although not necessary, delivery enhancers such as surfactants can be used to further enhance pulmonary delivery. A "surfactant" as used herein refers to a compound having a hydrophilic and lipophilic moiety, which promotes absorption of a
25 drug by interacting with an interface between two immiscible phases. Surfactants are useful in the dry particles for several reasons, e.g., reduction of particle agglomeration, reduction of macrophage phagocytosis, etc. When coupled with lung surfactant, a more efficient absorption of the compound can be achieved because surfactants, such as DPPC, will greatly facilitate diffusion of the compound. Surfactants are well known in
30 the art and include but are not limited to phosphoglycerides, e.g., phosphatidylcholines, L-alpha-phosphatidylcholine dipalmitoyl (DPPC) and diphosphatidyl glycerol (DPPG); hexadecanol; fatty acids; polyethylene glycol (PEG); polyoxyethylene-9-; auryl ether;

palmitic acid; oleic acid; sorbitan trioleate (Span 85); glycocholate; surfactin; poloxomer; sorbitan fatty acid ester; sorbitan trioleate; tyloxapol; and phospholipids.

IBD and Methods and Formulations Therefor

In one embodiment, a poly-PEGylated Kunitz domain that inhibits hNE is used to ameliorate at least one symptom of an inflammatory bowel disease, e.g., ulcerative colitis or Crohn's disease.

Inflammatory bowel diseases (IBD) are generally chronic, relapsing intestinal inflammation. IBD refers to two distinct disorders, Crohn's disease and ulcerative colitis (UC). Both diseases may involve either a dysregulated immune response to GI tract antigens, a mucosal barrier breach, and/or an adverse inflammatory reaction to a persistent intestinal infection (see, e.g., MacDermott, R. P., J Gastroenterology, 31:907:-916 (1996)).

In patients with IBD, ulcers and inflammation of the inner lining of the intestines lead to symptoms of abdominal pain, diarrhea, and rectal bleeding. Ulcerative colitis occurs in the large intestine, while in Crohn's, the disease can involve the entire GI tract as well as the small and large intestines. For most patients, IBD is a chronic condition with symptoms lasting for months to years. The clinical symptoms of IBD are intermittent rectal bleeding, crampy abdominal pain, weight loss and diarrhea. Diagnosis of IBD is based on the clinical symptoms, the use of a barium enema, but direct visualization (sigmoidoscopy or colonoscopy) is the most accurate test.

Symptoms of IBD include, for example, abdominal pain, diarrhea, rectal bleeding, weight loss, fever, loss of appetite, and other more serious complications, such as dehydration, anemia and malnutrition. A number of such symptoms are subject to quantitative analysis (e.g. weight loss, fever, anemia, etc.). Some symptoms are readily determined from a blood test (e.g. anemia) or a test that detects the presence of blood (e.g. rectal bleeding). A clinical index can also be used to monitor IBD such as the Clinical Activity Index for Ulcerative Colitis. See also, e.g., Walmsley et al. *Gut*. 1998 Jul;43(1):29-32 and Jowett et al. (2003) *Scand J Gastroenterol*. 38(2):164-71.

In one embodiment, administration of the compound to a subject having or predisposed to having ulcerative colitis causes amelioration of the index, e.g., a

statistically significant change in the index. The compound includes hNE inhibitor polypeptide that is physically associated with a moiety (e.g., a hydrophilic polymer)

In one embodiment, administration of the compound to a subject having or predisposed to having IBD causes amelioration of at least one symptom of IBD.

5 Crohn's disease, an idiopathic inflammatory bowel disease, is characterized by chronic inflammation at various sites in the gastrointestinal tract. While Crohn's disease most commonly affects the distal ileum and colon, it may manifest itself in any part of the gastrointestinal tract from the mouth to the anus and perianal area. The prognosis and diagnosis of Crohn's disease can be measured using a clinical index, e.g.,
10 Crohn's Disease Activity Index. See, e.g., American Journal of Natural Medicine, July/Aug 1997, and Best WR, et al., "Development of a Crohn's disease activity index." *Gastroenterology* 70:439-444, 1976. In one embodiment, administration of the compound to a subject having or predisposed to having Crohn's disease causes amelioration of the index, e.g., a statistically significant change in the index, or
15 amelioration of at least one symptom of Crohn's disease.

Accordingly, in one aspect, the invention provides a composition that includes poly-PEGylated Kunitz domain that inhibits hNE for treatment of a bowel disease (e.g., a colitis such as ulcerative colitis, Crohn's disease or IBP) or other gastrointestinal or rectal disease. The composition can be formulated as a suppository. Suppositories can
20 be formulated with base ingredients such as waxes, oils, and fatty alcohols with characteristics of remaining in solid state at room temperatures and melting at body temperatures. The active ingredients of this invention with or without optional therapeutic ingredients, like hydrocortisone (1.0%), topical anesthetics like benzocaine (1.0 to 6.0%) or others as already listed may be prepared at appropriate pH values; for
25 example pH 5 liquid fatty alcohols, such as oleyl alcohol (range 45% to 65%) or solid higher fatty alcohols like cetyl or stearyl alcohol (30% to 50%). The base ingredients are well known in the art of this industry. See, e.g., U.S. 4,945,084 and 5,196,405.

The composition may also be used as an active ingredient in creams, lotions, ointments, sprays, pads, patches, enemas, foams and suppositories and others or in
30 delivery vehicles such as micro-encapsulation in liposomes or glycospheres. Other delivery technologies include microsponges or the substitute cell membrane (Completech™) which entrap the active ingredients for both protection and for slower

release. Rectal foams can be prepared as topical aerosol compositions may also be used, e.g., to treat (ulcerative colitis, Crohns colitis, and others).

Diagnostic Uses

A poly-PEGylated Kunitz domain has diagnostic utilities.

5 In one aspect, the present invention provides a diagnostic method for detecting the presence of a elastase protein, *in vitro* (e.g., a biological sample, such as tissue, biopsy or *in vivo* (e.g., *in vivo* imaging in a subject). The method includes: (i) contacting a sample with a poly-PEGylated Kunitz domain, e.g., a Kunitz domain that binds to a target protease, e.g., elastase, plasmin, or kallikrein; and (ii) detecting
10 formation of a complex between the elastase ligand and the sample. The method can also include contacting a reference sample (e.g., a control sample) with the ligand, and determining the extent of formation of the complex between the ligand and the sample relative to the same for the reference sample. A change, e.g., a statistically significant change, in the formation of the complex in the sample or subject relative to the control
15 sample or subject can be indicative of the presence of elastase in the sample.

Another method includes: (i) administering the compound to a subject; and (iii) detecting formation of a complex between the compound, and the target protease. The detecting can include determining location or time of formation of the complex.

The compound can be directly or indirectly labeled with a detectable substance
20 to facilitate detection of the bound or unbound antibody. Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials.

Complex formation between the compound and target protease can be detected by measuring or visualizing either the ligand bound to the target protease or unbound
25 ligand. Conventional detection assays can be used, e.g., an enzyme-linked immunosorbent assays (ELISA), a radioimmunoassay (RIA) or tissue immunohistochemistry. Further to labeling the compound, the presence of target protease can be assayed in a sample by a competition immunoassay utilizing standards labeled with a detectable substance and an unlabeled protease ligand. In one example
30 of this assay, the biological sample, the labeled standards and the compound are combined and the amount of labeled standard bound to the unlabeled ligand is

determined. The amount of target protease in the sample is inversely proportional to the amount of labeled standard bound to the compound.

Fluorophore and chromophore labeled protein ligands can be prepared. A variety of suitable fluorescers and chromophores are described by Stryer (1968) *Science*, 162:526 and Brand, L. et al. (1972) *Annual Review of Biochemistry*, 41:843-868. The protein ligands can be labeled with fluorescent chromophore groups by conventional procedures such as those disclosed in U.S. Patent Nos. 3,940,475, 4,289,747, and 4,376,110. One group of fluorescers having a number of the desirable properties described above is the xanthene dyes, which include the fluoresceins and rhodamines. Another group of fluorescent compounds are the naphthylamines. Once labeled with a fluorophore or chromophore, the protein ligand can be used to detect the presence or localization of the a target protease in a sample, e.g., using fluorescent microscopy (such as confocal or deconvolution microscopy).

Protein Arrays. The compound can also be immobilized on a protein array. The protein array can be used as a diagnostic tool, e.g., to screen medical samples (such as isolated cells, blood, sera, biopsies, and the like). Methods of producing polypeptide arrays are described, e.g., above.

In vivo Imaging. In still another embodiment, the invention provides a method for detecting the presence of a target protease or a target protease-expressing tissue *in vivo*. The method includes (i) administering to a subject (e.g., a patient having a pulmonary or respiratory disorder) a compound that includes a Kunitz domain and that is polyPEGylated, conjugated to a detectable marker; (ii) exposing the subject to a means for detecting said detectable marker to the target protease-expressing tissues or cells. For example, the subject is imaged, e.g., by NMR or other tomographic means.

Examples of labels useful for diagnostic imaging in accordance with the present invention include radiolabels such as ^{131}I , ^{111}In , ^{123}I , $^{99\text{m}}\text{Tc}$, ^{32}P , ^{125}I , ^3H , ^{14}C , and ^{188}Rh , fluorescent labels such as fluorescein and rhodamine, nuclear magnetic resonance active labels, positron emitting isotopes detectable by a positron emission tomography ("PET") scanner, chemiluminescers such as luciferin, and enzymatic markers such as peroxidase or phosphatase. Short-range radiation emitters, such as isotopes detectable by short-range detector probes can also be employed. The compound that includes the Kunitz domain can be labeled with such reagents using known techniques. For example, see Wensel and Meares (1983) *Radioimmunoimaging and*

Radioimmunotherapy, Elsevier, New York for techniques relating to the radiolabeling of proteins and D. Colcher et al. (1986) *Meth. Enzymol.* 121: 802-816.

A radiolabeled compound of this invention can also be used for *in vitro* diagnostic tests. The specific activity of an isotopically-labeled compound depends
5 upon the half-life, the isotopic purity of the radioactive label, and how the label is incorporated into the compound.

Procedures for labeling polypeptides (e.g., the polypeptide portion of the compound) with the radioactive isotopes (such as ^{14}C , ^3H , ^{35}S , ^{125}I , ^{32}P , ^{131}I) are generally known. For example, tritium labeling procedures are described in U.S. Patent
10 No. 4,302,438. Iodinating, tritium labeling, and ^{35}S labeling procedures, e.g., as adapted for murine monoclonal antibodies, are described, e.g., by Goding, J.W. (*Monoclonal antibodies : principles and practice : production and application of monoclonal antibodies in cell biology, biochemistry, and immunology* 2nd ed. London ; Orlando : Academic Press, 1986. pp 124-126) and the references cited therein. Other
15 procedures for iodinating polypeptides, are described by Hunter and Greenwood (1962) *Nature* 144:945, David et al. (1974) *Biochemistry* 13:1014-1021, and U.S. Patent Nos. 3,867,517 and 4,376,110. Radiolabeling elements which are useful in imaging include ^{123}I , ^{131}I , ^{111}In , and $^{99\text{m}}\text{Tc}$, for example. Procedures for iodinating polypeptides are described by Greenwood, F. et al. (1963) *Biochem. J.* 89:114-123; Marchalonis, J.
20 (1969) *Biochem. J.* 113:299-305; and Morrison, M. et al. (1971) *Immunochemistry* 289-297. Procedures for $^{99\text{m}}\text{Tc}$ -labeling are described by Rhodes, B. et al. in Burchiel, S. et al. (eds.), *Tumor Imaging: The Radioimmunochemical Detection of Cancer*, New York: Masson 111-123 (1982) and the references cited therein. Procedures suitable for ^{111}In -labeling antibodies are described by Hnatowich, D.J. et al. (1983) *J. Immunol.*
25 *Methods*, 65:147-157, Hnatowich, D. et al. (1984) *J. Applied Radiation*, 35:554-557, and Buckley, R. G. et al. (1984) *F.E.B.S.* 166:202-204.

In the case of a radiolabeled compound, the compound is administered to the patient, is localized to the tissue the antigen with which the compound interacts, and is detected or "imaged" *in vivo* using known techniques such as radionuclear scanning
30 using e.g., a gamma camera or emission tomography. See e.g., A.R. Bradwell et al., "Developments in Antibody Imaging", *Monoclonal Antibodies for Cancer Detection and Therapy*, R.W. Baldwin et al., (eds.), pp 65-85 (Academic Press 1985). Alternatively, a positron emission transaxial tomography scanner, such as designated

Pet VI located at Brookhaven National Laboratory, can be used where the radiolabel emits positrons (e.g., ^{11}C , ^{18}F , ^{15}O , and ^{13}N).

MRI Contrast Agents. Magnetic Resonance Imaging (MRI) uses NMR to visualize internal features of living subject, and is useful for prognosis, diagnosis, treatment, and surgery. MRI can be used without radioactive tracer compounds for obvious benefit. Some MRI techniques are summarized in EP-A-0 502 814. Generally, the differences related to relaxation time constants T1 and T2 of water protons in different environments is used to generate an image. However, these differences can be insufficient to provide sharp high resolution images.

10 The differences in these relaxation time constants can be enhanced by contrast agents. Examples of such contrast agents include a number of magnetic agents paramagnetic agents (which primarily alter T1) and ferromagnetic or superparamagnetic (which primarily alter T2 response). Chelates (e.g., EDTA, DTPA and NTA chelates) can be used to attach (and reduce toxicity) of some paramagnetic
15 substances (e.g., Fe^{+3} , Mn^{+2} , Gd^{+3}). Other agents can be in the form of particles, e.g., less than 10 μm to about 10 nM in diameter). Particles can have ferromagnetic, antiferromagnetic or superparamagnetic properties. Particles can include, e.g., magnetite (Fe_3O_4), $\gamma\text{-Fe}_2\text{O}_3$, ferrites, and other magnetic mineral compounds of transition elements. Magnetic particles may include: one or more magnetic crystals
20 with and without nonmagnetic material. The nonmagnetic material can include synthetic or natural polymers (such as sepharose, dextran, dextrin, starch and the like.

The compounds can also be labeled with an indicating group containing of the NMR-active ^{19}F atom, or a plurality of such atoms inasmuch as (i) substantially all of naturally abundant fluorine atoms are the ^{19}F isotope and, thus, substantially all
25 fluorine-containing compounds are NMR-active; (ii) many chemically active polyfluorinated compounds such as trifluoroacetic anhydride are commercially available at relatively low cost, and (iii) many fluorinated compounds have been found medically acceptable for use in humans such as the perfluorinated polyethers utilized to carry oxygen as hemoglobin replacements. After permitting such time for incubation, a
30 whole body MRI is carried out using an apparatus such as one of those described by Pykett (1982) *Scientific American*, 246:78-88 to locate and image cancerous tissues.

Also within the scope of the invention are kits comprising the compound that binds to a target protease and instructions for use, e.g., the use of the compound (e.g.,

poly-PEGylated Kunitz domain) to detect the target protease, *in vitro*, e.g., in a sample, e.g., a biopsy or cells from a patient having a pulmonary disorder, or *in vivo*, e.g., by imaging a subject. The kit can further contain a least one additional reagent, such as a label or additional diagnostic agent. For *in vivo* use the compound can be formulated as
 5 a pharmaceutical composition.

An exemplary amino acid sequence of a human neutrophil elastase:

(Also listed in GenBank® under: gi|4503549|ref|NP_001963.1| elastase 2, neutrophil [Homo sapiens])

10

MTLGRRRLACLFLACVLPALLLGGTALASEIVGGRRARPHAWPFMVSLQLRGGHFCGATLIAPNFVMSAAH
 CVANVNVRAVRVVLGAHNLSRREPTRQVFAVQRI FENGYDPVNLLNDIVILQLNGSATINANVQVAQLPA
 QGRRRLGNGVQCLAMGWGLLGRNRGIASVLQELNVTVVTSLCRRSNVCTLVRGRQAGVCFGDSGSPLVCNG
 LIHGIA SFVRGGCASGLYPDAFAPVAQFVNWIDSIIQRSEDNPCPHPRDPDPASRTH

15

The following non-limiting examples further illustrate aspects of the invention:

Example

Peptides and small proteins are rapidly cleared from circulation *in vivo*. The rapid clearance often greatly limits therapeutic potency. High doses and frequent
 20 administration are needed to achieve therapeutic effects.

DX-890 consists of 56 amino acids, contains three intramolecular disulfide bonds, and has a molecular weight of 6,237 Da. For primary amine-based coupling, there are five potential PEGylation sites on DX-890, each of the four lysine residues and the N-terminus. Use of mPEG succinimidyl propionic acid can be used to couple
 25 PEG to each of these sites, e.g., at four lysine residues and the N-terminus. The PEG reagent that can be used may be mPEG that has an average molecular weight of about 5 kDa.

The reaction can be allowed to proceed to completion at a pH that permits modification of the amino groups on the lysine side chains and to the N-terminus. For
 30 example, the pH can be greater than 7.5, e.g., between 7.8 and 8.5. The reaction is quenched, e.g., with Tris. The reaction can be loaded onto an ion exchange or size exclusion column and fractions that contain PEGylated DX-890 are collected. These relevant fractions can be dialyzed, further purified, and then stored or analyzed.

DX-1000, a human plasmin inhibitor, is a Kunitz domain with fewer lysines than DX-890. It has a three available lysines and an N-terminus for modification with mPEG. DX-1000 can be combined with an mPEG succinimidyl propionic acid reagent having an average molecular weight of about 5 kDa or 7 kDa. DX-1000 can be
5 modified and purified, e.g., as described for DX-890. US 6,103,499 also describes other plasmin inhibitors, including DX-1000 related inhibitors. Kunitz domains having sequences or conforming to motifs described in US 6,103,499 can be modified as described herein.

10 DX-88, a kallikrein inhibitor is a Kunitz domain with fewer lysines than DX-890. It has a three available lysines and an N-terminus for modification with mPEG. DX-88 can be combined with an mPEG succinimidyl propionic acid reagent having an average molecular weight of about 5 kDa or 7 kDa. DX-88 can be modified and purified, e.g., as described for DX-890. US 6,333,402 also describes other kallikrein
15 inhibitors, including DX-88 related inhibitors. See, e.g., Tables 6 and 103 described therein. Kunitz domains having sequences or conforming to motifs described in US 6,333,402 can be modified as described herein.

The predicted or actual structures of DX-890, DX-88, and DX-1000 are shown
20 with the lysine residues indicated in FIG. 1, 2, and 3, respectively.

Example

The Example above is further detailed by the following methods for PEGylating a protein of interest at multiple or all possible reactive sites, in the following
25 implementations, the method is used to poly-PEGylate Kunitz domains at multiple or all possible primary amines.

A 5 kDa amino-reactive monofunctional PEG (mPEG-SPA) from NEKTAR Therapeutics (cat. no.: 2M4M0H01) was used as material of the PEGylation reactions.

We found that it is possible to poly-PEGylate DX-88, DX-890 and DX-1000
30 with four or five 5 kDa PEGs. Moreover, the poly-PEGylated proteins maintained the desired therapeutic activity while having increased circulating half-life. Additionally, reaction conditions were very efficient in terms of the conversion of unmodified protein to the desired PEGylated form. The reactions can be used, or scaled up, to provide

consistently homogenous preparations of poly-PEGylated product. Because of this great efficiency and few reaction side products, preparations of the poly-PEGylated products can be synthesized with higher yield and lower cost than Kunitz domains that include a single PEG moiety. This approach makes for easier manufacturability with more controlled batch-to-batch consistency and a final product which is easier to fully characterize.

Materials

- mPEG-SPA, MW 5,000 Da, NEKTAR Therapeutics, cat. no.: 2M4M0H01 (succinimidyl ester of methoxy-capped polyethylene glycol propionic acid)
- DX-88 API, MW 7,054 Da, ~10 mg/ml in PBS, pH 7.0
- DX-890 API, MW 6,231 Da, ~10 mg/ml in 10mM NaAc, pH 3.0
- DX-1000 API, MW 7,167 Da, ~10 mg/ml in PBS, pH 7.0
- 0.2-0.3M Hepes, pH 7.8-8.5
- 1M Tris, pH 8.0
- 1N HCl

PEGylation Reaction I:

- 1) Calculate the amount of PEG needed for reacting the Kunitz domain polypeptide at approximately 10:1 molar ratio of PEG:reactive group. For example, DX-890 has 5 total reactive groups, so a 50:1 molar ratio of PEG:DX-890 is used. Depending upon the Kunitz domain polypeptide and/or reaction conditions, a ratio of 25:1 to 50:1 is typically used. For example, for PEGylating 10 mg of DX-890 (MW 6,231 Da) at a 50:1 molar ratio of PEG:peptide, 401 mg of PEG (MW 5,000 Da) would be used.
- 2) Just prior to reacting the Kunitz domain polypeptide with the PEG, dilute the required volume of Kunitz domain polypeptide stock 1:1 with 0.2M Hepes, pH 7.8-8.5 buffer. The peptide stock is typically ~10.0 mg/ml. Therefore upon dilution, the concentration of the Kunitz domain polypeptide is ~ 5.0 mg/ml in 0.1M Hepes, pH 7.8-8.5 buffer. Both DX-88 and DX-1000 are relatively stable in terms of solubility upon dilution. DX-890, while initially soluble upon dilution, however, may precipitate over time. Reaction times can be chosen to minimize precipitation.

- 3) Immediately add the 1:1 diluted Kunitz domain polypeptide solution directly to the PEG powder and quickly dissolve the PEG by vortexing. Once completely dissolved, cap the tube, wrap in foil and allow to react while slowly rocking/tumbling for 2.5-3 hours at 2-8°C to 25°C.
- 5 4) Quench the reaction by adding 1/9th volume of 1 M Tris, pH 8.0 for 30-60 minutes at 2-8°C to 25°C while slowly rocking/tumbling.
- 5) Carefully and slowly adjust the pH of the quench reaction mixture to ~ pH 7 with small additions of 1N HCl while mixing.
- 6) The neutralized reaction can be stored at 2-8°C or frozen at -20°C to -80°C until
10 purification.

The direct addition of the Kunitz domain polypeptide solution to PEG powder can help simplify the number of steps in the reaction process and reduce hydrolysis prior to reaction.

15 PEGylation Reaction II:

The following is another method for poly-PEGylating a polypeptide.

- 1) PEG is weighed out, as described for Reaction I, and placed aside for use just prior to reaction.
- 2) Dilute the Kunitz domain polypeptide to 3-5 mg/ml in 0.3M Hepes, pH 7.8-8.5.
- 20 3) Just prior to reaction, quickly prepare a 200-250 mg/ml solution of PEG (in slight excess) in dH₂O that has been previously degassed and N₂-saturated. Add the water to the PEG and quickly and completely dissolve by vortexing.
- 4) Immediately add the required volume of PEG solution to the Kunitz domain polypeptide solution while mixing. Cap the tube, wrap in foil and allow to react
25 while slowly rocking/tumbling for 2.5-3 hours at 2-8C to 25C.
- 5) Continue with steps 4) through 6) above.

Example: Analytical Methods

Modified Kunitz domains can be analyzed and characterized by a variety of
30 methods. Exemplary methods include the following:

The unpurified reaction mix may be analyzed for the extent of PEGylation by both reducing/non-reducing SDS-PAGE analysis with both Coomassie and iodine staining as described in a separate protocol and size-exclusion high performance liquid chromatography (SEC-HPLC) by monitoring both refractive index (RI) and absorbance at 280 nm (UV). The SDS-PAGE analysis by Coomassie stain detects only the polypeptide component of the reaction mix (free and coupled) whereas staining with iodine preferentially detects the PEG (free and coupled). SEC-HPLC analysis by UV (abs. 280 nm) detects the peptide (free and coupled) and RI detects both peptide and PEG. Dynamic light scattering (LS) detection allows for determination of absolute MW and MW distribution.

SDS-PAGE and SEC-HPLC can show the distribution of PEGylated products, but the absolute molecular weights should be determined by MALDI-TOF or other methods. The reason is that PEGylated proteins run more slowly on gels and SEC-HPLC than do unPEGylated proteins, due to the PEG moieties large hydrodynamic radius, leading to overestimation of molecular weight. This could be overcome by using PEGylated Kunitz domains of known absolute molecular weight as standards.

Iodine Staining

Gels are loaded with approximately 2-3 µg of protein initially (for DX-1000, DX-88, and DX-890) for PEGylated samples that will resolve into one or two bands only. This loading is most often appropriate for the 25:1 and 50:1 PEG:protein reactions if the coupling was successful. However, for samples that were PEGylated at the lower reaction ratios (1:1, 5:1, and 10:1) and are expected to exhibit multiple PEGylated species, 10-15 µg of protein per lane is more appropriate (since 4-5 bands may appear). Samples are mixed with the appropriate amount of NuPAGE LDS Sample Buffer. Samples are vortexed and heated at 70°C for 10 minutes prior to loading.

Gels can be prepared and resolved according to standard methods, e.g. using the Invitrogen NuPAGE system with a 4-12% Bis-Tris gels. See, e.g., NuPAGE Novex Bis-Tris Gels Quick Reference Card, Invitrogen Life Technologies.

Gels are rinsed briefly in deionized water, then covered with a 5% barium chloride solution for 10 minutes on the shaker. The gel is rinsed again with deionized

water and then immersed in a 0.1N Iodine solution for at least 10 minutes on the shaker. Bands should be visible almost immediately. Full staining will be complete after 10 minutes. The gel is then photographed, for example, with UVP Epi Chem II Darkroom and the Ethidium bromide filter.

- 5 After iodine staining, the protein can be stained for proteins with Coomassie. The gel is first rinsed in water to destain then mixed with Coomassie and then destained in 300 mL methanol, 100 mL glacial acetic acid, and 600 mL water. UnPEGylated protein bands appear dark blue, and PEGylated protein may appear very light blue, if at all.

10 ***Chromatography***

- The chromatography system (Waters Corporation) used here was the 600 system (pump/controller) running EMPOWER™ software with 717 plus auto sampler, 996 photodiode array detector (PDA) and 2414 refractive index detector. In addition, a PD2010+ dynamic light scattering (LS) detector (Precision Detectors, Inc.) was also
15 run in series.

- SEC column chromatography can include the following features: SEC column: TSK G3000SW_{xl} (7.8 mm ID x 30 cm L) with guard (Tosoh Bioscience, cat. no.: 08541 and 08543).; Flowrate: 0.5 ml/minute; Run time: 35 minutes; Mobile phase: PBS, pH 7.2 with 0.05% NaN₃.; Sample injection volume: 25 – 100 µL; Sample load:
20 50-100 µg per injection; Detection: UV (280 nm), RI and LS; SEC Standards: BioRad, cat. no.: 151-1901

MALDI-TOF

- MALDI-TOF (matrix-assisted laser desorption ionization-time of flight) Mass
25 Spectrometry (ABI, Applied Biosystems Voyager-DE) can be used to evaluate actual mass of reaction products and subjects. For polypeptide analysis (e.g., prior to reaction), alpha-cyano-4-hydroxycinnamic acid can be used as a matrix. For analysis of reaction products or poly-PEGylated speices, 2,5-dihydroxybenzoic acid (DHB) can be used as a matrix. Chips can be spotted 1:1 (0.5 µL:0.5 µL) of sample:matrix, and air
30 dried prior to analysis

K_i measurement

The equilibrium inhibition constants (K_i) for a poly-PEGylated protein (e.g., a poly-PEGylated DX-890) can be determined according to the tight-binding inhibition model with formation of a reversible complex (1:1 stoichiometry). Reactions are set up with 100 pM enzyme (e.g., elastase) and a range of inhibitor concentrations (0-4 nM) at 30°C in 50 mM HEPES, pH 7.5, 150 mM NaCl, and 0.1% Triton X-100. Following a 24 h incubation, substrate is added (25 μM) to the enzyme-inhibitor solution and the rate of substrate hydrolysis is monitored at an excitation of 360 nm and an emission of 460 nm. Plots of the percent remaining activity versus active inhibitor concentration are fit by nonlinear regression analysis to Equation 1 to determine equilibrium dissociation constants. Unmodified protein and poly-PEGylated protein can be analyzed for comparison.

$$\%A = 100 - \left(\frac{(I + E + K_i) - \sqrt{(I + E + K_i)^2 - 4 \cdot E \cdot I}}{2 \cdot E} \right) \cdot 100 \quad \text{Equation 1}$$

Where:

%A = percent activity

I = Kunitz domain protein concentration (e.g., DX-890)

E = enzyme (e.g., HNE) concentration

K_i = equilibrium inhibition constant

Pharmacokinetics in Animals

The following methods can be used to evaluate the pharmacokinetics (PK) of proteins such as poly-PEGylated proteins in animals, e.g., mice and rabbits.

25

The protein to be tested is labeled with iodine (¹²⁵I) using the iodogen method (Pierce).

The reaction tube is rinsed with reaction buffer (25 mM Tris, 0.4 M NaCl, pH 7.5).

The tube is emptied and then replaced with 0.1 ml of reaction buffer and 12 μl of carrier free iodine-126, about 1.6 mCi. After six minutes, the activated iodine is

30

transferred to a tube containing the protein to be tested. After nine minutes, the reaction is terminated with 25 μl of saturated tyrosine solution. The reaction can be

purified on a 5 ml D-salt polyacrylamide 6000 column in Tris/NaCl. HSA can be used to minimize sticking to the gel.

A sufficient number of mice (about 36) are obtained. The weight of each animal is recorded. In the case of mice, the animals are injected in the tail vein with about 5 μ g of the protein to be tested. Samples are recovered at each time point per animal, with four animals per time point, at approximately 0, 7, 15, 30, and 90 minutes, 4 hours, 8 hours, 16 hours, and 24 hours post injection. Samples (about 0.5 ml) are collected into anti-coagulant (0.02 ml EDTA). Cells are spun down and separated from plasma/serum. Samples can be analyzed by radiation counting and SEC peptide column on HPLC with inline radiation detection.

For rabbits, the material is injected into the ear vein. Samples can be collected at 0, 7, 15, 30, 90 minutes, 4, 8, 16, 24, 48, 72, 96, 120, and 144 hours post-injection. Samples can be collected and analyzed as for mice.

Data can be fit to a bi-exponential (equation 2) or a tri-exponential (equation 3) decay curve describing “fast”, “slow”, and “slowest” phases of *in vivo* clearance:

Equation 2

$$y = Ae^{-\alpha t} + Be^{-\beta t}$$

Equation 3

$$y = Ae^{-\alpha t} + Be^{-\beta t} + Ce^{-\gamma t}$$

Where:

y	=	Amount of label remaining in plasma at time = t post-administration
A	=	Total label in “fast” clearance phase
B	=	Total label in “slow” clearance phase
C	=	Total label in “slowest” clearance phase
α	=	“Fast” clearance phase decay constant
β	=	“Slow” clearance phase decay constant
γ	=	“Slowest” clearance phase decay constant
t	=	Time post administration

The α , β , and γ phase decay constants can be converted to half-lives for their respective phases as:

α Phase Half-life	=	0.69 (1/ α)
β Phase Half-life	=	0.69 (1/ β)
γ Phase Half-life	=	0.69 (1/ γ)

In the case where the data are fit using the bi-exponential equation, the percentages of the total label cleared from *in vivo* circulation through the α and β phases are calculated as:

$$\begin{aligned} 5 \quad \% \alpha \text{ Phase} &= [A/(A+B)] \times 100 \\ \% \beta \text{ Phase} &= [B/(A+B)] \times 100 \end{aligned}$$

In the case where the data are fit using the tri-exponential equation, the percentages of the total label cleared from *in vivo* circulation through the α and β phases are calculated as:

$$\begin{aligned} \% \alpha \text{ Phase} &= [A/(A+B+C)] \times 100 \\ \% \beta \text{ Phase} &= [B/(A+B+C)] \times 100 \\ \% \gamma \text{ Phase} &= [C/(A+B+C)] \times 100 \end{aligned}$$

15

Table 4: Plasma Clearance in Mice

	$T_{1/2 \alpha}$ (min.)	Clearance (%)	$T_{1/2 \beta}$ (min.)	Clearance (%)
DX-890	1.3	79	59.2	21
DX-1000	1.5	87	26.9	13
	$T_{1/2 \alpha}$ (hrs.)	Clearance (%)	$T_{1/2 \beta}$ (hrs.)	Clearance (%)
5xPEG5-DX-890	1.1	33	20.2	67
4xPEG5-DX-1000	0.3	38	12.5	62

20

Table 5: Plasma Clearance in Rabbit

	$T_{1/2 \alpha}$ (min.)	Clearance (%)	$T_{1/2 \beta}$ (hrs.)	Clearance (%)	$T_{1/2 \gamma}$ (hrs.)	Clearance (%)
DX-890	1.7	83	3.4	17		
DX-1000	0.9	85	1	15		
5xPEG5-DX-890	2.8	28	4.5	34	97.6	38
4xPEG5-DX-1000	1.9	34	3	32	69.3	34

In the case of both the mouse (Table 4) and rabbit (Table 5), PEGylation of either DX-890 or of DX-1000 results in a decrease in the fraction of clearance through the alpha pathway. At the same time the fraction of clearance through the longer lived pathways (beta and gamma) increases.

5

The poly-PEGylated proteins also showed good *in vivo* stability by SEC analysis.

Purification:

One exemplary purification method is as follows:

- 10 1) Purification of polyPEGylated-protein from excess/unreacted PEG and trace amounts of both high molecular weight and lower molecular weight PEGylated species may be accomplished by ion-exchange chromatography on an AKTA Basic 10/100 chromatography system (Amersham).
- 15 2) For example, a column of appropriate size and capacity may be packed with a strong cation exchange resin (i.e.: Poros 50HS, Applied Biosystems, prod. code: 1-3359-11) in the case of at least PEGylated DX-88 and DX-1000.
- 3) Briefly, a volume of the PEGylation reaction mix is diluted 5-15 fold or as necessary, with water followed by pH adjustment to pH~3.0 with 1 M acetic acid (100-200 mM final) and conductivity <3 mS/cm.
- 20 4) The column is first equilibrated with 100 mM acetic acid, pH 3.0. Linear flowrate of 100 cm/hr.
- 5) Loaded and washed with same for ~5 column volumes. Linear flowrate during loading is 50 cm/hr.
- 6) The PEGylated protein is eluted from the column in a series of step gradients.
- 25 7) The first step elution is 100 mM acetic acid, with 20 mM NaCl, pH 3.2 to help remove HMW components (~20 CV at 100 cm/hr).
- 8) The second step elution is 100 mM acetic acid with 50 mM NaCl, pH 3.8 (~10 CV at 100 cm/hr) elutes the main product (i.e.: 4 x 5kDa PEG/peptide for DX-88 and DX-1000).
- 30 9) The third and final step elution is PBS, pH 7.2 (~5 CV 100 cm/hr) to help remove trace amounts of LMW PEGylated species.
- 10) Followed by 0.2M NaOH cleaning (~5 CV with contact time of 30 minutes).
- 11) Followed by column storage in 20% ethanol (~10 CV).

- 12) Fractions are collected across the profile and analyzed by SDS-PAGE prior to pooling.
- 13) The final pool of purified PEGylated protein is then UF/DF into PBS, pH 7.2 using conventional means available. The final material is then 0.22 μ m filtered, quantitated by abs. 280 nm (as previously described), aliquoted and frozen at -20°C to -80°C until use.

Another exemplary purification method, and one that can be used to purify poly-PEGylated DX-88, is as follows.

- Reaction products are loaded on a cation exchange column. Poros 50HS was found to have a fair binding capacity (~3 mg DX88-PEG5K/ml resin) at this small scale that would allow for separation of free PEG and a fairly concentrated eluate that includes the poly-PEGylated species. Conductivity can be maintained below 2 mS/cm. For example, a 9.5 cm AKTA Poros 50HS column (1.1cm w. x 10cm h.) can be used.
- The column is washed and cleaned to remove endotoxin and other contaminants. The column can be equilibrated and loaded in 10mM sodium acetate pH 3.5.

UF/DF and Final DX-88-PEG5K Pool Analysis

- Fractions containing poly-PEGylated DX-88 were pooled for a total sample volume of ~6 mL. The sample was buffer exchanged into 1X DPBS pH 7.2 (unmodified) from Invitrogen (endotoxin specification <0.25 EU/ml) and concentrated using two Amicon Ultra-15 Centrifugal Filter Devices with a molecular weight cut-off of 10,000 kDa and a centrifugal force of 4500 x g. The CENTRICONs™ were washed with 0.1 N NaOH (diluted from 1 N NaOH, Acros, for low endotoxin production) for one hour followed by several rinses with HyClone water prior to use. The final exchange factor was 300 fold into 1X DPBS. A total of 3 mL of concentrated and purified DX-88-PEG5K were recovered from the centricons. The final sample concentration, 4.97 mg/mL, was determined by diluting the sample 1:10 and measuring the O.D. 280 nm against 1X DPBS using an extinction coefficient for DX-88 of 0.954. The final sample pH was ~2 measured using Whatman pH Indicator strips (pH 0-14). All filtrates (33 mL total) were analyzed for protein content and had an O.D. 280 nm of 0.003 or less measured against 1X DPBS. The purified DX-88-PEG5K sample was aliquoted into 0.5 mL fractions (2.5 mg each) in sterile tubes and frozen at -80°C. The

1:10 diluted sample was analyzed by SDS-PAGE in a dilution series to estimate the purity of the main product of interest, 4-PEG5K-DX-88. The purity of 4-PEG5K-DX-88 is approximately 90%.

5 **DX-890.** We prepared a poly-PEGylated DX-890. Gel electrophoresis and chromatographic analysis indicated that a reaction with a 1:50 or 1:63 ratio of DX-890 to 5K PEG reagent produced a reaction product that was predominantly (>85%) a modified DX-890 with five attached PEG moieties. DX-890 pegylated under a variety of ratios maintained its specific activity relative to a control (about 10 U/mg).

10 Poly-pegylated DX-890 is predicted to have five PEG moieties (each having about 5,266 Daltons molecular weight) plus the mass of DX-890 (6,237 Daltons, theoretical; 6,229 Daltons, observed). The predicted total mass is 34,682 Daltons. The mass of the species observed by MALDI-TOF was about 34,219 Daltons, in agreement with the theoretical prediction, as the mass of individual PEG moieties can vary.

15

20 **DX-88.** We prepared a poly-PEGylated DX-88. Gel electrophoresis and chromatographic analysis indicated that a reaction with a 1:50 ratio of DX-890 to 5K PEG reagent at pH 7.8 produced a reaction product that was predominantly (>85%) a modified DX-88 with four attached PEG moieties.

25 Poly-pegylated DX-88 is predicted to have four PEG moieties (each having about 5,266 Daltons molecular weight) plus the mass of DX-88 (7,054 Daltons). The predicted total mass is 28,126 Daltons. The mass of the species observed by MALDI-TOF was about 29,680 Daltons, in agreement with the theoretical prediction, as the mass of individual PEG moieties can vary.

30 Other embodiments are within the following claims.

What is claimed is:

1. A compound comprising:
 - (i) a polypeptide that comprises a Kunitz domain that specifically binds to and inhibits a protease, wherein the Kunitz domain is selected from the group consisting of:
 - (a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;
 - (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24; and
 - (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25;
 - and
 - (ii) a plurality of polyethylene glycol moieties, wherein the average molecular weight of each of the polyethylene glycol moieties is less than 12 kDa, and wherein each of at least four primary amines of the polypeptide is attached to one of the moieties of the plurality.
2. The compound of claim 1 wherein the average molecular weight of each of the polyethylene glycol moieties is less than 8 kDa.
3. The compound of claim 1 wherein the average molecular weight of each of the polyethylene glycol moieties is between 3 and 8 kDa.
4. The compound of claim 1 wherein the polypeptide has a molecular weight less than 8 kDa, and the compound has a molecular weight greater than 16 kDa.
5. The compound of claim 1 wherein the plurality of polyethylene glycol moieties consists of five moieties, each attached to a different primary amine.
6. The compound of claim 1 wherein each lysine of the Kunitz domain is attached to a different polyethylene glycol moiety.

7. The compound of claim 6 wherein the polypeptide comprises an N-terminal primary amine, and each lysine of the Kunitz domain and the N-terminal primary amine is attached to a different polyethylene glycol moiety.

8. The compound of claim 1 wherein the protease is elastase.

9. The compound of claim 8 wherein the polypeptide comprises the amino acid sequences of the two binding loops of DX-890.

10. The compound of claim 1 wherein the protease is kallikrein.

11. The compound of claim 10 wherein the polypeptide comprises the amino acid sequences of the two binding loops of DX-88.

12. The compound of claim 1 wherein the protease is plasmin.

13. The compound of claim 12 wherein the polypeptide comprises the amino acid sequences of the two binding loops of DX-1000.

14. A preparation that comprises

(i) polypeptides that specifically bind and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of:

(a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;

(b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24; and

(c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25,

wherein the at least 80% of the polypeptides that comprise said Kunitz domain comprise a polyethylene glycol moiety attached to each of at least four primary amines in each of said

polypeptides, and wherein the average molecular weight of each of the attached polyethylene glycol moieties is less than 12 kDa, and

(ii) a pharmaceutically acceptable carrier.

15. The preparation of claim 14 wherein the average molecular weight of each of the attached polyethylene glycol moieties is less than 10 kDa.

16. The preparation of claim 15 wherein the average molecular weight of each of the attached polyethylene glycol moieties is less than 8 kDa.

17. The preparation of claim 14 wherein at least 95% of the polypeptides in the preparation have a polyethylene glycol moiety attached to at least four primary amines in each of the polypeptides.

18. The preparation of claim 14 wherein the at least 80% of the polypeptides in the preparation have a polyethylene glycol moiety attached to at least five primary amines in each of the polypeptides.

19. The preparation of claim 14 wherein the at least 80% of the polypeptides in the preparation have a polyethylene glycol moiety attached to each of the primary amines in each of the polypeptides.

20. The preparation of claim 14 wherein the at least 80% of the polypeptides in the preparation have a polyethylene glycol moiety attached to each of the primary amines in each of the polypeptides, and to the N-terminal amine of each of the polypeptides.

21. The preparation of claim 14 wherein each of the polypeptides in the preparation binds and inhibits the protease.

22. The preparation of claim 14 wherein 95% of the polypeptides in the preparation bind and inhibit the protease.

23. The preparation of claim 14 wherein the protease is elastase.
24. The preparation of claim 23 wherein at least 80% of the polypeptides in the preparation comprise the amino acid sequence of the binding loops of DX-890.
25. The preparation of claim 14 wherein the protease is kallikrein.
26. The preparation of claim 25 wherein at least 80% of the polypeptides in the preparation comprise the amino acid sequence of the binding loops of DX-88.
27. The preparation of claim 14 wherein the protease is plasmin.
28. The preparation of claim 27 wherein at least 80% of the polypeptides in the preparation comprise the amino acid sequence of the binding loops of DX-1000.
29. A preparation that comprises
- (i) polypeptides that specifically bind and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of:
 - (a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;
 - (b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24; and
 - (c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25,and wherein the at least 80% of the polypeptides comprise a polyethylene glycol moiety attached to each of at least five primary amines in each of the polypeptides and wherein the average molecular weight of each of the attached polyethylene glycol moieties is less than 12 kDa, and
 - (ii) a pharmaceutically acceptable carrier.

30. The preparation of claim 29, wherein the at least 80% of the polypeptides in the preparation comprise a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23.

31. The preparation of claim 29, wherein the at least 80% of the polypeptides in the preparation comprise a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24.

32. The preparation of claim 29, wherein the at least 80% of the polypeptides in the preparation comprise a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25.

33. A method of preparing a pegylated Kunitz domain, the method comprising:
providing a polypeptide that comprises a Kunitz domain that has at least one lysine in the framework region of the Kunitz domain, wherein the Kunitz domain is selected from the group consisting of:

(a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;

(b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24;

and

(c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25;

and

contacting the polypeptide with activated polyethylene glycol, of average molecular weight less than 12 kDa, under conditions in which a polyethylene glycol moiety is attached to each of at least four primary amines in each of said Kunitz domain, and wherein at least one polyethylene glycol moiety is attached to the N-terminal primary amine.

34. A method of preparing a pegylated Kunitz domain, the method comprising:
providing a polypeptide that comprises a Kunitz domain that has at least two primary amine groups in the framework region of the Kunitz domain wherein the Kunitz domain is selected from the group consisting of:

(a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;

(b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24; and

(c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25;

and

contacting the polypeptide with activated polyethylene glycol, of average molecular weight less than 12 kDa, under conditions in which a polyethylene glycol moiety is attached to each of at least four primary amines in each Kunitz domain.

35. The method of claim 34 wherein the primary amine of each lysine of each polypeptide is attached to a polyethylene glycol moiety.

36. The method of claim 34 wherein the yield of pegylated Kunitz domains is greater than 40%.

37. The method of claim 34 wherein at least five primary amines are attached to polyethylene glycol (PEG) moieties.

38. The method of claim 34 wherein the conditions for contacting are a pH greater than 7.5.

39. The method of claim 34 wherein a polyethylene glycol moiety is attached to the N-terminal primary amine.

40. The method of claim 34 wherein the conditions are such that at least 70% of the Kunitz domains are pegylated on at least five primary amines.

41. The method of claim 34 wherein the conditions are such that at least 85% of the Kunitz domains are pegylated on at least five primary amines.

42. The method of claim 34 wherein the conditions are such that at least 70% of the pegylated Kunitz domains have the same number of attached polyethylene glycol moieties, the moieties being attached at the same positions.

43. The method of claim 42 wherein the conditions are such that at least 85% of the pegylated Kunitz domains have the same number of attached polyethylene glycol moieties, the moieties being attached at the same positions.

44. The method of claim 34 further comprising formulating the pegylated polypeptide as a pharmaceutical composition.

45. A composition for treating a disorder in a subject having or suspected of having the disorder, the composition comprising the preparation of claim 14 or claim 29,

wherein the preparation of claim 14 or claim 29 comprises polypeptides comprising the amino acid sequence of DX-890 and the disorder is selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), and an inflammatory disorder,

wherein the preparation of claim 14 or claim 29 comprises polypeptides comprising the amino acid sequence of DX-88 and the disorder is selected from the group consisting of hemophilia, post-operative bleeding, peri-operative bleeding, and hereditary angioedema, or

wherein the preparation of claim 14 or claim 29 comprises polypeptides comprising the amino acid sequence of DX-1000 and the disorder is selected from the group consisting of fibrinolysis, fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis.

46. The composition of claim 45 wherein the protease is elastase and the polypeptides comprise the amino acid sequence of DX-890.

47. The composition of claim 46 wherein the disorder is cystic fibrosis, COPD, or an inflammatory disorder.

48. The composition of claim 45, wherein the protease is kallikrein and the polypeptides comprise the amino acid sequence of DX-88.

49. The composition of claim 48 wherein the disorder is hemophilia, post-operative bleeding, peri-operative bleeding, or hereditary angioedema.

50. The composition of claim 45, wherein the protease is plasmin and the polypeptides comprise the amino acid sequence of DX-1000.

51. The composition of claim 50 wherein the disorder is fibrinolysis, fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, or inappropriate androgenesis.

52. A preparation comprising:

(i) polypeptides that specifically bind to and inhibit a protease, wherein at least 80% of the polypeptides in the preparation comprise a Kunitz domain selected from the group consisting of:

(a) a peptide comprising the DX-890 amino acid sequence set forth in SEQ ID NO: 23;

(b) a peptide comprising the DX-88 amino acid sequence set forth in SEQ ID NO: 24; and

(c) a peptide comprising the DX-1000 amino acid sequence set forth in SEQ ID NO: 25,

and wherein the at least 80% of the polypeptides comprise a polyethylene glycol moiety attached to each lysine and the N-terminal primary amine of the polypeptides, and wherein the average molecular weight of each of the moieties is less than 12 kDa, and

(ii) a pharmaceutically acceptable carrier.

53. The preparation of claim 52 wherein the average molecular weight of each of the polyethylene glycol moieties is between 3 kDa and 10 kDa.

54. The preparation of claim 53 wherein the average molecular weight of each of the polyethylene glycol moieties is between 3 kDa and 8 kDa.

55. A composition for treating a disorder in a subject having the disorder or suspected of having the disorder, the composition comprising the preparation of claim 52;

wherein the preparation of claim 52 comprises polypeptides comprising the amino acid sequence of DX-890 and the disorder is selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), and an inflammatory disorder,

wherein the preparation of claim 52 comprises polypeptides comprising the amino acid sequence of DX-88 and the disorder is selected from the group consisting of hemophilia, post-operative bleeding, peri-operative bleeding, and hereditary angioedema, or

wherein the preparation of claim 52 comprises polypeptides comprising the amino acid sequence of DX-1000 and the disorder is selected from the group consisting of fibrinolysis, fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis.

56. Use of a composition for treating a disorder in a subject having the disorder or suspected of having the disorder, the composition comprising the preparation of claim 14, claim 29, or claim 52;

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-890 and the disorder is selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), and an inflammatory disorder,

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-88 and the disorder is selected from the group consisting of hemophilia, post-operative bleeding, peri-operative bleeding, and hereditary angioedema, or

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-1000 and the disorder is selected from the group consisting of fibrinolysis, fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis.



57. Use of a composition for the manufacture of a medicament for treating a disorder in a subject having the disorder or suspected of having the disorder, the composition comprising the preparation of claim 14, claim 29 or claim 52;

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-890 and the disorder is selected from the group consisting of cystic fibrosis, Chronic Obstructive Pulmonary Disease (COPD), and an inflammatory disorder,

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-88 and the disorder is selected from the group consisting of hemophilia, post-operative bleeding, peri-operative bleeding, and hereditary angioedema, or

wherein the preparation of claim 14, claim 29, or claim 52 comprises polypeptides comprising the amino acid sequence of DX-1000 and the disorder is selected from the group consisting of fibrinolysis, fibrinogenolysis, excessive bleeding associated with thrombolytics, post-operative bleeding, peri-operative bleeding, and inappropriate androgenesis.

FIG. 1








FIG. 2

5	10	15	20	25	30	35	40	45	50	55	60
.	.	.	.	.	.	.	.	.	.	.	.
E	A	M	H	S	E	C	A	F	K	A	D
D	G	P	C	R	A	H	P	R	W	F	N
I	F	T	R	Q	C	E	E	E	F	I	Y
G	G	C	E	G	N	Q	N	R	F	E	S
L	E	E	C	K	K	M	C	T	R	D	

**FIG. 3**

5	10	15	20	25	30	35	40	45	50	55	60
EAMHSECA	FKAETG	PCRARF	DRWFFN	IFTRQ	CEEF	IYGG	CEGN	QNR	FESL	EECK	KMCTRD



PEGylation Scheme

FIG. 4




FIG. 5




FIG. 6A

GP-HPLC

Refractive Index




FIG. 6B




FIG. 7


FIG. 8a




FIG. 8b


FIG. 8c

Allometric Extrapolation for $t_{1/2\beta}$





DX-890



5	10	15	20	25	30	35	40	45	50	55																																													
.	.	.	.	.	.	.	.	.	.	.																																													
E	A	C	N	L	P	I	V	R	G	P	C	I	A	F	F	P	R	W	A	F	D	A	V	K	G	K	C	V	L	F	P	Y	G	G	C	Q	G	N	G	N	K	F	Y	S	E	K	E	C	R	E	Y	C	G	V	P