



(86) Date de dépôt PCT/PCT Filing Date: 2014/05/01

(87) Date publication PCT/PCT Publication Date: 2014/11/06

(45) Date de délivrance/Issue Date: 2021/11/16

(85) Entrée phase nationale/National Entry: 2015/10/29

(86) N° demande PCT/PCT Application No.: AU 2014/000483

(87) N° publication PCT/PCT Publication No.: 2014/176636

(30) Priorité/Priority: 2013/05/01 (AU2013901516)


(51) Cl.Int./Int.Cl. *C07C 279/02* (2006.01),
A61K 31/12 (2006.01), *A61K 31/155* (2006.01),
A61K 31/341 (2006.01), *A61K 31/381* (2006.01),
A61K 31/4045 (2006.01), *A61K 31/4192* (2006.01),
A61K 31/44 (2006.01), *A61K 31/498* (2006.01),
A61P 31/04 (2006.01), *C07C 241/04* (2006.01),
C07C 249/14 (2006.01), ...

(72) Inventeurs/Inventors:
PAGE, STEPHEN, AU;
GARG, SANJAY, AU;
KEENAN, MARTINE, AU;
MCCLUSKEY, ADAM, AU;
STEVENS, ANDREW, AU

(73) Propriétaire/Owner:

(54) Titre : COMPOSES ET METHODES DE TRAITEMENT D'INFECTIONS

(54) Title: COMPOUNDS AND METHODS OF TREATING INFECTIONS



Formula I

(57) **Abrégé/Abstract:**

The invention provides compounds of Formula (I), and methods of treating or preventing a bacterial infection in a subject using a compound of Formula (I). The invention also provides the use of a compound of Formula (I) in the manufacture of a medicament

(51) Cl.Int./Int.Cl. (suite/continued) *C07C 275/68* (2006.01), *C07C 281/08* (2006.01), *C07C 281/18* (2006.01), *C07C 337/08* (2006.01), *C07D 401/12* (2006.01), *C07D 403/12* (2006.01), *C07D 407/12* (2006.01), *C07D 409/12* (2006.01)

(73) Propriétaires(suite)/Owners(continued):NEOCULI PTY LTD, AU

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(57) Abrégé(suite)/Abstract(continued):

for the treatment of a bacterial infection in a subject. The invention further provides a medical device when used in a method of treating or preventing a bacterial infection in a subject and to a medical device comprising the composition of the invention.

ABSTRACT

5 The invention provides compounds of Formula (I), and methods of treating or preventing a bacterial infection in a subject using a compound of Formula (I). The invention also provides the use of a compound of Formula (I) in the manufacture of a medicament for the treatment of a bacterial infection in a subject. The invention further provides a medical device when used in a method of treating or preventing a bacterial infection in a subject and to a medical device comprising the composition of the invention.

COMPOUNDS AND METHODS OF TREATING INFECTIONS

TECHNICAL FIELD

[0001] This invention relates to compounds of Formula I, methods of treating or preventing a bacterial infection in a subject using a compound of Formula I, the use of a compound of Formula I in the manufacture of a medicament for the treatment of a bacterial infection in a subject, and medical devices when used in a method of treating or preventing a bacterial infection in a subject.

BACKGROUND ART

[0002] A marked increase in prevalence of multi-drug resistance in disease-causing Gram-positive (G+ve) (*Staphylococcus aureus*, *Enterococcus* spp. and *Streptococcus pneumoniae*) and Gram negative (G-ve) pathogens (*Escherichia coli*, *Enterobacter* spp., *Salmonella* spp., *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*) has coincided with an unprecedented global decline in investment in new anti-infective drugs. There are few currently registered alternatives for multidrug resistant (MDR) bacterial infections, forcing clinicians to consider older generation drugs such as colistin with narrow spectrum and considerable potential for toxic side-effects. In addition, there are fewer novel classes of anti-infective therapeutics moving through the drug development pipeline.

[0003] Since the year 2000, a period of almost 15 years, only 5 novel mode of action (MOA) antibacterial agents have been approved by the US FDA – linezolid (an oxazolidinone) in 2000, daptomycin (a lipopeptide) in 2003, retapamulin (a pleuromutilin) in 2007, fidaxomicin (a macrolide tiacumicin) in 2011, and bedaquiline (a diarylquinoline) in 2012. Notably, none of these agents has significant activity against gram negative bacteria. No novel MOA antibacterial agents were approved in 2013 and to date in 2014 only tedizolid and dalbavancin, both analogs of existing classes, have been recommended for approval in the US. While there are more than 300 anti-infective medicines in various stages of development, the large majority of these medicines are previously approved antibacterial compounds or their derivatives that are undergoing studies for new indications.

[0004] Furthermore, the prevalence of multidrug-resistance in animal-specific pathogens together with greater regulation of the registration and usage of antimicrobials in animals, has caused veterinarians to become increasingly reliant on the traditional classes of antimicrobial agents. The risk of transfer of MDR zoonotic organisms from animals to humans has also led to calls for further restrictions on the usage of some recently registered antibacterial drugs such as the fluoroquinolones and the third and fourth generation cephalosporins.

Epidemiology of antibacterial resistance development in pathogens of humans and animals

[0005] Much of the evolution in resistance development is driven by changes in the epidemiology of key MDR organisms. Once only restricted to human hospitals and aged care facilities, methicillin resistant *Staphylococcus aureus* (MRSA) strains are now being isolated from the community in alarming proportions. Furthermore, community-acquired MRSA strains are more likely to carry the Pantone–Valentine leukocidin (PVL) toxin, a virulence factor linked to skin and soft tissue lesions as well as a rapid, fulminating, necrotizing pneumonia with significant associated mortality. Recently MRSA strains have become host-adapted in several key animal species including livestock, horses and companion animals and regular cases of human-to-animal and animal-to-human transfer are being documented. This has important consequences for strain transmission and public health. A recent survey of 751 Australian veterinarians for MRSA nasal carriage found that a remarkable 21.4% of equine veterinarians were MRSA-positive compared to 4.9% of small animal veterinarians and 0.9% of veterinarians with little animal contact. These ecological shifts of MRSA together with the emergence of resistance to new drugs developed specifically for MRSA such as linezolid, confirm that new MRSA anti-infectives are urgently needed. Furthermore, hospitals that use vancomycin for treating MRSA then have to contend with outbreaks of vancomycin-resistant enterococci (VRE) infections in their patients, once again with limited alternative antimicrobial choices.

[0006] The global emergence and spread within the community of highly virulent MDR Gram-negative (G-ve) bacteria such as *E. coli* O25b:ST131 confirms that bacterial pathogens can simultaneously evolve both virulence and resistance determinants. Echoing recent MRSA epidemiology, *E. coli* O25b:ST131, a major cause of urinary tract and bloodstream infections in humans, has now been isolated from extraintestinal infections in companion animals, and poultry. The increasing significance of *E. coli* O25b:ST131 and other MDR Enterobacteriaceae with combined resistance to fluoroquinolones and extended spectrum β -lactams and carbapenems is another worrying trend, especially considering there have been few recent breakthroughs in the development of G-ve spectrum anti-infectives apart from incremental advances in the carbapenem family.

[0007] The World Health Organisation has identified antibiotic resistance as one of the three major future threats to global health. A recent report from the US Centers for Disease Control and Prevention (CDC) estimated that “in the United States, more than two million people are sickened every year with antibiotic-resistant infections, with at least 23,000 dying as a result.” The extra medical costs, in the USA alone, associated with treating and managing a single case of antibiotic-resistant infection are estimated to be between US\$18,588 and US\$29,069 per year resulting in an overall direct cost to the US health system of over US\$20 billion annually. In addition, the cost to US households in terms of lost productivity is estimated at over US\$35

billion per annum. Twenty five thousand patients in the European Union (EU) still die annually from infection with MDR bacteria despite many EU countries having world's best practice hospital surveillance and infection control strategies. The EU costs from health care expenses and lost productivity associated with MDR infections are estimated to be at least €1.5 billion per year.


[0008] There is an unmet clinical need for antibacterial agents with novel mechanisms of action to supplement and replace currently available antibacterial agents, the efficacy of which is increasingly undermined by antibacterial resistance mechanisms. There additionally remains a need for alternative antibacterials in the treatment of infection by multi-resistant bacteria. However, as reported by the Infectious Diseases Society of America and the European Centre for Disease Control and Prevention, few new drugs are being developed that offer promising results over existing treatments (Infectious Diseases Society of America 2010, *Clinical Infectious Diseases*, 50(8):1081-1083).

[0009] It is an object of the present invention to overcome at least one of the failings of the prior art.

[0010] The discussion of the background art set out above is intended to facilitate an understanding of the present invention only. The discussion is not an acknowledgement or admission that any of the material referred to is or was part of the common general knowledge as at the priority date of the application.


SUMMARY OF INVENTION

[0011] According to one aspect of the invention, there is provided a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof:




Formula I

[0012] In one preferred embodiment, R₁ is H, cycloalkyl, Formula II, or Formula III;




Formula II




Formula III


wherein R_3 is H, NH_2 , NHNH_2 , $\text{O-CH}_2\text{-CH}_3$, NH-C(O)-phenyl , NH-chlorophenyl , $\text{NH-CH}_2\text{-chlorophenyl}$, $\text{NH-N=CH-cycloalkyl}$, Formula IV, Formula V or Formula VI;



Formula IV




Formula V



Formula VI

wherein A_0 is N, C, CH, or A_0 is C and A_0 is bonded to R_4 , via R_2 , to form a triazole ring;

wherein A_1 is N, C, NH, $=\text{CH-CH=N-}$, $=(\text{C}_6\text{H}_5)\text{C-CH=N-}$, or Formula VII;



Formula VII

A_2 is N, C, NH, N-C(O)-phenyl , or Formula VII;

wherein A_3 , A_4 , A_5 , A_6 , A_7 , A_8 , A_{11} , A_{12} , A_{13} , A_{14} , A_{15} , A_{16} , A_{17} , A_{18} , A_{19} , A_{20} , A_{21} , A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are independently C, O, N, NH, S;

wherein A_9 is C, O, N, NH, N-C(O)-O-CH₂-CH₃, N-C(O)-O-CH(CH₃)₂, N-C(O)-NH-CH₂-CH₃, N-C(O)-NH-CH₂-phenyl, N-C(O)-CH₂-CH₂-CH₂-CH₂-CH₂-CH₃, N-C(O)-CH₂-furan-2-yl;

wherein A_{10} is C, NH, -N=CH-CH=, -N=CH-C(C₆H₅)=;

wherein A_{22} is -CH(CH₃)-, -N-CH-, -N-C(CH₃)-, N-C(CH₂OH)-;

R_2 is H, COOH, CH₂NH₂, CH₂OH, CH₂NHNH₂, methyl, ethyl, propyl, butyl, cyclopentyl, or Formula VII and R_2 and R_3 are bonded together to form a pyrimidine, pyrazine or triazine ring, or R_2 and R_9 are bonded together to form a pyrrolidinyloxindole ring;

wherein R_4 is N, NH, O, S, or R_4 and A_6 are bonded, via R_2 , to form a triazole ring, or R_4 is N and R_4 and R_2 are bonded together to form a pyrimidine ring;

wherein R_7 is H, Cl, Br, F, OH, CH₃, OCH₃, SCH₃, CN, CCH, CF₃, OCF₃, SCF₃, NO₂, butyl, *t*-butyl, dimethylamino, phenyl, *n*-propyl, *i*-propyl, -NH-C(O)-CH₃, -CH=CH-COOH, piperazin-1-yl, or R_7 and R_8 are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring;

wherein R_8 , R_9 , R_{14} , R_{16} , R_{25} and R_{27} are independently H, OH, Cl, F, Br, CH₃, CN, OCH₃, COOH, NO₂, CF₃, R_8 and R_7 bond together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring, or benzene ring, R_{14} and R_{15} are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring, R_8 and R_9 are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring, or R_{14} and R_{13} are bonded together to form a substituted or unsubstituted saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring;

wherein R_5 , R_6 , R_{17} , R_{24} and R_{28} are independently H, O, OH, Cl, F, Br, NH₂, CH₃, CF₃, OCH₃, CN, NO₂, phenyl, -NH-CH(OH)-CH₃, -NH-C(O)-CH₃, or R_6 and R_8 are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring, or R_{13} and R_{14} are bonded together to form a substituted or unsubstituted saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring;

wherein R_{10} , R_{11} , R_{19} , R_{20} , R_{22} and R_{23} are independently H, Cl, or Br, or R_{10} and R_{11} are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring, or R_{19} and R_{20} are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring, or R_{22} and R_{23} are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring;

wherein R_{12} , R_{18} and R_{21} are independently H, COOH, CH_2NH_2 , CH_2OH , methyl, ethyl, propyl, butyl, cyclopentyl, or R_{12} and R_{13} are bonded together to form a pyrrolidinyloxindole ring;

wherein R_{15} and R_{26} are independently H, Cl, Br, F, OH, CH_3 , OCH_3 , SCH_3 , CN, CF_3 , OCF_3 , SCF_3 , NO_2 , CCH, *n*-butyl, *t*-butyl, dimethylamino, phenyl, *n*-propyl, *i*-propyl, $-NH-C(O)-CH_3$, $-CH=CH-COOH$, piperazin-1-yl, or R_{15} and R_{14} are bonded together to form a substituted or unsubstituted, saturated or unsaturated aliphatic ring, heterocyclic ring or benzene ring; and

wherein "----" is a double bond or a single bond.

[0013] The compound of Formula I is preferably a chloride salt.

[0014] In another aspect of the invention, there is provided a compound, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the list of compounds presented in Example 13. Where a salt is presented in Example 13, the invention covers both the salt as presented and the freebase of that salt, and stereoisomers, tautomers, other pharmaceutically acceptable salts, and also other prodrugs of the freebase.

[0015] Preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof,

wherein A_0 is C;

wherein A_1 is N; or Formula VII;

wherein A_2 is N; or NH;

wherein A_3 , A_4 , A_6 , A_7 , A_{11} , A_{12} , A_{14} , A_{15} , are N; or C;

wherein A_5 , A_{13} , A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C;

wherein A_8 and A_{21} are S;

wherein A_9 is NH;

wherein A_{10} is N;

wherein A_{22} is $-N-CH-$; $-N-C(CH_3)-$; or $-N-C(CH_2OH)-$;

wherein R_1 is H; Formula II; Formula III; cycloalkyl;

wherein R_2 is H; methyl; ethyl; CH_2NHNH_2 ; CH_2OH ; butyl; cyclopentyl; or Formula VII and R_2 is bonded to R_4 , to form a pyrimidine ring;

wherein R_3 is NH_2 ; Formula IV; Formula V; Formula VI; NH_2 ; $NH-N=CH-$ cycloalkyl; or $O-CH_2-CH_3$;

wherein R_4 is NH; O; S; or R_4 is N and R_4 and R_2 are bonded together to form a pyrimidine ring;;

wherein R_7 is H; F; Cl; CF_3 ; methyl; R_7 and R_8 are bonded together to form an unsubstituted, benzene ring; OH; *t*-butyl; phenyl; dimethylamino; *i*-propyl; *n*-propyl; CN; CCH; *n*-butyl; SCH_3 ; R_7 and R_8 are bonded together to form an unsubstituted, unsaturated heterocyclic ring; OCH_3 ; Br; OCF_3 ; piperazin-1-yl; or SCF_3 ;

wherein R_6 , R_8 , R_{14} , and R_{16} are independently H; OH; F; OCH_3 ; CF_3 ; methyl; Cl; CN; Br; R_8 and R_7 are bonded together to form an unsubstituted, benzene ring; R_8 and R_7 are bonded together to form an unsubstituted, unsaturated heterocyclic ring; R_{14} and R_{15} are bonded together to form an unsubstituted, benzene ring; or R_{14} and R_{15} are bonded together to form an unsubstituted, unsaturated heterocyclic ring;

wherein R_5 , R_9 , R_{13} , and R_{17} are independently H; OH; NH_2 ; Cl; F; OCH_3 ; OH; $-NH-CH(OH)-CH_3$;

wherein R_{12} is H; methyl; ethyl; CH_2OH ; or cyclopentyl;

wherein R_{15} is H; F; Cl; CF_3 ; methyl; R_7 and R_8 are bonded together to form an unsubstituted, benzene ring; OH; *t*-butyl; phenyl; dimethylamino; *i*-propyl; *n*-propyl; CN; CCH; *n*-butyl; SCH_3 ; R_{15} and R_{14} are bonded together to form an unsubstituted, unsaturated heterocyclic ring; OCH_3 ; Br; OCF_3 ; piperazin-1-yl; or SCF_3 ;

wherein R_{24} and R_{26} are independently H; OH; or Cl;

wherein R_{25} and R_{27} are independently H; or OH;

wherein R_{26} is H; CH_3 ; Br; Cl; OH; dimethylamino; $-O-P(O)(OEt)_2$; CF_3 ; or F; and

wherein "—" is independently a single or a double bond.

[0016] More preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the group comprising: NCL008; NCL009; NCL023; NCL025; NCL026; NCL029; NCL036; NCL037; NCL039; NCL040; NCL050; NCL061; NCL064; NCL065; NCL068; NCL075; NCL076; NCL078; NCL079; NCL080; NCL081; NCL084; NCL085; NCL086; NCL088; NCL089; NCL090; NCL092; NCL094; NCL095; NCL097; NCL098; NCL099; NCL101; NCL104; NCL105; NCL106; NCL108; NCL111; NCL112; NCL114; NCL115; NCL116; NCL118; NCL119; NCL121; NCL122; NCL123; NCL124; NCL125; NCL126; NCL130; NCL131; NCL132; NCL133; NCL135; NCL136; NCL137; NCL138; NCL139; NCL140; NCL141; NCL144; NCL145; NCL146; NCL147; NCL148; NCL150; NCL152; NCL153; NCL154; NCL156; NCL157; NCL158; NCL159; NCL161; NCL162; NCL164; NCL165; NCL166; NCL167; NCL168; NCL169; NCL170; NCL171; NCL172; NCL173; NCL174; NCL176; NCL177; NCL178; NCL179; NCL180; NCL181; NCL183; NCL184; NCL185; NCL186; NCL187; NCL188; NCL189; NCL190; NCL193; NCL194; NCL195; NCL196; NCL197; NCL198; NCL199; NCL200; NCL201; NCL202; NCL203; NCL204; NCL205; NCL206; NCL207; NCL208; NCL209; NCL210; NCL211; NCL212; NCL213; NCL215; NCL216; NCL217; NCL218; NCL219; NCL220; NCL221; NCL222; NCL223; NCL224; NCL225; NCL226; NCL227; NCL228; NCL229; and NCL230.

[0017] Even more preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the group comprising: NCL040; NCL078; NCL079; NCL080; NCL081; NCL084; NCL088; NCL089; NCL097; NCL099; NCL123; NCL146; NCL157; NCL158; NCL177; NCL179; NCL188; NCL193; NCL195; NCL196; NCL197; NCL199; NCL202; NCL204; NCL205; NCL215; NCL216; NCL217; NCL219; and NCL221.

[0018] Even more preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the group comprising: NCL078; NCL079; NCL080; NCL081; NCL084; NCL088; NCL097; NCL157; NCL158; NCL179; NCL188; NCL193; NCL195; NCL196; NCL199; NCL204; NCL216; NCL217; NCL219; and NCL221.


[0019] Even more preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the group

comprising: NCL089; NCL097; NCL157; NCL179; NCL188; NCL193; NCL195; NCL196; NCL216; NCL219; and NCL221.


[0020] Most preferably, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, selected from the group comprising: NCL097; NCL157; NCL179; NCL188; NCL195; and NCL196.

[0021] In one preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein the compound is not a compound selected from the group consisting of: NCL812, NCL001, NCL002, NCL003, NCL004, NCL005, NCL006, NCL007, NCL010, NCL011, NCL012, NCL013, NCL014, NCL015, NCL016, NCL017, NCL018, NCL019, NCL020, NCL021, NCL022, NCL024, NCL027, NCL028, NCL030, NCL031, NCL032, NCL033, NCL034, NCL035, NCL038, NCL041, NCL042, NCL043, NCL044, NCL045, NCL046, NCL047, NCL048, NCL049, NCL051, NCL052, NCL053, NCL054, NCL055, NCL056, NCL057, NCL058, NCL059, NCL060, NCL062, NCL063, NCL066, NCL067, NCL069, NCL070, NCL071, NCL072, NCL073, NCL074, NCL077, NCL082, NCL083, NCL087, NCL091, NCL093, NCL096, NCL100, NCL102, NCL103, NCL107, NCL109, NCL110, NCL113, NCL117, NCL120, NCL127, NCL128, NCL129, NCL134, NCL142, NCL143, NCL149, NCL151, NCL155, NCL160, NCL163, NCL175, NCL182, NCL191, NCL192, and NCL214.


[0022] In a preferred aspect of the invention, the compound of Formula I is not robenidine (also referenced in this specification as NCL812 and also known as 1,3-bis[(*E*)-(4-chlorophenyl)methyleneamino]guanidine), which has a structure as follows:




[0023] In one preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_7 , R_8 , R_{12} , R_{13} , R_{15} , R_{16} , R_{17} are H; R_4 is O; R_9 and R_{14} are CF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and all Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0024] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N and A_2 is NH; A_0 , A_3 , A_4 , A_5 , A_6 , and A_7 are C; R_2 , R_5 , R_6 , R_7 , and R_8 are H; R_4 is NH; R_9 is Cl; and "—" in Formula I between A_0 and A_1 and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0025] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{15} , R_{17} are H; R_4 is NH; R_7 and R_{16} are F; and "—" in Formula I between A_0 and A_1 , all Formula II and all Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0026] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_7 , R_8 , R_{12} , R_{13} , R_{15} , R_{16} , R_{17} are H; R_4 is NH; R_9 and R_{14} are F; and "—" in Formula I between A_0 and A_1 , all Formula II and all Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0027] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_7 , R_8 , R_{12} , R_{14} , R_{15} , R_{16} , R_{17} are H; R_4 is NH; R_9 and R_{13} are OCH_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and all Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0028] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_7, R_8, R_{12}, R_{13}, R_{15}, R_{16}, R_{17}$ are H; R_4 is NH; R_3 and R_{14} are OCH_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and all Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0029] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 , and A_7 are C; R_2, R_5, R_6, R_8 and R_9 are H; R_4 is NH; R_7 is Cl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0030] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}, R_{17}$ are H; R_4 is NH; R_7 and R_{15} are CF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0031] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}, R_{17}$ are H; R_4 is NH; R_7 and R_{15} are methyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0032] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_7, R_8, R_{12}, R_{14}, R_{15}, R_{16}, R_{17}$ are H; R_4 is NH; R_9 and R_{13} are methyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0033] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_7, R_9, R_{12}, R_{13}, R_{15}, R_{16}$ and R_{17} are H; R_4 is NH; R_8 and R_{14} are methyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0034] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 , and A_7 are C; R_2, R_5, R_6, R_8 , and R_9 are H; R_4 is NH; R_7 is CF_3 ; and "—" in Formula I between A_0 and A_1 , and all Formula


II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0035] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0 , A_3 , A_4 , A_5 , A_6 , and A_7 are C; R_2 , R_5 , R_6 , R_7 , and R_9 are H; R_4 is NH; R_8 is CF_3 ; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0036] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0 , A_3 , A_4 , A_5 , A_6 , and A_7 are C; R_2 , R_5 , R_6 , R_7 , and R_9 are H; R_4 is NH; R_8 is methyl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0037] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0 , A_3 , A_4 , A_5 , A_6 , and A_7 are C; R_2 , R_5 , R_6 , R_7 , and R_9 are H; R_4 is NH; R_8 is Cl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0038] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_7, R_8, R_{12}, R_{13}, R_{15}, R_{16}, R_{17}$ are H; R_4 is NH; R_9 and R_{14} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0039] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}, R_{17}$ are H; R_4 is NH; R_7 and R_{15} are CF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0040] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0041] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is $NHNH_2$; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 , and A_7 are C; R_2 is methyl; R_5, R_6, R_8 and R_9 are H; R_4 is NH; R_7 is Cl; and "—" in Formula I between A_0 and A_1 ,


and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0042] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; $R_2, R_5, R_6, R_7, R_9, R_{12}, R_{13}, R_{15}, R_{16}$, and R_{17} are H; R_4 is S; R_8 and R_{14} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0043] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 , and A_7 are C; R_2, R_5, R_6, R_8 , and R_9 are H; R_4 is NH; R_7 is Cl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0044] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is $NHNH_2$; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 , and A_7 are C; R_2 is methyl; R_5, R_6, R_8 , and R_9 are H; R_4 is NH; R_7 is CF_3 ; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0045] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , and R_{16} are H; R_4 is NH; R_7 , R_{15} and R_{17} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0046] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , and R_{16} are H; R_4 is NH; R_7 is Cl; R_{15} is CF_3 ; R_{17} is F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL078):




[0047] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} and R_{17} are H; R_4 is NH; R_7 is Cl; R_{15} is F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL079):




[0048] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof,


wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 is Cl; R_{12} is methyl; R_{15} is CF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL080):



[0049] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are Cl; R_{12} is methyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0050] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{15}$, and R_{16} are H; R_4 is NH; R_7 and R_{17} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0051] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}$, and R_{16} are H; R_4 is NH; R_7 and R_{15} are Cl; R_{17} is F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula


IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL084):




[0052] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{15}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 is Cl; R_{14} is CN; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0053] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{15}$, and R_{16} are H; R_4 is NH; R_7 is Cl; R_{17} is F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0054] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; R_2 and R_{12} are methyl; $R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 is Cl; R_{15} is CF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL089):




[0055] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_9, R_{12}, R_{13}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 and R_8 are bonded together to form an unsubstituted, benzene ring; R_{14} and R_{15} are bonded together to form an unsubstituted, benzene ring; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0056] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; A_1 is N; A_2 is NH; R_1 is cyclohexyl; R_3 is $\text{NH}-\text{N}=\text{CH}-\text{cyclohexyl}$; R_4 is NH; R_2 is H; and "—" in Formula I between A_0 and A_1 is a double bond. An example of a compound of this embodiment of the invention includes:




[0057] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_{12}, R_{13}$ and R_{17} are H; R_4 is NH; $R_8, R_7, R_9, R_{14}, R_{15}$, and R_{16} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL097):




[0058] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} and R_{17} are H; R_4 is NH; R_7 and R_{15} are *t*-butyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0059] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_7 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , and R_{16} are H; R_4 is NH; R_5 , R_6 , R_{10} , and R_{17} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0060] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_6 , R_8 , R_{12} , R_{14} , and R_{17} are H; R_4 is NH; R_5 , R_7 , R_9 , R_{13} , R_{15} , and R_{16} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0061] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof,


wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_8, R_{12}, R_{16}$, and R_{17} are H; R_4 is NH; $R_7, R_8, R_9, R_{13}, R_{14}$, and R_{15} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL097):



[0062] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_7, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{15}$, and R_{16} are H; R_4 is NH; R_6 and R_{17} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0063] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_7, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{15}$, and R_{17} are H; R_4 is NH; R_6 and R_{16} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0064] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_8, R_9, R_{12}, R_{13}, R_{14}$, and R_{17} are H; R_4 is NH; R_6 ,


R_7 , R_{15} , and R_{16} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0065] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are phenyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0066] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are dimethylamino; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0067] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_{12} , R_{13} , R_{14} , and R_{17} are H; R_4 is NH; R_8 and R_{16} are OCH_3 ; R_7 and R_{15} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II


and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0068] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are *i*-propyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0069] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are *n*-propyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0070] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , and R_{17} are H; R_4 is NH; R_7 , R_{15} , and R_{16} are F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0071] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are CCH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0072] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_7 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{15} , and R_{17} are H; R_4 is NH; R_6 and R_{16} are Br; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0073] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are butyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0074] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 is $-C(C_6H_5)-CH-N-$ and A_{10} is $-N=CH-C(C_6H_5)=$; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; $R_2, R_5, R_6, R_7, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{15}, R_{16}$, and R_{17} are H; R_4 is NH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0075] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 and R_{15} are CH_2S ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0076] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula III; R_3 is Formula VI; A_0 is C; R_2 and R_{21} are H; A_1 and A_{20} are N; A_2 and A_{19} are NH; A_8 and A_{21} are S; R_4 is NH; R_{10} and R_{11} are bonded together to form a substituted benzene ring; R_{22} and R_{23} are bonded together to form a substituted benzene ring; and "—" in Formula I between A_0 and A_1 , and all Formula III and Formula VI "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0077] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_5, R_6, R_7, R_8, R_9, R_{13}, R_{14}, R_{15}, R_{16}$, and R_{17} are H; R_4 is NH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0078] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_{12}, R_{13}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 and R_8 are bonded together to form an unsubstituted, heterocyclic ring; R_{14} and R_{15} are bonded together to form an unsubstituted, unsaturated heterocyclic ring; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0079] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 is $=CH-CH=N-$ and A_{10} is $-N-(CH)_2-$; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}$, and R_{17} are H; R_4 is NH; R_7 and R_{15} are OCH_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0080] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are OH; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0081] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 and R_{12} are ethyl; R_5 , R_6 , R_8 , R_9 , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0082] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 and R_{12} are methyl; R_5 , R_6 , R_8 , R_9 , R_{13} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are Br; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0083] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_5 , R_6 , R_8 , R_{12} , R_{14} , R_{16} , and R_{17} are H; R_4 is NH; R_7 and R_{15} are Cl; R_9 and R_{13} are NH_2 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL157):



[0084] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 and R_{12} are ethyl; R_5 and R_{17} are OH; R_6 , R_8 , R_9 , R_{13} , R_{14} , and R_{16} are H; R_4 is NH; R_7 and R_{15} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL158):




[0085] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 and R_{12} are cyclopentyl; R_5 and R_{17} are OH; R_6 , R_8 , R_9 , R_{13} , R_{14} , and R_{16} are H; R_4 is NH; R_7 and R_{15} are Cl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0086] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof,


wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are OCF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:



[0087] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_5, R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are piperazin-1-yl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0088] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is $\text{O-CH}_2\text{-CH}_3$; A_1 is N; A_2 is NH; A_0, A_3, A_4, A_5, A_6 and A_7 are C; R_2 is methyl; R_5, R_6, R_8 , and R_9 are H; R_4 is NH; R_7 is Cl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0089] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_5, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are SCF_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula


IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0090] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , A_7 , A_{11} , A_{12} , A_{13} , A_{14} and A_{15} , are C; R_2 , R_6 , R_8 , R_9 , R_{12} , R_{13} , R_{14} , and R_{18} are H; R_4 is NH; R_7 and R_{16} are Cl; R_5 and R_{17} are $-NH-CH(OH)-CH_3$; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes:




[0091] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-N-CH-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{20} is Cl. An example of a compound of this embodiment of the invention includes (NCL179):




[0092] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is NH_2 ; A_1 is N; A_2 and R_4 are NH; A_0 , A_3 , A_4 , A_5 , A_6 , and A_7 are C; R_5 , R_6 , R_8 , and R_9 are H; R_2 is butyl; R_7 is Cl; and "—" in Formula I between A_0 and A_1 , and all Formula II "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL188):




[0093] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-\text{N}-\text{CH}-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{28} is CH_3 . An example of a compound of this embodiment of the invention includes (NCL195):




[0094] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-\text{N}-\text{CH}-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{28} is OH. An example of a compound of this embodiment of the invention includes (NCL196):




[0095] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-N-CH-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{26} is Br. An example of a compound of this embodiment of the invention includes (NCL193):




[0096] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-N-CH-$; R_{24} , R_{25} , R_{26} , R_{27} and R_{28} are H; and A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C. An example of a compound of this embodiment of the invention includes (NCL199):




[0097] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_2 is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_0 , and between A_1 and A_2 are double bonds; A_{22} is $-N-C(CH_3)-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{26} is Cl. An example of a compound of this embodiment of the invention includes (NCL204):




[0098] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; $R_2, R_6, R_8, R_9, R_{12}, R_{13}, R_{14}$, and R_{16} are H; R_4 is NH; R_7 and R_{15} are Cl; R_5 and R_{17} are F; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL216):



[0099] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are CH_3 ; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL217):




[00100] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein R_1 is Formula II; R_3 is Formula IV; A_1 and A_{10} are N; A_2 and A_9 are NH; $A_0, A_3, A_4, A_5, A_6, A_7, A_{11}, A_{12}, A_{13}, A_{14}$ and A_{15} , are C; R_2 and R_{12} are methyl; $R_6, R_8, R_9, R_{13}, R_{14}, R_{16}$ and R_{17} are H; R_4 is NH; R_7 and R_{15} are *t*-butyl; and "—" in Formula I between A_0 and A_1 , all Formula II and Formula IV "—" are double bonds. An example of a compound of this embodiment of the invention includes (NCL219):



[00101] In another preferred embodiment of the invention, the compound is a compound of Formula I, or a stereoisomer, tautomer, pharmaceutically acceptable salt, or prodrug thereof, wherein A_0 is C; R_1 is H; A_2 and R_4 are N; R_3 is NH_2 ; A_1 is Formula VII; R_2 is Formula VII and R_5

is bonded to R_4 , forming a pyrimidine ring; "—" in Formula I between R_2 and A_6 , and between A_1 and A_2 are double bonds; A_{22} is $-N-CH-$; R_{24} , R_{25} , R_{27} and R_{28} are H; A_{23} , A_{24} , A_{25} , A_{26} and A_{27} are C; and R_{28} is CF_3 . An example of a compound of this embodiment of the invention includes (NCL221):



[00102] According to another aspect of the invention, there is provided a method of treating or preventing a bacterial colonisation or infection in a subject, the method comprising the step of administering a therapeutically effective amount of a compound of Formula I, or a therapeutically acceptable salt thereof, to the subject. In this aspect, the bacterial infection is caused by a bacterial agent. The method of treating or preventing a bacterial infection or colonisation in a subject, may also comprise the administration of the pharmaceutical or veterinary compositions of the invention.

[00103] According to a further aspect of the invention, there is provided the use of a compound of Formula I, or a therapeutically acceptable salt thereof, in the manufacture of medicament the treatment of a bacterial colonisation or infection in a subject. In this aspect, the bacterial infection is caused by a bacterial agent.

[00104] The subject may be any subject capable of colonisation and infection by bacteria. The subject may be mammalian, or may be piscine or avian. Preferably, the subject is selected from the group comprising, but not limited to, human, canine, feline, bovine, ovine, caprine, other ruminant species, porcine, equine, avian, or piscine.

[00105] The compound of Formula I may be administered to the subject in a dose selected from the group comprising 0.1 mg/kg to 250 mg/kg body weight, preferably 1 mg/kg to 100 mg/kg body weight, and more preferably 5 mg/kg to 50 mg/kg body weight. The compound of Formula I may be administered to the subject using a dosing schedule selected from the group consisting of: hourly, 3 times daily; twice daily; daily; every second day; twice weekly; once weekly; once fortnightly; once monthly; once every two months or by constant rate or variable rate infusion. Preferably, the compound of Formula I is administered until colonisation or the signs and symptoms of infection or colonisation have at least been partially treated or alleviated.

[00106] In one embodiment, the concentration of compound of Formula I (or a metabolite) in the subject's blood after treatment is within a range selected from the group comprising, but not limited to: between 0.1 and 10 ug/mL at 2 hours, 1 and 200 ug/mL after 12 hours; between 0.1 and 5 ug/mL after 24 h; between 0.01 and 2 ug/mL after 48 hours; between 0.0001 and 1 ug/mL after 72 h. Preferably, the concentration is selected from the group comprising, but not limited to: less than 200 ug/mL after 12 hours; less than 5 ug/mL after 24 hours; less than 1 ug/L after 48 hours and less than 0.5 ug/mL after 72 hours.

[00107] The agent causing the bacterial infection is a bacterial agent. In one preferred embodiment, the agent is not a protozoan species. In one preferred embodiment, the agent is not a coccidian protozoan. More preferably, the agent is not *Clostridium perfringens* nor a heterotrophic bacterial species present in soil samples collected by Hansen et al from Jyndevad Denmark as discussed in the following papers: Hansen et al. 2012, *Chemosphere*, 86:212-215; and Hansen et al. 2009, *Environmental Pollution* 157:474-480.

[00108] In another embodiment, the bacterial agent is gram negative. In another embodiment, the bacterial agent is gram positive. In another embodiment, the bacterial agent has no cell wall. In another embodiment, the bacterial infection is caused by a mixture of at least two agents selected from the group consisting of: gram negative, gram positive and bacterial agents with no cell wall.

[00109] The bacterial agent causing the bacterial infection may be a gram positive bacterial agent selected from the group comprising, but not limited to, *Staphylococcus* spp, Streptococci, *Enterococcus* spp, *Leuconostoc* spp, *Corynebacterium* spp, *Arcanobacteria* spp, *Trueperella* spp, *Rhodococcus* spp, *Bacillus* spp, Anaerobic Cocci, Anaerobic Gram-Positive Nonsporulating Bacilli, *Actinomyces* spp, *Clostridium* spp, *Nocardia* spp, *Erysipelothrix* spp, *Listeria* spp, *Kytococcus* spp, *Mycoplasma* spp, *Ureaplasma* spp, and *Mycobacterium* spp.

[00110] In one embodiment, the bacterial agent is gram positive and is selected from the group comprising, but not limited to, *Staphylococcus* spp. Examples of *Staphylococcus* spp include *Staphylococcus epidermidis*, *Staphylococcus haemolyticus*, *Staphylococcus lugdunensis*, *Staphylococcus saprophyticus*, *Staphylococcus auricularis*, *Staphylococcus capitis*, *Staphylococcus caprae*, *Staphylococcus carnosus*, *Staphylococcus cohnii*, *Staphylococcus hominis*, *Staphylococcus pasteurii*, *Staphylococcus pettenkoferi*, *Staphylococcus pulvereri*, *Staphylococcus saccharolyticus*, *Staphylococcus simulans*, *Staphylococcus schleiferi*, *Staphylococcus warneri*, *Staphylococcus xylosus*, *Staphylococcus arlettae*, *Staphylococcus caseolyticus*, *Staphylococcus chromogenes*, *Staphylococcus condimentii*, *Staphylococcus delphini*, *Staphylococcus equorum*, *Staphylococcus felis*, *Staphylococcus fleurettii*, *Staphylococcus gallinarum*, *Staphylococcus hyicus*, *Staphylococcus*

intermedius, *Staphylococcus kloosii*, *Staphylococcus lentus*, *Staphylococcus lutrae*, *Staphylococcus muscae*, *Staphylococcus nepalensis*, *Staphylococcus piscifermentans*, *Staphylococcus pseudintermedius*, *Staphylococcus sciuri*, *Staphylococcus simiae*, *Staphylococcus succinus*, and *Staphylococcus vitulinus*.

[00111] In another embodiment, the bacterial agent is gram positive and is selected from the group comprising, but not limited to, *Streptococcus* spp. Examples of *Streptococcus* spp include *Streptococcus agalactiae*, *Streptococcus alactolyticus*, *Streptococcus anginosus*, *Streptococcus canis*, *Streptococcus constellatus*, *Streptococcus cricetus*, *Streptococcus cristatus*, *Streptococcus downei*, *Streptococcus dysgalactiae* subsp. *dysgalactiae*, *Streptococcus dysgalactiae* subsp. *equisimilis*, *Streptococcus equi* subsp. *equi*, *Streptococcus equi* subsp. *zooepidemicus*, *Streptococcus ferus*, *Streptococcus gallolyticus* subsp. *gallolyticus* (formerly *Streptococcus bovis* biotype i), *Streptococcus gallolyticus* subsp. *pasteurianus* (formerly *Streptococcus bovis* biotype ii/2), *Streptococcus gordonii*, *Streptococcus hyointestinalis*, *Streptococcus hyovaginalis*, *Streptococcus infantarius*, *Streptococcus infantarius* subsp. *infantarius*, *Streptococcus infantis*, *Streptococcus iniae*, *Streptococcus intermedius*, *Streptococcus lutetiensis* (formerly *Streptococcus bovis* biotype ii.1), *Streptococcus macaccae*, *Streptococcus mitis*, *Streptococcus mutans*, *Streptococcus oralis*, *Streptococcus orisratii*, *Streptococcus parasanguinis*, *Streptococcus peroris*, *Streptococcus pneumoniae*, *Streptococcus porcinus*, *Streptococcus pseudintermedius*, *Streptococcus pyogenes*, *Streptococcus rattii*, *Streptococcus salivarius*, *Streptococcus sanguinis*, *Streptococcus sobrinus*, *Streptococcus suis*, *Streptococcus thermophilus*, *Streptococcus vestibularis*, and Nutritionally Variant (Deficient) Streptococci (*Abiotrophia defectiva*, *Granulicatella adiacens*, *Granulicatella elegans*, and *Granulicatella para-adiacens*) and related species such as *Rothia mucilaginosa* (formerly *Stomatococcus mucilaginosus*) and *Pediococcus*.

[00112] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Enterococcus* spp. Examples of *Enterococcus* spp include *Enterococcus faecalis*, *Enterococcus faecium*, *Enterococcus gallinarum*, *Enterococcus durans*, *Enterococcus avium*, *Enterococcus raffinosus*, *Enterococcus pallens*, *Enterococcus gilvus*, *Enterococcus cecorum*, *Enterococcus malodoratus*, *Enterococcus italicus*, *Enterococcus sanguinicola*, *Enterococcus mundtii*, *Enterococcus casseliflavus/flavescens*, *Enterococcus dispar*, *Enterococcus hirae*, *Enterococcus pseudoavium*, and *Enterococcus bovis*.

[00113] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Leuconostoc* spp. Examples of *Leuconostoc* spp include *Leuconostoc mesenteroides*, *Leuconostoc pseudomesenteroides*, *Leuconostoc paramesenteroides*, *Leuconostoc citreum*, and *Leuconostoc lactis*.

[00114] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Corynebacterium* spp. Examples of *Corynebacterium* spp include nonlipophilic, fermentative *Corynebacterium* spp such as *Corynebacterium ulcerans*, *Corynebacterium pseudotuberculosis*, *Corynebacterium xerosis*, *Corynebacterium striatum*, *Corynebacterium minutissimum*, *Corynebacterium amycolatum*, *Corynebacterium glucuronolyticum*, *Corynebacterium argentoratense*, *Corynebacterium matruchotii*, *Corynebacterium riegellii*, *Corynebacterium confusum*, *Corynebacterium cystidis*, *Corynebacterium diphtheria*, *Corynebacterium simulans*, *Corynebacterium sundvallense*, *Corynebacterium thomssensii*, *Corynebacterium freneyi*, and *Corynebacterium aurimucosum*, nonlipophilic, nonfermentative *Corynebacterium* spp such as *Corynebacterium afermentans*, *Corynebacterium auris*, *Corynebacterium pseudodiphtheriticum*, and *Corynebacterium propinquum* and lipophilic *Corynebacterium* spp such as *Corynebacterium jeikeium*, *Corynebacterium urealyticum*, *Corynebacterium afermentans lipophilum*, *Corynebacterium accolens*, *Corynebacterium macginleyi*, *Corynebacterium tuberculostearum*, *Corynebacterium kroppenstedtii*, *Corynebacterium kutscheri*, *Corynebacterium pilosum*, *Corynebacterium bovis*, CDC coryneform groups F-1 and G, and *Corynebacterium lipophiloflavum*, and other *Corynebacterium* spp such as *Turicella*, *Arthrobacter*, *Brevibacterium*, *Dermabacter*, *Rothia*, *Oerskovia*, *Microbacterium*, and *Leifsonia aquatica*.

[00115] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Arcanobacteria* spp. Examples of *Arcanobacteria* spp include *A. haemolyticum*, *A. pyogenes* (now known as *Trueperella pyogenes*, originally known as *Actinomyces pyogenes*), and *A. bernardiae*.

[00116] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Rhodococcus* spp. Examples of *Rhodococcus* spp include *Rhodococcus equi*, *Rhodococcus erythropolis*, *Rhodococcus fasciens*, and *Rhodococcus rhodochrous*.

[00117] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Gordonia* spp.

[00118] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Tsukamurella* spp.

[00119] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Acholeplasma* spp.

[00120] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Actinobacteria* such as *Crossiella equi*.

[00121] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Bacillus* spp. Examples of *Bacillus* spp include *Bacillus anthracis*, *Bacillus cereus*, *Bacillus circulans*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus sphaericus*, *Bacillus subtilis*, *Brevibacillus brevis*, *Brevibacillus laterosporus*, and *Paenibacillus alvei*.

[00122] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, Anaerobic Cocci. Examples of Anaerobic Cocci include *Anaerococcus murdochii*, *Anaerococcus prevotii*, *Anaerococcus tetradius*, *Anaerococcus octavius*, *Anaerococcus hydrogenalis*, *Anaerococcus lactolyticus*, *Anaerococcus vaginalis*, *Atopobium parvulum*, *Fingoldia magna*, *Gallicola barnesae*, *Gemella asaccharolytica*, *Gemella bergeri*, *Gemella cuniculi*, *Gemella haemolysans*, *Gemella morbillorum*, *Gemella palaticanis*, *Gemella sanguinis*, *Parvimonas micra*, *Peptococcus niger*, *Peptoniphilus asaccharolyticus*, *Peptoniphilus gorbachii*, *Peptoniphilus indolicus*, *Peptoniphilus harei*, *Peptoniphilus ivorii*, *Peptoniphilus lacrimalis*, *Peptoniphilus olsenii*, *Peptostreptococcus stomatis*, *Peptostreptococcus anaerobius*, *Ruminococcus productus*, *Slackia heliotrinireducens*, and *Staphylococcus saccharolyticus*.

[00123] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, Anaerobic Gram-Positive Nonsporulating Bacilli. Examples of Anaerobic Gram-Positive Nonsporulating Bacilli include *Alloscardovia omnicolens*, *Atopobium* species (such as *Atopobium minutum*, *Atopobium rimae*, *Atopobium parvulum*, and *Atopobium vaginae*), *Bifidobacteria* (such as *Bifidobacteria adolescentis*, *Bifidobacteria dentium*, *Bifidobacteria scardovii*), *Catabacter hongkongensis*, *Collinsella aerofaciens*, *Eggerthella* (such as *Eggerthella lenta*, *Eggerthella hongkongensis* and *Eggerthella sinensis*), *Eubacterium* and related species (such as *Eubacterium nodatum*, *Eubacterium tenue*, *Eubacterium brachy*, *Eubacterium infirmum*, *Eubacterium minutum*, *Eubacterium nodatum*, *Eubacterium saphenum*, *Eubacterium sulci*, *Filifactor alocis*, *Mogibacterium timidum*, *Mogibacterium vescum*, *Pseudoramibacter alactolyticus*, *Bulleidia extracta*, and *Solobacterium moorei*), *Lactobacillus* species (such as *Lactobacillus rhamnosus*, *Lactobacillus casei*, *Lactobacillus fermentum*, *Lactobacillus gasseri*, *Lactobacillus plantarum*, *Lactobacillus acidophilus*, *Lactobacillus iners* and *Lactobacillus ultunensis*), *Mobiluncus* species (such as *Mobiluncus curtisii*, *Mobiluncus mulieris*), *Moryella indoligenes*, *Olsenella* oral species (such as *Olsenella uli* and *Olsenella profuse*), *Oribacterium sinus*, *Propionibacterium* (such as *Propionibacterium acnes* and *Propionibacterium propionicum*), *Slackia exigua*, and *Turicibacter sanguine*.

[00124] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Actinomyces* spp. Examples of *Actinomyces* spp include *Actinomyces israelii*, *Actinomyces naeslundii*, *Actinomyces viscosus*, *Actinomyces*

odontolyticus, *Actinomyces meyeri*, and *Actinomyces gerencseriae* (formerly *Actinomyces israelii* serotype II), *Actinomyces europaeus*, *Actinomyces neuii*, *Actinomyces radingae*, *Actinomyces graevenitzii*, *Actinomyces hordeovulneris*, *Actinomyces turicensis*, *Actinomyces georgiae*, *Arcanobacterium (Actinomyces) pyogenes*, *Arcanobacterium (Actinomyces) bernardiae*, *Actinomyces funkei*, *Actinomyces lingnae*, *Actinomyces houstonensis*, and *Actinomyces cardiffensis*.

[00125] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Clostridium* spp. Examples of *Clostridium* spp include *Clostridium baratii*, *Clostridium bifermentans*, *Clostridium botulinum*, *Clostridium botulinum* (types A, B, C, D, E, F, G), *Clostridium butyricum*, *Clostridium difficile*, *Clostridium histolyticum*, *Clostridium novyi* (type A), *Clostridium novyi* (type B), *Clostridium perfringens*, *Clostridium perfringens* (types A-E), *Clostridium ramosum*, *Clostridium septicum*, *Clostridium sordelli*, *Clostridium sphenoides*, *Clostridium tertium*, and *Clostridium tetani*.

[00126] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Nocardia* spp. Examples of *Nocardia* spp include *Nocardia asteroides*, *Nocardia brasiliensis*, *Nocardia farcinica*, *Nocardia nova*, *Nocardia otitidiscaviarum*, and *Nocardia transvalensis*.

[00127] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Erysipelothrix* spp, such as *Erysipelothrix rhusiopathiae*.

[00128] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Listeria* spp, such as *Listeria monocytogenes*.

[00129] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Kytococcus* spp, such as *Kytococcus schroeteri*.

[00130] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Mycobacterium* spp. Examples of *Mycobacterium* spp include *Mycobacterium abscessus*, *Mycobacterium arupense*, *Mycobacterium asiaticum*, *Mycobacterium aubagnense*, *Mycobacterium avium* complex, *Mycobacterium bolletii*, *Mycobacterium bolletii*, *Mycobacterium branderi*, *Mycobacterium canettii*, *Mycobacterium caprae*, *Mycobacterium celatum*, *Mycobacterium chelonae*, *Mycobacterium chimaera*, *Mycobacterium colombiense*, *Mycobacterium conceptionense*, *Mycobacterium conspicuum*, *Mycobacterium elephantis*, *Mycobacterium farcinogenes*, *Mycobacterium florentinum*, *Mycobacterium fortuitum* group, *Mycobacterium genavense*, *Mycobacterium goodii*, *Mycobacterium haemophilum*, *Mycobacterium heckeshomense*, *Mycobacterium heidelbergense*, *Mycobacterium houstonense*, *Mycobacterium immunogenum*, *Mycobacterium*

interjectum, *Mycobacterium intracellulare*, *Mycobacterium senegalense*, *Mycobacterium africanum*, *Mycobacterium avium* subsp *paratuberculosis*, *Mycobacterium kansasii*, *Mycobacterium lacus*, *Mycobacterium lentiflavum*, *Mycobacterium leprae*, *Mycobacterium lepraemurium*, *Mycobacterium mageritense*, *Mycobacterium malmoeense*, *Mycobacterium marinum*, *Mycobacterium massiliense*, *Mycobacterium microti*, *Mycobacterium montefiorensis* (eels), *Mycobacterium moracense*, *Mycobacterium mucogenicum*, *Mycobacterium nebraskense*, *Mycobacterium neoaurum*, *Mycobacterium novocastrense*, *Mycobacterium palustre*, *Mycobacterium parmensis*, *Mycobacterium phlei*, *Mycobacterium phocaicum*, *Mycobacterium pinnipedii*, *Mycobacterium porcinum*, *Mycobacterium pseudoshottsii* (fish), *Mycobacterium pseudotuberculosis*, *Mycobacterium saskatchewanense*, *Mycobacterium scrofulaceum*, *Mycobacterium sensuense*, *Mycobacterium septicum*, *Mycobacterium simiae*, *Mycobacterium smegmatis*, *Mycobacterium szulgai*, *Mycobacterium terrae/chromogenicum* complex, *Mycobacterium triplex*, *Mycobacterium tuberculosis*, *Mycobacterium tusciae*, *Mycobacterium ulcerans*, *Mycobacterium wolinskyi*, and *Mycobacterium xenopi*.

[00131] In another embodiment, the bacterial agent is gram positive and selected from the group comprising, but not limited to, *Trueperella* spp. Examples of *Trueperella* spp include *Trueperella abortusovis*, *Trueperella bernardiae*, *Trueperella bialowiezensis*, *Trueperella bonasi*, *Trueperella pyogenes* (*Arcanobacterium pyogenes*).

[00132] In another embodiment, the bacterial agent is gram positive, gram negative or does not have a cell wall and selected from the group comprising, but not limited to, livestock pathogens. Examples of livestock pathogens include *Actinobaculum suis*, *Actinomyces bovis*, *Arcanobacterium pyogenes*, *Bacillus anthracis*, *cereus*, *licheniformis*, *pumilus*, *melaninogenicus*, *subtilis*, *Clostridium botulinum*, *chauvoei*, *haemolyticum*, *novyi*, *perfringens*, *septicum*, *sordellii*, *tetani*, *colinum*, *Corynebacterium pseudotuberculosis*, *renale*, *Dermatophilus congolensis*, *Enterococcus* spp (such as *E. faecalis*, *E. faecium*, *E. durans*, *E. avium*, *E. hirae*), *Erysipelothrix rhusiopathiae*, *Listeria ivanovii*, *grayi*, *innocua*, *seeligeri*, *welshimeri*, *monocytogenes*, *Mycobacterium avium*, *bovis*, *paratuberculosis* (John's Disease), *Mycoplasma* (such as *capricolum* subsp. *capripneumoniae*, subsp. *capricolum*, *M. mycoides* subsp *mycoides*, *M. agalactiae*, *M. ovipneumoniae*, *M. conjunctivae*, *M. arginini*, *M. bovis*, and *M. putrefaciens*) *Mycoplasma bovis*, *dispar*, *mycoides* subsp. *mycoides* (such as Contagious bovine pleuropneumonia CBPP) *Mycoplasma gallisepticum* (MG), *iowae meleagridis* (MM), *synoviae* (MS) *Mycoplasma haemosuis* (formerly *Eperythrozoon suis*), *alkalescens*, *bovigenitalum*, *bovirhinis*, *bovoculi*, *californicum*, *canadense*, *cynos*, *equigenitalum*, *gateae*, *haemocanis*, *haemofelis*, *hyopneumoniae*, *hyorhinis*, *hyosynoviae*, *iowae*, *leachii*, *meleagridis*, *mycoides* subsp *capri*, *weryonii*, *suis*, *Rhodococcus equi*, *Staphylococcus epidermidis*, *Staphylococcus simulans*, *Staphylococcus felis*, *Staphylococcus xylosus*, *Staphylococcus chromogenes*,

Staphylococcus warneri, *Staphylococcus haemolyticus*, *Staphylococcus sciuri*, *Staphylococcus saprophyticus*, *Staphylococcus hominis*, *Staphylococcus caprae*, *Staphylococcus cohnii* subsp. *cohnii*, *Staphylococcus cohnii* subsp. *urealyticus*, *Staphylococcus capitis* subsp. *capitis*, *Staphylococcus capitis* subsp. *urealyticus*, *Staphylococcus hyicus*, *Staphylococcus aureus*, *Staphylococcus pseudintermedius*, *Staphylococcus delphini*, *Staphylococcus schleiferi* subsp. *coagulans*, *Staphylococcus aureus* subsp. *anaerobius*, *Streptococcus uberis*, *Streptococcus canis*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus pyogenes*, *Streptococcus bovis*, *Streptococcus equi* subsp. *Zooepidemicus*, *Streptococcus equinus*, *Streptococcus equi* (*Streptococcus equi* subsp. *equi*), *Streptococcus equisimilis* (*Streptococcus dysgalactiae* subsp. *equisimilis*), *porcinus*, *suis*, *zooepidemicus*, *Streptococcus zooepidemicus* (*Streptococcus equi* subsp. *zooepidemicus*), *Streptococcus dysgalactiae* subsp. *equisimilis*, *Propionibacterium acnes*, *Propionibacterium granulosum*, *Eubacterium*, *Peptococcus indolicus*, and *Peptostreptococcus anaerobius*; and various species of the following Gram negative genera: *Actinobacillus*, *Aeromonas*, *Anaplasma*, *Arcobacter*, *Avibacterium*, *Bacteroides*, *Bartonella*, *Bordetella*, *Borrelia*, *Brachyspira*, *Brucella*, *Campylobacter*, *Capnocytophaga*, *Chlamydia*, *Chlamydophila*, *Chryseobacterium*, *Coxiella*, *Cytophaga*, *Dichelobacter*, *Edwardsiella*, *Ehrlichia*, *Escherichia*, *Flavobacterium*, *Francisella*, *Fusobacterium*, *Gallibacterium*, *Haemophilus*, *Histophilus*, *Klebsiella*, *Lawsonia*, *Leptospira*, *Mannheimia*, *Megasphaera*, *Moraxella*, *Neorickettsia*, *Nicoletella*, *Ornithobacterium*, *Pasteurella*, *Photobacterium*, *Piscichlamydia*, *Piscirickettsia*, *Porphyromonas*, *Prevotella*, *Proteus*, *Pseudomonas*, *Rickettsia*, *Riemerella*, *Salmonella*, *Streptobacillus*, *Tenacibaculum*, *Vibrio*, and *Yersinia*.

[00133] In another embodiment, the bacterial agent is selected from the group comprising, but not limited to, pathogens of companion animal species such as cats, dogs and horses. Examples of such pathogens include equine pathogens such as *Streptococcus equi*, *Streptococcus zooepidemicus*, *Rhodococcus equi*, *Clostridium difficile*, *Clostridium perfringens*, *Corynebacterium pseudotuberculosis*, *Clostridium piliforme*, *Actinomyces bovis*, *Staphylococcus aureus*, β -haemolytic *Streptococcus* spp, *Dermatophilus congolense*, *Clostridium tetani*, and *Clostridium botulinum*. Further examples include pathogens of dogs and cats such as *Staphylococcus* spp, *Streptococcus* spp, *Clostridium* spp, *Actinomyces* spp, *Enterococcus* spp, *Nocardia* spp, *Mycoplasma* spp, and *Mycobacterium* spp.

[00134] In another embodiment, the bacterial agent is gram negative and selected from the group consisting of the following representative families and species: *Acetobacteraceae*:- *Roseomonas cervicalis*; *Roseomonas fauriae*; *Roseomonas gilardii*. -- *Aeromonadaceae*:- *Aeromonas allosaccharophila*; *Aeromonas aquariorum*; *Aeromonas caviae*; *Aeromonas hydrophila* (and subspecies); *Aeromonas salmonicida*; *Aeromonas shubertii*; *Aeromonas veronii*

biovar *sobria* (*Aeromonas sobria*). -- *Alcaligenaceae*:- *Achromobacter xylosoxidans*; *Alcaligenes faecalis*; *Bordetella ansorpii*; *Bordetella avium*; *Bordetella bronchiseptica*; *Bordetella hinzii*; *Bordetella holmesii*; *Bordetella parapertussis*; *Bordetella pertussis*; *Bordetella petrii*; *Bordetella trematum*; *Oligella ureolytica*; *Oligella urethralis*. -- *Anaplasmataceae*:- *Anaplasma phagocytophilum*; *Anaplasma platys*; *Anaplasma bovis*; *Anaplasma centrale*; *Anaplasma marginale*; *Anaplasma odocoilei*; *Anaplasma ovis*; *Ehrlichia canis*; *Ehrlichia chaffeensis*; *Ehrlichia ewingii*; *Ehrlichia muris*; *Ehrlichia ovina*; *Ehrlichia ruminantium*; *Neoehrlichia lotoris*; *Neoehrlichia mikurensis*; *Neorickettsia helminthoeca*; *Neorickettsia risticii*; *Neorickettsia sennetsu*; *Wolbachia pipientis*. -- *Armatimonadaceae*:- *Armatimonas rosea*. -- *Bacteroidaceae*:- *Bacteroides forsythus*; *Bacteroides fragilis*; *Bacteroides melaninogenicus*; *Bacteroides ruber*; *Bacteroides urealyticus*. -- *Bartonellaceae*:- *Bartonella alsatica*; *Bartonella australis*; *Bartonella bacilliformis*; *Bartonella birtlesii*; *Bartonella bovis*; *Bartonella capreoli*; *Bartonella chomelii*; *Bartonella clarridgeiae*; *Bartonella doshiae*; *Bartonella elizabethae*; *Bartonella grahamii*; *Bartonella henselae*; *Bartonella koehlerae*; *Bartonella peromysci*; *Bartonella phoceensis*; *Bartonella quintana*; *Bartonella rattimassiliensis*; *Bartonella rochalimae*; *Bartonella schoenbuchensis*; *Bartonella talpae*; *Bartonella tamiae*; *Bartonella taylorii*; *Bartonella tribocorum*; *Bartonella vinsonii* subsp. *berkhoffii*; *Bartonella vinsonii* subsp. *arupensis*; *Bartonella vinsonii* subsp. *vinsonii*. -- *Bdellovibrionaceae*:- *Bdellovibrio* spp. -- *Brachyspiraceae*:- *Brachyspira* spp including *Brachyspira hampsonii*, *Brachyspira hyodysenteriae*, *Brachyspira murdochii*, *Brachyspira pilosicoli*. -- *Brucellaceae*:- *Brucella abortus*; *Brucella canis*; *Brucella ceti*; *Brucella melitensis*; *Brucella ovis*; *Brucella pinnipedialis*; *Brucella suis*; *Ochrobactrum anthropi*; *Ochrobactrum intermedium*. -- *Burkholderiaceae*:- *Burkholderia aboris*; *Burkholderia ambifaria* (genomovar VII); *Burkholderia anthina* (genomovar VIII); *Burkholderia cenocepacia* (genomovar III); *Burkholderia cepacia* (genomovar I); *Burkholderia diffusa*; *Burkholderia dolosa* (genomovar VI); *Burkholderia latens*; *Burkholderia mallei*; *Burkholderia metallica*; *Burkholderia multivorans* (genomovar II); *Burkholderia pseudomallei*; *Burkholderia pyrrocinia* (genomovar IX); *Burkholderia seminalis*; *Burkholderia stabilis* (genomovar IV); *Burkholderia ubonensis* (genomovar X); *Burkholderia vietnamiensis* (genomovar V); *Cupriavidus pauculus*; *Cupriavidus gilardii*; *Ralstonia pickettii*; *Ralstonia mannitolilytica*; *Sphaerotilus hippel*; *Sphaerotilus moritanus*; *Sphaerotilus natans*. -- *Campylobacteraceae*:- *Arcobacter* spp including *Arcobacter skirrowii*; *Campylobacter coli*; *Campylobacter concisus*; *Campylobacter curvus*; *Campylobacter fetus*; *Campylobacter gracilis*; *Campylobacter helveticus*; *Campylobacter hominis*; *Campylobacter hyointestinalis*; *Campylobacter insulaenigrae*; *Campylobacter jejuni*; *Campylobacter lanienae*; *Campylobacter lari*; *Campylobacter laridis*; *Campylobacter mucosalis*; *Campylobacter rectus*; *Campylobacter showae*; *Campylobacter sputorum*; *Campylobacter upsaliensis*. -- *Candidatus*:- *Piscichlamydia salmonis*. -- *Cardiobacteriaceae*:- *Cardiobacterium hominis*; *Cardiobacterium valvarum*; *Dichelobacter nodosus*. -- *Chlamydiaceae*:- *Chlamydia* spp including *Chlamydia avium*, *Chlamydia gallinacea*, *Chlamydia muridarum*, *Chlamydia suis*,

Chlamydia trachomatis; *Chlamydophila* spp including *Chlamydophila pneumoniae*, *Chlamydophila pecorum*, *Chlamydophila psittaci*, *Chlamydophila abortus*, *Chlamydophila caviae*, and *Chlamydophila felis*. -- *Chthonomonadaceae*:- *Chthonomonas calidirosea*. -- *Comamonadaceae*:- *Comamonas testosteroni*; *Verminephrobacter* spp. -- *Coxiellaceae*:- *Coxiella burnetii*. -- *Cytophagaceae*:- *Cytophaga columnaris*; *Cytophaga hutchinsonii*; *Flexibacter echinica*; *Flexibacter elegans*; *Flexibacter flexilis*; *Flexibacter litoralis*; *Flexibacter polymorphus*; *Flexibacter roseolus*; *Flexibacter ruber*. -- *Desulfovibrionaceae*:- *Bilophila wadsworthia*; *Lawsonia intracellularis*. -- *Enterobacteriaceae*:- *Cedecea davisae*; *Cedecea lapagei*; *Cedecea neteri*; *amalonaticus*; *Citrobacter diversus*; *Citrobacter freundii*; *Citrobacter koseri*; *Cronobacter condimentii*; *Cronobacter dublinensis*; *Cronobacter helveticus*; *Cronobacter malonaticus*; *Cronobacter muytjensii*; *Cronobacter pulveris*; *Cronobacter sakazakii*; *Cronobacter turicensis*; *Cronobacter universalis*; *Cronobacter zurichensis*; *Edwardsiella ictaluri*; *Edwardsiella tarda*; *Enterobacter aerogenes*; *Enterobacter agglomerans*; *Enterobacter cloacae*; *Enterobacter cowanii*; *Escherichia albertii*; *Escherichia coli*, including AIEC = adherent invasive *E. coli*, EaggEC = enteroaggregative *E. coli*; EHEC = enterohemorrhagic *E. coli*; EIEC = enteroinvasive *E. coli*; EPEC = enteropathogenic *E. coli*; ETEC = enterotoxigenic *E. coli*; ExPEC = extraintestinal pathogenic *E. coli*, NMEC = neonatal meningitis *E. coli*, NTEC = necrotogenic *E. coli*, UPEC = uropathogenic *E. coli*; *Escherichia fergusonii*; *Ewingella americana*; *Hafnia alvei*; *Hafnia paralvei*; *Klebsiella granulomatis*; *Klebsiella oxytoca*; *Klebsiella pneumoniae*; *Kluyvera ascorbata*; *Kluyvera cryocrescens*; *Morganella morganii*; *Pantoea* (formerly *Enterobacter*) *agglomerans*; *Photorhabdus asymbiotica*; *Plesiomonas shigelloides*; *Proteus mirabilis*; *Proteus penneri*; *Proteus vulgaris*; *Providencia alcalifaciens*; *Providencia rettgeri*; *Providencia stuartii*; *Raoultella electrica*; *Raoultella ornithinolytica*; *Raoultella planticola*; *Raoultella terrigena*; *Salmonella bongori*, *Salmonella enterica* subspecies *enterica* (many serotypes); *Serratia liquefaciens*; *Serratia marcescens*; *Shigella boydii*; *Shigella dysenteriae*; *Shigella flexneri*; *Shigella sonnei*; *Yersinia enterocolitica*; *Yersinia pestis*; *Yersinia pseudotuberculosis*; *Yersinia ruckeri*. -- *Fimbrimonadaceae*:- *Fimbrimonas ginsengisoli*. -- *Flavobacteriaceae*:- *Bergeyella zoohelcum*; *Capnocytophaga canimorsus*; *Capnocytophaga cynodegmi*; *Capnocytophaga gingivalis*; *Capnocytophaga granulosa*; *Capnocytophaga haemolytica*; *Capnocytophaga leadbetteri*; *Capnocytophaga ochracea*; *Capnocytophaga sputigena*; *Chryseobacterium indologenes*; *Chryseobacterium piscicola*; *Elizabethkingia meningoseptica*; *Flavobacterium branchiophilum*; *Flavobacterium columnare*; *Flavobacterium oncorhynchi*; *Flavobacterium piscicida*; *Flavobacterium psychrophilum*; *Myroides odoratus*; *Myroides odoratimimus*; *Ornithobacterium rhinotracheale*; *Riemerella anatipestifer*; *Riemerella columbina*; *Riemerella columbipharyngis*; *Tenacibaculum dicentrarchi*; *Tenacibaculum discolour*; *Tenacibaculum gallaicum*; *Tenacibaculum maritimum*; *Tenacibaculum soleae*; *Weeksella virosa*. -- *Francisellaceae*:- *Francisella tularensis* subsp. *tularensis*; *Francisella tularensis* subsp. *holarctica*; *Francisella tularensis* subsp. *novicida*; *Francisella philomiragia*;

Francisella noatunensis; *Francisella noatunensis* subsp. *orientalis* (also termed *Francisella asiatica*). -- *Fusobacteriaceae*:- *Fusobacterium* spp. including *Fusobacterium necrophorum*, *Fusobacterium nucleatum*, *Fusobacterium polymorphum*. -- *Helicobacteraceae*:- *Helicobacter cinaedi*; *Helicobacter fennelliae*; *Helicobacter pylori*. -- *Legionellaceae*:- *Legionella pneumophila* and other species including; *Legionella anisa*; *Legionella birminghamensis*; *Legionella bozemannii*; *Legionella cincinnatiensis*; *Legionella dumoffii*; *Legionella feeleyi*; *Legionella gormanii*; *Legionella hackeliae*; *Legionella jordanis*; *Legionella lansingensis*; *Legionella longbeachae*; *Legionella maceachernii*; *Legionella micdadei*; *Legionella oakridgensis*; *Legionella parisiensis*; *Legionella sainthelens*; *Legionella tusconensis*; *Legionella wadsworthii*; *Legionella waltersii*. -- *Leptospiraceae*:- *Leptospira alexanderi* (including *Leptospira alexanderi* serovar *Hebdomadis*, *Leptospira alexanderi* serovar *Manhao 3*); *Leptospira alstoni* (including *Leptospira alstoni* serovar *Pingchang*, *Leptospira alstoni* serovar *Sichuan*); *Leptospira biflexa* (including *Leptospira biflexa* serovar *Ancona*, *Leptospira biflexa* serovar *Canela*); *Leptospira borgpetersenii* (including *Leptospira borgpetersenii* serovar *Hardjo*, *Leptospira borgpetersenii* serovar *Hardjo-bovis*, *Leptospira borgpetersenii* serovar *Pomona*, *Leptospira borgpetersenii* serovar *Tarassovi*); *Leptospira broomii* (including *Leptospira broomii* serovar *Hurstbridge*); *Leptospira fainei* (including *Leptospira fainei* serovar *Hurstbridge*); *Leptospira idonii*; *Leptospira inadai* (including *Leptospira inadai* serovar *Lyme*, *Leptospira inadai* serovar *Malaya*); *Leptospira interrogans* (including *Leptospira interrogans* serovar *Australis*, *Leptospira interrogans* serovar *Autumnalis*, *Leptospira interrogans* serovar *Bratislava*, *Leptospira interrogans* serovar *Canicola*, *Leptospira interrogans* serovar *Grippotyphosa*, *Leptospira interrogans* serovar *Hardjo*, *Leptospira interrogans* serovar *Hardjo-bovis*, *Leptospira interrogans* serovar *Icterohaemorrhagiae*, *Leptospira interrogans* serovar *Pomona*, *Leptospira interrogans* serovar *Pyrogenes*, *Leptospira interrogans* serovar *Tarassovi*); *Leptospira kirschneri* (including *Leptospira kirschneri* serovar *Bulgarica*, *Leptospira kirschneri* serovar *Cynopteri*, *Leptospira kirschneri* serovar *Grippotyphosa*); *Leptospira kmetyi*; *Leptospira licerasiae*; *Leptospira meyeri* (including *Leptospira meyeri* serovar *Sofia*); *Leptospira noguchii* (including *Leptospira noguchii* serovar *Panama*, *Leptospira noguchii* serovar *Pomona*); *Leptospira santarosai*; *Leptospira terpstrae*; *Leptospira vanthielii*; *Leptospira weilii* (including *Leptospira weilii* serovar *Celledoni*, *Leptospira weilii* serovar *Sarmin*); *Leptospira wolbachii*; *Leptospira wolffii*; *Leptospira yanagawae*. -- *Leptotrichiaceae*:- *Leptotrichia buccalis*; *Streptobacillus moniliformis*. -- *Methylobacteriaceae*:- *Methylobacterium extorquens* group; *Methylobacterium fujisawaense*; *Methylobacterium mesophilicum*; *Methylobacterium zatmanii*. -- *Moraxellaceae*:- *Acinetobacter baumannii* (genomic species 2); *Acinetobacter baylyi*; *Acinetobacter bouvetii*; *Acinetobacter calcoaceticus* (genomic species 1); *Acinetobacter gerneri*; *Acinetobacter grimontii*; *Acinetobacter haemolyticus* (genomic species 4); *Acinetobacter johnsonii* (genomic species 7); *Acinetobacter junii* (genomic species 5); *Acinetobacter lwoffii* (genomic species 8/9); *Acinetobacter parvus*; *Acinetobacter radioresistens* (genomic species 12); *Acinetobacter*

schindleri; *Acinetobacter tandoii*; *Acinetobacter tjernbergiae*; *Acinetobacter towneri*; *Acinetobacter ursingii*; *Acinetobacter venetianus*; *Moraxella atlantae*; *Moraxella boevrei*; *Moraxella bovis*; *Moraxella bovoculi*; *Moraxella canis*; *Moraxella caprae*; *Moraxella catarrhalis*; *Moraxella caviae*; *Moraxella cuniculi*; *Moraxella equi*; *Moraxella lacunata*; *Moraxella lincolnii*; *Moraxella macacae*; *Moraxella nonliquefaciens*; *Moraxella oblonga*; *Moraxella osloensis*; *Moraxella ovis*; *Moraxella phenylpyruvica*; *Moraxella pluranimalium*; *Moraxella porci*. --
Moritellaceae:- *Moritella abyssi*; *Moritella dasanensis*; *Moritella japonica*; *Moritella marina*; *Moritella pro-funda*; *Moritella viscosa*; *Moritella yayanosii*. -- *Neisseriaceae*:- *Chromobacterium violaceum*; *Eikenella corrodens*; *Kingella denitrificans*; *Kingella kingae*; *Kingella oralis*; *Kingella potus*; *Neisseria cinerea*; *Neisseria elongata*; *Neisseria flavescens*; *Neisseria gonorrhoeae*; *Neisseria lactamica*; *Neisseria meningitidis*; *Neisseria mucosa*; *Neisseria polysaccharea*; *Neisseria sicca*; *Neisseria subflava*; *Neisseria weaver*; *Vitreoscilla* spp. -- *Nitrosomonadaceae*:- *Nitrosomonas eutropha*; *Nitrosomonas halophila*; *Nitrosomonas oligotropha*. --
Pasteurellaceae:- *Actinobacillus actinomycetemcomitans*; *Actinobacillus equuli*; *Actinobacillus lignieresii*; *Actinobacillus pleuropneumoniae*; *Actinobacillus seminis*; *Actinobacillus succinogenes*; *Actinobacillus ureae*; *Aggregatibacter actinomycetemcomitans*; *Aggregatibacter segnis*; *Aggregatibacter aphrophilus*; *Avibacterium avium*; *Avibacterium endocarditidis*; *Avibacterium gallinarum*; *Avibacterium paragallinarum*; *Avibacterium volantium*; *Bibersteinia trehalose*; *Gallibacterium anatis*; *Gallibacterium genomospecies 1*; *Gallibacterium genomospecies 2*; *Gallibacterium genomospecies 3*; *Gallibacterium group V*; *Gallibacterium melopsittaci*; *Gallibacterium salpingitidis*; *Gallibacterium trehalosifermentans*; *Haemophilus aegyptius*; *Haemophilus avium*; *Haemophilus ducreyi*; *Haemophilus haemolyticus*; *Haemophilus influenzae*; *Haemophilus parahaemolyticus*; *Haemophilus parainfluenzae*; *Haemophilus parasuis*; *Histophilus somni*; *Mannheimia caviae*; *Mannheimia glucosida*; *Mannheimia granulomatis*; *Mannheimia haemolytica*; *Mannheimia ruminalis*; *Mannheimia varigena*; *Nicotetella semolina*; *Pasteurella aerogenes*; *Pasteurella bettyae*; *Pasteurella caballi*; *Pasteurella canis*; *Pasteurella dagmatis*; *Pasteurella multocida* (subspecies *multocida*, *septicum*, *gallicida*); *Pasteurella pneumotropica*; *Pasteurella stomatis*; *Pasteurella trehalosi*. --
Piscirickettsiaceae:- *Piscirickettsia salmonis*. -- *Plesiomonadaceae*:- *Plesiomonas shigelloides*. --
Polyangiaceae:- *Sorangium cellulosum*. -- *Porphyromonadaceae*:- *Dysgonomonas capnocytophagoides*; *Dysgonomonas gadei*; *Dysgonomonas hofstadii*; *Dysgonomonas mossii*; *Dysgonomonas oryzarvi*; *Dysgonomonas wimpennyi*; *Porphyromonas gingivalis*. --
Prevotellaceae:- *Prevotella* spp. including *Prevotella intermedia*, *Prevotella melaninogenica*. --
Pseudomonadaceae:- *Chryseomonas luteola*; *Pseudomonas aeruginosa*; *Pseudomonas luteola*; *Pseudomonas fluorescens*; *Pseudomonas putida*; *Pseudomonas stutzeri*; *Pseudomonas oryzae*. -- *Rhizobiaceae*:- *Agrobacterium tumefaciens*; *Rhizobium radiobacter*. -- *Rickettsiaceae*:- *Orientia chuto*; *Orientia tsutsugamushi*; *Rickettsia aeschlimannii*; *Rickettsia africae*; *Rickettsia akari*; *Rickettsia argasii*; *Rickettsia asiatica*; *Rickettsia australis*;

Rickettsia bellii; *Rickettsia canadensis*; *Rickettsia conorii*; *Rickettsia cooley*; *Rickettsia felis*; *Rickettsia heilongjiangensis*; *Rickettsia helvetica*; *Rickettsia honei*; *Rickettsia hoogstraalii*; *Rickettsia hulinensis*; *Rickettsia hulinii*; *Rickettsia japonica*; *Rickettsia marmionii*; *Rickettsia martinii*; *Rickettsia massiliae*; *Rickettsia monacensis*; *Rickettsia montanensis*; *Rickettsia monteiroi*; *Rickettsia moreli*; *Rickettsia parkeri*; *Rickettsia peacockii*; *Rickettsia philipii*; *Rickettsia prowazekii*; *Rickettsia raoultii*; *Rickettsia rhipicephali*; *Rickettsia rickettsii*; *Rickettsia sibirica* subgroup; *Rickettsia slovaca*; *Rickettsia tamurae*; *Rickettsia typhi*. -- Shewanellaceae:- *Shewanella putrefaciens*. -- Sphingomonadaceae:- *Sphingobacterium multivorum*; *Sphingobacterium spiritivorum*; *Sphingomonas paucimobilis*. -- Spirillaceae:- *Spirillum minus*; *Spirillum volutans*; *Spirillum winogradskyi*. -- Spirochaetaceae:- *Borrelia afzelii*; *Borrelia anserina*; *Borrelia bissettii*; *Borrelia burgdorferi*; *Borrelia coriaceae*; *Borrelia duttonii*; *Borrelia garinii*; *Borrelia hermsii*; *Borrelia hispanica*; *Borrelia japonica*; *Borrelia lonestari*; *Borrelia lusitanae*; *Borrelia miyamotoi*; *Borrelia parkeri*; *Borrelia persica*; *Borrelia recurrentis*; *Borrelia spielmanii*; *Borrelia turicatae*; *Borrelia turicatae*; *Borrelia valaisiana*; *Treponema carateum*; *Treponema pallidum* ssp. *endemicum*; *Treponema pallidum* ssp. *pallidum*; *Treponema pallidum* ssp. *pertenue*. -- Succinivibrionaceae:- *Anaerobiospirillum* spp. -- Sutterellaceae:- *Sutterella* spp including *Sutterella wadsworthia*. -- Thermaceae:- *Meiothermus* spp. -- Thermotogaceae:- *Thermotoga neapolitana*. -- Veillonellaceae:- *Dialister* spp; *Megamonas* spp; *Megasphaera* spp; *Pectinatus* spp; *Pelosinus* spp; *Propionispora* spp; *Sporomusa* spp; *Veillonella* spp; *Zymophilus* spp. -- Vibrionaceae:- *Photobacterium damsela*; *Vibrio adaptatus*; *Vibrio alginolyticus*; *Vibrio azasii*; *Vibrio campbellii*; *Vibrio cholera*; *Vibrio damsela*; *Vibrio fluvialis*; *Vibrio furnisii*; *Vibrio hollisae*; *Vibrio metchnikovii*; *Vibrio mimicus*; *Vibrio parahaemolyticus*; *Vibrio vulnificus*. -- Wolbachiae:- *Wolbachia* spp. -- Xanthomonadaceae:- *Luteimonas aestuarii*; *Luteimonas aquatica*; *Luteimonas composti*; *Luteimonas lutimaris*; *Luteimonas marina*; *Luteimonas mephitis*; *Luteimonas vadosa*; *Pseudoxanthomonas broegbernensis*; *Pseudoxanthomonas japonensis*; *Stenotrophomonas maltophilia*; *Stenotrophomonas nitritireducens*.

[00135] Most preferably, the bacterial agent causing the bacterial infection is gram negative and is selected from the group comprising: *Acinetobacter* species, *Aeromonas hydrophila*, *Citrobacter* species, *Enterobacter* species, *Escherichia coli*, *Klebsiella pneumoniae*, *Morganella morganii*, *Pseudomonas aeruginosa*, and *Stenotrophomonas maltophilia*.

[00136] In another preferred embodiment, the bacteria agent causing the bacterial colonisation or infection is resistant to a conventional antibiotic used to treat the colonisation or infection. In one preferred embodiment, the bacterial agent is resistant to a compound selected from the group comprising: one or more of aminoglycosides (for example gentamicin, tobramycin, amikacin, or netilmicin); anti-MRSA cephalosporins (for example ceftaroline); antipseudomonal penicillins + β -lactamase inhibitors (for example ticarcillin-clavulanic acid or

piperacillin-tazobactam); carbapenems (for example ertapenem, imipenem, meropenem or doripenem); non-extended spectrum cephalosporins; 1st and 2nd generation cephalosporins (for example cefazolin or cefuroxime); extended-spectrum cephalosporins; 3rd and 4th generation cephalosporins (for example cefotaxime or ceftriaxone); cephamycins (for example cefoxitin or cefotetan); fluoroquinolones (for example ciprofloxacin); folate pathway inhibitors (for example trimethoprim-sulphamethoxazole); glycylicyclines (for example tigecycline); monobactams (for example aztreonam); penicillins (for example ampicillin); penicillins + β -lactamase inhibitors (for example amoxicillin-clavulanic acid or ampicillin-sulbactam); phenicols (for example chloramphenicol); phosphonic acids (for example fosfomycin); polymyxins (for example colistin); and tetracyclines (for example tetracycline, doxycycline or minocycline). Preferably, the bacterial agent resistant to these compounds is gram negative.

[00137] Preferably, the bacterial agent is resistant to a compound selected from the group comprising: penicillins, cephalosporins, carbapenems, monobactams and other β -lactam antibiotics, fusidanes, aminoglycosides, fluoroquinolones, streptogramins, tetracyclines, glycylicyclines, chloramphenicol and other phenicols, macrolides and ketolides, lincosamides, oxazolidinones, aminocyclitols, polymyxins, glycopeptides, lipopeptides, bacitracin, mupirocin, pleuromutilins, rifamycins, sulphonamides and trimethoprim. Preferably, the compound is selected from the group comprising: beta lactams, glycopeptides, lipopeptides, macrolides, oxazolidinones and tetracyclines. Preferably, the bacterial agent is resistant to the compound when the compound is at a concentration range selected from the following: 0.001 $\mu\text{g/mL}$ - 10,000 $\mu\text{g/mL}$; 0.01 $\mu\text{g/mL}$ - 1000 $\mu\text{g/mL}$; 0.10 $\mu\text{g/mL}$ - 100 $\mu\text{g/mL}$; and 1 $\mu\text{g/mL}$ - 50 $\mu\text{g/mL}$.

[00138] In another preferred embodiment, the bacterial agent causing the bacterial infection is selected from the group comprising, but not limited to, gram positive bacteria. The microbe is preferably a gram positive bacterial agent selected from the group comprising *Staphylococcus aureus*, *Staphylococcus pseudintermedius*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus uberis*, *Enterococcus faecium*, *Enterococcus faecalis*, and *Clostridium difficile*.

[00139] In one preferred embodiment, the bacterial agent has no cell wall. Preferably, the bacterial agent is selected from the group comprising: *Mycoplasma* spp, *Mycoplasma agalactiae*, *Mycoplasma alkalescens*, *Mycoplasma amphoriforme*, *Mycoplasma arginini*, *Mycoplasma bovigenitalum*, *Mycoplasma bovirhinis*, *Mycoplasma bovis*, *Mycoplasma bovoculi*, *Mycoplasma buccale*, *Mycoplasma californicum*, *Mycoplasma canadense*, *Mycoplasma capricolum* subsp. *capricolum*, *Mycoplasma capricolum* subsp. *capripneumoniae*, *Mycoplasma conjunctivae*, *Mycoplasma cynos*, *Mycoplasma dispar*, *Mycoplasma equigenitalum*, *Mycoplasma faucium*, *Mycoplasma felis*, *Mycoplasma fermentans* (*incognitus* str.), *Mycoplasma gallisepticum* (MG), *Mycoplasma gateae*, *Mycoplasma genitalium*, *Mycoplasma haemocanis*,

Mycoplasma haemofelis, *Mycoplasma haemosuis* (formerly *Eperythrozoon suis*), *Mycoplasma hominis*, *Mycoplasma hyopneumoniae*, *Mycoplasma hyorhinis*, *Mycoplasma hyosynoviae*, *Mycoplasma iowae meleagridis* (MM), *Mycoplasma iowae*, *Mycoplasma leachii*, *Mycoplasma lipophilum*, *Mycoplasma meleagridis*, *Mycoplasma mycoides subsp capri*, *Mycoplasma mycoides subsp mycoides*, *Mycoplasma mycoides subsp. mycoides* (such as Contagious bovine pleuropneumonia CBPP), *Mycoplasma orale*, *Mycoplasma ovipneumoniae*, *Mycoplasma ovis*, *Mycoplasma penetrans*, *Mycoplasma pirum*, *Mycoplasma pneumoniae*, *Mycoplasma primatum*, *Mycoplasma putrefaciens*, *Mycoplasma salivarium*, *Mycoplasma spermatophilum*, *Mycoplasma suis*, *Mycoplasma synoviae* (MS), *Mycoplasma wenyonii*, *Mycoplasma*, *Ureaplasma spp*, *Ureaplasma parvum*, *Ureaplasma urealyticum*, *Ureaplasma*, and *Ureoplasma diversum*.

[00140] In another most preferred embodiment, the bacterial agent is *Staphylococcus aureus*.

[00141] In another preferred embodiment, the bacterial agent is resistant to a compound selected from the group comprising: one or more of aminoglycosides (for example gentamicin); ansamycins (for example rifampicin); anti-MRSA cephalosporins (for example ceftaroline); anti-staphylococcal β -lactams (or cephamycins) (for example oxacillin or ceftoxitin); carbapenems (for example ertapenem, imipenem, meropenem or doripenem); non-extended spectrum cephalosporins; 1st and 2nd generation cephalosporins (for example cefazolin or cefuroxime); extended-spectrum cephalosporins; 3rd and 4th generation cephalosporins (for example cefotaxime or ceftriaxone); cephamycins (for example ceftoxitin or cefotetan); fluoroquinolones (for example ciprofloxacin or moxifloxacin); folate pathway inhibitors (for example trimethoprim-sulphamethoxazole); fucidanes (for example fusidic acid); glycopeptides (for example vancomycin, teicoplanin or telavancin); glycyclines (for example tigecycline); lincosamides (for example clindamycin); lipopeptides (for example daptomycin); macrolides (for example erythromycin); oxazolidinones (for example linezolid or tedizolid); phenicols (for example chloramphenicol); phosphonic acids (for example fosfomycin); streptogramins (for example quinupristin-dalfopristin); and tetracyclines (for example tetracycline, doxycycline or minocycline). Preferably, the bacterial agent resistant to these compounds is gram positive.

[00142] In another most preferred embodiment, the bacterial agent is *Streptococcus pneumoniae*. The *Streptococcus pneumoniae* may be a strain that is resistant to one or more of β -lactams and macrolides.

[00143] In another most preferred embodiment, the bacterial agent is *Streptococcus pyogenes*.

[00144] In another most preferred embodiment, the bacterial agent is *Streptococcus agalactiae*.

[00145] In another most preferred embodiment, the bacterial agent is either *Enterococcus faecium* or *Enterococcus faecalis*. The *Enterococcus faecium* or *Enterococcus faecalis* may be a strain that is resistant to aminoglycosides (for example gentamicin (high level) or streptomycin (for example streptomycin (high level)); carbapenems (for example imipenem, meropenem or doripenem); fluoroquinolones (for example ciprofloxacin, levofloxacin or moxifloxacin); glycopeptides (for example vancomycin or teicoplanin); glycyclines (for example tigecycline); lipopeptides (for example daptomycin); oxazolidinones (for example linezolid); penicillins (for example ampicillin); streptogramins (for example quinupristin-dalfopristin); tetracycline (for example doxycycline or minocycline).

[00146] In another most preferred embodiment, the bacterial agent is *Clostridium difficile*.

[00147] The bacterial infection in the subject may cause a disease selected from the group comprising, but not limited to, nosocomial pneumonia caused by *Staphylococcus aureus* (MDR, XDR, PDR or methicillin-susceptible or -resistant strains), or invasive pneumococcal diseases such as pneumonia, bronchitis, acute sinusitis, otitis media, conjunctivitis, meningitis, bacteremia, sepsis, osteomyelitis, septic arthritis, endocarditis, peritonitis, pericarditis, cellulitis, and brain abscess caused by *Streptococcus pneumoniae* (including multi-drug resistant strains [MDRSP] such as those resistant to β -lactams and macrolides), complicated skin and skin structure infections, including diabetic foot infections, with or without concomitant osteomyelitis, caused by *Staphylococcus aureus* (methicillin-susceptible and -resistant strains), *Streptococcus pyogenes*, or *Streptococcus agalactiae*, uncomplicated skin and skin structure infections caused by *Staphylococcus aureus* (methicillin-susceptible and -resistant strains) or *Streptococcus pyogenes*, community-acquired pneumonia caused by *Streptococcus pneumoniae* (including multi-drug resistant strains [MDRSP], including cases with concurrent bacteraemia, or *Staphylococcus aureus* (methicillin-susceptible and -resistant strains) and *Staphylococcus aureus* bloodstream infections (bacteraemia), including those with right-sided infective endocarditis, caused by methicillin-susceptible and methicillin-resistant isolates, vancomycin-resistant enterococcus infections, including cases with concurrent bacteraemia, and treatment of *Clostridium difficile*-associated diarrhea (CDAD).

[00148] Gram negative organisms are important causes of many infectious diseases in humans and other animal species. Bone and joint infections (Gram-negative organisms or mixed bacteria, are an important cause of vertebral osteomyelitis and septic arthritis), cardiovascular system infections (including endocarditis caused by the HACEK group - *Haemophilus parainfluenzae*, *Haemophilus aphrophilus*, *Aggregatibacter*

actinomyces comitans, *Cardiobacterium hominis*, *Eikenella corrodens*, *Kingella kingae*), central nervous system infections (the commonest causes of bacterial meningitis are *Neisseria meningitidis*, *Streptococcus pneumoniae* and, in nonvaccinated young children, *Haemophilus influenzae* type b (Hib), in neonates and infants less than 3 months of age, *Streptococcus agalactiae* (group B streptococcus), *Escherichia coli* and other aerobic Gram-negative rods are important pathogens, brain abscess or subdural empyema, the infecting organism(s) vary with the underlying predisposing cause but where the likely site of origin is the ear, enteric Gram-negative bacilli are commonly involved), eye infections (common pathogens include *Haemophilus influenza*, *Neisseria gonorrhoeae* or *Chlamydia trachomatis*), gastrointestinal tract infections (a wide range of pathogens are implicated including enterotoxigenic *Escherichia coli* (ETEC), *Salmonella*, *Campylobacter*, *Shigella*, *Vibrio cholera* and *Yersinia enterocolitica*), genital infections (bacterial vaginosis is a polymicrobial clinical syndrome with high concentrations of anaerobic (eg *Mobiluncus* species) and other fastidious bacteria (including *Gardnerella vaginalis* and *Atopobium vaginae*), and *Mycoplasma hominis*; non-sexually acquired pelvic inflammatory disease (PID) is usually caused by mixed vaginal flora, including anaerobes, facultative Gram-negative bacteria and *Mycoplasma hominis*, while sexually acquired PID is usually initiated by *C. trachomatis* or *N. gonorrhoeae* with growing evidence that *M. genitalium* infection is involved in a significant minority of cases), intra-abdominal infections (peritonitis due to perforated viscus is usually a polymicrobial infection with aerobic and anaerobic bowel flora while spontaneous bacterial peritonitis (SBP) is usually caused by enteric Gram-negative bacilli, such as *Escherichia coli* and *Klebsiella* species, *Klebsiella pneumoniae* is an increasingly identified cause of liver abscess), community-acquired pneumonia (*Mycoplasma pneumoniae*, *Chlamydia* (*Chlamydia*) *pneumoniae*, *Chlamydia* (*Chlamydia*) *psittaci*, *Haemophilus influenza*, aerobic Gram-negative bacilli including *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Burkholderia pseudomallei*), otitis externa (including acute diffuse) (bacterial cultures commonly yield *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Proteus* and *Klebsiella* species), otitis media (including acute) (common bacterial pathogens include *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis*), sepsis (including severe) (including *Acinetobacter baumannii*, disseminated gonococcal sepsis, Gram-negative enteric bacteria, *Neisseria meningitidis* (meningococcal sepsis) and *Pseudomonas aeruginosa*), Systemic infections (Spotted fevers (*Rickettsia*) and scrub typhus (*Orientia*), Brucellosis, Cat-scratch disease and other *Bartonella* infections, Leptospirosis, Lyme disease, Melioidosis, Q fever, Typhoid and paratyphoid fevers (enteric fevers), urinary tract infections (acute cystitis, acute pyelonephritis, recurrent urinary tract infections and atheter-associated bacteriuria and urinary tract infections).

[00149] In humans gram negative bacteria are common causes of intra-abdominal infections (IAIs), urinary tract infections (UTIs), hospital acquired pneumonia, and bacteraemia.

Escherichia coli (*E. coli*), *Klebsiella pneumoniae* (*K. pneumoniae*), and *Pseudomonas aeruginosa* (*P. aeruginosa*) are important pathogens in the hospital setting, accounting for 27% of all pathogens and 70% of all Gram-negative pathogens causing healthcare-associated infections [Sievert DM, Ricks P, Edwards JR, et al. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2009-2010. *Infect Control Hosp Epidemiol.* 2013;34:1-14.].

[00150] Gram negative bacteria are showing rising rates of resistance to current therapies. The production of extended-spectrum β -lactamase (ESBL) enzymes is a common mechanism of resistance. Rates of ESBL-producing *E. coli* and *K. pneumoniae* have risen substantially, with the result that these bacteria are increasingly resistant to widely used antimicrobials.

[00151] *P. aeruginosa* is the most common Gram-negative cause of nosocomial pneumonia and the second most common cause of catheter-related UTIs in the U.S.

[00152] *E. coli* is the most common cause of UTIs. Cases of UTI caused by ESBL-producing *E. coli* and *K. pneumoniae* as well as *P. aeruginosa*, including MDR strains, are increasing. ESBL-producing *E. coli* and *K. pneumoniae* are also frequently isolated in patients with complicated IAI (cIAI).

[00153] *P. aeruginosa* is a clinically challenging and virulent pathogen that can be a cause of common infections in humans such as nosocomial pneumonia, UTI, IAI, and bloodstream infections. *P. aeruginosa* is the most common Gram-negative organism causing ventilator associated pneumonia and the second most common cause of catheter-associated UTIs.

[00154] The increase in the number of infections caused by Gram-negative bacteria is being accompanied by rising rates of resistance. Treatment options to meet this challenge are increasingly limited. There is a critical need for new antibiotics to meet the needs of patients now and in the future.

[00155] In one preferred aspect, more than one compound of the invention is administered to the subject.

[00156] In another preferred embodiment, a compound of the invention, or a therapeutically acceptable salt thereof, is administered together with a compound or agent that removes or substantially removes or reduces the integrity of the cell wall of the bacterial agent. As an example, the compound is selected from the group consisting of: β lactams, fosfomycin,

lysozyme, polymyxins and chelating agents such as ethylenediaminetetraacetic acid (EDTA). As an example, the agent is an immunological agent (such as an antibody or vaccine) that reduces the integrity of the cell wall. In one preferred embodiment, the compound, or a therapeutically acceptable salt thereof, is administered together with a compound that removes or substantially removes or weakens the integrity of the outer cell wall of a gram negative bacterial agent.

[00157] According to another aspect of the invention, there is provided an antibacterial pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I, or a therapeutically acceptable salt thereof. Preferably, the composition is an antibacterial pharmaceutical composition.

[00158] According to another aspect of the invention, there is provided an antibacterial veterinary composition comprising a therapeutically effective amount of a compound of Formula I, or a therapeutically acceptable salt thereof. Preferably, the composition is an anti-bacterial veterinary composition.

[00159] The pharmaceutical composition may optionally include a pharmaceutically acceptable excipient or carrier. The veterinary composition may optionally include a veterinary acceptable excipient or carrier.

[00160] The pharmaceutical or veterinary composition of the invention preferably contains a compound of Formula I, or a pharmaceutically acceptable salt, at a concentration of selected from the group consisting of: 1 mg/g to 500 mg/g; 5 mg to 400 mg/g; 10 mg/g to 200 mg/g; 20 mg/g to 100 mg/g; 30 mg/g to 70 mg/g; and 40 mg/g to 60 mg/g.

[00161] In another embodiment, the pharmaceutical or veterinary composition comprises impurities, wherein the quantity of impurities as a percentage of the total weight of the composition is selected from the group consisting of: less than 20% impurities (by total weight of the composition); less than 15% impurities; less than 10% impurities; less than 8% impurities; less than 5% impurities; less than 4% impurities; less than 3% impurities; less than 2% impurities; less than 1% impurities; less than 0.5% impurities; less than 0.1% impurities. In one embodiment, the pharmaceutical or veterinary composition comprises microbial impurities or secondary metabolites, wherein the quantity of microbial impurities as a percentage of the total weight of the composition is selected from the group consisting of: less than 5%; less than 4%; less than 3%; less than 2%; less than 1%; less than 0.5%; less than 0.1%; less than 0.01%; less than 0.001%. In one embodiment, the pharmaceutical or veterinary composition is sterile and stored in a sealed and sterile container. In one embodiment, the pharmaceutical or veterinary composition contains no detectable level of microbial contamination.

[00162] The pharmaceutical or veterinary composition of the invention may comprise a further antimicrobial agent. The further antimicrobial agent may be an antifungal agent or antibacterial agent. The method of treating or preventing a bacterial infection or colonisation in a subject, may also comprise the administration of a compound of the invention with a further antimicrobial agent.

[00163] The pharmaceutical or veterinary composition of the invention may comprise more than one compound of the invention. For example, a combination of compounds. The method of treating or preventing a bacterial infection or colonisation in a subject, may also comprise the administration of more than one compound of the invention.

[00164] In one embodiment, the antifungal agent is selected from the group comprising, but not limited to naturally occurring agents including Echinocandins (Anidulafungin, Caspofungin, Micafungin), Polyenes (Amphotericin B, Candicidin, Filipin, Fungichromin (Pentamycin), Hachimycin, Hamycin, Lucensomycin, Mepartricin, Natamycin, Nystatin, Pecilocin, Perimycin), and other naturally occurring antifungal agents including Griseofulvin, Oligomycins, Pyrrolnitrin, Siccanin, and Viridin. The antifungal agent may be a synthetic compound selected from the group comprising, but not limited to Allylamines (Butenafine, Naftifine, Terbinafine) Imidazoles (Bifonazole, Butoconazole, Chlormidazole, Climbazole, Croconazole (Cloconazole), Clotrimazole, Eberconazole, Econazole, Enilconazole, Fenticonazole, Flutrimazole, Fosfluconazole, Isoconazole, Ketoconazole, Lanoconazole, Luliconazole, Miconazole, Neticonazole, Omoconazole, Oxiconazole Nitrate, Parconazole, Sertaconazole, Sulconazole, Tioconazole), Thiocarbamates (Liranaftate, Tolciclate, Tolindate, Tolnaftate), Triazoles (Fluconazole, Isavuconazole, Itraconazole, Posaconazole, Ravuconazole, Saperconazole, Terconazole, Voriconazole), and other synthetic agents such as Acrisorcin, Amorolfine, Bromosalicylchloranilide (Bromochlorosalicylanilide), Buclosamide, Calcium Propionate, Chlorphenesin, Ciclopirox, Cloxyquin (Cloxiquine), Coparaffinate, Exalamide, Flucytosine, Haloprogin, Hexetidine, Loflucarban, Nifuratel, Nifuroxime, Piroctone, Potassium Iodide, Propionic Acid, Pyrithione, Salicylanilide, Sodium Parachlorobenzoate, Sodium Propionate, Sulbentine, Tenonitroazole, Triacetin, Trimetrexate, Undecylenic Acid (Undecenoic Acid), and Zinc Propionate.

[00165] The composition of the invention may comprise an antibiotic adjunct selected from the group comprising, but not limited to, β -Lactamase Inhibitors (Avibactam, Clavulanic Acid, Sulbactam, Sulfamicillin, Tazobactam), Renal Dipeptidase Inhibitors (Cilastatin), and Renal Protectant (Betamipron).

[00166] In one embodiment, the composition of the invention comprises a further antibiotic selected from the group comprising, but not limited to, 2,4-DIAMINOPYRIMIDINES,

including Baquiloprim, Brodimoprim, Iclaprim, Ormetoprim, Pyrimethamine, Tetroxoprim, Trimethoprim; AMINOCOUMARINS, including Novobiocin; AMINOCYCLITOLS, including Spectinomycin; AMINOGLYCOSIDES, including Amikacin, Apramycin, Arbekacin, Bekanamycin, Butirosin, Dibekacin, Dihydrostreptomycin, Etimicin, Fortimicins (Astromicin), Framycetin, Gentamicin, Hygromycin B, Isepamicin, Kanamycin, Micronomicin, Neomycin, Netilmicin, Paromomycin, Plazomicin, Ribostamycin, Sisomicin, Streptomycin, Tobramycin, Verdamicin; AMINOMETHYLCYCLINES, including Omadacycline; AMPHENICOLS, including Azidamfenicol, Chloramphenicol, Florfenicol, Thiamphenicol; ANSAMYCINS, including Rifabutin, Rifamide, Rifampin (Rifampicin), Rifamycin, Rifapentine, Rifaximin; ANTISEPTIC AGENTS, including Acridine derivatives (including acriflavine, aminoacridine, ethacridine, proflavine), Bispyridines (including octenidine dihydrochloride), Brominated salicylanilides (including bromsalans), Chlorhexidine, Phenol derivatives (including thymol and triclosan), Quarternary ammonium compounds (including Alkyldimethylethylbenzyl Ammonium Chloride, benzalkonium chloride, cetylpyridinium chloride, benzethonium chloride, cetrimonium); ANTITUBERCULAR AGENTS, including Cycloserine, Delamanid, Ethambutol, Ethionamide, Isoniazid (Ftivazide), Morinamide, *p*-Aminosalicylic Acid (PAS), Protionamide, Pyrazinamide, Terizidone, Thioacetazone, Tiocarlide; ARSENICALS, including Arsanilic Acid, Roxarsone; BACTERIOCINS, including Nisin, Brilacidin (PMX-30063); β -LACTAM CARBACEPHEMS, including Loracarbef; β -LACTAM CARBAPENEMS, including Biapenem, Doripenem, Ertapenem, Faropenem, Imipenem, Meropenem, Panipenem, Razupenem, Ritipenem, Sulopenem, Tebipenem, Tomopenem; β -LACTAM CEPHALOSPORINS, including Cefacetile, Cefaclor, Cefadroxil, Cefalexin, Cefaloglycin, Cefalonium, Cefaloridine, Cefalothin, Cefamandole, Cefapirin, Cefatrizine, Cefazaflur, Cefazedone, Cefazolin, Cefcapene, Cefdinir, Cefditoren, Cefepime, Cefetamet, Cefixime, Cefmenoxime, Cefodizime, Cefonicid, Cefoperazone, Ceforanide, Cefoselis, Cefotaxime, Cefotiam, Cefovecin, Cefozopran, Cefpimizole, Cefpiramide, Cefpirome, Cefpodoxime, Cefprozil, Cefquinome, Cefradine, Cefroxadine, Cefsulodin, Ceftaroline, Ceftazidime, Cefteram, Ceftezole, Ceftibuten, Cefitofur, Ceftizoxime, Ceftobiprole, Ceftolozane, Ceftradine, Ceftrezole, Ceftriaxone, Cefroxadine, Cefuroxime, Cefuzonam, Pivcefalexin; β -LACTAM CEPHAMYCINS, including Cefbuperazone, Cefmetazole, Cefminox, Cefotetan, Cefoxitin; β -LACTAM MONOBACTAMS, including Aztreonam, Carumonam, Tigemonam; β -LACTAM OXACEPHEMS, including Flomoxef, Latamoxef, Moxalactam; β -LACTAM PENICILLINS, including Amdinocillin (Mecillinam), Amoxicillin, Ampicillin, Apalcillin, Aspoxicillin, Azidocillin, Azlocillin, Bacampicillin, Carbenicillin, Carindacillin, Ciclacillin, Clemizole Penicillin, Clometocillin, Cloxacillin, Cyclacillin, Diclloxacillin, Epicillin, Fenbenicillin, Floxacillin (Flucloxacillin), Hetacillin, Lenampicillin, Mecillinam, Metampicillin, Methicillin Sodium, Mezlocillin, Nafcillin, Oxacillin, Penamecillin, Penethamate Hydriodide, Penicillin G, Penicillin G Benzathine, Penicillin G Procaine, Penicillin N, Penicillin O, Penicillin V, Phenethicillin Potassium, Piperacillin, Pivampicillin, Pivmecillinam, Propicillin,

Quinacillin, Sulbenicillin, Sultamicillin, Talampicillin, Temocillin, Ticarcillin; BICYCLOMYCINS, including Bicozamycin; BORON CONTAINING ANTIBACTERIAL AGENTS, including AN3365 (aminomethylbenzoxaboroles), GSK2251052 (leucyl-tRNA synthetase inhibitors); CYCLIC ESTERS, including Fosfomycin; FATTY ACID SYNTHESIS INHIBITORS (Fabi), AFN-1252, MUT056399, FAB-001; FLUOROQUINOLONES, including Avarofloxacin, Balofloxacin, Besifloxacin, Chinoxacin, Cinoxacin, Ciprofloxacin, Clinafloxacin, Danofloxacin, Delafloxacin, Difloxacin, Enoxacin, Enrofloxacin, Finafloxacin, Fleroxacin, Flumequine, Garenoxacin, Gatifloxacin, Gemifloxacin, Grepafloxacin, Ibafoxacin, Levofloxacin, Lomefloxacin, Marbofloxacin, Miloxacin, Moxifloxacin, Nadifloxacin, Norfloxacin, Ofloxacin, Orbifloxacin, Pazufloxacin, Pefloxacin, Pradofloxacin, Prulifloxacin, Rosoxacin, Rufloxacin, Sarafloxacin, Sitafloxacin, Sparfloxacin, Terafloxacin, Tosufloxacin, Trovafloxacin, Zabofloxacin; FUSIDANES, including Fusidic Acid; GLYCOLIPODEPSIPEPTIDE, including Ramoplanin; GLYCOPEPTIDES, including Avoparcin, Dalbavancin, Norvancomycin, Oritavancin, Teicoplanin, Telavancin, Vancomycin; GLYCOPHOSPHOLIPIDS, including Bambermycins (bambermycin, moenomycins, flavophospholipol); GLYCYLCYCLINES, including Tigecycline; HYBRIDS, Cadazolid (Oxazolidinone-quinolone), TD-1792 (glycopeptide-cephalosporin); LINCOSAMIDES, including Clindamycin, Lincomycin, Pirlimycin; LIPOPEPTIDES, including Daptomycin, Surotomycin; MACROLIDES, including Azithromycin, Carbomycin, Cethromycin, Clarithromycin, Dirithromycin, Erythromycin, Fidaxomicin, Flurithromycin, Gamithromycin, Josamycin, Kitasamycin, Leucomycin, Meleumycin, Midecamycins, Miodamycin, Miosamycin, Oleandomycin, Primycin, Rokitamycin, Rosaramicin, Roxithromycin, Sedecamycin, Solithromycin, Spiramycin, Telithromycin, Terdecamycin, Tildipirosin, Tilmicosin, Troleandomycin, Tulathromycin, Tylosin, Tylvalosin; NITROFURANS, including Furalfadone, Furazidin, Furazolidone, Furazolum Chloride, Nifuratel, Nifurfoline, Nifuroxazide, Nifurpirinol, Nifurtinol, Nifurzide, Nitrofuril, Nitrofurantoin, Nitrofurazone; NITROIMIDAZOLES, including Dimetridazole, Metronidazole, Ornidazole, Ronidazole, Secnidazole, Tinidazole; OLIGOSACCHARIDES, including Avilamycin, Everminomicin; OTHER ANTIBACTERIAL AGENTS, including Auriclosene, Chloroxine, Chlorquinaldol, Clioquinol, Clofoctol, Halquinol, Lotilibcin, Mandelic Acid, Methenamine (hexamine), Nitazole, Nitroxoline, Perchlozone, Taurolidine, Thenoic Acid, Xibornol; OXAZOLIDINONES, including Eperezolid, Linezolid, Posizolid, Radezolid, Sutezolid, Tedizolid (Torezolid); PEPTIDE DEFORMYLASE INHIBITORS, including GSK1322322; PEPTIDES, including Omiganan, Pexiganan; PLEUROMUTILINS, including Retapamulin, Tiamulin, Valnemulin; POLYETHER IONOPHORES, including Laidlomycin, Lasalocid, Maduramicin, Monensin, Narasin, Salinomycin, Semduramicin; POLYMYXINS, including Colistin, Polymyxin B; POLYPEPTIDES, including Amphomycin, Bacitracin, Capreomycin, Enduracidin, Enramycin, Enviomycin, Fusafungine, Gramicidin(s), Isegran, Magainins, Nosiheptide, Ristocetin, Thiostrepton, Tuberactinomycin, Tyrocidine, Tyrothricin, Viomycin; PSEUDOMONIC ACIDS, including Mupirocin; QUINOLONES, including

Nalidixic Acid, Nemonoxacin, Oxolinic Acid, Ozenoxacin, Pipemidic Acid, Piromidic Acid; QUINOXALINES, including Carbadox, Olaquinox; RIMINOFEAZINES, including Clofazimine; STATINS, including Atorvastatin, Fluvastatin, Lovastatin, Mevastatin, Pitavastatin, Pravastatin, Rosuvastatin, Simvastatin; STREPTOGRAMINS, including Dalfopristin, Flopristin, Linopristin, Pristinamycin, Quinupristin, Virginiamycin; STREPTOTHRICINS, including Nourseothricin; SULFONAMIDES, including Acetyl Sulfamethoxypyrazine, Chloramine-B, Chloramine-T, Dichloramine T, Formosulfathiazole, Mafenide, N4-Sulfanilysulfanilamide, Noprysulfamide, N-Sulfanilyl-3,4-xylamide, Ormaosulfathiazole, Phthalylsulfacetamide, Phthalylsulfathiazole, Salazosulfadimidine, Succinylsulfathiazole, Sulfabenzamide, Sulfacarbamide, Sulfacetamide, Sulfachlorpyridazine, Sulfachrysoidine, Sulfaclozine, Sulfacytine, Sulfadiazine, Sulfadicramide, Sulfadimethoxine, Sulfadimidine, Sulfadoxine, Sulfathiazole, Sulfaguanidine, Sulfaguanole, Sulfalene, Sulfaloximic Acid, Sulfamerazine, Sulfameter, Sulfamethazine, Sulfamethazole, Sulfamethomidine, Sulfamethoxazole, Sulfamethoxypyridazine, Sulfamethylthiazole, Sulfametopyrazine, Sulfamethoxazole, Sulfamidochrysoidine, Sulfamonomethoxine, Sulfamoxole, Sulfanilamide, Sulfanilurea, Sulfaperine, Sulfaphenazole, Sulfaproxyline, Sulfapyrazine, Sulfapyridine, Sulfapyrimidine, Sulfathiazole, Sulfathiourea, Sulfatiazole, Sulfisomidine, Sulfisoxazole (Sulfafurazole); SULFONES, including Acediasulfone, Dapsone, Glucosulfone Sodium, p-Sulfanilylbenzylamine, Succisulfone, Sulfanilic Acid, Sulfoxone Sodium, Thiazolsulfone; TETRACYCLINES, including Chlortetracycline, Clomocycline, Demeclocycline, Doxycycline, Eravacycline, Guamecycline, Lymecycline, Meclocycline, Methacycline, Minocycline, Oxytetracycline, Penimepicycline, Pipacycline, Rolitetracycline, Sarecycline, and Tetracycline.

[00167] The composition of the invention may further comprise an excipient selected from the group comprising, but not limited to, binders and compression aids, coatings and films, colouring agents diluents and vehicles disintegrants, emulsifying and solubilising agents, flavours and sweeteners, repellents, glidants and lubricants, plasticisers, preservatives, propellants, solvents, stabilisers, suspending agents and viscosity enhancers.

[00168] According to a further aspect of the invention, there is provided a medical device when used in a method of treating or preventing a bacterial infection in the subject.


[00169] According to further aspect of the invention, there is provided a medical device comprising the composition of the invention. The composition of the invention may be any slow release form, and/or in the form of a coating of the medical device.

[00170] The medical device may be in a form selected from the group comprising: an implant, a plaster, a bandage, and other dressing applied to a bacterial infection in a subject.

[00171] According to further aspect of the invention, there is provided a method of killing bacteria, the method including the step of contacting the bacteria with a compound of the invention, or a therapeutically acceptable salt thereof.

[00172] According to further aspect of the invention, there is provided the use of a compound of the invention, or a therapeutically acceptable salt thereof, to kill bacteria, said use comprising the step of contacting the bacteria with a compound of the invention, or a therapeutically acceptable salt thereof.

[00173] Terms used herein will have their customary meanings in the art unless specified. As used herein, the term robenidine, NCL812 (also known as 1,3-bis[(*E*)-(4-chlorophenyl)methyleneamino]guanidine) refers to a compound having the following chemical structure:



BRIEF DESCRIPTION OF THE DRAWINGS

[00174] Further features of the present invention are more fully described in the following description of several non-limiting embodiments thereof. This description is included solely for the purposes of exemplifying the present invention. It should not be understood as a restriction on the broad summary, disclosure or description of the invention as set out above. The description will be made with reference to the accompanying drawings in which:

Figure 1 shows a graph of the effect of NCL812 on DNA macromolecular synthesis in *Staphylococcus aureus* according to example 2;

Figure 2 shows a graph of the effect of NCL812 on RNA macromolecular synthesis in *Staphylococcus aureus* according to example 2;

Figure 3 shows a graph of the effect of NCL812 on protein macromolecular synthesis in *Staphylococcus aureus* (ATCC29213) according to example 2;

Figure 4 shows a graph of the effect of NCL812 on cell wall macromolecular synthesis in *Staphylococcus aureus* (ATCC29213) according to example 2;

Figure 5 shows a graph of the effect of NCL812 on lipid macromolecular synthesis in *Staphylococcus aureus* (ATCC29213) according to example 2;

Figure 6 shows a graph summarising the effect of NCL812 on macromolecular synthesis in *Staphylococcus aureus* (ATCC29213) according to example 2;

Figure 7 shows a graph of the effect of NCL812 on ATP release from *Staphylococcus aureus* (ATCC29213) according to example 3;

Figure 8 shows a graph showing the average melting point peaks for the negative derivative dF/dT after real-time polymerase chain reaction of the *mecA* gene in methicillin-resistant *S. aureus* isolates grouped by *mec* gene complexes, A (n=4), B (n=10), C2 (n=4) and unclassified (n=2). Groups indicated with different superscripts are significantly different ($P < 0.05$), according to example 4;

Figure 9 shows a graph of the optical densities of the unsupplemented growth control, ampicillin and different concentrations of antibacterial agent NCL812 against methicillin-susceptible *S. aureus* ATCC 49775 using broth microdilution methodology according to example 4. The concentrations of NCL812 tested were at the MIC and four times the MIC determined under test conditions, up to 24 h incubation. Ampicillin was tested at the MIC. Bactericidal activity was tested at 0, 1, 2, 4, 8, 12, and 24 h for antibacterials;

Figure 10 shows a graph of kill kinetic curves for methicillin-susceptible *S. aureus* ATCC 49775 demonstrating bactericidal activity of NCL812 using the Clinical and Laboratory Standards Institute macrodilution methodology in a 10 ml vial according to example 4. The concentrations of antibacterials tested were at 1× and 4× the MIC determined under test conditions. Bactericidal activity was determined at 0, 1, 2, 4, 8, 12 and 24 h after antibacterial addition. Bactericidal activity was defined as a $3\log_{10}$ (99.9%) decrease in the number viable bacteria from the initial inoculum size;

Figure 11 shows a graph indicating the change of pH during macro-broth dilution assay for *S. pneumoniae* strain D39 exposed to 4 µg/mL in NCL812 and 0.0023 µg/mL ampicillin according to example 5;

Figure 12 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with NCL812 according to example 5;

Figure 13 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with NCL062 according to example 5;

Figure 14 shows a graph illustrating in the 14-hour time-kill of *S. pneumoniae* strain D39 treated with NCL812 according to example 5;

Figure 15 shows a graph illustrating in the 14-hour time-kill of *S. pneumoniae* strain D39 treated with NCL062 according to example 5;

Figure 16 shows a graph illustrating the 14-hour time-kill of *S. pneumoniae* strain D39 treated with ampicillin according to example 5;

Figure 17 shows a graph illustrating the 12-hour time-kill of *S. pneumoniae* strain D39 treated with NCL812, according to example 5;

Figure 18 shows a graph illustrating the 12-hour time-kill of *S. pneumoniae* strain D39 treated with NCL062, according to example 5;

Figure 19 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with ampicillin according to example 5;

Figure 20 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with erythromycin according to example 5;

Figure 21 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with NCL812 and 5% choline chloride;

Figure 22 shows a graph illustrating the 12-hour time-kill of *S. pneumoniae* strain D39 treated with NCL812 and 5% choline chloride according to example 5;

Figure 23 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with NCL062 and 5% choline chloride according to example 5;

Figure 24 shows a graph illustrating the 12-hour time-kill of *S. pneumoniae* strain D39 treated with NCL062 and 5% choline chloride according to example 5;

Figure 25 shows a graph illustrating the 48-hour time-kill of *S. pneumoniae* strain D39 treated with ampicillin and choline chloride according to example 5;

Figure 26 shows a graph of the relative MBC of D39 treated with NCL812 or NCL062 for 48 hours according to example 5;

Figure 27 shows a graph illustrating the relative minimal bactericidal concentration (MBC) of *S. pneumoniae* strain D39 treated with ampicillin over a 48 h time period according to example 5;

Figure 28 shows a graph illustrating the relative MBC for *S. pneumoniae* strain D39 treated with erythromycin over a 48 h time period according to example 5;

Figure 29 shows a graph illustrating the viable count ($\log_{10}$ CFU/ml) of *S. pneumoniae* strain D39 treated with NCL812 from a macro-broth dilution of time-kill over 24 hours according to example 5;

Figure 30 shows a graph illustrating the viable count ($\log_{10}$ CFU/ml) of *S. pneumoniae* strain D39 treated with ampicillin from a macro-broth dilution of time-kill over 24 hours according to example 5;

Figure 31 is a bar graph illustrating the mean cell membrane thickness of treated and untreated D39 according to example 5;

Figure 32 is a bar graph illustrating the mean width of periplasmic space of treated (16 μ g/mL NCL812) and untreated D39 samples according to example 5;

Figure 33 shows the kill kinetics of MRSA 580 isolate obtained at different concentrations of NCL812 over a period of 8 hours according to example 7;

Figure 34 shows the kill kinetics of MRSA 580 in different concentrations of NCL812 over a period of 24 h according to example 7;

Figure 35 shows the kill kinetics of MRSA 698 in different concentrations of NCL812 over a period of 24 h according to example 7;

Figure 36 shows the kill kinetics of VRE 26c(dc) at different concentrations of NCL812 over a period of 24 h according to example 7;

Figure 37 shows the kill kinetics of VRE 16c(dc) at different concentrations of NCL812 over a period of 24 h according to example 8;

Figure 38 shows the kill kinetics assay of *Staphylococcus aureus* KC01 at different concentrations of NCL812, up to 24 h incubation according to example 8;

Figure 39 shows the kill kinetics assay of *Enterococcus faecalis* USA01 at different concentrations of NCL812, up to 24 h incubation according to example 8; and

Figure 41 is a graph illustrating the cumulative release of NCL812 and NCL099 from Formulation B according to example 10.

DESCRIPTION OF EMBODIMENTS

General

[00175] Before describing the present invention in detail, it is to be understood that the invention is not limited to particular exemplified methods or compositions disclosed herein. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

[00176] The applications that are mentioned herein are referred to simply for the purpose of describing and disclosing the procedures, protocols, and reagents referred to in the publication which may have been used in connection with the invention. The citation of any publications referred to herein is not to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

[00177] In addition, the carrying out of the present invention makes use of, unless otherwise indicated, conventional microbiological techniques within the skill of the art. Such conventional techniques are known to the skilled worker.

[00178] As used herein, and in the appended claims, the singular forms “a”, “an”, and “the” include the plural unless the context clearly indicates otherwise.

[00179] Unless otherwise indicated, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any materials and methods similar to, or equivalent to, those described herein may be used to carry out the present invention, the preferred materials and methods are herein described.

[00180] The invention described herein may include one or more ranges of values (e.g. size, concentration, dose etc). A range of values will be understood to include all values within the range, including the values defining the range, and values adjacent to the range that lead to the same or substantially the same outcome as the values immediately adjacent to that value which define the boundary of the range.

[00181] The pharmaceutical or veterinary compositions of the invention may be administered in a variety of unit dosages depending on the method of administration, target site,

physiological state of the patient, and other medicaments administered. For example, unit dosage form suitable for oral administration include solid dosage forms such as powder, tablets, pills, and capsules, and liquid dosage forms, such as elixirs, syrups, solutions and suspensions. The active ingredients may also be administered parenterally in sterile liquid dosage forms. Gelatin capsules may contain the active ingredient and inactive ingredients such as powder carriers, glucose, lactose, sucrose, mannitol, starch, cellulose or cellulose derivatives, magnesium stearate, stearic acid, sodium saccharin, talcum, magnesium carbonate, and the like.

[00182] The phrase "therapeutically effective amount" as used herein refers to an amount sufficient to inhibit bacterial growth associated with a bacterial infection or colonisation. That is, reference to the administration of the therapeutically effective amount of a compound of Formula I according to the methods or compositions of the invention refers to a therapeutic effect in which substantial bacteriocidal or bacteriostatic activity causes a substantial inhibition of bacterial infection. The term "therapeutically effective amount" as used herein, refers to a sufficient amount of the composition to provide the desired biological, therapeutic, and/or prophylactic result. The desired results include elimination of bacterial infection or colonisation or reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. An effective amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation. In relation to a pharmaceutical or veterinary composition, effective amounts can be dosages that are recommended in the modulation of a diseased state or signs or symptoms thereof. Effective amounts differ depending on the composition used and the route of administration employed. Effective amounts are routinely optimized taking into consideration pharmacokinetic and pharmacodynamic characteristics as well as various factors of a particular patient, such as age, weight, gender, etc and the area affected by disease or disease causing microbes.

[00183] As referred to herein, the terms "treatment" or "treating" refers to the full or partial removal of the symptoms and signs of the condition. For example, in the treatment of a bacterial infection or colonisation, the treatment completely or partially removes the signs of the infection. Preferably in the treatment of infection, the treatment reduces or eliminates the infecting bacterial pathogen leading to microbial cure.

[00184] As referred to herein, the term "bacteria" refers to members of a large domain of prokaryotic microorganisms. Typically a few micrometres in length, bacteria have a number of shapes, ranging from spheres to rods and spirals and can be present as individual cells or present in linear chains or clusters of variable numbers and shape. Preferably the terms "bacteria" and its adjective "bacterial" refer to bacteria such as the Gram positive *Staphylococcus* spp, *Streptococcus* spp, *Bacillus* spp, *Enterococcus* spp, *Listeria* spp, and

anaerobic bacteria; Gram negative *Escherichia coli*, *Enterobacter* spp, *Klebsiella* spp and *Pseudomonas* spp; and the cell wall free bacteria such as *Mycoplasma* spp and *Ureaplasma* spp. The terms may refer to an antibiotic-sensitive strain or an antibiotic-resistant strain. In a preferred embodiment, the terms refer to MRSA or MRSP. In another preferred embodiment, the terms refer to MDR *Staphylococcus* spp, *Streptococcus* spp, *Enterococcus* spp, *Clostridium difficile*, *Escherichia coli*, *Enterobacter* spp, *Klebsiella* spp and *Pseudomonas* spp.

[00185] Referred to herein, the term "methicillin-resistant bacteria" (such as methicillin-resistant *Staphylococcus*) refers a bacteria isolate that demonstrates resistance at any dose to all β -lactams including penicillins, carbapenems and first to fourth generation cephalosporins, but not to the fifth generation anti-MRSA cephalosporins (for example ceftaroline). Multidrug-resistant (MDR) is defined as acquired non-susceptibility to at least one agent in three or more antimicrobial categories, extensively drug-resistant (XDR) is defined as non-susceptibility to at least one agent in all but two or fewer antimicrobial categories (i.e. bacterial isolates remain susceptible to only one or two categories) and pandrug-resistant (PDR) is defined as non-susceptibility to all agents in all antimicrobial categories currently available.

[00186] An example of susceptible, MDR, XDR and PDR bacteria includes the following. Wild type, antibacterial unexposed isolates of *Staphylococcus aureus* that are likely to be susceptible to all of the following antibacterial categories (and agents): aminoglycosides (for example gentamicin); ansamycins (for example rifampicin); anti-MRSA cephalosporins (for example ceftaroline); anti-staphylococcal β -lactams (for example oxacillin or cefoxitin); carbapenems (for example ertapenem, imipenem, meropenem or doripenem); non-extended spectrum cephalosporins; 1st and 2nd generation cephalosporins (for example cefazolin or cefuroxime); extended-spectrum cephalosporins; 3rd and 4th generation cephalosporins (for example cefotaxime or ceftriaxone); cephamycins (for example cefoxitin or cefotetan); fluoroquinolones (for example ciprofloxacin or moxifloxacin); folate pathway inhibitors (for example trimethoprim-sulphamethoxazole); fucidanes (for example fusidic acid); glycopeptides (for example vancomycin, teicoplanin or telavancin); glycyclines (for example tigecycline); lincosamides (for example clindamycin); lipopeptides (for example daptomycin); macrolides (for example erythromycin); oxazolidinones (for example linezolid or tedizolid); phenicols (for example chloramphenicol); phosphonic acids (for example fosfomycin); streptogramins (for example quinupristin-dalfopristin); and tetracyclines (for example tetracycline, doxycycline or minocycline). Isolates that are non-susceptible to more than one agent in more than three antimicrobial categories are classified as MDR (all MRSA, for example, meet the definition of MDR). Isolates that are non-susceptible to more than one agent in all but one or two antimicrobial categories are classified as XDR. Isolates that are non-susceptible to all listed antibacterial agents are PDR.

[00187] Pharmaceutically and veterinary acceptable salts include salts which retain the biological effectiveness and properties of the compounds of the present disclosure and which are not biologically or otherwise undesirable. In many cases, the compounds disclosed herein are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto. Acceptable base addition salts can be prepared from inorganic and organic bases. Salts derived from inorganic bases, include by way of example only, sodium, potassium, lithium, ammonium, calcium and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary and tertiary amines, such as by way of example only, alkyl amines, dialkyl amines, trialkyl amines, substituted alkyl amines, di(substituted alkyl) amines, tri(substituted alkyl) amines, alkenyl amines, dialkenyl amines, trialkenyl amines, substituted alkenyl amines, di(substituted alkenyl) amines, tri(substituted alkenyl) amines, cycloalkyl amines, di(cycloalkyl) amines, tri(cycloalkyl) amines, substituted cycloalkyl amines, disubstituted cycloalkyl amines, trisubstituted cycloalkyl amines, cycloalkenyl amines, di(cycloalkenyl) amines, tri(cycloalkenyl) amines, substituted cycloalkenyl amines, disubstituted cycloalkenyl amines, trisubstituted cycloalkenyl amines, aryl amines, diaryl amines, triaryl amines, heteroaryl amines, diheteroaryl amines, triheteroaryl amines, heterocyclic amines, diheterocyclic amines, triheterocyclic amines, mixed di- and tri-amines where at least two of the substituents on the amine are different and are selected from the group consisting of alkyl, substituted alkyl, alkenyl, substituted alkenyl, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, aryl, heteroaryl, heterocyclic, and the like. Also included are amines where the two or three substituents, together with the amino nitrogen, form a heterocyclic or heteroaryl group.

[00188] Pharmaceutically and veterinary acceptable acid addition salts may be prepared from inorganic and organic acids. The inorganic acids that can be used include, by way of example only, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. The organic acids that can be used include, by way of example only, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like.

[00189] The pharmaceutically or veterinary acceptable salts of the compounds useful in the present disclosure can be synthesized from the parent compound, which contains a basic or acidic moiety, by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack

Publishing Company, Easton, Pa. (1985), p. 1418. Examples of such acceptable salts are the iodide, acetate, phenyl acetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, γ -hydroxybutyrate, β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, hexyne-1,6-dioate, caproate, caprylate, chloride, cinnamate, citrate, decanoate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, terephthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzenesulfonate, p-bromophenylsulfonate, chlorobenzenesulfonate, propanesulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, merhanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like.

[00190] The pharmaceutical or veterinary compositions of the invention may be formulated in conventional manner, together with other pharmaceutically acceptable excipients if desired, into forms suitable for oral, parenteral, or topical administration. The modes of administration may include parenteral, for example, intramuscular, subcutaneous and intravenous administration, oral administration, topical administration and direct administration to sites of infection such as intraocular, intraaural, intrauterine, intranasal, intramammary, intraperitoneal, intralesional, etc.

[00191] The pharmaceutical or veterinary compositions of the invention may be formulated for oral administration. Traditional inactive ingredients may be added to provide desirable colour, taste, stability, buffering capacity, dispersion, or other known desirable features. Examples include red iron oxide, silica gel, sodium laurel sulphate, titanium dioxide, edible white ink, and the like. Conventional diluents may be used to make compressed tablets. Both tablets and capsules may be manufactured as sustained-release compositions for the continual release of medication over a period of time. Compressed tablets may be in the form of sugar coated or film coated tablets, or enteric-coated tablets for selective disintegration in the gastrointestinal tract. Liquid dosage forms for oral administration may contain colouring and/or flavouring to increase patient compliance. As an example, the oral formulation comprising compounds of the invention may be a tablet comprising any one, or a combination of, the following excipients: calcium hydrogen phosphate dehydrate, microcrystalline cellulose, lactose, hydroxypropyl methyl cellulose, and talc.

[00192] The compositions described herein may be in the form of a liquid formulation. Examples of preferred liquid compositions include solutions, emulsions, injection solutions,

solutions contained in capsules. The liquid formulation may comprise a solution that includes a therapeutic agent dissolved in a solvent. Generally, any solvent that has the desired effect may be used in which the therapeutic agent dissolves and which can be administered to a subject. Generally, any concentration of therapeutic agent that has the desired effect can be used. The formulation in some variations is a solution which is unsaturated, a saturated or a supersaturated solution. The solvent may be a pure solvent or may be a mixture of liquid solvent components. In some variations the solution formed is an *in situ* gelling formulation. Solvents and types of solutions that may be used are well known to those versed in such drug delivery technologies.

[00193] The composition described herein may be in the form of a liquid suspension. The liquid suspensions may be prepared according to standard procedures known in the art. Examples of liquid suspensions include micro-emulsions, the formation of complexing compounds, and stabilising suspensions. The liquid suspension may be in undiluted or concentrated form. Liquid suspensions for oral use may contain suitable preservatives, antioxidants, and other excipients known in the art functioning as one or more of dispersion agents, suspending agents, thickening agents, emulsifying agents, wetting agents, solubilising agents, stabilising agents, flavouring and sweetening agents, colouring agents, and the like. The liquid suspension may contain glycerol and water.

[00194] The composition described herein may be in the form of an oral paste. The oral paste may be prepared according to standard procedures known in the art.

[00195] The composition is described herein may be in the form of a liquid formulation for injection, such as intra-muscular injection, and prepared using methods known in the art. For example, the liquid formulation may contain polyvinylpyrrolidone K30 and water.

[00196] The composition is described herein may be in the form of topical preparations. The topical preparation may be in the form of a lotion or a cream, prepared using methods known in the art. For example, a lotion may be formulated with an aqueous or oily base and may include one or more excipients known in the art, functioning as viscosity enhancers, emulsifying agents, fragrances or perfumes, preservative agents, chelating agents, pH modifiers, antioxidants, and the like. For example, the topical formulation comprising one or more compounds of the invention may be a gel comprising anyone, or a combination of, the following excipients: PEG 8000, PEG 4000, PEG 200, glycerol, propylene glycol. The NCL812 compound may further be formulated into a solid dispersion using SoluPlus (BASF, www.soluplus.com) and formulated with anyone, or a combination of, the following excipients: PEG 8000, PEG 4000, PEG 200, glycerol, and propylene glycol.

[00197] For aerosol administration, the composition of the invention is provided in a finely divided form together with a non-toxic surfactant and a propellant. The surfactant is preferably soluble in the propellant. Such surfactants may include esters or partial esters of fatty acids.

[00198] The compositions of the invention may alternatively be formulated for delivery by injection. As an example, the compound is delivered by injection by any one of the following routes: intravenous, intramuscular, intradermal, intraperitoneal, and subcutaneous.

[00199] The compositions of the invention may alternatively be formulated using nanotechnology drug delivery techniques such as those known in the art. Nanotechnology-based drug delivery systems have the advantage of improving bioavailability, patient compliance and reducing side effects.

[00200] The formulation of the composition of the invention includes the preparation of nanoparticles in the form of nanosuspensions or nanoemulsions, based on compound solubility. Nanosuspensions are dispersions of nanosized drug particles prepared by bottom-up or top-down technology and stabilised with suitable excipients. This approach may be applied to the compounds of the invention which can have poor aqueous and lipid solubility, in order to enhance saturation solubility and improve dissolution characteristics. An example of this technique is set out in Sharma and Garg (2010) (Pure drug and polymer-based nanotechnologies for the improved solubility, stability, bioavailability, and targeting of anti-HIV drugs. *Advanced Drug Delivery Reviews*, 62: p. 491-502). Saturation solubility will be understood to be a compound-specific constant that depends on temperature, properties of the dissolution medium, and particle size (<1–2 μm).

[00201] The composition of the invention may be provided in the form of a nanosuspension. For nanosuspensions, the increase in the surface area may lead to an increase in saturation solubility. Nanosuspensions are colloidal drug delivery systems, consisting of particles below 1 μm . Compositions of the invention may be in the form of nanosuspensions including nanocrystalline suspensions, solid lipid nanoparticles (SLNs), polymeric nanoparticles, nanocapsules, polymeric micelles and dendrimers. Nanosuspensions may be prepared using a top-down approach where larger particles may be reduced to nanometre dimensions by a variety of techniques known in the art including wet-milling and high-pressure homogenisation. Alternatively, nanosuspensions may be prepared using a bottom-up technique where controlled precipitation of particles may be carried out from solution.

[00202] The composition of the invention may be provided in the form of a nanoemulsion. Nanoemulsions are typically clear oil-in-water or water-in-oil biphasic systems, with a droplet size in the range of 100–500 nm, and with compounds of interest present in the hydrophobic phase. The preparation of nanoemulsions may improve the solubility of the compounds of the

invention described herein, leading to better bioavailability. Nanosized suspensions may include agents for electrostatic or steric stabilisation such as polymers and surfactants. Compositions in the form of SLNs may comprise biodegradable lipids such as triglycerides, steroids, waxes and emulsifiers such as soybean lecithin, egg lecithin, and poloxamers. The preparation of a SLN preparation may involve dissolving/dispersing drug in melted lipid followed by hot or cold homogenisation. If hot homogenisation is used, the melted lipidic phase may be dispersed in an aqueous phase and an emulsion prepared. This may be solidified by cooling to achieve SLNs. If cold homogenisation is used, the lipidic phase may be solidified in liquid nitrogen and ground to micron size. The resulting powder may be subjected to high-pressure homogenisation in an aqueous surfactant solution.

[00203] The Compounds of Formula I as described herein may be dissolved in oils/liquid lipids and stabilised into an emulsion formulation. Nanoemulsions may be prepared using high- and low-energy droplet reduction techniques. High-energy methods may include high-pressure homogenisation, ultrasonication and microfluidisation. If the low-energy method is used, solvent diffusion and phase inversion will generate a spontaneous nanoemulsion. Lipids used in nanoemulsions may be selected from the group comprising triglycerides, soybean oil, safflower oil, and sesame oil. Other components such as emulsifiers, antioxidants, pH modifiers and preservatives may also be added.

[00204] The composition may be in the form of a controlled-release formulation and may include a degradable or non-degradable polymer, hydrogel, organogel, or other physical construct that modifies the release of the compound. It is understood that such formulations may include additional inactive ingredients that are added to provide desirable colour, stability, buffering capacity, dispersion, or other known desirable features. Such formulations may further include liposomes, such as emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. Liposomes for use in the invention may be formed from standard vesicle-forming lipids, generally including neutral and negatively charged phospholipids and a sterol, such as cholesterol.

[00205] The formulations of the invention may have the advantage of increased solubility and/or stability of the compounds, particularly for those formulations prepared using nanotechnology techniques. Such increased stability and/or stability of the compounds of Formula I may improve bioavailability and enhance drug exposure for oral and/or parenteral dosage forms.

[00206] Throughout this specification, unless the context requires otherwise, the word "comprise" or variations such as "comprises" or "comprising", will be understood to imply the

inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.

EXAMPLES

EXAMPLE 1: The minimum inhibitory concentrations (MIC) for NCL812 in methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant *Enterococcus* spp. (VRE) and *Streptococcus pneumoniae*.

Specific

[00207] As is apparent from the preceding summary of the invention, the invention relates to compounds of Formula I, methods of treatment of a bacterial infection, uses and medical devices.

[00208] This study was undertaken to determine minimum inhibitory concentrations (MIC) for antibacterial agent NCL812. The antibacterial agent represents a potentially new class of drug with a perceived narrow spectrum of activity against bacteria and a novel mechanism of action. This study focused on recent isolates of three major opportunistic pathogens of humans where the development of antibacterial resistance to existing antibacterial classes is problematic: methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant *Enterococcus* spp. (VRE) and *Streptococcus pneumoniae*.

[00209] In this example, NCL812 minimum inhibitory concentrations (MICs) were determined for 61 Australian clinical isolates (comprised of 21 MRSA, 20 putative VRE and 20 *S. pneumoniae* isolates). The MIC profiles for NCL812 were found to be remarkably consistent, with MIC₅₀ and MIC₉₀ values of 4 µg/mL recorded for each of the species tested.

Materials and Methods

Bacterial Isolate Collection and Identification

[00210] Sixty one test isolates were sourced from clinical diagnostic microbiology laboratories. The MRSA isolates were originally cultured on selective Brilliance MRSA Chromogenic Agar (Oxoid). Suspect colonies were selected on the basis of their colony appearance on this agar and identification as *Staphylococcus aureus* was determined using colony characteristics on non-selective Sheep Blood Agar (SBA) and phenotypic characteristics such as Gram stain, positive catalase test, positive coagulase test (tube coagulase test using rabbit plasma) and clumping factor (agglutination with the Oxoid Staphylect latex test), positive Voges Proskauer test, and the ability to produce acid from trehalose. A positive cefoxitin resistance screen confirmed the isolates as MRSA. All *Enterococcus* isolates underwent a standard biochemical identification. Biochemical profiling provisionally identified four of the VRE

isolates as *Enterococcus faecalis* and the remainder as *Enterococcus faecium*. All *S. pneumoniae* isolates were identified on the basis of standard biochemical profiling.

Preparation of Antibacterials

[00211] Analytical grade NCL812 (batch 20081214) with a defined potency of 1000 mg/g (ie 100%) was obtained. The powder was stored at a temperature of -20 °C. Aliquots (1 mL) of stock solution (25.6 mg/mL) were prepared in DMSO and stored at -80 °C and defrosted immediately before use.

Preparation of 96 well microtitre plates for broth microdilution MIC test using NCL812

[00212] Cation adjusted Mueller Hinton Broth (CAMHB) was prepared using 100 mL sterile Mueller Hinton broth (pH adjusted). To each 100 mL volume, 125 µL of calcium stock solution (10 mg Ca²⁺ per mL) and 43 µL magnesium stock solution (10 mg Mg²⁺ per mL) was aseptically added. Sufficient broth is made up for daily use, with unused portions being stored at 4 °C overnight.

[00213] Microdilution trays with 4% lysed horse blood in CAMHB was prepared by lysing horse blood (Oxoid) by repeated freezing and thawing (3-4 times) and aseptically mixing the lysed horse blood (LHB) 50:50 with sterile distilled water. A cell free suspension was obtained by centrifuging 50% LHB at 16,000×g (7000 rpm) for 20 min. The supernatant was decanted, recentrifuged and stored frozen. 50% LHB was diluted with CAMHB to obtain a final concentration of 4% (7 mL LHB into 93 mL CAMHB). 4% LHB-CAMHB was used instead of CAMHB in all steps in the preparation of the microdilution trays and preparation of antimicrobial solutions for *Streptococcus* species.

[00214] A stock antibiotic working solution of NCL812 was prepared to a concentration of 25.60 mg/mL. Potency was described as 1000 mg/g or 100%. The powder was dissolved in 10 mL DMSO and 1 mL volumes were aliquoted into eppendorf tubes and stored at -80 °C. When added to CAMHB, a fine cloudy precipitate formed, and was shaken well before and during aliquoting.

[00215] A stock solution of ampicillin was prepared to the concentration of 25.60 mg/mL. Ampicillin was used for internal quality control. The powdered ampicillin was dissolved in 4 mL of phosphate buffer pH 8.0, 0.1 mol/L, then diluted in 6 mL phosphate buffer pH 6.0, 0.1 mol/L. 1 mL volumes were aliquoted into eppendorf tubes and stored at -80 °C.

[00216] For *Staphylococcus aureus*, a working solution of 256 µg/mL was prepared by diluting stock solutions as described above 1:100 in CAMHB (100 µL into 9.9 mL). When 90 µL

was added to each well 12, there was a 1:2 dilution so that well 12 had 128 µg/mL of antibiotic. The range of antimicrobial was calculated as 0.25 µg/mL (well 3) to 128 µg/mL (well 12).

[00217] For *Enterococcus* species, a working solution of 64 µg/mL was prepared by diluting stock solutions as described above 1:400 in CAMHB (100 µL into 9.9 mL, then further dilute this 1:4). When 90 µL was added to well 12 there was a 1:2 dilution so that well 12 had 32 µg/mL of antibiotic.

[00218] For *Streptococcus pneumoniae*, a working solution of 64 µg/mL was prepared by diluting stock solutions as described above 1:400 in 4% LHB-CAMHB (100 µL into 9.9 mL, then further diluting this 1:4). When 90 µL was added to well 12 there was a 1:2 dilution so that well 12 had 32 µg/mL of antibiotic.

[00219] Serial dilutions were prepared in 96 well plates were set up in a safety cabinet according to methods standard in the art. Briefly: 90 µL of the working antibiotic solution was added to each well in Column 12 of the plate, and mixed well, before 90 µL was transferred to column 11. The solutions were mixed again, and then transferred to the next column as before, continuing the dilutions through to column 3. Mixing the well requires the pick up and expulsion of 90 µL in each well 3-4 times before picking up and transferring the 90 µL to the next well. Column 2 (bacterial positive control) and column 1 (negative control) did not form part of the serial dilution. The trays were set up as follows: 2 strains were tested in duplicate in one tray, such that strain 1 was located in rows A to D, strain 2 was located in rows E to H, etc. The MIC (µg/mL) Interpretive Standard for Ampicillin using Control strains is shown in Table 1 below. *Staphylococcus aureus* ATCC 29213 Acceptable MIC range for Ampicillin = 0.5 to 2 µg/mL, *Enterococcus faecalis* ATCC 29212 Acceptable MIC range for Ampicillin = 0.5 to 2 µg/mL, *Streptococcus pneumoniae* ATCC 49619 Acceptable MIC range for Ampicillin = 0.06 to 0.25 µg/mL.

Table 1: MIC (µg/mL) Interpretive Standard for Ampicillin using Control strains according to Example 1.

	Sensitive	Intermediate	Resistant
<i>Staphylococcus aureus</i>	≤ 0.25		≥ 0.5
<i>Enterococcus species</i>	≤ 8		≥ 16
<i>Streptococcus pneumoniae</i>	≤ 2	4	≥ 8

Preparation of bacterial suspension for broth microdilution MIC method

[00220] Fresh cultures of bacteria were prepared for testing on sheep blood agar (SBA), and overnight incubation at 37 °C as follows; 2-3 colonies of each strain in 7 mL sterile saline, and the OD₆₀₀ measured as an indication of the density (approx. 0.5×10⁸ CFU/mL or 0.5

McFarland Standard). The bacterial suspension was adjusted to a final absorbance of 0.08 to 0.100, using saline to achieve correct density, and as the blank. Within 15 minutes of preparation, the adjusted bacterial suspension with sterile saline 1:20 (1 mL into 19 mL sterile saline) to achieve a final bacterial concentration of 4 to 5×10^5 CFU/mL. The bacterial solution was placed into a sterile trough and 10 μ L of bacterial solution added into wells 2 through to 12 on each required row (dilution of 1:10, with final concentration of bacteria in wells = 5×10^5 CFU/mL). The tray was sealed and incubated at 37 °C for 18-24 h. The purity of bacterial suspension was confirmed by streaking out 50 μ L of the 1:20 dilution onto a SBA plate, which was incubated for 37 °C for 18 h and examined. Viable counts were carried out to ensure that the correct concentration of bacteria had been added to the wells. The diluted bacterial solution (4 to 5×10^5 CFU/mL) was diluted 1:10 downwards by adding 100 μ L to 900 μ L of sterile saline in sterile tubes, and the serial dilutions continued 1:10 for 5 tubes. 100 μ L (4-5 drops) of the 4th and 5th dilutions (tube 4 = 105 and tube 5 = 106 CFU/ml) was plated around in duplicate on pre-dried PCA agar plates and incubated at 37°C overnight. The following day the number of colonies on the plates was counted and the average count in 100 μ L obtained. The study was multiplied by 10 to obtain a viable bacterial count per mL.

Description and identification of isolates

[00221] The MRSA isolates were originally cultured on selective Brilliance MRSA Chromogenic Agar (Oxoid). Suspect colonies were selected on the basis of their colony appearance on this agar and identification as *Staphylococcus aureus* was determined using colony characteristics on non-selective SBA and phenotypic characteristics such as Gram stain, positive catalase test, positive coagulase test (tube coagulase test using rabbit plasma) and clumping factor (agglutination with the Oxoid Staphylect latex test), positive Voges Proskauer test, and the ability to produce acid from trehalose. A positive cefoxitin resistance screen confirmed the isolates as MRSA.

[00222] MRSA clonal complexes were determined by rapid molecular typing. Two of the strains could not be typed using the rapid method, as shown in Table 2 below.

Table 2: a table showing the MRSA clonal complexes according to Example 1.

Organism / Sample no.	Clonal complex
<i>Staph. aureus</i> ATCC 29213	NA
MRSA 718	CC8
MRSA 741	CC8
MRSA 580	CC8
MRSA 622	CC22
MRSA 815	CC88

MRSA 844	CC22
MRSA 806	CC22
MRSA 786	CC88
MRSA 570	CC8
MRSA 773	TBD
MRSA 898	CC1
MRSA 787	CC8
MRSA 728	CC88
MRSA 713	CC8
MRSA 747	CC8
MRSA 616	CC5
MRSA 734	CC8
MRSA 516	TBD
MRSA 823	CC8
MRSA 778	CC8
MRSA 610	CC22

NA: Not Applicable; TBD: Isolates could not be typed using the rapid method and are currently being identified using traditional methodology.

[00223] All *Enterococcus* isolates underwent a simplified biochemical identification based on Quinn et al. (1994, Clinical Veterinary Microbiology, Mosby Ltd, New York). Biochemical profiling provisionally identified four of the VRE isolates as *Enterococcus faecalis* and the remainder as possibly *Enterococcus faecium*. All *S. pneumoniae* isolates were identified on the basis of standard biochemical profiling.

Test product and storage

[00224] Analytical grade NCL812 (batch 20081214) with a defined potency of 1000 mg/g (ie 100%) was obtained and the powder was stored at a temperature of -20 °C. Aliquots (1 mL) of stock solution (25.6 mg/mL) were prepared in DMSO and stored at -80 °C and defrosted immediately before use.

Minimum inhibitory concentration determination

[00225] Minimum inhibitory concentrations (µg/mL) were determined using the broth microdilution method recommended by the Clinical and Laboratory Standards Institute (CLSI) (Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard – Seventh Edition. CLSI M7-A7, 2006; Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals; Approved Standard – Second Edition. CLSI M31-A2, 2002; Performance Standards for Antimicrobial Susceptibility Testing; CLSI M2-A9, 2006).

[00226] The MIC was regarded as the the lowest concentration of an antimicrobial agent that completely inhibited growth of the organism in the microdilution wells as detected by the unaided eye. MIC breakpoints were determined by visual assessment and then confirmed using an ELISA plate reader, measuring absorbance levels at 450 nm. Bacterial growth (turbidity) in the wells with antimicrobial was compared with the amount of growth (turbidity) in the growth-control well (containing no antimicrobial). All isolates were tested in duplicate, if there was a difference of greater than one two-fold dilution in the results, the test was repeated a third time. The purity of the isolates was closely monitored during testing by subculturing the prepared bacterial inoculum onto SBA. Control organisms (*Enterococcus faecalis* strain ATCC 29212, *S. aureus* strain ATCC 29213 and *S. pneumoniae* strain ATCC 49619) were used throughout the testing to monitor quality control. The MICs of the control strains for the antimicrobial ampicillin (range 1.0, 2.0 and 0.06 µg/mL, respectively) were determined for each testing run as an internal quality control. The MIC₅₀, MIC₉₀ and MIC range (minimum and maximum) were calculated for each of the bacterial groups.

Results

[00227] Ampicillin MIC values obtained for the ATCC control strains were within the normal range expected on the basis of CLSI recommendations. The NCL812 and ampicillin MIC values for each isolate are indicated in Table 3 (MRSA isolates), Table 4 (VRE isolates) and Table 5 (*S. pneumoniae* isolates) below. MIC₅₀, MIC₉₀, MIC mode and MIC range for each of the species of bacteria tested are shown in Table 6 below. The MIC₅₀ is considered to be the lowest concentration which inhibits visible growth for 50% of the isolates. The MIC₉₀ is considered to be the lowest concentration which inhibits visible growth for 90% of the isolates. The MIC mode is the most commonly occurring MIC value and MIC range the minimum and maximum MIC values obtained.

Table 3: The Minimum Inhibitory Concentrations for the individual *Staphylococcus aureus* isolates according to Example 1.

MIC Testing - AMP/NCL812		Test 1	Test 2	Test 1	Test 2
Organism / Sample no.	AMP	AMP	NCL812	NCL812	
<i>Staph. aureus</i> ATCC 29213	2 µg/ml	2 µg/ml	4 µg/ml	4 µg/ml	
MRSA 748	>128 µg/ml	>128 µg/ml	4 µg/ml	8 µg/ml	
MRSA 741	8 µg/ml	8 µg/ml	4 µg/ml	4 µg/ml	
MRSA 580	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 822	64 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 815	64 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 844	128 µg/ml	128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 806	32 µg/ml	32 µg/ml	4 µg/ml	4 µg/ml	
MRSA 788	32 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 870	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 773	128 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 898	64 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 787	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 726	64 µg/ml	64 µg/ml	4 µg/ml	4 µg/ml	
MRSA 743	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 747	>128 µg/ml	>128 µg/ml	8 µg/ml	4 µg/ml	
MRSA 616	64 µg/ml	64 µg/ml	8 µg/ml	4 µg/ml	
MRSA 734	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 819	32 µg/ml	32 µg/ml	4 µg/ml	4 µg/ml	
MRSA 823	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 778	>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml	
MRSA 810	128 µg/ml	128 µg/ml	4 µg/ml	4 µg/ml	
AMP Ampicillin					
MIC90		>128 µg/ml	>128 µg/ml	4 µg/ml	4 µg/ml

Table 4: The Minimum Inhibitory Concentrations for the individual *Enterococcus* isolates according to Example 1.

MIC Testing - AMP/NCL812		Test 1	Test 2	Test 1	Test 2
Organism / Sample no.	AMP	AMP	NCL812	NCL812	
<i>Enterococcus</i> spp. ATCC 29212	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
VRE 001	0.5 µg/ml	0.5 µg/ml	2 µg/ml	2 µg/ml	
VRE 002	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
VRE 003	0.5 µg/ml	0.5 µg/ml	2 µg/ml	2 µg/ml	
VRE 004	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
VRE 005	1 µg/ml	1 µg/ml	2 µg/ml	2 µg/ml	
VRE 006	1 µg/ml	1 µg/ml	2 µg/ml	2 µg/ml	
VRE 007	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
<i>Enterococcus</i> 008	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
<i>Enterococcus</i> 009	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
<i>Enterococcus</i> 010	2 µg/ml	2 µg/ml	4 µg/ml	4 µg/ml	
<i>Enterococcus</i> 011	1 µg/ml	1 µg/ml	2 µg/ml	4 µg/ml	
<i>Enterococcus</i> 012	0.5 µg/ml	0.5 µg/ml	2 µg/ml	2 µg/ml	
<i>Enterococcus</i> 013	2 µg/ml	2 µg/ml	2 µg/ml	2 µg/ml	
<i>Enterococcus</i> 014	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
B09/ 2192	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
B09/ 2121	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
B09/ 2086	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
B09/ 2087	2 µg/ml	1 µg/ml	2 µg/ml	4 µg/ml	
B09/ 2032	2 µg/ml	2 µg/ml	4 µg/ml	4 µg/ml	
B09/ 2021	1 µg/ml	1 µg/ml	4 µg/ml	4 µg/ml	
AMP Ampicillin					
MIC90		2 µg/ml	2 µg/ml	4 µg/ml	4 µg/ml

Table 5: The Minimum Inhibitory Concentrations for the individual *Streptococcus pneumoniae* isolates according to Example 1.

MIC Testing - AMP/NCL812		Test 1	Test 2	Test 1	Test 2
Organism / Sample no.		AMP	AMP	NCL812	NCL812
<i>Strep. pneumoniae</i> ATCC 49619		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 1		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 2		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 3		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	2 µg/ml
<i>Strep. pneumoniae</i> 4		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 5		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 6		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 7		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 8		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 9		< 0.25 µg/ml	< 0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 10		< 0.06 µg/ml	< 0.06 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 12		< 0.06 µg/ml	< 0.06 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 13		< 0.06 µg/ml	< 0.06 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 14		< 0.06 µg/ml	< 0.06 µg/ml	8 µg/ml	8 µg/ml
<i>Strep. pneumoniae</i> 15		0.25 µg/ml	0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 16		2 µg/ml	2 µg/ml	0.5 µg/ml	1 µg/ml
<i>Strep. pneumoniae</i> 18		0.25 µg/ml	0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 19		0.25 µg/ml	0.25 µg/ml	4 µg/ml	4 µg/ml
<i>Strep. pneumoniae</i> 20		2 µg/ml	2 µg/ml	4 µg/ml	4 µg/ml
AMP Ampicillin					
MIC ₅₀		0.25 µg/ml	0.25 µg/ml	4 µg/ml	4 µg/ml

Table 6: The NCL812 MIC₅₀, MIC₉₀, MIC mode and MIC range for Australian isolates of MRSA, VRE and *S. pneumoniae*.

Bacteria	No. Isolates	MIC ₅₀ (µg/ml)	MIC ₉₀ (µg/ml)	MIC mode (µg/ml)	MIC range (µg/ml)
MRSA	21	4 (64)§	4 (>128)	4 (>128)	4 – 8 (8->128)
VRE	20	4 (1)	4 (2)	4 (1)	2 – 4 (0.5-2)
<i>E. faecium</i>	16	4 (1)	4 (2)	4 (1)	2-4 (1)
<i>E. faecalis</i>	4	NA	NA	NA	2-4 (1)
<i>Str. pneumoniae</i>	20	4 (<0.25)	4 (0.25)	4 (<0.25)	0.5 – 8 (<0.06-2)

* Mode – the most commonly occurring MIC value.

§ Comparative susceptibility MICs are shown in parenthesis.

[00228] NCL812 MIC values were consistent within and between each of the three species. MIC₅₀ and MIC₉₀ values were both equal (4 µg/ml) for MRSA, VRE and *S. pneumoniae* isolates, with less than 10% of isolates showing MIC values either 1-2 dilutions below or only one dilution above this figure.

[00229] On the basis of these results, NCL812 represents a new antibacterial.

EXAMPLE 2: Effect of NCL812 on *Staphylococcus aureus* Macromolecular Synthesis

Materials and Methods

Test compounds

[00230] Test compound NCL812 was transported to the experimental facility under conditions of ambient temperature and then stored at 2-8 °C until assayed. Stock solutions were made by dissolving NCL812 dry powder in 100% DMSO to a concentration of 6,400 µg/mL. Vancomycin (Cat. # 1134335), Rifampicin (Cat. # R-7382) and Cerulenin (Cat. # C-2389) were all obtained from Sigma, Ciprofloxacin was obtained from USP (Cat. # 1134335) and Linezolid was obtained from ChemPacific (Cat. # 35710).

Minimal Inhibitory Concentration Testing

[00231] The MIC assay method followed the procedure described by the Clinical and Laboratory Standards Institute, or CLSI (Clinical and Laboratory Standards Institute. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard—Eighth Edition*. CLSI document M07-A8 [ISBN 1-56238-689-1]. Clinical and Laboratory Standards Institute, 940 West Valley Road, Suite 1400, Wayne, Pennsylvania 19087-19898 USA, 2009), and employed automated liquid handlers to conduct serial dilutions and liquid transfers. The medium employed for the MIC assay was Mueller Hinton II Broth (MHB II- Becton Dickinson, Sparks, MD; Cat No 212322; Lot 9044411). *S. aureus* ATCC 29213 served as the quality control strain, and linezolid was utilized as the quality control antibiotic to validate the assay. NCL812 and linezolid were both dissolved in 100% DMSO before addition to the growth medium.

Macromolecular Synthesis Assays

Bacteria and growth conditions

[00232] The effect of NCL812 on whole cell DNA, RNA, cell wall, protein and lipid synthesis was investigated using *S. aureus* ATCC 29213. Cells were grown at 35 °C overnight on Trypticase Soy agar. A colony from the plate was used to inoculate 10 ml of Mueller Hinton broth II (MHBII), and the culture was grown to early exponential growth phase ($OD_{600} = 0.2$ to 0.3) while incubating in a shaker at 35 °C and 200 rpm.

DNA, RNA, and protein synthesis

[00233] When cells reached early exponential phase, 100 µL of culture was added to triplicate wells containing various concentrations of test compound or control antibiotics (5 µL) at 20× the final concentration in 100% DMSO. A 5% DMSO treated culture served as the "no drug" control for all experiments. Cells were added in MHBII at 105% to account for the volume of

drug added to each reaction or in M9 minimal medium for protein synthesis reactions. Following 15 min incubation at room temperature, either [^3H] thymidine (DNA synthesis), [^3H] uridine (RNA synthesis) or [^3H] leucine (protein synthesis) was added at 0.5-1.0 μCi per reaction, depending on the experiment. Reactions were allowed to proceed at room temperature for 15-30 min and then stopped by adding 12 μL of cold 5% trichloroacetic acid (TCA) or 5% TCA/2% casamino acids (protein synthesis only). Reactions were incubated on ice for 30 min and the TCA precipitated material was collected on a 25 mm GF/A filter. After washing three times with 5 ml of cold 5% TCA, the filters were rinsed two times with 5 mL 100% ethanol, allowed to dry, and then counted using a Beckman LS3801 liquid scintillation counter.

Cell wall synthesis

[00234] Bacterial cells in early exponential growth phase were transferred to M9 minimal medium and added to 1.5 mL eppendorf tubes (100 μL /tube) containing various concentrations of test compound or control antibiotics (5 μL) at 20 $\times$ the final concentration in 100% DMSO as described above. Following a 5 min incubation at 37 $^{\circ}\text{C}$, [^{14}C]N-acetylglucosamine (0.4 μCi /reaction) was added to each tube and incubated for 45 min in a 37 $^{\circ}\text{C}$ heating block. Reactions were stopped through the addition of 100 μL of 8% SDS to each tube. Reactions were then heated at 95 $^{\circ}\text{C}$ for 30 min in a heating block, cooled, briefly centrifuged, and spotted onto pre-wet HA filters (0.45 μM). After washing three times with 5 mL of 0.1% SDS, the filters were rinsed two times with 5 ml of deionized water, allowed to dry, and then counted using a Beckman LS3801 liquid scintillation counter.

Lipid synthesis

[00235] Bacterial cells were grown to early exponential growth phase in MHBII broth and added to 1.5 mL eppendorf tubes (in triplicate) containing various concentrations of test compound or control antibiotics as described above. Following a 5 min incubation at room temperature, [^3H] glycerol was added at 0.5 μCi per reaction.

[00236] Reactions were allowed to proceed at room temperature for 15 min and then stopped through the addition of 375 μL chloroform/methanol (1:2) followed by vortexing for 20 seconds after each addition. Chloroform (125 μL) was then added to each reaction, vortexed, followed by the addition of 125 μL dH_2O and vortexing. Reactions were centrifuged at 13,000 rpm for 10 min, and then 150 μL of the organic phase was transferred to a scintillation vial and allowed to dry in a fume hood for at least 1 hr. Samples were then counted via liquid scintillation counting.

Results

[00237] Susceptibility testing was conducted with NCL812 and *S. aureus* ATCC 29213 to determine the concentrations of drug needed in the macromolecular synthesis assays.

[00238] **Table 7** shows that the MIC for NCL812 was 4 µg/mL, while the quality control agent linezolid was within the CLSI-established quality control range (Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing; Nineteenth Informational Supplement*. CLSI document M100-S20 [ISBN 1-56238-716-2]. Clinical and Laboratory Standards Institute, 940 West Valley Road, Suite 1400, Wayne, Pennsylvania 19087-1898 USA, 2010). Precipitation of NCL812 was observed at ≥8 µg/mL in plates that were prepared in an identical fashion, but did not receive an inoculum of *S. aureus*. Macromolecular synthesis inhibition studies were performed using concentrations of NCL812 that were equivalent to 0, 0.25, 0.5, 1, 2, 4 or 8-fold the MIC value (4 µg/ml) for *S. aureus* ATCC 29213 (**Figures 10-15**).

Table 7: Minimum Inhibitory Concentrations values for NCL812 (robenidine) and linezolid against *Staphylococcus aureus* ATCC29213 according to Example 2.

Compound	<i>Staphylococcus aureus</i> ATCC ¹ 29213
NCL812	4
Linezolid	4
Quality control range for linezolid ²	1 - 4

[00239] **Figure 1** shows the effect of NCL812 on DNA synthesis. NCL812 demonstrated no inhibition at 0.25 fold the MIC, 40% inhibition at 0.5 fold, and approximately 95% inhibition at the MIC. This is compared to the control ciprofloxacin which showed approximately 51% at 8 fold the MIC (0.5 µg/mL). The results for NCL812 inhibition of RNA synthesis were very similar to the DNA synthesis study, with rifampicin serving as the positive control (**Figure 2**). It should be noted that precipitation was observed at 4 to 8 fold the MIC in the Mueller Hinton broth II utilized in the DNA and RNA synthesis assays.

[00240] Protein synthesis was inhibited in a dose-dependent manner at 0.25, 0.5, and 1 fold the MIC value of NCL812 showing up to 97% inhibition at the MIC (**Figure 3**). Linezolid demonstrated approximately 61% inhibition of protein synthesis at 8 fold the MIC (2 µg/mL). Precipitation of NCL812 occurred at 4 and 8 fold the MIC in the protein synthesis assay.

[00241] In **Figure 4** NCL812 also showed a somewhat dose-dependent inhibition of cell wall synthesis, though there was a large increase in inhibition from 1 to 2 fold the MIC.

However, inhibition dropped to approximately 68% and 52% at 4 fold and 8 fold the MIC, respectively. Precipitation of NCL812 occurred at 2, 4, and 8 fold the MIC in the M9 minimal medium used for the cell wall synthesis assay, and that is the likely cause of the decline in inhibition. In comparison, the positive control vancomycin showed 96% inhibition at 8 fold the MIC (2 µg/mL). NCL812 demonstrated a similar inhibition profile against lipid synthesis as that shown for DNA and RNA synthesis, reaching approximately 90% inhibition at the MIC (**Figure 5**). The positive control inhibitor cerulenin demonstrated 72% inhibition at 8 fold the MIC (32 µg/mL).

[00242] **Figure 6** represents a composite of all five macromolecular synthesis reactions. It can be observed that the inhibition curves were similar for each pathway, suggesting a global inhibition of several pathways simultaneously by NCL812. It is possible that NCL812 targets the cell membrane, causing leakage of essential ions and/or metabolites, thereby leading to a global shutdown of the cell synthesis pathways.

[00243] In summary, NCL812 inhibited DNA, RNA, protein, cell wall, and lipid pathways in a growing culture of *S. aureus*. Though some instances of dose-dependent inhibition of pathways was observed, all five macromolecular synthesis reactions were similarly sensitive to NCL812.

EXAMPLE 3: Effect of NCL812 on ATP Release from *Staphylococcus aureus*

Materials and Methods

Test compounds

[00244] The test compound NCL812 was shipped under conditions of ambient temperature and then stored at 2-8 °C until assayed. Stock solutions were made by dissolving NCL812 dry powder in 100% DMSO to a concentration of 1,600 µg/mL. Polymyxin B was obtained from Sigma (Cat. # P-4932).

Test Organism

[00245] *S. aureus* ATCC 29213 was originally acquired from the American Type Culture Collection (Manassas, VA).

ATP Release Assay

[00246] The CellTiter-Glo Luminescent Cell Viability Assay (Promega) was utilized to measure the leakage of ATP from bacteria. Cultures were grown to early exponential phase (0.2 – 0.3 OD₆₀₀) in Mueller-Hinton Broth II and then treated with seven different concentrations of either NCL812 or polymyxin B (positive control) utilizing the MIC for each compound as a guide

(0, 0.25, 0.5, 1, 2, 3, 4, or 8 fold the MIC). The negative control received 2% DMSO, which represented the final DMSO concentration in each assay. After a 30 min exposure to drug, cells were sedimented by centrifugation and the supernatant was analyzed for the presence of ATP. Results were expressed as ATP concentration released to the medium (μM).

Results

[00247] The MIC for NCL812 has been previously determined to be 4 $\mu\text{g/mL}$. The ATP release assay is conducted by growing *S. aureus* to exponential phase and then adding drug at multiples of the MIC in an effort to detect a dose-dependent response.

[00248] As shown in Figure 8 the positive control polymyxin B released ATP from *S. aureus* cells in a dose-dependent fashion with maximal release of approximately 0.34 μM ATP at 8 fold the MIC (256 $\mu\text{g/mL}$). ATP release in the presence of NCL812 was dose-dependent at 0.5–1 fold the MIC, resulting in maximal release (0.33 μM) observed at the MIC (4 $\mu\text{g/mL}$). ATP release actually decreased thereafter at 2 to 8 fold the MIC. It should be noted that in previous studies precipitation of NCL812 was observed at 4 to 8 fold the MIC in Mueller Hinton broth II.

[00249] In summary, NCL812 demonstrated dose-dependent release of ATP from actively growing *S. aureus* cells. ATP release from the cells into the growth medium reached maximum levels at the MIC value, and this was followed by a decrease in ATP release at higher doses. The data indicated that NCL812 may interact with the cell membrane of *S. aureus*, causing leakage of vital metabolites such as ATP.

EXAMPLE 4: *In vitro* antibacterial activity of NCL812 against methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*

Materials and Methods

Antimicrobial agents

[00250] Aliquots of stock solution of NCL812 (25.6 mg/ml) was prepared in DMSO, stored at $-80\text{ }^{\circ}\text{C}$ and defrosted immediately before use. Ampicillin stock was obtained from Sigma-Aldrich (Australia). Antimicrobial discs were obtained from Thermo Fisher Scientific (Australia).

Microorganisms

[00251] Clinical isolates of MRSA that represented the most common sequence types of both hospital-acquired (HA) and community-associated (CA) MRSA in Australia were obtained and are described in Table 8 below. The *S. aureus* control organism ATCC 49775 was used. Isolate identification was confirmed by conventional phenotypic methodologies, including the

slide coagulase test, Vogues-Proskauer test, polymyxin B sensitivity (300 units), and Staphytest Plus Protein A latex slide agglutination (Thermo Fisher Scientific Australia). Bacteria were stored at -80 °C in 40% glycerol broth and routinely grown from stock on sheep blood agar (SBA) incubated at 37 °C. In subsequent experiments, only fresh cultures <24 h were used.

Table 8: *Staphylococcus aureus* clone/isolate name, type, source, antibiogram, clindamycin resistance status, multi-locus sequence type (MLST), staphylococcal cassette chromosome (SCCmec) type, clonal complex, Panton-Valentine leukocidin status (PVL), and spa type for isolates used according to example 4

Name/Clone	Type	Source	Epi Number	Antibiogram	Clindamycin	MLST	SCC mec	Clonal Complex	PVL	spa
MSSA CN-55	Cat MSSA	M.B	-	-	-	-	-	-	-	-
MSSA DE-25	Dog MSSA	M.B	-	-	-	-	-	-	-	-
MSSA DO-27	Dog MSSA	M.B	-	-	-	-	-	-	-	-
MSSA COW-1	Cow MSSA	G	-	-	-	-	-	-	-	-
MSSA COW-2	Cow MSSA	J.P.	-	-	-	-	-	-	-	-
MSSA H-D	MSSA	VIMP	-	-	-	-	-	-	-	-
MSSA H-50	MSSA	VIMP	-	-	-	-	-	-	POS	-
MSSA ATCC 49775	MSSA	S.P.	-	-	-	-	-	-	-	-
MSSA H-46	MSSA	VIMP	-	-	-	-	-	-	-	-
MSSA H-48	MSSA	VIMP	-	-	-	-	-	-	-	-
MRSA H-79	HA-MRSA	VIMP	-	-	-	-	-	-	-	-
MRSA H-32	HA-MRSA	VIMP	-	-	-	-	-	-	-	-
UK EMRSA-15	HA-MRSA	G.C.	01-16337	Em, Cl	Inducible	ST22	IV	22	NEG	1022
UK EMRSA-15 PVL pos	HA-MRSA	G.C.	07-16386	Gn, Tm	-	ST22	IVb	22	POS	1891
UK EMRSA-16	HA-MRSA	G.C.	02-17800	Em, Cl	Constitutive resistance	ST36	II	30	NEG	1018
UK EMRSA-17	HA-MRSA	G.C.	03-15701	Gn, Em, Cl, Te, FA, Rf	Constitutive resistance	ST247	I	8	NEG	1051
Irish EMRSA-1	HA-MRSA	G.C.	04-16780	Gn, Em, Cl, Mp, Tm	Inducible	ST8	II	8	NEG	1498
CLASSIC MRSA	HA-MRSA	G.C.	03-17580	Em	Inducible	ST250	I	8	NEG	1008
NY/JAPAN	HA-MRSA	G.C.	03-16981	Em, Cl	Constitutive resistance	ST5	II	5	NEG	1242

Queensland clone (PVL pos)	CA-MRSA	G.C.	03-16790	Fully S	-	ST93	IVa	Singleton	POS	1202
Taiwan cMRSA	CA-MRSA	G.C.	03-16672	Em, Te	Constitutive resistance	ST59	5 (C2& 5)	59	POS	1437
WA MRSA-1	CA-MRSA	G.C.	WSG 8287	Em, FA	Inducible	ST1	IVa	1	NEG	1127
WA MRSA-2	CA-MRSA	G.C.	03-16928	Em	Inducible	ST78	IVa	88	NEG	1186
WA MRSA-3	CA-MRSA	G.C.	WSG 8376	Em	Inducible	ST5	IVa	5	NEG	1002
USA 300 MRSA	CA-MRSA	G.C.	04-15086	Te		ST8	IVc	8	POS	1008
Bengal Bay clone	CA-MRSA	G.C.	07-17048	Gn, Em, Cl, Tm	Non-inducible	ST172	V	1	POS	1387
WSPR MRSA (PVL neg)	CA-MRSA	G.C.	08-19231	Mp, Tm	-	ST30	IVa	30	NEG	15074
ST398-MRSA-V	CA-MRSA	G.C.	09-16670	Em, Te, Tm	Constitutive resistance	ST398	V	398	NEG	1034
WA MRSA-84	CA-MRSA	G.C.	07-16502	Cl	-	ST45	V	45	NEG	11081

MSSA; methicillin-susceptible *S. aureus*. HA-MRSA; hospital-acquired methicillin-resistant *S. aureus*. CA-MRSA; community-associated methicillin-resistant *S. aureus*. Em; Erythromycin. Cl; Ciprofloxacin. Gn; Gentamicin. Tm; Trimethoprim. Te; Tetracycline. FA; Fusidic Acid. Rf; Rifampicin. Mp; Mupirocin

Isolate resistotyping

[00252] Antibiotic-susceptibility profiling of the isolate collection was undertaken using Kirby-Bauer disc diffusion, as recommended by the Clinical and Laboratory Standards Institute (CLSI) on Mueller-Hinton agar. Isolates were grown overnight on SBA at 37 °C. Colonies were suspended in physiological saline. Turbidity was adjusted to a 0.5 McFarland standard and suspensions were spread over the medium. Antibiotic discs according to Table 9 below were transferred onto the inoculated medium and analysed after 24 h incubation at 37 °C. Isolates labelled as MRSA that were not β -lactam-resistant on the basis of the Kirby-Bauer test were grown from stock on plate count agar supplemented with 5 μ g/ml ampicillin and subject to repeat testing, as PBP2a expression can be induced by exposure to β -lactam antimicrobials.

Table 9: Antibacterial agent zone diameter interpretive sizes for Kirby-Bauer disc diffusion, as used in Example 4.

Antimicrobial Class	Agent	Disk content	Zone diameter interpretive sizes (mm)		
			Resistant	Intermediate	Susceptible
Aminoglycoside	Gentamicin	10 μ g	≤ 12	13-14	≥ 15
Ansamycin	Rifampicin	5 μ g	≤ 16	17-19	≥ 20
Cephalosporin	Cephalexin	30 μ g	≤ 14	15-17	≥ 18
Cepharmycin	Cefotetan	30 μ g	≤ 12	13-15	≥ 16
Folate pathway inhibitor	Trimethoprim-sulfamethoxazole	1.25/23.75 μ g	≤ 10	11-15	≥ 16
Glycopeptide	Vancomycin	30 μ g	-	-	≥ 15
Lincosamide	Clindamycin	2 μ g	≤ 14	15-20	≥ 21
Macrolide	Erythromycin	15 μ g	≤ 13	14-22	≥ 23
Fluoroquinolone	Ciprofloxacin	5 μ g	≤ 15	16-20	≥ 21
Tetracycline	Tetracycline	30 μ g	≤ 14	15-18	≥ 19
β -lactam/penicillin	Penicillin-G	10 units	≤ 28	-	≥ 29

β -lactam/penicillin-stable penicillin	Oxacillin	1 μ g	≤ 10	11-12	≥ 13
β -lactam/ β -lactamase inhibitor combination	Amoxicillin-clavulanic acid	20/10 μ g	≤ 19	-	≥ 20

Molecular detection of the protein A and mecA genes to confirm MRSA status

[00253] Isolate identities were confirmed genotypically using a duplex polymerase chain reaction (PCR) test targeting the *spa* (protein A) and *mecA* (methicillin resistance) genes. In addition, the isolates were tested in a *mecA* and *spa* Sybr green real-time PCR. Approximately ten colonies of each overnight bacterial subculture was suspended in 1× phosphate buffered saline (pH 7.4) and vortexed. Isolates were subject to DNA extraction using the QIAamp® DNA Mini Kit (Qiagen, Australia) following the manufacturers protocols. Template DNA was eluted in 50 μ L of elution buffer and either used directly in PCR, or stored at -20 °C prior to DNA amplification using the *spa* forward (5'-TGATACAGTAAATGACATTG-3') and reverse (5'-TTCTTATCAACAACAAGTTC-3') primers and *mecA* forward (5'-TTCGTGTCTTTTAATAAGTGAGG-3') and reverse (5'-ATGAAGTGGTAAATGGTAATATCG-3') primers (Invitrogen, Australia). Conventional PCR amplification was performed in a 20 μ L volume containing 10 μ L HotStarTaq Plus Master Mix (Qiagen, Australia), 0.5 μ M of each *spa* primer, 0.2 μ M of each *mecA* primer, and 3 μ L of extracted DNA. An automated thermal cycler (T100 Thermal Cycler, Bio-Rad) was used for PCR amplification of the *spa* and *mecA* genes according to the following conditions: PCR Stage (Enzyme activation at 95 °C for 300 s, followed by 38 amplification rounds of 94 °C for 30 s (denaturation), 50 °C for 30 s (annealing) and 72 °C for 38 s (extension) and then a cooling stage of 20 °C until required); Real-time PCR Stage (Enzyme activation at 95 °C for 300 s, followed by 40 amplification rounds of 95 °C for 15 s (denaturation), 50 °C for 20 s (annealing) and 70 °C for 40 s (extension), a single round at 95 °C for 5 s, a single round at 55 °C for 20 s, continuous melting curve from 95 °C to 0 °C and a cooling period of 40 °C for 30 s. The *mecA* and *spa* amplified products of 325 and 120 bp, respectively, were detected by GelRed staining followed by electrophoresis in 2% agarose gels.

Minimum inhibitory concentration testing

[00254] The *in vitro* activities of NCL812 and ampicillin as a positive control were determined by broth microdilution as recommended by the CLSI in cation-adjusted Mueller-Hinton II broth. Microtiter plates containing two-fold dilutions of each antimicrobial agent were inoculated with $\sim 10^6$ CFU/ml of each isolate in a 100 μ L final volume. Plates were incubated for

24 h at 37 °C. Turbidity (absorbance at OD₆₀₀) was measured using a Bio-Rad Benchmark Plus microplate spectrophotometer in Microplate Manager® version 5.2.1 (Bio-Rad). Minimum inhibitory concentration (MIC) endpoints were defined as the lowest antimicrobial concentration assessed by the spectrophotometer that inhibited bacterial growth. ATCC 49775 was included in the isolate collection as a control organism using breakpoints defined by the CLSI. The MIC₅₀, MIC₉₀ (concentrations that inhibited growth of the lower 50% and 90% of total organisms, respectively), and MIC range (minimum and maximum) were calculated to profile the antimicrobial susceptibility of the isolate collection.

Bactericidal activity

[00255] The bactericidal activity of NCL812 was established by determination of the minimum bactericidal concentration (MBC) and time-kill analyses using CLSI guidelines. The MBC was defined as the lowest drug concentration at which 99.95% of the original inoculum was eliminated.

[00256] Time-kill assays for ATCC 49775 were performed in cation-adjusted Mueller-Hinton II broth in Microtiter plates and again in 10 ml volumes for macrodilution assays at antimicrobial concentrations equivalent to 1× and 4× the MIC. Bactericidal activity in macrodilution assays was identified as a 3log₁₀ decrease from the initial inoculum size. Bacteria were cultured overnight at 37 °C on SBA. Colonies were suspended in broth and the turbidity was adjusted to a 0.5 McFarland standard to obtain a bacterial suspension of ~10⁸ CFU/ml. Bacterial suspensions were incubated at 37 °C with shaking. Aliquots were removed at 0, 1, 2, 4, 8, 12, and 24 h after antimicrobial addition, diluted, plated onto SBA and incubated for 48 h at 37 °C for viable count determination. Turbidimetric growth curves for *S. aureus* were obtained for Microtiter plate assays by monitoring optical density changes using a Bio-Rad Benchmark Plus microplate spectrophotometer at 600 nm. Optical densities were measured at 0, 1, 2, 4, 8, 12, and 24 h after antimicrobial addition.

Statistical methodology

[00257] Microbiological data was interpreted using CLSI guidelines. Data was examined using the student's t-test, Fisher's exact test, analysis of variance, and a generalized linear model for tests of between-subjects effects where appropriate. Differences were considered significant at the 0.05 level in IBM SPSS® version 19.0.

Results

Confirmation of Staphylococcus aureus identity and mecA status

[00258] Latex agglutination tests confirmed that all 30 isolates were protein A positive. The isolates tested positive for coagulase activity using slide agglutination. Voges-Proskauer and polymyxin B resistance tests confirmed that all isolates were *S. aureus* except for a single methicillin-susceptible isolate; MSSA DE-25, as shown in Table 10 below. Based on *spa* gene PCR amplification, this isolate was not identified as a *S. aureus* isolate despite testing positive in the protein A latex agglutination and slide coagulase tests. This canine-origin *Staphylococcus* spp. was identified as *Staphylococcus pseudintermedius* based on biochemical characteristics. *mecA* conventional and real-time PCR results confirmed that 66.66% of the isolates were classified as methicillin-resistant on the basis of possession of the *mecA* gene. There were no significant differences between the ability of conventional and real-time PCR to detect the *mecA* gene ($P>0.05$).

Table 10: Percentage of presumptively identified *S. aureus* isolates reporting positive to selected phenotypic and genotypic tests according to Example 4.

Organism	Biochemical tests				PCR		Real-time PCR	
	Protein A	Slide coagulase	Voges-Proskauer	Polymyxin B	spa gene	mecA gene	spa gene	mecA gene
Methicillin-susceptible staphylococci (n=10)	100%	100%	90%	90%	90%	0%	90%	0%
HA-MRSA (n=10)	100%	100%	100%	100%	100%	100%	100%	100%
CA-MRSA (n=10)	100%	100%	100%	100%	100%	90%	100%	100%
Total (n=30)	100%	100%	96.67%	96.67%	96.67%	63.33%	96.67%	66.66%

HA-MRSA; hospital-acquired *S. aureus*. CA-MRSA; community-associated *S. aureus*. *S. aureus* isolates were identified as testing positive to protein A latex agglutination (Protein A), slide coagulase, Voges-Proskauer and polymyxin B resistance tests, as well as testing positive for polymerase chain reaction (PCR) and real-time PCR amplification of the spa gene. Methicillin-resistant *S. aureus* isolates were identified as isolates testing positive to the criteria described above, as well as positive for PCR and real-time PCR of the mecA gene.

Table 11: Resistance of *S. aureus* isolates to antibacterial agents using the Kirby-Bauer disc diffusion method according to example 4

Antimicrobial agent(s)	HA-MRSA (n=10)	CA-MRSA (n=10)	Methicillin-susceptible staphylococci (n=10)	Total isolates resistant (n=30)
Penicillin-G	100.00%	100.00%	50.00%	83.33%
Erythromycin	80.00%	100.00%	40.00%	73.33%
Amoxicillin-Clavulanic acid	50.00%	30.00%	0.00%	36.67%
Cefotetan	80.00%	20.00%	0.00%	33.33%
Cephalexin	80.00%	20.00%	10.00%	40.00%
Oxacillin	80.00%	10.00%	0.00%	30.00%
Ciprofloxacin	60.00%	20.00%	20.00%	33.33%
Clindamycin	50.00%	50.00%	30.00%	43.33%
Gentamicin	40.00%	20.00%	0.00%	20.00%
Tetracycline	30.00%	30.00%	0.00%	20.00%
Rifampicin	10.00%	0.00%	0.00%	3.33%
Trimethoprim-	10.00%	0.00%	0.00%	3.33%
Sulfamethoxazole				
Vancomycin	0.00%	0.00%	0.00%	0.00%

HA-MRSA; hospital-acquired methicillin-resistant *S. aureus*. CA-MRSA; community-associated methicillin-resistant *S. aureus*

Staphylococcus aureus antimicrobial susceptibility profiles

[00259] Antimicrobial susceptibility assays revealed that HA-MRSA isolates had the highest mean prevalence of resistance to multiple antimicrobial classes ($P < 0.000$). CA-MRSA isolates were next most resistant ($P < 0.007$), followed by methicillin-susceptible staphylococci ($P < 0.037$), as shown in Table 11 above. Oxacillin resistance was expressed in only 80.00% and 10.00% of HA-MRSA and CA-MRSA isolates, respectively. Cefotetan resistance was expressed in 80.00% and 20.00% of HA-MRSA and CA-MRSA isolates, respectively. Although oxacillin and cefotetan did not significantly differ in their ability to detect MRSA ($P > 0.05$), detection was significantly improved when using the *mecA* PCR when compared to disc diffusion ($P < 0.013$). The majority of HA-MRSA isolates expressed resistance to amoxicillin-clavulanic acid, cefotetan, cephalexin, clindamycin, erythromycin, oxacillin, and penicillin-G, whereas the majority of CA-MRSA isolates were resistant to only clindamycin, erythromycin, and penicillin-G. None of the isolates tested were vancomycin resistant. Overall, the most prevalent resistance phenotypes were penicillin-G (83.33%), erythromycin (73.33%), and clindamycin (43.33%), whilst only single isolates (3.33%) were resistant to trimethoprim-sulfamethoxazole and rifampicin.

mec gene complex interactions

[00260] All MRSA isolates belonging to *mec* gene complex A expressed resistance to both oxacillin and cefotetan, as shown in Table 12 below. However, only 20% of *mec* gene complex B MRSA isolates were phenotypically resistant to these antimicrobials. Of the MRSA isolates belonging to *mec* gene complex C2, only a single isolate expressed methicillin resistance to oxacillin and only two isolates expressed resistance to cefotetan. Unclassified MRSA isolates expressed full resistance to oxacillin and cefotetan.

Table 12: Number and percentage of identified *mec* gene complexes in 20 *S. aureus* strains classified as methicillin-resistant according to example 4

<i>mec</i> gene complex (number of isolates)	SCC <i>mec</i> types	Oxacillin resistant isolates	Cefotetan resistant isolates
A (n=4)	II	3 (100%)	3 (100%)
	III	1 (100%)	1 (100%)
	Overall	4 (100%)	4 (100%)
B (n=10)	I	1 (50%)	1 (50%)
	IV	1 (12.5%)	1 (12.5%)

	Overall	2 (20%)	2 (20%)
C2 (n=4)	V	1 (25%)	2 (50%)
	Overall	1 (25%)	2 (50%)
Unclassified (n=2)	Overall	2 (100%)	2 (100%)

Respective staphylococcal cassette chromosome (*SCCmec*) complexes and types expressing phenotypic resistance to oxacillin and cefotetan are indicated as well as real-time *mecA* status, and the average negative dF/dT peak obtained from melting point analysis from real-time PCR of the *mecA* gene

[00261] Melting point peaks for the *mecA* real-time PCR negative derivative plot $-dF/dT$ differed between *mec* gene complex ($P<0.003$) (**Figure 10**). On average, *mec* gene complex B and unclassified isolates demonstrated higher melting point peaks than other *SCCmec* types ($P<0.012$).

Physical properties of test antimicrobials and comparison of minimum inhibitory concentration results from initial analogue testing

[00262] Test antimicrobials were selected on the basis of solubility and antimicrobial activity from preliminary studies. Cloudy precipitates were observed when both NCL812 were dissolved in cation-adjusted Mueller-Hinton II broth, as shown in Table 13 below. Following initial structure-activity testing on each synthesized analogue, NCL812 was found to have consistent MIC values in this present study.

Table 13: Characteristics of antibacterial NCL812 and the β -lactam antibacterial ampicillin according to Example 4.

Compound	Solubility		Original MIC ($\mu\text{g/ml}$) at 24-h ¹		Observed MIC ($\mu\text{g/ml}$) at 24-h		
	DMSO	CAMHB	MRSA5 80	MRSA698	MRSA5 80	MRSA6 98	ATCC 49775
NCL812	Good	Cloudy	4	4	4	4	4
Ampicillin	Good	Good	>128	16	>128	16	0.25

Detailing antibacterial solubility in dimethyl sulfoxide (DMSO), solubility in cation-adjusted Mueller-Hinton II broth (CAMHB), and average minimum inhibitory concentrations (MIC) ($\mu\text{g/ml}$ at 24 h) against methicillin-resistant *S. aureus* (MRSA) determined from preliminary studies and those determined during this present study. ATCC 49775; methicillin-susceptible *S. aureus*

isolate and ATCC control strain. MRSA580; methicillin-resistant *S. aureus* isolate #580. MRSA898; methicillin-resistant *S. aureus* isolate #898

In vitro antibacterial activities: minimum inhibitory concentrations

[00263] MIC₅₀ and MIC₉₀ values for lead compound NCL812 (4- and 4-8 µg/mL) are shown in Table 14 below. MIC values differed by *S. aureus* classification (susceptible, HA- or CA-MRSA) ($P < 0.005$). In many cases, NCL812 had significantly increased activity against CA-MRSA and methicillin-susceptible staphylococci by one dilution when compared to HA-MRSA ($P < 0.002$ and $P < 0.020$, respectively), however there were no significant differences between MIC values for methicillin-susceptible staphylococci and CA-MRSA ($P > 0.05$). Ampicillin MIC values obtained for the ATCC control strain were within the normal range expected on the basis of CLSI guidelines

Table 14: *In vitro* activities of the novel antibacterial NCL812 and the β-lactam antibacterial ampicillin against *S. aureus* clinical isolates according to Example 4.

Organism and antimicrobial agent (no. of isolates tested)	MIC (µg/ml)			MBC (µg/ml)		
	MIC _{range}	MIC ₅₀	MIC ₉₀	MBC _{range}	MBC ₅₀	MBC ₉₀
Methicillin-susceptible staphylococci (n=10)						
NCL812	2-8	4	4	2-8	4	8
Ampicillin [†]	0.25-16	0.25	8	-	-	-
HA-MRSA (n=10)						
NCL812	4-8	4	8	4-8	8	8
Ampicillin [†]	16->128	64	>128	-	-	-
CA-MRSA (n=10)						
NCL812	2-4	4	4	2-4	4	4
Ampicillin [†]	8-128	64	128	-	-	-
Total bacterial isolates (n=30)						
NCL812	2-8	4	8	2-8	4	8
Ampicillin [†]	0.25->128	16	128	-	-	-

HA-MRSA; hospital-acquired methicillin-resistant *S. aureus*. CA-MRSA; community-associated methicillin-resistant *S. aureus*. MIC; minimum inhibitory concentration (µg/ml). MBC; minimum bactericidal concentration (µg/ml). MIC/MBC_{range}; minimum and maximum MIC/MBC for all isolates. MIC/MBC₅₀; MIC/MBC at which 50% of isolates are inhibited. MIC/MBC₉₀; MIC/MBC at which 90% of isolates are inhibited

In vitro antibacterial activities: minimum bactericidal concentrations

[00264] The MBCs determined from NCL812 were equivalent to the MIC for 93.33% and 83.33% of *S. aureus* isolates, respectively (**Table 14**). In all remaining cases, MBCs were one dilution higher. For NCL812, MBCs ranged from 2-8 µg/mL and 4-16 µg/mL, respectively.

Time-kill studies

[00265] In comparison to the turbidimetric growth curve of ATCC 49775, no visible bacterial growth was observed when ATCC 49775 was inoculated into cation-adjusted Mueller Hinton II broth supplemented with NCL812 at 1× and 4× the MIC in microdilution assays ($P < 0.033$ and $P < 0.038$, respectively) (**Figure 9**).

[00266] When analysed in 10 mL macrodilution assays, broth supplemented with antimicrobials at 1× and 4× the MIC and inoculated with ATCC 49775 displayed significantly reduced viable counts for both NCL812 concentrations when compared to the growth control ($0.000 < P < 0.008$) (**Figure 10**). Additionally, the time-kill profiles of each concentration of NCL812 did not significantly differ ($P > 0.05$). Both concentrations remained bactericidal until approximately 8-12 h after antimicrobial addition, where bacterial regrowth was observed. Considerable variation in the killing activity of NCL812 was observed from 8-24 h. Although NCL812 was no longer bactericidal by 24 h, viable counts observed at 1× the MIC remained significantly lower than those obtained from unsupplemented broth ($P < 0.046$).

[00267] In summary, the example set out above demonstrates bactericidal activity against both methicillin-susceptible staphylococci and MRSA. MIC and MBC values were consistently low across the selection of isolates (MIC_{range} 2-8 µg/mL). NCL812 retained good *in vitro* antimicrobial activity against common, multidrug-resistant MRSA isolates, including the epidemic UK EMRSA-15, EMRSA-16, and EMRSA-17, Irish EMRSA-1, AUS EMRSA-3, NY/JAPAN HA-MRSA, and predominant CA-MRSA clones. NCL812 was also active against one *S. pseudintermedius* isolate that was originally identified as a *S. aureus* strain.

[00268] Preliminary testing suggests that NCL812 targets the *S. aureus* cell membrane, causing dose-dependent release of vital metabolites such as ATP. Disruption of the bacterial membrane bilayer or proteins that are integral to membrane function in bacteria is a target for numerous large antimicrobials which are ubiquitous in nature; including glycolipids, lipopeptides, lipoproteins, fatty acids, neutral lipids, phospholipids, and biosurfactants. Although NCL812 is a low molecular mass (≤ 500 Da) synthetic compound, it does appear to exert bactericidal activity in a similar manner to other antimicrobials which target the Gram-positive cell membrane, including the high molecular weight cyclic lipodepsipeptide antimicrobial agent daptomycin, or

the low molecular mass quinolone-derived HT61, whose chemical structure is not currently available. Many of these lipophilic antibacterial agents are also not effective against Gram-negative microorganisms due to the presence of the outer lipid bilayer membrane, which contains narrow porin channels reducing the net penetration of some compounds into the cell.

[00269] The insolubility of NCL812 at even low concentrations in microbiological media may reflect the amphipathic and oligomeric nature of this antimicrobial and suggests that the real MIC may be much lower than observed, as it is likely that it is only NCL812 in solution that is biologically active. In time-kill studies, NCL812 exerted rapid *in vitro* bactericidal activity against ATCC 49775. Again, these findings are consistent with a time-kill profile of cell membrane function inhibitors such as daptomycin and HT61.

[00270] Importantly, the apparent short *in vitro* half-life of this antimicrobial resulted in bacterial regrowth observed at 12 h after antimicrobial addition. This suggests that if a viable bacterial population survives the initial exposure to NCL812 prior to antimicrobial inactivation, bacterial regrowth will occur. The development of resistance to NCL812 in these studies was ruled out as test bacteria remained susceptible to NCL812 following harvesting, washing and MIC testing. Whilst the apparent short *in vitro* half-life of NCL812 may be a desirable characteristic for future *in vivo* application, it does suggest that NCL812 should be administered every 8 h in future *in vivo* safety and efficacy experiments to maintain adequate systemic concentrations, though it would appear from the time-kill profile that the NCL compound series are concentration-dependent rather than time-dependent antimicrobials.

[00271] To overcome the methicillin-susceptible phenotype, extending disc diffusion incubation time from 24 to 48 h compensates for the slow derepression of the *mecR* gene. Although the effects of longer incubation were not examined, and the small sample size of MRSA isolates prevented further investigation into *mec* complex interactions; genetic techniques were of significantly improved sensitivity when compared to phenotypic methods for confirmation of the *mecA* status of the isolates in this study. Although genetic techniques are not always employed as a routine method for detecting MRSA, real-time PCR identification of the presence of the *mecA* gene in a *Staphylococcus* spp. isolate remains the diagnostic gold standard.

EXAMPLE 5: *In vitro* pharmacodynamics of a new antimicrobial agent for *Streptococcus pneumoniae*.

Materials and Methods

Pneumococcal antimicrobial susceptibility

Pneumococcal strains and growth conditions

[00272] Twenty pneumococcal isolates that comprised six characterised laboratory strains and 14 clinical isolates were the subject of this study (P9/6A, P21/3, WCH16/6A, WCH43/4, WCH46/4, WCH57/8, WCH77/5, WCH86/4, WCH89/7, WCH92/4, WCH137/6A, WCH158/19F, WCH184/19F and WCH211/11; strain/serotype, respectively). Other isolates used in this example were: A66.1/3 (Francis *et al.*, 2001. *Infect Immun.* 69: 3350-2358); EF3030/19F (Briles *et al.*, 2003 *J. Infect. Diseases.* 188:339-348); L82016/6B (Briles *et al.*, 2000 *Infect Immun.* 68:796-800); TIGR4/4 (Tettelin *et al.*, 2001 *Science* 293:498-506); and WU2/3 (Briles *et al.*, 1981 *J. Exp. Med.* 153:694-705). See Table 15 below for the phenotypic characteristics of the isolates used in this study. The National Collection of Type Cultures (NCTC) control strain D39 (Avery *et al.*, 2010 *Nature Reviews Microbiology* 8:260-271) was used as a growth control for all MIC and MBC assays. D39 was later designated for kill kinetics, point of resistance assays and transmission electron microscopy (TEM) studies as it is a well documented laboratory strain with a defined *in vivo* pathogenesis (Table 15) that displayed consistent NCL812 MICs and MBCs.

Table 15: Pneumococcal isolates and their phenotypic description according to Example 5.

Strain	Phenotypic description	ID ₅₀
D39 (NCTC 7486)	Flat, round, 1mm wide, dark green, α-haemolysis	10 ²
A66.1	Slightly raised, undulate, irregular in shape, 2mm wide colonies, glossy, dark green, mucoid, α-haemolysis	8 x 10 ² (10, 98)
EF3030	Slightly raised, undulate, irregular in shape, 2mm wide colonies, glossy, dark green, mucoid, α-haemolysis	≥ 10 ⁵
L82016	Flat, round, 1mm wide, pale green, α-haemolysis	≥ 10 ²
P9	Flat, circular, 0.5mm wide colonies, glossy, light green, α-haemolysis	10 ⁴
P21	Slightly raised, undulate, irregular in shape, 4mm wide colonies, glossy, dark green, mucoid, α-haemolysis	≤ 10 ¹
TIGR4	Flat, round, 1mm wide, pale green, α-haemolysis	10 ³
WU2	Slightly raised, undulate, irregular in shape, 4mm wide colonies, glossy, dark green, mucoid, α-haemolysis	5x10 ¹²
WCH16	Flat, round, 1mm wide, pale green, α-haemolysis	5 x 10 ⁴
WCH43	Flat, round, 1mm wide, pale green, α-haemolysis	10 ²
WCH46	Flat, round, 1mm wide, pale green, α-haemolysis	10 ³
WCH57	Slightly raised, undulate, irregular in shape, 2mm wide colonies, glossy, dark green, mucoid, α-haemolysis	10 ³
WCH77	Slightly raised, round in shape, <1mm wide colonies, glossy, dark green,	10 ⁴

	slight α -haemolysis.	
WCH86	Flat, round, 1mm wide, pale green, α -haemolysis	10^4
WCH89	Flat, round, > 1mm wide, pale green, α -haemolysis	$\geq 10^5$
WCH92	Flat, round, 1mm wide, dark green, α -haemolysis	$\leq 10^8$
WCH137	Slightly raised, round in shape, <1mm wide colonies, glossy, dark green, slight α -haemolysis.	ND
WCH158	Round, sunken, 1mm wide, dark green, α -haemolysis	10^5
WCH184	Flat, round, 1mm wide, dark green, α -haemolysis	10^8 (14)
WCH211	Flat, round, > 1mm wide, dark green, α -haemolysis, irregular shape	5×10^5

ND = Not determined

[00273] For all *in vitro* assays, fresh pneumococcal isolates were grown overnight (O/N) on horse blood agar (HBA) plates (39 g/L Columbia blood agar base [Oxoid] 5% [v/v] defibrinated horse blood [Oxoid] at 37 °C with 5% supplemented CO₂). Mueller-Hinton blood agar with 5% defibrinated sheep blood (MHSBA Roseworthy Media and Blood Service) was used for disk diffusion analysis as directed by Clinical Laboratory Standards Institute (CLSI) standards. Pneumococci were routinely grown in broth consisting of 4% lysed horse blood (LHB) with Cation Adjusted Mueller Hinton Broth (CAHMB, [Difco]) at 37 °C with 5% supplemented CO₂. Horse serum broth (HSB, 10% (v/v) donor horse serum in nutrient broth [10 g/L peptone, 10 g/L Lab Lemco (Oxoid) and 5 g/L NaCl]) was also used in some MIC assays. Isolates were stored in HSB at -80 °C.

Antibiotic stocks and reagents

[00274] NCL812 was provided in dry powder form. A total of 256 mg was dispensed into 10 mL of 100% DMSO to make a stock of 25.6 mg/mL, which was then diluted 1:100 in CAHMB to make a final working stock of 256 μ g/mL. Ampicillin dry powder was from Sigma A0166. The original 25.6 mg/mL stock was diluted in saline 1:100, 1:4, 1:20 and finally 1:16 in CAMHB to make a final working stock of 0.18 μ g/mL. Erythromycin was purchased from Sigma Aldrich and choline chloride was from Roche Diagnostics. Twenty micro litres of 0.05 μ g/mL erythromycin was diluted 1:25 in 4.980 mL of CAMHB to give a final working stock of 0.2 μ g/mL. Choline chloride (0.5%) was added to 4% LHB:CAMHB for specific kill kinetic assays.

Defining antimicrobial susceptibility of pneumococcal isolates

[00275] Isolate susceptibility to 12 different antimicrobials (Table 16) was determined by CLSI and European Committee on Antimicrobial Susceptibility Testing (EUCAST) methods. Antimicrobials were selected based upon the CLSI and EUCAST guidelines. Standardised bacterial suspensions were spread onto MHSBA using a sterile cotton swab. Bacterial suspensions from of *Streptococcus pneumoniae* were standardised to an OD₆₀₀ between 0.08 and 0.1 using a spectrophotometer and then diluted 1:20. Bacterial colonies were taken from an O/N horse blood agar plate. To ensure the purity of the 1:20 bacterial suspension, 50 μ L was

spread plated onto horse blood agar and incubated O/N at 37 °C with 5% CO₂. The CFU was calculated and compared to the initial plate counts. Antibiotic disks (Purchased from Sigma Aldrich) were placed using a disk dispenser (Purchased from Oxoid) according to CLSI standards. MHSBA plates were incubated for 16 h – 24 h at 37 °C in 5% CO₂. Zones of complete inhibition were measured in triplicate to the nearest millimetre using a ruler on natural light-reflected growth, and the mode was represented as the diameter for each isolate. Pneumococcal isolates were categorised as sensitive, intermediate (I) or resistant (R) by CLSI standards and quality control (QC) ranges (Table 16).

Table 16: Antibacterials used for disc diffusion analysis with interpretive standards of zone diameters (mm) according to Example 5.

Antibiotic Class	Antimicrobial (µg)	Interpretive Standards for Zone Diameters (mm) (98)		
		Resistant (R)	Intermediate (I)	Sensitive
<i>β-lactam</i>	Oxacillin (1 µg) [*]	≤ 20	≤ 20	≥ 20
	Ampicillin (10 µg) [*]	≤ 20	≤ 20	≥ 20
	Amoxicillin-clavulanate (20/10 µg) [*]	≤ 20	≤ 20	≥ 20
<i>Fluoroquinolone</i>	Ciprofloxacin (5 µg) [*]	≤ 22	≤ 22	≥ 22
<i>Folate pathway inhibitor</i>	Trimethoprim-sulphamethoxazole (1.25/23.75 µg) [*]	≤ 15	16-18	≥ 19
<i>Glycopeptide</i>	Vancomycin (30 µg) [*]	-	-	≤ 17
<i>Lincosamide</i>	Clindamycin (2 µg) [*]	≤ 15	16-18	≥ 19
<i>Macrolide</i>	Erythromycin (15 µg) [*]	≤ 15	16-20	≥ 21
	Clarithromycin (15 µg) [*]	≤ 16	17-20	≥ 21
<i>Phenocoi</i>	Chloramphenicol (30 µg) [*]	≤ 20	-	≥ 21
<i>Rifamycin</i>	Rifampin (5 µg) [*]	≤ 16	17-18	≥ 19
<i>Tetracycline</i>	Tetracycline (30 µg) [*]	≤ 18	19-22	≥ 23

[00276] ^{*} Zone diameters for antimicrobials other than Ciprofloxacin for *S. pneumoniae* were determined by CLSI standards.

^{*}Zone diameters for Ciprofloxacin antimicrobial susceptibility to *S. pneumoniae* were determined by EUCAST.

Determination of NCL812 MIC₅₀, MIC₉₀, MIC range and MBC₅₀, MBC₉₀, MBC range

[00277] MICs for NCL812 for all isolates listed in Table 15 were determined by measuring OD₆₀₀ (Spectramax spectrophotometer, Molecular Devices Corporation) as an indicator of bacterial growth using 96 well microtitre trays after incubation for 24 h at 37 °C in 5% CO₂. [Micro-broth dilutions and 96 well trays are prepared by the following method: 90 µL of 4% LHB:CAMHB is aliquotted into all wells using a multichannel pipette. 90 µL of working antimicrobial stocks were serially diluted down the tray by a 1:2 dilution. Negative broth controls and dilution control were taken into account when planning the set up of a 96 well tray.]

10 μ L of bacterial suspension was then added to the appropriate wells in the 96 well tray. Appropriate positive (no antimicrobial), negative (no antimicrobial or bacteria) and negative dilution (a serial dilution control of antimicrobial and broth) controls were included in each assay. MBC and plate counts for kill kinetic assays were determined by aliquotting 20 μ L from each well of the 96 well microtitre tray onto HBA, and incubating at 37 °C with 5% CO₂. The MBC was determined by a 99.95% inhibition of *S. pneumoniae*, taking into account the dilution factor. MICs and MBCs were determined in quadruplicate and the mode was taken as the representative value. The MIC₅₀, MIC₉₀ and MIC range and MBC₅₀, MBC₉₀ and MBC range were determined according to CLSI standards. The MIC₅₀ and MIC₉₀, or MBC₅₀ and MBC₉₀, are defined by the lowest concentrations which, when all the MICs and MBCs of the isolates are arranged from lowest to highest, inhibited the 50th and 90th percentile of the total amount of isolates, respectively.

Micro-broth dilution time kill studies with NCL812 using strain D39

[00278] Bacterial suspensions were added in triplicate to a 96 well microtitre tray containing NCL812 with a starting concentration of 128 μ g/mL and serially diluted 1:2 sequentially to a concentration of 0.25 μ g/mL. Negative dilution controls were subtracted from the median growth value to obtain a suitable indicator of overall bacterial production. The 96 well tray was incubated at 37 °C in 5% CO₂ and OD₆₀₀ read every 2 h for the first 12 h followed by final reads at 24 and 48 h. To further supplement this data, a separate experiment in which a 96 well tray was read automatically at half hourly intervals using a spectrophotometer (Spectramax spectrophotometer, Molecular Devices Corporation) for 14 h was performed to confirm the trends in growth curves observed from original micro-broth dilution studies.

MBC time kill studies with NCL812 using strain D39

[00279] MBC kill kinetics assays involved the preparation of three 96 well microtitre trays. At specific time points, aliquots obtained from these trays provided viable counts following incubation at 37 °C in 5% CO₂ on HBA, and the MBC was determined after 24 h of growth.

Macro-broth dilution time kill studies of D39 with NCL812

[00280] Bacterial suspensions and working antibiotic stocks were prepared as described above. [For preparing macro-broth dilutions, 20 mL tubes were filled each with 9 mL of 4% LHB:CAMHB. 9 mL of a working antimicrobial stock was diluted 1:2 when added to one of the tubes, and then serially diluted down from a high to low concentration of antimicrobial. 1 mL of *S. pneumoniae* bacterial suspension was added to the appropriate tubes, including the positive control. Tubes were incubated at 37 °C with 5% CO₂ with gentle manual tilting of the tubes

treated with NCL812 every 10 min for the first 12 h. At every 2-3 h during the first 12 h of growth and then at 24 h and 48 h, 50 μ L of each bacterial suspension was spread plated onto HBA and incubated at 37 °C with 5% CO₂ for 16-24 h.]

[00281] Table 17 below indicates the concentrations used for each antimicrobial. Cultures were incubated at 37 °C in 5% CO₂ with gentle manual tilting every 10 min for the first 12 h. Viable counts from 50 μ L aliquots of each concentration were read following incubation at 37 °C in 5% CO₂ for 24 h. The pH of each sample was measured at specific time points using pH indicator strips. Confluent growth was defined when more than 1000 colonies were counted per plate. A bactericidal effect was defined as a 1000 fold reduction (99.9%) of the original cell suspension determined at 24 h for each concentration.

Table 17: Antibacterial agent concentrations used in macro-broth dilution assays according to Example 5.

Serial dilution	NCL812 (μ g/ml)	NCL062 (μ g/ml)	Ampicillin (μ g/ml)
1	128	128	0.09
2	64	64	0.045
3	32	32	0.023
4	16	16	0.011
5	8	8	0.0065
6	4		
7	2		

Point of resistance assay for NCL812

[00282] Macro-broth dilutions were prepared as above. Broth cultures of strain D39 (10 mL) were incubated in the presence of 2 μ g/mL and 4 μ g/mL of NCL812, and 0.022 μ g/mL of Ampicillin for 6 h at 37 °C in 5% CO₂. Samples were centrifuged at a relative centrifugal force (RCF) of 101.45 $\times$ g for 10 min and washed in 50 mL of phosphate buffered saline (PBS) twice to remove any residual antimicrobial, and/or bacterial end products and media. Washed bacteria were resuspended and MICs were performed.

Effect of NCL812 on D39 cell membrane ultra-structure

Transmission Electron Microscopy

[00283] Morphological appearance and morphometric analysis of the cell membrane was determined using transmission electron microscopy (TEM). Bacterial suspensions and 10 mL cultures of D39 were prepared as before. Samples were incubated at 37 °C in 5% CO₂ with gentle manual tilting of the cultures every 10 min. Cultures were exposed to either 1 μ g/mL, 4 μ g/mL or 16 μ g/mL of NCL812 and harvested at 6 or 12 h by centrifugation at RCF of 101.45 $\times$ g

for 20 min and washed twice in 50 mL of PBS. Critical time points for TEM work were determined by analysing trends in the growth curves produced from the kill kinetics studies. Samples were resuspended in PBS containing 20% glycerol and stored at -80 °C until required. Before fixation, 20% glycerol was removed by centrifugation and washing on ice three times in 50 mL of PBS.

[00284] Samples were fixed using modified protocols defined by a previous study examining cell wall ultrastructure of *S. pneumoniae* (Hammerschmidt, S. *et al.* 2005, *Infect Immun* 73:4653-4667). A lysine acetate-based formaldehyde-glutaraldehyde ruthenium red-osmium fixation procedure involved fixing the bacterial pellets with a cacodylate buffer solution containing 2% formaldehyde, 2.5% glutaraldehyde, 0.075% ruthenium red and 0.075 M of lysine acetate for 1 h. After washing with cacodylate buffer containing 0.075% ruthenium red three times, a second fixation in cacodylate buffer solution containing 2% formaldehyde, 2.5% glutaraldehyde and 0.075% ruthenium red was undertaken for 1.5 h. Cells were subsequently washed three times with cacodylate buffer containing 0.075% ruthenium red and underwent a final fixation in 1% osmium tetroxide in cacodylate containing 0.075% ruthenium red for 1 h. The samples were then washed three times in cacodylate buffer containing 0.075% ruthenium red only.

[00285] Samples were washed and dehydrated using a graded series of ethanol (70, 90, 95 and 100%) for 10-20 min, two times for each step. Samples were infiltrated using 50:50 LR White resin in 100% ethanol for 1 h, and subsequently washed with 100% LR White resin for 1 h and left O/N in a third change of 100% LR white to ensure adequate infiltration of resin. The samples were then embedded in fresh LR White resin and incubated at 50 °C for 48 h. Sections were cut to 1 µm using a glass knife, stained with Toluidene Blue and viewed under a light microcope at 400× to identify the presence of stained pneumococci. At least four ultra-thin sections were then cut to 90 nm using a diamond knife and placed on matrix grids, one section per grid. Ultra-thin sections were then stained with uranyl acetate and lead citrate alternatively at 5 min intervals, followed by three washes with distilled water in-between each exposure. Stained sections were then placed on grids and viewed between 25000× and 130000× on a Philips CM100 Transmission Electron Microscope. Images were obtained at 130000× magnification and analysed using analySIS [Olympus Soft Imaging Systems].

Statistical analysis

[00286] Statistical analyses were conducted using statistics program GraphPad Prism (5th ed, GraphPad Software Inc.) for Windows. For growth curves, data presented were the mean and standard error of mean (SEM) (represented as error bars) for each data point except

for macro-broth dilution studies where multiple replicates could not be obtained due to the high costs involved in this assay. Two tailed, unpaired *t*-tests were performed.

Results

Pharmacodynamics of NCL812 in S. pneumoniae

Quality control disk diffusion analysis for 20 S. pneumoniae isolates

[00287] Although nine out of the twelve antimicrobials used for disk diffusion analysis had established QC ranges by EUCAST, QC ranges were not defined for amoxicillin-clavulanate, clarithromycin and clindamycin (Table 18 and Table 19). WCH16 and WCH184 were both resistant to at least two antimicrobials whereas EF3030 and WCH137 were intermediate and resistant to trimethoprim-sulphamethoxazole respectively (Table 19). The other remaining sixteen isolates were sensitive to all twelve antimicrobials. Sensitivity to ampicillin was confirmed for each isolate, enabling the use of ampicillin as a positive control in later micro-broth dilution assays (Table 18).

Table 18: Antibacterial susceptibility of 20 *S. pneumoniae* isolates for six different antibacterials according to Example 5.

Antimicrobial	Range of disk diffusion zone diameters (nearest whole mm)					
	Ampicillin ≥ 20mm	Amoxicillin- clavulanate ≥ 20mm	Clarithromycin ≥ 21mm	Clindamycin ≥ 19mm	Chloramphenicol ≥ 21mm	Ciprofloxacin ≥ 22mm
Sensitivity standard	25-31mm	Not defined	Not defined	Not defined	24-30mm	22-28mm
QC range	37	43	30	34	34	25
A66.1	37	47	32	27	31	26
D39	38	43	32	26	29	22
EF3030	41	39	34	32	26	26
L62016	33	42	40	31	31	27
P9	40	47	36	25	35	23
P21	37	42	35	28	32	24
TIGR4	38	42	31	27	30	28
WU2	41	46	33	28	11 (R)	25
WCH16	36	42	28	22	26	21
WCH43	37	41	38	30	28	27
WCH46	43	53	38	34	40	32
WCH57	43	48	37	28	36	26
WCH77	38	35	36	30	34	36
WCH86	38	40	31	27	32	26
WCH89	38	40	27	24	32	21
WCH92	40	43	33	26	31	24
WCH137	42	46	38	32	34	28
WCH158	33	38	10 (R)	10 (R)	29	22
WCH184	39	42	33	27	32	24

Blue cells represent sensitive isolates; orange cells represent intermediate (I) isolates; green cells represent resistant (R) isolates.

Table 19: Antibacterial susceptibility of 20 *S. pneumoniae* isolates for six different antibacterials according to example 5

Antimicrobial	Range of disk diffusion zone diameters (nearest whole mm)				
	Erythromycin	Oxacillin	Rifampin (Rifampicin)	Tetracycline	Trimethoprim- sulphamethoxazole
Vancomycin					

Sensitivity standard	≥ 21mm	≥ 20mm	≥ 23 mm	≥ 23 mm	ole	≥ 17mm
QC range	26-32mm	6-14mm	26-32mm	28-34mm	20-26mm	17-23mm
A66.1	33	28	32	32	23	25
D39	35	26	28	35	25	22
EF3030	30	18	32	34	17 (I)	22
L82016	29	29	26	32	23	23
P9	32	26	27	32	21	21
P21	35	28	32	38	25	21
TIGR4	34	28	30	35	24	22
WU2	34	28	31	34	22	21
WCH16	34	30	30	16 (R)	21	23
WCH43	29	26	27	31	22	20
WCH46	27	27	29	30	24	21
WCH57	41	32	38	48	32	28
WCH77	37	26	33	39	29	24
WCH86	30	34	31	31	27	22
WCH89	35	28	32	35	25	25
WCH92	28	25	22	29	23	22
WCH137	34	29	30	37	10 (R)	24
WCH158	34	28	34	35	27	22
WCH184	10 (R)	11	32	11 (R)	22	23
WCH211	33	25	32	36-37	34	23

Blue cells represent sensitive isolates; orange cells represent intermediate (I) isolates; green cells represent resistant (R) isolates

Solubility and activity of NCL812 and NCL062 in different media

[00288] NCL812 visually appeared to have higher solubility in 100% DMSO compared to NCL062 and only developed turbidity when it was further diluted into CAMHB or PBS (Table 20). Although a CAMHB diluent for NCL062 appeared to be transparent by visual inspection (Table 20), further studies on NCL062 with a CAMHB diluent resulted in complete confluence in microbroth dilution assays for six *S. pneumoniae* isolates in comparison to growth with the DMSO diluents (Table 21 and Table 22).

Table 20: Visual analysis of NCL812 and NCL062 and ampicillin solubility according to Example 5.

Diluent	NCL812	NCL062	Ampicillin
CAMHB	Turbid	Transparent	Transparent
DMSO	Transparent	Precipitate	Transparent
PBS	Precipitate	Precipitate	Transparent
Media			
4% LHB: CAMHB	Turbid	Transparent	Transparent
10% horse serum-supplemented broth	Precipitate	Transparent	Transparent

Table 21: Individual MICs of NCL062 for each pneumococcal isolate according to Example 5

NCL062	MIC ($\mu\text{g.mL}^{-1}$)	MBC ($\mu\text{g.mL}^{-1}$)
D39	2	4
EF3030	16	16
A66.1	8	32
TIGR4	8	8
WU2	16	64
L82016	8	32
P9	4	16
P21	4	4
WCH158	16	16
WCH89	4	4
WCH57	4	8
WCH77	8	8
WCH46	32	32
WCH86	4	8
WCH137	2	4
WCH184	16	16
WCH16	32	32
WCH43	2	4
WCH92	8	8
WCH211	2	4

Table 22: Difference in activity of NCL812 and NCL062 in different media using micro-broth dilution to obtain an MIC as a predictor according to Example 5.

	NCL062	Ampicillin
--	--------	------------

Isolate	Resistance status to Ampicillin	MIC ($\mu\text{g/ml}$)	MBC ($\mu\text{g/ml}$)	MIC ($\mu\text{g/ml}$)	MBC ($\mu\text{g/ml}$)
D39	Sensitive	Confluent	*	0.0225	*
A66.1	Sensitive	Confluent	*	0.045	*
WU-2	Sensitive	Confluent	*	0.0225	*
P21	Sensitive	Confluent	*	0.0225	0.045
WCH158	Sensitive	Confluent	64	0.0225	0.022
WCH57	Sensitive	Confluent	*	0.09	0.09

[00289] Growth of *S. pneumoniae* strain D39 in an MIC assay for NCL812 and NCL062 using 10% HSB (220 mL of horse serum is filtered to 10% in 180 mL of Lemco nutrient broth) resulted in a threefold increase in the MIC for D39 treated with NCL812 and NCL062 (Table 23) with a twofold increase for the positive ampicillin control. There was no notable change in MIC for D39 with differing storage conditions of pre-prepared 96 well microtitre trays (Table 24). During macro-broth dilutions, the pH of the media did not change compared to appropriate controls (Figure 11).

Table 23: Growth of *S. pneumoniae* strain D39 in an MIC assay for NCL812 and NCL062 using horse serum supplemented broth

Antimicrobial	Relative MIC with media type for D39 ($\mu\text{g.mL}^{-1}$)		
	4%LHB:CAMHB	10% horse serum-supplemented broth	Fold-increase
NCL812	4	32	3
NCL062	4	32	3
Ampicillin	0.023	0.09	2

[00290] Table 24: Storage of prepared micro-titer trays for micro-broth dilution does not change MIC of D39 according to Example 5.

Antimicrobial	Storage condition	
	-2°C	4°C
NCL812	8 $\mu\text{g.mL}^{-1}$	8 $\mu\text{g.mL}^{-1}$
NCL062	4 $\mu\text{g.mL}^{-1}$	4 $\mu\text{g.mL}^{-1}$
Ampicillin	0.023 $\mu\text{g.mL}^{-1}$	0.023 $\mu\text{g.mL}^{-1}$

Determination of S. pneumoniae in vitro susceptibility to NCL812 and NCL062

Determination of NCL812 and NCL062 MIC₅₀, MIC₉₀, MIC range

[00291] NCL812 exhibited a MIC₅₀ and MIC₉₀ of 8 $\mu\text{g/mL}$ and MIC range of 4-8 $\mu\text{g/mL}$ whereas for NCL062 these values were higher and more variable (Table 25 and Table 26). The MIC for ampicillin was comparable to recent published findings using micro-broth

dilution as an endpoint for antimicrobial resistance in pneumococcal isolates, thus confirming the accuracy of MICs obtained for NCL812 and NCL062 (**Tables 25 to 26 and Figure 12**).

Table 25: MIC₅₀, MIC₉₀, MBC₅₀, MBC₉₀ and MIC range for all isolates treated with NCL812, NCL062, and ampicillin according to Example 5.

	NCL812 (µg/mL)	NCL062 (µg/mL)	Ampicillin (µg/mL)
MIC₅₀	8	8	0.023
MIC₉₀	8	32	0.023
MIC Range	4-8	2-32	0.011 - 0.09
MBC₅₀	8	16	0.023
MBC₉₀	8	32	0.023
MBC Range	4-8	4-64	0.011-0.09

Table 26: MICs of NCL812 of each pneumococcal isolate according to Example 5.

	NCL812		Ampicillin	
	MIC (µg.mL ⁻¹)	MBC (µg.mL ⁻¹)	MIC (µg.mL ⁻¹)	MBC (µg.mL ⁻¹)
D39	4	8	0.023	0.023
EF3030	8	8	0.023	0.023
A66.1	8	8	0.045	0.045
TIGR4	4	8	0.023	0.023
WU2	4	8	0.023	0.023
L82016	8	8	0.023	0.023
P9	8	8	0.023	0.023
P21	4	8	0.023	0.023
WCH158	4	8	0.023	0.023
WCH89	4	4	0.023	0.023
WCH57	8	8	0.023	0.023
WCH77	4	8	0.023	0.023
WCH46	4	4	0.023	0.045
WCH86	4	8	0.023	0.023
WCH137	4	8	0.023	0.023
WCH184	4	4	0.045	0.045
WCH16	8	autolysis	0.023	Autolysis
WCH43	4	8	0.023	0.023
WCH92	8	8	0.09	0.09
WCH211	4	8	0.023	0.023

Determination of NCL812 and NCL062 MBC₅₀, MBC₉₀, MBC range

[00292] Minimum bactericidal concentrations (MBC₅₀, MBC₉₀ and MBC range respectively) were determined for NCL812 and ampicillin for all twenty isolates (**Tables 25 to Table 26**). The MBC₅₀, MBC₉₀, and MBC range was lower and more consistent for NCL812 compared with NCL062 (**Table 25**).

Micro-broth dilution time kill studies of D39 treated with NCL812 and NCL062

[00293] D39 exposed to sub-inhibitory concentrations ($\leq 2 \mu\text{g/ml}$) of NCL812 or NCL062 grew similarly to unexposed controls over a 48 h period (**Figure 12 and 13**). Higher concentrations of NCL812 and NCL062 ($\geq 16 \mu\text{g/mL}$) resulted in no bacterial growth for 48 h (**Figure 13 and 14**). These growth characteristics were validated by a micro-broth kill kinetic study using a Spectramax spectrophotometer, which measured growth (represented as OD_{600}) at half-hourly intervals for 14 h for NCL812, NCL062 and ampicillin (**Figures 14 to 16**). There was an approximate six hour difference between the commencement of exponential growth for D39 treated with NCL812 and D39 treated with NCL062 (**Figures 12, 13, 17 and 18**).

[00294] The growth of D39 treated with NCL812 or NCL062 was compared to D39 treated with ampicillin or erythromycin over 48 h (**Figures 19 and 20**). D39 treated with ampicillin exhibited similar growth to D39 exposed to NCL812 or NCL062 over 48 h (**Figure 19**). Erythromycin-treated D39 produced very different growth curves from NCL812 and NCL062 where a larger difference in growth between concentrations was observed (**Figure 20**). The addition of 5% choline chloride to the media over a 48 h period resulted in no significant difference in growth for NCL812 and NCL062 compared to positive and growth controls (**Figures 21 to 25**).

Point of resistance testing

[00295] D39 treated with $\leq 4 \mu\text{g/mL}$ NCL812 entered a log phase of growth at 6 h (**Figure 12 and 17**), as shown in four independent experiments. The possibility of antimicrobial resistance to NCL812 between 5 and 6 h was investigated by determining further MICs on D39 exposed to $2 \mu\text{g/mL}$ NCL812, $4 \mu\text{g/mL}$ NCL812 and $0.0225 \mu\text{g/mL}$ ampicillin for 6 h. Results showed no significant increase in MIC for all samples of D39 exposed to NCL812 compared to growth controls, and ampicillin (**Table 27**).

Table 27: MICs of D39 exposed to $2 \mu\text{g/mL}$ or $4 \mu\text{g/mL}$ of NCL812 for 6 h according to Example 5.

	Original MIC of D39	MIC of D39 following exposure to NCL-812 for 6 hrs.	Original MBC of D39	MBC of D39 following exposure to NCL-812 for 6 hrs
D39 exposed to $2 \mu\text{g/ml}$ NCL812	$4 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$
D39 exposed to $4 \mu\text{g/ml}$ NCL812	$4 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$
D39 exposed $0.023 \mu\text{g/ml}$ Ampicillin	$0.023 \mu\text{g.mL}^{-1}$	$0.045 \mu\text{g.mL}^{-1}$	$0.023 \mu\text{g.mL}^{-1}$	$0.023 \mu\text{g.mL}^{-1}$
D39 growth *	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$	$8 \mu\text{g.mL}^{-1}$

D39 growth2 **	8 µg.mL ⁻¹	8 µg.mL ⁻¹	8 µg.mL ⁻¹	8 µg.mL ⁻¹
-----------------------	-----------------------	-----------------------	-----------------------	-----------------------

*D39 growth control: *S. pneumoniae* strain D39 grown for 6 hrs in 4%LHB:CAMHB.

** D39 growth2 control: *S. pneumoniae* strain D39 on HBA O/N, resuspended in saline (0.1 OD₆₀₀) and diluted 1/20 in sterile saline.

Micro-broth dilutions by measuring relative MBC at specific time points

[00296] Relative MBCs were determined at specific time intervals from using broth dilution assays incubated for 48 h for NCL812 and NCL062 (**Figure 26**) and control antimicrobials ampicillin and erythromycin (**Figures 19 and 20**). MICs of ampicillin and erythromycin for D39 were determined (**Tables 26 and 28**). The comparative features of the growth of ampicillin, and erythromycin are described (**Figures 27 and 28**). Ampicillin and erythromycin demonstrated a time-dependent reduction in bacteria. NCL062 exhibited rapid bactericidal action, with an immediate (within the first 10 min of administration) MBC of 8 µg/mL (**Figure 26**). Although there were inconsistencies in the MBCs for NCL062 between 5 and 12 h, NCL062 maintained a constant bactericidal concentration (4 µg/mL) between 24 and 48 h. NCL812 exhibited fast bactericidal action, evidenced by an approximate 3 fold decrease in MBC within 5 h (**Figure 26**). A consistent bactericidal concentration (8 µg/mL) was maintained for the full 48 h for NCL812.

[00297] Table 28: MIC and MBC for erythromycin with D39 according to Example 5.

Isolate	Resistance status to Erythromycin	MIC (µg/ml)	MBC (µg/ml)
D39	Sensitive	0.00275	0.00275

Macro-broth dilution time kill studies of D39 with NCL812 and NCL062

[00298] Viable counts for each time point were represented as a log₁₀ CFU/mL reduction for NCL812 (**Figure 29**) and ampicillin (**Figure 30**). Consistent confluent growth (determined by a limit of 2×10⁴ CFU) was observed for unexposed controls and 2 µg/mL NCL812. Complete bactericidal activity (defined by a 3log₁₀ reduction in CFU) for 128 µg/mL of NCL812 was observed by a 4log₁₀ reduction of CFU in 3 h and concentrations between 16 µg/mL and 64 µg/mL NCL812 were effective at eliminating bacterial growth within 8 h (**Figure 29**). NCL812 at 4 µg/mL and 8 µg/mL appeared to be inactivated at 11 h post-exposure, as increased growth of strain D39 after this time point was observed (**Figure 19 and 30**). The viable counts of strain D39 treated with ampicillin demonstrated consistency for this particular assay by showing a constant diminished time-dependant killing over 48 h (**Figure 30**).

Transmission electron microscopy

[00299] Morphometric analysis revealed significant changes to the cell membrane in strain D39 exposed to 16 µg/mL NCL812 for 6 h compared to growth controls. Samples treated with 4 µg/mL as well as 12 h cultures were not considered for morphometric analysis due to the lack of bacterial cells available in each section. Treated samples possessed significantly thicker cell membranes (6.43 ± 0.29 nm) compared to untreated samples (4.35 ± 0.24 nm) ($p < 0.0001$) (**Figures 31 and 28**). The periplasmic space (intracellular space between the cell membrane and the cell wall) of D39 treated with 16 µg/mL NCL812 was significantly wider (4.54 ± 0.096 nm) compared to untreated samples (3.91 ± 0.14 nm) ($p < 0.001$) (**Figures 28 and 32**).

[00300] Table 29: Morphometric studies on the ultra structures of D39 treated with NCL812 for 6 hours according to Example 5.

	Growth control	Treatment (16µg/ml NCL812 for 6 h)	
Statistical test	Mean ± SEM	Mean ± SEM	Unpaired t-test (P value)
Cell membrane	4.35 ± 0.24 nm, n = 12	6.43 ± 0.29 nm, n = 13	$P < 0.0001$
Periplasmic space	3.91 ± 0.14 nm, n = 11	4.54 ± 0.096 nm, n = 11	$P < 0.001$

[00301] In summary, NCL812 produced highly consistent MICs and equivalent MBCs for the *S. pneumoniae* strain collection, confirming that it is bactericidal against this organism. In kill kinetics experiments, which measured the relative MBC over a 48 h period, a consistent bactericidal effect was elicited in D39 after 6 h from initial exposure to NCL812.

[00302] This demonstration of bactericidal activity is the first to be observed in *S. pneumoniae*. This demonstrates that NCL812 is effective against pneumococcal in vitro.

[00303] Competitive binding between components in blood, serum or broth decreased the antimicrobial activity of NCL. This was reflected in the increase of MIC observed between different broth types and diluents. Following the completion of these studies, recent independent research confirmed precipitation of NCL812 in PBS and reported complete solubility in water containing 4% DMSO, following initial dilution in 100% DMSO. A water-soluble NCL812 will greatly improve *in vivo* bioavailability and negative interaction between blood or serum proteins.

[00304] Based on the findings of this study, NCL812 exhibits a mechanism of action against *S. pneumoniae* that is different from β-lactam or macrolide classes, as it appears to exhibit concentration-dependent bactericidal activity as opposed to time-dependant qualities. Identifying the maximum pharmacokinetic serum concentration of NCL812 in vivo will assist

confirmation of its concentration-dependant pharmacodynamic activity. Furthermore, the addition of choline chloride to the media confirmed that the mechanism of action for NCL is not associated with the affinity to cell wall choline binding proteins, and therefore may not be cell wall associated.

[00305] Morphometric analysis of the cell membrane and periplasmic space of D39 treated with 16 µg/mL NCL812 for 6 h showed that the cell membrane and periplasmic space was larger in treated samples, compared to control samples. The apparent increase in membrane size could be due to an accumulation of electron dense intracellular material beneath the cell membrane. The increase in the size of the periplasmic space may be have been due to disruption of the cell membrane, potentially by depolarisation or ATP inhibition. The mechanism of action of NCL812 may not be calcium-dependant as it appears that no competitive binding between NCL812 and ruthenium red, a calcium channel inhibitor of lipid bilayers, was observed in electron micrographs.

[00306] In conclusion, this *in vitro* study has demonstrated that NCL812 has many desirable characteristics as a fast-acting concentration-dependent bactericidal antimicrobial that appears to target the cell membrane of *S. pneumoniae*. These characteristics are desirable to treat acute pneumococcal infections. As NCL812 may possess a mechanism of action that targets the cell membrane, it will act much more quickly than time-dependent antimicrobials such as β-lactams and macrolides and potentially more effective than other bactericidal concentration-dependent antimicrobials such as fluoroquinolones which have intracellular targets.

EXAMPLE 6: Characterization of methicillin-susceptible and methicillin-resistant isolates of *Staphylococcus pseudintermedius* from Australia and preliminary *in vitro* efficacy of a new anti-staphylococcal compound

Materials and Methods

Sample collection and identification of methicillin susceptible Staphylococcus pseudintermedius (MSSP) and methicillin resistant Staphylococcus pseudintermedius (MRSP)

[00307] A total of 23 *Staphylococcus pseudintermedius* isolates were obtained from dogs (Table 30).

Table 30: *Staphylococcus pseudintermedius* isolates tested according to Example 6.

Adelaide #	GLY	BOX	SITE OF ISOLATION	Species	BREED	MRSP / MSSP	mec gene Part	mecA by RT-PCR Adl	Cefoxitin ZD (mm)	Cefoxitin ZD, Adl (mm)	oxacillin ZD (mm)	oxacillin ZD, Adl (mm)	oxacillin Etest MIC (mg/L)
S1P1	191	8	AXILLA	<i>S. pseudintermedius</i>		MRSP	POS	POS	0	0	0	0	>256
S2P2	193	8	TISSUE	<i>S. pseudintermedius</i>	Shar pei X	MRSP	POS	POS	21	21	0	0	>256
S3P3	194	8	SKIN SWAB	<i>S. pseudintermedius</i>	Mastiff X	MRSP	POS	POS	21	24	0	0	>256
S4P4	214	8	SKIN SWAB	<i>S. pseudintermedius</i>	CKCS	MRSP	POS	POS	29	20	0	0	2
S5P5	215	8	SKIN SWAB	<i>S. pseudintermedius</i>	Shar pei	MRSP	POS	POS	26	30	0	0	4
S6P6	216	8	PAW SWAB	<i>S. pseudintermedius</i>	Dachshund	MRSP	POS	POS	22	28	0	0	>256
S7P7	219	8	SKIN SWAB	<i>S. pseudintermedius</i>	British Bulldog	MRSP	POS	POS	21	19	0	0	>256
S8P8	220	8	SKIN SWAB	<i>S. pseudintermedius</i>	British Bulldog	MRSP	POS	POS	21	22	0	0	>256
S9P9	96	9	SKIN SWAB	<i>S. pseudintermedius</i>	Akita	MRSP	POS	POS	23	26	0	0	4
S10P10	190	9	SKIN SWAB	<i>S. pseudintermedius</i>	Akita	MRSP	POS	POS	21	25	0	0	>256
S11P11	187	9	TISSUE	<i>S. pseudintermedius</i>	Bull terrier	MSSP	POS	POS	22	21	0	0	>256
S12P12	188	9	SKIN SWAB	<i>S. pseudintermedius</i>	Great Dane	MRSP	POS	POS	24	24	13	16	1.5
S13P13	189	9	R EAR SWAB	<i>S. pseudintermedius</i>	CKCS	MSSP	NEG	NEG	36	40	26	36	0.125
S14P14	185	9	TISSUE	<i>S. pseudintermedius</i>	Labrador Retriever	MSSP	NEG	NEG	34	32	22	29	0.25
S15P15	191	9	TISSUE	<i>S. pseudintermedius</i>	Maltese X	MSSP	NEG	NEG	33	34	22	25	0.25
S16P16	194	9	SKIN SWAB	<i>S. pseudintermedius</i>	Maltese	MSSP	NEG	NEG	38	42	26	34	0.19
S17P17	195	9	SKIN SWAB	<i>S. pseudintermedius</i>	Shar pei X	MSSP	NEG	NEG	36	38	22	26	0.25
S18P18	196	9	TISSUE	<i>S. pseudintermedius</i>	JRT	MSSP	NEG	NEG	25	40	22	30	0.25
S19P19	197	9	SKIN SWAB	<i>S. pseudintermedius</i>	Labrador Retriever	MSSP	NEG	NEG	38	38	25	32	0.25
S20P20	198	9	SKIN SWAB	<i>S. pseudintermedius</i>	Fox Terrier	MSSP	NEG	NEG	38	36	23	30	0.25
S21P21	199	9	R EAR SWAB	<i>S. pseudintermedius</i>	Labrador Retriever	MSSP	NEG	NEG	36	40	27	37	0.125
S22P22	200	9	SKIN SWAB	<i>S. pseudintermedius</i>	Maltese	MSSP	NEG	NEG	34	35	21	26	0.25
S23P23	203	9						POS	29	26	13	14	1.5

[00308] Ten methicillin susceptible and 13 methicillin resistant *Staphylococcus pseudintermedius* were collected for the study. Isolates were phenotypically classified as methicillin resistant on the basis of in vitro resistance to oxacillin and genetically for the presence of *mecA* gene according to standard procedures.

[00309] Oxacillin and ceftiofur susceptibility testing using disk diffusion technique and Epsilon meter testing were performed. Identification of *mecA* gene was performed using polymerase chain reaction (PCR)

[00310] CLSI disk diffusion susceptibility testing was performed on the 23 *Sp.* isolates for the following antimicrobials: penicillin, amoxicillin, erythromycin, gentamicin, clindamycin, ciprofloxacin, cephalexin, chloramphenicol, tetracycline, oxytetracycline, vancomycin, cefotetan, moxifloxacin and rifampin.

[00311] Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) testing was undertaken using CLSI methodology for NCL812 and included ampicillin as a control. Anti-staphylococcal compounds were then tested against all 23 isolates and minimum inhibitory concentrations (MIC) were determined according to standard protocols. After the MICs were determined, the minimum bactericidal concentrations were performed to determine if these compounds are bacteriostatic or bacteriocidal.

Results

[00312] The *mecA* gene was present in 13 isolates of MRSP and negative in 10 MSSP (Tables 30 and 31). All MRSP isolates were resistant to oxacillin based on disc diffusion (≤ 17 mm) and E-test MIC (≥ 0.5 mg/L).

Table 31: *Staphylococcus pseudintermedius* isolates tested according to Example 6.

	mechA by RT- PCR	Penicillin	Ampicillin	Amoxi- cillin	Erythro- mycin	Genta- mycin	Clinda- mycin	Cipro- floxacin	Cepha- lothin	Chloram- phenicol	Tetra- cyclin	Oxyetra- cycline	Vanco- mycin	Cefotetan	Maxi- floxacin	Rifampin
S1P1	POS	R	R	R	R	R	S	R	R	S	R	R	R	S	S	S
S2P2	POS	R	R	R	R	R	I	R	S	R	S	S	R	S	I	S
S3P3	POS	R	R	R	R	R	R	R	S	R	I	R	S	R	S	S
S4P4	POS	R	R	R	R	R	R	R	I	R	R	R	S	S	S	S
S5P5	POS	R	R	R	R	R	R	R	S	R	R	R	S	S	S	S
S6P6	POS	R	R	R	R	R	R	R	S	R	I	R	S	S	S	S
S7P7	POS	R	R	R	R	R	R	R	R	R	R	R	R	S	S	S
S8P8	POS	R	R	R	R	R	R	R	I	R	R	R	S	S	S	S
S9P9	POS	R	R	S	R	R	R	R	S	R	R	I	S	S	S	S
S10P10	POS	R	R	R	R	R	R	R	S	R	I	R	S	S	S	S
S11P11	POS	R	R	R	R	S	I	S	S	S	S	S	S	S	S	S
S12P12	POS	R	R	S	S	R	S	R	S	S	R	R	S	S	S	S
S13P13	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S14P14	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S15P15	NEG	R	R	S	S	S	S	S	S	S	I	I	S	S	S	S
S16P16	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S17P17	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S18P18	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S19P19	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S20P20	NEG	R	R	S	S	S	S	S	S	S	R	I	S	S	S	S
S21P21	NEG	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
S22P22	NEG	R	R	S	S	S	S	S	S	S	S	S	S	S	S	S
S23P23	POS	R	R	S	R	S	I	S	S	S	S	S	S	S	S	S

[00313] When cefoxitin resistance breakpoint was set at ≤ 24 mm, 3/13 (23%) and 5/13 (38%) of MRSP tested respectively were susceptible to cefoxitin. When cefoxitin resistance break point was set at ≤ 30 mm, only 1/13 (7.7%) of MRSP tested was susceptible (Tables 30 and 31).

[00314] The MRSP isolates were resistant to multiple antibiotic classes. Of the 13 MRSP isolates, all 13 were susceptible to rifampin. 3/13 (23%) were susceptible to chloramphenicol; 10/13 (77%) were susceptible to vancomycin (Tables 30 and 31).

[00315] Interestingly, 3/13 (23%) of the MRSP isolates were susceptible to amoxicillin; 8/13 (62%) were susceptible to cephalothin; 12/13 (92%) susceptible to cefotetan and 12/13 (92%) susceptible to moxifloxacin (Tables 30 and 31).

[00316] All 23 isolates were susceptible to NCL812 based on MICs. In addition, NCL812 has been shown to be bactericidal based on minimal bactericidal concentrations (MBC).

[00317] The MIC range of NCL812 against the *Staphylococcus pseudintermedius* isolates was found to be between 1 $\mu\text{g/mL}$ and 4 $\mu\text{g/mL}$ (Table 32). The MIC₅₀ and MIC₉₀ of NCL812 against the *Staphylococcus pseudintermedius* isolates was found to be 2 $\mu\text{g/mL}$ and 4 $\mu\text{g/mL}$ respectively (Table 33). The MIC mode and MIC range of NCL812 against the *Staphylococcus pseudintermedius* isolates was found to be 2 $\mu\text{g/mL}$ and 1-4 $\mu\text{g/mL}$ respectively (Table 33).

Table 32: MICs of NCL812 and ampicillin against *Staphylococcus pseudintermedius* isolates according to Example 6.

<i>Staphylococcus pseudintermedius</i> isolate	Ampicillin MIC ($\mu\text{g.mL}^{-1}$)	NCL812 MIC ($\mu\text{g.mL}^{-1}$)
S1P1	128	4
S2P2	128	2
S3P3	128	2
S4P4	128	1
S5P5	16	2
S6P6	64	2
S7P7	128	2
S8P8	128	2
S9P9	32	2
S10P10	64	2
S11P11	128	4
S12P12	32	2
S13P13	0.25	2
S14P14	1	2

S15P15	4	4
S16P16	0.25	2
S17P17	1	2
S18P18	4	4
S19P19	0.5	4
S20P20	4	4
S21P21	0.1	2
S22P22	8	4
S23P23	32	2

Table 33: MIC50, MIC90, the MIC mode, and MIC range of NCL812 against *Staphylococcus pseudintermedius* isolates according to Example 6.

Effectiveness against <i>Staphylococcus pseudintermedius</i>	Ampicillin	NCL812
MIC50 (µg/ml)	32	2
MIC90 (µg/ml)	128	4
MIC mode (µg/ml)	128	2
MIC range (µg/ml)	0.1-128	1 - 4

[00318] Methicillin resistant *Staphylococcus pseudintermedius* (MRSP) is an emerging problem in dogs, cats and horses. Two major clonal MRSP lineages have been reported from dogs in Europe (ST 71) and North America (ST 68). There were also reports of MRSP affecting dogs in Japan and a single case of MRSP in veterinary personnel in Hong Kong.

[00319] In this study, MRSP isolates were determined using a combination of presence of *mecA* gene and in vitro resistance to oxacillin. Cefoxitin susceptibility has been used as a substitute for oxacillin for methicillin resistant *Staphylococcus aureus*. However, cefoxitin disk diffusion tests using interpretive guidelines recommended for human isolates of methicillin resistant *Staphylococcus aureus* and coagulase negative staphylococci are unreliable in identifying MRSP. A cefoxitin breakpoint resistance of ≤ 30 mm=resistant and ≥ 31 =susceptible has been proposed by Bemis *et al*, 2012[Bemis, D. A., R. D. Jones, *et al.* (2012). "Evaluation of cefoxitin disk diffusion breakpoint for detection of methicillin resistance in *Staphylococcus pseudintermedius* isolates from dogs." *Journal of Veterinary Diagnostic Investigation* 24(5): 964-967]. This study is in agreement that this breakpoint may be more reliable in predicting methicillin resistant *Staphylococcus pseudintermedius*.

[00320] MRSP isolates are generally resistant to multiple antibiotic classes. Bacterial culture and antibiotic susceptibilities are therefore recommended for all suspect MRSP

infections to allow appropriate selection of antibiotics. A limitation noted in this study is the apparent in vitro susceptibility of MRSP isolates to amoxicillin and cephalosporins (cephalothin and cefotetan).

[00321] NCL812 was effective against all 23 isolates of both MSSP and MRSP. A larger scale study is warranted to confirm the effectiveness of NCL812 against *Staphylococcus pseudintermedius* as it may provide a safe alternative antibiotic option for emerging MRSP infections in domestic animals.

EXAMPLE 7: Preparation and testing of NCL812 analogues (also known as compounds of the invention).

Materials and Methods

NCL812

Analytical grade NCL812 with a defined potency of 960 mg/g (i.e. 96%) was obtained. The powder was stored in a sealed sample container out of direct sunlight and at room temperature at the study site. Aliquots (1 mL) of stock solution (containing 25.6 mg/mL of NCL812 in DMSO) were prepared and stored at -80 °C and defrosted immediately before use.

Synthesising and Testing of NCL812 Analogues

[00322] Analogues NCL001 to NCL230, as identified in Example 13, were synthesised using standard methods in the art. As an example, the methods used to manufacture compounds NCL097; NCL157; NCL179; NCL188; NCL195; and NCL196 are as follows:

NCL 097 (2,2'-bis[(3,4,5-trihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride)

[00323] A suspension of 3,4,5-trihydroxybenzaldehyde (412.0 mg, 2.673 mmol, 2.21 eq.) and *N,N'*-diaminoguanidine hydrochloride (152.0 mg, 1.211 mmol) in EtOH (5 mL) was subjected to microwave irradiation (150 W) at 100 °C for 10 min. The reaction was then allowed to cool to ambient temperature. The resulting precipitate was collected and washed with chilled EtOH (5 mL) and Et₂O (5 mL) to afford the carbonimidicdihydrazide (369.0 mg, 77%) as a pale brown solid. M.P. 292°C (Decomp.). ¹H NMR (300 MHz, DMSO-d₆) δ 9.06 (br s, 6H), 8.25 – 8.01 (m, 4H), 6.83 (s, 4H). ¹³C NMR (75 MHz, DMSO-d₆) δ 152.2, 149.7, 146.2, 136.5, 123.7, 107.4. LRMS(ESI⁺): 361.95 [M + 1]⁺.

NCL157 (2,2'-bis[(2-amino-4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride)

[00324] *Synthesis of 2-amino-4-chloro-N-methoxy-N-methylbenzamide.* To a solution of 2-amino-4-chlorobenzoic acid (5.6691 g, 33.041 mmol), *N,O*-dimethylhydroxylamine hydrochloride (5.7504 g, 58.954 mmol, 1.78 eq.), *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (7.7925 g, 40.649 mmol, 1.23 eq.) and *N*-hydroxybenzotriazole hydrate (5.2371 g, 38.793 mmol (anhydrous basis), 1.17 eq.) in DMF (100 mL) was added diisopropylethylamine (18.0 mL, 13.4 g, 104 mmol, 3.15 eq.) and the brown solution stirred at ambient temperature for 7 h. The reaction was then concentrated *in vacuo* before dilution with 1M NaOH (100 mL) and extracting with CH₂Cl₂ (3 × 100 mL). The combined organic extracts were washed with 1M HCl (100 mL) before drying over MgSO₄ and concentrating *in vacuo* to afford a brown syrup. This oil was then further dried at 60 °C under high vacuum to afford the crude Weinreb amide (7.021 g, 99%) as a brown syrup that crystallised on standing. The crude material was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.24 (d, *J* = 8.4 Hz, 1H), 6.62 (d, *J* = 18 Hz, 1H), 6.54 (dd, *J* = 8.4, 1.9 Hz, 1H), 4.75 (s, 2H), 3.48 (s, 3H), 3.24 (s, 3H). ¹³C NMR (101 MHz, CDCl₃) δ 169.2, 148.4, 137.1, 130.6, 116.6, 116.1, 115.0, 61.1, 34.0.

[00325] *Synthesis of 2-amino-4-chlorobenzaldehyde.* Crude 2-amino-4-chloro-*N*-methoxy-*N*-methylbenzamide (751.1 mg, 3.532 mmol) was broken up into ca. 120 mg batches and each dissolved in THF (10 mL) and cooled to 0 °C before LiAlH₄ (2M in THF, 0.5 mL) was added to each and the solutions stirred for 16 h, allowing the reactions to achieve room temperature. The reactions were quenched with saturated NH₄Cl (1 mL) before being combined, diluted with saturated NaHCO₃ (160 mL) and extracted with CHCl₃ (2 × 150 mL, 1 × 75 mL). The combined organics were dried over MgSO₄ and concentrated *in vacuo* to afford the crude benzaldehyde (463.3 mg, 85%) as yellow/orange crystals. The material was used without further purification. ¹H (400 MHz, CD₃OD) 9.77 (d, *J* = 0.7 Hz, 1H), 7.46 (d, *J* = 8.3 Hz, 1H), 6.63 – 6.71 (m, 1H), 6.63 (dd, *J* = 8.4, 1.9 Hz, 1H). ¹³C NMR (101 MHz, CD₃OD) δ 194.6, 153.0, 142.5, 138.4, 118.3, 116.8, 116.1.

[00326] *Synthesis of 2,2'-bis[(2-amino-4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride.* A suspension of 2-amino-4-chlorobenzaldehyde (128.0 mg, 0.823 mmol, 1.78 eq.) and *N,N*-diaminoguanidine hydrochloride (58.0 mg, 0.462 mmol) in EtOH (2 mL) was subjected to microwave irradiation (100 W) at 60 °C for 5 minutes. Most solvent was then removed *in vacuo*, EtOH (1 mL) was added and the flask was transferred to the freezer to effect crystallisation. The resulting precipitate was collected and washed with EtOH (1 mL) to afford the carbonimidicdihydrazide (21.0 mg, 13%) as a pale yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.71 (br s, 2H), 8.40 (s, 2H), 8.37 (s, 2H), 7.29 (d, *J* = 8.4 Hz, 2H), 6.87 (d, *J* = 2.0 Hz, 2H), 6.73 (br s, 4H), 6.59 (dd, *J* = 8.3, 2.0 Hz, 2H). ¹³C NMR (101 MHz, DMSO-*d*₆) δ 152.1, 151.5, 148.9, 136.0, 134.7, 115.1, 114.5, 112.8.

NCL179 (4,6-bis(2-((*E*)-4-chlorobenzylidene)hydrazinyl)pyrimidin-2-amine)

[00327] A suspension of 2-amino-4,6-dihydrazinylpyrimidine (67.3 mg, 0.434 mmol) and 4-chlorobenzaldehyde (198.8 mg, 1.414 mmol, 3.26 eq.) in EtOH (25 mL) was heated at reflux for 16 h. After this time, the condenser was removed and the solution concentrated to approx. 1 mL and the resulting precipitate filtered hot and washed with Et₂O (10 mL) to afford the aminopyrimidine (42.8 mg, 25%) as an off-white amorphous powder. M.P. 275 °C (Decomp.). ¹H NMR (400 MHz, DMSO) δ 10.70 (s, 2H), 8.02 (s, 2H), 7.67 (d, *J* = 8.4 Hz, 4H), 7.52 (d, *J* = 8.4 Hz, 4H), 6.28 (s, 1H), 5.85 (s, 2H). ¹³C NMR (101 MHz, DMSO) δ 162.8, 162.6, 138.8, 134.1, 133.1, 128.9, 127.6, 73.5.

NCL188 ((*E*)-2-(1-(4-chlorophenyl)pentylidene)hydrazine-1-carboximidamide hydrochloride)

[00328] A suspension of 1-(4-chlorophenyl)pentanone (1.8319 g, 9.3146 mmol, 1.95 eq.) and aminoguanidine hydrochloride (527.6 mg, 4.773 mmol) in EtOH (15 mL) was heated at 65 °C for 16 h. The crude was cooled to ambient temperature before being diluted with Et₂O (60 mL) and cooled to 0 °C to precipitate unreacted aminoguanidine hydrochloride (174.5 mg). The mother liquors were then concentrated *in vacuo* and the residue dissolved in Et₂O (20 mL). The solution was then boiled and hexanes (10 mL) added to afford the caboximidamide as a cream solid. ¹H NMR (400 MHz, DMSO) δ 11.54 (s, 1H), 7.99 (d, *J* = 8.7 Hz, 2H), 7.90 (s, 3H), 7.47 (d, *J* = 8.6 Hz, 2H), 2.91 – 2.82 (m, 2H), 1.48 – 1.32 (m, 4H), 0.89 – 0.84 (m, 3H). ¹³C NMR (101 MHz, DMSO) δ 156.2, 153.8, 134.8, 134.4, 128.7, 128.4, 28.1, 26.6, 22.0, 13.8

NCL195 (4,6-bis(2-((*E*)-4-methylbenzylidene)hydrazinyl)pyrimidin-2-amine)

A suspension of 2-amino-4,6-dihydrazinopyrimidine (58.9 mg, 0.380 mmol) and 4-methylbenzaldehyde (0.10 mL, 100 mg, 0.832 mmol, 2.19 eq.) in EtOH (4 mL) was heated at reflux for 16 h. The reaction mixture was cooled to ambient temperature before collecting the pellet-like precipitate, washing with Et₂O (20 mL). The 'pellets' were then crushed and the solid further washed with Et₂O (10 mL) to afford the pyrimidine (85.8 mg, 63%) as a white 'fluffy' powder. M.P. 274-276 °C. ¹H NMR (400 MHz, DMSO) δ 10.51 (s, 2H), 8.00 (s, 2H), 7.54 (d, *J* = 8.0 Hz, 4H), 7.26 (d, *J* = 7.9 Hz, 4H), 6.26 (s, 1H), 5.77 (s, 2H), 2.34 (s, 6H). ¹³C NMR (101 MHz, DMSO) δ 162.8, 162.6, 140.1, 138.4, 132.5, 129.4, 126.0, 73.3, 21.0.

NCL196 (4,4'-((1*E*,1'*E*)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol)

[00329] A suspension of 2-amino-4,6-dihydrazinopyrimidine (70.4 mg, 0.454 mmol) and 4-hydroxybenzaldehyde (140.3 mg, 1.149 mmol, 2.53 eq.) in EtOH (3 mL) was heated at reflux for 16 h. The reaction mixture was cooled to ambient temperature before collecting the precipitate, washing with Et₂O (25 mL), to afford the pyrimidine (91.4 mg, 55%) as an off-white powder. M.P. 298 °C (Decomp.). ¹H NMR (400 MHz, DMSO) δ 10.31 (s, 2H), 9.74 (s, 2H), 7.94

(s, 2H), 7.48 (d, $J = 8.6$ Hz, 4H), 6.83 (d, $J = 8.6$ Hz, 4H), 6.20 (s, 1H), 5.70 (s, 2H). ^{13}C (101 MHz, DMSO) δ 162.7, 162.5, 158.3, 140.5, 127.7, 126.3, 115.7, 73.0.

MIC tests

[00330] Minimum inhibitory concentrations ($\mu\text{g/ml}$) were determined using the broth microdilution method recommended by the Clinical and Laboratory Standards Institute (CLSI). MIC breakpoints were determined by visual assessment and then confirmed using an ELISA plate reader, measuring absorbance levels at 600 nm. Bacterial growth (turbidity) in the wells with antimicrobial was compared with the amount of growth (turbidity) in the growth-control well (containing no antimicrobial). All isolates were tested in duplicate, if there was a difference of greater than one two-fold dilution in the results, the test was repeated a third time. The purity of the isolates was closely monitored during testing by subculturing the prepared bacterial inoculum onto SBA (sheep blood agar). The MICs of the control strains for the antimicrobial ampicillin were determined for each testing run as an internal quality control. The MIC_{50} , MIC_{90} and MIC range (minimum and maximum) were calculated for each of the bacterial groups.

Activity of NCL812 and MIC against Gram-negative bacteria

[00331] The activity of NCL812 against Gram-negative bacteria was assessed using the broth microdilution method recommended by the Clinical and Laboratory Standards Institute (CLSI), and the MICs ($\mu\text{g/mL}$) for NCL812 and ampicillin were determined.

Determination of Minimum Bactericidal Concentration (MBC)

CLSI methodology

[00332] Briefly, 10 μL of the contents of each well starting at the MIC was inoculated on to a Columbia SBA plate and incubated at 37 °C for 48 h. Plates were examined at 24 and 48 h and the MBC was recorded as the lowest concentration of NCL812 at which no colonies of bacteria were observed on the plate (or significant inhibition of growth was observed compared to the control) (CLSI 2005).

Kill kinetics assays (MRSA & VRE)

[00333] MRSA/VRE were grown overnight on Columbia SBA at 37 °C. A few colonies of bacteria were then suspended in CAMHB and adjusted to an optical density of 0.08 to 0.10. The bacterial suspension was diluted 1:10. One millilitre of the bacteria was added to 9 mL of CAMHB containing various concentrations (up to 4 x MIC) of NCL812, to achieve a final bacterial concentration of 1 to 3×10^8 CFU/ml. The tubes were incubated at 37 °C. In order to determine the number of viable bacteria present at various time points, a 100 μL aliquot was

removed from each tube and diluted in normal saline. Then, 100 µL of each dilution was spread onto colony count agar, in duplicate, and incubated for 48 h at 37 °C. After 24 h the numbers of colonies present on each plate were counted and therefore the number of viable bacteria present in the original suspension enumerated. Plates were re-checked after 48 hours.

Synergy studies with other classes of antimicrobial agent.

[00334] The checkerboard method (Gunics et al., 2000 *Int. J. Antimicrob. Agents*. 14:239-42) was used to find interactions (synergy, antagonism, no effect) of NCL812 in combination with tetracycline, chloramphenicol, erythromycin (macrolide), ampicillin (β-lactam broad-spectrum), gentamicin (aminoglycoside), ciprofloxacin (fluoroquinolone), sulfamethoxazole (sulphonamide), or penicillin G (β-lactam narrow-spectrum). For initial experiments, a laboratory strain of *Staphylococcus aureus* T3-129 was used, however this strain gave inconsistent results for some of the antimicrobials and a new strain of *Staphylococcus* spp. designated MK1 (definitive species identification currently in progress) that was sensitive to all tested antimicrobials was used in subsequent tests.

[00335] Firstly, the MIC of each antibiotic alone was determined in accordance to CLSI standard guidelines. Secondly, the combination of NCL812 with each of above antibiotics was tested in duplicate. To evaluate the effect of the combination the fractional inhibitory concentration (FIC) was calculated for each antibiotic as follows:

FIC of tested antibiotic = MIC of tested antibiotic in combination / MIC of antibiotic alone.

FIC of NCL812 = MIC of NCL812 in combination / MIC of NCL812 alone.

FIC_i = FIC index = FIC of NCL812 + FIC of each tested antibiotic.

[00336] According to the checkerboard guidelines, Synergy (S) was defined as an $FIC_i < 0.5$. No effect (NE) was defined as $0.5 < FIC_i < 4$. Antagonism (A) was defined as a $4 < FIC_i$.

Testing of NCL812 Analogues

[00337] NCL812 analogues were stored at 4 °C until assayed. MICs were determined against two MRSA strains, two VRE strains and one strain each of *E. coli* and *Pseudomonas aeruginosa*.

Results

Determination of Minimum inhibitory Concentration (MIC)

[00338] The comparative NCL812 and ampicillin MIC values ($\mu\text{g/mL}$) for 21 MRSA isolates were obtained. The results for the original experiments (phase I), and repeat testing (phase II) are shown in Table 34. Each MIC test was performed in duplicate.

Table 34: NCL812 and ampicillin MIC values ($\mu\text{g/ml}$) for 21 MRSA isolates obtained according to Example 7.

Organism	Phase I					Phase II				
	NCL812 1st	NCL812 2nd	Ampicillin 1st	Ampicillin 2nd		NCL812 1st	NCL812 2nd	Ampicillin 1st	Ampicillin 2nd	
MRSA 516	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	16 $\mu\text{g/ml}$	
MRSA 570	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	
MRSA 580	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	
MRSA 606	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 610	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 616	8 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 622	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 638	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	
MRSA 713	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	
MRSA 718	4 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 728	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 734	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 741	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	
MRSA 747	8 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 773	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 776	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	
MRSA 786	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 787	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	
MRSA 815	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	
MRSA 823	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	
MRSA 844	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$		4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	

Table 35: Comparative NCL812 and ampicillin MIC values ($\mu\text{g/ml}$) for 13 VRE isolates obtained according to Example 7.

Organism ¹	Phase I				Phase II			
	NCL812 1st	NCL812 2nd	Ampicillin 1st	Ampicillin 2nd	NCL812 1st	NCL812 2nd	Ampicillin 1st	Ampicillin 2nd
VRE 002 [H12c]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 003 [H14i]	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 004 [H16c]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 008 [H16c(dc)]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$
VRE 005 [H17c]	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 006 [H19i]	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 007 [H23c]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$
VRE 009 [H25c]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$
VRE 011 [H26c(dc)]	2 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$
VRE 010 [H26c(w)]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$
VRE 012 [H35i]	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	0.25 $\mu\text{g/ml}$	0.5 $\mu\text{g/ml}$
VRE 014 [H37c(g)]	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$
VRE 013 [H37c(w)]	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	2 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$

[00339] The comparative NCL812 and ampicillin MIC values ($\mu\text{g/mL}$) for 13 VRE isolates were obtained. The results for the original experiments (phase I), and repeat testing (phase II) are shown in Table 35. Each MIC test was performed in duplicate.

[00340] NCL812 MIC₅₀, MIC₉₀, MIC mode and MIC range were obtained for Australian isolates of MRSA and VRE, as shown in Table 6. Comparative MIC values for ampicillin are shown in parentheses.

Activity of NCL812 and MIC against Gram-negative bacteria

[00341] Comparative NCL812 and ampicillin MIC values ($\mu\text{g/mL}$) for *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella arizonae* were obtained, as shown in Table 36. Each MIC test was performed in duplicate.

Table 36: Comparative NCL812 and ampicillin MIC values ($\mu\text{g/ml}$) for *Escherichia coli*, *Pseudomonas aeruginosa* and *Salmonella arizonae* obtained according to Example 7.

Organism	NCL812 1 st	NCL812 2 nd	Ampicillin 1 st	Ampicillin 2 nd
<i>E. coli</i>	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$
<i>P. aeruginosa</i>	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$
<i>S. arizonae</i>	>128 $\mu\text{g/ml}$	>128 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$	1 $\mu\text{g/ml}$

[00342] The antimicrobial activity of NCL812 against the selected Gram-negative bacteria was >128 $\mu\text{g/ml}$.

Determination of Minimum Bactericidal Concentration (MBC)

[00343] MBC results for MRSA isolates are shown in Table 37 which shows NCL812 MBC values ($\mu\text{g/mL}$) for 20 MRSA isolates. Each MBC test was performed in duplicate starting from the NCL812 MIC concentration to 16 times the MIC. For all isolates, the MBC was equal to the MIC. However, inconsistent growth on agar plates was recorded for some concentrations.

Table 37: NCL812 MBC values ($\mu\text{g/ml}$) for 20 MRSA isolates according to Example 7.

Organism / Sample No.		NCL812 MBC				
		4 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	16 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$
MRSA 1	- 1 st	0	0	0	0	N**
	- 2 nd	0	0	0	0	N
MRSA 2	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 3	- 1 st	0	GB*	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 4	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N

MRSA 5	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 6	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 7	- 1 st	0	0	0	0	N
	- 2 nd	0	GB	0	0	N
MRSA 8	- 1 st	GB	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 9	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 10	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 11	- 1 st	0	0	0	0	N
	- 2 nd	GB	0	0	0	N
MRSA 12	- 1 st	0	0	0	0	N
	- 2 nd	0	GB	0	0	N
MRSA 13	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 14	- 1 st	0	0	0	0	N
	- 2 nd	0	0	0	0	N
MRSA 516	- 1 st	0	0	0	0	0
	- 2 nd	0	0	0	0	0
MRSA 570	- 1 st	0	0	0	0	0
	- 2 nd	0	0	0	0	0
MRSA 580	- 1 st	0	0	0	0	0
	- 2 nd	0	0	0	0	0
MRSA 606	- 1 st	0	0	0	0	GB
	- 2 nd	0	0	0	0	0
MRSA 610	- 1 st	0	0	0	0	0
	- 2 nd	0	GB	0	0	0

GB = Bacterial Growth on Sheep Blood Agar

N** = Not cultured on Sheep Blood Agar

[00344] The results for 10 VRE isolates are shown in Table 38. Each MBC test was performed in duplicate starting from NCL812 MIC concentration to 32 times the MIC. As with the MRSA isolates tested, the MBC appears to be equal to the MIC. However, with the VRE isolates, an anomaly was observed at higher NCL812 concentrations. There is significant inhibition of growth at concentrations near to the MIC, but as the NCL812 concentration increases, bacteria appear to be less subject to inhibition. High numbers of bacteria were observed on the plates at NCL812 concentrations ≥ 16 $\mu\text{g/mL}$.

Table 38: NCL812 MBC values ($\mu\text{g/ml}$) for 10 VRE isolates according to Example 7

Organism / Sample No.		NCL812 MBC						
		2 $\mu\text{g/ml}$	4 $\mu\text{g/ml}$	8 $\mu\text{g/ml}$	16 $\mu\text{g/ml}$	32 $\mu\text{g/ml}$	64 $\mu\text{g/ml}$	128 $\mu\text{g/ml}$
VRE 26c(dc)	- 1 st	90*	20	4000	M	M	M	M
	- 2 nd	0	70	3500	M	M	M	M
VRE 37c	- 1 st	500	100	20	250	M	M	M
	- 2 nd	M	50	100	1100	1400	M	M
VRE 36t	- 1 st	0	0	0	720	0	0	0
	- 2 nd	0	0	0	0	10	20	10
VRE 16c(dc)	- 1 st	90	330	0	M	M	M	M

	- 2 nd	200	0	20	M	M	M	M
VRE 23c	- 1 st	0	120	20	10	M	M	M
	- 2 nd	0	0	0	0	570	M	M
VRE 25c	- 1 st	0	0	M	M	M	M	M
	- 2 nd	20	20	M	M	M	M	M
VRE 16c	- 1 st	10	820	980	M	M	M	M
	- 2 nd	M	790	890	M	M	M	M
VRE 19t	- 1 st	0	0	0	180	10	110	M
	- 2 nd	30	0	0	70	40	M	M
VRE 14t	- 1 st	10	0	10	0	180	970	M
	- 2 nd	0	0	0	40	780	M	M
VRE 12c	- 1 st	0	0	0	M	M	M	M
	- 2 nd	0	M	300	M	M	M	M

*Number of bacteria growing after 24 hours per ml of sample (CFU/ml); M = many bacteria growing on the plate (too many to count)

[00345] NCL812 was found to be bactericidal against Gram-positive bacteria at concentrations equivalent to the MIC.

Kill kinetics assays (MRSA & VRE)

[00346] In preliminary experiments, colony counts were performed at $t = 15, 30, 45$ and 60 min. No significant changes in the bacterial concentration were observed at these time points, suggesting that NCL812 is not rapidly bactericidal (by comparison, the lipoglycopeptide oritavancin (McKay et al. (2009) *J. Antimicrob. Chemother.* 63 (6): 1191-1199) caused a $3\log_{10}$ reduction in viable count within an hour of exposure to a concentration equivalent to the C_{max}). Therefore, for future experiments, sampling time points were extended out to one and then two hour intervals.

[00347] In the initial experiments, for MRSA, at 4 h a reduction of at least $2.5\log_{10}$ CFU/mL was observed in comparison to the growth control. At 8 h there was at least a $3.5\log_{10}$ CFU/mL difference between the control and the bacteria exposed to NCL812. After 24 h the numbers of bacteria present in all NCL812 concentrations was not significantly different to the control. There was a consistent reduction in bacterial numbers at NCL812 concentrations from $4-16$ $\mu\text{g/mL}$, up to 8 h, but the same was not observed for concentrations greater than 16 $\mu\text{g/mL}$. By comparison, most bactericidal antimicrobial agents, used or being developed for the treatment of MRSA and VRE (oritavancin, daptomycin, vancomycin) are rapidly bactericidal achieving similar log reductions within 1 h of exposure in a concentration dependent manner (McKay et al., 2009). In kill kinetics experiments bacteriostatic antimicrobials recommended for the treatment of MRSA and VRE infections (teicoplanin and linezolid only marginally decrease the viable count and growth).

[00348] For VRE the observed decrease in the CFU/mL of bacteria exposed to NCL812 was less than for MRSA. At 4 h there was approximately a $2\log_{10}$ reduction in the viable count

compared to the control, and at 8 h there was approximately a $2.5\log_{10}$ reduction. However, at 24 h the growth of bacteria was no longer as significantly reduced in comparison to the control. Bacterial numbers increased after 8 h incubation and this affect appeared to be more pronounced with increasing concentrations of NCL812.

[00349] The kill kinetics of MRSA 580 were obtained at different concentrations of NCL812 over a period of 8 h, as shown in **Figure 33** The kill kinetics of MRSA 580 in different concentrations of NCL812 over a period of 24 h are shown in **Figure 34** After 4 h of incubation the media was changed to fresh media containing the same concentration of NCL812.

[00350] The kill kinetics of MRSA 698 in different concentrations of NCL812 over a period of 24 h is shown in **Figure 35** After 4 h of incubation the media was changed to fresh media containing the same concentration of NCL812.

[00351] The kill kinetics of VRE 26c(dc) at different concentrations of NCL812 over a period of 24 h are shown in **Figure 36**.

[00352] The kill kinetics of VRE 16c(dc) at different concentrations of NCL812 over a period of 24 h are shown in **Figure 37**.

Test for bacterial resistance to NCL812

[00353] Preliminary tests were carried out to determine whether bacterial resistance may account for the observations of bacterial growth at higher concentrations of NCL812 and the increase in bacterial numbers in kill kinetics experiments at 24 h incubation. Bacteria (MRSA) growing at high concentrations of NCL812, in the 96 well micro titre tray, were subcultured on to SBA and incubated for 24 h, then MIC testing was performed. There was no change in the MIC of these bacteria. Bacteria growing in broth used for kill kinetics experiments was also tested for any change in the MIC. No change was observed.

[00354] Additionally, bacteria exposed to NCL812 at high concentrations were then subcultured on to plate count agar containing NCL812 (64 $\mu\text{g/mL}$ and 128 $\mu\text{g/mL}$) and incubated for 24 h at 37 °C. Bacteria growing on the plate were then used for running an MIC test. There was still no change in the MIC of the bacteria.

[00355] In summary, NCL812 has bactericidal activity against MRSA which is less pronounced against VRE strains. The bactericidal effect is not rapid in comparison to bactericidal antimicrobials developed for MRSA and VRE infections (daptomycin, oratovancin, vancomycin). Aberrant bactericidal results at higher concentrations of NCL812 are not indicative of resistance development, but may be suggestive of loss of activity. Stability testing of the compound in broth media should therefore be undertaken before exploring these interesting but

currently unexplained results. This will include a detailed examination of the literature to determine if this phenomenon is observed in other classes of antimicrobial agent. In this case, closer examination of the kill kinetics between 8 and 24 h will be required. NCL812 kill curves for MRSA and VRE suggest more bactericidal activity in comparison to bacteriostatic antimicrobials (linezolid, teicoplanin). Kill curves should now be generated for *Streptococcus pneumoniae* once stability issues of NCL812 are investigated, as for example with linezolid, as some antibacterial agents can be bacteriostatic against some bacteria and bactericidal against others.

Synergy studies with other classes of antimicrobial agent.

[00356] MICs, FICs, FIC_i and the interaction between NCL812 and eight antibiotics is shown in Table 39. None of the eight tested compounds, representing distinct classes of antimicrobial agent showed either positive (synergism) or negative (antagonism) interaction with NCL812 consistent with an additive effect when antibacterial agents are added to NCL812.

Table 39: MICs, FICs, FIC_i and the interaction between NCL812 and eight antibiotics according to Example 7.

Antibiotic Name	Experiment	Repeat	Antibiotic			NCL812			FIC _i	Result
			MIC (µg/ml)		FIC ₁	MIC (µg/ml)		FIC ₂		
			With NCL812	Alone		With Antibiotic	Alone			
Tetracycline ¹	1	1 st	0.25	0.5	0.5	1	4	0.25	0.75	NE
		2 nd	0.25	0.5	0.5	1	8	0.125	0.62	NE
	2	1 st	0.031	0.25	0.125	4	8	0.5	0.625	NE
		2 nd	0.031	0.25	0.125	4	8	0.5	0.625	NE
Chloramphenicol ¹	1	1 st	4	8	0.5	1	4	0.25	0.75	NE
		2 nd	2	4	0.5	2	8	0.25	0.75	NE
	2	1 st	4	8	0.5	2	8	0.25	0.75	NE
		2 nd	0.5	8	0.0625	4	8	0.5	0.562	NE
Erythromycin ¹	1	1 st	0.031	0.125	0.25	2	4	0.5	0.75	NE
		2 nd	0.007	0.125	0.063	2	4	0.5	0.562	NE
	2	1 st	0.007	0.25	0.25	2	8	0.25	0.5	NE
		2 nd	0.007	0.25	0.031	4	8	0.5	0.531	NE
Ampicillin ¹	1	1 st	0.125	0.25	0.5	1	4	0.25	0.75	NE
		2 nd	0.25	0.5	0.5	0.125	4	0.031	0.53	NE
	2	1 st	0.062	0.125	0.5	2	8	0.25	0.75	NE
		2 nd	0.125	0.25	0.5	2	8	0.25	0.75	NE
Gentamicin ²	1	1 st	0.062	0.125	0.5	0.5	4	0.125	0.625	NE
		2 nd	0.062	0.125	0.5	1	4	0.25	0.75	NE
	2	1 st	0.5	1	0.5	1	4	0.25	0.75	NE
		2 nd	0.007	0.5	0.0156	2	4	0.5	0.515	NE
Ciprofloxacin ²	1	1 st	0.062	0.125	0.5	2	4	0.5	0.75	NE
		2 nd	0.003	0.125	0.025	4	2	0.5	0.525	NE
	2	1 st	0.125	0.25	0.5	0.5	4	0.125	0.625	NE
		2 nd	0.125	0.25	0.5	0.25	4	0.0625	0.562	NE
Sulfamethoxazole ²	1	1 st	4	8	0.5	1	4	0.25	0.75	NE

		2 nd	4	8	0.5	2	4	0.5	1	NE
	2	1 st	4	8	0.5	1	4	0.25	0.75	NE
		2 nd	4	8	0.5	2	4	0.5	1	NE
Penicillin G ²	1	1 st	0.062	0.125	0.5	2	4	0.5	1	NE
		2 nd	0.062	0.125	0.5	2	4	0.5	1	NE
	2	1 st	0.062	0.125	0.5	2	4	0.5	1	NE
		2 nd	0.031	0.25	0.125	2	4	0.5	0.625	NE

¹ *S. aureus* strain T3-29

² *Staphylococcus* spp. Strain MK1

FIC₁ = MIC of antibiotic in combination with NCL812/MIC of antibiotic alone

FIC₂ = MIC of NCL812 in combination with antibiotic/MIC of NCL812 alone

FIC_i = FIC index

Testing of NCL812 Analogues

[00357] The chemical structures of analogues NCL001 to NCL230 are shown in **Example 13**.

[00358] MICs for NCL812 and analogues NCL001-070 are shown in **Table 40**.

[00359] MICs for analogues NCL071 to 171 are shown in **Table 41**.

[00360] MICs for analogues NCL171 to 230 are shown in **Table 42**.

Table 40: MICs for NCL812 and analogues NCL001-070 according to Example 7.

Compound		MIC (µg/ml) at 24 hours					
		MRSA 1	MRSA 2	VRE 1	VRE 2	E. coli	P. aeruginosa
Ampicillin	Amp	>128	16	0.5	0.25	4	>128
NCL812	1 st	4	4	2	2	>128	>128
	2 nd	4	2	2	1	>128	>128
NCL001	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL002	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL003	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL004	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL005	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL006	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL007	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL008	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL009	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL010	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL011	1 st	>128	>128	>128	32	>128	>128
	2 nd	>128	>128	>128	32	>128	>128
NCL012	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL013	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL014	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL015	1 st	>128	>128	>128	128	>128	>128
	2 nd	>128	>128	>128	128	>128	>128
NCL016	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL017	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL018	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128

NCL019	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL020	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL021	1 st	16	16	16	16	8	8	8	8	>128	>128
	2 nd	16	16	16	16	8	8	8	8	>128	>128
NCL022	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL023	1 st	16	16	16	16	32	32	16	16	>128	>128
	2 nd	16	16	16	16	32	32	16	16	>128	>128
NCL024	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL025	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL026	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL027	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL028	1 st	>128	>128	>128	>128	>128	>128	128	128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	128	128	>128	>128
NCL029	1 st	32	32	32	32	32	32	32	32	>128	>128
	2 nd	32	32	32	32	32	32	32	32	>128	>128
NCL030	1 st	128	128	128	128	64	64	64	64	>128	128
	2 nd	128	128	128	128	64	64	64	64	>128	128
NCL031	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL032	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL033	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL034	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL035	1 st	16	16	16	16	8	8	4	4	>128	>128
	2 nd	16	16	16	16	8	8	4	4	>128	>128
NCL036	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL037	1 st	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128	>128	>128
NCL038	1 st	8**	8**	8**	8**	8**	8**	8**	8**	>128	>128
	2 nd	8**	8**	8**	8**	8**	8**	8**	8**	>128	>128
NCL039	1 st	32	32	32	32	32	32	8	8	>128	>128
	2 nd	64	64	32	32	32	32	8	8	>128	>128
NCL040	1 st	8	8	8	8	8	8	8	8	>128	>128
	2 nd	8	8	8	8	8	8	8	8	>128	>128
NCL041	1 st	64	64	64	64	64	64	64	64	128	128

	2 nd	64	64	64	64	64	128
NCL042	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL043	1 st	128	128	128	128	128	128
	2 nd	128	128	128	64	128	128
NCL044	1 st	128	128	128	128	>128	128
	2 nd	128	128	128	128	>128	128
NCL045	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL046	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL047	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL048	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL049	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL050	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL051	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL052	1 st	128	128	128	128	128	128
	2 nd	128	128	128	128	128	128
NCL053	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL054	1 st	4	2	4	4	4	>128
	2 nd	2	4	4	4	4	>128
NCL055	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL056	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL057	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL058	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL059	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL060	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128
NCL061	1 st	>128	>128	>128	>128	8	>128
	2 nd	>128	>128	>128	>128	8	>128
NCL062	1 st	2	4	2	2	1	>128
	2 nd	2	4	2	2	1	>128
NCL063	1 st	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128

NCL064	1 st	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128
NCL065	1 st	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128
NCL066	1 st	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128
NCL067	1 st	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128
NCL068	1 st	>128	>128	>128	>128	>128	>128	>128	>128
	2 nd	>128	>128	>128	>128	>128	>128	>128	>128
NCL069	1 st	128	128	128	128	64	>128	>128	>128
	2 nd	128	128	128	128	64	>128	>128	>128
NCL070	1 st	>128	>128	>128	>128	32	>128	>128	>128
	2 nd	>128	>128	>128	>128	32	>128	>128	>128

Table 41: MICs for analogues NCL071-170 according to Example 7.

NCL Compound Code	MIC (µg/ml) at 24 hours					
	MRSA 1	MRSA 2	VRE 1	VRE 2	<i>E. coli</i>	<i>P. aeruginosa</i>
NCL071	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL072	32	64	64	64	>128	>128
	32	64	64	64	>128	>128
NCL073	128	128	16	64	>128	>128
	128	128	16	64	>128	>128
NCL074	16	16	2	8	>128	>128
	16	16	1	4	>128	>128
NCL075	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL076	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL077	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL078	4	16	4	4	>128	>128
	4	8	2	2	>128	>128
NCL079	4	4	2	4	>128	>128
	4	8	2	4	>128	>128
NCL080	4	4	2	4	>128	>128
	4	8	2	4	>128	>128
NCL081	4	8	1	4	>128	>128
	4	4	1	4	>128	>128
NCL082	8	8	4	2	>128	>128
	4	8	4	4	>128	>128
NCL083	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL084	2	8	2	2	>128	>128
	2	4	2	2	>128	>128
NCL085	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL086	>128	>128	>128	128	>128	>128
	>128	>128	>128	128	>128	>128
NCL087	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL088	16	8	4	4	>128	>128
	8	8	4	4	>128	>128
NCL089	4	4	0.5	1	>128	>128
	8	4	0.5	1	>128	>128
NCL090	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL091	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL092	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL093	64	128	64	>128	>128	>128
	64	128	64	>128	>128	>128
NCL094	128	128	64	128	>128	>128
	128	128	64	128	>128	>128
NCL095	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL096	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL097	128	32	32	32	>128	128
	128	32	32	32	>128	>128
NCL098	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL099	8	4	8	8	>128	>128
	8	4	8	8	>128	>128

NCL100	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL101	32	32	64	128	>128	>128
	32	32	64	128	>128	>128
NCL102	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL103	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL104	64	64	128	>128	>128	>128
	64	64	128	>128	>128	>128
NCL105	64	>128	64	>128	>128	>128
	64	>128	64	>128	>128	>128
NCL106	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL107	6	16	32	128	>128	>128
	8	16	32	128	>128	>128
NCL108	64	64	64	128	>128	>128
	64	64	64	128	>128	>128
NCL109	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL110	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL111	64	64	64	>128	>128	>128
	64	64	64	>128	>128	>128
NCL112	>128	128	64	32	>128	>128
	>128	128	64	32	>128	>128
NCL113	16	8	8	16	>128	>128
	16	8	8	16	>128	>128
NCL114	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 115	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 116	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 117	64	128	64	64	>128	>128
	64	128	64	64	>128	>128
NCL 118	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 119	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 120	8	4	2	8	>128	>128
	8	8	2	8	>128	>128
NCL 121	32	32	8	32	>128	>128
	32	32	8	64	>128	>128
NCL 122	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL123	16	4	4	4	>128	>128
	32	8	4	4	>128	>128
NCL 124	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 125	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 126	>128	>128	4	>128	>128	>128
	>128	>128	8	>128	>128	>128
NCL 127	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 128	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 129	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 130	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 131	>128	>128	4	>128	>128	>128

	>128	>128	4	>128	>128	>128
NCL 132	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 133	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 134	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 135	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 136	64	128	32	128	>128	>128
	64	128	32	128	>128	>128
NCL 137	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 138	>128	32	>128	>128	>128	>128
	>128	32	>128	>128	>128	>128
NCL 139	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 140	>128	16	2	8	>128	>128
	>128	32	2	8	>128	>128
NCL 141	>128	>128	16	>128	>128	>128
	>128	>128	16	>128	>128	>128
NCL 142	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 143	16	16	8	8	>128	>128
	16	16	8	8	>128	>128
NCL 144	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 145	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 146	8	16	16	16	>128	>128
	8	16	16	16	>128	>128
NCL 147	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 148	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 149	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 150	128	128	>128	>128	>128	>128
	128	128	>128	>128	>128	>128
NCL 151	64	64	128	64	>128	>128
	64	64	128	64	>128	>128
NCL 152	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 153	32	16	8	8	>128	>128
	32	16	8	8	>128	>128
NCL 154	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 155	8	8	16	8	>128	>128
	8	8	16	8	>128	>128
NCL 156	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 157	2	2	2	2	>128	>128
	2	2	2	4	>128	>128
NCL 140	8	16	16	16	>128	>128
	8	16	16	16	>128	>128
NCL 038	8	8	8	8	>128	>128
	8	8	8	8	>128	>128
NCL 158	4	4	4	4	>128	>128
	4	4	4	4	>128	>128
NCL 159	64	128	16	128	>128	>128
	64	128	16	128	>128	>128
NCL 160	32	128	16	64	>128	>128
	64	128	16	64	>128	>128

NCL 161	64	128	128	128	>128	>128
	128	128	128	128	>128	>128
NCL 162	>128	128	>128	>128	>128	>128
	>128	128	>128	>128	>128	>128
NCL 163	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 164	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 165	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 166	32	64	8	128	>128	>128
	32	64	8	128	>128	>128
NCL 167	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 168	32	32	64	32	>128	>128
	32	32	64	32	>128	>128
NCL 169	32	64	32	32	>128	>128
	32	64	32	32	>128	>128
NCL 170	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128

Table 42: MICs for analogues NCL 171-230 according to Example 7.

NCL Compound Code	MIC (µg/ml) at 24 hours					
	MRSA 1	MRSA 2	VRE 1	VRE 2	<i>E. coli</i>	<i>P. aeruginosa</i>
NCL 171	128	64	32	>128	>128	>128
	128	64	32	>128	>128	>128
NCL 172	128	64	>128	>128	>128	>128
	128	64	>128	>128	>128	>128
NCL 173	16	16	4	32	>128	>128
	16	16	4	32	>128	>128
NCL 174	32	32	4	32	>128	>128
	32	32	4	32	>128	>128
NCL 175	64	>128	32	>128	>128	>128
	64	>128	32	>128	>128	>128
NCL 176	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL 177	8	8	4	8	>128	>128
	8	8	4	8	>128	>128
NCL Compound Code	MIC (µg/ml) at 24 hours					
	<i>S. aureus</i> ATCC29213	<i>E. faecalis</i> #32	<i>E. coli</i> ATCC25922	<i>Pseudomonas</i> #008		
NCL 178	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 179	4	8	>64	>64		
	4	8	>64	>64		

NCL 180	>64	64	>64	>64		
	>64	64	>64	>64		
NCL 181	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 182	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 183	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 184	>64	64	>64	>64		
	>64	64	>64	>64		
NCL 185	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 186	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 187	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 188	32	32	64	>64		
	32	32	64	>64		
NCL 189	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 190	32	64	>64	>64		
	64	64	>64	>64		
NCL 191	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL 192	>64	>64	>64	>64		
	>64	>64	>64	>64		
NCL Compound Code	MIC (µg/ml) at 24 hours					
	MRSA 1	MRSA 2	VRE 1	VRE 2	<i>E. coli</i>	<i>P. aeruginosa</i>
NCL193	2	2	4	4	>128	>128
	2	4	4	8	>128	>128
NCL194	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL195	2	2	4	2	>128	>128
	2	2	4	4	>128	>128
NCL196	32	64	32	64	64	64
	32	32	64	64	64	128
NCL197	8	8	8	8	>128	>128
	8	8	4	8	>128	>128
NCL198	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL199	2	4	>128	>128	>128	>128
	4	4	>128	>128	>128	>128

NCL200	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL201	32	16	>128	>128	>128	>128
	64	32	>128	>128	>128	>128
NCL202	8	8	8	8	>128	>128
		8	8	8	>128	>128
NCL203	32	64	32	32	>128	>128
	32	64	64	32	>128	>128
NCL204	4	4	64	>128	>128	>128
	8	8	64	>128	>128	>128
NCL205	4	4	64	>128	>128	>128
	8	8	64	>128	>128	>128
NCL206	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL207	16	32	>128	>128	>128	>128
	32	32	>128	>128	>128	>128
NCL208	>128	>128	>128	>128	>128	>128
	>128		>128	>128	>128	>128
NCL209	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL210	>128	>128	128	32	>128	>128
	>128	>128	128	64	>128	>128
NCL211	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL212	>128	>128	32	16	>128	>128
	>128	128	32	16	>128	>128
NCL213	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL214	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL215	8	8	2	4	>128	>128
	16	8	4	8	>128	>128
NCL216	2	2	2	4	>128	>128
	4		2	4	>128	>128
NCL217	4	4	2	4	>128	>128
	4	4	2	4	>128	>128
NCL218	>128	>128	16	>128	>128	>128
	>128	>128	16	>128	>128	>128
NCL219	2	2	16	16	>128	>128
	2	2	16	16	>128	>128
NCL220	16	16	32	32	>128	>128
	16	16	32	32	>128	>128
NCL221	4	2	64	64	>128	>128
	4	2	64	64	>128	>128

NCL222	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL223	>128	>128	>128	>128	>128	>128
	>128	>128	>128	>128	>128	>128
NCL224	>128	>128	>128	>128	>128	>128
NCL225	>128	>128	>128	>128	>128	>128
NCL226	>128	>128	>128	>128	>128	>128
NCL227	>128	>128	>128	>128	>128	>128
NCL228	>128	>128	>128	>128	>128	>128
NCL229	>128	>128	>128	>128	>128	>128
NCL230	>128	>128	>128	>128	>128	>128

[00361] The NCL analogues showing the highest level of Gram-negative antibacterial activity included NCL030 (especially *Pseudomonas*), NCL041, NCL043, NCL044 (especially *Pseudomonas*), NCL052 (especially *Pseudomonas*), and NCL053 (especially *Pseudomonas*), NCL097 (especially *Pseudomonas*), NCL196 and NCL188 (especially *E. coli*).

[00362] The NCL analogues showing greatest activity against MRSA included: NCL021; NCL023; NCL029; NCL030; NCL035; NCL038; NCL039; NCL040; NCL041; NCL043; NCL044; NCL052; NCL054; NCL062; NCL069; NCL072; NCL073; NCL074; NCL078; NCL079; NCL080; NCL081; NCL082; NCL084; NCL088; NCL089; NCL093; NCL094; NCL097; NCL099; NCL101; NCL104; NCL107; NCL108; NCL111; NCL113; NCL117; NCL120; NCL121; NCL123; NCL136; NCL138; NCL140; NCL143; NCL146; NCL150; NCL151; NCL153; NCL155; NCL157; NCL158; NCL159; NCL160; NCL161; NCL166; NCL168; NCL169; NCL171; NCL172; NCL173; NCL174; NCL177; NCL178; NCL179; NCL180; NCL181; NCL182; NCL183; NCL184; NCL185; NCL186; NCL187; NCL188; NCL189; NCL190; NCL191; NCL192; NCL193; NCL195; NCL196; NCL197; NCL199; NCL201; NCL202; NCL203; NCL204; NCL205; NCL207; NCL215; NCL216; NCL217; NCL219; NCL220; and NCL221.

[00363] The NCL analogues showing greatest activity against VRE included: NCL011; NCL021; NCL023; NCL029; NCL030; NCL035; NCL038; NCL039; NCL040; NCL041; NCL043; NCL044; NCL052; NCL054; NCL061; NCL062; NCL069; NCL070; NCL072; NCL073; NCL074; NCL078; NCL079; NCL080; NCL081; NCL082; NCL084; NCL088; NCL093; NCL094; NCL097; NCL099; NCL101; NCL105; NCL107; NCL108; NCL111; NCL112; NCL113; NCL117; NCL120; NCL121; NCL123; NCL126; NCL131; NCL136; NCL140; NCL141; NCL143; NCL146; NCL151; NCL153; NCL155; NCL157; NCL158; NCL159; NCL160; NCL161; NCL166; NCL168; NCL169; NCL171; NCL173; NCL174; NCL175; NCL177; NCL178; NCL179; NCL180; NCL181; NCL182; NCL183; NCL184; NCL185; NCL186; NCL187; NCL188; NCL189; NCL190; NCL191; NCL192; NCL193; NCL195; NCL196; NCL197; NCL202; NCL203; NCL204; NCL205; NCL210; NCL212; NCL215; NCL216; NCL217; NCL218; NCL219; NCL220; and NCL221.

[00364] The bioassay ranking of the analogues tested is shown in Table 43.

Table 43: The bioassay ranking of the analogues tested according to Example 7.

NCL ID	G-Rank	NCL ID	M+R mcg/ml	M+R rank	NCL ID	MRSA mcg/ml	MRSA rank	NCL ID	VRE mcg/ml	VRE rank
NCL097	1	NCL062	2.25	1	NCL157	2	1	NCL089	0.75	1
NCL196	1	NCL157	2.25	2	NCL195	2	2	NCL062	1.5	2
NCL030	1	NCL812	2.375	3	NCL179	2	3	NCL812	1.75	3
NCL041	1	NCL195	2.75	4	NCL219	2	4	NCL084	2	4
NCL043	1	NCL216	2.75	5	NCL216	2.5	5	NCL157	2.5	5
NCL044	1	NCL089	2.875	6	NCL193	2.5	6	NCL081	2.5	6
NCL052	1	NCL084	3	7	NCL062	3	7	NCL216	3	7

NCL053	1	NCL179	3	8	NCL812	3	8	NCL217	3	8
		NCL054	3.5	9	NCL054	3	9	NCL079	3	9
		NCL217	3.5	10	NCL221	3	10	NCL080	3	10
		NCL081	3.75	11	NCL199	3.5	11	NCL078	3	11
		NCL193	3.75	12	NCL084	4	12	NCL195	3.5	12
		NCL079	4	13	NCL217	4	13	NCL082	3.5	13
		NCL080	4	14	NCL158	4	14	NCL074	3.75	14
		NCL158	4	15	NCL089	5	15	NCL179	4	15
		NCL082	5.25	16	NCL081	5	16	NCL054	4	16
		NCL078	5.5	17	NCL079	5	17	NCL158	4	17
		NCL120	6	18	NCL080	5	18	NCL088	4	18
		NCL088	7	19	NCL099	6	19	NCL123	4	19
		NCL099	7	20	NCL204	6	20	NCL215	4.5	20
		NCL177	7	21	NCL205	6	21	NCL193	5	21
		NCL215	7.25	22	NCL082	7	22	NCL120	5	22
		NCL197	7.5	23	NCL120	7	23	NCL035	5	23
		NCL040	8	24	NCL078	8	24	NCL140	5	24
		NCL038	9	25	NCL177	8	25	NCL177	6	25
		NCL202	9	26	NCL197	8	26	NCL197	7	26
		NCL219	9	27	NCL040	8	27	NCL099	8	27
		NCL123	9.5	28	NCL038	8	28	NCL040	8	28
		NCL074	9.875	29	NCL155	8	29	NCL202	8	29
		NCL155	10	30	NCL088	10	30	NCL021	8	30
		NCL035	10.5	31	NCL215	10	31	NCL143	8	31
		NCL021	12	32	NCL202	10	32	NCL153	8	32
		NCL113	12	33	NCL113	12	33	NCL038	10	33
		NCL143	12	34	NCL146	12	34	NCL155	12	34
		NCL146	14	35	NCL107	12	35	NCL113	12	35
		NCL153	16	36	NCL123	15	36	NCL219	16	36
		NCL188	16	37	NCL074	16	37	NCL146	16	37
		NCL173	17	38	NCL035	16	38	NCL188	16	38
		NCL023	20	39	NCL021	16	39	NCL173	18	39
		NCL220	24	40	NCL143	16	40	NCL174	18	40
		NCL174	25	41	NCL188	16	41	NCL039	20	41
		NCL190	28	42	NCL173	16	42	NCL023	24	42
		NCL039	30	43	NCL023	16	43	NCL212	24	43
		NCL121	30	44	NCL220	16	44	NCL121	28	44
		NCL029	32	45	NCL153	24	45	NCL220	32	45
		NCL221	33.5	46	NCL190	24	46	NCL190	32	46
		NCL168	40	47	NCL207	28	47	NCL029	32	47
		NCL169	40	48	NCL174	32	48	NCL169	32	48
		NCL203	44	49	NCL121	32	49	NCL180	32	49
		NCL107	46	50	NCL029	32	50	NCL184	32	50

EXAMPLE 8: The effects of NCL812 on antimicrobial sensitive isolates of *Staphylococcus aureus* and *Enterococcus faecalis*

Materials and Methods

Strain information

[00365] Two *Staphylococcus aureus* isolates were used in the following experiments; *S. aureus* MK01 a human skin strain, and *S. aureus* KC01 an equine skin strain. These isolates were identified by Gram stain and biochemical methods, including the Remel Staphaurex commercial kit. One *Enterococcus faecalis* isolate (USA01), was not identified as a VRE strain. As this isolate has previously been speciated, it was not subjected to further testing, except for observation of pure, characteristic growth on blood agar.

Investigation of Minimum Bactericidal Concentration (MBC)

CLSI methodology

[00366] As in previous experiments, 10 μ L of the contents of each well starting at the MIC was inoculated on to a Columbia SBA plate and incubated at 37 °C for 48 h. Plates were examined at 24 and 48 h and the MBC was recorded as the lowest concentration of NCL812 at which no colonies of bacteria were observed on the plate (or significant inhibition of growth was observed compared to the control) (CLSI 2005).

Kill kinetics assays for S. aureus KC01 & E. faecalis USA01 Method

[00367] *S. aureus* KC01 and *E. faecalis* USA01, not determined to be MRSA or VRE, respectively, were grown overnight on Columbia SBA at 37 °C. A few colonies of bacteria were then suspended in CAMHB (cation-adjusted Mueller Hinton broth) and adjusted to OD₆₀₀ of 0.08 to 0.10. The bacterial suspension was diluted 1:10. One millilitre of the bacteria were added to 9 mL of CAMHB containing various concentrations (up to 4×MIC) of NCL, to achieve a final bacterial concentration of 1 to 3×10⁶ CFU/mL. The tubes were incubated at 37 °C, with constant shaking. In order to determine the number of viable bacteria present at various time points, a 100 μ L aliquot was removed from each tube and diluted. Then, 100 μ L of each dilution were spread onto colony count agar, in duplicate, and incubated for 48 h at 37 °C. After 24 h the numbers of colonies present on each plate were counted and therefore the number of viable bacteria present in the original suspension enumerated. Plates were re-checked after 48 hours.

Results

Minimum Inhibitory Concentration (MIC)

[00368] The NCL812 MIC for isolates *S. aureus* MK01 and KC01, and *E. faecalis* USA01 was investigated. The results were: *S. aureus* MK01 = 4-8 μ g/mL, *S. aureus* KC01 = 2 μ g/mL, *E. faecalis* USA 01 = 4 μ g/mL.

[00369] *S. aureus* isolates MK01 and KC01 were investigated and no growth, or growth only at low concentrations of NCL812 (2 μ g/ml), was observed, indicating that NCL812 is

bactericidal against *S. aureus*. For the *E. faecalis* isolate tested (USA01) however, growth of bacteria was observed at all concentrations of NCL812 tested. There was an obvious reduction in the number of bacteria with increasing concentration, but growth was present compared with no growth for *S. aureus*. A summary of these results can be seen in **Table 45**. **Table 45** shows the results for NCL812 MBC tests on two non-MRSA *S. aureus* isolates and one non-VRE *E. faecalis* isolate. Each MBC test was performed in duplicate. No change in the results was observed at 48 h. **Table 37** shows NCL812 MBC values ($\mu\text{g/mL}$) for 20 MRSA isolates. Each MBC test was performed in duplicate starting from NCL812 MIC concentration to 16 times of MIC. **Table 38** shows NCL812 MBC values ($\mu\text{g/mL}$) for 10 VRE isolates. Each MBC test was performed in duplicate starting from NCL812 MIC concentration to 32 times the MIC.

Table 45: NCL812 MBC tests on two non-MRSA *Staphylococcus aureus* isolates and one non-VRE *Enterococcus faecalis* isolate according to Example 8.

Organism/Sample No.	NCL812 MBC						
	2 $\mu\text{g/mL}$	4 $\mu\text{g/mL}$	8 $\mu\text{g/mL}$	16 $\mu\text{g/mL}$	32 $\mu\text{g/mL}$	64 $\mu\text{g/mL}$	128 $\mu\text{g/mL}$
<i>S. aureus</i> (KC01)	1 st	+	0	0	0	0	0
	2 nd	+	+	+	0	0	0
<i>S. aureus</i> (MK01)	1 st	0 (5)	0	0	0 (N)	0 (N)	0 (N)
	2 nd	0	0	0	0	0	0
<i>E. faecalis</i> (USA01)	1 st	N	+ (488)	+	+	+ (7)	+
	2 nd	N	+	+	+	+	+

+ = Growth on Sheep Blood Agar; 0 = No Growth on Sheep Blood Agar; N = Not Cultured; Numbers in Parenthesis are the Number of Bacteria Growing after 24 hours per ml of sample (CFU/ml)

Kill kinetics assays for S. aureus KC01 & E. faecalis USA01 Method

[00370] Colony counts were performed at $t = 0, 120, 240$, and 360 min, then again at 24 h. At the 2 h time point *S. aureus* KC01 showed a minimum of a $2.5\log_{10}$ reduction in bacterial numbers from initial numbers, and greater than a $3\log_{10}$ reduction in comparison to the control at the same time point. A minimum of a $2\log_{10}$ reduction was still evident at 6 h incubation, however after 24 h the numbers of bacteria present had increased and this was not significantly different to the control.

[00371] Similar results were obtained with *E. faecalis* USA01, however the reduction in bacterial numbers observed was less than for *S. aureus* KC01. A $2\log_{10}$ reduction in CFU/mL was observed at 2 h, compared to the growth control. However, the reduction in CFU/mL compared to the original bacterial numbers was only just greater than $1\log_{10}$. At concentrations of 4 - 16 $\mu\text{g/mL}$ of NCL812 this reduction in bacterial numbers remained consistent until the 6 h time point. At concentrations of 32 and 64 $\mu\text{g/mL}$ however, there was approximately a $1\log_{10}$

rise in bacterial numbers over the same time period. At 24 h bacterial numbers at all concentrations had increased to almost the same level as the growth control.

[00372] The results observed with these strains of *S. aureus* and *E. faecalis* are consistent with the results observed for the kill kinetics assay for all MRSA and VRE isolates tested. The kill kinetics assay of *Staphylococcus aureus* KC01 at different concentrations of NCL812, up to 24 h incubation are shown in **Figure 38**. The kill kinetics assay of *Enterococcus faecalis* USA01 at different concentrations of NCL812, up to 24 h incubation are shown in **Figure 39**.

EXAMPLE 9: Formulations of Compounds

[00373] The following formulations were prepared using standard methods in the art.

Formulation A – Topical Formulation - PEG-based Gel with compounds of the invention

4.0g PEG 4000;
3.5g PEG 200;
0.6g propylene glycol;
1.9g water; and
0.204g of Compound (for example, NCL099)

[00374] PEG 4000, PEG 200 and propylene glycol were mixed and heated to 150 °C and until all solid crystals were dissolved. Compound was added to water and sonicated for 30 minutes until fully suspended. The Compound solution and gel solutions were mixed and allowed to cool and solidify. Formulation A will likely demonstrate acceptable viscosity, ease of skin application, uniform suspension and consistent and fine texture.

Formulation B – Topical Formulation - PEG-based Gel with compounds of the invention

3.0g PEG 4000;
1.0g PEG 8000;
3.0g PEG 200;
1.0g propylene glycol;
1.9g water; and
0.202g of Compound (for example, NCL099)

[00375] PEG 4000, PEG 8000, PEG 200 and propylene glycol were mixed and heated to 150 °C and until all solid crystals were dissolved. Compound (for example, NCL099) was added to water and sonicated for 30 minutes until fully suspended. The Compound solution and gel solutions were mixed and allowed to cool and solidify. Formulation B demonstrated acceptable viscosity, ease of skin application, uniform suspension and consistent and fine texture.

Formulation C – Topical Formulation - PEG-based Gel with Compound-Soluplus

2.5g PEG 4000;
4.0g PEG 200;
2.5g propylene glycol;
1.0g water; and
1.8g solid dispersion of Compound-SoluPlus.

[00376] Soluplus was purchased from BASF (www.soluplus.com). Compound-SoluPlus was prepared using standard methods in the art. PEG 4000, PEG 200, Compound-SoluPlus and propylene glycol were mixed and heated to 150 °C and until all solid crystals were dissolve. Water was added and then the solution was sonicated. The solution was allowed to cool and solidify. Formulation C demonstrated acceptable viscosity, ease of skin application, uniform suspension and consistent and fine texture.

Formulation D – Tablet Formulation

30mg Calcium hydrogen phosphate dehydrate;
80mg Microcrystalline cellulose;
50mg Lactose;
8mg Hydroxypropyl methyl cellulose
1.5mg Talc
10mg of compound (for example NCL099)

[00377] The excipients were weighed and mixed for 5 minutes. The mixture was fed into a feed hopper of a tablet press machine and the machine was operated according to standard procedures in the art. Formulation D demonstrated acceptable tablet hardness, disintegration and friability.

Formulation E – Oral Suspension

2.0 ml Glycerol;
1.5ml Absolute ethanol;
600mg NCL812; and
To 60ml Vehicle (Ora Sweet and Ora Plus, 1:1).

[00378] NCL 812 powder was sieved through a 75 µm sieve. 600 mg of sieved NCL 812 was mixed with 2.0 ml glycerol and 1.5 ml absolute ethanol. The mixture was placed in a mortar and manually milled until all NCL 812 was suspended uniformly. The suspension was sonicated for 30 minutes. Vehicle (55 ml of Ora Sweet and Ora Plus mixture) was then added to the suspension and milled for another 10 minutes. Volume was made up with the Ora plus and Ora sweet mixture to 60 ml by transferring to a measuring cylinder

[00379] Formulation E demonstrated acceptable suspension and demonstrated acceptable short term stability.

Formulation F – Intramuscular Injection

20mg/ml	Polyvinylpyrrolidone K30 (PVPK30);
0.09mg/ml	NCL812; and
50ml	water.

[00380] Two percent of w/v PVP K30 solution was prepared by the addition of 1.0 g of PVP K30 to 50 ml of MilliQ water. The solution was then placed in a sonicator for 30 minutes to equilibrate and 4.5 mg of NCL 812 was added to the PVP solution and placed on an incubator shaker at a maximum speed of 10 rpm over a period of 24 hours, with controlled temperature of $25\pm1^{\circ}\text{C}$. Solution was transferred to 5 ml vials and checked for clarity, appearance, pH and short-term stability. The pH of solution was 7.25.

[00381] Formulation F demonstrated acceptable transparency and short term stability.

EXAMPLE 10: Release of NCL812 and NCL099 from Formulation B.

[00382] The objective of this study was to measure the release of NCL812 and NCL099 from Formulation B prepared in Example 9.

[00383] Franz diffusion cells were utilized to quantify the release rate of NCL 812 and NCL099 from its topical formulations. Five millilitres of absolute ethanol, which was chosen as the desired release medium, was loaded into the receptor chamber. Temperature of the receptor fluid was kept constant, at $32\pm1^{\circ}\text{C}$ using a water jacket. Acetyl cellulose membranes, with pore size of $0.45\ \mu\text{m}$ (Pall Corporation) was selected and placed between donor and receptor chamber. Followed by that, a number of test samples (Formulation B) was loaded into the donor chamber. One millilitre of receptor fluid was collected at regular time intervals of 0.25, 0.50, 0.75, 1, 2, 3, 4, 5, 6, 7, 8 and 24 hours through the sampling port. One millilitre of fresh absolute ethanol was immediately returned to the receptor chamber. UV-HPLC was utilized to analyse the content of the receptor fluids attained.

[00384] Figure 40 presents the cumulative release of NCL812 and NCL099 over time. This study demonstrates that Formulation B provides an acceptable release profile for NCL812 and NCL099.

EXAMPLE 11: NMR Spectroscopy Lists of Compounds NCL812, NCL001-NCL230

[00385] NMR Spectroscopy was performed on compounds NCL812, NCL001-NCL230 using standard methods in the art. The lists of the NMR spectroscopy are presented in **Table 46**.

Table 46: NMR Spectroscopy Lists of Compounds NCL812, NCL001-NCL230

NCL Code	NMR
NCL812	¹ H NMR (400 MHz, DMSO) δ 12.04 (br. s, 2H), 8.48 (br. s, 1H), 8.37 (br. s, 2H), 7.97 (d, J = 8.6 Hz, 4H), 7.57 (d, J = 8.6 Hz, 4H)
NCL001	¹ H NMR (400 MHz, DMSO) δ 10.84 (br. s, 2H), 8.17 (br. s, 2H), 7.77 (d, J = 8.2 Hz, 4H), 7.50 (d, J = 8.6 Hz, 4H)
NCL002	¹ H NMR (400 MHz, DMSO) δ 11.06 (s, 2H), 8.58 (br. s, 2H), 8.17 (br. s, 2H), 7.50 - 7.52 (m, 2H), 7.41 - 7.45 (m, 4H)
NCL003	¹ H NMR (400 MHz, DMSO) δ 10.71 (s, 2H), 8.17 (br. s, 2H), 7.73 - 7.88 (m, 4H), 7.28 (t, J = 8.8 Hz, 4H)
NCL004	¹ H NMR (400 MHz, DMSO) δ 10.89 (br. s, 2H), 8.19 (br. s, 2H), 7.65 (br. s, 2H), 7.43 - 7.56 (m, 4H), 7.19 - 7.27 (m, 2H)
NCL005	¹ H NMR (400 MHz, DMSO) δ 10.94 (br. s, 2H), 8.43 (br. s, 2H), 8.10 (br. s, 2H), 7.39 - 7.52 (m, 2H), 7.21 - 7.35 (m, 4H)
NCL006	¹ H NMR (400 MHz, DMSO) δ 10.55 (s, 2H), 8.11 (br. s, 2H), 7.88 (d, J = 8.6 Hz, 4H), 8.99 (d, J = 8.6 Hz, 4H), 3.80 (s, 6H)
NCL007	¹ H NMR (400 MHz, DMSO) δ 11.10 (br. s, 2H), 8.24 (br. s, 2H), 7.81 - 8.03 (m, 8H)
NCL008	¹ H NMR (400 MHz, DMSO) δ 11.24 (br. s, 2H), 8.51 (br. s, 2H), 8.18 - 8.29 (m, 2H), 7.90 (d, J = 7.4 Hz, 2H), 7.80 (t, J = 7.6 Hz, 2H), 7.59 (t, J = 7.0 Hz, 2H)
NCL009	¹ H NMR (400 MHz, DMSO) δ 11.02 (s, 2H), 8.26 (br. s, 4H), 8.07 (d, J = 7.8 Hz, 2H), 7.85 (d, J = 7.8 Hz, 2H), 7.65 (t, J = 7.8 Hz, 2H)
NCL010	¹ H NMR (400 MHz, DMSO) δ 10.74 (br. s, 2H), 8.15 (br. s, 2H), 7.25 - 7.39 (m, 6H), 6.94-7.01 (m, 2H), 3.82 (s, 6H)
NCL011	¹ H NMR (400 MHz, DMSO) δ 11.02 (s, 2H), 8.28 (br. s, 2H), 8.13 (s, 2H), 8.04 (d, J = 7.4 Hz, 2H), 7.75 (d, J = 8.0 Hz, 2H), 7.68 (t, J = 8.0 Hz, 2H)
NCL012	¹ H NMR (400 MHz, DMSO) δ 11.04 (br. s, 2H), 8.27 (br. s, 2H), 7.97 (d, J = 7.8 Hz, 4H), 7.80 (d, J = 8.2 Hz, 4H)
NCL013	¹ H NMR (400 MHz, DMSO) δ 11.22 (br. s, 2H), 8.55 (br. s, 2H), 8.35 (d, J = 7.0 Hz, 2H), 7.73 - 7.82 (m, 4H), 7.57 - 7.65 (m, 2H)
NCL014	¹ H NMR (400 MHz, DMSO) δ 10.30 (s, 1H), 7.81 (s, 1H), 7.76 (d, J = 8.6 Hz, 2H), 7.43 (d, J = 8.6 Hz, 2H), 6.53 (br. s, 2H)
NCL015	¹ H NMR (400 MHz, DMSO) δ 12.02 (br. s, 1H), 8.55 (s, 1H), 8.27-8.33 (m, 1H), 7.79 (br. s, 3H), 7.51 - 7.56 (m, 1H), 7.39 - 7.51 (m, 2H)
NCL016	¹ H NMR (400 MHz, DMSO) δ 11.98 (br. s, 1H), 8.39 (s, 1H), 8.19-8.26 (m, 1H), 7.80 (br. s, 3H), 7.46 - 7.53 (m, 1H), 7.20 - 7.38 (m, 2H)
NCL017	¹ H NMR (400 MHz, DMSO) δ 11.79 (br. s, 1H), 8.17 (s, 1H), 7.87 (d, J = 9.0 Hz, 1H), 7.71 (br. s, 3H), 7.62 (d, J = 7.4 Hz, 1H), 7.45 - 7.54 (m, 1H), 7.25-7.32 (m, 1H)
NCL018	¹ H NMR (400 MHz, DMSO) δ 10.66 (s, 2H), 8.47 (br. s, 2H), 7.91-8.00 (m, 2H), 7.19 - 7.32 (m, 6H), 2.42 (s, 6H)
NCL019	¹ H NMR (400 MHz, DMSO) δ 10.66 (br. s, 2H), 8.15 (br. s, 2H), 7.57 (s, 2H), 7.52 (d, J = 7.4 Hz, 2H), 7.32 (t, J = 7.6 Hz, 2H), 7.21 (d, J = 7.4 Hz, 2H), 2.38 (s, 6H)
NCL020	¹ H NMR (400 MHz, DMSO) δ 12.37 (br. s, 2H), 8.83 (br. s, 2H), 8.63 (br. s, 2H), 8.39 - 8.44 (m, 2H), 7.55 - 7.60 (m, 2H), 7.44 - 7.55 (m, 4H)
NCL021	¹ H NMR (400 MHz, DMSO) δ 12.11 (br. s, 1H), 8.52 (br. s, 2H), 8.02 (t, J = 8.6 Hz, 4H), 7.35 (t, J = 8.8 Hz, 4H)
NCL022	¹ H NMR (400 MHz, DMSO) δ 12.19 (br. s, 2H), 8.85 (br. s, 2H), 8.58 (br. s, 1H), 8.34 (t, J = 7.6 Hz, 2H), 7.51 - 7.60 (m, 2H), 7.34 (t, J = 8.2 Hz, 4H)
NCL023	¹ H NMR (400 MHz, DMSO) δ 12.08 (br. s, 2H), 8.36 (br. s, 2H), 7.92-8.00 (m, 2H), 7.66-7.71 (m, 2H), 7.50 - 7.56 (m, 2H), 7.29 - 7.37 (m, 2H)

NCL024	¹ H NMR (400 MHz, DMSO) δ 12.32 (br. s, 2H), 8.67 (br. s, 2H), 8.44 (br. s, 2H), 8.15 (d, J = 8.6 Hz, 4H), 7.98 (d, J = 8.6 Hz, 4H)
NCL025	¹ H NMR (400 MHz, DMSO) δ 8.75 (br. s, 2H), 8.50 (d, J = 8.2 Hz, 2H), 7.97 (d, J = 7.4 Hz, 2H), 7.85 (t, J = 7.6 Hz, 2H), 7.68 (t, J = 7.6 Hz, 2H)
NCL026	¹ H NMR (400 MHz, DMSO) δ 12.26 (br. s, 1H), 8.66 (br. s, 1H), 8.55 (s, 2H), 8.43 (br. s, 2H), 8.21 (d, J = 7.8 Hz, 2H), 7.94 (d, J = 7.8 Hz, 2H), 7.71 (t, J = 7.8 Hz, 2H)
NCL027	¹ H NMR (400 MHz, DMSO) δ 11.78 (br. s, 2H), 8.31 (br. s, 3H), 7.87 (d, J = 8.6 Hz, 4H), 7.04 (d, J = 8.6 Hz, 4H), 3.83 (s, 6H)
NCL028	¹ H NMR (400 MHz, DMSO) δ 12.00 (br. s, 2H), 8.75 (br. s, 2H), 8.39 (br. s, 2H), 8.22 (d, J = 6.7 Hz, 2H), 7.44 - 7.52 (m, 2H), 7.14 (d, J = 8.2 Hz, 2H), 7.05 (t, J = 7.6 Hz, 2H), 3.89 (s, 6H)
NCL029	¹ H NMR (400 MHz, DMSO) δ 11.98 (br. s, 2H), 8.48 (br. s, 2H), 8.36 (br. s, 2H), 7.56 (s, 2H), 7.35 - 7.49 (m, 4H), 7.04-7.10 (m, 2H), 3.84 (s, 6H)
NCL030	¹ H NMR (400 MHz, DMSO) δ 11.83 (br. s, 1H), 8.16 (s, 1H), 7.91 (d, J = 8.2 Hz, 2H), 7.75 (br. s, 1H), 7.53 (d, J = 8.2 Hz, 2H)
NCL031	¹ H NMR (400 MHz, DMSO) δ 11.91 (br. s, 1H), 8.22 (s, 1H), 8.09 (d, J = 8.2 Hz, 2H), 7.93 (d, J = 8.2 Hz, 2H)
NCL032	¹ H NMR (400 MHz, DMSO) δ 12.12 (s, 1H), 8.48 (s, 1H), 8.38 (d, J = 7.8 Hz, 1H), 7.94 (d, J = 7.8 Hz, 1H), 7.86 (br. s, 2H), 7.80 (t, J = 7.8 Hz, 2H), 7.64 (t, J = 7.6 Hz, 1H)
NCL033	¹ H NMR (400 MHz, DMSO) δ 11.93 (s, 1H), 8.50 (s, 1H), 8.20 (s, 1H), 8.14 (d, J = 7.8 Hz, 1H), 7.90 (d, J = 7.8 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H)
NCL034	¹ H NMR (400 MHz, DMSO) δ 11.87 (br. s, 1H), 8.48 (s, 1H), 8.09 (d, J = 7.8 Hz, 1H), 7.70 (br. s, 2H), 7.38 - 7.49 (m, 1H), 7.11 (d, J = 8.6 Hz, 1H), 7.01 (t, J = 7.4 Hz, 1H), 3.86 (s, 3H)
NCL035	¹ H NMR (400 MHz, DMSO) δ 12.32 (br. s, 2H), 8.69 (br. s, 2H), 8.49 (br. s, 2H), 8.18 (d, J = 7.8 Hz, 4H), 7.86 (d, J = 8.2 Hz, 4H)
NCL036	¹ H NMR (400 MHz, DMSO) δ 12.51 (br. s, 1H), 8.80 (br. s, 2H), 8.72 (br. s, 1H), 8.59 (d, J = 7.8 Hz, 2H), 7.78 - 7.91 (m, 4H), 7.71 (t, J = 8.0 Hz, 2H)
NCL037	¹ H NMR (400 MHz, DMSO) δ 12.28 (br. s, 2H), 8.70 (br. s, 2H), 8.50 (br. s, 2H), 8.38 (s, 2H), 8.22 (d, J = 7.8 Hz, 2H), 7.85 (d, J = 7.8 Hz, 2H), 7.74 (t, J = 7.8 Hz, 2H)
NCL038	¹ H NMR (400 MHz, DMSO) δ 11.92 (br. s, 2H), 8.41 (br. s, 2H), 8.36 (br. s, 2H), 7.83 (d, J = 8.2 Hz, 4H), 7.31 (d, J = 7.8 Hz, 4H), 2.37 (s, 6H)
NCL039	¹ H NMR (400 MHz, DMSO) δ 11.99 (br. s, 2H), 8.73 (br. s, 2H), 8.41 (br. s, 2H), 8.19 (d, J = 7.8 Hz, 2H), 7.37 (t, J = 8.0 Hz, 2H), 7.30 (t, J = 7.8 Hz, 4H), 2.46 (s, 6H)
NCL040	¹ H NMR (400 MHz, DMSO) δ 11.97 (br. s, 2H), 8.44 (br. s, 2H), 8.37 (br. s, 2H), 7.71 (d, J = 7.8 Hz, 2H), 7.38 (t, J = 7.8 Hz, 2H), 7.31 (d, J = 7.8 Hz, 2H), 2.38 (s, 6H)
NCL041	¹ H NMR (400 MHz, DMSO) δ 11.94 (s, 1H), 8.25 (s, 1H), 8.11 (d, J = 7.8 Hz, 2H), 7.71 - 7.91 (m, 4H)
NCL042	¹ H NMR (400 MHz, DMSO) δ 12.04 (s, 1H), 8.46 - 8.56 (m, 2H), 7.70 - 7.93 (m, 5H), 7.66 (t, J = 7.8 Hz, 1H)
NCL043	¹ H NMR (400 MHz, DMSO) δ 11.88 (s, 1H), 8.33 (s, 1H), 8.26 (s, 1H), 8.14 (d, J = 7.8 Hz, 1H), 7.80 (d, J = 7.8 Hz, 1H), 7.69 (t, J = 7.8 Hz, 1H)
NCL044	¹ H NMR (400 MHz, DMSO) δ 11.71 (br. s, 1H), 8.13 (s, 1H), 7.76 (d, J = 8.2 Hz, 2H), 7.27 (d, J = 7.8 Hz, 2H), 2.35 (s, 3H)
NCL045	¹ H NMR (400 MHz, DMSO) δ 11.69 (br. s, 1H), 8.45 (s, 1H), 8.06 (d, J = 7.4 Hz, 1H), 7.67 (br. s, 2H), 7.30 - 7.39 (m, 1H), 7.20 - 7.29 (m, 2H), 2.42 (s, 3H)
NCL046	¹ H NMR (400 MHz, DMSO) δ 11.64 (br. s, 1H), 8.12 (s, 1H), 7.53 - 7.77 (m, 4H), 7.34 (t, J = 7.8 Hz, 1H), 7.27 (d, J = 7.8 Hz, 1H), 2.35 (s, 3H)
NCL047	¹ H NMR (400 MHz, DMSO) δ 10.47 (s, 1H), 8.23 (s, 1H), 8.16 - 8.21 (m, 1H), 7.42 - 7.50 (m, 1H), 7.30 - 7.40 (m, 2H), 6.57 (br. s, 2H)
NCL048	¹ H NMR (400 MHz, DMSO) δ 10.40 (s, 1H), 8.08 - 8.15 (m, 1H), 8.05 (s, 1H), 7.34 - 7.44 (m, 1H), 7.17 - 7.28 (m, 2H), 6.54 (br. s, 2H)
NCL049	¹ H NMR (400 MHz, DMSO) δ 10.51 (s, 1H), 7.93 (d, J = 8.2 Hz, 1H), 7.86 (s, 1H), 7.83 (d, J = 8.2 Hz, 1H), 6.63 (br. s, 2H)

NCL050	¹ H NMR (400 MHz, DMSO) δ 10.86 (s, 1H), 8.13 (d, J = 8.0 Hz, 1H), 8.11 (s, 1H), 7.85 (d, J = 7.4 Hz, 1H), 7.71 (t, J = 7.8 Hz, 1H), 7.52 (t, J = 7.0 Hz, 1H), 6.60 (br, s, 2H)
NCL051	¹ H NMR (400 MHz, DMSO) δ 10.43 (s, 1H), 8.37 (s, 1H), 7.98 (d, J = 7.8 Hz, 1H), 7.84 (s, 1H), 7.77 (d, J = 7.8 Hz, 1H), 7.57 (t, J = 7.8 Hz, 1H)
NCL052	¹ H NMR (400 MHz, DMSO) δ 11.95 (s, 1H), 8.16 (s, 1H), 8.10 (s, 1H), 7.80 (br, s, 2H), 7.76 (d, J = 7.0 Hz, 1H), 7.40 - 7.55 (m, 2H)
NCL053	¹ H NMR (400 MHz, DMSO) δ 11.88 (s, 1H), 8.17 (s, 1H), 7.95 (dd, J = 8.8, 5.7 Hz, 2H), 7.76 (br, s, 1H), 7.30 (t, J = 9.0 Hz, 2H)
NCL054	¹ H NMR (400 MHz, DMSO) δ 12.17 (br, s, 2H), 8.61 (br, s, 2H), 8.39 (br, s, 2H), 8.16 (s, 2H), 7.83 (d, J = 7.0 Hz, 2H), 7.45 - 7.61 (m, 4H)
NCL055	¹ H NMR (400 MHz, DMSO) δ 10.34 (s, 1H), 7.93 (s, 1H), 7.80 (s, 1H), 7.54 - 7.67 (m, 1H), 7.32 - 7.46 (m, 2H), 6.56 (br, s, 1H)
NCL056	¹ H NMR (400 MHz, DMSO) δ 10.23 (s, 1H), 7.82 (s, 1H), 7.78 (dd, J = 8.8, 5.7 Hz, 2H), 7.21 (t, J = 8.8 Hz, 1H), 6.49 (br, s, 2H)
NCL057	¹ H NMR (400 MHz, DMSO) δ 10.33 (s, 1H), 7.82 (s, 1H), 7.71 (d, J = 9.8 Hz, 1H), 7.35 - 7.53 (m, 2H), 7.10 - 7.23 (m, 1H), 6.57 (br, s, 1H)
NCL058	¹ H NMR (400 MHz, DMSO) δ 10.08 (s, 1H), 7.78 (s, 1H), 7.65 (d, J = 9.0 Hz, 2H), 6.94 (d, J = 8.6 Hz, 2H), 6.40 (br, s, 2H), 3.78 (s, 3H)
NCL059	¹ H NMR (400 MHz, DMSO) δ 10.23 (s, 1H), 7.80 (s, 1H), 7.34 (s, 1H), 7.29 (t, J = 7.8 Hz, 1H), 7.21 (d, J = 7.8 Hz, 1H), 6.91 (dd, J = 7.8, 2.0 Hz, 1H), 6.51 (br, s, 2H), 3.79 (s, 3H)
NCL060	¹ H NMR (400 MHz, DMSO) δ 10.56 (s, 1H), 8.41 (d, J = 7.6 Hz, 1H), 8.19 (br, s, 1H), 7.74 (d, J = 7.8 Hz, 1H), 7.67 (t, J = 7.6 Hz, 1H), 7.54 (t, J = 7.8 Hz, 1H), 6.82 (br, s, 2H)
NCL061	¹ H NMR (400 MHz, DMSO) δ 11.71 (br, s, 2H), 8.91 (br, s, 1H), 8.28 (d, J = 7.8 Hz, 4H), 7.82 (d, J = 8.2 Hz, 4H), 2.49 (br, s, 6H)
NCL062	¹ H NMR (400 MHz, DMSO) δ 11.68 (br, s, 2H), 8.78 (br, s, 1H), 8.10 (d, J = 8.6 Hz, 4H), 7.52 (d, J = 8.6 Hz, 4H), 2.43 (s, 6H)
NCL063	¹ H NMR (400 MHz, DMSO) δ 7.99 (br, s, 4H), 7.83 (d, J = 7.6 Hz, 6H)
NCL064	¹ H NMR (400 MHz, DMSO) δ 7.94 (d, J = 7.6 Hz, 4H), 7.84 (t, J = 7.6 Hz, 4H), 7.64 (t, J = 7.6 Hz, 4H)
NCL065	¹ H NMR (400 MHz, DMSO) δ 12.20 (br, s, 1H), 11.86 (br, s, 1H), 8.65 (br, s, 1H), 8.50 (br, s, 1H), 8.03-8.27 (m, 3H), 7.90 (d, J = 7.8 Hz, 2H), 7.68 (t, J = 7.6 Hz, 2H)
NCL066	¹ H NMR (400 MHz, DMSO) δ 7.43 - 7.67 (m, 7H), 7.23 - 7.34 (m, 2H)
NCL067	¹ H NMR (400 MHz, DMSO) δ 10.40 (s, 1H), 8.16 (s, 1H), 7.98 (d, J = 7.4 Hz, 1H), 7.90 (s, 1H), 7.67 (d, J = 7.3 Hz, 1H), 7.60 (t, J = 7.4 Hz, 1H)
NCL068	¹ H NMR (400 MHz, DMSO) δ 8.47 (br, s, 1H), 8.18 (d, J = 7.8 Hz, 3H), 8.07 (d, J = 8.6 Hz, 2H), 7.85 (d, J = 7.8 Hz, 2H), 7.53 (d, J = 8.6 Hz, 2H), 2.40 (s, 3H)
NCL069	¹ H NMR (400 MHz, DMSO) δ 9.44 (br, s, 1H), 8.03 (d, J = 8.2 Hz, 2H), 7.94 (br, s, 2H), 7.46 (d, J = 8.6 Hz, 2H), 2.31 (s, 3H)
NCL070	¹ H NMR (400 MHz, DMSO) δ 8.12 (br, s, 1H), 7.71 (br, s, 3H), 7.49 (br, s, 6H)
NCL071	¹ H NMR (400 MHz, DMSO) δ 11.43 (s, 1H), 8.40 (s, 1H), 8.16 (br, s, 1H), 8.10 (d, J = 7.4 Hz, 1H), 7.95 (br, s, 1H), 7.57 (t, J = 7.8 Hz, 1H), 7.06 (d, J = 8.2 Hz, 1H), 6.95 (t, J = 7.4 Hz, 1H), 3.82 (s, 3H)
NCL072	¹ H NMR (400 MHz, DMSO) δ 7.91 (br, d, J = 8.2 Hz, 3H), 7.52 (d, J = 8.6 Hz, 2H)
NCL073	¹ H NMR (400 MHz, DMSO) δ 9.51 (br, s, 1H), 8.22 (d, J = 8.3 Hz, 2H), 8.01 (br, s, 2H), 7.77 (d, J = 8.3 Hz, 2H), 2.36 (s, 3H)
NCL074	¹ H NMR (400 MHz, DMSO) δ 12.46 (br, s, 1H), 8.79 (br, s, 1H), 8.66 (br, s, 2H), 8.46 (d, J = 8.2 Hz, 2H), 7.99 (d, J = 7.0 Hz, 2H), 7.77 (br, s, 1H), 7.58 (d, J = 6.7 Hz, 3H)
NCL075	¹ H NMR (400 MHz, DMSO) δ 8.48 (br, s, 1H), 8.26 (d, J = 7.4 Hz, 2H), 8.19 (d, J = 7.4 Hz, 2H), 7.75-7.93 (m, 4H), 2.46 (s, 3H)

NCL076	¹ H NMR (400 MHz, DMSO) δ 8.53 (br. s, 2H), 8.29 - 8.46 (m, 2H), 7.99 (d, J = 7.8 Hz, 2H), 7.83 (d, J = 7.8 Hz, 2H), 7.57 (d, J = 7.8 Hz, 2H), 7.31 (d, J = 7.8 Hz, 2H), 2.37 (s, 3H)
NCL077	¹ H NMR (400 MHz, DMSO) δ 12.28 (br. s, 1H), 8.67 (br. s, 1H), 8.49 (br. s, 1H), 8.42 (br. s, 1H), 8.18 (d, J = 7.8 Hz, 2H), 8.00 (d, J = 8.2 Hz, 2H), 7.86 (d, J = 8.2 Hz, 2H), 7.58 (d, J = 7.8 Hz, 2H)
NCL078	¹ H NMR (400 MHz, DMSO) δ 12.33 (br. s, 1H), 8.74 (br. s, 2H), 8.58 (t, J = 7.6 Hz, 1H), 8.42 (br. s, 1H), 8.00 (d, J = 8.2 Hz, 2H), 7.87 (d, J = 10.2 Hz, 1H), 7.74 (d, J = 8.2 Hz, 1H), 7.58 (d, J = 8.2 Hz, 2H)
NCL079	¹ H NMR (400 MHz, DMSO) δ 12.08 (br. s, 1H), 8.54 (br. s, 1H), 8.39 (br. s, 2H), 7.93 - 8.09 (m, 4H), 7.57 (d, J = 8.2 Hz, 2H), 7.35 (t, J = 8.6 Hz, 2H)
NCL080	¹ H NMR (400 MHz, DMSO) δ 12.04 (br. s, 1H), 11.36 (br. s, 1H), 8.40 (br. s, 1H), 8.25 (d, J = 8.2 Hz, 2H), 8.00 (d, J = 7.0 Hz, 2H), 7.82 (d, J = 7.4 Hz, 2H), 7.57 (d, J = 6.7 Hz, 2H), 2.44 (s, 3H)
NCL081	¹ H NMR (400 MHz, DMSO) δ 8.39 (br. s, 1H), 8.07 (d, J = 8.6 Hz, 2H), 7.99 (d, J = 8.6 Hz, 2H), 7.57 (d, J = 8.2 Hz, 2H), 7.53 (d, J = 8.6 Hz, 2H), 2.39 (s, 3H)
NCL082	¹ H NMR (400 MHz, DMSO) δ 12.40 (br. s, 1H), 8.84 (br. s, 1H), 8.63 (br. s, 2H), 8.42 (d, J = 7.8 Hz, 2H), 7.99 (d, J = 8.6 Hz, 2H), 7.58 (d, J = 8.2 Hz, 3H), 7.43 - 7.54 (m, 2H)
NCL083	¹ H NMR (400 MHz, DMSO) δ 12.30 (br. s, 1H), 8.64 (br. s, 2H), 8.41 (br. s, 2H), 8.17 (s, 1H), 7.99 (d, J = 8.6 Hz, 2H), 7.82 (d, J = 6.7 Hz, 1H), 7.45 - 7.64 (m, 4H)
NCL084	¹ H NMR (400 MHz, DMSO) δ 12.38 (br. s, 1H), 8.64 (br. s, 2H), 8.30 - 8.50 (m, 2H), 7.99 (d, J = 8.6 Hz, 2H), 7.62 (d, J = 10.4, 1.4 Hz, 1H), 7.57 (d, J = 8.6 Hz, 2H), 7.46 (d, J = 8.6 Hz, 1H)
NCL085	¹ H NMR (400 MHz, DMSO) δ 8.77 (br. s, 1H), 8.68 (br. s, 2H), 8.52 (d, J = 8.2 Hz, 1H), 8.46 (br. s, 1H), 7.92 - 8.06 (m, 3H), 7.84 (t, J = 7.6 Hz, 1H), 7.68 (t, J = 7.6 Hz, 1H), 7.58 (d, J = 8.2 Hz, 2H)
NCL086	¹ H NMR (400 MHz, DMSO) δ 12.39 (br. s, 1H), 8.68 (br. s, 2H), 8.57 (s, 1H), 8.43 (br. s, 2H), 8.21 (d, J = 7.4 Hz, 1H), 8.00 (d, J = 8.2 Hz, 2H), 7.94 (d, J = 7.0 Hz, 1H), 7.70 (t, J = 7.2 Hz, 1H), 7.58 (d, J = 7.8 Hz, 2H)
NCL087	¹ H NMR (400 MHz, DMSO) δ 8.71 (br. s, 2H), 8.48 (br. s, 1H), 8.43 (br. s, 1H), 8.16 (d, J = 8.2 Hz, 2H), 7.99 (t, J = 8.0 Hz, 4H), 7.58 (d, J = 8.6 Hz, 2H)
NCL088	¹ H NMR (400 MHz, DMSO) δ 12.33 (br. s, 1H), 8.68 (br. s, 1H), 8.61 (br. s, 2H), 8.42 (br. s, 1H), 8.35 (t, J = 7.4 Hz, 1H), 7.99 (d, J = 8.2 Hz, 2H), 7.49 - 7.64 (m, 3H), 7.27 - 7.41 (m, 2H)
NCL089	¹ H NMR (400 MHz, DMSO) δ 11.45 (br. s, 1H), 8.76 - 8.94 (m, 1H), 8.11 (d, J = 8.0 Hz, 4H), 7.53 (d, J = 8.0 Hz, 4H), 2.42 (br. s, 6H)
NCL090	¹ H NMR (400 MHz, DMSO) δ 10.60 (br. s, 1H), 8.68 (s, 1H), 8.09 (br. s, 1H), 7.95 (br. s, 4H), 7.32 - 7.71 (m, 10H)
NCL091	¹ H NMR (400 MHz, DMSO) δ 9.61 (br. s, 1H), 8.19 (d, J = 6.3 Hz, 2H), 7.90 (br. s, 2H), 7.56 - 7.73 (m, 2H), 7.26 - 7.40 (m, 2H)
NCL092	¹ H NMR (400 MHz, DMSO) δ 12.38 (br. s, 1H), 7.82 - 7.94 (m, 2H), 7.18 - 7.63 (m, 7H), 4.39 (br. s, 2H)
NCL 093	¹ H NMR (300 MHz, DMSO) δ 8.64 (s, 4H), 8.35 - 8.24 (m, 4H), 8.06 - 7.93 (m, 6H), 7.64 - 7.54 (m, 4H).
NCL 094	¹ H NMR (300 MHz, CDCl ₃) δ 7.16 (d, J = 4.7 Hz, 2H), 6.10 (br. s, 3H), 2.27 - 2.14 (m, 2H), 1.84 - 1.61 (m, 10H), 1.37 - 1.13 (m, 10H).
NCL 095	¹ H NMR (300 MHz, DMSO) δ 8.39 (s, 2H), 8.36 - 8.11 (m, 4H), 7.78 (s, 2H), 7.12 (d, J = 1.4 Hz, 2H).
NCL 096	¹ H NMR (300 MHz, MeOD) δ 8.08 (d, J = 8.2 Hz, 2H), 7.61 - 7.53 (m, 4H), 7.45 - 7.30 (m, 6H), 7.17 - 6.97 (m, 4H).
NCL 097	¹ H NMR (300 MHz, DMSO) δ 9.06 (br. s, 6H), 8.25 - 8.01 (m, 4H), 6.83 (s, 4H). ¹³ C NMR (75 MHz, DMSO) δ 152.2, 149.7, 146.2, 136.5, 123.7, 107.4.

NCL 098	¹ H NMR (300 MHz, DMSO) δ 8.65 (s, 2H), 8.53 (s, 2H), 8.40 (s, 2H), 8.24 (d, J = 7.3 Hz, 2H), 8.09 – 7.98 (m, 2H), 7.62 (t, J = 7.7 Hz, 2H).
NCL 099	¹ H NMR (300 MHz, DMSO) δ 8.56 – 8.32 (m, 4H), 7.85 (d, J = 8.3 Hz, 4H), 7.49 (d, J = 8.3 Hz, 4H), 1.31 (s, 18H). ¹³ C NMR (75 MHz, DMSO) δ 153.7, 152.7, 148.8, 130.7, 127.8, 125.6, 34.7, 31.0.
NCL 100	¹ H NMR (300 MHz, DMSO) δ 12.39 (br s, 2H), 8.55 (s, 2H), 8.46 (s, 2H), 8.01 – 7.88 (m, 4H), 7.55 – 7.41 (m, 6H).
NCL 101	¹ H NMR (300 MHz, DMSO) δ 12.06 (br s, 2H), 9.71 (br s, 2H), 8.70 (s, 2H), 8.30 (s, 2H), 7.50 (d, J = 7.9 Hz, 2H), 6.90 (d, J = 7.7 Hz, 2H), 6.70 (t, J = 7.7 Hz, 2H).
NCL 102	¹ H NMR (300 MHz, DMSO) δ 12.86 (br s, 2H), 8.89 (s, 2H), 8.77 (s, 2H), 8.52 (d, J = 7.9 Hz, 2H), 8.11 (d, J = 8.1 Hz, 2H), 7.91 – 7.76 (m, 2H), 7.77 – 7.65 (m, 2H).
NCL 103	¹ H NMR (300 MHz, DMSO) δ 11.81 (br s, 2H), 10.32 – 9.86 (m, 4H), 8.52 (s, 2H), 8.12 (s, 2H), 7.85 (d, J = 8.4 Hz, 2H), 6.43 (s, 2H), 6.33 (d, J = 8.5 Hz, 2H).
NCL 104	¹ H NMR (300 MHz, DMSO) δ 11.72 (br s, 2H), 9.78 (br s, 2H), 9.45 (s, 2H), 8.48 (s, 2H), 8.34 (br s, 2H), 8.04 (s, 2H), 7.33 (s, 2H), 6.44 (s, 2H).
NCL 105	¹ H NMR (300 MHz, DMSO) δ 11.75 (br s, 2H), 9.71 (br s, 2H), 9.15 (br s, 2H), 8.66 – 8.40 (m, 4H), 8.13 (s, 2H), 7.33 (d, J = 8.6 Hz, 2H), 6.42 (d, J = 8.6 Hz, 2H).
NCL 106	¹ H NMR (300 MHz, DMSO) δ 9.12 (br s, 4H), 8.29 (s, 2H), 8.20 (s, 2H), 7.10 (s, 2H), 6.93 (s, 2H), 3.84 (s, 6H).
NCL 107	¹ H NMR (300 MHz, DMSO) δ 12.19 (s, 2H), 10.25 (s, 2H), 8.70 (s, 2H), 8.34 (s, 2H), 8.06 (d, J = 7.6 Hz, 2H), 7.35 – 7.23 (m, 2H), 7.00 (d, J = 8.2 Hz, 2H), 6.87 (t, J = 7.5 Hz, 2H).
NCL 108	¹ H NMR (300 MHz, DMSO) δ 8.00 (s, 2H), 7.26 – 7.08 (m, 6H), 6.98 – 6.43 (m, 4H).
NCL 109	¹ H NMR (300 MHz, DMSO) δ 6.60 (d, J = 1.2 Hz, 2H), 8.33 – 8.19 (m, 4H), 8.15 (d, J = 8.0 Hz, 2H), 7.74 – 7.61 (m, 2H), 7.12 (s, 2H).
NCL 110	¹ H NMR (300 MHz, DMSO) δ 8.95 (s, 2H), 8.56 (s, 2H), 8.31 (d, J = 8.7 Hz, 4H), 8.23 (d, J = 8.9 Hz, 4H).
NCL 111	¹ H NMR (300 MHz, DMSO) δ 11.80 (br s, 2H), 10.30 – 9.80 (m, 4H), 8.52 (s, 2H), 8.12 (s, 2H), 7.84 (d, J = 8.6 Hz, 2H), 6.42 (d, J = 8.6 Hz, 2H), 6.33 (d, J = 8.6 Hz, 2H).
NCL 112	¹ H NMR (300 MHz, DMSO) δ 12.48 (br s, 2H), 8.62 (s, 2H), 8.51 (s, 2H), 8.04 (d, J = 7.5 Hz, 4H), 7.85 – 7.69 (m, 8H), 7.54 – 7.36 (m, 6H).
NCL 113	¹ H NMR (400 MHz, DMSO) δ 11.92 (br s, 2H), 8.24 (s, 2H), 8.16 (s, 2H), 7.71 (d, J = 8.9 Hz, 4H), 6.74 (d, J = 8.9 Hz, 4H), 2.98 (s, 12H).
NCL 114	¹ H NMR (400 MHz, DMSO) δ 12.67 (br s, 2H), 8.31 (s, 2H), 8.40 (s, 2H), 8.06 (d, J = 1.8 Hz, 4H), 7.68 (t, J = 1.8 Hz, 2H).
NCL 115	¹ H NMR (400 MHz, DMSO) δ 12.09 (br s, 2H), 8.44 (s, 2H), 8.34 (s, 2H), 7.63 (d, J = 1.4 Hz, 2H), 7.33 (dd, J = 8.3, 1.6 Hz, 2H), 7.03 (d, J = 8.4 Hz, 2H), 3.86 (s, 6H), 3.81 (s, 6H).
NCL 116	¹ H NMR (400 MHz, DMSO) δ 12.27 (br s, 2H), 8.49 (s, 2H), 8.44 – 8.27 (m, 4H), 7.57 – 7.44 (m, 10H), 7.41 – 7.33 (m, 6H).
NCL 117	¹ H NMR (400 MHz, DMSO) δ 9.77 (br s, 2H), 8.36 (s, 2H), 8.28 (s, 2H), 7.58 (d, J = 1.5 Hz, 2H), 7.23 (dd, J = 8.1, 1.2 Hz, 2H), 6.87 (d, J = 8.1 Hz, 2H), 3.86 (s, 6H).
NCL 118	¹ H NMR (400 MHz, DMSO) δ 12.83 (br s, 2H), 8.76 (s, 2H), 8.69 (s, 2H), 8.31 – 8.21 (m, 2H), 7.45 – 7.31 (m, 4H).
NCL 119	¹ H NMR (400 MHz, DMSO) δ 12.13 (br s, 2H), 10.35 (s, 2H), 8.39 (s, 2H), 8.35 (s, 2H), 7.65 (d, J = 8.7 Hz, 4H), 7.72 (d, J = 8.6 Hz, 4H), 2.08 (s, 6H).
NCL 120	¹ H NMR (400 MHz, DMSO) δ 8.82 – 8.28 (m, 4H), 7.85 (d, J = 8.2 Hz, 4H), 7.33 (d, J = 8.2 Hz, 4H), 2.98 – 2.87 (m, 2H), 1.21 (d, J = 6.9 Hz, 12H).
NCL 121	¹ H NMR (400 MHz, DMSO) δ 8.60 – 8.30 (m, 4H), 7.64 (d, J = 8.1 Hz, 4H), 7.28 (d, J = 8.1 Hz, 4H), 2.59 (t, J = 7.5 Hz, 4H), 1.64 – 1.54 (m, 4H), 0.88 (t, J = 7.3 Hz, 6H).
NCL 122	¹ H NMR (400 MHz, DMSO) δ 8.56 (s, 2H), 8.42 (d, J = 2.0 Hz, 2H), 8.38 (s, 2H), 8.09 (dd, J = 8.7, 2.0 Hz, 2H), 7.29 (d, J = 8.7 Hz, 2H).

- NCL 123 ¹H NMR (400 MHz, DMSO) δ 8.68 (s, 2H), 8.43 (s, 2H), 8.28 – 8.16 (m, 2H), 7.77 – 7.64 (m, 2H), 7.58 – 7.46 (m, 2H).
¹³C NMR (101 MHz, DMSO) δ 153.0, 150.8 (dd, J = 250.6, 13.0 Hz), 149.9 (dd, J = 245.9, 13.2 Hz), 146.6, 131.2 (dd, J = 6.4, 3.4 Hz), 126.0 (dd, J = 6.4, 2.8 Hz), 117.8 (d, J = 17.7 Hz), 115.7 (d, J = 18.5 Hz).
- NCL 124 ¹H NMR (400 MHz, DMSO) δ 10.84 (br s, 2H), 9.29 (s, 2H), 8.80 (d, J = 8.6 Hz, 2H), 8.43 (s, 2H), 7.94 (d, J = 9.0 Hz, 2H), 7.87 (d, J = 7.9 Hz, 2H), 7.64 – 7.56 (m, 2H), 7.43 – 7.38 (m, 2H), 7.34 (d, J = 8.9 Hz, 2H).
- NCL 125 ¹H NMR (400 MHz, DMSO) δ 9.21 (br s, 2H), 8.42 – 8.17 (m, 4H), 7.43 (d, J = 1.9 Hz, 2H), 7.25 (dd, J = 8.4, 1.9 Hz, 2H), 6.99 (d, J = 8.4 Hz, 2H), 3.83 (s, 6H).
- NCL 126 ¹H NMR (400 MHz, DMSO) δ 12.27 (br s, 2H), 8.58 (s, 2H), 8.42 (s, 2H), 7.96 (d, J = 8.3 Hz, 4H), 7.58 (d, J = 8.2 Hz, 4H), 4.39 (s, 2H).
- NCL 127 ¹H NMR (400 MHz, DMSO) δ 12.84 (br s, 2H), 8.84 (s, 2H), 8.74 (s, 2H), 8.46 (d, J = 8.6 Hz, 2H), 7.71 (d, J = 1.6 Hz, 2H), 7.54 (dd, J = 8.6, 1.2 Hz, 2H).
- NCL 128 ¹H NMR (400 MHz, DMSO) δ 8.64 (s, 2H), 8.36 (s, 2H).
- NCL 129 ¹H NMR (400 MHz, DMSO) δ 8.84 (s, 2H), 8.69 (s, 2H), 8.41 (dd, J = 7.8, 1.7 Hz, 2H), 7.70 (dd, J = 8.0, 1.0 Hz, 2H), 7.48 (t, J = 7.3 Hz, 2H)*, 7.44 – 7.37 (m, 2H)*.
- NCL 130 ¹H NMR (400 MHz, DMSO) δ 8.66 (s, 2H), 8.37 (s, 2H), 7.79 (d, J = 1.6 Hz, 2H), 7.68 – 7.62 (m, 2H), 3.92 (s, 6H), 3.78 (s, 6H).
- NCL 131 ¹H NMR (400 MHz, DMSO) δ 12.45 (br s, 2H), 8.68 (s, 2H), 8.41 (s, 2H), 8.29 (s, 2H), 7.87 (d, J = 7.8 Hz, 2H), 7.70 – 7.63 (m, 2H), 7.44 (t, J = 7.9 Hz, 2H).
- NCL 132 ¹H NMR (400 MHz, DMSO) δ 12.69 (br s, 2H), 8.61 (s, 4H), 7.34 (dd, J = 9.0, 2.8 Hz, 2H), 7.22 (td, J = 8.6, 2.9 Hz, 2H), 7.03 (dd, J = 8.8, 4.6 Hz, 2H), 5.29 (s, 4H).
- NCL 133 ¹H NMR (400 MHz, DMSO) δ 12.51 (br s, 2H), 8.49 – 8.28 (m, 4H), 8.16 (s, 2H), 7.35 (s, 2H).
- NCL 134 ¹H NMR (400 MHz, DMSO) δ 12.42 (br s, 2H), 8.60 (s, 2H), 8.42 (s, 2H), 7.91 (d, J = 8.5 Hz, 4H), 7.69 (d, J = 8.5 Hz, 4H).
- NCL 135 ¹H NMR (400 MHz, DMSO) δ 12.24 (br s, 2H), 8.66 (s, 2H), 8.49 (br s, 2H), 7.79 (s, 2H), 7.24 (s, 2H), 3.88 (s, 6H), 3.85 (s, 6H).
- NCL 136 ¹H NMR (400 MHz, DMSO) δ 12.21 (br s, 2H), 8.44 (s, 2H), 8.39 (s, 2H), 7.83 (d, J = 8.2 Hz, 4H), 7.30 (d, J = 8.2 Hz, 4H), 2.63 (t, J = 7.7 Hz, 4H), 1.61 – 1.52 (m, 4H), 1.36 – 1.26 (m, 4H), 0.90 (t, J = 7.3 Hz, 6H).
- NCL 137 ¹H NMR (400 MHz, DMSO) δ 12.94 (br s, 2H), 8.68 (s, 2H), 8.33 (s, 2H), 7.60 (d, J = 7.9 Hz, 4H)*, 7.49 (dd, J = 8.7, 7.4 Hz, 2H)*.
- NCL 138 ¹H NMR (400 MHz, DMSO) δ 12.12 (br s, 2H), 8.21 (s, 2H), 7.97 (s, 2H), 7.54 – 7.47 (m, 6H), 7.42 – 7.36 (m, 6H), 7.31 – 7.21 (m, 8H), 6.84 (d, J = 9.8 Hz, 2H).
- NCL 139 ¹H NMR (400 MHz, DMSO) δ 12.65 (br s, 2H), 9.63 (d, J = 1.8 Hz, 2H), 8.88 – 8.58 (m, 6H), 8.13 – 8.01 (m, 4H), 7.88 – 7.79 (m, 2H), 7.68 (t, J = 7.4 Hz, 2H).
- NCL 140 ¹H NMR (400 MHz, DMSO) δ 12.17 (br s, 2H), 8.46 (s, 2H), 8.37 (s, 2H), 7.86 (d, J = 8.2 Hz, 4H), 7.34 (d, J = 8.2 Hz, 4H), 2.53 (s, 6H).
- NCL 141 ¹H NMR (400 MHz, DMSO) δ 8.66 (s, 2H), 8.60 (d, J = 1.9 Hz, 2H), 8.52 (br s, 2H), 8.49 (s, 2H), 8.09 (d, J = 8.6 Hz, 2H), 7.50 (dd, J = 8.6, 2.0 Hz, 2H).
- NCL 142 ¹H NMR (400 MHz, DMSO) δ 8.80 (s, 2H), 7.41 – 7.20 (m, 12H), 5.56 (s, 2H), 3.79 (d, J = 3.4 Hz, 4H).
- NCL 143 ¹H NMR (400 MHz, DMSO) δ 11.91 (br s, 2H), 8.66 (br s, 2H), 8.10 – 8.00 (m, 4H), 7.51 – 7.41 (m, 6H), 2.45 (s, 6H).
- NCL 144 ¹H NMR (400 MHz, DMSO) δ 12.36 (br s, 2H), 8.38 – 8.23 (m, 4H), 7.16 (d, J = 3.5 Hz, 2H), 6.82 (d, J = 3.5 Hz, 2H).
- NCL 145 ¹H NMR (400 MHz, DMSO) δ 12.41 (br s, 2H), 8.41 – 8.21 (m, 4H), 7.20 (d, J = 3.5 Hz, 2H), 6.73 (d, J = 3.5 Hz, 2H).
- NCL 146 ¹H NMR (400 MHz, DMSO) δ 11.97 (br s, 2H), 11.45 (s, 2H), 8.47 (s, 2H), 8.30 (s, 2H), 8.02 (s, 2H), 7.80 (dd, J = 8.6, 0.9 Hz, 2H), 7.48 (d, J = 8.5 Hz, 2H), 7.45 – 7.40 (m, 2H), 6.53 (s, 2H).
¹³C NMR (101 MHz, DMSO) δ 152.4, 150.6, 137.4, 127.6, 126.7, 124.5, 121.8, 120.3, 111.9, 102.0.
- NCL 147 ¹H NMR (400 MHz, DMSO) δ 13.09 (br s, 2H), 9.97 (s, 2H), 9.01 (s, 2H), 8.73 (s, 2H), 8.19 – 8.09 (m, 4H), 7.95 – 7.84 (m, 4H).

NCL 148	¹ H NMR (400 MHz, DMSO) δ 12.44 (s, 4H), 8.60 (s, 2H), 7.99 (d, J = 8.3 Hz, 4H), 7.81 (d, J = 8.3 Hz, 4H), 7.63 (d, J = 16.0 Hz, 2H), 6.68 (d, J = 16.0 Hz, 2H).
NCL 149	¹ H NMR (400 MHz, DMSO) δ 8.75 (d, J = 6.1 Hz, 4H), 8.68 (s, 2H), 8.47 (s, 2H), 8.07 (d, J = 6.1 Hz, 4H).
NCL 150	¹ H NMR (400 MHz, DMSO) δ 12.10 (br s, 2H), 8.23 – 8.07 (m, 4H), 7.55 (d, J = 8.7 Hz, 4H), 7.13 (d, J = 15.0 Hz, 2H), 6.99 (d, J = 8.7 Hz, 4H), 6.81 (dd, J = 15.0, 9.4 Hz, 2H), 3.79 (s, 6H).
NCL 151	¹ H NMR (400 MHz, DMSO) δ 11.92 (br s, 2H), 10.13 (br s, 2H), 8.28 (s, 4H), 7.75 (d, J = 8.5 Hz, 4H), 6.86 (d, J = 8.5 Hz, 4H).
NCL 152	¹ H NMR (400 MHz, DMSO) δ 12.06 (br s, 2H), 9.35 (s, 2H), 8.66 (s, 2H), 8.48 (s, 2H), 7.67 (d, J = 7.2 Hz, 2H), 7.25 (d, J = 7.2 Hz, 2H), 6.88 (t, J = 7.6 Hz, 2H), 2.23 (s, 6H).
NCL 153	¹ H NMR (400 MHz, DMSO) δ 11.68 (s, 2H), 8.78 (s, 2H), 8.09 (d, J = 8.4 Hz, 4H), 7.52 (d, J = 8.6 Hz, 4H), 2.92 (q, J = 7.5 Hz, 4H), 1.12 (t, J = 7.4 Hz, 6H).
NCL 154	¹ H NMR (400 MHz, DMSO) δ 12.09 (s, 2H), 8.69 (s, 2H), 8.07 (d, J = 8.8 Hz, 4H), 7.50 (d, J = 8.6 Hz, 4H), 3.01 – 2.88 (m, 4H), 1.49 – 1.39 (m, 8H), 0.88 (t, J = 6.8 Hz, 6H).
NCL 155	¹ H NMR (400 MHz, DMSO) δ 11.63 (s, 2H), 8.76 (s, 2H), 8.01 (d, J = 8.6 Hz, 4H), 7.65 (d, J = 8.6 Hz, 4H), 2.41 (s, 6H).
NCL 156	¹ H NMR (400 MHz, DMSO) δ 12.16 (s, 2H), 8.71 (s, 2H), 8.06 (d, J = 8.0 Hz, 4H), 7.50 (d, J = 8.3 Hz, 4H), 3.01 – 2.87 (m, 4H), 1.58 – 1.46 (m, 4H), 1.01 (t, J = 7.1 Hz, 6H).
NCL 157	¹ H NMR (400 MHz, DMSO) δ 11.71 (br s, 2H), 8.40 (s, 2H), 8.37 (s, 2H), 7.29 (d, J = 8.4 Hz, 2H), 6.87 (d, J = 2.0 Hz, 2H), 6.73 (br s, 4H), 6.58 (dd, J = 8.3, 2.0 Hz, 2H).
NCL 158	¹³ C NMR (101 MHz, DMSO) δ 152.1, 151.5, 148.9, 136.0, 134.7, 115.1, 114.5, 112.8. ¹ H NMR (400 MHz, DMSO) δ 13.03 (br s, 1H), 10.49 (br s, 1H), 9.35 (br s, 1H), 7.24 (d, J = 7.6 Hz, 2H), 7.11 – 6.61 (m, 8H), 2.5 (Contains CH ₂ groups determined by COSY, however eclipsed by solvent signal), 0.93 (s, 6H).
NCL 159	¹³ C NMR (101 MHz, DMSO) δ 157.8, 155.4, 152.9, 134.4, 130.1, 118.7, 115.8, 29.9, 10.8. *COSY was used to determine that the signal due to the methylene protons appears under the DMSO signal. Line broadening is apparent in the ¹³ C-NMR (due to tautomerisation effects) making carbon allocation difficult.
NCL 160	¹ H NMR (400 MHz, DMSO) δ 12.69 (br s, 1H), 10.44 (br s, 1H), 9.15 (br s, 1H), 7.21 (s, 2H), 7.09 – 6.60 (m, 8H), 2.93 (p, J = 7.7 Hz, 2H), 1.84 – 1.01 (m, 16H).
NCL 161	¹ H NMR (400 MHz, DMSO) δ 12.49 (br s, 2H), 8.65 (s, 2H), 8.49 (s, 2H), 8.10 (d, J = 8.7 Hz, 4H)*, 7.47 (d, J = 8.3 Hz, 4H)*.
NCL 162	¹ H NMR (400 MHz, DMSO) δ 7.73 (d, J = 8.8 Hz, 2H), 6.92 (d, J = 8.9 Hz, 2H), 3.22 – 3.13 (m, 4H), 2.99 – 2.86 (m, 4H), 2.24 (s, 3H).
NCL 163	¹ H NMR (400 MHz, CDCl ₃) δ 7.76 – 7.70 (m, 2H), 7.35 – 7.30 (m, 2H), 4.99 (s, 2H), 4.34 (q, J = 7.1 Hz, 2H), 2.39 (s, 3H), 1.35 (t, J = 7.1 Hz, 3H).
NCL 164	¹ H NMR (400 MHz, DMSO) δ 13.88 (s, 1H), 13.21 (s, 1H), 11.50 (s, 2H), 9.70 (s, 2H), 7.70 (s, 2H), 7.43 (t, J = 7.7 Hz, 2H), 7.14 (t, J = 7.5 Hz, 2H), 7.00 (d, J = 7.8 Hz, 2H).
NCL 165	¹ H NMR (400 MHz, DMSO) δ 10.85 (s, 2H), 9.38 (s, 2H), 8.64 (s, 3H), 7.90 (s, 3H), 7.34 (d, J = 8.5 Hz, 2H), 7.19 (s, 3H), 6.83 (d, J = 2.1 Hz, 2H), 6.57 (dd, J = 8.5, 2.1 Hz, 2H).
NCL 166	¹ H NMR (400 MHz, DMSO) δ 10.87 (br s, 1H), 9.53 (br s, 1H), 8.77 (br s, 3H), 7.61–7.44 (m, 8H), 7.40 (br s, 3H), 7.27 (br s, 3H), 4.07 (br s, 2H).
NCL 167	¹ H NMR (400 MHz, DMSO) δ 12.70 (br s, 2H), 8.75 (br s, 2H), 8.55 (br s, 2H), 8.13 (d, J = 8.2 Hz, 4H), 7.81 (d, J = 8.1 Hz, 4H).
NCL 168	¹ H NMR (400 MHz, DMSO) δ 14.09 (s, 1H), 8.12 (d, J = 6.9 Hz, 2H), 7.97 (dd, J = 6.4, 3.0 Hz, 2H), 7.56 (odd, J = 8.2, 6.7, 2.9 Hz, 7H).
NCL 169	¹ H NMR (400 MHz, DMSO) δ 11.85 (br s, 2H), 6.42 (br s, 4H), 7.35 (d, J = 5.8 Hz, 4H), 6.75 (s, 2H), 6.67 (d, J = 7.9 Hz, 2H), 3.44 (d, J = 6.9 Hz, 2H)*, 1.24 (t, J = 5.3 Hz, 6H). *Signal partly eclipsed by H ₂ O in DMSO

NCL170	¹ H NMR (400 MHz, DMSO) δ 12.29 (s, 2H), 10.29 (s, 2H), 8.55 (s, 2H), 8.13 (d, J = 8.5 Hz, 2H), 7.71 (s, 2H), 7.35 (dd, J = 8.5, 1.8 Hz, 2H), 2.12 (s, 6H).
NCL171	¹ H NMR (400 MHz, DMSO) δ 11.63 (s, 2H), 9.84 (s, 2H), 8.46 (s, 2H), 8.02 (s, 2H), 7.74 (d, J = 8.8 Hz, 2H), 6.30 (d, J = 7.4 Hz*, 2H), 6.17 (s, 2H), 2.94 (s, 12H). *Poorly resolved doublet gives reduced coupling constant.
NCL172	¹ H NMR (400 MHz, DMSO) δ 12.00 (s, 2H), 8.92 (s, 2H), 8.64 (dd, J = 4.7, 0.6 Hz, 2H), 8.57 (d, J = 8.1 Hz, 2H), 7.89 (td, J = 8.0, 1.6 Hz, 2H), 7.50 – 7.41 (m, 2H), 2.52 (s, 6H).
NCL 173	¹ H NMR (400 MHz, DMSO) δ 11.49 (br s, 2H), 10.62 (br s, 2H), 8.55 (br s, 2H), 7.59 (d, J = 8.3 Hz, 2H), 6.98 (d, J = 1.9 Hz, 2H), 6.92 (dd, J = 8.4, 2.0 Hz, 2H), 2.37 (s, 6H).
NCL174	¹ H NMR (400 MHz, DMSO) δ 12.02 (s, 2H), 10.81 (s, 2H), 8.63 (s, 2H), 8.38 (s, 2H), 8.12 (d, J = 8.3 Hz, 2H), 7.13 – 6.84 (m, 4H).
NCL 175	¹ H NMR (400 MHz, DMSO) δ 12.63 (br s, 2H), 8.90 (d, J = 2.1 Hz, 2H), 8.74 (s, 2H), 8.56 – 8.42 (m, 4H), 7.66 (d, J = 8.4 Hz, 2H).
NCL 176	¹ H NMR (400 MHz, DMSO) δ 11.94 (br s, 2H), 8.44 (s, 2H), 8.36 (s, 2H), 8.07 (dd, J = 4.8, 1.7 Hz, 2H), 7.72 (dd, J = 7.6, 1.4 Hz, 2H), 7.19 (s, 4H), 6.67 (dd, J = 7.5, 4.9 Hz, 2H).
NCL 177	¹ H NMR (400 MHz, DMSO) δ 8.67 (s, 2H), 7.97 (s, 4H), 7.50 (d, J = 8.6 Hz, 4H), 4.81 (s, 4H).
NCL178	¹ H NMR (400 MHz, DMSO) δ 10.17 (s, 2H), 8.24 (s, 1H), 7.83 (d, J = 8.6 Hz, 4H), 7.50 (d, J = 8.6 Hz, 4H), 6.97 (s, 1H), 2.32 (s, 6H).
NCL179	¹ H NMR (400 MHz, DMSO) δ 10.70 (s, 2H), 8.02 (s, 2H), 7.67 (d, J = 8.4 Hz, 4H), 7.52 (d, J = 8.4 Hz, 4H), 6.28 (s, 1H), 5.85 (s, 2H). ¹³ C NMR (101 MHz, DMSO) δ 162.8, 162.6, 138.8, 134.1, 133.1, 128.9, 127.6, 73.5.
NCL180	¹ H NMR (400 MHz, DMSO) δ 10.62 (s, 2H), 8.22 (d, J = 0.9 Hz, 1H), 7.82 – 7.74 (m, 4H), 7.53 – 7.47 (m, 4H), 6.93 (d, J = 0.6 Hz, 1H), 5.85 (t, J = 5.3 Hz, 2H), 4.74 (d, J = 5.2 Hz, 4H).
NCL181	¹ H NMR (400 MHz, DMSO) δ 11.20 (s, 2H), 8.17 (s, 1H), 8.09 (s, 2H), 7.72 (d, J = 7.4 Hz, 4H), 7.54 (d, J = 7.6 Hz, 4H), 6.83 (s, 1H).
NCL182	¹ H NMR (400 MHz, CDCl ₃) δ 7.44 – 7.19 (m, 6H), 5.67 (s, 1H), 5.42 (s, 1H), 5.06 (s, 2H), 4.95 – 4.63 (m, 1H), 1.52 (d, J = 6.8 Hz, 3H).
NCL183	¹ H NMR (400 MHz, CDCl ₃) δ 8.28 (s, 1H), 7.43 – 7.13 (m, 10H), 6.19 (s, 1H), 5.94 (s, 1H), 4.95 – 4.48 (m, 2H), 1.56 (d, J = 6.8 Hz, 4H), 1.50 (d, J = 6.5 Hz, 2H).
NCL184	¹ H NMR (400 MHz, MeOD) δ 7.44 – 7.04 (m, 10H), 5.26 – 4.53 (m, 7H), 1.51 – 1.35 (m, 6H).
NCL188	¹ H NMR (400 MHz, DMSO) δ 11.54 (s, 1H), 7.99 (d, J = 8.7 Hz, 2H), 7.90 (s, 3H), 7.47 (d, J = 8.6 Hz, 2H), 2.91 – 2.82 (m, 2H), 1.48 – 1.32 (m, 4H), 0.89 – 0.84 (m, 3H).
NCL190	¹³ C NMR (101 MHz, DMSO) δ 156.2, 153.8, 134.8, 134.4, 128.7, 128.4, 28.1, 26.6, 22.0, 13.8.
NCL191	¹ H NMR (400 MHz, DMSO) δ 11.42 (s, 1H), 8.06 (d, J = 8.7 Hz, 3H), 8.01 – 7.71 (m, 5H), 7.53 (d, J = 8.7 Hz, 3H), 4.90 (s, 2H), 3.69 (br s, 3H).
NCL191	¹ H NMR (400 MHz, DMSO) δ 11.51 (s, 1H), 8.85 (s, 3H), 7.99 (d, J = 8.6 Hz, 7H), 7.46 (d, J = 8.6 Hz, 4H), 2.35 (s, 3H).
NCL192	¹ H NMR (400 MHz, DMSO) δ 11.65 (s, 1H), 8.21 (s, 4H), 7.82 (dd, J = 7.6, 1.9 Hz, 2H), 7.53 – 7.39 (m, 3H).
NCL193	¹ H NMR (400 MHz, DMSO) δ 10.71 (s, 2H), 8.00 (s, 2H), 7.66 (d, J = 8.5 Hz, 4H), 7.60 (d, J = 8.6 Hz, 4H), 6.27 (s, 1H), 5.86 (s, 2H). ¹³ C NMR (101 MHz, DMSO) δ 162.7, 162.6, 138.8, 134.5, 131.8, 127.9, 121.7, 73.5.
NCL194	¹ H NMR (400 MHz, DMSO) δ 12.01 (br s, 1H), 10.84 (s, 2H), 9.98 (br s, 1H), 9.79 (br s, 1H), 7.96 (br s, 3H), 7.79 (br s, 2H), 7.67 – 7.37 (m, 6H), 5.37 (br s, 1H).
NCL195	¹ H NMR (400 MHz, DMSO) δ 10.51 (s, 2H), 8.00 (s, 2H), 7.54 (d, J = 8.0 Hz, 4H), 7.26 (d, J = 7.9 Hz, 4H), 6.26 (s, 1H), 5.77 (s, 2H), 2.34 (s, 6H). ¹³ C NMR (101 MHz, DMSO) δ 162.8, 162.6, 140.1, 138.4, 132.5, 129.4, 126.0, 73.3, 21.0.

NCL196	¹ H NMR (400 MHz, CDCl ₃) δ 10.31 (s, 2H), 9.74 (s, 2H), 7.48 (d, J = 8.8 Hz, 4H), 6.83 (d, J = 8.6 Hz, 4H), 6.20 (s, 1H), 5.70 (s, 2H). ¹³ C NMR (101 MHz, CDCl ₃) δ 162.7, 162.5, 158.3, 140.5, 127.7, 126.3, 115.7, 73.0.
NCL197	¹ H NMR (400 MHz, DMSO) δ 10.51 (s, 2H), 9.55 (s, 2H), 7.95 (s, 2H), 7.22 (t, J = 7.9 Hz, 2H), 7.11 – 7.04 (m, 4H), 6.76 (d, J = 8.4 Hz, 2H), 6.23 (s, 1H), 5.80 (s, 2H). ¹³ C NMR (101 MHz, DMSO) δ 162.6, 162.6, 157.7, 140.4, 136.4, 129.9, 117.4, 116.1, 112.4, 73.3.
NCL199	¹ H NMR (400 MHz, DMSO) δ 10.50 (s, 2H), 8.04 (s, 2H), 7.86 (d, J = 7.5 Hz, 4H), 7.45 (t, J = 7.1 Hz, 4H), 7.40 – 7.34 (m, 2H), 6.30 (s, 1H), 5.82 (s, 2H). ¹³ C NMR (101 MHz, DMSO) δ 163.3, 163.1, 140.5, 135.7, 129.3, 129.2, 126.5, 73.9.
NCL200	¹ H NMR (400 MHz, DMSO) δ 10.51 (s, 2H), 8.02 (s, 2H), 7.88 (d, J = 8.3 Hz, 4H), 7.47 (d, J = 8.3 Hz, 4H), 6.25 (s, 1H), 5.77 (s, 2H), 1.31 (s, 18H).
NCL201	¹ H NMR (400 MHz, DMSO) δ 10.19 (s, 2H), 7.91 (s, 2H), 7.47 (d, J = 8.8 Hz, 4H), 6.77 (d, J = 8.9 Hz, 4H), 6.18 (s, 1H), 5.63 (s, 2H).
NCL202	¹ H NMR (600 MHz, DMSO) δ 13.43 (s, 1H), 11.45 (s, 2H), 10.28 (br. s, 1H), 9.47 (s, 1H), 8.38 (s, 2H), 7.71 (dd, J = 7.7, 1.3 Hz, 1H), 7.67 (br d, J = 7.0 Hz, 2H), 7.53 – 7.48 (m, 1H), 7.28 – 7.23 (m, 2H), 7.03 – 6.98 (m, 2H), 6.96 – 6.90 (m, 4H), 6.56 (br s, 1H). *Due to tautomers and rotamers associated with proximity of the phenol to the hydrazone >17 protons are observed. ¹³ C NMR (151 MHz, DMSO) δ 167.7, 163.2, 162.6, 158.7*, 141.4*, 135.3, 133.9, 131.0, 130.5, 120.7*, 119.95, 119.8, 118.9, 117.6, 116.7, 116.6*. *Line broadening due to tautomers and rotamers made signals difficult to assign.
NCL203	¹ H NMR (400 MHz, DMSO) δ 12.60 – 10.94 (m, 3H), 7.79 – 7.39 (m, 3H), 2.25 (s, 2H), 1.82 – 1.59 (m, 10H), 1.35 – 1.15 (m, 10H).
NCL204	¹ H NMR (400 MHz, DMSO) δ 9.58 (d, J = 7.6 Hz, 2H), 7.82 (s, 4H), 7.50 (s, 4H), 6.45 (d, J = 9.6 Hz, 1H), 5.75 (s, 2H), 2.28 (d, J = 9.3 Hz, 6H). ¹³ C NMR (101 MHz, DMSO) δ 163.48, 162.11, 143.34, 137.68, 132.68, 128.34, 127.16, 75.15, 13.01.
NCL205	¹ H NMR (600 MHz, DMSO) δ 10.92 (s, 2H), 8.42 (s, 2H), 7.98 (d, J = 7.5 Hz, 2H), 7.50 (d, J = 7.8 Hz, 2H), 7.44 (t, J = 7.4 Hz, 2H), 7.38 (t, J = 7.3 Hz, 2H), 6.35 (s, 1H), 5.95 (s, 2H). ¹³ C NMR (151 MHz, DMSO) δ 162.75, 162.68, 135.0, 132.4, 132.0, 130.1, 129.9, 127.6, 126.2, 73.7.
NCL206	¹ H NMR (400 MHz, DMSO) δ 10.75 (s, 2H), 8.09 (s, 2H), 7.52 (d, J = 8.1 Hz, 4H), 7.24 (d, J = 8.0 Hz, 4H), 6.71 (s, 1H), 2.33 (s, 6H).
NCL207	¹ H NMR (400 MHz, DMSO) δ 11.43 (s, 2H), 11.05 (s, 2H), 8.31 (s, 2H), 7.42 (s, 2H), 7.29 – 7.18 (m, 2H), 6.89 (t, J = 7.7 Hz, 4H), 6.74 (s, 2H).
NCL208	¹ H NMR (400 MHz, DMSO) δ 9.63 (s, 2H), 7.82 (d, J = 8.5 Hz, 4H), 7.46 (d, J = 8.5 Hz, 4H), 6.73 (s, 2H), 2.29 (s, 6H).
NCL209	¹ H NMR (400 MHz, DMSO) δ 10.75 (s, 2H), 9.55 (s, 2H), 8.04 (s, 2H), 7.21 (t, J = 7.6 Hz, 2H), 7.06 (s, 2H), 7.02 (d, J = 7.6 Hz, 2H), 6.84 – 6.57 (m, 4H).
NCL210	¹ H NMR (400 MHz, DMSO) δ 11.16 (s, 2H), 8.22 (s, 2H), 7.85 (d, J = 8.3 Hz, 4H), 7.83 (d, J = 8.5 Hz, 4H), 6.92 (s, 2H).
NCL211	¹ H NMR (400 MHz, DMSO) δ 10.56 (s, 2H), 9.78 (s, 2H), 8.02 (s, 2H), 7.45 (d, J = 8.6 Hz, 4H), 6.80 (d, J = 8.6 Hz, 4H), 6.62 (s, 2H).
NCL212	¹ H NMR (400 MHz, DMSO) δ 10.94 (s, 2H), 8.10 (s, 2H), 7.67 – 7.60 (m, 4H), 7.61 – 7.54 (m, 4H), 6.80 (s, 2H).
NCL213	¹ H NMR (400 MHz, DMSO) δ 10.23 (br s, 2H), 7.27 (s, 2H), 2.27 – 1.98 (m, 2H), 1.78 – 1.42 (m, 10H), 1.33 – 1.00 (m, 10H).
NCL214	¹ H NMR (400 MHz, DMSO) δ 10.84 (s, 2H), 8.14 (s, 2H), 7.63 (d, J = 7.4 Hz, 4H), 7.46 – 7.33 (m, 3H), 6.76 (s, 2H).
NCL215	¹ H NMR (600 MHz, DMSO) δ 12.06 (s, 2H), 8.72 (s, 2H), 7.97 (t, J = 8.4 Hz, 2H), 7.56 (dd, J = 11.1, 1.9 Hz, 2H), 7.38 (dd, J = 8.5, 1.9 Hz, 2H), 2.42 (d, J = 2.9 Hz, 6H).

NCL216	¹ H NMR (400 MHz, DMSO) δ 12.43 (br s, 2H), 8.66 (br s, 2H), 8.62 (br s, 2H), 8.38 (t, J = 8.3 Hz, 2H), 7.61 (dd, J = 10.5, 1.9 Hz, 2H), 7.45 (dd, J = 8.6, 1.6 Hz, 2H). ¹³ C NMR (101 MHz, DMSO) δ 160.7 (d, J = 254.5 Hz), 152.8*, 140.8*, 136.3 (d, J = 10.8 Hz), 128.5, 125.3, 120.2 (d, J = 10.0 Hz), 116.7 (d, J = 24.7 Hz). *Broad signals
NCL217	¹ H NMR (400 MHz, DMSO) δ 11.66 (s, 2H), 8.61 (s, 2H), 7.94 (d, J = 8.2 Hz, 4H), 7.25 (d, J = 8.1 Hz, 4H), 2.41 (s, 6H), 2.35 (s, 6H).
NCL218	¹³ C NMR (101 MHz, DMSO) δ 154.0, 153.3, 139.7, 133.9, 129.0, 127.0, 21.0, 14.9. ¹ H NMR (400 MHz, DMSO) δ 10.73 (s, 2H), 8.05 (s, 2H), 7.70 (d, J = 8.58 Hz, 4H), 7.31 – 7.27 (m, 4H), 6.24 (s, 1H), 5.92 (s, 2H), 4.17 (dq, J = 7.06, 8.70 Hz, 8H), 1.28 (td, J = 1.01, 7.05 Hz, 12H).
NCL219	¹ H NMR (400 MHz, DMSO) δ 11.74 (s, 2H), 8.60 (s, 2H), 7.95 (d, J = 8.6 Hz, 4H), 7.45 (d, J = 8.6 Hz, 4H), 2.42 (s, 6H), 1.31 (s, 18H).
NCL220	¹³ C NMR (101 MHz, DMSO) δ 154.1, 153.3, 152.6, 134.0, 126.8, 125.0, 34.5, 31.0, 14.9. ¹ H NMR (400 MHz, DMSO) δ 10.61 (s, 2H), 8.03 (s, 2H), 7.70 (dd, J = 8.7, 5.6 Hz, 4H), 7.29 (t*, J = 8.9 Hz, 4H), 6.27 (s, 1H), 5.82 (s, 2H).
NCL221	¹ H NMR (600 MHz, DMSO) δ 10.89 (s, 2H), 8.11 (s, 2H), 7.86 (d, J = 8.2 Hz, 4H), 7.81 (d, J = 8.4 Hz, 4H), 6.34 (s, 1H), 5.94 (s, 2H).
NCL222	¹ H NMR (400 MHz, DMSO) δ 10.76 (s, 2H), 7.99 (s, 2H), 7.70 (td, J = 1.7, 9.2 Hz, 2H), 7.54 – 7.41 (m, 4H), 6.30 (s, 1H), 5.87 (s, 2H).
NCL223	¹ H NMR (400 MHz, DMSO) δ 10.48 (s, 2H), 10.07 (s, 2H), 7.98 (s, 2H), 7.65 (d, J = 8.7 Hz, 4H), 7.58 (d, J = 8.7 Hz, 4H), 6.24 (s, 1H), 5.76 (s, 2H), 2.07 (s, 6H).
NCL224	¹ H NMR (400 MHz, DMSO) δ 8.36 (s, 1H), 7.96 – 7.90 (m, 2H), 7.86 (s, 1H), 7.79 – 7.73 (m, 2H), 7.54 – 7.47 (m, 6H), 4.26 (q, J = 7.1 Hz, 2H), 1.27 (t, J = 7.1 Hz, 3H).
NCL225	¹ H NMR (400 MHz, DMSO) δ 8.35 (s, 1H), 7.98 – 7.91 (m, 2H), 7.87 (s, 1H), 7.79 – 7.74 (m, 2H), 7.51 (dd, J = 2.8, 8.5 Hz, 6H), 3.99 (d, J = 6.5 Hz, 2H), 1.95 (hept, J = 6.7 Hz, 1H), 0.93 (d, J = 6.7 Hz, 6H).
NCL226	¹ H NMR (400 MHz, DMSO) δ 10.95 (s, 1H), 9.18 (t, J = 5.6 Hz, 1H), 8.38 (d, J = 10.1 Hz, 2H), 8.32 (s, 1H), 7.99 – 7.92 (m, 2H), 7.83 (d, J = 8.5 Hz, 2H), 7.57 – 7.46 (m, 5H), 3.30 – 3.20 (m, 2H), 1.14 (t, J = 7.2 Hz, 3H).
NCL227	¹ H NMR (400 MHz, DMSO) δ 11.00 (s, 1H), 9.68 (s, 1H), 8.52 (s, 1H), 8.41 (s, 1H), 8.32 (s, 1H), 8.01 – 7.80 (m, 4H), 7.56 – 7.46 (m, 4H), 7.36 (d, J = 6.00 Hz, 4H), 7.30 – 7.21 (m, 1H), 4.56 – 4.33 (m, 2H).
NCL228	¹ H NMR (400 MHz, DMSO) δ 10.95 (s, 1H), 9.22 (t, J = 5.6 Hz, 1H), 8.40 (s, 1H), 8.34 (s, 1H), 8.32 (s, 1H), 7.98 – 7.90 (m, 2H), 7.87 – 7.80 (m, 2H), 7.55 – 7.48 (m, 4H), 3.22 (q, J = 6.6 Hz, 2H), 1.56 – 1.46 (m, 2H), 1.39 – 1.25 (m, 6H), 0.88 (t, J = 6.6 Hz, 3H).
NCL229	¹ H NMR (400 MHz, DMSO) δ 10.98 (s, 1H), 9.60 (s, 1H), 8.54 (s, 1H), 8.40 (s, 1H), 8.30 (s, 1H), 8.01 – 7.89 (m, 2H), 7.85 (d, J = 8.4 Hz, 2H), 7.64 – 7.59 (m, 1H), 7.54 – 7.47 (m, 4H), 6.42 (dd, J = 1.8, 3.3 Hz, 1H), 6.31 (m, 1H), 4.44 (d, J = 5.5 Hz, 2H).
NCL230	¹ H NMR (400 MHz, DMSO) δ 10.41 (s, 2H), 7.98 (s, 2H), 7.59 (d, J = 8.9 Hz, 4H), 7.02 (d, J = 8.9 Hz, 4H), 6.23 (s, 1H), 5.73 (s, 2H), 3.80 (s, 6H).

EXAMPLE 12: Treatment of a bacterial infection *in vivo* by the administration of NCL812 or NCL099.

[00386] The objective of this study was to determine the efficacy of an Investigational Veterinary Product containing NCL812 or NCL099 in the treatment of a skin infection in mice

[00387] Summary of the Model: A useful animal model system should be clinically relevant, experimentally robust, ethically acceptable, convenient to perform and should provide reliable and reproducible results. There are many animal models of topical skin infection that have been described including the croton oil-inflamed skin model (Akiyama, H., H. Kanzaki, Y. Abe, J. Tada and J. Arata (1994). "Staphylococcus aureus infection on experimental croton oil-inflamed skin in mice." *Journal of Dermatological Science* 8(1): 1-10), the burnt skin model (Stieritz, D. D., A. Bondi, D. McDermott and E. B. Michaels (1982). "A burned mouse model to evaluate anti-pseudomonas activity of topical agents." *Journal of Antimicrobial Chemotherapy* 9(2): 133-140), the skin suture-wound model (McRipley, R. J. and R. R. Whitney (1976). "Characterization and Quantitation of Experimental Surgical-Wound Infections Used to Evaluate Topical Antibacterial Agents." *Antimicrobial Agents and Chemotherapy* 10(1): 38-44), the skin tape-stripping model (Kugelberg, E., T. Norström, T. K. Petersen, T. Duvold, D. I. Andersson and D. Hughes (2005). "Establishment of a Superficial Skin Infection Model in Mice by Using *Staphylococcus aureus* and *Streptococcus pyogenes*." *Antimicrobial Agents and Chemotherapy* 49(8): 3435-3441) and the linear full thickness scalpel cut method (Guo, Y., R. I. Ramos, J. S. Cho, N. P. Donegan, A. L. Cheung and L. S. Miller (2013). "In Vivo Bioluminescence Imaging To Evaluate Systemic and Topical Antibiotics against Community-Acquired Methicillin-Resistant *Staphylococcus aureus*-Infected Skin Wounds in Mice." *Antimicrobial Agents and Chemotherapy* 57(2): 855-863)

[00388] Preliminary studies prior to the conduct of the current study established a new method of skin infection arising from a detailed study of the models mentioned above. Briefly, study mice are anaesthetised, a patch of dorsal skin is clipped to reveal the skin and a circular area of skin is removed with a hand held punch, leaving a wound on the dorsum with a central cavity. The wound is infected with a known number of the challenge organism. Approximately four to six hours after infection, the wound is either treated topically with a vehicle formulation or an active formulation. The infected skin wound is retreated every 12 hours for a total of 14 treatments. Mice are humanely euthanased, the area of the original infected wound is dissected and removed and its bacterial content quantified by standard microbiologic tests. In this way, the change in bacterial concentration due to treatment with the active formulation can be readily determined by examining the reduction in bacterial burden compared with the vehicle control.

Materials and Methods

Preparation of Infection Inoculum

[00389] Fresh cultures of bacteria (*Staphylococcus aureus*) were grown on Sheep Blood Agar at 37 °C for 16 - 18 hours. A few typical colonies were selected and suspended in 10 ml of Tryptic Soy Broth and incubated overnight in a shaking incubator (240 rpm) at 37 °C. The overnight suspension was vortexed and diluted (1:100) in fresh Tryptic soy broth (100 µl [0.1 ml] in 9.9 ml broth). The fresh suspension was incubated for 3 hours in a shaking incubator (as above) in order to obtain mid-logarithmic phase bacteria. Bacteria were pelleted through centrifugation at 7,500 rpm for 10 mins. Broth supernatant was removed and bacteria suspended in 10 ml Phosphate Buffered Saline (PBS). These steps were repeated a further two times. The density of the suspension was checked by measuring absorbance at 600 nm, using a spectrophotometer with saline as a blank, to confirm the target density at a reading of approximately 0.100, consistent with a bacterial density of 2.5×10^7 CFU/ml. The suspension was placed into a rack placed into a lockable transport box with an ice brick to maintain refrigeration during transport, followed by storage in cool room upon arrival at the mouse skin infection laboratory. Final suspension was mixed thoroughly before inoculating the skin wounds created in mice.

[00390] In order to ensure the purity and accuracy of the suspension, the following steps were performed prior to placement into lock box.

[00391] Purity of bacterial suspension ensured by spreading 100 µl of the final suspension onto a SBA (sheep blood agar) plate which was incubated at 37 °C for 18 hours and examined to confirm uniform growth of one colony type. Viable counts were performed on final suspension by prepping saline in Eppendorf tubes (approximately 900 µl per tube), removing 100 µl sample and adding to first Eppendorf tube, vortexing the mixture and repeating using 2nd Eppendorf tube containing saline. This process was continued for 5 - 6 tubes. Finally, 100 µl of 5th and 6th dilutions were plated out on plate count agar, incubated at 37 °C for 18 hours and colony counts performed to confirm that the CFU/ml was approximately 2.5×10^7 . Following inoculation of the wounds, this process was repeated to ensure that no contamination or decrease in viable counts had occurred during the time of the surgery.

Skin Wound Surgical Procedure

[00392] Each mouse was placed into induction chamber and anaesthesia induced using 2% isoflurane. Eyes of each anaesthetised mouse were covered with veterinary eye lubricant in order to prevent corneal dehydration. Each mouse removed from induction chamber and placed onto surgical area, in front of individual aesthetic nose cone. While under anaesthesia each

mouse was monitored for assessment of depth of anaesthesia (response to pain, blink reflex, skeletal muscle tone) and respiratory and cardiac function. Back skin hair was shaved from surgical area with mechanical clippers. Shaved area was cleaned using 70% ethanol applied to paper towel followed by 10% w/v povidone-iodine solution. Once the iodine solution was dry, a subcutaneous injection of the nonsteroidal anti-inflammatory agent meloxicam was administered. Dorsal skin was pinched gently to allow creation of a circular full-thickness wound using ear punch/biopsy punch. Vehicle control and NCL812 and NCL099 mice had wounds inoculated with 10 µl of bacterial suspension using a micropipette (2.5×10^5 CFU/10 µl). Once the bacterial suspension was dry, mice were placed into individual recovery boxes labelled with the mouse number. The time of inoculation was recorded. Initial body weights of each mouse were recorded on the appropriate score sheet. Mice recovered to full consciousness within 5 minutes. Recovered mice were returned to individual housing and monitored hourly for post-surgical or anaesthetic complications.

Post-Surgical Care (4 hours post-surgery)

[00393] Mice were assessed for post-surgical complications and observations were recorded on clinic record sheet. Each mouse was carefully removed from IVC and placed into an assessment container, avoiding excessive handling or touching of the surgical site. Once the mouse was inside assessment container, it was assessed and the observations recorded on the post-surgical clinical record sheet. Whenever the suggested wellness breakpoints were reached, post-operative analgesia was administered and recorded on the clinical record sheet.

Animal Monitoring and Daily Care

[00394] Antibiotic Administration (7am and 6pm). The first administration of vehicle or NCL812 or NCL099 ointment occurred 4 hours post-surgically. Each ointment container was weighted prior to administration and the weight recorded. Each mouse was carefully restrained. Ointment (vehicle or NCL812 or NCL099) was applied to the lesion area and the treated mouse was returned to IVC where each mouse was observed to ensure ointment was not immediately removed by grooming. The weight of the ointment container post-administration was recorded. The vehicle and active NCL products were applied to the skin wound each 12 hours following the first administration for a total of 14 consecutive treatments. Both the NCL812 and NCL099 products (Formulation B, as presented in Example 9) contained their respective active ingredients at a concentration of 20 mg/g. Approximately 0.1-0.2 g of ointment was applied on each occasion, delivering a total topical dose of NCL812 or NCL099 between 28 and 56 mg to mice weighing between 18 g and 25 g.

[00395] Daily Monitoring. Monitoring of each mouse took place once daily at around 12pm. Each mouse carefully removed from IVC and placed into observation container, avoiding excessive handling or touching surgical site. The coat, posture, eyes, behaviour, vocalisation and activity whilst in the container were carefully assessed and observations recorded on assessment sheet. Mouse faeces (either on floor of cage or in container) were checked for consistency and observations recorded. The weight of each mouse was determined whilst it was in the container and change in body weight calculated and recorded. The observation container was disinfected with ethanol and set aside to dry while a fresh container was used for the next mouse. For every second day, mice were again anaesthetised using 2% isoflurane and photographed using a ruler for size referencing. These photos were used to assess lesion size and infection progression during the trial period.

Tissue analysis and assessment of antibacterial efficacy

[00396] At the end of the 7 day skin wound assessment period, all test mice were euthanased prior to wound collection for post mortem examination. The skin wound was dissected from the dorsum of each mouse. The sample was placed in a sample tube and weighed before 1 ml PBS and sterile tissue homogenisation beads were added. Tissue samples were homogenised for 10 mins using a tissue homogeniser (Next Advance Bullet Blender) and then vortexed for approximately 30 seconds. 100 µl of supernatant was removed and placed into an Eppendorf tube containing 900 µl of PBS. This procedure was repeated using serial dilutions for a total of 8 dilutions. Finally, 100 µl of each dilution was pipetted onto a plate count agar in duplicate and incubated overnight at 37 °C. Ten microlitres of original suspension was placed onto sheep blood agar to assess culture purity and incubated overnight at 37 °C. The following day, viable counts were performed using incubated plate count agar plates and the identity of *Staphylococcus aureus* (the challenge organisms) as the harvested strain was confirmed.

Results

[00397] The mean colony count per g of tissue observed in vehicle treated group was 5,888,436 (6.77 log₁₀). The mean colony count per g of tissue observed in NCL812 group was 141,254 (5.15 log₁₀). The mean colony count per g of tissue observed in NCL099 treated mice was 1,318 (3.12log₁₀). The log₁₀ colony forming units per gram of tissue and % reduction are summarised in the following table.










Table 47: Log₁₀ colony forming units per gram of tissue and percentage reduction following topical administration of vehicle and treatment.

Treatment	Log ₁₀ (CFU/g)	%reduction
Vehicle	6.77	
NCL812	5.15	97.6
NCL099	3.12	99.98









[00398] It is clear from this table that treatment with either NCL812 or NCL099 leads to high level reductions in the number of infecting *Staphylococcus aureus*. These results demonstrate effective treatment of a bacterial colonisation or infection *in vivo* by administration of compounds of the invention.










EXAMPLE 13: List of Compounds and NCL Numbers.

[00399] Examples of compounds of the present disclosure are provided below with their corresponding NCL numbers.


NCL Code	Compound Name	Structure
NCL812	2,2'-bis[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL001	2,2'-bis[(4-chlorophenyl)methylene]carbonic dihydrazide	
NCL002	2,2'-bis[(2-chlorophenyl)methylene]carbonic dihydrazide	
NCL003	2,2'-bis[(4-fluorophenyl)methylene]carbonic dihydrazide	
NCL004	2,2'-bis[(3-fluorophenyl)methylene]carbonic dihydrazide	
NCL005	2,2'-bis[(2-fluorophenyl)methylene]carbonic dihydrazide	
NCL006	2,2'-bis[(4-methoxyphenyl)methylene]carbonic dihydrazide	
NCL007	2,2'-bis[(4-cyanophenyl)methylene]carbonic dihydrazide	
NCL008	2,2'-bis[(2-cyanophenyl)methylene]carbonic dihydrazide	

NCL009	2,2'-bis[(3-cyanophenyl)methylene]carbonic dihydrazide	
NCL010	2,2'-bis[(3-methoxyphenyl)methylene]carbonic dihydrazide	
NCL011	2,2'-bis[(3-(trifluoromethyl)phenyl)methylene]carbonic dihydrazide	
NCL012	2,2'-bis[(4-(trifluoromethyl)phenyl)methylene]carbonic dihydrazide	
NCL013	2,2'-bis[(2-(trifluoromethyl)phenyl)methylene]carbonic dihydrazide	
NCL014	2-[(4-chlorophenyl)methylene]hydrazinecarboxamide	
NCL015	2-[(2-chlorophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL016	2-[(2-fluorophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL017	2-[(3-fluorophenyl)methylene]hydrazinecarboximidamide hydrochloride	


NCL018	2,2'-bis[(2-methylphenyl)methylene]carbonic dihydrazide	
NCL019	2,2'-bis[(3-methylphenyl)methylene]carbonic dihydrazide	
NCL020	2,2'-bis[(2-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL021	2,2'-bis[(4-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL022	2,2'-bis[(2-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL023	2,2'-bis[(3-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL024	2,2'-bis[(4-cyanophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL025	2,2'-bis[(2-cyanophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL026	2,2'-bis[(3-cyanophenyl)methylene]carbonimidic dihydrazide hydrochloride	

NCL027	2,2'-bis[(4-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL028	2,2'-bis[(2-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL029	2,2'-bis[(3-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL030	2-[(4-chlorophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL031	2-[(4-cyanophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL032	2-[(2-cyanophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL033	2-[(3-cyanophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL034	2-[(2-methoxyphenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL035	2,2'-bis[4-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride	


NCL036 2,2'-bis[[2-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride




NCL037 2,2'-bis[[3-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride




NCL038 2,2'-bis[[4-methylphenyl]methylene]carbonimidic dihydrazide hydrochloride




NCL039 2,2'-bis[[2-methylphenyl]methylene]carbonimidic dihydrazide hydrochloride




NCL040 2,2'-bis[[3-methylphenyl]methylene]carbonimidic dihydrazide hydrochloride




NCL041 2-[[4-(trifluoromethyl)phenyl]methylene]hydrazinecarboximidamide hydrochloride




NCL042 2-[[2-(trifluoromethyl)phenyl]methylene]hydrazinecarboximidamide hydrochloride




NCL043 2-[[3-(trifluoromethyl)phenyl]methylene]hydrazinecarboximidamide hydrochloride




NCL044 2-[[4-methylphenyl]methylene]hydrazinecarboximidamide hydrochloride




NCL045 2-[(2-methylphenyl)methylene]
hydrazinecarboximidamide hydrochloride




NCL046 2-[(3-methylphenyl)methylene]
hydrazinecarboximidamide hydrochloride




NCL047 2-[(2-chlorophenyl)methylene]
hydrazinecarboxamide




NCL048 2-[(2-fluorophenyl)methylene]
hydrazinecarboxamide




NCL049 2-[(4-cyanophenyl)methylene]
hydrazinecarboxamide




NCL050 2-[(2-cyanophenyl)methylene]
hydrazinecarboxamide




NCL051 2-[(3-cyanophenyl)methylene]
hydrazinecarboxamide












NCL052 2-[(3-chlorophenyl)methylene]
hydrazinecarboximidamide hydrochloride





















NCL053 2-[(4-fluorophenyl)methylene]
hydrazinecarboximidamide hydrochloride





















NCL054	2,2'-bis[(3-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL055	2-[(3-chlorophenyl)methylene]hydrazinecarboxamide	
NCL056	2-[(4-fluorophenyl)methylene]hydrazinecarboxamide	
NCL057	2-[(3-fluorophenyl)methylene]hydrazinecarboxamide	
NCL058	2-[(4-methoxyphenyl)methylene]hydrazinecarboxamide	
NCL059	2-[(3-methoxyphenyl)methylene]hydrazinecarboxamide	
NCL060	2-[[2-(trifluoromethyl)phenyl]methylene]hydrazinecarboxamide	
NCL061	2,2'-bis[1-[4-(trifluoromethyl)phenyl]ethylidene]carbonimidic dihydrazide hydrochloride	
NCL062	2,2'-bis[1-[4-chlorophenyl]ethylidene]carbonimidic dihydrazide hydrochloride	

C1=CC=C(C=C1)C(=O)N1C(=O)N(C1)C(=O)N2C(=O)N(C2)C(=O)N3C(=O)N(C3)C(=O)N4C(=O)N(C4)C(=O)N5C(=O)N(C5)C(=O)N6C(=O)N(C6)C(=O)N7C(=O)N(C7)C(=O)N8C(=O)N(C8)C(=O)N9C(=O)N(C9)C(=O)N10C(=O)N(C10)C(=O)N11C(=O)N(C11)C(=O)N12C(=O)N(C12)C(=O)N13C(=O)N(C13)C(=O)N14C(=O)N(C14)C(=O)N15C(=O)N(C15)C(=O)N16C(=O)N(C16)C(=O)N17C(=O)N(C17)C(=O)N18C(=O)N(C18)C(=O)N19C(=O)N(C19)C(=O)N20C(=O)N(C20)C(=O)N21C(=O)N(C21)C(=O)N22C(=O)N(C22)C(=O)N23C(=O)N(C23)C(=O)N24C(=O)N(C24)C(=O)N25C(=O)N(C25)C(=O)N26C(=O)N(C26)C(=O)N27C(=O)N(C27)C(=O)N28C(=O)N(C28)C(=O)N29C(=O)N(C29)C(=O)N30C(=O)N(C30)C(=O)N31C(=O)N(C31)C(=O)N32C(=O)N(C32)C(=O)N33C(=O)N(C33)C(=O)N34C(=O)N(C34)C(=O)N35C(=O)N(C35)C(=O)N36C(=O)N(C36)C(=O)N37C(=O)N(C37)C(=O)N38C(=O)N(C38)C(=O)N39C(=O)N(C39)C(=O)N40C(=O)N(C40)C(=O)N41C(=O)N(C41)C(=O)N42C(=O)N(C42)C(=O)N43C(=O)N(C43)C(=O)N44C(=O)N(C44)C(=O)N45C(=O)N(C45)C(=O)N46C(=O)N(C46)C(=O)N47C(=O)N(C47)C(=O)N48C(=O)N(C48)C(=O)N49C(=O)N(C49)C(=O)N50C(=O)N(C50)C(=O)N51C(=O)N(C51)C(=O)N52C(=O)N(C52)C(=O)N53C(=O)N(C53)C(=O)N54C(=O)N(C54)C(=O)N55C(=O)N(C55)C(=O)N56C(=O)N(C56)C(=O)N57C(=O)N(C57)C(=O)N58C(=O)N(C58)C(=O)N59C(=O)N(C59)C(=O)N60C(=O)N(C60)C(=O)N61C(=O)N(C61)C(=O)N62C(=O)N(C62)C(=O)N63C(=O)N(C63)C(=O)N64C(=O)N(C64)C(=O)N65C(=O)N(C65)C(=O)N66C(=O)N(C66)C(=O)N67C(=O)N(C67)C(=O)N68C(=O)N(C68)C(=O)N69C(=O)N(C69)C(=O)N70C(=O)N(C70)C(=O)N71C(=O)N(C71)C(=O)N72C(=O)N(C72)C(=O)N73C(=O)N(C73)C(=O)N74C(=O)N(C74)C(=O)N75C(=O)N(C75)C(=O)N76C(=O)N(C76)C(=O)N77C(=O)N(C77)C(=O)N78C(=O)N(C78)C(=O)N79C(=O)N(C79)C(=O)N80C(=O)N(C80)C(=O)N81C(=O)N(C81)C(=O)N82C(=O)N(C82)C(=O)N83C(=O)N(C83)C(=O)N84C(=O)N(C84)C(=O)N85C(=O)N(C85)C(=O)N86C(=O)N(C86)C(=O)N87C(=O)N(C87)C(=O)N88C(=O)N(C88)C(=O)N89C(=O)N(C89)C(=O)N90C(=O)N(C90)C(=O)N91C(=O)N(C91)C(=O)N92C(=O)N(C92)C(=O)N93C(=O)N(C93)C(=O)N94C(=O)N(C94)C(=O)N95C(=O)N(C95)C(=O)N96C(=O)N(C96)C(=O)N97C(=O)N(C97)C(=O)N98C(=O)N(C98)C(=O)N99C(=O)N(C99)C(=O)N100C(=O)N(C100)C(=O)N101C(=O)N(C101)C(=O)N102C(=O)N(C102)C(=O)N103C(=O)N(C103)C(=O)N104C(=O)N(C104)C(=O)N105C(=O)N(C105)C(=O)N106C(=O)N(C106)C(=O)N107C(=O)N(C107)C(=O)N108C(=O)N(C108)C(=O)N109C(=O)N(C109)C(=O)N110C(=O)N(C110)C(=O)N111C(=O)N(C111)C(=O)N112C(=O)N(C112)C(=O)N113C(=O)N(C113)C(=O)N114C(=O)N(C114)C(=O)N115C(=O)N(C115)C(=O)N116C(=O)N(C116)C(=O)N117C(=O)N(C117)C(=O)N118C(=O)N(C118)C(=O)N119C(=O)N(C119)C(=O)N120C(=O)N(C120)C(=O)N121C(=O)N(C121)C(=O)N122C(=O)N(C122)C(=O)N123C(=O)N(C123)C(=O)N124C(=O)N(C124)C(=O)N125C(=O)N(C125)C(=O)N126C(=O)N(C126)C(=O)N127C(=O)N(C127)C(=O)N128C(=O)N(C128)C(=O)N129C(=O)N(C129)C(=O)N130C(=O)N(C130)C(=O)N131C(=O)N(C131)C(=O)N132C(=O)N(C132)C(=O)N133C(=O)N(C133)C(=O)N134C(=O)N(C134)C(=O)N135C(=O)N(C135)C(=O)N136C(=O)N(C136)C(=O)N137C(=O)N(C137)C(=O)N138C(=O)N(C138)C(=O)N139C(=O)N(C139)C(=O)N140C(=O)N(C140)C(=O)N141C(=O)N(C141)C(=O)N142C(=O)N(C142)C(=O)N143C(=O)N(C143)C(=O)N144C(=O)N(C144)C(=O)N145C(=O)N(C145)C(=O)N146C(=O)N(C146)C(=O)N147C(=O)N(C147)C(=O)N148C(=O)N(C148)C(=O)N149C(=O)N(C149)C(=O)N150C(=O)N(C150)C(=O)N151C(=O)N(C151)C(=O)N152C(=O)N(C152)C(=O)N153C(=O)N(C153)C(=O)N154C(=O)N(C154)C(=O)N155C(=O)N(C155)C(=O)N156C(=O)N(C156)C(=O)N157C(=O)N(C157)C(=O)N158C(=O)N(C158)C(=O)N159C(=O)N(C159)C(=O)N160C(=O)N(C160)C(=O)N161C(=O)N(C161)C(=O)N162C(=O)N(C162)C(=O)N163C(=O)N(C163)C(=O)N164C(=O)N(C164)C(=O)N165C(=O)N(C165)C(=O)N166C(=O)N(C166)C(=O)N167C(=O)N(C167)C(=O)N168C(=O)N(C168)C(=O)N169C(=O)N(C169)C(=O)N170C(=O)N(C170)C(=O)N171C(=O)N(C171)C(=O)N172C(=O)N(C172)C(=O)N173C(=O)N(C173)C(=O)N174C(=O)N(C174)C(=O)N175C(=O)N(C175)C(=O)N176C(=O)N(C176)C(=O)N177C(=O)N(C177)C(=O)N178C(=O)N(C178)C(=O)N179C(=O)N(C179)C(=O)N180C(=O)N(C180)C(=O)N181C(=O)N(C181)C(=O)N182C(=O)N(C182)C(=O)N183C(=O)N(C183)C(=O)N184C(=O)N(C184)C(=O)N185C(=O)N(C185)C(=O)N186C(=O)N(C186)C(=O)N187C(=O)N(C187)C(=O)N188C(=O)N(C188)C(=O)N189C(=O)N(C189)C(=O)N190C(=O)N(C190)C(=O)N191C(=O)N(C191)C(=O)N192C(=O)N(C192)C(=O)N193C(=O)N(C193)C(=O)N194C(=O)N(C194)C(=O)N195C(=O)N(C195)C(=O)N196C(=O)N(C196)C(=O)N197C(=O)N(C197)C(=O)N198C(=O)N(C198)C(=O)N199C(=O)N(C199)C(=O)N200C(=O)N(C200)C(=O)N201C(=O)N(C201)C(=O)N202C(=O)N(C202)C(=O)N203C(=O)N(C203)C(=O)N204C(=O)N(C204)C(=O)N205C(=O)N(C205)C(=O)N206C(=O)N(C206)C(=O)N207C(=O)N(C207)C(=O)N208C(=O)N(C208)C(=O)N209C(=O)N(C209)C(=O)N210C(=O)N(C210)C(=O)N211C(=O)N(C211)C(=O)N212C(=O)N(C212)C(=O)N213C(=O)N(C213)C(=O)N214C(=O)N(C214)C(=O)N215C(=O)N(C215)C


NCL072	2-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL073	2-[[1-(4-trifluoromethyl)phenyl]ethylidene] carbonimidic dihydrazide hydrochloride	
NCL074	2-[(2,4-dichlorophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL075	2-{1-[4-(trifluoromethyl)phenyl]ethylidene}-2'-[[4-(trifluoromethyl)phenyl]methylene] carbonimidic dihydrazide hydrochloride	
NCL076	2-[(4-chlorophenyl)methylene]-2'-[(4-methylphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL077	2-[(4-chlorophenyl)methylene]-2'-[[4-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride	
NCL078	2-[[2-fluoro-4-(trifluoromethyl)phenyl]methylene]-2'-[[4-chlorophenyl]methylene] carbonimidic dihydrazide hydrochloride	
NCL079	2-[(4-chlorophenyl)methylene]-2'-[(4-fluorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL080	2-{1-[4-(trifluoromethyl)phenyl]ethylidene}-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	

NCL081	2-[1-(4-chlorophenyl)ethylidene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL082	2-[(2-chlorophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL083	2-[(3-chlorophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL084	2-[(2-fluoro-4-chlorophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL085	2-[(2-cyanophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL086	2-[(3-cyanophenyl)methylene]-2'-[(4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL087	2-[(4-chlorophenyl)methylene]-2'-[(4-cyanophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL088	2-[(2-fluorophenyl)methylene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL089	2-[1-(4-chlorophenyl)ethylidene]-2'-[1-[4-(trifluoromethyl)phenyl]ethylidene] carbonimidic dihydrazide hydrochloride	


NCL090	<i>N</i> -benzoyl-1-benzoyl-2-[(2-chlorophenyl)methylene]hydrazinecarboximidamide hydrochloride	
NCL091	<i>N</i> -(4-chlorophenyl)-3-[4-(trifluoromethyl)phenyl]-1 <i>H</i> -1,2,4-triazol-5-amine	
NCL092	<i>N</i> -(4-chlorobenzyl)-3-(4-chlorophenyl)-1 <i>H</i> -1,2,4-triazol-5-amine	
NCL093	2,2'-bis(2-naphthalenylmethylene)carbonimidic dihydrazide hydrochloride	
NCL094	2,2'-bis(cyclohexylmethylene)carbonimidic dihydrazide hydrochloride	
NCL095	2,2'-bis(3-furanylmethylene)carbonimidic dihydrazide hydrochloride	
NCL096	2,2'-bis(3-phenyl-2-propenylidene)carbonimidic dihydrazide hydrochloride	
NCL097	2,2'-bis[(3,4,5-trihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride	
NCL098	2,2'-bis[(3-carboxyphenyl)methylene]carbonimidic dihydrazide hydrochloride	

NCL099	2,2'-bis[4-(1,1-dimethylethyl)phenyl]methylene] carbonimidic dihydrazide hydrochloride	
NCL100	2,2'-bis(phenylmethylene) carbonimidic dihydrazide hydrochloride	
NCL101	2,2'-bis[(2,3-dihydroxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL102	2,2'-bis[(2-nitrophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL103	2,2'-bis[(2,4-dihydroxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL104	2,2'-bis[(2,4,5-trihydroxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL105	2,2'-bis[(2,3,4-trihydroxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL106	2,2'-bis[(4,5-dihydroxy-3-methoxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL107	2,2'-bis[(2-hydroxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	


NCL108 2,2'-bis[(3-hydroxyphenyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL109 2,2'-bis[(3-nitrophenyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL110 2,2'-bis[(4-nitrophenyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL111 2,2'-bis[(3,4-dihydroxyphenyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL112 2,2'-bis[(1,1'-biphenyl)-4-ylmethylene]
carbonimidic dihydrazide hydrochloride




NCL113 2,2'-bis[4-(dimethylamino)phenyl]methylene]
carbonimidic dihydrazide hydrochloride




NCL114 2,2'-bis[(3,5-dichlorophenyl)methylene]
carbonimidic dihydrazide hydrochloride





















NCL 115 2,2'-bis[(3,4-dimethoxyphenyl)methylene]
carbonimidic dihydrazide hydrochloride












NCL 116 2,2'-bis[(1,1'-biphenyl)-2-ylmethylene]
carbonimidic dihydrazide hydrochloride




NCL 117	2,2'-bis[(4-hydroxy-3-methoxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 118	2,2'-bis[(2,5-fluorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 119	2,2'-bis[(4-acetamidophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 120	2,2'-bis[(4-(1-methylethyl)phenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 121	2,2'-bis[(4-propylphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 122	2,2'-bis[(4-hydroxy-3-nitrophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 123	2,2'-bis[(3,4-difluorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 124	2,2'-bis[(2-hydroxy-1-naphthalenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 125	2,2'-bis[(3-hydroxy-4-methoxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	


NCL 126	2,2'-bis[(3-ethynylphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 127	2,2'-bis[(2,4-dichlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 128	2,2'-bis[(2,3,4,5,6-pentafluorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 129	2,2'-bis[(2-bromophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 130	2,2'-bis[(3-bromo-4,5-dimethoxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 131	2,2'-bis[(3-bromophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 132	2,2'-bis[(4-chloro-6-fluoro-2H-1-benzopyran-3-yl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 133	2,2'-bis[(4-bromo-2-furanyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 134	2,2'-bis[(4-bromophenyl)methylene] carbonimidic dihydrazide hydrochloride	

NCL 135	2,2'-bis[(2-bromo-4,5-dimethoxyphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 136	2,2'-bis[(4-butylphenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 137	2,2'-bis[(2,6-dichlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 138	2,2'-bis(2,3-diphenyl-2-propenylidene) carbonimidic dihydrazide hydrochloride	
NCL 139	2,2'-bis(3-quinolyl)methylene carbonimidic dihydrazide hydrochloride	
NCL 140	2,2'-bis[4-(methylsulfanyl)phenyl]methylene carbonimidic dihydrazide hydrochloride	
NCL 141	2,2'-bis[(5-chlorobenzo[b]thien-3-yl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 142	1,3-bis(benzylamino)guanidine hydrochloride	
NCL 143	2,2'-bis[1-phenylethylidene] carbonimidic dihydrazide hydrochloride	


NCL 144 2,2'-bis[(5-bromo-2-furanyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL 145 2,2'-bis[(5-chloro-2-furanyl)methylene]
carbonimidic dihydrazide hydrochloride




NCL 146 2,2'-bis(1*H*-indol-5-yl)methylene
carbonimidic dihydrazide hydrochloride




NCL 147 2,2'-bis(2-quinoxaliny)methylene
carbonimidic dihydrazide hydrochloride




NCL 148 2,2'-bis[[4-(carboxypropenyl)phenyl]methylene]
carbonimidic dihydrazide hydrochloride




NCL 149 2,2'-Bis(4-pyridinyl)methylene
Carbonimidic dihydrazide hydrochloride




NCL 150 2,2'-bis[3-(4-methoxyphenyl)-2-
propenylidene]carbonimidic dihydrazide
hydrochloride




NCL 151 2,2'-bis[(4-hydroxyphenyl)methylene]
carbonimidic dihydrazide hydrochloride






















NCL 152 2,2'-bis[(2-hydroxy-3-methylphenyl)methylene]
carbonimidic dihydrazide hydrochloride



NCL 153 2,2'-bis[1-(4-chlorophenyl)propylidene]
carbonimidic dihydrazide hydrochloride



NCL 154	2,2'-bis[1-(4-chlorophenyl)pentylidene] carbonimidic dihydrazide hydrochloride	
NCL 155	2,2'-bis[1-(4-bromophenyl)ethylidene] carbonimidic dihydrazide hydrochloride	
NCL 156	2,2'-bis[1-(4-chlorophenyl)butylidene] carbonimidic dihydrazide hydrochloride	
NCL 157	2,2'-bis[(2-amino-4-chlorophenyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 158	2,2'-bis[1-(2-hydroxy-4-chlorophenyl)propylidene] carbonimidic dihydrazide hydrochloride	
NCL 159	2,2'-bis[(2-hydroxy-4-chlorophenyl)(cyclopentyl)methylene] carbonimidic dihydrazide hydrochloride	
NCL 160	2,2'-bis[[4-(trifluoromethoxy)phenyl]methylene] carbonimidic dihydrazide hydrochloride	
NCL 161	2,2'-bis[1-(4-piperazinylphenyl)ethylidene] carbonimidic dihydrazide hydrochloride	
NCL 162	ethyl (E)-N'-((E)-1-(4-chlorophenyl)ethylidene) carbamohydrazone	









NCL 163	2,2'-bis(2-Oxo-1,2-dihydro-3H-indol-3-ylidene)carbonimidic dihydrazide hydrochloride	
NCL 164	2,2'-bis[1-(2-amino-4-chlorophenyl)ethylidene]carbonimidic dihydrazide hydrochloride	
NCL 165	2,2'-bis(1-phenyl-2-aminoethylidene)carbonimidic dihydrazide trihydrochloride	
NCL 166	2,2'-bis[[4-(trifluoromethylsulfanyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride	
NCL 167	2,2'-bis(phenylcarboxymethylene)carbonimidic dihydrazide hydrochloride	
NCL 168	2,2'-bis[[2-(1-hydroxyethylamino)-4-chlorophenyl]methylene]carbonimidic dihydrazide hydrochloride	
NCL 169	2,2'-bis[(2-amino-4-chlorophenyl)methylene]carbonimidic dihydrazide	
NCL 170	2,2'-bis[(2-acetamido-4-chlorophenyl)methylene]carbonimidic dihydrazide	
NCL 171	2,2'-bis[[4-(dimethylamino)-2-hydroxyphenyl]methylene]carbonimidic dihydrazide	
NCL 172	2,2'-Bis[1-(2-pyridinyl)ethylidene]Carbonimidic dihydrazide hydrochloride	

NCL 173	2,2'-bis[1-(4-chloro-2-hydroxyphenyl)ethylidene] carbonimidic dihydrazide hydrochloride	
NCL 174	2,2'-bis(4-chloro-2-hydroxyphenylmethylene) carbonimidic dihydrazide hydrochloride	
NCL 175	2,2'-Bis(4-chloropyridin-3-ylmethylene)Carbonimidic dihydrazide hydrochloride	
NCL 176	2,2'-Bis(2-aminopyridin-3-ylmethylene) Carbonimidic dihydrazide hydrochloride	
NCL 177	2,2'-bis[1-(4-chlorophenyl)-2-hydroxyethylidene] carbonimidic dihydrazide hydrochloride	
NCL 178	4,6-bis(2-((E)-1-(4-chlorophenyl)ethylidene)hydrazinyl)pyrimidine	
NCL 179	4,6-bis(2-((E)-4-chlorobenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL 180	(2Z,2'Z)-2,2'-(pyrimidine-4,6-diylbis(hydrazin-2-yl-1-ylidene))bis(2-(4-chlorophenyl)ethan-1-ol)	
NCL 181	4,6-bis(2-((E)-4-chlorobenzylidene)hydrazinyl)pyrimidine	

NCL 182	6-chloro-N ⁴ -(1-phenylethyl)pyrimidine-2,4-diamine	
NCL 183	N ⁴ ,N ⁶ -bis(1-phenylethyl)pyrimidine-4,6-diamine	
NCL 184	N ⁴ ,N ⁶ -bis(1-phenylethyl)pyrimidine-2,4,6-triamine	
NCL 185	4,6-bis(2-((E)-1-(4-chlorophenyl)ethylidene)hydrazinyl)pyrimidine hydrochloride	
NCL 186	N ² ,N ⁵ -bis(1-(4-chlorophenyl)ethyl)pyrazine-2,5-diamine	
NCL 187	4,6-bis(2-((E)-4-chlorobenzylidene)hydrazinyl)pyrimidine hydrochloride	
NCL 188	(E)-2-(1-(4-chlorophenyl)pentylidene)hydrazine-1-carboximidamide hydrochloride	
NCL 189	(2Z,2'Z)-2,2'-(pyrimidine-4,6-diylbis(hydrazin-2-yl-1-ylidene))bis(2-(4-chlorophenyl)ethan-1-ol) hydrochloride	
NCL 190	(Z)-2-(1-(4-chlorophenyl)-2-hydrazinylethylidene)hydrazine-1-carboximidamide hydrochloride	


NCL 191	(E)-2-(1-(4-chlorophenyl)ethylidene)hydrazine-1-carboximidamide hydrochloride	
NCL 192	(Z)-2-(2-carbamimidoylhydrazono)-2-phenylacetic acid hydrochloride	
NCL193	4,6-bis(2-((E)-4-bromobenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL194	N',N'''-(2-aminopyrimidine-4,6-diyl)di(benzohydrazide)	
NCL195	4,6-bis(2-((E)-4-methylbenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL196	4,4'-((1E,1'E)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	
NCL197	3,3'-((1E,1'E)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	
NCL198	4,6-bis(2-((E)-4-(tert-butyl)benzylidene)hydrazinyl)pyrimidin-2-amine dihydrochloride	
NCL199	4,6-bis(2-((E)-benzylidene)hydrazinyl)pyrimidin-2-amine	

NCL200	4,6-bis(2-((E)-4-(tert-butyl)benzylidene)hydrazinyl)pyrimidin-2-amine	
NCL201	4,4'-((1E,1'E)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))bis(N,N-dimethylaniline)	
NCL202	2,2'-((1E,1'E)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	
NCL203	4,6-bis(2-((E)-cyclohexylmethylene)hydrazinyl)pyrimidin-2-amine dihydrochloride	
NCL204	4,6-bis(2-((E)-1-(4-chlorophenyl)ethylidene)hydrazinyl)pyrimidin-2-amine	
NCL205	4,6-bis(2-((E)-2-chlorobenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL206	4,6-bis(2-((E)-4-methylbenzylidene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL207	2,2'-((1E,1'E)-((6-amino-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	
NCL208	4,6-bis(2-((E)-1-(4-chlorophenyl)ethylidene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL209	3,3'-((1E,1'E)-((6-amino-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	


NCL210	4,6-bis(2-((E)-4-(trifluoromethyl)benzylidene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL211	4,4'-((1E,1'E)-((6-amino-1,3,5-triazine-2,4-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))diphenol	
NCL212	4,6-bis(2-((E)-4-bromobenzylidene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL213	4,6-bis(2-((E)-cyclohexylmethylene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL214	4,6-bis(2-((E)-benzylidene)hydrazinyl)-1,3,5-triazin-2-amine	
NCL215	(E)-N'-((E)-1-(4-chloro-2-fluorophenyl)ethylidene)-2-(1-(4-chloro-2-fluorophenyl)ethylidene)hydrazine-1-carboximidhydrazide hydrochloride	
NCL216	N',2-bis((E)-4-chloro-2-fluorobenzylidene)hydrazine-1-carboximidhydrazide hydrochloride	
NCL217	N',2-bis((E)-1-(p-tolyl)ethylidene)hydrazine-1-carboximidhydrazide hydrochloride	

NCL218	4-((E)-(2-(2-amino-6-(2-((E)-4-((diethoxyphosphoryl)oxy)benzylidene)hydrazinyl)pyrimidin-4-yl)hydrazono)methyl)phenyl diethyl phosphate	
NCL219	(E)-N'-((E)-1-(4-(tert-butyl)phenyl)ethylidene)-2-(1-(4-(tert-butyl)phenyl)ethylidene)hydrazine-1-carboximidhydrazide hydrochloride	
NCL220	4,6-bis(2-((E)-4-fluorobenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL221	4,6-bis(2-((E)-4-(trifluoromethyl)benzylidene)hydrazinyl)pyrimidin-2-amine	
NCL222	4,6-bis(2-((E)-3,4-difluorobenzylidene)hydrazinyl)pyrimidin-2-amine	
NCL223	N,N'-(((1E,1'E)-((2-aminopyrimidine-4,6-diyl)bis(hydrazin-2-yl-1-ylidene))bis(methanylylidene))bis(4,1-phenylene))diacetamide	
NCL224	ethyl 2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)hydrazine-1-carboxylate	
NCL225	isobutyl 2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)hydrazine-1-carboxylate	
NCL226	2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-ethylhydrazine-1-carboxamide	


NCL227 N-benzyl-2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)hydrazine-1-carboxamide




NCL228 2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-hexylhydrazine-1-carboxamide



NCL229 2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-(furan-2-ylmethyl)hydrazine-1-carboxamide



NCL230 4,6-bis(2-((E)-4-methoxybenzylidene)hydrazinyl)pyrimidin-2-amine



CLAIMS

1. A compound selected from the group consisting of:

NCL023	2,2'-bis[(3-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL025	2,2'-bis[(2-cyanophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL026	2,2'-bis[(3-cyanophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL029	2,2'-bis[(3-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL036	2,2'-bis[[2-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride;
NCL037	2,2'-bis[[3-(trifluoromethyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride;
NCL039	2,2'-bis[(2-methylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL040	2,2'-bis[(3-methylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL061	2,2'-bis{1-[4-(trifluoromethyl)phenyl]ethylidene}carbonimidic dihydrazide hydrochloride;
NCL068	2-[1-(4-chlorophenyl)ethylidene]-2'-{1-[4-(trifluoromethyl)phenyl]methylene}carbonimidic dihydrazide hydrochloride;
NCL075	2-{1-[4-(trifluoromethyl)phenyl]ethylidene}-2'-{[4-(trifluoromethyl)phenyl]methylene}carbonimidic dihydrazide hydrochloride;
NCL076	2-[(4-chlorophenyl)methylene]-2'-[(4-methylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL078	2-{[2-fluoro-4-(trifluoromethyl)phenyl]methylene}-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL079	2-[(4-chlorophenyl)methylene]-2'-[(4-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL080	2-{1-[4-(trifluoromethyl)phenyl]ethylidene}-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL081	2-[1-(4-chlorophenyl)ethylidene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL084	2-[(2-fluoro-4-chlorophenyl)methylene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL085	2-[(2-cyanophenyl)methylene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL086	2-[(3-cyanophenyl)methylene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL088	2-[(2-fluorophenyl)methylene]-2'-[(4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL089	2-[1-(4-chlorophenyl)ethylidene]-2'-{1-[4-(trifluoromethyl)phenyl]ethylidene}carbonimidic dihydrazide hydrochloride;
NCL090	N-benzoyl-1-benzoyl-2-[(2-chlorophenyl)methylene]hydrazine carboximidamide hydrochloride;
NCL094	2,2'-bis(cyclohexylmethylene)carbonimidic dihydrazide hydrochloride;
NCL095	2,2'-bis(3-furanylmethylene)carbonimidic dihydrazide hydrochloride;
NCL097	2,2'-bis[(3,4,5-trihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL098	2,2'-bis[(3-carboxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL099	2,2'-bis{[4-(1,1-dimethylethyl)phenyl]methylene}carbonimidic dihydrazide hydrochloride;
NCL101	2,2'-bis[(2,3-dihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL104	2,2'-bis[(2,4,5-trihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;

NCL105	2,2'-bis[(2,3,4-trihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL106	2,2'-bis[(4,5-dihydroxy-3-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL108	2,2'-bis[(3-hydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL111	2,2'-bis[(3,4-dihydroxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL112	2,2'-bis[(1,1'-biphenyl]-4-ylmethylene)carbonimidic dihydrazide hydrochloride;
NCL114	2,2'-bis[(3,5-dichlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL115	2,2'-bis[(3,4-dimethoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL116	2,2'-bis[(1,1'-biphenyl]-2-ylmethylene)carbonimidic dihydrazide hydrochloride;
NCL118	2,2'-bis[(2,5-fluorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL119	2,2'-bis[(4-acetamidophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL121	2,2'-bis[(4-propylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL122	2,2'-bis[(4-hydroxy-3-nitrophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL123	2,2'-bis[(3,4-difluorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL124	2,2'-bis[(2-hydroxy-1-naphthalenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL125	2,2'-bis[(3-hydroxy-4-methoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL126	2,2'-bis[(3-ethynylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL130	2,2'-bis[(3-bromo-4,5-dimethoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL131	2,2'-bis[(3-bromophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL132	2,2'-bis[(4-chloro-6-fluoro-2H-1-benzopyran-3-yl)methylene]carbonimidic dihydrazide hydrochloride;
NCL133	2,2'-bis[(4-bromo-2-furanyl)methylene]carbonimidic dihydrazide hydrochloride
NCL135	2,2'-bis[(2-bromo-4,5-dimethoxyphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL136	2,2'-bis[(4-butylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL137	2,2'-bis[(2,6-dichlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL138	2,2'-bis(2,3-diphenyl-2-propenylidene)carbonimidic dihydrazide hydrochloride;
NCL139	2,2'-bis(3-quinolinylmethylene)carbonimidic dihydrazide hydrochloride;
NCL140	2,2'-bis[[4-(methylsulfanyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride;
NCL141	2,2'-bis[(5-chlorobenzo[b]thien-3-yl)methylene]carbonimidic dihydrazide hydrochloride;
NCL144	2,2'-bis[(5-bromo-2-furanyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL145	2,2'-bis[(5-chloro-2-furanyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL146	2,2'-bis(1H-indol-5-ylmethylene)carbonimidic dihydrazide hydrochloride;
NCL147	2,2'-bis(2-quinoxalinylmethylene)carbonimidic dihydrazide hydrochloride;
NCL148	2,2'-bis[[4-(carboxypropenyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride;
NCL150	2,2'-bis[3-(4-methoxyphenyl)-2-propenylidene]carbonimidic dihydrazide hydrochloride;
NCL152	2,2'-bis[(2-hydroxy-3-methylphenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL153	2,2'-bis[1-(4-chlorophenyl)propylidene]carbonimidic dihydrazide hydrochloride;
NCL154	2,2'-bis[1-(4-chlorophenyl)pentylidene]carbonimidic dihydrazide hydrochloride;
NCL156	2,2'-bis[1-(4-chlorophenyl)butylidene]carbonimidic dihydrazide hydrochloride;

NCL157	2,2'-bis[(2-amino-4-chlorophenyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL158	2,2'-bis[1-(2-hydroxy-4-chlorophenyl)propylidene]carbonimidic dihydrazide hydrochloride;
NCL159	2,2'-bis[(2-hydroxy-4-chlorophenyl)(cyclopentyl)methylene]carbonimidic dihydrazide hydrochloride;
NCL161	2,2'-bis[1-(4-piperazinylphenyl)ethylidene]carbonimidic dihydrazide hydrochloride;
NCL164	2,2'-bis[1-(2-amino-4-chlorophenyl)ethylidene]carbonimidic dihydrazide hydrochloride;
NCL165	2,2'-bis(1-phenyl-2-aminoethylidene)carbonimidic dihydrazidetrihydrochloride;
NCL166	2,2'-bis[[4-(trifluoromethylsulfanyl)phenyl]methylene]carbonimidic dihydrazide hydrochloride;
NCL167	2,2'-bis(phenylcarboxymethylene)carbonimidic dihydrazide hydrochloride;
NCL168	2,2'-bis{[2-(1-hydroxyethylamino)-4-chlorophenyl]methylene}carbonimidic dihydrazide hydrochloride;
NCL169	2,2'-bis[(2-amino-4-chlorophenyl)methylene]carbonimidic dihydrazide;
NCL170	2,2'-bis[(2-acetamido-4-chlorophenyl)methylene]carbonimidic dihydrazide;
NCL171	2,2'-bis{[4-(dimethylamino)-2-hydroxyphenyl]methylene}carbonimidic dihydrazide;
NCL172	2,2'-Bis[1-(2-pyridinyl)ethylidene]Carbonimidic dihydrazide hydrochloride;
NCL173	2,2'-bis[1-(4-chloro-2-hydroxyphenyl)ethylidene]carbonimidic dihydrazide hydrochloride;
NCL174	2,2'-bis(4-chloro-2-hydroxyphenylmethylene)carbonimidic dihydrazide hydrochloride;
NCL176	2,2'-Bis(2-aminopyridin-3-ylmethylene)Carbonimidic dihydrazide hydrochloride;
NCL188	(E)-2-(1-(4-chlorophenyl)pentylidene)hydrazine-1-carboximidamidehydrochloride;
NCL190	(Z)-2-(1-(4-chlorophenyl)-2-hydrazinylethylidene)hydrazine-1-carboximidamidehydrochloride;
NCL192	(Z)-2-(2-carbamimidoylhydrazono)-2-phenylaceticacidhydrochloride;
NCL193	4,6-bis(2-((E)-4-bromobenzylidene)hydrazinyl)pyrimidin-2-amine;
NCL215	(E)-N'-((E)-1-(4-chloro-2-fluorophenyl)ethylidene)-2-(1-(4-chloro-2-fluorophenyl)ethylidene)hydrazine-1-carboximidhydrazidehydrochloride;
NCL216	N',2-bis((E)-4-chloro-2-fluorobenzylidene)hydrazine-1-carboximidhydrazidehydrochloride;
NCL217	N',2-bis((E)-1-(p-tolyl)ethylidene)hydrazine-1-carboximidhydrazidehydrochloride;
NCL219	(E)-N'-((E)-1-(4-(tert-butyl)phenyl)ethylidene)-2-(1-(4-(tert-butyl)phenyl)ethylidene)hydrazine-1-carboximidhydrazidehydrochloride;
NCL226	2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-ethylhydrazine-1-carboxamide;
NCL227	N-benzyl-2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)hydrazine-1-carboxamide;
NCL228	2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-hexylhydrazine-1-carboxamide;
NCL229	2-((E)-4-chlorobenzylidene)-1-((E)-N'-((E)-4-chlorobenzylidene)carbamohydrazonoyl)-N-(furan-2-ylmethyl)hydrazine-1-carboxamide;



a stereoisomer, tautomer, pharmaceutically acceptable salt and prodrug thereof.

- The compound of claim 1, wherein the compound is selected from the group consisting of NCL146, NCL157, NCL158, NCL193, NCL216, NCL217, and NCL219.



3. The compound of claim 1 or 2, wherein the compound is a chloride salt.
4. An antibacterial pharmaceutical composition comprising a compound of any one of claims 1 to 3, or a therapeutically acceptable salt thereof, and a pharmaceutically acceptable excipient or carrier.
5. The antibacterial pharmaceutical composition of claim 4 that is an antibacterial veterinary composition.
6. The composition of claim 4 or claim 5, wherein the composition comprises a further antimicrobial agent selected from the group consisting of antibacterial agents and antifungal agents.
7. The composition of any one of claims 4 to 6, wherein the composition is formulated for oral administration, parenteral administration or topical administration.
8. Use of a compound of any one of claims 1 to 3, or a therapeutically acceptable salt thereof, in the manufacture of a medicament for the treatment of a bacterial colonisation or infection in a subject.
9. Use of a compound of any one of claims 1 to 3 in the treatment of a bacterial colonisation or infection in a subject in need thereof.
10. The use of claim 8 or 9, wherein the bacterial colonisation or infection is caused by a bacterial agent that is Gram-positive.
11. The use of claim 10, wherein the bacterial agent is selected from the group consisting of *Staphylococcus aureus*, *Staphylococcus pseudintermedius*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Enterococcus faecium*, *Enterococcus faecalis*, and *Clostridium difficile*.
12. The use of claim 8 or 9, wherein the bacterial colonisation or infection is caused by a bacterial agent that is Gram-negative.
13. The use of claim 12, wherein the bacterial agent is selected from the group consisting of *Acinetobacter* species, *Aeromonas hydrophila*, *Citrobacter* species, *Enterobacter* species, *Escherichia coli*, *Klebsiella pneumoniae*, *Morganella morganii*, *Pseudomonas aeruginosa*, *Neisseria* species and *Stenotrophomonas maltophilia*.
14. The use of claim 8 or 9, wherein the bacterial colonisation or infection is caused by a bacterial agent that has no cell wall.

15. The use of claim 14, wherein the bacterial agent is selected from the group consisting of *Mycoplasma* spp, *Mycoplasma agalactiae*, *Mycoplasma alkalescens*, *Mycoplasma amphoriforme*, *Mycoplasma arginini*, *Mycoplasma bovigenitalum*, *Mycoplasma bovirhinis*, *Mycoplasma bovis*, *Mycoplasma bovoculi*, *Mycoplasma buccale*, *Mycoplasma californicum*, *Mycoplasma canadense*, *Mycoplasma capricolum* subsp. *capricolum*, *Mycoplasma capricolum* subsp. *capripneumoniae*, *Mycoplasma conjunctivae*, *Mycoplasma cynos*, *Mycoplasma dispar*, *Mycoplasma equigenitalum*, *Mycoplasma faucium*, *Mycoplasma felis*, *Mycoplasma fermentans* (*incognitus* str.), *Mycoplasma gallisepticum*, *Mycoplasma gateae*, *Mycoplasma genitalium*, *Mycoplasma haemocanis*, *Mycoplasma haemofelis*, *Mycoplasma haemosuis* (formerly *Eperythrozoon suis*), *Mycoplasma hominis*, *Mycoplasma hyopneumoniae*, *Mycoplasma hyorhinis*, *Mycoplasma hyosynoviae*, *Mycoplasma iowae meleagridis* (MM), *Mycoplasma iowae*, *Mycoplasma leachii*, *Mycoplasma lipophilum*, *Mycoplasma meleagridis*, *Mycoplasma mycoides* subsp *capri*, *Mycoplasma mycoides* subsp *mycoides*, *Mycoplasma mycoides* subsp. *mycoides*, *Mycoplasma orale*, *Mycoplasma ovipneumoniae*, *Mycoplasma ovis*, *Mycoplasma penetrans*, *Mycoplasma pirum*, *Mycoplasma pneumoniae*, *Mycoplasma primum*, *Mycoplasma putrefaciens*, *Mycoplasma salivarium*, *Mycoplasma spermatophilum*, *Mycoplasma suis*, *Mycoplasma synoviae*, *Mycoplasma wenyonii*, *Mycoplasma*, *Ureaplasma* spp, *Ureaplasma parvum*, *Ureaplasma urealyticum*, and *Ureaplasma diversum*.
16. The use of claim 8 or 9, wherein the bacterial colonisation or infection is caused by at least two bacterial agents selected from the group consisting of Gram-positive bacteria, Gram-negative bacteria, and bacterial with no cell wall.
17. The use of claim 12 or 13, wherein the compound of any one of claims 1 to 3 is for use together with a compound that reduces the integrity of bacterial cell wall.
18. The use of any one of claims 8 to 17, wherein the subject is an animal.
19. The use of claim 18, wherein the animal is selected from the group consisting of human, canine, feline, bovine, ovine, caprine, porcine, avian, piscine and equine species.
20. The use of any one of claims 8 to 19, wherein the compound of any of claims 1 to 3 is for use orally, parenterally, or topically.



1/20



**Figure 1****Figure 2**

2/20



**Figure 3****Figure 4**

3/20



**Figure 5****Figure 6**

**Figure 7****Figure 8**



5/20

**Figure 9****Figure 10**



6/20

**Figure 11****Figure 12**



7/20

**Figure 13****Figure 14**



8/20

**Figure 15****Figure 16**



9/20

**Figure 17****Figure 18**



10/20



**Figure 19****Figure 20**

11/20



**Figure 21****Figure 22**



12/20

**Figure 23****Figure 24**



**Figure 25****Figure 26**



14/20



**Figure 27****Figure 28**



**Figure 29****Figure 30**



16/20


**Figure 31****Figure 32**

**Figure 33****Figure 34**

**Figure 35****Figure 36**

**Figure 37****Figure 38**

**Figure 39****Figure 40**



Formula 1