



(51) International Patent Classification:

*A61K 9/16* (2006.01) *A61K 48/00* (2006.01)  
*A61K 47/30* (2006.01)

(21) International Application Number:

PCT/US2014/036699

(22) International Filing Date:

2 May 2014 (02.05.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/819,517	3 May 2013 (03.05.2013)	US
61/881,851	24 September 2013 (24.09.2013)	US
61/881,921	24 September 2013 (24.09.2013)	US
61/881,913	24 September 2013 (24.09.2013)	US
61/907,177	21 November 2013 (21.11.2013)	US
61/948,313	5 March 2014 (05.03.2014)	US
61/948,384	5 March 2014 (05.03.2014)	US

(71) Applicant: **SELECTA BIOSCIENCES, INC.** [US/US];  
480 Arsenal Street, Building One, Watertown, MA 02472  
(US).

(72) Inventor: **MALDONADO, Roberto, A.**; 44 Gartland  
Street #1, Jamaica Plain, MA 02130 (US).

(74) Agent: **VATLAND, Janice, A.**; Wolf, Greenfield & Sacks,  
P.C., 600 Atlantic Avenue, Boston, MA 02210-2206 (US).

(81) Designated States (unless otherwise indicated, for every  
kind of national protection available): AE, AG, AL, AM,  
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,  
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,  
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,  
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,  
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,  
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,  
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,  
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,  
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,  
ZW.

(84) Designated States (unless otherwise indicated, for every  
kind of regional protection available): ARIPO (BW, GH,  
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,  
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,  
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,  
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,  
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,  
KM, ML, MR, NE, SN, TD, TG).

Published:

— with declaration under Article 17(2)(a); without abstract;  
title not checked by the International Searching Authority

(54) Title: TOLEROGENIC SYNTHETIC NANOCARRIERS TO REDUCE OR PREVENT ANAPHYLAXIS IN RESPONSE TO  
A NON-ALLERGENIC ANTIGEN

(57) Abstract:



WO 2014/179772 A2

**TOLEROGENIC SYNTHETIC NANOCARRIERS TO REDUCE OR PREVENT  
ANAPHYLAXIS IN RESPONSE TO A NON-ALLERGENIC ANTIGEN**

**RELATED APPLICATIONS**

5           This application claims the benefit under 35 U.S.C. §119 of United States provisional applications 61/819517, filed May 3, 2013; 61/881851, filed September 24, 2013; 61/881913, filed September 24, 2013; 61/881921, filed September 24, 2013; 61/907177, filed November 21, 2013; 61/948313, filed March 5, 2014; and 61/948384, filed March 5, 2014, the entire contents of each of which are incorporated herein by reference.

**FIELD OF THE INVENTION**

10           This invention relates to methods for reducing or preventing anaphylaxis to non-allergenic antigen with compositions comprising immunosuppressants, and related compositions. The methods and compositions allow for reduction or prevention of anaphylaxis  
15       development as a result of non-allergenic antigens, such as non-allergenic therapeutic macromolecules. Thus, methods and compositions are provided that can be used to reduce or prevent anaphylaxis due to the administration of a non-allergenic antigen, such as a non-allergenic therapeutic macromolecule.

**BACKGROUND OF THE INVENTION**

20           Conventional immunosuppressant drugs are broad-acting. Additionally, in order to maintain immunosuppression, immunosuppressant drug therapy is generally a life-long proposition. Unfortunately, the use of broad-acting immunosuppressants are associated with a risk of severe side effects, such as tumors, infections, nephrotoxicity and metabolic disorders.  
25       Accordingly, new antigen-specific tolerogenic therapies would be beneficial.

**SUMMARY OF THE INVENTION**

30           In one aspect, a method for reducing or preventing an anaphylactic reaction to a non-allergenic antigen, comprising providing the non-allergenic antigen, the non-allergenic antigen being unattached to synthetic nanocarriers; providing a composition comprising synthetic nanocarriers, in some embodiments, attached to an immunosuppressant; and administering to a subject the composition concomitantly with the non-allergenic antigen. In one embodiment of any one of the methods provided, the concomitant administration to the subject is according to a protocol that has been demonstrated to reduce or prevent an anaphylactic reaction to the non-

allergenic antigen. In another embodiment of any one of the methods provided, the method further comprises determining the protocol.

In another embodiment of any one of the methods provided, the method further comprises assessing the anaphylactic reaction in the subject prior to and/or after the administration.

In another embodiment of any one of the methods provided, the administering is by intravenous, intraperitoneal or subcutaneous administration.

In another embodiment of any one of the methods provided, the method further comprises recording a reduction or prevention of an anaphylactic reaction to the non-allergenic antigen.

In another embodiment of any one of the methods provided, the immunosuppressant comprises a statin, an mTOR inhibitor, a TGF- $\beta$  signaling agent, a corticosteroid, an inhibitor of mitochondrial function, a P38 inhibitor, an NF- $\kappa$ B inhibitor, an adenosine receptor agonist, a prostaglandin E2 agonist, a phosphodiesterase 4 inhibitor, an HDAC inhibitor or a proteasome inhibitor. In another embodiment of any one of the methods provided, the mTOR inhibitor is rapamycin.

In another embodiment of any one of the methods provided, the non-allergenic comprises a therapeutic macromolecule. In another embodiment of any one of the methods provided, the therapeutic macromolecule is a therapeutic protein or a therapeutic polynucleotide. In another embodiment of any one of the methods provided, the therapeutic protein is for protein replacement of protein supplementation therapy. In another embodiment of any one of the methods provided, the therapeutic protein comprises a/an infusible or injectable therapeutic protein, enzyme, enzyme cofactor, hormone, blood or blood coagulation factor, cytokine, interferon, growth factor, monoclonal antibody, polyclonal antibody, or protein associated with Pompe's disease. In another embodiment of any one of the methods provided, the infusible or injectable therapeutic protein comprises Tocilizumab, alpha-1 antitrypsin, Hematide, albinterferon alfa-2b, Thucin, tesamorelin, ocrelizumab, belimumab, pegloticase, taliglucerase alfa, agalsidase alfa, or velaglucerase alfa. In another embodiment of any one of the methods provided, the enzyme comprises an oxidoreductase, transferase, hydrolase, lyase, isomerase or ligase. In another embodiment of any one of the methods provided, the enzyme comprises an enzyme for enzyme replacement therapy for a lysosomal storage disorder. In another embodiment of any one of the methods provided, the enzyme for replacement therapy for a lysosomal storage disorder comprises imiglucerase,  $\alpha$ -galactosidase

A (a-gal A), agalsidase beta, acid  $\alpha$ -glucosidase (GAA), alglucosidase alfa, LUMIZYME, MYOZYME, arylsulfatase B, laronidase, ALDURAZYME, idursulfase, ELAPRASE, arylsulfatase B, pegloticase, pegsiticase or NAGLAZYME. In another embodiment of any one of the methods provided, the cytokine comprises a lymphokine, interleukin, chemokine, type 1  
5 cytokine or a type 2 cytokine. In another embodiment of any one of the methods provided, the blood or blood coagulation factor comprises Factor I, Factor II, tissue factor, Factor V, Factor VII, Factor VIII, Factor IX, Factor X, Factor Xa, Factor XII, Factor XIII, von Willebrand factor, prekallikrein, high-molecular weight kininogen, fibronectin, antithrombin III, heparin cofactor II, protein C, protein S, protein Z, protein Z-related protease inhibitor (ZPI),  
10 plasminogen, alpha 2-antiplasmin, tissue plasminogen activator (tPA), urokinase, plasminogen activator inhibitor-1 (PAI1), plasminogen activator inhibitor-2 (PAI2), cancer procoagulant or epoetin alfa.

In another embodiment of any one of the methods provided, a load of immunosuppressant attached to the synthetic nanocarriers, on average across the synthetic  
15 nanocarriers, is between 0.1% and 50%. In another embodiment of any one of the methods provided, the load is between 0.1% and 20%.

In another embodiment of any one of the methods provided, the synthetic nanocarriers comprise lipid nanoparticles, polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles or peptide or protein  
20 particles. In another embodiment of any one of the methods provided, the synthetic nanocarriers comprise lipid nanoparticles. In another embodiment of any one of the methods provided, the synthetic nanocarriers comprise liposomes. In another embodiment of any one of the methods provided, the synthetic nanocarriers comprise metallic nanoparticles. In another embodiment of any one of the methods provided, the metallic nanoparticles comprise gold  
25 nanoparticles. In another embodiment of any one of the methods provided, the synthetic nanocarriers comprise polymeric nanoparticles. In another embodiment of any one of the methods provided, the polymeric nanoparticles comprise polymer that is a non-methoxy-terminated, pluronic polymer. In another embodiment of any one of the methods provided, the polymeric nanoparticles comprise a polyester, polyester attached to a polyether, polyamino  
30 acid, polycarbonate, polyacetal, polyketal, polysaccharide, polyethyloxazoline or polyethyleneimine. In another embodiment of any one of the methods provided, the polyester comprises a poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid) or polycaprolactone. In another embodiment of any one of the methods provided, the polymeric

nanoparticles comprise a polyester and a polyester attached to a polyether. In another embodiment of any one of the methods provided, the polyether comprises polyethylene glycol or polypropylene glycol.

5 In another embodiment of any one of the methods provided, the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers is a diameter greater than 100nm. In another embodiment of any one of the methods provided, the diameter is greater than 150nm. In another embodiment of any one of the methods provided, the diameter is greater than 200nm. In another embodiment of any one of the methods provided, the diameter is greater than 250nm. In another embodiment of any one of the methods  
10 provided, the diameter is greater than 300nm.

In another embodiment of any one of the methods provided, an aspect ratio of the synthetic nanocarriers is greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7 or 1:10.

In another aspect, a composition or kit comprising immunosuppressants, in some embodiments attached to synthetic nanocarriers; and a non-allergenic antigen, the non-  
15 allergenic antigen being unattached to synthetic nanocarriers, is provided. In one embodiment, the composition or kit is for use in any one of the methods provided herein. In another embodiment, the composition or kit further comprises a pharmaceutically acceptable carrier. In another embodiment, the immunosuppressants and non-allergenic antigen are contained in the same container. In another embodiment, the immunosuppressants and non-allergenic  
20 antigen are contained in separate containers. In another embodiment, the composition or kit further comprises instructions for use.

In another aspect, a method of manufacturing any one of the compositions or kits provided herein is provided, the method comprising combining the elements of the composition or kit. In one embodiment, one or more of the elements of the composition or kit  
25 are prepared. Accordingly, any one of the methods of manufacturing can comprise a step of preparing a composition of immunosuppressants and preparing a composition of non-allergenic antigen. In one embodiment, the step of preparing a composition of immunosuppressants can comprise attaching the immunosuppressants to synthetic nanocarriers.

In another embodiment, the method of manufacturing comprises producing a dose or  
30 dosage form of a non-allergenic antigen and producing a dose or dosage form of an immunosuppressant. In another embodiment, the step of producing a dose or dosage form of an immunosuppressant comprises attaching the immunosuppressant to synthetic nanocarriers. In another embodiment of any one of the methods of manufacturing provided, the method

further comprises combining the dose or dosage form of the immunosuppressant and dose or dosage form of non-allergenic antigen in a kit.

In another aspect, a use of any of the compositions or kits provided herein for the manufacture of a medicament for reducing or preventing an anaphylactic reaction to a non-allergenic antigen, in a subject is provided. In one embodiment, the composition or kit  
5 comprises an immunosuppressant and a non-allergenic antigen. In another embodiment of any one of the uses provided herein, the immunosuppressant is attached to synthetic nanocarriers.

In another aspect, any one of the compositions or kits provided herein can be for use in any one of the methods provided herein.

10 In another aspect, a method of manufacturing a medicament intended for reducing or preventing an anaphylactic reaction to a non-allergenic antigen is provided. In one embodiment, the medicament comprises an immunosuppressant and a non-allergenic antigen. In another embodiment of any one of the methods of manufacturing provided herein, the immunosuppressant is attached to synthetic nanocarriers.

15 In another aspect, a method of manufacturing a composition or kit comprising preparing or obtaining immunosuppressant, in some embodiments attached to synthetic nanocarriers, and a non-allergenic antigen, the non-allergenic antigen being unattached to synthetic nanocarriers is provided.

## BRIEF DESCRIPTION OF FIGURES

**Fig. 1** shows IgG titers are reduced as a result of an inventive treatment.

**Fig. 2** shows the reduction in IgG titers and anaphylactic score as a result of an inventive treatment.

25 **Fig. 3** shows the reduction in anaphylactic score as a result of an inventive treatment.

**Fig. 4** shows the reduction in IgG titers as a result of an inventive treatment.

## DETAILED DESCRIPTION OF THE INVENTION

Before describing the present invention in detail, it is to be understood that this  
30 invention is not limited to particularly exemplified materials or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting of the use of alternative terminology to describe the present invention.

All publications, patents and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety for all purposes.

As used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the content clearly dictates otherwise. For example, reference to "a polymer" includes a mixture of two or more such molecules or a mixture of differing molecular weights of a single polymer species, reference to "a synthetic nanocarrier" includes a mixture of two or more such synthetic nanocarriers or a plurality of such synthetic nanocarriers, reference to "a RNA molecule" includes a mixture of two or more such RNA molecules or a plurality of such RNA molecules, reference to "an immunosuppressant" includes a mixture of two or more such materials or a plurality of such immunosuppressant molecules, and the like.

As used herein, the term "comprise" or variations thereof such as "comprises" or "comprising" are to be read to indicate the inclusion of any recited integer (e.g. a feature, element, characteristic, property, method/process step or limitation) or group of integers (e.g. features, element, characteristics, properties, method/process steps or limitations) but not the exclusion of any other integer or group of integers. Thus, as used herein, the term "comprising" is inclusive and does not exclude additional, unrecited integers or method/process steps.

In embodiments of any one of the compositions and methods provided herein, "comprising" may be replaced with "consisting essentially of" or "consisting of". The phrase "consisting essentially of" is used herein to require the specified integer(s) or steps as well as those which do not materially affect the character or function of the claimed invention. As used herein, the term "consisting" is used to indicate the presence of the recited integer (e.g. a feature, element, characteristic, property, method/process step or limitation) or group of integers (e.g. features, element, characteristics, properties, method/process steps or limitations) alone.

## A. INTRODUCTION

Anaphylaxis is generally associated with allergic responses to allergens. However, anaphylaxis can also occur in response to administration of a non-allergenic antigen, such as a non-allergenic therapeutic macromolecule. Surprisingly, it has been found that synthetic nanocarriers attached to immunosuppressants administered concomitantly with non-allergenic antigens can be successfully used to block the immune pathways responsible for anaphylaxis to a non-allergenic antigen. Thus, the methods and compositions provided herein are useful for

treating subject in need of therapy with a non-allergenic therapeutic macromolecule, particularly where the subject is at risk of anaphylaxis as a result of such treatment.

It has been found that delivering doses of immunosuppressants concomitantly with non-allergenic antigens can effectively reduce or prevent anaphylaxis related to the non-allergenic antigens. The doses of such immunosuppressants when concomitantly administered with a non-allergenic antigen were surprisingly found to reduce or prevent anaphylaxis even when non-allergenic antigen is subsequently administered without further administration of the immunosuppressants. Thus, the doses of immunosuppressants concomitantly administered with non-allergenic antigen provide long-term protection from anaphylaxis. As such immune responses can be beneficial in countering anaphylaxis during therapeutic treatment with non-allergenic therapeutic macromolecules, the invention can be useful for promoting tolerogenic immune responses in subjects in need of treatment with a non-allergenic therapeutic macromolecule.

The inventors have unexpectedly and surprisingly discovered that the problems and limitations noted above can be overcome by practicing the invention disclosed herein. The present invention is illustrated in the Examples below.

The invention will now be described in more detail below.

## B. DEFINITIONS

"Administering" or "administration" or "administer" means providing a material to a subject in a manner that is pharmacologically useful. The term is intended to include "causing to be administered" in some embodiments. "Causing to be administered" means causing, urging, encouraging, aiding, inducing or directing, directly or indirectly, another party to administer the material.

"Amount effective" in the context of a composition or dosage form for administration to a subject refers to an amount of the composition or dosage form that produces one or more desired responses in the subject, for example, the generation of a tolerogenic immune response (e.g., a reduction in or prevention of anaphylaxis). This amount can refer to a single dose or multiple doses. Therefore, in some embodiments, an amount effective is the amount of any composition provided herein given as multiple doses that produces one or more of these desired immune responses and/or therapeutic effects as provided herein. The amount effective can be for *in vitro* or *in vivo* purposes. For *in vivo* purposes, the amount can be one that a



clinician would believe may have a clinical benefit for a subject in need of reducing or preventing an anaphylactic reaction in a subject as a result of a non-allergenic antigen.

Amounts effective can involve reducing the level of an undesired immune response, although in some embodiments, it involves preventing an undesired immune response altogether. Amounts effective can also involve delaying the occurrence of an undesired immune response. An amount that is effective can also be an amount of a composition provided herein that produces a desired therapeutic endpoint or a desired therapeutic result. Amounts effective, preferably, result in a tolerogenic immune response in a subject to an antigen, such as a non-allergenic antigen. The achievement of any of the foregoing can be monitored by routine methods.

In some embodiments of any one of the compositions and methods provided, the amount effective is one in which the desired immune response persists in the subject for at least 1 week, at least 2 weeks or at least 1 month. In other embodiments of any of the compositions and methods provided, the amount effective is one which produces a measurable desired immune response, for example, a measurable decrease in an immune response (e.g., to a specific non-allergenic antigen), for at least 1 week, at least 2 weeks or at least 1 month.

Amounts effective will depend, of course, on the particular subject being treated; the severity of a condition, disease or disorder; the individual patient parameters including age, physical condition, size and weight; the duration of the treatment; the nature of concurrent therapy (if any); the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reason.

In general, doses of the immunosuppressants and/or non-allergenic antigens in the compositions of the invention can refer to the amount of the immunosuppressants and/or non-allergenic antigens. Alternatively, the dose can be administered based on the number of synthetic nanocarriers, such as synthetic nanocarriers that provide the desired amount of immunosuppressants.

“Antigen-specific” refers to any immune response that results from the presence of the antigen, or portion thereof, or that generates molecules that specifically recognize or bind the

antigen. When the antigen is a non-allergenic antigen, antigen-specific may mean non-allergenic antigen-specific. For example, where the immune response is antigen-specific antibody production, antibodies are produced that specifically bind the antigen.

“Assessing an immune response” refers to any measurement or determination of the level, presence or absence, reduction, increase in, etc. of an immune response in vitro or in vivo. Such measurements or determinations may be performed on one or more samples obtained from a subject. Such assessing can be performed with any of the methods provided herein or otherwise known in the art. The assessing may be assessing the reduction, prevention, presence or absence of an anaphylactic reaction to a non-allergenic antigen.

“Attach” or “Attached” or “Couple” or “Coupled” (and the like) means to chemically associate one entity (for example a moiety) with another. In some embodiments, the attaching is covalent, meaning that the attachment occurs in the context of the presence of a covalent bond between the two entities. In non-covalent embodiments, the non-covalent attaching is mediated by non-covalent interactions including but not limited to charge interactions, affinity interactions, metal coordination, physical adsorption, host-guest interactions, hydrophobic interactions, TT stacking interactions, hydrogen bonding interactions, van der Waals interactions, magnetic interactions, electrostatic interactions, dipole-dipole interactions, and/or combinations thereof. In embodiments, encapsulation is a form of attaching. In embodiments, non-allergenic antigens and immunosuppressants are not attached to one another, meaning that the antigens and immunosuppressants are not subjected to a process specifically intended to chemically associate one with another. In embodiments, non-allergenic antigens are not attached to synthetic nanocarriers, meaning that the antigens and synthetic nanocarriers are not subjected to a process specifically intended to chemically associate one with another.

“Average”, as used herein, refers to the arithmetic mean unless otherwise noted.

“Combination”, as applied to two or more materials and/or agents (also referred to herein as the *components*), is intended to define material in which the two or more materials/agents are associated. Components may be separately identified, e.g. first component, second component, third component, etc. The terms “combined” and “combining” in this context are to be interpreted accordingly.

The association of the two or more materials /agents in a combination may be physical or non-physical. Examples of physically associated combined materials/agents include:

- compositions (e.g. unitary formulations) comprising the two or more materials/agents in admixture (for example within the same unit dose);

- compositions comprising material in which the two or more materials/agents are chemically/physicochemically linked (for example by crosslinking, molecular agglomeration or binding to a common vehicle moiety);
- compositions comprising material in which the two or more materials/agents are chemically/physicochemically co-packaged (for example, disposed on or within lipid vesicles, particles (e.g. micro- or nanoparticles) or emulsion droplets);
- pharmaceutical kits, pharmaceutical packs or patient packs in which the two or more materials/agents are co-packaged or co-presented (e.g. as part of an array of unit doses);

Examples of non-physically associated combined materials/agents include:

- material (e.g. a non-unitary formulation) comprising at least one of the two or more materials/agents together with instructions for the extemporaneous association of the at least one compound/agent to form a physical association of the two or more materials/agents;
- material (e.g. a non-unitary formulation) comprising at least one of the two or more materials/agents together with instructions for combination therapy with the two or more materials/agents;
- material comprising at least one of the two or more materials/agents together with instructions for administration to a patient population in which the other(s) of the two or more materials/agents have been (or are being) administered;
- material comprising at least one of the two or more materials/agents in an amount or in a form which is specifically adapted for use in combination with the other(s) of the two or more materials/agents.

As used herein, the term “combination therapy” is intended to define therapies which comprise the use of a combination of two or more materials/agents (as defined above). Thus, references to “combination therapy”, “combinations” and the use of materials/agents “in combination” in this application may refer to materials/agents that are administered as part of the same overall treatment regimen. As such, the posology of each of the two or more materials/agents may differ: each may be administered at the same time or at different times. It will therefore be appreciated that the materials/agents of the combination may be administered sequentially (e.g. before or after) or simultaneously, either in the same pharmaceutical formulation (i.e. together), or in different pharmaceutical formulations (i.e.

separately). Simultaneously in the same formulation is as a unitary formulation whereas simultaneously in different pharmaceutical formulations is non-unitary. The posologies of each of the two or more materials/agents in a combination therapy may also differ with respect to the route of administration.

5           “Concomitantly” means administering two or more materials/agents to a subject in a manner that is correlated in time, preferably sufficiently correlated in time so as to provide a modulation in a physiologic or immunologic response, and even more preferably the two or more materials/agents are administered in combination. In embodiments, concomitant administration may encompass administration of two or more materials/agents within a  
10 specified period of time, preferably within 1 month, more preferably within 1 week, still more preferably within 1 day, and even more preferably within 1 hour. In embodiments, the materials/agents may be repeatedly administered concomitantly; that is concomitant administration on more than one occasion, as may be provided in the Examples.

          “Determining” or “determine” means to ascertain a factual relationship. Determining  
15 may be accomplished in a number of ways, including but not limited to performing experiments, or making projections. For instance, a dose of an immunosuppressant or antigen may be determined by starting with a test dose and using known scaling techniques (such as allometric or isometric scaling) to determine the dose for administration. Such may also be used to determine a protocol as provided herein. In another embodiment, the dose may be  
20 determined by testing various doses in a subject, i.e. through direct experimentation based on experience and guiding data. In embodiments, “determining” or “determine” comprises “causing to be determined.” “Causing to be determined” means causing, urging, encouraging, aiding, inducing or directing or acting in coordination with an entity for the entity to ascertain a factual relationship; including directly or indirectly, or expressly or impliedly.

25           “Dosage form” means a pharmacologically and/or immunologically active material in a medium, carrier, vehicle, or device suitable for administration to a subject. Any one of the compositions or doses provided herein may be in a dosage form.

          “Dose” refers to a specific quantity of a pharmacologically and/or immunologically active material for administration to a subject for a given time.

30           “Encapsulate” means to enclose at least a portion of a substance within a synthetic nanocarrier. In some embodiments, a substance is enclosed completely within a synthetic nanocarrier. In other embodiments, most or all of a substance that is encapsulated is not exposed to the local environment external to the synthetic nanocarrier. In other embodiments,

no more than 50%, 40%, 30%, 20%, 10% or 5% (weight/weight) is exposed to the local environment. Encapsulation is distinct from absorption, which places most or all of a substance on a surface of a synthetic nanocarrier, and leaves the substance exposed to the local environment external to the synthetic nanocarrier.

5           “Generating” means causing an action, such as a physiologic or immunologic response (e.g., a tolerogenic immune response) to occur, either directly oneself or indirectly.

          “Identifying a subject” is any action or set of actions that allows a clinician to recognize a subject as one who may benefit from the methods, compositions or kits provided herein. Preferably, the identified subject is one who is in need of therapeutic benefit from a non-  
10   allergenic antigen as provided herein. The action or set of actions may be either directly oneself or indirectly. In one embodiment of any one of the methods provided herein, the method further comprises identifying a subject in need of a method, composition or kit as provided herein.

          “Immunosuppressant” means a compound that causes an APC to have an  
15   immunosuppressive effect (e.g., tolerogenic effect) or a T cell or a B cell to be suppressed. An immunosuppressive effect generally refers to the production or expression of cytokines or other factors by the APC that reduces, inhibits or prevents an undesired immune response or that promotes a desired immune response, such as a regulatory immune response. When the APC acquires an immunosuppressive function (under the immunosuppressive effect) on  
20   immune cells that recognize an antigen presented by this APC, the immunosuppressive effect is said to be specific to the presented antigen. Without being bound by any particular theory, it is thought that the immunosuppressive effect is a result of the immunosuppressant being delivered to the APC, preferably in the presence of an antigen. In one embodiment, the immunosuppressant is one that causes an APC to promote a regulatory phenotype in one or  
25   more immune effector cells. For example, the regulatory phenotype may be characterized by the inhibition of the production, induction, stimulation or recruitment of antigen-specific CD4+ T cells or B cells, the inhibition of the production of antigen-specific antibodies, the production, induction, stimulation or recruitment of Treg cells (e.g., CD4+CD25<sup>high</sup>FoxP3+ Treg cells), etc. This may be the result of the conversion of CD4+ T cells or B cells to a  
30   regulatory phenotype. This may also be the result of induction of FoxP3 in other immune cells, such as CD8+ T cells, macrophages and iNKT cells. In one embodiment, the immunosuppressant is one that affects the response of the APC after it processes an antigen. In another embodiment, the immunosuppressant is not one that interferes with the processing of

the antigen. In a further embodiment, the immunosuppressant is not an apoptotic-signaling molecule. In another embodiment, the immunosuppressant is not a phospholipid.

Immunosuppressants include, but are not limited to, statins; mTOR inhibitors, such as rapamycin or a rapamycin analog; TGF- $\beta$  signaling agents; TGF- $\beta$  receptor agonists; histone  
5 deacetylase inhibitors, such as Trichostatin A; corticosteroids; inhibitors of mitochondrial function, such as rotenone; P38 inhibitors; NF- $\kappa$ B inhibitors, such as 6Bio, Dexamethasone, TCPA-1, IKK VII; adenosine receptor agonists; prostaglandin E2 agonists (PGE2), such as Misoprostol; phosphodiesterase inhibitors, such as phosphodiesterase 4 inhibitor (PDE4), such as Rolipram; proteasome inhibitors; kinase inhibitors; G-protein coupled receptor agonists; G-  
10 protein coupled receptor antagonists; glucocorticoids; retinoids; cytokine inhibitors; cytokine receptor inhibitors; cytokine receptor activators; peroxisome proliferator-activated receptor antagonists; peroxisome proliferator-activated receptor agonists; histone deacetylase inhibitors; calcineurin inhibitors; phosphatase inhibitors; PI3KB inhibitors, such as TGX-221; autophagy inhibitors, such as 3-Methyladenine; aryl hydrocarbon receptor inhibitors; proteasome inhibitor  
15 I (PSI); and oxidized ATPs, such as P2X receptor blockers. Immunosuppressants also include IDO, vitamin D3, cyclosporins, such as cyclosporine A, aryl hydrocarbon receptor inhibitors, resveratrol, azathiopurine (Aza), 6-mercaptopurine (6-MP), 6-thioguanine (6-TG), FK506, sangliferin A, salmeterol, mycophenolate mofetil (MMF), aspirin and other COX inhibitors, niflumic acid, estriol, methotrexate and triptolide. In embodiments, the immunosuppressant  
20 may comprise any of the agents provided herein.

The immunosuppressant can be a compound that directly provides the immunosuppressive effect on APCs or it can be a compound that provides the immunosuppressive effect indirectly (i.e., after being processed in some way after  
administration). Immunosuppressants, therefore, include prodrug forms of any of the  
25 compounds provided herein.

In embodiments of any one of the methods, compositions or kits provided herein, the immunosuppressants provided herein are attached to synthetic nanocarriers. In preferable  
embodiments, the immunosuppressant is an element that is in addition to the material that makes up the structure of the synthetic nanocarrier. For example, in one embodiment, where  
30 the synthetic nanocarrier is made up of one or more polymers, the immunosuppressant is a compound that is in addition and attached to the one or more polymers. As another example, in one embodiment, where the synthetic nanocarrier is made up of one or more lipids, the immunosuppressant is again in addition and attached to the one or more lipids. In

embodiments, such as where the material of the synthetic nanocarrier also results in an immunosuppressive effect, the immunosuppressant is an element present in addition to the material of the synthetic nanocarrier that results in an immunosuppressive effect.

Other exemplary immunosuppressants include, but are not limited, small molecule drugs, natural products, antibodies (e.g., antibodies against CD20, CD3, CD4), biologics-based drugs, carbohydrate-based drugs, nanoparticles, liposomes, RNAi, antisense nucleic acids, aptamers, methotrexate, NSAIDs; fingolimod; natalizumab; alemtuzumab; anti-CD3; tacrolimus (FK506); cytokines and growth factors, such as TGF- $\beta$  and IL-10; etc. Further immunosuppressants, are known to those of skill in the art, and the invention is not limited in this respect.

In embodiments of any one of the methods, compositions or kits provided herein, the immunosuppressant is in a form, such as a nanocrystalline form, whereby the form of the immunosuppressant itself is a particle or particle-like. In embodiments, such forms mimic a virus or other foreign pathogen. Many drugs have been nanonized and appropriate methods for producing such drug forms would be known to one of ordinary skill in the art. Drug nanocrystals, such as nanocrystalline rapamycin are known to those of ordinary skill in the art (Katteboinaa, et al. 2009, International Journal of PharmTech Resesarch; Vol. 1, No. 3; pp682-694. As used herein a “drug nanocrystal” refers to a form of a drug (e.g., an immunosuppressant) that does not include a carrier or matrix material. In some embodiments, drug nanocrystals comprise 90%, 95%, 98%, or 99% or more drug. Methods for producing drug nanocrystals include, without limitation, milling, high pressure homogenization, precipitation, spray drying, rapid expansion of supercritical solution (RESS), Nanoedge® technology (Baxter Healthcare), and Nanocrystal Technology™ (Elan Corporation). In some embodiments, a surfactant or a stabilizer may be used for steric or electrostatic stability of the drug nanocrystal. In some embodiments the nanocrystal or nanocrytalline form of an immunosuppressant may be used to increase the solubility, stability, and/or bioavailability of the immunosuppressant, particularly immunosuppressants that are insoluble or labile. In some embodiments, an anaphylactic reaction to a non-allergenic antigen is reduced or prevented to a comparable extent following administering a non-allergic antigen concomitantly with an immunosuppressant in nanocrystalline form as achieved following administration of a non-allergenic antigen concomitantly with a composition comprising synthetic nanocarriers attached to an immunosuppressant.

“Load” of the immunosuppressant, when attached to a synthetic nanocarrier, is the amount of the immunosuppressant attached to a synthetic nanocarrier based on the total dry recipe weight of materials in an entire synthetic nanocarrier (weight/weight). Generally, such a load is calculated as an average across a population of synthetic nanocarriers. In one embodiment, the load of the immunosuppressant on average across a population of synthetic nanocarriers is between 0.0001% and 99%. In another embodiment, the load of the immunosuppressant is between 0.01% and 50%. In another embodiment, the load of the immunosuppressant is between 0.01% and 20%. In a further embodiment, the load of the immunosuppressant is between 0.1% and 10%. In still a further embodiment, the load of the immunosuppressant is between 1% and 10%. In still a further embodiment, the load is between 7% and 20%. In yet another embodiment, the load of the immunosuppressant is at least 0.1%, at least 0.2%, at least 0.3%, at least 0.4%, at least 0.5%, at least 0.6%, at least 0.7%, at least 0.8%, at least 0.9%, at least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least at least 7%, at least 8%, at least 9%, at least 10%, at least 11%, at least 12%, at least 13%, at least 14%, at least 15%, at least 16%, at least 17%, at least 18%, at least 19% at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% on average across the population of synthetic nanocarriers. In yet a further embodiment, the load of the immunosuppressant is 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% on average across the population of synthetic nanocarriers. In some embodiments of the above embodiments, the load of the immunosuppressant is no more than 25% on average across a population of synthetic nanocarriers. In embodiments, the load is calculated as may be described in the Examples or as otherwise known in the art.

In some embodiments, when the form of the immunosuppressant is itself a particle or particle-like, such as a nanocrystalline immunosuppressant, the load of immunosuppressant is the amount of the immunosuppressant in the particles or the like (weight/weight). In such embodiments, the load can approach 97%, 98%, 99% or more..

“Maximum dimension of a synthetic nanocarrier” means the largest dimension of a nanocarrier measured along any axis of the synthetic nanocarrier. “Minimum dimension of a synthetic nanocarrier” means the smallest dimension of a synthetic nanocarrier measured along any axis of the synthetic nanocarrier. For example, for a spheroidal synthetic nanocarrier, the maximum and minimum dimension of a synthetic nanocarrier would be substantially identical,



and would be the size of its diameter. Similarly, for a cuboidal synthetic nanocarrier, the minimum dimension of a synthetic nanocarrier would be the smallest of its height, width or length, while the maximum dimension of a synthetic nanocarrier would be the largest of its height, width or length. In an embodiment, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100 nm. In an embodiment, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or less than 5  $\mu\text{m}$ . Preferably, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is greater than 110 nm, more preferably greater than 120 nm, more preferably greater than 130 nm, and more preferably still greater than 150 nm. Aspects ratios of the maximum and minimum dimensions of synthetic nanocarriers may vary depending on the embodiment. For instance, aspect ratios of the maximum to minimum dimensions of the synthetic nanocarriers may vary from 1:1 to 1,000,000:1, preferably from 1:1 to 100,000:1, more preferably from 1:1 to 10,000:1, more preferably from 1:1 to 1000:1, still more preferably from 1:1 to 100:1, and yet more preferably from 1:1 to 10:1. Preferably, a maximum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample is equal to or less than 3  $\mu\text{m}$ , more preferably equal to or less than 2  $\mu\text{m}$ , more preferably equal to or less than 1  $\mu\text{m}$ , more preferably equal to or less than 800 nm, more preferably equal to or less than 600 nm, and more preferably still equal to or less than 500 nm. In preferred embodiments, a minimum dimension of at least 75%, preferably at least 80%, more preferably at least 90%, of the synthetic nanocarriers in a sample, based on the total number of synthetic nanocarriers in the sample, is equal to or greater than 100 nm, more preferably equal to or greater than 120 nm, more preferably equal to or greater than 130 nm, more preferably equal to or greater than 140 nm, and more preferably still equal to or greater than 150 nm. Measurement of synthetic nanocarrier dimensions (e.g., effective diameter) may be obtained, in some embodiments, by suspending the synthetic nanocarriers in a liquid (usually aqueous) media and using dynamic light scattering (DLS) (e.g. using a Brookhaven ZetaPALS instrument). For example, a suspension of synthetic nanocarriers can be diluted from an aqueous buffer into purified water to achieve a final synthetic nanocarrier suspension

concentration of approximately 0.01 to 0.1 mg/mL. The diluted suspension may be prepared directly inside, or transferred to, a suitable cuvette for DLS analysis. The cuvette may then be placed in the DLS, allowed to equilibrate to the controlled temperature, and then scanned for sufficient time to acquire a stable and reproducible distribution based on appropriate inputs for viscosity of the medium and refractive indices of the sample. The effective diameter, or mean of the distribution, is then reported. Determining the effective sizes of high aspect ratio, or non-spheroidal, synthetic nanocarriers may require augmentative techniques, such as electron microscopy, to obtain more accurate measurements. “Dimension” or “size” or “diameter” of synthetic nanocarriers means the mean of a particle size distribution, for example, obtained using dynamic light scattering.

A “non-allergenic antigen” means an antigen that induces non-allergic immune responses and/or would not be considered likely to result in an allergic response. Non-allergenic antigens can result in an anaphylactic or anaphylactoid response and such a response is characterized, in some embodiments, by production of primarily non-IgE antigen-specific antibodies, such as antigen-specific IgG antibodies. Such anaphylactic or anaphylactoid responses may result in the production of antigen-specific IgE antibodies but the production of such antibodies is not the primary response and/or is less than the production of non-IgE antibodies. Accordingly, in some embodiments, a non-allergenic antigen is one that can cause an anaphylactic response or anaphylactoid response generally characterized by one or more (or all) of the following: the presence of or increased concentration of antigen-specific IgG antibodies (such as IgG1 antibodies); the absence of or reduced concentration of antigen-specific IgE antibodies; presence of IgG immune complexes; decreased expression of FcγRIII, for example on neutrophils and/or basophils; increased concentration of platelet activating factor (PAF), for example, by macrophages and/or basophils, and the like. In some embodiments, such anaphylactic responses can be monitored by assessing the quantity and/or specificity of antibodies; expression of FcγRIII; detection of IgG immune complexes, vasoactive mediators; cutaneous inflammation or irritation (*e.g.*, a rash, hives, etc.), inflammation or swelling of mucosal tissues, respiratory compromise, organ dysfunction, gastrointestinal symptoms, measuring vascular permeability, smooth muscle contraction, cardiac output, or temperature. In some embodiments, a non-allergenic antigen comprises a non-allergenic therapeutic macromolecule or fragment or derivative thereof.

“Non-methoxy-terminated polymer” means a polymer that has at least one terminus that ends with a moiety other than methoxy. In some embodiments, the polymer has at least

two termini that ends with a moiety other than methoxy. In other embodiments, the polymer has no termini that ends with methoxy. “Non-methoxy-terminated, pluronic polymer” means a polymer other than a linear pluronic polymer with methoxy at both termini. Polymeric nanoparticles as provided herein can comprise non-methoxy-terminated polymers or non-methoxy-terminated, pluronic polymers.

“Pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” means a pharmacologically inactive material used together with a pharmacologically active material to formulate the compositions. Pharmaceutically acceptable excipients comprise a variety of materials known in the art, including but not limited to saccharides (such as glucose, lactose, and the like), preservatives such as antimicrobial agents, reconstitution aids, colorants, saline (such as phosphate buffered saline), and buffers.

“Protocol” means a pattern of administering to a subject and includes any dosing regimen of one or more substances to a subject. Protocols are made up of elements (or variables); thus a protocol comprises one or more elements. Such elements of the protocol can comprise dosing amounts, dosing frequency, routes of administration, dosing duration, dosing rates, interval between dosing, combinations of any of the foregoing, and the like. In some embodiments, such a protocol may be used to administer one or more compositions of the invention to one or more test subjects. Immune responses in these test subjects can then be assessed to determine whether or not the protocol was effective in generating a desired or desired level of an immune response or therapeutic effect. Any therapeutic and/or immunologic effect may be assessed. One or more of the elements of a protocol may have been previously demonstrated in test subjects, such as non-human subjects, and then translated into human protocols. For example, dosing amounts demonstrated in non-human subjects can be scaled as an element of a human protocol using established techniques such as allometric scaling or other scaling methods. Whether or not a protocol had a desired effect can be determined using any of the methods provided herein or otherwise known in the art. For example, a sample may be obtained from a subject to which a composition provided herein has been administered according to a specific protocol in order to determine whether or not specific immune cells, cytokines, antibodies, etc. were reduced, generated, activated, etc. Useful methods for detecting the presence and/or number of immune cells include, but are not limited to, flow cytometric methods (e.g., FACS), ELISpot, proliferation responses, cytokine production, and immunohistochemistry methods. Antibodies and other binding agents for specific staining of immune cell markers, are commercially available. Such kits typically

include staining reagents for antigens that allow for FACS-based detection, separation and/or quantitation of a desired cell population from a heterogeneous population of cells. In embodiments, a number of compositions as provided herein are administered to another subject using one or more or all or substantially all of the elements of which the protocol is comprised.

5           “Providing” means an action or set of actions that an individual performs that supply a needed item or set of items or methods for practicing of the present invention. The action or set of actions may be taken either directly oneself or indirectly.

          “Providing a subject” is any action or set of actions that causes a clinician to come in contact with a subject and administer a composition provided herein thereto or to perform a  
10       method provided herein thereupon. Preferably, the subject is one who is in need of antigen-specific tolerance or reduction or prevention of anaphylaxis to a non-allergenic antigen. The action or set of actions may be taken either directly oneself or indirectly. In one embodiment of any one of the methods provided herein, the method further comprises providing a subject.

          “Recording” means noting, or causing directly or indirectly activities in the expectation  
15       that such noting would take place, in any written or electronic form, that a method or composition provided herein achieved a reduction in or prevention of anaphylaxis to a non-allergenic antigen. In some embodiments, the recording occurs when an immunosuppressant in combination with a non-allergenic antigen are administered to a subject according to a method provided herein or at some point thereafter. “Written form”, as used herein, refers to  
20       any recordation on a medium such as paper. “Electronic form”, as used herein, refers to any recordation on electronic media. Any one of the methods provided herein can further comprise a step of recording a reduction on prevention of an anaphylactic or anaphylactoid response to a non-allergenic antigen in a subject receiving a treatment according to a method provided herein.

25       “Subject” means animals, including warm blooded mammals such as humans and primates; avians; domestic household or farm animals such as cats, dogs, sheep, goats, cattle, horses and pigs; laboratory animals such as mice, rats and guinea pigs; fish; reptiles; zoo and wild animals; and the like.

          “Synthetic nanocarrier(s)” means a discrete object that is not found in nature, and that  
30       possesses at least one dimension that is less than or equal to 5 microns in size. Albumin nanoparticles are generally included as synthetic nanocarriers, however in certain embodiments the synthetic nanocarriers do not comprise albumin nanoparticles. In embodiments, synthetic nanocarriers do not comprise chitosan. In other embodiments, synthetic nanocarriers are not

lipid-based nanoparticles. In further embodiments, synthetic nanocarriers do not comprise a phospholipid.

A synthetic nanocarrier can be, but is not limited to, one or a plurality of lipid-based nanoparticles (also referred to herein as lipid nanoparticles, i.e., nanoparticles where the majority of the material that makes up their structure are lipids), polymeric nanoparticles, 5 metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles (i.e., particles that are primarily made up of viral structural proteins but that are not infectious or have low infectivity), peptide or protein-based particles (also referred to herein as protein particles, i.e., particles where the majority of the material that makes up their structure are peptides or proteins) (such as albumin nanoparticles) and/or nanoparticles that are 10 developed using a combination of nanomaterials such as lipid-polymer nanoparticles.

Synthetic nanocarriers may be a variety of different shapes, including but not limited to spheroidal, cuboidal, pyramidal, oblong, cylindrical, toroidal, and the like. Synthetic

nanocarriers according to the invention comprise one or more surfaces. Exemplary synthetic

15 nanocarriers that can be adapted for use in the practice of the present invention comprise: (1)

the biodegradable nanoparticles disclosed in US Patent 5,543,158 to Gref et al., (2) the

polymeric nanoparticles of Published US Patent Application 20060002852 to Saltzman et al.,

(3) the lithographically constructed nanoparticles of Published US Patent Application

20090028910 to DeSimone et al., (4) the disclosure of WO 2009/051837 to von Andrian et al.,

(5) the nanoparticles disclosed in Published US Patent Application 2008/0145441 to Penades et al., (6) the protein nanoparticles disclosed in Published US Patent Application 20090226525 to

de los Rios et al., (7) the virus-like particles disclosed in published US Patent Application

20060222652 to Sebbel et al., (8) the nucleic acid attached virus-like particles disclosed in

published US Patent Application 20060251677 to Bachmann et al., (9) the virus-like particles

25 disclosed in WO2010047839A1 or WO2009106999A2, (10) the nanoprecipitated

nanoparticles disclosed in P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853

(2010), (11) apoptotic cells, apoptotic bodies or the synthetic or semisynthetic mimics

disclosed in U.S. Publication 2002/0086049, or (12) those of Look et al., Nanogel-based

30 delivery of mycophenolic acid ameliorates systemic lupus erythematosus in mice" *J. Clinical Investigation* 123(4):1741-1749(2013). In embodiments, synthetic nanocarriers may possess

an aspect ratio greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7, or greater than 1:10.

Synthetic nanocarriers according to the invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface with hydroxyl groups that activate complement or alternatively comprise a surface that consists essentially of moieties that are not hydroxyl groups that activate complement. In a preferred embodiment, synthetic nanocarriers according to the invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface that substantially activates complement or alternatively comprise a surface that consists essentially of moieties that do not substantially activate complement. In a more preferred embodiment, synthetic nanocarriers according to the invention that have a minimum dimension of equal to or less than about 100 nm, preferably equal to or less than 100 nm, do not comprise a surface that activates complement or alternatively comprise a surface that consists essentially of moieties that do not activate complement. In embodiments, synthetic nanocarriers exclude virus-like particles. In embodiments, synthetic nanocarriers may possess an aspect ratio greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7, or greater than 1:10.

A “therapeutic macromolecule” refers to any protein, carbohydrate, lipid or nucleic acid that may be administered to a subject and have a therapeutic effect. In some embodiments, administration of the therapeutic macromolecule to a subject may result in an undesired immune response. As described herein, administration of a therapeutic macromolecule as provided herein, in some embodiments, can enhance the therapeutic effectiveness of the therapeutic macromolecule, such as by reducing undesired immune responses thereto. In some embodiments, the therapeutic macromolecule may be a therapeutic polynucleotide or therapeutic protein. As provided herein, the therapeutic macromolecule is a non-allergenic therapeutic macromolecule.

“Therapeutic polynucleotide” means any polynucleotide or polynucleotide-based therapy that may be administered to a subject and have a therapeutic effect. Such therapies include gene silencing. Examples of such therapy are known in the art, and include, but are not limited to, naked RNA (including messenger RNA, modified messenger RNA, and forms of RNAi). Examples of other therapeutic polynucleotides are provided elsewhere herein. Therapeutic polynucleotides may be produced in, on or by cells and also may be obtained using cell free or fully synthetic from in vitro methods. Subjects, therefore, include any subject that is in need of treatment with any of the foregoing. Such subject include those that will

receive any of the foregoing. As provided herein, the therapeutic polynucleotide is a non-allergenic therapeutic polynucleotide.

A “therapeutic protein” refers to any protein or protein-based therapy that may be administered to a subject and have a therapeutic effect. Such therapies include protein replacement and protein supplementation therapies. Such therapies also include the administration of exogenous or foreign proteins, antibody therapies, and cell or cell-based therapies. Therapeutic proteins comprise, but are not limited to, enzymes, enzyme cofactors, hormones, blood clotting factors, cytokines, growth factors, monoclonal antibodies, antibody-drug conjugates, and polyclonal antibodies. Examples of other therapeutic proteins are provided elsewhere herein. Therapeutic proteins may be produced in, on or by cells and may be obtained from such cells or administered in the form of such cells. In embodiments, the therapeutic protein is produced in, on or by mammalian cells, insect cells, yeast cells, bacteria cells, plant cells, transgenic animal cells, transgenic plant cells, etc. The therapeutic protein may be recombinantly produced in such cells. The therapeutic protein may be produced in, on or by a virally transformed cell. Subjects, therefore, include any subject that is in need of treatment with any of the foregoing. Such subject include those that will receive any of the foregoing. As provided herein, the therapeutic protein is a non-allergenic therapeutic protein.

“Undesired immune response” refers to any undesired immune response that results from exposure to an antigen, promotes or exacerbates a disease, disorder or condition provided herein (or a symptom thereof), or is symptomatic of a disease, disorder or condition provided herein. Such immune responses generally have a negative impact on a subject’s health or is symptomatic of a negative impact on a subject’s health. Undesired immune responses include anaphylaxis to a non-allergenic antigen.

## C. COMPOSITIONS AND METHODS

Provided herein are compositions comprising immunosuppressants, in some embodiments attached to synthetic nanocarriers, and related methods or kits. Such compositions, methods, or kits are useful for reducing the generation of undesired immune responses and promoting the generation of tolerogenic immune responses that are specific to non-allergenic antigens. The compositions can be administered to subjects in which a tolerogenic immune response to a non-allergenic therapeutic macromolecule is desired. Such subjects include those that are in need of therapy with the therapeutic macromolecule. In

embodiments, multiple doses of the immunosuppressants are administered concomitantly as described herein.

A wide variety of synthetic nanocarriers can be used according to the invention. In some embodiments, synthetic nanocarriers are spheres or spheroids. In some embodiments, synthetic nanocarriers are flat or plate-shaped. In some embodiments, synthetic nanocarriers are cubes or cubic. In some embodiments, synthetic nanocarriers are ovals or ellipses. In some embodiments, synthetic nanocarriers are cylinders, cones, or pyramids.

In some embodiments, it is desirable to use a population of synthetic nanocarriers that is relatively uniform in terms of size or shape so that each synthetic nanocarrier has similar properties. For example, at least 80%, at least 90%, or at least 95% of the synthetic nanocarriers, based on the total number of synthetic nanocarriers, may have a minimum dimension or maximum dimension that falls within 5%, 10%, or 20% of the average diameter or average dimension of the synthetic nanocarriers.

Synthetic nanocarriers can be solid or hollow and can comprise one or more layers. In some embodiments, each layer has a unique composition and unique properties relative to the other layer(s). To give but one example, synthetic nanocarriers may have a core/shell structure, wherein the core is one layer (e.g. a polymeric core) and the shell is a second layer (e.g. a lipid bilayer or monolayer). Synthetic nanocarriers may comprise a plurality of different layers.

In some embodiments, synthetic nanocarriers may optionally comprise one or more lipids. In some embodiments, a synthetic nanocarrier may comprise a liposome. In some embodiments, a synthetic nanocarrier may comprise a lipid bilayer. In some embodiments, a synthetic nanocarrier may comprise a lipid monolayer. In some embodiments, a synthetic nanocarrier may comprise a micelle. In some embodiments, a synthetic nanocarrier may comprise a core comprising a polymeric matrix surrounded by a lipid layer (e.g., lipid bilayer, lipid monolayer, etc.). In some embodiments, a synthetic nanocarrier may comprise a non-polymeric core (e.g., metal particle, quantum dot, ceramic particle, bone particle, viral particle, proteins, nucleic acids, carbohydrates, etc.) surrounded by a lipid layer (e.g., lipid bilayer, lipid monolayer, etc.).

In other embodiments, synthetic nanocarriers may comprise metal particles, quantum dots, ceramic particles, etc. In some embodiments, a non-polymeric synthetic nanocarrier is an aggregate of non-polymeric components, such as an aggregate of metal atoms (e.g., gold atoms).



In some embodiments, synthetic nanocarriers may optionally comprise one or more amphiphilic entities. In some embodiments, an amphiphilic entity can promote the production of synthetic nanocarriers with increased stability, improved uniformity, or increased viscosity. In some embodiments, amphiphilic entities can be associated with the interior surface of a lipid membrane (e.g., lipid bilayer, lipid monolayer, etc.). Many amphiphilic entities known in the art are suitable for use in making synthetic nanocarriers in accordance with the present invention. Such amphiphilic entities include, but are not limited to, phosphoglycerides; phosphatidylcholines; dipalmitoyl phosphatidylcholine (DPPC); dioleoylphosphatidyl ethanolamine (DOPE); dioleoyloxypropyltriethylammonium (DOTMA);

10 dioleoylphosphatidylcholine; cholesterol; cholesterol ester; diacylglycerol; diacylglycerolsuccinate; diphosphatidyl glycerol (DPPG); hexanodecanol; fatty alcohols such as polyethylene glycol (PEG); polyoxyethylene-9-lauryl ether; a surface active fatty acid, such as palmitic acid or oleic acid; fatty acids; fatty acid monoglycerides; fatty acid diglycerides; fatty acid amides; sorbitan trioleate (Span®85) glycocholate; sorbitan monolaurate (Span®20);

15 polysorbate 20 (Tween®20); polysorbate 60 (Tween®60); polysorbate 65 (Tween®65); polysorbate 80 (Tween®80); polysorbate 85 (Tween®85); polyoxyethylene monostearate; surfactin; a poloxomer; a sorbitan fatty acid ester such as sorbitan trioleate; lecithin; lysolecithin; phosphatidylserine; phosphatidylinositol; sphingomyelin; phosphatidylethanolamine (cephalin); cardiolipin; phosphatidic acid; cerebrosides;

20 dicetylphosphate; dipalmitoylphosphatidylglycerol; stearylamine; dodecylamine; hexadecylamine; acetyl palmitate; glycerol ricinoleate; hexadecyl stearate; isopropyl myristate; tyloxapol; poly(ethylene glycol)5000-phosphatidylethanolamine; poly(ethylene glycol)400-monostearate; phospholipids; synthetic and/or natural detergents having high surfactant properties; deoxycholates; cyclodextrins; chaotropic salts; ion pairing agents; and combinations thereof.

25 An amphiphilic entity component may be a mixture of different amphiphilic entities. Those skilled in the art will recognize that this is an exemplary, not comprehensive, list of substances with surfactant activity. Any amphiphilic entity may be used in the production of synthetic nanocarriers to be used in accordance with the present invention.

In some embodiments, synthetic nanocarriers may optionally comprise one or more

30 carbohydrates. Carbohydrates may be natural or synthetic. A carbohydrate may be a derivatized natural carbohydrate. In certain embodiments, a carbohydrate comprises monosaccharide or disaccharide, including but not limited to glucose, fructose, galactose, ribose, lactose, sucrose, maltose, trehalose, cellbiose, mannose, xylose, arabinose, glucuronic

acid, galactonic acid, mannuronic acid, glucosamine, galatosamine, and neuramic acid. In certain embodiments, a carbohydrate is a polysaccharide, including but not limited to pullulan, cellulose, microcrystalline cellulose, hydroxypropyl methylcellulose (HPMC), hydroxycellulose (HC), methylcellulose (MC), dextran, cyclodextran, glycogen, hydroxyethylstarch, carageenan, glycon, amylose, chitosan, N,O-carboxymethylchitosan, algin and alginic acid, starch, chitin, inulin, konjac, glucomannan, pustulan, heparin, hyaluronic acid, curdlan, and xanthan. In embodiments, the synthetic nanocarriers do not comprise (or specifically exclude) carbohydrates, such as a polysaccharide. In certain embodiments, the carbohydrate may comprise a carbohydrate derivative such as a sugar alcohol, including but not limited to mannitol, sorbitol, xylitol, erythritol, maltitol, and lactitol.

In some embodiments, synthetic nanocarriers can comprise one or more polymers. In some embodiments, the synthetic nanocarriers comprise one or more polymers that is a non-methoxy-terminated, pluronic polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated, pluronic polymers. In some embodiments, all of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated, pluronic polymers. In some embodiments, the synthetic nanocarriers comprise one or more polymers that is a non-methoxy-terminated polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated polymers. In some embodiments, all of the polymers that make up the synthetic nanocarriers are non-methoxy-terminated polymers. In some embodiments, the synthetic nanocarriers comprise one or more polymers that do not comprise pluronic polymer. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or 99% (weight/weight) of the polymers that make up the synthetic nanocarriers do not comprise pluronic polymer. In some embodiments, all of the polymers that make up the synthetic nanocarriers do not comprise pluronic polymer. In some embodiments, such a polymer can be surrounded by a coating layer (e.g., liposome, lipid monolayer, micelle, etc.). In some embodiments, various elements of the synthetic nanocarriers can be attached to the polymer.

The immunosuppressants can be attached to the synthetic nanocarriers by any of a number of methods. Generally, the attaching can be a result of bonding between the

immunosuppressants and the synthetic nanocarriers. This bonding can result in the immunosuppressants being attached to the surface of the synthetic nanocarriers and/or contained (encapsulated) within the synthetic nanocarriers. In some embodiments, however, the immunosuppressants are encapsulated by the synthetic nanocarriers as a result of the structure of the synthetic nanocarriers rather than bonding to the synthetic nanocarriers. In preferable embodiments, the synthetic nanocarrier comprises a polymer as provided herein, and the immunosuppressants are attached to the polymer.

When attaching occurs as a result of bonding between the immunosuppressants and synthetic nanocarriers, the attaching may occur via a coupling moiety. A coupling moiety can be any moiety through which an immunosuppressant is bonded to a synthetic nanocarrier. Such moieties include covalent bonds, such as an amide bond or ester bond, as well as separate molecules that bond (covalently or non-covalently) the immunosuppressant to the synthetic nanocarrier. Such molecules include linkers or polymers or a unit thereof. For example, the coupling moiety can comprise a charged polymer to which an immunosuppressant electrostatically binds. As another example, the coupling moiety can comprise a polymer or unit thereof to which it is covalently bonded.

In preferred embodiments, the synthetic nanocarriers comprise a polymer as provided herein. These synthetic nanocarriers can be completely polymeric or they can be a mix of polymers and other materials.

In some embodiments, the polymers of a synthetic nanocarrier associate to form a polymeric matrix. In some of these embodiments, a component, such as an immunosuppressant, can be covalently associated with one or more polymers of the polymeric matrix. In some embodiments, covalent association is mediated by a linker. In some embodiments, a component can be noncovalently associated with one or more polymers of the polymeric matrix. For example, in some embodiments, a component can be encapsulated within, surrounded by, and/or dispersed throughout a polymeric matrix. Alternatively or additionally, a component can be associated with one or more polymers of a polymeric matrix by hydrophobic interactions, charge interactions, van der Waals forces, etc. A wide variety of polymers and methods for forming polymeric matrices therefrom are known conventionally.

Polymers may be natural or unnatural (synthetic) polymers. Polymers may be homopolymers or copolymers comprising two or more monomers. In terms of sequence, copolymers may be random, block, or comprise a combination of random and block sequences. Typically, polymers in accordance with the present invention are organic polymers.

In some embodiments, the polymer comprises a polyester, polycarbonate, polyamide, or polyether, or unit thereof. In other embodiments, the polymer comprises poly(ethylene glycol) (PEG), polypropylene glycol, poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid), or a polycaprolactone, or unit thereof. In some embodiments, it is preferred that the polymer is biodegradable. Therefore, in these embodiments, it is preferred that if the polymer comprises a polyether, such as poly(ethylene glycol) or polypropylene glycol or unit thereof, the polymer comprises a block-co-polymer of a polyether and a biodegradable polymer such that the polymer is biodegradable. In other embodiments, the polymer does not solely comprise a polyether or unit thereof, such as poly(ethylene glycol) or polypropylene glycol or unit thereof.

Other examples of polymers suitable for use in the present invention include, but are not limited to polyethylenes, polycarbonates (e.g. poly(1,3-dioxan-2-one)), polyanhydrides (e.g. poly(sebacic anhydride)), polypropylfumerates, polyamides (e.g. polycaprolactam), polyacetals, polyethers, polyesters (e.g., polylactide, polyglycolide, polylactide-co-glycolide, polycaprolactone, polyhydroxyacid (e.g. poly( $\beta$ -hydroxyalkanoate))), poly(orthoesters), polycyanoacrylates, polyvinyl alcohols, polyurethanes, polyphosphazenes, polyacrylates, polymethacrylates, polyureas, polystyrenes, and polyamines, polylysine, polylysine-PEG copolymers, and poly(ethyleneimine), poly(ethylene imine)-PEG copolymers.

In some embodiments, polymers in accordance with the present invention include polymers which have been approved for use in humans by the U.S. Food and Drug Administration (FDA) under 21 C.F.R. § 177.2600, including but not limited to polyesters (e.g., polylactic acid, poly(lactic-co-glycolic acid), polycaprolactone, polyvalerolactone, poly(1,3-dioxan-2-one)); polyanhydrides (e.g., poly(sebacic anhydride)); polyethers (e.g., polyethylene glycol); polyurethanes; polymethacrylates; polyacrylates; and polycyanoacrylates.

In some embodiments, polymers can be hydrophilic. For example, polymers may comprise anionic groups (e.g., phosphate group, sulphate group, carboxylate group); cationic groups (e.g., quaternary amine group); or polar groups (e.g., hydroxyl group, thiol group, amine group). In some embodiments, a synthetic nanocarrier comprising a hydrophilic polymeric matrix generates a hydrophilic environment within the synthetic nanocarrier. In some embodiments, polymers can be hydrophobic. In some embodiments, a synthetic nanocarrier comprising a hydrophobic polymeric matrix generates a hydrophobic environment within the synthetic nanocarrier. Selection of the hydrophilicity or hydrophobicity of the

polymer may have an impact on the nature of materials that are incorporated (e.g. attached) within the synthetic nanocarrier.

In some embodiments, polymers may be modified with one or more moieties and/or functional groups. A variety of moieties or functional groups can be used in accordance with the present invention. In some embodiments, polymers may be modified with polyethylene glycol (PEG), with a carbohydrate, and/or with acyclic polyacetals derived from polysaccharides (Papisov, 2001, ACS Symposium Series, 786:301). Certain embodiments may be made using the general teachings of US Patent No. 5543158 to Gref et al., or WO publication WO2009/051837 by Von Andrian et al.

In some embodiments, polymers may be modified with a lipid or fatty acid group. In some embodiments, a fatty acid group may be one or more of butyric, caproic, caprylic, capric, lauric, myristic, palmitic, stearic, arachidic, behenic, or lignoceric acid. In some embodiments, a fatty acid group may be one or more of palmitoleic, oleic, vaccenic, linoleic, alpha-linoleic, gamma-linoleic, arachidonic, gadoleic, arachidonic, eicosapentaenoic, docosahexaenoic, or erucic acid.

In some embodiments, polymers may be polyesters, including copolymers comprising lactic acid and glycolic acid units, such as poly(lactic acid-co-glycolic acid) and poly(lactide-co-glycolide), collectively referred to herein as "PLGA"; and homopolymers comprising glycolic acid units, referred to herein as "PGA," and lactic acid units, such as poly-L-lactic acid, poly-D-lactic acid, poly-D,L-lactic acid, poly-L-lactide, poly-D-lactide, and poly-D,L-lactide, collectively referred to herein as "PLA." In some embodiments, exemplary polyesters include, for example, polyhydroxyacids; PEG copolymers and copolymers of lactide and glycolide (e.g., PLA-PEG copolymers, PGA-PEG copolymers, PLGA-PEG copolymers, and derivatives thereof. In some embodiments, polyesters include, for example, poly(caprolactone), poly(caprolactone)-PEG copolymers, poly(L-lactide-co-L-lysine), poly(serine ester), poly(4-hydroxy-L-proline ester), poly[ $\alpha$ -(4-aminobutyl)-L-glycolic acid], and derivatives thereof.

In some embodiments, a polymer may be PLGA. PLGA is a biocompatible and biodegradable co-polymer of lactic acid and glycolic acid, and various forms of PLGA are characterized by the ratio of lactic acid:glycolic acid. Lactic acid can be L-lactic acid, D-lactic acid, or D,L-lactic acid. The degradation rate of PLGA can be adjusted by altering the lactic acid:glycolic acid ratio. In some embodiments, PLGA to be used in accordance with the present invention is characterized by a lactic acid:glycolic acid ratio of approximately 85:15,

approximately 75:25, approximately 60:40, approximately 50:50, approximately 40:60, approximately 25:75, or approximately 15:85.

In some embodiments, polymers may be one or more acrylic polymers. In certain embodiments, acrylic polymers include, for example, acrylic acid and methacrylic acid  
5 copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, aminoalkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamide copolymer, poly(methyl methacrylate), poly(methacrylic acid anhydride), methyl methacrylate, polymethacrylate, poly(methyl methacrylate) copolymer, polyacrylamide, aminoalkyl methacrylate copolymer, glycidyl methacrylate copolymers,  
10 polycyanoacrylates, and combinations comprising one or more of the foregoing polymers. The acrylic polymer may comprise fully-polymerized copolymers of acrylic and methacrylic acid esters with a low content of quaternary ammonium groups.

In some embodiments, polymers can be cationic polymers. In general, cationic polymers are able to condense and/or protect negatively charged strands of nucleic acids  
15 (Zauner et al., 1998, Adv. Drug Del. Rev., 30:97; and Kabanov et al., 1995, Bioconjugate Chem., 6:7), poly(ethylene imine) (PEI; Boussif et al., 1995, Proc. Natl. Acad. Sci., USA, 1995, 92:7297), and poly(amidoamine) dendrimers (Kukowska-Latallo et al., 1996, Proc. Natl. Acad. Sci., USA, 93:4897; Tang et al., 1996, Bioconjugate Chem., 7:703; and Haensler et al., 1993, Bioconjugate Chem., 4:372) are positively-charged at physiological pH, form ion pairs  
20 with nucleic acids. In embodiments, the synthetic nanocarriers may not comprise (or may exclude) cationic polymers.

In some embodiments, polymers can be degradable polyesters bearing cationic side chains (Putnam et al., 1999, Macromolecules, 32:3658; Barrera et al., 1993, J. Am. Chem. Soc., 115:11010; Kwon et al., 1989, Macromolecules, 22:3250; Lim et al., 1999, J. Am. Chem.  
25 Soc., 121:5633; and Zhou et al., 1990, Macromolecules, 23:3399). Examples of these polyesters include poly(L-lactide-co-L-lysine) (Barrera et al., 1993, J. Am. Chem. Soc., 115:11010), poly(serine ester) (Zhou et al., 1990, Macromolecules, 23:3399), poly(4-hydroxy-L-proline ester) (Putnam et al., 1999, Macromolecules, 32:3658; and Lim et al., 1999, J. Am. Chem. Soc., 121:5633), and poly(4-hydroxy-L-proline ester) (Putnam et al., 1999,  
30 Macromolecules, 32:3658; and Lim et al., 1999, J. Am. Chem. Soc., 121:5633).

The properties of these and other polymers and methods for preparing them are well known in the art (see, for example, U.S. Patents 6,123,727; 5,804,178; 5,770,417; 5,736,372; 5,716,404; 6,095,148; 5,837,752; 5,902,599; 5,696,175; 5,514,378; 5,512,600; 5,399,665;

5,019,379; 5,010,167; 4,806,621; 4,638,045; and 4,946,929; Wang et al., 2001, J. Am. Chem. Soc., 123:9480; Lim et al., 2001, J. Am. Chem. Soc., 123:2460; Langer, 2000, Acc. Chem. Res., 33:94; Langer, 1999, J. Control. Release, 62:7; and Uhrich et al., 1999, Chem. Rev., 99:3181). More generally, a variety of methods for synthesizing certain suitable polymers are described in Concise Encyclopedia of Polymer Science and Polymeric Amines and Ammonium Salts, Ed. by Goethals, Pergamon Press, 1980; Principles of Polymerization by Odian, John Wiley & Sons, Fourth Edition, 2004; Contemporary Polymer Chemistry by Allcock et al., Prentice-Hall, 1981; Deming et al., 1997, Nature, 390:386; and in U.S. Patents 6,506,577, 6,632,922, 6,686,446, and 6,818,732.

10 In some embodiments, polymers can be linear or branched polymers. In some embodiments, polymers can be dendrimers. In some embodiments, polymers can be substantially cross-linked to one another. In some embodiments, polymers can be substantially free of cross-links. In some embodiments, polymers can be used in accordance with the present invention without undergoing a cross-linking step. It is further to be understood that the synthetic nanocarriers may comprise block copolymers, graft copolymers, blends, mixtures, and/or adducts of any of the foregoing and other polymers. Those skilled in the art will recognize that the polymers listed herein represent an exemplary, not comprehensive, list of polymers that can be of use in accordance with the present invention.

15 In some embodiments, synthetic nanocarriers do not comprise a polymeric component. In some embodiments, synthetic nanocarriers may comprise metal particles, quantum dots, ceramic particles, etc. In some embodiments, a non-polymeric synthetic nanocarrier is an aggregate of non-polymeric components, such as an aggregate of metal atoms (e.g., gold atoms).

20 Compositions according to the invention can comprise elements, such as immunosuppressants, in combination with pharmaceutically acceptable excipients, such as preservatives, buffers, saline, or phosphate buffered saline. The compositions may be made using conventional pharmaceutical manufacturing and compounding techniques to arrive at useful dosage forms. In an embodiment, compositions, such as those comprising synthetic nanocarriers are suspended in sterile saline solution for injection together with a preservative.

30 In embodiments, when preparing synthetic nanocarriers as carriers, methods for attaching components to the synthetic nanocarriers may be useful. If the component is a small molecule it may be of advantage to attach the component to a polymer prior to the assembly of the synthetic nanocarriers. In embodiments, it may also be an advantage to prepare the

synthetic nanocarriers with surface groups that are used to attach the component to the synthetic nanocarrier through the use of these surface groups rather than attaching the component to a polymer and then using this polymer conjugate in the construction of synthetic nanocarriers.

In certain embodiments, the coupling can be with a covalent linker. In embodiments, immunosuppressants according to the invention can be covalently attached to the external surface via a 1,2,3-triazole linker formed by the 1,3-dipolar cycloaddition reaction of azido groups on the surface of the nanocarrier with immunosuppressant containing an alkyne group or by the 1,3-dipolar cycloaddition reaction of alkynes on the surface of the nanocarrier with immunosuppressants containing an azido group. Such cycloaddition reactions are preferably performed in the presence of a Cu(I) catalyst along with a suitable Cu(I)-ligand and a reducing agent to reduce Cu(II) compound to catalytic active Cu(I) compound. This Cu(I)-catalyzed azide-alkyne cycloaddition (CuAAC) can also be referred as the click reaction.

Additionally, the covalent coupling may comprise a covalent linker that comprises an amide linker, a disulfide linker, a thioether linker, a hydrazone linker, a hydrazide linker, an imine or oxime linker, an urea or thiourea linker, an amidine linker, an amine linker, and a sulfonamide linker.

An amide linker is formed via an amide bond between an amine on one component such as an immunosuppressant with the carboxylic acid group of a second component such as the nanocarrier. The amide bond in the linker can be made using any of the conventional amide bond forming reactions with suitably protected amino acids and activated carboxylic acid such N-hydroxysuccinimide-activated ester.

A disulfide linker is made via the formation of a disulfide (S-S) bond between two sulfur atoms of the form, for instance, of R1-S-S-R2. A disulfide bond can be formed by thiol exchange of a component containing thiol/mercaptan group(-SH) with another activated thiol group on a polymer or nanocarrier or a nanocarrier containing thiol/mercaptan groups with a component containing activated thiol group.



A triazole linker, specifically a 1,2,3-triazole of the form  $R_1$  and  $R_2$ , wherein R1 and R2 may be any chemical entities, is made by the 1,3-dipolar cycloaddition reaction of an azide attached to a first component such as the nanocarrier with a terminal alkyne attached to a second component such as the immunosuppressant. The 1,3-dipolar cycloaddition reaction is



performed with or without a catalyst, preferably with Cu(I)-catalyst, which links the two components through a 1,2,3-triazole function. This chemistry is described in detail by Sharpless et al., *Angew. Chem. Int. Ed.* 41(14), 2596, (2002) and Meldal, et al, *Chem. Rev.*, 2008, 108(8), 2952-3015 and is often referred to as a “click” reaction or CuAAC.

5 In embodiments, a polymer containing an azide or alkyne group, terminal to the polymer chain is prepared. This polymer is then used to prepare a synthetic nanocarrier in such a manner that a plurality of the alkyne or azide groups are positioned on the surface of that nanocarrier. Alternatively, the synthetic nanocarrier can be prepared by another route, and subsequently functionalized with alkyne or azide groups. The component is prepared with the  
10 presence of either an alkyne (if the polymer contains an azide) or an azide (if the polymer contains an alkyne) group. The component is then allowed to react with the nanocarrier via the 1,3-dipolar cycloaddition reaction with or without a catalyst which covalently attaches the component to the particle through the 1,4-disubstituted 1,2,3-triazole linker.

A thioether linker is made by the formation of a sulfur-carbon (thioether) bond in the  
15 form, for instance, of R1-S-R2. Thioether can be made by either alkylation of a thiol/mercaptan (-SH) group on one component with an alkylating group such as halide or epoxide on a second component. Thioether linkers can also be formed by Michael addition of a thiol/mercaptan group on one component to an electron-deficient alkene group on a second component containing a maleimide group or vinyl sulfone group as the Michael acceptor. In  
20 another way, thioether linkers can be prepared by the radical thiol-ene reaction of a thiol/mercaptan group on one component with an alkene group on a second component.

A hydrazone linker is made by the reaction of a hydrazide group on one component with an aldehyde/ketone group on the second component.

A hydrazide linker is formed by the reaction of a hydrazine group on one component  
25 with a carboxylic acid group on the second component. Such reaction is generally performed using chemistry similar to the formation of amide bond where the carboxylic acid is activated with an activating reagent.

An imine or oxime linker is formed by the reaction of an amine or N-alkoxyamine (or aminoxy) group on one component with an aldehyde or ketone group on the second  
30 component.

An urea or thiourea linker is prepared by the reaction of an amine group on one component with an isocyanate or thioisocyanate group on the second component.

An amidine linker is prepared by the reaction of an amine group on one component with an imidoester group on the second component.

An amine linker is made by the alkylation reaction of an amine group on one component with an alkylating group such as halide, epoxide, or sulfonate ester group on the second component. Alternatively, an amine linker can also be made by reductive amination of an amine group on one component with an aldehyde or ketone group on the second component with a suitable reducing reagent such as sodium cyanoborohydride or sodium triacetoxyborohydride.

A sulfonamide linker is made by the reaction of an amine group on one component with a sulfonyl halide (such as sulfonyl chloride) group on the second component.

A sulfone linker is made by Michael addition of a nucleophile to a vinyl sulfone. Either the vinyl sulfone or the nucleophile may be on the surface of the nanocarrier or attached to a component.

The component can also be conjugated to the nanocarrier via non-covalent conjugation methods. For example, a negative charged immunosuppressant can be conjugated to a positive charged nanocarrier through electrostatic adsorption. A component containing a metal ligand can also be conjugated to a nanocarrier containing a metal complex via a metal-ligand complex.

In embodiments, the component can be attached to a polymer, for example polylactic acid-block-polyethylene glycol, prior to the assembly of the synthetic nanocarrier or the synthetic nanocarrier can be formed with reactive or activatable groups on its surface. In the latter case, the component may be prepared with a group which is compatible with the attachment chemistry that is presented by the synthetic nanocarriers' surface. In other embodiments, a peptide component can be attached to VLPs or liposomes using a suitable linker. A linker is a compound or reagent that is capable of coupling two molecules together. In an embodiment, the linker can be a homobifunctional or heterobifunctional reagent as described in Hermanson 2008. For example, an VLP or liposome synthetic nanocarrier containing a carboxylic group on the surface can be treated with a homobifunctional linker, adipic dihydrazide (ADH), in the presence of EDC to form the corresponding synthetic nanocarrier with the ADH linker. The resulting ADH linked synthetic nanocarrier is then conjugated with a peptide component containing an acid group via the other end of the ADH linker on nanocarrier to produce the corresponding VLP or liposome peptide conjugate.

For detailed descriptions of available conjugation methods, see Hermanson G T “Bioconjugate Techniques”, 2nd Edition Published by Academic Press, Inc., 2008. In addition to covalent attachment the component can be attached by adsorption to a pre-formed synthetic nanocarrier or it can be attached by encapsulation during the formation of the synthetic nanocarrier.

Any immunosuppressant as provided herein can be used in the methods or compositions provided and can be, in some embodiments, attached to synthetic nanocarriers. Immunosuppressants include, but are not limited to, statins; mTOR inhibitors, such as rapamycin or a rapamycin analog; TGF- $\beta$  signaling agents; TGF- $\beta$  receptor agonists; histone deacetylase (HDAC) inhibitors; corticosteroids; inhibitors of mitochondrial function, such as rotenone; P38 inhibitors; NF- $\kappa$ B inhibitors; adenosine receptor agonists; prostaglandin E2 agonists; phosphodiesterase inhibitors, such as phosphodiesterase 4 inhibitor; proteasome inhibitors; kinase inhibitors; G-protein coupled receptor agonists; G-protein coupled receptor antagonists; glucocorticoids; retinoids; cytokine inhibitors; cytokine receptor inhibitors; cytokine receptor activators; peroxisome proliferator-activated receptor antagonists; peroxisome proliferator-activated receptor agonists; histone deacetylase inhibitors; calcineurin inhibitors; phosphatase inhibitors and oxidized ATPs. Immunosuppressants also include IDO, vitamin D3, cyclosporine A, aryl hydrocarbon receptor inhibitors, resveratrol, azathiopurine, 6-mercaptopurine, aspirin, niflumic acid, estriol, triplide, interleukins (e.g., IL-1, IL-10), cyclosporine A, siRNAs targeting cytokines or cytokine receptors and the like.

Examples of statins include atorvastatin (LIPITOR<sup>®</sup>, TORVAST<sup>®</sup>), cerivastatin, fluvastatin (LESCOL<sup>®</sup>, LESCOL<sup>®</sup> XL), lovastatin (MEVACOR<sup>®</sup>, ALTOCOR<sup>®</sup>, ALTOPREV<sup>®</sup>), mevastatin (COMPACTIN<sup>®</sup>), pitavastatin (LIVALO<sup>®</sup>, PIAVA<sup>®</sup>), rosuvastatin (PRAVACHOL<sup>®</sup>, SELEKTINE<sup>®</sup>, LIPOSTAT<sup>®</sup>), rosuvastatin (CRESTOR<sup>®</sup>), and simvastatin (ZOCOR<sup>®</sup>, LIPEX<sup>®</sup>).

Examples of mTOR inhibitors include rapamycin and analogs thereof (e.g., CCL-779, RAD001, AP23573, C20-methylrapamycin (C20-Marap), C16-(S)-butylsulfonamidrapamycin (C16-BSrap), C16-(S)-3-methylindolerapamycin (C16-iRap) (Bayle et al. Chemistry & Biology 2006, 13:99-107)), AZD8055, BEZ235 (NVP-BEZ235), chrysophanic acid (chrysophanol), deforolimus (MK-8669), everolimus (RAD0001), KU-0063794, PI-103, PP242, temsirolimus, and WYE-354 (available from Selleck, Houston, TX, USA).

Examples of TGF- $\beta$  signaling agents include TGF- $\beta$  ligands (e.g., activin A, GDF1, GDF11, bone morphogenic proteins, nodal, TGF- $\beta$ s) and their receptors (e.g., ACVR1B, ACVR1C, ACVR2A, ACVR2B, BMPR2, BMPR1A, BMPR1B, TGF $\beta$ RI, TGF $\beta$ RII), R-SMADS/co-SMADS (e.g., SMAD1, SMAD2, SMAD3, SMAD4, SMAD5, SMAD8), and

5 ligand inhibitors (e.g., follistatin, noggin, chordin, DAN, lefty, LTBP1, THBS1, Decorin).

Examples of inhibitors of mitochondrial function include atractyloside (dipotassium salt), bongkreikic acid (triammonium salt), carbonyl cyanide m-chlorophenylhydrazone, carboxyatractyloside (e.g., from *Atractylis gummifera*), CGP-37157, (-)-Deguelin (e.g., from *Mundulea sericea*), F16, hexokinase II VDAC binding domain peptide, oligomycin, rotenone,

10 Ru360, SFK1, and valinomycin (e.g., from *Streptomyces fulvissimus*) (EMD4Biosciences, USA).

Examples of P38 inhibitors include SB-203580 (4-(4-Fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)1H-imidazole), SB-239063 (trans-1-(4hydroxycyclohexyl)-4-(fluorophenyl)-5-(2-methoxy-pyrimidin-4-yl) imidazole), SB-220025

15 (5-(2amino-4-pyrimidinyl)-4-(4-fluorophenyl)-1-(4-piperidinyl)imidazole)), and ARRY-797.

Examples of NF (e.g., NK- $\kappa$ B) inhibitors include IFRD1, 2-(1,8-naphthyridin-2-yl)-Phenol, 5-aminosalicylic acid, BAY 11-7082, BAY 11-7085, CAPE (Caffeic Acid Phenethyl ester), diethylmaleate, IKK-2 Inhibitor IV, IMD 0354, lactacystin, MG-132 [Z-Leu-Leu-Leu-CHO], NF $\kappa$ B Activation Inhibitor III, NF- $\kappa$ B Activation Inhibitor II, JSH-23,

20 parthenolide, Phenylarsine Oxide (PAO), PPM-18, pyrrolidinedithiocarbamic acid ammonium salt, QNZ, RO 106-9920, rocaglamide, rocaglamide AL, rocaglamide C, rocaglamide I, rocaglamide J, rocaglaol, (R)-MG-132, sodium salicylate, triptolide (PG490), and wedelolactone.

Examples of adenosine receptor agonists include CGS-21680 and ATL-146e.

25 Examples of prostaglandin E2 agonists include E-Prostanoid 2 and E-Prostanoid 4.

Examples of phosphodiesterase inhibitors (non-selective and selective inhibitors) include caffeine, aminophylline, IBMX (3-isobutyl-1-methylxanthine), paraxanthine, pentoxifylline, theobromine, theophylline, methylated xanthines, vinpocetine, EHNA (erythro-9-(2-hydroxy-3-nonyl)adenine), anagrelide, enoximone (PERFAN<sup>TM</sup>), milrinone,

30 levosimendon, mesembrine, ibudilast, piclamilast, luteolin, drotaverine, roflumilast (DAXAS<sup>TM</sup>, DALIRESP<sup>TM</sup>), sildenafil (REVATION<sup>®</sup>, VIAGRA<sup>®</sup>), tadalafil (ADCIRCA<sup>®</sup>, CIALIS<sup>®</sup>), vardenafil (LEVITRA<sup>®</sup>, STAXYN<sup>®</sup>), udenafil, avanafil, icariin, 4-methylpiperazine, and pyrazolo pyrimidin-7-1.

Examples of proteasome inhibitors include bortezomib, disulfiram, epigallocatechin-3-gallate, and salinosporamide A.

Examples of kinase inhibitors include bevacizumab, BIBW 2992, cetuximab (ERBITUX<sup>®</sup>), imatinib (GLEEVEC<sup>®</sup>), trastuzumab (HERCEPTIN<sup>®</sup>), gefitinib (IRESSA<sup>®</sup>),  
5 ranibizumab (LUCENTIS<sup>®</sup>), pegaptanib, sorafenib, dasatinib, sunitinib, erlotinib, nilotinib, lapatinib, panitumumab, vandetanib, E7080, pazopanib, and mubritinib.

Examples of glucocorticoids include hydrocortisone (cortisol), cortisone acetate, prednisone, prednisolone, methylprednisolone, dexamethasone, betamethasone, triamcinolone, beclometasone, fludrocortisone acetate, deoxycorticosterone acetate (DOCA), and aldosterone.

10 Examples of retinoids include retinol, retinal, tretinoin (retinoic acid, RETIN-A<sup>®</sup>), isotretinoin (AC CUTANE<sup>®</sup>, AMNESTEEM<sup>®</sup>, CLARAVIS<sup>®</sup>, SOTRET<sup>®</sup>), alitretinoin (PANRETIN<sup>®</sup>), etretinate (TEGISON<sup>™</sup>) and its metabolite acitretin (SORIATANE<sup>®</sup>), tazarotene (TAZORAC<sup>®</sup>, AVAGE<sup>®</sup>, ZORAC<sup>®</sup>), bexarotene (TARGRETIN<sup>®</sup>), and adapalene (DIFFERIN<sup>®</sup>).

15 Examples of cytokine inhibitors include IL1ra, IL1 receptor antagonist, IGFBP, TNF-BF, uromodulin, Alpha-2-Macroglobulin, Cyclosporin A, Pentamidine, and Pentoxifylline (PENTOPAK<sup>®</sup>, PENTOXIL<sup>®</sup>, TRENTAL<sup>®</sup>).

Examples of peroxisome proliferator-activated receptor antagonists include GW9662, PPAR $\gamma$  antagonist III, G335, and T0070907 (EMD4Biosciences, USA).

20 Examples of peroxisome proliferator-activated receptor agonists include pioglitazone, ciglitazone, clofibrate, GW1929, GW7647, L-165,041, LY 171883, PPAR $\gamma$  activator, Fmoc-Leu, troglitazone, and WY-14643 (EMD4Biosciences, USA).

Examples of histone deacetylase inhibitors include hydroxamic acids (or hydroxamates) such as trichostatin A, cyclic tetrapeptides (such as trapoxin B) and  
25 depsipeptides, benzamides, electrophilic ketones, aliphatic acid compounds such as phenylbutyrate and valproic acid, hydroxamic acids such as vorinostat (SAHA), belinostat (PXD101), LAQ824, and panobinostat (LBH589), benzamides such as entinostat (MS-275), CI994, and mocetinostat (MGCD0103), nicotinamide, derivatives of NAD, dihydrocoumarin, naphthopyranone, and 2-hydroxynaphthaldehydes.

30 Examples of calcineurin inhibitors include cyclosporine, pimecrolimus, voclosporin, and tacrolimus.

Examples of phosphatase inhibitors include BN82002 hydrochloride, CP-91149, calyculin A, cantharidic acid, cantharidin, cypermethrin, ethyl-3,4-dephostatin, fostriecin

sodium salt, MAZ51, methyl-3,4-dephostatin, NSC 95397, norcantharidin, okadaic acid ammonium salt from prorocentrum concavum, okadaic acid, okadaic acid potassium salt, okadaic acid sodium salt, phenylarsine oxide, various phosphatase inhibitor cocktails, protein phosphatase 1C, protein phosphatase 2A inhibitor protein, protein phosphatase 2A1, protein phosphatase 2A2, and sodium orthovanadate.

The non-allergenic antigen may be delivered in the form of the antigen itself, or as fragments or derivatives thereof. Non-allergenic antigens include non-allergenic therapeutic macromolecules, such as non-allergenic therapeutic proteins or non-allergenic therapeutic polynucleotides.

Therapeutic proteins include, but are not limited to, infusible therapeutic proteins, enzymes, enzyme cofactors, hormones, blood clotting factors, cytokines and interferons, growth factors, monoclonal antibodies, and polyclonal antibodies (e.g., that are administered to a subject as a replacement therapy), and proteins associated with Pompe's disease (e.g., acid glucosidase alfa, rhGAA (e.g., Myozyme and Lumizyme (Genzyme))). Therapeutic proteins also include proteins involved in the blood coagulation cascade. Therapeutic proteins include, but are not limited to, Factor VIII, Factor VII, Factor IX, Factor V, von Willebrand Factor, von Heldebrant Factor, tissue plasminogen activator, insulin, growth hormone, erythropoietin alfa, VEGF, thrombopoietin, lysozyme, antithrombin and the like. Therapeutic proteins also include adipokines, such as leptin and adiponectin. Other examples of therapeutic proteins are as described below and elsewhere herein.

Examples of therapeutic proteins used in enzyme replacement therapy of subjects having a lysosomal storage disorder include, but are not limited to, imiglucerase for the treatment of Gaucher's disease (e.g., CEREZYME<sup>TM</sup>), a-galactosidase A (a-gal A) for the treatment of Fabry disease (e.g., agalsidase beta, FABRYZYME<sup>TM</sup>), acid  $\alpha$ -glucosidase (GAA) for the treatment of Pompe disease (e.g., acid glucosidase alfa, LUMIZYME<sup>TM</sup>, MYOZYME<sup>TM</sup>), arylsulfatase B for the treatment of Mucopolysaccharidoses (e.g., laronidase, ALDURAZYME<sup>TM</sup>, idursulfase, ELAPRASE<sup>TM</sup>, arylsulfatase B, NAGLAZYME<sup>TM</sup>), pegloticase (KRYSTEXXA) and pegsiticase.

Examples of enzymes include oxidoreductases, transferases, hydrolases, lyases, isomerases, asparaginases, uricases, glycosidases, asparaginases, uricases, proteases, nucleases, collagenases, hyaluronidases, heparinases, heparanases, lysins, and ligases.

Therapeutic proteins may also include any enzyme, toxin, or other protein or peptide isolated or derived from a bacterial, fungal, or viral source.

Examples of hormones include Melatonin (N-acetyl-5-methoxytryptamine), Serotonin, Thyroxine (or tetraiodothyronine) (a thyroid hormone), Triiodothyronine (a thyroid hormone), Epinephrine (or adrenaline), Norepinephrine (or noradrenaline), Dopamine (or prolactin inhibiting hormone), Antimüllerian hormone (or müllerian inhibiting factor or hormone),

5 Adiponectin, Adrenocorticotrophic hormone (or corticotropin), Angiotensinogen and angiotensin, Antidiuretic hormone (or vasopressin, arginine vasopressin), Atrial-natriuretic peptide (or atriopeptin), Calcitonin, Cholecystokinin, Corticotropin-releasing hormone, Erythropoietin, Follicle-stimulating hormone, Gastrin, Ghrelin, Glucagon, Glucagon-like peptide (GLP-1), GIP, Gonadotropin-releasing hormone, Growth hormone-releasing hormone,

10 Human chorionic gonadotropin, Human placental lactogen, Growth hormone, Inhibin, Insulin, Insulin-like growth factor (or somatomedin), Leptin, Luteinizing hormone, Melanocyte stimulating hormone, Orexin, Oxytocin, Parathyroid hormone, Prolactin, Relaxin, Secretin, Somatostatin, Thrombopoietin, Thyroid-stimulating hormone (or thyrotropin), Thyrotropin-releasing hormone, Cortisol, Aldosterone, Testosterone, Dehydroepiandrosterone,

15 Androstenedione, Dihydrotestosterone, Estradiol, Estrone, Estriol, Progesterone, Calcitriol (1,25-dihydroxyvitamin D<sub>3</sub>), Calcidiol (25-hydroxyvitamin D<sub>3</sub>), Prostaglandins, Leukotrienes, Prostacyclin, Thromboxane, Prolactin releasing hormone, Lipotropin, Brain natriuretic peptide, Neuropeptide Y, Histamine, Endothelin, Pancreatic polypeptide, Renin, and Enkephalin.

Examples of blood or blood coagulation factors include Factor I (fibrinogen), Factor II

20 (prothrombin), tissue factor, Factor V (proaccelerin, labile factor), Factor VII (stable factor, proconvertin), Factor VIII (antihemophilic globulin), Factor IX (Christmas factor or plasma thromboplastin component), Factor X (Stuart-Prower factor), Factor Xa, Factor XI, Factor XII (Hageman factor), Factor XIII (fibrin-stabilizing factor), von Willebrand factor, prekallikrein (Fletcher factor), high-molecular weight kininogen (HMWK) (Fitzgerald factor), fibronectin,

25 fibrin, thrombin, antithrombin III, heparin cofactor II, protein C, protein S, protein Z, protein Z-related protease inhibitor (ZPI), plasminogen, alpha 2-antiplasmin, tissue plasminogen activator (tPA), urokinase, plasminogen activator inhibitor-1 (PAI1), plasminogen activator inhibitor-2 (PAI2), cancer procoagulant, and epoetin alfa (Epogen, Procrit).

Examples of cytokines include lymphokines, interleukins, and chemokines, type 1

30 cytokines, such as IFN- $\gamma$ , TGF- $\beta$ , and type 2 cytokines, such as IL-4, IL-10, and IL-13.

Examples of growth factors include Adrenomedullin (AM), Angiopoietin (Ang), Autocrine motility factor, Bone morphogenetic proteins (BMPs), Brain-derived neurotrophic factor (BDNF), Epidermal growth factor (EGF), Erythropoietin (EPO), Fibroblast growth

factor (FGF), Glial cell line-derived neurotrophic factor (GDNF), Granulocyte colony-stimulating factor (G-CSF), Granulocyte macrophage colony-stimulating factor (GM-CSF), Growth differentiation factor-9 (GDF9), Hepatocyte growth factor (HGF), Hepatoma-derived growth factor (HDGF), Insulin-like growth factor (IGF), Migration-stimulating factor, 5 Myostatin (GDF-8), Nerve growth factor (NGF) and other neurotrophins, Platelet-derived growth factor (PDGF), Thrombopoietin (TPO), Transforming growth factor alpha(TGF- $\alpha$ ), Transforming growth factor beta(TGF- $\beta$ ), Tumour\_necrosis\_factor-alpha(TNF- $\alpha$ ), Vascular endothelial growth factor (VEGF), Wnt Signaling Pathway, placental growth factor (PlGF), (Foetal Bovine Somatotrophin) (FBS), IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, and IL-7.

10 Examples of monoclonal antibodies include Abagovomab, Abciximab, Adalimumab, Adecatumumab, Afelimomab, Afutuzumab, Alacizumab pegol, ALD, Alemtuzumab, Altumomab pentetate, Anatumomab mafenatox, Anrukinzumab, Anti-thymocyte globin, Apolizumab, Arcitumomab, Aselizumab, Atlizumab (tocilizumab), Atorolimumab, Bapineuzumab, Basiliximab, Bavituximab, Bectumomab, Belimumab, Benralizumab, 15 Bertilimumab, Besilesomab, Bevacizumab, Biciromab, Bivatuzumab mertansine, Blinatumomab, Brentuximab vedotin, Briakinumab, Canakinumab, Cantuzumab mertansine, Capromab pendetide, Catumaxomab, Cedelizumab, Certolizumab pegol, Cetuximab, Citatuzumab bogatox, Cixutumumab, Clenoliximab, Clivatuzumab tetraxetan, Conatumumab, Dacetuzumab, Daclizumab, Daratumumab, Denosumab, Detumomab, Dorlimomab aritox, 20 Dorlixizumab, Echromeximab, Eculizumab, Edobacomab, Edrecolomab, Efalizumab, Efungumab, Elotuzumab, Elsilimomab, Enlimomab pegol, Epitumomab cituxetan, Epratuzumab, Erlizumab, Ertumaxomab, Etaracizumab, Exbivirumab, Fanolesomab, Faralimomab, Farletuzumab, Felvizumab, Fezakinumab, Figitumumab, Fontolizumab, Foravirumab, Fresolimumab, Galiximab, Gantenerumab, Gavilimumab, Gemtuzumab 25 ozogamicin, GC1008, Girentuximab, Glembatumumab vedotin, Golimumab, Gomiliximab, Ibalizumab, Ibritumomab tiuxetan, Igovomab, Imciromab, Infliximab, Intetumumab, Inolimomab, Inotuzumab ozogamicin, Ipilimumab, Iratumumab, Keliximab, Labetuzumab, Lebrikizumab, Lemalesomab, Lerdelimomab, Lexatumumab, Libivirumab, Lintuzumab, Lorvotuzumab mertansine, Lucatumumab, Lumiliximab, Mapatumumab, Maslimomab, 30 Matuzumab, Mepolizumab, Metelimomab, Milatuzumab, Minretumomab, Mitumomab, Morolimumab, Motavizumab, Muromonab-CD3, Nacolomab tafenatox, Naptumomab estafenatox, Natalizumab, Nebacumab, Necitumumab, Nerelimomab, Nimotuzumab, Nofetumomab merpentan, Ocrelizumab, Odulimumab, Ofatumumab, Olaratumab,



Omalizumab, Oportuzumab monatox, Oregovomab, Otelixizumab, Pagibaximab, Palivizumab, Panitumumab, Panobacumab, Pascolizumab, Pemtumomab, Pertuzumab, Pexelizumab, Pintumomab, Priliximab, Pritumumab, Rafivirumab, Ramucirumab, Ranibizumab, Raxibacumab, Regavirumab Reslizumab, Rilotumumab, Rituximab, Robatumumab, 5 Rontalizumab, Rovelizumab, Ruplizumab, Satumomab pendetide, Sevirumab, Sibrotuzumab, Sifalimumab, Siltuximab, Siplizumab, Solanezumab, Sonepcizumab, Sontuzumab, Stamulumab, Sulesomab, Tacatuzumab tetraxetan, Tadocizumab, Talizumab, Tanezumab, Taplitumomab paptox, Tefibazumab, Telimomab aritox, Tenatumomab, Teneliximab, Teplizumab, Ticilimumab (tremelimumab), Tigatuzumab, Tocilizumab (atlizumab), 10 Toralizumab, Tositumomab, Trastuzumab, Tremelimumab, Tucotuzumab celmoleukin, Tuvirumab, Urtioxazumab, Ustekinumab, Vapaliximab, Vedolizumab, Veltuzumab, Vepalimomab, Visilizumab, Volociximab, Votumumab, Zalutumumab, Zanolimumab, Ziralimumab, and Zolimomab aritox. Monoclonal antibodies further include anti-TNF-antibodies.

15 Examples of infusion therapy or injectable therapeutic proteins include, for example, Tocilizumab (Roche/Actemra®), alpha-1 antitrypsin (Kamada/AAT), Hematide® (Affymax and Takeda, synthetic peptide), albinterferon alfa-2b (Novartis/Zalbin™), Rhucin® (Pharming Group, C1 inhibitor replacement therapy), tesamorelin (Theratechnologies/Egrifta, synthetic growth hormone-releasing factor), ocrelizumab (Genentech, Roche and Biogen), belimumab 20 (GlaxoSmithKline/Benlysta®), pegloticase (Savient Pharmaceuticals/Krystexxa™), pegsiticase, taliglucerase alfa (Protalix/Uplyso), agalsidase alfa (Shire/Replagal®), velaglucerase alfa (Shire), and Keyhole Limpet Hemocyanin (KLH).

Additional therapeutic proteins include, for example, engineered proteins, such as Fc fusion proteins, bispecific antibodies, multi-specific antibodies, nanobodies, antigen-binding 25 proteins, antibody fragments, and protein conjugates, such as antibody drug conjugates.

Therapeutic polynucleotides include, but are not limited to nucleic acid aptamers such as Pegaptanib (Macugen, a pegylated anti-VEGF aptamer), antisense therapeutics such as antisense poly- or oligonucleotides (e.g., antiviral drug Fomivirsen, or Mipomersen, an antisense therapeutic that targets the messenger RNA for apolipoprotein B for reduction of 30 cholesterol level); small interfering RNAs (siRNAs) (e.g., dicer substrate siRNA molecules (DsiRNAs) which are 25-30 base pair asymmetric double-stranded RNAs that mediate RNAi with extremely high potency); or modified messenger RNAs (mmRNAs) such as those

disclosed in US Patent application 2013/0115272 to de Fougerolles et al. and in Published US Patent application 2012/0251618 to Schrum et al.

Additional therapeutic macromolecules useful in accordance with aspects of this invention will be apparent to those of skill in the art, and the invention is not limited in this respect.

In some embodiments, a substance as provided herein, such as a non-allergenic antigen or immunosuppressant, may be isolated. Isolated refers to the element being separated from its native environment and present in sufficient quantities to permit its identification or use. This means, for example, the element may be (i) selectively produced by expression cloning or (ii) purified as by chromatography or electrophoresis. Isolated elements may be, but need not be, substantially pure. Because an isolated element may be admixed with a pharmaceutically acceptable excipient in a pharmaceutical preparation, the element may comprise only a small percentage by weight of the preparation. The element is nonetheless isolated in that it has been separated from the substances with which it may be associated in living systems, i.e., isolated from other lipids or proteins. Any of the elements provided herein may be isolated and included in the compositions or used in the methods in isolated form.

#### D. METHODS OF MAKING AND USING THE METHODS AND RELATED COMPOSITIONS

Aspects of the invention relate to determining a protocol for the methods of concomitant administration as provided herein. A protocol can be determined by varying the frequency, dosage amount and other aspects of administration of the non-allergenic antigen and the composition of immunosuppressant and subsequently assessing an anaphylactic reaction based on such variation. A preferred protocol for practice of the invention reduces or prevents anaphylaxis to a non-allergenic antigen.

Synthetic nanocarriers may be prepared using a wide variety of methods known in the art. For example, synthetic nanocarriers can be formed by methods such as nanoprecipitation, flow focusing using fluidic channels, spray drying, single and double emulsion solvent evaporation, solvent extraction, phase separation, milling, microemulsion procedures, microfabrication, nanofabrication, sacrificial layers, simple and complex coacervation, and other methods well known to those of ordinary skill in the art. Alternatively or additionally, aqueous and organic solvent syntheses for monodisperse semiconductor, conductive, magnetic, organic, and other nanomaterials have been described (Pellegrino et al., 2005, Small, 1:48;

Murray et al., 2000, *Ann. Rev. Mat. Sci.*, 30:545; and Trindade et al., 2001, *Chem. Mat.*, 13:3843). Additional methods have been described in the literature (see, e.g., Doubrow, Ed., "Microcapsules and Nanoparticles in Medicine and Pharmacy," CRC Press, Boca Raton, 1992; Mathiowitz et al., 1987, *J. Control. Release*, 5:13; Mathiowitz et al., 1987, *Reactive Polymers*, 6:275; and Mathiowitz et al., 1988, *J. Appl. Polymer Sci.*, 35:755; US Patents 5578325 and 6007845; P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853 (2010)).

Various materials may be encapsulated into synthetic nanocarriers as desirable using a variety of methods including but not limited to C. Astete et al., "Synthesis and characterization of PLGA nanoparticles" *J. Biomater. Sci. Polymer Edn*, Vol. 17, No. 3, pp. 247–289 (2006); K. Avgoustakis "Pegylated Poly(Lactide) and Poly(Lactide-Co-Glycolide) Nanoparticles: Preparation, Properties and Possible Applications in Drug Delivery" *Current Drug Delivery* 1:321-333 (2004); C. Reis et al., "Nanoencapsulation I. Methods for preparation of drug-loaded polymeric nanoparticles" *Nanomedicine* 2:8– 21 (2006); P. Paolicelli et al., "Surface-modified PLGA-based Nanoparticles that can Efficiently Associate and Deliver Virus-like Particles" *Nanomedicine*. 5(6):843-853 (2010). Other methods suitable for encapsulating materials into synthetic nanocarriers may be used, including without limitation methods disclosed in United States Patent 6,632,671 to Unger issued October 14, 2003.

In certain embodiments, synthetic nanocarriers are prepared by a nanoprecipitation process or spray drying. Conditions used in preparing synthetic nanocarriers may be altered to yield particles of a desired size or property (e.g., hydrophobicity, hydrophilicity, external morphology, "stickiness," shape, etc.). The method of preparing the synthetic nanocarriers and the conditions (e.g., solvent, temperature, concentration, air flow rate, etc.) used may depend on the materials to be attached to the synthetic nanocarriers and/or the composition of the polymer matrix.

If synthetic nanocarriers prepared by any of the above methods have a size range outside of the desired range, such synthetic nanocarriers can be sized, for example, using a sieve.

Elements (i.e., components) of the synthetic nanocarriers may be attached to the overall synthetic nanocarrier, e.g., by one or more covalent bonds, or may be attached by means of one or more linkers. Additional methods of functionalizing synthetic nanocarriers may be adapted from Published US Patent Application 2006/0002852 to Saltzman et al., Published US Patent

Application 2009/0028910 to DeSimone et al., or Published International Patent Application WO/2008/127532 A1 to Murthy et al.

Alternatively or additionally, synthetic nanocarriers can be attached to components directly or indirectly via non-covalent interactions. In non-covalent embodiments, the non-covalent attaching is mediated by non-covalent interactions including but not limited to charge interactions, affinity interactions, metal coordination, physical adsorption, host-guest interactions, hydrophobic interactions, TT stacking interactions, hydrogen bonding interactions, van der Waals interactions, magnetic interactions, electrostatic interactions, dipole-dipole interactions, and/or combinations thereof. Such attachments may be arranged to be on an external surface or an internal surface of a synthetic nanocarrier. In embodiments, encapsulation and/or absorption is a form of attaching. In embodiments, the synthetic nanocarriers can be combined with an antigen by admixing in the same vehicle or delivery system.

Compositions provided herein may comprise inorganic or organic buffers (e.g., sodium or potassium salts of phosphate, carbonate, acetate, or citrate) and pH adjustment agents (e.g., hydrochloric acid, sodium or potassium hydroxide, salts of citrate or acetate, amino acids and their salts) antioxidants (e.g., ascorbic acid, alpha-tocopherol), surfactants (e.g., polysorbate 20, polysorbate 80, polyoxyethylene9-10 nonyl phenol, sodium desoxycholate), solution and/or cryo/lyo stabilizers (e.g., sucrose, lactose, mannitol, trehalose), osmotic adjustment agents (e.g., salts or sugars), antibacterial agents (e.g., benzoic acid, phenol, gentamicin), antifoaming agents (e.g., polydimethylsiloxane), preservatives (e.g., thimerosal, 2-phenoxyethanol, EDTA), polymeric stabilizers and viscosity-adjustment agents (e.g., polyvinylpyrrolidone, poloxamer 488, carboxymethylcellulose) and co-solvents (e.g., glycerol, polyethylene glycol, ethanol).

Compositions according to the invention may comprise pharmaceutically acceptable excipients. The compositions may be made using conventional pharmaceutical manufacturing and compounding techniques to arrive at useful dosage forms. Techniques suitable for use in practicing the present invention may be found in Handbook of Industrial Mixing: Science and Practice, Edited by Edward L. Paul, Victor A. Atiemo-Obeng, and Suzanne M. Kresta, 2004 John Wiley & Sons, Inc.; and Pharmaceuticals: The Science of Dosage Form Design, 2nd Ed. Edited by M. E. Auten, 2001, Churchill Livingstone. In an embodiment, compositions are in a sterile saline solution for injection together with a preservative.

It is to be understood that the compositions of the invention can be made in any suitable manner, and the invention is in no way limited to compositions that can be produced using the

methods described herein. Selection of an appropriate method of manufacture may require attention to the properties of the particular moieties being associated.

In some embodiments, compositions are manufactured under sterile conditions or are terminally sterilized. This can ensure that resulting compositions are sterile and non-infectious, thus improving safety when compared to non-sterile compositions. This provides a valuable safety measure, especially when subjects receiving the compositions have immune defects, are suffering from infection, and/or are susceptible to infection. In some embodiments, the compositions may be lyophilized and stored in suspension or as lyophilized powder depending on the formulation strategy for extended periods without losing activity.

Administration according to the present invention may be by a variety of routes, including but not limited to subcutaneous, intravenous, intraperitoneal, intramuscular, transmucosal, transdermal, transcutaneous or intradermal routes. In a preferred embodiment, administration is via a subcutaneous route of administration. The compositions referred to herein may be manufactured and prepared for administration, preferably in some embodiments concomitant administration, using conventional methods.

The compositions of the invention can be administered in effective amounts, such as the effective amounts described elsewhere herein. Doses of dosage forms may contain varying amounts of populations of varying amounts of immunosuppressants and/or non-allergenic antigens, according to the invention. The amount of immunosuppressants and/or non-allergenic antigens present in the inventive dosage forms can be varied according to the nature of the non-allergenic antigen, and/or immunosuppressants, the therapeutic benefit to be accomplished, whether or not the immunosuppressants are attached to synthetic nanocarriers and other such parameters. In embodiments, dose ranging studies can be conducted to establish optimal therapeutic amount of the amount of immunosuppressants and/or non-allergenic antigens to be present in dosage forms. In embodiments, the immunosuppressants and/or non-allergenic antigens (when administered concomitantly) are present in dosage forms in an amount effective to generate a tolerogenic immune response to the non-allergenic antigens upon administration to a subject. Preferably the immunosuppressive immune response is a reduction or prevention of anaphylaxis to the non-allergenic antigen. It may be possible to determine amounts of the immunosuppressants and/or non-allergenic antigens effective to generate a tolerogenic immune response using conventional dose ranging studies and techniques in subjects. Inventive dosage forms may be administered at a variety of

frequencies. In a preferred embodiment, a number of administrations are utilized to ensure a pharmacologically relevant response.

In some embodiments, administration of immunosuppressants with non-allergenic antigens is undertaken e.g., prior to subsequent further administration of the non-allergenic antigens. In exemplary embodiments, immunosuppressants are administered multiple times with concomitant administration of non-allergenic antigen prior to subsequent further administration of the non-allergenic antigen. In some embodiments, the immunosuppressants are administered multiple times with concomitant administration of non-allergenic antigen over a period of at least 7, 14 or 21 days prior to subsequent further administration of the non-allergenic antigen alone.

Another aspect of the disclosure relates to kits. In some embodiments, the kit comprises an immunosuppressant, in some embodiments attached to synthetic nanocarriers, and a non-allergenic antigen. The immunosuppressant and non-allergenic antigen can be contained within separate containers or within the same container in the kit. In some embodiments, the container is a vial or an ampoule. In some embodiments, the non-allergenic antigen and/or immunosuppressant are contained within a solution separate from the container, such that the non-allergenic antigen and/or immunosuppressant may be added to the container at a subsequent time. In some embodiments, the non-allergenic antigen and/or immunosuppressant are in lyophilized form each in a separate container or in the same container, such that they may be reconstituted at a subsequent time. In some embodiments, the kit further comprises instructions for reconstitution, mixing, administration, etc. In some embodiments, the instructions include a description of the methods described herein. Instructions can be in any suitable form, e.g., as a printed insert or a label. In some embodiments, the kit further comprises one or more syringes.

## EXAMPLES

### **Example 1: Evaluating Tolerogenic Immune Responses with Synthetic Nanocarriers Comprising Immunosuppressant *In Vivo* (Prophetic)**

#### **Method for Synthetic Nanocarrier Containing Rapamycin**

A primary water-in-oil emulsion is prepared first. W1/O1 is prepared by combining 0.13 M hydrochloric acid solution (0.2 mL), solution 2 (0.75 mL), solution 3 (0.25 mL), and

solution 4 (0.2 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) is then prepared by combining solution 5 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250.

5           The W1/O1/W2 emulsion is added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the synthetic nanocarriers to form. A portion of the synthetic nanocarriers are washed by transferring the synthetic nanocarrier suspension to a centrifuge tube and centrifuging at 21,000×g and 4 °C for one hour, removing the supernatant, and re-  
10       suspending the pellet in phosphate buffered saline. The washing procedure is repeated, and the pellet is re-suspended in phosphate buffered saline for a final synthetic nanocarrier dispersion of about 10 mg/mL.

          The amount of rapamycin in the synthetic nanocarrier is determined by HPLC analysis. The total dry-synthetic nanocarrier mass per mL of suspension is determined by a gravimetric  
15       method.

### **Method for Measuring Rapamycin Load**

          Approximately 3 mg of synthetic nanocarriers are collected and centrifuged to separate supernatant from synthetic nanocarrier pellet. Acetonitrile is added to the pellet, and the  
20       sample is sonicated and centrifuged to remove any insoluble material. The supernatant and pellet are injected on RP-HPLC and absorbance is read at 278nm. The µg found in the pellet are used to calculate % entrapped (load), µg in supernatant and pellet are used to calculate total µg recovered.

### **25       Measurement of IgG**

          The level of IgG antibodies are measured. Blocker Casein in PBS (Thermo Fisher, Catalog #37528) is used as diluent. 0.05% Tween-20 in PBS is used as wash buffer, prepared by adding 10 ml of Tween-20 ((Sigma, Catalog #P9416-100mL) to 2 liters of a 10x PBS stock (PBS: OmniPur® 10X PBS Liquid Concentrate, 4L, EMD Chemicals, Catalog #6505) and 18  
30       Liters of deionized water.

          Keyhole Limpet Hemocyanin (KLH) at a stock concentration of 5 mg/ml is used as a coating material. A 1:1000 dilution to 5 µg/ml is used as a working concentration. Each well of the assay plates is coated with 100 µl diluted OVA per well, plates are sealed with sealing

film (VWR catalog #60941-120), and incubated overnight at 4°C. Costar 9017 96-well Flat bottom plates are used as assay plates (Costar 9017).

Low-binding polypropylene 96-well plate or tubes are used as set-up plates, in which samples are prepared before being transferred to the assay plate. The setup plates did not contain any antigen and, therefore, serum antibodies did not bind to the plate during the setup of the samples. Setup plates are used for sample preparation to minimize binding that might occur during preparation or pipetting of samples if an antigen-coated plate is used to prepare the samples. Before preparing samples in the setup plate, wells are covered with diluent to block any non-specific binding and the plate is sealed and incubated at 4°C overnight.

Assay plates are washed three times with wash buffer, and wash buffer is completely aspirated out of the wells after the last wash. After washing, 300 µl diluent are added to each well of assay plate(s) to block non-specific binding and plates are incubated at least 2 hours at room temperature. Serum samples are prepared in the setup plate at appropriate starting dilutions. Starting dilutions are sometimes also prepared in 1.5 ml tubes using diluent and then transferred to the set-up plate. Appropriate starting dilutions are determined based on previous data, where available. Where no previous data is available, the lowest starting dilution is 1:40. Once diluted, 200 µl of the starting dilution of the serum sample is transferred from the tube to the appropriate well of the setup plate.

Once all samples were prepared in the setup plate, the plate is sealed and stored at 4°C until blocking of the assay plates is complete. Assay plates are washed three times with wash buffer, and wash buffer is completely aspirated after the last wash. After washing, 100 µL of diluent is added to wells in of the assay plates. A pipet is used to transfer samples from the setup plate to the assay plate. Samples are mixed prior to transfer by pipetting 150 µl of diluted serum up and down 3 times. After mixing, 150µl of each sample is transferred from the setup plate and added to the respective assay plate.

Once the starting dilutions of each sample are transferred from the setup plate to the assay plate, serial dilutions are pipetted on the assay plate as follows: 50 µl of each serum sample is removed using a pipet and mixed with the 100 µl of diluent previously added. This step is repeated down the entire plate. After pipetting the dilution of the final row, 50 µl of fluid is removed from the wells in the final row and discarded, resulting in a final volume of 100 µl in every well of the assay plate. Once sample dilutions are prepared in the assay plates, the plates are incubated at room temperature for at least 2 hours.



After the incubation, plates are washed three times with wash buffer. Detection antibody (Goat anti-mouse anti-IgG, HRP conjugated) is diluted 1:1500 (0.33 µg/mL) in diluent and 100 µl of the diluted antibody is added to each well. Plates are incubated for 1 hour at room temperature and then washed three times with wash buffer, with each washing step including a soak time of at least 30 seconds.

After washing, detection substrate is added to the wells. Equal parts of substrate A and substrate B (BD Biosciences TMB Substrate Reagent Set, catalog #555214) are combined immediately before addition to the assay plates, and 100 µl of the mixed substrate solution are added to each well and incubated for 10 minutes in the dark. The reaction is stopped by adding 50 µl of stop solution (2N H<sub>2</sub>SO<sub>4</sub>) to each well after the 10 minute period. The optical density (OD) of the wells is assessed immediately after adding the stop solution on a plate reader at 450 nm with subtraction at 570 nm. Data analysis is performed using Molecular Device's software SoftMax Pro v5.4. A four-parameter logistic curve-fit graph is prepared with the dilution on the x-axis (log scale) and the OD value on the y-axis (linear scale), and the half maximum value (EC<sub>50</sub>) for each sample is determined. The plate template at the top of the layout is adjusted to reflect the dilution of each sample (1 per column).

### **Example 2: Polymeric Nanocarrier Containing Polymer-Rapamycin Conjugate (Prophetic)**

Preparation of PLGA-rapamycin conjugate:

PLGA polymer with acid end group (7525 DLG1A, acid number 0.46 mmol/g, Lakeshore Biomaterials; 5 g, 2.3 mmol, 1.0 eq) is dissolved in 30 mL of dichloromethane (DCM). N,N-Dicyclohexylcarbodiimide (1.2 eq, 2.8 mmol, 0.57 g) is added followed by rapamycin (1.0 eq, 2.3 mmol, 2.1 g) and 4-dimethylaminopyridine (DMAP) (2.0 eq, 4.6 mmol, 0.56 g). The mixture is stirred at rt for 2 days. The mixture is then filtered to remove insoluble dicyclohexylurea. The filtrate is concentrated to ca. 10 mL in volume and added to 100 mL of isopropyl alcohol (IPA) to precipitate out the PLGA-rapamycin conjugate. The IPA layer is removed and the polymer is then washed with 50 mL of IPA and 50 mL of methyl t-butyl ether (MTBE). The polymer is then dried under vacuum at 35 C for 2 days to give PLGA-rapamycin as a white solid (ca. 6.5 g).

Nanocarrier containing PLGA-rapamycin is prepared according to the procedure described in Example 1 as follows:

Solutions for nanocarrier formation are prepared as follows:

Solution 1: PLGA-rapamycin @ 100 mg/mL in methylene chloride. The solution is prepared by dissolving PLGA-rapamycin in pure methylene chloride. Solution 2: PLA-PEG @ 100 mg/mL in methylene chloride. The solution is prepared by dissolving PLA-PEG in pure methylene chloride. Solution 3: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer.

A primary water-in-oil emulsion is prepared first. W1/O1 is prepared by combining solution 1 (0.75 mL), and solution 2 (0.25 mL) in a small pressure tube and sonicating at 50% amplitude for 40 seconds using a Branson Digital Sonifier 250. A secondary emulsion (W1/O1/W2) is then prepared by combining solution 3 (3.0 mL) with the primary W1/O1 emulsion, vortexing for 10 s, and sonicating at 30% amplitude for 60 seconds using the Branson Digital Sonifier 250. The W1/O1/W2 emulsion is added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers is washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g and 4 °C for 35 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure is repeated, and the pellet is re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

### **Example 3: Preparation of Gold Nanocarriers (AuNCs) Containing Rapamycin**

#### **(Prophetic)**

Preparation of HS-PEG-rapamycin:

A solution of PEG acid disulfide (1.0 eq), rapamycin (2.0-2.5 eq), DCC (2.5 eq) and DMAP (3.0 eq) in dry DMF is stirred at rt overnight. The insoluble dicyclohexylurea is removed by filtration and the filtrate is added to isopropyl alcohol (IPA) to precipitate out the PEG-disulfide-di-rapamycin ester and washed with IPA and dried. The polymer is then treated with tris(2-carboxyethyl)phosphine hydrochloride in DMF to reduce the PEG disulfide to thiol PEG rapamycin ester (HS-PEG-rapamycin). The resulting polymer is recovered by precipitation from IPA and dried as previously described and analyzed by <sup>1</sup>H NMR and GPC.

Formation of Gold NCs (AuNCs):

An aq. solution of 500 mL of 1 mM HAuCl<sub>4</sub> is heated to reflux for 10 min with vigorous stirring in a 1 L round-bottom flask equipped with a condenser. A solution of 50 mL of 40 mM of trisodium citrate is then rapidly added to the stirring solution. The resulting deep wine red solution is kept at reflux for 25-30 min and the heat is withdrawn and the solution is

cooled to room temperature. The solution is then filtered through a 0.8  $\mu$ m membrane filter to give the AuNCs solution. The AuNCs are characterized using visible spectroscopy and transmission electron microscopy. The AuNCs are ca. 20 nm diameter capped by citrate with peak absorption at 520 nm.

5        AuNCs conjugate with HS-PEG-rapamycin:

          A solution of 150  $\mu$ l of HS-PEG-rapamycin (10  $\mu$ M in 10 mM pH 9.0 carbonate buffer) is added to 1 mL of 20 nm diameter citrate-capped gold nanocarriers (1.16 nM) to produce a molar ratio of thiol to gold of 2500:1. The mixture is stirred at room temperature under argon for 1 hour to allow complete exchange of thiol with citrate on the gold nanocarriers. The  
10    AuNCs with PEG-rapamycin on the surface is then purified by centrifuge at 12,000g for 30 minutes. The supernatant is decanted and the pellet containing AuNC-S-PEG-rapamycin is then pellet washed with 1x PBS buffer. The purified Gold-PEG-rapamycin nanocarriers are then resuspend in suitable buffer for further analysis and bioassays.

15    **Example 4: Synthetic Nanocarriers for use in the Inventive Methods and Related Compositions**

**Materials**

          Rapamycin was purchased from TSZ CHEM (185 Wilson Street, Framingham, MA  
20    01702; Product Catalogue # R1017). PLGA with 76% lactide and 24% glycolide content and an inherent viscosity of 0.69 dL/g was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL 35211. Product Code 7525 DLG 7A.) PLA-PEG block co-polymer with a PEG block of approximately 5,000 Da and PLA block of approximately 40,000 Da was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL  
25    35211; Product Code 100 DL mPEG 5000 5CE). Polyvinyl alcohol (85-89% hydrolyzed) was purchased from EMD Chemicals (Product Number 1.41350.1001).

**Method**

          Solutions were prepared as follows:

30        Solution 1: PLGA at 75 mg/mL and PLA-PEG at 25 mg/mL in methylene chloride. The solution was prepared by dissolving PLGA and PLA-PEG in pure methylene chloride.

          Solution 2: Rapamycin at 100 mg/mL in methylene chloride. The solution was prepared by dissolving rapamycin in pure methylene chloride.

Solution 3: Polyvinyl alcohol at 50 mg/mL in 100 mM pH 8 phosphate buffer.

An oil-in-water emulsion was used to prepare the nanocarriers. The O/W emulsion was prepared by combining solution 1 (1 mL), solution 2 (0.1 mL), and solution 3 (3 mL) in a small pressure tube and sonicating at 30% amplitude for 60 seconds using a Branson Digital Sonifier 250. The O/W emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution (30 mL) and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers was washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,000×g and 4 °C for 35 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

Nanocarrier size was determined by dynamic light scattering. The amount rapamycin in the nanocarrier was determined by HPLC analysis. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.

Effective Diameter (nm)	Rapamycin Content (% w/w)
227	6.4

**Example 5: Evaluating Tolerogenic Immune Responses Following Concomitant Administration of Synthetic Nanocarriers Comprising Immunosuppressant and Therapeutic Proteins**

**Tolerogenic Responses to Keyhole Limpet Hemocyanin**

C57BL/6 age-matched (5-6 weeks) female were injected with 200µg Keyhole Limpet Hemocyanin (KLH) in 200µl PBS i.v. and administered according to the schedule below. Injections for select groups were mixed with the synthetic nanocarriers just prior to injection. Anti-KLH IgG antibody titer was determined. The anaphylactic score induced by repetitive i.v. injections was also determined.

Group	Immunization KLH i.v. 200ug	Treatments tSIP Rapa i.v.	Description	# mice
1	d0, 7, 14, 21		Control immunization	5
2	d7, 14, 21		Control immunization	5
3	d0, 7, 14, 21	100ug d0-21	Test whether highest dose to inhibits the response	5
4	same	50ug d0-21	Test high dose to inhibits the response	5
5	same	30ug d0-21	Test lowest dose to inhibits the response	5
6	same	10ug d0-21	Test ineffective dose to inhibits the response	5
7	same	100ug d0	Test whether highest dose tolerized towards an antigen	5
8	same	50ug d0	Test high dose to inhibits the response	5
9	same	30ug d0	Test lowest dose to inhibits the response	5
10	same	10ug d0	Test ineffective dose to inhibits the response	5
				50

The results in **Fig. 1** show that multiple doses of synthetic nanocarriers administered concomitantly with KLH were effective in reducing the antibody response as observed at day 26. The animals presenting high antibody responses were eliminated from subsequent boosts.

Only animals with low or undetectable titers were kept as well as the control. These consisted mainly of the animals having received multiple doses of synthetic nanocarriers at 100µg, 50µg and 30µg. After day 21, all mice received 5 further injections of just KLH i.v. As seen in **Fig. 2**, untreated mice experienced a significant boost in their antibody titers while synthetic nanocarrier-treated mice had little to no antibody increase between days 21 to 54 (black symbols). Injection of KLH i.v. induces an anaphylactic shock that can be evaluated by observing the following symptoms: 0 score represented normal behavior, 1 represented decreased spontaneous activity, 2 represented decreased activity with falling over and a loss of the self-righting reflex, and 3 was assigned to mice that became moribund with cool, cyanotic extremities (Liu, E., et al. (2002). "Anti-peptide autoantibodies and fatal anaphylaxis in NOD mice in response to insulin self-peptides B:9-23 and B:13-23." *Journal of Clinical Investigation* 110(7): 1021-1027.). Similar to the titers, anaphylaxis was observed only in control animals, while little to no anaphylaxis was observed in synthetic nanocarrier-treated animals. Thus, it is shown that compositions provided herein when administered over a period of time concomitantly with a protein can reduce or prevent antibody responses and the resulting anaphylaxis as a result of administration of the protein in a subject.

**Example 6: Inhibition of Anaphylaxis in an OVA-Alum Model****Materials**

Rapamycin was purchased from TSZ CHEM (185 Wilson Street, Framingham, MA 01702; Product Code R1017). PLGA with a lactide:glycolide ratio of 3:1 and an inherent viscosity of 0.75 dL/g was purchased from SurModics Pharmaceuticals (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 7525 DLG 7A). PLA-PEG-OMe block copolymer with a methyl ether terminated PEG block of approximately 5,000 Da and an overall inherent viscosity of 0.5 DL/g was purchased from Lakeshore Biochemicals (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 100 DL mPEG 5000 5CE). EMPROVE® Polyvinyl Alcohol 4-88, USP (85-89% hydrolyzed, viscosity of 3.4-4.6 mPa·s) was purchased from EMD Chemicals Inc. (480 South Democrat Road Gibbstown, NJ 08027. Product Code 1.41350).

**Method**

Solutions were prepared as follows:

Solution 1: PLGA at 75 mg/mL, PLA-PEG-OMe at 25 mg/mL, and rapamycin at 12.5 mg/mL in methylene chloride. The solution was prepared by dissolving PLGA, PLA-PEG-OMe, and rapamycin in pure methylene chloride. Solution 2: Polyvinyl alcohol @ 50 mg/mL in 100 mM pH 8 phosphate buffer.

An oil-in-water emulsion was used to prepare the nanocarriers. The O/W emulsion was prepared by combining Solution 1 (1.0 mL) and Solution 2 (3.0 mL) in a small pressure tube and sonicating at 30% amplitude for 60 seconds using a Branson Digital Sonifier 250. The O/W emulsion was added to a beaker containing 70 mM pH 8 phosphate buffer solution and stirred at room temperature for 2 hours to allow the methylene chloride to evaporate and for the nanocarriers to form. A portion of the nanocarriers was washed by transferring the nanocarrier suspension to a centrifuge tube and centrifuging at 75,600×g and 4 °C for 50 min, removing the supernatant, and re-suspending the pellet in phosphate buffered saline. The washing procedure was repeated, and the pellet was re-suspended in phosphate buffered saline for a final nanocarrier dispersion of about 10 mg/mL.

Nanocarrier size was determined by dynamic light scattering. The amount of rapamycin in the nanocarrier was determined by HPLC analysis. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.

Effective Diameter (nm)	Rapamycin Content (% w/w)
241	11.5

### **Suppression of Intravenously-Induced Anaphylactic Reactions using Tolerogenic Nanocarriers**

Control Balb/c age-matched (5-6 weeks) female mice (d0) were left naïve or injected twice intraperitoneously (i.p., d0 and d5) with 5µg of whole Ovalbumin protein with 40µg of Alum (OVA+Alum). Other groups received a similar injection but 100µg of Rapamycin were injected free in solution i.v. or contained in 0.9mg of synthetic nanocarriers i.p. (admixed to the OVA+Alum), intragastrically (i.g.) intravenously (i.v.) (tail vein) or subcutaneously (hind limb). Fifteen days later all animals were challenged with a bolus of OVA i.v. (25µg). As expected, animals that were not immunized did not react to the OVA infusion whereas a strong anaphylaxis reaction was recorded in untreated immunized animals as shown in **Fig. 3**. Treatment of the animals with free rapamycin or synthetic nanocarriers i.g. did not reduce the anaphylactic reaction whereas providing the synthetic nanocarriers i.v., i.p. or s.c. abolished it.

These results show that compositions provided herein when administered concomitantly with a protein can reduce or prevent anaphylaxis induced by an infusion of the protein. Anaphylaxis scores were determined by three independent observers as follows: 0=no symptom, 1=lethargy, 2=lethargy and inability to right, 3=moribund. The bars represent the mean +/- standard error.

### **Example 7: Antigen-Specific Tolerogenic Responses to Chicken Ovalbumin with Encapsulated Rapamycin**

#### **NP[Rapa] Materials and Methods**

##### **Materials**

Rapamycin was purchased from TSZ CHEM (185 Wilson Street, Framingham, MA 01702), product code R1017. PLGA with a lactide:glycolide ratio of 1:1 and an inherent viscosity of 0.24 dL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211), product code 5050 DLG 2.5A. PLA-PEG-OMe block co-polymer

with a methyl ether terminated PEG block of approximately 5,000 Da and an overall inherent viscosity of 0.50 DL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211), product code 100 DL mPEG 5000 5CE. EMPROVE® Polyvinyl Alcohol 4-88, USP (85-89% hydrolyzed, viscosity of 3.4-4.6 mPa·s) was purchased from EMD Chemicals Inc. (480 South Democrat Road Gibbstown, NJ 08027), product code 1.41350. Cellgro phosphate buffered saline 1X (PBS 1X) was purchased from Corning (9345 Discovery Blvd. Manassas, VA 20109), product code 21-040-CV.

## Method

Solutions were prepared as follows:

Solution 1: A polymer and rapamycin mixture was prepared by dissolving PLGA at 75 mg per 1 mL, PLA-PEG-Ome at 25 mg per 1 mL, and rapamycin as 12.5 mg per 1 mL in dichloromethane. Solution 2: Polyvinyl alcohol was prepared at 50 mg/mL in 100 mM pH 8 phosphate buffer.

An O/W emulsions was prepared by combining Solution 1 (1.0 mL) and Solution 2 (3.0 mL) in a small glass pressure tube and sonicating at 30% amplitude for 60 seconds using a Branson Digital Sonifier 250. The O/W emulsion was added to an open beaker containing 70 mM pH 8 phosphate buffer solution (60 mL). Three additional, identical O/W emulsions were prepared and added to the same beaker as the first. These were then stirred at room temperature for 2 hours to allow the dichloromethane to evaporate and for the nanocarriers to form. A portion of the nanocarriers was washed by transferring the nanocarrier suspension to centrifuge tubes and centrifuging at 75,600×g and 4 °C for 35 minutes, removing the supernatant, and re-suspending the pellet in PBS 1X. The wash procedure was repeated and then the pellet was re-suspended in PBS 1X to achieve a nanocarrier suspension having a nominal concentration of 10 mg/mL on a polymer basis. An identical formulation was prepared as above in a separate beaker, and combined with the first after the wash step. The mixed nanocarrier solution was then filtered using 1.2µm PES membrane syringe filters from Pall part number 4656, and stored at -20°C.

Nanocarrier size was determined by dynamic light scattering. The amount of rapamycin in the nanocarrier was determined by HPLC analysis. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.



Effective Diameter (nm)	Rapamycin Content (% w/w)
220	11.85

## NP[OVA] Materials and Methods

### Materials

5 Ovalbumin protein, was purchased from Worthington Biochemical Corporation (730 Vassar Avenue, Lakewood, NJ 08701), product code LS003054). PLGA with 54% lactide and 46% glycolide content and an inherent viscosity of 0.24 dL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211), product code 5050 DLG 2.5A). PLA-PEG block co-polymer with a methyl ether terminated PEG block of  
10 approximately 5,000 Da and Mw of 28,000 Da, inherent viscosity of 0.38 dL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211), product code 100 DL mPEG 5000 4CE. EMPROVE® Polyvinyl Alcohol 4-88, USP, 85-89% hydrolyzed, viscosity of 3.4-4.6 mPa.s, was purchased from EMD Chemicals Inc. (480 South Democrat Road Gibbstown, NJ 08027), product code 1.41350.1001. Cellgro Phosphate-buffered saline  
15 1X (PBS 1X) was purchased from Corning (9345 Discovery Blvd. Manassas, VA 20109), product code 21-040-CV.

### Method

Solutions were prepared as follows:

20 Solution 1: Ovalbumin protein @ 50 mg/mL was prepared in 10mM phosphate buffer pH 8 with 10% by weight sucrose. Solution 2: PLGA was prepared by dissolving PLGA at 100 mg per 1 mL of dichloromethane in the chemical fume hood. Solution 3: PLA-PEG-OMe was prepared by dissolving PLA-PEG-OMe at 100 mg per 1 mL of dichloromethane in the chemical fume hood. Solution 4: Polyvinyl alcohol @ 65 mg/mL in 100mM phosphate buffer,  
25 pH 8.

A primary (W1/O) emulsion was first created by mixing Solutions 1 through 3. Solution 1 (0.2 mL), Solution 2 (0.75 mL), and Solution 3 (0.25mL) were combined in a small glass pressure tube which was pre-chilled >4 minutes in an ice water bath, and sonicated at 50% amplitude for 40 seconds over an ice bath using a Branson Digital Sonifier 250. A  
30 secondary (W1/O/W2) emulsion was then formed by adding Solution 4 (3 mL) to the primary

emulsion, vortex mixing to create a milky dispersion, and then sonicating at 30% amplitude for 60 seconds over an ice bath using the Branson Digital Sonifier 250. The secondary emulsion was added to an open 50 mL beaker containing PBS 1X (30 mL). A second identical double emulsion formulation was prepared as described above, and added to the same 50 mL beaker as the first. The two preparations were stirred at room temperature for 2 hours to allow the dichloromethane to evaporate and the nanocarriers to form in suspension. A portion of the suspended nanocarriers was washed by transferring the nanocarrier suspension to a centrifuge tube, spinning at 75,600 rcf for 50 minutes, removing the supernatant, and re-suspending the pellet in PBS 1X. This washing procedure was repeated and then the pellet was re-suspended in PBS 1X to achieve a nanocarrier suspension having a nominal concentration of 10 mg/mL on a polymer basis. The suspension was stored frozen at -20C until use.

Effective Diameter (nm)	Ovalbumin Content (% w/w)
164	5.81

## NP[GSK1059615] Materials and Methods

### Materials

GSK1059615 was purchased from MedChem Express (11 Deer Park Drive, Suite 102D Monmouth Junction, NJ 08852), product code HY-12036. PLGA with a lactide:glycolide ratio of 1:1 and an inherent viscosity of 0.24 dL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211), product code 5050 DLG 2.5A. PLA-PEG-OMe block co-polymer with a methyl ether terminated PEG block of approximately 5,000 Da and an overall inherent viscosity of 0.26 DL/g was purchased from Lakeshore Biomaterials (756 Tom Martin Drive, Birmingham, AL 35211; Product Code 100 DL mPEG 5000 5K-E). Cellgro phosphate buffered saline 1X pH 7.4 (PBS 1X) was purchased from Corning (9345 Discovery Blvd. Manassas, VA 20109), product code 21-040-CV.

## Method

Solutions were prepared as follows:

Solution 1: PLGA (125 mg), and PLA-PEG-OMe (125 mg), were dissolved in 10 mL of acetone. Solution 2: GSK1059615 was prepared at 10 mg in 1 mL of N-methyl-2-pyrrolidinone (NMP).

Nanocarriers were prepared by combining Solution 1 (4 mL) and Solution 2 (0.25 mL) in a small glass pressure tube and adding the mixture drop wise to a 250 mL round bottom flask containing 20 mL of ultra-pure water under stirring. The flask was mounted onto a rotary evaporation device, and the acetone was removed under reduced pressure. A portion of the nanocarriers was washed by transferring the nanocarrier suspension to centrifuge tubes and centrifuging at 75,600 rcf and 4 °C for 50 minutes, removing the supernatant, and re-suspending the pellet in PBS 1X. The washing procedure was repeated, and the pellet was re-suspended in PBS 1X to achieve a nanocarrier suspension having a nominal concentration of 10 mg/mL on a polymer basis. The washed nanocarrier solution was then filtered using 1.2µm PES membrane syringe filters from Pall, part number 4656. An identical nanocarrier solution was prepared as above, and pooled with the first after the filtration step. The homogenous suspension was stored frozen at -20°C.

Nanocarrier size was determined by dynamic light scattering. The amount of GSK1059615 in the nanocarrier was determined by UV absorption at 351nm. The total dry-nanocarrier mass per mL of suspension was determined by a gravimetric method.

Effective Diameter (nm)	GSK1059615 Content (% w/w)
143	1.02

C57BL/6 age-matched (5-6 weeks) female mice were injected i.v. in the tail vein on days -21 and -14 with saline (No Treatment), 1.1mg of whole Ovalbumin-loaded nanocarriers (NP[OVA]) combined to either 1.2mg of rapamycin-containing nanocarriers (NP[Rapa]) or 8mg of GSK1059615-loaded nanocarriers (NP[GSK1059615]).

At day 0 all animals were injected s.c. in the hind limbs with 25µg of particulate OVA (pOVA) admixed to 2µg of CpG followed by injections of just 25µg pOVA on days 7 and 14. Antibody titers were measured on day 21. In absence of any treatment, the animals developed a robust immune response against OVA that can be measured by the anti-OVA IgG antibody

titers. The antibody titers at day 21 shown in **Fig. 4** demonstrate that 2 doses of synthetic tolerogenic nanocarriers administered concomitantly with encapsulated OVA in the same solution (NP[OVA]+NP[Rapa] or NP[GSK1059615]) were effective in reducing antibody formation to OVA even after 1 injection of OVA+CpG and 2 injections of OVA alone. These results show that encapsulated immunosuppressants (such as rapamycin and GSK1059615) when concomitantly delivered with a protein can prevent antibody formation to that protein for multiple challenges and periods of time, such an effect is helpful in reducing or preventing the occurrence of anaphylaxis.

#### **Example 8: Allergenic IgE-Mediated Anaphylaxis versus IgG-Mediated Anaphylaxis (Prophetic)**

Subjects that have allergies to latex and subjects that have presented with anaphylaxis following infusion of von Willebrand's factor ("vWF") are recruited for a set of pharmacodynamic studies. Anaphylaxis is induced in the two sets of subjects, and blood samples are taken before, during, and following anaphylactic episodes.

The blood samples are assayed for increased T cell IL4R- $\alpha$  expression, and decreased blood neutrophil Fc $\gamma$ RIII expression. In latex allergenic subjects, increased T cell IL4R- $\alpha$  expression, without decreased blood neutrophil Fc $\gamma$ RIII expression is characterized as being indicative of allergenic IgE-mediated anaphylaxis. In vWF sensitized subjects, decreased blood neutrophil Fc $\gamma$ RIII expression, without T cell IL4R- $\alpha$  expression is characterized as being indicative of IgG-mediated anaphylaxis.

#### **Example 9: Monoclonal Antibody-Induced Anaphylaxis (Prophetic)**

Subjects are repeatedly exposed to infliximab (REMICADE), and any presentation of anaphylaxis in the subjects is noted. Blood samples are taken from the subjects before, during, and following anaphylactic episodes.

The blood samples are assayed for increased T cell IL4R- $\alpha$  expression, and decreased blood neutrophil Fc $\gamma$ RIII expression. Decreased blood neutrophil Fc $\gamma$ RIII expression, without T cell IL4R- $\alpha$  expression is characterized as being indicative of IgG-mediated anaphylaxis.

#### **Example 10: Messenger RNA-Induced Anaphylaxis (Prophetic)**

Modified mRNA encoding erythropoietin is prepared according to US Patent application 2013/0115272 to de Fougères et al. ("mmRNA"). Subjects are repeatedly

exposed to mmRNA, and any presentation of anaphylaxis in the subjects is noted. Blood samples are taken from the subjects before, during, and following anaphylactic episodes.

The blood samples are assayed for increased T cell IL4R- $\alpha$  expression, and decreased blood neutrophil Fc $\gamma$ RIII expression. Decreased blood neutrophil Fc $\gamma$ RIII expression, without  
5 T cell IL4R- $\alpha$  expression is characterized as being indicative of IgG-mediated anaphylaxis.

**Example 11: Antigen-Specific Tolerogenic Responses to Chicken Ovalbumin with Nanocrystalline Rapamycin (Prophetic)**

C57BL/6 age-matched (5-6 weeks) female mice are injected i.v. in the tail vein on days  
10 -21 and -14 with saline (No Treatment) or 1.1mg of whole Ovalbumin and 1.2mg of nanocrystalline rapamycin. At day 0 all animals are injected s.c. in the hind limbs with 25 $\mu$ g of particulate OVA (pOVA) admixed to 2 $\mu$ g of CpG followed by injections of just 25 $\mu$ g pOVA on days 7 and 14. Antibody titers are measured on day 21. In the absence of any treatment, the animals develop a robust immune response against OVA that can be measured by the anti-  
15 OVA IgG antibody titers.

A reduction in OVA-specific IgG antibodies in the animals that received OVA in combination with nanocrystalline rapamycin indicates that the nanocrystal-form of the immunosuppressant when concomitantly delivered with antigen can prevent antibody formation to the antigen for multiple challenges and periods of time and can be useful in  
20 reducing or preventing anaphylaxis.

**Example 12: Reduction of Monoclonal Antibody-Induced Anaphylaxis with Nanocarriers (Prophetic)**

Subjects are repeatedly exposed to infliximab (REMICADE). Control groups are left  
25 untreated (No Treatment) whereas three other groups are treated with synthetic nanocarriers attached to GSK1059615 (NP[GSK]). Blood samples are taken from the subjects before, during, and following anaphylactic episodes. The extent of anaphylaxis in response to REMICADE is assessed and the titer of REMICADE-specific IgG antibody titers in the blood of the animals is quantified. The blood samples are also assayed for increased T cell IL4R- $\alpha$   
30 expression, and decreased blood neutrophil Fc $\gamma$ RIII expression.

A reduction in the symptoms of anaphylaxis and/or the titer of anti-REMICADE IgG antibodies in the animals that were exposed to REMICADE concomitantly with nanocarriers attached to GSK1059615 as compared to animals that did not received the nanocarrier

compositions indicates that the nanocarriers attached to GSK1059615 when concomitantly delivered with an antigen, such as REMICADE, is able to reduce or prevent anaphylaxis.

**Example 13: Allergenic IgE-Mediated Anaphylaxis versus IgG-Mediated Anaphylaxis**

**(Prophetic)**

Subjects that have allergies to latex and subjects that have presented with anaphylaxis following infusion of von Willebrand's factor ("vWF") are recruited for a set of pharmacodynamic studies. Anaphylaxis is induced in the two sets of subjects in the presence or absence of nanocrystalline rapamycin, and blood samples are taken before, during, and following anaphylactic episodes.

The blood samples are assayed for increased T cell IL4R- $\alpha$  expression, and decreased blood neutrophil Fc $\gamma$ RIII expression. In latex allergenic subjects, increased T cell IL4R- $\alpha$  expression, without decreased blood neutrophil Fc $\gamma$ RIII expression is characterized as being indicative of allergenic IgE-mediated anaphylaxis. In vWF sensitized subjects, decreased blood neutrophil FRIII expression, without T cell IL4R expression is characterized as being indicative of IgG-mediated anaphylaxis. A reduction in vWF-specific responses in the subjects that received nanocrystalline rapamycin as compared to the subjects that did not, indicates that the nanocrystalline immunosuppressant is able to reduce or prevent anaphylaxis.

**Example 14: Messenger RNA-Induced Anaphylaxis (Prophetic)**

Modified mRNA encoding erythropoietin is prepared according to US Patent application 2013/0115272 to de Fougérolles et al. ("mmRNA"). Subjects are repeatedly exposed to mmRNA in the presence or absence of nanocrystalline rapamycin, and any presentation of anaphylaxis in the subjects is noted. Blood samples are taken from the subjects before, during, and following anaphylactic episodes.

The blood samples are assayed for increased T cell IL4R- $\alpha$  expression, and decreased blood neutrophil Fc $\gamma$ RIII expression. Decreased blood neutrophil Fc $\gamma$ RIII expression, without T cell IL4R- $\alpha$  expression is characterized as being indicative of IgG-mediated anaphylaxis. A reduction in mmRNA-specific responses in the subjects that received nanocrystalline rapamycin as compared to the subjects that did not, indicates that the nanocrystalline immunosuppressant is able to reduce or prevent anaphylaxis.

What is claimed is:

### CLAIMS

1. A method for reducing or preventing an anaphylactic reaction to a non-allergenic  
5 antigen, comprising:  
    providing the non-allergenic antigen, the non-allergenic antigen being unattached to  
synthetic nanocarriers;  
    providing a composition comprising synthetic nanocarriers attached to an  
immunosuppressant; and  
10 administering to a subject the composition concomitantly with the non-allergenic  
antigen.
2. The method of claim 1, wherein the concomitant administration to the subject is  
according to a protocol that has been demonstrated to reduce or prevent an anaphylactic  
15 reaction.
3. The method of claim 1 or 2, wherein the method further comprises determining the  
protocol.
- 20 4. The method of any one of claims 1-3, wherein the method further comprises assessing  
the anaphylactic reaction in the subject prior to and/or after the administration.
5. The method of any one of claims 1-4, wherein the administering is by intravenous,  
intra-peritoneal or subcutaneous administration.  
25
6. The method of any one of claims 1-5, wherein the method further comprises recording  
a reduction or prevention of an anaphylactic reaction to the non-allergenic antigen.
7. The method or composition or kit of any one of the preceding claims, wherein the  
30 immunosuppressant comprises a statin, an mTOR inhibitor, a TGF- $\beta$  signaling agent, a  
corticosteroid, an inhibitor of mitochondrial function, a P38 inhibitor, an NF- $\kappa$ B inhibitor, an  
adenosine receptor agonist, a prostaglandin E2 agonist, a phosphodiesterase 4 inhibitor, an  
HDAC inhibitor or a proteasome inhibitor.

8. The method or composition or kit of claim 7, wherein the mTOR inhibitor is rapamycin.

5 9. The method or composition or kit any of the preceding claims, wherein the non-allergenic comprises a therapeutic macromolecule.

10. The method or composition or kit of claim 9, wherein the therapeutic macromolecule is a therapeutic protein or a therapeutic polynucleotide.

10

11. The method or composition or kit of claim 10, wherein the therapeutic protein is for protein replacement of protein supplementation therapy.

12. The method or composition or kit of claim 10, wherein the therapeutic protein  
15 comprises a/an infusible or injectable therapeutic protein, enzyme, enzyme cofactor, hormone, blood or blood coagulation factor, cytokine, interferon, growth factor, monoclonal antibody, polyclonal antibody, or protein associated with Pompe's disease.

13. The method or composition or kit of claim 12, wherein the infusible or injectable  
20 therapeutic protein comprises Tocilizumab, alpha-1 antitrypsin, Hematide, albinterferon alfa-2b, Thucin, tesamorelin, ocrelizumab, belimumab, pegloticase, taliglucerase alfa, agalsidase alfa, or velaglucerase alfa.

14. The method or composition or kit of claim 12, wherein the enzyme comprises an  
25 oxidoreductase, transferase, hydrolase, lysase, isomerase or ligase.

15. The method or composition or kit of claim 12, wherein the enzyme comprises an enzyme for enzyme replacement therapy for a lysosomal storage disorder.

30 16. The method or composition or kit of claim 15, wherein the enzyme for replacement therapy for a lysosomal storage disorder comprises imiglucerase, a-galactosidase A (a-gal A), agalsidase beta, acid  $\alpha$ -glucosidase (GAA), alglucosidase alfa, LUMIZYME, MYOZYME,



arylsulfatase B, laronidase, ALDURAZYME, idursulfase, ELAPRASE, arylsulfatase B, pegloticase, pegsiticase or NAGLAZYME.

17. The method or composition or kit of claim 12, wherein the cytokine comprises a lymphokine, interleukin, chemokine, type 1 cytokine or a type 2 cytokine.
18. The method or composition or kit of claim 12, wherein the blood or blood coagulation factor comprises Factor I, Factor II, tissue factor, Factor V, Factor VII, Factor VIII, Factor IX, Factor X, Factor Xa, Factor XII, Factor XIII, von Willebrand factor, prekallikrein, high-molecular weight kininogen, fibronectin, antithrombin III, heparin cofactor II, protein C, protein S, protein Z, protein Z-related protease inhibitor (ZPI), plasminogen, alpha 2-antiplasmin, tissue plasminogen activator (tPA), urokinase, plasminogen activator inhibitor-1 (PAI1), plasminogen activator inhibitor-2 (PAI2), cancer procoagulant or epoetin alfa.
19. The method or composition or kit of any one of the preceding claims, wherein a load of immunosuppressant attached to the synthetic nanocarriers, on average across the synthetic nanocarriers, is between 0.1% and 50%.
20. The method or composition or kit of claim 19, wherein the load is between 0.1% and 20%.
21. The method or composition or kit of any one of the preceding claims, wherein the synthetic nanocarriers comprise lipid nanoparticles, polymeric nanoparticles, metallic nanoparticles, surfactant-based emulsions, dendrimers, buckyballs, nanowires, virus-like particles or peptide or protein particles.
22. The method or composition or kit of claim 21, wherein the synthetic nanocarriers comprise lipid nanoparticles.
23. The method or composition or kit of claim 21, wherein the synthetic nanocarriers comprise liposomes.

24. The method or composition or kit of claim 21, wherein the synthetic nanocarriers comprise metallic nanoparticles.
25. The method or composition or kit of claim 24, wherein the metallic nanoparticles  
5 comprise gold nanoparticles.
26. The method or composition or kit of claim 21, wherein the synthetic nanocarriers comprise polymeric nanoparticles.
- 10 27. The method or composition or kit of claim 26, wherein the polymeric nanoparticles comprise polymer that is a non-methoxy-terminated, pluronic polymer.
28. The method or composition or kit of claim 26 or 27, wherein the polymeric nanoparticles comprise a polyester, polyester attached to a polyether, polyamino acid,  
15 polycarbonate, polyacetal, polyketal, polysaccharide, polyethyloxazoline or polyethyleneimine.
29. The method or composition or kit of claim 28, wherein the polyester comprises a poly(lactic acid), poly(glycolic acid), poly(lactic-co-glycolic acid) or polycaprolactone.
- 20 30. The method or composition or kit of claim 28 or 29, wherein the polymeric nanoparticles comprise a polyester and a polyester attached to a polyether.
31. The method or composition or kit of any one of claims 28-30, wherein the polyether comprises polyethylene glycol or polypropylene glycol.  
25
32. The method or composition or kit of any one of the preceding claims, wherein the mean of a particle size distribution obtained using dynamic light scattering of the synthetic nanocarriers is a diameter greater than 100nm.
- 30 33. The method or composition or kit of claim 32, wherein the diameter is greater than 150nm.

34. The method or composition or kit of claim 33, wherein the diameter is greater than 200nm.
35. The method or composition or kit of claim 34, wherein the diameter is greater than 250nm.
36. The method or composition or kit of claim 35, wherein the diameter is greater than 300nm.
37. The method or composition or kit of any one of the preceding claims, wherein an aspect ratio of the synthetic nanocarriers is greater than 1:1, 1:1.2, 1:1.5, 1:2, 1:3, 1:5, 1:7 or 1:10.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

# PATENT COOPERATION TREATY

# PCT

## DECLARATION OF NON-ESTABLISHMENT OF INTERNATIONAL SEARCH REPORT (PCT Article 17(2)(a), Rules 13ter.1(c) and (d) and 39)

Applicant's or agent's file reference <b>S1681.70064</b>	<b>IMPORTANT DECLARATION</b>	Date of mailing ( <i>day/month/year</i> ) <b>09 September 2014 (09.09.2014)</b>
International application No. <b>PCT/US2014/036699</b>	International filing date ( <i>day/month/year</i> ) <b>02 May 2014 (02.05.2014)</b>	(Earliest) Priority date ( <i>day/month/year</i> ) <b>03 May 2013 (03.05.2013)</b>
International Patent Classification (IPC) or both national classification and IPC <b>A61K 9/16(2006.01)i, A61K 47/30(2006.01)i, A61K 48/00(2006.01)i</b>		
Applicant <b>SELECTA BIOSCIENCES, INC. et al</b>		

This International Searching Authority hereby declares, according to Article 17(2)(a), that **no international search report will be established** on the international application for the reasons indicated below.

1. ☒ The subject matter of the international application relates to:
  - a. ☐ scientific theories
  - b. ☐ mathematical theories
  - c. ☐ plant varieties
  - d. ☐ animal varieties
  - e. ☐ essentially biological processes for the production of plants and animals, other than microbiological processes and the products of such processes
  - f. ☐ schemes, rules or methods of doing business
  - g. ☐ schemes, rules or methods of performing purely mental acts
  - h. ☐ schemes, rules or methods of playing games
  - i. ☒ methods for treatment of the human body by surgery or therapy
  - j. ☐ methods for treatment of the animal body by surgery or therapy
  - k. ☐ diagnostic methods practised on the human or animal body
  - l. ☐ mere presentation of information
  - m. ☐ computer programs for which this International Searching Authority is not equipped to search prior art
2. ☒ The failure of the following parts of the international application to comply with prescribed requirements prevents a meaningful search from being carried out:
 

☐ the description
 ☒ the claims
 ☐ the drawings
3. ☐ A meaningful search could not be carried out without the sequence listing; the applicant did not, within the prescribed time limit:
 

☐ furnish a sequence listing on paper complying with the standard provided for in Annex C of the Administrative Instructions, and such listing was not available to the International Searching Authority in a form and manner acceptable to it.  
☐ furnish a sequence listing in electronic form complying with the standard provided for in Annex C of the Administrative Instructions, and such listing was not available to the International Searching Authority in a form and manner acceptable to it.  
☐ pay the required late furnishing fee for the furnishing of a sequence listing in response to an invitation under Rule 13ter.1(a) or (b).

4. Further comments:

Name and mailing address of ISA/KR International Application Division Korean Intellectual Property Office 189 Cheongsu-ro, Seo-gu, Daejeon Metropolitan City, 302-701, Republic of Korea Facsimile No. +82-42-472-7140	Authorized officer  CHOI, Sung Hee  Telephone No. +82-42-481-8740
--	---

