



- (51) **International Patent Classification:**
C09B 29/00 (2006.01) *C11D 3/42* (2006.01)
C09B 69/00 (2006.01)
- (21) **International Application Number:** PCT/US2012/039537
- (22) **International Filing Date:** 25 May 2012 (25.05.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/492,935 3 June 2011 (03.06.2011) US
13/479,340 24 May 2012 (24.05.2012) US
- (71) **Applicant (for all designated States except US):** MIL-LIKEN & COMPANY [US/US]; 920 Milliken Road, M-495, Spartanburg, South Carolina 29303 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** TORRES, Eduardo [US/US]; 145 Lewis Burnett Drive, Boiling Springs, South Carolina 29316 (US). BRUHNKE, John D. [US/US]; 125 Hillbrook Drive, Spartanburg, South Carolina 29307 (US). DEY, Sanjeev K. [IN/US]; 8308 Meridian River Run, Spartanburg, South Carolina 29301 (US). MIRACLE, Gregory S. [US/US]; 7063 Francis Drive, Liberty Township, Ohio 45044 (US).
- (74) **Agent:** WENTZ, Brenda D.; 920 Milliken Road, M-495, Spartanburg, South Carolina 29303 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report (Art. 21(3))



WO 2012/166584 A1

(54) **Title:** THIOPHENE AZO CARBOXYLATE DYES AND LAUNDRY CARE COMPOSITIONS CONTAINING THE SAME

(57) **Abstract:** This application relates to thiophene azo carboxylate dyes for use as hueing agents, laundry care compositions comprising such dyes that may serve as hueing agents, processes for making such dyes and laundry care compositions and methods of using the same. The aforementioned dyes contain a formally charged moiety and are generally comprised of at least two components: at least one chromophore component and at least one polymeric component. Suitable chromophore components generally fluoresce blue, red, violet, or purple color when exposed to ultraviolet light, or they may absorb light to reflect these same shades. Such dyes are advantageous in providing a hueing effect, for example, a whitening effect to fabrics, while not building up over time and causing undesirable blue discoloration to the treated fabrics. Such dyes are also generally stable to bleaching agents used in laundry care compositions.

THIOPHENE AZO CARBOXYLATE DYES AND LAUNDRY CARE
COMPOSITIONS CONTAINING THE SAME

5

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority to US Provisional Patent Application Number 61/492,935, entitled "Thiophene Azo Carboxylate Dyes and Laundry Care Compositions Containing the Same," which was filed on June 3, 2011, and is
10 entirely incorporated by reference herein.

TECHNICAL FIELD

This application relates to thiophene azo carboxylate dyes for use as hueing agents, laundry care compositions comprising such dyes, processes for making
15 such dyes and laundry care compositions and methods of using the same. The aforementioned dyes contain a formally charged moiety and are generally comprised of at least two components: at least one chromophore component and at least one polymeric component. Suitable chromophore components generally fluoresce blue, red, violet, or purple color when exposed to ultraviolet
20 light, or they may absorb light to reflect these same shades. These dyes are advantageous in providing a hueing effect, for example, a whitening effect to fabrics, while not building up over time and causing undesirable discoloration to the treated fabrics. Such dyes are also generally stable to bleaching agents used in laundry care compositions.

25

BACKGROUND

As textile substrates age, their color tends to fade or yellow due to exposure to light, air, soil, and natural degradation of the fibers that comprise the substrates. Thus, the purpose of hueing agents is generally to visually brighten these textile
30 substrates and counteract the fading and yellowing of the substrates. Typically, hueing agents may be found in laundry detergents, fabric softeners, or rinse aids and are therefore applied to textile substrates during the laundering process. However, it is important that hueing agents function to visually brighten treated

textile substrates without causing undesirable staining of the textile substrates. Cellulosic substrates, in particular, tend to exhibit a yellow hue after exposure to light, air, and/or soiling. This yellowness is often difficult to reverse by normal laundering procedures. As a result, there exists a need for improved hueing agents which are capable of eliminating the yellowness exhibited by ageing
5 cellulosic substrates. By utilizing such improved hueing agents, the life of the textile substrates, such as clothing articles, table linens, etc., may be extended. Unfortunately, current thiophene azo dyes, are not as physically stable as desired. Thus, they require special solvents/handling which can limit liquid dye
10 product and laundry care composition formulation flexibility. In addition, current thiophene azo carboxylate dyes disclosed for use in laundry compositions, either do not provide the desired hueing benefit at the levels that can be used in products, are too costly and/or they build up to an undesirable level, thus overhueing the treated situs over multiple treatment cycles.

15

The thiophene azo carboxylate dyes disclosed herein and the laundry care compositions comprising same offer advantages over previous efforts in this area, as, unlike previous thiophene azo hueing agents, the present thiophene azo carboxylate dyes comprise a formally charged carboxylate moiety (above its
20 pKa) that allows such dyes to provide enhanced deposition, removal and hue angle, without the negatives of pretreat spot staining on polyurea fibers such as Lycra® and/or excessive cost. While not being bound by theory, Applicants believe that such negatives are avoided as the instant thiophene azo carboxylate dyes interacts with metal ions, for example, in the wash water in a more
25 favorable manner. The aforementioned benefits are offered over a range of fabric types. In short, Applicants recognized the source of the current hueing deficiencies and herein provide the solution to such problem. The hueing compounds disclosed herein also absorb light at a wavelength appropriate to visually neutralize the yellowness of substrates, including textile substrates.
30 These compounds function ideally as hueing agents for substrates, including textile substrates, and may be incorporated into laundry care compositions for use by consumers.

SUMMARY OF INVENTION

This application relates to thiophene azo carboxylate dyes for use as hueing agents, laundry care compositions comprising such dyes that may serve as hueing agents, processes for making such dyes and laundry care compositions and methods of using the same. The aforementioned dyes contain a formally charged moiety and are generally comprised of at least two components: at least one chromophore component and at least one polymeric component. Suitable chromophore components generally fluoresce blue, red, violet, or purple color when exposed to ultraviolet light, or they may absorb light to reflect these same shades. Such dyes are advantageous in providing a hueing effect, for example, a whitening effect to fabrics, while not building up over time and causing undesirable blue discoloration to the treated fabrics. Such dyes are also generally stable to bleaching agents used in laundry care compositions.

15

DETAILED DESCRIPTION

As used herein, the term "alkoxy" is intended to include C₁-C₈ alkoxy and alkoxy derivatives of polyols having repeating units such as butylene oxide, glycidol oxide, ethylene oxide or propylene oxide.

20

As used herein, the terms "alkyl" and "alkyl capped" are intended to include C₁-C₁₈ alkyl groups, and in one aspect, C₁-C₆ alkyl groups.

As used herein, the term "aryl" is intended to include C₃-C₁₂ aryl groups.

25

As used herein, the term "arylalkyl" is intended to include C₁-C₁₈ alkyl groups and, in one aspect, C₁-C₆ alkyl groups.

30

As used herein, the term "formally charged moiety" means a moiety having at least one formal positive charge or at least one formal negative charge in aqueous solution at a pH in the range from 7 to 11.

The terms "ethylene oxide," "propylene oxide" and "butylene oxide" may be shown herein by their typical designation of "EO," "PO" and "BO," respectively.

5 As used herein, the term "laundry care composition" includes, unless otherwise indicated, granular, powder, liquid, gel, paste, unit dose bar form and/or flake type washing agents and/or fabric treatment compositions.

10 As used herein, the term "fabric treatment composition" includes, unless otherwise indicated, fabric softening compositions, fabric enhancing compositions, fabric freshening compositions and combinations thereof. Such compositions may be, but need not be rinse added compositions.

15 As used herein, "cellulosic substrates" are intended to include any substrate which comprises at least a majority by weight of cellulose. Cellulose may be found in wood, cotton, linen, jute, and hemp. Cellulosic substrates may be in the form of powders, fibers, pulp and articles formed from powders, fibers and pulp. Cellulosic fibers, include, without limitation, cotton, rayon (regenerated cellulose), acetate (cellulose acetate), triacetate (cellulose triacetate), and mixtures thereof. Articles formed from cellulosic fibers include textile articles such as fabrics.
20 Articles formed from pulp include paper.

As used herein, the articles including "the", "a" and "an" when used in a claim, are understood to mean one or more of what is claimed or described.

25 As used herein, the terms "include", "includes" and "including" are meant to be non-limiting.

As used herein, the term "maximum extinction coefficient" is intended to describe the molar extinction coefficient at the maximum wavelength in the range of 400
30 nanometers to 750 nanometers.

As a consequence of their manufacturing process, the thiophene azo carboxylate dyes disclosed herein may contain a distribution of repeating units in their polymeric moiety. Accordingly, in one aspect, the molecular weight of the thiophene azo carboxylate dyes disclosed herein may be reported as an average
5 molecular weight, as determined by its molecular weight distribution.

The test methods disclosed in the Test Methods Section of the present application should be used to determine the respective values of the parameters of Applicants' inventions.

10

Unless otherwise noted, all component or composition levels are in reference to the active portion of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources of such components or compositions.

15

All percentages and ratios are calculated by weight unless otherwise indicated. All percentages and ratios are calculated based on the total composition unless otherwise indicated.

20

It should be understood that every maximum numerical limitation given throughout this specification includes every lower numerical limitation, as if such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly
25 written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

30


All documents cited are, in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present invention.

Laundry Care Composition Comprising Suitable Hueing Agents

In one aspect, a laundry care composition comprising a hueing agent and one or more laundry care ingredients is disclosed. Suitable hueing agents include all the hueing agents listed below.


5

In one aspect, a composition comprising a thiophene azo carboxylate dye containing a carboxylic acid moiety, with the proviso that the thiophene azo carboxylate dye does not have the following structures



10

and



is disclosed.

15

In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has, in the wavelength range of about 400 nm to about 750 nm


in methanol solution, a maximum extinction coefficient greater than about 1000 liter/mol/cm.

In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has, in the wavelength range of about 540 nm to about 630 nm,
5 a maximum extinction coefficient from about 20,000 to about 100,000 liter/mol/cm.

In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has, in the wavelength range of about 560 nm to about 610 nm, a maximum extinction coefficient from about 20,000 to about 65,000 liter/mol/cm.

10 In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has a molecular weight from greater than 232 Daltons, from about 233 Daltons to about 5000 Daltons, from about 365 Daltons to about 2500 Daltons, or even from about 423 Daltons to about 1000 Daltons.

In one aspect of the aforementioned composition, said thiophene azo
15 carboxylate dye is represented by general Formula (I):




Formula I

wherein:

- 20 a.) R₁, R₂ and R₃ are each independently selected from hydrogen, electron-withdrawing moieties, and electron-donating moieties, provided that at least one of R₁, R₂ and R₃ is an electron-withdrawing moiety; and
- 25 b.) wherein X is an organic moiety having a molecular weight from about 104 Daltons to about 4871 Daltons, from about 236 Daltons to about 2371 Daltons from about 294 Daltons to about 871 Daltons.

In one aspect of the aforementioned composition, for said thiophene azo carboxylate dye having Formula I, each R_1 , R_2 and R_3 is independently selected from hydrogen, (C₁-C₄)-alkyl, (C₃-C₁₀)-aryl, carboxylate, cyano, sulfonate, phosphonate, sulfate, acetate, nitro, (C₁-C₄)-alkyl ester, halogen or amino moiety.

In one aspect of the aforementioned composition, for said thiophene azo carboxylate dye having Formula I, X is a moiety having Formula (II) below:




10

Formula II

wherein:

- i.) R_4 is selected from a moiety having Formula (III) below

15



Formula III


wherein:

- each R_8 is independently selected from hydrogen, C₁-C₈ alkyl optionally substituted with a hydroxy, or acetyl;
- m is an integer from 0 to 10;
- Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted

25

9


succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IV,



Formula IV

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

10 ii.) R_4 is selected from a moiety having Formula (V) below:




Formula V

wherein

15 each R_{11} and R_{12} is independently selected from hydrogen, C_1 - C_8 alkyl, aryl, acetyl or hydroxyl moiety; m and n are independent and are integers from 0 to 10,

20 Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VI,




25

Formula VI

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons; or

5

- iii.) R_4 is selected from a moiety having Formula (VII) below:



Formula VII

10

wherein


R_{13} is selected from a C_6 - C_{10} aryl moiety, C_7 - C_{12} arylalkyl moiety, C_1 - C_{18} alkyl moiety, or a siloxane moiety;

15

each R_{14} is independently selected from hydrogen, C_1 - C_4 alkyl; m is an integer from 0 to 10;

20

Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VIII,



Formula VIII

25

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons


to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

5 R_5 is an independently selected R_4 moiety as defined above or R_5 is selected from C_1 - C_{12} alkyl moiety, a C_2 - C_{12} ether moiety; C_6 - C_{10} aryl moiety or C_7 - C_{12} arylalkyl moiety; wherein the index a is an integer from 0 to 4, and each R_6 may be independently selected from a C_1 - C_6 alkyl, a C_1 - C_4 alkoxy, a nitro, a hydroxyl, a halogen, or $-NHC(O)R_{22}$ wherein R_{22} is selected from H, $-NH_2$, C_1 - C_6 alkyl, phenyl, -
10 $(CH_2)_sOR_{23}$ where the index s is 1 or 2 and R_{23} is selected from Me, phenyl, and $-CO_2CH_2CN$; $-NHSO_2R_{24}$ wherein R_{24} is C_1 - C_4 alkyl or phenyl; said alkyl, alkoxy and acetamido moieties may be optionally substituted with a carboxylate moiety;

15 or, X is a moiety having Formula II above:

wherein each R_4 and R_5 can independently be selected from:

- a) $[(CH_2CR'HO)_x(CH_2CR''HO)_yR_{15}]$;
- b) C_1 - C_{12} alkyl, C_6 - C_{10} aryl or C_7 - C_{12} aryl alkyl;
- c) $[CH_2CH(OR_{16})CH_2OR_{17}]$;
- 20 d) the amino addition product of styrene oxide, glycidyl methyl ether, isobutyl glycidyl ether, isopropylglycidyl ether, t-butyl glycidyl ether, 2-ethylhexylglycidyl ether, and glycidylhexadecyl ether, followed by the addition of from 1 to 10 alkylene oxide units wherein at least one such alkyleneoxide unit is substituted with R_{15} that is not $-H$;
- 25 wherein R' is selected from the group consisting of H, CH_3 , $CH_2O(CH_2CH_2O)_zR_{15}$, and mixtures thereof; R'' is selected from the group consisting of H, $CH_2O(CH_2CH_2O)_zR_{15}$, and mixtures thereof; $x + y \leq 20$; $y \geq 1$; $z = 0$ to 10; each R_{15} is independently selected from $-H$ and Y wherein Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IX,
- 30




Formula IX

5 said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

10 R_{16} is selected from the group consisting of H, $(CH_2CH_2O)_zR_{15}$ wherein $z = 0$ to 10, and mixtures thereof; R_{17} is selected from the group consisting of C_1 - C_{16} alkyl, C_6 - C_{10} aryl groups, and mixtures thereof; the index m is an integer from 0 to 4 and each R_6 is as defined above; further provided the molecule contains at least one non-H R_{15} group.


15 In one aspect of the aforementioned composition, for said thiophene azo carboxylate dye having Formula I X is a moiety having Formula (II) below:



Formula II

20 wherein:

- i.) R_4 is selected from a moiety having Formula (III) below




Formula III

25

wherein:

R₈ is a hydrogen, C₁-C₄ alkyl moiety or an C₆-C₁₀ aryl moiety;

Y is -OC(O)GCO₂M wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IV,




Formula IV

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

or, R₄ is selected from a moiety having Formula (V)

below:



Formula V


wherein

each R₁₁ and R₁₂ is independently selected from hydrogen, C₁-C₄ alkyl or C₆-C₁₀ aryl moiety; m and n are independent and are integers from 0 to 5,

Y is -OC(O)GCO₂M wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted

14


succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VI,



Formula VI

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

or, R_4 is selected from a moiety having Formula (VII) below:



Formula VII


wherein

R_{13} is selected from an C_6 - C_{10} aryl moiety, benzyl moiety, or a C_1 - C_{18} alkyl moiety;

each R_{14} is independently selected from hydrogen or $-CH_3$; m is an integer from 0 to 10,

Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VIII,

25



Formula VIII

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

5

ii.) R_5 is an independently selected R_4 moiety as defined above or R_5 is R_5 selected from C_1 - C_6 alkyl moiety or benzyl moiety;

10


iii.) wherein the index a is an integer from 0 to 2, and each R_6 may be independently selected from methyl, methoxy, or acetamido moiety.

In one aspect of the aforementioned composition, for said thiophene azo carboxylate dye having Formula I, each R_1 , R_2 and R_3 is independently selected from hydrogen, $(C_1$ - C_4)-alkyl, $(C_6$ - $C_{10})$ -aryl, carboxylate, cyano, sulfonate, acetate, nitro, $(C_1$ - C_4)-alkyl ester, halogen or amino moiety, in one aspect, each R_1 , R_2 and R_3 is independently selected from hydrogen, $(C_1$ - C_4)-alkyl, carboxylate, cyano, sulfonate, nitro, $(C_1$ - C_4)-alkyl ester or halogen moiety.

15

In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has Formula (X):

20




Formula X

wherein the A moiety is selected from the group consisting of Table 1 A Moieties Nos. 1 – 118 and wherein X is selected from the group consisting of Table 4 X Moieties Nos. 1 – 24 wherein G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula XI,

30


16



Formula XI


said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons.

In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has Formula (XII):



Formula XII


wherein the A moiety is selected from the group consisting of Table 1 A Moieties Nos. 9 – 11, 15, 23, 34 – 35, 37 – 39, 41, 43, 47, 50 – 51, 57 – 58, 77, 83, 89, 95, 106, and 110 – 118; and wherein the X moiety is selected from the group consisting of Table 4 X Moieties Nos. 1 – 8, 13, 14, and 18 – 21, wherein G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula XIII,



Formula XIII

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons. In one aspect of the aforementioned composition, said G moiety of said thiophene azo carboxylate dye is selected from Table 5 G Moieties Nos. 1 – 48.


In one aspect of the aforementioned composition, said thiophene azo carboxylate dye has the following formula:



wherein the moiety A is selected from Table 1 A Moieties Nos. 1 – 118; a = 0 to
 5 2; when a = 1 or 2, R₆ is selected from Table 2 R₆ Substituent Identity and
 Position Nos. 1 – 40; and R₄ and R₅ grouping is selected from Table 3 R₄ and R₅
 Groupings Nos. 1 – 83.

In one aspect of the aforementioned composition, said thiophene azo
 carboxylate dye has the following formula:

10





















wherein the moiety A is selected from Table 1 A Moieties Nos. 9 – 11, 15, 23, 34
 – 35, 37 – 39, 41, 43, 47, 50 – 51, 57 – 58, 77, 83, 89, 95, 106, and 110 – 118; a
 15 = 0 to 2; when a = 1 or 2, R₆ is selected from Table 2 R₆ Substituent Identity and
 Position Nos. 1, 3, 5, 7, 12, 13, 14, 31, 36 and 40; and R₄ and R₅ grouping is
 selected from Table 3 R₄ and R₅ Groupings Nos. 3 – 8, 14, 15, 18 – 23, 29, 30,
 33 – 38, 44, 45, 48 – 53, 59, 60, 63
 -66, 71-76, 82 and 83.



















20 In one aspect of the aforementioned composition, said thiophene azo
 carboxylate dye has a formula selected from Table 6 Formula Nos. 1 – 48, in
 one aspect, said thiophene azo carboxylate dye has a formula selected from
 Table 6 Formula Nos. 1 – 36, 43 and 44.





















25



















A moieties may be selected from the moieties shown in Table 1:

















Table 1: A Moieties























No.	A	No.	A
1		60	
2		61	
3		62	
4		63	
5		64	
6		65	
7		66	
8		67	
9		68	







10		69	
11		70	
12		71	
13		72	
14		73	
15		74	
16		75	
17		76	
18		77	

19		78	
20		79	
21		80	
22		81	
23		82	
24		83	
25		84	
26		85	
27		86	
28		87	

29		88	
30		89	
31		90	
32		91	
33		92	
34		93	
35		94	
36		95	
37		96	

38		97	
39		98	
40		99	
41		100	
42		101	
43		102	
44		103	
45		104	


46		105	
47		106	
48		107	
49		108	
50		109	
51		110	
52		111	
53		112	
54		113	
55		114	
56		115	

57		116	
58		117	
59		118	

R₆ may be selected from the substituents shown in Table 2:

5

Table 2: R₆ Substituent – Identity and Position



R ₆ Substituent – Identity and Position on Aniline Ring				
No.	1	2	3	4
1				-OH
2		-Cl		-OH
3		-OMe		-OH
4		-On-Bu		-OH
5				-Me
6		-Me		-Me
7		-OMe		-Me




8		-OEt		-Me
9				-Cl
10				-Br
11				-NO ₂
12				-NH ₂
13				-NHAc
14		-OMe		-NHAc
15				-NHC(O)Et
16		-OMe		-NHC(O)Et
17		-OEt		-NHC(O)Et
18				-NHC(O)n-Pr
19		-Me		-NHC(O)n-Pr
20				-NHC(O)i-Pr
21				-NHC(O)n-Bu
22		-OMe		-NHC(O)n-Bu
23				-NHC(O)c-C ₆ H ₁₁
24				-NHC(O)Ph
25		-OMe		-NHC(O)Ph
26				-NHC(O)CH ₂ OMe
27				-NHC(O)CH ₂ OPh
28				-NHC(O)CH ₂ CH ₂ OMe
29				-NHC(O)CH ₂ CH ₂ OEt
30				- NHC(O)(CH ₂) ₂ CO ₂ CH ₂ CN
31				-NHC(O)NH ₂
32				-NHSO ₂ Me

33				-NHSO ₂ Et
34		-OMe		-NHSO ₂ Et
35				-NHSO ₂ Ph
36		-OMe		
37		-Me		
38		-Cl		
39		-NO ₂		
40		-OMe		-OMe

R₄ and R₅ groupings may be selected from the groupings shown in Table 3:







5







Table 3: R₄ and R₅ Groupings








R ₄ and R ₅ Groupings		
N o.	R ₄	R ₅
1	Ethyl	
2	Ethyl	
3	Ethyl	







4	Ethyl	
5	Ethyl	
6	Ethyl	
7	Ethyl	
8	Ethyl	
9	Ethyl	








<p>10</p>	<p>Ethyl</p>	
<p>11</p>	<p>Ethyl</p>	
<p>12</p>	<p>Ethyl</p>	
<p>13</p>	<p>Ethyl</p>	
<p>14</p>	<p>Ethyl</p>	
<p>15</p>	<p>Ethyl</p>	







16	Benzyl	
17	Benzyl	
18	Benzyl	
19	Benzyl	
20	Benzyl	
21	Benzyl	







<p>22</p>	<p>Benzyl</p>	
<p>23</p>	<p>Benzyl</p>	
<p>24</p>	<p>Benzyl</p>	
<p>25</p>	<p>Benzyl</p>	
<p>26</p>	<p>Benzyl</p>	
<p>27</p>	<p>Benzyl</p>	












28	Benzyl	
29	Benzyl	
30	Benzyl	
31	Hexyl	
32	Hexyl	
33	Hexyl	
34	Hexyl	











<p>35</p>	<p>Hexyl</p>	
<p>36</p>	<p>Hexyl</p>	
<p>37</p>	<p>Hexyl</p>	
<p>38</p>	<p>Hexyl</p>	
<p>39</p>	<p>Hexyl</p>	
<p>40</p>	<p>Hexyl</p>	








41	Hexyl	
42	Hexyl	
43	Hexyl	
44	Hexyl	
45	Hexyl	
46	Isopropyl	
47	Isopropyl	





48	Isopropyl	
49	Isopropyl	
50	Isopropyl	
51	Isopropyl	
52	Isopropyl	
53	Isopropyl	

<p>54</p>	<p>Isopropyl</p>	
<p>55</p>	<p>Isopropyl</p>	
<p>56</p>	<p>Isopropyl</p>	
<p>57</p>	<p>Isopropyl</p>	
<p>58</p>	<p>Isopropyl</p>	
<p>59</p>	<p>Isopropyl</p>	

<p>60</p>	<p>Isopropyl</p>	 <p>Chemical structure showing an isopropyl ester group (MO₂C) attached to a long polyether chain consisting of four repeating units of -CH₂-CH₂-O-CH₂-CH₂-.</p>
<p>61</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M) and a polyether chain (-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring.</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M) and a shorter polyether chain (-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring, ending in an asterisk (*).</p>
<p>62</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M), a long alkyl chain, and a polyether chain (-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring.</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M), a long alkyl chain, and a shorter polyether chain (-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring, ending in an asterisk (*).</p>
<p>63</p>	 <p>Chemical structure showing a benzene ring with a methyl ester group (CO₂M) and a polyether chain (-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring.</p>	 <p>Chemical structure showing a benzene ring with a methyl ester group (CO₂M) and a shorter polyether chain (-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring, ending in an asterisk (*).</p>
<p>64</p>	 <p>Chemical structure showing a cyclohexene ring with a methyl ester group (CO₂M) and a polyether chain (-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring.</p>	 <p>Chemical structure showing a cyclohexene ring with a methyl ester group (CO₂M) and a shorter polyether chain (-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring, ending in an asterisk (*).</p>
<p>65</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M) and a polyether chain (-CH₂-CH₂-O-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring.</p>	 <p>Chemical structure showing a cyclohexane ring with a methyl ester group (CO₂M) and a shorter polyether chain (-CH₂-CH₂-O-CH₂-CH₂-) attached to the ring, ending in an asterisk (*).</p>

<p>66</p>		
<p>67</p>		
<p>68</p>		
<p>69</p>	<p>Methoxyethyl</p>	
<p>70</p>	<p>Methoxyethyl</p>	
<p>71</p>	<p>Methoxyethyl</p>	
<p>72</p>	<p>Methoxyethyl</p>	

73	Methoxyethyl	
74	Methoxyethyl	
75	Methoxyethyl	
76	Methoxyethyl	
77	Methoxyethyl	
78	Methoxyethyl	
79	Methoxyethyl	


80	Methoxyethyl	
81	Methoxyethyl	
82	Methoxyethyl	
83	Methoxyethyl	







The “*” denotes point of attachment to additional moieties.
 “M” denotes H or the appropriate charge balancing counterion.







5






X moieties may be selected from the moieties shown in Table 4:






Table 4: X Moieties


No.	G moieties may be selected from moieties 1-48 shown in Table 5.
1	

<p>2</p>	
<p>3</p>	
<p>4</p>	
<p>5</p>	
<p>6</p>	
<p>7</p>	

<p>8</p>	
<p>9</p>	
<p>10</p>	
<p>11</p>	
<p>12</p>	
<p>13</p>	

<p>14</p>	
<p>15</p>	
<p>16</p>	
<p>17</p>	
<p>18</p>	

<p>19</p>	
<p>20</p>	
<p>21</p>	
<p>22</p>	
<p>23</p>	



























The “*” denotes point of attachment to the azo moiety.













G moieties in the X moieties in Table 4 above may be selected from moieties 1-48 shown in Table 5:

5

Table 5: G Moieties

No.	G moiety	No.	G moiety
1		25	
2		26	
3		27	
4		28	
5		29	
6		30	

7		31	
8		32	
9		33	
10		34	
11		35	
12		36	
13		37	
14		38	
15		39	
16		40	
17		41	
18		42	

19		43	
20		44	
21		45	
22		46	
23		47	
24		48	




Anhydrides suitable for use in creating the Y moiety in the instant invention include, but are not limited to, the following: 3-(2E)-2-dodecen-1-yl-dihydro-2,5-Furandione; (3aR,7aS)-rel-3a,4,7,7a-tetrahydro-4,7-Epoxyisobenzofuran-1,3-dione; 3-hexyl-dihydro-3-methyl-2,5-Furandione; 3-hexyl-dihydro-2,5-Furandione; (3aR,8aS)-rel-hexahydro-1H-Cyclohepta[c]furan-1,3(3aH)-dione; (3aR,6aS)-rel-tetrahydro-1H-Cyclopenta[c]furan-1,3(3aH)-dione; 5-(1,1-dimethylethyl)-1,3-Isobenzofurandione; dihydro-3-(phenylmethyl)-2,5-Furandione; 3-decyl-dihydro-2,5-Furandione; dihydro-3,3-dimethyl-2,5-Furandione; (3aR,4S,7R,7aS)-rel-hexahydro-4,7-Methanoisobenzofuran-1,3-dione; 3-ethyl-dihydro-2,5-Furandione; (3aR,7aS)-rel-3a,4,7,7a-tetrahydro-5,6-dimethyl-1,3-Isobenzofurandione; 3-(2-hexen-1-yl)-dihydro-2,5-Furandione; dihydro-3-(2-propen-1-yl)-2,5-Furandione; hexahydro-4,7-Methanoisobenzofuran-1,3-dione; 5,6-dimethyl-1,3-Isobenzofurandione; tetrahydro-1H-Cyclopenta[c]furan-1,3(3aH)-dione; 5-nitro-







1,3-Isobenzofurandione; hexahydro-4,7-Epoxyisobenzofuran-1,3-dione;
3a,4,7,7a-tetrahydro-4,7-Epoxyisobenzofuran-1,3-dione; 5,6-dimethoxy-1,3-
Isobenzofurandione; dihydro-3-octyl-2,5-Furandione; dihydro-3-methyl-2,5-
Furandione; 1,3-dihydro-1,3-dioxo-5-Isobenzofurancarboxylic acid, methyl
5 ester; 3-dodecyldihydro-2,5-Furandione; dihydro-3-phenyl-2,5-Furandione;
3a,4,7,7a-tetrahydro-4,7-Methanoisobenzofuran-1,3-dione; 4-nitro-1,3-
Isobenzofurandione; 1,3-dihydro-1,3-dioxo-5-Isobenzofurancarboxylic acid;
(3aR,4S,7R,7aS)-rel-3a,4,7,7a-tetrahydro-4,7-Methanoisobenzofuran-1,3-dione;
4,5,6,7-tetrachloro-1,3-Isobenzofurandione; dihydro-2,5-Furandione; 1,3-
10 Isobenzofurandione; 3a,4,7,7a-tetrahydro-1,3-Isobenzofurandione; hexahydro-
1,3-Isobenzofurandione; (3aR,4S,7R,7aS)-rel-hexahydro-3a,7a-dimethyl-4,7-
Epoxyisobenzofuran-1,3-dione; 4-hexyldihydro-2H-Pyran-2,6(3H)-dione; dihydro-
4-(2-methylpropyl)-2H-Pyran-2,6(3H)-dione; 4-[[[(1,1-
dimethylethyl)dimethylsilyl]oxy]dihydro-2H-Pyran-2,6(3H)-dione; 4,4-dimethyl-
15 1H-2-Benzopyran-1,3(4H)-dione; 4-ethyldihydro-4-methyl-2H-Pyran-2,6(3H)-
dione; 8-Oxaspiro[4.5]decane-7,9-dione; dihydro-4,4-dimethyl-2H-Pyran-2,6(3H)-
dione; dihydro-3-phenyl-2H-Pyran-2,6(3H)-dione; dihydro-3,3-dimethyl-2H-
Pyran-2,6(3H)-dione; 1H-2-Benzopyran-1,3(4H)-dione; dihydro-2H-Pyran-
2,6(3H)-dione; 1H,3H-Naphtho[1,8-cd]pyran-1,3-dione; 3-
20 Oxabicyclo[3.2.1]octane-2,4-dione; Dibenz[c,e]oxepin-5,7-dione; 4,5-dihydro-2-
Benzoxepin-1,3-dione; and 2,7-Oxepanedione. Such suitable anhydrides may
be obtained from one or more of the following: 3B Scientific Corporation,
Libertyville, IL, 60048; Accel Pharmtech, LLC, East Brunswick, NJ, 08816; A
Chemtek, Worcester, MA, 01606; AK Scientific, Inc., Union City, CA, 94587;
25 AKos Consulting and Solutions Deutschland GmbH, Steinen, D-79585, Germany;
Alfa Aesar, Ward Hill, MA, 01835; Allichem LLC, Baltimore, MD, 21224;
American Custom Chemicals Corp., San Diego, CA, 92196-2527; Aurora Fine
Chemicals LLC, San Diego, CA, 92126; Beta Pharma Scientific, Inc., Branford,
CT, 06405; City Chemical LLC, West Haven, CT, 06516; Ryan Scientific, Inc.,
30 Mt. Pleasant, SC, 29465; Sigma-Aldrich, St. Louis, MO, 63178; TCI America,
Portland, OR, 97203; Waterstone Technology, Carmel, IN, 46032; and
Youchemicals Limited, Minhang, Shanghai, 201100, People's Republic of China.






It is contemplated to be within the scope of this invention that the thiophene azo carboxylate dye may comprise any one of the A moieties selected from Table 1,
 5 any one of the R₆ substituents selected from Table 2, any one of the R₄ and R₅ groupings selected from Table 3, any one of the X moieties selected from Table 4, and any one of the G moieties selected from Table 5.






In yet another aspect of the invention, suitable thiophene azo carboxylate dyes
 10 include, but are not limited to, the structures shown in Table 6:





Table 6: Thiophene Azo Carboxylate Dyes





No.	Formula
Example 1	
Example 2	
Example 3	



Example 4	
Example 5	
Example 6	
Example 7	
Example 8	
Example 9	

Example 10	
Example 11	
Example 12	
Example 13	
Example 14	

Example 15	
Example 16	
Example 17	
Example 18	
Example 19	






Example 20	
Example 21	
Example 22	
Example 23	

Example 24	
Example 25	
Example 26	
Example 27	


Example 28	 <p>The chemical structure of Example 28 consists of a cyclohexane ring substituted with a methyl ester group (-CO₂M) and a side chain. The side chain is a polyoxyethylene chain (-(CH₂)₂O)₄-(CH₂)₂-) connected to a nitrogen atom. This nitrogen atom is also bonded to a pentyl group and a 4-methylphenyl ring. The 4-methylphenyl ring is further substituted with a diazo group (-N=N-) which is attached to a 2,4-dicyano-5-methylthiophene ring.</p>
Example 29	 <p>The chemical structure of Example 29 is similar to Example 28, but the cyclohexane ring is replaced by a long, zigzag alkyl chain. The rest of the molecule, including the polyoxyethylene chain, the nitrogen atom with its pentyl and 4-methylphenyl substituents, and the diazo-linked 2,4-dicyano-5-methylthiophene group, remains identical to Example 28.</p>

<p>Example 30</p>	
<p>Example 31</p>	
<p>Example 32</p>	
<p>Example 33</p>	

<p>Example 34</p>	
<p>Example 35</p>	
<p>Example 36</p>	
<p>Example 37</p>	
<p>Example 38</p>	

<p>Example 39</p>	
<p>Example 40</p>	
<p>Example 41</p>	
<p>Example 42</p>	
<p>Example 43</p>	

<p>Example 44</p>	
<p>Example 45</p>	
<p>Example 46</p>	
<p>Example 47</p>	



As indicated previously, the hueing agents described in the present specification may be incorporated into laundry care compositions including but not limited to laundry detergents and fabric care compositions. The laundry care compositions including laundry detergents may be in solid or liquid form, including a gel form, and/or unit dose forms, including multi-compartment unit dose forms. Such compositions may comprise one or more of said hueing agents and a laundry care ingredient.

- 5
- 10 In one aspect, said laundry care composition may comprise, based on total laundry care composition weight, less than 15% builder, less than 10% builder, or even less than 5% builder.

- 15 In one aspect, said laundry care composition may comprise, based on total laundry care composition weight, a total of no more than 20% water; a total of no more than 15% water; a total of no more than 10% water; or even a total of no more than 5% water.

- 20 In one aspect, said laundry care composition may comprise, based on total laundry care composition weight, from about 10% to about 70% of a water-miscible organic solvent having a molecular weight of greater than 70 Daltons.

- 25 In one aspect, said laundry care composition may comprise, based on total laundry care composition weight, a perfume microcapsule comprising a core and a shell that encapsulates said core, said perfume microcapsule having a D[4,3]

average particle of from about 0.01 microns to about 200 microns and optionally a formaldehyde scavenger that is supplied via the addition of the microcapsules (contained in a perfume microcapsule slurry that is added to the laundry care ingredient) and/or added directly to the laundry care composition. In one aspect, 5 the shell of said perfume microcapsules may be made of any material, including materials selected from the group consisting of polyethylenes, polyamides, polystyrenes, polyisoprenes, polycarbonates, polyesters, polyacrylates, polyureas, polyurethanes, polyolefins, polysaccharides, epoxy resins, vinyl polymers, and mixtures thereof. In one aspect, useful shell materials include 10 materials that are sufficiently impervious to the core material and the materials in the environment in which the perfume microcapsule will be employed, to permit the delivery perfume to be obtained. Suitable impervious shell materials include materials selected from the group consisting of reaction products of one or more amines with one or more aldehydes, such as urea cross-linked with 15 formaldehyde or gluteraldehyde, melamine cross-linked with formaldehyde; gelatin-polyphosphate coacervates optionally cross-linked with gluteraldehyde; gelatin-gum Arabic coacervates; cross-linked silicone fluids; polyamine reacted with polyisocyanates and mixtures thereof. In one aspect, the shell material comprises melamine cross-linked with formaldehyde and/or a polyacrylate. 20 Suitable perfume microcapsules may be obtained from Appleton Papers of Appleton Wisconsin, USA.

In one aspect, suitable formaldehyde scavengers include materials selected from the group consisting of sodium bisulfite, urea, ethylene urea, cysteine, 25 cysteamine, lysine, glycine, serine, carnosine, histidine, glutathione, 3,4-diaminobenzoic acid, allantoin, glycouril, anthranilic acid, methyl anthranilate, methyl 4-aminobenzoate, ethyl acetoacetate, acetoacetamide, malonamide, ascorbic acid, 1,3-dihydroxyacetone dimer, biuret, oxamide, benzoguanamine, pyroglutamic acid, pyrogallol, methyl gallate, ethyl gallate, propyl gallate, 30 triethanol amine, succinamide, thiabendazole, benzotriazol, triazole, indoline, sulfanilic acid, oxamide, sorbitol, glucose, cellulose, poly(vinyl alcohol), partially hydrolyzed poly(vinylformamide), poly(vinyl amine), poly(ethylene imine),

poly(oxyalkyleneamine), poly(vinyl alcohol)-co-poly(vinyl amine), poly(4-aminostyrene), poly(L-lysine), chitosan, hexane diol, ethylenediamine-N,N'-bisacetoacetamide, N-(2-ethylhexyl)acetoacetamide, 2-benzoylacetoacetamide, N-(3-phenylpropyl)acetoacetamide, lillial, helional, melonal, triplal, 5,5-dimethyl-1,3-cyclohexanedione, 2,4-dimethyl-3-cyclohexenecarboxaldehyde, 2,2-dimethyl-1,3-dioxan-4,6-dione, 2-pentanone, dibutyl amine, triethylenetetramine, ammonium hydroxide, benzylamine, hydroxycitronellol, cyclohexanone, 2-butanone, pentane dione, dehydroacetic acid, or a mixture thereof. These formaldehyde scavengers may be obtained from Sigma/Aldrich/Fluka of St. Louis, MO. U.S.A. or PolySciences, Inc. of Warrington, PA U.S.A.

Such formaldehyde scavengers are typically combined with a slurry containing said perfume microcapsules, at a level, based on total slurry weight, of from about 2 wt.% to about 18 wt.%, from about 3.5 wt.% to about 14 wt.% or even from about 5 wt.% to about 13 wt.%.

In one aspect, such formaldehyde scavengers may be combined with a product containing a perfume microcapsule, said scavengers being combined with said product at a level, based on total product weight, of from about 0.005% to about 0.8%, alternatively from about 0.03% to about 0.5%, alternatively from about 0.065% to about 0.25% of the product formulation.

In another aspect, such formaldehyde scavengers may be combined with a slurry containing said perfume microcapsules, at a level, based on total slurry weight, of from about 2 wt.% to about 14 wt.%, from about 3.5 wt.% to about 14 wt.% or even from about 5 wt.% to about 14 wt.% and said slurry may be added to a product matrix to which addition an identical or different scavenger may be added at a level, based on total product weight, of from about 0.005% to about 0.5%, alternatively from about 0.01% to about 0.25%, alternatively from about 0.05% to about 0.15% of the product formulation.

In one aspect, one or more of the aforementioned formaldehyde scavengers may be combined with a liquid fabric enhancing product containing perfume microcapsules at a level, based on total liquid fabric enhancing product weight, of from 0.005% to about 0.8%, alternatively from about 0.03% to about 0.4%,
5 alternatively from about 0.06% to about 0.25% of the product formulation.

In one aspect, such formaldehyde scavengers may be combined with a liquid laundry detergent product containing perfume microcapsules, said scavengers being selected from the group consisting of sodium bisulfite, urea, ethylene urea,
10 cysteine, cysteamine, lysine, glycine, serine, carnosine, histidine, glutathione, 3,4-diaminobenzoic acid, allantoin, glycouril, anthranilic acid, methyl anthranilate, methyl 4-aminobenzoate, ethyl acetoacetate, acetoacetamide, malonamide, ascorbic acid, 1,3-dihydroxyacetone dimer, biuret, oxamide, benzoguanamine, pyroglutamic acid, pyrogallol, methyl gallate, ethyl gallate, propyl gallate,
15 triethanol amine, succinamide, thiabendazole, benzotriazol, triazole, indoline, sulfanilic acid, oxamide, sorbitol, glucose, cellulose, poly(vinyl alcohol), partially hydrolyzed poly(vinylformamide), poly(vinyl amine), poly(ethylene imine), poly(oxyalkyleneamine), poly(vinyl alcohol)-co-poly(vinyl amine), poly(4-aminostyrene), poly(l-lysine), chitosan, hexane diol, ethylenediamine-N,N'-
20 bisacetoacetamide, N-(2-ethylhexyl)acetoacetamide, 2-benzoylacetoacetamide, N-(3-phenylpropyl)acetoacetamide, lillial, helional, melonal, triplal, 5,5-dimethyl-1,3-cyclohexanedione, 2,4-dimethyl-3-cyclohexenecarboxaldehyde, 2,2-dimethyl-1,3-dioxan-4,6-dione, 2-pentanone, dibutyl amine, triethylenetetramine, ammonium hydroxide, benzylamine, hydroxycitronellol, cyclohexanone, 2-
25 butanone, pentane dione, dehydroacetic acid and mixtures thereof, and combined with said liquid laundry detergent product at a level, based on total liquid laundry detergent product weight, of from about 0.003 wt.% to about 0.20 wt.%, from about 0.03 wt.% to about 0.20 wt.% or even from about 0.06 wt.% to about 0.14 wt.%.

30

The hueing agents may be added to substrates using a variety of application techniques. For instance, for application to cellulose-containing textile

substrates, the hueing agent may be included as a component of a laundry detergent. Thus, application to a cellulose-containing textile substrate actually occurs when a consumer adds laundry detergent to a washing machine. The hueing agent may be present in the laundry detergent composition in an amount
5 from about 0.000001% to about 10% by weight of the composition, from about 0.00001% to about 10% by weight of the composition, from about 0.0001% to about 5% by weight of the composition, and even from about 0.0001% to about 1% by weight of the composition.

10 The laundry detergent composition typically comprises a surfactant in an amount sufficient to provide desired cleaning properties. In one aspect, the laundry detergent composition may comprise, based on total laundry detergent composition weight, from about 0.5% to about 99% of the surfactant; from about 1% to about 95% of the surfactant; from about 5% to about 90% of the
15 surfactant, from about 5% to about 70% of the surfactant, or even from about 5% to about 40% of the surfactant. The surfactant may comprise anionic, nonionic, cationic, zwitterionic and/or amphoteric surfactants. In one aspect, the detergent composition comprises anionic surfactant, nonionic surfactant, or mixtures thereof.

20 Fabric care compositions are typically added in the rinse cycle, which is after the detergent solution has been used and replaced with the rinsing solution in typical laundering processes. The fabric care compositions disclosed herein may be comprise a rinse added fabric softening active and a suitable hueing agent as
25 disclosed in the present specification. The fabric care composition may comprise, based on total fabric care composition weight, from about 1% to about 90%, or from about 5% to about 50% fabric softening active. The hueing agent may be present in the fabric care composition in an amount from about 0.5 ppb to about 50 ppm, or from about 0.5 ppm to about 30 ppm.

30

Suitable Laundry Care Ingredients

While not essential for the purposes of the present invention, the non-limiting list of laundry care ingredients illustrated hereinafter are suitable for use in the laundry care compositions and may be desirably incorporated in certain aspects of the invention, for example to assist or enhance performance, for treatment of the substrate to be cleaned, or to modify the aesthetics of the composition as is the case with perfumes, colorants, dyes or the like. It is understood that such ingredients are in addition to the components that were previously listed for any particular aspect. The total amount of such adjuncts may range, once the amount of dye is taken into consideration from about 90% to about 99.99999995% by weight of the laundry care composition.

The precise nature of these additional components, and levels of incorporation thereof, will depend on the physical form of the composition and the nature of the operation for which it is to be used. Suitable laundry care ingredients include, but are not limited to, fabric softening actives, polymers, for example cationic polymers, surfactants, builders, chelating agents, dye transfer inhibiting agents, dispersants, enzymes, and enzyme stabilizers, catalytic materials, bleach activators, polymeric dispersing agents, clay soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, perfume(s) including quadrant perfumes and additional perfume delivery systems including perfume loaded zeolites, starch encapsuled accords, and Schiff base pro-perfumes, structure elasticizing agents, fabric softeners, carriers, hydrotropes, processing aids and/or pigments. In addition to the disclosure below, suitable examples of such other adjuncts and levels of use are found in U.S. Patent Nos. 5,576,282, 6,306,812 B1 and 6,326,348 B1 that are incorporated by reference.

As stated, the laundry care ingredients are not essential to Applicants' laundry care compositions. Thus, certain aspects of Applicants' compositions do not contain one or more of the following adjuncts materials: fabric softening actives, bleach activators, surfactants, builders, chelating agents, dye transfer inhibiting

agents, dispersants, enzymes, and enzyme stabilizers, catalytic metal complexes, polymeric dispersing agents, clay and soil removal/anti-redeposition agents, brighteners, suds suppressors, dyes, additional perfumes and perfume delivery systems, structure elasticizing agents, fabric softeners, carriers, 5 hydrotropes, processing aids and/or pigments. However, when one or more adjuncts are present, such one or more adjuncts may be present as detailed below:

Surfactants

- 10 Suitable anionic surfactants useful herein can comprise any of the conventional anionic surfactant types typically used in liquid detergent products. These include the alkyl benzene sulfonic acids and their salts as well as alkoxyated or non-alkoxyated alkyl sulfate materials.
- 15 Exemplary anionic surfactants are the alkali metal salts of C₁₀-C₁₆ alkyl benzene sulfonic acids, or C₁₁-C₁₄ alkyl benzene sulfonic acids. In one aspect, the alkyl group is linear and such linear alkyl benzene sulfonates are known as "LAS". Alkyl benzene sulfonates, and particularly LAS, are well known in the art. Such surfactants and their preparation are described for example in U.S. Pat. Nos. 20 2,220,099 and 2,477,383. Especially useful are the sodium and potassium linear straight chain alkylbenzene sulfonates in which the average number of carbon atoms in the alkyl group is from about 11 to 14. Sodium C₁₁-C₁₄, e.g., C₁₂, LAS is a specific example of such surfactants.
- 25 Another exemplary type of anionic surfactant comprises ethoxylated alkyl sulfate surfactants. Such materials, also known as alkyl ether sulfates or alkyl polyethoxylate sulfates, are those which correspond to the formula: R'-O-(C₂H₄O)_n-SO₃M wherein R' is a C₈-C₂₀ alkyl group, n is from about 1 to 20, and M is a salt-forming cation. In one aspect, R' is C₁₀-C₁₈ alkyl, n is from about 1 to 30 15, and M is sodium, potassium, ammonium, alkylammonium, or alkanolammonium. In one aspect, R' is a C₁₂-C₁₆, n is from about 1 to 6 and M is sodium.

The alkyl ether sulfates will generally be used in the form of mixtures comprising varying R' chain lengths and varying degrees of ethoxylation. Frequently such mixtures will inevitably also contain some non-ethoxylated alkyl sulfate materials, 5 i.e., surfactants of the above ethoxylated alkyl sulfate formula wherein $n=0$. Non-ethoxylated alkyl sulfates may also be added separately to the compositions of this invention and used as or in any anionic surfactant component which may be present. Specific examples of non-alkoxylated, e.g., non-ethoxylated, alkyl ether sulfate surfactants are those produced by the sulfation of higher C_8 - C_{20} fatty 10 alcohols. Conventional primary alkyl sulfate surfactants have the general formula: $ROSO_3-M^+$ wherein R is typically a linear C_8 - C_{20} hydrocarbyl group, which may be straight chain or branched chain, and M is a water-solubilizing cation. In one aspect, R is a C_{10} - C_{15} alkyl, and M is alkali metal, more specifically R is C_{12} - C_{14} and M is sodium.

15

Specific, non-limiting examples of anionic surfactants useful herein include: a) C_{11} - C_{18} alkyl benzene sulfonates (LAS); b) C_{10} - C_{20} primary, branched-chain and random alkyl sulfates (AS); c) C_{10} - C_{18} secondary (2,3) alkyl sulfates having formulae (I) and (II): wherein M in formulae (I) and (II) is hydrogen or a cation 20 which provides charge neutrality, and all M units, whether associated with a surfactant or adjunct ingredient, can either be a hydrogen atom or a cation depending upon the form isolated by the artisan or the relative pH of the system wherein the compound is used, with non-limiting examples of suitable cations including sodium, potassium, ammonium, and mixtures thereof, and x is an 25 integer of at least about 7, or at least about 9, and y is an integer of at least 8, or at least about 9; d) C_{10} - C_{18} alkyl alkoxy sulfates (AE_xS) wherein x is from 1-30; e) C_{10} - C_{18} alkyl alkoxy carboxylates in one aspect, comprising 1-5 ethoxy units; f) mid-chain branched alkyl sulfates as discussed in U.S. Pat. No. 6,020,303 and U.S. Pat. No. 6,060,443; g) mid-chain branched alkyl alkoxy sulfates as 30 discussed in U.S. Pat. No. 6,008,181 and U.S. Pat. No. 6,020,303; h) modified alkylbenzene sulfonate (MLAS) as discussed in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO

00/23549, and WO 00/23548; i) methyl ester sulfonate (MES); and j) alpha-olefin sulfonate (AOS).

5 Suitable nonionic surfactants useful herein can comprise any of the conventional nonionic surfactant types typically used in liquid detergent products. These include alkoxyated fatty alcohols and amine oxide surfactants. In one aspect, for use in the liquid detergent products herein are those nonionic surfactants which are normally liquid.

10 Suitable nonionic surfactants for use herein include the alcohol alkoxyate nonionic surfactants. Alcohol alkoxyates are materials which correspond to the general formula: $R^1(C_mH_{2m}O)_nOH$ wherein R^1 is a C_8 - C_{16} alkyl group, m is from 2 to 4, and n ranges from about 2 to 12. In one aspect, R^1 is an alkyl group, which may be primary or secondary, that comprises from about 9 to 15 carbon atoms,
15 or from about 10 to 14 carbon atoms. In one aspect, the alkoxyated fatty alcohols will also be ethoxylated materials that contain from about 2 to 12 ethylene oxide moieties per molecule, or from about 3 to 10 ethylene oxide moieties per molecule.

20 The alkoxyated fatty alcohol materials useful in the liquid detergent compositions herein will frequently have a hydrophilic-lipophilic balance (HLB) which ranges from about 3 to 17 from about 6 to 15, or from about 8 to 15. Alkoxyated fatty alcohol nonionic surfactants have been marketed under the tradenames Neodol and Dobanol by the Shell Chemical Company.

25

Another suitable type of nonionic surfactant useful herein comprises the amine oxide surfactants. Amine oxides are materials which are often referred to in the art as "semi-polar" nonionics. Amine oxides have the formula:
30 $R(EO)_x(PO)_y(BO)_zN(O)(CH_2R')_2 \cdot qH_2O$. In this formula, R is a relatively long-chain hydrocarbonyl moiety which can be saturated or unsaturated, linear or branched, and can contain from 8 to 20, 10 to 16 carbon atoms, or is a C_{12} - C_{16} primary alkyl. R' is a short-chain moiety, in one aspect R' may be selected from

hydrogen, methyl and $-\text{CH}_2\text{OH}$. When $x+y+z$ is different from 0, EO is ethyleneoxy, PO is propyleneoxy and BO is butyleneoxy. Amine oxide surfactants are illustrated by C_{12-14} alkyldimethyl amine oxide.

- 5 Non-limiting examples of nonionic surfactants include: a) C_{12-18} alkyl ethoxylates, such as, NEODOL® nonionic surfactants from Shell; b) $\text{C}_6\text{-C}_{12}$ alkyl phenol alkoxyates wherein the alkoxyate units are a mixture of ethyleneoxy and propyleneoxy units; c) C_{12-18} alcohol and $\text{C}_6\text{-C}_{12}$ alkyl phenol condensates with ethylene oxide/propylene oxide block polymers such as Pluronic® from BASF; d)
- 10 $\text{C}_{14}\text{-C}_{22}$ mid-chain branched alcohols, BA, as discussed in U.S. Pat. No. 6,150,322; e) $\text{C}_{14}\text{-C}_{22}$ mid-chain branched alkyl alkoxyates, BAE_x , wherein x if from 1-30, as discussed in U.S. Pat. No. 6,153,577, U.S. Pat. No. 6,020,303 and U.S. Pat. No. 6,093,856; f) Alkylpolysaccharides as discussed in U.S. Pat. No. 4,565,647 to Llenado, issued Jan. 26, 1986; specifically alkylpolyglycosides as
- 15 discussed in U.S. Pat. No. 4,483,780 and U.S. Pat. No. 4,483,779; g) Polyhydroxy fatty acid amides as discussed in U.S. Pat. No. 5,332,528, WO 92/06162, WO 93/19146, WO 93/19038, and WO 94/09099; and h) ether capped poly(oxyalkylated) alcohol surfactants as discussed in U.S. Pat. No. 6,482,994 and
- 20 WO 01/42408.

In the laundry detergent compositions herein, the deterative surfactant component may comprise combinations of anionic and nonionic surfactant materials. When this is the case, the weight ratio of anionic to nonionic will typically range from 10:90 to 90:10, more typically from 30:70 to 70:30.

- 25 Cationic surfactants are well known in the art and non-limiting examples of these include quaternary ammonium surfactants, which can have up to 26 carbon atoms. Additional examples include a) alkoxyate quaternary ammonium (AQA) surfactants as discussed in U.S. Pat. No. 6,136,769; b) dimethyl hydroxyethyl quaternary ammonium as discussed in U.S. Pat. No. 6,004,922; c) polyamine cationic surfactants as discussed in WO 98/35002, WO 98/35003, WO 98/35004, WO 98/35005, and WO 98/35006; d) cationic ester surfactants as
- 30

discussed in U.S. Pat. Nos. 4,228,042, 4,239,660 4,260,529 and U.S. Pat. No. 6,022,844; and e) amino surfactants as discussed in U.S. Pat. No. 6,221,825 and WO 00/47708, specifically amido propyldimethyl amine (APA).

5 Non-limiting examples of zwitterionic surfactants include derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See U.S. Pat. No. 3,929,678 to Laughlin et al.,
10 issued Dec. 30, 1975 at column 19, line 38 through column 22, line 48, for examples of zwitterionic surfactants; betaine, including alkyl dimethyl betaine and cocodimethyl amidopropyl betaine, C₈ to C₁₈ (in one aspect C₁₂ to C₁₈) amine oxides and sulfo and hydroxy betaines, such as N-alkyl-N,N-dimethylamino-1-propane sulfonate where the alkyl group can be C₈ to C₁₈, or C₁₀ to C₁₄.

15

Non-limiting examples of ampholytic surfactants include aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents comprises at least about 8 carbon
20 atoms, typically from about 8 to about 18 carbon atoms, and at least one comprises an anionic water-solubilizing group, e.g. carboxy, sulfonate, sulfate. See U.S. Pat. No. 3,929,678 to Laughlin et al., issued Dec. 30, 1975 at column 19, lines 18-35, for examples of ampholytic surfactants.

25 Aqueous, non-surface active liquid carrier

As noted, the laundry care compositions may be in the form of a solid, either in tablet or particulate form, including, but not limited to particles, flakes, sheets, or the like, or the compositions may be in the form of a liquid. The liquid detergent compositions may comprise an aqueous, non-surface active liquid carrier.
30 Generally, the amount of the aqueous, non-surface active liquid carrier employed in the compositions herein will be effective to solubilize, suspend or disperse the composition components. For example, the liquid detergent compositions may

comprise, based on total liquid detergent composition weight, from about 5% to about 90%, from about 10% to about 70%, or from about 20% to about 70% of the aqueous, non-surface active liquid carrier.

- 5 The most cost effective type of aqueous, non-surface active liquid carrier is typically water. Accordingly, the aqueous, non-surface active liquid carrier component will generally be mostly, if not completely, comprised of water. While other types of water-miscible liquids, such alkanols, diols, other polyols, ethers, amines, and the like, have been conventionally been added to liquid detergent
- 10 compositions as co-solvents or stabilizers, for purposes of the present invention, the utilization of such water-miscible liquids typically is minimized to hold down composition cost. Accordingly, the aqueous liquid carrier component of the liquid detergent products herein will generally comprise water present in concentrations ranging from about 5% to about 90%, or from about 5% to about
- 15 70%, by weight of the liquid detergent composition.

Bleaching Agents

- Bleaching Agents – The cleaning compositions of the present invention may comprise one or more bleaching agents. Suitable bleaching agents other than
- 20 bleaching catalysts include photobleaches, bleach activators, hydrogen peroxide, sources of hydrogen peroxide, pre-formed peracids and mixtures thereof. In general, when a bleaching agent is used, the compositions of the present invention may comprise from about 0.1% to about 50% or even from about 0.1% to about 25% bleaching agent by weight of the subject cleaning
- 25 composition. Examples of suitable bleaching agents include:
- (1) photobleaches for example sulfonated zinc phthalocyanine;
 - (2) preformed peracids: Suitable preformed peracids include, but are not limited to, compounds selected from the group consisting of percarboxylic acids and salts, percarbonic acids and salts, perimidic acids and salts,
- 30 peroxymonosulfuric acids and salts, for example, Oxzone ®, and mixtures thereof. Suitable percarboxylic acids include hydrophobic and hydrophilic peracids having the formula $R-(C=O)O-O-M$ wherein R is an alkyl group,

optionally branched, having, when the peracid is hydrophobic, from 6 to 14 carbon atoms, or from 8 to 12 carbon atoms and, when the peracid is hydrophilic, less than 6 carbon atoms or even less than 4 carbon atoms; and M is a counterion, for example, sodium, potassium or hydrogen;

5 (3) sources of hydrogen peroxide, for example, inorganic perhydrate salts, including alkali metal salts such as sodium salts of perborate (usually mono- or tetra-hydrate), percarbonate, persulphate, perphosphate, persilicate salts and mixtures thereof. In one aspect of the invention the inorganic perhydrate salts are selected from the group consisting of sodium salts of
10 perborate, percarbonate and mixtures thereof. When employed, inorganic perhydrate salts are typically present in amounts of from 0.05 to 40 wt%, or 1 to 30 wt% of the overall composition and are typically incorporated into such compositions as a crystalline solid that may be coated. Suitable coatings include, inorganic salts such as alkali metal silicate, carbonate or borate salts or
15 mixtures thereof, or organic materials such as water-soluble or dispersible polymers, waxes, oils or fatty soaps; and

(4) bleach activators having R-(C=O)-L wherein R is an alkyl group, optionally branched, having, when the bleach activator is hydrophobic, from 6 to 14 carbon atoms, or from 8 to 12 carbon atoms and, when the bleach activator is
20 hydrophilic, less than 6 carbon atoms or even less than 4 carbon atoms; and L is leaving group. Examples of suitable leaving groups are benzoic acid and derivatives thereof - especially benzene sulphonate. Suitable bleach activators include dodecanoyl oxybenzene sulphonate, decanoyl oxybenzene sulphonate, decanoyl oxybenzoic acid or salts thereof, 3,5,5-trimethyl hexanoyloxybenzene
25 sulphonate, tetraacetyl ethylene diamine (TAED) and nonanoyloxybenzene sulphonate (NOBS). Suitable bleach activators are also disclosed in WO 98/17767. While any suitable bleach activator may be employed, in one aspect of the invention the subject cleaning composition may comprise NOBS, TAED or mixtures thereof.

30


When present, the peracid and/or bleach activator is generally present in the composition in an amount of from about 0.1 to about 60 wt%, from about 0.5 to

about 40 wt % or even from about 0.6 to about 10 wt% based on the composition. One or more hydrophobic peracids or precursors thereof may be used in combination with one or more hydrophilic peracid or precursor thereof.

- 5 The amounts of hydrogen peroxide source and peracid or bleach activator may be selected such that the molar ratio of available oxygen (from the peroxide source) to peracid is from 1:1 to 35:1, or even 2:1 to 10:1.

Bleach Boosting Compounds - The compositions herein may comprise one or
 10 more bleach boosting compounds. Bleach boosting compounds provide increased bleaching effectiveness in lower temperature applications. The bleach boosters act in conjunction with conventional peroxygen bleaching sources to provide increased bleaching effectiveness. This is normally accomplished through in situ formation of an active oxygen transfer agent such as a dioxirane,
 15 an oxaziridine, or an oxaziridinium. Alternatively, preformed dioxiranes, oxaziridines and oxaziridiniums may be used.

Among suitable bleach boosting compounds for use in accordance with the present invention are cationic imines, zwitterionic imines, anionic imines and/or
 20 polyionic imines having a net charge of from about +3 to about -3, and mixtures thereof. These imine bleach boosting compounds of the present invention include those of the general structure:



25 [I]

where R¹ - R⁴ may be a hydrogen or an unsubstituted or substituted radical selected from the group consisting of phenyl, aryl, heterocyclic ring, alkyl and cycloalkyl radicals.

Suitable bleach boosting compounds include zwitterionic bleach boosters zwitterionic bleach boosters, which are described in U.S. Patent Nos. 5,576,282 and 5,718,614. Other bleach boosting compounds include cationic bleach boosters described in U.S. Patent Nos. 5,360,569; 5,442,066; 5,478,357; 5,370,826; 5,482,515; 5,550,256; and WO 95/13351, WO 95/13352, and WO 95/13353.

Peroxygen sources are well-known in the art and the peroxygen source employed in the present invention may comprise any of these well known sources, including peroxygen compounds as well as compounds, which under consumer use conditions, provide an effective amount of peroxygen in situ. The peroxygen source may include a hydrogen peroxide source, the in situ formation of a peracid anion through the reaction of a hydrogen peroxide source and a bleach activator, preformed peracid compounds or mixtures of suitable peroxygen sources. Of course, one of ordinary skill in the art will recognize that other sources of peroxygen may be employed without departing from the scope of the invention. The bleach boosting compounds, when present, are typically employed in conjunction with a peroxygen source in the bleaching systems of the present invention.

Enzyme Bleaching - Enzymatic systems may be used as bleaching agents. The hydrogen peroxide may also be present by adding an enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generating hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in EP Patent Application 91202655.6 filed October 9, 1991.

The present invention compositions and methods may utilize alternative bleach systems such as ozone, chlorine dioxide and the like. Bleaching with ozone may be accomplished by introducing ozone-containing gas having ozone content from about 20 to about 300 g/m³ into the solution that is to contact the fabrics.

The gas:liquid ratio in the solution should be maintained from about 1:2.5 to about 1:6. U.S. Patent No. 5,346, 588 describes a process for the utilization of ozone as an alternative to conventional bleach systems and is herein incorporated by reference.

5


In one aspect, the fabric softening active ("FSA") is a quaternary ammonium compound suitable for softening fabric in a rinse step. In one aspect, the FSA is formed from a reaction product of a fatty acid and an aminoalcohol obtaining mixtures of mono-, di-, and, in one aspect, triester compounds. In another aspect, the FSA comprises one or more softener quaternary ammonium compounds such, but not limited to, as a monoalkylquaternary ammonium compound, a diamido quaternary compound and a diester quaternary ammonium compound, or a combination thereof.

10 In one aspect, the FSA comprises one or more softener quaternary ammonium compounds such, but not limited to, as a monoalkylquaternary ammonium compound, a diamido quaternary compound and a diester quaternary ammonium compound, or a combination thereof.

15 In one aspect of the invention, the FSA comprises a diester quaternary ammonium (hereinafter "DQA") compound composition. In certain aspects of the present invention, the DQA compounds compositions also encompasses a description of diamido FSAs and FSAs with mixed amido and ester linkages as well as the aforementioned diester linkages, all herein referred to as DQA.

20

A first type of DQA ("DQA (1)") suitable as a FSA in the present CFSC includes a compound comprising the formula:




wherein each R substituent is either hydrogen, a short chain C₁-C₆, for example C₁-C₃ alkyl or hydroxyalkyl group, e.g., methyl, ethyl, propyl, hydroxyethyl, and the like, poly (C₂₋₃ alkoxy), for example. polyethoxy, group, benzyl, or mixtures thereof; each m is 2 or 3; each n is from 1 to about 4, or 2; each Y is -O-(O)C-, -C(O)-O-, -NR-C(O)-, or -C(O)-NR- and it is acceptable for each Y to be the same or different; the sum of carbons in each R¹, plus one when Y is -O-(O)C- or -NR-C(O) -, is C₁₂-C₂₂, or C₁₄-C₂₀, with each R¹ being

30

a hydrocarbyl, or substituted hydrocarbyl group; it is acceptable for R¹ to be unsaturated or saturated and branched or linear and in one aspect it is linear; it is acceptable for each R¹ to be the same or different and typically these are the same; and X⁻ can be any softener-compatible anion, suitable anions include, chloride, bromide, methylsulfate, ethylsulfate, sulfate, phosphate, and nitrate, in one aspect the anions are chloride or methyl sulfate. Suitable DQA compounds are typically made by reacting alkanolamines such as MDEA (methyldiethanolamine) and TEA (triethanolamine) with fatty acids. Some materials that typically result from such reactions include N,N-di(acyl-oxyethyl)-N,N-dimethylammonium chloride or N,N-di(acyl-oxyethyl)-N,N-methylhydroxyethylammonium methylsulfate wherein the acyl group is derived from animal fats, unsaturated, and polyunsaturated, fatty acids, e.g., tallow, hardened tallow, oleic acid, and/or partially hydrogenated fatty acids, derived from vegetable oils and/or partially hydrogenated vegetable oils, such as, canola oil, safflower oil, peanut oil, sunflower oil, corn oil, soybean oil, tall oil, rice bran oil, palm oil, etc.

Non-limiting examples of suitable fatty acids are listed in US Patent No. 5,759,990 at column 4, lines 45-66. In one aspect, the FSA comprises other actives in addition to DQA (1) or DQA. In yet another aspect, the FSA comprises only DQA (1) or DQA and is free or essentially free of any other quaternary ammonium compounds or other actives. In yet another aspect, the FSA comprises the precursor amine that is used to produce the DQA.

In another aspect of the invention, the FSA comprises a compound, identified as DTTMAC comprising the formula:



wherein each m is 2 or 3, each R¹ is a C₆-C₂₂, or C₁₄-C₂₀, but no more than one being less than about C₁₂ and then the other is at least about 16, hydrocarbyl, or substituted hydrocarbyl substituent, for example, C₁₀-C₂₀ alkyl

or alkenyl (unsaturated alkyl, including polyunsaturated alkyl, also referred to sometimes as "alkylene"), in one aspect C₁₂-C₁₈ alkyl or alkenyl, and branch or unbranched. In one aspect, the Iodine Value (IV) of the FSA is from about 1 to 70; each R is H or a short chain C₁-C₆, or C₁-C₃ alkyl or hydroxyalkyl group, 5 e.g., methyl, ethyl, propyl, hydroxyethyl, and the like, benzyl, or (R² O)₂₋₄H where each R² is a C₁₋₆ alkylene group; and A⁻ is a softener compatible anion, suitable anions include chloride, bromide, methylsulfate, ethylsulfate, sulfate, phosphate, or nitrate; in one aspect the anions are chloride or methyl sulfate.

10 Examples of these FSAs include dialkyldimethylammonium salts and dialkylenedimethylammonium salts such as ditallowdimethylammonium and ditallowdimethylammonium methylsulfate. Examples of commercially available dialkylenedimethylammonium salts usable in the present invention are di-
15 hydrogenated tallow dimethyl ammonium chloride and ditallowdimethyl ammonium chloride available from Degussa under the trade names Adogen[®] 442 and Adogen[®] 470 respectively. In one aspect, the FSA comprises other actives in addition to DTTMAC. In yet another aspect, the FSA comprises only compounds of the DTTMAC and is free or essentially free of any other quaternary ammonium compounds or other actives.

20

In one aspect, the FSA comprises an FSA described in U.S. Pat. Pub. No. 2004/0204337 A1, published Oct. 14, 2004 to Corona et al., from paragraphs 30 – 79. In another aspect, the FSA is one described in U.S. Pat. Pub. No. 2004/0229769 A1, published Nov. 18, 2005, to Smith et al., on paragraphs 26 –
25 31; or U.S. Pat. No. 6,494,920, at column 1, line 51 *et seq.* detailing an "esterquat" or a quaternized fatty acid triethanolamine ester salt.

In one aspect, the FSA is chosen from at least one of the following: ditallowoyloxyethyl dimethyl ammonium chloride, dihydrogenated-
30 tallowoyloxyethyl dimethyl ammonium chloride, ditallow dimethyl ammonium

chloride, ditallowoyloxyethyl dimethyl ammonium methyl sulfate, dihydrogenated-tallowoyloxyethyl dimethyl ammonium chloride, dihydrogenated-tallowoyloxyethyl dimethyl ammonium chloride, or combinations thereof.

- 5 In one aspect, the FSA may also include amide containing compound compositions. Examples of diamide comprising compounds may include but not limited to methyl-bis(tallowamidoethyl)-2-hydroxyethylammonium methyl sulfate (available from Degussa under the trade names Varisoft 110 and Varisoft 222). An example of an amide-ester containing compound is N-[3-
10 (stearoylamino)propyl]-N-[2-(stearoyloxy)ethoxy]ethyl]-N-methylamine.

Another aspect of the invention provides for a rinse added fabric softening composition further comprising a cationic starch. Cationic starches are disclosed in US 2004/0204337 A1. In one aspect, the rinse added fabric softening
15 composition comprises from about 0.1% to about 7% of cationic starch by weight of the fabric softening composition. In one aspect, the cationic starch is HCP401 from National Starch.

Builders - The compositions of the present invention can comprise one or more
20 detergent builders or builder systems. When present, the compositions will typically comprise at least about 1% builder, or from about 5% or 10% to about 80%, 50%, or even 30% by weight, of said builder. Builders include, but are not limited to, the alkali metal, ammonium and alkanolammonium salts of polyphosphates, alkali metal silicates, alkaline earth and alkali metal carbonates,
25 aluminosilicate builders polycarboxylate compounds. ether hydroxypolycarboxylates, copolymers of maleic anhydride with ethylene or vinyl methyl ether, 1,3,5-trihydroxybenzene-2,4,6-trisulphonic acid, and carboxymethyl-oxysuccinic acid, the various alkali metal, ammonium and substituted ammonium salts of polyacetic acids such as ethylenediamine
30 tetraacetic acid and nitrilotriacetic acid, as well as polycarboxylates such as mellitic acid, succinic acid, oxydisuccinic acid, polymaleic acid, benzene 1,3,5-tricarboxylic acid, carboxymethyloxysuccinic acid, and soluble salts thereof.

Chelating Agents - The compositions herein may also optionally contain one or more copper, iron and/or manganese chelating agents. If utilized, chelating agents will generally comprise from about 0.1% by weight of the compositions herein to about 15%, or even from about 3.0% to about 15% by weight of the compositions herein.

Dye Transfer Inhibiting Agents - The compositions of the present invention may also include one or more dye transfer inhibiting agents. Suitable polymeric dye transfer inhibiting agents include, but are not limited to, polyvinylpyrrolidone polymers, polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof. When present in the compositions herein, the dye transfer inhibiting agents are present at levels from about 0.0001%, from about 0.01%, from about 0.05% by weight of the cleaning compositions to about 10%, about 2%, or even about 1% by weight of the cleaning compositions.

Dispersants - The compositions of the present invention can also contain dispersants. Suitable water-soluble organic materials are the homo- or copolymeric acids or their salts, in which the polycarboxylic acid may comprise at least two carboxyl radicals separated from each other by not more than two carbon atoms.

Enzymes - The compositions can comprise one or more detergent enzymes which provide cleaning performance and/or fabric care benefits. Examples of suitable enzymes include, but are not limited to, hemicellulases, peroxidases, proteases, cellulases, xylanases, lipases, phospholipases, esterases, cutinases, pectinases, keratanases, reductases, oxidases, phenoloxidases, lipoxygenases, ligninases, pullulanases, tannases, pentosanases, malanases, β -glucanases, arabinosidases, hyaluronidase, chondroitinase, laccase, and amylases, or mixtures thereof. A typical combination is a cocktail of conventional applicable

enzymes like protease, lipase, cutinase and/or cellulase in conjunction with amylase.

5 Enzyme Stabilizers - Enzymes for use in compositions, for example, detergents can be stabilized by various techniques. The enzymes employed herein can be stabilized by the presence of water-soluble sources of calcium and/or magnesium ions in the finished compositions that provide such ions to the enzymes.

10 Processes of Making Laundry Care Compositions

The laundry care compositions of the present invention can be formulated into any suitable form and prepared by any process chosen by the formulator, non-limiting examples of which are described in Applicants' examples and in U.S. 5,879,584; U.S. 5,691,297; U.S. 5,574,005; U.S. 5,569,645; U.S. 5,565,422; 15 U.S. 5,516,448; U.S. 5,489,392; U.S. 5,486,303 all of which are incorporated herein by reference.

The liquid detergent compositions may be in the form of an aqueous solution or uniform dispersion or suspension of surfactant, hueing agent, and certain 20 optional other ingredients, some of which may normally be in solid form, that have been combined with the normally liquid components of the composition, such as the liquid alcohol ethoxylate nonionic, the aqueous liquid carrier, and any other normally liquid optional ingredients. Such a solution, dispersion or suspension will be acceptably phase stable and will typically have a viscosity 25 which ranges from about 100 to 600 cps, or from about 150 to 400 cps. For purposes of this invention, viscosity is measured with a Brookfield LVDV-II+ viscometer apparatus using a #21 spindle.

The liquid detergent compositions herein can be prepared by combining the 30 components thereof in any convenient order and by mixing, e.g., agitating, the resulting component combination to form a phase stable liquid detergent composition. In a process for preparing such compositions, a liquid matrix is

formed containing at least a major proportion, or even substantially all, of the liquid components, e.g., nonionic surfactant, the non-surface active liquid carriers and other optional liquid components, with the liquid components being thoroughly admixed by imparting shear agitation to this liquid combination. For example, rapid stirring with a mechanical stirrer may usefully be employed. While shear agitation is maintained, substantially all of any anionic surfactants and the solid form ingredients can be added. Agitation of the mixture is continued, and if necessary, can be increased at this point to form a solution or a uniform dispersion of insoluble solid phase particulates within the liquid phase. After some or all of the solid-form materials have been added to this agitated mixture, particles of any enzyme material to be included, e.g., enzyme prills, are incorporated. As a variation of the composition preparation procedure hereinbefore described, one or more of the solid components may be added to the agitated mixture as a solution or slurry of particles premixed with a minor portion of one or more of the liquid components. After addition of all of the composition components, agitation of the mixture is continued for a period of time sufficient to form compositions having the requisite viscosity and phase stability characteristics. Frequently this will involve agitation for a period of from about 30 to 60 minutes.

20

In one aspect of forming the liquid detergent compositions, the hueing agent is first combined with one or more liquid components to form a hueing agent premix, and this hueing agent premix is added to a composition formulation containing a substantial portion, for example more than 50% by weight, more specifically, more than 70% by weight, and yet more specifically, more than 90% by weight, of the balance of components of the laundry detergent composition. For example, in the methodology described above, both the hueing agent premix and the enzyme component are added at a final stage of component additions. In another aspect, the hueing agent is encapsulated prior to addition to the detergent composition, the encapsulated hueing agent is suspended in a structured liquid, and the suspension is added to a composition formulation

30

containing a substantial portion of the balance of components of the laundry detergent composition.

As noted previously, the detergent compositions may be in a solid form. Suitable
5 solid forms include tablets and particulate forms, for example, granular particles, flakes or sheets. Various techniques for forming detergent compositions in such solid forms are well known in the art and may be used herein. In one aspect, for example when the composition is in the form of a granular particle, the hueing agent is provided in particulate form, optionally including additional but not all
10 components of the laundry detergent composition. The hueing agent particulate is combined with one or more additional particulates containing a balance of components of the laundry detergent composition. Further, the hueing agent, optionally including additional but not all components of the laundry detergent composition, may be provided in an encapsulated form, and the hueing agent
15 encapsulate is combined with particulates containing a substantial balance of components of the laundry detergent composition.

The compositions of this invention, prepared as hereinbefore described, can be used to form aqueous washing solutions for use in the laundering of fabrics.
20 Generally, an effective amount of such compositions is added to water, for example in a conventional fabric laundering automatic washing machine, to form such aqueous laundering solutions. The aqueous washing solution so formed is then contacted, typically under agitation, with the fabrics to be laundered therewith. An effective amount of the liquid detergent compositions herein
25 added to water to form aqueous laundering solutions can comprise amounts sufficient to form from about 500 to 7,000 ppm of composition in aqueous washing solution, or from about 1,000 to 3,000 ppm of the detergent compositions herein will be provided in aqueous washing solution.

30 Method of Use

Certain of the consumer products disclosed herein can be used to clean or treat a situs *inter alia* a surface or fabric. Typically at least a portion of the situs is

contacted with an embodiment of Applicants' consumer product, in neat form or diluted in a liquor, for example, a wash liquor and then the situs may be optionally washed and/or rinsed. In one aspect, a situs is optionally washed and/or rinsed, contacted with an aspect of the consumer product and then optionally washed and/or rinsed. For purposes of the present invention, washing includes but is not limited to, scrubbing, and mechanical agitation. The fabric may comprise most any fabric capable of being laundered or treated in normal consumer use conditions. Liquors that may comprise the disclosed compositions may have a pH of from about 3 to about 11.5. Such compositions are typically employed at concentrations of from about 500 ppm to about 15,000 ppm in solution. When the wash solvent is water, the water temperature typically ranges from about 5 °C to about 90 °C and, when the situs comprises a fabric, the water to fabric ratio is typically from about 1:1 to about 30:1. Employing one or more of the aforementioned methods results in a treated situs.

15

In one aspect, a method of treating and/or cleaning a surface or fabric comprising the steps of optionally washing and/or rinsing said surface or fabric, contacting said surface or fabric with any laundry care composition disclosed in this specification, then optionally washing and/or rinsing said surface and/or fabric then optionally letting said surface or fabric to dry and/or actively drying said surface or fabric, is disclosed.

20

Examples


The following examples are provided to further illustrate the hueing agents of the present invention; however, they are not to be construed as limiting the invention as defined in the claims appended hereto. In fact, it will be apparent to those skilled in the art that various modifications and variations can be made in this invention without departing from the scope or spirit of the invention. All parts and percents given in these examples are by weight unless otherwise indicated.

30

Sample Preparation and Test Methods

A. Sample Preparation

The following general procedures were used to prepare the thiophene azo dyes
5 of the present invention.

Preparation of Glycidol Materials

10


18 parts N-ethyl-m-toluidine, 52 parts isopropyl-glycidyl ether, and 50 parts
toluene were charged into a 200 mL round bottom flask at room temperature.
The mixture was refluxed overnight. The solvent was removed and the resulting
material was used crude for the next step. These materials were then
15 alkoxyated as described herein.

Preparation of Alkoxyated Intermediates

General procedure for alkoxylation – Method A:

20


Intermediate Type 1, initial single site polymer



25 18 parts N-ethyl-m-toluidine, 14 parts acetic acid, and 20 parts water were
charged into a 200 mL stainless steel autoclave at room temperature. 11 parts


ethylene oxide was added over several hours. After the reaction was kept for 24 hours at room temperature, the vessel was charged with 0.2 parts NaOH and heated up to 125°C. Then, ethylene oxide was added over about 1 hour. After continuing the reaction for another 3 hours at 125°C, residual EO was removed by vacuum. Then it was cooled to room temperature and the intermediate obtained was an amber-colored viscous liquid. The constant average number of EO could be achieved every time by controlling the amount of ethylene oxide in the synthesis.

10 Intermediate Type 2, initial dual chain polymer




18 parts 2-Methoxy-5-methylaniline, 14 parts acetic acid, and 20 parts water were charged into a 200 mL stainless steel autoclave at room temperature. 22 parts ethylene oxide was added over several hours. After the reaction was kept for 24 hours at room temperature, the vessel was charged with 0.2 parts NaOH and heated up to 125°C. Then, 40 parts ethylene oxide was added over about 1 hour. After continuing the reaction for another 3 hours at 125°C, residual EO was removed by vacuum. Then it was cooled to room temperature and the intermediate obtained was an amber-colored viscous liquid. The constant average number of EO could be achieved every time by controlling the amount of ethylene oxide in the synthesis.

25 General procedure for alkoxylation – Method B:



18 parts aniline, 60 parts mono-bromo-polyethylene glycol-200, 20 parts NaHCO₃ and 50 parts toluene were charged into a 200 mL round bottom flask at room temperature. The reaction mixture was heated to 80°C for 5 hours. The salt was filtered and the solvent removed from the filtrate via low pressure distillation. The crude product was used with no further purification for the next step.



10


It is also possible to use a mono-bromo-mono-chloro glycol in order to directly make the halogenated product using this same procedure above.

15

General Procedures for Color Synthesis

All colors were prepared following the same general procedure.

Color intermediate synthesis for Example 1



20


2 parts amino-thiophene, and 30 parts phosphoric acid, were charged into 200 mL glass flask and cooled to 0-5°C. 1 part NaNO₂ was slowly added as a solid, maintaining the temperature below 10°C. When addition was completed for

diazotization, the mixture was stirred for 30 minutes and excess sodium nitrite was consumed by adding 0.3 parts sulfamic acid. Enough sulfamic acid was added until starch iodide paper provided a negative result. 5 parts of the m-toluidine ethoxylate coupler was added to a flask with 100 mL of water and the temperature lowered to 5°C. The prepared diazonium salt solution was slowly added into the above solution for coupling reaction. Care was taken to not allow the temperature to rise above 10°C. After complete addition of diazonium salt solution, the reaction was allowed to slowly reach room temperature over an hour. The mixture was then neutralized with sodium hydroxide and phase separated. The product layer was then dissolved with methanol and filtered to remove any excess salts. The filtrate was evaporated and the product of this reaction is ready to use at this point.

Addition of anhydride to color

15


Example 1



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of succinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

87


Example 3



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of succinic
 5 anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to
 room temperature. The pH of the mixture is adjusted to 7 using sodium
 hydroxide, and the water is removed from the mixture via rotary evaporation. The
 desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 4




To 25 parts of the thiophene azo polymeric colorant is added 8 parts of maleic
 15 anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to
 room temperature. The pH of the mixture is adjusted to 7 using sodium

hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

5


Example 5



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of phthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.


15

Example 7



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of hexahydrophthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

Example 6




10

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of allyl succinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

15


Example 8



- 5 To 25 parts of the thiophene azo polymeric colorant is added 8 parts of octylsuccinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.


10

Example 10



- 15 To 25 parts of the thiophene azo polymeric colorant is added 8 parts of succinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

Example 11




5

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of maleic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 12




15

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of phthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to

room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

5


Example 14



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of hexahydrophthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

15


Example 13



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of allyl succinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 15




15

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of octylsuccinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

20


Example 17



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of succinic
 5 anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to
 room temperature. The pH of the mixture is adjusted to 7 using sodium
 hydroxide, and the water is removed from the mixture via rotary evaporation. The
 desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 18




To 25 parts of the thiophene azo polymeric colorant is added 8 parts of maleic
 15 anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to
 room temperature. The pH of the mixture is adjusted to 7 using sodium

hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

5

Example 19




10

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of phthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

15


Example 21



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of hexahydrophthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 24




15

To 25 parts of the thiophene azo polymeric colorant is added 8 parts of succinic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

20


Example 25



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of maleic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

10

Example 34




To 25 parts of the thiophene azo polymeric colorant is added 8 parts of phthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

5


Example 44



To 25 parts of the thiophene azo polymeric colorant is added 8 parts of hexahydrophthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.

15

Example 45



- 5 To 25 parts of the thiophene azo polymeric colorant is added 16 parts of hexahydro-phthalic anhydride. The reaction is heated to 85°C for 12 hours and then cooled down to room temperature. The pH of the mixture is adjusted to 7 using sodium hydroxide, and the water is removed from the mixture via rotary evaporation. The desired product is obtained in quantitative conversion as assessed by HPLC.
- 10

B. Test Methods

I. Method for Determining Extinction Coefficient

15

A sample of 0.1000 grams of color is diluted to volume in a 100mL volumetric flask using methanol. 2mL of this solution are transferred to a second 100mL volumetric flask and diluted to volume with methanol. The absorbance and lambda max of the diluted colorant are measured using a spectrophotometer in a 1cm cell.

20

Lambda max is taken directly from the spectra.

Extinction coefficient = ((absorbance/mass used)X5)/ MW of colorant

Exemplary Detergent Formulations

5 Formulations 1a – 1l: Liquid Detergent Formulations

Tables 7A and 7B provide examples of liquid detergent formulations which include at least one thiophene azo dye of the present invention as a hueing agent. The formulations are shown in Table 7A as Formulations 1a through 1f and in Table 7B as Formulations 1g through 1l.

Table 7A – Liquid Detergent Formulations Comprising the Inventive Hueing Agent

Ingredient	1a	1b	1c	1d	1e	1f ⁵
sodium alkyl ether sulfate	wt % 14.4%	wt % 14.4%	wt %	wt % 9.2%	wt % 5.4%	wt %
linear alkylbenzene sulfonic acid	4.4%	4.4%	12.2%	5.7%	1.3%	22.0%
alkyl ethoxylate	2.2%	2.2%	8.8%	8.1%	3.4%	18.0%
amine oxide	0.7%	0.7%	1.5%			
citric acid	2.0%	2.0%	3.4%	1.9%	1.0%	1.6%
fatty acid	3.0%	3.0%	8.3%			16.0%
protease	1.0%	1.0%	0.7%	1.0%		2.5%
amylase	0.2%	0.2%	0.2%			0.3%
lipase				0.2%		
borax	1.5%	1.5%	2.4%	2.9%		
calcium and sodium formate	0.2%	0.2%				
formic acid						1.1%
amine ethoxylate polymers	1.8%	1.8%	2.1%			3.2%
sodium polyacrylate					0.2%	
sodium polyacrylate copolymer				0.6%		
DTPA ¹	0.1%	0.1%				0.9%
DTPMP ²			0.3%			
EDTA ³					0.1%	
fluorescent whitening agent	0.15%	0.15%	0.2%	0.12%	0.12%	0.2%
ethanol	2.5%	2.5%	1.4%	1.5%		

propanediol	6.6%	6.6%	4.9%	4.0%		15.7%
sorbitol				4.0%		
ethanolamine	1.5%	1.5%	0.8%	0.1%		11.0%
sodium hydroxide	3.0%	3.0%	4.9%	1.9%	1.0%	
sodium cumene sulfonate			2.0%			
silicone suds suppressor			0.01%			
perfume	0.3%	0.3%	0.7%	0.3%	0.4%	0.6%
Non-tinting dyes ⁴	0.0001%	0.001%	0.008%	0.03%	0.015%	0.05%
Hueing Agent ⁶	0.01%				0.005%	
Hueing Agent ⁷		0.01%	0.02%	0.003%		0.012%
water	balance	balance	balance	balance	balance	balance
	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

Table 7B – Liquid Detergent Formulations Comprising the Inventive Hueing Agent

5

Ingredient	1g	1h	1i	1j	1k	1l ⁵
	wt %	wt %	wt %	wt %	wt %	wt %
sodium alkyl ether sulfate	14.4%	14.4%		9.2%	5.4%	
linear 1benzene sulfonic acid	4.4%	4.4%	12.2%	5.7%	1.3%	22.0%
alkyl ethoxylate	2.2%	2.2%	8.8%	8.1%	3.4%	18.0%
amine oxide	0.7%	0.7%	1.5%			
citric acid	2.0%	2.0%	3.4%	1.9%	1.0%	1.6%
fatty acid	3.0%	3.0%	8.3%			16.0%
protease	1.0%	1.0%	0.7%	1.0%		1.7%
amylase	0.2%	0.2%	0.2%			0.6%
lipase				0.2%		0.2%
borax	1.5%	1.5%	2.4%	2.9%		
calcium and sodium formate	0.2%	0.2%				
formic acid						1.1%
amine ethoxylate polymers	1.8%	1.8%	2.1%			3.2%
sodium polyacrylate					0.2%	
sodium polyacrylate copolymer				0.6%		
DTPA ¹	0.1%	0.1%				0.9%
DTPMP ²			0.3%			
EDTA ³					0.1%	
fluorescent whitening agent	0.15%	0.15%	0.2%	0.12%	0.12%	0.2%

ethanol	2.5%	2.5%	1.4%	1.5%		
propanediol	6.6%	6.6%	4.9%	4.0%		15.7%
sorbitol				4.0%		
ethanolamine	1.5%	1.5%	0.8%	0.1%		11.0%
sodium hydroxide	3.0%	3.0%	4.9%	1.9%	1.0%	
sodium cumene sulfonate			2.0%			
silicone suds suppressor			0.01%			
perfume	0.3%	0.3%	0.7%	0.3%	0.4%	0.6%
Non-tinting dyes ⁴	0.0001%	0.001 %	0.008 %	0.03%	0.015%	0.05%
Hueing Agent ⁶	0.01%				0.005%	
Hueing Agent ⁷		0.01%	0.02%	0.003 %		0.012 %
opacifier ⁸		0.5%				
water	balance	balance	balance	balance	balance	balance
	100.0%	100.0 %	100.0 %	100.0 %	100.0%	100.0 %

Footnotes for Formulations 1a-l:

¹ diethylenetriaminepentaacetic acid, sodium salt

5 ² diethylenetriaminepentakis(methylenephosphonic acid, sodium salt

³ ethylenediaminetetraacetic acid, sodium salt

⁴ a non-tinting dyes used to adjust formula color

⁵ compact formula, packaged as a unitized dose in polyvinyl alcohol film

10 ⁶ thiophene azo carboxylate hueing agent from Table 6, Examples 1-21, preferably with hueing efficiency >10 and wash removability 30-85%

⁷ thiophene azo carboxylate hueing agent from Table 6, Examples 22-48, preferably with hueing efficiency >10 and wash removability 30-85%

⁸ Acusol OP301

15

Formulations 2a – 2e: Granular Detergent Formulations

Table 8 provides examples of granular detergent formulations which include at least one thiophene azo carboxylate dye of the present invention as a hueing agent. The formulations are shown in Table 8 as Formulations 2a through 2e.

20

Table 8 – Granular Detergent Formulations Comprising the Inventive Hueing Agent

Ingredient	2a	2b	2c	2d	2e
	wt %	wt %	wt %	wt %	wt %
Na linear alkylbenzene sulfonate	3.4%	3.3%	11.0%	3.4%	3.3%
Na alkylsulfate	4.0%	4.1%		4.0%	4.1%
Na alkyl sulfate (branched)	9.4%	9.6%		9.4%	9.6%
alkyl ethoxylate			3.5%		
type A zeolite	37.4%	35.4%	26.8%	37.4%	35.4%
sodium carbonate	22.3%	22.5%	35.9%	22.3%	22.5%
sodium sulfate	1.0%		18.8%	1.0%	
sodium silicate			2.2%		
protease	0.1%	0.2%		0.1%	0.2%
sodium polyacrylate	1.0%	1.2%	0.7%	1.0%	1.2%
carboxymethylcellulose			0.1%		
PEG 600		0.5%			0.5%
PEG 4000		2.2%			2.2%
DTPA	0.7%	0.6%		0.7%	0.6%
fluorescent whitening agent	0.1%	0.1%	0.1%	0.1%	0.1%
sodium percarbonate		5.0%			5.0%
sodium nonanoyloxybenzenesulfonate		5.3%			5.3%
silicone suds suppressor	0.02%	0.02%		0.02%	0.02%
perfume	0.3%	0.3%	0.2%	0.3%	0.3%
Hueing Agent ¹	0.004%				0.002%
Hueing Agent ²		0.006%	0.002%	0.004%	
water and miscellaneous	balance	balance	balance	balance	balance
	100.0%	100.0%	100.0%	100.0%	100.0%

5 Footnotes for Formulations 2a-e:

¹thiophene azo carboxylate hueing agent from Table 6, Examples 1-21, preferably with hueing efficiency >10 and wash removability 30-85%

²thiophene azo carboxylate hueing agent from Table 6, Examples 22-48, preferably with hueing efficiency >10 and wash removability 30-85%

10

Exemplary Fabric Care Compositions

Formulations 3a – 3d: Liquid Fabric Care Compositions

Table 9 provides examples of liquid fabric care compositions which include at least one thiophene azo carboxylate dye of the present invention as a hueing agent. The compositions are shown in Table 9 as Formulations 3a through 3d.

5 Table 9 – Liquid Fabric Care Compositions
Comprising the Inventive Hueing Agent

Ingredients	3a	3b	3c	3d
Fabric Softening Active ^a	13.70%	13.70%	13.70%	13.70%
Ethanol	2.14%	2.14%	2.14%	2.14%
Cationic Starch ^b	2.17%	2.17%	2.17%	2.17%
Perfume	1.45%	1.45%	1.45%	1.45%
Phase Stabilizing Polymer ^c	0.21%	0.21%	0.21%	0.21%
Calcium Chloride	0.147%	0.147%	0.147%	0.147%
DTPA ^d	0.007%	0.007%	0.007%	0.007%
Preservative ^e	5 ppm	5 ppm	5 ppm	5 ppm
Antifoam ^f	0.015%	0.015%	0.015%	0.015%
Hueing Agent ^g	30 ppm			15 ppm
Hueing Agent ^h		30 ppm		
Hueing Agent ⁱ			30 ppm	15 ppm
Tinopal CBS-X ^j	0.2	0.2	0.2	0.2
Ethoquad C/25 ^k	0.26	0.26	0.26	0.26
Ammonium Chloride	0.1%	0.1%	0.1%	0.1%
Hydrochloric Acid	<u>0.012 %</u>	<u>0.012 %</u>	<u>0.012 %</u>	<u>0.012 %</u>
Deionized Water	Balance	Balance	Balance	Balance

Footnotes for Formulations 3a-d:

- 10 ^a N,N-di(tallowoyloxyethyl)-N,N-dimethylammonium chloride.
- ^b Cationic starch based on common maize starch or potato starch, containing 25% to 95% amylose and a degree of substitution of from 0.02 to 0.09, and having a viscosity measured as Water Fluidity having a value from 50 to 84.
- 15 ^c Copolymer of ethylene oxide and terephthalate having the formula described in US 5,574,179 at col.15, lines 1-5, wherein each X is methyl, each n is 40, u is 4, each R¹ is essentially 1,4-phenylene moieties, each R² is essentially ethylene, 1,2-propylene moieties, or mixtures thereof.
- ^d Diethylenetriaminepentaacetic acid.
- ^e KATHON[®] CG available from Rohm and Haas Co.
- 20 ^f Silicone antifoam agent available from Dow Corning Corp. under the trade name DC2310.
- ^g thiophene azo carboxylate hueing agent from Table 6, Examples 1-21, preferably with hueing efficiency >10 and wash removability 30-85%

^h thiophene azo carboxylate hueing agent from Table 6, Examples 22-48, preferably with hueing efficiency >10 and wash removability 30-85%

ⁱ thiophene azo carboxylate hueing agent from Table 6, Examples 1-48, preferably with hueing efficiency >10 and wash removability 30-85%

5 ^j Disodium 4,4'-bis-(2-sulfostyryl) biphenyl, available from Ciba Specialty Chemicals.

^k Cocomethyl ethoxylated [15] ammonium chloride, available from Akzo Nobel.

* * *

10 The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm".

15

All documents cited in the Detailed Description of the Invention are, in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present invention. To the extent that any meaning or definition of a term in this document conflicts with any meaning or definition of the same term in a document incorporated by reference, the meaning or definition assigned to that term in this document shall govern.

20


While particular aspects of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

25

CLAIMS


We claim:

- 5 1. A composition comprising a thiophene azo carboxylate dye containing a carboxylic acid moiety, with the proviso that the thiophene azo carboxylate dye does not have the following structures




10

and



2. The composition of Claim 1, wherein said thiophene azo carboxylate dye has, in the wavelength range of about 400 nm to about 750 nm in methanol solution, a maximum extinction coefficient greater than about 1000 liter/mol/cm.
- 5 3. The composition of Claim 2, wherein said thiophene azo carboxylate dye has, in the wavelength range of about 540 nm to about 630 nm, a maximum extinction coefficient from about 20,000 to about 100,000 liter/mol/cm.
4. The composition of Claim 3, wherein said thiophene azo carboxylate dye has, in the wavelength range of about 560 nm to about 610 nm, a maximum extinction coefficient from about 20,000 to about 65,000 liter/mol/cm.
- 10 5. The composition of Claim 1, wherein said thiophene azo carboxylate dye has a molecular weight from greater than 232 Daltons, from about 233 Daltons to about 5000 Daltons, from about 365 Daltons to about 2500 Daltons, or even from about 423 Daltons to about 1000 Daltons.
- 15 6. The composition of Claim 1, wherein said thiophene azo carboxylate dye is represented by general Formula (I):
- 20



Formula I

wherein:

- 25 a.) R₁, R₂ and R₃ are each independently selected from hydrogen, electron-withdrawing moieties, and electron-donating moieties, provided that at least one of R₁, R₂ and R₃ is an electron-withdrawing moiety; and


b.) wherein X is an organic moiety having a molecular weight from about 104 Daltons to about 4871 Daltons, from about 236 Daltons to about 2371 Daltons from about 294 Daltons to about 871 Daltons.

5

7. The composition of Claim 6, wherein for said thiophene azo carboxylate dye each R_1 , R_2 and R_3 is independently selected from hydrogen, (C₁-C₄)-alkyl, (C₃-C₁₀)-aryl, carboxylate, cyano, sulfonate, phosphonate, sulfate, acetate, nitro, (C₁-C₄)-alkyl ester, halogen or amino moiety.

10

8. The composition of Claim 6, wherein for said thiophene azo carboxylate dye X is a moiety having Formula (II) below:




Formula II

wherein:

20

i.) R_4 is selected from a moiety having Formula (III) below



Formula III


25

wherein:

each R_8 is independently selected from hydrogen, C_1 - C_8 alkyl optionally substituted with a hydroxy, or acetyl;

m is an integer from 0 to 10;


Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IV,



Formula IV

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

ii.) R_4 is selected from a moiety having Formula (V) below:



Formula V


wherein

each R_{11} and R_{12} is independently selected from hydrogen, C_1 - C_8 alkyl, aryl, acetyl or hydroxyl moiety; m and n are independent and are integers from 0 to 10,

110

Y is $-\text{OC}(\text{O})\text{GCO}_2\text{M}$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VI,

5



Formula VI


said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons; or

10

iii.) R_4 is selected from a moiety having Formula (VII)

15

below:



Formula VII

wherein

R_{13} is selected from a C_6 - C_{10} aryl moiety, C_7 - C_{12} arylalkyl moiety, C_1 - C_{18} alkyl moiety, or a siloxane moiety;


20

each R_{14} is independently selected from hydrogen, C_1 - C_4 alkyl; m is an integer from 0 to 10;

25

Y is $-\text{OC}(\text{O})\text{GCO}_2\text{M}$ wherein M is H or a charge balancing cation; G is an organic group

derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VIII,



Formula VIII

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;


R_5 is an independently selected R_4 moiety as defined above or R_5 is selected from C_1 - C_{12} alkyl moiety, a C_2 - C_{12} ether moiety; C_6 - C_{10} aryl moiety or C_7 - C_{12} arylalkyl moiety; wherein the index a is an integer from 0 to 4, and each R_6 may be independently selected from a C_1 - C_6 alkyl, a C_1 - C_4 alkoxy, a nitro, a hydroxyl, a halogen, or $-NHC(O)R_{22}$ wherein R_{22} is selected from H, $-NH_2$, C_1 - C_6 alkyl, phenyl, $-(CH_2)_sOR_{23}$ where the index s is 1 or 2 and R_{23} is selected from Me, phenyl, and $-CO_2CH_2CN$; $-NHSO_2R_{24}$ wherein R_{24} is C_1 - C_4 alkyl or phenyl; said alkyl, alkoxy and acetamido moieties may be optionally substituted with a carboxylate moiety;

or, X is a moiety having Formula II above:

wherein each R_4 and R_5 can independently be selected from:

- a) $[(CH_2CR'HO)_x(CH_2CR''HO)_yR_{15}]$;
- b) C_1 - C_{12} alkyl, C_6 - C_{10} aryl or C_7 - C_{12} aryl alkyl;
- c) $[CH_2CH(OR_{16})CH_2OR_{17}]$;

- d) the amino addition product of styrene oxide, glycidyl methyl ether, isobutyl glycidyl ether, isopropylglycidyl ether, t-butyl glycidyl ether, 2-ethylhexylglycidyl ether, and glycidylhexadecyl ether, followed by the addition of from 1 to 10 alkylene oxide units wherein at least one such alkyleneoxide unit is substituted with R_{15} that is not $-H$; wherein R' is selected from the group consisting of H , CH_3 , $CH_2O(CH_2CH_2O)_z R_{15}$, and mixtures thereof; R'' is selected from the group consisting of H , $CH_2O(CH_2CH_2O)_z R_{15}$, and mixtures thereof; $x + y \leq 20$; $y \geq 1$; $z = 0$ to 10 ; each R_{15} is independently selected from $-H$ and Y wherein Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IX,




Formula IX

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

R_{16} is selected from the group consisting of H , $(CH_2CH_2O)_z R_{15}$ wherein $z = 0$ to 10 , and mixtures thereof; R_{17} is selected from the group consisting of C_1 - C_{16} alkyl, C_6 - C_{10} aryl groups, and mixtures thereof; the index m is an integer from 0 to 4 and each R_6 is as defined above; further provided the molecule contains at least one non- H R_{15} group.

9. The composition of Claim 8, wherein for said thiophene azo carboxylate dye X is a moiety having Formula (II) below:

113




Formula II

5

wherein:

- i.) R_4 is selected from a moiety having Formula (III) below



10

Formula III


wherein:

R_8 is a hydrogen, C_1 - C_4 alkyl moiety or an C_6 - C_{10} aryl moiety;

15

Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula IV,

20




Formula IV

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons

114

to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

or, R_4 is selected from a moiety having Formula (V) below:




Formula V

wherein

each R_{11} and R_{12} is independently selected from hydrogen, C_1 - C_4 alkyl or C_6 - C_{10} aryl moiety; m and n are independent and are integers from 0 to 5,

Y is $-\text{OC}(\text{O})\text{GCO}_2\text{M}$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VI,




Formula VI

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

or, R_4 is selected from a moiety having Formula (VII) below:


115



Formula VII

wherein

- 5 R_{13} is selected from an C_6 - C_{10} aryl moiety, benzyl moiety, or a C_1 - C_{18} alkyl moiety;
- each R_{14} is independently selected from hydrogen or $-CH_3$; m is an integer from 0 to 10,
- 10 Y is $-OC(O)GCO_2M$ wherein M is H or a charge balancing cation; G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having Formula VIII,




Formula VIII

- 15 said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 Daltons;

- 20 ii.) R_5 is an independently selected R_4 moiety as defined above or R_5 is R_5 selected from C_1 - C_6 alkyl moiety or benzyl moiety;
- iii.) wherein the index a is an integer from 0 to 2, and each
- 25 R_6 may be independently selected from methyl, methoxy, or acetamido moiety.


10. The composition of Claim 9, wherein for said thiophene azo carboxylate dye each R₁, R₂ and R₃ is independently selected from hydrogen, (C₁-C₄)-alkyl, (C₆-C₁₀)-aryl, carboxylate, cyano, sulfonate, acetate, nitro, (C₁-C₄)-alkyl ester, 5 halogen or amino moiety, in one aspect, each R₁, R₂ and R₃ is independently selected from hydrogen, (C₁-C₄)-alkyl, carboxylate, cyano, sulfonate, nitro, (C₁-C₄)-alkyl ester or halogen moiety.

11. The composition of Claim 1 wherein said thiophene azo carboxylate dye has 10 Formula (X):



Formula X

15 wherein the A moiety is selected from the group consisting of Table 1 A Moieties Nos. 1 – 118 and wherein X is selected from the group consisting of Table 4 X Moieties Nos. 1 – 24 wherein G is an organic group derived from a substituted or unsubstituted succinic, maleic, glutaric, adipic or phthalic anhydride having 20 Formula XI,




Formula XI

said anhydride having a molecular weight less than about 353 Daltons, from about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325 25 Daltons.

12. The composition of Claim 1 wherein said thiophene azo carboxylate dye has Formula (XII):


117



Formula XII

wherein the A moiety is selected from the group consisting of Table 1 A Moieties
 5 Nos. 9 – 11, 15, 23, 34 – 35, 37 – 39, 41, 43, 47, 50 – 51, 57 – 58, 77, 83, 89,
 95, 106, and 110 – 118; and wherein the X moiety is selected from the group
 consisting of Table 4 X Moieties Nos. 1 – 8, 13, 14, and 18 – 21, wherein G is an
 organic group derived from a substituted or unsubstituted succinic, maleic,
 glutaric, adipic or phthalic anhydride having Formula XIII,

10




Formula XIII

said anhydride having a molecular weight less than about 353 Daltons, from
 15 about 98 Daltons to 352 Daltons, or even from about 100 Daltons to about 325
 Daltons.

13. The composition of Claim 12 wherein said G moiety of said thiophene azo
 carboxylate dye is selected from Table 5 G Moieties Nos. 1 – 48.

20

14. The composition of Claim 1, wherein said thiophene azo carboxylate dye has
 the following formula:




25

wherein the moiety A is selected from Table 1 A Moieties Nos. 1 – 118; a = 0 to
 2; when a = 1 or 2, R₆ is selected from Table 2 R₆ Substituent Identity and

Position Nos. 1 – 40; and R₄ and R₅ grouping is selected from Table 3 R₄ and R₅ Groupings Nos. 1 – 83.

15. The composition of Claim 1, wherein said thiophene azo carboxylate dye has the following formula:



wherein the moiety A is selected from Table 1 A Moieties Nos. 9 – 11, 15, 23, 34 – 35, 37 – 39, 41, 43, 47, 50 – 51, 57 – 58, 77, 83, 89, 95, 106, and 110 – 118; a = 0 to 2; when a = 1 or 2, R₆ is selected from Table 2 R₆ Substituent Identity and Position Nos. 1, 3, 5, 7, 12, 13, 14, 31, 36 and 40; and R₄ and R₅ grouping is selected from Table 3 R₄ and R₅ Groupings Nos. 3 – 8, 14, 15, 18 – 23, 29, 30, 33 – 38, 44, 45, 48 – 53, 59, 60, 63-66, 71-76, 82 and 83.

15

16. The composition of Claim 1, wherein said thiophene azo carboxylate dye has a formula selected from Table 6 Formula Nos. 1 – 48, in one aspect, said thiophene azo carboxylate dye has a formula selected from Table 6 Formula Nos. 1 – 36, 43 and 44.

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2012/039537

A. CLASSIFICATION OF SUBJECT MATTER INV. C09B29/00 C09B69/00 C11D3/42 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) C09B		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, CHEM ABS Data, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 2 262663 A (FUJI PHOTO FILM CO LTD) 25 October 1990 (1990-10-25) compound 31 -----	1-7
X	DE 23 33 828 A1 (ICI LTD) 17 January 1974 (1974-01-17) example s -----	1-7
X	GB 1 465 941 A (ICI LTD) 2 March 1977 (1977-03-02) the whole document -----	1-7
X	EP 0 474 402 A2 (ICI PLC [GB]) 11 March 1992 (1992-03-11) 2nd formula on page 6 -----	1
	-/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 24 August 2012	Date of mailing of the international search report 31/08/2012	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Friebel, Friedrich	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2012/039537

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DEL AMO V ET AL: "Two-colour screening in combinatorial chemistry: prospecting for enantioselectivity in a library of steroid-based receptors", TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 65, no. 32, 8 August 2009 (2009-08-08), pages 6370-6381, XP026281500, ISSN: 0040-4020, DOI: 10.1016/J.TET.2009.06.018 [retrieved on 2009-06-10] Scheme 3: comp.18,19; Scheme 4: comp.24,25 -----	1-7
Y	US 2008/177090 A1 (VALENTI MICHAEL A [US] ET AL) 24 July 2008 (2008-07-24) the whole document -----	1-16
Y	WO 2011/017719 A2 (MILLIKEN & CO [US]; TORRES EDUARDO [US]; HONG XIAOYONG [US]; MAHAFFEY) 10 February 2011 (2011-02-10) claims 9,10; examples 14,28,46,62-64 -----	1-16
A	DE 10 2007 040928 A1 (CLARIANT INT LTD [CH]) 5 March 2009 (2009-03-05) the whole document -----	1-16
A	US 2003/212172 A1 (DANIELSON TODD D [US] ET AL) 13 November 2003 (2003-11-13) the whole document -----	1-16
A	US 5 043 013 A (KLUGER EDWARD W [US] ET AL) 27 August 1991 (1991-08-27) the whole document -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2012/039537

Patent document cited in search report	Publication date	Patent family member(s)	Publication date	
JP 2262663	A	25-10-1990	JP 2262663 A	25-10-1990
			JP 2670843 B2	29-10-1997

DE 2333828	A1	17-01-1974	AR 197409 A1	05-04-1974
			AU 468842 B2	22-01-1976
			AU 5698673 A	19-12-1974
			BE 801411 A1	26-12-1973
			CA 997754 A1	28-09-1976
			CH 575981 A5	31-05-1976
			DE 2333828 A1	17-01-1974
			ES 416529 A1	16-02-1976
			FR 2236900 A1	07-02-1975
			GB 1425029 A	18-02-1976
			IT 994896 B	20-10-1975
			JP 49063736 A	20-06-1974
			NL 7309179 A	07-01-1974

GB 1465941	A	02-03-1977	NONE	

EP 0474402	A2	11-03-1992	CA 2050716 A1	06-03-1992
			DE 69109980 D1	29-06-1995
			DE 69109980 T2	02-11-1995
			EP 0474402 A2	11-03-1992
			ES 2072552 T3	16-07-1995
			JP 5032904 A	09-02-1993
			US 5290630 A	01-03-1994

US 2008177090	A1	24-07-2008	CN 101600760 A	09-12-2009
			EP 2104709 A1	30-09-2009
			JP 2010516907 A	20-05-2010
			US 2008177090 A1	24-07-2008
			US 2010025633 A1	04-02-2010
			US 2011289698 A1	01-12-2011
			US 2012178665 A1	12-07-2012
			WO 2008091524 A1	31-07-2008

WO 2011017719	A2	10-02-2011	NONE	

DE 102007040928	A1	05-03-2009	DE 102007040928 A1	05-03-2009
			WO 2009030344 A2	12-03-2009

US 2003212172	A1	13-11-2003	AU 2003228715 A1	11-11-2003
			US 2003212172 A1	13-11-2003
			WO 03095546 A1	20-11-2003

US 5043013	A	27-08-1991	NONE	
