



- (51) **International Patent Classification:**
C07D 487/04 (2006.01) A61P 31/12 (2006.01)
A61K 31/519 (2006.01)
- (21) **International Application Number:**
PCT/US20 12/07 1065
- (22) **International Filing Date:**
20 December 2012 (20.12.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/579,625 22 December 2011 (22.12.2011) US
61/618,510 30 March 2012 (30.03.2012) US
- (71) **Applicant (for all designated States except US):** GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, California 94404 (US).
- (72) **Inventor; and**
(71) **Applicant (for US only):** BOOJAMRA, Constantine G. [US/US]; 1 Worth Street, San Francisco, California 94114 (US).
- (72) **Inventor; and**
(71) **Applicant :** HUI, Hon Chung [US/US]; 853 Woodside Way, #229, San Mateo, California 94401 (US).
- (72) **Inventors; and**
(71) **Applicants (for US only):** JANSA, Petr [CZ/US]; 3952 Casanova Drive, San Mateo, California 94403 (US). MACKMAN, Richard L. [GB/US]; 360 Ashton Avenue, Millbrae, California 94030 (US). PARRISH, Jay, P. [US/US]; 720 Bair Island Road #106, Redwood City, California 94063 (US).
- (72) **Inventor; and**
(71) **Applicant :** SANGI, Michael [US/US]; 5507 College Ave., Apt. #7, Oakland, California 94618 (US).
- (72) **Inventors; and**
(71) **Applicants (for US only):** SIEGEL, Dustin [US/US]; 758 Bounty Dr. Apt. 5803, Foster City, California 94404 (US). SPERANDIO, David [US/US]; 3804 Nathan Way, Palo Alto, California 94303 (US).
- (72) **Inventor; and**
(71) **Applicant :** YANG, Hai [US/US]; 32 Jody Court, San Mateo, California 94402 (US).
- (74) **Agents:** MICKELSON, John, W. et al; VIKSNINS HARRIS & PADYS PLLP, 7900 International Drive, Suite 670, Bloomington, Minnesota 55425 (US).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report (Art. 21(3))



5 PYRAZOLO[1,5-A]PYRIMIDINES AS ANTIVIRAL AGENTS

Cross Reference to Related Applications

This patent application claims the benefit of priority of U.S. Application Serial No. 61/579625, filed December 22, 2011 and of U.S. application serial No. 61/618510, filed March 10 30, 2012. The content of each of these provisional applications is hereby incorporated herein in its entirety.


Background

Pneumovirinae viruses are negative-sense, single-stranded, RNA viruses that are 15 responsible for many prevalent human and animal diseases. The *Pneumovirinae* sub-family of viruses is a part of the family *Paramyxoviridae* and includes human respiratory syncytial virus (HRSV). Almost all children will have had an HRSV infection by their second birthday. HRSV is the major cause of lower respiratory tract infections in infancy and childhood with 0.5% to 2% of those infected requiring hospitalization. The elderly and adults with chronic heart, lung 20 disease or those that are immunosuppressed also have a high risk for developing severe HRSV disease (<http://www.cdc.gov/rsv/index.html>). No vaccine to prevent HRSV infection is currently available. The monoclonal antibody palivizumab is available for immunoprophylaxis, but its use is restricted to infants at high risk, e.g., premature infants or those with either congenital heart or lung disease, and the cost for general use is often prohibitive. In addition, 25 nucleoside analog ribavirin has been approved as the only antiviral agent to treat HRSV infections but has limited efficacy. Therefore, there is a need for *anti-Pneumovirinae* therapeutics.

Summary

30 Provided herein are methods and compounds for the treatment of infections caused by the *Pneumovirinae* virus family.

Accordingly, one embodiment provides a compound of formula I:



5

or a salt or ester, thereof;

wherein:

A is $-(C(R^4)_2)_n-$ wherein any one $C(R^4)_2$ of said $-(C(R^4)_2)_n-$ may be optionally replaced with $-O-$, $-S-$, $-S(O)_p-$, NH or NR^a ;

10

n is 3, 4, 5 or 6;

each p is 1 or 2;

Ar is a C_2-C_{20} heterocyclyl group or a C_6-C_{20} aryl group, wherein the C_2-C_{20} heterocyclyl group or the C_6-C_{20} aryl group is optionally substituted with 1, 2, 3, 4 or 5 R^6 ;

each R^3 , R^4 and R^6 is independently H, oxo, OR^{11} , $NR^{11}R^{12}$, $NR^{11}C(O)R^{11}$,

15

$NR^{11}C(O)OR^{11}$, $NR^{11}C(O)NR^{11}R^{12}$, N_3 , CN , NO_2 , SR^{11} , $S(O)_pR^a$, $NR^{11}S(O)_pR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)_p(OR^{11})$, $-SO_2NR^{11}R^{12}$, $-NR^{11}S(O)_p(OR^{11})$, $NR^{11}SO_pNR^{11}R^{12}$, $NR^{11}C(=NR^{11})NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, C_6-C_{20} aryl, C_2-C_{20} heterocyclyl, (C_3-C_7) cycloalkyl or (C_4-C_8) carbocyclalkyl;

20

or two R^4 on adjacent carbon atoms, when taken together, may optionally form a double bond between the two carbons to which they are attached or may form a (C_3-C_7) cycloalkyl ring wherein one carbon atom of said (C_3-C_7) cycloalkyl ring may be optionally replaced by $-O-$, $-S-$, $-S(O)_p-$, $-NH-$ or $-NR^a$;

25

or four R^4 on adjacent carbon atoms, when taken together, may optionally form an optionally substituted C_6 aryl ring;

or two R^4 on the same carbon atom, when taken together, may optionally form a (C_3-C_7) cycloalkyl ring wherein one carbon atom of said (C_3-C_7) cycloalkyl ring may be optionally replaced by $-O-$, $-S-$, $-S(O)_p-$, $-NH-$ or $-NR^a$;

30


or two R^6 on adjacent carbon atoms, when taken together, may optionally form a (C_3-C_7) cycloalkyl ring wherein one carbon atom of said (C_3-C_7) cycloalkyl ring may be optionally replaced by $-O-$, $-S-$, $-S(O)_p-$, $-NH-$ or $-NR^a$;

5 each R^a is independently (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl wherein any (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀heterocyclyl, and wherein any aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R^a is optionally substituted with one or more (e.g. 1, 2, 3, 4 or 5) OH, NH₂, CO₂H, C₂-C₂₀heterocyclyl or (C₁-C₈)alkyl;

10 each R¹¹ or R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclylalkyl, -C(=O)R^a, -S(O)_pR^a or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S-, -S(O)_p-, -NH-, -NR^a- or -C(O)-; and

15 wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(d-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of each R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more (e.g. 1, 2, 3, 4 or 5) oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(O)_pR^a, OR^a, (d-C₈)alkyl, (C₁-C₈)haloalkyl, -C(O)R^a, -C(O)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(O)_pR^a, NR^aS(O)_pR^a, NHC(O)R^a, NR^aC(O)R^a, NHC(O)OR^a, NR^aC(O)OR^a, NR^aC(O)NHR^a, NR^aC(O)N(R^a)₂, NR^aC(O)NH₂, NHC(O)NHR^a, NHC(O)N(R^a)₂, NHC(O)NH₂, =NH, =NOH, =NOR^a, NR^aS(O)_pNHR^a, NR^aS(O)_pN(R^a)₂, NR^aS(O)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(O)(OH)₂ or R^a;

provided the compound is not:



Another embodiment provides a compound of formulas 1-11 (i.e., compounds 1-11), as described in examples 1 and 2, or a salt or ester thereof.

5 Another embodiment provides a compound of formula I (e.g., compounds 112-209) or a stereoisomer (e.g. enantiomer, diastereomer, atropisomer) or a salt or ester thereof.

Another embodiment provides a compound of formulas 1-1 11 or a stereoisomer (e.g., enantiomer, diastereomer, atropisomer) or a salt or ester thereof.

10 Another embodiment provides a pharmaceutical composition comprising a compound of formula I or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Another embodiment provides a pharmaceutical composition comprising a compound of formulas 1-1 11 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

15 Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt or ester thereof.

20 Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a compound of formulas 1-1 11, or a pharmaceutically acceptable salt or ester thereof.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of a compound of formula I, or a pharmaceutically acceptable salt or ester thereof.

25 Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of formulas 1-1 11, or a pharmaceutically acceptable salt or ester thereof.

30 Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a compound of formula I, or a pharmaceutically acceptable salt or ester thereof.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically

5 effective amount of a compound of formulas 1-1 11, or a pharmaceutically acceptable salt or ester thereof.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g., a human) in need thereof comprising administering a therapeutically effective amount of a tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or
10 solvate of a compound of formula I, or a pharmaceutically acceptable salt or ester thereof.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof comprising administering a therapeutically effective amount of a tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of formulas 1-1 11, or a pharmaceutically acceptable salt or ester thereof.

15 Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof comprising administering a therapeutically effective amount of a compound of formula I or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable diluent or carrier.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a
20 mammal (e.g. a human) in need thereof comprising administering a therapeutically effective amount of a compound of formulas 1-1 11 or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable diluent or carrier.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof comprising administering a therapeutically effective
25 amount of a compound of formula I or a pharmaceutically acceptable salt or ester thereof, in combination with at least one additional therapeutic agent.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof comprising administering a therapeutically effective amount of a compound of formulas 1-1 11 or a pharmaceutically acceptable salt or ester thereof,
30 in combination with at least one additional therapeutic agent.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

a) a first pharmaceutical composition comprising a compound of formula I or a
35 pharmaceutically acceptable salt or ester thereof; and

- 5 b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious *Pneumovirinae* viruses.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 10 a) a first pharmaceutical composition comprising a compound of formulas 1-1 11 or a pharmaceutically acceptable salt or ester thereof; and
- b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious *Pneumovirinae* viruses.

Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 15 a) a therapeutic agent selected from a compound a of formula I and pharmaceutically acceptable salts and esters thereof; and
- b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

20 Another embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a therapeutic agent selected from a compound of formulas 1-1 11 and pharmaceutically acceptable salts and esters thereof; and
- 25 b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 30 a) a first pharmaceutical composition comprising a compound of formula I or a pharmaceutically acceptable salt or ester thereof; and
- b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious respiratory syncytial viruses.

5 Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a first pharmaceutical composition comprising a compound of formulas 1-111 or a pharmaceutically acceptable salt or ester thereof; and
- 10 b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious respiratory syncytial viruses.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 15 a) a therapeutic agent selected from a compound of formula I and pharmaceutically acceptable salts and esters thereof; and
- b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

Another embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, comprising administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 20 a) a therapeutic agent selected from a compound of formulas 1-111 and pharmaceutically acceptable salts and esters thereof; and
- b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

Another embodiment provides a compound of formula I or a pharmaceutically acceptable salt or ester thereof, for use in medical therapy.

Another embodiment provides a compound of formulas 1-111 or a pharmaceutically acceptable salt or ester thereof for use in medical therapy.

Another embodiment provides a compound of formula I or a pharmaceutically acceptable salt or ester thereof, for use in the prophylactic or therapeutic treatment of a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

Another embodiment provides a compound of formulas 1-111 or a pharmaceutically acceptable salt or ester thereof, for use in the prophylactic or therapeutic treatment of a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

Another embodiment provides the use of a compound of formula I or a pharmaceutically acceptable salt or ester thereof, for the manufacture of a medicament useful for the treatment of

5 a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus in a mammal (e.g. a human).

Another embodiment provides the use of a compound of formulas 1-111 or a pharmaceutically acceptable salt or ester thereof for the manufacture of a medicament useful for the treatment of a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial
10 virus in a mammal (e.g. a human).

Another embodiment provides processes and novel intermediates disclosed herein which are useful for preparing a compound of formula I or a compound of formulas 1-111.

Another embodiment provides novel methods for synthesis, analysis, separation, isolation, purification, characterization, and testing of the compounds as disclosed herein.
15

Detailed Description

Definitions

Unless stated otherwise, the following terms and phrases as used herein are intended to have the following meanings:

20 When trade names are used herein, applicants intend to independently include the tradename product and the active pharmaceutical ingredient(s) of the tradename product.

The term "alkyl" refers to a straight or branched hydrocarbon. For example, an alkyl group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ alkyl), 1 to 8 carbon atoms (*i.e.*, C₁-C₈ alkyl), or 1 to 6 carbon atoms (*i.e.*, C₁-C₆ alkyl). Examples of suitable alkyl groups include, but are not
25 limited to, methyl (Me, -CH₃), ethyl (Et, -CH₂CH₃), 1-propyl (n-Pr, n-propyl, -CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, -CH(CH₃)₂), 1-butyl (n-Bu, n-butyl, -CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (i-Bu, i-butyl, -CH₂CH(CH₃)₂), 2-butyl (s-Bu, s-butyl, -CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (t-Bu, t-butyl, -C(CH₃)₃), 1-pentyl (n-pentyl, -CH₂CH₂CH₂CH₂CH₃), 2-pentyl (-CH(CH₃)CH₂CH₂CH₃), 3-pentyl (-CH(CH₂CH₃)₂), 2-methyl-2-butyl
30 (-C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (-CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (-CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (-CH₂CH(CH₃)CH₂CH₃), 1-hexyl (-CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (-CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (-CH(CH₂CH₃)(CH₂CH₂CH₃)), 2-methyl-2-pentyl (-C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-pentyl (-CH(CH₃)CH(CH₃)CH₂CH₃), 4-methyl-2-pentyl (-CH(CH₃)CH₂CH(CH₃)₂), 3-methyl-3-pentyl
35 (-C(CH₃)(CH₂CH₃)₂), 2-methyl-3-pentyl (-CH(CH₂CH₃)CH(CH₃)₂), 2,3-dimethyl-2-butyl

5 (-C(CH₃)₂CH(CH₃)₂), 3,3-dimethyl-2-butyl (-CH(CH₃)C(CH₃)₃), and octyl (-CH₂)₇CH₃).

The term "alkoxy" refers to a group having the formula -O-alkyl, in which an alkyl group, as defined above, is attached to the parent molecule via an oxygen atom. The alkyl portion of an alkoxy group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ alkoxy), 1 to 12 carbon atoms (*i.e.*, C₁-C₁₂ alkoxy), or 1 to 6 carbon atoms (*i.e.*, C₁-C₆ alkoxy). Examples of suitable alkoxy groups include, but are not limited to, methoxy (-O-CH₃ or -OMe), ethoxy (-OCH₂CH₃ or -OEt), t-butoxy (-O-C(CH₃)₃ or -OtBu) and the like.

The term "haloalkyl" refers to an alkyl group, as defined above, in which one or more hydrogen atoms of the alkyl group is replaced with a halogen atom. The alkyl portion of a haloalkyl group can have 1 to 20 carbon atoms (*i.e.*, C₁-C₂₀ haloalkyl), 1 to 12 carbon atoms (*i.e.*, C₁-C₁₂ haloalkyl), or 1 to 6 carbon atoms (*i.e.*, C₁-C₆ haloalkyl). Examples of suitable haloalkyl groups include, but are not limited to, -CF₃, -CHF₂, -CFH₂, -CH₂CF₃, and the like.

The term "alkenyl" refers to a straight or branched hydrocarbon with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp*² double bond. For example, an alkenyl group can have 2 to 20 carbon atoms (*i.e.*, C₂-C₂₀ alkenyl), 2 to 8 carbon atoms (*i.e.*, C₂-C₈ alkenyl), or 2 to 6 carbon atoms (*i.e.*, C₂-C₆ alkenyl). Examples of suitable alkenyl groups include, but are not limited to, ethylene or vinyl (-CH=CH₂), allyl (-CH₂CH=CH₂), cyclopentenyl (-C₅H₇), and 5-hexenyl (-CH₂CH₂CH₂CH₂CH=CH₂).

The term "alkynyl" refers to a straight or branched hydrocarbon with at least one site of unsaturation, *i.e.* a carbon-carbon, *sp* triple bond. For example, an alkynyl group can have 2 to 20 carbon atoms (*i.e.*, C₂-C₂₀ alkynyl), 2 to 8 carbon atoms (*i.e.*, C₂-C₈ alkyne), or 2 to 6 carbon atoms (*i.e.*, C₂-C₆ alkynyl). Examples of suitable alkynyl groups include, but are not limited to, acetylenic (-C≡CH), propargyl (-CH₂C≡CH), and the like.

The term "halogen" or "halo" refers to F, Cl, Br, or I.

The term "aryl" refers to an aromatic hydrocarbon radical derived by the removal of one hydrogen atom from a single carbon atom of a parent aromatic ring system. For example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 10 carbon atoms. Typical aryl groups include, but are not limited to, radicals derived from benzene (e.g., phenyl), substituted benzene, naphthalene, anthracene, biphenyl, and the like.


The term "arylalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or *sp*³ carbon atom, is replaced with an aryl

5 radical. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like. The arylalkyl group can comprise 7 to 20 carbon atoms, *e.g.*, the alkyl moiety is 1 to 6 carbon atoms and the aryl moiety is 6 to 14 carbon atoms.

10 The term "prodrug" as used herein refers to any compound that when administered to a biological system generates the drug substance, *i.e.*, active ingredient, as a result of spontaneous chemical reaction(s), enzyme catalyzed chemical reaction(s), photolysis, and/or metabolic chemical reaction(s). A prodrug is thus a covalently modified analog or latent form of a therapeutically active compound.


15 One skilled in the art will recognize that substituents and other moieties of the compounds of formula I should be selected in order to provide a compound which is sufficiently stable to provide a pharmaceutically useful compound which can be formulated into an acceptably stable pharmaceutical composition. Compounds of formula I which have such stability are contemplated as falling within the scope of the present invention.

20 The term "heterocycle" or "heterocyclyl" as used herein includes by way of example and not limitation those heterocycles described in Paquette, Leo A.; Principles of Modern Heterocyclic Chemistry (W.A. Benjamin, New York, 1968), particularly Chapters 1, 3, 4, 6, 7, and 9; The Chemistry of Heterocyclic Compounds, A Series of Monographs (John Wiley & Sons, New York, 1950 to present), in particular Volumes 13, 14, 16, 19, and 28; and *J. Am. Chem. Soc.* (1960) 82:5566. In one specific embodiment of the invention "heterocycle" includes a "carbocycle" as defined herein, wherein one or more (*e.g.* 1, 2, 3, or 4) carbon atoms have been replaced with a heteroatom (*e.g.* O, N, or S). The terms "heterocycle" or "heterocyclyl" includes saturated rings, partially unsaturated rings, and aromatic rings (*i.e.*, heteroaromatic rings). Substituted heterocyclyls include, for example, heterocyclic rings substituted with any of the substituents disclosed herein including carbonyl groups. A non-limiting example of a
30 carbonyl substituted heterocyclyl is:



Examples of heterocycles include by way of example and not limitation pyridyl,

5 dihydropyridyl, tetrahydropyridyl (piperidyl), thiazolyl, tetrahydrothiophenyl, sulfur oxidized tetrahydrothiophenyl, pyrimidinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, tetrazolyl, benzofuranyl, thianaphthalenyl, indolyl, indolenyl, quinolinyl, isoquinolinyl, benzimidazolyl, piperidinyl, 4-piperidonyl, pyrrolidinyl, 2-pyrrolidonyl, pyrrolinyl, tetrahydrofuranlyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, octahydroisoquinolinyl, 10 azocinyl, triazinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, thienyl, thianthrenyl, pyranlyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxathinyl, 2H-pyrrolyl, isothiazolyl, isoxazolyl, pyrazinyl, pyridazinyl, indolizinyl, isoindolyl, 3H-indolyl, 1H-indazolyl, purinyl, 4H-quinolizinyl, phthalazinyl, naphthyridinyl, quinoxalynyl, quinazolynyl, cinnolynyl, pteridinyl, 4aH-carbazolyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, pyrimidinyl, 15 phenanthrolinyl, phenazinyl, phenothiazinyl, furazanyl, phenoxazinyl, isochromanyl, chromanyl, imidazolidinyl, imidazolynyl, pyrazolidinyl, pyrazolynyl, piperazinyl, indolynyl, isoindolynyl, quinuclidinyl, morpholinyl, oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolynyl, isatinoyl, and bis-tetrahydrofuranlyl:



20 By way of example and not limitation, carbon bonded heterocycles are bonded at position 2, 3, 4, 5, or 6 of a pyridine, position 3, 4, 5, or 6 of a pyridazine, position 2, 4, 5, or 6 of a pyrimidine, position 2, 3, 5, or 6 of a pyrazine, position 2, 3, 4, or 5 of a furan, tetrahydrofuran, thiofuran, thiophene, pyrrole or tetrahydropyrrole, position 2, 4, or 5 of an oxazole, imidazole or thiazole, position 3, 4, or 5 of an isoxazole, pyrazole, or isothiazole, 25 position 2 or 3 of an aziridine, position 2, 3, or 4 of an azetidine, position 2, 3, 4, 5, 6, 7, or 8 of a quinoline or position 1, 3, 4, 5, 6, 7, or 8 of an isoquinoline. Still more typically, carbon bonded heterocycles include 2-pyridyl, 3-pyridyl, 4-pyridyl, 5-pyridyl, 6-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl, 6-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl, 2-pyrazinyl, 3-pyrazinyl, 5-pyrazinyl, 6-pyrazinyl, 2-thiazolyl, 4-thiazolyl, or 5- 30 thiazolyl.

By way of example and not limitation, nitrogen bonded heterocycles are bonded at position 1 of an aziridine, azetidine, pyrrole, pyrrolidine, 2-pyrroline, 3-pyrroline, imidazole,

5 imidazolidine, 2-imidazoline, 3-imidazoline, pyrazole, pyrazoline, 2-pyrazoline, 3-pyrazoline, piperidine, piperazine, indole, indoline, 1H-indazole, position 2 of a isoindole, or isoindoline, position 4 of a morpholine, and position 9 of a carbazole, or β -carboline. Still more typically, nitrogen bonded heterocycles include 1-aziridyl, 1-azetetyl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, and 1-piperidinyl.

10 The term "heteroaryl" refers to an aromatic heterocyclyl having at least one heteroatom in the ring. Non-limiting examples of suitable heteroatoms which can be included in the aromatic ring include oxygen, sulfur, and nitrogen. Non-limiting examples of heteroaryl rings include all of those aromatic rings listed in the definition of "heterocyclyl", including pyridinyl, pyrrolyl, oxazolyl, indolyl, isoindolyl, purinyl, furanyl, thienyl, benzofuranyl, benzothiophenyl, 15 carbazolyl, imidazolyl, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, quinolyl, isoquinolyl, pyridazyl, pyrimidyl, pyrazyl, etc.

The term "carbocycle" or "carbocyclyl" refers to a saturated (i.e., cycloalkyl), partially unsaturated (e.g., cycloalkenyl, cycloalkadienyl, etc.) or aromatic ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic carbocycles have 3 to 7 ring atoms, still more typically 5 or 6 ring 20 atoms. Bicyclic carbocycles have 7 to 12 ring atoms, e.g., arranged as a bicyclo [4,5], [5,5], [5,6] or [6,6] system, or 9 or 10 ring atoms arranged as a bicyclo [5,6] or [6,6] system, or spiro-fused rings. Non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, and phenyl. Non-limiting examples of bicyclo carbocycles includes naphthyl, tetrahydronaphthalene, and decaline.

The term "cycloalkyl" refers to a saturated or partially unsaturated ring having 3 to 7 carbon atoms as a monocycle, 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic cycloalkyl groups have 3 to 7 ring atoms, still more typically 30 5 or 6 ring atoms. Bicyclic cycloalkyl groups have 7 to 12 ring atoms, e.g., arranged as a bicyclo (4,5), (5,5), (5,6) or (6,6) system, or 9 or 10 ring atoms arranged as a bicyclo (5,6) or (6,6) system. Cycloalkyl groups include hydrocarbon mono-, bi-, and poly-cyclic rings, whether fused, bridged, or spiro. Non-limiting examples of monocyclic carbocycles include cyclopropyl, cyclobutyl, cyclopentyl, 1-cyclopent-1-enyl, 1-cyclopent-2-enyl, 1-cyclopent-3-enyl, 35 cyclohexyl, 1-cyclohex-1-enyl, 1-cyclohex-2-enyl, 1-cyclohex-3-enyl, bicyclo[3.1.0]hex-6-yl

5 and the like.

The term "carbocyclalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom is replaced with a carbocyclyl radical as described herein. Typical, but non-limiting, examples of carbocyclalkyl groups include cyclopropylmethyl, cyclopropylethyl, cyclobutylmethyl, cyclopentylmethyl and
10 cyclohexylmethyl.

Selected substituents comprising the compounds of formula I may be present to a recursive degree. In this context, "recursive substituent" means that a substituent may recite another instance of itself. The multiple recitations may be direct or indirect through a sequence of other substituents. Because of the recursive nature of such substituents, theoretically, a large
15 number of compounds may be present in any given embodiment. One of ordinary skill in the art of medicinal chemistry understands that the total number of such substituents is reasonably limited by the desired properties of the compound intended. Such properties include, by way of example and not limitation, physical properties such as molecular weight, solubility or log P, application properties such as activity against the intended target, and practical properties such
20 as ease of synthesis. Recursive substituents may be an intended aspect of the invention. One of ordinary skill in the art of medicinal chemistry understands the versatility of such substituents. To the degree that recursive substituents are present in an embodiment of the invention, they may recite another instance of themselves, 0, 1, 2, 3, or 4 times.

"Protecting group" refers to a moiety of a compound that masks or alters the properties
25 of a functional group or the properties of the compound as a whole. The chemical substructure of a protecting group varies widely. One function of a protecting group is to serve as an intermediate in the synthesis of the parental drug substance. Chemical protecting groups and strategies for protection/deprotection are well known in the art. See: "Protective Groups in Organic Chemistry", Theodora W. Greene (John Wiley & Sons, Inc., New York, 1991).

30 Protecting groups are often utilized to mask the reactivity of certain functional groups, to assist in the efficiency of desired chemical reactions, e.g. making and breaking chemical bonds in an ordered and planned fashion. Protection of functional groups of a compound alters other physical properties besides the reactivity of the protected functional group, such as the polarity, lipophilicity (hydrophobicity), and other properties which can be measured by common
35 analytical tools. Chemically protected intermediates may themselves be biologically active or

5 inactive.

Protected compounds may also exhibit altered, and in some cases, optimized properties *in vitro* and *in vivo*, such as passage through cellular membranes and resistance to enzymatic degradation or sequestration. In this role, protected compounds with intended therapeutic effects may be referred to as prodrugs. Another function of a protecting group is to convert the parental
10 drug into a prodrug, whereby the parental drug is released upon conversion of the prodrug *in vivo*. Because active prodrugs may be absorbed more effectively than the parental drug, prodrugs may possess greater potency *in vivo* than the parental drug. Protecting groups are removed either *in vitro*, in the instance of chemical intermediates, or *in vivo*, in the case of prodrugs. With chemical intermediates, it is not particularly important that the resulting
15 products after deprotection, e.g. alcohols, be physiologically acceptable, although in general it is more desirable if the products are pharmacologically innocuous.

"Prodrug moiety" means a labile functional group which separates from the active inhibitory compound during metabolism, systemically, inside a cell, by hydrolysis, enzymatic cleavage, or by some other process (Bundgaard, Hans, "Design and Application of Prodrugs" in
20 Textbook of Drug Design and Development (1991), P. Krosgaard-Larsen and H. Bundgaard, Eds. Harwood Academic Publishers, pp. 113-191). Enzymes which are capable of an enzymatic activation mechanism with, for example any phosphate or phosphonate prodrug compounds of the invention, include but are not limited to, amidases, esterases, microbial enzymes, phospholipases, cholinesterases, and phosphases. Prodrug moieties can serve to enhance
25 solubility, absorption and lipophilicity to optimize drug delivery, bioavailability and efficacy. A prodrug moiety may include an active metabolite or drug itself.

It is to be noted that all tautomers, atropisomers, polymorphs, pseudopolymorphs of compounds disclosed herein and pharmaceutically acceptable salts and esters thereof are embraced by the present invention.

30 A compound disclosed herein and its pharmaceutically acceptable salts may exist as different polymorphs or pseudopolymorphs. As used herein, crystalline polymorphism means the ability of a crystalline compound to exist in different crystal structures. The crystalline polymorphism may result from differences in crystal packing (packing polymorphism) or differences in packing between different conformers of the same molecule (conformational
35 polymorphism). As used herein, crystalline pseudopolymorphism means the ability of a hydrate

5 or solvate of a compound to exist in different crystal structures. The pseudopolymorphs of the instant invention may exist due to differences in crystal packing (packing pseudopolymorphism) or due to differences in packing between different conformers of the same molecule (conformational pseudopolymorphism). The instant invention comprises all polymorphs and pseudopolymorphs of the compounds disclosed herein

10 A compound disclosed herein and its pharmaceutically acceptable salts may also exist as an amorphous solid. As used herein, an amorphous solid is a solid in which there is no long-range order of the positions of the atoms in the solid. This definition applies as well when the crystal size is two nanometers or less. Additives, including solvents, may be used to create the amorphous forms of the instant invention. The instant invention comprises all amorphous forms
15 of the compounds disclosed herein and their pharmaceutically acceptable salts.

The modifier "about" used in connection with a quantity is inclusive of the stated value and has the meaning dictated by the context (e.g., includes the degree of error associated with measurement of the particular quantity).

20 The term "treating", as used herein, unless otherwise indicated, means reversing, alleviating, inhibiting the progress of, or preventing the disorder or condition to which such term applies, or one or more symptoms of such disorder or condition. The term "treatment", as used herein, refers to the act of treating, as "treating" is defined immediately above.

The term "therapeutically effective amount", as used herein, is the amount of compound disclosed herein present in a composition described herein that is needed to provide a desired
25 level of drug in the secretions and tissues of the airways and lungs, or alternatively, in the bloodstream of a subject to be treated to give an anticipated physiological response or desired biological effect when such a composition is administered by the chosen route of administration. The precise amount will depend upon numerous factors, for example the particular disclosed herein, the specific activity of the composition, the delivery device employed, the physical
30 characteristics of the composition, its intended use, as well as patient considerations such as severity of the disease state, patient cooperation, etc., and can readily be determined by one skilled in the art and in reference to the information provided herein.

The term "normal saline" means a water solution containing 0.9% (w/v) NaCl.

35 The term "hypertonic saline" means a water solution containing greater than 0.9% (w/v) NaCl. For example, 3% hypertonic saline would contain 3% (w/v) NaCl.

5 Physiologically acceptable salts (e.g. pharmaceutically acceptable salt) of the compounds of the invention include salts derived from an appropriate base, such as an alkali metal or an alkaline earth (for example, Na^+ , Li^+ , K^+ , Ca^{+2} and $\text{Mg}^{+3/4}$ ammonium and NR^+)
10 Physiologically acceptable salts of a nitrogen atom or an amino group include (a) acid addition salts formed with inorganic acids, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, sulfamic acids, phosphoric acid, nitric acid and the like; (b) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid, malic acid, ascorbic acid, benzoic acid, isethionic acid, lactobionic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, benzenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, malonic acid, sulfosalicylic acid, glycolic acid, 2-hydroxy-3-naphthoate, pamoate, salicylic acid, stearic acid, phthalic acid, mandelic acid, lactic acid, ethanesulfonic acid, lysine, arginine, glutamic acid, glycine, serine, threonine, alanine, isoleucine, leucine and the like; and (c) salts formed from elemental anions for example, chlorine, bromine, and iodine. Physiologically acceptable salts of a compound of a hydroxy group include the anion of said compound in combination with a suitable cation such as Na^+ and NR_4^+ . Each R is independently selected from H and $(\text{C}_1\text{-C}_8)\text{alkyl}$.

For therapeutic use, salts of active ingredients of the compounds of the invention will be physiologically acceptable, i.e. they will be salts derived from a physiologically acceptable acid or base. However, salts of acids or bases which are not physiologically acceptable may also find use, for example, in the preparation or purification of a physiologically acceptable compound.
25 All salts, whether or not derived from a physiologically acceptable acid or base, are within the scope of the present invention.

It is to be understood that the compositions herein comprise compounds disclosed herein in their un-ionized, as well as zwitterionic form, and combinations with stoichiometric amounts of water as in hydrates.
30

Stereochemical definitions and conventions used herein generally follow S. P. Parker, Ed., McGraw-Hill Dictionary of Chemical Terms (1984) McGraw-Hill Book Company, New York; and Eliel, E. and Wilen, S., Stereochemistry of Organic Compounds (1994) John Wiley & Sons, Inc., New York. Many organic compounds exist in optically active forms, i.e., they have the ability to rotate the plane of plane-polarized light. In describing an optically active
35

5 compound, the prefixes D and L or R and S are used to denote the absolute configuration of the molecule about its chiral center(s). The prefixes d and l, D and L, or (+) and (-) are employed to designate the sign of rotation of plane-polarized light by the compound, with S, (-), or l meaning that the compound is levorotatory while a compound prefixed with R, (+), or d is dextrorotatory. For a given chemical structure, these stereoisomers are identical except that they are mirror
10 images of one another. A specific stereoisomer may also be referred to as an enantiomer, and a mixture of such isomers is often called an enantiomeric mixture. A 50:50 mixture of enantiomers is referred to as a racemic mixture or a racemate, which may occur where there has been no stereoselection or stereospecificity in a chemical reaction or process. The terms "racemic mixture" and "racemate" refer to an equimolar mixture of two enantiomeric species,
15 devoid of optical activity.

The compounds disclosed herein have chiral centers, e.g. chiral carbon. The compounds of the invention include enriched or resolved optical isomers at any or all asymmetric, chiral atoms. In other words, the chiral centers apparent from the depictions are provided as the chiral isomers. Individual enantiomers or diastereomers, isolated or synthesized, substantially free of
20 their enantiomeric or diastereomeric partners, are all within the scope of the invention. The stereoisomeric mixtures can be separated into their individual, substantially optically pure isomers through well-known techniques such as, for example, the separation of diastereomeric salts formed with optically active adjuncts, e.g., acids or bases followed by conversion back to the optically active substances. Typically, the desired optical isomer is synthesized by means of
25 stereospecific reactions, beginning with the appropriate stereoisomer of the desired starting material.

The term "chiral" refers to molecules which have the property of non-superimposability of the mirror image partner, while the term "achiral" refers to molecules which are superimposable on their mirror image partner.

30 The term "stereoisomers" refers to compounds which have identical chemical constitution, but differ with regard to the arrangement of the atoms or groups in space.


"Diastereomer" refers to a stereoisomer with two or more centers of chirality and whose molecules are not mirror images of one another. Diastereomers have different physical properties, e.g. melting points, boiling points, spectral properties, and reactivities. Mixtures of
35 diastereomers may separate under high resolution analytical procedures such as electrophoresis

5 and chromatography. Enantiomers" refer to two stereoisomers of a compound which are non-superimposable mirror images of one another.

It is to be understood that for compounds disclosed herein when a bond is drawn in a non-stereochemical manner (e.g. flat) the atom to which the bond is attached includes all stereochemical possibilities. It is also to be understood that when a bond is drawn in a
 10 stereochemical manner (e.g. bold, bold-wedge, dashed or dashed-wedge) the atom to which the stereochemical bond is attached has the stereochemistry as shown unless otherwise noted.

Accordingly, in one embodiment, the compounds disclosed herein are greater than 50% a single enantiomer. In another embodiment, the compounds disclosed herein are at least 80% a single enantiomer. In another embodiment, the compounds disclosed herein are at least 90% a
 15 single enantiomer. In another embodiment, the compounds disclosed herein are at least 98% a single enantiomer. In another embodiment, the compounds disclosed herein are at least 99% a single enantiomer. In another embodiment, the compounds disclosed herein are greater than 50% a single diastereomer. In another embodiment, the compounds disclosed herein are at least 80% a single diastereomer. In another embodiment, the compounds disclosed herein are at least
 20 90% a single diastereomer. In another embodiment, the compounds disclosed herein are at least 98% a single diastereomer. In another embodiment, the compounds disclosed herein are at least 99% a single diastereomer.


In one embodiment compounds are represented by formula I and Ia (and salts and esters, thereof) as shown below wherein one position of chirality is marked with an asterisk.



The stereochemistry at the carbon marked with an asterisk as shown above for formula I is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk
 30 carbon following the Cahn-Ingold-Prelog system (March, J., Advanced Organic Chemistry, 4th Edition, John Wiley and Sons, pages 109-111). The stereochemistry at the carbon marked with

5 an asterisk as shown above for formula Ia is the (S) stereochemistry. In one embodiment, the compounds of formula I and Ia are greater than 50% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula I and Ia are at least 60% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula I and Ia are at least 70% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula I and Ia are at least 80% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula I and Ia are at least 90% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula I and Ia are at least 95% a single stereoisomer at the asterisk position.


15 In one embodiment compounds are represented by formula I and Ia (and salts and esters, thereof) as shown below wherein three positions of chirality are marked with an asterisk.



20 The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula I is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula Ia is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 2 (*2) as shown above for formula I and formula Ia is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 3 (*3) as shown above for formula I and formula Ia is the (R) stereochemistry. In one embodiment, the compounds of formula I and Ia are greater than 50% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula I and Ia are at least 60% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula I and Ia are at least 70% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula I and Ia are at least 80% a single


5 stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula I and Ia are at least 90% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds formula I and Ia are at least 95% a single stereoisomer at each of the asterisk positions.

10 In one embodiment compounds are represented by formulas II or IIa or salts or esters thereof, wherein one position of chirality is marked with an asterisk.



The stereochemistry at the carbon marked with an asterisk as shown above for formula II is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the carbon marked with an asterisk as shown above for formula IIa is the (S) stereochemistry. In one embodiment, the compounds of formula II and IIa are greater than 50% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula II and IIa are at least 60% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula II and IIa are at least 70% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula II and IIa are at least 80% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula II and IIa are at least 90% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula II and IIa are at least 95% a single stereoisomer at the asterisk position.

One embodiment provides compounds of formula II and IIa (and salts and esters, thereof) as shown below wherein three positions of chirality are marked with an asterisk.



5 The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula II is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the


10 carbon marked with an asterisk 1 (*1) as shown above for formula IIa is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 2 (*2) as shown above for formula II and formula IIa is the (R) stereochemistry. The stereochemistry at the carbon marked with an asterisk 3 (*3) as shown above for formula I and formula Ia is the (S) stereochemistry. In one embodiment, the compounds of formula II and IIa are greater than 50% a single stereoisomer at

15 each of the asterisk positions. In another embodiment, the compounds of formula II and IIa are at least 60% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula I and Ia are at least 70% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula II and IIa are at least 80% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of

20 formula II and IIa are at least 90% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula II and IIa are at least 95% a single stereoisomer at each of the asterisk positions.

One embodiment provides for compounds of formulas III or IIIa or salts or esters thereof. Compounds of formula III and IIIa are shown below wherein one position of chirality is

25 marked with an asterisk.




The stereochemistry at the carbon marked with an asterisk as shown above for formula III is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system or the (R)

30 stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the carbon marked

5 with an asterisk as shown above for formula IIia is the (S) stereochemistry. In one embodiment, the compounds of formula III and IIia are greater than 50% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula III and IIia are at least 60% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula III and IIia are at least 70% a single stereoisomer at the asterisk position. In another

10 embodiment, the compounds of formula III and IIia are at least 80% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula III and IIia are at least 90% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula III and IIia are at least 95% a single stereoisomer at the asterisk position.

15 One embodiment provides compounds of formula III and IIia (and salts and esters, thereof) as shown below wherein three positions of chirality are marked with an asterisk.




20 The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula III is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number two of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula IIia is the (S)

25 stereochemistry. The stereochemistry at the carbon marked with an asterisk 2 (*2) as shown above for formula III and formula IIia is the (R) stereochemistry. The stereochemistry at the carbon marked with an asterisk 3 (*3) as shown above for formula III and formula IIia is the (R) stereochemistry. In one embodiment, the compounds of the invention of formula III and IIia are greater than 50% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of the invention of formula III and IIia are at least 60% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of the invention of

30 formula III and IIia are at least 70% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of the invention of formula III and IIia are at least 80% a


5 single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of the invention of formula III and IIIa are at least 90% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of the invention of formula III and IIIa are at least 95% a single stereoisomer at each of the asterisk positions.

One embodiment provides for compounds of formulas IV or IVa or salts or esters thereof. Compounds of formula IV and IVa are shown wherein one position of chirality is marked with an asterisk.



The stereochemistry at the carbon marked with an asterisk as shown above for formula IV is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk carbon following the Cahn-Ingold-Prelog system. The stereochemistry at the carbon marked with an asterisk as shown above for formula IVa is the (S) stereochemistry. In one embodiment, the compounds of formula IV and IVa are greater than 50% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula IV and IVa are at least 60% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula IV and IVa are at least 70% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula IV and IVa are at least 80% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula IV and IVa are at least 90% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formula IV and IVa are at least 95% a single stereoisomer at the asterisk position.


One embodiment provides compounds of formula IV and IVa (and salts and esters, thereof) as shown below wherein three positions of chirality are marked with an asterisk.



5

The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula IV is the (S) stereochemistry provided that A is ranked the lowest (3) or highest (1) of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog system or the (R) stereochemistry provided that A is ranked number 2 of the three substituents of the asterisk 1 (*1) carbon following the Cahn-Ingold-Prelog System. The stereochemistry at the carbon marked with an asterisk 1 (*1) as shown above for formula IVa is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 2 (*2) as shown above for formula IV and formula IVa is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 3 (*3) as shown above for formula IV and formula IVa is the (S) stereochemistry. In one embodiment, the compounds of formula IV and IVa are greater than 50% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula IV and IVa are at least 60% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula IV and IVa are at least 70% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula IV and IVa are at least 80% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula IV and IVa are at least 90% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formula IV and IVa are at least 95% a single stereoisomer at each of the asterisk positions.

Each of the compounds of formulas 1-24 described herein have a core represented by formula V wherein a position of chirality is marked with an asterisk (wherein R represents an amine).




5

The stereochemistry at the carbon marked with an asterisk as shown for formula V is the (S) stereochemistry. In one embodiment, the compounds of formulas 1-24 are greater than 50% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formulas 1-24 are at least 60% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formulas 1-24 are at least 70% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formulas 1-24 are at least 80% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formulas 1-24 are at least 90% a single stereoisomer at the asterisk position. In another embodiment, the compounds of formulas 1-24 are at least 95% a single stereoisomer at the asterisk position.

10

Each of the compounds of formula 25-11 described herein have a core represented by formula VI wherein two positions of chirality are marked with an asterisk.



The stereochemistry at the carbon marked with an asterisk 1 (1) as shown for formula VI is the (S) stereochemistry. The stereochemistry at the carbon marked with an asterisk 2 (2) as shown for formula VI is the (S) stereochemistry. In one embodiment, the compounds of formulas 25-11 are greater than 50% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formulas 25-11 are at least 60% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formulas 25-11 are at least 70% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formulas 25-11 are at least 80% a single stereoisomer at each of the asterisk

25

5 positions. In another embodiment, the compounds of formulas 25-1 11 are at least 90% a single stereoisomer at each of the asterisk positions. In another embodiment, the compounds of formulas 25-1 11 are at least 95% a single stereoisomer at each of the asterisk positions.

The compounds disclosed herein also include molecules that incorporate isotopes of the atoms specified in the particular molecules. Non-limiting examples of these isotopes include D,
10 T, ^{14}C , ^{13}C and ^{15}N .

Whenever a compound described herein is substituted with more than one of the same designated group, e.g., "R" or "R¹", then it will be understood that the groups may be the same or different, i.e., each group is independently selected. Wavy lines, \sim , indicate the site of covalent bond attachments to the adjoining substructures, groups, moieties, or atoms.


15 The compounds of the invention can also exist as tautomeric isomers in certain cases. Although only one delocalized resonance structure may be depicted, all such forms are contemplated within the scope of the invention.

Detailed description of exemplary embodiments.

20 Reference will now be made in detail to certain embodiments, examples of which are illustrated in the accompanying description, structures and formulas. While the invention will be described in conjunction with the enumerated embodiments, it will be understood that they are not intended to limit the invention to those embodiments. On the contrary, the invention is intended to cover all alternatives, modifications, and equivalents, which may be included within
25 the full scope of the present invention as described herein.

Specific values listed below for radicals, substituents, and ranges, are for illustration only; they do not exclude other defined values or other values within defined ranges for the radicals and substituents. Specific values listed are values for compounds of formula I, II, III, IV as well as sub-formulas of these formulas (e.g. formula Ia, IIa, IIIa and IVa).

30 A specific group of compounds of formula I are compounds of formula Ia.



and salts and esters, thereof.

5 A specific value for R³ is H.

A specific value for n is 3.

A specific value for R⁴ is H.

A specific value for A is -(CH₂)₃-.

10 A specific value for Ar is phenyl, pyridyl, 1,2,3,4-tetrahydronaphthyl, indazolyl, 1,6-naphthyridyl, 2,3-dihydroindanyl, quinolyl, indolyl, 4H-benzo[d][1,3]dioxanyl, pyrazolo[1,5-a]pyridinyl, imidazo[1,2-a]pyridinyl, 1, 2,3,4-tetrahydroquinolyl, benzo[d][1,3]dioxolyl, quinoxalyl, isoquinolyl, naphthyl, thiophenyl, pyrazolyl, 4,5,6,7-tetrahydrobenzothiophenyl or pyrazolo[3,4,b]pyridinyl, wherein any phenyl, pyridyl, 1,2,3,4-tetrahydronaphthyl, indazolyl, 1,6-naphthyridyl, 2,3-dihydroindanyl, quinolyl, indolyl, 4H-benzo[d][1,3]dioxanyl, pyrazolo[1,5-a]pyridinyl, imidazo[1,2-a]pyridinyl, 1, 2,3,4-tetrahydroquinolyl, benzo[d][1,3]dioxolyl, quinoxalyl, isoquinolyl, naphthyl, thiophenyl, pyrazolyl, 4,5,6,7-tetrahydrobenzothiophenyl or pyrazolo[3,4,b]pyridinyl of A is optionally substituted with 1 to 5 R⁶.

20 A specific value for Ar is phenyl, monocyclic-heterocycle or bicyclic-heterocycle, wherein any phenyl, monocyclic-heterocycle or bicyclic-heterocycle of A is optionally substituted with 1 to 5 R⁶.

25 A specific value for Ar is phenyl, monocyclic-heterocycle or bicyclic-heterocycle, wherein the monocyclic-heterocycle or bicyclic-heterocycle consists of 1-10 carbon atoms and 1-5 heteroatoms within the ring system and wherein any phenyl, monocyclic-heterocycle or bicyclic-heterocycle of A is optionally substituted with 1 to 5 R⁶.

A specific value for Ar is phenyl optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups.

A specific value for Ar is phenyl optionally substituted by 1, 2, 3, or 4 R⁶ groups.

A specific value for Ar is phenyl optionally substituted by 1, 2, or 3 R⁶ groups.

A specific value for Ar is naphthyl optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups.

30 A specific value for Ar is naphthyl optionally substituted by 1, 2, 3, or 4 R⁶ groups.

A specific value for Ar is naphthyl optionally substituted by 1, 2, or 3 R⁶ groups.

35 A specific value for Ar is phenyl optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -SO₂NH(C₁-C₆ alkyl), -SO₂N(C₁-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or

5 two R⁶ groups on the phenyl ring together with the atoms to which they are attached form a benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is phenyl substituted by 1, 2, 3, or 4 R⁶ groups selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -
10 SO₂NH(d-C₆ alkyl), -SO₂N(d-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or two R⁶ groups on the phenyl ring together with the atoms to which they are attached form a benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is phenyl optionally substituted by 1, 2, or 3 R⁶ groups selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -
15 SO₂NH(d-C₆ alkyl), -SO₂N(d-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or two R⁶ groups on the phenyl ring together with the atoms to which they are attached form a benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is naphthyl optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups
20 selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -SO₂NH(d-C₆ alkyl), -SO₂N(d-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or two R⁶ groups on the phenyl ring together with the atoms to which they are attached form a benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is naphthyl optionally substituted by 1, 2, 3, or 4 R⁶ groups
25 selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -SO₂NH(Ci-C₆ alkyl), -SO₂N(d-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or two R⁶ groups on the phenyl ring together with the atoms to which they are attached form a
30 benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is naphthyl optionally substituted by 1, 2, or 3 R⁶ groups selected from halogen, alkyl, alkoxy, -CH₂OH, CF₃, -O-CF₃, -O-CF₂, -O-(C₃-C₆ cycloalkyl), -SO₂NH₂, -
SO₂NH(d-C₆ alkyl), -SO₂N(d-C₆ alkyl)₂, -SO₂-d-C₆ alkyl, -S(=O)-d-C₆ alkyl, an oxadiazole optionally substituted by CH₃, and a triazole ring optionally substituted by CH₃; or two R⁶ groups

5 on the phenyl ring together with the atoms to which they are attached form a benzodioxole, benzodioxine, or dihydroindene ring.

A specific value for Ar is pyridine optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups.

A specific value for Ar is pyridine optionally substituted by 1, 2, 3, or 4 R⁶ groups.

A specific value for Ar is pyridine optionally substituted by 1, 2, or 3 R⁶ groups.

10 A specific value for Ar is pyridine optionally substituted by 1, 2, 3, 4, or 5 R⁶ groups independently selected from halogen, alkyl, alkoxy, -CH₂OH, phenyl and CF₃.

A specific value for Ar is pyridine optionally substituted by 1, 2, 3, or 4 R⁶ groups independently selected from halogen, alkyl, alkoxy, -CH₂OH, phenyl and CF₃.

15 A specific value for Ar is pyridine optionally substituted by 1, 2, or 3 R⁶ groups independently selected from halogen, alkyl, alkoxy, -CH₂OH, phenyl and CF₃.

Individual embodiments provide compounds of the formula I wherein Ar is, respectively, indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine, naphthyridine, in which, in each embodiment the Ar group is either unsubstituted or substituted by 1, 2, 3, 4, or 5
20 R⁶ groups.

Individual embodiments provide compounds of the formula I wherein Ar is, respectively, indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine, naphthyridine, in which, in each embodiment the Ar group is either unsubstituted or substituted by 1, 2, 3, or 4 R⁶
25 groups.

Individual embodiments provide compounds of the formula I wherein Ar is, respectively, indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine, naphthyridine, in which, in each embodiment the Ar group is either unsubstituted or substituted by 1, 2, or 3 R⁶
30 groups.

Individual embodiments provide compounds of the formula I wherein Ar is, respectively, indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine, naphthyridine, in which, in each embodiment the Ar group is either unsubstituted or substituted by 1, 2, or 3
35 substituents selected from halogen, alkyl, alkoxy, CF₃, -O-CF₃, and -O-CF₂.

5 A specific value for Ar is indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyndine or naphthyndine, wherein indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyndine or naphthyndine are not substituted.

10 A specific value for Ar is indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyndine or naphthyridine, wherein indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine or naphthyridine are each optionally substituted by 1, 2, 3,
15 4, or 5 R⁶ groups.

A specific value for Ar is indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine or naphthyridine, wherein indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole,
20 triazolopyrimidine, pyrazolopyridine or naphthyridine are each optionally substituted by 1, 2, 3, or 4 R⁶ groups.

A specific value for Ar is indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine or naphthyridine, wherein indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole,
25 triazolopyrimidine, pyrazolopyridine or naphthyridine are each optionally substituted by 1, 2, or 3, or 4 R⁶ groups.


A specific value for Ar is indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine or naphthyridine, wherein indazole, imidazopyridazine, benzothiophene, benzoimidazole, indazole, quinoline, isoquinoline, quinoxaline, tetrahydroquinoline, indole, triazolopyrimidine, pyrazolopyridine or naphthyridine are each optionally substituted by 1, 2, or 3 substituents selected from halogen, alkyl, alkoxy, CF₃, -O-CF₃, and -O-CF₂.
30

A specific value for R⁶ is OR¹¹, CN, S(0)_pR^a, halogen, (C₁-C₈)alkyl, C₆-C₂₀ aryl or C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclalkyl, wherein any (C₁-C₈)alkyl,
35


5 (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R⁶ is optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(O)_pR^a, OR^a, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, -C(O)R^a, -C(O)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(O)_pR^a, NR^aS(O)_pR^a, NHC(O)R^a, NR^aC(O)R^a, NHC(O)OR^a, NR^aC(O)OR^a, NR^aC(O)NHR^a, NR^aC(O)N(R^a)₂, NR^aC(O)NH₂, NHC(O)NHR^a, NHC(O)N(R^a)₂, NHC(O)NH₂, =NH, =NOH, =NOR^a, NR^aS(O)_pNHR^a, NR^aS(O)_pN(R^a)₂, NR^aS(O)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(O)(OH)₂ or R^a.



Another specific value for R⁶ is OR¹¹, CN, S(O)_pR^a, halogen, (C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclylalkyl, NR¹¹C(O)R¹¹ or NR¹¹S(O)_pR^a, wherein any (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R⁶ is optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(O)_pR^a, OR^a, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, -C(O)R^a, -C(O)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(O)_pR^a, NR^aS(O)_pR^a, NHC(O)R^a, NR^aC(O)R^a, NHC(O)OR^a, NR^aC(O)OR^a, NR^aC(O)NHR^a, NR^aC(O)N(R^a)₂, NR^aC(O)NH₂, NHC(O)NHR^a, NHC(O)N(R^a)₂, NHC(O)NH₂, =NH, =NOH, =NOR^a, NR^aS(O)_pNHR^a, NR^aS(O)_pN(R^a)₂, NR^aS(O)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(O)(OH)₂ or R^a.

A specific value for Ar is:






25







5





10










5








10









5

Another specific value for Ar is:








5







10




15




5



10



In one embodiment the invention provides a compound of formula I:



or a salt or ester, thereof;

10 wherein:

A is $-(C(R^4)_2)_n$ - wherein any one $C(R^4)_2$ of said $-(C(R^4)_2)_n$ - may be optionally replaced with -O-, -S-, -S(O)_p-, NH or NR^a;

n is 3, 4, 5 or 6;

each p is 1 or 2;

15 Ar is a C₂-C₂₀ heterocyclyl group or a C₆-C₂₀ aryl group, wherein the C₂-C₂₀ heterocyclyl group or the C₆-C₂₀ aryl group is optionally substituted with 1, 2, 3, 4 or 5 R⁶;

each R³, R⁴ or R⁶ is independently H, oxo, OR¹¹, NR¹¹R¹², NR¹¹C(O)R¹¹, NR¹¹C(O)OR¹¹, NR¹¹C(O)NR¹¹R¹², N₃, CN, NO₂, SR¹¹, S(O)_pR^a, NR¹¹S(O)_pR^a, -C(=O)R¹¹, -C(=O)OR¹¹, -C(=O)NR¹¹R¹², -C(=O)SR¹¹, -S(O)_p(OR¹¹), -SO₂NR¹¹R¹², -NR¹¹S(O)_p(OR¹¹),

20 NR¹¹SO_pNR¹¹R¹², NR¹¹C(=NR¹¹)NR¹¹R¹², halogen, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclalkyl;

5 or two R⁴ on adjacent carbon atoms, when taken together, may optionally form a double bond between the two carbons to which they are attached or may form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a-;

10 four R⁴ on adjacent carbon atoms, when taken together, may optionally form an optionally substituted C₆ aryl ring;

two R⁴ on the same carbon atom, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a-;

15 two R⁶ on adjacent carbon atoms, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a-;


20 each R^a is independently (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl wherein any (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀ heterocyclyl, and wherein any aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀ heterocyclyl or (d-C₈)alkyl;

25 each R¹¹ or R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclylalkyl, -C(=O)R^a, -S(O)_pR^a, or aryl(CrC₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S-, -S(O)_p-, -NH-, -NR^a- or -C(O)-; and

30 wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of each R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(O)_pR^a, OR^a, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, -C(O)R^a, -C(O)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(O)_pR^a,


- 5 $\text{NR}^a\text{S}(\text{O})_p\text{R}^a$, $\text{NHC}(\text{O})\text{R}^a$, $\text{NR}^a\text{C}(\text{O})\text{R}^a$, $\text{NHC}(\text{O})\text{OR}^a$, $\text{NR}^a\text{C}(\text{O})\text{OR}^a$, $\text{NR}^a\text{C}(\text{O})\text{NHR}^a$,
 $\text{NR}^a\text{C}(\text{O})\text{N}(\text{R}^a)_2$, $\text{NR}^a\text{C}(\text{O})\text{NH}_2$, $\text{NHC}(\text{O})\text{NHR}^a$, $\text{NHC}(\text{O})\text{N}(\text{R}^a)_2$, $\text{NHC}(\text{O})\text{NH}_2$, $=\text{NH}$, $=\text{NOH}$,
 $=\text{NOR}^a$, $\text{NR}^a\text{S}(\text{O})_p\text{NHR}^a$, $\text{NR}^a\text{S}(\text{O})_p\text{N}(\text{R}^a)_2$, $\text{NR}^a\text{S}(\text{O})_p\text{NH}_2$, $\text{NHS}(\text{O})_p\text{NHR}^a$, $\text{NHS}(\text{O})_p\text{N}(\text{R}^a)_2$,
 $\text{NHS}(\text{O})_p\text{NH}_2$, $-\text{OC}(=\text{O})\text{R}^a$, $-\text{OP}(\text{O})(\text{OH})_2$ or R^a ;

provided the compound is not:




10

A specific group of compounds of formula I are compounds of formula Ia:




and salts and esters, thereof; provided the compound does not include:






15



In one embodiment the compounds of formula I or Ia do not include:





In one embodiment a compound of formula I is selected from:







5







5







5



5.

















5




and salts and esters, thereof.

In one embodiment a compound of formula I is selected from:







and



5

and salts and esters, thereof.


One embodiment provides a compound of formula II or IIa:




or a salt or ester, thereof, provided the compound is not

10


5




In one embodiment the compounds of formula II or IIa do not include:




One embodiment provides a compound of formula III or IIIa:




10 or a salt or ester, thereof, provided the compound is not:



In one embodiment the compounds of formula III or IIIa do not include:




One embodiment provides a compound of formula IV or IVa:




5

or a salt or ester, thereof, provided the compound is not



In one embodiment the compounds of formula IV or IVa do not include:



10


One embodiment provides for a mixture of a compound of formula III with a corresponding compound of formula IV. The mixture of the compound of formula III with the corresponding compound of formula IV is thus a mixture of trans diastereomers wherein the trans substituents are the substituents connected to the carbon marked with an asterisk 2 (*2) and the carbon marked with an asterisk 2 (*3). The invention also provides a mixture of a

15


compound of formula IIa with a corresponding compound of formula IVa. The mixture of the compound of formula IIa with the corresponding compound of formula IVa is thus a mixture of trans diastereomers wherein the trans substituents are the substituents connected to the carbon marked with an asterisk 2 (*2) and the carbon marked with an asterisk 2 (*3).

One embodiment does not include:

5






In one embodiment a compound is selected from:






and

and salts and esters, thereof, or a mixture of a compound of formula IIa and a corresponding compound of formula IVa or a salt or ester, thereof; wherein Ar is selected from:







5





10









5




10





5


In one embodiment a compound is selected from:





and

and salts and esters, thereof, or a mixture of a compound of formula IIa and a corresponding compound of formula IVa or a salt or ester, thereof; wherein Ar is selected from:




10






5






10




5





10





5



One embodiment provides a pharmaceutical composition comprising a compound of formula II, IIa, III, IIIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable carrier.

One embodiment provides a pharmaceutical composition comprising a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof, and a pharmaceutically acceptable carrier.

One embodiment provides a pharmaceutical composition comprising a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, and a pharmaceutically acceptable carrier.

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a compound of formula II, IIa, III, IIIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof.

5 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof.

10 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof.

15 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of formula II, IIa, III, IIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof.

20 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof.

25 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a, tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof.

30 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a compound of formula II, IIa, III, IIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof.

One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof.

5 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof.

10 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a compound of formula II, IIa, III, IIia, IV or IVa or a pharmaceutically acceptable salt or ester thereof.

15 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof.

20 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a tautomer, polymorph, pseudopolymorph, amorphous form, hydrate or solvate of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof.

25 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a compound of formula II, IIa, III, IIia, IV or IVa or a pharmaceutically acceptable salt or ester thereof, and a pharmaceutically acceptable diluent or carrier.

30 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof, and a pharmaceutically acceptable diluent or carrier.

35 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, and a pharmaceutically acceptable diluent or carrier.

5

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a compound of formula II, IIa, III, IIIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof, in combination with at least one additional therapeutic agent.

10

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof, in combination with at least one additional therapeutic agent.

15

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof by administering a therapeutically effective amount of a mixture of a compound of formula IIIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, in combination with at least one additional therapeutic agent.

20

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a first pharmaceutical composition comprising a compound of formula II, IIa, III, IIIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof,; and
 - b) a second pharmaceutical composition comprising at least one additional
- 25 therapeutic agent active against infectious *Pneumovirinae* viruses.

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a first pharmaceutical composition comprising a mixture of a compound of
- 30 formula III and a corresponding compound of formula IV, or pharmaceutically acceptable salts or esters thereof; and
- b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious *Pneumovirinae* viruses.

5 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

a) a first pharmaceutical composition comprising a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof; and
10

b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious *Pneumovirinae* viruses.

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a
15 combination pharmaceutical agent comprising:

a) a compound of formula II, IIa, III, IIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof; and

b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal
20 (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

a) a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof; and

b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

25 One embodiment provides a method of treating a *Pneumovirinae* infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

a) a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof; and

30 b) a therapeutic agent active against infectious *Pneumovirinae* viruses.

One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

a) a first pharmaceutical composition comprising a compound of formula II, IIa, III, IIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof; and
35

- 5 b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious respiratory syncytial viruses.

One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 10 a) a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof; and
- b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious respiratory syncytial viruses.

15 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a mixture of a compound of formula IIa and a corresponding compound of formula IVa or a pharmaceutically acceptable salt thereof a pharmaceutically acceptable salt or ester thereof; and
- 20 b) a second pharmaceutical composition comprising at least one additional therapeutic agent active against infectious respiratory syncytial viruses.

One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- 25 a) a compound of formula II, IIa, III, IIa, IV or IVa or a pharmaceutically acceptable salt or ester thereof; and
- b) a therapeutic agent active against infectious respiratory syncytial viruses.

30 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof; and
- b) a therapeutic agent active against infectious respiratory syncytial viruses.

5 One embodiment provides a method of treating a respiratory syncytial virus infection in a mammal (e.g. a human) in need thereof, by administering a therapeutically effective amount of a combination pharmaceutical agent comprising:

- a) a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof; and
- 10 b) a therapeutic agent active against infectious respiratory syncytial viruses.

One embodiment provides a compound of formula II, IIa, III, IIia, IV or IVa or a pharmaceutically acceptable salt or ester thereof for use in medical therapy.

One embodiment provides a mixture of a compound of formula III and a corresponding compound of formula IV a pharmaceutically acceptable salts or esters thereof, for use in medical therapy.

One embodiment provides a mixture of a compound of formula IIia and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, for use in medical therapy.

One embodiment provides a compound of formula II, IIa, III, IIia, IV or IVa or a pharmaceutically acceptable salt or ester thereof, for use in the prophylactic or therapeutic treatment of a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

One embodiment provides a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof, for use in the prophylactic or therapeutic treat a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

One embodiment provides a mixture of a compound of formula IIia and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, for use in the prophylactic or therapeutic treatment of a viral infection caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

One embodiment provides the use of a compound of formula II, IIa, III, IIia, IV or IVa or a pharmaceutically acceptable salt or ester thereof, for the manufacture of a medicament useful for the treatment of a viral infection in a mammal (e.g. a human) caused by a *Pneumovirinae* virus or a respiratory syncytial virus.


One embodiment provides the use of a mixture of a compound of formula III and a corresponding compound of formula IV or pharmaceutically acceptable salts or esters thereof,


5 for the manufacture of a medicament useful for the treatment of a viral infection in a mammal (e.g. a human) caused by a *Pneumovirinae* virus or a respiratory syncytial virus.

One embodiment provides the use of a mixture of a compound of formula IIa and a corresponding compound of formula IVa or pharmaceutically acceptable salts or esters thereof, for the manufacture of a medicament useful for the treatment of a viral infection in a mammal


10 (e.g. a human) caused by a *Pneumovirinae* virus or a respiratory syncytial virus.



In one embodiment a compound is selected from:











5






5












5





5















5













5












5






5






and



5




and salts and esters thereof.

One embodiment provides a compound selected from:





10

wherein W is:







5










or a salt, or stereoisomer thereof.

10 In another embodiment the compound is selected from:



5






and

5


and salts and esters thereof.


One embodiment provides a compound of formula:






10

wherein Y is selected from:








5




10



5






5


or a salt, or stereoisomer thereof.


In another embodiment the compound is selected from:






10



5








5









5




















5





5





5

and salts and esters thereof.

Esters of compounds of the invention.


The compounds disclosed herein also include "esters" of the compounds of the invention. Accordingly, one example of esters of the compounds of the invention include esters wherein a hydroxyl group of the compound of the invention is an ester. These esters of the invention are typically labile and thus the ester may be converted to the corresponding hydroxyl group *in vivo* (e.g. after administration). Esters include those esters based on carbon and phosphorus.

15 Typical esters include: $(R^aO)_2P(=O)O-$, $(HO)_2P(=O)O-$, $(C_1-C_8)alkyl(C=O)O-$, $C_6-C_{20}aryl(C=O)O-$, $C_2-C_{20}heterocyclyl(C=O)O-$ or $(C_4-C_8)carbocyclylalkyl(C=O)O-$ wherein each $(C_1-C_8)alkyl(C=O)O-$, $C_6-C_{20}aryl(C=O)O-$, $C_2-C_{20}heterocyclyl(C=O)O-$ or $(C_4-C_8)carbocyclylalkyl(C=O)O-$, is independently, optionally substituted with one or more oxo, halogen, hydroxy, NH_2 , CN , N_3 , $N(R^a)_2$, NHR^a , SH , SR^a , $S(O)_pR^a$, OR^a , $(C_1-C_8)alkyl$, $(C_1-C_8)haloalkyl$, $-C(O)R^a$, $-C(O)H$, $-C(=O)OR^a$, $-C(=O)OH$, $-C(=O)N(R^a)_2$, $-C(=O)NHR^a$, $-C(=O)NH_2$, $NHS(O)_pR^a$, $NR^aS(O)_pR^a$, $NHC(O)R^a$, $NR^aC(O)R^a$, $NHC(O)OR^a$, $NR^aC(O)OR^a$, $NR^aC(O)NHR^a$, $NR^aC(O)N(R^a)_2$, $NR^aC(O)NH_2$, $NHC(O)NHR^a$, $NHC(O)N(R^a)_2$, $NHC(O)NH_2$, $=NH$, $=NOH$, $=NOR^a$, $NR^aS(O)_pNHR^a$, $NR^aS(O)_pN(R^a)_2$, $NR^aS(O)_pNH_2$, $NHS(O)_pNHR^a$, $NHS(O)_pN(R^a)_2$, $NHS(O)_pNH_2$, $-OC(=O)R^a$, $-OP(O)(OH)_2$ or R^a ;

5 each R^a is independently (C_1-C_8) alkyl, (C_1-C_8) haloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, C_6-C_{20} aryl, C_2-C_{20} heterocyclyl, (C_3-C_7) cycloalkyl or (C_4-C_8) carbocyclalkyl wherein any (C_1-C_8) alkyl, (C_1-C_8) haloalkyl, (C_2-C_8) alkenyl or (C_2-C_8) alkynyl of R^a is optionally substituted with one or more OH, NH_2 , CO_2H , **C2-C20** heterocyclyl, and wherein any aryl(C_1-C_8)alkyl, C_6-C_{20} aryl, C_2-C_{20} heterocyclyl, (C_3-C_7) cycloalkyl or (C_4-C_8) carbocyclalkyl of R^a is optionally substituted with one or more OH, NH_2 , CO_2H , C_2-C_{20} heterocyclyl or (C_1-C_8) alkyl; and each p is 1 or 2.

It is to be understood that the point of connection of the esters $(R^aO)2P(=O)O-$, $(HO)_2P(=O)O-$, $(C_1-C_8)alkyl(C=O)O-$, $C_6-C_{20}aryl(C=O)O-$, $C_2-C_{20}heterocyclyl(C=O)O-$ and $(C_4-C_8)carbocyclalkyl(C=O)O-$ to the compound of the invention is through the oxygen of the ester.

In one embodiment the compounds of formula I include compounds of formula Ib



Ib

or a salt or ester, thereof;

20 wherein:

A is $-(C(R^4)_2)_n-$ wherein any one $C(R^4)_2$ of said $-(C(R^4)_2)_n-$ may be optionally replaced with $-O-$, $-S-$, $-S(O)_p-$, NH or NR^a ;

n is 3, 4, 5 or 6;

each p is 1 or 2;

25 Ar is a C_2-C_{20} heterocyclyl group or a C_6-C_{20} aryl group, wherein the C_2-C_{20} heterocyclyl group or the C_6-C_{20} aryl group is optionally substituted with 1, 2, 3, 4 or 5 R^6 ;

each R^3 , R^4 or R^6 is independently H, oxo, OR^{11} , $NR^{11}R^{12}$, $NR^{11}C(O)R^{11}$, $NR^{11}C(O)OR^{11}$, $NR^{11}C(O)NR^{11}R^{12}$, N_3 , CN, NO_2 , SR^{11} , $S(O)_pR^a$, $NR^{11}S(O)_pR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)_p(OR^{11})$, $-SO_2NR^{11}R^{12}$, $-NR^{11}S(O)_p(OR^{11})$,

30 $-NR^{11}SO_pNR^{11}R^{12}$, $NR^{11}C(=NR^{11})NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl,

5 (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl;

two R⁴ on adjacent carbon atoms, when taken together, may optionally form a double bond between the two carbons to which they are attached or may form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-,
10 -S(O)_p-, -NH- or -NR^a;

four R⁴ on adjacent carbon atoms, when taken together, may optionally form an optionally substituted C₆ aryl ring;

two R⁴ on the same carbon atom, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally
15 replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a;

two R⁶ on adjacent carbon atoms, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a;

each R^a is independently (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl,
20 aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl wherein any (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀ heterocyclyl, and wherein any aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R^a is optionally substituted with one or more OH,
25 NH₂, CO₂H, C₂-C₂₀ heterocyclyl or (C₁-C₈)alkyl;

each R¹¹ or R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl,
aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclylalkyl,
-C(=O)R^a, -S(O)_pR^a, or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which
30 they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S-, -S(O)_p-, -NH-, -NR^a- or -C(O)-;

wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of each R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃,

- 5 $N(R^a)_2$, NHR^a , SH , SR^a , $S(O)_p R^a$, OR^a , $(C_1-C_8)alkyl$, $(C_1-C_8)haloalkyl$, $-C(O)R^a$, $-C(O)H$,
 $-C(=O)OR^a$, $-C(=O)OH$, $-C(=O)N(R^a)_2$, $-C(=O)NHR^a$, $-C(=O)NH_2$, $NHS(O)_p R^a$,
 $NR^a S(O)_p R^a$, $NHC(O)R^a$, $NR^a C(O)R^a$, $NHC(O)OR^a$, $NR^a C(O)OR^a$, $NR^a C(O)NHR^a$,
 $NR^a C(O)N(R^a)_2$, $NR^a C(O)NH_2$, $NHC(O)NHR^a$, $NHC(O)N(R^a)_2$, $NHC(O)NH_2$, $=NH$, $=NOH$,
 $-NOR^a$, $NR^a S(O)_p NHR^a$, $NR^a S(O)_p N(R^a)_2$, $NR^a S(O)_p NH_2$, $NHS(O)_p NHR^a$, $NHS(O)_p N(R^a)_2$,
10 $NHS(O)_p NH_2$, $-OC(=O)R^a$, $-OP(O)(OH)_2$ or R^a ; and

R^x is H , $(R^a O)_2 P(=O)-$, $(HO)_2 P(=O)O-$, $(C_1-C_8)alkyl(C=O)-$, $C_6-C_{20}aryl(C=O)-$,


$C_2-C_{20}heterocyclyl(C=O)-$ or $(C_4-C_8)carbocyclylalkyl(C=O)-$, wherein each $(C_1-C_8)alkyl(C=O)-$,

$C_6-C_{20}aryl(C=O)-$, $C_2-C_{20}heterocyclyl(C=O)-$ or $(C_4-C_8)carbocyclylalkyl(C=O)-$ is

independently, optionally substituted with one or more oxo, halogen, hydroxy, NH_2 , CN , N_3 ,

- 15 $N(R^a)_2$, NHR^a , SH , SR^a , $S(O)_p R^a$, OR^a , $(d-C_8)alkyl$, $(C_1-C_8)haloalkyl$, $-C(O)R^a$, $-C(O)H$,
 $-C(=O)OR^a$, $-C(=O)OH$, $-C(=O)N(R^a)_2$, $-C(=O)NHR^a$, $-C(=O)NH_2$, $NHS(O)_p R^a$,
 $NR^a S(O)_p R^a$, $NHC(O)R^a$, $NR^a C(O)R^a$, $NHC(O)OR^a$, $NR^a C(O)OR^a$, $NR^a C(O)NHR^a$,
 $NR^a C(O)N(R^a)_2$, $NR^a C(O)NH_2$, $NHC(O)NHR^a$, $NHC(O)N(R^a)_2$, $NHC(O)NH_2$, $=NH$, $=NOH$,
 $=NOR^a$, $NR^a S(O)_p NHR^a$, $NR^a S(O)_p N(R^a)_2$, $NR^a S(O)_p NH_2$, $NHS(O)_p NHR^a$, $NHS(O)_p N(R^a)_2$,
20 $NHS(O)_p NH_2$, $-OC(=O)R^a$, $-OP(O)(OH)_2$ or R^a ;

provided the compound is not:



A specific value for R^x is H , $(HO)_2 P(=O)-$, $(C_1-C_8)alkyl(C=O)-$ or



$C_2-C_{20}heterocyclyl(C=O)-$, wherein each $(C_1-C_8)alkyl(C=O)-$ or $C_2-C_{20}heterocyclyl(C=O)-$ is

- 25 independently, optionally substituted with one or more oxo, halogen, hydroxy, NH_2 , CN , N_3 ,
 $N(R^a)_2$, NHR^a , SH , SR^a , $S(O)_p R^a$, OR^a , $(Q-CiO)haloalkyl$, $-C(O)R^a$, $-C(O)H$,
 $-C(=O)OR^a$, $-C(=O)OH$, $-C(=O)N(R^a)_2$, $-C(=O)NHR^a$, $-C(=O)NH_2$, $NHS(O)_p R^a$,
 $NR^a S(O)_p R^a$, $NHC(O)R^a$, $NR^a C(O)R^a$, $NHC(O)OR^a$, $NR^a C(O)OR^a$, $NR^a C(O)NHR^a$,
 $NR^a C(O)N(R^a)_2$, $NR^a C(O)NH_2$, $NHC(O)NHR^a$, $NHC(O)N(R^a)_2$, $NHC(O)NH_2$, $=NH$, $=NOH$,
30 $=NOR^a$, $NR^a S(O)_p NHR^a$, $NR^a S(O)_p N(R^a)_2$, $NR^a S(O)_p NH_2$, $NHS(O)_p NHR^a$, $NHS(O)_p N(R^a)_2$,

5 NHS(O)_pNH₂, -OC(=O)R^a, -OP(O)(OH)₂ or R^a.

Another specific value for R^x is H, (HO)₂P(=O)-, (C₁-C₈)alkyl(C=O)- or C₂-C₂₀heterocyclyl(C=O)-, wherein each (C₁-C₈)alkyl(C=O)- or C₂-C₂₀heterocyclyl(C=O)- is independently, optionally substituted with one or more NH₂, -C(=O)OH or NR^aC(O)R^a.

Another specific value for R^x is:



10

Preparation of compounds of the invention.

The compounds of formulas I and Ia were prepared by the procedures described in examples 3, 4 and 6-31. The compounds of formulas 1-24 (i.e. compounds of formula V) were prepared by the procedures described in example 1 and example 4. The compounds of formulas 25-111 (i.e. compounds of formula VI) were prepared by the procedures described in examples 2, 4 and 5.

The compounds of formula II and IIa can be prepared following the procedures described in examples 3, 4, 5 and 6 by using intermediate 9b instead of intermediate 9a.

20 The compounds of formulas III, IIIa, IV and IVa can be prepared following the procedures described in examples 3, 4, 5 and 6 by using the enantiomers of trans-3-cyano-4-hydroxypyrrolidine instead of the cis enantiomers. The enantiomers of trans-3-cyano-4-hydroxypyrrolidine can be prepared following literature procedures (Schauss, S.E., et al., Organic Letters, 2(7), 2000, pages 1001-1004).

5 Pharmaceutical Formulations

The compounds of this invention are formulated with conventional carriers and excipients, which will be selected in accord with ordinary practice. Tablets will contain excipients, glidants, fillers, binders and the like. Aqueous formulations are prepared in sterile form, and when intended for delivery by other than oral administration generally will be isotonic. All formulations will optionally contain excipients such as those set forth in the "Handbook of Pharmaceutical Excipients" (1986). Excipients include ascorbic acid and other antioxidants, chelating agents such as EDTA, carbohydrates such as dextran, hydroxyalkylcellulose, hydroxyalkylmethylcellulose, stearic acid and the like. The pH of the formulations ranges from about 3 to about 11, but is ordinarily about 7 to 10.

15 While it is possible for the active ingredients to be administered alone it may be preferable to present them as pharmaceutical formulations. The formulations, both for veterinary and for human use, of the invention comprise at least one active ingredient, as above defined, together with one or more acceptable carriers and optionally other therapeutic ingredients, particularly those additional therapeutic ingredients as discussed herein. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and physiologically innocuous to the recipient thereof.

The formulations include those suitable for the foregoing administration routes. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Techniques and formulations generally are found in Remington's Pharmaceutical Sciences (Mack Publishing Co., Easton, PA). Such methods include the step of bringing into association the active ingredient with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

30 Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be administered as a bolus, electuary or paste.

35 A tablet is made by compression or molding, optionally with one or more accessory

5 ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered active ingredient moistened with an inert liquid diluent. The tablets may optionally be coated or scored and
10 optionally are formulated so as to provide slow or controlled release of the active ingredient therefrom.

For infections of the eye or other external tissues e.g. mouth and skin, the formulations are preferably applied as a topical ointment or cream containing the active ingredient(s) in an amount of, for example, 0.075 to 20% w/w (including active ingredient(s) in a range between
15 0.1% and 20% in increments of 0.1% w/w such as 0.6% w/w, 0.7% w/w, etc.), preferably 0.2 to 15% w/w and most preferably 0.5 to 10% w/w. When formulated in an ointment, the active ingredients may be employed with either a paraffinic or a water-miscible ointment base. Alternatively, the active ingredients may be formulated in a cream with an oil-in-water cream base.

20 If desired, the aqueous phase of the cream base may include, for example, at least 30% w/w of a polyhydric alcohol, i.e. an alcohol having two or more hydroxyl groups such as propylene glycol, butane 1,3-diol, mannitol, sorbitol, glycerol and polyethylene glycol (including PEG 400) and mixtures thereof. The topical formulations may desirably include a compound which enhances absorption or penetration of the active ingredient through the skin or
25 other affected areas. Examples of such dermal penetration enhancers include dimethyl sulphoxide and related analogs.

The oily phase of the emulsions of this invention may be constituted from known ingredients in a known manner. While the phase may comprise merely an emulsifier (otherwise known as an emulgent), it desirably comprises a mixture of at least one emulsifier with a fat or
30 an oil or with both a fat and an oil. Preferably, a hydrophilic emulsifier is included together with a lipophilic emulsifier which acts as a stabilizer. It is also preferred to include both an oil and a fat. Together, the emulsifier(s) with or without stabilizer(s) make up the so-called emulsifying wax, and the wax together with the oil and fat make up the so-called emulsifying ointment base which forms the oily dispersed phase of the cream formulations.

35 Emulgents and emulsion stabilizers suitable for use in the formulation of the invention

5 include Tween® 60, Span® 80, cetostearyl alcohol, benzyl alcohol, myristyl alcohol, glyceryl mono-stearate and sodium lauryl sulfate.

The choice of suitable oils or fats for the formulation is based on achieving the desired cosmetic properties. The cream should preferably be a non-greasy, non-staining and washable product with suitable consistency to avoid leakage from tubes or other containers. Straight or
10 branched chain, mono- or dibasic alkyl esters such as di-isoadipate, isocetyl stearate, propylene glycol diester of coconut fatty acids, isopropyl myristate, decyl oleate, isopropyl palmitate, butyl stearate, 2-ethylhexyl palmitate or a blend of branched chain esters known as Crodamol CAP may be used, the last three being preferred esters. These may be used alone or in combination depending on the properties required. Alternatively, high melting point lipids such as white soft
15 paraffin and/or liquid paraffin or other mineral oils are used.

Pharmaceutical formulations according to the present invention comprise a combination according to the invention together with one or more pharmaceutically acceptable carriers or excipients and optionally other therapeutic agents. Pharmaceutical formulations containing the active ingredient may be in any form suitable for the intended method of administration. When
20 used for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents and preserving
25 agents, in order to provide a palatable preparation. Tablets containing the active ingredient in admixture with non-toxic pharmaceutically acceptable excipient which are suitable for manufacture of tablets are acceptable. These excipients may be, for example, inert diluents, such as calcium or sodium carbonate, lactose, calcium or sodium phosphate; granulating and disintegrating agents, such as maize starch, or alginic acid; binding agents, such as starch,
30 gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

5 Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

10 Aqueous suspensions of the invention contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally-occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a
15 condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate). The aqueous suspension may also contain one or more preservatives such as ethyl or n-propyl p-hydroxy-benzoate, one or more coloring agents, one or more flavoring agents and one or more
20 sweetening agents, such as sucrose or saccharin.

Oil suspensions may be formulated by suspending the active ingredient in a vegetable oil, such as arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oral suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth above, and flavoring agents may be
25 added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

Dispersible powders and granules of the invention suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, a suspending agent, and one or more preservatives. Suitable dispersing or
30 wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include
35 naturally-occurring gums, such as gum acacia and gum tragacanth, naturally-occurring

5 phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids and
hexitol anhydrides, such as sorbitan monooleate, and condensation products of these partial
esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may
also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with
sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a
10 demulcent, a preservative, a flavoring or a coloring agent.

The pharmaceutical compositions of the invention may be in the form of a sterile
injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This
suspension may be formulated according to the known art using those suitable dispersing or
wetting agents and suspending agents which have been mentioned above. The sterile injectable
15 preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally
acceptable diluent or solvent, such as a solution in 1,3-butane-diol or prepared as a lyophilized
powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's
solution and isotonic sodium chloride solution. In addition, sterile fixed oils may conventionally
be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be
20 employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid
may likewise be used in the preparation of injectables.

The amount of active ingredient that may be combined with the carrier material to
produce a single dosage form will vary depending upon the host treated and the particular mode
of administration. For example, a time-release formulation intended for oral administration to
25 humans may contain approximately 1 to 1000 mg of active material compounded with an
appropriate and convenient amount of carrier material which may vary from about 5 to about
95% of the total compositions (weightweight). The pharmaceutical composition can be
prepared to provide easily measurable amounts for administration. For example, an aqueous
solution intended for intravenous infusion may contain from about 3 to 500 μg of the active
30 ingredient per milliliter of solution in order that infusion of a suitable volume at a rate of about
30 mL/hr can occur.

Formulations suitable for topical administration to the eye also include eye drops
wherein the active ingredient is dissolved or suspended in a suitable carrier, especially an
aqueous solvent for the active ingredient. The active ingredient is preferably present in such
35 formulations in a concentration of 0.5 to 20%, advantageously 0.5 to 10%, and particularly

5 about 1.5% w/w.

Formulations suitable for topical administration in the mouth include lozenges comprising the active ingredient in a flavored basis, usually sucrose and acacia or tragacanth; pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid
10 carrier.

Formulations for rectal administration may be presented as a suppository with a suitable base comprising for example cocoa butter or a salicylate.

Formulations suitable for intrapulmonary or nasal administration have a particle size for example in the range of 0.1 to 500 microns, such as 0.5, 1, 30, 35 etc., which is administered by
15 rapid inhalation through the nasal passage or by inhalation through the mouth so as to reach the alveolar sacs. Suitable formulations include aqueous or oily solutions of the active ingredient. Formulations suitable for aerosol or dry powder administration may be prepared according to conventional methods and may be delivered with other therapeutic agents such as compounds heretofore used in the treatment or prophylaxis of *Pneumovirinae* infections as described below.

20 In another aspect, the invention is a novel, efficacious, safe, nonirritating and physiologically compatible inhalable composition comprising a compound of formula I or a compound of formulas 1-1 11, or a pharmaceutically acceptable salt thereof, suitable for treating *Pneumovirinae* infections and potentially associated bronchiolitis. Preferred pharmaceutically acceptable salts are inorganic acid salts including hydrochloride, hydrobromide, sulfate or
25 phosphate salts as they may cause less pulmonary irritation. Preferably, the inhalable formulation is delivered to the endobronchial space in an aerosol comprising particles with a mass median aerodynamic diameter (MMAD) between about 1 and about 5 μm . Preferably, the compound of formula I or formulas 1-1 11 is formulated for aerosol delivery using a nebulizer, pressurized metered dose inhaler (pMDI), or dry powder inhaler (DPI).

30 Non-limiting examples of nebulizers include atomizing, jet, ultrasonic, pressurized, vibrating porous plate, or equivalent nebulizers including those nebulizers utilizing adaptive aerosol delivery technology (Denyer, *J Aerosol medicine Pulmonary Drug Delivery* 2010, 23 Supp 1, SI-SI0). A jet nebulizer utilizes air pressure to break a liquid solution into aerosol droplets. An ultrasonic nebulizer works by a piezoelectric crystal that shears a liquid into small
35 aerosol droplets. A pressurized nebulization system forces solution under pressure through

5 small pores to generate aerosol droplets. A vibrating porous plate device utilizes rapid vibration to shear a stream of liquid into appropriate droplet sizes.

In a preferred embodiment, the formulation for nebulization is delivered to the endobronchial space in an aerosol comprising particles with a MMAD predominantly between about 1 μm and about 5 μm using a nebulizer able to aerosolize the formulation of the
10 compound of formula I or formulas 1-11 into particles of the required MMAD. To be optimally therapeutically effective and to avoid upper respiratory and systemic side effects, the majority of aerosolized particles should not have a MMAD greater than about 5 μm . If an aerosol contains a large number of particles with a MMAD larger than 5 μm , the particles are deposited in the upper airways decreasing the amount of drug delivered to the site of
15 inflammation and bronchoconstriction in the lower respiratory tract. If the MMAD of the aerosol is smaller than about 1 μm , then the particles have a tendency to remain suspended in the inhaled air and are subsequently exhaled during expiration.

When formulated and delivered according to the method of the invention, the aerosol formulation for nebulization delivers a therapeutically efficacious dose of the compound of
20 formula I or formulas 1-11 to the site of *Pneumovirinae* infection sufficient to treat the *Pneumovirinae* infection. The amount of drug administered must be adjusted to reflect the efficiency of the delivery of a therapeutically efficacious dose of the compound of formula I or formulas 1-11. In a preferred embodiment, a combination of the aqueous aerosol formulation with the atomizing, jet, pressurized, vibrating porous plate, or ultrasonic nebulizer permits,
25 depending on the nebulizer, about, at least, 20, to about 90%, typically about 70% delivery of the administered dose of the compound of formula I or formulas 1-11 into the airways. In a preferred embodiment, at least about 30 to about 50% of the active compound is delivered. More preferably, about 70 to about 90% of the active compound is delivered.

In another embodiment of the instant invention, a compound of formula I or formulas 1-
30 111 or a pharmaceutically acceptable salt thereof, is delivered as a dry inhalable powder. The compounds of the invention are administered endobronchially as a dry powder formulation to efficaciously deliver fine particles of compound into the endobronchial space using dry powder or metered dose inhalers. For delivery by DPI, the compound of formula I or formulas 1-11 is processed into particles with, predominantly, MMAD between about 1 μm and about 5 μm by
35 milling spray drying, critical fluid processing, or precipitation from solution. Media milling, jet

5 milling and spray-drying devices and procedures capable of producing the particle sizes with a MMAD between about 1 μm and about 5 μm are well known in the art. In one embodiment, excipients are added to the compound of formula I or formulas 1-111 before processing into particles of the required sizes. In another embodiment, excipients are blended with the particles of the required size to aid in dispersion of the drug particles, for example by using lactose as an
10 excipient.

Particle size determinations are made using devices well known in the art. For example a multi-stage Anderson cascade impactor or other suitable method such as those specifically cited within the US Pharmacopoeia Chapter 601 as characterizing devices for aerosols within metered-dose and dry powder inhalers.

15 In another preferred embodiment, a compound of formula I or formulas 1-111 is delivered as a dry powder using a device such as a dry powder inhaler or other dry powder dispersion devices. Non-limiting examples of dry powder inhalers and devices include those disclosed in US5,458,135; US5,740,794; US5,775,320; US5,785,049; US3,906,950; US4,013,075; US4,069,819; US4,995,385; US5,522,385; US4,668,218; US4,667,668;
20 US4,805,811 and US5,388,572. There are two major designs of dry powder inhalers. One design is a metering device in which a reservoir for the drug is placed within the device and the patient adds a dose of the drug into the inhalation chamber. The second design is a factory-metered device in which each individual dose has been manufactured in a separate container. Both systems depend on the formulation of the drug into small particles of MMAD from 1
25 μm and about 5 μm , and often involve co-formulation with larger excipient particles such as, but not limited to, lactose. Drug powder is placed in the inhalation chamber (either by device metering or by breakage of a factory-metered dosage) and the inspiratory flow of the patient accelerates the powder out of the device and into the oral cavity. Non-laminar flow characteristics of the powder path cause the excipient-drug aggregates to decompose, and the
30 mass of the large excipient particles causes their impaction at the back of the throat, while the smaller drug particles are deposited deep in the lungs. In preferred embodiments, a compound of formula I or formulas 1-111, or a pharmaceutically acceptable salt thereof, is delivered as a dry powder using either type of dry powder inhaler as described herein, wherein the MMAD of the dry powder, exclusive of any excipients, is predominantly in the range of 1 μm to about 5
35 μm .

5 In another preferred embodiment, a compound of formula I or formulas 1-11 is delivered as a dry powder using a metered dose inhaler. Non-limiting examples of metered dose inhalers and devices include those disclosed in US5,261,538; US5,544,647; US5,622,163; US4,955,371; US3,565,070; US3,361,306 and US6,116,234. In preferred embodiments, a compound of formula I or formulas 1-11, or a pharmaceutically acceptable salt thereof, is delivered as a dry powder using a metered dose inhaler wherein the MMAD of the dry powder, exclusive of any excipients, is predominantly in the range of about 1-5 μm .

Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams or spray formulations containing in addition to the active ingredient such carriers as are known in the art to be appropriate.

15 Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents.

The formulations are presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions are prepared from sterile powders, granules and tablets of the kind previously described. Preferred unit dosage formulations are those containing a daily dose or unit daily sub-dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient.

It should be understood that in addition to the ingredients particularly mentioned above the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavoring agents.

30 The invention further provides veterinary compositions comprising at least one active ingredient as above defined together with a veterinary carrier therefor.

Veterinary carriers are materials useful for the purpose of administering the composition and may be solid, liquid or gaseous materials which are otherwise inert or acceptable in the veterinary art and are compatible with the active ingredient. These veterinary compositions may be administered orally, parenterally or by any other desired route.

5 Compounds of the invention are used to provide controlled release pharmaceutical formulations containing as active ingredient one or more compounds of the invention ("controlled release formulations") in which the release of the active ingredient are controlled and regulated to allow less frequency dosing or to improve the pharmacokinetic or toxicity profile of a given active ingredient.

10 Effective dose of active ingredient depends at least on the nature of the condition being treated, toxicity, whether the compound is being used prophylactically (lower doses) or against an active viral infection, the method of delivery, and the pharmaceutical formulation, and will be determined by the clinician using conventional dose escalation studies. It can be expected to be from about 0.0001 to about 100 mg/kg body weight per day; typically, from about 0.01 to about 15 10 mg/kg body weight per day; more typically, from about .01 to about 5 mg/kg body weight per day; most typically, from about .05 to about 0.5 mg/kg body weight per day. For example, the daily candidate dose for an adult human of approximately 70 kg body weight will range from 1 mg to 1000 mg, preferably between 5 mg and 500 mg, and may take the form of single or multiple doses.

20 Routes of Administration

 One or more compounds of the invention (herein referred to as the active ingredients) are administered by any route appropriate to the condition to be treated. Suitable routes include oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), vaginal and parenteral (including subcutaneous, intramuscular, intravenous, intradermal, intrathecal and epidural), and 25 the like. It will be appreciated that the preferred route may vary with for example the condition of the recipient. An advantage of the compounds of this invention is that they are orally bioavailable and can be dosed orally.

Combination Therapy

 Compositions of the invention are also used in combination with other active ingredients. 30 For the treatment of *Pneumovirinae* virus infections, preferably, the other active therapeutic agent is active against *Pneumovirinae* virus infections, particularly respiratory syncytial virus infections. Non-limiting examples of these other active therapeutic agents are ribavirin, palivizumab, motavizumab, RSV-IGIV (RespiGam[®]), MEDI-557, A-60444 (also known as RSV604), MDT-637, BMS-433771, ALN-RSV0, ALX-0171 and mixtures thereof.

5 Many of the infections of the *Pneumovirinae* viruses are respiratory infections. Therefore, additional active therapeutics used to treat respiratory symptoms and sequelae of infection may be used in combination with the compounds of formula I or formulas 1-1 11. The additional agents are preferably administered orally or by direct inhalation. For example, other preferred additional therapeutic agents in combination with the compounds of formula I or
10 formulas 1-1 11 for the treatment of viral respiratory infections include, but are not limited to, bronchodilators and corticosteroids.

Glucocorticoids, which were first introduced as an asthma therapy in 1950 (Carrier, Journal of Allergy, 21, 282-287, 1950), remain the most potent and consistently effective therapy for this disease, although their mechanism of action is not yet fully understood (Morris, J. Allergy Clin. Immunol., 75 (1 Pt) 1-13, 1985). Unfortunately, oral glucocorticoid therapies are associated with profound undesirable side effects such as truncal obesity, hypertension, glaucoma, glucose intolerance, acceleration of cataract formation, bone mineral loss, and psychological effects, all of which limit their use as long-term therapeutic agents (Goodman and Gilman, 10th edition, 2001). A solution to systemic side effects is to deliver steroid drugs
15 directly to the site of inflammation. Inhaled corticosteroids (ICS) have been developed to mitigate the severe adverse effects of oral steroids. Non-limiting examples of corticosteroids that may be used in combinations with the compounds of formula I or compounds of formulas 1-11 are dexamethasone, dexamethasone sodium phosphate, fluorometholone, fluorometholone acetate, loteprednol, loteprednol etabonate, hydrocortisone, prednisolone, fludrocortisone,
20 triamcinolone, triamcinolone acetonide, betamethasone, beclomethasone dipropionate, methylprednisolone, flucinolone, flucinolone acetonide, flunisolide, flucortin-21-butylate, flumethasone, flumetasone pivalate, budesonide, halobetasol propionate, mometasone furoate, fluticasone propionate, ciclesonide; or a pharmaceutically acceptable salts thereof.

Other anti-inflammatory agents working through anti-inflammatory cascade mechanisms are
30 also useful as additional therapeutic agents in combination with the compounds of formula I or the compounds of formulas 1-1 11 for the treatment of viral respiratory infections. Applying "anti-inflammatory signal transduction modulators" (referred to in this text as AISTM), like phosphodiesterase inhibitors (e.g. PDE-4, PDE-5, or PDE-7 specific), transcription factor inhibitors (e.g. blocking NF κ B through IKK inhibition), or kinase inhibitors (e.g. blocking P38
35 MAP, JNK, PI3K, EGFR or Syk) is a logical approach to switching off inflammation as these

5 small molecules target a limited number of common intracellular pathways - those signal transduction pathways that are critical points for the anti-inflammatory therapeutic intervention (see review by P.J. Barnes, 2006). These non-limiting additional therapeutic agents include: 5-(2,4-Difluoro-phenoxy)-1-isobutyl-1H-indazole-6-carboxylic acid (2-dimethylamino-ethyl)-amide (P38 Map kinase inhibitor ARRY-797); 3-Cyclopropylmethoxy-N-(3,5-dichloro-pyridin-4-yl)-4-difluoromethoxy-benzamide (PDE-4 inhibitor Roflumilast); 4-[2-(3-cyclopentyloxy-4-methoxyphenyl)-2-phenyl-ethyl]-pyridine (PDE-4 inhibitor CDP-840); N-(3,5-dichloro-4-pyridinyl)-4-(difluoromethoxy)-8-[(methylsulfonyl)amino]-1-dibenzofurancarboxamide (PDE-4 inhibitor Oglemilast); N-(3,5-Dichloro-pyridin-4-yl)-2-[1-(4-fluorobenzyl)-5-hydroxy-1H-indol-3-yl]-2-oxo-acetamide (PDE-4 inhibitor AWD 12-281); 8-Methoxy-2-trifluoromethyl-quinoline-5-carboxylic acid (3,5-dichloro-1-oxy-pyridin-4-yl)-amide (PDE-4 inhibitor Sch 351 591); 4-[5-(4-Fluorophenyl)-2-(4-methanesulfinyl-phenyl)-1H-imidazol-4-yl]-pyridine (P38 inhibitor SB-203850); 4-[4-(4-Fluoro-phenyl)-1-(3-phenyl-propyl)-5-pyridin-4-yl-1H-imidazol-2-yl]-but-3-yn-1-ol (P38 inhibitor RWJ-67657); 4-Cyano-4-(3-cyclopentyloxy-4-methoxy-phenyl)-cyclohexanecarboxylic acid 2-diethylamino-ethyl ester (2-diethyl-ethyl ester prodrug of Cilomilast, PDE-4 inhibitor); (3-Chloro-4-fluorophenyl)-[7-methoxy-6-(3-morpholin-4-yl-propoxy)-quinazolin-4-yl]-amine (Gefitinib, EGFR inhibitor); and 4-(4-Methyl-piperazin-1-ylmethyl)-N-[4-methyl-3-(4-pyridin-3-yl-pyrimidin-2-ylamino)-phenyl]-benzamide (Imatinib, EGFR inhibitor).

Combinations comprising inhaled β_2 -adrenoreceptor agonist bronchodilators such as formoterol, albuterol or salmeterol with the compounds of formula I or formulas 1-11 are also suitable, but non-limiting, combinations useful for the treatment of respiratory viral infections.

Combinations of inhaled β_2 -adrenoreceptor agonist bronchodilators such as formoterol or salmeterol with ICS's are also used to treat both the bronchoconstriction and the inflammation (Symbicort® and Advair®, respectively). The combinations comprising these ICS and β_2 -adrenoreceptor agonist combinations along with the compounds of formula I or formulas 1-11 are also suitable, but non-limiting, combinations useful for the treatment of respiratory viral infections.

For the treatment or prophylaxis of pulmonary broncho-constriction, anticholinergics are of potential use and, therefore, useful as an additional therapeutic agents in combination with the compounds of formula I or formulas 1-11 for the treatment of viral respiratory infections.

5 These anticholinergics include, but are not limited to, antagonists of the muscarinic receptor (particularly of the M3 subtype) which have shown therapeutic efficacy in man for the control of cholinergic tone in COPD (Witek, 1999); 1-{4-Hydroxy-1-[3,3,3-tris-(4-fluoro-phenyl)-propionyl]-pyrrolidine-2-carbonyl}-pyrrolidine-2-carboxylic acid (1-methyl-piperidin-4-ylmethyl)-amide; 3-[3-(2-Diethylamino-acetoxy)-2-phenyl-propionyloxy]-8-isopropyl-8-methyl-
 10 8-azonia-bicyclo[3.2.1]octane (Ipratropium-N,N-diethylglycinate); 1-Cyclohexyl-3,4-dihydro-1H-isoquinoline-2-carboxylic acid 1-aza-bicyclo[2.2.2]oct-3-yl ester (Solifenacin); 2-Hydroxymethyl-4-methanesulfinyl-2-phenyl-butyric acid 1-aza-bicyclo[2.2.2]oct-3-yl ester (Revatropate); 2-[1-[2-(2,3-Dihydro-benzofuran-5-yl)-ethyl]-pyrrolidin-3-yl]-2,2-diphenyl-acetamide (Darifenacin); 4-Azepan-1-yl-2,2-diphenyl-butyramide (Buzepide); 7-[3-(2-
 15 Diethylamino-acetoxy)-2-phenyl-propionyloxy]-9-ethyl-9-methyl-3-oxa-9-azonia-tricyclo[3.3.1.0^{2,4}]nonane (Oxotropium-N,N-diethylglycinate); 7-[2-(2-Diethylamino-acetoxy)-2,2-di-thiophen-2-yl-acetoxy]-9,9-dimethyl-3-oxa-9-azonia-tricyclo[3.3.1.0^{2,4}]nonane (Tiotropium-N,N-diethylglycinate); Dimethylamino-acetic acid 2-(3-diisopropylamino-1-phenyl-propyl)-4-methyl-phenyl ester (Tolterodine-N,N-dimethylglycinate); 3-[4,4-Bis-(4-
 20 fluoro-phenyl)-2-oxo-imidazolidin-1-yl]-1-methyl-1-(2-oxo-2-pyridin-2-yl-ethyl)-pyrrolidinium; 1-[1-(3-Fluoro-benzyl)-piperidin-4-yl]-4,4-bis-(4-fluoro-phenyl)-imidazolidin-2-one; 1-Cyclooctyl-3-(3-methoxy-1-aza-bicyclo[2.2.2]oct-3-yl)-1-phenyl-prop-2-yn-1-ol; 3-[2-(2-Diethylamino-acetoxy)-2,2-di-thiophen-2-yl-acetoxy]-1-(3-phenoxy-propyl)-1-azonia-bicyclo[2.2.2]octane (Aclidinium-N,N-diethylglycinate); or (2-Diethylamino-acetoxy)-di-
 25 thiophen-2-yl-acetic acid 1-methyl-1-(2-phenoxy-ethyl)-piperidin-4-yl ester.

The compounds of formula I or formulas 1-11 may also be combined with mucolytic agents to treat both the infection and symptoms of respiratory infections. A non-limiting example of a mucolytic agent is ambroxol. Similarly, the compounds of formula I or formulas 1-11 may be combined with expectorants to treat both the infection and symptoms of
 30 respiratory infections. A non-limiting example of an expectorant is guaifenesin.

Nebulized hypertonic saline is used to improve immediate and long-term clearance of small airways in patients with lung diseases (Kuzik, *J Pediatrics* 2007, 266). The compounds of formula I or formulas 1-11 may also be combined with nebulized hypertonic saline particularly when the *Pneumovirinae* virus infection is complicated with bronchiolitis. The
 35 combination of the compounds of formula I or formulas 1-11 with hypertonic saline may also

5 comprise any of the additional agents discussed above. In a preferred aspect, nebulized about 3% hypertonic saline is used.

It is also possible to combine any compound of the invention with one or more additional active therapeutic agents in a unitary dosage form for simultaneous or sequential administration to a patient. The combination therapy may be administered as a simultaneous or sequential
10 regimen. When administered sequentially, the combination may be administered in two or more administrations.

Co-administration of a compound of the invention with one or more other active therapeutic agents generally refers to simultaneous or sequential administration of a compound of the invention and one or more other active therapeutic agents, such that therapeutically
15 effective amounts of the compound of the invention and one or more other active therapeutic agents are both present in the body of the patient.

Co-administration includes administration of unit dosages of the compounds of the invention before or after administration of unit dosages of one or more other active therapeutic agents, for example, administration of the compounds of the invention within seconds, minutes,
20 or hours of the administration of one or more other active therapeutic agents. For example, a unit dose of a compound of the invention can be administered first, followed within seconds or minutes by administration of a unit dose of one or more other active therapeutic agents. Alternatively, a unit dose of one or more other therapeutic agents can be administered first, followed by administration of a unit dose of a compound of the invention within seconds or
25 minutes. In some cases, it may be desirable to administer a unit dose of a compound of the invention first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of one or more other active therapeutic agents. In other cases, it may be desirable to administer a unit dose of one or more other active therapeutic agents first, followed, after a period of hours (e.g., 1-12 hours), by administration of a unit dose of a compound of the
30 invention.

The combination therapy may provide "synergy" and "synergistic", i.e. the effect achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a
35 combined formulation; (2) delivered by alternation or in parallel as separate formulations; or (3)

5 by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are administered or delivered sequentially, e.g. in separate tablets, pills or capsules, or by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e. serially, whereas in combination therapy, effective dosages of two or more active ingredients are
10 administered together. A synergistic anti-viral effect denotes an antiviral effect which is greater than the predicted purely additive effects of the individual compounds of the combination.

Another embodiment provides for methods of treating *Pneumovirinae* virus infection in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of formula I or formulas 1-11, or a pharmaceutically acceptable salt, solvate, and/or
15 ester thereof.

Another embodiment provides for methods of treating *Pneumovirinae* virus infection in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of formula I or formulas 1-11, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent.

20 Another embodiment provides for methods of treating Human respiratory syncytial virus infection in a patient, comprising: administering to the patient a therapeutically effective amount of a compound of formula I or formulas 1-11, or a pharmaceutically acceptable salt, solvate, and/or ester thereof, and at least one additional active therapeutic agent.

Metabolites of the Compounds of the Invention

25 Also falling within the scope of this invention are the *in vivo* metabolic products of the compounds described herein, to the extent such products are novel and unobvious over the prior art. Such products may result for example from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the invention includes novel and unobvious compounds produced by a process
30 comprising contacting a compound of this invention with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products typically are identified by preparing a radiolabeled (e.g. ^{14}C or ^3H) compound of the invention, administering it parenterally in a detectable dose (e.g. greater than about 0.5 mg/kg) to an animal such as rat, mouse, guinea pig, monkey, or to man, allowing sufficient time for metabolism to occur

5 (typically about 30 seconds to 30 hours) and isolating its conversion products from the urine, blood or other biological samples. These products are easily isolated since they are labeled (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, e.g. by MS or NMR analysis. In general, analysis of metabolites is done in the same way as conventional drug metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for therapeutic dosing of the compounds of the invention even if they possess no HSV antiviral activity of their own.

10 Recipes and methods for determining stability of compounds in surrogate gastrointestinal secretions are known. Compounds are defined herein as stable in the gastrointestinal tract where less than about 50 mole percent of the protected groups are deprotected in surrogate intestinal or gastric juice upon incubation for 1 hour at 37°C. Simply because the compounds are stable to the gastrointestinal tract does not mean that they cannot be hydrolyzed *in vivo*. The prodrugs of the invention typically will be stable in the digestive system but may be substantially hydrolyzed to the parental drug in the digestive lumen, liver, lung or other metabolic organ, or within cells

20 in general.

Tissue Distribution

It has also been discovered that certain compounds of the invention show high lung to plasma ratios which may be beneficial for therapy. One particular group of compounds of the invention that demonstrate this property are compounds that include an amine functional group.

25

Examples.

Certain abbreviations and acronyms are used in describing the experimental details. Although most of these would be understood by one skilled in the art, Table 1 contains a list of many of these abbreviations and acronyms.

30

Table 1. List of abbreviations and acronyms.


Abbreviation	Meaning
Ac ₂ O	acetic anhydride

AIBN	2,2'-azobis(2-methylpropionitrile)
Bn	benzyl
BnBr	benzylbromide
BSA	bis(trimethylsilyl)acetamide
BzCl	benzoyl chloride
CDI	carbonyl diimidazole
DABCO	1,4-diazabicyclo [2.2.2] octane
DBN	1,5-diazabicyclo [4.3.0]non-5-ene
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DBU	1,5-diazabicyclo [5.4.0]undec-5-ene
DCA	dichloroacetamide
DCC	dicyclohexylcarbodiimide
DCM	dichloromethane
DIPEA	<i>N,N</i> -diisopropylethylamine
DMA	dimethylacetamide
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMTC1	dimethoxytrityl chloride
DMSO	dimethylsulfoxide
DMTr	4, 4'-dimethoxytrityl
DMF	dimethylformamide
EtOAc	ethyl acetate
ESI	electrospray ionization
HATU	2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate Methanaminium
HMDS	hexamethyldisilazane
HPLC	High pressure liquid chromatography
IPA	isopropyl alcohol
LDA	lithium diisopropylamide
LRMS	low resolution mass spectrum

MCPBA	meta-chloroperbenzoic acid
MeCN	acetonitrile
MeOH	methanol
MMTC	mono methoxytrityl chloride
m/z or m/e	mass to charge ratio
MH ⁺	mass plus 1
MH ⁻	mass minus 1
MsOH	methanesulfonic acid
MS or ms	mass spectrum
NBS	N-bromosuccinimide
Ph	phenyl
rt or r.t.	room temperature
TBAF	tetrabutylammonium fluoride
TMSCl	chlorotrimethylsilane
TMSBr	bromotrimethylsilane
TMSI	iodotrimethylsilane
TMSOTf	(trimethylsilyl)trifluoromethylsulfonate
TEA	triethylamine
TBA	tributylamine
TBAP	tributylammonium pyrophosphate
TBSCl	t-butyldimethylsilyl chloride
TEAB	triethylammonium bicarbonate
TFA	trifluoroacetic acid
TLC or tic	thin layer chromatography
Tr	triphenylmethyl
Tol	4-methylbenzoyl
Turbo Grignard	1:1 mixture of isopropylmagnesium chloride and lithium chloride
δ	parts per million down field from tetramethylsilane

5 The invention will now be illustrated by the preparation of the following non-limiting compounds of the invention. It is to be understood that individual steps described herein may be combined. It is also to be understood that separate batches of a compound may be combined and then carried forth in the next synthetic step.

10 Example 1. Procedure for the preparation of compounds of formulas 1-24.











In 50 mL, singled necked, round bottomed flask was placed (s)-N-(4-chloro-2-(2-(5-chloro-6-methylpyrazolo[1,5-a]pyrimidine-2-yl)piperidine-1-





15 carbonyl)phenyl)methanesulfonamide (**A1**), (1200 mg, 2.49 mmol) in DMA (10.8 mL). The amines (R-(Boc)) (0.12 mmol) were placed in separate 2-ml vials. Then, into each vial was dispensed a solution of **A1** (0.2 mL, 0.041 mmol) followed by 0.1 mL of K_2CO_3 (2M) or DIPEA. The resulting reaction mixtures were placed on a hot plate at 85 °C for 16 h. Then, to each reaction mixture was added EtOAc (4 mL), washed with saturated NaHCO_3 (2 mL x 2),






20 and concentrated in Genevac to give A2 as a solid. The crude product A2 was redissolved in dichloromethane (0.5 mL) followed by the addition of TFA (0.2 mL). After the reaction mixture was stirred at room temperature for 1 h, it was loaded onto the CUBCX column. The mixture






- 5 was washed with MeOH:EtOAc (1:4, 4 mL) and MeOH:dichloromethane (1:4, 4 mL), eluted with 7 N NH_4OMe :EtOAc (3:7, 4 mL), and concentrated to afford the final compound (i.e. compounds 1-24).



Compound Formula	Compound	calculated MW	observed MW
1		520.056	520.5
2		532.067	532.5
3		582.127	582.6

<p>4</p>		<p>608.165</p>	<p>608.6</p>
<p>5</p>		<p>520.056</p>	<p>520.2</p>
<p>6</p>		<p>532.067</p>	<p>532.5</p>
<p>7</p>		<p>546.094</p>	<p>546.6</p>
<p>8</p>		<p>582.127</p>	<p>582.5</p>

<p>9</p>		<p>590.103</p>	<p>590.6</p>
<p>10</p>		<p>546.094</p>	<p>546.6</p>
<p>11</p>		<p>546.094</p>	<p>546.6</p>
<p>12</p>		<p>590.103</p>	<p>590.6</p>


<p>13</p>		<p>652.218</p>	<p>652.7</p>
<p>14</p>		<p>576.076</p>	<p>576.5</p>
<p>15</p>		<p>560.121</p>	<p>560.6</p>
<p>16</p>		<p>558.105</p>	<p>558.6</p>
<p>17</p>		<p>622.192</p>	<p>622.7</p>

18		544.078	544.6
19		576.076	576.6
20		560.121	560.6
21		558.105	558.5
22		574.148	574.6

23		589.119	589.6
24		560.121	560.5

5

Example 2. General procedure for the preparation of compounds of formulas 25-111.












10






In 50 mL, singled necked, round bottomed flask was placed tert-butyl (S)-1-(6-methyl-2-((S)-piperidin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-ylcarbamate (**B1**) (2640 mg, 6.59 mmol) and TEA (1.83 mL, 13.2 mmol) in DMF (8.8 mL). The carboxylic acids **B2** (between 0.10 mmol and 0.50 mmol) were placed in 132 separate 2-ml vials. Then, into each vial was






5 dispensed a solution of B1 (0.050 mmol) followed by the addition of HATU (38 mg, 0.10 mmol). The resulting reaction mixtures were placed on an orbital shaker at room temperature for 16 h. Then, to each reaction mixture was added EtOAc (4 mL), washed with sat. NaHCO₃ (2 mL x 2), and concentrated in Genevac to give B3 as a solid. The crude product B3 was redissolved in dichloromethane (0.5 mL) followed by the addition of TFA (0.2 mL). After the
10 reaction mixture was stirred at room temperature for 1 h, it was loaded onto the CUBCX column. The mixture was washed with MeOH:EtOAc (1:4, 4 mL) and MeOH:dichloromethane (1:4, 4 mL), eluted with 7 N NH₄OMe :EtOAc (3:7, 4 mL), and concentrated to afford the final compound (i.e. compounds 25-1 11).






5






Compound formula	Structure	calculated MW	observed MW
25		436.535	437.6
26		423.496	424.5
27		434.544	435.6
28		492.933	493.5






29		513.621	514.6
30		456.953	457.5
31		458.61	459.6
32		447.518	448.6
33		454.525	455.5






34		490.652	491.7
35		486.542	487.6
36		486.542	487.6
37		506.504	507.6
38		444.543	445.6






39		454.525	455.5
40		478.988	479.6
41		456.554	457.6
42		481.604	482.6
43		444.543	445.5






44		490.505	491.5
45		483.595	484.6
46		455.566	456.6
47		486.58	487.6
48		480.544	481.6






54		444.543	445.6
55		474.573	475.6
56		478.988	479.6
57		509.561	510.6
58		496.634	497.2






<p>59</p>		<p>497.622</p>	<p>498.6</p>
<p>60</p>		<p>459.598</p>	<p>460.6</p>
<p>61</p>		<p>448.527</p>	<p>449.5</p>
<p>62</p>		<p>502.598</p>	<p>503.6</p>
<p>63</p>		<p>497.043</p>	<p>497.6</p>






64		458.57	459.6
65		475.553	476.5
66		488.636	489.6
67		514.624	515.6
68		455.566	456.6






69		438.963	439.5
70		435.532	436.6
71		423.496	424.5
72		454.525	455.6
73		440.498	441.5






74		474.396	474.5
75		454.525	455.5
76		484.407	484.5
77		436.535	437.5
78		436.535	437.6






79		440.498	441.5
80		481.604	482.6
81		423.496	424.2
82		456.953	457.5
83		439.951	440.5




84		448.571	449.6
85		462.598	463.6
86		473.503	474.6
87		432.572	433.6
88		432.572	433.6

89		452.99	453.5
90		497.446	497.5
91		436.535	437.6
92		448.571	449.6
93		439.951	440.5

94		484.604	485.6
95		504.038	504.6
96		455.566	456.6
97		484.604	485.6
98		458.57	459.6


99		539.505	539.5
100		474.396	474.5
101		445.531	446.5
102		506.014	506.6
103		466.608	467.5

104		507.948	508.5
105		507.948	508.5
106		506.96	507.5
107		453.522	454.6
108		444.543	445.2

109		490.505	491.6
110		433.56	434.5
111		452.534	453.5

5

Example 3. General procedure for the preparation of compounds 112-191.













10






In 50 mL, single necked, round bottomed flask was placed tert-butyl (3S,4R)-4-hydroxy-1-(6-methyl-2-((S)-piperidin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidine-3-carbonitrile (**C1**) (1932 mg, 5.90 mmol) and TEA (1.64 mL, 11.8 mmol) in DMF (13 mL). The carboxylic acids **C2** (0.040 mmol) were placed in separate 2-ml vials. Then, into each vial was






- 5 dispensed a solution of **CI** (0.037 mmol) followed by the addition of HATU (15.2 mg, 0.04 mmol). The resulting reaction mixtures were placed on an orbital shaker at room temperature for 16 h. Then, to each reaction mixture was added EtOAc (4 mL), washed with sat. NaHCO₃ (2 mL x 2), and concentrated in Genevac. Then, it was loaded onto the CUSIL column, washed with EtOAc:Hexane (1:1, 4 mL), eluted with MeOH:EtOAc (5:95, 3 mL), and concentrated in
- 10 Genevac to give the final compound (i.e. compounds 112-191).






Compound formula	Compound	calculated MW	observed MW
112		523.44	523.4
113		460.538	461.5
114		518.927	519.4






<p>115</p>		<p>465.945</p>	<p>466.4</p>
<p>116</p>		<p>482.947</p>	<p>483.4</p>
<p>117</p>		<p>473.512</p>	<p>474.5</p>
<p>118</p>		<p>480.519</p>	<p>481.5</p>
<p>119</p>		<p>516.646</p>	<p>517.6</p>






<p>120</p>		<p>512.536</p>	<p>513.5</p>
<p>121</p>		<p>512.536</p>	<p>513.5</p>
<p>122</p>		<p>532.498</p>	<p>533.5</p>
<p>123</p>		<p>470.537</p>	<p>471.5</p>
<p>124</p>		<p>480.519</p>	<p>481.5</p>






125		507.598	508.5
126		478.528	479.5
127		482.548	483.5
128		507.598	508.5
129		512.574	513.5






<p>130</p>		<p>506.538</p>	<p>507.5</p>
<p>131</p>		<p>478.984</p>	<p>479.5</p>
<p>132</p>		<p>500.603</p>	<p>501.5</p>
<p>133</p>		<p>499.55</p>	<p>500.5</p>
<p>134</p>		<p>511.63</p>	<p>512.5</p>






<p>135</p>		<p>470.537</p>	<p>471.5</p>
<p>136</p>		<p>504.982</p>	<p>505.4</p>
<p>137</p>		<p>535.555</p>	<p>536.5</p>
<p>138</p>		<p>522.628</p>	<p>523.5</p>
<p>139</p>		<p>485.592</p>	<p>486.5</p>






140		474.521	475.5
141		528.592	528.5
142		523.037	523.5
143		484.564	485.5
144		514.63	515.6






<p>145</p>		<p>540.618</p>	<p>541.5</p>
<p>146</p>		<p>464.957</p>	<p>465.4</p>
<p>147</p>		<p>461.526</p>	<p>462.4</p>
<p>148</p>		<p>499.497</p>	<p>500.5</p>
<p>149</p>		<p>480.519</p>	<p>481.5</p>






150		466.492	467.5
151		469.549	470.5
152		500.39	500.5
153		480.519	481.5
154		510.401	510.4






155		462.529	463.5
156		462.529	463.5
157		466.492	467.2
158		507.598	508.5
159		445.527	446.2






<p>160</p>		<p>449.49</p>	<p>450.2</p>
<p>161</p>		<p>557.396</p>	<p>558.1</p>
<p>162</p>		<p>484.604</p>	<p>485.3</p>
<p>163</p>		<p>481.56</p>	<p>482.2</p>
<p>164</p>		<p>456.51</p>	<p>457.2</p>



165		474.565	475.2
166		460.538	462.2
167		488.592	489.5
168		499.497	500.5
169		458.566	459.5

170		478.984	479.5
171		523.44	523.4
172		462.529	463.5
173		474.565	475.5
174		510.598	511.5

<p>175</p>		<p>470.537</p>	<p>471.5</p>
<p>176</p>		<p>445.527</p>	<p>446.5</p>
<p>177</p>		<p>499.402</p>	<p>499.4</p>
<p>178</p>		<p>466.564</p>	<p>467.5</p>
<p>179</p>		<p>445.527</p>	<p>446.5</p>

180		506.61	507.5
181		436.538	437.4
182		482.976	483.5
183		490.63	491.5
184		498.509	499.5

185		483.576	484.5
186		482.947	483.4
187		484.564	485.5
188		494.983	495.5
189		502.501	503.5

190		497.003	497.5
191		513.606	514.5

5

Example 4: Procedure for the preparation of intermediate A1.

Intermediate 1:



10

N-Boc-(5)-piperidine-2-carboxylic acid (5.0 g, 22 mmol) in DMF (100 mL) was treated with Cs_2CO_3 (3.5 g, 10.9 mmol) and MeI (1.5 mL, 24 mmol). The mixture was stirred for 4 hours and diluted with MTBE (250 mL). The mixture was washed with water (2 x 100 mL) and saturated sodium chloride solution (1 x 100 mL). The solution was dried over anhydrous sodium sulfate and concentrated to afford the ester intermediate **1** which was used without further purification.

15


^1H NMR (CDCl_3 , 300MHz): δ 4.80 (m, 1H), 3.97 (m, 1H), 3.73 (s, 3H), 2.93 (m, 1H), 2.18 (app d, $J = 13.2$ Hz, 1H), 1.67 (m, 2H), 1.45 (br s, 10H), 1.20 (app t, $J = 13.5$ Hz, 1H).

$R_f = 0.90$ (30% EtOAc-hexanes).

20

Intermediate 2:

5



10

(S)-1-Boc-piperidine-2-carboxylic acid (25 g, 109 mmol, Sigma-Aldrich) in DMF (500 mL) was treated sequentially with MeNHOMe •HCl (11.2 g, 115 mmol), N-methylmorpholine (36 mL, 327 mmol), HOBt (16.2 g, 120 mmol), and EDCI (23 g, 120 mmol) and stirred for 18 h.

15

The solution was diluted with EtOAc (1000 mL) and washed with H₂O (2 x 500 mL) and saturated NaCl solution (500 mL). The solution was dried over MgSO₄, filtered and concentrated. The residue was subjected to a 330 g SiO₂ Combiflash High Performance Gold column (0-100% EtOAc-hexanes gradient) to afford the Weinreb amide intermediate **2**:

¹H NMR (CDCl₃, 300MHz): δ 5.06 (br m, 1H), 3.93 (br m, 1H), 3.77 (br s, 3H), 3.18 (s, 3H), 2.01 (app d, *J* = 13.5 Hz, 1H), 1.71 (m, 4H), 1.45 (s, 9H).


20

LCMS (ESI) *m/z* 273 [M + H]⁺, *t_R* = 2.31 min.

HPLC (RP: 6-98% MeCN-H₂O gradient, 0.05% TFA modifier) *t_R* = 4.423 min.

R_f = 0.60 (50% EtOAc-hexanes).

20 Intermediate 3 :



25

To a solution of acetonitrile (5 ml, 93.8 mmol) in dry THF (50 ml) at -78 °C was added dropwise NaN(TMS)₂ (34 ml, 68 mmol, 2M in hexanes). The solution was warmed up to -40 °C and stirred for 20 min. The solution was then cooled down to -78 °C and a solution of the ester (Intermediate **1**) (7.6 g, 31.1 mmol) in THF (20 ml) was added dropwise. The solution was warmed up to -40 °C and stirred for 2 h. The solution was then cooled down to -78 °C and a solution of acetic acid (4.8 ml, 80 mmol) in THF (20 ml) added dropwise. The solution was then warmed to RT and volatiles were removed under reduced pressure at 40 °C. The resulting

5 residue was dissolved in EtOAc (300 mL) and the organic phase was washed with brine twice. Volatiles were removed under reduced pressure at 40 °C.

¹H NMR (DMSO, 300 MHz) δ 4.63 (br s, 1H), 4.18-4.13 (m, 1H), 3.82-3.78 (m, 1H), 3.65 (s, 2H), 2.85-2.63 (m, 1H), 1.65-1.52 (m, 9H), 1.38 (s, 9H).

LCMS *m/z* : 153 [M-Boc group+H], *t_R* = 2.50 min.

10

The residue was dissolved in EtOH (150 ml) and hydrazine acetate (4.5 g, 47 mmol) was added. The solution was stirred for 16 h at RT. Volatiles were removed under reduced pressure at 40 °C, EtOAc added (200 ml) and the organic phase washed with aqueous dilute NaHCO₃, then H₂O followed by brine. Volatiles were removed under reduced pressure at 40 °C, the

15 resulting residue was purified by silica gel column (DCM/ MeOH, gradient from 0% to 20%) to afford the product intermediate.

LCMS *m/z* [M+H]⁺ C₁₃H₂₂N₄O₂ requires: 266.34. Found 266.84.


HPLC (min, purity) *t_R* = 2.13, 100%.

¹H NMR (DMSO, 300 MHz) 11.20 (br s, 1 H), 5.09 (m, 1H), 5.07 (s, 1H), 4.67 (br s, 2H), 3.81 (app d, *J* = 12.0 Hz, 1H), 2.72 (app br t, *J* = 12.0 Hz, 1H), 2.08 (app d, *J* = 12.9 Hz, 1H), 1.57 (m, 4H), 1.39 (s, 9H); MS (ESI) *m/z* 267 [M + H]⁺, *t_R* = 1.97 min. (3.5min method).; HPLC (Chiral: Chiralpak AD-H, isocratic «-heptane-isopropanol 70:30). *t_R* (desired) = 22.42 min, *t_R* (enantiomer of desired isomer) = 25.67 min; %ee = 93.

20

25

Intermediate 3 via Weinreb amide:



MeCN (3.20 mL, 60.7 mmol) in THF (50 mL) was cooled to -78 °C under Ar. NaHMDS

30 solution (1.0 M in THF, 36.8 mL, 36.8 mmol) was added dropwise over 5 min, during which time an off-white suspension had formed. The suspension was warmed to -20°C and stirred for 20 min. The suspension was cooled to -78 °C and transferred via cannula to the Weinreb amide

5 intermediate **2** (5.02 g, 18.4 mmol) in THF (50 mL) at -78 °C over 5 min. The suspension is warmed to -45 °C and stirred for 3 h, during which time the suspension became a yellow solution. The solution was cooled to -78 °C and AcOH (4.2 mL in 10 mL THF, 73.6 mmol) was added dropwise. The solution was warmed to room temperature and diluted with EtOAc (100 mL). The solution was washed with H₂O (50 mL) and saturated NaCl solution (50 mL). The solution was dried over MgSO₄ and concentrated to afford the cyano ketone which was used
 10 without further purification.


The crude α -cyano ketone was used in the next reaction with hydrazine acetate to synthesize desired amino pyrazole intermediate **3** as described above.

MS (ESI) *m/z* 267 [M + H]⁺, *t_R* = 1.81 min.

15 HPLC (RP: 6-98% MeCN-H₂O gradient, 0.05% TFA modifier) *t_R* = 3.212 min (>95% purity @ 254 nM).

HPLC (Chiral: Chiralpak AD-H 250 4.6 mm, 5 micron; isocratic n-heptane-isopropanol 70:30) *t_R* (a isomer, desired) = 22.35 min, *t_R* (b isomer) = 25.78 min; α = 1.15; %ee = >90%.

20 Intermediate 4:



(E)-ethyl-3-Ethoxy-2-methylacrylate (Intermediate **32**) (11.8 g, 67.6 mmol) and Cs₂CO₃
 25 (22.0 g, 67.6 mmol) were added to a solution of intermediate **3** (12.0 g, 45.1 mmol) at room temperature and the reaction mixture was heated to 130 °C. After 17 h, the reaction mixture was allowed to cool to room temperature and was concentrated under reduced pressure. The crude residue was diluted with ethyl acetate (250 mL) and was filtered. The resulting filtrate was concentrated under reduced pressure and the residue was purified via SiO₂ column
 30 chromatography (330 g SiO₂ Combiflash HP Gold Column, 0-100% ethyl acetate/hexanes) to afford intermediate **4**.

5 ¹H NMR (CDCl₃, 400MHz): δ 12.01 (br s, 1H), 7.99 (s, 1H), 5.73 (s, 1H), 5.42 (br s, 1H), 4.01 (br d, *J* = 12.2 Hz, 1H), 2.81 (br t, *J* = 11.2 Hz, 1H), 2.29 (d, *J* = 13.5 Hz, 1H), 2.07 (d, *J* = 1.1 Hz, 3H), 1.87 - 1.69 (m, 1H), 1.68 - 1.41 (m, 4H), 1.48 (s, 9H).

¹³C NMR (CDCl₃, 100MHz): δ 162.87, 156.34, 155.43, 140.16, 135.00, 113.29, 86.50, 79.75, 28.41, 27.79, 25.27, 21.00, 19.88, 13.38.

10 LCMS (ESI) *m/z* 333.0 [M + H]⁺, *t_R* = 2.24 min.


HPLC *t_R* (min), purity % : 3.969, 99%.

R_f = 0.50 (EtOAc).

Chiral HPLC, 98%ee (Chiralpak IC 5 mM, 4.6 150 mm, 10 -95% MeCN/ H₂O, 0.05% trifluoroacetic acid modifier) (<S)-isomer *t_R* = 22.234 min, (i?)-isomer *t_R* = 20.875 min.

15

Intermediate 5:



20

POCl₃ (5.60 mL, 59.8 mmol) was added to intermediate 4 (993.4 mg, 2.99 mmol) at room temperature and the reaction mixture was heated to 100 °C. After 2 h, the reaction mixture was allowed to cool to room temperature and was concentrated under reduced pressure to afford intermediate 5 which was used directly in the following step.


25 ¹H NMR (DMSO-J₆, 400MHz): δ 9.40 (br d, *J* = 7.6 Hz, 1H), 9.27-9.16 (m, 2H), 6.85 (s, 1H), 4.54 (t, *J* = 112.4 Hz, 1H), 3.32 (d, *J* = 12.8 Hz, 1H), 3.08 (q, *J* = 8.81 Hz, 1H), 2.33 (s, 3H), 2.23-2.14 (m, 1H), 1.92-1.61 (m, 5H).

LCMS (ESI) *m/z* 251.1 [M + H]⁺, *t_R* = 0.21 min.

HPLC *t_R* = 2.35 min.

30

Intermediate A1:



5

HATU (1.37 g, 3.59 mmol) was added to a solution of 5-chloro-2-(methylsulfonamido) benzoic acid (823 mg, 3.29 mmol) in DMF (15.0 mL), and the reaction mixture was stirred at room temperature. After 1 h, a solution of crude intermediate **5** (220 mg, 2.99 mmol) in DMF (1 mL) was added followed by the addition of triethylamine (2.00 mL, 14.3 mmol), and the reaction mixture was stirred at room temperature for 19 h. The reaction mixture was partitioned between ethyl acetate (250 mL) and saturated aqueous sodium bicarbonate solution (200 mL), and the layers were separated. The organic layer was washed with saturated aqueous sodium bicarbonate solution (200 mL) and saturated sodium chloride solution (200 mL), was dried over Na_2SO_4 , and was concentrated under reduced pressure. The crude residue was purified via SiO_2 column chromatography (12 g SiO_2 Combiflash HP Gold Column, 0-100% ethyl acetate/hexanes) to afford intermediate **A1** (736.2 mg, 51% (2-steps)) as a white solid.

^1H NMR (CDCl_3 , 400MHz): δ 10.05 (br s, 0.2H), 9.13 (br s, 1H), 8.95 (br s, 1H), 8.81 (br s, 0.2H), 7.70 (d, $J = 8.8$ Hz, 1H), 7.56 (d, $J = 8.8$ Hz, 0.2H), 7.40 (dd, $J = 8.8, 2.4$ Hz, 1H), 7.33 (d, $J = 2.4$ Hz, 1H), 7.31 (d, $J = 4.4$ Hz, 0.2H), 6.45 (s, 1H), 6.40 (br s, 0.2H), 6.28 (br d, $J = 4.4$ Hz, 1H), 5.01 (br s, 0.2H), 4.54 (br d, $J = 14.0$ Hz, 0.2H), 3.35 (br d, $J = 13.2$ Hz, 1H), 3.15-3.03 (m, 1H), 2.92 (s, 3H), 2.39 (s, 3H), 2.13-1.98 (m, 1H), 1.90-1.59 (m, 2H), 1.59-1.31 (m, 3H).

^{13}C NMR (CDCl_3 , 100MHz): δ 167.09, 156.12, 153.13, 147.86, 135.68, 131.79, 131.66, 131.38, 130.12, 125.91, 125.44, 117.08, 93.74, 47.65, 44.07, 39.81, 27.83, 25.47, 19.78, 16.90.

LCMS (ESI) m/z 482.1 $[\text{M} + \text{H}]^+$, $t_R = 2.79$ min.


HPLC t_R (min), purity %: 5.438, 99%.

$R_f = 0.47$ (50% EtOAc/hexanes).

Chiral HPLC, 99%ee (Chiralpak IC 5 mM, 4.6 150 mm, 10-95% MeCN/ H_2O , 0.05% trifluoroacetic acid modifier) (5)-isomer $t_R = 29.739$ min, (if)-isomer $t_R = 29.495$ min.

5

Example 5: Procedure for the preparation of intermediate B1.



10 To a solution of intermediate **5** (100.0 mg, 0.35 mmol) in MeOH (1.74 mL) was added (5)-tert-butyl pyrrolidin-3-ylcarbamate (648 mg, 3.48 mmol) and triethylamine (970 μ L, 6.96 mmol) at room temperature, and the reaction mixture was heated to 70 $^{\circ}$ C. After 4 h, the reaction mixture was allowed to cool to room temperature and was concentrated under reduced pressure. The crude residue was purified by preparatory HPLC (5-100% MeCN/H₂O, 0.1% trifluoroacetic acid modifier) to afford intermediate **B1**.


15

LCMS (ESI) m/z 401.23 [M + H]⁺, t_R = 1.86 min.

Example 6: Procedure for the preparation of intermediate CI.

20

Intermediate 7:




25

A mixture of cis/ trans tert-butyl 3-cyano-4-hydroxypyrrolidine-1-carboxylate was separated on a silica column (200-300) eluting with ethyl acetate:petroleum ether=1:10, ethyl acetate:petroleum ether = 1:5 to give intermediate 7 (earlier eluting peak, 30 g, 46%) as white solid.

TLC (Eluent: ethyl acetate:petroleum ether =1:1): Starting material cis/ trans mixture (R_f = 0.4 and 0.45).

5 Column: ChiralPak IC-H, 250 50mmI.D, mobile Phase: CO₂/ iPrOH (35% isocratic), retention time (**9a**) 1.94 min, retention time (**9b**): 2.73 min.

Intermediate 10a:



A solution of intermediate **9a** (16.3g, 0.036 mol) in CH₂Cl₂ (200mL) at r.t was added TBAF (8.0 g, 0.025 mol). The reaction mixture was stirred at r.t for 30 min, then diluted with CH₂Cl₂ (500 mL), and washed with saturated aq. NH₄Cl and brine, dried over MgSO₄, filtered, and concentrated. The crude product was purified by silica gel chromatography (petroleum ether: ethyl acetate =10:1 to 2:1) to afford intermediate **10a**.


15 TLC Information (10a) (Eluent: petroleum ether: ethyl acetate =1:1).

1. Starting material (R_f = 0.90).

2. Reaction Mixture (Product: R_f = 0.4).

20 ¹H NMR (**10a**): 400 MHz DMSO δ 4.60-4.58 (m, 1H), 3.87-3.79 (m, 1H), 3.69-3.64 (m, 1H), 3.56-3.49 (m, 2H), 2.9-3.1 (m, 1H), 1.4-1.5 (s, 9H).


Intermediate CI:



5 A solution of intermediate **17** (Example 18: prepared from 1g BOC intermediate **10a**) was dissolved in MeOH (10 ml), to the solution was added intermediate **5** (944 mg, 3.76 mmol) and NEt₃ (2 ml). The reaction mixture was heated at 70° overnight. The solvent was evaporated and the residue was purified with purified by combi-flash column chromatography (0-100% MeOH/DCM) to afford intermediate Cl.


10 LCMS (*m/z*) 327.40[M + H]⁺
MW 326.19.

Example 7: Preparation of phosphate (-P(O)OH₂) compound **192**.



15 Compound **192** can be prepared from intermediate **D1** (preparation described directly below) by treating a solution of **D1** in THF at about 0°C with POCl₃ and triethylamine. The reaction mixture can be stirred at about 0 °C for a period of time and quenched with triethylammonium bicarbonate buffer (1M). The mixture can then be concentrated and purified
20 by HPLC to give the desired product.

Compound **192** can also be prepared by the following protocol.



25 Intermediate **D1** (100mg, 0.18 mmol) was dissolved in THF (2 ml), and the reaction mixture was cooled to 0°C using an ice bath. To the above solution was added POCl₃ (110mg, 0.72 mmol) followed by pyridine (71mg, 0.9 mmol). The reaction mixture was stirred at 0°C for 30mins before it was quenched with saturated NaHCO₃. Washed the aqueous layer with DCM twice (20 mL), the aqueous layer was lyophilized and the residue was then purified by prep

5 HPLC (Gemini C18, 100 30 mm, 5 micron column) using a gradient of water/acetonitrile 0-100 to afford the title compound **192**.


LCMS (m/z) 638.12 [M+H], Tr = 2.87 min.

CALC. MW 638.03.

Accordingly one embodiment includes compound **192** and salts thereof, as well as
10 methods and intermediates that are useful for the preparation of compound **192** and intermediate **D1**.


In a similar manner the corresponding phosphate prodrugs (-P(O)OH₂) of the compounds
of formulas I, Ia, II, IIa, III, IIIa, IV and IVa can also be prepared. Accordingly, one
embodiment includes the phosphate compounds (-P(O)OH₂) of the compounds of formulas I, Ia,
15 II, IIa, III, IIIa, IV and IVa and salts thereof.

Preparation of intermediate D1.




The starting chloride intermediate **A1** (0.92g, 1.9 mmol) was dissolved in iPrOH (10 mL)
20 and treated with triethylamine (0.45 mL, 0.33 g) and the intermediate **17** (0.45 g) and then
heated to a gentle reflux under nitrogen. After heating overnight the solution was cooled and
diluted with ethyl acetate and brine. The organic layer was separated and concentrated under
reduced pressure. The crude product was purified by silica gel chromatography (50-100% ethyl
acetate in hexanes) to afford the product as a white solid (1.15 g). LCMS (ESI) m/z 558.2 [M +
25 H]⁺, t_R = 2.17 min.


Using similar procedures the corresponding compounds 193, 194 and 195 can also be
prepared. Accordingly, one embodiment includes compounds 193, 194 and 195 and salts
thereof.



193




194



195

5

Example 8: Preparation of compound 196.




10

Compound **196** was prepared according to the general procedure cited below LCMS (m/z)

446.04 $[M + H]^+$

MW 445.22

15 General Procedure




5

In 50 mL, singled necked, round bottomed flask was placed intermediate **CI** (1932 mg, 5.90 mmol) and TEA (1.64 mL, 11.8 mmol) in DMF (13 mL). The carboxylic acid (0.040 mmol) was placed in a separate 2-ml vial. Then, into the vial was dispensed a solution of intermediate **CI** (0.037 mmol) followed by the addition of HATU (15.2 mg, 0.04 mmol). The resulting reaction mixture was placed on an orbital shaker at room temperature for 16 h. Then, to the reaction mixture was added EtOAc (4 mL), washed with sat. NaHCO₃ (2 mL x 2), and concentrated in Genevac. The crude residue was loaded onto the CUSIL column, washed with EtOAc:Hexane (1:1, 4 mL), eluted with MeOH:EtOAc (5:95, 3 mL), and concentrated in Genevac to give the final compound **196**.

15

Example 9: Preparation of intermediate **11**.




To an oven dried 50 mL round-bottom flask, methyl 2-bromo-5-methylbenzoate (352 mg, 1.54 mmol), sultam (236 mg, 1.95 mmol), cesium carbonate (732 mg, 2.25 mmol), palladium acetate (40.4 mg, 0.18 mmol), and Xanphos (136 mg, 0.235 mmol) were added and flask was placed under argon. Reagents were suspended in 8 mL of anhydrous dioxane and mixture was heated at 100 °C overnight. After cooling to room temperature, reaction mixture was filtered, washing with ethyl acetate. Combined filtrate was concentrated under reduced pressure and resulting film was purified by silica gel column chromatography (25-100% Ethyl Acetate in Hexanes) to yield intermediate **11**.


25

¹H-NMR (DMSO, 400 MHz): δ 7.75 (d, 1H), 7.44 (m, 1H), 7.35 (m, 1H), 3.89 (s, 3H), 3.81 (t, 2H), 3.28 (t, 2H), 2.55 (m, 2H), 2.39 (s, 3H).

LCMS m/z [M+H]⁺ C₁₂H₁₅N₀₄S requires: 270.07. Found 270.12.

5 Example 10: Preparation of intermediate 12.

- Lithium hydroxide monohydrate (496 mg, 11.8 mmol) was added to a solution of intermediate **11** (316 mg, 1.17 mmol) in 22 mL of THF and 12 mL of water at room temperature. Reaction mixture was heated at 60 °C for two hours. After cooling to room temperature, reaction mixture was acidified with 40 mL of 1N HCl_(aq) and extracted with ethyl acetate (3 x 30 mL). The combined organic layers were washed with 50 mL of Brine, separated, dried (MgSO₄), filtered, and concentrated under reduced pressure to yield intermediate **12**.
- ¹H-NMR (DMSO, 400 MHz): δ 12.9 (s, 1H), 7.57 (d, J=1.6 Hz, 1H), 7.41-7.34 (m, 2H), 3.66 (t, J=6.8 Hz, 2H), 3.28 (m, 2H), 2.37 (m, 2H), 2.33 (s, 3H).
- 15 LCMS m/z [M+H]⁺ C₁₁H₁₃N₄O₄S requires: 254.06. Found 254.18.

Example 11: Preparation of compound 197.


- 20 HATU (70 mg, 0.185 mmol) was added to a solution of intermediate **12** (40 mg, 0.16 mmol) in 3 mL of anhydrous DMF at room temperature. After 60 minutes of stirring, intermediate **CI** (40 mg, 0.123 mmol) was added followed immediately by triethylamine (0.030 mL, 0.213 mmol). Reaction mixture stirred at room temperature overnight under argon. Mixture was then poured into 30 mL of H₂O and extracted three times with 30 mL of ethyl acetate. The combined organic layers were washed with 50 mL brine, dried (MgSO₄), filtered, and concentrated under reduced pressure leaving a residue. Product was purified by prep HPLC
- 25

5 (15-100% Acetonitrile (with 0.1% trifluoroacetic acid) in water (with 0.1% trifluoroacetic acid))
to yield compound **197** as a trifluoroacetic acid salt, after lyophilization.

LCMS m/z [M+H]⁺ C₂₈H₃₃N₇O₄S requires: 564.23. Found 564.13.

HPLC Tr (min), purity % : 5.33, 99%

10 Example 12: Preparation of intermediate **13**.




Step 1: Sodium azide (158 mg, 2.43 mmol) was added to a solution of methyl 2-(bromomethyl)-5-chlorobenzoate (518 mg, 1.97 mmol) in 3 mL of DMF at room temperature. After stirring overnight, reaction mixture was quenched with 25 mL of water. Aqueous was
15 extracted with ethyl acetate (3x30 mL) and combined organics were washed with water (2x40 mL) and 50 mL of brine. Organics were dried (Na₂SO₄), filtered, and concentrated under reduced pressure to yield methyl 2-(azidomethyl)-5-chlorobenzoate which was used in the next step without further purification.

Step 2: Lithium hydroxide monohydrate (794 mg, 18.9 mmol) was added to a solution of
20 methyl 2-(azidomethyl)-5-chlorobenzoate (426 mg, 1.88 mmol), from the previous step, in 27 mL of 1:1:1 THF:methanol: water at room temperature. After stirring overnight, reaction mixture was quenched with 20 mL of 2N HCl_(aq), and extracted with ethyl acetate (3x30 mL). Combined organics were washed with brine, dried (MgSO₄), filtered, and concentrated under reduced pressure to yield intermediate **13**.

25 ¹H-NMR (DMSO, 400 MHz): 7.88 (m, 1H), 7.70-7.65 (m, 1H), 7.54 (m, 1H), 4.78 (s, 2H).

Example 13: Preparation of compound **198**.




Following the procedure for the synthesis of compound **197**, beginning with intermediate **13** (36.2 mg, 0.171 mmol) and intermediate **CI** (40 mg, 0.123 mmol), compound **198** was synthesized as a trifluoroacetic acid salt after lyophilization.

LCMS m/z $[M+H]^+$ $C_{25}H_{26}ClN_9O_2$ requires: 520.19. Found 520.03.

10 HPLC Tr (min), purity % : 6.34, 97%.

Example 14: Preparation of intermediate 14.




To a solution of methyl 2-amino-5-bromobenzoate (7.38 g, 32.0 mmol) and pyridine (6.3 mL, 81.5 mmol) in 100 mL of anhydrous CH_2Cl_2 , was added slowly methane sulfonylchloride (6.5 mL, 79.9 mmol). After stirring overnight, reaction mixture was quenched with 100 mL of 1N HCl (aq). Aqueous mixture was extracted with ethyl acetate (3x120 mL) and combined organic layers were washed 200 mL brine. Organics were dried ($MgSO_4$), filtered, and concentrated under reduced pressure to yield intermediate **14**. Silica gel column chromatography (0-30% Ethyl Acetate in Hexanes), yielded intermediate **14**.

1H -NMR ($CDCl_3$, 300 MHz): δ 10.4 (s, 1H), 8.22 (s, 1H), 7.63 (s, 2H), 3.96 (s, 3H), 3.05 (s, 3H)

LCMS m/z $[M+H]^+$ $C_9H_{10}BrNO_4S$ requires: 307.95. Found 308.06.

25

Example 15: Preparation of intermediate 15.





10 A 2.65 M solution of NaOH in water (2.65 mL, 7.02 mmol) was added to a solution of intermediate **14** in 9 mL of THF with strong stirring. The reaction mixture was stirred at room temperature over night. The mixture was then acidified with 10 mL of 1N HCl and extracted with ethyl acetate (3 x 20 mL). The combined organic layers were washed 30 mL of brine, separated, dried (MgSO₄), filtered, and concentrated under reduced pressure to yield intermediate **15**.

1H-NMR (DMSO, 300 MHz): δ 10.6 (s, 1H), 8.05 (s, 1H), 7.79 (d, 1H), 7.55 (d, 1H), 3.18 (s, 3H).

15 LCMS m/z $[M+H]^+$ C₈H₈BrNO₄S requires: 291.94. Found 291.90.

Example 16: Preparation of compound **199**.






5

DMF (0.070 mL, 0.908 mmol) was added slowly to a suspension of 5-methyl-2-(methylsulfonamido)benzoic acid (1.01 g, 4.59 mmol) and oxalyl chloride (1.6 mL, 18.3 mmol) in 11 mL of anhydrous dichloromethane. After 3 hours, reaction mixture was concentrated and dried *in-vacuo* to yield intermediate **16** which was used in the next step without further purification.

10

¹H-NMR (CDCl₃, 400 MHz): δ 10.2 (s, 1H), 7.92 (s, 1H), 7.64 (m, 1H), 7.39 (m, 1H), 3.03 (s, 3H), 2.35 (s, 3H).

Example 18: Preparation of intermediate 17.




15

The BOC pyrrolidine intermediate **10a** (1g, 4.7 mmol) was added to HCOOH (5 ml) and was heated at 40°C for 2h. The solvent was evaporated under reduced pressure and preheated IPA (100°C) was added to dissolve the residue, white precipitate formed after the IPA solution cooled down. The product was filtered and washed with IPA to give intermediate **17** that was used without further purification in subsequent reactions.

20

Example 19: Preparation of intermediate 18.



5


Triethylamine (0.58 mL, 4.16 mmol) was added slowly to a mixture of intermediate **17** (479 mg, 2.01 mmol) and intermediate **5** (573 mg, 2.00 mmol) in 10 mL of dichloromethane under argon at 0 °C. After 3 hours, LC/MS indicated full conversion to desired product. The reaction mixture was concentrated and dried *in-vacuo* to yield intermediate **18** that was used in the next steps without further purification.

10

LCMS m/z $[M+H]^+$ $C_{21}H_{24}ClN_5O_3S$ requires: 462.13. Found 462.32.

Example 20: Preparation of compound 200.

15




20

Triethylamine (0.100 mL, 0.717 mmol) was added to a mixture of intermediate **18** (102 mg, 0.221 mmol) and intermediate **17** (50 mg, 0.316 mmol) in 3 mL of methanol at room temperature. After heating at 75 °C overnight, reaction mixture was cooled to room temperature and concentrated under reduced pressure. The remaining residue was purified by silica gel column chromatography (10-75% ethyl acetate in hexanes) to yield compound **200**.

LCMS m/z $[M+H]^+$ $C_{26}H_{31}N_7O_4S$ requires: 538.22. Found 538.01.

HPLC Tr (min), purity % : 6.10, 97%.


5

Example 21: Preparation of compound 201.

2-Methoxy-5-chlorobenzoic acid (29mg, 0.17 mmol) and HATU (76mg, 0.2 mmol) were dissolved in DMF (2 ml). The reaction mixture was stirred at room temperature for 10mins. To the above solution was added intermediate CI (35mg, 0.1 mmol) and NEt_3 (55 μl). The reaction was stirred at room temperature for 30mins and was quenched with brine (10 ml) and then extracted with EtOAc (20 ml). The organic layer was washed with brine twice (10 ml) and then was evaporated under reduced pressure. The residue was purified with prep HPLC (0-100% $\text{CH}_3\text{CN}/\text{H}_2\text{O}$) to afford compound 201.

LCMS (m/z) 495.17 $[\text{M} + \text{H}]^+$.

MW 493.97.


Example 22: Preparation of compound 202.

The title compound was prepared according to the procedure for compound 201 starting from intermediate CI and 2-methyl-5-chlorobenzoic acid.

LCMS (m/z) 479.20 $[\text{M} + \text{H}]^+$.

MW 477.97.

Example 23: Preparation of intermediate 19.




2- Amino-5-methylbenzoic acid (316 mg, 2.09 mmol), HATU (992mg, 2.61 mmol) were dissolved in anhydrous DMF (2 ml). After activation for 1 hour, intermediate **5** (500 mg, 1.74 mmol) and triethylamine (0.7 ml) was added to the above solution. The reaction was stirred under nitrogen for 2 hours. Solvents were removed by rotary evaporation. The residue was purified with silica gel column chromatography to provide intermediate **19**.

LCMS m/z $[M+H]^+$ $C_{20}H_{22}ClN_5O$ requires: 384.15. Found 383.99.

HPLC Tr (min), purity % : 2.00, 98%.

15 Example 24: Preparation of intermediate 20.




Intermediate **19** (320mg, 0.84mmol) was dissolved in pyridine (2 ml). Then acetyl chloride (78mg, 1.0 mmol) was added to the above solution. The reaction was stirred under nitrogen for 30mins. Solvents were removed by rotary evaporation. The residue was purified with silica gel column chromatography to provide intermediate **20**.

LCMS m/z $[M+H]^+$ $C_{22}H_{24}ClN_5O_2$ requires: 426.16. Found 425.89.

HPLC Tr (min), purity % : 2.40, 98%.

25

Example 25: Preparation of compound 203.



5

The title compound was prepared in 25% yield according to the procedure for compound **201** starting from intermediate **20** and the cis and trans mixture of 3-cyano-4-hydroxypyrrolidine. Compound **203** was obtained as a mixture of all 4 isomers at the


10

pyrrolidine.
LCMS (m/z) 501.87 [M + H]⁺.

MW 500.58.

Example 26: Preparation of compound 204.

15




20

Intermediate **D1** (Example 7) (0.100g) was dissolved in DMF (2 ml) and Boc-L-valine (0.075g), DMAP (0.02g) and HATU (0.096g) added with stirring at room temperature. After 5 h, the solution was diluted with MeCN (2 ml) and water (4 ml) and was purified with preparatory HPLC to yield the corresponding ester. The ester was dissolved in dioxane (2 ml) and HCl (4M in dioxane, 2 ml) was added with stirring. After 2h, volatiles were removed at room temperature and the crude product was purified with preparatory HPLC to yield compound **204**.

25 LCMS (m/z) 657.23, Tr = 1.64 min.

MW 657.18.

5 Example 27: Preparation of compound 205.


10 The title compound was prepared in an analogous way as described for compound **204** above utilizing proline to afford the product compound **205**.

LCMS (m/z) 655.21, Tr = 1.58 min.

CALC. MW 655.17.

Example 28: Preparation of compound 206.

15




Intermediate **DI** (Example 7) (200mg, 0.36 mmol) was dissolved in DCM (2 ml), to the above solution was added Boc-Glycine (90mg, 0.6 mmol) and EDCI (111mg, 0.58 mmol) followed by DMAP (23mg, 0.18 mmol). The reaction mixture was stirred at room temperature for 3 hours. To the above reaction mixture was added TFA (0.1 ml) and stirred at room temperature for 1 h. The solvent was evaporated under reduced pressure and purified with prep HPLC (Gemini C18, 100 30 mm, 5 micron column) using a gradient of water/acetonitrile 0-100 to afford the title compound **206**.

25 LCMS (m/z) 615.24 [M+H], Tr = 2.67 min.

CALC. MW 615.22.

Example 29: Preparation of compound 207.

5



Into oven-dried, argon purged flask were placed intermediate **DI** (Example 7) (140 mg, 0.25 mmol), succinic anhydride (55 mg, 0.55 mmol) and catalytic amount (1 mg) of 4-dimethylaminopyridine. The flask was sealed with septa and repurged with argon three times.

10 Dry tetrahydrofuran (20 mL) was added into the reaction mixture via syringe, followed by diisopropylethylamine (72 mg, 0.56 mmol). This reaction mixture was heated for 14 hours at 70 °C to achieve the full conversion. The solvent was evaporated, the residue was dissolved in ethyl acetate (30 mL) and this solution was washed twice with 10% solution of citric acid (30 mL), brine (30 mL), dried over magnesium sulfate, filtered and evaporated. The residue was

15 purified by silica gel chromatography using gradient of dichloromethane/methanol (from 10/0 to 9/1) to afford the title compound **207**.

TLC R_f = 0.49 (10% methanol in dichloromethane, silica gel).


LCMS (m/z) 658.2 (100%) and 660.2 (44%) [$M + H$]⁺; 655.8 (100%) and 657.9 (40%) [$M - H$]⁺, Tr = 3.96 min., purity >99%.

20 For $C_{29}H_{32}ClN_7O_7S$ Calc. MW 657.2 (100%) and 659.2 (37%).

¹H-NMR (CDCl₃, 400 MHz): 9.19 (s, 1H), 8.73 (s, 1H), 7.61 (d, J = 8.8 Hz, 1H), 7.31 (d, J = 8.8 Hz, 1H), 7.25 (s, 1H), 6.14 (s, 1H), 6.03 (s, 1H), 5.51 (s, 1H), 5.23 (s, 1H), 4.04 (m, 3H), 3.75 (m, 1H), 3.34 (m, 1H), 3.19 (m, 1H), 3.06 (m, 1H), 2.84 (m, 2H), 2.66 (m, 3H), 2.29 (m, 3H), 2.00-1.22 (m, 6H).

25

Example 30: Preparation of compound 208.



5

Into oven-dried, argon purged flask were placed intermediate **D1** (Example 7) (56 mg, 0.1 mmol), niacin (19 mg, 0.15 mmol) and 4-dimethylaminopyridine (18 mg, 0.15 mmol). The flask was sealed with septa and repurged with argon three times. Dry dichloromethane (20 mL) was added into the reaction mixture via syringe, followed by 1-ethyl-3-(3-

10 dimethylaminopropyl)carbodiimide hydrochloride (67 mg, 0.3 mmol). This reaction mixture was repurged with argon three times and stirred at room temperature for 10 minutes to achieve the full conversion. The solvent was evaporated, the residue was dissolved in ethyl acetate (30 mL) and this solution was washed twice with 5% solution of citric acid (30 mL), saturated solution of NaHCO₃ (30 mL), water (30 mL), brine (30 mL), dried over magnesium sulfate,

15 filtered and evaporated. The residue was purified by silica gel chromatography using gradient of dichloromethane/methanol (from 10/0 to 95/5) to afford the title compound.

TLC R_f = 0.78 (5% methanol in dichloromethane, silica gel).


LCMS (*m/z*) 663.3 (100%) and 665.2 (43%) [M + H]⁺; 661.2 (100%) and 663.2 (38%) [M - H]⁺,
T_R = 4.19 min., purity >99%.

20 For C₃₁H₃₁C₁N₈O₅S Calc. MW 662.2 (100%) and 664.2 (38%).

¹H-NMR (CDCl₃, 400 MHz): 9.21 (s, 1H), 8.77 (d, *J* = 4.2 Hz, 1H), 8.74 (s, 1H), 8.31 (d, *J* = 7.9 Hz, 1H), 7.61 (d, *J* = 8.8 Hz, 1H), 7.39 (dd, *J* = 7.9 Hz, *J* = 4.2 Hz, 1H), 7.31 (d, *J* = 8.8 Hz, 1H), 7.24 (s, 1H), 6.14 (m, 1H), 6.02 (s, 1H), 5.73 (m, 1H), 4.16 (m, 4H), 3.88 (m, 1H), 3.47 (m, 1H), 3.00 (s, 1H), 3.06 (m, 1H), 2.82 (s, 3H), 2.31 (s, 3H), 1.60-1.22 (m, 6H).


25

Example 31: Preparation of Compound 209.



- 5 Into oven-dried, argon purged flask were placed compound **207** (123 mg, 0.19 mmol) and 2-morpholinoethanamine (27 mg, 0.21 mmol). The flask was sealed with septa and repurged with argon three times. Reaction flask was placed in an ice-bath. 6 mL of dry acetonitrile were added into the reaction mixture via syringe, followed by diisopropylethylamine (97 mg, 0.75 mmol) and 0-(7-azabenzotriazol-1-yl)-*N,N,N',iV*-tetramethyluronium hexafluorophosphate (107 mg, 0.28 mmol). This reaction mixture was repurged three times with argon, the ice-bath was removed and the reaction mixture was stirred at room temperature for 5 minutes to achieve the full conversion. The solvent was evaporated and the residue was purified by silica gel chromatography (triethylamine neutralized silica gel) using gradient of dichloromethane/methanol (from 10/0 to 9/1) to afford the title compound **209**.
- 10 TLC $R_f = 0.57$ (10% methanol in dichloromethane, silica gel).
 LCMS (m/z) 770.3 (100%) and 772.2 (51%) $[M + H]^+$; 768.2 (100%) and 770.1 (43%) $[M - H]^+$, Tr = 2.95 min., purity >99%.
 For $C_{35}H_{44}ClN_9O_7S$ Calc. MW 769.3 (100%) 771.3 (37%).
- 20 1H -NMR ($CDCl_3$, 400 MHz): 9.23 (s, 1H), 8.72 (s, 1H), 7.62 (d, $J = 8.7$ Hz, 1H), 7.31 (d, $J = 8.7$ Hz, 1H), 7.25 (m, 1H), 6.14 (m, 1H), 6.00 (s, 2H), 5.51 (m, 1H), 5.23 (s, 2H), 4.03 (m, 1H), 3.75 (m, 1H), 3.64 (m, 6H), 3.26 (m, 2H), 3.19 (m, 1H), 3.06 (m, 2H), 2.80 (s, 3H), 2.73 (m, 2H), 2.45 (m, 2H), 2.38 (m, 3H), 2.28 (s, 3H), 1.94-1.22 (m, 4H).

25 Example 32. Preparation of intermediate 32.



- 30 A solution of 1-ethoxy-propene (5.1 mL, 46 mmol) in pyridine (3.4 mL) was added slowly via addition funnel (~1 drop/sec) to neat trichloroacetyl chloride (4.7 mL, 42 mmol) at -10 °C under an argon atmosphere. The reaction mixture was then allowed to slowly warm to 23 °C. After 20 h, the reaction mixture was diluted with dichloromethane (50 mL) and the resulting mixture was washed with 0.01N HCl (3 50 mL) and brine (50 mL), was dried over

5 anhydrous sodium sulfate, and was concentrated under reduced pressure. To the crude residue was added sodium ethoxide (21 wt% in ethanol, 7.1 g, 44 mmol) slowly via syringe. After 30 min, the reaction mixture was partitioned between dichloromethane (500 mL) and water (500 mL). The phases were split and the aqueous layer was extracted with dichloromethane (500 mL). The combined organic extracts were dried over anhydrous sodium sulfate, and were
10 concentrated to afford intermediate **32**.

¹H-NMR (CDCl₃, 400 MHz): 7.28 (app s, 1H), 4.09 (q, *J* = 7.1 Hz, 2H), 3.96 (q, *J* = 7.1 Hz, 2H), 1.66 (s, 3H), 1.25 (t, *J* = 7.1 Hz, 3H), 1.20 (t, *J* = 7.1 Hz, 3H).

Antiviral Activity

15 Another embodiment relates to methods of inhibiting viral infections, comprising the step of treating a sample or subject suspected of needing such inhibition with a composition of the invention.

Samples suspected of containing a virus include natural or man-made materials such as living organisms; tissue or cell cultures; biological samples such as biological material samples
20 (blood, serum, urine, cerebrospinal fluid, tears, sputum, saliva, tissue samples, and the like); laboratory samples; food, water, or air samples; bioproduct samples such as extracts of cells, particularly recombinant cells synthesizing a desired glycoprotein; and the like. Typically the sample will be suspected of containing an organism which induces a viral infection, frequently a pathogenic organism such as a tumor virus. Samples can be contained in any medium including
25 water and organic solvent/water mixtures. Samples include living organisms such as humans, and manmade materials such as cell cultures.

If desired, the anti-virus activity of a compound described herein after application of the composition can be observed by any method including direct and indirect methods of detecting such activity. Quantitative, qualitative, and semi-quantitative methods of determining such
30 activity are all contemplated. Typically one of the screening methods described above are applied, however, any other method such as observation of the physiological properties of a living organism are also applicable.

The antiviral activity of a compound described herein can be measured using standard screening protocols that are known. For example, the antiviral activity of a compound can be
35 measured using the following general protocols.

5

Respiratory syncytial virus (RSV) antiviral activity and cytotoxicity assays

Anti-RSV activity

Antiviral activity against RSV was determined using an in vitro cytoprotection assay in
10 Hep2 cells. In this assay, compounds inhibiting the virus replication exhibit cytoprotective effect
against the virus-induced cell killing were quantified using a cell viability reagent. The method
used was similar to methods previously described in published literature (Chapman et al.,
Antimicrob Agents Chemother. **2007**, 57(9):3346-53.)

Hep2 cells were obtained from ATCC (Manassas, VI) and maintained in MEM media
15 supplemented with 10% fetal bovine serum and penicillin/streptomycin. Cells were passaged
twice a week and kept at subconfluent stage. Commercial stock of RSV strain A2 (Advanced
Biotechnologies, Columbia, MD) was titered before compound testing to determine the
appropriate dilution of the virus stock that generated desirable cytopathic effect in Hep2 cells.

For antiviral tests, Hep2 cells were seeded into 96-well plates 24 hours before the assay
20 at a density of 3,000 cells/well. On a separate 96well plate, compounds to be tested were
serially diluted in cell culture media. Eight concentrations in 3-fold serial dilution increments
were prepared for each tested compound and 100 uL/well of each dilution was transferred in
duplicate onto plates with seeded Hep2 cells. Subsequently, appropriate dilution of virus stock
previously determined by titration was prepared in cell culture media and 100 uL/well was
25 added to test plates containing cells and serially diluted compounds. Each plate included three
wells of infected untreated cells and three wells of uninfected cells that served as 0% and 100%
virus inhibition control, respectively. Following the infection with RSV, testing plates were
incubated for 4 days in a tissue culture incubator. After the incubation, RSV-induced cytopathic
effect was determined using a Cell TiterGlo reagent (Promega, Madison, WI) followed by a
30 luminescence read-out. The percentage inhibition was calculated for each tested concentration
relative to the 0% and 100% inhibition controls and the EC50 value for each compound was
determined by non-linear regression as a concentration inhibiting the RSV-induced cytopathic
effect by 50%. Ribavirin (purchased from Sigma, St. Louis, MO) was used as a positive control
for antiviral activity.

5 Compounds were also tested for antiviral activity against RSV in Hep2 cells using a 384 well format. Compounds were diluted in DMSO using a 10-step serial dilution in 3-fold increments via automation in 4 adjacent replicates each. Eight compounds were tested per dilution plate. 0.4uL of diluted compounds were then stamped via Biomek into 384-well plates (Nunc 142761 or 164730 w/lid 264616) containing 20uL of media (Mediatech Inc. MEM
10 supplemented with Glutamine, 10% FBS and Pen/Strep). DMSO and a suitable positive control compound, such as 80 μM GS-329467 or 10 μM 427346 was used for the 100% and 0% cell killing controls, respectively.

Hep2 cells (1.0×10^5 cells/ml) were prepared as above in batch to at least 40 mis excess of the number of sample plates (8 mis cell mix per plate) and infected with vendor supplied
15 (ABI) RSV strain A2 to arrive at an MOI of 1:1000 (virusxell #) or 1:3000 (vol virus: cell vol). Immediately after addition of virus, the RSV infected Hep2 cell suspension was added to each stamped 384-well plate at 20 μl per well using a uFlow dispenser, giving a final volume of 40 μL/well, each with 2000 infected cells. The plates were then incubated for 5 days at 37°C and 5% CO₂. Following incubation, the plates were equilibrated to room temperature in a biosafety
20 cabinet hood for 1.5 hrs and 40μL of Cell-Titer Glo viability reagent (Promega) was added to each well via uFlow. Following a 10-20 minute incubation, the plates were read using an EnVision or Victor Luminescence plate reader (Perkin-Elmer). The data was then uploaded and analyzed on the Bioinformatics portal under the RSV Cell Infectivity and 8-plate EC50-Hep2-384 or 8-plate EC50-Hep2-Envision protocols.

25 Multiple point data generated in the assay was analysed using Pipeline Pilot (Accelrys, Inc., Version 7.0) to generate a dose response curve based on least squares fit to a 4-parameter curve. The generated formula for the curve was then used to calculate the % inhibition at a given concentration. The % inhibition reported in the table was then adjusted based on the normalization of the bottom and top of the curve % inhibition values to 0% and 100%
30 respectively.

Representative activities for the compounds of the invention against RSV-induced cytopathic effects are shown in the Table below.

Compound formula	Percent inhibition at 0.5 μ M
1	100
2	100
3	100
4	100
5	100
6	100
7	100
8	90
9	99
10	95
11	92
12	89
13	82
14	81
15	82
16	71
17	70
18	70
19	70
20	63
21	58
22	55
23	48
24	30
25	100
26	84
27	100
28	100
29	71
30	100
31	100
32	79
33	99
34	13
35	100
36	98
37	100
38	100
39	100
40	95
41	4
42	9

Compound formula	Percent inhibition at 0.5 μ M
43	88
44	100
45	45
46	100
47	90
48	100
49	99
50	15
51	100
52	71
53	25
54	100
55	5
56	84
57	13
58	61
59	52
60	100
61	100
62	11
63	100
64	100
65	97
66	95
67	21
68	37
69	100
70	56
71	98
72	100
73	99
74	100
75	100
76	99
77	100
78	100
79	100
80	14
81	96
82	100
83	99
84	100

Compound formula	Percent inhibition at 0.5 μ M
85	100
86	80
87	100
88	100
89	100
90	100
91	100
92	100
93	44
94	88
95	21
96	65
97	100
98	97
99	100
100	100
101	16
102	16
103	100
104	56
105	31
106	100
107	100
108	99
109	100
110	36
111	n.d.
112	100
113	100
114	100
115	98
116	100
117	46
118	99
119	n.d.
120	100
121	92
122	100
123	98
124	100
125	n.d.
126	82

Compound formula	Percent inhibition at 0.5 μ M
127	n.d.
128	n.d.
129	87
130	100
131	96
132	97
133	56
134	35
135	100
136	83
137	n.d.
138	65
139	100
140	100
141	n.d.
142	100
143	100
144	80
145	60
146	100
147	n.d.
148	77
149	100
150	99
151	100
152	100
153	100
154	90
155	100
156	100
157	98
158	n.d.
159	91
160	71
161	61
162	100
163	100
164	n.d.
165	92
166	100
167	100
168	44

Compound formula	Percent inhibition at 0.5 μ M
169	100
170	100
171	100
172	100
173	98
174	58
175	91
176	97
177	99
178	100
179	93
180	94
181	83
182	93
183	100
184	100
185	100
186	100
187	100
188	100
189	89
190	100
191	n.d.
192	100
193	n.d.
194	n.d.
195	n.d.
196	96
197	100
198	100
199	100
200	100
201	100
202	100
203	100
204	100
205	100
206	100
207	100
208	100
209	100

(n.d. not determined)

5 Cytotoxicity

Cytotoxicity of tested compounds was determined in uninfected Hep2 cells in parallel with the antiviral activity using the cell viability reagent in a similar fashion as described before for other cell types (Cihlar et al., *Antimicrob Agents Chemother.* **2008**,52(2):655-65.). The same protocol as for the determination of antiviral activity was used for the measurement of compound cytotoxicity except that the cells were not infected with RSV. Instead, fresh cell culture media (100 uL/well) without the virus was added to tested plates with cells and prediluted compounds. Cells were then incubated for 4 days followed by a cell viability test using CellTiter Glo reagent and a luminescence read-out. Untreated cell and cells treated with 50 ug/mL puromycin (Sigma, St. Louis, MO) were used as 100% and 0% cell viability control, respectively. The percent of cell viability was calculated for each tested compound concentration relative to the 0% and 100% controls and the CC50 value was determined by non-linear regression as a compound concentration reducing the cell viability by 50%.

To test for compound cytotoxicity in Hep2 cells using a 384 well format, compounds were diluted in DMSO using a 10-step serial dilution in 3-fold increments via automation in 4 adjacent replicates each. Eight compounds were tested per dilution plate. 0.4uL of diluted compounds were then stamped via Biomek into 384-well plates (Nunc 142761 or 164730 w/lid 264616) containing 20µL of media (Mediatech Inc. MEM supplemented with Glutamine, 10% FBS and Pen/Strep). 50 µg/mL puromycin and DMSO were used for the 100% and 0% cytotoxicity controls, respectively.

Hep2 cells (1.0×10^5 cells/ml) were added to each stamped plate at 20 ul per well to give a total of 2000 cells/well and a final volume of 40 µL/well. Usually, the cells were batch prediluted to 1.0×10^5 cells/mL in excess of the number of sample plates and added at 20 ul per well into each assay plate using a uFlow dispenser. The plates were then incubated for 4 days at 37°C and 5% CO₂. Following incubation, the plates were equilibrated to room temperature in a biosafety cabinet hood for 1.5 hrs and 40µL of Cell-Titer Glo viability reagent (Promega) was added to each well via uFlow. Following a 10-20 minute incubation, the plates were read using an EnVision or Victor Luminescence plate reader (Perkin-Elmer). The data was then uploaded and analyzed on the Bioinformatics portal (Pipeline Pilot) under the Cytotoxicity assay using the 8-plate CC50-Hep2 or 8-plate CC50-Hep2 Envision protocols. Compounds tested for Anti-RSV activity were also tested in this cytotoxicity assay.

5


All publications, patents, and patent documents cited herein above are incorporated by reference herein, as though individually incorporated by reference.

The invention has been described with reference to various specific and preferred embodiments and techniques. However, one skilled in the art will understand that many variations and modifications may be made while remaining within the spirit and scope of the invention.

10

5 What is claimed is

1. A compound of formula I:



10 or a salt or ester, thereof;

wherein:

A is $-(C(R^4))_n-$ wherein any one $C(R^4)_2$ of said $-(C(R^4))_n-$ may be optionally replaced with $-O-$, $-S-$, $-S(O)_p-$, NH or NR^a ;

n is 3, 4, 5 or 6;

15 each p is 1 or 2;

Ar is a C_2-C_{20} heterocyclyl group or a C_6-C_{20} aryl group, wherein the C_2-C_{20} heterocyclyl group or the C_6-C_{20} aryl group is optionally substituted with 1, 2, 3, 4 or 5 R^6 ;

20 each R^3 , R^4 or R^6 is independently H, oxo, OR^{11} , $NR^{11}R^{12}$, $NR^{11}C(O)R^{11}$, $NR^{11}C(O)OR^{11}$, $NR^{11}C(O)NR^{12}$, N_3 , CN, NC-2, SR^{11} , $S(O)_pR^a$, $NR^{11}S(O)_pR^a$, $-C(=O)R^{11}$, $-C(=O)OR^{11}$, $-C(=O)NR^{11}R^{12}$, $-C(=O)SR^{11}$, $-S(O)_p(OR^{11})$, $-SO_2NR^{11}R^{12}$, $-NR^{11}S(O)_p(OR^{11})$, $-NR^{11}SO_pNR^{11}R^{12}$, $NR^{11}C(=NR^{11})NR^{11}R^{12}$, halogen, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, aryl(C_1-C_8)alkyl, C_6-C_{20} aryl, C_2-C_{20} heterocyclyl, (C_3-C_7) cycloalkyl or (C_4-C_8) carbocyclylalkyl;

25 or two R^4 on adjacent carbon atoms, when taken together, may optionally form a double bond between the two carbons to which they are attached or may form a (C_3-C_7) cycloalkyl ring wherein one carbon atom of said (C_3-C_7) cycloalkyl ring may be optionally replaced by $-O-$, $-S-$, $-S(O)_p-$, $-NH-$ or $-NR^a-$;

or four R^4 on adjacent carbon atoms, when taken together, may optionally form an optionally substituted C_6 aryl ring;

5 or two R⁴ on the same carbon atom, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a-;


or two R⁶ on adjacent carbon atoms, when taken together, may optionally form a (C₃-C₇)cycloalkyl ring wherein one carbon atom of said (C₃-C₇)cycloalkyl ring may be optionally
10 replaced by -O-, -S-, -S(O)_p-, -NH- or -NR^a-;

each R^a is independently (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclalkyl wherein any (C₁-C₈)haloalkyl, (C₂-C₈)alkenyl or (C₂-C₈)alkynyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀heterocyclyl, and wherein any aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclalkyl of R^a is optionally substituted with one or more OH, NH₂, CO₂H, C₂-C₂₀heterocyclyl or (C₁-C₈)alkyl;

each R¹¹ or R¹² is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclalkyl, -C(=O)R^a, -S(O)_pR^a or aryl(C₁-C₈)alkyl; or R¹¹ and R¹² taken together with a nitrogen to which they are both attached form a 3 to 7 membered heterocyclic ring wherein any one carbon atom of said heterocyclic ring can optionally be replaced with -O-, -S-, -S(O)_p-, -NH-, -NR^a- or -C(O)-; and

wherein each (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, aryl(C₁-C₈)alkyl, C₆-C₂₀aryl, C₂-C₂₀heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclalkyl of each R⁶, R¹¹ or R¹² is, independently, optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(O)_pR^a, OR^a, (d-C₈)alkyl, (C₁-C₈)haloalkyl, -C(O)R^a, -C(O)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(O)_pR^a, NR^aS(O)_pR^a, NHC(O)R^a, NR^aC(O)R^a, NHC(O)OR^a, NR^aC(O)OR^a, NR^aC(O)NHR^a, NR^aC(O)N(R^a)₂, NR^aC(O)NH₂, NHC(O)NHR^a, NHC(O)N(R^a)₂, NHC(O)NH₂, =NH, =NOH, =NOR^a, NR^aS(O)_pNHR^a, NR^aS(O)_pN(R^a)₂, NR^aS(O)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(O)(OH)₂ or R^a;

provided the compound is not:



5

2. The compound of claim 1 wherein each R^3 is H.

3. The compound of claim 1 or claim 2 wherein n is 3.


10

4. The compound of any one of claims 1-3 wherein each R^4 is H.

5. The compound of any one of claims 1-4 wherein A is $-(CH_2)_3-$.

15

6. The compound of claim 1 wherein the compound of formula I is a compound of formula Ia:



or a salt or ester, thereof.




20

7. The compound of any one of claims 1-6 wherein Ar is phenyl, pyridyl, 1,2,3,4-tetrahydronaphthyl, indazolyl, 1,6-naphthyridyl, 2,3,-dihydroindanyl, quinolyl, indolyl, 4H-benzo[d][1,3]dioxanyl, pyrazolo[1,5-a]pyridinyl, imidazo[1,2-a]pyridinyl, 1, 2,3,4-tetrahydroquinolyl, benzo[d][1,3]dioxolyl, quinoxalyl, isoquinolyl, naphthyl, thiophenyl, pyrazolyl, 4,5,6,7-tetrahydrobenzothiophenyl or pyrazolo[3,4,b]pyridinyl, wherein any phenyl,



25



pyridyl, 1,2,3,4-tetrahydronaphthyl, indazolyl, 1,6-naphthyridyl, 2,3,-dihydroindanyl, quinolyl, indolyl, 4H-benzo[d][1,3]dioxanyl, pyrazolo[1,5-a]pyridinyl, imidazo[1,2-a]pyridinyl, 1, 2,3,4-tetrahydroquinolyl, benzo[d][1,3]dioxolyl, quinoxalyl, isoquinolyl, naphthyl, thiophenyl,

- 5 pyrazolyl, 4,5,6,7-tetrahydrobenzothiophenyl or pyrazolo[3,4,b]pyridinyl of A is optionally substituted with 1 to 5 R⁶.
8. The compound of any one of claims 1-7 wherein each R⁶ is independently OR¹¹, CN, S(0)_pR^a, halogen, (C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl, (C₄-C₈)carbocyclylalkyl, NR¹¹C(0)R¹¹ or NR¹¹S(0)_pR^a, wherein any (C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R⁶ is optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(0)_pR^a, OR^a, (C₁-C₈)alkyl, (d-CiOhaloalkyl, -C(0)R^a, -C(0)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(0)_pR^a, NR^aS(0)_pR^a, NHC(0)R^a, NR^aC(0)R^a, NHC(0)OR^a, NR^aC(0)OR^a, NR^aC(0)NHR^a, NR^aC(0)N(R^a)₂, NR^aC(0)NH₂, NHC(0)NHR^a, NHC(0)N(R^a)₂, NHC(0)NH₂, =NH, =NOH, =NOR^a, NR^aS(0)_pNHR^a, NR^aS(0)_pN(R^a)₂, NR^aS(0)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(0)(OH)₂ or R^a.
- 10
- 15
- 20 9. The compound of any one of claims 1-7 wherein each R⁶ is independently OR¹¹, CN, S(0)_pR^a, halogen, (C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl, wherein any (C₁-C₈)alkyl, C₆-C₂₀ aryl, C₂-C₂₀ heterocyclyl, (C₃-C₇)cycloalkyl or (C₄-C₈)carbocyclylalkyl of R⁶ is optionally substituted with one or more oxo, halogen, hydroxy, NH₂, CN, N₃, N(R^a)₂, NHR^a, SH, SR^a, S(0)_pR^a, OR^a, (C₁-C₈)alkyl, (C₁-C₈)haloalkyl, -C(0)R^a, -C(0)H, -C(=O)OR^a, -C(=O)OH, -C(=O)N(R^a)₂, -C(=O)NHR^a, -C(=O)NH₂, NHS(0)_pR^a, NR^aS(0)_pR^a, NHC(0)R^a, NR^aC(0)R^a, NHC(0)OR^a, NR^aC(0)OR^a, NR^aC(0)NHR^a, NR^aC(0)N(R^a)₂, NR^aC(0)NH₂, NHC(0)NHR^a, NHC(0)N(R^a)₂, NHC(0)NH₂, =NH, =NOH, =NOR^a, NR^aS(0)_pNHR^a, NR^aS(0)_pN(R^a)₂, NR^aS(0)_pNH₂, NHS(O)_pNHR^a, NHS(O)_pN(R^a)₂, NHS(O)_pNH₂, -OC(=O)R^a, -OP(0)(OH)₂ or R^a.
- 25
- 30 10. The compound of claim 1 selected from:







5







5







5











5






5














and









and salts and esters, thereof.

11. The compound of claim 1 selected from:

10






5




and salts and esters, thereof.

12. A compound of formula:





10

wherein W is:







5









or a salt, or stereoisomer thereof.



10

13. The compound of claim 12 selected from:






5






5






and



5


and salts thereof.


14. A compound of formula:






10

wherein Y is:








5




10



5






5


or a salt, or stereoisomer thereof.



15. The compound of claim 14 selected from:

10






5







5








5







10














5













5






5






5



and



and salts thereof.

- 10 16. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 1-15 or a pharmaceutically acceptable salt or ester thereof and a pharmaceutically acceptable carrier.

5 17. The pharmaceutical composition of claim 16 further comprising a therapeutic agent
selected from ribavirin, palivizumab, motavizumab, RSV-IGIV, MEDI-557, A-60444, MDT-
637, BMS-433771, ALN-RSVO and ALX-0171 and mixtures thereof.

10 18. A method of treating a *Pneumovirinae* virus infection in a mammal in need thereof
comprising administering to the mammal a therapeutically effective amount of a compound of
any one of claims 1-15, or a pharmaceutically acceptable salt or ester thereof.

19. The method of claim 18 wherein the *Pneumovirinae* virus infection is caused by a
respiratory syncytial virus.

15 20. The method of claim 18 or claim 19 further comprising administering a therapeutically
effective amount of a therapeutic agent or composition thereof selected from the group
consisting of ribavirin, palivizumab, motavizumab, RSV-IGIV, MEDI-557, A-60444, MDT-
637, BMS-433771, ALN-RSVO and ALX-0171 and mixtures thereof.

20 21. A compound of any one of claims 1-15, or a pharmaceutically acceptable salt or ester
thereof, for use in the therapeutic or prophylactic treatment of a *Pneumovirinae* virus infection
or a respiratory syncytial virus infection.

25 22. The compound of claim 21 wherein the use in the therapeutic or prophylactic treatment
of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection further comprises
administering a therapeutically effective amount of a therapeutic agent or composition thereof
selected from the group consisting of ribavirin, palivizumab, motavizumab, RSV-IGIV, MEDI-
557, A-60444, MDT-63, BMS-433771, ALN-RSVO and ALX-0171 and mixtures thereof.

30 23. A compound of any one of claims 1-15, or a pharmaceutically acceptable salt or ester
thereof, for use in medical therapy.

- 5 24. The use of a compound of any one of claims 1-15 or a pharmaceutically acceptable salt or ester thereof, for the manufacture of a medicament for the treatment of a *Pneumovirinae* virus infection or a respiratory syncytial virus infection in a mammal.

INTERNATIONAL SEARCH REPORT

International application No PCT/US2012/071065

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D487/04 A61K31/519 A61P31/12
 ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
EPO-Internal , CHEM ABS Data, WPI Data, BEI LSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	wo 03/095455 A2 (SMITHKLINE BEECHAM CORP [US] ; GUDMUNDSSON KRISTJAN S [US] ; JOHNS BRIAN) 20 November 2003 (2003-11-20) See formul a (1) i n c l a i m 1 -----	1-24
A	wo 00/42043 AI (UNIV MICHIGAN [US] ; TOWNSEND LEROY B [US] ; DRACH JOHN C [US]) 20 July 2000 (2000-07-20) c l a i m 1 -----	1-24
X,P	wo 2011/163518 AI (GI LEAD SCI ENCES INC [US] ; BABA OGLU KERIM [US] ; B00JAMRA CONSTANTINE G) 29 December 2011 (2011-12-29) the whol e document -----	1-24

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
---	---

Date of the actual completion of the international search 4 March 2013	Date of mailing of the international search report 11/03/2013
--	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Menchaca, Roberto
--	--

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2012/071065
--

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 03095455	A2	20-11-2003	AT 365740 T 15-07-2007
		AU 2003228770 A1	11-11-2003
		DE 60314623 T2	28-02-2008
		EP 1504004 A2	09-02-2005
		ES 2287476 T3	16-12-2007
		JP 2005529919 A	06-10-2005
		US 2005203106 A1	15-09-2005
		WO 03095455 A2	20-11-2003

WO 0042043	A1	20-07-2000	AU 2962800 A 01-08-2000
		US 6342501 B1	29-01-2002
		WO 0042043 A1	20-07-2000

WO 2011163518	A1	29-12-2011	AR 082275 A1 28-11-2012
		AU 2011270798 A1	24-01-2013
		CA 2800834 A1	29-12-2011
		TW 201213332 A	01-04-2012
		US 2012003215 A1	05-01-2012
		UY 33467 A	31-01-2012
		WO 2011163518 A1	29-12-2011
