

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
19 June 2003 (19.06.2003)


PCT

(10) International Publication Number
WO 03/049703 A2

- (51) International Patent Classification⁷: **A61K**
- (21) International Application Number: PCT/US02/39646
- (22) International Filing Date:
11 December 2002 (11.12.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/340,775 12 December 2001 (12.12.2001) US
60/424,840 8 November 2002 (08.11.2002) US
10/314,512 9 December 2002 (09.12.2002) US
- (71) Applicants: **VIRGINIA TECH INTELLECTUAL PROPERTIES, INC.** [US/US]; 1872 Pratt Dr., Blacksburg, VA 24060 (US). **IOWA STATE UNIVERSITY RESEARCH FOUNDATION, INC.** [US/US]; 310 Lab of Mechanics, Ames, IA 50011-2131 (US).
- (72) Inventors: **MENG, Xiang-Jin**; 401 Craig Drive, Blacksburg, VA 24060 (US). **FENAUX, Martijn**; 305 N. Knollwood Drive, Apt. 5204, Blacksburg, VA 24060 (US). **HALBUR, Patrick, G.**; 3211 Kingman Road, Ames, IA 50010 (US).
- (74) Agent: **ROSENBLUM, Anne, M.**; 163 Delaware Avenue, Suite 212, Delmar, NY 12054 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

[Continued on next page]

(54) Title: CHIMERIC INFECTIOUS DNA CLONES, CHIMERIC PORCINE CIRCOVIRUSES AND USES THEREOF



(57) Abstract: The present invention relates to infectious DNA clones, infectious chimeric DNA clones of porcine circovirus (PCV), vaccines and means of protecting pigs against viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2). The new chimeric infectious DNA clone and its derived, a virulent chimeric virus are constructed from the nonpathogenic PCV1 in which the immunogenic ORF gene of the pathogenic PCV2 replaces a gene of the nonpathogenic PCV1, preferably in the same position. The chimeric virus advantageously retains the nonpathogenic phenotype of PCV1 but elicits specific immune responses against the pathogenic PCV2. The invention further embraces the immunogenic polypeptide expression products.



European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

5 CHIMERIC INFECTIOUS DNA CLONES, CHIMERIC
PORCINE CIRCOVIRUSES AND USES THEREOF

10 CROSS-REFERENCE TO RELATED U.S. APPLICATIONS

This nonprovisional application claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 60/424,840, filed on November 8, 2002, which claims the benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application No. 60/340,775, filed on December 12, 2001. The two prior applications are incorporated herein by reference in their
15 entirety.

STATEMENT REGARDING FEDERALLY SPONSORED
RESEARCH OR DEVELOPMENT

Not Applicable

20 REFERENCE TO A "Sequence Listing"

The material on a single compact disc containing a Sequence Listing file provided in this application is incorporated by reference. The date of creation is _____, 2003 and the size is approximately ____.

25 BACKGROUND OF THE INVENTION

Field of the Invention

The present invention concerns infectious porcine circovirus type-1 (PCV1) and type-2 (PCV2) DNA clones, chimeric PCV1-2 infectious DNA clones and live chimeric viruses
30 derived from the chimeric DNA clones, useful as vaccines.

Description of the Related Art

All patents and publications cited in this specification are hereby incorporated by reference in their entirety.

Porcine circovirus (PCV) was originally isolated as a cell culture contaminant of a porcine kidney cell line PK-15 (I. Tischer *et al.*, "A very small porcine virus with circular single-stranded DNA," *Nature* **295**:64-66 (1982); I. Tischer *et al.*, "Characterization of papovavirus and picornavirus-like particles in permanent pig kidney cell lines," *Zentralbl. Bakteriol. Hyg. Otg. A.* **226**(2):153-167 (1974)). PCV is a small icosahedral non-enveloped virus that contains a single stranded circular DNA genome of about 1.76 kb. PCV is classified in the family of *Circoviridae*, which consists of three other animal circoviruses (chicken anemia virus (CAV), psittacine beak and feather disease virus (PBFDV) and the recently discovered columbid circovirus (CoCV) from pigeons) and three plant circoviruses (banana bunchy top virus, coconut foliar decay virus and subterranean clover stunt virus) (M. R. Bassami *et al.*, "Psittacine beak and feather disease virus nucleotide sequence analysis and its relationship to porcine circovirus, plant circoviruses, and chicken anemia virus," *Virology* **249**:453-459 (1998); J. Mankertz *et al.*, "Transcription analysis of porcine circovirus (PCV)," *Virus Genes* **16**:267-276 (1998); A. Mankertz *et al.*, "Cloning and sequencing of columbid circovirus (CoCV), a new circovirus from pigeons," *Arch. Virol.* **145**:2469-2479 (2000); B. M. Meehan *et al.*, "Sequence of porcine circovirus DNA: affinities with plant circoviruses," *J. Gen. Virol.* **78**:221-227 (1997); B. M. Meehan *et al.*, "Characterization of novel circovirus DNAs associated with wasting syndromes in pigs," *J. Gen. Virol.* **79**:2171-2179 (1998); D. Todd *et al.*, "Comparison of three animal viruses with circular single-stranded DNA genomes," *Arch. Virol.* **117**:129-135 (1991)). Members of the three previously recognized animal circoviruses (PCV, CAV, and PBFDV) do not share nucleotide sequence homology or antigenic determinants with each other (M. R. Bassami *et al.*, 1998, *supra*; D. Todd *et al.*, 1991, *supra*). The genome of the newly identified CoCV shared about 40% nucleotide sequence identity with that of PCV (A. Mankertz *et al.*, "Cloning and sequencing of columbid circovirus (CoCV), a new circovirus from pigeons," *Arch. Virol.* **145**:2469-2479 (2000)). Recently, a novel human circovirus with a circular genome, designated as transfusion transmitted virus or TT virus (TTV), was identified from individuals associated with post-transfusion hepatitis (H. Miyata *et al.*, "Identification of a novel GC-rich 113-

nucleotide region to complete the circular, single-stranded DNA genome of TT virus, the first human circovirus," J. Virol. **73**:3582-3586 (1999); T. Nishizawa *et al.*, "A novel DNA virus (TTV) associated with elevated transaminase levels in posttransfusion hepatitis of unknown etiology," Biochem. Biophys. Res. Commun. **241**:92-97 (1997)). Additionally, a
5 human TTV-like mini virus (TLMV) was identified from normal blood donors (P. Biagini *et al.*, "Genetic analysis of full-length genomes and subgenomic sequences of TT virus-like mini virus human isolates," J. Gen. Virol. **82**: 379-383 (2001); K. Takahashi *et al.*, "Identification of a new human DNA virus (TTV-like mini virus, TLMV) intermediately related to TT virus and chicken anemia virus," Arch. Virol. **145**:979-93 (2000)) and a third
10 novel human circovirus, known as SEN virus (SENV), was also discovered from humans with post-transfusion hepatitis (T. Umemura *et al.*, "SEN virus infection and its relationship to transfusion-associated hepatitis," Hepathology **33**:1303-1311 (2001)). The genomic organization of both human TTV and TLMV is similar to that of the CAV (P. Biagini *et al.*, 2001, *supra*; H. Miyata *et al.*, 1999, *supra*; K. Takahashi *et al.*, 2000, *supra*). Although
15 antibodies to PCV were found in various animal species including humans, mice, cattle and pigs (G. M. Allan *et al.*, "Production, preliminary characterization and applications of monoclonal antibodies to porcine circovirus," Vet. Immunol. Immunopathol. **43**:357-371 (1994); G. C. Dulac and A. Afshar, "Porcine circovirus antigens in PK-15 cell line (ATCC CCL-33) and evidence of antibodies to circovirus in Canadian pigs," Can. J. Vet. Res.
20 **53**:431-433 (1989); S. Edwards and J. J. Sands, "Evidence of circovirus infection in British pigs," Vet. Rec. **134**:680-1 (1994); J. C. Harding and E.G. Clark, "Recognizing and diagnosing postweaning multisystemic wasting syndrome (PMWS)," Swine Health and Production **5**:201-203 (1997); R. K. Hines and P. D. Lukert, "Porcine circovirus: a serological survey of swine in the United States," Swine Health and Production **3**:71-73
25 (1995); G. P. Nayar *et al.*, "Evidence for circovirus in cattle with respiratory disease and from aborted bovine fetuses," Can. Vet. J. **40**:277-278 (1999); I. Tischer *et al.*, "Distribution of antibodies to porcine circovirus in swine populations of different breeding farms," Arch. Virol. **140**:737-743 (1995); I. Tischer *et al.*, "Presence of antibodies reacting with porcine circovirus in sera of humans, mice, and cattle," Arch. Virol. **140**:1427-1439 (1995)), little is
30 known regarding the pathogenesis of PCV in these animal species. Experimental infection of pigs with the PK-15 cells-derived PCV did not produce clinical disease and thus, this virus is

not considered to be pathogenic to pigs (G. M. Allan *et al.*, "Pathogenesis of porcine circovirus; experimental infections of colostrum deprived piglets and examination of pig foetal material," Vet. Microbiol. **44**:49-64 (1995); I. Tischer *et al.*, "Studies on epidemiology and pathogenicity of porcine circovirus," Arch. Virol. **91**:271-276 (1986)). The
5 nonpathogenic PCV derived from the contaminated PK-15 cell line was designated as porcine circovirus type 1 or PCV1.

Postweaning multisystemic wasting syndrome (PMWS), first described in 1991 (J. C. Harding and E.G. Clark, 1997, *supra*), is a complex disease of weaning piglets that is becoming increasingly more widespread. With the threat of a potential serious economic
10 impact upon the swine industry, it has become urgent to develop a vaccine against PCV2, the primary causative agent of PMWS. PMWS mainly affects pigs between 5-18 weeks of age. Clinical PMWS signs include progressive weight loss, dyspnea, tachypnea, anemia, diarrhea, and jaundice. Mortality rate may vary from 1% to 2%, and up to 40% in some complicated cases in the U.K. (M. Muirhead, "Sources of information on PMWS/PDNS," Vet. Rec.
15 **150**:456 (2002)). Microscopic lesions characteristic of PMWS include granulomatous interstitial pneumonia, lymphadenopathy, hepatitis, and nephritis (G. M. Allan and J. A. Ellis, "Porcine circoviruses: a review," J. Vet. Diagn. Invest. **12**:3-14 (2000); J. C. Harding and E.G. Clark, 1997, *supra*). PMWS has now been recognized in pigs in Canada, the United States (G. M. Allan *et al.*, "Novel porcine circoviruses from pigs with wasting disease
20 syndromes," Vet. Rec. **142**:467-468 (1998); G. M. Allan *et al.*, "Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe," J. Vet. Diagn. Invest. **10**:3-10 (1998); G. M. Allan and J. A. Ellis, 2000, *supra*; J. Ellis *et al.*, "Isolation of circovirus from lesions of pigs with postweaning multisystemic wasting syndrome," Can. Vet. J. **39**:44-51 (1998); A. L. Hamel *et al.*, "Nucleotide sequence of
25 porcine circovirus associated with postweaning multisystemic wasting syndrome in pigs," J. Virol. **72**:5262-5267 (1998); M. Kiupel *et al.*, "Circovirus-like viral associated disease in weaned pigs in Indiana," Vet. Pathol. **35**:303-307 (1998); R. Larochelle *et al.*, "Identification and incidence of porcine circovirus in routine field cases in Quebec as determined by PCR," Vet. Rec. **145**:140-142 (1999); B. M. Meehan *et al.*, 1998, *supra*; I. Morozov *et al.*,
30 "Detection of a novel strain of porcine circovirus in pigs with postweaning multisystemic wasting syndrome," J. Clin. Microbiol. **36**:2535-2541 (1998)), most European countries (G.

M. Allan *et al.*, "Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe," J. Vet. Diagn. Invest. 10:3-10 (1998); G. M. Allan and J. A. Ellis, 2000, *supra*; S. Edwards and J. J. Sands, 1994, *supra*; S. Kennedy *et al.*, "Porcine circovirus infection in Northern Ireland," Vet. Rec. 142:495-496 (1998); A. Mankertz *et al.*,
5 "Characterization of PCV-2 isolates from Spain, Germany and France," Virus Res. 66:65-77 (2000); C. Rosell *et al.*, "Identification of porcine circovirus in tissues of pigs with porcine dermatitis and nephropathy syndrome. Vet. Rec. 146:40-43 (2000); P. Spillane *et al.*, "Porcine circovirus infection in the Republic of Ireland," Vet. Rec. 143:511-512 (1998); G. J. Wellenberg *et al.*, "Isolation and characterization of porcine circovirus type 2 from pigs
10 showing signs of post-weaning multisystemic wasting syndrome in the Netherlands," Vet. Quart. 22:167-72 (2000)) and some countries in Asia (C. Choi *et al.*, "Porcine postweaning multisystemic wasting syndrome in Korean pig: detection of porcine circovirus 2 infection by immunohistochemistry and polymerase chain reaction," J. Vet. Diagn. Invest. 12:151-153 (2000); A. Onuki *et al.*, "Detection of porcine circovirus from lesions of a pig with wasting
15 disease in Japan," J. Vet. Med. Sci. 61:1119-1123 (1999)). PMWS potentially has a serious economic impact on the swine industry worldwide.

The primary causative agent of PMWS is a pathogenic strain of PCV designated as porcine circovirus type 2 or PCV2 (G. M. Allan *et al.*, "Novel porcine circoviruses from pigs with wasting disease syndromes," Vet. Rec. 142:467-468 (1998); G. M. Allan *et al.*,
20 "Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe," J. Vet. Diagn. Invest. 10:3-10 (1998); G. M. Allan *et al.*, "Isolation and characterisation of circoviruses from pigs with wasting syndromes in Spain, Denmark and Northern Ireland," Vet. Microbiol. 66:115-23 (1999); G. M. Allan and J. A. Ellis, 2000, *supra*; J. Ellis *et al.*, 1998, *supra*; A. L. Hamel *et al.*, 1998, *supra*; B. M. Meehan *et al.*, 1998, *supra*; I. Morozov *et al.*, 1998, *supra*). The complete genomic sequence of the PMWS-associated PCV2 has been determined (M. Fenaux *et al.*, "Genetic characterization of type 2
25 porcine circovirus (PCV-2) from pigs with postweaning multisystemic wasting syndrome in different geographic regions of North America and development of a differential PCR-restriction fragment length polymorphism assay to detect and differentiate between infections
30 with PCV-1 and PCV-2," J. Clin. Microbiol. 38:2494-503 (2000); A. L. Hamel *et al.*, 1998,

supra; J. Mankertz *et al.*, 1998, *supra*; B. M. Meehan *et al.*, 1997, *supra*; B. M. Meehan *et al.*, 1998, *supra*; I. Morozov *et al.*, 1998, *supra*).

PCV1 is ubiquitous in pigs but is not pathogenic to pigs. In contrast, the genetically related PCV2 is pathogenic and causes PMWS in pigs. Sequence analyses reveals that the
5 PMWS-associated PCV2 shares only about 75% nucleotide sequence identity with the nonpathogenic PCV1. The ORF2 gene of both the nonpathogenic PCV1 and the pathogenic PCV2 encodes for the major immunogenic viral capsid protein (P. Nawagitgul *et al.*, "Modified indirect porcine circovirus (PCV) type 2-based and recombinant capsid protein (ORF2)-based ELISA for the detection of antibodies to PCV," Immunol. Clin. Diagn. Lab
10 Immunol. 1:33-40 (2002); P. Nawagitgul *et al.*, "Open reading frame 2 of porcine circovirus type 2 encodes a major capsid protein," J. Gen. Virol. 81:2281-2287 (2000)).

Initial attempts to reproduce clinical PMWS in conventional pigs by PCV2 inoculation were unsuccessful (M. Balasch *et al.*, "Experimental inoculation of conventional pigs with tissue homogenates from pigs with post-weaning multisystemic wasting
15 syndrome," J. Comp. Pathol. 121:139-148 (1999); M. Fenaux *et al.*, "Cloned Genomic DNA of Type 2 Porcine Circovirus (PCV-2) Is Infectious When Injected Directly into the Liver and Lymph Nodes of SPF Pigs: Characterization of Clinical Disease, Virus Distribution, and Pathologic Lesions," J. Virol. 76:541-551 (2002)). Experimental reproduction of clinical PMWS in gnotobiotic pigs and conventional pigs with tissue homogenates from pigs with
20 naturally occurring PMWS and with cell culture propagated PCV2 produced mixed results. Clinical PMWS was reproduced in gnotobiotic (SPF) pigs and colostrum-deprived and caesarian-derived pigs co-infected with PCV2 and porcine parvovirus (PPV) (G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," J. Comp. Pathol. 121:1-11 (1999); S. Krakowka
25 *et al.*, "Viral wasting syndrome of swine: experimental reproduction of postweaning multisystemic wasting syndrome in gnotobiotic swine by coinfection with porcine circovirus 2 and porcine parvovirus," Vet. Pathol. 37:254-263 (2000)), and in PCV2-inoculated gnotobiotic pigs when their immune system was activated by keyhole hemocyanin in incomplete Freund's adjuvant (S. Krakowka *et al.*, "Activation of the immune system is the
30 pivotal event in the production of wasting disease in pigs infected with porcine circovirus-2 (PCV-2)," Vet. Pathol. 38:31-42 (2001)).

Clinical PMWS was also reproduced in cesarean derived/colostrum deprived pigs (CD/CD) inoculated with PCV2 alone (P. A. Harms *et al.*, "Experimental reproduction of severe disease in CD/CD pigs concurrently infected with type 2 porcine circovirus and porcine reproductive and respiratory syndrome virus," *Vet. Pathol.* **38**:528-539 (2001)) and
5 in conventional pigs co-infected with PCV2 and either porcine parvovirus (PPV) or porcine reproductive and respiratory syndrome virus (PRRSV) (A. Rivora *et al.*, "Experimental inoculation of conventional pigs with porcine reproductive and respiratory syndrome virus and porcine circovirus 2," *J. Virol.* **76**: 3232-3239 (2002)). In cases of the PRRSV/PCV2 co-infection, the PMWS characteristic pathological signs such as lymphoid depletion,
10 granulomatous inflammation and necrotizing hepatitis are induced by PCV2 and not by PRRSV (P. A. Harms *et al.*, 2001, *supra*). However, clinical PMWS was not reproduced in gnotobiotic pigs infected with PCV2 alone (G. M. Allan *et al.*, "Experimental infection of colostrums deprived piglets with porcine circovirus 2 (PCV2) and porcine reproductive and respiratory syndrome virus (PRRSV) potentiates PCV2 replication," *Arch. Virol.* **145**:2421-
15 2429 (2000); G. M. Allan *et al.*, "A sequential study of experimental infection of pigs with porcine circovirus and porcine parvovirus: immunostaining of cryostat sections and virus isolation, *J. Vet. Med.* **47**:81-94 (2000); G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," *J. Comp. Pathol.* **121**:1-11 (1999); M. Balasch *et al.*, 1999, *supra*; J. Ellis *et al.*,
20 "Reproduction of lesions of postweaning multisystemic wasting syndrome in gnotobiotic piglets," *J. Vet. Diagn. Invest.* **11**:3-14 (1999); S. Kennedy *et al.*, "Reproduction of lesions of postweaning multisystemic wasting syndrome by infection of conventional pigs with porcine circovirus type 2 alone or in combination with porcine parvovirus" *J. Comp. Pathol.* **122**:9-24 (2000); S. Krakowka *et al.*, 2001, *supra*; S. Krakowka *et al.*, 2000, *supra*; R. M. Pogranichnyy *et al.*, "Characterization of immune response of young pigs to porcine circovirus type 2 infection," *Viral. Immunol.* **13**:143-153 (2000)). The virus inocula used in these studies were either homogenates of tissues from pigs with naturally occurring PMWS, or virus propagated in PK-15 cell cultures (G. M. Allan *et al.*, "Experimental infection of colostrums deprived piglets with porcine circovirus 2 (PCV2) and porcine reproductive and
30 respiratory syndrome virus (PRRSV) potentiates PCV2 replication," *Arch. Virol.* **145**:2421-2429 (2000); G. M. Allan *et al.*, "A sequential study of experimental infection of pigs with

porcine circovirus and porcine parvovirus: immunostaining of cryostat sections and virus isolation, J. Vet. Med. **47**:81-94 (2000); G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," J. Comp. Pathol. **121**:1-11 (1999); M. Balasch *et al.*, 1999, *supra*; J. Ellis *et al.*, 5 1999, *supra*; S. Kennedy *et al.*, 2000, *supra*; S. Krakowka *et al.*, 2001, *supra*; S. Krakowka *et al.*, 2000, *supra*; R. M. Pogranichnyy *et al.*, 2000, *supra*). Since tissue homogenates may contain other common swine agents such as PPV and porcine reproductive and respiratory syndrome virus (PRRSV) (G. M. Allan *et al.*, "Experimental infection of colostrums deprived piglets with porcine circovirus 2 (PCV2) and porcine reproductive and respiratory syndrome virus (PRRSV) potentiates PCV2 replication," Arch. Virol. **145**:2421-2429 (2000); 10 G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," J. Comp. Pathol. **121**:1-11 (1999); G. M. Allan and J. A. Ellis, 2000, *supra*; J. A. Ellis *et al.*, "Coinfection by porcine circoviruses and porcine parvovirus in pigs with naturally acquired postweaning multisystemic wasting syndrome," J. Vet. Diagn. Invest. **12**:21-27 (2000); C. Rosell *et al.*, 2000, *supra*), and since 15 the ATCC PK-15 cell line used for PCV2 propagation was persistently infected with PCV1 (G. C. Dulac and A. Afshar, 1989, *supra*), the clinical disease and pathological lesions reproduced in those studies may not be solely attributable to PCV2 infection (G. M. Allan *et al.*, "Experimental infection of colostrums deprived piglets with porcine circovirus 2 (PCV2) and porcine reproductive and respiratory syndrome virus (PRRSV) potentiates PCV2 replication," Arch. Virol. **145**:2421-2429 (2000); G. M. Allan *et al.*, "A sequential study of experimental infection of pigs with porcine circovirus and porcine parvovirus: immunostaining of cryostat sections and virus isolation, J. Vet. Med. **47**:81-94 (2000); G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," J. Comp. Pathol. **121**:1-11 (1999); G. M. Allan and J. A. Ellis, 2000, *supra*; J. A. Ellis *et al.*, 2000, *supra*). 20 25

Clinical PMWS has also been reproduced in PCV2-inoculated CDCD pigs when vaccinated with *Mycoplasma hyopneumoniae* (G. M. Allan *et al.*, "Immunostimulation, PCV-2 and PMWS," Vet. Rec. **147**:171-172 (2000)). Two recent field studies by G. M. Allan *et al.*, "Neonatal vaccination for *Mycoplasma hyopneumoniae* and postweaning multisystemic wasting syndrome: a field trial," Pig J. **48**:34-41 (2001), and S. C. Kyriakis *et al.*, "The 30

effects of immuno-modulation on the clinical and pathological expression of postweaning multisystemic wasting syndrome," J. Comp. Pathol. 126:38-46 (2002), tested the effect of immuno-modulation by *Mycoplasma hyopneumoniae* vaccine on the development of PMWS in endemic herds, and showed a significant decrease in PMWS cases in unvaccinated groups compared to the vaccinated animals. However, another recent study using conventional SPF piglets under controlled laboratory conditions could not reproduce such an effect, suggesting that vaccinations with *M. hyopneumoniae* may potentially influence the development of clinical PMWS but it is clearly a secondary role to a PCV2 infection. Based on these and other studies, PCV2 is nevertheless considered to be the primary but not the exclusive causative agent of PMWS.

The lack of an infectious virus stock of a biologically pure form of PCV2 has impeded the understanding of PCV2 pathogenesis and the etiological role of PCV2 in PMWS. Vaccinations against PPV and possibly PRRSV have not consistently been shown to prevent the onset of PMWS in PCV2 infected pigs. Consequently, finding a safe yet potent vaccine that specifically targets PMWS has been difficult. There is a definite art-recognized need in the veterinary field to produce an efficacious, safe vaccine against PCV2 infections and PMWS.

U.S. Patent No. 6,287,856 (Poet *et al.*) and WO 99/45956 concern nucleic acids from psittacine beak and feather disease virus (BFDV), a circovirus that infects avian species, and from porcine circovirus (PCV). The patent proposes vaccine compositions comprising naked DNA or mRNA and discloses a nucleic acid vector for the transient expression of PCV in a eukaryotic cell comprising a *cis*-acting transcription or translation regulatory sequence derived from the human cytomegalovirus immediate or early gene enhancer or promoter functionally linked to a nucleic acid of the sequence. However, since the PCV DNA is derived solely from the PK-15 cell line, it is likely to comprise the nonpathogenic PCV1 discovered nearly 30 years ago by I. Tischer *et al.*, 1974, *supra*, and, therefore, it is not likely to be effective in eliciting an immune reaction to PCV2 or infections caused by PCV2. Subunit vaccines of recombinant proteins made from vectors comprising open reading frames are also suggested in the patent but the open reading frames from PCV are not well characterized or distinguished from each other. Since the source of the PCV DNA is PK-15

cells, the proteins made from those vectors comprising the open reading frames of PCV1 would not possess reliable immunogenic properties, if any, against PCV2.

U.S. Patent No. 6,217,883 (Allan *et al.*) and French Patent No. 2,781,159B relate to the isolation of five PCV strains from pulmonary or ganglionic samples taken from pigs infected with PMWS in Canada, California and France (Brittany), and their use in combination with at least one porcine parvovirus antigen in immunogenic compositions. Proteins encoded by PCV2 open reading frames (ORF) consisting of ORF1 to ORF13 are broadly described in the patent but there is no exemplification of any specific protein exhibiting immunogenic properties. The patent further discloses vectors consisting of DNA plasmids, linear DNA molecules and recombinant viruses that contain and express *in vivo* a nucleic acid molecule encoding the PCV antigen. Several other references, for example, U.S. Patent No. 6,391,314 B1; U.S. Patent No. 6,368,601 B1; French Patent No. 2,769,321; French Patent No. 2,769,322; WO 01/96377 A2; WO 00/01409; WO 99/18214; WO 00/77216 A2; WO 01/16330 A2; WO 99/29871; *etc.*, also describe the administration of PCV1 or PCV2 polypeptides or the nucleic acids encoding the polypeptides of various strains.

However, the nonpathogenic PCV1 will not be useful against PCV2 infections and the pathogenic PCV2 strains described in the art, even if attenuated, are likely to be of limited value due to the usual tendency of a live virus to revert to its virulent state. Therefore, there is still a long-standing need in the art for a live, infectious, nonpathogenic antigen for the inoculation of pigs against serious infection or PMWS caused by PCV2 that is efficacious and remains safe in veterinary vaccines. These goals are met by the construction of the new live chimeric porcine circovirus described herein, which is based upon the genomic backbone of the nonpathogenic PCV1 isolated by I. Tischer *et al.* almost 30 years ago. The novel chimeric porcine circovirus of the present invention is able to satisfy that long-standing need by uniquely and advantageously retaining the nonpathogenic phenotype of PCV1 but eliciting specific immune response against pathogenic PCV2.

BRIEF SUMMARY OF THE INVENTION

The present invention concerns infectious chimeric DNA clones of porcine circovirus (PCV) and live chimeric viruses derived from the DNA clones that are useful as vaccines.

The new live chimeric, genetically avirulent viruses are made from the nonpathogenic PCV1 genomic structure in which an immunogenic gene of a pathogenic PCV2 strain replaces a gene of the PCV1, typically in the same corresponding position. The invention encompasses the biologically functional plasmids, viral vectors and the like that contain the new recombinant nucleic acid molecules described herein, suitable host cells transfected by the vectors comprising the DNA and the immunogenic polypeptide expression products. Also included within the scope of the present invention is a novel method of protecting pigs against viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2 comprising administering to a pig in need of such protection an immunologically effective amount of a vaccine comprising, for example, the cloned chimeric DNA in a plasmid, a chimeric virus derived from the chimeric DNA clone, the polypeptide products expressed from the DNA described herein, *etc.* The invention further provides new infectious PCV2 molecular DNA and reciprocal chimeric DNA clones of PCV that find use as experimental models in obtaining or characterizing the novel avirulent viral vaccines.

BRIEF DESCRIPTION OF THE DRAWINGS

The background of the invention and its departure from the art will be further described hereinbelow with reference to the accompanying drawings, wherein:

Figure 1 represents the construction of an infectious PCV2 molecular DNA clone. The relative positions of the primer pair used to amplify the complete PCV2 genome are indicated by the arrows (reverse primer PCVSAC2, forward primer PCVSAC2). The PCV2 genomic DNA amplified by PCR is digested with *SacII* restriction enzyme, and purified. The *purified* and *SacII*-digested genomic DNA is ligated to form concatemers. Ligated concatemers are separated by gel electrophoresis, the tandem genome dimer of PCV2 is purified and cloned into pSK vector that is pre-digested with *SacII* enzyme to produce a molecular PCV2 DNA clone.

Figures 2A and 2B illustrate that the cloned PCV2 plasmid DNA is infectious when transfected *in vitro* in PK-15 cells. Figure 2A shows the detection of PCV2 antigen by immunofluorescence assay (IFA) in PK-15 cells transfected with the cloned PCV2 plasmid DNA. Intense immunolabeling of PCV2 antigen is visualized in the nucleus, and to a lesser degree, cytoplasm of the transfected cells. Figure 2B shows mock-transfected PK-15 cells.

Figure 3A shows the lungs from a pig inoculated by intralymphoid route with PCV2 DNA at 21 DPI. The lungs are rubbery, failed to collapse, and are mottled tan-red. Tracheobronchial lymph nodes are markedly enlarged and tan (arrows). Figure 3B represents a microscopic section of a normal lung from a control pig (25X). Figure 3C represents a microscopic section of the lung from the pig in Figure 3A. Note the peribronchiolar lymphohistiocytic inflammation and mild necrotizing bronchiolitis (25X). Figure 3D illustrates the immunohistochemical staining of the lung in Figure 3A. Note the PCV2 antigen in macrophages (arrows) and fibroblast-like cells (arrow heads) around airways (64X).

Figure 4A shows a normal lymph node from a control pig. Note the well-defined lymphoid follicles (arrows) (25X). Figure 4B represents a microscopic section of the tracheobronchial lymph node from the pig in Figure 3A inoculated 21 days previously by intralymphoid route with cloned PCV2 genomic DNA. Lymphoid follicles are poorly defined, there is mild-to-moderate lymphoid depletion, and mild multifocal granulomatous inflammation (25X). Figure 4C represents a microscopic section of the lymph node in Figure 4B in a larger magnification focusing on one follicle. Note the poorly defined follicle with macrophages and giant cells (arrow) replacing follicular lymphocytes (64X). Figure 4D illustrates the immunohistochemical detection of PCV2 antigen in the same lymph node as Figure 4B in macrophages (arrows) and giant cells (small arrowheads), and dendritic-like cells (large arrowheads) in the follicles (64X).

Figure 5 illustrates the construction of a chimeric PCV1-2 (PCV1/PCV2) DNA clone with the nonpathogenic PCV1 genome carrying the immunogenic ORF2 capsid gene of the pathogenic PCV2. The dimmerized DNA clone is used for *in vitro* transfection of PK-15 cells to produce live chimeric virus expressing ORF2 protein of PCV2, and *in vivo* animal experiments to confirm activity.

Figure 6 represents the construction and organization of the infectious PCV1, PCV2, chimeric PCV1-2 and reciprocal chimeric PCV2-1 molecular DNA clones. The PCV2 DNA clone is constructed by ligating two full-length linear PCV2 genomes in tandem into the Bluescript SK vector (pSK) by the general methods described previously (M. Fenaux *et al.*, 2002, *supra*). PCV1 DNA clone is constructed by ligating two full-length linear PCV1 genomes in tandem into pSK vector. Chimeric PCV1-2 DNA clone is constructed by

replacing the ORF2 capsid gene of PCV1 with that of the PCV2 in the nonpathogenic PCV1 genomic backbone in pSK vector. Reciprocal chimeric PCV2-1 DNA clone is constructed by replacing the ORF2 capsid gene of the pathogenic PCV2 with that of the nonpathogenic PCV1 in the pathogenic PCV2 genomic backbone in pSK vector. Both chimeric clones are dimmers in pSK vector. The arrows represent the relative locations of the PCR primers for the detection of PCV1, PCV2, PCV1-2 and PCV2-1 viremia in inoculated animals.

Figures 7A-7J demonstrate that the PCV1, PCV2, chimeric PCV1-2 and reciprocal chimeric PCV2-1 DNA clones are infectious and express respective viral antigens when transfected *in vitro* in PK-15 cells. The left panel (7A, 7C, 7E, 7G and 7I) is stained with monoclonal antibody against the PCV1 ORF2. The right panel (7B, 7D, 7F, 7H and 7J) is stained with antibody against PCV2. Panels 7A and 7B are mock transfected PK-15 cells. Panels 7C and 7D are PK-15 cells transfected with the PCV1 DNA clone. Panels 7E and 7F are PK-15 cells transfected with the PCV2 DNA clone. Panels 7G and 7H are PK-15 cells transfected with the chimeric PCV1-2 DNA clone. Panels 7I and 7J are PK-15 cells transfected with the reciprocal chimeric PCV2-1 DNA clone.

Figure 8 represents the full-length DNA sequence of the cloned PCV2 molecular DNA (which corresponds to SEQ ID NO:1).

Figure 9 represents the full-length DNA sequence of the cloned chimeric PCV1-2 DNA (which corresponds to SEQ ID NO:2).

Figure 10 represents the DNA sequence of the immunogenic ORF2 capsid gene of the cloned chimeric PCV1-2 DNA (which corresponds to SEQ ID NO:3).

Figure 11 represents the putative amino acid translation of the immunogenic ORF2 capsid gene of the chimeric PCV1-2 DNA (which corresponds to SEQ ID NO:4).

DETAILED DESCRIPTION OF THE INVENTION

In accordance with the present invention, there are provided infectious molecular and chimeric nucleic acid molecules of porcine circovirus (PCV), live chimeric viruses produced from the chimeric nucleic acid molecule and veterinary vaccines to protect pigs from viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2. The invention further provides immunogenic polypeptide expression products that may be used as vaccines.

The new avirulent, infectious chimeric DNA molecule of PCV (PCV1-2) comprises a nucleic acid molecule encoding an infectious, nonpathogenic PCV1 that contains an immunogenic open reading frame (ORF) gene of a pathogenic PCV2 in place of an ORF gene in the PCV1 genome. The infectious chimeric PCV1-2 DNA clone preferably contains the immunogenic capsid gene (ORF2) of the PCV2 DNA cloned in the genomic backbone of the infectious, nonpathogenic PCV1 DNA clone. Generally, the capsid gene of the PCV2 DNA replaces the ORF2 gene of the PCV1 DNA in the nonpathogenic PCV1 genomic structure but it is contemplated that a variety of positional permutations may be constructed through genetic engineering to obtain other avirulent or attenuated chimeric DNA clones. The reciprocal chimeric infectious PCV2-1 DNA clone between PCV1 and PCV2 is disclosed as a control to analyze the chimeric PCV1-2 clone of the invention and is constructed by replacing the capsid gene of PCV2 with that of PCV1 in the backbone of the pathogenic PCV2 infectious DNA clone. In addition to being an experimental model, the reciprocal chimeric PCV2-1 DNA clone may find use in making specially tailored vaccines.

The cloned genomic DNA of PCV2 described herein is shown to be *in vitro* and *in vivo* infectious when transfected into PK-15 cells and given to pigs. The infectious PCV2 DNA clone produces pathological lesions characteristic of PMWS in pigs allowing for an improved characterization of clinical disease and understanding of virus distribution in the tissue cells. This new, readily reproducible pathogenic agent lends itself to the development of a suitable vaccination program to prevent PMWS in pigs.

The novel chimeric PCV1-2 DNA clone is also infectious by both *in vitro* transfection of PK-15 cells and *in vivo* administration to pigs. In transfected PK-15 cells, the chimeric PCV1-2 DNA clone expresses the PCV2 capsid antigen (the immunogenic capsid protein of PCV2) whereas the reciprocal chimeric PCV2-1 DNA clone expresses the PCV1 capsid antigen, which is demonstrated by immunofluorescence assay (IFA) using antibodies specific to PCV1 or PCV2 capsid antigen. Seroconversion to PCV2-specific antibody is detected in pigs inoculated with the infectious PCV2 clone as well as the chimeric PCV1-2 clone. Detecting the seroconversion to PCV2-specific antibody establishes that the chimeric PCV1-2 DNA clone induces the PCV2-specific antibody in infected pigs and, consequently, acts to protect inoculated pigs from infection with PCV2.

The below examples describe the evaluation of the immunogenicity and pathogenicity of the chimeric DNA clones in inoculated pigs in more detail. Basically, seroconversions to antibodies against PCV2 ORF2 antigen are detected in pigs inoculated with the PCV2 DNA clone (Group 3) and the chimeric PCV1-2 DNA clone (Group 4). All of the pigs inoculated with the PCV1 clone and the reciprocal chimeric PCV2-1 DNA clone (Groups 2 and 5, respectively) seroconvert to the PCV1 antibody. The viruses recovered from selected pigs in each group are partially sequenced and confirmed to be the authentic respective infectious DNA clones used in the inoculation. Gross and microscopic lesions in various tissues of animals inoculated with the PCV2 DNA clone are significantly more severe than those found in pigs inoculated with PCV1, chimeric PCV1-2 and reciprocal chimeric PCV2-1 DNA clones.

Surprisingly and advantageously, the chimeric PCV1-2 infectious DNA clone having the immunogenic capsid gene (ORF2) of the pathogenic PCV2 cloned into the nonpathogenic PCV1 genomic backbone induces a specific antibody response to the pathogenic PCV2 capsid antigen while it uniquely retains the nonpathogenic nature of PCV1 in pigs. Animals inoculated with the chimeric PCV1-2 infectious DNA clone develop a mild infection resembling that of PCV1 inoculated animals while seroconverting to the antibody against the ORF2 capsid protein of the pathogenic PCV2. The average length of viremia observed in PCV1 and chimeric PCV1-2 inoculated animals is shorter, 0.625 weeks and 1 week respectively, than that in pathogenic PCV2 inoculated animals which is about 2.12 weeks. The lack of detectable chimeric PCV1-2 viremia in some inoculated animals does not affect seroconversion to antibody against PCV2 ORF2 capsid protein in the PCV1-2 inoculated pigs (Group 4). The results indicate that, even though the chimeric PCV1-2 viremia is short or undetectable in some inoculated animals, the chimeric PCV1-2 virus is able to induce antibody response against PCV2 ORF2 capsid protein. The special ability of the chimeric PCV1-2 infectious DNA clone to induce the immune response specific to the pathogenic PCV2 immunogenic ORF2 capsid protein yet remain nonpathogenic to pigs makes the chimeric PCV1-2 clone particularly useful as a genetically engineered live-attenuated vaccine and other types of vaccines.

The novel, purified and isolated nucleic acid molecules of this invention comprise the full-length DNA sequence of the cloned chimeric PCV1-2 DNA set forth in SEQ ID NO:2,

shown in Figure 9 and deposited in the American Type Culture Collection under Patent Deposit Designation PTA-3912; its complementary strand (*i.e.*, reverse and opposite base pairs) or the nucleotide sequences having at least 95% homology to the chimeric nucleotide sequence (*i.e.*, a significant active portion of the whole gene). Conventional methods that are well known in the art can be used to make the complementary strands or the nucleotide sequences possessing high homology, for instance, by the art-recognized standard or high stringency hybridization techniques. The purified and isolated nucleic acid molecule comprising the DNA sequence of the immunogenic capsid gene of the cloned chimeric PCV1-2 DNA is also set forth in SEQ ID NO:3 and Figure 10.

Suitable cells containing the chimeric nucleic acid molecule uniquely produce live, infectious chimeric porcine circoviruses. The live, infectious chimeric virus is derived from the chimeric DNA clone by transfecting PK-15 cells via *in vitro* and *in vivo* transfections as illustrated herein. A preferred example of the cloned chimeric PCV1-2 DNA is the nucleotide sequence set forth in SEQ ID NO:2 and Figure 9. The invention further envisions that the chimeric virus is derived from the complementary strand or the nucleotide sequences having a high homology, at least 95% homology, to the chimeric nucleotide sequence.

Also included within the scope of the present invention are biologically functional plasmids, viral vectors and the like that contain the new recombinant nucleic acid molecules described herein, suitable host cells transfected by the vectors comprising the chimeric and molecular DNA clones and the immunogenic polypeptide expression products. A particularly preferred immunogenic protein has the amino acid sequence set forth in SEQ ID NO:4 and Figure 11. The biologically active variants thereof are further encompassed by the invention. One of ordinary skill in the art would know how to modify, substitute, delete, *etc.*, amino acid(s) from the polypeptide sequence and produce biologically active variants that retain the same, or substantially the same, activity as the parent sequence without undue effort.

To produce the immunogenic polypeptide products of this invention, the process may include the following steps: growing, under suitable nutrient conditions, prokaryotic or eucaryotic host cells transfected with the new recombinant nucleic acid molecules described herein in a manner allowing expression of said polypeptide products, and isolating the desired polypeptide products of the expression of said nucleic acid molecules by standard methods

known in the art. It is contemplated that the immunogenic proteins may be prepared by other techniques such as, for example, biochemical synthesis and the like.

Vaccines of the chimeric viral and molecular DNA clones, and methods of using them, are also included within the scope of the present invention. Inoculated pigs are protected from serious viral infection and PMWS caused by PCV2. The novel method protects pigs in need of protection against viral infection or PMWS by administering to the pig an immunologically effective amount of a vaccine according to the invention, such as, for example, a vaccine comprising an immunogenic amount of the chimeric PCV1-2 DNA, the cloned chimeric virus, a plasmid or viral vector containing the chimeric DNA of PCV1-2, the polypeptide expression products, the recombinant PCV2 DNA, *etc.* Other antigens such as PRRSV, PPV, other infectious swine agents and immune stimulants may be given concurrently to the pig to provide a broad spectrum of protection against viral infections.

The vaccines comprise, for example, the infectious chimeric PCV1-2 DNA, the cloned PCV chimeric DNA genome in suitable plasmids or vectors such as, for example, the pSK vector, an avirulent, live chimeric virus, an inactivated chimeric virus, *etc.* in combination with a nontoxic, physiologically acceptable carrier and, optionally, one or more adjuvants. The vaccine may also comprise the infectious PCV2 molecular DNA clone described herein. The infectious chimeric PCV1-2 DNA, the plasmid DNA containing the infectious chimeric viral genome and the live chimeric virus are preferred with the live chimeric virus being most preferred. The avirulent, live viral vaccine of the present invention provides an advantage over traditional viral vaccines that use either attenuated, live viruses which run the risk of reverting back to the virulent state or killed cell culture propagated whole virus which may not induce sufficient antibody immune response for protection against the viral disease.

The adjuvant, which may be administered in conjunction with the vaccine of the present invention, is a substance that increases the immunological response of the pig to the vaccine. The adjuvant may be administered at the same time and at the same site as the vaccine, or at a different time, for example, as a booster. Adjuvants also may advantageously be administered to the pig in a manner or at a site different from the manner or site in which the vaccine is administered. Suitable adjuvants include, but are not limited to, aluminum hydroxide (alum), immunostimulating complexes (ISCOMS), non-ionic block polymers or

copolymers, cytokines (like IL-1, IL-2, IL-7, IFN- α , IFN- β , IFN- γ , *etc.*), saponins, monophosphoryl lipid A (MLA), muramyl dipeptides (MDP) and the like. Other suitable adjuvants include, for example, aluminum potassium sulfate, heat-labile or heat-stable enterotoxin isolated from *Escherichia coli*, cholera toxin or the B subunit thereof, diphtheria toxin, tetanus toxin, pertussis toxin, Freund's incomplete or complete adjuvant, *etc.* Toxin-based adjuvants, such as diphtheria toxin, tetanus toxin and pertussis toxin may be inactivated prior to use, for example, by treatment with formaldehyde.

The vaccines may further contain additional antigens to promote the immunological activity of the infectious chimeric PCV DNA clones such as, for example, porcine reproductive and respiratory syndrome virus (PRRSV), porcine parvovirus (PPV), other infectious swine agents and immune stimulants.

The new vaccines of this invention are not restricted to any particular type or method of preparation. The cloned viral vaccines include, but are not limited to, infectious DNA vaccines (*i.e.*, using plasmids, vectors or other conventional carriers to directly inject DNA into pigs), live vaccines, modified live vaccines, inactivated vaccines, subunit vaccines, attenuated vaccines, genetically engineered vaccines, *etc.* These vaccines are prepared by standard methods known in the art.

The live viral vaccine is generally the most desirable vaccine in that all possible immune responses are activated in the recipient of the vaccine, including systemic, local, humoral and cell-mediated immune responses. A killed vaccine, on the other hand, can only induce humoral immune response. Albeit the most desirable, however, live viral vaccines have several disadvantages, such as the potential risk of contamination with live adventitious viral agents or the risk that the virus may revert to virulence in the field. Remarkably, the unique PCV1-2 chimeric DNA of the present invention overcomes those disadvantages. Using only the immunogenic genes of the pathogenic PCV2, the chimeric DNA constructs a live, replicating chimeric virus that is nonpathogenic yet elicits the complete, beneficial immune responses of live viral vaccines against the pathogenic PCV2 virus. The live virus vaccine based on the chimeric virus will have little chance, if any, for reversion to a pathogenic phenotype. Thus, the new chimeric virus based on the structure of the nonpathogenic PCV1 has a huge advantage over any recombinant PCV2 DNA virus, any

live, attenuated PCV2 vaccine or any other type of vaccine predicated solely on PCV2 for immunity against the PCV2 infections.

Although the live viral vaccine is most preferred, other types of vaccines may be used to inoculate pigs with the new chimeric virus and other antigens described herein. To prepare
5 inactivated virus vaccines, for instance, the virus propagation from the infectious DNA clone is done by methods known in the art or described herein. Serial virus inactivation is then optimized by protocols generally known to those of ordinary skill in the art.

Inactivated virus vaccines may be prepared by treating the chimeric virus derived from the cloned PCV DNA with inactivating agents such as formalin or hydrophobic
10 solvents, acids, *etc.*, by irradiation with ultraviolet light or X-rays, by heating, *etc.* Inactivation is conducted in a manner understood in the art. For example, in chemical inactivation, a suitable virus sample or serum sample containing the virus is treated for a sufficient length of time with a sufficient amount or concentration of inactivating agent at a sufficiently high (or low, depending on the inactivating agent) temperature or pH to
15 inactivate the virus. Inactivation by heating is conducted at a temperature and for a length of time sufficient to inactivate the virus. Inactivation by irradiation is conducted using a wavelength of light or other energy source for a length of time sufficient to inactivate the virus. The virus is considered inactivated if it is unable to infect a cell susceptible to infection.

20 The preparation of subunit vaccines typically differs from the preparation of a modified live vaccine or an inactivated vaccine. Prior to preparation of a subunit vaccine, the protective or antigenic components of the vaccine must be identified. Such protective or antigenic components include certain amino acid segments or fragments of the viral capsid proteins which raise a particularly strong protective or immunological response in pigs;
25 single or multiple viral capsid proteins themselves, oligomers thereof, and higher-order associations of the viral capsid proteins which form virus substructures or identifiable parts or units of such substructures; oligoglycosides, glycolipids or glycoproteins present on or near the surface of the virus or in viral substructures such as the lipoproteins or lipid groups associated with the virus, *etc.* Preferably, a capsid protein, such as the protein encoded by the
30 ORF2 gene, is employed as the antigenic component of the subunit vaccine. Other proteins encoded by the infectious DNA clone may also be used. These immunogenic components are

readily identified by methods known in the art. Once identified, the protective or antigenic portions of the virus (*i.e.*, the "subunit") are subsequently purified and/or cloned by procedures known in the art. The subunit vaccine provides an advantage over other vaccines based on the live virus since the subunit, such as highly purified subunits of the virus, is less toxic than the whole virus.

If the subunit vaccine is produced through recombinant genetic techniques, expression of the cloned subunit such as the ORF2 (capsid) gene, for example, may be optimized by methods known to those in the art (see, for example, Maniatis *et al.*, "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory, Cold Spring Harbor, MA., 1989). If the subunit being employed represents an intact structural feature of the virus, such as an entire capsid protein, the procedure for its isolation from the virus must then be optimized. In either case, after optimization of the inactivation protocol, the subunit purification protocol may be optimized prior to manufacture.

To prepare attenuated vaccines from pathogenic clones, the tissue culture adapted, live, pathogenic PCV2 is first attenuated (rendered nonpathogenic or harmless) by methods known in the art, typically made by serial passage through cell cultures. Attenuation of pathogenic clones may also be made by gene deletions or viral-producing gene mutations. Then, the attenuated PCV2 viruses may be used to construct additional chimeric PCV1-2 viruses that retain the nonpathogenic phenotype of PCV1 but can vary in the strength of the immunogenicity traits selected from the PCV2 genome through recombinant technology.

The most preferred vaccine employs the live chimeric virus DNA clone, in particular, the clone containing the immunogenic genes of PCV2 cloned in the backbone of the nonpathogenic PCV1. Advantageously, the live chimeric virus, which is naturally avirulent when constructed through genetic engineering, does not require time-consuming attenuation procedures. The virus uniquely serves as a live but nonpathogenic replicating virus that produces immunogenic proteins against PCV2 during virus replication, which can then elicit a full range of immune responses against the pathogenic PCV2.

As a further benefit, the preferred live chimeric virus of the present invention provides a genetically stable vaccine that is easier to make, store and deliver than other types of attenuated vaccines. Avirulent or attenuated vaccines based upon chimeric viruses are generally considered as safe as, if not safer than, the traditionally modified live vaccines (J.

Arroyo *et al.*, "Molecular basis for attenuation of neurovirulence of a yellow fever Virus/Japanese encephalitis virus chimera vaccine (ChimeriVax-JE)," J. Virol. **75**(2):934-942 (2001); F. Guirakhoo *et al.*, "Recombinant chimeric yellow fever-dengue type 2 virus is immunogenic and protective in nonhuman primates," J. Virol. **74**(12):5477-5485 (2000); S. Tang *et al.*, "Toward a poliovirus-based simian immunodeficiency virus vaccine: correlation between genetic stability and immunogenicity," J. Virol. **71**(10):7841-7850 (1997)). For example, the ChimeriVax-JE vaccine against Japanese encephalitis virus (JEV), which is a genetically engineered derivative of the yellow fever virus vaccine YFV17D in which the genes encoding the structural proteins prM and E of YFV17D are replaced with the corresponding genes of the attenuated JEV SA14-14-2 strain, has been shown to be genetically stable after prolonged passages both *in vitro* and *in vivo* (J. Arroyo *et al.*, 2001, *supra*). Another chimeric virus vaccine ChimeriVax-D2 against Dengue virus type 2, which is an attenuated chimeric yellow fever (YF)-dengue type 2 (dengue-2) virus, has also been found to be genetically stable; its sequences were reported to be unchanged after 18 passages in Vero cells (F. Guirakhoo *et al.*, 2000, *supra*).

Another preferred vaccine of the present invention utilizes suitable plasmids for delivering the nonpathogenic chimeric DNA clone to pigs. In contrast to the traditional vaccine that uses live or killed cell culture propagated whole virus, this invention provides for the direct inoculation of pigs with the plasmid DNA containing the infectious chimeric viral genome.

Additional genetically engineered vaccines, which are desirable in the present invention, are produced by techniques known in the art. Such techniques involve, but are not limited to, further manipulation of recombinant DNA, modification of or substitutions to the amino acid sequences of the recombinant proteins and the like.

Genetically engineered vaccines based on recombinant DNA technology are made, for instance, by identifying alternative portions of the viral gene encoding proteins responsible for inducing a stronger immune or protective response in pigs (*e.g.*, proteins derived from ORF3, ORF4, *etc.*). Such identified genes or immuno-dominant fragments can be cloned into standard protein expression vectors, such as the baculovirus vector, and used to infect appropriate host cells (see, for example, O'Reilly *et al.*, "Baculovirus Expression Vectors: A Lab Manual," Freeman & Co., 1992). The host cells are cultured, thus expressing

the desired vaccine proteins, which can be purified to the desired extent and formulated into a suitable vaccine product.

If the clones retain any undesirable natural abilities of causing disease, it is also possible to pinpoint the nucleotide sequences in the viral genome responsible for the virulence, and genetically engineer the virus avirulent through, for example, site-directed mutagenesis. Site-directed mutagenesis is able to add, delete or change one or more nucleotides (see, for instance, Zoller *et al.*, DNA 3:479-488, 1984). An oligonucleotide is synthesized containing the desired mutation and annealed to a portion of single stranded viral DNA. The hybrid molecule, which results from that procedure, is employed to transform bacteria. Then double-stranded DNA, which is isolated containing the appropriate mutation, is used to produce full-length DNA by ligation to a restriction fragment of the latter that is subsequently transfected into a suitable cell culture. Ligation of the genome into the suitable vector for transfer may be accomplished through any standard technique known to those of ordinary skill in the art. Transfection of the vector into host cells for the production of viral progeny may be done using any of the conventional methods such as calcium-phosphate or DEAE-dextran mediated transfection, electroporation, protoplast fusion and other well-known techniques (*e.g.*, Sambrook *et al.*, "Molecular Cloning: A Laboratory Manual," Cold Spring Harbor Laboratory Press, 1989). The cloned virus then exhibits the desired mutation. Alternatively, two oligonucleotides can be synthesized which contain the appropriate mutation. These may be annealed to form double-stranded DNA that can be inserted in the viral DNA to produce full-length DNA.

Genetically engineered proteins, useful in vaccines, for instance, may be expressed in insect cells, yeast cells or mammalian cells. The genetically engineered proteins, which may be purified or isolated by conventional methods, can be directly inoculated into pigs to confer protection against viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2.

An insect cell line (like HI-FIVE) can be transformed with a transfer vector containing nucleic acid molecules obtained from the virus or copied from the viral genome which encodes one or more of the immuno-dominant proteins of the virus. The transfer vector includes, for example, linearized baculovirus DNA and a plasmid containing the

desired polynucleotides. The host cell line may be co-transfected with the linearized baculovirus DNA and a plasmid in order to make a recombinant baculovirus.

Alternatively, DNA from a pig suffering from PMWS, which encode one or more capsid proteins, the infectious PCV2 molecular DNA clone or the cloned PCV chimeric DNA genome can be inserted into live vectors, such as a poxvirus or an adenovirus and used
5 as a vaccine.

An immunologically effective amount of the vaccines of the present invention is administered to a pig in need of protection against viral infection or PMWS. The immunologically effective amount or the immunogenic amount that inoculates the pig can be
10 easily determined or readily titrated by routine testing. An effective amount is one in which a sufficient immunological response to the vaccine is attained to protect the pig exposed to the virus which causes PMWS. Preferably, the pig is protected to an extent in which one to all of the adverse physiological symptoms or effects of the viral disease are significantly reduced, ameliorated or totally prevented.

The vaccine can be administered in a single dose or in repeated doses. Dosages may
15 range, for example, from 1 to 1,000 micrograms of the plasmid DNA containing the infectious chimeric DNA genome (dependent upon the concentration of the immuno-active component of the vaccine), but should not contain an amount of virus-based antigen sufficient to result in an adverse reaction or physiological symptoms of viral infection.
20 Methods are known in the art for determining or titrating suitable dosages of active antigenic agent based on the weight of the pig, concentration of the antigen and other typical factors. Preferably, the infectious chimeric viral DNA clone is used as a vaccine, or a live infectious chimeric virus can be generated *in vitro* and then the live chimeric virus is used as a vaccine. In that case, 100 to 200 micrograms of cloned chimeric PCV DNA or about 10,000 50%
25 tissue culture infective dose (TCID₅₀) of live chimeric virus can be given to a pig.

Desirably, the vaccine is administered to a pig not yet exposed to the PCV virus. The vaccine containing the chimeric PCV1-2 infectious DNA clone or other antigenic forms thereof can conveniently be administered intranasally, transdermally (*i.e.*, applied on or at the skin surface for systemic absorption), parenterally, *etc.* The parenteral route of administration
30 includes, but is not limited to, intramuscular, intravenous, intraperitoneal, intradermal (*i.e.*, injected or otherwise placed under the skin) routes and the like. Since the intramuscular and

intradermal routes of inoculation have been successful in other studies using viral infectious DNA clones (E. E. Sparger *et al.*, "Infection of cats by injection with DNA of feline immunodeficiency virus molecular clone," *Virology* **238**:157-160 (1997); L. Willems *et al.*, "*In vivo* transfection of bovine leukemia provirus into sheep," *Virology* **189**:775-777 (1992)),
5 these routes are most preferred, in addition to the practical intranasal route of administration. Although less convenient, it is also contemplated that the vaccine is given to the pig through the intralymphoid route of inoculation. A unique, highly preferred method of administration involves directly injecting the plasmid DNA containing PCV1-2 chimera into the pig intramuscularly, intradermally, intralymphoidly, *etc.*

10 When administered as a liquid, the present vaccine may be prepared in the form of an aqueous solution, syrup, an elixir, a tincture and the like. Such formulations are known in the art and are typically prepared by dissolution of the antigen and other typical additives in the appropriate carrier or solvent systems. Suitable carriers or solvents include, but are not limited to, water, saline, ethanol, ethylene glycol, glycerol, *etc.* Typical additives are, for
15 example, certified dyes, flavors, sweeteners and antimicrobial preservatives such as thimerosal (sodium ethylmercurithiosalicylate). Such solutions may be stabilized, for example, by addition of partially hydrolyzed gelatin, sorbitol or cell culture medium, and may be buffered by conventional methods using reagents known in the art, such as sodium hydrogen phosphate, sodium dihydrogen phosphate, potassium hydrogen phosphate,
20 potassium dihydrogen phosphate, a mixture thereof, and the like.

Liquid formulations also may include suspensions and emulsions that contain suspending or emulsifying agents in combination with other standard co-formulants. These types of liquid formulations may be prepared by conventional methods. Suspensions, for example, may be prepared using a colloid mill. Emulsions, for example, may be prepared
25 using a homogenizer.

Parenteral formulations, designed for injection into body fluid systems, require proper isotonicity and pH buffering to the corresponding levels of porcine body fluids. Isotonicity can be appropriately adjusted with sodium chloride and other salts as needed. Suitable solvents, such as ethanol or propylene glycol, can be used to increase the solubility of the
30 ingredients in the formulation and the stability of the liquid preparation. Further additives that can be employed in the present vaccine include, but are not limited to, dextrose,

conventional antioxidants and conventional chelating agents such as ethylenediamine tetraacetic acid (EDTA). Parenteral dosage forms must also be sterilized prior to use.

Another embodiment of the present invention involves a new method of preparing an infectious, nonpathogenic chimeric nucleic acid molecule of PCV1-2, which comprises removing an open reading frame (ORF) gene of a nucleic acid molecule encoding an infectious nonpathogenic PCV1, replacing the same position with an immunogenic ORF gene of a nucleic acid molecule encoding an infectious pathogenic PCV2, and recovering the chimeric nucleic acid molecule. The nucleic acid molecule is typically DNA. A preferred method replaces the ORF2 gene of the nonpathogenic PCV1 DNA with the immunogenic ORF2 capsid gene of the infectious pathogenic molecular DNA of PCV2 described herein. It is contemplated that other ORF positions or immunogenic fragments thereof can be exchanged between the PCV1 and PCV2 DNA to construct the attenuated infectious chimeric DNA clones according to the methods described herein.

The recombinant nucleic acid molecule is then used to construct the live, infectious, replicating chimeric virus of the present invention that advantageously retains the nonpathogenic nature of PCV1 yet expresses the immunogenic ORF2 protein of the pathogenic PCV2 and elicits a complete immune response against the pathogenic PCV2. Desirably, the PCV1-2 DNA clone serves as a genetically engineered avirulent, live vaccine against PCV2 infection and PMWS in pigs.

An infectious DNA clone of PCV2 is constructed, as described herein, so that a biologically pure and homogeneous infectious virus stock can be generated for pathogenesis studies and the development of nonpathogenic, chimeric vaccines. The course of clinical disease, virus distribution and pathological lesions associated with PCV2 infection are more definitively characterized by using this molecular DNA clone and a biologically pure and homogeneous infectious PCV2 virus stock derived from the molecular DNA clone than have been observed in the past, which lends itself to the development of the desired vaccine products of the present invention.

The PCV2 molecular clone is generated by ligating two copies of the complete PCV2 genome in tandem into the pSK vector. In sharp contrast to the single copy genome disclosed in the art, the infectious DNA PCV2 clone made by the methods described herein contains two complete copies of the PCV2 genome ligated together in tandem repeat.

Ligating two copies of genome in tandem provides a similar circular genome that mimics the usual circular genome of PCV2. The advantage of having two copies of the genome in tandem in the infectious DNA PCV2 clone is to be able to maximize replication when the infectious DNA clone is transfected *in vitro* and *in vivo*. Thus, the clone of the invention
5 operates more efficiently and effectively than the prior single copy genome.

Infection of animals with the molecular viral clone is extremely useful to studying the genetic determinants of viral replication and virulence in the host. Type-2 porcine circovirus (PCV2) has been incriminated as the causative agent of postweaning multisystemic wasting syndrome (PMWS). PMWS is a complex disease syndrome in swine and multiple factors
10 may be involved in the clinical presentation of PMWS. However, the difficulty in producing a biologically pure form of PCV2 due to the presence of other common swine agents in the tissue homogenates of diseased pigs has impeded a definitive characterization of the clinical disease and pathological lesions solely attributable to PCV2 infection. This is the first time an infectious molecular DNA clone of PCV2 has been constructed and used to characterize
15 the disease and pathological lesions associated with PCV2 infection by direct *in vivo* transfection of pigs with the molecular clone.

The homogeneous PCV2 live virus stock derived from the molecular clone is shown to be infectious *in vitro* when transfected into PK-15 cells. The cloned PCV2 genomic DNA is also infectious when directly injected into the livers and superficial iliac lymph nodes of
20 specific-pathogen-free (SPF) pigs. Animals injected with the cloned PCV2 plasmid DNA develop an infection and disease resembling that induced by intranasal inoculation with a homogenous, infectious PCV2 live virus stock. Seroconversion to PCV2-specific antibody is detected in the majority of pigs from the inoculated groups at 35 days postinoculation (DPI).

The onset and duration of viremia in pigs inoculated with the chimeric PCV1-2 DNA
25 clone are similar to those of the pigs inoculated with the nonpathogenic PCV1 DNA clone, whereas viremia in pigs inoculated with the PCV2 clone appears earlier and lasted longer. Beginning at 14 DPI and lasting about 2-4 weeks, viremia is detected in the majority of the PCV2-inoculated animals. Similarly, the majority of inoculated pigs necropsied at 35 DPI seroconverted to PCV2-antibodies. PCV2 antigen is detected in various tissues and organs in
30 inoculated pigs. Gross lesions are limited to the lungs and lymph nodes, and are characterized by systematically enlarged tan colored lymph nodes, lungs that failed to

collapse and mild multifocal tan-colored foci of consolidation. Gross lesions affecting the lymph nodes in both the nonpathogenic PCV1 and the chimeric PCV1-2 inoculated pigs are mild and limited to only a few animals, whereas the pathogenic PCV2 inoculated pigs all have moderate-to-severe swelling and discoloration of lymphoid tissues (Table 9, below).

5 Statistical analysis reveals that the scores of the gross lesions in the lymph nodes of the chimeric PCV1-2 inoculated animals are similar to those in nonpathogenic PCV1 inoculated pigs. At 21 DPI, PCV2 inoculated pigs have gross lesions that are statistically more severe than those of the PCV1 and the chimeric PCV1-2 inoculated pigs. Histopathological lesions and PCV2-specific antigen are detected in numerous tissues and organs including brain, lung,
10 heart, kidney, tonsil, lymph nodes, spleen, ileum and liver of the inoculated (infected) pigs. The histopathological lesions in multiple tissues and organs similar to those of PMWS are reproduced with the PCV2 molecular DNA clone as well as with the infectious virus prepared *in vitro* from the molecular DNA clone. Microscopically, at both 21 and 49 DPIs, the chimeric PCV1-2 inoculated animals have statistically less microscopic lesions than the
15 PCV2 inoculated animals. The microscopic lesion scores in lymph nodes of the chimeric PCV1-2 inoculated pigs are similar to those of the nonpathogenic PCV1, the reciprocal chimeric PCV2-1 and the uninoculated control animals. Moderate to severe microscopic lesions are found in multiple tissues of pathogenic PCV2 inoculated animals including lung, liver, lymphoid, spleen, brain, heart, kidney and tonsil tissue. However, in chimeric PCV1-2
20 inoculated animals, mild to moderate microscopic lesions are limited only to liver, lymph nodes and kidney tissues (see Table 10, below).

There are no remarkable clinical signs of PMWS in the control or any of the inoculated pigs. Although the characteristic clinical symptoms of PMWS are not observed with the cloned PCV2 plasmid DNA (the infectious PCV2 DNA clone) or with a biologically
25 pure PCV2 infectious virus stock, PCV2 is clearly responsible for the PMWS-like histopathological lesions reproduced in the below illustrative examples. It is generally believed that PCV2 is the primary but not the sole pathogenic agent responsible for the onset of clinical PMWS.

This invention more definitively characterizes the clinical course and pathological
30 lesions exclusively attributable to PCV2 infection. The present data in the below illustrative examples indicate that the readily reproduced, cloned PCV2 genomic DNA is available to

replace infectious virus for the PCV2 pathogenesis and immunization studies. While PCV2 is shown as essential for development of PMWS, other factors or agents such as PRRSV, PPV, *etc.* may be required to induce the full spectrum of clinical signs and lesions associated with advanced cases of PMWS. However, with the knowledge that PCV2 is a key factor, the novel infectious, replicating viral clone of the present invention can be further modified or genetically engineered to achieve the desired optimal immunogenic effect through methods known to those of ordinary skill in immunology and molecular genetics.

The availability of the infectious DNA clone of PCV2 described herein makes it feasible to develop the genetically engineered attenuated vaccine for preventing PCV2 infection and PMWS in pigs. It is known that PCV2 replicates in the lymph nodes, lungs and liver during natural infection, and one of the major pathogenic effects is the impairment of the immune system by degradation of the lymphoid structures (S. Krakowka *et al.*, 2001, *supra*; G. M. Allan and J. A. Ellis, 2000, *supra*; S. Kennedy *et al.*, 2000, *supra*; G. J. Wellenberg *et al.*, 2000, *supra*; G. M. Allan *et al.*, "Experimental reproduction of severe wasting disease by co-infection of pigs with porcine circovirus and porcine parvovirus," J. Comp. Pathol. **121**:1-11 (1999); J. Ellis *et al.*, "Reproduction of lesions of postweaning multisystemic wasting syndrome in gnotobiotic piglets," J. Vet. Diagn. Invest. **11**:3-14 (1999); J. C. Harding and E.G. Clark, 1997, *supra*). By using this novel infectious PCV2 molecular DNA clone, the clinical disease, pathological lesions and virus distribution exclusively attributable to PCV2 infection are more definitively characterized.

The structural and functional relationships of the PCV genes are better understood due to the availability of the PCV2, PCV1, chimeric PCV1-2, and reciprocal chimeric PCV2-1 infectious DNA clones described herein. Will *et al.*, "Cloned HBV DNA causes hepatitis in chimpanzees," Nature **299**:740-742 (1982), first demonstrated the feasibility of using a cloned hepatitis B virus DNA to infect chimpanzees by direct *in vivo* injection. This approach has since been used to study viral replication and pathogenesis of several other viruses (T. W. Dubensky *et al.*, "Direct transfection of viral and plasmid DNA into the liver or spleen of mice," Proc. Natl. Acad. Sci. USA **81**:7529-7533 (1984); R. Girones *et al.*, "Complete nucleotide sequence of a molecular clone of woodchuck hepatitis virus that is infectious in the natural host," Proc. Natl. Acad. Sci. USA **86**:1846-1849 (1989); N. L. Letvin *et al.*, "Risks of handling HIV," Nature **349**:573 (1991); C. Seeger *et al.*, "The cloned

genome of ground squirrel hepatitis virus is infectious in the animal. Proc. Natl. Acad. Sci. USA. **81**:5849-5852 (1984); E. E. Sparger *et al.*, "Infection of cats by injection with DNA of feline immunodeficiency virus molecular clone," Virology **238**:157-160 (1997); R. Sprengel *et al.*, "Homologous recombination between hepadnaviral genomes following *in vivo* DNA transfection: implications for studies of viral infectivity," Virology **159**:454-456 (1987); H. Will *et al.*, 1982, *supra*; L. Willems *et al.*, "In vivo transfection of bovine leukemia provirus into sheep," Virology **189**:775-777 (1992)).

The construction of an infectious PCV2 molecular DNA clone, and the demonstration of infection by direct injection of cloned PCV2 plasmid DNA into the liver and lymph nodes of pigs in the context of the present invention are advantageous for PCV2 studies. This *in vivo* transfection system will enhance the study of the structural and functional relationship of PCV2 genes using recombinant plasmids constructed *in vitro* to test different regions or genes of PCV2 for their roles in virus replication and pathogenesis in the host. The replication and pathogenesis of PCV2 can be studied *in vivo* without having to produce infectious virus stocks by propagating PCV2 in cell cultures. This is advantageous as serial cell culture passages may select for viral variants. Another advantage of using cloned PCV2 genomic DNA, instead of live virus, for animal studies is its relative ease for quantitation of the inoculation dose. The amount of the cloned PCV2 DNA used for animal inoculation can be easily determined by a spectrophotometer, whereas the dose of live PCV2 virus requires infectivity titration in cell cultures and confirmation of infection by IFA. Direct injection of animals with cloned PCV2 plasmid DNA eliminates the problems associated with the presence of other indigenous swine agents in tissue homogenate inocula in animal studies.

In the present invention, the immunogenic ORF2 capsid gene is switched between the pathogenic PCV2 and the nonpathogenic PCV1 to produce the unique structure of the chimeric PCV1-2 infectious DNA clone. Surprisingly and advantageously, the chimeric PCV1-2 infectious clone replicated, expressed the immunogenic ORF2 capsid antigen *in vitro* and *in vivo*, and induced a specific antibody response against PCV2 ORF2 but retained the nonpathogenic nature of PCV1. The chimeric PCV1-2 infectious DNA clone has the ability to induce a strong immune response against PCV2 while inducing only a limited infection with mild pathologic lesions similar to that of the nonpathogenic PCV1. For vaccine development, the relatively easy storage and stability of cloned DNA, and the

economy of large-scale recombinant PCV2 plasmid DNA and chimeric PCV1-2 DNA clone production provides an attractive means of delivering a live, infectious viral DNA vaccine or genetically engineered, attenuated viral vaccines to pigs. Therefore, the chimeric PCV1-2 infectious DNA clone taught in this invention is a useful vaccine candidate against PCV2 infection and PMWS.

It should be appreciated that all scientific and technological terms used herein have the same meaning as commonly understood by those of ordinary skill in the art. For purposes of this invention, the term "infectious" means that the virus replicates in pigs, regardless of whether or not the virus causes any diseases. "SPF" refers to Specific-pathogen-free pigs. The "gnotobiotic" pigs intend germ-free pigs. The terms "PCV2 plasmid DNA," "PCV2 genomic DNA" and "PCV2 molecular DNA" are being used interchangeably to refer to the same cloned nucleotide sequence.

The infectious PCV1/PCV2 chimeric DNA clone (strain designation "PCV1-2 chimera"), the infectious PCV2 molecular DNA clone (strain designation "PCV2 clone") and the biologically pure and homogeneous PCV2 stock derived from an Iowa sample of PCV2 that had been isolated from a pig with severe PMWS and identified as isolate number 40895 (strain designation "PCV2 #40895") are deposited under the conditions mandated by 37 C.F.R. § 1.808 and maintained pursuant to the Budapest Treaty in the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Virginia 20110-2209, U.S.A. The DNA sequences described herein are contained within 6,490 bp plasmids cloned into pBluescript SK(+) vector (pSK) (Stratagene Inc., La Jolla, CA) and transformed into *Escherichia coli* DH5 α competent cells. The plasmids containing the infectious chimeric PCV1-2 DNA clone (identified as "chimeric porcine circovirus Type 1 (PCV1) and Type 2 (PCV2) infectious DNA clone") and the infectious PCV2 molecular DNA clone (identified as "infectious DNA clone of Type 2 porcine circovirus (PCV2)") have been deposited in the ATCC on December 7, 2001 and have been assigned ATCC Patent Deposit Designations PTA-3912 and PTA-3913, respectively. It should be appreciated that other plasmids, which may be readily constructed using site-directed mutagenesis and the techniques described herein, are also encompassed within the scope of the present invention. The biologically pure and homogeneous PCV2 sample of isolate number 40895 (identified as "Type 2 porcine circovirus (PCV2)") has also been deposited in the ATCC on December 7, 2001 and has been assigned

ATCC Patent Deposit Designation PTA-3914. The genomic (nucleotide) sequence of the PCV2 isolate number 40895 has been deposited with the Genbank database and has been publicly available since July 23, 2000 under accession number AF264042.

The following examples demonstrate certain aspects of the present invention. However, it is to be understood that these examples are for illustration only and do not purport to be wholly definitive as to conditions and scope of this invention. It should be appreciated that when typical reaction conditions (*e.g.*, temperature, reaction times, *etc.*) have been given, the conditions both above and below the specified ranges can also be used, though generally less conveniently. The examples are conducted at room temperature (about 23°C to about 28°C) and at atmospheric pressure. All parts and percents referred to herein are on a weight basis and all temperatures are expressed in degrees centigrade unless otherwise specified.

A further understanding of the invention may be obtained from the non-limiting examples that follow below.

EXAMPLE 1

Generation of a PK-15 Cell Line Free of PCV1 Contamination

The source of the PCV2 isolate was from a spleen tissue sample of a pig with naturally occurring PMWS (PCV2 serial identification number 40895, referred to as "isolate 40895") (M. Fenau *et al.*, 2000, *supra*). Immunohistochemical staining (IHC) with PCV2-specific antibody confirmed the presence of PCV2 antigen in the tissue. The spleen tissues were stored at -80°C until use.

The PK-15 cell line purchased from the American Type Culture Collection (ATCC accession number CCL-33) was persistently infected with PCV1 (G. C. Dulac and A. Afshar, 1989, *supra*). Since only a subpopulation of PK-15 cells was persistently infected (*id.*), a PK-15 cell line that is free of PCV1 contamination by end-point dilution was generated. Protocol proceeded as follows: PK-15 cells were grown in MEM with Earle's salts and L-glutamine (Life Technologies, Inc., Grand Island, NY) supplemented with 10% fetal bovine serum (FBS) and 1X antibiotic (Life Technologies, Inc.). Confluent cell monolayers were trypsinized, and the cells were then counted and serially diluted to an end point with one cell per 0.2 ml. The end point dilution was plated in 96-well plates and allowed to grow into a

monolayer starting from a single cell. Cells from each well were tested for PCV1 DNA using a PCR-RFLP assay capable of detecting and differentiating PCV1 and PCV2 (M. Fenaux *et al.*, 2000, *supra*). PK-15 cells from wells that were tested negative for PCV1 by the PCR-RFLP assay were subsequently expanded. The PCV1 free PK-15 cell line was subcultured five additional passages and was found negative for PCV1 DNA by PCR at each passage.

Four cell lines that were negative for PCV1 contamination were produced by the end-point dilution of the persistently infected PK-15 cells from ATCC. The cell lines remained negative for PCV1 by PCR after the five additional passages. One of the cell lines was subsequently expanded and was shown to be able to support PCV2 replication when the cells were transfected with the PCV2 molecular DNA clone (Fig. 2) and infected with PCV2 virus. The cloned cells were further used for the *in vitro* transfection of PCV2 molecular DNA clone to generate a biologically pure PCV2 infectious virus stock for the animal inoculation experiment.

EXAMPLE 2

Construction of the PCV2 Infectious DNA Clone

To construct a PCV2 molecular DNA clone, a pair of PCR primers was designed according to the published sequence of the PCV2 isolate 40895 (M. Fenaux *et al.*, 2000, *supra*): forward primer F-PCVSAC2 (5'-GAACCGCGGGCTGGCTGAACTTTTGAAAGT-3'), set forth in SEQ ID NO:5, and reverse primer R-PCVSAC2 (5'-GCACCGCGGAAATTTCTGACAAACGTTACA-3'), set forth in SEQ ID NO:6. This pair of primers amplifies the complete genome of PCV2 with an overlapping region containing the unique *SacII* restriction enzyme site (Fig. 1). DNA was extracted using the QIAamp DNA Minikit (Qiagen, Inc., Valencia, CA) from a spleen tissue sample of a pig with naturally occurring PMWS (isolate 40895) (M. Fenaux *et al.*, 2000, *supra*). The extracted DNA was amplified by PCR with AmpliTaq Gold polymerase (Perkin-Elmer, Norwalk, CT). The PCR reaction consisted of an initial enzyme activation step at 95°C for 9 min, followed by 35 cycles of denaturation at 94°C for 1 min, annealing at 48°C for 1 min, extension at 72°C for 3 min, and a final extension at 72°C for 7 min. The PCR product of expected size

was separated by gel electrophoresis and purified with the glassmilk procedure with a GeneClean Kit (Bio 101, Inc., La Jolla, CA).

To construct a molecular DNA clone containing a tandem dimer of PCV2 genome, the PCR product containing the complete PCV2 genome was first ligated into the advanTAge plasmid vector (Clontech, Palo Alto, CA). *E. Coli* DH5 α competent cells were transformed. The recombinant plasmids were verified by restriction enzyme digestion. The full length PCV2 genomic DNA was excised from the advanTAge vector by digestion with *Sac*II restriction enzyme. The digested PCV2 genomic DNA was ligated with T4 DNA ligase at 37°C for only 10 min, which favors the production of tandem dimers. The tandem dimers were subsequently cloned into pBluescript SK(+) vector (pSK) (Stratagene Inc., La Jolla, CA) (Fig. 1). Recombinant plasmids containing tandem dimers of PCV2 genome (herein referred to as PCV2 molecular DNA clone) were confirmed by PCR, restriction enzyme digestion, and DNA sequencing. The DNA concentration of the recombinant plasmids was determined spectrophotometrically.

Specifically, the complete genome of the PCV2 (isolate 40895) was amplified by PCR to construct the infectious PCV2 molecular DNA clone. Two copies of the complete PCV2 genome were ligated in tandem into the pSK vector to produce the PCV2 molecular DNA clone (Fig. 1). The infectivity of the PCV2 molecular DNA clone was determined by *in vitro* transfection of the PK-15 cells. IFA with PCV2-specific antibody confirmed that the molecular DNA clone is infectious *in vitro* and that about 10-15% of the PK-15 cells were transfected. PCV2-specific antigen was visualized by IFA in the nucleus, and to a lesser degree, cytoplasm of the transfected cells (Fig. 2). The cells mock-transfected with the empty pSK vector remained negative for PCV2 antigen.

EXAMPLE 3

In Vitro Transfection with the PCV2 Molecular DNA Clone and Generation of a Biologically Pure and Homogenous PCV2 Infectious Virus Stock

To test the infectivity of the molecular DNA clone *in vitro*, PK-15 cells free of PCV1 contamination were grown in 8-well LabTek chamber slides. When the PK-15 cells reached about 85% confluency, cells were transfected with the molecular DNA clone using Lipofectamine Plus Reagents according to the protocol supplied by the manufacturer (Life

Technologies, Inc). Mock-transfected cells with empty pSK vector were included as controls. Three days after transfection, the cells were fixed with a solution containing 80% acetone and 20% methanol at 4° C for 20 min., and an immunofluorescence assay using a PCV2-specific rabbit polyclonal antisera was performed to determine the *in vitro* infectivity of the molecular DNA clone (see below).

To generate a biologically pure and homogeneous PCV2 infectious virus stock for the animal inoculation experiment, PK-15 cells free of PCV1 contamination were cultivated in T-25 culture flasks and transfected with the PCV2 molecular DNA clone. PK-15 cells were grown to about 85% confluency in T-25 flasks. The cells were washed once with sterile PBS buffer before transfection. For each transfection reaction in a T-25 flask, 12 µg of the PCV2 plasmid DNA was mixed with 16 µl of Plus Reagent in 0.35 ml of MEM media. A flask of mock-transfected cells with empty pSK vector was included as the negative control. After incubation at room temperature for 15 min., 50 µl of Lipofectamine Reagent diluted in 0.35 ml of MEM media was added to the mixture and incubated at room temperature for another 15 min. The transfection mixture was then added to a T-25 flask of PK-15 cells containing 2.5 ml of fresh MEM. After incubation at 37° C for 3 hrs, the media was replaced with fresh MEM media containing 2 % FBS and 1 X antibiotics. The transfected cells were harvested at 3 days post transfection and stored at -80° C until use. The infectious titer of the virus stock was determined by IFA (see below).

Basically, biologically pure and homogenous PCV2 infectious virus stock was generated by transfection of PK-15 cells with the PCV2 molecular DNA clone. PCV2 virions produced by *in vitro* transfection were infectious since the transfected cell lysates were successfully used to infect PK-15 cells. Thus, the PCV2 molecular DNA clone is capable of producing infectious PCV2 virions when transfected *in vitro*. The infectious titer of the homogenous PCV2 virus stock prepared from transfected cells was determined to be $1 \times 10^{4.5}$ TCID₅₀/ml. This virus stock was used to inoculate pigs in Group 2. Lysates of cells mock-transfected with the empty pSK vector were unable to infect PK-15 cells.

EXAMPLE 4

Virus Titration by Immunofluorescence Assay (IFA)

To determine the infectious titer of the homogenous PCV2 virus stock, PK-15 cells were cultivated on 8-well LabTek chamber slides. The virus stock was serially diluted 10-fold in MEM, and each dilution was inoculated onto 10 wells of the monolayers of the PK-15 cells growing on the LabTek chamber slides. Wells of non-inoculated cells were included as controls. The infected cells were fixed at 3 days post inoculation with a solution containing 80% acetone and 20% methanol at 4°C for 20 min. After washing the cells with PBS buffer, the infected cells were incubated with a 1:1,000 diluted PCV2-specific rabbit polyclonal antibody (S. D. Sorden *et al.*, "Development of a polyclonal-antibody-based immunohistochemical method for the detection of type 2 porcine circovirus in formalin-fixed, paraffin-embedded tissue," J. Vet. Diagn. Invest. 11:528-530 (1999)) at 37°C for 1 hr. The cells were then washed three times with PBS buffer, and incubated with a secondary FITC-labeled goat anti-rabbit IgG (Kirkegaard & Perry Laboratories Inc, Gaithersburg, MD) at 37°C for 45 min. After washing the slides three times with PBS buffer, and the slides were mounted with fluoromount-G, cover-slipped and examined under a fluorescence microscope. The 50% tissue culture infectious dose per ml (TCID₅₀/ml) was calculated. Initially, cells were transfected with a plasmid construct containing a single copy of PCV2 genome but the infectious PCV2 titer from the single genome construct is much lower than the one containing the tandem genome. Therefore, the plasmid construct containing the dimeric form of PCV2 genome was used for the *in vitro* and *in vivo* transfection experiments.

EXAMPLE 5

In Vivo Transfection of Pigs With the PCV2 Molecular DNA Clone and ExperimentalInoculation of Pigs With the Homogeneous PCV2 Infectious Virus Stock

Forty specific-pathogen-free (SPF) swine of 4 weeks of age were randomly assigned into 4 rooms of 10 animals each. Prior to inoculation, the SPF pigs were tested for antibodies to PCV, PRRSV, PPV and swine hepatitis E virus. Pigs in Group 1 were uninoculated and served as negative controls. Pigs in Group 2 were each inoculated intranasally with about 1.9×10^5 TCID₅₀ of the PCV2 infectious virus stock derived from the PCV2 molecular DNA clone. Pigs in Group 3 received direct intrahepatic injection of the recombinant plasmid DNA

of the PCV2 molecular clone. Each pig was injected with a total of 200 µg of recombinant plasmid DNA (the cloned PCV2 plasmid DNA), through an ultrasound-guided technique, into 6 different sites of the liver. Pigs in Group 4 were each injected with a total of 200 µg of the recombinant PCV2 plasmid DNA directly into the superficial iliac lymph nodes, and each lymph node received two separate injections. The animals were monitored daily for clinical signs of disease. Serum samples were collected from each animal at 0, 7, 14, 21, 28, 35 days post inoculation (DPI). At 21 DPI, five pigs were randomly selected from each group and necropsied. The remaining five animals in each group were necropsied at 35 DPI. Various tissues and organs were collected during necropsy and processed for histological examination and immunohistochemical staining (see below).

The results are shown in Table 1 below. All inoculated pigs from Groups 2, 3 and 4 were negative for PCV2 antibodies at 0 DPI. Two pigs in the uninoculated control Group 1 had detectable PCV2 maternal antibody at 0 DPI. The maternal antibody in these two piglets waned by 7 DPI. No seroconversion to PCV2 antibody was detected in any of the 10 uninoculated control pigs. In Group 2 pigs intranasally inoculated with PCV2 infectious virus, 1 piglet seroconverted to PCV2 antibody at 21 DPI. By 35 DPI, 4 of the 5 remaining Group 2 pigs had seroconverted. Seroconversion in transfected animals from Groups 3 and 4 first appeared at 28 DPI. By 35 DPI, 5 of 5 remaining pigs from Group 3 and 3 of 5 remaining pigs from Group 4 had seroconverted to PCV2 antibody.

PPV antibodies were tested at 3 and 21 DPI for all pigs, and at 35 DPI for the remaining pigs. Maternal antibodies to the ubiquitous swine agent PPV were detected in the SPF piglets. The PPV HI antibody titers in all piglets but one decreased significantly from 3 DPI (an average titer of 1:2,665) to 21 DPI (an average titer of 1:246), indicating the antibody detected in these piglets was passively derived. One piglet had a slightly increased PPV HI titer from 1:32 at 3 DPI to 1:64 at 21 DPI, which is likely due to testing variation. Serum samples collected from all pigs at 0, 21, and 35 DPI were further tested for PPV DNA with a published PCR assay (J. M. Soucie *et al.*, "Investigation of porcine parvovirus among persons with hemophilia receiving Hyate: C porcine factor VIII concentrate," *Transfusion* 40:708-711 (2000)). No PPV viremia was detected from any pigs at any DPI, further indicating the pigs were not infected by PPV.

Table 1. Seroconversion to PCV2 Specific Antibodies in Pigs Inoculated With PCV2 Live Virus or Directly Injected With Cloned PCV2 Plasmid DNA

Group	Inocula	Route of Inoculation	Days Postinoculation					
			0	7	14	21	28	35
1	None		2/10 ^a	0/10	0/10	0/10	0/5	0/5
2	PCV2 live virus ^b	Intranasal	0/10	0/10	0/10	1/10	1/5	4/5
3	PCV2 DNA ^c	Intrahepatic	0/10	0/10	0/10	0/10	1/5	5/5
4	PCV2 DNA ^c	Intralymphoid	0/10	0/10	0/10	0/10	1/5	3/5

^aPCV2 antibody was measured with an ELISA, number positive/number tested.

^bA biologically pure and homogeneous PCV2 virus stock generated by transfection of PK-15 cells with PCV2 molecular DNA clone.

^cCloned PCV2 genomic DNA in pSK plasmid.

EXAMPLE 6

PCR-RFLP Analyses

To measure PCV2 viremia in pigs transfected with PCV2 molecular DNA clone and in pigs infected with PCV2 infectious virus stock, serum samples collected at different DPIs were tested for the presence of PCV2 DNA by the general methods of a PCR-RFLP assay previously described (M. Fenaux *et al.*, 2000, *supra*). Viral DNA was extracted from 50 µl of each serum sample using the DNAzol® reagent according to the protocol supplied by the manufacturer (Molecular Research Center, Cincinnati, OH). The extracted DNA was resuspended in DNase-, RNase-, and proteinase-free water and tested for PCV2 DNA by PCR-RFLP (*id.*). PCR products from selected animals were sequenced to verify the origin of the virus infecting pigs.

Serum samples were collected from all control and inoculated animals at 0, 7, 14, 21, 28, and 35 DPIs and assayed for PCV2 viremia by detection of PCV2 DNA (*id.*). The results are shown in Table 2 below. PCV2 DNA was not detected in the Group 1 uninoculated control pigs at any DPI. Viremia was detected in 7/10 pigs from Group 2 at 14 DPI and 8/10 by 35 DPI. Viremia lasted only a few weeks as the PCV2 DNA was not detectable at 28 DPI and 35 DPI in all 5 remaining pigs from Group 2. In Group 3 pigs that were intrahepatically injected with PCV2 molecular DNA clone, 8/10 pigs were viremic at 14 DPI, and 9/10 pigs had detectable viremia by 35 DPI. Group 4 pigs were injected with PCV2 molecular DNA clone into the lymph nodes. Two of 10 pigs at 14 DPI and 8 of 10 pigs at 21 DPI from Group 4 were viremic. The results show that PCV2 molecular DNA clone is infectious when injected directly into the liver and superficial iliac lymph nodes of SPF pigs. PCR products amplified from selected animals were sequenced. The sequence of the PCR products amplified from selected animals was identical to the corresponding region of the PCV2 molecular DNA clone.

Table 2. Detection of Viremia (PCV2 DNA) by PCR in Sera of Inoculated and Control Pigs

Group	Inocula	Route of Inoculation	Days Postinoculation						Total
			0	7	14	21	28	35	
1	None		0/10 ^a	0/10	0/10	0/10	0/5	0/5	0/10
2	PCV2 live virus ^b	Intranasal	0/10	0/10	7/10	5/10	0/5	0/5	8/10
3	PCV2 DNA ^c	Intrahepatic	0/10	0/10	8/10	6/10	3/5	3/5	9/10
4	PCV2 DNA ^c	Intralymphoid	0/10	0/10	2/10	8/10	2/5	0/5	8/10

^a10 pigs in each group, number positive/number tested.

^bA biologically pure and homogeneous PCV2 virus stock generated by transfection of PK-15 cells with PCV2 molecular DNA clone.

^cCloned PCV2 genomic DNA in pSK plasmid.

EXAMPLE 7

Clinical Evaluation

Pigs were weighed on 0 DPI and at the time of necropsy. Rectal temperatures and clinical respiratory disease scores, ranging from 0 to 6 (0 = normal, 6 = severe) (P. G. Halbur *et al.*, "Comparison of the pathogenicity of two U.S. porcine reproductive and respiratory syndrome virus isolates with that of the Lelystad virus," Vet. Pathol. **32**:648-660 (1995)), were recorded every other day from 0 to 35 DPI. Clinical observations including evidence of central nervous system disease, liver disease (icterus), musculoskeletal disease, and changes in body condition, were also recorded daily.

To evaluate the gross pathology and histopathology, five pigs from each group were randomly selected for necropsies at 21 and 35 DPI. The necropsy team was blinded to infection status of the pigs at necropsy. Complete necropsies were performed on all pigs. An estimated percentage of the lung with grossly visible pneumonia was recorded for each pig based on a previously described scoring system (*id.*). The scoring system is based on the approximate volume that each lung lobe contributes to the entire lung: the right cranial lobe, right middle lobe, cranial part of the left cranial lobe, and the caudal part of the left cranial lobe each contribute 10% of the total lung volume, the accessory lobe contributes 5%, and the right and left caudal lobes each contribute 27.5%. Other lesions such as enlargement of lymph nodes were noted separately. Sections for histopathologic examination were taken from nasal turbinate, lungs (seven sections) (*id.*), heart, brain, lymph nodes (tracheobronchial, iliac, mesenteric, subinguinal), tonsil, thymus, liver, gall bladder, spleen, joints, small intestine, colon, pancreas, and kidney. The tissues were examined in a blinded fashion and given a subjective score for severity of lung, lymph node, and liver lesions. Lung scores ranged from 0 (normal) to 3 (severe lymphohistiocytic interstitial pneumonia). Liver scores ranged from 0 (normal) to 3 (severe lymphohistiocytic hepatitis). Lymph node scores were for an estimated amount of lymphoid depletion of follicles ranging from 0 (normal or no lymphoid depletion) to 3 (severe lymphoid depletion and histiocytic replacement of follicles).

The serology protocol involved collecting blood on arrival at 11 to 12 days of age, and from all pigs at 0, 7, 14, 21, 28, and 35 DPIs. Serum antibodies to PRRSV were assayed using Herd Check PRRSV ELISA (IDEXX Laboratories, Westbrook, MA). Serum

antibodies to PPV were detected by a hemagglutination inhibition (HI) assay (H. S. Joo *et al.*, "A standardized haemagglutination inhibition test for porcine parvovirus antibody," Aust. Vet. J. **52**:422-424 (1976)). Serum antibodies to PCV2 were detected by a modified indirect ELISA based on the recombinant ORF2 protein of PCV2 (P. Nawagitgul *et al.*, "Modified indirect porcine circovirus (PCV) type 2-based and recombinant capsid protein (ORF2)-based ELISA for the detection of antibodies to PCV," Immunol. Clin. Diagn. Lab Immunol. **1**:33-40 (2002)). A partially purified PCV2 antigen was prepared from Hi Five cells (Invitrogen, Carlsbad, CA) infected with recombinant baculovirus containing the major capsid ORF2 protein of PCV2 (P. Nawagitgul *et al.*, "Open reading frame 2 of porcine circovirus type 2 encodes a major capsid protein," J. Gen. Virol. **81**:2281-2287 (2000)). Cell lysates of Hi Five cells infected with wild-type baculovirus were prepared similarly and served as negative control antigen. The Immulon 2 HB polystyrene microtiter plates (Dynex Technologies Inc, Chantilly, VA) were coated with optimal concentrations of positive and negative antigens at 4°C for 36 hrs. One hundred µl of each serum sample diluted 1:100 in 5% milk diluent (Kirkegaard & Perry Laboratories, Inc.) was added into each well. The serum samples were tested in quadruplicate: 2 wells for negative control antigen and 2 parallel wells for PCV2 antigen. Positive control and negative control sera were included in each plate. The sera were incubated at 37°C for 30 min. and then washed 5 times with 0.1 M PBS buffer containing 0.1% Tween-20. A peroxidase-labeled secondary anti-swine IgG (Sigma Co, St. Louis, MO) was incubated at 37°C for 30 min. The plates were washed again and incubated with 2,2'-azino-di-(3-ethylbenzthiazoline-6-sulfonate) (Kirkegaard & Perry Laboratories Inc) at 37°C for 15 min. for color development. The optical density (OD) was read at 405 nm. The corrected OD of each tested and control sera was calculated by subtraction of mean OD value of the wells containing negative antigen from that of the parallel wells containing PCV2 antigen. The data was normalized by dividing the corrected OD value of a tested serum sample (S) with that of the positive control serum (P) and reported as S/P ratios. The samples with S/P ratios ≤ 0.12 , 0.12 to 0.2, and > 0.2 were considered as negative, equivocal and positive, respectively.

From the results of the clinical evaluation, none of the control and inoculated pigs showed obvious signs of disease resembling those of clinical PMWS. There was no difference in weight gain or mean rectal temperatures between any of the four groups. The

control pigs of Group 1 remained normal throughout the experiment. There was mild transient respiratory disease observed in the majority of the pigs in PCV2 DNA-transfected and PCV2 virus-infected groups from 8 to 14 DPI. This was characterized by mild dyspnea (clinical respiratory scores of 1 to 2) of one-to-two days duration in individual pigs and 5-6 days duration for the group.

There were no gross lesions observed in the control pigs at necropsy. Pigs in the three inoculated groups had gross lesions limited to the lungs and lymph nodes (see Table 3, below). The lesions were similar among pigs in the PCV2 plasmid DNA-transfected and PCV2 virus-infected groups. Lungs failed to collapse and had random, multifocal, moderately well-demarcated areas of tan-to-purple consolidation involving 0-2% of the lung (Fig. 3) at 21 DPI, and 0-13% of the lung at 35 DPI. Lymph nodes were systemically enlarged 2 to 5 times normal size, firm, and tan (Fig. 3) at both 21 and 35 DPI in most of the pigs from all three PCV2-inoculated groups.

Microscopic examination revealed no lesions in any tissues of the control pigs except for the livers. Eight of ten control pigs had very mild multifocal lymphoplasmacytic inflammation predominately in the periportal regions of the liver as is commonly observed in normal pigs and considered normal background (P. G. Halbur *et al.*, 2001, *supra*).

Pigs from the two PCV2 plasmid DNA-transfected groups (intrahepatic and intralymphoid) and the PCV2 virus-infected group (intranasal) had similar lesions in brain, lung, heart, kidney, lymphoid tissues (tonsil, lymph nodes, spleen), ileum, and liver (see Table 4, below). Brain lesions were observed in 23/30 of the pigs from the three inoculated groups and were characterized as mild-to-moderate multifocal lymphoplasmacytic meningoencephalitis with perivascular cuffing and gliosis. Lung lesions were observed in 28/30 PCV2-inoculated pigs and characterized as mild-to-moderate peribronchiolar lymphoplasmacytic and histiocytic bronchointerstitial pneumonia (Fig. 3C). One pig from the PCV2 virus-infected Group 2 necropsied at 21 DPI, and one pig each from the two PCV2 plasmid DNA-transfected groups necropsied at 35 DPI had ulcerative and proliferative bronchiolitis with fibroplasia and granulomatous inflammation in the lamina propria and peribronchiolar regions of bronchi. Mild multifocal lymphoplasmacytic myocarditis was also observed in 18/30 PCV2-inoculated pigs. In 14/30 of the PCV2-inoculated pigs, mild-to-moderate multifocal lymphoplasmacytic interstitial nephritis was observed. No lesions were observed

in the thymuses. Mild-to-moderate lymphoid depletion (Fig. 4B) and histiocytic replacement of follicles was observed in the tonsil of 8/30, in the spleen of 7/30, and in the lymph nodes of 26/30 of the PCV2-inoculated pigs. Moderate granulomatous lymphadenitis with giant cells (Fig. 4C) was observed at 21 DPI in three pigs inoculated intranasally with PCV2 virus, and in one pig at 35 DPI in each of the PCV2 plasmid DNA-transfected groups. Mild lymphoplasmacytic and histiocytic enterocolitis were observed in 3/5 pigs in the PCV2 virus-infected group, in 3/5 pigs in the PCV2 plasmid DNA intrahepatically-transfected group, and 1/5 pigs in the PCV2 plasmid DNA intralymphoid-transfected group at 35 DPI. One pig in each of the PCV2 plasmid DNA-transfected groups had mild lymphoid depletion with histiocytic replacement and low numbers of giant cells in the Peyer's patches. Mild-to-moderate lymphoplasmacytic hepatitis was observed in 29/30 of the three PCV2-inoculated pigs. Low numbers of widely scattered individually necrotic hepatocytes surrounded by lymphohistiocytic inflammation was observed in one pig in each of the PCV2 plasmid DNA-transfected groups at 21 DPI. Lesions in other tissues were unremarkable.

Microscopic lesions in the lung, liver and lymph nodes were scored according to published scoring systems (Table 4, below) (P. G. Halbur *et al.*, 2001, *supra*; P. G. Halbur *et al.*, 1995, *supra*). There were no acceptable scoring systems for other tissues and organs. The average scores of lesions in lung and lymph nodes in pigs of the three PCV2-inoculated groups were statistically different from those in the control pigs of Group 1. The average scores of the liver lesions in pigs of the three PCV2-inoculated groups are not statistically different from those of control pigs.

Table 3. Gross Lesions of Lung and Lymph Nodes in Control and PCV2-Inoculated Pigs

Group	Inocula	Route of Inoculation	21 DPI		35 DPI	
			Lymph Nodes	Lung	Lymph Nodes	Lung
1	None		0/5 ^a	0/5	0/5	0/5
2	PCV2 live virus ^b	Intranasal	5/5	1/5(0-1) ^c	5/5	4/5(0-5)
3	PCV2 DNA ^d	Intrahepatic	2/5	2/5(0-2)	5/5	2/5(0-13)
4	PCV2 DNA ^d	Intralymphoid	4/5	5/5(0-1)	3/5	1/5(0-9)

^aFive pigs from each group were necropsied at 21 DPI, and the remaining 5 pigs were necropsied at 35 DPI. Number positive / number tested.

^bA biologically pure and homogeneous PCV2 virus stock generated by transfection of PK-15 cells with PCV2 molecular DNA clone.

^cNumber with lesions / number tested (range of the estimated percent of the lung affected by grossly visible pneumonia lesions, 0-100%)

^dCloned PCV2 genomic DNA in pSK plasmid.

Table 4. Distribution of Histopathological Lesions in Control and PCV2-Inoculated Pigs

Route of													
Group	Inocula	Inoculation	DPI ^a	Lung ^b	Liver ^c	LN ^d	Spleen	Thymus	Ileum	Brain	Heart	Kidney	Tonsil
1	None		21	0/5(0.0)	4/5(0.8)	0/5(0.0)	0/5	0/5	0/5	0/5	0/5	0/5	0/5
2	PCV2 virus	Intranasal	35	0/5(0.0)	4/5(0.8)	0/5(0.0)	0/5	0/5	0/5	0/5	0/5	0/5	0/5
			21	5/5(1.6)	5/5(1.2)	3/5(1.2)	1/5	0/5	0/5	4/5	3/5	1/5	0/5
3	PCV2 DNA	Intrahepatic	35	3/5(0.6)	4/5(1.0)	4/5(0.8)	3/5	0/5	3/5	4/5	0/5	1/5	3/5
			21	5/5(1.0)	5/5(1.0)	5/5(1.0)	1/5	0/5	0/5	5/5	4/5	1/5	0/5
4	PCV2 DNA	Intralymphoid	35	5/5(1.2)	5/5(1.0)	4/5(1.0)	2/5	0/5	3/5	3/5	4/5	5/5	3/5
			21	5/5(1.2)	5/5(1.0)	5/5(0.8)	0/5	0/5	0/5	4/5	4/5	3/5	0/5
			35	5/5(1.0)	5/5(1.2)	5/5(1.4)	0/5	0/5	1/5	3/5	3/5	3/5	2/5

^aDays postinoculation (DPI): 5 animals from each group were necropsied at 21 DPI and the remaining 5 animals from each group were necropsied at 35 DPI

^bNumber positive/number tested (Average histological lung score: 0=normal, 1=mild interstitial pneumonia, 2=moderate, 3=severe)

^cNumber positive/number tested (Average histological liver score: 0=normal, 1=mild hepatitis, 2=moderate, 3=severe)

^dNumber positive/number tested (Average histological lymphoid (LN) depletion score: 0=normal, 1=mild, 2=moderate, 3=severe)

EXAMPLE 8

Immunohistochemistry

Immunohistochemistry (IHC) detection of PCV2-specific antigen was performed on all tissues collected during necropsies at DPIs 21 and 35. A rabbit polyclonal PCV2-specific antiserum was used for the IHC, and the general procedures have been previously described (S. D. Sorden *et al.*, 1999, *supra*).

For the detection and tissue distribution of PCV2 antigen, IHC staining of PCV2 antigen was done on brain, lungs, turbinate, heart, kidneys, tonsil, lymph nodes, spleen, thymus, ileum, liver, gall bladder and pancreas of all pigs necropsied at 21 and 35 DPI. All tissues from the control pigs were negative for PCV2 antigen. Tissue distribution of PCV2 antigen in the three PCV2-inoculated groups was similar (see Table 5, below). In the brain, the PCV2 antigen was found predominately in mononuclear cells, fibroblast-like cells, and endothelial cells in the meninges and choroid plexus and less often in endothelial cells and perivascular mononuclear cells in the cerebrum and cerebellum. In the lungs, PCV2 antigen was detected within alveolar and septal macrophages and in fibroblast-like cells in the lamina propria of airways (Fig. 3D). In the heart, PCV2 antigen was detected in widely scattered macrophages and endothelial cells. In kidneys, PCV2 antigen was detected within tubular epithelial cells and mononuclear cells in the interstitium. In the lymphoid tissues (lymph nodes, spleen, tonsil, and Peyer's patches), PCV2 antigen was detected primarily within macrophages and dendritic-like cells and giant cells within follicles (Fig. 4D). PCV2 antigen was also detected within macrophages in the lamina propria of the small intestine. In the liver, PCV2 antigen was detected within mononuclear cells and Kupffer cells. PCV2 antigen was not detected in turbinate, thymus, or gall bladder.

Table 5. Detection and Distribution of PCV2-Specific Antigen by Immunohistochemistry in Control and PCV2-Inoculated Pigs

Route of													
Group	Inocula	Inoculation	DPI ^a	Lung	Liver	LN	Spleen	Thymus	Ileum	Brain	Heart	Kidney	Tonsil
1	None		21	0/5 ^b	0/5	0/5	0/5	0/5	0/5	0/5	0/5	0/5	0/5
2	PCV2 virus	Intranasal	35	0/5	0/5	0/5	0/5	0/5	0/5	0/5	0/5	0/5	0/5
			21	4/5	5/5	5/5	3/5	0/5	3/5	3/5	1/5	1/5	2/5
3	PCV2 DNA	Intrahepatic	35	1/5	2/5	3/5	2/5	0/5	0/5	2/5	0/5	0/5	0/5
			21	5/5	5/5	5/5	5/5	0/5	0/5	5/5	1/5	0/5	2/5
4	PCV2 DNA	Intralymphoid	35	4/5	4/5	3/5	4/5	0/5	3/5	4/5	2/5	2/5	3/5
			21	4/5	4/5	5/5	4/5	0/5	3/5	3/5	0/5	0/5	3/5
			35	3/5	4/5	5/5	4/5	0/5	2/5	3/5	1/5	0/5	4/5

^aDays postinoculation (DPI): 5 animals from each group were necropsied at 21 DPI and the remaining 5 animals from each group were necropsied at 35 DPI

^bNumber positive/number tested.

EXAMPLE 9

Construction of the Nonpathogenic PCV1 Infectious DNA Clone

The procedure used to construct a PCV1 infectious DNA clone is essentially the same as that described herein for PCV2. A pair of PCR primers, KPNPCV1.U set forth in SEQ ID NO:7 and KPNPCV1.L set forth in SEQ ID NO:8 (see Table 6, below), was designed based on the published sequence of PCV1. This pair of primers amplifies the complete genome of PCV1 with an overlapping region containing the unique *KpnI* restriction enzyme site. The DNA of the PCV1 virus was extracted from the contaminated ATCC PK-15 cell line that was obtained from the American Type Culture Collection (ATCC accession number CCL-33). The PCV1 DNA was extracted from the ATCC PK-15 cells persistently infected with PCV1, using the QIAmp DNA minikit (Qiagen, Inc., Valencia, CA.). The extracted DNA was amplified by PCR with AmpliTaq Gold Polymerase (Perkin-Elmer, Norwalk, CT). The PCR cycles consisted of an initial step of 95°C for 10 min., followed by 35 cycles of denaturation at 94°C for 1 min., annealing at 48°C for 1 min., extension at 72°C for 2 min., and a final extension at 72°C for 7 min. The PCR product of expected size was separated by gel electrophoresis and purified by the glassmilk procedure using a GeneClean Kit (Bio 101, Inc., La Jolla, CA). The purified PCR product containing the complete PCV1 genome was first ligated into the advanTAge plasmid vector (Clontech, Palo Alto, CA). *Escherichia coli* DH5 α competent cells were used for transformation. The recombinant plasmids were verified by restriction enzyme digestion. The full length PCV1 genomic DNA was excised from the advanTAge vector by digestion with *KpnI* restriction enzyme. The full-length PCV1 genomic DNA was ligated into pBluescript SK(+) (pSK) vector (Stratagene, La Jolla, CA) with T4 DNA ligase at 37°C overnight. Recombinant plasmids containing the full-length PCV1 genome were isolated with a Qiagen plasmid mini kit (Qiagen, Valencia, CA) and were verified by restriction enzyme digestion and DNA sequencing. The full-length PCV1 genomic DNA was excised from the pSK vector by *KpnI* digestion, and dimmerized to make the PCV1 infectious DNA clone as described above in Example 2 for the PCV2 infectious clone. These tandem dimers were made because the dimmerized tandem DNA clones are advantageously found to be more efficient to transfect cells and produce infectious virions. To make the tandem dimer of the PCV1 DNA, the digested PCV1 genomic DNA was ligated with T4 DNA ligase at 37°C for only 10 min., which favors the production of

tandem dimers. The tandem dimers were subsequently cloned into pBluescript SK(+) (pSK) vector (Stratagene, La Jolla, CA). Recombinant plasmids containing tandem dimers of PCV1 genome (herein referred to as "PCV1 DNA clone") were confirmed by PCR, restriction enzyme digestion, and DNA sequencing. The DNA concentration of the recombinant plasmids was determined spectrophotometrically.

5

Table 6. Oligonucleotide Primers Employed in this Invention

Primer	Direction	Primer Sequence	Application
<i>Construction primers:</i>			
KPNPCV1.U.	> ^a	5'-TTTGGTACCCGAAGCCGATT-3' (corresponds to SEQ ID NO:7)	PCV1 DNA clone construction
KPNPCV1.L.	<	5'-ATTGGTACCTCCGTGGATTGTTCT-3' (corresponds to SEQ ID NO:8)	PCV1 DNA clone construction
Hpa I-2	<	5'-GAAGTTAACCCCTAAATGAATAAAATAAACCATTCAG-3' (corresponds to SEQ ID NO:9)	PCV1-2 DNA clone construction
Nar I-3	>	5'-GGTGGCGCCTCCTTGGATACGTCATCCTATAAAAGTG-3' (corresponds to SEQ ID NO:10)	PCV1-2 DNA clone construction
Psi I-5	>	5'-AGGTTATAAGTGGGGGTCTTTAAGATTAA-3' (corresponds to SEQ ID NO:11)	PCV1-2 DNA clone construction
Acl I-6	<	5'-GGAAACGTTACCGCAGAAAGACACC-3' (corresponds to SEQ ID NO:12)	PCV1-2 DNA clone construction
Bgl II-ORF2	>	5'-ACTATAGATCTTTATTCATTTAGAGGGTCTTTCAG-3' (corresponds to SEQ ID NO:13)	PCV2-1 DNA clone construction
SpH I-ORF2	<	5'-TACGGGCATGCATGACGTGGCCAGGAGG-3' (corresponds to SEQ ID NO:14)	PCV2-1 DNA clone construction
Bgl II-PCV2	<	5'-AGACGAGATCTATGAATAATAAAACCATTCAGAAAG-3' (corresponds to SEQ ID NO:15)	PCV2-1 DNA clone construction
SpH I-PCV2	>	5'-CGTAAGCATGCAGCTGAAACGAAAGAAAGTG-3' (corresponds to SEQ ID NO:16)	PCV2-1 DNA clone construction
<i>Detection primers:</i>			
MCV1	>	5'-GCTGAACCTTTTGAAAGTGAGCGGG-3' (corresponds to SEQ ID NO:17)	PCV1 and PCV2 detection
MCV2	<	5'-TCACACAGTCTCAGTAGATCATCCCA-3' (corresponds to SEQ ID NO:18)	PCV1 and PCV2 detection
Orf.PCV1	<	5'-CCAACCTTTGTAAACCCCTCCA-3' (corresponds to SEQ ID NO:19)	PCV1 and PCV2-1 detection
Gen.PCV1	>	5'-GTGGACCCACCCTGTGCC-3' (corresponds to SEQ ID NO:20)	PCV1 and PCV1-2 detection
Nested.Orf.PCV1	<	5'-CCAGCTGTGGCTCCATTAA-3' (corresponds to SEQ ID NO:21)	PCV1 and PCV2-1 detection
Nested.Gen.PCV1	>	5'-TTCCCATATAAAATAAATTACTGATGTCCTT-3' (corresponds to SEQ ID NO:22)	PCV1 and PCV1-2 detection
Orf.PCV2	<	5'-CAGTCAGAACGCCCTCCTG-3' (corresponds to SEQ ID NO:23)	PCV2 and PCV1-2 detection
Gen.PCV2	>	5'-CCTAGAAACAAGTGTGGGATG-3' (corresponds to SEQ ID NO:24)	PCV2 and PCV2-1 detection
Nested.Orf.PCV2	<	5'-TTGTAAACAAGGCCACAGC-3' (corresponds to SEQ ID NO:25)	PCV2 and PCV1-2 detection
Nested.Gen.PCV2	>	5'-GTGTGATCGATATCCATTGACTG-3' (corresponds to SEQ ID NO:26)	PCV2 and PCV2-1 detection

^aPrimer direction.

EXAMPLE 10

Evaluation of Infectivity of the PCV1 DNA Clone When
Transfected into PK-15 Cells Free of Virus Contamination

The infectivity of the PCV1 molecular DNA clone was determined by *in vitro* transfection of the PK-15 cells. IFA with PCV1 specific monoclonal antibody (a gift from Dr. Gordon Allan, Belfast, U.K.) confirmed that the PCV1 molecular DNA clone is infectious in PK-15 cells. PCV1-specific antigen was visualized by IFA in the nucleus, and to a lesser degree cytoplasm of the transfected cells. The cells mock-transfected with the empty pSK vector remained negative for PCV1 antigen.

EXAMPLE 11

Construction of a Chimeric PCV1-2 Viral DNA Clone

A chimeric virus was constructed between the nonpathogenic PCV1 and the PMWS-associated PCV2 by using infectious DNA clones of PCV1 and PCV2. To construct a chimeric PCV1-2 DNA clone, the ORF2 capsid gene of the nonpathogenic PCV1 was removed from the PCV1 infectious DNA clone, and replaced with the immunogenic ORF2 capsid gene of the pathogenic PCV2 in the genome backbone of PCV1 (see Figs. 5 and 6). Two pairs of PCR primers were designed. The first primer pair for PCV2 ORF2, Psi I-5 set forth in SEQ ID NO:11 and Acl I-6 set forth in SEQ ID NO:12, was designed with point mutations at the 5' ends of the primers to create restriction enzyme sites *AcII* and *PsiI* to amplify the ORF2 gene of PCV2 and introduce flanking *PsiI* and *AcII* restriction enzyme sites by point mutation. The PCR reaction for the PCV2 ORF2 amplification consisted of an initial step at 95°C for 9 min., followed by 38 cycles of denaturation at 95°C for 1 min., annealing at 48°C for 1 min., extension at 72°C for 1 min., and a final extension at 72°C for 7 min.

A second pair of PCR primers, Hpa I-2 set forth in SEQ ID NO:9 and Nar I-3 set forth in SEQ ID NO:10, was designed for the amplification of the pSK+ vector and its PCV1 genome insert. Point mutations were introduced at the 5' ends of the PCR primers to create flanking restriction enzyme sites *NarI* and *HpaI*. This primer pair amplified the pSK+ vector and its insert PCV1 genomic DNA lacking the ORF2 capsid gene, that is, the PCV1 genome minus the PCV1 ORF2 (pSK-PCV1 Δ ORF2) by using the PCV1 infectious DNA clone as

the PCR template. The PCR reaction consisted of an initial step at 95°C for 9 min., followed by 38 cycles of denaturation at 95°C for 1 min., annealing at 50°C for 1min., extension at 72°C for 3.5 min., and a final extension at 72°C for 7min. The PCV2 ORF2 PCR product was digested with the *AcII* and *PsiI* to remove the introduced point mutations. The pSK-PCV1 Δ ORF2 product (the pSK vector-PCV1 genome PCR product lacking ORF2 gene of PCV1) was digested with the *NarI* and *HpaI* to remove the PCR introduced point mutations. The latter digestion produced a sticky end and a blunt end complementary to the PCV2 ORF2 PCR product digested by the *AcII* and *PsiI* restriction enzymes. The digested PCV2 ORF2 product and the ORF2-deleted pSK-PCV1 product were ligated with T4 DNA ligase to form the chimeric PCV1-2 genomic DNA clone, in which the ORF2 gene of PCV1 is replaced with the ORF2 gene of PCV2. Once the two PCR products were digested and religated, all the PCR introduced point mutations used to facilitate cloning were removed in the resulting chimeric clone. *Escherichia coli* DH5 α competent cells were transformed. The recombinant plasmids containing the chimeric DNA clone were isolated and confirmed by PCR, restriction enzyme digestion and partial DNA sequencing. The full-length chimeric PCV1-2 genome was excised from the pSK+ vector (the recombinant plasmid) with *KpnI* digestion. The chimeric DNA genome was then dimmerized by a short 10-minute ligation reaction with T4 DNA ligase that favors the formation of linear dimers to produce the PCV1-2 chimeric infectious DNA clone (Fig. 6). The recombinant plasmids containing two copies of the chimeric viral genome were confirmed by PCR, restriction enzyme digestion and DNA sequencing.

EXAMPLE 12

Evaluation of *In Vitro* Infectivity of PCV1-2 Chimeric DNA Clone

The viability of the chimeric PCV DNA clone (nonpathogenic PCV1 with the immunogenic capsid gene of PCV2) was tested in PK-15 cells. When PK-15 cells were transfected with the chimeric viral DNA clone, viral antigen specific for PCV2 ORF2 capsid was detected by IFA at about 2 days post-transfection. The PCV1 capsid antigen was not detected in transfected cells. This experiment indicated that the chimeric DNA clone is infectious *in vitro*, is replicating in PK-15 cells and producing the immunogenic capsid protein of PCV2.

EXAMPLE 13

Construction of a Reciprocal Chimeric PCV2-1 DNA Clone

To construct a reciprocal PCV2-1 chimeric DNA clone, the ORF2 capsid gene of PCV2 is replaced by that of the non-pathogenic PCV1 in the genome backbone of the pathogenic PCV2 (Fig. 6). Two PCR primer pairs were designed: the pair, Bgl-II-ORF2 set forth in SEQ ID NO:13 and SpH-I-ORF2 set forth in SEQ ID NO:14, amplifies the PCV1 ORF2 gene and introduces flanking *Bgl*II and *Sp*HI restriction enzyme sites by point mutation. The second PCR primer pair, Bgl-II-PCV2 set forth in SEQ ID NO:15 and SpH-I-PCV2 set forth in SEQ ID NO:16, amplified the pSK vector and the PCV2 genome minus the ORF2 gene (pSK-PCV2 ΔORF2) by using the PCV2 infectious DNA clone as the PCR template, and introduced flanking restriction enzymes sites *Bgl*II and *Sp*HI by point mutation. The pSK-PCV2 ΔORF2 product and the PCV1 ORF2 PCR product were digested by *Bgl*II and *Sp*HI restriction enzymes to produce complementary sticky and blunt ends ligated together. After transformation into E. Coli cells, the authentic recombinant plasmids were isolated and confirmed by enzyme digestion and partial DNA sequencing. The full-length reciprocal chimeric PCV2-1 genome was excised from the recombinant plasmid by *Sac*II digestion, and dimmerized as described herein to produce the reciprocal chimeric PCV2-1 infectious clone.

EXAMPLE 14

In vitro Transfection of PK-15 Cells with PCV1, PCV2, PCV1-2 and PCV2-1 DNA Clones

The infectivity of PCV2 clone *in vitro* and *in vivo* has been demonstrated in the above Examples 3-5. To test the infectivity of the PCV1 and two chimeric clones *in vitro*, PK-15 cells free of PCV1 contamination prepared per the method of Example 1 were grown in 8-well LabTek chambers slides (Nalge Nunc Int., Denmark). When the PK-15 cells reached about 80% confluency, cells were transfected with PCV1, PCV2, PCV1-2 and PCV2-1 DNA clones respectively, using the Lipofectamine Plus Reagent according to the protocols supplied by the manufacturer (Life Technologies, Inc.). Mock-transfected cells with empty pSK vector were included as controls. Three days after transfection, the cells were fixed

with a solution containing 80% acetone and 20% methanol at 4°C for 20 min. Evidence of infectivity and virus replication in cells transfected with the PCV1 and PCV2-1 DNA clones were confirmed by indirect immunofluorescence assay (IFA) using monoclonal antibody against PCV1 ORF2 capsid gene, kindly provided by Dr. G. M. Allan (G. M. Allan *et al.*,
5 "Production, preliminary characterization and applications of monoclonal antibodies to porcine circovirus," Vet. Immunol. Immunopathol. 43:357-371 (1994)). The fixed cells were washed with phosphate buffered saline (PBS) and incubated with 1:20 diluted PCV1 monoclonal antibody at 37°C for 1 hour. The cells were then washed three times with PBS buffer and incubated with fluorescein isothiocyanate (FITC) labeled goat anti-mouse
10 immunoglobulin G (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Md.) at 37°C for 45 min. After washing three times with PBS buffer, the slides were mounted with fluoromount-G, coverslipped, and examined under a fluorescence microscope. The infectivity of cells transfected with the PCV2 and the chimeric PCV1-2 DNA clones were confirmed by IFA using antibody specific for PCV2, as previously described in Example 4.

15 The infectivity of the PCV1, the chimeric PCV1-2 DNA and the reciprocal chimeric PCV2-1 DNA clones were substantiated by the *in vitro* transfection of PK-15 cells. Two identical copies of the complete PCV1 genome were ligated in tandem into the pSK vector to produce the PCV1 DNA clone (Fig. 6). The chimeric PCV1-2 DNA clone had the ORF2 capsid gene of PCV1 replaced by that of the pathogenic PCV2 in the backbone of the
20 nonpathogenic PCV1 genome. The reciprocal chimeric PCV2-1 DNA clone had the ORF2 capsid gene of PCV2 replaced by that of the nonpathogenic PCV1 in the backbone of the pathogenic PCV2 genome. If infectious *in vitro*, the chimeric PCV1-2 DNA clone will produce the ORF2 capsid antigen of PCV2 and the reciprocal chimeric PCV2-1 DNA clone will express PCV1 ORF2 capsid antigen in transfected PK-15 cells. The results showed that
25 the PCV1, the chimeric PCV1-2 and the reciprocal chimeric PCV2-1 DNA clones were all surprisingly shown to be infectious when transfected into PK-15 cells and expressed respective viral capsid antigen proteins as demonstrated by IFA using antibodies specific for PCV1 or PCV2. IFA using monoclonal antibodies against PCV1 ORF2 and antibodies against PCV2 confirmed that the PCV1 DNA and the PCV1-2 DNA clones were infectious.
30 IFA using PCV1 ORF2-specific monoclonal antibody showed that the PCV1-2 chimeric DNA clone was also infectious. About 10-20% of the transfected PK-15 cells were positive

for PCV1 capsid antigen and PCV2 antigen, and expressed PCV1 ORF2 antigen, within the nucleus of transfected cells (Fig. 7).

EXAMPLE 15

Experimental Inoculation of Pigs with PCV1, PCV2, Chimeric PCV1-2 and Reciprocal Chimeric PCV2-1 DNA Clones

To evaluate the immunogenicity and pathogenicity of the chimeric DNA clones, forty specific-pathogen-free (SPF) pigs of 4-6 weeks of age were randomly assigned into five rooms of 8 animals each. Prior to inoculation, animals were tested for antibodies to PCV, PRRSV, PPV, and swine hepatitis E virus. In addition, pre-inoculation serum samples were tested by PCR for PCV1 and PCV2 nucleic acid to confirm that the pigs are not naturally infected by either of the viruses. The PCV1, PCV2, PCV1-2 and PCV2-1 DNA clones were all inoculated by direct injection of the cloned plasmid DNA into the superficial iliac lymph nodes of pigs. Pigs in Group 1 received phosphate buffered saline (PBS buffer) and served as the negative control. Group 2 pigs were each injected into the superficial iliac lymph nodes with 200 µg of infectious PCV1 DNA clone. Group 3 pigs were each injected with 200 µg of infectious PCV2 DNA clone. Group 4 pigs each received injections of 200 µg of infectious chimeric PCV1-2 DNA clone. Group 5 pigs each received 200 µg of the infectious reciprocal chimeric PCV2-1 DNA clone. All animals were monitored daily for clinical signs of disease. Serum samples were collected from each animal at -2, 7, 14, 21, 28, 35, 42 and 49 days post-inoculation (DPI). At 21 DPI, four randomly selected animals from each group were necropsied. The remaining four animals in each group were necropsied at 49 DPI. Various tissues and organs were collected during necropsy as previously described in Example 7, and processed for histological examination.

The immunogenicity of the PCV1, the PCV2 and the chimeric infectious DNA clones was examined in the pigs. Serum samples collected from all control and inoculated animals at -2 (0), 7, 14, 21, 28, 35, 42 and 49 DPIs were assayed for PCV1, PCV2, PCV1-2 and PCV2-1 viremia by PCR detection of clone-specific DNA sequences, for anti-PCV1 antibody by IFA and for anti-PCV2 ORF2 antibody by ELISA. Prior to inoculation at -2 DPI, animals from all five groups tested negative by PCR for both PCV1 and PCV2 DNA.

Negative control animals were negative for both PCV1 and PCV2 viremia throughout the study (see Table 7, below). Five pigs in the uninoculated control group had detectable PCV2 maternal antibody at -2 DPI and 2 pigs had detectable PCV1 maternal antibodies at 7 DPI (see Table 8, below). The maternal antibodies to both PCV1 and PCV2 in these piglets
5 waned by 21 DPI. No seroconversion to either PCV1 or PCV2 was detected in any of the 8 uninoculated control pigs throughout the study.

In the PCV1 inoculated group, viremia was first detected in an inoculated pig at 7 DPI (Table 7, below), and was last detected at 35 DPI. Five out of 8 animals inoculated with PCV1 infectious DNA clone were positive for PCV1 viremia. Average length of continuous
10 PCV1 viremia was 0.625 weeks. By 21 DPI, all animals in the PCV1 inoculated group had seroconverted to PCV1 and remained positive to PCV1 antibodies until the end of the study at 49 DPI.

The PCV2 DNA clone is shown herein to be infectious in pigs. In the PCV2 DNA clone inoculated group, PCV2 viremia was first detected at 7 DPI (Table 7, below). By 21
15 DPI, all PCV2 inoculated Group 3 animals were positive for PCV2 viremia. The average length of PCV2 viremia was 2.12 weeks. Two pigs in the PCV2 inoculated group had detectable levels of maternal PCV2 antibodies at 7 DPI (Table 8, below), and the maternal antibodies in these piglets waned by 14 DPI. Seroconversion to PCV2, assayed by a PCV2-specific ELISA, was first detected at 35 DPI. By 42 DPI, all pigs inoculated with PCV2
20 infectious DNA clone had seroconverted to PCV2.

In Group 4 pigs inoculated with PCV1-2 chimeric infectious DNA clone, viremia specific for the chimeric virus was first detected at 14 DPI (Table 7, below). Four out of 7 inoculated animals became viremic to PCV1-2 between 14 DPI and 42 DPI. The average
length of chimeric PCV1-2 viremia was 1 week. One pig had detectable levels of maternal
25 PCV2 antibodies at 7 and 14 DPI, but the maternal antibody waned by 21 DPI (Table 8, below). Seroconversion to PCV2 ORF2-specific antibody first occurred at 28 DPI. By 49 DPI, all pigs inoculated with chimeric PCV1-2 DNA clone had seroconverted to PCV2 ORF2-specific antibody.

In pigs inoculated with the reciprocal chimeric PCV2-1 clone, viral DNA specific for
30 PCV2-1 chimeric virus was not detected in serum samples (Table 7, below). However, by 21 DPI all animals in Group 5 seroconverted to PCV1 antibody. PCR products amplified from

selected pigs in each group were sequenced and confirmed to be the authentic respective infectious clones used in the inoculation in each group.

Table 7. Detection of Viremia by Nested PCR in Sera of Inoculated and Control Pigs

Group	Inoculum	DPI								Total
		-2	7	14	21	28	35	42	49	
1	PBS ^a	0/8 ^b	0/8	0/8	0/8	0/4	0/4	0/4	0/4	0/8
2	PCV1 DNA ^c	0/8	1/8	1/8	2/8	0/4	2/4	0/4	0/4	5/8
3	PCV2 DNA ^c	0/8	3/8	6/8	7/8	1/4	2/4	2/4	0/4	8/8
4	PCV1-2 DNA ^c	0/7	0/7	1/7	2/7	2/4	2/4	2/4	0/4	4/7
5	PCV2-1 DNA ^c	0/8	0/8	0/8	0/8	0/4	0/4	0/4	0/4	0/8

^aPhosphate buffered saline (PBS) used as negative control

^bEight pigs in each group; number positive/number tested

^cCloned genomic PCV or chimeric PCV DNA in pSK plasmid

Table 8. Seroconversion to Antibodies Against PCV2 in Pigs Inoculated with PCV2 or Chimeric PCV1-2 DNA Clones and Seroconversion to Antibodies Against PCV1 in Pigs Inoculated with PCV1 or PCV2-1 DNA Clones

Group	Inoculum ^a	Antibody Tested For ^b	-2	7	14	21	28	35	42	49
1	PBS	PCV1	NA	2/8	2/8	1/8	0/4	0/4	0/4	0/4
		PCV2	5/8	5/8	2/8	0/8	0/4	0/4	0/4	0/4
2	PCV1 DNA	PCV1 ORF2	NA	3/8	2/8	8/8	4/4	4/4	4/4	4/4
3	PCV2 DNA	PCV2 complete virus	3/8	2/8	0/8	0/8	0/4	3/4	4/4	4/4
4	PCV1-2 DNA	PCV2 complete virus	2/8	1/8	1/8	0/8	1/4	1/4	3/4	4/4
5	PCV2-1 DNA	PCV1 ORF2	NA	3/8	3/8	8/8	3/4	3/4	4/4	4/4

^aPhosphate buffered saline (PBS) used as negative control. The inocula were cloned genomic PCV and PCV chimeric DNA in pSK plasmid

^bPCV1 antibody to ORF2 was measured with an indirect immunofluorescence assay specific for PCV1 antigen. PCV2 antibody was measured with an enzyme-linked immunosorbent assay.

^cDays post inoculation, number positive/number tested.

EXAMPLE 16

Clinical Evaluation

Pigs were weighed on 0 DPI and at the time of necropsies. Rectal temperatures and clinical respiratory scores, ranging from 0 to 6 (0 = normal; 6 = severe) (P. G. Halbur *et al.*,
5 "Comparison of the pathogenicity of two U.S. porcine reproductive and respiratory syndrome virus isolates with that of the Lelystad virus," Vet. Pathol. 32:648-660 (1995)), were recorded every other day from 0 to 49 DPI. Clinical observations, including evidence of central nervous system disease, liver disease (icterus), musculoskeletal disease and changes in body condition, were also recorded daily. A team of two people performed all clinical evaluations.

10 None of the control or inoculated pigs showed obvious signs of the full-spectrum clinical PMWS. There were no differences in weight gain or mean rectal temperatures between any of the groups. One of the pigs from PCV1-2 inoculated Group 3 died one day after inoculation. After necropsy and clinical analysis, no pathogenic agent was detected and death was not associated with the inoculation procedure or the chimeric PCV1-2 virus.

EXAMPLE 17

Gross Pathology and Histopathology

Four pigs from each group were necropsied at 21 and 49 DPI, respectively. The necropsy team was blinded to infection status of the pigs at necropsy. Complete necropsies
20 were performed on all pigs. An estimated percentage of the lung with grossly visible pneumonia was recorded for each pig based on a previously described scoring system (P. G. Halbur *et al.*, 1995, *supra*). Other lesions such as enlargement of lymph nodes were noted separately. Sections for histopathologic examination were taken from nasal turbinate, lungs (seven sections) (*id.*), heart, brain, lymph nodes (tracheobronchial, iliac, mesenteric,
25 subinguinal), tonsil, thymus, liver, gall bladder, spleen, joints, small intestine, colon, pancreas, and kidney. The tissues were examined in a blinded fashion and given a subjective score for severity of lung, lymph node, and liver lesions as described in Example 7. Lung scores ranged from 0 (normal) to 3 (severe lymphohistiocytic interstitial pneumonia). Liver scores ranged from 0 (normal) to 3 (severe lymphohistiocytic hepatitis). Lymph node scores
30 were for an estimated amount of lymphoid depletion of follicles ranging from 0 (normal or

no lymphoid depletion) to 3 (severe lymphoid depletion and histiocytic replacement of follicles).

To determine the pathogenicity of PCV1, PCV2, chimeric PCV1-2 and reciprocal chimeric PCV2-1 infectious DNA clones in pigs, gross lesions were examined first. The results are shown in Table 9 below. Lymph nodes of animals from the uninoculated control Group 1 were normal at both 21 and 49 DPIs. Pigs in the four inoculated groups had variable degrees of gross lesions limited to the lymph nodes. In PCV1 inoculated Group 2 pigs, lymph nodes were grossly normal at 21 DPI, however, mild to moderate swelling and discoloration of lymph nodes was detected at 49 DPI. All PCV2 inoculated Group 3 pigs had enlarged lymph nodes two to five times the normal size, that were firm and tan colored at both 21 and 49 DPIs. Lymph nodes from chimeric PCV1-2 inoculated animals were mild to moderately swollen and discolored at both 21 and 49 DPIs in 5 out of 7 pigs. In Group 5 pigs, inoculated with the PCV2-1 clone, 1 out of 8 animals had mild swelling and discoloration of the lymph nodes at 21 DPI. The average scores of gross lesions of the lymph nodes in pigs inoculated with chimeric PCV1-2 clone were not statistically different from those in Groups 1, 2, and 5, but were statistically different from those of the pathogenic PCV2 inoculated Group 3 pigs at both 21 and 49 DPIs. Average lymphoid gross lesion scores on 49 DPI from the PCV1, PCV2, and PCV1-2 inoculated animals were not statistically different from each other, but were all statistically different from the average gross lesion scores of Groups 1 and 5.

Next, microscopic lesions were examined. The results are shown in Table 10 below. No microscopic lesions were detected in either uninoculated control Group 1 pigs or PCV1 inoculated Group 2 pigs at any DPI. Microscopic lung lesions characterized as mild peribronchiolar lymphoplasmacytic and histiocytic bronchointerstitial pneumonia, were observed in 1 out of 8 of the PCV2 inoculated pigs. In PCV1-2 and PCV2-1 inoculated animals, no microscopic lesions were observed in the lungs. No lesions were observed in the thymuses of any inoculated pigs. Mild multifocal lymphoplasmacytic myocarditis was observed in 2 of 8 pigs in the PCV2 inoculated group. Heart tissues from PCV1-2 and PCV2-1 inoculated animals were free of microscopic lesions. Mild multifocal lymphoplasmacytic interstitial nephritis was observed in 4 out of 8 pigs in PCV2 inoculated group, in 2 out of 7 PCV1-2 inoculated pigs and in 1 out of 8 PCV2-1 inoculated pigs. Mild-to-moderate

lymphoid depletion and histiocytic replacement of follicles were observed in the tonsil in 5 out of 8 pigs, in the spleen in 3 out of 8 pigs, and in the lymph nodes in 8 out of 8 pigs in the PCV2-inoculated group. In the chimeric PCV1-2 inoculated animals, mild lymphoid depletion and histiocytic replacement of follicles were observed in the lymph nodes of 2 out of 7 pigs but were not detected in either the spleen or tonsils. No lymphoid depletion and histiocytic replacement of follicles were observed in the lymph nodes, spleen or tonsils of the reciprocal chimeric PCV2-1 inoculated animals. Mild-to-moderate lymphoplasmacytic hepatitis was observed in 7 out of the 8 PCV2-inoculated pigs. Mild lymphoplasmacytic hepatitis was observed in 2 out of the 7 chimeric PCV1-2 inoculated pigs. No lymphoplasmacytic hepatitis was observed in reciprocal chimeric PCV2-1 inoculated pigs. Lesions in other tissues were unremarkable.

Microscopic lesions in the lung, liver, and lymph nodes were scored according to a published scoring system (P. G. Halbur *et al.*, 1995, *supra*). The results are shown in Table 10 below. Average scores of lesions in lymph nodes in pigs from the chimeric PCV1-2 inoculated Group 4 were similar to those from Groups 1, 2 and 5 but were statistically different from those of the pathogenic PCV2 inoculated Group 3 pigs, at both 21 and 49 DPIs. Average microscopic liver lesion scores from the chimeric PCV1-2 inoculated group at 21 DPI were statistically different from those of PCV2 inoculated Group 3 animals but were similar to those of Groups 1, 2 and 5 pigs at 21 DPI. At 49 DPI, the average microscopic liver scores from Group 4 chimeric PCV1-2 inoculated pigs were not statistically different from those of Groups 1, 2, 3 and 5 pigs. There were no acceptable scoring systems for other tissues or organs.

Table 9. Gross Lesions of Lymph Nodes in Control and Inoculated Pigs

Group	Inoculum ^a	DPI ^b	
		21	49
1	PBS	0/4(0.0)	0/4(0.0)
2	PCV1 DNA	0/4(0.0)	4/4(1.5)
3	PCV2 DNA	4/4(2.5)	4/4(2.25)
4	PCV1-2 DNA	2/3(0.66)	3/4(1.25)
5	PCV2-1 DNA	1/4(0.25)	0/4(0.0)

^aPhosphate buffered saline (PBS) used as negative control. The inocula were cloned genomic PCV or chimeric PCV DNA in pSK plasmid.

^bFour pigs from each group were necropsied at 21 DPI and the remaining pigs were necropsied at 49 DPI; number positive/number tested. Number with lesions/number tested (range of estimated severity lymph node enlargement)

Table 10. Distribution of Histopathological Lesions in Different Tissues/Organs from Control and Inoculated Pigs

Group	Inoculum ^a	DPI ^b	Lung ^c	Liver ^d	Lymph Nodes ^e	Spleen	Thymus	Ileum	Brain	Heart	Kidney	Tonsil
1	PBS	21 49	0/4(0.0) 0/4(0.0)	0/4(0.0) 0/4(0.0)	0/4(0.0) 0/4(0.0)	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4
2	PCV1 DNA	21 49	0/4(0.0) 0/4(0.0)	0/4(0.0) 0/4(0.0)	0/4(0.0) 0/4(0.0)	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4
3	PCV2 DNA	21 49	0/4(0.0) 1/4(0.25)	0/4(0.0) 3/4(0.75)	0/4(0.0) 4/4(1.0)	0/4 0/4	0/4 0/4	0/4 0/4	1/4 1/4	1/4 1/4	2/4 2/4	3/4 2/4
4	PCV1-2 DNA	21 49	0/3(0.0) 0/4(0.0)	1/3(0.33) 1/4(0.25)	1/3(0.33) 1/4(0.25)	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	2/4 2/4	0/4 0/4
5	PCV2-1 DNA	21 49	0/4(0.0) 0/4(0.0)	0/4(0.0) 0/4(0.0)	0/4(0.0) 1/4(0.25)	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4	0/4 0/4

^aPhosphate buffered saline (PBS) used as negative control. The inocula were cloned genomic PCV or chimeric PCV DNA in pSK plasmid.

^bFour pigs from each group were necropsied at 21 DPI and the remaining pigs were necropsied at 49 DPI.

^cNumber positive/number tested (average histological lung score: 0, normal; 1, mild interstitial pneumonia; 2, moderate; 3, severe).

^dNumber positive/number tested (Average histological liver score: 0, normal; 1, mild hepatitis; 2, moderate; 3, severe.)

^eNumber positive/number tested (Average histological lymphoid (lymph nodes) depletion score: 0, normal; 1, mild; 2, moderate; 3, severe.)

EXAMPLE 18

Serology

Blood was collected from all pigs at -2, 7, 14, 21, 28, 35, 42 and 49 DPIs. Serum antibodies to PRRSV were assayed using Herd Check PRRSV ELISA (IDEXX Laboratories, Westbrook, MA). Serum antibodies to PPV were detected by a hemagglutination inhibition (HI) assay (H. S. Joo *et al.*, "A standardized haemagglutination inhibition test for porcine parvovirus antibody," Aust. Vet. J. 52:422-424 (1976)). Serum antibodies to PCV2 were detected by a modified indirect ELISA based on the recombinant ORF2 capsid protein of PCV2 as described hereinabove (see also P. Nawagitgul *et al.*, "Modified indirect porcine circovirus (PCV) type 2-based and recombinant capsid protein (ORF2)-based ELISA for the detection of antibodies to PCV," Immunol. Clin. Diagn. Lab Immunol. 1:33-40 (2002)). Serum antibodies to PCV1 were detected by an indirect immunofluorescence assay (IFA). PK-15 cells infected with PCV1 were grown on eight-well LabTek chamber slides. When the infected PK-15 cells reach about 95-100% confluency, the infected cells were fixed with a solution containing 80% acetone and 20% methanol at 4°C for 20 min. The fixed cells were washed once with PBS buffer. One hundred microliters of 1:10 diluted pig serum sample in PBS was added to the chambers, and incubated for 1 hour at 37°C. The cells were then washed three times with PBS and incubated for 45 min. at 37°C with FITC-labeled goat anti-swine secondary antibody. The slides were subsequently washed three times with PBS, mounted with fluoromount-G, coverslipped and examined under a fluorescent microscope. For the positive control, PCV1 infected cells were incubated with a diluted PCV1 specific monoclonal antibody (gift of Dr. G. M. Allan), followed by an incubation with FITC-labeled goat anti-mouse IgG (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Md.). For the negative control, PCV1 infected cells were incubated with 1:10 diluted swine serum free of PCV1 and PCV2 antibody, followed by an incubation with FITC-labeled goat anti-swine IgG (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, Md.).

EXAMPLE 19

PCR Detection

To detect PCV1, PCV2, chimeric PCV1-2 and reciprocal chimeric PCV2-1 viremia in sera from inoculated pigs, serum samples collected at different DPIs were tested by PCR.

Viral DNA was extracted from 100 µl of each serum sample using DNazol reagent according to the manufacturer's protocol (Molecular Research Center, Cincinnati, OH). The extracted DNA was resuspended in DNase, RNase and proteinase-free water. To amplify clone-specific genomic sequences of PCV1, PCV2, chimeric PCV1-2 and chimeric reciprocal PCV2-1, two sets of nested PCR primer pairs were designed (Table 6, above). The first set of nested primers was designed based on published PCV1 sequences. Primers Gen.PCV1 set forth in SEQ ID NO:20 and Orf.PCV1 set forth in SEQ ID NO:19 amplified a 400 bp fragment in the presence of the PCV1 genome. The nested primers, nested.Gen.PCV1 set forth in SEQ ID NO:22 and nested.Orf.PCV1 set forth in SEQ ID NO:21, amplified a 220 bp fragment.

To detect PCV2 viremia, PCV2 primer pair Gen.PCV2 set forth in SEQ ID NO:24 and Orf.PCV2 set forth in SEQ ID NO:23 amplified a 900 bp fragment in the presence of PCV2 in the first round of PCR. Primers nested.Gen.PCV2 set forth in SEQ ID NO:26 and nested.Orf.PCV2 set forth in SEQ ID NO:25 amplified a 600 bp fragment in the nested PCR.

To detect chimeric PCV1-2 viremia, the first round of PCR reaction employed the PCV1-specific primer Gen.PCV1 set forth in SEQ ID NO:20 and the PCV2 ORF2-specific primer Orf.PCV2 set forth in SEQ ID NO:23 to amplify a chimeric fragment of 580 bp. For the nested PCR, PCV1-specific primer nested.Gen.PCV1 set forth in SEQ ID NO:22 and the PCV2 ORF2-specific primer nested.Orf.PCV2 set forth in SEQ ID NO:25 were used to amplify a chimeric fragment of 370 bp.

To detect reciprocal chimeric PCV2-1 viremia, the first round of PCR employed the PCV2-specific primer Gen.PCV2 set forth in SEQ ID NO:24 and the PCV1 ORF2-specific primer Orf.PCV1 set forth in SEQ ID NO:19 to amplify a chimeric fragment of 700 bp. For the nested PCR, the PCV2-specific primer nested.Gen.PCV2 set forth in SEQ ID NO:26 and the PCV1 ORF2-specific primer nested.Orf.PCV1 set forth in SEQ ID NO:21 were used to amplify a 460 bp chimeric fragment. All PCR parameters were essentially the same, consisting of 38 cycles of denaturation at 94°C for 1 min., annealing at 45°C for 1 min., and extension at 72°C for 1.5 min. The serum samples from negative control pigs were tested by a PCR-RFLP diagnostic assay, which can detect and differentiate both PCV1 and PCV2 as described previously (M. Fenaux *et al.*, "Genetic characterization of type 2 porcine circovirus (PCV-2) from pigs with postweaning multisystemic wasting syndrome in different

geographic regions of North America and development of a differential PCR-restriction fragment length polymorphism assay to detect and differentiate between infections with PCV-1 and PCV-2," J. Clin. Microbiol. **38**: 2494-503 (2000)). PCR products from selected animals in each group were sequenced to verify the origin of the virus infecting pigs.

5

EXAMPLE 20

Immunohistochemistry (IHC)

IHC detection of PCV2-specific antigen was performed on lymph node tissues collected from all pigs necropsied at 21 and 49 DPIs. A rabbit polyclonal antiserum against
10 PCV2 was used for the IHC, according to the general procedures described previously (S. D. Sorden *et al.*, "Development of a polyclonal-antibody-based immunohistochemical method for the detection of type 2 porcine circovirus in formalin-fixed, paraffin-embedded tissue," J. Vet. Diagn. Invest. **11**:528-530 (1999)).

Based on the IHC staining of PCV2-specific antigen, lymphoid tissues from the
15 uninoculated control, PCV1 and PCV2-1 inoculated pigs were negative for PCV2 antigen. PCV2 antigen was detected in lymphoid tissues of 7 out of 8 animals in the PCV2 inoculated group. PCV2 antigen was also detected in lymphoid tissue of 1 out of 7 pigs from the chimeric PCV1-2 inoculated group.

20 In the foregoing, there has been provided a detailed description of particular embodiments of the present invention for the purpose of illustration and not limitation. It is to be understood that all other modifications, ramifications and equivalents obvious to those having skill in the art based on this disclosure are intended to be included within the scope of the invention as claimed.

25

Applicant's or agent's file reference	AM100878	International Publication No.
--	----------	-------------------------------

(Sheet 1/3)

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 30, line 22-27 (chimeric PCV1-2 DNA)	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet ☒	
Name of depositary institution American Type Culture Collection (ATCC)	
Address of depositary institution (including postal code and country) 10801 University Blvd., Manassas, VA 20110-2209 US	
Date of deposit 07 December 2001 (07/12/01)	Accession Number PTA-3912
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet ☐	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only ☒ This sheet was received with the international application Authorized officer S. HOSTAD	For International Bureau use only ☐ This sheet was received by the International Bureau on: Authorized officer
--	--

Applicant's or agent's file reference AM100878	International Application No.
--	-------------------------------

(Sheet 2/3)

INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 30, line 22-27 (PCV2 DNA clone)	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet ☒	
Name of depositary institution American Type Culture Collection (ATCC)	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 07 December 2001 (07/12/01)	Accession Number PTA-3913
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet ☐	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

☒ For receiving Office use only This sheet was received with the international application	☐ For International Bureau use only This sheet was received by the International Bureau on:
Authorized officer S. HOSTAD	Authorized officer

Applicant's or agent's file reference AM100878	International Application No.
--	-------------------------------

(Sheet 3/3)

**INDICATIONS RELATING TO DEPOSITED MICROORGANISM
OR OTHER BIOLOGICAL MATERIAL**

(PCT Rule 13bis)

A. The indications made below relate to the deposited microorganism or other biological material referred to in the description on page 30 line 29-31 (PCV2 sample #40895)	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet ☒	
Name of depositary institution American Type Culture Collection (ATCC)	
Address of depositary institution (including postal code and country) 10801 University Blvd. Manassas, VA 20110-2209 US	
Date of deposit 07 December 2001 (07/12/01)	Accession Number PTA-3914
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet ☐	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
☒ This sheet was received with the international application	☐ This sheet was received by the International Bureau on:
Authorized officer S. HOSTAD	Authorized officer

CLAIMS

What is claimed is:

5

1. An infectious chimeric nucleic acid molecule of porcine circovirus (PCV1-2) characterized by having a nucleic acid molecule encoding an infectious, nonpathogenic PCV1 which contains an immunogenic open reading frame (ORF) gene of a pathogenic PCV2 in place of an ORF gene of the PCV1 nucleic acid molecule.

10

2. The chimeric nucleic acid molecule according to Claim 1, characterized in that the immunogenic PCV2 ORF gene replaces the same ORF gene position in the PCV1 nucleic acid molecule.

3. The chimeric nucleic acid molecule according to Claim 2, characterized in that the immunogenic ORF gene is the ORF2 capsid gene.

15

4. The chimeric nucleic acid molecule according to Claim 3, characterized in that the chimeric nucleic acid molecule has a nucleotide sequence set forth in SEQ ID NO:2, its complementary strand or a nucleotide sequence which has at least 95% homology to the nucleotide sequence of SEQ ID NO:2.

20

5. A biologically functional plasmid or viral vector characterized by containing the chimeric nucleic acid molecule according to Claim 4.

6. The plasmid according to Claim 5 characterized by having ATCC Patent Deposit Designation PTA-3912.

7. A suitable host cell transfected by a vector characterized by comprising the chimeric nucleic acid molecule according to Claim 4.

25

8. An avirulent, infectious chimeric porcine circovirus characterized by being produced by cells containing the chimeric nucleic acid molecule according to Claim 4.

9. The infectious chimeric porcine circovirus according to Claim 8, characterized in that said cells containing the chimeric nucleic acid molecule are contained in or derived from a plasmid having ATCC Patent Deposit Designation PTA-3912.

30

10. A process for the production of an immunogenic polypeptide product, said process characterized by comprising: growing, under suitable nutrient conditions, prokaryotic or

eucaryotic host cells transfected with a nucleic acid molecule according to Claim 4 in a manner allowing expression of said polypeptide product, and isolating the desired polypeptide product of the expression of said nucleic acid molecule.

11. An immunogenic polypeptide product of the expression according to Claim 10.

5 12. An immunogenic polypeptide characterized by having the amino acid sequence set forth in SEQ ID NO:4 or a biologically active variant thereof.

13. A viral vaccine that protects a pig against viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2 characterized by comprising a nontoxic, physiologically acceptable carrier and an immunogenic amount of a member selected from
10 the group consisting of:

(a) a chimeric nucleic acid molecule having a nucleotide sequence set forth in SEQ ID NO:2, its complementary strand or a nucleotide sequence having at least 95% homology to the nucleotide sequence of SEQ ID NO:2;

(b) a biologically functional plasmid or viral vector containing the chimeric nucleic
15 acid molecule, the complementary strand or the nucleotide sequence having at least 95% homology to the nucleotide sequence of SEQ ID NO:2;

(c) an avirulent, infectious chimeric porcine circovirus;

(d) a polypeptide having the amino acid sequence set forth in SEQ ID NO:4 or a biologically active variant thereof;

20 (e) an infectious PCV2 molecular DNA clone having ATCC Patent Deposit Designation PTA-3913 or a PCV2 DNA clone derived therefrom; and


(f) a biologically functional plasmid or viral vector containing the infectious PCV2 molecular DNA clone having ATCC Patent Deposit Designation PTA-3913 or the PCV2 DNA clone derived therefrom.

25 14. The viral vaccine according to Claim 13, characterized in that said vaccine contains live chimeric porcine circovirus.

15. A method of protecting a pig against viral infection or postweaning multisystemic wasting syndrome (PMWS) caused by PCV2 characterized by comprising administering to the pig in need of protection an immunologically effective amount of the vaccine according to
30 Claim 13.

16. The method according to Claim 15, characterized by comprising administering the chimeric nucleic acid molecule or live chimeric porcine circovirus to the pig.
17. The method according to Claim 16, characterized by comprising administering the vaccine parenterally, intranasally, intradermally or transdermally to the pig.
- 5 18. The method according to Claim 17, characterized by comprising administering the vaccine intralymphoidly or intramuscularly to the pig.
19. A method of preparing the infectious chimeric nucleic acid molecule of PCV1-2 according to Claim 1, characterized by comprising removing an open reading frame (ORF) gene of a nucleic acid molecule encoding an infectious, nonpathogenic PCV1; replacing the
- 10 ORF gene position of the PCV1 with an immunogenic ORF gene from a pathogenic PCV2; and recovering the chimeric nucleic acid molecule.
20. The method according to Claim 19, characterized in that said immunogenic PCV2 ORF gene replaces the same ORF gene position of the PCV1 nucleic acid molecule.
21. The method according to Claim 20, characterized in that said immunogenic ORF gene
- 15 is ORF2.
22. The method according to Claim 21, characterized in that said ORF2 gene of PCV2 is obtained from the molecular nucleic acid molecule of PCV2 contained in an expression vector having ATCC Patent Deposit Designation PTA-3913.
23. An infectious PCV2 molecular DNA clone characterized by having ATCC Patent
- 20 Deposit Designation PTA-3913 or a PCV2 DNA clone derived therefrom.
24. A biologically functional plasmid or viral vector characterized by containing the infectious PCV2 molecular DNA clone according to Claim 23.
25. An infectious reciprocal chimeric nucleic acid molecule of PCV2-1 characterized by comprising a nucleic acid molecule encoding an infectious, pathogenic PCV2 which has an
- 25 immunogenic ORF2 gene from a nonpathogenic PCV1 in place of an ORF2 gene of the PCV2 nucleic acid molecule.

Fig. 1






Fig. 2A





Fig. 2B


Figs. 3A - 3D




3A



3B



3C



3D

Figs. 4A - 4D







Fig. 5





Fig. 6



Figs. 7A - 7J




Fig. 8

AAATTTCTGACAAACGTTACAGGGTGCTGCTCTGCAACGGTCACCAGACTCCCGCTCTCC
AACAAAGGTACTCACAGCAGTAGACAGGTCCTCCGTTGTCCTTGAGATCGAGGAGCTCCA
CATTCATAAGTAAGTTGCCTTCTTTACTGCAATATTCTTTATTCTGCTGATCAGTTCCCT
TTGGCTTTCTCGATATGGCAGCGGGCACCCAAATACCACTTCACTTTATTAAAAGTTTGC
TTCTTCACAAAATTAGCGAACCCCTGGAGGTGAGGTGTTTCGTCTCTCCTCATTACCCTCC
TCGCCAACATAAAATAATCAAATAGGGAGATTGGGAGCTCCCGTATTTTCTTGCGCTCG
TCTTCGGAAGGATTATTACAGCGTGAACACCCACCTTTTATGTGGTTGGGGTCCGCTTCTT
CCATTCTTCTTGCTGGGCATGTTGCTGCTGAGGTGCTGCCGAGGTGCTGCCGCTGCCGAA
GTGCGCTGGTAATACTTACAGCGCACTTCTTTCTGTTTTTCAGCTATGACGTATCCAAGGAG
GCGTTACCGCAGAAGAAGACACCGCCCCCGCAGCCATCTTGGCCAGATCCTCCGCCGCCG
CCCCTGGCTCGTCCACCCCCGCCACCGCTACCGTTGGAGAAGGAAAAATGGCATCTTCAA
CACCCGCCTCTCCCGCACCTTCGGATATACTGTCAAGGCTACCACAGTCAGAACGCCCTC
CTGGGCGGTGGACATGATGAGATTTAATATTGACGACTTTGTTCCCCCGGGAGGGGGGAC
CAACAAAATCTCTATAACCCTTTGAATACTACAGAATAAGAAAGGTTAAGGTTGAATTCTG
GCCCTGCTCCCCCATCACCCAGGGTGATAGGGGAGTGGGCTCCACTGCTGTTATTCTAGA
TGATAACTTTGTAAACAAAGGCCACAGCCCTAACCTATGACCCATATGTAACTACTCCTC
CCGCCATACAATCCCCCAACCTTCTCCTACCACTCCCGTTACTTCACACCCAAACCTGT
TCTTGACTCCACCATTGATTACTTCCAACCAAATAACAAAAGGAATCAGCTTTGGATGAG
GCTACAAACCTCTAGAAATGTGGACCACGTAGGCCTCGGCACTGCGTTCGAAAACAGTAT
ATACGACCAGGACTACAATATCCGTGTAACCATGTATGTACAATTCAGAGAATTTAATCT
TAAAGACCCCCCACTTAAACCTAAATGAATAATAAAAAACCATTACGAAGTGATAAAAA
GACTCAGTAATTTATTTTCATATGGAAATTCAGGGCATGGGGGGGAAAGGGTGACGAACTG
GCCCCCTTCTCCTCGTGGATTGTTCTGTAGCATTCTTCCAAAATACCAAGAAAGTAATCCT
CCGATAGAGAGCTTCTACAGCTGGGACAGCAGTTGAGGAGTACCATTCCAACGGGGTCTG
ATTGCTGGTAATCAGAATACTGCGGGCCAAAAAAGGTACAGTTCCACCTTTAGTCTCTAC
AGTCAATGGATATCGATCACACAGTCTCAGTAGATCATCCACGGCAGCCAGCCATAAAA
GTCATCAATAACAACCACTTCTTCACCATGGTAACCATCCCACCACTTGTTTCTAGGTGG
TTTCCAGTATGTGGTTTCCGGGTCTGCAAAATTAGCAGCCCATTTGCTTTTACCACACCC
AGGTGGCCCCACAATGACGTGTACATTGGTCTTCCAATCACGCTTCTGCATTTTCCCGCT
CACTTTCAAAAGTTCAGCCAGCCCCGCG

Fig. 9

GGTACCTCCGTGGATTGTTCTCCAGCAGTCTTCCAAAATTGCAAAGTAGTAATCCTCCGA
TAGAGAGCTTCTACAGCTGGGACAGCAGTTGAGGAGTACCATTCTGGGGGGCCTGATTG
CTGGTAATCAAAATACTGCGGGCCAAAAAAGGAACAGTACCCCTTTAGTCTCTACAGTC
AATGGATACCGGTCACACAGTCTCAGTAGATCATCCCAAGGTAACCAGCCATAAAAATCA
TCCAAAACAACAACCTTCTTCTCCATGATATCCATCCCACCACTTATTTCTACTAGGCTTC
CAGTAGGTGTCCCTAGGCTCAGCAAAATTACGGGGCCCACTGGCTCTTCCCACAACCGGGC
GGGCCCCTATGACGTGTACAGCTGTCTTCCAATCACGCTGCTGCATCTTCCCGCTCACT
TTCAAAAAGTTCAGCCAGCCCCGCGGAAATTTCTCACATACGTTACAGGAACTGCTCGGCT
ACAGTCACCAAAGACCCCGTCTCCAAAAGGGTACTCACAGCAGTAGACAGGTGCGTGC GC
TTCCCCTGGTTCCGCGGAGCTCCACACTCGATAAGTATGTGGCCTTCTTTACTGCAGTAT
TCTTTATTTCTGCTGGTCGGTTCCCTTTCGCTTTCTCGATGTGGCAGCGGGCACAAAATAC
CACTTCACCTTGTTAAAAGTCTGCTTCTTAGCAAAATTTCGCAAAACCCCTGGAGGTGAGGA
GTTCTACCCTCTTCCAAACCTTCTCGCCACAAACAAAATAATCAAAAAGGGAGATTGGA
AGCTCCCGTATTTTGTTTTTCTCCTCCTCGGAAGGATTATTAAGGGTGAACACCCACCTC
TTATGGGGTTGCGGGCCGCTTTTCTTGCTTGGCATTTCCTCACTTTTATAGGATG
ACGTATCCAAGGAGGCGTTACCGCAGAAGAAGACACCGCCCCCGCAGCCATCTTGGCCAG
ATCCTCCGCCCGCCGCCCTGGCTCGTCCACCCCCGCCACCGCTACCGTTGGAGAAGGAAA
AATGGCATCTTCAACACCCGCCTCTCCCGCACCTTCGGATATACTGTCAAGGCTACCACA
GTCAGAACGCCCTCCTGGGCGGTGGACATGATGAGATTTAATATTGACGACTTTGTTCCC
CCGGGAGGGGGGACCAACAAAATCTCTATACCCTTTGAATACTACAGAATAAGAAAGGTT
AAGGTTGAATTCTGGCCCTGCTCCCCCATCACCCAGGGTGATAGGGGAGTGGGCTCCACT
GCTGTTATTCTAGATGATAACTTTGTAACAAAGGCCACAGCCCTAACCTATGACCCATAT
GTAAACTACTCCTCCCGCCATACAATCCCCCAACCCTTCTCCTACCACTCCCGTTACTTC
ACACCCAAACCTGTTCTTGACTCCACCATTGATTACTTCCAACCAAATAACAAAAGGAAT
CAGCTTTGGATGAGGCTACAAACCTCTAGAAATGTGGACCACGTAGGCCTCGGCACTGCG
TTCGAAAACAGTATATACGACCAGGACTACAATATCCGTGTAACCATGTATGTACAATTC
AGAGAAATTTAATCTTAAAGACCCCCCACTTAAACCCTAAATGAATAAAAATAAAAACCAT
TACGATGTGATAACAAAAAAGACTCAGTAATTTATTTTATATGGGAAAAGGGCACAGGGT
GGGTCCACTGCTTCAAATCGGCCTTCGGGTACC

Fig. 10

ATGACGTATCCAAGGAGGCGTTACCGCAGAAGAAGACACCGCCCCCGCAGCCATCTTGGC
CAGATCCTCCGCCGCCGCCCTGGCTCGTCCACCCCGCCACCGCTACCGTTGGAGAAGG
AAAAATGGCATCTTCAACACCCGCCTCTCCCGCACCTTCGGATATACTGTCAAGGCTACC
ACAGTCAGAACGCCCTCCTGGGCGGTGGACATGATGAGATTTAATATTGACGACTTTGTT
CCCCCGGGAGGGGGGACCAACAAAATCTCTATACCCTTTGAATACTACAGAATAAGAAAG
GTTAAGGTTGAATTCTGGCCCTGCTCCCCCATCACCAGGGTGATAGGGGAGTGGGCTCC
ACTGCTGTTATTCTAGATGATAACTTTGTAACAAAGGCCACAGCCCTAACCTATGACCCA
TATGTAACTACTCCTCCCGCCATACAATCCCCCAACCCTTCTCCTACCACTCCCGTTAC
TTCACACCCAAACCTGTTCTTGACTCCACCATTGATTACTTCCAACCAAATAACAAAAGG
AATCAGCTTTGGATGAGGCTACAAACCTCTAGAAATGTGGACCACGTAGGCCTCGGCACT
GCGTTCGAAAACAGTATATACGACCAGGACTACAATATCCGTGTAACCATGTATGTACAA
TTCAGAGAATTTAATCTTAAAGACCCCCCACTTAAACCCTAA

Fig. 11

MTYPRRRYRRRRHRPRSHLGQILRRRPWLVHPRHRYRWRRKNGIFNTRLSTFGYTVKAT
TVRTPSWAVDMMRFNIDDFVPPGGGTNKISIPFEYYRIRKVKVEFWPCSPITQGDRGVGS
TAVILDDNFVTKATALTYDPYVNYSSRHTIPQPFSYHSRYFTP KPVL DSTIDYFQPNNKR
NQLWMRLQTSRNV DHVGLGTAFENSIYDQDYNIRVTMYVQFREFNLKDPPLKP*

*translation termination codon