

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 August 2006 (24.08.2006)

PCT

(10) International Publication Number
WO 2006/088541 A2

(51) International Patent Classification:
A61K 31/551 (2006.01) **C07D 403/02** (2006.01)

(21) International Application Number:
PCT/US2005/044978

(22) International Filing Date:
12 December 2005 (12.12.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/654,207 17 February 2005 (17.02.2005) US

(71) Applicant (for all designated States except US): **STATE OF OREGON ACTING BY & THROUGH THE STATE BOARD OF HIGHER EDU. ON BEHALF OF PORTLAND STATE UNV.** [US/US]; P.o. Box 751, Portland, Oregon 97207 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **PEYTON, David, H.** [US/US]; 2410 Sw 6th Avenue, Portland, Oregon 97201 (US). **BURGESS, Steven** [US/US]; 19580 Sw 56th Court, Tualatin, Oregon 97062 (US).

(74) Agent: **YOUNG, Travis; KLARQUIST SPARKMAN, LLP**, ONE WORLD TRADE CENTER, SUITE 1600, 121 Sw Salmon Street, Portland, Oregon 97204 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: QUINOLINE DERIVATIVES AND USES THEREOF

(57) Abstract: This disclosure provides a new class of compounds referred to as "reversed chloroquinines" (RCQs), which are highly effective against CQ^R and CQ^S malaria parasites. RCQs are hybrid molecules, which include an antimalarial quinoline analog (such as chloroquine) moiety and a CQ^R reversal moiety. Exemplary RCQ chemical structures are provided. Also provided are pharmaceutical compositions including the disclosed RCQ compounds, and methods of using such compounds and compositions for the treatment of malaria and inhibition of CQ^R or CQ^S *Plasmodium* sp. (such as *P. falciparum*).



WO 2006/088541 A2

- 1 -

Quinoline Derivatives and Uses Thereof

CROSS REFERENCE TO RELATED APPLICATIONS

This claims the benefit of U.S. Provisional Application No. 60/654,207, filed
5 February 17, 2005, which is incorporated herein in its entirety.

STATEMENT OF GOVERNMENT SUPPORT

This invention was made with United States government support pursuant to
grant no. AI051509-01A2 from the National Institutes of Health; the United States
10 government has certain rights in the invention.

FIELD OF THE DISCLOSURE

This disclosure relates to quinoline derivatives (such as chloroquine
derivatives) useful in the treatment of chloroquine-resistant (CQ^R) and
15 chloroquine-sensitive (CQ^S) malaria parasites.

BACKGROUND

Malaria is a disease caused by various species of hemosporidian blood
parasites of the genus *Plasmodium* (including *P. falciparum*, *P. vivax*, *P. ovale*, and
20 *P. malariae*). The human disease process begins with the bite of an infected female
Anopheline mosquito. *Plasmodium* sp. sporozoites released from the salivary glands
of the mosquito enter the bloodstream and quickly (within about 30 minutes) invade
liver cells (hepatocytes). The liver-stage parasites differentiate and undergo asexual
multiplication resulting in tens of thousands of merozoites which burst from the
25 hepatocyte. Merozoites then invade red blood cells (erythrocytes) where they
undergo an additional round of multiplication. The clinical signs of malaria, fever
and chills, are associated with the synchronous rupture of infected erythrocytes.

Malaria is a worldwide public health problem. Up to 3 million people die of
malaria each year, and the total number of people infected with malaria worldwide
30 approaches half a billion. The impact of this disease, in terms of suffering and
economics, is huge. In the past, the inexpensive and orally administered antimalarial
drug, chloroquine (CQ), was the "gold standard" treatment.

- 2 -

During its blood stage, the malaria parasite metabolizes (in its digestive vacuole (DV)) hemoglobin present in erythrocytes. A by-product of this metabolic process is heme, which would be toxic to the parasite except for heme detoxification mechanisms developed by the organism. CQ is believed to inhibit heme
5 detoxification in the DV by binding to heme and/or to hemozoin (Ginsburg *et al.*, *Parasitol. Today*, 15:357, 1999; Chong and Sullivan, *Biochem. Pharmacol.*, 66(11):2201-2212, 2003; Leed *et al.*, *Biochem.*, 41(32):10245-10255, 2002; Egan *et al.*, *Biochem.*, 40(1):204-213, 2001; Raynes, *Int. J. Parasitol.*, 29(3):367-379, 1999; Egan *et al.*, *J. Med. Chem.*, 43(2):283-291, 2000). This binding, in part, is thought
10 to drive the thermodynamics that concentrates CQ in the parasite DV. Chloroquine-dependent heme accumulation in the DV may inhibit or kill the parasites.

Unfortunately, certain *Plasmodium sp.* strains have evolved resistance to CQ. In fact, the spread of chloroquine-resistant (CQ^R) *Plasmodium sp.* parasites has
15 rendered CQ almost useless for malaria treatment. In addition, *Plasmodium sp.* resistance to other antimalarial drugs, such as artemisinin and its derivatives, has been reported (Xiao *et al.*, *Parasitol. Res.*, 92(3):215-219, 2004). These are particularly devastating problems in many impoverished parts of the world where such drugs are most needed (Krogstad, *Epidemiol. Rev.*, 18(1):77-89, 1996).

20 *Plasmodium sp.* CQ^R is correlated with mutations in the parasite's DV membrane transporter protein (PfCRT). PfCRT is thought to enhance CQ export from the DV (Zhang *et al.*, *Biochem.*, 43(26):8290-8296, 2004; Bennett *et al.*, *Mol. Biochem. Parasitol.*, 133(1):99-114, 2004; Martin and Kirk, *Mol. Biol. Evol.*, 21(10):1938-1949, 2004). A particularly well-studied PfCRT mutation is K76T,
25 which is correlated with CQ^R (Martin and Kirk, *Mol. Biol. Evol.*, 21(10):1938-1949, 2004; Johnson *et al.*, *Mol. Cell*, 15(6):867-877, 2004; Durand *et al.*, *Mol. Biochem. Parasitol.*, 136(2):273-285, 2004; Ranjit *et al.*, *Trop. Med. Int. Health*, 9(8):857-861, 2004; Durand *et al.*, *Antimicrob. Agents Chemother.*, 46(8):2684-2686, 2002; Durand *et al.*, *Mol. Biochem. Parasitol.*, 114(1):95-102, 2001; Cooper *et al.*, *Mol.*
30 *Pharmacol.*, 61(1):35-42, 2002; Djimde *et al.*, *N. Engl. J. Med.*, 344(4):257-263, 2001).

- 3 -

Alternative therapies for treatment of chloroquine-resistant (CQ^R) malaria parasites have been developed, including combination therapies (Kumar *et al.*, *Curr. Med. Chem.*, 10(13):1137-1150, 2003). However, none of these therapies meet CQ's simplicity of use and low cost.

5 One approach to treatment of CQ^R parasites involves the use of chemicals known as "reversal agents." Reversal agents are chemicals that have been found to quash CQ resistance (Krogstad *et al.*, *Science*, 238(4831):1283-1285, 1987; Martin *et al.*, *Science*, 235(4791):899-901, 1987; Ryall, *Parasitol. Today*, 3(8):256, 1987); thus, making CQ^R strains sensitive again to CQ. CQ^R reversal agents do not appear
10 to have independent therapeutic value against *Plasmodium sp.* because the therapeutically useful dose for CQ applied to a CQ^S strain in the presence of a reversal agent is nearly the same as in the absence of the reversal agent.

 At least some reversal agents are believed to act by inhibiting CQ export from the DV (Krogstad *et al.*, *Science*, 238(4831):1283-1285, 1987; Martin *et al.*,
15 *Science*, 235(4791):899-901, 1987). However, despite their effectiveness against CQ^R *Plasmodium sp.* strains *in vitro*, therapeutically effective doses of many reversal agents would have to be quite high (with associated side effects) to reverse CQ^R *in vivo* (van Schalkwyk *et al.*, *Antimicrob. Agents Chemother.*, 45(11):3171-3174, 2001; Millet *et al.*, *Antimicrob. Agents Chemother.*, 48(7):2753-2756, 2004).

20 New compositions and methods for the treatment of malaria, particularly for the treatment of CQ^R malaria parasites, are needed.

SUMMARY OF THE DISCLOSURE

 This disclosure concerns the discovery of a class of compounds referred to as
25 "reversed chloroquines" (RCQs), which are highly effective against CQ^R and CQ^S malaria parasites. RCQs are hybrid molecules, which include an antimalarial quinoline analog (such as chloroquine) moiety and a CQ^R reversal moiety. An antimalarial quinoline analog moiety of a RCQ is believed to inhibit the growth of *Plasmodium sp.* (the malaria pathogens), and a CQ^R reversal moiety of a RCQ is
30 believed to sensitize CQ^R *Plasmodium sp.* strains to the RCQ inhibitory moiety. Without being limited to one theory, covalently linking these moieties may increase the local concentration of growth-inhibitory and/or CQ^R-reversal functionalities in

- 4 -

the pathogen to effectively inhibit their growth and/or treat CQ^R and CQ^S malaria strains.

Exemplary RCQ chemical structures are provided throughout the disclosure. By way of example, such compounds are represented by the following general

5 tripartite formula



including an antimalarial quinoline analog "Q" covalently linked via a linker "L" to a CQ^R reversal moiety designated by the group "G":

Pharmaceutical compositions containing, and methods of using, the disclosed
10 RCQs are also provided. For example, the disclosed RCQs are effective to treat malaria and/or inhibit the growth of CQ^S and CQ^R *Plasmodium sp.* strains.

The foregoing and other features and advantages will become more apparent from the following detailed description of several embodiments, which proceeds with reference to the accompanying figures.

15

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows electrostatic potential ranges (from negative (dark) to positive (light)) for chloroquine (top left) and RCQ_02 (top right) mapped onto a surface that is defined by a density of 0.002 of an electron (also referred to as a 0.002 density
20 surface).

FIG. 2 shows a series of UV-VIS spectra of heme in the presence of increasing amounts of RCQ_02 at pH 4.7. RCQ_02 absorbance is shown by the open up-arrow. A decreased value of heme absorbance in the Soret region (open down-arrow) indicated RCQ_02 heme binding. These spectra were corrected for the
25 slight change in sample volume (2.5% at the last addition) during the course of the titration.

FIG. 3 shows a series of graphs relating to IC₅₀ determinations for chloroquine (CQ) and RCQ_02 in CQ^R (Dd2) and CR^S (D6) *P. falciparum* strains. FIG. 3A shows dose response curves for CQ as applied to D6 (closed diamonds) and
30 Dd2 (closed squares) strains. FIG. 3B shows dose response curves for RCQ_02 as applied to D6 (closed diamonds) and Dd2 (closed squares) strains. Y-axes in panels A and B are "Relative Fluorescence Units" (*i.e.*, from 0 to 100; see, *e.g.*, De *et al.*, *J.*

- 5 -

Med. Chem., 41(25):4918-4926, 1998; De *et al.*, *Am. J. Trop. Med. Hyg.*, 55(6):579-583, 1996; Smilkstein *et al.*, *Antimicrob. Agents Chemother.*, 48(5):1803-1806, 2004). The x-axes in FIGs. 3A and 3B are log(concentration); hence negative numbers are expected. FIG. 3C is a bar graph consolidating the data from FIGs. 3A and 3B.

FIG. 4 shows a comparison between selected reversal agent aromatic ring ends (Series 1, 2, and 3). These are energy-minimized representative conformers at the HF(6-31G) level in Gaussian03.

10


DETAILED DESCRIPTION

I. Introduction

Disclosed herein are compounds that have been recently discovered that are highly effective against CQ^R and CQ^S malaria parasites. Such compounds may be used, at least, in pharmaceutical compositions, and to treat malaria and inhibit the growth of *Plasmodium sp.*

15

In certain examples, the compounds RCQ compounds include compounds of the general, tripartite formula



20

Such disclosed compounds include a quinoline analog Q (such as a chloroquine derivative) covalently linked to a CQ^R reversal moiety G via a linker L. In some examples, the C₄ dimethylamine side chain of chloroquine is covalently linked to the CQ^R reversal moiety. In other examples, the CQ^R reversal moiety is amitriptyline, amlodipine, azatadine, chlorpheniramine, citalopram, cyclosporine, cyproheptadine, cyproheptadine, desipramine, diethyl-{3-[3-(4-methoxy-benzylidene)-pyrrolidin-1-yl]-propyl}-amine, erythromycin, fantofarone, fluoxetine, haloperidol, icajine, imipramine, isoretuline, ivermectin, ketotefin, ketotifen, nomifensine, NP30: C₉H₁₉-Phenyl-(O-CH₂CH₂)₃₀OH, oxaprotiline, probenecid, progesterone, promethazine, strychnobrasiline, BG958, trifluoperazine, verapamil, or WR 268954.

30

- 6 -

In yet other embodiments, any of the foregoing or other disclosed compounds may be incorporated into pharmaceutical compositions that include a therapeutically effective amount of the compound or extract, and a pharmaceutically acceptable carrier. In some instances, a disclosed pharmaceutical composition
5 further includes at least one additional antimalarial therapeutic agent (such as artesunate, mefloquine, sulfadoxine, or pyrimethamine, or combinations thereof).

The present disclosure also concerns methods of treating a subject for malaria or *Plasmodium sp.* infection by administering to the subject a therapeutically effective amount (such as from about 1 to about 50 mg/kg) of any of the compounds
10 disclosed herein. In some cases, the compound is administered prophylactically. In other embodiments, the malarial pathogen is *P. falciparum*.

Also disclosed are methods for inhibiting the growth of *Plasmodium sp.* involving contacting at least one *Plasmodium sp.* parasite with a growth inhibitory amount (such as from about 1 to about 50 nM) of at least one disclosed compound.
15 In some method embodiments, the *Plasmodium sp.* is *P. falciparum*, *P. vivax*, *P. ovale*, or *P. malariae*, or a combination thereof.

II. Abbreviations and Terms

CQ	chloroquine
CQ ^R	chloroquine resistant (or chloroquine resistance)
CQ ^S	chloroquine sensitive
DV	digestive vacuole
IC ₅₀	concentration of an agent resulting in 50% inhibition of parasite growth
RCQ	reversed chloroquine

20 Unless otherwise noted, technical terms are used according to conventional usage. Definitions of common terms in chemistry terms may be found in *The McGraw-Hill Dictionary of Chemical Terms*, 1985, and *The Condensed Chemical Dictionary*, 1981. The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. "Comprising" means "including"; hence,
25 "comprising A or B" means "including A or B" or "including A and B."

- 7 -

In order to facilitate review of the various embodiments of the invention, the following explanations of specific terms are provided:

Acyl: A group of the formula $RC(O)-$ wherein R is an organic group.

Aliphatic: An acyclic, straight- or branched-chain alkane, alkene, or alkyne, which, unless expressly stated otherwise, contains from one to twenty-five carbon atoms; for example, from one to fifteen, from one to ten, from one to six, or from one to four carbon atoms. The term "lower aliphatic" refers to an aliphatic group containing from one to ten carbon atoms. An aliphatic chain may be substituted or unsubstituted. Unless expressly referred to as an "unsubstituted aliphatic," an aliphatic groups can either be unsubstituted or substituted. An aliphatic group can be substituted with one or more substituents (up to two substituents for each methylene carbon in an aliphatic chain, or up to one substituent for each carbon of a $-C=C-$ double bond in an aliphatic chain, or up to one substituent for a carbon of a terminal methine group). Exemplary aliphatic substituents include, for instance, amine, amide, sulfonamide, halogen, cyano, carboxy, hydroxy, mercapto, trifluoromethyl, alkyl, alkoxy, alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, or other functionality.

Alkenyl: An acyclic, straight- or branched-chain aliphatic radical containing at least two carbon atoms and having at least one carbon-carbon double bond; for example, containing from two to six carbon atoms, including (but not limited to) vinyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, and 5-hexenyl.

Alkoxy: A radical (or substituent) having the structure $-O-R$, where R is a substituted or unsubstituted alkyl. Methoxy ($-OCH_3$) is an exemplary alkoxy group. In a substituted alkoxy, R is alkyl substituted with a non-interfering substituent. "Thioalkoxy" refers to $-S-R$, where R is substituted or unsubstituted alkyl. "Haloalkyloxy" means a radical $-OR$ where R is a haloalkyl.

Alkyl: An acyclic, saturated, branched- or straight-chain hydrocarbon radical, which, unless expressly stated otherwise, contains from one to fifteen carbon atoms; for example, from one to ten, from one to six, or from one to four carbon atoms. This term includes, for example, groups such as methyl, ethyl, n-propyl, isopropyl, isobutyl, t-butyl, pentyl, heptyl, octyl, nonyl, decyl, or dodecyl. The term "lower alkyl" refers to an alkyl group containing from one to ten carbon atoms.

- 8 -

Unless expressly referred to as an "unsubstituted alkyl," alkyl groups can either be unsubstituted or substituted. An alkyl group can be substituted with one or more substituents (for example, up to two substituents for each methylene carbon in an alkyl chain). Exemplary alkyl substituents include, for instance, amine, amide, sulfonamide, halogen, cyano, carboxy, hydroxy, mercapto, trifluoromethyl, alkyl, alkoxy (such as methoxy), alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, alkylsulfano, keto, or other functionality.

Alkylamino: An alkyl group where at least one hydrogen is substituted with an amino, mono-substituted amino or di-substituted amino group.

Amino: A chemical functionality $-N(R)R'$ where R and R' are independently hydrogen, alkyl, heteroalkyl, haloalkyl, aliphatic, heteroaliphatic, aryl (such as optionally substituted phenyl or benzyl), heteroaryl, alkylsulfano, or other functionality. A "primary amino" group is $-NH_2$. "Mono-substituted amino" means a radical $-N(H)R$ substituted as above and includes, *e.g.*, methylamino, (1-methylethyl)amino, phenylamino, and the like. "Di-substituted amino" means a radical $-N(R)R'$ substituted as above and includes, *e.g.*, dimethylamino, methylethylamino, di(1-methylethyl)amino, and the like.

Analog: A molecule that differs in chemical structure from a parent compound, for example a homolog (differing by an increment in the chemical structure, such as a difference in the length of an alkyl chain), a molecular fragment, a structure that differs by one or more functional groups, or a change in ionization. Structural analogs are often found using quantitative structure activity relationships (QSAR), with techniques such as those disclosed in *Remington: The Science and Practice of Pharmacy*, 19th Edition, Chapter 28, 1995.

Animal: Living multi-cellular vertebrate organisms, a category that includes, for example, mammals and birds. The term mammal includes both human and non-human mammals. Similarly, the term "**subject**" includes all animals, including humans, simians, dogs, cats, horses, cows, rodents, *etc.* Likewise, the term "**mammal**" includes both human and non-human mammals.

Aralkyl: an aryl group (such as a phenyl group) appended to an alkyl radical including, but not limited to, benzyl, ethylbenzene, propylbenzene, butylbenzene, pentylbenzene, and the like. Conversely the term "phenylalkyl" refers to a phenyl

- 9 -

- group appended to an alkyl radical. Aralkyl groups, such as benzyl groups, may be unsubstituted or substituted with one, two or three substituents, with substituent(s) independently selected from alkyl, heteroalkyl, aliphatic, heteroaliphatic, thioalkoxy, haloalkyl (such as $-\text{CF}_3$), halo, nitro, cyano, $-\text{OR}$ (where R is hydrogen or alkyl),
- 5 $-\text{N(R)R'}$ (where R and R' are independently of each other hydrogen or alkyl), $-\text{COOR}$ (where R is hydrogen or alkyl) or $-\text{C(O)N(R')R''}$ (where R' and R'' are independently selected from hydrogen or alkyl). Non-limiting examples, include o-, m-, and/or p-chlorobenzyl, o-, m-, and/or p-methoxybenzyl, and o-, m-, and/or p-(trifluoromethyl)benzyl.
- 10 **Aryl:** The term aryl is used to refer to monocyclic and polycyclic aromatic ring systems. Certain examples of aryl groups include a non-aromatic ring fused with one or more aromatic rings. Aryl groups also may be optionally substituted independently with one or more substituents, such as one, two or three substituents per ring selected from alkyl, haloalkyl, alkylthio, heteroalkyl, aliphatic,
- 15 heteroaliphatic, alkoxy, cycloalkyl, hydroxy, halo, cyano, nitro, optionally substituted phenyl, optionally substituted aralkyl (such as phenylalkyl), optionally substituted heteroaryl, optionally substituted heteroaralkyl, amino, monosubstituted amino, disubstituted amino, hydroxylamino, $-\text{OR}$ (where R is hydrogen, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally
- 20 substituted heteroaryl or optionally substituted heteroaralkyl), $-\text{S(O)}_n\text{R}$ (where n is an integer from 0 to 2 and R is alkyl, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, amino, monosubstituted or disubstituted amino),
- 25 $-\text{N(R)C(O)R'}$ (where R is hydrogen or alkyl and R' is hydrogen, alkyl, heteroalkyl, haloalkyl, optionally substituted phenyl, or monosubstituted or disubstituted amino), $-\text{N(R)SO}_2\text{R'}$ (where R is hydrogen or alkyl and R' is alkyl, amino, monosubstituted or disubstituted amino), $-\text{C(O)R}$ (where R is hydrogen, alkyl, haloalkyl, optionally substituted phenyl or optionally substituted heteroaryl), $-\text{COOR}$ (where R is hydrogen, alkyl, optionally substituted phenyl, optionally substituted phenylalkyl,
- 30 optionally substituted heteroaryl or optionally substituted heteroaralkyl), $-\text{C(O)N(R')R''}$ (where R' and R'' are independently selected from hydrogen, alkyl, haloalkyl, optionally substituted phenyl, optionally substituted phenylalkyl,

- 10 -

optionally substituted heteroaryl or optionally substituted heteroaralkyl). In specific examples, the term aryl includes, for example, phenyl, benzyl, phenol, naphthyl, anthracenyl, phenanthrenyl, and substituted derivatives thereof.

Carbocycle: A saturated or unsaturated cyclic radical of 3 to 8 ring atoms in which each of the ring atoms are carbon. Such carbocyclic groups may be optionally substituted independently with one, two or three substituents selected from alkyl, heteroalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aliphatic, heteroaliphatic, aryl, aralkyl, heteroaryl, heteroaralkyl, halo, cyano, acyl, acylamino, amino, monosubstituted amino, disubstituted amino, -COOR (where R is hydrogen or alkyl), -XR (where X is O or S(O)_n, where n is an integer from 0 to 2, and R is hydrogen, alkyl, haloalkyl, cycloalkyl, aralkyl, aryl, heteroaryl or heteroaralkyl), or -C(O)N(R')R'' (where R' and R'' are independently selected from hydrogen or alkyl). Representative examples include, but are not limited to, cyclopentyl, cyclohexyl, cycloheptyl, cycloheptenyl, or cycloheptyl-2, 3, or 4-one, and the like.

Carbonyl: A radical of the formula -C(O)-. Carbonyl-containing groups include any substituent containing a carbon-oxygen double bond (C=O), including acyl groups, amides, carboxy groups, esters, ureas, carbamates, carbonates and ketones and aldehydes, such as substituents based on -COR or -RCHO where R is an aliphatic, heteroaliphatic, alkyl, heteroalkyl, hydroxyl, or a secondary, tertiary, or quaternary amine.

Carboxyl: A -COOH radical. Substituted carboxyl refers to -COOR where R is aliphatic, heteroaliphatic, alkyl, heteroalkyl, or a carboxylic acid or ester.

Cycloalkyl: A saturated monovalent cyclic hydrocarbon radical of three to seven ring carbons, *e.g.*, cyclopentyl, cyclohexyl, cycloheptyl and the like.

Derivative: A chemical substance that differs from another chemical substance by one or more functional groups. Preferably, a derivative (such as a RCQ) retains a biological activity of a molecule from which it was derived (such as a quinoline analog, like chloroquine, or a CQ^R reversal agent). In some cases, a derivative may be a hybrid of at least two otherwise separate chemical substances (such as a hybrid molecule containing a chloroquine moiety and a CQ^R reversal moiety).

- 11 -

Haloalkyl: An alkyl group substituted with one or more same or different halo atoms, *e.g.*, -CH₂Cl, -CF₃, -CH₂CF₃, -CF₂CF₃, -CH₂CCl₃, and the like.

Halogen: The elements fluorine, bromine, chlorine, and/or iodine. The term "halo" refers to fluoro, bromo, chloro and/or iodo substituents.

5 **Heteroalkyl:** An alkyl or cycloalkyl radical as defined above containing at least one heteroatom, such as N, O, S, or S(O)_n (where n is 1 or 2).

Heteroaliphatic: An aliphatic group as defined above containing at least one heteroatom, such as N, O, S, or S(O)_n (where n is 1 or 2).

10 **Heteroaryl:** A monovalent monocyclic or bicyclic aromatic radical of 5 to 10 ring atoms with each ring containing one, two, or three ring heteroatoms selected from N, O, or S, the remaining ring atoms being C. The aromatic radical is optionally fused to a phenyl or an optionally substituted heteroaryl ring or it is optionally substituted independently with one or more substituents, such as one or two substituents selected from alkyl, haloalkyl, heteroalkyl, aliphatic,
15 heteroaliphatic, alkoxy, halo, cyano, nitro, aryl, optionally substituted heteroaryl, amino, monosubstituted amino, disubstituted amino, hydroxyamino, -OR (where R is hydrogen, haloalkyl, or optionally substituted phenyl), -S(O)_nR (where n is an integer from 0 to 2 and R is alkyl, haloalkyl, optionally substituted phenyl, amino, mono or disubstituted amino), -C(O)R (where R is hydrogen, alkyl, haloalkyl or
20 optionally substituted phenyl), -COOR (where R is hydrogen, alkyl or optionally substituted phenyl), -C(O)N(R')R'' (where R' and R'' are independently selected from hydrogen, alkyl, haloalkyl, or optionally substituted phenyl). In specific examples, the term heteroaryl includes, but is not limited to pyridyl, pyrrolyl, thiophene, pyrazolyl, thiazolyl, imidazolyl, pyrimidinyl, thiadiazolyl, indolyl,
25 carbazolyl, azaindolyl, benzofuranyl, benzimidazolyl, benzthiazolyl, quinoxaliny, benzotriazolyl, benzisoxazolyl, purinyl, quinolinyl, isoquinolinyl, benzopyranyl, and derivatives thereof.

Heterocycle: A saturated or unsaturated cyclic radical of 3 to 8 ring atoms in which one or two ring atoms are heteroatomic groups selected from N, O, or
30 S(O)_n (where n is an integer from 0 to 2), the remaining ring atoms being C. The heterocyclic ring may be optionally substituted independently with one, two or three substituents selected from alkyl, heteroalkyl, haloalkyl, cycloalkyl, cycloalkylalkyl,

- 12 -

aliphatic, heteroaliphatic, aryl, aralkyl, heteroaryl, heteroaralkyl, halo, cyano, acyl, acylamino, amino, monosubstituted amino, disubstituted amino, -COOR (where R is hydrogen or alkyl), -XR (where X is O or S(O)_n, where n is an integer from 0 to 2, and R is hydrogen, alkyl, haloalkyl, cycloalkyl, aralkyl, aryl, heteroaryl or heteroaralkyl), or -C(O)N(R')R'' (where R' and R'' are independently selected from hydrogen or alkyl). Representative examples include, but are not limited to, tetrahydropyranyl, piperidino, piperazino, pyrrolidino, and the like.

Hydroxyalkyl: An alkyl group as defined above substituted with at least one hydroxyl group, provided that if two or more hydroxyl groups are present no two hydroxyl groups are on the same carbon atom.

Inhibiting parasite growth: The phrase "inhibiting parasite growth" (and analogous phrases, such as inhibition of parasite growth or inhibition of parasite proliferation) conveys a wide-range of inhibitory effects that an agent (*e.g.*, an antimalarial composition) may have on a normal (for example, untreated or control) rate of parasite growth. The phrase "inhibiting parasite growth" (or like terminology) may be considered relative to a normal (for example, untreated or control) rate of growth of a particular parasite species (such as *Plasmodium sp.*) or a particular life stage of a parasite (such as a blood-stage and/or liver-stage *Plasmodium sp.* parasite). Thus, inhibiting parasite growth includes situations wherein the normal growth rate of a parasite or a particular life-stage of a parasite has slowed (such that parasite number increases over time, but not as rapidly as in a control population), equals zero (such that there is substantially no change in number of parasites in the population over time, *e.g.*, parasite growth is approximately equal to parasite death), or becomes negative (such that the number of parasites decreases over time, *e.g.*, parasite death exceeds parasite growth). A negative rate of parasite growth can (but need not) result in the death of all parasites in a population. For example, an antimalarial agent may inhibit the rate of parasite growth in a manner such that it may or may not result over time in a decrease in the absolute numbers of parasites in the overall population relative to the numbers of parasites present just prior to administering the agent.

Oxygen-containing group: An R-group containing at least one oxygen atom. Exemplary, non-limiting oxygen containing groups include oxygen alone

- 13 -

(which may be attached to the molecule by a single or double bond), hydroxyl, hydroxylalkyl, ether, ester, or any group containing a carbonyl moiety.

Optional: Means that the subsequently described event or circumstance may but need not occur, and that the description includes instances where the event or
5 circumstance occurs and instances in which it does not. For example, "heterocyclo group optionally mono- or di-substituted with an alkyl group" means that the alkyl may but need not be present, and the description includes situations where the heterocyclo group is mono- or disubstituted with an alkyl group and situations where the heterocyclo group is not substituted with the alkyl group.

10 **Phenyl:** Phenyl groups may be unsubstituted or substituted with one, two or three substituents, with substituent(s) independently selected from alkyl, heteroalkyl, aliphatic, heteroaliphatic, thioalkoxy, halo, haloalkyl (such as -CF₃), nitro, cyano, -OR (where R is hydrogen or alkyl), -N(R)R' (where R and R' are independently of each other hydrogen or alkyl), -COOR (where R is hydrogen or alkyl) or –
15 C(O)N(R')R'' (where R' and R'' are independently selected from hydrogen or alkyl). Non-limiting examples, include o-, m-, and/or p-chlorophenyl, o-, m-, and/or p-methoxyphenyl, and o-, m-, and/or p-(trifluoromethyl)phenyl.

Sulfonyl: refers to the radical -SO₂-. The sulfonyl group can be further substituted with a variety of groups to form, for example, sulfonic acids,
20 sulfonamides, sulfonate esters and sulfones.

Thioalkoxyalkyl: A thioalkoxy group appended to an alkyl radical.

Treating or treatment: With respect to disease, either term includes (1) preventing the disease, *e.g.*, causing the clinical symptoms of the disease not to develop in an animal that may be exposed to or predisposed to the disease but does
25 not yet experience or display symptoms of the disease, (2) inhibiting the disease, *e.g.*, arresting the development of the disease or its clinical symptoms, or (3) relieving the disease, *e.g.*, causing regression of the disease or its clinical symptoms.

Tricyclic ring system: A ring system containing three independently carbocyclic or heterocyclic rings including a central ring and two peripheral rings.
30 Each peripheral ring is fused to the central ring, but neither peripheral ring is fused to the other. Each peripheral ring is aromatic (*e.g.*, phenyl) and can be unsubstituted or substituted as described above for a phenyl group. The central ring can be

- 14 -

aromatic or aliphatic, unsaturated or partially saturated (for example, with bridgehead carbons being unsaturated and one or more other ring bonds being saturated), and can be unsubstituted or substituted as described for carbocyclic or heterocyclic rings (and as valence rules allow). Non-limiting examples of a tricyclic ring system include *N*-xanthenyl, *N*-phenothiazinyl, *N*-phenoxazinyl, *N*-carbazolyl; 10H-acridin-9-one-10-yl; 5,11-dihydro-dibenzo[a,d]cyclohepten-10-one-5-yl; 10,11-dihydro-5H-dibenzo[a,d]cycloheptene-5-yl; 10H-benzo[b][1,8]naphthyridin-5-one-*N*-yl; 6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2 b]pyridine-11-yl.

Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers." When a compound has an asymmetric center, for example, if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (*i.e.*, as (+) or (-) isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture."

The compounds described herein may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see, *e.g.*, March, *Advanced Organic Chemistry*, 4th edition, New York: John Wiley and Sons, 1992, Chapter 4).

It is further to be understood that any molecular weight or molecular mass values are approximate and are provided only for description. All publications,

- 15 -

patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including explanations of terms, will control.

Except as otherwise noted, the methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. See, e.g., Loudon, *Organic Chemistry*, Fourth Edition, New York:Oxford University Press, 2002, pp. 360-361, 1084-1085; Smith and March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, Fifth Edition, Wiley-Interscience, 2001; or Vogel, *A Textbook of Practical Organic Chemistry, Including Qualitative Organic Analysis*, Fourth Edition, New York:Longman, 1978.

III. RCQ Compounds

Disclosed herein are compounds (referred to as RCQs) useful for the treatment of malaria and *Plasmodium sp.* infection. Such compounds are equally effective against CQ^R and CQ^S strains of *Plasmodium sp.*, and provide another weapon in the arsenal against malaria, which cumulatively has killed more people than any other infectious disease in history (e.g., Resolution 61-02, Association of American Physicians and Surgeons, Inc., Tucson, AZ, 2004; available at the website having the address www.aapsonline.org/resolutions/2004-2.htm). The disclosed compounds can be prepared by any methods known to those of skill in the art; for example by chemical synthesis. Non-limiting exemplary methods for synthesis of particular RCQ embodiments are provided herein.


- 16 -

A. Chemical Formulas

Examples of the disclosed RCQ compounds include compounds of the general, tripartite formula:



5 wherein Q represents a quinoline derivative of the formula




wherein X_1 and X_2 are independently alkoxy, optionally substituted amino, halo, haloalkyl, hydroxy; n is from 0 to 2; m is from 0 to 4; and R_1 is H, optionally substituted alkyl, heteroalkyl, alkylamino, sulfonyl (such as alkyl or aryl sulfonyl),
 10 haloalkyl, or carbonyl.

With continued reference to the general formula above, L represents a linker, typically constituting a hydrocarbon chain, optionally branched and/or including heteroatoms, such as nitrogen and/or oxygen. Where the linker group is a hydrocarbon chain, the group typically includes from 2 to about 12 carbon atoms,
 15 optionally interrupted by one or more heteroatom. For example, the L can include one or more ethylene glycol groups, ethanolamine groups and/or diaminoethane moieties. Similarly, linkers can include a propyl derivative optionally containing one or more heteroatoms, such as a 1,3-diaminopropyl group. The chain can comprise aliphatic and aryl groups and can comprise straight chain, branched chain
 20 and/or cyclic groups, such as heterocyclic groups.


With reference to the general formula above, G represents a reversal agent moiety. Typically, suitable G groups include derivatives of reversal agents having one or more cyclic groups.

- 17 -

In some embodiments, a disclosed RCQ conforms to the chemical structure:



wherein G is



- 5 (X₁)_m and (X₂)_n are independently halo, haloalkyl (such as trifluoromethyl), amino, hydroxyl, alkoxy (such as methoxy), alkylamino, or arylamino; m is 0 to 4 (such as 0 to 2, 1 or 2, or 1); n is 0 to 2 (all ring valences are filled by H where m and/or n is 0);

- 10 R₁ is H, alkyl, heteroalkyl, alkylamino (such as methylamino), haloalkyl (such as trifluoromethyl), amide (such as -C(O)NH₂), alkylamide, or alkylsulfonamide;

- 15 R₂ is H, alkyl (such as methyl or ethyl), heteroalkyl, alkylamino (such as methylamino), haloalkyl (such as trifluoromethyl), alkylsulfonamide, amide, alkylamide (such as -(CH₂)_n-C(O)NH₂, where n is 1, 2, 3, or 4), cyano, furyl, or alkylfuran (such as -(CH₂)_n-furan, where n is 1, 2, 3, or 4); L₁ and L₂ are independently alkyl or heteroalkyl;

Y and Y₁ independently are N; CH; CH₂CH or C(O)CH;

- 20 R₃ and R₄ are independently alkyl, cycloalkyl, heterocyclyl, aryl (such as substituted or unsubstituted phenyl or benzyl), heteroaryl (such as pyridyl, pyridylalkyl, pyrazinyl, furyl, or indolyl) or together with Y₁ form a polycyclic ring system, such as a bicyclic or tricyclic ring system; and

ring 1 is a five-, six-, or seven-membered heterocyclic ring.







- 25 In some examples, (X₁)_m and (X₂)_n are independently halo (such as chloro), haloalkyl (such as trifluoromethyl), or alkoxy (such as methoxy). In other examples, (X₁)_m are independently halo (such as chloro), or haloalkyl (such as trifluoromethyl)

- 18 -

and n is 0 (*i.e.*, all ring valences satisfied by H). In more particular examples, n is 0, m is 2, and $(X_1)_1$ and $(X_1)_2$ are C_7 and C_6 substituents, respectively, which are independently selected from halo (such as chloro), or haloalkyl (such as trifluoromethyl). In even more particular examples, n is 0, m is 1, and $(X_1)_1$ is

5 chloro at position C_7 . Some non-limiting representative chemical structures illustrating exemplary variations of $(X_1)_m$ and/or $(X_2)_n$ are shown in the following Table 1. The Table 1 structures can also be described as representative antimalarial quinoline derivatives.

Table 1. Exemplary Antimalarial Quinoline Moieties

10 In some embodiments, R_1 and R_2 are independently H, alkyl having from 1 to 4 carbon atoms (such as from 1 to 2, or 1 carbon atom), or heteroalkyl having from 1 to 4 carbon atoms and at least one heteroatom (such as N, S, or O). In particular examples, R_1 and R_2 are independently selected from H, methyl, trifluoromethyl, $-CH_2CH_2CF_3$, $-CH_2CF_3$, alkylamine (such as $-CH_2CH_2NH_2$,

15 $-CH_2NH_2$), or alkylamide (such as $-(CH_2)_n-C(O)NH_2$, where n is 1, 2, 3, or 4). In more particular examples, R_1 and R_2 are independently selected from H, methyl or methylamine. In even more particular examples, R_1 and R_2 are each H. In some embodiments, R_2 is further selected from cyano, furyl, or alkylfuran (such as

- 19 -

-(CH₂)_n-furan, where n is 1, 2, 3, or 4). In specific embodiments, R₁ is H and R₂ is H, methyl, amine, alkylamine (such as -(CH₂)_n-NH₂, where n is 1, 2, 3, or 4), amide, alkylamide (such as -(CH₂)_n-C(O)NH₂, where n is 1, 2, 3, or 4), furyl, alkylfuran (such as -(CH₂)_n-furan, where n is 1, 2, 3, or 4), or cyano. In other specific
5 embodiments, R₁ is H and R₂ is H, methyl, methylamine, ethylamine, amide, methylamide, ethylamide, methylfuran, or cyano.



L₁ and L₂ are each linkers, which are believed to tolerate considerable variation in structure without substantial effect on a biological activity of the RCQ molecule (such as inhibition of CQ^R and/or CQ^S *Plasmodium sp.* growth). For
10 example, it is known that CQ with only 3 methylene units (at a position analogous to L₁ in Formula I) and no branching methyl between the CQ aliphatic amines (at a position analogous to R₁ in Formula I) is effective against CQ^R *P. falciparum* strains (De *et al.*, *J. Med. Chem.*, 41(25):4918-4926, 1998; De *et al.*, *Am. J. Trop. Med. Hyg.*, 55(6):579-583, 1996).

15 In some embodiments, L₁ and L₂ are alkyl or heteroalkyl independently having from 1 to 12 atoms (such as from 1 to 7 carbons, from 2 to 5 carbons, or from 3 to 4 carbons). In other embodiments, each carbon in L₁ and L₂ has two substituents, each of which substituents is independently selected from H, alkyl (such as methyl or ethyl), haloalkyl (such as trifluoromethyl), alkoxy (such as
20 methoxy), amino (such as -NH₂), or halo (such as -F or -Cl). In particular embodiments, substituents bonded to one carbon atom are the same so that neither L₁ nor L₂ include any chiral carbons. In other particular embodiments, L₁ and L₂ are independently from 1 to 12 atoms in length (such as from 2 to 6 atoms, for example ethyl, propyl, hexyl or the like), and each position of L₁ and L₂ is independently
25 selected from methylene, dihalomethylene (such as difluoromethylene or dichloromethylene), di(methylamino), -NH-, and -O-. In specific embodiments, L₁ and L₂ are independently, ethyl, propyl, butyl, -(CH₂)_{n1}-O-(CH₂)_{n2}- (where n₁ and n₂ are independently 1, 2, 3, or 4), or -(CH₂)_n-NH- (where n is 1, 2, 3, or 4). In even more specific embodiments, L₁ is ethyl, propyl, or butyl, and L₂ is ethyl, propyl,
30 butyl, -CH₂-O-CH₂-, or -(CH₂)₂-NH-. In still more specific embodiments, L₁ and L₂ are independently, ethyl, propyl, or butyl moieties.

- 20 -

Some non-limiting representative chemical structures illustrating exemplary variations of alkyl L_1 and/or L_2 are shown in the following table.

Table 2. Exemplary Alkyl Linkers


L_1:  L_2: 					
R7	R8	R9	R10	o	p
H	H	H	H	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
-F	-F	-F	-F	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
H	H	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
H	H	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
H	H	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
H	H	-F	-F	2, 3, or 4	2, 3, or 4
H	H	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	H	H	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	H	H	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	H	H	2, 3, or 4	2, 3, or 4
-F	-F	H	H	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	H	H	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	-F	-F	2, 3, or 4	2, 3, or 4
-CH ₃	-CH ₃	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
-F	-F	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	-CH ₃	-CH ₃	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	-F	-F	2, 3, or 4	2, 3, or 4
-CH ₂ CH ₃	-CH ₂ CH ₃	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
-F	-F	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	-CH ₂ CH ₃	-CH ₂ CH ₃	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	-F	-F	2, 3, or 4	2, 3, or 4
-CF ₃	-CF ₃	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
-F	-F	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	-CF ₃	-CF ₃	2, 3, or 4	2, 3, or 4
-F	-F	-NH ₂	-NH ₂	2, 3, or 4	2, 3, or 4
-NH ₂	-NH ₂	-F	-F	2, 3, or 4	2, 3, or 4

Exemplary R_3 and R_4 groups include aryl groups and substituted alkyl groups, such as cyclohexyl, methylcyclohexyl ($-\text{CH}_2\text{C}_6\text{H}_{11}$), adamantyl, methyladamantyl and ethyl adamantyl. Typically, at least one of R_3 and R_4 is a sterically

- 21 -


bulky group. Suitable sterically bulky groups for incorporation as R_3 and/or R_4 will be readily apparent to those of ordinary skill in the art upon consideration of the present disclosure.

Other exemplary compounds disclosed herein conform to the chemical
 5 structure:




wherein $(X_1)_m$, $(X_2)_n$, R_1 , R_2 , L_1 , L_2 , Y_1 , R_3 and R_4 are as described for above.

In one aspect of this formula R_3 and R_4 can represent arylalkyl groups, which may be the same or different. For example, a specific group of RCQ compound
 10 structures have the formula:




wherein Ar^1 and Ar^2 are independently selected aryl groups. In one embodiment, such compounds have an oxygen-containing linker group, for example:



- 22 -

In some embodiments, R_3 and R_4 independently have a chemical structure as follows:




where each X_3 is independently halo, haloalkyl (such as trifluoromethyl), amino, hydroxyl, hydroxyalkyl, alkylamino, cyano, alkoxy, sulfonamide, mercapto, keto, or carboxyl; s is 0 to 5 as valence requirements permit (for example, when Y_2 is nitrogen, s is from 0 to 4); Y_2 is C, S, or N; L_3 is a linker, which is alkyl, heteroalkyl, -N-, -O-, or -S-.

In some examples, $(X_3)_s$ are independently fluoro, chloro, trifluoromethyl, methoxy, amine (*e.g.*, -NH₂), carboxyl (*e.g.*, -COOH), sulfonamide (*e.g.*, -SO₂NH₂), hydroxylalkyl (*e.g.*, -(CH₂)_n-OH, where n is 1, 2, 3, or 4), and/or s is 0 to 3, 0 to 2, or 1. In other examples, Y_2 is nitrogen. In particular examples, L_3 is alkyl (such as methyl, ethyl, propyl, i-propyl, or butyl), alkoxy (such as methoxy or ethoxy), alkylamino (such as methylamino or ethylamino), -N-, or -O-. In some instances, R_3 and R_4 are the same (referred to as, symmetric) and, in other instances, R_3 and R_4 are different, such compounds are referred to herein as being asymmetric.

In other examples, X_3 is independently fluoro, chloro, trifluoromethyl, or methoxy. X_3 can be independently selected from these groups for each occurrence "s." In still other examples, L_3 is methyl, ethyl, methoxy, -N-, or -O-.

In some embodiments, R_3 and R_4 together with Y_1 form the central ring of a polycyclic ring system, such as a bicyclic or tricyclic ring system. In particular embodiments, the central ring formed by R_3 , R_4 , and Y_1 is a 5-, 6-, or 7-membered ring. Examples of suitable structures for a tricyclic ring system formed by R_3 , R_4 , and Y_1 include, without limitation:

- 23 -



where Y_2 is NR_{14} , O, or S; R_{13} is H, alkyl, heteroalkyl, =O, amino, amine, amide, sulfonamide, halo (such as F, Cl), cyano, hydroxy, mercapto, haloalkyl (such as trifluoromethyl), alkoxy (such as methoxy), alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, or alkylsulfano; $(X_4)_q$ and $(X_5)_t$ are independently H, alkyl, halo (such as F, Cl), haloalkyl (such as trifluoromethyl), alkoxy (such as methoxy), amino, hydroxyl, alkylamino, cyano, or mercapto; R_{14} is H, lower alkyl or acyl; q and t are independently 0 to 4 as valence requirements permit (such as 0 to 2, 1 or 2, or 1); and each of the peripheral rings can independently include at least one heteroatom (such as N) at any position that valence requirements permit. In some examples, $(X_4)_q$ and $(X_5)_t$ are independently H, methyl, fluoro, chloro, trifluoromethyl, methoxy, or cyano. In other examples, Y_2 is N or S. In particular examples, R_{13} is H or =O (and the ring carbon, which is double bonded to oxygen is otherwise single bonded to adjacent atoms).



















- 24 -

Some specific embodiments of RCQ compounds are provided in the following table.

Table 3. Exemplary RCQ Compounds

RCQ_05	RCQ_06	RCQ_07
RCQ_08	RCQ_05(desMe)	RCQ_06(desMe)
RCQ_07(desMe)	RCQ_08(desMe)	RCQ_09
RCQ_10	RCQ_11	RCQ_12
RCQ_13	RCQ_23mPoP	RCQ_14

Table 3. Exemplary RCQ Compounds

 RCQ_14(pP)2	 RCQ_21	 RCQ_22
 RCQ_23	 RCQ_23(oA)2	 RCQ_24
 RCQ_25	 RCQ_101ca	 RCQ_101fm
 RCQ_102 (a chlorpheniramine derivative)	 RCQ_104	
		
		 RCQ_150 (a verapamil derivative)

- 26 -

Table 3. Exemplary RCQ Compounds








 RCQ_33	 RCQ_26	
		
 RCQ_101 (a chlorpheniramine derivative)	 RCQ_101 (a chlorpheniramine derivative)	 RCQ_103 (a fluoxetine derivative)
 RCQ_61		

Table 3. Exemplary RCQ Compounds













		
 RCQ_01	 RCQ_02	 RCQ_03
 RCQ_04	 RCQ_105	

Table 3. Exemplary RCQ Compounds

		
 RCQ_130 (a promethazine derivative)	 RCQ_131 (a promethazine derivative)	 RCQ_132
 RCQ_133		


Also disclosed herein are compounds that conform to the chemical structure:




where $(X_1)_m$, $(X_2)_n$, R_1 , and L_1 are as described above. Ring 1 is a five-, six- or
 5 seven-membered heterocyclic ring; Y_1 is carbon; R_3 and R_4 are independently aryl or heteroaryl, or together with Y_1 form a tricyclic ring system.

- 29 -

In some embodiments, Ring 1 has the following structure:




In some embodiments of this formula, and as described above, R_3 and R_4 are aryl groups. Examples of suitable aryl groups for R_3 and R_4 include, without limitation those of the formula:



wherein $(X_3)_s$, s , Y_2 , and L_3 are as described above.

In other examples, Y_1 , R_3 and R_4 form the center ring of a tricyclic ring system. Non-limiting exemplary structures for a tricyclic ring system formed by Y_1 , R_3 and R_4 include:



where Y_2 is NR_{14} , O, CH (when Y_2 double bonded to CR_{13}), CH_2 or S; R_{13} is H, alkyl, heteroalkyl, =O, amino, amine, amide, sulfonamide, halo (such as F, Cl), cyano, hydroxy, mercapto, haloalkyl (such as trifluoromethyl), alkoxy (such as methoxy), alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, or alkylsulfano; $(X_4)_q$ and $(X_5)_t$ are independently alkyl, halo (such as F or Cl), haloalkyl (such as trifluoromethyl), alkoxy (such as methoxy), amino, hydroxyl, alkylamino, cyano, or mercapto; R_{14} is H, acyl or lower alkyl; q and t are independently 0 to 4 as valence requirements permit (hydrogen satisfies all ring

- 30 -

valence requirements where q or t is 0); and each of the peripheral rings can independently include at least one heteroatom (such as N) at any position that valence requirements permit. In some examples, $(X_4)_q$ and $(X_5)_t$ are independently methyl, fluoro, chloro, trifluoromethyl, methoxy, or cyano. In other examples, Y_2 is N or S. In particular examples, R_{13} is H or =O (and the ring carbon, which is double bonded to oxygen is otherwise single bonded to adjacent atoms).

Some specific embodiments of disclosed tetrasubstituted ethylene-based RCQ compounds are listed in Table 4.


Table 4. Exemplary Tetrasubstituted Ethylene-Based Embodiments

RCQ_42	RCQ_41H2	RCQ_110 (a cyproheptadine derivative)
RCQ_111 (a ketotifen derivative)	RCQ_112 (an azatadine derivative)	RCQ_113 10H-anthracen-9-one derivative
RCQ_114 10H-Anthracen-9-ylidene derivative	RCQ_115 4-thioxanthen-9-ylidene-piperidin-1-yl derivative	RCQ_116 4-Benzhydrylidene derivative

Table 4. Exemplary Tetrasubstituted Ethylene-Based Embodiments

RCQ_117 Dipyridin-2-yl-methylene derivative	


Also disclosed herein are compounds that conform to the chemical structure:



where $(X_1)_m$, $(X_2)_n$, R_1 , and L_1 are as described above. In some embodiments, L_1 is
 5 also $-(CH_2)_{n1}-N(R')-(CH_2)_{n2}-N(R'')-(CH_2)_{n3}-$ (where n_1 , n_2 , and n_3 are
 independently 1, 2, 3, or 4, and R' and R'' are independently H or alkyl (such as
 methyl, ethyl or propyl). Ring 1 is a five-, six- or seven-membered heterocyclic
 ring; Y represents $-N-$; $-CH-$; $-CH_2CH-$ or $-C(O)CH-$; R_3 and R_4 are
 independently cycloalkyl, aryl or heteroaryl. In certain examples R_3 and R_4 and Y_1
 10 together form a ring system, such as a bicyclic or tricyclic ring system, which can
 include aliphatic rings, aromatic rings, or both.

In particular examples, a ring system formed by Y_1 , R_3 and R_4 has the
 following structure:

- 32 -



where $(X_4)_q$ and $(X_5)_t$ are independently lower alkyl (such as methyl), halo (such as fluoro or chloro), haloalkyl (such as trifluoromethyl), alkoxy (such as methoxy) or cyano. In some examples, q and t are each 0 and all ring valence requirements are

5 satisfied by hydrogen.


Table 5. Exemplary Tricyclic RCQ Compounds

- 33 -

Table 5. Exemplary Tricyclic RCQ Compounds


RCQ_131	RCQ_132	RCQ_133
RCQ_134	RCQ_135	RCQ_136

In one embodiment, Y₁, R₃ and R₄ together form an aliphatic ring system, for example a sterically bulky ring system, such as




- 34 -

In one aspect, disclosed RCQ compounds are represented by the formula:



wherein X_1 is a halogen; X represents $-N-$ or $-CH-$; Y_1 represents $-N-$; $-CH-$; $-CH_2CH-$ or $-C(O)CH-$ and u is from 0 to 2. Embodiments of such compounds can have the formula:























and specific examples of such compounds are recited in Table 6.

Table 6. Exemplary Piperazine-based RCQ Compounds





RCQ_31	RCQ_31(pAc)2	RCQ_31(pCH2OH)2
RCQ_32	RCQ_34	RCQ_35
RCQ_46H2	RCQ_46H2	RCQ_46H2

Table 6. Exemplary Piperazine-based RCQ Compounds


		
 RCQ_37F2	 RCQ_31(oP)2	
		
		
		
		
		
		

- 36 -

Table 6. Exemplary Piperazine-based RCQ Compounds

Additional RCQ compounds have the aminopiperidine-based formula:



and specific examples of such compounds are recited in Table 7.

5

Table 7. Exemplary Aminopiperidine RCQ Compounds





		
 RCQ_81	 RCQ_91	 RCQ_91pPNH2

Table 7. Exemplary Aminopiperidine RCQ Compounds





























		
 RCQ_91pPCH2OH	 RCQ_91pPfm	 RCQ_91(fm)2
 RCQ_91(pP)2	 RCQ_91_pNpSA	 RCQ_91(pP)2
 RCQ_91pNIm	 RCQ_91pNpSA	 RCQ_91(pP)2
 RCQ_91pNIm	 RCQ_91pNpSA	 RCQ_91(pP)2

Table 7. Exemplary Aminopiperidine RCQ Compounds

Additional RCQ compounds have a diazaphane-based formula:



- 39 -

and specific examples of such compounds are recited in Table 8.

Table 8. Exemplary Diazapane RCQ Compounds











		
 RCQ_35(pP)2	 RCQ_36	 RCQ_72
 RCQ_73	 RCQ_74	 RCQ_76

Table 9. Exemplary Pyrrolidine RCQ Compounds


		
 RCQ_151 (a WR268954 derivative)		

- 40 -

Also disclosed herein are compounds having a cyclic linker group linking the quinoline derivative and the reversal agent moiety. Such compounds can have the formula



- 5 wherein R_1 and L , together form a cyclic group. Examples of such compounds can conform to the chemical structure:



wherein the variable groups are as set forth above. Particular examples of such compounds have the structures recited in Table 10.

Table 10. Exemplary RCQ Compounds Having a Cyclic Linker



















		
 RCQ_51	 RCQ_51(pCH₂OH)₂	 RCQ_51(pAc)₂

Table 10. Exemplary RCQ Compounds Having a Cyclic Linker

		
 RCQ_51(pP)2	 RCQ_161	 RCQ_161(pN)2
 RCQ_251	 RCQ_251(pOH)2	 RCQ_251(pCOOH)2
 RCQ_251(pP)2	 RCQ_261	 RCQ_261(pP)2
 RCQ_263	 RCQ_263(pP)2	 RCQ_351
 RCQ_351(pP)2		

- 42 -

B. Additional RCQ Compounds

In some embodiments, an RCQ compound includes a quinoline analog (such as chloroquine) covalently linked to a CQ^R reversal moiety. Typically, a quinoline analog in a disclosed RCQ compound has a C₄ substitution, where the C₄ substituent (or a modified form thereof) can be used to covalently link a CQ^R reversal moiety to the quinoline-analog core. Exemplary C₄ substituents include, for example, alkylamino, alkylhalo (such as alkylchloro), alkoxy, or thioalkyl. Others have chemically modified the CQ molecular framework and found that the C₄ side chain can be modified without substantial loss of *in vitro* antimalarial efficacy (De *et al.*, *J. Med. Chem.*, 41(25):4918-4926, 1998; Egan *et al.*, *J. Med. Chem.*, 43(2):283-291, 2000); thus, it is believed that a wide range of linkers between a quinoline analog (such as CQ) and a CQ^R reversal moiety can be used with minimal impact on the antimalarial efficacy of an RCQ. In addition, the chiral methyl of the CQ alkylamino side chain is known to be unnecessary for action against malaria (De *et al.*, *J. Med. Chem.*, 41(25):4918-4926, 1998; De *et al.*, *Am. J. Trop. Med. Hyg.*, 55(6):579-583, 1996); thus, in particular examples, RCQ molecules do not have any chiral centers. Advantageously, production of non-chiral RCQ molecules may be more cost effective than production of chiral molecules, which may further the goal of providing low-cost drugs for malaria treatment (especially in impoverished countries).

A quinoline analog as envisioned herein also can be substituted at other carbons of the quinoline ring system. In some examples, a quinoline analog is a quinoline ring substituted at C₆, C₇, or C₈, or combinations thereof, with, for example, halo, alkylhalo (such as trifluoromethyl), amino, alkylamino, or hydroxyl substituents. In particular examples, quinoline analogs include, for instance, chloroquine, 7-chloroquinoline, 8-aminoquinoline, amodiaquine, primaquine, 7-(trifluoromethyl) quinoline, or 6-(trifluoromethyl) chloroquine, 6,7-dichloroquinoline, and as illustrated in the foregoing chemical formulas.

CQ^R reversal agents suitable for incorporation as, for example "G" in the general structures presented above, are known in the art (*e.g.*, Bhattacharjee *et al.*, *J. Chem. Inf. Comput. Sci.*, 42:1212-1220, 2002; Guan *et al.*, *J. Med. Chem.*, 45(13):2741-2748, 2002). CQ^R reversal agents cause CQ^R *Plasmodium sp.* strains

- 43 -

(such as *P. falciparum* CQ^R strains) to revert to CQ sensitivity. An exemplary pharmacophore for a CQ^R reversal agent has recently been identified (e.g., Bhattacharjee *et al.*, *J. Chem. Inf. Comput. Sci.*, 42:1212-1220, 2002). This representative pharmacophore is characterized by two hydrophobic groups (such as aromatic rings or fused ring systems) located somewhat near each other and an aliphatic chain terminated by a base (Bhattacharjee *et al.*, *Antimicrob. Agents Chemother.*, 45(9):2655-2657, 2001; Bhattacharjee *et al.*, *J. Chem. Inf. Comput. Sci.*, 42(5):1212-1220, 2002). Without being limited to one theory, CQ^R reversal agents are believed to exert their biological effect, at least in part, by decreasing the efflux of CQ (or CQ-like drugs) from the plasmodial DV, which efflux is thought to be mediated by a membrane pump (e.g., PfCRT; see, Zhang *et al.*, *Biochem.*, 43(26):8290-8296, 2004; Bennett *et al.*, *Mol. Biochem. Parasitol.*, 133(1):99-114, 2004; Martin and Kirk, *Mol. Biol. Evol.*, 21(10):1938-1949, 2004). CQ^R reversal agents are not thought to have a CQ-independent toxic effect on plasmodial parasites. Exemplary reversal agents include, without limitation, amitriptyline, amlodipine, azatadine, chlorpheniramine, citalopram, cyclosporine, cyproheptadine, cyproheptadine, desipramine, dibenzosuberanylpiperazine derivatives (Osa *et al.*, *J. Med. Chem.*, 46(10):1948-1956, 2003), diethyl-{3-[3-(4-methoxy-benzylidene)-pyrrolidin-1-yl]-propyl}-amine, erythromycin, fantofarone (Adovelande *et al.*, *J. Biochem. Pharmacol.*, 55(4):433-440, 1998), fluoxetine, haloperidol, icajine, imipramine, isoretuline, ivermectin, ketotefin, ketotifen, nomifensine, nonylphenoethoxylates (including, e.g., NP30: C₉H₁₉-Phenyl-(O-CH₂CH₂)₃₀OH; Crandall *et al.*, *Antimicrob. Agents Chemother.*, 44(9):2431-2434, 2000), oxaprotiline, probenecid, progesterone, promethazine, strychnobrasiline, BG958 (a 9,10-dihydroethanoanthracene derivative; Millet *et al.*, *Antimicrob. Agents Chemother.*, 48(12):4869-4872, 2004), trifluoperazine, verapamil, and WR 268954 (De *et al.*, *Am. J. Trop. Med. Hyg.*, 49(1):113-120, 1993).

A CQ^R reversal moiety as described throughout the specification includes any known CQ^R reversal agent or derivative thereof that retains CQ^R reversal function, or any chemical structure that satisfies a CQ^R reversal agent pharmacophore (such as the Bhattacharjee *et al.* pharmacophore described above) and confers CQ^R reversal function. As discussed above, CQ^R reversal function is the

- 44 -

ability of a compound (or moiety) to sensitize at least one CQ^R *Plasmodium sp.* strain to CQ. A CQ^R *Plasmodium sp.* strain is sensitized to CQ if a combination of CQ and a CQ^R reversal agent/moiety (or in the presence of a RCQ) inhibits parasite growth as compared to parasite growth in the presence of CQ alone. For example,
5 parasite growth in the presence of a combination of CQ and a CQ^R reversal agent/moiety (or in the presence of a RCQ) may be at least about 10%, at least about 25%, at least about 40%, at least about 50%, at least about 75%, at least about 80%, at least about 90%, or at least about 95% less than parasite growth in the presence of CQ only.

10 A quinoline analog (such as chloroquine) can be covalently linked to a CQ^R reversal agent moiety via commonly known chemical syntheses. Several chemical syntheses useful for such purposes are described throughout this specification. In one example, the C₄ side chain of a quinoline analog (such as CQ) may contain (or be modified to contain) a leaving group (such as a leaving group located at the
15 terminus of the C₄ side chain). A leaving group, as is well known in the art, is an atom or group capable of being displaced by a nucleophile. Suitable leaving groups include halogen, alkanesulfonyloxy, arenesulfonyloxy, ester, or amino; more particular non-limiting examples of leaving groups are chloro, bromo, iodo, mesyloxy, tosyloxy, trifluorosulfonyloxy, methoxy, N,O-dimethylhydroxylamino,
20 and the like. When reacted with a CQ^R reversal agent having an aliphatic chain terminated by a base (which acts as a nucleophile under the reaction conditions), the basic residue (*e.g.*, a nitrogen) of the reversal agent is substituted for the leaving group by a commonly known nucleophilic substitution (S_N2) reaction to form a quinoline analog covalently linked to the CQ^R reversal agent (see, *e.g.*, Loudon,
25 *Organic Chemistry*, Fourth Edition, New York:Oxford University Press, 2002, pp. 360-361, 1084-1085). Conversely, the terminal base of the CQ^R reversal agent can be modified to a leaving group, and the C₄ side chain of a quinoline analog (such as CQ) may contain (or be modified to contain) an nucleophile (such as an amine, alkoxide, or enolate). In this latter example, the nucleophile present on the C₄ side
30 chain of the quinoline analog can be substituted (via S_N2 reaction) for the leaving group on the CQ^R reversal agent (see, *e.g.*, Loudon, *Organic Chemistry*, Fourth

- 45 -

Edition, New York:Oxford University Press, 2002, pp. 360-361, 1084-1085); thereby, forming the combined RCQ molecule.

Non-limiting representative RCQs having a quinoline analog (such as chloroquine) covalently linked to a CQ^R reversal agent moiety include

5 N-(7-chloro-quinolin-4-yl)-N'-(2-phenothiazin-10-yl-ethyl-propane-1,3-diamine (a promethazine derivative; see RCQ_100); N-[3-(4-chloro-phenyl)-3-pyridin-2-yl-propyl]-N'-(7-chloro-quinolin-4-yl)-N-methyl-propane-1,3-diamine (a chlorpheniramine derivative; see RCQ_101); N-(7-chloro-quinolin-4-yl)-N'-[4-phenyl-4-(4-trifluoromethyl-phenoxy-butyl)-propane-1,3-diamine (a

10 fluoxetine derivative; see RCQ_103); (7-chloro-quinolin-4-yl)-[3-(4-dibenzo[a,d]cyclohepten-5-ylidene-piperidin-1-yl)-propyl]-amine (a cyproheptadine derivative; see RCQ_110); 5-{1-[3-(7-chloro-quinolin-4-ylamino)-propyl]-piperidin-4-ylidene}-5,11-dihydro-dibenzo[a,d]cyclohepten-10-one (a ketotifen derivative; see RCQ_111); and (7-chloro-quinolin-4-yl)-

15 {3-[4-(5,6-dihydro-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-piperidin-1-yl]-propyl}-amine (a azatadine derivative; see RCQ_112); 5-{2-[3-(7-Chloro-quinolin-4-ylamino)-propylamino]-ethyl}-[2-(3,4-dimethoxy-phenyl)-ethyl]-amino}-2-(3,4-dimethoxy-phenyl)-2-isopropyl-pentanenitrile (a verapamil derivative; see

20 RCQ_150); N-(7-Chloro-quinolin-4-yl)-N'-[2-(ethyl-{3-[3-(4-methoxy-benzyl)-pyrrolidin-1-yl]-propyl}-amino)-ethyl]-propane-1,3-diamine (a WR268954 derivative; see RCQ_151).

C. Methods of RCQ Synthesis

The disclosed RCQ compounds can be synthesized by any method known in the art. Many general references providing commonly known chemical synthetic

25 schemes and conditions useful for synthesizing the disclosed compounds are available (see, e.g., Smith and March, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, Fifth Edition, Wiley-Interscience, 2001; or Vogel, *A Textbook of Practical Organic Chemistry, Including Qualitative Organic Analysis*, Fourth Edition, New York:Longman, 1978).


30 Compounds as described herein may be purified by any of the means known in the art, including chromatographic means, such as HPLC, preparative thin layer chromatography, flash column chromatography and ion exchange chromatography.

- 46 -

Any suitable stationary phase can be used, including normal and reversed phases as well as ionic resins. Most typically the disclosed compounds are purified via silica gel and/or alumina chromatography. See, *e.g.*, *Introduction to Modern Liquid Chromatography*, 2nd Edition, ed. L.R. Snyder and J.J. Kirkland, John Wiley and Sons, 1979; and *Thin Layer Chromatography*, ed E. Stahl, Springer-Verlag, New York, 1969.


Suitable exemplary syntheses of RCQ compounds are provided below:

1. One Method for Synthesis of RCQ_02 and Structurally Related RCQs




10

2. Representative Nucleophilic Substitution (S_N2) Reaction




15

3. Exemplary Synthesis of Intermediates Useful for RCQ Synthesis





4. Representative Synthesis of Asymmetrically Substituted RCOs

Schemes 4 and 5 demonstrate how even unsymmetrical aralkyl amines can be prepared. This method is exceedingly flexible (Ar^1 and Ar^2 may be derived from a variety of aryl groups as described above, including, without limitation phenyl groups, pyridines, furans, imidazoles, etc.). These syntheses are amenable to execution via parallel synthesis in a carousel reactor. As will be apparent to those of skill in the art upon consideration of Schemes 4 and 5, certain functional groups may require protection during the synthesis (e.g., carboxyl groups as esters, which can be hydrolyzed to acids, or reduced to alcohols).




- 48 -

5. Second Representative Synthesis of Asymmetrically Substituted RCQs5 6. Synthesis of Hydrophilic RCQs


7. General Synthesis of Tetrasubstituted Ethylene-based RCO compounds

Tetrasubstituted ethylene-based compounds, including those illustrated in Table 4, can be prepared according to the following general method. As is known to those of skill in the art the desired tetrasubstituted ethylene compound can be prepared by selecting the appropriate ketone starting material. Examples of suitable ketones include, without limitation, those illustrated below.




10 8. *General Synthesis of RCO Compounds*


The following synthesis can be used to prepare many of the disclosed RCQ compounds, including, without limitation, compounds recited in Tables 3, 5 and 6.




- 50 -

9. Route to RCQ 5110. Route to RCQ 81

5 This example illustrates a specific route using the general method of general Method 8 above.

11. Route to various linker-modified RCQs.

10 The synthesis of various linker-modified RCQs, such as the RCQ 101 compounds recited in Table 3 can be accomplished according to the following general scheme.




15 As will be readily apparent to those of skill in the art upon consideration of the above scheme, the R group may require protection to ensure the synthesis proceeds smoothly.

- 51 -

12. Examples of alternative syntheses for RCQ molecules

An alternative to the nucleophilic displacement method 8 described above, this method uses the chloroquine-linker moiety as the nucleophilic component and the reversal agent derivative as the electrophilic component. As will be apparent to those of skill in the art, this scheme can be used to prepare a variety of disclosed RCQ compounds.



10

D. **Optional Considerations for RCQ Molecules**

In some embodiments RCQ compounds are included in pharmaceutical compositions and/or administered to subjects for the treatment of malaria and/or *Plasmodium sp.* infection. Accordingly, it is advantageous for some RCQ

embodiments to have characteristics suitable for *in vivo* administration (particular formulations are discussed in more detail below). One rule of thumb for drug design is Lipinski's "Rule of 5" (Lipinski *et al.*, *Adv. Drug Deliv. Rev.*, 23(1-3):3-25, 1997), which suggests that absorption and permeation of a drug candidate (which is not a biological transporter substrate) is likely to be increased if at least three of the following parameters is satisfied: (i) there are no more than about 5 H-bond donors (expressed as the sum of hydroxyl and amine groups); (ii) the molecular weight is not substantially greater than 500 g/mole (but can be anything less); (iii) the logP (logarithm of the octanol-water partition coefficient of the drug candidate) is no more than about 5; and (iv) there are no more than about 10 H-bond acceptors (the sum of nitrogens and oxygens).

25

- 52 -

As described above, some RCQ compound embodiments can be 'decorated' by adding polar and/or charged groups. See, for example, RCQ_13, which is drawn above with appropriate amines protonated (as they would be expected to exist in the DV). Similar protonated representations can be made for each of the disclosed RCQ structures.

RCQ_13 illustrates methods to increase water solubility of RCQs without dramatically increasing molecular size or perturbing the CQ end of the molecule. Coincidentally, although not bound by theory, RCQ molecules having at least one (such as two, three or four) protonated amines should target to the parasite acidic DV even more efficiently than CQ because the pH of the DV is lower than the surrounding, favoring the protonated state of each nitrogen, thereby thermodynamically driving the equilibrium toward the interior of the DV.

If desirable, water solubility of an RCQ can be increased by modifying (among other positions) the central nitrogen of the aliphatic portion of a disclosed RCQ (as described above; and see, *e.g.*, RCQ_14). For example, adding methylamine or diethylmethylamine at this exemplary position introduces a positive charge. According to Lipinski's "Rule of 5," such positive charge may increase the likelihood for good absorption and permeation of the molecule. RCQ solubility may be measured under a variety of conditions, *e.g.*, at various pHs and in various solvents. Aqueous solvents (such as, water, normal saline solution or other buffered salt solutions) are useful for solubility determinations. For oral administration of a drug (such as, an RCQ), one optional consideration is the solubility of such drug in the digestive system, which has an acid pH. For example, the stomach has a pH from about 1.5 to about 4 and the small intestine has a pH from about 2.5 to about 5.5. In some embodiments, an RCQ preferably will be soluble in aqueous solution (*e.g.*, water) at a pH of less than about 5.5 (such as, from about pH 1.5 to about pH 4, from about pH 2.5 to about pH 5.5, or from pH 1.5 to about pH 5.0).

E. Optional Characterization of RCQ Functional Properties

The disclosed RCQ compounds are useful, at least, in the treatment of malaria and *Plasmodium sp.* infection. CQ is believed to exert its inhibitory effect on *Plasmodium* parasites by binding heme in the DV of the parasite. Accordingly, if

- 53 -

it is desirable to do so, non-limiting methods useful to functionally characterize RCQ compounds are heme binding assays and *in vitro* and *in vivo Plasmodium sp.* bioassays.

1. Heme Binding

5 Optionally, heme binding of RCQ compounds can be determined by any method known in the art, including, for example, spectrophotometric methods, like UV-VIS spectroscopy or NMR spectroscopy. Methods of measuring heme binding to a variety of molecules using the UV-VIS spectrum have been described (see, *e.g.*, Xu *et al.*, *J. Inorg. Biochem.*, 86(2-3):617-625, 2001; Xu *et al.*, *Antimicrob. Agents*
10 *Chemother.*, 46(1):144-150, 2002). Generally, a heme-binding compound (such as a disclosed RCQ) is mixed in increasing concentration with heme (or a heme model compound, such as porphyrin or metalloporphyrin) and the absorbance spectrum is measured from the ultraviolet region well into the visible region (from about 300 nm to about 750 nm). This spectral range includes the so-called Soret and Q-band
15 regions that arise from porphyrin or metalloporphyrin absorbance. Upon interaction between a heme-binding compound and heme (or a heme model compound, such as porphyrin or metalloporphyrin), decreased absorbance in the Soret region is observed. It is believed that such decrease arises from π - π stacking interactions (White, Aggregation of porphyrins and metalloporphyrins, In: *The Porphyrins*. ed.
20 by Dolphin, New York:Academic Press, 1978, pp. 303-339).

 In particular examples, the formation of heme-RCQ complexes can be determined by UV-VIS spectrophotometric titration, using a variety of commercially available spectrophotometers (such as a Varian-Cary 3E spectrophotometer). The approximate pH of the plasmodial DV, where CQ (or RCQ) is believed to bind to
25 heme *in vivo*, is pH 4.7; thus, it is useful (though not obligatory) to conduct heme-binding reactions at a pH (*e.g.*, 4.7 or 7) and temperature (*e.g.*, 25°C) similar to DV conditions *in vivo*. Titrations of heme (or a heme model) with a RCQ compound can be performed, for example, by successive addition of aliquots of a 1 mM RCQ solution to a 10 μ M heme solution. Optionally, pH can be monitored
30 throughout the procedure to ensure that it remains unchanged. In some circumstances, UV-VIS spectral data can be analyzed digitally, and absorbance readings and concentrations corrected for dilution effects. The amount of

- 54 -

heme-RCQ complex versus amount of RCQ added to the reaction can be plotted. Such titration curves can be analyzed with Hill plot and non-linear curve fitting methods as described previously (Xu *et al.*, *Antimicrob. Agents Chemother.*, 46(1):144-150, 2002) to obtain association constants for the heme-RCQ binding
5 reaction.

NMR is another optional method that can be used, for example, to provide information about stoichiometry, interaction affinities, and structural details of complexes between RCQ molecules and heme (or heme models, such porphyrins or metalloporphyrins) (White, Aggregation of porphyrins and metalloporphyrins, In:
10 *The Porphyrins*. ed. by Dolphin, New York: Academic Press, 1978, pp. 303-339). NMR parameters useful to measure include changes in chemical shift, line-width, and relaxation times. For example, NMR relaxation times give information about proximity to paramagnetic centers (*e.g.*, the Fe in heme), or chemical exchange between species (*e.g.*, bound and free RCQ). A fourth useful parameter to measure
15 is peak area ratio, which is convenient if exchange between complexes is unexpectedly slow. NMR studies can be performed by adding aliquots of heme to RCQ solutions, then monitoring chemical shifts, linewidths, and relaxation times of the RCQ NMR signals. Such techniques are well known in the art (see, for instance, Xu *et al.*, *J. Inorg. Biochem.*, 86(2-3):617-625, 2001).

20 If desirable, fluorescence confocal microscopy can also be used to determine whether an RCQ is localized to the DV (see, for instance, Xu *et al.*, *Mol. Biochem. Parasitol.*, 123(1):47-54, 2002).

2. In vitro and In Vivo Bioassays

One exemplary method for measuring inhibitory effects of RCQ compound
25 *in vitro* has been reported by Smilkstein *et al.* (*Antimicrob. Agents Chemother.*, 48(5):1803-1806, 2004). This method employs fluorescent detection of the level of parasitemia in cultured red blood cells and can be automated for high-throughput screening of antimalarial RCQs. This exemplary method allows the inhibitory agent (such as a RCQ) to be in contact with the parasite for an entire developmental cycle
30 without purine starvation, which is in contrast to the traditional ³H-hypoxanthine method (Desjardins *et al.*, *Antimicrob. Agents Chemother.*, 16(6):710-718, 1979). Briefly, to perform the Smilkstein *et al.* method, an initial parasitemia of

approximately 0.2% is attained by adding uninfected red blood cells (RBCs) to a stock culture of *Plasmodium sp.*-infected RBCs. Infected cells (2% v/v) are combined with RCQs (for example, dissolved in DMSO), *e.g.*, at a final concentration of 10^{-11} to 10^{-4} M. Chloroquine may be added (instead of a RCQ) to some infected RBCs as a control. After a period of incubation (such as about 72 hours), a sufficient amount (*e.g.*, 100 μ L) of a fluorescent DNA binding dye-detergent mixture (*e.g.*, Sybr Green I dye-detergent mixture) to lyse cells and bind to DNA is added to the reactions. Because uninfected RBCs have no DNA, only parasite DNA (in infected RBCs) is available for binding to the fluorescent dye (*e.g.*, Sybr Green I). Samples are then incubated in the dark for a period of time (*e.g.*, one hour), followed by measurement of levels of fluorescence. For Sybr Green I, excitation and emission wavelength are 485 and 530 nm, respectively. Fluorescence readings, which represent the amount of parasite DNA in the sample, can be plotted against the Log[drug] and fit to a curve by nonlinear regression to obtain the IC₅₀ value for the tested compound.

Various *Plasmodium sp.* parasites are available for testing *in vitro* efficacy of antimalarial compounds (such as a disclosed RCQ). It is expected that the disclosed RCQ will be effective against both CQ^R and CQ^S *Plasmodium sp.* strains (such as CQ^R and CQ^S *P. falciparum* strains). Exemplary CQ^S and CQ^R *P. falciparum* strains are shown in the following table:

Table 11. Drug Susceptibility of *P. falciparum* strains

Drug	<i>P. falciparum</i> strain					
	D6	FCR3	W2	Dd2	3D7	TM91C235
CQ	S	R	R	R	S	R
Quinine	S	S	R	R	NC	R
Mefloquine	S/R	S	S	R	NC	R
Pyrimethamine	S	S	R	R	NC	R
Cycloguanil	S	S	R	S	NC	NC
Sulfadoxine	S	S	R	S	NC	R

S=sensitive; R=resistant; S/R=intermediate level of resistance; NC=not characterized

- 56 -

Antimalarial compounds can also be tested for efficacy *in vivo*. One non-limiting exemplary method for performing such testing is the 4-day suppressive test (Peters, *Ann. Trop. Med. Parasitol.*, 69(2):155-171, 1975). Briefly, such test involves collecting *P. chabaudi* (Mackinnon and Read, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 359:965-986, 2004; Mackinnon *et al.*, *Exp. Parasitol.*, 101(2-3):121-128, 2002) from a donor animal harboring ~20% parasitemia. Male CF-1 mice at 4-5 weeks of age (~20 g) are infected with ~10⁷ infected red blood cells *via* 100 µL tail vein injection and randomly sorted into groups of 4 mice each on day 0. One hour after infection, the mice receive the test agent *via* oral gavage, or by the intraperitoneal route (at approximately 1-10 times the *in vitro* IC₅₀ values). Drugs are administered once daily for at least 4 days beginning at day 0. Animals are tested weekly for levels of parasitemia, for example, by microscopic analysis of Giemsa-stained blood smears. Efficacy of a tested compound in this bioassay is measured by inhibition of parasite growth as defined previously. In some examples, subjects are expected to be freed of measurable parasites after a period of treatment (such as 30 days).

IV. *Pharmaceutical Compositions*

The disclosed RCQ compounds are useful, at least, for the treatment of malaria and/or inhibiting the growth of malarial parasites, such as *P. falciparum*. Accordingly, pharmaceutical compositions comprising at least one disclosed RCQ compound are also described herein.

Formulations for pharmaceutical compositions are well known in the art. For example, *Remington's Pharmaceutical Sciences*, by E. W. Martin, Mack Publishing Co., Easton, PA, 15th Edition, 1975, describes exemplary formulations (and components thereof) suitable for pharmaceutical delivery of disclosed RCQ compounds. Pharmaceutical compositions comprising at least one of the disclosed RCQ compounds can be formulated for use in human or veterinary medicine. Particular formulations of a disclosed pharmaceutical composition may depend, for example, on the mode of administration (*e.g.*, oral or parenteral) and/or on the location of the infection to be treated (*e.g.*, liver-stage and/or blood-stage malaria parasites). In some embodiments, formulations include a pharmaceutically

- 57 -

acceptable carrier in addition to at least one active ingredient, such as a RCQ compound. In other embodiments, other medicinal or pharmaceutical agents, for example, with similar, related or complementary effects on the affliction being treated (such as malaria or *Plasmodium sp.* infection), can also be included as active ingredients in a pharmaceutical composition.

Pharmaceutically acceptable carriers useful for the disclosed methods and compositions are conventional in the art. The nature of a pharmaceutical carrier will depend on the particular mode of administration being employed. For example, parenteral formulations usually comprise injectable fluids that include pharmaceutically and physiologically acceptable fluids such as water, physiological saline, balanced salt solutions, aqueous dextrose, glycerol or the like as a vehicle. For solid compositions (e.g., powder, pill, tablet, or capsule forms), conventional non-toxic solid carriers can include, for example, pharmaceutical grades of mannitol, lactose, starch, or magnesium stearate. In addition to biologically neutral carriers, pharmaceutical compositions to be administered can optionally contain minor amounts of non-toxic auxiliary substances (e.g., excipients), such as wetting or emulsifying agents, preservatives, and pH buffering agents and the like; for example, sodium acetate or sorbitan monolaurate. Other non-limiting excipients include, nonionic solubilizers, such as cremophor, or proteins, such as human serum albumin or plasma preparations.

The disclosed pharmaceutical compositions may be formulated as a pharmaceutically acceptable salt of a disclosed RCQ compound. Pharmaceutically acceptable salts are non-toxic salts of a free base form of a compound that possesses the desired pharmacological activity of the free base. These salts may be derived from inorganic or organic acids. Non-limiting examples of suitable inorganic acids are hydrochloric acid, nitric acid, hydrobromic acid, sulfuric acid, hydriodic acid, and phosphoric acid. Non-limiting examples of suitable organic acids are acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, methyl sulfonic acid, salicylic acid, formic acid, trichloroacetic acid, trifluoroacetic acid, gluconic acid, asparagic acid, aspartic acid,

- 58 -

benzenesulfonic acid, p-toluenesulfonic acid, naphthalenesulfonic acid, and the like. Lists of other suitable pharmaceutically acceptable salts are found in *Remington's Pharmaceutical Sciences*, 17th Edition, Mack Publishing Company, Easton, PA, 1985. A pharmaceutically acceptable salt may also serve to adjust the osmotic pressure of the composition.

The dosage form of a disclosed pharmaceutical composition will be determined by the mode of administration chosen. For example, in addition to injectable fluids, topical or oral dosage forms may be employed. Topical preparations may include eye drops, ointments, sprays and the like. Oral formulations may be liquid (e.g., syrups, solutions or suspensions), or solid (e.g., powders, pills, tablets, or capsules). Methods of preparing such dosage forms are known, or will be apparent, to those skilled in the art.

Certain embodiments of the pharmaceutical compositions comprising a disclosed RCQ compound may be formulated in unit dosage form suitable for individual administration of precise dosages. The amount of active ingredient (e.g., RCQ compound) administered will depend on the subject being treated, the severity of the affliction, and the manner of administration, and is known to those skilled in the art. Within these bounds, the formulation to be administered will contain a quantity of the extracts or compounds disclosed herein in an amount effective to achieve the desired effect in the subject being treated.

V. *Methods of Use*

The present disclosure includes methods of treating malaria and/or inhibiting *Plasmodium sp.* parasite growth (including CQ^R and CQ^S parasites). In some examples, the pathogen is *P. falciparum*, *P. vivax*, *P. ovale*, *P. malariae*, or a combination thereof. In other examples, the disease or *Plasmodium sp.* to be treated is resistant to CQ or other traditional malaria treatment.

Disclosed methods includes administering a disclosed RCQ compounds (and, optionally, one or more other pharmaceutical agents) to a subject in a pharmaceutically acceptable carrier and in an amount effective to treat malaria or *Plasmodium sp.* infection. The treatment can be used prophylactically in any subject in a demographic group at substantial risk for such diseases; for example, subjects

- 59 -

who are traveling to areas where malaria (such as CQ^R malaria) is endemic (including, *e.g.*, Southeast Asia, Africa, Papua New Guinea, Indonesia, Thailand, and India). Notably, pregnant women are twice as likely to attract malaria-carrying mosquitoes as non-pregnant women (perhaps due to a greater volume of exhaled air and a warmer skin surface), and, therefore, are especially vulnerable to malaria. Alternatively, subjects can be selected using more specific criteria, such as a probable or definitive diagnosis of malaria or *Plasmodium sp.* infection based on, for example, clinical signs and symptoms and/or laboratory evidence of parasite infection. An example of such a subject would be a person who presents clinically with symptoms resembling the flu (including periods of chills and fever lasting several hours and occurring every few days). In more severe cases, an infected subject may present with enlarged spleen and/or liver, anemia, and jaundice. Other subjects may be identified based on positive tests for parasite-specific proteins, including plasmodial histidine rich protein-2 (HRP-2) or parasite-specific lactate dehydrogenase (pLDH) or parasite DNA. A number of antibodies specific for *Plasmodium* parasites are available and are useful for diagnostic immunoassays or immunofluorescence techniques. PCR can also be used to diagnosis malaria in a subject (*Am. J. Trop. Med. Hyg.*, 65(4):355-363, 2001).

Routes of administration useful in the disclosed methods include but are not limited to oral and parenteral routes, such as intravenous (iv), intraperitoneal (ip), rectal, topical, ophthalmic, nasal, and transdermal. Formulations for these dosage forms are described above.

An effective amount of a RCQ will depend, at least, on the particular method of use, the subject being treated, the severity of the affliction, and the manner of administration of the therapeutic composition. A “therapeutically effective amount” of a composition is a quantity of a specified compound sufficient to achieve a desired effect in a subject (host) being treated. For example, this may be the amount of a RCQ necessary to prevent, inhibit, reduce or relieve *Plasmodium sp.* infection and/or one or more symptoms of malaria in a subject. Specific anti-*Plasmodium sp.* effects that may result from RCQ administration are described herein. Ideally, a therapeutically effective amount of a RCQ is an amount sufficient to prevent, inhibit, reduce or relieve *Plasmodium sp.* infection and/or one or more symptoms of

- 60 -

5 malaria without causing a substantial cytotoxic effect on host cells. It is anticipated that disclosed RCQs will be well tolerated in humans because, at least some, quinoline analogs (such as CQ) and reversal agents (such as Imipramine) have been demonstrated safe for administration to humans in combination therapies (albeit at much higher dosages of reversal agent).

Therapeutically effective doses (or growth inhibitory amounts) of a disclosed RCQ compound or pharmaceutical composition can be determined by one of skill in the art, with a goal of achieving local (*e.g.*, tissue) concentrations that are at least as high as the IC_{50} of the applicable compound disclosed in the examples herein. An
10 example of a dosage range is from about 0.1 to about 200 mg/kg body weight orally in single or divided doses. In particular examples, a dosage range is from about 1.0 to about 100 mg/kg body weight orally in single or divided doses, including from about 1.0 to about 50 mg/kg body weight, from about 1.0 to about 25 mg/kg body weight, from about 1.0 to about 10 mg/kg body weight (assuming an average body
15 weight of approximately 70 kg; values adjusted accordingly for persons weighing more or less than average). For oral administration, the compositions are, for example, provided in the form of a tablet containing from about 50 to about 1000 mg of the active ingredient, particularly about 75 mg, about 100 mg, about 200 mg, about 400 mg, about 500 mg, about 600 mg, about 750 mg, or about 1000 mg of the
20 active ingredient for the symptomatic adjustment of the dosage to the subject being treated. In one exemplary oral dosage regimen, a tablet containing from about 500 mg to about 1000 mg active ingredient is administered once (*e.g.*, a loading dose) followed by administration of 1/2 dosage tablets (*e.g.*, from about 250 to about 500 mg) each 6 to 24 hours for at least 3 days.

25 The specific dose level and frequency of dosage for any particular subject may be varied and will depend upon a variety of factors, including the activity of the specific RCQ compound, the metabolic stability and length of action of that compound, the age, body weight, general health, sex and diet of the subject, mode and time of administration, rate of excretion, drug combination, and severity of the
30 condition of the host undergoing therapy.

The present disclosure also contemplates combinations of one or more disclosed RCQs with one or more other agents or therapies useful in the treatment of

- 61 -

malaria and/or *Plasmodium sp.* parasitemia. For example, one or more disclosed RCQs may be administered in combination with effective doses of other medicinal and pharmaceutical agents, or in combination other non-medicinal therapies, such as hormone or radiation therapy. The term "administration in combination with" refers to both concurrent and sequential administration of the active agents. In some examples, the one or more other antimalarial agents or therapies include artesunate and mefloquine (either individually or in an artesunate-mefloquine combination), or sulfadoxine and pyrimethamine (either individually or in a sulfadoxine-pyrimethamine combination (commercially available as Fanisdar™)). In particular examples, the one or more other antimalarial agents or therapies have at least one different mode of action than is proposed (although not binding) for a disclosed RCQ; thus, for instance, a combination agent or therapy may target mitochondria and/or dihydrofolate reductase.

Other method embodiments involve inhibiting the growth of at least one *Plasmodium sp.* (such as *P. falciparum*, *P. vivax*, *P. ovale*, or *P. malariae*, or a combination thereof) by contacting the parasite(s) with a growth inhibitory amount of a disclosed RCQ. Inhibition of *Plasmodium sp.* growth was defined above. Contact between a RCQ and a parasite may occur *in vitro* (such as in culture conditions) or *in vivo* (such as in a subject infected with at least one *Plasmodium sp.*). In certain method embodiments, growth inhibitory amounts include amounts described in the Examples (for example, LC₅₀ concentrations). In other examples, a growth inhibitory amount is from about 1 nM to about 1 μM of a disclosed RCQ (such as from about 5 nM to about 50 nM, or from about 5 nM to about 25 nM).

EXAMPLES


The following examples are provided to illustrate certain particular features and/or embodiments. These examples should not be construed to limit the invention to the particular features or embodiments described.

- 62 -

Example 1**[RCQ_02] IS HIGHLY EFFECTIVE AGAINST CQ^R and CQ^S*****P. FALCIPARIUM***

This Example illustrates the synthesis of an exemplary RCQ, referred to as RCQ_02 (formally, N'-(7-Chloro-quinolin-4-yl)-N-[3-(10,11-dihydro-dibenzo[b,f]azepin-5-yl)-propyl]-N-methyl-propane-1,3-diamine). RCQ_02 has a 7-chloroquinoline CQ core with the reversal agent, imipramine, substituted for the diethylamino end of the chloroquine side chain. Imipramine was selected as the reversal agent moiety in RCQ_02 because imipramine is synthetically (in fact, commercially) available, symmetric, simple, and conforms to a published reversal agent pharmacophore (Bhattacharjee *et al.*, *J. Chem. Inf. Comput. Sci.*, 42(5):1212-1220, 2002), having the two hydrophobic rings and an aliphatic chain terminating in a positively charged nitrogen. CQ analogs with 3 and 4 carbons in the aliphatic chain are known (*e.g.*, De *et al.*, *J. Med. Chem.*, 41(25):4918-4926, 1998; De *et al.*, *Am. J. Trop. Med. Hyg.*, 55(6):579-583, 1996; Ramanathan-Girish *et al.*, *Int. J. Toxicol.*, 23(3):179-189, 2004), and the 3-carbon methylene analog, as well as significantly longer methylene chain molecules have activity against CQ^R *P. falciparum*.

RCQ_02 was synthesized as follows:



Reactions were monitored by thin layer chromatography and/or ^1H nuclear magnetic resonance spectroscopy (NMR) as appropriate. All products were fully characterized by ^1H NMR, and were deemed to be >95% pure after column chromatography using an alumina matrix and ethyl acetate/hexane (30:70 by volume) as the eluting solvent. Reaction yield was approximately 10%; however,

25

- 63 -

higher yields would be expected with minor modifications of the reaction conditions, including isolation of intermediates or varying reaction temperature, reactant concentrations and/or reaction solvent(s). Such minor modifications are easily performed by those of ordinary skill in the art.

5

Example 2

RCQ_02 AND CQ HAVE SUBSTANTIALLY SIMILAR ELECTROSTATIC POTENTIALS AND ELECTRON DENSITIES, AND RCQ_02 AND RCQ_31 EFFECTIVELY BIND HEME

10 This Example illustrates that the covalent coupling of a reversal agent (*e.g.*, imipramine) to CQ does not substantially affect the electrostatic potential or electron density of the CQ moiety, nor affect the ability of the hybrid molecule to interact with heme.

The GAUSSIAN-03 program (which is publicly available at the website
15 having the address www.gaussian.com) was used to ascertain the charge distribution around CQ and the CQ moiety of RCQ_02. Calculations were performed at the B3LYP/6-311+g(d) level of theory, and results were visualized with the program gOpenMol (which is publicly available at the website having the address www.csc.fi/gopenmol).

20 FIG. 1 shows that the electrostatic potentials and electron densities are nearly identical for the CQ portion of each of these molecules. The same result was obtained for deprotonated quinoline ring nitrogens. Because the electrical properties of chloroquine are believed to underlie its binding to heme, the results shown in FIG. 1 strongly suggested that RCQ_02 would also bind with heme or with
25 hemozoin.


Heme binding by RCQ_02, in fact, was demonstrated by measuring the optical spectrum of heme in the absence and presence of increasing amounts of RCQ_02 under conditions (*e.g.*, pH 4.7) similar to *Plasmodium sp.* DV, as previously described (Xu *et al.*, *J. Inorg. Biochem.*, 86(2-3):617-625, 2001; Xu *et al.*, *Antimicrob. Agents Chemother.*, 46(1):144-150, 2002). Briefly, 2-10 μ l aliquots
30 of a 1 mM RCQ_02 solution (pH 4.8) were successively added to a 10 μ M heme solution (pH 4.8) in a 1 cm (1-3 ml) quartz cuvette. The initial volume of heme

- 64 -

solution in the cuvette was 2000 μ l. pH was monitored to ensure that it remained unchanged throughout the procedure. The UV-Vis spectrum (from 200 nm to 800 nm) was measured at each concentration of RCQ-02 using a Varian-Cary 3E spectrophotometer with a temperature-controlled cell holder. All UV-VIS spectral data were analyzed digitally, and absorbance readings and concentrations were corrected for dilution effects.

FIG. 2 shows the expected lowering of the Soret band, which is known to accompany binding or complexation of heme by an aromatic molecule (such as CQ or a RCQ). This interaction is thought to involve π - π stacking of the molecules (e.g., Shelnutt, *Inorg. Chem.*, 22:2535–2544, 1983).

RCQ_02 titration curves were with Hill plot and non-linear curve fitting methods as described previously (Xu *et al.*, *Antimicrob. Agents Chemother.*, 46(1):144-150, 2002). With the assumption that the heme is dimeric (μ -oxo), the following equilibrium expression was applied to the RCQ/heme reaction:



Accordingly, a $\text{heme}_2 \cdot \text{RCQ_02}$ association constant of approximately $7.5 \times 10^5 \text{ M}^{-1}$ at pH 5.7 as calculated, which is comparable to (albeit slightly lower than) the association constant reported for $\text{CQ} \cdot \text{heme}_2$ ($K \sim 4 \times 10^5 \text{ M}^{-1}$ at pH 5.7; Dorn *et al.*, *Biochem. Pharmacol.*, 55(6):727-736, 1998; Xu *et al.*, *J. Inorg. Biochem.*, 86(2-3):617-625, 2001).

Similar methodologies as described for RCQ_02 were used to measure a heme association constant for RCQ_31 of $5 \times 10^5 \text{ M}^{-1}$.

Example 3

RCQ_02 IS HIGHLY EFFECTIVE AGAINST CQ^R AND CQ^S *P. FALCIPARUM* STRAINS

This Example demonstrates that RCQ_02 is a highly effective inhibitor of the growth of CQ^R and CQ^S *P. falciparum* strains *in vitro*.

An *in vitro* fluorescence assay for determining IC₅₀ of antimalarial drugs was performed as previously described by Smilkstein *et al.* (*Antimicrob. Agents Chemother.*, 48(5):1803-1806, 2004). This relatively low-cost procedure can be automated for high-throughput screening of the antimalarial activities of RCQs

- 65 -

described herein. Advantageous, this method also allows the tested agent (e.g, a RCQ) to be in contact with the parasite for an entire developmental cycle without purine starvation (which occurs with the commonly used ^3H -hypoxanthine method; see, e.g., Desjardins *et al.*, *Antimicrob. Agents Chemother.*, 16,(6):710-7188, 1979).

5 Briefly, an initial parasitemia of approximately 0.2% was attained by addition of uninfected red blood cells to a stock culture of cells infected with Dd2 (CQ^R) or D6 (CQ^S) *P. falciparum* strains. A 10 mM RCQ_02 solution (in DMSO) was prepared. RCQ_02 (at a final concentration from 10^{-11} to 10^{-4} M) and infected red blood cells (at a final concentration of 2% (v/v)) were mixed in 100 μL samples in individual wells of a 96-well plate. Each sample was prepared in triplicate. A triplicate set of reactions containing CQ (instead of RCQ_02) was included as a control. After a 72 hour incubation period, Sybr Green I dye-detergent mixture (100 μL) was added to each reaction. The samples were further incubated for an hour in the dark, and then placed in a 96-well fluorescence plate reader (Gemini-EM, Molecular Diagnostics) and analyzed with excitation and emission wavelength bands at 485 and 530 nm, respectively. Fluorescence readings were plotted against the Log[drug], and curve fitting performed by nonlinear regression to find the IC₅₀ value.

FIG. 3A shows the effect of CQ concentrations on the growth of D6 and Dd2 *P. falciparum* strains. FIG. 3B shows the effect of RCQ_02 concentrations on the growth of D6 and Dd2 *P. falciparum* strains. IC₅₀ values were calculated from these data, as described above, and are summarized in FIG. 3C and in the following Table 12:

Table 12. IC₅₀ Values for RCQ_02 and CQ Against CQ^R and CQ^S *P. falciparum*

Molecule	<i>P. falciparum</i> cell line	Characteristic	IC ₅₀ (nM)
RCQ_02	D6	CQ ^S	2.9
CQ	D6	CQ ^S	6.5
RCQ_02	Dd2	CQ ^R	5.3
CQ	Dd2	CQ ^R	102

- 66 -

FIG. 3 and Table 12 demonstrate that RCQ_02 has an *in vitro* activity that is better than CQ against either CQ^R or CQ^S *P. falciparum*. In fact, RCQ_02 is as, or nearly as, active against CQ^R and CQ^S *P. falciparum* as some of the most effective drugs reported to date, including bisquinoline derivatives (Vennerstrom *et al.*, *J. Med. Chem.*, 41(22):4360-4364, 1998; Vennerstrom *et al.*, *J. Med. Chem.*, 35(11):2129-2134, 1992; Basco *et al.*, *Am. J. Trop. Med. Hyg.*, 50(2):200-205, 1994).

As compared to CQ, RCQ_02 had a lower heme binding constant and a lower IC₅₀ against CQ^R and CQ^S *P. falciparum*. Taken together, these findings suggest at least one non-binding theory of RCQ_02 action, which follows: RCQ_02 may obtain higher intra-DV concentrations (than does CQ) by the reversal agent moiety reducing the efflux of the RCQ_02 molecule from the DV. Higher intra-DV concentrations of RCQ_02 may permit more heme to be retained in the DV (as mediated by the CQ moiety of the RCQ), which further thwarts *Plasmodium sp.* heme-detoxification mechanism and thereby inhibits parasite growth. Since low-level CQ efflux occurs even in CQ^S malaria (Krogstad *et al.*, *Science*, 238(4831):1283-1285, 1987; Sanchez *et al.*, *Biochem.*, 43(51):16365-16373, 2004; Sanchez *et al.*, *Biochem.*, 42(31):9383-9394, 2003), RCQ_02 (and RCQs, in general) may be more effective than CQ even against CQ^S strains by inhibiting low-level drug efflux in the CQ^S strains.

Example 4

OTHER RCQs ALSO ARE HIGHLY EFFECTIVE AGAINST CQ^R AND CQ^S *P. FALCIPARUM* STRAINS

This Example demonstrates that numerous of the disclosed RCQs strongly inhibit the growth of CQ^R and CQ^S *P. falciparum* strains. Using the same methods as described in Example 3, the IC₅₀ values for *P. falciparum* growth inhibition for a variety of RCQs were determined and are summarized below:


Table 13. Growth Inhibition of Various Strains by Exemplary RCQs

Molecule	IC ₅₀ (nM)		
	D6 (CQ ^S)	Dd2 (CQ ^R)	W2 (CQ ^R)
CQ	6.9	102	250
RCQ_02	2.9	5.3	

- 67 -

Molecule	IC ₅₀ (nM)		
	D6 (CQ ^S)	Dd2 (CQ ^R)	W2 (CQ ^R)
RCQ_05(desMe)	36	27	
RCQ_22	45	79	
RCQ_26	7.6	10.6	
RCQ_31	1.0	3.6	4.3
RCQ_32	2.6	3.4	3.4
RCQ_33	8.1	9.3	9.3
RCQ_34	2.4	3.7	2.5
RCQ_35	4.0	6.0	4.0
RCQ_36	11.7	98	
RCQ_37F2	15	19	
RCQ_42	2.1	1.8	
RCQ_72	9.2	9.6	
RCQ_73	4.8	4.1	
RCQ_74	15	9.8	
RCQ_76	21.1	15.3	
RCQ_351	141	164	
Q_38a	280	400	

The compound named Q_38a has the following structure:



- This Example validates the effectiveness of using hybrid molecules (RCQs) such as those described herein to impair the growth of CQ^R and CQ^S *Plasmodium* sp. parasites. All of the RCQs tested inhibited the growth of CQ^S strains near to (or significantly less than) the effective concentration of CQ. In contrast to CQ, however, each of the foregoing RCQs advantageously also inhibited the growth of at least one (and in some cases, two) CQ^R strains.
- The results in this Example and Examples 2 and 3, indicate (among other things) that (i) K_{assoc} for heme binding is not simply correlated with IC₅₀; (ii) RCQs work nearly as well against CQ^R *P. falciparum* strains (e.g., W2 or Dd2) as they do against CQ^S strains (e.g., D6); (iii) many RCQs are significantly better than CQ even against CQ^S *P. falciparum* (e.g., D6) by as much as a factor of nearly 7;
- (iv) substitution of Cl at the p-position of an aromatic ring at the reversal agent end

- 68 -

of an RCQ molecule has no negative effect on the activity of the molecule (compare RCQ_34 to RCQ_35); thus, comparable substitutions in a variety of RCQs should be well tolerated; and (v) a piperazine ring system in the linker between the reversal agent and quinoline ring is one exemplary structural feature that appears to decrease
5 the IC₅₀ of the resultant compound.

Example 5

RCQs ARE NON-TOXIC TO MOUSE SPLEEN LYMPHOCYTES

This Example illustrates that a variety of RCQs have no significant
10 cytotoxicity when introduced into mouse spleen lymphocytes cultures *in vitro*.

RCQ cytotoxicity was determined against mitogen-induced murine spleen lymphocytes *in vitro* using the Alamar Blue assay (as described by Ahmed *et al.*, *J. Immunol. Meth.*, 170(2):211-224, 1994). Mouse spleen lymphocytes were isolated from C57B1/6J mice by teasing tissues on a metallic sieve screen. Cells thus
15 obtained were washed in RPMI 1640 media, then resuspended in complete RPMI media containing 10% FBS, 50 µg/ml penicillin/streptomycin, 50 µM β-mercaptoethanol, and 1 µM/mL concanavalin A for 1 minute. Two hundred (200) µl isolated cells were then seeded in separate wells of a 96-well flat-bottom tissue culture plate at a density of 2x10⁵ cells per well. Test drug was added to the
20 individual wells at concentrations of 0 to 62 µM. After 72 hours incubation in a humidified atmosphere at 37°C and 5% CO₂, a solution of resazurin PBS was added to a final concentration 10 µM, and the plates were returned to the incubator for another 24 hours. Resazurin is a substrate which changes color in response to metabolic activity (such as, living cells). The fluorescence of the cell-containing
25 samples was measured with a Gemini EM plate reader with excitation at 560 nM and emission at 590 nM. LC₅₀ values were determined from plots of fluorescence versus drug concentration using non-linear regression by Prism software.

As shown in the following table, the tested RCQs had cytotoxicity values in the range of 700 nM (*e.g.*, RCQ_02, RCQ_22, and RCQ_73) to 5300 nM (*e.g.*,
30 RCQ_36).

- 69 -

Table 14. Cytotoxicity of Exemplary RCQs

Molecule	Cytotoxicity (nM)	Therapeutic Index	
		D6	Dd2
CQ	12400	1797	122
RCQ_02	700	240	132
RCQ_22	700	16	9
RCQ_31	1900	1900	528
RCQ_32	1400	538	412
RCQ_33	1400	173	151
RCQ_34	1100	458	297
RCQ_35	800	200	133
RCQ_36	5300	453	54
RCQ_37F2	4100	273	215
RCQ_42	1100	523	611
RCQ_72	2500	272	260
RCQ_73	700	145	171
RCQ_74	900	60	92
RCQ_76	2500	118	163
Q_38a	43000	153	108

Although chloroquine (CQ) had a higher cytotoxic concentration than all of the tested RCQs, the *in vitro* therapeutic index (ratio of cytotoxicity/IC₅₀ for *P. falciparum*) for each of the RCQs was still quite large. Thus, the toxic dose for nearly all of the tested RCQs is hundreds of times higher or even thousands of times higher (for RCQ_31) than the therapeutic dose. These results indicate that there is a good margin of safety for each RCQ in the treatment of *Plasmodium* parasites.

10

Example 6**RCQs ARE NOT READILY UPTAKEN BY CNS NEUROTRANSMITTER TRANSPORTERS**

Some known reversal agents, such as imipramine, have been reported to have neurological side effects. Although some side effects are acceptable (in quantity and/or quality) for a drug that otherwise effectively treats a disease of concern, such as malaria, this Example demonstrates that tested RCQs do not effectively compete with the natural ligands of the human serotonin (hSERT), dopamine (HEK-hDAT), and norepinephrine (hNET) transporters. Thus, such compounds are expected to have minimum impact on the *in vivo* function of such transporters and, correspondingly, few related side effects.

20

- 70 -

Selected RCQs were tested for inhibition of [^3H]neurotransmitter uptake by the recombinant human serotonin (hSERT), dopamine (HEK-hDAT), and norepinephrine (hNET) transporters by a modification of the method of Eshleman *et al.* (*J. Pharmacol. Exp. Ther.*, 289(2):877-885, 1999). HEK-hDAT, hSERT, and –

5 hNET expressing cells were grown on tissue culture dishes. After removing the medium and washing the plates with Ca^{2+} -, Mg^{2+} -free phosphate buffered saline (PBS), fresh medium was added and the plates were placed in a 25°C water bath for 5 minutes. Cells were gently lifted from the plates and cell clusters separated by trituration. 50 μL aliquots of the suspended cells were added to assay tubes

10 containing drugs and Krebs-HEPES assay buffer in a final volume of 0.5 mL.

Competition experiments were conducted in triplicate. After a 10 minute preincubation period at 25°C, [^3H]neurotransmitter (20 nM final concentration; [^3H]DA, [^3H]5-HT, or [^3H]NE) were added, and the mixtures incubated for 10 minutes. Reactions were terminated by filtration through Wallac filtermat A filters

15 (presoaked in 0.05% polyethylenimine), using a Tomtec cell harvester. Scintillation fluid was added to each filtered spot, and radioactivity remaining on the filters determined using a Wallac 1205 Betaplate scintillation counter.

Specific uptake was defined as the difference in uptake observed in the absence and presence of 5 μM mazindol (hDAT and hNET) or 5 μM imipramine

20 (hSERT). Inhibition of uptake was determined by non-linear regression using Prism software.

Table 15. Inhibition of Neurotransmitter Uptake by Exemplary RCQs

Molecule	Neurotransmitter Transporter Uptake (nM)		
	hSERT	HEK-hDAT	hNET
RCQ 02	230	2200	350
RCQ 73	1160	1400	660
RCQ 76	965	1080	550

IC₅₀ values for representative RCQs neurotransmitter transporter uptake are

25 shown in the above table. These values are orders of magnitude higher than the IC₅₀ for these same compounds against even CQ^R *P. falciparum* parasites (see Example 4). This indicates that significant neurological side effects may not be an issue for the disclosed RCQs, even those that have imipramine-like moieties.

Example 7**RCQs CLEAR PARASITEMIA *IN VIVO***

This Example demonstrates the *in vivo* effectiveness of a representative RCQs in the treatment of *Plasmodium* sp. parasitemia.

- 5 A variation of the 4-day suppressive test described by Peters (*Ann. Trop. Med. Parasitol.*, 69(2):155-171, 1975) was used to test the *in vivo* effectiveness of RCQ_02 as a treatment for *Plasmodium* sp. infection. Briefly, *P. chabaudi* parasites (Mackinnon and Read, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 359:965-986, 2004) were collected from a donor animal harboring approximately 20% parasitemia.
- 10 Sixteen (16) male CF-1 mice at 4-5 weeks of age (approximately 20 g) were infected with approximately 10^7 infected red blood cells via 100 μ L tail vein injection. The infected mice were randomly sorted into groups of 4 mice each on day 0. One hour after infection, each group of four mice received 64, 32, 16, or 0 mg/kg RCQ_02 via oral gavage once per day for four days. After the treatment course, the animals were
- 15 weighed and blood collected for the determination of parasite burden via direct microscopic analysis of Giemsa-stained blood smears.

- Mice treated with 64 mg/kg RCQ_02 were all cleared of parasites to >99% by day 4. In addition, none of the treated mice showed any adverse side effects from RCQ_02 (*e.g.*, weight loss, reduced eating, reduced activity, or ruffled ungroomed
- 20 fur). This result was obtained even though RCQ_02 has relatively low water solubility; nonetheless, RCQ_02 was sufficiently orally available to produce a malaria cure *in vivo* without apparent adverse side effects.


- While this disclosure has been described with an emphasis upon particular embodiments, it will be obvious to those of ordinary skill in the art that variations of
- 25 the particular embodiments may be used and it is intended that the disclosure may be practiced otherwise than as specifically described herein. Accordingly, this disclosure includes all modifications encompassed within the spirit and scope of the disclosure as defined by the following claims:

- 72 -


We claim:

1. An antimalarial compound having the formula:

5



wherein Q represents



X_1 and X_2 are independently alkoxy, optionally substituted amino, halo, haloalkyl or hydroxy;

10

n is from 0 to 2;


m is from 0 to 4;

R_1 is H, alkyl, heteroalkyl, alkylamino, sulfonyl, haloalkyl, or carbonyl;

L is a hydrocarbon chain containing from 1 to 12 carbon atoms, optionally substituted, branched or both and optionally interrupted by one or more heteroatom;

15

wherein G is



R_2 is H, alkyl, heteroalkyl, alkylamino, haloalkyl, alkylsulfonamide, amide, alkylamide, cyano, furyl, or alkylfuran; L_1 and L_2 are independently alkyl or heteroalkyl;

20

Y and Y_1 independently represent N; CH; CH_2CH or $C(O)CH$;

R_3 and R_4 are independently cycloalkyl, heterocyclyl, aryl, heteroaryl;


L_1 and L_2 are independently alkyl or heteroalkyl; and

ring 1 is a five-, six-, or seven-membered heterocyclic ring.


25

2. The compound of claim 1, having the formula:

- 73 -



3. The compound of claim 2, having the formula




5

wherein X_1 is a halogen;

X represents $-N-$ or $-CH-$;


Y_1 represents $-N-$; $-CH-$; $-CH_2CH-$ or $-C(O)CH-$ and u is from 0 to 2.

4. The compound of claim 2, having the formula



10

5. The compound of claim 2, having the formula




15

6. The compound of claim 2, wherein L_1 is a hydrocarbon chain having from 2 to 6 carbon atoms.

- 74 -


7. The compound of claim 2, wherein L_1 is an ethyl or propyl chain.

8. The compound of claim 2, having the formula




5

9. The compound of claim 2, having the formula



10

10. The compound of claim 2, having the formula




11. The compound of claim 1, wherein R_3 and R_4 together with Y, Y_1 or Y'_1 form a tricyclic ring system.


15

12. The compound of claim 1, having the formula


- 75 -



13. The compound of claim 1, having the formula:



5 wherein G is



$(X_1)_m$ and $(X_2)_n$ are independently H, halo, haloalkyl, amino, hydroxyl, alkoxy, alkylamino, or arylamino;

m is 0 to 4;

10 n is 0 to 2;

R_1 and R_2 are independently H, alkyl, heteroalkyl, alkylamino, haloalkyl, or alkylsulfonamide;

L_1 and L_2 are independently alkyl or heteroalkyl;

Y_1 is carbon or nitrogen;

15 Y'_1 is carbon;

R_3 and R_4 are independently aryl, or heteroaryl, or together with Y_1 or Y'_1 form a tricyclic ring system; and

- 76 -

Ring 1 is a five- to six-membered heterocyclic ring.

14. The compound of claim 1, wherein m is 1 or 2; n is 0 to 1; and $(X_1)_m$ and $(X_2)_n$ are independently H, chloro, fluoro, or trifluoromethyl.

5

15. The compound of claim 13, wherein $(X_1)_m$ are independently H, chloro, or trifluoromethyl; and $(X_2)_n$ are H.

16. The compound of claim 15, wherein m is 1 and $(X_1)_1$ is chloro at position C₇.

10

17. The compound of claim 13, wherein R₁ and R₂ are independently H, alkyl having from 1 to 4 carbon atoms, or heteroalkyl having from 1 to 4 carbon atoms.

15

18. The compound of claim 17, wherein R₁ and R₂ are independently H, methyl, trifluoromethyl, -CH₂CH₂CF₃, -CH₂CF₃, -CH₂CH₂NH₂, -CH₂NH₂, -CH₂CH₂C(O)NH₂, or -CH₂C(O)NH₂.

19. The compound of claim 1, wherein L₁ and L₂ are independently from 1 to 12 atoms in length, and each position of L₁ and L₂ is independently methylene, difluoromethylene, dichloromethylene, di(methylamino), -NH-, or -O-.

20


20. The compound of claim 1, wherein m is 1; $(X_1)_m$ is chloro at position C₇; $(X_2)_n$ are H; R₁ and R₂ are independently H, methyl, trifluoromethyl, -CH₂CH₂CF₃, -CH₂CF₃, -CH₂CH₂NH₂, -CH₂NH₂, -CH₂CH₂C(O)NH₂, or -CH₂C(O)NH₂; and L₁ and L₂ are independently from 1 to 12 atoms in length, and each position of L₁ and L₂ is independently methylene, difluoromethylene, dichloromethylene, di(methylamino), -NH-, or -O-.

25

30

21. The compound of claim 1, having the formula:


- 77 -



wherein R_1 and R_2 are independently H, alkyl, or heteroalkyl; L_1 and L_2 are independently alkyl or heteroalkyl; Y_1 is carbon or nitrogen; and R_3 and R_4 are independently aryl, or heteroaryl, or together with Y_1 form a tricyclic ring system.

5

22. The compound of claim 21, wherein R_3 and R_4 independently have the formula:



where $(X_3)_s$ are independently H, halo, haloalkyl, amino, hydroxyl, alkylamino, cyano, alkoxy, sulfonamide, mercapto, or keto; s is 0 to 4; Y_2 is C, S or N; and L_3 is alkyl, heteroalkyl, -N-, -O-, or -S-.

10

23. The compound of claim 22, wherein $(X_3)_s$ are independently H, fluoro, chloro, trifluoromethyl, methoxy, mercapto, or keto; and s is 1 to 2.

15

24. The compound of claim 22, wherein Y_2 is nitrogen.

25. The compound of claim 22, wherein L_3 is methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, -N-, or -O-.

20

26. The compound of claim 22, wherein s is 1 to 2; $(X_3)_s$ are independently H, fluoro, chloro, trifluoromethyl, methoxy, mercapto, or keto; Y_2 is nitrogen; and L_3 is methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, -N-, or -O-.

25

- 78 -


27. The compound of claim 22, wherein R_3 and R_4 are the same.

28. The compound of claim 27, wherein the compound is selected from RCQ_05, RCQ_06, RCQ_07, RCQ_08, RCQ_09, RCQ_10, RCQ_11, RCQ_12, and
5 RCQ_104.

29. The compound of claim 22, wherein R_3 and R_4 are different.


30. The compound of claim 29, wherein the compound is selected from
10 RCQ_14, RCQ_101, RCQ_102, RCQ_103, RCQ_105, and RCQ_150.

31. The compound of claim 1, having the formula




15 wherein Ar^1 and Ar^2 are independently selected aryl groups.

32. The compound of claim 31, having the formula




20 33. The compound of claim 31, having the formula

- 79 -



34. The compound of claim 11, wherein the tricyclic ring system comprises a central ring and two peripheral rings, having a formula selected from:



5

wherein Y_2 is N, O, CH, CH_2 or S; R_{13} is H, alkyl, heteroalkyl, =O, amino, amine, amide, sulfonamide, halo, cyano, hydroxy, mercapto, haloalkyl, alkoxy, alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, or alkylsulfano; $(X_4)_q$ and $(X_5)_t$ are independently H, alkyl, halo, haloalkyl, alkoxy, amino, hydroxyl, alkylamino, cyano, or mercapto; q and t are independently 0 to 4 as valence requirements permit; and each of peripheral rings can independently include at least one heteroatom at any position that valence requirements permit.

10

35. The compound of claim 34, wherein q and t are independently 0 to 2; $(X_4)_q$ and $(X_5)_t$ are independently H, methyl, fluoro, chloro, trifluoromethyl, methoxy, or cyano; Y_2 is N or S; and R_{13} is H or =O.

15


36. The compound of claim 34, wherein Y_1 is nitrogen; Y_2 is CH, CH_2 or S; R_{13} is H or =O; $(X_4)_q$ and $(X_5)_t$ are independently H, fluoro, chloro, trifluoromethyl, cyano, or alkoxy; and q and t are independently 0 to 2.

20

- 80 -

37. The compound of claim 34, wherein the compound is selected from RCQ_1, RCQ_2, RCQ_3, RCQ_4, and RCQ_100.

38. The compound of claim 1, having the formula:




5

wherein m is 1; $(X_1)_m$ is chloro at position C_7 ; $(X_2)_n$ are each H; R_1 and R_2 are independently H, alkyl, or heteroalkyl; L_1 is alkyl or heteroalkyl; Y'_1 is carbon; and R_3 and R_4 are independently aryl, or heteroaryl, or together with Y'_1 form a tricyclic ring system.

10

39. The compound of claim 38, wherein R_3 and R_4 independently have the formula:



where $(X_3)_s$ are independently H, halo, haloalkyl, amino, hydroxyl, alkylamino, cyano, alkoxy, sulfonamide, mercapto, or keto; s is 0 to 4; Y_2 is CH, S or N; and L_3 is alkyl, heteroalkyl, -N-, -O-, or -S-.

15

40. The compound of claim 39, wherein s is 1 to 2; $(X_3)_s$ are independently H, fluoro, chloro, trifluoromethyl, methoxy, mercapto, or keto.

20


41. The compound of claim 41, wherein Y_2 is nitrogen.

42. The compound of claim 39, wherein L_3 is methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, -N-, or -O-.

- 81 -

43. The compound of claim 39, wherein s is 1 to 2; $(X_3)_s$ are independently H, fluoro, chloro, trifluoromethyl, methoxy, mercapto, or keto; Y_2 is nitrogen; and L_3 is methyl, ethyl, methoxy, ethoxy, methylamino, ethylamino, -N-,
 5 or -O-.

44. The compound of claim 13, wherein Y'_1 , R_3 , and R_4 form a tricyclic ring system having a formula selected from:



10 wherein Y_2 is NR_{14} , O, or S; R_{13} is H, alkyl, heteroalkyl, =O, amino, amine, amide, sulfonamide, halo, cyano, hydroxy, mercapto, haloalkyl, alkoxy, alkylthio, thioalkoxy, arylalkyl, heteroaryl, alkylamino, dialkylamino, or alkylsulfano; $(X_4)_q$ and $(X_5)_t$ are independently H, alkyl, halo, haloalkyl, alkoxy, amino, hydroxyl, alkylamino, cyano, or mercapto; q and t are independently 0 to 4 as valence
 15 requirements permit; R_{14} is H, lower alkyl or acyl and each peripheral ring can independently include at least one heteroatom at any position that valence requirements permit.


45. The compound of claim 44, wherein Y_2 is C or N; R_{13} is H or =O;
 20 $(X_4)_q$ and $(X_5)_t$ are independently H, fluoro, chloro, trifluoromethyl, cyano, or alkoxy; and q and t are independently 0 to 2.

46. The compound of claim 45, wherein the compound is selected from RCQ_110, RCQ_111, RCQ_112, RCQ_113, RCQ_114, RCQ_115, RCQ_116, and
 25 RCQ_117.

- 82 -

47. A compound comprising an antimalarial quinoline analog covalently linked to a CQ^R reversal moiety.

5 48. The compound of claim 47 having the formula:



wherein G is a CQ^R reversal moiety; m is 0 to 4; n is 0 to 2; (X₁)_m and (X₂)_n are independently H, fluoro, chloro, or trifluoromethyl; R₁ is H, methyl, trifluoromethyl, -CH₂CH₂CF₃, -CH₂CF₃, -CH₂CH₂NH₂, -CH₂NH₂, -CH₂CH₂C(O)NH₂, or
 10 -CH₂C(O)NH₂; and L₁ is alkyl or heteroalkyl.

49. The compound of claim 48, wherein m is 1; (X₁)_m is chloro at position C₇; (X₂)_n are H; and L₁ is from 1 to 12 atoms in length, and each position of L₁ is independently methylene, difluoromethylene, dichloromethylene),
 15 di(methylamino), -NH-, or -O-.

50. The compound of claim 43, wherein the antimalarial quinoline analog is chloroquine.







20 51. The compound of claim 50, wherein the C4 dimethylamine side chain of chloroquine is covalently linked to the CQ^R reversal moiety.

52. The compound of claim 47, wherein the CQ^R reversal moiety is amitriptyline, amlodipine, azatadine, chlorpheniramine, citalopram, cyclosporine,
 25 cyproheptadine, cyproheptadine, desipramine, diethyl-{3-[3-(4-methoxy-benzylidene)-pyrrolidin-1-yl]-propyl}-amine, erythromycin, fantofarone, fluoxetine, haloperidol, icajine, imipramine, isoretuline, ivermectin, ketotefin, ketotifen, nomifensine, NP30: C₉H₁₉-Phenyl-(O-CH₂CH₂)₃₀OH, oxaprotiline, probenecid,



















- 83 -

progesterone, promethazine, strychnobrasiline, BG958, trifluoperazine, verapamil, or WR 268954.












53. The compound of claim 1, having a formula selected from
























- 84 -


























- 85 -


















- 86 -















- 87 -




- 88 -




- 89 -



- 90 -



- 91 -



54. A pharmaceutical composition, the composition comprising a therapeutically effective amount of the compound of claim 1, 13, 21, 31, 34, 38, 47 or 53; and a pharmaceutically acceptable carrier.

- 92 -

55. The pharmaceutical composition of claim 54 further comprising at least one additional antimalarial therapeutic agent.

56. The pharmaceutical composition of claim 55, wherein the at least one additional antimalarial therapeutic agent is artesunate, mefloquine, sulfadoxine, or pyrimethamine, or combinations thereof.

57. A method for inhibiting or treating malaria in a subject, comprising administering to a subject a therapeutically effective amount of the compound of claim 1, 13, 21, 31, 34, 38, 47 or 53.

58. The method of claim 57, wherein the compound is administered prophylactically.

59. The method of claim 57, wherein the therapeutically effective amount is from about 1 to about 50 mg/kg administered at least one time.

60. The method of claim 57, wherein at least one of the causes of the malaria is *P. falciparum* infection.

61. A method for inhibiting the growth of *Plasmodium sp.*, comprising contacting at least one *Plasmodium sp.* parasite with a growth inhibitory amount of the compound of claim 1, 13, 21, 31, 34, 38, 47 or 53.

62. The method of claim 61, wherein the *Plasmodium sp.* is *P. falciparum*, *P. vivax*, *P. ovale*, or *P. malariae*, or a combination thereof.

63. The method of claim 62, wherein the *Plasmodium sp.* is *P. falciparum*.

64. The method of claim 61, wherein the growth inhibitory amount is from about 1 to about 50 nM.

FIG. 1




FIG. 2






FIG. 3

A



B



C




FIG. 4

