



(86) Date de dépôt PCT/PCT Filing Date: 1994/07/21

(87) Date publication PCT/PCT Publication Date: 1995/02/02

(45) Date de délivrance/Issue Date: 2002/07/02

(85) Entrée phase nationale/National Entry: 1996/01/19

(86) N° demande PCT/PCT Application No.: US 1994/008335

(87) N° publication PCT/PCT Publication No.: 1995/003330

(30) Priorité/Priority: 1993/07/21 (08/095,760) US

(51) Cl.Int.<sup>6</sup>/Int.Cl.<sup>6</sup> C07K 14/655, A61K 38/31, A61K 51/08,  
A61K 38/08, C07K 1/13, C07K 7/06

(72) Inventeurs/Inventors:  
McBride, William, US;  
Dean, Richard T., US

(73) Propriétaire/Owner:  
DIATIDE, INC., US

(74) Agent: MBM & CO.

(54) Titre : DERIVES DE LA SOMATOSTATINE ET LEURS PRODUITS RADIOMARQUES

(54) Title: RADIOACTIVE PEPTIDES

(57) **Abrégé/Abstract:**

This invention relates to therapeutic reagents and peptides, radiodiagnostic agents and peptides, and methods for producing labelled radiodiagnostic agents. Specifically, the invention relates to linear peptides derivatives and analogs of somatostatin, and embodiments of such peptides radiolabelled with a radioisotope, as well as methods and kits for making, radiolabelling and using such peptides for radiodiagnostic and radiotherapeutic purposes. This invention specifically relates to linear peptide derivatives and analogues of somatostatin radiolabelled with technetium-99m and used thereof as scintigraphic imaging agents. The invention also specifically relates to linear peptide derivatives and analogues of somatostatin radiolabelled with cytotoxic isotopes such as rhenium-186 (<sup>186</sup>Re) and rhenium-188 (<sup>188</sup>Re) for use as radiotherapeutic agents. Methods and kits for making, radiolabelling and using peptidesdiagnostically and therapeutically in a mammalian body are also provided.

## RADIOACTIVE PEPTIDES

2167678

### Abstract:

This invention relates to therapeutic reagents and peptides, radiodiagnostic agents and peptides, and methods for producing labelled radiodiagnostic agents. Specifically, the invention relates to linear peptides derivatives and analogs of somatostatin, and embodiments of such peptides radiolabelled with a radioisotope, as well as methods and kits for making, radiolabelling and using such peptides for radiodiagnostic and radiotherapeutic purposes. This invention specifically relates to linear peptide derivatives and analogues of somatostatin radiolabelled with technetium-99m and used thereof as scintigraphic imaging agents. The invention also specifically relates to linear peptide derivatives and analogues of somatostatin radiolabelled with cytotoxic isotopes such as rhenium-186 ( $^{186}\text{Re}$ ) and rhenium-188 ( $^{188}\text{Re}$ ) for use as radiotherapeutic agents. Methods and kits for making, radiolabelling and using peptides diagnostically and therapeutically in a mammalian body are also provided.



## Radioactive Peptides

### BACKGROUND OF THE INVENTION

#### 5      1.      Field of the Invention

10                      This invention relates to therapeutic agents and peptides, radiotherapeutic agents and peptides, radi. diagnostic agents and peptides, and methods for producing such labeled radiodiagnostic and radiotherapeutic agents. Specifically, the invention relates to linear peptide derivatives and analogues of somatostatin, and embodiments of such peptides labeled with gamma radiation-emitting radioisotopes such as technetium-99m (Tc-99m), as well as methods and kits for making, radiolabeling and using such peptides to image sites in a mammalian body. The invention also relates to linear peptide derivatives and analogues of somatostatin labeled with cytotoxic radioisotopes such as rhenium-186 (<sup>186</sup>Re) and rhenium-188 (<sup>188</sup>Re), and methods and  
15                      kits for making, radiolabeling and using such peptides therapeutically in a mammalian body.

#### 2.      Description of the Prior Art

20                      Somatostatin is a tetradecapeptide that is endogenously produced by the hypothalamus and pancreas in humans and other mammals. The peptide has the formula:

##### Formula I

25                      Ala-Gly-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Thr-Ser-Cys



30                      (Single letter abbreviations for amino acids can be found in G. Zubay, Biochemistry (2d ed.), 1988, (MacMillan Publishing: New York), p.33). This peptide exerts a wide variety of biological effects *in vivo*. It is known to act physiologically on the central nervous system, the hypothalamus, the pancreas, and the gastrointestinal tract.

35                      Somatostatin inhibits the release of insulin and glucagon from the pancreas, inhibits growth hormone release from the hypothalamus, and reduces gastric secretions. Thus, somatostatin has clinical and therapeutic applications for the

alleviation of a number of ailments and diseases, both in humans and other animals. Native somatostatin is of limited utility, however, due to its short half-life *in vivo*, where it is rapidly degraded by peptidases. For this reason, somatostatin analogues having improved *in vivo* stability have been developed in the prior art.

5 Freidinger, U.S. Patent No. 4,235,886 disclose cyclic hexapeptide somatostatin analogues useful in the treatment of a number of diseases in humans.

Coy and Murphy, U.S. Patent No. 4,485,101 disclose synthetic dodecapeptide somatostatin analogues.

10 Freidinger, U.S. Patent No. 4,611,054 disclose cyclic hexapeptide somatostatin analogues useful in the treatment of a number of diseases in humans.

Nutt, U.S. Patent No. 4,612,366 disclose cyclic hexapeptide somatostatin analogues useful in the treatment of a number of diseases in humans.

Coy *et al.*, U.S. Patent No. 4,853,371 disclose synthetic octapeptide somatostatin analogues.

15 Coy and Murphy, U.S. Patent No. 4,871,717 disclose synthetic heptapeptide somatostatin analogues.

Coy *et al.*, U.S. Patent No. 4,904,642 disclose synthetic octapeptide somatostatin analogues.

20 Taylor *et al.*, U.S. Patent No. 5,073,541 disclose a method of treating small cell lung cancer.

Brady, European Patent Application No. 83111747.8 discloses dicyclic hexapeptide somatostatin analogues useful in the treatment of a number of human diseases.

25 Bauer *et al.*, European Patent Application No. 85810617.2 disclose somatostatin derivatives useful in the treatment of a number of human diseases.

Eck and Moreau, European Patent Application No. 90302760.5 disclose therapeutic octapeptide somatostatin analogues.

Coy and Murphy, European Patent Application Serial No. 90304551.6 disclose linear somatostatin analogues.

30 Coy and Murphy, International Patent Application Serial No. PCT/US90/07074 disclose somatostatin analogues for therapeutic uses.

Schally *et al.*, European Patent Application Serial No. EPA 911048445.2

disclose cyclic peptides for therapeutic use.

Bodgen and Moreau, International Patent Application Serial No. PCT/US92/01027 disclose compositions and methods for treating proliferative skin disease.

5 Somatostatin exerts its effects by binding to specific receptors expressed at the cell surface of cells comprising the central nervous system, the hypothalamus, the pancreas, and the gastrointestinal tract. These high-affinity somatostatin binding sites have been found to be abundantly expressed at the cell surface of most endocrine-active tumors arising from these tissues. Expression of high-affinity binding sites for  
10 somatostatin is a marker for these tumor cells, and specific binding with somatostatin can be exploited to locate and identify tumor cells *in vivo*.

Methods for radiolabeling somatostatin analogues that have been modified so as to contain a tyrosine amino acid (Tyr or Y) are known in the prior art.

15 Albert *et al.*, UK Patent Application 892725.3 disclose radioimaging using somatostatin derivatives such as octreotide labeled with  $^{123}\text{I}$ .

Bakker *et al.*, 1990, J. Nucl. Med. 31: 1501-1509 describe radioactive iodination of a somatostatin analog and its usefulness in detecting tumors *in vivo*.

Bakker *et al.*, 1991, J. Nucl. Med. 32: 1184-1189 teach the usefulness of radiolabeled somatostatin for radioimaging *in vivo*.

20 Bomanji *et al.*, 1992, J. Nucl. Med. 33: 1121-1124 describe the use of iodinated (Tyr-3) octreotide for imaging metastatic carcinoid tumors.

Alternatively, methods for radiolabeling somatostatin by covalently modifying the peptide to contain a radionuclide-chelating group have been disclosed in the prior art.

25 Albert *et al.*, UK Patent Application 8927255.3 disclose radioimaging using somatostatin derivatives such as octreotide labeled with  $^{111}\text{In}$  via a chelating group bound to the amino-terminus.

30 Albert *et al.*, European Patent Application No. WO 91/01144 disclose radioimaging using radiolabeled peptides related to growth factors, hormones, interferons and cytokines and comprised of a specific recognition peptide covalently linked to a radionuclide chelating group.

Albert *et al.*, European Patent Application No. 92810381.1 disclose

somatostatin peptides having amino-terminally linked chelators.

Faglia *et al.*, 1991, J. Clin. Endocrinol. Metab. 73: 850-856 describe the detection of somatostatin receptors in patients.

5 Kwekkeboom *et al.*, 1991, J. Nucl. Med. 32: 981 Abstract #305 relates to radiolabeling somatostatin analogues with  $^{111}\text{In}$ .

Albert *et al.*, 1991, Abstract LM10, 12th American Peptide Symposium: 1991 describe uses for  $^{111}\text{In}$ -labeled diethylene-triaminopentaacetic acid-derivatized somatostatin analogues.

10 Krenning *et al.*, 1992, J. Nucl. Med. 33: 652-658 describe clinical scintigraphy using ( $^{111}\text{In}$ )(DTPA)octreotide.

These methods can be readily adapted to enable detection of tumor cells *in vivo* by radioimaging, based on the expression of high affinity binding sites for somatostatin on tumor cells. Radionuclides which emit gamma radiation can be readily detected by scintigraphy after injection into a human or an animal. A variety  
15 of radionuclides are known to be useful for radioimaging, including  $^{67}\text{Ga}$ ,  $^{68}\text{Ga}$ ,  $^{99\text{m}}\text{Tc}$  (Tc-99m),  $^{111}\text{In}$ ,  $^{123}\text{I}$  or  $^{125}\text{I}$ . The sensitivity of imaging methods using radioactively-labeled peptides is much higher than other techniques known in the art, since the specific binding of the radioactive peptide concentrates the radioactive signal over the cells of interest, for example, tumor cells. This is particularly important for  
20 endocrine-active gastrointestinal tumors, which are usually small, slow-growing and difficult to detect by conventional methods. Labeling with technetium-99m (Tc-99m) is advantageous because the nuclear and radioactive properties of this isotope make it an ideal scintigraphic imaging agent. Tc-99m has a single photon energy of 140 keV and a radioactive half-life of about 6 hours, and is readily available from a  
25  $^{99}\text{Mo}$ - $^{99\text{m}}\text{Tc}$  generator. Other radionuclides have effective half-lives which are much longer (*for example*,  $^{111}\text{In}$ , which has a half-life of 60-70 h) or are toxic (*for example*,  $^{125}\text{I}$ ). Although Tc-99m is an ideal radiolabeling reagent, it has not been widely used in the art prior to the present invention (*see, for example*, Lamberts, J. Nucl. Med. 32: 1189-1191 (1991)).

30 Somatostatin and radiolabeled somatostatin analogues can also be used therapeutically. For these applications, cytotoxic radioisotopes are advantageous, such as scandium-47, copper-67, gallium-72, yttrium-90, iodine-125, iodine-131,

samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211, and bismuth-212. The rhenium isotopes  $^{186}\text{Re}$  and  $^{188}\text{Re}$  are particularly advantageous.

5 The use of chelating agents for radiolabeling proteins are known in the prior art, and methods for labeling peptides Tc-99m are disclosed in U.S. Patent Nos. 5,225,180 and 5,620,675 and PCT International Applications WO92/13572, WO93/10747, WO93/17719, WO93/21962, WO93/23085, WO94/00489 and WO94/07918.

10

Fritzberg, U.S. Patent No. 4,444,690 describes a series of technetium-  
15 chelating agents based on 2,3-bis(mercaptoacetamido) propanoate.

Gansow *et al.*, U.S. Patent No. 4,472,509 teach methods of manufacturing and purifying Tc-99m chelate-conjugated monoclonal antibodies.

Reno and Bottino, European Patent Application 87300426.1 disclose radiolabeling antibodies with Tc-99m.

20 Pak *et al.*, European Patent Application No. WO 88/07382 disclose a method for labeling antibodies with Tc-99m.

Cox, International Patent Application WO92/21383 discloses radiolabeled somatostatin derivatives containing two cysteine residues.

25 Rhodes, 1974, Sem. Nucl. Med. 4: 281-293 teach the labeling of human serum albumin with technetium-99m.

Khaw *et al.*, 1982, J. Nucl. Med. 23: 1011-1019 disclose methods for labeling biologically active macromolecules with Tc-99m.

Byrne and Tolman, *supra*, disclose a bifunctional thiolactone chelating agent for coupling Tc-99m to biological molecules.

30 Cox *et al.*, 1991, Abstract, 7th International Symposium on Radiopharmacology, p. 16, disclose the use of, Tc-99m-,  $^{131}\text{I}$ - and  $^{111}\text{In}$ -labeled somatostatin analogues in biolocalization of endocrine tumors *in vivo* by

scintigraphy.

5 Methods for directly labeling somatostatin, derivatives of somatostatin, analogues of somatostatin or peptides that bind to the somatostatin receptor and contain at least 2 cysteine residues that form a disulfide or wherein the disulfide is reduced to the sulfhydryl form, are disclosed in co-pending U.S. Patent Application Serial No. 07/807,062, now U.S. Patent No. 5,225,180, issued July 6, 1993.

10 There remains a need for synthetic (to make routine manufacture practicable and to ease regulatory acceptance) somatostatin analogues having increased *in vivo* stability, to be used therapeutically, as scintigraphic agents when radiolabeled with Tc-99m or other detectable radioisotopes for use in imaging tumors *in vivo*, and as radiotherapeutic agents when radiolabeled with a cytotoxic radioisotope such as rhenium-188. Small synthetic somatostatin analogues are provided by this invention that specifically fulfill this need.

15

### SUMMARY OF THE INVENTION

20 The present invention provides somatostatin analogues that are linear peptides for therapeutic applications, including radiotherapeutic applications, and diagnostic applications, including radiodiagnostic applications, in particular scintigraphic imaging applications. Distinct from native somatostatin and somatostatin analogues known in the prior art, the linear peptides of the invention are not constrained within a cyclic structure. The invention also provides linear peptide reagents comprised of the linear peptide somatostatin analogues of the invention, wherein such peptides are covalently linked to a radiolabel-binding moiety. The invention provides such linear  
25 peptides, linear peptide reagents and radiolabeled linear peptide reagents that are scintigraphic imaging agents, radiodiagnostic agents and radiotherapeutic agents. Scintigraphic imaging agents of the invention comprise linear peptide reagents radiolabeled with a radioisotope, preferably technetium-99m. Radiotherapeutic agents of the invention comprise linear peptide reagents radiolabeled with a cytotoxic radioisotope, preferably rhenium-186 or rhenium-188. Methods for making and  
30 using such linear peptides, linear peptide reagents and radiolabeled embodiments thereof are also provided.

The present invention also provides scintigraphic imaging agents comprised of a linear peptide that is a somatostatin analogue and that is labeled with iodine-123, iodine-125 or iodine-131. Similarly, the invention provides alternative embodiments of the linear somatostatin peptide analogues radiolabeled with iodine-125, iodine-131 or astatine-211 for use as therapeutic agents.

The somatostatin analogues provided by the invention are somatostatin-receptor binding peptides having the following formula:

Formula II



wherein  $X^1$  is a hydrophilic moiety which is not greater than 1500 Daltons in formula weight;  $A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, S-alkylated cysteine, penicillamine (Pen), homocysteine (Hcy) or homohomocysteine (Hhc; 3-mercaptopropyl) glycine;  $B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-2-naphthylalanine (Nal), or 2-amino-indane-2-carboxylic acid (Ain) or substituted derivatives thereof;  $B^2$  is D- or L-Trp or substituted derivatives thereof;  $B^3$  is D- or L-Lys, or homolysine (Hly), 4-amino-cyclohexylalanine (Achxa), 4-aminomethyl-phenylalanine (Amf), S-(2-aminoethyl)cysteine (Aec), S-(3-aminopropyl)cysteine (Apc), O-(2-aminoethyl) serine (Aes), O-(3-aminopropyl)serine (Aps) or substituted derivatives thereof;  $B^4$  is Thr, Ser, Val, Phe, Leu, Ile, 2-amino-isobutyric acid (Aib), 2-aminobutyric acid (Abu), norvaline (Nva), or norleucine (Nle);  $C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof;  $X^2$  is a hydrophilic moiety which is not more than 1500 Daltons in formula weight. In a preferred embodiment,  $X^1$  is a hydrophilic moiety that comprises an amino acid, or a peptide having an amino acid sequence of no more than 10 residues, or a monosaccharide, or an oligosaccharide comprising 10 or fewer saccharide units, or a poly(N-carboxyalkyl)amine, or a polyoxyanion. In another preferred embodiment,  $X^2$  is a hydrophilic moiety that comprises a poly(N-carboxyalkyl)amine or polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than 10 residues (including peptides wherein the carboxyl group of the carboxyl-terminal amino acid is reduced to an alcohol), or a monosaccharide or an oligosaccharide comprising 10 or fewer saccharide units. In another preferred

embodiment, B<sup>1</sup> is phenylalanine or tyrosine, B<sup>2</sup> is tryptophan, most preferably D-tryptophan, B<sup>3</sup> is lysine and B<sup>4</sup> is threonine or valine.

The invention also provides linear peptide reagents comprising a linear peptide of Formula II covalently linked to a radiolabel-binding moiety, wherein X<sup>1</sup> is H, lower alkyl or substituted alkyl, aryl or substituted aryl, alkanoyl or substituted alkanoyl, aroyl or substituted aroyl, or a hydrophilic moiety which is not greater than 1500 Daltons in formula weight; A<sup>1</sup>, A<sup>2</sup> and C<sup>1</sup> are each independently a lipophilic D- or L-amino acid, S-alkylated cysteine, penicillamine (Pen), homocysteine (Hcy) or homohomocysteine (Hhc; 3-mercaptopropyl) glycine; B<sup>1</sup> is D- or L-Phe, or D- or L-Tyr, or D- or L-2-naphthylalanine (Nal), or 2-amino-indane-2-carboxylic acid (Ain) or substituted derivatives thereof; B<sup>2</sup> is D- or L-Trp or substituted derivatives thereof; B<sup>3</sup> is D- or L-Lys, or homolysine (Hly), 4-amino-cyclohexylalanine (Achxa), 4-aminomethyl-phenylalanine (Amf), S-(2-aminoethyl)cysteine (Aec), S-(3-aminopropyl)cysteine (Apc), O-(2-aminoethyl) serine (Aes), O-(3-aminopropyl)serine (Aps) or substituted derivatives thereof; B<sup>4</sup> is Thr, Ser, Val, Phe, Leu, Ile, 2-amino-isobutyric acid (Aib), 2-aminobutyric acid (Abu), norvaline (Nva), or norleucine (Nle); C<sup>2</sup> is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof; X<sup>2</sup> is -COOR<sup>9</sup>, -CH<sub>2</sub>OH, CH<sub>2</sub>COOR<sup>9</sup>, or -CON(R<sup>9</sup>)<sub>2</sub>, where each R<sup>9</sup> is independently H, lower linear or cyclic alkyl or substituted derivatives thereof, or substituted with a hydrophilic moiety which is not more than 1500 Daltons in formula weight. In a preferred embodiment, when X<sup>1</sup> is a hydrophilic moiety that moiety comprises an amino acid, or a peptide having an amino acid sequence of no more than 10 residues, or a monosaccharide, or an oligosaccharide comprising 10 or fewer saccharide units, or a poly(N-carboxyalkyl)amine, or a polyoxyanion. In another preferred embodiment, when X<sup>2</sup> is a hydrophilic moiety that moiety comprises a poly(N-carboxyalkyl)amine or polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than 10 residues (including peptides wherein the carboxyl group of the carboxyl-terminal amino acid is reduced to an alcohol), or a monosaccharide or an oligosaccharide comprising 10 or fewer saccharide units. In another preferred embodiment, B<sup>1</sup> is phenylalanine or tyrosine, B<sup>2</sup> is tryptophan, most preferably D-tryptophan, B<sup>3</sup> is lysine and B<sup>4</sup> is threonine or valine.

The present invention provides peptides that are linear somatostatin peptide analogues as described herein having increased *in vivo* stability compared with native somatostatin, and that are therapeutically useful in the alleviation of diseases or other ailments in humans or other animals.

5           The invention also provides scintigraphic imaging agents comprising the linear peptide reagents of the invention wherein the radiolabel-binding moiety is stably complexed with a radioisotope. In one such embodiment is provided a scintigraphic imaging agent wherein the linear somatostatin peptide analogue reagents of the invention are radiolabeled with technetium-99m. In other embodiments of the  
10           scintigraphic imaging agents of the invention the radioisotope is indium-111 or gallium-68. In still other embodiments, the scintigraphic imaging agents of the invention are linear peptides that are radiolabeled with iodine-123 or iodine-125.

          The invention also provides radiotherapeutic agents that are the linear peptide reagents of the invention radiolabeled with a cytotoxic radioisotope that is selected  
15           from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211 and bismuth-212. In preferred embodiments, the radioisotope is rhenium-186 or rhenium-188. In additional preferred embodiments, the cyclic peptides of the  
20           invention are radiolabeled with iodine-125, iodine-131 or astatine-211.

          In another embodiment, the invention provides therapeutic agents comprising the linear somatostatin analogue peptide reagents of the invention complexed with a non-radioactive metal such as rhenium. Combination embodiments, wherein such a complex is also radiolabeled, either directly or *via* a radiolabel-binding moiety, are  
25           also provided by the invention and are within its scope.


          The invention also provides pharmaceutical compositions comprising the somatostatin receptor-binding peptides of the invention in a pharmaceutically acceptable carrier.

          The invention also provides a method for alleviating somatostatin-related  
30           diseases in animals, preferably humans, comprising administering a therapeutically effective amount of the somatostatin analogues of the invention to the animal. In preferred embodiments, the amount of the somatostatin analogue administered is

from about 0.1 to about 50 mg/kg body weight/day.

It is an advantage of the somatostatin analogues provided by this invention that the peptides retain high affinity for somatostatin receptors even though they are linear peptides. As the preferred embodiments lack intramolecular disulfide bonding, the advantageous feature of the linear somatostatin peptide analogues of this invention is that their stability is not dependent on the formation or persistence of intramolecular disulfide bonds. This feature is in turn advantageous because the high affinity of the peptides of this invention for somatostatin receptors is thus not a function of the integrity of labile intramolecular crosslinks such as disulfide bonds. Additionally, the peptide reagents of the invention retain their high affinity for somatostatin receptors after being subjected to radiolabeling *via* covalently-linked radiolabel binding moieties. In contrast, for example, Tc-99m conjugation to a Tc-99m binding moiety covalently linked to native somatostatin, or to a somatostatin analogue having a disulfide bond, can result in reduction of the disulfide accompanied by a loss of biological activity. Such loss of biological activity can also occur *in vivo* using native somatostatin, or to any somatostatin analogue having a disulfide bond. The present invention is not subject to similar losses in biological activity *in vivo* because the somatostatin analogues of the invention are active as linear peptides.

A first aspect of the reagents provided by the invention for preparing radiolabeled agents of the invention are reagents, each comprised of a peptide that is a somatostatin analogue that is covalently linked to a radiolabel-binding moiety having formula:



wherein  $(pgp)^S$  is H or a thiol protecting group and (aa) is any  $\alpha$ - or  $\beta$ -amino acid. In a preferred embodiment, the amino acid is glycine. In another preferred embodiment, the agent is a scintigraphic imaging agent. In yet another preferred embodiment, the agent is a radiotherapeutic agent.

In a second embodiment, the invention provides peptide reagents capable of being radiolabeled for use as scintigraphic imaging agents or radiotherapeutic agents, each comprising a somatostatin analogue that is covalently linked to a radiolabel-binding moiety of formula:



wherein A is H, HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC or R''''; B is H, SH or -NHR''', -N(R''')-(amino acid or peptide) or R''''; X is SH or -NHR''', -N(R''')-(amino acid or peptide) or R''''; Z is H or R''''; R', R'', R''' and R'''' are independently H or straight or branched chain or cyclic lower alkyl; n is 0, 1 or 2; and: (1) where B is -NHR''' or -N(R''')-(amino acid or peptide), X is SH and n is 1 or 2; (2) where X is -NHR''' or -N(R''')-(amino acid or peptide), B is SH and n is 1 or 2; (3) where B is H or R'''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC, X is SH and n is 0 or 1; (4) where A is H or R'''', then where B is SH, X is -NHR''' or -N(R''')-(amino acid or peptide) and where X is SH, B is -NHR''' or -N(R''')-(amino acid or peptide); (5) where X is H or R'''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC and B is SH; (6) where Z is methyl, X is methyl, A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC and B is SH and n is 0; and (7) where Z is SH and X is SH, n is not 0; and wherein the thiol moiety is in the reduced form.

Preferred embodiments of this radiolabel-binding moiety have a chemical formula that is:




wherein (amino acid)<sup>1</sup> and (amino acid)<sup>2</sup> are each independently any primary  $\alpha$ - or  $\beta$ -amino acid that does not comprise a thiol group, Z is a thiol-containing moiety that is cysteine, homocysteine, isocysteine, penicillamine, 2-mercaptoethylamine or 3-mercaptopropylamine, and R<sup>1</sup> is lower (C<sup>1</sup>-C<sup>4</sup>) alkyl, an amino acid or a peptide comprising 2 to 10 amino acids. When Z is cysteine, homocysteine, isocysteine or penicillamine, the carbonyl group of said moiety is covalently linked to a hydroxyl group, a NR<sup>3</sup>R<sup>4</sup> group, wherein each of R<sup>3</sup> and R<sup>4</sup> are independently H or lower (C<sup>1</sup>-C<sup>4</sup>) alkyl, an amino acid or a peptide comprising 2 to 10 amino acids; or




wherein Y is a thiol-containing moiety that is cysteine, homocysteine, isocysteine, penicillamine, 2-mercaptoacetate or 3-mercaptopropionate, (amino acid)<sup>1</sup> and (amino acid)<sup>2</sup> are each independently any primary  $\alpha$ - or  $\beta$ -amino acid that does not comprise a thiol group, and R<sup>2</sup> is H or lower (C<sup>1</sup>-C<sup>4</sup>) alkyl, an amino acid or a peptide

comprising 2 to 10 amino acids. When Y is cysteine, homocysteine, isocysteine or penicillamine, the amino group of said moiety is covalently linked to -H, an amino acid or a peptide comprising 2 to 10 amino acids; or



or


III.



IV.

wherein n, m and p are each integers that are independently 0 or 1; each R' is independently H, lower alkyl, C<sub>2</sub>-C<sub>4</sub> hydroxyalkyl, or C<sub>2</sub>-C<sub>4</sub> alkoxyalkyl, and each R is independently H or R'', where R'' is substituted or unsubstituted lower alkyl or phenyl not comprising a thiol group, and one R or R' is L, where L is a functionality covalently linked to the somatostatin receptor binding peptide.

In particular embodiments of this aspect of the invention, the radiolabel-binding moiety has a formula that is:




$-\{A-CZ(B)-\{C(R^1R^2)\}_n-X\}-(\text{amino acid})^1-(\text{amino acid})^2,$   
 $-(\text{a primary } \alpha,\omega\text{- or } \beta,\omega\text{-diamino acid})-(\text{amino acid})^1-\{A-CZ(B)-\{C(R^1R^2)\}_n-$   
 $X\}, \text{ or}$

$-\{A-CZ(B)-\{C(R^1R^2)\}_n-X\}-(\text{amino acid})^1-(\text{a primary } \alpha,\omega\text{- or } \beta,\omega\text{-diamino acid})$   
 5 wherein  $(\text{amino acid})^1$  and  $(\text{amino acid})^2$  are each independently any naturally-  
 occurring, modified, substituted or altered  $\alpha$ - or  $\beta$ -amino acid not containing a thiol  
 group; A is H, HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or  
 peptide)-OOC or R<sup>4</sup>; B is H, SH or -NHR<sup>3</sup>, -N(R<sup>3</sup>)-(amino acid or peptide) or R<sup>4</sup>;  
 Z is H or R<sup>4</sup>; X is SH or -NHR<sup>3</sup>, -N(R<sup>3</sup>)-(amino acid or peptide) or R<sup>4</sup>; R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>  
 10 and R<sup>4</sup> are independently H or straight or branched chain or cyclic lower alkyl; n is  
 an integer that is either 0, 1 or 2; (peptide) is a peptide of 2 to about 10 amino acids;  
 and: (1) where B is -NHR<sup>3</sup> or -N(R<sup>3</sup>)-(amino acid or peptide), X is SH and n is 1 or  
 2; (2) where X is -NHR<sup>3</sup> or -N(R<sup>3</sup>)-(amino acid or peptide), B is SH and n is 1 or  
 2; (3) where B is H or R<sup>4</sup>, A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC,  
 15 (amino acid or peptide)-OOC, X is SH and n is 0 or 1; (4) where A is H or R<sup>4</sup>, then  
 where B is SH, X is -NHR<sup>3</sup> or -N(R<sup>3</sup>)-(amino acid or peptide) and where X is SH,  
 B is -NHR<sup>3</sup> or -N(R<sup>3</sup>)-(amino acid or peptide); (5) where X is H or R<sup>4</sup>, A is HOOC,  
 H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC and B is SH;  
 (6) where Z is methyl, X is methyl, A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-  
 20 NHOC, (amino acid or peptide)-OOC and B is SH and n is 0; and (7) where Z is SH  
 and X is SH, n is not 0; and wherein the thiol group is in the reduced form.

Additional preferred embodiments include radiolabel binding moieties having  
 the formula: -Gly-Gly-Cys-, Cys-Gly-Gly-, Gly-Gly-Cys-, -( $\epsilon$ -Lys)-Gly-Cys-, ( $\delta$ -  
 Orn)-Gly-Cys-, -( $\gamma$ -Dab)-Gly-Cys-, and -( $\beta$ -Dap)-Gly-Cys-. (In these formulae, it  
 25 will be understood that  $\epsilon$ -Lys represents a lysine residue in which the  $\epsilon$ -amino group,  
 rather than the typical  $\alpha$ -amino group, is covalently linked to the carboxyl group of  
 the adjacent amino acid to form a peptide bond;  $\delta$ -Orn represents an ornithine residue  
 in which the  $\delta$ -amino group, rather than the typical  $\alpha$ -amino group, is covalently  
 linked to the carboxyl group of the adjacent amino acid to form a peptide bond;  $\gamma$ -  
 30 Dab represents a 2,4-diaminobutyric acid residue in which the  $\gamma$ -amino group is  
 covalently linked to the carboxyl group of the adjacent amino acid to form a peptide  
 bond; and  $\beta$ -Dap represents a 1,3-diaminopropionic acid residue in which the  $\beta$ -


amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond.)

In another embodiment, the invention provides peptide reagents capable of being radiolabeled with a radioisotope, for radiotherapy or for imaging sites within a mammalian body, each comprising a somatostatin analogue that is covalently linked to a radiolabel-binding moiety of formula:




(for purposes of this invention, radiolabel-binding moieties having this structure will be referred to as picolinic acid (Pic)-based moieties)

or




wherein X is H or a protecting group; (amino acid) is any amino acid and the radiolabel-binding moiety is covalently linked to the peptide. For purposes of this invention, radiolabel-binding moieties having this structure will be referred to as picolylamine (Pica)-based moieties. In a preferred embodiment, the amino acid is glycine and X is an acetamidomethyl protecting group.

Yet another embodiment of the invention provides peptide reagents capable of being radiolabeled with a radioisotope, for imaging sites within a mammalian body or for use as a radiotherapeutic agent, each comprising a somatostatin analogue that is covalently linked to a radiolabel-binding moiety that is a bisamino bisthiol radiolabel-binding moiety. The bisamino bisthiol radiolabel-binding moiety in this embodiment of the invention has the formula:



wherein each R can be independently H, CH<sub>3</sub> or C<sub>2</sub>H<sub>5</sub>; each (pgp)<sup>s</sup> can be independently a thiol protecting group or H; m, n and p are independently 2 or 3; A is linear or cyclic lower alkyl, aryl, heterocyclyl, combinations or substituted derivatives thereof; and X is peptide; or



5 wherein each R is independently H, CH<sub>3</sub> or C<sub>2</sub>H<sub>5</sub>; m, n and p are independently 2 or 3; A is linear or cyclic lower alkyl, aryl, heterocyclyl, combinations or substituted derivatives thereof; V is H or CO-peptide; R' is H or peptide; provided that when V is H, R' is peptide and when R' is H, V is peptide. For purposes of this invention, radiolabel-binding moieties having these structures will be referred to as  
10 "BAT" moieties.

This invention provides methods for preparing peptide reagents of the invention by chemical synthesis *in vitro*. In a preferred embodiment, peptides are synthesized by solid phase peptide synthesis.

15 This invention provides reagents for preparing a radiolabeled somatostatin receptor-binding agent comprising the somatostatin receptor-binding peptides of the invention covalently linked to a radiolabel-binding moiety. In a preferred embodiment, the reagent is radioactively labeled with Tc-99m. In another preferred embodiment, the reagent is radioactively labeled with <sup>186</sup>Re or <sup>188</sup>Re.

20 The invention also provides complexes of the linear peptide reagents of the invention with a radioisotope, as well as methods for radiolabeling the peptide reagents of the invention. For example, in one embodiment scintigraphic imaging agents provided by the invention comprise Tc-99m labeled complexes formed by reacting the peptide reagents of the invention with Tc-99m in the presence of a reducing agent. Preferred reducing agents include but are not limited to dithionite  
25 ion, stannous ion and ferrous ion. Such Tc-99m complexes of the invention are also formed by labeling the peptide reagents of the invention with Tc-99m by ligand exchange of a prereduced Tc-99m complex as provided herein.

30 The invention also provides kits for preparing radiolabeled linear somatostatin analogue peptides from the peptide reagents of the invention. Kits for radiolabeling the peptide reagents of the invention are comprised of a sealed vial containing a predetermined quantity of a peptide reagent of the invention and a sufficient amount of reducing agent to radiolabel the peptide. In a preferred embodiment, the

radiolabeled somatostatin analogue is a scintigraphic imaging agent. Also preferred is radiolabeling the peptide reagents of the invention with Tc-99m. Kits for preparing radiotherapeutic agents are also provided, wherein the preferred radioisotopes are rhenium-186 and rhenium-188.

5           This invention provides methods for using the radiolabeled peptide reagents of the invention diagnostically and therapeutically. In one embodiment of the invention, methods are provided for using scintigraphic imaging agents that are Tc-99m labeled peptide reagents for imaging sites within a mammalian body by obtaining *in vivo* gamma scintigraphic images. These methods comprise  
10           administering an effective diagnostic amount of Tc-99m labeled peptide reagents of the invention and detecting the gamma radiation emitted by the Tc-99m label localized at the site within the mammalian body.

          The invention also provides methods for alleviating somatostatin-related diseases in animals, preferably humans, comprising administering a therapeutically  
15           effective amount of the radiolabeled somatostatin-binding peptide reagents of the invention to the animal. In preferred embodiments, the reagent is radioactively labeled with  $^{186}\text{Re}$  or  $^{188}\text{Re}$ .

          The peptides and peptide reagents of the invention may also be comprised of a polyvalent linking moiety. Polyvalent linking moieties of the invention are  
20           comprised of at least 2 identical linker functional groups capable of covalently bonding to somatostatin analogue peptides or Tc-99m binding moieties. Preferred linker functional groups are primary or secondary amines, hydroxyl groups, carboxylic acid groups or thiol-reactive groups. In preferred embodiments, the polyvalent linking moieties are comprised of *bis*-succinimidylmethylether (BSME),  
25           4-(2,2-dimethylacetyl)benzoic acid (DMBA), *N*-[2-(*N'*,*N'*-*bis*(2-succinimidoethyl)aminoethyl)]-*N*<sup>6</sup>,*N*<sup>9</sup>-*bis*(2-methyl-2-mercapto-propyl)-6,9-diazanonanamide (BAT-BS), *tris*(succinimidylethyl)amine (TSEA), *bis*-succinimidohexane (BSH), 4-(O-CH<sub>2</sub>CO-Gly-Gly-Cys.amide)-2-methylpropiophenone (ETAC),  
          *tris*(acetamidoethyl)amine, *bis*-acetamidomethyl ether, *bis*-acetamidoethyl ether,  $\alpha,\epsilon$ -  
30           *bis*-acetyllysine, lysine and 1,8-*bis*-acetamido-3,6-dioxo-octane, or a derivative thereof.

          Specific preferred embodiments of the present invention will become evident

from the following more detailed description of certain preferred embodiments and the claims.

### DETAILED DESCRIPTION OF THE INVENTION

5           The present invention provides linear peptide reagents for preparing radiolabeled agents for radiodiagnostic and radiotherapeutic uses. The present invention provides linear peptides that are somatostatin analogues and that are not constrained within a cyclic structure. Such somatostatin analogues thereby possess increased *in vivo* stability compared with native somatostatin. These linear peptides  
10           are themselves therapeutic agents for alleviating diseases and other ailments in animals including humans.

          Also provided by the invention are linear peptides that may be radioiodinated or radioastatinated and which are thereby useful in radiotherapeutic and radiodiagnostic applications.

15           Another embodiment of these linear peptides that is provided by this invention are linear peptide reagents wherein the linear peptides of the invention are covalently linked to a radiolabel-binding moiety. Such linear peptide reagents are capable of being radiolabeled to provide radiodiagnostic or radiotherapeutic agents. One example of a radiodiagnostic application using the radiolabeled agents of the  
20           invention is scintigraphic imaging, wherein the location and extent of somatostatin receptor-bearing tumors may be determined.

          The linear peptide reagents of the invention can also advantageously be radiolabeled with cytotoxic radioisotopes such as rhenium-186 or rhenium-188 for radiotherapeutic uses. The linear peptide reagents of the invention are also useful  
25           in preparing complexes with non-radioactive metals, said complexes being useful therapeutically.

          The invention provides a method for using the somatostatin analogues of the invention to alleviate diseases or other ailments in animals, preferably humans. These diseases and ailments include but are not limited to diabetes and diabetes-related retinopathy, cirrhosis of the liver and hepatitis infection, bleeding ulcers and  
30           other gastrointestinal bleeding, pancreatitis, central nervous system disorders, endocrine disorders, Alzheimer's disease, acromegaly and other diseases and

disorders related to the production of inappropriate levels of growth hormone *in vivo*, and cancer, particularly those cancers whose growth is dependent or influenced by growth hormone production. Dosages of the somatostatin analogues provided by the invention may be the same as those dosages of native somatostatin routinely used for treatment of the above or other diseases, or less of the compounds of the invention may be administered due to their longer *in vivo* half-life.

In embodiments of the invention useful as scintigraphic imaging agents, labeling with Tc-99m is an advantage of the present invention because the nuclear and radioactive properties of this isotope make it an ideal scintigraphic imaging agent. This isotope has a single photon energy of 140 keV and a radioactive half-life of about 6 hours, and is readily available from a  $^{99}\text{Mo}$ - $^{99\text{m}}\text{Tc}$  generator. Other radionuclides may also be used in the practice of the invention as disclosed herein.

Radiotherapeutic embodiments of the invention, on the other hand, are advantageously labeled with cytotoxic radioisotopes including but not limited to scandium-47, copper-67, gallium-72, yttrium-90, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211 and bismuth-212, most preferably  $^{186}\text{Re}$  or  $^{188}\text{Re}$ . Such embodiments are useful in the treatment of somatostatin-related diseases or other ailments in animals, preferably humans, including but not limited to cancer and other diseases characterized by the growth of malignant or benign tumors capable of binding somatostatin or somatostatin analogues *via* the expression of somatostatin receptors on the cell surface of cells comprising such tumors.

In the radiolabel-binding moieties and linear peptides covalently linked to such moieties that contain a thiol covalently linked to a thiol protecting groups ((pgp)<sup>S</sup>) provided by the invention, the thiol-protecting groups may be the same or different and may be but are not limited to:

- CH<sub>2</sub>-aryl (aryl is phenyl or alkyl or alkyloxy substituted phenyl);
- CH-(aryl)<sub>2</sub>, (aryl is phenyl or alkyl or alkyloxy substituted phenyl);
- C-(aryl)<sub>3</sub>, (aryl is phenyl or alkyl or alkyloxy substituted phenyl);
- CH<sub>2</sub>-(4-methoxyphenyl);
- CH-(4-pyridyl)(phenyl)<sub>2</sub>;
- C(CH<sub>3</sub>)<sub>3</sub>

- 9-phenylfluorenyl;
- CH<sub>2</sub>NHCOR (R is unsubstituted or substituted alkyl or aryl);
- CH<sub>2</sub>-NHCOOR (R is unsubstituted or substituted alkyl or aryl);
- CONHR (R is unsubstituted or substituted alkyl or aryl);
- CH<sub>2</sub>-S-CH<sub>2</sub>-phenyl

Preferred protecting groups have the formula -CH<sub>2</sub>-NHCOR wherein R is a lower alkyl having 1 and 8 carbon atoms, phenyl or phenyl-substituted with lower alkyl, hydroxyl, lower alkoxy, carboxy, or lower alkoxycarbonyl. The most preferred protecting group is an acetamidomethyl group.

- Each somatostatin receptor-binding linear peptide-containing embodiment of the invention is comprised of a sequence of amino acids. The term amino acid as used in this invention is intended to include all L- and D- amino acids, naturally occurring and otherwise. Reagents comprising somatostatin receptor-binding peptides provided by the invention include but are not limited to the following illustrative examples of the peptide embodiments of the invention:

- C<sub>Acm</sub>GC<sub>Acm</sub>GGGF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 (DTPA).F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 maGGGF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 20 Ac.C<sub>Acm</sub>GC<sub>Acm</sub>F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFTC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 (DTPA).D-Nal.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 AKCGGGF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 (DTPA).D-Nal.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 25 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.GGGC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 (DTPA).Aca.F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 (DTPA).(ε-K)GCF<sub>D</sub>.FYW<sub>D</sub>KTFT.amide  
 Ac.CGCF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFTCGC.amide  
 30 (DTPA).(D-Nal.CYW<sub>D</sub>KVCT)<sub>2</sub>  
 Ac.F<sub>D</sub>.FYW<sub>D</sub>KTFT(ε-K)GC.amide  
 Ac.F<sub>D</sub>FYW<sub>D</sub>KTFTGGG(ε-K)GC.amide

- $F_D$ .Cpa.YW<sub>D</sub>KTC.Nal.amide  
 K(BAT).D-Nal.C<sub>Me</sub>YW<sub>D</sub>KVC<sub>Me</sub>T.amide  
 Ac.F<sub>D</sub>FYW<sub>D</sub>KTFGGG(ε-K)KC.amide  
 Pic.GC<sub>Acm</sub>GGGF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 5 (DTPA).D-Nal.CYW<sub>D</sub>KVCT.amide  
 (2-ketogulonyl)D-NalFYW<sub>D</sub>KVCT.amide  
 $F_D$ .Cpa.YW<sub>D</sub>K.Abu.Nal.T(ε-K)GC.amide  
 (DTPA).K(BAT).D-Nal.C<sub>Me</sub>YW<sub>D</sub>KVC<sub>Me</sub>T.amide  
 $F_D$ .Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 10 (DTPA).F<sub>D</sub>FYW<sub>D</sub>KTFT(ε-K)GC.amide  
 AF<sub>D</sub>CFW<sub>D</sub>KTC<sub>Me</sub>T(CH<sub>2</sub>OH)  
 (DTPA).F<sub>D</sub>GYW<sub>D</sub>KTCT(CH<sub>2</sub>OH)  
 (DTPA).Nal.SYW<sub>D</sub>KVT.K(BAT).amide  
 (DTPA).Nal.SYW<sub>D</sub>KVCT.amide  
 15 F<sub>D</sub>FYW<sub>D</sub>KTFTGGCK.amide  
 DDD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 Ac.DDD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 Hca.G.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 F<sub>D</sub>FYW<sub>D</sub>KTFTC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 20 F<sub>D</sub>FYW<sub>D</sub>KTFTGGC.amide  
 F<sub>D</sub>FYW<sub>D</sub>KTFT(ε-K)GC.amide  
 (Trc.imide)<sub>2</sub>K.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCRR.amide  
 Trc(Trc.imide)K.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCRR.amide  
 (Trc.imide)Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCR.amide  
 25 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCR.amide  
 K<sub>D</sub>KKK.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKDKD.amide  
 K<sub>D</sub>KKK.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCDD.amide  
 (Trc)<sub>2</sub>K.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 Hca.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 30 (2-ketogulonyl)F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 KKKK.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCDDDD.amide  
 Ac.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide

2167678

Ac.KKKKK.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 (2-ketogulonyl)F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 DDDD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKKKK.amide  
 5 (DTPA)Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 (DTPA)Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFTC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 Ac.KKKKK.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 KDKD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKDKD.amide.

As used herein, the following amino acids and amino acid analogues are  
 10 intended to be represented by the following abbreviations: Ac is an acetyl group;  
 ma is mercaptoacetic acid group; Aca is 6-aminocaproic acid; Hcy is homocysteine;  
 Hhc is homohomocysteine, which is (3-mercaptopropyl)glycine; Pen is penicillamine;  
 Mob is the sulfhydryl protecting group 4-methoxybenzyl; Acm is the sulfhydryl  
 15 protecting group acetamidomethyl; Aib is aminoisobutyric acid; Nal is 2-  
 naphthylalanine; Ain is 2-amino-indan-2-carboxylic acid; Hly is homolysine; Achxa  
 is 4-amino-cyclohexylalanine; Amf is 4-aminomethylphenylalanine; Aec is S-(2-  
 aminoethyl)cysteine; Apc is S-(3-aminopropyl)cysteine; Aes is O-(2-  
 aminoethyl)serine; Aps is O-(3-aminopropyl)serine; Abu is 2-aminobutyric acid; Nva  
 20 is norvaline; Aca is 6-aminocaproic acid; F<sub>D</sub> is D-phenylalanine; W<sub>D</sub> is D-tryptophan;  
 Y<sub>D</sub> is D-tyrosine; Cpa is L-(4-chlorophenyl)alanine; Thp is 4-amino-  
 tetrahydrothiopyran-4-carboxylic acid; D-Nal is D-2-naphthylalanine; Dpg is  
 dipropylglycine; Abu is α-aminobutyric acid; Trc is tricarboalkylic acid; Hca is  
 hexacarboxy-cyclohexane; and Nle is norleucine. All naturally-occurring amino  
 25 acids are abbreviated using standard abbreviations (which can be found in G. Zubay,  
*Biochemistry* (2d. ed.), 1988 (MacMillen Publishing: New York) p.33.

For the purposes of this invention, the naturally-occurring amino acids are  
 characterized as lipophilic (alanine, isoleucine, leucine, methionine, phenylalanine,  
 tyrosine, proline, tryptophan and valine, as well as S-alkylated derivatives of  
 cysteine), hydrophilic (asparagine, glutamine, threonine, serine), acidic (glutamic  
 30 acid and aspartic acid), basic (arginine, histidine and lysine). T(CH<sub>2</sub>OH) represents a  
 threoninol residue, wherein the carboxyl group of the amino acid is reduced to a  
 primary alcohol, incorporated into the peptide using the procedure of Neugebauer *et*

al. (1990, Peptides: Proceedings of the 11th American Peptide Symposium, pp. 1020-21).  $\epsilon$ -K is intended to represent a covalent linkage *via* the  $\epsilon$ -amino group on the sidechain of a lysine residue.  $\delta$ -Orn represents an ornithine residue in which the  $\delta$ -amino group, rather than the typical  $\alpha$ -amino group, is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond.  $\gamma$ -Dab represents a 2,4-diaminobutyric acid residue in which the  $\gamma$ -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond.  $\beta$ -Dap represents a 1,3-diaminopropionic acid residue in which the  $\beta$ -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond. Pic is picolinoyl (pyridine-2-carbonyl); Pica is picolylamine (2-(aminomethyl)pyridine); (BAT) represents  $N^6, N^9$ -bis(2-mercapto-2-methyl-propyl)-6,9-diazanonanoic acid; K.(BAT) and Lys.(BAT) represent the amino acid lysine, acylated at the  $\epsilon$ -amino group on the amino acid sidechain to (BAT); (BAM) is ( $N^1, N^4$ -bis(2-mercapto-2-methylpropyl)-1,4,10-triazadecane; E.(BAM) and Glu.(BAM) represent the amino acid glutamic acid having a  $\gamma$ -amide linkage between the sidechain carboxylic acid group of glutamic acid and a (BAM)-derived primary amino group; (BAT-BM) is  $N$ -(2-( $N', N'$ -bis(2-maleimidoethyl)aminoethyl)- $N^9$ -(*t*-butoxycarbonyl)- $N^6, N^9$ -bis(2-methyl-2-triphenylmethylthiopropyl)-6,9-diazanonanamide; (BAT-BS) is  $N$ -(2-( $N', N'$ -bis(2-succinimidoethyl)aminoethyl)- $N^6, N^9$ -bis(2-mercapto-2-methylpropyl)-6,9-diazanonanamide; (BMME) is *bis*-maleimidomethylether; (BSME) is *bis*-succinimidomethylether; and (DTPA) is diethylenetriaminepentaacetic acid.

For the purposes of this invention the term "poly(*N*-carboxyalkyl)amine" is intended to describe a series of compounds exemplified by nitrilotriacetic acid, iminodiacetic acid, ethylenediaminetetraacetic acid (EDTA) and diethylenetriaminepentaacetic acid (DTPA).

For the purposes of this invention the term "polyoxyanion" is intended to encompass sulfates, phosphates, sulfonates, phosphonates and like compounds.

Linear somatostatin analogue peptides of the present invention can be chemically synthesized *in vitro*. Peptides of the present invention can generally advantageously be prepared on a peptide synthesizer. The peptides of this invention can be synthesized wherein the radiolabel-binding moiety is covalently linked to the

peptide during chemical synthesis *in vitro*, using techniques well known to those with skill in the art. Such peptides covalently-linked to the radiolabel-binding moiety during synthesis are advantageous because specific sites of covalent linkage can be determined.

5 Radiolabel binding moieties of the invention may be introduced into the target linear somatostatin analogue peptides during peptide synthesis. For embodiments comprising picolinic acid ((Pic-); *e.g.*, Pic-Gly-Cys(protecting group)-), the radiolabel-binding moiety can be synthesized as the last (*i.e.*, amino-terminal) residue in the synthesis. In addition, the picolinic acid-containing radiolabel-binding moiety  
10 may be covalently linked to the  $\epsilon$ -amino group of lysine to give, for example,  $\alpha$ N(Fmoc)-Lys- $\epsilon$ N(Pic-Gly-Cys(protecting group)), which may be incorporated at any appropriate position in the peptide chain. This sequence is particularly advantageous as it affords an easy mode of incorporation into the target somatostatin analogue peptide.

15 Similarly, the picolylamine (Pica)-containing radiolabel-binding moiety (-Cys(protecting group)-Gly-Pica) can be prepared during peptide synthesis by including the sequence (-Cys(protecting group)-Gly-) at the carboxyl terminus of the peptide chain. Following cleavage of the peptide from the resin the carboxyl terminus of the peptide is activated and coupled to picolylamine. This synthetic  
20 route requires that reactive side-chain functionalities remain masked (protected) and do not react during the conjugation of the picolylamine.

This invention also provides small linear synthetic peptides that are somatostatin analogues and incorporate bisamine bithiol (BAT) chelators that may be labeled with Tc-99m.

25 This invention provides for the incorporation of these chelators into virtually any position in the peptide, *via* covalently linkage to any appropriate functional group of the peptide, except that the chelating moieties of the invention are not covalently linked to functional groups comprising the amino acid side chains of the amino acids B<sup>1</sup>, B<sup>2</sup>, B<sup>3</sup> or B<sup>4</sup>.

30 In forming a complex of radioactive technetium with the reagents of this invention, the technetium complex, preferably a salt of Tc-99m pertechnetate, is reacted with the reagent in the presence of a reducing agent. Preferred reducing

agents are dithionite, stannous and ferrous ions; the most preferred reducing agent is stannous chloride. Means for preparing such complexes are conveniently provided in a kit form comprising a sealed vial containing a predetermined quantity of a reagent of the invention to be labeled and a sufficient amount of reducing agent to label the reagent with Tc-99m. Alternatively, the complex may be formed by reacting a reagent of this invention with a pre-formed labile complex of technetium and another compound known as a transfer ligand. This process is known as ligand exchange and is well known to those skilled in the art. The labile complex may be formed using such transfer ligands as tartrate, citrate, gluconate or mannitol, for example. Among the Tc-99m pertechnetate salts useful with the present invention are included the alkali metal salts such as the sodium salt, or ammonium salts or lower alkyl ammonium salts.

In a preferred embodiment of the invention, a kit for preparing technetium-labeled peptides is provided. An appropriate amount of the peptide reagent is introduced into a vial containing a reducing agent, such as stannous chloride, in an amount sufficient to label the peptide with Tc-99m. An appropriate amount of a transfer ligand as described (such as tartrate, citrate, gluconate or mannitol, for example) can also be included. The kit may also contain conventional pharmaceutical adjunct materials such as, for example, pharmaceutically acceptable salts to adjust the osmotic pressure, buffers, preservatives and the like. The components of the kit may be in liquid, frozen or dry form. In a preferred embodiment, kit components are provided in lyophilized form.

Technetium-99m labeled imaging reagents according to the present invention may be prepared by the addition of an appropriate amount of Tc-99m or Tc-99m complex into the vials and reaction under conditions described in Example 2 hereinbelow.

Radioactively-labeled scintigraphic imaging agents provided by the present invention are provided having a suitable amount of radioactivity. In forming Tc-99m radioactive complexes, it is generally preferred to form radioactive complexes in solutions containing radioactivity at concentrations of from about 0.01 millicurie (mCi) to 100 mCi per mL.

The imaging reagents provided by the present invention can be used for

visualizing organs such as the kidney for diagnosing disorders in these organs, and tumors, in particular gastrointestinal tumors, myelomas, small cell lung carcinoma and other APUDomas, endocrine tumors such as medullary thyroid carcinomas and pituitary tumors, brain tumors such as meningiomas and astrocytomas, and tumors of the prostate, breast, colon, and ovaries can also be imaged. In accordance with this invention, the Tc-99m labeled peptide reagents are administered in a single unit injectable dose. The Tc-99m labeled peptide reagents provided by the invention may be administered intravenously in any conventional medium for intravenous injection such as an aqueous saline medium, or in blood plasma medium. Generally, the unit dose to be administered has a radioactivity of about 0.01 mCi to about 100 mCi, preferably 1 mCi to 20 mCi. The solution to be injected at unit dosage is from about 0.01 mL to about 10 mL. After intravenous administration, imaging *in vivo* can take place in a matter of a few minutes. However, imaging can take place, if desired, in hours or even longer, after the radiolabeled peptide is injected into a patient. In most instances, a sufficient amount of the administered dose will accumulate in the area to be imaged within about 0.1 of an hour to permit the taking of scintiphotos. Any conventional method of scintigraphic imaging for diagnostic purposes can be utilized in accordance with this invention.

The somatostatin receptor-binding linear peptides and non-radioactive metal complexes of the linear peptide reagents of the invention may be used clinically to promote regression of certain types of tumors, particularly those that express somatostatin receptors. The linear somatostatin analogue peptides of the invention can also be used to reduce the hormonal hypersecretion that often accompanies certain cancers, such as the APUDomas. Peptides of the invention used as therapeutic agents may be administered by any appropriate route, including intravenous, intramuscular or by mouth, and in any acceptable pharmaceutical carrier, in doses ranging from about 0.1 to about 49 mg/kgbody weight/day.

This invention also provides peptides radiolabeled with a cytotoxic radioisotope such as rhenium-186 or rhenium-188 that may be used for radiotherapy of certain tumors as described above. For this purpose, an amount of radioactive isotope from about 10mCi to about 200mCi may be administered *via* any suitable clinical route, preferably by intravenous injection.

The methods for making and labeling these compounds are more fully illustrated in the following Examples. These Examples illustrate certain aspects of the above-described method and advantageous results, and are shown by way of illustration and not limitation.

5

### EXAMPLE 1

#### Solid Phase Peptide Synthesis

10

Solid phase peptide synthesis (SPPS) was carried out on a 0.25 millimole (mmole) scale using an Applied Biosystems Model 431A Peptide Synthesizer and using 9-fluorenylmethyloxycarbonyl (Fmoc) amino-terminus protection, coupling with dicyclohexylcarbodiimide/hydroxybenzotriazole or 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate/ hydroxybenzotriazole (HBTU/HOBT), and using *p*-hydroxymethylphenoxy-methylpolystyrene (HMP) resin for carboxyl-terminus acids or Rink<sup>TM</sup> amide resin for carboxyl-terminus amides.

15

Where appropriate, the following amino acid derivatives were synthesized. Homocysteine was prepared by alkaline hydrolysis of L-homocysteine lactone. Threoninol residues, wherein the carboxyl group of the amino acid is reduced to a primary alcohol, can be introduced into the peptides of the invention where appropriate using the procedure of Neugebauer *et al.* (1990, Peptides: Proceedings of the 11th American Peptide Symposium, pp. 1020-21). Fmoc.Hcy(Trt) and Fmoc.Pen(Trt) were prepared from the appropriate amino acids by tritylation with triphenylmethanol in TFA, followed by Fmoc derivitization as described by Atherton *et al.* (1989, Solid Phase Peptide Synthesis, IRL Press: Oxford). Fmoc.homohomocysteine(Trt) was prepared by reducing *N,N*-bis-Boc-glutamic acid- $\alpha$ -methyl ester with borane-THF, followed by mesylation and reaction with trityl-mercaptide, followed by removal of the Boc groups with BF<sub>3</sub>OEt in acetic acid, and then Fmoc derivitization as described above. PhCH<sub>2</sub>CHBrCOOH was prepared by treating phenylalanine (in a solution of water and TFA/ saturated with NaBr) with sodium nitrite, followed by distillation to recover the pure product.

20

25

30

Where appropriate, 2-chloroacetyl, 2-bromoacetyl and 2-bromo-3-phenylpropionyl groups were introduced either by using the appropriate 2-halo acid as the last residue coupled during SPPS, or by treating the N-terminus free amino

acid peptide bound to the resin with either 2-halo acid/ diisopropylcarbodiimide/*N*-hydroxysuccinimide/NMP or 2-halo acid anhydride/ diisopropylethylamine/NMP.

5 Where appropriate, HPLC-purified 2-haloacylated peptides were cyclized by stirring an 0.1-1.0 mg/mL solution in phosphate or bicarbonate buffer or dilute ammonium hydroxide (pH 8.0), optionally containing 0.5-1.0 mM EDTA, or acetonitrile or THF for 1-48 h followed optionally by acidification with acetic acid, lyophilization and HPLC purification.

10 Where appropriate, (BAM) (*N*<sup>1</sup>,*N*<sup>4</sup>-bis(2-mercapto-2-methylpropyl)-1,4,10-triazadecane) was conjugated to the peptide by first activating the peptide carboxylate with a mixture of diisopropylcarbodiimide/ *N*-hydroxysuccinimide or HBTU/HOBt in DMF, NMP or methylene chloride, followed by coupling in the presence of diisopropylethylamine. After coupling, the conjugates were deprotected as described above.

15 Where appropriate, (BAT) (*N*<sup>6</sup>,*N*<sup>9</sup>-bis(2-mercapto-2-methylpropyl)-6,9-diazanonanoic acid) was incorporated into peptide as (*N*α(Fmoc)-*N*ε(*N*-Boc)-*S*,*S*' -bistrityl-BAT)lysine (prepared from *N*α(Fmoc)-lysine and *N*ε(*N*-Boc)-*S*,*S*' -bistrityl-BAT as described elsewhere during peptide synthesis and then deprotected after cleavage of the completed peptide from the synthetic resin.

20 Where appropriate, BSME adducts were prepared by reacting single thiol-containing peptides (5 to 50 mg/mL in DMF buffered to pH 7 with *N*-methylemorpholine or *N*-ethyl-morpholine, or 50mM sodium phosphate buffer, pH 7-8, optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.5 molar equivalents of BMME (*bis*-maleimidomethylether) pre-dissolved in acetonitrile  
25 at room temperature for approximately 1-18 hours. The solution was concentrated and the product was purified by HPLC.

30 Where appropriate, TSEA adducts were prepared by reacting single thiol-containing peptide (at concentrations of 10 to 100 mg/mL peptide in DMF buffered to pH 7 with *N*-methylemorpholine or *N*-ethylmorpholine, or 5 to 50 mg/mL peptide in 50mM sodium phosphate, pH 7-8, optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.33 molar equivalents of TMEA (*tris*(2-maleimidoethyl)amine) pre-dissolved in acetonitrile or DMF, with or without 1 molar

equivalent of triethanolamine, at room temperature for approximately 1-18h. Such reaction mixtures containing adducts were concentrated and the adducts were then purified using HPLC.

5           Where appropriate, BAT-BS (*N*-(2-(*N'*,*N'*-bis(2-succinimidoethyl) aminoethyl))-*N*<sup>6</sup>,*N*<sup>9</sup>-bis(2-methyl-2-mercaptopropyl)-6,9-diazanonanamide) adducts were prepared by reacting single thiol-containing peptide (at concentrations of 2 to 50 mg/mL peptide in DMF buffered to pH 7 with *N*-methyl-morpholine or *N*-ethyl-morpholine, or in 50mM sodium phosphate (pH 7-8), optionally containing 0.5mM EDTA or DMF or THF or acetonitrile) with 0.5 molar equivalents of BAT-BM (*N*-(2-(*N'*,*N'*-bis(2-maleimidoethyl)aminoethyl))-*N*<sup>9</sup>-(*t*-butoxycarbonyl)-*N*<sup>6</sup>,*N*<sup>9</sup>-bis(2-methyl-2-triphenylmethylthiopropyl)-6,9-diazanonanamide) pre-dissolved in acetonitrile or THF, at room temperature for approximately 1-18h. The solution was then evaporated to dryness and (BAT-BS)-peptide conjugates deprotected by treatment with 10mL TFA and 0.2mL triethylsilane for 1h. The solution was  
10           concentrated, the product adducts precipitated with ether, and then purified by HPLC.  
15

Where appropriate, the (DTPA) moiety can be introduced using the method of Bakker *et al.* (1991, Life Sci. 49: 1583-1591).

20           Resin-bound products were routinely cleaved using a solution of trifluoroacetic acid or trifluoroacetic acid and methylene chloride, optionally containing water, thioanisole, ethanedithiol, and triethylsilane, prepared in ratios of 100 : 5 : 5 : 2.5 : 2 for 0.5 - 3 h at room temperature. Crude peptides were purified by preparative high pressure liquid chromatography (HPLC) using a Waters<sup>TM</sup> Delta Pak C18 column and gradient elution using 0.1% trifluoroacetic acid (TFA) in water modified with acetonitrile. Acetonitrile was evaporated from the eluted  
25           fractions which were then lyophilized. The identity of each product was confirmed by fast atom bombardment mass spectroscopy (FABMS) or by electrospray mass spectroscopy (ESMS).

30           Somatostatin analogues synthesized as provided herein, as well as the products of such synthesis identified by FABMS, are shown in Table I below.

2167678

## EXAMPLE 2

A General Method for Radiolabeling

0.1 mg of a peptide prepared as in Example 2 was dissolved in 0.1 mL of water or 50/50 ethanol/water or phosphate-buffered saline or 50 mM potassium phosphate buffer (pH = 5, 6 or 7.4). Tc-99m gluceptate was prepared by reconstituting a Glucoscan vial (E.I. DuPont de Nemours, Inc.) with 1.0 mL of Tc-99m sodium pertechnetate containing up to 200 mCi and allowed to stand for 15 minutes at room temperature. 25  $\mu$ l of Tc-99m gluceptate was then added to the peptide and the reaction allowed to proceed at room temperature or at 100°C for 15-30 min and then filtered through a 0.2  $\mu$ m filter.

The Tc-99m labeled peptide purity was determined by HPLC using the following conditions: a Waters<sup>TM</sup> Delta Pak RP-18, 5 $\mu$ , 4.6mm x 220mm analytical column was loaded with each radiolabeled peptide, and the peptides eluted at a solvent flow rate equal to 1 mL/min. Gradient elution was performed beginning with 100% solvent A (0.1% CF<sub>3</sub>COOH/H<sub>2</sub>O) and ending with 100% solvent B<sub>90</sub> (0.1% CF<sub>3</sub>COOH/90% CH<sub>3</sub>CN/H<sub>2</sub>O) over the course of 10-20 min.

Radioactive components were detected using an in-line radiometric detector linked to an integrating recorder. Tc-99m gluceptate and Tc-99m sodium pertechnetate elute between 1 and 4 minutes under these conditions, whereas the Tc-99m labeled peptides eluted after a much greater amount of time, as illustrated in Table I below.

2167678

TABLE I

| Peptide                                                                               | MH <sup>+</sup> | RCY (%)         | R <sub>t</sub> (min)    |
|---------------------------------------------------------------------------------------|-----------------|-----------------|-------------------------|
| C <sub>Acm</sub> GC <sub>Acm</sub> GGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide  | 1749            | 97 <sup>6</sup> | 15.7 <sup>2</sup>       |
| (DTPA).F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                          | 1837            | 97 <sup>7</sup> | 15.5 <sup>2</sup>       |
| ma.GGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                  | 1417            | 98 <sup>6</sup> | 12.2 <sup>3</sup>       |
| Ac.C <sub>Acm</sub> GC <sub>Acm</sub> F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide  | 1619            | 75 <sup>6</sup> | 17.1, 17.5 <sup>2</sup> |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTC <sub>Acm</sub> GC <sub>Acm</sub> .amide     | 1577            | 93 <sup>5</sup> | 12.1 <sup>3</sup>       |
| (DTPA).D-Nal.Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                                  | 2143            | N.D.            | N.D.                    |
| AKCGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                   | 1612            | 98 <sup>7</sup> | 15-16 <sup>2</sup>      |
| (DTPA).D-Nal.Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                                    | 1887            | 97 <sup>7</sup> | 16.2 <sup>2</sup>       |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.GGGC <sub>Acm</sub> GC <sub>Acm</sub> .amide | 1749            | 76 <sup>5</sup> | 17.7, 18.0 <sup>1</sup> |
| (DTPA).Aca.F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                      | 1950            | 97 <sup>3</sup> | 11.5 <sup>3</sup>       |
| (DTPA).(ε-K)GCF <sub>D</sub> .FYW <sub>D</sub> KTFT.amide                             | 1802            | 97 <sup>3</sup> | 11.5 <sup>3</sup>       |
| Ac.CGCF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                  | 1477            | 98 <sup>8</sup> | 18.1 <sup>2</sup>       |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTCGC.amide                                     | 1435            | 99 <sup>8</sup> | 16.8, 17.0 <sup>2</sup> |
| (DTPA).(D-Nal.CYW <sub>D</sub> KVCT) <sub>2</sub>                                     | 2554            | 97 <sup>8</sup> | 11.8-12.4 <sup>3</sup>  |
| Ac.F <sub>D</sub> .FYW <sub>D</sub> KTFT(ε-K)GC.amide                                 | 1469            | 96 <sup>3</sup> | 12.1, 12.6 <sup>3</sup> |
| Ac.F <sub>D</sub> FYW <sub>D</sub> KTFTGGG(ε-K)GC.amide                               | 1640            | 98 <sup>3</sup> | 11.9, 12.4 <sup>3</sup> |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC.Nal.amide                                     | 1224            | 88 <sup>8</sup> | 18.6, 20.4 <sup>2</sup> |

5

10

15

20

30

30

SUBSTITUTE SHEET (RULE 26)

2167678

TABLE I (cont'd.)

| Peptide                                                                               | MH <sup>+</sup> | RCY (%)          | R <sub>t</sub> (min)   |
|---------------------------------------------------------------------------------------|-----------------|------------------|------------------------|
| KDKD.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide                     | 2484            | 99 <sup>6</sup>  | 14.8                   |
| Ac.KKKKK.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide                   | 2450            | 98 <sup>6</sup>  | 14.2                   |
| Ac.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide                         | 1810            | 96 <sup>6</sup>  | 16.8                   |
| KKKK.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCDDDD.amide                     | 2485            | 98 <sup>6</sup>  | 14.6                   |
| (2-ketogulonyl)F <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide               | 1944            | 99 <sup>6</sup>  | 16.0                   |
| Hca.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide                        | 2097            | 98 <sup>6</sup>  | 15.8                   |
| (Trc) <sub>2</sub> K.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide       | 2212            | 98 <sup>6</sup>  | 15.7                   |
| K <sub>D</sub> KKK.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCDD.amide         | 2253            | 99 <sup>6</sup>  | 14.7                   |
| K <sub>D</sub> KKK.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide       | 2485            | 79 <sup>6</sup>  | 14.1                   |
| F <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCR.amide                               | 1617            | 99 <sup>6</sup>  | 15.4                   |
| (Trc.imide)Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCR.amide                  | 1808            | 99 <sup>6</sup>  | 17.3                   |
| Trc(Trc.imide)K.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCRR.amide            | 2250            | 99 <sup>6</sup>  | 16.2                   |
| (Trc.imide) <sub>2</sub> K.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCRR.amide | 2232            | 99 <sup>6</sup>  | 16.6                   |
| F <sub>D</sub> FYW <sub>D</sub> KTFT(ε-K)GC.amide                                     | 1427            | 99 <sup>6</sup>  | 15.1                   |
| F <sub>D</sub> FYW <sub>D</sub> KTFTGGC.amide                                         | 1356            | 99 <sup>6</sup>  | 15.2-16.2              |
| F <sub>D</sub> FYW <sub>D</sub> KTFTGGCK.amide                                        | 1484            | 99 <sup>6</sup>  | 14.9,15.6 <sup>3</sup> |
| DDD.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide                        | 1998            | 99 <sup>5</sup>  | 15.6 <sup>3</sup>      |
| Ac.DDD.Nal <sub>D</sub> .Cpa. YW <sub>D</sub> KTFT(ε-K)GCKK.amide                     | 2040            | 100 <sup>5</sup> | 16.0 <sup>3</sup>      |

2167678

TABLE I (cont'd)

| Peptide                                                                                    | MH <sup>+</sup> | RCY (%)         | R <sub>t</sub> (min)   |
|--------------------------------------------------------------------------------------------|-----------------|-----------------|------------------------|
| DDDD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.(ε-K).GCKKKK.amide                         | 2484            | 98 <sup>6</sup> | 15.1                   |
| (DTPA).Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.(ε-K).GCKK.amide                         | 2192            | 95 <sup>6</sup> | 15.8 <sup>3</sup>      |
| (DTPA).Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTC <sub>Acm</sub> GC <sub>Acm</sub> .amide | 2003            | 93 <sup>7</sup> | 16.4 <sup>3</sup>      |
| Ac.KKKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.(ε-K).GC.amide                         | 2192            | 94 <sup>2</sup> | 14.9 <sup>3</sup>      |
| Hca.G.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                            | 2136            | 93 <sup>5</sup> | 16.0 <sup>3</sup>      |
| (DTPA).F <sub>D</sub> FYW <sub>D</sub> KTFT(ε-K)GC.amide                                   | 1801            | 97 <sup>5</sup> | 11.3 <sup>3</sup>      |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                                      | 1461            | 98 <sup>7</sup> | 15.8 <sup>2</sup>      |
| (DTPA).K.(BAT).D-Nal.C <sub>Me</sub> YW <sub>D</sub> KVC <sub>Me</sub> T.amide             | 1949            | 96 <sup>8</sup> | 12.3 <sup>3</sup>      |
| F <sub>D</sub> .Cpa.YW <sub>D</sub> K.Abu.Nal.T(ε-K)GC.amide                               | 1495            | 95 <sup>7</sup> | 16.5 <sup>2</sup>      |
| (2-ketogulonyl)-D-NalFYW <sub>D</sub> KVCT.amide                                           | 1318            | 98 <sup>3</sup> | 12.4,13.0 <sup>3</sup> |
| (DTPA).D-Nal.CYW <sub>D</sub> KVCT.amide                                                   | 1473            | 97 <sup>8</sup> | 11.0 <sup>3</sup>      |
| Pic.GC <sub>Acm</sub> GGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                    | 1681            | 98 <sup>8</sup> | 13.8-16.8 <sup>1</sup> |
| Ac.F <sub>D</sub> FYW <sub>D</sub> KTFTGGG(ε-K)KC.amide                                    | 1710            | 98 <sup>8</sup> | 15.9 <sup>2</sup>      |
| K.(BAT).D-Nal.C <sub>Me</sub> YW <sub>D</sub> KVC <sub>Me</sub> T.amide                    | 1573            | 97 <sup>8</sup> | 12.5 <sup>3</sup>      |
| AF <sub>D</sub> CFW <sub>D</sub> KTC <sub>Me</sub> T(CH <sub>2</sub> OH)                   | 1106            | 99 <sup>2</sup> | 11.3-11.9 <sup>3</sup> |
| (DTPA).F <sub>D</sub> GYW <sub>D</sub> KTCT(CH <sub>2</sub> OH)                            | N.D.            | 96 <sup>3</sup> | 10.6 <sup>1</sup>      |
| (DTPA).Nal.SYW <sub>D</sub> KVTK.(BAT).amide                                               | 1801            | 96 <sup>3</sup> | 12.0 <sup>3</sup>      |
| (DTPA).Nal.SYW <sub>D</sub> KVCT.amide                                                     | 1457            | 95 <sup>8</sup> | 11.6 <sup>3</sup>      |

5

10

15

20

3 2

TABLE I (cont'd.)

| Peptide                                                                       | MH <sup>+</sup> | RCY (%)         | R <sub>t</sub> (min) |
|-------------------------------------------------------------------------------|-----------------|-----------------|----------------------|
| Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.(ε-K).GCKK.amide                   | 1767            | 98 <sup>6</sup> | 15.8                 |
| (2-ketogulonyl).F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.(ε-K).GC.amide       | 1636            | 99 <sup>2</sup> | 15.8                 |
| F <sub>D</sub> FYW <sub>D</sub> KTFTC <sub>Acm</sub> GC <sub>Acm</sub> .amide | 1544            | N.D.            | N.D.                 |

\* The following labeling conditions were used with the appropriate peptides:

1. The peptide is dissolved in 50 mM potassium phosphate buffer (pH 7.4) and labeled at room temperature.
2. The peptide is dissolved in water and labeled at room temperature.
3. The peptide is dissolved in water and labeled at 100°C.
4. The peptide is dissolved in 50% ethanol/water and labeled at 100°C.
5. The peptide is dissolved in 10% hydroxypropylcyclodextrin and labeled at room temperature.
6. The peptide is dissolved in 50% ethanol/water and labeled at room temperature.
7. The peptide is dissolved in water adjusted to pH 9 and labeled at 100°C.
8. The peptide is dissolved in water adjusted to pH 6.5 and labeled at 100°C.

\*\* HPLC methods:

20                    general:           solvent A       = 0.1% CF<sub>3</sub>COOH/H<sub>2</sub>O  
                                                                  solvent B<sub>90</sub>    = 0.1% CF<sub>3</sub>COOH/90% CH<sub>3</sub>CN/H<sub>2</sub>O  
                                                                  solvent flow rate = 1 mL/min

Columns: a. Vydak<sup>TM</sup> column = Vydak 218TP54 RP-18, 5 $\mu$  x 220mm x 4.6mm analytical column with guard column  
 b. Waters column = Waters Delta-Pak C18, 5 $\mu$ m, 39 x 150mm

5

|           |               |                                          |
|-----------|---------------|------------------------------------------|
| Method 1: | Vydak column  | 100% A to 100% B <sub>90</sub> in 10 min |
| Method 2: | Waters column | 100% A to 100% B <sub>90</sub> in 20 min |
| Method 3: | Waters column | 100% A to 100% B <sub>90</sub> in 10 min |

- 10 Single-letter abbreviations for amino acids can be found in G. Zubay, *Biochemistry* (2d. ed.), 1988 (MacMillen Publishing: New York) p.33; Ac = acetyl; Acn = acetamidomethyl; ma = mercaptoacetic acid; Aca = 6-aminocaproic acid; Hly = homolysine; Apc = L-(S-(3-aminopropyl)cysteine; F<sub>D</sub> = D-phenylalanine; W<sub>D</sub> = D-tryptophan; Y<sub>D</sub> = D-tyrosine; Cpa = L-(4-chlorophenyl)alanine; D-Nal = D-2-naphthylalanine; Nle = norleucine; Hcy = homocysteine; Hhc = homohomocysteine; Pen = penicillamine; Aib = aminoisobutyric acid; Nal = 2-naphthylalanine; D-Nal = D-2-naphthylalanine; Ain = 2-aminoindane-2-carboxylic acid; Achxa = 4-amino-cyclohexylalanine; Amf = 4-aminomethyl-phenylalanine; Aec = S-(2-aminoethyl)cysteine; Apc = S-(3-aminopropyl)cysteine; Aes = O-(2-aminoethyl)serine; Aps = O-(3-aminopropyl)serine; Abu = 2-aminobutyric acid; Trc = tricarboallylic acid; Hca = hexacarboxycyclohexane; Nva = norvaline; T(CH<sub>2</sub>OH) = threoninol (on which the carboxylic acid moiety has been reduced to a primary alcohol);  $\epsilon$ -K = a lysine residue in a peptide in which the peptide bond involves the  $\epsilon$ -amino group on the lysine sidechain rather than the  $\alpha$ -amino group;  $\delta$ -Orn = an ornithine residue in which the  $\delta$ -amino group, rather than the typical  $\alpha$ -amino group, is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond;  $\gamma$ -Dab = a 2,4-diaminobutyric acid residue in which the  $\gamma$ -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond;  $\beta$ -Dap = a 1,3-diaminopropionic acid residue in which the  $\beta$ -amino group is covalently linked to the carboxyl group of the adjacent amino acid to form a peptide bond; Pic = picolinoyl (pyridine-2-carbonyl); Pica = picolylamine (2-(aminomethyl)pyridine); BAT = N<sup>6</sup>,N<sup>9</sup>-bis(2-mercapto-2-methylpropyl)-6,9-diazanonanoic acid; BAT acid (protected) = N<sup>9</sup>-(*t*-butoxycarbonyl)-N<sup>6</sup>,N<sup>9</sup>-bis(2-methyl-2-triphenylmethylthiopropyl)-6,9-diazanonanoic acid; BAM = N<sup>1</sup>,N<sup>4</sup>-bis(2-mercapto-2-methylpropyl)-1,4,10-triazadecane; BAM (protected) = N<sup>1</sup>-(*t*-butoxycarbonyl)-N<sup>1</sup>,N<sup>4</sup>-bis(2-methyl-2-triphenylmethylthiopropyl)-1,4,10-triazadecane; (BAT-BM) = N-(2-(N',N'-bis(2-maleimidoethyl)aminoethyl)-N<sup>9</sup>-(*t*-butoxycarbonyl)-N<sup>6</sup>,N<sup>9</sup>-bis(2-methyl-2-triphenylmethylthiopropyl)-6,9-diazanonanamide; (BAT-BS) = N-(2-(N',N'-bis(2-succinimidoethyl)aminoethyl)-N<sup>6</sup>,N<sup>9</sup>-bis(2-mercapto-2-
- 15
- 20
- 25
- 30
- 35

methylpropyl)-6,9-diazanonanamide; (BMME) = *bis*-maleimidomethylether; (BSME) = *bis*-succinimidomethylether; (DTPA) = diethylenetriaminepentaacetic acid.  
RCY(%) = radiochemical yield (determined by HPLC)

5 Non-radioactive rhenium complexes were prepared by co-dissolving each of the peptide reagents of the invention with about one molar equivalent of tetrabutylammonium oxotetra-bromorhenate (+5), prepared as described by Cotton *et al.* (1966, *Inorg. Chem.* 5: 9-16) in dimethylformamide or acetonitrile/water and stirred for 0.5-5 days. The rhenium complexes were isolated by reverse phase  
10 HPLC as described above for Tc-99m labeled peptides and were characterized by FABMS or ESMS.

Radioactive rhenium complexes, using for example Re-186 or Re-188, are prepared from the appropriate perrhenate salts using the same protocol as for Tc-99m labeling, or by adding a reducing agent to a solution of the peptide and perrhenate,  
15 or optionally using a ligand transfer agent such as citrate and incubating the reaction at a temperature between room temperature and 100°C for between 5 and 60 min.

### EXAMPLE 3

#### 20 Inhibition of Binding of (<sup>125</sup>I-Tyr<sup>11</sup>)somatostatin-14 to AR42J Rat Pancreatic Tumor Cell Membranes

The ability of various somatostatin analogues of the invention to bind to somatostatin receptors *in vitro* was demonstrated by assaying the ability of such analogues to inhibit binding of a radiolabeled somatostatin analogue to somatostatin receptor-containing cell membranes. The rat pancreatic tumor cell line AR42J which  
25 expresses the somatostatin receptor was cultured in Dulbecco's minimal essential media (DMEM) supplemented with 10% fetal bovine serum (FBS) and 8mM glutamine in a humidified 5% CO<sub>2</sub> atmosphere at 37°C in T-flasks. Harvested cells were homogenized in cold 50mM Tris-HCl buffer (pH 7.4) and the homogenate then centrifuged at 39,000g for 10min at 4°C. Pellets were washed once with buffer and  
30 then resuspended in an ice-cold solution of 10mM Tris-HCl (pH 7.4). Equal aliquots of this cell membrane preparation were incubated with (<sup>125</sup>I-Tyr<sup>11</sup>)somatostatin-14 (at a final concentration of 0.5nM and 750,000cpm/mL, at a specific activity of 2000Ci/mmol, Amersham<sup>TM</sup>, Arlington Heights, IL) and peptide at a final

concentration of from  $10^{-11}$ M to  $10^{-6}$ M in a solution of 50mM HEPES (pH 7.4) containing 1% bovine serum albumin (BSA), 5mM  $\text{MgCl}_2$ , Trasylol<sup>TM</sup> (200,000 International Units), bacitracin (0.02mg/mL) and phenylmethanesulfonylfluoride (0.02mg/mL) for 25min at 30°C. Using a filtration manifold, this mixture was  
5 filtered through a polyethyleneimine-washed GC/F filter (Whatman<sup>TM</sup>, Maidstone, England), and the residue remaining on the filter washed thrice with 5mL cold HEPES buffer. The filter and a sample of the filter washings were then counted in a gamma counter. To assess non-specific binding, the assay was performed in the  
10 presence of unlabeled somatostatin-14 at 200nM. Data analysis including Hill plots of the data provided inhibition constants (see Bylund & Yamamura, "Methods of receptor binding", in Methods in Neurotransmitter Receptor Analysis, Yamamura *et al.*, eds., Raven Press: New York, 1990).

These results are presented in the following Table. The data show that the peptides of the instant invention have a high affinity of binding for somatostatin  
15 receptors.

It should be understood that the foregoing disclosure emphasizes certain specific embodiments of the invention and that all modifications or alternatives equivalent thereto are within the spirit and scope of the invention as set forth in the appended claims.

TABLE II

|    | Peptide                                                                            | K <sub>i</sub> (nM) |
|----|------------------------------------------------------------------------------------|---------------------|
|    | C <sub>Ac</sub> GC <sub>Ac</sub> GGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide | <0.01               |
|    | (DTPA)F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                        | 0.24                |
| 5  | maGGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                | 0.25                |
|    | <i>cyclo</i> (N-Me)FYW <sub>D</sub> KV.Hcy(CH <sub>3</sub> CO.GGCKK.amide)         | 0.26                |
|    | Ac.C <sub>Ac</sub> GC <sub>Ac</sub> F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide | 0.73                |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTC <sub>Ac</sub> GC <sub>Ac</sub> .amide    | 0.85                |
|    | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                    | 1.3                 |
| 10 | AKCGGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                               | 1.4                 |
|    | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                      | 2.0                 |
|    | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KT.Nal.T(ε-K)GCKK.amide                | 2.0                 |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTGGGC <sub>Ac</sub> GC <sub>Ac</sub> .amide | 2.4                 |
|    | (DTPA).Aca. <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                    | 2.6                 |
| 15 | KDKD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide                   | 2.6                 |
|    | (2-ketogulonyl)F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide               | 2.7                 |
|    | (DTPA).(ε-K)GCF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                       | 3.3                 |
|    | Ac.CGCF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                               | 4.4                 |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTCGC.amide                                  | 4.8                 |
| 20 | K <sub>D</sub> KKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide     | 4.9                 |
|    | Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                          | 5.6                 |
|    | Ac.KKKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                   | 6.5                 |
|    | KKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCDDDD.amide                   | 6.9                 |
|    | (DTPA)(Nal <sub>D</sub> .CYW <sub>D</sub> KVCT) <sub>2</sub>                       | 7.2                 |
| 25 | Ac.KKKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                 | 7.7                 |
|    | Ac.F <sub>D</sub> .FYW <sub>D</sub> KTFT(ε-K)GC.amide                              | 7.9                 |
|    | Ac.F <sub>D</sub> .FYW <sub>D</sub> KTFTGGG(ε-K)GC.amide                           | 8.2                 |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC.Nal.amide                                  | 8.2                 |
| 30 | K(BAT).Nal <sub>D</sub> .C <sub>Me</sub> YW <sub>D</sub> KVC <sub>Me</sub> T.amide | 9.9                 |

TABLE II (cont'd.)

|    | <u>(Re=O)-Complexed Peptides</u>                                                          | <u>MH<sup>+</sup></u> | <u>K<sub>i</sub> (nM)</u> |
|----|-------------------------------------------------------------------------------------------|-----------------------|---------------------------|
| 5  | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC(ε-K)GCKK.amide                                    | 1917                  | 0.13                      |
|    | Ac.D <sub>D</sub> F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC(ε-K)GCKK.amide                  | 2074                  | 0.20                      |
|    | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                           | 2343                  | 0.33                      |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC(ε-K)GC.amide                                      | N.D.                  | 0.36                      |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTC(ε-K)CGC.amide                                     | 1635                  | 0.37                      |
| 10 | F <sub>D</sub> FYW <sub>D</sub> KTFTGGC.amide                                             | 1683                  | 0.37                      |
|    | F <sub>D</sub> FYW <sub>D</sub> KTFTGGCK.amide                                            | 2032                  | 0.38                      |
|    | C <sub>Acm</sub> GC <sub>Acm</sub> GGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide      | 1807                  | 0.43                      |
|    | D <sub>D</sub> DF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                   | 2147                  | 0.50                      |
|    | F <sub>D</sub> FYW <sub>D</sub> KTFTC <sub>Acm</sub> GC <sub>Acm</sub> .amide             | 1601                  | 0.58                      |
| 15 | Ac.KKKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                        | 2649                  | 0.63                      |
|    | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                           | 2393                  | 0.67                      |
|    | AKCGGGF <sub>D</sub> FYW <sub>D</sub> KTFT.amide                                          | 1812                  | 0.76                      |
|    | KKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCDDDD.amide                          | 2683                  | 0.83                      |
|    | maGGGF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                       | 1618                  | 0.97                      |
| 20 | F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCR.amide                                    | 1817                  | 1.3                       |
|    | Ac.D <sub>D</sub> DF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                | 2188                  | 1.4                       |
|    | DDD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                             | 2197                  | 1.4                       |
|    | KDKD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide                          | 2083                  | 1.4                       |
|    | Ac.F <sub>D</sub> FYW <sub>D</sub> KTFT(ε-K)GC.amide                                      | 1688                  | 1.5                       |
| 25 | K <sub>D</sub> KK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCDDD.amide              | 2440                  | 1.5                       |
|    | K <sub>D</sub> KK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCDD.amide               | 2453                  | 1.6                       |
|    | Ac.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                              | 2008                  | 1.9                       |
|    | Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                                 | 1967                  | 2.2                       |
|    | AKCGGGF <sub>D</sub> FYW <sub>D</sub> KTFT.amide                                          | 1812                  | 2.9                       |
| 30 | (DTPA)Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFTC <sub>Acm</sub> GC <sub>Acm</sub> .amide | 2061                  | 3.1                       |
|    | F <sub>D</sub> .Cpa.YW <sub>D</sub> K.Abu.Nal.T(ε-K)GC.amide                              | 1695                  | 3.3                       |

TABLE II (cont'd.)

|    | <u>(Re=O)-Complexed Peptides</u>                                                     | <u>MH<sup>+</sup></u> | <u>K<sub>i</sub> (nM)</u> |
|----|--------------------------------------------------------------------------------------|-----------------------|---------------------------|
|    | (2-ketogulonyl)F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                 | 1837                  | 3.7                       |
| 5  | K <sub>D</sub> KKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKDKD.amide       | 2684                  | 3.8                       |
|    | Ac.CGCF <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT.amide                                 | 1677                  | 4.1                       |
|    | F <sub>D</sub> FYW <sub>D</sub> KTFT(ε-K)GC.amide                                    | 1637                  | 4.3                       |
|    | Ac.KKKKK.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                     | 2394                  | 4.4                       |
|    | (Trc.imide) <sub>2</sub> K.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCRR.amide | 2432                  | 4.9                       |
| 10 | (2-ketogulonyl)F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide               | 2143                  | 5.2                       |
|    | Ac.DDD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                     | 2239                  | 6.0                       |
|    | Ac.F <sub>D</sub> FYW <sub>D</sub> KTFTGGG(ε-K)KC.amide                              | 1911                  | 6.1                       |
|    | (DTPA).F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                         | 2036                  | 7.9                       |
|    | Hca.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKK.amide                        | 2298                  | 8.0                       |
| 15 | K <sub>D</sub> KKKF <sub>D</sub> .Cpa.YW <sub>D</sub> KTF.Nal.(ε-K)GCDDDD.amide      | 2730                  | 8.1                       |
|    | Ac.F <sub>D</sub> FYW <sub>D</sub> KTFTGGG(ε-K)GC.amide                              | 1840                  | 8.1                       |
|    | (DTPA).Aca.F <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                     | 2149                  | 8.2                       |
|    | DDDD.Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GCKKKK.amide                     | 2674                  | 9.8                       |
| 20 | (DTPA).Nal <sub>D</sub> .Cpa.YW <sub>D</sub> KTFT(ε-K)GC.amide                       | 2085                  | 11                        |
| 25 |                                                                                      |                       |                           |

**THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:**

1. Use of a composition of matter that is somatostatin receptor-binding peptide reagent having the formula:



wherein;  $X^1$  and  $X^2$  are each independently hydrophilic moieties having a molecular weight not greater than 1500 Daltons, wherein  $X^1$  is selected from the group consisting of an amino acid, a peptide having an amino acid sequence of no more than ten residues, a monosaccharide, an oligosaccharide comprising ten or fewer saccharide units, a poly(N-carboxyalkyl)amine, or a polyoxyanion, and  $X^2$  is selected from the group consisting of a poly(N-carboxyalkyl)amine, a polyoxyanion, an amino acid, a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxy-terminal amino acid is reduced to an alcohol), a mono-saccharide or an oligosaccharide comprising ten or fewer saccharide units;

$A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, or S-alkylated cysteine, penicillamine, homocysteine or homohomocysteine;

$B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;

$B^2$  is D- or L-Trp or substituted derivatives thereof;

$B^3$  is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

$B^4$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

$C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof,



for preparing a medicament for alleviating a somatostatin-related disease in an animal, wherein the medicament comprises a therapeutically effective amount of the somatostatin receptor binding peptide reagent.

2. A use according to Claim 1 wherein  $X^1$  is an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, or a poly(*N*-carboxyalkyl)amine, or a polyoxyanion, and  $X^2$  is a poly(*N*-carboxyalkyl)amine or polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxyl-terminal amino acid is reduced to an alcohol), or a monosaccharide or an oligosaccharide comprising ten or fewer saccharide units in the somatostatin receptor binding peptide reagent.
3. A use according to Claim 1 wherein  $B^1$  is phenylalanine or tyrosine,  $B^2$  is D-tryptophan,  $B^3$  is lysine and  $B^4$  is threonine or valine in the somatostatin receptor binding peptide reagent.
4. Use of a multimeric reagent having the formula:
  - a) at least two somatostatin receptor binding peptides, each of said peptides having a formula:



wherein;  $X^1$  and  $X^2$  are each independently hydrophilic moieties, having a molecular weight not greater than 1500 Daltons, wherein  $X^1$  is selected from the group consisting of an amino acid, a peptide, a monosaccharide, an oligosaccharide, a poly(*N*-carboxyalkyl)amine, and a polyoxyanion, and  $X^2$  is selected from the group consisting of a poly(*N*-carboxyalkyl)amine, a polyoxyanion, an amino acid, a peptide, a monosaccharide and an oligosaccharide;

A<sup>1</sup>, A<sup>2</sup> and C<sup>1</sup> are each independently a lipophilic D- or L-amino acid, or S-alkylated cysteine, penicillamine, homocysteine or homohomocysteine; B<sup>1</sup> is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;

B<sup>2</sup> is D- or L-Trp or substituted derivatives thereof;

B<sup>3</sup> is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

B<sup>4</sup> is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

C<sup>2</sup> is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof; and

- b) a polyvalent linker capable of covalently bonding to said peptides, for preparing a medicament for alleviating a somatostatin-related disease in an animal.

5. A use according to Claim 4 wherein the polyvalent linking moiety is *bis*-succinimidylmethylether, 4-(2,2-dimethylacetyl)benzoic acid, *N*-(2-(*N'*,*N'*-*bis*(2-succinimidoethyl)aminoethyl))-*N*<sup>6</sup>,*N*<sup>9</sup>-*bis*(2-methyl-2-mercaptopropyl)-6,9-diazanonanamide, *tris*(succinimidylethyl)amine *tris*(2-chloroacetamidoethyl)amine, 1,2-*bis*-(2-(chloroacetamido) ethoxy)ethane, *tris*(acetamidoethyl)amine, *bis*-acetamidomethyl ether, *bis*-acetamidoethyl ether,  $\alpha,\epsilon$ -*bis*-acetyllysine, lysine, 1,8-*bis*-acetamido-3,6-dioxaoctane, or a derivative thereof.
6. A pharmaceutical composition comprising a pharmaceutically-acceptable carrier or diluent and a somatostatin receptor binding peptide having the formula:



wherein; X<sup>1</sup> and X<sup>2</sup> are each independently hydrophilic moieties having a molecular

weight not greater than 1500 Daltons, wherein  $X^1$  is selected from the group consisting of an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, a poly(N-carboxyalkyl)amine, or a polyoxyanion, and  $X^2$  is selected from the group consisting of a poly(N-carboxyalkyl)amine, or a polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxy-terminal amino acid is reduced to an alcohol), or a mono-saccharide or an oligosaccharide comprising ten or fewer saccharide units;

$A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, or S-alkylated cysteine, penicillamine, homocysteine or homohomocysteine;  $B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;

$B^2$  is D- or L-Trp or substituted derivatives thereof;

$B^3$  is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

$B^4$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

$C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof.

7. A use according to Claim 1 wherein the animal is a human.
8. A composition of matter that is a somatostatin receptor-binding peptide reagent having the formula:



wherein;  $X^1$  is H, lower alkyl or substituted alkyl, aryl or substituted aryl, alkanoyl

or substituted alkanoyl, aroyl or substituted aroyl, or a hydrophilic moiety having a molecular weight not greater than 1500 Daltons, selected from the group consisting of an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, a poly(N-carboxyalkyl)amine, or a polyoxyanion;

A<sup>1</sup>, A<sup>2</sup> and C<sup>1</sup> are each independently a lipophilic D- or L-amino acid, or *S*-alkylated cysteine, penicillamine, homocysteine or homohomocysteine; B<sup>1</sup> is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Aib or substituted derivatives thereof;

B<sup>2</sup> is D- or L-Trp or substituted derivatives thereof;

B<sup>3</sup> is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;


B<sup>4</sup> is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof;

C<sup>2</sup> is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof;

X<sup>2</sup> is -COOR<sup>9</sup>, -CH<sub>2</sub>OH, CH<sub>2</sub>COOR<sup>9</sup>, or -CON(R<sup>9</sup>)<sub>2</sub>, where each R<sup>9</sup> is independently H, lower linear or cyclic alkyl or substituted derivatives thereof or a hydrophilic moiety having a molecular weight not greater than 1500 Daltons, selected from the group consisting of a poly(N-carboxyalkyl)amine, or a polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxy-terminal amino acid is reduced to an alcohol), or a mono-saccharide or an oligosaccharide comprising ten or fewer saccharide units;

and wherein the somatostatin receptor binding peptide is covalently linked to a radiolabel-binding moiety, wherein the radiolabel-binding moiety is covalently linked to the moieties A<sup>1</sup>, A<sup>2</sup>, C<sup>1</sup> or C<sup>2</sup> or is incorporated into either X<sup>1</sup> or X<sup>2</sup> of the peptide, and wherein the

radiolabel binding moiety has the formula:



wherein  $(pgp)^S$  is H or a thiol protecting group and (aa) is any  $\alpha$ - or  $\beta$ -amino acid; or



wherein; A is H, HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC or R'''';

B is H, SH, -NHR''', -N(R''')-(amino acid or peptide), or R'''';

X is H, SH, -NHR''', -N(R''')-(amino acid or peptide) or R'''';

Z is H or R'''';

R', R'', R''' and R'''' are independently H or lower straight or branched chain or cyclic alkyl;

n is 0, 1 or 2;

and where B is -NHR''' or -N(R''')-(amino acid or peptide), X is SH, and n is 1 or 2;

where X is -NHR''' or -N(R''')-(amino acid or peptide), B is SH, and n is 1 or 2;

where B is H or R''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC, X is SH, and n is 0 or 1;

where A is H or R''', then where B is SH, X is -NHR''' or -N(R''')-(amino acid or peptide) and where X is SH, B is -NHR''' or -N(R''')-(amino acid or peptide);

where X is H or R''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC and B is SH;

where Z is methyl, X is methyl, A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC, B is SH and n is 0;

where B is SH and X is SH, n is not 0.

9. The reagent of Claim 8 wherein X<sup>1</sup> is an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, or a poly(*N*-carboxyalkyl)amine, or a polyoxyanion and X<sup>2</sup> is a poly(*N*-carboxyalkyl)amine or a polyoxyanion, or an amino acid, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units.
10. The somatostatin receptor-binding peptide of Claim 8 wherein B<sup>1</sup> is phenylalanine or tyrosine, B<sup>2</sup> is D-tryptophan, B<sup>3</sup> is lysine and B<sup>4</sup> is threonine or valine.
11. A multimeric reagent having a molecular weight of less than about 20,000 Daltons comprising:
  - a) at least two somatostatin receptor binding peptides, each of said peptides having a formula:



wherein; X<sup>1</sup> and X<sup>2</sup> are each independently hydrophilic moieties, having a molecular weight not greater than 1500 Daltons, wherein X<sup>1</sup> is selected from the group consisting of an amino acid, a peptide, a monosaccharide, an oligosaccharide, a poly(*N*-carboxyalkyl)amine, and a polyoxyanion, and X<sup>2</sup> is selected from the group consisting of a poly(*N*-carboxyalkyl) amine, a polyoxyanion, an amino acid, a peptide, a monosaccharide and an

oligosaccharide;

$A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, or *S*-alkylated cysteine, penicillamine, homocysteine or homohomocysteine;

$B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;


$B^2$  is D- or L-Trp or substituted derivatives thereof;

$B^3$  is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

$B^4$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

$C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof; and

- b) at least two radiolabel-binding moieties, each of which is covalently linked to the moieties  $A^1$ ,  $A^2$ ,  $C^1$  or  $C^2$  or is incorporated into either  $X^1$  or  $X^2$  of the peptide, and wherein the radiolabel binding moiety has the formula:



wherein  $(\text{pgp})^S$  is H or a thiol protecting group and (aa) is any  $\alpha$ - or  $\beta$ -amino acid; or



wherein; A is H, HOOC,  $H_2NOC$ , (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC or  $R''''$ ;

B is H, SH,  $-NHR'''$ ,  $-N(R''')$ -(amino acid or peptide), or  $R''''$ ;

X is H, SH,  $-NHR'''$ ,  $-N(R''')$ -(amino acid or peptide) or  $R''''$ ;

Z is H or R'''';

R', R'', R''' and R'''' are independently H or lower straight or branched chain or cyclic alkyl;

n is 0, 1 or 2;

and where B is -NHR''' or -N(R''')-(amino acid or peptide), X is SH, and n is 1 or 2;

where X is -NHR''' or -N(R''')-(amino acid or peptide), B is SH, and n is 1 or 2;

where B is H or R''''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC, X is SH, and n is 0 or 1;

where A is H or R''''', then where B is SH, X is -NHR''' or -N(R''')-(amino acid or peptide) and where X is SH, B is -NHR''' or -N(R''')-(amino acid or peptide);

where X is H or R''''', A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC and B is SH;

where Z is methyl, X is methyl, A is HOOC, H<sub>2</sub>NOC, (amino acid or peptide)-NHOC, (amino acid or peptide)-OOC, B is SH and n is 0;

where B is SH and X is SH, n is not 0; and


c) a polyvalent linker capable of covalently bonding to said peptides, or to said radiolabel binding moieties.

12. The reagent of Claim 11 wherein the polyvalent linking moiety is *bis*-succinimidylmethylether, 4-(2,2-dimethylacetyl)benzoic acid, *N*-(2-(*N'*,*N'*-*bis*(2-succinimidoethyl)aminoethyl))-*N'*,*N'*-*bis*(2-methyl-2-mercaptopropyl)-6,9-diazanonanamide, *tris*(succinimidylethyl)amine *tris*(2-chloroacetamidoethyl)amine, 1,2-*bis*-(2-(chloroacetamido) ethoxy)ethane, *tris*(acetamidoethyl)amine, *bis*-acetamidomethyl ether, *bis*-acetamidoethyl ether,  $\alpha,\epsilon$ -*bis*-acetyllysine, lysine, 1,8-*bis*-acetamido-3,6-dioxaoctane, or a substituted derivative thereof.
13. A scintigraphic imaging agent comprising the reagent of Claim 8 radiolabeled with technetium-99m in combination with a conventional medium for intravenous injection.
14. A scintigraphic imaging agent comprising the reagent of Claim 8 radiolabeled with indium-111, gallium-67 or gallium-68 in combination with a conventional medium for intravenous injection.
15. A scintigraphic imaging agent comprising the somatostatin receptor binding peptide of Claim 8 radiolabeled with iodine-123 or iodine-125 in combination with a conventional medium for intravenous injection.
16. A radiotherapeutic agent comprising the reagent of Claim 8 radiolabeled with a cytotoxic radioisotope selected from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, and bismuth-212 in combination with a conventional medium for intravenous injection.
17. A radiotherapeutic agent comprising the somatostatin receptor binding peptide of Claim 8 radiolabeled with iodine-125, iodine-131 or astatine-131 in combination with a conventional medium for intravenous injection.

18. A complex formed by reacting the reagent of Claim 8 with technetium-99m in the presence of a reducing agent.
19. The complex of Claim 18 ,wherein the reducing agent is selected from the group consisting of a dithionite ion, a stannous ion and a ferrous ion.
20. A complex formed by labeling the reagent of Claim 8 with technetium-99m by ligand exchange of a prereduced technetium-99m complex.
21. A composition of matter comprising the reagent of Claim 8 and a stannous ion.
22. A kit for preparing a radiopharmaceutical preparation, said kit comprising a sealed vial containing a predetermined quantity of the reagent of Claim 8 and a sufficient amount of reducing agent to label the reagent with technetium-99m.
23. A method for labeling a reagent according to Claim 8 comprising reacting the reagent with technetium-99m in the presence of a reducing agent.
24. The method of Claim 23, wherein the reducing agent is selected from the group consisting of a dithionite ion, a stannous ion and a ferrous ion.
25. Use of the agent of Claim 13 for preparing a medicament for imaging a site within a mammalian body, wherein the medicament comprises an effective diagnostic amount of

the agent of Claim 13.

26. The reagent according to Claim 8 wherein the somatostatin receptor-binding peptide is chemically synthesized *in vitro*.
27. The reagent according to Claim 26 wherein the somatostatin receptor-binding peptide is synthesized by solid phase peptide synthesis.
28. The reagent according to Claim 26 wherein the radiolabel-binding moiety is covalently linked to the somatostatin receptor-binding peptide during *in vitro* chemical synthesis.
29. The reagent according to Claim 28 wherein the radiolabel-binding moiety is covalently linked to the somatostatin receptor-binding peptide during solid phase peptide synthesis.
30. The reagent of Claim 8 wherein the cysteine of the radiolabel-binding moiety having formula




has a protecting group of the formula




wherein; R is a lower alkyl having 1 to 6 carbon atoms, 2-,3-,4-pyridyl, phenyl, or phenyl substituted with lower alkyl, hydroxy, lower alkoxy, carboxy, or lower alkoxycarbonyl.

31. The reagent of Claim 8 wherein the radiolabel-binding moiety  $\text{C(pgp)}^{\text{S}}\text{-(aa)}\text{-C(pgp)}^{\text{S}}$  has

the formula:



32. The reagent of Claim 8 radiolabeled with technetium-99m.
33. The reagent of Claim 8 radiolabeled with indium-111, gallium-67 or gallium-68.
34. The reagent of Claim 8 radiolabeled with iodine-123.
35. The reagent of Claim 8 radiolabeled with a radioisotope selected from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, astatine-211, and bismuth-212.
36. The reagent of Claim 8 radiolabeled with iodine-125, iodine-131 or astatine-211.
37. A composition of matter comprising a somatostatin receptor-binding peptide reagent selected from the group consisting of reagents having the formula:



Ac.C<sub>Acm</sub>GC<sub>Acm</sub>F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFTC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 (DTPA).D-Nal.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 AKCGGGF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 (DTPA).D-Nal.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.GGGC<sub>Acm</sub>GC<sub>Acm</sub>.amide  
 (DTPA).Aca.F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 (DTPA).(ε-K)GCF<sub>D</sub>.FYW<sub>D</sub>KTFT.amide  
 Ac.CGCF<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFTCGC.amide  
 (DTPA).(D-Nal.CYW<sub>D</sub>KVCT)<sub>2</sub>  
 Ac.F<sub>D</sub>.FYW<sub>D</sub>KTFT(ε-K)GC.amide  
 Ac.F<sub>D</sub>FYW<sub>D</sub>KTFTGGG(ε-K)GC.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTC.Nal.amide  
 Ac.F<sub>D</sub>FYW<sub>D</sub>KTFGGG(ε-K)KC.amide  
 (DTPA).D-Nal.CYW<sub>D</sub>KVCT.amide  
 (2-ketogulonyl)-D-NalFYW<sub>D</sub>KVCT.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>K.Abu.Nal.T(ε-K)GC.amide  
 F<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GC.amide  
 (DTPA).F<sub>D</sub>FYW<sub>D</sub>KTFT(ε-K)GC.amide  
 AF<sub>D</sub>CFW<sub>D</sub>KTC<sub>Me</sub>T(CH<sub>2</sub>OH)  
 (DTPA).F<sub>D</sub>GYW<sub>D</sub>KTCT(CH<sub>2</sub>OH)  
 (DTPA).Nal.SYW<sub>D</sub>KVCT.amide  
 F<sub>D</sub>FYW<sub>D</sub>KTFTGGCK.amide  
 DDD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 Ac.DDD.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide  
 Hca.G.Nal<sub>D</sub>.Cpa.YW<sub>D</sub>KTFT(ε-K)GCKK.amide

$F_D FYW_D KTFT C_{Acm} GC_{Acm}.amide$   
 $F_D FYW_D KTFT GGC.amide$   
 $F_D FYW_D KTFT(\epsilon-K)GC.amide$   
 $(Trc.imide)_2 K.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCRR.amide$   
 $Trc(Trc.imide)K.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCRR.amide$   
 $(Trc.imide)Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCR.amide$   
 $F_D.Cpa.YW_D KTFT(\epsilon-K)GCR.amide$   
 $K_D KKK.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKDKD.amide$   
 $K_D KKK.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCDD.amide$   
 $(Trc)_2 K.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $Hca.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $(2-ketogulonyl)F_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $KKKK.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCDDDD.amide$   
 $Ac.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $Ac.KKKKK.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $(2-ketogulonyl)F_D.Cpa.YW_D KTFT(\epsilon-K)GC.amide$   
 $Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $DDDD.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKKKK.amide$   
 $(DTPA)Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKK.amide$   
 $(DTPA)Nal_D.Cpa.YW_D KTFT C_{Acm} GC_{Acm}.amide$   
 $Ac.KKKKK.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GC.amide$   
 $KDKD.Nal_D.Cpa.YW_D KTFT(\epsilon-K)GCKDKD.amide$

38. The composition of matter of Claim 37 radiolabeled with a radioisotope selected from the group consisting of gallium-68, technetium-99m, indium-111, and iodine-123.
39. The composition of matter of Claim 37 radiolabeled with a radioisotope selected from the group consisting of scandium-47, copper-67, gallium-72, yttrium-90, iodine-125, iodine-131, samarium-153, gadolinium-159, dysprosium-165, holmium-166, ytterbium-175, lutetium-177, rhenium-186, rhenium-188, bismuth-212 and astatine-211.
40. The use according to Claim 1 wherein the therapeutically effective amount of the composition is from about 0.1 to about 49 mg/kg body weight/day.
41. Use of the radiotherapeutic agent of Claim 16 for preparing a medicament for alleviating a somatostatin-related disease in an animal wherein the medicament comprises a therapeutically effective amount of the agent of Claim 16.
42. The use according to Claim 41 wherein the animal is a human.
43. The use according to Claim 41 wherein the therapeutically effective amount of the composition is from about 10 to about 200mCi.
44. A pharmaceutical composition comprising the radiolabeled reagent of Claim 8 in a pharmaceutically-acceptable carrier.
45. A composition of matter comprising a complex formed by reacting the reagent of Claim 8 with a non-radioactive metal.

46. The complex of Claim 45 wherein the non-radioactive metal is rhenium.
47. A composition of matter comprising a complex formed by reacting the agent of Claim 13 with a non-radioactive metal.
48. A composition of matter comprising a complex formed by reacting the scintigraphic imaging agent of Claim 13 with a non-radioactive metal.
49. A composition of matter comprising a complex formed by reacting the scintigraphic imaging agent of Claim 14 with a non-radioactive metal.
50. A composition of matter comprising a complex formed by reacting the scintigraphic imaging agent of Claim 15 with a non-radioactive metal.
51. A composition of matter comprising a complex formed by reacting the radiotherapeutic agent of Claim 16 with a non-radioactive metal.
52. A composition of matter comprising a complex formed by reacting the radiotherapeutic agent of Claim 17 with a non-radioactive metal.
53. Use of the reagent of Claim 8 for preparing a medicament for imaging a site within a mammalian body wherein the medicament comprises an effective diagnostic amount of the reagent of Claim 8 radiolabeled with a detectable radioisotope.

54. A scintigraphic imaging agent comprising a somatostatin receptor-binding peptide reagent having the formula:



wherein;  $X^1$  and  $X^2$  are each independently hydrophilic moieties having a molecular weight not greater than 1500 Daltons, wherein  $X^1$  is selected from the group consisting of an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, a poly(N-carboxyalkyl)amine, or a polyoxyanion, and  $X^2$  is selected from the group consisting of a poly(N-carboxyalkyl)amine, or a polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxy-terminal amino acid is reduced to an alcohol), or a mono-saccharide or an oligosaccharide comprising ten or fewer saccharide units;

$A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, or *S*-alkylated cysteine, penicillamine, homocysteine or homohomocysteine;

$B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;

$B^2$  is D- or L-Trp or substituted derivatives thereof;

$B^3$  is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

$B^4$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

$C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof,

radiolabeled with iodine-123 or iodine-125.

55. A radiotherapeutic agent comprising a somatostatin receptor-binding peptide reagent having the formula:



wherein;  $X^1$  and  $X^2$  are each independently hydrophilic moieties having a molecular weight not greater than 1500 Daltons, wherein  $X^1$  is selected from the group consisting of an amino acid, or a peptide having an amino acid sequence of no more than ten residues, or a monosaccharide, or an oligosaccharide comprising ten or fewer saccharide units, a poly(N-carboxyalkyl)amine, or a polyoxyanion, and  $X^2$  is selected from the group consisting of a poly(N-carboxyalkyl)amine, or a polyoxyanion, or an amino acid, or a peptide having an amino acid sequence of no more than ten residues (including peptides wherein the carboxyl group of the carboxy-terminal amino acid is reduced to an alcohol), or a mono-saccharide or an oligosaccharide comprising ten or fewer saccharide units;

$A^1$ ,  $A^2$  and  $C^1$  are each independently a lipophilic D- or L-amino acid, or S-alkylated cysteine, penicillamine, homocysteine or homohomocysteine;

$B^1$  is D- or L-Phe, or D- or L-Tyr, or D- or L-Nal, or Ain or substituted derivatives thereof;

$B^2$  is D- or L-Trp or substituted derivatives thereof;

$B^3$  is D- or L-Lys, or Hly, Achxa, Amf, Aec, Apc, Aes, Aps or substituted derivatives thereof;

$B^4$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva or Aib or substituted derivatives thereof; and

$C^2$  is D- or L-Thr, Ser, Val, Phe, Ile, Abu, Nle, Leu, Nva, Nal or Aib or substituted derivatives thereof,

radiolabeled with iodine-125, iodine-131 or astatine-131.