




(86) Date de dépôt PCT/PCT Filing Date: 1999/03/26
(87) Date publication PCT/PCT Publication Date: 1999/09/30
(45) Date de délivrance/Issue Date: 2007/05/15
(85) Entrée phase nationale/National Entry: 2000/09/25
(86) N° demande PCT/PCT Application No.: EP 1999/002337
(87) N° publication PCT/PCT Publication No.: 1999/048500
(30) Priorité/Priority: 1998/03/26 (EP98400723.7)

(51) Cl.Int./Int.Cl. *A61K 31/485* (2006.01),
A61K 31/185 (2006.01)
(72) Inventeurs/Inventors:
DAOUST, MARTINE, FR;
BONHOMME, YVES, FR;
DURBIN, PHILIPPE, FR
(73) Propriétaire/Owner:
MERCK SANTE, FR
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : COMBINAISON DESTINEE AU TRAITEMENT DE LA DEPENDANCE A L'ALCOOL ET AUX DROGUES
CONTENANT UN ANTAGONISTE D'OPIOIDES ET UN MODULATEUR DU COMPLEXE DU RECEPTEUR NMDA
(54) Title: COMBINATION FOR THE TREATMENT OF ALCOHOL AND DRUG DEPENDENCE CONTAINING AN OPIOID
ANTAGONIST AND A NMDA RECEPTOR COMPLEX MODULATOR

ALCOHOL INTAKE



(57) **Abrégé/Abstract:**

The present invention relates to a pharmaceutical composition for the treatment of alcohol and drug dependence comprising a therapeutically effective amount of a combination of: (i) an opioid antagonist; and (ii) a NMDA receptor complex modulator, and a pharmaceutical kit comprising: (i) an opioid antagonist; and (ii) a NMDA receptor complex modulator.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/485 // (A61K 31/485, 31:185)	A1	(11) International Publication Number: WO 99/48500 (43) International Publication Date: 30 September 1999 (30.09.99)
(21) International Application Number: PCT/EP99/02337 (22) International Filing Date: 26 March 1999 (26.03.99) (30) Priority Data: 98400723.7 26 March 1998 (26.03.98) EP (71) Applicant (for all designated States except US): LIPHA [FR/FR]; 34, rue Saint Romain, F-69008 Lyon (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): DAOUST, Martine [FR/FR]; 7, allée Alphonse Daudet, F-76690 Notre Dame de Bondeville (FR). BONHOMME, Yves [FR/FR]; Le Buclay, 21, avenue de la Paix, F-69260 Charbonnières Les Bains (FR). DURBIN, Philippe [FR/FR]; 83, rue Paul Verlaine, F-69100 Villeurbanne (FR). (74) Agent: MONCHENY, Michel; Cabinet Lavoix, 2, place d'Estiennes d'Orves, F-75441 Paris Cedex 09 (FR).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: COMBINATION FOR THE TREATMENT OF ALCOHOL AND DRUG DEPENDENCE CONTAINING AN OPIOID ANTAGONIST AND A NMDA RECEPTOR COMPLEX MODULATOR (57) Abstract The present invention relates to a pharmaceutical composition for the treatment of alcohol and drug dependence comprising a therapeutically effective amount of a combination of: (i) an opioid antagonist; and (ii) a NMDA receptor complex modulator, and a pharmaceutical kit comprising: (i) an opioid antagonist; and (ii) a NMDA receptor complex modulator.		

WO 99/48500

PCT/EP99/02337

COMBINATION FOR THE TREATMENT OF ALCOHOL AND DRUG DEPENDENCE CONTAINING AN OPIOID ANTAGONIST AND A NMDA RECEPTOR COMPLEX MODULATOR

The present invention relates to a method for the treatment of alcohol and
5 drug dependence.

More specifically the present invention relates to a method for the treatment of the alcohol dependence.

It relates also to a method for the treatment of a dependence to drugs such as opioid derivatives, cannabinoids, nicotin derivatives, amphetamines
10 and tranquilizers.

The present invention also relates to compositions and kits for such treatment.

A number of studies indicated a relationship between alcohol intake and endogenous opioid activity (Schulz R., Wuster M., Duka T., Herz A., Acute and
15 chronic ethanol treatment changes endorphin levels in brain and pituitary, Psychopharmacology, 68, 221-227, 1980 ; Hoffman P., Melchior C., Ritzmann R.F., Tabakoff B., Structural requirements for neurohypophyseal peptide effects of ethanol tolerance, Alcohol Exp. Clin. Res., 5, 154, 1981). Ethanol intake increases synthesis and release of beta endorphin in the pituitary gland and the
20 stimulation of opiate receptors seems to be weakly involved in the regulation of positive reinforcement properties of ethanol. These findings have suggested the use of opiates antagonists (such as naloxone or naltrexone) to prevent ethanol-induced analgesia, intoxication, coma (Kiianmaa K., Tabakoff B.
Neurochemical correlates of tolerance and strain differences in the
25 neurochemical effects of ethanol, Pharm. Biochem. Behav., 18, 383-388, 1983) or more recently ethanol dependence (Davidson D., Amit Z., Naltrexone blocks acquisition of voluntary ethanol intake in rats, Alcohol Clin. Exp. Res., 21, 677-683, 1997). Naltrexone, when used together with supportive therapy, has been shown to reduce the rate of relapse to heavy drinking in alcohol dependent
30 patients (O'Malley S.S., Opioid antagonists in the treatment of alcohol dependence : clinical efficacy and prevention of relapse, Alcohol Alcoholism 31, 77-91, 1996).

The present invention provides a method for the treatment of alcohol and drug dependence comprising administering to a patient a therapeutically effective amount of a combination of : (i) an opioid antagonist, such as naltrexone or naloxone ; and (ii) a NMDA receptor complex modulator, in particular a spermidine site modulator. The present invention is also directed to compositions and pharmaceutical kits containing the same. Such combination therapy provides surprisingly efficient and effective methodology for use in the treatment of alcohol and drug dependence.

The NMDA subtype of the glutamate receptor is a ligand-gated ion channel involved in excitatory neurotransmission in the mammalian CNS. Activation of the NMDA receptor-channel complex has been implicated in several physiological phenomena important to higher order CNS functions. Overstimulation of this receptor results in an inflow of Ca^{++} and neuronal excitotoxicity. This ligand-gated ionotropic glutamate receptor is subject to complex regulation by a number of ligands. Separate regulatory sites include the binding site for the agonist L-glutamate ; a high affinity binding site for the obligatory co-agonist glycine ; a site where Zn^{++} acts to allosterically inhibit the agonist-induced response independently of membrane potential ; a site within the channel where Mg^{++} and phencyclidine (PCP), dizocilpine and ketamine bind to produce a voltage-dependent open channel block ; and a distinct binding site for the endogenous polyamines, spermine and spermidine, which modulate NMDA receptor function (Bergeron et al., J. Med. Chem., 39, 5257, 1996).

Spermidine can modulate certain NMDA receptor subtypes by either glycine-independent or glycine dependent mechanisms (or both). This action of polyamines to potentiate agonist-mediated responses may reside in its shielding of the NMDA receptor proton sensor from the extracellular pH. The influence of polyamines *in vivo* will depend on effective extracellular levels and the presence of suitable polyamine sensitive NMDA receptors sub-units. These *in vivo* effects of polyamines on NMDA receptor may explain the excessive activation of NMDA receptors observed during some pathological situations in which polyamine synthesis is not preserved.

Acamprosate is an example of a compound which is able to induce a modulation of the NMDA receptor complex by interactions on a spermidine sensitive site.

Other examples of compounds which induce a modulation of the NMDA receptor complex are :

- glycine antagonists such as L 701 324 (J. Kotlinska et al., Psychopharmacology, 127, 238, 1886) ;
- ibogaine : (Popick P. et al., J. Pharmacol. Exp. Ther., 275, 753, 1995),
- memantine and derivatives : (Popick P. et al., Pharmacol. Biochem. Behav., 53, 791, 1996),
- ifenprodil and eliprodil : (Schoemaker II. et al., Eur. J. Pharmacol., 176, 240, 1990),
- modulators of the NMDA receptor complex on the glycine site (Bienkowski P., Alcohol, 15, 87, 1998).

By "administering a combination", or the like, when referring to component (i), and component (ii), of the present invention, it is meant that the components are administered concurrently to a patient being treated. By concurrently, it is meant that each component may be administered at the same time or sequentially in any order at different points in time. However if not administered at the same time, they should be administered sufficiently closely in time so as to provide the desired treatment effect. Suitable dosing intervals and dosing order with such compounds will be readily apparent to those skilled in the art, once armed with the present disclosure. Preferably, all components are administered at the same time, and if not administered at the same time, preferably they are all administered less than one hour apart from one another.

The present invention also includes pharmaceutical compositions comprising or consisting essentially of, in combination, an opioid antagonist (such as naltrexone), and a NMDA receptor complex modulator. Such compositions may be in solid, liquid, transdermal, transnasal, or depot dosage units and may further include a suitable pharmaceutical carrier. Examples of compositions for the oral route are coated tablets and capsules.

20497-759(S)

4

The present invention also includes pharmaceutical kits comprising or consisting essentially of an opioid antagonist (such as naltrexone), together with a NMDA receptor complex modulator. In the kit, the opioid antagonist and the NMDA receptor complex modulator may each be presented in separate vials as compounds, and/or in separate vials as compounds in combination with a pharmaceutically acceptable carrier. Alternatively, the opioid antagonist and the NMDA receptor complex modulator may be combined together in one or more vials, with or without a carrier. Thus, for example, the invention includes pharmaceutical kits comprising a separate vial comprising the opioid antagonist and a separate vial comprising the NMDA receptor complex modulator, each vial also containing, if desired, a carrier.

The compositions and kits of the present invention may be employed in the treatment of alcohol and drug dependence.

For use in the treatment of diseases characterized by abnormally high consumption of alcohol, a daily dosage of active ingredients can be about 0.5 to 20 mg/kg of body weight for the opioid antagonist such as naltrexone and about 10 to 400 mg/kg of body weight for a NMDA receptor complex modulator such as acamprosate.

In one aspect, the invention provides a pharmaceutical composition for the treatment of alcohol dependence comprising a therapeutically effective amount of a combination of: (i) an opioid antagonist which is either naloxone or naltrexone; and (ii) a NMDA receptor complex modulator which is acamprosate.

The invention also provides uses of the above

20497-759(S)

4a

composition for: (i) preparing a medicament for the treatment of alcohol dependence, and (ii) treating alcohol dependence.

The invention also provides a commercial package
5 comprising the above composition and associated therewith instructions for the use thereof in the treatment of alcohol dependence.

Brief description of the drawings:

Fig. 1 is a graph showing the alcohol intake in a
10 first series of tests in rats.

The unexpected effect of the combinations is shown in the following tests.

Material and methods

Ethanol dependent rats were selected as previously
15 described (Boismare F., Daoust M., Moore N., Saligaut C., Chadelaud M., Chrétien P., Durlach J., Lhuintre J.P., A homotaurine derivative reduces the voluntary intake of ethanol by rats : are cerebral GABA receptors involved?, Pharm. Biochem. Behav., 21, 787-789, 1984)

- **Alcoholization**

Adult male Long Evans rats weighing 180 ± 20 g at the beginning of the experiment were obtained from "Centre d'élevage Janvier" (France). The rats were housed in individual cages and had free access to food (UAR, France standard diet). They were kept with an ambient temperature of 21° C and a 12 hr/12 hr light-dark photoperiod. During the initial selection period (14 days), they only had access to a 10% (v/v) ethanol solution, prepared from 95% ethanol and water as drinking fluid for 14 days. This period was followed by another two-week period during which they has a free choice between ethanol solution and water. The two fluids three bottles method was used to prevent fluid selection on the basis of bottle situation. Every other day, fluid intakes and body weights were measured, the drinking bottles were refilled and randomly rotated. Food consumption was also measured each week be weighting food pellets. Animals preferring ethanol 4-5 g/kg per day during these last two seeks ($\approx 30\%$ of the rats) were selected as drinking rats and used for the different treatments.

- **Treatments**

All the drugs were prepared as saline (NaCl 0.9%) solutions and animals were daily i.p. injected with : acamprosate 100 mg/kg, naltrexone 10 mg/kg, acamprosate 100 mg/kg + naltrexone 10 mg/kg or saline 1 ml/200 g body weight. Five groups of six or seven alcohol preferring rats were used. Each group receiving its own treatment. Rats were treated for 14 days (treatment period) and after that, they were kept one week more without treatment for the post-treatment analysis. Alcohol intakes were expressed as gram of absolute alcohol drunk per kilo of body weight per day (g.kg.d) for alcohol.

Results

Body weights and ethanol consumption were the same in both groups during the pre-treatment period ($p < 0.1$).

Ethanol intake significantly decreased in acamprosate ($p < 0.008$) and naltrexone ($p = 0.092$) groups. Acamprosate and naltrexone when are given together, present a significant effect on ethanol intake ($p < 0.001$).

The results are shown in Figure 1.

5 Figure 1 represents means $\pm$ sem of ethanol intake, expressed as gram of absolute alcohol during each period of pre-treatment, treatment and post-treatment in all groups of rats. 6 or 7 rats per group for : Control (Cont), Acamprosate 100 mg/kg/d (AC), Naltrexone 10 mg/kg/d (NTX), Acamprosate 100 mg/kg/d + Naltrexone 10 mg/kg/d (AC + NTX).

10 The most interesting finding is that the acamprosate and naltrexone have an unexpected effect on alcohol intake when they are administered together.

Examples of pharmaceutical compositions

15 Example 1 - tablet

A tablet is prepared with the following composition (mg/tablet) :

core :

	acamprosate	350
	naltrexone	25 to 50
20	microcristalline cellulose	25 to 75 (for example 50)
	PVP 30	10 to 25 (for example 15)
	croscarmellose	10 to 20 (for example 17)

external phase :

	croscarmellose	10-20 (for example 17)
25	microcristalline cellulose	25-75 (for example 50)
	lactose	75-125 (for example 100)
	magnesium stearate	9-15 (for exemple 12)
	anhydrous colloidal silica	1-3 (for example 2)

coating :

30	sepifilm (HPMC, TiO ₂ , stearic acid)	25-45 (for example 30).
----	--	-------------------------

The core of this tablet is obtained by wet granulation.

20497-759 (S)

7

Example 2 - capsule with immediate release

The capsule has the following composition (mg/capsule):

	acamprosate	350
	naltrexone	25
5	gelucire™ (polyglycosyl glyceride)	250-330 (for example 290)
	soya bean lecithine	1-10 (for example 7)
	sorbitanne trioleate	10-35 (for example 35).

Example 3- capsule with extended release

The capsule has the following composition (mg/capsule):

10	acamprosate	350
	naltrexone	25
	gelucire™ (polyglycosyl glyceride)	200-300 (for example 250)
	soya bean lecithine	1-10 (for example 7)
	sorbitanne trioleate	10-50 (for example 35)
15	precirol (glycerol stearate)	20-60 (for example 40).

20497-759 (S)

8


CLAIMS:

1. A pharmaceutical composition for the treatment of alcohol dependence comprising a therapeutically effective amount of a combination of:
 - 5 (i) an opioid antagonist which is either naloxone or naltrexone; and
 - (ii) a NMDA receptor complex modulator which is acamprosate.
2. A pharmaceutical composition as claimed in claim 1 in which said opioid antagonist is naltrexone.
- 10 3. Use of a pharmaceutical composition as claimed in claim 1 or 2 for preparing a medicament for the treatment of alcohol dependence.
4. Use of a pharmaceutical composition as claimed in claim 1 or 2 for the treatment of alcohol dependence.
- 15 5. A commercial package comprising a pharmaceutical composition as claimed in claim 1 or 2 and associated therewith instructions for the use thereof in the treatment of alcohol dependence.

FETHERSTONHAUGH & CO.
OTTAWA, CANADA
PATENT AGENTS

1/1

ALCOHOL INTAKE



ALCOHOL INTAKE

