



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2019/07/09
(87) Date publication PCT/PCT Publication Date: 2020/01/16
(85) Entrée phase nationale/National Entry: 2020/12/29
(86) N° demande PCT/PCT Application No.: IB 2019/055853
(87) N° publication PCT/PCT Publication No.: 2020/012365
(30) Priorité/Priority: 2018/07/09 (US62/695,388)

(51) Cl.Int./Int.Cl. *A01H 5/00* (2018.01),
A01H 5/10 (2018.01), *A01H 6/20* (2018.01),
C07K 14/415 (2006.01), *C12N 15/82* (2006.01)
(71) Demandeur/Applicant:
CELLECTIS, FR
(72) Inventeurs/Inventors:
ZHANG, WENZHENG, US;
ZHANG, FENG, US
(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : COLZA A FORTE TENEUR EN ACIDE OLEIQUE
(54) Title: CANOLA WITH HIGH OLEIC ACID

(57) **Abrégé/Abstract:**

Materials and methods for creating canola (e.g., Brassica napus) lines having oil with increased oleic acid content are provided herein.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date

16 January 2020 (16.01.2020)



(10) International Publication Number

WO 2020/012365 A1

(51) International Patent Classification:

A01H 5/00 (2018.01) C07K 14/415 (2006.01)

A01H 5/10 (2018.01) C12N 15/82 (2006.01)

A01H 6/20 (2018.01)

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))
- in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE

(21) International Application Number:

PCT/IB2019/055853

(22) International Filing Date:

09 July 2019 (09.07.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/695,388 09 July 2018 (09.07.2018) US

(71) Applicant: CELLECTIS [FR/FR]; 8 rue del Croix Jarry,
75013 Paris (FR).

(72) Inventors: ZHANG, Wenzheng; 9666 Dunning Circle,
Highlands Ranch, Colorado 80126 (US). ZHANG, Feng;
11780 62nd Court, Maple Grove, Minnesota 55311 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

(54) Title: CANOLA WITH HIGH OLEIC ACID

(57) Abstract: Materials and methods for creating canola (e.g., *Brassica napus*) lines having oil with increased oleic acid content are provided herein.



WO 2020/012365 A1

CANOLA WITH HIGH OLEIC ACID

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. Provisional Application Serial No.
5 62/695,388, filed July 9, 2018. The disclosure of the prior application is considered part of (and is incorporated by reference in) the disclosure of this application.

TECHNICAL FIELD

This document relates to materials and methods for generating novel canola plants
10 with improved oil characteristics. For example, this document relates to *Brassica* plants with novel mutations in fatty acid desaturase 2 (FAD2) gene copies.

BACKGROUND

Oilseed rape (*Brassica napus*) is an important crop that produces considerable
15 amounts of edible oil – with about 40% of the seed being oil. Oilseed rape is grown for animal feed, biodiesel, and edible vegetable oils. The edible and processing qualities of the seed oil is determined primarily by the fatty acid composition of the triglycerol lipids. Oilseed rape with low erucic acid (canola) has five predominant fatty acids within the seed oil: 16:0 (palmitic acid), 18:0 (stearic acid), 18:1 (oleic acid), 18:2 (linoleic acid),
20 and 18:3 (linolenic acid). Canola oil is valued, in part, for its saturated fatty acid content, as it has the lowest palmitic acid and stearic acid content of any vegetable oil, with about 6% of total fatty acids being saturated fatty acids.

The quality of canola oil also is determined by the relative composition of the mono- and polyunsaturated fatty acids, specifically, the relative composition of oleic acid,
25 linoleic acid, and linolenic acid. Conventional canola cultivars normally produce oil with ~60% oleic acid, ~20% linoleic acid, and ~10% linolenic acid. Oil with high oleic acid can have a longer shelf life and greater heat stability during frying (Warner et al., *J Am Oil Chem Soc* 74:1317-1322, 1997; Matthaus, *Eur J Lipid Sci Technol* 108:200-211, 2006). Oil with low linolenic acid also can be beneficial for storage and extending shelf
30 life (Scarth and Tang, *Crop Sci* 46:1225-1236).

SUMMARY

Genes involved within the fatty acid synthesis pathway in *Brassica* include the fatty acid desaturase genes. FAD2 catalyzes the conversion of oleic acid (18:1) to linoleic

acid (18:2). In *B. napus*, four copies of FAD2 genes have been identified (Yang et al., *Theor Appl Genet* 125:715-729, 2012). Due to limited studies on high oleic mutants, the individual contribution of most FAD2 gene copies on oil composition was not previously known. This document is based, at least in part, on the development of materials and methods for generating novel canola plants with improved oil characteristics. The resulting canola plants can be valuable to canola growers and the food industry. Thus, this document provides methods and compositions for using genome editing to generate modified Brassica plants that can be selected for enhanced oil characteristics, such as increased oleic acid content and decreased linolenic acid content. As described herein, Brassica plants with enhanced oil characteristics can be generated using sequence-specific nucleases to inactivate or attenuate fatty acid desaturase 2 (FAD2) gene copies.

In a first aspect, this document features a Brassica plant, plant part, or plant cell containing an induced mutation in one or more FAD2 gene copies, wherein the plant, plant part, or plant cell produces oil that has increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell. Each of the induced mutations can be within a nucleic acid sequence as set forth in any of SEQ ID NOS:1-3, or within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. The induced mutation can include a frameshift, where the frameshift includes a deletion selected from the group consisting of a deletion of the guanine at position 253 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 416 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 99 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and a deletion of the guanine at position 322 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. In some cases, the induced mutation can be an in-frame or frameshift mutation, where the in-frame or frameshift mutation includes an insertion or deletion selected from the group consisting of a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 3 bp deletion at positions 99-101 and a 1 bp insertion at

position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. Each induced mutation can have been induced by a rare-cutting endonuclease. In some embodiments, the plant, plant part, or plant cell may not contain a transgene. In some embodiments, the plant, plant part, or plant cell contains a transgene encoding a protein, where the protein is selected from the group consisting of a plant 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an aryloxyalkanoate dioxygenase (AAD) protein, a phosphinothricin N-acetyltransferase (PAT) protein, an acetohydroxyacid synthase large subunit protein, a p-hydroxyphenylpyruvate dioxygenase (hppd) protein, and a dicamba monooxygenase (DMO) protein. The Brassica plant, plant part, or plant cell can be a seed. In some cases, a mutated FAD2 gene copy can include the sequence set forth in any of SEQ ID NOS:53900-53975, or a sequence at least 90% identical to the sequence set forth in any of SEQ ID NOS:53900-53975.

In another aspect, this document features a Brassica plant or plant part that is a progeny of the Brassica plant or plant part described above, where the progeny plant or plant part has a mutation in at least one FAD2 gene copy.

In another aspect, this document features a method for producing a Brassica plant having increased oleic acid content and decreased linolenic acid content, where the method includes providing a population of Brassica cells containing functional FAD2 gene copies, contacting the population of Brassica cells with one or more rare-cutting endonucleases targeted to the one or more FAD2 gene copies, regenerating Brassica plants from the population of Brassica cells contacted with the one or more rare-cutting endonucleases, and selecting a Brassica plant with a mutation in one or more FAD2 gene copies, where oil produced by the selected Brassica plant has increased oleic acid content

and decreased linolenic acid content as compared to oil produced by a corresponding wild type Brassica plant. The Brassica cells can be selected from the group consisting of protoplast cells, embryo cells, callus cells, leaf cells, and petiole explant cells. The method can include transforming the Brassica cells with one or more vectors encoding the one or more rare-cutting endonucleases. The one or more rare-cutting endonucleases can be transcription activator-like effector (TALE) nucleases. Each of the one or more TALE nucleases can be targeted to a sequence within SEQ ID NOS:1-3, or to a sequence having at least 90% identity to a sequence within SEQ ID NOS:1-3. Each of the one or more TALE nucleases can be targeted to a sequence within any of the sequences set forth in SEQ ID NOS:9-29, or to a sequence having at least 90% identity to any of SEQ ID NOS:9-29. In some cases, a mutated FAD2 gene copy can include the sequence set forth in any of SEQ ID NOS:53900-53975, or a sequence at least 90% identical to the sequence set forth in any of SEQ ID NOS:53900-53975.

In another embodiment, this document features a method for generating a Brassica plant that produces oil having increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell, where the method includes providing a first Brassica plant having at least one induced mutation in an endogenous FAD2 gene copy; providing a second Brassica plant having wild type FAD2 gene copies; and crossing the first Brassica plant with the second Brassica plant, thereby producing a plurality of progeny seed, wherein the progeny seed produce plants that contain an induced mutation in at least one endogenous FAD2 gene copy. Each of the induced mutations can be at a sequence within any of SEQ ID NOS:1-3. The induced mutation can include a frameshift, where the frameshift includes a deletion selected from the group consisting of a deletion of the guanine at position 253 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 416 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 99 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and a deletion of the guanine at position 322 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. In some cases, the induced mutation can be an in-frame or frameshift mutation, where the in-frame or frameshift mutation includes an insertion or deletion selected from the group consisting of

a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. Each induced mutation can have been induced by a rare-cutting endonuclease. In some embodiments, the first Brassica plant and the second Brassica plant do not contain a transgene. In some embodiments, the first Brassica plant and/or the second Brassica plant contains a transgene encoding a protein, where the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid synthase large subunit protein, a hppd protein, and a DMO protein. In some cases, a mutated FAD2 gene copy can include the sequence set forth in any of SEQ ID NOS:53900-53975, or a sequence at least 90% identical to the sequence set forth in any of SEQ ID NOS:53900-53975.

In yet another aspect, this document features a method for generating a Brassica plant that produces seed oil having increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell, where the method includes providing a first Brassica plant having at least a first induced mutation in a first endogenous FAD2 gene copy, providing a second Brassica plant having at least a second induced mutation in a second endogenous FAD2 gene copy, and crossing the first Brassica plant with the second Brassica plant, thereby producing a plurality of progeny seed, wherein the progeny seed produce plants that contain the first and second induced mutations in the first and second endogenous

FAD2 gene copies. Each of the induced mutations can be at a sequence within any of SEQ ID NOS:1-3. The induced mutation can include a frameshift, where the frameshift includes a deletion selected from the group consisting of a deletion of the guanine at position 253 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 416 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 99 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and a deletion of the guanine at position 322 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. In some cases, the induced mutation can be an in-frame or frameshift mutation, where the in-frame or frameshift mutation includes an insertion or deletion selected from the group consisting of a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. Each induced mutation can have been induced by a rare-cutting endonuclease. In some cases, the first Brassica plant and the second Brassica plant do not contain a transgene. In some embodiments, the first Brassica plant and/or the second Brassica plant contains a transgene encoding a protein, wherein the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid

synthase large subunit protein, a hppd protein, and a DMO protein. In some cases, a mutated FAD2 gene copy can include the sequence set forth in any of SEQ ID NOS:53900-53975, or a sequence at least 90% identical to the sequence set forth in any of SEQ ID NOS:53900-53975.

- 5 In another aspect, this document features a method for producing Brassica oil having increased oleic acid content and reduced linoleic acid content, where the method includes (a) providing a Brassica plant or plant part having an induced mutation in one or more FAD2 gene copies, and (b) producing oil from the plant or plant part. The induced mutation can include a frameshift, where the frameshift includes a deletion selected from
10 the group consisting of a deletion of the guanine at position 253 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 416 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a deletion of the adenine at position 99 of any of SEQ ID
15 NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and a deletion of the guanine at position 322 of any of SEQ ID NOS:1-3 or at the corresponding position within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. In some cases, the induced mutation can be an in-frame or frameshift mutation, where the in-frame or frameshift mutation includes an
20 insertion or deletion selected from the group consisting of a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 3 bp deletion at positions 99-
25 101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding
30 positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions

within a sequence having at least 90% identity to any of SEQ ID NOS:1-3. Each induced mutation can have been induced by a rare-cutting endonuclease. In some embodiments, the first Brassica plant and the second Brassica plant do not contain a transgene. In some embodiments, the first Brassica plant and/or the second Brassica plant contains a

5 transgene encoding a protein, wherein the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid synthase large subunit protein, a hppd protein, and a DMO protein. In some cases, the mutated FAD2 gene copy can include the sequence set forth in any of SEQ ID

10 NOS:53900-53975, or a sequence at least 90% identical to the sequence set forth in any of SEQ ID NOS:53900-53975.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those

15 described herein can be used to practice the invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

20 The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

25 DESCRIPTION OF DRAWINGS

FIG. 1 is an illustration of the *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* gene structures. Position of nucleotides encoding conserved amino acid sequences is shown.

FIG. 2 shows the genotype of T0 plant c02 at the individual *BnaA.FAD2.a*,

30 *BnaC.FAD2.a*, and *BnaC.FAD2.b* TALE nuclease target sites.

FIG. 3 shows the genotype of T0 plant e01 at the individual *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* TALE nuclease target sites.

FIG. 4 shows the genotype of T0 plant e02 at the individual *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* TALE nuclease target sites.

FIG. 5 shows the genotype of T0 plant f02 at the individual *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* TALE nuclease target sites.

FIG. 6 shows sequences from the FAD2 alleles of mutant T0 plants Bn432-a and Bn432-b. Dashes indicate deletions, and underlining indicates insertions.

5 FIG. 7 is a diagram illustrating a scheme for generating F3 seeds to be analyzed for oil composition, showing the FAD2 genotypes of T0 plants Bn432-a and Bn432-b, T1 plants resulting from selfed T0 plants, and F2 plants.

DETAILED DESCRIPTION

10 Studies on canola lines with high oleic acid have been limited, providing incomplete information on the types and combinations of mutations which can result in specific oil compositions. This document provides *B. napus* plants that produce oil with desirable oleic and linolenic levels. More specifically, this document describes novel mutations, alone and in combination, that result in significant improvement in *B. napus*
15 oil composition.

This document also provides novel methods for generating *B. napus* plants that produce oil with desirable oleic and linolenic levels. For example, this document relates to the finding that novel mutations within genes of canola plants can be created using sequence-specific nucleases. These sequence-specific nucleases can be used to inactivate
20 or attenuate genes involved in the fatty acid synthesis pathway.

In some embodiments, the target gene for inactivation or attenuation is a member of the fatty acid synthesis pathway. For example, the target gene can be FAD2, which catalyzes the conversion of oleic acid (18:1) to linoleic acid (18:2). In *B. napus*, four copies of FAD2 genes have been identified (Yang et al., *supra*). As used herein, the four
25 gene copies are named *BnaA.FAD2.a*, *BnaA.FAD2.b-like*, *BnaC.FAD2.a*, and *BnaC.FAD2.b*.

The FAD2 gene copy *BnaA.FAD2.a* is located on chrA05. A representative example of the coding sequence for *BnaA.FAD2.a* is shown in SEQ ID NO:1:

30 ATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAAAGTCT
GAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCACT
GTCGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCG
ATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCT
GCTTCTACTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTC
TCCTACTTCGCCTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAA

CCGGCGTCTGGGTCATAGCCACGAGTGCGGCCACCACGCCTTCAGCG
 ACTACCAGTGGCTGGACGACACCGTCGGCCTCATCTTCCACTCCTTCCT
 CCTCGTCCCTTACTTCTCCTGGAAGTACAGTCATCGACGCCACCATTCC
 AACACTGGCTCCCTCGAGAGAGACGAAGTGTTTGTCCCAAGAAGAA
 5 GTCAGACATCAAGTGGTACGGCAAGTACCTCAACAACCCTTTGGGACG
 CACCGTGATGTTAACGGTTCAGTTCCTCTCGGCTGGCCTTTGTACTTA
 GCCTTCAACGTCTCGGGGAGACCTTACGACGGCGGCTTCGCTTGCCAT
 TTCCACCCCAACGCTCCCATCTACAACGACCGTGAGCGTCTCCAGATA
 TACATCTCCGACGCTGGCATCCTCGCCGCTGCTACGGTCTCTACCGCT
 10 ACGCTGCTGTCCAAGGAGTTGCCTCGATGGTCTGCTTCTACGGAGTTCC
 TCTTCTGATTGTCAACGGGTTCTTAGTTTTGATCACTTACTTGCAGCAC
 ACGCATCCTTCCCTGCCTCACTATGACTCGTCTGAGTGGGATTGGTTGA
 GGGGAGCTTTGGCCACCGTTGACAGAGACTACGGAATCTTGAACAAG
 GTCTTCCACAATATCACGGACACGCACGTGGCGCATCACCTGTTCTCG
 15 ACCATGCCGCATTATCATGCGATGGAAGCTACGAAGGCGATAAAGCC
 GATACTGGGAGAGTATTATCAGTTCGATGGGACGCCGGTGGTTAAGGC
 GATGTGGAGGGAGGCGAAGGAGTGTATCTATGTGGAACCGGACAGGC
 AAGGTGAGAAGAAAGGTGTGTTCTGGTACAACAATAAGTTATGA (SEQ
 ID NO:1)

20 The *BnaA.FAD2.a* gene produces a protein with conserved domains and signaling motifs. A first conserved domain includes the histidine-rich motif HECGHH (SEQ ID NO:35), encoded by nucleotides 313-330 of SEQ ID NO:1. A second conserved domain includes the histidine-rich motif HRRHH (SEQ ID NO:36), encoded by nucleotides 421-435 of SEQ ID NO:1. A third conserved domain includes the histidine-rich motif
 25 HVAHH (SEQ ID NO:37), encoded by nucleotides 946-960 of SEQ ID NO:1. A signaling motif responsible for allowing FAD2 to selectively bind to and embed in the endoplasmic reticulum (YNNKL; SEQ ID NO:38) is encoded by nucleotides 1138-1152 of SEQ ID NO:1.

The FAD2 gene copy *BnaC.FAD2.a* is located on chrC05. A representative
 30 example of the coding sequence for *BnaC.FAD2.a* is shown in SEQ ID NO:3:

ATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGAAGTCT
 GAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCACT
 GTCGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCG
 ATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCT

GCTTCTACTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCCCTCTC
 TCCTACTTTCGCCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCTTAA
 CCGGCGTCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGCG
 ACTACCAGTGGCTTGACGACACCGTTCGGTCTCATCTTCCACTCCTTCCT
 5 CCTCGTCCCTTACTTCTCCTGGAAGTACAGTCATCGACGCCACCATTCC
 AACACTGGCTCCCTCGAGAGAGACGAAGTGTTTGTCCCCAAGAAGAA
 GTCAGACATCAAGTGGTACGGCAAGTACCTCAACAACCCTTTGGGACG
 CACCGTGATGTAAACGGTTCAGTTCACTCTCGGCTGGCCGTTGTACTTA
 GCCTTCAACGTCTCGGGAAGACCTTACGACGGCGGCTTCGCTTGCCAT
 10 TTCCACCCCAACGCTCCCATCTACAACGACCGCGAGCGTCTCCAGATA
 TACATCTCCGACGCTGGCATCCTCGCCGTCTGCTACGGTCTCTTCCGTT
 ACGCCGCCGCGCAGGGAGTGGCCTCGATGGTCTGCTTCTACGGAGTCC
 CGCTTCTGATTGTCAATGGTTTCCTCGTGTTGATCACTTACTTGCAGCA
 CACGCATCCTTCCCTGCCTCACTACGATTTCGTCGAGTGGGATTGGTTG
 15 AGGGGAGCTTTGGCTACCGTTGACAGAGACTACGGAATCTTGAACAAG
 GTCTTCCACAATATTACCGACACGCACGTGGCGCATCATCTGTTCTCCA
 CGATGCCGCATTATCACGCGATGGAAGCTACCAAGGCGATAAAGCCG
 ATACTGGGAGAGTATTATCAGTTCGATGGGACGCCGGTGGTTAAGGCG
 ATGTGGAGGGAGGCGAAGGAGTGTATCTATGTGGAACCGGACAGGCA
 20 AGGTGAGAAGAAAGGTGTGTTCTGGTACAACAATAAGTTATGA (SEQ
 ID NO:3)

The *BnaC.FAD2.a* gene produces a protein with conserved domains and signaling motifs. A first conserved domain includes the histidine-rich motif HECGHH (SEQ ID NO:35), encoded by nucleotides 313-330 of SEQ ID NO:3. A second conserved domain
 25 includes the histidine-rich motif HRRHH (SEQ ID NO:36), encoded by nucleotides 421-435 of SEQ ID NO:3. A third conserved domain includes the histidine-rich motif HVAHH (SEQ ID NO:37), encoded by nucleotides 946-960 of SEQ ID NO:3. A signaling motif responsible for allowing FAD2 to selectively bind to and embed in the endoplasmic reticulum (YNNKL; SEQ ID NO:38) is encoded by nucleotides 1138-1152
 30 of SEQ ID NO:3.

The FAD2 gene copy *BnaA.FAD2.b-like* is located on chrA01 based on synteny of *B. napus* and its two progenitor species, *B. rapa* and *B. oleracea*. A representative example of the coding sequence for *BnaA.FAD2.b-like* is shown in SEQ ID NO:4:

ATGGGCGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCTCCCC
 GGAACCAACACCCTCAAACGCGTCCCCTGCGAGACACCACCATTCACT
 CTCGGAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCC
 ATCCCACGCTCCTTCTCCTCTTCGACATCATCATCTCCTCCTCGGCTCCT
 5 CCCTCTACCACCTCTCCACAGCCTACTTCCCTCTCCCTTACCTCGCCTG
 ACCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGT
 CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACCACCAGTGGCT
 GGACGACGCCGCCGGCCTCGTCTTCCACTCCTTCCTCCTCGTCCCGTAC
 TTCTCCTGGAAGTACATCCATGACGCCACCATTCCAACACCGGATCCC
 10 TCGATAGGGACGAAGTGTTTCGTCCCCAAGAAGAAATCCGACATCAAGT
 GGTACGGCAAGTACCTCAACAACCCGCTAGGACGCACGGTGATGCTA
 ACCGTCCAGTTCAAGCTCGGCTGGCCGTTGTACTTAGCCTTCAACGTCT
 CGGGAAGACCTTACAGCGACGGTTTCGCTTGCCATTTCCACCCGAACG
 CTCCCATCTACAACGACCGCGAGCGTCTCCAGATATACATCTCTGACG
 15 CTGGCGTCCTCTCCGTATGTTACGGTCTCTACCGTTACGCTGCTTCGCG
 AGGAGTAGCCTCTGTGGTCTGTGTCTACGGAGTTCCGCTTCTAATTGTC
 AACTGTTTCCTCGTCTTGATCACTTACTTGCAGCACACGCACCCTTCGC
 TGCCTCACTATGATTCTTCCGAGTGGGATTGGTTGAGAGGAGCTTTGG
 CTACTGTGGATAGAGACTATGGAATCTTGAACAAGGTGTTCCATAACA
 20 TCACGGACACGCACGTGGCGCATCATCTGTTCTCGACGATGCCGCATT
 ATAACGCGATGGAAGCGACCAAGGCGATAAAGCCGATACTTTGGAGA
 GTATTACCAGTTTGATGGAACGCCGGCGGTTAAGGCGATGTGGAGGGA
 GGCGAAGGAGTGTATCTATGTTGAACCGGATAGGCAAGGTGAGAAGA
 AAGGTGTGTTCTGGTACAACAATAAGTTATGA (SEQ ID NO:4)

25 *BnaA.FAD2.b-like* is predicted to be a pseudogene, as there is a premature stop
 codon located at position 409-411 of SEQ ID NO:4.

The FAD2 gene copy *BnaC.FAD2.b* is located on chrC01 based on synteny of *B.*
napus and its two progenitor species, *B. rapa* and *B. oleracea*. A representative example
 of the coding sequence for *BnaC.FAD2.b* is shown in SEQ ID NO:2:

30 ATGGGCGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCTCCCC
 GAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCACT
 CTCGGAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCC
 ATCCCTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTC
 CCTCTACCACCTCTCCACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCC

CTTACCTCGCCTGGCCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAAC
 GGGCCTCTGGGTCATCGCCACGAATGCGGCCACCACGCCTTCAGCGA
 CCACCAGTGGCTGGACGACGCCGTGGGCCTCGTCTTCCACTCCTTCCTC
 CTCGTCCCTTACTTCTCCTGGAAGTACGGCCATCGACGCCACCATTCCA
 5 ACACCGGATCCCTCGAGAGGGATGAAGTGCTCGTCCCCAAGAAGAAA
 TCCGACATCAAGTGGTACGGAAAGTACCTCAACAACCCGCTAGGACGC
 ACGGTGATGCTAACCGTCCAGTTCACGCTCGGCTGGCCGTTGTACTTA
 GCCTTCAACGTCTCTGGAAGACCTTACAGCGACGGTTTCGCTTGCCATT
 TCCACCCGAACGCTCCCATCTACAACGACCGCGAGCGTCTCCAGATAT
 10 ACATCTCTGACGCTGGCGTCCTCTCCGTATGTTACGGTCTCTACCGCTA
 CGCTGGTTCGCGAGGAGTGGCCTCGATGGTCTGTGTCTACGGAGTTCC
 GCTTATGATTGTCAACTGTTTCCTCGTCTTGATCACTTACTTGCAGCAC
 ACGCACCTTCGCTGCCTCACTATGATTCTTCGGAGTGGGATTGGTTGA
 GAGGAGCTTTGGCTACTGTGGATAGAGACTATGGAATCTTGAACAAGG
 15 TGTTTCATAACATCACGGACACGCACGTGGCGCATCATCTGTTCTCGA
 CGATGCCGCATTATAACGCGATGGAAGCGACCAAGGCGATAAAGCCG
 ATACTTGGAGAGTATTACCAGTTTGATGGAACGCCGGTGGTTAAGGCG
 ATGTGGAGGGAGGCGAAGGAGTGTATCTATGTTGAACCGGATAGGCA
 AGGTGAGAAGAAAGGTGTGTTCTGGTACAACAATAAGTTATGA (SEQ
 20 ID NO:2)

The *BnaC.FAD2.b* gene produces a protein with conserved domains and signaling motifs. A first conserved domain includes the histidine-rich motif HECGHH (SEQ ID NO:35), encoded by nucleotides 313-330 of SEQ ID NO:2. A second conserved domain includes the histidine-rich motif HRRHH (SEQ ID NO:36), encoded by nucleotides 421-
 25 435 of SEQ ID NO:2. A third conserved domain includes the histidine-rich motif HVAHH (SEQ ID NO:37), encoded by nucleotides 946-960 of SEQ ID NO:2. A signaling motif responsible for allowing FAD2 to selectively bind to and embed in the endoplasmic reticulum (YNNKL; SEQ ID NO:38) is encoded by nucleotides 1138-1152 of SEQ ID NO:2.

30 Induced mutations within FAD2 gene copies, which can result in inactivation or attenuation of gene function, can include deletions, insertions, and substitutions. Deletions and insertions can range in size from one nucleotide (nt) to 200 or more nt (e.g., one to five nt, five to 10 nt, 10 to 15 nt, 15 to 20 nt, 20 to 30 nt, 30 to 40 nt, 40 to 50 nt, 50 to 75 nt, 75 to 100 nt, 100 to 150 nt, 150 to 200 nt, or more than 200 nt). A deletion

may result in removal of 0.1% to 100% of a gene's coding sequence (e.g., 0.1 to 0.5%, 0.5 to 1%, 1 to 5%, 5 to 10%, 10 to 15%, 15 to 20%, 20 to 30%, 30 to 40%, 40 to 50%, 50 to 60%, 60 to 70%, 70 to 80%, 80 to 90%, or 90 to 100% of the coding sequence).

In some embodiments, deletions that result in inactivation or attenuation of gene function are frameshift mutations (i.e., $-3N+1$ and $-3N+2$, where N is a whole number that is greater than or equal to 1). In some embodiments, insertions that result in inactivation or attenuation are frameshift mutations (i.e., $3N-1$ and $3N-2$, where N is a whole number that is greater than or equal to 1). An inactivating/attenuating frameshift may occur within an exon, after the start codon and before the codon that encodes the last amino acid that is important for protein function. For the *B. napus* FAD2 gene copies, an inactivating/attenuating frameshift can occur before or within the nucleotides corresponding to nucleotides 313-330 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating frameshift can occur before or within the nucleotides corresponding to nucleotides 421-435 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating frameshift can occur before or within the nucleotides corresponding to nucleotides 946-960 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating frameshift can occur before or within the nucleotides corresponding to nucleotides 1138-1152 of SEQ ID NO:1-3.

In some embodiments, deletions that can result in inactivation or attenuation of FAD2 gene function also may include in-frame deletions that remove codons encoding amino acids that are important for protein function. In some cases, for the *B. napus* FAD2 gene copies, an inactivating/attenuating in-frame deletion can be within or include the nucleotides corresponding to nucleotides 313-330 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating in-frame deletion can be within or include the nucleotides corresponding to nucleotides 421-435 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating in-frame deletion can be within or include the nucleotides corresponding to nucleotides 946-960 of SEQ ID NO:1-3. In some cases, an inactivating/attenuating in-frame deletion can be within or include the nucleotides corresponding to nucleotides 1138-1152 of SEQ ID NO:1-3.

To generate FAD2 mutants, *B. napus* cells can be transformed with one or more sequence-specific nucleases targeted to one or more of the FAD2 gene copies. Numerous independent mutants typically can be generated using each of one or more sequence-specific nucleases. Plants with specific FAD2 mutations also can be generated by crossing a first plant containing one or more sequence-specific nuclease-induced FAD2

mutations with a second plant that may or may not have one or more sequence-specific nuclease-induced FAD2 mutations. The resulting FAD2-mutant plants can be evaluated in any suitable way, including (1) phenotyping the desired trait (e.g., oil composition), (2) molecular characterization of the resulting plant(s) to ensure no off-target mutations or
 5 random integration of foreign DNA, (3) segregation of the mutation(s) and phenotype, and (4) agronomic performance of the FAD2-mutant plant(s).

Mutations in the different FAD2 gene copies can be combined. Different combination of mutations can result in different oil characteristics. In some embodiments, Brassica plants with one or more mutations in *BnaA.FAD2.a* can be generated. In some
 10 embodiments, Brassica plants with one or more mutations in *BnaC.FAD2.a* can be generated. In some embodiments, Brassica plants with one or more mutations in *BnaC.FAD2.b* can be generated. In some embodiments, Brassica plants with one or more mutations in *BnaA.FAD2.a* and *BnaC.FAD2.a* can be generated. In some embodiments, Brassica plants with one or more mutations in *BnaA.FAD2.a* and *BnaC.FAD2.b* can be
 15 generated. In some embodiments, Brassica plants with one or more mutations in *BnaC.FAD2.a* and *BnaC.FAD2.b* can be generated. In some embodiments, Brassica plants with one or more mutations in *BnaC.FAD2.a*, *BnaC.FAD2.b*, and *BnaA.FAD2.a* can be generated.

In some embodiments, a plant with one or more mutations in a FAD2 gene copy
 20 also can contain a transgene. The transgene can be integrated into the *Brassica* genome using standard transformation protocols. In some cases, the transgene can result in the expression of a protein that confers tolerance or resistance to one or more herbicides (e.g., glufosinate, mesotrione, imidazolinone, isoxaflutole, glyphosate, 2,4-D, hydroxyphenylpyruvate dioxygenase-inhibiting herbicides, or dicamba). In some cases,
 25 the transgene can encode a plant 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein or a modified plant EPSPS protein, a bacterial EPSPS protein, where the modified plant EPSPS contains an amino acid substitution within the conserved TAMRP (SEQ ID NO:33) sequence. The substitution can include, for example, a threonine to isoleucine substitution, a proline to serine substitution, or a proline to adenine substitution. In some
 30 cases, the transgene can encode a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an aryloxyalkanoate dioxygenase (AAD) protein, a phosphinothricin N-acetyltransferase (PAT) protein, an acetohydroxyacid synthase large subunit protein, a p-hydroxyphenylpyruvate dioxygenase (hppd) protein, or a dicamba monooxygenase (DMO) protein.

In some cases, a nucleic acid can have a nucleotide sequence with at least about 75 percent sequence identity to a representative FAD2 nucleotide sequence. For example, a nucleotide sequence can have at least 75, at least 80, at least 85, at least 90, at least 91, at least 92, at least 93, at least 94, at least 95, at least 96, at least 97, at least 98, or at least 99 percent sequence identity to a representative FAD2 nucleotide sequence as set forth in any of SEQ ID NOS:1-4.

The percent sequence identity between a particular nucleic acid or amino acid sequence and a sequence referenced by a particular sequence identification number is determined as follows. First, a nucleic acid or amino acid sequence is compared to the sequence set forth in a particular sequence identification number using the BLAST 2 Sequences (B12seq) program from the stand-alone version of BLASTZ containing BLASTN version 2.0.14 and BLASTP version 2.0.14. This stand-alone version of BLASTZ can be obtained online at fr.com/blast or at ncbi.nlm.nih.gov. Instructions explaining how to use the B12seq program can be found in the readme file accompanying BLASTZ. B12seq performs a comparison between two sequences using either the BLASTN or BLASTP algorithm. BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. To compare two nucleic acid sequences, the options are set as follows: -i is set to a file containing the first nucleic acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second nucleic acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastn; -o is set to any desired file name (e.g., C:\output.txt); -q is set to -1; -r is set to 2; and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastn -o c:\output.txt -q -1 -r 2. To compare two amino acid sequences, the options of B12seq are set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two amino acid sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

Once aligned, the number of matches is determined by counting the number of positions where an identical nucleotide or amino acid residue is presented in both sequences. The percent sequence identity is determined by dividing the number of matches either by the length of the sequence set forth in the identified sequence (e.g.,
 5 SEQ ID NO:1), or by an articulated length (e.g., 100 consecutive nucleotides or amino acid residues from a sequence set forth in an identified sequence), followed by multiplying the resulting value by 100. For example, an amino acid sequence that has 1100 matches when aligned with the sequence set forth in SEQ ID NO:1 is 95.2 percent identical to the sequence set forth in SEQ ID NO:1 (i.e., $1100 / 1155 \times 100 = 95.2$). It is
 10 noted that the percent sequence identity value is rounded to the nearest tenth. For example, 75.11, 75.12, 75.13, and 75.14 is rounded down to 75.1, while 75.15, 75.16, 7.17, 75.18, and 7.19 is rounded up to 7.2. It also is noted that the length value will always be an integer.

In some embodiments, the methods described herein involve the delivery of
 15 genome engineering reagents to canola (e.g., *B. napus*) plant cells. Any suitable method can be used to introduce the nucleic acid into the plant cell. In some embodiments, for example, a method as provided herein can include contacting a plant cell with an organism that is capable of horizontal gene transfer (e.g., a bacterium, such as an *Agrobacterium*), where the organism contains a Ti or Ri plasmid, or T-DNA plasmid
 20 having a T-DNA region that includes the promoter, UTRs, coding sequence, and a poly-A tail. Methods for *Agrobacterium*-mediated transformation in canola or *B. napus* are described elsewhere (see, e.g., Block et al., *Plant Physiol.* 91:694-701, 1989). In other embodiments, a method for introducing genome editing reagents as provided herein can include biolistic transformation, electroporation-mediated transformation, or polyethylene
 25 glycol-mediated transformation of Brassica plant cells (e.g., protoplasts). The protoplasts can be obtained from hypocotyl or leaf tissue. Plants containing mutations or TALE nuclease DNA can be regenerated using standard plant regeneration protocols. See, for example, Ali et al., *Pakistan Journal of Botany*, 39:1251, 2007; Lu et al., *Zeitschrift für Pflanzenphysiologie*, 107:59-63, 1994; Hu et al., *Plant cell, tissue and organ culture*,
 30 59:189-196, 1999. Nucleic acid can be delivered to Brassica protoplasts, where the nucleic acid is integrated into the host genome. Alternatively, nucleic acid can be delivered to Brassica protoplasts, where the nucleic acid does not integrate into the host genome, but instead remains extrachromosomal. Such transient delivery can result in protein expression from the nucleic acid sequence. If a sequence-specific nuclease is

encoded by the delivered nucleic acid sequence, targeted mutations can be introduced into the Brassica cell without integration of the foreign nucleic acid sequence.

In some embodiments, the methods described herein can include determining the fatty acid composition within *B. napus* seed oil. Fatty acid composition can be determined using, for example, fatty acid methyl esters (FAME) gas chromatography (Beuselinck et al., *Crop Sci.* 47:747-750, 2006). FAME can be used to determine the levels of palmitic, stearic, oleic, linoleic and linolenic acid within *B. napus* seed oil.

In some embodiments, the methods provided herein can include identifying the intended gene edit. Any appropriate means can be employed to identify the desired targeted insertion. One suitable means is polymerase chain reaction (PCR), in which primers are designed to amplify DNA encompassing the TALE nuclease target site. The PCR product can be cloned and sequenced using standard DNA sequencing techniques. The sequencing results can be used to determine if the TALE nuclease target site includes mutations.

As used herein, the amino acid sequences follow the standard single letter or three letter nomenclature. All protein or peptide sequences are shown in conventional format where the N-terminus appears on the left and the carboxyl group at the C-terminus on the right. Amino acid nomenclature, both single letter and three letter, for naturally occurring amino acids are as follows: alanine (Ala; A), asparagine (Asn; N), aspartic acid (Asp; D), arginine (Arg; R), cysteine (Cys; C), glutamic acid (Glu; E), glutamine (Gln; Q), glycine (Gly; G), histidine (His; H), leucine (Leu; L), isoleucine (Ile; I), lysine (Lys; K), methionine (Met; M), phenylalanine (Phe; F), proline (Pro; P), serine (Ser; S), threonine (Thr; T), tryptophan (Trp; W), tyrosine (Tyr; Y), and valine (Val; V).

As used herein, “progeny” includes any plant, seed, plant cell, and/or a regenerated plant part having a non-naturally occurring mutation in the FAD2 gene copies derived from an ancestor plant. Progeny may contain non-naturally occurring mutations in one, two, three, or four of the FAD2 gene copies.

The term “allele(s)” means any of one or more alternative forms of a gene at a particular locus. In a diploid (or amphidiploid) cell of an organism, alleles of a given gene are located at a specific location or locus on a chromosome. One allele is present on each chromosome of the pair of homologous chromosomes. “Heterozygous” alleles are two different alleles residing at a specific locus, positioned individually on corresponding pairs of homologous chromosomes. “Homozygous” alleles are two identical alleles residing at a specific locus, positioned individually on corresponding pairs of homologous

chromosomes in the cell. As used herein, “backcrossing” refers to a repetitive crossing of hybrid plants, where, for example, a first-generation hybrid is crossed back to one of the parents of the hybrid progeny. Backcrossing can be used to transfer one or more loci from one genetic background to a different genetic background.

- 5 As used herein, “crossing” refers to the mating of two parent plants, wherein the pollen of one parent is transferred to the stigma of the second parent. Crossing can be used to transfer one or more specific alleles to a different plant with a different genetic background. Crossing can be used to create a population of *B. napus* plants with a desired trait, where the population includes *B. napus* plants with different genetic backgrounds.
- 10 Here, the *B. napus* plants of different genetic backgrounds are crossed to plants having an allele that produces a desired trait. Crossing, backcrossing and breeding can be carried out via known procedures. DNA fingerprinting, SNP or similar technologies may be used in a marker-assisted selection (MAS) breeding program to transfer or breed mutant FAD2 gene copies into other *B. napus* plants. For example, a breeder can create segregating
- 15 populations from hybridizations of a genotype containing a mutant allele with an agronomically desirable genotype. Plants in the F2 or backcross generations can be screened using markers developed from FAD2 gene copy sequences or fragments thereof. Plants identified as possessing the mutant allele can be backcrossed or self-pollinated to create a second population to be screened. Depending on the expected inheritance pattern
- 20 or the MAS technology used, it may be necessary to self-pollinate the selected plants before each cycle of backcrossing to aid identification of the desired individual plants. Backcrossing or other breeding procedure can be repeated until the desired phenotype of the recurrent parent is recovered. The result of a plant breeding program using the mutant *B. napus* plants described herein can be novel and useful lines and varieties. As used
- 25 herein, the term “variety” refers to a population of plants that share constant characteristics which separate them from other plants of the same species. With respect to *B. napus*, a variety can refer to a population of *B. napus* plants with different genetic backgrounds. A variety is often, although not always, sold commercially. While possessing one or more distinctive traits, a variety can be further characterized by a very
- 30 small overall variation between individuals within that variety. A “pure line” variety may be created by several generations of self-pollination and selection, or vegetative propagation from a single parent using tissue or cell culture techniques. A variety can be essentially derived from another line or variety. As defined by the International Convention for the Protection of New Varieties of Plants (December 2, 1961, as revised

at Geneva on November 10, 1972, on October 23, 1978, and on March 19, 1991), a variety is “essentially derived” from an initial variety if: a) it is predominantly derived from the initial variety, or from a variety that is predominantly derived from the initial variety, while retaining the expression of the essential characteristics that result from the genotype or combination of genotypes of the initial variety; b) it is clearly distinguishable from the initial variety; and c) except for the differences which result from the act of derivation, it conforms to the initial variety in the expression of the essential characteristics that result from the genotype or combination of genotypes of the initial variety. Essentially derived varieties can be obtained, for example, by the selection of a natural or induced mutant, a somaclonal variant, a variant individual from plants of the initial variety, backcrossing, or transformation. A “line” as distinguished from a variety most often denotes a group of plants used non-commercially, for example in plant research. A line typically displays little overall variation between individuals for one or more traits of interest, although there may be some variation between individuals for other traits.

As used herein, the term “explant” refers to a section of plant cells or plant parts that are taken from a donor plant and used for culturing.

The term “increased oleic acid content” and “increased oleic content” refers to any increase in the oleic acid content from *B. napus* seed oil, as compared to the oleic acid content in a corresponding wild type *B. napus* seed oil. In some embodiments, the oleic acid content can be increased by at least 3% (e.g., at least 5%, at least 10%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%), as compared to the oleic acid content in a corresponding wild type *B. napus* seed oil.

The term “reduced linoleic acid content” and “reduced linoleic content” refers to any reduction in the linoleic acid content from *B. napus* seed oil, as compared to the linoleic acid content in a corresponding wild type *B. napus* seed oil. In some embodiments, the linoleic acid content can be reduced by at least 3% (e.g., at least 5%, at least 10%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%), as compared to the linoleic acid content in a corresponding wild type *B. napus* seed oil.

The term “reduced linolenic acid content” and “reduced linolenic content” refers to any reduction in the linolenic acid content from *B. napus* seed oil, as compared to the linolenic acid content in a corresponding wild type *B. napus* seed oil. In some

embodiments, the linolenic acid content can be reduced by at least 3% (e.g., at least 5%, at least 10%, at least 20%, at least 25%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90%), as compared to the linolenic acid content in a corresponding wild type *B. napus* seed oil.

5 The term “induced mutation” as used herein refers to a mutation that was introduced by human intervention. An “induced mutation” can be a mutation that was introduced using one or more sequence-specific nucleases for example. The sequence-specific nuclease can be a meganuclease, TALE nuclease, zinc-finger nuclease, or CRISPR/Cas. An “induced mutation” also can be a mutation that was introduced using a
10 chemical substance (e.g., ethylmethylsulfonate (EMS) or ethylnitrosourea (ENU)). In some embodiments, an “induced mutation” can be a mutation that was introduced using ionizing radiation such as neutrons (fast neutron mutagenesis), gamma rays, or X-rays.

“Wild type” as used herein refers to a typical form of a plant or a gene as it most commonly occurs in nature. For example, a “wild type FAD2 gene copy” is a naturally
15 occurring FAD2 gene copy (e.g., as found within naturally occurring *B. napus* plants) that encodes a functional FAD2 protein, while a “mutant FAD2 gene copy” is a FAD2 gene copy that does not encode a functional FAD2 protein or encodes an attenuated FAD2 protein. Such a “mutant FAD2 gene copy” can include one or more mutations in its nucleic acid sequence, where the mutation(s) result in no detectable amount of the
20 corresponding FAD2 protein activity in the plant, plant part or plant cell in vivo.

As used herein, the term “functional variant” is intended to refer to a catalytically active mutant of a protein or a protein domain. Such a mutant can have the same level of activity, or a higher or lower level of activity as compared to the parent protein or protein domain.

25 As used herein, “nuclear localization sequence” and “NLS” and “NLS tag” refer to an amino acid sequence that facilitates trafficking to the plant cell nucleus. Nuclear localization sequence and NLS and NLS tag can also refer to the nucleotide sequence that codes for an amino acid sequence that facilitates trafficking to the plant cell nucleus. An NLS tag can be located within the a protein sequence, or can be added to the C-terminus
30 or N-terminus of a protein. Localization of an NLS tag near the N-terminus of a protein can be particularly useful.

As used herein, “deoxyribonucleic acid” and “DNA” refer to a biopolymer that includes nucleotides linked together by phosphodiester bridges. The four nucleotides include dAMP (2'-deoxyadenosine-5- monophosphate), dGMP (2'-deoxyguanosine-5-

monophosphate), dCMP (2'-deoxycytosine-5-monophosphate) and dTMP (2'-deoxythymosine-5- monophosphate).

As used herein, the term “codon” refers to nucleotide triplets which code for amino acids. Due to the redundancy of the genetic code, the same amino acid can be coded for by different codons. The following is a list of amino acids and their respective
 5 codons: Met (ATG); Glu (GAA, GAG); Val (GTA, GTC, GTG, GTT); Arg (CGA, CGC, CGG, CGT, AGA, AGG); Leu (CTA, CTC, CTG, CTT, TTA, TTG); Ser (TCA, TCC, TCG, TCT, AGC, AGT); Thr (ACA, ACC, ACG, ACT); Pro (CCA, CCC, CCG, CCT); Ala (GCT, GCA, GCC, GCG); Gly (GGA, GGC, GGG, GGT); Ile (ATA, ATC, ATT);
 10 Lys (AAA, AAG); Asn (AAC, AAT); Gln (CAG, CAA); His (CAC, CAT); Asp (GAC, GAT); Tyr (TAC, TAT); Cys (TGC, TGT); Phe (TTC, TTT); and Trp (UGG).

As used herein, the terms “plant” and “plant part” refer to cells, tissues, organs, seeds, and severed parts (e.g., roots, leaves, and flowers) that retain the distinguishing characteristics of the parent plant. “Seed” refers to any plant structure that is formed by
 15 continued differentiation of the ovule of the plant, following its normal maturation point at flower opening, irrespective of whether it is formed in the presence or absence of fertilization and irrespective of whether or not the seed structure is fertile or infertile.

As referred to herein, “coding sequence” or “CDS” refers to DNA that harbors the necessary information that is required to produce a functional RNA or protein. Coding
 20 sequence or CDS can include a DNA sequence starting with ATG and ending with a stop codon. The coding sequence or CDS usually does not contain introns, if no introns are required to produce the functional RNA or protein.

The term “rare-cutting endonucleases” herein refer to natural or engineered proteins having endonuclease activity directed to nucleic acid sequences having a
 25 recognition sequence (target sequence) about 12-40 bp in length (e.g., 14-40 bp in length). Typical rare-cutting endonucleases cause cleavage inside their recognition site, leaving 4 nt staggered cut with 3'OH or 5'OH overhangs. These rare-cutting endonucleases may be meganucleases, such as wild type or variant proteins of homing endonucleases, more particularly belonging to the dodecapeptide family (LAGLIDADG
 30 (SEQ ID NO:34; see, WO 2004/067736) or may result from fusion proteins that associate a DNA binding domain and a catalytic domain with cleavage activity. TALE nucleases and zinc finger nucleases (ZFN) are examples of fusions of DNA binding domains with the catalytic domain of the endonuclease *FokI*. Customized TALE nucleases are

commercially available under the trade name TALENTM (Collectis, Paris, France). For a review of rare-cutting endonucleases, see Baker, *Nature Methods* 9:23-26, 2012.

As referred to herein, “plant” can refer to crop plants, or monocots and dicots. Examples of a crop plants include soybean, wheat, alfalfa, canola, potato, rice, corn, millet, barley, tomato, apple, pear, strawberry, orange, watermelon, pepper, carrot, sugar beets, yam, lettuce, spinach, sunflower, and rape seed, a flowering plant, such as petunia, rose, and chrysanthemum, conifers and pine trees, a plant used in phytoremediation (e.g., heavy metal-accumulating plants), and a plant used for experimental purposes (e.g., Arabidopsis). The plant can be a monocot or a dicot. Examples of monocots include, without limitation, wheat, maize, rice, orchids, onion, aloe, true lilies, grasses (e.g., setaria), woody shrubs and trees (e.g., palms and bamboo), and food plants such as pineapple and sugar cane. Examples of dicots include, without limitation, tomato, cassava, soybean, tobacco, canola, potato, Arabidopsis, rose, pansy, sunflower, grape, strawberry, squash, bean, pea, and peanut. Orders of dicots include Magnoliales, Illiciales, Laurales, Piperales, Aristochiales, Nymphaeales, Ranunculales, Papaverales, Sarraceniales, Trochodendrales, Hamamelidales, Eucomiales, Leitneriales, Myricales, Fagales, Casuarinales, Caryophyllales, Batales, Polygonales, Plumbaginales, Dilleniales, Theales, Malvales, Urticales, Lecythidales, Violales, Salicales, Capparales, Ericales, Diapensales, Ebenales, Primulales, Rosales, Fabales, Podostemales, Haloragales, Myrtales, Cornales, Proteales, Santales, Rafflesiales, Celastrales, Euphorbiales, Rhamnales, Sapindales, Juglandales, Geraniales, Polygalales, Umbellales, Gentianales, Polemoniales, Lamiales, Plantaginales, Scrophulariales, Campanulales, Rubiales, Dipsacales, and Asterales. Genera of dicots include Atropa, Alseodaphne, Anacardium, Arachis, Beilschmiedia, Brassica, Carthamus, Cocculus, Croton, Cucumis, Citrus, Citrullus, Capsicum, Catharanthus, Cocos, Coffea, Cucurbita, Daucus, Duguetia, Eschscholzia, Ficus, Fragaria, Galucium, Glycine, Gossypium, Helianthus, Hevea, Hyoscyamus, Lactuca, Landolphia, Linum, Litsea, Lycopersicon, Lupinus, Manihot, Majorana, Malus, Medicago, Nicotiana, Olea, Parthenium, Papaver, Persea, Phaseolus, Pistacia, Pisum, Pyrus, Prunus, Raphanus, Ricinus, Senecio, Sinomenium, Stephania, Sinapsis, Solanum, Theobroma, Trifolium, Trigonella, Vicia, Vinca, Vilis, and Vigna. Orders of monocots include Alismatales, Hydrocharitales, Najadales, Triuridales, Commelinales, Eriocaulales, Restionales, Poales, Juncales, Cyperales, Typhales, Bromeliales, Zingiberales, Arecales, Cyclanthales, Pandanales, Arales, Lilliales, and Orchidales. Genera of monocots include Allium, Andropogon, Aragrostis, Asparagus,

Avena, Cynodon, Elaeis, Festuca, Festulolium, Heterocallis, Hordeum, Lemna, Lolium, Musa, Oryza, Panicum, Pannesetum, Phleum, Poa, Secale, Sorghum, Triticum, and Zea. Other plants include Gymnospermae, such as the orders Pinales, Ginkgoales, Cycadales, and Gnetales, such as the genera Abies, Cunninghamia, Picea, Pinus, and Pseudotsuga, such as fir and pine.

“Mutagenesis” as used herein refers to processes in which mutations are introduced into a selected DNA sequence. Mutations induced by endonucleases generally are obtained by a double-strand break, which results in insertion/deletion mutations (“indels”) that can be detected by deep-sequencing analysis. Such mutations typically are deletions of several base pairs, and have the effect of inactivating the mutated allele. In the methods described herein, for example, mutagenesis occurs via double-stranded DNA breaks made by nucleases targeted to selected DNA sequences in a plant cell. Such mutagenesis results in “nuclease-induced mutations” (e.g., nuclease-induced knockouts, such as TALE-nuclease-induced knockouts) and reduced expression of the targeted gene. Following mutagenesis, plants can be regenerated from the treated cells using known techniques (e.g., planting seeds in accordance with conventional growing procedures, followed by self-pollination).

Methods for selecting endogenous target sequences and generating TALE nuclease pairs targeted to such sequences can be performed as described elsewhere. See, for example, PCT Publication No. WO 2011/072246, which is incorporated herein by reference in its entirety. In some embodiments, software that specifically identifies TALE nuclease recognition sites, such as TALE-NT 2.0 (Doyle et al., *Nucl Acids Res* 40:W117-122, 2012) can be used.

Transcription activator-like (TAL) effectors are found in plant pathogenic bacteria of the genus *Xanthomonas*. These proteins play important roles in disease, or trigger defense, by binding host DNA and activating effector-specific host genes (see, e.g., Gu et al., *Nature* 435:1122-1125, 2005; Yang et al., *Proc Natl Acad Sci USA* 103:10503-10508, 2006; Kay et al., *Science* 318:648-651, 2007; Sugio et al., *Proc Natl Acad Sci USA* 104:10720-10725, 2007; and Römer et al., *Science* 318:645-648, 2007). Specificity depends on an effector-variable number of imperfect, typically 34 amino acid repeats (Schornack et al., *J Plant Physiol* 163:256-272, 2006; and WO 2011/072246). Polymorphisms are present primarily at repeat positions 12 and 13, which are referred to as the repeat variable-diresidue (RVD).

The RVDs of TALEs correspond to the nucleotides in their target sites in a direct, linear fashion, one RVD to one nucleotide, with some degeneracy and no apparent context dependence. This mechanism for protein-DNA recognition enables target site prediction for new target specific TALEs, as well as target site selection and engineering of new TALEs with binding specificity for the selected sites.

TALE DNA binding domains can be fused to other sequences, such as endonuclease sequences, resulting in chimeric endonucleases targeted to specific, selected DNA sequences, and leading to subsequent cutting of the DNA at or near the targeted sequences. Such cuts (i.e., double-stranded breaks) in DNA can induce mutations into the wild type DNA sequence via non-homologous end joining (NHEJ) or homologous recombination, for example. In some cases, TALE nucleases can be used to facilitate site directed mutagenesis in complex genomes, knocking out or otherwise altering gene function with great precision and high efficiency. As described in the Examples below, TALE nucleases targeted to *B. napus* FAD2 gene copies can be used to mutagenize the endogenous alleles, resulting in plants without detectable activity of the corresponding FAD2 protein, or attenuated activity. Some endonucleases (e.g., *FokI*) function as dimers and can be used to enhance the target specificity of the TALE nuclease. For example, in some cases a pair of TALE nuclease monomers targeted to different DNA sequences can be used. When the two TALE nucleases recognition sites are in close proximity, the inactive monomers can come together to create a functional enzyme that cleaves the DNA. By requiring DNA binding to activate the nuclease, a highly site-specific restriction enzyme can be created.

Methods for using TALE nucleases to generate Brassica plants, plant cells, or plant parts having mutations in endogenous genes include, for example, those described in the Examples herein. For example, one or more nucleic acids encoding TALE nucleases targeted to conserved nucleotide sequences present in one or more FAD2 gene copies can be transformed into plant cells or plant parts, where they can be expressed. In some cases, one or more TALE nuclease proteins can be introduced into plant cells or plant parts. The cells or plant parts, or a plant cell line or plant part generated from the cells, can subsequently be analyzed to determine whether mutations have been introduced at the target site(s), through next-generation sequencing techniques (e.g., 454 pyrosequencing or illumina sequencing) or conventional sequencing methods (e.g., Sanger sequencing). The template for sequencing can be, for example, the TALE nuclease target site within a FAD2 gene copy.

RNA-guided systems also can be used in the methods provided herein. For example, the clustered regularly interspaced short palindromic repeats/CRISPR-associated (CRISPR/Cas) systems use RNA to direct DNA cleavage (see, e.g., Belahj et al., *Plant Methods* 9:39, 2013). This system consists of a Cas9 endonuclease and a guide RNA (either a complex between a CRISPR RNA [crRNA] and trans-activating crRNA [tracrRNA], or a synthetic fusion between the 3' end of the crRNA and 5' end of the tracrRNA). The guide RNA directs Cas9 binding and DNA cleavage to sequences that are adjacent to a proto-spacer adjacent motif (PAM; e.g., NGG for Cas9 from *Streptococcus pyogenes*). Once at the target DNA sequence, Cas9 generates a DNA double-strand break at a position three nucleotides from the 3' end of the crRNA sequence that is complementary to the target sequence. As there are several PAM motifs present in the nucleotide sequence of the FAD2 gene copies, the CRISPR/Cas system may be employed to introduce mutations within the FAD2 gene copies within *B. napus* plant cells in which the Cas9 endonuclease and the guide RNA are transfected and expressed. This approach can be used as an alternative to TALE nucleases in some instances, to obtain plants and plant parts as described herein.

The term "expression" as used herein refers to the transcription of a particular nucleic acid sequence to produce sense or antisense RNA or mRNA, and/or the translation of an mRNA molecule to produce a polypeptide, with or without subsequent post-translational events.

The term "modulating" as used herein refers to increasing or decreasing translational efficiency of an mRNA. This can be accomplished by inserting, removing, or altering a 5' UTR sequence, a 3' UTR sequence, or 5' and 3' UTR sequences.

As used herein, the term "nucleic acid" refers to a polymer made up of nucleotide monomers. A nucleic acid can be single stranded or double stranded, and can be linear or circular. Where single-stranded, a nucleic acid can be a sense strand or an antisense strand. A nucleic acid can be composed of DNA (e.g., cDNA, genomic DNA, synthetic DNA, or a combination thereof), RNA, or DNA and RNA. Further, nucleic acids can contain information for gene expression, including, but not limited to, promoters, 5' UTRs, 3' UTRs, coding sequences, and terminators.

A "vector" is a replicon, such as a plasmid, phage, or cosmid, into which another DNA segment may be inserted so as to bring about the replication of the inserted segment. Generally, a vector is capable of replication when associated with the proper control elements. Suitable vector backbones include, for example, those routinely used in

the art such as plasmids, viruses, artificial chromosomes, BACs, YACs, or PACs. The term “vector” includes cloning and expression vectors, as well as viral vectors and integrating vectors. An “expression vector” is a vector that includes one or more expression control sequences, and an “expression control sequence” is a DNA sequence that controls and regulates the transcription and/or translation of another DNA sequence. Suitable expression vectors include, without limitation, plasmids and viral vectors derived from, for example, bacteriophage, baculoviruses, tobacco mosaic virus, herpes viruses, cytomegalovirus, retroviruses, vaccinia viruses, adenoviruses, and adeno-associated viruses. Numerous vectors and expression systems are commercially available from such corporations as Novagen (Madison, WI), Clontech (Palo Alto, CA), Stratagene (La Jolla, CA), and Invitrogen/Life Technologies (Carlsbad, CA).

The terms “regulatory region,” “control element,” and “expression control sequence” refer to nucleotide sequences that influence transcription or translation initiation and rate, and stability and/or mobility of the transcript or polypeptide product. Regulatory regions include, without limitation, promoter sequences, enhancer sequences, response elements, protein recognition sites, inducible elements, promoter control elements, protein binding sequences, 5' and 3' UTRs, transcriptional start sites, termination sequences, polyadenylation sequences, introns, and other regulatory regions that can reside within coding sequences, such as secretory signals, Nuclear Localization Sequences (NLS) and protease cleavage sites.

As used herein, “operably linked” means incorporated into a genetic construct so that expression control sequences effectively control expression of a coding sequence of interest. A coding sequence is “operably linked” and “under the control” of expression control sequences in a cell when RNA polymerase is able to transcribe the coding sequence into RNA, which if an mRNA, then can be translated into the protein encoded by the coding sequence. Thus, a regulatory region can modulate, e.g., regulate, facilitate or drive, transcription in the plant cell, plant, or plant tissue in which it is desired to express a modified target nucleic acid.

In addition, in some embodiments in which a plant part or plant cell is used, the methods provided herein can include regenerating a plant from the plant part or plant cell. The methods also can include breeding the plant (e.g., the plant into which the nucleic acids were introduced, or the plant obtained after regeneration of the plant part or plant cell used as a starting material) to obtain a genetically desired plant lineage. Methods for regenerating and breeding plants are well established in the art.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

EXAMPLES

5 Example 1 - Design of TALE nucleases targeting FAD2 genes in *B. napus*

To identify potential target sequences for TALE nucleases, the FAD2 genes were sequenced. To this end, publicly available coding sequences for the FAD2 loci in canola were downloaded from NCBI. See, SEQ ID NOS:1-4 for the coding sequences of *BnaA.FAD2.a*, *BnaC.FAD2.b*, *BnaC.FAD2.a* and *BnaA.FAD2.b*-like. Primers were
 10 designed to amplify the coding sequences of *BnaA.FAD2.a*, *BnaC.FAD2.b*, and *BnaC.FAD2.a* in two *B. napus* spring cultivars, Westar and Topas. Primers are shown in TABLE 1.

TABLE 1
Primers for sequencing the *Brassica napus* FAD2 genes

Primer name	Description	Sequence	SEQ ID NO:
BnFAC1Fa	FAD2.a forward	CAGGATCCATGGGtGCAGGTGGAAGAAT	5
BnFAC1Fb	FAD2.b forward	CAGGATCCATGGGcGCAGGTGGAAGAAT	6
BnFAC1R	FAD2 reverse	CAGAGCTCTCATAACTTATTGTTGTACCAG	7

15

DNA was extracted from *B. napus* spring cultivars, Westar and Topas, using standard CTAB-based methods (Murray and Thompson, *Nucl Acids Res*, 8:4321- 4325, 1980). Primers shown in TABLE 1 were used to amplify the FAD2 gene copies. Resulting PCR amplicons were cloned and sequenced. Sequences were aligned and a
 20 consensus sequence for the FAD2 gene copies was elucidated.

Seven TALE nucleases were designed to target sequence near the 5' end of the FAD2 coding sequence. TALE nucleases were designed to target sequence upstream of one or more sites which encode conserved amino acids (e.g., HECGHH [SEQ ID NO:35], HRRHH [SEQ ID NO:36], HVAHH [SEQ ID NO:37], YNNKL [SEQ ID NO:38]). Due
 25 to differences in nucleotide sequences between the copies of the FAD2 genes, TALE nucleases could not be designed to target a conserved sequence between the FAD2 gene copies. Instead, TALE nucleases were designed to target either one or two of the copies of the FAD2 gene copies.

The seven TALE nuclease pairs were named BnFAD2_T01, BnFAD2_T02, BnFAD2_T03, BnFAD2_T04, BnFAD2_T05, BnFAD2_T06, and BnFAD2_T07. A listing of the TALE nucleases and predicted binding sites are shown in TABLE 2.

5

TABLE 2
Individual TALE nuclease target sequences

TALE nuclease target	Target sequence	SEQ ID NO:
BnFAD2_T01	TCCCTCACCCTCTCTCCTACTTCGCCTGGCCTCTCTACTGG GCCTGCCA	8
BnFAD2_T02	TCGTCCCTTACTTCTCCTGGAAGTACAGTCATCGACGCCAC CATTCCAA	9
BnFAD2_T03	TGCGAGACACCGCCCTTCACTGTCTGGAGAACTCAAGAAAG CAATCCCA	10
BnFAD2_T04	TACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA CGTCGCCA	11
BnFAD2_T05	TCTGGGTCATAGCCCACGAGTGC GGCCACCACGCCTTCAG CGACTACCA	12
BnFAD2_T06	TCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCC CTCTACCA	13
BnFAD2_T07	TCTGGGTCATCGCCCACGAATGCGGCCACCACGCCTTCAG CGACCACCA	14

Differences in nucleotide sequences between the FAD2 gene copies prevented the design of TALE nucleases which target a conserved sequence across all FAD2 gene copies. A list of the predicted TALE nuclease binding sequences within each FAD2 gene copy for each TALE nuclease pair is shown in TABLES 3-9.

TABLE 3
Predicted binding sequences for BnFAD2_T01 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TCCCTCACCCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGG CCTGCCA	8
<i>BnaC.FAD2.a</i>	TCCCTCACCCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGG CCTGCCA	8
<i>BnaC.FAD2.b</i>	TCCCCCACCCTCTCCCTTACCTCGCCTGGCCCCCTCTACTGGG CCTGCCA	15
<i>BnaA.FAD2.b</i> -like	CTTACCTCGCCTGACCCCTCTACTGGGCCTGCCA	16

TABLE 4

Predicted binding sequences for BnFAD2_T02 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TCGTCCCTTACTTCTCCTGGAAGTACAGTCATCGACGCCAC CATTCCAA	9
<i>BnaC.FAD2.a</i>	TCGTCCCTTACTTCTCCTGGAAGTACAGTCATCGACGCCAC CATTCCAA	9
<i>BnaC.FAD2.b</i>	TCGTCCCTTACTTCTCCTGGAAGTACGGCCATCGACGCCAC CATTCCAA	17
<i>BnaA.FAD2.b</i> -like	TCGTCCCGTACTTCTCCTGGAAGTACATCCATGACGCCACC ATTCCAA	18

5

TABLE 5

Predicted binding sequences for BnFAD2_T03 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TGCGAGACACCGCCCTTCACTGTCGGAGAACTCAAGAAAG CAATCCCA	10
<i>BnaC.FAD2.a</i>	TGCGAGACACCGCCCTTCACTGTCGGAGAACTCAAGAAAG CAATCCCA	10
<i>BnaC.FAD2.b</i>	TGCGAGACACCACTTCACTCTCGGAGACCTCAAGAAAG CAATCCCA	19
<i>BnaA.FAD2.b</i> -like	TGCGAGACACCACTTCACTCTCGGAGACCTCAAGAAAG CAATCCCA	20

TABLE 6

Predicted binding sequences for BnFAD2_T04 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA CGTCGCCA	11
<i>BnaC.FAD2.a</i>	TACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA CGTCGCCA	11
<i>BnaC.FAD2.b</i>	TACCTCCTCTCGACATCCTCGTCTCCTCCTCCCTCTACCAC CTCTCCA	21
<i>BnaA.FAD2.b</i> -like	TCTTCGACATCATCTCCTCCTCGGCTCCTCCCTCTACCA CCTCTCCA	22

10

TABLE 7

Predicted binding sequences for BnFAD2_T05 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACTACCA	12
<i>BnaC.FAD2.a</i>	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACTACCA	12
<i>BnaC.FAD2.b</i>	TCTGGGTCATCGCCCACGAATGCGGCCACCACGCCTTCAGC GACCACCA	14
<i>BnaA.FAD2.b</i> -like	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACCACCA	23

TABLE 8

5 Predicted binding sequences for BnFAD2_T06 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGC TTCTACTA	24
<i>BnaC.FAD2.a</i>	TCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGC TTCTACTA	25
<i>BnaC.FAD2.b</i>	TCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCC CTCTACCA	13
<i>BnaA.FAD2.b</i> -like	TCCTTCTCCTCTTCGACATCATCATCTCCTCCTCGGCTCCTC CCTCTACCA	26

TABLE 9

Predicted binding sequences for BnFAD2_T07 within the four FAD2 gene copies

FAD2 target	Predicted binding sequence for BnFAD2_T01	SEQ ID NO:
<i>BnaA.FAD2.a</i>	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACTACCA	27
<i>BnaC.FAD2.a</i>	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACTACCA	28
<i>BnaC.FAD2.b</i>	TCTGGGTCATCGCCCACGAATGCGGCCACCACGCCTTCAGC GACCACCA	14
<i>BnaA.FAD2.b</i> -like	TCTGGGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGC GACCACCA	29

Example 2 – Assessing the activity of TALE nucleases targeting FAD2 gene copies

To assess the activity of the FAD2-targeting TALE nuclease pairs, a protoplast transformation experiment was performed. Brassica protoplasts were isolated from leaves of plants grown *in vitro*. Plants (*B. napus* cv. Westar) were grown *in vitro* for 3-4 weeks before leaves were collected. Leaves were digested in an enzyme solution containing 0.25% cellulase and 0.05% macerozyme. The enzyme solution was placed in a 25°C incubator and shaken at 25 rpm for approximately 16 hours. Following digestion, protoplasts were isolated using conventional protoplast isolation techniques with some modification. Specifically, protoplasts were filtered through a 100 um cell strainer into a 50 mL conical tube. Cells were pelleted by centrifugation at 100 g for 5 minutes and washed by W5 solution for three times. After washing, cell pellet was resuspended in an MMG solution, then protoplasts (100,000 per sample) were transformed in a solution containing 20% polyethylene glycol and 30 ug of TALE nuclease-encoded plasmid DNA (i.e., 15 ug of plasmid encoding each TALE nuclease monomer). Following transformation, cells were washed one time in W5. After the wash, cells were resuspended in K8P solution and incubated at 25°C. Two days post transformation, protoplasts were collected and DNA was extracted. TALE nuclease target sites were amplified by PCR, and the resulting amplicons were deep sequenced using illumina sequencing. Primers used to amplify the TALE nuclease target sequences are shown in TABLE 10.

TABLE 10
Oligo names and binding sequences for illumina sequencing

Oligo name	Binding sequence	SEQ ID NO:
BnFAC1Fa-HTS	CAGGATCCATGGGTGCAGGTGGAAGAAT	5
BnFA353R-HTS	GGTAGTCGCTGAAGGCGTGGT	30
BnFA231F-HTS	CCTCTCCTCCCTCACCCCTCTC	31
BnFA591R-HTS	GCTAAGTACAAAGGCCAGCCGAG	32

Mutation frequencies of TALE nuclease pairs were assessed for activity at the FAD2 gene copies. Mutation frequencies were calculated by dividing the number of mutant sequences by the total number of sequences. Reads were grouped by the corresponding FAD2 gene copy before mutation frequencies were calculated. The mutation frequencies for all TALE nuclease pairs are shown in TABLE 11.

TABLE 11
TALE nuclease mutation frequencies at FAD2

TALE nuclease pair	FAD2 target	# of events	Total reads	Mutagenesis frequency (%)
BnFAD2_T01	<i>BnaA.FAD2.a</i>	8065	96878	8.3
BnFAD2_T01	<i>BnaC.FAD2.b</i>	3062	159046	1.9
BnFAD2_T01	<i>BnaC.FAD2.a</i>	7936	78825	10
BnFAD2_T02	<i>BnaA.FAD2.a</i>	10003	102684	9.7
BnFAD2_T02	<i>BnaC.FAD2.b</i>	5030	91948	5.5
BnFAD2_T02	<i>BnaC.FAD2.a</i>	5870	57266	10.3
BnFAD2_T03	<i>BnaA.FAD2.a</i>	9229	102038	9
BnFAD2_T03	<i>BnaC.FAD2.b</i>	15617	156185	10
BnFAD2_T03	<i>BnaC.FAD2.a</i>	6614	67230	9.8
BnFAD2_T04	<i>BnaA.FAD2.a</i>	ND	ND	ND
BnFAD2_T04	<i>BnaC.FAD2.b</i>	ND	ND	ND
BnFAD2_T04	<i>BnaC.FAD2.a</i>	ND	ND	ND
BnFAD2_T05	<i>BnaA.FAD2.a</i>	3010	78379	3.8
BnFAD2_T05	<i>BnaC.FAD2.b</i>	1549	67387	2.3
BnFAD2_T05	<i>BnaC.FAD2.a</i>	2114	48613	4.4
BnFAD2_T06	<i>BnaA.FAD2.a</i>	ND	ND	ND
BnFAD2_T06	<i>BnaC.FAD2.b</i>	ND	ND	ND
BnFAD2_T06	<i>BnaC.FAD2.a</i>	ND	ND	ND
BnFAD2_T07	<i>BnaA.FAD2.a</i>	1904	85910	2.2
BnFAD2_T07	<i>BnaC.FAD2.b</i>	1492	57404	2.6
BnFAD2_T07	<i>BnaC.FAD2.a</i>	1154	41932	2.8

FAD2 mutations within Brassica cells were further analyzed. Both insertions and deletions were observed, with the majority being deletions. With respect to TALE nuclease pair BnFAD2_T01, the majority of mutations resulted in deletion of a specific nucleotide within the TALE nuclease target sequence. For TALE nuclease pair BnFAD2_T01, a deletion of the guanine at position 253 of SEQ ID NOS:1, 2 and 3 was observed in the majority of mutations.

With respect to TALE nuclease pair BnFAD2_T02, the majority of mutations resulted in deletion of a specific nucleotide within the TALE nuclease target sequence. For TALE nuclease pair BnFAD2_T02, a deletion of the adenine at position 416 of SEQ ID NOS:1, 2 and 3 was observed in the majority of mutations.

With respect to TALE nuclease pair BnFAD2_T03, the majority of mutations resulted in deletion of a specific nucleotide within the TALE nuclease target sequence.

For TALE nuclease pair BnFAD2_T03, a deletion of the adenine at position 99 of SEQ ID NOS:1, 2 and 3 was observed in the majority of mutations.

With respect to TALE nuclease pair BnFAD2_T05, the majority of mutations resulted in deletion of a specific nucleotide within the TALE nuclease target sequence.

- 5 For TALE nuclease pair BnFAD2_T05, a deletion of the guanine at position 322 of SEQ ID NOS:1, 2 and 3 was observed in the majority of mutations.

With respect to TALE nuclease pair BnFAD2_T07, the majority of mutations resulted in deletion of a specific nucleotide within the TALE nuclease target sequence.

- 10 For TALE nuclease pair BnFAD2_T07, a deletion of the guanine at position 322 of SEQ ID NOS:1, 2 and 3 was observed in the majority of mutations.

A list of FAD2 sequences containing TALE nuclease-induced mutations, both insertions and deletions, identified within Brassica cells is provided within SEQ ID NOS:39-53891.

- 15 Examples of induced mutations within *BnaA.FAD2.a* using TALE nuclease pair BnFAD2_T01 are shown within SEQ ID NOS:2533-7234.

Examples of induced mutations within *BnaC.FAD2.a* using TALE nuclease pair BnFAD2_T01 are shown within SEQ ID NOS:7235-11755.

Examples of induced mutations within *BnaC.FAD2.b* using TALE nuclease pair BnFAD2_T01 are shown within SEQ ID NOS:39-2532.

- 20 Examples of induced mutations within *BnaA.FAD2.a* using TALE nuclease pair BnFAD2_T02 are shown within SEQ ID NOS:11756-18467.

Examples of induced mutations within *BnaC.FAD2.a* using TALE nuclease pair BnFAD2_T02 are shown within SEQ ID NOS:18468-22172.

- 25 Examples of induced mutations within *BnaC.FAD2.b* using TALE nuclease pair BnFAD2_T02 are shown within SEQ ID NOS:22173-25734.

Examples of induced mutations within *BnaA.FAD2.a* using TALE nuclease pair BnFAD2_T03 are shown within SEQ ID NOS:25735-31174.

Examples of induced mutations within *BnaC.FAD2.a* using TALE nuclease pair BnFAD2_T03 are shown within SEQ ID NOS:31175-35193.

- 30 Examples of induced mutations within *BnaC.FAD2.b* using TALE nuclease pair BnFAD2_T03 are shown within SEQ ID NOS:35194-46067.

Examples of induced mutations within *BnaA.FAD2.a* using TALE nuclease pair BnFAD2_T05 are shown within SEQ ID NOS:46068-48123.

Examples of induced mutations within *BnaC.FAD2.a* using TALE nuclease pair BnFAD2_T05 are shown within SEQ ID NOS:48124-49475.

Examples of induced mutations within *BnaC.FAD2.b* using TALE nuclease pair BnFAD2_T05 are shown within SEQ ID NOS:49476-50588.

5 Examples of induced mutations within *BnaA.FAD2.a* using TALE nuclease pair BnFAD2_T07 are shown within SEQ ID NOS:50589-52024.

Examples of induced mutations within *BnaC.FAD2.a* using TALE nuclease pair BnFAD2_T07 are shown within SEQ ID NOS:52025-52787.

10 Examples of induced mutations within *BnaC.FAD2.b* using TALE nuclease pair BnFAD2_T07 are shown within SEQ ID NOS:52788-53891.

Example 3 – Regenerating *B. napus* plants with mutations in FAD2 gene copies

To regenerate *Brassica* plants having mutations within the FAD2 gene, protoplasts were transformed with DNA encoding functional TALE nuclease
 15 BnFAD2_T03 and regenerated into whole plants. *Brassica* protoplasts were isolated from leaves of plants grown *in vitro*. Plants (*B. napus* cv. Westar) were grown *in vitro* for 3-4 weeks before leaves were collected. Leaves were digested in an enzyme solution containing 0.25% cellulase and 0.05% macerozyme. The enzyme solution was placed in a 25°C incubator and shaken at 25 rpm for about 16 hours. Following digestion, protoplasts
 20 were isolated using conventional protoplast isolation techniques with some modification. Specifically, protoplasts were filtered through a 100 µm cell strainer into a 50 mL conical tube. Cells were pelleted by centrifugation at 100 g for 5 minutes and washed by W5 solution for three times. After washing, cell pellet was resuspended in an MMG solution, then protoplasts (100,000 per sample) were transformed in a solution containing 20%
 25 polyethylene glycol and 30 µg of TALE nuclease-encoded plasmid DNA (15 µg of plasmid encoding each TALE nuclease monomer). Following transformation, cells were washed one time in W5. After the wash, cells were resuspended in K8P solution and incubated at 25°C.

Plants were regenerated from protoplasts using conventional *B. napus*
 30 regeneration techniques; see, for example, Ali et al., *Pakistan Journal of Botany*, 39:1251, 2007; Lu et al., *Zeitschrift für Pflanzenphysiologie*, 107:59-63, 1994; Hu et al., *Plant Cell, Tissue and Organ Culture*, 59:189-196, 1999.

Individual *B. napus* plants regenerated from protoplasts were then advanced to molecular screening.

Example 4 – Screening Brassica plants for mutations in FAD2 gene copies

DNA was extracted from Brassica plants using standard CTAB-based methods (Murray and Thompson, *Nucl Acids Res*, 8:4321- 4325, 1980). Target sites for

5 BnFAD2_T03 TALE nuclease pair were amplified by PCR, and the resulting amplicons were deep sequenced using Illumina sequencing. Primers used to amplify the BnFAD2_T03 TALE nuclease target sequences are shown in TABLE 10.

Plant e02 had a 4 bp deletion in *BnaC.FAD2.a* (FIG. 2). The 4 bp deletion corresponded to nucleotides 252-255 of SEQ ID NO:3; the region with the 4 bp deletion
10 is shown in FIG. 2 (SEQ ID NO:53894).

Plant e01 had a 4 bp deletion in *BnaC.FAD2.a* and a 5 bp deletion in *BnaC.FAD2.a* (FIG. 3). The 4 bp deletion corresponded to nucleotides 252-255 of SEQ ID NO:3, and the 5 bp deletion corresponded to nucleotides 251-255 of SEQ ID NO:3. The region with the 5 bp and 4 bp deletions is shown in FIG. 3 (SEQ ID NOS:53897 and
15 53894, respectively).

Plant e02 had a 5 bp deletion in *BnaC.FAD2.a* (FIG. 4). The 5 bp deletion corresponded to nucleotides 251-255 of SEQ ID NO:3; the region with the 5 bp deletion is shown in FIG. 4 (SEQ ID NO:53897).

Plant f02 had a 5 bp deletion in *BnaC.FAD2.a* (FIG. 5). The 5 bp deletion
20 corresponded to nucleotides 251-255 of SEQ ID NO:3; the region with the 5 bp deletion is shown in FIG. 5 (SEQ ID NO:53897).

In further studies, thirty-two (32) fertile *B. napus* T0 plants transformed with TALE nuclease pair BnFAD2_T03 were identified as containing mutations within FAD2 gene copies (TABLE 12). In total, 66 novel alleles containing deletions or insertions in
25 the FAD2 genes *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* were generated in the T0 population, including 26 alleles in the *BnaA.FAD2.a* gene with indels ranging from -25 bp to +96 bp, 27 alleles in the *BnaC.FAD2.a* gene with indels ranging from -69 bp to +101 bp, and 13 alleles in the *BnaC.FAD2.b* gene with indels ranging from -16 bp to +44 bp. Sequences for the mutant *BnaA.FAD2.a* alleles are provided in TABLE 13A,
30 sequences for the mutant *BnaC.FAD2.a* alleles are provided in TABLE 13B, and sequences for the mutant *BnaC.FAD2.b* alleles are provided in TABLE 13C.

T0 plant Bn432-a had mutations in the genes *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* genes:

- A 3 bp deletion (SEQ ID NO:53900) and a 6 bp deletion (SEQ ID NO:53901) in the *BnaA.FAD2.a* gene. The 3 bp deletion corresponded to nucleotides 98-100 of SEQ ID NO:1, and the 6 bp deletion corresponded to nucleotides 97-102 of SEQ ID NO:1.
- 5 - A 3 bp deletion and a 1 bp insertion in one allele of the *BnaC.FAD2.a* gene (SEQ ID NO:53902) and a 7 bp deletion (SEQ ID NO:53903) in the other allele of the *BnaC.FAD2.a* gene. The 3 bp deletion corresponded to nucleotides 99-101 of SEQ ID NO:3, and the 1 bp insertion was at nucleotide 99. The 7 bp deletion corresponded to nucleotides 98-104 of SEQ ID NO:3.
- 10 - A 13 bp deletion (SEQ ID NO:53904) in one allele of the *BnaC.Fad2.b* gene. The 13 bp deletion corresponded to nucleotides 92-104 of SEQ ID NO:2.

Sequences containing the deletions are shown in FIG. 6.

T0 plant Bn432-b had mutations in the *BnaA.FAD2.a* and *BnaC.FAD2.a* genes:

- A 6 bp deletion and a 50 bp insertion (SEQ ID NO:53905) in both alleles of the *BnaA.FAD2.a* gene. The 6 bp deletion corresponded to nucleotides 96-101 of SEQ ID NO:1, with the 50 bp fragment inserted at position 96.
- 15 - A 6 bp deletion and a 50 bp insertion (SEQ ID NO:53905) in one allele of the *BnaC.FAD2.a* gene and an 18 bp deletion and a 166 bp insertion (SEQ ID NO:53906) in the other allele of the *BnaC.FAD2.a* gene. The 6 bp deletion corresponded to nucleotides 96-101 of SEQ ID NO:3, with the 50 bp fragment inserted at position 96. The 18 bp deletion corresponded to nucleotides 101-118 of SEQ ID NO:3 with the 166 bp fragment inserted at position 101.
- 20 - A 6 bp deletion and a 50 bp insertion (SEQ ID NO:53905) in one allele of the *BnaC.FAD2.a* gene and an 18 bp deletion and a 166 bp insertion (SEQ ID NO:53906) in the other allele of the *BnaC.FAD2.a* gene. The 6 bp deletion corresponded to nucleotides 96-101 of SEQ ID NO:3, with the 50 bp fragment inserted at position 96. The 18 bp deletion corresponded to nucleotides 101-118 of SEQ ID NO:3 with the 166 bp fragment inserted at position 101.

Sequences showing the deletions and insertions are shown in FIG. 6.

As illustrated in FIG. 7, T1 Brassica plants from T0 plant Bn432-a (Bn432-01, Bn432-02, and Bn432-03) and T1 Brassica plants from T0 plant Bn432-b (Bn432-05 and Bn432-07) containing mutations in FAD2 gene copies (TABLE 14) were crossed in order to combine knockout mutations and inactivate one, two, or all three FAD2 genes (*BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b*). F2 mutants were generated and advanced to phenotypic analysis. Mutant F2 plants were classified in different groups for analysis, based on the mutation profile of the *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b* genes:

- Group 1 (3-gene knockout: *BnaA.FAD2.a*, *BnaC.FAD2.a*, and *BnaC.FAD2.b*);
- 21 lines
- Group 2a (2-gene knockout: *BnaA.FAD2.a* and *BnaC.FAD2.a*); 7 lines

- Group 2b (2-gene knockout: *BnaC.FAD2.a* and *BnaC.FAD2.b*); 4 lines
- Group 3 (1-gene knockout: *BnaC.FAD2.a*); 5 lines
- Group 4 (cv Westar wild type control); 2 lines

5 Example 5 – Phenotyping FAD2-mutant *B. napus* plants for oil composition

F3 seeds derived from 37 F2 *B. napus* lines with mutations in one, two, or all three FAD2 genes, or seeds from wild type control cv Westar, were isolated and analyzed for fatty acid composition. Fatty acid composition was determined using FAME gas chromatography (Beuselinek et al., *Crop Sci.* 47:747-750, 2006). Results are presented in
10 TABLES 15 and 16.

Samples with 3-gene knockout mutations (Group 1) demonstrated increased oleic acid (18:1), reduced linoleic acid (18:2), and reduced linolenic acid (18:3) content as compared to wild type Brassica cv Westar. In Group 1 mutants on average, oleic acid content was increased by 24%, from 67.3% to 83.5%, linoleic acid content was decreased
15 by 85%, from 16.5% to 2.4%, and linolenic acid content was decreased by 47%, from 5.9% to 3.1%, as compared to wild type cv Westar. Samples in Group 1 with different knockout mutation profiles in the *BnaC.FAD2.a* gene (-18,+166/-18,+166; -3,+1/-3,+1; -7/-7; -18,+166/-3,+1; and -18,+166/-7) exhibited comparable oleic, linoleic, and linolenic acid composition profiles.

20 Samples with 2-gene knockout mutations in the *BnaA.FAD2.a* and *BnaC.FAD2.a* genes (Group 2a) had increased oleic acid (18:1), reduced linoleic acid (18:2), and reduced linolenic acid (18:3) content as compared to wild type Brassica cv Westar. In Group 2a mutants on average, oleic acid content was increased by 21.9%, from 67.3% to 82%, linoleic acid content was decreased by 81.5%, from 16.5% to 3%, and linolenic acid
25 content was decreased by 35%, from 5.9% to 3.8%, as compared to wild type cv Westar. Samples in Group 2a with different knockout mutation profiles in the *BnaC.FAD2.a* gene (-18,+166/-18,+166; -3,+1/-3,+1; and -18,+166/-7) demonstrated comparable oleic, linoleic, and linolenic acid composition profiles.

30 Samples with 2-gene knockout mutations in the *BnaC.FAD2.a* and *BnaC.FAD2.b* genes (Group 2b) exhibited increased oleic acid (18:1), reduced linoleic acid (18:2), and reduced linolenic acid (18:3) content as compared to wild type Brassica cv Westar. In Group 2b mutants on average, oleic acid content was increased by 23.4%, from 67.3% to 83%, linoleic acid content was decreased by 84.5%, from 16.5% to 3.2%, and linolenic acid content was decreased by 44.9%, from 5.9% to 3.2%, as compared to wild type cv

Westar. Samples in Group 2b with different in-frame mutations in the *BnaA.FAD2.a* gene and different knockout mutation profiles in the *BnaC.FAD2.a* gene (-18,+166/-7 and -3,+1/-3,+1) had comparable oleic, linoleic, and linolenic acid composition profiles.

Samples with one-gene knockout of the *BnaC.FAD2.a* gene (Group 3)

- 5 demonstrated increased oleic acid (18:1), reduced linoleic acid (18:2), and reduced linolenic acid (18:3) content as compared to wild type Brassica cv Westar. In Group 3 mutants on average, oleic acid content was increased by 20.9%, from 67.3% to 81.3%, linoleic acid content was decreased by 79.7%, from 16.5% to 3.3%, and linolenic acid content was decreased by 38.8%, from 5.9% to 3.6%, as compared to wild type cv
- 10 Westar. Samples in Group 3 with different in-frame mutations in the *BnaA.FAD2.a* gene and different knockout mutation profiles in the *BnaC.FAD2.a* gene (-18,+166/-18,+166; -18,+166/-7; and -7/-7) had comparable oleic, linoleic, and linolenic acid composition profiles.

TABLE 12*T0 plants*

T0 plant	BnaA.Fad2a	BnaC.Fad2a	BnaC.Fad2b
Bn432-a	-3/-6	-3,+1/-7	-13/wt
Bn432-b	-6,+50/-6,+50	-6,+50/-18,+166	wt/wt
Bn432-c	-36,+13/-36,+13	-36,+13/wt	wt/wt
Bn432-d	-29,+4/wt	-29,+4/-29,+4	-4,+1/-4,+1
Bn432-e	-10/wt	-4,+106/-10	-9/wt
Bn432-f	-3/-5,+1	-18/-18	-4/+44
Bn432-g	-3/-3	-11/-3	-8/-8
Bn432-h	wt/wt	-2,+55/-2,+55	wt/wt
Bn468-a	-3,+1/-6	+1/-3	wt/wt
Bn468-b	-8/-16	-3/-3	wt/wt
Bn468-c	wt/wt	wt/wt	-7/wt
Bn468-d	wt/wt	wt/wt	-7/wt
Bn468-e	-7,+20/wt	wt/wt	wt/wt
Bn468-f	wt/wt	-2/wt	wt/wt
Bn468-g	-17,+14/-17,+14	-5/-5	wt/wt
Bn468-h	wt/wt	wt/wt	-4,+44/wt
Bn471-a	-4/-9	-3,+90/-3,+90	wt/wt
Bn471-b	-4,+1/-4,+1	wt/wt	wt/wt
Bn476-a	-6/-6	-8/-8	+1/wt
Bn476-b	-2,+1/-3	wt/wt	-2/wt
Bn476-c	wt/wt	wt/wt	-16/wt
Bn482-a	-3/-3	-5/-18	-4/wt
Bn482-b	-3/-74,+5	-74,+5/-74,+5	wt/wt
Bn482-c	-4/-4	-3/-3	-3/+29
Bn482-d	-3/-3	wt/wt	wt/wt
Bn482-e	-3/-3	-3/-3	wt/wt
Bn482-f	-4/wt	wt/wt	-3/wt
Bn482-g	wt/wt	-5/-5	wt/wt
Bn482-h	-11/-11	-6/-10	wt/wt
Bn482-i	-3/-3	-2/+77	wt/wt
Bn482-j	-7/-3+99	-7/-7	-3/-3
Bn482-k	-3,+1/+4	-3/+95	-4/-4
No. of novel alleles	26	27	13

TABLE 13A*Sequences of mutant T0 BnaA.Fad2a alleles*

5

Bn432-a (-3):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC

TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACC GGCGTCTGGGT
CATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53907)

Bn432-a (-6):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAACCTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTC
GCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA
CGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCT
GGCCTCTCTACTGGGCCTGCCAGGGTGCGTCCTAACC GGCGTCTGGGTCA
TAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53908)

Bn432-b (-6,+50):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CAGTTTCCAAAAGGTCGTTGATCAAAGCTCGCCGCGTTGTTTCATCAAGCC
AGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCG
CTCTTTCTCCTACCTCATCTGGGACCTCATCATAGCCTCCTGCTTCTACTACG
TCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGG
CCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACC GGCGTCTGGGTCATA
GCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53909)

Bn432-c (-36,+13):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAACCGACAACAACAAGAACTCAAGAAAGC
AATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTACCTC
ATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACTTACT
TCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGGCC
TGCCAGGGCTGCTTCCTAACC GGCGTCTGGGTCATAGCCACGAGTTCGGC
CACCACGCCTTCAGCGACTAC (SEQ ID NO:53910)

Bn432-d (-29,+4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTGAGAACTGTTTCAAACGCACGATCCCTCGCTCTTTCTCCTACCTCATC
TGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACTTACTTCC
CTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGGCCTGC
CAAGGCTGCGTCCTAACC GGCGTCTGGGTCATAGCCACGAGTGCGGCCAC
CACGCCTTCAGCGACTAC (SEQ ID NO:53911)

Bn432-e (-10):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTGA
ACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTC
TTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTC
GCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCC

TCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTCATAGC
CCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53912)

Bn432-f (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTTGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53913)

Bn432-f (-5,+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCC
TCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53914)

Bn432-g (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53915)

Bn468-a (-3,+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATC
CCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTA
CTACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCG
CCTGGCCTCTCTACTGGGCCTGCCAGGGTGCGTCCTAACCGGCGTCTGGG
TCATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53916)

Bn468-a (-6):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTC
GCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA

CGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCT
GGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCA
TAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53917)

Bn468-b (-8):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGCCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTCCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGC
TCTTTCTCCTACCTCATCTGGGACATCATATAGCCTCCTGCTTCTACTACGT
CGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGC
CTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCAATCG
CCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53918)

Bn468-b (-16):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCC
TACCTCATCTGGGACATCATATAGCCTCCTGCTTCTACTACGTGCGCCACCA
CTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTAC
TGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCAATAGCCACGAG
TGCAGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53919)

Bn468-c (-7,+20):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
ATCGCCAGCAATAATGGTGGAGAACTCAAGGAAGCAATCCCACCGCACTG
TTTCAAACGCTCGATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATC
ATAGCCTCCTGCTTCTACTACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCA
CCCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTC
CTAACCGGCGTCTGGGTCAATAGCCACGAGTGCGGCCACCACGCCTTCAGC
GACTAC (SEQ ID NO:53920)

Bn468-g (-17,+14):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACAACCTCTTTTC
TCGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGT
CATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53921)

Bn471-a (-4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCC
TCGCTCTTTCTCCTACCTCATCTGGGACATCATATAGCCTCCTGCTTCTACT
ACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC

TGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53922)

Bn471-a (-9):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTAG
AACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCT
CTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGT
CGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGC
CTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCATAG
CCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53923)

Bn471-b (-4,+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTAGTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGCCCACCCCGCCTTCGGCGACTCC (SEQ ID
NO:53924)

Bn476-a (-6):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTC
GCTCTTTCTCCTACCGCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA
CGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCT
GGCCTCTCTACTGGGACTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCA
TAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53908)

Bn476-b (-2,+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTCTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGAT
CCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCT
ACTACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTC
GCCTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGG
GTCATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53925)

Bn476-b (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC

CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53913)

Bn482-a (-3):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53913)

Bn482-b (-3):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53926) 53929

Bn482-b (-74,+5):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCATCAAGA
AAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTA
CCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACT
TACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTG
GGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTCATAGCCCACGAGTG
CGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53927)

Bn482-c (-4):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCC
TCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53922)

Bn482-d (-3):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC

CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACC GGCGTCTGGGT
CATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53926)

Bn482-e (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACC GGCGTCTGGGT
CATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53913)

Bn482-f (-4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCC
TCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACC GGCGTCTGGGTC
ATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53922)

Bn482-h (-11):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTT
TCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTGCG
CACCCTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTC
TCTACTGGGCCTGCCAGGGCTGCGTCCTAACC GGCGTCTGGGTTCATAGCCC
ACGAGTGCGGGCCACCACGCCTTCAGCGACTAA (SEQ ID NO:53928)

Bn482-i (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACC GGCGTCTGGGT
CATAGCCCACGAGTGCGGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53913)

Bn482-j (-3,+99):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTATGGCTTGACCCCGGAGCAGGTGGTGGCCATCGCCAGCAATCTTGGC
GGCAAGCAGGCGCTGGAGACAATGCAGGCGCTGTTGCCGGTGCTGTGCCA

GGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCCAACGCTCGATCCCT CGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT ACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC TGGCCTCTCTACTGGGCCTGCCAAGGGTGCCTCCTAACGGGCGTCTGGGTC ATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53929)
Bn482-j (-7): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCG CTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTAC GTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTG GCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACGGGCTCTGGGTCAT AGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53930)
Bn482-k (+4): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTGTGCGTTCGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGC TCGATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCT GCTTCTACTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCC TACTTCGCCTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACGGGCG TCTGGGTCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53931)
Bn482-k (-3,+1): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA AGTCTGAAACCGACAACATCAAGCGCGTACCCCGCGAGACACCGCCCTTCA CTGTGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATC CCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTA CTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCG CCTGGCCTCTCTACTGGGCCTGCCAGGGCTGCGTCCTAACGGGCGTCTGGG TCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53932)

TABLE 13B*Sequences of mutant T0 BnaC.Fad2a alleles*

Bn432-a (-3,+1): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTGTTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATC CCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTA CTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCG CCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCCTCCTAACGGGCGTCTGGG TCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53933)
--

TCATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53933)
Bn432-a (-7): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTGACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCG CTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTAC GTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTG GCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTGTCAT AGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53934)
Bn432-b (-6,+50): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CAGTTTCCAAAAGGTCGTTGATCAAAGCTCGCCGCGTTGTTTCATCAAGCC AGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCG CTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTAC GTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTG GCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTGTCAT AGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53935)
Bn432-b (-18,+166): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTGTGCGCTGGAGACGGTCCAGCGGCTGTTGCCGGTGCTGTGCCAGGCCCAC GGCTTGACCCCGGAGCAGGTGGTGGCCATCGCCTTTTTCGCTTTCTACAAAC TCTTCCTGGCTAGCGGTACGCGTATTAATTGCGTTGCGCTCACTGCCCGCTT TCCAGTCGGGAAACCTGTCGTGCCACCGCACTGTTTCAAACGCTCGATCCC TCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT ACGTCGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53936)
Bn432-c (-36,+13): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAACCGACAACATCAAGAACTCAAGAAAGC AATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTACCTC ATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACTTACT TCCCTCTCCTCCCTCACCTCTCCCTTACCTCGCCTGGCCCCCTCTCCTGGGCC TGCCAAGGCTGCGTCCTAACGGGCTCTGGGTGTCATCGCCCACGAATGCGGC CACCACGCCTTCAGCGACTAC (SEQ ID NO:53937)
Bn432-d (-29,+4): CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA CTGTGAGAACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTACCTCATC TGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACTTACTTCC CTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTGGGCCTGC

CAAGGGTGCGTCCTAACCGGCGTCTGGGTCATAGCCCACGAGTGCGGCCAC
CACGCCTTCAGCGACTAC (SEQ ID NO:53938)

Bn432-e (-4,+106):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGCCAGGCCCACGGCTTGACCCCTCAGCAGGTGGTGGCCATCGCCAGCAA
TGGCGGCGGCAGGCAGGCGCTGGAGACGGTCCAGCGGCTGTTGCCGGTGC
TGTGCGGGGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGA
TCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTC
TACTACGTGCGCACCACTTACTTCCCTCTCCTCCCTCACCCTCTCTCCTACTT
CGCCTGGCCTCTCTACTGGGCTGCCAAGGGTGCGTCCTAACCGGCGTCTG
GGTCATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53939)

Bn432-e (-10):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACAACATCAAGCGCGTACCCTGCGAGACACCGCCCTTGA
ACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTC
TTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTC
GCCACCTCTTACTTCCCTCTCCTCCCTCACCCTCTCTCCTACTTCGCCTGGCC
TCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCATAGC
CCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53940)

Bn432-f (-18):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCAAG
AAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCT
ACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTGCGCCACCAC
TACTTCCCTCTCCTCCCTCACCCTCTCTCCTACTTCGCCTGGCCTCTCTACT
GGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCATAGCCCACGAGT
GCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53941)

Bn432-g (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53942)

Bn432-g (-11):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTT
TCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTGCG

CACCACTTACTTCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTC
TCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCATAGCCC
ACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53943)

Bn432-h (-2,+55):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTCCGGCAACAGCCGCTGGACCGTCTCCAGCGCCTGCTTGCCGCCATCG
TGGCTGGCGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTC
GATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGC
TTCTACTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTA
CTTCGCCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGCTC
TGGGTTCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53944)

Bn468-a (+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTCCGGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCG
ATCCCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTT
CTACTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACT
TCGCCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCT
GGGTTCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53945)

Bn468-a (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53946)

Bn468-b (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGT
CATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53946)

Bn468-f (-2):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA

CTGTCAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATC
CCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTA
CTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCG
CCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGG
TCATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53947)

Bn468-g (-5):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCT
CGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53948)

Bn471-a (-3,+90):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTTGACCCACGGCGTAACGCGCTTGCTGCTTGGATGCCCCGAGGCATAG
ACTGTACCCCAAAAAACAGTCATAACAAGCCATGAAAACCGGAGAACTC
AAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCT
CCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCAC
CACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCT
ACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTCATAGCCCACG
AGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53949)

Bn476-a (-8):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
GAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGC
TCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGT
CGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGC
CTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTCATAG
CCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53950)

Bn482-a (-5):
CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAAGTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCT
CGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53951)

Bn482-a (-18):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
AAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTA
CCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACT
TACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTG
GGCCTGCCAAGGGTGCCTTAACCGGCGTCTGGGTCATAGCCCACGAGTG
CGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53952)

Bn482-b (-74,+5):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCATCAAGA
AAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTA
CCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACT
TACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGCCTCTCTACTG
GGCCTGCCAAGGGTGCCTTAACCGGCGTCTGGGTCATAGCCCACGAGTG
CGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53953)

Bn482-c (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCTGCCAAGGGTGCCTTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53946)

Bn482-e (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
TACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
CTGGCCTCTCTACTGGGCTGCCAAGGGTGCCTTAACCGGCGTCTGGGT
CATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53946)

Bn482-g (-5):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGGAAGTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCT
CGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACT
ACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCC
TGGCCTCTCTACTGGGCTGCCAAGGGTGCCTTAACCGGCGTCTGGGTC
ATAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53954)

Bn482-h (-6):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTC
GCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTA
CGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCT
GGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTCA
TAGCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAA (SEQ ID NO:53955)

Bn482-h (-10):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTGA
ACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTC
TTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGTC
GCCACCACTTACTTCCCTCTCCTCCCTCACCACTCTCCTACTTCGCCTGGCC
TCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCATAGC
CCACGAGTGCGGCCACCACGCCTTCAGCGACTAA (SEQ ID NO:53940)

Bn482-i (+77):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTCGGCGCTGCTTGCCGGTGCTGTGCCAGGCCCACGGCTTGACCCCGGA
GCAGGTGGTGGCCATCGCCAGCCACGATGGCGGAGAACTCAAGAAAGCAA
TCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCTCTTTCTCCTACCTCAT
CTGGGACATCATCATAGCCTCCTGCTTCTACTACGTCGCCACCACTTACTTC
CCTCTCCTCCCTCACCTCTCTCCTTCTTCGCCTGGCCTCTCTACTGGGCCTG
CCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCATAGCCAACGAGTGCGGCCA
CCACGCCTGCAGCGACTAC (SEQ ID NO:53956)

Bn482-i (-2):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGGGAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATC
CCTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTA
CTACGTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCG
CCTGGCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGG
TCATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53957)

Bn482-j (-7):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAAA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCG
CTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTAC
GTCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTG
GCCTCTCTACTGGGCCTGCCAAGGGTGCGTCCTAACCGGCGTCTGGGTTCAT
AGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53958)

Bn482-k (+95):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
 AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
 CTGTGCGGAATTTATGGAACGTCAGTGGAGCATTTTAAAGACTGAGCCTTTTCG
 TTTTATCTGTTGTTTGTGCGGTGAACGCTCTCCTGAGTAGGACAAATCCGGAG
 AACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCCCTCGCT
 CTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTACTACGT
 CGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGCCTGGC
 CTCTCTACTGGGCCTGCCAAGGGTGCCTTAACCGGCGTCTGGGTCATAG
 CCCACGAGTGCGGCCACACGCCTTCAGCGACTAC (SEQ ID NO:53959)

Bn482-k (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
 AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
 CTGTAGAACTCAAGAAAGCAATCCCACCGCACTGTTTCAAACGCTCGATCC
 CTCGCTCTTTCTCCTACCTCATCTGGGACATCATCATAGCCTCCTGCTTCTAC
 TACGTGCGCCACCACTTACTTCCCTCTCCTCCCTCACCTCTCTCCTACTTCGC
 CTGGCCTCTCTACTGGGCCTGCCAAGGGTGCCTTAACCGGCGTCTGGGT
 CATAGCCACGAGTGCGGCCACACGCCTTCAGCGACTAC (SEQ ID
 NO:53946)

TABLE 13C

Sequences of mutant T0 BnaC.Fad2b alleles

Bn432-a (-13):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
 CCCCCGAAACCAAACCCTCAAACGCGTCCCCTGCGAGACACCACCTCCT
 CAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCTCCTTC
 TCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTCTCCAC
 AGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGGCCCCCTCT
 ACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCATAGCCACG
 AGTGCGGCCACACGCCTTCAGCGACTAC (SEQ ID NO:53960)

Bn432-d (-4,+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
 CCCCCGAAACCAAACCCTCAAACGCGTCCCCTGCGAGACACCACCTTCA
 CTAGAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCC
 CTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTAC
 CACCTCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGC
 CTGGCCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGT
 CATCGCCACGAATGCGGCCACACGCCTTCAGCGACTAC (SEQ ID
 NO:53961)

Bn432-e (-9):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
 CCCCCGAAACCAAAGCCCTCAAACGCGTCCCCTGCGAGACACCACCTTAG
 ACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCT
 CCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTC

TCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGGCC
CCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCATCGC
CCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53962)

Bn432-f (-4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTCACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCT
CGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCA
CCTCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCT
GGCCCCCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCA
TCGCCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53963)

Bn432-f (+44):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCGGAGAATCTCGCTCTCTCCAGGGGAAGCCGAAGTTTCAAAGGTGCG
AGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCG
CTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACC
TCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGG
CCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCAATC
GCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53964)

Bn432-g (-8):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCT
CCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTC
TCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGGCC
CCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCAATAGC
CCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53965)

Bn468-c (-7):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCATCG
CTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACC
TCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGG
CCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCAATC
GCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53966)

Bn468-d (-7):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCG
CTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACC
TCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGG

CCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCATA
GCCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53966)

Bn468-h (-4,+44):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTGGCTGGCGATGGCCACCACCTGCTCCGGGGTCAAGCCGTGGGCGGAC
CTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCTCCT
TCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTCTCC
ACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACCTCGCCTGGCCCCCT
CTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCATCGCCCA
CGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID NO:53967)

Bn476-a (+1):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTCGGGAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCC
ATCCCTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCT
CTACCACCTCTCCACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACC
TCGCCTGGCCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCT
GGGTCATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53968)

Bn476-b (-2):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTGAGAACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCAT
CCCTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCT
ACCACCTCTCCACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACCTC
GCCTGGCCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGG
GTCATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53969)

Bn476-c (-16):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTCAA
GAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCTCCTTCTCC
TACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTCTCCACAGC
CTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACCTCGCCTGACCCCTCTACT
GGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTCATAGCCCACGAGT
GCGGCCACCACGCCTTCAGCGACTAA (SEQ ID NO:53970)

Bn482-a (-4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTGAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCC
TCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACC
ACCTCTCCACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACCTCGCC

TGGCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTC
ATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53971)

Bn482-c (+29):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTCGGCAAGCAGGCGCTGGAGACGGTCCAGCGGCAGACCTCAAGAAAG
CAATCCCACCTCACTGCTTCAAACGCTCCATCCCTCGCTCCTTCTCCTACCT
CCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACCACCTCTCCACAGCCTACT
TCCCTCTCCTCCCCACCCTCTCCCTTACCTCGCCTGGCCCCTCTACTCGGCC
TGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTATCGCCACGAATGCGGC
CACCACGCCTTCAGCGACTAC (SEQ ID NO:53972)

Bn482-c (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTGTCTCCTCCCTCCAAGA
AGTCTGAAACCGACACCATCAAGCGCGTACCCTGCGAGACACCGCCCTTCA
CTGTAGAACTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCC
CTCGCTCCTTCTCCTACCTCCTCTCCGACCTCCTCGTCTCATCCTCCCTCTAC
CACCTCTCCACAGCCCCCTTCCCTCTCCTCCCCACCCTCCCCCTTCCCCCGC
CTGGCCCCTCTTCTGAGCCTGCCAAGGCTGCGTCCTAAAGGGCCTCTGGGT
CATAGCCACGAGTGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53973)

Bn482-f (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCC
CTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTAC
CACCTCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGC
CTGGCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGT
CATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53974)

Bn482-j (-3):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCTAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCC
CTCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTAC
CACCTCTCCACAGCCTACTTCCCTCTCCTCCCCACCCTCTCCCTTACCTCGC
CTGGCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGT
CATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
NO:53974)

Bn482-k (-4):

CAGGATCCATGGGTGCAGGTGGAAGAATGCAAGTCTCTCCTCCCTCCAGCT
CCCCGAAACCAAAACCCTCAAACGCGTCCCCTGCGAGACACCACCCTTCA
CTCAGACCTCAAGAAAGCAATCCCACCTCACTGCTTCAAACGCTCCATCCC

TCGCTCCTTCTCCTACCTCCTCTTCGACATCCTCGTCTCCTCCTCCCTCTACC
 ACCTCTCCACAGCCTACTTCCCTCTCCTCCCCCACCCTCTCCCTTACCTCGCC
 TGGCCCCCTCTACTGGGCCTGCCAAGGCTGCGTCCTAACGGGCCTCTGGGTC
 ATCGCCACGAATGCGGCCACCACGCCTTCAGCGACTAC (SEQ ID
 NO:53975)

TABLE 14

T1 plants used in crosses

T1 plant	T0 parent	<i>BnaA.FAD2.a</i>	<i>BnaC.FAD2.a</i>	<i>BnaC.FAD2.b</i>
Bn432-01	Bn432-a	-3/-6	-3,+1/-7	-13/-13
Bn432-02	Bn432-a	-3/-6	-3,+1/-3,+1	-13/-13
Bn432-03	Bn432-a	-3/-6	-7/-7	-13/-13
Bn432-05	Bn432-b	-6,+50/-6,+50	-18,+166/-18,+166	wt/wt
Bn432-07	Bn432-b	-6,+50/-6,+50	-18,+166/-18,+166	wt/wt

TABLE 15
Mutation profile and oil composition of F2 plants

F2 plant	Group	Mutation profile of FAD2 genes			Oil composition (%)		
		<i>BnaA.FAD2.a</i>	<i>BnaC.FAD2.a</i>	<i>BnaC.FAD2.b</i>	Oleic acid (18:1)	Linoleic acid (18:2)	Linolenic acid (18:3)
Bn432-01 x Bn432-07 F2-04	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	82.70%	2.60%	3.10%
Bn432-01 x Bn432-07 F2-02	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	83.70%	2.60%	3.20%
Bn432-02 x Bn432-05 F2-02	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	83.80%	2.10%	2.80%
Bn432-01 x Bn432-07 F2-03	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	83.80%	2.40%	3.10%
Bn432-01 x Bn432-07 F2-01	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	83.90%	2.40%	3.30%
Bn432-02 x Bn432-05 F2-01	1	-6,+50 / -6,+50	-18,+166 / -18,+166	-13 / -13	84.00%	2.30%	3.00%
Bn432-02 x Bn432-05 F2-05	1	-6,+50 / -6,+50	-3,+1 / -18,+166	-13 / -13	82.00%	2.40%	3.00%
Bn432-02 x Bn432-05 F2-07	1	-6,+50 / -6,+50	-3,+1 / -18,+166	-13 / -13	83.00%	2.30%	3.10%
Bn432-02 x Bn432-05 F2-08	1	-6,+50 / -6,+50	-3,+1 / -18,+166	-13 / -13	83.70%	2.20%	3.10%
Bn432-02 x Bn432-05 F2-06	1	-6,+50 / -6,+50	-3,+1 / -18,+166	-13 / -13	84.50%	2.10%	3.00%
Bn432-02 x Bn432-05 F2-04	1	-6,+50 / -6,+50	-3,+1 / -18,+166	-13 / -13	84.90%	2.10%	2.90%

Bn432-02 x Bn432-05 F2-03	1	-6,+50 / -6,+50	-3,+1 / -3,+1	-13 / -13	85.00%	2.10%	3.20%
Bn432-01 x Bn432-07 F2-12	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	81.30%	2.90%	3.10%
Bn432-01 x Bn432-07 F2-09	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	82.60%	2.70%	3.00%
Bn432-01 x Bn432-07 F2-11	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	83.00%	2.40%	3.10%
Bn432-01 x Bn432-07 F2-08	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	83.50%	2.60%	2.90%
Bn432-01 x Bn432-07 F2-10	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	83.50%	2.70%	3.20%
Bn432-01 x Bn432-07 F2-07	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	83.70%	2.60%	3.00%
Bn432-01 x Bn432-07 F2-13	1	-6,+50 / -6,+50	-7 / -18,+166	-13 / -13	83.80%	2.60%	3.30%
Bn432-01 x Bn432-07 F2-06	1	-6,+50 / -6,+50	-7 / -7	-13 / -13	83.00%	2.70%	3.20%
Bn432-01 x Bn432-07 F2-05	1	-6,+50 / -6,+50	-7 / -7	-13 / -13	83.30%	2.50%	3.20%
Bn432-02 x Bn432-05 F2-09	2a	-6,+50 / -6,+50	-18,+166 / -18,+166	wt/wt	82.30%	3.00%	3.80%
Bn432-01 x Bn432-07 F2-14	2a	-6,+50 / -6,+50	-18,+166 / -18,+166	wt/wt	82.20%	3.10%	3.60%
Bn432-02 x Bn432-05 F2-10	2a	-6,+50 / -6,+50	-3,+1 / -3,+1	wt/wt	82.30%	2.60%	3.60%

Bn432-02 x Bn432-05 F2-11	2a	-6,+50 / -6,+50	-3,+1 / -3,+1	wt/wt	81.00%	3.50%	3.90%
Bn432-02 x Bn432-05 F2-12	2a	-6,+50 / -6,+50	-3,+1 / -3,+1	wt/wt	82.60%	3.00%	3.90%
Bn432-02 x Bn432-05 F2-13	2a	-6,+50 / -6,+50	-3,+1 / -3,+1	wt/wt	81.70%	3.00%	4.20%
Bn432-01 x Bn432-07 F2-15	2a	-6,+50 / -6,+50	-7 / -18,+166	wt/wt	81.80%	3.10%	3.60%
Bn432-02 x Bn432-05 F2-14	2b	-3 / -3	-3,+1 / -3,+1	-13 / -13	83.80%	2.30%	3.10%
Bn432-02 x Bn432-05 F2-15	2b	-3 / -3	-3,+1 / -3,+1	-13 / -13	84.00%	2.20%	3.30%
Bn432-01 x Bn432-07 F2-16	2b	-6 / -6	-7 / -18,+166	-13 / -13	82.00%	2.80%	3.10%
Bn432-01 x Bn432-07 F2-17	2b	-6 / -6	-7 / -18,+166	-13 / -13	82.20%	2.90%	3.40%
Bn432-01 x Bn432-07 F2-18	3	-6 / -6	-7 / -18,+166	wt/wt	81.90%	3.10%	3.70%
Bn432-02 x Bn432-05 F2-17	3	-3 / -3	-18,+166 / -18,+166	wt/wt	80.80%	3.50%	3.40%
Bn432-02 x Bn432-05 F2-16	3	-3 / -3	-18,+166 / -18,+166	wt/wt	80.70%	3.30%	3.80%
Bn432-01 x Bn432-07 F2-20	3	-6 / -6	-7 / -7	wt/wt	81.70%	3.40%	3.70%
Bn432-01 x Bn432-07 F2-19	3	-6 / -6	-7 / -18,+166	wt/wt	81.40%	3.40%	3.30%

Wild type cv Westar	4	wt/wt	wt/wt	wt/wt	68.40%	15.60%	5.20%
Wild type cv Westar	4	wt/wt	wt/wt	wt/wt	66.10%	17.30%	6.50%

TABLE 16*Average oil composition of F2 plants, by group*

Group	Oil composition (%)			Increase/decrease (%) [*]		
	Oleic acid (18:1)	Linoleic acid (18:2)	Linolenic acid (18:3)	Oleic acid (18:1)	Linoleic acid (18:2)	Linolenic acid (18:3)
1	83.46%	2.44%	3.09%	24.1%	-85.1%	-47.3%
2a	81.99%	3.04%	3.80%	21.9%	-81.5%	-35.0%
2b	83.00%	2.55%	3.23%	23.4%	-84.5%	-44.9%
3	81.30%	3.34%	3.58%	20.9%	-79.7%	-38.8%
4	67.25%	16.45%	5.85%	NA	NA	NA

*Compared to wild type (group 4)

5

OTHER EMBODIMENTS

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A Brassica plant, plant part, or plant cell comprising an induced mutation in one or more FAD2 gene copies, oil produced from the plant, plant part, or plant cell has increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell.
2. The Brassica plant, plant part, or plant cell of claim 1, wherein each of the induced mutations is within a nucleic acid sequence as set forth in any of SEQ ID NOS:1-3, or within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.
3. The Brassica plant, plant part, or plant cell of claim 1 or claim 2, wherein the induced mutation comprises an in-frame or frameshift mutation, and wherein the in-frame or frameshift mutation comprises an insertion or deletion selected from the group consisting of:
 - a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
 - a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
 - a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
 - a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
 - a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
 - a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and
 - an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.
4. The Brassica plant, plant part, or plant cell of any one of claims 1 to 3, wherein each induced mutation was induced by a rare-cutting endonuclease.

5. The Brassica plant, plant part, or plant cell of any one of claims 1 to 4, wherein the plant, plant part, or plant cell does not contain a transgene.
6. The Brassica plant, plant part, or plant cell of any one of claims 1 to 4, wherein the plant, plant part, or plant cell comprises a transgene encoding a protein, and wherein the protein is selected from the group consisting of a plant 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an aryloxyalkanoate dioxygenase (AAD) protein, a phosphinothricin N-acetyltransferase (PAT) protein, an acetohydroxyacid synthase large subunit protein, a p-hydroxyphenylpyruvate dioxygenase (hppd) protein, and a dicamba monooxygenase (DMO) protein.
7. The Brassica plant, plant part, or plant cell of any one of claims 1 to 6, wherein the plant part is a seed.
8. A Brassica plant or plant part that is a progeny of the Brassica plant or plant part of claim 1, wherein the progeny plant or plant part comprises a mutation in at least one FAD2 gene copy.
9. A method for producing a Brassica plant having increased oleic acid content and decreased linolenic acid content, the method comprising:
 - providing a population of Brassica cells comprising functional FAD2 gene copies,
 - contacting the population of Brassica cells with one or more rare-cutting endonucleases targeted to the one or more FAD2 gene copies,
 - regenerating Brassica plants from the population of Brassica cells contacted with the one or more rare-cutting endonucleases, and
 - selecting a Brassica plant with a mutation in one or more FAD2 gene copies,
 wherein the selected Brassica plant produces oil having increased oleic acid content and decreased linolenic acid content as compared to oil produced by a corresponding wild type Brassica plant.
10. The method of claim 9, wherein the Brassica cells are selected from the group consisting of protoplast cells, embryo cells, callus cells, leaf cells, and petiole explant cells.
11. The method of claim 9 or claim 10, comprising transforming the Brassica cells with one or more vectors encoding the one or more rare-cutting endonucleases.

12. The method of claim 11, wherein the one or more rare-cutting endonucleases are transcription activator-like effector (TALE) nucleases.
13. The method of claim 12, wherein each of the one or more TALE nucleases is targeted to a sequence within SEQ ID NOS:1-3, or to a sequence having at least 90% identity to a sequence within SEQ ID NOS:1-3.
14. The method of claim 12, wherein each of the one or more TALE nucleases is targeted to a sequence within any of the sequences set forth in SEQ ID NOS:9-29, or to a sequence having at least 90% identity to any of SEQ ID NOS:9-29.
15. A method for generating a Brassica plant that produces oil having increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell, the method comprising:
providing a first Brassica plant comprising at least one induced mutation in an endogenous FAD2 gene copy;
providing a second Brassica plant comprising wild type FAD2 gene copies; and
crossing the first Brassica plant with the second Brassica plant;
thereby producing a plurality of progeny seed, wherein the progeny seed produce plants that comprise an induced mutation in at least one endogenous FAD2 gene copy.
16. The method of claim 15, wherein each of the induced mutations is at a sequence within any of SEQ ID NOS:1-3, or within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.
17. The method of claim 15 or claim 16, wherein the induced mutation comprises an in-frame or frameshift mutation, and wherein the in-frame or frameshift mutation comprises an insertion or deletion selected from the group consisting of:
a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;
a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and

an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.

18. The method of any one of claims 15 to 17, wherein each induced mutation was induced by a rare-cutting endonuclease.

19. The method of any one of claims 15 to 18, wherein the first Brassica plant and the second Brassica plant do not contain a transgene.

20. The method of any one of claims 15 to 18, wherein the first Brassica plant or the second Brassica plant comprises a transgene encoding a protein, and wherein the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid synthase large subunit protein, a hppd protein, and a DMO protein.

21. A method for generating a Brassica plant that produces seed oil having increased oleic acid content and decreased linolenic acid content as compared to oil produced from a corresponding wild type Brassica plant, plant part, or plant cell, the method comprising:

providing a first Brassica plant comprising at least a first induced mutation in a first endogenous FAD2 gene copy;

providing a second Brassica plant comprising at least a second induced mutation in a second endogenous FAD2 gene copy; and

crossing the first Brassica plant with the second Brassica plant;

thereby producing a plurality of progeny seed, wherein the progeny seed produce plants that comprise the first and second induced mutations in the first and second endogenous FAD2 gene copies.

22. The method of claim 21, wherein each of the induced mutations is at a sequence within any of SEQ ID NOS:1-3.

23. The method of claim 21 or claim 22, wherein at least one of the first and second induced mutations comprises an in-frame or frameshift mutation, and wherein the in-frame or frameshift mutation comprises an insertion or deletion selected from the group consisting of:

a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and

an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.

24. The method of any one of claims 21 to 23, wherein each induced mutation was induced by a rare-cutting endonuclease.

25. The method of any one of claims 21 to 24, wherein the first Brassica plant and the second Brassica plant do not contain a transgene.

26. The method of any one of claims 21 to 24, wherein the first Brassica plant or the second Brassica plant comprises a transgene encoding a protein, and wherein the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid synthase large subunit protein, a hppd protein, and a DMO protein.

27. A method for producing Brassica oil having increased oleic acid content and reduced linoleic acid content, comprising:

- (a) providing a Brassica plant or plant part comprising an induced mutation in one or more FAD2 gene copies, and
- (b) producing oil from the plant or plant part.

28. The method of claim 27, wherein the induced mutation comprises an in-frame or frameshift mutation, and wherein the in-frame or frameshift mutation comprises an insertion or deletion selected from the group consisting of:

- a 3 bp deletion at positions 98-100 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

- a 6 bp deletion at positions 97-102 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

- a 3 bp deletion at positions 99-101 and a 1 bp insertion at position 99 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

- a 7 bp deletion at positions 98-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

- a 13 bp deletion at positions 92-104 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3;

- a 6 bp deletion at positions 96-101 and a 50 bp insertion at position 96 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3; and

- an 18 bp deletion at positions 101-118 and a 166 bp insertion at position 101 of any of SEQ ID NOS:1-3 or at the corresponding positions within a sequence having at least 90% identity to any of SEQ ID NOS:1-3.

29. The method of claim 27, wherein each induced mutation was induced by a rare-cutting endonuclease.
30. The method of claim 27 or claim 28, wherein the first Brassica plant and the second Brassica plant do not contain a transgene.
31. The method of claim 27 or claim 28, wherein the first Brassica plant or the second Brassica plant comprises a transgene encoding a protein, and wherein the protein is selected from the group consisting of a plant EPSPS protein, a modified plant EPSPS protein, a bacterial EPSPS protein, an agrobacterium CP4 EPSPS protein, an AAD protein, a PAT protein, an acetohydroxyacid synthase large subunit protein, a hppd protein, and a DMO protein.




FIG. 1

Plant: c02

	(SEQ ID NO)
BnaA.FAD2.a	WT (53892)
TCC TCC C C T C A C C C T C T C T C C T A C T T C G C C T G G C C T C T C T A C T G G G C C T G C C A G G G C T	
BnaC.FAD2.a	WT (53893)
T C C T C C C C T C A C C C T C T C T C C T A C T T C G C C T G G C C T C T C T A C T G G G C C T G C C A A G G G T	
	-4bp (53894)
T C C T C C C C T C A C C C T C T C T C C T A C T T - - - - T G G C C T C T C T A C T G G G C C T G C C A A G G G T	
BnaC.FAD2.b	WT (53895)
T C C T C C C C C A C C C T C T C C C T T A C C T C G C C T G G C C C C T C T A C T G G G C C T G C C A A G G C T	
BnaA.FAD2.b-like	WT (53896)
T A C T T C C C C T C T C C C T T A C C T C G C C T G A C C C C T C T A C T G G G C C T G C C A A G G C T G C G	

FIG. 2

Plant: e01		(SEQ ID NO)
BnaA.FAD2.a	TCCTCCCTCACCCTCTCTCCTACTTCGGCTGGCCCTCTCTACTGGGCCTGCCAGGGCT	WT (53892)
BnaC.FAD2.a	TCCTCCCTCACCCTCTCTCCTACT-----TGGCCTCTCTACTGGGCCTGCCAAGGGT	-5bp (53897)
	TCCTCCCTCACCCTCTCTCCTACTT-----TGGCCTCTCTACTGGGCCTGCCAAGGGT	-4bp (53894)
BnaC.FAD2.b	TCCTCCCCCACCCTCTCCCTTACCTCGCCTGGCCCCCTCTACTGGGCCTGCCAAGGGT	WT (53895)
BnaA.FAD2.b-like	TACTTCCCTCTCCCTTACCTCGCCTGACCCCTCTACTGGGCCTGCCAAGGCTGG	WT (53896)

FIG. 3

Plant: e02		(SEQ ID NO)
BnaA.FAD2.a	TCCTCCCTCACCCCTCTCTCCTACTTCGCCCTGGCCCTCTCTACTGGGCCCTGCCAGGGCT	WT (53892)
BnaC.FAD2.a	TCCTCCCTCACCCCTCTCTCCTACTTCGCCCTGGCCCTCTCTACTGGGCCCTGCCAAGGGT	WT (53893)
	TCCTCCCTCACCCCTCTCTCCTACT-----TGGCCCTCTCTACTGGGCCCTGCCAAGGGT	-5bp (53897)
BnaC.FAD2.b	TCCTCCCCCACCCCTCTCTCCTTACCTCGCCCTGGCCCTCTCTACTGGGCCCTGCCAAGGGT	WT (53895)
BnaA.FAD2.b-like	TACTCCCTCTCTCCCTTACCTCGCCCTGACCCCTCTCTACTGGGCCCTGCCAAGGCTGCG	WT (53896)

FIG. 4

Plant: f02		(SEQ ID NO)
BnaA.FAD2.a	TCCTCCCTCACCCTCTCTCCTACTTCGCCCTGGCCCTCTCTACTGGGCCCTGCCAGGGCT	WT (53892)
BnaC.FAD2.a	TCCTCCCTCACCCTCTCTCCTACTTCGCCCTGGCCCTCTCTACTGGGCCCTGCCAAGGGT	WT (53893)
	TCCTCCCTCACCCTCTCTCCTACT-----TGGCCCTCTCTACTGGGCCCTGCCAAGGGT	-5bp (53897)
BnaC.FAD2.b	TCCTCCCTCACCCTCTCTCCTTACCCTGCCCTGGCCCTCTCTACTGGGCCCTGCCAAGGGT	WT (53895)
BnaA.FAD2.b-like	TACTTCCCTCTCCCTTACCCTGCCCTGACCCCTCTACTGGGCCCTGCCAAGGCTGCG	WT (53896)

FIG. 5

FIG. 6

WT			(SEQ ID NO)
	BnaA.FAD2.a	CCCTGCGAGACACCGCCCTTCACTGTCTGGAGAACTCAAGAAAGCAATCCCACCGCACT	(53898)
	BnaC.FAD2.a	CCCTGCGAGACACCGCCCTTCACTGTCTGGAGAACTCAAGAAAGCAATCCCACCGCACT	(53898)
	BnaC.FAD2.b	CCCTGCGAGACACCAACCCCTTCACTGTCTGGAGACCTCAAGAAAGCAATCCCACCTCACT	(53899)
Plant:Bn432-a			
	BnaA.FAD2.a	CCCTGCGAGACACCGCCCTTCACTG---GAGAACTCAAGAAAGCAATCCCACCGCACT	-3bp (53900)
		CCCTGCGAGACACCGCCCTTCACTG-----AACTCAAGAAAGCAATCCCACCGCACT	-6bp (53901)
	BnaC.FAD2.a	CCCTGCGAGACACCGCCCTTCACTGTT--AGAACTCAAGAAAGCAATCCCACCGCACT	-3bp, +1bp (53902)
		CCCTGCGAGACACCGCCCTTCACTG-----ACTCAAGAAAGCAATCCCACCGCACT	-7bp (53903)
	BnaC.FAD2.b	CCCTGCGAGACACCAACCCCTTCACTGTCTCTGGAGACCTCAAGAAAGCAATCCCACCTCACT	WT (53899)
		CCCTGCGAGACACCAACCCCT-----CCTCAAGAAAGCAATCCCACCTCACT	-13bp (53904)
Plant:Bn432-b			
	BnaA.FAD2.a	CCCTGCGAGACACCGCCCTTCACTGTTTCCAAAAGTGCTGTGATCAAAAGCTCGCCGCG <u>TTGTTTCATCAAGCCAGAACTCAAGAAAGCAATCCCACCGCACT</u>	+50bp, (53905) -6bp
	BnaC.FAD2.a	CCCTGCGAGACACCGCCCTTCACTGTTTCCAAAAGTGCTGTGATCAAAAGCTCGCCGCG <u>TTGTTTCATCAAGCCAGAACTCAAGAAAGCAATCCCACCGCACT</u>	+50bp, (53905) -6bp
		CCCTGCGAGACACCGCCCTTCACTGTCTGCTGGAGACGGTCCAGCGGCTGTGCCGGTG <u>CTGTGCCAGGCCACGGCTTGACCCCGGAGCAGGTGGTGGCCATCGCCTTTTTCGGTT</u> <u>TCTACAAACTCTTCCTGGCTAGCGGTACGGTATTAAATTGCTGCGCTCACTGCCCG</u> <u>CTTCCAGTCGGGAAACCTGTCTGTCGCCACCGCACT</u>	-18bp, (53906) +166bp
	BnaC.FAD2.b	CCCTGCGAGACACCAACCCCTTCACTGTCTCTGGAGACCTCAAGAAAGCAATCCCACCTCACT	WT (53899)

FIG. 7

