

A&A Ref: 153244

PUBLICATION PARTICULARS AND ABSTRACT  
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2005/07614

20 September 2005

12.4.06

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

A61K A61P

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

AstraZeneca AB

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

GRIND, Margaretha

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
	33 GB	31 0306615.6	32 22 March 2003

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

Use of low molecular weight thrombin inhibitors in cholesterol-lowering therapy

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	58
------------------	----

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.  
~~The figure of the drawing to which the abstract refers is attached.~~

## REVISED VERSION

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
30 September 2004 (30.09.2004)

PCT

(10) International Publication Number  
WO 2004/082702 A1

- (51) International Patent Classification<sup>7</sup>: A61K 38/05, A61P 7/02, 3/06, 3/10
- (21) International Application Number: PCT/SE2004/000417
- (22) International Filing Date: 19 March 2004 (19.03.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0306615.6 22 March 2003 (22.03.2003) GB
- (71) Applicant (for all designated States except US): ASTRAZENECA AB [SE/SE]; S- 151 85 Södertälje (SE).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): GRIND, Margaretha [SE/GB]; AstraZeneca R & D Charnwood, Bakersell Road, Loughborough, Leicestershire LE11 5RH (GB).
- (74) Agent: GLOBAL INTELLECTUAL PROPERTY; AstraZeneca AB, S- 151 85 Södertälje (SE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declaration under Rule 4.17:**  
— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- Published:**  
— with international search report
- (88) Date of publication of the revised international search report: 24 March 2005
- (15) Information about Correction: see PCT Gazette No. 12/2005 of 24 March 2005, Section II
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF LOW MOLECULAR WEIGHT THROMBIN INHIBITORS IN CHOLESTEROL-LOWERING THERAPY

**Abstract:** According to the invention there is provided the use of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, for the manufacture of a medicament for use in cholesterol-lowering therapy and/or modification of lipid (triglyceride), lipoprotein, and apolipoprotein, profiles associated with an increased risk of cardiovascular complications.

## USE OF LOW MOLECULAR WEIGHT THROMBIN INHIBITORS IN CHOLESTEROL-LOWERING THERAPY

### Field of the Invention

5

This invention relates to a new use of low molecular weight thrombin inhibitors.

### Background and Prior Art

10

It is well known that high levels of cholesterol are associated with heart disease. More than half of all US citizens are understood to have cholesterol levels that exceed those recommended, and one in five has cholesterol levels that are considered high.

15

Cholesterol is involved in the production and maintenance of cell membranes, as well as the production of sex hormones (including progesterone, testosterone, estradiol and cortisol), bile salts and Vitamin D. It is formed primarily in the liver, but also in other parts of the body, such as  
20 the small intestine.

25

In healthy individuals, all of the cholesterol that is needed to perform the above-mentioned functions is produced naturally. However, in a typical blood test, of the amount of cholesterol circulating in blood, about 85% is endogenous, the other 15% arising from external sources. Dietary  
cholesterol usually originates from meat, poultry, fish, seafood and dairy products. In this respect, high consumption levels of these foodstuffs may give rise to increased cholesterol levels in the bloodstream.

Increased cholesterol levels in serum have been associated with atherosclerosis, which is known to increase significantly the risk of blood vessel blockage (stenosis), and thus the likelihood of angina pectoris, myocardial infarction and other cardiovascular complications, such as  
5 stroke.

Cholesterol is insoluble in aqueous environments and thus needs to be transported within the bloodstream by apolipoproteins (Apos). When apolipoproteins are associated with cholesterol, complexes known as  
10 lipoproteins are formed. The density of these lipoproteins is determined by the amount of protein in the molecule and, in this respect, low-density lipoproteins (LDLs), which are the major cholesterol carrier in the blood, are known to have more of the negative effects mentioned herein than protective high-density lipoproteins (HDLs). High levels of LDLs are thus  
15 associated with atherosclerosis, whereas greater levels of HDLs are understood to provide some protection against stenosis, and hence coronary risk, by way of removal of excess cholesterol (transporting it to the liver for disposal).

20 A third group of carrier molecules, very low-density lipoproteins (VLDLs) are converted to LDLs following the delivery of triglycerides to the muscles and adipose tissue. Triglycerides are a mixture of fatty acids and glycerol and are the major components of lipids circulating in blood. Like cholesterol, triglycerides are substances that are found endogenously in the  
25 bloodstream, and may be deposited in adipose tissue. Triglycerides contain high-energy fatty acids which provide much of the fuel needed for normal cellular function. However, an excessive amount of triglycerides, or VLDLs, in the bloodstream can result in similar problems to those

associated with high cholesterol and LDL levels, as well as obesity and diabetes.

Thus, levels of HDLs, LDLs, total cholesterol and triglycerides are all key indicators in determining the risk of atherosclerosis and associated cardiovascular disorders, such as coronary artery diseases (e.g. angina pectoris, myocardial infarction, etc.), stroke (including cerebro-vascular accident and transient ischaemic attack), peripheral arterial occlusive disease, obesity and diabetes. Patients with high overall cholesterol and/or triglycerides levels are at a significant risk, irrespective of whether or not they also have a favourable HDL level. Patients with normal cholesterol levels but low HDL levels are also at increased risk. Recently, it has also been noted that the level of risk of cardiovascular disease associated with high levels of apolipoprotein B (ApoB; which carries lipids in VLDLs and LDLs), and/or low levels of apolipoprotein A-I (ApoA-I; which carries lipids in HDLs), is extremely high.

There are numerous factors that influence cholesterol and triglyceride levels, including diet, age, weight, gender, genetics, diseases (such as diabetes) and lifestyle.

Positive changes in relation to diet, lifestyle and exercise are often insufficient to decrease the risk of cardiovascular problems. In such instances, cholesterol- and/or triglyceride-lowering medication may be prescribed.

Drugs that reduce LDL levels in serum can prevent or reduce the build-up of artery blocking plaques, and can reduce the risk of plaque rupture and associated thrombo-embolic complications. There are several types of drugs

that can help reduce blood cholesterol levels. The most commonly prescribed are the statins, HMG-CoA reductase inhibitors, such as lovastatin, pravastatin, fluvastatin, simvastatin, atorvastatin, pitavastatin and rosuvastatin (e.g. rosuvastatin-calcium). These drugs prevent directly the formation of cholesterol in the liver and thus reduce the risk of cardiovascular disease. Other prescribed drug categories include resins (such as cholestyramine and colestipol), which act by binding bile acids, so causing the liver to produce more of the latter, and using up cholesterol in the process. Further, the B vitamin niacin has been reported at high doses to lower triglycerides and LDL levels in addition to increasing HDL levels. Fibrates (such as gemfibrozil and fenofibrate) are known to lower triglycerides and can increase HDL levels.

However, some of these drugs are known to have side effects, including liver damage. Hence, there is a need for alternative and/or more effective drugs for use in cholesterol-lowering therapy.

The early development of low molecular weight inhibitors of thrombin has been described by Claesson in *Blood Coagul. Fibrinol.* (1994) 5, 411. Low molecular weight thrombin inhibitors (and prodrugs thereof) have been described more recently in US Patent N° 4,346,078; International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/17860, WO 96/24609, WO 96/25426, WO 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/23499, WO 97/30708, WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO 97/11693, WO 97/24135, WO 97/47299, WO 98/01422, WO 98/57932, WO 99/29664, WO 98/06741, WO 99/37668, WO 99/37611, WO 98/37075, WO 99/00371, WO 99/28297, WO 99/29670, WO 99/40072, WO 99/54313, WO 96/31504,

WO 00/01704, WO 00/08014, WO 00/35859, WO 00/35869, WO 00/42059,  
WO 00/61577, WO 00/61608, WO 00/61609, WO 01/87879, WO 02/14270,  
WO 02/44145, WO 03/000653, WO 04/000818 and WO 04/014894; and  
European Patent Applications 648 780, 468 231, 559 046, 641 779, 185 390,  
5 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642,  
669 317, 601 459 and 623 596.

In particular, international patent application WO 94/29336 discloses a  
group of compounds, including HOOC-CH<sub>2</sub>-(*R*)Cgl-(*S*)Aze-Pab-H (in which  
10 Cgl represents cyclohexylglycyl, Aze represents azetidine-2-carboxyl and  
Pab-H represents 4-amidinobenzylamino), which is also known as  
melagatran (see Example 1 of WO 94/29336). International Patent  
Application WO 97/23499 discloses prodrugs of *inter alia* melagatran.

15 More recently, international patent application WO 02/44145 discloses  $\alpha$ -  
hydroxy acid-based low molecular weight thrombin inhibitors and prodrugs  
thereof.

To the applicant's knowledge, none of the above-mentioned documents  
20 disclose or suggest the direct use a low molecular weight thrombin inhibitor  
or a prodrug thereof in cholesterol-lowering therapy and/or modifications of  
lipid (including triglyceride), lipoprotein, or apolipoprotein, profiles.

### Disclosure of the Invention

25

We have found, surprisingly, that administration of a low molecular weight  
thrombin inhibitor may give rise to reduced levels of lipids, such as total  
cholesterol, LDLs (i.e. LDL-cholesterol) and triglycerides in the  
bloodstream, in addition to increasing HDL (i.e. HDL-cholesterol) levels.

According to a first aspect of the invention there is provided the use of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, for the manufacture of a medicament for use in  
5 cholesterol-lowering therapy.

When employed in the context of the present invention and disclosure, the term "cholesterol-lowering therapy" includes any therapy that results in beneficial modifications of serum profiles of total cholesterol, lipids  
10 (including triglycerides), lipoproteins or apolipoproteins, and will thus be understood to encompass the terms "lipid-modifying therapy" and "lipid-(and triglyceride-) lowering therapy", as well as the treatment of hyperlipidaemias (i.e. the elevation of lipids in the bloodstream), including hypercholesterolaemia (high cholesterol levels in the blood; including  
15 primary and secondary (combined) hypercholesterolaemia), hyperlipoproteinemia (elevated plasma lipoproteins levels) and/or hypertriglyceridemia (high triglyceride levels in the blood). The term will thus be understood to include types I, II (IIa and IIb), III, IV and/or V hyperlipoproteinaemia, as well as secondary hypertriglyceridaemia and/or  
20 familial lecithin cholesterol acyltransferase deficiency, but in principle includes any treatment of a patient which results in a decrease in serum levels of cholesterol, LDLs, VLDLs, triglycerides and/or ApoB, and/or an increase in serum levels of HDLs and/or ApoA-I.

25 According to a second aspect of the invention there is provided a cholesterol-lowering therapy method, which method comprises the administration of a low molecular weight thrombin inhibitor, or a pharmaceutically acceptable derivative thereof, to a patient in need of such therapy.

For the avoidance of doubt, in the context of this disclosure, the terms “treatment”, “therapy” and “therapy method” include the therapeutic and/or prophylactic treatment of patients in need of modifications of cholesterol, lipid (including triglyceride), lipoprotein and/or apolipoprotein profiles.

“Pharmaceutically acceptable derivatives” of thrombin inhibitors includes salts (e.g. pharmaceutically-acceptable non-toxic organic or inorganic acid addition salts) and solvates. It will be appreciated that the term further includes derivatives that have, or provide for, the same biological function and/or activity as any relevant inhibitor. Thus, for the purposes of this invention, the term also includes prodrugs of thrombin inhibitors.

The term “low molecular weight thrombin inhibitor” will be understood by those skilled in the art. The term may also be understood to include any composition of matter (e.g. chemical compound) which inhibits thrombin to an experimentally determinable degree in *in vivo* and/or in *in vitro* tests, and which possesses a molecular weight of below 2,000, preferably below 1,000.

Preferred low molecular weight thrombin inhibitors include low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors.

The term “low molecular weight peptide-based, amino acid-based, and/or peptide analogue-based, thrombin inhibitors” will be well understood by one skilled in the art to include low molecular weight thrombin inhibitors with one to four peptide linkages, and includes those described in the review paper by Claesson in *Blood Coagul. Fibrin.* (1994) 5, 411, as well as those disclosed in

US Patent N° 4,346,078; International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/17860, WO 96/24609, WO 96/25426, WO 96/32110, WO 97/01333, WO 97/02284, 5 WO 97/15190, WO 97/30708, WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO 97/11693, WO 97/24135, WO 97/47299, WO 98/01422, WO 98/57932, WO 99/29664, WO 98/06741, WO 99/37668, WO 99/37611, WO 98/37075, WO 99/00371, WO 99/28297, WO 99/29670, WO 99/40072, WO 99/54313, WO 96/31504, WO 00/01704, WO 00/08014, WO 00/35859, 10 WO 00/35869, WO 00/42059, WO 00/61577, WO 00/61608, WO 00/61609, WO 01/87879, WO 02/14270, WO 02/44145, WO 03/000653 and WO 04/000818; and European Patent Applications 648 780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642, 669 317, 601 459 and 623 596, the specific and generic 15 disclosures in all of which documents are hereby incorporated by reference.

Preferred low molecular weight peptide-based thrombin inhibitors include those described generically and specifically in international patent application WO 98/37075, including the compound that is the subject of Claim 8 of that 20 application as published (1-methyl-2-[N-(4-amidinophenyl)aminomethyl]-benzimidazol-5-yl-carboxylic acid, N-(2-pyridyl)-N-(2-hydroxycarbonyl-ethyl)amide) and prodrugs thereof, such as the compound that is the subject of Claim 10 of that application as published (1-methyl-2-[N-[4-(N-*n*-hexyloxycarbonylamidino)phenyl]aminomethyl]benzimidazol-5-yl-carboxylic 25 acid, N-(2-pyridyl)-N-(2-ethoxycarbonyl-ethyl)amide) and those described in international patent application WO 04/014894.


Preferred low molecular weight peptide-based thrombin inhibitors also include HOOC-CH<sub>2</sub>-(*R*)Cha-Pic-Nag-H (wherein Cha represents

9

cyclohexylalanine, Pic represents (*S*)-pipecolinic acid and Nag represents noragmatine; known as inogatran; see International Patent Application WO 93/11152) and, especially, HOOC-CH<sub>2</sub>-(*R*)Cgl-(*S*)Aze-Pab-H (known as melagatran; see above and International Patent Application WO 94/29336).

5

Further thrombin inhibitors include those of the formula I,



10 wherein

R<sup>a</sup> represents -OH or -CH<sub>2</sub>OH;


R<sup>1</sup> represents at least one optional halo substituent;

R<sup>2</sup> represents one or two C<sub>1-3</sub> alkoxy substituents, the alkyl parts of which substituents are themselves substituted with one or more fluoro substituents


15 (i.e. R<sup>2</sup> represents one or two fluoroalkoxy(C<sub>1-3</sub>) groups);

Y represents -CH<sub>2</sub>- or -(CH<sub>2</sub>)<sub>2</sub>-; and

R<sup>3</sup> represents a structural fragment of formula I(i) or I(ii):



I(i)



I(ii)

20 wherein

R<sup>4</sup> represents H or one or more fluoro substituents; and


10

one or two of  $X_1$ ,  $X_2$ ,  $X_3$  and  $X_4$  represent -N- and the others represent -CH-.


Preferred compounds of formula I include:

5

(a) Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab:




10 (b) Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF):



(c) Ph(3-Cl)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab:

15




The term "prodrug" of a low molecular weight thrombin inhibitor includes any compound that, following oral or parenteral administration, is metabolised *in vivo* to form a low molecular weight thrombin inhibitor (as defined herein), in an experimentally-detectable amount, and within a predetermined time (e.g. within a dosing interval of between 6 and 24 hours (i.e. once to four times daily)). For the avoidance of doubt, the term "parenteral" administration includes all forms of administration other than oral administration.

10

Prodrugs of the thrombin inhibitor melagatran that may be mentioned include those disclosed in international patent application WO 97/23499. Preferred prodrugs are those of the formula  $R^1O_2C-CH_2-(R)Cgl-(S)Aze-Pab-OH$  (see the list of abbreviations above or in WO 97/23499), wherein  $R^1$  represents  $C_{1-10}$  alkyl or benzyl, such as linear or branched  $C_{1-6}$  alkyl (e.g.  $C_{1-4}$  alkyl, especially methyl, propyl and, particularly, ethyl) and the OH group replaces one of the amidino hydrogens in Pab.


Preferred prodrugs of compounds of formula I that may be mentioned include those of formula Ia,



Ia

wherein  $R^{3a}$  represents a structural fragment of formula I(iii) or I(iv):

25



wherein  $R^5$  represents  $OR^6$  or  $C(O)OR^7$ ;

5  $R^6$  represents H,  $C_{1-10}$  alkyl,  $C_{1-3}$  alkylaryl or  $C_{1-3}$  alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also optionally substituted by one  
10 or more halo substituents);


$R^7$  represents  $C_{1-10}$  alkyl (which latter group is optionally interrupted by one or more oxygen atoms), or  $C_{1-3}$  alkylaryl or  $C_{1-3}$  alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally  
15 substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also optionally substituted by one or more halo substituents); and

$R^a$ ,  $R^1$ ,  $R^2$ ,  $Y$ ,  $R^4$ ,  $X_1$ ,  $X_2$ ,  $X_3$  and  $X_4$  are as hereinbefore defined.


20 Preferred prodrugs of compounds of formula I are methoxyamidine prodrugs thereof. Hence preferred compounds of formula Ia include:

(i)  $Ph(3-Cl)(5-OCHF_2)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe)$ :

13




(ii) Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe):



5

(iii) Ph(3-Cl)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe):



10

Compounds of formulae I and Ia may be made in accordance with techniques described in international patent application WO 02/44145.

In accordance with the invention, thrombin inhibitors and derivatives thereof may be administered orally, intravenously, subcutaneously, buccally, rectally, dermally, nasally, tracheally, bronchially, topically, by any other parenteral route, or *via* inhalation, in the form of a pharmaceutical preparation comprising the thrombin inhibitor or prodrug in a pharmaceutically acceptable dosage form. Depending on the disorder, and the patient, to be treated, as well as the route of administration, the compositions may be administered at varying doses.

Preferred modes of delivery are systemic. For melagatran and derivatives thereof, preferred modes of administration are parenteral, more preferably intravenous, and especially subcutaneous. For prodrugs of melagatran and compounds of formula Ia, preferred modes of administration are oral.

In the therapeutic treatment of mammals, and especially humans, thrombin inhibitors, prodrugs of thrombin inhibitors, and derivatives of either will generally be administered as a pharmaceutical formulation in admixture with a pharmaceutically acceptable adjuvant, diluent or carrier, which may be selected with due regard to the intended route of administration and standard pharmaceutical practice.

Suitable formulations for use in administering thrombin inhibitors are known in the art, and include those known from US Patent N° 4,346,078; International Patent Applications WO 93/11152, WO 93/18060, WO 93/05069, WO 94/20467, WO 94/29336, WO 95/35309, WO 95/23609, WO 96/03374, WO 96/06832, WO 96/06849, WO 96/17860, WO 96/24609, WO 96/25426, WO 96/32110, WO 97/01338, WO 97/02284, WO 97/15190, WO 97/30708, WO 97/40024, WO 97/46577, WO 98/06740, WO 97/49404, WO 97/11693, WO 97/24135, WO 97/47299, WO 98/01422, WO 98/57932, WO

99/29664, WO 98/06741, WO 99/37668, WO 99/37611, WO 98/37075, WO 99/00371, WO 99/28297, WO 99/29670, WO 99/40072, WO 99/54313, WO 96/31504, WO 00/01704, WO 00/08014, WO 00/35859, WO 00/35869, WO 00/42059, WO 00/61577, WO 00/61608, WO 00/61609, WO 01/37879, WO 5 02/14270, WO 02/44145, WO 03/000653 and WO 04/000818; and European Patent Applications 648 780, 468 231, 559 046, 641 779, 185 390, 526 877, 542 525, 195 212, 362 002, 364 344, 530 167, 293 881, 686 642, 669 317, 601 459 and 623 596, the disclosures in all of which documents are hereby incorporated by reference.

10

Suitable formulations for use with melagatran, derivatives and prodrugs thereof are described in the literature, for example as described in *inter alia* international patent applications WO 94/29336, WO 96/14084, WO 96/16671, WO 97/23499, WO 97/39770, WO 97/45138, WO 98/16252, WO 15 99/27912, WO 99/27913, WO 00/12043 and WO 00/13671, the disclosures in which documents are hereby incorporated by reference. Otherwise, the preparation of suitable formulations may be achieved non-inventively by the skilled person using routine techniques.

20 The amounts of thrombin inhibitor or derivative in the formulation will depend on the severity of the condition, and on the patient, to be treated, as well as the compound(s) which is/are employed, but may be determined non-inventively by the skilled person.

25 Suitable doses of thrombin inhibitors and derivatives thereof in the therapeutic and/or prophylactic treatment of mammalian, especially human, patients may be determined routinely by the medical practitioner or other skilled person, and include the respective doses discussed in the relevant

prior art documents that are mentioned hereinbefore, the relevant disclosures in which are hereby incorporated by reference.

In the case of melagatran, suitable doses of active compound, prodrugs and derivatives thereof, in the therapeutic and/or prophylactic treatment of mammalian, especially human, patients include those which give a mean plasma concentration of up to 5  $\mu\text{mol/L}$ , for example in the range 0.001 to 5  $\mu\text{mol/L}$  over the course of treatment of the relevant condition. Suitable doses may thus be in the range 0.1 mg once daily to 25 mg three times daily, and/or up to 100 mg infused parenterally over a 24 hour period, for melagatran, and in the range 0.1 mg once daily to 100 mg three times daily for prodrugs of melagatran (see also the specific doses mentioned hereinafter for the prodrug of melagatran, ximelagatran).

In the case of compounds of formulae I and Ia, suitable daily doses of compounds in the therapeutic treatment of humans are about 0.001-100 mg/kg body weight at peroral administration and 0.001-50 mg/kg body weight at parenteral administration.

In any event, the physician, or the skilled person, will be able to determine the actual dosage which will be most suitable for an individual patient, which is likely to vary with the severity of the condition that is to be treated, as well as the age, weight, sex and response of the particular patient to be treated. The above-mentioned dosages are exemplary of the average case; there can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

Low molecular weight thrombin inhibitors may be employed in the method described herein by way of co-administration along with other cholesterol-

lowering, or lipid-lowering/modifying, drugs/therapies that are mentioned hereinbefore, such as the statins (HMG-CoA reductase inhibitors) and particularly any one of those statins specifically mentioned hereinbefore, in combination therapy.

5

According to a further aspect of the invention, there is provided a combination product comprising:

(A) a low molecular weight thrombin inhibitor, or a pharmaceutically-acceptable derivative thereof; and

10 (B) another cholesterol-lowering, or lipid-lowering/modifying, therapeutic agent,

wherein each of components (A) and (B) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

15 Such combination products provide for the administration of low molecular weight thrombin inhibitor/derivative in conjunction with the other therapeutic agent, and may thus be presented either as separate formulations, wherein at least one of those formulations comprises thrombin inhibitor/derivative, and at least one comprises the other therapeutic agent,  
20 or may be presented (i.e. formulated) as a combined preparation (i.e. presented as a single formulation including thrombin inhibitor/derivative and the other therapeutic agent).

Thus, there is further provided:

25

(1) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically-acceptable derivative thereof; another cholesterol-lowering, or lipid lowering/modifying, therapeutic agent; and a pharmaceutically-acceptable adjuvant, diluent or carrier; and

- (2) a kit of parts comprising components:
- (a) a pharmaceutical formulation including a low molecular weight thrombin inhibitor, or a pharmaceutically-acceptable derivative thereof, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and
  - (b) a pharmaceutical formulation including another cholesterol-lowering, or lipid lowering/modifying, therapeutic agent in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier,
- 10 which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

The method described herein may have the advantage that, in cholesterol-lowering therapy, it may be more convenient for the physician and/or patient than, be more efficacious than, be less toxic than, have a broader range of activity than, be more potent than, produce fewer side effects than, or that it may have other useful pharmacological properties over, similar methods (treatments) known in the prior art for use in such therapy.

20 The invention is illustrated, but in no way limited, by the following examples, in which:

Figure 1 illustrates the difference in mean values (with 95% Confidence Intervals) of total cholesterol levels in serum as between patients receiving ximelagatran (36 mg bid) or warfarin (INR levels between 2 and 3) during 25 the course of a clinical trial over a 21 month period.

Figure 2 illustrates the difference in mean values (with 95% Confidence Intervals) of levels of total triglycerides in serum as between patients

receiving ximelagatran (36 mg bid) or warfarin (INR levels between 2 and 3) during the course of a clinical trial over a 21 month period.

Figure 3 illustrates the difference in mean values (with 95% Confidence Intervals) of LDL (i.e. LDL-cholesterol) levels in serum as between patients receiving ximelagatran (36 mg bid) or warfarin (INR levels between 2 and 3) during the course of a clinical trial over a 21 month period.

Figure 4 illustrates the difference in mean values (with 95% Confidence Intervals) of HDL (i.e. HDL-cholesterol) levels in serum as between patients receiving ximelagatran (36 mg bid) or warfarin (INR levels between 2 and 3) during the course of a clinical trial over a 21 month period.

#### Example 1

#### 15 Lipid Measurements in Patients Undergoing Thrombin Inhibition Therapy in a Clinical Trial

A large-scale Phase III clinical trial was set up to establish the efficacy of the study compound  $\text{EtO}_2\text{C}-\text{CH}_2-(R)\text{Cgl}-(S)\text{Aze-Pab-OH}$  (ximelagatran; see Example 17 of international patent application WO 97/23499) in the prevention of stroke in patients with non-valvular atrial fibrillation, as compared to the current frontline treatment for this indication, warfarin.

Ximelagatran is a prodrug of the low molecular weight thrombin inhibitor, melagatran (see Example 1 of international patent application WO 94/29226).

The clinical trial protocol was similar to that described in international patent application WO 02/36157, with the following major differences:

- (a) the study objective was to show that the efficacy of ximelagatran is non-inferior to that of dose-adjusted warfarin, aiming for an INR 2.0-3.0 (with INR measurements taken at least every  $28 \pm 3$  days), in the prevention of all strokes (fatal and non-fatal) and systemic embolic events in patients with chronic non-valvular atrial fibrillation;
- (b) the dosage of ximelagatran was fixed at 36 mg bid;
- (c) in the exclusion criteria, subjects who had experienced stroke within the previous 30 days or transient ischaemic attack within the previous 3 days were excluded (as opposed to 2 years in the study described in WO 02/36157);
- (d) the duration of treatment was long term (between 12 and 26 months); and
- (e) the total number of patients in the trial was 3407 (as opposed to 220 in the study described in WO 02/36157). The study was a multicentre, multinational, IVRS-randomised, open-label, parallel-group study carried out across approximately 300 centres in approximately 25 countries.

Over the first 21 months of the trial, blood samples were taken from all patients in a standard manner at the following intervals: beforehand, and at, 1, 1.5, 2, 3, 4, 5, 6, 8, 10, 12, 15, 18, and 21, months. These samples were routinely analysed for total cholesterol, LDL (i.e. LDL-cholesterol), HDL (i.e. HDL-cholesterol), and total triglycerides, content using standard techniques for the detection of these lipids in serum.

A comparison was made between patients on ximelagatran ( $n = 1704$ ) and warfarin ( $n = 1703$ ). When the raw data were analysed, an unexpected statistically-significant difference in favour of the ximelagatran group was observed. As of the second month of treatment, marked mean differences


21

were observed for cholesterol, triglycerides and LDL serum concentrations (consistently significantly lower in the ximelagtran group over the entire 21 month period), and for the HDL serum concentration (consistently significantly higher in the ximelagtran group over the entire 21 month  
5 period), as illustrated in Figures 1 to 4, respectively.

These data clearly demonstrate the potential utility of melagatran and derivatives thereof (e.g. prodrugs, such as ximelagtran), as well as, potentially, low molecular weight thrombin inhibitors and  
10 derivatives/prodrugs thereof, in cholesterol-lowering therapy.

**Claims**

1. The use of melagatran or a pharmaceutically acceptable derivative thereof, for the manufacture of a medicament for use in cholesterol-lowering therapy.
- 5
2. The use of melagatran or a pharmaceutically acceptable derivative thereof, for the manufacture of a medicament for the treatment of hypercholesterolaemia, hyperlipoproteinaemia and/or hyper-triglyceridaemia.
- 10
3. The use as claimed in Claim 1 or Claim 2, wherein the therapy/treatment results in a decrease in serum levels of cholesterol, low-density lipoproteins, very low-density lipoproteins, triglycerides and/or apolipoprotein B; and/or an increase in serum levels of high-density lipoproteins and/or apolipoprotein A-I.
- 15
4. The use as claimed in any one of Claims 1 to 3, wherein the derivative of melagatran is a prodrug of melagatran.
5. The use as claimed in Claim 4, wherein the prodrug is of the formula
- $$R^1O_2C-CH_2-(R)Cgl-(S)Aze-Pab-OH,$$
- 20 wherein  $R^1$  represents linear or branched  $C_{1-6}$  alkyl and the OH group replaces one of the amidino hydrogens in Pab.
6. The use as claimed in Claim 5, wherein  $R^1$  represents methyl, ethyl or propyl.
- 25
7. The use as claimed in Claim 6, wherein  $R^1$  represents ethyl.
8. The use of a thrombin inhibitor of formula I,



wherein

$R^a$  represents -OH or -CH<sub>2</sub>OH;


5  $R^1$  represents at least one optional halo substituent;

$R^2$  represents one or two C<sub>1-3</sub> alkoxy substituents, the alkyl parts of which substituents are themselves substituted with one or more fluoro substituents;


Y represents -CH<sub>2</sub>- or -(CH<sub>2</sub>)<sub>2</sub>-; and

$R^3$  represents a structural fragment of formula I(i) or I(ii):

10



I(i)



I(ii)

wherein

$R^4$  represents H or one or more fluoro substituents; and

15 one or two of X<sub>1</sub>, X<sub>2</sub>, X<sub>3</sub> and X<sub>4</sub> represent -N- and the others represent -CH-, for the manufacture of a medicament for use in cholesterol-lowering therapy.

9. The use of a thrombin inhibitor of formula I, as defined in Claim 8, or a pharmaceutically acceptable derivative thereof, for the manufacture of a medicament for the treatment of hypercholesterolaemia, hyperlipoproteinaemia and/or  
20 hypertriglyceridaemia.

10. The use as claimed in Claim 8 or Claim 9, wherein the therapy/treatment results in a decrease in serum levels of cholesterol, low-density lipoproteins, very low-density lipoproteins, triglycerides and/or apolipoprotein B; and/or an increase in serum levels of high-density lipoproteins and/or apolipoprotein A-I.

5

11. The use as claimed in Claim 10, wherein the thrombin inhibitor is:

Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab;

Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF); or


Ph(3-Cl)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab.

10


12. The use as claimed in Claim 10 or Claim 11, wherein the derivative of the thrombin inhibitor is a prodrug of that inhibitor.

13. The use as claimed in Claim 12, wherein the prodrug is of formula Ia,

15



wherein R<sup>3a</sup> represents a structural fragment of formula I(iii) or I(iv):



20

wherein R<sup>5</sup> represents OR<sup>6</sup> or C(O)OR<sup>7</sup>;

$R^6$  represents H,  $C_{1-10}$  alkyl,  $C_{1-3}$  alkylaryl or  $C_{1-3}$  alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also optionally substituted by one or more halo substituents);

$R^7$  represents  $C_{1-10}$  alkyl (which latter group is optionally interrupted by one or more oxygen atoms), or  $C_{1-3}$  alkylaryl or  $C_{1-3}$  alkyloxyaryl (the alkyl parts of which latter two groups are optionally interrupted by one or more oxygen atoms, and the aryl parts of which latter two groups are optionally substituted by one or more substituents selected from halo, phenyl, methyl or methoxy, which latter three groups are also optionally substituted by one or more halo substituents); and

$R^a$ ,  $R^1$ ,  $R^2$ , Y,  $R^4$ ,  $X_1$ ,  $X_2$ ,  $X_3$  and  $X_4$  are as defined in Claim 8.

14. The use as claimed in Claim 13, wherein the prodrug is:

15  $\text{Ph}(3\text{-Cl})(5\text{-OCHF}_2)\text{-}(R)\text{CH}(\text{OH})\text{C}(\text{O})\text{-}(S)\text{Aze-Pab}(\text{OMe})$ ;  
 $\text{Ph}(3\text{-Cl})(5\text{-OCHF}_2)\text{-}(R)\text{CH}(\text{OH})\text{C}(\text{O})\text{-}(S)\text{Aze-Pab}(2,6\text{-diF})(\text{OMe})$ ; or  
 $\text{Ph}(3\text{-Cl})(5\text{-OCH}_2\text{CH}_2\text{F})\text{-}(R)\text{CH}(\text{OH})\text{C}(\text{O})\text{-}(S)\text{Aze-Pab}(\text{OMe})$ .

15. A substance or composition for use in a method of cholesterol-lowering  
 20 therapy, said substance or composition comprising melagatran or a pharmaceutically acceptable derivative thereof, and said method comprising the administration of said substance or composition to a patient in need of such therapy.


16. A substance or composition for use in a method of treatment of  
 25 hypercholesterolaemia, hyperlipoproteinaemia and/or hypertriglyceridaemia, said substance or composition comprising melagatran or a pharmaceutically acceptable derivative thereof, and said method comprising the administration of said substance or composition to a patient in need of such treatment.

30 17. A substance or composition for use in a method of therapy/treatment as claimed in Claim 15 or Claim 16 wherein the therapy/treatment results in a decrease

in serum levels of cholesterol, low-density lipoproteins, very low-density lipoproteins, triglycerides and/or apolipoprotein B; and/or an increase in serum levels of high-density lipoproteins and/or apolipoprotein A-I.

5 18. A substance or composition for use in a method of therapy/treatment as claimed in any one of Claims 15 to 17, wherein the derivative of melagatran is a prodrug of melagatran.

19. A substance or composition for use in a method of therapy/treatment as  
10 claimed in Claim 18, wherein the prodrug is of the formula



wherein  $R^1$  represents linear or branched  $C_{1-6}$  alkyl and the OH group replaces one of the amidino hydrogens in Pab.

15 20. A substance or composition for use in a method of therapy/treatment as claimed in Claim 19, wherein  $R^1$  represents methyl, ethyl or propyl.

21. A substance or composition for use in a method of therapy/treatment as  
20 claimed in Claim 20, wherein  $R^1$  represents ethyl.

22. A substance or composition for use in a method of cholesterol-lowering  
therapy said substance or composition comprising a thrombin inhibitor of formula I,  
as defined in Claim 8, and said method comprising the administration of said  
substance or composition to a patient in need of said therapy.

25 23. A substance or composition for use in a method of treatment of  
hypercholesterolaemia, hyperlipoproteinaemia and/or hypertriglyceridaemia, said  
substance or composition comprising a thrombin inhibitor of formula I, as defined in  
Claim 8, or a pharmaceutically acceptable derivative thereof, and said treatment  
30 comprising the administration of said substance or composition to a patient in need of  
said treatment.

24. A substance or composition for use in a method of therapy/treatment as claimed in Claim 22 or Claim 23, wherein the therapy/treatment results in a decrease in serum levels of cholesterol, low-density lipoproteins, very low-density lipoproteins, triglycerides and/or apolipoprotein B; and/or an increase in serum levels of high-  
5 density lipoproteins and/or apolipoprotein A-I.

25. A substance or composition for use in a method of therapy/treatment as claimed in Claim 24, wherein the thrombin inhibitor is:

Ph(3-C1)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab;

10 Ph(3-C1)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF); or

Ph(3-C1)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab.

26. A substance or composition for use in a method of therapy/treatment as claimed in Claim 24 or Claim 25, wherein the derivative of the thrombin inhibitor is a  
15 prodrug of that inhibitor.

27. A substance or composition for use in a method of therapy/treatment or prevention as claimed in Claim 26, wherein the prodrug is of formula Ia as defined in  
20 Claim 13.

28. A substance or composition for use in a method of therapy/treatment as claimed in Claim 27, wherein the prodrug is:

Ph(3-C1)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe);

Ph(3-C1)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe); or

25 Ph(3-C1)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe).

29. A combination product comprising:

(A) a low molecular weight thrombin inhibitor, or a pharmaceutically-acceptable derivative thereof; and

30 (B) another cholesterol-lowering, or lipid-lowering/modifying, therapeutic agent,

wherein each of components (A) and (B) is formulated in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier.

5 30. A combination product as claimed in Claim 29, which comprises a pharmaceutical formulation including the thrombin inhibitor or derivative, the other therapeutic agent and a pharmaceutically-acceptable adjuvant, diluent or carrier.

31. A combination product as claimed in Claim 29, which comprises a kit of parts comprising components:

10 (a) a pharmaceutical formulation including the thrombin inhibitor, or derivative, in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier; and

(b) a pharmaceutical formulation including the other therapeutic agent in admixture with a pharmaceutically-acceptable adjuvant, diluent or carrier,


15 which components (a) and (b) are each provided in a form that is suitable for administration in conjunction with the other.

32. A combination product as claimed in any one of Claims 29 to 31, wherein the thrombin inhibitor is melagatran.

20

33. A combination product as claimed in Claim 32, wherein the derivative of melagatran is a prodrug of melagatran.

25 34. A combination product as claimed in Claim 33, wherein the prodrug is of the formula




wherein  $R^1$  represents linear or branched  $C_{1-6}$  alkyl and the OH group replaces one of the amidino hydrogens in Pab.

30 35. A combination product as claimed in Claim 34, wherein  $R^1$  represents methyl, ethyl or propyl.


36. A combination product as claimed in Claim 35, wherein R<sup>1</sup> represents ethyl.
37. A combination product as claimed in any one of Claims 29 to 31, wherein the thrombin inhibitor is a compound of formula I as defined in Claim 8.
- 5
38. A combination product as claimed in Claim 37, wherein the thrombin inhibitor is:
- Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab;  
Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF); or  
10 Ph(3-Cl)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab.
39. A combination product as claimed in Claim 37 or Claim 38, wherein the derivative of the thrombin inhibitor is a prodrug of that inhibitor.
- 15 40. A combination product as claimed in Claim 39, wherein the prodrug is a compound of formula Ia as defined in Claim 13.
41. A combination product as claimed in Claim 40, wherein the prodrug is:
- Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe);  
20 Ph(3-Cl)(5-OCHF<sub>2</sub>)-(R)CH(OH)C(O)-(S)Aze-Pab(2,6-diF)(OMe); or  
Ph(3-Cl)(5-OCH<sub>2</sub>CH<sub>2</sub>F)-(R)CH(OH)C(O)-(S)Aze-Pab(OMe).
42. A combination product as claimed in any one of Claims 29 to 41, wherein the other therapeutic agent is a statin.
- 25
43. A combination product as claimed in Claim 42 wherein the statin is lovastatin, pravastatin, fluvastatin, simvastatin, atorvastatin, pitavastatin or rosuvastatin.
44. Use as claimed in Claim 1 or Claim 2 or Claim 8 or Claim 9, substantially as  
30 herein described with reference to and as illustrated in any of the examples and accompanying drawings.

45. A substance or composition for use in a method of treatment as claimed in Claim 15 or Claim 16 or Claim 22 or Claim 23, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.
- 5 46. A product as claimed in Claim 29, substantially as herein described with reference to and as illustrated in any of the examples and accompanying drawings.




Mean at baseline: ximelagatran 5.27, warfarin 5.32

FIG. 1




Mean at baseline: ximelagatran 1.86, warfarin 1.95

FIG. 2



Mean at baseline: ximelagatran 3.14, warfarin 3.18

FIG. 3



Mean at baseline: ximelagatran 1.31, warfarin 1.28

FIG. 4