



(86) Date de dépôt PCT/PCT Filing Date: 2010/08/09
(87) Date publication PCT/PCT Publication Date: 2011/02/17
(45) Date de délivrance/Issue Date: 2019/04/09
(85) Entrée phase nationale/National Entry: 2012/02/10
(86) N° demande PCT/PCT Application No.: US 2010/002194
(87) N° publication PCT/PCT Publication No.: 2011/019375
(30) Priorité/Priority: 2009/08/10 (US61/273,920)

(51) Cl.Int./Int.Cl. *A61K 31/4704* (2006.01),
A61P 25/00 (2006.01), *A61P 25/22* (2006.01),
A61P 25/24 (2006.01), *C07D 215/56* (2006.01)
(72) Inventeur/Inventor:
HAYARDENY, LIAT, IL
(73) Propriétaire/Owner:
ACTIVE BIOTECH AB, SE
(74) Agent: AITKEN KLEE LLP

(54) Titre : TRAITEMENT DE TROUBLES LIES AU FNDC AU MOYEN DE LAQUINIMOD
(54) Title: TREATMENT OF BDNF-RELATED DISORDERS USING LAQUINIMOD

(57) **Abrégé/Abstract:**

A method of increasing brain-derived neurotrophic factor (BDNF) serum level in a human subject comprising periodically administering to the subject an amount of laquimmod or pharmaceutically acceptable salt thereof The method can further comprise periodically administering to the subject an amount of a second BDNF-increasing agent A method for treating a human subject suffering from a BDNF-related disease comprising periodically administering laquimmod or a pharmaceutically acceptable salt thereof in an amount effective to treat the human subject Use of laquimmod in the manufacture of a medicament for increasing BDNF serum level in a human subject A pharmaceutical preparation comprising an amount of laquimmod and an amount of a second BDNF-increasing agent effective for use in increasing BDNF serum level in a human subject.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
17 February 2011 (17.02.2011)

(10) International Publication Number
WO 2011/019375 A1

(51) International Patent Classification:
C07D 401/00 (2006.01)

(21) International Application Number:
PCT/US2010/002194

(22) International Filing Date:
9 August 2010 (09.08.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/273,920 10 August 2009 (10.08.2009) US

(71) Applicant (for all designated States except MT, US):
TEVA PHARMACEUTICAL INDUSTRIES LTD.
[IL/IL]; 5 Basel Street, P.O. Box 3190, Petach-Tikva
49131 (IL).

(71) Applicant (for BB only): **TEVA PHARMACEUTI-
CALS USA, INC.** [US/US]; 1090 Horsham Road, North
Wales, PA 19454 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **HAYARDENY, Liat**
[IL/IL]; 37 Yehuda Hanasie St., Tel Aviv 69391 (IL).

(74) Agent: **WHITE, John, P.**; Cooper & Dunham Llp, 30
Rockefeller Plaza, New York, NY 10112 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG,
ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: TREATMENT OF BDNF-RELATED DISORDERS USING LAQUINIMOD

(57) Abstract: A method of increasing brain-derived neurotrophic factor (BDNF) serum level in a human subject comprising periodically administering to the subject an amount of laquimmod or pharmaceutically acceptable salt thereof The method can further comprise periodically administering to the subject an amount of a second BDNF-increasing agent A method for treating a human subject suffering from a BDNF-related disease comprising periodically administering laquimmod or a pharmaceutically acceptable salt thereof in an amount effective to treat the human subject Use of laquimmod in the manufacture of a medicament for increasing BDNF serum level in a human subject A pharmaceutical preparation comprising an amount of laquimmod and an amount of a second BDNF-increasing agent effective for use in increasing BDNF serum level in a human subject.



WO 2011/019375 A1

TREATMENT OF BDNF-RELATED DISORDERS USING LAQUINIMOD

5

10 Background

Neurotrophic factors have a profound influence on developmental events such as naturally occurring cell death, differentiation and process outgrowth (Snider, 1989). The brain-derived neurotrophic factor (BDNF) is a neurotrophic factor which belongs to the neurotrophin family of growth factors. BDNF acts on certain neurons of the central nervous system (CNS) and the peripheral nervous system (PNS), helping to support the survival of existing neurons and encourage the growth and differentiation of new neurons and synapses (Acheson, 1995; Huang, 2001). Increasing BDNF has been associated with the treatment of a number of disorders including Parkinson's disease, Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), depressive disorders, retinitis pigmentosa, erectile dysfunction, memory disorders, Rett syndrome, Alzheimer's Disease, bipolar disorder and acute mania.

Parkinson's disease is a chronic and progressive degenerative disease of the brain that impairs motor control, speech, and other functions. One of the most striking features of Parkinson's disease is that it primarily affects a restricted neuronal population in the brain. Although other neurons are also affected, the dopaminergic neurons of the substantia

nigra pars compacta are the most vulnerable to the disease process (Chesselet, 2003). BDNF has potent effects on survival and morphology of mesencephalic dopaminergic neurons, increasing their survival, and thus its loss could contribute
5 to death of these cells in Parkinson's disease (PD) (Hyman, 1991; Howell, 2000).

Huntington's disease (HD) is a neurodegenerative disorder characterized by motor, cognitive, and psychiatric symptoms and by a progressive degeneration of neurons in basal ganglia
10 in brain cortex. Patients suffering from HD have significantly lower BDNF levels in serum compared to healthy controls (Ciammola, 2007).

Amyotrophic lateral sclerosis (ALS) is a chronic and debilitating neurodegenerative disease which involves
15 degeneration of cortical, bulbar and medullar motor neurons. Riluzole (2-amino-6-[trifluoromethoxy]benzothiazole) is an antagonist of glutamatergic neurotransmission that prolongs survival in ALS (Riviere, 1998). Riluzole has also been shown to significantly increase BDNF levels in the rat brain,
20 thereby promoting precursor proliferation (Kato-Semba, 2002).

Depression is another indication in which BDNF has been shown to have an effect. In a meta-analysis which encompasses many studies, depressed patients were shown to have lower BDNF levels than healthy control subjects, and anti-depressant
25 therapy has been shown to increase BDNF levels in depressed patients after treatment (Sen, 2008).

Retinitis pigmentosa is a disease associated with retinal photoreceptor cell loss. It has been shown that BDNF culturing of retina explants from *rd* mice, who suffer from a mutation in
30 the same gene that has been found mutated in human autosomal recessive retinitis pigmentosa, showed an increase in number of photoreceptor nuclei in the outer nuclear layer (Caffe Romeo, 2001). This suggests that increasing BDNF levels in humans suffering from retinitis pigmentosa may slow the
35 progression of the disease.

Erectile dysfunction is another disease which has been shown to be associated with BDNF. In U.S. Patent No. 7,485,311, example 3, it was shown that treatment of rats in the bilateral cavernous nerve freezing model using intracavernous injection of AAV-BDNF improves maximum intracavernous pressure in response to bilateral cavernous nerve electrostimulation.

BDNF has also been associated with learning and memory. In a transgenic mouse model of BDNF knockout mice, impairment of BDNF production has been shown to cause impairments in learning and memory in the adult stage and especially during early development (Monteggia, 2007). In a rat model, it was shown that endogenous BDNF in the hippocampus is involved in memory formation. Whereas infusion of function-blocking anti-BDNF antibody was shown to impair short term and long term memory in a model of fear-motivated learning, infusion of recombinant human BDNF facilitated long-term memory retention (Alonsa, 2005). Increasing BDNF levels in human patients in need of learning or memory improvement may also improve short and long term memory.

Rett Syndrome (RTT) is an X-linked neurodevelopmental disorder and the leading cause of severe mental retardation in females, affecting 1:10,000-15,000 births worldwide (Amaral, 2007). The disease is associated with mutations in the gene MeCP2. One of the targets of MeCP2 is the BDNF gene. This suggests that deregulation of BDNF expression in Rett Syndrome may be the cause of structural anomalies observed in patients, especially the reduced dendritic branching and loss of dendritic spines (Amaral, 2007). This suggests that increasing BDNF levels in Rett syndrome may be a viable therapy in treating the disease.

In bipolar disorder patients suffering from acute mania, levels of BDNF were shown to vary depending upon whether the patient was treated or untreated. Before treatment with known mania treatments, BDNF levels were found to be lower than in healthy controls, but upon treatment, the difference between BDNF levels in serum of treated patients and controls was no

longer significant (Tramontina, 2009). It may be possible to treat acute mania by increasing BDNF levels in patients in need thereof.

Several agents have been identified to increase BDNF levels including riluzole and antidepressants such as fluoxetine (Prozac®) (Kato-Semba, 2002; Molteni, 2006).

Summary of the Invention

Disclosed herein is another agent, i.e., Laquinimod, which has been shown to increase BDNF in humans. Laquinimod is a novel synthetic compound with high oral bioavailability, which has
5 been suggested as an oral formulation for Relapsing Remitting Multiple Sclerosis (RRMS). The relationship between laquinimod and BDNF has not been reported. Laquinimod is a compound whose chemical name is N-ethyl-N-phenyl-1,2-dihydro-4-hydroxy-5-chloro-1-methyl-2-oxoquinoline-3-carboxamide, and its Chemical
10 Registry number is 248281-82-7. The process of synthesis of laquinimod and the preparation of its sodium salt are disclosed in U.S. Patent No. 6,077,851. Additional process of synthesis of laquinimod is disclosed in U.S. Patent No. 6,875,869 and in U.S. Patent Application Publication No. 2007-
15 0088050. Pharmaceutical compositions comprising laquinimod sodium are disclosed in PCT International Application Publication No. WO 2005/074899 as well as in U.S. Patent Application Publication Nos. 2007-0293537 and 2009-0162432.

This application provides for a method of increasing brain-
20 derived neurotrophic factor (BDNF) serum level in a human subject comprising periodically administering to the subject an amount of laquinimod or pharmaceutically acceptable salt thereof effective to increase BDNF serum level in the human subject. The method can further comprise periodically
25 administering to the subject an amount of a second BDNF-increasing agent.

This application also provides for a method for treating a human subject suffering from a BDNF-related disease selected
30 from the group consisting of Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, depressive disorders, anxiety disorders, retinitis pigmentosa, erectile dysfunction, memory disorders, Rett syndrome, Alzheimer's disease, bipolar disorder and acute mania comprising periodically administering
35 laquinimod or a pharmaceutically acceptable salt thereof in an amount effective to treat the human subject.

This application also provides for use of laquinimod in the manufacture of a medicament for increasing BDNF serum level in a human subject.

5 This application also provides for a pharmaceutical composition comprising an amount of laquinimod effective for use in increasing BDNF serum level in a human subject.

10 This application also provides for a pharmaceutical preparation comprising an amount of laquinimod and an amount of a second BDNF-increasing agent effective for use in increasing BDNF serum level in a human subject.

Brief Description of the Figures

Figure 1: Shows one-way analysis of BDNF comparing the placebo group and 0.6 mg laquinimod group at baseline.

Figure 2: Shows one-way analysis of BDNF comparing the placebo group and 0.6 mg laquinimod group at week 12 (V3).

Figure 3: Shows one-way analysis of BDNF comparing the placebo group and 0.6 mg laquinimod group at week 36 (V9).

Figure 4: Shows Forced Swim Test results conducted in Example 2.1. (N = 5/group) Laquinimod (1-25 mg/kg/dx3d po) showed antidepressant activity with the 1 mg/kg p.o. dose being the most active.

4A: Effect of Laquinimod (1;5;10;25 mg/kg/dx4, p.o., -90) and Fluoxetine (10 mg/kg/dx4, -90 min, p.o.) in the FST on Distance moved [cm] on Balb C mice [4 min]

4B: Effect of Laquinimod (1;5;10;25 mg/kg/dx4, p.o., -90) and Fluoxetine (10 mg/kg/dx4, -90 min, p.o.) in the FST on Velocity (cm/s) on Balb C mice [4min]

4C: Effect of Laquinimod (1;5;10;25 mg/kg/dx4, p.o., -90) and Fluoxetine (10 mg/kg/dx4, -90 min, p.o.) in the FST on Movement (Moving duration) on Balb C mice [4min]

4D: Effect of Laquinimod (1;5;10;25 mg/kg/dx4, p.o., -90) and Fluoxetine (10 mg/kg/dx4, -90 min, p.o.) in the FST on Immobility total duration [10%] on Balb C mice [4min]

Figure 5: Shows Forced Swim Test results conducted in Example 2.2. (*p<0.01 vs. cont) Laquinimod (0.5-25 mg/kg/dx3d po) showed antidepressant activity with doses at 5 and 25 mg/kg inducing significant antidepressant activity.

5A: Effect of Laquinimod (0.5;1;5;25 mg/kg, p.o., -90) and Fluoxetine (10 mg/kg,-90 min, p.o.) in the FST on Distance moved [cm] on Balb C mice [4 min]

5B: Effect of Laquinimod (0.5;1;5;25 mg/kg, p.o., -90) and Fluoxetine (10 mg/kg,-90 min, p.o.) in the FST on velocity Balb C mice [4 min]

5C: Effect of Laquinimod (0.5;1;5;25 mg/kg, p.o., -90) and Fluoxetine (10 mg/kg,-90 min, p.o.) in the FST on movement (total duration) on Balb C mice [4 min].

5D: Effect of Laquinimod (0.5;1;5;25 mg/kg, p.o., -90) and Fluoxetine (10 mg/kg,-90 min, p.o.) in the FST on Immobility total duration [10%] on Balb C mice [4 min]

Figure 6: Shows Open Field Test - motility parameter results conducted in Example 3.1. (N = 5/group) Laquinimod (1-25 mg/kg/dx3d po) did not modify general mobility parameters in open field in Balb/c mice.

6A: Effect of Laquinimod (1;5;10;25 mg/kg, po; -90min) and Fluoxetine (10 mg/kg, po; -90min) and combination in the Open field test on **Distance moved [cm]** on BALB/c mice [20 min]

6B: Effect of Laquinimod (1;5;10;25 mg/kg, po; -90min) and Fluoxetine (10 mg/kg, po; -90min) and combination in the Open field test on **Velocity mean [cm/s]** on BALB/c mice [20 min]

6C: Effect of Laquinimod (1;5;10;25 mg/kg, po; -90min) and Fluoxetine (10 mg/kg, po; -90min) and combination in the Open field test on **Immobility total duration** on BALB/c mice [20 min]

6D: Effect of Laquinimod (1;5;10;25 mg/kg, po; -90min) and Fluoxetine (10 mg/kg, po; -90min) and combination in the Open field test on **Strong mobility total duration [s]** on BALB/c mice [20 min]

Figure 7: Shows Open Field Test - anxiety parameter results (in zone 2) conducted in Example 3.1. (N = 5/group) Laquinimod (1-25 mg/kg/dx3d po) showed tendency toward anxiolytic effect with animals being more active in the center (zone 2).

Figure 8: Shows Open Field Test - anxiety parameter results (in zone 2 + 3) conducted in Example 3.1. (N = 5/group) Laquinimod (1-25/kg/dx3d po) showed tendency toward anxiolytic effect with animals being more active in the center (zones 2 + 3). 1 mg/kg was shown to be most potent dose.

8C: Effect of Laquinimod (1;5;10;25 mg/kg, po; -90min) and Fluoxetine (10 mg/kg, po; -90min) and combination in the Open field test on **Rearing Frequency** on BALB/c mice [20 min]

Figure 9: Shows Open Field Test - motility parameter results conducted in Example 3.2. (N= 5/group; *p < 0.05 vs. cont) Laquinimod (0.5-25 mg/kg/dx3d po) induced a slight increase in motility (0.5-5 mg/kg) in the open field in Balb/c mice.

Figure 10: Shows Open Field Test - anxiety parameter results (in zones 2 + 3) conducted in Example 3.2. Laquinimod (0.5-25mg/kg/dx3d po) shows significant anxiolytic effect with animals showing more

activity in the center (zones 2 + 3). Both fluoxetine and laquinimod (0.5-5 mg/kg) showed significant anxiolytic effect vs. the control. The most potent dose was laquinimod at 5 mg/kg.

Figure 11: Shows Elevated Plus Maze model results conducted in Example 4.1. Laquinimod (5 and 25 mg/kg/dx3d po) showed significant anxiolytic effect. Laquinimod showed a dose dependent increase in activity on the open arm. The dose at 25 mg/kg p.o. was the most active.

Figure 12: Shows Elevated Plus Maze model results conducted in Example 4.2. (Open arm parameters in Balb/c mice - 5 min; N=5/group) Laquinimod (5-25 mg/kg/dx3d po) showed significant anxiolytic effect.

Detailed Description of the Invention

This application provides for a method of increasing brain-derived neurotrophic factor (BDNF) serum level in a human subject comprising periodically administering to the subject an amount of laquinimod or pharmaceutically acceptable salt thereof effective to increase BDNF serum level in the human subject.

In one embodiment, the amount of laquinimod or pharmaceutically acceptable salt thereof is administered to the human subject once daily. In another embodiment, the periodic administration continues for at least 3 days.

In one embodiment, the amount of laquinimod administered is 0.1 mg/day-40.0 mg/day. In another embodiment, the amount of laquinimod administered is 0.6 mg/day. In another embodiment, the amount of laquinimod is administered orally.

In one embodiment, the subject is suffering from a BDNF-related disease. In another embodiment, the BDNF-related disease is Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, Alzheimer's disease, bipolar disorder or acute mania.

In an embodiment, the depressive disorder is depression, depression in cancer patients, depression in Parkinson's disease patients, postmyocardia infarction depression, depression in patients with human immunodeficiency virus (HIV), subsyndromal symptomatic depression, depression in infertile women, pediatric depression, major depression, single episode depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, neuropathic pain, manic depressive illness including manic depressive illness with mixed episodes and manic depressive illness with

depressive episodes, seasonal affective disorder, bipolar depression BP I, bipolar depression BP II, or major depression with dysthymia.

5 In another embodiment, the anxiety disorder is generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, or childhood anxiety.

10 In one embodiment, the method further comprises periodically administering to the subject an amount of a second BDNF-increasing agent.

In one embodiment, the amount of the second BDNF-increasing agent when taken alone is not effective to increase BDNF serum level in the subject.

15 In one embodiment, the administration of the laquinimod substantially precedes the administration of the second BDNF-increasing agent. In another embodiment, the administration of the second BDNF-increasing agent substantially precedes the administration of the laquinimod.

20 In one embodiment, the effect on the subject is greater than the effect on a subject treated with the second BDNF-increasing agent alone. In another embodiment, the increase in BDNF serum level in the subject is greater than the increase in BDNF serum level in a subject treated with the second BDNF-increasing agent alone.

25 In one embodiment, the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.

30 In one embodiment, the diagnosis of the subject prior to administration excluded multiple sclerosis, insulin-dependent diabetes mellitus, systemic lupus erythematosus, rheumatoid arthritis, inflammatory bowel disease, psoriasis, asthma, atherosclerosis, stroke and Alzheimer's disease.

This application also provides for a method for treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, depressive disorders, anxiety disorders, retinitis pigmentosa, erectile dysfunction, memory disorders, Rett syndrome, Alzheimer's disease, bipolar disorder and acute mania comprising periodically administering laquinimod or a pharmaceutically acceptable salt thereof in an amount effective to treat the human subject.

10 This application also provides for use of laquinimod in the manufacture of a medicament for increasing BDNF serum level in a human subject.

This application also provides for a pharmaceutical composition comprising an amount of laquinimod effective for use in increasing BDNF serum level in a human subject.

This application also provides for a pharmaceutical preparation comprising an amount of laquinimod and an amount of a second BDNF-increasing agent effective for use in increasing BDNF serum level in a human subject.

20 For the foregoing embodiments, each embodiment disclosed herein is contemplated as being applicable to each of the other disclosed embodiment.

Terms

As used herein, and unless stated otherwise, each of the following terms shall have the definition set forth below.

As used herein, "administering to the subject" means the giving of, dispensing of, or application of medicines, drugs, or remedies to a subject to relieve or cure a pathological condition. Oral administration is one way of administering the instant compounds to the subject.

As used herein, "BDNF" means brain-derived neurotrophic factor, a neurotrophic factor belonging to the neurotrophin family of growth factors.

As used herein, a "BDNF-related disease" is a disease in which
5 a patient suffering from the disease has BDNF serum levels which are lower than those of a corresponding healthy individual and/or a disease in which the elevation of BDNF in a patient suffering from the disease can be associated with amelioration of the disease or of symptoms thereof.

10 As used herein, a "BDNF-increasing agent" is any agent which directly or indirectly elevates BDNF level in a subject. For example, a BDNF-increasing agent can be riluzole or an antidepressant such as fluoxetine. As used herein, the term "agent" includes any molecule, compound, protein, peptide,
15 polypeptide, nucleic acid, antibody, or drug or any combination thereof.

As used herein, an "amount" or "dose" of laquinimod as measured in milligrams refers to the milligrams of laquinimod acid present in a preparation, regardless of the form of the
20 preparation. For example, 0.6 mg of laquinimod means the amount of laquinimod acid in a preparation is 0.6 mg, regardless of the form of the preparation. Thus, when in the form of a salt, e.g. a laquinimod sodium salt, the weight of the salt form necessary to provide a dose of 0.6 mg laquinimod
25 would be greater than 0.6 mg due to the presence of the additional salt ion, but would be a molar equivalent amount.

As used herein, "effective" as in an amount effective to achieve an end means the quantity of a component that is sufficient to yield an indicated therapeutic response without
30 undue adverse side effects (such as toxicity, irritation, or allergic response) commensurate with a reasonable benefit/risk ratio when used in the manner of this disclosure. For example, an amount effective to treat a symptom of a disorder or disease without causing undue adverse side effects. The
35 specific effective amount will vary with such factors as the

particular condition being treated, the physical condition of the patient, the type of mammal being treated, the duration of the treatment, the nature of concurrent therapy (if any), and the specific formulations employed and the structure of the compounds or its derivatives.

A "salt" is salt of the instant compounds which have been modified by making acid or base salts of the compounds. The term "pharmaceutically acceptable salt" in this respect, refers to the relatively non-toxic, inorganic and organic acid or base addition salts of compounds of the present invention.

A pharmaceutically acceptable salt of laquinimod can be used. A pharmaceutically acceptable salt of laquinimod as used in this application includes lithium, sodium, potassium, magnesium, calcium, manganese, copper, zinc, aluminum and iron. Salt formulations of laquinimod and the process for preparing the same are described, e.g., in U.S. Patent Application Publication No. 2005-0192315 and PCT International Application Publication No. WO 2005/074899.

As used herein, to "treat" or "treating" encompasses, e.g., inducing inhibition, regression, or stasis of the disorder and/or disease. As used herein, "inhibition" of disease progression or disease complication in a subject means preventing or reducing the disease progression and/or disease complication in the subject.

As used herein, "pharmaceutically acceptable carrier" refers to a carrier or excipient that is suitable for use with humans and/or animals without undue adverse side effects (such as toxicity, irritation, and allergic response) commensurate with a reasonable benefit/risk ratio. It can be a pharmaceutically acceptable solvent, suspending agent or vehicle, for delivering the instant compounds to the subject.

A dosage unit as used herein may comprise a single compound or mixtures of compounds thereof. A dosage unit can be prepared

for oral dosage forms, such as tablets, capsules, pills, powders, and granules.

Laquinimod can be administered in admixture with suitable pharmaceutical diluents, extenders, excipients, or carriers (collectively referred to herein as a pharmaceutically acceptable carrier) suitably selected with respect to the intended form of administration and as consistent with conventional pharmaceutical practices. The unit can be in a form suitable for oral administration. Laquinimod can be administered alone but is generally mixed with a pharmaceutically acceptable carrier, and co-administered in the form of a tablet or capsule, liposome, or as an agglomerated powder. Examples of suitable solid carriers include lactose, sucrose, gelatin and agar. Capsule or tablets can be easily formulated and can be made easy to swallow or chew; other solid forms include granules, and bulk powders. Tablets may contain suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents flow-inducing agents, and melting agents.

Specific examples of the techniques, pharmaceutically acceptable carriers and excipients that may be used to formulate oral dosage forms of the present invention are described, e.g., in U.S. Patent Application Publication No. 2005/0192315, PCT International Application Publication Nos. WO 2005/074899, WO 2007/047863, and WO/2007/146248.

General techniques and compositions for making dosage forms useful in the present invention are described in the following references: 7 Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, Editors, 1979); Pharmaceutical Dosage Forms: Tablets (Lieberman et al., 1981); Ansel, Introduction to Pharmaceutical Dosage Forms 2nd Edition (1976); Remington's Pharmaceutical Sciences, 17th ed. (Mack Publishing Company, Easton, Pa., 1985); Advances in Pharmaceutical Sciences (David Ganderton, Trevor Jones, Eds., 1992); Advances in

Pharmaceutical Sciences Vol 7. (David Ganderton, Trevor Jones, James McGinity, Eds., 1995); Aqueous Polymeric Coatings for Pharmaceutical Dosage Forms (Drugs and the Pharmaceutical Sciences, Series 36 (James McGinity, Ed., 1989);
5 Pharmaceutical Particulate Carriers: Therapeutic Applications: Drugs and the Pharmaceutical Sciences, Vol 61 (Alain Rolland, Ed., 1993); Drug Delivery to the Gastrointestinal Tract (Ellis Horwood Books in the Biological Sciences. Series in Pharmaceutical Technology; J. G. Hardy, S. S. Davis, Clive G.
10 Wilson, Eds.); Modern Pharmaceutics Drugs and the Pharmaceutical Sciences, Vol. 40 (Gilbert S. Banker, Christopher T. Rhodes, Eds.).

15 Tablets may contain suitable binders, lubricants, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. For instance, for oral administration in the dosage unit form of a tablet or capsule, the active drug component can be combined with an
20 oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, gelatin, agar, starch, sucrose, glucose, methyl cellulose, dicalcium phosphate, calcium sulfate, mannitol, sorbitol, microcrystalline cellulose and the like. Suitable binders include starch, gelatin, natural sugars such
25 as glucose or beta-lactose, corn starch, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, povidone, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, sodium benzoate, sodium acetate,
30 sodium chloride, stearic acid, sodium stearyl fumarate, talc and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, croscarmellose sodium, sodium starch glycolate and the like.

It is understood that where a parameter range is provided, all
35 integers within that range, and tenths thereof, are also

provided by the invention. For example, "0.1 mg -40.0 mg" includes 0.1 mg, 0.2 mg, 0.3 mg, 0.4 mg, etc. up to 40.0 mg.

This invention will be better understood by reference to the Experimental Details which follow, but those skilled in the art will readily appreciate that the specific experiments detailed are only illustrative of the invention as described more fully in the claims which follow thereafter.

Experimental Details

EXAMPLE 1: Clinical Trial Showing Effect of Laquinimod on BDNF Levels in Human Subjects Afflicted With Relapsing Remitting Multiple Sclerosis

A study was initiated in relapsing remitting multiple sclerosis (RRMS) patients using laquinimod, 0.6mg daily, in the form of the sodium salt, in an oral, once-daily tablet formulation. The study was a multinational, multicenter, randomized, double-blind, parallel-group, placebo controlled study assessing the efficacy, tolerability and safety of two doses of laquinimod in subjects with RRMS. Eligible subjects were randomized into the following three groups:

1. 0.6 mg of laquinimod per os (p.o.) once daily;
2. 0.3 mg of laquinimod per os (p.o.) once daily; and
3. Matching placebo, per os (p.o.) once daily.

Subjects must meet the following inclusion criteria to be included in the study:

1. a confirmed MS diagnosis as defined by the McDonald criteria;
2. a RRMS disease course;

3. at least one documented relapse n the 12 months prior to screening;
4. at least one Gadolinium-enhanced lesion on their screening MRI scan;
- 5 5. ambulatory with a Kurtzke EDSS score of 1.0-5.0 (Converted);
6. between 18-50 years of age inclusive;
7. relapse-free and off corticosteroids or ACTH for at least 30 days prior to the MRI scan at screening;
- 10 8. relapse-free and off corticosteroids between screening and randomization;
9. women of child-bearing potential practiced a reliable method of birth control;
10. willing and able to comply with the protocol requirements for the duration of the study; and
- 15 11. able to give signed, written and informed consent prior to entering the study.

Subjects meeting any one of the following exclusion criteria were excluded from the study:

- 20 1. suffered from any form of progressive MS;
2. use of immunosuppressive or cytotoxic treatments within 6 months prior to the screening visit (including azathioprine, cyclophosphamide and methotrexate);
- 25 3. use of experimental drugs, and/or participation in drug clinical studies within the 6 months prior to screening;

4. previous treatment with immunomodulators (including IFN- β 1a and 1b, glatiramer acetate, laquinimod and IVIG) within the 2 months prior to screening;
5. use of potent inhibitors of CYP3A4, for example, oral ketoconazole and erythromycin within 2 weeks prior to baseline visit;
6. previous use of amiodarone;
7. use of fluoxetine one month prior to baseline visit;
8. use of substrates of CYP1A2 such as theophylline and warfarin within 2 weeks prior to screening;
9. previous treatment with cladribine within the last 2 years prior to screening visit;
10. subjects for whom potential immunosuppression would be contraindicated, for example, Hepatitis B/C or HIV.
11. previous total body irradiation or total lymphoid irradiation;
12. chronic corticosteroid treatment (30 or more consecutive days) within the 2 months prior to screening;
13. pregnancy or breastfeeding;
14. subjects with a clinically significant or unstable medical or surgical condition that would preclude safe and complete study participation as determined by medical history, physical exams, ECG, abnormal laboratory tests and chest X-ray.
15. inability to give informed consent, or to complete the study, or if the subject is considered by the

investigator to be, for any reason, an unsuitable candidate for the study;

16. a known history of sensitivity to Gd; or

17. inability to successfully undergo MRI scanning.

5 The study duration was 40 weeks and consisted of 2 periods: 4 weeks of screening period (week -4 [screening] to week 0 [baseline]) and 36 weeks of double blind treatment (week 0 [baseline] to week 36 [termination]).

10 Subjects were evaluated at study sites at weeks -4, 0, 4, 8, 12, 16, 20, 24, 28, 32 and 36. Samples for inflammatory markers were collected at all visits between 0 and 36 weeks.

15 BDNF concentration in serum was measured in patients in the placebo group (102 patients were in the group) and in the 0.6 mg laquinimod group (106 patients were in the group) at week 0 (V0), week 12 (v3) and week 36 (V9). The analysis of BDNF content in the plasma was performed using the sandwich ELISA method. The concentration of BDNF in serum is expressed in terms of pg/mL.

20 **Results**

The results in the tables below represent mean BDNF levels in both the laquinimod 0.6 mg/day group and the placebo group.

Baseline Values (at V0)

25 Table 1: Mean and Std Deviations

Level	Number	Mean	Std Dev	Std Mean	Err Lower 95%	Err Upper 95%
Placebo	95	14110.7	6432.61	659.97	12800	15421
Laquinimod 0.6 mg	96	13892.7	7267.89	741.78	12420	15365

Table 2: t Test

Laquinimod 0.6 mg-Placebo - Assuming unequal variances

Difference	-218.0	t-Ratio	-0.2196
Std Err Dif	922.9	DF	186.6984
Upper CL Dif	1740.7	Prob > t	0.8264
Lower CL Dif	-2176.7	Prob > t	0.5868
Confidence	0.95	Prob < t	0.4132

5 **Week 12 Values (at V3)**

Table 3: Mean and Std Deviations

Level	Number	Mean	Std Dev	Std Mean	Err Lower 95%	Err Upper 95%
Placebo	96	13073.9	5712.02	582.98	11916	14231
Laquinimod 0.6 mg	100	15519.6	7204.47	720.45	14090	16949

Table 4: t Test

Laquinimod 0.6 mg-Placebo - Assuming unequal variances

Difference	2445.70	t-Ratio	2.638935
Std Err Dif	926.77	DF	187.3752
Upper CL Dif	4273.95	Prob > t	0.0090*
Lower CL Dif	617.44	Prob > t	0.0045*
Confidence	0.95	Prob < t	0.9955

10

Week 36 Values (at V9)

Table 5: Mean and Std Deviations

Level	Number	Mean	Std Dev	Std Mean	Err Lower 95%	Err Upper 95%
Placebo	92	12783.6	6017.17	627.33	11538	14030
Laquinimod 0.6 mg	92	15335.8	6699.89	698.51	13948	16723

Table 6: t Test

Laquinimod 0.6 mg-Placebo - Assuming unequal variances

Difference	2552.21	t-Ratio	2.718403
Std Err Dif	938.86	DF	179.9376
Upper CL Dif	4404.81	Prob > t	0.0072*
Lower CL Dif	699.61	Prob > t	0.0036*
Confidence	0.95	Prob < t	0.9964

15

Starting from V3 there is a statistically significant elevation of BDNF (at Confidence Level of 95%) between the placebo group and the laquinimod 0.6 mg group. The difference was also statistically significant at V9.

5 Laquinimod is effective in experimental autoimmune encephalomyelitis (aEAE) model, and is currently being tested in MS patients (on phase III clinical trial) thus suggesting the possibility of its use in treatment of multiple sclerosis (MS). Laquinimod is also effective in
10 increasing BDNF levels in the patients.

The ability of laquinimod to increase BDNF levels is unexpected. A per se connection between treatment of MS and increase in BDNF levels has not been established. Not all agents which are effective in treating MS are
15 effective in elevating BDNF levels. A recent study showed that treatment with interferon- β 1a and immunoglobulins, even after 12 months, and even in MS patients benefitting from the treatment, did not show an increase in plasma BDNF levels (Sarchielli, 2007). The
20 finding that laquinimod elevates BDNF levels in patients after only 12 weeks of treatment is unexpected.

EXAMPLE 2: LAQUINIMOD SHOWS ANTIDEPRESSANT ACTIVITY IN MICE MODELS- THE FORCED SWIM TEST (FST)

Depression and anxiety disorders are burdensome
25 conditions with lifetime prevalence rates of approximately 7-20%. Animal models are indispensable tools in the search to identify new antidepressant drugs. Various paradigms have been developed and are instrumental in detecting the antidepressant-like potential of novel compounds in
30 preclinical settings (Cryan, 2002; Ganbarana, 2001).

The Forced Swim Test (FST) is one of the most widely used tools for screening antidepressant activity pre-

clinically. The acute test was first described by Porsolt et al (1977).

The test is based on the observation that rats and mice develop an immobile posture when placed in an inescapable cylinder of water. This behavior is considered to be a behavioral despair as opposed to active form of coping with stressful conditions. An antidepressant will reduce immobility and increase motivated behavior of the rodent to escape from the despaired conditions. This is evidenced by increase in time of swimming, distance moved, velocity and attempts to climb the walls (strong mobility).

FST is considered a screening tool with high reliability and predictive validity. The test can be performed in mice and rats.

In example 2.1 and 2.2 described below, the FST was conducted in male mice (Balb/c Harlan IL) after 3 days of drug administration, 90 minutes after the last drug administration. Round glass cylinders 18 cm diameter and 20 cm depth were used. Water temperature was 24-28°C. Motivated behavior was defined by immobility, swimming and strong mobility.

Immobility in the animals was defined by activity lower than 10% movement of the center of gravity of the animal as determined by the Noldus system. Swimming was defined by the distance and the velocity of the animal, and climbing was related to strong mobility (movement of center of gravity more than 30%). Animals were released in the cylinder for 6 min and scoring was performed in the last 4 min after 2 min of adaptation.

All results were analyzed by the Noldus (Holland) system including a camera and software for animal behavior analyses.

Example 2.1

5 Balb/c mice were divided to 6 groups (5/group) and administered daily for 3 days laquinimod (1, 5, 10 and 25 mg/kg per os in the form of laquinimod sodium solution via gavage), fluoxetine (positive control 10mg/kg per os) or vehicle. On the third day, 90 min following
10 administration, mice were exposed to the forced swim test. Behavior was video recorded and analyzed using Ethovision software (Noldus Holland). The result is shown in **Fig. 4**.

Conclusion:

The results show that laquinimod showed a significant
15 antidepressant activity as expressed by the increased mobility and reduced immobility at 1mg/kg, and a trend to significance at higher doses (Laquinimod at 25mg/kg). In this experiment, the positive control fluoxetine did not show significant effect, possibly due to non optimal
20 conditions of time and dose.

Example 2.2

Balb/c mice were divided to 6 groups (5 mice/group) and administered laquinimod daily for 3 days with (0.5, 1, 5 and 25mg/kg p.o. in the form of laquinimod sodium
25 solution via gavage), fluoxetine (positive control 10mg/kg po) or vehicle. On the third day, 90 minutes following administration, mice were exposed to the forced swim test. Behavior was video recorded and analyzed using Ethovision software (Noldus Holland). The result is shown
30 in **Fig. 5**.

Conclusion:

The results show that laquinimod showed a significant antidepressant activity as expressed by the increased mobility and reduced immobility at 5 and 25 mg/kg. The positive control fluoxetine showed a trend to the same activity. This is possibly due to non optimal conditions of time and dose.

EXAMPLE 3: LAQUINIMOD SHOWS ANXIOLYTIC ACTIVITY IN MICE MODELS - THE OPEN FIELD TEST

10 Anxiety disorders are blanket terms covering several different forms of abnormal and pathological fear and anxiety. Current psychiatric diagnostic criteria recognize a wide variety of anxiety disorders. Recent surveys have found that as many as 18% of Americans may
15 be affected by one or more of them (Kessler et al). The disorders are divided to several classes including: Generalized Anxiety, Panic disorders. Phobias, Post Traumatic Stress Disorders (PTSD), Obsessive Compulsive Disorder (OCD), Separation Anxiety and Childhood Anxiety.

20 Chronically administered antidepressant drugs, particularly selective serotonin (5-HT) reuptake inhibitors (SSRIs), are clinically effective in the treatment of all anxiety disorders, including post traumatic stress disorder (PTSD) and obsessive compulsive
25 disorder (OCD). While the clinical effectiveness of traditional anxiolytics, such as benzodiazepines (BDZs), is limited to generalized anxiety disorder or acute panic attacks (Borsini et al). Thus the potential anxiolytic effect of antidepressants is of great relevance.

30 The Open Field Test - Exploratory locomotor activity used in Examples 3.1 and 3.2 described below is one of the

most popular in evaluation of animals' behavior. It tests both motility parameters and anxiety (Prut et al). An individual mouse is placed in a novel plexiglass arena of 50x50cm the floor of which is divided into 3 digital zones: the outer peripheral zone 1, the medial zone 2 and the most central zone 3. The animal behavior in the open field is recorded by videotaping for 20 min and is analyzed subsequently digitally using Noldus software for animal behavior. The measurements include general motility (distance moved, velocity and strong mobility) and anxiety parameters (including frequency of visits to the central area, time spent in the inner field, and number of rearing events in the center). The more the animal stays and performs in the center, the less anxious it is.

Example 3.1

Balb/c male mice were divided to 6 groups (5/group) and administered laquinimod (1, 5, 10 and 25 mg/kg p.o. in the form of laquinimod sodium solution via gavage), fluoxetine (positive control 10mg/kg po) or vehicle daily for 3 days. On the third day, 90 min following administration, mice were exposed to the open field for 20 min. Behavior was video recorded and analyzed using Ethovision software (Noldus Holland). The results with respect to motility parameters are shown in **Fig. 6**. The results with respect to anxiety parameters are shown in **Figs. 7 and 8**.

Conclusion:

Motility Parameters - Laquinimod and fluoxetine did not modify significantly the motility parameters in the field in this model.

Anxiety Parameters (Zone 2) - Animals treated with laquinimod and with fluoxetine tended to perform more in the center in terms of frequency, time spent, distance moved and number of rearings. The results show 1 mg/kg and 5 mg/kg were the most potent.

Anxiety Parameters (Zones 2 and 3) - Animals treated with laquinimod tended to perform more in the center as evidenced by time spent, distance moved and number of rearings. The results show 1 mg/kg was the most potent dosage of laquinimod in this experiment.

This model shows that laquinimod shows anxiolytic effects.

Example 3.2

Balb/c male mice were divided to 6 groups (5/group) and administered laquinimod (0.5, 1, 5 and 25mg/kg p.o. in the form of laquinimod sodium solution via gavage), fluoxetine (positive control 10mg/kg po) or vehicle daily for 3 days. On the third day, 90 min following administration, mice were exposed to the open field test for 20 minutes. Behavior was video recorded and analyzed using Ethovision software (Noldus Holland). The motility parameter results are shown in **Fig. 9**. The anxiety parameter results are shown in **Fig. 10**.

Conclusion:

Motility Parameters - Laquinimod (0.5-25 mg/kg/dx3d po) induced a slight increase in motility (0.5-5 mg/kg) in the open field in Balb/c mice.

Anxiety Parameters - Animals treated with laquinimod (0.5-5mg/kg po) and with Fluoxetine (10mg/kg po) performed significantly more in the center as judged by

frequency, time spent and number of rearings. 5mg/kg was the most potent.

This model shows that laquinimod shows anxiolytic effects.

EXAMPLE 4: LAQUINIMOD SHOWS ANXIOLYTIC ACTIVITY IN MICE

5 MODELS - ELEVATED PLUS MAZE (EPM)

The EPM model used in Examples 4.1 and 4.2 described below utilizes the natural fear of rodents to avoid open and elevated places. The apparatus consists of a plus-maze with two enclosed and two opposite open arms,
10 elevated above the floor. Naive animals spend only about 30% of the test time on open arms, while treatment with benzodiazepines significantly increases open-arm exploration (Pellow et al). This is one of the most widely used models to study effects of anxiety-like
15 behavior.

The maze consisted of two opposing open arms (40 x 10cm) and two opposing closed arms (40 x 10 cm, with 40 cm walls) on a platform 50 cm above the ground. Mice were placed in the center square (10 x 10cm) facing an open
20 arm and videotaped during a 5 min exploration. Arm entries and duration were scored when all four paws enter the arm.

Example 4.1

Balb/c male mice were divided to 6 groups (5/group) and
25 administered laquinimod (0.5, 1, 5 and 25mg/kg p.o. in the form of laquinimod sodium solution via gavage), fluoxetine (positive control 10mg/kg po) or vehicle daily for 3 days. On the third day, 90 min following administration, mice were exposed to the EPM for 5
30 minutes. Mice were placed in the center square (10 x 10cm) facing an open arm and videotaped during a 5 minute

exploration. Arm entries and duration was scored when all four paws enter the arm. . Behavior was video recorded and analyzed using Ethovision software (Noldus Holland). The results are shown in **Fig. 11**.

5 Conclusion:

Mice treated with laquinimod show anxiolytic activity in a dose dependent manner. Maximal effect was obtained with the dose of 25mg/kg po.

Example 4.2

10 Balb/c male mice were divided to 6 groups (5/group) and administered laquinimod (0.5, 1, 5 and 25mg/kg p.o. in the form of laquinimod sodium solution via gavage), fluoxetine (positive control 10mg/kg po) or vehicle daily for 3 days. On the third day, 90 min following
15 administration, mice were exposed to the EPM for 5 min. Behavior was video recorded and open arms parameters were analyzed using Ethovision software (Noldus Holland). The results are shown in **Fig. 12**.

Conclusion:

20 Mice treated with laquinimod show anxiolytic activity in a dose dependent manner. Maximal effect was obtained with 5-25mg/kg po.

References

1. PCT International Application Publication No. WO 2007/047863, published April 26, 2007, international filing date October 18, 2006.
- 5 2. PCT International Application Publication No. WO 2007/146248, published December 21, 2007, international filing date June 12, 2007.
3. Acheson, A, et al. (1995). "A BDNF autocrine loop in adult sensory neurons prevents cell death". Nature,
10 374(6521):450-3.
4. Alonsa, M, et al. (2005) "Endogenous BDNF is required for long-term memory formation in the rat parietal cortex". Learning & Memory, 12:504-510.
5. Amaral, MD, et al. (2007) "TRPC channels as novel
15 effectors of BDNF signaling: Potential implications for Rett syndrome". Pharmacol Ther, 113(2):394-409.
18. Borsini F, et al. (2002) "Do animal models of anxiety predict anxiolytic-like effects of antidepressants?" Psychopharmacology. 163:121-41.
- 20 6. Caffè Romeo, A, et al. (2001) "A combination of CNTF and BDNF rescues rd photoreceptors but changes rod differentiation in the presence of RPE in retinal explants". Investigative Ophthalmology & Visual Science, 42:275-82.
- 25 7. Chesselet, MF (2003) "Dopamine and Parkinson's disease: is the killer in the house?" Molecular Psychiatry, 8:369-370.
8. Ciammola, A, et al. (2007) "Low brain-derived neurotrophic factor (BDNF) levels in serum of

- Huntington's disease patients". Am J Med Gent Part B, 144b:574-577.
9. Cryan JF, et al. (2002) "Assessing antidepressant activity in rodents: recent developments and future needs". Trend in Pharmacological Science. 23:238-45.
10. Gambarana, C., et al. (2001) "Animal models for the study of antidepressant activity". Brain Res. Protocol. 7:11-20.
11. Howells, DW, et al. (2000) "Reduced BDNF mRNA expression in the Parkinson's disease substantia nigra". Experimental Neurology, 166(1):127-135.
12. Huang, EJ and Reichardt, LF (2001) "Neurotrophins: roles in neuronal development and function". Annu. Rev. Neurosci, 24:677-736.
13. Hyman, C. et al., (1991) "BDNF is a neurotrophic factor for dopaminergic neurons of the substantia nigra". Nature, 350(6315):230-2.
14. Kato-Semba, R, et al. (2002) "Riluzole enhances expression of brain-derived neurotrophic factor with consequent proliferation of granule precursor cells in the rat hippocampus". FASEB J, 16:1328-30.
19. Kessler RC, et al. (June 2005) "Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication". Arch. Gen. Psychiatry. 62(6):617-27.
15. Molteni, R, et al. (2006) "Abstract: Chronic treatment with fluoxetine [Prozac®] up-regulates cellular BDNF mRNA expression in rat dopaminergic regions". Int J Neuropsychopharmacol. 9(3):307-17.

16. Monteggia, LM (2007) "Elucidating the role of brain-derived neurotrophic factor in the brain". Am J Psychiatry, 164:1790.
20. Pellow S, et al. (1985) "Validation of open:closed arm entries in an elevated plus-maze as a measure of anxiety in the rat". J Neurosci Methods. 14:149-167.
21. Porsolt R.D., et al. (1977) "Behavioral despair in mice: a primary screening test for antidepressants". Arch. Int. Pharmacodyn. Ther. **229**: 327-336.
- 10 22. Prut L, and Belzung C. (2003) "The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review". Eur J Pharmacol. 463:3-33.
17. Riviere, M (1998) "An analysis of extended survival in patients with amyotrophic lateral sclerosis treated with riluzole". Arch Neurol, 55:526-8.
- 15
18. Sarchielli, P, et al. (2007) "Production of Brain-derived neurotrophic factor by mononuclear cells of patients with multiple sclerosis treated with glatiramer acetate, interferon- β 1a, and high doses of immunoglobulins". Multiple Sclerosis, 13:313-331.
- 20
19. Sen, S, et al. (2008) "Serum brain-derived neurotrophic factor, depression, and antidepressant medications: meta-analyses and implications". Biol Psychiatry, 64:527-532.
20. Snider, et al., (1989) "Neurotrophic molecules". Ann Neurol, 26(4):489-506.
- 25
21. Tramontina, JF, et al. (2009) "brain-derived neurotrophic factor serum levels before and after treatment for acute mania". Neuroscience Letters, 452:111-3.

What is claimed is:

1. Use of laquinimod or a pharmaceutically acceptable salt thereof for treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, bipolar disorder, and acute mania, wherein the laquinimod or pharmaceutically acceptable salt thereof is formulated for periodic administration to the subject.
2. The use of claim 1, wherein the periodic administration comprises once daily administration.
3. The use of claims 1 or 2, wherein the periodic administration continues for at least 3 days.
4. The use of any one of claims 1-3, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-40.0 mg/day.
5. The use of claim 4, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-0.6 mg/day.
6. The use of claim 4, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.5 mg/day.
7. The use of claim 4, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.6 mg/day.
8. The use of claim 4, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 1.0 mg/day.

9. The use of any one of claims 1-8, wherein the laquinimod or pharmaceutically acceptable salt thereof is formulated for oral administration.
10. The use of any one of claims 1-9, wherein the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.
11. The use of any one of claims 1-10, wherein the BDNF-related disease is Huntington's disease.
12. The use of any one of claims 1-10, wherein the BDNF-related disease is a depressive disorder selected from the group consisting of depression, depression in a cancer patient, depression in a Parkinson's Disease patient, postmyocardia infarction depression, depression in a patient with human immunodeficiency virus, subsyndromal symptomatic depression, depression in an infertile woman, pediatric depression, major depression, single episode depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, manic depressive illness, manic depressive illness with mixed episodes, manic depressive illness with depressive episodes, seasonal affective disorder, bipolar depression BP I, bipolar depression BP II, and major depression with dysthymia.
13. The use of any one of claims 1-10, wherein the BDNF-related disease is an anxiety disorder and is selected from the group consisting of generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, and childhood anxiety.
14. The use of any one of claims 1-13, wherein the subject is also being treated with a second BDNF-increasing agent which is fluoxetine or riluzole.
15. The use of claim 14, wherein the administration of the laquinimod precedes the administration of the second BDNF-

increasing agent.

16. The use of claim 14, wherein the administration of the second BDNF-increasing agent precedes the administration of the laquinimod.
17. Use of laquinimod or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, bipolar disorder, and acute mania, wherein the medicament is formulated for periodic administration.
18. The use of claim 17, wherein the periodic administration comprises once daily administration.
19. The use of claims 17 or 18, wherein the periodic administration continues for at least 3 days.
20. The use of any one of claims 17-19, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-40.0 mg/day.
21. The use of claim 20, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-0.6 mg/day.
22. The use of claim 20, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.5 mg/day.
23. The use of claim 20, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.6 mg/day.
24. The use of claim 20, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 1.0 mg/day.

25. The use of any one of claims 17-24, wherein the laquinimod or pharmaceutically acceptable salt thereof is formulated for oral administration.
26. The use of any one of claims 17-25, wherein the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.
27. The use of any one of claims 17-26, wherein the BDNF-related disease is Huntington's disease.
28. The use of any one of claims 17-26, wherein the BDNF-related disease is a depressive disorder selected from the group consisting of depression, depression in a cancer patient, depression in a Parkinson's Disease patient, postmyocardia infarction depression, depression in a patient with human immunodeficiency virus, subsyndromal symptomatic depression, depression in an infertile woman, pediatric depression, major depression, single episode depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, manic depressive illness, manic depressive illness with mixed episodes, manic depressive illness with depressive episodes, seasonal affective disorder, bipolar depression BP I, bipolar depression BP II, and major depression with dysthymia.
29. The use of any one of claims 17-26, wherein the BDNF-related disease is an anxiety disorder and is selected from the group consisting of generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, and childhood anxiety.
30. The use of any one of claims 17-29, wherein the subject is also being treated with a second BDNF-increasing agent which is fluoxetine or riluzole.
31. The use of claim 30, wherein the administration of the

laquinimod precedes the administration of the second BDNF-increasing agent.

32. The use of claim 30, wherein the administration of the second BDNF-increasing agent precedes the administration of the laquinimod.
33. Laquinimod or a pharmaceutically acceptable salt thereof for use in treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, bipolar disorder, and acute mania, wherein the laquinimod or pharmaceutically acceptable salt thereof is formulated for periodic administration.
34. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claim 33, wherein the periodic administration comprises once daily administration.
35. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claims 33 or 34, wherein the periodic administration continues for at least 3 days.
36. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-35, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-40.0 mg/day.
37. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claim 36, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-0.6 mg/day.
38. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claim 36, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.5 mg/day.

39. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claim 36, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.6 mg/day.
40. Laquinimod or a pharmaceutically acceptable salt thereof for the use of claim 36, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 1.0 mg/day.
41. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-40, wherein the laquinimod or pharmaceutically acceptable salt thereof is formulated for oral administration.
42. Laquinimod or a pharmaceutically acceptable salt thereof for use of any one of claims 33-41, wherein the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.
43. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-42, wherein the BDNF-related disease is Huntington's disease.
44. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-42, wherein the BDNF-related disease is a depressive disorder selected from the group consisting of depression, depression in a cancer patient, depression in a Parkinson's Disease patient, postmyocardia infarction depression, depression in a patient with human immunodeficiency virus, subsyndromal symptomatic depression, depression in an infertile woman, pediatric depression, major depression, single episode depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, neuropathic pain, manic depressive illness, manic depressive illness with mixed episodes, manic depressive illness with depressive episodes, seasonal affective disorder,

bipolar depression BP I, bipolar depression BP II, and major depression with dysthymia.

45. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-42, wherein the BDNF-related disease is an anxiety disorder and is selected from the group consisting of generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, and childhood anxiety.
46. Laquinimod or a pharmaceutically acceptable salt thereof for the use of any one of claims 33-45, wherein the subject is also being treated with a second BDNF-increasing agent which is fluoxetine or riluzole.
47. Laquinimod or a pharmaceutically acceptable salt thereof for use of claim 46, wherein the administration of the laquinimod precedes the administration of the second BDNF-increasing agent.
48. Laquinimod or a pharmaceutically acceptable salt thereof for use of claim 46, wherein the administration of the second BDNF-increasing agent precedes the administration of the laquinimod.
49. A pharmaceutical composition comprising laquinimod and a pharmaceutically acceptable carrier, for treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, bipolar disorder, and acute mania.
50. The pharmaceutical composition of claim 49, wherein the pharmaceutical composition is formulated for periodic administration.
51. The pharmaceutical composition of claim 50, wherein the periodic administration comprises once daily administration.

52. The pharmaceutical composition of claims 50 or 51, wherein the periodic administration continues for at least 3 days.
53. The pharmaceutical composition of any one of claims 49-52, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-40.0 mg/day.
54. The pharmaceutical composition of claim 53, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-0.6 mg/day.
55. The pharmaceutical composition of claim 53, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.5 mg/day.
56. The pharmaceutical composition of claim 53, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.6 mg/day.
57. The pharmaceutical composition of claim 53, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 1.0 mg/day.
58. The pharmaceutical composition of any one of claims 49-57, wherein laquinimod or pharmaceutically acceptable salt thereof is formulated for oral administration.
59. The pharmaceutical composition of any one of claims 49-58, wherein the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.
60. The pharmaceutical composition of any one of claims 49-59, wherein the BDNF-related disease is Huntington's disease.
61. The pharmaceutical composition of any one of claims 49-59, wherein the BDNF-related disease is a depressive disorder selected from the group consisting of depression, depression in a cancer patient, depression in a Parkinson's Disease patient, postmyocardia infarction depression, depression in a

patient with human immunodeficiency virus, subsyndromal symptomatic depression, depression in an infertile woman, pediatric depression, major depression, single episode depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, manic depressive illness, manic depressive illness with mixed episodes, manic depressive illness with depressive episodes, seasonal affective disorder, bipolar depression BP I, bipolar depression BP II, and major depression with dysthymia.

62. The pharmaceutical composition of any one of claims 49-59, wherein the BDNF-related disease is an anxiety disorder and is selected from the group consisting of generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, and childhood anxiety.
63. A pharmaceutical preparation comprising laquinimod and a second BDNF-increasing agent which is fluoxetine or riluzole, for treating a human subject suffering from a BDNF-related disease selected from the group consisting of Parkinson's disease, Huntington's disease, a depressive disorder, an anxiety disorder, retinitis pigmentosa, erectile dysfunction, a memory disorder, Rett syndrome, bipolar disorder, and acute mania.
64. The pharmaceutical preparation of claim 63, wherein the pharmaceutical preparation is formulated for periodic administration.
65. The pharmaceutical preparation of claim 64, wherein the periodic administration comprises once daily administration.
66. The pharmaceutical preparation of claims 64 or 65, wherein the periodic administration continues for at least 3 days.

67. The pharmaceutical preparation of any one of claims 63-66, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-40.0 mg/day.
68. The pharmaceutical preparation of claim 67, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.1 mg/day-0.6 mg/day.
69. The pharmaceutical preparation of claim 67, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.5 mg/day.
70. The pharmaceutical preparation of claim 67, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 0.6 mg/day.
71. The pharmaceutical preparation of claim 67, in which the laquinimod or pharmaceutically acceptable salt thereof is formulated for administration at 1.0 mg/day.
72. The pharmaceutical preparation of any one of claims 63-71, wherein laquinimod or pharmaceutically acceptable salt thereof is formulated for oral administration.
73. The pharmaceutical preparation of any one of claims 63-72, wherein the pharmaceutically acceptable salt of laquinimod is laquinimod sodium.
74. The pharmaceutical preparation of any one of claims 63-73, wherein the BDNF-related disease is Huntington's disease.
75. The pharmaceutical preparation of any one of claims 63-73, wherein the BDNF-related disease is a depressive disorder selected from the group consisting of depression, depression in a cancer patient, depression in a Parkinson's Disease patient, postmyocardia infarction depression, depression in a patient with human immunodeficiency virus, subsyndromal symptomatic depression, depression in an infertile woman, pediatric depression, major depression, single episode

depression, recurrent depression, child abused-induced depression, post-partum depression, DSM-IV major depression, treatment-refractory major depression, severe depression, psychotic depression, post-stroke depression, manic depressive illness, manic depressive illness with mixed episodes, manic depressive illness with depressive episodes, seasonal affective disorder, bipolar depression BP I, bipolar depression BP II, and major depression with dysthymia.

76. The pharmaceutical preparation of any one of claims 63-73, wherein the BDNF-related disease is an anxiety disorder and is selected from the group consisting of generalized anxiety, panic disorder, phobia, post traumatic stress disorder, obsessive compulsive disorder, separation anxiety, and childhood anxiety.

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D

FIGURE 5




Fig. 5A




Fig. 5B




Fig. 5C




Fig. 5D

FIGURE 6




Fig. 6A




Fig. 6B




Fig. 6C




Fig. 6D

FIGURE 7

In zone frequency (zone 2)




Fig. 7A

In zone total duration (zone 2)




Fig. 7B

Rearing frequency (zone 2)




Fig. 7C




Fig. 7D

In zone total duration (zone 3)




FIGURE 8

Fig. 8A

Distance Moved in the center (zone 3)




Fig. 8B




Fig. 8C

FIGURE 9




Fig. 9A




Fig. 9B




Fig. 9C




Fig. 9D

FIGURE 10

In zone frequency (zone 2+3)




Fig. 10A

In zone total duration (zone 2+3)




Fig. 10B

Rearing (zone 2+3)




Fig. 10C

FIGURE 11




Fig. 11A




Fig. 11B

Strong mobility Frequency (Open Arm)




Fig. 11C




Fig. 11D

FIGURE 12




Fig. 12A




Fig. 12B




Fig. 12C




Fig. 12D