



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:
14.07.2021 Bulletin 2021/28

(21) Application number: **21158464.4**

(22) Date of filing: **11.12.2016**

(51) Int Cl.:
A61K 31/506 ^(2006.01) **C07D 405/12** ^(2006.01)
A61P 25/00 ^(2006.01) **C07D 201/00** ^(2006.01)
A61K 31/00 ^(2006.01) **A61K 33/00** ^(2006.01)
C07D 401/14 ^(2006.01) **C07D 413/14** ^(2006.01)
C07D 237/20 ^(2006.01) **C07D 417/04** ^(2006.01)
C07D 417/14 ^(2006.01) **C07D 471/10** ^(2006.01)
C07D 487/10 ^(2006.01) **C07D 403/12** ^(2006.01)

(84) Designated Contracting States:
AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR

(30) Priority: **10.12.2015 US 201562265652 P**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
16874024.9 / 3 386 511

(71) Applicant: **PTC Therapeutics, Inc.**
South Plainfield, NJ 07080 (US)

(72) Inventors:
• **Babu, Suresh**
Pennington, NJ 08534 (US)
• **Bhattacharyya, Anuradha**
Edison, NJ 08820 (US)


- **Hwang, Seongwoo**
Edison, NJ 08817 (US)
- **Jani, Minakshi**
Iselin, NJ 08830 (US)
- **Moon, Young-choon**
Belle Mead, NJ 08502 (US)
- **Sydorenko, Nadiya**
Princeton, NJ 08540 (US)

(74) Representative: **Kolster Oy Ab**
Salmisaarenaukio 1
P.O. Box 204
00181 Helsinki (FI)

Remarks:
This application was filed on 22-02-2021 as a
divisional application to the application mentioned
under INID code 62.

(54) **1,4-DISUBSTITUTED PYRIDAZINE COMPOUNDS FOR TREATING HUNTINGTON'S DISEASE**

(57) The present description relates to compounds, forms, and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease.



In particular, the present description relates to substituted monocyclic heteroaryl compounds of Formula (I), forms and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease.

Description

[0001] The present description relates to compounds, forms, and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease. In particular, the present description relates to substituted monocyclic heteroaryl compounds, forms and pharmaceutical compositions thereof and methods of using such compounds, forms, or compositions thereof for treating or ameliorating Huntington's disease.


BACKGROUND

[0002] Huntington's disease (HD) is a progressive, autosomal dominant neurodegenerative disorder of the brain, having symptoms characterized by involuntary movements, cognitive impairment, and mental deterioration. Death, typically caused by pneumonia or coronary artery disease, usually occurs 13 to 15 years after the onset of symptoms. The prevalence of HD is between three and seven individuals per 100,000 in populations of western European descent. In North America, an estimated 30,000 people have HD, while an additional 200,000 people are at risk of inheriting the disease from an affected parent. The disease is caused by an expansion of uninterrupted trinucleotide CAG repeats in the "mutant" huntingtin (Htt) gene, leading to production of HTT (Htt protein) with an expanded poly-glutamine (polyQ) stretch, also known as a "CAG repeat" sequence. There are no current small molecule therapies targeting the underlying cause of the disease, leaving a high unmet need for medications that can be used for treating or ameliorating HD. Consequently, there remains a need to identify and provide small molecule compounds for treating or ameliorating HD.

[0003] All other documents referred to herein are incorporated by reference into the present application as though fully set forth herein.

SUMMARY

[0004] The present description relates to methods for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I):




wherein W, X, A and B are as defined herein, or forms and compositions thereof.

[0005] In particular, the present description relates to a use of a compound of Formula (I) or a form or composition thereof in a method for treating or ameliorating HD in a subject in need thereof comprising, administering an effective amount of the compound or a form or composition thereof, to the subject.

[0006] The present description further relates to the use of a compound of Formula (I) or a form thereof in combination with agents having additive or synergistic activity, thus providing a combination product for the treatment of HD.

DETAILED DESCRIPTION

[0007] The present description relates to a method or use of a compound for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I):



or a form thereof, wherein

W is CH=CH or S;

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy;


R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino; and

R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0008] An embodiment of the present description further relates to a method or use of a compound for treating or

ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) selected from a compound of Formula (Ia) and Formula (Ib):



10 or a form thereof, wherein

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

15 wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

20 wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

25 B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

30 R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

35 wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

40 wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

45 R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl;

50 R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy;


55 R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkylamino, (C₁₋₄alkyl)₂-amino

or hydroxyl-C₁₋₄alkyl-amino; and

R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0009] Another embodiment of the present description further relates to methods for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I) selected from a compound of Formula (Ia) and Formula (Ib):




or a form thereof, wherein


X is O, NH, N(CH₃) or a bond;


A is aryl, heteroaryl or heterocyclyl,

wherein aryl is selected from the group consisting of




wherein heteroaryl is selected from the group consisting of







a12




a13




a14




a15




a16




a17




a18




a19




a20




a21




a22




a23




a24




a25




a26




a27




a28




a29




a30




a31




a32




a33



a34




a35, and




a36


wherein heterocyclyl is selected from the group consisting of




a37




a38




a39




a40




a41




a42




a43




a44



a45




a46, and




a47


B is heterocyclyl selected from the group consisting of




b1




b2



b3



b4



R_{1a}, R_{1b} and R_{1c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-ami-

no-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R_{2a}, R_{2b} and R_{2c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and


wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy; and


R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f} and R_{4g} are independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.



[0010] Another aspect of the present description relates to a compound of Formula (I) selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1):



(Ia11),



(Ia15),



or a form thereof, wherein (when present),

X is selected from O, NR₅, or a bond;

A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy,

wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a};

B is selected from 1H-pyrazolyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, (1*R*,5*S*)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl, 2,6-diazaspiro[3.4]octyl or 2,7-diazaspiro[3.5]nonyl, each optionally substituted with 1 or 2 substituents each selected from R_{4a};

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or heteroaryl, wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S optionally substituted with 1 or 2 substituents each selected from R_{3a};

R_{3a} is selected from nitro or C₁₋₄alkyl; and,

R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0011] Another aspect of the present description relates to a compound of Formula (I) selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1): or a form thereof, wherein (when present),

R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy or 1H-pyrazolyl or 1H-imidazol-1-yl,

wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a};


R_{3a} is selected from nitro or methyl or amino; and,

R_{4a} is methyl or ethyl;

R_{5a} is hydrogen or methyl;


wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0012] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia1) or a form thereof, wherein substituents R_{1a}, R_{1b}, and X, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a}, R_{1b}, and X substituents are not present:



Cpd	R _{1a}	R _{1b}	X
1	--	--	NH
8	2-OH	--	N(CH ₃)
40	1-CH ₂ CH=CH ₂ , 2-OH	--	N(CH ₃)
106	1-Br, 2-OH	7-OH	N(CH ₃)
107	1-Cl, 2-OH	7-OH	N(CH ₃)

[0013] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia2) or a form thereof, wherein substituents R_{1a}, R_{1b}, and R_{4a}, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a}, R_{1b}, and R_{4a} substituents are not present:




(Ia2)

Cpd	R _{1a}	R _{1b}	R _{4a}
13	--	--	--
207	--	7-OH	--
208	--	7-OH	CH ₃
210	2-CH ₃	7-OH	--
222	3-Cl	7-OH	--
223	3-Br	7-OH	--
224	3-CN	7-OH	--
225	3-(1-CH ₃ -1H-imidazol-4-yl)	7-OH	--
226	3-(1H-imidazol-1-yl)	7-OH	--
227	3-OH	7-OH	--
228	3-CH ₂ CH ₃	7-OH	--
229	3-CH(CH ₃) ₂	7-OH	--
232	2-CH ₃ , 4-OCH ₃	7-OH	--
233	2-CH ₃ , 4-(pyrrolidin-1-yl)	7-OH	--

(continued)

Cpd	R _{1a}	R _{1b}	R _{4a}
234	2-CH ₃ , 4-(morpholin-4-yl)	7-OH	-
235	2-CH ₃ , 4-N(CH ₃) ₂	7-OH	-
236	2-CH ₃ , 4-OCH ₂ CH ₃	7-OH	-
237	2-CH ₃ , 4-(1-CH ₃ -1H-pyrazol-4-yl)	7-OH	-
240	3-(tetrahydro-2H-pyran-4-yl)	7-OH	-
249	4-OCH ₃	7-OH	-
250	2-CH ₃ , 4-(azetidin-1-yl)	7-OH	-
251	2-CH ₃ , 4-CN	7-OH	-
252	2-CH ₃ , 4-cyclopropyl	7-OH	-
253	2-CH ₃ , 4-(3,6-dihydro-2H-pyran-4-yl)	7-OH	-
254	2-CH ₃ , 4-(tetrahydro-2H-pyran-4-yl)	7-OH	-
255	2-CH ₃ , 4-(oxetan-3-yl)	7-OH	-
256	4-N(CH ₃) ₂	7-OH	-
262	2-CN	7-OH	-
265	2-C(O)NH ₂	7-OH	-
293	3-Cl	7-OH	-
294	3-CH(CH ₃) ₂	7-OH	-
296	2-CH ₃ , 4-Cl	7-OH	-


[0014] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia3) or a form thereof, wherein substituents R_{1a}, R_{1b} and X, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a}, R_{1b} and X substituents are not present:



(Ia3)

Cpd	R _{1a}	R _{1b}	X
11	--	--	O
15	--	--	N(CH ₃)
218	--	7-OH	N(CH ₃)
261	1-CN	7-OH	N(CH ₃)
272	1-CH ₃	7-OH	N(CH ₃)
275	1-CN, 3-CH ₃	7-OH	N(CH ₃)
292	3-(OCH ₂ -phenyl)	--	N(CH ₃)

[0015] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia4) or a form thereof, wherein substituents X, R_{1a}, R_{1b} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a}, R_{1b} and R_{4a} substituents are not present:




(Ia4)

Cpd	R _{1a}	R _{1b}	X	R _{4a}
10	--	--	O	--
14	--	--	N(CH ₃)	--
159	1-(OCH ₂ -phenyl)	--	N(CH ₃)	--
211	--	6-OH	N(CH ₃)	CH ₃
212	--	6-OH	N(CH ₃)	--
213	--	6-OH	O	--
215	1-cyclopropyl	6-OH	N(CH ₃)	--
216	1-OH	6-OH	N(CH ₃)	--
217	1-CN	6-OH	N(CH ₃)	--
264	1-C(O)NH ₂	6-OH	N(CH ₃)	--
273	1-CH ₃	6-OH	N(CH ₃)	--
274	1,3-(CH ₃) ₂	6-OH	N(CH ₃)	--
276	1-NH ₂	6-OH	N(CH ₃)	--
283	1-OCH ₂ CH ₃	6-OH	N(CH ₃)	--
284	1-OH	6-OH	O	--
285	3-phenyl	6-OH	N(CH ₃)	--

(continued)

Cpd	R _{1a}	R _{1b}	X	R _{4a}
286	3-CH ₃	6-OH	N(CH ₃)	--
287	3-cyclopropyl	6-OH	N(CH ₃)	--
288	3-CH(CH ₃) ₂	6-OH	N(CH ₃)	--
289	3-(CH ₂) ₂ CH ₃	6-OH	N(CH ₃)	--
290	3-CH(CH ₃) ₂	6-OH	O	--


[0016] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia5) or a form thereof, wherein substituents R_{1a} and R_{1b}, when present, are indicated in the table below with multiple substituents separated by a comma; and, "--" indicates that one or more R_{1a} and R_{1b} substituents are not present:



(Ia5)


Cpd	R _{1a}	R _{1b}
12	--	--
220	--	6-OH
221	2-CH ₃	6-OH
238	4-OCH ₃	6-OH
241	3-Cl	6-OH
242	3-Br	6-OH
243	3-CH ₃	6-OH
244	3-CH ₃	5-Br, 6-OH
263	2-CN	6-OH
266	2-C(O)NH ₂	6-OH
267	2-CO ₂ CH ₃	6-OH
297	4-Cl	6-OH
300	--	6-OH

[0017] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia6) or a form thereof, wherein substituents R_{1a}, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:




Cpd	R _{1a}
239	--
246	2,3-(CH ₃) ₂
247	2-CH ₃
248	3-CH ₃

[0018] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia7) or a form thereof, wherein substituents R_{1a}, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:




Cpd	R _{1a}
258	--
260	2-CH ₃

[0019] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia8) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:




Cpd	R _{1a}	B
209	--	6-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
269	2-CN	piperazin-1-yl

[0020] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia9) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:




Cpd	R _{1a}	B
214	--	6-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
270	--	piperazin-1-yl
291	3-CH ₃	piperazin-1-yl

[0021] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia10) or a form thereof, wherein substituents R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and B substituents are not present:



Cpd	R _{1a}	B
268	2-CN	piperazin-1-yl
271	--	1,2,3,6-tetrahydropyridin-4-yl

[0022] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia11) or a form thereof, wherein substituents A, X and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more A, X and R_{4a} substituents are not present:



(Ia11)

Cpd	A	X	R _{4a}
2	benzo[b]thiophen-2-yl	N(CH ₃)	--
4	5-CN-benzo[b]thiophen-2-yl	N(CH ₃)	--
5	quinolin-3-yl	NH	--
6	benzo[b]thiophen-2-yl	O	--
9	benzo[b]thiophen-2-yl	NH	--
16	imidazo[1,2-a]pyridin-6-yl	N(CH ₃)	--
17	6-phenyl-pyridin-3-yl	N(CH ₃)	--
18	6-(1H-pyrrol-1-yl)-pyridin-3-yl	N(CH ₃)	--
19	6-(1H-pyrazol-1-yl)-pyridin-3-yl	N(CH ₃)	--
20	quinoxalin-2-yl	N(CH ₃)	--
21	quinolin-3-yl	N(CH ₃)	--
22	phthalazin-6-yl	N(CH ₃)	--
23	benzo[c][1,2,5]oxadiazol-5-yl	NH	--
24	benzo[d]thiazol-5-yl	NH	--
25	2-CH ₃ -benzo[d]oxazol-6-yl	NH	--
30	2-(4-CN-phenol)	N(CH ₃)	--
32	2-(4-CF ₃ -phenol)	N(CH ₃)	--
33	6-(2-F-phenol)	N(CH ₃)	--
34	2-[3,5-(OCH ₃) ₂ -phenol]	N(CH ₃)	--
35	2-[4,5-(OCH ₃) ₂ -phenol]	N(CH ₃)	--
37	2-(4,5-F ₂ -phenol)	N(CH ₃)	--
41	benzo[b]thiophen-2-yl	NH	CH ₃
53	2-[4-(1H-pyrazol-1-yl)-phenol]	N(CH ₃)	--
115	2-[3-OH-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
116	2-[3-OCH ₃ -5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
117	2-[5-(1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	NH	--
118	2-[5-(1-CH ₃ -1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	N(CH ₃)	--
119	2-[5-(1H-pyrazol-4-yl)-3-OCF ₃ -phenol]	N(CH ₃)	--
120	2-[5-(1-CH ₃ -pyridin-2(1H)-one)-3-OCF ₃ -phenol]	N(CH ₃)	--
121	2-[3-OCH ₃ -5-(1-CH ₃ -1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
122	2-[3-OCH ₃ -5-(5,6,7,8-tetrahydroimidazo-[1,2-a]pyridin-3-yl)-phenol]	N(CH ₃)	--
123	2-[3-OCH ₃ -5-(pyridin-3-yl)-phenol]	N(CH ₃)	--

EP 3 848 035 A1

(continued)

5

10

15

20

25

30

35

40

45

50


55

Cpd	A	X	R _{4a}
124	2-[3-OCH ₃ -5-(1-cyclopentyl-1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
125	2-[5-(3-OCH ₃ -phenyl)-3-OCH ₃ -phenol]	N(CH ₃)	--
126	2-[3-benzyloxy-5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
127	2-[3-OCH ₂ CH ₃ -5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
128	2-[3-(OCH ₂ -cyclopropyl)-5-(5-CH ₃ -oxazol-2-yl)-phenol]	N(CH ₃)	--
129	5-(2-CH ₃ -1H-benzo[d]imidazol-6-ol)	N(CH ₃)	--
134	2-[4-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
135	2-[4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)-phenol]	N(CH ₃)	--
136	2-[4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)-phenol]	N(CH ₃)	--
137	2-[4-(1H-indol-2-yl)-phenol]	N(CH ₃)	--
138	2-[4-(cyclopent-1-en-1-yl)-phenol]	N(CH ₃)	--
139	2-[4-(1H-pyrazol-3-yl)-phenol]	N(CH ₃)	--
140	2-[4-(2-OH-pyridin-4-yl)-phenol]	N(CH ₃)	--
141	2[4-(1-CH ₃ -pyridin-2(1H)-one)-phenol]	O	--
142	2-[4-(2-OH-pyridin-4-yl)-phenol]	O	--
144	2-[4-Cl-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
145	2-[4-F-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
146	2-[5-F-4-(1H-imidazol-4-yl)-phenol]	N(CH ₃)	--
147	2-[5-F-4-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
148	2-[5-F-(1H-pyrazol-5-yl)-phenol]	N(CH ₃)	--
149	6-OH-1-oxo-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
150	6-(1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol)	N(CH ₃)	--
151	6-OH-1-OH-imino-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
152	6-OH-1-OH-2,3-dihydro-1H-inden-5-yl	N(CH ₃)	--
153	6-(2-NH ₂ -8H-indeno[1,2-d]thiazol-5-ol)	N(CH ₃)	--
154	9-(5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol)	N(CH ₃)	--
155	2-{ 4-[C(O)NHCH ₂ -(1-CH ₃ -1H-pyrazol-4-yl)]-phenol}	N(CH ₃)	--
156	2-[4-(4-CH ₂ OH-1H-pyrazol-1-yl)-phenol]	N(CH ₃)	--
158	3-(OCH ₂ -phenyl)-isoquinolin-6-yl	N(CH ₃)	--
160	2-[3-F-5-(2-OCH ₃ -pyridin-4-yl)-phenol]	N(CH ₃)	--
161	4-[1-(4-pyridin-2(1H)-one)-3-F-5-OH-phenyl]	N(CH ₃)	--
162	4-{1-[4-(1-CH ₃ -pyridin-2(1H)-one)]-3-F-5-OH-phenyl}	N(CH ₃)	--
163	4-{1-[5-51-CH ₃ -pyridin-2(1H)-one)]-3-F-5-OH-phenyl}	N(CH ₃)	--
164	2-[3-F-5-(1H-pyrazol-4-yl)-phenol]	O	--
165	2-(5-Cl-3-F-phenol)	N(CH ₃)	--
166	2-[3-F-5-(1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
167	2-[3-F-5-(1-CH ₃ -1H-pyrazol-4-yl)-phenol]	N(CH ₃)	--
219	8-(quinolin-7-ol)	N(CH ₃)	--

(continued)

Cpd	A	X	R _{4a}
230	6-(7-OH-quinolin-2(1H)-one)	N(CH ₃)	--
231	6-(7-OH-1-CH ₃ -quinolin-2(1H)-one)	N(CH ₃)	--
245	7-(6-OH-1-CH ₃ -quinolin-4(1H)-one)	N(CH ₃)	--
257	6-(7-OH-quinazolin-4(1H)-one)	N(CH ₃)	--
259	6-(7-OH-1-CH ₃ -3,4-dihydroquinolin-2(1H)-one)	N(CH ₃)	--
277	7-OH-1,3-(CH ₃) ₂ -quinazolin-6-yl-2,4(1H,3H)-dione	N(CH ₃)	--
278	6-OH-benzo[d]oxazol-5-yl-2(3H)-one	N(CH ₃)	--
279	2-CH ₃ -6-OH-2H-indazol-5-yl	N(CH ₃)	--
280	1-CH ₃ -6-OH-1H-indazol-5-yl	N(CH ₃)	--
281	7-(6-OH-2-CH ₃ -isoquinolin-1(2H)-one)	N(CH ₃)	--
282	7-(6-OH-2-CH ₂ CH ₃ -isoquinolin-1(2H)-one)	O	--

[0023] Another embodiment of the present description includes a compound of Formula (Ia) form thereof selected from a compound of Formula (Ia11) or a form thereof, wherein substituents A, X and R_{4a}, when present, are indicated in the table below; and, "--" indicates one or more A, X and R_{4a} substituents are not present:



(Ia11)


Cpd	A	X
420	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	N(CH ₃)
428	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
430	2,3-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
431	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
434	2-OCH ₃ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
435	4-(1H-pyrazol-4-yl)phenyl	O
437	2-F-4-(1H-pyrazol-4-yl)phenyl	O
438	4-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
440	2-F-4-OH-phenyl	N(CH ₃)
442	2-CH ₃ -2H-indazol-5-yl	N(CH ₃)
443	2-CH ₃ -2H-indazol-5-yl	O
444	4-Cl-2-OCH ₃ -phenyl	O
445	2-CH ₃ -pyrazolo[1,5-a]pyridin-3-yl	N(CH ₃)
446	imidazo[1,2-a]pyridin-6-yl	O
447	2-OCH ₃ -4-(1H-pyrazol-1-yl)phenyl	O
448	5-(1H-pyrazol-4-yl)thiophen-2-yl	O

EP 3 848 035 A1

(continued)

Cpd	A	X
449	5-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
450	4-(1H-pyrazol-4-yl)thiophen-2-yl	O
451	2-OH-4-[3,5-(CH ₃) ₂ -1H-pyrazol-4-yl]phenyl	O
452	2-F-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
453	2-OCH ₃ -4-OH-phenyl	O
454	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	O
455	2,4-(OH) ₂ -phenyl	O
456	2-Cl-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
457	5-amino-2-(1H-pyrazol-4-yl)pyrimidin-4-yl	O
458	2,6-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
464	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	O
465	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)

[0024] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia1) or a form thereof, wherein substituents A, X and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more A, X and R_{4a} substituents are not present:




(Ia1)

Cpd	A	X
420	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	N(CH ₃)
428	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
430	2,3-F ₂ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
431	2,5-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
434	2-OCH ₃ -4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
435	4-(1H-pyrazol-4-yl)phenyl	O
437	2-F-4-(1H-pyrazol-4-yl)phenyl	O
438	4-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
440	2-F-4-OH-phenyl	N(CH ₃)
442	2-CH ₃ -2H-indazol-5-yl	N(CH ₃)
443	2-CH ₃ -2H-indazol-5-yl	O
444	4-Cl-2-OCH ₃ -phenyl	O
445	2-CH ₃ -pyrazolo[1,5-a]pyridin-3-yl	N(CH ₃)
446	imidazo[1,2-a]pyridin-6-yl	O

(continued)

Cpd	A	X
447	2-OCH ₃ -4-(1H-pyrazol-1-yl)phenyl	O
448	5-(1H-pyrazol-4-yl)thiophen-2-yl	O
449	5-(1-CH ₃ -1H-pyrazol-4-yl)thiophen-2-yl	O
450	4-(1H-pyrazol-4-yl)thiophen-2-yl	O
451	2-OH-4-[3,5-(CH ₃) ₂ -1H-pyrazol-4-yl]phenyl	O
452	2-F-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
453	2-OCH ₃ -4-OH -phenyl	O
454	2-OCH ₃ -4-(4-NO ₂ -1H-pyrazol-1-yl)phenyl	O
455	2,4-(OH) ₂ -phenyl	O
456	2-Cl-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)
457	5-amino-2-(1H-pyrazol-4-yl)pyrimidin-4-yl	O
458	2,6-F ₂ -4-(1H-pyrazol-4-yl)phenyl	O
464	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	O
465	2-(CHF ₂)-4-(1H-pyrazol-4-yl)phenyl	N(CH ₃)

[0025] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia12) or a form thereof, wherein substituents X, R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and B substituents are not present:




(Ia12)

Cpd	R _{1a}	X	B
66	H	NH	azetidin-3-yl
82	OH	--	piperazin-1-yl
85	H	--	1,2,3,6-tetrahydropyridin-4-yl
86	OH	--	1,2,3,6-tetrahydropyridin-4-yl
87	OH	--	2,2,6,6-tetramethyl-(1,2,3,6-tetrahydropyridin-4-yl)
88	OH	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
89	OH	--	piperidin-4-yl
99	H	CH ₂	piperidin-4-yl


[0026] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia13) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not

present:



Cpd	X	R _{1a}	R _{4a}
26	N(CH ₃)	H	--
28	NH	H	--
31	O	H	--
90	O	OH	--
91	N(CH ₃)	OH	--
92	NH	OH	--
93	N(CH ₃)	O(CH ₂) ₃ NHCO ₂ C(CH ₃) ₃	--
94	N(CH ₃)	O(CH ₂) ₃ NH ₂	--
95	N(CH ₃)	O(CH ₂) ₃ NHCO ₂ CH ₃	--
96	N(CH ₃)	O(CH ₂) ₃ OH	--
97	N(CH ₃)	O(CH ₂) ₃ OCH ₃	--
98	O	O(CH ₂) ₃ -morpholin-4-yl	--
103	N(CH ₃)	CN	--
104	N(CH ₃)	CH ₂ -1-piperidinyI	--
105	N(CH ₃)	CH ₂ -pyrrolidin-1-yl	--
108	N(CH ₃)	OCH ₃	--
109	N(CH ₃)	OCH ₃	CH ₃
110	N(CH ₃)	3,6-dihydro-2H-pyran-4-yl	--
111	N(CH ₃)	tetrahydro-2H-pyran-4-yl	--
112	N(CH ₃)	CHF ₂	--
113	N(CH ₃)	OC(CH ₃) ₂ (CH ₂) ₂ OH	--
114	N(CH ₃)	O(CH ₂) ₂ C(CH ₃) ₂ OH	--


[0027] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia14) or a form thereof, wherein substituents X and B, when present, are indicated in the table below; and, "--" indicates that one or more X and B substituents are not present:



(Ia14)

Cpd	X	B
55	O	piperidin-4-yl
56	O	(2S,4R,6R)-2,6-(CH ₃) ₂ -piperidin-4-yl
57	O	2,6-(CH ₃) ₂ -piperidin-4-yl
58	O	pyrrolidin-3-yl
59	O	2-CH ₃ -piperidin-4-yl
60	OCH ₂	1H-pyrrolidin-3-yl
61	O	3-F-piperidin-4-yl
65	--	piperazin-1-yl
67	NH	azetidin-3-yl
68	--	3,5-(CF ₃) ₂ -piperazin-1-yl
69	--	7-CH ₃ -2,7-diazaspiro[4.4]non-2-yl
70	--	[1,4]diazepan-1-yl
71	--	4-CH ₂ CH ₂ OH-piperazin-1-yl
72	--	2,7-diazaspiro[3.5]non-7-yl
73	--	2,7-diazaspiro[3.5]non-7-yl
74	--	3-CH ₂ OH-piperazin-1-yl
75	--	1,7-diazaspiro[4.4]non-7-yl
76	--	4-NH ₂ -4-CH ₃ -piperidin-1-yl
77	--	3-N(CH ₃) ₂ -piperidin-1-yl
79	--	3,3-(CF ₃) ₂ -piperazin-1-yl
80	--	7-CH ₂ CH ₂ OH-2,7-diazaspiro[4.4]non-2-yl
83	--	1,2,3,6-tetrahydropyridin-4-yl
84	--	piperidin-4-yl
102	O	(6S)-6-[(S)-CH(OH)CH ₃]-2,2-(CH ₃) ₂ -piperidin-4-yl
133	O	2,2-(CH ₃) ₂ -piperidin-4-yl

[0028] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:




(Ia15)

Cpd	X	R _{1a}	R _{4a}
3	NH	H	--
7	N(CH ₃)	H	--
27	N(CH ₃)	Cl	CH ₃
29	NH	Cl	CH ₃
36	N(CH ₃)	OCH ₃	--
38	N(CH ₃)	F	--
39	N(CH ₃)	CN	--
42	N(CH ₃)	C(O)NHCH ₂ CH=CH ₂	--
43	N(CH ₃)	1H-pyrazol-1-yl	--
44	N(CH ₃)	5-CH ₃ -oxazol-2-yl	--
45	N(CH ₃)	4-CH ₂ OH-1H-pyrazole-1-yl	--
46	N(CH ₃)	1H-imidazole-1-yl	--
47	N(CH ₃)	4-NH ₂ -1H-pyrazol-1-yl	--
48	N(CH ₃)	1H-pyrazol-4-yl	--
49	N(CH ₃)	3-NH ₂ -1H-pyrazol-1-yl	--
50	N(CH ₃)	1-(CH ₂ CH ₂ -morpholin-4-yl)-1H-pyrazol-4-yl	--
51	N(CH ₃)	1-CH ₃ -1H-pyrazol-4-yl	--
52	N(CH ₃)	5-NH ₂ -1H-pyrazol-1-yl	--
54	N(CH ₂ CH ₂ OH)	1H-pyrazol-1-yl	--
62	O	1H-pyrazol-1-yl	CH ₃
63	O	1H-pyrazol-1-yl	--
64	O	1H-pyrazol-4-yl	--
78	NH	1H-pyrazol-1-yl	CH ₃
100	CH ₂	1H-pyrazol-1-yl	--
130	N(CH ₃)	Cl	--
131	NH	1H-pyrazol-1-yl	--
132	NH	CN	--
143	N(CH ₃)	1H-indazol-7-yl	--
157	CH ₂	1H-pyrazol-4-yl	--
168	N(CH ₃)	5-OCH ₃ -pyridin-3-yl	--
169	N(CH ₃)	5-pyridin-2-ol	--

(continued)

Cpd	X	R _{1a}	R _{4a}
170	N(CH ₃)	4-pyridin-2-yl	--
171	N(CH ₃)	6-OCH ₃ -pyridin-3-yl	--
172	N(CH ₃)	5-(3-CF ₃ -pyridin-2-yl)	--
173	N(CH ₃)	5-(1-CH ₃ -pyridin-2(1H)-one)	--
174	N(CH ₃)	4-(1-CH ₃ -pyridin-2(1H)-one)	--
175	N(CH ₃)	2-OCH ₃ -pyridin-4-yl	--
176	O	4-pyridin-2-yl	--
177	N(CH ₃)	6-N(CH ₃) ₂ -pyridin-3-yl	--
178	O	4-(1-CH ₃ -pyridin-2(1H)-one)	--
179	N(CH ₃)	pyrimidin-5-yl	--
180	N(CH ₃)	5-pyridin-3-ol	--
181	N(CH ₃)	4-(1-cyclopropyl-pyridin-2(1H)-one)	--
182	N(CH ₃)	1,2,3,6-tetrahydropyridin-4-yl	--
183	N(CH ₃)	cyclopent-1-en-1-yl	--
184	N(CH ₃)	3,6-dihydro-2H-pyran-4-yl	--
185	N(CH ₃)	imidazo[1,5-a]pyridin-7-yl	--
186	N(CH ₃)	imidazo[1,2-a]pyridin-7-yl	--
187	N(CH ₃)	2-CH ₃ -pyridin-4-yl	--
188	N(CH ₃)	1H-imidazol-2-yl	--
189	N(CH ₃)	1H-imidazol-4-yl	--
190	N(CH ₃)	imidazo[1,2-a]pyrazin-3-yl	--
191	N(CH ₃)	5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl	--
192	N(CH ₃)	4-CH ₃ -1H-imidazol-2-yl	--
193	N(CH ₃)	1-CH ₃ -1H-imidazol-4-yl	--
194	N(CH ₃)	1-CH ₃ -1H-imidazol-5-yl	--
195	N(CH ₃)	4-NO ₂ -1H-imidazol-2-yl	--
196	N(CH ₃)	2-CH ₃ -1H-imidazol-4-yl	--
197	N(CH ₃)	1,2-(CH ₃) ₂ -1H-imidazol-4-yl	--
198	N(CH ₃)	4-C(O)NH ₂ -1H-pyrazol-1-yl	--
206	N(CH ₃)	H	--


[0029] Another embodiment of the present description includes a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:



(Ia15)

Cpd	X	R _{1a}	R _{4a}
413	NH	1H-pyrazol-4-yl	--
414	O	1-CH ₃ -1H-pyrazol-4-yl	--
416	N(CH ₃)	5-CH ₃ -1H-pyrazol-4-yl	--
417	O	1H-imidazol-1-yl	--
418	O	5-CH ₃ -1H-pyrazol-4-yl	--
419	N(CH ₃)	4-NO ₂ -1H-pyrazol-1-yl	--
421	O	4-NH ₂ -1H-pyrazol-1-yl	--
423	O	4-NO ₂ -1H-pyrazol-1-yl	--
460	N(CH ₃)	1H-pyrazol-4-yl	--
461	O	1H-pyrazol-4-yl	--


[0030] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia15) or a form thereof, wherein substituents X, R_{1a} and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and R_{4a} substituents are not present:



(Ia15)


Cpd	X	R _{1a}
413	NH	1H-pyrazol-4-yl
414	O	1-CH ₃ -1H-pyrazol-4-yl
416	N(CH ₃)	5-CH ₃ -1H-pyrazol-4-yl
417	O	1H-imidazol-1-yl
418	O	5-CH ₃ -1H-pyrazol-4-yl
419	N(CH ₃)	4-NO ₂ -1H-pyrazol-1-yl
421	O	4-NH ₂ -1H-pyrazol-1-yl
423	O	4-NO ₂ -1H-pyrazol-1-yl
460	N(CH ₃)	1H-pyrazol-4-yl
461	O	1H-pyrazol-4-yl

[0031] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia16) or a form thereof, wherein substituents R_{1a} and R_{4a} , when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} and R_{4a} substituents are not present:




Cpd	R_{1a}	R_{4a}
81	1H-pyrazol-1-yl	--
199	1H-pyrazol-4-yl	$(CH_2)_2OH$
200	1H-pyrazol-4-yl	--
201	1H-pyrazol-4-yl	CH_3
202	4-(1- CH_3 -pyridin-2(1H)-one)	CH_3
203	4-(1- CH_3 -pyridin-2(1H)-one)	CH_3

[0032] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia17) or a form thereof, wherein substituent R_{1a} , when present, is indicated in the table below; and, "--" indicates that one or more R_{1a} substituents are not present:



Cpd	R_{1a}
204	1H-pyrazol-4-yl
205	4-(1- CH_3 -pyridin-2(1H)-one)


[0033] Another embodiment of the present description includes a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia18) or a form thereof, wherein substituents X and B, when present, are indicated in the table below; and, "--" indicates that one or more X and B substituents are not present:



(Ia18)

Cpd	X	B
411	N(CH ₃)	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
412	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
415	O	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
422	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
424	--	1,2,3,6-tetrahydropyridin-4-yl
425	--	1-CH ₃ CH ₂ -(1,2,3,6-tetrahydropyridin-4-yl)
426	N(CH ₃)	piperidin-4-yl
427	NH	piperidin-4-yl
429	--	8-azabicyclo[3.2.1]oct-2-en-3-yl
432	O	piperidin-4-yl
433	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
436	O	2,6-(CH ₃) ₂ -piperidin-4-yl
439	--	2,7-diazaspiro[3.5]non-2-yl
441	O	2,6-(CH ₃) ₂ -piperidin-4-yl
459	--	2,6-diazaspiro[3.4]oct-2-yl

[0034] Another embodiment of the method of the present description includes the use of a compound of Formula (Ia) or a form thereof selected from a compound of Formula (Ia18) or a form thereof, wherein substituents X, R_{1a} and B, when present, are indicated in the table below; and, "--" indicates that one or more X, R_{1a} and B substituents are not present:




(Ia18)

Cpd	X	B
411	N(CH ₃)	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
412	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
415	O	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl

(continued)

Cpd	X	B
422	--	1-CH ₃ -(1,2,3,6-tetrahydropyridin-4-yl)
424	--	1,2,3,6-tetrahydropyridin-4-yl
425	--	1-CH ₃ CH ₂ -(1,2,3,6-tetrahydropyridin-4-yl)
426	N(CH ₃)	piperidin-4-yl
427	NH	piperidin-4-yl
429	--	8-azabicyclo[3.2.1]oct-2-en-3-yl
432	O	piperidin-4-yl
433	NH	(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl
436	O	2,6-(CH ₃) ₂ -piperidin-4-yl
439	--	2,7-diazaspiro[3.5]non-2-yl
441	O	2,6-(CH ₃) ₂ -piperidin-4-yl
459	--	2,6-diazaspiro[3.4]oct-2-yl

[0035] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:



(Ib1)

Cpd	A
302	6-(naphthalen-2-ol)
320	6-(naphthalen-2,7-diol)
331	7-OCH ₃ -quinolin-6-yl
332	7-OH-quinolin-6-yl
337	2-CN-7-OCH ₃ -quinolin-6-yl
355	3-F-5-(1H-pyrazol-4-yl)-pyridin-2-yl
364	2-(6-OCH ₃ -3,4-dihydroisoquinolin-1(2H)-one)
392	6-OH-1-oxo-2,3-dihydro-1H-inden-5-yl
401	3-(4-OCH ₃ -1-CH ₃ -quinolin-2(1H)-one)
402	3-(4-OH-1-CH ₃ -quinolin-2(1H)-one)
403	3-(quinolin-2(1H)-one)
404	3-(1-OCH ₃ -quinolin-2(1H)-one)
408	5-CN-benzo[b]thiophen-2-yl


(continued)

Cpd	A
409	3-Cl-benzo[b]thiophen-2-yl

[0036] Another embodiment of the present description includes a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:

Cpd	A
462	3-(1H-pyrazol-4-yl)phenoxy
463	4-(1H-pyrazol-4-yl)phenoxy


[0037] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib1) or a form thereof, wherein substituent A is indicated in the table below:



(Ib1)

Cpd	A
462	3-(1H-pyrazol-4-yl)phenoxy
463	4-(1H-pyrazol-4-yl)phenoxy


[0038] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib2) or a form thereof, wherein substituent A is indicated in the table below:



(Ib2)

Cpd	A
321	6-naphthalen-2,7-diol


[0039] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib3) or a form thereof, wherein substituents R_{1a} , R_{1b} and B, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a} , R_{1b} and B substituents are not present:



(Ib3)

Cpd	R _{1a}	R _{1b}	B
329	1H-pyrazol-1-yl	OCH ₃	1,2,3,6-tetrahydropyridin-4-yl
330	1H-pyrazol-1-yl	OH	piperazin-1-yl
381	1H-pyrazol-1-yl	Cl	5-((3aR,6aR)-1-CH ₃ -hexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)
382	1H-pyrazol-1-yl	Cl	2-NHCH(CH ₃) ₂ -morpholin-4-yl
383	1H-pyrazol-1-yl	Cl	2-OCH ₃ -2,7-diazaspiro[4.5]decan-7-yl
385	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	5-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
394	1-CH ₃ -1H-pyrazol-4-yl	OH	5-((3aR,6aS)-5-CH ₃ -hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)
406	1H-pyrazol-1-yl	Cl	2,7-diazaspiro[4.5]decan-2-yl
407	1H-pyrazol-1-yl	Cl	(3R)-(3-(R)-CH ₂ OH)-piperazin-1-yl

[0040] Another embodiment of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib4) or a form thereof, wherein substituents R_{1a}, R_{1b}, R_{1c}, R_{1d} (each representative of the scope of R₁) and X, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a}, R_{1b}, R_{1c}, R_{1d} and X substituents are not present:



(Ib4)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
301	1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
305	1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
306	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
307	1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
308	4-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
309	5-pyridin-2-ol	OCH ₃	H	H	N(CH ₃)
310	5-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
311	1-CH ₃ -1H-pyrazol-4-yl	CH ₃	H	H	N(CH ₃)


(continued)

	Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
5	312	4-(1-CH ₃ -pyridin-2(1H)-one)	OCF ₃	H	H	N(CH ₃)
	313	3,5-(CH ₃) ₂ -1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)
	314	1-CH ₃ -1H-pyrazol-4-yl	CF ₃	H	H	N(CH ₃)
	315	1-CH ₃ -1H-pyrazol-4-yl	OH	H	H	N(CH ₃)
10	316	1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
	317	5-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	H	N(CH ₃)
	318	4-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	H	N(CH ₃)
15	319	5-pyridin-2-ol	OH	H	H	N(CH ₃)
	324	H	OH	1H-pyrazol-1-yl	H	N(CH ₃)
	325	1-CH ₃ -1H-pyrazol-4-yl	H	H	Cl	N(CH ₃)
	326	1-CH ₃ -1H-pyrazol-4-yl	OH	H	Cl	N(CH ₃)
20	327	1-CH ₃ -1H-pyrazol-4-yl	H	H	Cl	N(CH ₃)
	328	5-CH ₃ -oxazol-2-yl	OH	H	OCH ₃	N(CH ₃)
	333	CN	OCH ₃	H	H	N(CH ₃)
25	334	CN	F	H	H	N(CH ₃)
	335	CO ₂ CH ₃	F	H	H	N(CH ₃)
	336	3-NHCH ₃ -1H-pyrazol-1-yl	OCH ₃	H	H	N(CH ₃)
	338	4-(1-CH ₃ -pyridin-2(1H)-one)	OCH ₃	H	H	N(CH ₃)
30	339	4-(1-CH ₃ -pyridin-2(1H)-one)	Cl	H	H	N(CH ₃)
	340	1H-pyrazol-4-yl	Cl	H	H	N(CH ₃)
	341	4,5,6,7-tetrahydropyrazolo[1, 5-a] pyridin-3-yl	Cl	H	H	N(CH ₃)
35	343	1-CH ₃ -1H-pyrazol-4-yl	Cl	H	H	O
	344	6-OCH ₃ -pyridin-3-yl	Cl	H	H	N(CH ₃)
	345	6-NH ₂ -pyridin-3-yl	F	H	H	N(CH ₃)
	346	3-CH ₃ -1H-pyrazol-5-yl	F	H	H	N(CH ₃)
40	347	1H-pyrazol-5-yl	F	H	H	N(CH ₃)
	348	1H-pyrazol-4-yl	H	F	F	N(CH ₃)
	349	1H-pyrazol-5-yl	H	F	F	N(CH ₃)
45	350	1H-pyrazol-4-yl	F	F	H	N(CH ₃)
	351	1H-pyrazol-5-yl	F	F	H	N(CH ₃)
	352	1H-pyrazol-4-yl	F	H	F	N(CH ₃)
	354	1H-pyrazol-4-yl	Cl	F	H	N(CH ₃)
50	356	2-NH ₂ -pyrimidin-4-yl	Cl	H	H	N(CH ₃)
	357	H	Cl	2-NH ₂ -pyrimidin-4-yl	H	N(CH ₃)
	358	2,4-(CH ₃) ₂ -thiazol-5-yl	F	F	H	N(CH ₃)
55	359	2,4-(CH ₃) ₂ -thiazol-5-yl	H	F	F	N(CH ₃)
	360	4-(1-CH ₃ -pyridin-2(1H)-one)	OH	H	OCF ₃	N(CH ₃)
	361	1H-pyrazol-4-yl	OCH ₃	H	F	N(CH ₃)

(continued)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	X
363	1H-pyrazol-4-yl	OCH ₃	F	F	N(CH ₃)
365	1H-pyrazol-1-yl	Cl	H	H	N(CH ₃)
366	1H-1,2,3-triazol-1-yl	Cl	H	H	N(CH ₃)
367	2H-1,2,3-triazol-2-yl	Cl	H	H	N(CH ₃)
368	1H-1,2,4-triazol-1-yl	Cl	H	H	N(CH ₃)
369	3-NH ₂ -1H-pyrazol-1-yl	Cl	H	H	N(CH ₃)
371	1H-imidazol-1-yl	Cl	H	H	N(CH ₃)
372	1H-imidazol-1-yl	F	H	H	N(CH ₃)
373	1H-pyrazol-5-yl	OCH ₃	H	H	N(CH ₃)
374	2,4-(CH ₃) ₂ -thiazol-5-yl	OCH ₃	H	H	N(CH ₃)
375	pyridin-3-yl	OCH ₃	H	H	N(CH ₃)
376	1H-pyrazol-4-yl	F	H	H	N(CH ₃)
377	2-OCH ₃ -pyridin-4-yl	OCH ₃	H	H	N(CH ₃)
378	6-OCH ₃ -pyridin-3-yl	OCH ₃	H	H	N(CH ₃)
387	1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
388	5-(pyridin-2(1H)-one)	Cl	H	H	N(CH ₃)
389	3-NHCH ₃ -1H-pyrazol-1-yl	OH	H	H	N(CH ₃)
390	1H-pyrazol-4-yl	OH	H	F	N(CH ₃)
391	1H-pyrazol-4-yl	OH	F	F	N(CH ₃)
393	1H-pyrazol-4-yl	OH	H	H	N(CH ₃)
397	1H-pyrazol-4-yl	OH	H	Cl	N(CH ₃)
398	1H-pyrazol-1-yl	OCH ₃	H	H	CH ₂
410	1H-pyrazol-4-yl	OCH ₃	H	H	N(CH ₃)

[0041] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib5) or a form thereof, wherein substituents R_{1a}, R_{1b}, R_{1c}, R_{1d} (each representative of the scope of R₁) and R_{4a}, when present, are indicated in the table below; and, "--" indicates that one or more R_{1a}, R_{1b}, R_{1c}, R_{1d} and R_{4a} substituents are not present:




(Ib5)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	R _{4a}
353	1H-pyrazol-4-yl	F	F	H	--

(continued)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}	R _{4a}
362	1H-pyrazol-4-yl	OCH ₃	H	F	CH ₃
370	1H-imidazol-1-yl	Cl	H	H	CH ₃
379	1-CH ₃ -1H-pyrazol-4-yl	Cl	H	H	CH ₃
380	1H-pyrazol-4-yl	Cl	H	H	CH ₃
384	1H-pyrazol-4-yl	F	H	H	CH ₃
396	1H-pyrazol-4-yl	F	H	OH	--
405	1H-pyrazol-4-yl	Cl	H	H	--


[0042] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib6) or a form thereof, wherein substituents R_{1a}, R_{1b}, R_{1c} and R_{1d} (each representative of the scope of R_i), when present, are indicated in the table below; and, "--" indicates that one or more R_{1a}, R_{1b}, R_{1c} and R_{1d} substituents are not present:



(Ib6)

Cpd	R _{1a}	R _{1b}	R _{1c}	R _{1d}
386	1-CH ₃ -1H-pyrazol-4-yl	OCH ₃	H	H
395	1-CH ₃ -1H-pyrazol-4-yl	OH	H	H
399	1H-pyrazol-4-yl	H	F	F
400	1H-pyrazol-4-yl	OH	H	F


[0043] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib7) or a form thereof, wherein substituent R_{1b}, when present, is indicated in the table below:



(Ib7)

Cpd	R _{1b}
304	OCH ₃
322	OH


[0044] Another embodiment of the method of the present description includes the use of a compound of Formula (Ib) or a form thereof selected from a compound of Formula (Ib8) or a form thereof, wherein substituent R_{1b}, when present, is indicated in the table below:




(Ib8)

Cpd	R _{1b}
303	OCH ₃
323	OH


[0045] An embodiment of the use of a compound of Formula (I) or a form thereof includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject, selected from the group consisting of:




1















2




3




4


**5****6****7****8****9****10****11****12****13****14****15****16**




17




18




19




20




21




22




23




24




25















26




27




28







41




42




43




44




45




46




47




48




49



50




51




52

10




53




54

15

20




55




56

25

30




57




58

35

40



59




60


45

50


55




61




62




63




64




65




66




67




68




69




70




71




72




73




74




75




76




77




78




79




80




81




82




83




84




85




86




87











88




89




90







99




100




101




102




103




104




105




106




107




108




109




110




111




112




113




114




115




116




117




118




119




120




121




122




123




124




125




126




127




128




129




130




131




132




133




134




135




136




137













138




139




140







151




152




153




154




155




156




157













158














159




160









181




182




183




184




185




186




187




188




189




190




191




192




193




194




195




196




197




198




199




200




201




202




203




204




205




206




207















208



























209
















210














259




260




261




262




263




264




265




266




267




268




269




270




271




272




273




274




275




276




277




278




279



280




281




282


5




10




15




20




25




30




35




40





45














50




55









307




308




309




310




311




312




313




314




315




316




317




318




319











320












321




322









339




340




341




342




343











344




345




346







355




356




357




358




359




360




361




362




363




364




365




366




367




368




369




370




371




372




373









374




375




376







383




384




385




386




387




388




389




390




391



392



393




394

5

10

15




395

20

25

30




397

35

40


45




399

50


55











396



398




400




5

10




409

15




410

20




411

25




412

30




413

35




414

40




415

45




416

50













417


55



418




5




429

10




430

15




431

20




432

25




433

30




434

35




435

40




436

45




437

50




438


55




439




440




441




442




443




444




445




446




447




448




449




450




451




452




453




454




455




456




457




458




459




460




461




462



463













464














465

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.


[0046] In another aspect, the compound of Formula (I) or a form thereof includes a compound selected from the group consisting of:









431




432




433




434




435




436




437




438




439




440




441




442




443




444




445




446




447




448




449




450




451




452




453




454




455




456




457




458




459




460




461




462



463



464



465

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0047] Another embodiment of the use of a compound of Formula (I) or a form thereof includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof (wherein compound number (#1) indicates that the salt form was isolated) to the subject, selected from the group consisting of:

Cpd	Name
1	6-(naphthalen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
2	6-(benzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
3	2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)-pyridazin-3-yl)phenol
4	2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[b]-thiophene-5-carbonitrile
5	6-(quinolin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
6	3-(benzo[b]-thiophen-2-yl)-6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)pyridazine
7	2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)phenol
8	6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)naphthalen-2-ol
9	6-(benzo[b]-thiophen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
10	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
11	6-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
12	N-methyl-6-(quinolin-7-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
13	N-methyl-6-(quinolin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
14	6-(isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
15	6-(isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

(continued)

Cpd	Name
16	6-(imidazo[1,2-a]pyridin-6-yl-pyridazin-3-yl)-methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
5 17	N-methyl-6-(6-phenylpyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
18	6-(6-(1H-pyrrol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
19	6-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
20	methyl-(6-quinoxalin-2-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
10 21	methyl-(6-quinolin-3-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
22	N-methyl-6-(phthalazin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
23	6-(benzo[c][1,2,5]oxa-diazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
24	6-(benzo[d]thiazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
25	6-(2-methylbenzo[d]oxazol-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
15 26	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
27	5-chloro-2-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
28	3-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)pyridazin-3-yl)naphthalen-2-ol
29	5-chloro-2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)pyridazin-3-yl)phenol
30	4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
20 31	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
32 ¹	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol
33	2-fluoro-6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
34	3,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
25 35	4,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
36	5-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
37	4,5-difluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
38	5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
39	3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
30 40	1-allyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
41	6-(benzo[b]thiophen-2-yl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyridazin-3-amine
42	N-allyl-3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzamide
43	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
35 44	5-(5-methyl-oxazol-2-yl)-2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
45	5-(4-hydroxymethyl)-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
46	5-(1H-imidazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
47	5-(4-amino-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
40 48	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
49	5-(3-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
50	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-(2-morpholino-ethyl)-1H-pyrazol-4-yl)phenol
51	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
45 52	5-(5-amino-1H-pyrazol-1-yl)-2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)pyridazin-3-yl)phenol
53 ¹	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol
54	2-((6-((2-hydroxy-ethyl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)-5-pyrazol-1-yl)phenol
55	2-(6-(piperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
50 56	2-(6-(((2S,4R,6R)-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
57	2-(6-((-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
58	5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-yl-oxy)pyridazin-3-yl)phenol
59	2-(6-(((2S,4S)-2-methylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
60	(5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-ylmethoxy)pyridazin-3-yl)phenol
55 61	2-(6-((3-fluoropiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
62	2-(6-(1,2,2,6,6-pentamethyl-piperidin-4-yl-oxy)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
63	5-1H-pyrazol-1-yl-2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)-pyridazin-3-yl)phenol

(continued)

Cpd	Name
64	5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
5 65 ¹	2-(6-piperazin-1-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
66	3-(6-(azetidin-3-ylamino)-pyridazin-3-yl)naphthalen-2-ol
67	2-(6-(azetidin-3-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
68	2-(6-(3,5-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
10 69	2-(6-(7-methyl-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
70	2-(6-(1,4-diazepan-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
71	2-(6-(4-(2-hydroxyethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
72	2-(6-(3,6-diazabicyclo[3.2.1]octan-3-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
73	2-(6-(2,7-diazaspiro[3.5]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
15 74	2-(6-(3-(hydroxymethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
75	2-(6-(1,7-diazaspiro[4.4]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
76	2-(6-(4-amino-4-methylpiperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
77	2-(6-(3-(dimethylamino)piperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
20 78	2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
79	2-(6-(3,3-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
80	2-(6-(7-(2-hydroxyethyl)-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
81	2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
82 ¹	3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol
25 83	5-(1H-pyrazol-1-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol
84	2-(6-piperidin-4-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
85	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalen-2-ol
86 ¹	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
87	3-(6-(2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
30 88 ¹	3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
89 ¹	3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
90	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalene-2,7-diol
91	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
35 92	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
93	tert-butyl (3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)carbamate
94	7-(3-amino-propoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)naphthalen-2-ol
40 95	N-(3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)acetamide
96	7-(3-hydroxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
97	7-(3-methoxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
98	7-(2-morpholinoethoxy)-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
99	3-(6-(piperidin-4-ylmethyl)pyridazin-3-yl)naphthalen-2-ol
45 100	5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
101	3-methoxy-2-(6-(methyl(2,2,6-trimethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
102	2-(6-((6S)-6-((S)-1-hydroxyethyl)-2,2-dimethylpiperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
103	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2-naphthonitrile
50 104	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(piperidinylmethyl)naphthalen-2-ol
105	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(pyrrolidinylmethyl)naphthalen-2-ol
106	1-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
107	1-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
108	7-methoxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2-ol
55 109	7-methoxy-3-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
110	7-(3,6-dihydro-2H-pyran-4-yl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol

(continued)

Cpd	Name
5	111 3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(tetrahydro-2H-pyran-4-yl)naphthalene-2-ol
	112 7-(difluoromethyl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	113 7-((4-hydroxy-2-methylbutan-2-yl)oxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
10	114 7-(3-hydroxy-3-methylbutoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	115 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)benzene-1,3-diol
	116 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
15	117 5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(trifluoromethoxy)phenol
	118 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
	119 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
20	120 4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one
	121 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
25	122 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyridin-3-yl)phenol
	123 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyridine-3-yl)phenol
	124 5-(1-cyclopentyl-1H-pyrazol-4-yl)-3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
30	125 3',5-dimethoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-(1,1'-biphenyl)-3-ol
	126 3-(benzyloxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
	127 3-ethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
35	128 3-(cyclopropylmethoxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
	129 2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-benzo[d]imidazol-6-ol
	130 5-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	131 5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
40	132 3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
	133 2-(6-((2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	134 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
	135 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenol
45	136 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyrazin-3-yl)phenol
	137 4-(1H-indol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	138 4-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
50	139 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-3-yl)phenol
	140 4-(4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
	141 4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	142 4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol
	143 5-(1H-indazol-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
55	144 4-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	145 4-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	146 5-fluoro-4-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

(continued)

Cpd	Name
5	147 5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
	148 5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-5-yl)phenol
	149 6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one
	150 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol
10	151¹ 6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime
	152 5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-indene-1,6-diol
	153¹ 2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol
15	154¹ 9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol
	155 4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)benzamide
	156 4-(4-(hydroxymethyl)-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
20	157 5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
	158 6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	159 6-(1-(benzyloxy)isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	160¹ 3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
25	161¹ 4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one
	162¹ 4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
30	163¹ 5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	164¹ 3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
	165¹ 5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
35	166¹ 3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	167¹ 3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
	168 5-(5-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	169 5-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
40	170 4-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
	171 5-(6-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	172 5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-(trifluoromethyl)pyridin-2-ol
45	173 5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	174 4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	175 5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	176 4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol
50	177 5-(6-(dimethylamino)pyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	178 4-(3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	179 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyrimidin-5-yl)phenol
55	180 5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-3-ol
	181 1-cyclopropyl-4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one

(continued)

Cpd	Name
5	182 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)phenol
	183 5-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	184 5-(3,6-dihydro-2H-pyran-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	185 5-(imidazo[1,5-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
10	186 5-(imidazo[1,2-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	187 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methylpyridin-4-yl)phenol
	188 5-(1H-imidazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	189 5-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	190 5-(imidazo[1,2-a]pyrazin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
15	191 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl)phenol
	192 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-methyl-1H-imidazol-2-yl)phenol
	193 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-4-yl)phenol
	194 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-5-yl)phenol
20	195 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-imidazol-2-yl)phenol
	196 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methyl-1H-imidazol-4-yl)phenol
	197 5-(1,2-dimethyl-1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
25	198 1-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1H-pyrazole-4-carboxamide
	199 2-(6-((3aR,6aS)-5-(2-hydroxyethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	200 2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
30	201 2-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	202 4-(3-hydroxy-4-(6-(5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	203 4-(3-hydroxy-4-(6-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
35	204 2-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	205 4-(4-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-3-hydroxyphenyl)-1-methylpyridin-2(1H)-one
	206 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
	207 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
40	208 6-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	209 6-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)quinolin-7-ol
	210 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	211 7-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	212 7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
45	213 7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-6-ol
	214 7-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)isoquinolin-6-ol
	215 1-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	216 7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1,6-diol
50	217 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
	218 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
	219 8-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	220 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	221 2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
55	222 3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	223 3-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	224 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3-carbonitrile

(continued)

Cpd	Name
5	225 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(1-methyl-1H-imidazol-4-yl)quinolin-7-ol
	226¹ 3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	227 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol
	228 3-ethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
10	229 3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	230 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one
	231¹ 7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one
	232 4-methoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
15	233 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(pyrrolidin-1-yl)quinolin-7-ol
	234 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-morpholinoquinolin-7-ol
	235 4-(dimethylamino)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	236 4-ethoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
20	237 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1-methyl-1H-pyrazol-4-yl)quinolin-7-ol
	238¹ 4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	239¹ 7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
25	240 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
	241 3-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	242 3-bromo-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	243 3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
30	244 5-bromo-3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	245 6-hydroxy-1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-4(1H)-one
	246 2,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	247 2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
	248 3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
35	249 4-methoxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	250 4-(azetidin-1-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	251 7-hydroxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolone-4-carbonitrile
40	252 4-cyclopropyl-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
	253 4-(3,6-dihydro-2H-pyran-4-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	254¹ 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
45	255 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(oxetan-3-yl)quinolin-7-ol
	256¹ 4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol
	257 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-4(1H)-one
	258 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
50	259 7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3,4-dihydroquinolin-2(1H)-one
	260 2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol
	261 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
	262 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carbonitrile
55	263 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carbonitrile
	264 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carboxamide
	265 7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carboxamide

(continued)

Cpd	Name
5	266 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxamide methyl 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-
	267 carboxylate
	268 6-hydroxy-7-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
	269 7-hydroxy-6-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile
10	270 7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
	271 7-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)quinolin-6-ol
	272 1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
	273 1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	274 1,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
15	7-hydroxy-3-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-
	275 carbonitrile
	276 1-amino-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	7-hydroxy-1,3-dimethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazoline-2, 4
	277 (1H,3H)-dione
20	278 6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[d]oxazol-2(3H)-one
	279 2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2H-indazol-6-ol
	280 1-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-indazol-6-ol
	6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1
25	281 ¹ (2H)-one
	282 2-ethyl-6-hydroxy-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinolin-1(2H)-one
	283 1-ethoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	284 7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-1,6-diol
	285 7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-3-phenylisoquinolin-6-ol
30	286 3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	287 3-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	288 3-isopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	289 3-propyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
	290 3-isopropyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
35	291 3-methyl-7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
	292 6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	293 3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	294 3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
40	295 3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
	296 4-chloro-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	297 4-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	300 7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
45	5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-
	301 amine
	302 6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
	303 5-(2-methoxyquinolin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	304 5-(3-methoxy-naphthalen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
50	305 5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-
	306 thiadiazol-2-amine
	5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-
	307 amine
55	4-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-
	308 methylpyridin-2(1H)-one

(continued)

Cpd	Name
5	309 5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
	310 5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
	311 N-methyl-5-(2-methyl-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
10	312 1-methyl-4-(4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3-(trifluoromethoxy)phenyl)pyridin-2(1H)-one
	313 5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
15	314 5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	315 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
	316 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
20	317 5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
	318 4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
	319 5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
25	320 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
	321 3-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
	322¹ 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
	323 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2-ol
30	324 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-4-(1H-pyrazol-1-yl)phenol
	325 5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	326 3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
35	327 5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	328 3-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-methyloxazol-2-yl)phenol
	329 2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-(1,2,3,6-tetrahydropyridin-4-yl)-1,3,4-thiadiazole
40	330 2-(5-(piperazin-1-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
	331 5-(7-methoxyquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazole-2-amine
	332 6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-7-ol
	333 3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile
45	334 3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile
	335 methyl-3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzoate
	336 5-(2-methoxy-4-(3-(methylamino)-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
50	337 7-methoxy-6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinoline-2-carbonitrile
	338 4-(3-methoxy-4-(5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
	339 4-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
55	340 5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

(continued)

Cpd	Name
5	341 5-(2-chloro-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	342¹ N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	343 2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazole
10	344 5-(2-chloro-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	345 5-(4-(6-aminopyridin-3-yl)-2-fluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	346 5-(2-fluoro-4-(3-methyl-1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
15	347 5-(2-fluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	348 5-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
20	349 5-(2,3-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	350 5-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
25	351 5-(2,5-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	352 5-(2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	353 2-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
30	354 5-(2-chloro-5-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	355 5-(3-fluoro-5-(1H-pyrazol-4-yl)pyridin-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
35	356 5-(4-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	357 5-(5-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	358 5-(4-(2,4-dimethylthiazol-5-yl)-2,5-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
40	359 5-(4-(2,4-dimethylthiazol-5-yl)-2,3-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	360 4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one
45	361 5-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	362 2-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
50	363 5-(2,3-difluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	364 6-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3,4-dihydroisoquinolin-1-(2H)-one
	365 5-(2-chloro-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
55	366 5-(2-chloro-4-(1H-1,2,3-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

(continued)

Cpd	Name
5	367 5-(2-chloro-4-(2H-1,2,3-triazol-2-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	368 5-(2-chloro-4-(1H-1,2,4-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	369 5-(4-(3-amino-1H-pyrazol-1-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
10	370 2-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
	371 5-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
15	372 5-(2-fluoro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	373 5-(2-methoxy-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
20	374 5-(4-(2,4-dimethylthiazol-5-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	375 5-(2-methoxy-4-(pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	376 5-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
25	377 5-(2-methoxy-4-(2-methoxypyridin-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	378 5-(2-methoxy-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
30	379 2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
	380 2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
	381 2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-1,3,4-thiadiazole
35	382 1-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)morpholin-2-yl)-N,N-dimethylmethanamine
	383 2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2-methyl-2,7-diazaspiro[4.5]decan-7-yl)-1,3,4-thiadiazole
40	384 2-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
	385 2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
	386 2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
	387 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
45	388 5-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2(1H)-one
	389 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(3-(methylamino)-1H-pyrazol-1-yl)phenol
50	390 3-fluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	391 3,4-difluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	392 6-hydroxy-5-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1H-inden-1-one
55	393 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	394 2-(5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
	395 2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

(continued)

Cpd	Name
5	396¹ 3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	397 3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	398 2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)-1,3,4-thiadiazole
10	399 2-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
	400 2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-3-fluoro-5-(1H-pyrazol-4-yl)phenol
	401 4-methoxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
	402 4-hydroxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
15	403 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
	404 1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
	405¹ 2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
20	406¹ 2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole
	407¹ (R)-4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol
	408 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzo[b]thiophene-5-carbonitrile
	409 5-(3-chlorobenzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
25	410 5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	411¹ 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	412¹ 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	413¹ 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
30	414 5-(1-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	415¹ 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yloxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	416 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
	417 5-(1H-imidazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	418 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
35	419¹ 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol
	420 6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	421 5-(4-amino-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
40	422¹ 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	423 5-(4-nitro-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	424¹ 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
	425¹ 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	426¹ 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
45	427¹ 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	428¹ 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	429¹ 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	430¹ 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
50	431¹ 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	432 2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	433¹ 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	434¹ 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
55	435 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	436¹ 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	437¹ 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

(continued)

Cpd	Name
5	438 3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	439¹ 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	440 3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
	441 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol
	442 N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
10	443 2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole
	444 3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	445 N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	446 6-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]imidazo[1,2-a]pyridine
	447 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
15	448¹ 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	449 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	450¹ 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	451 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
20	452 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	453 3-methoxy-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	454 3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	455 4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]benzene-1,3-diol
	456¹ 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
25	457 2-(1H-pyrazol-4-yl)-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyrimidin-5-amine
	458¹ 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	459 2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	460¹ 3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol
	461 6-(1H-pyrazol-4-yl)-3-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyridin-2-ol
30	462¹ N,2,2,6,6-pentamethyl-N-[5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine
	463¹ N,2,2,6,6-pentamethyl-N-[5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine
	464¹ 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and
35	465¹ 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.	
40	[0048] In one aspect, the compound of Formula (I) or a form thereof includes a compound selected from the group consisting of:
	411¹ 2-[6-[(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	412¹ 2-[6-[(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	413¹ 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
45	414 5-(1-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	415¹ 2-[6-[(1 <i>R</i> ,5 <i>S</i>)-8-azabicyclo[3.2.1]oct-3-yloxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	416 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
	417 5-(1H-imidazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
50	418 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	419¹ 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol
	420 6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	421 5-(4-amino-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
55	422¹ 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	423 5-(4-nitro-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	424¹ 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol

(continued)

	425¹	2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	426¹	2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
5	427¹	2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	428¹	6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	429¹	2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	430¹	6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
10	431¹	3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	432	2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	433¹	2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	434¹	6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
15	435	3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	436¹	2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	437¹	3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	438	3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	439¹	2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
20	440	3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
	441	2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol
	442	N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	443	2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole
25	444	3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	445	N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	446	6-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]imidazo[1,2-a]pyridine
	447	3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	448¹	3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
30	449	3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	450¹	3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	451	5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	452	6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
35	453	3-methoxy-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
	454	3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	455	4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]benzene-1,3-diol
	456¹	6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	457	2-(1H-pyrazol-4-yl)-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyrimidin-5-amine
40	458¹	3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
	459	2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
	460¹	3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol
	461	6-(1H-pyrazol-4-yl)-3-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyridin-2-ol
45	462¹	N,2,2,6,6-pentamethyl-N-[5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine
	463¹	N,2,2,6,6-pentamethyl-N-[5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine
	464¹	3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and
	465¹	6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0049] Another embodiment of the use of a compound salt of Formula (I) or a form thereof includes a method of use of a compound salt of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject, selected from the group consisting of:

	Cpd	Name
5	32	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol hydrochloride
	53	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol hydrochloride
	65	2-(6-piperazin-1-yl-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol hydrochloride
10	82	3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
	86	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
	88	3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate
15	89	3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-dioltrifluoroacetate
	151	6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime hydrochloride
	153	2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol hydrochloride
20	154	9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol hydrochloride
	160	3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride
	161	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one hydrochloride
25	162	4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride
	163	5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride
	164	3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol hydrochloride
30	165	5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride
	166	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol hydrochloride
	167	3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol hydrochloride
40	226	3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol hydrochloride
	227	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol formate
	231	7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one hydrochloride
45	238	4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol formate
	239	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol hydrochloride
50	254	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol formate
	256	4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol formate
	281	6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one hydrochloride
55	322	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol hydrobromide

(continued)

	Cpd	Name
5	342	N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine hydrochloride
	396	3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol dihydrochloride
10	405	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole hydrochloride
	406	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole hydrochloride
	407	(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol hydrochloride
15	411	2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
	412	2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
20	413	5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol hydrochloride
	415	2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
	419	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
25	422	2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
	424	5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
30	425	2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
	426	2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
	427	2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
35	428	6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
	429	2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
40	430	6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
	431	3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
45	433	2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
	434	6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
	436	2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride
50	437	3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
	439	2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
55	448	3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
	450	3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride

(continued)

	Cpd	Name
5	456	6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
	458	3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
10	460	3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
	462	N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
	463	N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
15	464	3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
	465	6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
20	wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.	
	In one aspect, the compound salt of Formula (I) or a form thereof includes a compound salt selected from the group consisting of:	
25	411	2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
	412	2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
	413	5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol hydrochloride
30	415	2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
	419	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
	422	2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
35	424	5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
	425	2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
40	426	2-{6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
	427	2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
45	428	6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
	429	2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
50	430	6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
	431	3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
	433	2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol hydrochloride
55	434	6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
	436	2-{6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol trihydrochloride

(continued)

	Cpd	Name
5	437	3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
	439	2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
	448	3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
10	450	3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
	456	6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
15	458	3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
	460	3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
20	462	N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
	463	N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
	464	3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
25	465	6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0050] Another embodiment of the present description includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

[0051] Another embodiment of the present description includes a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject.

[0052] Another embodiment of the present description includes a use of the compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

[0053] Another embodiment of the present description includes a use of the compound salt of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound salt of Formula (I) or a form thereof to the subject.

Chemical Definitions

[0054] As used herein, the term "C₁₋₄alkyl" generally refers to saturated hydrocarbon radicals having from one to four carbon atoms in a straight or branched chain configuration, including, without limitation, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, and the like. In some embodiments, C₁₋₄alkyl includes C₁₋₃alkyl, C₁₋₂alkyl, and the like. A C₁₋₄alkyl radical may be optionally substituted where allowed by available valences.

[0055] As used herein, the term "C₂₋₆alkenyl" generally refers to partially unsaturated hydrocarbon radicals having from two to five carbon atoms in a straight or branched chain configuration and one or more carbon-carbon double bonds therein, including, without limitation, ethenyl, allyl, propenyl and the like. In some embodiments, C₂₋₆alkenyl includes C₂₋₄alkenyl, C₂₋₃alkenyl, and the like. A C₂₋₆alkenyl radical may be optionally substituted where allowed by available valences.

[0056] As used herein, the term "C₁₋₄alkoxy" generally refers to saturated hydrocarbon radicals having from one to four carbon atoms in a straight or branched chain configuration of the formula: -O-C₁₋₄alkyl, including, without limitation, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, and the like. In some embodiments, C₁₋₄alkoxy includes C₁₋₃alkoxy, C₁₋₂alkoxy and the like. A C₁₋₄alkoxy radical may be optionally substituted where

allowed by available valences.

[0057] As used herein, the term "C₃₋₁₄cycloalkyl" generally refers to a saturated monocyclic, bicyclic or polycyclic hydrocarbon radical, including, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, 1H-indanyl, indenyl, tetrahydro-naphthalenyl and the like. In some embodiments, C₃₋₁₄cycloalkyl includes C₃₋₁₀cycloalkyl, C₃₋₈cycloalkyl, C₃₋₇cycloalkyl, C₅₋₈cycloalkyl, C₉₋₁₀cycloalkyl and the like. A C₃₋₁₄cycloalkyl radical may be optionally substituted where allowed by available valences.

[0058] As used herein, the term "C₃₋₁₄cycloalkenyl" generally refers to a partially unsaturated monocyclic, bicyclic or polycyclic hydrocarbon radical having one or more chemically stable carbon-carbon double bonds therein, including, without limitation, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl and the like. In some embodiments, C₃₋₁₄cycloalkenyl includes C₃₋₇cycloalkenyl, C₃₋₈cycloalkenyl, C₅₋₈cycloalkenyl, C₃₋₁₀cycloalkenyl and the like. A C₃₋₁₄cycloalkenyl radical may be optionally substituted where allowed by available valences.

[0059] As used herein, the term "aryl" generally refers to a monocyclic, bicyclic or polycyclic aromatic carbon atom ring structure radical, including, without limitation, phenyl, naphthyl, anthracenyl, fluorenyl, azulenyl, phenanthrenyl and the like. An aryl radical may be optionally substituted where allowed by available valences.

[0060] As used herein, the term "heteroaryl" generally refers to a monocyclic, bicyclic or polycyclic aromatic carbon atom ring structure radical in which one or more carbon atom ring members have been replaced, where allowed by structural stability, with one or more heteroatoms, such as an O, S or N atom, including, without limitation, furanyl, thienyl (also referred to as thiophenyl), pyrrolyl, pyrazolyl, imidazolyl, isoxazolyl, isothiazolyl, oxazolyl, thiazolyl, triazolyl, oxadiazolyl, thiadiazolyl, tetrazolyl, pyranlyl, thiopyranlyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, indolyl, indazolyl, indolizynyl, benzofuranyl, benzothienyl, benzimidazolyl, benzothiazolyl, benzooxazolyl, 9H-puranyl, quinoxalynyl, isoindolyl, quinolynyl, isoquinolynyl, quinazolynyl, acridinyl, phthalazinyl, imidazo[1,2-a]pyridinyl, imidazo[1,5-a]pyridinyl, imidazo[5,1-a]isoquinolynyl, 1,4-dihydroindeno[1,2-c]-1H-pyrazolyl, 2,3-dihydro-1H-inden-1-one, 2,3-dihydro-1H-indenyl, 3,4-dihydroquinolin-2(1H)-one, 5,6-dihydroimidazo[5,1-a]isoquinolynyl, 8H-indeno[1,2-d]thiazolyl, benzo[c][1,2,5]oxadiazolyl, benzo[d]oxazol-2(3H)-one, quinolin-2(1H)-one, quinazolin-4(1H)-one, quinazoline-2,4(1H,3H)-dione, benzo[d]oxazolyl, pyrazolo[1,5-a]pyridinyl, and the like. A heteroaryl radical may be optionally substituted on a carbon or nitrogen atom ring member where allowed by available valences.

[0061] As used herein, the term "heterocyclyl" generally refers to a saturated or partially unsaturated monocyclic, bicyclic or polycyclic carbon atom ring structure radical in which one or more carbon atom ring members have been replaced, where allowed by structural stability, with a heteroatom, such as an O, S or N atom, including, without limitation, oxiranyl, oxetanyl, azetidynyl, dihydrofuranyl, tetrahydrofuranyl, dihydrothienyl, tetrahydrothienyl, pyrrolinyl, pyrrolidinyl, dihydropyrazolyl, pyrazolinyl, pyrazolidinyl, dihydroimidazolyl, imidazolynyl, imidazolidinyl, isoxazolynyl, isoxazolidinyl, isothiazolynyl, isothiazolidinyl, oxazolynyl, oxazolidinyl, thiazolynyl, thiazolidinyl, triazolynyl, triazolidinyl, oxadiazolynyl, oxadiazolidinyl, thiadiazolynyl, thiadiazolidinyl, tetrazolynyl, tetrazolidinyl, dihydro-2H-pyranlyl, dihydro-pyridinyl, tetrahydro-pyridinyl, 1,2,3,6-tetrahydropyridinyl, hexahydro-pyridinyl, dihydro-pyrimidinyl, tetrahydro-pyrimidinyl, 1,4,5,6-tetrahydropyrimidinyl, dihydro-pyrazinyl, tetrahydro-pyrazinyl, dihydro-pyridazinyl, tetrahydro-pyridazinyl, piperazinyl, piperidinyl, morpholinyl, thiomorpholinyl, dihydro-triazinyl, tetrahydro-triazinyl, hexahydro-triazinyl, 1,4-diazepanyl, dihydro-indolyl, indolynyl, tetrahydro-indolyl, dihydro-indazolyl, tetrahydro-indazolyl, dihydro-isoindolyl, dihydro-benzofuranyl, tetrahydro-benzofuranyl, dihydro-benzothienyl, tetrahydro-benzothienyl, dihydro-benzimidazolyl, tetrahydro-benzimidazolyl, dihydro-benzooxazolyl, 2,3-dihydrobenzo[d]oxazolyl, tetrahydro-benzooxazolyl, dihydro-benzooxazinyl, 3,4-dihydro-2H-benzo[b][1,4]oxazinyl, tetrahydro-benzooxazinyl, benzo[1,3]dioxolyl, benzo[1,4]dioxanyl, dihydro-puranyl, tetrahydro-puranyl, dihydro-quinolynyl, tetrahydro-quinolynyl, 1,2,3,4-tetrahydroquinolynyl, dihydro-isoquinolynyl, 3,4-dihydro-isoquinolin-(1H)-yl, tetrahydro-isoquinolynyl, 1,2,3,4-tetrahydroisoquinolynyl, dihydro-quinazolynyl, tetrahydro-quinazolynyl, dihydro-quinoxalynyl, tetrahydro-quinoxalynyl, 1,2,3,4-tetrahydroquinoxalynyl, 1,3-dioxolanyl, 2,5-dihydro-1H-pyrrolyl, 4,5-dihydro-1H-imidazolyl, tetrahydro-2H-pyranlyl, hexahydropyrrolo[3,4-b][1,4]oxazin-(2H)-yl, (4aR,7aS)-hexahydropyrrolo[3,4-b][1,4]oxazin-(4aH)-yl, 3,4-dihydro-2H-pyrrolo[3,2-b][1,4]oxazinyl, (cis)-octahydrocyclopenta[c]pyrrolyl, hexahydropyrrolo[3,4-b]pyrrol-(1H)-yl, (3aR,6aR)-hexahydropyrrolo[3,4-b]pyrrol-(1H)-yl, (3aR,6aS)-hexahydropyrrolo[3,4-b]pyrrol-(1H)-yl, 5H-pyrrolo[3,4-b]pyridin-(7H)-yl, 5,7-dihydro-6H-pyrrolo[3,4-b]pyridinyl, tetrahydro-1H-pyrrolo[3,4-b]pyridin-(2H,7H,7aH)-yl, hexahydro-1H-pyrrolo[3,4-b]pyridin-(2H)-yl, (4aR,7aR)-hexahydro-1H-pyrrolo[3,4-b]pyridin-(2H)-yl, octahydro-6H-pyrrolo[3,4-b]pyridinyl, 2,3,4,9-tetrahydro-1H-carbazolyl, 1,2,3,4-tetrahydropyrazino[1,2-a]indolyl, 2,3-dihydro-1H-pyrrolo[1,2-a]indolyl, (3aR,6aR)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, (3aR,4R,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, (3aR,4S,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, (3aR,5r,6aS)-hexahydrocyclopenta[c]pyrrol-(1H)-yl, 1,3-dihydro-2H-isoindolyl, octahydro-2H-isoindolyl, (3aS)-1,3,3a,4,5,6-hexahydro-2H-isoindolyl, (3aR,4R,7aS)-1H-isoindol-(3H,3aH,4H,5H,6H,7H,7aH)-yl, (3aR,7aS)-octahydro-2H-isoindolyl, (3aR,4R,7aS)-octahydro-2H-isoindolyl, (3aR,4S,7aS)-octahydro-2H-isoindolyl, 2,5-diazabicyclo[2.2.1]heptanyl, 2-azabicyclo[2.2.1]heptenyl, 3-azabicyclo[3.1.0]hexanyl, 3,6-diazabicyclo[3.1.0]hexanyl, (1R,5S)-3-azabicyclo[3.1.0]hexanyl, (1S,5R)-3-azabicyclo[3.2.0]heptanyl, 5-azaspiro[2.4]heptanyl, 2,6-diazaspiro[3.3]heptanyl, 2,5-diazaspiro[3.4]octanyl, 2,6-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,7-diazaspiro[4.4]nonanyl, 2-azaspiro[4.5]decanyl, 2,8-diazaspiro[4.5]decanyl, 3,6-diazabicyclo[3.2.1]octyl, 1,4-dihydroindeno[1,2-c]pyrazolyl, dihydropyranlyl, dihydropyridinyl, di-

hydroquinolinyl, 8H-indeno[1,2-d]thiazolyl, tetrahydroimidazo[1,2-a]pyridinyl, pyridin-2(1H)-one, (1R,5S)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl and the like. A heterocyclyl radical may be optionally substituted on a carbon or nitrogen atom ring member where allowed by available valences.

- [0062]** As used herein, the term "C₂₋₄alkenyl-amino-carbonyl" refers to a radical of the formula: -C(=O)-NH-C₂₋₄alkenyl.
- [0063]** As used herein, the term "C₁₋₄alkoxy-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-O-C₁₋₄alkyl.
- [0064]** As used herein, the term "C₁₋₄alkoxy-carbonyl" refers to a radical of the formula: -C(=O)-O-C₁₋₄alkyl.
- [0065]** As used herein, the term "C₁₋₄alkoxy-carbonyl-amino" refers to a radical of the formula: -NH-C(=O)-O-C₁₋₄alkyl.
- [0066]** As used herein, the term "C₁₋₄alkoxy-carbonyl-amino-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-NH-C(=O)-O-C₁₋₄alkyl.
- [0067]** As used herein, the term "C₁₋₄alkyl-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-C₁₋₄alkyl.
- [0068]** As used herein, the term "C₁₋₄alkyl-amino" refers to a radical of the formula: -NH-C₁₋₄alkyl.
- [0069]** As used herein, the term "(C₁₋₄alkyl)₂-amino" refers to a radical of the formula: -N(C₁₋₄alkyl)₂.
- [0070]** As used herein, the term "C₁₋₄alkyl-amino-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-NH-C₁₋₄alkyl.
- [0071]** As used herein, the term "(C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-N(C₁₋₄alkyl)₂.
- [0072]** As used herein, the term "C₁₋₄alkyl-amino-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-NH-C₁₋₄alkyl.
- [0073]** As used herein, the term "(C₁₋₄alkyl)₂-amino-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-N(C₁₋₄alkyl)₂.
- [0074]** As used herein, the term "C₁₋₄alkyl-amino-carbonyl" refers to a radical of the formula: -C(=O)-NH-C₁₋₄alkyl.
- [0075]** As used herein, the term "(C₁₋₄alkyl)₂-amino-carbonyl" refers to a radical of the formula: -C(=O)-N(C₁₋₄alkyl)₂.
- [0076]** As used herein, the term "C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-NH-C₁₋₄alkyl.
- [0077]** As used herein, the term "(C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-N(C₁₋₄alkyl)₂.
- [0078]** As used herein, the term "C₁₋₄alkyl-carbonyl" refers to a radical of the formula: -C(=O)-C₁₋₄alkyl.
- [0079]** As used herein, the term "C₁₋₄alkyl-carbonyl-amino" refers to a radical of the formula: -NH-C(=O)-C₁₋₄alkyl.
- [0080]** As used herein, the term "C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl.
- [0081]** As used herein, the term "C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl.
- [0082]** As used herein, the term "amino" refers to a radical of the formula: -NH₂.
- [0083]** As used herein, the term "amino-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-NH₂.
- [0084]** As used herein, the term "amino-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-NH₂.
- [0085]** As used herein, the term "amino-carbonyl" refers to a radical of the formula: -C(=O)-NH₂.
- [0086]** As used herein, the term "cyano" refers to a radical of the formula: -CN.
- [0087]** As used herein, the term "C₃₋₇cycloalkyl-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-C₃₋₇cycloalkyl.
- [0088]** As used herein, the term "halo-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-halo, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more halogen atoms. In some embodiments, halo-C₁₋₄alkoxy includes halo-C₁₋₆alkoxy, halo-C₁₋₄alkoxy and the like.
- [0089]** As used herein, the term "halo-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-halo, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more halogen atoms. In some embodiments, halo-C₁₋₄alkyl includes halo-C₁₋₆alkyl, halo-C₁₋₄alkyl and the like.
- [0090]** As used herein, the term "heteroaryl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-heteroaryl.
- [0091]** As used herein, the term "heteroaryl-C₁₋₄alkyl-amino" refers to a radical of the formula: -NH-C₁₋₄alkyl-heteroaryl.
- [0092]** As used herein, the term "heteroaryl-C₁₋₄alkyl-amino-carbonyl" refers to a radical of the formula: -C(=O)-NH-C₁₋₄alkyl-heteroaryl.
- [0093]** As used herein, the term "heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-C(=O)-NH-C₁₋₄alkyl-heteroaryl.
- [0094]** As used herein, the term "heteroaryl-C₁₋₄alkyl-carbonyl-amino" refers to a radical of the formula: -NH-C(=O)-C₁₋₄alkyl-heteroaryl.
- [0095]** As used herein, the term "heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-NH-C(=O)-C₁₋₄alkyl-heteroaryl.
- [0096]** As used herein, the term "heterocyclyl-C₁₋₄alkoxy" refers to a radical of the formula: -C₁₋₄alkoxy-heterocyclyl.
- [0097]** As used herein, the term "heterocyclyl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-heterocyclyl.
- [0098]** As used herein, the term "hydroxyl" refers to a radical of the formula: -OH.
- [0099]** As used herein, the term "hydroxyl-C₁₋₄alkoxy" refers to a radical of the formula: -O-C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more hydroxy

radicals.

[0100] As used herein, the term "hydroxyl-C₁₋₄alkyl" refers to a radical of the formula: -C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more hydroxy radicals.

[0101] As used herein, the term "hydroxyl-C₁₋₄alkyl-amino" refers to a radical of the formula: -NH-C₁₋₄alkyl-OH, wherein C₁₋₄alkyl may be partially or completely substituted where allowed by available valences with one or more hydroxyl radicals.

[0102] As used herein, the term "hydroxyl-imino" refers to the =NOH radical of the formula: C(=NOH).

[0103] As used herein, the term "oxo" refers to the radical of the formula: C=O.

[0104] As used herein, the term "phenyl-C₁₋₄alkoxy" refers to a radical of the formula: -C₁₋₄alkoxy-phenyl.

[0105] As used herein, the term "substituent" means positional variables on the atoms of a core molecule that are substituted at a designated atom position, replacing one or more hydrogens on the designated atom, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. A person of ordinary skill in the art should note that any carbon as well as heteroatom with valences that appear to be unsatisfied as described or shown herein is assumed to have a sufficient number of hydrogen atom(s) to satisfy the valences described or shown. In certain instances one or more substituents having a double bond (e.g., "oxo" or "=O") as the point of attachment may be described, shown or listed herein within a substituent group, wherein the structure may only show a single bond as the point of attachment to the core structure of Formula (I). A person of ordinary skill in the art would understand that, while only a single bond is shown, a double bond is intended for those substituents.

[0106] As used herein, the term "and the like," with reference to the definitions of chemical terms provided herein, means that variations in chemical structures that could be expected by one skilled in the art include, without limitation, isomers (including chain, branching or positional structural isomers), hydration of ring systems (including saturation or partial unsaturation of monocyclic, bicyclic or polycyclic ring structures) and all other variations where allowed by available valences which result in a stable compound.

[0107] For the purposes of this description, where one or more substituent variables for a compound of Formula (I) or a form thereof encompass functionalities incorporated into a compound of Formula (I), each functionality appearing at any location within the disclosed compound may be independently selected, and as appropriate, independently and/or optionally substituted.

[0108] As used herein, the terms "independently selected," or "each selected" refer to functional variables in a substituent list that may occur more than once on the structure of Formula (I), the pattern of substitution at each occurrence is independent of the pattern at any other occurrence. Further, the use of a generic substituent variable on any formula or structure for a compound described herein is understood to include the replacement of the generic substituent with species substituents that are included within the particular genus, e.g., aryl may be replaced with phenyl or naphthalenyl and the like, and that the resulting compound is to be included within the scope of the compounds described herein.

[0109] As used herein, the terms "each instance of" or "in each instance, when present," when used preceding a phrase such as "...C₃₋₁₄cycloalkyl, C₃₋₁₄cycloalkyl-C₁₋₄alkyl, aryl, aryl-C₁₋₄alkyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heterocyclyl and heterocyclyl-C₁₋₄alkyl," are intended to refer to the C₃₋₁₄cycloalkyl, aryl, heteroaryl and heterocyclyl ring systems when each are present either alone or as a substituent.

[0110] As used herein, the term "optionally substituted" means optional substitution with the specified substituent variables, groups, radicals or moieties.

Compound Forms

[0111] As used herein, the term "form" means a compound of Formula (I) having a form selected from the group consisting of a free acid, free base, prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

[0112] In certain embodiments described herein, the form of the compound of Formula (I) is a free acid, free base or salt thereof.

[0113] In certain embodiments described herein, the form of the compound of Formula (I) is a salt thereof.

[0114] In certain embodiments described herein, the form of the compound of Formula (I) is an isotopologue thereof.

[0115] In certain embodiments described herein, the form of the compound of Formula (I) is a stereoisomer, racemate, enantiomer or diastereomer thereof.

[0116] In certain embodiments described herein, the form of the compound of Formula (I) is a tautomer thereof.

[0117] In certain embodiments described herein, the form of the compound of Formula (I) is a pharmaceutically acceptable form.

[0118] In certain embodiments described herein, the compound of Formula (I) or a form thereof is isolated for use.

[0119] As used herein, the term "isolated" means the physical state of a compound of Formula (I) or a form thereof after being isolated and/or purified from a synthetic process (e.g., from a reaction mixture) or natural source or combination

thereof according to an isolation or purification process or processes described herein or which are well known to the skilled artisan (e.g., chromatography, recrystallization and the like) in sufficient purity to be characterized by standard analytical techniques described herein or well known to the skilled artisan.

[0120] As used herein, the term "protected" means that a functional group in a compound of Formula (I) or a form thereof is in a form modified to preclude undesired side reactions at the protected site when the compound is subjected to a reaction. Suitable protecting groups will be recognized by those with ordinary skill in the art as well as by reference to standard textbooks such as, for example, T.W. Greene et al, *Protective Groups in organic Synthesis* (1991), Wiley, New York. Such functional groups include hydroxy, phenol, amino and carboxylic acid. Suitable protecting groups for hydroxy or phenol include trialkylsilyl or diarylalkylsilyl (e.g., t-butyltrimethylsilyl, t-butylphenylsilyl or trimethylsilyl), tetrahydropyranyl, benzyl, substituted benzyl, methyl, methoxymethanol, and the like. Suitable protecting groups for amino, amidino and guanidino include t-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for carboxylic acid include alkyl, aryl or arylalkyl esters. In certain instances, the protecting group may also be a polymer resin, such as a Wang resin or a 2-chlorotriethylchloride resin. Protecting groups may be added or removed in accordance with standard techniques, which are well-known to those skilled in the art and as described herein. It will also be appreciated by those skilled in the art, although such protected derivatives of compounds described herein may not possess pharmacological activity as such, they may be administered to a subject and thereafter metabolized in the body to form compounds described herein which are pharmacologically active. Such derivatives may therefore be described as "prodrugs". All prodrugs of compounds described herein are included within the scope of the use described herein.

[0121] As used herein, the term "prodrug" means a form of an instant compound (e.g., a drug precursor) that is transformed *in vivo* to yield an active compound of Formula (I) or a form thereof. The transformation may occur by various mechanisms (e.g., by metabolic and/or non-metabolic chemical processes), such as, for example, by hydrolysis and/or metabolism in blood, liver and/or other organs and tissues. A discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, "Pro-drugs as Novel Delivery Systems," Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

[0122] In one example, when a compound of Formula (I) or a form thereof contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a functional group such as alkyl and the like. In another example, when a compound of Formula (I) or a form thereof contains a hydroxyl functional group, a prodrug form can be prepared by replacing the hydrogen atom of the hydroxyl with another functional group such as alkyl, alkylcarbonyl or a phosphonate ester and the like. In another example, when a compound of Formula (I) or a form thereof contains an amine functional group, a prodrug form can be prepared by replacing one or more amine hydrogen atoms with a functional group such as alkyl or substituted carbonyl. Pharmaceutically acceptable prodrugs of compounds of Formula (I) or a form thereof include those compounds substituted with one or more of the following groups: carboxylic acid esters, sulfonate esters, amino acid esters, phosphonate esters and mono-, di- or triphosphate esters or alkyl substituents, where appropriate. As described herein, it is understood by a person of ordinary skill in the art that one or more of such substituents may be used to provide a compound of Formula (I) or a form thereof as a prodrug.

[0123] One or more compounds described herein may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and the description herein is intended to embrace both solvated and unsolvated forms.

[0124] As used herein, the term "solvate" means a physical association of a compound described herein with one or more solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. As used herein, "solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanlates, methanlates, and the like.

[0125] As used herein, the term "hydrate" means a solvate wherein the solvent molecule is water.

[0126] The compounds of Formula (I) can form salts, which are intended to be included within the scope of this description. Reference to a compound of Formula (I) or a form thereof herein is understood to include reference to salt forms thereof, unless otherwise indicated. The term "salt(s)", as employed herein, denotes acidic salts formed with inorganic and/or organic acids, as well as basic salts formed with inorganic and/or organic bases. In addition, when a compound of Formula (I) or a form thereof contains both a basic moiety, such as, without limitation an amine moiety, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein.

[0127] The term "pharmaceutically acceptable salt(s)", as used herein, means those salts of compounds described herein that are safe and effective (*i.e.*, non-toxic, physiologically acceptable) for use in mammals and that possess biological activity, although other salts are also useful. Salts of the compounds of the Formula (I) may be formed, for example, by reacting a compound of Formula (I) or a form thereof with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

[0128] Pharmaceutically acceptable salts include one or more salts of acidic or basic groups present in compounds described herein. Embodiments of acid addition salts include, and are not limited to, acetate, ascorbate, benzoate, benzenesulfonate, bisulfate, bitartrate, borate, bromide, butyrate, chloride, citrate, camphorate, camphorsulfonate, ethanesulfonate, formate, fumarate, gentisinate, gluconate, glucuronate, glutamate, iodide, isonicotinate, lactate, maleate, methanesulfonate, naphthalenesulfonate, nitrate, oxalate, pamoate, pantothenate, phosphate, propionate, saccharate, salicylate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate (also known as tosylate), trifluoroacetate salts and the like. Certain embodiments of acid addition salts include chloride, dichloride, trichloride, bromide, acetate, formate or trifluoroacetate salts.

[0129] Additionally, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by P. Stahl et al, Camille G. (eds.) Handbook of Pharmaceutical Salts. Properties, Selection and Use. (2002) Zurich: Wiley-VCH; S. Berge et al, Journal of Pharmaceutical Sciences (1977) 66(1) 1-19; P. Gould, International J. of Pharmaceutics (1986) 33, 201-217; Anderson et al, The Practice of Medicinal Chemistry (1996), Academic Press, New York; and in The Orange Book (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference thereto.

[0130] Suitable basic salts include, but are not limited to, aluminum, ammonium, calcium, lithium, magnesium, potassium, sodium and zinc salts.

[0131] All such acid salts and base salts are intended to be included within the scope of pharmaceutically acceptable salts as described herein. In addition, all such acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of this description.

[0132] Compounds of Formula (I) and forms thereof, may further exist in a tautomeric form. All such tautomeric forms are contemplated and intended to be included within the scope of the compounds of Formula (I) or a form thereof as described herein.

[0133] The compounds of Formula (I) or a form thereof may contain asymmetric or chiral centers, and, therefore, exist in different stereoisomeric forms. The present description is intended to include all stereoisomeric forms of the compounds of Formula (I) as well as mixtures thereof, including racemic mixtures.

[0134] The compounds described herein may include one or more chiral centers, and as such may exist as racemic mixtures (*R/S*) or as substantially pure enantiomers and diastereomers. The compounds may also exist as substantially pure (*R*) or (*S*) enantiomers (when one chiral center is present). In one embodiment, the compounds described herein are (*S*) isomers and may exist as enantiomerically pure compositions substantially comprising only the (*S*) isomer. In another embodiment, the compounds described herein are (*R*) isomers and may exist as enantiomerically pure compositions substantially comprising only the (*R*) isomer. As one of skill in the art will recognize, when more than one chiral center is present, the compounds described herein may also exist as a (*R,R*), (*R,S*), (*S,R*) or (*S,S*) isomer, as defined by IUPAC Nomenclature Recommendations.

[0135] As used herein, the term "substantially pure" refers to compounds consisting substantially of a single isomer in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100% of the single isomer.

[0136] In one aspect of the description, a compound of Formula (I) or a form thereof is a substantially pure (*S*) enantiomer form present in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100%.

[0137] In one aspect of the description, a compound of Formula (I) or a form thereof is a substantially pure (*R*) enantiomer form present in an amount greater than or equal to 90%, in an amount greater than or equal to 92%, in an amount greater than or equal to 95%, in an amount greater than or equal to 98%, in an amount greater than or equal to 99%, or in an amount equal to 100%.

[0138] As used herein, a "racemate" is any mixture of isometric forms that are not "enantiomerically pure", including mixtures such as, without limitation, in a ratio of about 50/50, about 60/40, about 70/30, or about 80/20.

[0139] In addition, the present description embraces all geometric and positional isomers. For example, if a compound of Formula (I) or a form thereof incorporates a double bond or a fused ring, both the *cis*- and *trans*-forms, as well as mixtures, are embraced within the scope of the description. Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods well known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by use of chiral HPLC column or other chromatographic methods known to those skilled in the art. Enantiomers can also be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Also, some of the compounds of Formula (I) may be atropisomers (e.g., substituted biaryls) and are considered as part of this description.

[0140] All stereoisomers (for example, geometric isomers, optical isomers and the like) of the present compounds

(including those of the salts, solvates, esters and prodrugs of the compounds as well as the salts, solvates and esters of the prodrugs), such as those which may exist due to asymmetric carbons on various substituents, including enantiomeric forms (which may exist even in the absence of asymmetric carbons), rotameric forms, atropisomers, and diastereomeric forms, are contemplated within the scope of this description, as are positional isomers (such as, for example, 4-pyridyl and 3-pyridyl). Individual stereoisomers of the compounds described herein may, for example, be substantially free of other isomers, or may be present in a racemic mixture, as described supra.

[0141] The use of the terms "salt", "solvate", "ester", "prodrug" and the like, is intended to equally apply to the salt, solvate, ester and prodrug of enantiomers, stereoisomers, rotamers, tautomers, positional isomers, racemates or isotopologues of the instant compounds.

[0142] The term "isotopologue" refers to isotopically-enriched compounds described herein which are identical to those recited herein, but for the fact that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds described herein include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{35}Cl and ^{36}Cl , respectively, each of which are also within the scope of this description.

[0143] Certain isotopically-enriched compounds described herein (e.g., those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (i.e., ^3H) and carbon-14 (i.e., ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (i.e., ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (e.g., increased in vivo half-life or reduced dosage requirements) and hence may be preferred in some circumstances.

[0144] Polymorphic crystalline and amorphous forms of the compounds of Formula (I) and of the salts, solvates, hydrates, esters and prodrugs of the compounds of Formula (I) are further intended to be included in the present description.

Compound Uses

[0145] The present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound or a form thereof to the subject.

[0146] The present description further relates to use of the compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof.

[0147] The present description further relates to use of the compound of Formula (I) or a form thereof having activity toward HD

[0148] The present description further relates to use of the compound of Formula (I) or a form thereof in a combination therapy to provide additive or synergistic activity, thus enabling the development of a combination product for treating or ameliorating HD

[0149] In addition to monotherapeutic use, the instant compounds are useful in a combination therapy with current standard of agents, having additive or synergistic activity with one or more known agents.

[0150] A combination therapy comprising compounds described herein in combination with one or more known drugs may be used to treat HD regardless of whether HD is responsive to the known drug.

[0151] Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a combination therapy for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s).

[0152] Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a combination therapy for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s).

[0153] In an embodiment of a use or method provided herein, compounds of Formula (I) or a form thereof used in combination with one or more additional agents can be administered to a subject or contacted with a subject or patient cell(s) prior to, concurrently with, or subsequent to administering to the subject or patient or contacting the cell with an additional agent(s). A compound(s) of Formula (I) or a form thereof and an additional agent(s) can be administered to a subject or contacted with a cell in single composition or different compositions. In a specific embodiment, a compound(s) of Formula (I) or a form thereof is used in combination with gene therapy to inhibit HTT expression (using, e.g., viral delivery vectors) or the administration of another small molecule HTT inhibitor. In another specific embodiment, a compound(s) of Formula (I) or a form thereof are used in combination with cell replacement using differentiated non-mutant HTT stem cells. In another specific embodiment, a compound(s) of Formula (I) or a form thereof are used in combination with cell replacement using differentiated HTT stem cells.

[0154] In one embodiment, provided herein is the use of compounds of Formula (I) or a form thereof in combination with supportive standard of care therapies, including palliative care.

[0155] An embodiment of the present description includes the use of a compound of Formula (I) or a form thereof in the preparation of a kit comprising the compound of Formula (I) or a form thereof and instructions for administering an effective amount of the compound of Formula (I) or a form thereof and an effective amount of one or more agent(s) in a combination therapy for treating or ameliorating HD in a subject in need thereof.

[0156] Accordingly, the present description relates to use of a compound of Formula (I) or a form thereof for treating or ameliorating HD. In accordance with the use of the present description, compounds that are useful in selectively treating or ameliorating HD, have been identified and use of these compounds for treating or ameliorating HD has been provided.

[0157] One embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof to the subject.

[0158] One embodiment of the use of the present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound to the subject.

[0159] An embodiment of the use of the present description relates to a method of use of a compound of Formula (I) or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound to the subject.

[0160] An embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof in the manufacture of a medicament for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the medicament to the subject.

[0161] An embodiment of the use of the present description relates to use of a compound of Formula (I) or a form thereof in the preparation of a kit comprising the compound of Formula (I) or a form thereof and instructions for administering the compound for treating or ameliorating HD in a subject in need thereof.

[0162] In one respect, for each of such embodiments, the subject is treatment naive. In another respect, for each of such embodiments, the subject is not treatment naive.

[0163] As used herein, the term "treating" refers to: (i) preventing a disease, disorder or condition from occurring in a subject that may be predisposed to the disease, disorder and/or condition but has not yet been diagnosed as having the disease, disorder and/or condition; (ii) inhibiting a disease, disorder or condition, *i.e.*, arresting the development thereof; and/or (iii) relieving a disease, disorder or condition, *i.e.*, causing regression of the disease, disorder and/or condition.

[0164] As used herein, the term "subject" refers to an animal or any living organism having sensation and the power of voluntary movement, and which requires oxygen and organic food. Nonlimiting examples include members of the human, primate, equine, porcine, bovine, murine, rattus, canine and feline specie. In some embodiments, the subject is a mammal or a warm-blooded vertebrate animal. In other embodiments, the subject is a human. As used herein, the term "patient" may be used interchangeably with "subject" and "human".

[0165] As used herein, the terms "effective amount" or "therapeutically effective amount" mean an amount of compound of Formula (I) or a form, composition or medicament thereof effective in inhibiting the above-noted diseases and thus producing the desired therapeutic, ameliorative, inhibitory or preventative effect in a subject in need thereof.

[0166] The dose administered to achieve an effective target plasma concentration may also be administered based upon the weight of the subject or patient. Doses administered on a weight basis may be in the range of about 0.001 mg/kg/day to about 3500 mg/kg/day, or about 0.001 mg/kg/day to about 3000 mg/kg/day, or about 0.001 mg/kg/day to about 2500 mg/kg/day, or about 0.001 mg/kg/day to about 2000 mg/kg/day, or about 0.001 mg/kg/day to about 1500 mg/kg/day, or about 0.001 mg/kg/day to about 1000 mg/kg/day, or about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.001 mg/kg/day to about 250 mg/kg/day, or about 0.001 mg/kg/day to about 100 mg/kg/day, or about 0.001 mg/kg/day to about 75 mg/kg/day, or about 0.001 mg/kg/day to about 50 mg/kg/day, or about 0.001 mg/kg/day to about 40 mg/kg/day, or about 0.001 mg/kg/day to about 30 mg/kg/day, or about 0.001 mg/kg/day to about 20 mg/kg/day, or about 0.001 mg/kg/day to about 10 mg/kg/day, or about 0.01 mg/kg/day to about 2000 mg/kg/day, or about 0.01 mg/kg/day to about 1500 mg/kg/day, or about 0.01 mg/kg/day to about 1000 mg/kg/day, or about 0.01 mg/kg/day to about 600 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 300 mg/kg/day, or about 0.015 mg/kg/day to about 200 mg/kg/day, or about 0.02 mg/kg/day to about 100 mg/kg/day, or about 0.025 mg/kg/day to about 100 mg/kg/day, or about 0.03 mg/kg/day to about 100 mg/kg/day, wherein said amount is orally administered once (once in approximately a 24 hour period), twice (once in approximately a 12 hour period) or thrice (once in approximately an 8 hour period) daily according to subject weight.

[0167] In certain embodiments, the effective amount will be in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.1 mg to about 500 mg/kg/day, or about 1.0 mg/day to about 500 mg/kg/day, in single, divided, or a continuous dose for a patient or subject having a weight in a range of between about 40 to about 200 kg (which dose may be adjusted for patients or subjects above or below this range, particularly children under 40 kg). The typical adult subject is expected to have a median weight in a range of

about 70 kg.

[0168] In another embodiment, where daily doses are adjusted based upon the weight of the subject or patient, compounds described herein may be formulated for delivery at about 0.02, 0.025, 0.03, 0.05, 0.06, 0.075, 0.08, 0.09, 0.10, 0.20, 0.25, 0.30, 0.50, 0.60, 0.75, 0.80, 0.90, 1.0, 1.10, 1.20, 1.25, 1.50, 1.75, 2.0, 3.0, 5.0, 10, 20, 30, 40, 50, 100, 150, 200, 250, 300, 400 or 500 mg/kg/day. Daily doses adjusted based upon the weight of the subject or patient may be administered as a single, divided, or continuous dose. In embodiments where a dose of compound is given more than once per day, the dose may be administered twice, thrice, or more times per day.

[0169] Within the scope of the present description, the "effective amount" of a compound of Formula (I) or a form thereof for use in the manufacture of a medicament, for use in the preparation of a pharmaceutical kit or in a method of use for treating or ameliorating HD in a subject in need thereof is intended to include an amount in a range of from about 0.001 mg/kg/day to about 3500 mg/kg/day, or about 0.001 mg/kg/day to about 3000 mg/kg/day, or about 0.001 mg/kg/day to about 2500 mg/kg/day, or about 0.001 mg/kg/day to about 2000 mg/kg/day, or about 0.001 mg/kg/day to about 1500 mg/kg/day, or about 0.001 mg/kg/day to about 1000 mg/kg/day, or about 0.001 mg/kg/day to about 500 mg/kg/day, or about 0.001 mg/kg/day to about 250 mg/kg/day, or about 0.001 mg/kg/day to about 100 mg/kg/day, or about 0.001 mg/kg/day to about 75 mg/kg/day, or about 0.001 mg/kg/day to about 50 mg/kg/day, or about 0.001 mg/kg/day to about 40 mg/kg/day, or about 0.001 mg/kg/day to about 30 mg/kg/day, or about 0.001 mg/kg/day to about 20 mg/kg/day, or about 0.001 mg/kg/day to about 10 mg/kg/day, or about 0.01 mg/kg/day to about 2000 mg/kg/day, or about 0.01 mg/kg/day to about 1500 mg/kg/day, or about 0.01 mg/kg/day to about 1000 mg/kg/day, or about 0.01 mg/kg/day to about 600 mg/kg/day, or about 0.01 mg/kg/day to about 500 mg/kg/day, or about 0.01 mg/kg/day to about 300 mg/kg/day, or about 0.015 mg/kg/day to about 200 mg/kg/day, or about 0.02 mg/kg/day to about 100 mg/kg/day, or about 0.025 mg/kg/day to about 100 mg/kg/day, or about 0.03 mg/kg/day to about 100 mg/kg/day, wherein said amount is administered once (once in approximately a 24 hour period; i.e., "q.d."), twice (once in approximately a 12 hour period; i.e., "b.i.d." or "q. 12h"), thrice (once in approximately an 8 hour period; i.e., "t.i.d." or "q.8h"), or four times (once in approximately a 6 hour period; i.e., "q.d.s.", "q.i.d." or "q.6h") daily according to subject weight.

[0170] Such amounts may further include an amount in a range of from about 0.001 mg to about 3500 mg administered daily; 0.001 mg to about 3000 mg administered daily; 0.001 mg to about 2500 mg administered daily; 0.001 mg to about 2000 mg administered daily; 0.001 mg to about 1500 mg administered daily; 0.001 mg to about 1000 mg administered daily; 0.001 mg to about 500 mg administered daily; 0.001 mg to about 250 mg administered daily; 1.0 mg to about 3500 mg administered daily; 1.0 mg to about 1500 mg administered daily; 1.0 mg to about 1000 mg administered daily; 10.0 mg to about 600 mg administered daily; 0.5 mg to about 2000 mg administered daily; or, an amount in a range of from about 5.0 mg to about 300 mg administered daily.

[0171] For example, the effective amount may be the amount required to treat HD in a subject or, more specifically, in a human. The effective amount for a subject will depend upon various factors, including the subject's body weight, size and health. Effective amounts for a given patient can be determined by routine experimentation that is within the skill and judgment of the clinician.

[0172] For any compound, the effective amount can be estimated initially either in cell culture assays or in relevant animal models, such as a mouse, chimpanzee, marmoset or tamarin animal model. Relevant animal models may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED₅₀ (the dose therapeutically effective in 50% of the population) and LD₅₀ (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is therapeutic index, and can be expressed as the ratio, LD₅₀/ED₅₀. In some embodiments, the effective amount is such that a large therapeutic index is achieved. In further embodiments, the dosage is within a range of circulating concentrations that include an ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[0173] More specifically, the concentration-biological effect relationships observed with regard to a compound of Formula (I) or a form thereof indicate a target plasma concentration ranging from approximately 0.001 µg/mL to approximately 50 µg/mL, from approximately 0.01 µg/mL to approximately 20 µg/mL, from approximately 0.05 µg/mL to approximately 10 µg/mL, or from approximately 0.1 µg/mL to approximately 5 µg/mL. To achieve such plasma concentrations, the compounds described herein may be administered at doses that vary, such as, for example, without limitation, from 0.1 ng to 10,000 mg, depending upon the route of administration in single, divided, or continuous doses for a patient weighing between about 10 to about 100 kg (which dose may be adjusted for patients within this weight range, particularly for children under 40 kg).

[0174] The exact dosage will be determined by the practitioner, in light of factors related to the subject. Dosage and administration may be adjusted to provide sufficient levels of the active agent(s) or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, ethnicity, age, weight, gender, diet, time of day and frequency of administration, drug combination(s), reaction sensitivities, experience with other therapies, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered

every 2, 3 or 4 days, once every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

[0175] The compounds and compositions described herein may be administered to the subject via any drug delivery route known in the art. Nonlimiting examples include oral, ocular, rectal, buccal, topical, nasal, sublingual, transdermal, subcutaneous, intramuscular, intravenous (bolus and infusion), intracerebral, and pulmonary routes of administration.

[0176] In one aspect, provided herein are methods for modulating the amount of HTT (huntingtin protein), comprising contacting a human cell with a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein are methods for modulating the amount of HTT, comprising contacting a human cell with a compound of Formula (I) or a form thereof that modulates the expression of HTT. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, e.g., in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the Htt gene, resulting in a loss of HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

[0177] In a specific embodiment, provided herein is a method for enhancing the inhibition of mutant HTT transcribed from the Htt gene, comprising contacting a human cell with a compound of Formula (I) or a form thereof. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, e.g., in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the Htt gene, resulting in a loss of wild-type "normal" HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

[0178] In another aspect, provided herein is a method for modulating the inhibition of mutant HTT transcribed from the Htt gene, comprising administering to a non-human animal model for HD a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein is a method for modulating the inhibition of mutant HTT transcribed from the Htt gene, comprising administering to a non-human animal model for HD a compound of Formula (I) or a form thereof. In a specific embodiment, the compound is a form of the compound of Formula (I).

[0179] In another aspect, provided herein is a method for decreasing the amount of mutant HTT, comprising contacting a human cell with a compound of Formula (I) or a form thereof. In a specific embodiment, provided herein is a method for decreasing the amount of mutant HTT, comprising contacting a human cell with a compound of Formula (I) that inhibits the transcription of mutant *HTT* (huntingtin mRNA) from the Htt gene. In another specific embodiment, provided herein is a method for decreasing the amount of HTT, comprising contacting a human cell with a compound of Formula (I) that inhibits the expression of mutant HTT transcribed from the Htt gene. The human cell can be contacted with a compound of Formula (I) or a form thereof *in vitro*, or *in vivo*, e.g., in a non-human animal or in a human. In a specific embodiment, the human cell is from or in a human. In another specific embodiment, the human cell is from or in a human with HD. In another specific embodiment, the human cell is from or in a human with HD, caused by a CAG repeat in the Htt gene, resulting in a loss of HTT expression and/or function. In another embodiment, the human cell is from a human with HD. In another embodiment, the human cell is in a human with HD. In one embodiment, the compound is a form of the compound of Formula (I).

[0180] In certain embodiments, treating or ameliorating HD with a compound of Formula (I) or a form thereof (alone or in combination with an additional agent) has a therapeutic effect and/or beneficial effect. In a specific embodiment, treating HD with a compound of Formula (I) or a form thereof (alone or in combination with an additional agent) results in one, two or more of the following effects: (i) reduces or ameliorates the severity of HD; (ii) delays onset of HD; (iii) inhibits the progression of HD; (iv) reduces hospitalization of a subject; (v) reduces hospitalization length for a subject; (vi) increases the survival of a subject; (vii) improves the quality of life for a subject; (viii) reduces the number of symptoms associated with HD; (ix) reduces or ameliorates the severity of a symptom(s) associated with HD; (x) reduces the duration of a symptom associated with HD; (xi) prevents the recurrence of a symptom associated with HD; (xii) inhibits the development or onset of a symptom of HD; and/or (xiii) inhibits of the progression of a symptom associated with HD.

Metabolites of the Compounds

[0181] Also included within the scope of the present description are the use of *in vivo* metabolic products of the compounds described herein. Such products may result, for example, from the oxidation, reduction, hydrolysis, amidation, esterification and the like of the administered compound, primarily due to enzymatic processes. Accordingly, the description includes the use of compounds produced by a process comprising contacting a compound described herein with a mammalian tissue or a mammal for a period of time sufficient to yield a metabolic product thereof.

[0182] Such products typically are identified by preparing a radio-labeled isotopologue (e.g., ^{14}C or ^3H) of a compound

described herein, administering the radio-labeled compound in a detectable dose (e.g., greater than about 0.5 mg/kg) to a mammal such as a rat, mouse, guinea pig, dog, monkey or human, allowing sufficient time for metabolism to occur (typically about 30 seconds to about 30 hours), and identifying the metabolic conversion products from urine, bile, blood or other biological samples. The conversion products are easily isolated since they are "radiolabeled" by virtue of being isotopically-enriched (others are isolated by the use of antibodies capable of binding epitopes surviving in the metabolite). The metabolite structures are determined in conventional fashion, e.g., by MS or NMR analysis. In general, analysis of metabolites may be done in the same way as conventional drug metabolism studies well-known to those skilled in the art. The conversion products, so long as they are not otherwise found *in vivo*, are useful in diagnostic assays for therapeutic dosing of the compounds described herein even if they possess no biological activity of their own.

Pharmaceutical Compositions

[0183] Embodiments of the present description include the use of a compound of Formula (I) or a form thereof in a pharmaceutical composition for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the compound of Formula (I) or a form thereof in admixture with one or more pharmaceutically acceptable excipient(s).

[0184] An embodiment of the present description includes the use of a pharmaceutical composition of the compound of Formula (I) or a form thereof in the preparation of a kit comprising the pharmaceutical composition of the compound of Formula (I) or a form thereof and instructions for administering the compound for treating or ameliorating HD in a subject in need thereof.

[0185] As used herein, the term "composition" means a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

[0186] The pharmaceutical composition may be formulated to achieve a physiologically compatible pH, ranging from about pH 3 to about pH 11. In some embodiments, the pharmaceutical composition is formulated to achieve a pH of from about pH 3 to about pH 7. In other embodiments, the pharmaceutical composition is formulated to achieve a pH of from about pH 5 to about pH 8.

[0187] The term "pharmaceutically acceptable excipient" refers to an excipient for administration of a pharmaceutical agent, such as the compounds described herein. The term refers to any pharmaceutical excipient that may be administered without undue toxicity. Pharmaceutically acceptable excipients may be determined in part by the particular composition being administered, as well as by the particular mode of administration and/or dosage form. Nonlimiting examples of pharmaceutically acceptable excipients include carriers, solvents, stabilizers, adjuvants, diluents, *etc.* Accordingly, there exists a wide variety of suitable formulations of pharmaceutical compositions for the instant compounds described herein (see, e.g., Remington's Pharmaceutical Sciences).

[0188] Suitable excipients may be carrier molecules that include large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive antibodies. Other exemplary excipients include antioxidants such as ascorbic acid; chelating agents such as EDTA; carbohydrates such as dextrin, hydroxyalkylcellulose, hydroxyalkylmethylcellulose (e.g., hydroxypropylmethylcellulose, also known as HPMC), stearic acid; liquids such as oils, water, saline, glycerol and ethanol; wetting or emulsifying agents; pH buffering substances; and the like. Liposomes are also included within the definition of pharmaceutically acceptable excipients.

[0189] The pharmaceutical compositions described herein may be formulated in any form suitable for the intended use described herein. Suitable formulations for oral administration include solids, liquid solutions, emulsions and suspensions, while suitable inhalable formulations for pulmonary administration include liquids and powders. Alternative formulations include syrups, creams, ointments, tablets, and lyophilized solids which can be reconstituted with a physiologically compatible solvent prior to administration.

[0190] When intended for oral use for example, tablets, troches, lozenges, aqueous or oil suspensions, non-aqueous solutions, dispersible powders or granules (including micronized particles or nanoparticles), emulsions, hard or soft capsules, syrups or elixirs may be prepared. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions, and such compositions may contain one or more agents including sweetening agents, flavoring agents, coloring agents, and preserving agents, in order to provide a palatable preparation.

[0191] Pharmaceutically acceptable excipients suitable for use in conjunction with tablets include, for example, inert diluents, such as celluloses, calcium or sodium carbonate, lactose, calcium or sodium phosphate; disintegrating agents, such as croscarmellose sodium, cross-linked povidone, maize starch, or alginic acid; binding agents, such as povidone, starch, gelatin or acacia; and lubricating agents, such as magnesium stearate, stearic acid, or talc. Tablets may be uncoated or may be coated by known techniques including microencapsulation to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay

material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

[0192] Formulations for oral use may be also presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example celluloses, lactose, calcium phosphate, or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with non-aqueous or oil medium, such as glycerin, propylene glycol, polyethylene glycol, peanut oil, liquid paraffin, or olive oil.

[0193] In other embodiments, pharmaceutical compositions described herein may be formulated as suspensions comprising a compound of Formula (I) or a form thereof in admixture with one or more pharmaceutically acceptable excipient(s) suitable for the manufacture of a suspension. In yet other embodiments, pharmaceutical compositions described herein may be formulated as dispersible powders and granules suitable for preparation of a suspension by the addition of one or more excipient(s).

[0194] Excipients suitable for use in connection with suspensions include suspending agents, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropyl methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth, gum acacia, dispersing or wetting agents such as a naturally occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g., heptadecaethyleneoxycethanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan monooleate); and thickening agents, such as carbomer, beeswax, hard paraffin, or cetyl alcohol. The suspensions may also contain one or more preservatives such as acetic acid, methyl and/or n-propyl p-hydroxy-benzoate; one or more coloring agents; one or more flavoring agents; and one or more sweetening agents such as sucrose or saccharin.

[0195] The pharmaceutical compositions described herein may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachis oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include naturally-occurring gums, such as gum acacia and gum tragacanth; naturally occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids; hexitol anhydrides, such as sorbitan monooleate; and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, such as glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavoring or a coloring agent.

[0196] Additionally, the pharmaceutical compositions described herein may be in the form of a sterile injectable preparation, such as a sterile injectable aqueous emulsion or oleaginous suspension. Such emulsion or suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, such as a solution in 1,2-propanediol. The sterile injectable preparation may also be prepared as a lyophilized powder. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile fixed oils may be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

[0197] The compounds described herein may be substantially insoluble in water and sparingly soluble in most pharmaceutically acceptable protic solvents and vegetable oils, but generally soluble in medium-chain fatty acids (e.g., caprylic and capric acids) or triglycerides and in propylene glycol esters of medium-chain fatty acids. Thus, contemplated in the description are compounds which have been modified by substitutions or additions of chemical or biochemical moieties which make them more suitable for delivery (e.g., increase solubility, bioactivity, palatability, decrease adverse reactions, etc.), for example by esterification, glycosylation, PEGylation, etc.

[0198] In some embodiments, the compound described herein is formulated for oral administration in a lipid-based composition suitable for low solubility compounds. Lipid-based formulations can generally enhance the oral bioavailability of such compounds. As such, pharmaceutical compositions described herein may comprise an effective amount of a compound of Formula (I) or a form thereof, together with at least one pharmaceutically acceptable excipient selected from medium chain fatty acids or propylene glycol esters thereof (e.g., propylene glycol esters of edible fatty acids such as caprylic and capric fatty acids) and pharmaceutically acceptable surfactants, such as polysorbate 20 or 80 (also referred to as Tween® 20 or Tween® 80, respectively) or polyoxyl 40 hydrogenated castor oil.

[0199] In other embodiments, the bioavailability of low solubility compounds may be enhanced using particle size optimization techniques including the preparation of nanoparticles or nanosuspensions using techniques known to those skilled in the art. The compound forms present in such preparations include amorphous, partially amorphous, partially crystalline or crystalline forms.

[0200] In alternative embodiments, the pharmaceutical composition may further comprise one or more aqueous solubility enhancer(s), such as a cyclodextrin. Nonlimiting examples of cyclodextrin include hydroxypropyl, hydroxyethyl, glucosyl, maltosyl and maltotriosyl derivatives of α -, β -, and γ -cyclodextrin, and hydroxypropyl- β -cyclodextrin (HPBC). In some embodiments, the pharmaceutical composition further comprises HPBC in a range of from about 0.1% to about 20%, from about 1% to about 15%, or from about 2.5% to about 10%. The amount of solubility enhancer employed may

depend on the amount of the compound in the composition.

Preparation of Compounds

[0201] Compounds provided herein can be prepared by those skilled in the art, such as, by the synthetic methods set forth in International Application Publication Number WO2014/028459 A1, published February 20, 2014, International Application Publication Number WO2014/116845 A1, published July 31, 2014, and International Application Publication Number WO2015/017589 A1, published February 5, 2015, each of which are herein incorporated by reference.


General Schemes

[0202] The following reaction schemes illustrate methods to make compounds described herein. It is understood that one skilled in the art would be able to make these compounds by similar methods or by methods known to one skilled in the art.

[0203] In general, starting components and reagents may be commercially obtained from various sources such as commercial vendors, synthesized according to methodology known to those skilled in the art, or prepared as described herein. It is understood that in the following description, combinations of substituents and/or variables of the depicted formulae are permissible only if such contributions result in stable compounds.

[0204] In general, compounds of Formula (I) described herein can be synthesized following the general procedure for Scheme 1.

Scheme 1



General Procedure for Scheme 1

[0205] The starting materials for the above reaction scheme are commercially available or can be prepared according to methods known to one skilled in the art or by methods disclosed herein. In general, the compounds described herein are prepared in the above reaction Scheme 1 as follows: a halogenated Compound 1a (wherein Hal is a halogen selected from bromine or chlorine) is reacted with a Compound 1b (wherein H is a reactive hydrogen atom in an amine or alcohol X functional group, wherein X and B for a compound of Formula (I) are as described herein) using a displacement or a metal-mediated cross coupling reaction, such as a Buchwald reaction, to provide an intermediate Compound 1c. Compound 1c is carried forward and reacted with a boronate acid or boronate ester Compound 1d (where B represents a boron atom, R represents an acid or ester functional group or, when taken together with boron, form a ring system and A for a compound of Formula (I) is as described herein) using a transition metal-mediated cross coupling reaction, such as a Suzuki reaction, to provide a Compound of Formula (I).

[0206] In a complementary manner, compounds of Formula (I) may also be synthesized in reverse order wherein the boronate Compound 1d is reacted first with Compound 1a followed by reaction of the intermediate with amine Compound 1b. Other variations representing different combinations for substituting either or both boronate Compound 1d and amine Compound 1b on Compound 1a are intended to be included within the scope of the synthetic methodologies described herein.

Specific Examples

[0207] To assist in understanding the present description, the following specific examples are included. The experiments relating to this description should not, of course, be construed as specifically limiting the description and such variations of the description, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the description as described herein and hereinafter claimed.

[0208] Other than in the working examples, unless indicated to the contrary, all numbers expressing quantities of materials, reagents, reaction conditions, experimental data, and so forth used in the specification and claims are to be understood as being modified by the term "about". Accordingly, all such numbers represent approximations that may

vary depending upon the desired properties sought to be obtained by a reaction or as a result of variable experimental conditions. Therefore, within an expected range of experimental reproducibility, the term "about" in the context of the resulting data, refers to a range for data provided that may vary according to a standard deviation from the mean. As well, for experimental results provided, the resulting data may be rounded up or down to present data consistently, without loss of significant figures. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding techniques.

[0209] While the numerical ranges and parameters setting forth the broad scope of the description are approximations, the numerical values set forth in the working examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0210] The column chromatography system used for product purification was an ISCO CombiFlash™ brand chromatography system (manufactured by Teledyne Isco).

Synthetic Examples

[0211] Greater details of the present description are provided with reference to the following non-limiting examples, which are offered to more fully illustrate the description, but are not to be construed as limiting the scope thereof. The examples illustrate the preparation of certain compounds described herein, and the testing of these compounds *in vitro* and/or *in vivo*. Those of skill in the art will understand that the techniques described in these examples represent techniques described by the inventors to function well in the practice of the description, and as such constitute preferred modes for the practice thereof. However, those of skill in the art should appreciate in light of the present disclosure that many changes can be made to the specific methods that are disclosed and still obtain a like or similar result without departing from the spirit and scope of the description.

[0212] As used above, and throughout this description, the following abbreviations, unless otherwise indicated, shall be understood to have the following meanings:

Abbreviation	Meaning
AcOH or HOAc	acetic acid
ACN or MeCN	acetonitrile
Bn	benzyl
BnBr	benzyl bromide
BnO or OBn	benzyloxy
Boc	<i>tert</i> -butoxycarbonyl
Boc ₂ O or (Boc) ₂ O	di- <i>tert</i> -butyl dicarbonate
<i>t</i> BuXPhos-Pd-G3	[(2-di- <i>tert</i> -butylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)-2-(2'-amino-1,1'-biphenyl)] palladium (II) methanesulfonate
Cbz	benzyloxycarbonyl
DCE	dichloroethane
DCM	dichloromethane (CH ₂ Cl ₂)
DIBAL-H	diisobutylaluminium hydride
DIPEA	N,N-diisopropylethylamine
DMF	dimethyl formamide
DMA	dimethylacetamide
DMAP	4-dimethylaminopyridine
DMB	2,4-dimethoxybenzyl
DMSO	dimethylsulfoxide
EA or EtOAc	ethyl acetate
EtOH	ethanol
Et ₂ O	diethyl ether
HCl	hydrochloric acid
HPLC	high performance liquid chromatography
h/hr/hrs/min/s	hour(s)(h, hr or hrs)/minute(s)(min/mins)/second(s)

(continued)

	Abbreviation	Meaning
	K ₂ CO ₃	potassium carbonate
5	KOAc	potassium acetate
	K ₃ PO ₄	potassium phosphate
	LAH	lithium aluminium hydride
	LC/MS, LCMS or LC-MS	liquid chromatographic mass spectroscopy
10	LDA	lithium diisopropylamide
	LiHMDS	lithium bis(trimethylsilyl)amide
	Me ₄ tBu- XPhos	di-tert-butyl(2',4',6'-triisopropyl-3,4,5,6-tetramethyl-[1,1'-biphenyl]-2-yl)phosphine
15	MeI	methyl iodide
	MeOH	methanol
	Me ₂ NH or NHMe ₂	dimethyl amine
	MS	mass spectroscopy
20	N ₂	nitrogen
	Na ₂ SO ₄	sodium sulfate
	NaBH(OAc) ₃	sodium triacetoxyborohydride
	NaHMDS	sodium bis(trimethylsilyl)amide
25	NBS	N-bromosuccinimide
	NH ₄ OH	ammonium hydroxide
	NMO	N-methylmorpholine-N-oxide
	NMP	N-methyl-2-pyrrolidone
	n-BuLi	n-butyl lithium
30	NMR	nuclear magnetic resonance
	Pd/C	palladium on carbon
	Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
	Pd(dppf)Cl ₂	[1,1'-bis(diphenylphosphino)ferrocene] dichloropalladium(II)
35	Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium
	Pin	pinacol
	psi	pounds per square inch pressure
	S-Phos	2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl
	TBDPSCI	tert-butyldiphenylchlorosilane
40	TBS	tert-butyldimethylsilyl
	TBSCl	tert-butyldimethylsilyl chloride
	t-BuOK	potassium tert-butoxide
	TEA or NEt ₃	triethylamine
45	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
	THP	tetrahydro-2H-pyranyl
	THPO or OTHP	tetrahydro-2H-pyran-2-yl-oxy
50	TIPS-H or TIPSH	triisopropyl silane
	TLC	thin layer chromatography
	TMP	2,2,6,6-tetramethylpiperidinyl
55	TMPMgCl- LiCl	2,2,6,6-tetramethylpiperidinylmagnesium chloride lithium chloride
	TMSI	trimethylsilyl iodide
	TMSOK	potassium trimethylsilanolate

Example 1 (Compound 411)

2-[6-[8-Azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

[0213] Step 1: 3,6-Dibromopyridazine (133.7 mg, 0.56 mmol), tert-butyl (1R,5S)-3-(methylamino)-8-azabicyclo[3.2.1]octane-8-carboxylate (85 mg, 0.28 mmol) and DIPEA (0.15 mL, 0.84 mmol) were mixed in 1 mL of ACN and heated to 100 °C for 1 h until LC-MS showed complete consumption of the starting material. The reaction mixture was concentrated and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 4g to afford tert-butyl (1S,5R)-3-[(6-bromopyridazin-3-yl)-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (55 mg, 49.25% yield).

[0214] Step 2: tert-Butyl 3-[(6-bromopyridazin-3-yl)-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (55 mg, 0.14 mmol), 4-[3-(methoxymethoxy)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1-tetrahydropyran-2-ylpyrazole (68.82 mg, 0.17 mmol), Pd(dppf)Cl₂ (10.34 mg, 0.01 mmol), K₂CO₃ (57.98 mg, 0.42 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, purified via column chromatography: eluting with gradient CH₂Cl₂/EtOAc (0% to 80%), column: silica 4g, to provide tert-butyl (1S,5R)-3-[[6-[2-(methoxymethoxy)-4-(1-tetrahydropyran-2-yl)pyrazol-4-yl]phenyl]pyridazin-3-yl]-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (52 mg, 62%) as a tan solid.

[0215] Step 3: To a solution of tert-butyl (1S,5R)-3-[[6-[2-(methoxymethoxy)-4-(1-tetrahydropyran-2-yl)pyrazol-4-yl]phenyl]pyridazin-3-yl]-methyl-amino]-8-azabicyclo[3.2.1]octane-8-carboxylate (52 mg, 0.086 mmol) in a 1 mL mixture of CH₂Cl₂ and 0.5 mL of MeOH was added HCl (4 mol/L) in 1,4-dioxane (0.06 mL). The reaction was stirred overnight. The precipitate was filtered, then dried under vacuum to provide 2-[6-[8-azabicyclo[3.2.1]octan-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol; hydrochloride (15 mg, 42.25% yield) as a yellow solid.

LC-MS: 377 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.37-9.51 (m, 1H), 8.87-8.96 (m, 1H), 8.31 (d, *J*=9.8 Hz, 1H), 8.12 (s, 2H), 7.74 (d, *J*=8.2 Hz, 2H), 7.19-7.27 (m, 2H), 4.80-4.92 (m, 1H), 4.08-4.14 (m, 3H), 3.07 (s, 3H), 2.30 (td, *J*=12.6, 2.8 Hz, 2H), 1.99-2.15 (m, 4H), 1.74-1.87 (m, 2H)

[0216] Using the procedure described for Example 1 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
412	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.24-9.34 (m, 1H), 9.09-9.21 (m, 1H), 8.30 (d, <i>J</i> =9.8 Hz, 1H), 8.09 (s, 2H), 7.64-7.67 (m, 1H), 7.63 (d, <i>J</i> =8.2 Hz, 1H), 7.25-7.27 (m, 1H), 7.24 (s, 1H), 5.42 (br s, 2H), 4.23-4.31 (m, 1H), 4.00-4.13 (m, 2H), 2.17 (s, 4H), 1.97-2.02 (m, 2H), 1.94 (td, <i>J</i> =12.6, 2.5 Hz, 2H)
413	LC-MS: 393 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.34-9.49 (m, 1H), 8.43-8.56 (m, 1H), 8.30 (d, <i>J</i> =9.8 Hz, 1H), 8.10 (s, 2H), 7.63 (s, 2H), 7.19-7.30 (m, 2H), 5.77 (br s, 2H), 4.41-4.62 (m, 1H), 2.08 (dd, <i>J</i> =11.7, 2.8 Hz, 2H), 1.72 (td, <i>J</i> =12.1, 2.8 Hz, 2H), 1.53 (s, 6H), 1.49 (s, 6H)
415	LC-MS: 364 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.61 (d, <i>J</i> =9.5 Hz, 1H), 8.24 (s, 2H), 7.85 (d, <i>J</i> =8.2 Hz, 1H), 7.72 (d, <i>J</i> =9.5 Hz, 1H), 7.38 (dd, <i>J</i> =8.2, 1.6 Hz, 1H), 7.32 (d, <i>J</i> =1.6 Hz, 1H), 5.65 (spt, <i>J</i> =5.7 Hz, 1H), 4.25 (dd, <i>J</i> =3.6, 2.8 Hz, 2H), 4.17-4.32 (m, 2H), 2.24 (s, 4H), 2.11 (tt, <i>J</i> =11.2, 2.8 Hz, 2H) (3 hydrogens corresponding to OH and NH unobserved)
433	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.42 (d, <i>J</i> =9.5 Hz, 1H), 8.34 (s, 2H), 7.80 (d, <i>J</i> =9.8 Hz, 1H), 7.74 (d, <i>J</i> =8.5 Hz, 1H), 7.40 (dd, <i>J</i> =8.2, 2.5 Hz, 1H), 7.32 (d, <i>J</i> =1.6 Hz, 1H), 4.18 (td, <i>J</i> =6.9, 0.9 Hz, 1H), 4.11-4.16 (m, 2H), 2.42-2.52 (m, 4H), 2.32-2.35 (m, 1H), 2.28-2.32 (m, 1H), 2.15-2.25 (m, 2H), (4 hydrogens corresponding to OH and NH unobserved)

Example 2 (Compound 416)

5-(5-Methyl-1H-pyrazol-4-yl)-2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol

[0217] Step 1: 3-Methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl trifluoromethanesulfonate (80 mg, 0.16 mmol), (tert-butyl 3-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (59 mg, 0.19 mmol), Pd(dppf)Cl₂ (7 mg, 0.01 mmol), K₂CO₃ (66 mg, 0.48 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction

was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, then purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 4g to provide tert-butyl 4-(3-methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-methyl-1H-pyrazole-1-carboxylate (46 mg, 54%) as light brown solid.

[0218] Step 2: A solution of tert-butyl 4-(3-methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-methyl-1H-pyrazole-1-carboxylate (46 mg, 0.09mmol) in 2 mL of dry CH₂Cl₂ was cooled in ice-water bath. Boron tribromide 1.0 M in CH₂Cl₂ (0.45 mL, 0.45 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (2.5% NH₄OH) (0% to 30% MeOH/NH₄OH), column: silica 4g, to provide tert-butyl 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(3-methyl-1H-pyrazol-4-yl)phenol (20 mg, 55%) as a yellow solid.

LC-MS: 421 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 13.74 (s, 1H), 12.63-12.71 (m, 1H), 8.20 (d, *J*=9.5Hz, 1H), 8.03-8.04 (m, 1H), 7.86 (d, *J*=8.0Hz, 1H), 7.73-7.75 (m, 1H), 7.36 (d, *J*=9.5Hz, 1H), 7.03-7.06 (m, 2H), 4.90-5.05 (m, 1H), 2.92 (s, 3H), 2.43 (s, 3H), 1.43-1.54 (m, 2H), 1.15-1.43 (m, 2H), 1.14 (s, 6H), 1.09 (s, 6H)

[0219] Using the procedure described for Example 2 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
414	LC-MS: 408 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 13.84 (s, 1H), 8.10-8.29 (m, 2H), 7.92 (s, 1H), 7.84 (d, <i>J</i> =8.2 Hz, 1H), 7.36 (d, <i>J</i> =9.8 Hz, 1H), 7.08-7.19 (m, 2H), 4.87-5.03 (m, 1H), 3.87 (s, 3H), 2.96 (s, 3H), 1.48-1.59 (m, 2H), 1.38-1.48 (m, 2H), 1.26 (s, 6H), 1.09 (s, 6H)
418	LC-MS: 422 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 12.89 (s, 1H), 12.29-12.48 (m, 1H), 8.41 (dd, <i>J</i> =8.8, 0.9 Hz, 1H), 7.95 (dd, <i>J</i> =7.9, 0.9 Hz, 1H), 7.37 (dd, <i>J</i> =9.1, 1.3 Hz, 1H), 6.85-6.94 (m, 2H), 5.61-5.70 (m, 1H), 2.25 (br s., 6H), 2.09 (dd, <i>J</i> =11.8, 3.9 Hz, 2H), 1.25-1.33 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)
451	LC-MS: 408 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 13.00 (s, 1H), 12.59-12.80 (m, 1H), 8.42 (d, <i>J</i> =9.5 Hz, 1H), 7.94 (d, <i>J</i> =8.8 Hz, 1H), 7.38 (d, <i>J</i> =9.5 Hz, 1H), 7.03-7.18 (m, 2H), 5.55-5.78 (m, 1H), 2.39 (br s., 3H), 2.03-2.13 (m, 2H), 1.26-1.41 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)

Example 3 (Compound 419)

2-{6-[Methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride

[0220] Step 1: 3-Methoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl trifluoromethanesulfonate (335 mg, 0.67 mmol), 4-nitro-1H-pyrazole (340 mg, 4.5 mmol), Pd₂(dba)₃ (67 mg, 0.07 mmol), Me₄tBu-XPhos (35 mg, 0.07 mmol), K₃PO₄ (368 mg, 1.73 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (8 mL) was added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, filtered through celite, concentrated, purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 12g, to provide 6-(2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (202 mg, 65%).

LC-MS: 466 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.81-9.85 (m, 1H), 8.61 (d, *J*=0.6 Hz, 1H), 7.91 (d, *J*=8.5 Hz, 1H), 7.81 (d, *J*=9.8 Hz, 1H), 7.73 (d, *J*=1.9 Hz, 1H), 7.68 (dd, *J*=8.2, 2.2 Hz, 1H), 7.10-7.21 (m, 1H), 5.08-5.26 (m, 1H), 3.96 (s, 3H), 2.95 (s, 3H), 1.59-1.75 (m, 4H), 1.40 (s, 6H), 1.33, (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)

[0221] Step 2: 6-(2-Methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (150 mg, 0.32 mmol), benzenethiol (33 μL, 0.32 mmol) and K₂CO₃ (44 mg, 0.32 mmol) were mixed in a microwave tube. The reaction was degassed with N₂ for 15 min and dry NMP (1.5 mL) was added. The reaction was heated in a Biotage microwave at 190 °C for 20 min, then diluted with 10 mL EtOAc, and washed with water and brine. The product was dried over Na₂SO₄, concentrated under vacuum, then purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30% MeOH), column: silica 4g, to provide 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-pyrazol-1-yl)phenol (60 mg, 42%) as a brown solid.

LC-MS: 452 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.77 (s, 1H), 8.68 (d, *J*=9.8 Hz, 1H), 8.62 (s, 1H), 8.27 (d, *J*=8.8 Hz, 1H), 7.87-7.94 (m, 1H), 7.66 (dd, *J*=8.5, 2.6 Hz, 1H), 7.64 (d, *J*=2.6 Hz, 1H), 5.03-5.22 (m, 1H), 3.01 (s, 3H), 1.84-1.94

(m, 2H), 1.70-1.77 (m, 2H), 1.53 (s, 6H), 1.43 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)

[0222] Using the procedure described for Example 3 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
417	LC-MS: 394 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 13.49 (br s, 1H), 8.49 (d, <i>J</i> =9.8 Hz, 1H), 8.39 (s, 1H), 8.10 (d, <i>J</i> =8.8 Hz, 1H), 7.79-7.92 (m, 1H), 7.43 (d, <i>J</i> =9.5 Hz, 1H), 7.34 (d, <i>J</i> =2.5 Hz, 1H), 7.30 (dd, <i>J</i> =8.5, 2.5 Hz, 1H), 7.12 (s, 1H), 5.61-5.71 (m, 1H), 2.09 (dd, <i>J</i> =12.0, 3.8 Hz, 2H), 1.25-1.32 (m, 2H), 1.24 (s, 6H), 1.11 (s, 6H), (1 hydrogen corresponding to OH or NH unobserved)
421	LC-MS: 409 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.31 (d, <i>J</i> =9.5 Hz, 1H), 7.92 (d, <i>J</i> =9.1 Hz, 1H), 7.79 (d, <i>J</i> =0.6 Hz, 1H), 7.44 (d, <i>J</i> =0.9 Hz, 1H), 7.24-7.33 (m, 3H), 5.77 (tt, <i>J</i> =11.2, 4.1 Hz, 1H), 2.28 (dd, <i>J</i> =12.3, 4.2 Hz, 2H), 1.49 (t, <i>J</i> =12.3 Hz, 2H), 1.42 (s, 6H), 1.30 (s, 6H), (4 hydrogens corresponding to OH and NH unobserved)
423	LC-MS: 439 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.71 (d, <i>J</i> =0.6 Hz, 1H), 8.58 (d, <i>J</i> =0.6 Hz, 1H), 8.47 (d, <i>J</i> =9.5 Hz, 1H), 8.14 (d, <i>J</i> =8.5 Hz, 1H), 7.59-7.64 (m, 2H), 7.44 (d, <i>J</i> =9.5 Hz, 1H), 5.67-5.73 (m, 1H), 2.18-2.28 (m, 2H), 1.45-1.56 (m, 2H), 1.37 (s, 6H), 1.27 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)

Example 4 (Compound 424)

5-(1H-Pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride

[0223] Step 1: 3-Bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazine (100 mg, 0.24 mmol), (1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)boronic acid (66 mg, 0.29 mmol), PddppfCl₂ (8 mg, 0.012 mmol), K₂CO₃ (99 mg, 0.72 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (2 mL) and water (0.5 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, and partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, then concentrated under vacuum, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 20% MeOH), column: silica 4g, to provide 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (102 mg, 82%) as an off-white solid.

[0224] Step 2: A solution of 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (102 mg, 0.196 mmol) in 2 mL of dry CH₂Cl₂ was cooled in ice-water bath. Boron tribromide 1.0 M in CH₂Cl₂ (0.98 mL, 0.98 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified using preparative HPLC to provide 5-(1H-pyrazol-4-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol (41 mg, 66%) as an off-white solid.

LC-MS: 320 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.19 (br s, 2H), 8.54 (d, *J*=9.6 Hz, 1H), 8.22 (d, *J*=9.6 Hz, 1H), 8.18 (s, 1H), 8.05 (d, *J*=8.8 Hz, 1H), 7.25-7.29 (m, 2H), 6.92 (s, 1H), 3.85-3.92 (m, 2H), 3.38-3.41 (m, 2H), 2.88-2.98 (m, 2H), (1 hydrogen corresponding to OH or NH unobserved)

[0225] Using the procedure described for Example 4 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
422	LC-MS: 334 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 11.06 (br s, 1H), 8.56 (d, <i>J</i> =9.2 Hz, 1H), 8.25 (d, <i>J</i> =9.2 Hz, 1H), 8.20 (s, 2H), 8.05 (d, <i>J</i> =8.8 Hz, 1H), 7.22-7.31 (m, 2H), 6.92 (s, 1H), 4.06-4.12 (m, 1H), 3.87-3.94 (m, 1H), 3.62-3.70 (m, 1H), 3.21-3.30 (m, 1H), 3.07-3.12 (m, 2H), 3.03 (s, 3H), (1 hydrogen corresponding to OH or NH unobserved)
425	LC-MS: 348 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 10.94 (br s, 1H), 8.56 (d, <i>J</i> =9.6 Hz, 1H), 8.26 (d, <i>J</i> =9.6 Hz, 1H), 8.19-8.23 (m, 2H), 8.05 (d, <i>J</i> =8.4 Hz, 1H), 7.26-7.31 (m, 2H), 6.93 (s, 1H), 4.07-4.13 (m, 1H), 3.85-3.91 (m, 1H), 3.70-3.74 (m, 1H), 3.22-3.30 (m, 3H), 3.08-3.14 (m, 2H), 1.33 (t, <i>J</i> =7.2 Hz, 3H), (1 hydrogen corresponding to OH or NH unobserved)

(continued)

Cpd	Data
429	LC-MS: 346 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 1.79 - 1.89 (m, 1H) 1.98 - 2.11 (m, 1H) 2.11 - 2.33 (m, 2H) 2.65 - 2.80 (m, 1H) 3.09 - 3.21 (m, 1H) 4.02 - 4.16 (m, 2H) 7.02 - 7.14 (m, 1H) 7.24 (s, 2H) 7.85 - 7.97 (m, 1H) 8.04 (d, <i>J</i> =9.46 Hz, 3H) 8.33 (d, <i>J</i> =9.46 Hz, 1H)

Example 5 (Compound 427)

2-[6-(Piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride

[0226] Step 1: 3,6-Dibromopyridazine (500 mg, 2.1 mmol), 4-(3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazole (970 mg, 2.52 mmol), Pd(dppf)Cl₂ (77 mg, 0.11 mmol), K₂CO₃ (870 mg, 6.3 mmol) were mixed in a Schlenk tube. The reaction was degassed with N₂ for 15 min and dioxane (8 mL) and water (1 mL) were added and the reaction was heated to 90 °C for 16 h. The reaction was cooled to room temperature, partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, purified via column chromatography: eluting with gradient hexanes/EtOAc (0% to 40% EtOAc), column: silica 4g, to provide 3-bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazine (690 mg, 79%) as an off-white fluffy solid.

[0227] Step 2: 3-Bromo-6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazine (50 mg, 0.12 mmol), tert-butyl 4-aminopiperidine-1-carboxylate (36 mg, 0.18 mmol), potassium *t*-butoxide (41 mg, 0.36 mmol), *t*BuX-Phos-Pd-G3 (10 mg, 0.012 mmol) were mixed in a Schlenk tube. The reaction was degassed with Ar then dry THF (2 mL) was added and the reaction was heated to 80 °C for 12 h. The reaction was cooled to room temperature, then partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated under vacuum, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 15% MeOH), column: silica 4g, to provide tert-butyl 4-((6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazin-3-yl)amino)piperidine-1-carboxylate (45 mg, 70%) as a tan solid.

[0228] Step 3: A solution of 4-((6-(2-methoxy-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)pyridazin-3-yl)amino)piperidine-1-carboxylate (45 mg, 0.084 mmol) in 2 mL of dry CH₂Cl₂ was cooled in an ice-water bath. Boron tribromide 1.0 M in CH₂Cl₂ (0.42 mL, 0.42 mmol) was added and the reaction mixture was stirred at room temperature for 16 hours. The reaction was quenched with 2 mL of MeOH, stirred for 30 min, then concentrated and purified using preparative HPLC to provide 2-(6-(piperidin-4-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol (21 mg, 73%) as a yellowish solid.

LC-MS: 377 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 8.95 (s, 2H), 8.25 (d, *J*=9.6 Hz, 1H), 8.09 (s, 2H), 7.69 (d, *J*=8.8 Hz, 1H), 7.49-7.41 (m, 1H), 7.21-7.24 (m, 2H), 4.11-4.15 (m, 1H), 3.34-3.39 (m, 4H), 3.01-3.08 (m, 2H), 2.11-2.17 (m, 2H), 1.79 (q, *J*=9.2 Hz, 2H)

[0229] Using the procedure described for Example 5 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
426	LC-MS: 351 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.22-9.28 (m, 2H), 8.34 (d, <i>J</i> =9.6Hz, 1H), 8.06-8.11 (m, 2H), 7.85-7.89 (m, 1H), 7.74 (d, <i>J</i> =8.2 Hz, 1H), 7.20-7.27 (m, 2H), 4.67-4.75 (m, 2H), 3.35-3.39 (m, 2H), 3.07-3.09 (m, 2H), 3.05 (s, 3H), 2.21 (q, <i>J</i> =10.2Hz, 2H), 1.83-1.88 (m, 2H)
432	LC-MS: 338 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.72 (d, <i>J</i> =9.6Hz, 1H), 8.41-8.54 (m, 2H), 7.91-8.08 (m, 1H), 7.82 (d, <i>J</i> =8.0Hz, 1H), 7.49 (d, <i>J</i> =8.2 Hz, 1H), 7.38 (s, 1H), 5.51 (quint, <i>J</i> =2.8Hz, 1H), 3.30-3.46 (m, 2H), 3.28-3.31 (m, 2H), 2.33-2.38 (m, 2H), 2.25-2.32 (m, 2H), (3 hydrogens corresponding to OH and NH unobserved)
436	LC-MS: 366 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.17-9.30 (m, 1H), 8.81-8.95 (m, 1H), 8.49 (d, <i>J</i> =9.8 Hz, 1H), 8.17 (s, 2H), 7.94 (d, <i>J</i> =8.8 Hz, 1H), 7.46 (d, <i>J</i> =9.8 Hz, 1H), 7.18-7.29 (m, 2H), 5.47 (spt, <i>J</i> =5.7 Hz, 1H), 3.31-3.52 (m, 2H), 2.41 (dd, <i>J</i> =12.9, 5.7 Hz, 2H), 1.68 (q, <i>J</i> =12.0 Hz, 2H), 1.34 (d, <i>J</i> =6.3 Hz, 6H), (1 hydrogen corresponding to OH or NH unobserved)
439	LC-MS: 363 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.07 (s, 2H), 8.28 (d, <i>J</i> =10Hz, 1H), 1.7 (s, 2H), 7.81 (d, <i>J</i> =8.8Hz, 1H), 7.73 (d, <i>J</i> =9.6Hz, 1H), 7.19-7.22 (m, 2H), 3.75-3.81 (m, 4H), 3.58-3.69 (m, 4H), 1.85-1.89 (m, 4H), (1 hydrogen corresponding to OH or NH unobserved)

(continued)

Cpd	Data
459	LC-MS: 349 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 10.53 (s, 1H), 10.40 (s, 1H), 9.79 (br s, 1H), 9.67 (br s, 1H), 8.28 (d, <i>J</i> =9.6Hz, 1H), 8.06 (s, 2H), 7.65 (d, <i>J</i> =9.6Hz, 1H), 7.58 (d, <i>J</i> =8.4Hz, 1H), 7.21-7.25 (m, 1H), 4.53 (q, <i>J</i> =14Hz, 2H), 3.64 (d, <i>J</i> =12.8Hz, 1H), 3.52 (d, <i>J</i> =12.8Hz, 1H), 3.37-3.40 (m, 2H), 3.15-3.24 (m, 2H), 1.92-2.02 (m, 2H)

Example 6 (Compound 430)

6-[2,3-Difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

[0230] Step 1: To a RBF, equipped with a N₂ inlet, were added: 6-bromo-N-methyl-N-(2,2,6,6-tetramethyl-4-piperidyl)pyridazin-3-amine (90 mg, 0.28 mmol), 4-[2,3-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]-1-tetrahydropyran-2-yl-pyrazole (128.8 mg, 0.33 mmol), Pd(PPh₃)₄ (31.8 mg, 0.03 mmol), Na₂CO₃ (87.43 mg, 0.83 mmol). The reaction was degassed with N₂ for 15 min then dioxane (8 mL) and water (2 mL) were added. The reaction was heated to 90 °C for 16 h, then cooled to room temperature, and partitioned between EtOAc and water. The organic layers were dried over Na₂SO₄, concentrated on a rotavap, and purified via column chromatography: eluting with gradient CH₂Cl₂/MeOH (0% to 30%), column: silica 4g to provide 6-(2,3-Difluoro-4-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (88 mg, 63%) as a grey solid.

[0231] Step 2: 6-[2,3-Difluoro-4-(1-tetrahydropyran-2-ylpyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethyl-4-piperidyl)pyridazin-3-amine (88 mg, 0.17 mmol) was dissolved in MeOH (2 mL) and 4M HCl in dioxane (90 μL, 0.34 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 30 min until LC-MS showed complete consumption of starting material. The reaction mixture was concentrated under reduced pressure to provide 6-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride salt. LC-MS: 427 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.21-9.33 (m, 1H), 8.27-8.43 (m, 1H), 8.20 (s, 2H), 8.01 (d, *J*=9.8 Hz, 1H), 7.60-7.79 (m, 3H), 5.01-5.14 (m, 1H), 3.05 (s, 3H), 2.10 (td, *J*=13.2, 1.6 Hz, 2H), 1.79 (dd, *J*=12.9, 4.1 Hz, 2H), 1.55 (s, 6H), 1.50 (s, 6H)

[0232] Using the procedure described for Example 6 above, additional compounds described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
426	LC-MS: 427 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.22-9.38 (m, 1H), 8.33-8.45 (m, 1H), 8.21 (d, <i>J</i> =1.9 Hz, 2H), 7.97 (dd, <i>J</i> =10.4, 2.8 Hz, 1H), 7.86 (d, <i>J</i> =5.7 Hz, 1H), 7.81 (dd, <i>J</i> =11.7, 6.6 Hz, 1H), 7.54-7.63 (m, 1H), 5.09-5.16 (m, 1H), 3.03 (s, 3H), 2.09 (t, <i>J</i> =12.9 Hz, 2H), 1.78 (dd, <i>J</i> =12.9, 3.8 Hz, 2H), 1.55 (s, 6H), 1.51 (s, 6H)
431	LC-MS: 414 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.24-9.35 (m, 1H), 8.43-8.58 (m, 1H), 8.22 (d, <i>J</i> =1.6 Hz, 2H), 8.06 (dd, <i>J</i> =9.1, 1.9 Hz, 1H), 7.87 (dd, <i>J</i> =12.1, 6.5 Hz, 1H), 7.83 (dd, <i>J</i> =11.5, 6.5 Hz, 1H), 7.37 (d, <i>J</i> =9.5 Hz, 1H), 5.77 (tt, <i>J</i> =10.7, 4.1 Hz, 1H), 2.33 (dd, <i>J</i> =13.1, 3.9 Hz, 2H), 1.84 (dd, <i>J</i> =12.9, 11.3 Hz, 2H), 1.54 (s, 6H), 1.52 (s, 6H)
435	LC-MS: 378 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.57-8.58 (m, 1H), 8.55 (d, <i>J</i> =5.0 Hz, 1H), 8.45 (s, 2H), 8.05-8.10 (m, 1H), 7.95-7.99 (m, 1H), 7.86 (s, 1H), 7.82 (d, <i>J</i> =9.5 Hz, 1H), 5.79 (tt, <i>J</i> =10.4, 4.4 Hz, 1H), 2.53 (dd, <i>J</i> =14.5, 4.5 Hz, 2H), 1.86-2.01 (m, 2H), 1.66 (s, 6H), 1.60 (s, 6H), (2 hydrogens corresponding to NH unobserved)
437	LC-MS: 396 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.28-9.44 (m, 1H), 8.47-8.61 (m, 1H), 8.25 (s, 2H), 8.01 (dd, <i>J</i> =9.3, 2.0 Hz, 1H), 7.92 (t, <i>J</i> =8.2 Hz, 1H), 7.69 (dd, <i>J</i> =12.8, 1.7 Hz, 1H), 7.65 (dd, <i>J</i> =8.0, 1.7 Hz, 1H), 7.34 (d, <i>J</i> =9.1 Hz, 1H), 5.76 (tt, <i>J</i> =10.7, 4.1 Hz, 1H), 2.33 (dd, <i>J</i> =13.2, 3.9 Hz, 2H), 1.85 (dd, <i>J</i> =13.2, 10.1 Hz, 2H), 1.54 (s, 6H), 1.53 (s, 6H)
440	LC-MS: 359 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: .06 (d, <i>J</i> =10.1 Hz, 1H), 7.78 (dd, <i>J</i> =9.5, 6.3 Hz, 1H), 7.32 (d, <i>J</i> =9.8 Hz, 1H), 6.58-6.76 (m, 2H), 5.11 (tt, <i>J</i> =12.0, 4.1 Hz, 1H), 3.02 (s, 3H), 1.73 (dd, <i>J</i> =12.9, 3.5 Hz, 2H), 1.63 (t, <i>J</i> =12.5 Hz, 2H), 1.43 (s, 6H), 1.27 (s, 6H), (1 hydrogen corresponding to NH unobserved)

(continued)

Cpd	Data
441	LC-MS: 364 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 1.33 (d, <i>J</i> =6.62 Hz, 6H) 1.65 (q, <i>J</i> =12.40 Hz, 2H) 2.42 (d, <i>J</i> =12.61 Hz, 2H) 3.38 - 3.50 (m, 2H) 5.48 (s, 1H) 6.51 - 6.63 (m, 1H) 7.42 - 7.57 (m, 3H) 7.80 (d, <i>J</i> =1.58 Hz, 1H) 8.09 (d, <i>J</i> =8.83 Hz, 1H) 8.50 (d, <i>J</i> =9.46 Hz, 1H) 8.61 (d, <i>J</i> =2.52 Hz, 1H) 8.76 (br s., 1H) 9.12 (br s., 1H) 13.18 (s, 1H)
442	LC-MS: 379 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 8.42 (s, 1H), 8.28 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 8.03 (dd, <i>J</i> =9.1, 1.6 Hz, 1H), 7.95 (s, 1H), 7.67 (d, <i>J</i> =9.1 Hz, 1H), 7.14 (d, <i>J</i> =9.1 Hz, 1H), 5.03-5.17 (m, 1H), 4.19 (s, 3H), 2.94 (s, 3H), 1.49-1.58 (m, 2H), 1.37-1.48 (m, 2H), 1.27 (s, 6H), 1.10 (s, 6H), (1 hydrogen corresponding to NH unobserved)
443	LC-MS: 366 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.35 (s, 1H), 8.31 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 8.11 (d, <i>J</i> =9.1 Hz, 1H), 7.99 (dd, <i>J</i> =9.1, 1.9 Hz, 1H), 7.75 (dt, <i>J</i> =9.1, 0.9 Hz, 1H), 7.21 (d, <i>J</i> =9.5 Hz, 1H), 5.81 (tt, <i>J</i> =11.2, 4.1 Hz, 1H), 4.27 (s, 3H), 2.28 (dd, <i>J</i> =12.6, 4.1 Hz, 2H), 1.48 (t, <i>J</i> =12.3 Hz, 2H), 1.41 (s, 6H), 1.30 (s, 6H), (1 hydrogen corresponding to NH unobserved)
445	LC-MS: 379 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 8.52 (dt, <i>J</i> =7.0, 1.1 Hz, 1H), 8.13 (dt, <i>J</i> =8.8, 1.3 Hz, 1H), 7.66 (d, <i>J</i> =9.8 Hz, 1H), 7.28 (ddd, <i>J</i> =9.0, 6.8, 1.3 Hz, 1H), 7.13 (d, <i>J</i> =9.5 Hz, 1H), 6.89 (td, <i>J</i> =6.9, 1.4 Hz, 1H), 5.22 (tt, <i>J</i> =12.3, 3.5 Hz, 1H), 3.04 (s, 3H), 2.63 (s, 3H), 1.67 (dd, <i>J</i> =12.3, 3.5 Hz, 2H), 1.56 (t, <i>J</i> =12.3 Hz, 2H), 1.36 (s, 6H), 1.18 (s, 6H), (1 hydrogen corresponding to NH unobserved)
446	LC-MS: 352 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 9.14 (dd, <i>J</i> =1.6, 0.9 Hz, 1H), 7.98 (dd, <i>J</i> =9.5, 1.9 Hz, 1H), 7.94 (t, <i>J</i> =0.9 Hz, 1H), 7.89 (d, <i>J</i> =9.5 Hz, 1H), 7.63 (dt, <i>J</i> =9.5, 0.8 Hz, 1H), 7.60 (d, <i>J</i> =1.3 Hz, 1H), 7.14 (d, <i>J</i> =9.5 Hz, 1H), 5.34 (tt, <i>J</i> =12.6, 3.5 Hz, 1H), 1.67 (dd, <i>J</i> =12.3, 3.8 Hz, 2H), 1.55 (t, <i>J</i> =12.1 Hz, 2H), 1.36 (s, 6H), 1.17 (s, 6H), (1 hydrogen corresponding to NH unobserved)
447	LC-MS: 394 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 8.47 (d, <i>J</i> =2.5 Hz, 1H), 8.05 (d, <i>J</i> =9.1 Hz, 1H), 7.98 (d, <i>J</i> =8.5 Hz, 1H), 7.76 (d, <i>J</i> =1.9 Hz, 1H), 7.70 (d, <i>J</i> =2.2 Hz, 1H), 7.59 (dd, <i>J</i> =8.5, 2.2 Hz, 1H), 7.06 (d, <i>J</i> =9.1 Hz, 1H), 6.57 (dd, <i>J</i> =2.4, 1.7 Hz, 1H), 5.85 (tt, <i>J</i> =11.3, 4.1 Hz, 1H), 2.21 (dd, <i>J</i> =12.1, 3.9 Hz, 2H), 2.04-2.08 (m, 2H), 1.35 (s, 6H), 1.18 (s, 6H), (2 hydrogens corresponding to OH and NH unobserved)
452	LC-MS: 409 [M+H] ⁺ . ¹ H NMR (500 MHz, acetone- <i>d</i> ₆) δ: 7.59 (dd, <i>J</i> =8.5, 6.0 Hz, 1H), 7.46 (s, 2H), 7.31 (dd, <i>J</i> =10.1, 2.5 Hz, 1H), 7.15 (td, <i>J</i> =8.5, 2.5 Hz, 1H), 7.04 (d, <i>J</i> =9.5 Hz, 1H), 6.91 (d, <i>J</i> =9.5 Hz, 1H), 5.23 (tt, <i>J</i> =12.3, 2.8 Hz, 1H), 3.00 (s, 3H), 1.65 (dd, <i>J</i> =12.3, 3.5 Hz, 2H), 1.51 (t, <i>J</i> =12.1 Hz, 2H), 1.33 (s, 6H), 1.15 (s, 6H), (2 hydrogens corresponding to NH unobserved)
455	LC-MS: 344 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.17 (d, <i>J</i> =9.5 Hz, 1H), 7.66 (d, <i>J</i> =8.8 Hz, 1H), 7.19 (d, <i>J</i> =9.5 Hz, 1H), 6.44 (dd, <i>J</i> =8.7, 2.4 Hz, 1H), 6.41 (d, <i>J</i> =2.5 Hz, 1H), 5.71 (tt, <i>J</i> =11.0, 3.8 Hz, 1H), 2.23 (dd, <i>J</i> =12.6, 4.1 Hz, 2H), 1.40-1.47 (m, 2H), 1.39 (s, 6H), 1.27 (s, 6H), (3 hydrogens corresponding to OH and NH unobserved)
456	LC-MS: 425 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 9.23-9.39 (m, 1H), 8.39-8.52 (m, 1H), 8.26 (s, 2H), 8.00-8.08 (m, 1H), 7.93 (s, 1H), 7.81-7.85 (m, 1H), 7.78 (d, <i>J</i> =7.9 Hz, 1H), 7.63 (d, <i>J</i> =8.2 Hz, 1H), 4.95-5.03 (m, 1H), 3.08 (s, 3H), 2.13 (t, <i>J</i> =12.9 Hz, 2H), 1.81 (dd, <i>J</i> =12.9, 3.5 Hz, 2H), 1.56 (s, 6H), 1.52 (s, 6H)
458	LC-MS: 414 [M+H] ⁺ . ¹ H NMR (500 MHz, DMSO- <i>d</i> ₆) δ: 8.70-9.36 (m, 1H), 8.29 (s, 2H), 7.85 (d, <i>J</i> =9.1 Hz, 1H), 7.60 (d, <i>J</i> =9.5 Hz, 1H), 7.33-7.39 (m, 1H), 5.59-5.68 (m, 1H), 2.24-2.33 (m, 2H), 1.72-1.82 (m, 2H), 1.50 (s, 6H), 1.48 (s, 6H), (2 hydrogens corresponding to NH unobserved)

Example 7 (Compound 434)

6-[2-Methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride

[0233] 6-(2-Methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (6 mg) was dissolved in 1 mL of 4N HCl/dioxane and the resulting mixture was stirred at RT for 20 minutes. The mixture was evaporated and triturated with ether to provide 6-(2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine 4 mg, 75%).

LC-MS: 422 [M+H]⁺. ¹H NMR (500 MHz, methanol-*d*₄) δ: 8.42 - 8.47 (m, 1H), 8.33 - 8.42 (m, 2H), 8.09 - 8.17 (m, 1H), 7.99 - 8.08 (m, 1H), 7.50 - 7.56 (m, 1H), 5.0 (m, 1H), 4.16 (s, 3H), 3.20 (s, 3H), 1.94 - 2.15 (m, 4H), 1.65 (s, 6H), 1.57 (s, 6H)

Example 8 (Compound 460)

3-{6-[Methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride

[0234] Step 1: 6-Chloro-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (280 mg, 1.0 mmol) was dissolved in 10 mL, a mixture of dioxane and water (4:1). To the solution was added sodium carbonate (190 mg, 3 eq), (6-chloro-2-methoxypyridin-3-yl)boronic acid (190 mg, 1 eq) and Pd(Ph₃P)₄ (70 mg, 0.1 eq). The reaction mixture was stirred at 90 °C for 16 h, then applied to a column and purified using DCM/MeOH to yield (133 mg, 55%) 6-(6-chloro-2-methoxypyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine.

LC-MS: 390 [M+H]⁺. ¹H NMR (500 MHz, DMSO-*d*₆) δ: 8.25 (d, J=7.88 Hz, 1H), 7.83 (d, J=9.77 Hz, 1H), 7.24 (d, J=7.88 Hz, 1H), 7.11 (d, J=9.77 Hz, 1H), 5.00 - 5.24 (m, 1H), 3.93 (s, 3 H), 2.87 - 2.96 (m, 3H), 1.40 - 1.57 (m, 4H), 1.21 - 1.29 (m, 7H), 1.10 (br s, 6H)

[0235] Step 2: 6-(6-Chloro-2-methoxypyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (90 mg) was dissolved in a mixture of dioxane and water (4:1, 5 mL). To the solution was added sodium carbonate (100 mg, 3 eq), 1-(tetrahydro-2H-pyran-2-yl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (70 mg, 1.1 eq) and Pd(Ph₃P)₄ (27 mg, 0.1 eq). The reaction mixture was stirred at 90 °C for 16 h, then purified by column chromatography using DCM/MeOH to yield 80 mg (68%) of 6-(2-methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine.

[0236] Step 3: 6-(2-Methoxy-6-(1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-4-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine (45 mg) was dissolved in 4 mL of 4N HCl/dioxane and the resulting mixture was heated for 2 hours at 60 °C. The resulting mixture was evaporated and triturated with ether to provide 3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-6-(1H-pyrazol-4-yl)pyridin-2-ol (24 mg, 61%).

LC-MS: 408 [M+H]⁺; ¹H NMR (500 MHz, methanol-*d*₄) δ: 8.62 - 8.69 (m, 1H), 8.55 - 8.61 (m, 1H), 8.39 (s, 2H), 7.95 - 8.06 (m, 1H), 7.04 - 7.14 (m, 1H), 5.02 - 5.24 (m, 1H), 3.18 (s, 3H), 2.03 (s, 4H), 1.67 (s, 6H), 1.57 (s, 6H)

[0237] Using the procedure described for Example 8 above, an additional compound described herein may be prepared by substituting the appropriate starting materials reagents, and reaction conditions, obtaining compounds such as those selected from:

Cpd	Data
461	LC-MS: 395 [M+H] ⁺ . ¹ H NMR (500 MHz, methanol- <i>d</i> ₄) δ: 8.46 (d, J=9.5 Hz, 1H), 8.33-8.42 (m, 1H), 8.29 (d, J=7.6 Hz, 1H), 8.10-8.23 (m, 1H), 7.20 (d, J=9.1 Hz, 1H), 6.86 (d, J=7.3 Hz, 1H), 5.84 (tt, J=10.4, 3.8 Hz, 1H), 2.49 (dd, J=13.9, 4.1 Hz, 2H), 1.86 (dd, J=13.9, 10.4 Hz, 2H), 1.65 (s, 6H), 1.56 (s, 6H), (3 hydrogens corresponding to OH and NH unobserved)

Biological Examples

[0238] The following *in vitro* biological examples demonstrate the usefulness of the compounds of the present description for treating Huntington's disease.

[0239] To describe in more detail and assist in understanding the present description, the following non-limiting biological examples are offered to more fully illustrate the scope of the description and are not to be construed as specifically limiting the scope thereof. Such variations of the present description that may be now known or later developed, which would be within the purview of one skilled in the art to ascertain, are considered to fall within the scope of the present description and as hereinafter claimed.

Example 1**Development and Validation of Meso Scale Discovery (MSD) Assay to Quantify Mutant and Total Huntingtin Protein (HTT) - Best Pair Analysis for Endogenous HTT Detection**

[0240] To identify orally bioavailable small molecules useful for HTT reduction, high throughput ELISA methods to detect and measure both total HTT (wild-type and mutant)(tHTT) and mutant HTT (mHTT) in cells and tissues from HD patients and animal models are needed. The method described herein permits relatively high throughput protein measurement of tHTT and mHTT levels in cells and tissues from HD patients and in HD animal models. The ELISA method described herein enables screening of molecules for their ability to decrease the level of HTT and to identify and optimize small molecules for use in treating HD

[0241] To determine the combination of antibodies that detect HTT (wild type and mutant) in a sensitive, high throughput

format, multiple commercially available antibodies (Table 1) were screened in capture/detection combinations.

Protocol for Best Pair Analysis for Antibody Comparison in Detecting Total and Mutant HTT in Cells

[0242]

1. Cells were cultured in a cell culture-compatible 96-well plate.
2. The medium was removed and 50-100 μ L of lysis buffer per well (composition described below) was added to cells to provide a "cell lysate". The plate was placed on a shaker at 4° C for 30 minutes, then stored at -20° C.
3. A Capture Antibody Diluent was prepared by diluting the capture antibody in PBS for standard plates as described in the plate layout, then 30 μ L of the Capture Antibody Diluent was added per well. The plate was sealed and incubated at 4° C overnight.
4. The plate was washed 3X with 200 μ L of Wash Buffer. Blocking Buffer (150 μ L per well) was added, then the plate was sealed and incubated on a plate shaker for 3-4 hours at room temperature (RT).
5. The Blocking Buffer was decanted and the plate was washed 3X with 200 μ L of Wash Buffer. The cell lysate (25 μ L per well) was then added. The plate was sealed and incubated at 4° C overnight.
6. A First Detection Antibody Diluent was prepared by diluting a first detection antibody for pairing in an Assay Antibody Diluent. The plate was washed 3X with 200 μ L of Wash Buffer, then 25 μ L of the First Detection Antibody Diluent was added per well. The plate was sealed and incubated on a plate shaker for 1 hour at RT.
7. A Second Detection Antibody Diluent was prepared by diluting a second detection antibody for pairing in an Assay Antibody Diluent. The plate was washed 3X with 200 μ L of Wash Buffer, then 25 μ L of the Second Detection Antibody Diluent was added per well. The plate was sealed and incubated on a plate shaker for 1 hour at RT.
8. The Read Buffer was diluted 4X. The plate was washed 3X with 200 μ L of Wash Buffer, then 150 μ L of the Read Buffer was added per well. The plate were read immediately following.

[0243] The various commercially available antibodies screened in the Best Pair Analysis, providing different capture/detection combinations are shown in Table 1. The capture/detection antibody pairs evaluated are shown in Table 2. The two antibody pairs that gave the maximum spread of values (higher signal to noise ratio) were selected for polyglutamine-expanded (mHTT) and total human HTT detection.

Table 1

Species	ID	Antibody Epitope
monoclonal mouse	MAB2166	Huntingtin fragment from aa 181-810 as a fusion protein
monoclonal rat	MAB2174	Huntingtin fusion protein 549-679, fused to Glutathione S-transferase
monoclonal mouse	MAB5374	GST fusion protein from the first 256 amino acids from human huntingtin with the deletion of the polyglutamine tract
goat polyclonal	sc-8767	Peptide mapping near the N-terminus of Huntingtin of human origin (N18)
monoclonal mouse	P1874	GST human Huntingtin (N-terminal fragment of 171 amino acids containing 65Q)
monoclonal mouse	MAB1574	Homopolymeric glutamine stretch
monoclonal rabbit	#5656S	Synthetic peptide corresponding to residues surrounding Pro1220 of human huntingtin protein
monoclonal mouse	MW1	DRPLA-19Q

Table 2

Pair	HTT MSD	ID	Signal to Noise ratio
Capture/Detection	Total	MAB2166/MAB2174	95
Capture/Detection	Total	MAB2166/sc-8767	6
Capture/Detection	Total	sc-8767/#5656	5
Capture/Detection	Total	MAB2166/#5656S	483

(continued)

Pair	HTT MSD	ID	Signal to Noise ratio
Capture/Detection	Mutant	MAB5374/#5656S	8
Capture/Detection	Mutant	MAB1574/#5656S	25
Capture/Detection	Mutant	P1874/#5656S	20
Capture/Detection	Mutant	MW1/#5656S	45

[0244] Characteristics for Best Pair capture/detection antibody pairs are those combinations that give minimum background and maximum signal. Such highly sensitive ELISA assays would be useful to detect HTT in the context of drug discovery for screening agents (small and large molecules and gene therapies) and for use in detecting HTT as a biomarker in clinical studies. Such capture/detection antibody pairs for quantifying total HTT are selected from MAB2166/MAB2174 and MAB2166/#5656S. Such capture/detection antibody pairs for quantifying mutant HTT are selected from MAB 1574/#5656S, P1874/#5656S and MW1/#5656S. The antibody pair for mHTT (MW1/#5656S) and the other for total HTT (MAB2166/#5656S) from this analysis were further optimized and validated to provide high-throughput assays that enabled HTT detection in patient fibroblasts and lymphocytes.

[0245] The Endogenous Huntingtin Protein assay used in Example 2 was developed using the Best Pair antibodies identified in Table 2, providing highly sensitive, high-throughput detection assays for polyglutamine-expanded (mHTT) and total human HTT in multiple cell systems. The platform used for the assay is the ELISA-based Meso Scale Discovery (MSD) electrochemiluminescence assay platform.

Example 2

Endogenous Huntingtin Protein Assay

[0246] Meso Scale Discovery (MSD) 96-well or 384-well plates were coated overnight at 4°C with MW1 (expanded polyglutamine) or MAB2166 monoclonal antibody (for capture) at a concentration of 1 µg/mL in PBS (30 µL per well). Plates were then washed three times with 300 µL wash buffer (0.05% Tween-20 in PBS) and blocked (100 µL blocking buffer; 5% BSA in PBS) for 4-5 hours at room temperature with rotational shaking and then washed three times with wash buffer.

[0247] Samples (25 µL) were transferred to the antibody-coated MSD plate and incubated overnight at 4°C. After removal of the lysates, the plate was washed three times with wash buffer, and 25 µL of #5656S (Cell signaling; rabbit monoclonal) secondary antibody (diluted to 0.25 µg/mL in 0.05% Tween-20 in blocking buffer) was added to each well and incubated with shaking for 1 hour at room temperature. Following incubation with the secondary antibody, the wells were rinsed with wash buffer after which 25 µL of goat anti-rabbit SULFO TAG secondary detection antibody (required aspect of the MSD system) (diluted to 0.25 µg/mL in 0.05% Tween-20 in blocking buffer) was added to each well and incubated with shaking for 1 hour at room temperature. After rinsing three times with wash buffer, 150 µL of read buffer T with surfactant (MSD) were added to each empty well, and the plate was imaged on a SI 6000 imager (MSD) according to manufacturers' instructions provided for 96- or 384-well plates. The resulting IC₅₀ values (µM) for compounds tested are shown in Table 3.

[0248] As shown in Table 3, test compounds described herein had the following IC₅₀ values, an IC₅₀ value between > 3 µM and ≤ 9 µM is indicated by a single star (*), an IC₅₀ value between > 1 µM and ≤ 3 µM is indicated by two stars (**), an IC₅₀ value between > 0.5 µM and ≤ 1 µM is indicated by three stars (***), an IC₅₀ value between > 0.1 µM and ≤ 0.5 µM is indicated by four stars (****) and an IC₅₀ value of ≤ 0.1 µM is indicated by five stars (*****).

Table 3


	Cpd	IC ₅₀	Cpd	IC ₅₀	Cpd	IC ₅₀
5	16	*	413	*****	441	*
	26	*****	414	*****	442	*
	31	*****	415	*****	443	*
10	460	*	416	*****	444	*
	39	*	417	*****	445	*
	44	*****	418	*****	446	*
	46	*****	419	*****	447	*
15	47	*****	420	*****	448	*
	48	*****	421	*****	449	*
	51	*****	422	*****	450	*
20	63	*****	423	*****	451	*
	64	*****	424	*****	451	*
	170	*****	425	*****	452	*
25	176	*****	426	*****	454	*
	Cpd	IC ₅₀	Cpd	IC ₅₀	Cpd	IC ₅₀
	200	*****	427	*****	455	*
30	207	*****	428	*****	456	*
	212	*****	429	*****	456	*
	218	*****	430	*****	456	*
35	258	*****	431	*****	457	*
	307	*****	432	*****	458	*
	315	*****	433	*****	459	*
40	318	*****	434	*****	460	*
	348	*****	435	*****	461	*
	350	*****	436	*****	462	*
45	352	*****	437	*****	463	*
	393	*****	438	*****	464	*
	411	*****	439	*****	465	*
	412	*****	440	*		

[0249] Without regard to whether a document cited herein was specifically and individually indicated as being incorporated by reference, all documents referred to herein are incorporated by reference into the present application for any and all purposes to the same extent as if each individual reference was fully set forth herein.

[0250] Having now fully described the subject matter of the claims, it will be understood by those having ordinary skill in the art that the same can be performed within a wide range of equivalents without affecting the scope of the subject matter or embodiments described herein. It is intended that the appended claims be interpreted to include all such equivalents.

[0251] Further disclosed are the following embodiments:

1. A method of use of a compound for treating or ameliorating HD in a subject in need thereof comprising administering to the subject an effective amount of a compound of Formula (I):



(I)

or a form thereof, wherein

W is CH=CH or S;

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₂ is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-


C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy;

R₄ is independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkylamino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino; and

R₅ is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

2. The method of embodiment 1, wherein the compound of Formula (I) is selected from a compound of Formula (Ia) and Formula (Ib):



or a form thereof, wherein

X is CH₂, CH(C₁₋₄alkyl), C(C₁₋₄alkyl)₂, CH=CH, O, NR₅, or a bond;

A is aryl, heteroaryl, heterocyclyl, or C₉₋₁₀cycloalkyl,

wherein aryl is selected from phenyl and naphthyl, each optionally substituted with 1, 2, 3, or 4 substituents each selected from R₁,

wherein heteroaryl is a saturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₁,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or tricyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂, and

wherein C₉₋₁₀cycloalkyl is a saturated or partially unsaturated bicyclic ring system optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₂;

B is heterocyclyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic, bicyclic or polycyclic ring system having 1, 2, or 3 heteroatom ring members independently selected from N, O, or S, each optionally substituted with 1, 2, 3, 4, or 5 substituents each selected from R₄;

R₁ is halogen, hydroxyl, cyano, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R_2 is halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, (C_{1-4} alkyl) $_2$ -amino, amino- C_{1-4} alkyl, amino-carbonyl, hydroxyl- C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, C_{2-4} alkenyl, C_{3-7} cycloalkyl, or heterocyclyl- C_{1-4} alkyl;

R_3 is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, (C_{1-4} alkyl) $_2$ -amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl, (C_{1-4} alkyl) $_2$ -amino- C_{1-4} alkyl, amino-carbonyl, C_{1-4} alkyl-amino-carbonyl, C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl-amino, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, C_{1-4} alkyl-carbonyl, C_{1-4} alkoxy, halo- C_{1-4} alkoxy, amino- C_{1-4} alkoxy, hydroxyl- C_{1-4} alkoxy, C_{1-4} alkyl- C_{1-4} alkoxy, C_{1-4} alkyl-amino- C_{1-4} alkoxy, (C_{1-4} alkyl) $_2$ -amino- C_{1-4} alkoxy, C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkoxy, C_{1-4} alkoxy- C_{1-4} alkoxy, C_{1-4} alkoxy-carbonyl, C_{1-4} alkoxy-carbonyl-amino, C_{1-4} alkoxy-carbonylamino- C_{1-4} alkoxy, C_{2-4} alkenyl, C_{2-4} alkenyl-amino-carbonyl, C_{3-7} cycloalkyl, C_{3-7} cycloalkyl- C_{1-4} alkoxy, C_{3-7} cycloalkenyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino, heteroaryl- C_{1-4} alkyl-amino-carbonyl- C_{1-4} alkyl, heteroaryl- C_{1-4} alkyl-carbonyl-amino- C_{1-4} alkyl, heterocyclyl, heterocyclyl- C_{1-4} alkyl, phenyl, or phenyl- C_{1-4} alkoxy;

R_4 is independently selected from halogen, C_{1-4} alkyl, hydroxyl- C_{1-4} alkyl, amino, C_{1-4} alkylamino, (C_{1-4} alkyl) $_2$ -amino or hydroxyl- C_{1-4} alkyl-amino; and

R_5 is hydrogen, C_{1-4} alkyl, or hydroxyl- C_{1-4} alkyl;


wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

3. The method of embodiment 1, wherein:


X is O, NH, $N(CH_3)$ or a bond;


A is aryl, heteroaryl or heterocyclyl,


wherein, aryl is selected from the group consisting of:



wherein, heteroaryl is selected from the group consisting of:









a36; and


wherein, heterocyclyl is selected from the group consisting of:




a37




a38




a39




a40




a41




and
a42




a43




a44



a45




a46, and




a47


B is heterocyclyl selected from the group consisting of:




b1




b2



b3



b4



R_{1a} and R_{1b} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, C_{1-4} alkyl, halo- C_{1-4} alkyl, amino, C_{1-4} alkyl-amino, $(C_{1-4}alkyl)_2$ -amino, amino- C_{1-4} alkyl, C_{1-4} alkyl-amino- C_{1-4} alkyl, $(C_{1-4}alkyl)_2$ -amino- C_{1-4} alkyl, amino-carbonyl, C_{1-4} alkyl-amino-carbonyl,

(C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, heterocyclyl-C₁₋₄alkoxy, phenyl, or phenyl-C₁₋₄alkoxy,

wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and

wherein each instance of phenyl, heteroaryl or heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R_{2a}, R_{2b} and R_{2c} are each, where allowed by available valences, one or more substituents each selected from halogen, hydroxyl, cyano, oxo, hydroxyl-imino, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₂₋₄alkenyl, C₃₋₇cycloalkyl, or heterocyclyl-C₁₋₄alkyl,

wherein heterocyclyl is a saturated or partially unsaturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S, and


wherein each instance of heterocyclyl is optionally substituted with 1, or 2 substituents each selected from R₃;

R₃ is halogen, hydroxyl, nitro, oxo, hydroxyl-imino, C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino, amino-C₁₋₄alkyl, C₁₋₄alkyl-amino-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-C₁₋₄alkyl, amino-carbonyl, C₁₋₄alkyl-amino-carbonyl, (C₁₋₄alkyl)₂-amino-carbonyl, C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, (C₁₋₄alkyl)₂-amino-carbonyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl-amino, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, C₁₋₄alkyl-carbonyl, C₁₋₄alkoxy, halo-C₁₋₄alkoxy, amino-C₁₋₄alkoxy, hydroxyl-C₁₋₄alkoxy, C₁₋₄alkyl-C₁₋₄alkoxy, C₁₋₄alkyl-amino-C₁₋₄alkoxy, (C₁₋₄alkyl)₂-amino-C₁₋₄alkoxy, C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkoxy, C₁₋₄alkoxy-C₁₋₄alkoxy, C₁₋₄alkoxy-carbonyl, C₁₋₄alkoxy-carbonyl-amino, C₁₋₄alkoxy-carbonylamino-C₁₋₄alkoxy, C₂₋₄alkenyl, C₂₋₄alkenyl-amino-carbonyl, C₃₋₇cycloalkyl, C₃₋₇cycloalkyl-C₁₋₄alkoxy, C₃₋₇cycloalkenyl, heteroaryl, heteroaryl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino, heteroaryl-C₁₋₄alkyl-amino-carbonyl-C₁₋₄alkyl, heteroaryl-C₁₋₄alkyl-carbonyl-amino-C₁₋₄alkyl, heterocyclyl, heterocyclyl-C₁₋₄alkyl, phenyl, or phenyl-C₁₋₄alkoxy; and


R_{4a}, R_{4b}, R_{4c}, R_{4d}, R_{4e}, R_{4f} and R_{4g} are independently selected from halogen, C₁₋₄alkyl, hydroxyl-C₁₋₄alkyl, amino, C₁₋₄alkyl-amino, (C₁₋₄alkyl)₂-amino or hydroxyl-C₁₋₄alkyl-amino;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.



4. A compound selected from a compound of Formula (Ia11), Formula (Ia15), Formula (Ia18) or Formula (Ib1):



(Ia11),



(Ia15),



or a form thereof, wherein (when present),

X is selected from O, NR₅, or a bond;

A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy, wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a},

B is selected from 1H-pyrazolyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, (1*R*,5*S*)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl, 2,6-diazaspiro[3.4]octyl or 2,7-diazaspiro[3.5]nonyl, each optionally substituted with 1 or 2 substituents each selected from R_{4a};

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or heteroaryl, wherein heteroaryl is a saturated monocyclic or bicyclic ring system having 1, 2, or 3 heteroatom ring members selected from N, O, and S optionally substituted with 1 or 2 substituents each selected from R_{3a};

R_{3a} is selected from nitro or C₁₋₄alkyl; and,

R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

5. The compound of embodiment 4 wherein (when present),

R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy or 1H-pyrazolyl or 1H-imidazol-1-yl,

wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a};

R_{3a} is selected from nitro or methyl or amino; and,

R_{4a} is methyl or ethyl;

R_{5a} is hydrogen or methyl;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

6. A method of use of a compound or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering to the subject an effective amount of a compound selected from the group consisting of:

6-(naphthalen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

6-(benzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)-pyridazin-3-yl)phenol

2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[b]-thiophene-5-carbonitrile

6-(quinolin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

3-(benzo[b]-thiophen-2-yl)-6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)pyridazine

2-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)phenol

	6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)naphthalen-2-ol
	6-(benzo[b]-thiophen-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
5	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
	6-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline
10	N-methyl-6-(quinolin-7-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	N-methyl-6-(quinolin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
15	6-(isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(imidazo[1,2-a]pyridin-6-yl-pyridazin-3-yl)-methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
20	N-methyl-6-(6-phenylpyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(6-(1H-pyrrol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(6-(1H-pyrazol-1-yl)pyridin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
25	methyl-(6-quinoxalin-2-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
	methyl-(6-quinolin-3-yl-pyridazin-3-yl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amine
30	N-methyl-6-(phthalazin-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(benzo[c][1,2,5]oxa-diazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	6-(benzo[d]thiazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
35	6-(2-methylbenzo-[d]oxazol-6-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
40	5-chloro-2-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	3-(6-(2,2,6,6-tetramethylpiperidin-4-yl-amino)pyridazin-3-yl)naphthalen-2-ol
	5-chloro-2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)pyridazin-3-yl)phenol
45	4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
50	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol
	2-fluoro-6-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
	3,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
55	4,5-dimethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
	5-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

- 4,5-difluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5 3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
- 1-allyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
- 6-(benzo[b]thiophen-2-yl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)pyridazin-3-amine
- 10 N-allyl-3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzamide
- 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 15 5-(5-methyl-oxazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
- 5-(4-hydroxymethyl)-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 20 5-(1H-imidazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-(4-amino-1H-pyrazole-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
- 25 5-(3-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-(2-morpholinoethyl)-1H-pyrazol-4-yl)phenol
- 30 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
- 5-(5-amino-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 35 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol
- 2-((6-((2-hydroxy-ethyl)-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)-5-pyrazol-1-yl)phenol
- 2-(6-(piperidin-4-yloxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 40 2-(6-(((2S,4R,6R)-2,6-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 2-(6-(((2S,4S)-2-methylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 45 5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-yl-oxy)pyridazin-3-yl)phenol
- 2-(6-(((2S,4S)-2-methylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 50 (5-(1H-pyrazol-1-yl)-2-(6-(pyrrolidin-3-ylmethoxy)pyridazin-3-yl)phenol
- 2-(6-((3-fluoropiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 2-(6-(1,2,2,6,6-pentamethyl-piperidin-4-yl-oxy)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 55 5-1H-pyrazol-1-yl-2-(6-(2,2,6,6-tetramethylpiperidin-4-yl-oxy)-pyridazin-3-yl)phenol
- 5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol

EP 3 848 035 A1

	2-(6-piperazin-1-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	3-(6-(azetidin-3-ylamino)-pyridazin-3-yl)naphthalen-2-ol
5	2-(6-(azetidin-3-ylamino)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(3,5-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
10	2-(6-(7-methyl-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(1,4-diazepan-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(4-(2-hydroxyethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
15	2-(6-(3,6-diazabicyclo[3.2.1]octan-3-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(2,7-diazaspiro[3.5]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
20	2-(6-(3-(hydroxymethyl)piperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(1,7-diazaspiro[4.4]nonan-7-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(4-amino-4-methylpiperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
25	2-(6-(3-(dimethylamino)piperidin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(1,2,2,6,6-pentamethylpiperidin-4-ylamino)-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
30	2-(6-(3,3-dimethylpiperazin-1-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-(7-(2-hydroxyethyl)-2,7-diazaspiro[4.4]nonan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
35	3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol
	5-(1H-pyrazol-1-yl)-2-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)phenol
40	2-(6-piperidin-4-yl-pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalen-2-ol
	3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
45	3-(6-(2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
	3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
50	3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol
	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalene-2,7-diol
	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
55	3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
	tert-butyl 3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)carbamate

	7-(3-amino-propoxy)-3-(6-(methyl-(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)naphthalen-2-ol
5	N-(3-((7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-yl)oxy)propyl)acetamide
	7-(3-hydroxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	7-(3-methoxypropoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
10	7-(2-morpholinoethoxy)-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)naphthalen-2-ol
	3-(6-(piperidin-4-ylmethyl)pyridazin-3-yl)naphthalen-2-ol
15	5-(1H-pyrazol-1-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
	3-methoxy-2-(6-(methyl(2,2,6-trimethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
	2-(6-((6S)-6-((S)-1-hydroxyethyl)-2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
20	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2-naphthonitrile
	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(piperidinylmethyl)naphthalen-2-ol
25	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(pyrrolidinylmethyl)naphthalen-2-ol
	1-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
	1-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2,7-diol
30	7-methoxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalene-2-ol
	7-methoxy-3-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
35	7-(3,6-dihydro-2H-pyran-4-yl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-7-(tetrahydro-2H-pyran-4-yl)naphthalene-2-ol
40	7-(difluoromethyl)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	7-((4-hydroxy-2-methylbutan-2-yl)oxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
45	7-(3-hydroxy-3-methylbutoxy)-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)naphthalen-2-ol
	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)benzene-1,3-diol
50	3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(trifluoromethoxy)phenol
55	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol
	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)-3-(trifluoromethoxy)phenol

- 4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one
- 5 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
- 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimida-
zo[1,2-a]pyridin-3-yl)phenol
- 10 3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyridine-3-yl)phenol
- 5-(1-cyclopentyl-1H-pyrazol-4-yl)-3-methoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 15 3',5-dimethoxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-(1,1'-biphenyl)-3-ol
- 3-(benzyloxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
- 20 3-ethoxy-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5-methyloxazol-2-yl)phenol
- 3-(cyclopropylmethoxy)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-5-(5-methyloxa-
zol-2-yl)phenol
- 25 2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-benzo[d]imidazol-6-ol
- 5-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-(1H-pyrazol-1-yl)-2-(6-(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 30 3-hydroxy-4-(6-((2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzonitrile
- 2-(6-((2,2-dimethylpiperidin-4-yl)oxy)pyridazin-3-yl)-5-(1H-pyrazol-1-yl)phenol
- 35 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
- 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-
3-yl)phenol
- 40 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(4,5,6,7-tetrahydropyrazolo[1,5-
a]pyrazin-3-yl)phenol
- 4-(1H-indol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 45 4-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-3-yl)phenol
- 4-(4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
- 50 4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
- 4-(4-hydroxy-3-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol
- 55 5-(1H-indazol-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 4-chloro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol

- 4-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
- 5-fluoro-4-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-4-yl)phenol
- 5-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-5-yl)phenol
- 6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one
- 6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1,4-dihydroindeno[1,2-c]-1H-pyrazol-7-ol
- 6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime
- 5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-indene-1, 6-diol
- 2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol
- 9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol
- 4-hydroxy-3-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-N-((1-methyl-1H-pyrazol-4-yl)methyl)benzamide
- 4-(4-(hydroxymethyl)-1H-pyrazol-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)pyridazin-3-yl)phenol
- 6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 6-(1-(benzyloxy)isoquinolin-7-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one
- 4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
- 5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
- 3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol
- 5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
- 3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
- 5-(5-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol
- 5-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol
- 4-(3-hydroxy-4-(6-methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2-ol

5-(6-methoxypyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-3-(trifluoromethyl)pyridin-2-ol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(3-hydroxy-4-(6-(2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)pyridin-2-ol

5-(6-(dimethylamino)pyridin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

4-(3-hydroxy-4-(6-(2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(pyrimidin-5-yl)phenol

5-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-3-ol

1-cyclopropyl-4-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-2(1H)-one

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)phenol

5-(cyclopent-1-en-1-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(3,6-dihydro-2H-pyran-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,5-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,2-a]pyridin-7-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methylpyridin-4-yl)phenol

5-(1H-imidazol-2-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

5-(imidazo[1,2-a]pyrazin-3-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(5,6,7,8-tetrahydroimidazo[1,2-a]pyrazin-3-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-methyl-1H-imidazol-2-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-4-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-imidazol-5-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(4-nitro-1H-imidazol-2-yl)phenol

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(2-methyl-1H-imidazol-4-yl)phenol

5-(1,2-dimethyl-1H-imidazol-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol

EP 3 848 035 A1

	1-(3-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1H-pyrazole-4-carboxamide
5	2-(6-((3aR,6aS)-5-(2-hydroxyethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	2-(6-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
10	2-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
	4-(3-hydroxy-4-(6-(5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
15	4-(3-hydroxy-4-(6-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one
	2-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol
20	4-(4-(6-(2,7-diazaspiro[4.5]decan-2-yl)pyridazin-3-yl)-3-hydroxyphenyl)-1-methylpyridin-2(1H)-one
	2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)-amino)-pyridazin-3-yl)phenol
	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
25	6-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	6-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)quinolin-7-ol
30	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	7-(6-(methyl(1,2,2,6,6-pentamethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
35	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-6-ol
	7-(6-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)pyridazin-3-yl)isoquinolin-6-ol
40	1-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1,6-diol
	6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile
45	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol
	8-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
50	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
55	3-bromo-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3-carbonitrile

	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(1-methyl-1H-imidazol-4-yl)quinolin-7-ol
	3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
5	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol
	3-ethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
10	7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one
	7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one
15	4-methoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(pyrrolidin-1-yl)quinolin-7-ol
	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-morpholinoquinolin-7-ol
20	4-(dimethylamino)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	4-ethoxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
25	2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1-methyl-1H-pyrazol-4-yl)quinolin-7-ol
	4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
30	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
	6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol
	3-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
35	3-bromo-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
40	5-bromo-3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	6-hydroxy-1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-4(1H)-one
	2,3-dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
45	2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
50	4-methoxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	4-(azetidin-1-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
55	7-hydroxy-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolone-4-carbonitrile
	4-cyclopropyl-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

4-(3,6-dihydro-2H-pyran-4-yl)-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(oxetan-3-yl)quinolin-7-ol

4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol

7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-4(1H)-one

6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol

7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-3,4-dihydroquinolin-2(1H)-one

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazolin-7-ol

7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile

7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carbonitrile

6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carbonitrile

6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carboxamide

7-hydroxy-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2-carboxamide

6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxamide

methyl 6-hydroxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-2-carboxylate

6-hydroxy-7-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile

7-hydroxy-6-(6-(piperazin-1-yl)pyridazin-3-yl)quinoline-2-carbonitrile

7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol

7-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)quinolin-6-ol

1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-7-ol

1-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

1,3 -dimethyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

7-hydroxy-3-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinoline-1-carbonitrile

1-amino-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol

7-hydroxy-1,3-dimethyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinazoline-2,4(1H,3H)-dione

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)benzo[d]oxazol-2(3H)-one

2-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2H-indazol-6-ol

EP 3 848 035 A1

	1-methyl-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-1H-indazol-6-ol
	6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1 (2H)-one
5	2-ethyl-6-hydroxy-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinolin-1(2H)-one
	1-ethoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
10	7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)isoquinoline-1,6-diol
	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-pyridazin-3-yl)-3-phenylisoquinolin-6-ol
	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
15	3-cyclopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
	3-isopropyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-6-ol
20	3-propyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
	3-isopropyl-7-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-pyridazin-3-yl)isoquinolin-6-ol
	3-methyl-7-(6-(piperazin-1-yl)pyridazin-3-yl)isoquinolin-6-ol
25	6-(3-(benzyloxy)isoquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	3-chloro-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
30	3-isopropyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
	3-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol
	4-chloro-2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol
35	4-chloro-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
	7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol
40	5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
45	5-(2-methoxyquinolin-3-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	5-(3-methoxy-naphthalen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	5-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-N-(1,2,2,6,6-pentamethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
50	5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
55	4-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

- 5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
- 5-(3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
- 5 N-methyl-5-(2-methyl-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
- 10 1-methyl-4-(4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3-(trifluoromethoxy)phenyl)pyridin-2(1H)-one
- 5-(4-(3,5-dimethyl-1H-pyrazol-4-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
- 15 5-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
- 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
- 20 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
- 5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
- 25 4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one
- 5-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2-ol
- 30 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
- 3-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)naphthalene-2,7-diol
- 35 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol
- 3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2-ol
- 2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-4-(1H-pyrazol-1-yl)phenol
- 40 5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
- 3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol
- 45 5-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
- 3-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-methyloxazol-2-yl)phenol
- 50 2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-(1,2,3,6-tetrahydropyridin-4-yl)-1,3,4-thiadiazole
- 2-(5-(piperazin-1-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol
- 55 5-(7-methoxyquinolin-6-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazole-2-amine
- 6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-7-ol

3-methoxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile

3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzonitrile

5 methyl-3-fluoro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzoate

5-(2-methoxy-4-(3-(methylamino)-1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

10 7-methoxy-6-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinoline-2-carbonitrile

4-(3-methoxy-4-(5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

15 4-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)-1-methylpyridin-2(1H)-one

5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

20 5-(2-chloro-4-(4,5,6,7-tetrahydropyrazolo[1,5-a]pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

25 N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)-1,3,4-thiadiazole

30 5-(2-chloro-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(6-aminopyridin-3-yl)-2-fluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

35 5-(2-fluoro-4-(3-methyl-1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

40 5-(2,3-difluoro-4-(1-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,3-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

45 5-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

50 5-(2,5-difluoro-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

55 2-(2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2-chloro-5-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-

2-amine

5-(3-fluoro-5-(1H-pyrazol-4-yl)pyridin-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(5-(2-aminopyrimidin-4-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2,5-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2,3-difluorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

4-(3-hydroxy-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(trifluoromethoxy)phenyl)-1-methylpyridin-2(1H)-one

5-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-fluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2,3-difluoro-6-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

6-methoxy-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-3,4-dihydroisoquinolin-1(2H)-one

5-(2-chloro-4-(1H-pyrazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(1H-1,2,3-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(2H-1,2,3-triazol-2-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-chloro-4-(1H-1,2,4-triazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(3-amino-1H-pyrazol-1-yl)-2-chlorophenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

5-(2-chloro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1H-imidazol-1-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(1H-pyrazol-5-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(4-(2,4-dimethylthiazol-5-yl)-2-methoxyphenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(pyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(2-methoxypyridin-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

5-(2-methoxy-4-(6-methoxypyridin-3-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine

2-(2-chloro-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(H)-yl)-1,3,4-thiadiazole

2-(2-chloro-4-(H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aR)-1-methylhexahydropyrrolo[3,4-b]pyrrol-5(1H)-yl)-1,3,4-thiadiazole

1-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)morpholin-2-yl)-N,N-dimethylmethanamine

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2-methyl-2,7-diazaspiro[4.5]decan-7-yl)-1,3,4-thiadiazole

2-(2-fluoro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-5-methylhexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole

2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole

2-(2-methoxy-4-(1-methyl-1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-1-yl)phenol

5-(3-chloro-4-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)phenyl)pyridin-2(1H)-one

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(3-(methylamino)-1H-pyrazol-1-yl)phenol

3-fluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

3,4-difluoro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

6-hydroxy-5-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-2,3-dihydro-1H-inden-1-one

2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

2-(5-(2,6-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol

3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol

EP 3 848 035 A1

	3-chloro-2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol
	2-(2-methoxy-4-(1H-pyrazol-1-yl)phenyl)-5-((2,2,6,6-tetramethylpiperidin-4-yl)methyl)-1,3,4-thiadiazole
5	2-(2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazole
	2-(5-(2,7-diazaspiro[3.5]nonan-2-yl)-1,3,4-thiadiazol-2-yl)-3-fluoro-5-(1H-pyrazol-4-yl)phenol
10	4-methoxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
	4-hydroxy-1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
15	3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
	1-methyl-3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)quinolin-2(1H)-one
20	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole
	2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole
	(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol
25	2-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)benzo[b]thiophene-5-carbonitrile
	5-(3-chlorobenzo[b]thiophen-2-yl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
30	5-(2-methoxy-4-(1H-pyrazol-4-yl)phenyl)-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine
	2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
35	2-{6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
	5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
	5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
40	2-{6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol
45	5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
	5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
	2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol
50	6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
	5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol
55	2-{6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol
	5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

- 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol
- 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 5 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 10 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 15 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 20 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 25 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 30 3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol
- 3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol
- 35 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole
- 40 3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 45 6-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]imidazo[1,2-a]pyridine
- 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 50 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine
- 55 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol
- 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine
- 3-methoxy-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol

3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol

6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine

3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol

6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol

N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine

N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine

3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and

6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

7. A compound selected from the group consisting of:

2-{6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol

2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5-(1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

5-(1-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}phenol

5-(1H-imidazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

5-(5-methyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

2-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-5-(4-nitro-1H-pyrazol-1-yl)phenol

6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

5-(4-amino-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5-(4-nitro-1H-pyrazol-1-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol

5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol

2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

5 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

10 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

15 2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

20 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

25 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

30 3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol

2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol

35 N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole

3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

40 N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

6-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]imidazo[1,2-a]pyridine

45 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

50 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol

55 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

3-methoxy-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol

3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol

6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine

2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine

3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine

2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol

3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol

6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol

N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine

N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine

3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine and

6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;

wherein a form of the compound is selected from the group consisting of a prodrug, salt, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

8. A method of use of a compound salt or a form thereof for treating or ameliorating HD in a subject in need thereof, comprising administering to the subject an effective amount of a compound salt selected from the group consisting of:

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(trifluoromethyl)phenol hydrochloride

2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(1H-pyrazol-1-yl)phenol hydrochloride

2-(6-piperazin-1-yl-pyridazin-3-yl)-5-1H-pyrazol-1-yl-phenol hydrochloride

3-(6-(piperazin-1-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

3-(6-(piperidin-4-yl)pyridazin-3-yl)naphthalene-2,7-diol trifluoroacetate

6-hydroxy-5-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-2,3-dihydro-1H-inden-1-one oxime hydrochloride

2-amino-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-8H-indeno[1,2-d]thiazol-5-ol hydrochloride

9-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5,6-dihydroimidazo[5,1-a]isoquinolin-8-ol hydrochloride

3-fluoro-5-(2-methoxypyridin-4-yl)-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)pyridin-

2(1H)-one hydrochloride

4-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride

5-(3-fluoro-5-hydroxy-4-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenyl)-1-methylpyridin-2(1H)-one hydrochloride

3-fluoro-5-(1H-pyrazol-4-yl)-2-(6-((2,2,6,6-tetramethylpiperidin-4-yl)oxy)pyridazin-3-yl)phenol hydrochloride

5-chloro-3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)phenol hydrochloride

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1H-pyrazol-4-yl)phenol hydrochloride

3-fluoro-2-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-5-(1-methyl-1H-pyrazol-4-yl)phenol hydrochloride

3-(1H-imidazol-1-yl)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-7-ol hydrochloride

6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoline-3,7-diol formate

7-hydroxy-1-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-2(1H)-one hydrochloride

4-methoxy-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinolin-6-ol formate

7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)quinoxalin-6-ol hydrochloride

2-methyl-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-4-(tetrahydro-2H-pyran-4-yl)quinolin-7-ol formate

4-(dimethylamino)-6-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)-quinolin-7-ol formate

6-hydroxy-2-methyl-7-(6-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)pyridazin-3-yl)isoquinolin-1(2H)-one hydrochloride

3-(5-(methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino)-1,3,4-thiadiazol-2-yl)naphthalen-2-ol hydrobromide

N-methyl-5-(5-(1-methyl-1H-pyrazol-4-yl)pyridin-2-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)-1,3,4-thiadiazol-2-amine hydrochloride

3-fluoro-2-(5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazol-2-yl)-5-(1H-pyrazol-4-yl)phenol dihydrochloride

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-((3aR,6aS)-hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)-1,3,4-thiadiazole hydrochloride

2-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-5-(2,7-diazaspiro[4.5]decan-2-yl)-1,3,4-thiadiazole hydrochloride

(R)-(4-(5-(2-chloro-4-(1H-pyrazol-4-yl)phenyl)-1,3,4-thiadiazol-2-yl)piperazin-2-yl)methanol hydrochloride

2-[6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride

5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol hydrochloride

EP 3 848 035 A1

- 2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
- 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
- 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
- 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
- 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
- 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
- 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
- 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
- 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
- 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
- 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
- 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
- 3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
- N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
- N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine hydrochloride
- 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
- 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-

amine hydrochloride;

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

9. A compound salt selected from the group consisting of:

2-[6-[8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol hydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride
 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride
 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride
 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride
 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride
 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride
 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride
 3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride
 N,2,2,6,6-pentamethyl-N-[5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine hydrochloride
 N,2,2,6,6-pentamethyl-N-[5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine hydrochloride
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride;

wherein a form of the compound salt is selected from the group consisting of a prodrug, hydrate, solvate, clathrate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, polymorph and tautomer form thereof.

10. An assay for detecting HTT (Huntingtin protein) using capture/detection antibody pairs, wherein HTT is total HTT and the capture/detection antibody pairs are selected from MAB2166/MAB2174 and MAB2166/#5656S; and, wherein HTT is mutant HTT and the capture/detection antibody pairs are selected from MAB1574/#5656S, P1874/#5656S and MW1/#5656S.

11. The assay of embodiment 10, wherein HTT is total HTT and the capture/detection antibody pair is MAB2166/#5656S; and, wherein HTT is mutant HTT and the capture/detection antibody pair is MW1/#5656S.

12. The method of any of embodiments 1, 6 or 8, wherein the effective amount of the compound is in a range of

from about 0.001 mg/kg/day to about 500 mg/kg/day.

13. A use of the compound of any of embodiments 1, 6 or 8 for treating or ameliorating HD in a subject in need thereof, comprising administering to the subject an effective amount of the compound.

14. The use of embodiment 13, wherein the effective amount of the compound is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.

15. A use of the compound of any of embodiments 1, 6 or 8 in the manufacture of a medicament for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the medicament to the subject.


16. The use of embodiment 15, wherein the effective amount of the compound in the medicament is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.

17. A use of the compound of any of embodiments 1, 6 or 8 in admixture with one or more pharmaceutically acceptable excipient(s) in a pharmaceutical composition for treating or ameliorating HD in a subject in need thereof, comprising administering an effective amount of the pharmaceutical composition to the subject.

18. The use of embodiment 17, wherein the effective amount of the compound in the pharmaceutical composition is in a range of from about 0.001 mg/kg/day to about 500 mg/kg/day.

Claims

1. A compound of Formula (Ia11):



(Ia11),

or a form thereof, wherein (when present),

X is selected from O, NR_{5a}, or a bond;

A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy, wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a},

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or heteroaryl, wherein heteroaryl is a monocyclic, bicyclic, or polycyclic carbon atom ring structure radical having one or more heteroatoms selected from O, S, and N, optionally substituted with 1 or 2 substituents each selected from R_{3a}; wherein when A is phenyl, R_{1a} is not chloro or pyrazolyl;

R_{3a} is selected from nitro or C₁₋₄alkyl;

R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

2. The compound of claim 1 wherein (when present):

R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy, 1H-pyrazolyl or 1H-imidazol-1-yl,

wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a};

R_{3a} is selected from nitro or methyl; and,

R_{4a} is methyl or ethyl;

R_{5a} is hydrogen or methyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

3. A compound, or form thereof, selected from the group consisting of:

2-{6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol,
 2-[6-((1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol,
 5-(1-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-((1R,5S)-8-azabicyclo[3.2.1]oct-3-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol,
 5-(1H-imidazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol,
 6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 5-(4-amino-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(4-nitro-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol,
 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-(piperidin-4-yloxy)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol,
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol,
 N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole,
 3-(4-chloro-2-methoxyphenyl)-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 6-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]imidazo[1,2-a]pyridine,
 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-methoxy-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]benzene-1,3-diol,
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-(1H-pyrazol-4-yl)-4-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyrimidin-5-amine,
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,

2-[6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol,
 6-(1H-pyrazol-4-yl)-3-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]pyridin-2-ol,
 N,2,2,6,6-pentamethyl-N-[5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine,
 N,2,2,6,6-pentamethyl-N-[5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine,
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine, and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;

wherein the form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

4. The compound of claim 3, wherein the compound is a compound salt selected from the group consisting of:


2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride,
 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride,
 5-(1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol hydrochloride,
 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-yloxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride,
 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol dihydrochloride,
 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride,
 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol trihydrochloride,
 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride,
 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride,
 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride,
 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine tetrahydrochloride,
 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride,
 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride,
 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride,
 2-[6-[(1R,5S)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol hydrochloride,
 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride,
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol trihydrochloride,
 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride,
 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol tetrahydrochloride,
 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride,
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride,
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine trihydrochloride,
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine trihydrochloride,
 3-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-6-(1H-pyrazol-4-yl)pyridin-2-ol hydrochloride,
 N,2,2,6,6-pentamethyl-N-[5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine hydrochloride,
 N,2,2,6,6-pentamethyl-N-[5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl]piperidin-4-amine hydrochloride,
 3-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine hydrochloride, and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine hydrochloride;

wherein a form of the compound salt is selected from the group consisting of a hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

5. The compound of claim 1, wherein A is selected from thiophenyl, indazolyl, pyridinyl, pyrimidinyl and phenoxy.

6. The compound of claim 1, wherein when A is phenyl, R_{1a} is not hydroxyl.

7. A compound of Formula (Ia11):



or a form thereof, wherein (when present),

X is selected from O, NR_{5a}, or a bond;

A is selected from phenyl, thiophenyl, indazolyl, pyridinyl, pyrimidinyl or phenoxy, wherein phenyl and phenoxy are each optionally substituted with 1, 2 or 3 substituents each selected from R_{1a}, wherein thiophenyl, indazolyl, pyridinyl, pyrimidinyl are each optionally substituted with 1 or 2 substituents each selected from R_{1a};

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or heteroaryl, wherein heteroaryl is a monocyclic, bicyclic, or polycyclic carbon atom ring structure radical having one or more heteroatoms selected from O, S, and N, optionally substituted with 1 or 2 substituents each selected from R_{3a}; wherein when A is phenyl, R_{1a} is not hydroxyl;

R_{3a} is selected from nitro or C₁₋₄alkyl;

R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

8. The compound of claim 7, wherein A is selected from thiophenyl, indazolyl, pyridinyl, pyrimidinyl and phenoxy.

9. The compound of claim 7 wherein (when present):

R_{1a} is selected from fluoro, chloro, hydroxyl, methyl, difluoromethyl, amino, methoxy, 1H-pyrazolyl or 1H-imidazol-1-yl,

wherein 1H-pyrazolyl is optionally substituted with 1 or 2 substituents each selected from R_{3a};



R_{3a} is selected from nitro or methyl; and,

R_{4a} is methyl or ethyl;

R_{5a} is hydrogen or methyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

10. A compound selected from a compound Formula (Ia15) and Formula (Ia18):



or a form thereof, wherein (when present),

X is selected from O, NR_{5a}, or a bond;

B is selected from 1H-pyrazolyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, (1*R*,5*S*)-8-azabicyclo[3.2.1]octyl, 8-azabicyclo[3.2.1]oct-2-enyl, 2,6-diazaspiro[3.4]octyl or 2,7-diazaspiro[3.5]nonyl, each optionally substituted with 1 or 2 substituents each selected from R_{4a};

R_{1a} is selected from halogen, hydroxyl, C₁₋₄alkyl, halo-C₁₋₄alkyl, amino, C₁₋₄alkoxy, or heteroaryl, wherein heteroaryl is a monocyclic, bicyclic, or polycyclic carbon atom ring structure radical having one or more heteroatoms selected from O, S, and N, optionally substituted with 1 or 2 substituents each selected from R_{3a}; wherein R_{1a} is not chloro or pyrazole;

R_{3a} is selected from nitro or C₁₋₄alkyl; and,

R_{4a} is C₁₋₄alkyl;

R_{5a} is hydrogen, C₁₋₄alkyl, or hydroxyl-C₁₋₄alkyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

11. The compound of claim 10 wherein (when present):

R_{1a} is selected from fluoro, hydroxyl, methyl, difluoromethyl, amino, methoxy, or 1H-imidazol-1-yl,

R_{3a} is selected from nitro or methyl; and,

R_{4a} is methyl or ethyl;

R_{5a} is hydrogen or methyl;

wherein a form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue, racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.

12. The compound, or form thereof, of claim 3 selected from the group consisting of:

2-[6-[(1*R*,5*S*)-8-azabicyclo[3.2.1]oct-3-yl(methyl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-[(1*R*,5*S*)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(1-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-[(1*R*,5*S*)-8-azabicyclo[3.2.1]oct-3-yl]oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol,
 5-(1H-imidazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 5-(5-methyl-1H-pyrazol-4-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]-5-(4-nitro-1H-pyrazol-1-yl)phenol,
 6-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 5-(4-amino-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 2-[6-(1-methyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 5-(4-nitro-1H-pyrazol-1-yl)-2-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]phenol,
 5-(1H-pyrazol-4-yl)-2-[6-(1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]phenol,
 2-[6-(1-ethyl-1,2,3,6-tetrahydropyridin-4-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-[methyl(piperidin-4-yl)amino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-(piperidin-4-ylamino)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-[6-(8-azabicyclo[3.2.1]oct-2-en-3-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2,3-difluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-[2,5-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-(piperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 2-[6-[(1*R*,5*S*)-8-azabicyclo[3.2.1]oct-3-ylamino]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 6-[2-methoxy-6-(1H-pyrazol-4-yl)pyridin-3-yl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-[4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 3-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[4-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-[6-(2,7-diazaspiro[3.5]non-2-yl)pyridazin-3-yl]-5-(1H-pyrazol-4-yl)phenol,
 3-fluoro-4-[6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl]phenol,
 2-[6-[(2,6-dimethylpiperidin-4-yl)oxy]pyridazin-3-yl]-5-(1H-pyrazol-1-yl)phenol,
 N-methyl-6-(2-methyl-2H-indazol-5-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-methyl-5-[6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl]-2H-indazole,

3-(4-chloro-2-methoxyphenyl)-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 N-methyl-6-(2-methylpyrazolo[1,5-a]pyridin-3-yl)-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 6-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}imidazo[1,2-a]pyridine,
 3-[2-methoxy-4-(1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[5-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[5-(1-methyl-1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 3-[4-(1H-pyrazol-4-yl)thiophen-2-yl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 5-(3,5-dimethyl-1H-pyrazol-4-yl)-2-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol,
 6-[2-fluoro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 3-methoxy-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}phenol,
 3-[2-methoxy-4-(4-nitro-1H-pyrazol-1-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}benzene-1,3-diol,
 6-[2-chloro-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine,
 2-(1H-pyrazol-4-yl)-4-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyrimidin-5-amine,
 3-[2,6-difluoro-4-(1H-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine,
 2-{6-(2,6-diazaspiro[3.4]oct-2-yl)pyridazin-3-yl}-5-(1H-pyrazol-4-yl)phenol,
 3-{6-[methyl(2,2,6,6-tetramethylpiperidin-4-yl)amino]pyridazin-3-yl}-6-(1H-pyrazol-4-yl)pyridin-2-ol,
 6-(1H-pyrazol-4-yl)-3-{6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazin-3-yl}pyridin-2-ol,
 N,2,2,6,6-pentamethyl-N-{5-[3-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine,
 N,2,2,6,6-pentamethyl-N-{5-[4-(1H-pyrazol-4-yl)phenoxy]-1,3,4-thiadiazol-2-yl}piperidin-4-amine,
 3-[2-(difluoromethyl)-4-(1-pyrazol-4-yl)phenyl]-6-[(2,2,6,6-tetramethylpiperidin-4-yl)oxy]pyridazine, and
 6-[2-(difluoromethyl)-4-(1H-pyrazol-4-yl)phenyl]-N-methyl-N-(2,2,6,6-tetramethylpiperidin-4-yl)pyridazin-3-amine;
 wherein the form of the compound is selected from the group consisting of a salt, hydrate, solvate, isotopologue,
 racemate, enantiomer, diastereomer, stereoisomer, and tautomer form thereof.



EUROPEAN SEARCH REPORT

 Application Number
 EP 21 15 8464

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	US 2014/051672 A1 (CHEUNG ATWOOD KIM [US] ET AL) 20 February 2014 (2014-02-20) * [0024]-[0026]; [0039]; [0250]-[0251]; [0323]-[0328]; examples, e.g. [0422], [0445], [0448]-[0480]; [0794]-[0800]; [0846], "Biological Example 1"; [0847], "Activity Table"; claims, e.g. Claims 1, 7, 8-13, 14, 17, 18 *	1-12	INV. A61K31/506 C07D405/12 A61P25/00 C07D201/00 A61K31/00 A61K33/00 C07D401/14 C07D413/14 C07D237/20 C07D417/04 C07D417/14 C07D471/10 C07D487/10 C07D403/12
Y	WO 2007/130383 A2 (NEUROMEDIX INC [CA]; WATTERSON MARTIN [US]; VAN ELDIK LINDA [US]; HU W) 15 November 2007 (2007-11-15) * page 1, lines 7-15; page 8, line 26 - page 9, line 3; page 64, lines 2-3; page 65, lines 13-16; page 83, lines 8-15; pages 116-125, see compounds 108, 183, 184, 225; page 154, compound 178; claims *	1-12	
Y	WO 2015/017589 A1 (NOVARTIS AG [CH]; CHEUNG ATWOOD [US]; DALES NATALIE [US]; HURLEY TIMOT) 5 February 2015 (2015-02-05) * Examples, e.g. Example 1-1; pages 130-131, Biological Examples 1 and 2; pages 131-139, Activity Tables; claims *	1-12	TECHNICAL FIELDS SEARCHED (IPC) C07D A61K
Y	WO 2007/003604 A2 (NOVO NORDISK AS [DK]; HOHLWEG ROLF [DK]; ANDERSEN KNUD ERIK [DK]; SOER) 11 January 2007 (2007-01-11) * page 17, line 17 - page 20, line 9; page 20, line 22- page 21, line 2; page 40, line 18 - page 43, line 34; page 52, lines 15-19examples, e.g. Examples 15, 31, 197; claims *	1-12	
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 10 May 2021	Examiner Sen, Alina
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



EUROPEAN SEARCH REPORT

 Application Number
EP 21 15 8464

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
Y	US 2010/267721 A1 (HOHLWEG ROLF [DK] ET AL) 21 October 2010 (2010-10-21) * [0175]-[0176]; [0178], "narcolepsy, attention deficit disorders or reduced wakefulness"; [0186], "coronary heart disease"; [0197]; examples, e.g. Examples 13, 57, 59; claims *	1-12	
Y	WO 2007/133561 A2 (NEUROGEN CORP [US]; XU YUELIAN [US]; CALDWELL TIMOTHY M [US]; XIE LING) 22 November 2007 (2007-11-22) * page 1, lines 13-16; page 1, line 36 - page 2, line 8; page 41, line 6 - page 42, line 4; examples, e.g. at page 58, line 1 - page 59, line 1; starting from page 68, Tables, compounds 52-55, 150-153, 242-327, 387; claims, e.g. Claims 30-32 *	1-12	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (IPC)
Place of search Munich		Date of completion of the search 10 May 2021	Examiner Sen, Alina
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

1 EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 21 15 8464

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2021

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2014051672 A1	20-02-2014	AR 092108 A1	25-03-2015
		AU 2013302859 A1	19-02-2015
		BR 112015003004 A2	04-07-2017
		CA 2880273 A1	20-02-2014
		CL 2015000331 A1	28-08-2015
		CN 104583196 A	29-04-2015
		CR 20150078 A	29-05-2015
		CU 20150014 A7	30-07-2015
		EA 201590371 A1	30-06-2015
		EC SP15009862 A	29-03-2019
		EP 2885288 A1	24-06-2015
		EP 3564233 A1	06-11-2019
		GT 201500030 A	28-09-2017
		HK 1207862 A1	12-02-2016
		IL 237067 A	31-03-2019
		JO 3530 B1	05-07-2020
		JP 6190883 B2	30-08-2017
		JP 2015524842 A	27-08-2015
		KR 20150041655 A	16-04-2015
		KR 20200126016 A	05-11-2020
		MA 37834 A1	28-09-2018
		NZ 704738 A	26-02-2016
		PE 20151890 A1	13-01-2016
		PE 20201165 A1	28-10-2020
		PH 12015500236 A1	30-03-2015
		SG 10201701123Q A	27-04-2017
		SG 11201500507U A	30-03-2015
		TN 2015000038 A1	29-06-2016
		TW 201412721 A	01-04-2014
		UA 114726 C2	25-07-2017
		US 2014051672 A1	20-02-2014
		US 2014213570 A1	31-07-2014
		US 2017290828 A1	12-10-2017
		US 2019358226 A1	28-11-2019
		US 2020345732 A1	05-11-2020
		UY 34974 A	31-03-2014
		WO 2014028459 A1	20-02-2014
		ZA 201500531 B	28-11-2018
WO 2007130383 A2	15-11-2007	NONE	
WO 2015017589 A1	05-02-2015	AU 2014296255 A1	18-02-2016
		CA 2918805 A1	05-02-2015
		CN 105636953 A	01-06-2016
		EA 201690287 A1	29-07-2016
		EP 3027600 A1	08-06-2016

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 21 15 8464

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2021

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		JP 6461150 B2	30-01-2019
		JP 6856614 B2	07-04-2021
		JP 2016527272 A	08-09-2016
		JP 2019059783 A	18-04-2019
		KR 20160037199 A	05-04-2016
		US 2016184305 A1	30-06-2016
		US 2019160062 A1	30-05-2019
		WO 2015017589 A1	05-02-2015

WO 2007003604 A2	11-01-2007	AT 536344 T	15-12-2011
		AU 2006264966 A1	11-01-2007
		BR P10613564 A2	18-01-2011
		CA 2614116 A1	11-01-2007
		CN 103110635 A	22-05-2013
		EP 1902028 A2	26-03-2008
		EP 2233470 A1	29-09-2010
		EP 2386554 A1	16-11-2011
		ES 2375929 T3	07-03-2012
		JP 5121707 B2	16-01-2013
		JP 2009500372 A	08-01-2009
		KR 20080032069 A	14-04-2008
		RU 2011142654 A	27-04-2013
		US 2009312309 A1	17-12-2009
		US 2012232078 A1	13-09-2012
		WO 2007003604 A2	11-01-2007

US 2010267721 A1	21-10-2010	AR 066946 A1	23-09-2009
		CA 2690486 A1	18-12-2008
		CL 2008001705 A1	06-03-2009
		CN 101795567 A	04-08-2010
		EP 2014656 A2	14-01-2009
		EP 2166850 A1	31-03-2010
		JP 2010529200 A	26-08-2010
		TW 200918054 A	01-05-2009
		US 2010267721 A1	21-10-2010
		WO 2008154126 A1	18-12-2008

WO 2007133561 A2	22-11-2007	AU 2007249925 A1	22-11-2007
		CA 2651654 A1	22-11-2007
		EP 2021004 A2	11-02-2009
		JP 2009536651 A	15-10-2009
		KR 20090015956 A	12-02-2009
		US 2008247964 A1	09-10-2008
		WO 2007133561 A2	22-11-2007

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 2014028459 A1 [0201]
- WO 2014116845 A1 [0201]
- WO 2015017589 A1 [0201]

Non-patent literature cited in the description

- **T.W. GREENE et al.** Protective Groups in organic Synthesis. Wiley, 1991 [0120]
- Pro-drugs as Novel Delivery Systems. **T. HIGUCHI ; W. STELLA.** the A.C.S. Symposium Series, and in Bioreversible Carriers in Drug Design. American Pharmaceutical Association and Pergamon Press, 1987, vol. 14 [0121]
- Handbook of Pharmaceutical Salts. Properties, Selection and Use. Wiley-VCH, 2002 [0129]
- **S. BERGE et al.** *Journal of Pharmaceutical Sciences*, 1977, vol. 66 (1), 1-19 [0129]
- **P. GOULD.** *International J. of Pharmaceutics*, 1986, vol. 33, 201-217 [0129]
- **ANDERSON et al.** The Practice of Medicinal Chemistry. Academic Press, 1996 [0129]
- The Orange Book. Food & Drug Administration [0129]

摘要

本说明书涉及化合物、其形式及其药物组合物以及使用这些化合物、其形式或其组合物用于治疗或改善亨廷顿病的方法。

特别地，本说明书涉及式(I)的经取代的单环杂芳基化合物、其形式及其药物组合物以及使用这些化合物、其形式或其组合物用于治疗或改善亨廷顿病的方法。

