



(86) **Date de dépôt PCT/PCT Filing Date:** 2013/03/15
(87) **Date publication PCT/PCT Publication Date:** 2013/10/03
(85) **Entrée phase nationale/National Entry:** 2014/09/25
(86) **N° demande PCT/PCT Application No.:** US 2013/032180
(87) **N° publication PCT/PCT Publication No.:** 2013/148325
(30) **Priorités/Priorities:** 2012/03/26 (US61/615,819);
2012/11/20 (US61/728,685)

(51) **Cl.Int./Int.Cl. C07K 14/435** (2006.01),
A23J 3/04 (2006.01), **A23L 1/305** (2006.01),
C12N 1/21 (2006.01), **C12N 15/63** (2006.01)

(71) **Demandeur/Applicant:**
PRONUTRIA, INC., US

(72) **Inventeurs/Inventors:**
SILVER, NATHANIEL W., US;
BERRY, DAVID ARTHUR, US;
CHILLAKURU, RAJEEV, US;
VON MALTZAHN, GEOFFREY, US;
HAMILL, MICHAEL, J., US

(74) **Agent:** GOWLING LAFLEUR HENDERSON LLP

(54) **Titre : FRAGMENTS NUTRITIFS, PROTEINES NUTRITIVES ET PROCEDES**

(54) **Title: NUTRITIVE FRAGMENTS, PROTEINS AND METHODS**

(57) **Abrégé/Abstract:**

Nutritive proteins are provided. In some embodiments the nutritive proteins comprise a protein digestibility corrected amino acid score (PDCAAS) that exceeds a benchmark protein. Also provided are nucleic acids encoding the proteins, recombinant microorganisms that make the proteins, methods of making the proteins using recombinant microorganisms, compositions that comprise the proteins, and methods of using the proteins, among other things.



(51) **International Patent Classification:**
C07K 14/435 (2006.01) *C12N 15/63* (2006.01)
A23J 3/04 (2006.01) *C12N 1/21* (2006.01)
A23L 1/305 (2006.01)

(21) **International Application Number:**
PCT/US2013/032180

(22) **International Filing Date:**
15 March 2013 (15.03.2013)

(25) **Filing Language:** English

(26) **Publication Language:** English

(30) **Priority Data:**
61/615,819 26 March 2012 (26.03.2012) US
61/728,685 20 November 2012 (20.11.2012) US

(71) **Applicant:** PRONUTRIA, INC. [US/US]; 840 Memorial Drive, 3rd Floor, Cambridge, MA 02139 (US).

(72) **Inventors:** SILVER, Nathaniel, W.; 1218 Cambridge Street, #2, Cambridge, MA 02139 (US). BERRY, David, Arthur; 20 Cameron Street, #402, Brookline, MA 02445 (US). CHILLAKURU, Rajeev; 157 Sixth Street, Apt 204, Cambridge, MA 02142 (US). VON MALTZAHN, Geof-frey; 61 Salem St., Apt. 5, Boston, MA 02113 (US). HAMILL, Michael, J.; Michael J. Hamill, Pronutria, Inc., 840 Memorial Drive, 3rd Floor, Cambridge, MA 02139 (US).

(74) **Agents:** KABLER, Kevin, E. et al.; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Moun-tain View, California 94041 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *with amended claims and statement (Art. 19(1))*
- *with sequence listing part of description (Rule 5.2(a))*

(54) **Title:** NUTRITIVE FRAGMENTS, PROTEINS AND METHODS

(57) **Abstract:** Nutritive proteins are provided. In some embodiments the nutritive proteins comprise a protein digestibility corrected amino acid score (PDCAAS) that exceeds a benchmark protein. Also provided are nucleic acids encoding the proteins, recombinant microorganisms that make the proteins, methods of making the proteins using recombinant microorganisms, compositions that com-prise the proteins, and methods of using the proteins, among other things.

WO 2013/148325 A1

NUTRITIVE FRAGMENTS, PROTEINS AND METHODS**REFERENCE TO RELATED APPLICATION:**

[0001] This application claims priority to U.S. Provisional Patent Application No. 61/615,819, filed March 26, 2012, and Provisional Patent Application No. 61/728,685, filed November 20, 2012, both of which are hereby incorporated herein by reference in their entirety.

SEQUENCE LISTING:

[0002] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on March 12, 2013, is named 1005.014-PCT_SL.txt and is 10,802,713 bytes in size.

INTRODUCTION:

[0003] Dietary protein is an essential nutrient for human health and growth. The World Health Organization recommends that dietary protein should contribute approximately 10 to 15% of energy intake when in energy balance and weight stable. Average daily protein intakes in various countries indicate that these recommendations are consistent with the amount of protein being consumed worldwide. Meals with an average of 20 to 30% of energy from protein are representative of high-protein diets when consumed in energy balance.

[0004] The body cannot synthesize certain amino acids that are necessary for health and growth, and instead must obtain them from food. These amino acids, called “essential amino acids”, are Histidine (H), Isoleucine (I), Leucine (L), Lysine (K), Methionine (M), Phenylalanine (F), Threonine (T), Tryptophan (W), and Valine (V). Dietary proteins that provide all the essential amino acids are referred to as “high quality” proteins. Animal foods such as meat, fish, poultry, eggs, and dairy products are generally regarded as high quality protein sources that provide a good balance of essential amino acids. Casein (a protein commonly found in mammalian milk, making up 80% of the proteins in cow milk) and whey (the protein in the liquid that remains after milk has been curdled and strained) are major sources of high quality dietary protein. Foods that do not provide a good balance of essential amino acids are referred to as “low quality” proteins. Most fruits and vegetables are poor sources of protein. Some plants foods including beans, peas, lentils, nuts and grains (such as

wheat) are better sources of protein. Soy, a vegetable protein manufactured from soybeans, is considered by some to be a high quality protein.

[0005] The minimum fractions of each essential amino acid that make up a healthy adult diet provide a metric against which dietary protein sources can be measured. For example, the fractions can be used to measure how well a dietary protein source meets the nutritional protein needs of humans, by targeting nutritionally balanced amino acid percentages. The World Health Organization (WHO) has developed a method for ranking the nutritional quality of a protein that compares the amino acid profile of the specific food protein against the amino acid requirements for human health. This method assigns a nutritional value to the protein called the “Protein Digestibility Corrected Amino Acid Score” (PDCAAS). Recently the WHO has renamed and slightly revised the PDCAAS standard, giving the revised standard the name “Digestible Indispensable Amino Acid Score (DIAAS)”. (See “Dietary Protein Quality Evaluation in Human Nutrition: Report of an FAO Expert Consultation,” FAO Food and Nutrition Paper 92 (2013).) The new standard revises somewhat the digestibility assumptions used to calculate a DIAAS score versus a PDCAAS score. In many instances the two standards are largely interchangeable.

[0006] Studies of the acute effects of consuming high amounts of protein in humans have shown that inclusion and in some cases increasing protein content in the diet can have beneficial effects. For example, studies have shown that ingestion of protein can induce postprandial satiety (including by suppressing hunger), induce thermogenesis and reduce glycemic response in human subjects.

[0007] Studies of high protein diets for weight loss have shown that protein positively affects energy expenditure and lean body mass. Further studies have shown that overeating produces significantly less weight gain in diets containing at least 5% of energy from protein, and that a high-protein diet decreases energy intake.

[0008] Clinical studies provide evidence that protein prevents muscle loss due to aging or bed rest. In particular, studies have shown that protein supplementation increases muscle fractional synthetic rate (FSR) during prolonged bed rest, maintains leg mass and strength during prolonged bed rest increases lean body mass, improves functional measures of gait and balance, and may serve as a viable intervention for individuals at risk of sarcopenia due to immobility or prolonged bed rest.

[0009] Studies on increasing muscle protein anabolism in athletes have shown that protein provided following exercise promotes muscle hypertrophy to a greater extent than that achieved by exercise alone. It has also been shown that protein provided following exercise supports protein synthesis without any increase in protein breakdown, resulting in a net positive protein balance and muscle mass accretion. While muscle protein synthesis appears to respond in a dose-response fashion to essential amino acid supplementation, not all proteins are equal in building muscle. For example, milk proteins appear to be superior to soy in supporting muscle mass accretion with resistance training, while both are superior to carbohydrate alone. The amino acid leucine is an important factor in stimulating muscle protein synthesis.

[0010] Whole proteins commonly found in foods do not necessarily provide an amino acid composition that meets the amino acid requirements of a mammal, such as a human, in an efficient manner. The result is that, in order to attain the minimal requirements of each essential amino acid, a larger amount of total protein must be consumed in the diet than would be required if the quality of the dietary protein were higher. By increasing the quality of the protein in the diet it is possible to reduce the total amount of protein that must be consumed compared to diets that include lower quality proteins.

[0011] In general, proteins that have higher protein quality are considered more beneficial in a mammalian diet than other proteins that do not. Such proteins are useful, for example, as components of a mammalian diet. Under certain circumstances such proteins promote maintenance of muscle mass, a healthy body mass index, and glycemic balance, among other things. Accordingly, there is a need for sources of proteins that have high protein quality.

[0012] Traditionally, desirable mixtures of amino acids, such as mixtures comprising essential amino acids, have been provided by hydrolyzing a protein with relatively high levels of essential amino acids, such as whey protein, and/or by combining free amino acids in a mixture that optionally also includes a hydrolyzed protein such as whey. Mixtures of this type may have a bitter taste and may be deemed unsuitable or undesirable for certain uses. As a result, such mixtures sometimes include flavoring agents to mask the taste of the free amino acids and/or hydrolyzed protein. In some cases compositions in which a proportion of the amino acid content is provided by polypeptides or proteins are found to have a better taste than compositions with a high proportion of total amino acids provided as free amino acids

and/or certain hydrolyzed proteins. The availability of such compositions has been limited, however, because nutritional formulations have traditionally been made from protein isolated from natural food products, such as whey isolated from milk, or soy protein isolated from soy. The amino acid profiles of those proteins do not necessarily meet the amino acid requirements for a mammal. In addition, commodity proteins typically consist of mixtures of proteins and/or protein hydrolysates which can vary in their protein composition, thus leading to unpredictability regarding their nutritional value. Moreover, the limited number of sources of such high quality proteins has meant that only certain combinations of amino acids are available on a large scale for ingestion in protein form.

[0013] The agricultural methods required to supply high quality animal protein sources such as casein and whey, eggs, and meat, as well as plant proteins such as soy, also require significant energy inputs and have potentially deleterious environmental impacts. Accordingly, it would be useful in certain situations to have alternative sources and methods of supplying proteins for mammalian consumption.

[0014] In theory, synthetic polypeptide sequences comprising a desired mixture of amino acids could be designed and produced in a laboratory setting. This approach may raise various concerns, however, and is therefore not always applicable. First, skilled artisans are aware that obtaining high levels of production of such synthetic sequences may be very challenging. Second, even if such a synthetic protein were synthesized, its suitability for use in a nutritive product would be uncertain. For example, such a non-naturally occurring polypeptide could be an allergen or a toxin. Accordingly, in some embodiments this disclosure provides natural protein or polypeptide sequences, or variants thereof.

[0015] In this disclosure the inventors provide nutritive proteins that comprise or consist of naturally-occurring nutritive proteins, fragments of naturally-occurring nutritive proteins, or sequences derived from such proteins and fragments. The nutritive proteins and the fragments of which they are composed are made up of useful combinations of amino acids. The proteins may be produced by methods that do not rely solely on traditional agriculture for production. For example, the inventors have discovered and this disclosure provides naturally occurring nutritive proteins and fragments composed of combinations of amino acids that contain a useful balance of essential amino acids, as well as in some embodiments a useful balance of non-essential amino acids.

[0016] This disclosure also provides nucleic acids encoding the proteins, recombinant microorganisms that make the proteins, methods of making the proteins using recombinant microorganisms, compositions that comprise the proteins, and methods of using the proteins, among other things.

SUMMARY:

[0017] In a first aspect this disclosure provides nutritive proteins comprising a first polypeptide sequence, wherein at least 50% by weight of the amino acids in the first polypeptide sequence are essential amino acids, and wherein the first polypeptide sequence has a protein digestibility corrected amino acid score (PDCAAS) of at least about 1.4. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 1.8. In some embodiments the first polypeptide sequence comprises a weight ratio of at least 60% essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.2. In some embodiments the first polypeptide sequence comprises a weight ratio of at least 70% essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.5. In some embodiments the first polypeptide sequence comprises a weight ratio of at least 80% essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.9. In some embodiments the first polypeptide sequence comprises a weight ratio of at least 90% essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 3.2. In some embodiments the nutritive protein further comprises at least one of every non-essential amino acid. In some embodiments the nutritive protein further comprises at least one of every non-essential and conditionally-essential amino acid. In some embodiments the nutritive protein further comprises a nutritionally balanced ratio of non-essential amino acids. In some embodiments the first polypeptide sequence comprises a weight ratio of each non-essential amino acid to the total number of amino acids that is equivalent to the weight ratio of each non-essential amino acid to the total number of amino acids in a benchmark protein. In some embodiments the benchmark protein is selected from whey protein, egg protein and soy protein. In some embodiments the first polypeptide sequence comprises 100% essential and conditionally essential amino acids. In some embodiments wherein the first polypeptide sequence has a PDCAAS of from about 1.4 to about 3.6.

[0018] In some embodiments the first polypeptide sequence comprises at least 70% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof. In

some embodiments the first polypeptide sequence comprises at least 95% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof. In some embodiments the first polypeptide sequence comprises at least 70% homology to a naturally occurring protein or fragment thereof. In some embodiments the first polypeptide sequence comprises at least 95% homology to a naturally occurring protein or fragment thereof. In some embodiments the first polypeptide sequence comprises a naturally occurring protein or fragment thereof. In some embodiments the first polypeptide sequence comprises an altered amino acid sequence compared to the amino acid sequence of a naturally occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally occurring reference protein. In some embodiments the first polypeptide sequence comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally occurring reference protein.

[0019] In some embodiments the first polypeptide sequence is not an allergen. In some embodiments the first polypeptide sequence has less than 50% global homology to a known allergen.

[0020] In some embodiments the first polypeptide sequence is not a toxin. In some embodiments the first polypeptide sequence has less than 50% global homology to a known toxin.

[0021] In some embodiments the first polypeptide sequence has a simulated gastric digestion half-life of less than 60 minutes. In some embodiments the first polypeptide sequence has a simulated gastric digestion half-life of less than 30 minutes. In some embodiments the first polypeptide sequence has a simulated gastric digestion half-life of less than 10 minutes. In some embodiments the first polypeptide sequence is completely digested in simulated gastric fluid. In some embodiments the first polypeptide sequence comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site. In some embodiments the first polypeptide sequence comprises no cysteine residues. In some embodiments the first polypeptide sequence comprises no disulfide bonds. In some embodiments the first polypeptide sequence does not comprise N-linked glycosylation. In some embodiments the first polypeptide sequence does not comprise O-linked glycosylation.

[0022] In some embodiments the first polypeptide sequence is resistant to aggregation. In some embodiments the first polypeptide sequence is anionic at pH 7. In some embodiments the first polypeptide sequence has an aqueous solubility at pH 7 of at least 12.5 g/L. In some embodiments the first polypeptide sequence has a calculated solvation score of -20 or less. In some embodiments the first polypeptide sequence has a calculated aggregation score of 0.75 or less. In some embodiments the first polypeptide sequence has a calculated aggregation score of 0.5 or less.

[0023] In some embodiments the first polypeptide sequence comprises at least 20 amino acids. In some embodiments the first polypeptide sequence comprises at least 50 amino acids. In some embodiments the first polypeptide sequence is selected from: A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0024] In some embodiments the first polypeptide sequence consists of at least 20 amino acids. In some embodiments the first polypeptide sequence consists of at least 50 amino acids. In some embodiments the first polypeptide sequence is selected from: A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid

sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0025] In some embodiments the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from: A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0026] In some embodiments the first polypeptide sequence is selected from: A. an amino acid sequence consisting of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; and F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0027] In some embodiments the first polypeptide sequence is selected from: A. an amino acid sequence comprising the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; C. an amino acid sequence comprising a mutein of the amino acid sequence of a

sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; and F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0028] In some embodiments the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0029] In some embodiments the nutritive protein consists of the first polypeptide sequence.

[0030] In some embodiments the nutritive protein further comprises a polypeptide tag for affinity purification. In some embodiments the tag for affinity purification is a polyhistidine-tag.

[0031] In another aspect this disclosure provides nutritive proteins comprising at least 50% by weight essential amino acids, wherein the nutritive protein has a protein digestibility corrected amino acid score (PDCAAS) of at least about 1.4. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 1.8. In some embodiments the nutritive protein comprises a weight ratio of at least 60% essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.2. In some embodiments the nutritive protein comprises a weight ratio of at least 70% essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.5. In some embodiments the nutritive protein comprises a weight ratio of at least 80% essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.9. In some embodiments the nutritive protein comprises a weight ratio of at least 90% essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 3.2. In some embodiments the nutritive protein further comprises at least one of every non-essential amino acid. In some embodiments the nutritive protein further comprises at least one of every non-essential and conditionally-essential

amino acid. In some embodiments the nutritive protein further comprises a nutritionally balanced ratio of non-essential amino acids. In some embodiments the nutritive protein comprises a weight ratio of each non-essential amino acid to the total number of amino acids that is equivalent to the weight ratio of each non-essential amino acid to the total number of amino acids in a benchmark protein. In some embodiments the benchmark protein is selected from whey protein, egg protein and soy protein. In some embodiments the nutritive protein comprises 100% essential and conditionally essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 3.6.

[0032] In some embodiments the nutritive protein comprises at least 70% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof. In some embodiments the nutritive protein comprises at least 95% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof. In some embodiments the nutritive protein comprises at least 70% homology to a naturally occurring protein or fragment thereof. In some embodiments the nutritive protein comprises at least 95% homology to a naturally occurring protein or fragment thereof. In some embodiments the nutritive protein comprises a naturally occurring protein or fragment thereof. In some embodiments the nutritive protein comprises an altered amino acid sequence compared to the amino acid sequence of a naturally-occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally-occurring reference protein. In some embodiments the nutritive protein comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally occurring reference protein.

[0033] In some embodiments the nutritive protein is not an allergen. In some embodiments the nutritive protein has less than 50% global homology to a known allergen.

[0034] In some embodiments the nutritive protein is not a toxin. In some embodiments the nutritive protein has less than 50% global homology to a known toxin.

[0035] In some embodiments the nutritive protein has a simulated gastric digestion half-life of less than 60 minutes. In some embodiments the nutritive protein has a simulated gastric digestion half-life of less than 30 minutes. In some embodiments the nutritive protein has a simulated gastric digestion half-life of less than 10 minutes. In some embodiments the nutritive protein is completely digested in simulated gastric fluid. In some embodiments the nutritive protein comprises at least one protease recognition site selected from a pepsin

recognition site, a trypsin recognition site, and a chymotrypsin recognition site. In some embodiments the nutritive protein comprises no cysteine residues. In some embodiments the nutritive protein comprises no disulfide bonds. In some embodiments the nutritive protein does not comprise N-linked glycosylation. In some embodiments the nutritive protein does not comprise O-linked glycosylation.

[0036] In some embodiments the nutritive protein is resistant to aggregation. In some embodiments the nutritive protein is anionic at pH 7. In some embodiments the nutritive protein has an aqueous solubility at pH 7 of at least 12.5 g/L. In some embodiments the nutritive protein has a calculated solvation score of -20 or less. In some embodiments the nutritive protein has a calculated aggregation score of 0.75 or less. In some embodiments the nutritive protein has a calculated aggregation score of 0.5 or less.

[0037] In some embodiments the nutritive protein comprises at least 20 amino acids. In some embodiments the nutritive protein comprises at least 50 amino acids.

[0038] In some embodiments the nutritive protein comprises at least 20 amino acids. In some embodiments the nutritive protein comprises at least 50 amino acids. In some embodiments the nutritive protein is selected from: A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0039] In some embodiments the nutritive protein consists of at least 20 amino acids. In some embodiments the nutritive protein consists of at least 50 amino acids. In some

embodiments the nutritive protein is selected from: A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0040] In some embodiments the nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from: A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[0041] In some embodiments the nutritive protein is selected from: A. an amino acid sequence consisting of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; and F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0042] In some embodiments the nutritive protein is selected from: A. an amino acid sequence comprising the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; C. an amino acid sequence comprising a mutein of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426; and F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0043] In some embodiments the nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from the amino acid sequence of a sequence selected from SEQ ID NOS: 1-4406 and 4409-4426.

[0044] In some embodiments the nutritive protein further comprises a polypeptide tag for affinity purification. In some embodiments the tag for affinity purification is a polyhistidine-tag.

[0045] In some embodiments the nutritive protein of this disclosure is recombinant. In some embodiments the nutritive protein of this disclosure is isolated. In some embodiments the nutritive protein of this disclosure is purified.

[0046] In another aspect this disclosure provides isolated nucleic acids comprising a nucleic acid sequence that encodes a nutritive protein of this disclosure. In some embodiments the isolated nucleic acid is selected from genomic DNA, cDNA, sense RNA and antisense RNA. In some embodiments the isolated nucleic acid is genomic DNA. In some embodiments the isolated nucleic acid is cDNA. In some embodiments the isolated nucleic acid further comprises an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein.

[0047] In another aspect this disclosure provides a vector comprising a nucleic acid sequence that encodes a nutritive protein of this disclosure. In some embodiments the vector

further comprises an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein

[0048] In another aspect this disclosure provides a recombinant microorganism comprising at least one of a nucleic acid of this disclosure and a vector of this disclosure. In some embodiments the recombinant microorganism is a prokaryote. In some embodiments the prokaryote is heterotrophic. In some embodiments the prokaryote is autotrophic. In some embodiments the prokaryote is a bacteria.

[0049] In another aspect this disclosure provides methods of making a nutritive protein of this disclosure, the method comprising culturing a recombinant microorganism of this disclosure under conditions sufficient for production of the nutritive protein by the recombinant microorganism. In some embodiments the methods further comprise isolating the nutritive protein from the culture.

[0050] In another aspect this disclosure provides nutritive compositions comprising a nutritive protein of this disclosure and at least one second component. In some embodiments the second component is selected from a protein, a polypeptide, a peptide, a free amino acid, a carbohydrate, a fat, a mineral or mineral source, a vitamin, a supplement, an organism, a pharmaceutical, and an excipient. In some embodiments the second component is a protein. In some embodiments the protein is a nutritive protein. In some embodiments the second component is a free amino acid selected from essential amino acids, non-essential amino acids, branch chain amino acids, non-standard amino acids and modified amino acids. In some embodiments the at least one second component is a free amino acid selected from essential amino acids. In some embodiments the second component is a free amino acid selected from branch chain amino acids. In some embodiments the second component is a free amino acid selected from non-essential amino acids. In some embodiments the at least one second component is Leu. In some embodiments the at least one second component is a lipid. In some embodiments the lipid is selected from a fat, oil, triglyceride, cholesterol, phospholipid, and fatty acid. In some embodiments the at least one second component is selected from a mineral and a vitamin. In some embodiments the at least one second component is a supplement. In some embodiments the at least one second component is an organism. In some embodiments the at least one second component is a pharmaceutical. In some embodiments the second component is an excipient. In some embodiments the excipient is selected from a buffering agent, a preservative, a stabilizer, a binder, a

compaction agent, a lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent. In some embodiments the nutritive composition is formulated as a liquid solution, slurry, suspension, gel, paste, powder, or solid.

[0051] In another aspect this disclosure provides methods of making a nutritive composition, comprising providing a nutritive protein of this disclosure and combining the nutritive protein with the at least one second component.

[0052] In another aspect this disclosure provides methods of maintaining or increasing of muscle mass, muscle strength, and functional performance in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0053] In another aspect this disclosure provides methods of maintaining or achieving a desirable body mass index in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0054] In another aspect this disclosure provides methods of treating cachexia in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is of

elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0055] In another aspect this disclosure provides methods of providing protein to a subject with protein-energy malnutrition, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0056] In another aspect this disclosure provides methods of increasing thermogenesis in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0057] In another aspect this disclosure provides methods of inducing at least one of a satiation response and a satiety response in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of

exercise. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[0058] In another aspect this disclosure provides methods of treating at least one of cachexia, sarcopenia and frailty in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

BRIEF DESCRIPTION OF THE DRAWINGS:

[0059] **Figure 1** shows the essential amino acid content of the polypeptide fragment from amino acid 51 to amino acid 75 of the *Schizosaccharomyces pombe* ENTH domain-containing protein C19F8.03c protein (UniProt O60167) (SEQ ID NO: 415), which has a PDCAAS score of 2.5. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0060] **Figure 2** shows the essential amino acid content of the polypeptide fragment from amino acid 170 to amino acid 263 of the *Ovis aries* C-C chemokine receptor 9 protein (UniProt Q1WLP9) (SEQ ID NO: 2016), which has a PDCAAS score of 2.4. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0061] **Figure 3** shows the essential amino acid content of the polypeptide fragment from amino acid 307 to amino acid 369 of the *Gallus gallus* neuronal acetylcholine receptor subunit alpha-6 protein (UniProt P49581) (SEQ ID NO: 1366), which has a PDCAAS score of 2.4. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0062] **Figure 4** shows the essential amino acid content of the polypeptide fragment from amino acid 314 to amino acid 402 of the *Arabidopsis thaliana* putative cadmium/zinc-transporting ATPase HMA4 protein (UniProt O64474) (SEQ ID NO: 421), which has a PDCAAS score of 2.4. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0063] **Figure 5** shows the essential and non-essential amino acid content of the polypeptide fragment from amino acid 63 to amino acid 170 of the *Arabidopsis thaliana* oligoribonuclease protein (UniProt Q9ZVE0) (SEQ ID NO: 4401). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0064] **Figure 6** shows the essential and non-essential amino acid content of the polypeptide fragment from amino acid 7 to amino acid 116 of the *Saccharomyces cerevisiae* probable glutamine amidotransferase protein SNO3 (UniProt P43544) (SEQ ID NO: 1275). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0065] **Figure 7** shows the essential and non-essential amino acid content of the polypeptide fragment from amino acid 277 to amino acid 380 of the *Arabidopsis thaliana* cadmium/zinc-transporting ATPase HMA2 protein (UniProt Q9SZW4) (SEQ ID NO: 4199). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0066] **Figure 8** shows the essential and non-essential amino acid content of the polypeptide fragment from amino acid 234 to amino acid 340 of the *Arabidopsis thaliana* calcium transporting ATPase 3, endoplasmic reticulum-type protein (UniProt Q9SY55) (SEQ ID NO: 4190). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0067] **Figure 9** shows the essential amino acid content of the *Arabidopsis thaliana* reticulon-like protein B13 (UniProt O64837) (SEQ ID NO: 428), which has a PDCAAS score

of 1.9. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0068] **Figure 10** shows the essential amino acid content of the *Arabidopsis thaliana* mannose-1-phosphate guanylyltransferase 1 protein (UniProt O22287) (SEQ ID NO: 257), which has a PDCAAS score of 1.8. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0069] **Figure 11** shows the essential amino acid content of the *Acipenser transmontanus* somatolactin protein (UniProt O93262) (SEQ ID NO: 524), which has a PDCAAS score of 1.8. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0070] **Figure 12** shows the essential amino acid content of the *Vitis vinifera* chalcone synthase protein (UniProt P51090) (SEQ ID NO: 1391), which has a PDCAAS score of 1.7. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each

type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 (labeled “Ideal”) and for whey protein.

[0071] **Figure 13** shows the essential and non-essential amino acid content of the *Lactobacillus acidophilus* orotidine 5'-phosphate decarboxylase protein (UniProt Q5FJB3) (SEQ ID NO: 2530). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0072] **Figure 14** shows the essential and non-essential amino acid content of the *Gallus gallus* trifunctional purine biosynthetic protein adenosine-3 (UniProt P21872) (SEQ ID NO: 935). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0073] **Figure 15** shows the essential and non-essential amino acid content of the *Oryza sativa subsp. japonica* Peroxiredoxin-2C protein (UniProt Q9FR35) (SEQ ID NO: 3830). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars)

and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0074] **Figure 16** shows the essential and non-essential amino acid content of the *Ocimum basillicum* caffeic acid 3-O-methyltransferase 2 protein (UniProt Q9XGV9) (SEQ ID NO: 4313). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[0075] **Figure 17** is a heat map of the PDCAAS scores of fragments of the *Ovis aries* C-C chemokine receptor 9 protein (UniProt Q1WLP9) (SEQ ID NO: 4403). The fragment length is presented along the Y-axis and the fragment start position along the X-axis.

[0076] **Figure 18** is a heat map of the 60/40 EAA/NEAA distance to target distribution of fragments of the *Sus scrofa* serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A alpha isoform protein (UniProt P54612) (SEQ ID NO: 4404). The fragment length is presented along the Y-axis and the fragment start position along the X-axis.

[0077] **Figure 19** shows a two dimensional histogram indicating the relative likelihood (on a log scale) of a protein being expressed in an E. coli expression screen as a function of solvation score (y-axis) and aggregation score (x-axis).

[0078] **Figure 20** shows a two dimensional histogram indicating the relative likelihood (on a log scale) of a protein being solubly expressed in an E. coli expression screen as a function of solvation score (y-axis) and aggregation score (x-axis).

DETAILED DESCRIPTION:

[0079] Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include the plural and plural terms shall include the singular. Generally, nomenclatures used in connection with, and techniques of, biochemistry, enzymology, molecular and cellular biology, microbiology, genetics and protein and nucleic

acid chemistry and hybridization described herein are those well-known and commonly used in the art. Certain references and other documents cited herein are expressly incorporated herein by reference. Additionally, all UniProt/SwissProt records cited herein are hereby incorporated herein by reference. In case of conflict, the present specification, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

[0080] The methods and techniques of the present disclosure are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2001); Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992, and Supplements to 2002); Taylor and Drickamer, *Introduction to Glycobiology*, Oxford Univ. Press (2003); Worthington Enzyme Manual, Worthington Biochemical Corp., Freehold, N.J.; *Handbook of Biochemistry: Section A Proteins, Vol I*, CRC Press (1976); *Handbook of Biochemistry: Section A Proteins, Vol II*, CRC Press (1976); *Essentials of Glycobiology*, Cold Spring Harbor Laboratory Press (1999). Many molecular biology and genetic techniques applicable to cyanobacteria are described in Heidorn et al., "Synthetic Biology in Cyanobacteria: Engineering and Analyzing Novel Functions," *Methods in Enzymology*, Vol. 497, Ch. 24 (2011), which is hereby incorporated herein by reference.

[0081] This disclosure refers to sequence database entries (e.g., UniProt/SwissProt records) for certain protein and gene sequences that are published on the internet, as well as other information on the internet. The skilled artisan understands that information on the internet, including sequence database entries, is updated from time to time and that, for example, the reference number used to refer to a particular sequence can change. Where reference is made to a public database of sequence information or other information on the internet, it is understood that such changes can occur and particular embodiments of information on the internet can come and go. Because the skilled artisan can find equivalent information by searching on the internet, a reference to an internet web page address or a sequence database entry evidences the availability and public dissemination of the information in question.

[0082] Before the present proteins, compositions, methods, and other embodiments are disclosed and described, it is to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise.

[0083] The term “comprising” as used herein is synonymous with “including” or “containing”, and is inclusive or open-ended and does not exclude additional, unrecited members, elements or method steps.

[0084] This disclosure makes reference to amino acids. The full name of the amino acids is used interchangeably with the standard three letter and one letter abbreviations for each. For the avoidance of doubt, those are: Alanine (Ala, A), Arginine (Arg, R), Asparagine (Asn, N), Aspartic acid (Asp, D), Cysteine (Cys, C), Glutamic Acid (Glu, E), Glutamine (Gln, Q), Glycine (Gly, G), Histidine (His, H), Isoleucine (Ile, I), Leucine (Leu, L), Lysine (Lys, K), Methionine (Met, M), Phenylalanine (Phe, F), Proline (Pro, P), Serine (Ser, S), Threonine (Thr, T), Tryptophan (Trp, W), Tyrosine (Tyr, Y), Valine (Val, V).

[0085] As used herein, the term “in vitro” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, etc., rather than within an organism (e.g., animal, plant, or microbe).

[0086] As used herein, the term “in vivo” refers to events that occur within an organism (e.g., animal, plant, or microbe).

[0087] As used herein, the term “isolated” refers to a substance or entity that has been (1) separated from at least some of the components with which it was associated when initially produced (whether in nature or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated agents are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components.

[0088] As used herein, a “branch chain amino acid” is an amino acid selected from Leucine, Isoleucine, and Valine.

[0089] As used herein, an “essential amino acid” (EAA) is an amino acid selected from H, I, L, K, M, F, T, W, and V.

[0090] As used herein, a “non-essential amino acid” (NEAA) is an amino acid selected from A, R, N, D, Q, E, G, P, and S.

[0091] As used herein, a “conditionally essential amino acid” (CEAA) is an amino acid selected from C and Y.

[0092] As used herein, an “aromatic amino acid” (AAA) is an amino acid selected from F and Y.

[0093] As used herein, a “sulphur-containing amino acid” (SAA) is an amino acid selected from M and C.

[0094] The term “peptide” as used herein refers to a short polypeptide, e.g., one that typically contains less than about 50 amino acids and more typically less than about 30 amino acids. The term as used herein encompasses analogs and mimetics that mimic structural and thus biological function.

[0095] The term “polypeptide” encompasses both naturally-occurring and non-naturally occurring proteins, and fragments, mutants, derivatives and analogs thereof. A polypeptide may be monomeric or polymeric. Further, a polypeptide may comprise a number of different domains each of which has one or more distinct activities. For the avoidance of doubt, a “polypeptide” may be any length greater two amino acids.

[0096] The term “isolated protein” or “isolated polypeptide” is a protein or polypeptide that by virtue of its origin or source of derivation (1) is not associated with naturally associated components that accompany it in its native state, (2) exists in a purity not found in nature, where purity can be adjudged with respect to the presence of other cellular material (e.g., is free of other proteins from the same species) (3) is expressed by a cell from a different species, or (4) does not occur in nature (e.g., it is a fragment of a polypeptide found in nature or it includes amino acid analogs or derivatives not found in nature or linkages other than standard peptide bonds). Thus, a polypeptide that is chemically synthesized or synthesized in a cellular system different from the cell from which it naturally originates will be “isolated” from its naturally associated components. A polypeptide or protein may also be rendered substantially free of naturally associated components by isolation, using protein purification techniques well known in the art. As thus defined,

“isolated” does not necessarily require that the protein, polypeptide, peptide or oligopeptide so described has been physically removed from a cell in which it was synthesized.

[0097] The term “polypeptide fragment” as used herein refers to a polypeptide that has a deletion, e.g., an amino-terminal and/or carboxy-terminal deletion compared to a full-length polypeptide, such as a naturally occurring protein. In an embodiment, the polypeptide fragment is a contiguous sequence in which the amino acid sequence of the fragment is identical to the corresponding positions in the naturally-occurring sequence. Fragments typically are at least 5, 6, 7, 8, 9 or 10 amino acids long, or at least 12, 14, 16 or 18 amino acids long, or at least 20 amino acids long, or at least 25, 30, 35, 40 or 45, amino acids, or at least 50 or 60 amino acids long, or at least 70 amino acids long, or at least 100 amino acids long.

[0098] The term “fusion protein” refers to a polypeptide comprising a polypeptide or fragment coupled to heterologous amino acid sequences. Fusion proteins are useful because they can be constructed to contain two or more desired functional elements that can be from two or more different proteins. A fusion protein comprises at least 10 contiguous amino acids from a polypeptide of interest, or at least 20 or 30 amino acids, or at least 40, 50 or 60 amino acids, or at least 75, 100 or 125 amino acids. The heterologous polypeptide included within the fusion protein is usually at least 6 amino acids in length, or at least 8 amino acids in length, or at least 15, 20, or 25 amino acids in length. Fusions that include larger polypeptides, such as an IgG Fc region, and even entire proteins, such as the green fluorescent protein (“GFP”) chromophore-containing proteins, have particular utility. Fusion proteins can be produced recombinantly by constructing a nucleic acid sequence which encodes the polypeptide or a fragment thereof in frame with a nucleic acid sequence encoding a different protein or peptide and then expressing the fusion protein. Alternatively, a fusion protein can be produced chemically by crosslinking the polypeptide or a fragment thereof to another protein.

[0099] As used herein, a protein has “homology” or is “homologous” to a second protein if the nucleic acid sequence that encodes the protein has a similar sequence to the nucleic acid sequence that encodes the second protein. Alternatively, a protein has homology to a second protein if the two proteins have similar amino acid sequences. (Thus, the term “homologous proteins” is defined to mean that the two proteins have similar amino acid sequences.) As used herein, homology between two regions of amino acid sequence

(especially with respect to predicted structural similarities) is interpreted as implying similarity in function.

[00100] When “homologous” is used in reference to proteins or peptides, it is recognized that residue positions that are not identical often differ by conservative amino acid substitutions. A “conservative amino acid substitution” is one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of homology may be adjusted upwards to correct for the conservative nature of the substitution. Means for making this adjustment are well known to those of skill in the art. See, e.g., Pearson, 1994, *Methods Mol. Biol.* 24:307-31 and 25:365-89.

[00101] The following six groups each contain amino acids that are conservative substitutions for one another: 1) Serine, Threonine; 2) Aspartic Acid, Glutamic Acid; 3) Asparagine, Glutamine; 4) Arginine, Lysine; 5) Isoleucine, Leucine, Methionine, Alanine, Valine, and 6) Phenylalanine, Tyrosine, Tryptophan.

[00102] Sequence homology for polypeptides, which is also referred to as percent sequence identity, is typically measured using sequence analysis software. See, e.g., the Sequence Analysis Software Package of the Genetics Computer Group (GCG), University of Wisconsin Biotechnology Center, 910 University Avenue, Madison, Wis. 53705. Protein analysis software matches similar sequences using a measure of homology assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, GCG contains programs such as “Gap” and “Bestfit” which can be used with default parameters to determine sequence homology or sequence identity between closely related polypeptides, such as homologous polypeptides from different species of organisms or between a wild-type protein and a mutein thereof. See, e.g., GCG Version 6.1.

[00103] An exemplary algorithm when comparing a particular polypeptide sequence to a database containing a large number of sequences from different organisms is the computer program BLAST (Altschul et al., *J. Mol. Biol.* 215:403-410 (1990); Gish and States, *Nature Genet.* 3:266-272 (1993); Madden et al., *Meth. Enzymol.* 266:131-141 (1996); Altschul et al.,

Nucleic Acids Res. 25:3389-3402 (1997); Zhang and Madden, Genome Res. 7:649-656 (1997)), especially blastp or tblastn (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)).

[00104] Exemplary parameters for BLASTp are: Expectation value: 10 (default); Filter: seg (default); Cost to open a gap: 11 (default); Cost to extend a gap: 1 (default); Max. alignments: 100 (default); Word size: 11 (default); No. of descriptions: 100 (default); Penalty Matrix: BLOWSUM62. The length of polypeptide sequences compared for homology will generally be at least about 16 amino acid residues, or at least about 20 residues, or at least about 24 residues, or at least about 28 residues, or more than about 35 residues. When searching a database containing sequences from a large number of different organisms, it may be useful to compare amino acid sequences. Database searching using amino acid sequences can be measured by algorithms other than blastp known in the art. For instance, polypeptide sequences can be compared using FASTA, a program in GCG Version 6.1. FASTA provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. Pearson, Methods Enzymol. 183:63-98 (1990). For example, percent sequence identity between amino acid sequences can be determined using FASTA with its default parameters (a word size of 2 and the PAM250 scoring matrix), as provided in GCG Version 6.1, herein incorporated by reference.

[00105] In some embodiments, polymeric molecules (e.g., a polypeptide sequence or nucleic acid sequence) are considered to be “homologous” to one another if their sequences are at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% identical. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 99% similar. The term “homologous” necessarily refers to a comparison between at least two sequences (nucleotides sequences or amino acid sequences). In some embodiments, two nucleotide sequences are considered to be homologous if the polypeptides they encode are at least about 50% identical, at least about 60% identical, at least about 70% identical, at least about 80% identical, or at least about 90% identical for at least one stretch of at least about 20 amino acids. In some embodiments, homologous nucleotide sequences are characterized by the ability to encode a stretch of at least 4-5 uniquely specified amino acids.

Both the identity and the approximate spacing of these amino acids relative to one another must be considered for nucleotide sequences to be considered homologous. In some embodiments of nucleotide sequences less than 60 nucleotides in length, homology is determined by the ability to encode a stretch of at least 4-5 uniquely specified amino acids. In some embodiments, two protein sequences are considered to be homologous if the proteins are at least about 50% identical, at least about 60% identical, at least about 70% identical, at least about 80% identical, or at least about 90% identical for at least one stretch of at least about 20 amino acids.

[00106] As used herein, a “modified derivative” refers to polypeptides or fragments thereof that are substantially homologous in primary structural sequence to a reference polypeptide sequence but which include, e.g., in vivo or in vitro chemical and biochemical modifications or which incorporate amino acids that are not found in the reference polypeptide. Such modifications include, for example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labeling, e.g., with radionuclides, and various enzymatic modifications, as will be readily appreciated by those skilled in the art. A variety of methods for labeling polypeptides and of substituents or labels useful for such purposes are well known in the art, and include radioactive isotopes such as ^{125}I , ^{32}P , ^{35}S , and ^3H , ligands that bind to labeled antiligands (e.g., antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands that can serve as specific binding pair members for a labeled ligand. The choice of label depends on the sensitivity required, ease of conjugation with the primer, stability requirements, and available instrumentation. Methods for labeling polypeptides are well known in the art. See, e.g., Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992, and Supplements to 2002).

[00107] As used herein, “polypeptide mutant” or “mutein” refers to a polypeptide whose sequence contains an insertion, duplication, deletion, rearrangement or substitution of one or more amino acids compared to the amino acid sequence of a reference protein or polypeptide, such as a native or wild-type protein. A mutein may have one or more amino acid point substitutions, in which a single amino acid at a position has been changed to another amino acid, one or more insertions and/or deletions, in which one or more amino acids are inserted or deleted, respectively, in the sequence of the reference protein, and/or truncations of the amino acid sequence at either or both the amino or carboxy termini. A mutein may have the same or a different biological activity compared to the reference protein.

[00108] In some embodiments, a mutein has, for example, at least 85% overall sequence homology to its counterpart reference protein. In some embodiments, a mutein has at least 90% overall sequence homology to the wild-type protein. In other embodiments, a mutein exhibits at least 95% sequence identity, or 98%, or 99%, or 99.5% or 99.9% overall sequence identity.

[00109] As used herein, a “polypeptide tag for affinity purification” is any polypeptide that has a binding partner that can be used to isolate or purify a second protein or polypeptide sequence of interest fused to the first “tag” polypeptide. Several examples are well known in the art and include a His-6 tag, a FLAG epitope, a c-myc epitope, a Strep-TAGII, a biotin tag, a glutathione 5-transferase (GST), a chitin binding protein (CBP), a maltose binding protein (MBP), or a metal affinity tag.

[00110] As used herein, “recombinant” refers to a biomolecule, e.g., a gene or protein, that (1) has been removed from its naturally occurring environment, (2) is not associated with all or a portion of a polynucleotide in which the gene is found in nature, (3) is operatively linked to a polynucleotide which it is not linked to in nature, or (4) does not occur in nature. The term “recombinant” can be used in reference to cloned DNA isolates, chemically synthesized polynucleotide analogs, or polynucleotide analogs that are biologically synthesized by heterologous systems, as well as proteins and/or mRNAs encoded by such nucleic acids. Thus, for example, a protein synthesized by a microorganism is recombinant, for example, if it is synthesized from an mRNA synthesized from a recombinant gene present in the cell.

[00111] The term “polynucleotide”, “nucleic acid molecule”, “nucleic acid”, or “nucleic acid sequence” refers to a polymeric form of nucleotides of at least 10 bases in length. The term includes DNA molecules (e.g., cDNA or genomic or synthetic DNA) and RNA molecules (e.g., mRNA or synthetic RNA), as well as analogs of DNA or RNA containing non-natural nucleotide analogs, non-native internucleoside bonds, or both. The nucleic acid can be in any topological conformation. For instance, the nucleic acid can be single-stranded, double-stranded, triple-stranded, quadruplexed, partially double-stranded, branched, hairpinned, circular, or in a padlocked conformation. The nucleic acid (also referred to as polynucleotides) may include both sense and antisense strands of RNA, cDNA, genomic DNA, and synthetic forms and mixed polymers of the above. They may be modified chemically or biochemically or may contain non-natural or derivatized nucleotide bases, as

will be readily appreciated by those of skill in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), pendent moieties (e.g., polypeptides), intercalators (e.g., acridine, psoralen, etc.), chelators, alkylators, and modified linkages (e.g., alpha anomeric nucleic acids, etc.) Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Such molecules are known in the art and include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule. Other modifications can include, for example, analogs in which the ribose ring contains a bridging moiety or other structure such as the modifications found in "locked" nucleic acids.

[00112] A "synthetic" RNA, DNA or a mixed polymer is one created outside of a cell, for example one synthesized chemically.

[00113] The term "nucleic acid fragment" as used herein refers to a nucleic acid sequence that has a deletion, e.g., a 5'-terminal or 3'-terminal deletion compared to a full-length reference nucleotide sequence. In an embodiment, the nucleic acid fragment is a contiguous sequence in which the nucleotide sequence of the fragment is identical to the corresponding positions in the naturally-occurring sequence. In some embodiments fragments are at least 10, 15, 20, or 25 nucleotides long, or at least 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 nucleotides long. In some embodiments a fragment of a nucleic acid sequence is a fragment of an open reading frame sequence. In some embodiments such a fragment encodes a polypeptide fragment (as defined herein) of the protein encoded by the open reading frame nucleotide sequence.

[00114] As used herein, an endogenous nucleic acid sequence in the genome of an organism (or the encoded protein product of that sequence) is deemed "recombinant" herein if a heterologous sequence is placed adjacent to the endogenous nucleic acid sequence, such that the expression of this endogenous nucleic acid sequence is altered. In this context, a heterologous sequence is a sequence that is not naturally adjacent to the endogenous nucleic acid sequence, whether or not the heterologous sequence is itself endogenous (originating from the same host cell or progeny thereof) or exogenous (originating from a different host

cell or progeny thereof). By way of example, a promoter sequence can be substituted (e.g., by homologous recombination) for the native promoter of a gene in the genome of a host cell, such that this gene has an altered expression pattern. This gene would now become “recombinant” because it is separated from at least some of the sequences that naturally flank it.

[00115] A nucleic acid is also considered “recombinant” if it contains any modifications that do not naturally occur to the corresponding nucleic acid in a genome. For instance, an endogenous coding sequence is considered “recombinant” if it contains an insertion, deletion or a point mutation introduced artificially, e.g., by human intervention. A “recombinant nucleic acid” also includes a nucleic acid integrated into a host cell chromosome at a heterologous site and a nucleic acid construct present as an episome.

[00116] As used herein, the phrase “degenerate variant” of a reference nucleic acid sequence encompasses nucleic acid sequences that can be translated, according to the standard genetic code, to provide an amino acid sequence identical to that translated from the reference nucleic acid sequence. The term “degenerate oligonucleotide” or “degenerate primer” is used to signify an oligonucleotide capable of hybridizing with target nucleic acid sequences that are not necessarily identical in sequence but that are homologous to one another within one or more particular segments.

[00117] The term “percent sequence identity” or “identical” in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over a stretch of at least about nine nucleotides, usually at least about 20 nucleotides, more usually at least about 24 nucleotides, typically at least about 28 nucleotides, more typically at least about 32, and even more typically at least about 36 or more nucleotides. There are a number of different algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using FASTA, Gap or Bestfit, which are programs in Wisconsin Package Version 10.0, Genetics Computer Group (GCG), Madison, Wis. FASTA provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. Pearson, *Methods Enzymol.* 183:63-98 (1990). For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) or using Gap with its default parameters as

provided in GCG Version 6.1, herein incorporated by reference. Alternatively, sequences can be compared using the computer program, BLAST (Altschul et al., J. Mol. Biol. 215:403-410 (1990); Gish and States, Nature Genet. 3:266-272 (1993); Madden et al., Meth. Enzymol. 266:131-141 (1996); Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997); Zhang and Madden, Genome Res. 7:649-656 (1997)), especially blastp or tblastn (Altschul et al., Nucleic Acids Res. 25:3389-3402 (1997)).

[00118] The term “substantial homology” or “substantial similarity,” when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 76%, 80%, 85%, or at least about 90%, or at least about 95%, 96%, 97%, 98% or 99% of the nucleotide bases, as measured by any well-known algorithm of sequence identity, such as FASTA, BLAST or Gap, as discussed above.

[00119] Alternatively, substantial homology or similarity exists when a nucleic acid or fragment thereof hybridizes to another nucleic acid, to a strand of another nucleic acid, or to the complementary strand thereof, under stringent hybridization conditions. “Stringent hybridization conditions” and “stringent wash conditions” in the context of nucleic acid hybridization experiments depend upon a number of different physical parameters. Nucleic acid hybridization will be affected by such conditions as salt concentration, temperature, solvents, the base composition of the hybridizing species, length of the complementary regions, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. One having ordinary skill in the art knows how to vary these parameters to achieve a particular stringency of hybridization.

[00120] In general, “stringent hybridization” is performed at about 25°C below the thermal melting point (T_m) for the specific DNA hybrid under a particular set of conditions. “Stringent washing” is performed at temperatures about 5°C lower than the T_m for the specific DNA hybrid under a particular set of conditions. The T_m is the temperature at which 50% of the target sequence hybridizes to a perfectly matched probe. See Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989), page 9.51. For purposes herein, “stringent conditions” are defined for solution phase hybridization as aqueous hybridization (i.e., free of formamide) in

6xSSC (where 20xSSC contains 3.0 M NaCl and 0.3 M sodium citrate), 1% SDS at 65°C for 8-12 hours, followed by two washes in 0.2xSSC, 0.1% SDS at 65°C for 20 minutes. It will be appreciated by the skilled worker that hybridization at 65°C will occur at different rates depending on a number of factors including the length and percent identity of the sequences which are hybridizing.

[00121] As used herein, an “expression control sequence” refers to polynucleotide sequences which are necessary to affect the expression of coding sequences to which they are operatively linked. Expression control sequences are sequences which control the transcription, post-transcriptional events and translation of nucleic acid sequences. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (e.g., ribosome binding sites); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence. The term “control sequences” is intended to encompass, at a minimum, any component whose presence is essential for expression, and can also encompass an additional component whose presence is advantageous, for example, leader sequences and fusion partner sequences.

[00122] As used herein, “operatively linked” or “operably linked” expression control sequences refers to a linkage in which the expression control sequence is contiguous with the gene of interest to control the gene of interest, as well as expression control sequences that act in trans or at a distance to control the gene of interest.

[00123] As used herein, a “vector” is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid,” which generally refers to a circular double stranded DNA loop into which additional DNA segments may be ligated, but also includes linear double-stranded molecules such as those resulting from amplification by the polymerase chain reaction (PCR) or from treatment of a circular plasmid with a restriction enzyme. Other vectors include cosmids, bacterial artificial chromosomes (BAC) and yeast artificial chromosomes (YAC). Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome (discussed in more detail below). Certain vectors are capable of autonomous

replication in a host cell into which they are introduced (e.g., vectors having an origin of replication which functions in the host cell). Other vectors can be integrated into the genome of a host cell upon introduction into the host cell, and are thereby replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply “expression vectors”).

[00124] The term “recombinant host cell” (or simply “recombinant cell” or “host cell”), as used herein, is intended to refer to a cell into which a recombinant nucleic acid such as a recombinant vector has been introduced. In some instances the word “cell” is replaced by a name specifying a type of cell. For example, a “recombinant microorganism” is a recombinant host cell that is a microorganism host cell. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “recombinant host cell,” “recombinant cell,” and “host cell”, as used herein. A recombinant host cell may be an isolated cell or cell line grown in culture or may be a cell which resides in a living tissue or organism.

[00125] As used herein, the term “heterotrophic” refers to an organism that cannot fix carbon and uses organic carbon for growth.

[00126] As used herein, the term “autotrophic” refers to an organism that produces complex organic compounds (such as carbohydrates, fats, and proteins) from simple inorganic molecules using energy from light (by photosynthesis) or inorganic chemical reactions (chemosynthesis).

[00127] As used herein, “muscle mass” refers to the weight of muscle in a subject’s body. Muscle mass includes the skeletal muscles, smooth muscles (such as cardiac and digestive muscles) and the water contained in these muscles. Muscle mass of specific muscles can be determined using dual energy x-ray absorptiometry (DEXA) (Padden-Jones et al., 2004). Total lean body mass (minus the fat), total body mass, and bone mineral content can be measured by DEXA as well. In some embodiments a change in the muscle mass of a specific muscle of a subject is determined, for example by DEXA, and the change is used as a proxy for the total change in muscle mass of the subject. Thus, for example, if a subject

consumes a nutritive protein as disclosed herein and experiences an increase over a period of time in muscle mass in a particular muscle or muscle group, it can be concluded that the subject has experienced an increase in muscle mass.

[00128] As used herein, a “muscle strength” refers to the amount of force a muscle can produce with a single maximal effort. There are two types of muscle strength, static strength and dynamic strength. Static strength refers to isometric contraction of a muscle, where a muscle generates force while the muscle length remains constant and/or when there is no movement in a joint. Examples include holding or carrying an object, or pushing against a wall. Dynamic strength refers to a muscle generating force that results in movement. Dynamic strength can be isotonic contraction, where the muscle shortens under a constant load or isokinetic contraction, where the muscle contracts and shortens at a constant speed. Dynamic strength can also include isoinertial strength.

[00129] Unless specified, “muscle strength” refers to maximum dynamic muscle strength. Maximum strength is referred to as “one repetition maximum” (1RM). This is a measurement of the greatest load (in kilograms) that can be fully moved (lifted, pushed or pulled) once without failure or injury. This value can be measured directly, but doing so requires that the weight is increased until the subject fails to carry out the activity to completion. Alternatively, 1RM is estimated by counting the maximum number of exercise repetitions a subject can make using a load that is less than the maximum amount the subject can move. Leg extension and leg flexion are often measured in clinical trials (Borsheim et al., “Effect of amino acid supplementation on muscle mass, strength and physical function in elderly,” Clin Nutr 2008;27:189-195; Paddon-Jones, et al., “Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bed rest,” J Clin Endocrinol Metab 2004;89:4351-4358).

[00130] As used herein, “functional performance” refers to a functional test that simulates daily activities. “Functional performance” is measured by any suitable accepted test, including timed-step test (step up and down from a 4 inch bench as fast as possible 5 times), timed floor transfer test (go from a standing position to a supine position on the floor and thereafter up to a standing position again as fast as possible for one repetition), and physical performance battery test (static balance test, chair test, and a walking test) (Borsheim et al., “Effect of amino acid supplementation on muscle mass, strength and physical function in elderly,” Clin Nutr 2008;27:189-195).

[00131] As used herein, a “body mass index” or “BMI” or “Quetelet index” is a subject’s weight in kilograms divided by the square of the subject’s height in meters (kg/m^2).

[00132] For adults, a frequent use of the BMI is to assess how much an individual's body weight departs from what is normal or desirable for a person of his or her height. The weight excess or deficiency may, in part, be accounted for by body fat, although other factors such as muscularity also affect BMI significantly. The World Health Organization regards a BMI of less than 18.5 as underweight and may indicate malnutrition, an eating disorder, or other health problems, while a BMI greater than 25 is considered overweight and above 30 is considered obese. (World Health Organization. BMI classification. Accessed March 19, 2012 http://apps.who.int/bmi/index.jsp?introPage=intro_3.html.) As used herein a “desirable body mass index” is a body mass index of from about 18.5 to about 25. Thus, if a subject has a BMI below about 18.5, then an increase in the subject’s BMI is an increase in the desirability of the subject’s BMI. If instead a subject has a BMI above about 25, then a decrease in the subject’s BMI is an increase in the desirability of the subject’s BMI.

[00133] As used herein, an “elderly” mammal is one who experiences age related changes in at least one of body mass index and muscle mass (e.g., age related sarcopenia). In some embodiments an “elderly” human is at least 50 years old, at least 60 years old, at least 65 years old, at least 70 years old, at least 75 years old, at least 80 years old, at least 85 years old, at least 90 years old, at least 95 years old, or at least 100 years old. In some embodiments and an elderly animal, mammal, or human is a human who has experienced a loss of muscle mass from peak lifetime muscle mass of at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, or at least 60%. Because age related changes to at least one of body mass index and muscle mass are known to correlate with increasing age, in some embodiments an elderly mammal is identified or defined simply on the basis of age. Thus, in some embodiments an “elderly” human is identified or defined simply by the fact that their age is at least 60 years old, at least 65 years old, at least 70 years old, at least 75 years old, at least 80 years old, at least 85 years old, at least 90 years old, at least 95 years old, or at least 100 years old, and without recourse to a measurement of at least one of body mass index and muscle mass.

[00134] As used herein, a patient is “critically-medically ill” if the patient, because of medical illness, experiences changes in at least one of body mass index and muscle mass

(e.g., sarcopenia). In some embodiments the patient is confined to bed for at least 25%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or 100% of their waking time. In some embodiments the patient is unconscious. In some embodiments the patient has been confined to bed as described in this paragraph for at least 1 day, 2 days, 3 days, 4 days, 5 days, 10 days, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 10 weeks or longer.

[00135] As used herein, “protein-energy malnutrition” refers to a form of malnutrition where there is inadequate protein intake. Types include Kwashiorkor (protein malnutrition predominant), Marasmus (deficiency in both calorie and protein nutrition), and Marasmic Kwashiorkor (marked protein deficiency and marked calorie insufficiency signs present, sometimes referred to as the most severe form of malnutrition).

[00136] As used herein, “cachexia” refers to a multifaceted clinical syndrome that results in wasting and weight loss. It is a complex condition where protein catabolism exceeds protein anabolism, which makes muscle wasting a primary feature of the condition. In addition to the metabolic derangements in protein metabolism, it is also characterized by anorexia and inflammation. These derangements plus impaired protein metabolism are responsive to nutrition therapy to varying degrees.

[00137] As used herein, “thermogenesis” is the process of heat production in a mammal. Thermogenesis is accompanied by an increase in energy expenditure. Thermogenesis is specifically the energy burned following the metabolism of a food component (such as protein). This may also be referred to as the thermic effect of food. Total energy expenditure by an individual equals the sum of resting energy expenditure (energy consumed at rest in a fasting state to support basal metabolism), the thermic effect of food, and energy expenditure related to physical activity. Resting energy expenditure accounts for about 65-75% of total energy expenditure in humans. The amount and activity of muscle mass is one influencer of resting energy expenditure. Adequate protein consumption to support muscle also influences resting energy expenditure. The ingestion of protein tends to increase energy expenditure following a meal; this is the thermic effect of food. The thermic effect of food accounts for about 10% of total energy expenditure in humans. While this is a small proportion of total energy expenditure, small increases in this value can impact body weight. Protein has a higher thermic effect than fat or carbohydrate; this effect along with other metabolic influences of protein make it a useful substrate for weight control, diabetes management and other conditions.

[00138] As used herein, “satiation” is the act of becoming full while eating or a reduced desire to eat. This halts or diminishes eating.

[00139] As used herein, “satiety” is the act of remaining full after a meal which manifests as the period of no eating follow the meal.

[00140] As used herein, “exercise” is, most broadly, any bodily activity that enhances or maintains physical fitness and overall health and wellness. Exercise is performed for various reasons including strengthening muscles and the cardiovascular system, honing athletic skills, weight loss or maintenance, as well as for the purpose of enjoyment.

[00141] As used herein, a “sufficient amount” is an amount of a protein or polypeptide disclosed herein that is sufficient to cause a desired effect. For example, if an increase in muscle mass is desired, a sufficient amount is an amount that causes an increase in muscle mass in a subject over a period of time. A sufficient amount of a protein or polypeptide fragment can be provided directly, i.e., by administering the protein or polypeptide fragment to a subject, or it can be provided as part of a composition comprising the protein or polypeptide fragment. Modes of administration are discussed elsewhere herein.

[00142] As used herein, the term “mammal” refers to any member of the taxonomic class mammalia, including placental mammals and marsupial mammals. Thus, “mammal” includes humans, primates, livestock, and laboratory mammals. Exemplary mammals include a rodent, a mouse, a rat, a rabbit, a dog, a cat, a sheep, a horse, a goat, a llama, cattle, a primate, a pig, and any other mammal. In some embodiments, the mammal is at least one of a transgenic mammal, a genetically-engineered mammal, and a cloned mammal.

A. Nutritive Proteins

[00143] For the purposes of this disclosure, a “nutritive protein” is a protein that contains a desirable amount of essential amino acids. In some embodiments, the nutritive protein comprises at least 30% essential amino acids by weight. In some embodiments, the nutritive protein comprises at least 40% essential amino acids by weight. In some embodiments, the nutritive protein comprises at least 50% essential amino acids by weight. In some embodiments the nutritive protein comprises or consists of a protein or fragment of a protein that naturally occurs in an edible species. In its broadest sense, an “edible species” encompasses any species known to be eaten without deleterious effect by at least one type of mammal. A deleterious effect includes a poisonous effect and a toxic effect. In some embodiments an edible species is a species known to be eaten by humans without deleterious

effect. Some edible species are an infrequent but known component of the diet of only a small group of a type of mammal in a limited geographic location while others are a dietary staple throughout much of the world. In other embodiments an edible species is one not known to be previously eaten by any mammal, but that is demonstrated to be edible upon testing. Edible species include but are not limited to *Gossypium turneri*, *Pleurotus cornucopiae*, *Glycine max*, *Oryza sativa*, *Thunnus obesus*, *Abies bracteata*, *Acomys ignitus*, *Lathyrus aphaca*, *Bos gaurus*, *Raphicerus melanotis*, *Phoca groenlandica*, *Acipenser sinensis*, *Viverra tangalunga*, *Pleurotus sajor-caju*, *Fagopyrum tataricum*, *Pinus strobus*, *Ipomoea nil*, *Taxus cuspidata*, *Ipomoea wrightii*, *Mya arenaria*, *Actinidia deliciosa*, *Gazella granti*, *Populus tremula*, *Prunus domestica*, *Larus argentatus*, *Vicia villosa*, *Sargocentron punctatissimum*, *Silene latifolia*, *Lagenodelphis hosei*, *Spisula solidissima*, *Crossarchus obscurus*, *Phaseolus angularis*, *Lathyrus vestitus*, *Oncorhynchus gorboscha*, *Alligator mississippiensis*, *Pinus halepensis*, *Larus canus*, *Brassica napus*, *Silene cucubalus*, *Phoca fasciata*, *Gazella bennettii*, *Pinus taeda*, *Taxus canadensis*, *Zamia furfuracea*, *Pinus yunnanensis*, *Pinus wallichiana*, *Asparagus officinalis*, *Capsicum baccatum*, *Pinus longaeva*, *Taxus baccata*, *Pinus sibirica*, *Citrus sinensis*, *Sargocentron xantherythrum*, *Bison bison*, *Gazella thomsonii*, *Vicia sativa*, *Branta canadensis*, *Apium graveolens*, *Acer campestre*, *Coriandrum sativum*, *Silene conica*, *Lactuca sativa*, *Capsicum chinense*, *Abies veitchii*, *Capra hircus*, *Gazella spekei*, *Oncorhynchus keta*, *Ipomoea obscura*, *Cucumis melo* var. *conomon*, *Phoca hispida*, *Vulpes vulpes*, *Ipomoea quamoclit*, *Solanum habrochaites*, *Populus* sp., *Pinus rigida*, *Quercus lyrata*, *Phaseolus coccineus*, *Larus ridibundus*, *Sargocentron spiniferum*, *Thunnus thynnus*, *Vulpes lagopus*, *Bos gaurus frontalis*, *Acer opalus*, *Acer palmatum*, *Quercus ilex*, *Pinus mugo*, *Grus antigone*, *Pinus uncinata*, *Prunus mume*, *Oncorhynchus tshawytscha*, *Gazella subgutturosa*, *Vulpes zerda*, *Pinus coulteri*, *Gossypium barbadense*, *Acer pseudoplatanus*, *Oncorhynchus nerka*, *Sus barbatus*, *Fagopyrum esculentum* subsp. *Ancestrale*, *Cynara cardunculus*, *Phaseolus aureus*, *Populus nigra*, *Gossypium schwendimanii*, *Solanum chacoense*, *Quercus rubra*, *Cucumis sativus*, *Equus burchelli*, *Oncorhynchus kisutch*, *Pinus radiata*, *Phoca vitulina richardsi*, *Grus nigricollis*, *Abies grandis*, *Oncorhynchus masou*, *Spinacia oleracea*, *Solanum chilense*, *Addax nasomaculatus*, *Ipomoea batatas*, *Equus grevyi*, *Abies sachalinensis*, *Pinus pinea*, *Hipposideros commersoni*, *Crocus nudiflorus*, *Citrus maxima*, *Acipenser transmontanus*, *Gossypium gossypoides*, *Viverra zibetha*, *Quercus cerris*, *Anser indicus*, *Pinus balfouriana*, *Silene otites*, *Oncorhynchus* sp., *Viverra megaspila*, *Bos mutus grunniens*, *Pinus elliottii*,

Equus hemionus kulan, *Capra ibex ibex*, *Allium sativum*, *Raphanus sativus*, *Pinus echinata*,
Prunus serotina, *Sargocentron diadema*, *Silene gallica*, *Brassica oleracea*, *Daucus carota*,
Oncorhynchus mykiss, *Brassica oleracea* var. *alboglabra*, *Gossypium hirsutum*, *Abies alba*,
Citrus reticulata, *Cichorium intybus*, *Bos sauveli*, *Lama glama*, *Zea mays*, *Acorus gramineus*,
Vulpes macrotis, *Ovis ammon darwini*, *Raphicerus sharpei*, *Pinus contorta*, *Bos indicus*,
Capra sibirica, *Pinus ponderosa*, *Prunus dulcis*, *Solanum sogarandinum*, *Ipomoea aquatica*,
Lagenorhynchus albirostris, *Ovis canadensis*, *Prunus avium*, *Gazella dama*, *Thunnus*
alalunga, *Silene pratensis*, *Pinus cembra*, *Crocus sativus*, *Citrullus lanatus*, *Gazella*
rufifrons, *Brassica tournefortii*, *Capra falconeri*, *Bubalus mindorensis*, *Pinus palustris*,
Prunus laurocerasus, *Grus vipio*, *Ipomoea purpurea*, *Pinus leiophylla*, *Lagenorhynchus*
obscurus, *Raphicerus campestris*, *Brassica rapa* subsp. *Pekinensis*, *Acmella radicans*,
Ipomoea triloba, *Pinus patula*, *Cucumis melo*, *Pinus virginiana*, *Solanum lycopersicum*,
Pinus densiflora, *Pinus engelmannii*, *Quercus robur*, *Ipomoea setosa*, *Pleurotus djamor*,
Hipposideros diadema, *Ovis aries*, *Sargocentron microstoma*, *Brassica oleracea* var. *italica*,
Capra cylindricornis, *Populus kitakamiensis*, *Allium textile*, *Vicia faba*, *Fagopyrum*
esculentum, *Bison priscus*, *Quercus suber*, *Lagophylla ramosissima*, *Acrantophis*
madagascariensis, *Acipenser baerii*, *Capsicum annuum*, *Triticum aestivum*, *Xenopus laevis*,
Phoca sibirica, *Acipenser naccarii*, *Actinidia chinensis*, *Ovis dalli*, *Solanum tuberosum*,
Bubalus carabanensis, *Citrus jambhiri*, *Bison bonasus*, *Equus asinus*, *Bubalus*
depressicornis, *Pleurotus eryngii*, *Solanum demissum*, *Ovis vignei*, *Zea mays* subsp.
Parviglumis, *Lathyrus tingitanus*, *Welwitschia mirabilis*, *Grus rubicunda*, *Ipomoea coccinea*,
Allium cepa, *Gazella soemmerringii*, *Brassica rapa*, *Lama vicugna*, *Solanum peruvianum*,
Xenopus borealis, *Capra caucasica*, *Thunnus albacares*, *Equus zebra*, *Gallus gallus*,
Solanum bulbocastanum, *Hipposideros terasensis*, *Lagenorhynchus acutus*, *Hippopotamus*
amphibius, *Pinus koraiensis*, *Acer monspessulanum*, *Populus deltoides*, *Populus trichocarpa*,
Acipenser guldenstadti, *Pinus thunbergii*, *Brassica oleracea* var. *capitata*, *Abyssocottus*
korotneffi, *Gazella cuvieri*, *Abies homolepis*, *Abies holophylla*, *Gazella gazella*, *Pinus*
parviflora, *Brassica oleracea* var. *acephala*, *Cucurbita pepo*, *Pinus armandii*, *Abies mariesii*,
Thunnus thynnus orientalis, *Citrus unshiu*, *Solanum cheesmanii*, *Lagenorhynchus*
obliquidens, *Acer platanoides*, *Citrus limon*, *Acrantophis dumerili*, *Solanum commersonii*,
Gossypium arboreum, *Prunus persica*, *Pleurotus ostreatus*, *Abies firma*, *Gazella leptoceros*,
Salmo salar, *Homarus americanus*, *Abies magnifica*, *Bos javanicus*, *Phoca largha*, *Sus*
cebifrons, *Solanum melongena*, *Phoca vitulina*, *Pinus sylvestris*, *Zamia floridana*, *Vulpes*

corsac, Allium porrum, Phoca caspica, Vulpes chama, Taxus chinensis, Brassica oleracea var. botrytis, Anser anser anser, Phaseolus lunatus, Brassica campestris, Acer saccharum, Pinus pumila, Solanum pennellii, Pinus edulis, Ipomoea cordatotriloba, Populus alba, Oncorhynchus clarki, Quercus petraea, Sus verrucosus, Equus caballus przewalskii, Populus euphratica, Xenopus tropicalis, Taxus brevifolia, Lama guanicoe, Pinus banksiana, Solanum nigrum, Sus celebensis, Brassica juncea, Lagenorhynchus cruciger, Populus tremuloides, Pinus pungens, Bubalus quarlesi, Quercus gamelliflora, Ovis orientalis musimon, Bubalus bubalis, Pinus luchuensis, Sus philippensis, Phaseolus vulgaris, Salmo trutta, Acipenser persicus, Solanum brevidens, Pinus resinosa, Hippotragus niger, Capra nubiana, Asparagus scaber, Ipomoea platensis, Sus scrofa, Capra aegagrus, Lathyrus sativus, Sargocentron tiera, Hippoglossus hippoglossus, Acorus americanus, Equus caballus, Bos taurus, Barbarea vulgaris, Lama guanicoe pacos, Pinus pinaster, Octopus vulgaris, Solanum crispum, Hippotragus equinus, Equus burchellii antiquorum, Crossarchus alexandri, Ipomoea alba, Triticum monococcum, Populus jackii, Lagenorhynchus australis, Gazella dorcas, Quercus coccifera, Anser caerulescens, Acorus calamus, Pinus roxburghii, Pinus tabuliformis, Zamia fischeri, Grus carunculatus, Acomys cahirinus, Cucumis melo var. reticulatus, Gallus lafayettei, Pisum sativum, Pinus attenuata, Pinus clausa, Gazella saudiya, Capra ibex, Ipomoea trifida, Zea luxurians, Pinus krempfii, Acomys wilsoni, Petroselinum crispum, Quercus palustris, Triticum timopheevi, Meleagris gallopavo, Brassica oleracea, Brassica oleracea, Beta vulgaris, Solanum lycopersicum, Phaseolus vulgaris, Xiphias gladius, Morone saxatilis, Micropterus salmoides, Placopecten magellanicus, Sprattus sprattus, Clupea harengus, Engraulis encrasicolus, Cucurbita maxima, Agaricus bisporus, Musa acuminata x balbisiana, Malus domestica, Cicer arietinum, Anas platyrhynchos, Vaccinium macrocarpum, Rubus idaeus x strigosus, Vaccinium angustifolium, Fragaria ananassa, Rubus fruticosus, Cucumis melo, Ananas comosus, Cucurbita pepo, Cucurbita moschata, Sus scrofa domesticus, Ocimum basilicum, Rosmarinus officinalis, Foeniculum vulgare, Rheum rhabarbarum, Carica papaya, Mangifera indica, Actinidia deliciosa, Prunus armeniaca, Prunus avium, Cocos nucifera, Olea europaea, Pyrus communis, Ficus carica, Passiflora edulis, Oryza sativa subsp. Japonica, Oryza sativa subsp. Indica, Coturnix coturnix, Saccharomyces cerevisiae.

[00144] In some embodiments the nutritive protein comprises or consists of a derivative or mutein of a protein or fragment of a protein that naturally occurs in an edible species. Such a nutritive protein may be referred to as an “engineered nutritive protein.” In

such embodiments the natural protein or fragment thereof is a “reference” protein or polypeptide and the engineered nutritive protein or a first polypeptide sequence thereof comprises at least one sequence modification relative to the amino acid sequence of the reference protein or polypeptide. For example, in some embodiments the engineered nutritive protein or first polypeptide sequence thereof is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one reference nutritive protein amino acid sequence. Typically the ratio of at least one of branch chain amino acid residues to total amino acid residues, essential amino acid residues to total amino acid residues, and leucine residues to total amino acid residues, present in the engineered nutritive protein or a first polypeptide sequence thereof is greater than the corresponding ratio of at least one of branch chain amino acid residues to total amino acid residues, essential amino acid residues to total amino acid residues, and leucine residues to total amino acid residues present in the reference nutritive protein or polypeptide sequence.

[00145] In some embodiments the nutritive protein is an abundant protein in food or a derivative or mutein thereof, or is a fragment of an abundant protein in food or a derivative or mutein thereof. An abundant protein is a protein that is present in a higher concentration in a food relative to other proteins present in the food. The food can be a known component of the diet of only a small group of a type of mammal in a limited geographic location, or a dietary staple throughout much of the world. In some embodiments the abundant protein in food is selected from chicken egg proteins such as ovalbumin, ovotransferrin, and ovomucoid; meat proteins such as myosin, actin, tropomyosin, collagen, and troponin; cereal proteins such as casein, alpha1 casein, alpha2 casein, beta casein, kappa casein, beta-lactoglobulin, alpha-lactalbumin, glycinin, beta-conglycinin, glutelin, prolamine, gliadin, glutenin, albumin, globulin; chicken muscle proteins such as albumin, enolase, creatine kinase, phosphoglycerate mutase, triosephosphate isomerase, apolipoprotein, ovotransferrin, phosphoglucomutase, phosphoglycerate kinase, glycerol-3-phosphate dehydrogenase, glyceraldehyde 3-phosphate dehydrogenase, hemoglobin, cofilin, glycogen phosphorylase, fructose-1,6-bisphosphatase, actin, myosin, tropomyosin a-chain, casein kinase, glycogen phosphorylase, fructose-1,6-bisphosphatase, aldolase, tubulin, vimentin, endoplasmic reticulum chaperone, lactate dehydrogenase, desmin, transthyretin, fructose bisphosphate aldolase, carbonic anhydrase, aldehyde dehydrogenase, annexin, adenosyl homocysteinase; pork muscle proteins such as actin, myosin, enolase, titin, cofilin, phosphoglycerate kinase, enolase, pyruvate

dehydrogenase, glycogen phosphorylase, triosephosphate isomerase, myokinase; and fish proteins such as parvalbumin, pyruvate dehydrogenase, desmin, and triosephosphate isomerase.

[00146] Three natural sources of protein generally regarded as good sources of high quality amino acids are whey protein, egg protein, and soy protein. Each source comprises multiple proteins. Table 1A presents the weight proportional representation of each amino acid in the protein source (g AA / g protein) expressed as a percentage.

Table 1A

Amino Acid	Whey	Egg	Soy
Isoleucine	6.5%	5.5%	5.0%
Leucine	11.0%	8.6%	8.0%
Lysine	9.1%	7.2%	6.3%
Methionine	2.1%	3.1%	1.3%
Phenylalanine	3.4%	5.3%	1.2%
Threonine	7.0%	4.8%	3.7%
Tryptophan	1.7%	1.2%	1.3%
Valine	6.2%	6.1%	4.9%
Histidine	2.0%	2.4%	2.7%
<i>Other</i>	51.7%	49.5%	60.4%

[00147] Based on the percentages presented in Table 1, the weight proportion of each protein source that is essential amino acids, branched chain amino acids (L, I, and V), and leucine (L) is presented in Table 1B.

Table 1B

Protein Source	Essential Amino Acids	Branch Chain Amino Acids	Leucine
Whey	49.0%	23.7%	11.0%
Egg	50.5%	20.1%	8.6%
Soy	39.6%	17.9%	8.0%

[00148] The sources relied on to determine the amino acid content of Whey are: Belitz HD., Grosch W., and Schieberle P. Food Chemistry (4th Ed). Springer-Verlag, Berlin Heidelberg 2009; <http://www.gnc.com/product/index.jsp?productId=2986027>; http://www.nutrabio.com/Products/whey_protein_concentrate.htm; and http://www.nutrabio.com/Products/whey_protein_isolate.htm. The amino acid content values from those sources were averaged to give the numbers presented in Tables 1 and 2. The source for soy protein is Egg, National Nutrient Database for Standard Reference, Release 24 (<http://ndb.nal.usda.gov/ndb/foods/list>). The source for soy protein is Self Nutrition Data (<http://nutritiondata.self.com/facts/legumes-and-legume-products/4389/2>).

[00149] As used herein, “whey protein” or “whey” means a protein mixture comprising an amino acid composition according to Tables 1A and 1B. As used herein, whey protein comprises 49% essential amino acids, 24% branch chain amino acids, and 11% leucine, by weight.

[00150] As used herein, “egg protein” or “egg” means a protein mixture comprising an amino acid composition according to Tables 1A and 1B. As used herein, egg protein comprises 51% essential amino acids, 20% branch chain amino acids, and 9% leucine, by weight.

[00151] As used herein, “soy protein” or “soy” means a protein mixture comprising an amino acid composition according to Tables 1A and 1B. As used herein, soy protein comprises 40% essential amino acids, 18% branch chain amino acids, and 8% leucine, by weight.

[00152] While these proteins are relatively high in essential amino acids compared to the other proteins that make up traditional human diets, the proteins do not necessarily provide a proportion of total essential and conditionally essential amino acids for optimal nutrition and/or digestibility. Another feature of these protein sources is that the ratio of each essential amino acid and conditionally essential amino acid in the source to the optimum ratio for each amino acid is generally lower than it could be. In general, a protein source that provides each amino acid in a ratio that is close to the optimum for that amino acid is a more efficient protein source. Put another way, if a first protein source provides each amino acid at a ratio closer to its optimum ratio than a second protein source does, a diet comprising a significant contribution to total protein from the first protein source may provide a sufficient amount of each amino acid at a lower amount of total protein consumed than would be

required in a diet comprising a significant contribution to total protein from the second protein source.

[00153] As an example of the above point, the World Health Organization (WHO) recommends 58g/day of a protein diet with a protein digestibility corrected amino acid score (PDCAAS) of 1 for a person weighing 70kg. This requirement can be further broken down into the minimum amino acid requirements (in grams) shown in Table 2A.

Table 2A

H	I	L	K	SAA	AAA	T	W	V
0.87	1.74	3.42	2.61	1.28	2.20	1.33	0.35	2.26

[00154] By comparison, 58g of a protein with a PDCAAS of 2 translates into the minimum amino acid amounts (in grams) shown in Table 2B

Table 2B

H	I	L	K	SAA	AAA	T	W	V
1.74	3.49	6.86	5.23	2.56	4.42	2.67	0.70	4.53

[00155] As such, 29g of a protein with a PDCAAS of 2 provides the same amount of nutrition obtained in 58g of a protein with a PDCAAS of 1 (i.e. comparable amounts or more of each amino acid).

[00156] As another example of the above point, consider a low PDCAAS diet consisting only of wheat protein, which provides the follow amounts of each amino acid (in mg/g of total protein) and is notably deficient in L (**Table 2C**).

Table 2C

PDCAAS	H	I	L	K	SAA	AAA	T	W	V
0.15	23.0	40.9	85.3	6.9	22.7	107.4	20.3	7.8	42.9

[00157] In order to fortify 1g of this diet and bring the PDCAAS score up to a value of 1, (i.e. a diet containing the WHO recommended amino acid intake values), approximately 0.7g of whey protein, 1.8g of egg protein, or 4g of soy protein would be required. By comparison, only 0.3g of a protein with a nutritionally balanced ratio of essential and

conditionally essential amino acids (PDCAAS 3.61) is required to fortify 1 g of wheat protein. In some instances herein the portion of amino acid(s) of a particular type within a polypeptide, protein or a composition is quantified based on the weight ratio of the type of amino acid(s) to the total weight of amino acids present in the polypeptide, protein or composition in question. This value is calculated by dividing the weight of the particular amino acid(s) in the polypeptide, protein or a composition by the weight of all amino acids present in the polypeptide, protein or a composition.

[00158] The minimum daily requirements for essential amino acids in adult humans (>18 years of age) proposed by the WHO, in milligrams of each amino acid per gram of total protein, are as presented in **Table 3A**:

Table 3A

	H	I	L	K	SAA	AAA	T	W	V
>18	15	30	59	45	22	38	23	6.0	39

[00159] Note that in the WHO analysis Y (a conditionally-essential amino acid) is grouped with F and reported as a total value for aromatic amino acids (AAA), and C is grouped with M and reported as a total value for sulphur containing amino acids (SAA). The WHO adopts this convention because in certain groups under certain conditions C and Y can be essential.

[00160] Converting milligrams to grams yields numbers presented in **Table 3B** that correspond to the minimal fraction of essential amino acids required in a healthy, adult diet (i.e. grams of amino acid / grams of total protein):

Table 3B

	H	I	L	K	SAA	AAA	T	W	V
>18	0.015	0.03	0.059	0.045	0.022	0.038	0.023	0.006	0.039

[00161] Adding up these fractions yields a total of 0.277, meaning that the minimal required amount of EAAs and CEAAAs in the adult diet need only account for 27.7% of the total protein intake. If a protein is intended to provide only the essential/conditionally essential amino acid component of a diet it may comprise a higher proportion of EAA/CEAAAs, up to 100%. The relative percentages of each EAA and CEAA in such a

protein are obtained by multiplying the fractions shown in **Table 3B** by $(100/27.7 \approx 3.61)$.

The fraction of each EAA and CEAA in such a protein is then as presented in **Table 3C**:

Table 3C

	H	I	L	K	SAA	AAA	T	W	V
>18	0.054	0.108	0.213	0.162	0.079	0.137	0.083	0.022	0.141

[00162] The minimum fractions of each EAA that make up a healthy adult diet provide a metric against which dietary protein sources can be measured. For example, the fractions can be used to measure how well a dietary protein source meets the nutritional protein needs of humans, by targeting nutritionally balanced amino acid percentages. Based on the WHO recommendations, the nutritionally balanced amino acid percentages for the essential amino acids, along with the associated protein digestibility corrected amino acid score (PDCAAS), assuming 100% digestion, are shown below in **Table 4**.

Table 4: Nutritionally Ballanced EAA/CEAA Target Percentages for an Adult (>18 yo)

%EAA+%CEAA	PDCAAS	H	I	L	K	SAA	AAA	T	W	V
100	3.6	5.4	10.8	21.3	16.2	7.9	13.7	8.3	2.2	14.1
90	3.2	4.9	9.7	19.2	14.6	7.1	12.3	7.5	1.9	12.7
80	2.9	4.3	8.7	17.0	13.0	6.4	11.0	6.6	1.7	11.3
70	2.5	3.8	7.6	14.9	11.4	5.6	9.6	5.8	1.5	9.9
60	2.2	3.2	6.5	12.8	9.7	4.8	8.2	5.0	1.3	8.4
50	1.8	2.7	5.4	10.6	8.1	4.0	6.9	4.2	1.1	7.0
40	1.4	2.2	4.3	8.5	6.5	3.2	5.5	3.3	0.9	5.6
30	1.1	1.6	3.2	6.4	4.9	2.4	4.1	2.5	0.6	4.2
27.7	1	1.5	3	5.9	4.5	2.2	3.8	2.3	0.6	3.9
20	0.7	1.1	2.2	4.3	3.2	1.6	2.7	1.7	0.4	2.8
10	0.4	0.5	1.1	2.1	1.6	0.8	1.4	0.8	0.2	1.4

SAA – Sulfur containing amino acids (Methionine, Cysteine)

AAA – Aromatic amino acids (Phenylalanine, Tyrosine)

[00163] The PDCAAS score is a WHO suggested scoring metric for assessing nutritional protein values for humans. To compute the PCDAAS for a given protein, one

simply computes the relative fraction of EAAs and CEAAAs present and divides by the numbers shown in **Table 3B**. (Note: For the purpose of this disclosure, the calculation of a PDCAAS score assumes that the dietary protein under evaluation is completely digested and bioavailable). For each EAA and CEAA, this gives the fraction of the requirement provided by the protein. The PCDAAS value of a protein or polypeptide sequence is simply the smallest of these values for the protein or polypeptide sequence. It indicates the worst case fraction of EAA or CEAA requirements provided by the protein/polypeptide. Note that the above analysis shows why the highest possible PCDAAS is 3.61. The only way to increase the PCDAAS further would be to increase the total EAA+CEAA concentration > 100%, which of course is not possible.

[00164] Recently the WHO has renamed and slightly revised the PDCAAS standard, giving the revised standard the name “Digestible Indispensable Amino Acid Score (DIAAS)”. (See “Dietary Protein Quality Evaluation in Human Nutrition: Report of an FAO Expert Consultation,” FAO Food and Nutrition Paper 92 (2013).) The new standard revises somewhat the digestibility assumptions used to calculate a DIAAS score versus a PDCAAS score. Assuming the protein under consideration is 100% digestible and bioavailable, as is done herein, the two standards are largely interchangeable. For the avoidance of doubt, the disclosure uses the terms “PDCAAS” and “PDCAAS score” as defined herein to define properties of proteins and polypeptides. The PDCAAS values assigned to the nutritive proteins and polypeptides of this disclosure assume that the dietary protein under evaluation is completely digestible and bioavailable. As demonstrated in Example 13, for example, the nutritive proteins and polypeptides of this disclosure display useful levels of digestibility and in several embodiments are more easily digested than protein standards such as whey. Example 13 also presents data demonstrating that in some embodiments recombinant nutritive proteins and polypeptides of this disclosure have a higher digestibility than native forms of the proteins or polypeptides such as the forms that are present in whole foods and/or processed foods.

[00165] Digestibility of a protein source is an important prerequisite to the nutritional value of the protein. If a protein is not well digested its bioavailability will be lower and as a result the actual nutritional value of the protein may be lower than its amino acid composition would otherwise suggest. Moreover, a protein that is more rapidly digested than a second protein will deliver nutrition faster than the second protein and with less input energy. Rapid digestibility is also thought to be correlated with reduced allergenicity. (See Goodman R.E.,

Vieths S., Sampson H.A., Hill D., Ebisawa M., Taylor S., van Ree R. Nat. Biotech. (2008) 26: 73:81; Ofori-Anti A.O., Ariyaratna H., Chen L., Lee H.L., Pramod S.N., Goodman R.E. Reg. Toxicol. Pharmacol. (2008) 52: 94-103.) A protein that digests more rapidly than a second protein is around for shorter amount of time and thus less likely to interact with gut localized IgE and elicit an immune response before it is digested and/or absorbed. Whey is a good example of this. It is a known allergen and does not digest quickly, even though it is bioavailable over many hours. In some embodiments the nutritive proteins of this disclosure are more rapidly digested than protein standards such as whey, soy, and/or egg. In some embodiments that feature of the proteins of this disclosure contributes to their usefulness as protein sources for ingestion. In some embodiments the nutritive proteins of this disclosure have a lower allergenicity than protein standards such as whey, soy, and/or egg.

[00166] In general, the nutritive proteins provided by this disclosure comprise at least 50% by weight EAAs. In some embodiments the nutritive proteins comprise at least 55% by weight EAAs, at least 60% by weight EAAs, at least 65% by weight EAAs, at least 70% by weight EAAs, at least by weight 75% EAAs, at least 80% by weight EAAs, at least 85% by weight EAAs, at least 90% by weight EAAs, or at least 95% by weight EAAs, at least 96% by weight EAAs, at least 97% by weight EAAs, at least 98% by weight EAAs, at least 99% by weight EAAs, or 100% EAAs. In some embodiments the nutritive proteins of this disclosure comprise from 50 to 100% by weight EAAs, from 60 to 100% by weight EAAs, from 70 to 100% by weight EAAs, from 80 to 90% by weight EAAs, from 60 to 90% by weight EAAs, from 60 to 80% by weight EAAs, from 70 to 90% by weight EAAs, from 60 to 70% by weight EAAs, from 70 to 80% by weight EAAs, from 80 to 90% by weight EAAs, and from 90 to 100% by weight EAAs. In some embodiments the nutritive proteins comprise from 90% to 100% EAAs and CEAAAs (that is, the combined fraction of essential amino acids and conditionally essential amino acids in the nutritive proteins is from 90% to 100%. In some embodiments the nutritive proteins comprise 100% EAAs and CEAAAs.

[00167] In some embodiments the PDCAAS of the nutritive protein is at least about 1.4, at least about 1.5, at least about 1.6, at least about 1.7, at least about 1.8, at least about 1.9, at least about 2.0, at least about 2.1, at least about 2.2, at least about 2.3, at least about 2.4, at least about 2.5, at least about 2.6, at least about 2.7, at least about 2.8, at least about 2.9, at least about 3.0, at least about 3.1, at least about 3.2, at least about 3.3, at least about 3.4, at least about 3.5, or about 3.6. In some embodiments of the nutritive proteins, the PDCAAS of the nutritive protein is from about 1.4 to about 1.6, from about 1.4 to about 1.8,

from about 1.4 to about 2.0, from about 1.4 to about 2.2, from about 1.4 to about 2.4, from about 1.4 to about 2.6, from about 1.4 to about 2.8, from about 1.4 to about 3.0, from about 1.4 to about 3.2, from about 1.4 to about 3.4, or from about 1.4 to about 3.6.

[00168] In some embodiments the nutritive protein comprises at least 50% by weight EAAs and has a PDCAAS of at least about 1.4. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 1.8.

[00169] In some embodiments of the nutritive protein at least 60% by weight of the amino acids are essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.2.

[00170] In some embodiments of the nutritive protein at least 70% by weight of the amino acids are essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.5.

[00171] In some embodiments of the nutritive protein at least 80% by weight of the amino acids are essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 2.9.

[00172] In some embodiments of the nutritive protein at least 90% by weight of the amino acids are essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 3.2.

[00173] In some embodiments the nutritive protein further comprises at least one of every non-essential amino acid. In some embodiments the nutritive protein further comprises a nutritionally balanced ratio of non-essential amino acids.

[00174] In some embodiments the nutritive protein comprises 100% essential amino acids or 100% essential and conditionally-essential amino acids. In some embodiments the nutritive protein has a PDCAAS of from about 1.4 to about 3.6.

[00175] In some embodiments the nutritive protein comprises or consists of a naturally-occurring protein or fragment thereof.

[00176] In some embodiments the nutritive protein comprises an altered amino acid sequence compared to the amino acid sequence of a naturally-occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally-occurring reference protein. In some embodiments the nutritive protein comprises

at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally-occurring reference protein.

[00177] In some embodiments the nutritive protein is selected from: A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00178] In some embodiments the nutritive protein is selected from: A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00179] In some embodiments the nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from: A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00180] In some embodiments the nutritive protein comprises a first polypeptide sequence that comprises at least 20 amino acids. In some embodiments the first polypeptide sequence comprises at least 30 amino acids, at least 40 amino acids, at least 50 amino acids, at least 60 amino acids, at least 70 amino acids, at least 80 amino acids, at least 90 amino acids, at least 100 amino acids, at least 110 amino acids, at least 120 amino acids, at least 130 amino acids, at least 140 amino acids, at least 150 amino acids, at least 160 amino acids, at least 170 amino acids, at least 180 amino acids, at least 190 amino acids, at least 200 amino acids, at least 300 amino acids, at least 400 amino acids, at least 500 amino acids, at least 600 amino acids, at least 700 amino acids, at least 800 amino acids, at least 900 amino acids, or at least 1,000 amino acids. In some embodiments the first polypeptide sequence comprises from 20 to 50 amino acids, from 20 to 75 amino acids, from 20 to 100 amino acids, from 30 to 100 amino acids, from 40 to 100 amino acids, from 50 to 100 amino acids, from 50 to 200 amino acids, from 100 to 200 amino acids, from 200 to 300 amino acids, from 300 to 400 amino acids, or from 400 to 500 amino acids.

[00181] In some embodiments the first polypeptide sequence comprises at least 50% by weight EAAs. In some embodiments the first polypeptide sequence comprises at least 55% by weight EAAs, at least 60% by weight EAAs, at least 65% by weight EAAs, at least 70% by weight EAAs, at least by weight 75% EAAs, at least 80% by weight EAAs, at least 85% by weight EAAs, at least 90% by weight EAAs, or at least 95% by weight EAAs, at least 96% by weight EAAs, at least 97% by weight EAAs, at least 98% by weight EAAs, at least 99% by weight EAAs, or 100% EAAs. In some embodiments the first polypeptide sequence comprises from 50 to 100% by weight EAAs, from 60 to 100% by weight EAAs, from 70 to 100% by weight EAAs, from 80 to 90% by weight EAAs, from 60 to 90% by weight EAAs, from 60 to 80% by weight EAAs, from 70 to 90% by weight EAAs, from 60 to 70% by weight EAAs, from 70 to 80% by weight EAAs, from 80 to 90% by weight EAAs, and from 90 to 100% by weight EAAs. In some embodiments the first polypeptide sequence

comprises from 90% to 100% EAAs and CEAAAs. In some embodiments the first polypeptide sequence comprises 100% EAAs and CEAAAs.

[00182] In some embodiments the PDCAAS of the first polypeptide sequence is at least about 1.4, at least about 1.5, at least about 1.6, at least about 1.7, at least about 1.8, at least about 1.9, at least about 2.0, at least about 2.1, at least about 2.2, at least about 2.3, at least about 2.4, at least about 2.5, at least about 2.6, at least about 2.7, at least about 2.8, at least about 2.9, at least about 3.0, at least about 3.1, at least about 3.2, at least about 3.3, at least about 3.4, at least about 3.5, or about 3.6. In some embodiments of the nutritive proteins, the PDCAAS of the first polypeptide sequence is from about 1.4 to about 1.6, from about 1.4 to about 1.8, from about 1.4 to about 2.0, from about 1.4 to about 2.2, from about 1.4 to about 2.4, from about 1.4 to about 2.6, from about 1.4 to about 2.8, from about 1.4 to about 3.0, from about 1.4 to about 3.2, from about 1.4 to about 3.4, or from about 1.4 to about 3.6.

[00183] In some embodiments the first polypeptide sequence has a PDCAAS of at least about 1.4. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 1.8.

[00184] In some embodiments of the nutritive protein at least 60% by weight of the amino acids of the first polypeptide sequence are essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.2.

[00185] In some embodiments of the nutritive protein at least 70% by weight of the amino acids of the first polypeptide sequence are essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.5.

[00186] In some embodiments of the nutritive protein at least 80% by weight of the amino acids of the first polypeptide sequence are essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.9.

[00187] In some embodiments of the nutritive protein at least 90% by weight of the amino acids of the first polypeptide sequence are essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 3.2.

[00188] In some embodiments of the nutritive protein, the first polypeptide sequence further comprises at least one of every non-essential amino acid. In some embodiments of

the nutritive protein, the first polypeptide sequence further comprises a nutritionally balanced ratio of non-essential amino acids.

[00189] In some embodiments of the nutritive protein, the first polypeptide sequence comprises 100% essential and conditionally essential amino acids. In some embodiments the first polypeptide sequence has a PDCAAS of from about 1.4 to about 3.6.

[00190] In some embodiments of the nutritive protein, the first polypeptide sequence comprises or consists of a naturally-occurring protein or fragment thereof.

[00191] In some embodiments of the nutritive protein, the first polypeptide sequence comprises an altered amino acid sequence compared to the amino acid sequence of a naturally-occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally-occurring reference protein. In some embodiments the first polypeptide sequence comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally-occurring reference protein.

[00192] In some embodiments of the nutritive protein, the first polypeptide sequence is selected from: A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00193] In some embodiments of the nutritive protein, the first polypeptide sequence is selected from: A. an amino acid sequence consisting of the amino acid sequence of a protein

listed in at least one of **Tables 9-11**; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00194] In some embodiments of the nutritive protein, the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from: A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00195] In some embodiments the nutritive protein consists of the first polypeptide sequence.

[00196] In some embodiments the nutritive protein further comprises a second polypeptide sequence. In some embodiments the second polypeptide sequence comprises of from 3 to 10, 5 to 20, 10 to 30, 20 to 50, 25 to 75, 50 to 100 or 100 to 200 amino acids. In some embodiments the second polypeptide sequence is selected from a tag for affinity purification, a protein domain linker, and a protease recognition site. In some embodiments the tag for affinity purification is a polyhistidine-tag. In some embodiments the protein domain linker comprises at least one copy of the sequence GGSG. In some embodiments the protease is selected from pepsin, trypsin, and chymotrypsin.

[00197] In some embodiments the nutritive protein consists of the first polypeptide sequence and the second polypeptide sequence.

[00198] In some embodiments the nutritive protein further comprises a third polypeptide sequence. In some embodiments third polypeptide sequence comprises at least 20 amino acids, at least 30 amino acids, at least 40 amino acids, at least 50 amino acids, at least 60 amino acids, at least 70 amino acids, at least 80 amino acids, at least 90 amino acids, at least 100 amino acids, at least 110 amino acids, at least 120 amino acids, at least 130 amino acids, at least 140 amino acids, at least 150 amino acids, at least 160 amino acids, at least 170 amino acids, at least 180 amino acids, at least 190 amino acids, at least 200 amino acids, at least 300 amino acids, at least 400 amino acids, at least 500 amino acids, at least 600 amino acids, at least 700 amino acids, at least 800 amino acids, at least 900 amino acids, or at least 1,000 amino acids. In some embodiments the third polypeptide sequence comprises from 20 to 50 amino acids, from 20 to 75 amino acids, from 20 to 100 amino acids, from 30 to 100 amino acids, from 40 to 100 amino acids, from 50 to 100 amino acids, from 50 to 200 amino acids, from 100 to 200 amino acids, from 200 to 300 amino acids, from 300 to 400 amino acids, or from 400 to 500 amino acids

[00199] In some embodiments the third polypeptide sequence comprises at least 50% by weight EAAs. In some embodiments the third polypeptide sequence comprises at least 55% by weight EAAs, at least 60% by weight EAAs, at least 65% by weight EAAs, at least 70% by weight EAAs, at least by weight 75% EAAs, at least 80% by weight EAAs, at least 85% by weight EAAs, at least 90% by weight EAAs, or at least 95% by weight EAAs, at least 96% by weight EAAs, at least 97% by weight EAAs, at least 98% by weight EAAs, at least 99% by weight EAAs, or 100% EAAs. In some embodiments the third polypeptide sequence comprises from 50 to 100% by weight EAAs, from 60 to 100% by weight EAAs, from 70 to 100% by weight EAAs, from 80 to 90% by weight EAAs, from 60 to 90% by weight EAAs, from 60 to 80% by weight EAAs, from 70 to 90% by weight EAAs, from 60 to 70% by weight EAAs, from 70 to 80% by weight EAAs, from 80 to 90% by weight EAAs, and from 90 to 100% by weight EAAs. In some embodiments the third polypeptide sequence comprises from 90% to 100% EAAs and CEAAAs. In some embodiments the third polypeptide sequence comprises 100% EAAs and CEAAAs.

[00200] In some embodiments the PDCAAS of the third polypeptide sequence is at least about 1.4, at least about 1.5, at least about 1.6, at least about 1.7, at least about 1.8, at

least about 1.9, at least about 2.0, at least about 2.1, at least about 2.2, at least about 2.3, at least about 2.4, at least about 2.5, at least about 2.6, at least about 2.7, at least about 2.8, at least about 2.9, at least about 3.0, at least about 3.1, at least about 3.2, at least about 3.3, at least about 3.4, at least about 3.5, or about 3.6. In some embodiments of the nutritive proteins, the PDCAAS of the third polypeptide sequence is from about 1.4 to about 1.6, from about 1.4 to about 1.8, from about 1.4 to about 2.0, from about 1.4 to about 2.2, from about 1.4 to about 2.4, from about 1.4 to about 2.6, from about 1.4 to about 2.8, from about 1.4 to about 3.0, from about 1.4 to about 3.2, from about 1.4 to about 3.4, or from about 1.4 to about 3.6.

[00201] In some embodiments the third polypeptide sequence comprises at least 20 amino acids and at least 50% by weight of the amino acids of the third polypeptide sequence are essential amino acids. In some embodiments the third polypeptide sequence has a protein digestibility corrected amino acid score (PDCAAS) of at least about 1.4. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 1.8.

[00202] In some embodiments of the nutritive protein, the third polypeptide sequence comprises at least 60% by weight essential amino acids. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 2.2.

[00203] In some embodiments of the nutritive protein, the third polypeptide sequence comprises at least 70% by weight essential amino acids. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 2.5.

[00204] In some embodiments of the nutritive protein, the third polypeptide sequence comprises at least 80% by weight essential amino acids. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 2.9.

[00205] In some embodiments of the nutritive protein, the third polypeptide sequence comprises at least 90% by weight essential amino acids. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 3.2.

[00206] In some embodiments of the nutritive protein, the third polypeptide sequence comprises a weight ratio of each non-essential amino acid to the total number of amino acids that is equivalent to the weight ratio of each non-essential amino acid to the total number of amino acids in a benchmark protein. In some embodiments the benchmark protein is selected from whey protein, egg protein and soy protein.

[00207] In some embodiments of the nutritive protein, the third polypeptide sequence comprises at least 100% essential or conditionally essential amino acids. In some embodiments the third polypeptide sequence has a PDCAAS of from about 1.4 to about 3.6.

[00208] In some embodiments of the nutritive proteins the third polypeptide sequence comprises a naturally-occurring protein or fragment thereof.

[00209] In some embodiments of the nutritive protein, the third polypeptide sequence comprises an altered amino acid sequence compared to the amino acid sequence of a naturally-occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally-occurring reference protein. In some embodiments the third polypeptide sequence comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally-occurring reference protein.

[00210] In some embodiments of the nutritive protein, the third polypeptide sequence is selected from: A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00211] In some embodiments of the nutritive protein, the third polypeptide sequence is selected from: A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-**

11; C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; F.

an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00212] In some embodiments of the nutritive protein, the third polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to at least one amino acid sequence selected from: A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**; B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

[00213] In some embodiments the nutritive protein comprises first and third polypeptide sequences that are the same.

[00214] In some embodiments the nutritive protein comprises first and third polypeptide sequences that are different.

[00215] In some embodiments the nutritive protein comprises first and third polypeptide sequences and further comprise a second polypeptide sequence comprising at least 3 amino acids, wherein the second polypeptide sequence is flanked by the first and third polypeptide sequences.

[00216] In some embodiments the EAA content of the nutritive protein is less than 100%. In some embodiments the NEAAs present in such a nutritive protein may have any distribution of NEAAs. In some embodiments the nutritive protein does not comprise at least one NEAA.

[00217] In some embodiments the NEAAs present in such a nutritive protein have a distribution of NEAAs that provides a balanced mix of NEAAs. In some embodiments the NEAAs present in the nutritive protein are present in proportions similar to the proportions of NEAAs present in a benchmark dietary protein source. For example, if a nutritive protein comprises about 60% EAAs and about 40% NEAAs, the NEAA content of the nutritive protein may be selected or designed to be similar to the proportions of each NEAA present in a benchmark protein.

[00218] The NEAA distribution for Whey is shown in **Table 5A**. These values come from a basic tabulation of the average relative AA fractions for the proteins found in Bovine Whey. The fractions used for this analysis are: Beta-lactoglobulin (UniProtID: P02754), alpha-lactoglobulin (P00711), and serum albumin (P02769), assuming relative protein percentages of 65%, 25%, and 10%, respectively. See H.-D. Belitz, W. Grosch, and P. Schieberle. Food Chemistry 3rd revised Edition. Springer-Verlag, 2004. Note that Belitz et al. also point to a small contribution from immunoglobulins which are varied in amino acid composition. For the purposes of this analysis that fraction was ignored. Similarly, Belitz et al. point to a large proteose-peptone (digested proteins) component. Because that component is assumed to be primarily fragments of the three proteins listed here it was ignored.

Table 5A: Whey NEAA Target Percentages

% NEAA	A	R	N	D	Q	E	G	P	S
50	9.49	1.87	3.82	7.27	5.74	8.87	3.68	4.12	5.14
40	7.59	1.50	3.05	5.82	4.60	7.09	2.94	3.29	4.11
30	5.69	1.12	2.29	4.36	3.45	5.32	2.21	2.47	3.09
20	3.79	0.75	1.53	2.91	2.30	3.55	1.47	1.65	2.06
10	1.90	0.37	0.76	1.45	1.15	1.77	0.74	0.82	1.03

[00219] The NEAA distribution for Soy is shown in Table 5B. These values come from a basic tabulation of the average relative AA fractions for the proteins found in *Glycine Max*: Glycinin subunits 1-5 (UniProtID: P04776, P04405, P11828, P02858, and P04347, respectively), Beta-conglycinin A (P13916), Beta-conglycinin AP (P11827), and Beta-conglycinin B (P25974), assuming a glycinin/conglycinin breakdown of 64% to 34% and

equal contributions from each glycinin/Beta-conglycinin subunits with relative protein percentages of 0.128, 0.128, 0.128, 0.128, 0.128, 0.12, 0.12, and 0.12, respectively. See Adachi M, Kanamori J., Masuda T., Yagasaki K., Kitamura K., Mikami B., and S. Utsumi. Crystal structure of soybean 11S globulin: Glycinin A3B4 homohexamer. Proc. Natl. Acad. Sci. USA. (2003) 100: 7395-7400; and . J. H. Kuipers. Aggregation of peptides in soy protein isolate hydrolysates; The individual contributions of glycinin- and β -conglycinin-derived peptides. Ph.D. Thesis. Wageningen University, The Netherlands, 2007.

Table 5B: Soy NEAA Target Percentages

% NEAA	A	R	N	D	Q	E	G	P	S
50	2.77	7.61	6.07	3.72	8.65	9.17	2.67	4.44	4.89
40	2.22	6.09	4.86	2.97	6.92	7.34	2.14	3.55	3.92
30	1.66	4.57	3.64	2.23	5.19	5.50	1.60	2.66	2.94
20	1.11	3.04	2.43	1.49	3.46	3.67	1.07	1.78	1.96
10	0.55	1.52	1.21	0.74	1.73	1.83	0.53	0.89	0.98

[00220] The NEAA distribution for Egg is shown in Table 5C. These values come from a basic tabulation of the average relative AA fractions for the proteins found in *Gallus gallus*: Ovalbumin (UniProtID: P01012), ovotransferrin (P02789), and ovomucoid (P68390), assuming relative protein percentages of 0.72, 0.14, and 0.14, respectively. See H.-D. Belitz, W. Grosch, and P. Schieberle. Food Chemistry 3rd revised Edition. Springer-Verlag, 2004.

Table 5C: Egg NEAA Target Percentages

% NEAA	A	R	N	D	Q	E	G	P	S
50	5.62	5.75	5.28	4.81	4.08	10.40	3.14	3.51	7.41
40	4.50	4.60	4.23	3.85	3.26	8.32	2.51	2.81	5.93
30	3.37	3.45	3.17	2.89	2.45	6.24	1.88	2.11	4.45
20	2.25	2.30	2.11	1.92	1.63	4.16	1.25	1.40	2.96

10	1.12	1.15	1.06	0.96	0.82	2.08	0.63	0.70	1.48
----	------	------	------	------	------	------	------	------	------

[00221] In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 50% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 55% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least weight 60% of the ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 70% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 80% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 90% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 95% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the nutritive protein comprises each NEAA at a weight ratio that is at least 99% of the weight ratio of that NEAA present in a benchmark dietary protein source. In some embodiments the benchmark dietary protein source is whey protein. In some embodiments the benchmark dietary protein source is egg protein. In some embodiments the benchmark dietary protein source is soy protein. In some embodiments the benchmark dietary protein source is a blended average of at least two of from 25% to 75% whey protein, from 25% to 75% egg protein, and from 25% to 75% soy protein.

[00222] The Euclidean distance of a given amino acid distribution (p) to a target distribution (t) is calculated using the following formula

$$Distance = \sqrt{\sum_{i \in AA} (p_i - t_i)^2},$$

where $AA = \{H, I, L, K, SAA, AAA, T, W, V, A, R, N, D, Q, E, G, P, \text{ and } S\}$, p_i is the fraction of amino acid i in the given protein distribution, and t_i is the fraction of amino acid i in the target distribution. This metric corresponds to the square root of the sum of the squared error for each amino acid (EAAs, CEAAAs, and NEAAAs), and measures the percent distance between a given and target amino acid distribution.

[00223] In some embodiments herein the Euclidian distance of the amino acid distribution profile of a nutritive protein or fragment thereof is determined relative to a target amino acid distribution profile. In some embodiments the target amino acid distribution profile comprises a target distribution for EAAs. In some embodiments the target amino acid distribution profile comprises a target distribution for NEAAs. In some embodiments the target amino acid distribution profile comprises a target distribution for CEAAAs. In some embodiments the target amino acid distribution profile comprises a target distribution for at least two of EAAs, CEAAAs, and NEAAs. In some embodiments the target amino acid distribution profile comprises a target distribution for EAAs, CEAAAs, and NEAAs. In some embodiments the target amino acid distribution profile of EAAs and CEAAAs is a PDCAAS score. In some embodiments the target amino acid distribution profile of NEAAs is the NEAA amino acid distribution profile of NEAAs present in a benchmark dietary protein source.

[00224] In some embodiments the target amino acid distribution profile of EAAs and CEAAAs is a PDCAAS score of at least about 1.4, at least about 1.5, at least about 1.6, at least about 1.7, at least about 1.8, at least about 1.9, at least about 2.0, at least about 2.1, at least about 2.2, at least about 2.3, at least about 2.4, at least about 2.5, at least about 2.6, at least about 2.7, at least about 2.8, at least about 2.9, at least about 3.0, at least about 3.1, at least about 3.2, at least about 3.3, at least about 3.4, at least about 3.5, or about 3.6. In some embodiments the target amino acid distribution profile of EAAs and CEAAAs is a PDCAAS score of at least about 1.4 to about 1.6, from about 1.4 to about 1.8, from about 1.4 to about 2.0, from about 1.4 to about 2.2, from about 1.4 to about 2.4, from about 1.4 to about 2.6, from about 1.4 to about 2.8, from about 1.4 to about 3.0, from about 1.4 to about 3.2, from about 1.4 to about 3.4, or from about 1.4 to about 3.6. In some embodiments the target amino acid distribution profile comprises at least 50% by weight EAAs. In some embodiments the target amino acid distribution profile comprises at least 55% by weight EAAs, at least 60% by weight EAAs, at least 65% by weight EAAs, at least 70% by weight EAAs, at least by weight 75% EAAs, at least 80% by weight EAAs, at least 85% by weight EAAs, at least 90% by weight EAAs, or at least 95% by weight EAAs, at least 96% by weight EAAs, at least 97% by weight EAAs, at least 98% by weight EAAs, at least 99% by weight EAAs, or 100% EAAs. In some embodiments the target amino acid distribution profile comprises from 50 to 100% by weight EAAs, from 60 to 100% by weight EAAs, from 70 to 100% by weight EAAs, from 80 to 90% by weight EAAs, from 60 to 90% by weight EAAs, from 60 to 80%

by weight EAAs, from 70 to 90% by weight EAAs, from 60 to 70% by weight EAAs, from 70 to 80% by weight EAAs, from 80 to 90% by weight EAAs, and from 90 to 100% by weight EAAs. In some embodiments the target amino acid distribution profile comprises from 90% to 100% EAAs and CEAAAs (that is, the combined fraction of essential amino acids and conditionally essential amino acids in the nutritive proteins is from 90% to 100%.

[00225] In some embodiments herein a fragment of a naturally-occurring nutritive protein is produced and optionally isolated. In some embodiments the fragment comprises at least 50 amino acids. In some embodiments the fragment consists of at least 50 amino acids.

[00226] In some embodiments an isolated recombinant nutritive protein is provided. In some embodiments the nutritive protein comprises a first polypeptide sequence, and the first polypeptide sequence comprises a fragment of at least 50 amino acids of a naturally-occurring nutritive protein. In some embodiments the nutritive protein is isolated. In some embodiments the nutritive protein is recombinant. In some embodiments the nutritive protein comprises a first polypeptide sequence comprising a fragment of at least 50 amino acids of a naturally-occurring nutritive protein. In some embodiments the nutritive proteins are isolated recombinant nutritive proteins. In some embodiments the isolated recombinant nutritive proteins disclosed herein are provided in a non-isolated and/or non-recombinant form.

[00227] In some embodiments the nutritive protein is a nutritive protein other than at least one nutritive protein selected from egg proteins such as ovalbumin, ovotransferrin, and ovomucoid; meat proteins such as myosin, actin, tropomyosin, collagen, and troponin; milk proteins such as whey and casein; cereal proteins such as casein, alpha1 casein, alpha2 casein, beta casein, kappa casein, beta-lactoglobulin, alpha-lactalbumin, glycinin, beta-conglycinin, glutelin, prolamine, gliadin, glutenin, albumin, globulin; chicken muscle proteins such as albumin, enolase, creatine kinase, phosphoglycerate mutase, triosephosphate isomerase, apolipoprotein, ovotransferrin, phosphoglucomutase, phosphoglycerate kinase, glycerol-3-phosphate dehydrogenase, glyceraldehyde 3-phosphate dehydrogenase, hemoglobin, cofilin, glycogen phosphorylase, fructose-1,6-bisphosphatase, actin, myosin, tropomyosin a-chain, casein kinase, glycogen phosphorylase, fructose-1,6-bisphosphatase, aldolase, tubulin, vimentin, endoplasmic reticulum chaperone, lactate dehydrogenase, destrin, transthyretin, fructose bisphosphate aldolase, carbonic anhydrase, aldehyde dehydrogenase, annexin, adenosyl homocysteinase; pork muscle proteins such as actin, myosin, enolase, titin, cofilin, phosphoglycerate kinase, enolase, pyruvate dehydrogenase, glycogen phosphorylase,

triosephosphate isomerase, myokinase; and fish proteins such as parvalbumin, pyruvate dehydrogenase, desmin, and triosephosphate isomerase.

[00228] Phenylketonuria (PKU) is an autosomal recessive metabolic genetic disorder characterized by a mutation in the gene for the hepatic enzyme phenylalanine hydroxylase (PAH), rendering it nonfunctional. This enzyme is necessary to metabolize phenylalanine to tyrosine. When PAH activity is reduced, phenylalanine accumulates and is converted into phenylpyruvate (also known as phenylketone), which is detected in the urine. Untreated children are normal at birth, but fail to attain early developmental milestones, develop microcephaly, and demonstrate progressive impairment of cerebral function. Hyperactivity, EEG abnormalities and seizures, and severe learning disabilities are major clinical problems later in life. A characteristic odor of skin, hair, sweat and urine (due to phenylacetate accumulation); and a tendency to hypopigmentation and eczema are also observed. All PKU patients must adhere to a special diet low in Phe. Accordingly, nutritive proteins comprising a low number or no Phe residues are desirable for PKU patients. Such proteins can be obtained by selecting nutritive proteins provided herein that have few or no Phe residues. Accordingly, in some embodiments the nutritive protein comprises a ratio of Phe residues to total amino acid residues equal to or lower than 5%, 4%, 3%, 2%, or 1%. In some embodiments the nutritive protein comprises 10 or fewer Phe residues, 9 or fewer Phe residues, 8 or fewer Phe residues, 7 or fewer Phe residues, 6 or fewer Phe residues, 5 or fewer Phe residues, 4 or fewer Phe residues, 3 or fewer Phe residues, 2 or fewer Phe residues, 1 Phe residue, or no Phe residues. In some embodiments, the nutritive protein comprises no Phe residues.

[00229] Arginine is a conditionally nonessential amino acid, meaning most of the time it can be manufactured by the human body, and does not need to be obtained directly through the diet. Individuals who have poor nutrition, the elderly, or people with certain physical conditions (e.g., sepsis) may not produce sufficient amounts of arginine and therefore need to increase their intake of foods containing arginine. Arginine is believed to have beneficial health properties, including reducing healing time of injuries (particularly bone), and decreasing blood pressure, particularly high blood pressure during high risk pregnancies (pre-eclampsia). In addition, studies have shown that dietary supplementation with L-arginine is beneficial for enhancing the reproductive performance of pigs with naturally occurring intrauterine growth retardation, enhancing protein deposition and postnatal growth of milk-fed piglets, normalizing plasma glucose levels in streptozotocin-induced diabetic rats, reducing

fat mass in obese Zucker diabetic fatty (ZDF) rats, and improving vascular function in diabetic rats. In order to combine these benefits with at least one utility of the nutritive proteins disclosed herein, in some embodiments of the nutritive proteins disclosed herein the nutritive protein comprises a ration of Arginine residues to total amino acid residues in the nutritive protein of equal to or greater than 3%, equal to or greater than 4%, equal to or greater than 5%, equal to or greater than 6%, equal to or greater than 7%, equal to or greater than 8%, equal to or greater than 9%, equal to or greater than 10%, equal to or greater than 11%, or equal to or greater than 12%.

[00230] Digestibility is a parameter relevant to the nutritive benefits and utility of nutritive proteins. Information relating to the relative completeness of digestion can serve as a predictor of peptide bioavailability (Daniel, H., 2003. Molecular and Integrative Physiology of Intestinal Peptide Transport. *Annual Review of Physiology*, Volume 66, pp. 361-384). In some embodiments nutritive proteins disclosed herein are screened to assess their digestibility. Digestibility of nutritive proteins can be assessed by any suitable method known in the art. In some embodiments digestibility is assessed by a physiologically relevant *in vitro* digestion reaction that includes one or both phases of protein digestion, simulated gastric digestion and simulated intestinal digestion (see, e.g., Moreno, et al., 2005. Stability of the major allergen Brazil nut 2S albumin (Ber e 1) to physiologically relevant *in vitro* gastrointestinal digestion. *FEBS Journal*, pp. 341-352; Martos, G., Contreras, P., Molina, E. & Lopez-Fandino, R., 2010. Egg White Ovalbumin Digestion Mimicking Physiological Conditions. *Journal of Agricultural and food chemistry*, pp. 5640-5648; Moreno, F. J., Mackie, A. R. & Clare Mills, E. N., 2005. Phospholipid interactions protect the milk allergen α -Lactalbumin from proteolysis during *in vitro* digestion. *Journal of agricultural and food chemistry*, pp. 9810-9816). Briefly, test proteins are sequentially exposed to a simulated gastric fluid (SGF) for 120 minutes (the length of time it takes 90% of a liquid meal to pass from the stomach to the small intestine; see Kong, F. & Singh, R. P., 2008. Disintegration of Solid Foods in Human Stomach. *Journal of Food Science*, pp. 67-80) and then transferred to a simulated duodenal fluid (SDF) to digest for an additional 120 minutes. Samples at different stages of the digestion (e.g., 2, 5, 15, 30, 60 and 120 min) are analyzed by electrophoresis (e.g., chip electrophoresis or SDS-PAGE) to monitor the size and amount of intact protein as well as any large digestion fragments (e.g., larger than 4 kDa). The disappearance of protein over time indicates the rate at which the protein is digested in the assay. By monitoring the amount of intact protein observed over time, the

half-life ($\tau_{1/2}$) of digestion is calculated for SGF and, if intact protein is detected after treatment with SGF, the $\tau_{1/2}$ of digestion is calculated for SIF. This assay can be used to assess comparative digestibility (i.e., against a benchmark protein such as whey) or to assess absolute digestibility. In some embodiments the digestibility of the nutritive protein is higher (i.e., the SGF $\tau_{1/2}$ and/or SIF $\tau_{1/2}$ is shorter) than whey protein. In some embodiments the nutritive protein has a SGF $\tau_{1/2}$ of 30 minutes or less, 20 minutes or less, 15 minutes or less, 10 minutes or less, 5 minutes or less, 4 minutes or less, 3 minutes or less, 2 minutes or less or 1 minute or less. In some embodiments the nutritive protein has a SIF $\tau_{1/2}$ of 30 minutes or less, 20 minutes or less, 15 minutes or less, 10 minutes or less, 5 minutes or less, 4 minutes or less, 3 minutes or less, 2 minutes or less or 1 minute or less. In some embodiments the nutritive protein is not detectable in one or both of the SGF and SIF assays by 2 minutes, 5 minutes, 15 minutes, 30 minutes, 60 minutes, or 120 minutes. In some embodiments the nutritive protein is digested at a constant rate and/or at a controlled rate in one or both of SGF and SIF. In such embodiments the rate of digestion of the nutritive protein may not be optimized for the highest possible rate of digestion. In such embodiments the rate of absorption of the protein following ingestion by a mammal may be slower and the total time period over which absorption occurs following ingestion may be longer than for nutritive proteins of similar amino acid composition that are digested at a faster initial rate in one or both of SGF and SIF. In some embodiments the nutritive protein is completely or substantially completely digested in SGF. In some embodiments the nutritive protein is substantially not digested or not digested by SGF; in most such embodiments the protein is digested in SIF. Assessing protein digestibility can also provide insight into a protein's potential allergenicity, as proteins or large fragments of proteins that are resistant to digestive proteases can have a higher risk of causing an allergenic reaction (Goodman, R. E. et al., 2008. Allergenicity assessment of genetically modified crops - what makes sense? *Nature Biotechnology*, pp. 73-81). To detect and identify peptides too small for chip electrophoresis analysis, liquid chromatography and mass spectrometry can be used. In SGF samples, peptides can be directly detected and identified by LC/MS. SIF protein digestions may require purification to remove bile acids before detection and identification by LC/MS.

[00231] In some embodiments digestibility of a nutritive protein is assessed by identification and quantification of digestive protease recognition sites in the protein amino acid sequence. In some embodiments the nutritive protein comprises at least one protease

recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

[00232] As used herein, a “pepsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by pepsin. In some embodiments it is a peptide bond after (i.e., downstream of) an amino acid residue selected from Phe, Trp, Tyr, Leu, Ala, Glu, and Gln, provided that the following residue is not an amino acid residue selected from Ala, Gly, and Val.

[00233] As used herein, a “trypsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by trypsin. In some embodiments it is a peptide bond after an amino acid residue selected from Lys or Arg, provided that the following residue is not a proline.

[00234] As used herein, a “chymotrypsin recognition site” is any site in a polypeptide sequence that is experimentally shown to be cleaved by chymotrypsin. In some embodiments it is a peptide bond after an amino acid residue selected from Phe, Trp, Tyr, and Leu.

[00235] Disulfide bonded cysteine residues in a protein tend to reduce the rate of digestion of the protein compared to what it would be in the absence of the disulfide bond. For example, it has been shown that the rate of digestion of the protein b-lactoglobulin is increased when its disulfide bridges are cleaved (I. M. Reddy, N. K. D. Kella, and J. E. Kinsella. “Structural and Conformational Basis of the Resistance of B-Lactoglobulin to Peptic and Chymotryptic Digestion”. J. Agric. Food Chem. 1988, 36, 737-741). Accordingly, digestibility of a nutritive protein with fewer disulfide bonds tends to be higher than for a comparable nutritive protein with a greater number of disulfide bonds. In some embodiments the nutritive proteins disclosed herein are screened to identify the number of cysteine residues present in each and in particular to allow selection of a nutritive protein comprising a relatively low number of cysteine residues. For example, naturally occurring nutritive proteins or fragments may be identified that comprise a no Cys residues or that comprise a relatively low number of Cys residues, such as 10 or fewer Cys residues, 9 or fewer Cys residues, 8 or fewer Cys residues, 7 or fewer Cys residues, 6 or fewer Cys residues, 5 or fewer Cys residues, 4 or fewer Cys residues, 3 or fewer Cys residues, 2 or fewer Cys residues, 1 Cys residue, or no Cys residues. In some embodiments one or more Cys residues in a naturally occurring nutritive protein or fragment thereof is removed by deletion and/or by substitution with another amino acid. In some embodiments 1 Cys residue

is deleted or replaced, 1 or more Cys residues are deleted or replaced, 2 or more Cys residues are deleted or replaced, 3 or more Cys residues are deleted or replaced, 4 or more Cys residues are deleted or replaced, 5 or more Cys residues are deleted or replaced, 6 or more Cys residues are deleted or replaced, 7 or more Cys residues are deleted or replaced, 8 or more Cys residues are deleted or replaced, 9 or more Cys residues are deleted or replaced, or 10 or more Cys residues are deleted or replaced. In some embodiments the nutritive protein of this disclosure comprises a ratio of Cys residues to total amino acid residues equal to or lower than 5%, 4%, 3%, 2%, or 1%. In some embodiments the nutritive protein comprises 10 or fewer Cys residues, 9 or fewer Cys residues, 8 or fewer Cys residues, 7 or fewer Cys residues, 6 or fewer Cys residues, 5 or fewer Cys residues, 4 or fewer Cys residues, 3 or fewer Cys residues, 2 or fewer Cys residues, 1 Cys residue, or no Cys residues. In some embodiments, the nutritive protein comprises 1 or fewer Cys residues. In some embodiments, the nutritive protein comprises no Cys residues.

[00236] Alternatively or in addition, disulfide bonds that are or may be present in a nutritive protein may be removed. Disulfides can be removed using chemical methods by reducing the disulfide to two thiol groups with reducing agents such as beta-mercaptoethanol, dithiothreitol (DTT), or tris(2-carboxyethyl)phosphine (TCEP). The thiols can then be covalently modified or “capped” with reagents such as iodoacetamide, N-ethylmaleimide, or sodium sulfite (see, e.g., Crankshaw, M. W. and Grant, G. A. 2001. Modification of Cysteine. Current Protocols in Protein Science. 15.1.1–15.1.18).

[00237] Eukaryotic proteins are often glycosylated, and the carbohydrate chains that are attached to proteins serve various functions. N-linked and O-linked glycosylation are the two most common forms of glycosylation occurring in proteins. N-linked glycosylation is the attachment of a sugar molecule to a nitrogen atom in an amino acid residue in a protein. N-linked glycosylation occurs at Asparagine and Arginine residues. O-linked glycosylation is the attachment of a sugar molecule to an oxygen atom in an amino acid residue in a protein. O-linked glycosylation occurs at Threonine and Serine residues.

[00238] Glycosylated proteins are often more soluble than their un-glycosylated forms. In terms of protein drugs, proper glycosylation usually confers high activity, proper antigen binding, better stability in the blood, etc. However, glycosylation necessarily means that a protein “carries with it” sugar moieties. Such sugar moieties may reduce the usefulness of the nutritive proteins of this disclosure including recombinant nutritive proteins. For

example, as demonstrated in the examples, a comparison of digestion of glycosylated and non-glycosylated forms of the same proteins shows that the non-glycosylated forms are digested more quickly than the glycosylated forms. For these reasons, in some embodiments the nutritive proteins according to the disclosure comprise low or no glycosylation. For example, in some embodiments the nutritive proteins comprise a ratio of non-glycosylated to total amino acid residues of at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99%. In some embodiments the nutritive proteins do not comprise any glycosylation.

[00239] In some embodiments, the nutritive protein according to the disclosure is deglycosylated after it is produced or after it is isolated. Nutritive proteins of low or no glycosylation may be made by any method known in the art. For example, enzymatic and/or chemical methods may be used (Biochem. J. (2003) 376, p339-350.). Enzymes are produced commercially at research scales for the removal of N-linked and O-linked oligosaccharides. Chemical methods include use of trifluoromethanesulfonic acid to selectively break N-linked and O-linked peptide-saccharide bonds. This method often results in a more complete deglycosylation than does the use of enzymatic methods.

[00240] In other embodiments, the nutritive protein according to the disclosure is produced with low or no glycosylation by chemical synthesis or by a host organism. Most bacteria and other prokaryotes have very limited capabilities to glycosylate proteins, especially heterologous proteins. Accordingly, in some embodiments of this disclosure a nutritive protein is made recombinantly in a microorganism such that the level of glycosylation of the recombinant protein is low or no glycosylation. In some embodiments the level of glycosylation of the recombinant nutritive protein is lower than the level of glycosylation of the protein as it occurs in the organism from which it is derived.

[00241] In some embodiments a nutritive protein or polypeptide according to the disclosure comprises a ratio of amino acids selected from Asn, Arg, Ser, and Thr to total amino acids of 20% or less, 19% or less, 18 % or less, 17% or less, 16% or less, 15% or less, 14% or less, 13% or less, 12% or less, 11% or less, 10% or less, 9% or less, 8% or less, 7% or less, 6% or less, 5% or less, 4% or less, 3% or less, 2% or less, 1% or less. In some embodiments, a nutritive protein or polypeptide according to the disclosure comprises no amino acids selected from Asn, Arg, Ser, and Thr. In some embodiments a nutritive protein or polypeptide according to the disclosure comprises fewer than 20, fewer than 19, fewer than

18, fewer than 17, fewer than 16, fewer than 15, fewer than 14, fewer than 13, fewer than 12, fewer than 11, fewer than 10, fewer than 9, fewer than 8, fewer than 7, fewer than 6, fewer than 5, fewer than 4, fewer than 3, fewer than 2, fewer than 1, or no amino acids selected from Asn, Arg, Ser, and Thr.

[00242] In some embodiments of the nutritive proteins disclosed herein the nutritive protein is soluble. Solubility can be measured by any method known in the art. In some embodiments solubility is examined by centrifuge concentration followed by protein concentration assays. Samples of nutritive proteins in 20 mM HEPES pH 7.5 are tested for protein concentration according to protocols using two methods, Coomassie Plus (Bradford) Protein Assay (Thermo Scientific) and Bicinchoninic Acid (BCA) Protein Assay (Sigma-Aldrich). Based on these measurements 10 mg of protein is added to an Amicon Ultra 3 kDa centrifugal filter (Millipore). Samples are concentrated by centrifugation at 10,000 Xg for 30 minutes. The final, now concentrated, samples are examined for precipitated protein and then tested for protein concentration as above using two methods, Bradford and BCA.

[00243] In some embodiments the nutritive proteins have a final solubility limit of at least 5 g/L, 10 g/L, 20 g/L, 30 g/L, 40 g/L, 50 g/L, or 100 g/L at physiological pH. In some embodiments the nutritive proteins are greater than 50%, greater than 60%, greater than 70%, greater than 80%, greater than 90%, greater than 95%, greater than 96%, greater than 97%, greater than 98%, greater than 99%, or greater than 99.5% soluble with no precipitated protein observed at a concentration of greater than 5 g/L, or 10 g/L, or 20 g/L, or 30 g/L, or 40 g/L, or 50 g/L, or 100 g/L at physiological pH. In some embodiments, the solubility of the nutritive protein is higher than those typically reported in studies examining the solubility limits of whey (12.5 g/L; Pelegrine et al., *Lebensm.-Wiss. U.-Technol.* 38 (2005) 77-80) and soy (10 g/L; Lee et al., *JAOC* 80(1) (2003) 85-90).

[00244] As used herein, a “stable” protein is one that resists changes (e.g., unfolding, oxidation, aggregation, hydrolysis, etc.) that alter the biophysical (e.g., solubility), biological (e.g., digestibility), or compositional (e.g. proportion of Leucine amino acids) traits of the protein of interest.

[00245] Protein stability can be measured using various assays known in the art and nutritive proteins disclosed herein and having stability above a threshold can be selected. In some embodiments a protein is selected that displays thermal stability that is comparable to or better than that of whey protein. Thermal stability is a property that can help predict the

shelf life of a nutritive protein. In some embodiments of the assay stability of nutritive protein samples is determined by monitoring aggregation formation using size exclusion chromatography (SEC) after exposure to extreme temperatures. Aqueous samples of the protein to be tested are placed in a heating block at 90°C and samples are taken after 0, 1, 5, 10, 30 and 60 min for SEC analysis. Protein is detected by monitoring absorbance at 214nm, and aggregates are characterized as peaks eluting faster than the protein of interest. No overall change in peak area indicates no precipitation of protein during the heat treatment. Whey protein has been shown to rapidly form ~ 80% aggregates when exposed to 90°C in such an assay.

[00246] In some embodiments the thermal stability of a nutritive protein is determined by heating a sample slowly from 25°C to 95°C in presence of a hydrophobic dye (e.g., ProteoStat® Thermal shift stability assay kit, Enzo Life Sciences) that binds to aggregated proteins that are formed as the protein denatures with increasing temperature (Niesen, F. H., Berglund, H. & Vadadi, M., 2007. The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nature Protocols*, Volume 2, pp. 2212-2221). Upon binding, the dye's fluorescence increases significantly, which is recorded by an rtPCR instrument and represented as the protein's melting curve (Lavinder, J. J., Hari, S. B., Sullivan, B. J. & Magilery, T. J., 2009. High-Throughput Thermal Scanning: A General, Rapid Dye-Binding Thermal Shift Screen for Protein Engineering. *Journal of the American Chemical Society*, pp. 3794-3795). After the thermal shift is complete, samples are examined for insoluble precipitates and further analyzed by analytical size exclusion chromatography (SEC).

[00247] In some embodiments a nutritive protein of this disclosure shows resistance to aggregation, exhibiting, for example, less than 80% aggregation, 10% aggregation, or no detectable aggregation at elevated temperatures (e.g, 50°C, 60°C, 70°C, 80°C, 85°C, 90°C, or 95°C).

[00248] One benefit of stable nutritive proteins as disclosed herein is that they may be able to be stored for an extended period of time before use, in some instances without the need for refrigeration or cooling. In some embodiments, nutritive proteins are processed into a dry form (e.g., by lyophilization). In some embodiments, nutritive proteins are stable upon lyophilization. In some embodiments, such lyophilized nutritive proteins maintain their stability upon reconstitution (e.g., liquid formulation).

[00249] For most embodiments it is preferred that the nutritive protein not exhibit inappropriately high allergenicity. Accordingly, in some embodiments the potential allergenicity of the nutritive protein is assessed. This can be done by any suitable method known in the art. In some embodiments an allergenicity score is calculated. The allergenicity score is a primary sequence based metric based on WHO recommendations (<http://www.fao.org/ag/agn/food/pdf/allergygm.pdf>) for assessing how similar a protein is to any known allergen, the primary hypothesis being that high percent identity between a target and a known allergen is likely indicative of cross reactivity. For a given protein, the likelihood of eliciting an allergic response can be assessed via one or both of a complimentary pair of sequence homology based tests. The first test determines the protein's percent identity across the entire sequence via a global-global sequence alignment to a database of known allergens using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. It has been suggested that proteins with less than 50% global homology are unlikely to be allergenic (Goodman R. E. et al. Allergenicity assessment of genetically modified crops—what makes sense? Nat. Biotech. 26, 73-81 (2008); Aalberse R. C. Structural biology of allergens. J. Allergy Clin. Immunol. 106, 228-238 (2000)).

[00250] In some embodiments of a nutritive protein, the nutritive protein has less than 50% global homology to any known allergen in the database used for the analysis. In some embodiments a cutoff of less than 40% homology is used. In some embodiments a cutoff of less than 30% homology is used. In some embodiments a cutoff of less than 20% homology is used. In some embodiments a cutoff of less than 10% homology is used. In some embodiments a cutoff of from 40% to 50% is used. In some embodiments a cutoff of from 30% to 50% is used. In some embodiments a cutoff of from 20% to 50% is used. In some embodiments a cutoff of from 10% to 50% is used. In some embodiments a cutoff of from 5% to 50% is used. In some embodiments a cutoff of from 0% to 50% is used. In some embodiments a cutoff of greater than 50% global homology to any known allergen in the database used for the analysis is used. In some embodiments a cutoff of from 50% to 60% is used. In some embodiments a cutoff of from 50% to 70% is used. In some embodiments a cutoff of from 50% to 80% is used. In some embodiments a cutoff of from 50% to 90% is used. In some embodiments a cutoff of from 55% to 60% is used. In some embodiments a cutoff of from 65% to 70% is used. In some embodiments a cutoff of from 70% to 75% is used. In some embodiments a cutoff of from 75% to 80% is used.

[00251] The second test assesses the local allergenicity along the protein sequence by determining the local allergenicity of all possible contiguous 80 amino acid fragments via a global-local sequence alignment of each fragment to a database of known allergens using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. The highest percent identity of any 80 amino acid window with any allergen is taken as the final score for the protein of interest. The WHO guidelines suggest using a 35% identity cutoff with this fragment test. In some embodiments of a nutritive protein, all possible fragments of the nutritive protein have less than 35% local homology to any known allergen in the database used for the analysis using this test. In some embodiments a cutoff of less than 30% homology is used. In some embodiments a cutoff of from 30% to 35% homology is used. In some embodiments a cutoff of from 25% to 30% homology is used. In some embodiments a cutoff of from 20% to 25% homology is used. In some embodiments a cutoff of from 15% to 20% homology is used. In some embodiments a cutoff of from 10% to 15% homology is used. In some embodiments a cutoff of from 5% to 10% homology is used. In some embodiments a cutoff of from 0% to 5% homology is used. In some embodiments a cutoff of greater than 35% homology is used. In some embodiments a cutoff of from 35% to 40% homology is used. In some embodiments a cutoff of from 40% to 45% homology is used. In some embodiments a cutoff of from 45% to 50% homology is used. In some embodiments a cutoff of from 50% to 55% homology is used. In some embodiments a cutoff of from 55% to 60% homology is used. In some embodiments a cutoff of from 65% to 70% homology is used. In some embodiments a cutoff of from 70% to 75% homology is used. In some embodiments a cutoff of from 75% to 80% homology is used.

[00252] Skilled artisans are able to identify and use a suitable database of known allergens for this purpose. In some embodiments the database is custom made by selecting proteins from more than one database source. In some embodiments the custom database comprises pooled allergen lists collected by the Food Allergy Research and Resource Program (<http://www.allergenonline.org/>), UNIPROT annotations (<http://www.uniprot.org/docs/allergen>), and the Structural Database of Allergenic Proteins (SDAP, http://fermi.utmb.edu/SDAP/sdap_lnk.html). This database includes all currently recognized allergens by the International Union of Immunological Societies (IUIS, <http://www.allergen.org/>) as well as a large number of additional allergens not yet officially named. In some embodiments the database comprises a subset of known allergen proteins available in known databases; that is, the database is a custom selected subset of known

allergen proteins. In some embodiments the database of known allergens comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 1,100 proteins, at least 1,200 proteins, at least 1,300 proteins, at least 1,400 proteins, at least 1,500 proteins, at least 1,600 proteins, at least 1,700 proteins, at least 1,800 proteins, at least 1,900 proteins, or at least 2,000 proteins. In some embodiments the database of known allergens comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, or from 1,000 to 2,000 proteins.

[00253] In some embodiments all (or a selected subset) of contiguous amino acid windows of different lengths (e.g., 70, 60, 50, 40, 30, 20, 10, 8 or 6 amino acid windows) of a nutritive protein are tested against the allergen database and peptide sequences that have 100% identity, 95% or higher identity, 90% or higher identity, 85% or higher identity, 80% or higher identity, 75% or higher identity, 70% or higher identity, 65% or higher identity, 60% or higher identity, 55% or higher identity, or 50% or higher identity matches are identified for further examination of potential allergenicity.

[00254] Another method of predicting the allergenicity of a protein is to assess the homology of the protein to a protein of human origin. The human immune system is exposed to a multitude of possible allergenic proteins on a regular basis and has the intrinsic ability to differentiate between the host body's proteins and exogenous proteins. The exact nature of this ability is not always clear, and there are many diseases that arise as a result of the failure of the body to differentiate self from non-self (e.g. arthritis). Nonetheless, the fundamental hypothesis is that proteins that share a degree of sequence homology to human proteins are less likely to elicit an immune response. In particular, it has been shown that for some protein families with known allergenic members (tropomyosins, parvalbumins, caseins), those proteins that bear more sequence homology to their human counterparts relative to known allergenic proteins, are not thought to be allergenic (Jenkins J. A. et al. Evolutionary distance from human homologs reflects allergenicity of animal food proteins. *J. Allergy Clin Immunol.* 120 (2007): 1399-1405). For a given protein, a human homology score is measured by determining the maximum percent identity of the protein to a database of human proteins (e.g., the UNIPROT database) from a global-local alignment using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension

penalty of 2. According to Jenkins et al. (Jenkins J. A. et al. Evolutionary distance from human homologs reflects allergenicity of animal food proteins. *J. Allergy Clin Immunol.* 120 (2007): 1399-1405) proteins with a sequence identity to a human protein above about 62% are less likely to be allergenic. Skilled artisans are able to identify and use a suitable database of known human proteins for this purpose, for example, by searching the UNIPROT database (<http://www.uniprot.org>). In some embodiments the database is custom made by selecting proteins from more than one database source. Of course the database may but need not be comprehensive. In some embodiments the database comprises a subset of human proteins; that is, the database is a custom selected subset of human proteins. In some embodiments the database of human proteins comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 2,000 proteins, at least 3,000 proteins, at least 4,000 proteins, at least 5,000 proteins, at least 6,000 proteins, at least 7,000 proteins, at least 8,000 proteins, at least 9,000 proteins, or at least 10,000 proteins. In some embodiments the database comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, from 1,000 to 2,000 proteins, from 1,000 to 5,000 proteins, or from 5,000 to 10,000 proteins. In some embodiments the database comprises at least 90%, at least 95%, or at least 99% of all known human proteins.

[00255] In some embodiments of a nutritive protein, the nutritive protein is at least 20% homologous to a human protein. In some embodiments a cutoff of at least 30% homology is used. In some embodiments a cutoff of at least 40% homology is used. In some embodiments a cutoff of at least 50% homology is used. In some embodiments a cutoff of at least 60% homology is used. In some embodiments a cutoff of at least 70% homology is used. In some embodiments a cutoff of at least 80% homology is used. In some embodiments a cutoff of at least 62% homology is used. In some embodiments a cutoff of from at least 20% homology to at least 30% homology is used. In some embodiments a cutoff of from at least 30% homology to at least 40% homology is used. In some embodiments a cutoff of from at least 50% homology to at least 60% homology is used. In some embodiments a cutoff of from at least 60% homology to at least 70% homology is used. In some embodiments a cutoff of from at least 70% homology to at least 80% homology is used.

[00256] For most embodiments it is preferred that the nutritive protein not exhibit inappropriately high toxicity. Accordingly, in some embodiments the potential toxicity of the nutritive protein is assessed. This can be done by any suitable method known in the art. In some embodiments a toxicity score is calculated by determining the protein's percent identity to databases of known toxic proteins (e.g., toxic proteins identified from the UNIPROT database). A global-global alignment of the protein of interest against the database of known toxins is performed using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2. In some embodiments of a nutritive protein, the nutritive protein is less than 35% homologous to a known toxin. In some embodiments a cutoff of less than 35% homology is used. In some embodiments a cutoff of from 30% to 35% homology is used. In some embodiments a cutoff of from 25% to 35% homology is used. In some embodiments a cutoff of from 20% to 35% homology is used. In some embodiments a cutoff of from 15% to 35% homology is used. In some embodiments a cutoff of from 10% to 35% homology is used. In some embodiments a cutoff of from 5% to 35% homology is used. In some embodiments a cutoff of from 0% to 35% homology is used. In some embodiments a cutoff of greater than 35% homology is used. In some embodiments a cutoff of from 35% to 40% homology is used. In some embodiments a cutoff of from 35% to 45% homology is used. In some embodiments a cutoff of from 35% to 50% homology is used. In some embodiments a cutoff of from 35% to 55% homology is used. In some embodiments a cutoff of from 35% to 60% homology is used. In some embodiments a cutoff of from 35% to 70% homology is used. In some embodiments a cutoff of from 35% to 75% homology is used. In some embodiments a cutoff of from 35% to 80% homology is used. Skilled artisans are able to identify and use a suitable database of known toxins for this purpose, for example, by searching the UNIPROT database (<http://www.uniprot.org>). In some embodiments the database is custom made by selecting proteins identified as toxins from more than one database source. In some embodiments the database comprises a subset of known toxic proteins; that is, the database is a custom selected subset of known toxic proteins. In some embodiments the database of toxic proteins comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100, proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 2,000 proteins, at least 3,000 proteins, at least 4,000 proteins, at least 5,000 proteins, at least 6,000 proteins, at least 7,000 proteins, at

least 8,000 proteins, at least 9,000 proteins, or at least 10,000 proteins. In some embodiments the database comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, from 1,000 to 2,000 proteins, from 1,000 to 5,000 proteins, or from 5,000 to 10,000 proteins.

[00257] For some embodiments it is preferred that the nutritive protein not exhibit anti-nutritional activity (“anti-nutricity”), i.e., proteins that have the potential to prevent the absorption of nutrients from food. Examples of anti-nutritive factors include protease inhibitors, which inhibit the actions of trypsin, pepsin and other proteases in the gut, preventing the digestion and subsequent absorption of protein. Accordingly, in some embodiments the potential anti-nutricity of the nutritive protein is assessed. This can be done by any suitable method known in the art. In some embodiments an anti-nutricity score is calculated by determining the protein’s percent identity to databases of known protease inhibitors (e.g., protease inhibitors identified from the UNIPROT database). A global-global alignment of the protein of interest against the database of known protease inhibitors is performed using the FASTA algorithm with the BLOSUM50 substitution matrix, a gap open penalty of 10, and a gap extension penalty of 2, to identify whether the nutritive protein is homologous to a known anti-nutritive protein. In some embodiments of a nutritive protein, the nutritive protein has less than 35% global homology to any known anti-nutritive protein (e.g., any known protease inhibitor) in the database used for the analysis. In some embodiments a cutoff of less than 35% identify is used. In some embodiments a cutoff of from 30% to 35% is used. In some embodiments a cutoff of from 25% to 35% is used. In some embodiments a cutoff of from 20% to 35% is used. In some embodiments a cutoff of from 15% to 35% is used. In some embodiments a cutoff of from 10% to 35% is used. In some embodiments a cutoff of from 5% to 35% is used. In some embodiments a cutoff of from 0% to 35% is used. In some embodiments a cutoff of greater than 35% identify is used. In some embodiments a cutoff of from 35% to 40% is used. In some embodiments a cutoff of from 35% to 45% is used. In some embodiments a cutoff of from 35% to 50% is used. In some embodiments a cutoff of from 35% to 55% is used. In some embodiments a cutoff of from 35% to 60% is used. In some embodiments a cutoff of from 35% to 70% is used. In some embodiments a cutoff of from 35% to 75% is used. In some embodiments a cutoff of from 35% to 80% is used. Skilled artisans are able to identify and use a suitable database of known protease inhibitors for this purpose, for example, by searching the UNIPROT database (<http://www.uniprot.org>). In some embodiments the database is custom made by selecting

proteins identified protease-inhibitors as from more than one database source. In some embodiments the database comprises a subset of known protease inhibitors available in databases; that is, the database is a custom selected subset of known protease inhibitor proteins. In some embodiments the database of known protease inhibitor proteins comprises at least 10 proteins, at least 20 proteins, at least 30 proteins, at least 40 proteins, at least 50 proteins, at least 100 proteins, at least 200 proteins, at least 300 proteins, at least 400 proteins, at least 500 proteins, at least 600 proteins, at least 700 proteins, at least 800 proteins, at least 900 proteins, at least 1,000 proteins, at least 1,100 proteins, at least 1,200 proteins, at least 1,300 proteins, at least 1,400 proteins, at least 1,500 proteins, at least 1,600 proteins, at least 1,700 proteins, at least 1,800 proteins, at least 1,900 proteins, or at least 2,000 proteins. In some embodiments the database of known protease inhibitor proteins comprises from 100 to 500 proteins, from 200 to 1,000 proteins, from 500 to 1,000 proteins, from 500 to 1,000 proteins, or from 1,000 to 2,000 proteins, or from 2,000 to 3,000 proteins.

[00258] In other embodiments a nutritive protein that does exhibit some degree of protease inhibitor activity is used. For example, in some embodiments such a protein may be useful because it delays protease digestion when the nutritive protein is consumed such that the nutritive protein traverses a greater distance within the GI tract before it is digested, thus delaying absorption. For example, in some embodiments the nutritive protein inhibits gastric digestion but not intestinal digestion.

[00259] Delaney B. et al. (Evaluation of protein safety in the context of agricultural biotechnology. Food. Chem. Toxicol. 46 (2008: S71-S97)) suggests that one should avoid both known toxic and anti-nutritive proteins when assessing the safety of a possible food protein. In some embodiments of a nutritive protein, the nutritive protein has a favorably low level of global homology to a database of known toxic proteins and/or a favorably low level of global homology to a database of known anti-nutricity proteins (e.g., protease inhibitors), as defined herein.

[00260] One feature that can enhance the utility of a nutritive protein is its charge (or per amino acid charge). Nutritive proteins with higher charge can in some embodiments exhibit desirable characteristics such as increased solubility, increased stability, resistance to aggregation, and desirable taste profiles. For example, a charged nutritive protein that exhibits enhanced solubility can be formulated into a beverage or liquid formulation that includes a high concentration of nutritive protein in a relatively low volume of solution, thus

delivering a large dose of protein nutrition per unit volume. A charged nutritive protein that exhibits enhanced solubility can be useful, for example, in sports drinks or recovery drinks wherein a user (e.g., an athlete) wants to ingest nutritive protein before, during or after physical activity. A charged nutritive protein that exhibits enhanced solubility can also be particularly useful in a clinical setting wherein a subject (e.g., a patient or an elderly person) is in need of protein nutrition but is unable to ingest solid foods or large volumes of liquids.

[00261] For example, the net charge (Charge_P) of a polypeptide at pH 7 can be calculated using the following formula:

$$\text{Charge}_P = -0.002 - (C)(0.045) - (D)(0.999) - (E)(0.998) + (H)(0.091) + (K)(1.0) + (R)(1.0) - (Y)(-0.001)$$

where C is the number of cysteine residues, D is the number of aspartic acid residues, E is the number of glutamic acid residues, H is the number of histidine residues, K is the number of lysine residues, R is the number of arginine residues and Y is the number of tyrosine residues in the polypeptide. The per amino acid charge (Charge_A) of the polypeptide can be calculated by dividing the net charge (Charge_P) by the number of amino acid residues (N), i.e., $\text{Charge}_A = \text{Charge}_P/N$. (See Bassi S (2007), “A Primer on Python for Life Science Researchers.” *PLoS Comput Biol* 3(11): e199. doi:10.1371/journal.pcbi.0030199).

[00262] One metric for assessing the hydrophilicity and potential solubility of a given protein is the solvation score. Solvation score is defined as the total free energy of solvation (i.e. the free energy change associated with transfer from gas phase to a dilute solution) for all amino acid side chains if each residue were solvated independently, normalized by the total number of residues in the sequence. The side chain solvation free energies are found computationally by calculating the electrostatic energy difference between a vacuum dielectric of 1 and a water dielectric of 80 (by solving the Poisson-Boltzmann equation) as well as the non-polar, Van der Waals energy using a linear solvent accessible surface area model (D. Sitkoff, K. A. Sharp, B. Honig. “Accurate Calculation of Hydration Free Energies Using Macroscopic Solvent Models”. *J. Phys. Chem.* 98, 1994). For amino acids with ionizable sidechains (Arg, Asp, Cys, Glu, His, Lys and Tyr), an average solvation free energy is used based on the relative probabilities for each ionization state at the specified pH. Solvation scores start at 0 and continue into negative values, and the more negative the solvation score, the more hydrophilic and potentially soluble the protein is predicted to be. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or

less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at pH 7. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at pH 7.

[00263] The solvation score is a function of pH by virtue of the pH dependence of the molar ratio of undissociated weak acid ([HA]) to conjugate base ([A⁻]) as defined by the Henderson-Hasselbalch equation:

$$pH = pKa + \log \left(\frac{[A^-]}{[HA]} \right)$$

[00264] All weak acids have different solvation free energies compared to their conjugate bases, and the solvation free energy used for a given residue when calculating the solvation score at a given pH is the weighted average of those two values.

[00265] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at an acidic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at acidic pH.

[00266] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive

protein has a solvation score of -30 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at a basic pH. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at basic pH.

[00267] Accordingly, in some embodiments of a nutritive protein, the nutritive protein has a solvation score of -10 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -15 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -20 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -25 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -30 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -35 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12. In some embodiments of a nutritive protein, the nutritive protein has a solvation score of -40 or less at a pH range selected from 2-3, 3-4, 4-5, 5-6, 6-7, 7-8, 8-9, 9-10, 10-11, and 11-12.

[00268] The aggregation score is a primary sequence based metric for assessing the hydrophobicity and likelihood of aggregation of a given protein. Using the Kyte and Doolittle hydrophobicity scale (Kyte J, Doolittle RF (May 1982) "A simple method for displaying the hydropathic character of a protein". *J. Mol. Biol.* **157** (1): 105-32), which gives hydrophobic residues positive values and hydrophilic residues negative values, the average hydrophobicity of a protein sequence is calculated using a moving average of five residues. The aggregation score is drawn from the resulting plot by determining the area under the curve for values greater than zero and normalizing by the total length of the protein. The underlying hypothesis is that aggregation is the result of two or more hydrophobic patches coming together to exclude water and reduce surface exposure, and the likelihood that a protein will aggregate is a function of how densely packed its hydrophobic (i.e., aggregation prone) residues are. Aggregation scores start at 0 and continue into positive values, and the smaller the aggregation score, the less hydrophobic and potentially less prone to aggregation the protein is predicted to be. In some embodiments of a nutritive protein, the nutritive

protein has an aggregation score of 2 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 1.5 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 1 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.9 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.8 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.7 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.6 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.5 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.4 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.3 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.2 or less. In some embodiments of a nutritive protein, the nutritive protein has an aggregation score of 0.1 or less.

[00269] In some cases, soluble expression is desirable because it can increase the amount and/or yield of the nutritive protein and facilitate one or more of the isolation and purification of the nutritive protein. In some embodiments, the nutritive proteins of this disclosure are solubly expressed in the host organism. Solvation score and aggregation score can be used to predict soluble expression of recombinant nutritive proteins in a host organism. As shown in Example 8, this disclosure provides evidence suggesting that nutritive proteins with solvation scores of ≤ -20 and aggregation scores of ≤ 0.75 are more likely to be recombinantly expressed in a particular *E. coli* expression system. Moreover, the data also suggests that nutritive proteins with solvation scores of ≤ -20 and aggregation scores of ≤ 0.5 are more likely to be solubly expressed in this system. Therefore, in some embodiments the nutritive protein of this disclosure has a solvation score of -20 or less. In some embodiments the nutritive protein has an aggregation score of 0.75 or less. In some embodiments the nutritive protein has an aggregation score of 0.5 or less. In some embodiments the nutritive protein has a solvation score of -20 or less and an aggregation score of 0.75 or less. In some embodiments the nutritive protein has a solvation score of -20 or less and an aggregation score of 0.5 or less.

[00270] Certain free amino acids and mixtures of free amino acids are known to have a bitter or otherwise unpleasant taste. In addition, hydrolysates of common proteins (e.g., whey and soy) often have a bitter or unpleasant taste. In some embodiments, nutritive

proteins disclosed and described herein do not have a bitter or otherwise unpleasant taste. In some embodiments, nutritive proteins disclosed and described herein have a more acceptable taste as compared to at least one of free amino acids, mixtures of free amino acids, and/or protein hydrolysates. In some embodiments, nutritive proteins disclosed and described herein have a taste that is equal to or exceeds at least one of whey protein.

[00271] Proteins are known to have tastes covering the five established taste modalities: sweet, sour, bitter, salty and umami. The taste of a particular protein (or its lack thereof) can be attributed to several factors, including the primary structure, the presence of charged side chains, and the electronic and conformational features of the protein. In some embodiments, nutritive proteins disclosed and described herein are designed to have a desired taste (e.g., sweet, salty, umami) and/or not to have an undesired taste (e.g., bitter, sour). In this context “design” includes, for example, selecting naturally occurring proteins embodying features that achieve the desired taste property, as well as creating muteins of naturally-occurring proteins that have desired taste properties. For example, nutritive proteins can be designed to interact with specific taste receptors, such as sweet receptors (T1R2-T1R3 heterodimer) or umami receptors (T1R1-T1R3 heterodimer, mGluR4, and/or mGluR1). Further, nutritive proteins may be designed not to interact, or to have diminished interaction, with other taste receptors, such as bitter receptors (T2R receptors).

[00272] Nutritive proteins disclosed and described herein can also elicit different physical sensations in the mouth when ingested, sometimes referred to as “mouth feel”. The mouth feel of the nutritive proteins may be due to one or more factors including primary structure, the presence of charged side chains, and the electronic and conformational features of the protein. In some embodiments, nutritive proteins elicit a buttery or fat-like mouth feel when ingested.

[00273] In some embodiments the nutritive protein comprises from 20 to 5,000 amino acids, from 20-2,000 amino acids, from 20-1,000 amino acids, from 20-500 amino acids, from 20-250 amino acids, from 20-200 amino acids, from 20-150 amino acids, from 20-100 amino acids, from 20-40 amino acids, from 30-50 amino acids, from 40-60 amino acids, from 50-70 amino acids, from 60-80 amino acids, from 70-90 amino acids, from 80-100 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids, at least 45 amino acids, at least 50 amino acids, at least 55 amino acids, at least 60 amino acids, at least 65 amino acids, at least 70 amino acids, at least 75 amino acids,

at least 80 amino acids, at least 85 amino acids, at least 90 amino acids, at least 95 amino acids, at least 100 amino acids, at least 105 amino acids, at least 110 amino acids, at least 115 amino acids, at least 120 amino acids, at least 125 amino acids, at least 130 amino acids, at least 135 amino acids, at least 140 amino acids, at least 145 amino acids, at least 150 amino acids, at least 155 amino acids, at least 160 amino acids, at least 165 amino acids, at least 170 amino acids, at least 175 amino acids, at least 180 amino acids, at least 185 amino acids, at least 190 amino acids, at least 195 amino acids, at least 200 amino acids, at least 205 amino acids, at least 210 amino acids, at least 215 amino acids, at least 220 amino acids, at least 225 amino acids, at least 230 amino acids, at least 235 amino acids, at least 240 amino acids, at least 245 amino acids, or at least 250 amino acids. In some embodiments the nutritive protein consists of from 20 to 5,000 amino acids, from 20-2,000 amino acids, from 20-1,000 amino acids, from 20-500 amino acids, from 20-250 amino acids, from 20-200 amino acids, from 20-150 amino acids, from 20-100 amino acids, from 20-40 amino acids, from 30-50 amino acids, from 40-60 amino acids, from 50-70 amino acids, from 60-80 amino acids, from 70-90 amino acids, from 80-100 amino acids, at least 25 amino acids, at least 30 amino acids, at least 35 amino acids, at least 40 amino acids, at least 45 amino acids, at least 50 amino acids, at least 55 amino acids, at least 60 amino acids, at least 65 amino acids, at least 70 amino acids, at least 75 amino acids, at least 80 amino acids, at least 85 amino acids, at least 90 amino acids, at least 95 amino acids, at least 100 amino acids, at least 105 amino acids, at least 110 amino acids, at least 115 amino acids, at least 120 amino acids, at least 125 amino acids, at least 130 amino acids, at least 135 amino acids, at least 140 amino acids, at least 145 amino acids, at least 150 amino acids, at least 155 amino acids, at least 160 amino acids, at least 165 amino acids, at least 170 amino acids, at least 175 amino acids, at least 180 amino acids, at least 185 amino acids, at least 190 amino acids, at least 195 amino acids, at least 200 amino acids, at least 205 amino acids, at least 210 amino acids, at least 215 amino acids, at least 220 amino acids, at least 225 amino acids, at least 230 amino acids, at least 235 amino acids, at least 240 amino acids, at least 245 amino acids, or at least 250 amino acids.

B. Nucleic Acids

[00274] Also provided herein are nucleic acids encoding nutritive polypeptides or proteins. In some embodiments the nucleic acid is isolated. In some embodiments the nucleic acid is purified.

[00275] In some embodiments of the nucleic acid, the nucleic acid comprises a nucleic acid sequence that encodes a first polypeptide sequence disclosed in Section A above. In

some embodiments of the nucleic acid, the nucleic acid consists of a nucleic acid sequence that encodes a first polypeptide sequence disclosed in Section A above. In some embodiments of the nucleic acid, the nucleic acid comprises a nucleic acid sequence that encodes a nutritive protein disclosed in Section A above. In some embodiments of the nucleic acid, the nucleic acid consists of a nucleic acid sequence that encodes a nutritive protein disclosed in Section A above. In some embodiments of the nucleic acid the nucleic acid sequence that encodes the first polypeptide sequence is operatively linked to at least one expression control sequence. For example, in some embodiments of the nucleic acid the nucleic acid sequence that encodes the first polypeptide sequence is operatively linked to a promoter such as a promoter described in Section D below.

[00276] Accordingly, in some embodiments the nucleic acid molecule of this disclosure encodes a polypeptide or protein that itself is a nutritive polypeptide or protein. Such a nucleic acid molecule may be referred to as a “nutrative nucleic acid”. In some embodiments the nutritive nucleic acid encodes a polypeptide or protein that itself comprises at least one of: a) a ratio of branch chain amino acid residues to total amino acid residues of at least 24%; b) a ratio of Leu residues to total amino acid residues of at least 11%; and c) a ratio of essential amino acid residues to total amino acid residues of at least 49%. In some embodiments the nutritive nucleic acid comprises at least 10 nucleotides, at least 20 nucleotides, at least 30 nucleotides, at least 40 nucleotides, at least 50 nucleotides, at least 60 nucleotides, at least 70 nucleotides, at least 80 nucleotides, at least 90 nucleotides, at least 100 nucleotides, at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, at least 800 nucleotides, at least 900 nucleotides, at least 1,000 nucleotides. In some embodiments the nutritive nucleic acid comprises from 10 to 100 nucleotides, from 20 to 100 nucleotides, from 10 to 50 nucleotides, or from 20 to 40 nucleotides. In some embodiments the nutritive nucleic acid comprises all or part of an open reading frame that encodes a naturally occurring nutritive polypeptide or protein. In some embodiments the nutritive nucleic acid consists of an open reading frame that encodes a fragment of a naturally occurring nutritive protein, wherein the open reading frame does not encode the complete naturally occurring nutritive protein.

[00277] In some embodiments the nutritive nucleic acid is a cDNA.

[00278] In some embodiments nucleic acid molecules are provided that comprise a sequence that is at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 99.9% identical to a naturally occurring nutritive nucleic acid. In some embodiments nucleic acids are provided that hybridize under stringent hybridization conditions with at least one reference nutritive nucleic acid.

[00279] The nutritive nucleic acids and fragments thereof provided in this disclosure display utility in a variety of systems and methods. For example, the fragments may be used as probes in various hybridization techniques. Depending on the method, the target nucleic acid sequences may be either DNA or RNA. The target nucleic acid sequences may be fractionated (e.g., by gel electrophoresis) prior to the hybridization, or the hybridization may be performed on samples in situ. One of skill in the art will appreciate that nucleic acid probes of known sequence find utility in determining chromosomal structure (e.g., by Southern blotting) and in measuring gene expression (e.g., by Northern blotting). In such experiments, the sequence fragments are preferably detectably labeled, so that their specific hybridization to target sequences can be detected and optionally quantified. One of skill in the art will appreciate that the nucleic acid fragments of this disclosure may be used in a wide variety of blotting techniques not specifically described herein.

[00280] It should also be appreciated that the nucleic acid sequence fragments disclosed herein also find utility as probes when immobilized on microarrays. Methods for creating microarrays by deposition and fixation of nucleic acids onto support substrates are well known in the art. Reviewed in DNA Microarrays: A Practical Approach (Practical Approach Series), Schena (ed.), Oxford University Press (1999) (ISBN: 0199637768); Nature Genet. 21(1)(suppl):1-60 (1999); Microarray Biochip: Tools and Technology, Schena (ed.), Eaton Publishing Company/BioTechniques Books Division (2000) (ISBN: 1881299376), the disclosures of which are incorporated herein by reference in their entireties. Analysis of, for example, gene expression using microarrays comprising nucleic acid sequence fragments, such as the nucleic acid sequence fragments disclosed herein, is a well-established utility for sequence fragments in the field of cell and molecular biology. Other uses for sequence fragments immobilized on microarrays are described in Gerhold et al., Trends Biochem. Sci. 24:168-173 (1999) and Zweiger, Trends Biotechnol. 17:429-436 (1999); DNA Microarrays: A Practical Approach (Practical Approach Series), Schena (ed.), Oxford University Press (1999) (ISBN: 0199637768); Nature Genet. 21(1)(suppl):1-60 (1999); Microarray Biochip:

Tools and Technology, Schena (ed.), Eaton Publishing Company/BioTechniques Books Division (2000) (ISBN: 1881299376).

C. Vectors

[00281] Also provided are vectors, including expression vectors, which comprise at least one of the nucleic acid molecules disclosed herein, as described further herein. In some embodiments, the vectors comprise at least one isolated nucleic acid molecule encoding a nutritive protein as disclosed herein. In alternative embodiments, the vectors comprise such a nucleic acid molecule operably linked to one or more expression control sequence. The vectors can thus be used to express at least one recombinant protein in a recombinant microbial host cell.

[00282] Suitable vectors for expression of nucleic acids in microorganisms are well known to those of skill in the art. Suitable vectors for use in cyanobacteria are described, for example, in Heidorn et al., "Synthetic Biology in Cyanobacteria: Engineering and Analyzing Novel Functions," Methods in Enzymology, Vol. 497, Ch. 24 (2011). Exemplary replicative vectors that can be used for engineering cyanobacteria as disclosed herein include pPMQAK1, pSL1211, pFC1, pSB2A, pSCR119/202, pSUN119/202, pRL2697, pRL25C, pRL1050, pSG111M, and pPBH201.

[00283] Other vectors such as pJB161 which are capable of receiving nucleic acid sequences disclosed herein may also be used. Vectors such as pJB161 comprise sequences which are homologous with sequences present in plasmids endogenous to certain photosynthetic microorganisms (e.g., plasmids pAQ1, pAQ3, and pAQ4 of certain *Synechococcus* species). Examples of such vectors and how to use them is known in the art and provided, for example, in Xu et al., "Expression of Genes in Cyanobacteria: Adaptation of Endogenous Plasmids as Platforms for High-Level Gene Expression in *Synechococcus* sp. PCC 7002," Chapter 21 in Robert Carpentier (ed.), "Photosynthesis Research Protocols," Methods in Molecular Biology, Vol. 684, 2011, which is hereby incorporated herein by reference. Recombination between pJB161 and the endogenous plasmids in vivo yield engineered microbes expressing the genes of interest from their endogenous plasmids. Alternatively, vectors can be engineered to recombine with the host cell chromosome, or the vector can be engineered to replicate and express genes of interest independent of the host cell chromosome or any of the host cell's endogenous plasmids.

[00284] A further example of a vector suitable for recombinant protein production is the pET system (Novagen®). This system has been extensively characterized for use in *E. coli* and other microorganisms. In this system, target genes are cloned in pET plasmids under control of strong bacteriophage T7 transcription and (optionally) translation signals; expression is induced by providing a source of T7 RNA polymerase in the host cell. T7 RNA polymerase is so selective and active that, when fully induced, almost all of the microorganism's resources are converted to target gene expression; the desired product can comprise more than 50% of the total cell protein a few hours after induction. It is also possible to attenuate the expression level simply by lowering the concentration of inducer. Decreasing the expression level may enhance the soluble yield of some target proteins. In some embodiments this system also allows for maintenance of target genes in a transcriptionally silent un-induced state.

[00285] In some embodiments of using this system, target genes are cloned using hosts that do not contain the T7 RNA polymerase gene, thus alleviating potential problems related to plasmid instability due to the production of proteins potentially toxic to the host cell. Once established in a non-expression host, target protein expression may be initiated either by infecting the host with λ CE6, a phage that carries the T7 RNA polymerase gene under the control of the λ pL and pI promoters, or by transferring the plasmid into an expression host containing a chromosomal copy of the T7 RNA polymerase gene under lacUV5 control. In the second case, expression is induced by the addition of IPTG or lactose to the bacterial culture or using an autoinduction medium. Other plasmids systems that are controlled by the *lac* operator, but do not require the T7 RNA polymerase gene and rely upon *E. coli*'s native RNA polymerase include the pTrc plasmid suite (Invitrogen) or pQE plasmid suite (QIAGEN).

[00286] In other embodiments it is possible to clone directly into expression hosts. Two types of T7 promoters and several hosts that differ in their stringency of suppressing basal expression levels are available, providing great flexibility and the ability to optimize the expression of a wide variety of target genes.

[00287] Suitable vectors for expression of nucleic acids in mammalian cells typically comprise control functions provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma virus, Adenovirus 2, cytomegalovirus, or Simian Virus 40.

D. Promoters

[00288] Promoters useful for expressing the recombinant genes described herein include both constitutive and inducible/repressible promoters. Examples of inducible/repressible promoters include nickel-inducible promoters (e.g., PnrsA, PnrsB ; see, e.g., Lopez-Mauy et al., Cell (2002) v.43: 247-256) and urea repressible promoters such as PnirA (described in, e.g., Qi et al., Applied and Environmental Microbiology (2005) v.71: 5678-5684). Additional examples of inducible/repressible promoters include PnirA (promoter that drives expression of the nirA gene, induced by nitrate and repressed by urea) and Psuf (promoter that drives expression of the sufB gene, induced by iron stress). Examples of constitutive promoters include Pcpc (promoter that drives expression of the cpc operon), Prbc (promoter that drives expression of rubisco), PpsbAII (promoter that drives expression of PpsbAII), Pcro (lambda phage promoter that drives expression of cro). In other embodiments, a PaphII and/or a lacIq-Ptrc promoter can be used to control expression. Where multiple recombinant genes are expressed in an engineered microorganism, the different genes can be controlled by different promoters or by identical promoters in separate operons, or the expression of two or more genes may be controlled by a single promoter as part of an operon.

[00289] Further non-limiting examples of inducible promoters may include, but are not limited to, those induced by expression of an exogenous protein (e.g., T7 RNA polymerase, SP6 RNA polymerase), by the presence of a small molecule (e.g., IPTG, galactose, tetracycline, steroid hormone, abscisic acid), by absence of small molecules (e.g., CO₂, iron, nitrogen), by metals or metal ions (e.g., copper, zinc, cadmium, nickel), and by environmental factors (e.g., heat, cold, stress, light, darkness), and by growth phase. In some embodiments, the inducible promoter is tightly regulated such that in the absence of induction, substantially no transcription is initiated through the promoter. In some embodiments, induction of the promoter does not substantially alter transcription through other promoters. Also, generally speaking, the compound or condition that induces an inducible promoter is not naturally present in the organism or environment where expression is sought.

[00290] In some embodiments, the inducible promoter is induced by limitation of CO₂ supply to a cyanobacteria culture. By way of non-limiting example, the inducible promoter may be the promoter sequence of *Synechocystis* PCC 6803 that are up-regulated under the CO₂-limitation conditions, such as the *cmp* genes, *ntp* genes, *ndh* genes, *sbt* genes, *chp* genes, and *rbc* genes, or a variant or fragment thereof.

[00291] In some embodiments, the inducible promoter is induced by iron starvation or by entering the stationary growth phase. In some embodiments, the inducible promoter may be variant sequences of the promoter sequence of cyanobacterial genes that are up-regulated under Fe-starvation conditions such as *isiA*, or when the culture enters the stationary growth phase, such as *isiA*, *phrA*, *sigC*, *sigB*, and *sigH* genes, or a variant or fragment thereof.

[00292] In some embodiments, the inducible promoter is induced by a metal or metal ion. By way of non-limiting example, the inducible promoter may be induced by copper, zinc, cadmium, mercury, nickel, gold, silver, cobalt, and bismuth or ions thereof. In some embodiments, the inducible promoter is induced by nickel or a nickel ion. In some embodiments, the inducible promoter is induced by a nickel ion, such as Ni^{2+} . In another exemplary embodiment, the inducible promoter is the nickel inducible promoter from *Synechocystis* PCC 6803. In another embodiment, the inducible promoter may be induced by copper or a copper ion. In yet another embodiment, the inducible promoter may be induced by zinc or a zinc ion. In still another embodiment, the inducible promoter may be induced by cadmium or a cadmium ion. In yet still another embodiment, the inducible promoter may be induced by mercury or a mercury ion. In an alternative embodiment, the inducible promoter may be induced by gold or a gold ion. In another alternative embodiment, the inducible promoter may be induced by silver or a silver ion. In yet another alternative embodiment, the inducible promoter may be induced by cobalt or a cobalt ion. In still another alternative embodiment, the inducible promoter may be induced by bismuth or a bismuth ion.

[00293] In some embodiments, the promoter is induced by exposing a cell comprising the inducible promoter to a metal or metal ion. The cell may be exposed to the metal or metal ion by adding the metal to the microbial growth media. In certain embodiments, the metal or metal ion added to the microbial growth media may be efficiently recovered from the media. In other embodiments, the metal or metal ion remaining in the media after recovery does not substantially impede downstream processing of the media or of the bacterial gene products.

[00294] Further non-limiting examples of constitutive promoters include constitutive promoters from Gram-negative bacteria or a bacteriophage propagating in a Gram-negative bacterium. For instance, promoters for genes encoding highly expressed Gram-negative gene products may be used, such as the promoter for Lpp, OmpA, rRNA, and ribosomal proteins. Alternatively, regulatable promoters may be used in a strain that lacks the regulatory protein for that promoter. For instance P_{lac} , P_{tac} , and P_{trc} , may be used as constitutive promoters in

strains that lack LacI . Similarly, P_{22} , P_R and P_L may be used in strains that lack the lambda C2 repressor protein, and lambda P_R and P_L may be used in strains that lack the lambda C1 repressor protein. In one embodiment, the constitutive promoter is from a bacteriophage. In another embodiment, the constitutive promoter is from a *Salmonella* bacteriophage. In yet another embodiment, the constitutive promoter is from a cyanophage. In some embodiments, the constitutive promoter is a *Synechocystis* promoter. For instance, the constitutive promoter may be the P_{psbA} promoter or its variant sequences, the P_{rbc} promoter or its variant sequences, the P_{cpc} promoter or its variant sequences, and the P_{rnpB} promoter or its variant sequences.

E. Host Cells

[00295] Also provided are host cells transformed with the nucleic acid molecules or vectors disclosed herein, and descendants thereof. In some embodiments the host cells are microbial cells. In some embodiments, the host cells carry the nucleic acid sequences on vectors, which may but need not be freely replicating vectors. In other embodiments, the nucleic acids have been integrated into the genome of the host cells and/or into an endogenous plasmid of the host cells. The transformed host cells find use, e.g., in the production of recombinant nutritive proteins disclosed herein.

[00296] “Microorganisms” includes prokaryotic and eukaryotic microbial species from the Domains *Archaea*, *Bacteria* and *Eucarya*, the latter including yeast and filamentous fungi, protozoa, algae, or higher Protista. The terms “microbial cells” and “microbes” are used interchangeably with the term microorganism.

[00297] A variety of host microorganisms may be transformed with a nucleic acid sequence disclosed herein and may in some embodiments be used to produce a recombinant nutritive protein disclosed herein. Suitable host microorganisms include both autotrophic and heterotrophic microbes. In some applications the autotrophic microorganisms allows for a reduction in the fossil fuel and/or electricity inputs required to make a nutritive protein encoded by a recombinant nucleic acid sequence introduced into the host microorganism. This, in turn, in some applications reduces the cost and/or the environmental impact of producing the nutritive protein and/or reduces the cost and/or the environmental impact in comparison to the cost and/or environmental impact of manufacturing alternative nutritive proteins, such as whey, egg, and soy. For example, the cost and/or environmental impact of making a nutritive protein disclosed herein using a host microorganism as disclosed herein is

in some embodiments lower than the cost and/or environmental impact of making whey protein in a form suitable for human consumption by processing of cows milk.

[00298] Non-limiting examples of heterotrophs include *Escherichia coli*, *Salmonella typhimurium*, *Bacillus subtilis*, *Bacillus megaterium*, *Corynebacterium glutamicum*, *Streptomyces coelicolor*, *Streptomyces lividans*, *Streptomyces venezuelae*, *Streptomyces roseosporus*, *Streptomyces fradiae*, *Streptomyces griseus*, *Streptomyces calvuligerus*, *Streptomyces hygroscopicus*, *Streptomyces platensis*, *Saccharopolyspora erythraea*, *Corynebacterium glutamicum*, *Aspergillus niger*, *Aspergillus nidulans*, *Aspergillus oryzae*, *Aspergillus terreus*, *Aspergillus sojae*, *Penicillium chrysogenum*, *Trichoderma reesei*, *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *Clostridium thermocellum*, *Fusibacter paucivorans*, *Saccharomyces cerevisiae*, *Saccharomyces boulardii*, *Pichia pastoris*, and *Pichia stipitis*.

[00299] Photoautotrophic microorganisms include eukaryotic algae, as well as prokaryotic cyanobacteria, green-sulfur bacteria, green non-sulfur bacteria, purple sulfur bacteria, and purple non-sulfur bacteria.

[00300] Extremophiles are also contemplated as suitable organisms. Such organisms withstand various environmental parameters such as temperature, radiation, pressure, gravity, vacuum, desiccation, salinity, pH, oxygen tension, and chemicals. They include hyperthermophiles, which grow at or above 80°C such as *Pyrolobus fumarii*; thermophiles, which grow between 60-80°C such as *Synechococcus lividis*; mesophiles, which grow between 15-60°C; and psychrophiles, which grow at or below 15°C such as *Psychrobacter* and some insects. Radiation tolerant organisms include *Deinococcus radiodurans*. Pressure-tolerant organisms include piezophiles, which tolerate pressure of 130 MPa. Weight-tolerant organisms include barophiles. Hypergravity (*e.g.*, >1g) hypogravity (*e.g.*, <1g) tolerant organisms are also contemplated. Vacuum tolerant organisms include tardigrades, insects, microbes and seeds. Dessicant tolerant and anhydrobiotic organisms include xerophiles such as *Artemia salina*; nematodes, microbes, fungi and lichens. Salt-tolerant organisms include halophiles (*e.g.*, 2-5 M NaCl) *Halobacteriaceae* and *Dunaliella salina*. pH-tolerant organisms include alkaliphiles such as *Natronobacterium*, *Bacillus firmus* OF4, *Spirulina spp.* (*e.g.*, pH > 9) and acidophiles such as *Cyanidium caldarium*, *Ferroplasma sp.* (*e.g.*, low pH). Anaerobes, which cannot tolerate O₂ such as *Methanococcus jannaschii*; microaerophils, which tolerate some O₂ such as *Clostridium* and aerobes, which require O₂ are also

contemplated. Gas-tolerant organisms, which tolerate pure CO₂ include *Cyanidium caldarium* and metal tolerant organisms include metalotolerants such as *Ferroplasma acidarmanus* (e.g., Cu, As, Cd, Zn), *Ralstonia* sp. CH34 (e.g., Zn, Co, Cd, Hg, Pb). Gross, Michael. *Life on the Edge: Amazing Creatures Thriving in Extreme Environments*. New York: Plenum (1998) and Seckbach, J. "Search for Life in the Universe with Terrestrial Microbes Which Thrive Under Extreme Conditions." In Cristiano Batalli Cosmovici, Stuart Bowyer, and Dan Wertheimer, eds., *Astronomical and Biochemical Origins and the Search for Life in the Universe*, p. 511. Milan: Editrice Compositori (1997).

[00301] Mixotrophic organisms are also suitable organisms. Mixotrophic organisms can utilize a mix of different sources of energy and carbon, for example, photo- and chemotrophy, litho- and organotrophy, auto- and heterotrophy, and combinations thereof or a combination of it. Mixotrophs can be either eukaryotic or prokaryotic. Additionally, mixotrophs can be obligate or facultative. Suitable mixotrophic organisms include mixotrophic algae and mixotrophic bacteria.

[00302] Algae and cyanobacteria include but are not limited to the following genera: Acanthoceras, Acanthococcus, Acaryochloris, Achnanthes, Achnanthidium, Actinastrum, Actinochloris, Actinocyclus, Actinotaenium, Amphichrysis, Amphidinium, Amphikrikos, Amphipleura, Amphiprora, Amphithrix, Amphora, Anabaena, Anabaenopsis, Aneumastus, Ankistrodesmus, Ankyra, Anomoeoneis, Apatococcus, Aphanizomenon, Aphanocapsa, Aphanochaete, Aphanothece, Apiocystis, Apistonema, Arthrodesmus, Artherospira, Ascochloris, Asterionella, Asterococcus, Audouinella, Aulacoseira, Bacillaria, Balbiania, Bambusina, Bangia, Basichlamys, Batrachospermum, Binuclearia, Bitrichia, Blidingia, Botriopsis, Botrydium, Botryococcus, Botryosphaerella, Brachiomonas, Brachysira, Brachytrichia, Brebissonia, Bulbochaete, Bumilleria, Bumilleriopsis, Caloneis, Calothrix, Campylodiscus, Capsosiphon, Carteria, Catena, Cavinula, Centritractus, Centronella, Ceratium, Chaetoceros, Chaetochloris, Chaetomorpha, Chaetonella, Chaetonema, Chaetopeltis, Chaetophora, Chaetosphaeridium, Chamaesiphon, Chara, Characiochloris, Characiopsis, Characium, Charales, Chilomonas, Chlainomonas, Chlamydoublepharis, Chlamydopapsa, Chlamydomonas, Chlamydomonopsis, Chlamydomyxa, Chlamydonephris, Chlorangiella, Chlorangiopsis, Chlorella, Chlorobotrys, Chlorobrachis, Chlorochytrium, Chlorococcum, Chlorogloea, Chlorogloeopsis, Chlorogonium, Chlorolobion, Chloromonas, Chlorophysema, Chlorophyta, Chlorosaccus, Chlorosarcina, Choricystis, Chromophyton, Chromulina, Chroococcidiopsis, Chroococcus, Chroodactylon, Chroomonas, Chroothece,

Chrysamoeba, Chrysapsis, Chrysidiastrium, Chrysocapsa, Chrysocapsella, Chrysochaete,
 Chrysochromulina, Chrysococcus, Chrysocrinus, Chrysolepidomonas, Chrysolykos,
 Chrysonebula, Chrysophyta, Chrysopyxis, Chrysosaccus, Chrysophaerella,
 Chrysostephanosphaera, Clodophora, Clastidium, Closteriopsis, Closterium, Coccomyxa,
 Cocconeis, Coelastrella, Coelastrum, Coelosphaerium, Coenochloris, Coenococcus,
 Coenocystis, Colacium, Coleochaete, Collodictyon, Compsogonopsis, Compsopogon,
 Conjugatophyta, Conochaete, Coronastrum, Cosmarium, Cosmioneis, Cosmocladium,
 Crateriportula, Craticula, Crinalium, Crucigenia, Crucigeniella, Cryptoaulax, Cryptomonas,
 Cryptophyta, Ctenophora, Cyanodictyon, Cyanonephron, Cyanophora, Cyanophyta,
 Cyanothece, Cyanothomonas, Cyclonexis, Cyclostephanos, Cyclotella, Cylindrocapsa,
 Cylindrocystis, Cylindrospermum, Cylindrotheca, Cymatopleura, Cymbella,
 Cymbellonitzschia, Cystodinium Dactylococcopsis, Debarya, Denticula, Dermatochrysis,
 Dermocarpa, Dermocarpella, Desmatractum, Desmidium, Desmococcus, Desmonema,
 Desmosiphon, Diacanthos, Diacronema, Diadesmis, Diatoma, Diatomella, Dicellula,
 Dichothrix, Dichotomococcus, Dicranochaete, Dictyochloris, Dictyococcus,
 Dictyosphaerium, Didymocystis, Didymogenes, Didymosphenia, Dilabifilum,
 Dimorphococcus, Dinobryon, Dinococcus, Diplochloris, Diploneis, Diplostauron,
 Distrionella, Docidium, Draparnaldia, Dunaliella, Dysmorphococcus, Ecballocystis,
 Elakatothrix, Ellerbeckia, Encyonema, Enteromorpha, Entocladia, Entomoneis, Entophysalis,
 Epichrysis, Epipyxis, Epithemia, Eremosphaera, Euastropsis, Euastrum, Eucapsis,
 Eucocconeis, Eudorina, Euglena, Euglenophyta, Eunotia, Eustigmatophyta, Eutreptia,
 Fallacia, Fischerella, Fragilaria, Fragilariforma, Franceia, Frustulia, Curcilla, Geminella,
 Genicularia, Glaucocystis, Glaucophyta, Glenodiniopsis, Glenodinium, Gloecapsa,
 Gloeochaete, Gloeochrysis, Gloeococcus, Gloeocystis, Gloeodendron, Gloeomonas,
 Gloeoplax, Gloeotheca, Gloeotila, Gloeotrichia, Gloiodictyon, Golenkinia, Golenkiniopsis,
 Gomontia, Gomphocymbella, Gomphonema, Gomphosphaeria, Gonatozygon, Gongrosia,
 Gongrosira, Goniochloris, Gonium, Gonyostomum, Granulochloris, Granulocystopsis,
 Groenbladia, Gymnodinium, Gymnozyga, Gyrosigma, Haematococcus, Hafniomonas,
 Hallassia, Hammatoidea, Hannaea, Hantzschia, Hapalosiphon, Haplotaenium, Haptophyta,
 Haslea, Hemidinium, Hemitoma, Heribaudiella, Heteromastix, Heterothrix, Hibberdia,
 Hildenbrandia, Hillea, Holopedium, Homoeothrix, Hormanthonema, Hormotila,
 Hyalobranchion, Hyalocardium, Hyalodiscus, Hyalogonium, Hyalotheca, Hydrianum,
 Hydrococcus, Hydrocoleum, Hydrocoryne, Hydrodictyon, Hydrosera, Hydrurus, Hyella,

Hymenomonas, Isthmochloron, Johannesbaptistia, Juranyiella, Karayevia, Kathablepharis, Katodinium, Kephyrion, Keratococcus, Kirchneriella, Klebsormidium, Kolbesia, Koliella, Komarekia, Korshikoviella, Kraskella, Lagerheimia, Lagynion, Lamprothamnium, Lemanea, Lepocinclis, Leptosira, Lobococcus, Lobocystis, Lobomonas, Luticola, Lyngbya, Malleochloris, Mallomonas, Mantoniella, Marssoniella, Martyana, Mastigocoleus, Gastogloia, Melosira, Merismopedia, Mesostigma, Mesotaenium, Micractinium, Micrasterias, Microchaete, Microcoleus, Microcystis, Microglana, Micromonas, Microspora, Microthamnion, Mischococcus, Monochrysis, Monodus, Monomastix, Monoraphidium, Monostroma, Mougeotia, Mougeotiopsis, Myochloris, Myromecia, Myxosarcina, Naegeliella, Nannochloris, Nautococcus, Navicula, Neglectella, Neidium, Nephroclamys, Nephrocystium, Nephrodiella, Nephroselmis, Netrium, Nitella, Nitellopsis, Nitzschia, Nodularia, Nostoc, Ochromonas, Oedogonium, Oligochaetophora, Onychonema, Oocardium, Oocystis, Opephora, Ophiocystium, Orthoseira, Oscillatoria, Oxyneis, Pachycladella, Palmella, Palmodictyon, Pnadorina, Pannus, Paralia, Pascherina, Paulschulzia, Pediatrum, Pedinella, Pedinomonas, Pedinopera, Pelagodictyon, Penium, Peranema, Peridiniopsis, Peridinium, Peronia, Petroneis, Phacotus, Phacus, Phaeaster, Phaeodermatium, Phaeophyta, Phaeosphaera, Phaeothamnion, Phormidium, Phycopeltis, Phyllariochloris, Phyllocardium, Phyllomitas, Pinnularia, Pitophora, Placoneis, Planctonema, Planktosphaeria, Planothidium, Plectonema, Pleodorina, Pleurastrum, Pleurocapsa, Pleurocladia, Pleurodiscus, Pleurosigma, Pleurosira, Pleurotaenium, Pocillomonas, Podohedra, Polyblepharides, Polychaetophora, Polyedriella, Polyedriopsis, Polygoniochloris, Polyepidomonas, Polytaenia, Polytoma, Polytomella, Porphyridium, Posteriochromonas, Prasiniochloris, Prasinocladus, Prasinophyta, Prasiola, Prochlorophyta, Prochlorothrix, Protoderma, Protosiphon, Provasoliella, Prymnesium, Psammodictyon, Psammothidium, Pseudanabaena, Pseudenoclonium, Psuedocarteria, Pseudochate, Pseudocharacium, Pseudococcomyxa, Pseudodictyosphaerium, Pseudokephyrion, Pseudoncobyrsa, Pseudoquadrigula, Pseudosphaerocystis, Pseudostaurastrum, Pseudostaurosira, Pseudotetrastrum, Pteromonas, Punctastruata, Pyramichlamys, Pyramimonas, Pyrrophyta, Quadrichloris, Quadricoccus, Quadrigula, Radiococcus, Radiofilum, Raphidiopsis, Raphidocelis, Raphidonema, Raphidophyta, Peimeria, Rhabdoderma, Rhabdomonas, Rhizoclonium, Rhodomonas, Rhodophyta, Rhoicosphenia, Rhopalodia, Rivularia, Rosenvingiella, Rossithidium, Roya, Scenedesmus, Scherffelia, Schizochlamydella, Schizochlamys, Schizomeris, Schizothrix, Schroederia, Scolioneis, Scotiella, Scotiellopsis, Scourfieldia, Scytonema, Selenastrum, Selenochloris,

Sellaphora, Semiorbis, Siderocelis, Diderocystopsis, Dimonsenia, Siphononema, Sirocladium, Sirogonium, Skeletonema, Sorastrum, Spennatozopsis, Sphaerellocystis, Sphaerellopsis, Sphaerodinium, Sphaeroplea, Sphaerososma, Spiniferomonas, Spirogyra, Spirotaenia, Spirulina, Spondylomorum, Spondylosium, Sporotetras, Spumella, Staurastrum, Stauerodesmus, Stauroneis, Staurosira, Staurosirella, Stenopterobia, Stephanocostis, Stephanodiscus, Stephanoporus, Stephanosphaera, Stichococcus, Stichogloea, Stigeoclonium, Stigonema, Stipitococcus, Stokesiella, Strombomonas, Stylochrysalis, Stylodinium, Styloxyis, Stylosphaeridium, Surirella, Sykidion, Symploca, Synechococcus, Synechocystis, Synedra, Synochromonas, Synura, Tabellaria, Tabularia, Teilingia, Temnogametum, Tetmemorus, Tetrachlorella, Tetracyclus, Tetradesmus, Tetraedriella, Tetraedron, Tetraselmis, Tetraspora, Tetrastrium, Thalassiosira, Thamniochaete, Thorakochloris, Thorea, Tolypella, Tolypothrix, Trachelomonas, Trachydiscus, Trebouxia, Trentepohlia, Treubaria, Tribonema, Trichodesmium, Trichodiscus, Trochiscia, Tryblionella, Ulothrix, Uroglena, Uronema, Urosolenia, Urospora, Uva, Vacuolaria, Vaucheria, Volvox, Volvulina, Westella, Woloszynskia, Xanthidium, Xanthophyta, Xenococcus, Zygnema, Zygnemopsis, and Zygonium.

[00303] Additional cyanobacteria include members of the genus Chamaesiphon, Chroococcus, Cyanobacterium, Cyanobium, Cyanothece, Dactylococcopsis, Gloeobacter, Gloeocapsa, Gloeotheca, Microcystis, Prochlorococcus, Prochloron, Synechococcus, Synechocystis, Cyanocystis, Dermocarpella, Stanieria, Xenococcus, Chroococcidiopsis, Myxosarcina, Arthrospira, Borzia, Crinalium, Geitlerinemia, Leptolyngbya, Limnothrix, Lyngbya, Microcoleus, Oscillatoria, Planktothrix, Prochlorothrix, Pseudanabaena, Spirulina, Starria, Symploca, Trichodesmium, Tychonema, Anabaena, Anabaenopsis, Aphanizomenon, Cyanospira, Cylandrospermopsis, Cylandrospermum, Nodularia, Nostoc, Scydonema, Calothrix, Rivularia, Tolypothrix, Chlorogloeopsis, Fischerella, Geitleria, Iyengariella, Nostochopsis, Stigonema and Thermosynechococcus.

[00304] Green non-sulfur bacteria include but are not limited to the following genera: Chloroflexus, Chloronema, Oscillochloris, Heliothrix, Herpetosiphon, Roseiflexus, and Thermomicrobium.

[00305] Green sulfur bacteria include but are not limited to the following genera: *Chlorobium*, *Clathrochloris*, and *Prosthecochloris*.

[00306] Purple sulfur bacteria include but are not limited to the following genera: Allochromatium, Chromatium, Halochromatium, Isochromatium, Marichromatium, Rhodovulum, Thermochromatium, Thiocapsa, Thiorhodococcus, and Thiocystis.

[00307] Purple non-sulfur bacteria include but are not limited to the following genera: Phaeospirillum, Rhodobaca, Rhodobacter, Rhodomicrobium, Rhodopila, Rhodopseudomonas, Rhodothalassium, Rhodospirillum, Rhodovibrio, and Roseospira.

[00308] Aerobic chemolithotrophic bacteria include but are not limited to nitrifying bacteria such as *Nitrobacteraceae* sp., *Nitrobacter* sp., *Nitrospina* sp., *Nitrococcus* sp., *Nitrospira* sp., *Nitrosomonas* sp., *Nitrosococcus* sp., *Nitrosospira* sp., *Nitrosolobus* sp., *Nitrosovibrio* sp.; colorless sulfur bacteria such as, *Thiovulum* sp., *Thiobacillus* sp., *Thiomicrospira* sp., *Thiosphaera* sp., *Thermothrix* sp.; obligately chemolithotrophic hydrogen bacteria such as *Hydrogenobacter* sp., iron and manganese-oxidizing and/or depositing bacteria such as *Siderococcus* sp., and magnetotactic bacteria such as *Aquaspirillum* sp.

[00309] Archaeobacteria include but are not limited to methanogenic archaeobacteria such as *Methanobacterium* sp., *Methanobrevibacter* sp., *Methanothermus* sp., *Methanococcus* sp., *Methanomicrobium* sp., *Methanospirillum* sp., *Methanogenium* sp., *Methanosarcina* sp., *Methanlobus* sp., *Methanothrix* sp., *Methanococcoides* sp., *Methanoplanus* sp.; extremely thermophilic S-Metabolizers such as *Thermoproteus* sp., *Pyrodictium* sp., *Sulfolobus* sp., *Acidianus* sp. and other microorganisms such as, *Bacillus subtilis*, *Saccharomyces cerevisiae*, *Streptomyces* sp., *Ralstonia* sp., *Rhodococcus* sp., *Corynebacteria* sp., *Brevibacteria* sp., *Mycobacteria* sp., and oleaginous yeast.

[00310] Yet other suitable organisms include synthetic cells or cells produced by synthetic genomes as described in Venter et al. US Pat. Pub. No. 2007/0264688, and cell-like systems or synthetic cells as described in Glass et al. US Pat. Pub. No. 2007/0269862.

[00311] Still other suitable organisms include *Escherichia coli*, *Acetobacter aceti*, *Bacillus subtilis*, yeast and fungi such as *Clostridium ljungdahlii*, *Clostridium thermocellum*, *Penicillium chrysogenum*, *Pichia pastoris*, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pseudomonas fluorescens*, or *Zymomonas mobilis*. In some embodiments those organisms are engineered to fix carbon dioxide while in other embodiments they are not.

[00312] In some embodiments eukaryotic cells, such as insect cells or mammalian cells, such as human cells are used as host cells. Vectors and expression control sequences including promoters and enhancers are well known for such cells. Examples of useful

mammalian host cell lines for this purpose are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

F. Production of Nutritive Proteins

[00313] Skilled artisans are aware of many suitable methods available for culturing recombinant cells to produce (and optionally secrete) a recombinant nutritive protein as disclosed herein, as well as for purification and/or isolation of expressed recombinant proteins. The methods chosen for protein purification depend on many variables, including the properties of the protein of interest, its location and form within the cell, the vector, host strain background, and the intended application for the expressed protein. Culture conditions can also have an effect on solubility and localization of a given target protein. Many approaches can be used to purify target proteins expressed in recombinant microbial cells as disclosed herein, including without limitation ion exchange and gel filtration.

[00314] In some embodiments a peptide fusion tag is added to the recombinant protein making possible a variety of affinity purification methods that take advantage of the peptide fusion tag. In some embodiments, the use of an affinity method enables the purification of the target protein to near homogeneity in one step. Purification may include cleavage of part or all of the fusion tag with enterokinase, factor Xa, thrombin, or HRV 3C proteases, for example. In some embodiments, before purification or activity measurements of an expressed target protein, preliminary analysis of expression levels, cellular localization, and solubility of the target protein is performed. The target protein may be found in any or all of the following fractions: soluble or insoluble cytoplasmic fractions, periplasm, or medium. Depending on the intended application, preferential localization to inclusion bodies, medium,

or the periplasmic space can be advantageous, in some embodiments, for rapid purification by relatively simple procedures.

[00315] While *Escherichia coli* is widely regarded as a robust host for heterologous protein expression, it is also widely known that over-expression of many proteins in this host is prone to aggregation in the form of insoluble inclusion bodies. One of the most commonly used methods for either rescuing inclusion body formation, or to improve the titer of the protein itself, is to include an amino-terminal maltose-binding protein (MBP) [Austin BP, Nallamsetty S, Waugh DS. Hexahistidine-tagged maltose-binding protein as a fusion partner for the production of soluble recombinant proteins in *Escherichia coli*. *Methods Mol Biol*. 2009;498:157-72], or small ubiquitin-related modifier (SUMO) [Saitoh H, Uwada J, Azusa K. Strategies for the expression of SUMO-modified target proteins in *Escherichia coli*. *Methods Mol Biol*. 2009;497:211-21; Malakhov MP, Mattern MR, Malakhova OA, Drinker M, Weeks SD, Butt TR. SUMO fusions and SUMO-specific protease for efficient expression and purification of proteins. *J Struct Funct Genomics*. 2004;5(1-2):75-86; Panavas T, Sanders C, Butt TR. SUMO fusion technology for enhanced protein production in prokaryotic and eukaryotic expression systems. *Methods Mol Biol*. 2009;497:303-17] fusion to the protein of interest. These two proteins are expressed extremely well, and in the soluble form, in *Escherichia coli* such that the protein of interest is also effectively produced in the soluble form. The protein of interest can be cleaved by designing a site specific protease recognition sequence (such as the tobacco etch virus (TEV) protease) in-between the protein of interest and the fusion protein [1].

[00316] In some embodiments the recombinant protein is initially not folded correctly or is insoluble. A variety of methods are well known for refolding of insoluble proteins. Most protocols comprise the isolation of insoluble inclusion bodies by centrifugation followed by solubilization under denaturing conditions. The protein is then dialyzed or diluted into a non-denaturing buffer where refolding occurs. Because every protein possesses unique folding properties, the optimal refolding protocol for any given protein can be empirically determined by a skilled artisan. Optimal refolding conditions can, for example, be rapidly determined on a small scale by a matrix approach, in which variables such as protein concentration, reducing agent, redox treatment, divalent cations, etc., are tested. Once the optimal concentrations are found, they can be applied to a larger scale solubilization and refolding of the target protein.

[00317] In some embodiments the nutritive protein does not comprise a tertiary structure. In some embodiments less than half of the amino acids in the nutritive protein participate in a tertiary structure. In some embodiments the nutritive protein does not comprise a secondary structure. In some embodiments less than half of the amino acids in the nutritive protein participate in a secondary structure. Recombinant nutritive proteins may be isolated from a culture of cells expressing them in a state that comprises one or more of these structural features. In some embodiments the tertiary structure of a recombinant nutritive protein is reduced or eliminated after the protein is isolated from a culture producing it. In some embodiments the secondary structure of a recombinant nutritive protein is reduced or eliminated after the protein is isolated from a culture producing it.

[00318] In some embodiments a CAPS buffer at alkaline pH in combination with N-lauroylsarcosine is used to achieve solubility of the inclusion bodies, followed by dialysis in the presence of DTT to promote refolding. Depending on the target protein, expression conditions, and intended application, proteins solubilized from washed inclusion bodies may be > 90% homogeneous and may not require further purification. Purification under fully denaturing conditions (before refolding) is possible using His•Tag® fusion proteins and His•Bind® immobilized metal affinity chromatography (Novogen®). In addition, S•Tag™, T7•Tag®, and Strep•Tag® II fusion proteins solubilized from inclusion bodies using 6 M urea can be purified under partially denaturing conditions by dilution to 2 M urea (S•Tag and T7•Tag) or 1 M urea (Strep•Tag II) prior to chromatography on the appropriate resin. Refolded fusion proteins can be affinity purified under native conditions using His•Tag, S•Tag, Strep•Tag II, and other appropriate affinity tags (e.g., GST•Tag™, and T7•Tag) (Novogen®).

[00319] In some embodiments the recombinant nutritive protein is an endogenous protein of the host cell used to express it. That is, the cellular genome of the host cell comprises an open reading frame that encodes the recombinant nutritive protein. In some embodiments regulatory sequences sufficient to increase expression of the nutritive protein are inserted into the host cell genome and operatively linked to the endogenous open reading frame such that the regulatory sequences drive overexpression of the recombinant nutritive protein from a recombinant nucleic acid. In some embodiments heterologous nucleic acid sequences are fused to the endogenous open reading frame of the nutritive protein and cause the nutritive protein to be synthesized comprising a heterologous amino acid sequence that changes the cellular trafficking of the recombinant nutritive protein, such as directing it to an

organelle or to a secretion pathway. In some embodiments an open reading frame that encodes the endogenous host cell protein is introduced into the host cell on a plasmid that further comprises regulatory sequences operatively linked to the open reading frame. In some embodiments the recombinant host cell expresses at least 2 times, at least 3 times, at least 4 times, at least 5 times, at least 10 times, or at least 20 times, at least 30 times, at least 40 times, at least 50 times, or at least 100 times more of the recombinant nutritive protein than the amount of the nutritive protein produced by a similar host cell grown under similar conditions.

[00320] In some embodiments nutritive proteins of this disclosure are synthesized chemically without the use of a recombinant production system. Protein synthesis can be carried out in a liquid-phase system or in a solid-phase system using techniques known in the art (see, e.g., Atherton, E., Sheppard, R.C. (1989). *Solid Phase peptide synthesis: a practical approach*. Oxford, England: IRL Press; Stewart, J.M., Young, J.D. (1984). *Solid phase peptide synthesis* (2nd ed.). Rockford: Pierce Chemical Company).

G. Production of Recombinant Nutritive Proteins in Plants

[00321] The nucleic acid molecules comprising a nucleic acid sequence encoding a nutritive protein of this disclosure enable production of transgenic plants comprising the nucleic acid sequence. Accordingly, this disclosure also provides plant comprising a recombinant nucleic acid molecule comprising a nucleic acid sequence encoding a nutritive protein of this disclosure. The plant can be any plant that is subject to transformation and regeneration and include, but are not limited to, Acacia, alfalfa, aneth, apple, apricot, artichoke, arugula, asparagus, avocado, banana, barley, beans, beet, blackberry, blueberry, broccoli, brussels sprouts, cabbage, canola, cantaloupe, carrot, cassaya, cauliflower, celery, Chinese cabbage, cherry, cilantro, citrus, clementines, coffee, corn, cotton, cucumber, Douglas fir, eggplant, endive, escarole, eucalyptus, fennel, figs, forest trees, gourd, grape, grapefruit, honey dew, jicama, kiwifruit, lettuce, leeks, lemon, lime, Loblolly pine, mango, melon, mushroom, nut, oat, okra, onion, orange, an ornamental plant, papaya, parsley, pea, peach, peanut, pear, pepper, persimmon, pine, pineapple, plantain, plum, pomegranate, poplar, potato, pumpkin, quince, radiata pine, radicchio, radish, rapeseed, raspberry, rice, rye, sorghum, Southern pine, soybean, spinach, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet corn, sweet potato, sweetgum, tangerine, tea, tobacco, tomato, turf, a vine, watermelon, wheat, yams, and zucchini. In preferred embodiments, the plant is a bean, broccoli, cabbage, canola, carrot, cauliflower, celery, Chinese cabbage, corn, cotton

cucumber, eggplant, leek, lettuce, melon, pea, pepper, pumpkin, radish, spinach, soybean, squash, sugarcane, sweet corn, tomato, watermelon, and wheat plant. In some embodiments, the plant is a corn plant. In some embodiments, the plant is a soybean plant. In some embodiments, the plant is a cotton plant. In some embodiments, the plant is a canola plant. In some embodiments the plant is a member of a genus selected from *Arabidopsis*, *Beta*, *Glycine*, *Jatropha*, *Miscanthus*, *Panicum*, *Phalaris*, *Populus*, *Saccharum*, *Salix*, *Simmondsia* and *Zea*.

[00322] Numerous promoters that are active in plant cells have been described in the literature. These include promoters present in plant genomes as well as promoters from other sources, including nopaline synthase (NOS) promoter and octopine synthase (OCS) promoters carried on tumor-inducing plasmids of *Agrobacterium tumefaciens*, caulimovirus promoters such as the cauliflower mosaic virus. For instance, see U.S. Pat. Nos. 5,858,742 and 5,322,938, which disclose versions of the constitutive promoter derived from cauliflower mosaic virus (CaMV35S), U.S. Pat. No. 5,641,876, which discloses a rice actin promoter, U.S. Patent Application Publication 2002/0192813A1, which discloses 5', 3' and intron elements useful in the design of effective plant expression vectors, U.S. patent application Ser. No. 09/757,089, which discloses a maize chloroplast aldolase promoter, U.S. patent application Ser. No. 08/706,946, which discloses a rice glutelin promoter, U.S. patent application Ser. No. 09/757,089, which discloses a maize aldolase (FDA) promoter, and U.S. patent application Ser. No. 60/310,370, which discloses a maize nicotianamine synthase promoter. These and numerous other promoters that function in plant cells are known to those skilled in the art and available for use in recombinant nucleic acids to provide for expression of nutritive proteins in transgenic plants.

[00323] For some applications preferential expression in plant green tissues is desired. Promoters of interest for such uses include those from genes such as *Arabidopsis thaliana* ribulose-1,5-bisphosphate carboxylase (Rubisco) small subunit (Fischhoff et al. (1992) *Plant Mol. Biol.* 20:81-93), aldolase and pyruvate orthophosphate dikinase (PPDK) (Taniguchi et al. (2000) *Plant Cell Physiol.* 41(1):42-48).

[0036] Furthermore, the promoters may be altered to contain at least one enhancer sequence to assist in elevating gene expression. Such enhancers are known in the art. By including an enhancer sequence with such constructs, the expression of the nutritive protein may be enhanced. These enhancers often are found 5' to the start of transcription in a promoter that functions in eukaryotic cells, but can often be inserted upstream (5') or downstream (3') to

the coding sequence. In some instances, these 5' enhancing elements are introns. Particularly useful as enhancers are the 5' introns of the rice actin 1 (see U.S. Pat. No. 5,641,876) and rice actin 2 genes, the maize alcohol dehydrogenase gene intron, the maize heat shock protein 70 gene intron (U.S. Pat. No. 5,593,874) and the maize shrunken 1 gene.

[00324] For some applications expression in plant seed tissues is desired to effect modify seed composition. Exemplary promoters for use for seed composition modification include promoters from seed genes such as napin (U.S. Pat. No. 5,420,034), maize L3 oleosin (U.S. Pat. No. 6,433,252), zein Z27 (Russell et al. (1997) Transgenic Res. 6(2):157-166), globulin 1 (Belanger et al (1991) Genetics 129:863-872), glutelin 1 (Russell (1997) supra), and peroxiredoxin antioxidant (Per1) (Stacy et al. (1996) Plant Mol. Biol. 31(6):1205-1216)

[00325] Recombinant nucleic acid constructs prepared in accordance with the disclosure will also generally include a 3' element that typically contains a polyadenylation signal and site. Well-known 3' elements include those from *Agrobacterium tumefaciens* genes such as nos 3', tml 3', tnr 3', tms 3', ocs 3', tr7 3', for example disclosed in U.S. Pat. No. 6,090,627; 3' elements from plant genes such as wheat (*Triticum aestivum*) heat shock protein 17 (Hsp17 3'), a wheat ubiquitin gene, a wheat fructose-1,6-biphosphatase gene, a rice glutelin gene a rice lactate dehydrogenase gene and a rice beta-tubulin gene, all of which are disclosed in U.S. published patent application 2002/0192813 A1; and the pea (*Pisum sativum*) ribulose biphosphate carboxylase gene (rbs 3'), and 3' elements from the genes within the host plant.

[00326] Constructs and vectors may also include a transit peptide for targeting of a gene target to a plant organelle, particularly to a chloroplast, leucoplast or other plastid organelle. For descriptions of the use of chloroplast transit peptides see U.S. Pat. No. 5,188,642 and U.S. Pat. No. 5,728,925. For description of the transit peptide region of an *Arabidopsis* EPSPS gene, see Klee, H. J. et al (MGG (1987) 210:437-442).

[00327] Numerous methods for transforming plant cells with recombinant DNA are known in the art and may be used in the present disclosure. Two commonly used methods for plant transformation are *Agrobacterium*-mediated transformation and microprojectile bombardment. Microprojectile bombardment methods are illustrated in U.S. Pat. Nos. 5,015,580 (soybean); 5,550,318 (corn); 5,538,880 (corn); 5,914,451 (soybean); 6,160,208 (corn); 6,399,861 (corn) and 6,153,812 (wheat) and *Agrobacterium*-mediated transformation is described in U.S. Pat. Nos. 5,159,135 (cotton); 5,824,877 (soybean); 5,591,616 (corn); and

6,384,301 (soybean). For *Agrobacterium tumefaciens* based plant transformation system, additional elements present on transformation constructs will include T-DNA left and right border sequences to facilitate incorporation of the recombinant polynucleotide into the plant genome.

[00328] In general it is useful to introduce recombinant DNA randomly, i.e. at a non-specific location, in the genome of a target plant line. In special cases it may be useful to target recombinant DNA insertion in order to achieve site-specific integration, for example to replace an existing gene in the genome, to use an existing promoter in the plant genome, or to insert a recombinant polynucleotide at a predetermined site known to be active for gene expression. Several site specific recombination systems exist which are known to function in plants, including cre-lox as disclosed in U.S. Pat. No. 4,959,317 and FLP-FRT as disclosed in U.S. Pat. No. 5,527,695.

[00329] Transformation methods are generally practiced in tissue culture on media and in a controlled environment. "Media" refers to the numerous nutrient mixtures that are used to grow cells in vitro, that is, outside of the intact living organism. Recipient cell targets include, but are not limited to, meristem cells, callus, immature embryos and gametic cells such as microspores, pollen, sperm and egg cells. It is contemplated that any cell from which a fertile plant may be regenerated is useful as a recipient cell. Callus may be initiated from tissue sources including, but not limited to, immature embryos, seedling apical meristems, microspores and the like. Cells capable of proliferating as callus are also recipient cells for genetic transformation. Practical transformation methods and materials for making transgenic plants of this disclosure, for example various media and recipient target cells, transformation of immature embryo cells and subsequent regeneration of fertile transgenic plants are disclosed in U.S. Pat. Nos. 6,194,636 and 6,232,526.

[00330] The seeds of transgenic plants can be harvested from fertile transgenic plants and used to grow progeny generations of transformed plants that produce the recombinant nutritive protein of this disclosure. In addition to direct transformation of a plant with a recombinant DNA, transgenic plants can be prepared by crossing a first plant having a recombinant DNA with a second plant lacking the DNA. For example, recombinant DNA can be introduced into first plant line that is amenable to transformation to produce a transgenic plant which can be crossed with a second plant line to introgress the recombinant DNA into the second plant line. A transgenic plant with recombinant DNA encoding a nutritive protein

of this disclosure, can be crossed with transgenic plant line having other recombinant DNA that confers another trait, for example herbicide resistance or pest resistance, or production of a second nutritive product such as an oil, to produce progeny plants having recombinant DNA that confers both traits. Typically, in such breeding for combining traits the transgenic plant donating the additional trait is a male line and the transgenic plant carrying the base traits is the female line. The progeny of this cross will segregate such that some of the plants will carry the DNA for both parental traits and some will carry DNA for one parental trait; such plants can be identified by markers associated with parental recombinant DNA, e.g. marker identification by analysis for recombinant DNA or, in the case where a selectable marker is linked to the recombinant, by application of the selecting agent such as a herbicide for use with a herbicide tolerance marker, or by selection for the enhanced trait. Progeny plants carrying DNA for both parental traits can be crossed back into the female parent line multiple times, for example usually 6 to 8 generations, to produce a progeny plant with substantially the same genotype as one original transgenic parental line but for the recombinant DNA of the other transgenic parental line.

[00331] In the practice of transformation DNA is typically introduced into only a small percentage of target plant cells in any one transformation experiment. Marker genes are used to provide an efficient system for identification of those cells that are stably transformed by receiving and integrating a transgenic DNA construct into their genomes. Preferred marker genes provide selective markers which confer resistance to a selective agent, such as an antibiotic or herbicide. Any of the herbicides to which the transformed plants may be resistant are useful agents for selective markers. Potentially transformed cells are exposed to the selective agent. In the population of surviving cells will be those cells where, generally, the resistance-conferring gene is integrated and expressed at sufficient levels to permit cell survival. Cells may be tested further to confirm stable integration of the exogenous DNA. Commonly used selective marker genes include those conferring resistance to antibiotics such as kanamycin and paromomycin (nptII), hygromycin B (aph IV) and gentamycin (aac3 and aacC4) or resistance to herbicides such as glufosinate (bar or pat) and glyphosate (aroA or EPSPS). Examples of such selectable are illustrated in U.S. Pat. Nos. 5,550,318; 5,633,435; 5,780,708 and 6,118,047. Selectable markers which provide an ability to visually identify transformants can also be employed, for example, a gene expressing a colored or fluorescent protein such as a luciferase or green fluorescent protein (GFP) or a gene

expressing a beta-glucuronidase or uidA gene (GUS) for which various chromogenic substrates are known.

[00332] Plant cells that survive exposure to the selective agent, or plant cells that have been scored positive in a screening assay, may be cultured in regeneration media and allowed to mature into plants. Developing plantlets regenerated from transformed plant cells can be transferred to plant growth mix, and hardened off, for example, in an environmentally controlled chamber. Plants are regenerated from about 6 weeks to 10 months after a transformant is identified, depending on the initial tissue. Plants may be pollinated using conventional plant breeding methods known to those of skill in the art and seed produced, for example self-pollination is commonly used with transgenic corn. The regenerated transformed plant or its progeny seed or plants can be tested for expression of the recombinant DNA and selected for the presence of a heterologous nutritive protein.

[00333] Transgenic plants derived from the plant cells of this disclosure are grown to generate transgenic plants comprising the heterologous nucleic acid that encodes a nutritive protein of this disclosure and produce transgenic seed and haploid pollen comprising the heterologous nucleic acid sequence. Such plants with enhanced traits are identified by selection of transformed plants or progeny seed for the enhanced trait. Transgenic plants grown from transgenic seed provided herein demonstrate improved agronomic traits that contribute to increased yield or other traits that provide increased plant value, including, for example, improved protein quality such as increasing the content of at least one of essential amino acids, branch chain amino acids, or Leu.

[00334] The transgenic plants are useful as sources of nutritive proteins. For example, in some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of total protein compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of essential amino acids compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of branch chain amino acids compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises an increased weight fraction of Leu compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this

disclosure comprises at least one of: a) an increased ratio of branch chain amino acid residues to total amino acid residues compared to a control non-transgenic plant; b) an increased ratio of Leu residues to total amino acid residues compared to a control non-transgenic plant; and c) an increased ratio of essential amino acid residues to total amino acid residues compared to a control non-transgenic plant. In some embodiments a transgenic plant comprising a recombinant nutritive protein of this disclosure comprises: a) an increased ratio of branch chain amino acid residues to total amino acid residues compared to a control non-transgenic plant; b) an increased ratio of Leu residues to total amino acid residues compared to a control non-transgenic plant; and c) an increased ratio of essential amino acid residues to total amino acid residues compared to a control non-transgenic plant.

[00335] Accordingly, the transgenic plants are useful as sources of high quality protein. The plants may be harvested and used in mammalian diets with or without further processing. For example, flour made from transgenic wheat, cornmeal made from transgenic corn, or rice or rice flour derived from transgenic rice is enriched in at least one of protein, essential amino acids, branch chain amino acids, and Leu compared to similar products made from plants that do not comprise the recombinant nutritive protein. In some embodiments the recombinant nutritive protein is a plant protein or comprises a polypeptide sequence of a plant protein or a derivative or mutein thereof, such as but not necessarily a protein or polypeptide sequence of the same type of plant. In other embodiments the recombinant nutritive protein is not a plant protein or a derivative or mutein thereof.

[00336] In some embodiments the recombinant nutritive protein is recovered or partially recovered from the transgenic plant before it is consumed by a mammal.

H. Compositions

[00337] At least one nutritive protein disclosed herein may be combined with at least one second component to form a nutritive composition. In some embodiments the only source of amino acid in the composition is the at least one nutritive protein disclosed herein. In such embodiments the amino acid composition of the composition will be the same as the amino acid composition of the at least one nutritive protein disclosed herein. In some embodiments the composition comprises at least one nutritive protein disclosed herein and at least one second protein. In some embodiments the at least one second protein is a second nutritive protein disclosed herein, while in other embodiments the at least one second protein is not a nutritive protein disclosed herein. In some embodiments the composition comprises

1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nutritive proteins disclosed herein. In some embodiments the composition comprises 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more proteins that are not nutritive proteins disclosed herein. In some embodiments the composition comprises 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nutritive proteins and the composition comprises 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more proteins that are not nutritive proteins disclosed herein.

[00338] In some embodiments the nutritive composition as described in the preceding paragraph, further comprises at least one of at least one polypeptide, at least one peptide, and at least one free amino acid. In some embodiments the nutritive composition comprises at least one polypeptide and at least one peptide. In some embodiments the nutritive composition comprises at least one polypeptide and at least one free amino acid. In some embodiments the nutritive composition comprises at least one peptide and at least one free amino acid. In some embodiments the at least one polypeptide, at least one peptide, and/or at least one free amino acid comprises amino acids selected from 1) branch chain amino acids, 2) leucine, and 3) essential amino acids. In some embodiments the at least one polypeptide, at least one peptide, and/or at least one free amino acid consists of amino acids selected from 1) branch chain amino acids, 2) leucine, and 3) essential amino acids. In some embodiments, the nutritive composition comprises at least one modified amino acid or a non-standard amino acid. Modified amino acids include amino acids that have modifications to one or more of the carboxy terminus, amino terminus, and/or side chain. Non-standard amino acids may be selected from those that are formed by post-translational modification of proteins, for example, carboxylated glutamate, hydroxyproline, or hypusine. Other non-standard amino acids are not found in proteins. Examples include lanthionine, 2-aminoisobutyric acid, dehydroalanine, gamma-aminobutyric acid, ornithine and citrulline. In some embodiments, the nutritive composition comprises one or more D-amino acids. In some embodiments, the nutritive composition comprises one or more L-amino acids. In some embodiments, the nutritive composition comprises a mixture of one or more D-amino acids and one or more L-amino acids.

[00339] By adding at least one of a polypeptide, a peptide, and a free amino acid to a nutritive composition the proportion of at least one of branch chain amino acids, leucine, and essential amino acids, to total amino acid, present in the composition can be increased.

[00340] In some embodiments the composition comprises at least one carbohydrate. A “carbohydrate” refers to a sugar or polymer of sugars. The terms “saccharide,” “polysaccharide,” “carbohydrate,” and “oligosaccharide” may be used interchangeably. Most carbohydrates are aldehydes or ketones with many hydroxyl groups, usually one on each carbon atom of the molecule. Carbohydrates generally have the molecular formula $C_nH_{2n}O_n$. A carbohydrate may be a monosaccharide, a disaccharide, trisaccharide, oligosaccharide, or polysaccharide. The most basic carbohydrate is a monosaccharide, such as glucose, sucrose, galactose, mannose, ribose, arabinose, xylose, and fructose. Disaccharides are two joined monosaccharides. Exemplary disaccharides include sucrose, maltose, cellobiose, and lactose. Typically, an oligosaccharide includes between three and six monosaccharide units (e.g., raffinose, stachyose), and polysaccharides include six or more monosaccharide units. Exemplary polysaccharides include starch, glycogen, and cellulose. Carbohydrates may contain modified saccharide units such as 2'-deoxyribose wherein a hydroxyl group is removed, 2'-fluororibose wherein a hydroxyl group is replaced with a fluorine, or N-acetylglucosamine, a nitrogen-containing form of glucose (e.g., 2'-fluororibose, deoxyribose, and hexose). Carbohydrates may exist in many different forms, for example, conformers, cyclic forms, acyclic forms, stereoisomers, tautomers, anomers, and isomers.

[00341] In some embodiments the composition comprises at least one lipid. As used herein a “lipid” includes fats, oils, triglycerides, cholesterol, phospholipids, fatty acids in any form including free fatty acids. Fats, oils and fatty acids may be saturated, unsaturated (cis or trans) or partially unsaturated (cis or trans). In some embodiments the lipid comprises at least one fatty acid selected from lauric acid (12:0), myristic acid (14:0), palmitic acid (16:0), palmitoleic acid (16:1), margaric acid (17:0), heptadecenoic acid (17:1), stearic acid (18:0), oleic acid (18:1), linoleic acid (18:2), linolenic acid (18:3), octadecatetraenoic acid (18:4), arachidic acid (20:0), eicosenoic acid (20:1), eicosadienoic acid (20:2), eicosatetraenoic acid (20:4), eicosapentaenoic acid (20:5) (EPA), docosanoic acid (22:0), docosenoic acid (22:1), docosapentaenoic acid (22:5), docosahexaenoic acid (22:6) (DHA), and tetracosanoic acid (24:0). In some embodiments the composition comprises at least one modified lipid, for example a lipid that has been modified by cooking.

[00342] In some embodiments the composition comprises at least one supplemental mineral or mineral source. Examples of minerals include, without limitation: chloride, sodium, calcium, iron, chromium, copper, iodine, zinc, magnesium, manganese, molybdenum, phosphorus, potassium, and selenium. Suitable forms of any of the foregoing

minerals include soluble mineral salts, slightly soluble mineral salts, insoluble mineral salts, chelated minerals, mineral complexes, non-reactive minerals such as carbonyl minerals, and reduced minerals, and combinations thereof.

[00343] In some embodiments the composition comprises at least one supplemental vitamin. The at least one vitamin can be fat-soluble or water soluble vitamins. Suitable vitamins include but are not limited to vitamin C, vitamin A, vitamin E, vitamin B12, vitamin K, riboflavin, niacin, vitamin D, vitamin B6, folic acid, pyridoxine, thiamine, pantothenic acid, and biotin. Suitable forms of any of the foregoing are salts of the vitamin, derivatives of the vitamin, compounds having the same or similar activity of the vitamin, and metabolites of the vitamin.

[00344] In some embodiments the composition comprises at least one organism. Suitable examples are well known in the art and include probiotics (e.g., species of *Lactobacillus* or *Bifidobacterium*), spirulina, chlorella, and porphyra.

[00345] In some embodiments the composition comprises at least one dietary supplement. Suitable examples are well known in the art and include herbs, botanicals, and certain hormones. Non limiting examples include ginkgo, ginseng, and melatonin.

[00346] In some embodiments the composition comprises an excipient. Non-limiting examples of suitable excipients include a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent.

[00347] In some embodiments the excipient is a buffering agent. Non-limiting examples of suitable buffering agents include sodium citrate, magnesium carbonate, magnesium bicarbonate, calcium carbonate, and calcium bicarbonate.

[00348] In some embodiments the excipient comprises a preservative. Non-limiting examples of suitable preservatives include antioxidants, such as alpha-tocopherol and ascorbate, and antimicrobials, such as parabens, chlorobutanol, and phenol.

[00349] In some embodiments the composition comprises a binder as an excipient. Non-limiting examples of suitable binders include starches, pregelatinized starches, gelatin, polyvinylpyrrolidone, cellulose, methylcellulose, sodium carboxymethylcellulose, ethylcellulose, polyacrylamides, polyvinylloxazolidone, polyvinylalcohols, C₁₂-C₁₈ fatty acid

alcohol, polyethylene glycol, polyols, saccharides, oligosaccharides, and combinations thereof.

[00350] In some embodiments the composition comprises a lubricant as an excipient. Non-limiting examples of suitable lubricants include magnesium stearate, calcium stearate, zinc stearate, hydrogenated vegetable oils, sterotex, polyoxyethylene monostearate, talc, polyethyleneglycol, sodium benzoate, sodium lauryl sulfate, magnesium lauryl sulfate, and light mineral oil.

[00351] In some embodiments the composition comprises a dispersion enhancer as an excipient. Non-limiting examples of suitable dispersants include starch, alginic acid, polyvinylpyrrolidones, guar gum, kaolin, bentonite, purified wood cellulose, sodium starch glycolate, isoamorphous silicate, and microcrystalline cellulose as high HLB emulsifier surfactants.

[00352] In some embodiments the composition comprises a disintegrant as an excipient. In some embodiments the disintegrant is a non-effervescent disintegrant. Non-limiting examples of suitable non-effervescent disintegrants include starches such as corn starch, potato starch, pregelatinized and modified starches thereof, sweeteners, clays, such as bentonite, micro-crystalline cellulose, alginates, sodium starch glycolate, gums such as agar, guar, locust bean, karaya, pectin, and tragacanth. In some embodiments the disintegrant is an effervescent disintegrant. Non-limiting examples of suitable effervescent disintegrants include sodium bicarbonate in combination with citric acid, and sodium bicarbonate in combination with tartaric acid.

[00353] In some embodiments the excipient comprises a flavoring agent. Flavoring agents incorporated into the outer layer can be chosen from synthetic flavor oils and flavoring aromatics; natural oils; extracts from plants, leaves, flowers, and fruits; and combinations thereof. In some embodiments the flavoring agent is selected from cinnamon oils; oil of wintergreen; peppermint oils; clover oil; hay oil; anise oil; eucalyptus; vanilla; citrus oil such as lemon oil, orange oil, grape and grapefruit oil; and fruit essences including apple, peach, pear, strawberry, raspberry, cherry, plum, pineapple, and apricot.

[00354] In some embodiments the excipient comprises a sweetener. Non-limiting examples of suitable sweeteners include glucose (corn syrup), dextrose, invert sugar, fructose, and mixtures thereof (when not used as a carrier); saccharin and its various salts such as the sodium salt; dipeptide sweeteners such as aspartame; dihydrochalcone

compounds, glycyrrhizin; Stevia Rebaudiana (Stevioside); chloro derivatives of sucrose such as sucralose; and sugar alcohols such as sorbitol, mannitol, sylitol, and the like. Also contemplated are hydrogenated starch hydrolysates and the synthetic sweetener 3,6-dihydro-6-methyl-1,2,3-oxathiazin-4-one-2,2-dioxide, particularly the potassium salt (acesulfame-K), and sodium and calcium salts thereof.

[00355] In some embodiments the composition comprises a coloring agent. Non-limiting examples of suitable color agents include food, drug and cosmetic colors (FD&C), drug and cosmetic colors (D&C), and external drug and cosmetic colors (Ext. D&C). The coloring agents can be used as dyes or their corresponding lakes.

[00356] The weight fraction of the excipient or combination of excipients in the formulation is usually about 50% or less, about 45% or less, about 40% or less, about 35% or less, about 30% or less, about 25% or less, about 20% or less, about 15% or less, about 10% or less, about 5% or less, about 2% or less, or about 1% or less of the total weight of the amino acids in the composition.

[00357] The nutritive proteins and nutritive compositions disclosed herein can be formulated into a variety of forms and administered by a number of different means. The compositions can be administered orally, rectally, or parenterally, in formulations containing conventionally acceptable carriers, adjuvants, and vehicles as desired. The term “parenteral” as used herein includes subcutaneous, intravenous, intramuscular, or intrasternal injection and infusion techniques. In an exemplary embodiment, the nutritive protein or composition is administered orally.

[00358] Solid dosage forms for oral administration include capsules, tablets, caplets, pills, troches, lozenges, powders, and granules. A capsule typically comprises a core material comprising a nutritive protein or composition and a shell wall that encapsulates the core material. In some embodiments the core material comprises at least one of a solid, a liquid, and an emulsion. In some embodiments the shell wall material comprises at least one of a soft gelatin, a hard gelatin, and a polymer. Suitable polymers include, but are not limited to: cellulosic polymers such as hydroxypropyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose (HPMC), methyl cellulose, ethyl cellulose, cellulose acetate, cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropylmethyl cellulose phthalate, hydroxypropylmethyl cellulose succinate and carboxymethylcellulose sodium; acrylic acid polymers and copolymers, such as those formed from acrylic acid, methacrylic acid, methyl

acrylate, ammonio methacrylate, ethyl acrylate, methyl methacrylate and/or ethyl methacrylate (e.g., those copolymers sold under the trade name “Eudragit”); vinyl polymers and copolymers such as polyvinyl pyrrolidone, polyvinyl acetate, polyvinylacetate phthalate, vinylacetate crotonic acid copolymer, and ethylene-vinyl acetate copolymers; and shellac (purified lac). In some embodiments at least one polymer functions as taste-masking agents.

[00359] Tablets, pills, and the like can be compressed, multiply compressed, multiply layered, and/or coated. The coating can be single or multiple. In one embodiment, the coating material comprises at least one of a saccharide, a polysaccharide, and glycoproteins extracted from at least one of a plant, a fungus, and a microbe. Non-limiting examples include corn starch, wheat starch, potato starch, tapioca starch, cellulose, hemicellulose, dextrans, maltodextrin, cyclodextrins, inulins, pectin, mannans, gum arabic, locust bean gum, mesquite gum, guar gum, gum karaya, gum ghatti, tragacanth gum, funori, carrageenans, agar, alginates, chitosans, or gellan gum. In some embodiments the coating material comprises a protein. In some embodiments the coating material comprises at least one of a fat and an oil. In some embodiments the at least one of a fat and an oil is high temperature melting. In some embodiments the at least one of a fat and an oil is hydrogenated or partially hydrogenated. In some embodiments the at least one of a fat and an oil is derived from a plant. In some embodiments the at least one of a fat and an oil comprises at least one of glycerides, free fatty acids, and fatty acid esters. In some embodiments the coating material comprises at least one edible wax. The edible wax can be derived from animals, insects, or plants. Non-limiting examples include beeswax, lanolin, bayberry wax, carnauba wax, and rice bran wax. Tablets and pills can additionally be prepared with enteric coatings.

[00360] Alternatively, powders or granules embodying the nutritive proteins and nutritive compositions disclosed herein can be incorporated into a food product. In some embodiments the food product is be a drink for oral administration. Non-limiting examples of a suitable drink include fruit juice, a fruit drink, an artificially flavored drink, an artificially sweetened drink, a carbonated beverage, a sports drink, a liquid dairy product, a shake, an alcoholic beverage, a caffeinated beverage, infant formula and so forth. Other suitable means for oral administration include aqueous and nonaqueous solutions, creams, pastes, emulsions, suspensions and slurries, , each of which may optionally also containin at least one of suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, coloring agents, and flavoring agents.

[00361] In some embodiments the food product is a solid foodstuff. Suitable examples of a solid foodstuff include without limitation a food bar, a snack bar, a cookie, a brownie, a muffin, a cracker, a biscuit, a cream or paste, an ice cream bar, a frozen yogurt bar, and the like.

[00362] In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are incorporated into a therapeutic food. In some embodiments, the therapeutic food is a ready-to-use food that optionally contains some or all essential macronutrients and micronutrients. In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are incorporated into a supplementary food that is designed to be blended into an existing meal. In some embodiments, the supplemental food contains some or all essential macronutrients and micronutrients. In some embodiments, the nutritive proteins and nutritive compositions disclosed herein are blended with or added to an existing food to fortify the food's protein nutrition. Examples include food staples (grain, salt, sugar, cooking oil, margarine), beverages (coffee, tea, soda, beer, liquor, sports drinks), snacks, sweets and other foods.

[00363] The compositions disclosed herein can be utilized in methods to increase at least one of muscle mass, strength and physical function, thermogenesis, metabolic expenditure, satiety, mitochondrial biogenesis, weight or fat loss, and lean body composition for example.

I. Methods of Use

[00364] In some embodiments the nutritive proteins and nutritive compositions disclosed herein are administered to a patient or a user (sometimes collectively referred to as a "subject"). As used herein "administer" and "administration" encompasses embodiments in which one person directs another to consume a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose, and also situations in which a user uses a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose independently of or in variance to any instructions received from a second person. Non-limiting examples of embodiments in which one person directs another to consume a nutritive protein or nutritive composition in a certain manner and/or for a certain purpose include when a physician prescribes a course of conduct and/or treatment to a patient, when a trainer advises a user (such as an athlete) to follow a particular course of conduct and/or treatment, and when a manufacturer, distributor, or marketer recommends conditions of use to an end

user, for example through advertisements or labeling on packaging or on other materials provided in association with the sale or marketing of a product.

[00365] In some embodiments the nutritive proteins or nutritive compositions are provided in a dosage form. In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from 0.1g to 1g, 1g to 5g, from 2g to 10g, from 5g to 15g, from 10g to 20g, from 15g to 25g, from 20g to 40g, from 25-50g, and from 30-60g. In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from about 0.1g, 0.1g-1g, 1g, 2g, 3g, 4g, 5g, 6g, 7g, 8g, 9g, 10g, 15g, 20g, 25g, 30g, 35g, 40g, 45g, 50g, 55g, 60g, 65g, 70g, 75g, 80g, 85g, 90g, 95g, and 100g.

[00366] In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of essential amino acids administered is selected from 0.1g to 1g, from 1g to 5g, from 2g to 10g, from 5g to 15g, from 10g to 20g, and from 1-30 g. In some embodiments the dosage form is designed for administration of at least one nutritive protein disclosed herein, wherein the total amount of nutritive protein administered is selected from about 0.1g, 0.1-1g, 1g, 2g, 3g, 4g, 5g, 6g, 7g, 8g, 9g, 10g, 15g, 20g, 25g, 30g, 35g, 40g, 45g, 50g, 55g, 60g, 65g, 70g, 75g, 80g, 85g, 90g, 95g, and 100g.

[00367] In some embodiments the nutritive protein or nutritive composition is consumed at a rate of from 0.1g to 1g a day, 1g to 5 g a day, from 2g to 10g a day, from 5g to 15g a day, from 10g to 20g a day, from 15g to 30g a day, from 20g to 40g a day, from 25g to 50g a day, from 40g to 80g a day, from 50g to 100g a day, or more.

[00368] In some embodiments, of the total protein intake by the subject, at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or about 100% of the total protein intake by the subject over a dietary period is made up of at least one nutritive protein according to this disclosure. In some embodiments, of the total protein intake by the subject, from 5% to 100% of the total protein intake by the subject, from 5% to 90% of the total protein intake by the subject, from 5% to 80% of the total protein intake by the subject, from 5% to 70% of the total protein intake by the subject, from 5% to 60% of the total protein intake by the subject,

from 5% to 50% of the total protein intake by the subject, from 5% to 40% of the total protein intake by the subject, from 5% to 30% of the total protein intake by the subject, from 5% to 20% of the total protein intake by the subject, from 5% to 10% of the total protein intake by the subject, from 10% to 100% of the total protein intake by the subject, from 10% to 100% of the total protein intake by the subject, from 20% to 100% of the total protein intake by the subject, from 30% to 100% of the total protein intake by the subject, from 40% to 100% of the total protein intake by the subject, from 50% to 100% of the total protein intake by the subject, from 60% to 100% of the total protein intake by the subject, from 70% to 100% of the total protein intake by the subject, from 80% to 100% of the total protein intake by the subject, or from 90% to 100% of the total protein intake by the subject, over a dietary period, is made up of at least one nutritive protein according to this disclosure. In some embodiments the at least one nutritive protein of this disclosure accounts for at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, or at least 50% of the subject's calorie intake over a dietary period.

[00369] In some embodiments the at least one nutritive protein according to this disclosure comprises at least 2 nutritive proteins of this disclosure, at least 3 nutritive proteins of this disclosure, at least 4 nutritive proteins of this disclosure, at least 5 nutritive proteins of this disclosure, at least 6 nutritive proteins of this disclosure, at least 7 nutritive proteins of this disclosure, at least 8 nutritive proteins of this disclosure, at least 9 nutritive proteins of this disclosure, at least 10 nutritive proteins of this disclosure, or more.

[00370] In some embodiments the dietary period is 1 meal, 2 meals, 3 meals, at least 1 day, at least 2 days, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least 1 week, at least 2 weeks, at least 3 weeks, at least 4 weeks, at least 1 month, at least 2 months, at least 3 months, at least 4 months, at least 5 months, at least 6 months, or at least 1 year. In some embodiments the dietary period is from 1 day to 1 week, from 1 week to 4 weeks, from 1 month, to 3 months, from 3 months to 6 months, or from 6 months to 1 year.

[00371] Clinical studies provide evidence that protein prevents muscle loss due to aging or bed rest. In particular, studies have shown that protein supplementation increases muscle fractional synthetic rate (FSR) during prolonged bed rest, maintains leg mass and strength during prolonged bed rest, increases lean body mass, improves functional measures of gait and balance, and may serve as a viable intervention for individuals at risk of sarcopenia due to immobility or prolonged bed rest. (See, e.g., Paddon-Jones D, et al. *J Clin*

Endocrinol Metab 2004, 89:4351–4358; Ferrando, A et al. *Clinical Nutrition* 2009 1-6; Katsanos C et al. *Am J Physiol Endocrinol Metab.* 2006, 291: 381-387).

[00372] Studies on increasing muscle protein anabolism in athletes have shown that protein provided following exercise promotes muscle hypertrophy to a greater extent than that achieved by exercise alone. It has also been shown that protein provided following exercise supports protein synthesis without any increase in protein breakdown, resulting in a net positive protein balance and muscle mass accretion. While muscle protein synthesis appears to respond in a dose-response fashion to essential amino acid supplementation, not all proteins are equal in building muscle. For example, the amino acid leucine is an important factor in stimulating muscle protein synthesis. (See, e.g., Borsheim E et al. *Am J Physiol Endocrinol Metab* 2002, 283: E648-E657; Borsheim E et al. *Clin Nutr.* 2008, 27: 189-95; Esmarck B et al *J Physiol* 2001, 535: 301-311; Moore D et al. *Am J Clin Nutr* 2009, 89: 161-8).

[00373] In another aspect this disclosure provides methods of maintaining or increasing at least one of muscle mass, muscle strength, and functional performance in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00374] In another aspect this disclosure provides methods of maintaining or achieving a desirable body mass index in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a

nutritive composition made by a method of this disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00375] In another aspect this disclosure provides methods of providing protein to a subject with protein-energy malnutrition. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the nutritive protein of this disclosure, nutritive composition of this disclosure, or nutritive composition made by a method of this disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00376] The need for essential amino acid supplementation has been suggested in cancer patients and other patients suffering from cachexia. Dietary studies in mice have shown survival and functional benefits to cachectic cancer-bearing mice through dietary intervention with essential amino acids. Beyond cancer, essential amino acid supplementation has also shown benefits, such as improved muscle function and muscle gain, in patients suffering from other diseases who have difficulty exercising and therefore suffer from muscular deterioration, such as chronic obstructive pulmonary disease, chronic heart failure, HIV, and other disease states.

[00377] Studies have shown that specific amino acids have advantages in managing cachexia. A relatively high content of BCAAs and Leu in diets are thought to have a positive effect in cachexia by promoting total protein synthesis by signaling an increase in translation, enhancing insulin release, and inhibiting protein degradation. Thus, consuming increased dietary BCAAs in general and/or Leu in particular will contribute positively to reduce or reverse the effects of cachexia. Because nitrogen balance is important in countering the underlying cause of cachexia it is thought that consuming increased dietary glutamine and/or arginine will contribute positively to reduce or reverse the effects of cachexia. (See, e.g., Op den Kamp C, Langen R, Haegens A, Schols A. "Muscle atrophy in cachexia: can dietary protein tip the balance?" *Current Opinion in Clinical Nutrition and Metabolic Care* 2009, 12:611–616; Poon RT-P, Yu W-C, Fan S-T, et al. "Long-term oral branched chain amino acids in patients undergoing chemoembolization for hepatocellular carcinoma: a randomized trial." *Aliment Pharmacol Ther* 2004; 19:779–788; Tayek JA, Bistrian BR, Hehir DJ, Martin

R, Moldawer LL, Blackburn GL. "Improved protein kinetics and albumin synthesis by branched chain amino acid-enriched total parenteral nutrition in cancer cachexia." *Cancer*. 1986;58:147-57; Xi P, Jiang Z, Zheng C, Lin Y, Wu G "Regulation of protein metabolism by glutamine: implications for nutrition and health." *Front Biosci*. 2011 Jan 1;16:578-97).

[00378] Accordingly, also provided herein are methods of treating cachexia in a subject. In some embodiments a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure for a subject with cachexia is an amount such that the amount of protein of this disclosure ingested by the person meets or exceeds the metabolic needs (which are often elevated). A protein intake of 1.5 g/kg of body weight per day or 15-20% of total caloric intake appears to be an appropriate target for persons with cachexia. In some embodiments all of the protein consumed by the subject is a nutritive protein according to this disclosure. In some embodiments nutritive protein according to this disclosure is combined with other sources of protein and/or free amino acids to provide the total protein intake of the subject. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments the subject suffers from a disease that makes exercise difficult and therefore causes muscular deterioration, such as chronic obstructive pulmonary disease, chronic heart failure, HIV, cancer, and other disease states. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to this disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00379] Sarcopenia is the degenerative loss of skeletal muscle mass (typically 0.5-1% loss per year after the age of 25), quality, and strength associated with aging. Sarcopenia is a component of the frailty syndrome. The European Working Group on Sarcopenia in Older People (EWGSOP) has developed a practical clinical definition and consensus diagnostic criteria for age-related sarcopenia. For the diagnosis of sarcopenia, the working group has proposed using the presence of both low muscle mass and low muscle function (strength or performance). Sarcopenia is characterized first by a muscle atrophy (a decrease in the size of the muscle), along with a reduction in muscle tissue "quality," caused by such factors as replacement of muscle fibres with fat, an increase in fibrosis, changes in muscle metabolism,

oxidative stress, and degeneration of the neuromuscular junction. Combined, these changes lead to progressive loss of muscle function and eventually to frailty. Frailty is a common geriatric syndrome that embodies an elevated risk of catastrophic declines in health and function among older adults. Contributors to frailty can include sarcopenia, osteoporosis, and muscle weakness. Muscle weakness, also known as muscle fatigue, (or “lack of strength”) refers to the inability to exert force with one’s skeletal muscles. Weakness often follows muscle atrophy and a decrease in activity, such as after a long bout of bedrest as a result of an illness. There is also a gradual onset of muscle weakness as a result of sarcopenia.

[00380] The nutritive proteins of this disclosure are useful for treating sarcopenia or frailty once it develops in a subject or for preventing the onset of sarcopenia or frailty in a subject who is a member of an at risk groups. In some embodiments all of the protein consumed by the subject is a nutritive protein according to this disclosure. In some embodiments nutritive protein according to this disclosure is combined with other sources of protein and/or free amino acids to provide the total protein intake of the subject. In some embodiments the subject is at least one of elderly, critically-medically ill, and suffering from protein-energy malnutrition. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to this disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00381] Obesity is a multifactorial disorder associated with a host of comorbidities including hypertension, type 2 diabetes, dyslipidemia, coronary heart disease, stroke, cancer (eg, endometrial, breast, and colon), osteoarthritis, sleep apnea, and respiratory problems. The incidence of obesity, defined as a body mass index $>30 \text{ kg/m}^2$, has increased dramatically in the United States, from 15% (1976–1980) to 33% (2003–2004), and it continues to grow. Although the mechanisms contributing to obesity are complex and involve the interplay of behavioral components with hormonal, genetic, and metabolic processes, obesity is largely viewed as a lifestyle-dependent condition with 2 primary causes: excessive energy intake and insufficient physical activity. With respect to energy intake, there is evidence that modestly increasing the proportion of protein in the diet, while controlling total energy intake, may improve body composition, facilitate fat loss, and improve body weight

maintenance after weight loss. Positive outcomes associated with increased dietary protein are thought to be due primarily to lower energy intake associated with increased satiety, reduced energy efficiency and/or increased thermogenesis, positive effects on body composition (specifically lean muscle mass), and enhanced glycemic control.

[00382] Dietary proteins are more effective in increasing post-prandial energy expenditure than isocaloric intakes of carbohydrates or fat (see, e.g., Dauncey M, Bingham S. “Dependence of 24 h energy expenditure in man on composition of the nutrient intake.” *Br J Nutr* 1983, 50: 1-13; Karst H et al. “Diet-induced thermogenesis in man: thermic effects of single proteins, carbohydrates and fats depending on their energy amount.” *Ann Nutr Metab*.1984, 28: 245-52; Tappy L et al “Thermic effect of infused amino acids in healthy humans and in subjects with insulin resistance.” *Am J Clin Nutr* 1993, 57 (6): 912-6). This property along with other properties (satiety induction; preservation of lean body mass) make protein an attractive component of diets directed at weight management. The increase in energy expenditure caused by such diets may in part be due to the fact that the energy cost of digesting and metabolizing protein is higher than for other calorie sources. Protein turnover, including protein synthesis, is an energy consuming process. In addition, high protein diets may also up-regulate uncoupling protein in liver and brown adipose, which is positively correlated with increases in energy expenditure. It has been theorized that different proteins may have unique effects on energy expenditure.

[00383] Studies suggest that ingestion of protein, particularly proteins with high EAA and/or BCAA content, leads to distinct effects on thermogenesis and energy expenditure (see, e.g., Mikkelsen P. et al. “Effect of fat-reduced diets on 24 h energy expenditure: comparisons between animal protein, vegetable protein and carbohydrate.” *Am J Clin Nutr* 2000, 72:1135-41; Acheson K. et al. “Protein choices targeting thermogenesis and metabolism.” *Am J Clin Nutr* 2011, 93:525-34; Alfenas R. et al. “Effects of protein quality on appetite and energy metabolism in normal weight subjects” *Arg Bras Endocrinol Metabol* 2010, 54 (1): 45-51; Lorenzen J. et al. “The effect of milk proteins on appetite regulation and diet-induced thermogenesis.” *J Clin Nutr* 2012 66 (5): 622-7). Additionally, L-tyrosine has been identified as an amino acid that plays a role in thermogenesis (see, e.g., Belza A. et al. “The beta-adrenergic antagonist propranolol partly abolishes thermogenic response to bioactive food ingredients.” *Metabolism* 2009, 58 (8):1137-44). Further studies suggest that Leucine and Arginine supplementation appear to alter energy metabolism by directing substrate to lean body mass rather than adipose tissue (Dulloo A. “The search for compounds that

stimulate thermogenesis in obesity management: from pharmaceuticals to functional food ingredients.” *Obes Rev* 2011 12: 866-83).

[00384] Collectively the literature suggests that different protein types leads to distinct effects on thermogenesis. Because proteins or peptides rich in EAAs, BCAA, and/or at least one of Tyr, Arg, and Leu are believed to have a stimulatory effect on thermogenesis, and because stimulation of thermogenesis is believed to lead to positive effects on weight management, this disclosure also provides products and methods useful to stimulation thermogenesis and/or to bring about positive effects on weight management in general.

[00385] More particularly, this disclosure provides methods of increasing thermogenesis in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00386] At the basic level, the reason for the development of an overweight condition is due to an imbalance between energy intake and energy expenditure. Attempts to reduce food at any particular occasion (satiation) and across eating occasions (satiety) have been a major focus of recent research. Reduced caloric intake as a consequence of feeling satisfied during a meal and feeling full after a meal results from a complex interaction of internal and external signals. Various nutritional studies have demonstrated that variation in food properties such as energy density, content, texture and taste influence both satiation and satiety.

[00387] There are three macronutrients that deliver energy: fat, carbohydrates and proteins. A gram of protein or carbohydrate provides 4 calories while a gram of fat 9 calories. Protein generally increases satiety to a greater extent than carbohydrates or fat and therefore may facilitate a reduction in calorie intake. However, there is considerable evidence that indicates the type of protein matters in inducing satiety (see, e.g., W.L. Hall, et al.

“Casein and whey exert different effects on plasma amino acid profiles, gastrointestinal hormone secretion and appetite.” *Br J Nutr.* 2003 Feb, 89(2):239-48; R. Abou-Samra, et al. “Effect of different protein sources on satiation and short-term satiety when consumed as a starter.” *Nutr J.* 2011 Dec 23, 10:139; T. Akhavan, et al. “Effect of premeal consumption of whey protein and its hydrolysate on food intake and postmeal glycemia and insulin responses in young adults.” *Am J Clin Nutr.* 2010 Apr, 91(4):966-75, Epub 2010 Feb 17; MA Veldhorst “Dose-dependent satiating effect of whey relative to casein or soy” *Physiol Behav.* 2009 Mar 23, 96(4-5):675-82). Evidence indicates that protein rich in Leucine is particularly effective at inducing satiety (see, e.g., Fromentin G et al “Peripheral and central mechanisms involved in the control of food intake by dietary amino acids and proteins.” *Nutr Res Rev* 2012 25: 29-39).

[00388] In some embodiments a nutritive protein of this disclosure is consumed by a subject concurrently with at least one pharmaceutical or biologic drug product. In some embodiments the beneficial effects of the nutritive protein and the at least one pharmaceutical or biologic drug product have an additive effect while in some embodiments the beneficial effects of the nutritive protein and the at least one pharmaceutical or biologic drug product have a synergistic effect. Examples of pharmaceutical or biologic drug products that can be administered with the nutritive proteins of this disclosure are well known in the art. For example, when a nutritive protein of this disclosure is used to maintain or increase at least one of muscle mass, muscle strength, and functional performance in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to maintain or increase at least one of muscle mass, muscle strength, and functional performance in a subject, such as an anabolic steroid. When a nutritive protein of this disclosure is used to maintain or achieve a desirable body mass index in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to maintain or achieve a desirable body mass index in a subject, such as orlistat, lorcaserin, sibutramine, rimonabant, metformin, exenatide, or pramlintide. When a nutritive protein of this disclosure is used to induce at least one of a satiation response and a satiety response in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to induce at least one of a satiation response and a satiety response in a subject, such as rimonabant, exenatide, or pramlintide. When a nutritive protein of this disclosure is

used to treat at least one of cachexia, sarcopenia and frailty in a subject, the nutritive protein may be consumed by a subject concurrently with a therapeutic dosage regime of at least one pharmaceutical or biologic drug product indicated to treat at least one of cachexia, sarcopenia and frailty, such as omega-3 fatty acids or anabolic steroids. Because of the role of dietary protein in inducing satiation and satiety, the nutritive proteins and nutritive compositions disclosed herein can be used to induce at least one of a satiation response and a satiety response in a subject. In some embodiments the methods comprise providing to the subject a sufficient amount of a nutritive protein of this disclosure, a nutritive composition of this disclosure, or a nutritive composition made by a method of this disclosure. In some embodiments the subject is obese. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject in coordination with performance of exercise. In some embodiments, the nutritive protein according to disclosure, the nutritive composition according to disclosure, or the nutritive composition made by a method according to disclosure is consumed by the subject by an oral, enteral, or parenteral route.

[00389] In some embodiments incorporating a least one nutritive protein or nutritive composition of this disclosure into the diet of a subject has at least one effect selected from inducing postprandial satiety (including by suppressing hunger), inducing thermogenesis, reducing glycemic response, positively affecting energy expenditure positively affecting lean body mass, reducing the weight gain caused by overeating, and decreasing energy intake. In some embodiments incorporating a least one nutritive protein or nutritive composition of this disclosure into the diet of a subject has at least one effect selected from increasing loss of body fat, reducing lean tissue loss, improving lipid profile, and improving glucose tolerance and insulin sensitivity in the subject.

EXAMPLES

[00390] The following examples serve to more fully describe the manner of using the invention. These examples are presented for illustrative purposes and should not serve to limit the true scope of the invention.

Example 1: Identification of Naturally Occurring Proteins and Fragments

[00391] Naturally occurring proteins and fragments of naturally occurring proteins with useful PDCAAS values and/or similarity to a chosen EAA/NEAA (Whey) target amino

acid profile were identified by searching the full set of sequences in the reviewed UniProt database (also known as the SwissProt database). The full database set of proteins was trimmed to proteins that are non-human and non-viral in origin, and do not contain the words “toxic”, “toxin”, “virus”, or “viral” in either the protein or species name. The search was also limited to only those proteins that have been identified due to experimental observation of the protein itself or its mRNA transcript.

[00392] For each protein the weight fraction of each amino acid was computed and the PDCAAS value derived. The proteins were also evaluated for similarity to EAA/NEAA (Whey) target amino acid profiles of 70/30 and 60/40 using a Euclidean distance metric as described herein.

[00393] Additionally, fragments of each protein were identified and analyzed to compute the weight fraction of each amino acid and the PDCAAS value for the fragment. A deterministic, hierarchical search strategy employing tiered fragmentation frequencies and PDCAAS cutoffs was used to identify sets of 1,000 fragments from the protein database having the highest PDCAAS value or similarity to a chosen EAA/NEAA (Whey) target amino acid profile. While this approach could be applied to analyze every fragment of every possible size within every protein in the database the search was truncated for the purpose of the examples presented herein.

Example 2: Naturally Occurring Protein Fragments Ranked by PDCAAS Score

[00394] In this example the PDCAAS Score was used to rank protein fragments identified in the SwissProt database using the approach of Example 1. The 1000 fragments from the analysis with the highest PDCAAS scores were identified and are included in the Sequence Listing. The 100 fragments from the analysis with the highest PDCAAS scores were identified and are presented in **Table 6**. In **Table 6A** the proteins the fragments are derived from are identified by UniProtID number in the first column of the table. The fragment itself is further identified by the amino acids positions in the protein that appear at the N and C termini of the fragment (second column). For example (12:145) indicates that the N-terminus of the fragment is amino acid 12 of the protein and the C-terminus of the fragment is at amino acid 145 of the protein. The third column indicates whether the fragment contains every essential amino acid (1) or not (0). The fourth and fifth columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively.

The sixth column presents the PDCAAS score for each fragment. For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated and are presented in **Table 6B**.

[00395] Figures 1-4 present an analysis of the essential amino acid content of four of the identified fragments that have among the highest PCDAAS scores.

[00396] The essential amino acid content of the polypeptide fragment from amino acid 51 to amino acid 75 of the *Schizosaccharomyces pombe* ENTH domain-containing protein C19F8.03c protein (UniProt O60167) (SEQ ID NO: 415) is analyzed in **Figure 1**. That fragment has a PCDAAS value of 2.46. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00397] The essential amino acid content of the polypeptide fragment from amino acid 170 to amino acid 263 (SEQ ID NO: 2016) of the *Ovis aries* C-C chemokine receptor 9 protein (UniProt Q1WLP9) is presented in **Figure 2**. That fragment has a PCDAAS value of 2.41. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00398] The essential amino acid content of the polypeptide fragment from amino acid 307 to amino acid 369 of the *Gallus gallus* neuronal acetylcholine receptor subunit alpha-6

protein (UniProt P49581) (SEQ ID NO: 1366) is presented in **Figure 3**. That fragment has a PCDAAS value of 2.40. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00399] The essential amino acid content of the polypeptide fragment from amino acid 314 to amino acid 402 of the *Arabidopsis thaliana* putative cadmium/zinc-transporting ATPase HMA4 protein (UniProt O64474) (SEQ ID NO: 421) is presented in **Figure 4**. That fragment has a PCDAAS value of 2.39. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein. The PCDAAS scores of the set of 1000 fragments range from 2.61 to 1.93

Table 6A

SeqIDNo	UniProtID	Fragment Ends	EAA Complete	[EAA]	[NEAA]	PDCAAS
415	O60167	51:75	1	0.75	0.25	2.46
2016	Q1WLP9	170:263	1	0.67	0.33	2.41
2003	Q1LZF8	292:355	1	0.69	0.31	2.41
1366	P49581	307:369	1	0.75	0.25	2.40
421	O64474	314:402	1	0.63	0.37	2.39
4200	Q9SZW4	319:392	0	0.67	0.33	2.38
4104	Q9SA37	379:420	1	0.73	0.27	2.38
3975	Q9M008	219:281	1	0.68	0.32	2.36
1822	Q08213	177:230	1	0.65	0.35	2.36
4109	Q9SAK8	338:405	1	0.72	0.28	2.32

WO 2013/148325

PCT/US2013/032180

222	O13712	665:758	1	0.64	0.36	2.31
4334	Q9Y704	1186:1249	1	0.81	0.19	2.31
3984	Q9M175	380:406	1	0.78	0.22	2.31
666	P05363	246:272	1	0.81	0.19	2.30
4012	Q9N1R5	87:140	1	0.71	0.29	2.30
3513	Q92182	3:45	1	0.66	0.34	2.29
2715	Q5ZLF4	71:119	1	0.75	0.25	2.27
3793	Q9FFT9	289:345	1	0.71	0.29	2.27
3788	Q9FE38	458:555	1	0.65	0.35	2.27
1198	P38803	135:177	1	0.64	0.36	2.27
4025	Q9P7K5	750:815	1	0.69	0.31	2.27
2565	Q5N749	423:451	0	0.64	0.36	2.26
3967	Q9LYG9	209:381	1	0.68	0.32	2.26
3173	Q84YJ9	155:213	1	0.68	0.32	2.26
1323	P47077	526:582	1	0.64	0.36	2.26
3855	Q9GL50	119:161	1	0.68	0.32	2.26
3361	Q8LES0	199:257	1	0.64	0.36	2.26
2118	Q2HJ17	188:220	1	0.78	0.22	2.25
2062	Q28044	270:299	1	0.85	0.15	2.23
3300	Q8GXD6	257:300	1	0.72	0.28	2.23
3677	Q9CA25	249:277	1	0.73	0.27	2.23
274	O23413	50:162	1	0.63	0.37	2.23
88	A6QNL7	225:289	1	0.68	0.32	2.23
3163	Q84M97	167:299	1	0.75	0.25	2.22
2063	Q28132	2:67	1	0.61	0.39	2.22
2065	Q28132-2	2:67	1	0.61	0.39	2.22
1110	P32584	109:169	1	0.68	0.32	2.22
494	O81770	267:295	0	0.80	0.20	2.22
496	O81770-2	267:295	0	0.80	0.20	2.22
3844	Q9FZL4	264:292	0	0.80	0.20	2.22
194	O04722	302:347	1	0.70	0.30	2.22
762	P0CW78	328:423	1	0.66	0.34	2.21
185	O02697	195:223	1	0.63	0.37	2.21
3589	Q95M19	2:67	1	0.60	0.40	2.20
2466	Q58CW4	217:283	1	0.66	0.34	2.20

WO 2013/148325

PCT/US2013/032180

3806	Q9FIJ0	384:478	1	0.61	0.39	2.20
1628	P79368	2:67	1	0.61	0.39	2.20
1182	P38221	184:252	1	0.73	0.27	2.20
3595	Q95N03	166:288	1	0.64	0.36	2.19
2185	Q39079	328:359	1	0.70	0.30	2.19
3590	Q95MD2	41:70	0	0.62	0.38	2.19
1953	Q12276	525:592	1	0.67	0.33	2.19
1625	P79222-2	323:389	1	0.65	0.35	2.19
2131	Q2KTE1	218:251	1	0.79	0.21	2.18
31	A2PZE5	102:164	1	0.65	0.35	2.18
4162	Q9SUP0	14:154	1	0.65	0.35	2.18
3981	Q9M173	347:405	1	0.65	0.35	2.18
1688	P92949	91:150	1	0.69	0.31	2.18
1944	Q10M80	109:181	1	0.75	0.25	2.18
263	O22757	52:132	1	0.67	0.33	2.18
3450	Q8VYC2	329:358	1	0.66	0.34	2.18
1781	Q05005	63:109	1	0.78	0.22	2.17
3302	Q8GXR2	243:357	1	0.69	0.31	2.17
3445	Q8VXB5	123:205	1	0.62	0.38	2.17
1187	P38250	262:327	1	0.65	0.35	2.17
997	P25360	441:511	1	0.67	0.33	2.17
4212	Q9TTU3	158:231	0	0.60	0.40	2.17
342	O48901	982:1011	0	0.58	0.42	2.17
1909	Q0WPQ2	73:175	1	0.73	0.27	2.17
1230	P40456	607:652	1	0.78	0.22	2.16
2979	Q6R8G0	623:725	1	0.66	0.34	2.16
3835	Q9FUG4	412:474	1	0.61	0.39	2.16
269	O23278	382:454	1	0.67	0.33	2.16
1204	P38960	412:470	1	0.61	0.39	2.16
4194	Q9SYZ7	199:229	0	0.75	0.25	2.16
1959	Q12520	46:149	1	0.63	0.37	2.15
2318	Q41342	59:140	1	0.71	0.29	2.15
4107	Q9SAH8	407:438	1	0.75	0.25	2.14
486	O81016	482:572	1	0.68	0.32	2.14
1207	P39073	299:329	1	0.67	0.33	2.14

1192	P38352	488:534	0	0.60	0.40	2.14
798	P12394	294:339	1	0.64	0.36	2.14
4174	Q9SW17	439:499	1	0.72	0.28	2.14
1166	P36512	483:528	0	0.60	0.40	2.14
538	O94461	112:172	1	0.67	0.33	2.14
3926	Q9LK39	383:452	1	0.63	0.37	2.14
4179	Q9SWS1	50:105	1	0.73	0.27	2.13
2111	Q29RU8	508:600	1	0.62	0.38	2.13
2175	Q38707	142:215	1	0.57	0.43	2.13
1185	P38242	111:180	1	0.64	0.36	2.13
2079	Q28997	133:178	1	0.68	0.32	2.13
4342	Q9Y802	499:614	1	0.61	0.39	2.13
3882	Q9IAL7	5:67	1	0.62	0.38	2.13
3884	Q9IAL7-2	5:67	1	0.62	0.38	2.13
3606	Q96323	233:296	1	0.61	0.39	2.13
867	P16661	71:125	1	0.68	0.32	2.13
3372	Q8LPL8	459:492	0	0.73	0.27	2.13
2171	Q32PJ3	129:314	1	0.60	0.40	2.13
3974	Q9LZR3	418:463	1	0.69	0.31	2.12
3867	Q9H DU1	160:204	1	0.72	0.28	2.12

Table 6B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
415	-19.86	1.03	0.14	0.42	0.38	0.46
2016	-13.06	1.31	0.29	0.22	0.25	0.87
2003	-11.83	1.11	0.25	0.00	0.00	0.91
1366	-14.98	1.13	0.20	0.00	0.00	0.76
421	-11.24	1.23	0.29	0.00	0.29	0.35
4200	-11.84	1.25	0.24	0.00	0.25	0.42
4104	-14.53	1.17	0.15	0.26	0.19	0.29
3975	-13.90	1.08	0.21	0.31	0.29	0.00
1822	-20.36	0.77	0.23	0.31	0.22	0.33
4109	-13.02	1.12	0.22	0.25	0.22	0.30

WO 2013/148325

PCT/US2013/032180

222	-14.59	0.78	0.28	0.00	0.25	0.27
4334	-9.87	1.23	0.24	0.00	0.00	0.28
3984	-13.17	0.75	0.12	0.00	0.37	0.41
666	-9.89	1.63	0.12	0.37	0.29	0.96
4012	-12.94	1.24	0.19	0.31	0.00	0.96
3513	-14.66	1.15	0.18	0.29	0.29	0.41
2715	-10.97	1.06	0.20	0.32	0.31	0.94
3793	-17.61	0.71	0.20	0.24	0.00	0.32
3788	-11.14	0.90	0.30	0.30	0.28	0.25
1198	-16.72	0.59	0.16	0.35	0.33	0.36
4025	-14.82	0.84	0.23	0.24	0.25	0.00
2565	-15.51	0.87	0.14	0.34	0.34	0.45
3967	-16.45	0.90	0.30	0.21	0.23	0.22
3173	-14.87	0.74	0.18	0.27	0.26	0.32
1323	-19.83	0.42	0.19	0.28	0.32	0.33
3855	-14.21	1.06	0.21	0.28	0.30	0.95
3361	-12.31	0.91	0.23	0.24	0.25	0.34
2118	-8.63	1.61	0.15	0.33	0.27	0.88
2062	-8.99	1.50	0.13	0.40	0.37	0.97
3300	-14.39	1.03	0.16	0.00	0.31	0.41
3677	-12.31	1.21	0.13	0.00	0.31	0.41
274	-11.52	1.07	0.30	0.22	0.23	0.22
88	-15.67	1.14	0.22	0.26	0.28	0.82
3163	-16.23	1.15	0.28	0.22	0.00	0.00
2063	-16.26	0.62	0.24	0.24	0.22	0.91
2065	-16.26	0.62	0.24	0.24	0.22	0.91
1110	-13.94	0.81	0.19	0.25	0.28	0.39
494	-15.68	0.63	0.14	0.41	0.32	0.40
496	-15.68	0.63	0.14	0.41	0.32	0.40
3844	-15.68	0.63	0.14	0.41	0.32	0.40
194	-14.55	1.03	0.21	0.00	0.33	0.35
762	-13.63	0.95	0.29	0.00	0.25	0.38
185	-12.75	0.66	0.14	0.38	0.34	0.90
3589	-17.24	0.60	0.23	0.24	0.25	0.91
2466	-14.15	1.18	0.22	0.00	0.00	0.84

WO 2013/148325

PCT/US2013/032180

3806	-12.16	0.94	0.28	0.00	0.23	0.36
1628	-17.23	0.60	0.23	0.24	0.22	0.92
1182	-10.60	1.28	0.24	0.26	0.28	0.52
3595	-16.40	1.10	0.29	0.24	0.21	0.74
2185	-15.32	0.88	0.14	0.00	0.41	0.38
3590	-17.01	0.47	0.12	0.35	0.42	0.97
1953	-15.64	0.87	0.23	0.24	0.28	0.28
1625	-15.74	0.72	0.25	0.25	0.25	0.79
2131	-12.52	1.47	0.12	0.38	0.24	1.00
31	-9.20	1.02	0.21	0.31	0.29	0.33
4162	-13.57	1.11	0.32	0.25	0.23	0.00
3981	-9.72	0.89	0.21	0.00	0.00	0.38
1688	-16.78	1.02	0.25	0.27	0.30	0.28
1944	-13.57	0.92	0.27	0.00	0.00	0.35
263	-15.59	1.06	0.26	0.33	0.22	0.29
3450	-17.72	0.65	0.15	0.29	0.37	0.37
1781	-13.94	1.32	0.16	0.00	0.31	0.83
3302	-17.35	0.88	0.31	0.00	0.00	0.33
3445	-17.33	1.18	0.25	0.00	0.24	0.24
1187	-12.62	1.04	0.22	0.27	0.27	0.28
997	-15.05	1.12	0.25	0.24	0.00	0.28
4212	-11.70	0.99	0.27	0.31	0.25	0.99
342	-16.97	0.64	0.13	0.39	0.38	0.40
1909	-14.17	1.10	0.29	0.00	0.22	0.24
1230	-11.20	1.06	0.20	0.28	0.30	0.35
2979	-15.58	0.96	0.28	0.00	0.22	0.28
3835	-18.62	0.61	0.23	0.28	0.25	0.71
269	-21.06	0.75	0.22	0.00	0.26	0.00
1204	-19.20	0.44	0.20	0.24	0.30	0.29
4194	-16.03	1.31	0.15	0.32	0.32	0.42
1959	-12.87	0.93	0.29	0.00	0.26	0.29
2318	-12.83	1.09	0.26	0.26	0.26	0.27
4107	-8.94	1.07	0.13	0.41	0.38	0.41
486	-13.12	0.98	0.27	0.00	0.28	0.27
1207	-16.58	0.67	0.14	0.28	0.00	0.52

1192	-14.46	0.68	0.19	0.31	0.31	0.29
798	-19.06	0.68	0.17	0.00	0.25	0.63
4174	-13.67	1.08	0.23	0.00	0.00	0.34
1166	-13.75	1.21	0.18	0.29	0.35	0.63
538	-21.74	0.58	0.22	0.26	0.00	0.31
3926	-17.64	0.77	0.24	0.25	0.29	0.00
4179	-9.92	1.22	0.22	0.20	0.28	0.42
2111	-15.67	0.74	0.26	0.00	0.27	0.80
2175	-13.73	0.75	0.36	0.00	0.29	0.33
1185	-14.52	0.87	0.21	0.30	0.28	0.31
2079	-10.56	1.13	0.17	0.24	0.30	0.85
4342	-17.56	0.53	0.28	0.26	0.26	0.28
3882	-18.45	0.86	0.22	0.00	0.24	0.56
3884	-18.45	0.86	0.22	0.00	0.24	0.56
3606	-17.83	0.53	0.21	0.26	0.29	0.00
867	-15.72	0.80	0.22	0.24	0.33	0.31
3372	-8.74	1.51	0.16	0.26	0.32	0.32
2171	-17.28	0.64	0.29	0.00	0.22	0.88
3974	-19.93	0.71	0.17	0.31	0.00	0.30
3867	-11.83	1.43	0.16	0.35	0.29	0.30

Example 3: Protein Fragments Ranked by Euclidian Distance from a 60/40 EAA/NEAA (Whey) Profile or a 70/30 EAA/NEAA (Whey) Profile

[00400] In this example the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile or from a 70/30 EAA/NEAA (Whey) target amino acid profile was used to rank protein fragments identified in the SwissProt database using the approach of Example 1. After protein fragments were identified in this way the PDCAAS of each fragment was also determined. The 1000 fragments from the analysis with the smallest Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile are presented in the Sequence Listing. The 100 fragments from the analysis with the smallest Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile are presented in **Table 7**. In **Table 7A** the proteins the fragments are derived from are identified by UniProtID number in the first column of the table. The fragment itself is further identified by the amino acids positions in the protein that appear at the N and C termini of the fragment (second

column). For example (12:145) indicates that the N-terminus of the fragment is amino acid 12 of the protein and the C-terminus of the fragment is at amino acid 145 of the protein. The third column indicates whether the fragment contains every essential amino acid (1) or not (0). The fourth and fifth columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The sixth column presents the PDCAAS score for each fragment. For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated and are presented in **Table 7B**.

[00401] The 1000 fragments from the analysis with the smallest Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile were identified and are included in the Sequence Listing. The 100 fragments from the analysis with the smallest Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile are presented in **Table 8**. In each table the fragments were first ranked by PDCAAS score and then ranked by Euclidian distance from the target profile. In **Table 8A** the proteins the fragments are derived from are identified by UniProtID number in the first column of the table. The fragment itself is further identified by the amino acids positions in the protein that appear at the N and C termini of the fragment (second column). For example (12:145) indicates that the N-terminus of the fragment is amino acid 12 of the protein and the C-terminus of the fragment is at amino acid 145 of the protein. The third column indicates whether the fragment contains every essential amino acid (1) or not (0). The fourth and fifth columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The sixth column presents the PDCAAS score for each fragment. The seventh column presents the distance from the target distribution for each fragment. For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e.,

percent homology to known human proteins, as described herein) were calculated and are presented in **Table 8B**.

[00402] The Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of the set of 1000 fragments ranges from 0.0458 to 0.1524. The Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of the set of 1000 fragments ranges from 0.0380 to 0.117.

Table 7A

SeqIDNo	UniProtID	Fragment Ends	EAA Complete	[EAA]	[NEAA]	PDCAAS	Dist. Target
4199	Q9SZW4	277:380	1	0.58	0.42	2.20	0.0459
2017	Q1WLP9	170:279	1	0.65	0.35	2.11	0.0603
3836	Q9FUG4	412:476	1	0.59	0.41	2.10	0.0612
4343	Q9Y802	504:615	1	0.62	0.38	2.08	0.0754
4163	Q9SUP0	4:157	1	0.65	0.35	2.07	0.0779
87	A6QNL7	221:293	1	0.65	0.35	2.06	0.1091
4401	Q9ZVE0	63:170	1	0.56	0.44	2.01	0.0578
195	O04722	303:355	1	0.67	0.33	1.98	0.0616
1274	P43544	6:118	1	0.53	0.47	1.98	0.0622
2467	Q58CW4	227:284	1	0.64	0.36	1.97	0.0630
268	O23278	380:461	1	0.64	0.36	1.96	0.0651
2799	Q65AC2	315:513	1	0.60	0.40	1.96	0.0660
1433	P53666	343:479	1	0.59	0.41	1.96	0.0693
1837	Q08960	543:629	1	0.56	0.44	1.99	0.0694
4238	Q9UTL9	261:383	1	0.59	0.41	2.00	0.0697
2202	Q3E7D1	154:352	1	0.56	0.44	1.95	0.0703
1919	Q0WVW7	702:769	1	0.58	0.42	2.04	0.0713
3911	Q9LD37	224:300	1	0.62	0.38	1.96	0.0715
3569	Q94LW6	205:361	1	0.60	0.40	1.99	0.0717
89	A6QQL0	269:436	1	0.59	0.41	1.96	0.0720
4024	Q9P7D7	190:372	1	0.52	0.48	1.98	0.0733
2697	Q5ZIL2	19:105	1	0.56	0.44	1.96	0.0761
3825	Q9FLT4	306:360	1	0.69	0.31	2.04	0.0764
4178	Q9SWS1	22:81	1	0.63	0.37	1.99	0.0765
2156	Q32L23	343:479	1	0.58	0.42	1.96	0.0766

WO 2013/148325

PCT/US2013/032180

1867	Q09889	197:311	1	0.53	0.47	1.96	0.0774
1234	P40527	728:868	1	0.53	0.47	1.95	0.0775
2468	Q58D00	227:349	0	0.53	0.47	2.01	0.0781
3566	Q94KB7	394:462	1	0.61	0.39	1.96	0.0783
3514	Q92182	3:50	1	0.63	0.37	2.05	0.0790
4173	Q9SW17	332:483	1	0.61	0.39	1.97	0.0800
709	P08417	357:481	1	0.54	0.46	1.97	0.0807
3956	Q9LVQ5	53:111	1	0.63	0.37	1.97	0.0812
2110	Q29RU8	504:605	1	0.59	0.41	1.95	0.0812
54	A4IFA3	595:712	1	0.56	0.44	2.00	0.0815
3925	Q9LK39	366:485	1	0.57	0.43	1.97	0.0833
2910	Q6LA56	26:154	1	0.58	0.42	2.04	0.0839
1111	P32584	95:169	1	0.65	0.35	2.04	0.0846
1071	P30620	434:546	1	0.59	0.41	2.03	0.0868
1689	P93028	905:985	1	0.56	0.44	1.95	0.0880
469	O77480	292:356	1	0.55	0.45	1.97	0.0881
3976	Q9M008	224:290	1	0.65	0.35	1.98	0.0912
1203	P38960	408:472	1	0.57	0.43	1.96	0.0913
223	O13712	677:807	1	0.59	0.41	2.00	0.0926
1441	P53730	331:364	1	0.66	0.34	2.01	0.1053
1005	P25847	46:133	1	0.55	0.45	1.86	0.0608
344	O49160	246:319	1	0.62	0.38	1.85	0.0613
1832	Q08905	454:639	1	0.54	0.46	1.86	0.0615
2525	Q5F361	127:253	1	0.56	0.44	1.87	0.0632
4189	Q9SY55	234:338	1	0.59	0.41	1.91	0.0636
3816	Q9FKR3	144:245	1	0.59	0.41	1.85	0.0637
2108	Q29RL1	292:396	1	0.63	0.37	1.89	0.0642
1159	P36421	1:126	1	0.54	0.46	1.87	0.0651
534	O94265	97:269	1	0.55	0.45	1.91	0.0670
3299	Q8GXD6	255:338	1	0.60	0.40	1.94	0.0676
1325	P47082	386:521	1	0.60	0.40	1.88	0.0682
1814	Q07527	1305:1435	1	0.57	0.43	1.87	0.0687
3605	Q96323	219:285	1	0.56	0.44	1.94	0.0688
4132	Q9SJ73	323:417	1	0.60	0.40	1.93	0.0689
1776	Q04658	238:369	1	0.56	0.44	1.90	0.0690

WO 2013/148325

PCT/US2013/032180

357	O49621	14:205	1	0.62	0.38	1.91	0.0693
2702	Q5ZKB7	194:377	1	0.54	0.46	1.92	0.0699
3962	Q9LXW7	90:222	1	0.55	0.45	1.91	0.0700
3454	Q8VZZ4	791:964	1	0.54	0.46	1.91	0.0701
3983	Q9M175	261:435	1	0.59	0.41	1.86	0.0713
3591	Q95MD2	9:94	1	0.58	0.42	1.88	0.0721
2101	Q29449	706:835	1	0.52	0.48	1.94	0.0725
1213	P39535	411:492	1	0.63	0.37	1.92	0.0728
1119	P33333	28:198	1	0.59	0.41	1.85	0.0735
1197	P38803	112:228	1	0.56	0.45	1.95	0.0739
2693	Q5ZIE4	6:158	1	0.60	0.40	1.94	0.0741
484	O80970	97:179	1	0.54	0.46	1.85	0.0743
866	P16661	68:127	1	0.65	0.35	1.94	0.0746
475	O80580	1:62	1	0.59	0.41	1.95	0.0746
4150	Q9SNX8	43:233	1	0.54	0.46	1.87	0.0747
1882	Q0P5F2	163:266	1	0.56	0.44	1.92	0.0752
1317	P46682	341:510	1	0.58	0.42	1.91	0.0755
481	O80961	10:89	1	0.69	0.31	1.93	0.0768
3807	Q9FIJ0	396:498	1	0.55	0.45	1.87	0.0769
4026	Q9P7K5	783:939	1	0.55	0.45	1.90	0.0770
120	A8MQA3	127:245	1	0.53	0.47	1.95	0.0771
3594	Q95N03	149:315	1	0.62	0.38	1.89	0.0771
3968	Q9LYG9	240:435	1	0.60	0.40	1.88	0.0773
1773	Q04602	660:742	1	0.63	0.37	1.90	0.0774
1335	P48034	235:348	1	0.55	0.45	1.90	0.0776
3	A0A1F3	256:332	1	0.57	0.43	1.89	0.0778
1054	P29415	209:255	1	0.69	0.31	1.87	0.0779
2235	Q3MHL4	271:420	1	0.56	0.44	1.90	0.0779
4011	Q9N1R5	80:147	1	0.66	0.34	1.90	0.0786
2160	Q32LB9	472:609	1	0.58	0.42	1.87	0.0789
2699	Q5ZJ43	484:641	1	0.55	0.45	1.89	0.0797
1421	P53119	712:792	1	0.58	0.42	1.88	0.0801
2978	Q6R8G0	611:732	1	0.62	0.38	1.89	0.0809
4388	Q9ZQR8-2	66:167	1	0.57	0.43	1.91	0.0810
62	A5D7Q3	24:111	1	0.63	0.37	1.85	0.0812

WO 2013/148325

PCT/US2013/032180

251	O18784-2	566:658	1	0.60	0.40	1.89	0.0815
3370	Q8LPK2	288:379	1	0.59	0.41	1.85	0.0817
1229	P40395	602:693	1	0.54	0.46	1.91	0.0821
4229	Q9USR3	1131:1188	1	0.66	0.34	1.87	0.0823
3101	Q7XRA1	177:332	1	0.52	0.48	1.86	0.0826

Table 7B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
4199	-14.98	1.08	0.31	0.25	0.23	0.37
2017	-12.50	1.23	0.29	0.00	0.26	0.87
3836	-19.27	0.59	0.23	0.25	0.24	0.72
4343	-18.71	0.56	0.28	0.28	0.25	0.29
4163	-13.46	1.07	0.32	0.20	0.23	0.23
87	-15.30	1.05	0.25	0.26	0.29	0.84
4401	-19.75	0.65	0.29	0.24	0.00	0.45
195	-16.02	0.90	0.18	0.00	0.29	0.35
1274	-18.89	0.60	0.29	0.23	0.00	0.00
2467	-17.40	1.05	0.20	0.29	0.00	0.84
268	-19.84	0.69	0.25	0.00	0.26	0.00
2799	-13.62	0.94	0.30	0.00	0.23	0.95
1433	-21.81	0.48	0.33	0.24	0.21	0.98
1837	-19.50	0.59	0.30	0.20	0.28	0.63
4238	-17.25	0.54	0.32	0.28	0.21	0.30
2202	-17.51	0.59	0.31	0.20	0.00	0.00
1919	-17.71	0.62	0.24	0.26	0.29	0.27
3911	-16.41	0.77	0.26	0.00	0.23	0.34
3569	-13.40	0.95	0.33	0.23	0.21	0.27
89	-15.83	0.76	0.31	0.00	0.23	0.91
4024	-16.44	0.64	0.31	0.00	0.25	0.36
2697	-17.36	0.48	0.27	0.27	0.24	0.97
3825	-16.80	1.05	0.21	0.00	0.29	0.33
4178	-13.84	0.92	0.20	0.00	0.28	0.41
2156	-21.84	0.48	0.33	0.24	0.21	0.98

WO 2013/148325

PCT/US2013/032180

1867	-19.79	0.51	0.33	0.00	0.26	0.30
1234	-20.34	0.59	0.31	0.21	0.00	0.46
2468	-18.22	0.59	0.33	0.26	0.25	1.00
3566	-16.43	0.75	0.24	0.25	0.32	0.30
3514	-17.70	1.04	0.19	0.33	0.33	0.41
4173	-15.21	0.96	0.29	0.25	0.25	0.30
709	-17.14	0.58	0.31	0.20	0.25	0.63
3956	-17.30	0.70	0.20	0.00	0.29	0.32
2110	-15.86	0.71	0.27	0.23	0.25	0.81
54	-18.30	0.58	0.29	0.25	0.24	0.88
3925	-15.58	0.62	0.31	0.00	0.23	0.27
2910	-21.23	0.52	0.29	0.22	0.00	0.22
1111	-14.66	0.89	0.23	0.24	0.28	0.37
1071	-20.49	0.58	0.29	0.00	0.24	0.29
1689	-17.13	0.66	0.31	0.00	0.00	0.52
469	-16.59	0.57	0.22	0.28	0.25	0.86
3976	-14.46	0.99	0.22	0.31	0.25	0.28
1203	-22.02	0.30	0.24	0.28	0.24	0.28
223	-16.44	0.73	0.28	0.24	0.20	0.28
1441	-12.67	0.97	0.11	0.39	0.31	0.38
1005	-17.12	0.54	0.29	0.00	0.00	0.31
344	-18.02	0.58	0.25	0.28	0.22	0.33
1832	-17.14	0.51	0.31	0.00	0.25	0.25
2525	-19.16	0.61	0.32	0.23	0.22	0.79
4189	-15.25	0.92	0.28	0.27	0.00	0.74
3816	-19.02	0.58	0.28	0.00	0.25	0.29
2108	-18.89	0.61	0.29	0.00	0.25	0.74
1159	-18.00	0.51	0.29	0.23	0.24	0.60
534	-17.93	0.64	0.31	0.00	0.22	0.00
3299	-16.28	0.85	0.28	0.25	0.26	0.46
1325	-16.76	0.77	0.33	0.00	0.00	0.29
1814	-18.97	0.61	0.30	0.00	0.00	0.42
3605	-16.19	0.60	0.22	0.25	0.26	0.28
4132	-17.41	0.75	0.29	0.00	0.00	0.26
1776	-15.63	0.97	0.30	0.23	0.24	0.00

WO 2013/148325

PCT/US2013/032180

357	-17.98	0.87	0.31	0.00	0.22	0.00
2702	-18.32	0.62	0.30	0.00	0.22	0.73
3962	-18.81	0.69	0.33	0.24	0.00	0.38
3454	-18.24	0.86	0.29	0.00	0.24	0.30
3983	-16.22	0.81	0.29	0.00	0.22	0.22
3591	-18.96	0.53	0.29	0.26	0.26	0.88
2101	-17.25	0.61	0.27	0.26	0.25	0.98
1213	-12.68	1.25	0.27	0.21	0.23	0.31
1119	-15.92	0.69	0.31	0.00	0.25	0.34
1197	-18.25	0.59	0.32	0.23	0.26	0.25
2693	-19.37	0.67	0.30	0.23	0.23	0.77
484	-18.51	0.67	0.27	0.00	0.00	0.24
866	-17.20	0.71	0.23	0.25	0.30	0.30
475	-16.68	0.85	0.20	0.17	0.28	0.00
4150	-17.81	0.61	0.39	0.00	0.22	0.32
1882	-16.21	0.58	0.27	0.00	0.28	0.84
1317	-17.23	0.59	0.31	0.00	0.22	0.29
481	-14.92	1.28	0.26	0.27	0.26	0.31
3807	-16.59	0.79	0.30	0.27	0.27	0.32
4026	-20.21	0.52	0.31	0.00	0.25	0.00
120	-16.57	0.64	0.33	0.20	0.25	0.26
3594	-14.60	1.09	0.30	0.23	0.23	0.76
3968	-19.46	0.69	0.30	0.00	0.20	0.00
1773	-17.69	0.53	0.27	0.00	0.00	0.00
1335	-18.51	0.52	0.30	0.00	0.27	0.82
3	-20.64	0.56	0.25	0.28	0.00	0.92
1054	-14.97	1.12	0.20	0.31	0.29	0.91
2235	-19.14	0.47	0.31	0.23	0.00	0.97
4011	-10.82	1.15	0.23	0.00	0.26	0.94
2160	-18.47	0.59	0.33	0.25	0.26	0.85
2699	-21.38	0.62	0.30	0.00	0.00	0.94
1421	-19.03	0.60	0.29	0.00	0.25	0.34
2978	-16.80	0.90	0.30	0.00	0.00	0.27
4388	-20.52	0.61	0.29	0.25	0.25	0.29
62	-17.27	0.39	0.27	0.00	0.23	0.89

251	-15.12	0.92	0.29	0.29	0.27	0.97
3370	-13.90	0.84	0.30	0.28	0.25	0.34
1229	-18.50	0.55	0.27	0.00	0.26	0.30
4229	-16.46	0.71	0.19	0.35	0.29	0.25
3101	-17.11	0.55	0.32	0.24	0.22	0.24

Table 8A

SeqIDNo	UniProtID	Fragment Ends	EAA Complete	[EAA]	[NEAA]	PDCAAS	Dist. Target
4199	Q9SZW4	277:380	1	0.58	0.42	2.20	0.0533
3836	Q9FUG4	412:476	1	0.59	0.41	2.10	0.0593
4343	Q9Y802	504:615	1	0.62	0.38	2.08	0.0778
4163	Q9SUP0	4:157	1	0.65	0.35	2.07	0.0866
2017	Q1WLP9	170:279	1	0.65	0.35	2.11	0.0873
4401	Q9ZVE0	63:170	1	0.56	0.44	2.01	0.0410
1275	P43544	7:116	1	0.53	0.47	2.03	0.0477
4190	Q9SY55	234:340	1	0.58	0.42	2.01	0.0548
709	P08417	357:481	1	0.54	0.46	1.97	0.0574
1234	P40527	728:868	1	0.53	0.47	1.95	0.0652
2202	Q3E7D1	154:352	1	0.56	0.44	1.95	0.0661
4024	Q9P7D7	190:372	1	0.52	0.48	1.98	0.0677
89	A6QQL0	269:436	1	0.59	0.41	1.96	0.0701
1433	P53666	343:479	1	0.59	0.41	1.96	0.0701
2799	Q65AC2	315:513	1	0.60	0.40	1.96	0.0701
1919	Q0WVW7	702:769	1	0.58	0.42	2.04	0.0706
268	O23278	380:461	1	0.64	0.36	1.96	0.0708
1837	Q08960	543:629	1	0.56	0.44	1.99	0.0723
195	O04722	303:355	1	0.67	0.33	1.98	0.0747
2468	Q58D00	227:349	0	0.53	0.47	2.01	0.0756
1689	P93028	905:985	1	0.56	0.44	1.95	0.0770
2156	Q32L23	343:479	1	0.58	0.42	1.96	0.0771
3569	Q94LW6	205:361	1	0.60	0.40	1.99	0.0778
2910	Q6LA56	26:154	1	0.58	0.42	2.04	0.0787
1867	Q09889	197:311	1	0.53	0.47	1.96	0.0794

WO 2013/148325

PCT/US2013/032180

4238	Q9UTL9	261:383	1	0.59	0.41	2.00	0.0796
2697	Q5ZIL2	19:105	1	0.56	0.44	1.96	0.0806
3925	Q9LK39	366:485	1	0.57	0.43	1.97	0.0822
54	A4IFA3	595:712	1	0.56	0.44	2.00	0.0828
2110	Q29RU8	504:605	1	0.59	0.41	1.95	0.0834
3911	Q9LD37	224:300	1	0.62	0.38	1.96	0.0840
1071	P30620	434:546	1	0.59	0.41	2.03	0.0851
2467	Q58CW4	227:284	1	0.64	0.36	1.97	0.0876
4173	Q9SW17	332:483	1	0.61	0.39	1.97	0.0879
3566	Q94KB7	394:462	1	0.61	0.39	1.96	0.0883
4178	Q9SWS1	22:81	1	0.63	0.37	1.99	0.0893
3956	Q9LVQ5	53:111	1	0.63	0.37	1.97	0.0904
469	O77480	292:356	1	0.55	0.45	1.97	0.0906
1203	P38960	408:472	1	0.57	0.43	1.96	0.0965
3825	Q9FLT4	306:360	1	0.69	0.31	2.04	0.1020
223	O13712	677:807	1	0.59	0.41	2.00	0.1027
3514	Q92182	3:50	1	0.63	0.37	2.05	0.1076
1111	P32584	95:169	1	0.65	0.35	2.04	0.1082
3454	Q8VZZ4	791:964	1	0.54	0.46	1.91	0.0503
534	O94265	97:269	1	0.55	0.45	1.91	0.0520
1832	Q08905	454:639	1	0.54	0.46	1.86	0.0524
1814	Q07527	1305:1435	1	0.57	0.43	1.87	0.0532
2702	Q5ZKB7	194:377	1	0.54	0.46	1.92	0.0535
4150	Q9SNX8	43:233	1	0.54	0.46	1.87	0.0542
267	O23255	298:473	1	0.55	0.45	1.86	0.0546
2101	Q29449	706:835	1	0.52	0.48	1.94	0.0560
1335	P48034	235:348	1	0.55	0.45	1.90	0.0575
3962	Q9LXW7	90:222	1	0.55	0.45	1.91	0.0593
3605	Q96323	219:285	1	0.56	0.44	1.94	0.0594
1159	P36421	1:126	1	0.54	0.46	1.87	0.0597
1005	P25847	46:133	1	0.55	0.45	1.86	0.0604
344	O49160	246:319	1	0.62	0.38	1.85	0.0610
2108	Q29RL1	292:396	1	0.63	0.37	1.89	0.0612
2699	Q5ZJ43	484:641	1	0.55	0.45	1.89	0.0616
120	A8MQA3	127:245	1	0.53	0.47	1.95	0.0622

2525	Q5F361	127:253	1	0.56	0.44	1.87	0.0624
3101	Q7XRA1	177:332	1	0.52	0.48	1.86	0.0631
2235	Q3MHL4	271:420	1	0.56	0.44	1.90	0.0633
1776	Q04658	238:369	1	0.56	0.44	1.90	0.0652
3816	Q9FKR3	144:245	1	0.59	0.41	1.85	0.0655
2160	Q32LB9	472:609	1	0.58	0.42	1.87	0.0656
484	O80970	97:179	1	0.54	0.46	1.85	0.0665
475	O80580	1:62	1	0.59	0.41	1.95	0.0666
3983	Q9M175	261:435	1	0.59	0.41	1.86	0.0673
3792	Q9FFT9	138:337	1	0.55	0.45	1.90	0.0678
1317	P46682	341:510	1	0.58	0.42	1.91	0.0679
3	A0A1F3	256:332	1	0.57	0.43	1.89	0.0692
328	O43043	629:731	1	0.56	0.44	1.88	0.0703
985	P24719	9:132	1	0.51	0.49	1.90	0.0708
2693	Q5ZIE4	6:158	1	0.60	0.40	1.94	0.0709
4388	Q9ZQR8-2	66:167	1	0.57	0.43	1.91	0.0710
1229	P40395	602:693	1	0.54	0.46	1.91	0.0711
1197	P38803	112:228	1	0.56	0.45	1.95	0.0715
3968	Q9LYG9	240:435	1	0.60	0.40	1.88	0.0717
3299	Q8GXD6	255:338	1	0.60	0.40	1.94	0.0723
324	O42939	3:198	1	0.53	0.47	1.89	0.0733
357	O49621	14:205	1	0.62	0.38	1.91	0.0735
852	P15398	1465:1537	1	0.55	0.45	1.86	0.0738
4026	Q9P7K5	783:939	1	0.55	0.45	1.90	0.0743
4132	Q9SJ73	323:417	1	0.60	0.40	1.93	0.0743
1325	P47082	386:521	1	0.60	0.40	1.88	0.0745
3955	Q9LVF0	30:192	1	0.51	0.49	1.89	0.0756
3807	Q9FIJ0	396:498	1	0.55	0.45	1.87	0.0764
1421	P53119	712:792	1	0.58	0.42	1.88	0.0797
1119	P33333	28:198	1	0.59	0.41	1.85	0.0798
3370	Q8LPK2	288:379	1	0.59	0.41	1.85	0.0801
3919	Q9LFN5	229:428	1	0.56	0.44	1.87	0.0801
1827	Q08217	544:679	1	0.57	0.43	1.86	0.0807
1995	Q1JQA1	160:326	1	0.52	0.48	1.93	0.0810
3351	Q8L500	46:85	0	0.52	0.48	1.89	0.0812

3874	Q9HE16	395:464	1	0.56	0.44	1.94	0.0818
45	A2VDP5	206:306	1	0.54	0.46	1.90	0.0818
1213	P39535	411:492	1	0.63	0.37	1.92	0.0833
1882	Q0P5F2	163:266	1	0.56	0.44	1.92	0.0848
1468	P54612	422:589	1	0.55	0.45	1.83	0.0381

Table 8B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
4199	-14.98	1.08	0.31	0.25	0.23	0.37
3836	-19.27	0.59	0.23	0.25	0.24	0.72
4343	-18.71	0.56	0.28	0.28	0.25	0.29
4163	-13.46	1.07	0.32	0.20	0.23	0.23
2017	-12.50	1.23	0.29	0.00	0.26	0.87
4401	-19.75	0.65	0.29	0.24	0.00	0.45
1275	-18.77	0.61	0.30	0.25	0.21	0.00
4190	-15.00	0.94	0.28	0.27	0.00	0.74
709	-17.14	0.58	0.31	0.20	0.25	0.63
1234	-20.34	0.59	0.31	0.21	0.00	0.46
2202	-17.51	0.59	0.31	0.20	0.00	0.00
4024	-16.44	0.64	0.31	0.00	0.25	0.36
89	-15.83	0.76	0.31	0.00	0.23	0.91
1433	-21.81	0.48	0.33	0.24	0.21	0.98
2799	-13.62	0.94	0.30	0.00	0.23	0.95
1919	-17.71	0.62	0.24	0.26	0.29	0.27
268	-19.84	0.69	0.25	0.00	0.26	0.00
1837	-19.50	0.59	0.30	0.20	0.28	0.63
195	-16.02	0.90	0.18	0.00	0.29	0.35
2468	-18.22	0.59	0.33	0.26	0.25	1.00
1689	-17.13	0.66	0.31	0.00	0.00	0.52
2156	-21.84	0.48	0.33	0.24	0.21	0.98
3569	-13.40	0.95	0.33	0.23	0.21	0.27
2910	-21.23	0.52	0.29	0.22	0.00	0.22
1867	-19.79	0.51	0.33	0.00	0.26	0.30

WO 2013/148325

PCT/US2013/032180

4238	-17.25	0.54	0.32	0.28	0.21	0.30
2697	-17.36	0.48	0.27	0.27	0.24	0.97
3925	-15.58	0.62	0.31	0.00	0.23	0.27
54	-18.30	0.58	0.29	0.25	0.24	0.88
2110	-15.86	0.71	0.27	0.23	0.25	0.81
3911	-16.41	0.77	0.26	0.00	0.23	0.34
1071	-20.49	0.58	0.29	0.00	0.24	0.29
2467	-17.40	1.05	0.20	0.29	0.00	0.84
4173	-15.21	0.96	0.29	0.25	0.25	0.30
3566	-16.43	0.75	0.24	0.25	0.32	0.30
4178	-13.84	0.92	0.20	0.00	0.28	0.41
3956	-17.30	0.70	0.20	0.00	0.29	0.32
469	-16.59	0.57	0.22	0.28	0.25	0.86
1203	-22.02	0.30	0.24	0.28	0.24	0.28
3825	-16.80	1.05	0.21	0.00	0.29	0.33
223	-16.44	0.73	0.28	0.24	0.20	0.28
3514	-17.70	1.04	0.19	0.33	0.33	0.41
1111	-14.66	0.89	0.23	0.24	0.28	0.37
3454	-18.24	0.86	0.29	0.00	0.24	0.30
534	-17.93	0.64	0.31	0.00	0.22	0.00
1832	-17.14	0.51	0.31	0.00	0.25	0.25
1814	-18.97	0.61	0.30	0.00	0.00	0.42
2702	-18.32	0.62	0.30	0.00	0.22	0.73
4150	-17.81	0.61	0.39	0.00	0.22	0.32
267	-19.74	0.53	0.35	0.00	0.23	0.56
2101	-17.25	0.61	0.27	0.26	0.25	0.98
1335	-18.51	0.52	0.30	0.00	0.27	0.82
3962	-18.81	0.69	0.33	0.24	0.00	0.38
3605	-16.19	0.60	0.22	0.25	0.26	0.28
1159	-18.00	0.51	0.29	0.23	0.24	0.60
1005	-17.12	0.54	0.29	0.00	0.00	0.31
344	-18.02	0.58	0.25	0.28	0.22	0.33
2108	-18.89	0.61	0.29	0.00	0.25	0.74
2699	-21.38	0.62	0.30	0.00	0.00	0.94
120	-16.57	0.64	0.33	0.20	0.25	0.26

WO 2013/148325

PCT/US2013/032180

2525	-19.16	0.61	0.32	0.23	0.22	0.79
3101	-17.11	0.55	0.32	0.24	0.22	0.24
2235	-19.14	0.47	0.31	0.23	0.00	0.97
1776	-15.63	0.97	0.30	0.23	0.24	0.00
3816	-19.02	0.58	0.28	0.00	0.25	0.29
2160	-18.47	0.59	0.33	0.25	0.26	0.85
484	-18.51	0.67	0.27	0.00	0.00	0.24
475	-16.68	0.85	0.20	0.17	0.28	0.00
3983	-16.22	0.81	0.29	0.00	0.22	0.22
3792	-19.87	0.56	0.32	0.00	0.00	0.55
1317	-17.23	0.59	0.31	0.00	0.22	0.29
3	-20.64	0.56	0.25	0.28	0.00	0.92
328	-18.73	0.67	0.27	0.25	0.26	0.23
985	-20.85	0.48	0.30	0.25	0.24	0.29
2693	-19.37	0.67	0.30	0.23	0.23	0.77
4388	-20.52	0.61	0.29	0.25	0.25	0.29
1229	-18.50	0.55	0.27	0.00	0.26	0.30
1197	-18.25	0.59	0.32	0.23	0.26	0.25
3968	-19.46	0.69	0.30	0.00	0.20	0.00
3299	-16.28	0.85	0.28	0.25	0.26	0.46
324	-18.38	0.53	0.31	0.00	0.21	0.58
357	-17.98	0.87	0.31	0.00	0.22	0.00
852	-20.90	0.46	0.30	0.29	0.35	0.23
4026	-20.21	0.52	0.31	0.00	0.25	0.00
4132	-17.41	0.75	0.29	0.00	0.00	0.26
1325	-16.76	0.77	0.33	0.00	0.00	0.29
3955	-21.48	0.47	0.30	0.19	0.00	0.00
3807	-16.59	0.79	0.30	0.27	0.27	0.32
1421	-19.03	0.60	0.29	0.00	0.25	0.34
1119	-15.92	0.69	0.31	0.00	0.25	0.34
3370	-13.90	0.84	0.30	0.28	0.25	0.34
3919	-20.41	0.50	0.33	0.00	0.22	0.24
1827	-19.36	0.47	0.31	0.00	0.28	0.00
1995	-19.08	0.50	0.30	0.00	0.00	0.78
3351	-20.75	0.61	0.15	0.35	0.31	0.35

3874	-18.32	0.55	0.22	0.00	0.00	0.00
45	-21.38	0.46	0.29	0.25	0.27	0.79
1213	-12.68	1.25	0.27	0.21	0.23	0.31
1882	-16.21	0.58	0.27	0.00	0.28	0.84
1468	-17.90	0.57	0.31	0.23	0.00	1.00

[00403] Figures 5-8 present an analysis of the essential and non-essential amino acid content of four of the identified fragments that have among the highest PCDAAS scores.

[00404] The essential and non-essential amino acid content of the polypeptide fragment from amino acid 63 to amino acid 170 of the *Arabidopsis thaliana* oligoribonuclease protein (UniProt Q9ZVE0) (SEQ ID NO: 4401) is presented in **Figure 5**. The fragment has a PCDAAS value of 2.01 and a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.041, and a Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of 0.058. This increase in distance is due to a reduced match between both the NEAA and EAA/CEAA distributions. In particular, the EAA/CEAA distribution of this polypeptide is a far better match to the 60/40 EAA/NEAA (Whey) distribution (0.015) than the 70/30 EAA/NEAA (Whey) (0.033). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (right bar) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (left bar).

[00405] The essential and non-essential amino acid content of the polypeptide fragment from amino acid 7 to amino acid 116 of the *Saccharomyces cerevisiae* probable glutamine amidotransferase protein SNO3 (UniProt P43544) (SEQ ID NO: 1275) is presented in **Figure 6**. The fragment has a PCDAAS value of 1.98 and a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.048, and a Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of 0.062. This increase in distance is due to a reduced match between both the NEAA and EAA/CEAA distributions. In particular, the EAA/CEAA distribution of this polypeptide is a far better match to the 60/40 EAA/NEAA (Whey) distribution (0.010) than the 70/30 EAA/NEAA (Whey) (0.042). The top panel presents the fraction by weight of each type/category of essential amino acid in

the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[00406] The essential and non-essential amino acid content of the polypeptide fragment from amino acid 277 to amino acid 380 of the *Arabidopsis thaliana* cadmium/zinc-transporting ATPase HMA2 protein (UniProt Q9SZW4) (SEQ ID NO: 4199) is presented in **Figure 7**. The fragment has a PCDAAS value of 2.20, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.053, and a Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of 0.0459. This decrease in distance is primarily due to an improved match between the NEAA distribution of the polypeptide fragment and the 70/30 EAA/NEAA (Whey) NEAA distribution (0.036) versus that of the 60/40 EAA/NEAA (Whey) NEAA distribution (0.044). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[00407] The essential and non-essential amino acid content of the polypeptide fragment from amino acid 234 to amino acid 340 of the *Arabidopsis thaliana* calcium transporting ATPase 3, endoplasmic reticulum-type protein (UniProt Q9SY55) (SEQ ID NO: 4190) is presented in **Figure 8**. The fragment has a PCDAAS value of 1.91, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.054, and a Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of 0.0636. This increase in distance is due to a reduced match between the EAA/CEAA distributions. In particular, the EAA/CEAA distribution of this polypeptide is a far better match to the 60/40 EAA/NEAA (Whey) distribution (0.018) than the 70/30 EAA/NEAA (Whey) (0.039). The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding

to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

Example 4: Naturally Occurring Proteins Ranked by PDCAAS Score

[00408] In this example the PDCAAS Score was used to rank proteins identified in the SwissProt database using the approach of Example 1. The 1000 proteins from the analysis with the highest PDCAAS scores are included in the Sequence Listing. The 100 proteins from the analysis with the highest PDCAAS scores were identified and are presented in **Table 9**. In **Table 9A** the proteins are identified by UniProtID number in the first column of the table. The second column indicates whether the protein contains every essential amino acid (1) or not (0). The third and fourth columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The fifth column presents the PDCAAS score for each protein. The PCDAAS scores of the full set of 1000 identified proteins range from 1.92 to 1.57. It is noteworthy that the upper bound of this PDCAAS range is less than the lower bound of the PDCAAS range of the fragments analyzed in Example 2. Additionally, the average EAA percentage for the set of 1000 full length proteins in this list, 0.51, is smaller than the average EAA percentage of the fragments presented in Examples 2, 0.63. Together, these results suggest that one can find far more fragments with high PDCAAS scores compared to full length proteins, due in part to the fact that one can obtain higher EAA fractions, which dictate the theoretical maximum PDCAAS score. Furthermore, this intimates that the majority of natural proteins are not optimized for human nutritional needs.

[00409] For the set of proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated and are presented in **Table 9B**.

[00410] Figures 9-12 present an analysis of the essential amino acid content of four of the identified proteins that have among the highest PCDAAS scores.

[00411] The essential amino acid content of the *Arabidopsis thaliana* reticulon-like protein B13 (UniProt O64837) (SEQ ID NO: 428) is presented in **Figure 9**. That protein has a PCDAAS value of 1.92. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00412] The essential amino acid content of the *Arabidopsis thaliana* mannose-1-phosphate guanylyltransferase 1 protein (UniProt O22287) (SEQ ID NO: 257) is presented in **Figure 10**. That protein has a PCDAAS value of 1.84. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00413] The essential amino acid content of the *Acipenser transmontanus* somatolactin protein (UniProt O93262) (SEQ ID NO: 524) is presented in **Figure 11**. That protein has a PCDAAS value of 1.82. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

[00414] The essential amino acid content of the *Vitis vinifera* chalcone synthase protein (UniProt P51090) (SEQ ID NO: 1391) is presented in **Figure 12**. That protein has a PCDAAS value of 1.80. The top panel presents the fraction by weight of each type/category

of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 3.6 (black bars). The bottom panel presents the relative required amino acid fraction by weight of each type/category of essential amino acid that is provided by the fragment. For reference the hatched lines provide the corresponding values for a protein having a PCDAAS score of 3.6 and for whey protein.

Table 9A

SeqIDNo	UniProtID	EAA Complete	[EAA]	[NEAA]	PDCAAS
428	O64837	1	0.61	0.39	1.92
257	O22287	1	0.53	0.47	1.84
524	O93262	1	0.55	0.45	1.82
1391	P51090	1	0.50	0.50	1.80
2475	Q58DC7	1	0.53	0.47	1.79
3986	Q9M1T7	1	0.53	0.47	1.79
2343	Q43609	1	0.52	0.48	1.79
451	O74366	1	0.51	0.49	1.79
1184	P38242	1	0.59	0.41	1.79
1834	Q08959	1	0.62	0.38	1.78
1914	Q0WRB0	1	0.54	0.46	1.78
2215	Q3EBI7	1	0.50	0.50	1.78
2714	Q5ZL79	1	0.51	0.49	1.78
338	O48706	1	0.51	0.49	1.78
1771	Q04516	1	0.53	0.47	1.77
3607	Q96514	1	0.56	0.44	1.76
577	P00812	1	0.49	0.51	1.76
2207	Q3E8L3	1	0.55	0.45	1.75
2218	Q3EBI7-2	1	0.51	0.49	1.75
362	O49859	1	0.54	0.46	1.75
1487	P58050	1	0.55	0.45	1.75
3544	Q941T9	1	0.54	0.46	1.74
3786	Q9FE17	1	0.50	0.50	1.74
1739	Q00763	1	0.54	0.46	1.73
776	P11251	1	0.56	0.44	1.73

2009	Q1PE39	1	0.51	0.49	1.73
4233	Q9UT62	1	0.60	0.40	1.73
935	P21872	1	0.49	0.51	1.73
1262	P42734-2	1	0.51	0.49	1.73
1763	Q02774	1	0.54	0.46	1.73
4161	Q9SUP0	1	0.64	0.36	1.73
255	O22046	1	0.49	0.51	1.73
258	O22652	1	0.50	0.50	1.73
1705	P93834	1	0.51	0.49	1.72
361	O49696-3	1	0.55	0.45	1.72
1522	P62597	1	0.50	0.50	1.72
4393	Q9ZU07	1	0.54	0.46	1.72
473	O80580	1	0.61	0.39	1.72
3815	Q9FKR3	1	0.52	0.48	1.72
2339	Q43188	1	0.50	0.50	1.72
1843	Q08D83-2	1	0.54	0.46	1.72
2338	Q43163	1	0.50	0.50	1.72
966	P23418	1	0.50	0.50	1.72
2319	Q41436	1	0.50	0.50	1.72
4314	Q9XGW0	1	0.53	0.47	1.72
2244	Q3T099	1	0.51	0.49	1.72
407	O59667	1	0.49	0.51	1.72
3107	Q7YQE5	1	0.60	0.40	1.71
4313	Q9XGV9	1	0.53	0.47	1.71
4170	Q9SVQ1	1	0.51	0.49	1.71
1893	Q0V7T5-2	1	0.52	0.48	1.71
4320	Q9XJ57	1	0.50	0.50	1.71
3297	Q8GWI2	1	0.54	0.46	1.71
4130	Q9SIX1	1	0.51	0.49	1.71
136	B3VKQ2	1	0.48	0.52	1.71
3838	Q9FW44	1	0.52	0.48	1.71
844	P15122	1	0.50	0.50	1.71
3812	Q9FJK7	1	0.54	0.46	1.70
1117	P33333	1	0.55	0.45	1.70
3306	Q8H1D3	1	0.48	0.52	1.70

4225	Q9US20	1	0.53	0.47	1.70
3366	Q8LG50	1	0.54	0.46	1.70
3019	Q6Z9A3	1	0.54	0.46	1.69
4394	Q9ZU29	1	0.50	0.50	1.69
2123	Q2KJ64	1	0.51	0.49	1.69
3797	Q9FH66	1	0.53	0.47	1.69
1143	P35181	1	0.55	0.45	1.69
2242	Q3SZJ4	1	0.53	0.47	1.69
2251	Q3T148	1	0.50	0.50	1.69
1446	P53780-2	1	0.49	0.51	1.69
4119	Q9SEK3	1	0.49	0.51	1.69
4105	Q9SAG4	1	0.52	0.48	1.69
4197	Q9SZK1	1	0.52	0.48	1.69
2247	Q3T0M9	1	0.55	0.45	1.69
2208	Q3E9C3-2	1	0.53	0.47	1.69
1123	P34246	1	0.50	0.50	1.68
2122	Q2KJ53	1	0.49	0.51	1.68
3947	Q9LUL5	1	0.52	0.48	1.68
2798	Q65AC2	1	0.54	0.46	1.68
1639	P80276	1	0.49	0.51	1.68
3294	Q8GW80	1	0.50	0.50	1.68
2518	Q5E9H9	1	0.52	0.48	1.68
2180	Q38860	1	0.51	0.49	1.68
581	P00937	1	0.49	0.51	1.68
4154	Q9SRE4	1	0.59	0.41	1.68
412	O60053	1	0.50	0.50	1.68
3161	Q84JH5	1	0.53	0.47	1.68
467	O77480	1	0.47	0.53	1.68
1076	P30918	1	0.64	0.31	1.57
2310	Q3ZC84	1	0.49	0.51	1.68
3960	Q9LXN3-2	1	0.54	0.46	1.68
1871	Q09914	1	0.49	0.51	1.67
2898	Q6H501	1	0.57	0.43	1.67
2149	Q2YDM2	1	0.51	0.49	1.67
1420	P53101	1	0.50	0.50	1.67

1445	P53780	1	0.49	0.51	1.67
413	O60135	1	0.48	0.52	1.67
1890	Q0V7T5	1	0.52	0.48	1.67
4321	Q9XJ58	1	0.50	0.50	1.67
2163	Q32LH7	1	0.51	0.49	1.67

Table 9B

Seq ID No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
428	-20.21	0.69	0.33	0.00	0.22	0.24
257	-19.04	0.52	0.32	0.00	0.19	0.62
524	-21.14	0.53	0.33	0.19	0.17	0.27
1391	-18.99	0.54	0.33	0.00	0.20	0.00
2475	-20.15	0.52	0.30	0.00	0.00	0.76
3986	-18.98	0.52	0.56	0.20	1.00	0.29
2343	-17.21	0.54	0.32	0.00	0.21	0.24
451	-18.29	0.53	0.31	0.00	0.00	0.35
1184	-16.47	0.79	0.28	0.22	0.21	0.30
1834	-19.79	0.62	0.31	0.00	0.00	0.22
1914	-20.93	0.51	0.35	0.20	0.23	0.27
2215	-21.17	0.52	0.33	0.00	0.21	0.00
2714	-18.88	0.56	0.33	0.00	0.20	0.55
338	-20.01	0.47	0.55	0.00	1.00	0.30
1771	-18.13	0.55	0.35	0.00	0.00	0.30
3607	-19.85	0.59	0.33	0.00	0.21	0.29
577	-18.25	0.48	0.33	0.00	0.22	0.42
2207	-20.27	0.55	0.35	0.00	0.22	0.00
2218	-20.09	0.55	0.31	0.21	0.21	0.00
362	-18.81	0.56	0.35	0.00	0.22	0.25
1487	-20.85	0.52	0.34	0.00	0.22	0.28
3544	-19.07	0.56	0.33	0.00	0.21	0.60
3786	-21.96	0.40	0.34	0.00	0.22	0.35
1739	-18.25	0.53	0.34	0.00	0.21	0.24
776	-16.18	0.60	0.33	0.23	0.23	0.57

WO 2013/148325

PCT/US2013/032180

2009	-21.15	0.53	0.33	0.00	0.22	0.23
4233	-18.29	0.78	0.27	0.00	0.24	0.00
935	-17.64	0.56	0.35	0.22	0.21	0.74
1262	-16.24	0.58	0.43	0.00	0.21	0.27
1763	-16.09	0.83	0.32	0.20	0.21	0.00
4161	-11.99	1.15	0.32	0.00	0.21	0.00
255	-19.12	0.50	0.33	0.00	0.21	0.00
258	-19.92	0.50	0.34	0.00	0.23	0.00
1705	-19.82	0.57	0.33	0.21	0.21	0.36
361	-17.63	0.82	0.31	0.00	0.00	0.00
1522	-19.40	0.39	0.32	0.00	0.21	0.49
4393	-20.32	0.52	0.33	0.18	0.21	0.28
473	-16.54	0.78	0.32	0.00	0.22	0.00
3815	-20.76	0.45	0.36	0.00	0.20	0.20
2339	-18.85	0.49	0.34	0.00	0.21	0.00
1843	-14.34	0.85	0.32	0.22	0.24	0.96
2338	-19.13	0.52	0.33	0.19	0.21	0.00
966	-19.07	0.53	0.32	0.20	0.21	0.00
2319	-18.97	0.53	0.33	0.00	0.21	0.00
4314	-17.75	0.50	0.32	0.00	0.21	0.24
2244	-20.12	0.45	0.31	0.23	0.20	0.89
407	-21.22	0.44	0.33	0.22	0.21	0.00
3107	-14.14	0.65	0.30	0.20	0.22	0.40
4313	-17.88	0.51	0.33	0.00	0.20	0.24
4170	-19.83	0.46	0.31	0.00	0.22	0.26
1893	-18.40	0.58	0.46	0.00	0.20	0.29
4320	-19.20	0.49	0.33	0.19	0.22	0.00
3297	-19.91	0.49	0.33	0.21	0.22	0.21
4130	-18.00	0.51	0.33	0.19	0.20	0.00
136	-17.63	0.46	0.34	0.19	0.20	0.96
3838	-22.41	0.48	0.34	0.00	0.21	0.00
844	-19.96	0.38	0.32	0.00	0.00	0.89
3812	-21.57	0.48	0.31	0.22	0.00	0.39
1117	-17.54	0.59	0.35	0.00	0.21	0.28
3306	-23.23	0.40	0.35	0.20	0.21	0.00

WO 2013/148325

PCT/US2013/032180

4225	-17.22	0.69	0.30	0.20	0.21	0.31
3366	-16.91	0.71	0.32	0.00	0.23	0.33
3019	-19.20	0.55	0.32	0.00	0.21	0.60
4394	-21.05	0.45	0.33	0.20	0.21	0.19
2123	-20.08	0.43	0.33	0.00	0.20	0.91
3797	-19.97	0.52	0.33	0.21	0.21	0.26
1143	-22.02	0.48	0.32	0.22	0.00	0.54
2242	-18.24	0.53	0.36	0.00	0.24	0.80
2251	-22.36	0.50	0.31	0.00	0.20	1.00
1446	-18.31	0.48	0.33	0.00	0.22	0.37
4119	-20.69	0.52	0.33	0.22	0.22	0.37
4105	-19.69	0.50	0.34	0.00	0.21	0.00
4197	-18.54	0.56	0.35	0.00	0.21	0.00
2247	-20.96	0.45	0.31	0.00	0.21	1.00
2208	-20.17	0.60	0.33	0.00	0.00	0.27
1123	-19.77	0.42	0.31	0.00	0.00	0.00
2122	-18.77	0.52	0.33	0.00	0.23	0.96
3947	-19.63	0.48	0.32	0.00	0.20	0.00
2798	-15.64	0.77	0.33	0.00	0.20	0.89
1639	-21.12	0.38	0.32	0.00	0.20	0.87
3294	-19.60	0.45	0.34	0.18	0.21	0.00
2518	-18.85	0.55	0.34	0.00	0.00	0.85
2180	-17.85	0.55	0.32	0.00	0.20	0.00
581	-19.17	0.46	0.31	0.21	0.22	0.22
4154	-12.89	0.92	0.33	0.00	0.19	0.24
412	-18.68	0.41	0.34	0.00	0.21	0.00
3161	-18.80	0.58	0.33	0.00	0.22	0.61
467	-18.33	0.45	0.32	0.00	0.21	0.86
1076	-6.37	0.82	0.17	0.00	0.35	0.47
2310	-21.15	0.42	0.32	0.00	0.23	0.91
3960	-18.32	0.48	0.34	0.00	0.22	0.28
1871	-22.40	0.53	0.30	0.00	0.00	0.67
2898	-14.96	0.77	0.35	0.00	0.21	0.00
2149	-19.92	0.43	0.31	0.00	0.21	0.97
1420	-18.87	0.46	0.34	0.20	0.21	0.31

1445	-18.10	0.46	0.31	0.00	0.22	0.36
413	-20.75	0.40	0.33	0.00	0.21	0.35
1890	-17.86	0.62	0.46	0.00	0.21	0.27
4321	-19.47	0.50	0.33	0.00	0.21	0.00
2163	-21.02	0.40	0.31	0.21	0.21	0.88

Example 5: Naturally Occurring Proteins Ranked by Euclidian Distance from a 60/40 EAA/NEAA Profile and by Euclidian Distance from a 70/30 EAA/NEAA Profile

[00415] In this example the Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile and the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile were used to rank proteins identified in the SwissProt database using the approach of Example 1. After proteins were identified in this way the PDCAAS of each protein was also determined. The 1000 proteins from the analysis with the smallest Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile are included in the Sequence Listing. The 100 proteins from the analysis with the smallest Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile are presented in **Table 10**. The 1000 proteins from the analysis with the smallest Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile are included in the Sequence Listing. The 100 proteins from the analysis with the smallest Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile are presented in **Table 11**. In **Tables 10A** and **11A** the proteins were first ranked by PDCAAS score (to the nearest tenth) and then ranked by Euclidian distance from the target profile. The proteins are identified by UniProtID number in the first column of the tables. The second column indicates whether the fragment contains every essential amino acid (1) or not (0). The third and fourth columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The fifth column presents the PDCAAS score for each protein. The seventh column presents the distance from a 60/40 or 70/30 EAA/NEAA (Whey) target amino acid profile for each protein. The Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of the 1000 fragments ranges from 0.04834 to 0.0716. The Euclidian distance from a 70/30

EAA/NEAA (Whey) target amino acid profile of the fragments ranges from 0.0590 to 0.0936.

[00416] For each set of 100 proteins the solvation score at pH 7 (“SolvScore”), aggregation score at pH 7 (“AggScore”), allergenicity (i.e., percent local homology to known allergens, as described herein), toxicity (i.e., percent homology to known toxins, as described herein), anti-nutricity (i.e., percent homology to known protease inhibitors, as described herein), and human homology (i.e., percent homology to known human proteins, as described herein) were calculated and are presented in **Tables 10B and 11B**.

Table 10A

SeqIDNo	UniProtID	EAA Complete	[EAA]	[NEAA]	PDCAAS	Dist. Target
1391	P51090	1	0.50	0.50	1.80	0.0593
2343	Q43609	1	0.52	0.48	1.79	0.0610
2714	Q5ZL79	1	0.51	0.49	1.78	0.0626
362	O49859	1	0.54	0.46	1.75	0.0639
577	P00812	1	0.49	0.51	1.76	0.0652
935	P21872	1	0.49	0.51	1.73	0.0549
4313	Q9XGV9	1	0.53	0.47	1.71	0.0561
4320	Q9XJ57	1	0.50	0.50	1.71	0.0596
2242	Q3SZJ4	1	0.53	0.47	1.69	0.0624
255	O22046	1	0.49	0.51	1.73	0.0624
4314	Q9XGW0	1	0.53	0.47	1.72	0.0629
2234	Q3MHL4	1	0.50	0.50	1.65	0.0633
1362	P49440	1	0.50	0.50	1.66	0.0646
2339	Q43188	1	0.50	0.50	1.72	0.0647
2251	Q3T148	1	0.50	0.50	1.69	0.0649
1739	Q00763	1	0.54	0.46	1.73	0.0654
1380	P51071	1	0.50	0.50	1.66	0.0656
256	O22047	1	0.49	0.51	1.67	0.0659
844	P15122	1	0.50	0.50	1.71	0.0659
3100	Q7XRA1	1	0.50	0.50	1.65	0.0668
2338	Q43163	1	0.50	0.50	1.72	0.0670
3998	Q9MB33	1	0.50	0.50	1.67	0.0675

2319	Q41436	1	0.50	0.50	1.72	0.0678
4000	Q9MB37	1	0.50	0.50	1.67	0.0679
966	P23418	1	0.50	0.50	1.72	0.0687
1839	Q08975	1	0.47	0.53	1.66	0.0704
407	O59667	1	0.49	0.51	1.72	0.0708
3507	Q92093	1	0.49	0.51	1.67	0.0710
1871	Q09914	1	0.49	0.51	1.67	0.0713
4321	Q9XJ58	1	0.50	0.50	1.67	0.0715
2884	Q6EE31	1	0.49	0.51	1.57	0.0579
4198	Q9SZW4	1	0.50	0.50	1.56	0.0580
639	P04161	1	0.48	0.52	1.59	0.0585
2796	Q652P4	1	0.49	0.51	1.62	0.0592
544	O94564	1	0.51	0.49	1.59	0.0592
2313	Q3ZCI9	1	0.49	0.51	1.56	0.0600
221	O13639	1	0.51	0.49	1.58	0.0616
2330	Q42896	1	0.52	0.48	1.58	0.0617
1221	P39954	1	0.50	0.50	1.60	0.0622
1046	P28475	1	0.52	0.48	1.60	0.0626
2690	Q5ZI67	1	0.49	0.51	1.63	0.0628
814	P13491	1	0.55	0.45	1.56	0.0633
3813	Q9FK25	1	0.53	0.47	1.60	0.0641
2324	Q42586	1	0.52	0.48	1.59	0.0642
683	P07246	1	0.51	0.49	1.55	0.0642
3929	Q9LMU0	1	0.51	0.49	1.56	0.0645
3657	Q9BEC7	1	0.51	0.49	1.56	0.0646
2112	Q29RY4	1	0.51	0.49	1.55	0.0646
2495	Q59A28	1	0.50	0.50	1.56	0.0651
3097	Q7XJ81	1	0.51	0.49	1.58	0.0652
259	O22711	1	0.55	0.45	1.56	0.0654
2183	Q39069	1	0.49	0.51	1.58	0.0658
3540	Q93WU3	1	0.54	0.46	1.57	0.0659
365	O50008	1	0.48	0.52	1.61	0.0660
602	P02103	1	0.58	0.42	1.62	0.0661
4004	Q9MB41	1	0.49	0.51	1.60	0.0664
670	P06169	1	0.51	0.49	1.57	0.0665

4001	Q9MB38	1	0.49	0.51	1.60	0.0672
4003	Q9MB40	1	0.49	0.51	1.56	0.0675
2701	Q5ZJ64	1	0.51	0.49	1.59	0.0676
3923	Q9LK36	1	0.50	0.50	1.55	0.0678
1678	P87241	1	0.50	0.50	1.60	0.0681
266	O23255	1	0.50	0.50	1.58	0.0684
232	O14075	1	0.50	0.50	1.59	0.0685
1379	P51070	1	0.50	0.50	1.60	0.0686
1740	Q00775	1	0.51	0.49	1.64	0.0689
3539	Q93WU2	1	0.54	0.46	1.62	0.0689
3999	Q9MB36	1	0.49	0.51	1.55	0.0691
711	P08566	1	0.50	0.50	1.60	0.0691
4155	Q9SRZ4	1	0.55	0.45	1.57	0.0695
3872	Q9HE04	1	0.51	0.49	1.63	0.0695
3403	Q8RY79	1	0.48	0.52	1.57	0.0696
224	O13848	1	0.53	0.47	1.64	0.0699
114	A8E657	1	0.48	0.52	1.62	0.0699
1095	P31754	1	0.48	0.52	1.58	0.0704
1997	Q1KYT0	1	0.48	0.52	1.55	0.0705
1072	P30624	1	0.51	0.49	1.55	0.0706
1753	Q01781	1	0.49	0.51	1.55	0.0711
1017	P26690	1	0.50	0.50	1.57	0.0711
1176	P37829	1	0.51	0.49	1.56	0.0712
928	P21528	1	0.46	0.54	1.58	0.0716
479	O80944-3	1	0.52	0.48	1.51	0.0590
478	O80944-2	1	0.51	0.49	1.51	0.0611
233	O14134	1	0.48	0.52	1.51	0.0618
1747	Q01288	1	0.50	0.50	1.50	0.0636
703	P07991	1	0.51	0.49	1.54	0.0639
1217	P39676	1	0.51	0.49	1.55	0.0650
2527	Q5F418	1	0.47	0.53	1.46	0.0650
537	O94404	1	0.48	0.52	1.53	0.0652
1706	P93848	1	0.55	0.45	1.55	0.0653
477	O80944	1	0.52	0.48	1.47	0.0653
2496	Q59A32	1	0.48	0.52	1.53	0.0665

4176	Q9SWF5	1	0.51	0.49	1.55	0.0669
3315	Q8HYR6	1	0.54	0.46	1.50	0.0672
2522	Q5EA88	1	0.52	0.48	1.48	0.0673
4192	Q9SYZ7	1	0.52	0.48	1.52	0.0675
254	O22045	1	0.49	0.51	1.49	0.0680
565	P00337	1	0.53	0.47	1.48	0.0682
2530	Q5FJB3	1	0.51	0.49	1.22	0.0547
3830	Q9FR35	1	0.52	0.48	0.96	0.0553

Table 10B

Seq ID						Human
No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Homology
1391	-18.99	0.54	0.33	0.00	0.20	0.00
2343	-17.21	0.54	0.32	0.00	0.21	0.24
2714	-18.88	0.56	0.33	0.00	0.20	0.55
362	-18.81	0.56	0.35	0.00	0.22	0.25
577	-18.25	0.48	0.33	0.00	0.22	0.42
935	-17.64	0.56	0.35	0.22	0.21	0.74
4313	-17.88	0.51	0.33	0.00	0.20	0.24
4320	-19.20	0.49	0.33	0.19	0.22	0.00
2242	-18.24	0.53	0.36	0.00	0.24	0.80
255	-19.12	0.50	0.33	0.00	0.21	0.00
4314	-17.75	0.50	0.32	0.00	0.21	0.24
2234	-19.29	0.45	0.32	0.19	0.21	0.97
1362	-18.79	0.52	0.33	0.22	0.19	0.00
2339	-18.85	0.49	0.34	0.00	0.21	0.00
2251	-22.36	0.50	0.31	0.00	0.20	1.00
1739	-18.25	0.53	0.34	0.00	0.21	0.24
1380	-18.48	0.51	0.33	0.00	0.21	0.00
256	-19.07	0.50	0.32	0.00	0.21	0.00
844	-19.96	0.38	0.32	0.00	0.00	0.89
3100	-17.83	0.44	0.32	0.20	0.21	0.00
2338	-19.13	0.52	0.33	0.19	0.21	0.00
3998	-17.96	0.51	0.34	0.00	0.20	0.00

WO 2013/148325

PCT/US2013/032180

2319	-18.97	0.53	0.33	0.00	0.21	0.00
4000	-18.15	0.50	0.34	0.19	0.20	0.00
966	-19.07	0.53	0.32	0.20	0.21	0.00
1839	-19.71	0.41	0.33	0.00	0.21	0.00
407	-21.22	0.44	0.33	0.22	0.21	0.00
3507	-18.40	0.50	0.58	0.19	0.19	0.19
1871	-22.40	0.53	0.30	0.00	0.00	0.67
4321	-19.47	0.50	0.33	0.00	0.21	0.00
2884	-20.22	0.47	0.32	0.20	0.22	0.93
4198	-18.06	0.63	0.33	0.19	0.20	0.25
639	-19.93	0.57	0.31	0.24	0.24	0.34
2796	-19.96	0.46	0.35	0.00	0.21	0.37
544	-16.66	0.53	0.39	0.00	0.19	0.29
2313	-20.38	0.47	0.33	0.00	0.21	0.98
221	-19.46	0.51	0.32	0.20	0.22	0.77
2330	-17.96	0.56	0.33	0.00	0.21	0.26
1221	-18.95	0.51	0.36	0.00	0.21	0.70
1046	-20.92	0.40	0.34	0.00	0.20	0.40
2690	-18.40	0.48	0.34	0.00	0.20	0.75
814	-18.58	0.57	0.33	0.00	0.20	0.94
3813	-18.96	0.51	0.33	0.00	0.21	0.24
2324	-18.96	0.58	0.35	0.00	0.22	0.54
683	-18.18	0.46	0.86	0.00	0.20	0.25
3929	-17.75	0.43	0.31	0.00	0.22	0.34
3657	-17.48	0.55	0.34	0.00	0.21	0.92
2112	-20.30	0.48	0.35	0.18	0.20	0.94
2495	-20.70	0.49	0.31	0.00	0.24	0.94
3097	-17.73	0.57	0.34	0.00	0.22	0.26
259	-19.19	0.57	0.49	0.24	0.23	0.40
2183	-20.81	0.47	0.32	0.00	0.21	0.33
3540	-18.17	0.51	0.34	0.00	0.21	0.21
365	-19.90	0.42	0.95	0.00	0.22	0.00
602	-17.93	0.52	0.29	0.24	0.23	0.86
4004	-17.83	0.52	0.34	0.00	0.20	0.00
670	-16.92	0.48	0.33	0.00	0.22	0.22

WO 2013/148325

PCT/US2013/032180

4001	-17.81	0.52	0.34	0.00	0.21	0.20
4003	-17.78	0.51	0.34	0.20	0.22	0.00
2701	-22.19	0.51	0.32	0.00	0.20	0.96
3923	-19.50	0.50	0.33	0.20	0.21	0.55
1678	-20.35	0.47	0.32	0.00	0.19	0.00
266	-19.76	0.50	0.33	0.21	0.21	0.55
232	-20.63	0.47	0.32	0.21	0.00	0.29
1379	-18.20	0.51	0.33	0.00	0.22	0.00
1740	-18.49	0.50	0.33	0.00	0.22	0.00
3539	-18.39	0.50	0.32	0.00	0.21	0.22
3999	-17.81	0.52	0.34	0.00	0.21	0.00
711	-19.14	0.46	0.33	0.19	0.19	0.00
4155	-19.10	0.57	0.49	0.24	0.23	0.40
3872	-21.84	0.52	0.31	0.22	0.24	0.65
3403	-18.35	0.51	0.35	0.00	0.22	0.48
224	-20.40	0.40	0.31	0.00	0.00	0.33
114	-18.77	0.44	0.34	0.00	0.20	0.90
1095	-19.18	0.55	0.32	0.00	0.22	0.84
1997	-20.83	0.43	0.99	0.21	0.22	0.97
1072	-19.81	0.43	0.33	0.00	0.20	0.31
1753	-20.01	0.48	0.32	0.00	0.21	0.56
1017	-19.84	0.41	0.31	0.22	0.00	0.39
1176	-18.41	0.57	0.33	0.00	0.21	0.26
928	-18.74	0.45	0.33	0.18	0.22	0.35
479	-18.99	0.32	0.31	0.00	0.23	0.43
478	-19.19	0.32	0.31	0.00	0.22	0.43
233	-20.71	0.46	0.33	0.20	0.19	0.23
1747	-19.27	0.51	0.33	0.19	0.21	0.00
703	-17.28	0.56	0.31	0.00	0.21	0.52
1217	-19.88	0.43	0.32	0.00	0.19	0.00
2527	-20.93	0.52	0.36	0.00	0.20	0.97
537	-17.12	0.48	0.33	0.00	0.21	0.32
1706	-18.13	0.52	0.28	0.00	0.23	0.25
477	-20.15	0.30	0.31	0.00	0.24	0.42
2496	-18.27	0.53	0.36	0.20	0.22	0.88

4176	-19.62	0.48	0.31	0.00	0.22	0.55
3315	-18.36	0.60	0.33	0.23	0.00	0.87
2522	-17.51	0.57	0.31	0.00	0.20	0.93
4192	-22.14	0.51	0.33	0.00	0.00	0.20
254	-18.06	0.51	0.34	0.00	0.22	0.00
565	-17.65	0.58	0.31	0.00	0.18	0.90
2530	-19.87	0.47	0.33	0.20	0.00	0.23
3830	-18.09	0.52	0.46	0.00	0.00	0.43

Table 11A

SeqIDNo	UniProtID	EAA Complete	[EAA]	[NEAA]	PDCAAS	Dist. Target
1834	Q08959	1	0.62	0.38	1.78	0.0829
362	O49859	1	0.54	0.46	1.75	0.0830
2714	Q5ZL79	1	0.51	0.49	1.78	0.0835
2343	Q43609	1	0.52	0.48	1.79	0.0837
3607	Q96514	1	0.56	0.44	1.76	0.0837
2207	Q3E8L3	1	0.55	0.45	1.75	0.0861
1391	P51090	1	0.50	0.50	1.80	0.0909
3986	Q9M1T7	1	0.53	0.47	1.79	0.0915
1184	P38242	1	0.59	0.41	1.79	0.0919
1914	Q0WRB0	1	0.54	0.46	1.78	0.0926
577	P00812	1	0.49	0.51	1.76	0.0929
4313	Q9XGV9	1	0.53	0.47	1.71	0.0787
1117	P33333	1	0.55	0.45	1.70	0.0809
2242	Q3SZJ4	1	0.53	0.47	1.69	0.0824
473	O80580	1	0.61	0.39	1.72	0.0829
1739	Q00763	1	0.54	0.46	1.73	0.0838
4314	Q9XGW0	1	0.53	0.47	1.72	0.0849
2798	Q65AC2	1	0.54	0.46	1.68	0.0857
3960	Q9LXN3-2	1	0.54	0.46	1.68	0.0885
844	P15122	1	0.50	0.50	1.71	0.0886
3297	Q8GWI2	1	0.54	0.46	1.71	0.0886
361	O49696-3	1	0.55	0.45	1.72	0.0886

4233	Q9UT62	1	0.60	0.40	1.73	0.0889
935	P21872	1	0.49	0.51	1.73	0.0898
1763	Q02774	1	0.54	0.46	1.73	0.0904
2251	Q3T148	1	0.50	0.50	1.69	0.0905
4320	Q9XJ57	1	0.50	0.50	1.71	0.0906
2234	Q3MHL4	1	0.50	0.50	1.65	0.0914
1843	Q08D83-2	1	0.54	0.46	1.72	0.0915
3100	Q7XRA1	1	0.50	0.50	1.65	0.0916
4161	Q9SUP0	1	0.64	0.36	1.73	0.0929
1362	P49440	1	0.50	0.50	1.66	0.0929
1890	Q0V7T5	1	0.52	0.48	1.67	0.0936
4386	Q9ZQR8-2	1	0.57	0.43	1.56	0.0768
602	P02103	1	0.58	0.42	1.62	0.0799
814	P13491	1	0.55	0.45	1.56	0.0804
1046	P28475	1	0.52	0.48	1.60	0.0806
3959	Q9LXN3	1	0.55	0.45	1.63	0.0842
3813	Q9FK25	1	0.53	0.47	1.60	0.0847
259	O22711	1	0.55	0.45	1.56	0.0865
544	O94564	1	0.51	0.49	1.59	0.0865
683	P07246	1	0.51	0.49	1.55	0.0871
3872	Q9HE04	1	0.51	0.49	1.63	0.0877
2324	Q42586	1	0.52	0.48	1.59	0.0886
2796	Q652P4	1	0.49	0.51	1.62	0.0887
4155	Q9SRZ4	1	0.55	0.45	1.57	0.0890
2701	Q5ZJ64	1	0.51	0.49	1.59	0.0891
1453	P53970	1	0.54	0.46	1.56	0.0895
224	O13848	1	0.53	0.47	1.64	0.0899
4198	Q9SZW4	1	0.50	0.50	1.56	0.0900
3657	Q9BEC7	1	0.51	0.49	1.56	0.0901
2183	Q39069	1	0.49	0.51	1.58	0.0901
3665	Q9C1X2	1	0.50	0.50	1.58	0.0902
356	O49621	1	0.58	0.42	1.64	0.0902
2112	Q29RY4	1	0.51	0.49	1.55	0.0905
1072	P30624	1	0.51	0.49	1.55	0.0907
1740	Q00775	1	0.51	0.49	1.64	0.0910

2330	Q42896	1	0.52	0.48	1.58	0.0912
1678	P87241	1	0.50	0.50	1.60	0.0919
3539	Q93WU2	1	0.54	0.46	1.62	0.0920
607	P02112	1	0.56	0.44	1.58	0.0921
3453	Q8VZI2	1	0.54	0.46	1.57	0.0921
3540	Q93WU3	1	0.54	0.46	1.57	0.0921
608	P02114	1	0.56	0.44	1.58	0.0922
2690	Q5ZI67	1	0.49	0.51	1.63	0.0923
221	O13639	1	0.51	0.49	1.58	0.0926
3942	Q9LT02	1	0.53	0.47	1.60	0.0927
3929	Q9LMU0	1	0.51	0.49	1.56	0.0928
3646	Q9ATM1	1	0.59	0.41	1.55	0.0929
1305	P46202	1	0.58	0.42	1.57	0.0930
1221	P39954	1	0.50	0.50	1.60	0.0933
1470	P54781	1	0.53	0.47	1.56	0.0934
2495	Q59A28	1	0.50	0.50	1.56	0.0935
2884	Q6EE31	1	0.49	0.51	1.57	0.0935
1107	P32564	1	0.53	0.47	1.52	0.0783
3315	Q8HYR6	1	0.54	0.46	1.50	0.0805
479	O80944-3	1	0.52	0.48	1.51	0.0835
565	P00337	1	0.53	0.47	1.48	0.0844
478	O80944-2	1	0.51	0.49	1.51	0.0869
272	O23413	1	0.62	0.38	1.53	0.0870
2309	Q3ZC52	1	0.55	0.45	1.51	0.0870
2103	Q29563	1	0.54	0.46	1.46	0.0872
3461	Q8W245	1	0.58	0.42	1.51	0.0879
30	A2PZE5	1	0.62	0.38	1.49	0.0881
2485	Q58P71	1	0.56	0.44	1.47	0.0882
2246	Q3T0M0	1	0.54	0.46	1.48	0.0889
227	O13950	1	0.58	0.42	1.48	0.0891
3920	Q9LIK7	1	0.54	0.46	1.54	0.0896
765	P0DH60	1	0.55	0.45	1.50	0.0905
4309	Q9XEX2	1	0.54	0.46	1.52	0.0906
477	O80944	1	0.52	0.48	1.47	0.0910
1004	P25847	1	0.51	0.49	1.52	0.0911

2695	Q5ZIL2	1	0.54	0.46	1.48	0.0911
532	O94265	1	0.52	0.48	1.53	0.0913
1440	P53727	1	0.52	0.49	1.47	0.0913
3357	Q8L7M4	1	0.51	0.49	1.46	0.0915
4022	Q9P7C9	1	0.55	0.45	1.46	0.0918
1224	P39988	1	0.51	0.49	1.54	0.0923
703	P07991	1	0.51	0.49	1.54	0.0925
1747	Q01288	1	0.50	0.50	1.50	0.0927

Table 11B

Seq ID						Human
No	SolvScore	AggScore	Allergenicity	Toxicity	Antinutricity	Homology
1834	-19.79	0.62	0.31	0.00	0.00	0.22
362	-18.81	0.56	0.35	0.00	0.22	0.25
2714	-18.88	0.56	0.33	0.00	0.20	0.55
2343	-17.21	0.54	0.32	0.00	0.21	0.24
3607	-19.85	0.59	0.33	0.00	0.21	0.29
2207	-20.27	0.55	0.35	0.00	0.22	0.00
1391	-18.99	0.54	0.33	0.00	0.20	0.00
3986	-18.98	0.52	0.56	0.20	1.00	0.29
1184	-16.47	0.79	0.28	0.22	0.21	0.30
1914	-20.93	0.51	0.35	0.20	0.23	0.27
577	-18.25	0.48	0.33	0.00	0.22	0.42
4313	-17.88	0.51	0.33	0.00	0.20	0.24
1117	-17.54	0.59	0.35	0.00	0.21	0.28
2242	-18.24	0.53	0.36	0.00	0.24	0.80
473	-16.54	0.78	0.32	0.00	0.22	0.00
1739	-18.25	0.53	0.34	0.00	0.21	0.24
4314	-17.75	0.50	0.32	0.00	0.21	0.24
2798	-15.64	0.77	0.33	0.00	0.20	0.89
3960	-18.32	0.48	0.34	0.00	0.22	0.28
844	-19.96	0.38	0.32	0.00	0.00	0.89
3297	-19.91	0.49	0.33	0.21	0.22	0.21
361	-17.63	0.82	0.31	0.00	0.00	0.00

WO 2013/148325

PCT/US2013/032180

4233	-18.29	0.78	0.27	0.00	0.24	0.00
935	-17.64	0.56	0.35	0.22	0.21	0.74
1763	-16.09	0.83	0.32	0.20	0.21	0.00
2251	-22.36	0.50	0.31	0.00	0.20	1.00
4320	-19.20	0.49	0.33	0.19	0.22	0.00
2234	-19.29	0.45	0.32	0.19	0.21	0.97
1843	-14.34	0.85	0.32	0.22	0.24	0.96
3100	-17.83	0.44	0.32	0.20	0.21	0.00
4161	-11.99	1.15	0.32	0.00	0.21	0.00
1362	-18.79	0.52	0.33	0.22	0.19	0.00
1890	-17.86	0.62	0.46	0.00	0.21	0.27
4386	-18.04	0.67	0.30	0.00	0.23	0.00
602	-17.93	0.52	0.29	0.24	0.23	0.86
814	-18.58	0.57	0.33	0.00	0.20	0.94
1046	-20.92	0.40	0.34	0.00	0.20	0.40
3959	-20.03	0.48	0.34	0.00	0.22	0.27
3813	-18.96	0.51	0.33	0.00	0.21	0.24
259	-19.19	0.57	0.49	0.24	0.23	0.40
544	-16.66	0.53	0.39	0.00	0.19	0.29
683	-18.18	0.46	0.86	0.00	0.20	0.25
3872	-21.84	0.52	0.31	0.22	0.24	0.65
2324	-18.96	0.58	0.35	0.00	0.22	0.54
2796	-19.96	0.46	0.35	0.00	0.21	0.37
4155	-19.10	0.57	0.49	0.24	0.23	0.40
2701	-22.19	0.51	0.32	0.00	0.20	0.96
1453	-22.41	0.48	0.34	0.24	0.00	0.31
224	-20.40	0.40	0.31	0.00	0.00	0.33
4198	-18.06	0.63	0.33	0.19	0.20	0.25
3657	-17.48	0.55	0.34	0.00	0.21	0.92
2183	-20.81	0.47	0.32	0.00	0.21	0.33
3665	-20.13	0.45	0.34	0.22	0.22	0.46
356	-16.25	0.75	0.32	0.18	0.20	0.00
2112	-20.30	0.48	0.35	0.18	0.20	0.94
1072	-19.81	0.43	0.33	0.00	0.20	0.31
1740	-18.49	0.50	0.33	0.00	0.22	0.00

WO 2013/148325

PCT/US2013/032180

2330	-17.96	0.56	0.33	0.00	0.21	0.26
1678	-20.35	0.47	0.32	0.00	0.19	0.00
3539	-18.39	0.50	0.32	0.00	0.21	0.22
607	-17.37	0.55	0.32	0.21	0.22	0.75
3453	-19.14	0.77	0.33	0.19	0.21	0.32
3540	-18.17	0.51	0.34	0.00	0.21	0.21
608	-17.34	0.56	0.34	0.00	0.23	0.75
2690	-18.40	0.48	0.34	0.00	0.20	0.75
221	-19.46	0.51	0.32	0.20	0.22	0.77
3942	-18.32	0.61	0.33	0.18	0.20	0.43
3929	-17.75	0.43	0.31	0.00	0.22	0.34
3646	-13.05	0.87	0.33	0.24	0.23	0.27
1305	-24.17	0.49	0.33	0.00	1.00	0.32
1221	-18.95	0.51	0.36	0.00	0.21	0.70
1470	-19.35	0.56	0.32	0.19	0.21	0.24
2495	-20.70	0.49	0.31	0.00	0.24	0.94
2884	-20.22	0.47	0.32	0.20	0.22	0.93
1107	-14.06	0.74	0.31	0.00	0.24	0.00
3315	-18.36	0.60	0.33	0.23	0.00	0.87
479	-18.99	0.32	0.31	0.00	0.23	0.43
565	-17.65	0.58	0.31	0.00	0.18	0.90
478	-19.19	0.32	0.31	0.00	0.22	0.43
272	-13.56	1.09	0.31	0.00	0.21	0.00
2309	-20.15	0.47	0.32	0.00	0.23	0.96
2103	-18.59	0.60	0.32	0.00	0.20	0.89
3461	-12.98	0.92	0.33	0.23	0.22	0.22
30	-10.82	1.16	0.33	0.00	0.24	0.23
2485	-16.66	0.85	0.34	0.00	0.20	0.00
2246	-18.26	0.49	0.30	0.00	0.22	0.99
227	-19.78	0.61	0.33	0.00	0.00	0.27
3920	-19.75	0.64	0.34	0.19	0.19	0.36
765	-20.85	0.45	0.32	0.00	0.00	0.26
4309	-20.40	0.56	0.51	0.20	0.24	0.41
477	-20.15	0.30	0.31	0.00	0.24	0.42
1004	-21.93	0.45	0.33	0.19	0.20	0.40

2695	-18.37	0.49	0.30	0.00	0.22	0.97
532	-16.27	0.56	0.31	0.19	0.23	0.00
1440	-18.42	0.47	0.31	0.00	0.21	0.35
3357	-21.25	0.45	0.32	0.00	0.22	0.25
4022	-21.51	0.39	0.33	0.00	0.00	0.37
1224	-19.88	0.40	0.32	0.00	0.00	0.35
703	-17.28	0.56	0.31	0.00	0.21	0.52
1747	-19.27	0.51	0.33	0.19	0.21	0.00

[00417] The essential and non-essential amino acid content of the *Lactobacillus acidophilus* orotidine 5'-phosphate decarboxylase protein (UniProt Q5FJB3) (SEQ ID NO: 2530) is presented in **Figure 13**. The protein has a PCDAAS value of 1.22, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0547, and a Euclidian distance from a 70/30 EAA/NEAA (Whey) target amino acid profile of 0.0849. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[00418] The essential and non-essential amino acid content of the *Gallus gallus* trifunctional purine biosynthetic protein adenosine-3 (UniProt P21872) (SEQ ID NO: 935) is presented in **Figure 14**. The protein has a PCDAAS value of 1.73, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0549, and a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0898. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[00419] The essential and non-essential amino acid content of the *Oryza sativa subsp. japonica* Peroxiredoxin-2C protein (UniProt Q9FR35) (SEQ ID NO: 3830) is presented in **Figure 15**. The protein has a PCDAAS value of 0.96, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0553, and a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0851. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

[00420] The essential and non-essential amino acid content of the *Ocimum basillicum* caffeic acid 3-O-methyltransferase 2 protein (UniProt Q9XGV9) (SEQ ID NO: 4313) is presented in **Figure 16**. The protein has a PCDAAS value of 1.71, a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0561, and a Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile of 0.0787. The top panel presents the fraction by weight of each type/category of essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type/category of essential amino acid corresponding to the relative required amino acid fraction corresponding to a PCDAAS score of 2.2 (black bars). The bottom panel presents the fraction by weight of each type of non-essential amino acid in the fragment (white bars) and compares it to the fraction by weight of each type of non-essential amino acid present in whey protein (black bars).

Example 6: Analysis of Whey, Casein, Egg, and Soy Proteins and Protein Fragments

[00421] For comparison to the proteins and protein fragments identified and analyzed in Examples 2-5, abundant whey, casein, egg, and soy proteins were analyzed for essential amino acid content, non-essential amino acid content, and either PDCAAS value or Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile.

[00422] **Table 12** presents whole proteins ranked by PDCAAS value. The proteins are identified by UniProtID number in the first column of the table, their name in the second column, and the source of the protein in the third column. The fourth column indicates whether the protein contains every essential amino acid (1) or not (0). The fifth and sixth columns present the weight fraction of essential amino acids (EAA), and the total of non-

essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The seventh column presents the PDCAAS score for each protein. The proteins are ranked from highest to lowest PDCAAS value.

Table 12

SEQID	UniProtID	Name	Source	EAA		[NEAA]	PDCAAS
				Complete	[EAA]		
585	P01012	Ovalbumin	Egg	1	0.47	0.53	1.33
633	P02789	Ovotransferrin	Egg	1	0.45	0.55	1.31
574	P00711	Alpha-lactalbumin	Whey	1	0.53	0.47	1.25
623	P02666	Beta-casein	Casein	1	0.48	0.52	1.24
625	P02668	Kappa-casein	Casein	1	0.44	0.56	1.17
655	P04776	Glycinin G1	Soy	1	0.38	0.62	1.10
619	P02662	Alpha-S1-casein	Casein	1	0.44	0.56	1.08
621	P02663	Alpha-S2-casein	Casein	1	0.47	0.53	1.05
791	P11828	Glycinin G3	Soy	1	0.37	0.63	1.00
646	P04347	Glycinin	Soy	1	0.36	0.64	0.93
627	P02754	Beta-lactoglobulin	Whey	1	0.50	0.50	0.92
635	P02858	Glycinin G4	Soy	1	0.35	0.65	0.87
631	P02769	Serum albumin	Whey	1	0.46	0.54	0.82
789	P11827	Beta-conglycinin alpha prime chain	Soy	1	0.37	0.63	0.80
648	P04405	Glycinin G2	Soy	1	0.36	0.64	0.67
822	P13916	Beta-conglycinin alpha chain	Soy	1	0.34	0.66	0.59
1588	P68390	Ovomucoid	Egg	0	0.38	0.62	0.00

1010	P25974	Beta- conglycinin beta chain	Soy	0	0.41	0.59	0.00
------	--------	------------------------------------	-----	---	------	------	------

[00423] **Table 13** presents whole proteins ranked by the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile. The proteins are identified by UniProtID number in the first column of the table, their name in the second column, and the source of the protein in the third column. The fourth column indicates whether the protein contains every essential amino acid (1) or not (0). The fifth and sixth columns present the weight fraction of essential essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The seventh column presents the PDCAAS score for each protein. The eighth column presents the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile for each protein. The proteins are ranked from shortest to longest distance from the target profile.

[00424] The whole protein standards are all significantly less nutritious (as measured by their PDCAAS score) than either the whole protein or fragment candidates identified in Examples 2-5. Ovalbumin has the highest PDCAAS value, at 1.33, while the proteins and fragments presented in Examples 2-5 all have PDCAAS values > 1.8. Similarly, while some proteins have EAA/NEAA ratios similar to the WHO 60/40 targets (Beta-lactoglobulin has a distance of ~7%), they still lack high nutritional value (PDCAAS of ~0.9).

Table 13

			EAA					Distance from Target Distribution
SEQI D	UniProtI D	Name	Sourc e	Comple t	[EAA]	[NEAA]	PDCAA S	
633	P02789	Ovotransferri n	Egg	1	0.45	0.55	1.31	0.0862
574	P00711	Alpha- lactalbumin	Whey	1	0.53	0.47	1.25	0.0945
585	P01012	Ovalbumin	Egg	1	0.47	0.53	1.33	0.0969
625	P02668	Kappa- Casei	Casei	1	0.44	0.56	1.17	0.1308

		casein	n					
623	P02666	Beta-casein	Casei	1	0.48	0.52	1.24	0.1452
			n					
619	P02662	Alpha-S1-	Casei	1	0.44	0.56	1.08	0.1133
		casein	n					
621	P02663	Alpha-S2-	Casei	1	0.47	0.53	1.05	0.1299
		casein	n					
655	P04776	Glycinin G1	Soy	1	0.38	0.62	1.10	0.1412
791	P11828	Glycinin G3	Soy	1	0.37	0.63	1.00	0.1518
627	P02754	Beta-	Whey	1	0.50	0.50	0.92	0.0744
		lactoglobulin						
635	P02858	Glycinin G4	Soy	1	0.35	0.65	0.87	0.1452
646	P04347	Glycinin	Soy	1	0.36	0.64	0.93	0.1489
631	P02769	Serum	Whey	1	0.46	0.54	0.82	0.0948
		albumin						
789	P11827	Beta-	Soy	1	0.37	0.63	0.80	0.1572
		conglycinin						
		alpha prime						
		chain						
648	P04405	Glycinin G2	Soy	1	0.36	0.64	0.67	0.1509
822	P13916	Beta-	Soy	1	0.34	0.66	0.59	0.1688
		conglycinin						
		alpha chain						
1588	P68390	Ovomucoid	Egg	0	0.38	0.62	0.00	0.1388
1010	P25974	Beta-	Soy	0	0.41	0.59	0.00	0.1440
		conglycinin						
		beta chain						

[00425] **Table 14** presents protein fragments ranked by PDCAAS value. The protein fragments are identified by UniProtID number in the first column of the table, the beginning and ending amino acid positions relative to the protein amino acid sequence in the second column, their name in the third column, and the source of the protein in the fourth column. The fifth column indicates whether the protein fragment contains every essential amino acid (1) or not (0). The sixth and seventh columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e.,

including C and Y) (NEAA + CEAA), respectively. The eighth column presents the PDCAAS score for each protein fragment. The protein fragments are ranked from highest to lowest PDCAAS value.

Table 14

SEQID	UniProtID	Fragment Ends	Protein Name	Source	EAA		PDCAA S
					Complete	[EAA] [NEAA]	
586	P01012	250:332	Ovalbumin	Egg	1	0.52 0.48	1.65
624	P02666	38:159	Beta-casein	Casein	1	0.49 0.51	1.62
575	P00711	5:81	Alpha-lactalbumin	Whey	0	0.53 0.47	1.61
634	P02789	3:171	Ovotransferrin	Egg	0	0.46 0.54	1.52
632	P02769	205:294	Serum albumin	Whey	1	0.46 0.54	1.47
636	P02858	56:103	Glycinin G4	Soy	0	0.46 0.54	1.47
622	P02663	191:222	Alpha-S2-casein	Casein	0	0.55 0.45	1.45
656	P04776	221:420	Glycinin G1	Soy	1	0.40 0.60	1.38
792	P11828	228:379	Glycinin G3	Soy	1	0.41 0.59	1.34
626	P02668	69:133	Kappa-casein	Casein	1	0.45 0.55	1.32
628	P02754	57:177	Beta-lactoglobulin	Whey	1	0.48 0.52	1.28
647	P04347	53:115	Glycinin	Soy	1	0.47 0.53	1.23
620	P02662	87:211	Alpha-S1-casein	Casein	1	0.39 0.61	1.21
649	P04405	127:259	Glycinin G2	Soy	1	0.39 0.61	1.14
790	P11827	162:341	Beta-conglycinin alpha prime chain	Soy	1	0.37 0.63	0.56
823	P13916	63:216	Beta-conglycinin alpha chain	Soy	0	0.21 0.79	0.23

1589	P68390	1:20	Ovomucoid	Egg	0	0.39	0.61	0.00
1011	P25974	1:20	Beta- conglycinin beta chain	Soy	0	0.72	0.28	0.00

[00426] **Table 15** presents protein fragments ranked by the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile. The protein fragments are identified by UniProtID number in the first column of the table, the beginning and ending amino acid positions relative to the protein amino acid sequence in the second column, their name in the third column, and the source of the protein in the fourth column. The fifth column indicates whether the protein fragment contains every essential amino acid (1) or not (0). The sixth and seventh columns present the weight fraction of essential amino acids (EAA), and the total of non-essential and conditionally-essential amino acids (i.e., including C and Y) (NEAA + CEAA), respectively. The eighth column presents the PDCAAS score for each protein fragment. The ninth column presents the Euclidian distance from a 60/40 EAA/NEAA (Whey) target amino acid profile for each protein fragment. The protein fragments are ranked from shortest to longest distance from the target profile.

[00427] The whole protein standards are all significantly less nutritious (as measured by their PDCAAS score) than either the whole protein or fragment candidates identified in Examples 2-5. Ovalbumin has the highest PDCAAS value, at 1.33, while the proteins and fragments presented in Examples 2-5 all have PDCAAS values > 1.8. Similarly, while some proteins have EAA/NEAA ratios similar to the WHO 60/40 target (Beta-lactoglobulin has a distance of ~7%), they still lack high nutritional value (PDCAAS of ~0.9).

[00428] Even the best fragments of the whole protein standards are all significantly less nutritious (as measured by their PDCAAS score) than either the whole protein or fragment candidates identified in Examples 2-5. The highest scoring fragment, from ovalbumin, only has a PDCAAS score of 1.6, while the top fragment candidates have scores > 2.5. Similar to the full length standard proteins, there are fragments that have EAA/NEAA ratios similar to the WHO 60/40 target (~8%), but they are not nearly as nutritious (PDCAAS of 1.52). Again, for comparison, the top fragment candidates ranked by distance to the WHO 60/40 target show distances <5% with PDCAAS scores ~2.

Table 15

SEQID	UniProtID	Fragment		Source	EAA				Distance
		Ends	Name		Complete	[EAA]	[NEAA]	PDCAAS	from Target Distribution
586	P01012	250:332	Ovalbumin	Egg	1	0.52	0.48	1.65	0.1293
575	P00711	5:81	Alpha-lactalbumin	Whey	0	0.53	0.47	1.61	0.0984
624	P02666	38:159	Beta-casein	Casein	1	0.49	0.51	1.62	0.1493
634	P02789	3:171	Ovotransferrin	Egg	0	0.46	0.54	1.52	0.0801
632	P02769	205:294	Serum albumin	Whey	1	0.46	0.54	1.47	0.1160
636	P02858	56:103	Glycinin G4	Soy	0	0.46	0.54	1.47	0.1317
622	P02663	191:222	Alpha-S2-casein	Casein	0	0.55	0.45	1.45	0.2033
656	P04776	221:420	Glycinin G1	Soy	1	0.40	0.60	1.38	0.1253
628	P02754	57:177	Beta-lactoglobulin	Whey	1	0.48	0.52	1.28	0.1038
792	P11828	228:379	Glycinin G3	Soy	1	0.41	0.59	1.34	0.1188
626	P02668	69:133	Kappa-casein	Casein	1	0.45	0.55	1.32	0.1737
647	P04347	53:115	Glycinin	Soy	1	0.47	0.53	1.23	0.1277
620	P02662	87:211	Alpha-S1-casein	Casein	1	0.39	0.61	1.21	0.1395
649	P04405	127:259	Glycinin G2	Soy	1	0.39	0.61	1.14	0.1599
790	P11827	162:341	Beta-conglycinin alpha prime chain	Soy	1	0.37	0.63	0.56	0.1612
823	P13916	63:216	Beta-conglycinin alpha chain	Soy	0	0.21	0.79	0.23	0.3366
1589	P68390	1:20	Ovomucoid	Egg	0	0.39	0.61	0.00	0.2130
1011	P25974	1:20	Beta-conglycinin beta chain	Soy	0	0.72	0.28	0.00	0.2717

Example 7: Protein Fragmentation Analysis

[00429] When a protein is broken into overlapping fragments and the EAA and NEAA content of the fragments is analyzed, fragments that overlap with each other will tend to have similar EAA and NEAA content profiles and thus necessarily similar PDCAAS values and similar Euclidian distances from a 60/40 EAA/NEAA (Whey) target amino acid profile and a 60/40 EAA/NEAA (Whey) target amino acid profile. In **Tables 7, 8, and 10**, only the top fragment identified in each protein is listed for that protein. However, there are likely a multitude of variants that have very similar sequences that are thus very similar in PDCAAS value and distance from an EAA/NEAA (Whey) target profile. One way to visualize this is by performing a full fragmentation analysis and plotting the resulting fragment scores.

Figure 17 plots the PDCAAS values for all fragments of the protein Q1WLP9 (SEQ ID NO: 4403). Clearly, there is a hot spot of overlapping fragments starting around position 150, ending around 170, with lengths between 80 and 120 amino acids. **Figure 18** shows a similar fragmentation analysis for P54612 (SEQ ID NO: 4404) using the distance to the 60/40 EAA/NEAA distribution as the ranking metric.

Example 8: Protein Expression

[00430] Genes encoding nutritive proteins of this disclosure were codon optimized for expression in *Escherichia coli* and synthesized by either LifeTechnologies/GeneArt or DNA 2.0. Genes were designed to express the native protein or to contain one of two amino-terminal tags to facilitate purification:

[00431] MGSHHHHHHHH (SEQ ID NO: 4431)

[00432] MGSSHHHHHHSSGLVPRGSH (SEQ ID NO: 4430)

[00433] These gene constructs were inserted into the pET15b plasmid vector (Novagen) using *NcoI*-*Bam*HI restriction sites (in case of the first tag) or using the *NdeI*-*Bam*HI restriction sites (in the case of the second tag). All restriction enzymes were purchased from New England Biolabs. Plasmids were transformed into *Escherichia coli* T7 Express (New England Biolabs) and selected on lysogeny broth (LB) plates containing 100 mg/l carbenicillin. A single colony was picked, grown to $OD_{600nm} \approx 0.6$ in LB with 100 mg/l carbenicillin, and stored as a glycerol stock (in LB with 10% glycerol (v/v)) at $-80^{\circ}C$, to serve as a master cell stock.

[00434] 2ml LB with 100 mg/l carbenicillin (in a 14mm×100mm culture tube) was inoculated with a stab from the glycerol stock and grown overnight at $37^{\circ}C$ and 250rpm. The

next day, 2ml LB with 100 mg/l carbenicillin (in a 14mm×100mm culture tube) was inoculated with the overnight culture to $OD_{600nm} = 0.05$ and grown at 30°C or 37°C and 250rpm. At $OD_{600nm} \approx 0.8$, heterologous gene-expression was initiated with 1mM isopropyl β -D-1-thiogalactopyranoside (IPTG) and grown for another 2 hr (when grown at 37°C) or 4 hr (when grown at 30°C) until harvest. Upon harvesting, OD_{600nm} was measured, a 1ml aliquot was centrifuged, and the supernatant was decanted. Cells were re-suspended to $OD_{600nm} = 1.50$ for SDS-PAGE analysis to evaluate expression level. 10 μ l of resuspended culture was loaded onto either: 1) a Novex® NuPAGE® 12% Bis-Tris gel (Life Technologies), or 2) a Novex®16% Tricine gel (Life Technologies), and run using standard manufacturer's protocols. Gels were stained using SimplyBlue™ SafeStain (Life Technologies) using the standard manufacturer's protocol and imaged using the Molecular Imager® Gel Doc™ XR+ System (Bio-Rad). Over-expressed heterologous protein was identified by comparison against a molecular weight marker and control cultures.

[00435] Using this method, recombinant expression of the proteins listed in Tables 16A and 16B was observed.

Table 16A

Seq ID No	UniProt	FragEnds	EAA	BCAA	L	C	PDCAAS
4409	P02114	-	0.56	0.24	0.12	3	1.58
4410	P67975	-	0.50	0.25	0.13	5	1.40
4411	P12106	170:220	0.70	0.32	0.14	1	1.84
4412	P32191	112:163	0.58	0.31	0.15	1	1.27
4413	Q03262	225:277	0.65	0.30	0.13	1	1.51
4414	P12106	159:209	0.67	0.29	0.14	1	1.77
4415	P32492	1218:1319	0.67	0.25	0.14	1	1.47
4416	Q2XQV4	108:213	0.49	0.26	0.15	2	1.41
4417	P32492	1086:1287	0.58	0.29	0.15	1	1.20
4418	Q0WVK7	53:105	0.57	0.27	0.12	2	1.95
4419	Q94A52	149:261	0.54	0.26	0.12	2	2.01
4420	P38111	1093:1165	0.59	0.30	0.13	3	2.11
4421	P38111	1093:1182	0.57	0.27	0.13	3	1.76
4422	P38111	1093:1162	0.57	0.27	0.13	3	1.88
4423	P38111	1092:1166	0.59	0.29	0.13	3	2.07

4424	P38111	1093:1168	0.58	0.29	0.13	3	2.04
4425	P38111	1091:1164	0.59	0.30	0.13	3	1.79
4426	P38111	1089:1164	0.58	0.29	0.13	3	1.75

Table 16B

Seq ID No	SolveScore	AggScore	Allergenicity	Toxicity	Antinutricity	Human Homology
4409	-17.34	0.56	0.34	0.00	0.23	0.75
4410	-21.19	0.47	0.99	0.22	0.24	0.43
4411	-17.13	0.77	0.20	0.31	0.28	0.61
4412	-16.98	0.76	0.19	0.25	0.17	0.52
4413	-19.93	0.52	0.22	0.32	0.32	0.33
4414	-16.05	0.65	0.20	0.28	0.31	0.65
4415	-21.64	0.55	0.26	0.26	0.26	0.00
4416	-17.59	0.58	0.60	0.00	0.23	0.98
4417	-19.98	0.57	0.31	0.00	0.24	0.00
4418	-26.91	0.45	0.18	0.30	0.22	0.31
4419	-25.52	0.46	0.29	0.00	0.25	0.81
4420	-23.00	0.80	0.27	0.00	0.24	0.26
4421	-20.82	0.80	0.27	0.00	0.25	0.27
4422	-23.98	0.70	0.25	0.27	0.25	0.27
4423	-22.45	0.79	0.27	0.27	0.25	0.25
4424	-22.10	0.80	0.27	0.28	0.24	0.25
4425	-22.75	0.77	0.28	0.31	0.25	0.25
4426	-23.28	0.71	0.28	0.28	0.25	0.24

Example 9: Scaled Up Production of Recombinant Nutritive Proteins

[00436] A representative protocol for producing quantities of nutritive proteins as described in this disclosure is as follows.

[00437] 5ml LB with 100 mg/l carbenicillin (in a 50ml baffled Pyrex shake flask) is inoculated with a stab from the glycerol stock of a recombinant *E. coli* strain comprising a recombinant gene encoding a nutritive protein and grown until late exponential phase (OD_{600nm} $\approx$ 2) at 37°C and 250rpm. A 2.5l Ultra Yield Flask (Thomson Instrument Company) is inoculated with 500ml sterile water and enough EnBase EnPresso™ tablets

(BioSilta) to formulate 500ml growth medium. This medium is supplemented with 100 mg/l carbenicillin, 0.001% Industrol 204 antifoam, and 0.6 U/l EnzI'm (BioSilta). The shake flask is inoculated to OD_{600nm} = 0.05 and grown 16hr at 30°C and 250rpm. The growth medium is supplemented with EnPresso™ Booster tablets (BioSilta), 1.2 U/l EnzI'm, and 1mM IPTG to induce heterologous protein production. After another 8-24hr of shaking at 30°C and 250rpm, the flask is harvested by centrifugation, the supernatant is decanted, and the wet cell weight was measured. Approximately 20gWCW (grams wet cell weight)/l medium is typically recovered at this stage.

[00438] The harvested cells from each shake-flask fermentation are suspended in 25 mL of IMAC Equilibration Solution (30 mM Imidazole, 50 mM Phosphate, 0.5 M NaCl, pH 7.5). The suspended cells are then lysed by sonication on ice. The lysed cells are centrifuged for 60 minutes and decanted. The cell debris is discarded, and the supernatants are 0.2 µm filtered. Filters are then flushed with an additional 10 mL of IMAC Equilibration Solution. These filtered protein solutions are then purified by immobilized metal affinity chromatography (IMAC).

[00439] IMAC resin (GE Healthcare, IMAC Sepharose 6 Fast Flow) is charged with nickel and equilibrated. 30 mL of each protein solution is loaded onto a 5 mL IMAC column, and washed with additional equilibration solution to remove unbound impurities. The protein of interest is then eluted with 15 mL of 0.5 M NaCl, 0.2 M Imidazole, pH 7.5. At this stage, the purified proteins are typically shown to be at least 90% pure by SDS-PAGE. Approximately 20 to 60 mg of each protein is recovered in the IMAC elution fractions. Each IMAC elution fraction is buffer exchanged by dialysis into a formulation solution (20 mM HEPES, pH 7.5). After buffer exchange, the protein solutions are recovered for all downstream processing.

Example 10: Prediction of Soluble Expression of Nutritive Proteins

[00440] Open reading frames encoding a set of 292 nutritive proteins were cloned and introduced into E. coli to assess recombinant protein expression using the method of Example 6. In the system used, 163 proteins were identified as expressed while 129 were not. Of the 163 proteins that expressed, 125 were tested for soluble expression. It was found that 75 were solubly expressed while 50 were not.

[00441] Figure 19 shows a two dimensional histogram of protein expression in the E. coli expression screen. Figure 1 shows the relative likelihood (on a log scale) of a protein

being expressed as a function of solvation score (y-axis) and aggregation score (x-axis). A darker mark on the histogram indicates a higher number of proteins expressed, while a lighter mark indicates a fewer number of proteins expressed. Figure 19 shows that those proteins that were successfully expressed tend to cluster in the top left region of the plot, where the solvation score is more negative (≤ -20) and the aggregation score is smaller (≤ 0.75). There were few examples of proteins that were successfully expressed with less negative solvation scores (≥ -15) and large aggregation scores (≥ 1). This result suggests that nutritive proteins with solvation scores of -20 or less and aggregation scores of 0.75 or less are more likely to be expressed in this system.

[00442] Figure 20 shows a two dimensional histogram of the number of soluble protein expression in the E. coli expression screen. Figure 20 shows the relative likelihood (on a log scale) of a protein being solubly expressed as a function of solvation score (y-axis) and aggregation score (x-axis). Again, a darker mark on the histogram indicates a higher number of proteins expressed, while a lighter mark indicates a fewer number of proteins expressed. Figure 1 shows that those proteins that were expressed solubly tended to cluster in the top left region of the plot, where the solvation score is more negative (≤ -20) and the aggregation score is smaller (≤ 0.5). There were few examples of proteins that were expressed solubly with less negative solvation scores (≥ -15) and large aggregation scores (≥ 0.75). This result suggests that nutritive proteins with solvation scores of -20 or less and aggregation scores of 0.5 or less are more likely to be solubly expressed in this system.

Example 11: Solubility Screening

[00443] The solubility of one nutritive protein produced as described Examples 8 and 9 was examined by centrifuge concentration followed by protein concentration assays. Samples in 20 mM HEPES pH 7.5 were tested for protein concentration according to the protocol for Coomassie Plus (Bradford) Protein Assay (Thermo Scientific) and absorbance at 280 nm (if applicable). Based on these measurements 10 mg of protein was added to an Amicon Ultra 3 kDa centrifugal filter (Millipore). Samples were concentrated by centrifugation at 10,000 Xg for 30 minutes. The final concentrated samples were examined for precipitated protein and color, and then tested for protein concentration as described above. The results are shown in Table 17.

Table 17

Seq ID No	Appearance	Concentration (g/L)
4420	Clear Yellow	191

The solubilities of this nutritive protein was found to be significantly higher than concentrations typically found for whey (12.5 g/L) and soy (10 g/L) (Pelegrine, D. H. G. & Gasparetto, C. A., 2005. Whey proteins solubility as function of temperature and pH. *LWT - Food Science and Technology*, p. 77–80; Lee, K. H., Ryu, H. S. & Rhee, K. C., 2003. Protein solubility characteristics of commercial soy protein products. *Journal of the American Oil Chemists' Society*, pp. 85-90). This demonstrates the usefulness of the nutritive proteins disclosed herein. For example, the solubility of nutritive proteins may improve compliant delivery of high quality protein in as small of a volume as possible while avoiding the “chalkyness” that often characterizes proteins delivered in this manner. This may, for example, be useful to deliver proteins to the elderly or other subjects.

Example 12: Stability Screening

[00444] Thermal stability of nutritive proteins provides insight regarding whether the protein is likely to have a useful shelf life. Samples of a protein produced as described in Examples 8 and 9 were screened using a rapid thermal stability screening method. In this method proteins were heated slowly from 25°C to 95°C in two representative formulations in the presence of a hydrophobic dye (Enzo Life Sciences, ProteoStat® Thermal shift stability assay kit) that binds to aggregated proteins that form as the protein denatures with increasing temperature (Niesen, F. H., Berglund, H. & Vadadi, M., 2007. The use of differential scanning fluorimetry to detect ligand interactions that promote protein stability. *Nature Protocols*, Volume 2, pp. 2212-2221.). Upon binding, the dye’s fluorescence increases significantly, which is then recorded by the rtPCR instrument and represented as the protein’s melting curve (Lavinder, J. J., Hari, S. B., Sullivan, B. J. & Magilery, T. J., 2009. High-Throughput Thermal Scanning: A General, Rapid Dye-Binding Thermal Shift Screen for Protein Engineering. *Journal of the American Chemical Society*, pp. 3794-3795.). After the thermal shift is complete samples were examined for insoluble precipitates and further analyzed by analytical size exclusion chromatography (SEC).

[00445] Protein solutions (12.5 mg/ml) were prepared in both PBS and 20 mM HEPES pH 7.7 buffers, each containing 1X ProteoStat TS Detection Reagent. Samples of each

solution were heated slowly from 25°C – 95°C, 0.5°C/30 seconds using a real-time PCR (rtPCR) thermocycler while monitoring the fluorescence of the dye. From this thermal scan the temperature of aggregation was determined (T_{agg}) from the temperature with the strongest slope if an increase in fluorescence was observed. To supplement the assay, samples were taken before and after the thermal shift and analyzed by SEC (GE Healthcare - Superdex 75 5/150) which can detect large soluble aggregates. The results for one nutritive proteins of this disclosure and a whey standard are presented in Table 18. The presence of soluble aggregates detected by SEC is noted by a “yes” if observed or “no” if not observed, and the “n/a” entries for the whey standard indicate the production of insoluble precipitates such that no SEC analysis was performed.

Table 18

Seq ID No	HEPES - T_{agg}	PBS - T_{agg}	HEPES - SEC Agg?	PBS - SEC Agg?
4420	95	95	Yes	Yes
whey	79	81.5	n/a	n/a

[00446] As shown in Table 18, SEQ ID NO: 4420 had a much higher HEPES T_{agg} than whey (in fact, the protein did not form any insoluble aggregates at 95°C, which was the upper limit of the assay), and thus is expected to be more stable than whey.

Example 13: Digestibility Screening – Determination of Digestion Half-Life

[00447] The goal of screening for protein digestibility is to eliminate potentially unsafe allergenic proteins and to determine the relative completeness of digestion as a predictor of peptide bioavailability. This screening method utilizes a physiologically relevant *in vitro* digestion reaction that includes both phases of protein digestion, simulated gastric digestion and simulated intestinal digestion (Moreno, J. F. et al., 2005. Stability of the major allergen Brazil nut 2S albumin (Ber e 1) to physiologically relevant *in vitro* gastrointestinal digestion. FEBS Journal, pp. 341-352.). Samples can be taken throughout the reaction and analyzed for intact protein and peptide fragments using chip electrophoresis and LC-QTOF-MS. Proteins with allergenic properties can be assessed by identifying proteins or large fragments of proteins that are resistant to digestive proteases and thus have a higher risk of causing an allergenic reaction (Goodman, R. E. et al., 2008. Allergenicity assessment of genetically modified crops - what makes sense?. Nature Biotechnology, pp. 73-81.). Digestibility is measured by determining how efficiently the protein is broken down into peptides (Daniel,

H., 2003. Molecular and Integrative Physiology of Intestinal Peptide Transport. Annual Review of Physiology, Volume 66, pp. 361-384.).

[00448] The method used an automated assay for in vitro digestions of proteins wherein assay conditions and protease concentrations are physiologically relevant (Moreno, F. J., Mackie, A. R. & Clare Mills, E. N., 2005. Phospholipid interactions protect the milk allergen α -Lactalbumin from proteolysis during in vitro digestion. Journal of agricultural and food chemistry, pp. 9810-9816; Martos, G., Contreras, P., Molina, E. & Lopez-Fandino, R., 2010. Egg White Ovalbumin Digestion Mimicking Physiological Conditions. Journal of Agricultural and food chemistry, pp. 5640-5648; Moreno, J. F. et al., 2005. Stability of the major allergen Brazil nut 2S albumin (Ber e 1) to physiologically relevant in vitro gastrointestinal digestion. FEBS Journal, pp. 341-352.). The first phase of digestion is in simulated gastric fluid (SGF) and formulated at pH 1.5 and with a pepsin:substrate ratio of (1:10 w/w). The second phase of digestion is in simulated intestinal fluid (SIF) is formulated with bile salts at pH 6.5 and with an trypsin:chymotrypsin:substrate ratio of (1:4:400 w/w). The protein is treated for 120 mins in the simulated gastric fluid, which is how long it takes for 90% of a liquid meal to pass from the stomach to the small intestine (Kong, F. & Singh, R. P., 2008. Disintegration of Solid Foods in Human Stomach. *Journal of Food Science*, pp. 67-80), and then treated with simulated intestinal fluid for 120 mins. Sample time points are taken throughout both reactions and quenched for analysis. Bovine serum albumin, which is readily digested by pepsin, is the positive control for the SGF solution, and beta-lactoglobulin, which is naturally resistant to pepsin but digested in SIF, is the positive control for SIF solution. Intact protein and large fragments were detected using electrophoresis. For chip electrophoresis, a Caliper Labchip GXII equipped with a HT Low MW Protein Assay Kit was used to monitor the size and amount of intact protein as well as any digestion fragments larger than 4 kDa. By monitoring the amount of intact protein observed over time, the half-life ($\tau_{1/2}$) of digestion was calculated for SGF and, if intact protein is detected after SGF digestion, in SIF.

[00449] This method was used to analyze the digestion half-lives of two nutritive proteins of this disclosure (SEQ ID NOS: 4419 and 4420) produced as described in Examples 6 and 7, as well as native and recombinant ovalbumin (OVA and rOVA, respectively; SEQ ID NO: 4427) and beta-lactoglobulin (BLG and rBLG, respectively; SEQ ID NO: 4428) proteins and a whey standard. The results of these experiments are summarized in Table 19.

An “n/a” entry in the Simulated Intestinal Fluid field indicates that no intact protein was detected after SGF digestion.

Table 19

Seq ID No.	Digestion $\tau_{1/2}$ (min.)	
	Simulated Gastric Fluid	Simulated Intestinal Fluid
4419	1	0
4420	6	0
BLG (4428)	77	4
rBLG (4428)	50	0.7
OVA (4427)	18	1
rOVA (4427)	5	n/a
whey	99	4

[00450] The results shown in Table 19 indicate that the two nutritive proteins of this disclosure were completely digested by SGF and have SGF half lives of six minutes or less. By comparison whey is not completely digested by SGF and has an SGF half life of 99 minutes and a SIF half life of 4 minutes. This study suggests that the nutritive proteins of this disclosure are likely to be readily digested and not likely to elicit an allergic response when ingested.

[00451] The results in Table 19 also show that the recombinant beta-lactoglobulin and ovalbumin produced according to this disclosure were both more readily digested than their naturally-occurring counterparts. The speed in which a protein is broken down can be controlled by selecting for properties that improve or limit accessibility of the gastrointestinal proteases. This capability can be demonstrated for two typical protein properties, glycosylation and disulfide cross-linking. Like many naturally occurring proteins, naturally occurring OVA and BLG are glycosylated by their host organisms. In contrast, the recombinant proteins produced according to the present disclosure are not glycosylated because the host organism (*E. coli* in this case) does not glycosylate. The lack of glycosylation in recombinant nutritive proteins according to this disclosure may result in proteins that are more readily digested. Furthermore, BLG has four disulfide bonds that are known to slow down or interfere with digestion. When these disulfide bonds are disrupted, the rate of digestion increases (Reddy, I. M., Kella, N. K. D. & Kinsella, J. E., 1988.

Structural and conformational Basis of the Resistance of b-Lactoglobulin to Peptic and Chymotryptic Digestion. *J. Agric. Food Chem.*, Volume 36, pp. 737-741). A lack or disruption of disulfide bond formation in recombinant nutritive proteins according to this disclosure may result in proteins that are more readily digested.

Example 14: Digestibility Screening – Analysis of Digestion Products

[00452] Two nutritive proteins produced as described in Examples 8 and 9 (SEQ ID NOS: 4429 and 4420) were subjected to SGF and SIF digestion as described in Example 11. Both proteins were completely digested in SGF, and the SGF half lives are shown in Table 12.

Table 20

Seq ID No	Digestion $\tau_{1/2}$ (min.)	
	Simulated Gastric Fluid	Simulated Intestinal Fluid
4429	0.7	n/a
4420	6	n/a

[00453] To detect and identify peptides that were present after SGF and SIF digestion, samples of the SGF and SIF digests were analyzed by LC/Q-TOF MS/MS. Samples from the SGF digests were directly analyzed by LC/Q-TOF MS/MS, while SIF protein digestions required purification by SCX to remove bile acids before detection and identification by LC/Q-TOF MS/MS. Peptides were extracted from the chromatograms and identified using Bioconfirm Software (Aglient). The sequence assignment of peptides were based on accurate mass match (± 10 ppm) and further confirmed by MS/MS fragmentation. The results are shown in Tables 21 and 22 below.

Table 21
(SEQ ID NO: 4429)

SGF Peptides 120 min	SEQ ID NO	SIF Peptides 120 min	SEQ ID NO
LL		SE	
LAL		PSE	
HVL		HVL	
LEL		FKV	

LALA	4432	HQI	
LLLD	4433	PSEA	4434
IAEF	4435	REV	
IQQF	4436	FDK	
YDKL	4437	AEFK	4438
SNLTE	4439	LKHV	4440
ELLEA	4441	SSSEL	4442
EELAL	4443	FKVF	4444
DDLLL	4445	AELKH	4446
LAYDK	4447	LKHVL	4448
TKTRL	4449	FKEAF	4450
DLDHQ	4451	NGSISS	4452
GTLENL	4453	SLGLSPS	4454
EKLTD	4455	GEKLT	4456
AEVDDM	4457	ELATVM	4458
LDDLLL	4459	GSGEINI	4460
LDLDHQ	4461	RSLGLSP	4462
DLKKKL	4463	ELKHVL	4464
KTKTRL	4465	KLTD	4466
SQRLEE	4467	FKVFDK	4468
AEVDDML	4469	AEVDDML	4470
QQELDDL	4471	DAEVDDM	4472
LEKTKTR	4473	AELKHVL	4474
LEKTKTR	4475	HVLTSIGE	4476
GTLENLEE	4477	SIGEKLT	4478
QQELDDL	4479	RSLGLSPSE	4480
KLEKTKTRLQ	4481	VLT SIGEKL	4482
SRQLKSNDSEQ	4483	TSIGEKLT	4484
EKTKTRLQQEL	4485	HVLTSIGEK	4486
YDKLEKTKTRL	4487	AELKHVLT	4488
LAYDKLEKTKTRL	4489	RSLGLSPSEA	4490
EELKKKLLKDLEL	4491	SSNLTEEQIA	4492
EELKKKLLKDLELL	4493	RSLGLSPSEAE	4494
AELKHVLT SIGEKLTD	4495	KVFDKNGDGLISA	4496
MGSHHHHHHHSSNL	4497	FDKDNNGSISSSEL	4498

LREVSDGSGEINIQQF	4499	REVSDGSGEINIQQ	4500
AAELKHVLTSIGEKLTD	4501	DVDGNHQIEFSEF	4502
AELKHVLTSIGEKLTDAE	4503	LREVSDGSGEINIQQ	4504
AYDKLEKTKTRLQQEL	4505	REVSDGSGEINIQQF	4506
AAELKHVLTSIGEKLTDAE	4507	LREVSDGSGEINIQQFAALLS	4508
LENLEELKKKLLKDLEL	4509		
KLEKTKTRLQQELDDL	4510		
DKLEKTKTRLQQELDDL	4511		
LAYDKLEKTKTRLQQELDDL	4512		
LALAYDKLEKTKTRLQQELDDL	4513		

Table 22
(SEQ ID NO: 4420)

SGF Peptides 120 min	SEQ ID NO	SIF Peptides 120 min	SEQ ID NO
GVL		TKH	
ALL		INDI	4514
LVL		HLVL	4515
IGVL	4516	TIKF	4517
TIKF	4518	IGVLD	4519
TIKF	4520	RNLD	4521
EVYDL	4522	IGVLDV	4523
LNDSVQ	4524	QTIKF	4525
IWVIND	4526	VQTIKF	4527
DLNDSVQ	4528	SVQTIKF	4529
SVQTIKF	4530	KCAKCISMIGVL	4531
HHLVLGALLD	4532	EKCAKCISMIGV	4533
HHHHHHLVL	4534	HEFKRTTYSE	4535
HHHHHHHHLVL	4536	SHKFRNLDKDL	4537
DVTKHEFKRTTY	4538	ISMIGVLDVTKHE	4539
SHHHHHHHHHLVL	4540		
TKHEFKRTTYSEN	4541		
GSHHHHHHHHHLVLG	4542		
MGSHHHHHHHHHLVL	4543		
KRTTYSENEVYDLN	4544		

[00454] As can be seen in Tables 21 and 22, each protein was digested into multiple smaller peptide fragments ranging in size from 2 to 22 amino acids (SEQ ID NO: 762) or 2 to 13 amino acids (SEQ ID NO: 763). None of these peptide fragments was found to be homologous to any known allergen.

[00455] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

AMENDED CLAIMS
received by the International Bureau on
04 SEPT 2013
(04.09.2013)

CLAIMS:

1. An isolated nutritive protein comprising a first polypeptide comprising a first polypeptide sequence,
wherein at least 40% by weight of the amino acids in the first polypeptide sequence
5 are essential amino acids,
and wherein the first polypeptide sequence has a protein digestibility corrected amino acid score (PDCAAS) of at least about 1.4.
2. The nutritive protein of claim 1, wherein the first polypeptide sequence has a
10 PDCAAS of from about 1.4 to about 1.8.
3. The nutritive protein of claim 1, wherein the first polypeptide sequence comprises a weight ratio of at least 50% or 60% essential amino acids.
- 15 4. The nutritive protein of claim 3, wherein the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.2.
5. The nutritive protein of claim 1, wherein the first polypeptide sequence comprises a weight ratio of at least 70% essential amino acids.
20
6. The nutritive protein of claim 5, wherein the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.5.
7. The nutritive protein of claim 1, wherein the first polypeptide sequence
25 comprises a weight ratio of at least 80% essential amino acids.
8. The nutritive protein of claim 7, wherein the first polypeptide sequence has a PDCAAS of from about 1.4 to about 2.9.
- 30 9. The nutritive protein of claim 1, wherein the first polypeptide sequence comprises a weight ratio of at least 90% essential amino acids.

10. The nutritive protein of claim 9, wherein the first polypeptide sequence has a PDCAAS of from about 1.4 to about 3.2.

11. The nutritive protein of any one of claims 1 to 10, further comprising at least
5 one of every non-essential amino acid.

12. The nutritive protein of any one of claims 1 to 10, further comprising at least one of every non-essential and conditionally-essential amino acid.

10 13. The nutritive protein of claim 11 or 12, further comprising a nutritionally balanced ratio of non-essential amino acids.

14. The nutritive protein of claim 13, wherein the first polypeptide sequence comprises a weight ratio of each non-essential amino acid to the total number of amino acids
15 that is equivalent to the weight ratio of each non-essential amino acid to the total number of amino acids in a benchmark protein.

15. The nutritive protein of claim 14, wherein the benchmark protein is selected from whey protein, egg protein and soy protein.

20

16. The nutritive protein of claim 1, wherein the first polypeptide sequence comprises 100% essential and conditionally essential amino acids.

17. The nutritive protein of claim 16, wherein the first polypeptide sequence has a
25 PDCAAS of from about 1.4 to about 3.6.

18. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least 70% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof.

30

19. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least 95% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof.

20. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least 70% homology to a naturally occurring protein or fragment thereof.

5

21. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises at least 95% homology to a naturally occurring protein or fragment thereof.

10

22. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises a naturally occurring protein or fragment thereof.

15

23. The nutritive protein of any one of claims 1 to 17, wherein the first polypeptide sequence comprises an altered amino acid sequence compared to the amino acid sequence of a naturally occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally occurring reference protein.

20

24. The nutritive protein of claim 23, wherein the first polypeptide sequence comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally occurring reference protein.

25

25. The nutritive protein of any one of claims 1 to 24, wherein the first polypeptide sequence is not an allergen.

26. The nutritive protein of any one of claims 1 to 25, wherein the first polypeptide sequence has less than 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 75, 80, 85, or 90% global homology to a known allergen.

30

27. The nutritive protein of any one of claims 1 to 26, wherein the first polypeptide sequence is not a toxin.

28. The nutritive protein of claim 27, wherein the first polypeptide sequence has less than 50% global homology to a known toxin.

29. The nutritive protein of any one of claims 1 to 28, wherein the first
5 polypeptide sequence has a simulated gastric digestion half-life shorter than whey.

30. The nutritive protein of claim 29, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 60 minutes or less than 30 minutes.

10 31. The nutritive protein of claim 30, wherein the first polypeptide sequence has a simulated gastric digestion half-life of less than 10 minutes.

32. The nutritive protein of any one of claims 1 to 31, wherein the first
15 polypeptide sequence is completely digested in simulated gastric fluid.

33. The nutritive protein of any one of claims 1 to 32, wherein the first polypeptide sequence comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

20 34. The nutritive protein of any one of claims 1 to 33, wherein the first polypeptide sequence comprises 0 or 1 cysteine residues.

35. The nutritive protein of any one of claims 1 to 34, wherein the first
25 polypeptide sequence comprises no disulfide bonds.

36. The isolated nutritive protein of any one of claims 1 to 35, wherein the first polypeptide sequence does not comprise N-linked glycosylation.

37. The isolated nutritive protein of any one of claims 1 to 36, wherein the first
30 polypeptide sequence does not comprise O-linked glycosylation.

38. The nutritive protein of any one of claims 1 to 37, wherein the first polypeptide sequence is resistant to aggregation.

39. The nutritive protein of any one of claims 1 to 38, wherein the first polypeptide sequence is anionic at pH 7.

5 40. The nutritive protein of any one of claims 1 to 39, wherein the first polypeptide sequence has an aqueous solubility at pH 7 of at least 12.5 g/L.

41. The nutritive protein of any one of claims 1 to 40, wherein the first polypeptide sequence has a calculated solvation score of -20 or less.

10 42. The nutritive protein of any one of claims 1 to 41, wherein the first polypeptide sequence has a calculated aggregation score of 0.75 or less.

15 43. The nutritive protein of any one of claims 1 to 41, wherein the first polypeptide sequence has a calculated aggregation score of 0.5 or less.

44. The nutritive protein of any one of claims 1 to 43, wherein the first polypeptide sequence comprises at least 20 amino acids.

20 45. The nutritive protein of any one of claims 1 to 43, wherein the first polypeptide sequence comprises at least 50 amino acids.

46. The nutritive protein of any one of claims 1 to 45, wherein the first polypeptide sequence is selected from:

25 A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of Tables 9-11;

B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of Tables 9-11;

30 C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of Tables 9-11;

D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

5 F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of Tables 6-8;

H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of Tables 6-8; and

10 I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of Tables 6-8.

47. The nutritive protein of any one of claims 1 to 45, wherein the first polypeptide sequence is selected from:

15 A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of Tables 9-11;

B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of Tables 9-11;

20 C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of Tables 9-11;

D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

25 E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11;

G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of Tables 6-8;

30 H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of Tables 6-8; and

I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of Tables 6-8.

48. The nutritive protein of any one of claims 1 to 45, wherein the first polypeptide sequence is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from:

- A. the amino acid sequence of a protein listed in at least one of Tables 9-11;
- B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of Tables 9-11; and
- C. the amino acid sequence of a fragment listed in at least one of Tables 6-8.

49. The nutritive protein of any one of claims 1 to 48, wherein the nutritive protein consists of the first polypeptide sequence.

50. The nutritive protein of any one of claims 1 to 48, further comprising a polypeptide tag for affinity purification.

51. The nutritive protein of claim 50, wherein the tag for affinity purification is a polyhistidine-tag.

52. An isolated nutritive protein comprising at least 40% by weight essential amino acids, and

wherein the nutritive protein has a protein digestibility corrected amino acid score (PDCAAS) of at least about 1.4.

53. The nutritive protein of claim 52, wherein the nutritive protein has a PDCAAS of from about 1.4 to about 1.8.

54. The nutritive protein of claim 52, wherein the nutritive protein comprises a weight ratio of at least 50% or 60% essential amino acids.

55. The nutritive protein of claim 54, wherein the nutritive protein has a PDCAAS of from about 1.4 to about 2.2.

56. The nutritive protein of claim 52, wherein the nutritive protein comprises a weight ratio of at least 70% essential amino acids.

57. The nutritive protein of claim 56, wherein the nutritive protein has a PDCAAS
5 of from about 1.4 to about 2.5.

58. The nutritive protein of claim 52, wherein the nutritive protein comprises a weight ratio of at least 80% essential amino acids.

59. The nutritive protein of claim 58, wherein the nutritive protein has a PDCAAS
10 of from about 1.4 to about 2.9.

60. The nutritive protein of claim 52, wherein the nutritive protein comprises a weight ratio of at least 90% essential amino acids.

15

61. The nutritive protein of claim 60, wherein the nutritive protein has a PDCAAS of from about 1.4 to about 3.2.

62. The nutritive protein of any one of claims 52 to 61, further comprising at least
20 one of every non-essential amino acid.

63. The nutritive protein of any one of claims 52 to 61, further comprising at least one of every non-essential and conditionally-essential amino acid.

64. The nutritive protein of claim 62 or 63, further comprising a nutritionally
25 balanced ratio of non-essential amino acids.

65. The nutritive protein of claim 64, wherein the nutritive protein comprises a weight ratio of each non-essential amino acid to the total number of amino acids that is
30 equivalent to the weight ratio of each non-essential amino acid to the total number of amino acids in a benchmark protein.

66. The nutritive protein of claim 65, wherein the benchmark protein is selected from whey protein, egg protein and soy protein.

67. The nutritive protein of claim 52, wherein the nutritive protein comprises
5 100% essential and conditionally essential amino acids.

68. The nutritive protein of claim 67, wherein the nutritive protein has a PDCAAS of from about 1.4 to about 3.6.

10 69. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises at least 70% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof.

15 70. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises at least 95% homology to at least 50 amino acids of a naturally occurring protein or fragment thereof.

20 71. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises at least 70% homology to a naturally occurring protein or fragment thereof.

72. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises at least 95% homology to a naturally occurring protein or fragment thereof.

25 73. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises a naturally occurring protein or fragment thereof.

30 74. The nutritive protein of any one of claims 52 to 68, wherein the nutritive protein comprises an altered amino acid sequence compared to the amino acid sequence of a naturally-occurring reference protein and a weight ratio of essential amino acids to total amino acids present in the nutritive protein that is higher than the weight ratio of essential amino acids to total amino acids present in the naturally-occurring reference protein.

75. The nutritive protein of claim 74, wherein the nutritive protein comprises at least one essential amino acid residue substitution of a non-essential amino acid residue in the naturally occurring reference protein.

5 76. The nutritive protein of any one of claims 52 to 75, wherein the nutritive protein is not an allergen.

77. The nutritive protein of any one of claims 52 to 76, wherein the nutritive protein has less than 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 75, 80, 85, or 90% global
10 homology to a known allergen.

78. The nutritive protein of any one of claims 52 to 77, wherein the nutritive protein is not a toxin.

15 79. The nutritive protein of claim 78, wherein the nutritive protein has less than 50% global homology to a known toxin.

80. The nutritive protein of any one of claims 52 to 79, wherein the nutritive protein has a simulated gastric digestion half-life shorter than whey.

20 81. The nutritive protein of claim 80, wherein the nutritive protein has a simulated gastric digestion half-life of less than 60 minutes or less than 30 minutes.

82. The nutritive protein of claim 81, wherein the nutritive protein has a simulated
25 gastric digestion half-life of less than 10 minutes.

83. The nutritive protein of any one of claims 52 to 82, wherein the nutritive protein is completely digested in simulated gastric fluid.

30 84. The protein of any one of claims 52 to 83, wherein the nutritive protein comprises at least one protease recognition site selected from a pepsin recognition site, a trypsin recognition site, and a chymotrypsin recognition site.

85. The protein of any one of claims 52 to 84, wherein the nutritive protein comprises 0 or 1 cysteine residues.

5 86. The nutritive protein of any one of claims 52 to 85, wherein the nutritive protein comprises no disulfide bonds.

87. The nutritive protein of any one of claims 52 to 86, wherein the nutritive protein does not comprise N-linked glycosylation.

10 88. The nutritive protein of any one of claims 52 to 87, wherein the nutritive protein does not comprise O-linked glycosylation.

89. The nutritive protein of any one of claims 52 to 88, wherein the nutritive protein is resistant to aggregation.

15

90. The nutritive protein of any one of claims 52 to 89, wherein the nutritive protein is anionic at pH 7.

20 91. The nutritive protein of any one of claims 52 to 90, wherein the nutritive protein has an aqueous solubility at pH 7 of at least 12.5 g/L.

92. The nutritive protein of any one of claims 52 to 91, wherein the nutritive protein has a calculated solvation score of -20 or less.

25 93. The nutritive protein of any one of claims 52 to 92, wherein the nutritive protein has a calculated aggregation score of 0.75 or less.

94. The nutritive protein of any of claims 52 to 92, wherein the nutritive protein has a calculated aggregation score of 0.5 or less.

30

95. The nutritive protein of any one of claims 52 to 94, wherein the nutritive protein comprises at least 20 amino acids.

96. The nutritive protein of any one of claims 52 to 94, wherein the nutritive protein comprises at least 50 amino acids.

97. The nutritive protein of any one of claims 52 to 96, wherein the nutritive protein is selected from:

A. an amino acid sequence comprising the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

B. an amino acid sequence comprising a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

C. an amino acid sequence comprising a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

D. an amino acid sequence comprising a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

E. an amino acid sequence comprising a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

F. an amino acid sequence comprising a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

G. an amino acid sequence comprising the amino acid sequence of a fragment listed in at least one of **Tables 6-8**;

H. an amino acid sequence comprising a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and

I. an amino acid sequence comprising a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

98. The nutritive protein of any one of claims 52 to 96, wherein the nutritive protein is selected from:

A. an amino acid sequence consisting of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

B. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

C. an amino acid sequence consisting of a mutein of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

D. an amino acid sequence consisting of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

E. an amino acid sequence consisting of a modified derivative of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

F. an amino acid sequence consisting of a mutein of a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

G. an amino acid sequence consisting of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**;

H. an amino acid sequence consisting of a modified derivative of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**; and

I. an amino acid sequence consisting of a mutein of the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

99. The nutritive protein of any one of claims 52 to 96, wherein the nutritive protein is at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% homologous to an amino acid sequence selected from

A. the amino acid sequence of a protein listed in at least one of **Tables 9-11**;

B. a fragment of at least 20 amino acids of the amino acid sequence of a protein listed in at least one of **Tables 9-11**; and

C. the amino acid sequence of a fragment listed in at least one of **Tables 6-8**.

100. The nutritive protein of any one of claims 52 to 97 and 99, further comprising a polypeptide tag for affinity purification.

101. The nutritive protein of claim 100, wherein the tag for affinity purification is a polyhistidine-tag.

102. The nutritive protein of any one of claims 1 to 101, wherein the nutritive protein is recombinant.

103. An isolated nucleic acid comprising a nucleic acid sequence that encodes a nutritive protein of any one of claims 1 to 102.

5 104. The isolated nucleic acid of claim 103, wherein the isolated nucleic acid is selected from genomic DNA, cDNA, sense RNA and antisense RNA.

105. The isolated nucleic acid of claim 104, wherein the isolated nucleic acid is genomic DNA.

10 106. The isolated nucleic acid of claim 104, wherein the isolated nucleic acid is cDNA.

107. The isolated nucleic acid according to any one of claims 103 to 106, further comprising an expression control sequence operatively linked to the nucleic acid sequence
15 that encodes the nutritive protein.

108. A vector comprising a nucleic acid sequence that encodes a nutritive protein according to any one of claims 1 to 102.

20 109. The vector of claim 108, further comprising an expression control sequence operatively linked to the nucleic acid sequence that encodes the nutritive protein.

110. A recombinant microorganism comprising at least one of a nucleic acid of any one of claims 103 to 107 and a vector of any one of claims 108 and 109.
25

111. The recombinant microorganism according to claim 110, wherein the recombinant microorganism is a prokaryote.

112. The recombinant prokaryote according to claim 111, wherein the prokaryote is
30 heterotrophic.

113. The recombinant prokaryote according to claim 111, wherein the prokaryote is autotrophic.

114. The recombinant prokaryote according to claim 111, wherein the prokaryote is a bacteria.

5 115. A method of making a nutritive protein of any one of claims 1 to 102, the method comprising culturing a recombinant microorganism of any one of claims 110 to 114 under conditions sufficient for production of the nutritive protein by the recombinant microorganism.

10 116. The method of claim 115, further comprising isolating the nutritive protein from the culture.

117. A nutritive composition comprising a nutritive protein of any one of claims 1 to 102 and at least one second component.

15 118. The nutritive composition of claim 117, wherein the second component is selected from a protein, a polypeptide, a peptide, a free amino acid, a carbohydrate, a fat, a mineral or mineral source, a vitamin, a supplement, an organism, a pharmaceutical, and an excipient.

20 119. The nutritive composition according to claim 118, wherein the second component is a protein.

25 120. The nutritive composition according to claims 119, wherein the protein is a nutritive protein.

30 121. The nutritive composition according to claim 118, wherein the second component is a free amino acid selected from essential amino acids, non-essential amino acids, branch chain amino acids, non-standard amino acids and modified amino acids.

122. The nutritive composition of claim 121, wherein the at least one second component is a free amino acid selected from essential amino acids.

123. The nutritive composition according to claim 121, wherein the second component is a free amino acid selected from branch chain amino acids.

124. The nutritive composition according to claim 121, wherein the second
5 component is a free amino acid selected from non-essential amino acids.

125. The nutritive composition of claim 121, wherein the at least one second component is Leu.

10 126. The nutritive composition of claim 118, wherein the at least one second component is a lipid.

127. The nutritive composition of claim 126, wherein the lipid is selected from a fat, oil, triglyceride, cholesterol, phospholipid, and fatty acid.

15 128. The nutritive composition of claim 118, wherein the at least one second component is selected from a mineral and a vitamin.

129. The nutritive composition of claim 118, wherein the at least one second
20 component is a supplement.

130. The nutritive composition of claim 118, wherein the at least one second component is an organism.

25 131. The nutritive composition of claim 118, wherein the at least one second component is a pharmaceutical.

132. The nutritive composition according to claim 118, wherein the second component is an excipient.

30 133. The nutritive composition according to claim 132, wherein the excipient is selected from a buffering agent, a preservative, a stabilizer, a binder, a compaction agent, a

lubricant, a dispersion enhancer, a disintegration agent, a flavoring agent, a sweetener, a coloring agent.

134. A nutritive composition of any one of claims 117 to 133, wherein the nutritive
5 composition is formulated as a liquid solution, slurry, suspension, gel, paste, powder, or solid.

135. A method of making a nutritive composition of any one of claims 117 to 134,
comprising providing a nutritive protein according to any one of claims 1 to 102 and
10 combining the nutritive protein with the at least one second component.

136. A method of maintaining or increasing of muscle mass, muscle strength, and
functional performance in a subject, the method comprising providing to the subject a
sufficient amount of a nutritive protein of any one of claims 1 to 102, a nutritive composition
15 of any one of claims 117 to 134, or a nutritive composition made by the method of claim 135.

137. A method of maintaining or achieving a desirable body mass index in a
subject, the method comprising providing to the subject a sufficient amount of a nutritive
protein of any one of claims 1 to 102, a nutritive composition of any one of claims 117 to
20 134, or a nutritive composition made by the method of claim 135.

138. A method of treating cachexia in a subject, the method comprising providing
to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 102, a
nutritive composition of any one of claims 117 to 134, or a nutritive composition made by the
25 method of claim 135.

139. The method of any one of claims 136 to 138, wherein the subject is of elderly,
critically-medically ill, and suffering from protein-energy malnutrition.

30 140. A method of providing protein to a subject with protein-energy malnutrition,
the method comprising providing to the subject a sufficient amount of a nutritive protein of
any one of claims 1 to 102, a nutritive composition of any one of claims 117 to 134, or a
nutritive composition made by the method of claim 135.

141. A method of increasing thermogenesis in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 102, a nutritive composition of any one of claims 117 to 134, or a nutritive composition made
5 by the method of claim 135.

142. A method of inducing at least one of a satiation response and a satiety response in a subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 102, a nutritive composition of any one of claims
10 117 to 134, or a nutritive composition made by the method of claim 135.

143. The method of claim 141 or claim 142, wherein the subject is obese.

144. The method of any one of claims 136 to 143, wherein the nutritive protein of
15 any one of claims 1 to 102, a nutritive composition of any one of claims 117 to 134, or a nutritive composition made by the method of claim 135 is consumed by the subject in coordination with performance of exercise.

145. A method of treating at least one of cachexia, sarcopenia and frailty in a
20 subject, the method comprising providing to the subject a sufficient amount of a nutritive protein of any one of claims 1 to 102, a nutritive composition of any one of claims 117 to 134, or a nutritive composition made by the method of claim 135.

146. The method of any one of claims 136 to 145, wherein the nutritive protein of
25 any one of claims 1 to 102, a nutritive composition of any one of claims 117 to 134, or a nutritive composition made by the method of claim 135 is consumed by the subject by an oral, enteral, or parenteral route.

147. A method of making a nutritive protein of any one of claims 1 to 102, the method comprising chemically synthesizing the protein.
30

148. A method of making a nutritive protein of any one of claims 1 to 102, the method comprising isolating the protein.

149. The isolated nutritive protein of any one of claims 1 to 102, wherein the first polypeptide sequence is at least 25 amino acids in length.




FIG. 1




FIG. 2




FIG. 3





FIG. 4






FIG. 6

7/19





FIG. 7






FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17






FIG. 18

19/19

WHOLE PROTEINS/FRAGMENTS - RELATIVE LOG LIKELIHOOD OF EXPRESSION

**FIG. 19**

WHOLE PROTEINS/FRAGMENTS - RELATIVE LOG LIKELIHOOD OF SOLUBLE EXPRESSION

**FIG. 20**