

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>G01N 33/50</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/25872</b>
			<b>(43) International Publication Date:</b> 10 November 1994 (10.11.94)
<b>(21) International Application Number:</b> PCT/US94/04754 <b>(22) International Filing Date:</b> 2 May 1994 (02.05.94) <b>(30) Priority Data:</b> 08/056,456      3 May 1993 (03.05.93)      US <b>(71) Applicant:</b> THE GOVERNMENT OF THE UNITED STATES OF AMERICA, as represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES [US/US]; Office of Technology Transfer, National Institutes of Health, Box OTT, Bethesda, MD 20892 (US). <b>(72) Inventors:</b> ALKON, Daniel, L.; 6701 Bonaventure Court, Bethesda, MD 20817 (US). ETCHEBERRIGARAY, Rene; 12221 Village Square Terrace, Rockville, MD 20852 (US). ETSURO, Ito; 1-45-106, Minami 18-jo, Nishi 17-chome, Chuo-ku, Sapporo 060 (JP). GIBSON, Gary, E.; 60 Fernwood Road, Larchmont, NY 10538 (US). <b>(74) Agents:</b> FEILER, William, S. et al.; Morgan & Finnegan, 345 Park Avenue, New York, NY 10154 (US).			<b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>

**(54) Title:** CELL TEST FOR ALZHEIMER'S DISEASE**(57) Abstract**

The present invention is a method for the diagnosis of Alzheimer's disease using human cells. Specifically, the method detects differences between potassium channels in cells from Alzheimer's patient and normal donors, and differences in intracellular calcium concentrations between Alzheimer's and normal cells in response to chemicals known to increase intracellular calcium levels.



YC



AC



AD



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

CELL TEST FOR ALZHEIMER'S DISEASEFIELD OF THE INVENTION

5           The present invention relates to methods for  
diagnosing Alzheimer's disease. The technique utilizes  
newly discovered differences between cells from healthy  
donors and those with Alzheimer's disease. In one method,  
differences in the existence of functional potassium  
10 channels are assessed. In another method, differences in  
intracellular calcium levels in response to  
depolarization by a potassium channel blocker are  
assessed. In yet another method, differences in  
intracellular calcium levels in response to a chemical  
15 known to increase intracellular calcium levels by  
releasing calcium from intracellular stores are assessed.

BACKGROUND OF THE INVENTION

Alzheimer's disease is associated with extensive  
20 loss of specific neuronal subpopulations in the brain  
(Sims, N.R., et al. (1987) Annals of Neurology 21:451),  
with memory loss being the most universal symptom.  
(Katzman, R. (1986) New England Journal of Medicine  
314:964). Alzheimer's disease has been linked to a  
25 genetic origin. (Schellenberg, G.D., et al. (1992)  
Science 258:668; Li, G., et al. (1991) Psychiatric Clinics  
of North America 14:267; St. George-Hyslop, P.H., et al.  
(1989) Neurobiology of Aging 10:417; St. George-Hyslop,  
P.H., et al. (1987) Science 235:885). Early-onset  
30 familial forms of the disease exhibit a genetic defect on  
chromosome 21. (St. George-Hyslop, P.H., et al. (1987)).

Cellular changes, leading to neuronal loss and  
the underlying etiology of the disease, remain unknown.  
Proposed causes include environmental factors, (Perl, D.P.  
35 (1985) Environmental Health Perspective 63:149; Katzman,

- 2 -

o  
R. (1986)), including metal toxicity, (Perl, D.P., et al. (1980) Science 208:297), defects in  $\beta$ -amyloid protein metabolism, (Shoji, M., et al. (1992) Science 258:126; Joachim, C.L. and Selkoe, D.J. (1992) Alzheimer Disease Assoc. Disord. 6:7; Kosik, K.S. (1992) Science 256:780; Selkoe, D.J. (1991) Neuron 6:487; Hardy, H. and Allsop, D. (1991) Trends in Pharmacological Science 12:383), and abnormal calcium homeostasis and/or calcium activated kinases. (Mattson, M.P., et al. (1992) Journal of Neuroscience 12:376; Borden, L.A., et al. (1991) Neurobiology of Aging 13:33; Peterson, E., et al. (1989) Annals of New York Academy of Science 568:262; Peterson, C., et al. (1988) Neurobiology of Aging 9:261; Peterson, C., et al. (1986) Proceedings of the National Academy of Science 83:7999).

Alzheimer's disease is well characterized with regard to neuropathological changes. However, abnormalities have been reported in peripheral tissue supporting the possibility that Alzheimer's disease is a systemic disorder with pathology of the central nervous system being the most prominent. (Rizopoulos, E., et al. (1989) Neurobiology of Aging 10:717; Peterson (1986)).

Potassium channels have been found to change during memory storage. (Etcheberrigaray, R., et al. (1992) Proceeding of the National Academy of Science 89:7184; Sánchez-Andrés, J.V. and Alkon, D.L. (1991) Journal of Neurobiology 65:796; Collin, C., et al. (1988) Biophysics Journal 55:955; Alkon, D.L., et al. (1985) Behavioral and Neural Biology 44:278; Alkon, D.L. (1984) Science 226:1037). This observation, coupled with the almost universal symptom of memory loss in Alzheimer's patients, led to the investigation of potassium channel function as a possible site of Alzheimer's disease pathology and to the current invention.

- 3 -

°

The so-called patch clamp technique and improvements thereof, have been developed to study electrical currents in cells. The method is used to study ion transfer through channels. To measure these currents, the membrane of the cell is closely attached to the opening of the patch micropipette so that a very tight seal is achieved. This seal prevents current from leaking outside of the patch micropipette. The resulting high electrical resistance across the seal can be exploited to perform high resolution current measurements and apply voltages across the membrane. Different configurations of the patch clamp technique can be used. (Sakmann, B. and Neker, E. (1984) Annual Review of Physiology 46:455).

Currently, there is no laboratory diagnostic test for Alzheimer's disease. Therefore, there is a great need for a method to rapidly and clearly distinguish between Alzheimer's patients, normal aged people, and people suffering from other neurodegenerative diseases, such as Parkinson's, Huntington's chorea, Wernicke-Korsakoff or schizophrenia. Although some investigators have suggested that calcium imaging measurements in fibroblasts were of potential clinical use in diagnosing Alzheimer's disease (Peterson et al. 1986, 1988, supra), other researchers using similar cell lines and techniques, have shown no difference in calcium levels in Alzheimer's and normal control fibroblasts. (Borden et al. 1991, supra). Thus, the latter work refutes the findings of the former work.

The methods for diagnosing Alzheimer's disease of the present invention using cells isolated from patients are needed and will greatly improve the now very complicated clinical diagnostic process for Alzheimer's disease. These methods are especially important because they are able to distinguish patients with Alzheimer's

- 4 -

disease from patients with other neurodegenerative diseases.

#### SUMMARY OF THE INVENTION

5           The invention provides a method for assaying for Alzheimer's disease using cells isolated from patients. In one embodiment of the invention, the presence or absence of a particular potassium channel is measured. In a cell from a healthy control, potassium channels with  
10   slope conductances of 113 pS (pico Siemens) and 166 pS are present and functional. In Alzheimer's cells, the 113 pS potassium channel is missing or nonfunctional.

          In a second embodiment of the present invention, the effect of potassium channel blockers specific for the  
15   113 pS potassium channel on intracellular calcium levels is assessed. In this method, intracellular calcium levels are found to be elevated in response to potassium channel blockers in normal cells, but not in cells from donors with Alzheimer's disease. The preferred potassium channel  
20   blocker is tetraethylammonium ("TEA") at a final extracellular concentration of 100 mM. However, other potassium channel blockers which specifically block the 113 pS potassium channel may also be used. Furthermore, when TEA is used, other final concentrations of TEA may be  
25   used as long as the level of TEA causes intracellular calcium levels to be elevated in normal cells, but not in cells from donors with Alzheimer's disease.

          In a third embodiment of the invention, sample cells from a patient are contacted with an activator of  
30   intracellular calcium release, in an amount sufficient to release calcium from intracellular storage sites, and the resulting increase in intracellular calcium levels is measured. In this embodiment, both normal cells and cells from Alzheimer's patients exhibit an increase in  
35   intracellular calcium; however, the increase in

- 5 -

°

Alzheimer's patients is much greater. When an inositol-1,4,5,-trisphosphate ( $IP_3$ ) activator is used to increase intracellular calcium levels, the preferred embodiment utilizes bombesin added to a final extracellular  
5 concentration of 1  $\mu$ m. However, other final concentrations can be used.

As shown in the examples, the combination of the second and third embodiments of the invention can be used in series to provide a very accurate method of diagnosing  
10 AD, with no false positives or false negatives. Furthermore, these methods are able to distinguish patients with Alzheimer's disease from patients with other neurodegenerative diseases. Cells from patients with Parkinson's disease, schizophrenia, Huntington's chorea,  
15 and Wernicke-Korsakoff exhibit responses of normal cells when treated with either TEA or bombesin.

It is not known at the present time if the defects detected by the methods of this invention appear prior to or concurrently with the clinical onset of  
20 Alzheimer's disease. However, if the former is true, it is anticipated that the methods of this invention will have predictive as well as diagnostic utility in the detection of Alzheimer's disease.

#### 25 BRIEF DESCRIPTION OF THE FIGURES

Figures. 1A-1B. 113pS channel. (1A). Cell attached recordings from Alzheimer and control fibroblasts. A potassium channel of ~4.5 pA unitary current size (0 mV pipette potential), with identical  
30 kinetics appeared in age-matched control (AC) and young controls (YC) fibroblasts, but was entirely absent in the recording of AD fibroblasts (1A, bottom) Downward deflections represent the open state. (1B). I/V relationships and slope conductances. I/V relationships and slope conductances (determined by linear regression)  
35

- 6 -

were almost identical within the voltage range explored,  $113.2 \pm 0.9$  pS (mean  $\pm$  S.D.,  $n=8$ ) for YC and  $112.9 \pm 3.2$  pS ( $n=7$ ) for AC fibroblasts.

Figures 2A-2B. 166pS channel. (2A). Cell attached recordings from Alzheimer and control fibroblasts. A second channel (166 pS) was recorded under the same conditions from fibroblasts of all three groups (AD, YC and AC). (2B). I/V relations and slope conductances. I/V relations as well as slope conductances [YC=  $174 \pm 5.7$  pS,  $n=4$ ; AC=  $169.2 \pm 2.8$  pS,  $n=4$ ; AD=  $157.6 \pm 4.7$  pS,  $n=6$  (Mean  $\pm$  S.D.)] were approximately the same across groups. Membrane potential was similar in control ( $-42.6 \pm 5.4$ , Mean  $\pm$  S.D.,  $n=7$ ) and in AD ( $-45.4 \pm 6.9$ ,  $n=3$ ) fibroblasts.

Figures 3A-3C. (3A) and (3B). Percent of cells responding to the addition of 50 mM potassium chloride and average  $[Ca^{2+}]_i$  (nM) of responding cells. High potassium-induced depolarization caused  $[Ca^{2+}]_i$  elevation (at least 100% increase) in all three groups (AD  $N=13$  cell lines; AC  $N=10$ , YC  $N=6$ ). The proportion of responding cells and the  $[Ca^{2+}]_i$  peak values were significantly higher in YC ( $n=183$  cells) fibroblasts ( $\chi^2=14.22$ ,  $p < 0.001$ ), as compared to AC ( $n=299$ ) and AD ( $n=268$ ) fibroblasts (3A and 3B). (3C). Sample traces of time courses of the  $Ca^{2+}$  response in cells after the addition of 50 mM KCl. The  $[Ca^{2+}]_i$  peak occurs 10 to 15 seconds after stimulation, returning to basal levels after 100 seconds. No responses were observed if external  $[Ca^{2+}]$  was lowered ["nominally  $Ca^{2+}$  free" solution, 5 mM EGTA was added (estimated free  $Ca^{2+} = 0.04 \mu M$ )], or  $Ca^{2+}$  channel blockers (0.1 mM  $LaCl_3$ , 10 mM  $CoCl_2$ , 10 mM  $NiCl_2$ , 10 mM  $CdCl_2$  or 10  $\mu M$  nifedipine) were added before stimulation ("0  $Ca^{2+}$ ").

Figures 4A-4C.  $[Ca^{2+}]_i$  elevation in response to TEA. (4A) Percentage of cells responding to the addition of TEA and (4B) Average  $[Ca^{2+}]_i$  response in the cells after

- 7 -

TEA treatment. 1 mM TEA application elevated  $[Ca^{2+}]_i$  in YC fibroblasts (n= 130 cells) but not in AC (n= 184) or AD fibroblasts (n= 195). 10 mM TEA elevated  $[Ca^{2+}]_i$  in YC (n= 176 cells), AC (n= 231), but not in AD (n= 204) fibroblasts ( $\chi^2$  134.00,  $p < 0.001$ ). Similarly, 100 mM TEA elevated  $[Ca^{2+}]_i$  in YC (n= 532 cells), AC (n= 417), but not in AD (n= 738) fibroblasts,  $\chi^2$  231.44,  $p < 0.001$  (also see Table 2). Basal  $[Ca^{2+}]_i$  levels were virtually the same (S.E. < 2 nM), therefore, standard error bars are not distinguishable from the bar representing the arithmetic mean for those groups. (4C). Time course of  $Ca^{2+}$  responses. The  $[Ca^{2+}]_i$  peak occurs 20 to 30 seconds after 100 mM TEA addition in YC and AC fibroblasts, returning to basal levels after 100 seconds. Note that no response meeting criterion (10% of cells in a line with  $\geq 100\%$  elevation) was observed in AD cells. Similarly, the response was absent in control cells when external  $[Ca^{2+}]$  was lowered.

Figures 5A-5B. (5A).  $Ca^{2+}$  mobilization induced by 1  $\mu M$  bombesin in the absence of extracellular calcium. (5B).  $Ca^{2+}$  responses at 42 sec after 1  $\mu M$  bombesin application. The  $[Ca^{2+}]_i$  levels in AD cells are much larger than in AC and YC cells. The numbers of cell lines (N) are 9, 8 and 6 for AD, AC and YC, respectively. The values are means  $\pm$  S.E.M.

Figures 6A-6B. (6A).  $Ca^{2+}$  responses induced by 1  $\mu M$  bombesin in the presence of extracellular calcium. 1  $\mu M$  bombesin elicited a fast peak of  $[Ca^{2+}]_i$ , followed by a sustained phase for YC and AC cells, but not for AD cells, in the presence of extracellular 2.5 mM  $CaCl_2$ . The arrow indicates drug application. (6B). Bar graph illustrating differences evident 90 seconds after bombesin application. In the presence of normal extracellular calcium (2.5 mM), a sustained calcium entry follows the initial bombesin response in control cells but is completely absent in AD

- 8 -

fibroblasts. The difference evident 90 seconds after bombesin application is shown and has a significance level of  $p < 0.001$ .

#### DETAILED DESCRIPTION OF THE INVENTION

5           The invention concerns methods of diagnosing Alzheimer's disease (AD). These methods are based upon detecting the absence of a particular potassium ion channel in the cells of an AD patient; upon differences in intracellular calcium ion concentration in AD and non-AD  
10 cells in response to potassium channel blockers specific for the potassium ion channel that is absent in the cells of an AD patient; and differences between AD and non-AD cells in response to activators of intracellular calcium release such as activators of inositol-1,4,5-trisphosphate  
15 ( $IP_3$ ).

          The first embodiment of the invention is based upon the discovery by the inventors that cells from people not suffering from AD have (at least) two types of functional potassium channels, with conductances of 113 pS  
20 (pico Siemens) and 166 pS, as measured by the patch clamp technique (see Example 1). The 113 pS channel is either missing or not functioning in people with AD. The first embodiment of the invention involves diagnosing AD by determining whether cells of the patient have a  
25 functioning 113 pS potassium channel. The presence of a functioning 113 pS potassium channel indicates that the patient does not have AD. However, the absence of a functioning 113 pS potassium channel indicates that the patient does have AD.

30           In this embodiment of the invention, a suitable method of recording electrical conductances in the cells must be used to detect functional potassium channels in cells. Any technique which can measure electrical conductances in a cell can be used. Examples include  
35 intracellular microelectrode recording (indirect

- 9 -

o

measurement), two microelectrode voltage clamp, and single microelectrode voltage clamp. The patch clamp technique, as described herein, is a preferred method for measuring electrical conductance in small structures. In an embodiment of the invention, the cell attached mode of the patch clamp technique is used to record the existence of potassium channels and the inside-out and outside-out patch configurations are used to record the sensitivity of potassium channels to various chemicals.

10 The second embodiment of the invention concerns another method for diagnosing AD. In this second embodiment, the cells are contacted with a potassium channel blocker that blocks the 113 pS channel but not the 166 pS channel. This blocker may substantially block the 113 pS channel but not substantially block the 166 pS channel. An example of such a blocker is TEA, or tetraethylammonium. The blocker has the effect in non-AD cells of transiently increasing intracellular  $\text{Ca}^{2+}$  concentrations. In AD cells, the blocker has substantially no effect, allowing for variation within observational or technical error. In contrast, the intracellular calcium ion concentration increases several fold in non-AD cells after being exposed to 100 mM TEA (see Fig. 4B). The intracellular  $\text{Ca}^{2+}$  concentration can be measured in various ways, such as by adding fluorescent indicators or absorbance indicators or by using a  $\text{Ca}^{2+}$  electrode. Preferably, because of ease of operation, fluorescent indicators are used.

30 In this embodiment of the invention, the cells are first cultured with a  $\text{Ca}^{2+}$  indicator, such as quin or fura-2, that fluoresces with an intensity proportional to the calcium concentration. The cells are then contacted with a select potassium channel blocker that has the ability to block the 113 pS channel but not the 166 pS channel. The fluorescence intensity of the cells before

35

- 10 -

and after the addition of the potassium channel blocker is measured. In cells from people not suffering from AD the fluorescence intensity increases rapidly, peaks and then drops back down (Fig. 4C). This shows that the blocker has the effect of increasing, transiently, the calcium ion concentration. In cells from AD patients, the fluorescence intensity is substantially the same before and after the blocker is added. This is a reflection of the fact that the 113 pS channel is missing or non-functional in AD patients and thus potassium ion channel blockers that block the 113 pS channel, but not the 166 pS channel, do not have any effect on AD cells.

As mentioned above, the select potassium channel blocker used in this second embodiment of the invention is one that has the ability to block the 113 pS potassium channel but that has little or no effect on the 166 potassium channel. One example of such a blocker is TEA, with any biologically compatible counter anion. Preferably, the counterion is chloride. Other suitable potassium channel blockers can be easily found using the following method. Using the patch clamp technique described in Example 1, the 113 pS and 166 pS channels are detected in a viable human cell. The candidate potassium channel blocker is added to the culture containing the cells, and the patch clamp technique is used again. If the 166 pS channel is still functional, but the 113 pS channel is not, then the candidate blocker is suitable for use in this invention. Candidate potassium channel blockers include the known potassium channel blockers charybdotoxin, apamin, dendrotoxin, kalidotoxin, MCD-peptide, scyllatoxin, barium, cesium, leiurotoxin I and noxiustoxin. As shown in Example 2, TEA concentrations between 10 mM and 100 mM worked well. It is easy to extend this range of workable concentrations by using AD and non-AD control cells.

- 11 -

Example 2 exemplifies the second embodiment of the invention for diagnosing AD using a select potassium channel blocker, TEA, and measuring the effect on intracellular calcium ion. This method is so simple, with a yes or no answer, that the exemplified sophisticated apparatus is not required to make the diagnosis. Any method which will tell one if the intracellular calcium ion concentrations has increased or not as a result of contact with the select potassium ion channel blocker will suffice to give a diagnosis. In the preferred method, fluorescent calcium ion indicators are used. In this case, any method which will tell one if the fluorescence of the indicator has increased or not as a result of contact of the cells with the select potassium channel blockers will suffice. Any method used must be able to make the measurements in the short time available. The calcium ion influx peaks a short time after contact with the blocker, and then decreases to the baseline value. In Example 2, the time it takes to peak is less than one minute.

A simpler method for detecting a fluorescent calcium ion indicator would involve using a fluorimeter, a device with a light source for exciting the calcium ion indicator and a light meter for measuring the intensity of the fluorescence. Fluorimeters are well known and commercially available. At the simplest level, the calcium ion indicator is added to the cells taken from the patient (either fresh or expanded in culture). After an hour or so of being in contact with the indicator (at about 2 micromolar concentration) the cells in suspension are placed in the fluorimeter and the fluorescence intensity from the indicator is measured. Then the select potassium channel blocker is added; if TEA is used, it is added to a concentration of about 100 mM. The fluorescence is measured again. If the intensity, within

- 12 -

o

a time period between 20 seconds and 40 seconds, is substantially the same as before the TEA was added (taking account of changes in volume due to the addition of the TEA), then a positive diagnosis of AD is made. If the intensity increases within 30 seconds and subsides after another 30 seconds, then the patient does not have AD.

It is within the skill of the art to improve the simple scheme outlined above. For example, one could use a fluorimeter with dual sample holders, in which the difference in fluorescence from two samples is measured. Starting with identical samples of patient's cells (after incubation with the indicator) in each sample holder, the select potassium channel blocker is added to only one of the samples. If there is no change in the difference signal (that is, it remains as essentially zero), a diagnosis of AD is made. If the difference signal changes significantly, then the patient does not have AD. The advantage of the differences method is that it has a built in control which increases the accuracy of the measurement. It is still within the skill of the art to add the select potassium channel blocker automatically and to make more than one measurement at a time; i.e., to automate the method for a commercial medical laboratory. Before making any diagnoses using the methods taught here, the methods should be optimized for the particular apparatus and conditions in the laboratory by using non-AD and AD control cells, which are commercially available.

The third embodiment of the invention is yet another method of diagnosing AD. This method concerns the effect of agents that activate inositol-1,4,5,-trisphosphate ( $IP_3$ ) or otherwise induce the release of calcium from intracellular storage sites. Such storage sites include the endoplasmic reticulum and other organelles that have receptors for  $IP_3$ . The preferred  $IP_3$  activator is bombesin. Other agents that activate the

- 13 -

release of calcium from intracellular stores which are useful in the invention include thrombin, bradykinin, prostaglandin  $F_{2\alpha}$  and vasopressin. See, e.g., Berridge, M.J. and Irvine, R.F. (1984) Nature 312:135).

5           It has been discovered that cells from people not suffering from AD and cells from people suffering from AD both transiently release calcium ion in response to bombesin, but the resulting intracellular calcium concentration is much larger in AD cells than in non-AD  
10 cells. The determination is easily made using any method of measuring intracellular calcium ion concentration, as discussed above with respect to the second embodiment of the invention. Again, the use of fluorescent calcium indicators is the preferred method. The same experimental  
15 setup as described above for measuring fluorescence intensity can be used, i.e., a fluorimeter. In this method, it is also possible to standardize the fluorescence apparatus using non-AD and AD cells as controls. In this way, later measurements of just the  
20 patient's cells can provide a diagnosis. Alternatively, the patient's cells can be compared with non-AD cells as a control.

Example 3 exemplifies the third embodiment of the invention concerning the diagnosis of AD using  
25 activators of  $IP_3$  and measuring their effect on calcium ion release into the cytosol from intracellular storage sites after contact with said activators. The amount of released calcium is larger in AD cells compared to non-AD cells. The increase in intracellular calcium  
30 concentration is transient: the concentration peaks soon after contact with the activator and is back to baseline value with 90 seconds. This effect is enhanced when the extracellular calcium ion concentration is zero or near zero (which is generally accomplished by washing the cells  
35 with BSS nominally free of calcium, however, other methods

- 14 -

of tying up or negating the effect of the extracellular calcium ions can be used, such as adding EDTA, or adding a calcium channel blocker such as nifedipine, respectively). After contact with an  $IP_3$  activator, such as bombesin, the intracellular calcium ion concentration in AD cells reaches a higher peak value and takes longer to return to the baseline value than either young or aged control cells (Fig. 5A). In the experimental setup described in Example 3, it was found that 42 seconds after the bombesin was added to the cells that the difference between the intracellular calcium ion concentrations in AD cells and in control cells was at a maximum, and that at that time period, i.e., at 42 seconds after bombesin was applied, the concentration of calcium ions was always greater than 300 nM in AD cells and was always less than 300 nM in control non-AD cells (Fig. 5B). Basal levels of both AD and non-AD fibroblasts were at  $80 \text{ nM} \pm 0.5 \text{ nM}$ . However, it should be noted that control values might differ from 80 nM, necessitating a criterion level of calcium signal greater or less than 300 nM. Furthermore, differences in measuring conditions might require a time longer or briefer than 42 seconds to show maximal differences between the calcium signals of AD and non-AD fibroblasts.

Again, it is not necessary to use the sophisticated methods and apparatus exemplified herein. This method of diagnosing AD can be performed more simply. One need not measure the absolute concentration of intracellular calcium; a measurement of its relative value will also work. In Example 3, the basal level of intracellular calcium ion concentrations in resting (i.e., nonactivated) cells was the same for both AD and control non-AD cells,  $80 \text{ nM} \pm 0.5 \text{ nM}$ . Thus, at the time where the concentration differences between AD and non-AD cells was maximum (i.e., at 42 seconds using bombesin and the inventors' apparatus, but the time would need to be worked

- 15 -

out empirically for different activators and different setups) the intracellular calcium concentration in non-AD cells would be less than  $(300/80 =) 3.75$  times the basal level whereas the intracellular calcium concentration in AD cells would be greater than  $(300/80 =) 3.75$  times the basal level. Using commercially available AD and non-AD cells, one can easily determine the time at which the calcium concentrations are maximally different between AD and non-AD cells. This involves measuring relative intracellular calcium concentrations for resting cells, adding bombesin or another  $IP_3$  activator, following the relative calcium ion concentrations for a minute or so, and finding the time (after the activator is added) at which the difference in relative calcium ion concentrations is at its maximum. Then, for any real sample from a patient, one simply needs to measure the relative basal intracellular calcium concentration by any means known in the art, add the activator to its prescribed concentration (about 1 micromolar for bombesin), wait the predetermined time and again measure the relative intracellular calcium concentration. If the ratio of the intracellular calcium concentration "after" the addition of the activator to the intracellular calcium concentration "before" the addition of the activator is greater than 3.75, the patient has AD; if it is less than 3.75, the patient does not have AD. It is not necessary to determine the time of maximal difference in calcium concentrations -- any time where there is a reproducible difference between these ratios can be used. It is only necessary to work out the particular ratios for the time chosen from known AD and non-AD control cells.

The calcium ion indicators used in the second and third embodiments include any compounds which can enter the cell, are biocompatible, and which can bind to calcium ions to produce a species whose concentration is

- 16 -

°  
easily measured using any physico-chemical means and is proportional to the calcium ion concentration. Preferably the means is fluorescence or absorbance. Preferable fluorescent indicators are the commercially available  
5 indicators fura-2 AM, fura-2 pentapotassium salt, quin-2, and indo-1 from Molecular Probes (Eugene, OR). The Chemical Abstracts name for fura-2, AM is 5-oxazolecarboxylic acid, 2-(6-(bis(2-((acetyloxy)methoxy)-2-oxoethyl)amino)-5-(2-(2-(bis(2-((acetyloxy)methoxy)-2-oxoethyl)amino)-5-methylphenoxy)ethoxy)-2-benzofuranyl)-, (acetyloxy)methyl ester. The Chemical Abstracts name for  
10 fura-2, pentapotassium salt is 5-oxazolecarboxylic acid, 2-(6-(bis(carboxymethyl)amino)-5-(2-(2-(bis(carboxymethyl)amino)-5-methylphenoxy)ethoxy)-2-benzofuranyl)-. Other fluorescent calcium indicators  
15 include Fluo-3, Rhod-2, Calcium Green<sup>TM</sup>, Calcium Orange<sup>TM</sup>, Calcium Crimson<sup>TM</sup> Fura Red<sup>TM</sup> and Calcium Green Dextran<sup>TM</sup> (Molecular Probes (Eugene, OR)). Generally, the cells are incubated with the indicators at a concentration of about  
20 2 micromolar for about 60 minutes. An absorbance indicator which may be used is arsenazo. Finally, calcium levels could also be measured for this invention with calcium electrodes inserted into the cells.

In the exemplified embodiment of the invention,  
25 fluorescence was measured using an imaging system under the control of a personal computer. For excitation, 340 nm and 380 nm band path filters with a neutral-density filter were used. Images of fluorescence were obtained using a dichroic mirror, barrier filter and objective  
30 lens. The whole image can be recorded or portions thereof. A Hamamatsu Photonics Argus 50 Calcium Imaging system imaging 60 cells in a microscopic field at 10 x magnification was used. Fluorescence from the cells was quantified in ¼ of the field at 10x magnification. Such  
35 an imaging system (and other similar currently available

- 17 -

o

systems) with its microscope could be custom designed for everyday clinical laboratory analysis of cells' calcium signals. Other instrumentation and/or measurements would have to be adapted for the use of other calcium  
5 indicators.

In the methods of the invention, the cells that are taken from the patient can be any viable cells. Preferably they are fibroblasts; buccal mucosal cells; blood cells such as erythrocytes, lymphocytes, and  
10 lymphoblastoid cells; or nerve cells such as olfactory neurons. The cells may be fresh or may be cultured (as described in the examples). The fibroblast potassium channel dysfunction and resulting absence of TEA-induced calcium signals described herein suggest that AD, which  
15 primarily affects brain cells, is likely to alter potassium channel function in many different types of cells in the body. Similarly, AD is likely to alter calcium released by bombesin and related agents in many different types of cells in the body. The methods  
20 described herein to measure potassium channel function and calcium release, therefore, should be applicable for AD diagnosis using other cell types.

A punch skin biopsy could be used to obtain skin fibroblasts from a patient. These fibroblasts might be  
25 analyzed directly with the techniques described herein or be introduced into cell culture conditions. The resulting cultured fibroblasts would then be analyzed as described for the cultured fibroblasts obtained from the Coriell Cell Repositories described below. Other steps would be  
30 required to prepare other types of cells which might be used for analysis such as buccal mucosal cells, nerve cells such as olfactory cells, blood cells such as erythrocytes and lymphocytes, etc. For example, blood cells can be easily obtained by drawing blood from  
35 peripheral veins. Cells can then be separated by standard

- 18 -

procedures (e.g., by using a cell sorter, centrifugation, etc.) and later analyzed in suspension or on a solid support (e.g., in petri dishes).

The present invention will now be described by way of examples, which are meant to illustrate, but not limit, the scope of the invention.

#### Example 1

##### Patch-clamp Diagnostic Test

10 Cultured skin fibroblasts (described in Table 3) from the Coriell Cell Repositories (Camden, NJ) were grown under highly standardized conditions. Cristafallo, V.J. and Chapentier, R.J. (1980) Tissue Culture Methods 6:117. The following cell lines were used for the experiments:  
15 Young Control Fibroblasts ("YC") 3652, 3651, 2987, 4390, 3377, 8399 (21.5 $\pm$ 2.8 years, Mean  $\pm$ S.D); Age-matched Control Fibroblasts ("AC") 3524, 6010, 6842, 7603, 9878 (65.2 $\pm$ 6.0 years); and Alzheimer's Disease Fibroblasts ("AD") 6848, 7637, 5809, 8170, 6840, 8243, 6263 (60.6 $\pm$ 6.8  
20 years). Five AD lines were from familial patients. Some of the lines (2 AC and 4 AD) were from Canadian kindred.

In agreement with the literature, the data indicate the time to phase out does not vary between the AD and control lines (YC and AC). Cells were seeded  
25 (approximately 5 cells per mm<sup>2</sup>) in 35 mm Nunc petri dishes in Dulbecco's Modified Eagle Medium (DMEM, Gibco), supplemented with 10% fetal calf serum and used when cell density was equivalent for all cell lines, between days 2 and 4 after plating. On average, fibroblasts from AD  
30 patients and controls took the same time to reach erosion density (50 cells/mm<sup>2</sup>).

Patch-clamp experiments were performed at room temperature (21-23°C), following standard procedures set forth in Sakmann, B. and Neher, E. (1983) Single Channels Recordings (Plenum New York) and Kukuljan, M., et al.  
35

- 19 -

o  
(1991) J. Membrane Biol. 119:187. Before recordings, culture medium was replaced with the following solution: 150 mM NaCl, 5 mM KCl, 2 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 10 mM HEPES (NaCl) pH=7.4. Pipettes were made from Blue Tip capillary  
5 tubes (I.D. 1.1-1.2 mm) using a BB-CH Mecnex puller, and then filled with a high potassium solution of 140 mM KCl, 2 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 10 mM HEPES (NaOH), pH=7.4. Pipette resistances were approximately 6 MΩ. Records were obtained using an Axopatch-1C amplifier (dc-10 kHz),  
10 stored on tape (Toshiba PCM-video recorder), and later transferred to a personal computer using an Axolab interface. Only recordings lasting for at least 3 minutes were considered for final analysis. The pClamp suite of programs was used for single-channel data acquisition and  
15 analysis. Amplifier, interface and software were obtained from Axon Instruments (Foster City, CA).

In the cell-attached mode, two types of potassium channels were recorded from human skin fibroblasts. Since pipettes were filled with a high  
20 potassium solution, potassium currents were inward as expected, and their reversal potential approximately corresponded to the cell resting potential. A potassium channel (113 pS) of approximately 4.5 pA unitary current size (0 mV pipette potential), with identical kinetics  
25 appeared in YC and AC fibroblasts, but was entirely absent in the recording of AD fibroblasts (Fig. 1A). Downward deflections represent the open state. I/V relationships of the same channels in Fig. 1A (Fig. 1B) and slope conductances (determined by linear regression) were almost  
30 identical within the voltage range explored,  $113.2 \pm 0.9$  pS (Mean $\pm$ S.D., n=8) for YC and  $112.9 \pm 3.2$  pS (n=7) for AC fibroblasts.

A second channel (166 pS) was recorded under the same conditions from fibroblasts of all three groups (Fig.  
35 2A). I/V relations (Fig. 2B) as well as conductance

- 20 -

o

(YC=173.4±5.7 pS, n=4; AC= 169.2±2.8 pS, n=4; AD=157.6±4.7 pS, n=6 (Mean±S.D.)) were approximately the same across groups. Membrane potential was similar in control (-42.6±5.4, Mean±S.D., n=7) and in AD (-45.4±6.9, n=3) fibroblasts.

Both channels had linear voltage-current relationships, with slope conductances of 113 pS and 166 pS respectively (Figs. 1A-1B and 2A-2B). At 0 mV pipette potential, the channels could easily be identified by their unitary current size (Figs. 1A and 2A) and by their percentages of open time, approximately 60% for the 113 pS K<sup>+</sup> channel and approximately 10% for the 166 pS K<sup>+</sup> channel. For both channels, the percentages of open time showed no significant voltage-dependence (+60 to -40 mV pipette potential). The 113 pS K<sup>+</sup> channel was found in 47% of YC cells (n=30) and 94% of the AC cells (n=17), while it was never found in AD fibroblasts (n=24) ( $\chi^2 = 18.96$ ,  $p < 0.001$  (Table 1)). There were no AD cell lines (N=6) that had fibroblasts with an observable 113 pS channel. By contrast, all AC cell lines (N=5) and three of six YC cell lines had fibroblasts with observable 113 pS channels ( $\chi^2=11.93$ ,  $p<0.005$  (Table 2)). The 166 pS channel found was similar frequency in all three groups ( $\chi^2=0.89$ , N.S. (Tables 1 and 2)).

The 113 pS channel found to be "absent" in the AD fibroblasts, could be present but not functional. Such dysfunction could involve structural changes in the channel and/or alteration in processes involved in channel activity regulation.

Using cell-free patches, following the method described above, it was observed that both channels were sensitive to 50 mM Ba<sup>2+</sup> (inside-out, n=4 for each channel), but only the 113 pS channel was sensitive (outside-out, n=4 YC, n=3 AC) to the K<sup>+</sup> channel blocker tetraethylammonium (TEA). The TEA-blockade of the 113 pS

- 21 -

channels (possibly together with other channels) significantly affects membrane potential since control cells (n=4) depolarized 13-20 mV after 100 mM TEA addition.

5

Table 1  
Number of Cells

Condition	Total	113 pS K <sup>+</sup> Channel	166 pS K <sup>+</sup> Channel
Young Controls	30	14 (47%)	6 (20%)
Aged Controls	17	16 (94%)	6 (35%)
Alzheimer Patients	24	0 (0%)	8 (33%)

10

Table 2  
Number of Cell Lines

Condition	Total	113 pS K <sup>+</sup> Channel	166 pS K <sup>+</sup> Channel
Young Controls	6	3	4
Aged Controls	5	5	3
Alzheimer Patients	7	0	4

15

20

When using control cells, it is best to use age-matched control cells.

25

#### Example 2

##### TEA-Ca<sup>2+</sup> Diagnostic Test

Cultured skin fibroblasts (described in Table 3) from the Coriell Cell Repositories (Camden, NJ) were grown as described in Example 1.

30

Thirteen AD, ten AC, and six YC were used for the calcium-imaging experiments. Culture medium was replaced and washed three times with basal salt solution ("BSS") consisting of 140 mM NaCl, 5 mM KCl, 2.5 mM CaCl<sub>2</sub>, 1.5 mM MgCl<sub>2</sub>, 5 mM glucose, 10 mM HEPES (NaOH), pH 7.4.

35

- 22 -

Nominally  $\text{Ca}^{2+}$  free BSS was prepared as BSS without adding  $\text{CaCl}_2$ .

Fura-2 (acetyloxymethyl ester) (Fura-2AM) was purchased from Molecular Probes (Eugene, OR) and stored as a 1 mM solution in dimethylsulfoxide. Fura-2AM was added to a final concentration of 2  $\mu\text{M}$  and cells were incubated at room temperature (21°-23°C) for 60 minutes. After incubation, cells were washed at least three times with BSS at room temperature before  $[\text{Ca}^{2+}]$  determinations. Fluorescence was measured with a Hamamatsu ARGUS 50 imaging system (Hamamatsu Photonics, Japan) under the control of a personal computer (Hamamatsu imaging software package). Excitation at 340 nm and 380 nm was attenuated with neutral density filters. Fluorescent images were obtained with a 400 nm dichroic mirror and a 510 nm long-pass barrier filter. The objective lens was an X10 Nikon UV fluor. Fluorescence was measured within a uniformly illuminated fraction (1/4) of the whole image.

The averaged  $\text{Ca}^{2+}$  responses within 15 x 15 pixels in cytosolic and in nuclear cellular compartments obtained were quantified with ratios between emitted 510 nm fluorescence activated at 340 nm and fluorescence emitted at 510 nm with activation at 380 nm. These ratios were transformed to absolute values of  $[\text{Ca}^{2+}]$ , after calibration based on the following equation:

$$R = R_{\max} + (R_{\min} - R_{\max}) / (1 + ([\text{Ca}^{2+}]_i / K_d)^b).$$

Here R denotes fluorescence intensity illuminated by 340 nm divided by fluorescence intensity illuminated by 380 nm ( $F_{340}/F_{380}$ ), and  $R_{\max}$  and  $R_{\min}$  are the values of R when the concentration of calcium is at a maximum and a minimum (i.e., the maximum and minimum value measurable by the machine under the measuring conditions), respectively.  $K_d$  is a dissociation constant of fura-2 for  $\text{Ca}^{2+}$  and was determined as 240 nM. The value of b, which determined the degree of asymmetry, was 1.2. TEA

- 23 -

application caused a minimum of 100%  $[Ca^{2+}]_i$  elevation in at least 18% of cells in every control cell line except one young control. A response of 100%  $[Ca^{2+}]_i$  elevation in at least 10% of cells in a line was, therefore, considered to be a conservative criterion for a positive response. Only one AD cell line had cells with any response (100%  $[Ca^{2+}]_i$  elevation in 4% of cells), well below the criterion).

Depolarization of the fibroblasts by perfusion in elevated external potassium caused greater elevation of intracellular  $Ca^{2+}$  ( $[Ca^{2+}]_i$ ) in YC as compared to AC and AD cells (Fig. 3A-3C). This depolarization-induced  $[Ca^{2+}]_i$  elevation was eliminated by lowering external calcium or by adding calcium channel blockers (Fig. 3C). High  $K^+$ -induced depolarization caused a marked  $[Ca^{2+}]_i$  elevation (at least 100% increase) in all three groups (AD, n= 13 cell lines; AC, n= 10; YC, n= 6). The proportion of responding cells and the  $[Ca^{2+}]_i$  peak values were significantly higher in YC (n= 183 cells) fibroblasts ( $\chi^2=14.22$ ,  $p < 0.001$ ), as compared to AC (n= 299) and AD (n= 268) fibroblasts. The  $[Ca^{2+}]_i$  peak occurs 10 to 15 seconds after stimulation, returning to basal levels after 100 seconds. No responses were observed if external calcium was lowered by addition of "nominally  $Ca^{2+}$  free" solution or 5 mM EGTA (estimated free  $Ca^{2+}=0.04 \mu M$ ) or  $Ca^{2+}$  channel blockers (0.1 mM  $LaCl_3$ , 10 mM  $CoCl_2$ , 10 mM  $NiCl_2$ , 10 mM  $CdCl_2$  or 10  $\mu M$  nifedipine) before stimulation.

Depolarization of control fibroblasts by TEA also caused  $[Ca^{2+}]_i$  elevation, that was eliminated by lowering external calcium or by adding calcium channel blockers. AD fibroblasts, however, only showed  $[Ca^{2+}]_i$  elevation in elevated external potassium and had no  $[Ca^{2+}]_i$  response with addition of even 100 mM TEA. Every AC cell line (N=10) and all but one YC cell line (N=6) had cells responding to TEA, while none of the thirteen AD cell

- 24 -

lines examined had cells responding to 100 mM TEA ( $\chi^2=25.66$ ,  $p<0.001$ ) (Tables 3 and 5).

Table 3

Number of Cell Lines

Condition	Total	Increase in $[Ca^{2+}]_i$ with 100 mM TEA
Young Controls	6	5
Aged Controls	10	10
Alzheimer's Patients	13	0

1 mM TEA application elevated  $[Ca^{2+}]_i$  in YC fibroblasts (n=130 cells) but not in AC (n=184) or AD (n=195) fibroblasts. 10 mM TEA elevated  $[Ca^{2+}]_i$  in YC (n=176) and AC (n=231) but not in AD fibroblasts (n=204). Similarly 100 mM TEA elevated  $[Ca^{2+}]_i$  in YC (n=532) and AC (n=417), but not in AD fibroblasts (n=738) ( $\chi^2=231.44$ ,  $p<0.001$ ). At least 417 cells were explored in each experimental group (Table 4). The  $[Ca^{2+}]_i$  values of the responding cell were similar in YC and AC cells after 10 and 100 mM TEA addition. Basal  $[Ca^{2+}]_i$  levels were virtually the same (S.E. < 0.5 nM), therefore standard error bars are not distinguishable from the bar representing the arithmetic mean for those groups (Fig. 4B). Time courses of  $Ca^{+2}$  response shows that the  $[Ca^{2+}]_i$  peak occurs 20 to 30 seconds, after 100 mM TEA addition in YC and AC fibroblasts, returning to basal levels after 100 seconds. No response was observed in AD cells (10% of cells in a line with  $\geq 100\%$  elevation). Similarly, the response was absent in control cells when external  $[Ca^{2+}]$  was lowered (Fig. 4C).

- 25 -

Table 4  
Number of Cells

Condition	Total	Increase in $[Ca^{2+}]_i$ with 100 mM TEA
Young Controls	532	145 (27%)
Aged Controls	417	119 (29%)
Alzheimer's Patients	738	4 (0.5%)

TEA-induced  $[Ca^{2+}]_i$  elevations were repeated using a coded subsample that included Alzheimer's and control fibroblasts. Experiments and analyses were conducted without the experimenter's knowledge of the cell lines identity. The results were in complete agreement with the non-blind sample. None of the blindly examined AD cell lines (N=11) showed  $[Ca^{2+}]_i$  elevation in response to TEA and all but one of the control cell lines (4 AC and 6 YC) had TEA responses ( $\chi^2=17.33$ ,  $p < 0.001$  (Table 5)).

Since  $[Ca^{2+}]_i$  elevation in response to high potassium was virtually the same for AC and AD cells, the lack of AD cells response to TEA is almost certainly due to dysfunction of  $K^+$  channels and not to  $Ca^{2+}$  channel dysfunction.

The  $[Ca^{2+}]_i$  measurements are in agreement with the patch-clamp measurements insofar as they both indicate potassium channel dysfunction in the AD fibroblasts. See Table 5.

- 26 -

Table 5

	Line #	Age	Gender	Race	Diag. Criteria	113 K <sup>+</sup> Channel	TEA Response	
							Non Blind	Blind
5	Alzheimer's Disease Fibroblasts							
	AG06840 <sup>+</sup> <sup>1</sup>	56	M	W	Clinical - Fam. H.	-	-	-
	AG06848 <sup>+</sup> <sup>2</sup>	55	F	W	Clinical - Fam. H. <sup>*</sup>	-	-	N.T.
	AG07637 <sup>+</sup>	55	F	W	Clinical - Fam. H.	-	-	-
10	AG08170 <sup>+</sup>	56	M	W	Clinical - Fam. H.	-	-	-
	AG06844 <sup>+</sup>	59	M	W	Clinical - Fam. H. <sup>*</sup>	N.T.	N.T.	-
	AG04400 <sup>‡</sup>	61	F	W	Clinical - Fam. H.	N.T.	N.T.	-
	AG04401 <sup>‡</sup>	53	F	W	Clinical - Fam. H. <sup>*</sup>	N.T.	-	-
	AG05809	63	F	W	Clinical - Fam. H.	-	-	N.T.
15	AG08243	72	M	W	Clinical - No Fam. H.	-	-	-
	AG07375	71	M	W	Clinical - No Fam. H.	N.T.	-	-
	AG07376	59	M	W	Clinical - No Fam. H.	N.T.	-	-
	AG06263	67	F	W	Clinical - No Fam. H.	-	-	-
	AG07377	59	M	W	Clinical - No Fam. H.	N.T.	N.T.	-
20	Age-Matched Control Fibroblasts							
	GM03524	67	F	B	Normal	+	+	N.T.
	AG06010	62	F	W	Normal	+	+	+
	AG06842 <sup>+</sup>	75	M	W	Normal-Fam. H.	+	N.T.	N.T.
25	AG07603 <sup>+</sup>	61	F	W	Normal-Fam. H.	+	+	N.T.
	AG09878	61	F	B	Normal	+	+	+
	AG08044	58	F	B	Normal	N.T.	+	N.T.
	AG6241	61	M	W	Normal	N.T.	+	N.T.
	AG4560	59	M	W	Normal	N.T.	+	N.T.
30	GM04260	60	M	W	Normal	N.T.	+	N.T.
	AG07141	66	F	W	Normal	N.T.	N.T.	+
	AG11363	74	F	W	Normal	N.T.	N.T.	+
	Young Control Fibroblasts							
	GM03652	24	M	W	Normal	+	+	+
35	GM03651	25	F	W	Normal	+	+	+

- 27 -

	GM02987	19	M	W	Normal	-	-	-
	Line #	Age	Gender	Race	Diag. Criteria	113 K <sup>+</sup> Channel	TEA Response Non Blind	Blind
5	GM04390	23	F	W	Normal	+	+	+
	GM03377	19	M	W	Normal	-	+	+
	GM08399	19	F	?	Normal	-	+	+

Alzheimer's fibroblasts were from familial (N=8) and non-familial cases (N=5). Five (+) are members of the Canadian family 964, only 1 and 2 are immediate relatives (sibs). "+" are members (sibs) of family 747. Autopsy confirmed Alzheimer's Disease in three cases (\*). Two of the age-matched control (N=11) cell lines are unaffected members of the Canadian family (964). All young control lines (N=6) are from normal and without AD family history individuals. Criterion  $[Ca^{2+}]_i$  responses (to 100 mM TEA), indicates as +, were observed in all AC lines used and in all but one of the YC lines. The presence of the 113 pS K<sup>+</sup> channel is indicated by the "+" sign. None of the AD lines exhibited "positive" response. A blind protocol was conducted to measure TEA responses in Alzheimer's (N=11) and control (YC=6, AC=4) fibroblasts. The results exactly reproduced those of the non-blind sample: no AD cells line exhibited TEA responses and 9 out 10 control cells showed TEA responses,  $\chi^2=17.33$ ,  $p < 0.001$ . The notation "N.T." indicates cell line/conditions that were not tested.

### Example 3

#### Bombesin - $Ca^{2+}$ Diagnostic Test

Human skin fibroblasts listed in Table 3 were used. The average age for the AD cell lines used is  $60.5 \pm 5.9$  years; for the AC cell lines is  $62.3 \pm 9.6$  years; and for the YC cell lines is  $21.5 \pm 2.2$  years. The method

- 28 -

of maintenance for the cells was described in Example 1, i.e., maintained 3-5 days at 37°C in CO<sub>2</sub>/air (5%/95%) to reach a density of 50 cells/mm<sup>2</sup> before calcium measurements. The number of culture passages were less than 19.

Bombesin was purchased from Calbiochem (San Diego, CA). Bombesin was stored as a 1mM solution in distilled water. Fura-2 (acetyloxymethyl ester), fura-2 (pentapotassium salt) and omega-conotoxin ( $\omega$ -CgTX) GVIA were from Molecular Probes (Eugene, OR). Fura-2 AM was stored as a 1mM solution in dimethylsulfoxide; fura-2 pentapotassium salt was stored as a 6mM solution in potassium acetate, and  $\omega$ -CgTX was stored as a 100 $\mu$ M solution in distilled water. All of the chemicals except for phenytoin were maintained at -20°C and protected from light.

The cells were incubated with 2 $\mu$ M fura-2 AM in BSS (described in Example 1) at room temperature (21 - 23°C) for 60 min. After being washed at least three times with BSS, the cells were used for measurement of [Ca<sup>2+</sup>]<sub>i</sub> at room temperature. Cell fluorescence was measured as described in Example 2. Absolute calcium values were calculated as shown in Example 2.

Bombesin was added to the cells at a final concentration of 1 $\mu$ M. Calcium mobilization levels were measured from -30 seconds to 150 seconds after bombesin treatment. (Fig. 5A) The particular experimental set up resulted in a maximum difference in [Ca<sup>2+</sup>]<sub>i</sub> between AD cells and control cells at a time of 42 seconds after bombesin was added.

Forty two (42) seconds after bombesin treatment, in the absence of extracellular Ca<sup>2+</sup>, the [Ca<sup>2+</sup>]<sub>i</sub> levels in Alzheimer's disease cells are much larger (p<0.0001) than in age-matched and young controls. The numbers of cell lines (N) are 10, 8, and 6 for Alzheimer's disease, age-

- 29 -

matched and young cells, respectively. The values are means  $\pm$  S.E.M. (Fig. 5B)

Bombesin stimulated  $\text{IP}_3$ -induced  $\text{Ca}^{2+}$  release from intracellular storage sites in fibroblasts from all groups, but it caused a larger and more prolonged response in AD fibroblasts. This larger and prolonged response in AD cells was independent of extracellular  $\text{Ca}^{2+}$ . On the other hand, the  $\text{IP}_3$ -mediated  $\text{Ca}^{2+}$  responses in AC and YC cells were followed by  $\text{Ca}^{2+}$  entry. When this  $\text{Ca}^{2+}$  entry was diminished by removal of extracellular  $\text{Ca}^{2+}$ , or blocking with inorganic  $\text{Ca}^{2+}$  blockers, the bombesin-elicited  $\text{Ca}^{2+}$  responses in control cells were found to return to the basal level faster than in AD cells (Fig. 5A). The results shown in Fig. 5A are for cells washed with BSS nominally free of  $\text{Ca}^{2+}$ .

Since  $\text{Ca}^{2+}$  influx induced by bombesin was not observed in AD cells, this pathway of  $\text{Ca}^{2+}$  entry following the decrease of stored calcium seems to be altered. This test independently confirmed the diagnoses made by the previously described test based on potassium channel dysfunction. In particular, the  $\text{Ca}^{2+}$  responses at 42 sec after 1  $\mu\text{M}$  bombesin stimulation in AD fibroblasts in the absence of extracellular  $\text{Ca}^{2+}$  were always higher than 300 nM. In contrast, the  $[\text{Ca}^{2+}]_i$  in AC and YC were less than 300 nM and 200 nM, respectively (Fig. 5B).

In a variation on the above experiment,  $\text{Ca}^{2+}$  responses were induced by 1  $\mu\text{M}$  bombesin in the presence of extracellular calcium. In the presence of 2.5 mM extracellular  $\text{CaCl}_2$ , 1  $\mu\text{M}$  bombesin elicited a fast peak of  $[\text{Ca}^{2+}]_i$ , followed by a sustained phase for YC and AC cells, but not for AD cells. (Fig. 6A). This difference was evident 90 seconds after bombesin application and with a significance level of  $p < 0.001$ . (Fig. 6B). This difference in response of AD and non-AD cells to bombesin in the presence of extracellular calcium can be used to

- 30 -

provide a "yes or no" diagnosis of AD. Detection methods similar to those described above with respect to the second embodiment of the invention involving the diagnosis of AD by detecting differences between non-AD and AD cells in response to select potassium channel blockers (e.g., TEA) may be used. Furthermore, the combination of this diagnostic test with any one of the above diagnostic tests further increases the confidence level of a correct diagnosis as AD or non-AD.

10

#### Example 4

##### Responses In Neuropathological Non-AD Fibroblasts

Using the techniques described in Examples 2 and 3, cells from donors with other diseases were measured for intracellular calcium levels in response to either TEA or bombesin.

Fibroblasts from a Parkinson's disease donor had normal TEA (indicated as +) and bombesin responses ("N"), and did not significantly differ from responses observed in the age-matched control group. Fibroblasts from two schizophrenic patients also had normal TEA and bombesin responses. In addition, normal TEA responses were observed in five out of seven cases of Huntington's disease, and the bombesin response was normal in all Huntington's cases. Furthermore, normal TEA and bombesin responses were observed in four out of four cases of Wernicke-Korsakoff disease (Table 6). These responses are significantly different from those of AD fibroblasts to the level of  $p < 0.0001$  (Fisher's exact test). "\*" indicates autopsy confirmation.

35

- 31 -

Table 6

Line #	Age	Gender	Race	Condition	TEA	Bombesin
AG08395	85	F	W	Parkinson's*	+	N
GM01835	27	F	W	Schizophrenia	+	N
GM02038	22	M	W	Schizophrenia	+	N
GM06274	56	F	W	Huntington's	+	N
GM02165	55	M	W	Huntington's	+	N
GM00305	56	F	W	Huntington's	-	N
GM01085	44	M	W	Huntington's	+	N
GM01061	51	M	W	Huntington's	+	N
GM05030	56	M	W	Huntington's	-	N
GM04777	53	M	W	Huntington's	+	N
7504	50	M	W	Wernicke-Kors.	+	N
7505	52	F	W	Wernicke-Kors.	+	N
7507	63	M	W	Wernicke-Kors.	+	N
7508	64	M	W	Wernicke-Kors.	+	N

Every reference cited hereinbefore is hereby incorporated by reference in its entirety.

The invention has been described in detail with particular reference to the preferred embodiments thereof, but it will be understood that the invention is capable of other and different embodiments. As is readily apparent to those skilled in the art, variations and modifications can be affected within the spirit and scope of the invention. Accordingly, the foregoing disclosure and description are for illustrative purposes only, and do not in any way limit the invention, which is defined only by the claims.

- 32 -

We claim:

1. A method of diagnosing Alzheimer's disease (AD) in a patient, said method comprising the following steps:

- a. obtaining a sample of cells from said patient; and
- b. detecting the presence or absence of a functioning 113 pS potassium ion channel;

the absence of said potassium ion channel being indicative of AD.

2. The method of claim 1, wherein said cells are selected from the group consisting of fibroblasts, buccal mucosal cells, neurons, and blood cells.

3. The method of claim 2, wherein said cells are fibroblasts.

4. The method of claim 1, wherein said detecting step b is performed by the patch clamp technique.

5. A method of diagnosing Alzheimer disease (AD) in a patient, said method comprising the following steps:

- a. obtaining a sample of cells from said patient;
- b. measuring the intracellular calcium ion concentration in said cells;
- c. contacting said cells with a select potassium ion channel blocker, said blocker having the ability to block the 113 pS potassium ion channel but

- 33 -

not the 166 pS potassium ion channel;  
and

- d. within a time period less than one minute after said contacting step c, measuring the intracellular calcium ion concentration in said cells by the same method as in step b;

wherein if the intracellular calcium ion concentration does not increase within said one minute time period after said contacting step c, then said patient has AD, if said intracellular calcium ion concentration does increase then said patient does not have AD.

6. A method of diagnosing Alzheimer disease (AD) in a patient, said method comprising the following steps:

- a. obtaining a sample of cells from said patient;
- b. measuring the intracellular calcium ion concentration in said cells;
- c. contacting said cells with a select potassium ion channel blocker, said blocker having the ability to block the 113 pS potassium ion channel but not the 166 pS potassium ion channel;  
and
- d. measuring the intracellular calcium ion concentration in said cells by the same method as in step b at that time where there is a maximum difference in intracellular calcium ion concentration between AD cells and control cells;

wherein the absence of an increase of intracellular calcium ion concentration in the cells from said patient

- 34 -

in the time period between step c and step d indicates that the patient does have AD and wherein an increase of intracellular calcium ion concentration in the cells from said patient in the time period between step c and step d indicates that the patient does not have AD.

7. A method of diagnosing Alzheimer disease in a patient according to claim 6 wherein step d comprises measuring the intracellular calcium ion concentration in said cells within a time period less than one minute after said contacting step c.

8. The method of claim 5 wherein the select potassium ion channel blocker is tetraethylammonium.

9. The method of claim 5, wherein said cells are selected from the group consisting of fibroblasts, buccal mucosal cells, neurons, and blood cells.

10. The method of claim 9 wherein said cells are fibroblasts.

11. The method of claim 5 wherein said measuring steps b and d are performed using a fluorescent calcium ion indicator and a fluorimeter.

12. The method of claim 11 wherein said calcium ion indicator is selected from the group consisting of fura-2 AM, fura-2 pentapotassium salt and quin-2.

13. A method of diagnosing Alzheimer disease (AD) in a patient, said method comprising the following steps:

- a. obtaining a sample of cells from said patient;

- 35 -

- b. measuring the basal level concentration of intracellular calcium ion in said cells;
- c. contacting said cells with an activator of intracellular calcium release;
- d. measuring the intracellular calcium ion concentration in said cells at a predetermined time after the contacting step c; and
- e. comparing the ratio of measured concentrations of step d to step b to previously determined ratios for known AD cells and non-AD cells at said predetermined times;

wherein if the ratio of step e is the same as or greater than the previously determined ratio for known AD cells, the patient has AD, and if the ratio of step e is the same as or less than the previously determined ratio for known non-AD cells, the patient does not have AD.

14. The method of claim 13 wherein the predetermined time of step d is that time when the difference between relative intracellular calcium ion concentrations in AD cells and in control cells is at a maximum.

15. The method of claim 13 wherein the activator of release of intracellular calcium release is the  $IP_3$  activator bombesin.

16. The method of claim 13 wherein the extracellular calcium ion concentration is zero or near zero.

- 36 -

17. The method of claim 13, wherein said cells are selected from the group consisting of fibroblasts, buccal mucosal cells, neurons, and blood cells.

18. The method of claim 17, wherein said cells are fibroblasts.

19. The method of claim 13 wherein said measuring steps b and d are performed using a fluorescent calcium ion indicator and a fluorimeter.


20. The method of claim 19 wherein said calcium ion indicator is selected from the group consisting of fura-2 AM, fura-2 pentapotassium salt, and quin-2.

21. A method of diagnosing Alzheimer disease (AD) in a patient, said method comprising the following steps:

- a. obtaining a sample of cells from said patient;
- b. measuring the basal level concentration of intracellular calcium ion in said cells in the presence of extracellular calcium;
- c. contacting said cells with an activator of intracellular calcium release; and
- d. measuring the intracellular calcium ion concentration in said cells at a predetermined time after the contacting step c;


wherein if the intracellular calcium ion concentration is the same as that measured in step b, then said patient has AD, if said intracellular calcium ion concentration is greater than that measured in step b, then said patient does not have AD.

**FIG. 1A**




2/14

FIG. 1B




3/14

FIG. 2A




4/14

FIG. 2B




5/14

**FIG. 3A**




6/14

FIG. 3B




7/14

FIG. 3C




8/14

FIG. 4A




9/14

FIG. 4B




10/14

FIG. 4C




11/14

FIG. 5A




12/14

**FIG. 5B**


13/14

FIG. 6A



14/14

FIG. 6B



# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 94/04754

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 5 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 5 G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	NEUROSCIENCE RESEARCH, vol.18, December 1993 page 576 E.ITO ET AL. 'A laboratory diagnosis of Alzheimer's Disease with patch-clamp and Ca <sup>2+</sup> -imaging techniques' See abstract 615 ---	1-21
X,P	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES (USA), vol.90, no.17, 1 September 1993 pages 8209 - 8213 R.ETCHEBERRIGARAY ET AL. 'Potassium channel dysfunction in fibroblasts identifies patients with Alzheimer disease' see the whole document --- -/--	1-12



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

30 August 1994

Date of mailing of the international search report

25. 09. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Hitchen, C

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 94/04754

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES (USA), vol.91, no.2, 18 January 1994 pages 534 - 538 E.ITO ET AL. 'Internal Ca <sup>2+</sup> mobilization is altered in fibroblasts from patients with Alzheimer disease' see the whole document ---	13-21
A	NEUROBIOLOGY OF AGING, vol.12, 1991 pages 469 - 473 H-M.HUANG ET AL. 'Inositol Phosphates and Intracellular calcium After Bradykinin Stimulation in Fibroblasts From Young, Normal Aged and Alzheimer Donors' see page 469 - page 470 ---	13-21
A	NEUROBIOLOGY OF AGING, vol.14, no.2, April 1993 pages 177 - 185 A.GROSSMANN ET AL. 'Intracellular calcium Response Is Reduced in CD4+ Lymphocytes in Alzheimer's Disease and in Older Persons with Down's Syndrome' see page 177 - page 179 ---	5,6,9, 11,13, 16,17, 19,21
A	WO,A,89 09600 (MASSACHUSETTS INSTITUTE OF TECHNOLOGY) 19 October 1989 see claims ---	5,6,8
A	ANN. REV. PHYSIOL., vol.46, 1984 pages 455 - 472 B.SAKMANN ET AL. 'Patch Clamp techniques for Studying Ionic Channels in Excitable Membranes' cited in the application See the introduction ---	1,4
A	NEUROBIOLOGY OF AGING, vol.9, no.3, 1988 pages 261 - 266 C.PETERSON ET AL. 'Altered response of Fibroblasts From Aged and Alzheimer Donors to Drugs That Elevate Cytosolic Free Calcium' cited in the application see the whole document ---	5-7,9-12

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 94/04754

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	<p>GERONTOLOGY, vol.39, August 1993 pages 163 - 169 A.F.GHUYSEN-ITARD ET AL. 'Loss of calcium-Homeostatic Mechanisms in Polymorphonuclear Leukocytes of demented and Nondemented Elderly Subjects' see the whole document -----</p>	5-7,9-12

## Information on patent family members

PCT/US 94/04754

Form PCT/ISA:210 (patent family annex) (July 1992)