

(19) **DANMARK**

(10) **DK/EP 2964765 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 15/63 (2006.01)** **C 12 N 5/10 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2019-07-29**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2019-05-08**
- (86) Europæisk ansøgning nr.: **14759649.8**
- (86) Europæisk indleveringsdag: **2014-03-07**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-01-13**
- (86) International ansøgning nr.: **US2014022135**
- (87) Internationalt publikationsnr.: **WO2014138703**
- (30) Prioritet: **2013-03-08 US 201361774982 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Keck Graduate Institute of Applied Life Sciences, 535 Watson Drive, Claremont, CA 91711, USA**
- (72) Opfinder: **TOLSTORUKOV, Ilya I., 174 Marywood Avenue, Claremont, California 91711, USA**
CREGG, James M., 2692 San Andres Way, Claremont, California 91711, USA
- (74) Fuldmægtig i Danmark: **Larsen & Birkeholm A/S Skandinavisk Patentbureau, Banegårdspladsen 1, 1570 København V, Danmark**
- (54) Benævnelse: **GÆRPROMOTORER FRA PICHIA PASTORIS**
- (56) Fremdragne publikationer:
WO-A2-99/67398
WO-A2-2008/128701
WO-A2-2012/129036
US-A1- 2005 164 259
US-A1- 2008 108 108
US-A1- 2012 100 581
STADLMAYR G ET AL: "Identification and characterisation of novel Pichia pastoris promoters for heterologous protein production", JOURNAL OF BIOTECHNOLOGY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 150, no. 4, 1 December 2010 (2010-12-01), pages 519-529, XP027516642, ISSN: 0168-1656, DOI: 10.1016/J.JBIOTEC.2010.09.957 [retrieved on 2010-11-23]
GABRIEL POTVIN ET AL: "Bioprocess engineering aspects of heterologous protein production in Pichia pastoris: A review", BIOCHEMICAL ENGINEERING JOURNAL, vol. 64, 1 May 2012 (2012-05-01), pages 91-105, XP055312900, NL ISSN: 1369-703X, DOI: 10.1016/j.bej.2010.07.017
ANDREAS KÜBERL ET AL: "High-quality genome sequence of CBS7435", JOURNAL OF BIOTECHNOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 154, no. 4, 26 April 2011 (2011-04-26), pages 312-320, XP028240769, ISSN: 0168-1656, DOI: 10.1016/J.JBIOTEC.2011.04.014 [retrieved on 2011-05-06]
DE SCHUTTER, K.: 'Genome Sequence Of The Recombinant Protein Production Host Pichia Pastoris.' NAT

Fortsættes ...

DESCRIPTION

FIELD OF THE DISCLOSURE

[0001] The present invention is related to isolated nucleic acids, expression methods, host cells, and expression vectors. More particularly, the invention relates to nucleic acids isolated from *Pichia pastoris* wherein the nucleic acids have promoter activity. The invention also relates to expression methods, host cells, and expression vectors for using the *Pichia pastoris* promoters to produce proteins.

BACKGROUND AND SUMMARY OF THE INVENTION

[0002] Yeast expression systems can be used to effectively produce proteins, such as enzymes, hormones, and vaccine proteins, in part, because some yeast grow rapidly to high cell densities, are grown in simple and inexpensive media, and are eukaryotes so they can modify proteins in a manner similar to native proteins in mammals. Additionally, with a proper signal sequence, the expressed protein can be secreted into the culture medium for convenient isolation and purification. Some yeast expression systems are also accepted in the food and pharmaceutical industries as being safe for the production of pharmaceuticals and food products, unlike fungal and bacterial expression systems which may in some cases be unsafe, for example, for human food manufacturing.

[0003] Thus, it is beneficial for a variety of industries, such as the food and animal feed industries, the human and animal health industries, and the like, to develop or improve yeast expression systems that can be used to express high levels of proteins to increase yield, reduce the expense of isolation and purification of proteins, and reduce the costs of human and animal health products and food products.

[0004] A variety of types of yeast expression systems have been developed involving either the use of inducible or constitutive expression of proteins using nucleic acids encoding homologous or heterologous proteins, under the control of a yeast promoter. Promoters are regulatory elements that are linked to the 5' end of a nucleic acid encoding a protein, and may interact with various regulatory factors in the host cell (e.g., a yeast host cell) to control transcription of RNA from DNA. Promoters may also control the timing of transcription of RNA from DNA. For example, the AOX 1 promoter has been identified in the yeast *Pichia pastoris*, and is commonly used in yeast expression systems because it is a tightly regulated, strong promoter.

[0005] WO2012/129036A2 describes promoters for high level recombinant expression in fungal host cells.

[0006] US2008/108108 describes novel *P. pastoris pastoris* promoters, and the use thereof to direct expression of proteins in yeast, preferably using a haploid mating strategy.

[0007] Stadlmayer et al (2010), Journal of Biotechnology, vol. 150, no. 4, pages 519-529 describes the identification and characterization of novel *Pichia pastoris* promoters for heterologous protein production.

[0008] Gabriel Potvin et al (2012), Biochemical Engineering Journal, vol. 64, pages 91-105 describes bioprocess engineering aspects of heterologous protein production in *Pichia pastoris*.

[0009] WO2008/1287001A2 describes expression systems.

[0010] Andreas Küberl et al (2011), Journal of Biotechnology, vol. 154, pages 312-320 describes the high-quality genome sequence of *Pichia pastoris* CB7435.

[0011] Due to the importance of yeast expression systems for a variety of industries, including the human pharmaceuticals industry, and the human food and animal feed industries, the improvement of yeast expression systems is the focus of much research and development. Accordingly, the present inventors have identified promoters from *Pichia pastoris* that are particularly effective for use in expression of proteins in yeast. The promoters described herein can be used, for example, in systems for the constitutive expression of proteins.

[0012] The subject matter of the invention is as set out in the claims.

[0013] One aspect of the invention relates to an isolated nucleic acid wherein the isolated nucleic acid comprises a sequence at least 90% identical to SEQ ID NO:2, or at least 90% identical to a fragment thereof, wherein said sequence at least 90% identical to SEQ ID NO:2 or at least 90% identical to a fragment thereof exhibits promoter activity, wherein the nucleic acid comprises the sequence of a constitutive *Pichia pastoris* promoter and wherein the isolated nucleic acid is operably linked to a heterologous coding sequence.

[0014] A further aspect of the invention relates to an expression vector comprising the above isolated nucleic acid.

[0015] A further aspect of the invention relates to a host cell comprising the above isolated nucleic acid or the above expression vector.

[0016] A further aspect of the invention relates to a method of producing a protein, the method comprising the step of culturing in a culture medium a host cell comprising a first expression cassette comprising an isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to a heterologous coding sequence encoding a protein,

wherein the culturing is done under conditions permitting expression of the protein;

wherein the protein is expressed using the first expression cassette in combination with a second expression cassette, said second cassette comprising

1. (a) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising SEQ ID NO: 3 or SEQ ID NO: 4, wherein SEQ ID NO: 3 or SEQ ID NO: 4 have promoter activity; or
2. (b) the isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017]

FIGURE 1 shows the nucleotide sequence of the Chr1-0469 promoter (SEQ ID NO: 1).

FIGURE 2 shows the nucleotide sequence of the UPP promoter (SEQ ID NO: 2).

FIGURE 3 shows the nucleotide sequence of the AOX1 promoter sequence (SEQ ID NO:3).

FIGURE 4 shows the genomic sequence 1000bp upstream of the AOX1 promoter ATG start codon (SEQ ID NO:4).

FIGURE 5 shows the level of GFP expression for various transformants in replica plates.

FIGURE 6 shows the relative promoter activity for pUPP, TEF2, Chr1-0469, GAP, AOX1, and DAS promoters under various conditions.

FIGURE 7 shows the map of a reporter vector. Similar vectors were used with both the alpha and beta-Galactosidase ORF, as well as with, or without a secretion signal (alpha mating factor)

FIGURE 8 shows the nucleotide sequence of forward primers used for isolation of the Chr1-0469 promoter (SEQ ID NO: 1) and the UPP promoter (SEQ ID NO: 2) from *Pichia pastoris* genomic DNA by PCR. The corresponding DNA fragments - products of these PCR reactions - are introduced into the reporter vector with the alpha-Galactosidase ORF (figure 7).

FIGURE 9 (panels A -C) show the level of alpha-Galactosidase expression from isolated transformants with the indicated promoter-reporter constructs.

FIGURE 10 shows the activity of the *E.coli* appA2 phytase in *Pichia pastoris*.

DETAILED DESCRIPTION OF THE ILLUSTRATIVE EMBODIMENTS

[0018] In one aspect, the invention provides an isolated nucleic acid wherein the isolated nucleic acid comprises a sequence at least 90% identical to SEQ ID NO:2, or at least 90% identical to a fragment thereof, wherein said sequence at least 90% identical to SEQ ID NO:2 or at least 90% identical to a fragment thereof exhibits promoter activity, wherein the nucleic acid comprises the sequence of a constitutive *Pichia pastoris* promoter and wherein the isolated nucleic acid is operably linked to a heterologous coding sequence.

[0019] In one embodiment, the isolated nucleic acid sequence is SEQ ID NO:2, or a fragment thereof. In other embodiments, expression vectors, host cells, and DNA constructs comprising these promoter sequences are provided.

[0020] The above-described promoters have been isolated from a yeast strain (i.e., NRRL Y11430) currently classified as a *Pichia pastoris* yeast strain. However, the classification may change at some point to a *Komagataella* species (e.g., *Komagataella phaffii*).

[0021] In another embodiment, a method of producing a protein using these promoter sequences is provided. The method comprises the steps of culturing in a culture medium a host cell comprising a first expression cassette comprising an isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to a heterologous coding sequence encoding a protein, wherein the culturing is done under conditions permitting expression of the protein; wherein the protein is expressed using the first expression cassette in combination with a second expression cassette, said second cassette comprising

1. (a) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising SEQ ID NO: 3 or SEQ ID NO: 4, wherein SEQ ID NO: 3 or SEQ ID NO: 4 have promoter activity; or
2. (b) the isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

[0022] In another illustrative disclosure, an isolated protein produced according to this method is provided.

[0023] The phrase "consists of" or "consisting of" means that the sequence specified by the SEQ ID NO. has no additional nucleotide sequences other than those corresponding to the SEQ ID NO.

[0024] Any yeast expression system known to those skilled in the art can be used in accordance with the present invention. For example, various yeast expression systems are described in U.S. Patent Nos. 6,451,572, 6,841,370, 6,974,690, 7,320,876, 7,078,035, 7,138,260, and PCT Publication No. WO 2007/112739. In any of the embodiments described herein, any of these yeast expression systems can be used. Alternatively, any yeast species or yeast expression system suitable for expression of a protein can be used including yeast

species, such as *Saccharomyces* species (e.g., *Saccharomyces cerevisiae*), *Kluyveromyces* species (e.g., *Kluyveromyces lactis*), *Torulaspora* species, *Yarrowia* species (e.g., *Yarrowia lipolitica*), *Schizosaccharomyces* species (e.g., *Schizosaccharomyces pombe*). In another embodiment, methylotrophic yeast species such as *Pichia* species (e.g., *Pichia pastoris* or *Pichia methanolica*), *Hansenula* species (e.g., *Hansenula polymorpha*), *Torulopsis* species, *Komagataella* species, *Candida* species (e.g., *Candida boidinii*), and *Karwinskia* species can be used. In one embodiment the protein can be expressed in the methylotrophic yeast *Pichia pastoris*. Methylotrophic yeast are capable of utilizing methanol as a sole carbon source for the production of the energy resources necessary to maintain cellular function. Methylotrophic yeast contain genes encoding enzymes for methanol utilization such as the genes encoding alcohol oxidase. Any of these host cells can be a host cell strain that is heterologous to the promoter described herein (i.e., the host cell does not normally contain in nature the promoter described herein).

[0025] A yeast expression system can be used to produce a sufficient amount of the protein intracellularly, or secreted from the yeast cells so that the protein can be conveniently isolated and purified from the culture medium. As used herein, the term "expression" means transcription and/or translation of a nucleic acid in a host cell. A yeast expression system may include, but is not limited to, the yeast host cell and the expression vector (e.g., a DNA construct) used to express the protein. The expression vector can contain a promoter described herein and, as is known in the art, the promoter is heterologous to the expression vector (i.e., the combination does not occur in nature). In one embodiment, secretion of the protein into the culture medium is controlled by a signal peptide (e.g., the yeast α -factor signal peptide, the yeast KILM1 signal peptide, the yeast PHO1 signal peptide, or the yeast SUC2 signal peptide) incorporated into the expression vector and which is capable of directing the secretion of the expressed protein out of the yeast cell. In other embodiments, other signal peptides suitable for facilitating secretion of the protein from yeast cells are known to those skilled in the art. In one aspect, the signal peptide is typically cleaved from the protein after secretion.

[0026] In various embodiments, any expression vector known to the skilled artisan (e.g., a vector that replicates autonomously or integrates into the host genome) and compatible with a yeast expression system can be used. As used herein, the term "vector" means any plasmid, or other vector, in double-stranded or single-stranded form or in linear or circular form that can transform a yeast cell by integration into the yeast cell genome or by existing extrachromosomally (e.g., an autonomously replicating plasmid). As is known in the art, a vector (e.g., expression vector or expression cassette) is a nucleic acid construct used to transform a host cell for expression of a protein, polypeptide, or peptide and the vector is not found in nature in the host cell it transforms.

[0027] In one embodiment, the expression vector has restriction endonuclease cleavage sites for the insertion of DNA fragments (e.g., one or more cloning sites and/or a multiple cloning site), and genetic markers for selection of transformants. For example, the genetic markers for selection of transformants can include a selection marker that allows a transformed yeast to

grow on a medium devoid of a necessary nutrient that cannot be produced by a deficient strain, a selection marker that encodes an enzyme for which chromogenic substrates are known, or a selection marker that provides resistance to a drug, including, but not limited to, G418, Nourseothricin (Nat), Zeocin, Blasticidin, or Hygromycin. In another embodiment, the expression vector has a terminator sequence for transcription termination (e.g., the AOX 1 or HSP150 terminator). In another embodiment, the expression vector has a 3' untranslated region downstream from the protein coding sequence with a polyadenylation site. As used herein, "3' untranslated region" means nucleotide sequences that are not translated into protein and are located downstream from a coding sequence for a protein. Typically, a 3' untranslated region includes regulatory sequences for mRNA processing. In another embodiment, the expression vector has an origin of replication (e.g., a bacterial origin of replication) for use in synthesizing and amplifying the vector, for example, in a bacterial host. Various expression vectors are described in U.S. Patent Nos. 6,451,572, 6,841,370, 6,974,690, 7,320,876, 7,078,035, 7,138,260, and PCT Publication No. WO 2007/112739. The construction and use of expression vectors is described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001). In another embodiment, the expression vector, or a fragment thereof, can be synthesized *de novo* or PCR can be used to amplify and join sections of expression vectors.

[0028] As used herein, "regulatory sequences" means nucleotide sequences that are typically upstream or downstream from the 5' or 3' end, respectively, of a protein coding sequence. Regulatory sequences are not translated into protein. Regulatory sequences include, but are not limited to, sequences that affect RNA processing or stability, such as polyadenylation signal sequences, enhancers, repressor binding sites, and promoters.

[0029] In one embodiment, the protein coding sequence can be operably linked in the expression vector to the promoter sequence capable of directing the expression of the protein, for example, in yeast. As used herein, "operably linked" means functionally linked. As described herein, the promoter can be a "constitutive promoter". A "constitutive promoter" means a promoter that regulates expression of a gene of interest. The term "constitutive promoter" is known in the art. A constitutive promoter can have some inducible activity, but the maximal activity obtained with the promoter is not inducible.

[0030] As used herein "promoter" means a nucleotide sequence typically located upstream from the 5' end of a coding sequence for a protein that controls the transcription of RNA from DNA, in part, by interacting with various regulatory factors that control transcription. In one embodiment, the promoter may be derived from the same species of yeast as the yeast host cell used for protein expression. In another embodiment, the promoter may be derived from a different yeast species than the yeast host cell used for protein expression. In one embodiment, a promoter may include a TATA box sequence that acts as a recognition site to direct initiation of transcription, including, but not limited to one or more transcriptional enhancer elements. The enhancer elements may be proximal or distal to the TATA box sequence and may be in a normal 5' to 3' orientation or may be in a 3' to 5' orientation. In another embodiment, an enhancer element may be an enhancer element native to the

promoter sequence or it may be a heterologous enhancer element inserted into the expression vector construct. An "enhancer element" as used herein is a regulatory element that can stimulate promoter activity.

[0031] In various illustrative embodiments described herein, the promoter can be an isolated nucleic acid wherein the isolated nucleic acid comprises a sequence at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to SEQ ID NO:2, or at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to a fragment thereof, wherein the isolated nucleic acid comprises the sequence of a constitutive *Pichia pastoris* promoter. In another embodiment, the isolated nucleic acid sequence is SEQ ID NO:2, or a fragment thereof wherein the isolated nucleic acid comprises the sequence of a constitutive *Pichia pastoris* promoter.

[0032] As used herein "an isolated nucleic acid" means a nucleic acid that is substantially free of sequences that naturally flank the nucleic acid in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, an isolated nucleic acid in accordance with the invention can contain less than about 2 kb, less than about 1 kb, less than about 0.5 kb, less than about 0.1 kb of nucleotide sequences, less than about 0.05 kb, or no nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the organism from which the isolated nucleic acid is derived.

[0033] As used herein, "a fragment thereof" when referring to the isolated nucleic acid molecule means a fragment of the isolated nucleic acid of SEQ ID NO: 2. In various illustrative embodiments, the fragment can be about 50 nucleotides in length, about 100 nucleotides in length, about 200 nucleotides in length, about 300 nucleotides in length, about 400 nucleotides in length, about 500 nucleotides in length, about 600 nucleotides in length, about 700 nucleotides in length, about 800 nucleotides in length, or about 900 nucleotides in length. In other embodiments, the fragment can extend about 50 nucleotides, about 100 nucleotides, about 200 nucleotides, about 300 nucleotides, about 400 nucleotides, about 500 nucleotides, about 600 nucleotides, about 700 nucleotides, about 800 nucleotides, or about 900 nucleotides upstream from the 3' end of the isolated nucleic acid of SEQ ID NO: 2. In yet other embodiments, the fragment can include about 50 nucleotides, about 100 nucleotides, about 200 nucleotides, about 300 nucleotides, or about 350 nucleotides upstream and/or downstream from the TATA box sequence found in SEQ ID NO: 2.

[0034] In various embodiments, the isolated nucleic acids described herein may be purified by techniques for purification of nucleic acids (e.g., DNA) that are well-known in the art. For example, the nucleic acids may be separated from contaminants by physical methods including, but not limited to, centrifugation, pressure techniques, or by using a substance with affinity for nucleic acids (e.g., DNA), such as, for example, silica beads. After sufficient washing, the isolated nucleic acids may be suspended in either water or a buffer. In other embodiments, commercial kits are available, such as Qiagen™, Nuclisens™, and Wizard™ (Promega), and Promegam™. Methods for purifying nucleic acids are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory

Press, (2001). An isolated nucleic acid as described herein may be an isolated nucleic acid that is also purified, or the isolated nucleic acid may be impure. A "purified nucleic acid" is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

[0035] The isolated nucleic acids described herein are capable of specific hybridization, under appropriate hybridization conditions (e.g., appropriate buffer, ionic strength, temperature, formamide, and MgCl_2 concentrations), to a complementary nucleic acid. The isolated nucleic acids described herein can be modified by substitution, deletion, truncation, and/or can be fused with other nucleic acid molecules wherein the resulting isolated nucleic acids hybridize specifically to the complementary nucleic acids.

[0036] Also within the scope of the disclosure are nucleic acids complementary to the isolated nucleic acids, or fragments thereof, described herein, and those that hybridize to the isolated nucleic acids described herein or those that hybridize to their complements under highly stringent conditions. As used herein, the term "complementary" refers to the ability of purine and pyrimidine nucleotide sequences to associate through hydrogen bonding to form double-stranded nucleic acid molecules. Guanine and cytosine, adenine and thymine, and adenine and uracil are complementary and can associate through hydrogen bonding resulting in the formation of double-stranded nucleic acid molecules when two nucleic acid molecules have "complementary" sequences. The complementary DNA sequences are referred to as a "complement."

[0037] In accordance with the disclosure "highly stringent conditions" means hybridization at 65 °C in 5X SSPE and 50% formamide, and washing at 65 °C in 0.5X SSPE. Conditions for high stringency hybridization are described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001). In some illustrative aspects, hybridization can occur along the full-length of the isolated nucleic acid, or along part of its length, or to a fragment thereof.

[0038] Also included are isolated nucleic acid molecules having about 92%, about 95%, 96%, 97%, 98%, and 99% identity to the isolated nucleic acids described herein, or to a fragment thereof. Determination of percent identity or similarity between sequences can be done, for example, by using the GAP program (Genetics Computer Group, software; now available via Accelrys on <http://www.accelrys.com>), and alignments can be done using, for example, the ClustalW algorithm (VNTI software, InforMax Inc.). A sequence database can be searched using the isolated nucleic acid sequence of interest, or the sequence of a fragment thereof. Algorithms for database searching are typically based on the BLAST software (Altschul et al., 1990). In some embodiments, the percent identity can be determined along the full-length of the isolated nucleic acid.

[0039] Techniques for synthesizing the isolated nucleic acids described herein are well-known in the art and include chemical syntheses and recombinant methods. Such techniques are

described in Sambrook et al., "Molecular Cloning: A Laboratory Manual", 3rd Edition, Cold Spring Harbor Laboratory Press, (2001). Isolated nucleic acid molecules can also be made commercially. Techniques for synthesizing the isolated nucleic acids described herein are well-known in the art. The isolated nucleic acids described herein can be analyzed by techniques known in the art, such as restriction enzyme analysis or sequencing, to determine the sequence of the isolated nucleic acids. Thus, isolated nucleic acids described herein may be synthetic.

[0040] In illustrative aspects, yeast cells are transformed with an expression vector comprising a heterologous nucleic acid encoding the protein of interest operably linked to one of the promoters described herein using procedures well-known to those skilled in the art. The term "transformation" means the transfer of a nucleic acid, or a nucleic acid fragment, into a host cell. In illustrative embodiments, such transformation protocols include electroporation, lithium acetate methods, and use of spheroplasts. In illustrative aspects, the expressed nucleic acid coding sequence can be a heterologous nucleic acid coding sequence. As used herein, a heterologous coding sequence is defined as an artificial or synthetic nucleic acid or a nucleic acid originating from a different species than the species from which the promoter sequence was derived. Thus, a heterologous coding sequence linked to a promoter described herein does not occur in nature.

[0041] In various embodiments, the transformed yeast cells may be grown by techniques including batch and continuous fermentation in a liquid medium or on a semi-solid medium, or a solid medium. Typically, "conditions permitting expression of the protein" as used herein means conditions for batch or continuous fermentation of yeast in a liquid medium, but growth on a semi-solid medium, such as agar, is not excluded. Culture media for yeast cells are known in the art and are typically supplemented with a carbon source (e.g., glucose). A typical yeast culture medium is YPD broth (Sunrise Science Products, Inc.) comprising yeast extract (10 grams), Bacto peptone (20 grams), and dextrose (20 grams). In one illustrative aspect, the transformed yeast cells can be grown aerobically at 30°C in a controlled pH environment (a pH of about 6) and with the carbon source (e.g., glucose) maintained continuously at a predetermined level known to support growth of the yeast cells to a desired density within a specific period of time.

[0042] In one aspect, a method of producing a protein is provided. The method comprises the step of culturing in a culture medium a host cell comprising a first expression cassette comprising an isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to a heterologous coding sequence encoding a protein, wherein the culturing is done under conditions permitting expression of the protein; wherein the protein is expressed using the first expression cassette in combination with a second expression cassette, said second cassette comprising

1. (a) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising SEQ ID NO: 3 or SEQ ID NO: 4, wherein SEQ ID NO: 3 or SEQ ID NO: 4 have promoter activity; or

2. (b) the isolated nucleic acid of SEQ ID NOs: 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

The method can further comprise the step of purifying the protein from the medium of the cultured host cell. As used herein, an "expression cassette" means the elements of an expression vector that direct the yeast cell to make RNA. An expression cassette comprises at least regulatory sequences (e.g., a promoter) and a coding sequence for the RNA and protein (i.e., an open reading frame, ORF). In various illustrative aspects, the protein coding sequence can be from a bacterium, a yeast, a fungus, or a virus.

[0043] In this method embodiment, the protein is expressed using the first expression cassette in combination with a second expression cassette. The second expression cassette comprises 1) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising the sequence of SEQ ID NO:3 or SEQ ID NO:4 wherein SEQ ID NO:3 and SEQ ID NO:4 have promoter activity, or 2) the isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

[0044] In yet another embodiment, the protein can be expressed using the first expression cassette, the second expression cassette, and a third expression cassette. In another illustrative aspect, the third expression cassette can comprise 1) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising the sequence of SEQ ID NO:3 or SEQ ID NO:4 wherein SEQ ID NO:3 and SEQ ID NO:4 have promoter activity, or any other known methanol-regulated promoters, such as AOX 1, AOX 2, FLD, or DAS promoter sequences, or 2) the isolated nucleic acid of any one of SEQ ID NOS: 1 to 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

[0045] In another embodiment, any number of additional expression cassettes can be used and the expression cassettes can comprise 1) the heterologous coding sequence encoding the protein operably linked to an isolated nucleic acid having a sequence comprising the sequence of SEQ ID NO:3 or SEQ ID NO:4 wherein SEQ ID NO:3 and SEQ ID NO:4 have promoter activity, or any other known methanol-regulated promoters, such as AOX 1, AOX 2, FLD, or DAS promoter sequences, or 2) the isolated nucleic acid of any one of SEQ ID NOS: 1 to 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein. In another embodiment, the first expression cassette, the second expression cassette, and the third expression cassette as described above are used in the method, along with a fourth expression cassette, a fifth expression cassette, and a sixth expression cassette wherein all of the expression cassettes comprise the isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to the heterologous coding sequence encoding the protein.

[0046] In still another embodiment, a method of producing one or more proteins is provided. The method comprises the step of culturing in a culture medium a host cell comprising a first expression cassette, a second expression cassette, and, optionally, one or more additional

expression cassettes, wherein each of the expression cassettes comprises 1) a heterologous coding sequence encoding the one or more proteins operably linked to an isolated nucleic acid having a sequence comprising the sequence of SEQ ID NO:3 or SEQ ID NO:4 wherein SEQ ID NO:3 and SEQ ID NO:4 have promoter activity, or 2) the isolated nucleic acid of SEQ ID NO: 2, or a fragment thereof, operably linked to a heterologous coding sequence encoding the one or more proteins, wherein the culturing is done under conditions permitting expression of the one or more proteins. The method can further comprise the step of purifying one of the one or more proteins from the medium of the cultured host cell.

[0047] In any of the embodiments described in the preceding two paragraphs, the isolated nucleic acid can be at least 90%, 92%, 95%, 96%, 97%, 98%, or 99% homologous to the isolated nucleic acid of any of SEQ ID NO: 2, or a fragment thereof. In any of the embodiments described in the preceding two paragraphs, the expression cassettes can be included in one expression vector or in multiple expression vectors. In any of the embodiments described in the preceding two paragraphs, the expression vectors into which the expression cassettes are incorporated can be vectors that replicate autonomously or that integrate into the host cell genome.

[0048] In various illustrative embodiments, the protein encoded by the any of the heterologous coding sequences described herein operably linked to the promoter can be a protein selected from the group consisting of a toxin, an antibody, a hormone, an enzyme, a growth factor, a cytokine, a structural protein, an immunogenic protein (e.g., a vaccine antigen), and a cell signaling protein. In one embodiment, the protein is an enzyme for use in animal feed. In this embodiment, the protein can be selected from the group consisting of a phytase, a mannanase, a galactosidase, an amylase, a glucanase, a cellulase, a protease, and a xylanase. In another embodiment, the protein can be an enzyme useful for glycosylation. In another aspect, more than one protein can be expressed by using multiple expression cassettes, each with a heterologous coding sequence, operably linked to a promoter. However, these protein examples are non-limiting and any protein or peptide or polypeptide capable of being expressed in yeast can be expressed in accordance with the isolated nucleic acids, expression vectors, host cells, DNA constructs, methods, and isolated proteins described herein. The enzymes described above can be from any species (e.g., fungal species, such as a yeast species).

[0049] The yeast-expressed proteins for use in accordance with the present invention can be produced in purified form by conventional techniques. As used herein, "isolated protein" means a purified protein. A purified protein is substantially free from other yeast cell contaminants or contaminants from the culture medium. For example, "substantially free" from other yeast cell contaminants or contaminants from the culture medium means that the protein is at least about 60% pure, at least about 70% pure, at least about 80% pure, at least about 90% pure, at least about 95% pure, at least about 98% pure, about 60% pure, about 70% pure, about 80% pure, about 90% pure, about 95% pure, or about 98% pure (all based on dry weight). Typically, the protein is secreted into the yeast culture medium and is collected from the culture medium.

[0050] In one illustrative embodiment, for purification from the culture medium the protein can, for example, be subjected to ammonium sulfate precipitation followed by DEAE-Sepharose column chromatography. In other embodiments, conventional techniques known to those skilled in the art can be used such as ammonium sulfate or ethanol precipitation, acid extraction, gel filtration, anion or cation exchange chromatography, DEAE-Sepharose column chromatography, hydroxylapatite chromatography, lectin chromatography, affinity chromatography, solvent-solvent extraction, ultrafiltration, and HPLC.

[0051] Alternatively, purification steps may not be required because the protein may be present in such high concentrations in the culture medium that the protein is essentially pure in the culture medium (e.g., 70 to 80% pure). In one embodiment, the protein is collected from the culture medium without further purification steps by chilling the yeast culture (e.g., to about 4 °C to about 8°C) and removing the yeast cells using such techniques as centrifugation, microfiltration, and rotary vacuum filtration. The protein in the cell-free medium can then be concentrated by such techniques as, for example, ultrafiltration and tangential flow filtration.

[0052] In some embodiments where the protein is not secreted into the culture medium, the yeast cells can be lysed, for example, by sonication, heat, or chemical treatment, and the homogenate centrifuged to remove cell debris. The supernatant can then be subjected to ammonium sulfate precipitation, and additional fractionation techniques as required, such as gel filtration, ion exchange chromatography, DEAE-Sepharose column chromatography, affinity chromatography, solvent-solvent extraction, ultrafiltration, and HPLC to purify the protein. It should be understood that the purification methods described above for purification of proteins from the culture medium or from lysed yeast cells are non-limiting and any purification techniques known to those skilled in the art can be used to purify the yeast-expressed protein if such techniques are required to obtain a substantially pure protein.

[0053] Various formulations of the purified protein preparations may be prepared in accordance with the invention. In some embodiments, the proteins can be stabilized through the addition of other proteins (e.g., gelatin and skim milk powder), chemical agents (e.g., glycerol, polyethylene glycol, EDTA, potassium sorbate, sodium benzoate, and reducing agents and aldehydes), polysaccharides, monosaccharides, lipids (hydrogenated vegetable oils), and the like. In one embodiment, proteins for addition to food products or animal feed blends can be dried (e.g., spray drying, drum drying, and lyophilization) and formulated as powders, granules, pills, mineral blocks, liquids, and gels through known processes. In one embodiment, gelling agents such as gelatin, alginate, collagen, agar, pectin and carrageenan can be used.

[0054] In alternate embodiments, the protein expression can be for intracellular expression, such as for enzymatic action in the yeast in a biotransformation process, or for display on the yeast cell surface. For such embodiments, the protein, expressed as described herein, is not purified.

[0055] In various embodiments, the proteins described above are selected from the group consisting of a toxin, an antibody, a hormone, an enzyme, a growth factor, a cytokine, a

structural protein, an immunogenic protein (e.g., a vaccine antigen), and a cell signaling protein. In another embodiment, the proteins described above are selected from the group consisting of an antibody, a hormone, an enzyme, a growth factor, a cytokine, a structural protein, an immunogenic protein (e.g., a vaccine antigen), and a cell signaling protein. In yet another embodiment the coding sequence for a protein, a fragment thereof, a fusion protein (e.g., a chimeric protein), or a peptide, can be used in accordance with the invention. In another embodiment, a modified protein can be expressed, such as a mutated protein or a protein with non-natural amino acids.

[0056] In one embodiment, the toxins can be proteins such as, for example, botulinum toxin or verotoxin-1, and after preparation using the methods, isolated nucleic acids, expression vectors, host cells, and DNA constructs described herein, the toxins can be modified using a targeting agent so that they are directed specifically to diseased cells. In another illustrative aspect, the antibody can be a humanized antibody, an antibody that is not humanized, a nanobody, or an antibody fragment, such as an Fab fragment of an antibody or a single-chain antibody. In another embodiment, the hormone can be, for example, a gonadotropin, an adrenocorticotrophic hormone, a growth hormone, vasopressin, oxytocin, somatostatin, gastrin, or leptin. In another illustrative embodiment, the growth factor can be insulin, epidermal growth factor, fibroblast growth factor, vascular endothelial growth factor, erythropoietin, platelet-derived growth factor, thrombopoietin, or a bone morphogenic protein. In one aspect, the cytokine can be IL-2, IFN- α , IFN- γ , or GM-CSF. In one embodiment, the protein is not a silk toxin. In another illustrative aspect, the vaccine proteins can be any suitable vaccine proteins that are immunogenic in a patient or an animal, including, but not limited to, HPV proteins (e.g., HPV 16 and HPV 18), and tetanus vaccine proteins, as examples. In another illustrative embodiment, the enzymes can be, for example, enzymes for animal feeds as discussed herein, acetylcholinesterase, or cyclooxygenase, or any other useful enzyme that can be expressed in yeast. In another embodiment, structural proteins can be expressed, for example, netrins, actin-binding proteins, or myosin, and, in another embodiment, cell signaling proteins such as *ras* proteins, kinases, the ErbB2 protein (the Her-2 receptor) can be expressed using the methods, isolated nucleic acids, expression vectors, host cells, and DNA constructs described herein.

[0057] In one embodiment, the protein is an enzyme for use in animal feed. In this embodiment, the protein can be selected from the group consisting of a phytase, a mannanase, a galactosidase, an amylase, a glucanase, a cellulase, a protease, and a xylanase, or a combination thereof. For example, a variety of phytases may be expressed according to the methods described herein. Exemplary of phytase genes (i.e., a phytase coding sequence) that can be expressed in accordance with the invention are phytase genes derived from bacteria, filamentous fungi, plants, and yeast, such as the *appA* (Gene Bank accession number M58708) and *appA2* (Gene Bank accession number 250016) genes derived from *Escherichia coli* and the *phyA* and *phyB* genes derived from the fungus *Aspergillus niger*, or any mutant of these genes that retains or has improved myo-inositol hexakisphosphate phosphohydrolase activity (see, for example, Rodriguez et al., Arch. of Biochem. and Biophys. 382: 105-112 (2000)). Exemplary alpha- and beta-galactosidase genes are from *Aspergillus*

species (e.g., *Aspergillus alliaceus*, *Aspergillus oryzae*, *Aspergillus nidulans*, *Aspergillus parasiticus*, and *Aspergillus niger*). Substituted, deleted, and truncated phytase genes, or a fragment thereof, can also be expressed in accordance with the invention.

[0058] In one embodiment, the protein expressed using the methods described herein can be used in animal feed comprising an animal feed blend. In various embodiments, any animal feed blend known in the art can be used such as rapeseed meal, cottonseed meal, soybean meal, and cornmeal. Optional ingredients of the animal feed blend include sugars and complex carbohydrates such as both water-soluble and water-insoluble monosaccharides, disaccharides and polysaccharides. Optional amino acid ingredients that can be added to the feed blend are arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, tyrosine ethyl HCl, alanine, aspartic acid, sodium glutamate, glycine, proline, serine, cysteine ethyl HCl, and analogs, and salts thereof. Vitamins that can be optionally added are thiamine HCl, riboflavin, pyridoxine HCl, niacin, niacinamide, inositol, choline chloride, calcium pantothenate, biotin, folic acid, ascorbic acid, and vitamins A, B, K, D, E, and the like. Minerals, protein ingredients, including protein obtained from meat meal or fish meal, liquid or powdered egg, fish solubles, whey protein concentrate, oils (e.g., soybean oil), cornstarch, calcium, inorganic phosphate, copper sulfate, salt, and limestone can also be added. Antioxidants can also be added.

[0059] In another disclosure, a kit comprising an expression vector comprising the isolated nucleic acid of SEQ ID NO: 1 to SEQ ID NO: 2, or a fragment thereof, is provided. In one illustrative disclosure, the isolated nucleic acid, or the fragment, can be 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98%, or 99% identical to the isolated nucleic acid of SEQ ID NO: 1 to SEQ ID NO: 2, or to the fragment thereof. In various illustrative embodiments, the fragment can be about 50 nucleotides in length, about 100 nucleotides in length, about 200 nucleotides in length, about 300 nucleotides in length, about 400 nucleotides in length, about 500 nucleotides in length, about 600 nucleotides in length, about 700 nucleotides in length, about 800 nucleotides in length, or about 900 nucleotides in length. In other embodiments, and disclosures the fragment can extend about 50 nucleotides, about 100 nucleotides, about 200 nucleotides, about 300 nucleotides, about 400 nucleotides, about 500 nucleotides, about 600 nucleotides, about 700 nucleotides, about 800 nucleotides, or about 900 nucleotides upstream from the 3' end of the isolated nucleic acid of any of SEQ ID NOS: 1 to 2. In yet other embodiments, and disclosures the fragment can include about 50 nucleotides, about 100 nucleotides, about 200 nucleotides, about 300 nucleotides, or about 350 nucleotides upstream and/or downstream from the TATA box sequence found in each of SEQ ID NOS: 1 to 2.

[0060] In one disclosure, the expression vector (i.e., with the heterologous promoter) included in the kit can have any of the other elements described herein, such as a selection marker, a cloning site, such as a multiple cloning site, an enhancer, a termination sequence, a signal peptide sequence, and the like. In another aspect, the expression vector can be a vector that replicates autonomously or integrates into the host cell genome. In another embodiment, the expression vector can be circularized or linearized (i.e., digested with a restriction enzyme so that a gene of interest can easily be cloned into the expression vector). In another disclosure,

the kit can include an expression vector and a control ORF encoding a marker or control gene for expression (e.g., an ORF encoding a LacZ- α fragment) for use as a control to show that the expression vector is competent to be ligated and to be used with a gene of interest.

[0061] In another illustrative disclosure, the kit can contain a tube containing a circular expression plasmid. In another disclosure, the kit can contain a tube with a linear expression vector that is digested with a restriction enzyme so it is ready to clone a gene of interest. In one disclosure, the tube can be sterilized. In either of these disclosures, the kit can also include a circular or linear expression vector and a control ORF encoding a marker or control gene for expression (e.g., an ORF encoding a LacZ- α fragment) for use as a control in showing that the expression vector can be ligated with a gene of interest, and/or to show that the host cell is competent for transformation.

[0062] In yet another disclosure, the kit can contain multiple different expression vectors. In another disclosure, the multiple different expression vectors can contain the same promoter, but different selectable markers, such as genes for resistance to the drugs G418, Nourseothricin (Nat), Zeocin, Blasticidin, or Hygromycin. In this disclosure, the kit may also contain aliquots of the drugs (e.g., G418, Nourseothricin (Nat), Zeocin, Blasticidin, or Hygromycin) in tubes, or other containers, separate from the corresponding vectors.

[0063] In any of the kit disclosures described above, the isolated nucleic acid can consist of any one of SEQ ID NOS. 1 to 2, or a fragment thereof. The phrase "consists of" means that the sequence specified by the SEQ ID NO. has no additional nucleotide sequences other than those corresponding to the SEQ ID NO.

[0064] In another illustrative disclosure, the kit can include other components for use with the expression vector, such as components for transformation of yeast cells, restriction enzymes for incorporating a protein coding sequence of interest into the expression vector, ligases, components for purification of expression vector constructs, buffers (e.g., a ligation buffer), instructions for use (e.g., to facilitate cloning), and any other components suitable for use in a kit for making and using the expression vectors described herein. In another disclosure, the expression vector or any other component of the kit can be included in the kit in a sealed tube (e.g., sterilized or not sterilized) or any other suitable container or package (e.g., sterilized or not sterilized). The kits described in the preceding paragraphs that include the expression vector comprise the expression vector comprising a promoter described herein operably linked to the vector which is heterologous to the promoter (i.e., the combination does not occur in nature).

[0065] The following examples provide illustrative methods for carrying out the practice of the present invention. As such, these examples are provided for illustrative purposes only and are not intended to be limiting.

EXAMPLES

Example 1: Preparation of Promoter Sequences and Promoter-Containing Plasmids

[0066] Microarray "chips" to measure *Pichia pastoris* mRNA levels were created by Affymetrix (Affymetrix, Inc.) using *Pichia* genomic DNA sequence provided by Dr. James Cregg. Microarray chips were probed with RNA samples from yeast cultures grown on different carbon sources, including methanol, glycerol and ethanol. Cultures were harvested at various time points in mid-log phase and frozen for the subsequent RNA isolation as recommended by Affymetrix. Standard Affymetrix protocols for probing Affymetrix microarrays were followed.

[0067] Total RNA from each cell sample was extracted by a glass bead breakage/phenol extraction/LiCl precipitation method. RNA was converted to cDNA and subsequently amplified and labeled to probe Affymetrix chips. Labeled cDNAs were hybridized to Affymetrix *Pichia pastoris* microarray chips and evaluated using Affymetrix software. Data analysis identified genes induced by methanol (relative to glycerol) or genes expressed under most conditions (i.e., constitutively expressed) as shown in Table 1.

[0068] Table 1 lists the relative mRNA levels from the indicated genes comparing promoter activity from *Pichia pastoris* cells grown on media with methanol, glycerol or ethanol as the carbon source.

Table 1. Relative Activity of Promoters

Protein	Gene	Relative Promoter Activity			
		Methanol	MeOH starvation	Glycerol	Ethanol
Hypothetical protein	UPP	9942	8297	9512	10673
Protein Translation Elongation Factor 1A	TEF2	8910	5626	9309	9471
Hypothetical protein	Chr1-0469	6904	192	7673	7618
Glyceraldehyde 3-phosphate dehydrogenase	GAP	5685	4688	6701	6069
AOX1 Average	AOX1	9127	4974	610	361
Dihydroxy-acetone synthase	DAS	8533	2591	1211	796

[0069] Figure 6 shows the relative promoter activity for pUPP, TEF2, Chr1-0469, GAP, AOX1, and DAS using the carbon sources methanol (leftmost bar in each set of bars), methanol starvation (bar second from the left in each set of bars), glycerol (bar third from the left in each set of bars), and ethanol (rightmost bar in each set of bars). The hypothetical proteins are the ORFs for the UPP and Chr1-0469 promoters.

[0070] Promoters from two constitutively expressed genes were cloned and tested with reporter vectors. All of the reporter plasmids were missing promoter elements to drive expression of the reporter ORF; alternatively, a cloning scheme utilizing Type II restriction enzymes (i.e. Bsal) was put in place to introduce DNA fragments (promoters) upstream of the ORF. All constructs contained the same 3'AOX1 transcriptional terminator sequence after the stop codon of the reporter ORF.

[0071] One reporter plasmid, pTZ-GFP, is a derivative of pJAZ (zeocin resistance as a selection marker) without the AOX1 promoter, but with a green fluorescent protein (GFP) ORF as a reporter gene for intracellular expression. A second group of reporter plasmids contains selectable marker genes for zeocin or G418 resistance and the *Aspergillus* alpha-galactosidase or beta-galactosidase ORFs as reporters for intracellular expression. The third group of reporter plasmids has the *S. cerevisiae* pre-pro-alpha Mating Factor (aMF) signal for secretion fused in frame with the *E. coli* appA2 coding sequence for phosphatase (phytase), as the reporter ORF. None of the reporter plasmid sequences are expressed in *Pichia pastoris* without the introduction of a sequence that has promoter activity upstream of the reporter ORF.

[0072] Two of the most active constitutive promoters, from the UPP and Chr1-0469 genes, consisted of the DNA ~1000 bp "upstream" of ORF's of unknown function. These promoter sequences were amplified using PCR primers and *Pichia pastoris* genomic DNA. The sequence of the PCR products was confirmed, prior to ligation into reporter plasmids. The UPP promoter was cloned using strategically placed Bsal sites in the primers flanking the promoter. The Chr1-0469 promoter contains recognition sites for Bsal and BsmBI; therefore, cloning ends were created by a PCR-denaturation-annealing procedure. Briefly, the promoter was amplified with two sets of primers. The PCR products represent two nearly identical double-stranded DNA molecules except that the ends are off-set by 4 bp (one is four nucleotides shorter from the 5'-end and four nucleotides longer from the 3'-end, and the other four nucleotides longer from the 5'-end and four nucleotides shorter from the 3'-end). The two different PCR products are mixed, denatured and annealed to create promoter molecules with the proper 4 base single strand extensions at each end of the molecule ("sticky" ends to facilitate cloning). Cloning ends, generated by Bsal digestion or by PCR-denaturation-annealing, were complementary to ends created in the reporter plasmids. Promoters regulating the *Pichia pastoris* AOX1 and GAP1 genes were introduced into the same reporter constructs as controls.

[0073] The promoter specific plasmid constructs were linearized by digestion with specific restriction enzymes and introduced into wild type *Pichia pastoris* cells by electroporation. Transformants containing the plasmid DNA integrated into the genome of the host strain were selected on YPD agar with either zeocin or G418. Single colonies from the transformation were purified (again on selective plates), then, multiple clones were collected on YPD master plates. These master plates were replica-plated to a series of diagnostic plates with agar media containing one of the following compounds as the only carbon source for growth of the *Pichia pastoris* cells: glucose (i.e., dextrose), glycerol, methanol, or ethanol (all at 1% w/v). Plates were incubated at 30° C for 20 to 44 hours before scoring.

Example 2: Determination of GFP Expression in Transformants

[0074] The relative level of GFP expression between the various transformants was compared. In this example, replica plates were exposed to UV light using a transilluminator and the fluorescence intensity of the different strains was compared by direct visualization. *Pichia* cells without GFP expression, negative colonies, did not demonstrate GFP fluorescence under UV illumination. The intensity of the observed fluorescence correlated with an increase in GFP expression.

[0075] Figure 5 is an example of GFP expression for various transformants on replica plates with different carbon sources. The location of the different *Pichia* promoter strains in each panel is the same, and the "Master YPD" panel identifies the location of the strains by their promoter designation: AOX1, pUPP, GAP1, 0001, 0002, 0003, 0004, 0005, 0006, 0007, and Vector (mock control). The lower part of Figure 5, labeled as "Expression of GFP induced by Dex, Gly, MeOH, or EtOH" shows that the carbon source used in each panel is dextrose (D), glycerol (G), methanol (M) or ethanol (E), respectively. The data show that the UPP promoter is as strong or stronger than the GAP promoter under the conditions tested, and is stronger than the AOX promoter under most conditions tested.

Example 3: Determination of Relative Activity of Alpha- and Beta-Galactosidases of Transformants

[0076] The UPP promoter and the Chr1-0469 promoter sequences were amplified using PCR primers and *Pichia pastoris* genomic DNA (the sequences of the forward primers used for amplification of the promoter fragments are shown on FIGURE 8). The sequences of the PCR products were confirmed, prior to ligation into the alpha-GAL reporter plasmid (FIGURE 7). Promoters regulating the *Pichia pastoris* GAP, TEF2, DAS and AOX1 ORFs were introduced into the same reporter plasmid as controls. The sequences of the inserted promoter fragments were confirmed by sequencing the entire insertions, in both directions, using plasmid specific primers located upstream of the inserted fragment (forward primers) and in the 5' region of the alpha-Galactosidase ORF (reverse primers). The promoter specific plasmid constructs were linearized by digestion with specific restriction enzymes and introduced into wild type *Pichia pastoris* cells by electroporation, and transformants containing the plasmid DNA integrated into the genome of the host strain were selected. Cells from well isolated, single-colony transformants were purified and collected on a YPD master plate and incubated at 30°C. The master plate was subsequently replica-plated to a series of diagnostic plates with agar media containing one of the following compounds as the only carbon source for growth of the *Pichia pastoris* cells: glucose (i.e., dextrose), glycerol, methanol, or ethanol (all at 1% w/v) and alpha-X-Gal chromogenic substrate. These plates were then incubated at 30° C for 44 hours (FIGURE 9).

[0077] The expression level from the different promoter-alpha-galactosidase reporter gene constructs, was determined by the observation of the alpha-galactosidase from YPD agar plates made with 100 mM phosphate buffer, pH 6.5, and alpha-X-GAL as a chromogenic substrate. Reporter activity was detected as a blue colored product in, or around the cells.. Reporter activity was demonstrated with the UPP and Chr1-0469 promoters.

[0078] The promoter designations in FIG. 9 correspond to the Chr1-0469 promoter (SEQ ID NO: 1) and the UPP promoter (SEQ ID NO: 2). The length of each promoter fragment is indicated with the +1 position equal to the A in the ATG start codon of the corresponding gene in the *Pichia pastoris* genome (therefore the promoter sequences are from -1 to -1500). Fragments of other *Pichia pastoris* promoters as well as a reporter vector without any promoter sequences are used as controls. The transformants were incubated on agar plates containing X-alpha-GAL, a chromogenic substrate for alpha-Galactosidase, and one of four carbon sources as indicated. Plates A and B in Fig. 9, panel A, were replica-plated on different carbon sources to plates A and B, respectively, shown in Fig. 9, panels B and C, respectively. Thus, the designations in Fig. 9, panel A of plates A and B refer to Fig. 9, panels B and C, respectively. The notations of NO:1 and NO:2 refer to SEQ ID NOS: 1 and 2 (i.e., to the promoter used to express α -galactosidase). The level of α -galactosidase expression with the promoters is shown in the tables labeled Plate A and Plate B below on the various carbon sources. The level of α -galactosidase expression is shown as 0, +1, +2, +3, +4, and +5, with the level of expression increasing from 0 to +5.

[0079] Similarly, the beta-Galactosidase ORF was used for analysis of expression of the reporter constructs with the Chr1-0469 promoter and the UPP promoter sequences in the *Pichia pastoris* transformants. In these experiments the expression level from the beta-galactosidase reporter gene was determined on plates with beta-X-GAL as a chromogenic substrate. Reporter activity with beta-Galactosidase ORF was also demonstrated with both the UPP and Chr1-0469 promoters.

Relative Alpha-Galactosidase Expression, Plate A, from Figure 9.

Promoter (seq ID/name)	Fig 9 (Panel/row/column)	Carbon Source	Relative A-gal expression (1 - 5)
Control	A/C/1	Glucose	0
Control	A/F/1	Glucose	0
Control	A/E/2	Glucose	0
UPP-513 (No:2)	A/F/5	Glucose	4
UPP-354 (No:2)	A/G/1	Glucose	4
DAS	A/G/2	Glucose	0
1-0469 (No:1)	A/G/3	Glucose	4
GAP	A/G/4	Glucose	3
Control	A/C/1	Glycerol	0
Control	A/E/1	Glycerol	0

Promoter (seq ID/name)	Fig 9 (Panel/row/column)	Carbon Source	Relative A-gal expression (1 - 5)
Control	A/E/2	Glycerol	0
UPP-513 (No:2)	A/F/5	Glycerol	3
UPP-354 (No:2)	A/G/1	Glycerol	4
DAS	A/G/2	Glycerol	0
1-0469 (No:1)	A/G/3	Glycerol	3
GAP	A/G/4	Glycerol	2
Control	A/C/1	Methanol	1
Control	A/E/1	Methanol	0
Control	A/E/2	Methanol	1
UPP-513 (No:2)	A/F/5	Methanol	3
UPP-354 (No:2)	A/G/1	Methanol	4
DAS	A/G/2	Methanol	2
1-0469 (No:1)	A/G/3	Methanol	3
GAP	A/G/4	Methanol	3
Control	A/C/1	Ethanol	0
Control	A/E/1	Ethanol	0
Control	A/E/2	Ethanol	0
UPP-513 (No:2)	A/F/5	Ethanol	3
UPP-354 (No:2)	A/G/1	Ethanol	4
DAS	A/G/2	Ethanol	0
1-0469 (No:1)	A/G/3	Ethanol	3
GAP	A/G/4	Ethanol	2

Relative Alpha-Galactosidase Expression, Plate B, from Figure 9

Promoter (seq ID/name)	Fig 9 (Plate B/row/column)	Carbon Source	Relative A-gal expression (1 - 5)
UPP-513 (No:2)	B/F/1	Glucose	4
UPP-345 (No:2)	B/F/2	Glucose	4
DAS	B/F/3	Glucose	0
1-0469 (No:1)	B/F/4	Glucose	3
GAP	B/F/5	Glucose	4
UPP 222 (No:2)	B/G/2	Glucose	3
UPP-513 (No:2)	B/F/1	Glycerol	4
UPP-345 (No:2)	B/F/2	Glycerol	4
DAS	B/F/3	Glycerol	0

Promoter (seq ID/name)	Fig 9 (Plate B/row/column)	Carbon Source	Relative A-gal expression (1 - 5)
1-0469 (No:1)	B/F/4	Glycerol	3
GAP	B/F/5	Glycerol	4
UPP 222 (No:2)	B/G/2	Glycerol	3
UPP-513 (No:2)	B/F/1	Methanol	5
UPP-345 (No:2)	B/F/2	Methanol	5
DAS	B/F/3	Methanol	3
1-0469 (No:1)	B/F/4	Methanol	3
GAP	B/F/5	Methanol	3
UPP 222 (No:2)	B/G/2	Methanol	4
UPP-513 (No:2)	B/F/1	Ethanol	4
UPP-345 (No:2)	B/F/2	Ethanol	4
DAS	B/F/3	Ethanol	0
1-0469 (No:1)	B/F/4	Ethanol	3
GAP	B/F/5	Ethanol	4
UPP 222 (No:2)	B/G/2	Ethanol	3

Example 4: Determination of Phytase Activity of Transformants

[0080] The level of phytase activity was determined in the various transformants. Agar medium on plates was supplemented with 100 mM phosphate buffer, pH 6.5, and calcium phytate. The activity of phytase acts to convert the non-transparent colloid calcium phytate substance into transparent material, forming a transparent halo around colonies expressing and secreting a phosphatase (phytase). An increased expression of the phytase by the cells leads to increased secretion of the enzyme, as demonstrated by a wider halo observed around the cell colony. Reporter activity was demonstrated with the UPP and Chr1-0469 promoters (FIGURE 10). In figure 10, the transformants harboring promoter sequences GAP, Chr1-0469 promoter (SEQ ID NO: 1) and the UPP promoter (SEQ ID NO: 2) are cloned upstream of the ORF consisting of the *S.cerevisiae* pre-pro alpha factor fused in frame with DNA encoding the *E.coli* appA2 phytase. Cells were spotted on diagnostic plates with calcium phytate and either glucose or methanol and incubated for 27 hours at 30° C.. The level of phytase expression with the promoters is noted by 0, +1, +2, +3, +4, and +5, with the level of expression increasing from 0 to +5. The relative levels of phytase expression are shown in the table below with No:1 and No:2 representing SEQ ID NO: 1 and SEQ ID NO: 2, respectively.

Table X.

Relative Phytase Activity from Figure 10		
Promoter (seq:ID)	Carbon Source	Relative activity 1 - 5 (5 is highest activity)
GAP	glucose	2
No:1	glucose	2
No:2, c	glucose	5
No:2, d	glucose	5
No:2, e	glucose	5
GAP	methanol	1
No:1	methanol	2
No:2, c	methanol	2
No:2, d	methanol	2
No:2, e	methanol	2

SEQUENCE LISTING**[0081]**

<110> KECK GRADUATE INSTITUTE OF APPLIED LIFE SCIENCES

<120> YEAST PROMOTERS FROM PICHIA PASTORIS

<130> 62472-229032

<140> PCT/US2014/022135

<141> 2014-03-07

<150> 61/774,982

<151> 2013-03-08

<160> 13

<170> PatentIn version 3.5

<210> 1

<211> 1003

<212> DNA

<213> Pichia pastoris

<400> 1

tgccaggatc tgaccactac aagaagcgta attaatgtatt tattatgtag cttgaggaat	60
tgattttctaa aggccaaaga tgatgtagag caatagtaga gtgattagca agtttctgaa	120
catgttttgt ggtcgggtat taggcgccgt acctctgtat tttttttttt tggcagatgg	180

gatgaccact	gtgcttacat	aatcttcgat	gggccctttt	ggaaacgacc	actccttcg	240
atcacgtaat	actagggcag	tatcactacg	cgctgggtatc	cgcatcttag	ccgctttaac	300
gacaaggcca	gggtaaatac	tgtgttgcat	gggcgatgcc	gtaggatctg	gccaacttgg	360
aacagagcag	cccaccgcgt	aggcagcgct	tgggaccagt	ttgttagcca	atcagttccc	420
ccgtctgagt	ccctacacgc	tcaggcttgc	atggatatgc	atgaaaaatg	tcaactcagt	480
cgcacacagg	gatgctgttg	tcgtggtgag	aaaaagctaa	aatagcaaac	ccatagcaat	540
cgagagactg	tgattggaag	caaaagaacc	aggcattgac	taggaaagac	tgacaagaaa	600
attagacggg	gccagtggta	tttcattgaa	ttgcactgaa	cacgcctccc	atttccttcc	660
ctcaatccat	tcaaaagggt	aattaaacct	tatctcggcc	tacattagtt	gtcatggaaa	720
cgtctcactt	tctctctgtg	tcccacagta	gcgagtaggc	ggagagacca	aaaatggctt	780
ctcggcccta	atcaaaacat	cctttgtcgc	gttggtttca	gatcgattcg	caatctgaga	840
gaaaaacaat	aacaccaaaa	taaaaagctt	ggaaaaagaa	gacatgattc	acatccacta	900
gtaaaatata	aatactcctg	ccgcctcccc	ctttcctttc	ttcttattcc	tcaactcact	960
gtttcagttt	attccaacta	ctttcactca	cttatcaaaa	atg		1003

<210> 2

<211> 773

<212> DNA

<213> Pichia pastoris

<400> 2

gggtgaaagc	caaccatctt	tgtttcgggg	aaccgtgctc	gccccgtaaa	gttaattttt	60
ttttcccgcg	cagctttaat	ctttcggcag	agaaggcggt	ttcatcgtag	cgtgggaaca	120
gaataatcag	ttcatgtgct	atacaggcac	atggcagcag	tcactatctt	gctttttaac	180
cttaaagtcg	ttcatcaatc	attaactgac	caatcagatt	ttttgcattt	gccacttatt	240
taaaaatact	tttgatatct	gcagatacgt	tcagtgggtt	ccaggacaac	acccaaaaaa	300
aggatatcaat	gccactaggc	agtcgggttt	atcttttggtc	accacgcaa	agaagcacc	360
acctctttta	ggttttaagt	tgtgggaaca	gtaacaccgc	ctagagcttc	aggaaaaacc	420
agtacctgtg	accgcaattc	accatgatgc	agaatgttaa	tttaaacgag	tgccaaatca	480
agatttcaac	agacaaatca	atcgatccat	agttacccat	tccagccttt	tcgtcgtcga	540
gcctgcttca	ttctgcctc	aggcgcataa	ctttgcatga	aaagtccaga	ttagggcaga	600
ttttgagttt	aaaataggaa	atataaacaa	atataccgcg	aaaaagggtt	gtttatagct	660
tttcgcctgg	tgccgtacgg	tataaataca	tactctcttc	ccccccctgg	ttctcttttt	720
cttttggtac	ttacatttta	ccgttcgctc	actcgcttca	ctcaacaaca	aaa	773

<210> 3

<211> 939

<212> DNA

<213> Pichia pastoris

<400> 3

gatctaacat	ccaaagacga	aagggtgaat	gaaacctttt	tgccatccga	catccacagg	60
tcattctca	cacataagta	ccaaagcga	caggagggga	tacactagca	gcagaccatt	120

```

ccattctca cacatagtg ccaaacgca caggagggga cacactaga gcagaccgt 120
gcaaacgcag gacctccact cctcttctcc tcaacaccca cttttgccat cgaaaaacca 180
gccagttat tgggcttgat tggagctcgc tcattccaat tccttctatt aggctactaa 240
caccatgact ttattagcct gtctatcctg gccccctgg cgagggtcat gtttgtttat 300
ttccgaatgc aacaagctcc gcattacacc cgaacatcac tccagatgag ggctttctga 360
gtgtggggtc aaatagtttc atgttcccca aatggcccaa aactgacagt ttaaacgctg 420
tcttggaacc taatatgaca aaagcgtgat ctcatccaag atgaactaag tttggttcgt 480
tgaaatgcta acggccagtt ggtcaaaaag aaacttccaa aagtcggcat accgtttgtc 540
ttgtttggta ttgattgacg aatgctcaaa aataatctca ttaatgctta gcgcagtctc 600
tctatcgctt ctgaaccccg gtgcacctgt gccgaaacgc aaatggggaa acaccgctt 660
tttgatgat tatgcattgt ctccacattg tatgcttcca agattctggg ggggaatactg 720
ctgatagcct aacgttcatt atcaaaattt aactgttcta acccctactt gacagcaata 780
tataaacaga aggaagctgc cctgtcttaa accttttttt ttatcatcat tattagctta 840
ctttcataat tgcgactggg tccaattgac aagcttttga ttttaacgac ttttaacgac 900
aacttgagaa gatcaaaaaa caactaatta ttcgaaacg 939

```

<210> 4

<211> 1001

<212> DNA

<213> *Pichia pastoris*

<400> 4

```

catgttggtg ttgtgaaata gacgcagatc gggaaactg aaaaataaca gttattattc 60
gagatctaac atccaagac gaaaggttga atgaaacctt tttgccatcc gacatccaca 120
ggtccattct cacacataag tgccaaacgc aacaggaggg gatacactag cagcagaccg 180
ttgcaaacgc aggacctcca ctctcttctt cctcaacacc cacttttgcc atcgaaaaac 240
cagcccagtt attgggcttg attggagctc gctcattcca attccttcta ttaggctact 300
aacacatga ctttattagc ctgtctatcc tggccccctt ggcgagggtt atgtttgttt 360
atttccgaat gcaacaagct ccgcattaca cccgaacatc actccagatg agggctttct 420
gagtgtgggg tcaaatagtt tcatgttccc caaatggccc aaaactgaca gtttaaacgc 480
tgtcttgga cctaatatga caaaagcgtg atctcatcca agatgaacta agtttggttc 540
gttgaaatgc taacggccag ttggtcaaaa agaaacttcc aaaagtcggc ataccgtttg 600
tcttgtttgg tattgattga cgaatgetca aaaataatct cattaatgct tagcgcagtc 660
tctctatcgc ttctgaaccc cgggtgcacct gtgccgaaac gcaaatgggg aaacaccgc 720
tttttgatg attatgcatt gtctccacat tgtatgcttc caagattctg gtgggaatac 780
tgctgatagc ctaacgttca tgatcaaaat ttaactgttc taaccctac ttgacagcaa 840
tatataaaca gaaggaagct gccctgtctt aaaccttttt ttttatcatc attattagct 900
tactttcata attgcgactg gttccaattg acaagctttt gattttaacg acttttaacg 960
acaacttgag aagatcaaaa aacaactaat tattcgaaac g 1001

```

<210> 5

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<400> 5

atccctacac gctcaggctt 20

<210> 6

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<220>

<221> modified_base

<222> (10)..(10)

<223> a, c, t, g, unknown or other

<400> 6

nnnggtctcn atcctcgctg ggtgaaagcc aac 33

<210> 7

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<220>

<221> modified_base

<222> (10)..(10)

<223> a, c, t, g, unknown or other

<400> 7

nnnggtctcn atccgcagat acgttcagtg gtttcc 36

<210> 8

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<220>

<221> modified_base

<222> (10)..(10)

<223> a, c, t, g, unknown or other

<400> 8

nnnggtctcn atccagtacc tgtgaccgca attc 34

<210> 9

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<220>

<221> modified_base

<222> (10)..(10)

<223> a, c, t, g, unknown or other

<400> 9

nnnggtctcn atcctgcctc aggtgcataa ctt 33

<210> 10

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<220>

<221> modified_base

<222> (10)..(10)

<223> a, c, t, g, unknown or other

<400> 10

nnnggtctcn atccaatgga ccaaattgtt gcaaggt 37

<210> 11

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<400> 11

nnnggtctcc atcccttgt tgagcaaca 29

<210> 12

<211> 28

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic primer

<220>

<221> modified_base

<222> (1)..(3)

<223> a, c, t, g, unknown or other

<400> 12

nnnggtctcg atccgcccaa acgaacag 28

<210> 13
 <211> 34
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Description of Artificial Sequence: Synthetic primer

 <220>
 <221> modified_base
 <222> (1)..(3)
 <223> a, c, t, g, unknown or other

 <400> 13
 nnnggtctcc atccaaagac gaaaggttga atga 34

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2012129036A2 \[0005\]](#)
- [US2008108108A \[0006\]](#)
- [WO20081287001A2 \[0009\]](#)
- [US6451572B \[0024\] \[0027\]](#)
- [US6841370B \[0024\] \[0027\]](#)
- [US6974690B \[0024\] \[0027\]](#)
- [US7320876B \[0024\] \[0027\]](#)
- [US7078035B \[0024\] \[0027\]](#)
- [US7138260B \[0024\] \[0027\]](#)
- [WO2007112739A \[0024\]](#)
- [WO2007112739PCT \[0027\]](#)
- [WO62472229032A \[0081\]](#)
- [US2014022135W \[0081\]](#)
- [WO61774982A \[0081\]](#)

Non-patent literature cited in the description

- **STADLMAYER et al.**Journal of Biotechnology, 2010, vol. 150, 4519-529 [0007]
- **GABRIEL POTVIN et al.**Biochemical Engineering Journal, 2012, vol. 64, 91-105 [0008]
- **ANDREAS KÜBERL et al.**Journal of Biotechnology, 2011, vol. 154, 312-320 [0010]
- **SAMBROOK et al.**Molecular Cloning: A Laboratory ManualCold Spring Harbor Laboratory Press20010000 [0027] [0034] [0037] [0039]
- **RODRIGUEZ et al.**Arch. of Biochem. and Biophys., 2000, vol. 382, 105-112 [0057]

P A T E N T K R A V

1. Isoleret nukleinsyre, hvor den isolerede nukleinsyre omfatter en sekvens, som er mindst 90% identisk med SEQ ID NO: 2 eller mindst 90% identisk med et fragment deraf, hvor sekvensen, som er mindst 90% identisk med SEQ ID NO: 2 eller mindst 90% identisk med et fragment deraf, har promotoraktivitet, hvor nukleinsyren omfatter sekvensen af en konstitutiv *Pichia pastoris*-promotor, og hvor den isolerede nukleinsyre er funktionsmæssigt forbundet med en heterolog kodningssekvens.

2. Isoleret nukleinsyre ifølge krav 1, hvor sekvensen af den isolerede nukleinsyre er SEQ ID NO: 2 eller et fragment deraf.

3. Isoleret nukleinsyre ifølge et hvilket som helst af kravene 1 til 2, hvor sekvensen er SEQ ID NO: 2.

4. Isoleret nukleinsyre ifølge et hvilket som helst af kravene 1 til 3, hvor den heterologe kodningssekvens koder et protein valgt blandt gruppen bestående af et toksin, et antistof, et hormon, et enzym, en vækstfaktor, en cytokin, et strukturelt protein, et immunogent protein og et cellesignalerende protein.

5. Isoleret nukleinsyre ifølge krav 4, hvor proteinet er et enzym til anvendelse i dyrefoder.

6. Isoleret nukleinsyre ifølge krav 5, hvor proteinet er valgt blandt gruppen bestående af en mannanase, en amylase, en glucanase, en protease, en cellulase og en xylanase.

7. Isoleret nukleinsyre ifølge krav 5, hvor proteinet er en phytase.

8. Isoleret nukleinsyre ifølge krav 5, hvor proteinet er en galactosidase.

9. Expressionsvektor, som omfatter den isolerede nukleinsyre ifølge et hvilket som helst af kravene 1 til 8.

10. Værtscelle, som omfatter den isolerede nukleinsyre ifølge et hvilket som helst af kravene 1 til 8 eller ekspressionsvektoren ifølge krav 9.

11. Værtscelle ifølge krav 10, hvor værtscellen er en *Pichia* species.

12. Værtscelle ifølge krav 11, hvor *Pichia* species er *Pichia pastoris*.

13. Værtscelle ifølge krav 10, hvor værtscellen er valgt blandt gruppen bestående af *Hansenula* species, *Pichia* species, *Saccharomyces* species, *Yarrowia* species, *Candida* species, *Schizosaccharomyces* species, *Torulaspora* species og *Kluveromyces* species.

14. Fremgangsmåde til fremstilling af et protein, hvor fremgangsmåden omfatter trinnet i et dyrkningsmedium at dyrke en værtscelle, som omfatter en første ekspressionskassette, som omfatter en isoleret nukleinsyre af SEQ ID NO: 2 eller et fragment deraf, som er funktionsmæssigt forbundet med en heterolog kodningssekvens, som koder et protein, hvor dyrkningen foregår under betingelser, som tillader ekspression af proteinet; hvor proteinet eksprimeres ved anvendelse af den første ekspressionskassette i kombination med en anden ekspressionskassette, hvor den anden ekspressionskassette omfatter:

- (a) den heterologe kodningssekvens, som koder proteinet, som funktionsmæssigt er forbundet med en isoleret nukleinsyre med en sekvens, som omfatter SEQ ID NO: 3 eller SEQ ID NO: 4, hvor SEQ ID NO: 3 eller SEQ ID NO: 4 har promotoraktivitet; eller

- (b) den isolerede nukleinsyre af SEQ ID NO: 2 eller et fragment deraf, som funktionsmæssigt er forbundet med den heterologe kodningssekvens, som koder proteinet.

DRAWINGS

Chr1-0469 PROMOTER SEQUENCE (SEQ ID NO: 1)

TGCCAGGATCTGACCACTACAAGAAGCGTAATTAAGTATTTATTATGTAGCTT
GAGGAATTGATTTCTAAAGGCCAAAGATGATGTAGAGCAATAGTAGAGTGAT
TAGCAAGTTTCTGAACATGTTTTGTGGTCGGGTATTAGGCGCCGTACCTCTGT
ATTTTTTTTTTTTGGCAGATGGGATGACCACTGTGCTTACATAATTTTCGATGG
GCCCTTTTGGAAACGACCACTCCTTCCGATCACGTAATACTAGGCGAGTATCA
CTACGCGTCGGTATCCGCATTTTAGCCGCTTTAACGACAAGGTCAGGGTAAAT
ACTGTGTTGCATGGGCGATGCCGTAGGATCTGGCCAACTTGGAACAGAGCAG
CCCACCGCGTAGGCGAGCGCTTGGGACCAGTTTGTTAGCCAATCAGTTCCCCC
GTCTGAGTCCCTACACGCTCAGGCTTGCATGGATATGCATGAAAAATGTCAA
CTCAGTCGCACACAGGGATGCTGTTGTCGTGGTGAGAAAAAGCTAAAATAGC
AAACCCATAGCAATCGAGAGACTGTGATTGGAAGCAAAAGAACCAGGCATT
GACTAGGAAAGACTGACAAGAAAATTAGACGGTGCCAGTGGTATTTTCATTGA
ATTGCACTGAACACGCCTCCCATTTCCCTTCCCTCAATCCATTCAAAGGGCTAA
TAAACCTTATCTCGGCCTACATTAGTTGTCATGGAAACGTCTCACTTTCTCTC
TGTGTCCACAGTAGCGAGTAGGCGGAGAGACCAAAAATGGTCTCTCGGCCC
TAATCAAAACATCCTTTGTCGCGTTGTTTTAGATCGATTCCGAATCTGAGAG
AAAAACAATAACACCAAAATAAAAAGCTTGGAAGAAAGACATGATTCAC
ATCCACTAGTAAAATATAAATACTCCTGCCGCCTCCCCCTTTCCTTTCTTCTTA
TTCCTCAACTCACTGTTTCAGTTTATTCCAACACTTTCACTCACTTATCAAAA
ATG

FIG. 1

UPP PROMOTER SEQUENCE (SEQ ID NO: 2)

GGGTGAAAGCCAACCATCTTTGTTTTCGGGGAACCGTGCTCGCCCCGTAAAGTTAA
TTTTTTTTTCCCGCGCAGCTTTAATCTTTCGGCAGAGAAGGCGTTTTTCATCGTAGC
GTGGGAACAGAATAATCAGTTCATGTGCTATACAGGCACATGGCAGCAGTCACT
ATTTTGCTTTTTAACCTTAAAGTCGTTTCATCAATCATTAAGTACCAATCAGATTT
TTTGCATTTGCCACTTATCTAAAAATACTTTTGTATCTCGCAGATACGTTTCAGTGG
TTTCCAGGACAACACCCAAAAAAGGTATCAATGCCACTAGGCAGTCGGTTTTAT
TTTTGGTCACCCACGCAAAGAAGCACCCACCTCTTTTAGGTTTTAAGTTGTGGGA
ACAGTAACACCGCCTAGAGCTTCAGGAAAAACCAGTACCTGTGACCGCAATTCA
CCATGATGCAGAATGTTAATTTAAACGAGTGCCAAATCAAGATTTCAACAGACA
AATCAATCGATCCATAGTTACCCATTCCAGCCTTTTCGTCGTCGAGCCTGCTTCAT
TCCTGCCTCAGGTGCATAACTTTGCATGAAAAGTCCAGATTAGGGCAGATTTTGA
GTTTAAAATAGGAAATATAAACAAATATACCGCGAAAAAGGTTTGTATAGCTT
TTCGCCTGGTGCCGTACGGTATAAATACATACTCTCTCCCCCCCCCTGGTTCTCTT
TTTCTTTTGTTACTTACATTTTACCGTTCGGTCACTCGCTTCACTCAACAACAAAA

FIG. 2

AOX1 PROMOTER SEQUENCE (SEQ ID NO: 3)

GATCTAACATCCAAAGACGAAAGGTTGAATGAAACCTTTTTGCCATCCGACATCC
ACAGGTCCATTCTCACACATAAGTGCCAAACGCAACAGGAGGGGATACACTAGC
AGCAGACCGTTGCAAACGCAGGACCTCCACTCCTCTTCTCCTCAACACCCACTTT
TGCCATCGAAAAACCAGCCCAGTTATTGGGCTTGATTGGAGCTCGCTCATTCCAA
TTCCTTCTATTAGGCTACTAACACCATGACTTTATTAGCCTGTCTATCCTGGCCCC
CCTGGCGAGGTTTCATGTTTGTATTATTTCCGAATGCAACAAGCTCCGCATTACACC
CGAACATCACTCCAGATGAGGGCTTTCTGAGTGTGGGGTCAAATAGTTTCATGTT
CCCCAAATGGCCCCAAAACCTGACAGTTTAAACGCTGTCTTGGAACCTAATATGACA
AAAGCGTGATCTCATCCAAGATGAACTAAGTTTGGTTCGTTGAAATGCTAACGGC
CAGTTGGTCAAAAAGAACTTCCAAAAGTCGGCATAACCGTTTGTCTTGTGGTA
TTGATTGACGAATGCTCAAAAATAATCTCATTAAATGCTTAGCGCAGTCTCTCTAT
CGCTTCTGAACCCCGGTGCACCTGTGCCGAAACGCAAATGGGGAAACACCCGCT
TTTTGGATGATTATGCATTGTCTCCACATTGTATGCTTCCAAGATTCTGGTGGGAA
TACTGCTGATAGCCTAACGTTTCATGATCAAAATTAACTGTTCTAACCCCTACTTG
ACAGCAATATATAAACAGAAGGAAGCTGCCCTGTCTTAAACCTTTTTTTTATCA
TCATTATTAGCTTACTTTTCATAATTGCGACTGGTTCCAATTGACAAGCTTTTGATT
TTAACGACTTTTAACGACAACCTTGAGAAGATCAAAAAACAATAATTATTCGAA
ACG

FIG. 3

GENOMIC SEQUENCE, 1000BP UPSTREAM OF AOX1 PROMOTER ATG
(SEQ ID NO: 4)

CATGTTGGTATTGTGAAATAGACGCAGATCGGGAACACTGAAAAATAACAGTTA
TTATTCGAGATCTAACATCCAAAGACGAAAGGTTGAATGAAACCTTTTTGCCATC
CGACATCCACAGGTCCATTCTCACACATAAGTGCCAAACGCAACAGGAGGGGAT
ACACTAGCAGCAGACCGTTGCAAACGCAGGACCTCCACTCCTCTTCTCCTCAACA
CCCACTTTTGCCATCGAAAAACCAGCCCAGTTATTGGGCTTGATTGGAGCTCGCT
CATTCCAATTCTTCTATTAGGCTACTAACACCATGACTTTATTAGCCTGTCTATC
CTGGCCCCCCTGGCGAGGTTTCATGTTTGTGTTATTTCCGAATGCAACAAGCTCCGC
ATTACACCCGAACATCACTCCAGATGAGGGCTTTCTGAGTGTGGGGTCAAATAGT
TTCATGTTCCCCAAATGGCCCCAAAACCTGACAGTTTAAACGCTGTCTTGGAACCTA
ATATGACAAAAGCGTGATCTCATCCAAGATGAACTAAGTTTGGTTCGTTGAAATG
CTAACGGCCAGTTGGTCAAAAAGAACTTCCAAAAGTCGGCATAACGTTTGTCTT
GTTTGGTATTGATTGACGAATGCTCAAAAATAATCTCATTAAATGCTTAGCGCAGT
CTCTCTATCGCTTCTGAACCCCGGTGCACCTGTGCCGAAACGCAAAATGGGGAAAC
ACCCGCTTTTTGGATGATTATGCATTGTCTCCACATTGTATGCTTCCAAGATTCTG
GTGGGAATACTGCTGATAGCCTAACGTTTCATGATCAAAAATTTAACTGTTCTAACC
CCTACTTGACAGCAATATATAAACAGAAGGAAGCTGCCCTGTCTTAAACCTTTTT
TTTTATCATCATTATTAGCTTACTTTCATAATTGCGACTGGTTCCAATTGACAAGC
TTTTGATTTTAACGACTTTTAACGACAACCTTGAGAAGATCAAAAAACAATAATT
ATTCGAAACG

FIG. 4



Expression of GFP induced by Dex, Gly,
MeOH or EtOH




FIG. 5




FIG. 6




FIG. 7

Promoter	SEQ ID NO:	Forward primer	Primer sequence	Primer position on the Promoter sequence
Chr1-0469	1	095F-long	ATCCCTACACGCTCAGGCTT	-570
Upp	2	014-F778	NNNGGTCTCNATCCTCGCTGGGTGAAAGCCAAC	-778
Upp	2	014-F513	NNNGGTCTCNATCCGAGATACGTTAGTGGTTTCC	-513
Upp	2	014-F354	NNNGGTCTCNATCCAGTACCTGTGACCGCAATTC	-354
Upp	2	014-F4	NNNGGTCTCNATCCTGCCTCAGGTGCATAACTT	-221
GAP		GAP-F1	NNNGGTCTCNATCCAATGGACCAAATTGTTGCAAGGT	-619
DAS		226F620	NNNGGTCTCCATCCCTTTGTTGAGCAACA	-382
DAS		226F518	NNNGGTCTCGATCCGCCCAAACGAACAG	-483
AOX		AOXF1	NNNGGTCTCCATCCAAAGACGAAAGGTTGAATGA	-931

FIG. 8

Alpha-X-Gal plate assay for alpha-Galactosidase activity of *Pichia pastoris* transformants harboring a construct of a synthetic reporter alpha-gal ORF fused with an isolated promoter sequences (SEQ ID NO:). No promoter has been inserted into a control plasmid. All constructs are integrated into genome of the transformants.

Position	1	2	3	4	5
Row	Plate A				
A		xxx	xxx	xxx	
B	xxx	xxx	xxx	xxx	xxx
C	Control	xxx	xxx	xxx	xxx
D	xxx	xxx	xxx	xxx	xxx
E	Control	Control	xxx	xxx	xxx
F	xxx	xxx	xxx	xxx	NO:2(-513)
G	NO:2 (-354)	DAS	NO:1 (-570)	GAP	
H	xxx	xxx	xxx	xxx	xxx
I	xxx	xxx	xxx	xxx	
J		xxx	xxx		

Position	1	2	3	4	5
Row	Plate B				
A	xxx	xxx	xxx	xxx	
B	xxx	xxx	xxx	xxx	xxx
C	xxx	xxx	xxx	xxx	xxx
D	xxx	xxx	xxx	xxx	xxx
E	xxx	xxx	xxx	xxx	xxx
F	NO:2 (-513)	NO:2 (-354)	DAS	NO:1 (-570)	GAP
G	xxx	NO:2 (-221)	xxx	xxx	
H	xxx	xxx	xxx	xxx	
I		xxx	xxx	xxx	
J			xxx		

FIG. 9, Panel A




FIG. 9, Panel B




FIG. 9, Panel C

Phytase activity (plate assay). *Pichia pastoris* transformants harboring isolated promoter constructs regulating the yeast are pro alpha mazing factor fused, in frame with the Eco 1 appAZ phytase, integrated into genome of the host *Pichia* cells. The cultures were spotted on the surface of diagnostic plates containing calcium phytate (glucose, upper panel, and methanol, lower panel) and incubated for 27 hours.




FIG. 10