

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978
PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) – Regulation 22(1)(g) and 31)

OFFICIAL APPLICATION NO.

LODGING DATE

ACCEPTANCE DATE

21 2002/7148

22 5 SEP 2002

43 23.4.2003

INTERNATIONAL CLASSIFICATION

NOT FOR PUBLICATION

51 C07D

CLASSIFIED BY: WIPO

FULL NAMES OF APPLICANT

71 ~~H. LUNDBECK A/S~~
CAMBRIDGE ENTERPRISE LIMITED
11/03/08

FULL NAMES OF INVENTORS

72 1. PETERSEN, HANS
2. BØGESØ, KLAUS PETER
3. HOLM, PER

EARLIEST PRIORITY CLAIMED

COUNTRY

NUMBER

DATE

33 DK

31 PA200000402

32 13 MAR 2000

TITLE OF INVENTION

54 CRYSTALLINE BASE OF CITALOPRAM

57 ABSTRACT (NOT MORE THAN 150 WORDS)

NUMBER OF SHEETS

25

If no classification is finished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

Abstract


The present invention relates to the crystalline base of the well known antidepressant drug citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile, formulations of said base, a process for the preparation of purified salts of citalopram, such as the hydrobromide, using the base, the salts obtained by said process and formulations containing such salts.

Crystalline Base of Citalopram

The present invention relates to the crystalline base of the well known antidepressant drug citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuran-carbonitrile, formulations of said base, a process for the preparation of purified salts of citalopram, such as the hydrobromide, using the base, the salts obtained by said process and formulations containing such salts.

Background of the Invention

Citalopram is a well-known antidepressant drug that has now been on the market for some years and has the following structure:



It is a selective, centrally acting serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities. The antidepressant activity of the compound has been reported in several publications, eg. J. Hyttel, *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.*, **1982**, 6, 277-295 and A. Graven, *Acta Psychiatr. Scand.*, **1987**, 75, 478-486. The compound has further been disclosed to show effects in the treatment of dementia and cerebrovascular disorders. EP-A-474580.

Citalopram was first disclosed in DE 2,657,013, corresponding to US 4,136,193. This patent publication describes the preparation of citalopram by one method and outlines a further method, which may be used for preparing citalopram. The citalopram prepared was isolated as the oxalate, the hydrobromide and the hydrochloride salt, respectively. Furthermore, the citalopram base was obtained as an oil (B.P. 175 C/0.03 mmHg). Citalopram is marketed as the hydrobromide and the hydrochloride, respectively.

A number of processes for the preparation of citalopram have been disclosed. In many of these, the last step of the process is a conversion of a group different from cyano in the 5 position of the direct analogue of citalopram to a 5-cyano group. So citalopram has been prepared by:

Exchange of 5-halogen, or 5-CF₃-(CF₂)_n-SO₂-O- with cyano (DE 2,657,013 and co-pending WO 0011926 and WO 0013648)

Conversion of a 5-amido or 5-ester group to a 5-cyano group (WO 9819513)

5 Conversion of a 5-amino group to a 5-cyano group (WO 9819512)

Conversion of a 5-formyl group to a 5-cyano group (WO 9900548)

Conversion of a 5-oxazoliny or 5-thiazoliny group to a 5-cyano group (WO 0023431)

10 Other processes for the preparation of citalopram comprise exchange of the 5-bromo group of 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide with 5-cyano followed by alkylation with a 3-(N,N-dimethylamino)propyl-halogenide (DE 2,657,013 and WO 9819511).


Many of the processes mentioned above have the disadvantage that it is difficult to separate the intermediates formed during the process (the intermediates mentioned above or earlier
15 intermediates) from the end product and, accordingly, extensive purification procedures involving loss of citalopram are required in order to obtain the necessary quality of the end product.

It has now been found that the base of citalopram may be obtained as a very nice and pure crystalline product, which may easily be handled and conveniently be formulated into tablets and other
20 pharmaceutical forms. Furthermore, it has surprisingly been found that a very good and efficient purification of citalopram may be obtained during manufacture of citalopram (e.g. of the hydrobromide or the hydrochloride salt) by crystallising the base, and thereafter optionally forming a salt from the base.

25 This purification process is particularly useful for removing intermediates which are structurally closely related to citalopram, in particular compounds which only differ from citalopram by the substituent situated in position 5 on the isobenzofurane ring, and intermediates which have physical/chemical properties which are close to those of citalopram, e.g. the 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-isobenzofuranes having halogen (in
30 particular bromide and chloride), an amide or an ester in position 5 of the isobenzofurane ring, or 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, or -chloride.

Summary of the invention

35 The present invention provides the crystalline base of the compound




(I)

In a second aspect, the invention provides a process for the manufacture of a salt of citalopram, preferably the hydrobromide or hydrochloride in which the free base of citalopram is precipitated in crystalline form, optionally re-crystallised one or more times and then transferred to a pharmaceutically acceptable salt of citalopram.

In particular, the invention relates to a process for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free and precipitated in crystalline form from a solvent, and thereafter separated from the solvent and optionally re-crystallised one or more times, and then transferred into a salt thereof.

In particular, the invention relates to a process for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free from a crude salt or crude mixture of citalopram.

More particularly, the present invention relates to a process for the manufacture of citalopram base or a salt of citalopram characterised in that one or more impurities of the formula



(II)

wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^1$ wherein R^2 and R^1 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of

citalopram, by precipitating citalopram base in crystalline form, optionally re-crystallising said base one or more times and/or transferring said base into a salt thereof.

- 5 The crude mixture of citalopram containing the compound of formula II as an impurity may be prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source, or by subjecting 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranhalogenide, in particular the bromide, to a cyanide exchange reaction followed by alkylation with a 3-(N,N-dimethylamino)propyl-halogenide.

- 10 In a particular embodiment of the invention, Z is halogen, in particular bromide or chloride.

In a particularly preferred embodiment of the invention, the salt prepared is the hydrobromide or hydrochloride salt of citalopram.

- 15 The crude salt may be any convenient salt, such as the hydrobromide, hydrochloride, sulphate, oxalate, phosphate, nitrate or any other convenient salt. Other salts are salts of organic acids.

In a preferred embodiment of the invention, the crude salt is the sulphate, the hydrobromide or the hydrochloride salt.

20

In yet another aspect, a pharmaceutical formulation of the free base of citalopram is provided. Preferably the formulation is for oral administration.

The formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

In particular, the pharmaceutical composition of the invention contains the racemic mixture of citalopram base.

The crystalline base of citalopram is preferably more than 99.8% w/w pure, most preferably more than 99.9% w/w pure (peak area). The melting point is preferably a range within 90 – 93 °C, most

preferably 91 - 92 °C (DSC; onset, open capsule) or it is between 92 and 94°C, preferably 92.5 and 93.5 °C (DSC; onset, closed capsule). The crystalline base of citalopram is preferably in racemic form.

- 5 The terms "crude salt" and "crude mixture" refer to the fact that the salt and the mixture, respectively, comprise impurities, in particular impurities of formula II, which must be removed or which it is desired to remove.

10 The crude salt may be a salt separated directly from the reaction mixture, or the crude reaction mixture may have been subjected to some initial purification, e.g. one re-crystallisation, and /or treatment with activated carbon or silica gel, and the salt formed subsequently by treatment with an acid using methods known in the art. The salt may be isolated by precipitation or it may exist in a solvent, e.g. in the mixture resulting directly from the synthesis of the salt.

- 15 Similarly, the crude mixture comprising citalopram may be obtained directly from the synthesis of the compound according to any of the above mentioned processes or it may have been subjected to some initial or simultaneous purification, e.g. one re-crystallisation, treatment with activated carbon or silica gel.

- 20 The base of citalopram may be set free from the crude salt by dissolving the crude salt in a mixture of water and an organic solvent and then adding a base. The organic solvent may be toluene, ethyl acetate or any other suitable solvent and the base may be any convenient base, preferably NaOH or NH₃. Likewise, the base of citalopram may, if necessary, be set free from a crude mixture containing citalopram by treatment with a base.

25

Crude mixtures containing citalopram base may be subjected to further purification and extraction, before the base is precipitated in crystalline form. The base of citalopram may be isolated by separation of the organic phase, evaporation of the solvent in order to obtain the base most probably as an oil and then crystallisation of the base from an aprotic solvent, such as an alkane, including n-
30 heptane, hexane and isooctane, and high and low boiling petroleum ethers and substituted aromates, incl toluene and xylenes. Crystalline citalopram base may be re-crystallised from the same solvents.

- The pharmaceutically acceptable salt of citalopram, such as the hydrobromide or hydrochloride, may be prepared by methods known in the art. So, the base may be reacted with either the calculated
35 amount of acid in a water miscible solvent, such as acetone or ethanol, with subsequent isolation of the salt by concentration and cooling, or with an excess of the acid in a water immiscible solvent, such as ethylether, ethylacetate or dichloromethane, with the salt separating spontaneously. The

hydrobromide or hydrochloride of citalopram obtained by the method of the invention has a very high purity, preferably more than 99,8% pure, most preferably more than 99,9 % purity. Other salts of citalopram, e.g. the oxalate, may also be obtained in a very pure form by this process.

- 5 The cyanide exchange reactions mentioned above may be carried out as described in the patent applications mentioned above.

In particular, when Z is halogen, or $\text{CF}_3\text{-(CF}_2)_n\text{-SO}_2\text{-O-}$ wherein n is an integer in the range 0-8, incl., the conversion to a cyano group may be carried out by reaction with a cyanide source, for
10 example KCN, NaCN, CuCN, Zn(CN)_2 or $(\text{R}^4)_4\text{NCN}$ where R^4 indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a palladium catalyst and a catalytic amount of Cu^+ or Zn^{2+} , or with Zn(CN)_2 in the presence of a palladium catalyst.

- 15 The cyanide source is used in a stoichiometric amount or in excess, preferably 1-2 equivalents are used pr. equivalent starting material. R^4N^+ may conveniently be $(\text{Bu})_4\text{N}^+$. The cyanide compound is preferably NaCN or KCN or Zn(CN)_2 .


The palladium catalyst may be any suitable Pd(0) or Pd(II) containing catalyst, such as $\text{Pd(PPh}_3)_4$,
20 $\text{Pd}_2(\text{dba})_3$, $\text{Pd(PPh)}_2\text{Cl}_2$, etc. The Pd catalyst is conveniently used in an amount of 1-10, preferably 2-6, most preferably about 4-5 mol%.

Catalytic amounts of Cu^+ and Zn^{2+} , respectively, means substoichiometric amounts such as 0.1 - 5, preferably 1 - 3 %. Conveniently, about $\frac{1}{2}$ eq. is used per eq. Pd. Any convenient source of Cu^+
25 and Zn^{2+} may be used. Cu^+ is preferably used in the form of CuI and Zn^{2+} is conveniently used as the Zn(CN)_2 salt.

When Z is Br or I, the conversion to a cyano group may also be carried out by reaction with Cu(CN) without catalyst. In a preferred embodiment, the reaction is performed at elevated temperature.

30

In another aspect of the invention, the reaction is performed in an ionic liquid of the general formula $(\text{R}^5)_4\text{N}^+$, X^- , wherein R^5 are alkyl-groups or two of the R^5 groups together form a ring and X^- is the counterion. In one embodiment of the invention, $(\text{R}^5)_4\text{N}^+\text{X}^-$ represents



In another particular aspect, the reaction is conducted with apolar solvents such as benzene, xylene or mesitylene and under the influence of microwaves by using *i.e.* Synthewave 1000™ by Prolabo. In a particular aspect, the reaction is performed without added solvent.

5

The temperature ranges are dependent upon the reaction type. If no catalyst is present, preferred temperatures are in the range of 100-200°C. However, when the reaction is conducted under the influence of microwaves, the temperature in the reaction mixture may raise to above 300 °C. More preferred temperature ranges are between 120-170°C. The most preferred range is 130-150°C.

- 10 If catalyst is present, the preferred temperature range is between 0 and 100°C. More preferred are temperature ranges of 40-90°C. Most preferred temperature ranges are between 60-90°C.

Other reaction conditions, solvents, etc. are conventional conditions for such reactions and may easily be determined by a person skilled in the art.

15

When Z is Cl or Br, the conversion to a cyano group may also be carried out by reaction with a cyanide source, for example KCN, NaCN, CuCN, Zn(CN)₂ or (R⁴)₄NCN where (R⁴)₄ indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a nickel catalyst.

20

The nickel catalyst may be any suitable Ni(0) or Ni(II) containing complex which acts as a catalyst, such as Ni(PPh₃)₃, (σ-aryl)-Ni(PPh₃)₂Cl, etc. The nickel catalysts and their preparation are described in WO 96/11906, EP-A-613720 or EP-A-384392.

- 25 In one embodiment of the invention, the reaction is carried out in the presence of a catalytic amount of Cu⁺ or Zn²⁺.

In a particularly preferred embodiment, a Nickel(0) complex is prepared *in situ* before the cyanation reaction by reduction of a Nickel(II) precursor such as NiCl₂ or NiBr₂ by a metal, such as zinc, magnesium or mangan in the presence of excess of complex ligands, preferably triphenylphosphin.

30

The Ni-catalyst is conveniently used in an amount of 0.5-10, preferably 2-6, most preferably about 4-5 mol%.

Catalytic amounts of Cu^+ and Zn^{2+} , respectively, mean substoichiometric amounts such as 0.1 - 5, preferably 1 - 3 %. Any convenient source of Cu^+ and Zn^{2+} may be used. Cu^+ is preferably used in the form of CuI and Zn^{2+} is conveniently used as the $\text{Zn}(\text{CN})_2$ salt or formed in situ by reduction of a Nickel (II) compounds using zinc.

5

The Ni catalysts are *i.e.* Ni (0), Pd(0) or Pd(II) catalysts as described by Sakakibara et. al. in Bull. Chem. Soc. Jpn., **61**, 1985-1990, (1988). Preferred catalysts are $\text{Ni}(\text{PPh}_3)_3$ or $\text{Pd}(\text{PPh}_3)_4$, or $\text{Pd}(\text{PPh})_2\text{Cl}_2$.

- 10 The reactions may be performed in any convenient solvent as described in Sakakibara et. al. in Bull. Chem. Soc. Jpn., **61**, 1985-1990, (1988). Preferred solvents are acetonitril, ethylacetat, THF, DMF or NMP.

- When Z is CHO, the conversion to a cyano group may be carried out by conversion of the formyl group to an oxime or similar group by reaction with a reagent $\text{R}^6\text{-V-NH}_2$ wherein R^6 is hydrogen, optionally substituted alkyl, aryl or heteroaryl and V is O, N or S, followed by dehydration with a common dehydrating agent, for example thionylchloride, acetic anhydride/pyridine, pyridine/HCl or phosphor pentachloride. Preferred reagents $\text{R}^6\text{-V-NH}_2$ are hydroxylamin and compounds wherein R^6 is alkyl or aryl and V is N or O.

20

When Z is $-\text{COOH}$, the conversion to a cyano group may be carried out via the corresponding acid chloride, ester or amide.

- The acid chloride is conveniently obtained by treatment of the acid with POCl_3 , PCl_5 or SOCl_2 neat or in a suitable solvent, such as toluene or toluene comprising a catalytic amount of N,N-dimethylformamide. The ester is obtained by treatment of the acid with an alcohol, in the presence of an acid, preferably a mineral acid or a Lewis acid, such as HCl, H_2SO_4 , POCl_3 , PCl_5 or SOCl_2 . Alternatively, the ester may be obtained from the acid chloride by reaction with an alcohol. The ester or the acid chloride is then converted to an amide or by amidation with ammonia or an alkylamine, preferably t-butyl amine.

30

The conversion to amide may also be obtained by reaction of the ester with ammonia or an alkylamine under pressure and heating.

- 35 The amide group is then converted to a cyano group by dehydration. The dehydrating agent may be any suitable dehydrating agent, and the optimal agent may easily be determined by a person skilled in the art. Examples of suitable dehydrating agents are SOCl_2 , POCl_3 and PCl_5 , preferably SOCl_2 .

In a particularly preferred embodiment, the carboxylic acid is reacted with an alcohol, preferably ethanol, in the presence of POCl_3 , in order to obtain the corresponding ester, which is then reacted with ammonia thereby giving the corresponding amide, which in turn is reacted with SOCl_2 in
5 toluene comprising a catalytic amount of N,N-dimethylformamide.

Alternatively, a compound where Z is $-\text{COOH}$ may be reacted with chlorosulfonyl isocyanate in order to form the nitrile, or treated with a dehydrating agent and a sulfonamide.

10 When Z is $-\text{NHR}^1$, where R^1 is hydrogen, the conversion into cyano is preferably performed by diazotation and followed by reaction with CN^- . Most preferably NaNO_2 and CuCN and/or NaCN are used. When R^1 is alkylcarbonyl, it is initially subjected to hydrolysis thereby obtaining the corresponding compound wherein R^1 is H which is then converted as described above. The hydrolysis may be performed either in acidic or basic environment.

15

The compounds of formula (II) may be prepared as described in DE 2,657,013, WO 0011926 and WO 0013648, WO 9819513, WO 9819512 and WO 9900548.

Throughout this specification with claims halogen means chloro, bromo or iodo.

20

The term alkyl refers to a branched or unbranched alkyl group, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, and 2-methyl-1-propyl.

The term aryl refers to a carbocyclic aromatic group, in particular phenyl. Aralkyl refers to an
25 arylalkyl group wherein aryl and alkyl is as defined above. The aryl and aralkyl groups may optionally be substituted, e.g. with alkyl groups, forming for example tolyl.

The pharmaceutical compositions of the invention may be administered in any suitable way and in any suitable form, for example orally in the form of tablets, capsules, powders or syrups, or
30 parenterally in the form of usual sterile solutions for injection. Preferably the pharmaceutical compositions of the invention are administered orally.

The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active ingredient with ordinary adjuvants
35 and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: Corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive colourings, aroma, preservatives etc. may be used provided that they are compatible with the active ingredients.

In particular, the formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

Solutions for injections may be prepared by solving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilisation of the solution and filling in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

According to the present invention, the base of citalopram has been found to be crystalline with stable and nice white crystals and it has been found that the base may easily be crystallised in a very pure form. So for example more than 99.8% w/w pure citalopram base was obtained by crystallisation from up to 95% pure hydrobromide without further purification. Accordingly, the process of the invention for preparing salts of citalopram has been found to give the salts as very pure products of pharmaceutically acceptable quality. Accordingly, the yield may be improved substantially during the manufacture of citalopram.

Finally, it has been found that the crystalline citalopram base may be formulated into very good and stable solid formulations with good release properties.

The invention is further illustrated by the following examples.

Example 1

Crystallisation of R,S-Citalopram as the free base.

1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile.

1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile hydrobromide (101 grams, 0.25 mole) prepared from 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, is suspended in water (500 ml) and toluene (500 ml). NaOH (60 ml, 5 N (aq)) is added and the mixture (pH>10) is stirred for 15 min. before the phases are separated. The organic phase is washed with water (2x100 ml) and filtered through a pad of filter help. The volatiles are removed *in vacuo* and the title compound is obtained as an oil. n-Heptane (400 ml) is added and the mixture is heated to 70 °C. On cooling, crystals form. The white crystals of the title compound are filtered off and dried at ambient temperature over night in *vacuo*. Yield: 75.4 grams (93%). DSC(onset, open capsule): 91.3-91.8 °C DSC (onset, closed capsule): 92.8 °C. Purity: (> 99.8 % (peak area)).

Anal. calcd. for C₂₀H₂₁N₂F₁₀O₁; C, 74.04; H, 6.54; N, 8.64. Found C, 74.01; H, 6.49; N, 8.59.

¹H-NMR (DMSO-d₆, 500 MHz): 1.21 (1H, m), 1.29 (1H, m), 2.02 (6H, s), 2.09-2.23 (4 H, m), 5.15 (1H, d J=12.5 Hz), 5.22 (1H, d J=12.5 Hz), 7.16 (2H, t J=8.5 Hz), 7.60 (2H, dt J=8.5 Hz J=1.2 Hz), 7.76 (1H, d J= 8.5 Hz), 7.79 (1H, d J=8.5 Hz), 7.80 (1H, s). ¹³C-NMR (DMSO-d₆, 125 MHz):
21.8, 38.3, 45.0, 58.8, 71.0, 90.7, 110.5, 115.1 (d J=22 Hz), 118.8, 123.1, 125.1, 127.0 (d J=8 Hz),
132.0, 140.0 (d J=3 Hz), 140.5, 149.5, 161.3 (d J=245Hz).

Example 2

- 10 a) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.
- 15 b) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the very pure free base of citalopram is precipitated by cooling.
- 20 c) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.
- 25 d) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the extremely pure free base of citalopram is precipitated by cooling.
- 30

Example 3

Wet granulation and preparation of tablets

35

The batch size was 200 g and the granulation was performed in a small-scale laboratory high shear mixer (Micromixer).

Citalopram base was sieved through a sieve aperture of 0.3 mm. The ingredients of the intragranular phase (1 - 4 in Table 1) were mixed at 600 rpm. 25 ml of purified water (5) was added in 30 sec and the granulation terminated after a total processing time of 3 min. The granulate was wet sieved through a 0.7 mm sieve aperture and dried at 40 °C in 30 minutes to equilibrium relative humidity of 32 %. The dried granulate was finally sieved through a 0.7 mm sieve aperture.

The dried granulate was mixed for 3 minutes with the extragranular phase (6 – 7) in a Turbula mixer and finally mixed with the lubricant (8) for 30 sec.

10

	Materials	%
1	Citalopram (base)	16.00
2	Kollidon VA64	2.32
3	Lactose 350 mesh	38.98
4	Corn starch	20.00
5	Purified water	25
6	Avicel PH 200 (Microcrystalline cellulose)	20.00
7	Ac-Di-Sol (Croscarmellose sodium)	2.00
8	Magnesium stearate	0.7

Table 1. Composition of the tablets.

Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 2.

15

Parameter	Values
Tablet strength, mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating (special doomed)
Mean disintegration time, min	1.77
Mean chrushing strength, N	69.1
Mean tablet weight, mg	125.4
RSD tablet weight, %	0.42
Friability, %	0.3

Table 2. Tablet characteristics.

The tablets produced had satisfactory technical properties.

Example 4**Melt granulation**

- 5 The batch size was 200 g. Citalopram base was sieved through a sieve aperture of 0.3 mm. The granulation was performed in a small-scale laboratory high shear mixer (Micromixer)

The ingredients of the intra-granular phase (1 - 3 in Table 3) were mixed at 1200 rpm.

The jacket temperature was 80 °C. The granulation process was terminated after 3.5 min. The

- 10 granulate was sieved through a sieve aperture of 1.0 mm and mixed with the extra-granular phase (4, 5) for 3 min. and with the lubricant (6) for 30 sec.

	Materials	%
1	Citalopram (base)	16.00
2	Polyethyleneglycol 6000	9.14
3	Lactose 350 mesh	38.98
4	Avicel PH 200 (Microcrystalline cellulose)	30.00
5	Kollidon CL (Cross-linked povidone)	4.00
6	Magnesium stearate	0.7

Table 3. Composition of the tablet.

- 15 Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 4.


Parameter	Values
Tablet strength, 20 mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating , Special doomed
Mean disintegration time, min	1.0
Mean crushing strength, N	55.5
Mean tablet weight, mg	125.6
RSD tablet weight, %	0.5
Friability, %	0.4

Table 4. Tablet characteristics.

- 20 The tablets produced had satisfactory technical properties.

CLAIMS


1. A process for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free and precipitated in crystalline form from a solvent, and thereafter separated from the solvent and optionally re-crystallised one or more times, and then transferred into a salt thereof.
2. The process of Claim 1 for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free from a crude salt or a crude mixture of citalopram.
3. A process for the manufacture of citalopram base or a salt of citalopram characterised in that one or more impurities of the formula



wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form, optionally re-crystallising said base one or more times and/or transferring said base into a salt thereof.

4. The process according to Claim 3 wherein the crude mixture of citalopram containing the compound of formula II as an impurity, is prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source.
5. The process according to Claim 3, wherein Z is halogen.
6. The process according to Claim 5, wherein Z is bromide or chloride.
7. The process according to any of Claims 3 to 6 wherein the crude mixture of citalopram is subjected to initial purification before the base of citalopram is precipitated in crystalline form.

8. The process according to any of Claims 3 to 6 wherein the crude mixture of citalopram is subjected to initial purification before a crude salt is formed from said crude mixture.
9. The process according to any of Claims 3 to 8 wherein the base of citalopram is set free from a crude salt or a crude mixture of citalopram by treatment with a base and optionally subjected to further purification before the base of citalopram is precipitated in crystalline form.
10. The process according to any of Claims 1 to 9 characterised in that the citalopram base is transferred into the hydrobromide or the hydrochloride salt of citalopram.
11. The process according to any of Claims 2-3, characterised in that the crude salt is a hydrobromide, hydrochloride, sulphate, oxalate, phosphate or nitrate salt.
12. The process according to Claim 11, wherein the crude salt is the sulphate, hydrobromide, or hydrochloride salt.
13. Crystalline base of citalopram, characterised in that it has a purity of more than 99.8 %w/w.
14. The crystalline base of citalopram according to Claim 13, characterised in that it has a purity of more than 99.9 %w/w.
15. Crystalline base of citalopram, prepared by a process characterised in that the base of citalopram is set free and precipitated in crystalline form and optionally re-crystallised one or more times.
16. The crystalline base of citalopram according to claim 15 characterised in that the base of citalopram is set free from a crude salt or a crude mixture of citalopram.
17. Crystalline base of citalopram, prepared by a process which is characterised in that one or more impurities of the formula



wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form and optionally re-crystallising said base one or more times.

18. The crystalline base of citalopram according to claim 17 wherein the crude mixture of citalopram containing the compound of formula II as an impurity, is prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source.
19. The crystalline base of citalopram according to claim 17 wherein Z is halogen.
20. The crystalline base of citalopram according to claim 19 wherein Z is bromide or chloride.
21. The crystalline base of citalopram according to any one of claims 17 to 20 wherein the crude mixture of citalopram is subjected to initial purification before the base of citalopram is precipitated in crystalline form.
22. The crystalline base of citalopram according to any one of claims 17 to 20 wherein the crude mixture of citalopram is subjected to initial purification before a crude salt is formed from said crude mixture.
23. The crystalline base of citalopram according to any one of claims 17 to 21 wherein the base of citalopram is set free from a crude salt or a crude mixture of citalopram by treatment with a base and optionally subjected to further purification before the base of citalopram is precipitated in crystalline form.

24. The crystalline base of citalopram according to any one of claims 15 to 17 characterised in that the crude salt is a hydrobromide, hydrochloride, sulphate, oxalate, phosphate or nitrate salt.
25. The crystalline base of citalopram according to claim 24 wherein the crude salt is the sulphate, hydrobromide or hydrochloride salt.
26. The crystalline base of citalopram according to any one of claims 15 to 25, characterised in that it has a purity of more than 99.8 %w/w.
27. The crystalline base of citalopram according to claim 26, characterised in that it has a purity of more than 99.9% w/w.
28. A pharmaceutical composition containing the crystalline base of citalopram according to any one of Claims 13 to 27.
29. A pharmaceutical composition according to Claim 28 which is a tablet prepared by
- a) direct compression of citalopram, optionally in admixture with pharmaceutically acceptable adjuvants;
 - b) compression of a wet granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants; or
 - c) compression of a melt granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants.
30. The pharmaceutical composition according to claim 28 or 29, characterised in that it contains the racemic mixture of citalopram base.