



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2753481 A1 2010/09/02

(21) **2 753 481**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2010/02/25
(87) Date publication PCT/PCT Publication Date: 2010/09/02
(85) Entrée phase nationale/National Entry: 2011/08/23
(86) N° demande PCT/PCT Application No.: US 2010/025446
(87) N° publication PCT/PCT Publication No.: 2010/099342
(30) Priorité/Priority: 2009/02/25 (US61/155,364)

(51) Cl.Int./Int.Cl. *C07H 21/04* (2006.01),
C07H 21/00 (2006.01), *C12Q 1/68* (2006.01),
C40B 30/04 (2006.01), *C40B 40/06* (2006.01),
C12N 15/113 (2010.01)

(71) Demandeur/Applicant:
CEPHEID, US

(72) Inventeurs/Inventors:
MICHOT, BERNARD, FR;
DELFOUR, OLIVIER, FR;
PERSING, DAVID H., US

(74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : PROCÉDES DE DETECTION DU CANCER DU POUMON
(54) Title: METHODS OF DETECTING LUNG CANCER

(57) Abrégé/Abstract:

Methods of diagnosing lung cancer in a sample from a patient by detecting changes in expression of one or more target RNAs associated with lung cancer are provided.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 September 2010 (02.09.2010)

(10) International Publication Number
WO 2010/099342 A3

(51) International Patent Classification:
G01N 33/48 (2006.01)

(21) International Application Number:
PCT/US2010/025446

(22) International Filing Date:
25 February 2010 (25.02.2010)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/155,364 25 February 2009 (25.02.2009) US

(71) Applicant (*for all designated States except US*):
CEPHEID [US/US]; 904 Caribbean Drive, Sunnyvale,
CA 94089 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MICHOT, Bernard**
[FR/FR]; Cepheid Europe S.a., Vira Solelh, F-81470
Maurens-scopont (FR). **DELFOUR, Olivier** [FR/FR];
Cepheid Europe s.a., Vira Solelh, F-81470 Maurens-Sco-
pont (FR). **PERSING, David, H.** [US/US]; Cepheid, 904
Caribbean Drive, Sunnyvale, CA 94089 (US).

(74) Agent: **SCARR, Rebecca, B.**; Andrus, Sceales, Starke &
Sawall, LLP, 100 East Wisconsin Avenue, Suite 1100,
Milwaukee, WI 53202 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD,
SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR,
TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ,
TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))*

(88) Date of publication of the international search report:
28 October 2010

(54) Title: METHODS OF DETECTING LUNG CANCER

(57) Abstract: Methods of diagnosing lung cancer in a sample from a patient by detecting changes in expression of one or more target RNAs associated with lung cancer are provided.



WO 2010/099342 A3

METHODS OF DETECTING LUNG CANCER

[001] This application claims priority to U.S. Provisional Application No. 61/155,364, filed February 25, 2009, which is incorporated by reference herein in its entirety for any purpose.

1. BACKGROUND

[002] Lung cancer is the leading cause of death due to cancer in the world. Lung cancer is categorized into two types, small cell lung cancer (“SCLC”) and non-small cell lung cancer (“NSCLC”). SCLC is an extremely aggressive form of lung cancer with poor prognosis and a median length of survival after diagnosis of about 1 to 3 months. About 80% of lung cancer cases are categorized as NSCLC, which is categorized into three subtypes: adenocarcinoma, squamous cell carcinoma and large cell carcinoma. Greater than 85% of all NSCLCs are either adenocarcinoma or squamous-cell carcinoma.

[003] Lung cancer is difficult to diagnose in the early stages because it may manifest no outward symptoms. When symptoms do occur, they can vary depending on the type, location and spreading pattern of the cancer, and therefore, are not readily associated with cancer. Often, lung cancer is only correctly diagnosed when it has already metastasized.

[004] Current techniques for diagnosing lung cancer include chest x-ray and/or computed tomography (“CT”) scan. Diagnosis by one of these techniques is usually confirmed by a more invasive procedure, such as transthoracic needle biopsy or transbronchial biopsy, which may still result in misdiagnosis of lung cancer. (Butnor (2008) *Arch. Pathol. Lab. Med.* 132:1118-1132.)

[005] Despite advances in treatment (*e.g.*, by surgery, chemotherapy, radiation or a combination), the prognosis for lung cancer remains poor, with only 15% of patients surviving more than 5 years from the time of diagnosis. Of the most common NSCLCs, adenocarcinoma progresses more rapidly and therefore has a poorer prognosis than squamous-cell carcinoma, which takes several years to develop and is therefore more likely to be diagnosed in an early stage.

[006] One proposal for reducing the mortality and morbidity of lung cancer is to institute regular screening of high-risk individuals, *e.g.*, those who smoke or have smoked heavily for a certain period of time, in order to detect and treat lung cancer in asymptomatic individuals. In this way, early stage lung cancer can be eradicated by

surgical resection, which is thought to be the only realistic option for a cure. (Field *et al.* (2008) *Br. J. Cancer* 99:557-562).

[007] There remains a need for molecular markers in lung cancer.

2. SUMMARY

[008] In some embodiments, methods of detecting the presence of lung cancer in a subject are provided. In some embodiments, a method comprises detecting a level of at least one target RNA in a sample from the subject. In some embodiments, the at least one target RNA: (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a level of at least one target RNA in the sample that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer in the subject. In some embodiments, the method further comprises comparing the level of the at least one target RNA in the sample to a normal level of the at least one target RNA.

[009] In some embodiments, methods for facilitating the detection of lung cancer in a subject are provided. In some embodiments, a method comprises detecting a level of at least one target RNA in a sample from the subject. In some embodiments, the target RNA (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a method comprises communicating the results of the detection to a medical practitioner for the purpose of determining whether the subject has lung cancer.

[0010] In some embodiments, detecting a level of at least one target RNA in a sample comprises hybridizing nucleic acids of the sample with at least one polynucleotide that is complementary to a target RNA in the sample or to a complement thereof; and detecting at least one complex comprising a polynucleotide hybridized to at least one

nucleic acid selected from the target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA.

[0011] In some embodiments, a method for detecting the presence of lung cancer in a subject comprises (a) obtaining a sample from the subject; (b) providing the sample to a laboratory for detection of the level of at least one target RNA in the sample; and (c) receiving from the laboratory a communication indicating the level of at least one target RNA in the sample. In some embodiments, the at least one target RNA (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a level of at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer.

[0012] In some embodiments, levels of at least two, at least three, or at least five target RNAs are detected. In some embodiments, detection of levels of at least one, at least two, at least three, or at least five target RNAs that are greater than a normal level of the at least one target RNA indicates the presence of lung cancer.

[0013] In some embodiments, a method further comprises detecting a level of at least one target RNA that: (i) does not specifically hybridize to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (ii) does not comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; and (iii) does not comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

[0014] In some embodiments, a method comprises detection of at least one target RNA (a) is capable of specifically hybridizing to a polynucleotide sequence in Table 6; or (b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 6. In some such embodiments, detection of a level of

the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of non-small cell lung cancer.

[0015] In some embodiments, a method comprises detection of at least one target RNA (a) is capable of specifically hybridizing to a polynucleotide sequence in Table 7; or (b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 7. In some such embodiments, detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of squamous cell carcinoma.

[0016] In some embodiments, a method comprises detection of at least one target RNA (a) is capable of specifically hybridizing to a polynucleotide sequence in Table 8; or (b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 8. In some such embodiments, detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of adenocarcinoma.

[0017] In some embodiments, a method comprises detection of at least one target RNA (a) is capable of specifically hybridizing to a polynucleotide sequence in Table 9; or (b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 9. In some such embodiments, detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of aggressive lung cancer.

[0018] In some embodiments, a method comprises detection of at least one target RNA (a) is capable of specifically hybridizing to a polynucleotide sequence in Tables 32 or 33; or (b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 32 or Table 33.

[0019] In some embodiments, a method of detecting the presence of lung cancer is provided that comprises detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452. In some embodiments, a level of at least one target RNA in the sample that is reduced relative to a normal level of the at least one target RNA indicates the

presence of lung cancer in the subject. In some embodiments, a method comprises comparing the level of the at least one target RNA in the sample to a normal level of the at least one target RNA.

[0020] In some embodiments, a method of facilitating the detection of lung cancer in a subject is provided, comprising (a) detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from 1 to 397, 1363 to 1707, and 2312 to 2452; and (b) communicating the results of the detection to a medical practitioner for the purpose of determining whether the subject has lung cancer.

[0021] In some embodiments, a method of detecting the presence of lung cancer in a subject is provided, wherein the method comprises (a) obtaining a sample from the subject, (b) providing the sample to a laboratory for detection of the level of at least one target RNA in the sample, wherein the at least one target RNA (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; and (c) receiving from the laboratory a communication indicating the level of at least one target RNA in the sample. In some embodiments, a level of at least one target RNA that is reduced relative to a normal level of the at least one target RNA indicates the presence of lung cancer.

[0022] In some embodiments, a method further comprises detecting a level of at least one second target RNA in a sample from the subject, wherein the at least one second target RNA: (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or (iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a level of at least one second target RNA in the sample that is greater than a normal level of the at least one second target RNA indicates the presence of lung cancer in the subject.

[0023] In some embodiments, a method comprises isolating nucleic acids from a sample. In some embodiments, the nucleic acids comprise RNA that has been separated from DNA. In some embodiments, a target RNA in its mature form comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[0024] In some embodiments, synthetic polynucleotides are provided. In some embodiments, compositions comprising a plurality of synthetic polynucleotides are provided. In some embodiments, a synthetic polynucleotide comprises a first region, wherein the first region comprises a sequence of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15 contiguous nucleotides that is identical or complementary to a sequence of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15 contiguous nucleotides of one of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680. In some embodiments, a synthetic polynucleotide comprises a detectable label. In some embodiments, the detectable label is a FRET label. In some embodiments, the first region is identical or complementary to a region of a target RNA. In some embodiments, the polynucleotide further comprises a second region that is not identical or complementary to a region of the target RNA.

[0025] In some embodiments, kits are provided. In some embodiments, a kit comprises a synthetic polynucleotide. In some embodiments, a kit comprises a composition comprising a plurality of synthetic polynucleotides. In some embodiments, a kit comprises at least one polymerase. In some embodiments, a kit comprises dNTPs.

[0026] Further embodiments and details of the inventions are described below.

3. BRIEF DESCRIPTION OF THE FIGURES

[0027] FIG. 1 shows an electropherogram obtained on an Agilent Bioanalyser 2100 to assess the quality of total RNA purified as described in Example 1 from A549 human adenocarcinoma cell line.

4. DETAILED DESCRIPTION

4.1. Detecting lung cancer

4.1.1. General methods

[0028] Methods of detecting lung cancer by measuring levels of target RNAs are provided. In some embodiments, methods are presented for detecting non-small cell lung cancer in a human. In some embodiments, a method comprises detecting altered levels of

at least one target RNA relative to normal levels of the at least one target RNA. In some embodiments, elevated levels of one or more target RNAs are indicative of lung cancer. In some embodiments, reduced levels of one or more target RNAs are indicative of lung cancer. In some embodiments, the method comprises detecting an altered level of at least one target RNA that is capable of specifically hybridizing to a sequence selected from the sequences in Tables 1, 2, 6 to 9, 18, 20, 23, 27, 28, 30, and 32 to 34. In some embodiments, the method comprises detecting an altered level of at least one target RNA that is capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the method comprises detecting an altered level of at least one target RNA selected from the microRNAs in Tables 4, 5, and 38. In some embodiments, the method comprises detecting an altered level of at least one target RNA that comprises at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of a sequence selected from SEQ ID NOs.: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the method comprises detecting an altered level of at least one target RNA that comprises a sequence that is complementary to at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of a sequence selected from SEQ ID NO.: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, an altered level is an elevated level. In some embodiments, an altered level is a reduced level. In some embodiments, a method comprises detecting an elevated level of at least one target RNA and a reduced level of at least one other target RNA. In some embodiments, the target RNA, in its mature form, comprises fewer than 30 nucleotides. The target RNA, in some embodiments, is a microRNA.

[0029] In the present disclosure, “a sequence selected from” encompasses both “one sequence selected from” and “one or more sequences selected from.” Thus, when “a sequence selected from” is used, it is to be understood that one, or more than one, of the listed sequences may be chosen.

[0030] Detection of a level of target RNA that is greater than a normal level of target RNA indicates the presence of lung cancer in the patient from whom the sample is taken. In some embodiments, the detecting is done quantitatively. In other embodiments, the detecting is done qualitatively. In some embodiments, detecting a target RNA

comprises forming a complex comprising a polynucleotide and a nucleic acid selected from a target RNA, a DNA amplicon of a target RNA, and a complement of a target RNA. In some embodiments, the level of the complex is then detected and compared to a normal level of the same complex. The level of the complex, in some embodiments, correlates with the level of the target RNA in the sample.

[0031] “Non-small cell lung cancer” or “NSCLC” is one of two categories of lung cancer found in humans. About 80% of patients diagnosed with lung cancer have non-small cell lung cancer. NSCLC is further broken down into three sub-categories, depending on the cells in which they originate: (i) adenocarcinoma, which originates in the cells that line the alveoli and make substances such as mucus; (ii) squamous cell or epidermoid carcinoma, which originates in the squamous cells; and (iii) large cell carcinoma, which may originate in several different types of large cells. More than 50% of patients with NSCLC have either adenocarcinoma or squamous cell carcinoma. The histology class nonsquamous cell carcinoma includes both adenocarcinoma and large cell carcinoma.

[0032] Cancer can be divided into clinical and pathological stages. The clinical stage is based on all available information about a tumor, such as information gathered through physical examination, radiological examination, endoscopy, etc. The pathological stage is based on the microscopic pathology of a tumor.

[0033] The TNM (tumor, node, metastasis) system classifies a cancer by three parameters – the size of the tumor and whether it has invaded nearby tissues, involvement of lymph nodes, and metastases. T (tumor) is assigned a number from 1 to 4, according to the size and extent of the primary tumor. N (node) is assigned a number from 0 to 3, in which 0 means no spreading to the lymph nodes, 1 is spreading to the closest lymph nodes, and 3 is spreading to the most distant and greatest number of lymph nodes, and 2 is intermediate between 1 and 3. M (metastasis) is assigned 0 for no distant metastases, or 1 for distant metastases beyond regional lymph nodes.

[0034] For lung cancer, Overall Stage Grouping assigns a cancer a roman numeral of 0, I, II, III, and IV, and a letter, A or B, depending on the stage. Stage 0 is carcinoma in situ, which usually does not form a tumor. Stages IA (T1N0M0) and IB (T2N0M0) is cancer that is localized to one part of the body. Stage IIA (T1N1M0) and IIB (T2N1M0 and T3N0M0) is cancer that is localized, but more advanced. Stage IIIA (T1-3N2M0 or

T3N1M0) and IIIB (any T4 or any N3M0) cancer is also locally advanced. Stage IV (any M1) is cancer that has metastasized. As used herein, the term “early stage cancer” refers to Stages IA and IB and Stages IIA and IIB cancers.

[0035] As used herein, an “aggressive” form of lung cancer is a lung cancer that advances quickly from one stage to the next and/or metastasizes at an early stage, resulting in a poor prognosis for the patient.

[0036] Tables 1 and 2, below, list 397 hybridization probes that have been found to be complementary to, and hybridize with, target RNAs in lung cancer cells. These target RNAs were detected at elevated levels or at reduced levels in certain primary tumors and/or human lung cancer cell lines (respectively Examples 1 and 2). Two hundred seventy-five of the probes are complementary to, and hybridize with, novel target RNA species that are expressed in human cells. The other one hundred and twenty-two probes are complementary to, and hybridize with, publicly known microRNAs that have been accessioned by others into miRBase (<http://microrna.sanger.ac.uk/>; see Griffiths-Jones S. *et al.* (2007) *Nucl. Acids Res.* 36:154-158). These latter probes are designated by either “mir” or “let” in Tables 1 and 2.

[0037] Tables 18 to 21, below, list hybridization probes that have been found to be complementary to, and hybridize with, target RNAs that were detected at elevated levels in certain primary tumors (see Example 4). Certain probes listed in Tables 18 and 20 are complementary to, and hybridize with, novel target RNA species that are expressed in human cells. Other probes in Tables 18 and 20 are complementary to, and hybridize with, publicly known microRNAs that have been accessioned by others into miRBase (<http://microrna.sanger.ac.uk/>; see Griffiths-Jones S. *et al.* (2007) *Nucl. Acids Res.* 36:154-158). These latter probes are designated by either “mir” or “let.”

[0038] Tables 23 and 24, below, list hybridization probes that have been found to be complementary to, and hybridize with, target RNAs that were detected at reduced levels in certain primary tumors (see Example 4). Certain probes listed in Tables 23 and 24 are complementary to, and hybridize with, novel target RNA species that are expressed in human cells. Other probes in Tables 23 and 24 are complementary to, and hybridize with, publicly known microRNAs that have been accessioned by others into miRBase (<http://microrna.sanger.ac.uk/>; see Griffiths-Jones S. *et al.* (2007) *Nucl. Acids Res.* 36:154-158). These latter probes are designated by either “mir” or “let.”

[0039] Tables 27 and 28, below, list hybridization probes that have been found to be complementary to, and hybridize with, target RNAs that were detected at elevated levels in certain lung cancer cell lines (see Example 5). Certain probes listed in Tables 27 and 28 are complementary to, and hybridize with, novel target RNA species that are expressed in human cells. Other probes in Tables 27 and 28 are complementary to, and hybridize with, publicly known microRNAs that have been accessioned by others into miRBase (<http://microrna.sanger.ac.uk/>; see Griffiths-Jones S. *et al.* (2007) *Nucl. Acids Res.* 36:154-158). These latter probes are designated by either “mir” or “let.”

[0040] Table 30, below, lists hybridization probes that have been found to be complementary to, and hybridize with, target RNAs that were detected at reduced levels in certain lung cancer cell lines (see Example 5). Certain probes listed in Table 30 are complementary to, and hybridize with, novel target RNA species that are expressed in human cells. Other probes in Table 30 are complementary to, and hybridize with, publicly known microRNAs that have been accessioned by others into miRBase (<http://microrna.sanger.ac.uk/>; see Griffiths-Jones S. *et al.* (2007) *Nucl. Acids Res.* 36:154-158). These latter probes are designated by either “mir” or “let.”

[0041] In Tables 1 and 2, respectively, the expression levels of target RNAs measured for each of the identified primary tumors, and for each of the identified cell lines, are expressed as fold-changes in expression relative to expression levels measured in normal human lung total RNA (see Examples 1 and 2). Similarly, in Tables 18 to 21, 23, 24, 27, 28, and 30, the expression levels of target RNAs measured for each of the identified primary tumors, and for each of the identified cell lines, are expressed as fold-changes in expression relative to expression levels measured in normal human lung total RNA (see Examples 4 and 5).

[0042] Table 6 lists target RNAs from Table 1 that are present at increased levels in NSCLCs. Table 7 lists target RNAs that are more frequently present at elevated levels in squamous cell carcinoma. In some embodiments, a method comprises detecting an elevated level of at least one target RNA from Table 7. In some such embodiments, detection of an elevated level of at least one target RNA from Table 7 is indicative of squamous cell carcinoma. Table 8 lists target RNAs that are more frequently present at elevated levels in adenocarcinoma. In some embodiments, a method comprises detecting an elevated level of at least one target RNA from Table 8. In some such embodiments, detection of an elevated level of at least one target RNA from Table 8 is indicative of

adenocarcinoma. Table 9 lists target RNAs that are present at increased levels in aggressive forms of lung cancer. In some embodiments, a method comprises detecting an elevated level of at least one target RNA from Table 9. In some such embodiments, detection of an elevated level of at least one target RNA from Table 9 is indicative of an aggressive form of lung cancer.

[0043] In some embodiments, a method comprises detecting multiple isomirs with a single probe. Detection of an elevated level of one or multiple isomirs is considered to be indicative of lung cancer. When multiple microRNAs having the same sequence but are expressed from different genes, one or more of the genes may be upregulated in a lung cancer patient. Detection of a microRNA expressed from any one of the genes is considered to be indicative of lung cancer.

[0044] For convenience of reference herein, and not by way of limitation, some “target RNA” species are denominated “microRNAs” in the tables set forth herein and Example 1. In some embodiments, the target RNA is a single mature microRNA capable of specifically hybridizing to a hybridization probe set forth in any of Tables 1, 2, 6 to 9, 18, 20, 23, 27, 28, 30, and 32 to 34. In some embodiments, a target RNA is a single mature microRNA that comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NO.: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a target RNA is a single mature microRNA that comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, target RNA may include a plurality of target RNAs, all of which are capable of specifically hybridizing to a single complementary probe sequence (for example, when two or more target microRNAs are isomirs). In some embodiments, the so-denominated “microRNA” is one or more RNA species capable of specifically hybridizing to the respective hybridization probe, such that one or more target RNAs do not meet canonical definitions for mature microRNAs. In some embodiments, a target RNA is an mRNA. In some embodiments, the “target RNA” is a piwi-interacting RNA (piRNA), *i.e.*, a small RNA expressed in animal cells that is distinct in size (26-31 nt) from microRNA and that forms distinct complexes with Piwi proteins that are involved in transcriptional gene silencing.

[0045] Mature human microRNAs are typically composed of 17 to 27 contiguous ribonucleotides, and often are from 19 to 25 nucleotides in length, or 21 or 22 nucleotides

in length. The sequences of some target microRNAs that can be detected in accordance with the present disclosure can be found within the pre-microRNA sequences shown in Tables 3, 22, 25, 29, 31, and 37 (SEQ ID NOs: 398 to 793, 1211 to 1362, 1708 to 2063, 2184 to 2311, 2453 to 2575, and 2681 to 2688, 2690, and 2691). In some embodiments, more than one mature target RNA is derived from a single pre-microRNA shown in Tables 3, 22, 25, 29, 31, and 37. The sequences of some publicly known mature microRNAs are shown below in Tables 4 and 5 (SEQ ID NOs: 794 to 1043, and 2692). Further, in some embodiments, a microRNA comprises at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, or at least 26 contiguous nucleotides of a sequence in Table 38 (SEQ ID NOs: 2576 to 2672).

[0046] While not intending to be bound by theory, mammalian microRNAs mature as described herein. A gene coding for a microRNA is transcribed, leading to production of a microRNA precursor known as the “pri-microRNA” or “pri-miRNA.” The pri-miRNA can be part of a polycistronic RNA comprising multiple pri-miRNAs. In some circumstances, the pri-miRNA forms a hairpin with a stem and loop, which may comprise mismatched bases. The hairpin structure of the pri-miRNA is recognized by Drosha, which is an RNase III endonuclease protein. Drosha can recognize terminal loops in the pri-miRNA and cleave approximately two helical turns into the stem to produce a 60-70 nucleotide precursor known as the “pre-microRNA” or “pre-miRNA.” Drosha can cleave the pri-miRNA with a staggered cut typical of RNase III endonucleases yielding a pre-miRNA stem loop with a 5' phosphate and an approximately 2-nucleotide 3' overhang. Approximately one helical turn of the stem (about 10 nucleotides) extending beyond the Drosha cleavage site can be essential for efficient processing. The pre-miRNA is subsequently actively transported from the nucleus to the cytoplasm by Ran-GTP and the export receptor Exportin-5.

[0047] The pre-miRNA can be recognized by Dicer, another RNase III endonuclease. In some circumstances, Dicer recognizes the double-stranded stem of the pre-miRNA. Dicer may also recognize the 5' phosphate and 3' overhang at the base of the stem loop. Dicer may cleave off the terminal loop two helical turns away from the base of the stem loop leaving an additional 5' phosphate and an approximately 2-nucleotide 3' overhang. The resulting siRNA-like duplex, which may comprise mismatches, comprises the mature microRNA and a similar-sized fragment known as the microRNA*. The

microRNA and microRNA* may be derived from opposing arms of the pri-miRNA and pre-miRNA. The mature microRNA is then loaded into the RNA-induced silencing complex (“RISC”), a ribonucleoprotein complex. In some cases, the microRNA* also has gene silencing or other activity.

[0048] It is understood that where a sequence includes thymine (T) bases, a target RNA may contain uracil (U) bases instead.

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
266-R4-1	GTCGCCCCCTCCCCCAAGTTGAGACTTGACGCTAC	1	7.33	1.43	1.97	3.81	-3.74	-2.28	-3.72	-1.19	-11.61	-2.38
673-L4-1	GCTCCCCCTCACTGTGAACCTTTTCACCCAGCTAACCTGC TCCTCAC	2	3.55	1.45	-1.26	1.44	(nd) < -6.23	-2.70	-5.73	-1.21	-6.23	-4.03
836-R4-1	AAATAATCATTCCAAATGGTTCTCCCTGCTATGATTCAC	3	6.78	1.06	(nd) < -6.03	2.23	-1.23	-2.34	-3.54	-1.02	-6.03	-1.43
3249-L4-1	GCGGAGCCGCGCGCCATCCCCGGAGCCGCGCCGCGCGG CCGCC	4	1.46	1.91	2.34	-1.38	-2.37	-3.34	-3.73	-2.89	-6.52	-2.00
3371-L4-1	TTTCCTTTCTCCCTCCACACCCCATGACTCCCCCACA CTTGAG	5	7.78	1.44	(nd) < -3.63	3.28	(nd) < -3.63	-1.54	-2.45	-1.89	(nd) < -3.63	-1.75
3717-L2-1	CCGCCCTCCCCATAGCCTCACCCCAACCCACTCACA	6	5.45	-1.46	2.40	2.66	-3.24	-3.42	-6.97	-22.96	-22.96	-2.75
3799-R3-1	CCAGAGGCCCCCGCGGCC	7	4.81	1.62	1.19	3.07	-4.42	-2.84	-5.33	-1.46	-18.87	-2.91
3872-L1-1	TCATTTTCTTGCTCTTCTTCCCTTATGCAC	8	> 6.17	nd	nd	> 2.09	nd	nd	nd	nd	nd	nd
3875-R3-1	CTCTCTCCCACTTTAATAA	9	> 20.25	nd	nd	> 2.90	nd	nd	nd	nd	nd	nd
3897-R3-1	CAGCCGCCTCCCCCTCAGCGTTAA	10	5.07	1.62	1.80	1.97	-3.20	-3.17	-4.86	-1.07	-2.38	-1.89
3923-R3-1	GCCTCTCACAAAGGATCTCCTTCATCCCTCTCTCC	11	17.53	1.49	(nd) < -1.39	4.86	(nd) < -1.39	2.05	-1.32	(nd) < -1.39	(nd) < -1.39	-1.16
3953-R3-2	ACTCCAGCCTCCGCCGCCCTCAGCTTCCCAGC	12	1.50	1.86	2.21	1.28	-2.29	-3.82	-5.17	nd G3	nd G3	-2.73
3995-L2-2	CTATAAACTTCGAAAAGTCCCTCCTCCTCACGT	13	6.16	1.93	1.05	3.01	-3.51	-2.18	-4.19	1.21	-5.70	-1.85
4026-R3-1	GGCGAGAGAGAAAGCCCCCCT	14	6.37	-1.17	2.06	3.81	-3.69	-3.06	-5.40	-1.70	-18.38	-4.41
4037-R3-2	GCCTGTTCCTGGCATGTACTGTAATTTATCT	15	6.57	1.44	(nd) < -1.28	2.17	(nd) < -1.28	2.17	2.34	2.64	6.81	-1.06
4143-R3-1	TCAGCGTCTTGCTCTCCTCTGGTA	16	> 7.19	nd	nd	> 2.08	nd	nd	nd	nd	nd	nd
4203-R3-2	GCACATTCACACTTCCCCAGAGGCAGGCTCCATAT	17	> 4.30	nd	nd	nd	nd	nd	nd	nd	nd	nd
4205-R3-2	GCCCTACAACTTCATCCTCACCACCTCACACCAC	18	> 3.92	nd	nd	> 2.47	nd	nd	nd	nd	nd	nd
4303-R1-1	AGTGCCCGCTCCTCCGACCTCCCTGCGCACC	19	4.55	2.06	1.53	2.49	-3.30	-1.83	-3.64	-2.09	-10.57	-2.26

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
4315-R3-2	TCCCCGGCCCTCTCCATTCTCGGCTCCGGAGCA	20	5.42	1.69	1.44	2.31	-2.65	-2.17	-2.74	-1.64	-4.81	-1.87
4361-R3-1	CGTCTCCCTCCCTCATGTGC	21	13.35	3.46	(nd) < -1.73	4.34	(nd) < -1.73	-1.41	-1.74	2.20	(nd) < -1.73	1.27
4440-L3-2	TTTGACATTCAGAGCACTGGGCAGAAATCACA	22	6.02	-1.51	1.14	1.60	(nd) < -3.39	-1.05	-1.83	-1.14	1.57	-2.75
4440-R3-2	GTCATAGTTACTCCCGCGGTTTACCCGCATTTC	23	9.42	1.72	1.53	3.52	-3.11	-1.63	-3.21	-1.13	-1.70	-2.72
4448-R3-1	CCTACCCCCAGCATCTCCTCAGCCATTGCC	24	> 2.43	nd	nd	> 1.62	nd	nd	nd	nd	nd	nd
4479-R3-1	AGCCCCCTGCCCGGAAATTCAAAACAACCTGC	25	3.07	3.03	-1.01	2.30	-7.11	-3.04	-3.90	-2.34	-5.33	-2.81
4593-R3-1	AGCAGATGACATAACTCCCCCGGCATCAG	26	8.31	2.80	1.23	1.93	-4.73	-2.65	-5.00	1.79	-9.77	1.20
4666-R4-1	GCCGACTCCCCCACAACACCTGCGGGTGGCAC	27	24.77	2.27	3.49	7.40	-2.36	-1.70	-2.05	2.37	-3.62	-1.01
4790-L4-1	AACCTGTCTCCCTCATTACTAGAAATTTCTGGG	28	> 14.59	> 1.58	nd	> 4.07	nd	nd	nd	nd	nd	nd
4829-R2-1	TCCCITTTGTGCTGCCCCGAGTGCCITCCCCCTG	29	12.09	1.52	1.69	5.73	-6.10	-1.29	-4.20	-1.12	-9.92	-2.88
4855-R3-1	GGGCTGCCGGGTCTCCCCGCTTCC	30	8.26	1.91	1.44	1.78	-3.56	-1.50	-1.81	1.87	-2.80	1.20
4875-R2-2	CACAGCCCCCTTCTGTGACTTCACAC	31	4.28	-1.44	1.02	2.82	(nd) < -5.97	-2.46	-5.03	-1.17	(nd) < -5.97	-4.78
4988-R4-1	CTCCTCCTCCCCGTCTTTGGATACCAAAACACTGGAC	32	9.10	1.72	1.15	2.70	(nd) < -3.75	-2.04	-2.34	-1.34	(nd) < -3.75	-1.55
5006-L3-1	ACAGCTCCCTCTGCTGGCTCC	33	6.58	1.11	2.00	5.46	(nd) < -1.43	(nd) < -1.43	(nd)	(nd)	(nd)	(nd)
5080-R3-1	CTTGCAAAGGGTCTCCTTCATCCCTCTCTCCA	34	5.91	-1.01	-1.35	2.60	(nd) < -4.53	1.28	-1.87	1.18	(nd) < -4.53	-3.70
5192-L3-2	CATTTTCCCTTCTCTCTATATACAGAA	35	1.54	1.24	(nd) < -1.61	2.92	(nd) < -1.61	(nd) < -1.61	-1.79	1.39	(nd) < -1.61	(nd) < -1.61
5342-L3-1	CACCACCAAACCAAAATGCCGGCTGCTCTCCTTCCA	36	1.95	1.46	2.07	1.19	(nd) < -4.91	-2.70	-4.03	-1.81	(nd) < -4.91	-1.99

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
5521-L2-1	GTCTTGGGTGGCCCTCCCCAGAGCACACCCTCT	37	4.19	(nd) < -1.44	2.44	2.24	(nd) < -1.44	(nd) < -1.44	(nd) < -1.44	(nd) < -1.44	(nd) < -1.44	(nd) < -1.44
5554-R2-1	CCCCACCCCTCATCAGCTGCTCCCAGAT	38	3.45	1.79	1.00	1.73	(nd) < -6.36	-2.78	-2.03	-1.57	(nd) < -6.36	-1.91
5638-R2-1	GGCCCTCCCCCTGCCCTGTGATAGGCTGCTTG	39	5.54	-1.87	2.47	3.18	-2.79	-3.13	-9.55	1.18	-15.66	-3.29
5640-L3-1	GCCATGGAACACCGTGCCTGCCCTCTCGAGA	40	3.56	2.33	1.22	2.16	-4.15	-1.75	-2.54	-1.31	-9.19	-2.51
5726-L3-1	TAATAAAATATCTTCTCACTGTGCCCTTG	41	> 8.13	nd	nd	> 1.83	nd	nd	nd	nd	nd	nd
5782-L3-1	GATTCAGCCCCCTTCCCCC	42	6.87	1.80	1.13	3.16	(nd) < -3.21	-1.78	-3.13	(nd) < -3.21	(nd) < -3.21	-2.66
5795-R1-1	CTGCCCTCCAAGAAATAAATTACCCGCAATTACT	43	3.41	1.05	1.52	2.16	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02
5836-R3-2	CATTACCCCCATTATCACAGCAGCCCCCATTC	44	13.99	1.99	(nd) < -1.09	2.26	(nd) < -1.09	(nd) < -1.09	(nd) < -1.09	(nd) < -1.09	(nd) < -1.09	(nd) < -1.09
5854-R3-1	CCCTCCCTCTCCGAAAGAAATGTGTAC	45	22.51	-1.01	1.54	4.54	(nd) < -4.11	-1.03	-2.56	5.86	-1.93	-1.08
5971-R3-1	CTGCTCAGCCTCCACATCTGT	46	5.67	-1.01	(nd) < -1.78	2.33	(nd) < -1.78	-1.08	-1.64	2.13	(nd) < -1.78	-1.38
6008-R1-1	ACAATACCCCCACCTTTTTCTGTACCTTAC	47	> 3.56	nd	nd	> 2.06	nd	nd	nd	nd	nd	nd
6016-R2-1	AAACTCAGCAGCCCCCGTCAGCCTCCTGCT	48	4.31	1.91	1.65	2.09	(nd) < -2.53	(nd) < -2.53	-2.58	-1.41	(nd) < -2.53	-1.80
6037-R3-2	GCAATTCCCTTTCTCCATCTCCAATTTTCCTC	49	4.15	-1.55	1.15	1.38	(nd) < -2.24	(nd) < -2.24	(nd) < -2.24	(nd) < -2.24	(nd) < -2.24	(nd) < -2.24
6096-R3-1	TGTTTTAATCCTGCCCCCGT	50	8.12	2.89	2.83	1.81	(nd) < -1.12	(nd) < -1.12	(nd) < -1.12	(nd) < -1.12	(nd) < -1.12	(nd) < -1.12
6183-R3-1	GATTCACCTTTTCTTAATGACTTTCCCTCCT	51	> 2.90	nd	nd	> 3.58	nd	nd	nd	nd	nd	> 1.17

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
6192-L3-1	AGATAAAAAACCAACCCACCCAGCAC	52	6.12	1.20	(nd) < -3.75	1.36	(nd) < -3.75	-1.17	-3.42	(nd) < -3.75	(nd) < -3.75	(nd) < -3.75
6233-L3-1	AAAATTAGATTTCCACATTTATCCTTCTCCC	53	> 15.87	nd	nd	> 4.27	nd	nd	nd	> 2.53	nd	nd
6235-R3-1	GCTCCAAAAATCCATTAAATATATTGTCCTT	54	7.85	1.15	-2.40	2.29	-6.59	-2.77	-3.73	-1.08	-12.11	-2.01
6287-L3-2	GCCCCGCCCCACCTTTCGGGGCTCACCTGGC	55	5.41	2.01	1.64	3.04	(nd) < -3.75	-2.07	-2.75	(nd) < -3.75	(nd) < -3.75	-2.87
6409-L3-1	CGTTCCTCAACCGCACGCGCGCCTTCTGGAAC	56	2.22	1.37	2.40	1.28	-2.50	-3.09	-5.39	-2.53	-6.45	-2.76
6434-R3-1	AGCCCTCCACACGACGCTGCAGTGC	57	2.54	-1.65	1.71	2.49	-3.08	-2.94	-6.08	-3.25	-44.90	-7.25
6484-R3-2	CGCTTCGGGATCCTCTCCAACCTGCAACCACA	58	> 6.42	nd	> 2.40	> 4.37	nd	nd	nd	nd	nd	nd
6490-R4-1	CCCATCCCCCATATGACGCTTCCCCCTCCTAACCTCAC CACCCCCAGCA	59	3.28	1.04	1.30	2.00	-4.20	-3.75	-5.00	-2.07	-7.58	-2.49
6496-R3-1	CCCCTCCCCACCCACCACTTCCCCTAGAGTCC	60	14.40	1.47	1.69	4.18	-3.34	-2.65	-2.54	-2.65	-5.20	-1.59
6584-L1-1	TCGGCCCTGCCTCCTCCTCCT	61	8.46	1.93	1.09	2.11	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	(nd) < -2.02	-1.50
6602-R3-2	AGAGCCCCAGTGGAATCTCTCTCTCCAAATCCAT	62	> 4.95	nd	nd	> 1.86	nd	nd	nd	nd	nd	nd
6642-R3-1	CACGTCTCTCCCTCCCTCGAGGTGTACACACA	63	5.93	1.20	(nd) < -5.26	2.84	(nd) < -5.26	-2.73	-3.47	-1.15	(nd) < -5.26	-2.02
6681-R2-1	CCTGTTTTCTCCCCCTCTCTCTCTGCCCCCTCC	64	6.76	2.18	1.13	3.85	(nd) < -6.34	-1.90	-2.74	-1.12	(nd) < -6.34	-1.77
6683-R3-1	AAAATAAACTCTTCTGCTCAAG	65	> 10.55	> 1.57	nd	> 4.34	nd	nd	nd	nd	nd	nd
6752-R1-1	CCCTCCTTTCCCCCACCTCAGT	66	14.74	1.34	(nd) < -1.39	4.34	(nd) < -1.39	(nd) < -1.39	(nd) < -1.39	(nd) < -1.39	(nd) < -1.39	(nd) < -1.39
6795-R4-1	CTTCCCGAGGCCACATGCTTCTTTATATCCCCCATA	67	> 11.94	nd	nd	> 1.62	nd	> 1.60	nd	nd	nd	nd
6803-R3-1	GCTCCCTCTCTGTTGGACCTCACCCAAAGAT	68	19.16	1.39	1.71	4.21	(nd) < -4.36	-1.08	-3.03	1.87	(nd) < -4.36	-1.90

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
6839-L3-1	GCCCGCTGGGCCCTGCCACCCCAACCCCT	69	1.71	1.71	1.97	1.77	(nd) < -6.04	-3.04	-3.58	-2.14	(nd) < -6.04	-2.36
6880-L3-2	ACCTCCCCGCGAAGACATCCACATTCTGCA	70	7.16	1.64	1.21	3.97	-4.57	-2.33	-3.69	1.35	-12.07	-1.50
6906-L3-1	GTGTCCTTCTCCCCAACCCAGCCAGCCTCTCTGG	71	> 7.07	nd	nd	> 2.10	nd	nd	nd	nd	nd	nd
6930-R3-1	ATTAATCCTTCTCTCCCCCTCTG	72	34.08	2.60	2.04	9.78	(nd) < -1.68	(nd) < -1.68	(nd) < -1.68	2.54	(nd) < -1.68	1.50
6984-R4-1	CCCCCTGCCCAAGCATTTGCTTGGGCACCAAGTCCCT GCAA	73	17.80	2.11	2.35	4.74	(nd) < -1.93	(nd) < -1.93	(nd) < -1.93	1.69	(nd) < -1.93	(nd) < -1.93
7026-L3-2	CCTGATCGAAACCTCACCCACCAGATCCGGG	74	4.22	2.58	(nd) < -1.59	1.21	(nd) < -1.59	(nd) < -1.59	-1.35	(nd) < -1.59	(nd) < -1.59	(nd) < -1.59
7061-R3-1	ATGGAACCCCAACCTTCCC	75	3.76	2.16	-1.23	-1.09	-4.11	-2.69	-1.66	-1.31	-9.30	-1.73
7066-R4-1	CAAGGCCCGTAGCCTGAAAAAAGATGCCCCCACCAGC CCTGCC	76	2.80	1.32	1.45	1.90	(nd) < -7.42	-4.34	-4.43	-3.00	(nd) < -7.42	-3.99
7126-L3-1	GCACACCCGCTCTCCGGCCCGCCCGCCCTG	77	4.39	2.59	1.54	1.19	-3.23	-2.33	-3.21	-1.99	-3.87	-2.12
7182-L4-1	TCTGGGTAAGTAGCCGTTTCCGTCACCTTCCCCTGCC	78	8.66	1.60	(nd) < -7.09	3.08	(nd) < -7.09	-2.00	-2.46	-1.11	(nd) < -2.01	-1.82
7192-R4-1	GCAAAGCACTTCCCCCTCTAAGTCTGCCCTGGGCTCTTG GCAC	79	5.65	1.06	1.60	3.31	(nd) < -2.01	(nd) < -2.01	-2.07	1.24	(nd) < -2.01	-1.76
7292-L3-2	TAAATAGCTTCTGAACCTCCCTGCATTCTAATTGC	80	6.77	1.34	1.45	3.96	(nd) < -1.55	(nd) < -1.55	(nd) < -1.55	1.41	(nd) < -1.55	(nd) < -1.55
7352-R3-2	GCCCCCTGCCAGAATCCTCTAACAGCTCTAATTGG	81	2.28	2.91	(nd) < -5.74	1.79	(nd) < -5.74	-2.74	-3.58	-1.97	(nd) < -5.74	-2.31
7356-L2-1	ACCGCGACATAGCCTCGCCCCC	82	5.16	2.76	1.18	2.46	(nd) < -3.81	-1.64	-3.10	-1.59	(nd) < -3.81	-1.82
7356-R2-1	GAAGCTCCGGCGGACGTCCTCCGTTACTCC	83	> 15.13	nd	nd	> 1.62	nd	nd	nd	nd	nd	nd

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
7367-L1-1	AGGGTTAGAGCTGCCCCCTCTGGGACCG	84	2.18	-1.94	-1.30	1.62	(nd) < -3.55	(nd) < -3.55	-3.23	(nd) < -3.55	(nd) < -3.55	-1.80
7384-R3-1	CTCGCAAAGGATCTCCTTCATCCCTCCCCA	85	6.05	1.05	2.67	3.02	(nd) < -2.34	-1.86	(nd) < -2.34	1.20	(nd) < -2.34	-1.32
7411-R3-2	AGTCCCCCTGCCCTCATCTGCCACCCCTAATGAC	86	2.46	2.81	-1.22	1.74	(nd) < -5.03	-2.99	-3.82	-1.84	(nd) < -5.03	-2.12
7421-R2-1	TAAAGAGACTTCCTCCACTGCCAGAGATCT	87	3.81	(nd) < -1.11	(nd) < -1.11	3.90	(nd) < -1.11	(nd) < -1.11	(nd) < -1.11	(nd) < -1.11	(nd) < -1.11	(nd) < -1.11
7426-L3-1	TGGGAGACGAACACCTCCTGCTGTGCTTG	88	> 6.19	nd	nd	> 3.67	nd	nd	nd	> 2.02	nd	nd
7569-L3-1	TCAGGCCACAAAAGCTACCCCCAAGACAGGCC	89	1.43	1.15	-1.03	-1.03	(nd) < -3.88	(nd) < -3.88	(nd)	(nd)	(nd)	(nd)
7571-L1-1	AGGGCTCCCCCACCCTAAG	90	2.27	-1.15	-1.09	1.24	-8.82	-5.67	-5.94	-2.92	-13.36	-3.96
7572-R2-1	ATCACCCCTCCCCCTCCCAAATAAAG	91	11.01	1.19	1.42	4.86	(nd) < -4.50	-2.48	-3.89	-1.79	(nd) < -4.50	-1.63
7578-L3-1	CGCAGTGCACACCCCTGAGCTACAGCCCCCTC	92	-1.03	-1.06	-2.38	-1.25	-14.04	-6.24	-17.14	-2.93	-6.73	-2.44
7660-L2-1	CCCGGCCTCCGCCTGGCCCGAGCGATAA	93	3.94	2.11	2.13	1.41	(nd) < -3.09	(nd) < -3.09	-3.26	-1.11	(nd) < -3.09	-1.77
7702-L2-1	CCCAGAGAACCGGAATTCCTCCCCGCCCCC	94	6.99	2.64	3.18	3.26	(nd) < -3.36	-1.95	-2.34	-1.20	(nd) < -3.36	-1.44
7726-R3-2	CATCCCTCTCCAGAAGAGGAGGAGGAAACA	95	4.39	-1.31	(nd) < -3.13	2.45	(nd) < -3.13	1.32	-1.96	-1.06	(nd) < -3.13	(nd) < -3.13
7764-R3-2	CCCTCTCTGCCTCTCTCATCACCAATAACAGAC	96	9.88	1.35	1.56	3.58	(nd) < -1.65	-1.06	(nd) < -1.65	1.15	(nd) < -1.65	-1.46
8004-R3-2	GGAAGTCTTCTCCTTGCTCCAGTCATTGAAG	97	6.92	1.21	-2.65	3.26	(nd) < -9.22	-3.19	-4.22	(nd) < -9.22	(nd) < -9.22	-2.06

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
8016-L3-1	TCAGCGCAACAAGCCCCGCGAGTCACCCCTCT	98	4.04	2.25	1.06	1.40	(nd) < -4.39	-2.38	-3.04	-1.84	(nd) < -4.39	-2.39
8077-R3-1	CCATTCCCCACCCCTCAGGTAGTAAAAATA	99	4.61	2.89	(nd) < -1.93	1.35	(nd) < -1.93	(nd) < -1.93	-1.28	1.17	(nd) < -1.93	-1.04
8169-L3-1	AACAGAAATGATTATTTACCTCCCCACATG	100	8.55	1.42	1.85	5.76	(nd) < -1.59	(nd) < -1.59	(nd) < -1.59	1.97	(nd) < -1.59	-1.39
8250-R3-1	CAGCCGCCCTCTCCCTCAGCGTTAA	101	7.73	1.48	1.79	2.25	-2.24	-3.20	-5.50	1.35	-1.23	-1.26
8263-R3-1	GATTAAAAACAAGAAATCTATCTTCCCCCAGT	102	> 4.54	nd	nd	> 1.78	nd	nd	nd	nd	nd	nd
8281-L3-1	AGCCCCCTCCCCAGCTGCAGCTGAGGGCTGG	103	2.72	-1.03	1.97	3.10	-3.22	-2.88	-5.10	-2.09	-19.58	-4.66
8316-R3-1	ATCAGGGTATCCTCTCCCCA	104	> 39.96	nd	nd	> 7.56	nd	nd	nd	> 2.43	nd	nd
8394-L3-1	CCCCCGCCCTGCCCATCTCCGACT	105	3.70	2.46	1.62	2.36	-4.53	-3.13	-3.91	-2.65	-14.49	-2.52
8433-L3-1	AAATGGCTCCTTTCCCTTTCCCTCCACCG	106	11.25	1.11	1.90	4.23	(nd) < -2.16	-1.64	-2.20	1.18	(nd) < -2.16	-1.36
8564-L3-1	CCCTTCACCCCAGTTGCCAAACA	107	> 12.45	nd	nd	> 3.14	nd	nd	nd	nd	nd	nd
8587-R2-2	CCCGGCGCCCTCGCCGGCTCCAAACTTTCCCCAA	108	3.02	1.99	1.87	2.01	(nd) < -3.01	(nd) < -3.01	-3.33	(nd) < -3.01	(nd) < -3.01	-1.53
8724-R3-1	GCCAAGCTTGGAACCTCTCCCTGCCAGCATCAC	109	16.37	1.40	1.47	5.34	(nd) < -1.51	(nd) < -1.51	(nd) < -1.51	1.87	(nd) < -1.51	-1.00
8731-R3-1	TCATTGATGCCCCCATCTGCCCAG	110	3.79	3.51	(nd) < -1.34	3.09	(nd) < -1.34	(nd) < -1.34	(nd) < -1.34	(nd) < -1.34	(nd) < -1.34	(nd) < -1.34
8808-R3-1	CCAAAGACCCCTTTCTCCAGCCTGTTTCTGCAA	111	15.38	(nd) < -1.38	1.74	3.85	(nd) < -1.38	(nd) < -1.38	(nd) < -1.38	1.65	(nd) < -1.38	-1.15
8898-R3-1	CGGACGCCGCTCCGCCA	112	4.61	2.99	2.21	2.39	-3.44	-2.32	-3.50	-1.73	-8.22	-2.57
9021-L4-1	AAACAAACACCCAAGCTCCCCACACCATC	113	4.79	1.54	-1.06	1.48	(nd) < -3.02	(nd) < -3.02	-3.04	-1.15	(nd) < -3.02	(nd) < -3.02

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
9053-R3-1	TTC TTG C C C C T C C A A T C C C C G G G C T C C A C C A G C C	114	2.98	1.64	-1.13	1.16	-3.72	-2.19	-3.05	-1.88	-8.27	-2.12
9068-R2-1	C T G C C C T C C C T C T T G A T C A A G A C T G C T C T C C T A A	115	3.42	-1.69	2.79	2.64	-2.57	-3.10	-11.82	-1.57	-40.43	-4.16
9087-L4-1	G G A A A A G A A A C C C T C C C A G T C C A T T C C C T T C C T	116	2.03	-2.72	1.81	1.31	-2.99	-3.69	-4.84	-1.39	-9.51	-2.38
9217-L3-1	A C G A T C C C C G C C G T G A C T A A A G C C A A C A G T G G A	117	3.28	2.48	1.88	1.93	(nd) < -4.55	-2.32	-3.01	-1.46	(nd) < -4.55	-1.91
9245-R2-1	A A C C T C T C A T T A G C C A G C C A C T C G C T C C C A A G	118	5.25	4.01	1.82	1.83	(nd) < -1.69	(nd) < -1.69	(nd) < -1.69	(nd) < -1.69	(nd) < -1.69	(nd) < -1.69
9287-L4-1	G A T A T T C A G A G C C C T C C C A G C C C A C A C A T G C	119	4.16	-1.56	2.12	2.52	-2.91	-2.86	-6.49	-1.43	-32.89	-4.05
9347-L2-1	G C C C A A T A T G C A T T T T A C A T T T T A A C A A A G A	120	> 3.17	nd	nd	> 1.54	nd	nd	nd	nd	nd	nd
9349-R3-1	G T G A T G C A G A G G A C T T C C T G C T C C A G G T C T C	121	22.69	1.06	(nd) < -1.13	7.70	(nd) < -1.13	(nd) < -1.13	(nd) < -1.13	(nd) < -1.13	(nd) < -1.13	(nd) < -1.13
9387-R2-2	T C C A T C C T T G C C G T C G C C T T C A T C T C A A A G C C A T C	122	2.28	1.28	2.25	-1.06	-2.17	-3.48	-4.42	-2.26	-4.76	-1.91
9391-R3-1	C A G C T G C C A G G G A G A C A T A G A A A T T A A A A C A A	123	-1.34	1.31	1.27	-1.69	-1.57	-1.49	-2.45	nd	nd	-1.82
9507-L3-2	G T C T C C C T C A T C C A T C A T C C	124	> 8.07	nd	nd	> 3.35	nd	nd	nd	nd	nd	nd
9564-R1-1	G C C G C C C G C C G G G C A C C G G G C C	125	1.64	1.84	2.11	1.01	-2.74	-3.42	-4.08	-2.21	-7.78	-2.50
9594-R2-1	C T T A G A C T T C C T T C C C A C T C C C T G C A T	126	11.20	1.01	(nd) < -2.11	1.95	(nd) < -2.11	(nd) < -2.11	(nd) < -2.11	2.21	(nd) < -2.11	-1.03
9656-R3-1	G C C C T T A A A G T A C A T A C T G T G G A G A T T A A T G C T	127	> 5.47	nd	nd	> 2.33	nd	nd	nd	nd	nd	nd
9691-L4-1	A A T C A T C C A T T T C A T C C G C A T C T C C C T C T T G G C C C C T T G C	128	5.99	1.18	1.32	1.99	(nd) < -4.24	-1.97	-2.87	-1.57	(nd) < -4.24	-3.39
9733-L3-1	A A G G C T G T C C C T C A C C A G A C T T C C C C A C C C C T	129	7.61	3.56	(nd) < -1.48	2.44	(nd) < -1.48	(nd) < -1.48	1.07	1.69	(nd) < -1.48	1.33
9774-R2-2	C C G C C C C C T C A C C G C C T C C T G C T C C C A T C A G G C	130	3.72	1.83	2.45	2.54	-2.98	-2.83	-4.46	-1.84	-12.28	-2.00
9816-R2-1	C T G G C C C T T T A A G A G C C T C T C C G C G C T G C C G	131	2.42	2.46	2.03	2.04	-2.71	-2.17	-2.86	-1.67	-4.18	-2.08
9840-L3-2	T T C A G G T T T T T A T A A A T C A G G A T G T C A A C A A A T	132	-1.39	1.32	1.13	-2.15	-1.71	-1.66	-2.89	nd	nd	-1.70

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
10010-R2- 2	CCCGGCGCCCTCGCCGGCTCCAAACTTTCCCCAA	133	4.99	2.90	2.53	3.23	(nd) < -1.75	(nd) < -1.75	-1.85	(nd) < -1.75	(nd) < -1.75	-1.31
10030-R3- 1	AACTCAGCTGCCTTCGGCC	134	> 6.94	> 1.63	> 3.55	> 2.08	nd	nd	nd	nd	nd	nd
10138-L2- 1	AGCCTGCTCCGCTCTCCCTCCACCAGAAAAAGT	135	10.06	2.02	1.51	3.69	-3.13	-2.11	-3.65	1.01	-7.30	-1.76
10145-L2- 1	CTTTGCTCTCTCTTCTGTTATCTGGACC	136	> 5.20	nd	nd	> 1.95	nd	nd	nd	nd	nd	nd
10175-L1- 1	CCTGCTCCTAGCAACCAGGAGCCACAA	137	> 7.31	nd	nd	> 4.44	nd	nd	nd	nd	nd	nd
10209-L3- 1	AGAAAAATAAGTTAGCTATGTAAACAAATTGA	138	-1.56	1.29	-1.09	-2.17	-2.06	-1.65	-3.13	nd	nd	-1.86
10231-L3- 1	GTGCAGCAGCCCGCGCCAGCCTCCGCGCCGCC	139	1.76	1.50	3.56	-1.02	-1.97	-3.47	-5.87	-6.71	-3.38	-2.65
10231-R3- 1	TGAACTTTAGCTGGCGCGCGCCTGTCAGC	140	1.15	1.48	2.17	-1.45	-2.34	-4.21	-5.32	-3.26	-4.14	-2.20
10242-R3- 1	GGAAGAAAGCCCTTCGCTTCCACCCCCGAACAC	141	5.35	2.89	1.21	3.14	-2.60	-1.56	-3.46	-1.09	-6.84	-2.88
10333-L3- 1	TGTGCCCTGCCACCCCTCCCTCCCTGCCCCG	142	5.80	1.70	1.74	2.93	-3.43	-2.75	-4.34	-2.25	-9.80	-2.32
10335-L3- 1	CACTCCCCTCCTTTTAAATTAGAAAGCACTAAGA	143	10.97	1.56	2.36	6.06	(nd) < -2.08	(nd) < -2.08	(nd) < -2.08	(nd) < -2.08	(nd) < -2.08	-1.22
10342-R2- 2	CCCGCGCCGGAGCATCTCGAAGTTAATTAATA	144	1.75	2.08	2.13	-1.05	-2.55	-2.86	-2.31	-2.36	-10.37	-1.85
10366-R3- 2	AAACACCACCCACGCTCTTGCTACAAGACCCACAT	145	2.46	1.21	-1.12	-1.11	(nd) < -3.19	-2.45	-2.94	(nd) < -3.19	(nd) < -3.19	(nd) < -3.19
10374-R3- 2	GACACCGCCCGCTACTTTGTTAATGAAAAGCCCCC	146	2.59	2.62	2.48	1.28	-2.47	-2.10	-4.29	nd	nd	-1.97
10533-R3- 1	GCCCCCTCTTTTATATTGCCAAGA	147	4.36	(nd)	(nd)	2.33	(nd)	(nd)	(nd)	(nd)	(nd)	(nd)

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
				< -1.58	< -1.58		< -1.58	< -1.58	< -1.58	< -1.58	< -1.58	< -1.58
11370-L4-1	CCCTCGCCCCACACACTGCATCCTTGCCCCAGTTTGGCTG CCATCAGTATTGTCCCTGAGAACTGGAC	148	7.22	1.56	(nd) < -6.13	2.49	-2.48	-2.20	-6.13	-1.28	-3.12	-1.10
12184-L4-1	GACCTCAGCGTGCCCCCTTTCAACCCACAGACGAATATT GTGTACAA	149	2.50	1.10	-1.39	1.69	(nd) < -4.07	-2.20	-3.80	-1.92	(nd) < -4.07	-1.97
12223-L4-1	CCCAGAAAGACATCAGACAGAGTTGTTTCTTCTCCCTCTA	150	1.99	1.06	(nd) < -4.89	1.27	(nd) < -4.89	1.25	-1.49	1.03	-2.26	-3.14
4315_C-L4-1	GCAGCCCCCTCCTCCGAGAGGTTGGGGTGC GGCCG CCCGGCCCTCCCGTCCCTCCCTCC	151	-2.79	1.91	1.97	2.72	-3.38	-2.74	-3.87	-1.53	-9.33	-7.93
4315_D-R4-1	GGAAAGTCAGCCCCCAGCGCCCCCGGAGTTCTTGG	152	4.01	1.41	1.66	2.02	-4.14	-2.64	-7.06	-2.32	-9.62	-2.60
4315_E-R4-1	CCCCCACCAAACTATTCCCGCATCCTCCCCGGCTCTG G	153	7.07	1.72	1.30	2.47	-4.50	-2.77	-5.04	-2.24	-10.19	-2.01
4315_F-R4-1	AACCCGGGCTCCCCACCCGCTCCCTGAGC	154	3.58	1.70	1.23	1.44	-5.04	-3.19	-3.15	-2.24	-20.64	-2.42
4315_I-L4-1	ACACCTCTGCGCCCCCTCAGGCGCCCTGGGCCTCGGCG CCCCGCCGTCCAG	155	5.01	2.26	2.05	2.48	-3.25	-2.26	-2.87	-1.61	-6.36	-2.28
4315_K-L4-1	TCCCAGGGGGCCCTGAACTTGTCAAATCCTCGCCATCC TCCACCCCGAGCCCCG	156	4.24	1.87	2.15	1.72	-3.72	-2.41	-4.85	-1.91	-6.99	-1.83
10010_B-L4-1	GCCGAGCCCCCGCCCCCGGGGATGCTGCCCTCCG GAAGGAGGGGCGCTGCC	157	2.68	2.95	1.31	1.80	-4.25	-3.10	-3.43	-2.54	-5.76	-2.39
10010_D-L4-1	TGGCGCCCTCCCCCGCGGGGCTCAGCCTCTCACCT G	158	4.06	1.62	1.93	2.59	-3.56	-2.99	-4.26	-1.54	-15.24	-2.37
7356_A-R4-1	CAGAGCCCGCTCTCGCGACCGACCTGCCGCCGACCCG CACAG	159	1.71	1.65	3.03	-1.13	-1.99	-3.31	-4.25	-3.28	-3.88	-2.82
12722-L4-1	AATACGGACAAGCCCCACTCCCTCATTAGCATAAAAA CAAAGTACTTCCGACCTCCCCGCCGCCCG	160	3.15	2.05	2.31	2.04	-3.42	-2.39	-3.53	-2.00	-5.91	-2.63
99999-R4-1	CCCCTTGTCACCCCCAGCCCCCTTCTCTGGCCAGGACCC CAGCGAGGCCCCAGAGAA	161	10.71	2.08	1.97	3.33	(nd) < -2.88	-1.64	-2.59	2.58	(nd) < -2.88	-1.53

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
99997-R4-1	TCCTCACTGGGCCCCACCAAAACTGTGCCACCCCTCAAGCCCCAGGAGCTTCCTTAAC	162	6.84	-1.42	1.59	2.77	-3.46	-2.91	-4.06	-1.00	-5.37	-1.68
8433_B-L4-1	CATATTTTGTGTTGCTGAGTATTTGGGGTTTGCTCGCCCGT	163	> 5.03	> 3.56	> 2.45	> 2.44	nd	nd	nd	nd	nd	nd
8433_C-R4-1	AAACCAAAAAAAAAAATTAAAAAGCGACGAAAAATGCAATTGTGTGCCCTTCTCCCTCC	164	24.29	1.73	1.87	5.70	(nd) < -1.45	(nd) < -1.45	(nd) < -1.45	1.95	(nd) < -1.45	1.13
8433_D-R4-1	CCCGAGCCCCGGCGCCCTGTGTTGTGCTCCGCTCTCCGGAAATGCCATCACTAAT	165	8.52	2.31	1.63	2.00	(nd) < -4.10	-1.76	-3.76	-1.33	(nd) < -4.10	-2.68
let-7a	AACTATACAACCTACTACCTCA	166	-5.65	-1.33	1.33	-2.81	1.20	-1.63	-1.75	-1.18	1.20	-3.59
let-7b	AACCACACAACCTACTACCTCA	167	-2.30	-1.30	1.41	-2.86	1.19	-1.79	-2.26	1.05	1.51	-4.67
let-7c	AACCATACAACCTACTACCTCA	168	-2.52	-1.34	1.39	-2.93	1.22	-1.77	-1.97	1.17	1.38	-3.51
let-7d	AACTATGCAACCTACTACCTCT	169	-2.41	-1.35	1.23	-2.86	-1.03	-1.69	-2.05	1.31	1.64	-3.27
let-7e	AACTATACAACCTCCTACCTCA	170	-2.38	-1.32	-1.20	-2.76	-1.49	-1.83	-5.03	1.11	1.34	-8.91
let-7f	AACTATACAATCTACTACCTCA	171	-3.05	-1.28	1.19	-3.04	1.01	-1.73	-4.92	-1.20	1.06	-3.85
let-7g	AACTGTACAAACTACTACCTCA	172	-2.34	-1.12	1.13	-2.73	-3.04	-1.16	-1.34	1.28	2.07	-3.06
let-7i	AACAGCACAAACTACTACCTCA	173	-1.56	1.86	(nd) < -3.56	-1.68	1.05	2.20	-3.56	1.63	3.60	-1.90
miR-100	CACAAGTTCCGGATCTACGGGTT	174	-1.91	-1.56	1.24	-3.21	-1.72	-2.50	-2.63	1.39	2.17	-5.22
miR-1224-5p	CCACCTCCCGAGTCCTCAC	175	5.56	1.51	1.89	1.57	(nd) < -1.14	(nd) < -1.14	-1.11	(nd) < -1.14	(nd) < -1.14	(nd) < -1.14
miR-1225-5p	CCCCCCTACTGGCCCGTACCCAC	176	8.21	-1.18	2.95	4.25	(nd)	(nd)	-2.35	(nd)	(nd)	(nd)
miR-1228*	CACACACCTGCCCCCGCCAC	177	3.27	3.43	1.06	1.96	-6.69	-3.06	-3.39	-1.87	-12.52	-2.10
miR-125a-5p	TCACAGGTTAAAGGGTCTCAGGGA	178	(nd) < -10.19	-1.52	1.28	-3.29	-2.87	-1.81	-6.80	-1.99	-1.10	-6.52
miR-125b	TCACAAAGTTAGGGTCTCAGGGA	179	-3.83	-1.52	1.33	-3.15	-1.08	-1.77	-2.46	-1.78	1.01	-5.55

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
miR-126	CGCATTATTACTCACGGTACGA	180	-2.39	-1.40	5.58	-2.14	1.33	-1.45	-1.81	1.07	1.71	-1.42
miR-135a*	CGCCACGGCTCCAATCCCTATA	181	> 3.61	nd	nd	2.49	nd	nd	nd	nd	nd	nd
miR-142-3p	TCCATAAAGTAGGAAACACTACA	182	-2.17	1.15	1.09	-2.54	-4.28	1.14	1.45	-1.22	1.18	-3.10
miR-145	AGGGATTCCCTGGGAAAACCTGGAC	183	-4.94	-1.08	2.41	-2.67	1.13	1.01	-1.07	-2.03	1.31	-3.13
miR-146b-5p	AGCCTATGGAATTCAGTTCTCA	184	(nd) < -1.89	1.12	2.44	< -1.89	< -1.89	2.71	1.07	1.30	2.48	-1.21
miR-149*	GCACAGCCCCCGTCCCTCCCT	185	3.46	3.38	-1.02	2.16	-8.57	-3.25	-6.43	-2.34	-21.89	-2.17
miR-150*	CTGTCCCCCAGGCCTGTACCAG	186	> 4.16	nd	nd	> 1.76	nd	nd	nd	nd	nd	nd
miR-155	ACCCCTATCACGATTAGCATTA	187	nd	nd	nd	nd	nd	> 1.83	> 1.07	nd	nd	nd
miR-16	CGCCAATATTACGTGCTGCTA	188	-4.17	-1.04	(nd) < -10.19	-3.62	-1.53	-1.05	-1.04	1.73	2.03	-3.14
miR-181c	ACTCACCGACAGGTTGAATGT	189	(nd) < -3.23	-1.73	1.58	-2.37	-1.79	1.13	-2.25	-1.16	1.07	-3.23
miR-198	GAACCTATCTCCCCCTCTGGACC	190	> 33.49	> 2.21	nd	> 8.74	nd	nd	nd	nd	nd	nd
miR-199a-3p	TAACCAATGTGCAGACTACTGT	191	-3.52	1.40	1.43	-1.94	1.72	1.56	1.33	1.59	2.97	-4.20
miR-19b	TCAGTTTTGCATGGATTTCACA	192	-1.07	-1.01	1.41	-1.94	(nd) < -2.18	-1.10	1.48	1.28	4.82	(nd) < -2.18
miR-200b	TCATCATTACCAGGCAGTATTA	193	1.18	1.01	(nd) < -5.45	1.43	-1.58	-1.56	-1.38	1.72	2.60	-1.45
miR-200c	TCCATCATTAACCCGGCAGTATTA	194	1.45	1.11	-1.19	1.72	-1.88	-1.84	-1.47	1.92	2.85	-1.10
miR-205	CAGACTCCGGTGAATGAAGGA	195	> 11.24	> 24.11	nd	nd	nd	nd	nd	> 3.48	> 9.48	nd
miR-21	TCAACATCAGTCTGATAAGCTA	196	-2.52	1.77	-1.51	1.45	1.41	3.51	2.27	3.43	5.38	1.25

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
miR-23a	GGAAATCCCTGGCAATGTGAT	197	1.38	2.90	(nd) < -4.40	1.37	(nd) < -4.4	2.28	(nd) < -4.4	2.38	7.76	-2.09
miR-23a*	AAATCCCATCCCCAGGAACCCC	198	> 4.54	nd	nd	> 1.79	nd	nd	nd	nd	nd	nd
miR-23b	GGTAATCCCTGGCAATGTGAT	199	-1.43	1.45	(nd) < -9.07	-1.34	1.26	1.24	-2.02	1.27	4.35	-1.98
miR-24	CTGTTCTGCTGAAGCTGAGCCA	200	-2.08	1.26	2.49	-1.46	1.08	1.22	-1.17	1.09	2.86	-2.76
miR-25*	CAATTGCCCAAGTCTCCGCCT	201	> 4.17	nd	nd	> 1.38	nd	nd	nd	nd	nd	nd
miR-26a	AGCCTATCCTGGATTACTTGAA	202	-2.98	-1.73	2.04	-3.66	1.04	1.00	-1.33	1.07	1.67	-2.29
miR-26b	ACCTATCCTGAATTACTTGAA	203	-2.84	-1.67	1.20	-3.56	-1.49	-1.16	-1.18	1.13	1.74	-2.23
miR-27a	GCGGAACCTTAGCCACTGTGAA	204	-1.27	1.78	2.19	-1.78	-1.02	1.44	1.07	1.73	4.25	-2.26
miR-27b	GCAGAACTTAGCCACTGTGAA	205	-1.11	2.20	2.52	-1.91	1.18	1.64	1.24	2.05	6.50	-1.85
miR-298	TGGGAGAACCTCCCTGCTTCTGCT	206	24.71	2.84	(nd) < -1.29	10.70	(nd) < -1.29	(nd) < -1.29	-1.15	2.09	(nd) < -1.29	1.13
miR-29a	TAAACGATTTTCAGATGGTGCTA	207	-2.68	1.00	3.18	1.50	-1.90	2.13	1.41	1.23	2.18	-2.96
miR-29b	AACACTGATTTCAAATGGTGCTA	208	-5.79	-1.32	1.33	-1.17	-3.06	1.46	1.26	1.01	1.30	-5.30
miR-29c*	TAAACGATTTCAAATGGTGCTA	209	-4.12	-1.00	3.14	1.30	-3.52	2.01	1.49	1.66	2.61	-4.26
miR-30a	CTTCCAGTCGAGGATGTTTACA	210	-4.44	-3.14	3.50	-3.84	-1.50	-2.18	-3.97	-7.98	2.09	1.23
miR-30b	AGCTGAGTGTAGGATGTTTACA	211	(nd) < -6.56	-1.98	2.09	-4.38	-2.50	-1.73	-2.27	2.12	2.77	1.03
miR-30b*	GAAGTAAACATCCACCTCCCAG	212	3.38	-1.21	1.05	-1.36	-2.31	-2.53	-2.76	-1.56	-3.51	-3.51
miR-30c	GCTGAGAGTGTAGGATGTTTACA	213	-4.45	-2.14	(nd) < -6.05	(nd) < -6.05	(nd) < -6.05	-1.95	(nd) < -6.05	1.86	2.21	-2.40
miR-30c-1*	GGAGTAAACAACCCCTCTCCCAG	214	39.98	(nd) < -1.58	1.57	7.84	(nd) < -1.58	(nd) < -1.58	-1.42	2.67	(nd) < -1.58	(nd) < -1.58
miR-30c-2	GCTGAGAGTGTAGGATGTTTACA	215	-4.45	-2.14	(nd)	(nd)	(nd)	-1.95	(nd)	1.86	2.21	-2.40

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
					<-6.05	<-6.05	<-6.05		<-6.05			
miR-30d	CTTCCAGTCGGGGATGTTTACA	216	-6.68	-2.87	3.73	-3.55	-1.26	-2.03	-4.02	1.32	3.66	1.39
miR-30e	CTTCCAGTCAAGGATGTTTACA	217	(nd) < -4.99	-2.42	2.08	(nd) < -4.99	(nd) < -4.99	-2.00	(nd) < -4.99	1.45	1.71	-1.30
miR-320a	TCGCCCTCTCAACCCAGCTTTT	218	4.68	1.05	2.16	2.43	1.02	1.33	-1.39	nd	nd	-1.20
miR-331-3p	TTCTAGGATAGGCCAGGGGC	219	(nd) < -1.21	(nd) < -1.21	2.17	1.42	1.24	1.47	1.27	-1.21	-1.21	1.36
miR-371-5p	AGTGCCCCCACAGTTTGAGT	220	> 2.65	nd	nd	> 1.30	nd	nd	nd	nd	nd	nd
miR-373*	GGAAAGCGCCCCCATTTTGAGT	221	10.21	1.72	2.16	3.92	(nd) < -3.77	-1.66	-3.39	1.66	(nd) < -3.77	-1.73
miR-375	TCACGCGAGCCGAACGAACAAA	222	nd	nd	nd	nd	nd	nd	nd	nd	nd	> 2.88
miR-423-5p	AAAGTCTCGCTCTCTGCCCCCTCA	223	6.96	1.49	1.63	1.87	(nd) < -6.09	-1.38	-1.95	1.01	-1.06	-1.83
miR-424	TTCAAAACATGAATTGCTGCTG	224	nd	nd	nd	nd	nd	> 2.14	nd	> 2.99	> 11.35	nd
miR-483-5p	CTCCCTTCTTTCTCCTCCCGTCTT	225	24.38	1.51	1.68	5.17	(nd) < -1.54	1.11	-1.64	3.19	(nd) < -1.54	-1.03
miR-486-3p	ATCCTGTACTGAGCTGCCCCCG	226	3.44	2.92	2.61	1.43	(nd) < -1.91	(nd) < -1.91	-1.85	1.05	(nd) < -1.91	(nd) < -1.91
miR-491-3p	GTAGAAGGGAATCTTGATAAG	227	-2.17	1.13	4.21	-2.65	(nd) < -3.87	-1.40	-3.03	nd	nd	-3.87
miR-491-5p	CCTCATGAAGGGTTCCCCACT	228	> 2.09	nd	nd	nd	nd	nd	nd	nd	nd	nd
miR-513a-5p	ATGACACCTCCCTGTGAA	229	> 6.38	nd	nd	> 3.28	nd	nd	nd	nd	nd	nd

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
miR-513b	ATAAATGACACCTCCTTGTGAA	230	> 1.98	nd	nd	> 1.29	nd	nd	nd	nd	nd	nd
miR-516a-5p	GAAAGTGCTTCTTTCTCCTCGAGAA	231	> 23.45	nd	nd	> 5.80	nd	> 1.65	nd	> 2.27	nd	nd
miR-550	GGGCTCTTACTCCCTCAGGCACT	232	> 2.04	nd	nd	nd	nd	nd	nd	nd	nd	nd
miR-557	AGACAAGGGCCACCCGTGCAAAAC	233	4.19	1.38	2.01	1.78	(nd) < -1.39	(nd) < -1.39	-1.15	1.57	(nd) < -1.39	-1.21
miR-575	GCTCCTGTCCAACCTGGCTC	234	> 4.55	nd	nd	nd	nd	nd	nd	nd	nd	nd
miR-612	AAGGAGCTCAGAAGCCCTGCCCAGC	235	5.20	2.50	-1.18	1.81	(nd) < -5.94	-2.60	-2.93	-1.98	(nd) < -5.94	-2.85
miR-614	CCACCTGGCAAGAACAGGCGTTC	236	> 2.61	nd	nd	nd	nd	nd	nd	nd	nd	nd
miR-630	ACCTTCCCTGGTACAGAACTACT	237	> 4.81	nd	nd	> 1.88	nd	nd	nd	nd	nd	nd
miR-637	ACGCAGAGCCCCGAAAGCCCCCAGT	238	1.35	5.51	2.82	7.31	(nd) < -1.81	(nd) < -1.81	(nd) < -1.81	1.46	(nd) < -1.81	-1.52
miR-638	AGGCCGCCACCCGCCCGCATCCCT	239	2.50	2.06	3.13	1.32	-1.55	-3.45	-4.86	-2.17	-2.60	-2.40
miR-658	ACCAACGGACCTACTTCCCTCCGCC	240	3.65	3.18	1.67	3.41	(nd) < -3.56	-1.70	-1.93	1.04	(nd) < -3.56	-1.76
miR-663	GCGGTCCCCGCGCGCCCCCGCCT	241	3.51	2.57	2.02	1.44	(nd) < -3.06	(nd) < -3.06	(nd) < -3.06	(nd) < -3.06	(nd) < -3.06	(nd) < -3.06
miR-671-5p	CTCCAGCCCCCTCCAGGGCTTCCT	242	> 7.67	nd	nd	> 5.82	nd	nd	nd	> 1.94	nd	nd
miR-675	CACTGTGGGCCCTCTCCGCACCA	243	7.25	4.60	(nd) < -1.19	4.46	(nd) < -1.19	(nd) < -1.19	-1.07	1.88	(nd) < -1.19	(nd) < -1.19
miR-708	CCCAGCTAGATTGTAAGCTCCTT	244	nd	nd	nd	nd	nd	nd	< -1.01	nd	nd	> 2.16
miR-744	TGCTGTTAGCCCCTAGCCCCCGCA	245	4.02	1.97	1.40	1.40	(nd) < -3.47	-2.20	-3.66	-1.68	(nd) < -3.47	(nd) < -3.47
miR-765	CATCACCTTCCCTTCTCCTCCA	246	25.46	3.15	3.08	6.06	(nd)	3.70	1.82	3.00	(nd)	1.02

Table 1

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in primary tumors vs. normal Lung									
			Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
							< -1.58				< -1.58	
miR-920	TACTGCTTCCACAGCTCCCC	247	> 3.57	nd	nd	nd	nd	nd	nd	nd	nd	nd
miR-923	AGTTTCCTTTCTCCGCTGAC	248	6.70	1.19	1.24	1.35	-2.18	1.11	-1.43	1.19	2.00	-2.79
miR-92a-2*	GTAATGCAACAATAATCCCCACCC	249	3.79	2.45	1.23	1.20	(nd) < -1.93	(nd) < -1.93	(nd) < -1.93	1.18	(nd) < -1.93	-1.59
miR-92b*	CACTGCACCGCGTCCCGTCCCT	250	5.09	1.52	1.50	1.81	(nd) < -3.06	(nd) < -3.06	-3.14	(nd) < -3.06	(nd) < -3.06	(nd) < -3.06
miR-93	CTACCTGCACGAACAGCACTTTG	251	(nd) < -2.50	1.27	(nd) < -2.50	(nd) < -2.50	(nd) < -2.50	1.17	-1.22	1.14	2.52	1.03
miR-98	AACAATACAACCTTACTACCTCA	252	-4.47	-1.37	1.17	-2.84	-4.10	-1.83	-2.38	-1.22	1.21	-4.82
miR-99b	CGCAAGGTCGGTTCTACGGGTG	253	(nd) < -3.33	-1.36	1.59	-2.74	(nd) < -3.33	-2.23	(nd) < -3.33	1.30	-1.51	(nd) < -3.33

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
3717-L2-1	CCGCCCTCCCCATAGCCTCACCCCCAAACCCACTCACA	6	1.02	1.86	1.77	2.27
3758-R2-2	TGCAGGCTCCACTGACATTTTCACAAATTTAAATCAT	254	-1.85	2.07	1.89	2.26
3799-R3-1	CCAGAGGCCCCCCCGCCGCC	7	-1.63	1.14	1.33	(nd) <6,30
3820-R3-1	CCCCACCCCTCTGTGGGGCCATCCCTG	255	1.60	nd	nd	nd
3851-R3-4	GAGCTCCCAACCCCTCCTTTATGTTTTGTCTAAAGC	256	nd	nd	6.82	nd
3874-L3-1	TGAATATTATCCCTAATAACCTGCCACCCCA	257	2.75	10.71	10.14	10.68
3897-R3-1	CAGCCGCCTCCCCCTCAGCGTTAA	10	-1.97	1.50	1.79	1.75
3906-L3-1	AACATATGTAAACCCCTTTATTCCTCATTCTG	258	(nd) <4,59	1.25	1.70	2.14
3923-R3-1	GCCTCTCACAAGGATCTCCTTCATCCCTCTCC	11	-1.74	2.11	1.72	2.03
3952-L3-2	GCGCACAGAGCACTCAATCTGACACCCCTCGC	259	-1.66	1.90	1.73	2.13
3953-R3-2	ACTCCAGCCTCCGGCCCTCAGCTTCCCGAGC	12	-1.22	1.78	1.59	1.81
3976-L2-2	TCATACTCCTGCTTGCTGATCCACATCTGCTGGAA	260	1.58	5.83	7.55	8.31
3995-L2-2	CTATAAAACTTCGAAAAGTCCCTCCTCCTCAGCT	13	-1.17	1.41	1.31	1.36
4037-R3-2	GCCTGTCCCTGGCATGTACTGTAAATTTATCT	15	-2.05	2.26	1.66	1.88
4064-R3-1	CATAGCTGAATTCATCCAGCCCCAG	261	2.32	nd	nd	nd
4118-L2-2	TCATACTCCTGCTTGCTGATCCACATCTGCTGGAA	262	1.80	6.34	8.91	9.64
4130-L3-1	GCCAGCACGCCGTCCATGTCCACCAGCACCC	263	-1.51	1.55	1.91	(nd) <4,64
4143-R3-1	TCAGCGTCTTGCTCTCCTCCTGGTA	16	2.11	3.64	nd	nd
4155-R1-1	AGACCGGACTCGCCTCTTCCAACTCGAGTTCA	264	nd	3.56	nd	nd
4182-R2-2	AGTTCAGCAGCCCCAGTGGACATGCTGGGGTGGT	265	9.76	14.90	nd	nd
4203-R3-2	GCACATTCCCACTTCCCCAGAGGCAGGCTCCATAT	17	nd	5.60	7.52	nd

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
4205-R3-1	ACTTCATCCTCACCACCTCACACCACCCTAG	266	-1.94	1.83	2.35	(nd) <3,98
4216-R3-1	TCCCTCCCTTATACACAGATCAATTCCCCC	267	-1.53	1.78	1.86	2.51
4303-R1-1	AGTGCCCGCTCCTCCGACCTCCCTGCGCACC	19	-1.09	1.56	1.42	1.57
4315-R3-2	TCCCGGCCCTCTCCATTCTCGGCTCCGGAGCA	20	-1.54	1.63	1.53	1.61
4340-R3-1	ATTTTCCAGCCCCCTTGTCCTCCAGGCCAAAC	268	-1.75	1.98	1.79	2.09
4361-R3-1	CGTCTCCCTCCCTCATGTGC	21	-1.60	1.46	1.56	(nd) <4,18
4391-R2-1	GCCAAATTCTCAACCAATATAACTCTGTGA	269	2.65	7.30	10.84	nd
4413-L3-1	GGCACCTCCAGCTACAGTAAACAAAT	270	-1.65	1.74	1.74	1.99
4417-R1-1	GCTCATCAAAAAGTTCCCTGT	271	-1.75	1.40	1.49	1.66
4440-R3-1	TACTCCCGCCGTTTACCCGCAATTTCACTGAA	272	-1.71	1.75	1.76	2.23
4440-L3-1	TTCAGAGCACTGGGCAGAAATCACATCAC	273	-1.70	1.70	1.81	2.20
4448-R3-1	CCTACCCCCAGCATCTCCTCACGCCATTGCC	24	-1.76	2.12	1.76	2.36
4479-R3-1	AGCCCCCTGCCCGGAAATTCAAAACAACCTGC	25	-1.47	1.10	1.38	1.60
4498-L3-2	GAGATCCAGACGGCCGTGCGCCTGCTGCTGCCT	274	-1.95	2.00	1.73	2.11
4567-L1-1	ATCTGCCCAGTTCCAGCACACTCCC	275	2.50	6.95	8.90	nd
4579-L3-2	GGCTTCACTTGCCCTCCTGCAAAACACCAATAGC	276	-1.75	1.60	1.64	1.82
4593-R3-1	AGCAGATGACATAACTCCCCCGGCATCAG	26	-1.83	-1.24	1.68	1.74
4610-R3-1	GCCCTCTGGCCCCCTGCCCTAATTGGCTGC	277	1.04	1.55	1.43	1.66
4724-L3-2	CTTGGCATCTCTAGCACCTTCAGCTTTCTGTGCCT	278	-1.60	1.82	1.71	1.91
4754-R3-2	CCAAAGCCTTAGACAAGTGCCAAAGCCCATCTTT	279	-1.58	1.60	2.35	2.49
4801-L3-1	AACCTCTGCCCTCCTGTTTGCTACAAAAACATTAAT	280	-1.47	1.76	1.15	1.36
4829-R2-1	TCCCTTTGTGCTGCCCGAGTGCCCTTCCCCCTG	29	-1.59	1.81	1.62	1.92

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
4855-R3-1	GGGCTGCCGGGTCTCCCGCTTCC	30	2.95	3.52	3.97	nd
4964-L3-2	AGGGCTAACTTCTGAAAAACCCACCAAATTCCCCAA	281	-2.00	1.44	1.52	1.79
5006-L3-1	ACAGCTCCCTCTGCTGGCTCC	33	-2.37	1.57	1.68	1.78
5071-R2-1	CCCCAGTCCCAGCCCCCAATTAATAATGGG	282	-1.48	1.42	2.11	2.49
5080-R3-1	CTTGCAAAGGGTCTCCTTCATCCCTCTCCA	34	-1.69	2.01	1.67	1.96
5192-L3-1	CCCCCTTCCTCTCTATATCAGCAAAGAC	283	-1.45	1.75	1.73	2.15
5306-L3-2	CCTCTGACCCCCAGCTCTGGCCCTTTCTAGGG	284	-1.72	1.66	1.75	2.05
5327-L3-1	CTCAACCTCTGGGACTATGTCCTGTCCTCC	285	nd	6.36	8.44	nd
5342-L3-1	CACCACCAAAACCAATGCCGCTGCTCTCCTTCCA	36	2.47	nd	nd	nd
5372-R3-2	CCCTCTGTTTTCAACTTACACAAGATTCTTTTT	286	-2.07	2.01	1.61	2.10
5380-R2-2	GGCTCCCAGATGTGTCCCACATTGAAGAATTATC	287	-2.19	1.91	1.82	1.88
5441-L3-2	GCCAAGCTCCAAGTCAGTATAGCTAACAGAGCAG	288	nd	3.14	nd	nd
5474-L3-2	CTCCTCTCACCTGGCCAGACTCTGACCCACCTAC	289	2.77	4.11	7.78	8.58
5513-L3-1	CAACTGTTCTCCATGATGCCCTCAGAGCCCACCT	290	-2.43	2.15	1.58	1.91
5554-R2-1	CCCCACCCCTCATCAGCTGCTCCCAGAT	38	-1.67	1.88	1.71	2.23
5598-R2-2	CTCCCACCTCCGTGAAGCTATTTTTAACTGTGCA	291	-1.27	1.56	1.62	1.72
5618-R3-1	TCCCAGCCCCACCCAGTGCCACATTACAGCCCCA	292	-1.39	1.41	1.93	1.95
5619-L3-1	AATGCCAGTTCTGCTCAATCTTCCCTCAATGAG	293	-2.45	1.94	1.69	2.04
5638-R2-1	GGCCCTCCCCCTGCCCTGTGATAGGCTGCTTG	39	-1.06	1.87	1.63	1.59
5640-L3-1	GCCATGGAACACCGTGCCTGCCCTCTCGAGA	40	-1.37	1.75	1.80	1.84
5733-R3-2	CTCACCCAGCTCATCCTGCTTCTCAGTCCCAC	294	-1.58	1.47	1.34	1.76
5735-L3-1	AGAAAGTTGCTGTTTTCTCTGTGGCCTCAAGCCT	295	-1.64	1.94	1.78	2.10

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
5782-L3-1	GATTCCAGCCCCCTTCCCCC	42	3.41	5.68	7.06	9.69
5795-R1-1	CTGCCCTCCAAGAAATAAAATTACCCGCAATTACT	43	-1.04	2.08	1.83	2.32
5836-R3-2	CATTAAACCCCATATTATCACAGCACGCCCCCATTC	44	-1.60	1.58	2.26	2.56
5854-R3-1	CCCTCCCTCTCCGAAAGAATGTGCAC	45	-1.42	2.05	1.96	2.24
5863-L3-1	GCCGTTGCTGCTGGCAATTCCTGTGG	296	-2.05	2.24	1.56	2.05
5919-L3-1	AAGCAACACTGTCACITTTATCTCCCTAGA	297	-1.92	1.46	1.50	2.28
5971-R3-1	CTGCTCAGCCTCCCACATCTGT	46	-1.53	1.49	2.17	2.31
6008-R1-1	ACAATACCCCCACCTTTTTTCCCTGTACCTTAC	47	2.16	4.91	7.51	8.14
6016-R2-1	AAACTCCAGCAGCCCCGTCAGCCTCCTGCT	48	-1.45	1.75	1.66	1.86
6026-R3-1	CAGCCACCTTGTTTTGTGGTTTGCAAA	298	nd	5.34	7.28	7.92
6192-L3-1	AGATAAAAAACCCACCCACCCAGCAC	52	nd	4.46	nd	nd
6218-R3-1	AATCACATTACTGCCCTCTCATGTGACA	299	4.96	nd	nd	nd
6235-R3-1	GCTCCAAAAATCCATTTAATATATTGTCCCTT	54	-1.96	2.79	-1.84	(nd) <12,83
6253-L3-1	CCTGACAAATATCCTGGCTGCCATAATGCCAGC	300	-2.63	1.74	1.98	(nd) <6,02
6287-L3-2	GCCCCGCCCCACCTTTTCGGGGCTCACCTGGC	55	-2.16	1.72	1.47	1.68
6355-R3-1	TCCAGATCATCTGTTCCTTGAGGATTTACAGT	301	1.94	5.81	nd	nd
6409-L3-1	CGTTCCCAACCGCACGCGCCGCCCTTCTGGAAC	56	-1.32	2.14	1.60	2.05
6421-R3-2	CCCTCCTGTGAGAGTCTGAAGGACACTATTG	302	nd	10.40	nd	10.50
6434-R3-1	AGCCCTCCCACCCAGCCAGCTGCAGTGC	57	1.33	1.69	1.63	1.70
6450-R3-2	CTCCAATGGTGCTCTCCTGGTACTCATGGAAC	303	1.41	nd	nd	nd
6478-R2-2	GCCAAATTCTGCCCTTGATATGCATGCACAATT	304	-1.72	1.93	1.69	2.04
6496-R3-1	CCCCTCCCCACCCACCACTTCCCCTAGAGTCC	60	-1.36	1.95	2.28	2.20

Table 2

fold-changes in cell lines v. normal Lung						
Array probe	Array probe sequence (5' to 3' , without linker)	probe SEQ ID NO:	H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
6554-L3-2	TCCTTGTCATCTAGAACTACTTTGGTGCCCTCCATA	305	nd	3.79	nd	nd
6584-L1-1	TCGGCCCTGCCCTCCTCCTCCT	61	-1.36	1.82	1.33	1.52
6602-R3-2	AGAGCCCCAGTGGAATCTCTCTCCAAATCCAT	62	2.63	13.21	10.59	18.66
6642-R3-1	CACGTCCCTCCCTCCCTCGAGGTGTACACA	63	1.20	1.79	1.25	1.90
6647-R2-1	CTCAGCCCCAGCTGGAGAATTTTCCCCTCATTA	306	1.56	nd	nd	nd
6664-R2-1	GCCACCACCTCTCTTTTTTACAGGACATTACCA	307	nd	nd	5.29	nd
6681-R2-1	CCTGTTTTCTCCCTCTCTCTCTGCCCCCTCC	64	4.99	7.90	9.03	11.45
6712-L2-1	GCTGTGCTCTTGATATCAGTTGTCAGCCTG	308	1.83	4.86	4.41	nd
6718-L3-2	GCCTCCACCACCATAGGGCCAGAGCTTCTGCCT	309	-1.53	2.03	1.89	1.92
6718-R3-1	ATAGCCACCTTGTTTTGTGTTTGGCAAAG	310	1.66	7.03	nd	nd
6752-R1-1	CCCTCCTTTCCCCACCTCAGT	66	-2.15	1.89	1.60	2.23
6803-R3-1	GCTCCCTCTCTGTTGGACCTCACCCAAAGAT	68	-1.53	2.01	1.94	2.16
6839-L3-1	GCCCGCTGGCCCTTGCCACCCCCACCCCT	69	-1.88	1.57	1.43	1.81
6880-L3-2	ACCTCCCCCGGAAGACATCCACATTCTGCA	70	-2.04	1.49	1.48	2.07
6906-L3-1	GTGTCTTCTCCCCAACCCAGCCAGCTCTCCTGG	71	nd	3.61	nd	nd
6912-L3-1	GCCTTCAGCCTCTGGGTCCAGCAGTTAATTCT	311	-1.45	1.82	1.06	1.68
6930-R3-1	ATTAATCCTTCTCTCCCTCTG	72	2.11	nd	nd	nd
7019-R3-1	AGCATCAAACCTCCGTGCTAAATTTAAA	312	2.73	nd	nd	nd
7061-R3-1	ATGGAAACCCCAACCCTTCCC	75	nd	5.97	nd	nd
7070-R3-1	AGTCAACCTATACTGTCAGCACCAAGACCCAC	313	(nd) <3,72	1.57	2.14	2.63
7089-R1-1	CCTGAGCCAGCTCACATCACCCCTGACC	314	-2.12	1.57	1.68	2.43
7126-L3-1	GCACACCCGCTCTCCGGCCCCGGCCCCCTG	77	-1.46	1.95	1.68	2.00

Table 2

		fold-changes in cell lines v. normal Lung				
Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
7158-R3-1	TACATTTATAGATTCCCTCTTCAGCCATA	315	nd	nd	8.25	nd
7292-L3-4	GTCACCCAGTTAAATAGCTTCTGAACCTCCCTGCA	316	(nd) <4,32	1.40	1.39	(nd) <4,32
7304-L3-1	TTGATCCAAGCTCCACATTG	317	1.98	5.85	8.70	8.84
7340-R3-1	CTGCCACCAACTCTAATTGATT	318	(nd) <4,01	1.56	2.37	2.64
7352-R3-2	GCCCCTGCCAGAATCCTCTAACAGCTCTAATTGG	81	-1.43	1.57	1.61	2.40
7356-R2-1	GAAGCTCCGGCGGACGTCCCGTTACTCC	83	-1.80	2.04	2.41	3.11
7375-L3-1	AGCGCTGCTGTCTCCACAGTTACATACCTG	319	nd	7.15	9.97	12.05
7384-R3-1	CTCGCAAAGGATCTCCTTTCATCCCTCCCCA	85	-1.49	2.00	1.91	2.17
7411-R3-2	AGTCCCTGCCCTCATCTGCCACCCCTAATGAC	86	-1.61	1.70	1.62	1.87
7421-R2-1	TAAAGAGACTTCCTCCACTGCCAGAGATCT	87	2.12	4.83	8.07	nd
7426-L3-1	TGGGAGACGAACACCTCCTGCTGTGCTTG	88	4.29	nd	nd	nd
7435-L3-2	CTCCCCATCTGTTGCTAAGCCCCCATTAGCTGTGT	320	-1.79	2.09	1.86	2.33
7543-L3-2	TGCCGATGTCGTCCCTAATTACACAGGCCCCCGA	321	nd	3.71	6.97	nd
7571-L1-1	AGGGCTCCCCCACCCCTAAG	90	-1.73	1.43	1.50	1.43
7572-R2-1	ATCACCCCTTCCCCCTCCCAAATAAAG	91	-1.39	1.80	1.94	2.21
7578-L3-1	CGCAGTGCACACCCCTGAGCTACAGCCCCCTC	92	1.13	6.99	5.80	1.72
7597-L3-1	TTAATGGAACCTGGGCTCTGTGTC	322	1.75	4.17	8.05	8.93
7660-L2-1	CCCGGCCTCCGCCTGGCCCGAGCGATAA	93	-1.11	1.74	1.69	1.78
7702-L2-1	CCCAGAGAACCGGAATTCCTCCCCGCCCC	94	-2.10	1.41	1.47	1.48
7726-R3-2	CATCCCTCTCCAGAAGAGAGAGAGAAACA	95	-1.60	1.79	2.31	2.56
7763-R3-1	GTGCTCCCAATCCAGACGATCCATTA	323	-1.67	1.67	2.28	2.52
7764-R3-2	CCCTCTCTGCCTCTCTCATCACCAATAACAGAC	96	-1.09	1.93	1.74	2.01

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
7824-R3-1	TGGTGCCAGCTTCATCGCCG	324	nd	3.25	nd	nd
8004-R3-2	GGAAGTGTCTCTCTGCTCCAGTCATTGAAG	97	nd	3.51	nd	nd
8016-L3-1	TCAGCGCAACAAGCCCCGCGAGTCACCCCTCT	98	-1.99	1.68	1.62	2.30
8075-L3-1	CCGAGCTACACCTCCACGCA	325	-1.64	1.92	1.68	1.83
8077-R3-1	CCATTCCCACCCCTCAGGTAGTAAAAATA	99	-1.69	1.48	2.13	2.56
8169-L3-1	AACAGAAATGATTATTTACCTCCCCACATG	100	-2.46	1.52	1.79	(nd) <5,79
8250-R3-1	CAGCCGCTCTCCCTCAGCGTTAA	101	-1.41	1.55	1.60	1.84
8263-R3-1	GATTAAAAACAAGAAATCTATCTTCCCCCAGT	102	nd	nd	5.24	nd
8281-L3-1	AGCCCCTCCCCAGCTGCAGCTGAGGGCTGG	103	-1.34	1.78	2.04	2.78
8336-R3-1	CTCAGTCCCCACACCCCCCAGCCAGAGTC	326	-1.64	1.95	1.94	2.16
8394-L3-1	CCCCCGCCCTGCCCATCTCCGACT	105	-1.43	1.63	1.71	1.75
8433-L3-1	AAATGGCTCCTTTTCCCTTTTCCCTCCACCG	106	-1.35	1.78	1.76	2.47
8434-R3-1	CCTGAGGCTCCACTCCTAGAAAGATTGC	327	-1.61	2.16	1.98	2.42
8552-R3-1	CTCCCCAAGGGTCACCATATAAGAGGACACTATAA	328	nd	3.14	nd	nd
8564-L3-1	CCCTTCACCCCAGTTGCCAAACA	107	nd	nd	5.92	nd
8587-R2-1	CTCGCCGGCTCCAAACTTTCCCCAACTCCAGG	329	-2.38	1.99	1.58	2.15
8685-L3-1	CTGCTCTTTGCCCTCCTATAAGTGGAATGTCTCCC	330	-1.07	2.02	1.43	1.80
8719-L3-2	CTCACTTGCCCTCCTGCAAAAGCACCGTAGCTGC	331	nd	3.12	nd	nd
8724-R3-1	GCCAAGCTTGGAACTCTCCCTGCCAGCATCAC	109	nd	3.62	7.40	nd
8731-R3-1	TCATTGATGCCCCCATCTGCCCAG	110	-1.66	1.77	1.84	2.45
8760-L3-1	CTGGAGCCCCGAGGCAAAACTCACCCCCAGGCA	332	-1.85	1.57	2.29	2.59
8808-R3-1	CCAAAGACCCCTTTTCTCCCAGCCTGTTTCTGCAA	111	-1.69	2.22	1.94	2.25

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
8898-R3-1	CGGACGCCCCGCTCCCGCCA	112	-1.39	1.61	1.58	1.74
8898-L3-1	GAGTTGCCGGCGGCCGCCCGCGCGACAGCGCC	333	-1.51	1.99	1.88	2.15
9053-L3-1	GGCCCTGGGAATCAGAGACAGTGCCCTTCC	334	-1.75	2.11	1.90	2.20
9053-R3-1	TTCTTGCCCTCCAATCCCCGGGCTCCACAGCC	114	-1.57	1.95	1.65	2.13
9068-R2-1	CTGCCCTCCCTCTTGATCAAGACTGCTCTCTAA	115	nd	nd	nd	71.77
9092-R3-2	TCCTAGAAAGCCCATCAGTATCCACAGAGCCAG	335	-1.79	1.53	2.01	2.23
9217-L3-1	ACGATCCCCGCGGTGACTAAAGCCAAACAGTGGA	117	-1.38	2.00	1.87	2.15
9245-R2-1	AACCTCTCATTAGCCAGCCACTCGCTCCCAAG	118	1.91	4.41	6.66	nd
9387-R2-2	TCCATCCTTGCCGTCGCCCTTCATCTCAAAGCCATC	122	1.93	nd	nd	nd
9507-L3-2	GTCTCCCTCATCCATCATCC	124	-1.86	2.14	1.82	2.24
9557-R3-1	ACTGGCCCCAGTCCATTCTGCACCTCTTGCCCTA	336	-1.26	1.52	1.53	2.58
9582-R3-2	TACAAATCCTCAGATGTTTCCACAAAGGCTCCCTT	337	-1.80	1.69	2.19	2.19
9688-L2-1	GCTAAATGGCCCCAGACTGTTCTGCTGCA	338	nd	3.61	6.62	nd
9694-R3-1	GCCATCTGCCACCGACACTCATACTCTGT	339	-1.27	1.56	1.76	2.24
9733-L3-1	AAGGCTGTCCCTCACCAGACTTCCCCACCCCT	129	-1.55	2.36	2.12	2.58
9747-L3-1	GCACACCGCCTCCGGCAAACTGC	340	-1.27	1.88	1.37	1.56
9772-L3-1	ATTCTACAGCATTTTCCCATGACCTTTCCTGA	341	-2.02	2.12	1.72	2.14
9774-R2-2	CCGCCCCCTCACCGCCTCCTGCTCCCATCAGGC	130	-1.09	1.68	1.54	1.66
9798-R3-2	GTGCTTTTCATTTCCCCCAACAGAGGGCATT	342	-2.10	1.69	2.24	2.40
9812-L3-1	AAGCTCTATTTATCTGGGCTCCCCAGCTTGCT	343	-1.54	1.49	2.00	(nd) <4,17
9813-R3-2	GAAAGTAGAATTTGGCCCCCTCCAACTGTACAGGATGA	344	-1.08	1.68	1.65	1.87
9816-R2-1	CTGGCCCCCTTAAAGAGCCTCTCCGCGCTGCCG	131	-1.25	1.26	1.67	1.80

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
9987-R2-2	CAGGCTTCACCCCTCAGCCCACTTTGTTAAC	345	-1.73	2.09	1.92	2.17
10010-R2-1	CTCGCCGGCTCCAAACTTTCGCCAACTCCAGG	346	-1.90	2.06	1.75	2.07
10030-R3-1	AACTCAGCTGCCTTCCGCC	134	-1.41	1.98	1.48	1.79
10093-R2-2	ACCAGCCAGACCCCCCTGTAGTCTAACCCAAAGGT	347	-2.07	2.17	1.72	2.24
10120-R3-1	TTCCCCCTTGTTAAAAATTACAGCTGCACCA	348	-1.97	1.92	1.73	1.98
10133-R3-1	CTCCTCCCAATTCCTAATTTGATTTTAC	349	-1.45	1.90	1.88	2.28
10138-L2-1	AGCCTGCTCCGCTCTCCCTCCACCAGAAAAAGT	135	1.10	1.72	1.74	1.93
10154-R1-1	GATCTGTGCCCTTTGCCCTT	350	3.51	7.66	5.06	(nd) <8,10
10198-R3-1	AATTCTCTTTACCTGGCACCTTTAGGGCAAAGCA	351	1.64	3.88	7.36	8.75
10231-L3-1	GTGCAGCAGCCCGCGCCAGCCTCCGCAGCCGCC	139	-1.49	1.78	1.67	1.98
10231-R3-1	TGAACITTTAGCTGGGCGCGCCCTGTCAGC	140	-2.21	-1.70	-1.39	(nd) <8,10
10242-R3-1	GGAAGAGCCCTTCGCTTCCACCCCGAACAC	141	-1.11	1.49	1.33	1.40
10260-L3-1	AGGGCCCCCACCCGATGTCTCCAC	352	-1.85	2.24	1.79	2.27
10333-L3-1	TGTGCCCTGCCACCCCTCCCTGCCCCG	142	-1.21	1.48	1.74	1.85
10335-L3-1	CACTCCCCCTCCTTTTAAATTAGAAAGCACTAAGA	143	-1.31	1.94	1.97	2.22
10342-L2-1	CCACTTTCTTGGCGACCCCTCCGTGCGTGGG	353	-1.41	1.54	1.44	1.68
10342-R2-2	CCCGCCGCCGGAGCATCTCGAAGTTAATTAAA	144	-2.74	-2.11	-1.51	(nd) <13,60
10346-R3-2	AGCCTGTCTGTGCCCTCTGCAGCAGCTCACC	354	2.40	4.47	4.41	nd
10366-R3-2	AAACACCAACCACGCTCTTGCTACAAGACCCACAT	145	-1.64	1.88	1.91	2.29
10539-R3-1	TATTTGAGAAAAATTTACTATCCCCCAGCCT	355	-1.64	1.82	1.96	2.08
10543-R3-1	GCCTTCACCCCTTCCCATCC	356	-1.44	1.70	1.76	2.27
10553-R1-1	GCTGGCTCCATGCTCCAGTGGG	357	-2.27	2.01	1.64	2.01

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
10562-L1-2	GGGTCCCTGACTCCCACAGCCCTGTCATATCAAGCGC	358	2.42	6.15	8.86	8.95
10594-L3-1	ATTGTTACCCACACCAACACCCCACTCAACAG	359	-1.91	1.96	1.74	2.18
10639-R2-1	AGACCGGACTCGCCTCTTCCAACTCGAGTTCA	360	1.44	nd	6.00	nd
let-7a	AACATATACAAACCTACTACCTCA	166	-1.87	-1.06	1.39	-1.34
let-7b	AACCACACAACCTACTACCTCA	167	-1.85	-1.12	1.72	-1.38
let-7c	AACCATACAACCTACTACCTCA	168	-2.23	-1.18	1.47	-1.30
let-7d	AACATATGCAACCTACTACCTCT	169	-2.07	-1.13	1.38	-1.40
let-7e	AACATATACAACCTCCTACCTCA	170	-2.12	-1.05	1.46	-1.10
let-7f	AACATATACAATCTACTACCTCA	171	-1.94	1.00	1.29	-1.33
let-7g	AACGTGTACAAACTACTACCTCA	172	2.40	1.81	1.31	1.75
let-7i	AACAGCACAAACTACTACCTCA	173	3.29	1.99	1.31	2.02
miR-100	CACAAGTTCCGGATCTACGGGTT	174	-3.44	1.01	1.30	(nd) <7,51
miR-103	TCATAGCCCCTGTACAATGCTGCT	361	3.15	4.00	3.73	nd
miR-106a	CTACCTGCACTGTAAAGCACTTTT	362	-1.18	1.46	-1.32	(nd) <8,74
miR-106b	ATCTGCACTGTCAGCACTTTA	363	3.49	7.29	4.80	nd
miR-107	TGATAGCCCCTGTACAATGCTGCT	364	3.09	4.03	3.57	nd
miR-125a-5p	AGGGACTCTGGGAAATTGGACACT	365	nd	4.38	4.89	nd
miR-125b	TCACAAGTTAGGGTCTCAGGGA	179	nd	9.62	7.23	nd
miR-130a	ATGCCCTTTTAAACATTGCACTG	366	2.00	4.87	5.06	nd
miR-130b	ATGCCCTTTTCATCATTTGCACTG	367	1.85	4.20	4.06	nd
miR-134	CCCCCTCTGGTCAACCAGTCACA	368	-1.92	1.87	1.64	1.92
miR-138	CGGCCCTGATTCAACAACACCAGCT	369	1.56	nd	nd	nd

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
miR-155	ACCCCTATCACGATTAGCATTA	187	nd	nd	3.73	nd
miR-15a	CACAAACCATTATGTGCTGCTA	370	nd	nd	5.75	nd
miR-15b	TGTAAACCATGATGTGCTGCTA	371	nd	nd	6.48	nd
miR-16	CGCCAAATATTTACGTGCTGCTA	188	-1.03	2.33	3.10	(nd) <5,85
miR-17	CAAAGTGCTTACAGTGCAGGTAG	372	-1.18	1.52	-1.35	(nd) <8,91
miR-181a	ACTCACCACGACGCGTTGAATGTT	373	-1.45	(nd) <3,46	-1.05	(nd) <3,46
miR-181b	ACCCACCGACAGCAATGAATGTT	374	2.17	nd	nd	nd
miR-191	CAGCTGCTTTTGGGATTCCGTTG	375	1.89	6.53	3.50	nd
miR-195	GCCAATATTTCTGTGCTGCTA	376	nd	4.38	5.32	nd
miR-196b	CCCAACAACAGGAAACTACCTA	377	1.49	nd	nd	nd
miR-198	GAACCTATCTCCCCCTCTGGACC	190	nd	5.34	7.25	nd
miR-19a	TCAGTTTTGCATAGATTTGCACA	378	-1.38	1.29	-1.40	(nd) <7,33
miR-19b	TCAGTTTTGCATGGATTTGCACA	192	-1.42	1.30	-1.56	(nd) <10,00
miR-200b	TCATCATTACCAGGCAGTATTA	193	(nd) <8,612	(nd) <8,61	(nd) <8,61	(nd) <8,61
miR-200c	TCCATCATTACCCGGCAGTATTA	194	(nd) <18,47	(nd) <18,47	(nd) <18,47	(nd) <18,47
miR-205	CAGACTCCGGTGAATGAAGGA	195	(nd) <28,34	(nd) <28,34	(nd) <28,34	(nd) <28,34
miR-20a	CTACCTGCACATATAAGCACTTTA	379	-1.33	1.22	-1.87	(nd) <8,17
miR-20b	CTACCTGCACATATGAGCACTTTG	380	-1.53	1.33	-1.76	(nd) <6,30
miR-21	TCAACATCAGTCTGATAAGCTA	196	-7.45	1.45	1.38	1.98
miR-22	ACAGTTCTTCAACTGGCAGCTT	381	-1.31	-1.07	1.33	-1.71
miR-221	GAAACCCAGCAGACAATGTAGCT	382	-4.50	-1.92	-1.27	-2.19
miR-222	ACCCAGTAGCCAGATGTAGCT	383	-3.60	-2.07	-1.17	(nd) <11,32

Table 2

Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	fold-changes in cell lines v. normal Lung			
			H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
miR-23a	GGAAATCCCTGGCAATGTGAT	197	-2.44	-1.24	1.05	(nd) <14,57
miR-23b	GGTAATCCCTGGCAATGTGAT	199	-2.14	-1.20	1.22	-1.04
miR-24	CTGTTCTCTGCTGAACCTGAGCCA	200	-2.89	1.30	1.02	1.18
miR-25	TCAGACCCGAGACAAAGTGCAATG	384	3.98	8.33	5.07	nd
miR-26a	AGCCTATCCTGGATTACTTGAA	202	-3.08	1.23	1.24	(nd) <7,67
miR-26b	ACCTATCCTGAATTACTTGAA	203	nd	3.58	nd	nd
miR-27a	GCGGAACCTTAGCCACTGTGAA	204	-2.28	-1.28	-1.04	1.02
miR-27b	GCAGAACTTAGCCACTGTGAA	205	-1.92	1.27	1.14	1.38
miR-29a	TAACCGATTTTCAGATGGTGCTA	207	-1.77	1.26	-1.01	-1.07
miR-29b	AACACTGATTTTCAAATGGTGCTA	208	1.16	1.16	-1.30	(nd) <5,77
miR-29c	GAACACCAGGAGAAATCGGTCA	385	-1.29	1.29	-1.29	(nd) <7,48
miR-30a	ACATTTGTAGGAGCTGACCTTC	386	-1.62	3.78	2.06	(nd) <4,49
miR-30d	CTTCCAGTCGGGGATGTTTACA	216	2.12	16.27	7.18	nd
miR-31	AGCTATGCCAGCATCTTGCCT	387	-10.63	-2.12	-2.25	-1.74
miR-320a	TTTTCGACCCAACTCTCCCGCT	388	-1.22	-1.05	1.49	1.15
miR-335	ACATTTTTCGTTATTGCTCTTGA	389	nd	nd	3.55	nd
miR-342-3p	AGAGTGTGTCTTTAGCGTGGGCA	390	-3.37	1.81	1.25	1.76
miR-370	ACCAGGTTCCACCCCGCAGCGC	391	2.02	5.04	8.94	nd
miR-424	TTCAAAACATGAATTGCTGCTG	222	nd	nd	5.25	nd
miR-452	TCAGTTTCCCTCTGCAAAACAGTT	392	nd	nd	4.54	nd
miR-494	GAGGTTTCCCGTGTATGTTTCA	393	1.30	2.20	1.01	1.55
miR-513a-5p	AAGTGTCCTCCACAGTA	394	3.47	5.11	7.21	nd

Table 2

		fold-changes in cell lines v. normal Lung				
Array probe	Array probe sequence (5' to 3', without linker)	probe SEQ ID NO:	H460 (Large cell carcinoma)	H1703 (Adeno- carcinoma)	BEA2B (Immortalized bronchial epithelial cells – normal phenotype)	A549 (Adeno- carcinoma)
miR-614	CCACCTGGCAAGAACAGGCGTTC	236	-1.85	1.83	1.69	1.99
miR-638	AGGCCGCCACCCGCCCGCGATCCCT	239	-1.95	-1.02	1.09	-1.05
miR-658	ACCAACGGACCTACTTCCCTCCGCC	240	-1.35	1.50	1.30	1.60
miR-663	GCGGTCCCGCGGCCGCCCGCCT	241	-1.86	1.08	1.37	(nd) <6,22
miR-671-5p	TCCTTCGGGACCTCCCCGACCTC	242	1.85	nd	nd	nd
miR-7	ACAACAAAATCACTAGTCTTCCA	395	3.27	nd	nd	nd
miR-765	CATCACCTTCCTTCTCTCTCCA	246	nd	4.51	nd	nd
miR-92-a	ATAACGTGAACAGGGCCCGGACA	396	4.00	6.61	3.83	nd
miR-93	CTACCTGCACGAACAGCACTTTG	251	-1.07	1.75	-1.04	(nd) <5,78
miR-98	AACAATACAACCTTACTACCTCA	252	-1.90	-1.05	1.43	(nd) <5,20
miR-99a	CACAAGATCGGATCTACGGGT	397	-2.49	1.13	1.31	(nd) <5,34
miR-99b	CGCAAGGTCGGTTCTACGGGTG	253	2.05	4.50	3.79	nd

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
266-R4-1	12q14.3	GTTGCTATTTCCCTCAGTTGAGGGCGAAGTTAGCAAATCCGTAGCTGCAAGTCTCAACTTGGGGGAGGGGGCGAC	398
673-L4-1	01p36.21	GTGAGGAGCAGGTTAGCTGGGTGAAAAGTTCACAGTGAGGGGAGCTGTCTGTTCCCTCGCTTAATTTATCCACTATTTGGCTAACCTTGCTCTGAAC	399
836-R4-1	03q26.2	AAATAAGCCATTCCAAACCATTTCTGATTTGCTGTGAGTGGCAGAATCATTCAACCGTGGTGAATCATAGCAGGGAGAAACCAATTTGGAATGATTATTT	400
3249-L4-1	01q22	GGCGGCGGCGGGCGGCTCCGGGGGATGGCGGGGCTCCGCTGCTGCTGCTGCTGCTGCTGCCCGTGCCCGCTGCTGCCGCTGCTGCTGCTGCTGCT	401
3371-L4-1	18q21.33	CTCAAGTGTGGGAGTCATGGGGTGTGGAGGGGAGGAAAGGATTTTTGTTTTCTTTGTCTATACATTTTCCTAGATTTTCTATG CAGTTGGG	402
3717-L2-1	01q22	TGTGAGTGGGTTTGGGTGAGGCTATGGGAGGGCGGGGTGCCGCCCTTGCCACGCCCTGAGGGCCCCAGCCAGTACA	403
3799-R3-1	11q13.1	GAATTTGCCCTACGGTGTGACCCCCAGCCTCTCCCTCTGCCACAGCCAGGCCGCGGGGCCCTCTGGGAGCATCTTTCAGCAAGTTC	404
3872-L1-1	16p13.2	GTGCATAAGGGAAGACAAAGAAATGATATTGTCGTTTAATAGTTCACTTTAGATCTTCATCTCTTATCAC	405
3875-R3-1	05p15.1	GGCTCGGTTTCAAATCTCTCCTAATCCACTAATGAACCTTTATTAAAGTGGGAGAGAGGTTGAATCAGTC	406
3897-R3-1	09p11.2	CGAGCCGCCCGCCAGCCTCTCCATCTCGCAAGTTTTAATTAACGCTGAGGGGAGCGGCTGACGGCGGTCGG	407
3923-R3-1	19p12	GGCTCTGCACCAGGCGTTTCTTCTTGTTTCCCTCTTCTCTTCTGGAGGGGATGAAGGAGATCCTTTGTGAGAGGC	408
3953-R3-2	09q33.3	GCTCCTGCTCCGCCGCGGAGCTGCTCCGGCGCCGACGGGCTCGCTCGGGAAGCTGAGGCGCGGAGGCTGGAGT	409
3995-L2-2	07p21.1	TGGCCTGACGTGAGGAGGAGGACTTTTCGAAGTTTATAGGAAAGTTTCCGCTTTCCAGTCCCCCTCCCCGTCCCA	410
4026-R3-1	10q24.32	GGCTCTGGGAAAGCCTTCCTTTCCCGCTGGCCTGGCATTCAAAGCCAGACAAAGGGGGCTTTCTCTCTCGCC	411
4037-R3-2	03q13.31	GCCCATTTCCCTAATGGCAGCCGATTGCCATTTGCTATTCAAATCAGACTAGATAAAATTACAGTACATGCCAGGGAACAGGC	412
4143-R3-1	22q12.3	TCAGCGTCAGGAACCTTCATCCTGGCAGCCGACCTCATGAAGAGCATCTGGCTGCTGTTACCAGGAGGAGAGCAAGACGCTGA	413
4203-R3-2	11q23.3	AGCAATTCCAACTGCCCCCATTTATATTCCTAAGTAGAGGACTTGTTAATATGGAGCCCTGCCCTCTGGGAAAGTGGGAATGTGCT	414
4205-R3-2 4205-R3-1	15q26.2	GTCATAGATGGCCTCATTGTCTACCATGAAGCACAAATCAGAGTGCTCTAGGGTGGTGTGAGTGGTGAAGTTGTAGGGC	415
4303-R1-1	22q11.21	GGCTGGCCAGGCTCCGCCCCCGCCCTCCCTGCGCCCCGGCGGTGCGCAGGGAGGTCGGAGAGCGGGCACTGCCCACCC	416
4315-R3-2	01q22	GCTCCCCGGCTCCCTCACTGCGGCAGCCCGCGGCCCATAAATCGTGAGAGCGACGTGCTCCGGAGCCCGAGAAATGGAGAGGGC CGGGGAGC	417

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
4361-R3-1	Xp11.22	TGCTGGAGGTAAGGGTTTTCTGAAGCCTGGTGCCATGGCCACATGTGCACATGAGGGAGGAGACGCTGAGGCTAGCA	418
4440-L3-2 4440-L3-1 4440-R3-2 4440-R3-1	07q11.22	GTGATGTGATTTCTGCCCAGTGCTCTGAATGTCAAACCTGAAGAAATTCAGTGAAATCGGGTAAACGGCGGGAGTAACTATGAC	419
4448-R3-1	01p36.32	TTTCCTCTCTCCTTCTCCTCAAGCTGATTAGCGGGTCGGGCAATGGCGTGAGGAGATGCTGGGGGTAGGAAA	420
4479-R3-1	01p32	CCCAAGCTCCTTCCTGGAGGACTTAACACTGTGTTGAGCAGTTGTTTTGAATTTCCGGCAGGGGCTGCAAAAGGG	421
4593-R3-1	15q23	CAATCAATTAGCACATGAGTAATACCAAGCCCCATTAGGACAAACTGATGCCGGGGAGTTATGTCACTCTGCTATAGAAATGATTG	422
4666-R4-1	01q22	GCCGGCTCCAAACCAGAGGCCCGGAATAGCGCGGAGTTATAAATAGTGCCACCCCGCAGGTGTTGGGGGGAGTCGGC	423
4790-L4-1	02q14.1	CCCAGAATTCTAGTAATGAGGGAGACAGGTTATGCCAAGCCTGCTTCTCCAGGATGCACCTGGGAGCCTGGG	424
4829-R2-1	01q21.3	GGTGTGCTGCCCTCTCTTTCTGCCCCCTATACCCCTTGACCCCGGGGAAGGCACTCGGGCAGCACAAAAGGAGCAGATGCC	425
4855-R3-1	12q13.2	GGTCCGGGTCTCTACCGCGCCCTCATGCAGGAGGCCCTTGAGCAGGAGGGGAAGCGGGAGACCCCGCAGCCCC	426
4875-R2-2	03p22.1	TGTGAAGCCACAGGAAGGGCTCTGTGACATCACAGGTAGGGGCAGTGTGAAGTACAGGAAGGGCTGTGGGAAGTCACA	427
4988-R4-1	14q24.3	CTTTTCTCTCTGCTGGAAACCTTGCTTGACTTCATGTCCAGTGTGTTGGTATCCAAAGACGGGGAGGAGGAG	428
5006-L3-1	07q32.1	GGAGCCAGCAGAGGGAGCTGCTGCCAGAACTTTCTTAGAGCTGCTAAGAAATTCGTATTTTGAAAAAGATCTTCTAGGCTCC	429
5080-R3-1	11q23.3	GGCGTTTCTTCTTGTTTCTCTCTTCTCTCTCTGAGAGGGATGAAGGAGACCCCTTTGCAAGAGGCATGTT	430
5192-L3-2 5192-L3-1	05q34	GTCTTTGCTGATATAGAGGAAGGAAGGGGAAAAATGAGCGCATTAGTTCTCTTTTATTTAAAAAGAGTTATTTTCAGCATGAC	431
5342-L3-1	08p21.2	TGGAAGGAGAGCAGCGGCATTTGGTTTGGTGGTGCGAGATTTTCTTTTACGACTGCTAAATGCCTTTCCTCTTCTCCCA	432
5521-L2-1	05q31.3	AGAGGGTGTGCTCTGGGAGGGCCACCCAAAGACAGACACCTCATGGCCTTCAGTCCCAGCCTTCCTCAGGGTCCATCCTCT	433
5554-R2-1	01p34.3	CCCCACCAACCACAGTGCTCAGGACTTCTGCAAAATCCCATTCCGGAATCTGGGAGCAGCTGATGAGGGGGTGGGG	434
5638-R2-1	06q23.2	GGCTTTGCCCTTTTCGGTGACACAGGCTGTTGCTATTCCAAGCAGCCTATCACAGGCAGGGGAGGGGCC	435
5640-L3-1	01p34.1	TCTCGAGAGGGGCAGGCACGGTGTTCCATGGCAAGACGGCGGTTGATGTATAGGCGTGGCATGAAGCTGGGCTTGCTGCTCTC	436
5726-L3-1	03p14.3	CAAGGGCACAGTGAGAAAGATATTTTATTACCCGTGTATTTATTTATGGGTTTTTGGGGTTTTTAAAACTGGCAATTAACCTTG	437
5782-L3-1	05q35.1	GGGGGAAGGGCTGGAATCATCGTGGGTTGGAACAGTTAAAGGAACCTCTGTTCAGCCCCAGCCCCAAGGCTCCC	438
5795-R1-1	10q26.3	CTCTCTACCAACCAAAATAAAATCAATTACTAACTTTGAGTAATTTGCGGGTAATTTTATTTCTTTGGAGGGGCAGAGAG	439

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
5836-R3-2	11q23.3	GCCATGGGCCTCCATAGTTTCCTGTAGCCCCCTTGTTCCCAAGAAATAGTTTTGGAATGGGCGTGCTGTGATAATGGGGTTAA TGGT	440
5854-R3-1	11p14.1	GCCCTCCCTCCCACCGCACCTTACACCTGAACCTTGCTCTCCAGCACTGCGGACACCCGGGTGACACATTCTTTTCGGAGAGGGAGGG C	441
5971-R3-1	01q24.1	TGCCATCTGCTCTGAAGCCTCCCAAGCTGGGCCTCCCCCTCCCACTTCTGGAGCCCAAGAACAGATGTGGGAGGCTGAGCAGGC A	442
6008-R1-1	04p15	CGACTTTATCACCCATCGGTTATCTGTGCGCTGAAGGAACTCCGGGTAAAGGTACAGGAAAAAGGTGGGGTATTGTTG	443
6016-R2-1	01q23.3	TTTCTGTATATGTTTCTGGAGTCCTGAGCCTGAGCTAAACAAAAGCAGGAGGCTGACGGGGCTGCTGGAGTTTGCAGAGA	444
6037-R3-2	16q22.1	GCAGGATCCCTCTTTTCATCTGAAAAATTACCACTAATTGCAATTAGTTGAGGAAAAATTGGAGATGGAGGAAAGGGAATTGC	445
6096-R3-1	03q29	GCCATTTGTACCTGATGTATCGGGCTTTTTCCCTGCTGTGAAAAAACGGGCAGGATTAAACATAAGGGAAAGGTGGT	446
6183-R3-1	12q21.33	GATTCATCTATTCTTTTTCTCCTTCTCAAAGATAAAGTCTGTAGCACCTTAAGGAGGGGAAAGTCATTAGAAAAAGTGAATC	447
6192-L3-1	11q25	GTGCTGGGTGGTGTTTTTATCTTCACGGATTTATGGAGTCTCTTAAACATCTGTTCGGTTCTGATTCCCCCGCTCAGTAC	448
6233-L3-1	06q16.1	GGAAATGGGAGAAGGATAAAGTGGAAATCTAATTTTGAGAAATAAGGATTAAAGGTTCCATTATTCATGCTGTTTTTC	449
6235-R3-1	15q26.2	TCTGTTTTTATCAGTTTTAATATATGATACATCTTCTATCCAAAGGACAATATATTAAATGGATTTTGGAGCAGA	450
6287-L3-2	01p34.1	AGCAGCCAGGTAGCCCCCGAAAGGTGGGGCGGCAGGGGCGCTCCCAGCCCCACCCCGGATCTGGTGACGCT	451
6409-L3-1	11q13.1	GTTCCAGAAAGCGCGCGTGCGGTTGGGAACGCGGAGCGGACGGATTTCGATTCAACGGGGTTCGGGACCGCGCTGCGCTATG GAGC	452
6434-R3-1	15q25.2	GTGGGCTGCATGTTCCCGCATTTGCTGGTGAGGGTGCACGATCTGGCACTGCAGCTGGCTGGTGGAGGGCTGCATCCTAC	453
6484-R3-2	12q21.32	ACTGCTTAGCACTCCTCCACTTGCGAATGCACCTTAAGACAGTAGGTGTGGTTGCAGTTGGAGAGGATCCCCGAAGCGGT	454
6490-R4-1	01q22	TCCTTCCCCCTTCGTGGCTTCGGTCTCTCTTCCCCGCCTCGGCCCCCAAGAAAGTGTAGTGTGGGGTGGTGAGGTTAGGA GGGGAAGCGTCATATGGGGATGGGG	455
6496-R3-1	05q31.1	CTCCCTCAGGCCCCGCTGCACCTTTCCAGCCCCCAGGACTCTAGGGGAAGTGGTGGGGAGGGGG	456
6584-L1-1	12q24.23	GCTTGGTGAGAGGAGGAGGAGGCGGAGCCGCCACCCGCCCTGTCTGCCATCTGGTCCCCCTTCCCCCTCCTCTCATTTG C	457
6602-R3-2	03p13	CAGAGCTAAATGGAAGAGTCCTCCGTATTTACCCAGCTCATCTCCTGTGTAATGGATTTGGAGGAGAGATTTCCACTGGGGCTC TG	458
6642-R3-1	14q11.2	ACACTCTCCTCTTGTCCTTGTAAATCAATTCAATTGTATCATCAGAAATGTGTGACACCTCGAGGGGAGGGGACGTGT	459
6681-R2-1	11q12.2	TGTGCTCTCATTTGTTATTCAAAAAGTCTCTGTCTAGATCACTGGAGGGGCAGAGAGAGAGGGGAGAAAAACACGGGAGATACA	460

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
6683-R3-1	12q23.2	GTTCTAGTTCAGGATGCTGATACTTTAAGCCCGAGGCTCTAACTTGAGCAGGAAGAGTTTATTTTGGATGAAGAAT	461
6752-R1-1	Xq13.1	CCCTCCCAGTTCCCATAGCAACTGGGCTGTAGCAGCCAGAACTTGATTGAGCCCCAGCAGTGGCCCCGACTGAGGTGGGAAAGG AGGG	462
6795-R4-1	06p12.2	GGCCACGCGAGGAGGCACTTGCCGAAACACACACACTTCCTTACATTCCATAGCAAAGTAATCCATATGGGGATATAAAGAAGCAT GTGGCCTCGGGAAGCAGTGTC	463
6803-R3-1	22q12.3	GCCACCTTTTCATGGTGAGGATGCCCTGCCACCTTCAGGATCACATCTTTGGGTGAGGTCCAACCCAGAGAGGGAGC	464
6839-L3-1	03p21.31	AGGGGTGGGGGTGGCAGGGCCCCAGCGGGCTGGCAGGCCAAACCCCTGGTTTTTGGCCCCAGGGACCTATAATCAGCTCCTGCCCCCT	465
6880-L3-2	01q42.13	GGCTTGCAGAATGTGGATGTCTTCGCGGGGAGGTGGCCACGTTCTCCTGTGAGGTGTCTCACGCGGGTGGGCC	466
6906-L3-1	06p21.1	CCAGGAGAGCTGGCTGTTGGGGAGAAACACACTAACCCCTGTGAGTCTGACCTCAGCCAGCTAACCTGCCCTGG	467
6930-R3-1	09p21	TGTCATTTGTCCATTTTCTCTTGACCCAGTGGTATTCTGCAAGATCAGAGGGGAGAGAAAGGATTAAATGTCA	468
6984-R4-1	01q22	CCCCACTCCCTTGCAGGCTGCAGGCACTAGGGCTCTCAGGAATTGCAGGGACTTTGTGCCCCAAGCAAAATGCTTGGCAGGGG G	469
7026-L3-2	15q15.3	TCGTTCCCGGATCTGGTGAGGTTTTTCGATCAGGGCAAATACCTGATCACAGACCTTCACAGGATTCTGGATGA	470
7061-R3-1	01p13	TCATGGCAGCGACCCACCTCCAGTCCCCTGGACAATCGGGTACAAGAGACTTAAGGTTGGGCATGGGAAGGTGGGGTTTCCAT GA	471
7066-R4-1	15q23	CAAGGTCTTTGGTCTTGGAGGAAGGTGTGCTACTGGAAGAGGCCACCAGGCAGGGCTGGTGGGGCATCTTTTTTCAGGGCTAC GGCCCTTG	472
7126-L3-1	05q31.1	CAGGGGCGCGGGCGGAGAGCGGGGTGTGCAAAGTGGGCGCAGGGCCCTGGGGCCGCGCCCTTGCTCTGCCGGCTCGACTC TTG	473
7182-L4-1	12q13.3	GGGGCAGGGGAAGGTGACGGAACGGCTAGTTACCCAGAATTCTCTGGGGAACCCAGAAAAATCGGTTATCTAGAAATTTCTCCC	474
7192-R4-1	09q33.1	TGTAGCAAATCCCATCCATCTGTTTGGCTGCTCTTGCCCTCAGTGACAGTGCCAAAGAGCCCCAGGCAGACTTAGAGGGGAAAGTGC TTTTGCA	475
7292-L3-2	01p34.1	GCAATTAGAATGCAGGGAGGTTCAGAAGCTATTTAACTGGGTGACCCCTGAGGTCGCTGCATCTGACTCCCCATCCCTGGATAAAT ATTGT	476
7352-R3-2	01q25.2	GCCTCTGTGCGCATGGATATAATCAGCTTTGATAGGCAGAGGCTGAGGCTGTTTTTCCAAATTAGAGCTGTTAGAGGATTCTGGCA GGGGC	477
7356-L2-1 7356-R2-1	08q24.3	GGGGCGGAGGCTATGTCGCGGTGGCAGCCCCGGATGGGCCCGGCAGGGCCGGGAGTAACGGGACGTGCCCCGGGAGCTTCTTC CCCC	478
7367-L1-1	06p21.33	CGGTCCCCAGAGGGGCAGCTCTAACCCCTAAACAAGTGCTCAACCCCTTGAATGGGCCCTGGATGGCTCCCCCTGGGGACTG	479
7384-R3-1	12q12	GGCATTTCCTTCTGTGTTTCCTCTCTCTCCCTCTTCTGCGGAGGGATGAAGGAGATCCCTTTGCGAGAGGCGCATGTT	480

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
7411-R3-2	18q22.3	GAGTGTGAACTGGCTCCAGCTGTGACAAATAAAACAGCAGGTGGCTGCTGCTGCATTAGGGTGGCAGATGAGGCAGGGGACTAACA TTC	481
7421-R2-1	12p13.31	TGAAGAATTTCTTCTGGATGACTGACCAAGAGGGCTATTCAAGATCTCTGGCAGTGGAGGAAGTCTCTTTA	482
7426-L3-1	02q22.3	AAATCAAGCACAGCAGGAGGTTCGTCTCCAGGTAATGGGTAAATGATGAGCAGATGAGCCATCCTTCTATTGATTT	483
7569-L3-1	11q23.3	GGCCTGTCTTGGGGTAGCTTTGTGGCCTGAAAACAAATCATCCTTACAGCTTGCTCCCAAGTCCAATAAGCC	484
7571-L1-1	02p21	CTTAGGGGTGGGGAGCCCTGTTAGCCCTGTAAATAAAGTTTAAACGAGGTGAACAATGGCTGGCTCTGTCCCTGAG	485
7572-R2-1	11q12.1	ATCACCTTCCCCCTCCCATGTGCTTTCTTCATTTGAGATCTTTTGACCTTTGGCTTTATTTGGAGGGGAAGGGTGAT	486
7578-L3-1	02q12.1	GAGGGGCTGTAGCTCAGGGGTGCACCTGCGAGGCTGGACCTGTTGAGTCTGCAGTGGACATCCATTTAGCTTCAGGTTGTC	487
7660-L2-1	19q13.32	GAGCTTTATCGCTCGGGCCAGCGGAGCGCGGCCCGCTGGCTTCCGAGGCGCCCGGGGATGAGCTC	488
7702-L2-1	10q21.3	GGGGCGGGAGGAATTCCGGTTCTCTGGGACTTTCAAAAAAGCGGAAGATCCGGTGCCGGGGCTCCGCCCTCCCTAGCCCT	489
7726-R3-2	12p13.32	CATTTACATCCATGAAGTAGGAATTGGGGCTCTGCACCAGCGTTTCTTCTTGTTTTCTCTCTCTCTCTCTGGAGGGGATG	490
7764-R3-2	05q11.2	TGCTATCTCGCCTCACACATCAACACACAGTGCCAGACAGATTCTGACTGCAAAAGTCTGTTATTGGTGATGAGAGAGGCAGAGAGG GCA	491
8004-R3-2	Xq28	GGGGTGCCATCCTGCTGCCGTCACTGTGTGGTGGTCAACGGCCTTCAATGACTGGAGCAAGGAGAAAGCAGTTCC	492
8016-L3-1	12q21.1	AGAGGGTGACTGCGGGGCTTGTTCGCTGAAGATTTACAATGTACTTCTTGAGGGGCTCAGCAACCCCTCT	493
8077-R3-1	Xq22.3	CCAATTCTCACTTAGGTGTTAGGGATTTAATGATACTCCTCTGAAGAGTATTTTACTACCTGAGGGTGGGAATGG	494
8169-L3-1	06q16.1	CATGTGGGAGGTAATAATCATTTCTGTTAIGTCAGTGGAAAAATTTCTGGAGATCATAAAGAAATCAGTTTACCCCTGAAGCATG	495
8250-R3-1	09p11.2	CGAGCCGCCCGCCAGCCTCTCCATCTCGCAAAGTTTTAATTAAACGCTGAGGGAGAGCGGCTGACGGGCGGGTCGG	496
8263-R3-1	Xq25.1	ACAGACATCCAGTCTAATTACAGGTGCTCTTTTAAACCAGTGAGAGGAAACTGGGGAAGATAGATTCTTGTTTTTAATCTGCTATCT GT	497
8281-L3-1	11q13.4	CCAGCCCTCAGCTGCAGCTGGGGAGGGGCTGAAGGGTAGGGAGCCCTATCCACCTGCATCAGAGGCCTGG	498
8316-R3-1	14q24.3	GTCAGGCTGCTGTATTCTTTACACAGATGCCAGTAAGAACAAAGGCATCACGTGGGAGAGGATACCCTGAT	499
8394-L3-1	07p13	GGGGCCAGGGATAGTCGGAGATGGCAGGGCGGGGCCCCCACTGGCAGGGGCCCTCGGCTTCTGGGGTCCCTGAGCCCC	500
8433-L3-1	17q25.3	CGGTGGAGGGAAGGAGCCATTTTCTGCTGCACATCAGTCAGTGCCTGCGCCCTCCCTCCCTCCGCCG	501
8564-L3-1	05q13.2	AGGGGTGGAGGTCTGTTTGGCAAACCTGGGTGAAGGATTGCCCTTCCCTGCTGGGATTCCCCCAGCCCCCT	502
10010-R2-2 8587-R2-2 8587-R2-1	17q23.3	CCGGGCGCCCCGAGAGCCGCCCTCCCTTCCCTCCGTGACAGGTGGGCTGGAGTTGGGGAAGTTTGGAGCCGGCGAGGGG CGCCGGG	503

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
8724-R3-1	15q23	GGCCCCAGAAAGATGAAAAGCTGAAAGTCCTTTCCCTTCCAGCTGAAGCCAGGTGTGATGCTGGCAGGGAGAGGTTCCAAAGCTTGGC _C	504
8731-R3-1	Xq13.1	GTCAAAACCTGGTTTTCCCTGCCCTAACCCAGGCGCCCTTTGTTAAGGGCCTGGCAGGATGGGGCATGAATGAGGGTGAC	505
8808-R3-1	03p14.3	CCGATTATGGCTTTCTCTCTGCCCTTTTCAGTAGTGATTTGCAGAAACAGGCTGGGAGAAAGGGGTCTTTGG	506
8898-R3-1 9989-L3-1	17p13.3	GGCGCTGTCGGCCGGGGCGGCCCGGCAACTCGTCCGTCTTGATAACCATGGTGGCGGGAGCGGGCGTCCGCCCTCGGCTGT CCGCGCC	507
9021-L4-1	10q23.1	GATGTGTGGGAGCTTGGGTGTTTGTTCCTATTTACAAAACAAAGCAGCCAAACCTTACATTTCATC	508
9053-R3-1 9053-L3-1	Xq27.3	GGAAGGGCACTGTCTCTCTGATTTCCAGGGCCTGTCAATTTCCCGAGGGCTGGTGAGCCCCGGGATTGGAGGGCAAGAAGCCCC AGCC	509
9068-R2-1	14q24.3	GTCTGCCCTCTTTCTCTGCAGTAATTGCTTCCTGACATTTGTTTTATTTAATTAGGAGAGCAGTCTTGATCAAGAGGGAGGGCAGAC	510
9087-L4-1	03p13	AGGAAGGAATGGACTGGGAGGGTTCTTTTCCTGATGGAAGCCTATTTTTCTTATTGTGTTCCCTTTTCT	511
9217-L3-1	02q31.2	TCCACTGTTGGCTTTAGTCACGGCGGGGATCGTCAGTTTAGCGCGGCCCATCGCTAAAGGAGATCTGCACGCCGGGCAGAGTGGA AGTGA	512
9245-R2-1	05q21.1	AGCCTAAATACATTAGCGAGCTGGTAAAGCTTTTAAGGCCCTTCTTGGAGCGAGTGGCTGGCTAATGAGAGGTT	513
9287-L4-1	17q21.1	TGCATGTGTGGCTGGGAGGGCTCTGAATATCTCCTGGAACGGTACCCAGAGCCCTGTGGCTCTGCGCATGCG	514
9347-L2-1	06q15	TCITTTGTTAAATGTAAATGCATATTGGGCAATGCTCCAGGGCAATTTGCATAAAAAAGTGATGACAAAGA	515
9349-R3-1	21q22.11	GGACACTCTGAACCCCAAGTGGAATCCAACTGCCAGTCTTCATCCGAGACCTGGAGCAGGAAGTCCCTCTGCATCACTGTGTTC	516
9387-R2-2	03p21.2	TCTCCATCCTCTGTCTCCCTTGATCCCTCTGTTCTCCCTGATGGCTTTGAGATGAAGCGCAGCGCAAGGATGGAGG	517
9391-R3-1	02p14	TAGCTGCCCTCAGAGTAGAAAAATAAACTCAACAAGATTTATCTTGTTTTTAATTTCTATGTCTCCCTGGCAGCTG	518
9507-L3-2	16q22.3	GGTGTTTGGATGGATGAGGATGGTGGATGATGATGAGGGAGACGGAGGATTCCCTTATTAAAGCATCAAATTTCTTCCCTAAATA TC	519
9564-R1-1	09q33.2	GGCGCCCGCGGGCTGTCCGGAGCGGCCGATGGGGCCCGTGTGAGCGCGCCCCAGGCCCGGGTGGCCCGGGCGGGCGGC	520
9594-R2-1	02q12.1	TTCCAGCTATTTAGTAACTCTTCCAAAACACTGTCAAGCACCCTATGCTAGGATGCAGGGAGTGGGAAGAGTCTAAGTAGGGAA	521
9656-R3-1	03q25.33	GTCCTTTAAGACAGTGTGTTCTAAAAATGTAGCCGAGGATCTTTTGGCTTGCAGCATTAATCTCCACAGTATGTACTTTAAGGGC	522
9691-L4-1	14q24.3	GCAAGGGGCCAAGAGGGAGATCGGGATGAAATGGATGATTTAATGGTCACTCTCTCCTGTAGTTAATTTCTCTAGATCTCTTGT	523
9733-L3-1	15q23	AGGGGTGGGAAGTCTGGTGAGGGACAGCCCTTGAGTCAAAAGGATGGTCAACCGTCCCATGTGGTGGCTGCCACCCCT	524
9774-R2-2	13q13.3	GCTTGTCTCTAAAGATCTTCCCTCTGTTTCCCTGGGTTTATCCACTTGGTTGGCCTGATGGGAGCAGGCGGTGAGGGGGCGG GGC	525

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
9816-R2-1	17q12	CTGGCCCATTTTCATTCTGCATAAAATTTTAATGGTCTCTCTGGCTGATCCGGGACGGCAGCGCGGAGAGGCTCTTTAAAGGGC CAG	526
9840-L3-2	05q14.1	CTTGATTTTGTGACATCCTGATTTATAAAACCTGAACAAGTTCAGTTTCAATAATCTTTTGTTCAGGAACACAAAG	527
10010-R2-2 10010-R2-1	07q32.1	CCGGCGCCCCCGAGAGCCGGCCCTCCCTTCCTCCGTGACAGGTGGCCCTGGAGTTGGGAAAGTTTGGAGCCGGCGAGGGG CGCCGGG	528
10030-R3-1	10q24.1	GGATGCAACCGTGAAGCCCGTGCCGTTGAGGATCTGCCACAGGCGGAAGGCAGCTGAGTTGACATCCACGGGCATCC	529
10138-L2-1	17q22	ACTTTTCTGGTGGGAGGGGAGAGCGGAGCAGGCTCACGTGTAACCGCAGGAGCCTCCTCTGGCTTGAGCCCTTTCTTTGGTA AGT	530
10145-L2-1	05p15.33	GGTCAGATAACAGAAAGAGAGAGCAAAAGGAAAAAGAAATTTTTGAAGATCAAAAAGTGGCTGTTCAATTTTGTATCTGACC	531
10175-L1-1	Xq13.2	CATCTTTGTGGCTCCTGGTTGCTAGGAGCAGGTGCTTCTGTACTAAGCAACAGGAGCCTGTTGGATG	532
10209-L3-1	02q33.1	TCAATTTGTTACATAGCTAACTTATTTTCTAATAGACTATGTTGGTAATAAGAAAAATGAATTACATGCTGTTGGCAGAGTGA	533
10231-L3-1 10231-R3-1	09p11.2	GGCGGCTGCGGAGGCTGGCGGGCTGCTGCACCTTTAACGCTTTCTGGCGCTGACAGGGCGGGCCAGCTAAAGTTTCACAG CGCC	534
10242-R3-1	22q13.2	GGCAGGAAGGCCCTCCGGCTTCACAAAGTGGCCCTGGGCATCCAGGAAGTGTTCCGGGTGGAAGCGGAAAGGCTTCTTCC	535
10333-L3-1	11q12.2	CGGGCAGGGGAGGGGTGGCAGGGCACAAGCCTCCCACTGTGCCGTGTCCCACTCCCGTCCCGTTCCTCCCG	536
10335-L3-1	15q26.1	TCTTAGTGCTTTCTAATTAATAAAGGAGGGGAGTGGTGATCTTTTTGCTCTCTAAGTTCTGTTTCTCTGAGTGGAAAGCAGAGGG	537
10342-R2-2 10342-L2-1	19q12	CCCACGCACGGAGGGTCGCCAGGAAAGTGACATTACCGCTTAATACTTCGAGATGCTCCGGCGCGGG	538
10366-R3-2	Xq22.3	GAATACCATTAATCTGTTCACTAGGAGATTAAATTTGCAATTTGTTGGCAATCACATGTGGGTCTTTGTAGCAAGAGCGTGGTGTT GTTT	539
10374-R3-2	16q12.1	CAGGGGATTTGTTACCGCTGATGTGTGGCCCCGTCGGAATGAAGGGGGCTTTTCATTAACAAAGTAGCGGGCGGTGTCATCTTCC CCTG	540
10533-R3-1	20q12	TAATTGCCTGAATCGCCGGTTACATATCTGTAGGAAATCTCTTGGAATATAAAGAAGGGGCTCAGGACAGTTA	541
11370-L4-1	12q13.3	GTCAGTTCTCAGGGGACAATACTGATGGCAGCCAAACTGGGCAAGGATGCAGTGTGGGGCGGAGGGGCATGACCTCTATT CAAGTTCTGTGCTTGGCCCCCTGGCTGAGGTATTGAGTGTGAGGAAGGGAACACTGGGC	542
12184-L4-1	03p13	TTGTACACAAATATTCGTCTGTGGTTGAAAGGGGGCACGCTGAGGTCAAGTGATGTAGTGTTTTCCATTTTCCATATGAGTCTCAC AGTGTGCGA	543
12223-L4-1	04q27	TAGAGGGAGAAGAAACAACCTCTGTCTGATGTCTTCTGGATGGCCTTAATACAGATAGCATTTGTCTCTTCCATTCTG	544
4315_C-L4-1	01q22	GGGAGGGGACCGGGAGGGCGGGCGGACCCCAACCTCTCGGAGGAGGGGCTGCCGCTCGCCGCTCCGCTCTTTG TTGTTGGGCTCCGCGCTCCCCCTCTCTCCCTCCTC	545

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
4315_D-R4-1	01q22	GGGACGTGGCCCTCCCCCGGAGCGGACTCCAAGAACTCCGGGGGCGCTGGGGCTGACTTTCC	546
4315_E-R4-1	01q22	TCCCCGCCACCCCTTGAGCACCTCAGCGTTCTTAGGGAAAGCCAGAGCCGGGAGGATCGGGGAATAGGTTTGGTGGGG	547
4315_F-R4-1	01q22	GATCAGGTTCCCTCCCCCGCATACACCTGGCGCAGGTGAAGCTCAGGAGCGGGTGGGAGCCCCGGGTT	548
4315_I-L4-1	01q22	CTGGACGGGGGGCGCCGAGGCCAGGGCGCCTGAGGGGCGCAGAGGTGTCAGCGTGCAACCGCCGCCCCACAGCGTTC CCGCCACCCAGCCACCCCTCAAAGCCCCG	549
4315_K-L4-1	01q22	CCGGGCTGGGGTGGAGGATGGCGAGGATTTGACAAATTTCAGGGCCCCCTGGGATCCTTTCCCTACTCCCTGGTCTTGTGG ACACCCTGTTTACCTGCCCTAAATTGCCCCGG	550
10010_B-L4-1	07q32.1	GGCAGCGCCCTCCTTCCGGAGGGCAGCATCCCGCGGGGCGGGGCTCGGCTTTGATGCCAGGGCACCTTTTGTCCCTG GAGACGCTCTGCCAGCCAGGTGCGTGGAGGGAGTGCAGCCC	551
10010_D-L4-1	07q32.1	CAGGTGAGAGGCTGAGCCCCGGGGGGAGGGCGCCAGGCCCTGGGGCATTAAACCGTCCCCGGGACCCCTTTTGGCCTG	552
7356_A-R4-1	08q24.3	CGGAGTCTGCTCGCTCGCTCGCTCGCTGACTCGCCGGAGCGCTCTGTGGCGGTGCGCGCAGGTGCGTCTCGCAGAGC GGGCTCTG	553
12722-L4-1	13q31.3	GCGGGCGGGGGGAGGTCGGAAGTACTTTTGTATGCTAATGAGGGAGTGGGGCTTGTCGCTATTACGTTGAGGCGGGA GCCGCCGCCCTTCATTACCCACATGCTCTCGAGGTGCCCGCGCCGCCGACCTGC	554
999999-R4-1	17q25.3	CCCTTGCACCTCCCGGATTGGCGGTCAGGGCCAGGGCCCCCTTGAGAGTCTGGGAATCCCTTCTCTGGGCTCGCTGGGT CCTGGCCAGGAAGGGCTGGGGTGACAAAGGG	555
999997-R4-1	17q25.3	TCTTCCTCACCCCTGCCACCCCTAGTCTCTTTATTGATTCAAAGGTTAAGGAAGCTCCTGGGGCTTGAGGGGTGGCACAG TTTTGGTGGGCCCAAGTGAGGA	556
8433_B-L4-1	17q25.3	ACGGGCGAGCAACCCCAATACTCAGCAACACAAAAATATGCCTCCGTGTGTGTGTGTGCGTGTGCGCTGCGCGT	557
8433_C-R4-1	17q25.3	AAATTAAGAAAAAAATTCACCAAAAGCGCCGAGTGACAGTTCACCAATTTCTGCCCTTCTCTCCCTCCCTGCACCA CTAGGAGGAGAAAGGCACACAAATTCGCTTTTAAATTTTTTTTTTTTGGTTT	558
8433_D-R4-1	17q25.3	CCCGGCTCGGCCCGCGTCTCTCCAGCTCCTCCGGCTCCTTTTAGTGATAAATTAGTGATGGCATTTCCCGGAGAGCGGAGCA CAACACAGGGCGCGGGCTCGGG	559
3758-R2-2	18q21.31	TGCAGGAAAAATCAGTAACATTAGTCATCTTAAAGGGTTATTACTCAAGATGATTTAAATTGTGAAAAATGTCAGTGGAGCCTGCA	560
3820-R3-1	16p11.2	GGCCTTCACCCGGGAGCCACCTTGTCTCCAGGTCTGGCCAGTAGTCTCCTGAGGCAGGGATGGCCCCACAGAGGGGTGGGG GCC	561
3851-R3-4	16q21	ATGGAGATTTCTAGCCCTTCTAAAGTCTATTAAAAACCTTTAAATTCCTGCTTTAGACAAAACATAAAGGAGGGTTGGAGCTCTGT	562
3874-L3-1	07q35	TGGGTGGCAGGTATTAGGGATAATATTCAATTTCTGAGCTTTCTGGGAGAGACTTGGTGACCATGCCAGCTCCA	563
3906-L3-1	03p24.3	AGCTACACCAAGATGAGGAATAAAGGGTTTACATATGTTCCAATTTCTAAACCTATCAATCAAAACCCATGTTATTCCTTTGGAGCT	564
3952-L3-2	01p36.32	CCCCGGCGAGGGGTGTCAGATTGAGTCTGTGCGCATGTGCGAAGGTGTCCAAACTGACAATGCTGGGG	565

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
3976-L2-2	14q11.1	GTCCACCTTCCAGCAGATGTGGATCAGCAAGCAGGAGTATGATGAGTCAGGCCCTCCATTGTCCACCGCAAATGCTTCTAGGTG GAC	566
4064-R3-1	11q23.3	ACATGGCTGAACAAGATAAGGGTTTTATTCTTGTGTTAGGGACGTGCTGGGGCTGGGATGGAATTCAGCTATGT	567
4118-L2-2	02q21.1	GTCCACCTTCCAGCAGATGTGGATCAGCAAGCAGGAGTATGATGAGTCAGGCCCTCCATTGTCCACCGCAAATGCTTCTAGGTG GAC	568
4130-L3-1	17p11.2	GGGTGCTGGTGGACATGGACGGCGTGGCTGGCTGACTTCGAGGGCGGATTCCCTCAGGAAAGTTCCGCGCGGCTTTCCCGACCAG CCCT	569
4155-R1-1	03p12.3	TGACGACTGGCCCCGCCCTCTTCCCTCGTCCCATATTGAACTCGAGTTGGAAGAGCGAGTCCGGTCTCA	570
4182-R2-2	02q23.3	AGTTCTTGTAGTCCACATCGCTGACTAAGGTCTGGCACTTCTTGGCCAGCACACCCCCAGCATGTCCACTGGGCTGCTGAACT	571
4216-R3-1	04q24	TCCTTCCAAACCCAAATGATTATATGATTAAGATACCAGTACCCAGGGGAATTGATCTGTGTATAAGGGAGGGAAGAGA	572
4340-R3-1	10q22.1	GCTCTGCTTCCAGGCTGATTTTTAGGCTGGCATTTAGGTTTGGCCTGGGACAAAGGGCTGGAAAATGCAAGGC	573
4391-R2-1	Xp11.3	TAAGCCAGATTCTCAACTTACGTAATTCATGGGAATTCACAGAGTTATATTGGTTGAGAAATTTGGCTTA	574
4413-L3-1	11q23.2	ATTTGTTTACTGTAGCTGGAGGTGCCGAGTAATGAAATGCACCTTGTGTCTGCAGCTCACTGCTAAACAAAT	575
4417-R1-1	14q13.2	GCTGGGTTTCATCGGAGAAACTCCCTGCGATGAGCCACTAGGGTCAACGACAGGGAACCTTTTGTGATGAGCGCCGAGT	576
4498-L3-2	06p22.2	TTCCCCAGGCAGCAGCGCGCACGGCCGCTCTGGATCTCCCTGGAGGTGATGGTCGAGCGCTTGTCTCATATATGCGCCAGGCGGG A	577
4567-L1-1	Xq26.2	GGGAGTGTGCTGGAACTGGGCAGATAAAAAGGGCTGGAACTTATTTTGGCTAGACTCTCAATTCCC	578
4579-L3-2	06q14.1	AGCACAGCTATTGGTGTTTTGCAGGAGGCAAGTGAAGCCCACTCTGGTTGGCCTTTTTGAAGACACCCAAATGTGTGCT	579
4610-R3-1	08p12	GCCCAGTTAATTGGTCTCTCAACCTACATTAGCTGTTGCAATTGCAGCCAAATTAGGCAGGGGCCAGAGGGC	580
4724-L3-2	15q24.2	ACAGACCAGGCACAGAAAAGCTGAAGGTGCTAGAGATGCCAAAGTGAAACATGTGCATTTTTTGGTAACTGTGTACTTCTGTGACTG T	581
4754-R3-2	04q25	CCACCAACTTGACATACACAGGCTCATCACAGTTGGATGCAAGCACACAAAGATGGGCTTGGCAGCTTGCTAAAGGCTTTGG	582
4801-L3-1	01q32.2	ATTAATGTTTTGTAGCAAAACAGGAGGCAGAGTTCTCCAAAGGCTCTCATCTCTGTGCTTCCAGAAAAATATTGAT	583
4964-L3-2	02p14	TTGGGGAATTTGGTGGTTTTCAGAAAGTTAGCCCTTCTGGTGTAGGGTTTGTGATTTTCGATACACACAGATTATACTCGTCCCAA	584
5071-R2-1	17q12	GATTCCTGCTCCAGAGCCATAAAGTGGGAGCCCCCATTTATTAATTGGGCTGGGACTGGGGCGGGGGTC	585
5306-L3-2	12q13.13	GGACCCCTAGAAAGGGCCAGAGCTGGGGTCAGAGGCCACCTCTCCATTCTCTGCCCTGCTCTGCTGGGTCC	586
5327-L3-1	09q31.3	GGAGACAGGACATAGTCCCAGAGGTTGAGCTGGCTTATGGAGCCCCACAAAAGACTCAGCTGGGCTGAATCCCTCTCC	587
5372-R3-2	10q22.2	TGTTTTCCCTGTGGAAAAGATTACTCTAGGCAAAATTTAGAAAAATGCTTTTAAAAAGAAATCTTGTGTAAAGTTGAAAAACAGAGGGAAAG	588

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
		A	
5380-R2-2	1p36.22	GGCCCTGAGAGCAATACTCATATTGATTGCATTTATTTCACTCTAGGGAGGAGAGATAAATCTTCAATGTGGGACACATCTGGGAG CC	589
5441-L3-2	1q24.1	GTAGCTGCTGCTCTGTTAGCTATACTGACTTGGAGCTTGGCTGTAGGATCAAGTTTGGGGTGGGTAACTAGTGGGAGGCAGCTG C	590
5474-L3-2	9p13.3	GCTAGGTAGGTGGGTACAGAGTCTGGCCAGGTGAGAGGACCCCACTGCTTGGCCCTGACTCTGCCCCCTGGACACACCTTC TTCAGT	591
5513-L3-1	8p12	AAGTGGCTCTGAGGCATCATGGAGAACAGTTGAAAATCACATTGTCAGCTCGAAATGCATCAAAAGCTATTT	592
5598-R2-2	11q13.2	CTCCACGGCCTGAAGCTGCTGCCAAGCTATTTTGGTTCTGCACAGTTAAAAATAGCTTTCACGGAGGTGGGAG	593
5618-R3-1	8q21.11	TCCCCAACCCCATCCATTAGGCCAGCAACGCTTGTAGAGCTCACTGTGGGCTGTAATGTGCACTGGTGGCTGGGACACCCAGG GA	594
5619-L3-1	15q25.3	CTCATTGAGGGAAGATTGAGCAGAACTGGCATTGCTTTCGTCAAATTGATTGTGCCCGTCTGTTTGATCCAATTCAGTGAG	595
5733-R3-2	16q22.1	CTCTCCAGCCAGCCTGAGTCCTTGTGTATCGTGAATGGTGGGACTGAGAAGCAGGATGAGCTGGTGAG	596
5735-L3-1	5q31.1	CCCCCACAGGCTTGAGGCCAGAGGAACAGCAACTTCTTCGCTGGGAAAGTGTTGTGGGGCTCAAGCATTTGGGGG	597
5863-L3-1	9p13.3	GACATCTTCCTGGCGACAGGAATTGCCAGCAGCAACGGCCTTGCCGTTGCCATGGTGATACCCCTTGTCATTCGGATGAAGTC	598
5919-L3-1	7p21.3	TCTAGGGAGATAAAGTGACAGTGTTGCTTGTTCAGGCTATTGTTCAAAGAAGCACGTCCTTTTATTATAGA	599
6026-R3-1	18q22.3	TTGGAGGCAGAAAGCTCTGCCCCCTATGGTGGTGGAAAGCCAGTACTTTGCCAAACCACAAAACCAAGGTGGCTGGGTGTTCCA G	600
6218-R3-1	2p16.1	TTCTATTAATATGTATCATATTGGTATCCATTTTCACATAATGTGACATGAGAGGCAGTAATGTGATTACTTTTTTAGTGAA	601
6253-L3-1	4q31.23	GCTGGCATTATGGCAGCCAGGATATTGTCAGGAGGAACCTTGGCAGTGTCTCGTAGGTGTTATGATGCTGGC	602
6355-R3-1	2p14	AGCCTCCTTGTCTGGAGATTCTAACAAAATGGGTTTGCAGTGTGGAAACTGTAAATCCTCAGGGAACAGATGATCTGGACAATGTA GCT	603
6421-R3-2	11q13.1	CCCTCCTTGATCTGGGAACTATTTCTTCATTCCAGAAGGTCAGAGCTCTGGCAATAGTGTCTTCAGACTCTCACAGGAGGG	604
6450-R3-2	12q13.2	CTGCAGTGGGAAAGTCAAGCCTGGTATTACGTTTTGTCAAAGGGCAGTTCCTATGAGTACCAGGAGAGCACCATTTGGAG	605
6478-R2-2	2q22.3	GCTGGATTCTGCCCTTGGATACACACACAAAACCCCAATTGAAGTCAATGGAAATTGTGCATGCATATCCAGGGGCAGAAATTTG GC	606
6554-L3-2	15q26.1	TCCCAGAATATGGAGGCACCAAAAGTAGTTCTAGATGACAAAGGATTATTTCCCTATTTAGAGATGGTGACATTTCTTGA	607
6647-R2-1	1q23.3	CTCAGTATCTTCAGCTTGGGAAACTGACCTCGTTAATTTTAATGAGGGGAAAAAATCTCCAGCTGGGGCTGAG	608
6664-R2-1	14q31.1	GCACATTATAAACTCTAATTCATTAACGTCATCATAAATGGTAATGTCCTGTGAAAAAGAGAGGTGGTGGC	609

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
6712-L2-1	9q21.13	CAGGCTGACAACTGATATCACAAAGACCACAGCTAAGAGTGGTTTATTACTTTTAGTGGATTATTTAATTCAGTCTCACAGCCTG	610
6718-L3-2 6718-R3-1	3p22.1	CTTTGGAGGCAGAAAGCTCTGGCCCCCTATGTTGGTGGAGGCCAATACTTTGCCAAACCACAAAGGTGGCTATGGCGGTTC CAGTAG	611
6912-L3-1	13q22.3	AGAAATTAAGTCTGGACCCAGAGGCTGAAGGCACGTTGAAGGAATCCTTCAATTTCTTGAGCCCCTTGTTTGAAGGCTAGTGATTT T	612
7019-R3-1	18q22.3	TTCAAAAGAAATTACTTAGCAATTAAACCTTCAGCAAAATTGTAATTCATTTTAAATTTAGCACGGAGGTTTGATGCTGTTTCTGAGGA A	613
7070-R3-1	14q12	TCTCCTTCTTTCAAAACATGTCAGTGACCAATTTCTACAGTAAGTTTAGTGGTCTGCTGACAGTATAGGTTGACTTGAAGTA GA	614
7089-R1-1	6q16.2	ATGCACTCTGCTGTCAATTTGCAGCCTGACCTGCAGGTCAGGGGTGATGTGAGCTGGCTCAGGGATGCAT	615
7158-R3-1	3q13.31	ATTCAACACAGATTACAGGTGCTCTCAACAGCCCATGAAAATATATGGCTGAAGAGGGAATCTATAAATGTAATGAAT	616
7292-L3-4	1p34.1	GCAATTAGAATGCAGGGAGGTTTCAGAAGCTATTTAACTGGGTGACCCCTGAGGTCGCTGCATCTGACTCCCATCCCTGGATAAAAT ATTGT	617
7304-L3-1	1q24.2	CAAATGTGGGAGCTTGGATCAATGTTGAAGAATAATTTTCATCATAGTGAAAAATGTTGGTTCAAATAAAATTTCTACACTTG	618
7340-R3-1	6q16.1	GCTCACATGGGAATCCAAAACCTTCTAAATTTCTATACAGATTCAATTCAGAAATCAATTAGAGTTGGTGGCAGCCTTCTCCCTGTGT GC	619
7375-L3-1	6q23.3	CAGGTATGTAAGTGTGGAGACAGCAGCGCTGCAGAAAGAGCCCTCCTGATCAGAAAGAATGTCTGTCCCACAGTTGACTCTCTG	620
7435-L3-2	6q16.1	GTAGGAAAACACAGCTAATGGGGCTTAGCAACAGATGGGGAGCCAGCACAAAAGCTGTAGGTTCTTCTCCTCGCTCGGCTGTGACT TCCTGC	621
7543-L3-2	1p36.22	TCATCTCGGGCCCTGGTGAATTAGGACGACATCGGCATTTTTTATTGCTAAAGACGTCCAGATTGATCCAGCCCCTTGGCTGA	622
7597-L3-1	11p14.1	GGGATTGATCTGAGGGACACAGAGCCCCAGGTTCCATTAAAGTGTATTAATGCCAAGGTCTGTGCAATTAGTGAAAAAGTCAATTCC	623
7763-R3-1	8q11.21	GGCTTGGGCTTTGTTGGTCTTCCACTGCTCTGCTACATCATTTGCTAATGGATCGTCTGGATTGGGAGCACATAACAAGGCCTGG GTT	624
7824-R3-1	6q16.2	CCTGGATGCTGTTTCATTATGTAGAGTCAGGCAAAAGACAGACGGATGTGTGTGAGGCGGCGATGAAGCTGGCACCAGG	625
8075-L3-1	10q22.1	CAGCTGGCCTGGTGCCCTGGTGCCTGGAGGTGTAGCTGGGCTCTGACCCAGCTCCTCAAACAGGTTCCATATGGCCCCTCCCCGG CTG	626
8336-R3-1	Xq13.1	CCTCCCCAAGCAATCTGCTCACCCACTTGTGTGCTGACTCTGGCTGGGGGTGTGGGACTGAGTGGGGAGG	627
8434-R3-1	9q34.11	AGGAGGCCCTGAGTGTCTAGTAGCCACTTATAGAAAAGTCGCTTCCAGCAATTTCTTCTAGGAGTGGAGCCTCAGGGCTGCCT	628
8552-R3-1	13q21.33	CTCCTAGTGGTATCTGTGCATAAAGATACTGTTCAGGGCATAATTTTCATTATATTTTATAGTGTCCTCTTTATGGTGACCCCTGGGAG	629

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
8685-L3-1	17p13.3	GGGAGACATTCCACTTATAGGAGGCCAAAGAGCAGGATATCTGCACAGGAAGAGTTTCATCTTATATGACTTTTCGGGGATGGATTGTCTCCT	630
8719-L3-2	5q22.3	AGAGCGCAGCTACTGGTGCTTTGCAGGAGGCCAAGTGAGGCCCTATCTGGTTGGCCTTTTGAAGACACCAACCTGTGCGCT	631
8760-L3-1	3q27.3	GGTCATGCTATGCCTGGGTGAGTTTTGCCTCGGGCTCCAGCGGCACCTGGAGCAGAGCTGAGCTAATTTTACCCACAGGAGCTGACT	632
9092-R3-2	18q22.3	CTTCTTATAAGTCTGTGGTCACAGTAIGTTTTTGCAGTTCAGCACTCTGGCTCTGTGGGATACTGATGGCTTCTAGGAAG	633
9557-R3-1	22q12.1	TGCCCTGCTCTGGCCAGAACCAAGCACGTCAGCTTTTTTATTAGGGCAAGAGGTGCAGAATGGACTGGGCCAGTCCAGGCA	634
9582-R3-2	5q31.1	TTTTACACATTGTCAGCTGCATCTATTAAATTTTCTTACTGTCTCAGAGACAGTTAAGGGAGCCCTTTGTGAAACATCTGAGGATTTGTAAA	635
9688-L2-1	Xq26.2	TGCAGCAGAACAGTCTGGGGCCATTTAGCTTAGGGGCAATAGTTCCTCATACTTCAAAGAGCCCTAAGGACATTGCTGCA	636
9694-R3-1	7q21.2	TTATTATGCAACATCTGCTCTGAATTTCAAATGTGATAAACAGAGTATGAGTGTGCGTGGCAGATGGCTAATGA	637
9747-L3-1	18q11.2	GAGCTGCTTTGAATTGCTCGCAGTTTGCCGGAGGCGGTGTGCTGGTTGGACGCTCCGGGAAACAAAAGCAACCCAAAACAGCTC	638
9772-L3-1	10p15.2	TCAGGAAAGGTCATGGGAAAAATGCTGTAGAAATTTTCTAATGGTTCATCCATCAAAAAATGCAGCTCTCCATTGATTCCTCCCGA	639
9798-R3-2	5q12.3	GTGCTTTTCTTTCCCCCAAAGAAAGTCATACCAGGTATATATAGAGAGATCTATAATGCCCTTCTGTGGGGAATGAAAGCAC	640
9812-L3-1	5p13.3	AGCAAGCTGGGGAGCCAGATAAATAGAGCTTTCTGTTTCTTCTGGAGTCTAAAATATCTGATCTGGAGGTTCCCTCCCTGCT	641
9813-R3-2	12p13.32	GAAAGTGTGGGCCCTTTTGAAATTGTCAGAGAAACTGATTTTCTTGTCAGTCTCATCTGTACAGTTGGAGGGCCAAATTTCTACTTTC	642
9987-R2-2	13q13.3	CAGGCATTGCCCTGGCTCAGAGACCTTTGCTGCTGAGCAAACTGAGTTAACAAAAGTGGGCTGAGGGGTGAAGCCCTG	643
10093-R2-2	2p11.2	TGTTTCCAGCATTTCCAGGTAGGCCAAGGTGTCTCTACAGAAAAACCTTGGGTTAGACCTACAGGGGTCTGGCTGGTGTAAACA	644
10120-R3-1	7p14.1	CAAGTTTCTGAGCCCTGGGTTGATATACTGTACCATCTTTTGGTGCAGCTGTAATTTTAAACAAGGGGAACCGACTTG	645
10133-R3-1	Xp11.4	CTCTTCCCATATTTAATGTGGCTCTAATTTCTGACTTCATTGCAGAGCGGTGAAATCAAAATTAGGAAATGGGAGGAG	646
10154-R1-1	6p12.2	ATTCCAAAGGCTCTGACACTTGCTAAATCCTGTAGAAATGAGATCAAGGGCAAAAGGGCACAGATCCTGGGAAT	647
10198-R3-1	2p16.1	GAGCAGTTCTTTGTTGCCCATCACATTTTAGTGCAGGGAAGTCTTTGCCCTAAAGGTGCCAGGTAAAGAGAATTGCC	648
10260-L3-1	22q13.1	GTGGGAGACATCGGGTGGGGCCCTGGCGAAACAATAGGTGGGCCCCAGCTGGGGCCCCCTCCTGCCTGCCTCACCCGC	649
10346-R3-2	10q25.3	TGGGCAGAGCCAGGTCTGCACCCCTCGGAGCCCTGGAGGCCAGGTGAGCTGCTGCAGAGGGCACAGACAGGCTCATACTCA	650
10539-R3-1	2p22.3	TAGATGGGAATTTGAGTGGCATTGTTTCTCAGGCCCTGCTGGAGGCTGGGGGATAGTAAATTTTCTCAAAATACCATTTA	651
10543-R3-1	1p31.1	GCCTTAACCTTTTATCATCTTCTTGATTATGTCACCTGAATTATTAATTCATGAGCCAGGATGGGAAGGGTGAAGGC	652

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
10553-R1-1	1q24.2	GCTGGCTCCTGCTCAGAGTGACAGCACCCCTGTGGAGTCTGCTGGTACTGACCTAACCACCACAGAGCCCCACTGGAGCATGGAG CCAGC	653
10562-L1-2	8p12	TGATCAGGCGCTTGATATGACAGGCTGTGGAGTCAGGACCCCTTGAACAGAGTGGGCCCTTGCTGTACAGTCCGCCCTGACA	654
10594-L3-1	3p14.2	CTGTTGAGTGGGTGTTGGGTAAACAATAATGTTGTTTCATAAATATGTGCTGGCAGATTAGCACACTTCTCTACAG	655
10639-R2-1	2q31.2	TGACGACTGGCCCCCGCTCTTCCTCTCGGTCCCATATTGAACCTCGAGTTGGAAGAGGCGAGTCCGGTCTCA	656
let-7a	09q22.32	TGGGATGAGGTAGTAGGTTGTATAGTTTTAGGGTCACACCCACCACCTGGGAGATAACTATACAATCTACTGTCTTTCTCA	657
let-7a	11q24.1	AGGTTGAGGTAGTAGGTTGTATAGTTTAGAATTACATCAAGGGAGATAAAGTGTACAGCCTCCTAGCTTTTCCT	658
let-7a	22q13.31	GGGTGAGGTAGTAGGTTGTATAGTTTGGGGCTCTGCCCTGCTATGGGATAAAGTGTACATACTACTGTCTTTTCCT	659
let-7b	22q13.31	CGGGTGAGGTAGTAGGTTGTGGTTTCAGGGCAGTGATGTTGCCCTCGGAAGATAAAGTGTACATACTACTGCTTCCCTG	660
let-7c	21q21.1	GCATCCGGTTGAGGTAGTAGGTTGTATAGTTTAGAGTTACACCCCTGGGAGTTAAAGTGTACAAAGTGTCTTAGCTTTCCCTGGAGC	661
let-7d	09q22.32	CCTAGGAAGAGGTAGTAGGTTGCATAGTTTAGGGCAGGGATTTTGCCACAAAGGAGGTAAGTAACTATACGACCTGCTGCCTTTCTTA GG	662
let-7e	19q13.41	CCCCGGCTGAGGTAGGAGTTGTATAGTTGAGGAGGACACCCAAAGGAGATCACTATACGGCCTCCTAGCTTTTCCCCAGG	663
let-7f	09q22.32	TCAGAGTGAGGTAGTAGATTGTATAGTTGTGGGGTAGTGATTTTACCCTGTTTCAGGAGATAAAGTGTACATACTACTGCTTCCCTG A	664
let-7f	xp11.22	TGTGGGATGAGGTAGTAGATTGTATAGTTTAGGGTCATACCCCATCTTGGAGATAAAGTGTACATACTACTGCTTTCCCCACG	665
let-7g	03p21.2	AGGCTGAGGTAGTAGTTTGTACAGTTTGGGGTCTATGATACACCCGGTACAGGAGATAAAGTGTACAGGCCACTGCCCTTGCCA	666
let-7i	12q14.1	CTGGCTGAGGTAGTAGTTTGTGCTGTTGGTGGGTTGTGACATTGCCCGCTGTGGAGATAAAGTGTACGCAAGCTACTGCCCTTGCTA	667
miR-100	11q24.1	CCTGTTGCCACAAACCCGTAGATCCGAACCTTGTGGTATTAGTCCGCACAAGCTTGTATCTATAGGTATGTGTCTGTTAGG	668
miR-1224-5p	03q27.1	GTGAGGACTCGGGAGGTGGAGGGTGGTGCCCGCGGGCGCTGTTTCAGCTCGCTTCTCCCCCACCTCTCTCTCCTC AG	669
miR-1225-5p	16p13.3	GTGGGTACGGCCCGAGTGGGGGGAGAGGGACACGCCCTGGGCTCTGCCCAGGGTGCAGCCGGAGTGAAGCCCTGTGCC GCCCCAG	670
miR-1228*	12q13.3	GTGGCGGGGGCAGGTGTGTGGTGGTGGCTGCCCTGCCGTGAGCAGGGCCCTCACACCTGCCCTGCCCCCCCAG	671
miR-125a-5p	19q13.41	TGCCAGTCTCTAGGTCCTGAGACCCCTTAAACCTGTGAGGACATCCAGGGTCACAGGTGAGGTTCTTTGGGAGCCCTGGCGTCTGG CC	672
miR-125b	11q24.1	TGCGCTCCTCTCAGTCCCTGAGACCCCTAACTTGTGATGTTTACCGTTTAAATCCACGGGTTAGGCTCTTGGGAGCTGCGAGTCGT GCT	673

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
miR-125b	21q21.1	ACCAGACTTTTCCCTAGTCCCTGAGACCCCTAACTTGTGAGGTATTTTAGTAACATCACAAAGTCAGGCTCTTGGGACCTAGGCGGAG GGGA	674
miR-126 miR-126*	09q34.3	CGCTGGCGACGGGACATTATTACTTTTGGTACGCGCTGTGACACTTCAAACCTCGTACCGTGAGTAATAATGCGCCGTCACACGGCA	675
miR-135 miR-135a*	03p21.1	AGGCCTCGCTGTTCTCTATGGCTTTTATTCCCTATGTGATTCTACTGCTCACTCATATAGGGATTGGAGCCGTGGCGCACGGCGG GGACA	676
miR-142-3p	17q23.2	GACAGTGCAGTCACCCATAAAGTAGAAAAGCACTACTAACAGCACTGGAGGGTGTAGTGTGTTTCCCTACTTTTATGGATGAGTGTACTGT G	677
miR-145	05q32.1	CACCTTGTCCTCACGGTCCAGTTTTTCCAGGAATCCCTTAGATGCTAAGATGGGGATTCCTGGAAATACTGTTCTTGAGGTCATGG TT	678
miR-146b-5p	10q24.32	CCTGGCACTGAGAACTGAATTCATAGGCTGTGAGCTCTAGCAATGCCCTGTGGACTCAGTTCTGTGCCCGG	679
miR-149 miR-149*	02q37.3	GCCGGCGCCCGAGCTCTGGCTCCGTGCTTCACTCCCGTGCTTGTCCGAGGAGGGAGGACGGGGGCTGTGCTGGGGC AGCTGGA	680
miR-150 miR-150*	19q13.33	CTCCCCATGGCCCTGTCTCCCCAACCCCTTGTAACAGTGCTGGGCTCAGACCCCTGTACAGGCCCTGGGGGACAGGGACCTGGGGA C	681
miR-155	21q21.3	CTGTTAATGCTAATCGTGATAGGGGTTTTTGCCCTCCAACCTGACTCCTACATATTAGCATTAAACAG	682
miR-16	13q14.2	GTCAGCAGTGCCTTAGCAGCACGTAATAATTGGCGTTAAGATTCTAAAATTATCTCCAGTATTAACTGTGCTGCTGAAGTAAGGTT GAC	683
miR-16	03q25.33	GTTCCACTCTAGCAGCACGTAATAATTGGCGTAGTGAAATATATATTAAACACCAATATTACTGTGCTGCTTTAGTGTGAC	684
miR-181c	19p13.12	CGGAAAAATTTGCCAAGGGTTTGGGGGAACATTCAACCTGTCGGTGAGTTTGGGCAGCTCAGGCAAAACCATCGACCCGTTGAGTGG ACCCTGAGGCCTGGAATTGCCATCCT	685
miR-198	03q13.33	TCATTGGTCCAGAGGGGAGATAGGTTCCCTGTGATTTTCCCTTCTCTATAGAATAAATGA	686
miR-199a-3p	19p13.2	GCCAAACCAGTGTTTCAGACTACCTGTTCAGGAGGCTCTCAATGTGTACAGTAGTCTGCACATTGGTTAGGC	687
miR-199a-3p	01q24.3	AGGAAGCTTCTGGAGATCCTGCTCCGTCGCCCCCAGTGTTTCAGACTACCTGTTTCAGGACAAATGCCGTTGTACAGTAGTCTGCACAT TGGTTAGACTGGGCAAGGGAGAGCA	688
miR-199b-3p	09q34.11	CCAGAGGACACCTCCACTCCGTCCTACCCAGTGTTTAGACTATCTGTTTCAGGACTCCCCAAATTGTACAGTAGTCTGCACATTGGTTA GGCTGGGCTGGTTAGACCCCTCGG	689
miR-19b	13q31.3	CACTGTTCTATGGTTAGTTTTGCAGGTTTGCATCCAGCTGTGTGATATTCTGCTGTGCAAAATCCATGCAAAACTGACTGTGGTAGT G	690
miR-19b	Xq26.2	ACATTGCTACTTACAATTAGTTTTGCAGGTTTGCATTTTCAGCGGTATATATGTATATGTGGCTGTGCAAAATCCATGCAAAACTGATTG TGATAATGT	691

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
miR-200b	01p36.33	CCAGCTCGGGCAGCCGTGGCCATCTTACTGGGCAGCATTGGATGGAGTCAGGTCTCTAATACTGCCTGGTAATGATGACGGCGG AGCCCTGCACG	692
miR-200c	12p13.31	CCCTCGTCTTACCCAGCAGTGTTTGGGTGCGGTTGGGAGTCTCTAATACTGCCGGGTAATGATGGAGG	693
miR-205	01q32.2	AAAGATCCTCAGACAAATCCATGTGCTTCTCTTGTCTCTTCCACCGGAGTCTGTCTCATACCCAACAGATTTTCAGTGGAGTGA AGTTCAGGAGGCATGGAGCTGACA	694
miR-21	17q23.2	TGTCGGGTAGCTTATCAGACTGATGTTGACTGTTGAATCTCATGGCAACACCAGTCGATGGGCTGTCTTGACA	695
miR-23a miR-23a*	19p13.12	GGCCGGCTGGGGTTCCTGGGGATGGGATTTGCTTCCTGTCAAAATCACATTGCCAGGGATTTCCAACCGACC	696
miR-23b	09q22.32	CTCAGGTGCTCTGGCTGCTTGGGTTCCCTGGCATGCTGATTTGTGACTTAAGATTAAATACATTTGCCAGGGATTACCACGCAAC CACGACCTTGGC	697
miR-24-1	09q22.32	CTCCGGTGCCTACTGAGCTGATATCAGTTCTCATTTTTACACACTGGCTCAGTTCAGCAGGAACAGGAG	698
miR-24-2	19p13.12	CTCTGCCCTCCCGTGCCTACTGAGCTGAAACACAGTTGGTTTGTGTACACTGGCTCAGTTCAGCAGGAACAGGG	699
miR-25*	07q22.1	GGCCAGTGTTGAGAGGGCGGAGACTTGGGCAATTGCTGGACGCTGCCCTGGGCATTGCACCTTGTCTCGGTCTGACAGTGCCCGGC C	700
miR-26a-1	03p22.3	GTGGCCTCGTTCAAGTAATCCAGGATAGGCTGTGCAGGTCCCAATGGGCCTATTCTTGGTTACTTGCACGGGGACGC	701
miR-26a-2	12q14.1	GGCTGTGGCTGGATTCAAGTAATCCAGGATAGGCTGTTTCCATCTGTGAGGCCCTATTCTTGATTACTTGTCTTGAGGCGCAGCT	702
miR-26b	02q35	CCGGGACCCAGTTC AAGTAATTCAGGATAGTTGTGTGCTGTCCAGCCTGTTCTCCATTACTTGGCTCGGGGACCCGG	703
miR-27a	19p13.12	CTGAGGAGCAGGGCTTAGCTGCTTGTGAGCAGGGTCCACACCAAGTCGTGTTACAGTGGCTAAGTTCGCCCCCCCAG	704
miR-27b	09q22.32	ACCTCTCTAACAAGGTGCAGAGCTTAGCTGATTGGTGAACAGTGATTGGTTTCCGCTTTGTTTCACAGTGGCTAAGTTCTGACCCCTG AAGAGAAGGTG	705
miR-298	20q13.32	TCAGGTCTTCAGCAGAAAGCAGGGAGGTTCTCCCAAGTGGTTTTCCCTTGACTGTGAGGAACTAGCCTGCTGCTTTGCTCAGGAGTGA GCT	706
miR-29a	07q32.3	ATGACTGATTTCTTTTGGTTCAGAGTCAATATAATTTTCTAGCACCATCTGAAATCGGTTAT	707
miR-29b-2	07q32.3	CTTCAGGAAGCTGGTTTCATATGGTGGTTTAGATTTAAATAGTGATTGTCTAGCACCATTTGAAATCAGTGTTCTTGGGGG	708
miR-29b-1	01q32.2	CTTCTGGAAGCTGGTTTCACATGGTGGCTTAGATTTTCCATCTTTGTATCTAGCACCAATTTGAAATCAGTGTTTTTAGGAG	709
miR-29c*	01q32.2	ATCTCTTACACAGGCTGACCGATTTCTCCTGGTTCAGAGTCTGTTTTTGTCTAGCACCATTTGAAATCGGTTATGATGTAGGGG GA	710
miR-30a miR-30a-3p	06q13	GCGACTGTAAACATCCTCGACTGGAAGCTGTGAAGCCACAGATGGGCTTTCAGTCGGATGTTTGCAGCTGC	711

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
miR-30b miR-30b*	08q24.22	ACCAAGTTTCAGTTTCATGTAAACATCCTACACTCAGCTGTAAATACATGGATTGGCTGGGAGGTGGATGTTTACTTCAGCTGACTTG GA	712
miR-30c miR-30c-1*	01p34.2 06q13	ACCATGCTGTAGTGTGTGTAAACATCCTACACTCTCAGCTGTGAGCTCAAGGTGGCTGGGAGAGGGTTGTTTACTCCTTCTGCCA TGGA	713
miR-30c-2	06q13	AGATACTGTAAACATCCTACACTCTCAGCTGTGGAAGTAAGAAAGCTGGAGAAAGGCTGTTTACTCTTTCT	714
miR-30d	08q24.22	GTTGTTGTAAACATCCCCGACTGGAAAGCTGTAAAGACACAGCTAAGCTTTTCAGTCAGATGTTTGCTGCTAC	715
miR-30e	01p34.2	GGGCAGTCTTTTGCTACTGTAAACATCCTTGACTGGAAAGCTGTAAAGTGTTTCAGAGGAGCTTTTCAGTCGGATGTTTACAGCGGCAG GCTGCCA	716
miR-320a miR-320	08p21.3	GCTTCGCTCCCCCTCCGCCCTTCTCTTCCCGGTTCTTCCCGAGTCGGGAAAAGCTGGTTGAGAGGGCGGAAAAGGATGAGGT	717
miR-331-3p miR-331	12q22	GAGTTTGGTTTTGTTGGGTTTTGTTCTAGGTATGGTCCCAGGGATCCCAGATCAAACCAGGCCCTGGGCCCTATCCTAGAACC CCTAAGCTC	718
miR-371-5p miR-371	19q13.42	GTGGCACTCAAACCTGTGGGGGCACCTTCTGCTCTCTGGTGAAAGTGCCGCCATCTTTTGAGTGTTAC	719
miR-373*	19q13.42	GGGATACTCAAAATGGGGCGCTTTCCTTTTTGCTGTACTGGGAAGTGCTTCGATTTTGGGGTGTC	720
miR-375	02q35	CCCCGCGACGAGCCCCCTCGCACAAACCGGACCTGAGCGTTTTGTTTCGGCTCGCGTGAGGC	721
miR-423-5p	17q11.2	ATAAAGGAAGTTAGGCTGAGGGGCAGAGAGCGAGACTTTTCTATTTTCCAAAAGCTCGGTCTGAGGCCCTCAGTCTTGCTTCCT AACCCGCGC	722
miR-424	Xq26.3	CGAGGGGATACAGCAGCAATTTCATGTTTTGAAGTGTTCTAAATGGTTCAAAACGTGAGGCGCTGCTATACCCCTCGTGGGGAAG GTAGAAAGTGGG	723
miR-483-5p	11p15.5	GAGGGGAAGACGGGAGGAAAGAGGAGTGTTCCATCACGCCCTCCTCACTCCTCCCGTCTTCTCCTCTC	724
miR-486-3p	08p11.21	GCATCCTGTACTGAGCTGCCCGAGGCCCTTCATGTCTGCCAGCTCGGGCAGCTCAGTACAGGATAC	725
miR-491-3p miR-491-5p	09p21.3	TTGACTTAGCTGGGTAGTGGGGAACCCCTTCCATGAGGAGTAGAACACTCCTTATGCAAGATTCCCTTCTACCTGGCTGGGTTGG	726
miR-513a-5p	Xq27.3	GGGATGCCACATTCAGCCATTTCAGCGTACAGTGCCCTTTCACAGGGAGGTGTCATTTATGTGAACATAAAATATACCTTTCT TGAGAGGGTAATGTACAGCATGCACTGCATATGTGTGTCC	727
miR-513a-5p	Xq27.3	GGATGCCACATTCAGCCATTTCAGTGTCAGTGCCCTTTCACAGGGAGGTGTCATTTATGTGAACATAAAATATACCTTTCT GAGAAAGGTAATGTACAGCATGCACTGCATATGTGTGTCC	728
miR-513b	Xq27.3	GTGTACAGTGCCTTTCACAAAGGAGGTGTCATTTATGTGAACATAAAATATAAATGTACCTTTTTGAGAGGAGTAATGTACAGCA	729
miR-516a-5p	19q13.42	TCTCAGGCTGTGACCTTCTCGAGGAAAGAACACCTTCTGTTGTCTGAAAGAAAAGAGTCTTCTCCTTCAGAGGGTTACGGTTT	730

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
		GAGA	
miR-550	07p15.1	TGATGCTTTGGCTGGTGCAGTGCCTGAGGGAGTAAGAGCCCTGTTGTTGAAGATAGTGTCTTACTCCCTCAGGCACATCTC CAACAAGTCTCT	731
miR-550	07p14.3	TGATGCTTTGGCTGGTGCAGTGCCTGAGGGAGTAAGAGCCCTGTTGTTGTCAGATAGTGTCTTACTCCCTCAGGCACATCTC CAGCGAGTCTCT	732
miR-557	01q24.2	AGAAATGGGCAAAATGAACAGTAAATTTGAGGGCCTGGGGCCCTCCCTGCTGCTGGAGAAAGTGTTCACACGGGTGGCCCTTGCTTT TGAAAGGAGGTGGA	733
miR-575	04q21.22	AATTCAGCCCTGCCACTGGCTTATGTCATGACCTTGGGCTACTCAGGCTGTCTGCACAAATGAGCCAGTTGGACAGGAGCAGTGC CACTCAACTC	734
miR-612	11q13.1	TCCATCTGGACCCTGCTGGCAGGGCTTCTGAGCTCCTTAGCACTAGCAGGAGGGCTCCAGGGGCCCTCCCTCCATGGCAG CCAGGACAGGACTCTCA	735
miR-614	12p13.1	TCTAAGAAACGCAGTGGTCTCTGAAGCCTGCAGGGGCAGGCCCTGCACCTGAACGCCCTGTTCTTGCCAGGTGGCAGAAGGT TGCTGC	736
miR-630	15q24.1	AACCTAACATCATGCTACCTCTTTGTATCATATTTTGTATTCTGGTCACAGAAATGACCTAGTATTCTGTACCAGGGAAGGTAGTTC TTAACTATAT	737
miR-637	19p13.3	TGGCTAAGGTGTTGGCTCGGGCTCCCCACTGCAGTTACCCCTCCCTCGGCGTTACTGAGCACTGGGGGCTTTTCGGGCTCTGCGT CTGCACAGATACTTC	738
miR-638	19p13.2	GTAGCGGGCGCGGCAGGGATCGCGGGCGGGTGGCGGCCTAGGGCGCGGAGGGCGGACCGGGAATGGCGCGCCGTGCGGCC GCCGGCGTAACGCGCGCT	739
miR-658	22q13.1	GCTCGGTTGCCGTGTTGCGGGCCCTGCCCGCCCGCCAGCTCGCTGACAGCACGACTCAGGGCGGAGGGAAGTAGTCCGTT GGTCGGTCGGGAACGAGG	740
miR-663	20p11.1	CCTCCGGCGTCCCAGGCGGGCGCGGCCCTCGTGTCTGTGGCGGTGGGATCCCGCGGCCGTGTTTTCTGCTGGTG GCCCGGCCATG	741
miR-671-5p	07q36.1	GCAGGTGAACCTGGCAGGCCAGGAAGAGGAGGCCCTGGAGGGGCTGGAGGTGATGGATGTTTTCTCCGGTTCTCAGGGCT CCACCTCTTTCGGGCCGTAGAGCCAGGGCTGGTGC	742
miR-675	11p15.5	CCCAGGGTCTGGTGGGAGAGGGGCCACAGTGGACTTGTTGACGCTGTATGCCCTCACCGCTCAGCCCTGGG	743
miR-708	11q14.1	AACCTGCCCTCAAGGAGCTTACAATCTAGCTGGGGGTAAATGACTTGACATGAACACAACACTAGACTGTGAGCTTCTAGAGGGCAG GGA	744
miR-744	17p12	TTGGGCAAGGTGCGGGGCTAGGGCTAACAGCAGTCTTACTGAAGGTTTCTCGTGAACCAACGACACATGCTGTTGCCACTAACCTC AACCTTACTCGGC	745
miR-765	01q23.1	TTTAGGCGCTGATGAAGTGGAGTTTCTAGTACAGGCCCTTTTCAAGCCCTACGAGAACTGGGGTTTCTGGAGGAGAAGGAAGG TGATGAAGGATCTGTTCTCGTGAGCCTGAA	746

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
miR-920	12p12.1	GTAGTTGTTCTACAGAAACCTGGATGTGTAGGAGCTAAGACACACTCCAGGGGAGCTGTGGAAGCAGTAACACG	747
miR-923	17q12	TATTTGTCAGCGGAGGAAAGAACTAACAGGATTCCTCAGTAATGGCGAGTG	748
miR-92a-2*	Xq26.2	TCATCCCTGGTGGGATTTTGTGCAATTACTTGTGTTCTATATAAAGTATTGCACCTTGTCCTCCGGCCTGTGGAAGA	749
miR-92b*	01q22	CGGGCCCCGGGGCGGGAGGGACGGGACGGGTGCAGTGTTGTTTTTCCCCCGCCAATATTGCACCTCGTCCCCGGCCTCC GGCCCCCCCCGGCCC	750
miR-93	07q22.1	CTGGGGCTCCAAAGTGCTGTTCTGTCAGGTAGTGTGATTACCCAACTACTGCTGAGCTAGCACTTCCCAGCCCCCGG	751
miR-98	xp11.22	AGGATTCTGCTCATGCCAGGGGTGAGGTAGTAAGTTGTATTGTTGTGGGTAGGGATATTAGCCCCCAATTAGAAGATAACTATAC AACTTACTACTTTCCCTGGTGTGTGGCATAITCA	752
miR-99b	19q13.41	GGCACCCACCCGTAGAACCCGACCTTCGGGGGCCCTTCGCCGCACACAAGCTCGTGTCTGTGGGTCCGTGTC	753
miR-103	5q34	TACTGCCCTCGGCTTCTTTACAGTGCTGCCCTTGTTCATATGGATCAAGCAGCAATTGTACAGGGCTATGAAGGCATTG	754
miR-103	20p13	TTGTGCTTTCAGCTTCTTTACAGTGTCTGCCCTTGTAGCAATCAGGTCAAGCAGCAATTGTACAGGGCTATGAAAAGAACCA	755
miR-106a	Xq26.2	CCTTGCCCATGTAAAAGTGCTTACAGTGCAGGTAGCTTTTTGAGATCTACTGCAATGTAAGCACTTCTTACATTACCATGG	756
miR-106b	7q22.1	CCTGCCGGGGCTAAAGTGTGACAGTGCAGATAGTGGTCCTCTCCGTGCTACCGCACTGTGGGTACTTGTCTCCAGCAGG	757
miR-107	10q23.31	CTCTCTGCTTTCAGCTTCTTTACAGTGTTGCCCTTGTGGCATGGAGTTCAAGCAGCAATTGTACAGGGCTATCAAAGCACAGA	758
miR-130a	11q12.1	TGCTGCTGGCCAGAGCTCTTTTACACATTGTGCTACTGTCTGCACCTGTCTACTAGCAGTGCAATGTTAAAAGGGCAATTGGCCGTGT AGTG	759
miR-130b	22q11.21	GGCCTGCCCGACACICTTTCCCTGTGTGCACACTACTATAGGCCGGCTGGGAAGCAGTGCAATGATGAAAGGGCATCGGTCAAGTCTC	760
miR-134	14q32.31	CAGGGTGTGTGACTGGTTGACCAGAGGGGCATGCACTGTGTTCACTCCCTGTGGGCCACCTAGTCACCAACCCCTC	761
miR-138	3p21.33	CGTTGCTGCAGCTGGTGTGTGAATCAGGCCGACGAGCAGCGCATCCTCTTACCCGGCTATTTACGACACCAAGGGTTGCATCA	762
miR-138	16q13	CCCTGGCATGGTGTGGTGGGCAGCTGGTGTGAATCAGGCCGTTGCCAATCAGAGAACGGCTACTTCACAACACCCAGGGCC ACACCACACTACAGG	763
miR-15a	13q14.2	CCTTGAGTAAAGTAGCAGCACATAATGGTTTGTGGATTTTGAAAAGGTGCAGGCCATATTGTGCTGCCCTCAAAAAATACAAGG	764
miR-15b	03q25.33	TTGAGGCCCTTAAAGTACTGTAGCAGCACATCATGGTTTACATGCTACAGTCAAGATGCGAATCATTATTTGCTGCTCTAGAAAATTTA AGGAAATTCAT	765
miR-17	13q31.3	GTCAGAATAATGTCAAAGTGCTTACAGTGCAGGTAGTGATATGTGCATCTACTGCAGTGAAGGCACCTTGTAGCATTATGGTGAC	766
miR-181a-1	1q31.3	AGAAGGGCTATCAGGCCAGCCTTCAGAGGACTCCAAGGAACATTCAACGCTGTCGGTGAGTTTGGGATTTGAAAAAACCACTGAC CGTTGACTGTACCTTGGGGTCCTTA	767
miR-181a-2	9q33.3	TGAGTTTGTAGGTTGCTTTCAGTGAACATTCAACGCTGTGGTGAGTTTGGAATTAAATCAAAACCATCGACCGTTGATTGTACCC	768

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
		TATGGCTAACCATCATCTACTCCA	
miR-181b-1	1q31.3	CTGATGGCTGCACTCAACATTCATTGCTGTCGGTGGTTTGAGTCTGAATCAACTCACTGATCAATGAATGCAAACCTGCGGACCAA ACA	769
miR-181b-2	9q33.3	CCTGTGCAGAGATTATTTTTTAAAGGTCACAATCAACATTCATTGCTGTCGGTGGTTGAACCTGTGTGGACAAAGCTCACTGAACA ATGAATGCAACTGTGCCCCGCTT	770
miR-191	03p21.31	CGGCTGGACAGCGGGCAACGGAATCCCAAAAGCAGCTGTTGTCTCCAGAGCATTCAGCTGCGCTTGGATTTTCGTCCCCTGCTC TCCTGCCCT	771
miR-195	17p13.1	AGCTTCCCTGGCTCTAGCAGCACAGAAAATATTGGCACAGGGAAGCGAGTCTGCCAATATTGGCTGTGCTGCCAGGCAGGGGTG GTG	772
miR-196b	07p15.2	ACTGGTCGGTGATTTAGGTAGTTTCCTGTTGGGATCCACCTTTCTCTCGACAGCACACACTGCCTTCATTACTTCAGTTG	773
miR-19a	13q31.3	GCAGTCCTCTGTAGTTTTGCATAGTTGCACIACAGAAGAAATGTAGTTGTGCAAAATCTATGCAAAACTGATGGTGGCCTGC	774
miR-20a	13q31.3	GTAGCACTAAAGTGCTTATAGTGCAGGTAGTGTTTAGTTATCTACTGCATTATGAGCACTTAAAGTACTGC	775
miR-20b	Xq26.2	AGTACCAAAGTGCTCATAGTGCAGGTAGTTTTGGCATGACTCTACTGTAGTATGGGCACCTTCCAGTACT	776
miR-22	17p13.3	GGCTGAGCCCGCAGTAGTTCCTTCAGTGGCAAGCTTTATGTCTGACCCAGCTAAAGCTGCCAGTTGAAGAACTGTTGCCCTCTGCC	777
miR-221	Xp11.3	TGAACATCCAGGTCTGGGCATGAACCTGGCATACAATGTAGATTTCTGTGTTCTAGGCAACAGCTACATTGTCTGCTGGGTTT CAGGCTACCTGGAAACATGTTCTC	778
miR-222	Xp11.3	GCTGCTGGAAGGTGTAGGTACCCCTCAATGGCTCAGTAGCCAGTGTAGATCCTGTCTTTTCGTAATCAGCAGCTACATCTGGCTACT GGTCTCTGATGGCATCTTCTAGCT	779
miR-25	07q22.1	GGCCAGTGTTGAGAGGGCGGAGACTTGGGCCAATTGCTGGACGCTGCCCTGGGCATTGCACCTTGCTCGGCTGACAGTGCCCGGC C	780
miR-29c	01q32.2	ATCTCTTACACAGGCTGACCGATTTCTCCTGGTGTTTCAGAGTCTGTTTTTGTCTAGCACCATTTGAAATCGGTTATGATGTAGGGG GA	781
miR-31	09p21.3	GGAGAGGAGGCAAGATGCTGGCATAGCTGTTGAACTGGGAACCTGCTATGCCAACATATTGCCATCTTTCC	782
miR-335	07q32.2	TGTTTTGAGCGGGGTCAAGAGCAATAACGAAAAATGTTTGTCAAAACCGTTTTTCATTATTGCTCCTGACCTCCTCTCATTTGCT ATATTCA	783
miR-342-3p	14q32.2	GAAACTGGGCTCAAGGTGAGGGGTGCTATCTGTGATTGAGGGACATGTTAATGGAATTGTCTCACACAGAAAATCGCACCCGTCA CCTTGGCCTACTTA	784
miR-370	14q32.31	AGACAGAGAAAGCCAGGTCACGTCTCTGCAGTTACACAGCTCACGAGTGCCCTGCTGGGTGGAACCTGGTCTGTCT	785
miR-452	Xq28	GCTAAGCACTTACAACGTGTTTGCAGAGGAAACTGAGACTTTGTAACTATGTCTCAGTCTCATCTGCAAGAAGTAAGTGCTTTGC	786
miR-494	14q32.31	GATACTCGAAGGAGAGGTTGTCCGTGTTCTCTCTTTATTTATGATGAAACATACACGGGAAACCTCTTTTTTAGTATC	787

Table 3

Pre-target RNA Candidate	chrom. Location	Pre-microRNA sequences	pre-micro RNA SEQ ID NO:
miR-7-1	9q21.32	TTGGATGTTGGCCCTAGTTCTGTGTGGAAGACTAGTGATTTTGTGTTTTTAGATAAACTAAATCGACAACAAATCACAGTCTGCCATA TGGCACAGGCCCATGCCCTCTACAG	788
miR-7-3	19p13.3	AGATTAGAGTGGCTGTGGTCTAGTGTGTGGAAAGACTAGTGATTTTGTGTTCTGATGTACTACGACAACAAAGTCACAGCCCCGG CCTCATAGCGCAGACTCCCTTCGAC	789
miR-7-2	15q26.1	CTGGATACAGAGTGGACCCGGCTGGCCCCCATCTGGAAGACTAGTGATTTTGTGTTGTCTTACTGCGCTCAACAACAAATCCCAGT CTACCTAATGGTGCCAGCCCATCGCA	790
miR-92-a1	13q31.3	CTTTCTACACAGGTTGGGATCGGTTGCAATGCTGTGTTTTCTGTATGGTATTGCACTTGTCCCCGGCCTGTTGAGTTTGG	791
miR-92-a2	Xq26.2	TCATCCCTGGGTGGGATTTTGTTCATTACTTGTGTTCTATATAAAGTATTGCACCTTGTCCTCCGGCCTGTGGAAGA	792
miR-99a	21q21.1	CCCATTGGCATAAACCCGTAGATCCGATCTTGTGGTGAAGTGGACCCGCACAAAGCTCGCTTCTATGGGTCTGTGTCAGTGTG	793

Table 4		
Mature microRNA Sequences (5' to 3')		
microRNA	sequence	SEQ ID NO
let-7a	UGAGGUAGUAGGUUGUAUAGUU	794
let-7b	UGAGGUAGUAGGUUGUGUGGUU	795
let-7c	UGAGGUAGUAGGUUGUAUGGUU	796
let-7d	AGAGGUAGUAGGUUGCAUAGUU	797
let-7e	UGAGGUAGGAGGUUGUAUAGUU	798
let-7f	UGAGGUAGUAGAUUGUAUAGUU	799
let-7g	UGAGGUAGUAGUUUGUACAGUU	800
let-7i	UGAGGUAGUAGUUUGUGCUGUU	801
miR-100	AACCCGUAGAUCCGAACUUGUG	802
miR-103	AGCAGCAUUGUACAGGGCUAUGA	803
miR-106a	AAAAGUGCUUACAGUGCAGGUAG	804
miR-106b	UAAAGUGCUGACAGUGCAGAU	805
miR-107	AGCAGCAUUGUACAGGGCUAUCA	806
miR-1224-5p	GUGAGGACUCGGGAGGUGG	807
miR-1225-5p	GUGGGUACGGCCCAGUGGGGGG	808
miR-1228*	GUGGGCGGGGGCAGGUGUGUG	809
miR-125a-5p	UCCUGAGACCCUUUAACCUUGUGA	810
miR-125b	UCCUGAGACCCUAACUUGUGA	811
miR-126	UCGUACCGUGAGUAAUAAUGCG	812
miR-130a	CAGUGCAAUGUUAAAAGGGCAU	813
miR-130b	CAGUGCAAUGAUGAAAGGGCAU	814
miR-134	UGUGACUGGUUGACCAGAGGGG	815
miR-138	AGCUGGUGUUGUGAAUCAGGCCG	816
miR-135a*	UAUAGGGAUUGGAGCCGUGGCG	817
miR-142-3p	UGUAGUGUUUCCUACUUUAUGGA	818
miR-145	GUCCAGUUUCCCAGGAUCCCU	819
miR-146b-5p	UGAGAACUGAAUCCAUAGGCU	820
miR-149*	AGGGAGGGACGGGGGCUGUGC	821
miR-150*	CUGGUACAGGCCUGGGGGACAG	822
miR-155	UUAAUGCUAAUCGUGAUAGGGGU	823
miR-15a	UAGCAGCACAUAAUGGUUUUGUG	824
miR-15b	UAGCAGCACAUCAUGGUUUACA	825
miR-16	UAGCAGCACGUAAUAUUGGCG	826
miR-17	CAAAGUGCUUACAGUGCAGGUAG	827
miR-181a	AACAUUCAACGCUGUCGGUGAGU	828
miR-181b	AACAUUCAUUGCUGUCGGUGGGU	829
miR-181c	AACAUUCAACCUUGUCGGUGAGU	830
miR-191	CAACGGAAUCCCAAAAGCAGCUG	831
miR-195	UAGCAGCACAGAAUAUUGGC	832

Table 4		
Mature microRNA Sequences (5' to 3')		
microRNA	sequence	SEQ ID NO
miR-196b	UAGGUAGUUUCCUGUUGUUGGG	833
miR-198	GGUCCAGAGGGGAGAUAGGUUC	834
miR-199a-3p	ACAGUAGUCUGCACAUUGGUUA	835
miR-199b-3p	ACAGUAGUCUGCACAUUGGUUA	836
miR-19a	UGUGCAAUUCUAUGCAAAACUGA	837
miR-19b	UGUGCAAUCCAUGCAAAACUGA	838
miR-200b	UAAUACUGCCUGGUAAUGAUGA	839
miR-200c	UAAUACUGCCGGGUAAUGAUGGA	840
miR-205	UCCUUCAUUCCACCGGAGUCUG	841
miR-20a	UAAAGUGCUUAUAGUGCAGGUAG	842
miR-20b	CAAAGUGCUCAUAGUGCAGGUAG	843
miR-21	UAGCUUAUCAGACUGAUGUUGA	844
miR-22	AAGCUGCCAGUUGAAGAACUGU	845
miR-221	AGCUACAUUGUCUGCUGGGUUUC	846
miR-222	AGCUACAUCUGGCUACUGGGU	847
miR-23a	AUCACAUUGCCAGGGAUUUC	848
miR-23a*	GGGGUUCUGGGGAUGGGAUUU	849
miR-23b	AUCACAUUGCCAGGGAUUACC	850
miR-24	UGGCUCAGUUCAGCAGGAACAG	851
miR-25	CAUUGCACUUGUCUCGGUCUGA	852
miR-25*	AGGCGGAGACUUGGGCAAUUG	853
miR-26a	UUCAAGUAAUCCAGGAUAGGCU	854
miR-26b	UUCAAGUAAUUCAGGAUAGGU	855
miR-27a	UUCACAGUGGCUAAGUUCGCG	856
miR-27b	UUCACAGUGGCUAAGUUCUGC	857
miR-298	AGCAGAAGCAGGGAGGUUCUCCCA	858
miR-29a	UAGCACCAUCUGAAAUCGGUUA	859
miR-29b	UAGCACCAUUUGAAAUCAGUGUU	860
miR-29c	UAGCACCAUUUGAAAUCGGUUA	861
miR-29c*	UGACCGAUUUCUCCUGGUGUUC	862
miR-30a	UGUAAACAUCCUCGACUGGAAG	863
miR-30b	UGUAAACAUCCUACACUCAGCU	864
miR-30b*	CUGGGAGGUGGAUGUUUACUUC	865
miR-30c	UGUAAACAUCCUACACUCUCAGC	866
miR-30c-1*	CUGGGAGAGGGUUGUUUACUCC	867
miR-30c-2	UGUAAACAUCCUACACUCUCAGC	868
miR-30d	UGUAAACAUCCCGACUGGAAG	869
miR-30e	UGUAAACAUCCUUGACUGGAAG	870
miR-31	AGGCAAGAUGCUGGCAUAGCU	871

Table 4		
Mature microRNA Sequences (5' to 3')		
microRNA	sequence	SEQ ID NO
miR-320a	AAAAGCUGGGUUGAGAGGGCGA	872
miR-331-3p	GCCCCUGGGCCUAUCCUAGAA	873
miR-335	UCAAGAGCAAUAACGAAAAAUGU	874
miR-342-3p	UCUCACACAGAAAU CGCACCCGU	875
miR-370	GCCUGCUGGGGUGGAACCUGGU	876
miR-371-5p	ACUCAAACUGUGGGGGGCACU	877
miR-373*	ACUCAAAAUGGGGGCGCUUUCC	878
miR-375	UUUGUUCGUUCGGCUCGCGUGA	879
miR-423-5p	UGAGGGGCAGAGAGCGAGACUUU	880
miR-424	CAGCAGCAAUUCAUGUUUUGAA	881
miR-452	AACUGUUUGCAGAGGAAACUGA	882
miR-483-5p	AAGACGGGAGGAAAGAAGGGAG	883
miR-486-3p	CGGGGCAGCUCAGUACAGGAU	884
miR-491-3p	CUUAUGCAAGAUUCCCUUCUAC	885
miR-491-5p	AGUGGGGAACCCUCCAUGAGG	886
miR-494	UGAAACAUACACGGGAAACCUC	887
miR-513a-5p	UUCACAGGGAGGUGUCAU	888
miR-513b	UUCACAAGGAGGUGUCAUUUAU	889
miR-516a-5p	UUCUCGAGGAAAGAAGCACUUUC	890
miR-550	AGUGCCUGAGGGAGUAAGAGCCC	891
miR-557	GUUUGCACGGGUGGGCCUUGUCU	892
miR-575	GAGCCAGUUGGACAGGAGC	893
miR-612	GCUGGGCAGGGCUUCUGAGCUCCUU	894
miR-614	GAACGCCUGUUCUUGCCAGGUGG	895
miR-630	AGUAUUCUGUACCAGGGAAGGU	896
miR-637	ACUGGGGGCUUUCGGGCUCUGCGU	897
miR-638	AGGGAUCGCGGGCGGGUGGCGGCCU	898
miR-658	GGCGGAGGGAAGUAGGUCCGUUGGU	899
miR-663	AGGCGGGGCGCCGCGGGACCGC	900
miR-671-5p	AGGAAGCCCUGGAGGGGCUGGAG	901
miR-675	UGGUGCGGAGAGGGCCCACAGUG	902
miR-7	UGGAAGACUAGUGAUUUUGUUGU	903
miR-708	AAGGAGCUUACAAUCUAGCUGGG	904
miR-744	UGC GGGGCUAGGGCUAACAGCA	905
miR-765	UGGAGGAGAAGGAAGGUGAUG	906
miR-920	GGGGAGCUGUGGAAGCAGUA	907
miR-923	GUCAGCGGAGGAAAAGAAACU	908
miR-92a	UAUUGCACUUGUCCCGGCCUGU	909
miR-92a-2*	GGGUGGGGAUUUGUUGCAUUAC	910

Table 4		
Mature microRNA Sequences (5' to 3')		
microRNA	sequence	SEQ ID NO
miR-92b*	AGGGACGGGACGCGGUGCAGUG	911
miR-93	CAAAGUGCUGUUCGUGCAGGUAG	912
miR-98	UGAGGUAGUAAGUUGUAUUGUU	913
miR-99a	AACCCGUAGAUCCGAUCUUGUG	914
miR-99b	CACCCGUAGAACCGACCUUGCG	915
miR-200a	ACAUCGUUACCAGACAGUGUUA	916
miR-720	UGGAGGCCCCAGCGAGA	917
miR-1202	CUCCCCACUGCAGCUGGCAC	918
miR-1249	UGAAGAAGGGGGGGAAGGGCGU	919
miR-1275	GACAGCCUCUCCCCCAC	920
miR-129-3p	AUGCUUUUUGGGGUAAGGGCUU	921
miR-1321	AUCACAUUCACCUCCCUG	922
miR-1323	AGAAAUGCCCCUCAGUUUUGA	923
miR-376c	ACGUGGAAUUUCCUCUAUGUU	924
miR-429	ACGGUUUUACCAGACAGUAUUA	925

Table 5

Mature microRNA Sequences (5'-3')		
microRNA	sequence	SEQ ID NO
miR-1	UGGAAUGUAAAGAAGUAUGUA	926
miR-9	UCUUUGGUUAUCUAGCUGUAUGA	927
miR-9*	UAAAGCUAGAUAAACCGAAAGU	928
miR-10a	UACCCUGUAGAUCCGAAUUUGUG	929
miR-10b	UACCCUGUAGAACCGAAUUUGU	930
miR-17-3p	ACUGCAGUGAAGGCACUUGU	931
miR-18	UAAGGUGCAUCUAGUGCAGUA	932
miR-20	UAAAGUGCUUAUAGUGCAGGUA	933
miR-28	AAGGAGCUCACAGUCUAUUGAG	934
miR-30a-3p	CUUUCAGUCGGAUGUUUGCAGC	935
miR-32	UAUUGCACAUUACUAAGUUGC	936
miR-33	GUGCAUUGUAGUUGCAUUG	937
miR-34a	UGGCAGUGUCUAGCUGGUUGU	938
miR-34b	AGGCAGUGUCAUAGCUGAUUG	939
miR-34c	AGGCAGUGUAGUUAGCUGAUUG	940
miR-92	UAUUGCACUUGUCCCGGCCUGU	941
miR-95	UUCAACGGGUUUUAUUGAGCA	942
miR-96	UUUGGCACUAGCACAUUUUUGC	943
miR-101	UACAGUACUGUGAUAAACUGAAG	944
miR-105	UCAA AUGCUCAGACUCCUGU	945
miR-122a	UGGAGUGUGACAAUGGUGUUUGU	946
miR-124a	UUAAGGCACGCGGUGAAUGCCA	947
miR-126*	CAUUAUUACUUUUGGUACGCG	948
miR-127	UCGGAUCCGUCUGAGCUUGGCU	949
miR-128a	UCACAGUGAACCGGUCUCUUUU	950
miR-128b	UCACAGUGAACCGGUCUCUUUC	951
miR-129	CUUUUUGCGGUCUGGGCUUGC	952
miR-132	UACAGUCUACAGCCAUGGUCG	953
miR-133a	UUGGUCCCCUUAACCAGCUGU	954
miR-133b	UUGGUCCCCUUAACCAGCUA	955
miR-135a	UAUGGCUUUUUAUCCUAUGUGA	956
miR-135b	UAUGGCUUUUCAUCCUAUGUG	957
miR-136	ACUCCAUUUGUUUUGAUGAUGGA	958
miR-137	UAUUGCUUAAGAAUACGCGUAG	959
miR-139	UCUACAGUGCACGUGUCU	960
miR-140	AGUGGUUUUACCCUAUGGUAG	961

Table 5

Mature microRNA Sequences (5'-3')		
microRNA	sequence	SEQ ID NO
miR-141	AACACUGUCUGGUAAGAUGG	962
miR-142-5p	CAUAAAGUAGAAAGCACUAC	963
miR-143	UGAGAUGAAGCACUGUAGCUCA	964
miR-144	UACAGUAUAGAUGAUGUACUAG	965
miR-146	UGAGAACUGAAUUCCAUGGGUU	966
miR-147	GUGUGUGGAAAUGCUUCUGC	967
miR-148a	UCAGUGCACUACAGAACUUUGU	968
miR-148b	UCAGUGCAUCACAGAACUUUGU	969
miR-149	UCUGGCUCCGUGUCUUCACUCC	970
miR-150	UCUCCCAACCCUUGUACCAGUG	971
miR-151	ACUAGACUGAAGCUCCUUGAGG	972
miR-152	UCAGUGCAUGACAGAACUUGG	973
miR-153	UUGCAUAGUCACAAAAGUGA	974
miR-154	UAGGUUAUCCGUGUUGCCUUCG	975
miR-154*	AAUCAUACACGGUUGACCUAUU	976
miR-182	UUUGGCAAUGGUAGAACUCACA	977
miR-182*	UGGUUCUAGACUUGCCAACUA	978
miR-183	UAUGGCACUGGUAGAAUUCACUG	979
miR-184	UGGACGGAGAACUGAUAAGGGU	980
miR-185	UGGAGAGAAAGGCAGUUC	981
miR-186	CAAAGAAUUCUCCUUUUGGGCUU	982
miR-187	UCGUGUCUUGUGUUGCAGCCG	983
miR-188	CAUCCCUUGCAUGGUGGAGGGU	984
miR-189	GUGCCUACUGAGCUGAUAUCAGU	985
miR-190	UGAUAUGUUUGAUAUAUUAGGU	986
miR-192	CUGACCUAUGAAUUGACAGCC	987
miR-193	AACUGGCCUACAAAGUCCAG	988
miR-194	UGUACAGCAACUCCAUGUGGA	989
miR-196a	UAGGUAGUUUCAUGUUGUUGG	990
miR-197	UUCACCACCUUCUCCACCCAGC	991
miR-200a	UAACACUGUCUGGUAACGAUGU	992
miR-202	AGAGGUAUAGGGCAUGGGAAGA	993
miR-203	GUGAAAUGUUUAGGACCACUAG	994
miR-204	UUCCCUUUGUCAUCCUAUGCCU	995
miR-206	UGGAAUGUAAGGAAGUGUGUGG	996
miR-208	AUAAGACGAGCAAAAAGCUUGU	997
miR-210	CUGUGCGUGUGACAGCGGCUG	998

Table 5

Mature microRNA Sequences (5'-3')		
microRNA	sequence	SEQ ID NO
miR-211	UUCCCUUUGUCAUCCUUCGCCU	999
miR-212	UACAGUCUCCAGUCACGGCC	1000
miR-213	ACCAUCGACCGUUGAUUGUACC	1001
miR-214	ACAGCAGGCACAGACAGGCAG	1002
miR-215	AUGACCUAUGAAUUGACAGAC	1003
miR-216	UAAUCUCAGCUGGCAACUGUG	1004
miR-217	UACUGCAUCAGGAACUGAUUGGAU	1005
miR-218	UUGUGCUUGAUCUAACCAUGU	1006
miR-219	UGAUUGUCCAAACGCAAUUCU	1007
miR-220	CCACACCGUAUCUGACACUUU	1008
miR-223	UGUCAGUUUGUCAAAUACCCC	1009
miR-224	CAAGUCACUAGUGGUUCCGUUUA	1010
miR-296	AGGGCCCCCCCCUCAUCCUGU	1011
miR-299	UGGUUUACCGUCCCAUAACA	1012
miR-301	CAGUGCAAUAGUAUUGUCAAGC	1013
miR-302a	UAAGUGCUUCCAUGUUUUGGUGA	1014
miR-302b*	ACUUUAACAUGGAAGUGCUUUCU	1015
miR-302b	UAAGUGCUUCCAUGUUUUGAG	1016
miR-302c*	UUUAACAUGGGGUACCUGCUG	1017
miR-302c	UAAGUGCUUCCAUGUUUCAGUGG	1018
miR-302d	UAAGUGCUUCCAUGUUUGAGUGU	1019
miR-320	AAAAGCUGGGUUGAGAGGGCGAA	1020
miR-321	UAAGCCAGGGAUUGUGGGGUUC	1021
miR-323	GCACAUUACACGGUCGACCUCU	1022
miR-324-5p	CGCAUCCCCUAGGGCAUUGGUGU	1023
miR-324-3p	CCACUGCCCCAGGUGCUGCUGG	1024
miR-325	CCUAGUAGGUGUCCAGUAAGU	1025
miR-326	CCUCUGGGCCCUUCCUCCAG	1026
miR-328	CUGGCCCUCUCUGCCCUUCCGU	1027
miR-330	GCAAAGCACACGGCCUGCAGAGA	1028
miR-331	GCCCCUGGGCCUAUCCUAGAA	1029
miR-337	UCCAGCUCCUAUAUGAUGCCUUU	1030
miR-338	UCCAGCAUCAGUGAUUUUGUUGA	1031
miR-339	UCCCUGUCCUCCAGGAGCUCA	1032
miR-340	UCCGUCUCAGUUACUUUAUAGCC	1033
miR-342	UCUCACACAGAAUCGCACCCGUC	1034
miR-345	UGCUGACUCCUAGUCCAGGGC	1035

Table 5

Mature microRNA Sequences (5'-3')		
microRNA	sequence	SEQ ID NO
miR-346	UGUCUGCCCGCAUGCCUGCCUCU	1036
miR-367	AAUUGCACUUUAGCAAUGGUGA	1037
miR-368	ACAUAGAGGAAAUUCCACGUUU	1038
miR-369	AAUAAUACAUGGUUGAUCUUU	1039
miR-371	GUGCCGCCAUCUUUUGAGUGU	1040
miR-372	AAAGUGCUGCGACAUUUGAGCGU	1041
miR-373	GAAGUGCUUCGAUUUUGGGGUGU	1042
miR-374	UUAUAAUACAACCUGAUAAGUG	1043
miR-320c	AAAAGCUGGGUUGAGAGGGU	2692

Table 6
Up-regulated target RNAs for detection of non-small cell lung cancer

		Fold changes v. normal Lung									
Array Probe	Probe SEQ ID NO.	Epi4	Epi7	Epi5	Adk1	Adk3	Adk11	Adk8	Adk9	Adk2	Adk10
miR-21	196	-2.52	1.77	-1.51	1.45	1.41	3.51	2.27	3.43	5.38	1.25
miR-765	246	25.46	3.15	3.08	6.06	-1.58	3.70	1.82	3.00	-1.58	1.02
4037-R3-2	15	6.57	1.44	-1.28	2.17	-1.28	2.17	2.34	2.64	6.81	-1.06
miR-27b	205	-1.11	2.20	2.52	-1.91	1.18	1.64	1.24	2.05	6.50	-1.85
miR-29a	207	-2.68	1.00	3.18	1.50	-1.90	2.13	1.41	1.23	2.18	-2.96
miR-923	248	6.70	1.19	1.24	1.35	-2.18	1.11	-1.43	1.19	2.00	-2.79
miR-199a-3p	191	-3.52	1.40	1.43	-1.94	1.72	1.56	1.33	1.59	2.97	-4.20
miR-331-3p	219	-1.21	-1.21	2.17	1.42	1.24	1.47	1.27	-1.21	-1.21	1.36
miR-23a	197	1.38	2.90	-4.40	1.37	-4.40	2.28	-4.40	2.38	7.76	-2.09
miR-146b-5p	184	-1.89	1.12	2.44	-1.89	-1.89	2.71	1.07	1.30	2.48	-1.21
miR-483-5p	225	24.38	1.51	1.68	5.17	-1.54	1.11	-1.64	3.19	-1.54	-1.03
9733-L3-1	129	7.61	3.56	-1.48	2.44	-1.48	-1.48	1.07	1.69	-1.48	1.33
miR-30c-1*	214	39.98	-1.58	1.57	7.84	-1.58	-1.58	-1.42	2.67	-1.58	-1.58
4593-R3-1	26	8.31	2.80	1.23	1.93	-4.73	-2.65	-5.00	1.79	-9.77	1.20
8433_C-R4-1	164	24.29	1.73	1.87	5.70	-1.45	-1.45	-1.45	1.95	-1.45	1.13
4855-R3-1	30	8.26	1.91	1.44	1.78	-3.56	-1.50	-1.81	1.87	-2.80	1.20
4666-R4-1	27	24.77 *	2.27	3.49 *	7.40 *	-2.36	-1.70	-2.05	2.37	-3.62	-1.01

Table 7: Target RNAs more frequently present at elevated levels in squamous cell carcinoma (SCC)

Gene	Fold change average in SCC	Number of adenocarcinomas with increased levels	Number of SCC with increased levels	Probe SEQ ID NO	Pre-microRNA SEQ ID NO	microRNA SEQ ID NO
10366-R3-2	4.1	1	6	145	539	
12223-L5-1	3.2	2	5	2110	2234	2584
12907-L5-1	6.4	0	5	1106	1228	
12911-L5-1	3.2	2	5	2115	2239	2590
12917-R5-2	6.0	1	6	1108	1230	
13108-L5-2	3.3	1	5	2673	2681	2595
13122-L5-1	11.5	2	7	1066	1242	2597
13272-R5-2	4.0	1	5	2674	2682	2611
13316-R5-2	2.8	0	5	2675	2683	2616
13331-L5-2	4.1	1	5	2676	2684	2617
13499-R5-1	3.3	2	5	2677	2685	2634
3923-R5-1	5.3	3	7	1070	1273	2647
4261-R5-1	6.4	0	5	1148	1277	
4479-R3-1	4.3	1	6	25	421	
5232-L5-2	5.2	0	5	1154	1287	2653
5392-R5-1	7.0	2	5	1155	1288	2654
5971-R5-2	2.6	1	5	2678	2686	2657
6183-R5-1	2.8	0	5	2149	2274	
7026-L3-1	2.8	1	5	2679	2687	
7292-L3-2	3.5	2	6	80	476	
7471-L5-1	4.5	1	5	2680	2688	
8004-R3-2	11.0	2	6	97	492	
8316-R5-1	20.3	1	5	1177	1315	
9349-R5-2	24.4	3	6	1080	1325	2672
9594-R5-1	9.7	2	6	1081	1328	
miR-1323	4.9	2	6	1197	1338	923
miR-205	21.2	3	7	195	694	841
miR-675	2.9	2	5	243	743	902
miR-923	3.5	3	6	248	748	908

Table 8: Target RNAs more frequently present at elevated levels in adenocarcinoma

Gene	Fold change average in adenocarcinoma	Number of adenocarcinomas with increased levels	Number of SCC with increased levels	Probe SEQ ID NO	Pre-microRNA SEQ ID NO	microRNA SEQ ID NO
13252-L5-3	4.9	5	1	1123	1249	2609
13373-R5-2	5.1	5	3	1130	1257	2624
9798-L5-1	9.9	5	3	1188	1329	
miR-106b	4.5	5	2	363	757	805
miR-20b	4.8	6	2	1086	1349	843
miR-92a	4.0	4	0	2180	2307	909
miR-93	3.8	5	1	251	751	912

Table 9: Target RNAs present at increased levels in aggressive forms of lung cancer

Gene	Probe SEQ ID NO	Pre-microRNA SEQ ID NO(s)	microRNA SEQ ID NO
10083-L5-1	1090	1211	
10233-R5-1	1091	1212	2576
10455-L5-1	1063	1215	2578
11444-L5-3	1097	1219	
12729-R5-1	1103	1225	
12888-L5-2	1105	1227	
12907-L5-1	1106	1228	
12917-R5-2	1108	1230	
12947-L5-4	1064	1231	
12974-R5-2	1110	1233	2592
12979-R5-2	1111	1234	
13001-L5-1	1113	1236	
13070-R5-3	1115	1238	
13122-L5-1	1066	1242	2597
13185-L5-3	1118	1243	2603
13219-L5-1	1067	1245	
13245-L5-4	1122	1248	2607
13274-L5-3	1124	1251	2612
13357-L5-4	1128	1255	2621
13398-R5-4	1132	1259	2627
13467-L5-1	1069	1262	2630
13468-L5-1	1135	1263	2631
13470-R5-1	1136	1264	
13473-L5-3	1137	1265	2633
13500-L5-3	138	1266	2635, 2636
3744-R5-1	1143	1271	2645, 2646
3875-R5-2	1144	1272	
3992-R5-1	1146	1275	
4790-L5-2	1150	1282	
5080-R3-1	1073	1285	2650
5108-R5-2	1153	1286	
5392-R5-1	1155	1288	2654
6037-R3-2	1159	1292	2658
6181-L5-1	1160	1293	
6233-L5-2	1161	1295	
6235-R5-2	1075	1296	2661
6474-L5-1	1164	1299	
6602-R3-1	1165	1300	
6683-R5-1	1167	1302	
6906-L5-1	1172	1307	2666
6930-R5-1	1173	1308	2667
7764-R3-2	1175	1312	
8004-R3-2	97	492	
8316-R5-1	1177	1315	

836-R5-2	1079	1316	
8433 C-R4-1	1177	1317	
8808-R5-1	1183	1322	
9349-R5-2	1080	1325	2672
9594-R5-1	1081	1328	
miR-198	1202	1343	834
miR-298	1088	1351	858
miR-30c-1*	1204	1352	867
miR-320c	2689	2690, 2691	2692
miR-516a-5p	1209	1357	890
miR-765	246	746	906

[0049] In some embodiments, target RNAs can be measured in samples collected at one or more times from a patient to monitor the status or progress of lung cancer in the patient.

[0050] In some embodiments, a sample to be tested is obtained using one or more techniques commonly used for collecting lung tissue, *e.g.*, bronchoscopy, bronchial washing, brushing, or transbronchial needle aspiration. In some embodiments, the sample is obtained from a patient without lesions by bronchoalveolar lavage, *i.e.*, washing the airways with saline, to obtain cells. In some embodiments, the sample is obtained by biopsy, such as computed tomography (CT)-aided needle biopsy.

[0051] In some embodiments, the sample to be tested is a bodily fluid, such as blood, sputum, mucus, saliva, urine, semen, etc. In some embodiments, a sample to be tested is a blood sample. In some embodiments, the blood sample is whole blood. In some embodiments, the blood sample is a sample of blood cells. In some embodiments, the blood sample is plasma. In some embodiments, the blood sample is serum.

[0052] The clinical sample to be tested is, in some embodiments, freshly obtained. In other embodiments, the sample is a fresh frozen specimen. In some embodiments, the sample is a tissue sample, such as a formalin-fixed paraffin embedded sample. In some embodiments, the sample is a liquid cytology sample.

[0053] In some embodiments, the methods described herein are used for early detection of lung cancer in a sample of lung cells, such as those obtained by routine bronchoscopy. In some embodiments, the methods described herein are used for early detection of lung cancer in a sample of blood or serum.

[0054] In some embodiments, the clinical sample to be tested is obtained from individuals who have one or more of the following risk factors: history of smoking, over 45 years of age, exposure to radon gas, secondhand smoke or occupational carcinogens (*e.g.*, asbestos, radiation, arsenic, chromates, nickel, chloromethyl ethers, mustard gas, or coke-oven emissions), or lungs scarred by prior disease such as tuberculosis. In some embodiments, the clinical sample is obtained from individuals who have diagnostic signs

or clinical symptoms that may be associated with lung cancer, such as abnormal chest x-ray and/or computed tomography (“CT”) scan, cough, localized chest pain, or hoarseness.

[0055] Thus, in some embodiments, methods described herein can be used for routine screening of healthy individuals with no risk factors. In some embodiments, methods described herein are used to screen asymptomatic individuals having one or more of the above-described risk factors.

[0056] In some embodiments, the methods described herein can be used to assess the effectiveness of a treatment for lung cancer in a patient. In some embodiments, the target RNA expression levels are determined at various times during the treatment, and are compared to target RNA expression levels from an archival sample taken from the patient, *e.g.*, by bronchoscopy, before the manifestation of any signs of lung cancer or before beginning treatment. In some embodiments, the target RNA expression levels are compared to target RNA expression levels from an archival sample of normal tissue taken from the patient, *i.e.*, a sample of tissue taken from a tumor-free part of the patient’s lung by biopsy. Ideally, target RNA expression levels in the normal sample evidence no aberrant changes in target RNA expression levels. Thus, in such embodiments, the progress of treatment of an individual with lung cancer can be assessed by comparison to a sample of lung cells from the same individual when he was healthy or prior to beginning treatment, or by comparison to a sample of healthy lung cells from the same individual.

[0057] In embodiments in which the method comprises detecting expression of more than one target RNA, the expression levels of the plurality of target RNAs may be detected concurrently or simultaneously in the same assay reaction. In some embodiments, expression levels are detected concurrently or simultaneously in separate assay reactions. In some embodiments, expression levels are detected at different times, *e.g.*, in serial assay reactions.

[0058] In some embodiments, a method comprises detecting the level of at least one target RNA in a sample from a subject, wherein detection of a level of at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer in the subject. In some embodiments, a method comprises detecting the level of at least one target RNA in a sample from a subject and comparing

the level of the at least one target RNA in the sample to a normal level of the at least one target RNA, wherein a level of at least one target RNA in the sample that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer in the subject.

[0059] In some embodiments, a method of facilitating diagnosis of lung cancer in a subject is provided. Such methods comprise detecting the level of at least one target RNA in a sample from the subject. In some embodiments, information concerning the level of at least one target RNA in the sample from the subject is communicated to a medical practitioner. A “medical practitioner,” as used herein, refers to an individual or entity that diagnoses and/or treats patients, such as a hospital, a clinic, a physician’s office, a physician, a nurse, or an agent of any of the aforementioned entities and individuals. In some embodiments, detecting the level of at least one target RNA is carried out at a laboratory that has received the subject’s sample from the medical practitioner or agent of the medical practitioner. The laboratory carries out the detection by any method, including those described herein, and then communicates the results to the medical practitioner. A result is “communicated,” as used herein, when it is provided by any means to the medical practitioner. In some embodiments, such communication may be oral or written, may be by telephone, in person, by e-mail, by mail or other courier, or may be made by directly depositing the information into, e.g., a database accessible by the medical practitioner, including databases not controlled by the medical practitioner. In some embodiments, the information is maintained in electronic form. In some embodiments, the information can be stored in a memory or other computer readable medium, such as RAM, ROM, EEPROM, flash memory, computer chips, digital video discs (DVD), compact discs (CDs), hard disk drives (HDD), magnetic tape, etc.

[0060] In some embodiments, methods of detecting the presence lung cancer are provided. In some embodiments, methods of diagnosing lung cancer are provided. In some embodiments, the method comprises obtaining a sample from a subject and providing the sample to a laboratory for detection of at least one target RNA level in the sample. In some embodiments, the method further comprises receiving a communication from the laboratory that indicates the at least one target RNA level in the sample. In some embodiments, lung cancer is present if the level of at least one target RNA in the

sample is greater than a normal level of the at least one target RNA. A “laboratory,” as used herein, is any facility that detects the level of at least one target RNA in a sample by any method, including the methods described herein, and communicates the level to a medical practitioner. In some embodiments, a laboratory is under the control of a medical practitioner. In some embodiments, a laboratory is not under the control of the medical practitioner.

[0061] When a laboratory communicates the level of at least one target RNA to a medical practitioner, in some embodiments, the laboratory communicates a numerical value representing the level of at least one target RNA in the sample, with or without providing a numerical value for a normal level. In some embodiments, the laboratory communicates the level of at least one target RNA by providing a qualitative value, such as “high,” “elevated,” etc.

[0062] As used herein, when a method relates to detecting lung cancer, determining the presence of lung cancer, and/or diagnosing lung cancer, the method includes activities in which the steps of the method are carried out, but the result is negative for the presence of lung cancer. That is, detecting, determining, and diagnosing lung cancer include instances of carrying out the methods that result in either positive or negative results (e.g., whether target RNA levels are normal or greater than normal).

[0063] As used herein, the term “subject” means a human. In some embodiments, the methods described herein may be used on samples from non-human animals.

[0064] The common, or coordinate, expression of target RNAs that are physically proximal to one another in the genome permits the informative use of such chromosome-proximal target RNAs in methods herein.

[0065] Table 3 identifies the chromosomal location of each of the 397 target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence that is identically present in one of SEQ ID NOs: 1 to 397 in Tables 1 and 2. Table 22 identifies the chromosomal location of the target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence that is identically present in one of SEQ ID NOs: 1063 to 1210 in Tables 18 and 20. Table 25 identifies the chromosomal location of the target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence

that is identically present in one of SEQ ID NOs: 1363 to 1707 in Table 23. Table 29 identifies the chromosomal location of the target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence that is identically present in one of SEQ ID NOs: 2064 to 2183 in Tables 27 and 28. Table 31 identifies the chromosomal location of the target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence that is identically present in one of SEQ ID NOs: 2312 to 2452 in Table 30. Thus, in some embodiments, the level of expression of one or more target RNAs located within about 1 kilobase (kb), within about 2 kb, within about 5 kb, within about 10 kb, within about 20 kb, within about 30 kb, within about 40 kb, and even within about 50 kb of the chromosomal locations in Tables 3, 22, 25, 29, and 31 is detected in lieu of, or in addition to, measurement of expression of the respective tabulated target RNAs in the methods described herein. *See* Baskerville, S. and Bartel D.P. (2005) *RNA* 11:241-247.

[0066] In some embodiments, in combination with detecting one or more target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and/or detecting one or more target RNAs comprising at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 and/or detecting one or more target RNAs that comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, methods herein further comprise detecting the level(s) of expression of at least one microRNA from the human miRNome.

[0067] In some embodiments, at least one target RNA is capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least one target RNA comprises at least 15 contiguous nucleotides that are complementary to at least a portion of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least one target RNA comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and

2692. In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[0068] In some embodiments, more than one target RNA is detected simultaneously in a single reaction. In some embodiments, at least 2, at least 3, at least 5, or at least 10 target RNAs are detected simultaneously in a single reaction. In some embodiments, all target RNAs are detected simultaneously in a single reaction.

[0069] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken. In some embodiments, an increased level of one or more target RNAs that comprise at least 15 contiguous nucleotides that are complementary to at least a portion of a sequence selected from SEQ ID NO: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken. In some embodiments, an increased level of one or more target RNAs that comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken. In some embodiments, a decreased level of one or more target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken. In some embodiments, a decreased level of one or more target RNAs that comprise at least 15 contiguous nucleotides that are complementary to at least a portion of a sequence selected from SEQ ID NO: 1 to 397, 1363 to 1707, and 2312 to 2452 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken. In some embodiments, a decreased level of one or more target RNAs that comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 in a sample is indicative of the presence of lung cancer in an individual from whom the sample has been taken.

[0070] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 6 is indicative of the presence of non-small cell lung cancer.

[0071] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 7 is indicative of squamous cell carcinoma.

[0072] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 8 is indicative of adenocarcinoma.

[0073] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 9 is indicative of aggressive lung cancer.

[0074] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 32 or 33 is indicative of lung cancer.

[0075] In some embodiments, a decreased level of one or more target RNAs capable of specifically hybridizing to a polynucleotide sequence in Table 34 is indicative of lung cancer.

[0076] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 15, 26, 27, 30, 129, 164, 184, 191, 196, 197, 205, 207, 214, 219, 225, 246, and 248 is indicative of non-small cell lung cancer.

[0077] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 15, 26, 27 and 191 and decreased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 92 and 171 is indicative of squamous cell carcinoma or adenocarcinoma.

[0078] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 4, 36, 50,

93, 122, 125, 139, 140, 144, 146, 159, 226, 239 and 241 is indicative of squamous cell carcinoma or adenocarcinoma.

[0079] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 19, 27, 33, 48, 55, 72, 73, 94, 101, 105, 112, 117, 130, 131, 133, 134, 135, 143, 155, 158, 160, 161, 163, 165, 221, 238, 240 and 246 is indicative of squamous cell carcinoma.

[0080] In some embodiments, a decreased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 19, 27, 33, 48, 55, 72, 73, 94, 101, 105, 112, 117, 130, 131, 133, 134, 135, 143, 155, 158, 160, 161, 163, 165, 221, 238, 240 and 246 is indicative of adenocarcinoma.

[0081] In some embodiments, an increased level of one or more target RNAs capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 27, 72, 73, 161 or 246 is indicative of an aggressive form of adenocarcinoma.

4.1.2. Exemplary controls

[0082] In some embodiments, a normal level (a “control”) for each target RNA can be determined as an average level or range that is characteristic of normal human lung cells or other reference material, against which the level measured in the sample can be compared. The determined average or range of target RNA in normal subjects can be used as a benchmark for detecting above-normal or below-normal levels of target RNA indicative of lung cancer. In some embodiments, normal levels of target RNA can be determined using individual or pooled RNA-containing samples from one or more individuals, such as from normal lung tissue from patients undergoing surgical resection for stage I, II or IIIA non-small cell lung cancer.

[0083] In some embodiments, determining a normal level of expression of a target RNA comprises detecting a complex comprising a probe hybridized to a nucleic acid selected from a target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA. That is, in some embodiments, a normal level of expression can be determined by detecting a DNA amplicon of the target RNA, or a complement of the target RNA rather than the target RNA itself. In some embodiments, a normal level of such a complex is determined and used as a control. The normal level of the complex, in

some embodiments, correlates to the normal level of the target RNA. Thus, when a normal level of a target is discussed herein, that level can, in some embodiments, be determined by detecting such a complex.

[0084] In some embodiments, a control comprises RNA from cells of a single individual, *e.g.*, from normal tissue from a patient undergoing surgical resection for stage I, II or IIIA non-small cell lung cancer. In some embodiments, the control is drawn from anatomically and/or cytologically normal areas of the lung of the individual from whom the test sample was obtained. In some embodiments, a control comprises RNA from a pool of cells from multiple individuals. In some embodiments, a control comprises RNA from a pool of blood, such as whole blood or serum, from multiple individuals. In some embodiments, a control comprises commercially-available human RNA, such as, for example, human lung total RNA (Ambion; AM7968). In some embodiments, a normal level or normal range has already been predetermined prior to testing a sample for an elevated level.

[0085] In some embodiments, the normal level of target RNA can be determined from one or more continuous cell lines, typically cell lines previously shown to have expression levels of the at least one target RNA that approximate the level of expression in normal human lung cells.

[0086] In some embodiments, a method comprises detecting the level of expression of at least one target RNA. In some embodiments, a method further comprises comparing the level of expression of at least one target RNA to a normal level of expression of the at least one target RNA. In some embodiments, a method further comprises comparing the level of expression of at least one target RNA to a control level of expression of the at least one target RNA. A control level of expression of the at least one target RNA is, in some embodiments, the level of expression of the at least one target RNA in a normal cell. In some such embodiments, a control level may be referred to as a normal level. In some embodiments, a greater level of expression of the at least one target RNA relative to the level of expression of the at least one target RNA in a normal cell indicates lung cancer. In some embodiments, a reduced level of expression of the at least one target RNA relative to the level of expression of the at least one target RNA in a normal cell indicates lung cancer.

[0087] In some embodiments, the level of expression of the at least one target RNA is compared to a reference level of expression, e.g., from a patient with a confirmed lung cancer. In some such embodiments, a similar level of expression of the at least one target RNA relative to the reference sample indicates lung cancer.

[0088] In some embodiments, a level of expression of at least one target RNA that is at least about two-fold greater than a normal level of expression of the respective at least one target RNA indicates the presence of lung cancer. In some embodiments, a level of expression of at least one target RNA that is at least about two-fold greater than the level of the respective at least one target RNA in a control sample comprised of normal cells indicates the presence of a lung cancer. In various embodiments, a level of expression of at least one target RNA that is at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold greater than the level of expression of the respective at least one target RNA in a control sample comprised of normal cells indicates the presence of lung cancer. In various embodiments, a level of expression of at least one target RNA that is at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold greater than a normal level of expression of the at least one target RNA indicates the presence of lung cancer.

[0089] In some embodiments, a level of expression of at least one target RNA that is reduced by at least about two-fold relative to a normal level of expression of the respective at least one target RNA indicates the presence of lung cancer. In some embodiments, a level of expression of at least one target RNA that is reduced by at least about two-fold as compared to the level of the respective at least one target RNA in a control sample comprised of normal cells indicates the presence of a lung cancer. In various embodiments, a level of expression of at least one target RNA that is reduced by at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold as compared to the level of expression of the respective at least one target RNA in a control sample comprised of normal cells indicates the presence of lung cancer. In various embodiments, a level of expression of at least one target RNA that is reduced by at least

about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold as compared to a normal level of expression of the at least one target RNA indicates the presence of lung cancer.

[0090] In some embodiments, a control level of expression of a target RNA is determined contemporaneously, such as in the same assay or batch of assays, as the level of expression of the target RNA in a sample. In some embodiments, a control level of expression of a target RNA is not determined contemporaneously as the level of expression of the target RNA in a sample. In some such embodiments, the control level of expression has been determined previously.

[0091] In some embodiments, the level of expression of a target RNA is not compared to a control level of expression, for example, when it is known that the target RNA is expressed at very low levels, or not at all, in normal cells. In some such embodiments, detection of a high level of the target RNA in a sample is indicative of lung cancer. Alternatively, if the target RNA is known to be expressed at high levels in normal cells, then detection of a very low level of the target RNA in a sample is indicative of lung cancer.

4.1.3. Exemplary methods of preparing RNAs

[0092] Target RNA can be prepared by any appropriate method. Total RNA can be isolated by any method, including, but not limited to, the protocols set forth in Wilkinson, M. (1988) Nucl. Acids Res. 16(22):10,933; and Wilkinson, M. (1988) Nucl. Acids Res. 16(22): 10934, or by using commercially-available kits or reagents, such as the TRIzol® reagent (Invitrogen™), Total RNA Extraction Kit (iNtRON Biotechnology), Total RNA Purification Kit (Norgen Biotek Corp.), RNAqueous™ (Ambion), MagMAX™ (Ambion), RecoverAll™ (Ambion), RNeasy (Qiagen), etc.

[0093] In some embodiments, small RNAs are isolated or enriched. In some embodiments “small RNA” refers to RNA molecules smaller than about 200 nucleotides (nt) in length. In some embodiments, “small RNA” refers to RNA molecules smaller than about 100 nt, smaller than about 90 nt, smaller than about 80 nt, smaller than about 70 nt, smaller than about 60 nt, smaller than about 50 nt, or smaller than about 40 nt.

[0094] Enrichment of small RNAs can be accomplished by method. Such methods include, but are not limited to, methods involving organic extraction followed by adsorption of nucleic acid molecules on a glass fiber filter using specialized binding and wash solutions, and methods using spin column purification. Enrichment of small RNAs may be accomplished using commercially-available kits, such as mirVana™ Isolation Kit (Applied Biosystems), mirPremier™ microRNA Isolation Kit (Sigma-Aldrich), PureLink™ miRNA Isolation Kit (Invitrogen), miRCURY™ RNA isolation kit (Exiqon), microRNA Purification Kit (Norgen Biotek Corp.), miRNeasy kit (Qiagen), etc. In some embodiments, purification can be accomplished by the TRIzol® (Invitrogen) method, which employs a phenol/isothiocyanate solution to which chloroform is added to separate the RNA-containing aqueous phase. Small RNAs are subsequently recovered from the aqueous by precipitation with isopropyl alcohol. In some embodiments, small RNAs can be purified using chromatographic methods, such as gel electrophoresis using the flashPAGE™ Fractionator available from Applied Biosystems.

[0095] In some embodiments, small RNA is isolated from other RNA molecules to enrich for target RNAs, such that the small RNA fraction (e.g., containing RNA molecules that are 200 nucleotides or less in length, such as less than 100 nucleotides in length, such as less than 50 nucleotides in length, such as from about 10 to about 40 nucleotides in length) is substantially pure, meaning it is at least about 80%, 85%, 90%, 95% pure or more, but less than 100% pure, with respect to larger RNA molecules. Alternatively, enrichment of small RNA can be expressed in terms of fold-enrichment. In some embodiments, small RNA is enriched by about, at least about, or at most about 5X, 10X, 20X, 30X, 40X, 50X, 60X, 70X, 80X, 90X, 100X, 110X, 120X, 130X, 140X, 150X, 160X, 170X, 180X, 190X, 200X, 210X, 220X, 230X, 240X, 250X, 260X, 270X, 280X, 290X, 300X, 310X, 320X, 330X, 340X, 350X, 360X, 370X, 380X, 390X, 400X, 410X, 420X, 430X, 440X, 450X, 460X, 470X, 480X, 490X, 500X, 600X, 700X, 800X, 900X, 1000X, 1100X, 1200X, 1300X, 1400X, 1500X, 1600X, 1700X, 1800X, 1900X, 2000X, 3000X, 4000X, 5000X, 6000X, 7000X, 8000X, 9000X, 10,000X or more, or any range derivable therein, with respect to the concentration of larger RNAs in an RNA isolate or total RNA in a sample.

[0096] In yet other embodiments, expression is measured in a sample in which RNA has not first been purified from the cells.

[0097] In some embodiments, RNA is modified before target RNAs are detected. In some embodiments, the modified RNA is total RNA. In other embodiments, the modified RNA is small RNA that has been purified from total RNA or from cell lysates, such as RNA less than 200 nucleotides in length, such as less than 100 nucleotides in length, such as less than 50 nucleotides in length, such as from about 10 to about 40 nucleotides in length. RNA modifications that can be utilized in the methods described herein include, but are not limited to, the addition of a poly-dA or a poly-dT tail, which can be accomplished chemically or enzymatically, and/or the addition of a small molecule, such as biotin.

[0098] In some embodiments, one or more target RNAs are reverse transcribed. In some embodiments, where present, RNA is modified when it is reverse transcribed, such as when a poly-dA or a poly-dT tail is added to the cDNA during reverse transcription. In other embodiments, RNA is modified before it is reverse transcribed. In some embodiments, total RNA is reverse transcribed. In other embodiments, small RNAs are isolated or enriched before the RNA is reverse transcribed.

[0099] When a target RNA is reverse transcribed, a complement of the target RNA is formed. In some embodiments, the complement of the target RNA is detected rather than the target RNA itself (or a DNA copy thereof). Thus, when the methods discussed herein indicate that a target RNA is detected, or the level of a target RNA is determined, such detection or determination may be carried out on a complement of the target RNA instead of, or in addition to, the target RNA itself. In some embodiments, when the complement of the target RNA is detected rather than the target RNA, a probe is used that is complementary to the complement of the target RNA. In such embodiments, the probe comprises at least a portion that is identical in sequence to the target RNA, although it may contain thymidine in place of uridine, and/or comprise other modified nucleotides.

[00100] In some embodiments, the method of detecting one or more target RNAs comprises amplifying cDNA complementary to said target RNA. Such amplification can be accomplished by any method. Exemplary methods include, but are

not limited to, real time PCR, endpoint PCR, and amplification using T7 polymerase from a T7 promoter annealed to a cDNA, such as provided by the SenseAmp Plus™ Kit available at Implen, Germany.

[00101] When a target RNA or a cDNA complementary to a target RNA is amplified, in some embodiments, a DNA amplicon of a target RNA is formed. A DNA amplicon may be single stranded or double-stranded. In some embodiments, when a DNA amplicon is single-stranded, the sequence of the DNA amplicon is related to the target RNA in either the sense or antisense orientation. In some embodiments, the DNA amplicon of the target RNA is detected rather than the target RNA itself. Thus, when the methods discussed herein indicate that a target RNA is detected, or the level of a target RNA is determined, such detection or determination may be carried out on a DNA amplicon of the target RNA instead of, or in addition to, the target RNA itself. In some embodiments, when the DNA amplicon of the target RNA is detected rather than the target RNA, a probe is used that is complementary to the complement of the target RNA. In some embodiments, when the DNA amplicon of the target RNA is detected rather than the target RNA, a probe is used that is complementary to the target RNA. Further, in some embodiments, multiple probes may be used, and some probes may be complementary to the target RNA and some probes may be complementary to the complement of the target RNA.

[00102] In some embodiments, the method of detecting one or more target RNAs comprises RT-PCR, as described below. In some embodiments, detecting one or more target RNAs comprises real-time monitoring of an RT-PCR reaction, which can be accomplished by any method. Such methods include, but are not limited to, the use of TaqMan®, Molecular beacon, or Scorpion probes (i.e., FRET probes) and the use of intercalating dyes, such as SYBR green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc.

4.1.4. Exemplary analytical methods

[00103] As described above, methods are presented for detecting lung cancer in a sample from a patient. In some embodiments, the method comprises detecting a level of expression of at least one target RNA capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210,

2064 to 2183, 2673 to 2680, and 2689 that is greater in the sample than a normal level of expression of the at least one target RNA in a control sample, such as a sample derived from normal lung cells. In some embodiments, a method comprises detecting a level of one or more target RNAs that comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689 that is greater in the sample than a normal level of expression of the at least one target RNA in a control sample. In some embodiments, a method comprises detecting a level of one or more target RNAs that comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 that is greater in the sample than a normal level of expression of the at least one target RNA in a control sample.

[00104] In some embodiments, the method comprises detecting a level of expression of at least one target RNA capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452 that is reduced in the sample relative to a normal level of expression of the at least one target RNA in a control sample, such as a sample derived from normal lung cells. In some embodiments, a method comprises detecting a level of one or more target RNAs that comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452 that is reduced in the sample relative to a normal level of expression of the at least one target RNA in a control sample. In some embodiments, a method comprises detecting a level of one or more target RNAs that comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 that is reduced in the sample relative to a normal level of expression of the at least one target RNA in a control sample.

[00105] In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00106] In some embodiments, such as those described above, the method further comprises detecting a level of expression of at least one target RNA of the human miRNome that does not specifically hybridize to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to

2452, 2673 to 2680, and 2689 and does not comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692, that is altered in the sample relative to a normal level of expression of the at least one target RNA in a control sample. As used herein, the term “human miRNome” refers to all microRNA genes in a human cell and the mature microRNAs produced therefrom.

[00107] Any analytical procedure capable of permitting specific and quantifiable (or semi-quantifiable) detection of the desired at least one target RNA may be used in the methods herein presented. Such analytical procedures include, but are not limited to, the microarray methods set forth in Examples 1, 2, 4, and 5, the microbead methods set forth in Example 3, and methods known to those skilled in the art.

[00108] In some embodiments, detection of a target RNA comprises forming a complex comprising a polynucleotide that is complementary to a target RNA or to a complement thereof, and a nucleic acid selected from the target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA. Thus, in some embodiments, the polynucleotide forms a complex with a target RNA. In some embodiments, the polynucleotide forms a complex with a complement of the target RNA, such as a cDNA that has been reverse transcribed from the target RNA. In some embodiments, the polynucleotide forms a complex with a DNA amplicon of the target RNA. When a double-stranded DNA amplicon is part of a complex, as used herein, the complex may comprise one or both strands of the DNA amplicon. Thus, in some embodiments, a complex comprises only one strand of the DNA amplicon. In some embodiments, a complex is a triplex and comprises the polynucleotide and both strands of the DNA amplicon. In some embodiments, the complex is formed by hybridization between the polynucleotide and the target RNA, complement of the target RNA, or DNA amplicon of the target RNA. The polynucleotide, in some embodiments, is a primer or probe.

[00109] In some embodiments, a method comprises detecting the complex. In some embodiments, the complex does not have to be associated at the time of detection. That is, in some embodiments, a complex is formed, the complex is then dissociated or destroyed in some manner, and components from the complex are detected. An example of such a system is a TaqMan® assay. In some embodiments, when the

polynucleotide is a primer, detection of the complex may comprise amplification of the target RNA, a complement of the target RNA, or a DNA amplicon of a target RNA.

[00110] In some embodiments the analytical method used for detecting at least one target RNA in the methods set forth herein includes real-time quantitative RT-PCR. See Chen, C. et al. (2005) *Nucl. Acids Res.* 33:e179 and PCT Publication No. WO 2007/117256, which are incorporated herein by reference in its entirety. In some embodiments, the analytical method used for detecting at least one target RNA includes the method described in U.S. Publication No. US2009/0123912 A1, which is incorporated herein by reference in its entirety. In an exemplary method described in that publication, an extension primer comprising a first portion and second portion, wherein the first portion selectively hybridizes to the 3' end of a particular microRNA and the second portion comprises a sequence for universal primer, is used to reverse transcribe the microRNA to make a cDNA. A reverse primer that selectively hybridizes to the 5' end of the microRNA and a universal primer are then used to amplify the cDNA in a quantitative PCR reaction.

[00111] In some embodiments, the analytical method used for detecting at least one target RNA includes the use of a TaqMan® probe. In some embodiments, the analytical method used for detecting at least one target RNA includes a TaqMan® assay, such as the TaqMan® MicroRNA Assays sold by Applied Biosystems, Inc. In an exemplary TaqMan® assay, total RNA is isolated from the sample. In some embodiments, the assay can be used to analyze about 10 ng of total RNA input sample, such as about 9 ng of input sample, such as about 8 ng of input sample, such as about 7 ng of input sample, such as about 6 ng of input sample, such as about 5 ng of input sample, such as about 4 ng of input sample, such as about 3 ng of input sample, such as about 2 ng of input sample, and even as little as about 1 ng of input sample containing microRNAs.

[00112] The TaqMan® assay utilizes a stem-loop primer that is specifically complementary to the 3'-end of a target RNA. In an exemplary TaqMan® assay, hybridizing the stem-loop primer to the target RNA is followed by reverse transcription of the target RNA template, resulting in extension of the 3' end of the primer. The result of the reverse transcription is a chimeric (DNA) amplicon with the stem-loop primer sequence at the 5' end of the amplicon and the cDNA of the target RNA at the 3' end.

Quantitation of the target RNA is achieved by real time RT-PCR using a universal reverse primer having a sequence that is complementary to a sequence at the 5' end of all stem-loop target RNA primers, a target RNA-specific forward primer, and a target RNA sequence-specific TaqMan® probe.

[00113] The assay uses fluorescence resonance energy transfer (“FRET”) to detect and quantitate the synthesized PCR product. Typically, the TaqMan® probe comprises a fluorescent dye molecule coupled to the 5'-end and a quencher molecule coupled to the 3'-end, such that the dye and the quencher are in close proximity, allowing the quencher to suppress the fluorescence signal of the dye via FRET. When the polymerase replicates the chimeric amplicon template to which the TaqMan® probe is bound, the 5'-nuclease of the polymerase cleaves the probe, decoupling the dye and the quencher so that FRET is abolished and a fluorescence signal is generated. Fluorescence increases with each RT-PCR cycle proportionally to the amount of probe that is cleaved.

[00114] Additional exemplary methods for RNA detection and/or quantification are described, e.g., in U.S. Publication No. US 2007/0077570 (Lao et al.), PCT Publication No. WO 2007/025281 (Tan et al.), U.S. Publication No. US2007/0054287 (Bloch), PCT Publication No. WO2006/0130761 (Bloch), and PCT Publication No. WO 2007/011903 (Lao et al.), which are incorporated by reference herein in their entireties for any purpose.

[00115] In some embodiments, quantitation of the results of real-time RT-PCR assays is done by constructing a standard curve from a nucleic acid of known concentration and then extrapolating quantitative information for target RNAs of unknown concentration. In some embodiments, the nucleic acid used for generating a standard curve is an RNA (e.g., microRNA) of known concentration. In some embodiments, the nucleic acid used for generating a standard curve is a purified double-stranded plasmid DNA or a single-stranded DNA generated in vitro.

[00116] In some embodiments, where the amplification efficiencies of the target nucleic acids and the endogenous reference are approximately equal, quantitation is accomplished by the comparative Ct (cycle threshold, e.g., the number of PCR cycles required for the fluorescence signal to rise above background) method. Ct values are inversely proportional to the amount of nucleic acid target in a sample. In some

embodiments, Ct values of the target RNA of interest can be compared with a control or calibrator, such as RNA (e.g., microRNA) from normal tissue. In some embodiments, the Ct values of the calibrator and the target RNA samples of interest are normalized to an appropriate endogenous housekeeping gene.

[00117] In addition to the TaqMan® assays, other real-time RT-PCR chemistries useful for detecting and quantitating PCR products in the methods presented herein include, but are not limited to, Molecular Beacons, Scorpion probes and intercalating dyes, such as SYBR Green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc., which are discussed below.

[00118] In some embodiments, real-time RT-PCR detection is performed specifically to detect and quantify the expression of a single target RNA. The target RNA, in some embodiments, is selected from a target RNA capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the target RNA comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. The target RNA, in some embodiments, comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs.: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Table 6. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Table 7. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Table 8. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Table 9. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Tables 32 and 33. In some embodiments, the target RNA specifically hybridizes to a nucleic acid comprising a sequence selected from the probe sequences in Table 34.

[00119] In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00120] In various embodiments, real-time RT-PCR detection is utilized to detect, in a single multiplex reaction, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8 target RNAs. At least one target RNA, in some embodiments, is capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least one target RNA comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, at least one target RNA comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00121] In some embodiments, the method comprises detecting expression in a multiplex RT-PCR reaction of at least 2, at least 3, at least 5, at least 10, or at least 15 target RNAs, wherein each target RNA is capable of specifically hybridizing to a probe sequence in Table 6. In some embodiments, the method comprises detecting expression, using a single multiplex RT-PCR reaction, of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 12, at least 15, at least 20, or at least 25 target RNAs, wherein each target RNA is capable of specifically hybridizing to a probe sequence in Table 7. In some embodiments, the method comprises detecting expression, using a single multiplex RT-PCR reaction, of at least 2, at least 3, at least 4, at least 5, at least 6, or at least 7 target RNAs, wherein each target RNA is capable of specifically hybridizing to a probe sequence in Table 8. In some embodiments, the method comprises detecting expression in a multiplex RT-PCR reaction of at least 2, at least 3, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, or at least 50 target RNAs, wherein each target RNA is capable of specifically hybridizing to a probe sequence in Table 9. In some embodiments, the method comprises detecting expression in a multiplex RT-PCR reaction of at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, or at least target RNAs, wherein each target

RNA is capable of specifically hybridizing to a probe sequence in one of Tables 32 or 33. In some embodiments, the method comprises detecting expression in a multiplex RT-PCR reaction of at least two, at least five, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, or at least 60 target RNAs, wherein each target RNA is capable of specifically hybridizing to a probe sequence in Table 34.

[00122] In some multiplex embodiments, a plurality of probes, such as TaqMan® probes, each specific for a different RNA target, is used. In some embodiments, each target RNA-specific probe is spectrally distinguishable from the other probes used in the same multiplex reaction.

[00123] In some embodiments, quantitation of real-time RT PCR products is accomplished using a dye that binds to double-stranded DNA products, such as SYBR Green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc. In some embodiments, the assay is the QuantiTect SYBR Green PCR assay from Qiagen. In this assay, total RNA is first isolated from a sample. Total RNA is subsequently poly-adenylated at the 3'-end and reverse transcribed using a universal primer with poly-dT at the 5'-end. In some embodiments, a single reverse transcription reaction is sufficient to assay multiple target RNAs. Real-time RT-PCR is then accomplished using target RNA-specific primers and an miScript Universal Primer, which comprises a poly-dT sequence at the 5'-end. SYBR Green dye binds non-specifically to double-stranded DNA and upon excitation, emits light. In some embodiments, buffer conditions that promote highly-specific annealing of primers to the PCR template (e.g., available in the QuantiTect SYBR Green PCR Kit from Qiagen) can be used to avoid the formation of non-specific DNA duplexes and primer dimers that will bind SYBR Green and negatively affect quantitation. Thus, as PCR product accumulates, the signal from SYBR Green increases, allowing quantitation of specific products.

[00124] Real-time RT-PCR is performed using any RT-PCR instrumentation available in the art. Typically, instrumentation used in real-time RT-PCR data collection and analysis comprises a thermal cycler, optics for fluorescence excitation and emission collection, and optionally a computer and data acquisition and analysis software.

[00125] In some embodiments, the analytical method used in the methods described herein is a DASL® (cDNA-mediated Annealing, Selection, Extension, and Ligation) Assay, such as the MicroRNA Expression Profiling Assay available from Illumina, Inc. (See <http://www.illumina.com/downloads/MicroRNAAssayWorkflow.pdf>). In some embodiments, total RNA is isolated from a sample to be analyzed by any method. Additionally, in some embodiments, small RNAs are isolated from a sample to be analyzed by any method. Total RNA or isolated small RNAs may then be polyadenylated (> 18 A residues are added to the 3'-ends of the RNAs in the reaction mixture). The RNA is reverse transcribed using a biotin-labeled DNA primer that comprises from the 5' to the 3' end, a sequence that includes a PCR primer site and a poly-dT region that binds to the poly-dA tail of the sample RNA. The resulting biotinylated cDNA transcripts are then hybridized to a solid support via a biotin-streptavidin interaction and contacted with one or more target RNA-specific polynucleotides. The target RNA-specific polynucleotides comprise, from the 5'-end to the 3'-end, a region comprising a PCR primer site, region comprising an address sequence, and a target RNA-specific sequence.

[00126] In some DASL® embodiments, the target RNA-specific sequence comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides having a sequence identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the target RNA-specific sequence comprises a probe sequence that is complementary to at least a portion of a microRNA of the human miRNome.

[00127] After hybridization, the target RNA-specific polynucleotide is extended, and the extended products are then eluted from the immobilized cDNA array. A second PCR reaction using a fluorescently-labeled universal primer generates a fluorescently-labeled DNA comprising the target RNA-specific sequence. The labeled PCR products are then hybridized to a microbead array for detection and quantitation.

[00128] In some embodiments, the analytical method used for detecting and quantifying the expression of the at least one target RNA in the methods described

herein is a bead-based flow cytometric assay. See Lu J. et al. (2005) Nature 435:834-838, which is incorporated herein by reference in its entirety. An example of a bead-based flow cytometric assay is the xMAP® technology of Luminex, Inc. (See <http://www.luminexcorp.com/technology/index.html>). In some embodiments, total RNA is isolated from a sample and is then labeled with biotin. The labeled RNA is then hybridized to target RNA-specific capture probes (e.g., FlexmiR™ products sold by Luminex, Inc. at <http://www.luminexcorp.com/products/assays/index.html>) that are covalently bound to microbeads, each of which is labeled with 2 dyes having different fluorescence intensities. A streptavidin-bound reporter molecule (e.g., streptavidin-phycoerythrin, also known as “SAPE”) is attached to the captured target RNA and the unique signal of each bead is read using flow cytometry. In some embodiments, the RNA sample (total RNA or enriched small RNAs) is first polyadenylated, and is subsequently labeled with a biotinylated 3DNA™ dendrimer (i.e., a multiple-arm DNA with numerous biotin molecules bound thereto), such as those sold by Marligen Biosciences as the Vantage™ microRNA Labeling Kit, using a bridging polynucleotide that is complementary to the 3'-end of the poly-dA tail of the sample RNA and to the 5'-end of the polynucleotide attached to the biotinylated dendrimer. The streptavidin-bound reporter molecule is then attached to the biotinylated dendrimer before analysis by flow cytometry. See <http://www.marligen.com/vantage-microrna-labeling-kit.html>. In some embodiments, biotin-labeled RNA is first exposed to SAPE, and the RNA/SAPE complex is subsequently exposed to an anti-phycoerythrin antibody attached to a DNA dendrimer, which can be bound to as many as 900 biotin molecules. This allows multiple SAPE molecules to bind to the biotinylated dendrimer through the biotin-streptavidin interaction, thus increasing the signal from the assay.

[00129] In some embodiments, the analytical method used for detecting and quantifying the expression of the at least one target RNA in the methods described herein is by gel electrophoresis and detection with labeled probes (e.g., probes labeled with a radioactive or chemiluminescent label), such as by Northern blotting. In some embodiments, total RNA is isolated from the sample, and then is size-separated by SDS polyacrylamide gel electrophoresis. The separated RNA is then blotted onto a membrane and hybridized to radiolabeled complementary probes. In some embodiments, exemplary probes contain one or more affinity-enhancing nucleotide analogs as discussed below,

such as locked nucleic acid (“LNA”) analogs, which contain a bicyclic sugar moiety instead of deoxyribose or ribose sugars. See, e.g., Várallyay, E. et al. (2008) *Nature Protocols* 3(2):190-196, which is incorporated herein by reference in its entirety. In some embodiments, the total RNA sample can be further purified to enrich for small RNAs. In some embodiments, target RNAs can be amplified by, e.g., rolling circle amplification using a long probe that is complementary to both ends of a target RNA (“padlocked probes”), ligation to circularize the probe followed by rolling circle replication using the target RNA hybridized to the circularized probe as a primer. See, e.g., Jonstrup, S.P. et al. (2006) *RNA* 12:1-6, which is incorporated herein by reference in its entirety. The amplified product can then be detected and quantified using, e.g., gel electrophoresis and Northern blotting.

[00130] In alternative embodiments, labeled probes are hybridized to isolated total RNA in solution, after which the RNA is subjected to rapid ribonuclease digestion of single-stranded RNA, e.g., unhybridized portions of the probes or unhybridized target RNAs. In these embodiments, the ribonuclease treated sample is then analyzed by SDS-PAGE and detection of the radiolabeled probes by, e.g., Northern blotting. See mirVana™ miRNA Detection Kit sold by Applied Biosystems, Inc. product literature at <http://www.ambion.com/catalog/CatNum.php?1552>.

[00131] In some embodiments, the analytical method used for detecting and quantifying the at least one target RNA in the methods described herein is by hybridization to a microarray. See, e.g., Liu, C.G. et al. (2004) *Proc. Nat’l Acad. Sci. USA* 101:9740-9744; Lim, L.P. et al. (2005) *Nature* 433:769-773, each of which is incorporated herein by reference in its entirety, and Examples 1, 2, 4, and 5.

[00132] In some embodiments, detection and quantification of a target RNA using a microarray is accomplished by surface plasmon resonance. See, e.g., *Nanotech News* (2006), available at http://nano.cancer.gov/news_center/nanotech_news_2006-10-30b.asp. In these embodiments, total RNA is isolated from a sample being tested. Optionally, the RNA sample is further purified to enrich the population of small RNAs. After purification, the RNA sample is bound to an addressable microarray containing probes at defined locations on the microarray. Nonlimiting exemplary probes include probes comprising sequences

set forth in SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Exemplary probes also include, but are not limited to, probes comprising a region that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. Exemplary probes also include, but are not limited to, probes comprising at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the probes contain one or more affinity-enhancing nucleotide analogs as discussed below, such as locked nucleic acid (“LNA”) nucleotide analogs. After hybridization to the microarray, the RNA that is hybridized to the array is first polyadenylated, and the array is then exposed to gold particles having poly-dT bound to them. The amount of bound target RNA is quantitated using surface plasmon resonance.

[00133] In some embodiments, microarrays are utilized in a RNA-primed, Array-based Klenow Enzyme (“RAKE”) assay. See Nelson, P.T. et al. (2004) *Nature Methods* 1(2):1-7; Nelson, P.T. et al. (2006) *RNA* 12(2):1-5, each of which is incorporated herein by reference in its entirety. In some embodiments, total RNA is isolated from a sample. In some embodiments, small RNAs are isolated from a sample. The RNA sample is then hybridized to DNA probes immobilized at the 5'-end on an addressable array. The DNA probes comprise, in some embodiments, from the 5'-end to the 3'-end, a first region comprising a “spacer” sequence which is the same for all probes, a second region comprising three thymidine-containing nucleosides, and a third region comprising a sequence that is complementary to a target RNA of interest.

[00134] Exemplary target RNAs of interest include, but are not limited to, target RNAs capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689; target RNAs comprising a region that is identical to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and target RNAs comprising a region that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Target RNAs also include target RNAs in the miRNome that do not specifically hybridize to a

nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00135] After the sample is hybridized to the array, it is exposed to exonuclease I to digest any unhybridized probes. The Klenow fragment of DNA polymerase I is then applied along with biotinylated dATP, allowing the hybridized target RNAs to act as primers for the enzyme with the DNA probe as template. The slide is then washed and a streptavidin-conjugated fluorophore is applied to detect and quantitate the spots on the array containing hybridized and Klenow-extended target RNAs from the sample.

[00136] In some embodiments, the RNA sample is reverse transcribed. In some embodiments, the RNA sample is reverse transcribed using a biotin/poly-dA random octamer primer. When than primer is used, the RNA template is digested and the biotin-containing cDNA is hybridized to an addressable microarray with bound probes that permit specific detection of target RNAs. In some embodiments, the microarray includes at least one probe comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides identically present in, or complementary to a region of, a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. After hybridization of the cDNA to the microarray, the microarray is exposed to a streptavidin-bound detectable marker, such as a fluorescent dye, and the bound cDNA is detected. See Liu C.G. et al. (2008) Methods 44:22-30, which is incorporated herein by reference in its entirety.

[00137] In some embodiments, target RNAs are detected and quantified in an ELISA-like assay using probes bound in the wells of microtiter plates. See Mora J.R. and Getts R.C. (2006) BioTechniques 41:420-424 and supplementary material in BioTechniques 41(4):1-5; U.S. Patent Publication No. 2006/0094025 to Getts et al., each of which is incorporated by reference herein in its entirety. In these embodiments, a sample of RNA that is enriched in small RNAs is either polyadenylated, or is reverse

transcribed and the cDNA is polyadenylated. The RNA or cDNA is hybridized to probes immobilized in the wells of a microtiter plates, wherein each of the probes comprises a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, or a sequence such as one or more sequences of target RNAs (or the reverse complement thereof) of the human miRNome, depending on whether RNA or cDNA is hybridized to the array. In some embodiments, the hybridized RNAs are labeled using a capture sequence, such as a DNA dendrimer (such as those available from Genisphere, Inc., http://www.genisphere.com/about_3dna.html) that is labeled with a plurality of biotin molecules or with a plurality of horseradish peroxidase molecules, and a bridging polynucleotide that contains a poly-dT sequence at the 5'-end that binds to the poly-dA tail of the captured nucleic acid, and a sequence at the 3'-end that is complementary to a region of the capture sequence. If the capture sequence is biotinylated, the microarray is then exposed to streptavidin-bound horseradish peroxidase. Hybridization of target RNAs is detected by the addition of a horseradish peroxidase substrate such as tetramethylbenzidine (TMB) and measurement of the absorbance of the solution at 450nm.

[00138] In still other embodiments, an addressable microarray is used to detect a target RNA using quantum dots. See Liang, R.Q. et al. (2005) Nucl. Acids Res. 33(2):e17, available at <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=548377>, which is incorporated herein by reference in its entirety. In some embodiments, total RNA is isolated from a sample. In some embodiments, small RNAs are isolated from the sample. The 3'-ends of the target RNAs are biotinylated using biotin-X-hydrazide. The biotinylated target RNAs are captured on a microarray comprising immobilized probes comprising sequences that are identically present in, or complementary to a region of, one or more of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and/or probes comprising sequences other than those that are complementary to one or more microRNAs of the human miRNome. The hybridized target RNAs are then labeled with quantum dots via a biotin-streptavidin binding. A confocal laser causes the quantum dots to fluoresce and the signal can be quantified. In alternative embodiments, small RNAs can be detected using a colorimetric assay. In these embodiments, small RNAs are labeled with streptavidin-

conjugated gold followed by silver enhancement. The gold nanoparticles bound to the hybridized target RNAs catalyze the reduction of silver ions to metallic silver, which can then be detected colorimetrically with a CCD camera.

[00139] In some embodiments, detection and quantification of one or more target RNAs is accomplished using microfluidic devices and single-molecule detection. In some embodiments, target RNAs in a sample of isolated total RNA are hybridized to two probes, one which is complementary to nucleic acids at the 5'-end of the target RNA and the second which is complementary to the 3'-end of the target RNA. Each probe comprises, in some embodiments, one or more affinity-enhancing nucleotide analogs, such as LNA nucleotide analogs and each is labeled with a different fluorescent dye having different fluorescence emission spectra. The sample is then flowed through a microfluidic capillary in which multiple lasers excite the fluorescent probes, such that a unique coincident burst of photons identifies a particular target RNA, and the number of particular unique coincident bursts of photons can be counted to quantify the amount of the target RNA in the sample. See U.S. Patent Publication No. 2006/0292616 to Neely et al., which is hereby incorporated by reference in its entirety. In some alternative embodiments, a target RNA-specific probe can be labeled with 3 or more distinct labels selected from, e.g., fluorophores, electron spin labels, etc., and then hybridized to an RNA sample, such as total RNA, or a sample that is enriched in small RNAs. The target RNA/probe duplex is then passed through channels in a microfluidic device and that comprise detectors that record the unique signal of the 3 labels. In this way, individual molecules are detected by their unique signal and counted. See U.S. Patent Nos. 7,402,422 and 7,351,538 to Fuchs *et al.*, U.S. Genomics, Inc., each of which is incorporated herein by reference in its entirety.

[00140] Nonlimiting exemplary target RNA-specific probes include probes comprising sequences selected from of SEQ ID NOs: 1 to 397. Nonlimiting exemplary target RNA-specific probes include probes comprising sequences that are complementary to sequences selected from of SEQ ID NOs: 1 to 397. Nonlimiting exemplary target RNA-specific probes also include probes comprising at least 15 contiguous nucleotides of, or the complement of at least 15 contiguous nucleotides of, a sequence selected from

SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00141] Optionally, the sample RNA is modified before hybridization. The target RNA/probe duplex is then passed through channels in a microfluidic device and that comprise detectors that record the unique signal of the 3 labels. In this way, individual molecules are detected by their unique signal and counted. See U.S. Patent Nos. 7,402,422 and 7,351,538 to Fuchs et al., U.S. Genomics, Inc., each of which is incorporated herein by reference in its entirety.

[00142] In some embodiments, the detection and quantification of one or more target RNAs is accomplished by a solution-based assay, such as a modified Invader assay. See Allawi H.T. et al. (2004) RNA 10:1153-1161, which is incorporated herein by reference in its entirety. In some embodiments, the modified invader assay can be performed on unfractionated detergent lysates of cells. In other embodiments, the modified invader assay can be performed on total RNA isolated from cells or on a sample enriched in small RNAs. The target RNAs in a sample are annealed to two probes which form hairpin structures. A first probe has a hairpin structure at the 5' end and a region at the 3'-end that has a sequence that is complementary to the sequence of a region at the 5'-end of a target RNA. The 3'-end of the first probe is the "invasive polynucleotide". A second probe has, from the 5' end to the 3'-end a first "flap" region that is not complementary to the target RNA, a second region that has a sequence that is complementary to the 3'-end of the target RNA, and a third region that forms a hairpin structure. When the two probes are bound to a target RNA target, they create an overlapping configuration of the probes on the target RNA template, which is recognized by the Cleavase enzyme, which releases the flap of the second probe into solution. The flap region then binds to a complementary region at the 3'-end of a secondary reaction template ("SRT"). A FRET polynucleotide (having a fluorescent dye bound to the 5'-end and a quencher that quenches the dye bound closer to the 3' end) binds to a complementary region at the 5'-end of the SRT, with the result that an overlapping configuration of the 3'-end of the flap and the 5'-end of the FRET polynucleotide is created. Cleavase recognizes the overlapping configuration and cleaves the 5'-end of the

FRET polynucleotide, generates a fluorescent signal when the dye is released into solution.

4.1.5. Exemplary polynucleotides

[00143] In some embodiments, polynucleotides are provided. In some embodiments, synthetic polynucleotides are provided. Synthetic polynucleotides, as used herein, refer to polynucleotides that have been synthesized in vitro either chemically or enzymatically. Chemical synthesis of polynucleotides includes, but is not limited to, synthesis using polynucleotide synthesizers, such as OligoPilot (GE Healthcare), ABI 3900 DNA Synthesizer (Applied Biosystems), and the like. Enzymatic synthesis includes, but is not limited, to producing polynucleotides by enzymatic amplification, e.g., PCR.

[00144] In some embodiments, a polynucleotide is provided that comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and sequences complementary to SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the polynucleotide further comprises a region having a sequence that is not found in, or complementary to, any of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a polynucleotide is provided that comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692, and sequences complementary to SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the polynucleotide further comprises a region having a sequence that is not found in, or complementary to, any of SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

[00145] A “region” can comprise the full-length sequence, or the complement of the full-length sequence, of a particular sequence, such as any of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 794 to 1043, and 2576 to 2672 or it can comprise a subsequence, or the complement of a subsequence, of a particular sequence, such as any of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707,

2064 to 2183, 2312 to 2452, 794 to 1043, and 2576 to 2672. Such subsequences may comprise, in some embodiments, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or more contiguous nucleotides from a particular SEQ ID NO or its complement.

[00146] In various embodiments, a polynucleotide comprises fewer than 500, fewer than 300, fewer than 200, fewer than 150, fewer than 100, fewer than 75, fewer than 50, fewer than 40, or fewer than 30 nucleotides. In various embodiments, a polynucleotide is between 8 and 200, between 8 and 150, between 8 and 100, between 8 and 75, between 8 and 50, between 8 and 40, or between 8 and 30 nucleotides long.

[00147] In some embodiments, the polynucleotide is a primer. In some embodiments, the primer is labeled with a detectable moiety. In some embodiments, a primer is not labeled. A primer, as used herein, is a polynucleotide that is capable of specifically hybridizing to a target RNA or to a cDNA reverse transcribed from the target RNA or to an amplicon that has been amplified from a target RNA or a cDNA (collectively referred to as “template”), and, in the presence of the template, a polymerase and suitable buffers and reagents, can be extended to form a primer extension product.

[00148] In some embodiments, the polynucleotide is a probe. In some embodiments, the probe is labeled with a detectable moiety. A detectable moiety, as used herein, includes both directly detectable moieties, such as fluorescent dyes, and indirectly detectable moieties, such as members of binding pairs. When the detectable moiety is a member of a binding pair, in some embodiments, the probe can be detectable by incubating the probe with a detectable label bound to the second member of the binding pair. In some embodiments, a probe is not labeled, such as when a probe is a capture probe, e.g., on a microarray or bead. In some embodiments, a probe is not extendable, e.g., by a polymerase. In other embodiments, a probe is extendable.

[00149] In some embodiments, the polynucleotide is a FRET probe that in some embodiments is labeled at the 5'-end with a fluorescent dye (donor) and at the 3'-end with a quencher (acceptor), a chemical group that absorbs (i.e., suppresses) fluorescence emission from the dye when the groups are in close proximity (i.e., attached to the same probe). In other embodiments, the donor and acceptor are not at the ends of

the FRET probe. Thus, in some embodiments, the emission spectrum of the donor moiety should overlap considerably with the absorption spectrum of the acceptor moiety.

4.1.5.1. Exemplary polynucleotide modifications

[00150] In some embodiments, the methods of detecting at least one target RNA described herein employ one or more polynucleotides that have been modified, such as polynucleotides comprising one or more affinity-enhancing nucleotide analogs. Modified polynucleotides useful in the methods described herein include primers for reverse transcription, PCR amplification primers, and probes. In some embodiments, the incorporation of affinity-enhancing nucleotides increases the binding affinity and specificity of a polynucleotide for its target nucleic acid as compared to polynucleotides that contain only deoxyribonucleotides, and allows for the use of shorter polynucleotides or for shorter regions of complementarity between the polynucleotide and the target nucleic acid.

[00151] In some embodiments, affinity-enhancing nucleotide analogs include nucleotides comprising one or more base modifications, sugar modifications and/or backbone modifications.

[00152] In some embodiments, modified bases for use in affinity-enhancing nucleotide analogs include 5-methylcytosine, isocytosine, pseudoisocytosine, 5-bromouracil, 5-propynyluracil, 6-aminopurine, 2-aminopurine, inosine, diaminopurine, 2-chloro-6-aminopurine, xanthine and hypoxanthine.

[00153] In some embodiments, affinity-enhancing nucleotide analogs include nucleotides having modified sugars such as 2'-substituted sugars, such as 2'-O-alkyl-ribose sugars, 2'-amino-deoxyribose sugars, 2'-fluoro-deoxyribose sugars, 2'-fluoro-arabinose sugars, and 2'-O-methoxyethyl-ribose (2'MOE) sugars. In some embodiments, modified sugars are arabinose sugars, or d-arabino-hexitol sugars.

[00154] In some embodiments, affinity-enhancing nucleotide analogs include backbone modifications such as the use of peptide nucleic acids (PNA; *e.g.*, an oligomer including nucleobases linked together by an amino acid backbone). Other backbone modifications include phosphorothioate linkages, phosphodiester modified nucleic acids, combinations of phosphodiester and phosphorothioate nucleic acid,

methylphosphonate, alkylphosphonates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters, methylphosphorothioate, phosphorodithioate, p-ethoxy, and combinations thereof.

[00155] In some embodiments, a polynucleotide includes at least one affinity-enhancing nucleotide analog that has a modified base, at least nucleotide (which may be the same nucleotide) that has a modified sugar, and/or at least one internucleotide linkage that is non-naturally occurring.

[00156] In some embodiments, an affinity-enhancing nucleotide analog contains a locked nucleic acid (“LNA”) sugar, which is a bicyclic sugar. In some embodiments, a polynucleotide for use in the methods described herein comprises one or more nucleotides having an LNA sugar. In some embodiments, a polynucleotide contains one or more regions consisting of nucleotides with LNA sugars. In other embodiments, a polynucleotide contains nucleotides with LNA sugars interspersed with deoxyribonucleotides. *See, e.g.,* Frieden, M. *et al.* (2008) *Curr. Pharm. Des.* 14(11):1138-1142.

4.1.5.2. Exemplary primers

[00157] In some embodiments, a primer is provided. In some embodiments, a primer is identical or complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of a target RNA. In some embodiments, a primer may also comprise portions or regions that are not identical or complementary to the target RNA. In some embodiments, a region of a primer that is identical or complementary to a target RNA is contiguous, such that any region of a primer that is not identical or complementary to the target RNA does not disrupt the identical or complementary region.

[00158] In some embodiments, a primer comprises a portion that is identically present in a target RNA. In some such embodiments, a primer that comprises a region that is identically present in the target RNA is capable of selectively hybridizing to a cDNA that has been reverse transcribed from the RNA, or to an amplicon that has been produced by amplification of the target RNA or cDNA. In some embodiments, the

primer is complementary to a sufficient portion of the cDNA or amplicon such that it selectively hybridizes to the cDNA or amplicon under the conditions of the particular assay being used.

[00159] As used herein, “selectively hybridize” means that a polynucleotide, such as a primer or probe, will hybridize to a particular nucleic acid in a sample with at least 5-fold greater affinity than it will hybridize to another nucleic acid present in the same sample that has a different nucleotide sequence in the hybridizing region. Exemplary hybridization conditions are discussed in Example 1. In some embodiments, a polynucleotide will hybridize to a particular nucleic acid in a sample with at least 10-fold greater affinity than it will hybridize to another nucleic acid present in the same sample that has a different nucleotide sequence in the hybridizing region.

[00160] Nonlimiting exemplary primers include primers comprising sequences that are identically present in, or complementary to a region of, sequences selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Exemplary primers also include, but are not limited to, primers comprising regions that are identical or complementary to at least 15 contiguous nucleotides of sequences selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Exemplary primers also include, but are not limited to, primers comprising regions that are identical or complementary to at least 15 contiguous nucleotides of sequences selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

[00161] In some embodiments, a primer is used to reverse transcribe a target RNA, for example, as discussed herein. In some embodiments, a primer is used to amplify a target RNA or a cDNA reverse transcribed therefrom. Such amplification, in some embodiments, is quantitative PCR, for example, as discussed herein. In some embodiments, a primer comprises a detectable moiety.

4.1.5.3. Exemplary probes

[00162] In various embodiments, methods of detecting the presence of a lung cancer comprise hybridizing nucleic acids of a human sample with a probe. In some embodiments, the probe comprises a portion that is complementary to a target RNA. In some embodiments, the probe comprises a portion that is identically present in the target

RNA. In some such embodiments, a probe that is complementary to a target RNA is complementary to a sufficient portion of the target RNA such that it selectively hybridizes to the target RNA under the conditions of the particular assay being used. In some embodiments, a probe that is complementary to a target RNA is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of the target RNA. In some embodiments, a probe that is complementary to a target RNA comprises a region that is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of the target RNA. That is, a probe that is complementary to a target RNA may also comprise portions or regions that are not complementary to the target RNA. In some embodiments, a region of a probe that is complementary to a target RNA is contiguous, such that any region of a probe that is not complementary to the target RNA does not disrupt the complementary region.

[00163] In some embodiments, the probe comprises a portion that is identically present in the target RNA. In some such embodiments, a probe that comprises a region that is identically present in the target RNA is capable of selectively hybridizing to a cDNA that has been reverse transcribed from the RNA, or to an amplicon that has been produced by amplification of the target RNA or cDNA. In some embodiments, the probe is complementary to a sufficient portion of the cDNA or amplicon such that it selectively hybridizes to the cDNA or amplicon under the conditions of the particular assay being used. In some embodiments, a probe that is complementary to a cDNA or amplicon is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of the cDNA or amplicon. In some embodiments, a probe that is complementary to a target RNA comprises a region that is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides of the cDNA or amplicon. That is, a probe that is complementary to a cDNA or amplicon may also comprise portions or regions that are not complementary to the cDNA or amplicon.

In some embodiments, a region of a probe that is complementary to a cDNA or amplicon is contiguous, such that any region of a probe that is not complementary to the cDNA or amplicon does not disrupt the complementary region.

[00164] Nonlimiting exemplary probes include probes comprising sequences set forth in SEQ ID NOs: 1 to 397. Nonlimiting exemplary probes include probes comprising sequences that are identically present in, or complementary to a region of, sequences selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Exemplary probes also include, but are not limited to, probes comprising regions that are identical or complementary to at least 15 contiguous nucleotides of sequences selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00165] In some embodiments, the method of detectably quantifying one or more target RNAs comprises: (a) isolating total RNA; (b) reverse transcribing a target RNA to produce a cDNA that is complementary to the target RNA; (c) amplifying the cDNA from (b); and (d) detecting the amount of a target RNA using real time RT-PCR.

[00166] As described above, in some embodiments, the real time RT-PCR detection is performed using a FRET probe, which includes, but is not limited to, a TaqMan® probe, a Molecular beacon probe and a Scorpion probe. In some embodiments, the real time RT-PCR detection and quantification is performed with a TaqMan® probe, *i.e.*, a linear probe that typically has a fluorescent dye covalently bound at one end of the DNA and a quencher molecule covalently bound at the other end of the DNA. The FRET probe comprises a sequence that is complementary to a region of the cDNA such that, when the FRET probe is hybridized to the cDNA, the dye fluorescence is quenched, and when the probe is digested during amplification of the cDNA, the dye is released from the probe and produces a fluorescence signal. In such embodiments, the amount of target RNA in the sample is proportional to the amount of fluorescence measured during cDNA amplification.

[00167] The TaqMan® probe typically comprises a region of contiguous nucleotides having a sequence that is complementary to a region of a target RNA or its complementary cDNA that is reverse transcribed from the target RNA template (*i.e.*, the sequence of the probe region is complementary to or identically present in the target RNA

to be detected) such that the probe is specifically hybridizable to the resulting PCR amplicon. In some embodiments, the probe comprises a region of at least 6 contiguous nucleotides having a sequence that is fully complementary to or identically present in a region of a cDNA that has been reverse transcribed from a target RNA template, such as comprising a region of at least 8 contiguous nucleotides, at least 10 contiguous nucleotides, at least 12 contiguous nucleotides, at least 14 contiguous nucleotides, or at least 16 contiguous nucleotides having a sequence that is complementary to or identically present in a region of a cDNA reverse transcribed from a target RNA to be detected.

[00168] In some embodiments, the region of the cDNA that has a sequence that is complementary to the TaqMan® probe sequence is at or near the center of the cDNA molecule. In some embodiments, there are independently at least 2 nucleotides, at least 3 nucleotides, at least 4 nucleotides, at least 5 nucleotides of the cDNA at the 5'-end and at the 3'-end of the region of complementarity.

[00169] In some embodiments, Molecular Beacons can be used to detect and quantitate PCR products. Like TaqMan® probes, Molecular Beacons use FRET to detect and quantitate a PCR product via a probe having a fluorescent dye and a quencher attached at the ends of the probe. Unlike TaqMan® probes, Molecular Beacons remain intact during the PCR cycles. Molecular Beacon probes form a stem-loop structure when free in solution, thereby allowing the dye and quencher to be in close enough proximity to cause fluorescence quenching. When the Molecular Beacon hybridizes to a target, the stem-loop structure is abolished so that the dye and the quencher become separated in space and the dye fluoresces. Molecular Beacons are available, e.g., from Gene Link™ (see <http://www.genelink.com/newsite/products/mbintro.asp>).

[00170] In some embodiments, Scorpion probes can be used as both sequence-specific primers and for PCR product detection and quantitation. Like Molecular Beacons, Scorpion probes form a stem-loop structure when not hybridized to a target nucleic acid. However, unlike Molecular Beacons, a Scorpion probe achieves both sequence-specific priming and PCR product detection. A fluorescent dye molecule is attached to the 5'-end of the Scorpion probe, and a quencher is attached to the 3'-end. The 3' portion of the probe is complementary to the extension product of the PCR primer, and this complementary portion is linked to the 5'-end of the probe by a non-amplifiable

moiety. After the Scorpion primer is extended, the target-specific sequence of the probe binds to its complement within the extended amplicon, thus opening up the stem-loop structure and allowing the dye on the 5'-end to fluoresce and generate a signal. Scorpion probes are available from, e.g, Premier Biosoft International (see http://www.premierbiosoft.com/tech_notes/Scorpion.html).

[00171] In some embodiments, labels that can be used on the FRET probes include colorimetric and fluorescent labels such as Alexa Fluor dyes, BODIPY dyes, such as BODIPY FL; Cascade Blue; Cascade Yellow; coumarin and its derivatives, such as 7-amino-4-methylcoumarin, aminocoumarin and hydroxycoumarin; cyanine dyes, such as Cy3 and Cy5; eosins and erythrosins; fluorescein and its derivatives, such as fluorescein isothiocyanate; macrocyclic chelates of lanthanide ions, such as Quantum DyeTM; Marina Blue; Oregon Green; rhodamine dyes, such as rhodamine red, tetramethylrhodamine and rhodamine 6G; Texas Red; fluorescent energy transfer dyes, such as thiazole orange-ethidium heterodimer; and, TOTAB.

[00172] Specific examples of dyes include, but are not limited to, those identified above and the following: Alexa Fluor 350, Alexa Fluor 405, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 500, Alexa Fluor 514, Alexa Fluor 532, Alexa Fluor 546, Alexa Fluor 555, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 610, Alexa Fluor 633, Alexa Fluor 647, Alexa Fluor 660, Alexa Fluor 680, Alexa Fluor 700, and, Alexa Fluor 750; amine-reactive BODIPY dyes, such as BODIPY 493/503, BODIPY 530/550, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY 630/650, BODIPY 650/655, BODIPY FL, BODIPY R6G, BODIPY TMR, and, BODIPY-TR; Cy3, Cy5, 6-FAM, Fluorescein Isothiocyanate, HEX, 6-JOE, Oregon Green 488, Oregon Green 500, Oregon Green 514, Pacific Blue, REG, Rhodamine Green, Rhodamine Red, Renographin, ROX, SYPRO, TAMRA, 2', 4', 5', 7'-Tetrabromosulfonefluorescein, and TET.

[00173] Specific examples of fluorescently labeled ribonucleotides useful in the preparation of RT-PCR probes for use in some embodiments of the methods described herein are available from Molecular Probes (Invitrogen), and these include, Alexa Fluor 488-5-UTP, Fluorescein-12-UTP, BODIPY FL-14-UTP, BODIPY TMR-14-UTP, Tetramethylrhodamine-6-UTP, Alexa Fluor 546-14-UTP, Texas Red-5-UTP, and

BODIPY TR-14-UTP. Other fluorescent ribonucleotides are available from Amersham Biosciences (GE Healthcare), such as Cy3-UTP and Cy5-UTP.

[00174] Examples of fluorescently labeled deoxyribonucleotides useful in the preparation of RT-PCR probes for use in the methods described herein include Dinitrophenyl (DNP)-1'-dUTP, Cascade Blue-7-dUTP, Alexa Fluor 488-5-dUTP, Fluorescein-12-dUTP, Oregon Green 488-5-dUTP, BODIPY FL-14-dUTP, Rhodamine Green-5-dUTP, Alexa Fluor 532-5-dUTP, BODIPY TMR-14-dUTP, Tetramethylrhodamine-6-dUTP, Alexa Fluor 546-14-dUTP, Alexa Fluor 568-5-dUTP, Texas Red-12-dUTP, Texas Red-5-dUTP, BODIPY TR-14-dUTP, Alexa Fluor 594-5-dUTP, BODIPY 630/650-14-dUTP, BODIPY 650/665-14-dUTP; Alexa Fluor 488-7-OBEA-dCTP, Alexa Fluor 546-16-OBEA-dCTP, Alexa Fluor 594-7-OBEA-dCTP, Alexa Fluor 647-12-OBEA-dCTP. Fluorescently labeled nucleotides are commercially available and can be purchased from, e.g., Invitrogen.

[00175] In some embodiments, dyes and other moieties, such as quenchers, are introduced into polynucleotide used in the methods described herein, such as FRET probes, via modified nucleotides. A “modified nucleotide” refers to a nucleotide that has been chemically modified, but still functions as a nucleotide. In some embodiments, the modified nucleotide has a chemical moiety, such as a dye or quencher, covalently attached, and can be introduced into a polynucleotide, for example, by way of solid phase synthesis of the polynucleotide. In other embodiments, the modified nucleotide includes one or more reactive groups that can react with a dye or quencher before, during, or after incorporation of the modified nucleotide into the nucleic acid. In specific embodiments, the modified nucleotide is an amine-modified nucleotide, *i.e.*, a nucleotide that has been modified to have a reactive amine group. In some embodiments, the modified nucleotide comprises a modified base moiety, such as uridine, adenosine, guanosine, and/or cytosine. In specific embodiments, the amine-modified nucleotide is selected from 5-(3-aminoallyl)-UTP; 8-[(4-amino)butyl]-amino-ATP and 8-[(6-amino)butyl]-amino-ATP; N6-(4-amino)butyl-ATP, N6-(6-amino)butyl-ATP, N4-[2,2-oxy-bis-(ethylamine)]-CTP; N6-(6-Amino)hexyl-ATP; 8-[(6-Amino)hexyl]-amino-ATP; 5-propargylamino-CTP, 5-propargylamino-UTP. In some embodiments, nucleotides with different nucleobase moieties are similarly modified, for example, 5-(3-aminoallyl)-GTP instead of 5-(3-

aminoallyl)-UTP. Many amine modified nucleotides are commercially available from, e.g., Applied Biosystems, Sigma, Jena Bioscience and TriLink.

[00176] Exemplary detectable moieties also include, but are not limited to, members of binding pairs. In some such embodiments, a first member of a binding pair is linked to a polynucleotide. The second member of the binding pair is linked to a detectable label, such as a fluorescent label. When the polynucleotide linked to the first member of the binding pair is incubated with the second member of the binding pair linked to the detectable label, the first and second members of the binding pair associate and the polynucleotide can be detected. Exemplary binding pairs include, but are not limited to, biotin and streptavidin, antibodies and antigens, etc.

[00177] In some embodiments, multiple target RNAs are detected in a single multiplex reaction. In some such embodiments, each probe that is targeted to a unique cDNA is spectrally distinguishable when released from the probe. Thus, each target RNA is detected by a unique fluorescence signal.

[00178] One skilled in the art can select a suitable detection method for a selected assay, e.g., a real-time RT-PCR assay. The selected detection method need not be a method described above, and may be any method.

4.2. Exemplary compositions and kits

[00179] In another aspect, compositions are provided. In some embodiments, compositions are provided for use in the methods described herein.

[00180] In some embodiments, a composition comprises at least one polynucleotide. In some embodiments, a composition comprises at least one primer. In some embodiments, a composition comprises at least one probe. In some embodiments, a composition comprises at least one primer and at least one probe.

[00181] In some embodiments, compositions are provided that comprise at least one target RNA-specific primer. The term “target RNA-specific primer” encompasses primers that have a region of contiguous nucleotides having a sequence that is (i) identically present in one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, or (ii) complementary to the

sequence of a region of contiguous nucleotides found in one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00182] In some embodiments, compositions are provided that comprise at least one target RNA-specific probe. The term “target RNA-specific probe” encompasses probes that have a region of contiguous nucleotides having a sequence that is (i) identically present in one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, or (ii) complementary to the sequence of a region of contiguous nucleotides found in one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00183] In some embodiments, target RNA-specific primers and probes comprise deoxyribonucleotides. In other embodiments, target RNA-specific primers and probes comprise at least one nucleotide analog. Nonlimiting exemplary nucleotide analogs include, but are not limited to, analogs described herein, including LNA analogs and peptide nucleic acid (PNA) analogs. In some embodiments, target RNA-specific primers and probes comprise at least one nucleotide analog which increases the hybridization binding energy (e.g., an affinity-enhancing nucleotide analog, discussed above). In some embodiments, a target RNA-specific primer or probe in the compositions described herein binds to one target RNA in the sample. In some embodiments, a single primer or probe binds to multiple target RNAs, such as multiple isomirs.

[00184] In some embodiments, more than one primer or probe specific for a single target RNA is present in the compositions, the primers or probes capable of binding to overlapping or spatially separated regions of the target RNA.

[00185] It will be understood, even if not explicitly stated hereinafter, that in some embodiments in which the compositions described herein are designed to hybridize to cDNAs reverse transcribed from target RNAs, the composition comprises at least one target RNA-specific primer or probe (or region thereof) having a sequence that is identically present in a target RNA (or region thereof).

[00186] In some embodiments, a target RNA is capable of specifically hybridizing to at least one probe sequence in one of Tables 6, 7, 8, 9, 32, 33, or 34. In

some embodiments, a target RNA comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a target RNA comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00187] In some embodiments, the composition comprises a plurality of target RNA-specific primers and/or probes for each of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, or at least 8 target RNAs, the target RNAs comprising a region of contiguous nucleotides having a sequence that is identically present in one of SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the plurality includes a target RNA-specific primer and/or probe specific for each of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, or at least 12 target RNAs, the target RNAs comprising a region of contiguous nucleotides having a sequence that is identically present in one of SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the plurality includes a target RNA-specific primer and/or probe specific for each of at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 75, or at least 100 target RNAs comprising a region of contiguous nucleotides having a sequence that is identically present in one of SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. It will be understood that, in some embodiments, target RNAs described herein comprise a sequence identically present in a sequence set forth in at least one of Tables 4, 5, and 38. It is understood that where a sequence includes thymine (T) bases, a target RNA may contain uracil (U) bases instead.

[00188] In some embodiments, a composition is an aqueous composition. In some embodiments, the aqueous composition comprises a buffering component, such as phosphate, tris, HEPES, etc., and/or additional components, as discussed below. In some embodiments, a composition is dry, for example, lyophilized, and suitable for reconstitution by addition of fluid. A dry composition may include a buffering component and/or additional components.

[00189] In some embodiments, a composition comprises one or more additional components. Additional components include, but are not limited to, salts, such as NaCl, KCl, and MgCl₂; polymerases, including thermostable polymerases; dNTPs; RNase inhibitors; bovine serum albumin (BSA) and the like; reducing agents, such as β -mercaptoethanol; EDTA and the like; etc. One skilled in the art can select suitable composition components depending on the intended use of the composition.

[00190] In some embodiments, an addressable microarray component is provided that comprises target RNA-specific probes attached to a substrate.

[00191] Microarrays for use in the methods described herein comprise a solid substrate onto which the probes are covalently or non-covalently attached. In some embodiments, probes capable of hybridizing to one or more target RNAs or cDNAs are attached to the substrate at a defined location (“addressable array”). Probes can be attached to the substrate in a wide variety of ways, as will be appreciated by those in the art. In some embodiments, the probes are synthesized first and subsequently attached to the substrate. In other embodiments, the probes are synthesized on the substrate. In some embodiments, probes are synthesized on the substrate surface using techniques such as photopolymerization and photolithography.

[00192] In some embodiments, the solid substrate is a material that is modified to contain discrete individual sites appropriate for the attachment or association of the probes and is amenable to at least one detection method. Representative examples of substrates include glass and modified or functionalized glass, plastics (including acrylics, polystyrene and copolymers of styrene and other materials, polypropylene, polyethylene, polybutylene, polyurethanes, Teflon, etc.), polysaccharides, nylon or nitrocellulose, resins, silica or silica-based materials including silicon and modified silicon, carbon, metals, inorganic glasses and plastics. In some embodiments, the substrates allow optical detection without appreciably fluorescing.

[00193] In some embodiments, the substrate is planar. In other embodiments, probes are placed on the inside surface of a tube, such as for flow-through sample analysis to minimize sample volume. In other embodiments, probes can be in the wells of multi-well plates. In still other embodiments, probes can be attached to an addressable microbead array. In yet other embodiments, the probes can be attached to a

flexible substrate, such as a flexible foam, including closed cell foams made of particular plastics.

[00194] The substrate and the probe can each be derivatized with functional groups for subsequent attachment of the two. For example, in some embodiments, the substrate is derivatized with one or more chemical functional groups including, but not limited to, amino groups, carboxyl groups, oxo groups and thiol groups. In some embodiments, probes are attached directly to the substrate through one or more functional groups. In some embodiments, probes are attached to the substrate indirectly through a linker (*i.e.*, a region of contiguous nucleotides that space the probe regions involved in hybridization and detection away from the substrate surface). In some embodiments, probes are attached to the solid support through the 5' terminus. In other embodiments, probes are attached through the 3' terminus. In still other embodiments, probes are attached to the substrate through an internal nucleotide. In some embodiments the probe is attached to the solid support non-covalently, e.g., via a biotin-streptavidin interaction, wherein the probe biotinylated and the substrate surface is covalently coated with streptavidin.

[00195] In some embodiments, the compositions comprise a microarray having probes attached to a substrate, wherein at least one of the probes (or a region thereof) comprises a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least 2, at least 5, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, or at least 100 of the probes comprise a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the microarray comprises at least one target RNA-specific probe comprising a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and at least one target RNA-specific probe comprising a sequence that is identically present in, or complementary to a region of, a target RNA of the human miRNome. In some embodiments, the microarray comprises each target RNA-specific probe at only one location on the microarray. In

some embodiments, the microarray comprises at least one target RNA-specific probe at multiple locations on the microarray.

[00196] As used herein, the terms “complementary” or “partially complementary” to a target RNA (or target region thereof), and the percentage of “complementarity” of the probe sequence to that of the target RNA sequence is the percentage “identity” to the reverse complement of the sequence of the target RNA. In determining the degree of “complementarity” between probes used in the compositions described herein (or regions thereof) and a target RNA, such as those disclosed herein, the degree of “complementarity” is expressed as the percentage identity between the sequence of the probe (or region thereof) and the reverse complement of the sequence of the target RNA that best aligns therewith. The percentage is calculated by counting the number of aligned bases that are identical as between the 2 sequences, dividing by the total number of contiguous nucleotides in the probe, and multiplying by 100.

[00197] In some embodiments, the microarray comprises at least one probe having a region with a sequence that is fully complementary to a target region of a target RNA. In other embodiments, the microarray comprises at least one probe having a region with a sequence that comprises one or more base mismatches when compared to the sequence of the best-aligned target region of a target RNA.

[00198] As noted above, a “region” of a probe or target RNA, as used herein, may comprise or consist of 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or more contiguous nucleotides from a particular SEQ ID NO or the complement thereof. In some embodiments, the region is of the same length as the probe or the target RNA. In other embodiments, the region is shorter than the length of the probe or the target RNA.

[00199] In some embodiments, the microarray comprises at least one probe having a region of at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00200] In some embodiments, the microarray comprises at least one probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34. In some embodiments, a microarray further comprises at least one probe that does not have a region that is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34.

[00201] In some embodiments, the microarray comprises at least one, at least two, at least three, at least five, at least 10, or at least 15 probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 6. In some embodiments, the microarray comprises at least one, at least two, at least three, at least five, at least eight, at least 10, at least 12, at least 15, at least 20, or at least 25 probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 7. In some embodiments, the microarray comprises at least one, at least two, at least three, at least five, at least six, or at least seven probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 8. In some embodiments, the microarray comprises at least one, at least two, at least three, at least four, at least five, at least ten, at least 15, at least 20, at least 25, at least 30, at least 40, or at least 50 probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 9. In some embodiments, the microarray comprises at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, or at least 40 probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in one of Tables 32 or 33. In some embodiments, the microarray comprises at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, or at least 70 probes that each comprise a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 34.

[00202] In some embodiments, the microarrays comprise probes having a region with a sequence that is complementary to target RNAs that comprise a substantial portion of the human miRNome (*i.e.*, the publicly known microRNAs that have been

accessioned by others into miRBase (<http://microrna.sanger.ac.uk/> at the time the microarray is fabricated), such as at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% of the human miRNome. In some embodiments, the microarrays comprise probes that have a region with a sequence that is identically present in target RNAs that comprise a substantial portion of the human miRNome, such as at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% of the human miRNome.

[00203] In some embodiments, components are provided that comprise probes attached to microbeads, such as those sold by Luminex, each of which is internally dyed with red and infrared fluorophores at different intensities to create a unique signal for each bead. In some embodiments, the compositions useful for carrying out the methods described herein include a plurality of microbeads, each with a unique spectral signature. Each uniquely labeled microbead is attached to a unique target RNA-specific probe such that the unique spectral signature from the dyes in the bead is associated with a particular probe sequence. Nonlimiting exemplary probe sequences include SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. Nonlimiting exemplary probe sequences also include probes comprising a region that is identically present in, or complementary to, a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a probe sequence comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, or at least 24 contiguous nucleotides that are identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00204] In some embodiments, a uniquely labeled microbead has attached thereto a probe having a region with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In other embodiments, the uniquely labeled microbead has attached thereto a probe having a region with a sequence that comprises one or more base mismatches when compared to the most similar sequence

selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, and sequences complementary to SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00205] In some embodiments, a composition is provided that comprises a plurality of uniquely labeled microbeads, wherein at least one microbead has attached thereto a probe having a region of at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00206] In some embodiments, a composition is provided that comprises a plurality of uniquely labeled microbeads, wherein at least one microbead has attached thereto a probe having a region of at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides with a sequence that is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34. In some embodiments, the composition further comprises at least one uniquely labeled microbead having attached thereto a probe that does not have a region that is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34.

[00207] In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least 10, or at least 15 uniquely labeled microbeads that each have attached thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 6. In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least eight, at least 10, at least 12, at least 15, at least 20, or at least 25 uniquely labeled microbeads that each have attached thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 7. In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least six, or at least seven uniquely labeled microbeads that each have attached

thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 8. In some embodiments, the compositions comprise at least one, at least two, at least three, at least four, at least five, at least ten, at least 15, at least 20, at least 25, at least 30, at least 40, or at least 50 uniquely labeled microbeads that each have attached thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 9. In some embodiments, the compositions comprise at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, or at least 40 uniquely labeled microbeads that each have attached thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in one of Tables 32 or 33. In some embodiments, the compositions comprise at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, or at least 70 uniquely labeled microbeads that each have attached thereto a unique target RNA-specific probe having a region with a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 34.

[00208] In some embodiments, the compositions comprise a plurality of uniquely labeled microbeads, wherein the plurality comprises at least one microbead having attached thereto a probe having a region with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the plurality comprises at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, at least 75, or at least 100 microbeads each of which having attached thereto a probe having a region with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a composition comprises at least one uniquely labeled microbead having attached thereto a target RNA-specific probe having a region with a sequence that is not present in, or complementary to a region of, any of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00209] In some embodiments, the compositions comprise a plurality of uniquely labeled microbeads, at least one of which has attached thereto a probe having a region with a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and at least a second bead that has attached thereto a probe having a region with a sequence that is identically present in, or complementary to a region of, a target RNA from the human miRNome.

[00210] In some embodiments, the compositions comprise a plurality of uniquely labeled microbeads, each of which has attached thereto a unique probe having a region that is complementary to target RNAs that comprise a substantial portion of the human miRNome, such as at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% of the human miRNome. In some embodiments, the compositions comprise a plurality of uniquely labeled microbeads having attached thereto a unique probe having a region with a sequence that is identically present in target RNAs that comprise a substantial portion of the human miRNome, such as at least about 60%, at least about 70%, at least about 80%, at least about 90%, or at least about 95% of the human miRNome.

[00211] In some embodiments, compositions are provided that comprise at least one polynucleotide for detecting at least one target RNA. In some embodiments, the polynucleotide is used as a primer for a reverse transcriptase reaction. In some embodiments, the polynucleotide is used as a primer for amplification. In some embodiments, the polynucleotide is used as a primer for RT-PCR. In some embodiments, the polynucleotide is used as a probe for detecting at least one target RNA. In some embodiments, the polynucleotide is detectably labeled. In some embodiments, the polynucleotide is a FRET probe. In some embodiments, the polynucleotide is a TaqMan® probe, a Molecular Beacon, or a Scorpion probe.

[00212] In some embodiments, a composition comprises at least one FRET probe having a sequence that is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, a composition comprises at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50,

at least 60, at least 75, or at least 100 FRET probes, each of which has a sequence that is identically present in, or complementary to a region of, a different one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00213] In some embodiments, a FRET probe is labeled with a donor/acceptor pair such that when the probe is digested during the PCR reaction, it produces a unique fluorescence emission that is associated with a specific target RNA. In some embodiments, when a composition comprises multiple FRET probes, each probe is labeled with a different donor/acceptor pair such that when the probe is digested during the PCR reaction, each one produces a unique fluorescence emission that is associated with a specific probe sequence and/or target RNA. In some embodiments, the sequence of the FRET probe is complementary to a target region of a target RNA. In other embodiments, the FRET probe has a sequence that comprises one or more base mismatches when compared to the sequence of the best-aligned target region of a target RNA.

[00214] In some embodiments, a composition comprises a FRET probe consisting of at least 8, at least 9, at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 nucleotides, wherein at least a portion of the sequence is identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least 8, at least 9, at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 nucleotides of the FRET probe are identically present in, or complementary to a region of, one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, the FRET probe has a sequence with one, two or three base mismatches when compared to the sequence or complement of one of SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689.

[00215] In some embodiments, a composition comprises a FRET probe consisting of at least 8, at least 9, at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least

23, at least 24, or at least 25 nucleotides, wherein at least a portion of the sequence is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34. In some embodiments, a composition comprises at least one FRET probe that does not comprise a portion that is identically present in, or complementary to a region of, a probe sequence in one of Tables 6, 7, 8, 9, 32, 33, and 34.

[00216] In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least 10, or at least 15 uniquely labeled target RNA-specific FRET probes, each comprising a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 6. In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least eight, at least 10, at least 12, at least 15, at least 20, or at least 25 uniquely labeled target RNA-specific FRET probes, each of which comprises a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 7. In some embodiments, the compositions comprise at least one, at least two, at least three, at least five, at least six, or at least seven uniquely labeled target RNA-specific FRET probes, each of which comprises a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 8. In some embodiments, the compositions comprise at least one, at least two, at least three, at least four, at least five, at least ten, at least 15, at least 20, at least 25, at least 30, at least 40, or at least 50 uniquely labeled target RNA-specific FRET probes, each of which comprises a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 9. In some embodiments, the compositions comprise at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, or at least 40 uniquely labeled target RNA-specific FRET probes, each of which comprises a sequence that is identically present in, or complementary to a region of, a probe sequence in one of Tables 32 or 33. In some embodiments, the compositions comprise at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, or at least 70 uniquely labeled target RNA-specific FRET probes, each of which comprises a sequence that is identically present in, or complementary to a region of, a probe sequence in Table 34.

[00217] In some embodiments, a kit comprises a polynucleotide discussed above. In some embodiments, a kit comprises at least one primer and/or probe discussed above. In some embodiments, a kit comprises at least one polymerase, such as a thermostable polymerase. In some embodiments, a kit comprises dNTPs. In some embodiments, kits for use in the real time RT-PCR methods described herein comprise one or more target RNA-specific FRET probes and/or one or more primers for reverse transcription of target RNAs and/or one or more primers for amplification of target RNAs or cDNAs reverse transcribed therefrom.

[00218] In some embodiments, one or more of the primers and/or probes is “linear”. A “linear” primer refers to a polynucleotide that is a single stranded molecule, and typically does not comprise a short region of, for example, at least 3, 4 or 5 contiguous nucleotides, which are complementary to another region within the same polynucleotide such that the primer forms an internal duplex. In some embodiments, the primers for use in reverse transcription comprise a region of at least 4, at least 5, at least 6, at least 7 or more contiguous nucleotides at the 3'-end that has a sequence that is complementary to region of at least 4, at least 5, at least 6, at least 7 or more contiguous nucleotides at the 5'-end of a target RNA.

[00219] In some embodiments, a kit comprises one or more pairs of linear primers (a “forward primer” and a “reverse primer”) for amplification of a cDNA reverse transcribed from a target RNA. Accordingly, in some embodiments, a first primer comprises a region of at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 contiguous nucleotides having a sequence that is identical to the sequence of a region of at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 contiguous nucleotides at the 5'-end of a target RNA. Furthermore, in some embodiments, a second primer comprises a region of at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 contiguous nucleotides having a sequence that is complementary to the sequence of a region of at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 contiguous nucleotides at the 3'-end of a target RNA. In some embodiments, the kit comprises at least a first set of primers for amplification of a cDNA that is reverse transcribed from a target RNA capable of specifically hybridizing to a nucleic acid comprising a sequence identically present in one of SEQ ID NOs: 1 to 397,

1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and/or a cDNA that is reverse transcribed from a target RNA that comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

[00220] In some embodiments, the kit comprises at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, at least 75, or at least 100 sets of primers, each of which is for amplification of a cDNA that is reverse transcribed from a different target RNA capable of specifically hybridizing to a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and/or a cDNA that is reverse transcribed from a target RNA that comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, the kit comprises at least one set of primers that is capable of amplifying more than one cDNA reverse transcribed from a target RNA in a sample.

[00221] In some embodiments, probes and/or primers for use in the compositions described herein comprise deoxyribonucleotides. In some embodiments, probes and/or primers for use in the compositions described herein comprise deoxyribonucleotides and one or more nucleotide analogs, such as LNA analogs or other duplex-stabilizing nucleotide analogs described above. In some embodiments, probes and/or primers for use in the compositions described herein comprise all nucleotide analogs. In some embodiments, the probes and/or primers comprise one or more duplex-stabilizing nucleotide analogs, such as LNA analogs, in the region of complementarity.

[00222] In some embodiments, the compositions described herein also comprise probes, and in the case of RT-PCR, primers, that are specific to one or more housekeeping genes for use in normalizing the quantities of target RNAs. Such probes (and primers) include those that are specific for one or more products of housekeeping genes selected from U6 snRNA, RNU44, RNU48, U47, 7SL scRNA, U1 snRNA, 5.8S rRNA, and U87 scaRNA.

[00223] In some embodiments, the kits for use in real time RT-PCR methods described herein further comprise reagents for use in the reverse transcription and amplification reactions. In some embodiments, the kits comprise enzymes such as

reverse transcriptase, and a heat stable DNA polymerase, such as Taq polymerase. In some embodiments, the kits further comprise deoxyribonucleotide triphosphates (dNTPs) for use in reverse transcription and amplification. In further embodiments, the kits comprise buffers optimized for specific hybridization of the probes and primers.

4.2.1. Exemplary normalization of RNA levels

[00224] In some embodiments, quantitation of target RNA expression levels requires assumptions to be made about the total RNA per cell and the extent of sample loss during sample preparation. In order to correct for differences between different samples or between samples that are prepared under different conditions, the quantities of target RNAs in some embodiments are normalized to the expression of at least one endogenous housekeeping gene.

[00225] Appropriate genes for use as reference genes in the methods described herein include those as to which the quantity of the product does not vary between normal samples and samples from lung cancer patients, or between different cell lines or under different growth and sample preparation conditions. In some embodiments, endogenous housekeeping genes useful as normalization controls in the methods described herein include, but are not limited to, U6 snRNA, RNU44, RNU48, U47, 7SL scRNA, U1 snRNA, 5.8S rRNA, and U87 scaRNA. In typical embodiments, the at least one endogenous housekeeping gene for use in normalizing the measured quantity of microRNAs is selected from U6 snRNA, RNU44, RNU48, U47, 7SL scRNA, U1 snRNA, 5.8S rRNA, and U87 scaRNA. In some embodiments, one housekeeping gene is used for normalization. In some embodiments, more than one housekeeping gene is used for normalization.

4.2.2. Exemplary qualitative methods

[00226] In some embodiments, methods comprise detecting a qualitative change in a target RNA profile generated from a human sample as compared to a normal target RNA profile (in some exemplary embodiments, a target RNA profile of a control sample). Some qualitative changes in the expression profile are indicative of the presence of lung cancer in a sample from a subject. The term "target RNA profile" refers to a set of data regarding the concurrent expression of a plurality of target RNAs in the same sample.

[00227] In some embodiments, at least one, at least two, at least three, at least five, at least 10, or at least 15 of the target RNAs of the plurality of target RNAs are capable of specifically hybridizing to a probe sequence in Table 6. In some embodiments, at least one, at least two, at least three, at least five, at least eight, at least 10, at least 12, at least 15, at least 20, or at least 25 of the target RNAs of the plurality of target RNAs are capable of specifically hybridizing to a a probe sequence in Table 7. In some embodiments, at least one, at least two, at least three, at least five, at least six, or at least seven of the target RNAs of the plurality of target RNAs are capable of specifically hybridizing to a probe sequence in Table 8. In some embodiments, at least one, at least two, at least three, at least four, at least five, at least ten, at least 15, at least 20, at least 25, at least 30, at least 40, or at least 50 of the target RNAs of the plurality of target RNAs are capable of specifically hybridizing to a probe sequence in Table 9. In some embodiments, at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, or at least 40 of the target RNAs of the plurality of target RNAs are capable of specifically hybridizing to a probe sequence in one of Table 32 or 33. In some embodiments, at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 30, at least 40, at least 50, at least 60, or at least 70 of the target RNAs of the plurality of target RNAs is capable of specifically hybridizing to a probe sequence in Table 34.

[00228] In some embodiments, at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, or at least 75 of the plurality of target RNAs comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689. In some embodiments, at least one, at least two, at least five, at least 10, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, or at least 70 of the plurality of target RNAs comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692. In some embodiments, a target RNA, in its mature form, comprises fewer than 30 nucleotides. In some embodiments, a target RNA is a microRNA.

[00229] Qualitative expression data for use in preparing target RNA expression profiles is obtained using any suitable analytical method, including the analytical methods presented herein.

[00230] In some embodiments, for example, concurrent expression data are obtained using, e.g., a microarray, as described above. Thus, in addition to use for quantitative expression level assays of specific target RNAs as described above, a microarray comprising probes having sequences that are complementary to a substantial portion of the miRNome may be employed to carry out target RNA gene expression profiling, for analysis of target RNA expression patterns.

[00231] In some embodiments, distinct target RNA signatures are associated with established markers for lung cancer. In some embodiments, distinct target RNA signatures are associated with established markers for lung cancer caused by bacterial infection, such as for lung cancer caused by gram-positive bacterial infection, lung cancer caused by gram-negative bacterial infection or lung cancer caused by mycobacterial infection. In some embodiments, distinct target RNA signatures are associated with established markers for lung cancer caused by viral infection. In some embodiments, distinct target RNA signatures are associated with established markers for lung cancer caused by multiple infection, such as by co-infection with bacteria and viruses, or by co-infection with more than one viral or more than one bacterial strain. In some embodiments, distinct target RNA signatures are associated directly with the level of severity of the lung cancer.

[00232] According to the expression profiling method, in some embodiments, total RNA from a sample from a subject suspected of having lung cancer is quantitatively reverse transcribed to provide a set of labeled oligonucleotides complementary to the RNA in the sample. The oligonucleotides are then hybridized to a microarray comprising target RNA-specific probes to provide a hybridization profile for the sample. The result is a hybridization profile for the sample representing the expression pattern of target RNAs in the sample. The hybridization profile comprises the signal from the binding of the oligonucleotides reverse transcribed from the sample to the target RNA-specific probes in the microarray. In some embodiments, the profile is recorded as the presence or absence of binding (signal vs. zero signal). In some embodiments, the

profile recorded includes the intensity of the signal from each hybridization. The profile is compared to the hybridization profile generated from a normal, *i.e.*, nonseptic sample, or in some embodiments, a control sample. An alteration in the signal is indicative of the presence of lung cancer in the subject.

4.3. Exemplary additional target RNAs

[00233] In some embodiments, in combination with detecting one or more target RNAs that are capable of specifically hybridizing to a nucleic acid comprising a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689 and/or detecting one or more target RNAs comprising at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692 and/or detecting one or more target RNAs that comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 1363 to 1707, 2064 to 2183, 2312 to 2452, 2673 to 2680, and 2689, methods herein further comprise detecting the level(s) of expression of at least one other marker associated with lung cancer.

[00234] In some embodiments, the methods described herein further comprise detecting altered expression of lung cancer-associated small RNAs with non-canonical hairpins.

[00235] In alternative embodiments, the methods described herein further comprise detecting chromosomal codependents, *i.e.*, target RNAs clustered near each other in the human genome which tend to be regulated together. Accordingly, in further embodiments, the methods comprise detecting the expression of one or more target microRNAs, each situated within the chromosome no more than 50,000 bp from the chromosomal location of the pre-microRNA sequences in Tables 3, 22, 25, 29, and 31.

[00236] In some embodiments, the methods comprise detecting the expression of one or more target RNAs clustered on chromosome 8 at 8p21.3-21.2, at 8p12, at 8p11.21, at 8q11.21, at 8q21.11, or at 8q24.22-24.3; on chromosome 9 at 9p21.3, at 9p13.3, at 9p11.2, at 9q22.32, at 9q31.3-34.11, or at 9q34.3; on chromosome 10 at 10q21.3-23.31, at 10q24.1, at 10q24.32, at 10q25.3, or at 10q26.3; on chromosome 11 at 11p15.5, at 11p14.1, at 11q12.1-23.2, at 11q13.4-14.1, at 11q23.2-24.1, or at 11q25; on chromosome 12 at 12p13.32-13.31, at 12p13.1, at 12p12.1, at 12q12-14.1, at 12q14.3-

21.1, at 12q21.32-23.2, or at 12q24.23; on chromosome 13 at 13q13.3, at 13q14.2, at 13q21.33, at 13q22.3, or at 13q31.3; at chromosome 14 at 14q11.1-13.2, at 14q24.3-31.1, or at 14q32.2-32.31; on chromosome 15 at 15q3, at 15q23-24.32, or at 15q25.3-26.2; on chromosome 16 at 16p13.3-13.2, at 16p11.2, or at 16q12.1-22.3; on chromosome 17 at 17p13.3, at 17p13.1-11.2, at 17q12-21.1, at 17q22-23.3, or at 17q25.31; on chromosome 18 at 18q11.2, at 18q21.31-33, or at 18q22.3; on chromosome 19 at 19p13.3-13.2, at 19p13.12-12, at 19q12, or at 19q13.32-13.42; on chromosome 20 at 20p13, at 20p11.1, at 20q12, or at 20q13.32; on chromosome 21 at 21q21.1-22.11; on chromosome 22 at 22q11.21, or at 22q12.1-13.31; or on the X chromosome at Xp11.4-11.22, at Xq 13.1-13.3, at Xq22.3, at Xq25-27.1, or at Xq27.3-28.

4.4. Pharmaceutical compositions and methods of treatment

[00237] In some embodiments, the disclosure relates to methods of treating lung cancer in which expression of a target RNA is deregulated, *e.g.*, either down-regulated or up-regulated in the lung cancer cells of an individual. When at least one target RNA is up-regulated in the cancer cells, the method comprises administering to the individual an effective amount of at least one compound that inhibits the expression of the at least one target RNA, such that proliferation of lung cancer cells is inhibited. Alternatively, in some embodiments, when at least one target RNA is up-regulated in the cancer cells, the method comprises administering to the individual an effective amount of at least one compound that inhibits the activity of the at least one target RNA, such that proliferation of lung cancer cells is inhibited. Such a compound may be, in some embodiments, a polynucleotide, including a polynucleotide comprising modified nucleotides.

[00238] When at least one target RNA is down-regulated in the lung cancer cells, the method comprises administering an effective amount of an isolated target RNA (*i.e.*, in some embodiments, a target RNA that is chemically synthesized, recombinantly expressed or purified from its natural environment), or an isolated variant or biologically-active fragment thereof, such that proliferation of cancer cells in the individual is inhibited.

[00239] The disclosure further provides pharmaceutical compositions for treating lung cancer. In some embodiments, the pharmaceutical compositions comprise at

least one isolated target RNA, or an isolated variant or biologically-active fragment thereof, and a pharmaceutically-acceptable carrier. In some embodiments, the at least one isolated target RNA corresponds to a target RNA that exhibits a decreased level of expression in lung cancer cells relative to normal levels (in some exemplary embodiments, relative to the level of the target RNA in a control sample). In some embodiments, the isolated target RNA is a target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452. In some embodiments, the isolated target RNA that has a decreased level of expression in lung cancer relative to normal levels is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 92 or 171, or a sequence that is identically present in the probe sequences of Table 34.

[00240] In some embodiments, the pharmaceutical compositions are useful for treating adenocarcinoma and comprise an isolated target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 4, 36, 50, 93, 122, 125, 139, 140, 144, 146, 159, 226, 239 or 241. In some embodiments, the pharmaceutical compositions are useful for treating adenocarcinoma and comprise an isolated target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 19, 27, 33, 48, 55, 72, 73, 94, 101, 105, 112, 117, 130, 131, 133, 134, 135, 143, 155, 158, 160, 161, 163, 165, 221, 238, 240 or 246.

[00241] In some embodiments the isolated target RNA is identical to an endogenous wild-type target RNA gene product (such as a product of a gene set forth in Table 3) that is down-regulated in the cancer cell. In some embodiments, the isolated target RNA is a variant target RNA or biologically active fragment thereof. As used herein, a “variant” refers to a target RNA gene product that has less than 100% sequence identity to the corresponding wild-type target RNA, but still possesses one or more biological activities of the wild-type target RNA (*e.g.*, ability to inhibit expression of a target RNA molecule and cellular processes associated with lung cancer). A “biologically active fragment” of a target RNA is a fragment of the target RNA gene

product that possesses one or more biological activities of the wild-type target RNA. In some embodiments, the isolated target RNA can be administered with one or more additional anti-cancer treatments including, but not limited to, chemotherapy, radiation therapy and combinations thereof. In some embodiments, the isolated target RNA is administered concurrently with additional anti-cancer treatments. In some embodiments, the isolated target RNA is administered sequentially to additional anti-cancer treatments.

[00242] In some embodiments, the pharmaceutical compositions comprise at least one compound that inhibits expression of the target RNA. In some embodiments, the compound is specific for one or more target RNAs, the expression of which is increased in lung cancer cells relative to normal levels (in some exemplary embodiments, relative to the level of the target RNA in a control sample). In some embodiments, the target RNA expression inhibitor is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689. In some embodiments, the target RNA expression inhibitor is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 15, 26, 27 or 191. In some embodiments, the target RNA expression inhibitor is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 15, 26, 27, 30, 129, 164, 184, 191, 196, 205, 207, 214, 219, 225, 246 or 248. In some embodiments, the target RNA expression inhibitor is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe in one of Tables 6, 7, 8, 9, 32, and 33. In some embodiments, a target RNA expression inhibitor that is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe in Table 6 is useful for treating non-small cell lung cancer. In some embodiments, a target RNA expression inhibitor that is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe in Table 7 is useful for treating squamous cell carcinoma. In some embodiments, a target RNA expression inhibitor that is specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe in Table 8 is useful for treating adenocarcinoma. In some embodiments, a target RNA expression inhibitor that is

specific for at least one target RNA that is capable of selectively hybridizing to at least one nucleic acid probe in Table 9 is useful for treating aggressive forms of lung cancer.

[00243] In some embodiments, the pharmaceutical compositions are useful for treating squamous cell carcinoma and comprise a target RNA expression inhibitor that is specific to a target RNA capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 4, 36, 50, 93, 122, 125, 139, 140, 144, 146, 159, 226, 239 and 241. In some embodiments, the pharmaceutical compositions are useful for treating squamous cell carcinoma and comprise a target RNA expression inhibitor that is specific to a target RNA capable of selectively hybridizing to at least one nucleic acid probe comprising a sequence that is identically present in one of SEQ ID NOs: 19, 27, 19, 27, 33, 48, 55, 72, 73, 94, 101, 105, 112, 117, 130, 131, 133, 134, 135, 143, 155, 158, 160, 161, 163, 165, 221, 238, 240 or 246. In some embodiments, the pharmaceutical compositions are useful for treating squamous cell carcinoma and comprise a target RNA expression inhibitor that is specific to a target RNA capable of selectively hybridizing to at least one nucleic acid probe in Table 7. In some embodiments, the pharmaceutical compositions are useful for treating adenocarcinoma and comprise a target RNA expression inhibitor that is specific to a target RNA capable of selectively hybridizing to at least one nucleic acid probe in Table 8. In some embodiments, the pharmaceutical compositions are useful for treating aggressive forms of lung cancer and comprise a target RNA expression inhibitor that is specific to a target RNA capable of selectively hybridizing to at least one nucleic acid probe in Table 9.

[00244] In some embodiments, the target RNA inhibitor is selected from double-stranded RNA, antisense nucleic acids and enzymatic RNA molecules. In some embodiments, the target RNA inhibitor is a small molecule inhibitor. In some embodiments, the target RNA inhibitor can be administered in combination with other anti-cancer treatments, including but not limited to, chemotherapy, radiation therapy and combinations thereof. In some embodiments, the target RNA inhibitor is administered concurrently with other anti-cancer treatments. In some embodiments, the target RNA inhibitor is administered sequentially to other anti-cancer treatments.

[00245] In some embodiments, a pharmaceutical composition is formulated and administered according to Semple et al., Nature Biotechnology advance online publication, 17 January 2010 (doi:10.1038/nbt.1602)), which is incorporated by reference herein in its entirety for any purpose.

[00246] The terms “treat,” “treating” and “treatment” as used herein refer to ameliorating symptoms associated with lung cancer, including preventing or delaying the onset of symptoms and/or lessening the severity or frequency of symptoms of the lung cancer.

[00247] The term “effective amount” of a target RNA or an inhibitor of target RNA expression or activity is an amount sufficient to inhibit proliferation of cancer cells in an individual suffering from lung cancer. An effective amount of a compound for use in the pharmaceutical compositions disclosed herein is readily determined by a person skilled in the art, *e.g.*, by taking into account factors such as the size and weight of the individual to be treated, the stage of the disease, the age, health and gender of the individual, the route of administration and whether administration is localized or systemic.

[00248] In addition to an isolated target RNA or a target RNA inhibitor, or a pharmaceutically acceptable salt thereof, the pharmaceutical compositions disclosed herein further comprise a pharmaceutically acceptable carrier, including but not limited to, water, buffered water, normal saline, 0.4% saline, 0.3% glycine, and hyaluronic acid. In some embodiments, the pharmaceutical compositions comprise an isolated target RNA or a target RNA expression inhibitor that is encapsulated, *e.g.*, in liposomes. In some embodiments, the pharmaceutical compositions comprise an isolated target RNA or a target RNA expression inhibitor that is resistant to nucleases, *e.g.*, by modification of the nucleic acid backbone as described above in Section 4.1.5. In some embodiments, the pharmaceutical compositions further comprise pharmaceutically acceptable excipients such as stabilizers, antioxidants, osmolality adjusting agents and buffers. In some embodiments, the pharmaceutical compositions further comprise at least one chemotherapeutic agent, including but not limited to, alkylating agents, anti-metabolites, epipodophyllotoxins, anthracyclines, vinca alkaloids, plant alkaloids and terpenoids,

monoclonal antibodies, taxanes, topoisomerase inhibitors, platinum compounds, protein kinase inhibitors, and antisense nucleic acids.

[00249] Pharmaceutical compositions can take the form of solutions, suspensions, emulsions, tablets, pills, pellets, capsules, capsules containing liquids, powders, sustained-release formulations, suppositories, emulsions, aerosols, sprays, suspensions, or any other form suitable for use. Methods of administration include, but are not limited to, oral, parenteral, intravenous, oral, and by inhalation.

[00250] The following examples are for illustration purposes only, and are not meant to be limiting in any way.

5. EXAMPLES

5.1 Example 1: MicroRNAs from Primary Lung Tumors

[00251] Using microarray analysis, distinct microRNAs were demonstrated to be deregulated (*e.g.*, either over-expressed or under-expressed) in primary lung tumors.

[00252] Total RNA was extracted from each of the primary tumors in Table 10 by preparation from frozen tissue samples as described below.

RNA Preparation from frozen tissue samples

[00253] Archived or freshly snap-frozen tumor specimens were homogenized by mortar and pestle in TRIzol® Reagent, Invitrogen (Carlsbad, CA) and RNA was further extracted according to the manufacturer's protocol. All RNA samples were diluted in RNase-free water and stored at -80°C.

Table 10

Tumor designation	Sex	Age	Smoking Risk Factor	Cancer type	Stage
Adk-1	M	59	Smoker	Non Squamous, adenocarcinoma	YT2N2
Adk-2	M	67	Smoker	Non Squamous, adenocarcinoma	PT2N0
Adk-3	M	78	na	Non Squamous, adenocarcinoma	PT1N0
Adk-8	F	51	na	Non Squamous, adenocarcinoma	PT3N0
Adk-9	M	55	Smoker	Non Squamous, adenocarcinoma	PT1N0
Adk-10	M	49	Smoker	Non Squamous, adenocarcinoma	PT3N0
Adk-11	M	73	na	Non Squamous, adenocarcinoma	PT1N0

Table 10

Tumor designation	Sex	Age	Smoking Risk Factor	Cancer type	Stage
Epi-4	M	78	Smoker	Squamous, Epidermoid carcinoma	PT2NX
Epi-5	M	78	Smoker	Squamous, Epidermoid carcinoma	PT2NX
Epi-7	F	84	Non-smoker	Squamous, Epidermoid carcinoma	PT2NX
Normal-lung_1	M	81	na	Normal tissue from patient	PT2N0
Normal-lung_2	na	na	na	Normal tissue from Ambion	na

[00254] Epi4, Epi7, Epi5, Adk1, Adk3, Adk11, Adk8, Adk9 and Adk2 identify primary tumors resected from 10 patients. Epi (epidermoid) is squamous cell carcinoma, while Adk (adenocarcinoma) is non-squamous cell carcinoma.

[00255] In Table 10, above, “na” indicates that the information was not available. “Y” in the staging column indicates that the patient was treated before surgical resection. “P” indicates that the patient was not treated before surgical resection.

[00256] Staging is according to the “TNM” staging system for NSCLC. “T” categories for NSCLC indicate the following: (a) T1 indicates that the tumor is no larger than 3 centimeters across, has not reached the membranes that surround the lungs, and does not affect the main branches of the bronchi; (b) T2 tumors possess one or more of the following features: (i) larger than 3 centimeters across; (ii) involves a main bronchus, but is not closer than 2 cm to the carina (the point where the windpipe splits into the left and right main bronchi); (iii) has grown into the membranes that surround the lungs; or (iv) partially clogs the airways, but has not caused the entire lung to collapse or develop pneumonia; (c) T3 indicates a tumor of any size having one or more of the following features: (i) tumor has grown into the chest wall, the diaphragm, the membranes surrounding the space between the two lungs, or membranes of the sac surrounding the heart; (ii) tumor invades a main bronchus and is closer than 2 cm to the carina, but it does not involve the carina itself; or (iii) tumor has grown into the airways

enough to cause an entire lung to collapse or to cause pneumonia in the entire lung. The “N” number indicates involvement of the lymph nodes as follows: (a) N0 indicates that the cancer has not spread to nearby lymph nodes; (b) N2 indicates spread to lymph nodes around the carina or in the space behind the breastbone and in front of the mediastinum. Affected lymph nodes are on the same side as the primary tumor; and (c) NX indicates that nearby lymph nodes could not be accessed.

[00257] Total RNA from normal lung tissue resected from a patient (in this case from a location in the lung as distal as possible from the tumor) and normal lung tissue from a location adjacent to a lung tumor (Ambion) were used as the control.

Total RNA preparation and analysis

[00258] Total RNA was isolated by using standard TRIzol[®] protocol (Invitrogen). Cells from two confluent 75cm² flasks were harvested (= approx 10⁷ cells). Total RNA was prepared using TRIzol[®] Reagent, Invitrogen (Carlsbad, CA) according to the manufacturer's protocol. All RNA samples were diluted in RNase-free water and stored in -80°C (-112°F).

[00259] RNA quality was assessed by calculating OD 260/280 ratios. The quality of all RNA samples was high as assessed using an Agilent Bioanalyser 2100, as exemplified by the electropherogram shown in FIG. 1 obtained for total RNA from A549 human adenocarcinoma cell line. Similar electropherograms were obtained for total RNA from the other primary tumor samples as well.

MicroRNA Purification

[00260] MicroRNA purification was performed using a Flash PAGE Fractionator (Ambion). The Ambion gel purification protocol enriches for small RNAs less than 40 nucleotides (nt) long, including microRNAs. Briefly, a total RNA sample was loaded onto a pre-cast gel using the Flash PAGE Fractionator. The total RNA fraction smaller than 40 nt (the “microRNA fraction”) was recovered after gel migration and resuspended into nuclease free water.

Microarray Analysis

Probe design and spotting

[00261] The oligonucleotide probes used for microarray preparation had the configuration 5'-NH₂-(C)₆-(spacer)-(oligomer probe sequence)-3'. The 5'-amino group

allowed chemical bonding onto the array support. Each also included an identical spacer sequence of 15 nt, as shown below, to prevent non-specific interactions of the oligonucleotide probes with the array support:

5'AminoC6-TTGTAATACGACTCA – Oligo probe sequence (SEQ ID NO: 1044)

Probe sequences given in Table 1 and Table 2 omit the linker.

[00262] The probes were synthesized according to standard protocols by Eurofins MWG Operon (Ebersberg, Germany). Nexterion (Schott) microarray glass slides were used as the solid support for the microarray.

[00263] The oligonucleotide probe concentration used for the spotting was 25 μ mol. The probes were spotted in duplicate using the Nexterion spotting buffer provided with the array glass support by Schott with 1 % SDS (sodium dodecyl sulfate) added to allow larger spot sizes (*e.g.*, 100-150 microns compared to 70-100 microns without SDS). The spotter used was the QArray mini (Genetix) equipped with Stealth SMP3 pins (Telechem). After deposition of one series of spots, the spotting needle was washed 5 times with 60mM NaOH before spotting the next series of probes. Each slide is designed with 32 blocks of spotted probes, with each block being a 20x20 square of spotted probes. Each probe was spotted in duplicate. Spotted glass slides were stored at 4°C until use.

MicroRNA labelling

[00264] The labelling of the microRNA fraction was adapted from a published protocol developed at EMBL (Heidelberg, Germany) by the European Molecular Biology Group (Castoldi *et al.*, “A sensitive array for microRNA expression profiling (miChip) based on locked nucleic acids (LNA),” *RNA* 2006 May;12(5):913-20. Epub 2006 Mar 15, incorporated herein by reference in its entirety). Briefly, the microRNA fraction was incubated for 6 hours at 4°C with a mixture containing 10 μ M of dye-labelled tetra-nucleotide (5'-rUrUrUrU –Cy5- 3') (or alternatively, 5'-rUrUrUrU – Cy3-3') (Biospring, Germany) in Ambion buffer diluted to 1X with RNase free water, 8% polyethylene glycol (PEG), 2 mM adenosine triphosphate (ATP), and T4 RNA ligase (0.7U/ μ l). The labelling reaction was run by heating the mixture for 15 minutes at 65°C. This procedure ligated the poly-U dye-labelled tail to the 3' end of all the microRNAs. Labelled samples were stored at 4°C before hybridization.

Array Hybridization

[00265] The labelled microRNA fraction was hybridized to the spotted arrays using a Discovery hybridization station (Ventana, Tucson, AZ). Briefly, 2 mL of a mixture of 1% BSA, 2X SSC, and 0.2 % SDS was incubated with the chips for 30 min at 42°C. Then the chips were washed once using EZ Prep buffer (Ventana) and then three more times with Ribowash (Ventana). Next, 20 µl of the labelled microRNA mixture and 180µl of ChipHybe Reagent (Ventana) were added to the array. The arrays were heated for 6 minutes at 37°C, then were incubated at 42°C for 8 hours, after which the heating was stopped. The chips were washed once with Ribowash (Ventana) and then heated for 2 minutes at 37°C. The chips were washed again with Ribowash (Ventana) with one drop of CheapClean (Ventana) added, and incubated for 2 minutes at 37°C. The chips were washed two more times using Ribowash (Ventana). The chips were stored dry overnight. On the following day, the final washes were done according to Ventana's instructions for the Discovery hybridization station. The slides were washed twice with 2X SSC + 0.2X SDS buffer and then one more time with 0.1X SSC. All the slides were dried using a speed centrifuge from Arrayit (TeleChem International, Sunnyvale, CA) at room temperature and kept in the dark before scanning.

[00266] As an alternative to the ChipHybe Reagent solution (solution 1), the following solution may be used for array hybridization (solution 2) to form probe:target RNA hybrids by mixing 2 parts of the 1.5X TMAC Hybridization Solution to 1 part (v:v) sample, so that the final component concentrations are 3M TMAC, 0.10% Sarkosyl, 50 mM Tris, and 4 mM EDTA, and incubating on the array at 42°C for 8 hours:

1.5X TMAC Hybridization Solution

Reagent	Catalog Number	Final Conc	Amount/ 250 mL
5 M TMAC*	Sigma T3411	4.5 M	225 mL
20% Sarkosyl	-----	0.15%	1.88 mL
1 M Tris-HCl, pH 8.0	Sigma T3038	75mM	18.75 mL
0.5 M EDTA, pH 8.0	Invitrogen 15575-020	6mM	3.0 mL
H ₂ O	-----	-----	1.37 mL

*TMAC is tetramethyl ammonium chloride

[00267] The number of technical replicates hybridized for each biological sample is shown below in Table 11.

Table 11

	Normal		Squamous cell carcinoma			Non-squamous carcinoma						
Samples	Lung1	Lung2	Epi4	Epi5	Epi7	Adk1	Adk2	Adk3	Adk8	Adk9	Adk10	Adk11
G2	3	4	4	1	2	3		2	3		5	2
G3			2				2			1	2	

Array Image Acquisition

[00268] The arrays were scanned using an Axon™ scanner (Molecular Devices, Sunnyvale, CA) and their Genepix™ software. The image was formatted in tif format, defined by an image color depth of 16 bits/pixel (1600*1600). At such setting, pixels can assume intensity values ranging from 0 to 65,535. Pixels exhibiting the maximum intensity value are “saturated” and were assigned the value of 65,535. The resolution of the array scan was set at 10 μm/pixel. For hybridization experiments using different fluorescent dyes (e.g., Cy5 and Cy3) the photomultiplier tube (PMT) was adjusted to the higher intensity spot (Cy3 is scanned at lower PMT settings than Cy5).

Array Image Analysis

[00269] The PMT of the laser scanner digitized the captured fluorescence intensity for each given “point” of a slide and stored the numerical value as a pixel corresponding to that point. A picture composed of such pixels was then analyzed.

[00270] The first task for image analysis was to detect the spot position, using a process called segmentation. Spots were segmented by circles of adaptable or fixed radius. To be reliably segmented and quantified, the spot diameter was required to be more than 5 - 6 pixels. Before segmentation an indexing grid was provided giving the approximate positions of the spots. The segmentation itself detected the limits of spots near the grid circles. Briefly, the Genepix software assigns a circle to each spot on the array (segmentation). The segmentation had to be conducted in a somewhat flexible way

due to spotting imperfections and/or support deformation, as the spots were almost never on a perfectly rectangular grid.

[00271] After segmentation by the software, the circles were modified manually and adjusted onto the spots until all the spots on the array were clearly identified. At this stage, if the array presented high background noise preventing real spots from being distinguished from the background, the array was rejected for further analysis.

[00272] The second task of image analysis was to quantify spots and export the data into a result file. This was a relatively easy and well-defined task once the spots were located on the image. The statistical approach used most frequently to quantify spot intensity was the mean or median of pixels belonging to a spot. The median approach was more robust than the mean value in the presence of outlier pixels. In practice, however, there was little difference in the results obtained using mean or median.

Array Data Analysis

[00273] All the array data were analysed using the R bioconductor package (“Bioconductor: open software development for computational biology and bioinformatics,” *Genome Biol.* 2004;5(10):R80. Epub 2004 Sep 15, which is incorporated herein by reference in its entirety).

[00274] Array data were first tested for quality by comparing the spot intensities for the internal controls. (Table 12) One internal control (SEQ ID NO: 1046) was used as a labelling control (this synthetic RNA is added to the purified microRNA fraction before labelling), and 7 other internal controls (SEQ ID NOs: 1047 to 1052 and 1045) were used for the normalization of the data (these synthetic RNA controls are added to the total RNA fraction before hybridization at 520 fmol each/array).

Table 12
Internal controls added to total RNA or microRNA fraction

CGCGCGUCGCUUUAUCUACUGU	SEQ ID NO: 1046; CTL30_COMP
UUAUCGUUCGAUAAGUCGCGUU	SEQ ID NO: 1047; CTL11_COMP
GAAGUUACUAUGUAGGCAACCU	SEQ ID NO: 1048; CTL23_COMP
CGCGGGACUAAUUGUUACCGGG	SEQ ID NO: 1049; CTL26_COMP
UCGCGUCGAACUCCGCAACCGA	SEQ ID NO: 1050; CTL29_COMP
ACCGAACGCCGUACCCAUCGGG	SEQ ID NO: 1051; CTL31_COMP
CGAGGGUAAACGACUCUCGUGUC	SEQ ID NO: 1052; CTL36_COMP

Table 12
Internal controls added to total RNA or microRNA fraction

GCGUACCGACGCGUAGACGGAC	SEQ ID NO: 1045; CTL13_COMP
------------------------	-----------------------------

Table 13
Probes for hybridization of control sequences in microarray experiments

Sequence (5'-3')	Sequence identification number
TTGTAATACGACTCAACAGTAGATAAAGCGACGCGCG	SEQ ID NO: 1054; CTL30
TTGTAATACGACTCAAACGCGACTTATCGAACGATAA	SEQ ID NO: 1055; CTL11
TTGTAATACGACTCAAGGTTGCCTACATAGTAACTTC	SEQ ID NO: 1056; CTL23
TTGTAATACGACTCACCCGGTAACAATTAGTCCCGCG	SEQ ID NO: 1057; CTL26
TTGTAATACGACTCATCGGTTGCGGAGTTCGACGCGA	SEQ ID NO: 1058; CTL29
TTGTAATACGACTCACCCGATGGGTACGGCGTTCGGT	SEQ ID NO: 1059; CTL31
TTGTAATACGACTCAGACACGAGAGTCGTTACCCTCG	SEQ ID NO: 1060; CTL36
TTGTAATACGACTCACCCGGTAACAATTAGACCCGCG	SEQ ID NO: 1061; CTL26_MUT
TTGTAATACGACTCAGTCCGTCTACGCGTCGGTACGC	SEQ ID NO: 1062; CTL13
TTGTAATACGACTCAGGCCGTCTACGCGTCGGTACGC	SEQ ID NO: 1053; CTL13_MUT

[00275] All sequences for which the intensity of the spot was higher than the mean local background intensity plus 1.5 times its standard deviation were categorized as expressed microRNAs. The following criteria were required to be met in order consider the array intensity data valid for further analysis:

1. Specificity of the hybridization controls had to be within acceptance criteria (e.g. CTL26 vs. its corresponding single base mutant, CTL26_MUT, or CTL13 vs. its corresponding single base mutant, CTL13_MUT).
2. Approximate equality of the signal intensity of the replicates of the positive controls
3. Approximate equality between median block signal intensities based on the positive controls for each block
4. Approximate equality between median array signals based on all sequences detected
5. Signal intensity for the purification and labelling control (CTL30).

[00276] Statistical normalization of the data was done by computing the Log2ratio where the Log2ratio equals average intensity signal of the duplicated spots/median intensity of all positives controls for the block. The normalization was done per block to avoid non-homogenous labelling of all blocks of the array. This block-by-block normalization has been shown to be more efficient then using overall normalization of the slide. The obtained values are Log2 values.

[00277] The intensities of the spots for each oligonucleotide probe were compared in the sample from the primary lung tumors versus normal lung tissue, resulting in an evaluation of the relative expression for each microRNA.

[00278] The expression fold-change corresponds to $2^{(\text{Log2ratio})}$. The Log2ratio is the ratio between the two conditions compared, or $\log_2(\text{Xcell-line}/\text{Xnormal})$, which is the same as $(\log_2\text{Xcell-line} - \log_2\text{Xnormal})$, where X is the measured intensity value. In cases where there was no signal from the “normal” condition, the lowest measured intensity value in the experiment was used as the baseline from which a fold-change expression value was calculated. A fold-change value of less than zero corresponds to a down-regulation of $(1/\text{fold-change})$ times.

[00279] Data are tabulated in Table 1. The numerical values set forth in Table 1 indicate the fold-change calculated by comparing the signal intensity measured for hybridization to a particular probe in each tumor cell sample with the signal intensity measured for hybridization to the same probe in the control sample (NHBE cells). When the signal detected for hybridization to a particular probe in both the control sample and the tumor cell sample was below the threshold value, the indication “nd” is given in the Table and no fold-change was calculated. Likewise, when hybridization to a particular array probe was detected in the control sample but not in a tumor cell sample, the fold-change calculation is preceded by “nd” in the Table. When no signal was detected in the control sample, the threshold value was retained for the fold-change calculation. This results in an underestimation of the fold-change in the tumor cell sample as compared to the control sample, which is indicated respectively by a “>” or “<” sign in Table 1 for up-regulation and down-regulation of a target RNA.

5.2 Example 2: Analysis of target RNA from Lung Cancer Cell Lines

Cell lines

[00280] Total RNA was prepared from three different lung cell lines that are commonly used in studies of lung cancer. The RNA was used for target RNA array profiling.

[00281] As set forth in Table 14 below, cell lines were selected for diversity, deriving from adenocarcinomas, squamous cell carcinomas and large cell carcinomas. All cell lines were purchased from LGC Promochem (ATCC) and cultured according to ATCC's guidelines.

Table 14

Cell Line	ATCC Number	Cancer type
NHBE	CC-2540	Normal bronchial epithelial cells
BEA2B	CRL-9609	Immortalized bronchial epithelial cells - normal phenotype
A549	CCL-185	Adenocarcinoma
H1703	CRL-5889	Adenocarcinoma
H460	HTB-177	Carcinoma (big neuroendocrine cells)

[00282] Microarray data acquisition and analysis were conducted as described in Example 1.

[00283] Total RNA from normal bronchial epithelial cells (NHBE, ATCC no. CC-2540) was used as a control.

[00284] Data are tabulated in Table 2. The tabulated values correspond to the fold-changes calculated by comparing the signal intensity measured for hybridization to a particular array probe in a primary tumor sample with the mean signal intensity for hybridization to the same probe in the control samples. When the signal was below the threshold value for both the control sample and the primary tumor sample, no fold-change was calculated (indicated by “nd” in Table 2). When hybridization to a particular probe was detected in the control sample, but not in a primary tumor sample, the fold-change calculation is preceded by “nd”. When no signal was detected for hybridization to an array probe in the control samples, the threshold value was retained for the fold-change calculation, which resulted in an underestimation of the fold-change in the primary tumor

sample as compared to the control sample. This is indicated in Table 2 respectively by a “>” or “<” sign for up-regulated and down-regulated target RNAs.

5.3 Example 3: Analysis of target RNA on Luminex Platform

[00285] The Luminex technology (Luminex Corp., Austin, TX) is based on liquid phase hybridization to probe-labelled beads, followed by flow cytometry detection of beads with differing ratios of fluorescent dyes. Beads with up to 100 different dye ratios are available, making it possible to interrogate a single sample for up to 100 analytes simultaneously.

Coupling of Probes to Luminex Beads

[00286] Aliquots of each 5'-amino-modified probe having sequences as set forth in Example 1 and Table 1 are prepared at a concentration of 0.1nmol/ μ L in molecular biology grade water. The probes are coupled to the beads using carbodiimide chemistry according to the manufacturer's protocol (Luminex bead coupling protocol). The probe-coupled beads are stored at 4°C.

Total RNA Preparation for Luminex Analysis

[00287] Fifty fmoles of each of 7 internal controls (the same synthetic RNAs used for the array controls) are added to the total RNA fraction isolated from the biological samples. Prior to hybridization with Luminex beads, the total RNA preparation is treated to avoid the formation of dendrimers, which result from the circularization of a single RNA molecule, or concatenation to another RNA molecule. To avoid the formation of dendrimers, the RNA is pre-treated with calf intestinal phosphatase (CIP) to remove the 5'-phosphate groups. The CIP reagent can be obtained from Invitrogen (Carlsbad, CA) and the CIP reaction is run according to the manufacturer's protocol.

Bead Labelling and Hybridization

[00288] After CIP treatment, the total RNA fraction is then labelled with biotin using the Vantage microRNA Labelling Kit (Marligen). The labelled fraction is hybridized to the Luminex beads using the Marligen protocol. Briefly, the oligonucleotide beads are mixed with the Marligen hybridization solution (1.5 X TMAC) and the labelled total RNA. The hybridization is performed at 60°C for an hour in the dark. After

hybridization, the beads are washed using the Luminex standard 6X SSPET wash buffer (sodium phosphate, sodium chloride, EDTA, Triton X-100, pH 7.4).

Detection of Bead Hybridization

[00289] The detection of the Luminex beads is done using streptavidin phycoerythrin (SAPE) (Europa Bioproducts, Cambridge, UK). The SAPE is added to the washed beads according to the Luminex protocol. The beads are then read using the Luminex IS-200 instrument using the high gain setting for better resolution

Data Acquisition and Analysis

[00290] The Luminex IS-200 reads at least 25 beads of each dye-ratio in the reaction mix. Each dye-ratio bead corresponds to a particular probe sequence, and the intensity value is returned as an average value of all read beads. The mean fluorescence intensity (MFI) data is normalized using synthetic RNA controls, and fold changes between normal and diseased samples are computed using the Bioplex software (Bio-Rad, Hercules, CA) and the R bioconductor package (Bioconductor: open software development for computational biology and bioinformatics, Genome Biol. 2004;5(10):R80. Epub 2004 Sep 15).

5.4 Example 4: Analysis of target RNA from Additional Primary Lung Tumors

[00291] Three of the primary lung tumors analyzed in Example 1 were included in this analysis (Adk-9, Adk-10, and Epi-4). The microRNA fractions isolated from those tumors in Example 1 were used in the microarray experiment described below.

Patient cohort

[00292] In addition to the three tumors listed above (and included in Table 15 in the first three lines), fifteen patients diagnosed with lung cancer were included in the cohort, nine men and six women. Normal tissue from an additional patient (Ksarc-17) was also included in the study. Because squamous cell lung cancer and non-squamous cell lung cancer are two of the most frequent lung cancer types (more than 70% of all lung cancers), four patients diagnosed with squamous cell carcinoma (Kmalp-21, Kmalp-25, Epi-42, and Kmalp-44) and seven patients diagnosed with non-squamous cell carcinoma (Adk-40, Adk-41, Adk-48, Adk-49, Adk-15, Adk-23, and Adk-29) were included, in addition to the previous squamous cell carcinoma (Epi-4) and non-squamous

cell carcinoma (Adk-9, and Adk-10) patients. In addition, one patient diagnosed with a carcinoid (Car-13), one diagnosed with a carcinoma sarcomatoid (Ksarc-19), one diagnosed with a small cell lung cancer (Scc-27), and one diagnosed with a large cell neuroendocrine cancer (Lcnec-31) were also selected. Table 15 shows a list of the patients and various clinical characteristics of each patient.

Table 15. Clinical characteristics of patients in cohort.

Patient ID	histology	Tissue collected	gender	Birth year or Age	Smoking history	TNM staging	Stage group
Adk-9	adenocarcinoma	PT	M	1953	40 smokes / day	T1N0M0	IA
Adk-10	adenocarcinoma	PT	M	1959	25 smokes / day	T3N0M0	IIB
Epi-4	sqamous	PT	M	1930	30 smokes / day	T2NxM0	IIIA
Adk-29	adenocarcinome	PT & DNT	M	1932	? smokes / day	T4N2M1	IV
Adk-15	adenocarcinoma	PT & DNT	M	1947	? smokes / day	T2N0M0	IB
Adk-23	adenocarcinome	PT & DNT	F	1971	0	T3N0M0	IIB
Ksarc-19	carcinome sarcomatoide	PT & DNT	M	1949	0	T4N1M0	IIIB
Kmalp-25	Epidermoide	PT & DNT	M	1928	0	T2N0M0	IB
Kmalp-21	Epidermoide	PT & DNT	F	1942	0	T3N1M0	IIIA
Car-13	Carcinoide	PT & DNT	F	1952	0	T3N1M0	IIIA
Lcnec-31	neuroendocrine GC	PT & DNT	M	1946	0	T3N2M0	IIIA
Scc-27	petites cellules	PT & DNT	F	1943	0	T2N0M0	IB
Adk-40	Non Squamous, adenocarcinoma	PT	M	76	NA	T1aN0	IA
Adk-41	Non Squamous, adenocarcinoma	PT	F	61	Smoker	T2N0M0	IB
Adk-48	Non Squamous, adenocarcinoma	PT	F	56	NA	T1bN0M0	IA
Adk-49	Non Squamous, adenocarcinoma	PT	M	53	NA	T3N2	IIIA
Epi-42	Squamous, Epidermoid carcinoma	PT	M	59	NA	T3N1M0	IIIA

Kmalp-44	Squamous, Epidermoid carcinoma	PT	M	na	NA	T2bN0	IB
Ksarc-17	Carcinome Sarcomatoide	DNT	M	53	NA	T2N0M0	IB

PT = Primary Tumor

DNT = Distal Normal Tissue

NA= information not available

[00293] Five of the patients had very aggressive forms of lung cancer, Ksarc-19 (carcinoma sarcomatoide), Adk-9, Adk-10, Adk-29 (all adenocarcinoma), and Lcnec-31. Patient Ksarc-19 relapsed one month after surgery and died six months later. Patient Car-13 (carcinoide) had a slow growing form of lung cancer.

[00294] The primary tumors only were collected from six of the patients (as well as the four previous patients, for a total of 10 patients), while the primary tumor and distal normal tissue were collected from nine of the patients. Distal normal tissue only was used from one of the patients (Ksarc-17). The relative amount of tumor cells versus normal cells in primary tumors was determined to be between 90% and 100% for all patients. The distal normal tissue was collected from a location as far as possible from the primary tumor, and did not contain detectable tumor cells.

Tissue samples

[00295] Archived or freshly snap-frozen specimens from lung primary tumors were used. Tissue samples were homogenized by mortar and pestle in TRIzol® Reagent (Invitrogen; Carlsbad, CA) and RNA was extracted according to manufacturer's protocol. RNA samples were diluted in RNase-free water and stored in -80°C (-112°F).

MicroRNA Preparation:

[00296] All samples were enriched for the microRNA fraction using a Flash PAGE Fractionator (Ambion). Briefly, a total RNA sample was loaded onto a pre-cast gel using the Flash PAGE Fractionator. The total RNA fraction smaller than 40 nt (the "microRNA fraction") was recovered after gel migration and resuspended into nuclease free water.

Microarray Analysis

Probe design and spotting

[00297] The polynucleotide probes used for microarray preparation had the configuration 5'-NH₂-(C)₆-(spacer)-(oligomer probe sequence)-3'. The 5'-amino group allowed chemical bonding onto the array support. Each also included an identical spacer sequence of 15 nt, as shown below, to prevent non-specific interactions of the polynucleotide probes with the array support:

5'AminoC6-TTGTAATACGACTCA – Oligo probe sequence. (SEQ ID NO: 1044)

Probe sequences given in the Tables herein omit the linker.

[00298] The probes were synthesized according to standard protocols by Eurofins MWG Operon (Ebersberg, Germany). Nexterion (Schott) microarray glass slides were used as the solid support for the microarray.

[00299] The polynucleotide probe concentration used for the spotting was 25 µmol. The probes were spotted in duplicate using the Nexterion spotting buffer provided with the array glass support by Schott with 1 % SDS (sodium dodecyl sulfate) added to allow larger spot sizes (*e.g.*, 100-150 microns compared to 70-100 microns without SDS). The spotter used was the QArray mini (Genetix) equipped with Stealth SMP3 pins (Telechem). After deposition of one series of spots, the spotting needle was washed 5 times with 60mM NaOH before spotting the next series of probes. Each slide is designed with 48 blocks of spotted probes, with each block being a 20x18 square of spotted probes. Each probe was spotted in duplicate. Spotted glass slides were stored at 4°C until use.

MicroRNA labelling

[00300] The labelling of the microRNA fraction was adapted from a published protocol developed at EMBL (Heidelberg, Germany) by the European Molecular Biology Group (Castoldi *et al.*, “A sensitive array for microRNA expression profiling (miChip) based on locked nucleic acids (LNA),” *RNA* 2006 May;12(5):913-20. Epub 2006 Mar 15, incorporated herein by reference in its entirety). Briefly, the microRNA fraction was incubated for 6 hours at 4°C with a mixture containing 10 µM of

dye-labelled tetra-nucleotide (5'-rUrUrUrU -Cy5- 3') (or alternatively, 5'-rUrUrUrU -Cy3-3') (Biospring, Germany) in Ambion buffer diluted to 1X with RNase free water, 8% polyethylene glycol (PEG), 2 mM adenosine triphosphate (ATP), and T4 RNA ligase (0.7U/ μ l). The labelling reaction was run by heating the mixture for 15 minutes at 65°C. This procedure ligated the poly-U dye-labelled tail to the 3' end of all the microRNAs. Labelled samples were stored at 4°C before hybridization.

Array Hybridization

[00301] Table 16 shows the number of array replicates carried out for each tumor sample.

Table 16: Number of Replicates for Sample

Sample	Tumor type	Number replicates
Adk-9	Non-squamous	1
Adk-10	Non-squamous	2
Epi-4	Squamous	2
Adk-29	Non-squamous	3
Adk-15	Non-squamous	3
Adk-23	Non-squamous	3
Ksarc-19	Squamous	2
Kmalp-25	Squamous	1
Kmalp-21	Squamous	3
Car-13	Carcinoid	3
Lcnec-31	Large cell neuroendocrine	2
Scc-27	Small cell	2
Adk-40	Non-squamous	2
Adk-41	Non-squamous	3
Adk-48	Non-squamous	2
Adk-49	Non-squamous	3

Epi-42	Squamous	3
Kmalp-44	Squamous	3
Ksarc-17	Normal	3
Adk-29	Normal	1
Adk-15	Normal	2
Adk-23	Normal	3
Ksarc-19	Normal	1
Kmalp-25	Normal	1
Kmalp-21	Normal	1
Lcnec31	Normal	1
Scc27	Normal	1

[00302] The labelled microRNA fraction was hybridized to the spotted arrays using a Discovery hybridization station (Ventana, Tucson, AZ). Briefly, 2 mL of a mixture of 1% BSA, 2X SSC, and 0.2 % SDS was incubated with the chips for 30 min at 42°C. Then the chips were washed once using EZ Prep buffer (Ventana) and then three more times with Ribowash (Ventana). Next, 20 µl of the labelled microRNA mixture and 180µl of ChipHybe Reagent (Ventana) were added to the array. The arrays were heated for 6 minutes at 37°C, then were incubated at 42°C for 8 hours, after which the heating was stopped. The chips were washed once with Ribowash (Ventana) and then heated for 2 minutes at 37°C. The chips were washed again with Ribowash (Ventana) with one drop of CheapClean (Ventana) added, and incubated for 2 minutes at 37°C. The chips were washed two more times using Ribowash (Ventana). The chips were stored dry overnight. On the following day, the final washes were done according to Ventana's instructions for the Discovery hybridization station. The slides were washed twice with 2X SSC + 0.2X SDS buffer and then one more time with 0.1X SSC. All the slides were dried using a speed centrifuge from Arrayit (TeleChem International, Sunnyvale, CA) at room temperature and kept in the dark before scanning.

Array Image Acquisition

[00303] The arrays were scanned using an Axon™ scanner (Molecular Devices, Sunnyvale, CA) and their Genepix™ software. The image was formatted in tif format, defined by an image color depth of 16 bits/pixel (1600*1600). At such setting, pixels can assume intensity values ranging from 0 to 65,535. Pixels exhibiting the maximum intensity value are “saturated” and were assigned the value of 65,535. The resolution of the array scan was set at 10 μ m/pixel. For hybridization experiments using different fluorescent dyes (e.g., Cy5 and Cy3) the photomultiplier tube (PMT) was adjusted to the higher intensity spot (Cy3 is scanned at lower PMT settings than Cy5).

Array Image Analysis

[00304] The PMT of the laser scanner digitized the captured fluorescence intensity for each given “point” of a slide and stored the numerical value as a pixel corresponding to that point. A picture composed of such pixels was then analyzed.

The first task for image analysis was to detect the spot position, using a process called segmentation. Spots were segmented by circles of adaptable or fixed radius. To be reliably segmented and quantified, the spot diameter was required to be more than 5 - 6 pixels. Before segmentation an indexing grid was provided giving the approximate positions of the spots. The segmentation itself detected the limits of spots near the grid circles. Briefly, the Genepix software assigns a circle to each spot on the array (segmentation). The segmentation had to be conducted in a somewhat flexible way due to spotting imperfections and/or support deformation, as the spots were almost never on a perfectly rectangular grid.

[00305] After segmentation by the software, the circles were modified manually and adjusted onto the spots until all the spots on the array were clearly identified. At this stage, if the array presented high background noise preventing real spots from being distinguished from the background, the array was rejected for further analysis.

[00306] The second task of image analysis was to quantify spots and export the data into a result file. This was a relatively easy and well-defined task once the spots were located on the image. The statistical approach used most frequently to quantify spot intensity was the mean or median of pixels belonging to a spot. The median approach

was more robust than the mean value in the presence of outlier pixels. In practice, however, there was little difference in the results obtained using mean or median.

Array Data Analysis

[00307] All the array data were analysed using the R bioconductor package (“Bioconductor: open software development for computational biology and bioinformatics,” *Genome Biol.* 2004;5(10):R80. Epub 2004 Sep 15, which is incorporated herein by reference in its entirety).

[00308] Array data were first tested for quality by comparing the spot intensities for the internal controls. One internal control (SEQ ID NO: 1046; Table 17) was used as a labelling control (this synthetic RNA is added to the purified microRNA fraction before labelling), and 6 other internal controls (SEQ ID NOs: 1047 to 1052) were used for the normalization of the data (these synthetic RNA controls are added to the total RNA fraction before hybridization at 520 fmol each/array). The probe sequences that bind to the synthetic RNAs, and a mutant probe sequence, are also shown in Table 17 (SEQ ID NOs: 1054 to 1061).

Table 17
Control Sequences used in microarray experiments

Sequence (5'-3')	Sequence identification number
CGCGCGUCGCUUUAUCUACUGU	SEQ ID NO: 1046; CTL30_COMP
UUAUCGUUCGAUAAGUCGCGUU	SEQ ID NO: 1047; CTL11_COMP
GAAGUUACUAUGUAGGCAACCU	SEQ ID NO: 1048; CTL23_COMP
CGCGGGACUAAUUGUUACCGGG	SEQ ID NO: 1049; CTL26_COMP
UCGCGUCGAACUCCGCAACCGA	SEQ ID NO: 1050; CTL29_COMP
ACCGAACGCCGUACCCAUCGGG	SEQ ID NO: 1051; CTL31_COMP
CGAGGGUAACGACUCUCGUGUC	SEQ ID NO: 1052; CTL36_COMP
TTGTAATACGACTCAACAGTAGATAAAGCGACGCGCG	SEQ ID NO: 1054; CTL30
TTGTAATACGACTCAAACGCGACTTATCGAACGATAA	SEQ ID NO: 1055; CTL11
TTGTAATACGACTCAAGGTTGCCTACATAGTAACTTC	SEQ ID NO: 1056; CTL23
TTGTAATACGACTCACCCGGTAACAATTAGTCCCGCG	SEQ ID NO: 1057; CTL26
TTGTAATACGACTCATCGGTTGCGGAGTTCGACGCGA	SEQ ID NO: 1058; CTL29
TTGTAATACGACTCACCCGATGGGTACGGCGTTCGGT	SEQ ID NO: 1059; CTL31
TTGTAATACGACTCAGACACGAGAGTCGTTACCTCG	SEQ ID NO: 1060; CTL36
TTGTAATACGACTCACCCGGTAACAATTAGACCCGCG	SEQ ID NO: 1061; CTL26_MUT

[00309] All sequences for which the intensity of the spot was higher than the mean local background intensity plus 1.5 times its standard deviation were categorized as expressed microRNAs. The following criteria were required to be met in order consider the array intensity data valid for further analysis:

1. Specificity of the hybridization controls had to be within acceptance criteria (e.g. CTL26 vs. its corresponding single base mutant, CTL26_MUT).
2. Approximate equality of the signal intensity of the replicates of the positive controls
3. Approximate equality between median block signal intensities based on the positive controls for each block
4. Approximate equality between median array signals based on all sequences detected
5. Signal intensity for the purification and labelling control (CTL30).

[00310] Statistical normalization of the data was done by computing the Log2ratio where the Log2ratio equals average intensity signal of the duplicated spots/median intensity of all positives controls for the block. The normalization was done per block to avoid non-homogenous labelling of all blocks of the array. This block-by-block normalization has been shown to be more efficient then using overall normalization of the slide. The obtained values are Log2 values.

[00311] The intensities of the spots for each polynucleotide probe were compared in the sample from the tumors versus normal tissue, resulting in an evaluation of the relative expression for each microRNA.

[00312] The expression fold-change corresponds to $2^{(\text{Log2ratio})}$. The Log2ratio is the ratio between the two conditions compared, or $\log_2(\text{Xcell-line}/\text{Xnormal})$, which is the same as $(\log_2\text{Xcell-line} - \log_2\text{Xnormal})$, where X is the measured intensity value. In cases where there was no signal from the “normal” condition, the lowest measured intensity value in the experiment was used as the baseline from which a fold-

change expression value was calculated. A fold-change value of less than zero corresponds to a down-regulation of (1/fold-change) times.

Results

[00313] The microarray data was analyzed to identify target RNAs that were present at increased levels in more than half of the tumor samples tested. Tables 18 and 19 show the target RNAs, in what percentage of tumor samples they were present in increased levels, and the average fold-increase in those tumor samples in which they were present at increased levels. Data for six of the tumors are shown in Table 18 and data for the remaining 13 tumors are shown in Table 19. Table 18 also shows the probe sequences used to detect the target RNAs.

[00314] The microarray data was further analyzed to identify target RNAs that are present at levels at least 5-fold higher than normal levels in less than 50% of the tumor samples. Tables 20 and 21 show the target RNAs, in what percentage of tumor samples they were present in increased levels, and the average fold-increase in those tumor samples in which they were present at increased levels. Data for six of the tumors are shown in Table 20 and data for the remaining 13 tumors are shown in Table 21. Table 20 also shows the probe sequences used to detect the target RNAs. Finally, Table 22 shows the pre-microRNA sequences and chromosomal location of the pre-microRNA gene for each of the target RNAs in Tables 18 to 21.

[00315] The microarray data was next analyzed to identify target RNAs that were present at at least 5-fold decreased levels in more than half of the tumor samples tested. Tables 23 and 24 show the target RNAs, the probe sequences used to detect the target RNAs, and in what percentage of tumor samples they were present in increased levels. Data for six of the tumors are shown in Table 23 and data for the remaining 13 tumors are shown in Table 24. Table 25 shows the pre-microRNA sequences and chromosomal location of the pre-microRNA gene for each of the target RNAs in Tables 23 and 24.

[00316] Finally, Table 9, above, shows target RNAs that were present at higher levels in at least three of the five most aggressive cancers (Ksarc-19, Adk-9, Adk-10, Adk-29, and Lnec-31).

Table 18: Target RNAs present at increased levels in at least 50% of tumor samples

Gene	Probe sequence	SEQ ID NO	% tumors increased	Fold Chng Avg	ADK9	ADK10	Adk29	Adk15	Adk23	ADK48
10455-L5-1	TAACAACTGCAATTGCGCCAAACATCCCTCTCC	1063	58	6.4	2.05	-3.86	1.02	5.18	2.20	-3.86
12947-L5-4	GTTCCCTTCTCCACAGCCTCTAAATCTCCCATCCG	1064	63	5.0	3.17	2.58	-1.79	1.50	2.10	1.18
13098-L5-1	TGCCGGGGCGCTGTTGGCGGTTCCTCCGGGGGGT	1065	58	4.4	-2.86	-2.86	-2.86	1.70	-1.49	5.93
13122-L5-1	TTAGGAAATTCATCTCACCTGCTCCAGTCC	1066	58	11.5	1.32	1.17	1.40	1.68	1.56	2.09
13219-L5-1	CTCTGACTCCCTCACTCAGTCTCTCTGCTCCAGC	1067	68	7.8	1.54	-1.85	1.30	1.90	1.44	2.19
13254-R5-1	CAATGAACCACTGAACCACTCATGCACTGAACC	1068	63	5.0	2.21	6.72	2.48	2.40	1.52	4.02
13467-L5-1	GTGACAGTCAGACCCCTCCTTGCTCCAAGTCAAA	1069	53	17.7	2.83	3.05	-1.83	3.62	2.22	1.35
3923-R5-1	TCACAAAGGATCTCCTTCATCCCTCTCCAG	1070	58	5.3	-6.96	-3.57	-1.11	6.71	2.53	1.59
4440-L3-2*	TTTGACATTAGAGCACTGGGCAGAAATCACA	1071	68	6.9	2.32	-1.15	1.65	2.00	-1.34	1.29
4479-R3-1	AGCCCCCTGCCCGGAAATTCAAAACAACCTGC	1072	53	4.3	1.27	1.08	-1.38	1.11	1.55	1.75
5080-R3-1	CTTGCAAAAGGTCTCCTTCATCCCTCTCCCA	1073	68	7.2	2.60	-3.84	1.58	8.83	4.24	-1.12
6216-L1-1	GACATTGAGCACTGGGCAGAAATCACATG	1074	53	8.0	-1.89	-1.89	1.27	2.22	-1.89	-1.89
6235-R5-2	TCTGCTCCAAAAATCCATTATATATTGT	1075	74	13.5	2.51	1.63	1.29	3.07	2.15	3.39
7426-L5-1	AGACGAACACCTCCTGCTGTGCTTGATTT	1076	63	5.1	2.59	-1.45	3.22	2.64	1.21	-1.45
7726-R3-2	CATCCCTCTCCAGAAGAGGAGAAAGAGAAACA	1077	68	5.7	2.18	-2.52	1.66	7.34	3.64	-2.52
8004-R3-2	GGAAGTCTTCTCCTTGCTCCAGTCATTGAAG	1078	63	11.0	-5.32	1.60	1.51	1.80	1.80	2.37
836-R5-2	AAATAATCATTCCAAATGGTTCTCCCTGCTAT	1079	79	8.7	2.22	2.37	1.49	1.80	1.93	4.05
9349-R5-2	GAACACAGTGATGCAGAGGACTTCCTGCTCCA	1080	58	24.4	-1.44	4.53	1.39	7.28	1.65	-1.44
9594-R5-1	ACTTAGACTTCTTCCCACCTCCCTGCATCCT	1081	53	9.7	1.32	1.20	-1.03	1.68	1.46	1.95
miR-200a	ACATCGTTACCAGACAGTGTTA	1082	58	6.2	4.00	-1.98	-1.98	1.89	2.18	4.85
miR-200b	TCATCATTACCAGGCAGTATTA	1083	63	4.6	3.65	1.66	1.01	2.11	2.43	2.54
miR-200c	TCCATCATTAACCCGGCAGTATTA	1084	68	3.6	3.05	1.53	-1.02	2.00	3.71	2.39
miR-205	CAGACTCCGGTGAATGAAGGA	1085	53	21.2	6.16	-1.05	7.09	-1.05	-1.05	-1.05
miR-20b	CTACCTGCACTATGAGCACTTTG	1086	53	4.8	3.82	4.97	-2.24	5.93	1.03	2.35
miR-21	TCAACATCAGTCTGATAAGCTA	1087	68	6.7	10.37	4.28	-41.68	8.21	4.78	4.58
miR-298	TGGGAGAACCTCCCTGCTTCTGCT	1088	68	9.8	1.96	1.55	1.87	2.23	2.18	1.30
miR-720	TGGAGGCCCCAGCGAGA	1089	63	4.2	9.81	4.77	3.06	2.73	5.03	4.82

Table 19: Target RNAs present at increased levels in at least 50% of tumor samples (con't)

Gene	% tumors increased	Fold Chng Avg	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
10455-L5-1	58	6.4	-3.86	-3.86	4.88	6.15	11.22	8.61	2.50	20.65	3.92	1.74	1.71	-3.86
12947-L5-4	63	5.0	7.18	-2.89	1.75	2.00	10.83	4.19	2.17	15.66	4.39	3.14	2.39	-2.89
13098-L5-1	58	4.4	6.04	2.12	4.93	7.99	2.92	2.06	1.19	5.45	5.13	1.99	1.68	-2.86
13122-L5-1	58	11.5	-2.23	-3.30	11.06	3.00	45.07	8.66	12.05	25.75	4.32	8.92	4.14	-6.80
13219-L5-1	68	7.8	2.12	-3.09	4.87	2.37	29.25	8.14	10.45	23.80	3.38	7.05	3.81	-5.56
13254-R5-1	63	5.0	-5.53	-5.53	4.42	3.62	1.67	2.00	5.81	14.41	2.64	1.20	-1.16	-2.34
13467-L5-1	53	17.7	-3.86	-3.86	6.58	-1.54	47.10	19.28	11.18	87.67	-1.56	6.95	4.54	-3.86
3923-R5-1	58	5.3	1.69	1.36	5.55	6.66	7.94	6.74	2.87	12.31	3.49	1.89	1.78	-6.96
4440-L3-2*	68	6.9	14.22	-2.38	4.89	5.53	4.84	4.40	4.36	20.54	7.80	5.37	9.54	-2.38
4479-R3-1	53	4.3	-14.83	1.85	-1.15	5.75	5.60	6.31	-14.83	10.98	1.91	4.07	1.95	2.02
5080-R3-1	68	7.2	-3.84	-1.79	8.12	10.22	12.60	9.22	3.75	18.91	6.17	2.71	2.74	-3.84
6216-L1-1	53	8.0	15.02	-1.89	2.61	-1.89	5.37	4.69	3.76	22.08	9.22	5.16	9.56	-1.89
6235-R5-2	74	13.5	-2.30	-1.76	11.09	4.52	53.43	16.58	19.84	39.60	6.59	11.81	11.43	-8.31
7426-L5-1	63	5.1	2.23	-1.45	6.10	-1.45	6.82	4.16	3.33	11.69	5.48	5.61	7.41	-1.45
7726-R3-2	68	5.7	-2.52	-2.52	8.06	9.94	8.81	7.20	3.54	11.79	2.24	2.37	2.77	-2.52
8004-R3-2	63	11.0	-1.69	-5.32	6.36	3.58	48.62	7.86	12.44	28.62	4.41	9.48	5.21	-5.32
836-R5-2	79	8.7	5.02	-2.64	5.63	4.02	40.75	6.59	10.21	26.72	4.38	8.08	5.16	-2.64
9349-R5-2	58	24.4	-1.44	-1.44	10.49	3.33	49.05	33.90	19.27	112.09	8.79	10.70	8.85	-1.44
9594-R5-1	53	9.7	-6.58	-6.58	4.10	2.60	30.50	7.67	10.86	24.28	3.32	6.82	4.54	-6.58
miR-200a	58	6.2	-1.98	-1.98	5.20	4.55	-1.98	1.86	1.17	3.60	3.49	4.09	3.32	26.93
miR-200b	63	4.6	-1.76	2.73	3.27	2.23	-4.79	1.16	1.30	2.97	2.05	4.07	2.98	24.67
miR-200c	68	3.6	-8.29	2.84	1.30	2.38	-8.29	-1.03	2.11	3.38	2.05	4.09	4.89	11.97
miR-205	53	21.2	-1.05	-1.05	8.11	13.68	3.20	27.74	56.45	30.05	46.50	-1.05	-1.05	-1.05
miR-20b	53	4.8	-2.24	2.99	8.37	4.56	-1.45	-1.10	1.71	-2.24	-2.24	6.46	3.10	-2.24
miR-21	68	6.7	-6.36	6.76	17.86	8.91	3.81	-41.68	4.64	1.92	7.46	3.38	1.19	1.41
miR-298	68	9.8	-2.56	-2.56	6.60	5.82	36.81	9.96	10.86	31.32	5.07	7.72	4.43	-2.56
miR-720	63	4.2	6.70	-1.87	2.31	-1.87	2.72	1.99	1.95	-1.87	4.11	1.35	-1.87	-1.87

Table 20: Target RNAs present at increased levels in less than 50% of tumor samples, but with an average increase of at least 5-fold relative to normal levels

Gene	Probe sequence	SEQ ID NO	% tumors increased	Fold Change Average	ADK9	ADK10	Adk29	Adk15	Adk23
10083-L5-1	CCCTCTTCCTTTCTACCCCTCTCTCCACCC	1090	16	11.2	-1.64	-4.49	-10.22	-3.32	-1.50
10233-R5-1	ACCTCTCCCCCTTGATCTGGAGCAGCAGGTAGA	1091	16	13.1	-1.55	-7.07	-4.19	-1.46	-1.77
10335-L5-2	CCACTCCCCTCCTTTTAAATTAGAAAGCAC	1092	16	5.3	-1.72	-8.23	-8.23	-2.21	-3.36
10357-R5-1	CTCTCCACTGTCTGAATATTCATGTGTTGTTCCCTAG	1093	16	6.0	-1.13	-1.13	1.25	2.75	1.29
10520-L5-1	CAGCATCTCCTTGCTACCTCCTC	1094	26	5.9	-1.42	-1.42	-1.42	-1.42	1.14
10844-R4-1	AGGGGTAGCATATTTCAGGACTCTACTTTCAAA	1095	26	5.4	-2.58	-2.58	-2.58	1.36	-2.00
11358-R5-2	ATGGAGGGTGGGAACTAGGCTGACATGGCTC	1096	21	5.3	-2.77	-2.77	-2.77	1.77	-2.18
11444-L5-3	GCAC TTGTGTT CAGGCACCCACTCTCCAGCAAAG	1097	21	8.8	-1.20	-1.20	-1.20	-1.20	-1.20
11547-R4-1	GCCCCCTGCTCTGGCTGGTCAACGGAACCAAGTCCGTCT	1098	16	7.6	-2.09	-2.09	-2.09	-2.09	-1.66
11605-L5-4	TCCCTTCTTTAATTCCTTCCCCTCAATCCATAA	1099	26	5.0	-2.54	-2.54	-2.54	-2.54	-1.37
11688-R5-3	GAGACGCCGACTGGCTGAGACTCAAGCGGGG	1100	32	6.0	-1.33	-1.33	-1.33	-1.33	-1.33
12699-R5-2	AGGGTGGTGGCGGTGGTGGCGGAACGCTGGGGGG	1101	26	6.0	-1.96	-1.96	-1.96	1.14	-1.60
12707-L5-2	CATTCTCCTCTGCAATCCAACTCTAC	1102	42	5.0	-1.91	-1.91	-1.20	3.36	-1.40
12729-R5-1	CGACGAAATGCAATTGTGTGCTTCTCCCTCC	1103	16	11.8	-1.10	-5.55	-5.55	-1.51	-2.63
12730-R5-1	TTGTGCTCCGCTCTCCGGGAAATGCCATCACTAAT	1104	37	7.3	-3.66	-3.66	-2.24	1.55	-2.46
12888-L5-2	TTTCTACATTGTATGGTTCTCCAGCTCCT	1105	47	21.7	4.55	1.76	1.24	5.05	1.51
12907-L5-1	ATGCCCTCTCTACCACGTCCTAGACACTGAGCC	1106	26	6.4	-3.23	-3.23	-3.23	-1.53	-1.46
12912-L5-2	TCAAGTTTTCTGGCACCTTCCACCACAGAGGCT	1107	37	5.3	-2.26	-2.26	-1.29	2.16	-2.26
12917-R5-2	GGACCTATGGGCCCTTCCCTTCCCCCAACATTG	1108	47	6.0	1.03	-3.76	-1.02	1.54	2.26
12958-R5-1	TCTTCCCTGGGTGCATCTATAACGACTACAGA	1109	32	5.1	-1.32	-1.32	-1.32	-1.32	-1.32
12974-R5-2	ACCTCTGTGCCCCCAATTCCTAGTCCCTTGGT	1110	21	16.1	-1.94	-1.94	-1.94	-1.94	-1.30
12979-R5-2	TCCAGGTCCTTCCCCAGTAGCTAGATGAAGAAAA	1111	42	16.3	-1.39	-1.39	1.39	6.66	4.45
12992-L5-1	CATACCTGCTTCCCTCCACCCCATCTCTA	1112	16	6.7	-9.54	-2.35	-9.54	-2.64	-3.60
13001-L5-1	TTCTAGATACCACTCCAGCTCCA	1113	32	15.5	-1.83	-1.83	-1.09	1.61	1.27
13053-R5-4	GCCTTCCCTAAGATCAGGCTCATCAGGCAGCCCCA	1114	26	5.6	-1.32	-1.32	-1.32	-1.32	1.25
13070-R5-3	CTGTCTCCTCTCCCCAGTCCAAAGGACCTAATGC	1115	16	20.1	-1.13	-4.70	-4.70	-1.17	-2.22
13071-L5-3	AGAGAAAGGGTGAGGATCTCCAGGGGTGACTGCT	1116	32	7.2	-2.11	-2.11	-2.11	1.69	-2.11
13078-L5-1	GCCCTCATTTTCATAAGCCTGCTAGACAGGAGA	1117	32	5.0	-1.28	-1.28	-1.28	-1.28	1.18
13185-L5-3	TTCTGTTTTTTTCTTCCCTCTCTCCCTCTCT	1118	21	13.4	-1.02	-1.33	-6.52	-1.77	-1.16
13216-R5-2	ATATATTCTCCTTTCTCCTCTCTGCGAGCTGA	1119	26	5.1	-1.79	-1.79	-1.79	-1.79	-1.03
13225-L5-1	CTGCCCACTCTGGGACAGCCCCCTTCCATCTTC	1120	16	5.9	-3.37	-3.37	-3.37	-1.24	-3.37
13235-R5-1	CCTGGGCTTCTAGTCTCAGCTCTCCTCCAGCT	1121	42	5.1	1.75	2.72	-2.00	-2.00	-1.22
13245-L5-4	ATTCCAAATAGTCTTTCCTTCTACTCCCTTTAAA	1122	16	12.2	-2.87	-2.87	-2.87	-2.87	-2.25
13252-L5-3	ACGTGCCCTTCTGACTGTGAGCTCCTTGAGAGC	1123	32	5.0	3.73	2.50	-1.79	3.92	1.23
13274-L5-3	CCTTCTCTTCTCCCGTCTCCACCCCTCCCTCAGGG	1124	26	6.2	-1.53	-1.73	-2.51	-1.80	-1.69
13300-L5-1	TTACACACAGAAATTATCTCCCTCCTACTTACCC	1125	16	7.4	-1.04	-6.08	-6.08	-2.44	-3.08

Gene	Probe sequence	SEQ ID NO	% tumors increased	Fold Change Average	ADK9	ADK10	Adk29	Adk15	Adk23
13309-R5-1	TCCCCCTTGCTCTCAGCTACAAATTTTCTGTG	1126	16	8.2	1.66	-3.05	-3.05	-3.05	-3.05
13352-R5-1	CCGGAACCTCAGCGCTGCATCTGTGAAATGGGG	1127	37	5.8	-3.13	-3.13	-3.13	1.36	-2.38
13357-L5-4	AGCGGACCTTCTCCCCACACCTCCCTGCAGCCTC	1128	16	10.3	-1.89	-1.91	-6.27	-2.80	-2.38
13366-R5-3	CAGGCCTCACCCCAAGTGCCTCTCCTATTCCCAC	1129	16	6.5	-1.56	-5.82	-11.70	-1.40	-1.25
13373-R5-2	TGTGGGGGAGGGTGGTCAGCAGGTGGGCAG	1130	42	5.1	-1.86	-1.86	-1.86	3.87	-1.44
13375-R5-4	GTACCAGCTCTCAGACCCCTCACCACAGCCTGGGG	1131	32	5.8	-7.06	-7.06	-7.06	-2.47	-1.39
13398-R5-4	TTCTCTCTCCCTCCACCTTTTGTGCCACCACTT	1132	16	15.6	-5.01	-2.59	-5.01	-1.56	-2.35
13417-R5-4	TGCCCTTCCTTCTGTTAACACCCAGCCAGATCCCC	1133	32	6.4	-1.82	-1.82	-1.82	2.57	1.34
13436-L5-1	AGCTCAATGAAGCCCCCAGCTAGTGCATGCCTC	1134	21	5.3	-2.80	-8.31	-20.29	-2.23	-2.79
13468-L5-1	CAAGTGGCCATCAGACCCCTCTTTGCCCCCAAGTC	1135	21	25.5	-1.16	-1.16	-1.16	-1.16	1.07
13470-R5-1	TCCCACCCCTCTCCACCTCAGGGACCAGATCCT	1136	21	13.2	-1.29	-3.12	-6.50	-1.24	-1.23
13473-L5-3	TATGTTTGCCCTTCTACCACCTATCCTGATACA	1137	21	9.2	-1.57	-1.57	-1.57	1.32	1.04
13500-L5-3	CCAAGATCTTCCAGGCCTCTCTCTCCATTGAGT	1138	37	6.4	-1.38	-1.38	-1.38	2.14	1.20
13522-L5-4	CTGCCTCACCTGAGCTCCCGTGCCTGTGCACCTC	1139	16	5.3	-3.11	-3.11	-3.11	-3.11	-2.36
13523-R5-1	GTGGAATAGTAAGGCTTCTTCCCTGCCTGGC	1140	26	5.3	-3.75	-3.75	-1.20	1.11	1.61
13545-L5-1	GCCTGTGGCGAAGGGAGGCTGTGAGTCTGGGG	1141	21	5.3	-2.13	-2.13	-2.13	1.70	-2.13
227-L5-1	ACACCTGTCTCTCCCCAGTGTCTCCGCCCTCA	1142	16	5.0	-2.46	-3.09	-2.60	-3.09	-3.20
3744-R5-1	CTTCTCCTTCCCTCCCTGCTCCCCCTCCCAATAAT	1143	37	6.7	-1.38	-2.00	-2.67	-1.30	-1.23
3875-R5-2	GACTGATTCAACCTCTCTCTCCCACTT	1144	32	19.0	-2.72	-2.72	-2.72	1.47	-1.32
3926-L5-1	ACTGATAATTCAATTAGCATTTCTTGAACGTGCCCCACT	1145	16	5.1	-2.22	-2.22	-2.22	-2.22	-2.22
3992-R5-1	GCCCCTTTCTCTCTGTCTGTCAGTATCTGAGCTGATTG	1146	32	6.7	-1.27	-1.27	-1.27	-1.27	-1.27
4203-R3-2	GCACATTCCCACTTCCCCAGAGGCGAGGCTCCATAT	1147	16	6.0	-2.01	-2.01	-2.01	-1.45	-2.01
4261-R5-1	AATAACCCCTCCTTCTTGATAGATTTCAC	1148	21	6.4	-2.52	-2.52	-2.52	-2.52	-1.56
4291-R5-1	GATCAGGTAGTTACCAAGGAGCCATGGGGTGTA	1149	37	6.0	-1.67	-1.67	-1.67	-1.25	-1.67
4790-L5-2	TAACTGTCTCCCTCATTAAGTAAATCT	1150	21	12.6	-1.56	-1.56	-1.56	-1.56	-1.32
4958-L5-1	TAACTGTCTCCCTCATTAAGTAAATCT	1151	32	5.0	-2.14	-2.14	-2.14	-1.00	-2.14
4988-R5-2	GAGAAATTAATAAACCATGGCATCTTCTGAAAATGGGG	1152	16	5.3	-2.97	-3.81	-7.10	-2.52	-2.60
5108-R5-2	CTCCTCCTCCCGTCTTTGGATACCAACAC	1153	21	16.5	-1.14	-1.67	-4.77	-1.11	-1.10
5232-L5-2	CCCTCTCCTCCACACCGTCACTACAATAACCC	1154	32	5.2	-1.81	-1.81	1.81	-1.34	-1.44
5392-R5-1	TCCTCCTCCTTCATTTGCAGGACATCAAGG	1155	42	7.0	-2.12	-2.12	-2.12	2.25	-1.14
5723-R5-1	CCCTCTCCGACAGTATCTCATTAACAATAATTGCTAATCTC	1156	16	6.9	-1.96	-3.89	-8.51	-5.58	-3.53
5836-L5-1	CCTCCTCCCTTCTCCAGCAGTAGCCTTCTTAA	1157	26	5.0	-2.14	-2.14	-2.14	1.48	-2.14
5842-R5-1	TCTTGGGAACCAAGGGGCTACAG	1158	32	6.0	2.40	-5.08	-5.08	2.54	1.13
6037-R3-2	GCAATTCCCTTCTCCTCCATCTCCAATTTTCTCTC	1159	32	7.4	-1.71	-1.71	-1.71	-1.71	-1.32
6181-L5-1	CATGGTAACTTAGATGACTCTTCCCTGGATTT	1160	26	8.6	-1.40	-1.40	-1.40	-1.40	-1.40
6233-L5-2	CAAAATTAGATTTCACCTTTATCCTTCTCTCC	1161	16	10.1	-3.02	-3.02	-3.02	-1.89	-1.92
6395-L5-1	TCTCCCCCTGCAGAGAGGAGGAAAGCCTGCCTCGGAAC	1162	21	8.4	-3.73	-3.73	-3.73	1.09	-1.93
6405-R5-1	TCAAAGAAAAATAAGTGAAGCATATCTCATCATGGGG	1163	32	5.7	-2.10	-2.10	-2.10	-2.10	-2.10
6474-L5-1	CCCTAATTAAAGCCATCCCTCTTCCCTTCCACC	1164	16	11.9	-11.16	-5.02	-11.16	-1.96	-1.85
6602-R3-1	AGAGCCCCAGTGGAAATCTCTCTCCCAATCCAT	1165	32	8.8	-1.36	-1.36	-1.36	-1.36	-1.06
6681-R2-1	CCTGTTTTCTCCCCCTCTCTCTCTGCCCCCTCC	1166	16	5.0	-1.60	-2.15	-3.96	-1.87	-1.45

Gene	Probe sequence	SEQ ID NO	% tumors increased	Fold Change Average	ADK9	ADK10	Adk29	Adk15	Adk23
6683-R5-1	TCATCCCCAAATAAACTCTTCCTGCTCAAG	1167	37	11.9	-1.39	-1.39	-1.39	1.04	1.29
6752-R5-2	CCCTCCTTTCCCCACCTCAGTCGGCCACTGCT	1168	16	6.4	-7.01	-7.01	-4.90	-2.94	-2.85
6803-R5-2	GCTCCCTCTCTGTTGGACCTCACCCAAA	1169	21	12.2	1.30	-2.22	-1.17	-1.05	-1.19
6864-R4-1	GACACACAATTACACTCCCTGGAG	1170	21	12.5	-1.50	-1.50	-1.50	-1.50	-1.50
6872-L5-1	AAGACTCTGTCACAGTCTGTGACCCGGTGGGG	1171	42	5.2	-1.96	-1.96	-1.96	-1.00	-1.96
6906-L5-1	TCTTCTCCCCAACCAAGCCAGCTCTCCTGG	1172	21	8.0	-2.17	-2.17	-2.17	-2.17	-2.17
6930-R5-1	TTAATCCTTCTCTCCCTCTGTGATCTTGCGAG	1173	16	14.8	-1.27	-1.66	-4.45	-1.91	-1.65
7631-L3-2	CCTAGACGCCAGCTGCCTGCATGAAGCCTGCCAA	1174	26	5.2	-2.68	-2.68	-2.68	1.68	2.02
7764-R3-2	CCCTCTCTGCCCTCTCTCATCACCAATAACAGAC	1175	16	9.3	-1.03	-3.64	-3.64	-1.33	1.24
7849-L2-2	GAAATATCAAGACAAATAAGATGTTTGGGGCACA	1176	16	6.1	-2.04	-2.04	-2.04	-1.42	-2.04
8316-R5-1	CAGGGTATCCTCTCCCCACGTGATGCC	1177	42	20.3	2.51	-1.81	-1.81	1.36	1.12
8433_C-R4-1	AAACCAAAAAAAAAAATTAAAAAGCGACGAAAAATGCAATTGTGTGCCTTCTCCCTCC	1178	16	14.3	-1.03	-2.46	-5.42	-1.38	-1.21
8433-L3-1	AAATGGCTCCTTTCCCTTTCCCTCCACCG	1179	16	6.9	-1.70	-1.71	-5.48	-2.04	-1.62
8452-L3-1	ATGCCACATACATGGCTGGGTGTTGAA	1180	21	7.6	-1.08	-1.08	-1.08	-1.08	-1.08
8724-R5-2	GGCCAAAGCTTGGAACCTCTCCCTGCCAGCA	1181	16	6.0	-1.71	-2.12	-4.29	-1.92	-1.69
8746-R5-1	TAAGCGCAGGATGGGGTATTCAAGTCCAAAGC	1182	37	5.4	-2.03	-2.03	-2.03	1.54	-2.03
8808-R5-1	GACCCCTTTCTCCAGCCTGTTTCTGCAA	1183	21	8.4	-1.07	-2.31	-4.54	-4.54	-2.03
8832-R5-1	TGGAGTACCACCTGTTTTCCCCACTT	1184	32	6.0	-1.37	-1.37	-1.37	-1.37	-1.06
8879-L5-1	TGGGGCTGGGGCGCAGGTCGAGAGCCCGGG	1185	32	5.2	-1.79	-1.79	-1.79	-1.30	-1.79
9485-L5-1	TGCTGCACTCCATCCTCCACAATGGCCCCATTGTTCCCGAG	1186	26	5.2	-3.02	-3.02	-3.02	1.26	-3.02
9507-L5-1	GTCTCCCTCATCCATCATCCACCA	1187	26	6.0	-1.45	-1.45	-1.45	-1.45	1.16
9798-L5-1	CTCTATATATACCTGGTATGACTTCTTTGGGGGAAAGAAAGCAC	1188	42	9.9	-1.40	-1.40	-1.40	3.74	1.36
999996-L4-1	GGGAGGAGTCAGGTGTGTGCTGTGGGTTGGGGGAAGAC	1189	21	5.3	-4.63	-4.63	-4.63	1.35	-1.12
999999-R4-1	CCCCTTGTCACCCCCAGCCCCCTTCTGTGCCAGGACCCCCAGCGGAGGCCAGAGAA	1190	16	5.2	-1.71	-9.15	-23.66	-3.15	-2.48
miR-103	TCATAGCCCTGTACAATGCTGCT	1191	26	5.0	1.81	-3.09	1.10	1.01	1.33
miR-1202	CTCCCCCACTGCAGCTGGCAC	1192	16	5.4	-1.58	-7.52	-7.52	-5.40	-4.07
miR-1249	TGAAGAAAGGGGGGAAGGGCGT	1193	32	5.0	-2.26	-2.26	-2.26	1.48	-1.92
miR-1275	GACAGCCTCTCCCCCAC	1194	16	12.7	1.07	-1.12	-12.61	-1.07	-2.42
miR-129-3p	ATGCTTTTTGGGGTAAGGGCTT	1195	11	21.4	-1.00	-1.00	-1.00	-1.00	-1.00
miR-1321	ATCACATTACCTCCCTG	1196	16	5.9	-2.14	-2.14	-2.14	-1.05	-2.14
miR-1323	AGAAAAAGCCCCCTCAGTTTGA	1197	47	5.0	2.09	-1.57	-3.13	1.70	-1.05
miR-141	CCATCTTTACCAGACAGTGTTA	1198	42	5.6	4.01	-1.96	-1.37	2.58	3.17
miR-143	GAGCTACAGTGCTTCATCTCA	1199	42	5.8	1.65	-4.11	1.63	1.38	1.30
miR-182	AGGTGAGTTCTACCATTTGCCAAA	1200	37	8.7	-1.00	-1.00	-1.00	2.28	1.51
miR-183	AGTGAATTCTACCAGTGCCATA	1201	21	10.0	-1.00	-1.00	-1.00	-1.00	-1.00
miR-198	GAACCTATCTCCCTCTGGACC	1202	21	12.6	-4.73	-4.73	-4.73	-1.31	-2.54
miR-19a	TCAGTTTGCATAGATTGCACA	1203	47	5.0	2.88	-1.76	-1.23	2.71	-1.03
miR-30c-1*	GGAGTAAACAACCCCTCTCCCGAG	1204	26	11.2	-1.05	-8.25	-8.25	-1.14	-1.53
miR-375	TCACGCGAGCCGAACGAACAAA	1205	32	16.2	-1.60	4.80	-1.60	2.88	-1.60

Gene	Probe sequence	SEQ ID NO	% tumors increased	Fold Change Average	ADK9	ADK10	Adk29	Adk15	Adk23
miR-376c	ACGTGGAATTTCTCTATGTT	1206	16	62.7	-1.12	-1.12	-1.12	-1.12	-1.12
miR-429	ACGGTTTACCAGACAGATTATTA	1207	16	8.1	3.25	-1.00	-1.00	-1.00	-1.00
miR-483-5p	CTCCCTTCITTCCTCCCGTCTT	1208	42	10.1	2.71	-1.10	-3.38	2.43	1.23
miR-516a-5p	GAAAGTGCTTCTTTTCCTCGAGAA	1209	47	8.8	2.57	-1.65	-1.65	1.35	-1.19
miR-7	ACAACAAAATCACTAGTCTTCCA	1210	21	86.3	-1.00	-1.00	1.53	-1.00	1.56

Table 21: Target RNAs present at levels at least 5-fold higher than normal levels in less than 50% of tumor samples (con't)

Gene	% tumors increased	Fold Change Average	ADK48	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp 21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
10083-L5-1	16	11.2	-3.12	-10.22	-10.22	-4.85	1.60	14.20	6.04	-1.20	13.39	-1.15	1.20	-1.16	-10.22
10233-R5-1	16	13.1	-1.29	-15.77	-1.70	1.17	1.33	17.61	6.52	-1.39	15.23	1.27	1.51	1.10	-6.54
10335-L5-2	16	5.3	-2.42	-8.23	-8.23	-8.23	1.82	5.00	2.55	-1.85	8.20	-8.23	-1.15	-1.71	-8.23
10357-R5-1	16	6.0	-1.13	12.33	-1.13	1.73	-1.13	2.98	-1.13	1.86	-1.13	-1.13	1.37	-1.13	-1.13
10520-L5-1	26	5.9	-1.42	-1.42	-1.42	-1.42	-1.42	7.33	3.18	4.71	10.86	-1.42	3.49	1.34	-1.42
10844-R4-1	26	5.4	5.98	4.89	1.80	4.75	6.90	1.69	1.36	1.08	-2.58	4.49	1.09	-2.58	-2.58
11358-R5-2	21	5.3	4.38	-2.77	1.50	7.69	5.96	1.31	-1.16	1.17	-2.77	2.99	1.18	-2.77	-2.77
11444-L5-3	21	8.8	-1.20	-1.20	-1.20	-1.20	-1.20	13.50	5.64	-1.20	13.27	-1.20	2.62	-1.20	-1.20
11547-R4-1	16	7.6	-2.09	-2.09	-2.09	-2.09	1.98	2.11	5.05	1.30	15.60	-2.09	1.36	-1.23	-2.09
11605-L5-4	26	5.0	-2.54	-2.54	-2.54	-2.54	-2.54	7.43	3.53	1.60	9.69	-2.54	2.07	1.86	-2.54
11688-R5-3	32	6.0	6.88	2.76	1.49	5.01	7.47	1.72	1.11	-1.33	-1.33	4.67	-1.33	-1.33	-1.33
12699-R5-2	26	6.0	1.74	7.41	5.23	5.07	7.67	1.00	1.66	1.16	-1.96	4.62	-1.16	1.61	-1.96
12707-L5-2	42	5.0	-1.91	-1.91	-1.91	6.52	-1.91	7.16	4.03	2.78	-1.91	7.26	4.33	4.33	-1.91
12729-R5-1	16	11.8	-5.55	-5.55	-5.55	-5.55	-5.55	15.59	5.64	-1.24	14.17	1.67	1.74	1.06	-5.55
12730-R5-1	37	7.3	-1.10	-3.66	-3.66	3.20	-1.72	8.17	6.34	1.57	23.70	5.24	2.24	2.37	-3.66
12888-L5-2	47	21.7	-1.64	-1.64	-1.64	7.18	1.22	43.87	23.04	11.51	82.88	1.23	11.54	5.75	-1.64
12907-L5-1	26	6.4	-3.23	-3.23	-3.23	-1.68	4.63	10.17	3.99	1.05	11.02	2.09	1.42	1.32	-3.23
12912-L5-2	37	5.3	-2.26	-2.26	-2.26	-1.02	-2.26	7.22	2.14	2.86	14.96	-2.26	3.88	4.19	-2.26
12917-R5-2	47	6.0	-3.76	-3.76	-3.76	-3.76	2.93	14.76	7.83	2.46	16.79	2.54	2.64	2.10	-3.76
12958-R5-1	32	5.1	-1.32	-1.32	-1.32	-1.32	-1.32	6.38	4.19	1.98	12.26	-1.32	2.45	3.54	-1.32

Gene	% tumors increased	Fold Change Average	ADK48	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp 21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
12974-R5-2	21	16.1	-1.94	-1.94	-1.94	-1.94	-1.94	24.20	12.93	1.29	29.06	-1.94	1.81	1.64	-1.94
12979-R5-2	42	16.3	-1.39	-1.39	-1.39	-1.39	-1.39	10.78	36.28	4.84	58.47	1.46	5.16	4.12	-1.39
12992-L5-1	16	6.7	-9.54	-9.54	-9.54	-4.91	1.18	9.52	3.16	-2.51	7.29	-1.36	-1.30	-1.84	-3.31
13001-L5-1	32	15.5	-1.83	-1.83	-1.83	1.30	-1.83	42.81	17.02	13.68	4.38	-1.83	8.78	6.48	-1.83
13053-R5-4	26	5.6	-1.32	-1.32	-1.32	-1.32	-1.32	3.53	13.64	2.00	6.49	-1.32	2.42	1.51	-1.32
13070-R5-3	16	20.1	-1.56	-1.56	-4.70	-1.74	-2.22	28.78	10.28	-1.18	21.26	1.71	1.74	1.44	-4.70
13071-L5-3	32	7.2	9.95	11.19	-2.11	2.59	8.14	1.71	-1.39	1.41	-2.11	9.31	-1.31	-2.11	1.49
13078-L5-1	32	5.0	-1.28	-1.28	-1.28	-1.28	-1.28	5.67	3.96	2.12	9.20	-1.28	4.09	4.68	-1.28
13185-L5-3	21	13.4	1.19	-10.17	-4.63	-2.11	1.40	22.48	9.50	-1.19	19.78	1.34	1.88	1.66	-1.67
13216-R5-2	26	5.1	-1.79	-1.79	-1.79	-1.79	-1.79	9.13	3.36	1.72	8.24	-1.79	2.44	2.56	-1.79
13225-L5-1	16	5.9	-3.37	-3.37	-3.37	-1.14	-1.38	3.98	4.43	1.04	9.41	-3.37	1.69	1.75	-3.37
13235-R5-1	42	5.1	-2.00	-2.00	-2.00	-2.00	-2.00	4.04	10.73	2.12	9.58	1.76	2.43	2.20	-2.00
13245-L5-4	16	12.2	-2.87	-2.87	-2.87	-2.87	-2.87	14.29	12.06	1.04	10.37	-2.87	1.47	1.29	-2.87
13252-L5-3	32	5.0	7.58	-3.66	-3.66	3.75	-1.66	-1.06	-1.78	1.28	1.36	-3.66	1.14	1.12	-3.66
13274-L5-3	26	6.2	-1.00	-1.48	-2.00	1.50	1.33	13.34	3.11	2.88	9.05	1.27	2.80	1.56	-1.16
13300-L5-1	16	7.4	-1.75	-6.08	-6.08	-1.13	-2.47	5.40	3.67	-1.28	13.10	-6.08	1.07	-1.03	-6.08
13309-R5-1	16	8.2	-3.05	-3.05	-3.05	-3.05	-3.05	9.45	2.79	1.01	12.42	-3.05	1.39	-3.05	-3.05
13352-R5-1	37	5.8	6.85	7.65	-3.13	4.84	4.78	1.36	-1.04	-1.01	-3.13	7.39	1.08	-1.49	3.02
13357-L5-4	16	10.3	-4.08	-13.36	-13.36	-1.26	-1.04	13.07	4.68	-1.56	13.01	-1.17	1.17	-1.45	-13.36
13366-R5-3	16	6.5	-3.19	-11.70	-11.70	1.49	-4.14	6.00	3.36	-1.62	10.24	-4.94	-1.07	-1.49	-11.70
13373-R5-2	42	5.1	8.95	6.03	4.53	2.23	6.76	1.51	4.97	1.11	-1.86	3.80	-1.32	-1.06	-1.86
13375-R5-4	32	5.8	1.45	-1.36	3.34	5.09	3.16	2.32	-1.53	-1.93	4.20	16.42	-1.56	-3.11	-7.06
13398-R5-4	16	15.6	-1.64	-5.01	-5.01	-5.01	-5.01	24.60	7.38	-1.41	14.84	-5.01	1.57	1.33	-5.01
13417-R5-4	32	6.4	-1.82	-1.82	-1.82	2.73	11.74	7.23	4.35	-1.82	9.97	-1.82	-1.82	1.33	-1.82
13436-L5-1	21	5.3	10.40	-20.29	-20.29	3.70	-3.72	-1.06	2.55	-2.03	4.35	-1.04	-1.31	-2.32	-20.29
13468-L5-1	21	25.5	-1.16	-1.16	-1.16	-1.16	-1.16	25.58	42.79	-1.16	31.50	-1.16	2.25	-1.16	-1.16
13470-R5-1	21	13.2	-6.50	-6.50	-6.50	1.75	2.16	22.13	9.28	-1.38	19.17	1.62	1.31	1.04	-6.50
13473-L5-3	21	9.2	-1.57	-1.57	-1.57	-1.57	-1.57	12.99	5.85	1.57	15.53	-1.57	2.39	1.37	-1.57
13500-L5-3	37	6.4	-1.38	-1.38	-1.38	-1.38	-1.38	11.94	6.95	2.11	14.71	-1.38	2.73	3.87	-1.38

Gene	% tumors increased	Fold Change Average	ADK48	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp 21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
13522-L5-4	16	5.3	-3.11	-3.11	-3.11	3.37	-3.11	2.60	1.14	1.10	9.89	-3.11	1.73	1.65	-3.11
13523-R5-1	26	5.3	-3.75	-3.75	-3.75	1.01	-3.75	7.12	4.25	1.30	11.04	1.81	2.32	1.90	-3.75
13545-L5-1	21	5.3	1.56	5.75	4.10	4.33	7.14	1.80	-1.46	1.04	-2.13	-2.13	-2.13	-2.13	-2.13
227-L5-1	16	5.0	-1.10	-8.03	-2.54	-1.26	-1.22	3.33	5.25	-1.86	5.98	1.00	-1.14	-1.09	-4.63
3744-R5-1	37	6.7	1.22	-1.55	-1.88	2.15	1.70	17.73	4.94	4.32	12.79	1.63	2.72	1.99	-2.37
3875-R5-2	32	19.0	-2.72	-2.72	-2.72	-2.72	-2.72	47.37	21.41	1.80	36.52	3.32	2.67	2.60	-2.72
3926-L5-1	16	5.1	-2.22	-2.22	-2.22	-2.22	5.38	1.86	2.06	1.37	7.91	-2.22	1.57	-1.06	-2.22
3992-R5-1	32	6.7	-1.27	-1.27	-1.27	-1.27	-1.27	11.59	6.64	2.32	12.31	-1.27	3.77	3.55	-1.27
4203-R3-2	16	6.0	-2.01	-2.01	-2.01	-2.01	-2.01	7.42	3.12	-2.01	7.32	-2.01	-1.16	-2.01	-2.01
4261-R5-1	21	6.4	1.22	-2.52	-2.52	1.50	4.19	5.56	9.03	1.23	6.64	-2.52	1.69	1.33	-2.52
4291-R5-1	37	6.0	6.21	9.19	5.89	2.44	7.75	-1.03	1.01	-1.67	-1.67	4.73	-1.67	-1.67	-1.67
4790-L5-2	21	12.6	-1.56	-1.56	-1.56	-1.56	-1.56	21.19	7.09	1.61	19.81	-1.56	2.37	1.17	-1.56
4958-L5-1	32	5.0	4.94	6.00	4.13	4.23	6.24	-2.14	-1.19	-1.04	-2.14	4.24	-2.14	-2.14	-2.14
4988-R5-2	16	5.3	1.08	-25.28	-3.70	-1.17	1.20	5.54	3.48	-2.05	6.82	1.28	1.05	-1.34	-8.12
5108-R5-2	21	16.5	-2.27	-7.21	-7.21	1.49	1.77	31.12	11.82	1.05	21.13	1.91	2.06	1.54	-1.29
5232-L5-2	32	5.2	-1.81	-1.81	-1.81	-1.81	5.58	6.39	2.75	2.65	1.47	-1.81	2.30	-1.81	-1.81
5392-R5-1	42	7.0	-2.12	-2.12	-2.12	3.87	4.77	15.89	6.70	1.74	16.76	3.71	1.91	1.15	-2.12
5723-R5-1	16	6.9	-2.80	-8.51	-8.51	-2.09	1.62	8.94	2.94	-1.53	8.79	1.02	1.08	-1.42	-8.51
5836-L5-1	26	5.0	1.59	6.47	4.54	2.07	7.39	1.66	-1.54	-2.14	-2.14	4.43	-2.14	-2.14	-2.14
5842-R5-1	32	6.0	-1.05	-5.08	1.12	-1.02	2.33	2.71	9.50	1.02	16.41	-1.31	-1.32	-1.44	-5.08
6037-R3-2	32	7.4	-1.71	-1.71	-1.71	-1.71	-1.71	16.51	6.11	1.91	14.55	-1.71	2.97	2.07	-1.71
6181-L5-1	26	8.6	-1.40	-1.40	-1.40	-1.40	-1.40	16.60	5.10	1.57	15.61	-1.40	2.68	2.92	-1.40
6233-L5-2	16	10.1	-3.02	-3.02	-3.02	-3.02	-3.02	10.71	4.99	-1.05	14.68	-3.02	1.54	1.57	-3.02
6395-L5-1	21	8.4	-3.73	-3.73	-3.73	1.35	4.16	7.80	6.46	-1.04	15.00	-3.73	1.67	1.45	-3.73
6405-R5-1	32	5.7	6.13	7.07	4.77	5.11	6.86	-2.10	1.46	-2.10	-2.10	4.48	-2.10	-2.10	-2.10
6474-L5-1	16	11.9	-3.05	-11.16	-11.16	-11.16	-4.35	14.82	7.02	-1.48	13.75	-4.80	1.09	-1.19	-11.16
6602-R3-1	32	8.8	-1.36	-1.36	-1.36	-1.36	-1.36	19.18	6.22	3.09	17.33	-1.36	3.82	2.93	-1.36
6681-R2-1	16	5.0	-1.47	-1.37	-1.55	-1.23	1.40	5.10	2.76	-1.63	6.93	-1.55	1.43	-1.17	-3.04
6683-R5-1	37	11.9	-1.39	-1.39	-1.39	1.48	-1.39	26.81	6.23	10.02	25.16	5.08	6.03	4.31	-1.39

Gene	% tumors increased	Fold Change Average	ADK48	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp 21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
6752-R5-2	16	6.4	-2.32	-7.01	-1.56	-7.01	-1.38	7.39	3.11	-1.67	8.79	-7.01	1.18	-1.03	-7.01
6803-R5-2	21	12.2	1.33	1.11	-1.53	1.68	1.59	9.14	18.88	1.50	18.71	1.83	1.99	1.71	-11.71
6864-R4-1	21	12.5	2.22	-1.50	-1.50	1.54	1.44	2.95	1.88	1.78	42.80	-1.50	2.10	1.47	-1.50
6872-L5-1	42	5.2	5.15	7.34	4.47	4.25	5.82	1.59	6.06	-1.96	-1.96	3.74	-1.96	-1.96	-1.96
6906-L5-1	21	8.0	-2.17	-2.17	-2.17	-2.17	-2.17	14.47	3.55	1.13	12.00	-2.17	2.01	1.54	-2.17
6930-R5-1	16	14.8	-2.83	-10.13	-10.13	1.07	1.26	22.64	6.91	-1.19	14.99	1.28	1.74	1.04	-10.13
7631-L3-2	26	5.2	-2.68	-2.68	-2.68	-1.18	7.17	3.13	5.17	1.30	8.27	-2.68	1.40	1.50	-2.68
7764-R3-2	16	9.3	-3.64	-3.64	-3.64	-3.64	-3.64	12.47	4.37	-1.09	11.00	-3.64	1.53	1.54	-3.64
7849-L2-2	16	6.1	1.61	6.57	4.84	1.03	6.78	1.71	-1.05	-2.04	-2.04	-2.04	-1.29	-2.04	-2.04
8316-R5-1	42	20.3	-1.81	-1.81	-1.81	-1.81	-1.81	66.20	16.64	2.12	60.43	6.75	3.99	3.79	-1.81
8433_C-R4-1	16	14.3	-1.74	-5.42	-5.42	-1.36	-2.50	18.24	6.98	-1.37	17.79	1.52	1.68	1.01	-5.42
8433-L3-1	16	6.9	1.35	-8.08	-8.08	-4.05	1.17	8.10	3.41	-1.39	9.18	-1.06	1.19	1.05	-3.51
8452-L3-1	21	7.6	-1.08	11.39	4.06	4.04	11.10	-1.08	-1.08	-1.08	-1.08	-1.08	-1.08	-1.08	-1.08
8724-R5-2	16	6.0	-2.90	-9.03	-4.67	1.24	1.37	6.87	3.34	-1.55	7.87	1.04	1.14	-1.14	-9.03
8746-R5-1	37	5.4	6.65	6.64	4.26	4.91	6.70	1.49	1.68	1.04	-2.03	4.06	-2.03	-2.03	-2.03
8808-R5-1	21	8.4	-4.54	-4.54	-4.54	-4.54	-4.54	12.95	5.01	1.08	13.67	1.70	1.57	1.02	-4.54
8832-R5-1	32	6.0	-1.37	-1.37	-1.37	-1.37	-1.37	4.44	3.33	2.34	19.28	-1.37	3.13	3.30	-1.37
8879-L5-1	32	5.2	5.93	6.75	2.28	1.04	6.13	-1.79	-1.07	-1.79	-1.79	4.14	-1.79	-1.79	-1.79
9485-L5-1	26	5.2	-3.02	-3.02	-3.02	2.49	3.21	3.66	4.65	-1.18	11.90	-3.02	1.00	-1.56	-3.02
9507-L5-1	26	6.0	-1.45	-1.45	-1.45	-1.45	-1.45	8.15	1.99	2.56	14.45	-1.45	2.84	1.67	-1.45
9798-L5-1	42	9.9	14.66	15.37	4.13	9.21	16.73	1.77	4.26	-1.40	-1.40	10.93	-1.40	-1.40	-1.40
999996-L4-1	21	5.3	6.41	-1.05	5.20	4.33	1.88	1.23	-1.40	-1.10	-4.63	5.08	-1.06	-2.07	-1.67
999999-R4-1	16	5.2	9.26	-23.66	-1.25	2.35	-7.93	1.60	1.32	-2.72	3.99	-7.30	-1.89	-2.94	-23.66
miR-103	26	5.0	-1.99	-6.84	1.87	3.43	2.56	1.28	-1.00	1.19	1.66	1.80	3.59	1.79	12.88
miR-1202	16	5.4	-7.52	-7.52	-7.52	-7.52	1.62	6.96	2.14	-2.10	7.21	1.01	-1.32	-1.79	-7.52
miR-1249	32	5.0	5.38	5.84	2.27	4.52	8.04	1.49	-1.12	1.02	-2.26	4.17	1.10	-2.26	-2.26
miR-1275	16	12.7	-12.61	-4.87	-12.61	-1.62	1.01	9.83	7.58	-2.54	20.69	-1.44	-1.20	-1.52	-12.61
miR-129-3p	11	21.4	-1.00	-1.00	-1.00	-1.00	11.04	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	31.72

Gene	% tumors increased	Fold Change Average	ADK48	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp 21	Kmalp 25	EPI-4	Kmalp 44	Scc27	Lcnec31	Car13
miR-1321	16	5.9	-2.14	-2.14	-2.14	-2.14	-2.14	8.17	2.93	1.49	6.53	-2.14	1.63	-2.14	-2.14
miR-1323	47	5.0	1.07	1.16	-3.13	4.58	11.61	4.63	4.00	1.86	8.00	4.21	3.41	1.75	-3.13
miR-141	42	5.6	1.66	-1.96	1.02	5.85	1.14	-1.96	-1.96	1.49	-1.96	1.83	3.28	3.22	16.77
miR-143	42	5.8	4.04	-4.11	2.93	5.90	5.59	-4.11	4.76	-4.11	-4.11	4.40	-4.11	-4.11	7.28
miR-182	37	8.7	-1.00	-1.00	-1.00	11.30	-1.00	-1.00	3.30	2.12	-1.00	-1.00	4.01	7.56	30.62
miR-183	21	10.0	-1.00	-1.00	-1.00	8.60	-1.00	-1.00	1.85	-1.00	-1.00	-1.00	3.52	2.83	25.18
miR-198	21	12.6	-4.73	-4.73	-4.73	-2.51	1.98	21.91	7.44	-1.47	19.01	1.51	1.52	1.17	-4.73
miR-19a	47	5.0	1.80	-1.76	7.21	8.91	6.26	-1.04	1.14	1.68	4.45	-1.76	3.38	2.05	-1.76
miR-30c-1*	26	11.2	-2.57	-8.25	-3.89	1.91	1.94	22.43	8.60	-1.36	21.06	1.72	1.31	-1.01	-8.25
miR-375	32	16.2	-1.60	-1.60	-1.60	-1.60	-1.60	-1.60	-1.60	1.32	-1.60	-1.60	10.44	12.39	58.80
miR-376c	16	62.7	-1.12	-1.12	-1.12	-1.12	-1.12	-1.12	-1.12	2.50	-1.12	-1.12	15.41	-1.12	170.06
miR-429	16	8.1	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	3.48	1.76	17.65
miR-483-5p	42	10.1	1.50	-1.21	-3.38	1.14	2.56	9.82	7.11	1.75	52.02	2.12	1.94	1.66	-3.38
miR-516a-5p	47	8.8	6.97	-1.65	-1.65	1.17	-1.65	9.91	4.22	2.90	39.38	5.18	4.21	3.86	-1.65
miR-7	21	86.3	-1.00	-1.00	-1.00	4.57	-1.00	-1.00	-1.00	-1.00	-1.00	-1.00	11.46	7.18	321.94

Table 22: Chromosomal locations and pre-microRNA sequences for target RNAs in Tables 18 to 21.

Gene	Chr location	Pre-microRNA sequence	SEQ ID NO
10083-L5-1	14q12	GGGTGAGAGAGGGGTAGAAAGGAAGAGGGACATATGGAGCCTCTTCCCCCATGCCCGAAAGCTTCTCCCATTT	1211
10233-R5-1	9q34.13	ACCTTCTGCCCTTTGAGACCTGGAAGCCACAGCACTGGTGACATAAATTTCTACTGCTGCTGCCAGATCCAAGGGAGAGGGT	1212
10335-L5-2	15q26.1	TCTTAGTGCTTTCTAATTAAAGAGGGGAGTGGTGATCTTTTGGCTCTCTAAGTTCTGTTCCCTCTGAGTGGAAAGCAGAGGG	1213
10357-R5-1	3p25.3	CTCTCTTTGCTGAACCTCTGTGCCAGGGTTCTCTAGCCCTCTAGGGAAACAACACATGGAATATTCAGACAGTGGAGAG	1214
10455-L5-1	17q23.2	GGAGAGGGATGTTTGGCGCAATTGCAGTTGTAAATAGAACAGGCTTCTGAATTGTGGCCAGAACATAATGAATCCCATTTCT	1215
10520-L5-1	3p23	AGTCCACAGAGAAGGCTGGAGGAGGTAGCAAGGAGATGCTGTAAACAGCTGTACAGCCTTAAACAAAAGTTGGTGCTCTCTGGACT	1216
10844-R4-1	8p12	GGGTGGGTGGCAGATGCTTACTCTGCCCTTTAGAAACACATGTTTGAAAGTAGAGTCCCTGAATATGCTACCCCTT	1217
11358-R5-2	1q23.3	GTGAGAGGGTAAAGTAGTATGCTGTGGGTGAGAACTACTAGGGAAGTTCTGAGCCATGTGAGCCTAGTTTCCCACCCCTCCAT	1218
11444-L5-3	15q22.2	GAATAGTTGGAAGGTACCTTTTGCTGGAGGAGTGGTGCCCTGAACACAAAGTGCAGTCTCTATTAGTAAGCCAGGACCTTCAGCTCCTCA GTTGAAGGGAAATAACTATTC	1219
11547-R4-1	6p12.2	TGCCAGCCCTGCTGTAGCTGGTTGAAGGGGACCACCAACCTCTCAGGAAGACGGACTTGGTTCCGTTTGACCAGCCAGCAGGGGCA	1220

11605-L5-4	1p31.3	GGCCCTGTGCATAATAATCTTTATGGAATTGAGGGGAAGGGAATTAAAGAAAGGGAAGAGCAAAACCCACTACAGAGTTTATGA CCATCTATTCTTAATATATATTAGAACTGGGCC	1221
11688-R5-3	11q13.4	CGCCCACTAAACCGGATGTGACGTTGACCTACCTTAGTCACATTGTTAGGGAAGGAAGTGTGCCGCGCCTACCTATCTGCCCCGCGCTT GAGTCTCAGCCAGTCGGCGTCTCCATCCTGGCG	1222
12699-R5-2	1q22	CTGGACGGCGGGCGCGGAGGCCAGGGCGCCTGAGGGCGCGAGAGGTGTCAGCGTGCAAACCGCGCCCGCCAGCGTTCCCG CCACCACCGCCACCACCCCTCAAAGCCCGG	1223
12707-L5-2	7q32.1,17q2 3.2	GATTGTAGAGTTGTTGGATTGCAGAGGAGAATGTTTCGATTCACTTTTCAGTAACTTTAAAAATC	1224
12729-R5-1	17q25.3	AAATTAAAGAAAAAAATTTCTCCACCAAAAGCGCGCAGTGACAGTTCCTCAACATTTTCTGCCTTTCTTCTCCCTCCCTGCACCACCTA GGAGGAGAAGGCACACAATTGCAATTTTCGTGCTTTTAAATTTTTTTTTTTGGTTT	1225
12730-R5-1	17q25.3	CCGGCTCGGCCCGCGCTCTCCAGCTCCTCCGGCTCCTTTTAGTGCAATAAATTAGTGATGGCAATTTCCCGGAGAGCGGAGCACAA CACAGGCGCGGGCTCGG	1226
12888-L5-2	2q32.3	TCTCCAGGAGCTGGGAGAACCATACAATGTAGGAAAAATATAGTTTAATTGAATGGTACTCTGGTCTTCTGGAGGA	1227
12907-L5-1	17q12	GGCTCAGTGTCTAGGACGTGGTAGAGGGCATGAGCACAGTGGAAGCAAAAATGCCAGTTCAGTGCTTAGTGCTGACTTTTACCACC GTTCTTCATGGCAACTGAAC	1228
12912-L5-2	1p36.33	CCACCTACAGCCTCTGTGGGTGGAAGGTGCCAGAAAACTTGAAGAGTGGCTCTGGCCAGCTCTCTGGGCCAGTTGGCACCCAGGTG GTTGCAGAGAAAGGTGGG	1229
12917-R5-2	1p34.1	GGACCTGGGGCTTCTCTGACCCCTTGACAGCTTATACTATGAGACCTTGGAAACCTCCTCCATGCAGACACACAAGGCTCAATGTTG GGGAAGGGAAGGCCCATAGGTCC	1230
12947-L5-4	6p21.1	GCCTTGCTGAAGGGAGAGGCCCTGGCATCGGATGGGAGATTTAGAGGCTGTGGAGAAGGGAACCTTGGGGCTTTCCCTTCTCGTG GCCTCACTCCCTGGGCCCTCTCTCTATGGAGGGGC	1231
12958-R5-1	6q23.3	GTTAGGGCCTGGAATCCCTCTGTGCGGCATCTGTTAGAGTCTGTGGATGGCACTCTGTAGTCTGTTATAGATGCACCCAGGGGAAGATT CTCTCAGGCATGAT	1232
12974-R5-2	Xq27.3	ATTCTCTGCATGTTTACCTTGTTCTCTCTGTGTTTGGTGTTAAAAGTTAAATAACTCTGAACCAAGGGACTAGGAATTTGGGGGCACA GAGGGT	1233
12979-R5-2	3p21.1	TTTGGGTCCTTTGTCCAGAGTGTGAGGATGTTGGTGAAGGCCATGCCCTGACTAGATTCAAGGATCCCTTTCTTTCATCTAGCTACTGG GGAAGACCTGGA	1234
12992-L5-1	9q31.3	TAGAGATGGGGTGGAGGAAGCAGGTATGATTTACAGGGCAGCAGGAGATTCTCTTCTGTTTCCCTCTCTCCCAGCTCTA	1235
13001-L5-1	12q13.13	TGGAGCTGGGAGTGGTATCTAGGAATTTCTCCTTCTAGTTTTTACCTTCTTAGTTTTA	1236
13053-R5-4	19q13.33	ACCCATCTGTAAAGTGGGACAATCTTCTTGCTCCTAGGATGGCCATGGAGCTCAGTGCAGGGGTTGGGGCTGCCTGATGAGCCTGA TCTTAGGGGAAGGC	1237
13070-R5-3	5q35.1	CTGCTCTCCTTGACCTGTAGGAAGGTGATTTGATGGTTTCAAAACACATGCTGTTTCTCTCTTCATCAGGGTAGCTGGGTGGGTGGC ATTAGGTCTTTTGACTGGGAGAGAGACAG	1238
13071-L5-3	7p22.3	TACATGTCCCGAGATGTGAGCAGTCACCCCTGGAGATCCTCACCCCTTCTCTGTAGAACGAGGAGTTGGGCTGCCAGGATTGGGGT CTGGCGCGTTGCTGGTGTGTG	1239
13078-L5-1	2p13.1	TCTCCTGTCTAGCAGGCTTATGAAATGAGGGCCCATCCTTTGGTCAAGGCTTGTGTAAGGAGA	1240
13098-L5-1	11q24.1	ACCCCCGGAGGAACCGCCAACAGCGCCCCGGCACCATCAAGCATGGATCGGCGCTGGACGTGCTCTCTCCATGGGGT	1241
13122-L5-1	2p11.2	GGA CTGGAGCAGGTGAGATGGAATTTCTAAAGGTCCAGATATTAGGACCTGGACCCATCTCACCCGCTGCCCTCTGTCC	1242
13185-L5-3	3q13.31	GGGAAGAAAGAGCAAAAGGGGAGAGAGGGAAGAAAAACAGAAATAAGTTGGTCGTATTGAAGCTTCTCCAATCAACCCCTAGAACAA TTAGCTTCCACC	1243
13216-R5-2	1p21.1	TGAGTAAATAATCTTCAAATGTATGGCTTTTATTCTCTGGTGAGAGCATCATCTGAGAGAAACAGTGTCTCAGCTGCCAGAGAGGAGA AAGGAGGAATATATAGAAACATCCAAACTT	1244
13219-L5-1	11q22.1	GCTGGAGCAGAGAGACTGAGTGAGGGAGTAAAGAGAAATTAGTACAGGTGAGATTGTACTGATTATCTTAACTCTCTGACC CCCTCACTCAGTAAAGATCAGATTGTGCCAGGC	1245

13225-L5-1	11q23.3	GAAGATGGAAGGGGCTGTCCCAGAAAGTGGGCAGGAACACGAGCTGTTATGCCATGTGGGCATGTGGCACAACAGCCTCGTTCTTC	1246
13235-R5-1	11p11.2	CTGCCACCTTCTGGGGCACCTTGTTCCCTCT	1247
13245-L5-4	1q24.1	CTTCTGCTTGGCTGTTATATGAACCTGCCCTGGCTTCTAGTCTCAGCTCTCCTGACCAGCTGAGCTGGAGGAGAGCTGAGACTAG AAAGCCCAGGCAGGTCCAAC TGAAAA	1248
13252-L5-3	1q25.2	AAGACATTACTTTGAATATATTTAAAGGGAGTAGAAGGGAAGACTATTTGGAATGCTTTATTAAAAATAGTCTTTCCCTTCTACTCCTTT TAAATATATTCAAAGAAAGTT	1249
13254-R5-1	1q25.3	CTTTGGCACAGTCCGTGCTCTCAAGGAGCTCACAGTCAAGAACGACGTGGAATTTACGCCCTGGAGTTCCAAGTGTGCCCTCAGGG AGTGCTGGCCCTGAGCTGGGTGAGGCTGCAGGG	1250
13274-L5-3	12q13.13	CTCACACATGGTACGTTTCAATGAGCTGATTTGTCTTCTCCACTCAATGCAGTAATTGAGCTTCTTTGGTTCAAGTGATGAGTGGTTCA GTGGTTCAATTGGGCATCCTGTTGAGGG	1251
13300-L5-1	14q24.2	AGGTGGTGGTGGGAGACCTGAGGGAGGGTGGGAGCACGGGAGAGGAGATACCCAACCTGACCTACTTACCTGTCCCTT ACCCACAGAGGGCTTCCCTGGAGGCCGCAATTGC	1252
13309-R5-1	1p32.2	GGTAAAGTAGGAGGGGAGATAATTCTGTGTGTAACCTACAACCCCTTCTGCCCTTTATCTCTGGGGCCCCGAATTTCCATGTGGGGGAT AAAAGAAAGTCACCCCTCCTCATCCCCC	1253
13352-R5-1	19q12	ATAGATTTACACTCCCTGTCTCCACAAGCTCACAGAAAAAATTGTAGCTGAAGACAAGGGGGAAGAACCTA	1254
13357-L5-4	19q13.2	CACTTTTTATGACATCACATTGTTGACCTCAATCTGAGGAGATGGGGAGCTAGCCCCCATTTACACAGATGCAGCGCTGAGGTTCCGGG ATGTGAGGTGAGGTGA	1255
13366-R5-3	19p13.2	TCAGGGTTTGGTGACAAATTGTGGAGGCTGCAGGGAGGTGTGGGAGAGGTCGCTTTAGAGCTTGGTCCCTCTGTGTCAACCTC CCAGCCCCAAAGCCCCAGCACCGATCCTG	1256
13373-R5-2	20q13.33	GGCCGGTGAAGGCCCGCCTGGGTCCCATAACCCGGGTTGGGGTCAAGAGCCGCTCGGTCTCTGTGGGAATAGGAGAGGGCAC TGGGGTGAGGCCTG	1257
13375-R5-4	20q13.33	TCCCTGCACCCCTGCCTGTACAGGTGAGTGGGAGCCCGTGGGGCTGGAGTAAGGGGCACGCCGGGGCTGCCCCACCTGCTGACCA CCCTCCCCCACAGCACCCCTGTGCCGGGC	1258
13398-R5-4	2q35	ACCCTCTCAGGACCCCTCCTAAGGGGTAGGCAGGGGCTGGGGTTTCAGGTTCTCAGTCAGAAACCTTGCCCCCTCTCCCCAGACCCCC AGGCTGTGGTGAGGCTCTGAGAGCTGGTAC	1259
13417-R5-4	3q13.2	ATCCTCAATTCAATTCACAGGAGACCTACCGCAGAAAGAGACCCCTAGAACAATCCTTGGTTCAAAGTGAGTCTAGTCTGCAAAAGTGGTGGA CAAAGGTGGAGGAGAGAGAA	1260
13436-L5-1	3p12.1	CCATGATTCCCTTAAACAACATGTTTCCAGCATTTCCAGGTAGGCCAAGGTGTCTCTACAGAAAAAACCTTGGGTTAGACCTACAGGGGATC TGGCTGTTGTGTTAACAGAAAGGAAGGCA	1261
13467-L5-1	5q35.2	GAGGCATGACCTAGCTGGGGCTTCATTGAGCTGTGGTCAACCGGAGCTCAGAGGCTGCTCCTCCTAGGCACCCCTAGCTCTTAGGACT CAGTTCTCTGAAGCTCACCAAGGGCATCTATCTT	1262
13468-L5-1	5q35.2	TTTGACTTGGAGCAAGGAGGGTCTGACTGTCACTTGGAGCTAAACCAGTCTCCAAGTGGCCATCAGACCCCTCTTTGCCCAAGTCAGT GACTTGGGGCAAAGAGGGTCTGATGCCACTTGGAGACTGGTTTAGCTCCAAAGTGACAGTCAGACCCCTCCTTGCTCCCAAGTCAA	1263
13470-R5-1	5q13.2	GGTCAGTCTAGCTTGCTCTTGAAGCTCTCCAGAGACAAAGAACGCTGTAGCAAAATCAAAATGAGGATTCTGGTCCCTGAGGTGGAGAG GGTGGAGCCCCACTGTTTGCCC	1264
13473-L5-3	5q13.3	TATATAGAGAGACATATGTATCAGGATAGGTGGTAGAAGGGCAACACATAGAAAACACACCCTAGCCAGGTTGCCCTGAATGGCCCCA GTGCTAGCATAGTCTTTATG	1265
13500-L5-3	7q32.3	CTGGTGTGGACCTCACTCAATGGAGAGAGAGGCCCTGGAAGATCTTGGAACAGGAACATCCAGCCCTGCTTCTCTTAATCTCAGCCCC ACACTCAGAG	1266
13522-L5-4	8p21.1	AGTGCAGTGGTGTGATCATAGCCCAATAGAGGGTGCACAGGCACGGGAGCTCAGGTAGGCGAGGAGCTGAGCTCACCTGACCTCCC ATGCCTGTGCACCCCTCTATTATACACTCGTAT	1267
13523-R5-1	8p12	GGCCATAGCTGCTGGGTCCCTAAGGCACCTCTCATCAGTTAGGTCAAATGTGCCAGGCAGGGGAAGAACCTTACTATTCCACTGCATC TGTGGCCT	1268
13545-L5-1	Xq28	CCCCAGACTCACAGCCTCCCTTGGCGCCACAGGCCCTGTCTCCTGACTGGGACTCTTGGGACGCCAGCAAGCCCCGTTACCAATGCGCGA GAGGCCATGCAGCAGCGCGATGACTGGCTGGCA	1269

227-L5-1	3q27.2	TGAGGGGCGGAAGCACTGGGGAGAGACAGGTTGTGAGCTTCCCACGTGGTGATCAGCTCACACCTGTCTTGTGTTCTTGGTATTCACA GACTCTCA	1270
3744-R5-1	19p13.12	CTTCTCTATTCTCCCTGTTTTTCATCCTACTCTTTTAAGTAATAAAATTTGGCATTAGTGGAGGGAGCAGGGAGGAAG	1271
3875-R5-2	5p15.1	GGCTCGGTTTCAAATCTCTCCTAATCCACTAATGAACCTTTATTAAAGTGGGAGAGAGAGTTGAATCAGTC	1272
3923-R5-1	19p12	GGCTCTGCACACAGCGTTTCTTCTTGTTCCCTCTCTCTCTGAGAGGGATGAAGGAGATCCTTTTGTGAGAGGC	1273
3926-L5-1	3q27.3	AGTGGGGGCACGTTCAAGAATGCTAATGAATTATCAGTTTCTTTAAGCCTCATTAGATTTCTTGATAGCAAAACCTCCCATTT	1274
3992-R5-1	2p22.3	GTCCTGCGTCTCTGTTCTTATGCTGCTGTTGATTTTTTCCAACCATCAGCTCAGATACTGACAGAAGAGAAAGGGGC	1275
4203-R3-2	11q23.3	AGCAATTCCAACTGCCCCATTATATTCCTAAGTAGAGGACTTGTTAATATGGAGCCTGCCTCTGGGGAAGTGGGAATGTGCT	1276
4261-R5-1	1q23.3	GTAAATTTCAACCTATTTTTAAGGGTTATTTTCACTCAAGTGAAATCTATCAAGGAAGGAGGGTTATTTTTAC	1277
4291-R5-1	3p14.3	GGTTAAAATGGATTGCCAGGAACACCTGAAGGGTACAAAGTCATGTACACCCCCATGGCTCCTGGTAACCTACCTGATC	1278
4440-L3-2*	11q14.1	CATGTGATTTCTGCCCAGTCTCTGAATGTCAAAGTGAAGAAATTCATGAAGCACGGGTAAACGGCGGAGTAACTATG	1279
4440-L3-2*	7q11.22	GTGATGTGATTTCTGCCCAGTCTCTGAATGTCAAACCTGAAGAAATTCAGTGAATGCGGGTAAACGGCGGAGTAACTATGAC	1280
4479-R3-1	1p32.3	CCCAAGCTCCTTCCTGGAGGACTTAACACTGTGTGAGCAGTTGTTTTGAATTTCCGGCAGGGGCTGCAAAAGGG	1281
4790-L5-2	2q11.2, 2q14.1	CCCAGAAATTCAGTAATGAGGGAGACAGGTTATGCCAAGCCTGCTTCTCCCAGGATGCACTGGGAGCCTGGG	1282
4958-L5-1	6p21.1	CCCCATTTTCAGAAGATGCCATGGTTTATTAATTTCTCTGCAGAAGCAATAAGAACTGGGTATTGGCTTAATTGGGG	1283
4988-R5-2	14q24.3	CTTTTCTCTCTGCTGGGAAACCTTGCTTGACTTCATGTCCAGTGTGGTATCCAAAGACGGGAGGAGGAG	1284
5080-R3-1	11q23.3	GGCGTTTCTTCTTGTTCTCTCTTCTCTCTGAGAGGGATGAAGGAGACCCCTTTCGAAGAGGCATGTT	1285
5108-R5-2	2p13.1	CCAATGCGCTCTCACCTCTCACTTGTGGCACCTTCGCTTTTGATCCGAAGTGCGGGTATTGTGAGTGACGGTGTGGGAGGAGAGG G	1286
5232-L5-2	9q21.13,1q2 3.2	GGAGCCCTTGATGTCTGCAAAATGAAGGAGGAGGATGTCTTAAGTTCCCTTGCAAGCAGGAACCCACTTAGGTGGCACC	1287
5392-R5-1	2q24.2	CTCTCTCTCAGTTACTCACAAAACATGGCTGCTTATTCAGAGATTAGCAATTATTGTAATGAGATACTGTCTGGAGAGGG	1288
5723-R5-1	4p15.31	CCTCTGCCCTGGCTTTCTTTGTAAAGCCATTAAACTACATTAAGAAGGCTACTGCTGGAGAAAGGGAGGAGG	1289
5836-L5-1	11q23.2	GCCATGGGCGCTCCATAGTTTCTCTGTAGCCCCCTTGTTGCCAAGAAATAGTTTGGAAATGGGGCGTGTGTGATAATGGGGTTAATGG T	1290
5842-R5-1	1p33	GGCCACTTGGGTGCAATGGACTTGAACCTCTTTTTTGTCTCCCCCTTTAGGGGGATATAGATTTTCATTTCTCTTCATAACGGGCC	1291
6037-R3-2	16q21	GCAGGATCCCCTCTTTTCATCTGAAAATTACCACATAATTGCAATTAGTTGGAGGAAAATTGGAGATGGAGAAAGGGAATTGC	1292
6181-L5-1	1q41.5q23.1	AAATCCAGGGGGAAGAGTCATCTAAGTTACCATGCAGTTGTTACCAAAATAGAGGAGGAGAGTCTTAACCTTTTGTCTCTGGATTT	1293
6216-L1-1	11q14.1	CATGTGATTTCTGCCCAGTCTCTGAATGTCAAAGTGAAGAAATTCATGAAGCACGGGTAAACGGCGGAGTAACTATG	1294
6233-L5-2	6q16.2	GGAATGGGAGAAGGATAAAGTGGAATCTAAATTTTGAGAAAATAAGGATTAAAGGTTCCATTATTTCATGCTGTTTTTC	1295
6235-R5-2	15q26.2	TCTGTTTTTATCAGTTTAAATATATGATACATCTTCTATCCAAGGACAATATATTAATGGATTTTTTGAGGCAGA	1296
6395-L5-1	14q11.2	GTTCCGAGGCAGGCTTTCCTCCTCTCTGCAGGGGAGAGGCTCCCTCACACAAGAGGAGGATTACACTGGCTCTGAGC	1297
6405-R5-1	8q21.11	TCAAATGAGAAACTAAGTACATTTCTTTCAACAGGCAATTTACCCCATGATGAGATATGCTTTCACTTATTTTCTTTGA	1298
6474-L5-1	1p34.1	GGTGAAAGGGGGAAGAGGGGATGGCTTTAATTAGGGCTTCTTGCCCTCGTCAGTCTCGGGGTTGGTCGGGTAATCCAGTTAGAGTCC CAGCCTTTCATCTCAGCTGGCC	1299
6602-R3-1	3p14.1	CAGAGTCTAAATGGAAGAGTCCCTCCGTAATTTACCCAGCTCATCTCCTGTGTAATGGATTTGGAGGAGAGATTTCCACTGGGCTCTG	1300
6681-R2-1	11q12.2	TGTGCTCTCATTTGTTATTCCAAAGTCTCTGTCTAGATCACTGGAGGGGCAGAGAGAGAGGGGAGAAAACAGGGATACA	1301
6683-R5-1	12q23.2	GTTCTAGTCCAGGATGCTGATACTTTAAGCCCCGAGGCTCTAACTTGAGCAGGAAGAGTTTATTTTGGGATGAAGAAAT	1302
6752-R5-2	Xq13.1	CCCTCCCAGTTCCCATAGCAACTGGGCTGTAGCAGCCAGCAACTTGATTGAGCCCCAGCAGTGCCCCGACTGAGGTGGGAAAGGAGG G	1303
6803-R5-2	22q12.3	GCCACCTTTCATGGTGAGGATGCCCTGCCACCTTCAGGATCACATCTTTGGGTGAGGTCCAACCAGAGAGGGAGC	1304
6864-R4-1	3q26.2	ATCTGTGTTTGGCTACAGGTGGGCACCTTAAGGGGTCAAGTCTCCAGGGGAGTGTAATTGTGTCTCCTGGGCTTGTTCGAGAATG ACCTAAAGCTGAAGCCCCAGCT	1305

172

miR-182	7q32.2	GAGCTGCTTGCCCTCCCCCGTTTTTGGCAATGGTAGAACTCACACTGGTGAGGTAAACAGGATCCGGTGGTTCTAGACTTGCCAACTAT GGGCGAGACTCAGCCGGCAC	1341
miR-183	7q32.2	CCGACAGGTGTGACTCCTGTTCTGTGTATGGCACTGGTAGAATTCACCTGTGAACAGTCTCAGTCAGTGAAATTACCGAAGGGCCATAAA CAGAGCAGAGACAGATCCACGA	1342
miR-198	3q13.33	TCATTGGTCCAGAGGGGAGATAGGTTCCCTGTGATTTTTCCCTTCTCTATAGAATAATGA	1343
miR-19a	13q31.3	GCAGTCCCTCTGTTAGTTTTGCATAGTTGCACATAAAGAAGATGTAGTTGTGCAAAATCTATGCAAAACTGATGGTGGCCCTGC	1344
miR-200a	1p36.33	CCGGGCCCCCTGTGAGCATCTTACCGGACAGTGCTGGATTTCCAGCTTGACTCTAACACTGTCTGGTAACGATGTTCAAAGGTGACCCC GC	1345
miR-200b	1p36.33	CCAGCTCGGGCAGCCGTGGCCATCTTACTGGGCAGCATTTGGATGGAGTCAGGTCTCTAACTACTGCCTGGTAATGATGACGGCGGAGC CCTGCACG	1346
miR-200c	12p13.31	CCCTCGTCTTACCCAGCAGTGTTTGGGTGCGGTTGGAGTCTCTAATACTGCCGGTAATGATGGAGG	1347
miR-205	1q32.2	AAAGATCCCTCAGACAATCCATGTGCTTCTCTGTCCCTTCATTCCACCGGAGTCTGTCTCATACCCCAACCAGATTTTCAGTGGAGTGAAAGT TCAGGAGGCATGGAGCTGACA	1348
miR-20b	Xq26.2	AGTACCAAAGTGCTCATAGTGCAGGTAGTTTTGGCATGACTCTACTGTAGTATGGGCACITCCAGTACT	1349
miR-21	17q23.1	TGTCGGGTAGCTTATCAGACTGATGTTGACTGTTGAATCTCATGGCAACACCAGTCGATGGGCTGTCTGACA	1350
miR-298	20q13.32	TCAGGTCTTCAGCAGAGCAGGAGGTTCTCCAGTGGTTTCCTTGACTGTGAGGAACTAGCCTGCTGCTTTTGCTCAGGAGTGAGCT ACCATGCTGTAGTGTGTAAACATCCTACACTCTCAGCTGTGAGCTCAAGGTGGCTGGAGAGGGTTGTTTACTCCTTCTGCCATGG	1351
miR-30c-1*	1p34.2	A	1352
miR-375	2q35	CCCCGCGACGAGCCCCCTCGCACAAACCCGGACCTGAGCGTTTTGTTCCGTCGCGTGAGGC	1353
miR-376c	14q32.31	AAAAGGTGGATATTCCTTCTATGTTTATGTTATTTATGTTAAACATAGAGGAAATTCACCGTTTT	1354
miR-429	1p36.33	CGCCGGCCGATGGCGTCTTACCAGACATGGTTAGACCTGGCCCTCTGTCTAATACTGTCTGTTAAACCCGTCCATCCGCTGC	1355
miR-483-5p	11p15.5	GAGGGGGAAGACGGGAGGAAAGAGGAGTGGTTCCATCACGCCCTCCTCACCTCCTCCTCCGCTTCTCCTCTC	1356
miR-516a-5p	19q13.41	TCTCAGGTTGTGACCTTCTCGAGGAAAGAACGACATTTCTGTTGCTGAAAGAAAGAAAGTGCTTCCCTTCAGAGGGTTACGGTTTGAG A	1357
miR-516a-5p	19q13.41	TCTCAGGCTGTGACCTTCTCGAGGAAAGAACGACATTTCTGTTGCTGAAAGAAAGAAAGTGCTTCCCTTCAGAGGGTTACGGTTTGAG A	1358
miR-7	19p13.3	AGATTAGAGTGGCTGTGGTCTAGTGTGTGGAAGACTAGTGATTTTGTGTTCTGTACTACGACAACAAGTCACAGCCCGGCCCTC ATAGCGCAGACTCCCTTCGAC	1359
miR-7	9q21.32	TTGGATGTTGGCCCTAGTTCTGTGTGGAAGACTAGTGATTTTGTGTTTTTAGATAACTAAATCGACAACAAATCACAGTCTGCCATATGG CACAGGCCATGCCTCTACAG	1360
miR-7	15q26.1	CTGGATACAGAGTGGACCCGGCTGGCCCCCATCTGGAAGACTAGTGATTTTGTGTTGTTACTGCGCTCAACAACAAATCCCCAGTCTAC CTAATGGTGGCAGCCATCGCA	1361
miR-720	3q26.1	CCGGATCTCACACGGTGGTGTGTTAATATCTCGCTGGGGCCCTCCAAAATGTTGTGCCCAGGGGTGTAGAGAAACACACACTTTGAGA TGAATTAAAGAGTCCTTTATTAG	1362

Table 23: Target RNAs present at at least 5-fold decreased levels in at least 50% of tumor samples

Gene	Probe sequence	SEQ ID NO	% of down-expressed	ADK9	ADK10	Adk29	Adk15	Adk23	ADK48
10010_B-L4-1	GCCGAGCCCCCGCCCCCGGGGATGCTGCCCTCCGGAAGGAGG	1363	67	-4.54	-4.10	-6.89	-5.28	-3.68	-2.54
10010_D-L4-1	TGGCGCCCTCCCCCGCCGGGGCTCAGCCTCTCACCTG	1364	61	-4.69	-6.39	-10.13	-3.34	-2.75	-1.75
10010-R2-2	CTCGCCGGCTCCAAACTTTCCCCAACTCCAGG	1365	67	-10.92	-10.92	-7.27	-7.12	-4.44	-2.95
10030-R5-1	CCGTGGATGTCAACTCAGCTGCC TTCGGCC	1366	67	-6.81	-6.81	-6.81	-2.13	-3.82	-2.13
10145-L5-2	TCAAAAAATCTTTTTCCTTTGCTCTCTCTTCTGT	1367	56	-2.03	-2.03	-2.03	-2.03	-1.18	-2.03
10231-R3-1	TGAACTTTAGCTGGGCGCGCGCTGTCAGC	1368	72	-6.47	-4.83	-7.60	-4.34	-5.39	-1.70
10260-L5-2	ACCTATTGTTCCGAGGGCCCCCACCCGATGT	1369	67	-13.31	-13.31	-3.91	-3.96	-6.75	-4.01
10333-L5-1	TGCCCTGCCACCCCCCTCCCCTGCCCG	1370	78	-5.25	-5.14	-8.87	-5.02	-3.91	-2.93
10342-R2-2	CCCGCGCGGAGCATCTCGAAGTTAATTAAA	1371	56	-4.63	-3.69	-4.92	-3.66	-4.90	-1.74
10345-R5-1	ACCCTTCTCTCAAAAGCGACCCCTCCCATCCAGTCC	1372	72	-4.22	-4.59	-10.15	-5.45	-4.10	-2.11
10374-R3-2	GACACGCGCGCTACTTTGTTAATGAAAAGCCCCC	1373	67	-33.82	-4.71	-10.79	-4.92	-11.40	-1.26
10435-R5-1	TGCTAATTGTGCCCTGTTGTCTTTCTTAAACT	1374	56	-2.03	-2.03	-2.03	1.07	-1.10	-2.03
10533-R5-2	TAACTGTCTGAGCCCCCTTCTTTATATTGC	1375	56	-2.60	-2.60	-2.60	-1.21	-1.46	-2.60
10543-R5-2	GCCTTACCCCTTCCCATCTCTGGCTCATGAAT	1376	56	-2.16	-2.16	-2.16	-2.16	-1.24	-2.16
10578-R5-1	CCACCCCTCTACGGTCCCCACCAGCCCCG	1377	78	-5.22	-4.46	-6.46	-4.96	-4.51	-1.76
10818-L5-1	CTCAGTGATGACAAATGACCGCTGTCAGCCGCC	1378	83	-21.17	-21.17	-8.42	-4.40	-9.89	-1.24
11370-L5-5	CCCTCGCCCCCACCACACTGCATCCTTGCCCCAGTTTGG	1379	78	-6.80	-4.03	-7.05	-3.30	-2.15	-2.30
11605-L5-4	TCCCTTCTTTAATTCCTTCCCCTCAATTCATAA	1380	61	-2.54	-2.54	-2.54	-2.54	-1.37	-2.54
12184-L4-1	GACCTCAGCGTGCCCCCTTTCAACCACAGACGAATATTGTGTACAA	1381	56	-2.73	-2.21	-3.89	-3.08	-1.99	1.08
12184-L5-3	TGACCTCAGCGTGCCCCCTTTCAACCACAGACG	1382	61	-5.57	-5.57	-5.57	-5.57	-2.34	-1.75
12224-L4-1	CAAGGCGTCGCCATGAAAATCCACCCCATGGAGTAGCA	1383	78	-3.22	-3.22	-3.22	-3.22	-2.54	-3.22
12361-R5-1	ATGTAAATGTGCCATATTGCCGCCCATCT	1384	67	-3.34	-4.62	-6.42	-3.65	-3.60	-1.50
12691-R5-1	CCCCGCCCTGGCGCGCCCCCGACAGGC	1385	78	-67.73	-7.29	-9.93	-7.32	-3.57	-2.81
12692-L5-1	GACCCGGCCCCCGCAGCCAGCACCCGCCACCGCGC	1386	89	-7.80	-7.12	-6.21	-7.94	-5.87	-2.09
12693-L5-1	GTCGCGGCCCGCCGGCCCTCCCGGTCCCCTCCCC	1387	61	-84.98	-6.22	-8.38	-2.30	-2.98	-1.86
12694-R5-1	TCAGCCCCCAGCGCCCCCGGAGTTCITGGA	1388	78	-6.01	-5.88	-9.37	-6.86	-6.32	-2.41
12696-R5-2	AACCCGGGCTCCCCCACCCTCCCTGA	1389	61	-4.33	-3.38	-7.14	-5.25	-4.53	-1.41
12697-R5-1	CCGGTGTGCGCCCCCTCCTACCTCTGCCGGCC	1390	67	-4.28	-5.08	-27.65	-4.83	-3.72	-1.76

12699-L5-1	GGGCGCCTGGGCTCGGGCGCCCGCGCCCGTCCCAG	1391	50	-4.46	-29.74	-4.16	-3.65	-2.69	-1.18
12701-L5-1	AAATCCTGCGCATCCTCCACCCCCCAGCCCCGG	1392	78	-4.43	-12.74	-7.02	-6.42	-4.98	-1.97
12703-L5-3	AGCCGAGCCCCCGCCCCCGCCCGGATGCTGCCCTC	1393	61	-4.17	-3.97	-5.50	-4.94	-3.05	-1.64
12704-L5-2	CTGGCGCCCTCCCCCGCCCGGGCTCAGCCTC	1394	72	-4.44	-5.86	-10.21	-3.67	-2.76	-2.08
12713-R5-1	CTCTCGCGACCGACCTGCGCGCCGACCGCCACAG	1395	83	-7.39	-6.52	-7.89	-5.40	-7.13	-1.82
12722-L5-1	AAACAAAGTACTTCGACCTCCCGCGCCCGCCCGC	1396	56	-4.21	-4.38	-5.37	-2.81	-2.86	-2.02
12723-R5-2	CCCCAGCCCTTCCTGGCCAGGACCCCGAGG	1397	72	-18.13	-18.13	-18.13	-3.62	-3.52	-4.66
12725-R5-1	CCACCCCTCAAGCCCCCAGGAGCTTCCTTAAC	1398	67	-22.87	-22.87	-22.87	-4.18	-9.94	-1.79
12731-L5-1	CAGCCGCGCGCGCCCGCCAGCGCCCCGGCGCG	1399	72	-14.83	-14.83	-14.83	-4.58	-3.87	1.02
12900-R5-3	GTGCCCCGTGCCCTAAGCGACCCCTCCACTCCCTCAA	1400	56	-8.25	-1.80	-8.25	-1.90	-3.08	-2.33
12904-R5-2	CTGCCTCCCTGACCCCAAAACACACCTGACTGCC	1401	61	-2.67	-4.87	-4.22	-2.80	-2.26	-9.55
12910-R5-2	GCTCCATTTCCTCCACTGAAATGACAAATCCA	1402	61	-4.61	-4.61	-4.61	-4.61	-4.61	-1.30
12925-L5-3	GAACACCTCCTCCTTTTCCCTTTGGTGAATAAGA	1403	61	-2.02	-2.02	-2.02	-2.02	-1.68	-2.02
12932-L5-3	GACTCCCTTCACCCCCCATTTCCACCTACTAAT	1404	67	-8.22	-8.22	-8.22	-3.88	-4.15	-8.22
12939-L5-2	CTGATTCAAAACCCACCCTCTACAACTTAATAGCT	1405	72	-2.19	-2.19	-2.19	-2.19	-2.19	-2.19
12947-R5-3	GCCCCCTCCATAGAGAGAGGGCCCCAGGGGAGTGA	1406	72	-28.53	-28.53	-28.53	-5.22	-4.41	-2.39
12975-L5-1	GGAGTCTACAACATCCCTCCCGCCT	1407	50	-2.29	-3.08	-4.02	-2.31	-2.47	-1.53
12981-L5-1	AATGATGTCACCTTGGTCTGTTGCCCCACCCC	1408	61	-4.08	-4.08	-4.08	-4.08	-2.35	-1.27
12981-R5-1	GGGTGAAGACAGACAGAACCCAGACCCCTCCCATAG	1409	50	-2.22	-9.73	-9.73	-2.36	-3.84	1.96
12998-R5-1	CAGGCCCCAGTGCCACCCGCCAAGGCTATC	1410	89	-12.85	-12.85	-12.85	-12.85	-10.71	-3.87
13004-R5-1	CTCCAACCCCCGCAATTCTCGCTCCCTTCACCTGA	1411	56	-2.79	-4.11	-6.82	-4.51	-3.96	-2.17
13047-R5-2	GCACAGCAGACCCCATGCACTAGCCCCGGGCAC	1412	50	-2.69	-2.69	-2.69	1.15	-1.55	1.18
13050-R5-4	ACGCACCCCTATCTCCACCCCGCCGCAAGCCAAGCA	1413	78	-4.94	-6.60	-11.56	-5.96	-5.41	-2.49
13052-L5-1	TCCCCGGACCTAAGCATCTCCCCACCCCGCCAACC	1414	61	-2.01	-3.27	-4.41	-3.51	-3.73	-1.83
13066-R5-2	GGAGGTAGCCCCCAAAATCTAGTTGAACCAATTC	1415	61	-2.42	-2.42	-2.42	-2.42	-2.42	-2.42
13072-L5-2	GGATTTCAACCAACGTATAGCCCCCACCAGTA	1416	78	-6.78	-30.88	-30.88	-7.56	-5.70	-2.19
13075-L5-1	TGAAAGCTGAAGTCCAGCCCCAGCCCTCT	1417	56	-6.29	-6.29	-6.29	-2.93	-2.98	-2.05
13089-L5-2	ATCACCAACAACCTTGCCCCCACTCCAAGCCCTG	1418	61	-2.76	-2.76	-2.76	-2.76	-2.23	-2.76
13091-L5-2	CCTGAGAACCCCCGACCCCTCAAGTGTTCAGCAGGC	1419	78	-3.59	-8.53	-11.47	-12.31	-8.16	-1.70
13093-L5-2	CGACCCCGCAGAAACCCCAACCGCGCCCCGCGCAG	1420	67	-4.46	-10.00	-7.28	-7.97	-7.96	-1.06
13095-R5-1	GCAGGACCCACCTCCCTACTCTCTGGCCCCAGGT	1421	67	-10.16	-10.16	-4.31	-2.54	-5.00	-2.96
13097-L5-2	AGCCTCAGCCCCACCTCCAGCCCCACCCCTAGGG	1422	56	-3.53	-3.87	-4.76	-4.17	-3.23	-1.41

13110-R5-1	AACAGCAGACAGGGTCCCAGGGCCCTCCCTTGT	1423	56	-4.98	-4.98	-4.98	-1.26	-2.35	-1.54
13115-L5-3	ACCCCTCCAGGGCCGACCAACCCCAACCCAAAC	1424	78	-8.12	-18.66	-32.34	-7.11	-5.65	-2.64
13119-R5-2	GACTCTGCCGCTCCCGCCCGCCACCTCCCTGT	1425	61	-3.93	-3.70	-4.56	-3.71	-3.93	-1.95
13124-L5-1	TCCTCCCTCCGCGAAAGCCTAAACTTACCCCTCA	1426	50	-2.05	-2.84	-5.18	-3.12	-2.71	-1.10
13129-L5-3	AGCCTTCCTGTCCCTGGCCCCCGACCTGCTCCA	1427	78	-2.06	-2.16	-2.55	-2.91	-3.95	-6.35
13130-L5-1	AGCCCGCCCAACCCACCTCGATCTTTTCCTC	1428	56	-2.86	-5.27	-3.54	-2.26	-1.75	-2.84
13135-R5-1	CGAGGAGGCCCTGAGCACTGCCACCCCCACACAC	1429	72	-4.99	-4.99	-6.04	-4.00	-7.63	-1.74
13136-L5-3	ACACTTGCCCTGCCAGCCCTGCTCAGGGCCACCCCT	1430	78	-6.86	-6.86	-6.86	-6.86	-2.82	-2.13
13137-L5-1	GCTCTAACCCCGCAACCCACCTCCCATGCC	1431	61	-2.47	-3.81	-5.83	-3.96	-3.31	-1.66
13138-R5-1	TTCGCCACGCCCGCCACCCGAGCTGCCTCCC	1432	78	-5.42	-5.75	-7.42	-6.65	-4.73	-2.51
13163-R5-1	AGACCAAGCCCTCCACCCCTTCCCTCCCAGC	1433	72	-5.88	-16.87	-21.10	-2.25	-2.25	-2.13
13164-L5-1	GAAACAAACATCTAATAAACTCTCAGGATACC	1434	50	-5.10	-5.10	-5.10	-1.16	-2.10	-1.52
13166-L5-1	TCAGAGGACCCCGCAGCCACCCCGCGAGGC	1435	83	-6.08	-6.39	-9.71	-13.71	-5.78	-2.21
13181-L5-1	GAACAGTCACTGCCTGCCCAACCTCTTTCAGGA	1436	67	-7.92	-7.92	-4.10	-1.01	-3.99	-7.92
13184-L5-1	CAAAAAATAAACCAAAACCCACCACTAATG	1437	56	-2.31	-2.31	-2.31	-1.65	-2.31	-2.31
13186-R5-2	ACATTATCTCTGTGATCCCTCCCTCACCA	1438	67	-4.00	-9.93	-26.53	-3.44	-2.79	-1.55
13195-L5-3	TCTATATTCCTCCCTCCATGACCAATTGTATTAG	1439	78	-3.32	-11.15	-8.93	-3.70	-2.85	-5.82
13199-R5-1	GCTGGGGATTCTGCCCATGCTTACCTCCC	1440	50	-2.48	-8.02	-5.59	-2.40	-6.24	1.53
13202-L5-1	GGGTCTCTGCCACCTCCTCTGAGAAAGCCCAACCA	1441	61	-10.20	-10.20	-10.20	-4.67	-4.70	-3.14
13202-R5-2	GGTGGGAGACTGGAGAGAGCCCTCCTAATG	1442	56	-3.89	-14.39	-14.39	-2.07	-2.32	-1.06
13209-L5-2	TAACTCGCCTGTGCCCCCGCGGCTGCCCGCGG	1443	61	-45.23	-4.90	-5.49	-3.79	-3.36	-1.44
13209-R5-3	TGGTCGCCCGCGCAGCGCCTGAAGGCACGGCGG	1444	61	-12.46	-12.46	-3.05	-3.57	-6.27	1.13
13211-L5-1	ACGCGCCCCGCGCTCTCTGACCGACCGGAGCGC	1445	61	-7.25	-6.22	-5.47	-5.00	-4.38	-1.43
13220-L5-3	CCTGTAGCTGCCACTGCCCTTCTCCTCACTCAACC	1446	72	-5.92	-19.89	-9.41	-4.26	-3.98	-1.61
13229-L5-1	ACCCGTCCCTGCCCTTTACCCCTTGGGCCAGCA	1447	61	-5.10	-4.62	-6.00	-4.16	-3.60	-1.59
13230-L5-4	TAGTCCAGTGCCTCCAGCTCCAACCACTGAA	1448	50	-2.05	-2.05	-2.05	-2.05	-1.39	-2.05
13231-L5-2	GGGGCCGCTCCCCAGCACCCGACGCCAGCATCG	1449	56	-16.99	-16.99	-3.88	-4.81	-8.33	1.00
13237-L5-4	ACCCCTCCAGGGCCGACCAACCCCAACCCAAACAA	1450	72	-35.36	-16.82	-35.36	-6.88	-5.65	-2.41
13239-L5-2	CAGAGCTCCCCCATCTCCCCAGACTTACCCCT	1451	72	-4.10	-15.67	-39.77	-6.16	-5.51	-1.58
13240-L5-2	AATCGCGTCCCCCGCGGCAATCCCGGCCCCAA	1452	72	-60.67	-6.31	-7.99	-5.64	-4.93	-1.84
13241-L5-2	ACACCCCTCCAAACCAACACAGAGCAAGCAAGG	1453	72	-10.26	-10.26	-10.26	-3.74	-4.35	-10.26
13251-R5-2	TTGTGCCCCCTCCTCTGCCATGGCTGGTCCCTGG	1454	78	-5.70	-7.57	-14.60	-5.42	-4.16	-2.77

13259-R5-1	GCCAGAATTACCACTGTATCTGTCCCAACCCC	1455	72	-3.29	-3.29	-3.29	-3.29	-3.29	-2.05	-3.29
13267-L5-1	CACTCCCTGCTGGCCCCACCTCACCTATGGTG	1456	61	-1.87	-2.26	-13.85	-2.21	-3.68	1.35	-3.29
13281-L5-3	CACCCCAACCCACAGGACAGAGGAAGTGACGAG	1457	56	-3.70	-9.33	-5.75	-7.17	-4.05	-6.04	-3.29
13283-L5-3	GGACCCCTGCCTTCCCTTGCTGCGCACCCCTTTGCACA	1458	50	-3.60	-7.29	-3.33	-3.32	-2.19	-1.11	-3.29
13285-L5-3	GCTATGCACCCAGCCGCCAGCTCAGCCCCCTGC	1459	78	-6.84	-6.59	-7.59	-4.39	-4.35	-1.93	-3.29
13287-L5-3	GGAGCCACCCACCCCTCCTCCCAAGACCCACAT	1460	56	-3.34	-3.34	-4.88	-3.81	-2.82	-2.04	-3.29
13291-L5-1	CCCAAGCGCCCTTCCCTCCCTCCCTCCCTCCCG	1461	50	-2.07	-2.35	-3.92	-2.62	-2.12	-1.10	-3.29
13293-L5-1	TGCCTGGCCAGGCCCTCCGCCCTTGCGCTCGCCAC	1462	61	-3.68	-3.42	-6.95	-3.41	-3.34	-1.42	-3.29
13298-R5-1	CAGGGCTTCAGCCTCCCCACAGCCCCACACTT	1463	78	-14.82	-14.82	-14.82	-8.93	-8.41	-4.71	-3.29
13303-L5-3	CCTCCCCTGAAACCCAGTTGCCACAACTTTCCAC	1464	50	-2.57	-7.48	-4.39	-2.24	-5.82	-1.05	-3.29
13308-L5-1	CACTCACATCAGCCCCATCCACCTCCACCTCTCCAC	1465	50	-14.43	-3.76	-3.43	-2.63	-5.96	-3.88	-3.29
13310-R5-1	ATCCATTGCCACTACCACCACCAATAATTAAAA	1466	56	-6.22	-6.22	-6.22	-6.22	-4.23	-6.22	-3.29
13312-L5-2	TGCCACCCCAACCCCTCCCCACAGCCCCAGCCC	1467	72	-3.89	-4.71	-7.40	-5.54	-3.94	-2.64	-3.29
13313-L5-2	CCTGTCTCCCATGCCGTGTCCTCCCTCCCACTAACC	1468	61	-3.49	-4.79	-7.64	-2.95	-2.68	-1.55	-3.29
13316-R5-2	CCACCACCTTGCTGTCTGGCCCCACAGCACCAGGCC	1469	50	-4.30	-4.30	-2.50	-2.88	-4.30	-1.25	-3.29
13326-L5-2	CAGAGATTCCGGCTTCCCCCACCCGCCCTTC	1470	67	-4.02	-4.30	-8.40	-5.02	-4.30	-2.41	-3.29
13328-R5-2	TCACCTGCCCCCACATCTGCAACACACAAGAGT	1471	72	-14.53	-14.53	-14.53	-4.04	-7.65	-4.27	-3.29
13332-L5-1	CCATACTCTCCAGCTGTCTTCCCTCCCAA	1472	50	-4.39	-4.39	-3.09	-1.46	-2.15	-1.38	-3.29
13334-L5-3	TCCTGGCAGGACTCCCCTCCCCCTCCCACTGTG	1473	72	-3.05	-5.00	-8.54	-3.73	-3.16	-2.05	-3.29
13335-L5-3	ACCTCAGCCTCCACTGCCCTCCTCGCGCATCCTAT	1474	61	-4.66	-4.99	-9.25	-3.74	-2.81	-2.08	-3.29
13337-L5-2	TCTTTTCTGACATTCCCTCCCCCAACATGGAA	1475	67	-3.58	-4.94	-25.53	-3.71	-3.28	-1.08	-3.29
13339-L5-1	GACTGAGGGTTTAAAGAAAGATGGTGCCGCCGC	1476	78	-45.98	-6.02	-7.34	-3.40	-4.61	-1.22	-3.29
13343-L5-1	CCTCCACCCCTCCCGCAGCGCCCTCCCCCTCA	1477	50	-3.99	-5.54	-7.36	-3.68	-3.31	-1.90	-3.29
13349-L5-2	CACCACCTGTGTGGGCTATCCAGCCGCCCTCC	1478	83	-7.08	-7.21	-6.98	-4.08	-4.09	-1.92	-3.29
13353-L5-2	GCTTCGGCCACAGAAATGTTGCCCTCTGAAATC	1479	50	-2.24	-2.24	-2.24	-1.67	-1.24	-2.24	-3.29
13354-L5-1	CTGCCCCACAAAGCTCCTGTGAACCCCCCTCAGT	1480	72	-17.63	-17.63	-17.63	-4.43	-9.91	-1.86	-3.29
13355-L5-2	TGACCACACCCACCCCTATCCTTTCTGCAGTGTG	1481	78	-5.51	-5.51	-5.51	-5.51	-3.48	-5.51	-3.29
13356-L5-2	GGCCCCCTGGCCTCCCTGCCACCCAGCACGGTG	1482	56	-8.37	-4.34	-4.14	-2.41	-4.28	1.04	-3.29
13358-L5-2	CCTCCTCACCACCCCTCCACACTCCTGGGGAAGT	1483	78	-23.01	-23.01	-23.01	-5.08	-7.50	-6.25	-3.29
13361-L5-1	GGGAAGCAGAGTCAGTGACCCCAAGCCCTGCACAC	1484	83	-12.93	-12.93	-12.93	-9.37	-12.93	-3.94	-3.29
13363-L5-2	GGCCCCCACCCTCACCTGCTGACACCCCTCACATCC	1485	83	-20.98	-20.98	-20.98	-5.06	-8.92	-6.01	-3.29
13364-L5-2	TCCCAGGACCCCTTCCCTGAGCCTCAGCCCAT	1486	78	-7.47	-7.47	-7.47	-7.47	-7.47	-7.47	-3.29

13365-L5-3	TCCCCAGAGCCCGCCCCAACCCACCTCGATCTTTT	1487	56	-3.16	-5.55	-6.40	-2.46	-1.71	-2.63
13370-L5-2	CATGCCCCACCTCACCTCTGCTCCACATACT	1488	50	-3.37	-4.07	-4.87	-4.29	-3.66	-2.25
13373-L5-4	GCCCTTACTCCAGCCCCACCGGCTCCCACTCACCT	1489	56	-7.98	-7.98	-3.68	-2.29	-4.40	-1.07
13374-R5-1	AGCCTTCCTGTCCCTTGGCCCCCGACCTGC	1490	78	-5.54	-4.92	-5.77	-5.78	-4.29	-2.09
13375-L5-3	GAGAACCTGAAACCCACGCCCTGCCTACCCCTTAG	1491	72	-4.07	-4.48	-7.27	-6.24	-3.83	-1.92
13376-R5-1	TCTGCTGCAGGTAGTCTGAATGTCCCCCAACATC	1492	78	-4.80	-21.42	-28.03	-11.86	-5.27	-3.27
13380-R5-3	AACTCCCAGCCCCGGAAGAATGCACACGCAGGAG	1493	61	-10.07	-10.07	-10.07	-3.28	-5.31	1.13
13385-L5-1	AATGACTCCTCCGGCGCCACCTACAGT	1494	50	-3.98	-3.98	-3.98	-1.32	-1.86	-3.98
13396-L5-2	GACCTGCCCCGCCCCACTCGGGCTCCTTACCG	1495	78	-4.91	-4.88	-2.43	-5.34	-3.01	-2.26
13403-L5-1	TCCCCCAACCTGGGGCCAGGCCACCTGGT	1496	67	-2.41	-11.03	-7.77	-2.10	-2.32	-1.72
13412-L5-1	AGTCAAGATGGCGCCCCCTGGTCCCAG	1497	61	-2.15	-8.32	-5.84	-5.72	-4.18	-2.68
13423-L5-3	GCCCTCCCCTGACTCCCTGAAGCTATTTGTTT	1498	72	-5.62	-16.59	-20.12	-2.22	-2.23	-1.92
13425-L5-3	TGCCCTCCCAGTCATATGCGCGGCATCAAGGTT	1499	72	-5.55	-18.57	-19.05	-2.11	-2.19	-1.90
13430-L5-3	TTCTTACCTCCCCCAACCCCATCCAGTTACC	1500	72	-2.66	-5.22	-11.00	-5.21	-4.82	-2.37
13431-L5-3	CGACACCCACTCACTGCCGCTGCCGCACTCACAGC	1501	72	-6.02	-7.11	-7.16	-4.36	-6.77	-1.47
13432-R5-1	ACCTCCCCAAACTCCACGTAATCACCTACTATT	1502	50	-2.31	-4.47	-3.97	-1.94	-3.01	1.02
13456-R5-2	CAGATGCCCGCTATGAAATCTTTTCCAACC	1503	50	-1.49	-5.78	-14.18	-2.24	-1.86	1.34
13458-R5-2	CCTCCTCGGCACTCCCTCTACCTCACTGTCCAC	1504	56	-9.56	-4.35	-9.56	-2.05	-1.81	-2.58
13461-L5-4	TCCCAGCCCCGTCCTCCCAACCCCTAGAGAAAGTGAA	1505	78	-4.88	-5.69	-6.76	-5.00	-4.09	-2.33
13463-L5-2	ATCCAGGACTCGGACCAAGCCCCCAGCCTGAG	1506	78	-5.87	-22.46	-12.28	-4.92	-5.26	-2.41
13489-L5-1	GGGGAAAGCCTGTGTTCAAGCCCAAGGAACAAGA	1507	61	-2.32	-2.32	-2.32	-1.56	-2.32	-2.32
13497-L5-1	TGTGCCCCCCTCGCTCCCAAGCCCCCAGGGACCGC	1508	72	-6.04	-9.24	-13.05	-5.73	-5.31	-2.89
13513-L5-1	ATCCTCCCTGGTGGTGCTACTCTCACCAAAG	1509	50	-3.48	-3.48	1.08	-3.48	-3.48	-3.48
13519-L5-1	ACCTTAGCTCTACCCAAACCTCGCTTCCCACCCCC	1510	72	-8.34	-8.34	-8.34	-6.17	-4.60	-2.83
13525-L5-2	CCATGCTCTCGCGTGATCTCCCTACCGCCATCGT	1511	78	-3.26	-11.76	-11.76	-3.63	-4.94	-3.67
25-R5-1	TTCCCAGAGCCTCACCCCCCTCTTTTCTAACC	1512	50	-2.21	-2.37	-8.78	-2.46	-2.10	-1.26
266-R5-2	GTCGCCCCCTCCCCCAAGTTGAGACTTGCA	1513	78	-3.86	-6.49	-10.77	-4.19	-3.42	-2.49
2786-L5-3	CACCTCCCGCTCAACTGCCCATACTAATGCTTTT	1514	61	-7.14	-7.14	-7.14	-5.01	-4.23	-7.14
2811-R5-1	CCACAGCCACCCCGTGCCACTGTGTCCCAACCC	1515	67	-5.19	-5.19	-5.19	-5.19	-5.19	-5.19
2819-R5-4	CAGCCTGCCACCGCGCTTTTGAAAGAACCACTTCA	1516	78	-8.91	-8.06	-8.93	-4.59	-8.36	-1.88
3717-L5-2	CCGCCCCCTCCCATAGCCTCACCCCCAAACCCA	1517	61	161.90	-6.67	-8.36	-1.07	-3.17	-1.51
3732-R5-1	TCCCTTTCCCTGCCAGCAAAACCCCAACCCCTAAG	1518	50	-13.70	-5.71	-2.96	-2.29	-2.86	-1.31

3799-R5-1	CTGAAGATGCTCCAGAGGCCCCCGCGGGCC	1519	67	-3.62	-5.47	-7.63	-3.60	-3.29	-2.20
3897-R5-2	CCGACCCGCGCTCAGCCGCTCCCCCTCAG	1520	72	-6.31	-6.25	-5.66	-5.22	-5.39	-1.95
3942-L5-1	TCCCTTCACTCCAGTTGCCAAACAGATCCCCCACTCCC	1521	61	-2.19	-6.40	-7.64	-1.72	-2.42	-2.05
3952-L3-1	GAGCACTCAATCTGACACCCCTCGCCGGGG	1522	72	-7.09	-7.09	-7.09	-2.41	-4.47	1.53
3953-R3-2	ACTCCAGCCTCGCGCGCTCAGCTTCCCGAGC	1523	67	-	-	-	-	-	-
3966-L5-1	ACCCAGAGCTGTGCGCGCGCTGCGCGCTTCGCC	1524	78	-8.01	-6.99	-8.17	-5.56	-7.00	-1.82
3995-L2-2	CTATAAACTTCGAAAAGTCCCTCCTCCTCAGT	1525	50	-1.95	-3.65	-12.55	-2.00	-2.04	7.12
4013-L4-1	CTATGGCACTTGATGTTGAGCTATCAGCAGGGGACCAT	1526	67	-2.19	-2.19	-2.19	1.07	-2.19	-2.19
4026-R5-1	GAGAGAAAGCCCCCTTTGTCTGGCTTTG	1527	83	-81.69	-81.69	-18.32	-7.08	-6.50	-3.10
4026-R5-2	GGCGAGAGAGAAAAGCCCCCTTTGTCTGGC	1528	72	-7.22	-14.47	-19.24	-5.56	-6.21	-3.11
410-R5-1	CCAGCTCCACTACTCCGTCCCGAGGAAGCAAAACACGGCACC	1529	67	-5.18	-5.18	-5.18	-5.18	-3.55	-5.18
4130-L5-1	TCAGCCAGCAGCGCGTCCATGTCCACACGACCCC	1530	78	-18.42	-18.42	-13.52	-4.27	-4.12	-1.61
4143-R5-2	TCAGCGTCTTGCTCTCCTCCTGGTAACAGCAGCC	1531	67	-3.75	-3.75	-3.75	-3.75	-3.75	-3.75
4258-L5-1	CCTAGAGAACATATATCTGGTGCTCTCCTCTTTTCCCGT	1532	56	-2.02	-2.02	-2.02	-1.46	-1.05	-2.02
4315_C-L4-1	GCAGCCCTCTCCTCCGAGAGGTTGGGGTCGGCGCGCCCGGCC	1533	72	-3.12	-84.00	-7.08	-2.18	-2.69	-2.05
4315_D-R4-1	TCCCGTCCCCCTCCCC	1534	72	-6.78	-6.28	-7.97	-6.33	-6.20	-2.25
4315_E-R4-1	GGAAAGTCAGCCCCCAGCGCCCCCGGAGTTCTTGG	1535	50	-4.86	-4.04	-4.85	-3.23	-4.16	-1.61
4315_F-R4-1	CCCCACCAAACTATTCCCGCATCTCCCCGGCTCTGG	1536	67	-4.81	-4.70	-9.00	-5.38	-5.04	-2.05
4315_I-L4-1	AACCCGGGCTCCCCACCCGCTCCCTGAGC	1537	56	-3.26	-4.17	-7.04	-2.60	-2.59	-1.37
4315_K-L4-1	ACACCTCTGCGCCCCCTCAGGCGCCCTGGGCTCGGCGCCCCCGCC	1538	67	-5.56	-4.67	-6.69	-5.88	-4.81	-1.81
4315_L-L4-1	CGTCCCCAG	1539	56	-3.38	-3.19	-12.44	-3.43	-6.47	-1.41
4315-R3-2	TCCAGGGGGCCCTGAACCTTGTCAAATCCTCGCCATCCTCCACCCC	1540	50	-2.07	-2.07	-2.07	-1.43	-2.07	1.59
4338-L5-2	CCTGGGGTGGCGGGAGCTGGCCCCACTGC	1541	61	-5.28	-5.28	-5.28	-2.61	-3.21	-5.28
4340-R3-1	ATTTTCAGCCCCCTTGTCGCCAGGCCAAAC	1542	67	-3.82	-4.80	-7.88	-4.88	-3.22	-2.13
4346-L5-1	TTCTTCCCCCGCCCTCGCCGCGCGCCGACCCGG	1543	67	-2.26	-2.85	-16.06	-2.89	-2.23	-16.06
4361-R5-1	CTCAGCGTCTCCCTCCCTCATGTGCACATGT	1544	56	-2.36	-9.82	-2.18	-1.93	-6.91	1.37
4498-L3-2	GAGATCCAGACGGCCGTGCGCCTGCTGCCT	1545	72	-14.82	-14.82	-14.82	-4.16	-7.74	-1.36
4516-L5-1	TCCATCACCCCCCAGGCTGACTCTGGCTCCTGCCTGGCTCTGCC	1546	67	-2.62	-2.62	-2.62	-2.62	-2.62	-2.62
454-R5-1	CCAGCTCCACTACTCCGTCCCCGAGGAGGCCAAACACGGCACC	1547	50	-1.18	-1.22	-2.44	-1.60	-4.72	-1.09
4593-R5-1	CTATAGCAGATGACATAACTCCCCCGGCATCAG	1548	72	-7.69	-7.69	-7.69	-7.69	-4.10	-7.69
4610-R5-1	CCTCTGGCCCCCTGCCTAATTGGCTGC	1549	56	-3.23	-3.23	-3.23	-3.23	-1.70	-3.23
4610-R5-2	GCCCTCTGGCCCCCTGCCTAATTGGCTG								

4642-R3-2	CAGCGCTCCCTTCCTTATTTGAGATCTGTGCA	1550	67	-10.25	-10.25	-10.25	-10.25	-10.25	-5.81	1.30
4666-R5-1	GACTCCCCCAACACCTGCGGTGGCAC	1551	78	-3.26	-6.47	-13.10	-5.57	-5.25	-2.30	
4792-L5-2	AAGCCAGTTACAGCCCCCACTGCCCCCATAAAC	1552	61	-4.51	-4.61	-6.34	-4.01	-4.00	-1.49	
4801-L5-2	GAACTCTGCCTCCTGTTTGCTACAAAAACA	1553	61	-4.53	-4.53	-4.53	-4.53	-2.36	-4.53	
4813-R3-1	GATTTTACAATCGGGCTGTTAACCCCTCCCG	1554	61	-3.33	-3.33	-3.33	-3.33	-2.04	-1.06	
4875-R2-2	CACAGCCCCCTTCCTGTGACTTCACAC	1555	67	-2.71	-25.83	-7.24	-2.30	-2.06	-1.91	
4912-L5-2	TACCCCTGGCCCCCAGCTGTGATTGTCTAA	1556	78	-19.46	-19.46	-5.35	-5.07	-7.98	-1.65	
4929-R4-1	GACGCTGACCCACCGCAGCTGTGACTGT	1557	89	-11.74	-11.74	-11.74	-8.30	-7.20	-11.74	
5032-R5-1	CGCACCCGTCCCGTTTCGTCCCCGGACGT	1558	72	-13.13	-13.13	-13.13	-8.64	-6.34	-13.13	
5048-L5-1	CGCCGCTCCTTGACGCGTACTTCTGCTGCACACC	1559	61	-11.59	-11.59	-11.59	-3.55	-5.68	-1.14	
5071-R5-2	GACCCCGCCCCCAGTCCCAGCCCCAATTAATA	1560	72	-6.98	-7.41	-8.60	-7.82	-3.38	-3.14	
5107-L5-1	AACTCCCCCTTGACCCCCCAGTACAAACTG	1561	61	-2.41	-26.66	-26.66	-3.80	-3.15	1.08	
5210-L5-1	AAAGCTGCCCCATGGGGATCCAAACCCCA	1562	67	-1.97	-7.99	-7.99	-3.58	-4.09	-7.99	
5342-L5-1	CCACCAAACCAATGCCGCTGCTCTCCTTCCA	1563	56	-4.62	-3.77	-3.56	-3.08	-2.92	-1.28	
5491-R5-1	GCCGTCGCCCAACCATCATCTATCAGGCCATGTGGCA	1564	72	-5.64	-4.77	-6.07	-3.45	-4.72	-1.52	
5521-L5-2	AGGTCTGTCTTGGGTGGGCCCTCCCCAGAGCAC	1565	50	-7.60	-7.60	-7.60	-2.62	-1.22	-2.35	
554-R5-1	GAAAGAGAACAGAGAGAGCCCTCCAGCAGCC	1566	67	-5.86	-17.89	-18.96	-2.04	-2.19	-1.95	
5554-R5-2	CCCCACCCCTCATCATGCTGCTCCCAGATC	1567	61	-3.72	-3.98	-5.48	-4.08	-2.80	-1.59	
5638-R5-2	GGCCCTCCCCCTGCCTGTGATAGGCTG	1568	78	-5.39	-15.00	-20.91	-2.61	-2.48	-2.16	
5640-L3-1	GCCATGGAACACCGTGCCCTGCCCTCTCGAGA	1569	61	-2.75	-4.95	-5.84	-2.71	-2.38	-1.80	
5749-R5-1	ATCATAATCCCCCTTTTGACTTTACATGATC	1570	56	-3.50	-3.50	-3.50	-1.35	-1.84	-3.50	
5757-L5-1	TGCTTTATTGACTAATGAACCTACCGTGCCGCCAAC	1571	89	-22.78	-22.78	-22.78	-4.38	-4.58	-5.43	
5854-R5-2	GCCCTCCCTCTCCGAAAGAAATGTGTACCCGGG	1572	72	-4.59	-14.74	-16.09	-1.96	-1.94	-2.01	
5956-L5-1	GCCTGCTCTGCCAACCCCAATCCGTCAAGACGCATAG	1573	50	-1.50	-3.54	-8.83	-2.47	-1.20	-8.83	
5995-R5-1	CCCCCTTGAGGTTCCCTACTGAAATCTGACAAATCAG	1574	78	-2.29	-2.29	-2.29	-2.29	-2.29	-2.29	
6008-R5-1	ACCCCCACCTTTTTCCTGTACCTTACCCGGAG	1575	83	-9.71	-9.71	-9.71	-9.71	-5.88	-3.27	
6008-R5-2	CAACAATACCCCCACCTTTTTCCTGTACCTTA	1576	78	-5.41	-5.41	-5.41	-5.41	-5.41	-5.41	
6016-R2-1	AACTCCAGCAGCCCCCGTCAGCCTCCTGCT	1577	61	-3.58	-3.27	-3.65	-3.22	-2.34	-3.50	
6023-L5-1	AAGGCAGAGCCAGATGACTGGTCCAACCCCCAGAGACC	1578	72	-4.72	-19.11	-19.11	-5.65	-9.37	-1.46	
6087-L4-1	CTGTCTCCATTACTGCCCTGCCACCTTCTCCATC	1579	78	-8.97	-8.97	-8.97	-6.19	-5.07	-8.97	
6096-R5-1	CTTTCCTTATGTTTTTAATCCTGCCCCCGT	1580	89	-12.65	-12.65	-12.65	-2.15	-6.96	-12.65	
6192-L5-1	AATCCGTGAAGATAAAAAACCCACCCACCCAGCAC	1581	50	-13.47	-13.47	-2.70	-1.88	-6.10	1.34	

6198-R5-2	GCCGCCGCCGCCGGTCTTCCCGCGAAGCCT	1582	67	-4.03	-2.96	-5.71	-4.41	-4.65	-1.50
6242-R5-1	CCCCACAGTGGCATAATGTGACAAACCCAAAGCCCCCTGG	1583	67	-5.90	-5.90	-5.90	-5.90	-4.69	-5.90
6287-L3-2	GCCCCGCCCCACCTTTTCGGGGCTCACCTGGC	1584	72	-19.21	-19.21	-4.59	-4.31	-2.70	-1.67
6385-R5-2	GGCCCTGCCACGACCGACCGACTCCCGCGGCC	1585	67	-9.70	-9.70	-9.70	-9.70	-5.59	-3.26
6409-L3-1	CGTTCCCAACCGCACGCGCGCCTTCTGTGAAC	1586	72	-8.06	-7.09	-6.76	-4.68	-5.18	-1.54
6434-R5-1	TGCAGCCCTCCACACGACGCTGCAAGTGC	1587	67	-4.42	-12.82	-29.88	-2.13	-2.32	-1.24
6490-R5-3	CCCCATCCCCCATATGACGCTTCCCCCTCCTAAC	1588	67	-3.39	-4.20	-23.58	-4.26	-6.83	-1.35
6496-R5-2	CCCCCTCCCCACCCACCACTTCCCCTAGA	1589	67	-44.85	-14.21	-9.47	-4.89	-3.70	-2.49
6584-L5-1	GTCGGCCCTGCCTCCTCCTCCTCTCACCACAAGC	1590	56	-13.67	-5.50	-13.67	-2.42	-2.29	1.17
6590-L5-1	GTTAATGAGAAACCAATAAACAGCCTCTGCCCCAGCTG	1591	56	-4.65	-4.65	-4.65	-4.65	-2.56	-4.65
6642-R5-1	GTCCTCCCCTCCCCTCGAGGTGTACACA	1592	56	-4.01	-5.51	-6.59	-3.85	-2.99	-1.63
669-R5-2	CCAGCCCTGAGCCACCCCTCCCGGGAACCCACACA	1593	67	-10.57	-10.57	-10.57	-1.47	-6.18	-2.07
6718-L3-2	GCCTCCACCACCATAGGGGCCAGAGCTTCTGCCT	1594	72	-3.98	-3.98	-3.98	-3.98	-3.30	-3.98
6839-L3-1	GCCCGCTGGGCCCTGCCACCCCCACCCCT	1595	72	-5.32	-4.50	-6.48	-4.99	-4.55	-2.17
6880-L3-2	ACCTCCCCCGCGAAGACATCCACATTCTGCA	1596	61	-2.19	-4.09	-5.43	-2.08	-4.56	-1.52
6908-L3-2	AGAGGCTACTGGGGGAACAAGACTGGCAAGCC	1597	50	-2.84	-2.84	-2.84	-1.73	-2.84	4.40
6984-R4-1	CCCCCTGCCCCAAGCATTTGCTTGGGCACCAAGTCCCCTGCAA	1598	67	-3.05	-18.59	-18.59	-4.32	-7.50	-1.51
7029-R5-1	CAAGAGCCCTGCCCCAGCAGCAGCGCGAC	1599	67	-10.64	-10.64	-3.52	-2.90	-2.27	-3.48
7061-R5-2	TCATGGAAACCCACCCCTTCCCATGCCCAACC	1600	56	-3.93	-4.19	-5.53	-4.35	-3.52	-2.45
7066-R5-1	TAGCCTGAAAAAAGATGCCCCACACGACCCCTGCC	1601	78	-6.87	-6.68	-11.13	-7.24	-11.42	-2.31
7069-R5-1	CTCGGCCCGGCCCCCTCCGAGCTCAACAGGCTCCCA	1602	78	-6.84	-7.03	-9.92	-5.66	-4.77	-2.63
7113-R5-1	CGCCTAATTAGCCCCCCTGCTCCGGAGGCTCACC	1603	78	-10.08	-16.70	-18.94	-7.28	-6.96	-3.32
7126-L3-1	GCACACCCGCTCTCCGGCCCGCCCTG	1604	61	-3.58	-3.60	-3.88	-3.21	-4.09	-1.48
7141-R5-1	ACCCCAATCCTTGTTATGTAACCTACCACCTACCCCTT	1605	72	-2.31	-2.31	-2.31	-2.31	-2.31	-2.31
7221-R5-1	GGAGAGAAACCCCGGCCACTTCCCACACCCCTGTGGC	1606	56	-2.57	-2.97	-3.51	-3.96	-6.16	-1.10
7313-L5-2	CTGCCTCTGCCCTGATCATCAAGCCCTCAAGGA	1607	61	-2.26	-2.26	-2.26	-2.26	-1.58	-2.26
7352-R3-2	GCCCCTGCCAGAACTCTTAACAGCTCTAATTGG	1608	50	-2.04	-2.07	-2.22	-2.46	-1.45	-3.40
7356 A-R4-1	CAGAGCCCGCTCTCGCGACCGACCTGCCGCCGACCCACAG	1609	67	-7.83	-6.39	-7.11	-4.76	-6.17	-1.56
7356-L5-1	ATCCGGGCTGCCACCGCGACATAGCCTCGCCCC	1610	56	-3.91	-3.91	-6.19	-3.02	-2.48	-1.08
7367-L1-1	AGGGTTAGAGCTGCCCCCTCTGGGGACCG	1611	50	-5.61	-1.12	-5.61	-1.54	-2.24	-5.61
7384-R3-1	CTCGCAAAGGATCTCCTTCATCCCTCCCCA	1612	61	-3.15	-7.30	-4.15	-3.12	-2.40	-1.29
7411-R3-2	AGTCCCCCTGCCTCATCTGCCACCCCTAATGAC	1613	50	-2.55	-2.66	-13.01	-2.77	-1.83	-3.86

7569-L5-2	TTCAGGCCACAAAGCTACCCCCCAAGACAG	1614	67	-3.93	-3.93	-3.93	-3.93	-3.93	-3.93	-3.93	-3.93	-3.93	-3.93
7571-L5-1	CAGGGCTAACAGGGCTCCCCCACCCTAAG	1615	56	-3.75	-4.46	-6.71	-5.25	-4.30	-1.32				
7572-R5-2	ATCACCC TTCCCCTCCC AATAAAGCCAAA	1616	50	-22.49	-4.07	-6.89	-3.47	-2.12	-1.57				
7660-L5-1	GCCTCCG CCTGG CCGAGCGATAAAGCTC	1617	56	-3.55	-7.83	-4.16	-3.75	-3.08	-1.29				
7702-L2-1	CCAGAGA ACCGGA ATTC TCCCCGCC CC	1618	78	-6.38	-6.47	-8.59	-5.58	-4.12	-2.55				
7736-L5-1	ATCCCCACC ACCACCA GAAAGCGACGGTCTCC	1619	50	-10.35	-10.35	-2.65	-4.09	-5.59	-2.98				
7743-L5-1	GAGTCCCCA GCCAGCTCC AGTTCCGACCTGCCTT	1620	67	-7.13	-7.13	-7.13	-7.13	-5.90	-7.13				
7781-R5-2	AGCCTGTGCC TGCCGCTGTCTAGTACTGGT	1621	72	-22.71	-12.29	-6.84	-3.46	-3.67	-1.44				
7824-R5-1	TGCCAGCTTC ATCGCCG CCTCACACACACA	1622	72	-10.99	-8.91	-14.45	-5.40	-5.99	4.65				
7846-L5-2	GTCCTCTCCA TCCTCCATCC CTCCACCAGCCCT	1623	50	-2.66	-2.66	-2.66	-2.66	-1.45	-2.66				
7883-R5-1	CTCTCCCC TC CGCCCTCTC ACCCTCCCAACCTTATTAGAAAC	1624	50	-3.69	-4.72	-7.94	-3.56	-2.66	-2.03				
78-R4-1	CCAGAGCTTCTCAAGGGG GACTACA ACTCCCAAGGTACATACAA	1625	50	-2.39	-2.39	-2.39	1.35	-2.39	1.44				
7949-R5-1	GATGCGCGC GCGACCG CGCGCAGCTGCAATTCATAC	1626	72	-4.04	-3.23	-5.42	-3.02	-5.45	-1.05				
7971-L5-1	TGTCACTCCC CTGCCGCTCC CTGGTGCA GGCT	1627	61	-7.45	-7.45	-7.45	-3.26	-5.59	-2.17				
8016-L3-1	TCAGCGCAACAAGCCCC GCAGTCACCCCTCT	1628	50	-3.02	-3.60	-2.40	-2.98	-2.70	1.10				
8062-R5-1	GGCAAAGGTCA CGGGGTCCA AGGCCTTAAGCACCTCCGCCA	1629	50	-7.61	-7.61	-2.06	-1.46	-1.16	-2.34				
8077-R3-1	CCATTCCC ACCTCAGG TAGTAAAAATA	1630	67	-2.08	-10.66	-3.95	-3.94	-2.28	-10.66				
8089-L5-1	TAGAGACCCTTA TAAGCTCAGGGCCACCCCTCCC	1631	78	-12.58	-12.58	-12.58	-4.10	-6.46	-4.03				
8239-R5-1	TGATTTTCTT CCCACTTCA CCTCCCTCTGAGCTCTCCA	1632	56	-7.85	-7.85	-7.85	-2.34	-5.84	-2.52				
8250-R5-2	CCGACCCG CCGTCAG CCGCCTCTCCCTCAG	1633	72	-8.36	-6.52	-12.10	-5.21	-5.60	-2.31				
8281-L5-2	CAGCCCC TC CCGAGCTGC AGCTGAGGGC	1634	61	-2.82	-7.08	-8.59	-2.10	-2.55	-1.83				
8298-R5-1	GATGCTGGC GTCCGCCG CAGCCTCTCGCCCCATCCCGG	1635	72	-7.60	-6.38	-5.79	-4.53	-4.42	-1.51				
8329-L5-1	TCCTTCCA AACGCC CACCTGGGT CAGCTC	1636	72	-7.14	-7.14	-7.14	-4.59	-3.30	-7.14				
8336-R5-2	CCTCCCCA CTCAGTCCC CACACCCCCAGCCA	1637	56	-3.54	-3.08	-4.97	-3.00	-3.92	-1.33				
8394-L5-2	TGGGGCCC CCGCCCTG CCCATCTCCGACTATCC	1638	50	-3.26	-3.15	-4.42	-4.40	-2.96	-1.58				
8564-L5-1	ACCCAGTTGCC AAAACAG ACCTCCCACCCCT	1639	72	-9.90	-9.90	-9.90	-2.63	-4.92	-3.12				
8564-R5-2	AGGGGCTGG GGAATCC CAGCAGGGGAA	1640	50	-6.82	-6.82	-6.82	-1.12	-3.67	3.00				
8898-R5-1	CAGCCGAG CGGACG CCGCTCCCGCCACCATG	1641	72	-5.76	-6.58	-7.08	-4.99	-4.72	-2.23				
9021-L5-2	GAAACAAA CACCAA AGCTCCCCACACCA	1642	50	-9.46	-9.46	-9.46	-4.34	-5.86	1.27				
9068-R5-2	GTC TGCCCTCC CTCTTGATCA AAGACTGCTCT	1643	67	-6.33	-16.55	-18.60	-2.35	-2.21	-1.89				
9087-L5-2	AGGAAAAG AAACCC TC CCAAGTCC ATTCCCT	1644	72	-6.46	-10.39	-14.13	-5.66	-5.34	-1.11				
9134-R5-1	GAAAGGGT ATCCGC CTGGTTGCGGGGCTGC	1645	50	-2.19	-2.19	-2.19	-1.49	-2.19	1.49				

9217-L3-2	CGCTAAACTGACGATCCCCGCCGTGACTAAAGCCA	1646	72	-4.02	-10.70	-3.72	-4.17	-2.98	-1.46
9245-R5-1	TCTCATTAGCCAGCCACTCGTCCCAAG	1647	67	-7.70	-7.70	-7.70	-5.35	-4.29	-7.70
9287-L5-2	GATATTCAGAGCCCTCCCCAGCCACACA	1648	67	-4.67	-9.40	-29.04	-1.28	-2.93	-1.16
9369-L5-1	TCACACATTCTTCCACAGAGGGAATCAGGGGA	1649	50	-2.53	-2.53	-2.53	1.38	-2.53	1.22
9384-R5-2	AAACCAGCTAGCAAAACCGCTCCGTCCGTATTG	1650	78	-15.00	-15.00	-15.00	-15.00	-7.95	-4.59
9387-R2-2	TCCATCCTTGCCGTGCGCTTCATCTCAAAGCCATC	1651	67	-7.04	-4.60	-3.03	-4.14	-4.99	3.25
9564-R5-2	GCCGCCCGCCGGCACCCGGCCGGCCTGGGC	1652	50	-3.25	-3.61	-4.38	-3.31	-2.97	-1.18
9605-R5-1	GCCCCAAGTGAAAAACAGAGCAAAACTCATTTCTGA	1653	50	-2.11	-2.11	-2.11	-1.42	-2.11	-2.11
9691-L5-1	CATTTTCATCCGCATCTCCCTCTTGGCCCTTGC	1654	56	-3.33	-6.89	-4.05	-5.66	-2.65	-4.01
9770-R5-2	CACCTGTTGCCAACCCAGCCCTATAA	1655	67	-16.08	-16.08	-16.08	-3.95	-7.31	-1.78
9774-R2-2	CCGCCCCCTCACCGCCTCCTGCTCCCATCAGGC	1656	83	-7.02	-6.70	-13.75	-8.18	-5.97	-2.86
9812-L3-1	AAGCTCTATTTATCTGGGCTCCCCAGCTTGCT	1657	56	-1.83	-8.36	-8.36	-2.32	-2.02	-2.57
9866-L5-1	TAGGAGAGGGGCTCCATTGCCAGCCCCAGCCC	1658	94	-11.22	-11.22	-11.22	-11.22	-5.31	-3.72
999997-R4-1	TCCTCACTGGGCCCCACCAAACGTGCCACCCCTCAAGCCCCCA GGAGCTTCCTTAAC	1659	72	-4.04	-4.22	-40.37	-2.33	-5.94	-1.62
let-7b	TGAGGTAGTAGGTTGTGTGTT	1660	56	-1.48	-6.50	-2.74	-2.48	-2.61	-1.23
let-7c	TGAGGTAGTAGGTTGTATGTT	1661	50	-1.39	-5.08	-2.46	-2.14	-2.43	-1.26
let-7e	TGAGGTAGGAGTTGTATAGTT	1662	50	-1.27	-4.56	-2.29	-2.10	-2.25	-1.06
miR-100	AACCCGTAGATCCGAACTTGTG	1663	50	-1.16	-15.70	-1.03	-3.27	-2.27	-15.70
miR-101	TACAGTACTGTGATAACTGAA	1664	50	-1.54	-12.44	-3.65	-2.51	-5.38	1.25
miR-1182	GAGGGTCTTGGGAGGGAATGTGAC	1665	50	-2.49	-6.37	-2.91	-2.61	-4.75	1.04
miR-1207-5p	TGGCAGGGAGGCTGGGAGGGG	1666	50	-2.88	-9.91	-5.91	-3.63	-2.81	-1.28
miR-1224-5p	GTGAGGACTCGGAGGTGG	1667	61	-8.27	-8.27	-8.27	-2.05	-4.54	-8.27
miR-1225-5p	GTGGGTACGGCCCAGTGGGGG	1668	72	-32.61	-32.61	-32.61	-32.61	-14.04	-2.24
miR-1228*	GTGGCGGGGGCAGGTGTGTG	1669	67	-3.86	-3.61	-6.14	-5.78	-2.67	-2.95
miR-1234	TCGGCCTGACCAACCACCCAC	1670	61	-2.04	-2.04	-2.04	-1.47	-2.04	1.67
miR-125a-5p	TCCCTGAGACCCTTTAAACCTGTGA	1671	72	-3.80	-7.98	-2.23	-5.05	-2.99	-2.20
miR-126	TCGTACCGTGAGTAATAATGCG	1672	83	-6.57	-7.43	-7.29	-5.33	-20.83	-2.68
miR-1268	CGGGCGTGTGTGGGGG	1673	56	-10.67	-10.67	-10.67	-5.01	-5.42	-3.17
miR-130b	CAGTGCAATGATGAAAGGGCAT	1674	50	1.68	-2.21	-2.21	1.20	-1.89	-2.21
miR-140-3p	TACCACAGGGTAGAACCCACGG	1675	67	-4.44	-4.44	-4.44	-2.32	-2.82	-4.44
miR-145	GTCCAGTTTTCCCAGGAATCCCT	1676	61	-5.21	-7.10	-4.29	-4.12	-3.15	-1.88
miR-149*	AGGGAGGGACGGGGGCTGTGC	1677	56	-3.51	-2.89	-6.31	-3.80	-2.13	-2.15

184

Table 24: Target RNAs present at at least 5-fold decreased levels in at least 50% of tumor samples (con't)

Gene	Adk40	Adk41	Adk49	Epi42	Ksarc19	Kmalp21	Kmalp25	EPI-4	Kmalp44	Scc27	Lcnec31	Car13
10010_B-L4-1	-8.97	-3.24	-2.01	-1.22	-1.01	1.17	-4.47	1.82	-4.06	-1.50	-2.18	-10.58
10010_D-L4-1	-6.36	-2.67	1.05	1.34	1.42	-1.02	-4.33	1.72	-3.11	-1.85	-2.76	-2.21
10010-R2-2	-10.92	-10.92	-10.92	-10.92	1.41	1.26	-2.35	1.55	-1.11	-1.47	-1.43	-10.92
10030-R5-1	-1.67	-6.81	-6.81	-6.81	1.05	-1.33	-1.91	3.13	-6.81	-1.68	-1.42	-6.81
10145-L5-2	-2.03	-2.03	-1.03	-2.03	3.99	1.36	2.04	8.70	-2.03	2.51	2.49	-2.03
10231-R3-1	-5.36	-4.79	-1.71	-1.02	-2.00	-3.34	-3.16	-1.40	-2.97	-2.84	-3.01	-1.72
10260-L5-2	-13.31	-13.31	-1.39	-1.00	-1.36	-1.37	-2.68	-13.31	-1.73	-1.67	-1.91	-13.31
10333-L5-1	-6.06	-5.42	-2.91	-1.23	2.12	1.59	-3.76	3.00	-3.64	-2.40	-2.70	-2.39
10342-R2-2	-21.66	-4.09	-1.44	-1.12	-1.73	-1.64	-2.11	1.05	-2.44	-1.91	-1.99	-2.14
10345-R5-1	-38.49	-2.89	-2.15	-1.16	2.43	-1.11	-3.15	2.42	-2.93	-1.81	-2.89	-2.05
10374-R3-2	-33.82	-2.75	-1.86	1.01	-1.26	-1.38	-4.17	1.55	-2.29	-2.84	-2.81	-1.19
10435-R5-1	-2.03	-2.03	-2.03	-2.03	2.64	1.90	1.56	1.79	-2.03	1.98	2.99	-2.03
10533-R5-2	-2.60	-2.60	-2.60	4.07	3.00	2.42	-2.60	5.87	-2.60	1.93	1.78	-2.60
10543-R5-2	-2.16	-2.16	-2.16	-2.16	6.71	3.38	1.75	2.31	3.45	2.43	2.22	-2.16
10578-R5-1	-21.24	-21.24	-1.92	-1.12	1.03	-1.44	-3.78	-2.98	-2.45	-2.36	-2.85	-21.24
10818-L5-1	-21.17	-21.17	-4.88	1.05	-2.48	-3.84	-3.65	-2.42	-21.17	-3.59	-3.58	1.31
11370-L5-5	-5.19	-2.41	-1.94	1.06	2.73	1.42	-3.48	2.73	-2.57	-1.29	-2.09	-2.21
11605-L5-4	-2.54	-2.54	-2.54	-2.54	7.43	3.53	1.60	9.69	-2.54	2.07	1.86	-2.54
12184-L4-1	-10.83	-2.87	-1.07	1.72	1.44	1.16	-2.02	2.42	-10.83	-1.60	-1.80	-10.83
12184-L5-3	-5.57	-5.57	-1.45	2.35	1.86	-5.57	-1.13	-5.57	-5.57	-1.18	-1.11	-5.57
12224-L4-1	-3.22	-3.22	-3.22	-3.22	2.60	-1.30	-1.44	-1.08	-3.22	-2.02	-3.22	-3.22
12361-R5-1	-23.46	-2.62	-1.61	1.02	-1.78	-2.13	-3.81	1.16	-2.37	-2.85	-3.15	-1.06
12691-R5-1	-67.73	-4.80	-3.65	-1.52	-1.27	-1.61	-5.82	1.11	-3.99	-2.62	-3.78	-7.28
12692-L5-1	-44.23	-4.49	-2.70	-1.69	-2.39	-2.52	-4.11	-1.77	-4.02	-3.45	-3.57	-15.48
12693-L5-1	-7.71	-3.35	-1.78	1.31	1.05	-1.00	-4.98	2.24	-3.49	-2.55	-2.90	-1.64
12694-R5-1	-47.87	-4.06	-2.68	-1.06	-1.04	-1.38	-3.70	2.01	-3.35	-2.83	-3.35	-47.87
12696-R5-2	-3.88	-3.83	-1.63	-1.52	1.70	1.08	-2.08	2.73	-2.20	-1.93	-2.65	-1.28
12697-R5-1	-27.65	-2.65	-2.38	-1.00	1.40	2.28	-3.45	2.22	-2.90	-2.10	-3.18	-1.41
12699-L5-1	-29.74	-5.73	-1.65	1.05	1.25	1.02	-2.61	2.61	-1.59	-1.65	-1.88	-29.74

12701-L5-1	-3.53	-21.63	-2.75	-1.73	1.00	-1.43	-4.30	1.93	-3.70	-2.98	-3.18	-5.50
12703-L5-3	-5.66	-2.77	-1.61	-1.29	-1.04	1.10	-3.49	1.99	-2.38	-1.31	-2.16	-2.12
12704-L5-2	-23.11	-2.88	-1.96	1.02	1.23	-1.13	-3.66	1.95	-3.25	-1.73	-2.39	-3.04
12713-R5-1	-8.29	-6.75	-2.32	-1.25	-2.08	-3.21	-5.05	-1.31	-3.88	-3.28	-3.59	-2.05
12722-L5-1	-10.47	-4.23	-1.94	-1.16	1.08	1.06	-3.29	1.73	-2.48	-1.90	-1.51	-1.61
12723-R5-2	-18.13	-18.13	-1.57	1.12	-1.08	1.31	-3.01	3.13	-18.13	-2.21	-2.34	-18.13
12725-R5-1	-22.87	-10.61	-1.79	1.85	1.00	-1.07	-3.16	2.59	-1.96	-2.48	-4.02	-22.87
12731-L5-1	-14.83	-14.83	-3.85	1.07	-1.66	-1.01	-3.96	-1.24	-14.83	-3.53	-3.07	-2.48
12900-R5-3	-8.25	-8.25	-8.25	-8.25	3.50	1.81	-1.46	7.77	1.59	1.53	1.11	-8.25
12904-R5-2	-9.55	-9.55	-1.99	1.39	2.59	1.24	-1.19	3.91	-2.18	-1.08	-1.26	-9.55
12910-R5-2	-4.61	-4.61	-4.61	-4.61	3.61	3.13	-1.85	4.45	-4.61	-1.46	-1.18	-4.61
12925-L5-3	-2.02	-2.02	-2.02	-2.02	5.09	2.63	1.23	1.81	-2.02	2.16	1.67	-2.02
12932-L5-3	-2.89	-8.22	-8.22	1.29	4.62	1.65	-2.23	4.21	-8.22	-1.56	-1.84	-8.22
12939-L5-2	-2.19	-2.19	-2.19	-1.06	2.02	3.10	-1.04	-2.19	-2.19	1.30	-2.19	-2.19
12947-R5-3	-1.37	-13.13	-2.75	1.15	1.49	-1.39	-4.20	1.51	-28.53	-2.45	-3.99	-28.53
12975-L5-1	-2.38	-1.84	-1.17	1.10	1.87	-1.01	-2.24	3.23	-2.39	-1.76	-2.51	-1.73
12981-L5-1	-4.08	-4.08	-4.08	3.51	1.97	1.56	-1.29	-4.08	-4.08	-1.06	-1.09	-4.08
12981-R5-1	-9.73	1.26	1.11	3.25	1.24	-1.83	-2.07	1.61	-2.36	-1.79	-2.49	-1.21
12998-R5-1	-12.85	-12.85	-12.85	-1.18	-2.37	-4.23	-4.62	-12.85	-12.85	-4.10	-7.18	1.18
13004-R5-1	-1.80	-3.58	-1.64	-1.52	1.04	-1.21	-2.31	3.54	-1.93	-1.18	-1.69	-40.23
13047-R5-2	-2.69	-2.69	-2.69	-2.69	1.26	1.69	1.72	3.11	-2.69	1.66	1.70	-2.69
13050-R5-4	-15.79	-3.48	-3.24	-1.11	2.03	-1.04	-4.41	2.22	-3.65	-3.00	-3.58	-7.48
13052-L5-1	-26.92	-2.48	-1.98	-9.42	3.17	1.94	-3.50	4.57	-5.44	-1.79	-2.31	1.31
13066-R5-2	-2.42	-2.42	-1.04	12.45	2.07	-1.35	1.13	3.64	-2.42	1.57	-2.42	-2.42
13072-L5-2	-30.88	-3.39	-4.96	1.35	-1.14	-1.69	-4.40	1.45	-3.40	-2.95	-4.79	-30.88
13075-L5-1	-6.29	-6.29	-3.13	1.76	1.50	1.06	-1.59	3.33	1.36	-1.09	-1.21	-6.29
13089-L5-2	-2.76	-2.76	-2.76	1.48	2.89	1.67	-1.18	4.44	-2.76	1.25	-1.25	-2.76
13091-L5-2	-17.46	-17.46	-17.46	-1.31	-1.66	-2.34	-4.61	1.68	-2.00	-2.90	-3.09	-17.46
13093-L5-2	-20.22	-3.83	-1.31	1.02	-1.19	-1.65	-4.19	1.23	-2.08	-2.08	-2.36	-20.22
13095-R5-1	-10.16	-4.78	-2.40	1.37	3.83	-1.01	-2.18	2.72	-10.16	-1.67	-1.88	-10.16
13097-L5-2	-7.34	-1.77	-1.58	1.04	1.73	-1.12	-3.12	2.57	-1.98	-1.76	-2.44	-8.77
13110-R5-1	-4.98	-4.98	-4.98	4.37	1.73	1.02	-1.82	2.20	-4.98	-1.44	-2.75	-4.98
13115-L5-3	-32.34	-4.43	-16.43	-1.11	1.45	-1.55	-5.01	1.38	-4.43	-3.15	-4.39	-32.34

13119-R5-2	-6.15	-3.15	-2.12	-1.61	-1.06	-1.15	-2.81	1.90	-2.31	-1.75	-1.75	-1.55
13124-L5-1	-24.48	-1.59	-1.32	1.76	3.40	1.31	-2.81	9.22	-1.36	-1.46	-2.20	-24.48
13129-L5-3	-47.98	-6.80	-1.76	-15.40	-1.17	3.16	-3.45	7.03	-18.45	-2.26	-2.30	-13.71
13130-L5-1	-8.62	-2.94	-1.75	1.88	1.49	3.62	-2.87	1.74	-2.79	-1.89	-2.85	-1.53
13135-R5-1	-23.43	-23.43	-2.06	1.34	1.63	-1.41	-3.42	1.76	-2.53	-2.77	-3.89	-3.94
13136-L5-3	-6.86	-6.86	-6.86	-6.86	1.35	-2.00	-2.05	1.67	-6.86	1.09	-1.16	-6.86
13137-L5-1	-11.97	-2.58	-1.60	1.04	1.90	1.04	-2.52	3.20	-2.24	-1.54	-2.07	-5.87
13138-R5-1	-5.15	-3.33	-3.23	-1.50	-1.38	-1.56	-5.50	1.32	-4.04	-2.61	-3.13	-4.48
13163-R5-1	-10.02	-3.08	-1.87	2.52	1.68	-1.32	-3.55	1.41	-4.70	-2.94	-4.38	-3.99
13164-L5-1	-5.10	-5.10	-5.10	4.48	4.67	1.58	-1.49	3.96	-5.10	-1.16	-2.30	-5.10
13166-L5-1	-37.14	-3.17	-2.98	-1.24	-1.26	-2.36	-4.64	1.43	-3.21	-3.95	-4.48	-2.43
13181-L5-1	-7.92	-7.92	-7.92	1.65	1.40	2.36	-2.28	5.58	-2.57	-2.04	-1.63	-7.92
13184-L5-1	-2.31	-2.31	-2.31	4.42	1.67	-1.23	1.05	5.81	-2.31	1.90	-1.17	-2.31
13186-R5-2	-26.53	-4.05	-9.42	-1.04	2.59	1.05	-2.83	1.85	-9.17	-2.40	-4.24	1.98
13195-L5-3	-23.57	-2.07	-2.02	1.48	2.78	1.09	-2.90	1.89	-2.32	-2.10	-2.13	-2.45
13199-R5-1	-8.02	-2.26	1.36	1.62	3.08	1.12	-2.13	4.07	-1.08	-1.86	-1.57	-8.02
13202-L5-1	1.16	-10.20	1.03	1.38	1.62	-1.35	-2.12	2.03	-10.20	-1.77	-2.12	-10.20
13202-R5-2	-14.39	-1.40	1.14	3.25	1.30	-1.53	-2.85	1.55	-1.85	-2.70	-3.38	-14.39
13209-L5-2	-45.23	-3.07	-1.52	1.00	-1.08	-1.54	-3.29	1.39	-2.21	-1.88	-2.01	-4.56
13209-R5-3	-12.46	-6.49	1.10	1.57	-1.68	-2.00	-1.94	1.42	-1.36	-1.64	-2.04	-12.46
13211-L5-1	-31.07	-2.60	-1.64	-1.17	-1.32	-1.51	-4.08	1.30	-2.71	-1.84	-2.39	-10.90
13220-L5-3	-6.86	-10.00	-2.35	-1.04	-1.08	-2.44	-4.02	1.22	-2.77	-2.46	-2.77	1.00
13229-L5-1	-29.36	-2.91	-1.85	-1.07	1.36	1.32	-2.81	2.19	-2.21	-1.91	-2.31	-1.02
13230-L5-4	-2.05	-2.05	-1.17	-2.05	3.16	2.18	1.59	6.65	-1.14	2.50	-1.02	-2.05
13231-L5-2	-16.99	-3.66	-1.24	-1.09	-1.39	-1.43	-3.76	1.62	-1.61	-1.25	-2.31	-16.99
13237-L5-4	-35.36	-35.36	-3.55	-1.11	1.31	-1.52	-5.02	1.37	-4.63	-3.23	-4.11	-35.36
13239-L5-2	-39.77	-7.37	-3.11	-1.12	2.53	1.22	-3.59	3.98	-3.01	-2.21	-3.96	-39.77
13240-L5-2	-18.39	-4.67	-2.06	-1.28	-1.62	-1.59	-3.66	1.21	-3.23	-2.56	-2.62	-2.75
13241-L5-2	-10.26	-4.55	-4.91	1.77	1.77	-1.36	-2.23	3.91	-3.60	-1.33	-4.77	-10.26
13251-R5-2	-20.00	-3.70	-3.53	1.00	1.38	-1.26	-4.35	1.58	-4.26	-2.42	-3.46	-2.91
13259-R5-1	-3.29	-3.29	-3.29	-3.29	2.53	1.88	-1.19	-3.29	-3.29	1.28	1.08	-3.29
13267-L5-1	-30.85	-13.05	-1.69	-10.20	1.96	3.35	-1.63	8.62	-12.76	-1.63	-2.04	-9.14
13281-L5-3	1.06	-1.86	-1.81	-1.07	1.11	-1.64	-3.92	1.69	-1.24	-2.19	-3.54	-6.83

13283-L5-3	-13.68	-3.86	-2.82	-1.06	1.13	-1.43	-1.92	2.27	-1.49	-1.44	-1.67	-6.16
13285-L5-3	-11.84	-4.28	-2.47	-1.21	-1.65	-1.98	-4.63	-1.22	-3.72	-2.67	-3.38	-1.41
13287-L5-3	-27.01	-1.81	-1.63	-1.11	2.16	1.08	-3.42	2.88	-2.45	-1.53	-2.25	-1.07
13291-L5-1	-5.33	-8.08	-1.21	1.10	5.56	2.15	-2.09	6.10	-1.56	1.04	-1.83	-5.77
13293-L5-1	-18.63	-18.63	-1.65	1.07	1.04	-1.36	-3.93	2.07	-2.11	-1.84	-2.35	-18.63
13298-R5-1	-14.82	-14.82	-3.48	1.05	1.24	-1.67	-3.79	1.78	-14.82	-2.80	-3.15	-14.82
13303-L5-3	-14.35	-7.42	-1.08	1.87	3.44	1.82	-2.05	3.33	1.08	1.03	-1.75	-14.35
13308-L5-1	-14.43	-1.49	-1.10	1.30	3.61	1.10	-2.16	2.74	-1.69	-1.36	-1.79	-14.43
13310-R5-1	-6.22	-6.22	-6.22	1.86	-1.03	-1.50	-1.67	3.20	1.44	2.11	-1.28	-6.22
13312-L5-2	-14.33	-2.46	-2.40	-1.24	3.55	1.50	-3.72	3.22	-3.06	-2.08	-2.68	-1.53
13313-L5-2	-8.53	-2.34	-1.67	1.70	1.40	-1.33	-2.46	2.18	-2.74	-1.65	-2.66	-4.35
13316-R5-2	-4.30	-4.30	-2.20	2.94	1.96	3.07	-1.31	4.09	1.82	2.30	1.24	-4.30
13326-L5-2	-52.63	-3.83	-2.70	-1.22	1.72	1.01	-3.47	2.34	-3.63	-2.53	-3.07	-1.23
13328-R5-2	-14.53	-14.53	-3.23	1.52	-1.14	-1.34	-2.95	1.56	-1.83	-2.66	-2.68	-14.53
13332-L5-1	-4.39	-4.39	-2.11	2.76	2.77	1.33	-1.10	-1.46	-4.39	1.18	-1.07	-1.98
13334-L5-3	-43.43	-2.48	-4.73	1.14	3.67	1.66	-3.78	3.22	-3.11	-1.94	-2.34	1.24
13335-L5-3	-4.23	-3.07	-1.86	1.27	1.43	-1.36	-3.29	1.75	-2.97	-1.67	-2.59	-1.75
13337-L5-2	-42.63	-15.36	-1.48	-11.45	3.39	1.31	-3.13	2.96	-16.82	-2.38	-3.47	1.92
13339-L5-1	-3.11	-3.86	-2.52	-1.15	-2.17	-2.18	-3.10	-1.68	-3.33	-2.51	-2.85	-1.46
13343-L5-1	-5.44	-2.70	-1.77	1.12	2.31	1.34	-2.98	3.06	-2.38	-1.38	-1.82	-1.09
13349-L5-2	-10.66	-3.95	-2.27	-1.24	-1.33	-2.25	-4.40	1.03	-3.81	-2.98	-3.14	-1.13
13353-L5-2	-2.24	-2.24	-2.24	-2.24	1.70	5.74	1.55	1.27	2.75	2.47	1.48	-2.24
13354-L5-1	-17.63	-10.31	-17.63	1.04	-1.29	-1.42	-4.54	1.52	-3.25	-3.11	-4.57	-17.63
13355-L5-2	-5.51	-5.51	-5.51	1.73	1.33	2.51	-2.60	1.56	-5.51	-2.00	-5.51	-5.51
13356-L5-2	-8.37	-8.37	1.15	-8.37	1.19	-1.16	-1.71	2.41	-2.44	-1.56	-1.28	-8.37
13358-L5-2	-23.01	-4.99	-4.83	1.29	1.72	-1.44	-3.50	1.29	-10.96	-2.15	-3.31	-23.01
13361-L5-1	-12.93	-12.93	-12.93	-1.25	-1.51	-2.03	-3.61	1.42	-1.98	-2.17	-1.96	-12.93
13363-L5-2	-20.98	-20.98	-20.98	-1.40	-1.39	-2.08	-4.73	1.11	-2.88	-2.56	-2.49	-20.98
13364-L5-2	-7.47	-7.47	-3.52	-7.47	-1.24	-2.33	-2.43	3.10	-7.47	-1.78	-1.46	-7.47
13365-L5-3	-7.56	-3.41	-1.76	1.61	1.56	-1.00	-3.01	1.80	-2.88	-1.74	-2.58	-1.78
13370-L5-2	-27.25	-1.67	-1.39	1.09	1.33	-1.16	-2.69	2.19	-2.79	-1.56	-2.12	-1.10
13373-L5-4	-3.08	-7.98	-2.22	1.02	2.43	1.76	-2.02	3.28	-1.55	1.04	-1.31	-7.98
13374-R5-1	-35.28	-7.46	-2.24	-1.53	-1.13	-1.31	-3.05	1.69	-2.75	-2.15	-2.18	-5.68

13375-L5-3	-41.40	-3.21	-2.13	-1.40	-1.08	-1.32	-4.62	1.77	-3.47	-1.85	-2.99	-15.84
13376-R5-1	-46.25	-2.90	-3.40	1.26	1.17	-1.77	-3.98	1.62	-5.25	-4.65	-5.47	-46.25
13380-R5-3	-10.07	-10.07	-1.12	1.07	-1.25	-1.49	-2.40	1.95	-10.07	-2.00	-1.88	-10.07
13385-L5-1	-3.98	-3.98	-3.98	2.74	3.39	3.77	-1.37	-1.25	-3.98	1.45	1.01	-3.98
13396-L5-2	-51.56	-3.50	-3.03	-1.02	1.19	-1.18	-3.70	2.12	-3.31	-2.34	-2.84	-19.22
13403-L5-1	-21.42	-1.77	-1.59	2.39	2.56	-1.10	-2.29	1.93	-2.00	-2.82	-3.69	-21.42
13412-L5-1	-8.32	-8.32	1.16	1.31	-1.47	-1.43	-2.06	1.69	-8.32	-1.83	-1.40	-8.32
13423-L5-3	-9.37	-2.67	-1.70	2.74	1.65	-1.49	-3.52	1.30	-3.67	-2.44	-3.89	-4.22
13425-L5-3	-8.40	-2.50	-1.76	3.00	1.76	-1.51	-3.88	1.20	-4.15	-2.68	-4.80	-4.43
13430-L5-3	-3.90	-3.61	-2.27	-1.01	3.74	1.98	-3.74	5.44	-1.77	-2.17	-3.43	-18.09
13431-L5-3	-4.88	-6.00	-2.06	1.01	-2.24	-3.42	-4.62	-1.58	-3.62	-3.19	-4.73	-1.29
13432-R5-1	-17.04	-1.33	1.36	3.21	2.17	1.38	-2.00	2.50	1.55	-1.06	-2.15	-17.04
13456-R5-2	-14.18	-14.18	2.87	-6.32	1.05	1.63	-2.07	5.81	-4.94	-1.19	-1.58	-14.18
13458-R5-2	-9.56	-9.56	1.53	-4.22	4.25	-1.29	-1.67	10.75	-3.73	-1.02	-1.10	-9.56
13461-L5-4	-70.04	-4.10	-2.14	-1.39	1.18	-1.05	-3.94	2.59	-2.51	-2.43	-2.91	-2.77
13463-L5-2	-18.40	-3.50	-2.21	1.89	1.41	-1.11	-3.17	2.39	-3.17	-2.66	-4.28	-25.56
13489-L5-1	-2.32	-2.32	4.10	7.21	1.60	1.78	-1.10	-2.32	1.73	-2.32	-2.32	-2.32
13497-L5-1	-7.23	-4.09	-3.08	1.16	1.64	1.09	-4.05	1.93	-4.13	-2.92	-4.09	-3.23
13513-L5-1	-3.48	-3.48	4.44	-3.48	1.38	2.30	1.03	4.31	2.43	1.49	2.28	-3.48
13519-L5-1	-8.34	-8.34	-4.66	1.28	1.53	-1.40	-2.94	2.21	-8.34	-2.08	-1.85	-8.34
13525-L5-2	-11.76	-11.76	-2.97	1.16	1.18	-1.56	-3.63	1.43	-11.76	-1.93	-2.26	-4.62
25-R5-1	-13.46	-13.46	1.75	1.51	3.69	1.86	-1.81	3.93	-1.29	-1.53	-2.27	-13.46
266-R5-2	-5.40	-3.19	-2.35	1.20	3.10	1.60	-3.19	3.09	-2.31	-1.95	-3.02	-3.07
2786-L5-3	-7.14	-7.14	-1.80	1.48	1.86	-1.31	-2.37	2.47	-7.14	-1.82	-1.48	-7.14
2811-R5-1	-5.19	-5.19	-5.19	-5.19	1.07	-1.12	-1.90	2.46	-5.19	-1.53	-1.19	-5.19
2819-R5-4	-5.51	-5.80	-2.51	-1.25	-2.51	-3.80	-5.86	-1.79	-4.40	-3.96	-6.14	-1.09
3717-L5-2	-10.28	-3.32	-2.43	1.83	1.05	1.36	-5.47	2.60	-5.09	-3.62	-5.62	-1.15
3732-R5-1	-13.70	-1.52	-1.09	1.37	1.34	1.51	-2.32	-1.64	-1.65	-1.02	-2.09	-4.30
3799-R5-1	-6.12	-2.70	-1.95	1.54	1.37	1.25	-2.90	2.68	-2.69	-1.77	-2.48	-8.74
3897-R5-2	-8.76	-5.74	-2.75	-1.70	-1.02	1.56	-3.91	1.41	-3.43	-2.66	-2.84	-2.33
3942-L5-1	-4.60	-1.85	-1.50	2.56	2.76	1.21	-2.11	2.61	-2.13	-2.30	-3.87	-53.73
3952-L3-1	-7.09	-1.94	-1.94	1.81	-1.05	1.07	-2.42	1.66	-7.09	-2.27	-1.92	-7.09
3953-R3-2	-7.36	-4.86	-2.14	-1.41	-1.64	-1.73	-3.38	1.20	-3.51	-2.26	-3.09	-1.13

3966-L5-1	-9.29	-5.96	-2.17	-1.13	-1.86	-2.86	-3.59	-1.09	-3.21	-2.58	-3.27	-2.45
3995-L2-2	-25.12	-2.66	2.34	1.82	2.48	1.95	-2.10	4.87	4.02	-1.04	-1.85	-25.12
4013-L4-1	-2.19	-1.14	-1.19	5.14	1.52	-1.25	-2.19	-2.19	-2.19	-2.19	-2.19	-2.19
4026-R5-1	-5.56	-3.29	-2.88	1.85	1.58	-1.41	-3.53	-81.69	-5.46	-4.37	-5.89	-81.69
4026-R5-2	-7.79	-3.53	-2.44	2.12	1.69	-1.02	-3.26	1.98	-4.14	-3.33	-5.34	-145.60
410-R5-1	-5.18	-5.18	-5.18	-5.18	2.41	1.08	-1.67	4.07	-5.18	-1.60	-1.13	-5.18
4130-L5-1	-18.42	-18.42	-4.29	-1.07	-2.00	-3.97	-4.12	-1.60	-18.42	-3.45	-3.62	1.01
4143-R5-2	-3.75	-3.75	-3.75	-3.75	3.84	1.77	-1.16	6.22	-3.75	1.61	1.47	-3.75
4258-L5-1	-2.02	-2.02	-2.02	-2.02	5.96	3.36	1.46	8.78	-2.02	2.12	2.15	-2.02
4315_C-L4-1	-26.61	-2.74	-1.68	1.24	1.19	1.14	-4.07	-84.00	-3.03	-1.71	-2.46	-2.12
4315_D-R4-1	-52.47	-4.05	-1.59	-1.06	-1.09	-1.34	-4.58	1.76	-3.29	-2.93	-3.35	-16.76
4315_E-R4-1	-14.07	-3.05	1.21	-1.02	1.93	1.81	-2.20	4.06	-1.31	-1.04	1.16	-2.25
4315_F-R4-1	-6.50	-3.14	-1.75	-1.35	1.89	1.09	-3.03	2.07	-2.72	-1.88	-2.15	-2.00
4315_I-L4-1	-24.12	-2.69	-1.62	-1.14	1.34	1.53	-3.47	3.10	-2.18	-1.55	-1.78	-24.12
4315_K-L4-1	-11.95	-3.76	-1.84	-1.67	1.11	-1.40	-5.02	1.91	-2.86	-2.79	-3.01	-1.07
4315-R3-2	-18.48	-18.48	-1.57	-1.08	1.94	1.17	-2.31	3.55	-1.46	-1.77	-1.94	-18.48
4338-L5-2	-2.07	-1.04	-2.07	6.34	-1.05	2.04	1.12	-2.07	3.52	-1.27	-2.07	-2.07
4340-R3-1	-5.28	-5.28	-2.74	1.96	1.91	1.22	-1.42	3.91	-5.28	-1.09	-1.05	-5.28
4346-L5-1	-4.26	-3.59	-2.40	-1.31	1.61	1.08	-2.87	2.65	-2.47	-1.51	-1.60	-2.44
4361-R5-1	-16.06	-3.21	-2.18	1.15	2.56	1.70	-1.98	4.81	-16.06	-1.30	-1.94	-5.13
4498-L3-2	-9.82	-5.38	1.41	1.75	1.02	-1.42	-1.96	2.19	-1.12	-1.31	-1.02	-9.82
4516-L5-1	-14.82	-6.92	-14.82	1.39	1.36	-1.61	-3.67	1.30	-14.82	-3.07	-3.34	-14.82
454-R5-1	-2.62	-2.62	-2.62	-2.62	3.88	1.74	1.02	6.55	-2.62	1.18	-1.32	-2.62
4593-R5-1	-38.50	-3.52	-1.40	-1.57	1.21	3.25	-2.72	7.52	-2.16	-1.91	-2.36	-38.50
4610-R5-1	-7.69	-7.69	-4.23	-7.69	1.05	-1.38	-2.23	2.29	-7.69	-1.85	-1.57	-7.69
4610-R5-2	-3.23	-3.23	-1.64	-3.23	4.54	1.72	1.06	6.67	-3.23	1.85	1.59	-3.23
4642-R3-2	-10.25	-10.25	-5.37	1.32	-1.09	-1.37	-2.03	1.75	-10.25	-1.52	-1.92	-10.25
4666-R5-1	-52.31	-8.64	-2.97	-1.02	3.07	1.24	-3.80	4.13	-2.83	-3.00	-4.54	-20.59
4792-L5-2	-8.62	-2.62	-1.48	1.58	1.24	-1.03	-2.41	3.62	-1.95	-1.54	-2.22	-24.82
4801-L5-2	-4.53	-4.53	-2.30	2.06	1.80	1.04	-1.18	2.99	-4.53	1.18	1.17	-4.53
4813-R3-1	-3.33	-3.33	-3.33	1.39	1.45	-1.77	-1.46	-3.33	-3.33	-1.07	-3.33	-1.06
4875-R2-2	-25.83	-2.14	-1.33	1.90	1.88	1.16	-2.27	2.53	-1.98	-1.34	-2.73	-25.83
4912-L5-2	-19.46	-19.46	-1.91	-1.29	-1.45	-1.96	-2.67	1.56	-2.21	-2.67	-2.47	-19.46

4929-R4-1	-11.74	-11.74	-11.74	1.31	-2.37	-2.86	-4.16	-1.03	-11.74	-3.58	-2.44	-11.74
5032-R5-1	-13.13	-13.13	-6.46	-1.03	1.10	-1.14	-2.75	2.43	-13.13	-1.83	-1.91	-13.13
5048-L5-1	-11.59	-11.59	-1.39	1.05	-1.30	-1.54	-3.38	1.08	-6.02	-1.58	-2.26	-11.59
5071-R5-2	-7.99	-3.93	-3.06	-1.63	-1.09	-1.22	-4.17	1.01	-3.96	-1.64	-2.24	-3.88
5107-L5-1	-26.66	-1.96	-1.85	1.79	1.96	1.08	-3.02	3.02	-1.40	-2.20	-3.10	-26.66
5210-L5-1	-7.99	-7.99	1.22	1.47	-1.11	-1.13	-2.48	1.72	-7.99	-1.31	-2.28	-7.99
5342-L5-1	-1.33	-1.69	-1.29	1.18	-1.19	-2.25	-2.81	1.02	-1.76	-2.17	-2.61	-6.40
5491-R5-1	-8.97	-3.43	-1.87	1.26	-2.01	-3.08	-3.05	-1.22	-2.81	-3.09	-4.36	1.32
5521-L5-2	-7.60	-7.60	1.16	3.47	2.17	-1.21	-1.91	2.12	-7.60	-1.57	-1.79	-7.60
554-R5-1	-8.55	-2.59	-1.73	2.95	1.77	-1.55	-3.52	1.23	-4.27	-2.77	-4.42	-5.30
5554-R5-2	-28.10	-1.55	-1.48	1.18	1.34	-1.27	-3.63	1.83	-2.03	-2.07	-3.04	-2.37
5638-R5-2	-8.67	-2.30	-2.11	2.48	1.70	-1.46	-4.40	1.36	-3.47	-2.78	-4.60	-4.20
5640-L3-1	-11.16	-2.70	-1.45	1.39	1.59	1.12	-3.47	2.10	-2.10	-1.71	-2.50	-35.95
5749-R5-1	-3.50	-3.50	-1.65	7.88	3.53	2.22	1.19	-3.50	-3.50	-1.07	-3.50	-3.50
5757-L5-1	-22.78	-22.78	-4.13	1.16	-2.40	-6.68	-3.91	-22.78	-22.78	-4.48	-4.14	1.71
5854-R5-2	-8.37	-2.42	-1.59	3.10	1.90	-1.20	-2.79	1.86	-3.32	-2.33	-3.25	-4.60
5956-L5-1	-1.50	-1.19	-8.83	-3.12	1.38	1.74	-1.28	-1.96	-8.83	-1.21	-3.33	1.37
5995-R5-1	-2.29	-2.29	-2.29	-2.29	3.85	1.13	-2.29	4.79	-2.29	-1.51	-2.29	-2.29
6008-R5-1	-9.71	-9.71	-9.71	1.21	-1.27	-2.19	-3.80	1.01	-9.71	-2.70	-5.17	-9.71
6008-R5-2	-5.41	-5.41	-2.83	1.90	1.17	-5.41	-2.24	-5.41	-5.41	-1.58	-1.86	-5.41
6016-R2-1	-12.02	-12.02	-2.83	-1.03	1.09	-1.29	-2.43	2.59	-1.53	-1.37	-1.49	-12.02
6023-L5-1	-19.11	-4.43	-9.06	1.64	-1.55	-1.76	-4.04	1.19	-2.84	-3.49	-3.47	-19.11
6087-L4-1	-8.97	-8.97	-4.56	-8.97	1.20	-1.22	-2.10	1.95	-8.97	-1.82	-2.11	-8.97
6096-R5-1	-12.65	-12.65	-6.35	-12.65	-2.12	1.96	-4.37	3.74	-12.65	-3.78	-3.17	-12.65
6192-L5-1	-13.47	-13.47	2.69	2.33	1.51	-1.93	-1.74	4.14	-4.25	-1.11	-1.06	-13.47
6198-R5-2	-8.02	-4.03	-1.28	-1.07	-1.73	-2.33	-2.25	1.04	-2.46	-1.83	-2.17	-2.31
6242-R5-1	-5.90	-5.90	-5.90	-2.78	1.99	1.41	-2.49	3.61	-5.90	-2.04	-3.28	-5.90
6287-L3-2	-19.21	-10.03	-2.08	1.30	1.40	-1.38	-3.39	2.27	-2.02	-2.10	-2.68	-19.21
6385-R5-2	-9.70	-9.70	-2.46	1.00	-1.10	-1.39	-2.34	2.61	-9.70	-1.58	-1.52	-9.70
6409-L3-1	-2.84	-3.81	-1.89	-1.05	-1.55	-3.11	-4.36	-1.21	-2.13	-3.04	-2.46	-3.30
6434-R5-1	-4.10	-2.03	-1.40	3.14	1.59	-1.08	-3.58	2.01	-2.31	-2.55	-4.04	-2.74
6490-R5-3	-23.58	-2.43	-2.36	-1.08	2.40	-1.34	-2.61	3.20	-2.47	-1.71	-2.00	-4.00
6496-R5-2	-44.85	-3.00	-3.09	-1.17	4.58	2.18	-3.23	4.50	-2.86	-1.62	-2.20	-1.51

6584-L5-1	-13.67	-13.67	-5.86	1.18	3.62	2.49	-1.83	6.84	-13.67	-1.23	-1.55	-13.67
6590-L5-1	-4.65	-4.65	-1.25	2.00	2.52	1.66	-1.27	6.29	-2.59	1.13	-1.09	-4.65
6642-R5-1	-9.51	-1.99	-1.73	1.59	3.00	-1.02	-2.64	3.60	-1.98	-1.60	-2.83	-1.73
669-R5-2	-10.57	-10.57	-2.81	1.09	-1.32	1.39	-3.80	3.58	-1.86	-2.58	-2.81	-10.57
6718-L3-2	-3.98	-3.98	-2.18	2.26	1.29	-2.03	-1.58	-1.57	-3.98	1.66	-3.98	-3.98
6839-L3-1	-27.98	-2.76	-2.51	-1.37	1.05	-1.33	-4.23	2.43	-2.96	-2.29	-2.78	-1.52
6880-L3-2	-5.32	-2.54	-1.25	1.77	2.00	1.81	-2.74	3.05	-1.32	-1.35	-2.78	-2.32
6908-L3-2	-2.84	-2.84	3.97	6.90	1.33	1.57	-1.04	-2.84	3.32	-1.05	-2.84	-2.84
6984-R4-1	-18.59	-5.09	-1.67	1.15	3.57	1.76	-3.55	5.19	-1.99	-2.34	-3.28	-18.59
7029-R5-1	-10.64	-10.64	-2.56	1.54	1.33	-1.01	-2.23	3.16	-10.64	-1.40	-1.44	-10.64
7061-R5-2	-51.92	-1.69	-1.36	-1.17	1.39	1.10	-3.67	1.94	-2.85	-1.51	-2.19	-1.73
7066-R5-1	-31.19	-3.44	-2.16	1.08	-1.50	-1.75	-4.12	1.47	-3.39	-3.33	-4.27	-31.19
7069-R5-1	-42.68	-3.22	-2.52	1.02	1.34	-1.32	-3.98	1.59	-3.33	-2.47	-3.14	-18.77
7113-R5-1	-8.52	-4.48	-3.59	1.23	1.60	-1.25	-3.50	1.53	-5.21	-3.87	-5.23	-49.48
7126-L3-1	-21.53	-3.22	1.09	-1.52	1.21	1.28	-2.18	2.92	-1.66	-8.94	-3.88	-2.59
7141-R5-1	-2.31	-2.31	-2.31	-2.31	4.83	1.97	-1.07	5.66	-2.31	1.32	-2.31	-2.31
7221-R5-1	-11.70	-6.14	-1.36	-1.00	-1.61	-1.41	-2.93	1.25	-1.75	-1.65	-2.19	-11.70
7313-L5-2	-2.26	-2.26	-2.26	-2.26	2.68	2.29	1.59	5.74	-2.26	2.36	2.07	-2.26
7352-R3-2	-11.09	-5.02	1.09	1.44	1.44	1.64	-1.96	2.77	-1.31	1.60	-1.15	-11.09
7356_A-R4-1	-7.06	-5.54	-1.95	-1.05	-1.74	-2.76	-4.73	-1.20	-3.41	-3.22	-3.51	-1.25
7356-L5-1	-17.42	-17.42	-1.11	1.16	1.37	-1.26	-3.38	2.37	-1.70	-1.56	-2.28	-17.42
7367-L1-1	-5.61	-5.61	-2.76	2.85	2.74	1.81	-1.31	3.71	-5.61	-1.01	-1.26	-5.61
7384-R3-1	-16.46	-3.05	-1.45	1.34	3.85	1.12	-2.15	2.45	-1.99	-1.70	-2.75	-1.96
7411-R3-2	-13.01	-2.66	-1.16	1.17	1.26	1.15	-2.46	2.04	-1.61	1.08	-1.57	-13.01
7569-L5-2	-3.93	-3.93	-3.93	-3.93	-1.05	-1.16	-1.50	2.37	-3.93	-1.26	-3.93	-3.93
7571-L5-1	-32.04	-3.20	1.02	-1.14	1.30	-1.37	-2.52	2.12	-2.59	-1.83	-3.22	-1.07
7572-R5-2	-22.49	-2.34	-1.93	-1.03	6.19	2.41	-2.23	5.40	-2.12	-1.30	-1.54	-1.17
7660-L5-1	-18.40	-4.72	-1.14	1.08	1.23	-1.13	-2.11	2.32	-1.23	-1.17	-1.37	-18.40
7702-L2-1	-33.45	-17.84	-3.01	-1.64	1.18	-1.56	-5.68	1.73	-3.58	-3.35	-3.12	-14.37
7736-L5-1	-10.35	-10.35	1.09	2.15	1.62	1.20	-1.65	4.30	1.24	1.39	-1.23	-10.35
7743-L5-1	-7.13	-7.13	-7.13	-7.13	-1.31	-1.50	-2.01	2.34	-1.02	-1.50	-1.49	-7.13
7781-R5-2	-7.71	-5.45	-1.96	1.05	-1.75	-3.27	-3.53	-1.42	-2.53	-2.75	-2.78	-22.71
7824-R5-1	-2.36	-12.29	-1.52	-1.54	-2.01	-3.22	-4.60	-1.27	-24.95	-3.64	-4.64	-1.19

7846-L5-2	1.10	-2.66	-2.66	-2.66	5.00	2.15	1.24	4.64	-2.66	2.08	1.99	-1.18
7883-R5-1	-4.16	-2.58	-1.80	-1.08	3.79	1.59	-2.34	3.80	-1.95	-1.40	-1.82	-1.00
78-R4-1	-2.39	-2.39	-1.21	6.26	2.28	1.91	-1.02	1.00	3.43	1.26	-2.39	-2.39
7949-R5-1	-4.01	-4.36	-1.30	1.15	-2.21	-2.42	-2.64	1.16	-2.75	-2.23	-3.15	1.33
7971-L5-1	-7.45	-7.45	-1.61	-7.45	1.29	-1.04	-1.90	2.29	-7.45	1.31	-1.06	-7.45
8016-L3-1	-13.40	-1.65	1.05	1.08	1.03	2.34	-2.26	3.25	-1.63	-1.53	-2.21	-13.40
8062-R5-1	-7.61	-3.70	-1.70	2.18	2.59	1.56	-2.32	2.34	-7.61	1.60	-1.20	-7.61
8077-R3-1	-10.66	-10.66	-1.94	1.52	1.52	2.13	-2.53	4.22	-10.66	-1.39	-1.79	-10.66
8089-L5-1	-12.58	-12.58	-12.58	1.14	1.34	-1.19	-3.58	1.73	-12.58	-2.50	-2.89	-12.58
8239-R5-1	-7.85	-7.85	-1.71	1.38	3.30	1.25	-1.74	4.30	-7.85	-1.16	-1.28	-7.85
8250-R5-2	-8.86	-6.17	-3.27	-1.52	1.20	-1.96	-3.94	1.31	-3.73	-2.78	-2.56	-1.90
8281-L5-2	-4.41	-1.86	-1.49	2.18	1.25	-1.11	-2.92	2.00	-2.11	-1.90	-2.99	-5.52
8298-R5-1	-2.06	-3.00	-4.52	-1.25	1.18	-1.68	-4.67	1.41	-2.94	-3.38	-3.35	-2.74
8329-L5-1	-7.14	-7.14	-7.14	-1.04	1.64	-4.12	-1.88	-2.09	-7.14	-1.69	-1.50	-7.14
8336-R5-2	-7.05	-5.15	-1.26	1.44	1.78	-1.12	-2.29	2.53	-1.47	-1.45	-2.17	-26.66
8394-L5-2	-33.24	-2.09	-1.40	-1.37	1.26	1.30	-2.51	2.74	-2.20	-1.19	-1.38	-1.20
8564-L5-1	-9.90	-9.90	-9.90	-1.75	2.20	-1.21	-2.57	3.01	-9.90	-2.15	-2.29	-3.34
8564-R5-2	3.57	-3.01	2.65	4.26	-1.18	-1.95	-1.65	-6.82	-1.16	-1.87	-3.25	-6.82
8898-R5-1	-15.47	-9.30	-2.03	-1.44	1.04	-1.40	-4.53	1.61	-2.95	-1.90	-2.58	-1.63
9021-L5-2	-9.46	-4.89	-1.02	1.42	1.38	1.13	-1.84	4.21	-1.28	-1.21	-2.15	-9.46
9068-R5-2	-10.19	-2.29	-1.93	2.59	1.47	-1.63	-3.56	1.03	-4.24	-2.77	-4.29	-2.77
9087-L5-2	-5.65	-2.03	-2.17	1.21	-1.62	-4.32	-4.61	-1.82	-3.78	-3.28	-6.39	1.58
9134-R5-1	-2.19	1.85	-1.06	5.55	1.62	1.08	1.58	-2.19	-2.19	1.28	-2.19	-2.19
9217-L3-2	-25.69	-2.49	-2.14	-1.16	-1.07	1.71	-3.81	1.45	-2.41	-2.14	-2.41	-11.60
9245-R5-1	-7.70	-7.70	-7.70	-7.70	1.21	-1.21	-1.48	-7.70	1.20	-1.58	-1.57	-7.70
9287-L5-2	-8.66	-10.47	-2.11	1.95	1.22	1.28	-5.10	2.39	-5.82	-4.19	-6.03	-2.02
9369-L5-1	-2.53	2.54	3.26	5.80	-1.36	1.32	-1.20	-2.53	3.39	-2.53	-2.53	-2.53
9384-R5-2	-15.00	-15.00	-3.18	-1.02	-1.40	-1.71	-4.41	1.20	-4.03	-2.13	-2.58	-15.00
9387-R2-2	-2.27	-11.24	1.24	-1.66	-1.77	-2.13	-2.67	1.46	-18.94	-3.05	-3.88	-1.22
9564-R5-2	-21.32	-1.78	-1.27	1.30	-1.36	-1.75	-2.96	1.23	-1.50	-1.51	-2.02	-8.68
9605-R5-1	-2.11	-2.11	2.05	9.59	3.34	1.71	1.35	1.73	-2.11	2.19	1.04	-2.11
9691-L5-1	-13.91	-7.43	-3.19	-1.18	1.46	-1.16	-1.81	3.62	-1.45	-1.71	-1.56	-13.91
9770-R5-2	-16.08	-16.08	-1.38	-6.49	-1.54	-1.04	-3.18	2.79	-16.08	-2.24	-1.93	-16.08

9774-R2-2	-20.32	-5.37	-3.62	-1.47	-1.04	-2.24	-5.14	1.19	-4.78	-2.77	-3.92	-3.06
9812-L3-1	-8.36	-8.36	1.41	-3.81	-1.10	-1.30	-1.61	2.18	-8.36	-1.63	-1.54	-8.36
9866-L5-1	-11.22	-11.22	-11.22	-11.22	-1.94	-4.41	-3.65	1.13	-11.22	-2.98	-2.07	-11.22
999997-R4-1	-40.37	-3.22	-2.37	1.31	-1.44	1.32	-5.01	3.93	-3.66	-3.76	-5.46	-40.37
let-7b	-4.92	1.02	2.03	1.30	-4.60	-2.75	-2.04	-2.69	-1.28	-1.09	-2.68	1.40
let-7c	-6.31	-1.05	1.94	1.37	-4.31	-2.44	-1.65	-3.04	-1.21	-1.03	-2.43	1.67
let-7e	-22.08	1.03	2.47	1.47	-4.26	-2.47	-1.58	-2.23	-1.12	-1.04	-2.53	2.01
miR-100	-15.70	-1.06	2.47	1.30	-1.44	-2.47	-2.17	-1.90	-1.30	-1.64	-1.79	-15.70
miR-101	-12.44	-1.03	1.43	1.63	-3.70	-2.97	-2.13	-12.44	-1.45	-1.11	-1.92	1.97
miR-1182	-12.48	-2.69	1.05	1.43	2.25	-1.41	-1.78	-1.07	-12.48	-1.46	-2.26	2.67
miR-1207-5p	-7.81	-1.85	-1.68	1.60	1.80	-1.18	-2.41	2.42	-1.87	-1.78	-2.38	-2.36
miR-1224-5p	-8.27	-4.01	1.20	1.73	3.06	-1.10	-2.00	2.05	-8.27	-1.44	-1.68	-8.27
miR-1225-5p	-32.61	-3.21	-16.64	1.04	1.24	-1.74	-5.01	1.53	-3.91	-5.58	-5.48	-32.61
miR-1228*	-9.13	-2.28	-2.02	-1.49	1.07	1.35	-3.26	2.11	-3.56	-1.07	-1.87	-3.42
miR-1234	-2.04	3.94	4.09	6.04	1.42	1.35	-2.04	-2.04	-2.04	-2.04	-2.04	-2.04
miR-125a-5p	-12.05	-1.57	2.48	-1.11	-2.90	-3.80	-2.32	-36.39	-1.43	-2.16	-3.93	1.84
miR-126	-41.23	-1.60	-1.58	1.20	-9.24	-16.81	-5.80	-17.32	-3.33	-5.96	-5.54	-2.23
miR-1268	-10.67	-10.67	-10.67	1.26	1.67	-1.01	-1.65	4.86	-1.28	1.79	-1.47	-10.67
miR-130b	-2.21	-2.21	5.25	5.24	-1.32	-2.21	1.17	-2.21	-2.21	1.98	1.45	-2.21
miR-140-3p	-4.44	-4.44	1.97	2.44	-1.38	-2.55	-1.53	-4.44	-1.28	-1.52	-2.25	-4.44
miR-145	-31.48	-1.89	1.24	-1.28	-2.10	-1.49	-2.04	-13.08	1.01	-3.26	-2.80	1.20
miR-149*	-61.43	-1.79	-1.40	1.02	1.46	1.72	-2.47	3.20	-2.22	1.19	-1.22	-2.53
miR-150	-3.14	-3.14	3.09	4.65	1.17	-1.40	-1.27	-3.14	-3.14	2.18	-3.14	-3.14
miR-181b	-2.73	-2.73	4.73	-1.14	1.39	-1.40	-1.08	-2.73	-1.49	1.49	-1.51	-2.73
miR-181d	-2.47	-2.47	1.60	3.76	1.17	1.29	-2.47	-2.47	-2.47	1.49	1.28	-2.47
miR-185*	-10.19	-10.19	1.73	-3.97	-1.15	-2.49	-2.41	-10.19	-10.19	-1.56	-2.19	-10.19
miR-214	-4.43	-4.43	2.00	2.45	-1.27	1.26	-1.06	-4.43	1.76	-1.08	-2.47	-4.43
miR-23a*	-3.92	-3.92	-3.92	-3.92	2.27	1.12	1.28	-3.92	-3.92	-1.18	-3.92	-3.92
miR-30a	-19.76	-1.04	-1.44	1.58	-3.89	-3.05	-4.56	-7.27	-1.86	-1.24	-2.08	5.69
miR-30d	-1.08	-16.33	1.01	-5.75	-3.77	-3.17	-4.86	-16.33	-7.71	-1.17	-2.19	12.09
miR-320a	-3.29	-2.05	1.23	1.49	5.76	2.18	-1.27	-9.70	1.23	1.59	-1.65	-3.76
miR-320b	-10.09	-2.35	1.00	1.90	5.22	2.10	-1.35	4.46	-1.05	1.28	-1.58	-10.09
miR-335	-3.24	-3.24	1.22	3.13	-3.24	-3.24	-1.26	-3.24	-3.24	5.52	1.37	-3.24

miR-34a	-2.08	-2.08	5.16	-2.08	-1.23	-2.08	-2.08	1.19	-2.08	-2.08	-2.08	-1.15	1.18
miR-34b*	-3.29	-3.29	5.13	-3.29	-3.29	-3.29	-3.29	-1.55	-3.29	-3.29	-3.29	-1.06	-3.29
miR-34c-5p	-2.54	-2.54	5.54	-2.54	-2.54	-2.54	-2.54	-2.54	-2.54	-2.54	-2.54	-2.54	-2.54
miR-371-5p	-3.87	-3.87	-1.81	-1.51	1.98	1.65	1.06	-1.03	-1.06	-3.87	1.10	-1.83	-3.87
miR-373*	-34.48	-13.72	-1.65	-8.74	1.28	2.21	3.58	-3.72	5.31	-16.32	-2.81	-4.50	-34.48
miR-451	-14.34	-2.32	-4.25	-2.09	-10.15	-7.99	-3.72	-9.47	1.72	-9.88	-7.90	-2.86	-3.23
miR-486-3p	-10.16	-10.16	-2.27	1.10	-1.59	-1.36	-3.64	2.23	-2.34	-10.16	-2.14	-1.92	-10.16
miR-491-3p	-2.34	-2.34	-2.34	2.06	1.33	-2.34	2.23	2.23	-2.34	-2.34	1.67	-2.34	-2.34
miR-498	-28.12	-3.04	-1.96	1.11	-1.41	-2.01	-3.75	-3.75	1.36	-3.41	-2.64	-4.35	-28.12
miR-557	-10.00	-1.36	-1.92	1.93	1.16	-1.55	-2.17	1.60	1.60	-1.32	-2.10	-2.21	-1.67
miR-638	-5.13	-5.73	-2.92	-1.37	-1.87	-2.89	-5.04	-1.34	-1.34	-4.22	-3.54	-4.11	-1.97
miR-663	-15.30	-15.30	-2.65	-1.03	-1.31	-1.69	-3.07	1.69	1.69	-1.47	-1.68	-1.53	-15.30
miR-671-5p	-9.36	-9.36	1.12	2.50	1.56	1.06	-2.25	2.06	2.06	-1.39	-1.18	-2.06	-9.36
miR-744	-14.84	-14.84	-3.03	1.02	-1.26	-1.33	-3.71	2.36	2.36	-1.77	-2.34	-2.27	-14.84
miR-885-3p	-26.07	-26.07	-2.17	-1.16	-1.96	-2.54	-4.68	-1.16	-1.16	-3.53	-3.27	-3.46	-10.53
miR-92a-2*	-8.53	-2.11	1.23	1.61	1.34	1.10	-2.40	2.65	2.65	-1.15	-1.39	-1.82	-8.53
miR-92b*	-11.88	-11.88	-1.16	1.21	1.35	1.19	-2.46	3.28	3.28	-1.42	-1.41	-1.46	-11.88
miR-98	-45.32	1.35	2.78	1.82	-4.33	-2.12	-1.37	-9.42	1.07	1.07	1.08	-2.69	1.84
miR-99a	-10.35	1.22	3.06	1.55	-1.32	-4.39	-1.67	-10.35	-1.21	-1.21	-1.78	-2.15	-10.35

Table 25: Pre-microRNA sequences and chromosomal locations of target RNAs in Table 23 and 24

Gene	Chrom loc'n	Pre-microRNA sequence										SEQ ID NO
10010_B-L4-1	7q32.1	GGCAGCGCCCTCCTCCGGAGGGCAGCATCCCGCGGGCGGGGCTCGGCTTTGATGCCAGGGCACCTTTGTCCCTGGAGACGCTCTGCCAGCCAGGTGCGTGAGGGAGTGCAGCCC										1708
10010_D-L4-1	7q32.1	CAGGTGAGAGGCTGAGCCCCCGGGCGGGAGGGCGCCAGGCCCTGGGGCATTACCGTCCCGGGACCCCTTTTGGCCTGCCGGCGCCCCGAGAGCCGCCCTCCCTCCCTCCGTGACAGGTGGGCCTGGAGTTGGGAAAGTTTGAGCCCGCGAGGGGC										1709
10010-R2-2	7q32.1,17q23.2	GCCGGG										1710
10030-R5-1	10q24.1	GGATGCAACCGTGGAAGCCGGTGCCGTTGAGGATCTGCCACAGCGGAAGGCAGCTGAGTTGACATCCACGGGCATCC										1711
10145-L5-2	5p15.33	GGTCCAGATAACAGAAGAGAGAGCAAAAGAAATTTTTGAAGATCAAAAGTGCGCTGTTCAATTTGTTATCTGACCGCGGCTGCGGAGGCTGGCGGGCTGCTGCACCTTTAACGCTTCTGCGCTGACAGGCGGCGCCAGCTAAAGTTCACAGC										1712
10231-R3-1	9p11.2	GCC										1713

10260-L5-2	22q13.1	GTGGGAGACATCGGGTGGGGCCCTGGCGAACAAATAGGTGGGCCCAGCTGGGGCCCCCTCCTGCCTGCCTCACCGC	1714
10333-L5-1	11q12.2	CGGGCAGGGAGGGGGTGGGCAGGGCACAAAGCCTCCCACTGTCCGTGTCCCAACCTCCCCCGTTCCCGG	1715
10342-R2-2	19q12	CCCACGCACGAGGGTCGCCAGGAAAGTGGACATTACCGCTTTAATTAACCTTCGAGATGCTCCGGCGGGGG	1716
10345-R5-1	20p12.3	ACCTCAAAGCTCTCAAGTGGTGACCAACCGTCTTCCGGCCAGGTAGGACTGGATGGAGGGGTGCGCTTTTGAGAGAAAGGT	1717
10374-R3-2	16q12.1	CAGGGGATTTGTACCGCTGATGTGTGGCCCGTCCGAATGAAGGGGCTTTTCATTAACAAAGTAGCGGGCGGTGTCATCTTCCC	1718
10435-R5-1	5q35.1	GAGGCTGCTTAATGAGGTGCCCTTTTCAAAATGTCATCTTAATCTTTTATTAGTTTAAGAAAAGACAACAGGGGCACAATTAGCATGCAA	1719
10533-R5-2	20q12	TAATTGCCCTGAATCGCCGGGTACATATCTGTAGGAAATCTCTTGGCAATATAAAGAAAGGGGCTCAGGACAGTTA	1720
10543-R5-2	1p31.1	GCCTTAACCTTTTTATCATTTTATCTTCTTGATTAATGTCACCTGAATTAATTAATTCATGAGCCAGGATGGGAAGGTGAAGGC	1721
10578-R5-1	1p22.3	CCGCCAGCTTTGTAGCGGTTCCCTGCTTACAAAAGGGCTCTTCTTGAAACGGGGCTGGTGGGACCGTAGAGGGGTGG	1722
10818-L5-1	8q24.3	GGCGGCTGACAGCGGTCAATTGTCACTACTGAGCTGCCCAAACCTCCTCAGACTGCACCTGCGGATGGCTGCTTAGGGTGACTTATG	1723
11370-L5-5	12q13.2	GCCCTGTCCGGCTGCC	1724
11605-L5-4	1p31.3	GTCCAGTTCTCAGGGACAATACTGATGGCAGCCAAACTGGGCAAGGATGCAGTGTGGGGCGGAGGGGCGATGACCTCTATTCT	1725
12184-L4-1	3p14.1	AAGTTCTGTGCTTGGCCCTGGCTGAGGTATTGAGTGTGAGGAAGGGAACACTGGGC	1726
12184-L5-3	3p14.1	GGCCCTGTGCATAATAATCTTTATGGAATTGAGGGGAAGGGAATTAAAGAAAGGGAAGAGAAAGCAACCCACTACAGAGTTTAT	1727
12224-L4-1	4q27	GACCATCTATTCTTAATATTATAGAACTGGGCC	1728
12361-R5-1	6q23.3	TTGTACACAAATATTCTGCTGTGGTTGAAAGGGGCGCACGCTGAGGTCAAGTGATGTAGTGTGTTTCCATATGAGTCTCACA	1729
12691-R5-1	1q22	GTGTGCCA	1730
12692-L5-1	1q22	TTGCTACTCCATGGGGTGGATTTTCATGGCGACGCCCTTGTTATCTTGCTTAGGACTTTCCAAAGCACCCACATGGTT	1731
12693-L5-1	1q22	TGGTAA	1732
12694-R5-1	1q22	GGTCCACTGTCAGAGAGAGATGTGCCACTGTGCACCTCTCTGAGCAGATGGGGGCAATATGGCACATTTACATGGACC	1733
12696-R5-2	1q22	CCCACGCGTCGCGCGCTCCCGACCGGAGCGGGACGGGCCCTGTGCGGGCGCGCCAGGGCGGGG	1734
12697-R5-1	1q22	GC CGGTGCGCGGTGCTGGCTGCGGGCGGGTCTCATTTCTGCTCAGTCCCTTGCTGCCCTTCTCCTCCCGCCCAAGC	1735
12699-L5-1	1q22	CGCCGTGT	1736
12701-L5-1	1q22	GGGAGGGGACCGGGAGGGCCCGGGCGCGCCGACCCCAACCTCTCGGAGGAGGGGCTGCCGCTCGCCGCTCCGCTCTTTGT	1737
12703-L5-3	7q32.1	TGTTTGGGCTCCGGCTCCCGCTCCCGCTCTCTCCCTCCTC	1738
12704-L5-2	7q32.1	GGGACGTGGCCCCCTCCCCCGGAGCGGGACTCCAAGAACTCCGGGGGCGCTGGGGGCTGACTTTCC	1739
		GATCAGGTTCCTCCCTCCCGCATACACCTGGCGCAGGTGAAAGCTCAGGGAGCGGGTGGGGAGCCCGGGT	
		CGGAGCCTCCTTTCTGTCTCTCTACTCCGTGCGGGCCTGGCCCGCAGAGGTAGGAGGGGCGCACACCGGGCCAGGAGGC	
		TGCC	
		CTGGACGGCGGGCGCGGAGGCCAGGGCCCTGAGGGGCGCAGAGGTGTACAGGTGCAACCGCGCCCGCCAGCGTTCC	
		CGCCACCAACCGCACACCCCTCAAAGCCCG	
		CCGGGCTGGGGTGGAGATGGCAGGATTTGACAAGTTCAGGGCCCCCTGGGATCCCTTTCCCTACTCCCTGTCTTGTGGA	
		CACCTGTTTACCTGCCCTAATTGCCCGG	
		GGCAGCGCCCCCTCTCCGGAGGGCAGCATCCCGCGGGGGGCTCGGCTTTGATGCCAGGGCACCTTTTGTCCCTGG	
		AGACGCTCTGCCAGCCAGGTGCGTGGAGGGAGTGACGCC	
		CAGGTGAGAGGCTGAGCCCCCGGGCGGGGAGGGCGCCAGGCCATTAAACCGTCCCGGGGACCCCTTTTGGCCTG	

197

		TG		
13091-L5-2	9q34.11	TGCAGCCTGGGCCTGCTGAACACTTGAGGGTCGGGGTTCTCAGGAAATCCCTGAGAGCTCAAAGGATGGCCTGAGGGGCCCTGG TGGCA	1764	
13093-L5-2	11q13.1	CTGACTTCTGCGCGGGCGGTGGGGTTCTGCGGGGTGCGAAAGACTCCCCAGATCCCCGCGGCCCCAGACCCAG	1765	
13095-R5-1	1p36.11	GGCAGGGCAGCTTCACCTGGCTCTGCAGCTCCTGAGCCTGGGTGGCTAACCTGGGCCAGGAGTAGGGAGGTGGGTCCCTGCC ACTGCCCC	1766	
13097-L5-2	11q23.3	GTGGACAACCCCTAGGGTGGGGCTGGAGGTGGGGCTGAGGCTGAGTCTTCCCTCCCTCCCTGCCCCAGGGGTCCAC	1767	
13110-R5-1	16p13.3	AGTTGGGGTCCAGGAGGCTCCAGGCTGAGCCCACTCCTCCCCCTAGATACAAGGAGGGCCCTGGGACCCCTGTCTGCTGTTC TCCTGGACTCCTGCT	1768	
13115-L5-3	1q21.3	AGGAAAGGCAAGGTAGCCATTGTTGGGTTGGGTGGGTGCTCGGCCCTGGAGGGGTTTGTTGCTTATCCCCCTCTGTGCTTCAC CCCTACCCAGCTTCCCGCCAGAAAGCCCTG	1769	
13115-L5-3	1q21.3	GGCAAGGGTAGCCATTGTTGGGTTGGGTGCTCGGCCCTGGAGGGGTTTGTTGCTTATCCCCCTCTGTGCTTCACCCCTACC CAGCTTCCCGCC	1770	
13119-R5-2	17q21.32	TCTCCGTGCACGCTGCTGACCGGCTCGGCGACTGCCCTCCCTGCTGTGAGCAGGAGAAACAGGAAGTCTGCCCCGACAGGGAGGTGG CCGGCGGGAGCGGCAGAGTCGGCGTTGAGA	1771	
13124-L5-1	1q22	TGAGGGGTAAGTTTAGGCTTTCGCGGAGGGAGAGACATGGAGCCTGGGAACTCCTTGTTCTCCCCCTCTGCTGCCCTCTCCCCC CCCTTA	1772	
13129-L5-3	20q13.33	TGGCGCTGCTCTGCTGTTCCCTCTGCTCTCCAGACTCTGGGTGGATGGAGCAGGTGCGGGGCCAGGGGACAGGAAGGCTAGGGCC CCAGACTCTGGGTGGATGGAGCAGGTGCGGGGCCAGGGGACAGGAAGGCTAGGGCCCCAGAGACCTGTCTGGGCCCCCATGTC	1773	
13129-L5-3	20q13.33	CAGCTCTGCCCTTAGTCTTGG CCAAGCAGCTTATCGAGGAAAGATCGAGGTGGTTGGGCGGGCTCTGGGGAATTTGGTCTCACAGCCCCGATCCCAGCCCCACT TACCTTGGTTACTCTCTCTCTTAGGG	1774	
13130-L5-1	19p13.3	GAGGAAAAGATCGAGGTGGTTGGGCGGGCTCTGGGGAATTTGGTCTCACAGCCCCGGATCCCAGCCCCACTTACCCTTGTTACTCT CCTT	1775	
13130-L5-1	19p13.3	CATGAGTCTGCACCTCCAGGAGGCTCTGTGCAGTACTCACCCCCCTCCGTCTGCTCCCTCCAAGGTGTGGGGTGGGCAGTGCTC AGGCCCTCCTCGAGCCAAAGGACAGCATG	1776	
13135-R5-1	1q42.12	GAGGAGGAGCCCCCATAGGTGCCCCCTGACCAGGGCTGGCAGGGGCAAGTGTGGACGCCCGGCCCTTGCCCCAGGCTGGG ACCACTCTGAGGGACAACATCATCCTT	1777	
13136-L5-3	5p15.33	GGCATGGGAGGTGGGTTGCGGGGTTAGAGCTGTGGGTGATCAGAAAGGGAAGGGCTTCATTTCTACGCTCTGCCTCCGCTAC CTCTCCCCACCACCCCTAATCCC	1778	
13137-L5-1	5p15.2	ATCCGTGCGCAGTCGCTGCAGCCGCTGCAGTCCGAGCCGACTAAGGGCGGGAGGCAGCTCGGGTGGCGGGCGTGGCGAAC AGCGCGCCGGAT	1779	
13138-R5-1	5q22.3	GCTGGATCTCCGGGCTTGGTTCAGGCGGGGCTCTGGGTCCAAAGCACCAGTGAGGAGATGCTGGAGGGGAAGGGG TGGAGGGCTTTGGTCTTGTGGAGGAGAGAAAGGAGGGAGGT	1780	
13163-R5-1	19q13.32	GGTATCCTGAGAGTTTATAGATGTTTGTTTCATTCCTTTTGCAATGGCCTATGGTATATTATGTATCAACATTTAATGTATGAAAA CGTCTCACTTATATCTTCAGGATACT	1781	
13164-L5-1	10q23.33	GCCTCGCGGGTGGGTGCGGGGTCTCTGACGCGGCAGGCACCCCTCGCTCTCGCCCTCCTGTGGT TCCTGAAAAGAGGTTGGGCAGGCAGTGACTGTTTCAGACGTCCAATCTCTTTGGACGCCCTCTTCAGCGCTGTCTCCCTGCCTCTG CCTTAGGA	1782	
13166-L5-1	20p11.1	CATTAGGTGGTGGTTTTTGGTTTTTATTTTTTTCATATGGAATAAGCTTGTGGACTATCACTAACACCTCTGTG GCATGCACAGTCATTTTTGTTTAAGAGTAATATTTTTAATGTAAAGACGTGGTGAGGGAGGATCTGACAGAGATGAA TGTGCCAAGC	1783	
13181-L5-1	1p21.3	AAAAAAGATACTAATAACAAATGGTCAATGGAGGGGGAATATAGAGAAAGATCAATTTTGTACAGAAAAACCATTTGGTTAGTATT CCTTTAGGA	1784	
13184-L5-1	9q33.3	AAAAAAGATACTAATAACAAATGGTCAATGGAGGGGGAATATAGAGAAAGATCAATTTTGTACAGAAAAACCATTTGGTTAGTATT CCTTTAGGA	1785	
13186-R5-2	3q25.2	AAAAAAGATACTAATAACAAATGGTCAATGGAGGGGGAATATAGAGAAAGATCAATTTTGTACAGAAAAACCATTTGGTTAGTATT CCTTTAGGA	1786	
13195-L5-3	3p21.1	AAAAAAGATACTAATAACAAATGGTCAATGGAGGGGGAATATAGAGAAAGATCAATTTTGTACAGAAAAACCATTTGGTTAGTATT CCTTTAGGA	1787	

		TTTTTTCTTTT		
13199-R5-1	10q26.13	AGGTGCTGGAGAAAGCTGCCAGTGTCTCAACATGTCTGCCTCCCCTGGAGGTGAAGCATGGGCAGAAATCCCCAGCTTCCCA	1788	
13202-L5-1	10p15.3	TGGTGGGCTTCTCAGAGGAGGTGGCAGAGACCCGAGCCTGCCAAGTTGCACCTAAGGTCAACGGGCAGCATTAGGAGGGCTC	1789	
13202-R5-2	10p15.3	TCTCCAGTCTCCCAACCCCTCCATCCCCCTCCC	1790	
13209-L5-2	10q22.1	TGGTGGGCTTCTCAGAGGAGGTGGCAGGAGACCCGAGCCTGCCAAGTTGCACCTAAGGTCAACGGGCAGCATTAGGAGGGCTC	1791	
13209-R5-3	10q22.1	TCTCCAGTCTCCCAACCCCTCCATCCCCCTCCC	1792	
13211-L5-1	10q23.1	GGAGCCCGCGGGCAGGCCCGCGGGCAGCAGGGCAGTTACCTCAACTCCCGCGCTCCGGAGGTTGCCGGGCACCGA	1793	
13220-L5-3	1p13.3	GGAGCCCGCGGGCAGGCCCGCGGGCAGCAGGGCAGTTACCTCAACTCCCGCGCTCCGGAGGTTGCCGGGCACCGA	1794	
13229-L5-1	11p15.5	GGAGCCCGCGTGCCTTCAGGCGCTCGGGCGGACCA	1795	
13230-L5-4	11p13	GCGCTCCGGTCGGTCAGAGAGCGGGCGCGCTCGAGCCCTCCAAGCCCGCTCCTGCGCGCCGGTCCCCGCGCCCGC	1796	
13231-L5-2	11p13	TGCTGCTGCTGCTGCCCGCGCTGCGTGC	1797	
13237-L5-4	1q21.3	CCAGACCCAAACTAGTTGAGTGAGGAAGGGCAGTGGCAGCTACAGGAATAGCCTGTGATGCCTCTGTCCCTCTGAAGATGTCC	1798	
13239-L5-2	1q23.3	CAATGTCCCGTGGCTG	1799	
13240-L5-2	11q12.2	TGCTGGCCCAAGGGGTAAAGGGGCAGGGACGGGTGGCCCCAGGAAGGGCCTGGTGGAGCCGCTCTTCTCCCTGCCACAG	1800	
13241-L5-2	11q12.3	AGACTGGCGGAGCTGC	1801	
13251-R5-2	1q25.2	AGCTTTGTGTTTCGTAATGGCTCAGTTCAGGTGGTTGAGCTGGAGCACTGGAGCTAAGACTGTGGGCTCAGGGCCTCAAGTCCC	1802	
13259-R5-1	12q24.11	TAGAATCACACCACTTCCAATCATTAAGCTC	1803	
13267-L5-1	1q42.13	AGGAACAGGACGATGATGCTGGCTGCTGGGAGCGGCCCGGGTGGGCTCTGCTGGCCCCCTCCTGCGGCCCGCA	1804	
13281-L5-3	12p13.31	CTCTGCTCTGGCCGCTCCTTCC	1805	
13283-L5-3	1p36.11	AGGAAAGGCAAGGTAAGCTGGGGGAGATGGGGGAGCTCTGCTGAGGGTGCACAAGGCCCTGGCTCTACACACATCCCTGTCTT	1806	
13285-L5-3	1p36.32	ACAGAG	1807	
13287-L5-3	1p35.2	GCAGTGTGATTTGGGCGCGGAATGCCGCGCGGGACGGCGATTGGTCCGTATGTGTGCCACCGCCGGCGCTCCGGC	1808	
		CCGCCCCCGCCACACCGCGCAT	1809	
		CTCTGTTACCTTGTCTCTGTGGTTTTGGAGGGGTGTGGAAAGAGCGCAGAACATTCGTTACATTTCTTCTGCTTCTCTGTCAC		
		CAGCTAAAAGTTGCAGAGAC		
		GGCTGGACCACTCCTTGTGCCCCCTCCTCTGCCATGGCACCTGGCTGCCTGACCTCAGCCAGGACAGCCATGG		
		CAGAGGAGGGGCACAAAGGAGGTGGTCCAAAGGCCCA		
		CAGGTATGCTCTGTGACAAGATTTCAGCATTTTAGAGAGCCCCCTGGCCTGCTGGGTGGGACAGACATACAGTGGTAATTCTGGC		
		AGAAGATGAGGCCT		
		CACCATAGGTGAGGTGGGGCCAGCAGGGAGTGGGCTGGGCTGGGCCAAGGTACAAGGCCCTCACCCCTGCATCCCCGCAC		
		CCAGGCTTCAACGTGG		
		GCCGCCCCACCCCTGTCCCCTCGTCACTTCTCTGTCCCTGTGGGTGGGGTGCAGGCGCTTCTCCTTTAGCTGTGCCGCACTTCT		
		CCCTACAGCCAGGAGAAACAGAACACCCGTGTGCAC		
		GGCACAGGGGTGGGTGTGCAAGGGTGGCAGCAAGGAAGGCGGGTCTAAGGTGTCTCCTGCCCCCTCTTGTGTAGTA		
		CTTTGGCCTGAGCAAGAGGCC		
		CTGGGGGTGACCCCGTGACGGGGCTGAGCTGGGGCTGGGTGCATAGCCCATCTGTAGCCTAAGCAATGGTGCAAGCCCCCA		
		CGCCTTGGTTTCTCTCTCTGAAACGCGGCACACAGAG		
		TCCGCCGAATAACTCCATGTGGTCTTGGAGGAGGGTGGGTGGCTCCTCTGCAGTAGGTCTGCCTGGAGCTACTCCAC		
		CATCTCCCCAGCCCCCTGTATGGCTGGGAGGGAA		

13291-L5-1	1p34.3	CGGAGGGAAGGAGGAAGGGGCGCTTGGGCAGAACCAAGGTGGCAGATTATCCTAGGGACTCTTGGGGCAGAACCCAGA CGCCTCTGCGTCTCCCTCTCCCC	1810
13293-L5-1	14q32.2	GTGGCGAGCCAAAGGGCGGAGGCCCTGGCCAGGCAGAGGCTCCGAGGGCCAGCCAGGCCTCCTAACCCCTCCACC	1811
13298-R5-1	1p34.1	GTA TCCCTCACAGCTGCATTTCA GTGCCTGGCTCTGACCTGGCCACAGCACACACCAATTGCTAAGTGTGGGGCTGTGGGGAGGC TGAAGCCCTGCACCCACTCCAGGCTGAAGGCAGT	1812
13303-L5-3	14q32.13	TGGAAAGAGGGACATCGTGGAAAGTTGTGGCAACTGGGTTCAGGGGAGGCCAGGAGCCTGCCAGCTGTGGCCTCTAAAAAAATA AAACCCAGGTGCTCACTGGCCCTGCCTTCCAC	1813
13308-L5-1	15q23	GTGGAGAGGTGGAGGTGGATGGGCTGATGTGAGTGTGTGTGTAGACAGCCTTTCTGACAAACAAGGCTGTATTACACCAAGGAT CTCCTGGCGGGTATCTCAAATTCATCATTCATC	1814
13310-R5-1	15q23	TTTTGGTAATTTAGACTGGAAATGGCAATGGTAATTCAGAAAAAGAAATAAATTTTAATTATTGGTGGTAGTGGCAATGGATTTT CAAGA	1815
13312-L5-2	15q24.1	GTGAGTGGGGCTGGGCTGTGGGGAGGGGTGGGGTGGCAGGGAACAGGCAGACCATCCCTTCTACCCACAGGATCCTGTGCT GCAGACAG	1816
13313-L5-2	15q24.1	CTGATTAGGGTTAGTGGGAGGGACACGGCATGGGAGACAGGGAGATTTGCCTGTGGCCCTGAGCCTGACTGAGCTTCCTTTCTCC CTAGCAC	1817
13316-R5-2	16p12.3,16p13.11	GCTCAATGCCTCCAACGCAGTCAGCTGGTCTCAGCCACGTAACAACCTCACGGTGGAGGAGCCCCATCGTGGCCCTGGTGTGTG GGCAGCAGCAAGTGTGTGGCCCGGCA	1818
13316-R5-2	16p13.11	ATCCGGCTCAATGCCTCCAACGCAGTCAGCTGGTCTCAGCCACGTACAACCTCACGGTGGAGGAGCCCATCGTGGCCCTGGTG CTGTGGCCAGCAGCAAGTGTGTGGCCCGGGC	1819
13326-L5-2	17p12	TGCCATCGAAGGGCGGGTGGGGGAAGCCGGAATCTCTGTCCACATGCTCCAGGCACCTAGCTGTCTGAGGGGCAGAGAGCAG AGGTGGTGTCTCCCCCATCAGCATTTTGAGTTGGCT	1820
13328-R5-2	17p11.2	AGCTTGATCACCTGCCACTTGACCTGTATCACACCTGTGTGTACACCTGTGCTCCTGTGTGTGATGTCAGATG TGGGGCAGGTGAGCATCTGC	1821
13332-L5-1	17q21.31	TTGGGAGGGAAGACAGCTGGAGAGTATGGTCACAGCAGCATCCTCCTCTGTTTTCTTTCCCTAGAT	1822
13332-L5-1	17q21.31	TTGGGAGGGAAGACAGCTGGAGAGTATGGTCACAGCAGCATCCTCCTCTGTTTTCTTTCCCTAG	1823
13334-L5-3	17q21.31	TGGGCAGACAGGGCACAGTGGGAGGGGAGTGCTGCCAGGAGGGCCACCTGGTGACTCCACATCCTTTCTCACCCCCC AGA	1824
13335-L5-3	17p13.2	TGGCTGGGAGAGGAGCATAGGATGCGGCAGGAGGGCAGTGAGGGCTGAGGTACGGATTTCTAGCCCCGCCCTACCCTCCTCTCT GCCCTAGTGCCCGTGGCCAA	1825
13337-L5-2	17q24.3	ATCACGGGTTCCATGTTGGGGAGGGAATGTCAGAAAAAGAATGTCCGTTCAACATTCCTCTCCACCCTTAAATCTCTATGTGA GCGCGGACACCATCTTCTTTAAACCTCAGTCCGTATTGCTCTATGGCATCCATAGAGGCCATTGCGCTCTGAGGTCCTCAGT AAAGAACTTAGATGGTATTACTGTGT	1826
13339-L5-1	17p13.1	TGAGGGGAGGGCGCTCGGGAGGGGTGGAGGGCCAGGGAGGGTGAGGGCCGGGAGCCACTCTGCCCGGCACTCTCC GCCAGAAACAGCCCAACGCCCTTTCTTTCCCTTT	1827
13343-L5-1	17q25.3	AGAGGAAGGAGGGCGCTGGGATAGCCACACAGGTGGTGGAGTCTGTCTCTCCCTGCCAGGTGCGAGGACAGCAGGTCTCAA TCTCGCCCACTGGGACTCAGTGGTCCCTCATTCA	1828
13349-L5-2	18q23	ACAGGCTGGAATTTACAGAGGGCGAACAATTTCTGTGGCCGAAGCTCCCAGTGCGTTTGACAGTCCCTTGAGACGGCCGCAGACACAT TTGTTCTTGGCCGTCTCCAGGTCGTA	1829
13353-L5-2	19p13.3	ACTGAGGGGGTTCCAGGAGCTTTGTGGGGCAGCATTTGGGAGTGTGAACCCCCCATCCAGGTTCTCACCTCTCAAACCTTCCCTCAGAT GCTGAGCCCCACTTAT	1830
13354-L5-1	19q13.11	GTTTCCCATTCACACTGCAGGAAGGATAGGGGTGGTGTGTCAGAGGACCCCTCACATGGCTCCACTCTAGAAAGCTGGCATGGG CAGCTC	1831
13355-L5-2	19q13.12	TGGTGAGCACCGTGTGGTGGCAGGGAGGCCACCTAGTGCCACCTTTCCGTGACTCCTGTCTCCTCCCTGCAGCC CGTCGGGATGAGGCCAGG	1832
13356-L5-2	19q13.2		1833

13358-L5-2	19q13.32	CGGCCAGTGACTTCCCAGGAGTGTGGAGGGGGTGGTGAGGAGGAGCACCTGGGCTCTCTACCCCTCTCCTCACAGAAGTACCT GAAACTAGGTC	1834
13361-L5-1	19q13.33	GTGTGCAGGGCTGGGTCACTGACTCTGCTTCCCCTGCCCTGCATGGTGTCCCCACAGGGACAGCATGGACAGTGTCAAGCAGA GTGCGGCCCTGTGCCTCCTTCGACTGTACA	1835
13363-L5-2	19q13.42	AGTCGGGTGGGATGTGAGGGTGTACAGAGTACGGTGGGGGCCACGCTGACAGCCGCACCTGCCTCTCACCCACAGTGCAAC CAGAGACCCCTCTA	1836
13364-L5-2	19q13.43	AGGTGGAATGGGCTGAGGCTCAGGAAGGTCCTGGGATACAAATGCTCACTCTATGGGTCTCTCCCTGAGCAGGACATTGGTTAC C	1837
13365-L5-3	19p13.3	CCAAAGCAGCTTATCGAGGAAAAGATCGAGGTGGGTTGGGGCGGCTCTGGGGATTTGGTCTCACAGCCCCGGATCCCAGCCCCACT TACCTTGGTTACTCTCCTTCCTTCTAGGG	1838
13365-L5-3	19p13.3	GAGGAAAAGATCGAGGTGGGTTGGGGCGGCTCTGGGGATTTGGTCTCACAGCCCCGGATCCCAGCCCCACTTACCTTGGTTACTCT CCTT	1839
13370-L5-2	20q13.12	TACCGCAAGTATGTGGAGCAGAGGTGAGGGTGGGGCATGCATGTGTTAAGCCACCAACCCACAGACCTTTCGCGCCACACTCT CATCCTGCAGGTAC	1840
13373-L5-4	20q13.33	TCCCTGCACCCCTGCCTGTACAGGTGAGTGGGAGCCGGTGGGGCTGGAGTAAGGGCACGCCCGGGGCTGCCCCACCTGCTGAC CACCTCCCCCACAGCACCTGTGCCGGGGC	1841
13374-R5-1	20q13.33	TGGCGCTGCTCTGCTGTTCTCTCTCTCCAGACTCTGGGTGATGGAGCAGGTCGGGGCCAGGGGACAGGAAGGCTAGGGCC CCAGACTCTGGGTGATGGAGCAGGTGCGGGGCCAGGGACAGGAAGGCTAGGGCCACAGACCTGTCTCTGGCCCCCATGTC CAGCTCTGCCCCTTAGTGCTTGG	1842
13374-R5-1	20q13.33	ACCTCTCAGGACCCCTCCTAAGGGGTAGGCAGGGGCTGGGTTTCAGGTTCTCAGTCAGAAACCCTTGGCCCCCTCTCCCCCAGACCC CCAGGCTGTGTGAGGGTCTGAGAGCTGGTAC	1843
13375-L5-3	20q13.33	CCAGAGGAAAGCTGCAGGTGCTGATGTTGGGGGACATTCAGACTACCTGCAGCAGAGCCCCATGGCT TAATCTCCAGTGAAAAATAAGCATGAGTGTGAGGCTGAAAAGGCTGGAATCATCTCTCGGTGTGCATTTCTCCGGGGCTGGGAG TT	1844
13376-R5-1	2p25.1	CCAGAGGAAAGCTGCAGGTGCTGATGTTGGGGGACATTCAGACTACCTGCAGCAGAGCCCCATGGCT TAATCTCCAGTGAAAAATAAGCATGAGTGTGAGGCTGAAAAGGCTGGAATCATCTCTCGGTGTGCATTTCTCCGGGGCTGGGAG TT	1845
13380-R5-3	2q21.2	TT	1846
13385-L5-1	21q22.3	ACTGTAGGTGGCGCGGAGGAGTCAATTCCTCCATCACTAATGGCTTCTCTTGACACCCAG GGACACGCGGTAAAGAGCCCCAGTGGGGCGGCGAGTCCCTGCAGGGACTGTGACACTGAAGGACCTGCACCTTCGCCCCACA GAAAGACTCCTAGCCAGGAGGGCCTGC	1847
13396-L5-2	22q11.21	GAAAGACTCCTAGCCAGGAGGGCCTGC	1848
13403-L5-1	22q13.1	ACCAGGTGGGCCTGGCCCCAGGTTGGGGGACACGGGTGGGTCCCGGCACCCCTCCCCTGACCACCGTGCCTCTCCCCAGGA CTGGACACAGGGGGCGCCATCTTGACTCATGTCTCAAGATGGCGACCTGGAACGCTGAGCG	1849
13412-L5-1	2p16.3	CTGGACACAGGGGGCGCCATCTTGACTCATGTCTCAAGATGGCGACCTGGAACGCTGAGCG	1850
13423-L5-3	3p24.3	AACCTAGGCTCCTAAAAACAATAAGCTTCAGGGAGTCAGGGGAGGGCAGAAATAGATGGCCTTCCCCTGCTGGGAAGAAAGTGGGT CAAGGGGAAAGGGTG	1851
13425-L5-3	3q26.33	CCTGTTGATTCCCAAGAACCTTGATGCGGGCCATATGACTGGGAGGGCACCTAGAGCATTTTCTGGGTGAATAAGGCCCTCCATC CCTCAACAGCAGCACAAAGTGTGTGAAAAAGGA	1852
13430-L5-3	3p21.31	CCTAGGAAAGGCTGCTGGTAACCTGGGATGGGGTTGGGGGAGGTAAGAAGTCTCTGACTCCTCCTCTACCTCATCCCAGTTCCA TCACCTGAAGTGACCTCTTGGGAC	1853
13431-L5-3	3p21.31	GCCTCTTCCCTGCTCTGCTGTGAGTGGCGCAGCGGCAAGTGGGTGTCGACGGCGGGAATGCCCGTCGCTGCTGCTGCTGC TGCCCGACGGGCCTGGG	1854
13432-R5-1	3p14.2	TGCTTTCTGCATTTCTCTCCCTCCCGGCTCTTGTGACAAGCCATACTGTTAAATATCAGAATAGTAGGTGATTACGTGGAGTTTG GGAGGTGGTAGGAAGTGCCAGAAACTGTAA	1855
13456-R5-2	4q12	CCAGGAGCTACCAAGCAGAGGTTTATTCAAGTCTCCAGAAAGGCTATGCAGGTGGAAAAAGATTTCATAGCGGGGCATCTG GCTGGAGCCTTCCCCCTCCCTCCTCGGCACTTCCCCCACCCTCACTGCCCGGTGCCCAAGACTGTGGACAGTGAAGGTAGAGG GAGTGCCGAGGAGGGCACAGCTGTGCCTCAGA	1856
13458-R5-2	4p16.1	GAGTGCCGAGGAGGGCACAGCTGTGCCTCAGA	1857
13461-L5-4	5p15.33	TCCGGGAATCTGGAGGTTTCTAGCACCTTTCAGGCGGTGGGGACGGGGCTGGGGAGAGAAATCCCCCAGCCCTGTTCC TCCGGGAATCTGGAGGTTTCTAGCACCTTTCAGGCGGTGGGGACGGGGCTGGGGAGAGAAATCCCCCAGCCCTGTTCC	1858

		CTCCATCCTGGCTCCAAATCCCAGTTACTCCCCGGA	
13463-L5-2	5p15.33	GGGGACCTCAGGCTGGGGGGCTGTTCCGAGTCTTGATCCTCAGGGACTCTGGGAAGCTGGAGCTCTAGCACCCCTCCTCCC	1859
13489-L5-1	6p21.2	TCTTGTTCCCTTGGCTTGACCACACAGGCTTCCCCCAGCCAGGGGCTTCTCCCTCACTCCCCCTGGCTCTGCCCTCCCAATC	1860
13497-L5-1	7q22.1	AGAACCACTCGAGC	1861
13513-L5-1	7q11.22	GCGGTCCCCTGGGGCTGGGAGCGAGGGGGCACAGATCTGATGTGCCCCCACCCCTCTCACAGGACTGG	1862
13519-L5-1	8q24.21	CTTTGGTGAGAGTAGCACCAACCAGGGAGGATTACAGGCATTAATCATCCCTGGCAGAGTGTCCCTGGAAGTTTACTCTGAAGAC	1863
13525-L5-2	8q11.23	GGGGGTGGGAAGCGAGGTTGGGTAGAGCTAAGGTATCAGCAGGACATAGTAACAAATTAAATAATTATCTCTGGCAAATGATAATA	1864
25-R5-1	2q31.1	TAGATCTAAACTTTTCAGTCCCCAACTCAA	1865
266-R5-2	12q14.3	GGCCAGAAGACGATGGCGGTAGGGGAGATCACGCGAGAGCATGGCCTCCACCTCTCCTCCCCACCGCCCATCTCCACGTTCTCGC	1866
2786-L5-3	2q31.1	TCCCGCAGCCGGTGACTGGAGCCCACTCTGCAGAGACAAAGGTTAGAAAAAGAGGGGTGAGGCTCTGGGAAAGCAGAATGCG	1867
2811-R5-1	17q25.2	GGG	1868
2819-R5-4	15q22.2	GTTGCTATTTCCCTCAGTTGAGGGCGAAGTTAGCAAATCCGTAGCTGCAAGTCTCAACTTGGGGGAGGGGGCGAC	1869
3717-L5-2	1q21.3	AAGCCCTAGAGATCAAAAGCATTAGTATGGGCAGTTGAGCGGGAGGTGAATATTTAACGCTTTTGTTCATCAATAACTCGTTGGCTT	1870
3732-R5-1	Xp22.12	AGCATGGAACAAGCCAGAAATCCCTCTCTTACCCCGAGAGCTGTCCCTCCAGGTCAGGTTGGGACACACAGTGGCACGGGTGGC	1871
3799-R5-1	11q13.1	TGTGGTGGCTTGCACTGCCG	1872
3897-R5-2	9q12.9p11.2	AATGCCAGTGAGTTTGAAGGCACCTTTGTCCAATTAGAAAGTGTGGAGAAATATTTCATCCTGTCCAATGACAAAGATGAAGTGTCTTCTT	1873
3942-L5-1	1p31.3	TCAAAGCGCGGTGGCAGGCTG	1874
3952-L3-1	1p36.32	TGTAGTGGGTTTGGGTGAGGCTATGGGAGGGCGGGTGCCGCCCTTGCCCAGCCCCAGCCCCAGTACA	1875
3953-R3-2	9q33.3	CGGAGCGCCCGCGCCAGCCTCTCCATCTCGCAAGTTTTTAATTAAAGCTGAGGGGAGGGCGGTGACGGCGGGTCCGG	1876
3966-L5-1	3q27.2	GGGAGTGGGGGATCTGTTTGGCAACTGGAGTGAAGGATTGCCCTCCCCCTGCTGGGATCCCCCAGCCC	1877
3995-L2-2	7p21.1	CCCCGGCAGGGGTGTCAGATTGAGTGTCTGTGCGCATGTGCGAAGGTGTCCAAAGTGTCCAAACTGACAAATGCTGGGG	1878
4013-L4-1	15q22.2	GCTCCTGCTCCGCCCGGGAGCTGCTCCGGCGCGCAGGGCTCGCTCGGGAAGCTGAGCGCGGAGGCTGGAGT	1879
4026-R5-1	10q24.32	GGCGAAGCGCGCAGCGCGCGGACAGCTCTGGGTTTGCCTCGGGGTGTGTCGGCCCGCGCTGCTGTGGGCC	1880
4026-R5-2	10q24.32	TGGCCTGACGTGAGGAGGGGACTTTTCGAAGTTTTATAGGAAAGTTTCCGCTTCCAGTCCCCCTCCCCCGTCCCA	1881
410-R5-1	5q31.3	ATGGTCCCCCTGCTGATAGCTCAACCATGCAAGTGCCATAGGAAACACAGACTTTGGCAITTTGAGCTGCTCACAGTCTGGGTCAT	1882
4130-L5-1	17p11.2	GGCTCTGGGAAAGCCTTCCTTTCCGGCTGGCCTGGCATTCAAAGCCAGACAAAGGGGGCTTTTCTCTCTCGCC	1883
4143-R5-2	22q12.3	GGCTCTGGGAAAGCCTTCCTTTCCCGGCTGGCCTGGCATTCAAAGCCAGACAAAGGGGGCTTTTCTCTCTCGCC	1884
4258-L5-1	2q31.1	CCAGCTCCGCCAGCTCCAGCCCCAGGTCTCTGAGCGATGCGGCCCCACGAAGGTGCCGTGTTTGTCTTCCCTCGGGACGGAGTAGT	1885
4315 C-L4-1	1q22	GGAGCTGG	1886
		GGGTGCTGGTGGACATGGACGGCGTGCTGGCTGACTTCGAGGGCGGATTCTCTCAGGAAGTTCCGCGCGCGCTTTCCCGACCAGC	
		CCT	
		TCAGCGTCAGGAACCTTCATCCTGGCAGCCGACCTCATGAAGAGCATCTGGCTGCTGTTACCAGGAGAGAGCAAGACGCTGA	
		ACGGGAAAAGAGGAGAGGCACCAGATATATGTTCTCTAGGCCCTTTTAGAAAAACATGGGGTTTTTCCTTTGGCCACGT	
		GGGAGGGGACCGGGAGGGCGGGCGCGACCCCAACCTCTCGGAGGAGGGGCTGCCGCTCGCCGCTCCGCTCTTGT	

		TGTTTGGGCTCCGGCCTCCCCCTCTCTCCCTCCTC	
4315_D-R4-1	1q22	GGGACGTGGCCCCCTCCCCCCGAGCGGACTCCAAGAACTCCGGGGCGCTGGGGCTGACTTTCC	1887
4315_E-R4-1	1q22	TCCCCGCCACCCCTTGAGCACCTCAGCGTTCTTAGGGGAAGCCAGAGCCGGGAGGATCGGGAATAGTTTGTGGGG	1888
4315_F-R4-1	1q22	GATCAGGTTCCCTCCCCCGCATACACCTGGCGCAGGTGAAAGCTCAGGAGCGGGTGGGGAGCCCGGTT	1889
4315_I-L4-1	1q22	CTGGACGGCGGGCGCGGAGGCCAGGGCGCCTGAGGGCGCAGAGGTGTACGCGTGCACCGCGCCCGCCCGAGCGTTCC	1890
4315_K-L4-1	1q22	CGCCACCAACCGCCACACCCCTCAAAGCCCG	1891
4315-L4-1	1q22	CCGGGGCTGGGGTGGAGGATGGCGAGGATTTGACAAAGTTCAGGGCCCCCTGGGATCCTTTCCCTACTCCCTGGTCTTGTGGA	1892
4315-R3-2	1q22	CACCTGTTTACCTGCCCTAATTGCCCCGG	1893
4315-R3-2	1q22	GCTCCCCGGCTCCCTCACTGCGGCAGCCCGCGGCCCCATAAATCGTGAGAGCGACGTGCTCCGGAGCCGAGAAATGGAGAGGGCC	1894
4315-R3-2	1q22	GGGGAGC	1895
4338-L5-2	1p34.3	GCTGAGCAGTGGGGCCAGCTCCCCGCCACCCCCAGGAGACTGGTGAGGAGAGCTGTCCGGCTGAGCAGC	1896
4340-R3-1	10q21.3	GCTCTGCTTCCAGGCTGATTTTTTAGGCTGGCATTTAGGTTTGGCCCTGGGACAAAGGGCTGGAATAATGCAAGGC	1897
4346-L5-1	2p23.2	CCGGTGGCGGGCGCGGAGGGCGGGGGAAGAAACACCCCTGTTTCTCTCCGGCCCCCACCCTGATCATGTACCAGG	1898
4361-R5-1	Xp11.22	TGCTGGAGGTAAGGGTTTCTGAAGCCTGGTGCCATGGCCACATGTGCACATGAGGGAGGAGACGCTGAGGCTAGCA	1899
4498-L3-2	6p22.2	TTCCCCAGGCAGCAGCAGCGCCGCTCTGGATCTCCCTGGAGGTGATGGTCGAGCGCTTGTCTATAATGCCAGCGGGGA	1900
4516-L5-1	10q11.22,16q2 2.3	GGCAGAGCCAGGCAGGAGCCAGAGTCAGCCTGGGGGTGATGGAGCTGAGGCCATCACCGGTCTGACAGTGGACCATGATGGC	1901
4516-L5-1	2.3	ATGCT	1902
454-R5-1	5q31.3	CCAGCTCCGCCAGCTCCAGCCCCAGGTCTCGCGGATCCGGCCCCACGAAGGTGCCGTGTTTGGCCTCCTCGGGACGGAGTAGT	1903
454-R5-1	5q31.3	GGAGCTGG	1904
4593-R5-1	15q23	CAATCAATTAGCACATGAGTAAACCAAGCCCCATTAGGACAAACTGATGCCGGGGAGTTATGTCTCTATAGAAATGATTG	1905
4610-R5-1	8p12	GCCCAGTTAATTGGTCTCTCAACCTACATTAGCTGTTGCATTGCAGCCAATTAGGCAGGGGCCAGAGGGC	1906
4610-R5-2	8p12	GCCCAGTTAATTGGTCTCTCAACCTACATTAGCTGTTGCATTGCAGCCAATTAGGCAGGGGCCAGAGGGC	1907
4642-R3-2	11q24.1	CAGCCTGCCTTTCACATATCCGAGCCCTCTGTGACACATCAGATTTAATGAGACTGCACAGATCTCAATAAGGAAGGGAGCGCT	1908
4642-R3-2	11q24.1	G	1909
4666-R5-1	1q21.3	GCCGGCTCCAAACCCAGAGGCCCGGAATAGGCGCGGAGTTATAAATAGTGCCACCCCGCAGGTGTTGGGGGAGTCGGC	1910
4792-L5-2	2q31.1	GGCTTGTTATGGGGCAGTGGGGCTGTAACCTGGCTTCGATGTGGCCATTTCCCTGCCCTTGGTGCAGTTAAACCAAGCC	1911
4801-L5-2	1q32.2	ATTAATGTTTTTTGTAGCAAAACAGGAGGCAGAGTTCTCCAAAGGCTCTCATCTCTGTGCTTCCAGAAAATATTGAT	1912
4813-R3-1	16q22.3	ATGACATCTGATTGATAGCTTGGGCCAAAAGGTTGGCAGGCCCTGAAGGGCGGGAGGGTTAACAGCCCCGATTGTAAAATCAATG	1913
4813-R3-1	16q22.3	TCAT	1914
4875-R2-2	3p22.1	TGTGAAGCCACAGGAAGGGCTCTGTGACATCACAGTAGGGCAGTGTGAAGTCACAGGAAGGGCTGTGGGAAGTCACA	1915
4912-L5-2	12q13.2	CTTGTTAGACAAATCACAGCTGGGGCCAGGGGTAAATTTAGCCCTTTGTTACAGCCCCCAGTTATGTGGGATGCAGG	1916
4929-R4-1	17q25.3	GACGAACCTTGCCCGCAGTCCTTGCGAACTGTTCGCCAAAACAGTCCGGGCTGCGGTGGTTCAGCGTC	1917
5032-R5-1	14q32.31	GGGCGCTCCAGTCTCGACGTTCCCGGGGTAGAGAGCAACGTCCGGGGACGAACGGGACGGGTGCGCCC	1918
5048-L5-1	11q13.1	GGTGTGCAGCAGAAAGTACGCGTACAAGGAGCGGGGCCAAGCTGTACGTGTGTGAGGAGTGGGCTGCACATC	1919
5071-R5-2	17q12	GATTCCCTGCTCCAGAGCCATAAAGTGGGAGCCCCCATTTAATTTGGGCTGGGACTGGGCGGGGGTC	1920
5107-L5-1	11q23.3	CAGTTTGTAAGGGGTCAAAGGGGAGTTCTTTCCGAAGAATTTCTGGGAGAGACGCCCAACTAGAGCAAGCTG	1921

5210-L5-1	1p31.3	TGGGGTTGGATCCCCATGGGGCAGCTTTTGGACCACCAAGTGTAAACCCAGGAATTGGAGCGATCTAAACCCCTA	1915
5342-L5-1	8p21.2	TGGAAGGAGAGCAGCGCATTTGGTTTGGTGGGCAGATTTTCTTTTACGACTGCTAAATGCCCTTCTCCTCCCA	1916
5491-R5-1	6p21.33	GCCTCCCTTCCAGGTCATTACCTGCGGCCGCGGAGGCAACAGCTGCCACCATGGCCTGATGATGATCTGTTGGCGACGGC	1917
5521-L5-2	5q31.3	AGAGGGTGTCTCTGGGAGGGCCACCCAAAGACAGACCTCATGGCCCTTCCAGCCCTTCCATCAGGGTCCATCCTCT	1918
554-R5-1	14q24.3	GGAGAGGGCGCAGACAAGAGAAAATAAGCCTGCCCAGCAATCATGTCTGGGGCTGCTGGAGGGCTCTCTCTCTTTCC	1919
5554-R5-2	1p34.3	CCCCACCAACCAGTGCTCAGGACTTCTGCAATCCCATTCGGATCTGGGAGCAGCTGATGAGGGGTGGGG	1920
5638-R5-2	6q23.2	GGCTTTGCCCTTTTCGGTGACACAGGCTGTTGCTATTCCAAGCAGCCTATCACAGGCAGGGGAGGGCC	1921
5640-L3-1	1p34.1	TCTCGAGAGGGCAGGCACGGTGTTCATGGCAAGACGGCGTTGATGTATAGCGTGGCATGAAGCTGGGCTTGCTGCTCTCA GA	1922
5749-R5-1	18q23	ACAGGCTCATCCCTCTGAACAGATGAGATTAGTCGATCATGTAAAGTCAAAAGGGGATTATGATGCTTGT	1923
5757-L5-1	19q12	GTTGGCGCACGGTAGGTTTCAATTAGTCAATAAAGCATTGCCACGCGTTTTTATTGAATTGTAACCCATATTTGCCCTGAT	1924
5854-R5-2	11p14.1	GCCCTCCCTCCACCCGACATTACACCTGAACCTGTCTCCAGCACTGCCGACACCCGGGTGACACATTCTTTCCGAGAGGGAGGGC	1925
5956-L5-1	1q41	CTATGCGTCTTGACGGATTTGGGGTTGGCAGAGCAGGCTGCCCTGCTTTCTATCCCCATTCACTCCACTTATAG	1926
5995-R5-1	7q11.22	CCTGCTTGATTTGCTTAATGGAAGCTGACAGTGAAGTTCAACCCCTGATTGTCAGATTTCAAGTAACCTCAAGGGGG	1927
6008-R5-1	4p15.32	CGACTTTATCACCCATCGGTTATCTGTGTCGCTGAAGGAACCTCCGGTAAGGTACAGGAAAAAGGTGGGGTATTGTTG	1928
6008-R5-2	4p15.32	CGACTTTATCACCCATCGGTTATCTGTGTCGCTGAAGGAACCTCCGGTAAGGTACAGGAAAAAGGTGGGGTATTGTTG	1929
6016-R2-1	1q23.3	TTTCTGTATATGTTTCTGGAGTCTGAGCCTGAGCTAAACAAAAAGCAGGAGGCTGACGGGCTGCTGGAGTTTGCAGAGA	1930
6023-L5-1	Xp11.4	GGTCTCTGGGGGTTGGACCAGTCATCTGGCTCTGCCCTTCCTCAGGGCAGAGCTGGCTGCTGAGGCCAGGAGGCC	1931
6087-L4-1	9q33.1	GATGGAGAAAGTGGCAGGCAGTAATGGAGACAGAAATTTCTGTTAACTGCTGTAATTAATGTTATGTCATC	1932
6096-R5-1	3q29	GCCATTGGTACCTGATGTGATCGGGCTTTTCCCTGCTGTGAAAAAACGGGCGCAGGATTAAAAACATAAGGGAAGGTGGT	1933
6192-L5-1	11q25	GTGCTGGGTGGGTGTTTTTATCTTACCGGATTTATGGAGTCTTAAACATCTGTCCGTTCGTATCCCCCGCTCAGTAC	1934
6198-R5-2	1q25.3	GCCGCCAGCACCCCGGTGCCCGGGGGCGCTCCGAGGAGCCTGAGAGACCCACGGAGGCTTCGCGGGAAGACGCGCGCGCG	1935
6242-R5-1	8q21.13	CCCTCACGAACGTGCGGTAATGAGAAATCCAAACACCAGGTGCTCCAGGGGCTTTGGGTTTGTACATATGCCACTGTGGGGG	1936
6287-L3-2	1p34.1	AGCAGCCAGGTGAGCCCCGAAAGGTGGGCGGGGCAGGGCGCTCCCAGCCCCACCCCGGATCTGGTGACGCT	1937
6385-R5-2	11q13.1	GGCTCCGTCGACGTTCCGAGCCTGCTGGCCCGTCGGGCAGCTGTCCGGCCCGGGAGTCGGTCCGTGGGCAGGGCC	1938
6409-L3-1	11q13.1	GTTCCAGAAAGCGCGCGTGGGTTGGGAACGCGGAGCGGACGGATTTCGATTCAACGGGTTCCGGACCCGCTGCGCTATGG AGC	1939
6434-R5-1	15q25.2	GTGGGCTGCATGTTCCCGCATTTGCTGGTGAGGGTGACGATCTGGCAGCTGCAGCTGGCTGGTGGGAGGGCTGCATCCTAC	1940
6490-R5-3	1q21.3	TCCTTCCCCCTTCGTGGCTTGGGTCTCTCTTCCCCGCTCGGCCCTCGAGTGTGAGTGTGGGGTGGTGGGTTAGGAG GGGGAAGCGTCATATGGGGGATGGGG	1941
6496-R5-2	5q31.1	CTCCCTCAGGCCCCGCTGCACCTTTCCAGCCCCCAGGACTCTAGGGGAAGTGTGGGTGGGGGAGGGGG	1942
6584-L5-1	12q24.23	GCTTGGTGAGAGGAGGAGGAGGCGGCGGACCCGCTGTCTGCCATCTGGTCCCCCTTCCCCCTCCTCTCATTTGC	1943
6590-L5-1	Xq13.1	CAGCTGGGGCAGAGGCTGTTTATTGTTTCTCATTAACCAAGGAAGTGCCCTGGATCAGATGGAGCCTCTGCTTGACTG	1944
6642-R5-1	14q11.2	ACACTCTCCTCTTGCTCTCCTTGTAATCAATTCATTGTCATCAGAAATGTGTGACACCTCGAGGGGAGGGACGTGT	1945

205

7781-R5-2	17q12	AGCCTGTTCCGTGCTCGCTAACTATAAACTATCTGATTTATATTCATTAACCACTAGACAGCGGCAGGCACAGGCT	1974
7824-R5-1	6q16.2	CCTGGATGCTGTTTCATTATGTAGAGTCAGGCAAAAGACACAGACGGATGTGTGTGAGCGCGCATGAAGCTGGCACACAGG	1975
7846-L5-2	17q12	GCCCTGGCAGGGCTGGTGAAGGGATGGGATGGAGGAGACTCACATCCAGCCTCTGCCTTCCCTTCCCTCCCTCCCTCCCT	1976
7883-R5-1	11q14.2	CTCTCCGTGCTTCTCGGCCCCGCCGCCCTTCCCTGTCTCGGGGACACGGTTTCTAAATAAGGTTGGGAGGGTGAGAGGGCGGGAGG	1977
78-R4-1	1q21.3	GGAGAG CCAAATTGAGAGCTCAAGGACTACAACCTCCAGTGTGCACCACAATTGCTTTGTATGTACCTTGGGAGTTGTAGTCCCCCTTTGAGAA	1978
7949-R5-1	5q31.3	GGTGCGCGGGTGGACGCCGATTCTGGGGCTACAATGCGTGGCTTTCGTATGAATTGCAGCTGGCGGGCTCGGCGCGGCATC	1979
7971-L5-1	9q31.3	AGCCTGCACCCAGGAGCGGAGCGGAGGTGACACAGTGAGTCACAGACCACACAACAAAGCCCTTCCCTGGCATGTCTGGGCT	1980
8016-L3-1	12q21.1	AGAGGGGTGACTGCGGGGCTTGTTCGGCTGAAGATTTACAAATGTACTTCTTGACGGGCTCAGCAACCCCTCT	1981
8062-R5-1	11q23.2	GGGCAGGCGTCCCTGCGGGCTCCAGGCTTCTCTGCCACAGCTGGCGGAGGTGCTTAAGGCCTTGGACCCCGTGACCTTTTGCCC	1982
8077-R3-1	Xq22.3	CCAATTCTCACTTAGGTGTTAGGGATTTAATGATACTCTCTGAAGAGTATTTTACTACCTGAGGGTGGGAATGG	1983
8089-L5-1	4p16.2	GGAGGGGGTGGCCCTGAGCTTATAAGGGTCTCTAAGAGCCAGGAGACAGTCAGCCTGGCTACCTCTAATCTC	1984
8239-R5-1	Xp21.2	TGACTTTTCATCCCCCTTCACACCCCTCTGCATTTCCCTTTTGGAGAGCTCAGAGGGAGGTGAAGTGGGAAGAAAAATCA	1985
8250-R5-2	9p11.2	CGGAGCCGCCCGCGCCAGCCCTCTCCATCTCGCAAGTTTTTAATTAACGCTGAGGGAGAGCGGCTGACGGCGGGTCTGG	1986
8281-L5-2	11q13.4	CCAGCCCTCAGCTGCAGCTGGGAGGGGCTGAAGGGTAGGGAGCCCTATCCACCTGCATCAGAGGCCTGG	1987
8298-R5-1	22q13.1	GATGCCGGGCGCCCGCGCAGCCGCTGCCCGCGAGCCCGGATGGGCGAGAGGCTGCGGCGGACGCCAGCATC	1988
8329-L5-1	1p34.2	GAGCTGACCCAGGGTGGCGTTTGGAGGACTCAGCCTCCGTCCTGCCTGGTGGTGGATCAGCTT	1989
8336-R5-2	Xq13.1	CCTCCCCAAGCAATCTGCTCACCCACTTGTGTGCATGGTGAAGGCTGGGGTGTTGGGACTGAGTGGGAGG	1990
8394-L5-2	7p13	GGGGCCAGGGATAGTCGGAGATGGGAGGGCGGGGCCCCACTGGCGAGGGGCCCTCGGCTTCTGGGGTCCCTGAGCCCC	1991
8564-L5-1	5q13.2	AGGGGTGGGAGGTCTGTTTGGCAACTGGGTGAAGGATTGCCCTTCCCTGCTGGATTCCCCAGCCCCCT	1992
8564-R5-2	5q13.2	AGGGGTGGGAGGTCTGTTTGGCAACTGGGTGAAGGATTGCCCTTCCCTGCTGGATTCCCCAGCCCCCT	1993
8898-R5-1	17p13.3	GGCGCTGTCGGCCGGGCGGCCGGCAACTCGTCCGTCTTGATAAACCATGTGTGCGGAGCGGGCTCCGCCCTCGGCTGTC	1994
9021-L5-2	10q23.1	GATGGTGGGAGCTTGGGTGTTTGTGTTTCCCATTTACAAAAACAAGCAGCCAACCTTACATTTCATC	1995
9068-R5-2	14q24.3	GTCTGCCCTCTTTCTCTGCAGTAATTGCTTCTGCATTTGTGTTTATTTTAAATTAGGAGAGCAGTCTTGATCAAGAGGGAGGCAGAC	1996
9087-L5-2	3p14.1	AGGAAGGGAATGGACTGGAGGGTTCTTTTCCCTGATGGAAGCCCTATTTTCTTTATTGTGTTCCCTTTTCT	1997
9134-R5-1	8q24.3	GCTGCTTTGGGGTTTTCCGCCCTTCCGCGCGCTGCAGCTCCCCACGGCAGCCCCGCAACCAAGCGGATAAACCTTTCTCTCGG	1998
9217-L3-2	2q31.2	TCCACTGTTGGCTTTAGTCACGGCGGGGATCGTCAGTTTAGCGCGGCCCATCGCTAAAGGAGATCTGCACGCCCGGCAGAGTGGA	1999
9245-R5-1	5q21.1	GTGA AGCCTAAATACATTAGCGAGCTGGTAAAGCTTTTAAGGCCCTTCTTGGGAGCGAGTGGCTGGCTAATGAGAGGTT	2000
9287-L5-2	17q21.1	TGCATGTGTGGCTGGGAGGGCTCTGAATATCTCTGGAACGGTACCCAGAGCCCTGTGGCTCTGCGCATGCG	2001
9369-L5-1	Xq26.3	TCCCCTGATTTCCCTCTGTGTGGAAGAATGTGTGAATTCACATGCATCCTGTCTAAGTGTGAGGGGAAATTCAGTACGGGA	2002
9384-R5-2	16q22.3	AAACCAGTTAGTCAATTCTCAGGTCCCTAAGTGGCGATTGTGTGTAATGTCTACATCAATACGGACGGAGCGGTTTGCTAGCTG	2003

[illegible]

miR-149*	2q37.3	GCCGGCGCCGAGCTCTGGCTCCGTGTCTTCACTCCCCTGCTTGTCCGAGGAGGGAGGGGCTGTGCTGGGGCA GCTGGA	2031
miR-150	19q13.33	CTCCCCATGGCCCTGTCTCCCCAACCCCTTGACCAGTGTGGGCTCAGACCCCTGGGGGACAGGGACCTGGGGAC	2032
miR-181b	9q33.3	CTGATGGCTGCACTCAACATTCATTGCTGCGGTGGTTGAGTCTGAATCAACTCACTGATCAATGAATGCAAACTGCGGACCCAAA CA	2033
miR-181b	1q31.3	CCTGTGCAGAGATTATTTTTTAAAGGTCACAAATCAACATTCATTGCTGTGCGTGGTTGAACCTGTGTGGACAAGCTCACTGAACAA TGAATGCAACTGTGGCCCCGCTT	2034
miR-181d	19p13.12	GTCCCTCCCTAGGCCACAGCCGAGGTCACAAATCAACATTCATTGTTGTCGGTGGTTGTGAGGACTGAGGCCAGACCCACCGG GGGATGAATGTCACTGTGGCTGGCCAGACACGGCTTAAGGGGAATGGGGAC	2035
miR-185*	22q11.21	AGGGGCGAGGGATTGGAGAGAAAGGCAGTTCCTGATGTCCTCCCTCCCCAGGGCTGGCTTTCCCTCTGTCCTCCCTCCCA	2036
miR-214	1q24.3	GGCCTGGCTGGACAGAGTTGTCAATGTCTGCTGTCTACACTTGCTGTGCAGAACATCCGCTCACCTGTACAGCAGGCACAGAC AGGCAGTCACATGACACCCAGCCT	2037
miR-23a*	19p13.12	GGCCGGCTGGGTTCCCTGGGATGGGATTTTGCTTCTGTCACAAATCACATTTGCCAGGGATTTCCAAACCGACC	2038
miR-30a	6q13	GCGACTGTAAACATCCTCGACTGGAAAGCTGTGAAGCCACAGATGGGCTTTCACTCGGATGTTTGCAGCTGC	2039
miR-30d	8q24.22	GTTGTTGTAAACATCCCCGACTGGAAGCTGTAAGACACAGCTAAAGCTTTCACTCAGATGTTTGTCTGCTAC	2040
miR-320a	8p21.3	GCTTCGCTCCCTCCGCTTCTCTCCCGTCTTCTCCGAGTCGGGAAAGCTGGTTGAGAGGGCGGAAAAGGATGAGGT	2041
miR-320b	1q42.11	TGTTATTTTTTGTCTTCTACCTAAGAAATTCGTCTCTTAGGCTTTCTCTTCCAGATTTCCCAAAGTTGGGAAAAGCTGGTTGAGAG GGCAAAAGGAAAAAAGAAATTCGTCTCTGACATAATTAGATAGGGAA	2042
miR-320b	1p13.1	AATTAATCCCTCTCTTTCTAGTTCTTCTTAGAGTGAGGAAAAGCTGGTTGAGAGGGCAACAAATTAACATAATTAATT TGTTTTGAGCGGGGTCAAGAGCAATAACGAAAAATGTTTGTCAATAACCGTTTTTCATTATTGCTCCTGACCTCCTCTCATTTTGCTA	2043
miR-335	7q32.2	TATTCA	2044
miR-34a	1p36.23	GGCCAGCTGTGAGTGTTTCTTTGGCAGTGTCTTAGCTGGTTGTTGTGAGCAATAGTAAGGAAGCAATCAGCAAGTATACTGCCCTA GAAGTGTGCACGTTGTGGGGCCC	2045
miR-34b*	11q23.1	GTGCTCGGTTTGTAGGCAGTGTCAATTAGCTGATTGTACTGTGGTGGTTACAATCACTCACTCCACTGCCATCAAAACAAGGCAC	2046
miR-34c-5p	11q23.1	AGTCTAGTTACTAGGCAGTGTAGTTAGCTGATTGCTAATAGTACCAAATCACTAACCCACACGGCCAGGTAAAAAGATT	2047
miR-371-5p	19q13.41	GTGGCACTCAAACCTGTGGGGCACCTTCTGCTCTCTGTGTGAAAGTGCCGCCATCTTTTGAGTGTTAC	2048
miR-373*	19q13.41	GGGATACTCAAAATGGGGCGCTTTCCCTTTTGTCTGTACTGGGAAGTGCTTCGATTTTGGGGTGTCCC	2049
miR-451	17q11.2	CTTGGGAATGGCAAGGAAACCGTTACCATTAAGTTAGTAATGGTAATGGTTCTCTTGCTATACCCAGA	2050
miR-486-3p	8p11.21	GCATCCTGTACTGAGCTGCCCCGAGGCCCTTCATGTGCCAGCTCGGGCAGCTCAGTACAGGATAC	2051
miR-491-3p	9p21.3	TTGACTTAGCTGGGTAGTGGGGAACCCCTTCATGAGGAGTAGAACACTCCTTATGCAAGATTCCCCTTCTACCTGGCTGGTTGG	2052
miR-498	19q13.41	AACCCCTCCTTGGGAAGTGAAGCTCAGGCTGTGATTTCAAGCCAGGGGCGTTTTTCTATAACTGGATGAAAAGCACCTCCAGAGCT TGAAGCTCACAGTTTGAGAGCAATCGTCTAAGGAAATT	2053
miR-557	1q24.2	AGAAATGGCAAAATGAACAGTAAATTTGGAGGCCTGGGGCCCTCCCTGCTGCTGGAGAAAGTGTTTGCACGGGTGGGCCTTGTCTTT GAAAGGAGGTGGA	2054
miR-638	19p13.2	GTAGCGGGGCGCGGCAGGGATCGCGGGCGGGTGGCGGCCCTAGGGCGCGGAGGGCGGACCCGGGAATGGCGCGCCGTGCGCCG CCGGCGTAAGTCGGCGCT	2055
miR-663	20p11.1	CCTTCGGCGTCCAGCGCGGGCGCCCGGGACCCCTCGTGTGTGGCGGTGGGATCCCCGCGCCGTGTTTTCTCTGGTGG CCCGGCCATG	2056
miR-671-5p	7q36.1	GCAGGTGAAGTGCAGGCCAGGAAGAGGAGGAGCCCTGGAGGGGTGAGGTGATGATGTTTTCTCCGGTTCTCAGGGCTC CACCTCTTTCGGGCCGTAGAGCCAGGGCTGGTGC	2057

miR-744	17p12	TTGGGCAAGGTGCGGGGCTAGGGCTAACAGCAGTCTTACTGAAGGTTTCCTGGAACCCACGCACATGCTGTTGCCACTAACCTCA ACCTTACTCGGTC	2058
miR-885-3p	3p25.3	CCGCACCTCTCTCCATTACACTACCCCTGCCCTCTTCTCCATGAGAGGCAGCGGGGTGTAGTGGATAGAGCACGCGGT	2059
miR-92a-2*	Xq26.2	TCATCCCTGGGTGGGATTTGTTCATTACTTGTGTTCTATATAAAGTATTGCACCTTGTCCTCCGGCCTGTGGAAGA	2060
miR-92b*	1q22	CGGGCCCCGGGGCGGGGACGGGACGGGTGCAGTGTGTTTTTCCCCCGCCAATATTGCACTCGTCCCCGGCCTCCG GCCCCCCCCGCCC	2061
miR-98	Xp11.22	AGGATTCTGCTCATGCCAGGGTGAGGTAGTAAGTTGTATTGTGGGGTAGGGAATTAGCCCCCAATTAGAAGATAACTATACA ACTTACTACTTTCCCTGGTGTGTGGCATATTCA	2062
miR-99a	21q21.1	CCCATTGGCATAAACCCCGTAGATCCGATCTTGTGGTGAAGTGGACCCGCACAAGCTCGCTTCTATGGGTCTGTGTCAGTGTG	2063

5.5 Example 5: Analysis of target RNA from Additional Lung Cancer Cell Lines

[00317] Total RNA was prepared from the eight cell lines (seven lung cancer cell lines and one normal lung cell line) listed in Table 26. Cell lines were purchased from LGC promochem (ATCC) and cultured according to ATCC guidelines.

Table 26: Normal lung and lung tumor cell lines

Cell line	ATCC number	Cancer type
H520	HTB-182	squamous cell carcinoma
H69	HTB-119	small cell lung cancer
H146	HTB-173	small cell lung cancer
H23	CRL-5800	adenocarcinoma; non-small cell lung cancer
NHBE	CC-2540	normal Bronchial epithelial cells
H187	CRL-5804	small cell lung cancer
calu 1	HTB-54	epidermoid carcinoma
calu 3	HTB-55	anaplastic carcinoma

[00318] Microarray data acquisition and analysis were conducted as described in Example 4.

[00319] Total RNA from normal bronchial epithelial cells (NHBE) was used as a control.

[00320] The microarray data was analyzed to identify target RNAs that were present at increased levels in at least four of the cell lines tested. Table 27 shows the target RNAs, the probe sequences used to detect the target RNAs, and the fold-increase in each of the cell lines, relative to NHBE cells.

[00321] The microarray data was further analyzed to identify target RNAs that were present at increased levels in two or three of the cell lines, but for which the average increase in the level of the target RNA relative to NHBE cells was at least 5-fold. Table 28 shows the target RNAs, the probe sequences used to detect the target RNAs, and the fold-change in each of the cell lines, relative to NHBE cells. Table 29 shows the pre-

microRNA sequences and chromosomal location of the pre-microRNA gene for each of the target RNAs in Tables 27 and 28.

[00322] The microarray data was then analyzed to identify target RNAs that were present at at least 5-fold decreased levels in at least four of the cell lines. Table 30 shows the target RNAs, the probe sequences used to detect the target RNAs, and the fold-change in each of the cell lines relative to NHBE cells. Table 30 also shows the number of cell lines in which the same target RNA is present at at least 2-fold increased levels. Table 31 shows the pre-microRNA sequences and chromosomal location of the pre-microRNA gene for each of the target RNAs in Table 30.

Table 27: Target RNAs present at increased levels in at least four cell lines

Gene	Probe sequence	SEQ ID NO	Calu1 FC	calu3 FC	H146 FC	H187 FC	H23 FC	H520 FC	H69 FC
10083-L5-1	CCCTCTTCCTTTCTACCCCTCTCTCCACCC	2064	-1.00	-1.00	6.68	2.01	4.16	4.70	19.19
10333-L5-1	TGCCCTGCCACCCCTCCCTGCCCCG	2065	-8.68	-8.68	1.38	2.28	2.04	2.09	4.83
10398-R5-1	TTGCTACCCGTTTCCCTCCTCTTGAAGTTGCTT	2066	-1.00	-1.00	3.24	1.35	20.09	10.86	13.61
13122-L5-1	TTAGGAAATCCATCTCACCTGCTCCAGTCC	2067	-1.77	1.01	4.68	1.44	4.33	3.39	84.80
13124-L5-1	TCCTCCCTCCGCGAAAGCCTAAACTTACCCCTCA	2068	-1.00	-1.00	5.38	-1.00	5.68	2.02	19.83
13163-R5-4	ACCTCCCTCCTTCTCTCCTCCCAAGACCAA	2069	-1.18	-2.47	4.54	3.26	6.06	2.05	20.32
13185-L5-3	TTCTGTTTTTTTCTTCCCTCTCTCCCTCTCT	2070	-1.99	-1.99	5.80	3.21	12.07	3.25	16.10
13195-L5-3	TCTATATCCCTCCATGACCATTTGTATTAG	2071	-1.00	-1.00	2.03	-1.00	23.44	12.07	6.32
13219-L5-1	CTCTGACTCCCTCACTCAGTCTCTGCTCCAGC	2072	-1.00	1.77	7.06	2.04	9.36	3.89	89.50
13247-L5-2	CTTCCCTCCCGACCTCAGAGCCCTGTTCTTCCT	2073	-1.00	-1.00	2.45	1.33	13.02	3.99	10.40
13334-L5-2	AGGACTCCCTCCCTCCCTCCCACTGTGCCCTGTC	2074	-2.05	-2.05	3.45	-1.05	7.61	2.31	8.48
13335-L5-3	ACCTCAGCCTCCACTGCCCTCCTGCCGATCCTAT	2075	-2.19	-2.19	4.64	5.53	2.19	-1.17	7.44
25-R5-2	TTCTGCTTTCCAGAGCCTCACCCCTCTTTT	2076	-1.00	-1.00	3.62	-1.00	2.55	2.18	4.29
266-R5-2	GTCGCCCTCCCTCCCAAGTTGAGACTTGCA	2077	-2.31	-2.31	3.23	1.50	9.97	5.43	10.49
3249-L5-1	GCCGCCATCCCGGAGCGCGCGCGCGCGCC	2078	1.78	1.99	3.86	81.53	5.02	2.61	69.65
3371-L5-2	CTTTCCTTCTCCCTCCCTCCACACCCCATGACTCCC	2079	-1.00	-1.00	4.01	1.42	12.35	5.21	16.88
3744-R5-1	CTTCTCCTTCTCCTGCTCCCTCCCACTAATGCCAAAT	2080	-2.39	5.28	7.66	12.05	14.66	5.31	107.49
4855-R5-1	CGGGTCTCCCGCTTCCCTCCTGCTCCCAAGG	2081	-5.66	-8.69	3.89	3.18	3.61	2.49	9.85
5107-L5-1	AACTCCCTTTGACCCCCCAGTACAAACTG	2082	-1.00	-1.00	5.89	-1.00	6.60	3.56	35.94
6235-R5-2	TCTGCTCCAAAAATCCATTTAATATATTGT	2083	-1.00	1.77	10.45	7.18	13.80	16.55	207.50
6474-L5-1	CCCTAATTAAAGCCATCCCTCTTCCCTTCACC	2084	-1.00	-1.00	7.06	2.24	8.81	2.57	8.35
6490-R5-3	CCCCATCCCCCATAATGACGCTTCCCCCTCCTAAC	2085	-2.09	-2.09	2.58	-2.09	9.85	6.98	8.49
6681-R2-1	CCTGTTTTCTCCCTCTCTCTGCCCCCTCC	2086	-2.21	-2.21	4.83	2.74	6.51	4.82	19.47
7572-R5-2	ATCACCTTCCCCCTCCCAATAAAGCCAAA	2087	1.39	-1.00	7.08	4.90	27.15	15.82	29.39
7883-R5-1	CTCTCCCTCCCGCCTCTCACCTCCCAACCTATTAGAAAC	2088	-4.54	-2.60	2.34	3.18	3.80	1.61	6.57
8004-R3-2	GGAAGTCTTCTCCTTGCTCCAGTCATTGAAG	2089	-1.00	2.87	4.28	1.99	9.46	6.50	113.48
836-R4-1	AAATAATCATTCCAAATGGTTCTCCCTGCTATGATTCAC	2090	-1.00	-1.00	3.82	-1.00	2.16	2.67	104.03
9594-R5-1	ACTTAGACTTCCTTCCCACTCCCTGCATCCT	2091	-1.81	1.65	3.62	1.98	6.80	5.25	59.46

let-7c	AACCATACAACCTACTACTCA	2092	3.65	24.65	2.65	1.35	3.24	5.11	15.37
let-7d	AACATATGCAACCTACTACTCTCT	2093	2.02	13.30	1.78	-1.94	1.37	4.41	8.00
miR-103	TCATAGCCCTGTACAATGCTGCT	2094	-1.00	-1.00	2.97	3.07	1.27	3.15	3.05
miR-106a	CTACCTGCACTGTAAGCACTTTT	2095	-1.00	-1.00	13.51	23.92	4.32	6.14	17.48
miR-107	TGATAGCCCTGTACAATGCTGCT	2096	-1.00	1.59	4.64	2.94	1.85	7.68	8.76
miR-16	CGCCAATATTTACGTGCTGCTA	2097	-1.00	1.78	6.76	13.51	3.05	1.91	2.91
miR-17	CTACCTGCACTGTAAGCACTTTG	2098	-1.00	-1.00	13.68	23.68	3.51	8.86	18.29
miR-200b	TCATCATTACCAGGCAGTATTA	2099	-1.00	6.04	6.46	36.53	-1.00	7.67	2.72
miR-200c	TCCATCATTACCCGGCAGTATTA	2100	-3.97	5.00	4.38	7.25	-3.97	1.02	2.02
miR-20a	CTACCTGCACTATAAGCACTTTA	2101	512.00	-1.00	6.67	5.00	1.25	2.43	5.02
miR-20b	CTACCTGCACTATGAGCACTTTG	2102	-1.00	-1.00	15.49	7.38	1.90	2.68	8.04
miR-298	TGGGAGAACCTCCCTGCTTCTGCT	2103	-1.00	-1.00	4.26	-1.00	3.08	2.00	49.37
miR-320a	TCGCCCTCTCAACCCAGCTTTT	2104	-1.00	-1.00	5.63	6.24	4.27	-1.00	9.68
miR-320b	TTGCCCTCTCAACCCAGCTTTT	2105	-1.00	-1.00	3.91	3.97	3.28	-1.00	3.98
miR-320d	TCCTCTCAACCCAGCTTTT	2106	-1.00	-1.00	2.33	2.66	2.54	-1.00	3.22

Table 28: target RNAs present at increased levels in two or three cell lines, but with an average increase of at least 5-fold relative to

normal levels

Gene	Probe sequence	SEQ ID NO	Calu1 FC	calu3 FC	H146 FC	H187 FC	H23 FC	H520 FC	H69 FC
10233-R5-1	ACCTCTCCCCCTTGGATCTGGAGCAGCAGGAGTAGA	2107	-2.04	-2.04	3.75	1.08	-1.08	-2.04	10.41
10335-L5-2	CCACTCCCCCTCCTTTTAAATTAGAAAGCAC	2108	-1.00	-1.00	2.43	-1.00	2.56	1.35	12.59
12184-L4-1	GACCTCAGCGTGCCCCCTTTCAACCACAGACGAATATTGTGTACAA	2109	-1.98	-1.98	2.91	-1.03	-1.63	-1.51	9.45
12223-L5-1	GACATCAGACAGAGTTGTTTCTTCTCCCTCTA	2110	-4.85	-4.85	2.43	-1.45	-4.85	-3.28	7.69
12695-R5-2	CCCCCACCAAACCTATTCCCGCATCCTCCCCGG	2111	-2.87	-2.87	2.64	1.50	-1.43	-2.87	7.70
12721-R5-2	TCCACTCTGACCAGCTCACCCCTCACTGGACTA	2112	-1.00	-1.00	2.69	-1.00	-1.00	-1.00	19.11
12729-R5-1	CGACGAAATGCAATTGTGTGCCCTTCTCCCTCC	2113	-1.00	-1.00	2.17	-1.00	-1.00	-1.00	16.72
12888-L5-2	TTTCCTACATTGTATGGTTCTCCAGCTCCT	2114	-1.00	-1.00	1.29	-1.00	2.69	1.51	85.37
12911-L5-1	CAACTGACACCTCCTCCCTTTTCTCCTTCGT	2115	-1.00	-1.00	7.10	2.04	-1.00	-1.00	10.36
12917-R5-2	GGACCTATGGGCCCTTCCCTTCCCCCAACATTG	2116	-1.00	-1.00	4.25	-1.00	2.62	-1.00	10.54

WO 2010/099342

PCT/US2010/025446

12947-R5-3	GCCCCCTCCATAGAGAGAGGCCCCAGGGAGTGGA	2117	-1.00	-1.00	1.37	-1.00	-1.00	4.78	8.65
12992-L5-1	CATACCTGCTTCCCTCCACCCCCATCTCTA	2118	-1.00	-1.00	5.12	1.36	1.24	-1.00	10.93
13001-L5-1	TTCCTAGATACCACTCCAGCTCCA	2119	-1.00	-1.00	2.06	-1.00	1.28	2.46	11.10
13070-R5-3	CTGTCTCCTCTCCCCAGTCCAAAGGACCTAATGC	2120	-1.00	-1.00	5.17	1.43	1.89	-1.00	13.57
13115-L5-3	ACCCCTCCAGGGCCGACCAACCCCAACCCAAAC	2121	-1.00	-1.00	1.59	-1.00	10.07	13.09	12.60
13124-L5-2	GCTCCATGTCTCCTCCCTCCGCGAAAGCCTAAAC	2122	-2.76	-2.76	4.56	-1.18	7.69	-1.73	20.66
13195-L5-2	CCCCCTCCATGACCAATTTGTATTAGTATCTTTT	2123	-1.00	-1.00	-1.00	-1.00	12.25	13.17	10.79
13237-L5-4	ACCCCTCCAGGGCCGACCAACCCCAACCCAAACAA	2124	-1.00	-1.00	1.35	-1.00	8.49	13.59	14.13
13274-L5-3	CCTTCTCTTCTCCCGTGTCTCCACCCCTCCCTCAGGG	2125	-4.16	-3.37	1.81	1.66	3.05	2.03	18.48
13278-R5-2	TCAGACTGTGCTCTCTCCATTCCCCAGGACTCC	2126	-1.00	-1.00	4.29	1.86	-1.00	-1.00	22.61
13287-L5-4	ACTGCAGAGGAGCCACCCACCCCTCTCTCCCAAG	2127	-1.00	-1.00	4.08	-1.00	4.47	-1.00	8.90
13343-L5-4	CTCCCGGCCCTCACCCCTTCCCTGGGCCCTCCACCC	2128	-1.00	-1.00	4.67	1.44	-1.00	1.34	10.16
13357-L5-4	AGCGGACCTTCTCCCCACACCTCCCTGCAGCCTC	2129	-2.19	-2.19	3.44	1.44	-1.81	-2.19	15.06
13358-L5-2	CCTCCTCACCAACCCCTCCACACTCCTGGGAAGT	2130	-1.00	-1.00	1.41	-1.00	5.44	10.83	5.16
13366-R5-3	CAGGCCCTACCCCCAGTGCCCTCTCCTATTCCAC	2131	-1.00	-1.00	2.14	1.46	-1.00	-1.00	19.41
13395-R5-2	CGTCAGACTGTGCTCTCTCCATTCCCCAGGACTC	2132	-1.00	-1.00	4.13	-1.00	-1.00	-1.00	25.57
13403-L5-2	CCCGTGTCCCCCAACCTGGGGCCAGGCCCA	2133	-1.00	-1.00	1.56	-1.00	4.85	1.85	9.80
13467-L5-1	GTGACAGTCAGACCCCTCCTTGCTCCAAGTCAAA	2134	-1.00	-1.00	2.40	1.94	9.75	-1.00	105.32
13470-R5-1	TCCACCCCTCTCCACCTCAGGGACCAGAATCCT	2135	-1.00	-1.00	5.33	1.97	-1.00	-1.00	12.11
13472-L5-1	CTCAGAGCACTGTGTGACGTCTCTCTGTCTGTC	2136	9.44	30.30	-1.00	-1.00	-1.00	-1.00	-1.00
13508-L5-3	GACAGGCTGCTCTCCTTCTGTCTTAGCTTCCT	2137	-1.00	-1.00	2.96	-1.00	-1.00	-1.00	18.27
13530-L5-3	CCTTCCCTCCTTCCCTGGGTCCCCCTAAGTTCTCC	2138	-1.00	-1.00	8.37	5.70	1.56	1.34	16.94
13546-R5-2	TGCCTCCACCCCTCTCCATCCCGTCACCCCTCCCA	2139	-2.75	-2.75	3.08	6.44	-1.36	-2.04	8.01
3875-R5-2	GACTGATTCAACCTCTCTCTCCCACTTTA	2140	-1.00	-1.00	3.37	1.56	-1.00	-1.00	8.56
3923-R5-1	TCACAAAGGATCTCCTTCATCCCTCTCCAG	2141	-1.00	-1.00	1.18	3.17	-1.00	-1.00	8.48
5108-R5-2	CCCTCTCCTCCACACCCGTCACCTCACAATAACCC	2142	-1.94	-1.94	4.49	1.77	1.53	-1.94	12.32
5192-L3-2	CATTTTCCCCCTTCCCTTCTCTATATCAGCAA	2143	-1.00	-1.00	1.79	-1.00	24.42	13.63	4.95
5306_B-R4-1	TCCACTCTGACCAGCTCACCCCTCACTGGACTATACCCC	2144	-1.00	-1.00	3.27	-1.00	-1.00	-1.00	16.02
5633-L5-1	CACTCCCTCCCTGGCTCCTGGAGACCCCTCTCCAGACTTG	2145	-1.00	-1.00	2.22	2.39	1.27	-1.00	45.28
5638-R5-2	GGCCCTCCCCCTGCCTGTGATAGGCTG	2146	-1.57	-2.40	3.23	-1.15	5.85	-1.33	8.54
5723-R5-1	CCTCCTCCCCCTTCTCCAGCAGTAGCCTTCTTAA	2147	-1.00	-1.00	4.87	1.40	3.97	-1.00	11.99
5735-L5-1	TGCTGTTTCTCTGGCCTCAAGCCTGTGGGGG	2148	-1.00	-1.00	1.18	3.00	-1.00	-1.00	14.30

WO 2010/099342

PCT/US2010/025446

6183-R5-1	ACTTTTCTTAATGACTTTCCTCCCTCCTTAAG	2149	-1.00	-1.00	1.75	-1.00	18.55	12.12	10.01
6216-R5-1	TACTCCCGCGGTTTACCCGTGCTTCATTGAA	2150	-3.03	-3.03	2.63	13.19	1.06	-3.03	-1.10
6490-R5-3	CCCCATCCCCCATATGACGCTTCCCCCTCCTAAC	2151	-3.21	-4.42	-2.06	-1.29	5.92	3.87	5.51
6496-R5-2	CCCCCTCCCCCACCCACCACTTCCCCCTAGA	2152	-2.49	-5.45	1.82	-1.65	4.91	-1.83	10.29
6584-L5-1	GTCGGCCCTGCCCTCCTCCTCCTCTCACCAAGC	2153	-1.00	-1.00	3.33	-1.00	-1.00	-1.00	12.02
6647-R5-1	CCCCAGCTGGAGAAATTTTCCCCTCATTA	2154	-1.00	-1.00	-1.00	-1.00	12.90	5.66	6.72
6752-R5-2	CCCTCCTTTCCCCACCTCAGTCGGGCCACTGCT	2155	-1.00	-1.00	2.70	1.92	5.08	1.27	9.54
6803-R5-2	GCTCCCTCTCTGGTTGGACCTCACCCAAA	2156	-2.35	-2.35	5.77	3.27	-2.35	-2.35	13.06
6930-R5-1	TTAATCCTTCTCTCCCCCTCTGATCTTGCAAG	2157	-1.00	-1.00	7.92	1.50	2.40	-1.00	28.87
7113-R5-1	CGCCTAATTAGCCCCCCTGCTCCGGAGGCCCTCACC	2158	1.06	-2.16	2.60	-2.16	1.12	-1.18	12.56
7192-R5-1	TTCCCCCTCTAAGTCTGCCTGGGCTCTTTGGCACTG	2159	-1.00	-1.00	-1.00	-1.00	5.96	13.37	21.66
7384-R3-1	CTCGCAAAGGATCTCCTTTCATCCCTCCCCA	2160	-1.00	-1.00	1.30	-1.00	2.58	-1.00	9.23
7571-L5-1	CAGGGCTAACAGGGCTCCCCCACCCCTAAG	2161	-2.26	-2.26	4.35	-2.26	-1.81	-2.26	6.92
836-R5-2	AAATAATCATTCCAAATGTTCTCCCTGCTAT	2162	-1.00	-1.00	3.09	-1.00	2.40	1.86	51.66
8433_C-R4-1	AAACCAAAAAAAAAAAATTAAAAAGCGACGAAAATGCAATTGTGTGCCTTCTCCCTCC	2163	-1.00	-1.00	5.09	-1.00	-1.00	-1.00	16.86
8433-L3-1	AAATGGCTCCTTTCCCTCTTCCCTCCACCG	2164	-1.00	-1.00	6.04	1.95	5.50	1.32	11.98
8724-R5-2	GGCCAAAGCTTGGAACCTCTCCCTGCCAGCA	2165	-1.89	-1.89	2.12	1.04	-1.89	-1.89	7.98
8808-R5-1	GACCCCTTCTCCCGAGCCTGTTTCTGCAA	2166	-1.95	-1.95	2.65	1.01	-1.95	-1.95	9.58
8832-R5-1	TGGAGTACCACCTGTTTTTCCCCCACTT	2167	-1.00	-1.00	-1.00	-1.00	5.05	1.41	21.96
9349-R5-2	GAACACAGTGATGCAGAGGACTTCTGCTCCA	2168	-1.00	-1.00	1.84	-1.00	1.99	1.61	85.75
let-7b	AACCACACAACCTACTACCTCA	2169	-1.24	11.28	1.29	1.73	-1.24	2.39	6.09
let-7e	AACTATACAACCTCCTACCTCA	2170	-1.27	12.32	1.35	-1.94	1.66	1.71	7.50
miR-1268	CCCCCACCACCAAGCCCG	2171	-1.00	-1.00	2.84	-1.00	-1.00	1.37	11.23
miR-1274b	TGGCGCCCGAACAGGGA	2172	6.00	-1.00	9.59	-1.00	1.21	-1.00	-1.00
miR-15b	TGTAAACCATGATGTGCTGCTA	2173	-1.00	-1.00	2.40	1.66	-1.00	8.53	-1.00
miR-181a	ACTCACCAGACAGCGTTGAATGTT	2174	-1.00	4.65	1.33	-1.00	1.20	-1.00	6.04
miR-198	GAACCTATCTCCCTCTGGACC	2175	-1.00	-1.00	3.07	-1.00	-1.00	-1.00	17.81
miR-222	ACCCAGTAGCCAGATGTAGCT	2176	2.68	9.77	-1.96	-1.96	-1.00	-1.96	-1.96
miR-26a	AGCCTATCCTGGATTACTTGAA	2177	-1.00	4.43	1.73	6.73	1.20	-1.00	4.27
miR-30d	CTTCCAGTCGGGGATGTTTACA	2178	-1.00	9.69	1.71	7.03	-1.00	-1.00	-1.00
miR-720	TGGAGCCCCCAGCGAGA	2179	5.56	-1.00	4.48	-1.00	-1.00	-1.00	-1.00
miR-92a	ACAGGCCGGGACAAAGTGCAATA	2180	-1.00	-1.00	3.74	10.88	-1.00	-1.00	7.26

6474-L5-1	1p34.1	GGTGAAGGGGAAGAGGGGATGGCTTTAATTAGGGCTTCTTGGCCCTCGTCAGTCCTGGGGTTGGTCGGGGTAATCCAGTTAGAGTCCCAGCCTTTCATCTCAGCTGGCC	2204
6490-R5-3	1q21.3	TCCTTCCCCCTTCGTGGCTTGCGGTCTCTCTTCCCCGCCCTCGGCCCTCAGGAAAGTGTGAGTGTGGGGTGTGAGGTTAGGAGGGGAAGC	2205
6681-R2-1	11q12.2	GTCATATGGGGATGGGG	2206
7572-R5-2	11q12.1	TGTGCTCTCATTTGTTATTCCAAAGTCTCTGTCTAGATCACTGGAGGGCAGAGAGAGGGAGAAACAGGGAGATACA	2207
7883-R5-1	11q14.2	ATCACCTTTCCCCCTCCCATGTGCTTTCCCTTCATTTGAGATCTTTTGACCTTTGGCTTTATTTGGAGGGGAAGGTGAT	2208
8004-R3-2	Xq28	CTCTCCGTGCTTCTCGGCCCGCCGCCCTTCCCTGTCTCGGGACACGGTTTCTAAATAAGTTGGAGGGTGAGAGGGGAGGAGAG	2209
836-R4-1	3q26.2	GGGGCTGCCATCCTGCTGTCCGTGTCATCTGTGTGGTGTGCTGACGGCCTTCAATGACTGGAGCAAGGAGAACAGTTC	2210
9594-R5-1	2q12.1	AAATAAGCCATTCCAAACCATTCTCTGATTTGCTGTGAGTGGCAGAAATCATTCACCGTGGTGAATCATAGCAGGGAGAACCATTTTGAATGATTA	2211
let-7c	21q21.1	TTCCAGCTATTTAGTAACTCTTCCAAACACACTGTCAGCACCCCATGCTAGGATGCAGGGAGTGGGAAGAAAGTCTAAGTAGGGAA	2212
let-7d	9q22.32	GCATCCGGGTTGAGGTAGTAGGTTGTATGTTAGAGTTACACCCCTGGAGTTAACTGTACAACCTTCTAGCTTTCCCTTGGAGC	2213
miR-103	20p13	CCTAGGAAGAGGTAGTAGGTTGCATAGTTTTAGGGCAGGGATTTTGGCCACAAGGAGGTAACATAACGACCTGCTGCTTTCTTTAGG	2214
miR-103	5q35.1	TTGTGCTTTTACAGCTTCTTTACAGTGTGCTGCCCTGTAGCATTCAGGTCAAGCAGCATTTGTACAGGGCTATGAAAGAACCA	2215
miR-106a	Xq26.2	TACTGCCCTCGGCTTCTTTACAGTGTGCTGCCCTTGTTCATATGGATCAAGCAGCATTTGTACAGGGCTATGAAAGGCATTG	2216
miR-107	10q23.31	CCTTGGCCATGTAAAAGTGCTTACAGTGCAGGTAGCTTTTTGAGATCTACTGCAATGTAAAGCACCTTCTTACATTACCATGG	2217
miR-16	13q14.3	CTCTCTGCTTTTACAGCTTCTTTACAGTGTGCTTGGCATGGAGTTCAAGCAGCATTTGTACAGGGCTATCAAAGCACAGA	2218
miR-16	3q26.1	GTCAGCAGTGCCTTAGCAGCACGTAATAATTGGCGTTAAGATTCTAAAATTATCTCCAGTATTAAGTGTGCTGCTGCTGAAGTTGAC	2219
miR-17	13q31.3	GTCCACTCTAGCAGCAGCTAAATATTGGCGTAGTGAAATATATATAACACCAATATTACTGTGCTGCTTTAGTGTGAC	2220
miR-200b	1p36.33	GTCAGAAATAATGTCAAAGTGCTTACAGTGCAGGTAGTGATATGTGCATCTACTGCAGTGAAGGCACCTTGTAGCATTATGGTGAC	2221
miR-200c	12p13.31	CCAGCTCGGGCAGCCGTGGCCATCTTACTTGGCAGCATTTGGATGGAGTCAGGTCTCTAATACTGCCTGGTGAATGATGACGGCGGAGCCCTGCA	2222
miR-20a	13q31.3	CG	2223
miR-20b	Xq26.2	CCCTCGTCTTACCCAGCAGTGTTTGGGTGCGGTTGGGAGTCTCTAATACTGCCGGGTAAATGATGGAGG	2224
miR-298	20q13.32	GTAGCAGTAAAGTGCTTATAGTGCAGGTAGTGTAGTTATCTACTGCAATATGAGCAGCTTAAAGTACTGC	2225
miR-320a	8p21.3	AGTACCAAAGTGCTCATAGTGCAGGTAGTTTGGCATGACTCTACTGTAGTATGGGCACCTTCCAGTACT	2226
miR-320b	1q42.11	TCAGGTCTTCAGCAGAAAGCAGGGAGGTTCTCCAGTGTTTCCCTTACTGTGAGGAACTAGCCTGCTGCTTGTCTCAGGAGTGAGCT	2227
miR-320b	1p13.1	GCTTCGCTCCCCCTCCGCTTCTCTTCCCGGTTCTTCCCGGAGTCGGGAAAGCTGGGTTGAGAGGGCGGAAAAAGGATGAGGT	2228
miR-320d	13q14.11	TGTTATTTTTTGTCTTCTACCTAAGAATTCTGTCTCTTAGGCTTTCTCTTCCCAGATTTCCCAAGTTGGGAAAGCTGGGTTGAGAGGGCAAAAG	2229
10233-R5-1	9q34.13	GAAAAAAAAGAAATTCGTCTGACATAATTAGATAGGGAA	2230
10335-L5-2	15q26.1	AATTAATCCCCTCTCTTCTAGTTCTTCCCTAGAGTGAGGAAAGCTGGGTTGAGAGGGCAACAAATTAATAATT	2231
		TTCTCGTCCCAGTTCTTCCCAAAGTTGAGAAAAAGCTGGGTTGAGAGGA	2232
		TTCTCTTCCCAGTTCTTCTTGGAGTCAGGAAAAAGCTGGGTTGAGAGGA	2233
		ACCTTCTGCCCTTGTAGACCTTGAAGCCACAGCAGCTGGTGGACATAAATTTACTGCTGCTCCAGATCCAAAGGGGAGAGGGT	2234
		TCTTAGTGTCTTCTAATTAAAAAGGAGGGGAGTGGTGATCTTTTGTCTCTCTAAGTTCTGTTTCCCTCTGAGTGGAAAGCAGAGGG	2235

12184-L4-1	3p14.1	TTGTACACAATATTCGTCGTGGTTGAAAGGGGGCACGCTGAGGTCAAGTGAAGTAGTGTGTTTTCCATTTTCCATAATGAGTCTCACAGTGTGCGA	2233
12223-L5-1	4q27	TAGAGGGAGAAAGAAACAACCTCTGTCTGATGTCTTCTGGATGCGCTTAATACAGATAGCATTTGCTCTTCCATTTCG	2234
12695-R5-2	1q22	TCCCCGCCACCCCTTGAGCACCTCAGCGTTCTTAGGGGAAGCCAGAGCCGGGAGGATGCGGGAATAGGTTTGGTGGGG	2235
12721-R5-2	12q13.13	TTCAACCATGACCCCTTGCTCCTCACCCCCACCCAGGGTCCATAGTCCAGTGAGGCTGGTCAGAGTGGA	2236
12729-R5-1	17q25.3	AAATTAAAGAAAAAAATTTCTCCACCAAAAGCGCGCAGTGACAGTTTCCCAACATTTTCTGCCCTTCTTCTTCCCTCCCTGCACCACTAGGAGG	2237
12888-L5-2	2q32.3	GAGAAGGCACACAATTGCATTTTCGTCGCTTTTAAATTTTTTTTTTGTTT	2238
12911-L5-1	22q12.3	TCCTCCAGGAGCTGGGAGAACCATACAAATGTAGGAAAAATATAGTTTAAATTGAATGGTACTCTGGTCTTCTGGAGGA	2239
12917-R5-2	1p34.1	ACGAAGGAGGAAAAAGGAGGTGTCAGTTGGACTGCCCAAGGCAACCCCTTGAGCCATCAGGACAGCTCAACACCTCTCTGTCTTCTTCTT	2240
12947-R5-3	6p21.1	TGT GGACCTGGGGCTTCTCTGACCCCTTGAACAGCTTATACTATGAGACCTTGGGAACCTCCTCCATGCAGACACACAAGGCTCAATGTTGGGGA	2241
12992-L5-1	9q31.3	AGGAAAGGGCCCATAGGTCC	2242
13001-L5-1	12q13.13	GCCTTGTCTGAAGGGAGAGGCCCTGGCATGCGGATGGGAGATTIAGAGGCTGTGGAGAAGGGAACCTTGGGGCTTTCCTTCCCTTCGTGGCCTC	2243
13070-R5-3	5q35.1	ACTCCCCTGGGGCTCTCTCTATGGAGGGGC	2244
13115-L5-3	1q21.3	TAGAGATGGGGGTGGAGGGAAGCAGGTATGATTTACGGGGCAGCAGGAGATTCTCTTCTGTTTCCCTCTCTCCAGCTCTA	2245
13124-L5-2	1q22	TGGAGCTGGGAGTGGTATCTAGGAATTTCTCCCTTAGTTTTACCTTCTTAGTTTTA	2246
13195-L5-2	3p21.1	CTGCTCTCCTTGACCTGGTAGGAAGTGATTTGATGGTTTTCAAAACACATGCTGTTTCTCTCTTTTCATCAGGGTAGCTGGGTGGCATTAGG	2247
13237-L5-4	1q21.3	TCCTTTGGACTGGGAGAGGAGACAG	2248
13274-L5-3	12q13.13	AGGAAAGGCAAGGTAGCCATTGTTGGGTGGGTGTCGCCCCCTGGAGGGGTTTGTGTTGCTTATCCCCCTCTGTGCTTACCCCCCTACCCCTACCC	2249
13278-R5-2	22q11.21	AGCTCCCCGCCAGAAAGCCCTG	2250
13278-R5-2	12q13.13	TGAGGGGTAAAGTTTAGGCTTTCGCGAGGGGAGGAGACATGGAGCCTGGGAACCTCTGTTCTCCCCCTCTGCCCTCTCCCCACCCCTTA	2251
13287-L5-4	1p35.2	AAAAAAGGAAAGATACTAATAACAAATGGTCATGGAGGGGGAATATAGAGAAGATCAATTTGTACAGAAAAACCAATTGGTTAGTATTTTTTTTCT	2252
13343-L5-4	17q25.3	TTT GGCAAGGGTAGCCATTGTTTGGGTGGTGGCCCCTGAGAGGGGTTTGTGTTGCTTATCCCCCTCTGTGCTTACCCCCCTACCCAGCTTCC	2253
13357-L5-4	19q13.2	CGCC	2254
13358-L5-2	19q13.32	AGTGGTGGTGGGAGACCCCTGAGGGAGGGTGGAGCAGCAGTAGCAGTACCAGATGGTTCTGTGGAGTCCCTGGGGAATGGAGAGAGCA	2255
13366-R5-3	19p13.2	CAGTCTGACTCCCTGCCAAGTAGCCAG	2256
13395-R5-2	22q11.21	TCCGCCGAATAAATCTCCATGTGGGTCTGGGAGGGTGGGTGGCTCCTCTGCAGTGAGTAGGTCTGCCCTGGAGCTACTCCACCATCTCCCC	2257
		CAGCCCCGTATGGCTGGGAGGGG	
		TGAGGGGGAGGGGCGCTGCGGGAGGGGTGGAGGGCCCCAGGGAAGGGTGAGGGGCCGGGAGCCACTCTGCCCGGCACCTCTCCGCCCCAGAA	
		ACAGCCCCAACGCCCTTCTTTCCCCCTTT	
		TCAGGGTTTGGTGACAAATTGTGGGAGGCTGCAGGAGGTGTGGGAGAAAGGTCCGCTTTAGAGCTTGGTCCCTCTGTGCTCACCTCCCAGCC	
		CCAGCCCCCAGCACCGGATCTG	
		CGGCCAGTGACTTCCCCAGGAGTGTGAGGGGGTGTGAGGAGGAGCACCTGGGCTCTCTACCCCTCTCCTCACAGAAGTACCTGAAACTAG	
		GTC	
		GGCCCGGTGAAGGCCCCCGCTGGGTCCCATAACCCGGGTTGGGGTCCAGAACCCGCTCGGTCTCTGTGGGAATAGGAGAGGGCACTGGGGT	
		GAGGCCCTG	
		TGGTTCTCTGTGTTTTGTGGACTGTGCTCTCACTGTTACCCAGCACTAGCAGTACCAGACGGTTCTGTGGAGTCTCTGGGAATGGAGAGAGCA	

219

7571-L5-1	2p21	CTTAGGGGTGGGAGCCCTGTTAGCCCTGTAAATAAAGTTTAACGAGGTGAACAATGGCTGGCTCTGTCCCTGAG	2286
836-R5-2	3q26.2	AAATAAGCCATTCCAAACCATTCTCTGATTTGCTGTGAGTGGCAGAAATCATCCCGTGGTGAATCATAGCAGGGAGAACCACTTTGGAATGATTA TTT	2287
8433_C-R4-1	17q25.3	AAATTAAGAAAAAATTCTCCACCAAAAGCGCCGAGTGACAGTTCCTCAACATTTTCTGCCTTTCTTCTCCCTCCCTGCACCACTAGGAGG GAGAAGGCACACAATTGCATTTTCGTCGCTTTTAAATTTTTTTTTTTGGTTT	2288
8433-L3-1	17q25.3	CGGTGGAGGGAAAGGGGAAAGGAGCCCATTTTCTGCTGCACATCAGTCAGTGCCTCGCCCTCCCTCCCGCCG	2289
8724-R5-2	15q23	GGCCCAAGAGATGAAAAGCTGAAGTCCTTTCCCTTCCAGCTGAAGCCAGGTGTGATGCTGGCAGGAGAGGTTCCAAAGCTTGGCC	2290
8808-R5-1	3p14.3	CCGATTATGGCTTTCTTCTCTGCTGCCCCCTTTCAGTAGTGATTTGCAGAAACAGGCTGGGAGAAAGGGGTCTTTTGG	2291
8832-R5-1	9q33.2	TTCTGAGATATGATCTGTTGGATTCTCTACTACCAAAAGTGGGGGAAACACAGGTGGTACTCCAGAA	2292
9349-R5-2	21q22.11	GGACACTCTGAACCCCAAGTGGAAATCCAACTGCCAGTTCTTCATCCGAGACCTGGAGCAGGAAGTCTCTGCATCACTGTGTTT	2293
let-7b	22q13.31	CGGGTGAGGTAGTAGGTTGTGTGGTTTCAGGGCAGTGATGTTGCCCTCGGAAGATAACTATACAACCTACTGCCCTTCCCTG	2294
let-7e	19q13.33	CCCGGGCTGAGGTAGGAGGTTGTATAGTTGAGGAGGACACCCCAAGGAGATCACTATACGGCCTCCTAGCTTTCCCCAGG	2295
miR-1268	15q11.2	TAGCCGGCGGTGGTGGGGCCCTGTGGTCCCAGCTACTTTGGAGGGCTGAG	2296
miR-1274b	19q13.43	CTTCTTCACTCACGTCCCTGTTTCGGGGCCCACTTGTGGCTGCGGTTCCGGACTGAATGAAGAAGGA	2297
miR-15b	3q26.1	TTGAGGCCCTTAAAGTACTGTAGCAGCACATCATGTTTACATGCTACAGTCAAGATCGGAATCATTTATTGCTGCTCTAGAAAATTAAGGAAATTC AT	2298
miR-181a	1q31.3	TGAGTTTTGAGGTTGCTTTCAGTGAACATTCAACGCTGTCCGGTGAGTTTTGGAATTAATAAATCAAAACCATCGACCGTTGATTGTACCCCTATGGCTAA CCATCATCTACTCCA	2299
miR-181a	9q33.3	AGAAGGGCTATCAGGCCAGCCCTTCAGAGGACTCCAAGGAACATTCAACGCTGTCCGGTGAGTTTGGGATTTGAAAAAACCACTGACCCGTTGACTG TACCTTGGGGTCCCTTA	2300
miR-198	3q13.33	TCATTGGTCCAGAGGGGAGATAGGTTCCGTGATTTTTCTTCTTCTCTATAGAATAAATGA	2301
miR-222	Xp11.3	GCTGCTGGAAGGTGTAGGTACCCCTCAATGGCTCAGTAGCCAGTGTAGATCCTGTCTTTCGTAATCAGCAGCTACATCTGGCTACTGGGTCTCTG ATGCGATCTTCTAGCT	2302
miR-26a	3p22.2	GTGGCCTCGTTCAAGTAATCCAGGATAGGCTGTGCAGGTCCCAATGGGCCTATTCTTGGTTACTTGCACGGGACGC	2303
miR-26a	12q14.1	GGCTGTGGCTGGATTCAAGTAATCCAGGATAGGCTGTTTCCATCTGTGAGGCCCTATTCTTGATTACTTGTCTGGAGGCAGCT	2304
miR-30d	8q24.22	GTTGTTGTAACATCCCCGACTGGAAGCTGTAAAGACACAGCTAAGCTTTCAGTCAGATGTTTGTCTGCTAC	2305
miR-720	3q26.1	CCGGATCTCACACGGTGGTGTAAATATCTCGCTGGGGCCCTCCAAAATGTTGTGCCCAGGGGTGTAGAGAAAACACACACACTTTTGAGATGAATT AAGAGTCCTTTATTAG	2306
miR-92a	13q31.3	CTTTCTACACAGGTTGGGATCGGTTGCAATGCTGTGTTTCTGTATGTTGCATTGTCCCGGCCTGTTGAGTTTGG	2307
miR-92a	Xq26.2	TCATCCCTGGGTGGGATTTGTTGCATTACTTGTGTTCTATATAAAGTATTGCACCTGTGCCCGCCTGTGGAAGA	2308
miR-92b*	1q22	CGGGCCCCGGCGGGGACGGGACGGGTGCAGTGTGTTTTTCCCCCGCCAATATTGCACCTCGTCCCGGCCTCCGGCCCCCCCC GGCCC	2309
miR-936	10q25.1	TCAAGGCCCACTGGGACAGTAGAGGGGAGGAATCGCAGAAATCACTCCAGGAGCAACTGAGAGACCTTGCTTCTACTTTTACCAGGTCTGCTGGC CCAGA	2310
miR-98	Xp11.22	AGGATTCTGCTCATGCCAGGTGAGGTAGTAAGTTGTATTGTTGTGGGTAGGGATATTAGGCCCCCAATTAGAAAGATAACTATACAACCTTACTAC TTTCCCTGGTGTGGCATAATTCA	2311

Table 30: Target RNAs present at at least 5-fold decreased levels in at least four cell lines

Gene Down expressed in Cell lines	Probe sequence	SEQ ID NO	Calu1 FC	calu3 FC	H146 FC	H187 FC	H23 FC	H520 FC	H69 FC	Number of cell lines with at least a 2- fold increase
10010_B-L4-1	GCCGAGCCCCCGCCCCCGGGGATGCTGCCCTCCGGAA GGAGGGCGCTGCCC	2312	-16.79	-8.41	-4.11	-2.74	-7.08	-22.10	-2.85	0
10010_D-L4-1	TGGCGCCCTCCCCCGCCGGGCTCAGCCTCTCACCTG	2313	-30.70	-30.70	-2.08	-4.60	-3.76	-16.38	-1.22	0
10138-L2-1	AGCCTGCTCCGCTCTCCCTCCACCCAGAAAAAGT	2314	-7.46	-7.46	1.75	-1.81	1.87	-2.04	7.73	1
10231-R3-1	TGAACTTTAGCTGGCGCGCGCCTGTCAGC	2315	-29.14	-29.14	-3.13	-4.48	-29.14	-29.14	-5.84	0
10242-R5-1	GAAGCCCTTCCGCTTCCACCCCGAACAC	2316	-25.94	-25.94	-14.00	-25.94	-25.94	-25.94	-8.13	0
10342-R2-2	CCCGCCCGGAGCATCTCGAAGTTAATTAA	2317	-28.71	-3.68	-4.01	-2.19	-10.93	-50.54	-3.01	0
11370-L4-1	CCTCCGCCCCCACATGCATCCTTGCCCAGTTTGGCTGCCA TCAGTATTGTCCCTGAGAACTGGAC	2318	-5.20	-5.20	1.61	-2.49	-2.57	-2.80	3.46	1
11370-L5-5	CCCTCCGCCCCCACACTGCATCCTTGCCCCAGTTTGG	2319	-5.86	-5.86	1.74	1.95	1.55	-3.00	4.02	1
12223-L4-1	CCCAGAAACATCAGACAGAGTTGTTTCTTCTCCCTCTA	2320	-5.20	-5.20	2.99	-2.63	-5.20	-5.20	5.56	2
12691-R5-1	CCCCGCCCCTGCGCGCCCCCGACAGGC	2321	-46.43	-46.43	-6.02	-32.36	-46.43	-46.43	-3.50	0
12692-L5-1	GACCCGGCCCCGAGCCAGCACCCGGCCACCGCGC	2322	-7.57	-7.57	1.00	-5.34	-7.57	-7.57	1.21	0
12693-L5-1	GTCGGGGCGCCCGCCCTCCCGGTCCCCTCCCC	2323	-34.28	-34.28	-4.19	-6.29	-2.25	-14.44	-1.05	0
12694-R5-1	TCAGCCCCCAGCGCCCCCGGAGTCTTGGA	2324	-85.43	-85.43	-17.02	-29.82	-85.43	-85.43	-6.11	0
12695-R5-1	AAACCTATTCCCGCATCTCCCGGCTCTGGC	2325	-44.55	-44.55	-3.70	-3.25	-13.08	-32.36	-1.29	0
12696-R5-1	CCGGGCTCCCCACCCGCTCCCTGAGC	2326	-9.47	-9.47	1.52	-3.26	-4.38	-9.47	2.17	1
12696-R5-2	AACCCGGGCTCCCCCACCCGCTCCCCTGA	2327	-38.89	-38.89	-2.62	-3.23	-12.87	-26.21	1.03	0
12697-R5-1	CCGGTGTGCGCCCCCTCCTACCTCTGCCGGCC	2328	-26.89	-26.89	-5.74	-13.37	-13.11	-5.78	-1.74	0
12699-L5-1	GGCGCCCTGGGCCTCGGCGCCCCCGCCGTCCCAG	2329	-9.80	-9.80	-2.20	-4.84	-9.80	-9.80	-1.77	0
12701-L5-1	AAATCCTCGCCATCCTCCACCCCCAGCCCCCG	2330	-24.62	-24.62	-2.85	-2.72	-5.56	-12.45	-1.45	0
12703-L5-3	AGCCGAGCCCCCGCCCCCGCGGATGCTGCCCTC	2331	-64.38	-60.80	-7.33	-2.94	-20.92	-66.02	-9.82	0
12704-L5-1	CCTCCCCCGCCGGGCTCAGCCTCTCACCTG	2332	-8.28	-8.28	1.19	-1.16	-1.79	-4.47	2.96	1
12704-L5-2	CTGGCGCCCTCCCCCGCCGGGCTCAGCCTC	2333	-7.28	-7.28	-1.14	-1.77	-1.61	-5.51	2.57	1
12713-R5-1	CTCTCGCGACCGACCTGCCGCCGACCGCCACAG	2334	-43.69	-9.13	-3.01	-2.22	-23.87	-33.93	-2.33	0
12722-L5-1	AAACAAAGTACTTCGACCTCCCCGCCGCCCGC	2335	-34.34	-6.02	-2.48	-2.05	-6.64	-8.13	-1.75	0
13004-R5-1	CTCCAACCCCCGGCAATTCTGCTCCCTTCACCTGA	2336	-35.18	-35.18	-5.05	-10.62	-35.18	-35.18	-2.20	0

13006-R5-4	CCCGGCTCTCCCTCCCCTGCGCGCGCTCTCGCC	2337	-13.41	-29.15	-1.35	-3.15	-4.57	-3.39	1.17	0
13044-L5-3	CCCATCCAGCTGTCCCTTTCTTTGCTTTCATCA	2338	-23.71	-23.71	-17.84	-12.27	-23.71	-23.71	-4.70	0
13044-L5-4	CCAGGGCTTCTCCCATCCAGCTGTCCCTTCTTTG	2339	-30.11	-30.11	-4.72	-5.11	-30.11	-30.11	-2.34	0
13047-R5-2	GCACAGCAGACCCCATGCACTAGCCCCGGGCAC	2340	-19.29	-19.29	-19.29	-19.29	-19.29	-19.29	-19.29	0
13052-L5-1	TCCCCGACCTAAGCATCTCCCCACCCGCGCAACC	2341	-6.46	-6.46	1.30	-2.05	-2.52	-4.72	4.47	1
13093-L5-2	CGACCCCGCAGAAACCCACCGCGCCCCGCGCAG	2342	-45.57	-45.57	-16.39	-23.29	-45.57	-45.57	-10.53	0
13097-L5-2	AGCCTCAGCCCCACCTCCAGCCCCACCCTAGGG	2343	-7.94	-7.94	-1.16	-3.99	-5.23	-7.94	1.33	0
13106-R5-1	ATCAGAGCCGACCCCGCGGTCCAGGCCGGCA	2344	-37.32	-37.32	-26.17	-37.32	-37.32	-37.32	-24.68	0
13106-R5-2	TGACCCAATCAGAGGCCGACCCCGCGGTCCAG	2345	-42.08	-42.08	-35.48	-42.08	-42.08	-42.08	-9.62	0
13111-L5-3	TCTCCGCCGGGCTTACACCTGCCCTGCTCTTCT	2346	-8.41	-8.41	1.41	1.94	-4.26	-6.40	-1.19	0
13119-R5-2	GACTCTGCCGCTCCCGCCCGGCCACCTCCCTGT	2347	-6.27	-5.25	-3.10	-2.23	-8.08	-8.44	-2.59	0
13129-L5-2	GTCCCCTGGCCCCCGACCTGTCTCATCCACCCA	2348	-10.72	-10.72	1.69	-2.03	-7.75	-4.28	4.56	1
13129-L5-3	GTCCCCTGGCCCCCGACCTGTCTCATCCACCCA	2349	-10.72	-10.72	1.69	-2.03	-7.75	-4.28	4.56	1
13129-L5-3	AGCCTTCTGTCCCCTGCCCCCGGACCTGCTCCA	2350	-8.92	-15.69	-1.15	-1.02	-5.81	-15.69	3.68	1
13130-L5-1	AGCCCGCCCCAACCACCTCGATCTTTTCCCTC	2351	-2.96	-5.75	-1.02	-2.38	-7.72	-7.44	-1.49	0
13130-L5-2	TCCCCAGAGCCCCCGCCCCAACCCACCTCGATCT	2352	-19.17	-11.28	-2.91	-4.08	-19.00	-28.42	-1.70	0
13137-L5-1	GCTCTAACCCCCCGCAACCCACCTCCCCATGCC	2353	-6.42	-6.42	1.22	-1.92	-2.51	-6.42	2.70	1
13138-R5-1	TTGCCCAGCCCCCGCCACCCGAGCTGCCTCCC	2354	-22.78	-6.06	-3.57	-2.48	-3.75	-4.73	-2.04	0
13161-L5-4	TCCGCCAGGGTCCGCGTGTGAGTCCCCCTCTGGTGA	2355	-60.36	-106.72	-106.33	-4.55	-	-	-	0
13209-R5-3	TGGTCGCCCGCCGACGGCGCTGAAGGGCACGGCGG	2356	-23.10	-23.10	-23.10	-23.10	-23.10	-23.10	-14.65	0
13211-L5-1	ACGCGCCCCCGCGCTCTCTGACCGACCCGAGGCGC	2357	-33.55	-33.55	-13.40	-24.17	-33.55	-33.55	-8.31	0
13227-L5-1	CTCCTCGTCCCCCTTCCCACCTCGGTGCTTGCTT	2358	-25.42	-25.42	-3.92	-12.50	-2.75	-12.60	-4.39	0
13227-L5-2	GGCCCCAGTCTCTCTCGTCCCCCTTCCCACCTCGG	2359	-24.86	-38.26	-4.08	-9.37	-2.60	-5.71	-1.14	0
13229-L5-1	ACCGTCCCTGCCCTTTACCCCTTGGGCCAGCA	2360	-6.14	-6.14	1.13	-3.09	-3.21	-4.58	2.42	1
13229-L5-2	CCTGGGGCCACCCGTCCCTGCCCTTTACCCCTT	2361	-7.44	-7.44	-1.25	-7.44	-7.44	-7.44	3.22	1
13229-R5-3	GCAGCTCCGCCAGTCTCTGTGGCAGGGAGAAG	2362	-29.88	-18.41	-18.21	-1.79	-29.88	-29.88	-6.56	0
13239-L5-2	CAGAGCTCCCCCATCTCCCCAGACTTACCCCT	2363	-5.36	-5.36	-1.08	-3.59	-3.58	-5.36	3.91	1
13240-L5-2	AATCGCCGTCCCCCGCGCGGCATTCCCGGCCCCAA	2364	-28.53	-28.53	-2.82	-2.43	-28.53	-21.92	-1.98	0
13247-L5-3	TCCTGAGCCGCTTCCCCTCCCGACCTCAGAGCCCT	2365	-13.68	-19.46	-1.59	-1.37	-1.28	-2.19	1.48	0
13267-L5-1	CACTCCCTGCTGGCCCCCCCACCTCACCTATGGTG	2366	-58.86	-58.86	-9.66	-17.90	-7.03	-39.11	-1.08	0
13276-L5-4	CCCCCATTTGTGGCTGCTCCACCCCCACCTGCCTTCA	2367	-9.13	-3.05	-1.92	-2.52	-1.28	-2.58	-1.50	0
13281-L5-3	CACCCCCACCCACAGGACAGAGGAAGTACGAG	2368	-17.40	-17.40	-2.72	-9.25	-17.40	-17.40	-1.76	0

13283-L5-3	GGACCCCTGCGCTTCCTTGCTGCCACCCCTTTGCACA	2369	-5.08	-5.08	-1.49	-2.73	-5.08	-5.08	1.94	0
13285-L5-3	GCTATGCACCCAGCGGCCAGCTCAGCCCCCTGC	2370	-13.41	-13.41	-2.23	-7.03	-13.41	-13.41	-1.61	0
13289-L5-3	GCTCAAAAGTTTGCTCCCATGGCCCCCTCTGCCC	2371	-5.97	-5.97	-1.04	-2.79	-5.97	-5.97	1.72	0
13291-L5-2	TGGTTCTGCCCCAAGCGCCCTTCCTCCCTCCTT	2372	-5.00	-5.00	1.36	-3.54	1.37	-2.63	3.53	1
13312-L5-2	TGCCACCCACCCCTCCCCACAGCCAGCCCC	2373	-6.79	-15.24	-1.49	-6.97	-1.11	-3.12	1.59	0
13325-R5-1	CTGCCTGGCGCTGGGCTCTTCCCCATGAGCG	2374	-8.15	-8.15	-1.33	-3.29	-8.15	-8.15	2.04	1
13325-R5-2	TCCAACACTGCGCTGGCGCTGGGCTCTTCCCCA	2375	-8.85	-8.85	1.23	-1.20	-8.85	-8.85	2.07	1
13326-L5-1	TCCGGCTTCCCCCACCCGCCCTTCGATGGCA	2376	-5.52	-5.52	1.33	-1.60	2.24	-1.93	2.59	2
13326-L5-2	CAGAGATTCGGGCTTCCCCCACCCGCCCTTC	2377	-10.44	-14.87	-1.59	-3.94	-1.48	-5.00	1.20	0
13343-L5-1	CCTCCACCCCTCCCCGAGCGGCCCTCCCCCTCA	2378	-54.66	-54.66	-2.68	-1.95	-2.09	-4.50	-1.06	0
13349-L5-1	TGTGTGGCTATCCCCAGCGGCTCCTTCTCTCT	2379	-6.04	-6.04	1.05	-1.36	-6.04	-6.04	1.40	0
13349-L5-2	CACCACCTGTGTGGGCTATCCAGCCGCTCC	2380	-17.13	-17.13	-4.48	-12.67	-17.13	-17.13	-2.49	0
13365-L5-3	TCCCCAGAGCCCCCCCCAACCCACCTCGATCTTT	2381	-12.64	-27.91	-4.88	-2.98	-22.68	-21.98	-1.53	0
13374-R5-1	AGCCTTCTGTCCCCCTGGCCCCCGACCTGC	2382	-20.45	-20.45	-2.48	-5.82	-13.37	-11.72	-1.59	0
13374-R5-2	GGCCCTAGCCTTCTGTGCCCTGGCCCCCGA	2383	-5.79	-5.79	-1.35	-2.92	-4.80	-5.79	1.61	0
13375-L5-3	GAGAACCTGAACCCCCAGCCCCCTGCTACCCCTTAG	2384	-5.86	-5.86	1.24	-2.87	-5.86	-5.86	1.67	0
13396-L5-2	GACCTGCCCCCGCCCCACTCGGGCTCCTTACCG	2385	-5.03	-10.37	-1.25	-2.62	-8.50	-7.78	1.54	0
13431-L5-3	CGACACCCACTCACTGCCGCTGCCGCACTCACAGC	2386	-23.38	-23.38	-2.69	-4.01	-23.38	-23.38	-4.52	0
13432-R5-4	TTTACAGTTTCTGGCACTTCTTACCACCTCCCCA	2387	-5.41	-5.41	-1.42	-1.80	-5.41	-5.41	2.00	1
13456-R5-2	CAGATGCCCCCGCTATGAAATCTTTTCCAACC	2388	-16.80	-16.80	-13.79	-16.80	-16.80	-16.80	-16.80	0
13461-L5-4	TCCCCAGCCCCGTCCCCACCCCTAGAGAAAGTGAA	2389	-19.96	-28.08	-3.31	-5.34	-4.75	-16.77	-1.67	0
13497-L5-1	TGTGCCCCCCTCGCTCCCAAGCCCCCAGGGACCGC	2390	-7.88	-20.24	-4.39	-3.42	-3.34	-3.05	-1.79	0
227-L5-1	ACACCTGTCTCTCCCCAGTGCTTCCGCCCTCA	2391	-59.95	-93.40	-4.00	-10.53	-13.51	-33.21	-2.73	0
266-R4-1	GTCGCCCCCTCCCCCAAGTTGAGACTTGCAGCTAC	2392	-6.36	-14.52	-1.45	-2.41	1.12	-3.13	1.90	0
266-R5-1	CCCCCCCCCAAGTTGAGACTTGCAGCTAC	2393	-7.35	-7.35	1.06	-2.08	1.10	-4.92	5.32	1
2819-R5-4	CAGCCTGCCACCCGCCGCTTTTGAAAGAGCACTTCA	2394	-8.56	-8.56	1.00	-5.87	-8.56	-8.56	-1.66	0
3009-L5-1	ACCTCGGTCTCTCCACCAGACTTTAAACTCTC	2395	-17.19	-17.19	-5.92	-2.82	-7.48	-17.19	-4.70	0
3009-L5-2	CCAGTGATACTCGGTCTCCTCCACCAGACTTT	2396	-19.46	-19.46	-4.10	-9.16	-10.45	-19.46	-5.93	0
3009-L5-3	TGTAATAACCAGTGATACCTCGGTCTCCTCCAC	2397	-23.73	-23.73	-3.74	-2.02	-7.39	-23.73	-5.75	0
3249-L4-1	GCGGAGCCGCCGCCATCCCCGGAGCCGCCGCCGCCG	2398	-8.53	-6.22	-3.77	-2.41	-14.30	-15.31	-2.56	0
3249-L5-2	AGCGGAGCCGCCGCCATCCCCGGAGCCGCCGCCG	2399	-94.34	-18.11	-4.95	-2.32	-23.40	-29.20	-2.57	0
3799-R5-1	CTGAAGATGCTCCCCAGAGGCCCCCCCCCGGCC	2400	-7.96	-8.04	-1.30	1.45	-8.27	-13.22	1.59	0

3897-R5-2	CCGACCCGCGGTCAGCGCGCTCCCCCTCAG	2401	-11.04	-14.48	-1.65	-2.21	-3.50	-5.62	-1.55	0
3953-R3-2	ACTCCAGCCTCCGCGCCTCAGCTTCCCGAGC	2402	-34.43	-52.88	-3.53	-4.10	-15.19	-24.22	-5.59	0
3966-L5-1	ACCCAGAGCTGTGCGCGCGCTGCGCCTTCGCC	2403	-12.50	-9.10	-4.67	-3.33	-12.90	-27.25	-3.64	0
4303-R1-1	AGTGCCGCTCCTCCGACCTCCCTGCGCACC	2404	-16.94	-16.94	-1.82	-7.37	-13.90	-16.94	1.02	0
4315_C-L4-1	GCAGCCCCCTCCTCCGAGAGGTTGGGGTCGCGGCGCGCCC	2405	-10.99	-17.02	-1.97	-1.53	-1.97	-2.96	1.32	0
4315_D-R4-1	GAAAGTCAGCCCCCAGCGCCCCCGGAGTTCTTGG	2406	-88.80	-88.80	-12.25	-8.23	-57.77	-88.80	-6.18	0
4315_E-R4-1	CCCCACCAAACTATTCCCGCATCCTCCCCGGCTCTGG	2407	-7.17	-8.95	-1.28	1.62	-3.51	-8.09	2.47	1
4315_F-R4-1	AACCCGGGCTCCCCACCCGCTCCCTGAGC	2408	-19.29	-28.54	-1.97	-2.12	-6.85	-14.36	-1.32	0
4315_I-L4-1	ACACCTCTGCGCCCCCTCAGGCGCCCTGGGCCTCGGCGCCCC	2409	-30.88	-30.88	-4.30	-3.56	-30.88	-30.88	-1.71	0
4315_K-L4-1	TCCAGGGGGCCCTGAACCTTGTCAAATCCTCGCCATCCTCC	2410	-7.44	-7.44	1.35	-1.81	-2.45	-3.94	1.80	0
4440-R3-1	ACCCCGAGCCCCG	2411	-10.88	-1.67	-1.04	1.21	-4.04	-11.36	-1.05	0
4479-R3-1	AGCCCCCTGCCCCGAAATTCAAAACAACCTGC	2412	-14.50	-20.12	-3.55	-6.32	-16.37	-20.12	-3.97	0
4593-R5-1	CTATAGCAGATGACATAACTCCCCCGGCATCAG	2413	-12.46	-12.46	1.06	1.11	-12.46	-12.46	-1.12	0
4829-R2-1	TCCCTTTGTGCTGCCCCGAGTGCCCTTCCCCCTG	2414	-9.27	-9.27	2.07	-1.01	-3.71	-2.66	6.01	2
4855-R5-2	GGGTGCCGGGTCTCCCGCTTCCCGCTCCCTGC	2415	-10.65	-15.12	1.25	1.03	1.70	-1.51	1.72	0
4884-R5-1	TCCCCAGTCTCCCTGTTTCAGCACCTGCCTCA	2416	-17.58	-17.58	-1.81	-2.95	-7.44	-9.96	1.51	0
4988-R4-1	CTCCTCCTCCCGCTCTTTGGATACCAAACTGGAC	2417	-5.47	-5.47	1.61	-1.61	-1.44	-5.47	3.51	1
4988-R5-2	CTCCTCCTCCCGCTCTTTGGATACCAAAACAC	2418	-9.43	-9.43	1.16	-2.18	-1.17	-9.43	4.95	1
5071-R5-1	CGCCCCAGTCCAGCCCCCAATTAATAATGGG	2419	-6.09	-6.09	1.41	2.38	-6.09	-6.09	1.59	0
5071-R5-2	GACCCCGCCCCAGTCCAGCCCCAATTAATA	2420	-24.30	-13.23	-2.97	-3.00	-5.08	-18.13	-1.48	0
5640-L3-1	GCCATGGAACACCGTGCCCTGCCCTCTCGAGA	2421	-22.57	-22.57	-3.86	-4.16	-22.57	-22.57	-1.57	0
5707-L5-2	TCACCATGCGGCCCCGGTGCTCTTCACACAGCA	2422	-17.42	-17.42	-17.42	-17.42	-17.42	-17.42	-17.42	0
6198-R5-2	GCCGCGCGCGCGCGTCTTCCCGCGAAGCCT	2423	-8.01	-10.31	-4.02	-2.44	-21.45	-23.60	-3.04	0
6216-R5-2	CATAGTTACTCCCGCGGTTTACCCGTGCTTC	2424	-15.09	-10.39	-1.12	1.09	-5.30	-18.03	-1.30	0
6825-R5-1	TCCTTCTGCTCAGCTGTTCCTCCGGTGCCAG	2425	-28.95	-28.95	-21.84	-14.99	-28.95	-28.95	-6.06	0
6880-L3-2	ACCTCCCCCGCGAAGACATCCACATTCTGCA	2426	-20.08	-29.17	-1.83	-9.68	-3.55	-29.17	-3.36	0
7061-R5-2	TCATGGAACCCCAACCTTCCCATGCCCAACC	2427	-5.94	-5.94	-1.04	-4.28	-5.94	-5.94	1.37	0
7126-L3-1	GCACACCCGCTCTCCGGCCCCGCCCCCTG	2428	-25.95	-6.91	-4.10	-2.74	-11.10	-14.35	-2.85	0
7356_A-R4-1	CAGAGCCCGCTCTCGCGACCGACCTGCCGCCGACCGCCAC	2429	-18.01	-26.53	-3.41	-4.02	-22.46	-25.26	-3.07	0
7702-L2-1	CCCAGAGAACCGGAATTCTCCCCGCC	2430	-5.27	-5.27	-1.83	-2.91	-4.35	-5.27	2.04	1
7824-R5-1	TGCCAGCTTCATCGCCGCTCACACACACA	2431	-8.78	-8.78	-2.70	-8.78	-8.78	-8.78	-3.37	0

7949-R5-1	GATGCGCGCGCCGACCGCGCGCAGCTGCAATTCATAC	2432	-24.23	-24.23	-3.02	-1.84	-24.23	-24.23	-2.16	0
8016-L3-1	TCAGCGCAACAAGCCCCGAGTCACCCCTCT	2433	-54.19	-54.19	-16.08	-20.80	-54.19	-54.19	-6.74	0
8250-R5-2	CCGACCGCGCCGTCAGCCGCTCTCCCTCAG	2434	-17.68	-16.09	-1.77	-1.62	-3.89	-4.78	1.33	0
8394-L5-1	CCGCCCTGCCCATCTCCGACTATCCCTGGCCCC	2435	-7.06	-7.06	1.08	-1.28	-7.06	-7.06	2.40	1
8898-R5-1	CAGCCGAGCGGACGCCCCGCTCCGCAACCATG	2436	-10.42	-15.12	-2.07	-1.34	-12.35	-15.12	-1.31	0
9053-R3-1	TTCTTGCCCTCCAATCCCCGGGCTCCACACGCC	2437	-10.59	-10.59	-1.97	-1.65	-8.41	-10.59	-2.27	0
9387-R2-2	TCCATCCTTGCCGTCGCCCTTCATCTCAAAGCCATC	2438	-8.80	-8.80	-1.47	-8.80	-8.80	-8.80	-3.48	0
9691-L5-1	CATTTCATCCGCATCTCCCTCTTGCCCTTGC	2439	-8.33	-8.33	-2.45	-6.14	-8.33	-8.33	1.53	0
9774-R2-2	CCGCCCCCTCACCGCCTCCTGCTCCCATCAGGC	2440	-7.93	-16.14	-1.34	-2.55	-4.83	-3.47	1.40	0
miR-1228*	CACACACCTGCCCCCGCCAC	2441	-15.38	-12.90	-2.54	-1.74	-4.13	-8.82	-1.45	0
miR-1275	GACAGCCTCTCCCCCAC	2442	-8.09	-8.09	2.33	-1.30	-2.96	-8.09	2.04	2
miR-1308	CCACTGAACCAACCCATGC	2443	-18.92	-18.92	-5.22	-18.92	-4.03	-18.92	-5.64	0
miR-21	TCAACATCAGTCTGATAAGCTA	2444	-1.68	-23.57	-19.14	-23.57	1.39	-7.45	-2.50	0
miR-373*	GGAAAGCGCCCCCATTTTGAGT	2445	-48.87	-48.87	-48.87	-48.87	-48.87	-48.87	-21.32	0
miR-423-5p	AAAGTCTCGCTCTCTGCCCCCTCA	2446	-9.88	-4.20	-6.30	-4.12	-7.70	-16.50	-3.64	0
miR-486-3p	ATCCTGTACTGAGCTGCCCCG	2447	-40.84	-40.84	-36.13	-40.84	-40.84	-40.84	-7.70	0
miR-612	AAGGAGCTCAGAAGCCCTGCCCAGC	2448	-5.79	-5.79	-3.09	-2.94	-5.79	-5.79	-1.13	0
miR-638	AGCCCGCCACCCCGCCCGGATCCCT	2449	-11.00	-23.12	-2.25	-1.89	-19.12	-23.12	-1.53	0
miR-663	GCGGTCCCGCGCGCCCGCCGCT	2450	-45.02	-45.02	-10.68	-32.17	-45.02	-45.02	-11.52	0
miR-744	TGCTGTTAGCCCTAGCCCCGCA	2451	-37.67	-37.67	-10.59	-19.87	-37.67	-37.67	-4.41	0
miR-923	AGTTTCTTTTCTCCTCCGCTGAC	2452	-3.90	-3.72	1.48	1.30	-6.06	-8.95	2.30	1

Table 31: Pre-microRNA sequences and chromosomal locations of target RNAs in Table 30

Gene Down expressed in Cell lines	Chrom loc'n	Pre-microRNA sequence	SEQ ID NO
10010_B-L4-1	7q32.1	GGCAGCGCCCCCTCCTTCGGAGGCGAGCATCCCGCGGGGGGCTCGGCTTTGATGCCAGGGCACCTTTTGTCCCCTGGAG ACGCTCTGCCAGCCAGGTGCGTGGAGGGAGTGCAGCCC	2453
10010_D-L4-1	7q32.1	CAGGTGAGAGGCTAGCCCCCGGGGAGGCGCCAGGCTGGGGCATTAAACCGTCCCGGGGACCCCTTTTGGCCTG	2454
10138-L2-1	17q22	ACTTTTCTGTGGAGGGGAGAGCGGAGCAGGCTCACGTGTAAACCGCGCAGGAGCCTCCTCTGGCTTGAGCCCTTCTTGTTAAG T	2455
10231-R3-1	9p11.2	GGCGGCTGCGGAGGCTGGCGCGGGGCTGTCACCTTTAACGCTTTCGCGCTGACAGGCGGCGCCAGCTAAAGTTCACAGCG	2456

226

13106-R5-1, 13106-R5-2	14q11.2	TGGCTCTAGGCCACCCCTGGCAGCTGGGCCGCACTCTGCCGGCCTGGACGCCGGGCTCGGCTCTGATTGGTCCA	2481
13111-L5-3	16p13.3	AGCCTGTGGAAAGAGAGCAGGCGAGGTTGAAGCCCGCGGAGACACTCTGCCACCCACACACCCCTGCCTATGGGCCACAC AGCT	2482
13119-R5-2	17q21.3 2	TCTCCGTGCACGCTGCTGACCGGCTCGGCACTGCCTCCCTGCTGTGAGCAGGAGAAACAGGAAGTCTGCCCGACAGGGAGGTGGC CGGGCGGAGCGCAGAGTCGGCGTTGAGA	2483
13129-L5-2, 13129- L5-3	20q13.3 3	TGGCGCTGCTCTGCTGTTCCCTCTGCTCTCCAGACTCTGGGTGGATGGAGCAGGTGGGGGCCAGGGGACAGGAAGGCTAGGGCC	2484
13129-L5-2, 13129- L5-3	20q13.3 3	CCAGACTCTGGGTGGATGGAGCAGGTGGGGGCCAGGGGACAGGAAGGCTAGGGCCCCAGAGACCTGTCTCTGGGCCCCCATGTCCA GCTCTGCCCTTAGTGCTTGG	2485
13130-L5-1, 13130- L5-2	19p13.3	CCAAGCAGCTTATCGAGGAAAGATCGAGGTGGGTGGGCGGCTCTGGGATTTGGTCTCACAGCCCCGATCCCAGCCCCACTTA CCTTGGTTACTCTCTCTCTCTAGGG	2486
13137-L5-1	5p15.2	GGCATGGGAGGTGGGTTGCGGGGTTAGAGCTGTGGGTGATCAGAAAGGAGGCTTCATTTCTACGCTCTGCCTCCGCTACCT CTTCCCCACCAACCCTAATCCC	2487
13138-R5-1	5q22.3	ATCCGTCGCAGCAGTCGCTGCAGCCGCTGCAGTCCGAGCCGACTAAGGGCGGAGGCAGCTCGGGTGGCGGGCGGTGGCGAACA GCGCGCGCCGGAT	2488
13161-L5-4	8p21.3	GGAGAGAAAGGGCGCAGATCACACAGAGGGGACTGACACGCGGACCCCTGGCGGAACCCACCCCTTGCTTTCCGGTCCCCCA GTCGGTGCCTCCCTCCAGTGGCTCTCAGCTCCTCCCTCCC	2489
13209-R5-3	10q22.1	GGAGCCCGCGGCGCAGGCCCGCGGCGCAGGAGTTACCTCAACTCCCGCGCTCCGAGGTTGCCGGGCACCGAG GAGCCGCGTGCCCTTACGGCGCTCGCGCGCAGCA	2490
13211-L5-1	10q23.1	GCGCTCCGGTCGGTCAGAGAGCGGCGCGCTCTGCAGCCCTCCAAGCCGCTCCTGCGCGCGGTTCCCGCGCCCGCTG CTGCTGCTGCTGCCCGCCCTGCGTGC	2491
13227-L5-1, 13227- L5-2	11q25	AAGCAAGACACCGAGGTGGAAAGGGGACGAGGAGTGGGCCCTATTCTCCCATCTATGTAAAGGGAGGGATATCAGGGAAGT CTCTGTCTGTACTCAAGTTTGGGATGCT	2492
13229-L5-1, 13229- L5-2, 13229-R5-3	11p15.5	TGCTGGCCCCAAGGGTAAAGGGGCAGGGGACGGGTGGCCCCCAGGAAGGGCCTGGTGGAGCCGCTCTTCTCCCTGCCCCACAGAG ACTGGCGGAGCTGC	2493
13239-L5-2	1q23.3	CTGAGGACAGGGGTAAGTCTGGGAGATGGGGGAGCTCTGCTGAGGGTGACAAAGGCCCTGGCTCTACACACATCCCTGTCTTAC AGAG	2494
13240-L5-2	11q12.2	GCAGTGTGATTTGGGCCGGGAATGCCGCGGGGACGGCGATTGGTCCGTATGTGTGCCACCGCGCGCGCTCCGCC CGCCCCCGCCCCACACGCCGCAT	2495
13247-L5-3	1q24.2	ATCTACAGAGGAAGAACAGGGCTCTGAGGTCGGGAGGGGAAGCGGCTCAGGACTTCTGGCTCCAGAGCCCTCCTCTCCTTCCACC ATAGTGCTGCTCCAGAGGAGAC	2496
13267-L5-1	1q42.13	CACCATAGGTGAGGTGGGGCCAGCAGGGAGTGGGCTGGGCTGGGCCAAGGTACAAGGCCCTCACCCCTGCATCCCCGCACCC AGGCTTCAACGTGG	2497
13276-L5-4	12q13.1 3	GTGCTTTCTTGTCCAAGCAGTTGATGAAGGCAGGTGGGTGGAGCAGCCACAATGGGGGCTCTGCTGACCATCTGCCCTTCCACC CTACAGCACCCGGCAGACACCAAGT	2498
13281-L5-3	12p13.3 1	GCCGCCCCACCCCTGTCCCTCGTCACTTCTCTGTCCCTGTGGGTGGGGTGCAGGCGCTTCTCCTTTAGCTGTGCCGCACCTTCTCC CTACAGGCCAGGAGAAACAGAACACCGTGTGCAC	2499
13283-L5-3	1p36.11	GGCACGGGGTTGGGTGTGCAAGGGTGGCAGCAAGGAGGCGGTCTTAAGGTGTCTCCTGCCCTCCTTGTGTAGAC TTTGGCTGAGCAAGAGGCC	2500
13285-L5-3	1p36.32	CTGGGGGTGACCCCGTGCAGGGGCTGAGCTGGGCGGCTGGGTGCATAGCCCATCTGTAGCCTAAGCAATGGTGCAAGCCCCCAC GCCTTGGTTTCTCTCTCTGTAACGGGGCACACAGAG	2501
13289-L5-3	1p35.1	TGCTTGGAGTGGCAGAGGGCAGGGGCCATGGGAGGCAAACTTTGAGCGTGTCTCAGGAGCCCTAAAACATGAGAAGCCTCAGTTT CCCTCTCTAGACCATCTGCTTATCCTCTCCAGGC	2502

228

3897-R5-2	9q12.9p 11.2	CGGAGCCCGCCGCGCCTCTCCATCTCGCAAGTTTTAATTAACGCTGAGGGGAGGCGGCTGACGGGCGGGTCGG	2526
3953-R3-2	9q33.3	GCTCCTGCTCCGCGCGGAGCTGCTCCGGCGCGCAGGGCTCGCTCGGGAAGCTGAGCGCGGAGGCTGGAGT	2527
3966-L5-1	3q27.2	GGCGAAGCGCGCAGCGCGCGCACAGCTCTGGGGTTTGCCTCTCGGGTGTGTCGGCCCGCGCTGCTGCTTGGGCC	2528
4303-R1-1	22q11.2 1	GGCTGCCAGGCTCCGCCCGCCCTCCCTGCGCCCGCGGTGCGCAGGAGTGGAGGCGGACTGCCACCC	2529
4315_C-L4-1	1q22	GGGAGGGGACCGGAGGGCCGGCGCGCGACCCCAACCTCTCGGAGGAGGGCTGCCGCTCGCCGCTCTTTGTTG TTTGGGCTCCGCGCTCCCCCTCTCTCCCTCCTC	2530
4315_D-R4-1	1q22	GGGACGTGGCCCTCCCGCCGAGCGGACTCCAAGAACTCCGGGGCGCTGGGGCTGACTTTCC	2531
4315_E-R4-1	1q22	TCCCGCCACCCCTTGAGCACCTCAGCGTTCTTAGGGAAAGCCAGAGCCGGGAGGATCGGGAATAGTTTGGTGGGG	2532
4315_F-R4-1	1q22	GATCAGGTTCCCTCCCGCATACACCTGGCGCAGGTGAAAGCTCAGGAGCGGGTGGGAGCCCGGTT	2533
4315_I-L4-1	1q22	CTGGACGGCGGGCGCGAGGCCAGGGCGCGCTGAGGGCGCAGAGGTGTACGCTGCAACCGCGCCCGCCAGCGTTCCC GCCACACCGCCACACCTCAAGCCCCG	2534
4315_K-L4-1	1q22	CCGGGCTGGGGTGGAGGATGGCGAGGATTTGACAAGTTCAGGGCCCCCTGGGATCCTTTCCCTACTCCCTGGTCTTGTGGACA CCCTGTTACCTGCCCTAATTGCCCCG	2535
4440-R3-1	7q11.22	GTGATGTGATTTCTGCCCAGTGTCTGAATGTCAAACTGAAGAAATTCAGTGAATGCGGGTAAACGGCGGGAGTAACTATGAC	2536
4479-R3-1	1p32.3	CCCAAGCTCCTTCTGGAGGACTTAACACTGTGTGAGCAGTTGTTTTGAATTTCCGGCAGGGGCTGCAAAAGG	2537
4593-R5-1	15q23	CAATCAATTAGCACATGAGTAAACCAAGCCCCATTAGGACAAACTGATCCCGGGGAGTTATGTCACTCTGTATAGAAATGATTG	2538
4829-R2-1	1q21.3	GGTGTGCTGCCCTCTCTTTCTGCCCCCTATACCCCTTGACCCCAAGGGAAGGCACCTCGGGCAGCACAAAAGGAGCAGATGCC	2539
4855-R5-2	12q13.1 1	GGTCCGGGTCTCTACCGCGCCCTCATGCAGGAGGCCCTTGAGCAGGAGGGGAAGCGGAGACCCGGCAGCCC	2540
4884-R5-1	17q11.2	TCCACAGCTCCATAAATGTTTTAATGCCACCGCTGGCAGGAATGAGCAGGTGCTGAAACAGGGAGACTGGGA	2541
4988-R4-1, 4988-R5-2	14q24.3	CTTTTCTCTGCTGGAAACC TTGCTTGACTTCATGTCCAGTGTGTTGTTATCCAAAGACGGGAGGAGGAG	2542
5071-R5-1, 5071-R5-2	17q12	GATTCTGCTCCCAGAGCCATAAAGTGGAGCCCCCATTTATTAATTGGCTGGGACTGGGGCGGGGTC	2543
5640-L3-1	1p34.1	TCTGAGAGGGGCGAGGCACGGTGTTCATGGCAAGACGGCGTTGATGTATAGGCGTGGCATGAAGCTGGGCTTGCTCTCAGA	2544
5707-L5-2	9q34.3	GCTCCTCGTGTGTGAAGACCAACCGGGCGCATGGTGAAGCACCCAGTGTCTCCTGGCTGTGGCTACTTCTGCACAGCGGTAA GAGC	2545
6198-R5-2	1q25.3	GCCGCCAGCACCCCGGTGCCCGGGGCGCTCCGAGGAGCCTGAGAGACCCACGGAGGCTTCGCGGGAAGACGCGGGCGGG CGGC	2546
6216-R5-2	11q14.1	CATGTGATTTCTGCCCAGTGTCTGAATGTCAAAGTGAAGAAATTCATGAAGCACGGGTAAACGGCGGAGTAACTATG	2547
6825-R5-1	9q31.3	CAAAATTACATCTGTTATGCTTCTATTTGTTAGACAAATCTGGCACCGGGAACAGCTGAGCAGAAGGATTG	2548
6880-L3-2	1q42.13	GGCTTGAGAAATGTGGATGTCTTCGCGGGGAGGTGGCCACGTTCTCCTGTGAGGTGTCTCACGCGGGTGGCC	2549
7061-R5-2	1p13.3	TCATGGCAGCACCCACCTCCAGTCCCTGGACAAATCGGGTACAGAGACTTAAGTTGGGCATGGGAAGGTGGGTTTCCATGA	2550
7126-L3-1	5q31.1	CAGGGCGCGGGCGGAGAGCGGGTGTGCAAAGTGGCGCAGGGCCCTGGGGCCCGCCCTTGCTCTGCCGGCTCGACTCTTG CGGAGTCTGCTCGCTCGCTCGCTGCTGACTCGCCGAGCGCTCTGTGGCGGTGCGCGCAGGTCGTCGCGAGAGCGG	2551
7356_A-R4-1	8q24.3	GCTCTG	2552

7702-L2-1	10q21.2	GGGGCGGGAGGAATCCGGTTCTCTGGGACTTTCCAAAGGCGAAGATCCGGTGCCGGCTCCGCTCCCTAGCCCT	2553
7824-R5-1	6q16.2	CCTGGATGCTGTTTCATTATGAGATCAGGCAAAAGACAGACGGATGTGTGTGAGGCGCGCATGAAGCTGGCACCAGG	2554
7949-R5-1	5q31.3	GGTGCGCGGGTGGACGCCGATTCCGGCTACAAATGCGTGGCTTTCGATGAATTGCAGCTGGCGGCTCGCGCGCGCATC	2555
8016-L3-1	12q21.1	AGAGGGGTGACTGCGGGGCTTGTTCGCTGAAGATTTACAAATGTACTTCTTCAGGCGGCTCAGCAACCCCTCT	2556
8250-R5-2	9p11.2	CGAGCCGCCCGCCAGCCTCTCCATCTCGCAAGTTTTAATTAACGCTGAGGAGAGGCGGCTGACGGGCGGTCTGG	2557
8394-L5-1	7p13	GGGCCAGGGATAGTCGGAGATGGCAGGGCGGGGCCCCCACTGGCGAGGGGCCCTCGGCTTCTGGGTCCCTGAGCCCC	2558
8898-R5-1	17p13.3	GGCGCTGTCGGCCCGGGCGGCCGCGCAACTCGTCCGCTTGATAACCATGGTGGCGGAGCGGGGCTCCGCCCTCGGCTGTCC	2559
9053-R3-1	Xq27.3	GGAGGGCACTGTCTCTCTGATTCCAGGGCCTGTCAATTTCCGAGGGCTGGTGAGCCCCGGGATTGGAGGGCAAGAGCCAG	2560
9387-R2-2	3p21.1	TCTCCATCCTCTGTCTCCCTTGATCCTCTCTGTTCTCCCTGATGGCTTTGAGATGAAGCGCACGCAAGATGGAGG	2561
9691-L5-1	14q24.3	GCAAGGGGCCAAGAGGGAGATGCGGATGAATGGATGATTTAATGGGTCACTCTCTCTGATTTAATTTCTCTAGATCTTGT	2562
9774-R2-2	13q13.3	GCTTGTCCTAAAAGAATCTTCCCTCTGTTTCCCTGGGTTTATCCACTTGGTTGGCCTGATGGAGCAGGAGCGGTGAGGGGCGGGC	2563
miR-1228*	12q13.3	GTGGCGGGGGCAGGTGTGTGGTGGTGGCTGCGGTGAGCAGGGCCCTCACACCTGCCTCGCCCCCAG	2564
miR-1275	6p21.31	CCTCTGTGAGAAAGGGTGTGGGGAGAGGCTGTCTGTCTGTAAAGTATGCCAAACTTATTTTCCCCAAGGCAGAGGGA	2565
miR-1308	Xp22.1	CCCCGCATGGGTGTTTCTAGTGGCAGAAATCTCAAATGTAAATCCCCATAATCCC	2566
miR-21	17q23.1	TGTGGGTAGCTTATCAGACTGATGTTGACTGTTGAATCTCATGGCAACACCAGTCGATGGGCTGTCTGACA	2567
miR-373*	19q13.4	GGGATACATAAATGGGGCGCTTTCCTTTTGTCTGTACTGGGAAGTGCTTCGATTTTGGGTGTCCC	2568
miR-423-5p	17q11.2	ATAAAGGAAGTTAGGCTGAGGGGCAGAGAGCGAGACTTTTCTATTTTCCAAAGCTCGGTCTGAGGCCCTCAGTCTTGCTTCTTAAC	2569
miR-486-3p	8p11.21	GCATCCTGTACTGAGCTGCCCGAGGCCCTTCAATGCTGCCAGCTCGGGCAGCTCAGTACAGGATAC	2570
miR-612	11q13.1	TCCCATCTGGACCCCTGCTGGCAGGGCTTCTGAGCTCCTTAGCACTAGCAGGAGGGCTCCAGGGGCCCTCCCTCCATGGCAGCCA	2571
miR-638	19p13.2	GTAGCGGGCGCGGCGAGGATCGCGGGCGGGTGGCGGCCCTAGGGCGCGGAGGGCGGACCCGGGAATGGCGCGCGTGGCGCGC	2572
miR-663	20p11.1	CGCGTAACTGCGGCGCT	2573
miR-744	17p12	CCTTCCGCGTCCAGGCGGGCGCGGACCGCCCTCGTGTCTGTGGCGGTGGATCCCGCGCGCGTGTTCCTGTGTGGC	2574
miR-923	17q12	TTGGGCAAGGTGCGGGCTAGGGCTAACAGCAGTCTTACTGAAGGTTTCTGTGAAACACGACACATGCTGTTGCCACTAACCTCAAC	2575

5.6 Example 6: Target RNA Levels in Primary Tumors and Cell Lines

[00323] Data from Examples 4 and 5 were analyzed to identify target RNAs that were found to be up-expressed in both primary lung tumors and in lung cancer cell lines. Table 32 shows a list of the target RNAs from Table 27 (i.e., target RNAs that are present at increased levels in at least four lung cancer cell lines) that are also in Tables 18 to 21 (i.e., target RNAs that are present at increased levels in more than 50% of primary lung tumors tested, and target RNAs that are present at at least 5-fold increased levels in fewer than 50% of primary lung tumors tested). Table 33 shows a list of the target RNAs from Table 28 (i.e., target RNAs that are present at at least 5-fold increased levels in two or three cell lines) that are also in Tables 18 to 21.

Table 32: Target RNAs from Table 27 that are also in Tables 18 to 21

target RNA from Table 27 (increased in at least four cell lines)	probe SEQ ID NO	pre-microRNA SEQ ID NO	microRNA SEQ ID NO	Present in Tables 18 and 19 (increased in at least 50% tumors)?	Present in Tables 20 and 21 (increased at least 5-fold in <50% tumors)?
10083-L5-1	1090	1211			Yes
13122-L5-1	1066	1242	2587	Yes	
13185-L5-3	1118	1243	2603	Yes	
13219-L5-1	1067	1245		Yes	
3744-R5-1	1143	1271	2645, 2646		Yes
6235-R5-2	1075	1296	2661	Yes	
6474-L5-1	1164	1299			Yes
6681-R2-1	1166	1301			Yes
8004-R3-2	97	492		Yes	
836-R4-1	3	400		Yes	
9594-R5-1	1081	1328		Yes	
miR-103	361	754, 755	803		Yes
miR-200b	193	692	839	Yes	
miR-200c	194	693	840	Yes	
miR-20b	380	776	843	Yes	
miR-298	1088	1351	858	Yes	

Table 33: Target RNAs from Table 28 that are also in Tables 18 to 21

target RNA from Table 28 (increased 5-fold in 2 or 3 cell lines)	probe SEQ ID NO	pre-microRNA SEQ ID NO	microRNA SEQ ID NO	Present in Tables 18 and 19 (increased in at least 50% tumors)?	Present in Tables 20 and 21 (increased at least 5-fold in <50% tumors)?
10335-L5-2	1092	1213			Yes
12729-R5-1	1103	1186			Yes
12888-L5-2	1105	1188			Yes
12917-R5-2	1108	1230			Yes
12992-L5-1	1112	1235			Yes

13001-L5-1	1113	1236			Yes
13070-R5-3	1115	1238			Yes
13274-L5-3	1124	1251	2612		Yes
13357-L5-4	1128	1255	2621		Yes
13366-R5-3	1129	1256	2623		Yes
13467-L5-1	1069	1262	2630	Yes	
13470-R5-1	1136	1264			Yes
3875-R5-2	1144	1272			Yes
3923-R5-1	1070	1273	2647	Yes	
5108-R5-2	1153	1286			Yes
5723-R5-1	1156	1289			Yes
6752-R5-2	1168	1303			Yes
6803-R5-2	1169	1304			Yes
6930-R5-1	1173	1308	2667		Yes
8433_C-R4-1	1177	1317			Yes
8433-L3-1	1179	1318, 1319			Yes
8724-R5-2	1181	1320			Yes
8808-R5-1	1183	1322			Yes
8832-R5-1	1184	1323			Yes
9349-R5-2	1080	1325	2672	Yes	
miR-198	1202	1343	834		Yes
miR-720	1089	1362	917	Yes	

[00324] Finally, Table 34 shows target RNAs that are present at decreased levels in both primary tumors and cell lines relative to normal tissue or cell lines.

Table 34: Target RNAs present at decreased levels in primary tumors and cell lines

Gene	probe SEQ ID NO	pre-microRNA SEQ ID NO	microRNA SEQ ID NO
10010_B-L4-1	157	551	
10010_D-L4-1	158	552	
10231-R3-1	1368	1713	
10342-R2-2	1371	1716	
11370-L5-5	1379	1724	
12691-R5-1	1385	1730	
12692-L5-1	1386	1731	
12693-L5-1	1387	1732	
12694-R5-1	1388	1733	
12696-R5-2	1389	1734	
12697-R5-1	1390	1735	
12699-L5-1	1391	1736	2585
12701-L5-1	1392	1737	
12703-L5-3	1393	1738	
12704-L5-2	1394	1739	
12713-R5-1	1395	1740	

12722-L5-1	1396	1741	
13004-R5-1	1411	1756	
13047-R5-2	1412	1757	
13052-L5-1	1414	1759	
13093-L5-2	1419	1765	
13097-L5-2	1422	1767	
13119-R5-2	1425	1771	
13129-L5-3	1427	1773, 1774	
13130-L5-2	2352	2486	
13137-L5-1	1431	1779	
13138-R5-1	1432	1780	
13209-R5-3	1444	1792	
13211-L5-1	1445	1793	
13229-L5-1	1447	1795	
13239-L5-2	1451	1800	
13240-L5-2	1452	1801	
13267-L5-1	1456	1805	
13281-L5-3	1457	1806	
13283-L5-3	1458	1807	
13285-L5-3	1459	1808	
13291-L5-1	1461	1810	
13312-L5-2	1467	1816	
13326-L5-2	1470	1820	
13343-L5-1	1477	1828	2619
13349-L5-2	1478	1829	
13365-L5-3	1487	1838	
13374-R5-1	1490	1842, 1843	
13375-L5-3	1491	1844	2625
13396-L5-2	1495	1848	
13431-L5-3	1501	1854	
13432-R5-4	2387	2516	
13456-R5-2	1503	1856	
13461-L5-4	1505	1858	
13497-L5-1	1508	1861	
266-R5-2	1513	1866	
2819-R5-4	1516	1869	
3799-R5-1	1519	1872	
3897-R5-2	1520	1873	
3953-R3-2	1523	1876	
3966-L5-1	1524	1877	
4315_C-L4-1	1533	1886	
4315_D-R4-1	1534	1887	
4315_E-R4-1	1535	1888	
4315_F-R4-1	1536	1889	
4315_I-L4-1	1537	1890	
4315_K-L4-1	1538	1891	
4593-R5-1	1547	1900	
5071-R5-2	1560	1913	

5640-L3-1	1569	1922	
6198-R5-2	1582	1935	
6880-L3-2	1596	1949	
7126-L3-1	1604	1857	
7356 A-R4-1	1609	1962	
7702-L2-1	1618	1971	
7824-R5-1	1622	1975	
7949-R5-1	1626	1979	
8016-L3-1	1628	1981	
8250-R5-2	1633	1986	
8394-L5-2	1638	1991	
8898-R5-1	1641	1994	
9387-R2-2	1651	2004	
9691-L5-1	1654	2007	
9774-R2-2	1656	2009	
miR-373*	1693	2049	
miR-486-3p	1695	2051	
miR-638	1699	2055	
miR-663	1700	2056	
miR-744	1702	2058	

5.7 Example 7: Additional Target RNAs from Primary Tumors

[00325] In addition to the target RNAs identified in Example 4, certain additional target RNAs were identified in that experiment as being present at elevated levels in squamous cell carcinoma (SCC). Further, one additional target RNA (miR-320c) was identified as being elevated in at least three of the most aggressive tumors in Example 4. Those additional target RNAs, and miR-320c, are shown in Tables 35 and 36, along with the fold-change in each of the primary tumors. Data for six of the tumors are shown in Table 35 and data for the remaining 13 tumors are shown in Table 36. Table 35 also shows the probe sequences used to detect the target RNAs. Table 37 shows the pre-microRNA sequences and chromosomal location of the pre-microRNA gene for each of the target RNAs in Tables 36 and 37.

Table 35: Additional target RNAs more frequently present at elevated levels in squamous cell carcinoma (SCC), and miR-320c

Gene	Probes sequences	SEQ ID NO	ADK9	ADK10	Adk29	Adk15	Adk23	ADK48
13108-L5-2	CTGCTGCCCTTCCCTTGTTGAGGGGCCCTGAGCACG	2673	-5.44	-5.44	1.00	-1.07	-2.30	-1.55
13272-R5-2	TCACCGTGCCCTCCCTTTGGAAAGAGGTAGAGTCA	2674	-2.63	-2.63	-2.63	-2.63	-1.60	5.15
13316-R5-2	CCACCACCTTGCTGCTGGCCCCACAGCACACGGCC	2675	-4.30	-4.30	-2.50	-2.88	-4.30	-1.25
13331-L5-2	CCCGAACCCCACTCTGAGCACTCGGCCACCAAGCA	2676	1.69	-3.04	-1.84	1.35	-1.38	-3.04
13499-R5-1	ACAAAGTTTCTCAGGAAGTCTGAACACTGGGTTT	2677	2.23	-2.83	1.29	1.39	-1.37	1.07
5971-R5-2	TGCCTGCTCAGCCTCCACATCTGTTCCTGG	2678	1.36	-2.71	-5.23	-1.20	-2.32	2.33
7026-L3-1	GAAAACCTCACCCACCAAGATCCGGGAACGA	2679	-4.57	-4.57	-4.57	-3.02	-2.20	-4.57
7471-L5-1	AAATGCAAATGCCCCCTAAGGAGAAGAACTTC	2680	-6.11	-6.11	-6.11	-1.41	-3.03	3.29
miR-320c	ACCCTCTCAACCCAGCTTTT	2689	-6.09	-6.09	-3.75	-1.52	1.03	-6.09

Table 36: Additional target RNAs more frequently present at elevated levels in squamous cell carcinoma (SCC), and miR-320c (con't)

Gene	Adk40	Adk41	Adk49	Epi42	Epi43	Ksarc19	Kmalp21	Kmalp25	EPI-4	Kmalp44	Scc27	Lcnec31	Car13
13108-L5-2	-5.44	-5.44	3.75	2.76	2.52	2.92	1.32	1.28	6.13	2.43	1.71	2.26	-5.44
13272-R5-2	-2.63	-2.63	-1.32	-2.63	3.89	2.75	2.27	1.53	10.30	3.13	2.20	2.16	-2.63
13316-R5-2	-4.30	-4.30	-2.20	2.94	2.28	1.96	3.07	-1.31	4.09	1.82	2.30	1.24	-4.30
13331-L5-2	-3.04	-3.04	4.06	-3.04	4.60	2.19	1.84	2.46	8.67	4.20	1.80	2.26	-3.04
13499-R5-1	-2.83	-2.83	5.02	4.08	4.17	1.08	2.12	1.95	-2.83	3.61	1.34	1.21	-2.83
5971-R5-2	-5.23	-5.23	-2.44	2.15	2.01	2.54	1.93	1.48	4.62	-5.23	1.20	1.38	-5.23
7026-L3-1	-4.57	1.04	2.37	-1.93	2.63	2.98	2.00	-1.19	4.29	2.31	1.36	1.10	-4.57
7471-L5-1	-6.11	-6.11	-1.05	6.18	3.37	2.29	3.21	-1.10	8.46	-6.11	-1.14	-1.42	-6.11
miR-320c	-6.09	-6.09	1.74	-2.76	2.05	11.65	2.58	1.15	5.22	1.76	2.06	-1.01	-6.09

Table 37: Chromosomal locations and pre-microRNA sequences for target RNAs in Tables 36 and 37

Gene	Chromosomal Location	precursor sequence	SEQ ID NO
13108-L5-2	2p23.1	TTCCACACGTGCTCAGGCCCCCTCAACCAAGGAAGGCAGGCCCACTGGCCTCCTTATTACAGAGGGGCTGCAC GCACCCTAGGGAG	2681
13272-R5-2	12q13.11	GTCACACAGTTATGTTAGGCCATCACAGCATTTGGAATAGGGGATATCTCAGCATGTTGAGCCCTGTCTCTGGGAGC TGACTTCTACCTCTTCCAAAGGAGGCACGGTGATTGGAGTGC	2682
13316-R5-2	16p12.3,16p13.11	GCTCAATGCCCTCCAAACGAGTCAGCTGGGTCTCAGCCACGTACAACCTCACGGTGGAGGAGCCCCATCGTGGGCCTG GTGCTGTGGGCCAGCAGCAAGGTGGTGGCGCCCGGCA	2683
13331-L5-2	17q21.2	TGCGGAGTGCTGGTGCCGAGTGCTCAGAGTGGGTTCGGGTTCAGTCCCTGAACCCCAAGCATCCTCTGCACCCAGAT CCTGC	2684
13499-R5-1	7q32.2	AACAGTCCAGGGAAATGACAGTTGAGTTGCACAAACCCAGTGTTCAGACTTCCTGAGAAACTTGT	2685
5971-R5-2	1q24.1	TGCCATCTGCTCTGAAGCCTCCCAAGCTGGGCTCCCTCCCTCCACTTCTGGAGCCCCAGGAACAGATGTGGGAGGCTG AGCAGGCA	2686
7026-L3-1	15q15.3	TCGTTCCCGGATCTGGTGGGTGAGGTTTTCGATCAGGGCAAAATACCTGATCACAGACCTTCACAGGATTCTGGATGA	2687
7471-L5-1	1q21.1	GAAGTTCTTCTCCTTAGGGGGCATTTGCATTTTAAATGGGAATCTTAAAAACCCAAAGGAAATGTTCTCTAATGTGGG ATTTCT	2688
miR-320c	18q11.2	TTTGCAATTAAAAATGAGGCCCTTCTCTTCCAGTTCTTCCAGAGTCAGGAAAAAGCTGGGTTGAGAGGGTAGAAAAAA ATGATGTAGGCTTCTCTTCCAGTTCTTCCAGAAATGGGAAAAAGCTGGGTTGAGAGGGT	2690
miR-320c	18q11.2	CTTCTCTTCCAGTTCTTCCAGAAATGGGAAAAGCTGGGTTGAGAGGGT	2691

5.8 Example 8: Bioinformatic Analysis to Identify microRNAs

[00326] In order to identify the microRNAs detected with the probes shown, e.g., in Tables 1, 2, 6 to 9, 18 to 21, 23, 27, 28, 30, and 32 to 34, small RNA sequencing (smRNASeq) datasets were analysed using the probe sequences to identify expressed microRNAs detected by those sequences. The analysis identified 97 sequences with precise ends. Those 97 candidate microRNA sequences are show in Table 38.

Table 38: microRNA candidate sequences corresponding to probes

Name	Candidate sequences 5'->3'	SEQ ID NO
10233-R	TCTACTGCCTGCTGCTC	2576
10333-L	GGAGGGGGTGGGCAGGG	2577
10455-L	AGAGGGATGTTTGGCGC	2578
10520-L	CAGAGAAGGCTGGAGGAGG	2579
10844-R	AACACATGTTTGAAAGT	2580
11358-R	CATGTCAGCCTAGTTTCCC	2581
11605-L	AAAGAAGGGAAGAGAAGA	2582
11688-R	GTCGGCGTCTCCATCCTG	2583
12223-L	AGAGGGAGAAGAAACAA	2584
12699-R	CCACCACCGCCACCACCC	2585
12707-L	AGTTGTTGGATTGCAGA	2586
12723-R	CAGGAAGGGGCTGGGGG	2587
12730-R	CCCGGAGAGCGGAGCACAACACA	2588
12730-R	CCGGAGAGCGGAGCACAAC	2589
12911-L	GAGGAAAAGGAAGGAGG	2590
12912-L	CTGTGGGTGGAAGGTGCCAGAA	2591
12974-R	AAGTTAAATAACTCTGAACCA	2592
13071-L	ATGTGCCGAGATGTGAGCAGTC	2593
13071-L	ATGTGCCGAGATGTGAGCAGTCAC	2594
13108-L	CCAAGGAAGGCAGCAGGC	2595
13115-L	TTGGGGTGGTCGGCCCTGGAGG	2596
13122-L	GATGGAATTTCTAAAGG	2597
13124-L	GGAGGGGAGGAGACATG	2598
13124-L	GGAGGGGAGGAGACATG	2599
13163-R	GGGAGGGAGAGAAGGAG	2600
13163-R	GGGAGGGAGAGAAGGAGG	2601
13163-R	GGGAGGGAGAGAAGGAGGGG	2602
13185-L	AGAGGGAAGAAAAAAA	2603
13225-L	AGTGGGCAGGAAGAACCAGGCT	2604
13235-R	AGAGCTGAGACTAGAAAGCCCA	2605
13237-L	TTGGGGTGGTCGGCCCTGGAGG	2606

13245-L	GGGAGTAGAAGGGAAAGACTAT	2607
13247-L	GAGGTCGGGAGGGGAAGGCGGCT	2608
13252-L	TCAAGGAGCTCACAGTC	2609
13254-R	GCATGAGTGGTTCAGTGGT	2610
13272-R	TGACTTCTACCTCTTCCAAAG	2611
13274-L	TGAGGGAGGGTGGGAGC	2612
13278-R	GTGGAGTCCTGGGGAATGGAGA	2613
13287-L	AGGGTGGGGTGGCTCCTCTGCAG	2614
13309-L	TCCCTGTCCTCCACAAGCT	2615
13316-R	CAGCAGCAAGGTGGTGG	2616
13331-L	TGGTGCCGAGTGCTCAGAGTG	2617
13334-L	AGTGGGAGGGGAGGGGAGTCCTGCCA	2618
13343-L	TGAGGGGCCGGGAGCCA	2619
13352-R	TAGCCCCATTTCACAGAT	2620
13357-L	GCTGCAGGGAGGTGTGGGGA	2621
13358-L	AGTGTGGAGGGGGTGGTGA	2622
13366-R	TGGGAATAGGAGAGGGCACTG	2623
13373-R	TGCCCCACCTGCTGACCACCCTC	2624
13375-R	AACCTTGGCCCCTCTCCCCAG	2625
13395-R	GTGGAGTCCTGGGGAATGGAGA	2626
13398-R	CAAAGTGAGTCTAGTCTGCA	2627
13403-L	TTGGGGGGACACGGGTGG	2628
13436-L	AGCTGTGGTCACCGGAGCTCAGAGGC	2629
13467-L	GAGGGTCTGACTGTCACTTGGA	2630
13468-L	GAGGGTCTGATGGCCACTTGGA	2631
13472-L	CAGGAGGACGTCACACACAGTG	2632
13473-L	TAGGTGGTAGAAGGGCAAACA	2633
13499-R	CCCAGTGTTCAGACTTCCTG	2634
13500-L	TGGAGAGAGAGGCCTGGAAGA	2635
13500-L	TGGAGAGAGAGGCCTGGAAGAT	2636
13508-L	TGTGAGGAAGCTAAGAGCAG	2637
13522-L	AGTGCAGTGGTGTGATC	2638
13523-R	GCAGGGGAAGAAGCCTT	2639
13530-L	GGGGGACCCAGGGGAAGGAGG	2640
13545-L	ACAGCCTCCCTTGCGCCACAG	2641
25-R	TTAGAAAAAGAGGGGGTGAGG	2642
3249-L	GCGGCGGCGGCGGCGGC	2643
3371-L	TGGGGTGTGGAGGGGAGG	2644
3744-R	AGGGGAGCAGGGAGGAA	2645
3744-R	GCAGGGAGGAAGGAGAA	2646
3923-R	TGGAGAGGGATGAAGGAGAT	2647
4440-L	TCTGCCCAGTGCTCTGAATGTCA	2648

4440-L	CCAGTGCTCTGAATGTCAAA	2649
5080-R	TGGAGAGGGATGAAGGAGA	2650
5192-L	GAGGAAGGAAGGGGAAA	2651
5192-L	AGGAAGGAAGGGGAAAAA	2652
5232-L	CCTTGATGTCCTGCAAATGA	2653
5392-R	ATGAGATACTGTCGGAGA	2654
5723-R	CTGCTGGAGAAAGGGGAGGAGG	2655
5842-R	TTCATTTCTCTTTCATAA	2656
5971-R	ACTTCTGGAGCCCAGGAACA	2657
6037-R	TAGTTGGAGGAAAATTGGAG	2658
6216-L	TGCCCAGTGCTCTGAATGTC	2659
6216-R	ACGGCGGGAGTAACTATG	2660
6235-R	AAATGGATTTTTGGAGCAG	2661
6496-R	AGGGGAAGTGGTGGGTG	2662
6496-R	GTGGGTGGGGGAGGGGG	2663
6681-R	GGAGGGGCAGAGAGAGAG	2664
6864-R	AGGTGGGCACTCTAAGG	2665
6906-L	TGGTTGGGGAGAAGACA	2666
6930-R	TGCAAGATCAGAGGGGAGA	2667
7726-R	TCCTCTTCTCCTCTTCT	2668
8433-L	CGGTGGAGGGAAAGGGGAAA	2669
8452-L	AACACCCCAGCCATGTA	2670
8832-R	CAAAGTGGGGGAAAAACAGGTG	2671
9349-R	TCCGAGACCTGGAGCAG	2672

[00327] All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes.

[00328] While various specific embodiments have been illustrated and described, it will be appreciated that changes can be made without departing from the spirit and scope of the invention(s).

WHAT IS CLAIMED IS:

1. A method for detecting the presence of lung cancer in a subject, the method comprising detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692;

wherein a level of at least one target RNA in the sample that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer in the subject.

2. The method of claim 1, wherein the method further comprises comparing the level of the at least one target RNA in the sample to a normal level of the at least one target RNA.

3. A method for facilitating the detection of lung cancer in a subject, comprising:

(a) detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and

(b) communicating the results of the detection to a medical practitioner for the purpose of determining whether the subject has lung cancer.

4. The method of any one of the preceding claims, wherein detecting a level of at least one target RNA in a sample comprises:

- (a) hybridizing nucleic acids of the sample with at least one polynucleotide that is complementary to a target RNA in the sample or to a complement thereof; and
- (b) detecting at least one complex comprising a polynucleotide hybridized to at least one nucleic acid selected from the target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA.

5. A method for detecting the presence of lung cancer in a subject, comprising:

- (a) obtaining a sample from the subject,
 - (b) providing the sample to a laboratory for detection of the level of at least one target RNA in the sample, wherein the at least one target RNA:
 - (i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or
 - (ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or
 - (iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and
 - (c) receiving from the laboratory a communication indicating the level of at least one target RNA in the sample;
- wherein a level of at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer.

6. The method of any one of claims 1 to 4, wherein the method further comprises isolating nucleic acids from the sample.

7. The method of claim 6, wherein the nucleic acids comprise RNA that has been separated from DNA.

8. The method of any of the preceding claims, wherein at least one target RNA in its mature form comprises fewer than 30 nucleotides.

9. The method of any of the preceding claims, wherein at least one target RNA is a microRNA.

10. The method of any of the preceding claims, wherein levels of at least two target RNAs are detected, wherein at least two of the target RNAs:

(i) are capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and

wherein the at least two target RNAs are different.

11. The method of claim 10, wherein detection of a level of at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer.

12. The method of claim 10, wherein detection of levels of at least two target RNAs that are greater than normal levels of the at least two target RNAs indicates the presence of lung cancer.

13. The method of claim 10, wherein levels of at least three target RNAs are detected, wherein at least three of the target RNAs:

(i) are capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and

wherein the at least three target RNAs are different.

14. The method of claim 13, wherein detection of a level of at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of lung cancer.

15. The method of claim 13, wherein detection of levels of at least two target RNAs that are greater than normal levels of the at least two target RNAs indicates the presence of lung cancer.

16. The method of claim 13, wherein detection of levels of at least three target RNAs that are greater than normal levels of the at least three target RNAs indicates the presence of lung cancer.

17. The method of claim 10, wherein levels of at least five target RNAs are detected.

18. The method of any one of claims 10 to 17, wherein a level is detected of at least one target RNA that:

(i) does not specifically hybridize to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) does not comprise a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; and

(iii) does not comprise at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

19. The method of any one of the preceding claims, wherein at least one target RNA:

(a) is capable of specifically hybridizing to a polynucleotide sequence in Table 6; or

(b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 6.

20. The method of claim 19, wherein detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of non-small cell lung cancer.

21. The method of any one of claims 1 to 18, wherein at least one target RNA:

(a) is capable of specifically hybridizing to a polynucleotide sequence in Table 7; or

(b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 7.

22. The method of claim 21, wherein detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of squamous cell carcinoma.

23. The method of any one of claims 1 to 18, wherein at least one target RNA:

(a) is capable of specifically hybridizing to a polynucleotide sequence in Table 8; or

(b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 8.

24. The method of claim 23, wherein detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of adenocarcinoma.

25. The method of any one of claims 1 to 18, wherein at least one target RNA:

(a) is capable of specifically hybridizing to a polynucleotide sequence in Table 9; or

(b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 9.

26. The method of claim 25, wherein detection of a level of the at least one target RNA that is greater than a normal level of the at least one target RNA indicates the presence of aggressive lung cancer.

27. The method of any one of claims 1 to 18, wherein at least one target RNA:

(a) is capable of specifically hybridizing to a polynucleotide sequence in Table 32 or Table 33; or

(b) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a polynucleotide sequence in Table 32 or Table 33.

28. A method for detecting the presence of lung cancer in a subject, the method comprising detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452;

wherein a level of at least one target RNA in the sample that is reduced relative to a normal level of the at least one target RNA indicates the presence of lung cancer in the subject.

29. The method of claim 28, wherein the method further comprises comparing the level of the at least one target RNA in the sample to a normal level of the at least one target RNA.

30. The method of claim 28 or 29, wherein the method further comprises detecting a level of at least one second target RNA in a sample from the subject, wherein the at least one second target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692;

wherein a level of at least one second target RNA in the sample that is greater than a normal level of the at least one second target RNA indicates the presence of lung cancer in the subject.

31. A method for facilitating the detection of lung cancer in a subject, comprising:

(a) detecting a level of at least one target RNA in a sample from the subject, wherein the at least one target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from 1 to 397, 1363 to 1707, and 2312 to 2452; and

(b) communicating the results of the detection to a medical practitioner for the purpose of determining whether the subject has lung cancer.

32. The method of claim 31, wherein the method further comprises

(a) detecting a level of at least one second target RNA in a sample from the subject, wherein the at least one second target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692; and

(b) communicating the results of the detection to a medical practitioner for the purpose of determining whether the subject has lung cancer.

33. The method of any one of claims 28 to 32, wherein detecting a level of at least one target RNA in a sample comprises:

(a) hybridizing nucleic acids of the sample with at least one polynucleotide that is complementary to a target RNA in the sample or to a complement thereof; and

(b) detecting at least one complex comprising a polynucleotide hybridized to at least one nucleic acid selected from the target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA.

34. A method for detecting the presence of lung cancer in a subject, comprising:

(a) obtaining a sample from the subject;

(b) providing the sample to a laboratory for detection of the level of at least one target RNA in the sample, wherein the at least one target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1363 to 1707, and 2312 to 2452; and

(c) receiving from the laboratory a communication indicating the level of at least one target RNA in the sample;

wherein a level of at least one target RNA that is reduced relative to a normal level of the at least one target RNA indicates the presence of lung cancer.

35. The method of claim 34, wherein the sample is provided to a laboratory for detection of the level of at least one second target RNA in the sample, wherein the at least one second target RNA:

(i) is capable of specifically hybridizing to a nucleic acid having a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(ii) comprises a sequence that is complementary to at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 1 to 397, 1063 to 1210, 2064 to 2183, 2673 to 2680, and 2689; or

(iii) comprises at least 15 contiguous nucleotides of a sequence selected from SEQ ID NOs: 794 to 1043, 2576 to 2672, and 2692.

36. The method of any one of claims 28 to 33, wherein the method further comprises isolating nucleic acids from the sample.

37. The method of claim 36, wherein the nucleic acids comprise RNA that has been separated from DNA.

38. The method of any one of claims 28 to 37, wherein at least one target RNA in its mature form comprises fewer than 30 nucleotides.

39. The method of any one of claims 28 to 38, wherein at least one target RNA is a microRNA.

40. A synthetic polynucleotide comprising a first region, wherein the first region comprises a sequence of at least 8 contiguous nucleotides that is identical or complementary to a sequence of at least 8 contiguous nucleotides of one of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680.

41. The synthetic polynucleotide of claim 40, wherein the first region comprises a sequence of at least 9 contiguous nucleotides that is identical or complementary to a sequence of at least 9 contiguous nucleotides of one of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680.

42. The synthetic polynucleotide of claim 40, wherein the first region comprises a sequence of at least 10 contiguous nucleotides that is identical or complementary to a sequence of at least 10 contiguous nucleotides of one of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680.

43. The synthetic polynucleotide of claim 40, wherein the first region comprises a sequence of at least 12 contiguous nucleotides that is identical or complementary to a sequence of at least 12 contiguous nucleotides of one of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680.

44. The synthetic polynucleotide of any one of claims 40 to 43, wherein the polynucleotide comprises a detectable label.

45. The synthetic polynucleotide of claim 44, wherein the detectable label is a FRET label.

46. The synthetic polynucleotide of any one of claims 40 to 45, wherein the first region is identical or complementary to a region of a target RNA.

47. The synthetic polynucleotide of claim 46, wherein the polynucleotide further comprises a second region that is not identical or complementary to a region of the target RNA.

48. A composition comprising a plurality of synthetic polynucleotides, wherein at least one polynucleotide comprises a first region comprising a sequence of at least 8 contiguous nucleotides that is identical or complementary to a sequence of at least

8 contiguous nucleotides of one or more of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680.

49. The composition of claim 48, wherein at least two polynucleotides of the plurality of synthetic polynucleotides comprise a first region comprising a sequence of at least 9 contiguous nucleotides that is identical or complementary to a sequence of at least 9 contiguous nucleotides of one or more of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680, and wherein the first regions of the at least two polynucleotides are different.

50. The composition of claim 48, wherein at least three polynucleotides of the plurality of synthetic polynucleotides comprise a first region comprising a sequence of at least 10 contiguous nucleotides that is identical or complementary to a sequence of at least 10 contiguous nucleotides of one or more of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680, and wherein the first regions of the at least three polynucleotides are different.

51. The composition of claim 48, wherein at least five polynucleotides of the plurality of synthetic polynucleotides comprise a first region comprising a sequence of at least 12 contiguous nucleotides that is identical or complementary to a sequence of at least 12 contiguous nucleotides of one or more of SEQ ID NOs: 1 to 165, 254 to 360, 1063 to 1081, 1090 to 1190, 2064 to 2091, 2107 to 2168, 1363 to 1659, 2312 to 2440, and 2576 to 2680, and wherein the first regions of the at least five polynucleotides are different.

52. A kit comprising a synthetic polynucleotide of any one of claims 40 to 47.

53. A kit comprising a composition of any one of claims 48 to 51.

54. The kit of claim 52 or claim 53, wherein the kit further comprises at least one polymerase.

55. The kit of any one of claims 52 to 54, wherein the kit further comprises dNTPs.

1/1




FIG. 1