



(86) Date de dépôt PCT/PCT Filing Date: 2012/03/12
(87) Date publication PCT/PCT Publication Date: 2012/09/20
(85) Entrée phase nationale/National Entry: 2013/09/05
(86) N° demande PCT/PCT Application No.: US 2012/028706
(87) N° publication PCT/PCT Publication No.: 2012/125525
(30) Priorités/Priorities: 2011/03/14 (US61/452,390);
2011/07/21 (US61/510,226)

(51) Cl.Int./Int.Cl. *A61K 39/125* (2006.01),
A61K 39/145 (2006.01), *A61K 39/245* (2006.01)
(71) Demandeur/Applicant:
BOEHRINGER INGELHEIM VETMEDICA, INC., US
(72) Inventeurs/Inventors:
HAYES, PHILLIP WAYNE, US;
HENNESSY, KRISTINA J., US;
VIEL, LAURENT, CA;
DIAZ-MENDEZ, ANDRES, CA
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : VACCIN CONTRE LA RHINITE EQUINE
(54) Title: EQUINE RHINITIS VACCINE

(57) **Abrégé/Abstract:**

The disclosure provides for immunogenic compositions against Equine Rhinitis Virus, particularly Equine Rhinitis A and B Virus, and methods for their use and preparation. The immunogenic compositions, in alternate embodiments, also include other equine pathogens.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number

WO 2012/125525 A3

(43) International Publication Date
20 September 2012 (20.09.2012)

(51) International Patent Classification:

A61K 39/125 (2006.01) *A61K 39/245* (2006.01)
A61K 39/145 (2006.01)

(21) International Application Number:

PCT/US2012/028706

(22) International Filing Date:

12 March 2012 (12.03.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/452,390	14 March 2011 (14.03.2011)	US
61/510,226	21 July 2011 (21.07.2011)	US

(71) Applicant (*for all designated States except US*):

BOEHRINGER INGELHEIM VETMEDICA, INC.
[US/US]; 2621 North Belt Highway, St. Joseph, Missouri
64506-2002 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HAYES, Phillip Wayne** [US/US]; Boehringer Ingelheim Vetmedica, Inc., 2621 North Belt Highway, St. Joseph, Missouri 64506 (US). **HENNESSY, Kristina J.** [US/US]; 12700 Johnson Drive, Shawnee, Kansas 66216 (US). **VIEL, Laurent** [CA/CA]; RR#2 4176, Rockwood, Ontario NOB 2K0 (CA). **DIAZ-MENDEZ, Andres** [CA/CA]; 310 Middletown Road, Dundas, Ontario L9H 5E1 (CA).

(74) Agents: **MORRIS, Michael P.** et al.; Boehringer Ingelheim Pharmaceuticals, Inc., 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

(88) Date of publication of the international search report:

28 February 2013

(54) Title: EQUINE RHINITIS VACCINE

(57) Abstract: The disclosure provides for immunogenic compositions against Equine Rhinitis Virus, particularly Equine Rhinitis A and B Virus, and methods for their use and preparation. The immunogenic compositions, in alternate embodiments, also include other equine pathogens.



WO 2012/125525 A3

Equine Rhinitis Vaccine

Sequence Listing

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on January 26, 2012, is named 10-0144-PCT-SEQ.txt and is 33,877 bytes in size.

Background of the Invention

[0001] Equine viral respiratory infections are commonly associated with movement of horses and annually respiratory outbreaks are reported throughout the world. Equine Influenza 2 (AE2-H3N8), Equine Herpesvirus 1 and 4 (EHV1/4), and Equine Rhinitis A and B Viruses (ERAV and ERBV, respectively) are among the most important viruses isolated from the upper respiratory tract during respiratory outbreaks.

[0002] In particular, ERAV has been reported in acute febrile respiratory disease in horses (Li et al., J. Clin. Microbiol. 35:937-943; 1997, incorporated by reference). Similarly, a recent study in Ontario found ERBV and ERAV to be highly prevalent in the horse population (Diaz-Mendez et al., The Canadian Journal of Veterinary Research 74:271-278; 2010, incorporated by reference). Clinical signs of ERAV infection are non-specific and difficult to differentiate from other respiratory viral infections, including equine influenza and herpes virus infections. Non-cytopathic strains of this virus have been identified in equine respiratory outbreaks (Li et al., J. Clin. Microbiol. 35:937-943; 1997) making its diagnosis challenging. Moreover, ERAV and ERBV, being single stranded RNA viruses, have the potential for mutation, rendering the ability of the immune system to protect an animal against disease caused by a given ERAV/ERBV strain unclear.

[0003] A need exists for methods and medicaments for preventing respiratory diseases or for reducing the incidence or lessening the severity of clinical symptoms associated with such diseases, including those associated with Equine Rhinitis A and B Viruses.

Brief Summary of the Invention

[0004] The inventors have determined that immunogenic compositions comprising one or more strains of inactivated or live, attenuated ERAV, which when live and active and unattenuated are virulent (i.e., at least 50%, 60%, 70%, 80%, 90% or even 100% of seronegative horses, when purposely exposed to the virus, present with observable respiratory disease, particularly nasal and/or ocular discharge), can be grown to high titers in culture to yield a vaccine that is able to induce high titers of serum antibodies against ERAV when administered, for example, to an equine, for example resulting in a serum titer of at least 1:112, and preferably, at least 1:200, 1:500, 1:750 or 1:1000. In addition, the strains grow well in culture and are highly efficient to produce, for example, to a titer of at least 10^6 TCID₅₀/mL, more preferably at least 10^7 TCID₅₀/mL, 10^8 TCID₅₀/mL, or even 10^9 TCID₅₀/mL. Similarly, compositions comprising one or more strains of inactivated ERBV, which when live and active and not attenuated are also virulent (as defined above for ERAV), were also found to grow well in culture, to a titer of at least 10^6 TCID₅₀/mL, more preferably 10^7 TCID₅₀/mL, 10^8 TCID₅₀/mL or even 10^9 TCID₅₀/mL, to be highly efficient to produce, and to induce high titers of serum antibodies in animals following immunization, for example, resulting in a serum titer of at least 1:120 and preferably at least 1:200, 1:500, 1:750 or 1:1000. Both the ERAV and ERBV compositions, and combinations thereof, are capable of reducing the duration, severity, and incidence of disease in an animal such as a horse that has been immunized with the compositions and subsequently challenged.

[0005] Accordingly, the present invention provides an immunogenic composition comprising one or more strains of inactivated or live, attenuated ERAV or ERBV, wherein the ERAV strain or the ERBV strain, prior to inactivation or attenuation, causes detectable respiratory disease in at least 50% of seronegative horses exposed to the strain, or grows in cell culture to 10^6 TCID₅₀/mL or higher, or, when used as a vaccine in equines at a dose of 10^6 TCID₅₀ or higher results in a serum titer of at least 1:112.

[0006] In certain embodiments, the strain, when alive and unattenuated, causes detectable respiratory disease (e.g., detectable ocular and/or nasal discharge) in at least 50%, 60%, 70%, 80%, 90% or even 100% of seronegative horses upon exposure to the strain.

[0007] The strain can be grown in cell culture to a titer of at least 10^6 TCID₅₀/mL, more preferably 10^7 TCID₅₀/mL, 10^8 TCID₅₀/mL, or even 10^9 TCID₅₀/mL. Administration of an immunogenic composition containing the strain results in a serum titer of at least 1:120 and preferably at least 1:200, 1:500, 1:750 or 1:1000.

[0008] In the immunogenic compositions of the invention, the one or more strains of ERAV or ERBV preferably include ERAV/ON/05 (ATCC Accession No. PTA-11828) and/or ERBV strain 07-103042 (ATCC Accession PTA-11829). In addition, the immunogenic compositions of the invention contain at least an ERAV strain which comprises a genomic sequence whose reverse transcript has greater than 95% identity to SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to SEQ ID NO: 3, wherein said ERAV strain, when not inactivated, is active to infect and replicate in host cells. The immunogenic composition of the invention may also include an ERAV strain which comprises a genomic sequence whose reverse transcript has a nucleotide sequence comprising SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3.

[0009] The immunogenic compositions of the invention may also include an ERBV strain which comprises a genomic sequence whose reverse transcript has greater than 95% identity to the reverse transcript of the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829, wherein said ERBV strain, when not inactivated, is active to infect and replicate in host cells. The ERBV strain may also comprise a genomic sequence that is the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein that has the amino acid sequence of the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829.

[0010] In a specific embodiment, the immunogenic composition of the invention comprises Equine Rhinitis A Virus (ERAV) and Equine Rhinitis B Virus (ERBV), wherein the ERAV strain is ERAV/ON/05 and the ERBV strain has ATCC Accession No. PTA-11829.

[0011] In addition, the invention further includes multivalent immunogenic compositions comprising inactivated or live attenuated viruses or antigens from viruses other than ERAV or

ERBV that cause disease in Equidae. In particular, the invention provides immunogenic compositions comprising, in addition to inactivated or live, attenuated ERAV and/or ERBV, at least one antigen or one inactivated or live, attenuated strain of Equine Herpes Virus (EHV), and, in particular embodiments, the EHV is selected from the group consisting of EHV-1 and EHV-4, and a combination thereof, more specifically, the Equine Herpes Virus is selected from the group consisting of EHV-1, EHV-4, strains deposited with the ATCC under accession Nos. PTA-9525 and PTA-9526, and a combination thereof.

[0012] The invention further provides immunogenic composition, which, in addition to the inactivate or live, attenuated strain of ERAV and/or ERBV, at least one inactivated or live, attenuated strain of or at least one antigen of Equine Influenza Virus. In specific embodiments, the Equine Influenza Virus is selected from the group consisting of Clade 1 viruses, Clade 2 viruses, Influenza A/South Africa/2003, Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, Influenza A/equine-2/Richmond/1/2007, strains deposited with the ATCC under accession Nos. PTA-9522, PTA-9523, and PTA-9524, and combinations thereof. The immunogenic compositions may include, in addition to inactivated or live, attenuated ERAV and/or ERBV, at least one antigen or one inactivated or live, attenuated strain of Equine Herpes Virus and at least one antigen or one inactivated or live, attenuated strain of Equine Influenza Virus.

[0013] The immunogenic compositions of the invention may include, in addition to inactivated or live, attenuated ERAV and/or ERBV, at least one inactivated or live, attenuated virus or at least one antigen of one or more strains selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, and Tetanus Toxoid, and combinations thereof. Alternatively, the immunogenic composition, in addition to inactivated or live, attenuated ERAV and/or ERBV, comprises one or more inactivated or live, attenuated strains of or antigens of strains of Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis Virus, and Tetanus Toxoid. In specific embodiments, the West Nile Virus is one of the strains selected from the group consisting of Horse Origin 2005, deposited with the ATCC under accession number PTA-9409; NAEE159, deposited at the United States

Department of Agriculture Isolate under accession number 405330; NY2002Nassau; NY2002Clinton; NY2002Queens; GA20021; GA20022; TX20021; TX20022; IN2002; NY2003Albany; NY2003Suffolk; NY2003Chatauqua; CO20031; CO20032; TX2003; TX2003Harris4; TX2003Harris6; TX2003Harris7; TX2003Harris10; AZ2004; and TX2004Harris4; and combination thereof. In immunogenic compositions comprising Western Equine Encephalomyelitis Virus, the strain may be the strain deposited with the ATCC under accession number PTA-9410. In compositions comprising Venezuelan Equine Encephalomyelitis Virus, the strain may be the strain deposited with the ATCC under accession number PTA-9411. In immunogenic compositions comprising Eastern Equine Encephalomyelitis Virus, the strain may be the strain deposited with the ATCC under accession number PTA-9412. And, in immunogenic compositions comprising Equine Herpes Virus, the strain may be selected from the group consisting of the strains deposited with the ATCC under accession Nos. PTA-9525 or PTA-9526, and combinations thereof.

[0014] In specific embodiments, one or more of the strains in the immunogenic composition are present in an amount from about $10^{2.0}$ TCID₅₀/mL to about $10^{10.0}$ TCID₅₀/mL per dose. The composition may further include a suitable pharmaceutical carrier, such as a diluent, adjuvant, antimicrobial agent, preservative, inactivating agent, or combination thereof. In particular embodiments, the immunogenic composition comprises an adjuvant, specifically, HRA-5.

[0015] The invention further provides methods for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by Equine Rhinitis A Virus or Equine Rhinitis B Virus in an animal or a herd of animals comprising the step of administering an immunogenic composition that comprises one or more strains of inactivated or live, attenuated ERAV or ERBV, wherein the ERAV strain or the ERBV strain, prior to inactivation or attenuation, causes detectable respiratory disease in at least 50% of seronegative horses exposed to the strain, or grows in cell culture to 10^6 TCID₅₀/mL or higher, or, when used as a vaccine in equines at a dose of 10^6 TCID₅₀ or higher results in a serum titer of at least 1:112. In particular, the one or more strains of ERAV or ERBV preferably include ERAV/ON/05 (ATCC Accession No. PTA-11828) and/or ERBV strain 07-103042 (ATCC Accession PTA-11829). In addition,

the ERAV strain may comprise a genomic sequence whose reverse transcript has greater than 95% identity to SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to SEQ ID NO: 3, wherein said ERAV strain, when not inactivated or attenuated, is active to infect and replicate in host cells, or the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence comprising SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In addition, or alternatively, the ERBV strain may comprise a genomic sequence whose reverse transcript has greater than 95% identity to the reverse transcript of the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829, wherein said ERBV strain, when not inactivated or attenuated, is active to infect and replicate in host cells. The ERBV strain may also comprise a genomic sequence that is the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein that has the amino acid sequence of the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829.

[0016] In addition to providing methods for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV or ERBV in an animal or a herd of animals, the methods of the invention may further reduce the incidence or lessening the severity of clinical symptoms associated with or caused by one or more of the pathogens selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, and *Clostridium tetani* in an animal or a herd of animals by administering an immunogenic composition of the invention. The methods of the invention also include methods of reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV or ERBV in an animal or a herd of animals along with reducing the incidence or lessening the severity of clinical symptoms associated with or caused by one or more of the pathogens selected from the group consisting of: Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, Equine Herpes Virus, and *Clostridium*

tetani in an animal or a herd of animals by administering an immunogenic composition of the invention.

[0017] The invention also provides methods for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV and/or ERBV as well as one or more of the pathogens selected from the group consisting of: West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, Equine Herpes Virus, Equine Influenza Virus, and Clostridium tetani in an animal or a herd of animals, comprising the step of administering an immunogenic composition of the invention.

[0018] In connection with the methods of the invention, the incidence of clinical symptoms caused by one or more of said pathogens in a herd of animals is reduced from about 10% - 50% as compared to a herd not receiving the immunogenic composition. The methods of the invention, in particular embodiments, provide a duration of immunity of at least 12 months against one or more of the pathogens present in the immunogenic composition. In the methods of the invention, the immunogenic composition is administered to an Equidae, preferably a horse. The dosing scheme may include administration of the immunogenic composition in one or more doses. The doses for the methods of the invention may be formulated in 0.5 mL to 2.5 mL dosage forms. Preferably, the methods of the invention administer immunogenic compositions which are safe for use in foals or horses 4 months of age or older.

[0019] The invention also provides methods for producing an immunogenic composition comprising one or more strains of inactivated Equine Rhinitis A Virus (ERAV) or Equine Rhinitis B Virus (ERBV) as follows:

- a) infecting a susceptible cell line with ERAV or ERBV;
- b) growing the infected cell line in growth media until a cytopathic effect (CPE) is attained;
- c) harvesting the media;
- d) filtering the media to yield a filtered media; and
- e) contacting the filtered media with an inactivating agent to obtain the inactivated ERAV or ERBV.

Brief Description of the Drawings

- [0020] **Fig. 1** is a graphical representation of the proportion of virus positive across time.
- [0021] **Fig. 2** is a graphical representation of the proportion of buffy coat positive across time.
- [0022] **Fig. 3** is a graphical representation of the serum neutralization titers across time.
- [0023] **Fig. 4** is a graphical representation of the mean nasal scores across time.
- [0024] **Fig. 5** is a graphical representation of the mean ocular scores across time.
- [0025] **Fig. 6** is a ClustalW alignment of a portion of equine rhinitis A virus ERAV/ON/05 (5' UTR portion, SEQ ID NO: 1) with ERAV/PERV-1 (accession number: DQ272578 (SEQ ID NO: 14)). ERAV/ON/05 nucleotide insertions shown in bold and deletions in shading.
- [0026] **Fig. 7** is a graph of total clinical score means from control, infected, and re-infected groups from Example 4.
- [0027] **Fig. 8** is a graph of body temperature means from control, infected, and re-infected groups from Example 4.
- [0028] **Fig. 9** is a Table showing titers to equine rhinitis A virus (ERAV) and equine rhinitis B virus (ERBV) in control, infected, and re-infected groups of Example 4. The virus neutralization test (VN) was used to measure antibody titers on serum samples.

Detailed Description of the Invention

[0029] The inventors have determined that immunogenic compositions comprising one or more strains of inactivated ERAV, which when live and active are virulent (i.e., at least 50%, 60%, 70%, 80%, 90% or even 100% of seronegative horses, when purposely exposed to the virus, present with observable respiratory disease, particularly nasal and/or ocular discharge), can be grown to high titers in culture to yield a vaccine that is able to induce high titers of serum antibodies against ERAV when administered, for example, to an equine, for example resulting in a serum titer of at least 1:112, and preferably, at least 1:200, 1:500, 1:750 or 1:1000. In addition, the strains grow well in culture and are highly efficient to produce, for example, to a titer of at least 10^6 TCID₅₀/mL, more preferably 10^7 TCID₅₀/mL, 10^8 TCID₅₀/mL, or even 10^9 TCID₅₀/mL. Similarly, compositions comprising one or more strains of inactivated ERBV, which when live and active and not attenuated are also virulent (as defined above for ERAV), were also found to

grow well in culture, to a titer of at least 10^6 TCID₅₀/mL, more preferably 10^7 TCID₅₀/mL, 10^8 TCID₅₀/mL or even 10^9 TCID₅₀/mL, to be highly efficient to produce, and to induce high titers of serum antibodies in animals following immunization, for example, resulting in a serum titer of at least 1:120 and preferably at least 1:200, 1:500, 1:750 or 1:1000. Both the ERAV and ERBV compositions, and combinations thereof, are capable of reducing the duration, severity, and incidence of disease in an animal such as a horse that has been immunized with the compositions and subsequently challenged.

[0030] In one embodiment, provided is an immunogenic composition comprising one or more strains of inactivated ERAV and/or ERBV. Alternatively, the strains may be attenuated by routine means and the live, attenuated virus used in the vaccine composition. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In certain embodiments the ERAV strain is ERAV/ON/05 having ATCC Accession No. PTA-11828, deposited with the American Type Culture Collection on April 14, 2011, which is hereby incorporated by reference.

[0031] In some embodiments, the ERBV is strain 07-103042, having ATCC Accession No: PTA-11829, deposited with the American Type Culture Collection on April 14, 2011, which is hereby incorporated by reference. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence

of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0032] In one embodiment provided is an immunogenic composition comprising inactivated (or, alternatively, live, attenuated) ERAV and ERBV. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or

greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0033] In one embodiment, along with the inactivated or live, attenuated one or more strains of ERAV and/or ERBV, the immunogenic compositions provided herein further comprise at least one antigen or one additional inactivated or live, attenuated strain of Equine Herpes Virus (EHV). In some embodiments the compositions comprise at least one antigen of EHV. In some embodiments the EHV is selected from the group consisting of EHV-1, EHV-4, strains deposited with the ATCC under accession Nos. PTA-9525 and PTA-9526, and combinations thereof. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%,

greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0034] In one embodiment, along with the inactivated (or attenuated) one or more strains of ERAV and/or ERBV, the immunogenic compositions provided herein further comprise at least one antigen or one additional inactivated or attenuated strain of Equine Influenza Virus (EIV). In some embodiments the compositions comprise at least one antigen of EIV. In some embodiments the EIV is selected from the group consisting of Clade 1 viruses, Clade 2 viruses, Influenza A/South Africa/2003, Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, Influenza A/equine-2/Richmond/1/2007 and combinations thereof. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV

proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication

[0035] In one embodiment, along with the inactivated (or live, attenuated) one or more strains of ERAV and/or ERBV, the immunogenic compositions provided herein further comprise at least one antigen or one additional inactivated or live, attenuated strain of Equine Influenza Virus and at least one antigen or one additional inactivated or live, attenuated strain of Equine Herpes Virus. In some embodiments the compositions comprise at least one antigen of EHV and at least one antigen of EIV. In some embodiments the EHV is EHV-1 or EHV-4 or a combination thereof and the EIV is selected from the group consisting of Clade 1 viruses, Clade 2 viruses, Influenza A/South Africa/2003, Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, Influenza A/equine-2/Richmond/1/2007 and combinations thereof. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater

than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0036] In one embodiment, along with the inactivated (or live, attenuated) one or more strains of ERAV and/or ERBV, the immunogenic compositions provided herein further comprise at least one antigen or inactivated virus of one or more additional strains selected from the group consisting of Equine Influenza Virus, Equine Herpes Virus, West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, and Venezuelan Equine Encephalomyelitis Virus, and/or Tetanus Toxoid, and combinations thereof. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated is attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than

99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0037] In one embodiment, along with the inactivated (or live, attenuated) one or more strains of ERAV and/or ERBV, the immunogenic compositions provided herein further comprise at least one additional inactivated or live, attenuated virus of a strain selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, and Venezuelan Equine Encephalomyelitis Virus, and/or Tetanus Toxoid, and combinations thereof. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the

genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0038] In one embodiment, provided is a method of making the immunogenic composition of the present invention. The method generally comprises the steps of combining an inactivated or live, attenuated ERAV and/or ERBV and a pharmaceutically acceptable carrier. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828). In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater

than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication). In some embodiments the method further comprises the step of adding one or more additional equine virus antigens or inactivated or live, attenuated equine viruses. In another embodiment, the method further comprises the step of adding a suitable adjuvant to the composition.

[0039] In one embodiment, provided is a method for reducing the incidence of or lessening the severity of clinical symptoms associated with or caused by ERAV or ERBV in an animal or a herd of animals comprising administering an immunogenic composition as disclosed herein to an animal in need thereof. In some embodiments, the animal is a horse.

[0040] The aforementioned embodiments may further contain one or more of the following features described below.

[0041] In one embodiment, provided is a composition comprising at least one strain of inactivated (or, alternatively, live, attenuated) ERAV and/or ERBV and further containing many or all relevant antigenic components and proteins of pathogenic West Nile Virus (WNV) or an inactivated or live, attenuated strain of WNV.

[0042] In one embodiment, provided is a vaccine composition comprising one or more strains of inactivated (or live, attenuated) ERAV and/or ERBV in combination with one or more immunologically effective amounts of antigenic components or one or more inactivated or live, attenuated strains selected from the group consisting of West Nile Virus (WNV), Venezuelan Equine Encephalomyelitis (VEE), Eastern Equine Encephalomyelitis (EEE), Western Equine Encephalomyelitis (WEE), Tetanus toxoid (T), Equine herpes viruses (EHV) including types 1 and 4, Equine influenza viruses (EIV), and combinations thereof, along with a pharmaceutically acceptable carrier. Preferably such embodiments will include an adjuvant, such as a carbomer, and a pharmaceutically acceptable carrier. In other embodiments the adjuvant is HRA-5, a carbomer, or mineral oil.

[0043] In some embodiments, the compositions also include inactivated or live, attenuated ERAV and/or ERBV in combination with the following inactivated or live, attenuated viral strains or antigens and combinations of strains and antigens: West Nile Virus; Eastern Equine

Encephalomyelitis; Western Equine Encephalomyelitis; Venezuelan Equine Encephalomyelitis; Tetanus Toxoid; Eastern Equine Encephalomyelitis and Western Equine Encephalomyelitis; Eastern Equine Encephalomyelitis and Venezuelan Equine Encephalomyelitis; Eastern Equine Encephalomyelitis and Tetanus Toxoid; Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and Venezuelan Equine Encephalomyelitis; Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and Tetanus Toxoid; Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis and Tetanus Toxoid; Western Equine Encephalomyelitis and Venezuelan Equine Encephalomyelitis; Western Equine Encephalomyelitis and Tetanus Toxoid; Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, and Tetanus Toxoid; Venezuelan Equine Encephalomyelitis and Tetanus Toxoid; and Eastern Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis and Tetanus Toxoid, or antigens or antigenic components thereof. A preferred combination of these specified combinations includes ERAV and/or ERBV in combination with antigens or antigenic components of inactivated viruses of West Nile Virus, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, and Tetanus Toxoid. Another preferred combination includes ERAV and/or ERBV in combination with antigens or antigenic components of Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, and Tetanus Toxoid. Preferred ERAV include ERAV/ON/05(ATCC Accession No. PTA-11828), ERAV strain which comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1, which comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). The ERAV strain may also comprise a genomic sequence which, when reverse transcribed, has a nucleotide

sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. Preferred ERBV is a strain having ATCC Accession No: PTA-11829, or which comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication). In each such specified combination, an adjuvant or combination of adjuvants can be used such as, HRA-5, carbomer or with carbopol. The NJO strain of Eastern Equine Encephalomyelitis, the Fleming strain of Western Equine Encephalomyelitis strain, and the TC-83 strain of Venezuelan Equine Encephalomyelitis strain are all representative strains of these vaccine components.

[0044] Further preferred embodiments of the present invention include immunogenic compositions made using each of the specified combination vaccines listed above and adding antigens or inactivated or attenuated viruses from Equine Herpesvirus, preferably type 1, type 4, (EHV1 and/or EHV4) or combinations thereof.

[0045] Still further variations of each of the specified combination vaccines or immunogenic compositions listed above, including those that include EHV1 and/or EHV4 can be made by adding in antigens or inactivated or attenuated viruses from Equine influenza virus (EIV). Preferred embodiments incorporating Equine influenza virus include inactivated or live, attenuated: ERAV and/or ERBV and at least one inactivated or live, attenuated strain of each of WNV, Equine Influenza Virus, and Tetanus Toxoid; ERAV and/or ERBV and at least one inactivated or live, attenuated strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, and Eastern Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, Eastern Equine Encephalomyelitis, and Western Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine

Influenza Virus, Tetanus Toxoid, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis; and Venezuelan Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, and Eastern Equine Encephalomyelitis; ERAV and/or ERBV and, at least one strain of each of WNV, Equine Influenza Virus, and Western Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, and Venezuelan Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Eastern Equine Encephalomyelitis, and Western Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Eastern Equine Encephalomyelitis, and Venezuelan Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Western Equine Encephalomyelitis, and Venezuelan Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Western Equine Encephalomyelitis, and tetanus toxoid; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, and tetanus toxoid; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, Western Equine Encephalomyelitis, and tetanus toxoid; and ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, Eastern Equine Encephalomyelitis, and tetanus toxoid, wherein the aforementioned strains are inactivated or live, attenuated. In each specified embodiment one or more inactivated or live, attenuated strains of Equine Influenza Virus may be present. Preferred strains of Equine Influenza virus include Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, and combinations thereof. In all of the combinations listed above, it is preferred to use at least two inactivated or live, attenuated strains of Equine Influenza Virus and still more preferred to use at least 3 strains of Equine Influenza Virus.

[0046] Preferred embodiments incorporating Equine Herpes Virus include: ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, Eastern Equine Encephalomyelitis, and Equine Herpes Virus;

ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Tetanus Toxoid, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis; Venezuelan Equine Encephalomyelitis, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, and Eastern Equine Encephalomyelitis; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Western Equine Encephalomyelitis and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Eastern Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Western Equine Encephalomyelitis, Tetanus Toxoid, and Equine Herpes Virus; ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, Western Equine Encephalomyelitis, Tetanus Toxoid, and Equine Herpes Virus; and ERAV and/or ERBV and at least one strain of each of WNV, Equine Influenza Virus, Venezuelan Equine Encephalomyelitis, Eastern Equine Encephalomyelitis, Tetanus Toxoid, and Equine Herpes Virus, wherein the aforementioned strains are inactivated or live, attenuated. In all of the combinations listed above, it is preferred to use at least two strains of inactivated or live, attenuated Equine Influenza Virus and still more preferred to use at least 3 strains of inactivated or live, attenuated Equine Influenza Virus. Additionally, in all combinations above, the “at least one” strain of Equine Herpes virus is preferred to be selected from the group consisting of inactivated EHV-1 and EHV-4. In some preferred forms, both inactivated or live, attenuated strains, EHV-1 and EHV-4, will be included

in the immunogenic composition. In other preferred forms, just EHV-1 will be included. In a preferred combination, the inactivated or live, attenuated ERAV strain is ERAV/ON/05, an ERAV strain which comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1, which comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). The ERAV strain may also comprise a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In other preferred combinations, the ERBV is a strain having ATCC Accession No: PTA-11828, or which comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11828 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11828, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0047] The immunogenic compositions as disclosed herein can be administered in any immunogenically effective dose. In a preferred embodiment, the immunogenic composition is administered as a single dose. Preferably, the dose has a total volume between about 0.5 mL and about 2.5 mL, more preferably between about 0.6 mL and about 2.0 mL, even more preferably between about 0.7 mL and about 1.75 mL, still more preferably between about 0.8 mL and about

1.5 mL, even more preferably between about 0.9 mL and about 1.25 mL, with a single dose about 1.0 mL being the most preferred.

[0048] In another embodiment, the immunogenic composition is administered with a first dose being administered prior to the administration of a second (booster) dose. Preferably, the second dose is administered at least about 15 days after the first dose. More preferably, the second dose is administered between about 15 and about 28 days after the first dose. Even more preferably, the second dose is administered at least about 17 days after the first dose. Still more preferably, the second dose is administered between about 17 and about 25 days after the first dose. Even more preferably, the second dose is administered at least about 19 days after the first dose. Still more preferably, the second dose is administered between about 19 and about 23 days after the first dose. Most preferably the second dose is administered at least about 21 days after the first dose. In a preferred embodiment, both the first and second doses of the immunogenic composition are in the same amount. Preferably, each dose is in the preferred amounts specified above, with a dose of about 1 mL for the first and second dose being most preferred. In addition to the first and second dose regimen, an alternate embodiment comprises further subsequent doses. For example, a third, fourth, or fifth dose could be administered in these embodiments. Preferably, subsequent third, fourth, and fifth dose regimens are administered in the same amount as the first dose, with the time frame between the doses being consistent with the timing between the first and second doses mentioned above, although the timing may also vary.

[0049] In one embodiment the immunogenic composition is administered in three doses. In some embodiments the three doses are administered at three week intervals.

[0050] In an embodiment that comprises ERAV, preferably ERAV/ON/05, the amount of ERAV in the immunogenic composition is at least about $10^{2.0}$ TCID₅₀/dose. More preferably, the amount of ERAV is between about $10^{2.0}$ TCID₅₀/dose to about $10^{10.0}$ TCID₅₀/dose. Still more preferably, the amount of ERAV is at least about $10^{2.5}$ TCID₅₀/dose. Even more preferably, the amount of ERAV is between about $10^{2.5}$ TCID₅₀/dose to about $10^{9.5}$ TCID₅₀/dose. Still more preferably, the amount of ERAV is at least about $10^{3.0}$ TCID₅₀/dose. Even more preferably, the amount of ERAV is between about $10^{3.0}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Still more preferably, the amount of ERAV is at least about $10^{3.5}$ TCID₅₀/dose. Even more preferably, the

amount of ERAV is between about $10^{3.5}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Still more preferably, the amount of ERAV is between about $10^{6.5}$ TCID₅₀/dose and about $10^{8.5}$ TCID₅₀/dose. More preferably, the amount of ERAV is between about $10^{7.0}$ TCID₅₀/dose and about $10^{9.0}$ TCID₅₀/dose. The TCID₅₀ values of an inactivated or attenuated ERAV or any other inactivated or attenuated vaccine refer in general to the viral content in the final vaccine that however is equivalent to the viral content calculated for the vaccine composition prior to the inactivation of its virus. Preferably, the immunogenic composition of the present invention stimulates serum neutralizing antibodies to ERAV at a titer of at least 1:112, 1:300, 1:500, 1:700, 1:900, 1:1000, or 1:1500

[0051] In an embodiment that comprises ERBV, preferably a strain having ATCC Accession NO: PTA-11829, the amount of ERBV is at least about $10^{2.0}$ TCID₅₀/dose. More preferably, the amount of ERBV is between about $10^{2.0}$ TCID₅₀/dose to about $10^{10.0}$ TCID₅₀/dose. Still more preferably, the amount of ERBV is at least about $10^{2.5}$ TCID₅₀/dose. Even more preferably, the amount of ERBV is between about $10^{2.5}$ TCID₅₀/dose to about $10^{9.5}$ TCID₅₀/dose. Still more preferably, the amount of ERBV is at least about $10^{3.0}$ TCID₅₀/dose. Even more preferably, the amount of ERBV is between about $10^{3.0}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Still more preferably, the amount of ERBV is at least about $10^{3.5}$ TCID₅₀/dose. Even more preferably, the amount of ERBV is between about $10^{3.5}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Still more preferably, the amount of ERBV is between about $10^{6.5}$ TCID₅₀/dose and about $10^{8.5}$ TCID₅₀/dose. More preferably, the amount of ERBV is between about $10^{7.0}$ TCID₅₀/dose and about $10^{9.0}$ TCID₅₀/dose. The TCID₅₀ values of an inactivated ERBV or any other inactivated vaccine refer in general to the viral content in the final vaccine that however is equivalent to the viral content calculated for the vaccine composition prior to the inactivation of its virus. Preferably, the immunogenic composition of the present invention stimulates serum neutralizing antibodies to ERBV at a titer of at least 1:4 or higher. In some embodiments, the titer is at least 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:11; 1:12, 1:13, 1:14, or 1:15 or higher. In some embodiments, the titer is at least 1:64, 1:256, or 1:512 or higher. In some embodiments, the titer is at least 1:1024 or higher. In some embodiments, the titer is at least 1:2048, 1:1536, 1:3072,

1:4096, 1:6144, 1:8192, 1:12288, or 1:32769 or higher. In some embodiments, the titer is no more than 1:300, 1:1050, 1:32000, 1:70000, or 1:140000.

[0052] In one embodiment, in each dose of an embodiment of the present invention that comprises one or more additional equine antigens, the amount of Eastern Equine Encephalomyelitis or Venezuelan Equine Encephalomyelitis in any dose is preferably at least about $10^{5.5}$ TCID₅₀/dose. Even more preferably, the dose is between about $10^{5.5}$ TCID₅₀/dose and about $10^{9.5}$ TCID₅₀/dose. Still more preferably, the dose is at least about $10^{6.0}$ TCID₅₀/dose. Still more preferably, the dose is between about $10^{6.0}$ TCID₅₀/dose and about $10^{9.0}$ TCID₅₀/dose. Even more preferably, the dose is at least about $10^{6.5}$ TCID₅₀/dose. Still more preferably, the dose is between about $10^{6.5}$ TCID₅₀/dose and about $10^{9.5}$ TCID₅₀/dose. Even more preferably, the dose is at least about $10^{7.0}$ TCID₅₀/dose. Most preferably, the dose is between about $10^{6.7}$ TCID₅₀ and about $10^{9.2}$ TCID₅₀/dose.

[0053] In an embodiment that comprises inactivated or killed WNV or antigen, the amount of WNV or antigen is at least about $10^{2.0}$ TCID₅₀/dose. More preferably, the WNV or antigen is between about $10^{2.0}$ TCID₅₀/dose to about $10^{10.0}$ TCID₅₀/dose. Still more preferably, the WNV or antigen is at least about $10^{2.5}$ TCID₅₀/dose. Even more preferably, the WNV or antigen is between about $10^{2.5}$ TCID₅₀/dose to about $10^{9.5}$ TCID₅₀/dose. Still more preferably, the WNV or antigen is at least about $10^{3.0}$ TCID₅₀/dose. Even more preferably, the WNV or antigen is between about $10^{3.0}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Still more preferably, the WNV or antigen is at least about $10^{3.5}$ TCID₅₀/dose. Even more preferably, the WNV or antigen is between about $10^{3.5}$ TCID₅₀/dose to about $10^{9.0}$ TCID₅₀/dose. Most preferably, the WNV or antigen is between about $10^{7.0}$ TCID₅₀/dose and about $10^{9.0}$ TCID₅₀/dose. The TCID₅₀ values of an inactivated WNV vaccine or any other inactivated vaccine refer in general to the antigen content in the final vaccine that however is equivalent to the antigen content calculated for the vaccine composition prior to the inactivation of its antigen. Preferably, the immunogenic composition of the present invention stimulates serum neutralizing antibodies to WNV at a titer of at least 1:4 or higher. In some embodiments the titer is at least 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:11; 1:12, 1:13, 1:14, or 1:15 or higher. In some embodiments, the titer is no more than 1:300, 1:1050, 1:32000, 1:70000, or 1:140000. In a preferred embodiment, in each dose of an

embodiment of the present invention that comprises additional equine antigen, the amount of Eastern Equine Encephalomyelitis or Venezuelan Equine Encephalomyelitis in any dose is preferably at least about $10^{5.5}$ TCID₅₀/dose. Even more preferably, the dose is between about $10^{5.5}$ TCID₅₀/dose and about $10^{9.5}$ TCID₅₀/dose. Still more preferably, the dose is at least about $10^{6.0}$ TCID₅₀/dose. Still more preferably, the dose is between about $10^{6.0}$ TCID₅₀/dose and about $10^{9.0}$ TCID₅₀/dose. Even more preferably, the dose is at least about $10^{6.5}$ TCID₅₀/dose. Still more preferably, the dose is between about $10^{6.5}$ TCID₅₀/dose and about $10^{9.5}$ TCID₅₀/dose. Even more preferably, the dose is at least about $10^{7.0}$ TCID₅₀/dose. Most preferably, the dose is between about $10^{6.7}$ TCID₅₀ and about $10^{9.2}$ TCID₅₀/dose.

[0054] Preferably, the Western Equine Encephalomyelitis antigen, when present in the composition of the present invention, is in an amount of at least about $10^{6.2}$ PFU/mL. Even more preferably, the amount is between about $10^{6.2}$ PFU/mL and about $10^{10.2}$ PFU/mL. Still more preferably, the amount is at least about $10^{6.7}$ PFU/mL. Even more preferably, the amount is between about $10^{6.5}$ PFU/mL and about $10^{9.7}$ PFU/mL. Still more preferably, the amount is at least about $10^{7.2}$ PFU/mL. Even more preferably, the amount is between about $10^{7.2}$ PFU/mL and about $10^{9.2}$ PFU/mL. Still more preferably, the amount is at least about $10^{7.7}$ PFU/mL with between about $10^{6.5}$ PFU/dose and about $10^{9.0}$ PFU/mL being the most preferred.

[0055] In another preferred embodiment, the amount of tetanus toxoid, if present in the composition of the present invention, is in an amount of at least about 3 CPU, more preferably, between about 3 CPU and about 20 CPU, still more preferably, at least about 4 CPU, and most preferably, at least about 5 CPU but not more than about 20 CPU.

[0056] In an alternate embodiment, where one or more strains of Equine Influenza Virus is present, the amount of Equine Influenza present in the composition is in an amount of at least about $10^{5.0}$ TCID₅₀/mL. More preferably, the Equine Influenza is in an amount of between about $10^{5.0}$ TCID₅₀/mL to about $10^{9.0}$ TCID₅₀/mL, and, more preferably, at least about $10^{6.0}$ TCID₅₀/mL. Still more preferably, the amount is between about $10^{6.0}$ TCID₅₀/mL to about $10^{8.0}$ TCID₅₀/mL and, more preferably, the amount is at least about $10^{6.5}$ TCID₅₀/mL. Still more preferably, the amount is between about $10^{6.5}$ TCID₅₀/mL to about $10^{7.5}$ TCID₅₀/mL, with the most preferred amount being between about $10^{6.7}$ TCID₅₀/mL to about $10^{7.3}$ TCID₅₀/mL.

[0057] In an embodiment that comprises Equine Herpes Virus, the amount of Equine Herpes Virus in each dose is at least about $10^{6.0}$ TCID₅₀/mL. More preferably, Equine Herpes Virus is present in the composition in an amount of between about $10^{6.0}$ TCID₅₀/mL to about $10^{9.5}$ TCID₅₀/mL and, more preferably, in an amount of about $10^{7.0}$ TCID₅₀/mL. Still more preferably, Equine Herpes Virus is present in an amount between about $10^{7.5}$ TCID₅₀/mL to about $10^{9.0}$ TCID₅₀/mL and, more preferably, in an amount of about $10^{8.0}$ TCID₅₀/mL. Still more preferably, Equine Herpes Virus is present in an amount of between about $10^{8.0}$ TCID₅₀/mL to about $10^{9.0}$ TCID₅₀/mL and, most preferably, in an amount of about $10^{8.50}$ TCID₅₀/mL.

[0058] In yet another preferred embodiment, a vaccine composition comprising the chronologically contemporary and epidemiologically prevalent strains of ERAV and/or ERBV is provided. Preferably the composition comprises ERAV/ON/05 and/or a ERBV strain having ATCC Accession NO: PTA-11829. In specific embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising the nucleotide sequence of SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV

proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication). Such a composition will generally improve the efficacy of the composition.

[0059] The present invention additionally provides for a method of reduction of the incidence of and/or severity of clinical signs associated with, ERAV and/or ERBV infection in an animal, preferably a horse. Such methods generally comprise the step of administering a vaccine composition comprising an inactivated or live, attenuated strain of an ERAV and/or ERBV and a pharmaceutically acceptable carrier. In particular embodiments, the ERAV is ERAV/ON/05, or the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater

than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication). In some preferred embodiments, an adjuvant, particularly HRA-5, is added to the composition, and in other preferred forms, no adjuvant is provided.

[0060] In an alternate preferred embodiment, the method comprises administering a vaccine composition comprising one or more inactivated or live, attenuated strains of ERAV and/or ERBV in combination with immunologically effective amounts of antigenic components or inactivated strains from other equine pathogens. Preferably, the ERAV strain is ERAV/ON/05 (ATCC Accession No. PTA-11828) and the ERBV strain has ATCC Accession NO: PTA-11829. In certain such embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or live, attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of

the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

[0061] In some embodiments of the method, the pathogens in combination with the ERAV and/or ERBV strains, are selected from the group consisting of antigens or inactivated or attenuated strains of EHV and EIV and combinations thereof. In some embodiments the pathogens are antigens. In some embodiments the EHV is EHV-1 or EHV-4 or a combination thereof. In other embodiments the EIV is selected from the group consisting of Clade 1 viruses, Clade 2 viruses, Influenza A/South Africa/2003, Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, Influenza A/equine-2/Richmond/1/2007 and combinations thereof.

[0062] In still other embodiments of the method, the pathogens in combination with the ERAV and/or ERBV strains, are selected from the group consisting of antigens or inactivated or attenuated strains of WNV, Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and Venezuelan Equine Encephalomyelitis, and tetanus toxoid, and combinations thereof, and more preferably being those combinations described above. In another preferred embodiment, the vaccine of the present invention is combined with a suitable adjuvant and/or pharmaceutically acceptable carrier.

[0063] The present invention provides for reduction of the incidence of and/or severity of clinical symptoms associated with, ERAV and/or ERBV infection in a herd. Preferably, the severity and/or incidence of clinical symptoms in animals receiving the immunogenic composition of the present invention are reduced at least 10% in comparison to animals not receiving such an administration when both groups (animals receiving and animals not receiving the composition) are challenged with or exposed to infection by ERAV and/or ERBV. More preferably, the incidence or severity is reduced at least 20%, even more preferably, at least 30%, still more preferably, at least 40%, even more preferably, at least 50%, still more preferably, at least 60%, even more preferably, at least 70%, still more preferably, at least 80%, even more preferably, at least 90%, still more preferably, at least 95%, and most preferably, at least 100%, wherein the animals receiving the composition of the present invention exhibit no clinical symptoms, or alternatively exhibit clinical symptoms of reduced severity. Advantageously, the

present invention also provides protection from heterologous strains (relative to the strain used in the composition) of pathogens.

[0064] The present invention further provides a method of stimulating serum neutralizing or serum hemagglutination antibodies to a pathogen selected from the group consisting of ERAV, ERBV, WNV, WEE, VEE, EEE, EHV, EIV, and combinations thereof by administering a composition in accordance with the present invention described herein. In particular embodiments, the ERAV is ERAV/ON/05, or the ERAV strain comprises a genomic sequence whose reverse transcript has a 5'UTR comprising SEQ ID NO: 1. In some embodiments, the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of SEQ ID NO: 2 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERAV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of SEQ ID NO: 3, which polyprotein contains functional ERAV proteins (i.e., active in viral infection and replication). In some embodiments, the ERAV strain comprises a genomic sequence which, when reverse transcribed, has a nucleotide sequence of SEQ ID NO: 2 or which encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3. In some embodiments, the ERBV is a strain having ATCC Accession No: PTA-11829. In some embodiments, the ERBV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the nucleotide sequence of the reverse transcript of the genome of the ERBV strain having ATCC Accession No. PTA-11829 and, when not inactivated or attenuated, is active to infect and replicate in host cells and/or encodes functional ERBV proteins, or which encodes a polyprotein having an amino acid sequence with greater than 95%, greater than 96%, greater than 97%, greater than 98% or greater than 99% identity to the amino acid sequence of the polyprotein of the strain with ATCC Accession No. PTA-11829, which polyprotein contains functional ERBV proteins (i.e., active in viral infection and replication).

Preferably the compositions of the present invention stimulate serum neutralizing antibodies to ERAV and/or ERBV at a titer of at least 1:112, 1:120, 1:300, 1:500, 1:1000, or 1:1024, or higher.

[0065] The immunogenic composition of the present invention provides an extended duration of immunity against all strains present in the vaccine. Preferably, the duration of immunity against ERAV and/or ERBV is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4-24 months, still more preferably, the duration of immunity is at least 6-24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0066] The immunogenic composition of the present invention also provides an extended duration of immunity against all antigens present in the vaccine. Preferably, the duration of immunity against West Nile is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4-24 months, still more preferably, the duration of immunity is at least 6-24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0067] Preferably, the duration of immunity against EIV is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4-24 months, still more preferably, the duration of immunity is at least 6-24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24

months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0068] Preferably, the duration of immunity against EHV is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4- 24 months, still more preferably, the duration of immunity is at least 6 -24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0069] Preferably, the duration of immunity against Western Equine Encephalomyelitis is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4- 24 months, still more preferably, the duration of immunity is at least 6 -24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0070] Preferably, the duration of immunity against Eastern Equine Encephalomyelitis is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4- 24 months, still more preferably, the duration of immunity is at least 6 -24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0071] Preferably, the duration of immunity against Venezuelan Equine Encephalomyelitis is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of

immunity is at least 4- 24 months, still more preferably, the duration of immunity is at least 6 -24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0072] Preferably, the duration of immunity against Tetanus Toxoid is at least 1 month, more preferably, the duration of immunity is at least 2 months, still more preferably, the duration of immunity is at least 3 months, even more preferably, the duration of immunity is at least 4- 24 months, still more preferably, the duration of immunity is at least 6 -24 months, even more preferably, the duration of immunity is at least 7-24 months, still more preferably, the duration of immunity is at least 8-24 months, even more preferably, the duration of immunity is at least 9-24 months, still more preferably, the duration of immunity is at least 10-24 months, and most preferably, the duration of immunity is at least 12-24 months.

[0073] Preferably, the duration of immunity of at least 12 months further relates to any combination of antigens forming the immunogenic composition of the present invention.

[0074] In one embodiment comprising an inactivated (or, alternatively live attenuated) ERAV and/or ERBV as disclosed herein, the immunogenic composition ameliorates shedding of infectious ERAV and/or ERBV to prevent spread of the virus to other susceptible animals. In some embodiments the compositions prevent shedding of the virus.

[0075] In one embodiment comprising EIV and/or EHV antigen, as described above, the immunogenic composition ameliorates shedding of infectious EIV or EHV to prevent spread of the virus to other susceptible animals.

[0076] In one embodiment, compositions in accordance with the present invention described herein overcome interference from passively acquired maternal immunity and stimulate active immunity and a reduction in the incidence of or severity of clinical signs of EIV infection in vaccinated animals against EIV.

[0077] In one embodiment of the present invention, an immunogenic composition comprising ERAV and/or ERBV, VEE, WEE, EEE, tetanus, WNV, equine rhinopneumonitis and equine influenza, all as described herein, demonstrates efficacy against ERAV, ERBV, VEE,

WEE, EEE, tetanus, WNV, equine rhinopneumonitis and equine influenza after administration in accordance with the present invention. Preferably, such a composition will further include an adjuvant, preferably HRA-5, mineral oil and/or a carbomer, and a pharmaceutically acceptable carrier. In preferred forms, the composition will be administered in a single, 1 mL dose. In some embodiments composition is administered in two doses or preferably three doses, with each dose separated by 1, 2, 3, and 4 weeks.

[0078] Each of the immunogenic compositions described herein that include ERAV, particularly ERAV/ON/05 (ATCC Accessions NO: PTA-11828) and other strains as described *supra*, and/or ERBV, particularly a ERBV strain having ATCC Accession NO: PTA-11829, or others as also described *supra*, can be administered as described such that they reduce the incidence of or lessen the severity of clinical symptoms associated with ERAV and/or ERBV, such as pyrexia, elevations in temperature, increased lung sounds, lymphadenopathy, nasal discharge, ocular discharge, pharyngitis, edema of legs, cough, and in the case of ERAV, increased incidence of abortion in pregnant mares. In some aspects, the compositions lessen the amount or length of nasal or ocular discharge or the length of time that such symptoms are presented. In some aspects, animals inoculated with the compositions show no clinical symptoms of ERAV and/or ERBV infection one week or longer after exposure to ERAV and/or ERBV. In other aspects, animals inoculated with the compositions show no clinical symptoms of ERAV and/or ERBV infection when exposed to ERAV and/or ERBV. Clinical symptoms of ERAV and ERBV may be scored such as according to Table 3 in Example 2, Table 14 in Example 4, or Table 17 in Example 5.

[0079] Each of the immunogenic compositions described herein that include EIV antigen or inactivated EIV and can be administered as described such that they reduce the incidence of or lessen the severity of clinical symptoms associated with Equine Influenza Virus.

[0080] The present invention also provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Equine Herpes Virus comprising the step of administering any one of the immunogenic compositions described above containing EHV antigen or inactivated or attenuated EHV to an animal.

[0081] The present invention also provides a method for reducing the incidence of clinical symptoms associated with West Nile Virus comprising the step of administering any one of the immunogenic compositions that includes WNV antigen or inactivated or attenuated WNV, as described herein, to an animal.

[0082] The present invention also provides a method for reducing the incidence of clinical symptoms associated with Equine Influenza Virus comprising the step of administering any one of the immunogenic compositions described above, that includes an EIV antigen or inactivated or attenuated EIV, to an animal.

[0083] The present invention further provides a method for reducing the incidence of clinical symptoms associated with Equine Herpes Virus comprising the step of administering any one of the immunogenic compositions described above that includes an EHV antigen or inactivated or attenuated EHV, to an animal.

[0084] The present invention provides a method of reducing the incidence of viral infection in a herd comprising the step of administering any one of the immunogenic compositions described above to an animal, wherein the reduction of incidence of infection, compared to herds not receiving the immunogenic composition, is from about 10% to about 50% reduction. In one embodiment the compositions provided herein reduce ERAV infection by 10% to 50%. In other embodiments the compositions provided herein reduce ERBV infection by 10% to 50%.

[0085] The present invention also provides a method of reducing the incidence of clinical symptoms associated with Equine Influenza Virus comprising the step of administering any one of the immunogenic compositions described above to an animal, wherein the reduction in clinical signs, compared to animals not receiving the immunogenic composition, is at least a 10% reduction in clinical signs.

[0086] The present invention provides a method of reducing the incidence and severity of clinical symptoms of ERAV or ERBV in a herd, wherein the clinical symptoms are selected from the group consisting of pyrexia, elevations in temperature, increased lung sounds, lymphadenopathy, nasal discharge, ocular discharge, pharyngitis, cough, edema of legs, and in the case of ERAV, increased incidence of abortion in pregnant mares.

[0087] The present invention provides a method of reducing the incidence and severity of clinical symptoms of EHV in a herd, wherein the clinical symptoms are selected from the group consisting of respiratory disease, abortion, reproductive complications, neurological disease, central nervous system disease, and combinations thereof.

[0088] The present invention provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Equine Herpes Virus comprising the step of administering any one of the immunogenic compositions disclosed herein, that includes an EHV antigen or inactivated or attenuated EHV, to an animal.

[0089] The present invention provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Equine Influenza Virus in a herd, comprising the step of administering any one of the immunogenic compositions disclosed herein, that includes an EIV antigen or inactivated or attenuated EIV, to an animal.

[0090] The present invention provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with West Nile Virus in a herd, comprising the step of administering any one of the immunogenic compositions disclosed herein, that includes a WNV antigen or inactivated or attenuated WNV, to an animal.

[0091] The present invention provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Eastern Equine Encephalomyelitis in a herd, comprising the step of administering any one of the immunogenic compositions disclosed herein that includes an EEE virus antigen or an inactivated or attenuated EEE virus, to an animal.

[0092] The present invention further provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Western Equine Encephalomyelitis in a herd, comprising the step of administering any one of the immunogenic compositions disclosed herein, that includes a WEE virus antigen or an inactivated or attenuated WEE virus, to an animal.

[0093] The present invention further provides a method for reducing the incidence of or lessening the severity of clinical symptoms associated with Venezuelan Equine Encephalomyelitis in a herd, comprising the step of administering any one of the immunogenic

compositions disclosed herein, that includes a VEE virus antigen or an inactivated attenuated VEE virus, to an animal.

[0094] The aforementioned embodiments may be used in a combination therapy or as part of a immunization schedule in combination with other immunogenic agents and vaccines. In one embodiment, the compositions provided herein are used in combination with the immunogenic agents and vaccines described in WO 2010/025469, which is incorporated herein by reference in its entirety.

[0095] The present invention also provides a method of making any one of the immunogenic composition as described above and herein, comprising the steps of combining an inactivated or live, attenuated ERAV or ERBV with a suitable pharmaceutical carrier. In preferred forms, this method further comprises the step of adding one or more equine antigens or inactivated or attenuated viruses. A preferred group of equine antigens and viruses are selected from the group consisting of West Nile Virus, Western Equine Encephalomyelitis, Eastern Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, EHV, and EIV, and tetanus toxoid, and combinations thereof. In some preferred forms, the methods described herein can further comprise a filtration step, wherein the final product is in a more pure form.

[0096] “About” refers to $\pm 10\%$ of the specified quantity.

[0097] “Animals” as used herein includes domesticated animals including dogs and hooved animals including equidae, and specifically, horses. In some embodiments, the term also refers to a human.

[0098] “Adjuvants” as used herein, can include aluminum hydroxide and aluminum phosphate, saponins e.g., Quil A, QS-21 (Cambridge Biotech Inc., Cambridge MA), GPI-0100 (Galenica Pharmaceuticals, Inc., Birmingham, AL), non-metabolizable oil, mineral and/or plant/vegetable and/or animal oils, polymers, carbomers, surfactants, natural organic compounds, plant extracts, carbohydrates, cholesterol, lipids, water-in-oil emulsion, oil-in-water emulsion, water-in-oil-in-water emulsion, HRA-3 (acrylic acid saccharide cross-linked polymer), HRA-3 with cottonseed oil (CSO), or preferably HRA-5 (acrylic acid polyol cross-linked polymer). The emulsion can be based in particular on light liquid paraffin oil (European Pharmacopeia type); isoprenoid oil such as squalane or squalene; oil resulting from the oligomerization of alkenes, in

particular of isobutene or decene; esters of acids or of alcohols containing a linear alkyl group, more particularly plant oils, ethyl oleate, propylene glycol di-(caprylate/caprate), glyceryl tri-(caprylate/caprate) or propylene glycol dioleate; esters of branched fatty acids or alcohols, in particular isostearic acid esters. The oil is used in combination with emulsifiers to form the emulsion. The emulsifiers are preferably nonionic surfactants, in particular esters of sorbitan, of mannide (e.g. anhydromannitol oleate), of glycol, of polyglycerol, of propylene glycol and of oleic, isostearic, ricinoleic or hydroxystearic acid, which are optionally ethoxylated, and polyoxypropylene-polyoxyethylene copolymer blocks, in particular the Pluronic products, especially L121. See Hunter et al., *The Theory and Practical Application of Adjuvants* (Ed. Stewart-Tull, D. E. S.) John Wiley and Sons, NY, pp51-94 (1995) and Todd et al., *Vaccine* 15:564-570 (1997). In a preferred embodiment the adjuvant is at a concentration of about 0.01 to about 50%, preferably at a concentration of about 2% to 30%, more preferably at a concentration of about 5% to about 25%, still more preferably at a concentration of about 7% to about 22%, and most preferably at a concentration of about 10% to about 20% by volume of the final product.

[0099] As used herein, “a pharmaceutically acceptable carrier” or “pharmaceutical carrier” includes any and all excipients, solvents, growth media, dispersion media, coatings, adjuvants, stabilizing agents, diluents, preservatives, inactivating agents, antimicrobial, antibacterial and antifungal agents, isotonic agents, adsorption delaying agents, and the like. Such ingredients include those that are safe and appropriate for use in veterinary applications. In some preferred embodiments, and especially those that include lyophilized immunogenic compositions, stabilizing agents for use in the present invention include stabilizers for lyophilization or freeze-drying.

[0100] “Diluents” can include water, saline, dextrose, ethanol, glycerol, and the like. Isotonic agents can include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin and alkali salts of ethylenediaminetetracetic acid, among others.

[0101] In a preferred embodiment, the immunogenic composition of the present invention is prepared comprising a preservative and a stabilizer; and, more preferably, the immunogenic

composition of the present invention is prepared comprising Amphotericin, formaldehyde, gentamycin, EDTA, glycerol, and combinations thereof.

[0102] An “immunogenic or immunological composition” refers to a composition of matter that comprises at least one antigen, which elicits an immunological response in the host of a cellular and/or antibody-mediated immune response to the composition or vaccine of interest. Usually, an “immunological response” includes but is not limited to one or more of the following effects: the production or activation of antibodies, B cells, helper T cells, suppressor T cells, and/or cytotoxic T cells and/or gamma-delta T cells, directed specifically to an antigen or antigens included in the composition or vaccine of interest. Preferably, the host will display either a therapeutic or protective immunological response such that resistance to new infection will be enhanced and/or the clinical severity of the disease reduced. Such protection will be demonstrated by either a reduction or lack of clinical signs normally displayed by an infected host, a quicker recovery time and/or a lowered duration or bacterial titer in the tissues or body fluids or excretions of the infected host.

[0103] The term “in need of such administration” or “in need of such administration treatment,” as used herein means that the administration/treatment is associated with the boosting or improvement in health or any other positive medicinal effect on health of the animals which receive the immunogenic composition in accordance with the present invention, such as reducing the incidence or severity of a viral infection or disease.

[0104] “Equine rhinitis A virus (ERAV)” refers to an *Aphthovirus* in the family *Picornaviridae*, and was previously known as Equine rhinovirus 1. “ERAV” as used herein includes inactivated forms. In one embodiment, the ERAV is strain ERAV/ON/05 having accession number PTA-11829 deposited on April 14, 2011 with the ATCC (American Type Culture Collection, P.O. Box 1549 Manassas, VA 20108 USA) in accordance with the Budapest Treaty, and that was recovered from Rabbit-kidney-13 (RK-13) cell culture from a nasal swab from a horse in Ontario Canada in 2005. The ERAV/ON/05 when reverse transcribed and sequenced has SEQ ID NO: 1 in its 5' UTR region.

[0105] The reverse transcribed genomic sequence of ERAV/ON/05 is SEQ ID NO:2.

[0106] The polyprotein encoded by the genomic sequence of ERAV/ON/05 is SEQ ID NO:3.

[0107] The ERAV strains useful in the immunogenic compositions of the invention have reverse transcribed 5'UTR nucleotide sequences that have greater than 85%, 90%, 95%, 98%, 99% or 100% sequence identity to SEQ ID NO:1. In some embodiments, the ERAV strains have genomic sequences that when reverse transcribed into DNA are greater than 85%, 90%, 95%, 98%, 99% or 100% identical to SEQ ID NO:2 or have polyprotein coding sequences that are more than 97%, 98%, 99%, or 100% identical to the polyprotein coding sequence in SEQ ID NO:2. In some embodiments, the ERAV have polyproteins with an amino acid sequence greater than 95%, 95%, 97%, 98%, 99%, or 100% identical to SEQ ID NO:3 or to the VP1 sequence within SEQ ID NO:3. In yet other embodiments, the ERAV has an L protein, VP2, VP3, VP4, VP0, 2A, 2B, 2C, 3A, 3B, 3C or 3D, that have an amino acid sequence with greater than 80%, greater than 90%, greater than 99%, or 100% identity to the same protein found in SEQ ID NO:3. All of the ERAV strains are, when not inactivated or attenuated, infective and able to replicate in host cells.

[0108] "Equine rhinitis B virus (ERBV)" refers to an *Erbovirus* in the family *Picornaviridae*, and was previously known as Equine rhinovirus 2. "ERBV" as used herein includes inactivated forms. In one embodiment, the ERBV is a strain deposited with the ATCC that was recovered from Rabbit-kidney-13 (RK-13) cell culture from a nasal swab from a horse in Ontario Canada (ATCC Accession NO: PTA-11828) that was deposited with the ATCC (American Type Culture Collection, P.O. Box 1549 Manassas, VA 20108 USA) on April 14, 2011 under the Budapest Treaty. The ERBV strains useful in the immunogenic compositions of the invention have genomic sequences that when reverse transcribed into DNA are greater than 85%, 90%, 95%, 98%, 99% or 100% identical to the reverse transcript of the genomic sequence of the ERBV strain having ATCC Accession No. PTA-11829, or have polyprotein coding sequences that are more than 97%, 98%, 99%, or 100% identical to the polyprotein coding sequence of the ERBV strain having ATCC Accession No. PTA-11829. In some embodiments, the ERBV strains useful in the invention have polyproteins with an amino acid sequence greater than 95%, 95%, 97%, 98%, 99%, or 100% identical to the polyprotein sequence of the ERBV

strain having ATCC Accession No. PTA-11829. In yet other embodiments, the ERBV has an L protein, VP4, VP2, VP3, VP1, 2A, 2B, 2C, 3A (Vpg), 3B, 3Cpro, 3Dpol that have an amino acid sequence with greater than 80%, greater than 90%, greater than 99%, or 100% identity to the same protein found in the ERBV strain having ATCC Accession No. PTA-11829. All of the ERBV strains are, when not inactivated or attenuated, infective and able to replicate in host cells.

[0109] The term “West Nile Virus” antigen means, but is not limited to the components of the WNV virion that are immunogenic when present in an animal, and most particularly protein components, such as envelope and non-structural proteins, of the WNV that provoke humoral or cellular immune responses when present in an animal. Such antigens can include DNA, protein subunits, modified live virus, and inactivated virus. In preferred forms of the invention, the WNV antigen or antigens comprise inactivated or killed, and even more preferably, North American dominant, WNV strains.

[0110] The term “North American West Nile Virus (strains)” refers to, but is not limited to any West Nile Virus strain that has ever been discovered on the North American continent. Preferably, a North American West Nile Virus strain has a sequence identity to the NY99 strain (GenBank accession no. AF196835 or NCBI reference sequence NC_00942.1 of at least 97%, even more preferably, at least 98%, still more preferably, at least 98.5%, more preferably, at least 99%, even more preferably, at least 99.2%, and, most preferably of at least 99.4%. WN02 is a representative example of a WNV strain that can be referred to as a North American Dominant West Nile Virus strain. Specifically, North American Dominant strains are those having at least 1 nucleotide change resulting in an amino acid change from the WN99 isolates. Strain NY99 (GenBank accession no. AF196835) serves as a reference strain for determining if a strain is North American Dominant. In addition, these strains may have one or more silent amino acid changes. In some embodiments, the nucleotide change results in an amino acid change in an envelope protein of the strain and, more preferably, the nucleotide change results in an amino acid change from valine to alanine. Preferably, this amino acid change is associated with a greater ability to replicate in the intermediate host, namely, the mosquito. More preferably, North American Dominant strains include either (and preferably both) a U to C mutation and a C to U mutation at positions 1442 and 2466 (in comparison to a North American strain, e.g., NY

99), respectively. Still more preferably, North American Dominant strains further include a mutation in the nucleotide sequence encoding the E protein and the C to U mutation at position 9352 in the sequence encoding the NS5 protein (again in comparison to a North American strain, e.g., NY 99). These preferred mutations are shown in Phylogenetic Analysis of North American West Nile Virus Isolates, 2001-2004: Evidence For the Emergence of a Dominant Genotype, C. Todd Davis, et. al, Virology 342, p. 252-265 (2005), the teaching and content of which is hereby incorporated by reference. West Nile Virus strains, for purposes of the present invention, are not limited to horse and equine West Nile Virus strains but encompass, while not being limited to, those West Nile Virus strains of bird origin, donkey origin, pig origin, human origin, mammal origin, and equine origin.

[0111] “Sequence Identity” as it is known in the art refers to a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, namely a reference sequence and a given sequence to be compared with the reference sequence. Sequence identity is determined by comparing the given sequence to the reference sequence after the sequences have been optimally aligned to produce the highest degree of sequence similarity, as determined by the match between strings of such sequences. Upon such alignment, sequence identity is ascertained on a position-by-position basis, e.g., the sequences are “identical” at a particular position if at that position, the nucleotides or amino acid residues are identical. The total number of such position identities is then divided by the total number of nucleotides or residues in the reference sequence to give % sequence identity. Sequence identity can be readily calculated by known methods, including but not limited to, those described in Computational Molecular Biology, Lesk, A. N., ed., Oxford University Press, New York (1988), Biocomputing: Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York (1993); Computer Analysis of Sequence Data, Part I, Griffin, A.M., and Griffin, H. G., eds., Humana Press, New Jersey (1994); Sequence Analysis in Molecular Biology, von Heinge, G., Academic Press (1987); Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York (1991); and Carillo, H., and Lipman, D., SIAM J. Applied Math., 48: 1073 (1988), the teachings of which are incorporated herein by reference. Preferred methods to determine the sequence identity are designed to give the largest match between the sequences

tested. Methods to determine sequence identity are codified in publicly available computer programs which determine sequence identity between given sequences. Examples of such programs include, but are not limited to, the GCG program package (Devereux, J., et al., *Nucleic Acids Research*, 12(1):387 (1984)), BLASTP, BLASTN and FASTA (Altschul, S. F. et al., *J. Molec. Biol.*, 215:403-410 (1990)). The BLASTX program is publicly available from NCBI and other sources (BLAST Manual, Altschul, S. et al., NCVI NLM NIH Bethesda, MD 20894, Altschul, S. F. et al., *J. Molec. Biol.*, 215:403-410 (1990), the teachings of which are incorporated herein by reference). These programs optimally align sequences using default gap weights in order to produce the highest level of sequence identity between the given and reference sequences. As an illustration, by a polynucleotide having a nucleotide sequence having at least, for example, 85%, preferably 90%, even more preferably 95% "sequence identity" to a reference nucleotide sequence, it is intended that the nucleotide sequence of the given polynucleotide is identical to the reference sequence except that the given polynucleotide sequence may include up to 15, preferably up to 10, even more preferably up to 5 point mutations per each 100 nucleotides of the reference nucleotide sequence. In other words, in a polynucleotide having a nucleotide sequence having at least 85%, preferably 90%, even more preferably 95% identity relative to the reference nucleotide sequence, up to 15%, preferably 10%, even more preferably 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 15%, preferably 10%, even more preferably 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. Analogously, by a polypeptide having a given amino acid sequence having at least, for example, 85%, preferably 90%, even more preferably 95% sequence identity to a reference amino acid sequence, it is intended that the given amino acid sequence of the polypeptide is identical to the reference sequence except that the given polypeptide sequence may include up to 15, preferably up to 10, even more preferably up to 5 amino acid alterations per each 100 amino acids of the reference amino acid sequence. In other

words, to obtain a given polypeptide sequence having at least 85%, preferably 90%, even more preferably 95% sequence identity with a reference amino acid sequence, up to 15%, preferably up to 10%, even more preferably up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 15%, preferably up to 10%, even more preferably up to 5% of the total number of amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or the carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in the one or more contiguous groups within the reference sequence. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. However, conservative substitutions are not included as a match when determining sequence identity. In the present disclosure, it is understood that SEQ ID NO:1 (5' UTR) and SEQ ID NO:2 are the DNA sequences that result from reverse transcription of the 5' UTR and the entire genome of ERAV/ON/05, respectively. Likewise, SEQ ID NO: 3 refers to the amino acid sequence corresponding to the polyprotein encoded by the genomic sequence of ERAV/ON/05. Percent identity of a given ERAV strain in comparison to SEQ ID NO:1 or SEQ ID NO:2 is thus meant to refer to the corresponding DNA sequence resulting from reverse transcription and sequencing.

[0112] "TCID₅₀" refers to tissue culture infective dose infecting 50% of cells in a culture inoculated with the virus.

[0113] For purposes of the present invention the terms "strain" and "isolate" have the same meaning and are used interchangeably.

[0114] "Clinical signs" or "clinical symptoms" for ERAV and ERBV include but are not limited to pyrexia, elevation in temperature, increased lung sounds, lymphadenopathy, nasal discharge, ocular discharge, and cough, and pharyngitis. Still other signs or symptoms include anemia, anorexia, lymphadenitis of the head and neck, edema of the legs, lethargy, and pain. Additionally, clinical signs of ERAV and/or ERBV infections may include those associated with Equine Herpes virus and Equine Influenza virus. In one embodiment the clinical signs for ERAV include cough, pharyngitis, pyrexia, elevations in temperature, increased submandibular

lymph node size, nasal discharge, and ocular discharge. In certain embodiments, the clinical signs of ERAV include an increased incidence of abortion in pregnant mares. In another embodiment the clinical signs to be addressed by an immunological composition disclosed herein are those of respiratory infections, such as those cause by one or more of ERAV, ERBV, EIV, EHV-1, and EHV-4.

[0115] “Clinical signs” or “clinical symptoms” of West Nile Virus, for purposes of this invention, include, but are not limited to, symptoms or lesions associated with encephalitis, viremia, anorexia, depression, fever, weakness, abnormal gait, paralysis of hind limbs, impaired vision, ataxia, aimless wandering, convulsions, inability to swallow, coma, posterior weakness, paralysis, poor coordination, depression and related behavior, tremors, convulsions, paddling of the limbs, neurological problems, swelling of the central nervous system, death, and combinations thereof. The clinical signs exhibited by an infected animal vary depending on the severity of infection.

[0116] “Clinical Signs” or “clinical symptoms” of Equine Herpes virus, for purposes of this invention include, but are not limited to, abortion, neurological deficiencies, respiratory disease, reproductive system deficiencies and failure, and symptoms relating to the central nervous system. Additionally, clinical symptoms of EHV1 include, but are not limited to, the phenomenon of foals infected with EHV1, exhibiting respiratory complications, passing the virus to the older members of the herd who then exhibit reproductive deficiencies, including abortion, and neurological deficiencies, normally exhibited in the central nervous system.

[0117] “Clinical Signs” or “clinical symptoms” of Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, and Venezuelan Equine Encephalomyelitis, for purposes of the present invention are those symptoms normally known to be associated with encephalomyelitis, including, but are not limited to fever, nervous signs such as sensitivity to sound, periods of excitement, and restlessness, brain lesions, drowsiness, drooping ears, circling, abnormal gait, paralysis, loss of appetite, depression, head pressing, lack of coordination, long-term disability, brain damage, death, and combinations thereof. “Safety” as used herein, refers to the absence of adverse consequences in the vaccinated animal following vaccination, including but not limited to, potential reversion of vaccine virus to virulence and clinically significant side

effects, such as persistent systemic illness or unacceptable inflammation at the site of vaccine administration.

[0118] “Reduction of the incidence and/or severity of clinical signs” or “reduction in the incidence and/or severity of clinical symptoms,” as referred to herein, means reducing the number of infected animals in a group, reducing or eliminating the number of animals exhibiting clinical signs of infection, or reducing the severity of any clinical signs that are present in the animals, in comparison to wild-type infection. For example, such clinical signs included viremia, fever, antibody response, ocular discharge, nasal discharge, and histopathology. Preferably, these are reduced in animals receiving the composition of the present invention by at least 10% in comparison to animals not receiving the vaccination which may become infected. More preferably, clinical signs are reduced in animals receiving the composition of the present invention by at least 20%, more preferably by at least 30%, even more preferably by at least 40%, and even more preferably by at least 50%.

[0119] “Duration of Immunity,” as used herein, refers to the minimum number of days during which an animal produces an immunogenic response such that the animal will be relatively immune from contracting a virus and/or benefit from reduction of incidence and/or severity of clinical signs, as described herein.

[0120] The term “inactivated” and “inactivated virus” refers to a previously virulent virus that has been heated or chemically treated to inactivate, kill, or otherwise modify the virus to substantially eliminate its virulent properties while retaining its immunogenicity. In a preferred embodiment, the inactivated viruses disclosed herein are inactivated by treatment with an inactivating agent. Suitable inactivating agents include beta-propiolactone, binary ethyleneimine, glutaraldehyde, and formaldehyde. In some embodiments, the inactivating agent is formaldehyde.

[0121] The terms “vaccine” and “immunogenic composition”, when used herein, are meant to be used interchangeably.

[0122] Any West Nile Virus strain(s) or isolate(s) can be used in accordance with the present invention. In a preferred embodiment, the isolate is selected from one or more of the following: New York (Northeastern North American) Isolate (WN-NY 99), Horse Origin, 1999,

New York (Northeastern North American) Isolate (WN-NY 99), Crow Origin, 1999, United States Department of Agriculture Isolate 292206 (USDA 2004), Donkey Origin, United States Department of Agriculture Isolate 405330 (USDA 2005), Horse Origin, North American Isolate (WN-Texas-2002/2003), Southeast Texas Coastal Isolate 2002, Mexico (Tabasco) Isolate 2003, and combinations thereof, and in a more preferred embodiment the isolate is selected from one or more of the following: United States Department of Agriculture Isolate 292206 (USDA 2004), Donkey Origin, United States Department of Agriculture Isolate 405330 (USDA 2005), Horse Origin, North American Isolate (WN-Texas-2002/2003), Southeast Texas Coastal Isolate 2002, Mexico (Tabasco) Isolate 2003, and combinations thereof. In a most preferred embodiment, the isolate is United States Department of Agriculture Isolate 405330 (USDA 2005), Horse Origin singularly or in combination with one or more isolates as listed above. In an additionally preferred embodiment, those isolates which are part of the North American West Nile Virus isolates are included. In yet another preferred embodiment North American Dominant West Nile Virus isolates are included. In addition to those listed above, specific isolates include, but are not limited to, WN02 and isolates which have at least 1, preferably at least 2, and even more preferably at least 3 nucleotide changes resulting in at least one amino acid change from the WN NY99 isolates, and most preferred are strains with the amino acid change from valine to alanine at position 159 of the envelope protein. Most preferred North American Dominant strains include, but are not limited to: NY2002Nassau, NY2002Clinton, NY2002Queens, GA20021, GA20022, TX20021, TX20022, IN2002, NY2003Albany, NY2003Suffolk, NY2003Chatauqua, CO20031, CO20032, TX2003, TX2003Harris4, TX2003Harris6, TX2003Harris7, TX2003Harris10, AZ2004, and TX2004Harris4, and combinations thereof. The strains of West Nile Virus useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. In a preferred embodiment, the North American Dominant West Nile Virus strain used is either E-159 (Horse Origin) or E-159 (Donkey Origin). A representative strain of such a North American Dominant WNV strain includes the Horse Origin 2005 strain deposited with the ATCC (ATCC Accession NO: PTA-9409), located at 10801 University Boulevard, Manassas, VA, 20110-2209, on August 14, 2008, under the provisions of the Budapest Treaty. Equine Influenza strains useful in the vaccine or immunogenic composition of the present

invention can be any strain or isolate. Representative strains include Equi-2/Ohio/03, deposited as ATCC Accession NO: PTA-9522, Equi-2/Kentucky/95, deposited as ATCC Accession NO: PTA-9523, and Equi-2/New Market/2/93, deposited as ATCC Accession NO: PTA-9524. Representative strains ATCC Accession Nos. PTA-9522, PTA-9523, and PTA-9524 were each deposited with the ATCC at 10801 University Boulevard, Manassas, VA, 20110-2209 on September 23, 2008, under the provisions of the Budapest Treaty.

[0123] Equine Herpes Virus (“EHV”) strains useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. Representative strains include EHV Subtype 1, deposited as ATCC Accession NO: PTA-9525, and EHV Subtype 4, deposited as ATCC Accession NO: PTA-9526. Representative strains ATCC Accession Nos. PTA-9525 and PTA-9526 were each deposited with the ATCC at 10801 University Boulevard, Manassas, VA, 20110-2209 on September 23, 2008, under the provisions of the Budapest Treaty.

[0124] Western Equine Encephalomyelitis strains useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. A representative strain includes the Fleming Strain, deposited with the ATCC (ATCC Accession NO: PTA-9410), located at 10801 University Boulevard, Manassas, VA, 20110-2209, on August 14, 2008, under the provisions of the Budapest Treaty.

[0125] Venezuelan Equine Encephalomyelitis strains useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. A representative strain includes the TC-83 strain, deposited with the ATCC (ATCC Accession NO: PTA-9411), located at 10801 University Boulevard, Manassas, VA, 20110-2209, on August 14, 2008, under the provisions of the Budapest Treaty.

[0126] Eastern Equine Encephalomyelitis strains useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. A representative strain includes the NJO strain, deposited with the ATCC (ATCC Accession NO: PTA-9412), located at 10801 University Boulevard, Manassas, VA, 20110-2209, on August 14, 2008, under the provisions of the Budapest Treaty.

[0127] Tetanus toxoid strains useful in the vaccine or immunogenic composition of the present invention can be any strain or isolate. A representative strain is that taken from a master

seed of *Clostridium tetani* from The Massachusetts Department of Public Health Institute of Laboratories in Boston, Massachusetts.

[0128] The vaccine or immunogenic composition as disclosed herein is safe for administration in ERAV or ERBV susceptible species, particularly equidae, at any age. In a preferred embodiment, the present invention is safe for administration to foals 12 months of age or older, more preferably, it is safe for administration to foals 10 months of age or older, more preferably, it is safe for administration to foals 8 months or older, more preferably, it is safe for administration to foals 6 months of age or older, more preferably, is safe for administration to foals 4 months of age or older, more preferably, it is safe for administration to foals 2 months of age or older, more preferably, it is safe for administration to foals 1 month of age or older, even more preferably, it is safe for administration to foals between 1 day and 1 month of age, and, most preferably, it is safe for administration to foals 1 day of age or older.

[0129] The compositions as disclosed herein can be administered in any conventional manner. Examples of administration methods include any that afford access by cells of the immune system to the immunogenic composition including oral, transdermal/intradermal, intravenous, subcutaneous, intramuscular, intraocular, intraperitoneal, intrarectal, intravaginal, intranasal, intragastrical, intratracheal, intrapulmonarial, or any combination thereof. In a preferred embodiment, the vaccine is administered parenterally, preferably intranasally, subcutaneously, or intramuscularly, and in the most preferred embodiment the vaccine is administered intramuscularly.

[0130] In one embodiment, provided is a method for preparing and immunogenic composition comprising an ERAV and/or ERAB as disclosed herein. In one embodiment, the method comprises:

- a) infecting a susceptible cell line with ERAV or ERBV;
- b) growing the infected cell line in growth media until a cytopathic effect (CPE) is attained;
- c) harvesting the media;
- d) filtering the media to yield a filtered media; and

e) contacting the filtered media with an inactivating agent to obtain the inactivated ERAV or ERBV.

[0131] In one embodiment, the method comprises providing strain ERAV/ON/05 or a ERBV strain having ATCC Accession NO: PTA-11829. These strains are used to infect a susceptible cell line having advantageous growth and secretion properties useful for vaccine preparation. A preferred susceptible cell line is a Vero cell line such as a Vero76 or an E-Vero cell line.

[0132] Suitable growth media include E199. This media may be supplemented with L-glutamine solution and an antibiotic such as gentamicin sulfate solution. In some embodiments the growth media is E199 supplemented with L-glutamine solution (up to 2 mM), and gentamicin sulfate solution (up to 30 µg/mL). In some embodiments, viral fluids are harvested when the cytopathic effect CPE reaches 75% or greater. The harvested media are pooled and filtered such as through 5.0 micron pore-size rated polypropylene filter cartridges. The strains are inactivated chemically such as by treatment with 0.1-0.2% formaldehyde solution over a suitable period of time, for example 48 hours to 72 hours and the resulting fluids are concentrated. In some embodiments, preservatives and adjuvants are next added. Suitable preservatives include one or more of Amphotericin B, gentamicin sulfate, and formaldehyde. In other embodiments the adjuvant is HRA-5. In still other embodiments the adjuvant is HRA-3 with cottonseed oil (CSO).

[0133] In one embodiment and in accordance with the methods disclosed herein, provided is an immunogenic composition comprising ERAV/ON/05 and/or a ERBV strain having ATCC Accession NO: PTA-11829 and Amphotericin B, gentamicin sulfate, formaldehyde, and HRA-5 prepared according to the methods disclosed herein.

Examples

[0134] The following examples are set forth below to illustrate specific embodiments of the present invention. These examples are merely illustrative and are understood not to limit the scope or the underlying principles of the present invention.

Example 1

[0135] This example illustrates one embodiment of a Equine Rhinitis A Virus composition in accordance with the present invention.

Materials and Methods

[0136] Equine Rhinitis A Virus strain ERAV/ON/05 (ATCC Accession No. PTA-11828) was recovered from Rabbit-kidney-13 (RK-13) cell culture from a nasal swab from a horse in Ontario Canada. The virus was passaged once on E-Vero cells to produce a high pre-titered master stock, and was then diluted with cell culture media to produce the Master Seed Virus. Master Cell Stock is an E-Vero cell line grown and maintained using E199 supplemented with L-Glutamine solution (up to 2 mM) and Gentamicin sulfate solution (up to 30 µg/mL). Inactivated ERAV/ON/05 was produced according to the following general procedure.

[0137] Frozen Master Cell Stock is thawed at room temperature (18-26°C) and used to inoculate a range of pre-sterilized T-25 cm² up to T-150 cm² flasks or pre-sterilized 850 cm² or 1050 cm² PETG roller bottles. Thawed cells are suspended in growth medium at the rate of 0.15 to 0.40 mL per cm². Cells are incubated at 36-38°C for up to seven days. Cultures planted from frozen stock may be re-fed with medium to remove residual Dimethyl Sulfoxide (DMSO), to remove excessive debris, to stimulate the growth of cultures which have not reached confluency, or to maintain viability of confluent cultures. Cells are passaged by decanting the spent medium and adding 1-10 mL (depending on the size of the vessel), of 25% trypsin-EDTA solution to each vessel. The vessels are agitated gently until the cells slough from the surface. The cells are removed from the vessels by rinsing with growth medium and pooled together. The range of cell culture passages is one to twenty.

[0138] Prior to inoculation, cell growth medium is decanted from cells that are at least 90% confluent. The cell sheet is rinsed with 20-50 mL of virus infection media (E199 supplemented with L-Glutamine solution (up to 2 mM) and Gentamicin sulfate solution (up to 30 µg/mL), then re-fed with virus infection media at the rate of 0.15 to 0.40 mL/cm². Vessels are then inoculated at a Multiplicity of Infection (MOI) of 0.0005 to 0.005. Roller bottle cultures are incubated at 36-38°C for two to five days at 0.2-0.4 rpm.

[0139] During the growth period, cultures are checked for CPE microscopically and for gross contamination macroscopically. Unsuitable cultures are discarded after sterilization.

[0140] Virus fluids are harvested when CPE reaches 75% or greater. Roller bottles are swirled to remove loose cells, and fluids are pooled into sterile 2-20 L glass, plastic, or PETG bottles, 20 L sterile polypropylene containers or 2-200 L sterile stainless steel tanks appropriate for clarification. Pooled fluids may be held for up to seven days at 2-8°C prior to clarification.

[0141] Only fluids from monolayer cultures showing evidence of viral infection are harvested. Bottles indicating contamination are discarded. A sample of pooled, clarified fluids is collected before inactivation for titration by TCID₅₀. Fluids with a titer of 10^{6.2} TCID₅₀/mL or greater may be used in the preparation of final product. Multiple lots can be blended to achieve the minimum titers.

[0142] Harvested lots are clarified by filtration using 5.0 micron pore-size rated polypropylene filter cartridges. Post-clarified harvest lots may be stored for up to seven days at 2-8°C. Clarified fluids are then inactivated with formaldehyde solution, USP, 0.1-0.2% by volume, transferred to a secondary container, and held at room temperature (18-26°C) with agitation for a minimum of 48 to 72 hours. A sample of inactivated fluids is taken for inactivation assurance testing prior to concentration. Inactivated lot material is held at 2-8°C prior to concentration. Clarified, inactivated virus fluids are concentrated by a factor of 5X to 50X using tangential flow ultra-filtration membrane cartridges with molecular weight cut-off ratings of not more than 10,000 Dalton MW.

[0143] A number of suitable adjuvants may be added to the vaccine formulation, most preferably HRA-5. Other adjuvants include HRA-3 with cottonseed oil (CSO). Typical processing steps may be employed such as mixing, blending, microfluidization, and emulsification, of the adjuvant and/or the harvested virus antigens with other ingredients.

[0144] The product is then assembled to final formulation. In a batching method in one exemplary embodiment, the required amounts of adjuvant and MEME Diluent are combined in a sterile container and the pH is adjusted to approximately 6.3-6.7 with Sodium Hydroxide. ERAV, gentamicin sulfate, formaldehyde, and Amphotericin B are added one at a time during constant mixing. The pH is adjusted to 6.8-7.0 with sodium hydroxide or hydrochloric acid and

the serial is mixed for a minimum of 18 hours at 2-8°C. The completed bulk serial is then transferred to a suitable storage container and stored at 2-8°C. Amphotericin B may be added at a concentration of up to 2.5 µg/mL of the diluent volume as a preservative. Gentamicin sulfate is added at a concentration of up to 30 µg/mL of the diluent volume as a preservative. Formaldehyde is added at a concentration of 0.1% of the diluent volume as a preservative. An adjuvant such as HRA-5 is added at a concentration of 10% v/v of the final serial volume.

[0145] The vaccine is given by typical hypodermic injection, with booster vaccinations if desired.

Example 2

[0146] This investigation was carried out to obtain an efficacy evaluation of a Equine Rhinitis A Virus vaccine to protect horses from challenge with Equine Rhinitis A Virus.

Materials and Methods

[0147] A high dose A-9 ($10^{7.5}$ TCID₅₀/mL) and low dose A-10 ($10^{7.0}$ TCID₅₀/mL) ERAV vaccine were prepared according to Example 1.

Table 1: A-9 vaccine formulation (1 mL):

ERAV/ON/05	$10^{7.5}$ TCID ₅₀ /mL
HRA-5	100 µL
Diluent, MEM-E+ containing Gentamicin, 30 µg/mL of diluent volume and Formaldehyde, 0.1% of diluent volume	q.s.

Table 2: A-10 1 vaccine formulation (1 mL):

ERAV/ON/05	$10^{7.0}$ TCID ₅₀ /mL
HRA-5	100 µL
Diluent, MEM-E+ containing Gentamicin, 30 µg/mL of diluent volume and Formaldehyde, 0.1% of diluent volume	q.s.

[0148] A V-05 placebo 1 mL formulation was prepared as in the A-9 and A-10 vaccines but without the ERAV/ON/05 antigen.

[0149] A total of 44 horses were randomly divided into one of three treatment groups consisting of a 15 horse V-05 placebo control group, a 14 horse A-10 low-dose group, and a 15 horse A-9 high-dose group. Horses were vaccinated with a 1 mL dose product adjuvanted with HRA-5 for three total doses, with a 21 day interval in between doses. Horses were subsequently challenged 21 days after the third vaccination by intranasal aerosolization with a $10^{7.0}$ TCID₅₀/mL nebulized Rhinitis A dose over a four minute period. Horses were evaluated for clinical signs of signs of temperature, nasal exudate, ocular discharge daily. Blood for serum neutralization (SN) was taken weekly.

Challenge Virus

[0150] The challenge virus was produced in tissue culture on E-Vero cells. The titer of the challenge virus was determined to be $1 \times 10^{6.9}$ TCID₅₀/mL on the day of challenge. Challenge virus was diluted on the morning of challenge 1: 10 with tissue culture media to effect a titer of $1 \times 10^{5.9}$ TCID₅₀/mL.

Intranasal Challenge Method

[0151] Sedivet® (romifidine hydrochloride), a sedative and analgesic, was administered intravenously to each horse prior to challenge at a dosage of 50 µg/kg of body weight. Each horse was then challenged with approximately $10^{6.2}$ TCID₅₀ of equine rhinitis A virus. The challenge virus was administered intranasally as an aerosol produced by a nebulizer into an Equine AeroMask (Trudell Medical International, Ontario, Canada) by the following method.

[0152] Four milliliters of $10^{5.9}$ TCID₅₀/mL challenge virus were placed into the nebulizer cup in the AeroMask device. A pressure hose was fitted from an air compressor to the inlet port of the nebulizer. The outlet tube was then inserted into the Aero Mask attached to the head of the horse being challenged and approximately 10 psi of air pressure was applied to the inlet port for three minutes. During this time approximately two milliliters of challenge virus fluid was aerosolized directly into the nostrils of the horse being challenged.

Serum Neutralization

[0153] A standard microtiter serum neutralization test was employed in this study. A standard microtiter serum neutralization test was employed in this study. All sera were tested in sterile flat bottom microtiter plates using five wells per dilution and an 8 well dilution series for each of the 5 test wells. Each of the 5 test wells contained 25 μ l of serum dilution mixed with 25 μ l of the indicator virus and 150 μ l of a freshly planted eVero cell suspension containing approximately 5×10^4 cells. The test indicator virus used was either Equine Rhinitis Virus Type I. Serum neutralizing antibody titers are expressed as Reed-Muench ID₅₀ titers.

[0154] For performance of the test, two-fold dilutions of each test serum was made in a sterile flat bottom microtiter plate using five replicate wells per test serum and an 8 well dilution series. Dilutions were made with an adjustable volume single or multi-channel pipeting instrument using sterile microtiter tips. The volume of serum added each of 5 wells of the first row was 50 μ l. All other wells contained 25 μ l of DMEM (no FBS). Following serial dilution down the plate, 25 ml was discarded from the last row. 25 μ l of a pre-determined dilution of the indicator virus was added to each test well. Plates were then mixed and incubated for one hour at 37°C in 5% CO₂. On conclusion of the incubation period, 150 μ l of a suspension containing 4×10^6 /mL eVero cells was added to each test and cell control well. The plates were incubated at 37° C in a CO₂ incubator for 5 days, at which time plates were microscopically examined for CPE typical of equine rhinitis virus.

Nasal Exudate Evaluation

[0155] All nasal exudate observations were made prior to collection of nasopharyngeal swabs. On the Day of Challenge and for 10 days post-challenge, the nasal passages and muzzle of each of the 44 vaccinated and control horses were examined and graded using the grading and scoring description listed below.

[0156] The scoring grades of 0 through 6 were assigned on the basis of the severity of the disease indicated by each of the following classification:

Table 3: Scoring Grades

Score	Description of symptoms
0	Essentially normal indicates the horse was clean and essentially free of nasal exudate
1	Slight clear serous discharge that may be frequently observed in both diseased and normal horses
1.5	Very slight mucopurulent discharge indicates that mucus was definitely present in small amounts in either one or both nostrils
2	Moderate clear serous discharge is indicative of a definite increase in volume over that normally observed
2	Slightly mucopurulent is a discharge easily observed in one or both nostrils
3	Copious clear serous discharge that is generally observed only in diseased horses
4	Moderately mucopurulent indicates that mucoid discharges were present in large quantities in both nostrils
6	Heavy mucopurulent indicates that copious amounts of a mucoid discharge filled both nostrils

Ocular Evaluation

[0157] Ocular discharge was evaluated daily at the time of nasal exudate evaluation. Ocular discharge scores were recorded as 0=normal; 1=mild to moderate ocular discharge, and 2=severe ocular discharge.

Nasopharyngeal Viral Isolation

[0158] On each observation test day each nasal passage of each vaccinated and control horse was swabbed deeply by means of a sterile WECK-CEL™ surgical spear (Edward Weck and Company, Inc., Research Triangle Park, N.C. 27709) attached to an 11-inch long sterile plastic pipette. On collection, each of two surgical spears was immediately placed in a single tube containing 4 mL of chilled transport medium (E-199 supplemented with gentamicin, L-glutamine, 2X Pen/Strep, 2X Amphotericin B).

[0159] For isolation of virus, the tubes were mixed, the swabs aseptically removed, and the medium centrifuged at 1500 rpm for 10 to 15 minutes to remove particulates. Medium was filtered through a 0.2 µL syringe filter prior to inoculation on tissue culture cells. After filtration, 4-6% of sterile 85% sucrose solution was added to each sample for freezing at -80°C in order for all samples to be tested concurrently.

[0160] For isolation of virus, the tubes were mixed, the swabs aseptically removed, and the medium centrifuged at 1500 rpm for 10 minutes to remove particulates. Medium was filtered through a 0.2 μ L syringe filter prior to inoculation on tissue culture cells. One mL of the clarified transport medium was used to inoculate a 2 cm² two day old monolayer of E-Vero cells grown in a 24 well tissue culture plate from which the growth medium had been aseptically removed. Following inoculation, the inoculum was allowed to adsorb on the cell monolayer for one hour at 37°C in a humidified incubator containing a 5% CO₂ atmosphere. After the adsorption period, an additional 1 mL of re-feed medium (E-199 containing 7% fetal bovine serum (FBS), 2mM L-glutamine, Gentamicin 2X Pen-Strep and 2X Amphotericin B) was added to each well. Following addition of re-feed media the plates were then incubated at 37°C in a CO₂ incubator. Each test and control tissue culture well was examined microscopically for 7 days for signs of cytopathic effect (CPE) typical of the ERAV challenge virus. Wells that were negative at the end of the 7 day observation period were subcultured onto fresh cells and observed for an additional 7 days.

STATISTICAL EVALUATION METHODS

[0161] Data from all horses vaccinated with either A-9 or A-10 were combined for statistical evaluation. The influence of vaccination on the duration of disease (number of days with nasal scores > 0) was evaluated using the Kruskal-Wallis and Hodges-Lehmann test (the NPAR1WAY procedure in SAS, SAS Institute, Cary NC). Severity of disease was evaluated by comparing the maximum disease status between the vaccinated horses and the control horses. Nasal scores were dichotomized to ≤ 1.5 and > 1.5 based on the distribution of outcomes. The prevented fraction (PF) and 95% confidence intervals (CI) were estimated. Ocular scores were evaluated as present or absent and the prevented fraction and 95% confidence intervals were estimated (the FREQ procedure in SAS). Repeated measures analysis appropriate for continuous data was used to assess the effect of vaccination on body temperature and serum neutralization titers (the MIXED procedure in SAS). The proportion of horses which were virus positive over time and buffy coat positive over time was evaluated using Fisher's exact test at each time point (the Freq procedure). Nasal and ocular scores were assessed by Wilcoxon's rank sum test at each time point (the NPAR1WAY procedure).

Results and Conclusions

VIRUS ISOLATION FROM NASAL SWABS (Virus Shedding) and BUFFY COATS (Viremia)

[0162] The proportion of animals which were virus positive was significantly lower in the vaccinated groups on days 2 through 7 (Table 4 below and Figure 1) as compared to control group ($P < 0.05$).

Table 4. Proportion of nasal swab virus positive over time (* $P < 0.05$ versus control, Fisher's Exact test on each day)

Day	Control	A-9	A-10	A-9 + A-10
0	0	0	0	0
1	25	7.1	7.7	7.4
2	100	20*	23.1*	21.4*
3	100	46.7*	57.1*	51.7*
4	100	60*	57.1*	58.6*
5	93.3	53.3*	35.7*	44.8*
6	60	40	0*	20.7*
7	100	46.7*	78.6	62.1*
8	26.7	13.3	0	13.8
9	0	6.7	0	3.5
10	0	6.7	0	3.5
11	0	0	0	0
12	0	0	0	0
13	0	0	0	0
14	0	0	0	0

[0163] Buffy coat positive animals were less frequent in the vaccinated group on days 4-7 as compared to the control group (Table 5 below and Figure 2).

Table 5: Proportion of Buffy Coat positive over time (* $P < 0.05$ versus control, Fisher's Exact test on each day)

Day	Control	A-9	A-10	A-9 + A-10
0	0	0	0	0
1	0	0	0	0
2	0	0	0	0
3	6.7	0	7.1	3.5

4	73.3	0*	7.1*	3.5*
5	80	13.3*	0*	6.9*
6	80	0*	7.1*	3.5*
7	33.3	0*	0*	0*
8	0	0	7.1	3.5
9	0	0	0	0
10	0	13.3	0	0
11	0	0	0	0
12	0	0	0	0
13	0	0	0	0
14	0	0	7.1	3.5

Serum Neutralization

Table 6: Summary of body temperature and serum neutralization

Variable	P-Values		
	Vaccine	Day	Vaccine * Day
Temperature	0.9269	< 0.0001	0.0255
SN titer (ln transformed)	< 0.0001	< 0.0001	< 0.0001

[0164] Mean body temperature values (rectal temperature measured with calibrated GSA Electronics thermometer probe) on each challenge day were always within normal body temperature parameters. Large increases in SN titers resulting from vaccination were found (see also Fig. 3) following challenge.

Nasal And Ocular Evaluation

[0165] Mean ranks for nasal scores were lower on days 4 though 10 and days 12-13 in the vaccinated group compared to the control group (Table 7 below and Figure 4). The mean ranks for ocular scores were lower on day 7 in the vaccinated group as compared to the control group (see also Figure 5).

Table 7: Mean rank for nasal and ocular score across time (*P<0.05 versus control; Wilcoxon's rank sum test on each day, A-9 and A-10 combined).

Day	Ocular Score		Nasal Score	
	Control	A-9 + A-10	Control	A-9 + A-10
0	22.5	22.5	22.5	22.5
1	22.5	22.5	22.5	22.5
2	22.5	22.5	23.9	21.8
3	22.5	22.5	26.2	20.6
4	22.5	22.5	27.7	19.8*
5	21.8	23.9	20.2	19.5*

6	22.5	22.5	28.5	19.4*
7	29.2	19.0*	29.2	19.0*
8	21.8	23.9	30.1	18.5*
9	21.5	24.4	27.4	19.9*
10	22.5	22.5	28.1	19.6*
11	22.5	22.5	37.2	20.1
12	21.5	24.4	30.7	18.3*
13	22.0	23.5	28.8	19.2*
14	22.5	22.5	25.6	20.9

Disease Duration: Nasal Exudate Evaluation

Table 8: Summary of the effect of vaccination on the duration of disease (Number of days with nasal score > 0)

Group	Minimum	25 th quantile	50 th quantile	75 th quantile	Maximum
Control	8	9	10	11	13
Vaccinated	0	1	3	9	13

Table 9: Effect of vaccination on the duration of disease (nasal scores; *significantly lower than control group by Kruskal-Wallis test $P < 0.05$)

	Control	Vaccinate	Shift in days	95% confidence interval
Duration	10	3	7*	3, 8 days

[0166] Long duration of nasal discharge in controls was found following challenge (Table 8, see also Figure 4). Vaccinated group showed a significant reduction in duration of nasal discharge. Controls experienced a minimum of 8 days of nasal discharge vs 11 vaccinates (38%) with 1 day or less. Two thirds of vaccinates had shorter duration of nasal discharge than the minimum of 8 days in the control group. Duration was from first to last abnormal observation, even if horse had normal days in between. The number of days animals were sick with clinical signs of respiratory disease (nasal score > 0) was significantly shorter (shift of 7 days) in the vaccinated group compared to the control group.

Table 10: Maximum Nasal Scores -Significant difference between the two distributions of scores (Kruskal-Wallis test, $P = 0.0157$).

Group	Maximum Nasal Score
-------	---------------------

	0	1	1.5	2	3	4
Control	0	0	5 (33%)	9 (60%)	0	1 (7%)
Vaccinated	4 (14%)	2 (7%)	13 (45%)	10 (34%)	0	0

Table 11: Nasal scores (≤ 1.5 and > 1.5)

Group	% with score	P-value	Prevented Fraction	95% confidence interval
Control	66.7%	< 0.0588	0.48	0.042
Vaccinate	34.5%			0.72

[0167] Vaccination also reduced severity of nasal discharge (maximum score on any day post-challenge). Maximum nasal scores were compared between groups (Table 10). The minimum nasal score for horses in the control group was 1.5. Results were thus dichotomized to scores ≤ 1.5 and > 1.5 for the evaluation of disease severity (Table 11). The prevented fraction was 48% with a lower confidence limit greater than 0. The overall distribution of maximum nasal scores was significantly reduced by vaccination (Kruskal-Wallis test, $P=0.0157$).

Ocular Evaluation

Table 12: Ocular scores (present or absent)

Group	% with score	P-value	Prevented Fraction	95% confidence interval
Control	66.7%	< 0.0001	0.897	0.587
Vaccinate	6.9%			0.9741

[0168] Since only two values were reported for ocular scores (0 or 2), results were dichotomized to present or absent within an individual. The prevented fraction was then used to evaluate the effect of vaccination on the presence of ocular signs. The vaccine significantly reduced the severity of ocular discharge resulting from infection with ERAV. Mean ocular scores over time are also shown in Figure 5.

[0169] Data from this study demonstrate that administration of intramuscular doses of the inactivated virus is capable of immunizing an animal to high levels of antibody detection which

prevent ocular disease and reduce severity and duration of nasal discharge. Vaccine was highly effective in preventing any ocular discharge following ERAV challenge. Nasal discharge persisted for a relatively long period post-challenge, while ocular discharge peaked and abated quickly. Hence the vaccine showed a significant improvement in nasal discharge for a prolonged period (9 of 10 consecutive days), whereas vaccine significantly reduced ocular discharge for 1 day during which ocular signs were most severe. In addition, vaccination significantly reduced nasal virus shedding throughout the successive six days of peak viral shedding, and viremia also was significantly reduced during the four day of peak viremia within the challenge period.

[0170] No unacceptable adverse reactions either at the injection sites or by manifestation of signs of systemic illness were observed. The vaccine is safe and well tolerated for administration in Equine rhinitis A susceptible species, particularly equidae. This study thus demonstrates that 3 x 1 mL intramuscular injections of the ERAV compositions of the example significantly reduced severity and duration of respiratory disease caused by ERAV virulent challenge.

Example 3

[0171] This example illustrates sequencing and analytical studies carried out on an Equine Rhinitis A Virus strain as disclosed herein.

Cells And Virus

[0172] Rabbit-kidney-13 (RK-13) cells (passages 100-160) were grown in Dulbbeco's modified eagles medium nutrient mixture F12 HAM (DMEM F12) (Sigma-Aldrich Canada Ltd. Oakville, Ontario) with 2-5% fetal bovine serum (FBS) (Sigma). Cell growth and viral propagation were performed in a CO₂ (5%) incubator at 37 °C. The ERAV isolate ERAV/ON/05 was propagated in RK-13 cells and aliquots were stored at -70 °C for later work. RK-13 monolayers were inoculated at 90% confluence and RNA was extracted before cytopathic effects (CPE) was observed.

Virus Titration And Plaque Purification

[0173] RK-13 cells were grown on 3 cm round dishes, using DMEM F12 with 2% fetal calf serum. Cells were infected at 90% confluence and plates were incubated at 37 °C for 72 hours. After 24 hours, the medium was replaced and a 0.7 % agarose layer was added. Plaques

were counted and recorded every 24 hours. At 72 hours the agarose layer was removed and plaques were stained with crystal violet and counted.

[0174] For plaque purification RK-13 cells were infected with ERAV/ON/05, adsorption was allowed for 45 minutes and the inoculum was removed and replaced with a 0.7 % agarose layer. Plates were checked every 12 hours and plaques were classified as small, medium and large. Five plaques of each size were selected, picked into 300 μ L of DMEM F12, and frozen at -70 °C. Viruses from each plaque size were propagated in RK-13 cells and RNA was extracted from the small and large plaques for genome sequencing of the 5' UTR.

Viral Growth Kinetics

[0175] To study the growth characteristics of this strain, RK-13 cells were infected with ERAV/ON/05 and supernatant samples were collected at various times for titration. RK-13 cells were grown on 3 cm individual round plates and infected at 90% confluence. Plates were incubated at 37 °C and supernatant samples were taken every 4 hours starting at 0 hours for a period of 28 hours. All samples were titrated using the plaque forming unit (PFU) technique as previously described here.

Immunofluorescence

[0176] RK-13 cells were grown on glass tissue culture chamber/slides (Miles Scientific, Inc., Naperville, Illinois), and inoculated with ERAV/ON/05. Twenty eight hours post-infection, the cell culture medium was removed and cells were fixed in acetone. The slides were kept at 4 °C until processing. Sera from experimentally infected horses were used as a source of ERAV antibodies.

Rna Extraction And Sequencing

[0177] RNA was extracted from infected cell monolayers. Cells were treated with 1 mL of TRIzol (Invitrogen) 18-20 hours post-infection and extraction was performed according to manufacture's recommendations. RNA pellets were eluted in 30 μ L of RNase free water and kept at -70 °C for later use. First strand cDNA was synthesized using superscript II (Invitrogen) following manufacturer's recommendations. A 50 μ L PCR reaction was carried out using a set of sense and antisense primers. For genome sequencing, the primer walking approach was used,

and primers were designed based on eight ERAV sequences available on GenBank. PCR conditions were: 4 minutes at 94 °C, followed by 30 cycles of 30 seconds at 94 °C, 30 seconds at 55 °C and 30 seconds at 72 °C with a final extension at 72 °C for 10 minutes.

[0178] Sequencing of the 5' and 3' ends were completed with the 5' RACE and 3' RACE kits (Invitrogen) as recommended by the manufacturer. Several nested PCRs were required to amplify the 5' UTR end.

Sequence Analysis

[0179] The preliminary identification of the virus was completed by partial sequencing of the structural protein VP1 using primers designed based on other sequences available on GenBank.

Table 13: Primers used to amplify some of the ERAV regions.

NAME	PRIMER SEQUENCE	SEQ ID NO:	GENOMIC SITE
forwVP1last	5' tgaatagcaagggccgtgtt 3'	4	3087
revVP1last	5' accgttgtaaaagactggcaca 3'	5	3671
forwPC112	5' gtcagtaaaacgcaacaacat 3'	6	112
forwPC805	5' tgtgaagaatgtcctgaaggca 3'	7	805
revPC1749	5' accatccacctaaccagacga 3'	8	1749
forwPC5217	5' attggctttgtcaggtgttgaa 3'	9	5217
revPC5952	5' gtttctaactttgggaccgaa 3'	10	5952
forwPC6915	5' tggatttgagattggttctgca 3'	11	6915
revPC7511	5' gcgaacgaaactgaggattg 3'	12	7511

[0180] All primers were designed on Gene Runner version 3.05 (Hastings Software Inc.). Sequencing reactions were set and run by the Laboratory Services Division at the University of Guelph.

[0181] All sequences were assembled and edited with EditSeq and SeqMan DNASTAR Lasergene 8 (DNASTAR Inc., Madison, WI, USA). Sequencing results were entered into the BLAST software [National Center for Biotechnology Information, Bethesda, MD (NCBI)] and compared to similar entries on GenBank. ClustalW2 [European Bioinformatics institute, Dublin, Ireland (EBI)] was used for multiple sequence alignment and preliminary construction of the phylogenetic tree. The final phylogenetic tree was created on MEGA 4.0 by using the Maximum Composite Likelihood method and the reliability was evaluated by bootstrapping with 1000

replications. Analysis of the nucleotide sequences were plotted on SimPlot Version 3.5.1 (Baltimore, MD, USA). In order to investigate the possibility of viral recombination between ERAV isolates, we completed a Bootscan analysis on SimPlot (Version 3.5.1) comparing the genomic sequences of all reported ERAV available on GenBank. Polyprotein cleavage sites were predicted based on sequences reported on GenBank with accession numbers: DQ272578 and NC003982.

Results

Initial Characterization

[0182] The viral protein (VP1) of the virus was amplified, partially sequenced and compared to sequences available on GenBank (accession numbers: NC003982, DQ272577, DQ272128, DQ272127, DQ268580, DQ272578, L43052). Results from this initial comparison demonstrated a maximum identity of 95% to Equine rhinitis A virus VP1.

Virus Titration

[0183] The ERAV isolate was initially propagated in cell culture and titrated by the plaque forming unit method. The stock virus titre obtained was 5×10^7 PFU on the initial titration. Subsequent experiments showed no dramatic changes in viral titre after 3 years of initial freezing. For animal experiments the stock virus with the initial titre was employed.

GROWTH KINETICS

[0184] All supernatant samples were titrated using the PFU technique and results were entered on a spread sheet to construct a growth curve. As seen in other picornaviruses, the Ontario isolate showed an increased titre at 4 hours post-infection, reaching a plateau by 12 hours post-infection.

Immunofluorescence

[0185] Infected RK-13 cells were visualised under the immunofluorescence microscope. Bright green fluorescence signals were detected in the cytoplasm of ERAV infected cells compared to negative signals on mock infected cells.

Plaque Purification

[0186] Sequencing of a 426 nucleotide fragment in the 5' UTR on the small and large plaques showed no difference between plaque sizes at the nucleotide level. However, morphological characteristics were evidently different on RK-13 cell cultures.

Full Genome Sequencing

[0187] Genome sequencing of the ERAV isolate resulted in 7839 nucleotides in length with a GC content of 47% including the poly (A) tail. Four identical repeats (CTGTAGCGTCAGTAAAACGC SEQ ID NO: 13) separated by 18, 21 and 18 nucleotides were identified on the 5' UTR. The ERAV 5' UTR was composed of 940 nucleotide with a 54% GC content. A single polyprotein with 6747 nucleotides (2248 amino acids) composes the viral genome with a 46.8% GC content. The protein starting synthesis was detected on nucleotide 940 with the AUG start codon and the ending on nucleotide 7686 with the UAA stop codon. Of great interest, a second AUG sequence was identified following the initiation codon. A subsequent AUGAUG sequence was also identified 58-nucleotides downstream from the polyprotein start codon. Analysis with Blastx (NCBI) of the polyprotein on this isolate showed a 96% nucleotide identity with respect to other ERAV reported, however when the full-length genome sequence of the Ontario isolate was aligned and compared to others using ClustalW2 (EBI) and Blastx (NCBI), only a maximum identity of 80 % was observed. The amino acid composition showed an identical protein (structural and non structural) organization and length along the entire genome. These comparisons were made with PERV-1 and PERV reported genome sequences (accession numbers DQ272578, and NC003982).

[0188] The 3' UTR was composed of 110-nucleotides with a 24.3 % GC content, and a poly-A tail. Alignments of the 5' UTR of all ERAV available demonstrated a lower identity percentage, ranging from 73% to 81%. Similarly, the 3' UTR was analysed by the same method and the identity percentage ranged from 75% to 81%. Interestingly, various insertions (1-nucleotide, 2-nucleotides, and 13-nucleotides) and two small deletions (2-nucleotides, and 3-nucleotides) in the 5' UTR were identified (Fig. 4). No other major changes were observed throughout the entire genome.

Simplot Genome Analysis

[0189] SimPlot analysis showed a comparable similarity (percentage) among all ERAV isolates when compared to ERAV/ON/05. However, the mayor disparities were identified at nucleotide 1500 (59% similarity) and nucleotide 4700 (65% similarity). Nevertheless, this analysis showed that the similarity along the genome was between 70% and 82%. To further investigate the major divergence in the genomes, a scan (Bootskan) to identify possible recombination sites was performed. These analyses demonstrated that the Ontario isolate has not had predicted recombination sites with other ERAV isolates. Nevertheless, when the Ontario isolate was removed from the analysis, possible recombination between other ERAV isolates may have happened in the pass.

Discussion

[0190] ERAV is not routinely sought and recovered from clinical cases during equine respiratory outbreaks. Therefore ERAV isolation from clinical cases has been incidental and in most cases a challenge. For these reasons the recovery and sequencing rate of these viruses may be small.

[0191] During previous years, equine rhinitis viruses were classified within the *Picornaviridae* family, but were not clearly assigned to a specific genus. In 1996 Li and coworkers (Li F, Browning GF, Studdert MJ, and Crabb BS. Equine rhinovirus 1 is more closely related to foot-and-mouth disease virus than to other picornaviruses. Proc Natl Acad Sci U S A. 1996 6; 93:990-995) demonstrated that ERAV was closely related to foot and mouth disease virus (FMDV) based on the phylogenetic characteristics and nucleotide sequencing. In agreement with their findings and others (Wutz G, Auer H, Nowotny N, Grosse B, Skern T and Kuechler E. Equine rhinovirus serotypes 1 and 2: relationship to each other and to aphthoviruses and cardioviruses. J Gen Virol 1996, 77:1719-1730) the ERAV isolate has been found to be closely related to isolates L43052, DQ272578, and NC003982.

[0192] The ERAV/ON/05 isolate resulted in 7839 nucleotides including the 5' UTR, the polyprotein, 3' UTR and the poly-A tail. In comparison to previously reported ERAV isolates, the ERAV/ON/05 is one of the most complete ERAV sequences reported to date. Most of the other sequences lack some information either from the 5' UTR or are partial sequences of

individual viral proteins. Data from the 5' UTR revealed the presence of 3 repeats that have been previously reported in other ERAV and commonly found on the FMDV. It has been suggested that these repeats may be required in the formation of the secondary structures found in the 5' UTR, the internal ribosome entry site (IRES). The identity analysis showed that the ERAV/ON/05 isolate's polyprotein bears a highly conserved nucleotide sequence. As an RNA virus, the ERAV is prone to constant mutations due to the lack of proofreading by the polymerase. This may indicate, that even though this RNA virus has been under natural evolution and constant replication no significant genomic changes have been introduced since its first genomic sequence was reported. It is evident that the structural and non structural proteins, which are encoded within the polyprotein, represent the most conserved regions in the genome.

[0193] Interestingly, the viral replication rate in cell culture was rapid and efficient. A complete viral replication cycle was detected in less than 4 hours, reaching a plateau in 12 hours. These characteristics are typical of picornaviruses and may reflect the viral activity *in vivo*. Such features might explain the severity and speed of clinical signs in some equine respiratory viral infections. As observed in other viruses, such as poliovirus and FMDV, the presence of small deletions and/or insertions in the 5'UTR have been associated with differences in plaque size in cell culture, and excel virulence *in vivo*. We found that the ERAV/ON/05 isolate generates different plaque sizes when infecting RK-13 cells. To investigate and correlate the presence of these insertions and deletions in the 5' UTR in the ERAV/ON/05 isolate, we sequenced this region from plaque purified viruses from different plaque sizes, and found no discrepancy at the nucleotide level. This finding may indicate that the plaque size difference may be due to a growth and replication characteristic of the RK-13 cells and perhaps not associated with the presence or absence of the sequences we found in this region.

Example 4

[0194] This example shows results of a study designed to develop a reliable ERAV infection model for investigating the clinical characteristics of ERAV/ON/05.

Materials and Methods

[0195] On the first phase a pony foal was infected with ERAV/ON/05 to adjust the infectious dose and sampling collection techniques. A second pilot study was designed to

compare infected and control animals, and to master sampling techniques. The main infection study was conducted in two phases due to animal handling and time sampling constraints. One year after the first infection, four previously infected ponies were selected for a re-infection study based on their titre to ERAV. Animals with an intermediate and a high titre to ERAV were selected to be re-exposed to the Ontario isolate.

Experimental Animals

[0196] A total of 12 pregnant pony mares were selected and their foals were kept for the experimental ERAV infection. All mares and foals were kept in a separate group away from the main barn and biosecurity measurements were put into place to prevent viral respiratory infection. Blood samples from all foals were taken regularly and titers to ERAV were checked periodically. Foals were kept until they reached 12 months of age. These animals were not vaccinated against any respiratory viruses during this period and general practices were maintained to ensure healthy living conditions. All animals were de-wormed according to the herd management protocol. Animal socialization and handling was performed regularly. Ponies were trained for pulmonary function test (PFT). Once the animals were conditioned for the experiments, they were transported in groups to the isolation unit and acclimatized for at least one week prior to the viral infection.

Isolation unit

[0197] All infection trials were conducted at the animal isolation unit in the Department of Pathobiology, at the University of Guelph. This isolation unit contains individual stalls that are equipped with controlled temperature, humidity, airflow, and lighting. Access to the stalls was restricted to the researchers and care personnel.

Selection criteria

[0198] All foals were handled frequently, and health status was checked periodically. Based on health status and serological parameters, animals to be included in these experiments were chosen. At that time, all ponies remained seronegative to equine rhinitis A virus (ERAV), equine rhinitis B virus (ERBV), equine herpesvirus 1 and 4 (EHV1/4), and equine influenza 2 virus (AE2). Even though AE2 antibody titers were detected at birth, a steady decrease was

observed during the first 5 months and were not detectable by the single radial haemolysis test (SRH) at six months of age. For the ERAV experimental infection trial, one pony was selected to be used. Subsequently, two other ponies were selected for a second pilot study, and a group of 8 ponies (four infected and four controls) were selected for the main infection study. After a year from the first infection trial, 4 ponies with intermediate and high titers to ERAV were chosen for a re-infection trial.

Inoculum

[0199] ERAV isolate (ERAV/ON/05) recovered from a horse during an equine viral respiratory outbreak in Ontario 2005 was propagated in rabbit kidney-13 cells (RK-13) and aliquots were stored at -70 °C for viral characterization and animal viral infection experiments.

[0200] Briefly, for inoculum preparation, the isolate was propagated in RK-13 cells. 30 mL round dishes containing 90% confluent monolayers were infected with 500 µL of ERAV/ON/05 and incubated in the presence of CO₂ (5%) at 37 °C for 24–36 hours. All dishes were removed from the incubator and freeze/thawed four times to provoke cell rupture and viral release from intact cells. Supernatants from all dishes were pooled into one flask and centrifuged at 6000 RPMS for 15 minutes to clarify and discard cell debris. The clarified supernatant was aliquotted in 10 mL vials to be used as an inoculum in the animal viral infection experiments. Additionally, small 1 mL-aliquots were separated and kept for viral titration. The virus was titrated using the plaque forming unit method (PFU).

Animal model

[0201] Ponies were chosen as an animal model due to the nature of the experimental agent (ERAV), animal size, availability and handling.

Infection protocol

[0202] Ponies between 8 and 12 months of age were selected, trained and used in the infection experiments. Due to the large number of samples to be taken, these experiments were divided in five sections (two pilot studies, two main infection experiments, and one re-infection experiment). During the pilot studies, equipment, animal handling, inoculum dose, route of administration, and sample collection were optimized. The re-infection study was conducted one

year after the initial infection trial. Ponies in the re-infected group were exposed to the same viral strain at the same dose used during the first infection. Only clinical examination, blood sampling for titers assessment and nasopharyngeal swabs for virus isolation were collected from this group.

Face mask and nebulization

[0203] For viral infection, a small size Equine AeroMask™ was fitted with a rubber seal on the nostril ending. Size was adjusted depending on the pony's head and the breathing windows were not modified. The mask was fitted with an inhaler connector and a one way "T" valve for virus nebulization. Conventional 6 mL nebulizer cups were employed to deliver the inoculum. Nebulization was performed using a PM14 compressor (Precision Medical Inc. Northampton, PA) with a gas flow of 9 LPM (liters per minute). This flow rate and delivery cups allow for consistent delivery of breathable particles of more or less 5 microns. Each pony was nebulized for 45 min (taking a 5 minute break every 15 minutes) with 15 mL (total volume) of either inoculum or placebo. A nasopharyngeal swab per pony was collected post-infection to ensure viability of the inoculum when delivered. Ponies that were re-infected a year later were exposed to the virus by the same protocol.

Clinical examination

[0204] All ponies were clinically evaluated on a regular basis from birth. Prior to the pilot and infection studies, all ponies were evaluated daily and during trial experiments twice daily for the first 10 days and once daily from day 11 up to day 21 post-infection. Clinical examination included: Body temperature (temp) (Celsius degrees), heart rate (hr), respiratory rate (rr), capillary refill (ref), gastrointestinal motility (gi), lung sounds (ls), nasal discharge (nd), ocular discharge (od) [presence or absence and characteristics], lymph nodes (ln) [size and characteristics], ambulation, and general physical condition. Physical examination was performed at around the same times on each pony every day, commencing with the control animals and moving onto the infected group. Approximately 10 minutes were spent on each animal examination every time and all individual data were recorded on a daily check up form.

Pulmonary function test (PFT)

[0205] The PFT was carried out as previously described by Hare and coworkers (Hare JE, Viel L. Pulmonary eosinophilia associated with increased airway responsiveness in young racing horses. *J Vet Intern Med* 1998;12(3):163-70). The test was performed on all control and infected ponies prior to infection (day 0) and on days 1, 7, 14, and 21 post-infection. Briefly, on testing day, ponies were off feed in the morning and were mildly sedated as previously described here for endoscopy procedures. A rubber face mask was fashioned to snugly fit the ponies' muzzle. A pneumotachograp # 4 (Gould Electronics, Biltholven, The Netherlands) was attached to the face mask, and connected to a set of transducers that convert the flow and pressure signals into breath loops that are recorded on a computer (Pulmonary Mechanics Analyzer, Buxco Electronics Inc, Sharon, CT, USA). Flow was measure at the pneumotachograp level and the pleural pressure was assessed by an esophageal balloon (10cm long) that was placed in the mid thorax via esophageal tubing. A total volume of 3ml of air were introduced into the balloon. The pressure difference between the pleural pressure and the atmospheric pressure (measured at the nostril level) was considered as the transpulmonary pressure (ΔP_{pl}).

Bronchoprovocation testing

[0206] Bronchoprovocation challenge was carried out as part of the PFT testing. To determine the hyperresponsiveness of the airways post ERAV/ON/05 infection, control and infected animals were exposed to increasing histamine doses (doubling dose) by nebulization with a PM14 compressor with a gas flow of 9 LPM (Precision Medical Inc. Northampton, PA). Pulmonary physiological parameters were assessed initially by administering 0.9% physiological saline solution (base line), followed by increased histamine doses. After each administration (2 minutes) data was recorded for 3 minutes and respiratory physiology was later analyzed. Histamine doses were started at 0.5 mg/ml and were increased gradually to a maximum of 32 mg/ml. Dynamic compliance (C_{dyn}) and ΔP_{pl} were used as parameters to discontinue histamine nebulization. When C_{dyn} was decreased by two thirds or ΔP_{pl} was doubled during histamine administration, nebulization was suspended. Histamine triggering dose was later plotted and calculated. All ponies were mildly sedated and physically restrained during these procedures.

Sampling techniques

Blood collection

[0207] Blood samples were collected from either the right or left jugular vein. Approximately 10 ml of blood were collected on a red top tube (serum) from each pony according to the sample collection schedule. Additionally, 3 to 5 ml of blood were also collected for CBC (complete blood count) and profile. All samples were collected in the morning hours and processed within the same day.

[0208] Blood samples for serum separation were kept at room temperature for at least 30 minutes and then centrifuged at 3000 rpm on a table centrifuge. Serum was separated within 6 hours from the collection time and aliquots were labelled and frozen at -70° C for later analysis (antibodies to ERAV, ERBV, AE2, and EHV1/4). Blood samples for CBC and profile were processed within the same day.

Nasopharyngeal swabbing

[0209] Nasopharyngeal swabs were collected for virus isolation on days previous to the infection trial and on days 0, 1, 3, 5, 7, 10, 12, 14, 17 and 21. Each pony was restrained with a muzzle twitch and a 30 cm long swab (Kalayjian Industries, Inc. Signal Hill, California) was passed into either the right or left nostril until it reached the pharynx. Swabbing was performed by rotating the swab for about 5 to 10 seconds. The swab was removed carefully, and the tips were cut off into a vial containing 3ml of virus transport medium (VTM). The vials containing the swabs were shaken and kept on ice until processing. To release the viral particles and cells attached to the swabs, vials were vortexed for about 20 seconds and the medium was transfer to a 1.5 mL Eppendorf tube and frozen at -70 °C for later analysis.

Urine and fecal sampling

[0210] Virus isolation was attempted on urine and fecal samples pre and post-infection with ERAV/ON/05. Urine was collected in the morning after clinical examination and/or stall cleaning by holding a collection cup while the animals were urinating. When the sample could not be collected by hand, the animal was fitted with a plastic collection bag around the genitals. This bag was removed after the animal urinated and a 10 mL aliquot was saved for virus

isolation. Fecal samples were hand collected from fresh manure on the stall's floor. About 5 mL of manure were collected into a collection cup and 10 mL of sterile saline were added to dissolve the sample. Urine and fecal samples were kept on ice until processed.

Upper and lower endoscopy

[0211] Bronchoscopy and BAL were performed as previously described (Hare et al., 1998). A sterile flexible fiberoptic endoscope, 140 cm length with a 0.8 mm OD (Olympus, Corp., Tokyo, Japan) was advance through the right or left nostril into the nasal cavity to the larynx level. At this point the upper airways conformation was evaluated. Followed, the bronchoscope was advance into the trachea and the presence or absence of inflammation and/or mucus and its characteristics were recorded. Data were recorded on the daily evaluation sheet and all the endoscopies were video recorded for later analysis.

Pharyngeal and tracheal brush biopsies

[0212] In order to assess viral replication in the upper and lower airways, a brush biopsy was taken from the pharynx, mid trachea, and the carina of infected and control animals on days 0, 1, 3, 5, 7, 10, 14, and 21. During endoscopy examination, a 200 cm guarded (protective sleeve) cytology brush (Hobbs Medical Inc. Connecticut) was advance through the biopsy channel at the predetermined sample location and a sample was collected. Brushes were retracted into the protective sleeve and removed from the bronchoscope. To remove the sampled tissue from the brush, this was put into 600 μ L of VTM and vortexed for 10 to 20 seconds. All samples were kept on ice and transported to the laboratory for further analysis (virus isolation).

Bronchoalveolar lavage (BAL)

[0213] Briefly, all ponies were mildly sedated with Romifidine (0.04 mg/kg, IV) and a sterile 0.8 mm OD bronchoscope (Olympus, Corp., Tokyo, Japan) was advance through the right or left nostril into the trachea to the carina level. As the bronchoscope was advanced, a 0.2% warmed lidocaine solution was administered to reduce cough and distress. Once the cough reflex was reduced the bronchoscope was advanced and wedged into a proximal terminal bronchus. A total of 250 mL of warmed sterile saline solution was administered through the biopsy channel divided in two aliquots. BAL fluid was retrieved by manual suction with a 60 cc

syringe through the biopsy channel and placed on ice. The fluid was filtered and aliquots for virus isolation, cell count, and cytospin slides were made.

Clinical samples culturing

[0214] Collected samples were transported on ice and frozen at -70 °C for later inoculation on cell cultures.

[0215] RK-13 cells (passages 100-160) were grown in Dulbbeco's modified eagles medium nutrient mixture F12 HAM (DMEM F12) (Sigma-Aldrich Canada Ltd. Oakville, Ontario) with 2-5% fetal calf serum (Sigma). Cell growth and isolation were performed in a CO₂ (5%) incubator at 37 °C. RK-13 cells monolayers were grown on 6 well plates to a 90 % confluence and infected with the clinical samples collected from the infection trial. In brief, 90% of the growing medium was removed from each of the wells and 200 µL of the specimen to be tested were added to the monolayer. Plates were put on the rocker platform for one hour and 3 mL of medium were added after the time elapsed. Plates were incubated and checked every 24 hours for cytopathogenic effects (CPE). If CPE was detected, the supernatant was removed from the well and frozen for later analysis. Results from the virus isolation test were recorded on a spread sheet. Plates were checked for up to 7 days and if CPE was not developed, a second passage was attempted using 200 µL of supernatant from the first passage. After a second passage if CPE was not detected the sample was classified as negative. Supernatant from positive and negative samples was saved to be confirmed by reverse transcriptase polymerase chain reaction RT-PCR).

Statistical analysis

[0216] Results from virus isolation and all clinical scores (Table 7) were entered on a spread sheet.

Table 14. Scoring system for experimental infection with ERAV/ON05.

Clinical sign	Degree	Score
Cough	None	0
	Intermittent	1
	Frequent	2

Mucus membranes	Pink	0
	Pale	1
Gastro intestinal auscultation	Normal	0
	Abnormal	1
Feces/Urine	Normal	0
	Abnormal	1
Lung Sounds	Normal	0
	Slightly Increased	1
	Marked Increased throughout chest	2
	Crackles and wheezes	3
Nasal Discharge	None	0
	Moderate/severe serous	1
	Mucopurulent	2
Ocular Discharge	None	0
	Serous	1
	purulent	2
Adenitis	not palpable	0
	palpable (<1cm)	1
	Enlarged (>1 cm)	2
Anorexia	None	0
	Mild to Moderate	1
	Severe	2
Temperament	Bright, alert and responsive	0
	Dull (head down, disinterested)	1
Perfusion Time	Normal (2s)	0
	3-4s	1
	5s	2

[0217] Virus isolation results were categorized as positive or negative and results were compared between groups. To determine if there was a statistical difference between locations on virus isolation (in the infected group) an exact conditional logistic regression was used at each

day and site. The same test was used to determine if there was a statistical difference between groups depending on the isolation day. Clinical scores were summarized and the totals were analysed and compared between groups.

[0218] A generalized linear mixed-model was employed to analyze all clinical parameters. Factors included in the model were: pony, treatment, and time as well as their interactions. Since animals were measured over time, the AKAIKE information criterion (AIC) was used to determine an error structure for the auto-regression. The assumptions of the ANOVA were assessed by comprehensive residual analyses. A Shaprio-Wilk test, a Kolmogorov-Smirnov test, a Cramer-von Mises test, and an Anderson-Darling test were conducted to assess overall normality. Residuals were plotted against predicted values and explanatory variables (pony, treatment and time) to look for patterns that suggested outliers, unequal variance or other problems. If residual analyses suggested a need for data transformation or data was presented as a percent, analyses were done on a logit or log scale. If the overall f test was significant, a Dunnetts test going back to baseline within a treatment or a tukey test between treatments and sites at each time was applied.

[0219] Serological response was defined as a four fold increase in antibody levels from baseline (day 0) to any of the time points in sampling collection (days 7, 14, or 21).

Statistical analysis was carried out on SAS 9.1.3 (SAS institute Inc., 2004, Cary, NC). Statistical significance was set at $P < 0.05$.

Results

[0220] This study was designed to consistently reproduce ERAV clinical disease in ponies and to study its *in vivo* characteristics. Pilot studies demonstrated that nebulized ERAV/ON/05 was able to cause clinical respiratory disease in healthy ponies (age 10-12 months). Mild immunosuppression was induced in all ponies except in the re-infected animals to mimic natural conditions under stressful events (e.g. movement for sales, field relocation, vaccination programs, etc.). Results from the main infection study confirmed the results observed during the pilot studies. Ponies in the infected group (n=4) developed respiratory clinical disease that consisted with increased body temperature, lymphadenopathy, increased lung sounds, increased tracheal mucus, and increased nasal discharge (Fig. 7-8). None of these clinical signs were

significantly extreme to require additional animal care or supportive treatment. Neither respiratory rate, nor heart rate was significantly different between groups. None of the clinical signs developed by the infected group were observed on the control animals (n=4), which remained healthy throughout the extent of these trials. The etiological agent (ERAV) was only recovered from ponies in the infected group for up to seven days (Table 8). Ponies in the re-infected group (n=4) did not developed clinical respiratory signs and remained healthy throughout the experiments. The statistical analysis of the data collected during these trials demonstrated a significant difference between infected and control animals.

Table 15. Equine rhinitis A virus (ERAV) shedding summary. Results of virus isolation from samples collected during the ERAV/ON/05 experimental infection.

Sample location	Sample collection day						
	Day 0	Day 1	Day 3	Day 5	Day 7	Day 14	Day 21
Pharyngeal Swab	-	+	+	+	+	-	-
Pharynx brush biopsy	-	+	+	+	+	-	-
Mid trachea brush biopsy	-	+	+	+	+	-	-
Carina brush biopsy	-	+	+	+	-	-	-
BAL	-	+	+	-	-	-	-
Plasma	-	+	+	+	-	-	-
Urine	-	+	-	-	+	-	+
Feces	-	-	-	-	-	-	-

+ Virus was recovered from clinical sample on cell culture (RK-13 cells)

- Virus was not recovered from clinical sample

BAL Bronchoalveolar lavage

[0221] Neither depression, nor appetite lost was recorded in either group. As well hydration, urination, defecation, and gastrointestinal movements remained unchanged on all groups during the infection trials. There was no difference between re-infected and control animals on the clinical scores and virus isolation test, since no clinical disease was observed on the re-infected animals and the virus could not be recovered from this group.

Clinical findings

[0222] All ponies were considered to be healthy prior to these experiments. ERAV/ON/05 exposure induced clinical respiratory disease in infected animals compared to controls and re-infected animals. The main clinical signs detectable by physical examination were pyrexia, nasal discharge, and lymphadenopathy. Endoscopic examination also revealed the presence of large volumes of mucus and hyperaemia in the mid trachea and lower airways of the infected animals.

Rectal temperature

[0223] No significant differences in daily rectal temperature were found between groups prior to viral or placebo exposure. A significant treatment effect was identified when infected animals were compared to controls and re-infected ($p = 0.002$). When the body temperature from the three groups was compared at different times, the infected group was significantly different compared to controls and re-infected animals ($p = 0.028$). Increased body temperature was detected 24 hours post-infection only in the infected animals and was significantly different from day 2.5 to day 6 when compared to control animals at the same time points ($p = <0.005$). The body temperature increase peaked on day 4 with a mean body temperature of 38.45°C ($\text{SE}=0.15$) ($p = 0.001$) and persisted for two more consecutive days (Fig. 8). No statistical difference was found when body temperatures from control and re-infected groups were compared at different times. Animals in the control and re-infected groups did not have a significant change in their body's temperature when comparing baseline to individual points in time within the groups (days 1, 7, 14 and 21).

Lymph nodes

[0224] Submandibular and retropharyngeal lymph nodes were examined daily and classified as non palpable, palpable and enlarge. Palpability or enlargement of the lymph nodes was only recorded on the infected and re-infected animals. In all infected ponies, the submandibular area became sensitive to palpation on day two and in most cases persisted for up to two weeks. The submandibular lymph nodes size in the infected animals varied from 3 to 5 cm in length by 2 to 3 cm in thickness, and the analysis of the total scores demonstrated a statistical difference between groups ($p = <.0001$) (Fig.7). Interestingly, the retropharyngeal

lymph nodes were not consistently palpable in all infected animals, but a significant change in size was recorded in one pony. This lymphadenopathy did not appear to interfere with food or water consumption and the sensitivity to palpation became less pronounced as the days progressed. Submandibular lymph nodes were palpable in three animals from the re-infected animals with an average size of less than one cm in length and 0.5 cm in thickness approximately. Control animals had no detectable lymph node changes at palpation throughout the infection experiments.

Heart and respiratory rate

[0225] Respiratory rate (RR) and heart rate (HR) means were not significantly different between treatment groups ($P = 0.1$) at any time points. In general RR and HR were within the normal physiological parameters and small changes were only associated with handling and sample collection. The highest RR mean among all groups was identified on day 0 and the lowest mean from all the groups was recorded on day 21.

Endoscopic examination

[0226] On the endoscopic examination, infected animals had an increase amount of tracheal mucus detectable on day one that persisted up to day 21. Neither the control nor the re-infected animals had mucus secretions detectable at endoscopic examination throughout these experiments.

[0227] Characteristics of the mucus varied from clear and serous on day one to mucoid on days 7-21. Mucus patches were consistently distributed from the upper trachea to the bifurcation of the carina. Localized tracheal hyperaemia was observed in all infected and in some control animals throughout these experiments. The carina on all infected animals was blunted and in some cases hyperaemic starting on day three. Sensitivity to endoscopic examination was noticeably increased by day seven on infected animals and bronchoconstriction was observed during BAL. Nasal discharge varied from mild to moderate in all infected ponies and was not present in the control and re-infected groups. Serous nasal discharge was observed during clinical examination for about 8 days on the infected animals starting between 36 and 48 hours post-infection. However this discharge was not a reflection of the mucus (characteristics and

volume) observed during endoscopic examination. Mild ocular discharge was observed inconsistently in the infected animals.

Serology

[0228] A total of 12 ponies (age 10 to 12 months) were infected with ERAV/ON/05 or placebo by nebulization. All ponies were serologically negative to ERAV, ERBV, AE2, and EHV1/4 and in a healthy condition prior to the infection experiments. Following exposure to ERAV/ON/05 all infected animals (100%) seroconverted (four fold increase) to ERAV determined by the virus neutralization test (VN) (Fig. 9). A significant treatment by day interaction was observed in infected animals ($P = <0.0001$). A statistical difference between infected and re-infected animals was found on baseline and day 7 ($P = <0.001$). In the re-infected group no statistical differences were found when titers to ERAV on days 7, 14, and 21 were compared back to baseline.

[0229] Antibody titers against ERAV were significantly elevated in infected animals from day 7 and in most cases peaked by day 14 and maintained to day 21 ($P = <0.001$). In contrast, all control animals remained seronegative to ERAV and no detectable changes were identified by the VN test. Animals in all groups did not show an increase in antibody levels or serological conversion to any other respiratory viruses (ERBV, AE2, and EHV1/4) during these experiments (Fig. 9). Animals in the re-infected group ($n=4$) did not have a significant difference in antibody titers to ERAV between baseline and day 21 post-infection. However, a small change in antibody levels to ERAV was detected in 3 ponies and a four fold increase in one pony from the same group (Fig. 9).

Virus Isolation

[0230] Nasopharyngeal swabbing, laryngeal brushing, tracheal brushing, BAL, fecal and urine samples were negative for virus isolation (equine respiratory viruses) on all ponies prior to infection (Table 8). Swabs obtained from the nasopharynx from infected and control animals after completing nebulization were cultured on RK-13 cells and ERAV was recovered on a first passage from all infected animals only. RT-PCR using primers that targeted the VP1 gene confirmed the positive and negative diagnoses from both groups.

[0231] Virus isolation results from the infection and re-infection trials are summarized on Table 8. A significant difference between infected, control and re-infected animals was identified when comparing virus isolation between groups ($P = < 0.05$). ERAV was only recovered from animals in the infected group on specific days and specific areas of the respiratory tract (Table 8). No other viruses were recovered from samples collected during these experiments. All ponies in the control group were negative in the virus isolation tests throughout the study. A treatment by day interaction on infected animals was first detected on day 7 and persisted on days 14 and 21 ($P = < 0.05$).

[0232] Location for virus recovery varied from the lower airways on days 1, 3, and 5 to mid and upper airways on days 1, 3, 5, and 7. On day one ERAV was isolated from the pharynx and carina from all the ponies in the infected group and the mid trachea and BAL from 3 ponies from the same group. All attempts for virus recovery from feces in all groups were unsuccessful. Virus isolation from urine and plasma was achieved only in rare occasions (Table 8). Virus recovery was gradually decreased from day 1 up to day 7 when ERAV was consistently recovered from clinical samples. This last viral recovery was associated with an increase in antibody titre to ERAV and a decrease in the clinical signs scoring (Figs. 7 and 9).

Pulmonary function testing (PFT)

[0233] Hyperactivity of the airways was assessed based on the physiological and clinical response to histamine provocation. Data from infected and control animals were plotted and triggering histamine doses were calculated. Interestingly, ponies from both groups (infected and control) responded on day 0 to a low dose of histamine (< 6 mg of histamine). Overall, the triggering histamine doses did not go beyond 13 mg. The clinical histamine reaction (dose-dependant) was observed as hyperventilation associated with abdominal lift and breathing difficulty. The physiological reaction was detected in the PFT by a 35% drop in lung dynamic compliance (C_{dyn}) or a doubling in the transpulmonary pressure (ΔP_{pl}) when comparing saline and histamine administration. Ponies in the infected group showed a small reduction to the histamine triggering dose from day 0 to day 1. However, this was not significantly different between groups. A significant difference between infected and controls was detected on day 21 ($P = 0.02$).

BAL fluid differential cell counts

[0234] Differential cell counts were carried out on the cytospin slides prepared from BAL fluid aliquots. No significant differences in the cell counts were found among horses in the different treatment groups prior to the infection trial. No treatment by day effect was detected on the macrophage and epithelial cells percentages throughout the experiments. A significant increase in the number of neutrophils was observed on day 7 post-infection in the infected group ($P = <0.05$). These numbers were not significantly different on the control or re-infected animals when comparing base line to days 7, 14, and 21. The mean percentages of lymphocytes, eosinophils and mast cells were proportionally decreased on day 7 post-infection in the infected animals and a statistical difference was detected ($P = <0.05$). Ciliated epithelial cells were commonly observed on the slides from infected and control animals, however, no significant differences were detected, except on one of the infected animals that had a high count on day 7. In general, a non-septic suppurative inflammation with the presence of epithelial cells and sporadic giant cells was detected in the infected ponies. Interestingly, no major changes in the cytological examination were observed on the samples from the re-infected animals.

[0235] This study demonstrates that ERAV/ON/05 induced clinical respiratory disease in infected ponies. Serology demonstrated that no other respiratory viruses were present during these trials. The disease is characterized by pyrexia, nasal discharge, increased lung sounds, and increased submandibular lymph nodes size. Additionally, large volumes of mucus were endoscopically detected in the lower airways that persisted up to day 21. The virus was isolated from the lower and upper airways up to day 7 corresponding with the appearance of detectable equine rhinitis A virus (ERAV) antibodies. None of the re-infected animals developed clinical disease and only one pony from this group had a four fold increase in the antibody titers to ERAV. Ponies with pre-existing ERAV antibodies did not develop clinical disease when exposed to the virus.

Example 5

[0236] This example illustrates one embodiment of a Equine Rhinitis B Virus composition in accordance with the present invention.

Materials and Methods

[0237] Equine Rhinitis B Virus strain **07-10342** (ATCC Accession NoPTA-11829) was recovered from Rabbit-kidney-13 (RK-13) cell culture from a nasal swab from a horse in Ontario Canada. Inactivated **07-10342** was produced following the general procedure described in Example 1 for ERAV/ON/05.

[0238] The following compositions containing ERBV alone or in combination with ERAV were prepared.

Table 16. Vaccine Groups

Vaccine Group	Vaccine	Titers A /B	Adjuvant	Volume
1 (N=8)	Monovalent B - High	7.5	HRA-5	200 mL
2 (N=8)	Monovalent B - High	7.5	HRA-3 with CSO	100 mL
3 (N=8)	Monovalent B - Low	7.0	HRA-5	200 mL
4	Bivalent A and B - High	8.0 / 7.5	HRA-5	200 mL
5	Bivalent A and B - High	8.0 / 7.5	HRA-3 with CSO	100 mL

[0239] Eight horses from each of vaccine groups 1, 2, and 3, were challenged along with eight control horses with a $10^{6.6}$ TCID₅₀ challenge dose. Disease status based on nasal discharge and conjunctivitis scores as indicated in Table 10. Effect of vaccination on duration, severity, and incidence of disease is shown in Tables 17-22.

Table 17. Disease status scoring

Disease Status	Nasal Score	Conjunctivitis Score
Normal (0)	0 or 1	0 or 1
Mild (1)	0 or 1	2
Mild (1)	1.5, 2, or 3	any
Moderate (2)	4 or 6	any

[0240] Vaccinated group showed a significant reduction (Table 18) in duration of nasal discharge including in some cases no signs of mild respiratory disease.

Table 18. Number of days with mild, moderate, or severe respiratory disease

Vaccine Group	Minimum	25th quantile	50th quantile	75th quantile	Maximum
Control (N=8)	1	5	8	8	9
1 (N=8)	0	0	1	3.5	7
2 (N=8)	0	0.5	1	6.5	8
3 (N=8)	0	1	1	3	6

[0241] The number of days animals were sick with clinical signs of respiratory disease (nasal score > 0) was significantly shorter in the vaccinated group compared to the control group.

Table 19. Effect of vaccination on duration of disease

	Control	Vaccinate	Shift in days	95% confidence interval
Duration vs Group 1	8	1*	5	1, 8 days
Duration vs Group 2	8	1*	2	0, 8 days
Duration vs Group 3	8	1*	7	1, 7 days

*Significantly lower than the control group by Kruskal-Wallis test (P<0.05)

[0242] Immunization with ERBV also lessened the severity of the disease with a lower percentage of vaccinates than controls demonstrating mild and moderate clinical signs of respiratory disease throughout the study (Table 20).

Table 20. Summary of the severity of disease based on the maximum respiratory score

Vaccine Group	Normal	Mild	Moderate
Control (N=8)	0% (0/8)	87.5% (7/8)	12.5% (1/8)
1 (N=8)	37.5% (3/8)	50.0% (4/8)	12.5% (1/8)
2 (N=8)	25.0% (2/8)	75.0% (6/8)	0.0% (0/8)
3 (N=8)	12.5% (1/8)	87.5% (7/8)	0.0 % (0/8)

[0243] Immunization with ERBV was found to significantly reduce the incidence of disease, by 37.5%, 25%, and 12.5% in vaccinated groups 1, 2, and 3, respectively (Tables 21 and 22).

Table 21. Effect of vaccination on incidence of disease (disease statuses of mild and moderate were pooled for the purposes of this evaluation)

	Control (n=8)	Group (n=8)	Prevented fraction	95% confidence interval
Control vs. Group 1	1	62.5%	0.375	-0.069, 0.6346
Control vs. Group 2	1	75.0%	0.25	-0.119, 0.4973
Control vs. Group 3	1	87.5%	0.125	-0.137, 0.3266

[0244] Vaccination also reduced the shedding of virus from the nares indicating lesser clinical disease in the vaccinated horses as well as demonstrating the effectiveness of the vaccine in preventing spread of the infectious disease to other potentially susceptible horses (Table 22).

Table 22. Incidence of virus positive over time

Day	Control	Group 1	Group 2	Group 3
0	0.250	0.25	0.000	0
1	0.625	0.25	0.125	0
2	0.125	0.00	0.000	0
3	0.125	0.00	0.000	0
4	0.250	0.00	0.000	0
5	0.125	0.00	0.000	0
6	0.125	0.00	0.000	0
7	0.125	0.00	0.125	0

Results and Discussion

[0245] Inactivated ERBV vaccines were found to be capable of immunizing an animal to high levels of antibody detection. The vaccines reduced the duration, severity, and incidence of disease in immunized animals challenged with ERBV. Vaccination also reduced the shedding of infectious virus from ill horses. No unacceptable adverse reactions either at the injection sites or by manifestation of signs of systemic illness were observed. The vaccines are safe and well tolerated for administration in Equine rhinitis B susceptible species, particularly equidae.

Example 6

[0246] This example illustrates a guinea pig model for use in a release potency assay.

[0247] Vaccine 5 (A/B-1) and Vaccine 6 (A/B-2) were each administered to guinea pigs (five guinea pigs per vaccine, 0.5 mL per intramuscular vaccination). A booster vaccination shot was administered three weeks later. After 19 days, the pigs were bled and blood was analyzed using serum neutralization tests for ERAV and ERBV.

Table 23. Vaccine 5 (A/B-1) Batched with $10^{8.0}$ A/ $10^{7.5}$ B, with HRA-5 adjuvant

Guinea Pig	ERVA Titer	ERV B Titer
1	>1414	280
2	1300	180
3	448	120
4	1123	360
5	893	1120
Back titer	191	260

Table 24. Vaccine 6 (A/B-2) Batched with $10^{8.0}$ A/ $10^{7.5}$ B, with HRA-3 + cottonseed oil adjuvant

Guinea Pig	ERVA Titer	ERV B Titer
1	194	220
2	326	280
3	194	560
4	282	240
5	No Sample	No Sample
Back titer	191	260

Example 7

[0248] This example illustrates a challenge evaluation of Rhinitis Virus A after vaccination with a 2 dose equine Rhinitis A/Rhinopneumonitis/Influenza Killed Virus Vaccine and protection against Equine Rhinitis A respiratory infection after vaccination with the Rhinitis A/Rhinopneumonitis/Influenza Killed Virus Vaccine.

Objective

[0249] The objective of this vaccination-challenge study was to demonstrate efficacy of Equine Rhinitis A (ATCC Accession No. PTA-11828) for Equine Rhinitis A/Rhinopneumonitis/Influenza Killed Virus Vaccine. The primary outcome variable used to evaluate the efficacy of vaccination was reduction of respiratory disease caused by Equine Rhinitis A virus.

Material and Methods

[0250] The vaccine used in this study is an Equine Rhinitis A/Rhinopneumonitis/ Influenza Vaccine of the present invention, Killed Virus vaccine.

A. EQUINE RHINITIS A VIRUS

[0251] The original Equine Rhinitis Virus A Strain (ERhA V) was obtained from University of Guelph, Animal Health Laboratory, as Isolate number 04-54188, and was received on September 9, 2008 under Import Permit No. 106930. The virus was passed once on E-Vero cells to produce a high titered pre-master stock and was then diluted with cell culture media to produce the Master Seed Virus (MSV). The MSV is designated as ERhA V (04-54188), MSV, Lot 091508A-diluted, 24Sept08, and was approved for use for Establishment 597 by USDA on June 18, 2010.

[0252] The Equine Rhinitis A viral antigen used in vaccines evaluated in this study was a MSV+5 virus produced on the twentieth passage of APHIS approved E-Vero cells. Following growth, viral fluids were filtered, formalin inactivated, and concentrated in accordance with the Outline of Production for Product Code A522.20. The inactivated viral fluids were tested for residual live virus after inactivation. The results were satisfactory. Inactivated viral fluids were used to formulate vaccine at an antigen inclusion level of $10^{7.5}$ TCID₅₀/mL.

1. Experimental Vaccine Rhinitis Combo Lot 122110

[0253] Experimental vaccine Rhinitis Combo Lot 122110 was formulated based on pre-inactivation titers.

[0254] The final formulated vaccine contained the following ingredients per 1 mL dose:

Equine Rhinitis A	$10^{7.5}$ TCID ₅₀ /mL
EHV-1	$10^{7.0}$ TCID ₅₀ /mL
EHV-4	$10^{6.5}$ TCID ₅₀ /mL
Influenza A2/Ohio/03	$10^{7.0}$ TCID ₅₀ /mL
Influenza A2/KY/95	$10^{7.0}$ TCID ₅₀ /mL
Influenza A2/NewMarket/2/93	$10^{7.0}$ TCID ₅₀ /mL
Adjuvant (MVP Laboratories, S.O. #25)	100 μ L
Diluent, MEM-E containing	q.s.
Gentamicin, 30 μ g/mL of diluent volume	
Formaldehyde, 0.1% of diluent volume	
Amphotericin B, 2.5 μ g/mL	

B. Experimental Horses

1. Description of Experimental Horses

[0255] Forty (40), six to eight-month old, draft-cross horses purchased from Steve Waagen, Bottineau, North Dakota were microchipped upon arrival at Equine Resources, LLC, Butler, Mo. and were assigned to either IVP (Investigational Veterinary Product) or Control Product (CP) based on random number generator after being pre-screened for rhinitis A titers of $\leq 1:4$.

[0256] During the entire study, horses were quartered in a single large paddock with common feed bunks, waterers and hay racks. Upon challenge, each animal was assigned a 2 digit “barn code” by laboratory personnel that was attached to a halter worn throughout the post-challenge time period and used to identify horses when clinical signs and samples were taken each day. During each observation day, horses were corralled into holding pens and worked randomly through individual restraining chutes.

Table 25 summarizes the study design:

Table 25: Study Design

Group	No. Test Animals	Treatment	Doses (21 day intervals)	Route of Administration	Challenge (Study Day)
1	20	IVP	2 x 1 mL	IM	D46
2	20	CP	2 x 1 mL	IM	D46

2. Vaccination /Challenge and Sampling Schedule

[0257] On January 28, 2011 and February 18, 2011, experimental vaccine Rhinitis Combo Lot 122110 was administered intramuscularly in a 1 mL dose volume to each of 20 horses (vaccinate group, IVP). Twenty horses (control group, CP) received a 1 mL dose of adjuvanted MEM-E (Experimental product 005) containing the excipients used in the Lot 122110 vaccine (adjuvant, gentamicin, amphotericin B, and formaldehyde) but no antigens. All horses were challenged by intranasal aerosolization of virulent Equine Rhinitis A virus at Study Day 46 (25 days post-booster vaccination) on March 15, 2011. Table 26 shows the schedule of events.

Table 26: Schedule of Events:

Calendar Date	Study Day	Study Event
Jan 28, 2011	0	Randomize horses to groups Collected serum samples all horses IVP administered to Group 1 CP administered to Group 2
February 18, 2011	21	Collected serum samples all horses IVP administered to Group 1 CP administered to Group 2
March 15, 2011	46	Challenge Groups 1 and 2 Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations
March 16-21, 2011	47-52	Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations
March 22, 2011	53	Serum samples Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations
March 23-28, 2011	54-59	Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations
March 29, 2011	60	Serum samples Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding)

		Clinical observations
March 30-April 4, 2011	61-66	Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations
April 5, 2011	67	Serum samples Body Temperatures Whole blood samples (virus isolation) Nasal Swabs (virus shedding) Clinical observations End of Study

3. Intranasal Challenge Inoculation of Horses

a. Challenge Virus

[0258] The challenge virus Equine Rhinitis A lot 112108A was produced in tissue culture on E-Vero cells. The titer of the challenge virus was determined to be $1 \times 10^{7.3}$ TCID₅₀/mL on the day of challenge.

b. Intranasal Challenge Method

[0259] Sedivet® (romifidine hydrochloride), a sedative and analgesic, was administered intravenously to each horse prior to challenge at a dosage of 50 µg/kg of body weight. The challenge virus was administered intranasally as an aerosol produced by a nebulizer into an Equine AeroMask (Trudell Medical International, Ontario, Canada) by the following method: Four milliliters of $10^{7.3}$ TCID₅₀/mL challenge virus were placed into the nebulizer cup in the AeroMask device. A pressure hose was fitted from an air compressor to the inlet port of the nebulizer. The outlet tube was then inserted into the AeroMask attached to the head of the horse being challenged and approximately 10 psi of air pressure was applied to the inlet port for three minutes. During this time approximately two milliliters of challenge virus fluid was aerosolized directly into the nostrils of the horse being challenged. Challenge virus was administered to horses undiluted, effecting a challenge amount of $1 \times 10^{7.6}$ TCID₅₀ in a 2 mL dose.

C. PRE AND POST CHALLENGE EVALUATION PARAMETERS

1. Nasal Exudate Evaluation

[0260] All nasal exudate observations were made prior to collection of nasopharyngeal swabs. On the Day of Challenge (D46) and for 21 days post challenge, the nasal passages and muzzle of each of the 40 vaccinate and control horses were examined and graded using the grading and scoring description listed below.

The scoring grades of 0 through 6 were assigned on the basis of the severity of the disease indicated by each of the following classification:

Description of Symptoms	Score Sheet Designation	Score
Essentially normal indicates the horse was clean and essentially free of nasal exudate	EN	0
Slight clear serous discharge that may be frequently observed in both diseased and normal horses	C-1	1
Very slight mucopurulent discharge indicates that mucus was definitely present in small amounts in either one or both nostrils	VSM	1.5
Moderate clear serous discharge is indicative of a definite increase in volume over that normally observed/slightly mucopurulent is a discharge easily observed in one or both nostrils	C-2/SM	2
Copious clear serous discharge that is generally observed only in diseased horses	C-3	3
Moderately mucopurulent indicates that mucoid discharges were present in large quantities in both nostrils	MM	4
Heavy mucopurulent indicates that copious amounts of a mucoid discharge filled both nostrils	HM	6

2. Ocular Discharge

[0261] Ocular discharge was evaluated daily at the time of nasal exudate evaluation. Ocular discharge scores were recorded as 0 = normal; 1 = mild to moderate ocular discharge and 2 = severe ocular discharge.

3. Temperature

[0262] Daily rectal temperatures were recorded for each of the 40 vaccinate and control horses on Day of Challenge and for 21 days post challenge by means of a calibrated, electronic thermometer (GSA Electronics) probe. The daily rectal temperatures were recorded in degrees Fahrenheit (°F).

4. Nasopharyngeal Viral Isolation

[0263] On each observation test day each nasal passage of each vaccinated and control horse was swabbed deeply by means of a sterile WECK-CEL® surgical spear (Edward Weck and Company, Inc., Research Triangle Park, N.C. 27709) attached to an 11-inch long sterile plastic pipette. On collection, each of two surgical spears was immediately placed in a single tube containing 4 mL of chilled transport medium (E-199 supplemented with gentamicin, L-glutamine, 2X Pen/Strep, 2X Amphotericin B).

[0264] For isolation of virus, the tubes were mixed, the swabs aseptically removed, and the medium centrifuged at 1500 rpm for 10 to 15 minutes to remove particulates. Medium was filtered through a 0.2 µ syringe filter prior to inoculation on tissue culture cells. After filtration, 4-6% of sterile 85% sucrose solution was added to each sample for freezing at -80°C in order for all samples to be tested concurrently.

[0265] On the day of testing, one mL of the thawed, clarified transport medium was used to inoculate a 2 cm² two day old monolayer of E-Vero cells grown in a 24 well tissue culture plate from which the growth medium had been aseptically removed. Following inoculation, the inoculum was allowed to adsorb on the cell monolayer for at least one hour at 37°C in a humidified incubator containing a 5% CO₂ atmosphere. After the adsorption period, an additional 1 mL of re-feed medium (E-199 containing 2 mM L-glutamine, Gentamicin 2X Pen-Strep and 2X Amphotericin B) was added to each well. Following addition of re-feed media the plates were then incubated at 37°C in a CO₂ incubator. Each test and control tissue culture well was examined microscopically for 7 days for signs of cytopathic effect (CPE) typical of the

Equine Rhinitis A challenge virus. Wells that were negative at the end of the 7 day observation period were subcultured onto fresh cells and observed for an additional 7 days.

Statistical Evaluation Methods

[0266] Horses were classified to a range of respiratory disease status, on a daily basis. The classification to disease status included combining the nasal and ocular evaluations. The algorithm used is detailed below:

Disease Status	Nasal Score	Ocular Score	Discharge
Normal (0)	0 or 1	0 or 1	
Mild (1)	0 or 1	2	
Mild (1)	1.5	any	
Moderate (2)	2 or 3	any	
Severe (3)	4, 5 or 6	any	

[0267] The influence of vaccination on the duration of disease (number of days with at least moderate disease) was evaluated using the Hodges-Lehman (exact) estimate (the NPAR1WAY procedure in SAS, SAS Institute, Cary NC).

[0268] Severity of disease was evaluated by comparing the maximum disease status between the vaccinated horses and the placebo horses. The mitigated fraction and 95% confidence intervals (CI, asymmetric standard error) were calculated (the FREQ procedure in SAS).

[0269] Rectal temperature and serum neutralization titers were analyzed by repeated measures analysis appropriate for continuous data (temperature and log transformed SN titers; ANOVA). SN titers were assumed to be 0 if the titer was reported as < 4 and 709 if the titer was reported as > 709. SN titers were log transformed prior to the analysis. Buffy coat and nasal swab results on each day were compared using Fisher's exact test.

Results

[0270] One horse, #39 (IVP group) died on Day 62 of the study. Upon necropsy at University of Missouri College of Veterinary Medicine, the diagnosis was made of chronic-active, diffuse, severe necrotizing and emphysematous submucosal esophagitis, gastritis, and pharyngitis with intralesional coccobacilli bacterial colonies and plant material, as well as a diffuse severe fibrinosuppurative pleuropneumonia of the lung. The most likely explanation of this death suggests a previous mucosal breach, such as a mucosal ulcer in the cardia of the stomach which allowed seeding of plant material in the associated connective tissues (pathology report attached).

A. DURATION

[0271] The distribution of the duration of disease (number of days with moderate or severe respiratory disease) in days is summarized in Table 27. The median number of days animals in the placebo group were observed with disease was 14. In the vaccinated group, the median number of days with disease was 1. The duration of disease was significantly lower in the vaccinated animals as compared to the controls (Table 28; $P < 0.0001$).

Table 27: Summary of the effect of vaccination on the duration of disease (Number of days with moderate or severe respiratory disease)

Group	Minimum	25 th quantile	50 th quantile	75 th quantile	Maximum
Control (N = 20)	0	10	14	17	18
Vaccinated (N = 20)	0	0	1	4	12

Table 28: Effect of vaccination on the duration of disease (Number of days with moderate or severe respiratory disease)

	Control	Vaccinate	Shift in days	95% Confidence Interval
Duration (median days)	14	1	11	-14, -8

Severity

[0272] The distribution of the severity of disease is summarized in Table 29. The severity of disease was significantly lower in the vaccinated horses as compared to the placebo horses (Table 30; mitigated fraction = 0.7550, 95% CI = 0.5518, 0.9582). Individual outcomes are provided in Table 31.

[0273] In Table 31 the values for horse #39 which died on Day 62 of the study are reported as 0's from Day 62 through 67, but in the analysis only values > 1 are considered, therefore not affecting the outcome. Duration is 1 day for this horse and the max score is 3.

Table 29: Summary of the severity of disease based on the maximum nasal discharge score

Group	Normal	Mild	Moderate	Severe
Control (N = 20)	0% (0/20)	5% (1/20)	20% (4/20)	75% (15/20)
Vaccinated (N = 20)	5% (1/20)	35% (7/20)	55% (11/20)	5% (1/20)

Table 30: Effect of vaccination on the severity of disease

	Control n = 20	Vaccinate n = 20	Mitigated Fraction	95% Confidence Interval
Severity (mean rank)	28.05	12.95*	0.7550	0.5518, 0.9582

*Significantly lower than the placebo group by Wilcoxon's rank sum test ($P < 0.05$).

Table 31: Results from individual animals (shaded regions denote duration of moderate to severe disease.

Horse	Vacc	Day 46	Day 47	Day 48	Day 49	Day 50	Day 51	Day 52	Day 53	Day 54	Day 55	Day 56	Day 57	Day 58	Day 59	Day 60	Day 61	Day 62	Day 63	Day 64	Day 65	Day 66	Day 67
42849285	Combo	0	0	0	0	0	0	0	0	0	1	0	0	0	0	1	0	1	1	0	0	0	0
42856542	Combo	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	0	2	0	0	0	0
42857074	Combo	0	0	0	0	0	0	1	0	0	2	0	0	0	0	0	2	0	0	0	0	0	0
42860382	Combo	0	0	0	0	0	0	2	0	0	2	2	2	0	0	1	0	0	0	0	0	1	0
42861051	Combo	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0
42876062	Combo	0	0	0	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0	0
42878629	Combo	0	0	0	0	0	1	1	0	0	2	0	1	0	0	0	0	0	2	0	0	0	0
42889377	Combo	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	2	0	0	0
42889622	Combo	0	0	0	1	0	1	0	1	0	0	2	2	1	0	0	0	0	0	0	0	0	0
42890836	Combo	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
43315773	Combo	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	1
108356559	Combo	0	0	0	0	0	1	0	0	2	1	0	2	2	2	0	0	1	0	2	0	0	0
108357033	Combo	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	0
108361284	Combo	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
108363552	Combo	0	0	0	0	0	0	0	1	0	0	2	0	0	0	0	0	0	0	0	0	0	0
108366283	Combo	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0
108368584	Combo	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0
108512267	Combo	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0	0	0	0	2
108513075	Combo	0	0	0	0	0	0	0	0	0	0	0	0	3	1	0	0	0	0	0	0	0	0
108522069	Combo	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0

Table 31 (Continued): Results from individual animals (shaded regions denote duration of moderate to severe disease.

Horse	Vacc	Day 0	Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7	Day 8	Day 9	Day 10	Day 11	Day 12	Day 13	Day 14	Day 15	Day 16	Day 17	Day 18	Day 19	Day 20	Day 21	
18605089	Placebo	0	0	0	1	0	1	0	0	1	2	0	1	0	0	1	2	2	1	2	2	1	0	0
42849558	Placebo	0	0	1	0	0	1	1	2	0	0	1	1	1	0	1	3	2	0	0	0	1	1	1
42850873	Placebo	0	1	0	0	0	2	1	1	1	2	1	1	0	1	1	0	3	2	0	2	2	3	3
42861315	Placebo	0	0	0	1	1	1	2	0	0	2	3	1	3	0	2	3	0	2	3	0	1	0	0
42865808	Placebo	0	0	0	1	2	2	1	2	2	2	1	3	2	2	0	2	2	2	3	2	1	0	0
42867036	Placebo	0	0	0	1	2	1	2	3	2	2	3	2	0	3	2	0	0	1	0	1	0	3	3
42867892	Placebo	0	0	0	1	1	0	1	1	2	2	1	1	3	0	1	1	2	0	1	1	0	0	0
42869592	Placebo	0	0	1	2	2	1	2	1	2	2	0	0	2	0	2	0	2	1	2	2	1	0	0
42870316	Placebo	0	0	1	1	2	2	1	2	2	2	2	1	2	2	1	1	2	1	2	0	0	0	0
42874866	Placebo	0	0	1	0	2	2	2	2	3	3	3	3	0	2	1	3	2	1	0	1	0	1	1
42876021	Placebo	0	0	0	0	1	1	1	0	0	2	0	2	0	0	0	2	0	0	0	1	0	0	0
42879094	Placebo	0	0	0	0	0	0	1	0	0	3	0	1	2	3	1	0	2	0	0	1	0	0	0
42879533	Placebo	0	0	0	1	0	0	0	1	2	2	1	2	1	1	0	2	0	3	0	0	0	0	0
42889258	Placebo	0	0	0	0	0	1	1	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
42890047	Placebo	0	0	0	2	0	1	0	0	1	1	0	0	0	2	0	0	0	3	1	0	0	0	0
42891256	Placebo	0	0	0	0	2	1	1	0	1	3	3	2	1	2	0	2	2	1	0	0	1	2	2
42892606	Placebo	0	0	0	0	0	1	1	0	1	3	3	3	2	3	1	2	1	0	0	0	2	0	0
108339771	Placebo	0	1	0	1	0	2	1	0	0	3	2	2	2	3	0	2	2	1	2	0	1	3	3
108340611	Placebo	0	0	2	1	2	3	2	2	3	3	2	2	2	3	2	1	2	1	2	2	1	1	1
108341625	Placebo	0	0	0	1	0	0	2	2	3	1	2	2	1	0	2	0	3	2	0	2	2	1	1

B. RECTAL TEMPERATURE

[0274] The results of the statistical analysis are summarized in Table 32. The vaccine by day interaction was statistically significant. Within day analyses are provided in Table 33 (appendix). On Days 3, 4, and 5, horses in the vaccinated group had statistically significantly lower temperatures than those in the placebo group. On Day 20, horses in the vaccinated group had significantly higher temperatures than those in the placebo group. However, on all days any differences between treatment groups were not clinically meaningful and only one horse (#39) had a rectal temperature greater than 102°F (39°C) on any challenge study day.

Table 32: Summary of the effect of vaccination on rectal temperature and serum neutralization titers

Outcome	P-Values		
	Vaccine	Day	Vaccine * Day
Temperature	0.5754	< 0.0001	0.0006
SN titer	< 0.0001	< 0.0001	< 0.0001

1. Serum Neutralization Titers

[0275] The results of the statistical analysis are summarized in Table 32. On Days 46 and 53, horses in the vaccinated group had significantly higher titers than those in the placebo group. On Day 67, horses in the vaccinated group had significantly lower than those in the placebo group.

Discussion

[0276] This study was conducted to demonstrate efficacy of the Equine Rhinitis A virus component of an inactivated Equine Rhinitis A virus vaccine in combination with inactivated Equine Herpesvirus types 1 and 4 and Equine Influenza Viruses. Vaccination was given in a 2-dose format, and challenge of virulent Equine Rhinitis A virus was performed 25 days post-booster vaccination. Vaccinated and placebo treated control horses were evaluated to assess effect of vaccination on reduction of severity and duration of clinical respiratory disease. Clinical signs, nasal swabs and buffy coats were evaluated daily for evidence of Equine Rhinitis A disease and presence of the virus in the challenged horses.

[0277] Results from this challenge study show a statistically significant and clinically important reduction of both the severity and of the duration of respiratory disease in vaccinates following challenge. As confirmed by mitigated fraction analysis of respiratory disease scores in horses vaccinated with the IVP as compared to horses vaccinated with the control product, moderate/severe signs of respiratory disease were reduced 75.5% (95% confidence interval 96% to 55%). Duration of moderate/severe disease was also significantly reduced by 11 days in vaccinated horses as compared to control horses (95% confidence interval 14 days to 8 days shorter disease duration). In addition, vaccination significantly reduced nasal virus shedding throughout the successive 5 days of peak viral shedding, and viremia also was significantly reduced during the 4 days of peak viremia within the challenge period. Importantly, virus was recovered from buffy coat samples of control horses a total of 54 times over 5 study days in comparison to only 8 total days of viremia over 4 study days in vaccinated horses.

Conclusions

[0278] Data from this study clearly demonstrate that 2 x 1 mL intramuscular doses of Equine Rhinitis A/Rhinopneumonitis/Influenza Killed Virus Vaccine (Product Code TBD, unlicensed), administered at 21 day intervals between doses to horses, significantly reduced severity and duration of respiratory disease caused by Equine Rhinitis A virulent challenge.

[0279] The study results illustrate the use of this vaccine for the immunization of horses six months of age or older against respiratory disease caused by Equine Rhinitis A virus. These data also establish Vaccine Lot 122110 as an acceptable reference vaccine for potency testing of subsequent serials of Equine Rhinitis A/Rhinopneumonitis/ Influenza Vaccine, Killed Virus and of Equine Rhinitis A Vaccine, Killed Virus (Product Code 1522.21, unlicensed).

What is claimed is:

1. An immunogenic composition comprising one or more strains of inactivated or live, attenuated Equine Rhinitis A Virus (ERAV) or Equine Rhinitis B Virus (ERBV), wherein the ERAV strain or the ERBV strain, prior to inactivation or attenuation, causes detectable respiratory disease in at least 50% of seronegative horses exposed to the strain, or grows in cell culture to 10^6 TCID₅₀/mL or higher, or, when used as a vaccine in equines at a dose of 10^6 TCID₅₀ or higher results in a serum titer of at least 1:112.
2. An immunogenic composition comprising an ERAV strain comprising a genomic sequence whose reverse transcript has greater than 95% identity to SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to SEQ ID NO: 3, wherein said ERAV strain, prior to inactivation or attenuation, is active to infect and replicate in host cells.
3. The immunogenic composition according to claim 2, wherein the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence comprising SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3.
4. An immunogenic composition comprising an ERBV strain comprising a genomic sequence whose reverse transcript has greater than 95% identity to the reverse transcript of the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829, wherein said ERBV strain, prior to inactivation or attenuation, is active to infect and replicate in host cells.
5. The immunogenic composition according to claim 4, wherein the ERBV strain comprises a genomic sequence that is the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein that has the amino acid sequence of the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829.
6. The immunogenic composition according to any one of claims 1 to 5, in which the strain of ERAV or ERBV is inactivated.

7. The immunogenic composition according to any of claims 1-5 in which the strain of ERAV or ERBV is a live, attenuated strain.
8. The immunogenic composition of claims 1-7 in which the strain, prior to inactivation or attenuation, causes detectable respiratory disease in 100% of seronegative horses upon exposure to the strain.
9. The immunogenic composition of claims 1-8 in which the strain, prior to inactivation, grows in cell culture to at least 10^8 TCID₅₀/mL.
10. The immunogenic composition of claims 1-9 in which the strain results in a serum titer of at least 1:1000.
11. The immunogenic composition of claims 1-10 in which the strain of ERAV or ERBV is ERAV/ON/05 (ATCC Accession No. PTA-11828) or ERBV strain 07-103042 (ATCC Accession PTA-11829).
12. The immunogenic composition of claim 1 in which said ERAV strain comprises a genomic sequence whose reverse transcript has greater than 95% identity to SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to SEQ ID NO: 3, wherein said ERAV strain, prior to inactivation or attenuation, is active to infect and replicate in host cells.
13. The immunogenic composition according to claim 12, wherein the ERAV strain comprises a genomic sequence whose reverse transcript has a nucleotide sequence comprising SEQ ID NO: 2 or encodes a polyprotein with an amino acid sequence of SEQ ID NO: 3.
14. The immunogenic composition of claim 1 in which said ERBV strain comprises a genomic sequence whose reverse transcript has greater than 95% identity to the reverse transcript of the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein with an amino acid sequence with greater than 95% identity to the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829, wherein said ERBV strain, prior to inactivation or attenuation, is active to infect and replicate in host cells.

15. The immunogenic composition according to claim 14, wherein the ERBV strain comprises a genomic sequence that is the genomic sequence of the ERBV strain having ATCC Accession no. PTA-11829 or encodes a polyprotein that has the amino acid sequence of the polyprotein encoded by the genome of the ERBV strain having ATCC Accession no. PTA-11829.

16. The immunogenic composition according to claim 1, comprising ERAV and ERBV, wherein the ERAV strain is ERAV/ON/05 and the ERBV strain has ATCC Accession No. PTA-11829.

17. The immunogenic composition according to any one of claims 1 to 16, wherein said immunogenic composition comprises at least one antigen or one inactivated or live, attenuated strain of Equine Herpes Virus.

18. The immunogenic composition according to claim 17, wherein said Equine Herpes Virus is selected from the group consisting of EHV-1 and EHV-4, and a combination thereof.

19. The immunogenic composition according to claim 17, wherein said Equine Herpes Virus is selected from the group consisting of. EHV-1, EHV-4, strains deposited with the ATCC under accession Nos. PTA-9525 and PTA-9526, and a combination thereof.

20. The immunogenic composition according to any one of claims 17 to 19, comprising at least one antigen of Equine Herpes Virus.

21. The immunogenic composition according to any one of claims 1 to 20, wherein said immunogenic composition comprises at least one antigen or one inactivated or live, attenuated strain of Equine Influenza Virus.

22. The immunogenic composition according to claim 21, wherein said Equine Influenza Virus is selected from the group consisting of Clade 1 viruses, Clade 2 viruses, Influenza A/South Africa/2003, Influenza A/equine-2/Ohio/03, Influenza A/equine-2/New Market/2/93, Influenza A/equine-2/Kentucky/95, Influenza A/equine-2/Richmond/1/2007, strains deposited with the ATCC under accession Nos. PTA-9522, PTA-9523, and PTA-9524, and combinations thereof.

23. The immunogenic composition according to any one of claims 21 to 22, comprising at least one antigen of Equine Influenza Virus.

24. The immunogenic composition according to any one of claims 1 to 23, wherein said immunogenic composition comprises at least one antigen or one inactivated or live, attenuated strain of Equine Herpes Virus and at least one antigen or one inactivated or live, attenuated strain of Equine Influenza Virus.

25. The immunogenic composition according to claim 24, wherein the Equine Herpes Virus is EHV-1 or EHV-4 or a combination thereof.

26. The immunogenic composition according to any one of claims 1 to 25, wherein said immunogenic composition further comprises at least one antigen of one or more strains selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, and Tetanus Toxoid, and combinations thereof.

27. The immunogenic composition according to any one of claims 1 to 25, wherein said immunogenic composition comprises at least one inactivated or live, attenuated strain selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, and Tetanus Toxoid, and combinations thereof.

28. The immunogenic composition according to any one of claims 1 to 27, wherein said immunogenic composition comprises one or more inactivated or live, attenuated strains of Eastern Equine Encephalomyelitis, Western Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis Virus, and Tetanus Toxoid.

29. The immunogenic composition according to any one of claims 1 to 28, wherein said immunogenic composition comprises one or more inactivated or live, attenuated strains of Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, Equine Influenza Virus, Equine Herpes Virus, and Tetanus Toxoid.

30. The immunogenic composition according to any one of claims 26 to 27, wherein said West Nile Virus is one of the strains selected from the group consisting of Horse Origin 2005, deposited with the ATCC under accession number PTA-9409; NAEE159, deposit at the United

States Department of Agriculture Isolate under accession number 405330; NY2002Nassau; NY2002Clinton; NY2002Queens; GA20021; GA20022; TX20021; TX20022; IN2002; NY2003Albany; NY2003Suffolk; NY2003Chataqua; CO20031; CO20032; TX2003; TX2003Harris4; TX2003Harris6; TX2003Harris7; TX2003Harris10; AZ2004; and TX2004Harris4; and combination thereof.

31. The immunogenic composition according to any one of claims 26 to 29, wherein said Western Equine Encephalomyelitis Virus is the strain deposited with the ATCC under accession number PTA-9410.

32. The immunogenic composition according to any one of claims 26 to 29, wherein said Venezuelan Equine Encephalomyelitis Virus is the strain deposited with the ATCC under accession number PTA-9411.

33. The immunogenic composition according to any one of claims 26 to 29, wherein said Eastern Equine Encephalomyelitis Virus is the strain deposited with the ATCC under accession number PTA-9412.

34. The immunogenic composition according to any one of claims 28 to 29, wherein said Equine Herpes Virus is selected from the group consisting of the strains deposited with the ATCC under accession Nos. PTA-9525 or PTA-9526, and combinations thereof.

35. The immunogenic composition according to any one of claims 1 to 34, wherein any of the strains are present in an amount from about $10^{2.0}$ TCID₅₀/mL – $10^{10.0}$ TCID₅₀/mL per dose.

36. The immunogenic composition according to any one of claims 1 to 35, wherein said immunogenic composition further comprises a suitable pharmaceutical carrier.

37. The immunogenic composition according to claim 36, wherein said suitable pharmaceutical carrier is selected from the group consisting of a diluent, adjuvant, antimicrobial agent, preservative, inactivating agent, and combinations thereof.

38. The immunogenic composition according to claim 37, wherein said adjuvant is HRA-5.

39. A method for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV or ERBV in an animal or a herd of animals comprising the step of administering the immunogenic composition according to any one of claims 1 to 38 to an animal in need thereof.

40. A method for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV and/or ERBV and, one or more of the pathogens selected from the group consisting of West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, and Clostridium tetani in an animal or a herd of animals comprising the step of administering the immunogenic composition according to any one of claims 1 to 38 to an animal in need thereof.

41. A method for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV and/or ERBV and, one or more of the pathogens selected from the group consisting of: Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, Equine Herpes Virus, and Clostridium tetani in an animal or a herd of animals comprising the step of administering the immunogenic composition according to any one of claims 1 to 38 to an animal in need thereof.

42. A method for reducing the incidence or lessening the severity of clinical symptoms associated with or caused by ERAV and/or ERBV and one or more of the pathogens selected from the group consisting of: West Nile Virus, Eastern Equine Encephalomyelitis Virus, Western Equine Encephalomyelitis Virus, Venezuelan Equine Encephalomyelitis Virus, Equine Herpes Virus, Equine Influenza Virus, and Clostridium tetani in an animal or a herd of animals, comprising the step of administering the immunogenic composition according to any one of claims 1 to 38 to an animal in need thereof.

43. The method according to any one of claims 39 to 42, wherein the incidence of clinical symptoms caused by one or more of said pathogens in a herd of animals is reduced as compared to a herd not receiving said immunogenic composition.

44. The method according to any one of claims 39 to 42, wherein the administration of at least one dose of said immunogenic composition provides a duration of immunity of at least 12 months against one or more of said pathogens.

45. The method according to any one of claims 39 to 44, wherein said animal is a horse.

46. The method according to any one of claims 39 to 45, wherein any of the strains are present in an amount from about $10^{2.0}$ TCID₅₀/mL – $10^{10.0}$ TCID₅₀/mL per dose.

47. The method according to any one of claims 39 to 46, wherein said immunogenic composition further comprises a suitable pharmaceutical carrier.

48. The method according to claim 47, wherein said suitable pharmaceutical carrier is selected from the group consisting of a diluent, adjuvant, antimicrobial agent, preservative, inactivating agent, and combinations thereof.

49. The method according to claim 48, wherein said adjuvant is HRA-5.

50. The method according to any one of claims 39 to 49, wherein said immunogenic composition is administered in one or more doses.

51. The method according to any one of claims 39 to 50, wherein one dose of said immunogenic composition is formulated in a dosage form of 0.5 mL to 2.5 mL.

52. The method according to any one of claims 39 to 51, wherein said immunogenic composition is safe for use in foals or horses 4 months of age or older.

53. A method for producing an immunogenic composition comprising one or more strains of inactivated ERAV or ERBV, the method comprising:

- a) infecting a susceptible cell line with ERAV or ERBV;
- b) growing the infected cell line in growth media until a cytopathic effect (CPE) is attained;
- c) harvesting the media;
- d) filtering the media to yield a filtered media; and

e) contacting the filtered media with an inactivating agent to obtain the inactivated ERAV or ERBV.

54. The method of claim 53 further comprising adding an adjuvant and one or more preservatives.

55. The method of claim 54 comprising infecting cells with ERAV and ERBV.


56. The method of claim 54 wherein the ERAV is ERAV/ON/05 (ATCC Accession No. PTA-11828) and the ERBV is strain 07-103042C ATCC Accession NO: PTA-11829.

57. An immunogenic composition comprising ERAV/ON/05 and/or a ERBV strain having ATCC Accession NO: PTA-11829 and Amphotericin B, gentamicin sulfate, formaldehyde, and HRA-5.

1/9

FIG. 1


Proportion of virus positive across time



2/9

FIG. 2


Proportion of buffy coat positive across time



3/9

FIG. 3


SN Titers Across Time



4/9

FIG. 4


Mean Nasal Scores Across Time



5/9

FIG. 5

Mean Ocular Scores Across Time




PERV-1	ACTTTTTA-GGAGATGACCAAAACGCAGTAACCGCCAAGCAATTGGCTGTAGCGTCAGTAAAAA	111
ERAV/ON/05	ATGCCTGT T AGCGTCAGTAAAACGCGGTAAACATAGGCT-TTGACTGTAGCGTCAGTAAAA	119
	* * * *	
PERV-1	CGCAATA--CACAAAGAT-TTGAGCCTGTAGCGTCAGTAAAACGCTGCAACCACAAGCTAT	168
ERAV/ON/05	CGCAACA A CCATACGCT <u>G</u> TTGTGCCTGTAGCGTCAGTAAAACGCGCAAACGCAAGC-AT	178
	***** * ** * * * ***** **** * ***** **	
PERV-1	TGACTGTAGCGTCAGTAAAACGCAA-----ACATTCTTGTGGCGCTCGCGTA	215
ERAV/ON/05	TAACTGTAGCGTCAGTAAAACGCAA CAACCATACGCTA ATGTGCCTGAGGCGTCAGTAAA	238
	* ***** *	
PERV-1	-GCGCTCA--AGTGCAGAGCTTCCCGGCTTTAAGGGTTACTGCTCGTAATGAGAGCACAT	272
ERAV/ON/05	C GCATACAG C AAACCAGAGCTTCCCGGCTTTAAGGGTTACTGCTCGTAATGAGAGCACTT	298
	** ** *	
PERV-1	GACATTTTGTCCAAGATTTTCCTGGCAATTGTACGGGAGAGAGGCCCGTTCTCGGGCAC	332
ERAV/ON/05	GGCAATTTGTTCAGGATTTTCCTGGTGGTTGTACGGGAGAGAGGCCCGTTTTCGGGCAC	358
	* ** ***** ***** *****	
PERV-1	TTTTCTCTCAAACAATGTTGGCGCGCCTCGGCGCGCCCCCCTTTTTTCAGCCCCCTGTCA	392
ERAV/ON/05	TGTTCC---CAACAAACATTTGTGCGCTTCGGCGCACACCCCGCT---CAGCCCCCTGTCA	413
	* *** ** * * * ***** * ***** *	

7/9

FIG. 7

Total clinical score means from control, infected, and re-infected groups



8/9

FIG. 8

Body temperature means from control, infected, and re-infected groups.




FIG. 9

Titers to equine rhinitis A virus (ERAV) and equine rhinitis B virus (ERBV) in control, infected, and re-infected groups. The virus neutralization test (VN) was used to measure antibody titers on serum samples.

Control Group					Infected Group					Re-infected Group				
Pony # 1	ERAV	ERBV	AE2	EHV 1/4	Pony # 5	ERAV	ERBV	AE2	EHV 1/4	Pony # 9	ERAV	ERBV	AE2	EHV 1/4
Day 0	<1:2	1:6	ND		Day 0	<1:2	1:12	ND		Day 0	1:1536	1:48	ND	
Day 7	<1:2	1:4	ND	NT	Day 7	1:512	1:6	ND	NT	Day 7	1:1024	1:96	ND	NT
Day 14	<1:2	1:6	ND	NT	Day 14	1:1536	1:8	ND	NT	Day 14	1:1536	1:64	ND	NT
Day 21	<1:2	1:8	ND		Day 21	1:1536	1:16	ND		Day 21	1:1536	1:48	ND	
Pony # 2	ERAV	ERBV	AE2	EHV 1/4	Pony # 6	ERAV	ERBV	AE2	EHV 1/4	Pony # 10	ERAV	ERBV	AE2	EHV 1/4
Day 0	<1:2	1:24	ND		Day 0	<1:2	1:32	ND		Day 0	1:6144	1:64	ND	
Day 7	<1:2	1:32	ND	NT	Day 7	1:256	1:24	ND	NT	Day 7	1:12288	1:24	ND	NT
Day 14	<1:2	1:32	ND	NT	Day 14	1:2048	1:16	ND	NT	Day 14	1:8192	1:32	ND	NT
Day 21	<1:2	1:24	ND		Day 21	1:2048	1:12	ND		Day 21	1:32768	1:64	ND	
Pony # 3	ERAV	ERBV	AE2	EHV 1/4	Pony # 7	ERAV	ERBV	AE2	EHV 1/4	Pony # 11	ERAV	ERBV	AE2	EHV 1/4
Day 0	<1:2	1:12	ND		Day 0	<1:2	1:4	ND		Day 0	1:1024	1:24	ND	
Day 7	<1:2	1:16	ND	NT	Day 7	1:64	1:4	ND	NT	Day 7	1:1536	1:16	ND	NT
Day 14	<1:2	1:64	ND	NT	Day 14	1:3072	1:3	ND	NT	Day 14	1:2048	1:32	ND	NT
Day 21	<1:2	1:24	ND		Day 21	1:1536	1:4	ND		Day 21	1:1536	1:24	ND	
Pony # 4	ERAV	ERBV	AE2	EHV 1/4	Pony # 8	ERAV	ERBV	AE2	EHV 1/4	Pony # 12	ERAV	ERBV	AE2	EHV 1/4
Day 0	<1:2	1:16	ND		Day 0	<1:2	1:12	ND		Day 0	1:3072	1:4	ND	
Day 7	<1:2	1:32	ND	NT	Day 7	1:256	1:16	ND	NT	Day 7	1:2048	1:3	ND	NT
Day 14	<1:2	1:64	ND	NT	Day 14	1:2048	1:16	ND	NT	Day 14	1:2048	1:6	ND	NT
Day 21	<1:2	1:24	ND		Day 21	1:2048	1:16	ND		Day 21	1:4096	1:6	ND	

ERAV

ERBV

AE2

EHV1/4

ND

NT

Equine rhinitis A virus (virus neutralization test)

Equine rhinitis B virus (virus neutralization test)

Equine influenza 2 (H3N8) (hemagglutinin inhibition test)

Equine herpesvirus 1 & 4 (virus neutralization test)

Not detectable (single radial haemolysis test)

Not tested