



US00PP13695P29

(12) **United States Plant Patent**  
**Hofmann**

(10) **Patent No.:** **US PP13,695 P2**

(45) **Date of Patent:** **Apr. 1, 2003**

(54) **NEW GUINEA IMPATIENS PLANT NAMED**  
**'FISIMP 113'**

(76) **Inventor:** **Birgit Hofmann**, Gassenweg 29, 56170  
Bendorf (DE)

(\*) **Notice:** Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **10/087,918**

(22) **Filed:** **Mar. 4, 2002**

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./318**

(58) **Field of Search** ..... **Plt./318**

*Primary Examiner*—Bruce R. Campell

*Assistant Examiner*—Susan B. McCormick

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of New Guinea Impatiens plant  
named 'Fisimp 113', characterized by its upright, outwardly  
spreading, and uniformly mounded plant habit; freely  
branching and freely flowering habit; large rounded purple-  
colored flowers with red purple-colored eye that are posi-  
tioned above and beyond the foliage; and dark green-colored  
leaves.

**1 Drawing Sheet**

**1**

**BOTANICAL CLASSIFICATION/CULTIVAR**  
**DESIGNATION**

*Impatiens hawkeri* cultivar Fisimp 113.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct culti-  
var of New Guinea Impatiens plant, botanically known as  
*Impatiens hawkeri*, and hereinafter referred to by the name  
'Fisimp 113'.

The new Impatiens is a product of a planned breeding  
program conducted by the Inventor in Hillscheid, Germany.  
The objective of the breeding program is to develop new  
vigorous Impatiens cultivars that flower relatively early with  
large rounded flowers and attractive flower colors.

The new Impatiens originated from a cross made by the  
Inventor in July, 1998 of the *Impatiens hawkeri* cultivar  
Danharras, not patented, as the female, or seed, parent with  
the *Impatiens hawkeri* cultivar Toga, disclosed in U.S. Plant  
Pat. No. 10,304, as the male, or pollen, parent. The cultivar  
Fisimp 113 was discovered and selected by the Inventor as  
a flowering plant within the progeny of the stated cross in a  
controlled environment in Moncarapacho, Portugal in  
March, 1999.

Asexual reproduction of the new cultivar by terminal  
cuttings taken in Moncarapacho, Portugal, since March,  
1999, has shown that the unique features of this new  
Impatiens are stable and reproduced true to type in succes-  
sive generations.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of 'Fisimp  
113'. These characteristics in combination distinguish  
'Fisimp 113' as a new and distinct Impatiens cultivar:

1. Upright, outwardly spreading, and uniformly mounded  
plant habit.
2. Freely branching and freely flowering habit.
3. Large rounded purple-colored flowers with a red  
purple-colored eye that are positioned above and beyond the  
foliage.
4. Dark green-colored leaves.

**2**

Plants of the new Impatiens can be compared to plants of  
the female parent, the cultivar Danharras. In side-by-side  
comparisons conducted by the Inventor in Hillscheid,  
Germany, plants of the new Impatiens differed from plants  
of the cultivar Danharras in the following characteristics:

1. Plants of the new Impatiens were taller and more  
vigorous than plants of the cultivar Danharras.

2. Flowers of plants of the new Impatiens were purple in  
color whereas flowers of plants of the cultivar Danharras  
were white and cherry red bi-colored.

Plants of the new Impatiens can be compared to plants of  
the male parent, the cultivar Toga. In side-by-side compari-  
sons conducted by the Inventor in Hillscheid, Germany,  
plants of the new Impatiens differed from plants of the  
cultivar Toga in the following characteristics:

1. Plants of the new Impatiens were taller and more  
vigorous than plants of the cultivar Toga.

2. Plants of the new Impatiens flowered about one week  
earlier than plants of the cultivar Toga.

3. Flowers of plants of the new Impatiens were darker  
purple in color than flowers of plants of the cultivar Toga.

Plants of the new Impatiens and the cultivar Fisimp 535,  
U.S. Plant patent application Ser. No. 10/080,600 filed  
concurrently, differ primarily in flower coloration. In  
addition, plants of the new Impatiens are taller than plants of  
the cultivar Fisimp 535.

Plants of the new Impatiens can also be compared to  
plants of the cultivar Kipas, disclosed in U.S. Plant Pat. No.  
10,432. In side-by-side comparisons conducted by the  
Inventor in Hillscheid, Germany, plants of the new Impa-  
tiens differed from plants of the cultivar Kipas in the  
following characteristics:

1. Plants of the new Impatiens were more vigorous, taller  
and broader than plants of the cultivar Kipas.

2. Stems and peduncles of plants of the new Impatiens  
were reddish in color whereas stems and peduncles of plants  
of the cultivar Kipas were mostly green in color and overlain  
with anthocyanin.

3. Flowers of plants of the new Impatiens were more  
reddish purple in color than flowers of plants of the cultivar  
Kipas.

## BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Impatiens. The photograph comprises a side perspective view of a typical flowering plant of 'Fisimp 113' grown in a container.

## DETAILED BOTANICAL DESCRIPTION

The cultivar Fisimp 113 has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The aforementioned photograph, following observations and measurements describe plants grown in Langley, British Columbia, Canada, under commercial practice in a greenhouse. Rooted young plants were planted in 17.5-cm containers during the spring and the aforementioned photograph and following observations and measurements were taken during the summer about 17 weeks later. During the production of the plants, day temperatures ranged from 21 to 24° C. and night temperatures were about 17 to 18° C. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar Fisimp 113.

## Parentage:

*Female parent.*—*Impatiens hawkeri* cultivar Danharras, not patented.

*Male parent.*—*Impatiens hawkeri* cultivar Toga, disclosed in U.S. Plant Pat. No. 10,304.

## Propagation:

*Type cutting.*—Terminal tip cuttings.

*Time to initiate roots.*—Summer: About 8 to 9 days at 24° C. Winter: About 10 days at 21° C.

*Time to produce a rooted cutting.*—Summer: About 15 days at 24° C. Winter: About 18 days at 21° C.

*Root description.*—Numerous, fibrous, and freely branching; 179D in color.

## Plant description:

*General appearance.*—Upright, outwardly spreading, and uniformly mounded plant growth habit; freely branching habit, dense and bushy appearance; freely flowering. Vigorous.

*Crop time.*—From a rooted cutting, about 10 to 11 weeks are required to produce finished flowering plants in 12-cm containers.

*Plant height.*—About 26.6 cm.

*Plant diameter or spread.*—About 59 cm.

*Lateral branches.*—Quantity per plant: About 11. Length: About 25 cm. Diameter: About 9 mm. Internode length: About 4.25 cm. Texture: Smooth, glabrous. Color: 185A.

*Foliage description.*—Arrangement: Primarily in whorls. Length: About 14.2 cm. Width: About 4.9 cm. Shape: Elliptic. Apex: Acuminate. Base: Acute. Margin: Serrulate with ciliation. Texture: Smooth; glabrous. Color: Young foliage, upper surface: 139A.

Young foliage, lower surface: 185B. Fully expanded foliage, upper surface: 139A. Fully expanded foliage, lower surface: 139A with random marbling, 185B to 185C. Venation, upper surface: 53A to 53B. Venation, lower surface: 53A. Petiole: Length: About 2.75 cm. Diameter: About 3.5 mm. Texture: Smooth, glabrous. Color: Upper surface: 53B. Lower surface: 53A.

## Flower description:

*Flower type and flowering habit.*—Single; large rounded purple-colored flowers; mostly flat. Freely and continuously flowering; usually about 7 to 10 flowers and flower buds per lateral branch. Flowers positioned above and beyond the foliage and typically face parallel to the leaf canopy. Petals not persistent; gynoecium persistent. Flowers not fragrant.

*Flower longevity.*—Flowers last about 8 to 9 days on the plant.

*Flowering season.*—Year-round under greenhouse conditions. Plants begin flowering about 9 weeks after planting.

*Flower buds.*—Length: About 2.2 cm. Diameter: About 1.7 cm. Shape: Ovoid. Color: 74B.

*Flower length.*—About 7.3 cm.

*Flower width.*—About 6.8 cm.

*Flower depth.*—About 1.25 cm.

*Petals.*—Quantity: Five per flower, imbricate. Length: Banner petals: About 3.3 cm. Lateral and base petals: About 3.7 cm. Width: Banner petal: About 5.1 cm. Lateral and base petals: About 3.9 cm. Shape: Roughly cordate. Apex: Emarginate. Base: Attenuate. Margin: Entire. Aspect: Mostly flat. Texture: Smooth; satiny. Color: When opening, upper surface: Between 74B and 78A. When opening, lower surface: 67A. Fully opened, upper surface: 74B; towards the base or eye, 66B, faint; color fading to 72D with subsequent development. Fully opened, lower surface: 67A.

*Spur.*—Quantity: One per flower. Length: About 4.4 cm. Diameter: At apex: About 0.5 mm. At flower: About 3 mm. Aspect: Curved. Color: 53B.

*Peduncles.*—Length: About 5 cm. Diameter: About 2 mm. Texture: Smooth, glabrous. Strength: Strong, flexible. Angle: About 45° from the lateral branch. Color: 181A.

*Reproductive organs.*—Androecium: Stamen number: Five fused at anthers, hooded; filaments free. Anther length: About 5.5 mm. Anther shape: Obovate. Anther color: 66C. Pollen amount: Moderate. Pollen color: 8D. Gynoecium: Pistil quantity: One per flower. Pistil length: About 5 mm. Stigma color: 74C. Style color: 74C. Ovary: Five-celled. Ovary color: Initially 139A, becoming 187A with development.

*Seed/fruit.*—Seed and fruit development has not been observed.

Disease/pest resistance: Plants of the new *Impatiens* have not been observed to be resistant to pathogens and pests common to *Impatiens*.

It is claimed:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Fisimp 113', as illustrated and described.

\* \* \* \* \*

