



(51) International Patent Classification:
C11D 3/386 (2006.01)

(21) International Application Number:
PCT/US2016/035629

(22) International Filing Date:
3 June 2016 (03.06.2016)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
15170745.2 4 June 2015 (04.06.2015) EP

(71) Applicant: **THE PROCTER & GAMBLE COMPANY**
[US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio
45202 (US).

(72) Inventor: **LANT, Neil, Joseph**; Whitley Road, Long-
benton, P.O. Box: Forest Hall No. 2, Newcastle upon
Tyne, Northumberland NE12 9TS (GB).

(74) Agent: **KREBS, Jay A.**; c/o The Procter & Gamble Com-
pany, Global Patent Services, One Procter & Gamble
Plaza, C8-229, Cincinnati, Ohio 45202 (US).

(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: HAND DISHWASHING LIQUID DETERGENT COMPOSITION

(57) Abstract: A hand dishwashing detergent composition comprising a surfactant system a lipase at least 0.05% by weight of the composition of at least one monovalent, divalent or trivalent cation or a mixture thereof.



WO 2016/196874 A1

HAND DISHWASHING LIQUID DETERGENT COMPOSITION

FIELD OF THE INVENTION

5

The present invention relates to a hand dishwashing detergent composition comprising a surfactant system, a specific lipase and a stabilization system. The composition provides very good cleaning, it is stable in storage and it does not have malodour issues.

10

BACKGROUND OF THE INVENTION

Improved grease cleaning is an important need for manual dishwashing detergent users. While lipase enzymes have long been proposed as potential additives to improve the grease cleaning of manual dishwashing detergents, such systems have not been successfully practised due to three key challenges of (i) slow lipase kinetics in a fast manual dishwashing process, (ii) poor enzyme stability during storage and (iii) malodours arising from the action of lipase on short-chain fatty acid residues present in the dairy soil fats. The objective of the present invention is to provide a manual dishwashing detergent that provides effective grease cleaning in short wash processes, exhibits excellent storage stability and low risk of malodour generation during product usage.

20

SUMMARY OF THE INVENTION

According to a first aspect of the invention, there is provided a hand dishwashing detergent composition. The composition is preferably in liquid form. The composition comprises a surfactant system, a lipase and a cation that acts as stabilization system for the lipase.

The lipase is a variant of a parent lipase, which variant has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 1, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T and G38A of SEQ ID NO: 1. This lipase is very good in terms of cleaning and does not present malodour issues.

The surfactant system of the detergent of the invention can comprise any cleaning surfactant. Very good grease cleaning and at the same time very good suds profile have been found when

the surfactant system comprises: i) an anionic surfactant; and ii) an amphoteric and/or zwitterionic surfactant. Preferably the weight ratio of anionic surfactant to amphoteric and/or zwitterionic surfactant is less than 9:1, more preferably less than 5:1, more preferably less than 4:1, even more preferably from about 0.5:1 to about 3.5:1 and especially from about 1:1 to about 3:1.

Preferably the surfactant system comprises an anionic surfactant, the anionic surfactant can be any anionic cleaning surfactant, especially preferred are alkoxyated anionic surfactants, more preferably an alkyl alkoxy sulphate. Preferably the alkoxyated anionic surfactant has an average alkoxylation degree of from about 0.2 to about 3, preferably of from from about 0.3 to 2, most preferably from about 0.5 to 1. Also preferred are branched anionic surfactants having a weight average level of branching of from about 5% to about 40%.

Another preferred surfactant system for use herein is an anionic and amphoteric/zwitterionic system in which the amphoteric to zwitterionic weight ratio is preferably from about 2:1 to about 1:2, more preferably from about 1.5:1 to about 1:1.5. In particular a system in which the amphoteric surfactant is an amine oxide surfactant and the zwitterionic surfactant is a betaine and the weight ratio of the amine oxide to the betaine is about 1:1. Preferably the amine oxide is C12-14 alkyl dimethyl amine oxide, coco-alkyl dimethyl amine oxide or coco-alkyl amidopropyl dimethyl amine oxide (CAP dimethyl amine oxide). Preferably the betaine is coco-alkyl amidopropyl betaine (CAP-betaine).

Also preferred for use herein are surfactant systems comprising non-ionic surfactants. Preferably the non-ionic surfactant is an ethoxylated alcohol surfactant.

Especially preferred surfactant systems for the composition of the invention comprise an anionic surfactant preferably selected from the group consisting of alkyl sulphate, alkyl alkoxy sulphate and mixtures thereof, more preferably an alkoxyated sulphate, even more preferably an ethoxylated alkyl sulphate, and an amphoteric and/or zwitterionic surfactant, preferably an amino oxide and/or betaine surfactant, and a non-ionic surfactant, preferably an ethoxylated alcohol nonionic surfactant. In summary, the most preferred surfactant system for use herein comprises an ethoxylated alkyl sulfate surfactant, amine oxide and/or betaine, and ethoxylated alcohol non-ionic surfactant.

According to the second aspect of the invention, there is provided a method of manual dishwashing comprising the step of: delivering the detergent composition of the invention to a volume of water and immersing soiled dishware in the water. When the composition of the invention is used according to this method good and fast cleaning is achieved without malodours associated to it.

For the purpose of this invention “dishware” herein includes cookware and tableware.

According to the last aspect of the invention, there is provided a method of manual dishwashing comprising the step of: delivering the detergent composition of the invention directly onto dishware or onto a cleaning implement and using the cleaning implement to clean the dishware. Preferably the cleaning implement is a sponge and more preferably the sponge is wet. When the composition of the invention is used according to this method good and fast cleaning is achieved without malodours associated to it.

DETAILED DESCRIPTION OF THE INVENTION

The present invention envisages a hand dishwashing detergent composition. Preferably in liquid form. The detergent composition comprises a surfactant system, a lipase and a cation. It provides very good cleaning, especially grease cleaning even on plastic substrates that are the toughest substrates for grease removal.

The detergent composition

The detergent composition is a hand dishwashing detergent, preferably in liquid form. It typically contains from 30% to 95%, preferably from 40% to 90%, more preferably from 50% to 85% by weight of a liquid carrier in which the other essential and optional components are dissolved, dispersed or suspended. One preferred component of the liquid carrier is water.

Preferably the pH of the detergent is adjusted to between 4 and 13, preferably between 6 and 12 and most preferably between 8 and 10. The pH of the detergent can be adjusted using pH modifying ingredients known in the art.

Lipase

Preferably the lipase is present in the composition of the invention in a level of from 0.001-2%, more preferably from 0.005 to 1.5 and especially from 0.01 to 1% of pure enzyme, by weight of the composition.

5

Preferred lipase for use herein includes lipases in which the variant comprises substitutions of SEQ ID NO: 1 selected from the group consisting of:

a) D96E+T231R+N233R;

b) N33Q+D96E+T231R+N233R;

10

c) N33Q+D111A+T231 R+N233R;

d) N33Q+T231 R+N233R+P256T;

e) N33Q+G38A+G91T+G163K+T231R+N233R+D254S;

f) N33Q+G38A+G91T+D96E+D111A+G163K+T231R+N233R+D254S+P256T;

g) D27R+N33Q+G38A+D96E+D111A+G163K+T231R+N233R+D254S+P256T;

15

h) D27R+N33Q+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+P256T;

i) D27R+N33Q+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S;

j) D27R+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;

k) D96E+T231 R+N233R+D254S;

l) T231R+N233R+D254S+P256T;

20

m) G163K+T231 R+N233R+D254S;

n) D27R+N33Q+G38A+G91T+D96E+G163K+T231 R+N233R+D254S+P256T;

o) D27R+G91T+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;

- p) D96E+G163K+T231 R+N233R+D254S;
- q) D27R+G163K+T231 R+N233R+D254S;
- r) D27R+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S;
- s) D27R+G38A+G91T+D96E+G163K+T231 R+N233R+D254S+P256T;
- 5 t) D27R+G38A+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;
- u) D27R+D96E+G163K+T231R+N233R+D254S;
- v) D27R+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- w) D27R+G38A+D96E+G163K+T231 R+N233R+D254S+P256T;
- x) D111A+G163K+T231R+N233R+D254S+P256T;
- 10 y) D111A+T231R+N233R;
- z) D111A+T231 R+N233R+D254S+P256T;
- aa) D27R+D96E+D111A+G163K+T231R+N233R;
- bb) D27R+D96E+D111A+T231 R+N233R;
- cc) D27R+G38A+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- 15 dd) D27R+N33Q+G38A+D96E+D111A+T231R+N233R+D254S+P256T;
- ee) D27R+G38A+D96E+D111A+G163K+E210Q+T231 R+N233R+D254S+P256T;
- ff) D27R+T231 R+N233R+D254S+P256T;
- gg) D96E+D111A+G163K+T231R+N233R;
- hh) D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- 20 ii) D96E+D111A+G163K+T231R+N233R+P256T;

jj) D96E+D111A+T231R+N233R;

kk) D96E+D111A+T231R+N233R+D254S;

ll) D96E+D111A+T231R+N233R+D254S+P256T;

mm) D96E+D111A+T231R+N233R+P256T;

5 nn) D96E+G163K+T231R+N233R+D254S+P256T;

oo) D96E+T231R+N233R+D254S+P256T;

pp) D96E+T231R+N233R+P256T;

qq) G38A+D96E+D111A+T231 R+N233R;

rr) G91T+D96E+D111A+G163K+T231R+N233R+D254S+P256T;

10 ss) G91T+D96E+D111A+T231 R+N233R;

tt) G91T+D96E+T231 R+N233R;

uu) G91T+T231 R+N233R+D254S+P256T;

vv) N33Q+D96E+D111A+G163K+T231R+N233R+D254S+P256T;

ww) T231R+N233R+D254S+P256T; and

15 xx) T231R+N233R+P256T.

Additional enzymes

Additional enzyme(s) which may be comprised in the detergent composition include one or more enzymes such as protease, cutinase, amylase, carbohydrase, cellulase, pectinase, mannanase, 20 arabinase, galactanase, xylanase, perhydrolase, oxidase, e.g. laccase, and/or peroxidase.

A preferred combination of enzymes comprises, e.g., a protease, lipase and amylase. When present in a composition, the aforementioned additional enzymes may be present at levels from

0.00001 to 2wt%, from 0.0001 to 1wt% or from 0.001 to 0.5wt% enzyme protein by weight of the composition.

Lyases: The lyase may be a pectate lyase derived from *Bacillus*, particularly *B. licheniformis* or *B. agaradhaerens*, or a variant derived of any of these, e.g. as described in US 6124127, WO 99/27083, WO 99/27084, WO 02/006442, WO 02/092741 , WO 03/095638, Commercially available pectate lyases are XPect™; Pectawash™ and Pectaway™ (Novozymes A/S).

Mannanases: Suitable mannanases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. The mannanase may be an alkaline mannanase of Family 5 or 26. It may be a wild-type from *Bacillus* or *Humicola*, particularly *B. agaradhaerens*, *B. licheniformis*, *B. halodurans*, *B. clausii*, or *H. insolens*. Suitable mannanases are described in WO 1999/064619. A commercially available mannanase is Mannaway™ (Novozymes A/S).

Proteases: Suitable proteases include those of bacterial, fungal, plant, viral or animal origin e.g. vegetable or microbial origin. Microbial origin is preferred. Chemically modified or protein engineered mutants are included. It may be an alkaline protease, such as a serine protease or a metalloprotease. A serine protease may for example be of the S1 family, such as trypsin, or the S8 family such as subtilisin. A metalloproteases protease may for example be a thermolysin from e.g. family M4 or other metalloprotease such as those from M5, M7 or M8 families.

The term "subtilases" refers to a sub-group of serine protease according to Siezen et al., 1991, Protein Engng. 4: 719-737 and Siezen et al., 1997, Protein Science 6: 501-523. Serine proteases are a subgroup of proteases characterized by having a serine in the active site, which forms a covalent adduct with the substrate. The subtilases may be divided into 6 sub-divisions, i.e. the Subtilisin family, the Thermitase family, the Proteinase K family, the Lantibiotic peptidase family, the Kexin family and the Pyrolysin family.

Examples of subtilases are those derived from *Bacillus* such as *Bacillus lentus*, *B.*

alkalophilus, *B. subtilis*, *B. amyloliquefaciens*, *Bacillus pumilus* and *Bacillus gibsonii* described in; US 7,262,042 and WO 2009/021867, and subtilisin lentus, subtilisin Novo, subtilisin

Carlsberg, *Bacillus licheniformis*, subtilisin BPN', subtilisin 309, subtilisin 147 and subtilisin 168 described in WO 89/06279 and protease PD138 described in (WO 93/18140). Other useful proteases may be those described in WO 92/175177, WO 01/16285, WO 02/026024 and WO 02/016547. Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the *Fusarium* protease described in WO 89/06270, WO 94/25583 and WO 2005/040372, and the chymotrypsin proteases derived from *Cellulomonas* described in WO 2005/052161 and WO 2005/052146.

A further preferred protease is the alkaline protease from *Bacillus lentus* DSM 5483, as described for example in WO 95/23221, and variants thereof which are described in WO 92/21760, WO 95/23221, EP 1921 147 and EP 1921 148.

Examples of metalloproteases are the neutral metalloprotease as described in WO 2007/044993 (Genencor Int.) such as those derived from *Bacillus amyloliquefaciens*.

Examples of useful proteases are the variants described in: W092/19729, W096/034946, WO98/201 15, WO98/201 16, WO99/01 1768, WO01/44452, WO03/006602, WO2004/03186, WO2004/041979, WO2007/006305, WO201 1/036263, WO201 1/036264, especially the variants with substitutions in one or more of the following positions: 3, 4, 9, 15, 27, 36, 57, 68, 76, 87, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 106, 118, 120, 123, 128, 129, 130, 160, 167, 170, 194, 195, 199, 205, 206, 217, 218, 222, 224, 232, 235, 236, 245, 248, 252 and 274 using the BPN' numbering. More preferred the subtilase variants may comprise the mutations: S3T, V4I, S9R, A15T, K27R, *36D, V68A, N76D, N87S,R, *97E, A98S, S99G,D,A, S99AD, S101 G,M,R, S103A, V104I,Y,N, S106A, G1 18V,R, H120D,N, N123S, S128L, P129Q, S130A, G160D, Y167A, R170S, A194P, G195E, V199M, V205I, L217D, N218D, M222S, A232V, K235L, Q236H, Q245R, N252K, T274A (using BPN' numbering).

Suitable commercially available protease enzymes include those sold under the trade names Alcalase™, Duralase™, Durazym™, Relase™, Relase™ Ultra, Savinase™, Savinase™ Ultra, Primase™, Polarzyme™, Kannase™, Liquanase™, Liquanase™ Ultra, Ovozyme™, Coronase™, Coronase™ Ultra, Neutrase™, Everlase™ and Esperase™ (Novozymes A/S), those sold under the tradename Maxatase™, Maxacal™, Maxapem™, Purafect™, Purafect Prime™, Preferenz™, Purafect MA™, Purafect Ox™, Purafect OxP™, Puramax™, Properase™, Effectenz™, FN2™, FN3™, FN4™, Excellase™, Opticlean™ and Optimase™ (Danisco/DuPont), Axapem™ (Gist-Brocades N.V.), BLAP (sequence shown in Figure 29 of US5352604) and variants hereof (Henkel AG) and KAP {*Bacillus alkalophilus subtilisin*) from Kao.

Amylases: Suitable amylases include alpha-amylases and/or glucoamylases and may be of bacterial or fungal origin. Chemically modified or protein engineered mutants are included. Amylases include, for example, alpha-amylases obtained from *Bacillus*, e.g., a special strain of *Bacillus licheniformis*, described in more detail in GB 1,296,839.

Suitable amylases include amylases having SEQ ID NO: 2 in WO 95/10603 or variants having 90% sequence identity to SEQ ID NO: 3 thereof. Preferred variants are described in WO 94/02597, WO 94/18314, WO 97/43424 and SEQ ID NO: 4 of WO 99/019467, such as variants

with substitutions in one or more of the following positions: 15, 23, 105, 106, 124, 128, 133, 154, 156, 178, 179, 181, 188, 190, 197, 201, 202, 207, 208, 209, 211, 243, 264, 304, 305, 391, 408, and 444.

Different suitable amylases include amylases having SEQ ID NO: 6 in WO 02/010355 or
 5 variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a deletion in positions 181 and 182 and a substitution in position 193. Other amylases which are suitable are hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues 36-483 of the *B. licheniformis* alpha-amylase shown in SEQ ID NO: 4 of WO
 10 2006/066594 or variants having 90% sequence identity thereof. Preferred variants of this hybrid alpha-amylase are those having a substitution, a deletion or an insertion in one or more of the following positions: G48, T49, G107, H156, A181, N190, M197, I201, A209 and Q264. Most preferred variants of the hybrid alpha-amylase comprising residues 1-33 of the alpha-amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 6 of WO 2006/066594 and residues
 15 36-483 of SEQ ID NO: 4 are those having the substitutions:

M197T;

H156Y+A181T+N190F+A209V+Q264S; or

G48A+T49I+G107A+H156Y+A181T+N190F+I201F+A209V+Q264S.

Further amylases which are suitable are amylases having SEQ ID NO: 6 in WO99/019467 or
 20 variants thereof having 90% sequence identity to SEQ ID NO: 6. Preferred variants of SEQ ID NO: 6 are those having a substitution, a deletion or an insertion in one or more of the following positions: R181, G182, H183, G184, N195, I206, E212, E216 and K269. Particularly preferred amylases are those having deletion in positions R181 and G182, or positions H183 and G184.

Additional amylases which can be used are those having SEQ ID NO: 1, SEQ ID NO: 3, SEQ
 25 ID NO: 2 or SEQ ID NO: 7 of WO 96/023873 or variants thereof having 90% sequence identity to SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7. Preferred variants of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 7 are those having a substitution, a deletion or an insertion in one or more of the following positions: 140, 181, 182, 183, 184, 195, 206, 212, 243, 260, 269, 304 and 476, using SEQ ID 2 of WO 96/023873 for numbering. More
 30 preferred variants are those having a deletion in two positions selected from 181, 182, 183 and 184, such as 181 and 182, 182 and 183, or positions 183 and 184. Most preferred amylase variants of SEQ ID NO: 1, SEQ ID NO: 2 or SEQ ID NO: 7 are those having a deletion in

positions 183 and 184 and a substitution in one or more of positions 140, 195, 206, 243, 260, 304 and 476.

Other amylases which can be used are amylases having SEQ ID NO: 2 of WO

08/153815, SEQ ID NO: 10 in WO 01/66712 or variants thereof having 90% sequence identity to SEQ ID NO: 2 of WO 08/153815 or 90% sequence identity to SEQ ID NO: 10 in WO 01/66712. Preferred variants of SEQ ID NO: 10 in WO 01/66712 are those having a substitution, a deletion or an insertion in one of more of the following positions: 176, 177, 178, 179, 190, 201, 207, 211 and 264.

Further suitable amylases are amylases having SEQ ID NO: 2 of WO 09/061380 or variants having 90% sequence identity to SEQ ID NO: 2 thereof. Preferred variants of SEQ ID NO: 2 are those having a truncation of the C-terminus and/or a substitution, a deletion or an insertion in one of more of the following positions: Q87, Q98, S125, N128, T131, T165, K178, R180, S181, T182, G183, M201, F202, N225, S243, N272, N282, Y305, R309, D319, Q320, Q359, K444 and G475. More preferred variants of SEQ ID NO: 2 are those having the substitution in one of more of the following positions: Q87E,R, Q98R, S125A, N128C, T131 I, T165I, K178L, T182G, M201 L, F202Y, N225E,R, N272E,R, S243Q,A,E,D, Y305R, R309A, Q320R, Q359E, K444E and G475K and/or deletion in position R180 and/or S181 or of T182 and/or G183. Most preferred amylase variants of SEQ ID NO: 2 are those having the substitutions:

N128C+K178L+T182G+Y305R+G475K;
N128C+K178L+T182G+F202Y+Y305R+D319T+G475K;
S125A+N128C+K178L+T182G+Y305R+G475K; or
S125A+N128C+T131 I+T165I+K178L+T182G+Y305R+G475K

wherein the variants are C-terminally truncated and optionally further comprises a substitution at position 243 and/or a deletion at position 180 and/or position 181.

Further suitable amylases are amylases having SEQ ID NO: 1 of W013184577 or variants having 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one of more of the following positions: K176, R178, G179, T180, G181, E187, N192, M199, I203, S241, R458, T459, D460, G476 and G477. More preferred variants of SEQ ID NO: 1 are those having the substitution in one of more of the following positions: K176L, E187P, N192FYH, M199L, I203YF, S241 QADN, R458N, T459S, D460T, G476K and G477K and/or deletion in position R178 and/or S179 or of T180

and/or G181 . Most preferred amylase variants of SEQ ID NO: 1 are those having the substitutions:

E187P+I203Y+G476K

E187P+I203Y+R458N+T459S+D460T+G476K

- 5 wherein the variants optionally further comprises a substitution at position 241 and/or a deletion at position 178 and/or position 179.

Further suitable amylases are amylases having SEQ ID NO: 1 of W010104675 or variants having 90% sequence identity to SEQ ID NO: 1 thereof. Preferred variants of SEQ ID NO: 1 are those having a substitution, a deletion or an insertion in one of more of the following positions:

- 10 N21 , D97, V128 K177, R179, S180, 1181 , G182, M200, L204, E242, G477 and G478. More preferred variants of SEQ ID NO: 1 are those having the substitution in one of more of the following positions: N21 D, D97N, V128I K177L, M200L, L204YF, E242QA, G477K and G478K and/or deletion in position R179 and/or S180 or of 1181 and/or G182. Most preferred amylase variants of SEQ I D NO: 1 are those having the substitutions:

- 15 N21D+D97N+V128I

wherein the variants optionally further comprises a substitution at position 200 and/or a deletion at position 180 and/or position 181.

Other suitable amylases are the alpha-amylase having SEQ ID NO: 12 in WO01/66712 or a variant having at least 90% sequence identity to SEQ ID NO: 12. Preferred amylase variants are

- 20 those having a substitution, a deletion or an insertion in one of more of the following positions of SEQ ID NO: 12 in WO01/66712: R28, R1 18, N174; R181, G182, D183, G184, G186, W189, N195, M202, Y298, N299, K302, S303, N306, R310, N314; R320, H324, E345, Y396, R400, W439, R444, N445, K446, Q449, R458, N471 , N484. Particular preferred amylases include variants having a deletion of D183 and G184 and having the substitutions R1 18K, N195F, 25 R320K and R458K, and a variant additionally having substitutions in one or more position selected from the group: M9, G149, G182, G186, M202, T257, Y295, N299, M323, E345 and A339, most preferred a variant that additionally has substitutions in all these positions.

Other examples are amylase variants such as those described in WO201 1/098531 , WO2013/001078 and WO2013/001087.

- 30 Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™, Stainzyme™, Stainzyme Plus™, Natalase™, Liquozyme X™ and BAN™ (from Novozymes A S), and Rapidase™, Purastar™/Effectenz™, Powerase™, Preferenz S1000™, Preferenz S100™ and Preferenz S1 10™ (from Genencor International Inc./DuPont).

At least one cation

The “at least one cation” of the invention acts as a lipase stabilizing system. The composition of the invention comprises at least 0.05%, preferably at least 0.15%, more preferably at least 0.25% and most preferably at least 0.35% by weight of the composition of at least one monovalent, divalent or trivalent cation or a mixture thereof. The composition preferably comprises from 0.35 to 4%, more preferably from 0.35 to 3%, more preferably from 0.35 to 2% and especially from 0.35 to 1% by weight of the composition of the at least one cation.

Preferably, the cation source is selected from the inorganic or organic salts of alkali metals, alkaline earth metals, of aluminum, iron, copper and zinc, preferably of the alkali metals and alkaline earth metals, preferably selected from the halides, sulphates, sulphites, carbonates, bicarbonates, phosphates, nitrates, nitrites, phosphates, formates, acetates, propionates, citrates, malates, tartrates, succinates, oxalates, lactates, and mixtures thereof.

More preferably, the cation source is selected from sodium chloride, calcium chloride, potassium chloride, sodium sulfate, potassium sulfate, sodium acetate, potassium acetate, sodium formate, potassium formate, and mixtures thereof; more preferably the cation source is selected from calcium chloride, potassium chloride, potassium sulfate, sodium acetate, potassium acetate, sodium formate and potassium formate, and mixtures thereof and in particular from potassium chloride, potassium sulfate, potassium acetate, potassium formate, and mixtures thereof.

Surfactant system

The liquid detergent can comprise from about 1% to about 50%, preferably from about 5% to about 40% more preferably from about 8% to about 35% by weight thereof of a surfactant system. The surfactant system preferably comprises an anionic surfactant, more preferably an alkoxylated sulfate anionic surfactant. Most preferably the system further comprises an amphoteric and/or zwitterionic surfactant, and optionally a non-ionic surfactant.

Preferably, the anionic surfactant system comprises alkyl sulfates and/or alkyl ethoxy sulfates; more preferably a combination of alkyl sulfates and/or alkyl ethoxy sulfates with a combined

average ethoxylation degree of less than 5, preferably from about 0.2 to about 3, more preferably from about 0.3 to about 2, even more preferably from 0.5 to about 1. Preferably the anionic surfactant system has an average level of branching of from about 5% to about 40%.

5 Preferably, the composition of the present invention will further comprise amphoteric and/or zwitterionic surfactant, more preferably an amine oxide and/or betaine surfactant. The composition can comprise from about 0.01% to about 25%wt, preferably from about 0.2% to about 20%wt, more preferably from about 0.5% to about 15% by weight of the composition of amphoteric and/or zwitterionic surfactant, preferably amine oxide and/or betaine surfactant.

10

The composition can further comprise a nonionic surfactant, preferably an alkoxylated alcohol nonionic surfactant, even more preferably an ethoxylated nonionic surfactant.

The most preferred surfactant system for the detergent composition of the present invention will therefore comprise: (1) 1% to 40%, preferably 6% to 32%, more preferably 8% to 25% weight of the total composition of an anionic surfactant, preferably an alkoxylated sulfate surfactant (2) combined with 0.01% to 25%wt, preferably from 0.2% to 20%wt, more preferably from 0.5% to 15% by weight of the composition of amphoteric and/or zwitterionic surfactant, more preferably an amphoteric and even more preferred an amine oxide surfactant. It has been found that such surfactant system in combination with the lipase will provide the excellent cleaning required from a hand dishwashing detergent.

Anionic surfactant

Anionic surfactants include, but are not limited to, those surface-active compounds that contain an organic hydrophobic group containing generally 8 to 22 carbon atoms or generally 8 to 18 carbon atoms in their molecular structure and at least one water-solubilizing group preferably selected from sulfonate, sulfate, and carboxylate so as to form a water-soluble compound. Usually, the hydrophobic group will comprise a C 8-C 22 alkyl, or acyl group. Such surfactants are employed in the form of water-soluble salts and the salt-forming cation usually is selected from sodium, potassium, ammonium, magnesium and mono-, di- or tri-C 2-C 3 alkanolammonium, with the sodium, cation being the usual one chosen.

The anionic surfactant can be a single surfactant but usually it is a mixture of anionic surfactants. Preferably the anionic surfactant comprises a sulphate surfactant, more preferably a sulphate surfactant selected from the group consisting of alkyl sulphate, alkyl alkoxy sulphate and mixtures thereof. Preferred alkyl alkoxy sulphates for use herein are alkyl ethoxy sulphates.

5

Preferably the anionic surfactant is alkoxylation, more preferably, an alkoxylation branched anionic surfactant having an alkoxylation degree of from about 0.1 to about 4, even more preferably from about 0.2 to about 3, even more preferably from about 0.3 to about 2 and especially from about 0.5 to about 1. Preferably, the alkoxy group is ethoxy. When the branched anionic surfactant is a mixture of surfactants, the alkoxylation degree is the weight average alkoxylation degree of all the components of the mixture (weight average alkoxylation degree). In the weight average alkoxylation degree calculation the weight of anionic surfactant components not having alkoxylation groups should also be included.

10

Weight average alkoxylation degree = $(x_1 * \text{alkoxylation degree of surfactant 1} + x_2 * \text{alkoxylation degree of surfactant 2} + \dots) / (x_1 + x_2 + \dots)$

15

wherein $x_1, x_2, \dots$ are the weights in grams of each anionic surfactant of the mixture and alkoxylation degree is the number of alkoxy groups in each anionic surfactant.

Preferably the anionic surfactant to be used in the detergent of the present invention is a branched anionic surfactant having a level of branching of from about 5% to about 40%, preferably from about 10 to about 35% and more preferably from about 20% to about 30%. Preferably, the branching group is an alkyl. Typically, the alkyl is selected from methyl, ethyl, propyl, butyl, pentyl, cyclic alkyl groups and mixtures thereof. Single or multiple alkyl branches could be present on the main hydrocarbyl chain of the starting alcohol(s) used to produce the anionic surfactant used in the detergent of the invention. Most preferably the branched anionic surfactant is selected from alkyl sulphates, alkyl ethoxy sulphates, and mixtures thereof.

20

25

The branched anionic surfactant can be a single anionic surfactant or a mixture of anionic surfactants. In the case of a single surfactant the percentage of branching refers to the weight percentage of the hydrocarbyl chains that are branched in the original alcohol from which the surfactant is derived.

30

In the case of a surfactant mixture the percentage of branching is the weight average and it is defined according to the following formula:

Weight average of branching (%) = $[(x_1 * \text{wt\% branched alcohol 1 in alcohol 1} + x_2 * \text{wt\% branched alcohol 2 in alcohol 2} + \dots) / (x_1 + x_2 + \dots)] * 100$

5 wherein $x_1, x_2, \dots$ are the weight in grams of each alcohol in the total alcohol mixture of the alcohols which were used as starting material for the anionic surfactant for the detergent of the invention. In the weight average branching degree calculation the weight of anionic surfactant components not having branched groups should also be included.

10 Preferably, the anionic surfactant system comprises an alkyl ethoxylated sulphate having an average ethoxylation degree of from about 0.2 to about 3 and preferably a level of branching of from about 5% to about 40%.

Sulphate Surfactants

15 Suitable sulphate surfactants for use herein include water-soluble salts of C8-C18 alkyl or hydroxyalkyl, sulphate and/or ether sulfate. Suitable counterions include alkali metal cation or ammonium or substituted ammonium, but preferably sodium.

The sulphate surfactants may be selected from C8-C18 primary, branched chain and random alkyl sulphates (AS); C8-C18 secondary (2,3) alkyl sulphates; C8-C18 alkyl alkoxy sulphates (AExS) wherein preferably x is from 1-30 in which the alkoxy group could be selected from
20 ethoxy, propoxy, butoxy or even higher alkoxy groups and mixtures thereof.

Alkyl sulfates and alkyl alkoxy sulfates are commercially available with a variety of chain lengths, ethoxylation and branching degrees. Commercially available sulphates include, those
25 based on Neodol alcohols ex the Shell company, Lial – Isalchem and Safol ex the Sasol company, natural alcohols ex The Procter & Gamble Chemicals company.

Preferably, the branched anionic surfactant comprises at least 50%, more preferably at least 60% and especially at least 70% of a sulphate surfactant by weight of the branched anionic surfactant.
30 Especially preferred detergents from a cleaning view point are those in which the branched anionic surfactant comprises more than 50%, more preferably at least 60% and especially at least 70% by weight thereof of sulphate surfactant and the sulphate surfactant is selected from the group consisting of alkyl sulphate, alkyl ethoxy sulphates and mixtures thereof. Even more

preferred are those in which the branched anionic surfactant has a degree of ethoxylation of from about 0.2 to about 3, more preferably from about 0.3 to about 2, even more preferably from about 0.4 to about 1.5, and especially from about 0.5 to about 1 and even more preferably when the anionic surfactant has a level of branching of from about 10% to about 35%, %, more preferably from about 20% to 30%.

Sulphonate Surfactants

Suitable sulphonate surfactants for use herein include water-soluble salts of C8-C18 alkyl or hydroxyalkyl sulphonates; C11-C18 alkyl benzene sulphonates (LAS), modified alkylbenzene sulphonate (MLAS) as discussed in WO 99/05243, WO 99/05242, WO 99/05244, WO 99/05082, WO 99/05084, WO 99/05241, WO 99/07656, WO 00/23549, and WO 00/23548; methyl ester sulphonate (MES); and alpha-olefin sulphonate (AOS). Those also include the paraffin sulphonates may be monosulphonates and/or disulphonates, obtained by sulphonating paraffins of 10 to 20 carbon atoms. The sulfonate surfactant also include the alkyl glyceryl sulphonate surfactants.

Nonionic surfactants

Nonionic surfactant, when present, is comprised in a typical amount of from 0.1% to 30%, preferably 0.2% to 20%, more preferably 0.3% to 10%, most preferably 0.5-5% by weight of the composition. Suitable nonionic surfactants include the condensation products of aliphatic alcohols with from 1 to 25 moles of ethylene oxide. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 8 to 22 carbon atoms. Particularly preferred are the condensation products of alcohols having an alkyl group containing from 10 to 18 carbon atoms, preferably from 10 to 15 carbon atoms with from 2 to 18 moles, preferably 2 to 15, more preferably 5-12 of ethylene oxide per mole of alcohol. Highly preferred nonionic surfactants are the condensation products of guerbet alcohols with from 2 to 18 moles, preferably 2 to 15, more preferably 5-12 of ethylene oxide per mole of alcohol.

An alternative nonionic surfactant could be selected from the group of alkyl polyglucoside surfactants (APG's).

Amphoteric surfactant

Preferred amine oxides are alkyl dimethyl amine oxide or alkyl amido propyl dimethyl amine oxide, more preferably alkyl dimethyl amine oxide and especially coco dimethyl amino oxide.

Amine oxide may have a linear or branched alkyl moiety. Typical amine oxides include water-soluble amine oxides containing one R¹ C8-18 alkyl moiety and 2 R² and R³ moieties selected from the group consisting of C1-3 alkyl groups and C1-3 hydroxyalkyl groups. Preferably amine oxide is characterized by the formula R¹ – N(R²)(R³) O wherein R¹ is a C8-18 alkyl and R² and R³ are selected from the group consisting of methyl, ethyl, propyl, isopropyl, 2-hydroxyethyl, 2-hydroxypropyl and 3-hydroxypropyl. The linear amine oxide surfactants in particular may include linear C10-C18 alkyl dimethyl amine oxides and linear C8-C12 alkoxy ethyl dihydroxy ethyl amine oxides. Preferred amine oxides include linear C10, linear C10-C12, and linear C12-C14 alkyl dimethyl amine oxides. The amine oxide further comprises two moieties R² and R³, independently selected from a C1-3 alkyl, a C1-3 hydroxyalkyl group, or a polyethylene oxide group containing an average of from about 1 to about 3 ethylene oxide groups. Preferably the two moieties are selected from a C1-3 alkyl, more preferably both are selected as a C1 alkyl.

Zwitterionic surfactant

Other suitable surfactants include betaines, such as alkyl betaines, alkylamidobetaine, amidazoliniumbetaine, sulfobetaine (INCI Sultaines) as well as the Phosphobetaine and preferably meets formula I:

$R^1-[CO-X(CH_2)_n]_x-N^+(R^2)(R^3)-(CH_2)_m-[CH(OH)-CH_2]_y-Y$ (I) wherein

R¹ is a saturated or unsaturated C6-22 alkyl residue, preferably C8-18 alkyl residue, in particular a saturated C10-16 alkyl residue, for example a saturated C12-14 alkyl residue;

X is NH, NR⁴ with C1-4 Alkyl residue R⁴, O or S,

n a number from 1 to 10, preferably 2 to 5, in particular 3,

x 0 or 1, preferably 1,


R², R³ are independently a C1-4 alkyl residue, potentially hydroxy substituted such as a hydroxyethyl, preferably a methyl.

m a number from 1 to 4, in particular 1, 2 or 3,

y 0 or 1 and

Y is COO, SO₃, OPO(OR⁵)O or P(O)(OR⁵)O, whereby R⁵ is a hydrogen atom H or a C1-4 alkyl residue.

Preferred betaines are the alkyl betaines of the formula (Ia), the alkyl amido propyl betaine of the formula (Ib), the Sulfo betaines of the formula (Ic) and the Amido sulfobetaine of the formula (Id);



- 5 meaning as in formula I. Particularly preferred betaines are the Carbobetaine [wherein $Y^-=COO^-$], in particular the Carbobetaine of the formula (Ia) and (Ib), more preferred are the Alkylamidobetaine of the formula (Ib).

- Examples of suitable betaines and sulfobetaine are the following [designated in accordance with
- 10 INCI]: Almondamidopropyl of betaines, Apricotamidopropyl betaines, Avocamidopropyl of betaines, Babassuamidopropyl of betaines, Behenamidopropyl betaines, Behenyl of betaines, betaines, Canolamidopropyl betaines, Capryl/Capram idopropyl betaines, Carnitine, Cetyl of betaines, Cocamidoethyl of betaines, Cocamidopropyl betaines, Cocamidopropyl Hydroxysultaine, Coco betaines, Coco Hydroxysultaine, Coco/Oleamidopropyl betaines, Coco
- 15 Sultaine, Decyl of betaines, Dihydroxyethyl Oleyl Glycinate, Dihydroxyethyl Soy Glycinate, Dihydroxyethyl Stearyl Glycinate, Dihydroxyethyl Tallow Glycinate, Dimethicone Propyl of PG-betaines, Erucamidopropyl Hydroxysultaine, Hydrogenated Tallow of betaines, Isostearam idopropyl betaines, Lauramidopropyl betaines, Lauryl of betaines, Lauryl Hydroxysultaine, Lauryl Sultaine, Milkamidopropyl betaines, Minkamidopropyl of betaines, Myristamidopropyl
- 20 betaines, Myristyl of betaines, Oleamidopropyl betaines, Oleamidopropyl Hydroxysultaine, Oleyl of betaines, Olivamidopropyl of betaines, Palmamidopropyl betaines, Palm itamidopropyl betaines, Palmitoyl Carnitine, Palm Kernelamidopropyl betaines, Polytetrafluoroethylene Acetoxypentyl of betaines, Ricinoleamidopropyl betaines, Sesamidopropyl betaines, Soyamidopropyl betaines, Stearamidopropyl betaines, Stearyl of betaines,
- 25 Tallowamidopropyl betaines, Tallowamidopropyl Hydroxysultaine, Tallow of betaines, Tallow Dihydroxyethyl of betaines, Undecylenamidopropyl betaines and Wheat Germamidopropyl betaines.

A preferred betaine is, for example, Cocoamidopropylbetain.

- 30 The detergent composition herein may comprise a number of optional ingredients such as builders, chelants, conditioning polymers, cleaning polymers, surface modifying polymers, soil flocculating polymers, structurants, emmollients, humectants, skin rejuvenating actives, carboxylic acids, scrubbing particles, bleach and bleach activators, perfumes, pigments, dyes,

opacifiers, beads, pearlescent particles, microcapsules, diamines, antibacterial agents, preservatives and pH adjusters and buffering means. The composition is preferably free of malodour control agents.

5 Method of washing

Other aspects of the invention are directed to methods of washing dishware with the composition of the present invention. Said methods comprise the step of applying the composition, preferably in liquid form, onto the dishware surface, either in diluted or neat form and rinsing or leaving the composition to dry on the surface without rinsing the surface.

10

By “in its neat form”, it is meant herein that said composition is applied directly onto the surface to be treated and/or onto a cleaning device or implement such as a dish cloth, a sponge or a dish brush without undergoing any dilution (immediately) prior to the application. The cleaning device or implement is preferably wet before or after the composition is delivered to it. By “diluted form”, it is meant herein that said composition is diluted by the user with an appropriate solvent, typically water. By “rinsing”, it is meant herein contacting the dishware cleaned using a process according to the present invention with substantial quantities of appropriate solvent, typically water, after the step of applying the liquid composition herein onto said dishware. By “substantial quantities”, it is meant usually about 1 to about 10 liters.

20

The composition herein can be applied in its diluted form. Soiled dishes are contacted with an effective amount, typically from about 0.5 ml to about 20 ml (per about 25 dishes being treated), preferably from about 3ml to about 10 ml, of the detergent composition, preferably in liquid form, of the present invention diluted in water. The actual amount of detergent composition used will be based on the judgment of user, and will typically depend upon factors such as the particular product formulation of the composition, including the concentration of active ingredients in the composition, the number of soiled dishes to be cleaned, the degree of soiling on the dishes, and the like. Generally, from about 0.01 ml to about 150 ml, preferably from about 3ml to about 40ml of a liquid detergent composition of the invention is combined with from about 2000 ml to about 20000 ml, more typically from about 5000 ml to about 15000 ml of water in a sink having a volumetric capacity in the range of from about 1000 ml to about 20000 ml, more typically from about 5000 ml to about 15000 ml. The soiled dishes are immersed in the sink containing the diluted compositions then obtained, where contacting the soiled surface of

30

the dish with a cloth, sponge, or similar article cleans them. The cloth, sponge, or similar article may be immersed in the detergent composition and water mixture prior to being contacted with the dish surface, and is typically contacted with the dish surface for a period of time ranged from about 1 to about 10 seconds, although the actual time will vary with each application and user.

- 5 The contacting of cloth, sponge, or similar article to the dish surface is preferably accompanied by a concurrent scrubbing of the dish surface.

Another method of the present invention will comprise immersing the soiled dishes into a water bath or held under running water without any liquid dishwashing detergent. A device for
10 absorbing liquid dishwashing detergent, such as a sponge, is placed directly into a separate quantity of undiluted liquid dishwashing composition for a period of time typically ranging from about 1 to about 5 seconds. The absorbing device, and consequently the undiluted liquid dishwashing composition, is then contacted individually to the surface of each of the soiled dishes to remove said soiling. The absorbing device is typically contacted with each dish surface
15 for a period of time range from about 1 to about 10 seconds, although the actual time of application will be dependent upon factors such as the degree of soiling of the dish. The contacting of the absorbing device to the dish surface is preferably accompanied by concurrent scrubbing.

20 Alternatively, the device may be immersed in a mixture of the hand dishwashing composition and water prior to being contacted with the dish surface, the concentrated solution is made by diluting the hand dishwashing composition with water in a small container that can accommodate the cleaning device at weight ratios ranging from about 95:5 to about 5:95, preferably about 80:20 to about 20:80 and more preferably about 70:30 to about 30:70,
25 respectively, of hand dishwashing liquid:water respectively depending upon the user habits and the cleaning task.

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is
30 intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm".

Examples

Examples of Hand Dishwashing formulations comprising the lipase of the invention.

	1	2	3	4	5	6	7
	Wt%	Wt%	Wt%	Wt%	Wt%	Wt%	Wt%
Alkyl C ₁₀₋₁₄ Ethoxy Sulphate (AE0.6S)	26.9	21	-	-	-	5	15
Alkyl C ₁₀₋₁₄ Ethoxy Sulphate (AE2S)	-	-	18	14	13	-	-
Sodium alkyl benzene sulfonate	-	-	-	-	-	8	-
Sodium paraffin sulfonate	-	-	-	6	-	-	-
C12-14 dimethyl amine oxide	6.1	7	6	5	-	-	6
Cocamido propyl betaine	-	-	8	5	4	2	4
C12-13 EO7 nonionic	-	-	0.2	0.1	0.5	2	-
Branched Nonionic: 3-propyl heptanol EO8	1.0	0.5	-	-	-	-	1.0
PEI600-EO10-PO7 block polymer	-	0.5	-	-	-	0.4	0.8
Lipase ¹	0.02	0.02	0.001	0.03	0.1	0.01	0.02
Protease ²	-	0.04	-	-	-	-	-
Amylase ³	0.04	0.02	0.06	0.2	0.2	0.05	0.02
4-Formylphenylboronic acid	-	0.1	-	-	-	-	-
Potassium chloride	1.5	-	-	-	-	-	-
Calcium chloride	-	1	-	-	-	-	-
Sodium acetate	-	-	1.5	-	-	-	-
Potassium acetate	-	-	-	2	-	-	-
Sodium sulfate	-	-	-	-	1	-	-
Potassium sulfate	-	-	-	-	-	1.5	-
Potassium formate	-	-	-	-	-	-	2
Ethanol	4.0	5.0	3.0	3.0	2.0	-	3.0
Polypropylene glycol MW2000	1.1	0.8	1.1	1.1	1.1	0.5	1.1

22

Sodium chloride	1.3	0.8	1.3	0.5	0.8	1.3	1.3
Minors* and water	to balance up to 100%						

Notes

¹ Lipase is the D27R + G38A + D96E + D111A + G163K + T231R + N233R + D254S + P256T variant of SEQ ID: 1, supplied by Novozymes A/S, Bagsvaerd, Denmark

5 ² Protease is Savinase[®], supplied by Novozymes A/S, Bagsvaerd, Denmark

³ Amylase is Stainzyme[®] supplied by Novozymes A/S, Bagsvaerd, Denmark

CLAIMS

What is claimed is:-

1. A hand dishwashing liquid detergent composition comprising:
 - i) a surfactant system;
 - ii) at least one lipase wherein the lipase is a variant of a parent lipase, which variant has lipase activity, has at least 60% but less than 100% sequence identity with SEQ ID NO: 1, and comprises substitutions at positions corresponding to T231R+N233R and at least one or more (e.g., several) of D96E, D111A, D254S, G163K, P256T, G91T and G38A of SEQ ID NO: 1; and
 - iii) at least 0.05% by weight of the composition of at least one monovalent, divalent or trivalent cation or a mixture thereof.
2. A composition according to claim 1 wherein the variant further comprises substitutions at positions corresponding to D27R and/or N33Q of SEQ ID NO: 1.
3. A composition according to any of claims 1 or 2 wherein the variant comprises substitutions of SEQ ID NO: 1 selected from the group consisting of:
 - a) D96E+T231R+N233R;
 - b) N33Q+D96E+T231R+N233R;
 - c) N33Q+D111A+T231 R+N233R;
 - d) N33Q+T231 R+N233R+P256T;
 - e) N33Q+G38A+G91T+G163K+T231R+N233R+D254S;
 - f) N33Q+G38A+G91T+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
 - g) D27R+N33Q+G38A+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
 - h) D27R+N33Q+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+P256T;

- i) D27R+N33Q+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S;
- j) D27R+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;
- k) D96E+T231 R+N233R+D254S;
- l) T231R+N233R+D254S+P256T;
- m) G163K+T231 R+N233R+D254S;
- n) D27R+N33Q+G38A+G91T+D96E+G163K+T231 R+N233R+D254S+P256T;
- o) D27R+G91T+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;
- p) D96E+G163K+T231 R+N233R+D254S;
- q) D27R+G163K+T231 R+N233R+D254S;
- r) D27R+G38A+G91T+D96E+D111A+G163K+T231 R+N233R+D254S;
- s) D27R+G38A+G91T+D96E+G163K+T231 R+N233R+D254S+P256T;
- t) D27R+G38A+D96E+D111A+G163K+T231 R+N233R+D254S+P256T;
- u) D27R+D96E+G163K+T231R+N233R+D254S;
- v) D27R+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- w) D27R+G38A+D96E+G163K+T231 R+N233R+D254S+P256T;
- x) D111A+G163K+T231R+N233R+D254S+P256T;
- y) D111A+T231R+N233R;
- z) D111A+T231 R+N233R+D254S+P256T;
- aa) D27R+D96E+D111A+G163K+T231R+N233R;
- bb) D27R+D96E+D111A+T231 R+N233R;

- cc) D27R+G38A+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- dd) D27R+N33Q+G38A+D96E+D111A+T231R+N233R+D254S+P256T;
- ee) D27R+G38A+D96E+D111A+G163K+E210Q+T231 R+N233R+D254S+P256T;
- ff) D27R+T231 R+N233R+D254S+P256T;
- gg) D96E+D111A+G163K+T231R+N233R;
- hh) D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- ii) D96E+D111A+G163K+T231R+N233R+P256T;
- jj) D96E+D111A+T231R+N233R;
- kk) D96E+D111A+T231R+N233R+D254S;
- ll) D96E+D111A+T231R+N233R+D254S+P256T;
- mm) D96E+D111A+T231R+N233R+P256T;
- nn) D96E+G163K+T231R+N233R+D254S+P256T;
- oo) D96E+T231R+N233R+D254S+P256T;
- pp) D96E+T231R+N233R+P256T;
- qq) G38A+D96E+D111A+T231 R+N233R;
- rr) G91T+D96E+D111A+G163K+T231R+N233R+D254S+P256T;
- ss) G91T+D96E+D111A+T231 R+N233R;
- tt) G91T+D96E+T231 R+N233R;
- uu) G91T+T231 R+N233R+D254S+P256T;
- vv) N33Q+D96E+D111A+G163K+T231R+N233R+D254S+P256T;

ww) T231R+N233R+D254S+P256T; and

xx) T231R+N233R+P256T.

4. A composition according to any of the preceding claims comprising at least 0.35% by weight of the composition of the at least one cation.
5. A composition according to any of the preceding claims comprising from 0.35 to 4% by weight of the composition of the at least one cation.
6. A composition according to any of the preceding claims wherein the cation source is selected from the inorganic or organic salts of alkali metals, alkaline earth metals, of aluminum, iron, copper and zinc, preferably of the alkali metals and alkaline earth metals, with anion source preferably selected from the halides, sulphates, sulphites, carbonates, bicarbonates, phosphates, nitrates, nitrites, phosphates, formates, acetates, propionates, citrates, malates, tartrates, succinates, oxalates, lactates, and mixtures thereof.
7. A composition according to any of the preceding claims wherein the cation source is selected from sodium chloride, calcium chloride, potassium chloride, sodium sulfate, potassium sulfate, sodium acetate, potassium acetate, sodium formate, potassium formate, and mixtures thereof; more preferably the cation source is selected from calcium chloride, potassium chloride, potassium sulfate, sodium acetate, potassium acetate, sodium formate and potassium formate, and mixtures thereof and in particular from potassium chloride, potassium sulfate, potassium acetate, potassium formate, and mixtures thereof.
8. A composition according to any of the preceding claims having a pH of from 4 to 9 as measured in a 10% aqueous solution in distilled water at 20°C.
9. A composition according to any of the preceding claims comprising water and by weight of the composition from 5 to 15% of an anionic surfactant, preferably an alkyl ether sulfate, from 0.8 to 3% of an amphoteric surfactant, preferably an amine oxide surfactant, from 0.001-2% of a lipase, and at least 0.05% of a monovalent, divalent or trivalent cation and from 1 to 3% of a corresponding salt.

10. A composition according to any of the preceding claims further comprising an amphoteric and/or zwitterionic surfactant.
11. A composition according to any of the preceding claims further comprising a non-ionic surfactant.
12. A composition according to any of the preceding claims further comprising an amylase.
13. A composition according to any of the preceding claims further comprising a protease.
14. A composition according to at least one of the preceding claims comprising from 0.001 to 2% of lipase (pure enzyme) by weight of the composition.
15. A method of manually washing dishware comprising the step of: delivering a composition according to any of the preceding claims to a volume of water to form a wash liquor and immersing the dishware in the liquor.
16. A method of manually washing dishware comprising the step of: delivering a composition according to any of claims 1 to 14 directly onto the dishware or onto a cleaning implement and using the cleaning implement to clean the dishware.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/035629

A. CLASSIFICATION OF SUBJECT MATTER
INV. C11D3/386
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2014/184164 A1 (NOVOZYMES AS [DK]) 20 November 2014 (2014-11-20) page 77, line 34 page 79, line 1 - line 22; claims -----	1-16
Y	EP 2 623 586 A2 (PROCTER & GAMBLE [US]) 7 August 2013 (2013-08-07) paragraph [0318]; claims -----	1-16
Y	EP 2 365 050 A1 (PROCTER & GAMBLE [US]) 14 September 2011 (2011-09-14) paragraph [0045]; examples 5A-5T -----	1-16



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

22 July 2016

Date of mailing of the international search report

01/08/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Vernier, Frédéric

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/035629

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2014184164	A1	20-11-2014	AR 096270 A1 16-12-2015
		CN 105209612 A 30-12-2015	
		CN 105408475 A 16-03-2016	
		EP 2997139 A1 23-03-2016	
		EP 2997140 A1 23-03-2016	
		US 2016068787 A1 10-03-2016	
		US 2016076012 A1 17-03-2016	
		WO 2014184164 A1 20-11-2014	
		WO 2014186464 A1 20-11-2014	
EP 2623586	A2	07-08-2013	CN 104080902 A 01-10-2014
		EP 2623586 A2 07-08-2013	
		RU 2014127191 A 10-02-2016	
		US 2013203644 A1 08-08-2013	
		WO 2013116261 A2 08-08-2013	
EP 2365050	A1	14-09-2011	AR 080507 A1 11-04-2012
		BR 112012023014 A2 31-05-2016	
		CA 2791251 A1 15-09-2011	
		CA 2791252 A1 15-09-2011	
		EP 2365050 A1 14-09-2011	
		EP 2365051 A1 14-09-2011	
		JP 5571202 B2 13-08-2014	
		JP 5758412 B2 05-08-2015	
		JP 2013521401 A 10-06-2013	
		JP 2013521402 A 10-06-2013	
		US 2011220536 A1 15-09-2011	
		US 2011224455 A1 15-09-2011	
		US 2012151881 A1 21-06-2012	
		WO 2011112886 A1 15-09-2011	
		WO 2011112887 A1 15-09-2011	