



(86) Date de dépôt PCT/PCT Filing Date: 2009/05/26
(87) Date publication PCT/PCT Publication Date: 2009/11/26
(45) Date de délivrance/Issue Date: 2013/10/01
(85) Entrée phase nationale/National Entry: 2010/11/23
(86) N° demande PCT/PCT Application No.: US 2009/045183
(87) N° publication PCT/PCT Publication No.: 2009/143524
(30) Priorité/Priority: 2008/05/23 (US61/055,818)

(51) Cl.Int./Int.Cl. *A61K 9/107* (2006.01),
A61K 39/00 (2006.01), *A61K 39/02* (2006.01),
A61K 39/07 (2006.01), *A61K 39/08* (2006.01),
A61K 39/116 (2006.01), *A61K 39/29* (2006.01),
A61P 31/00 (2006.01), *A61P 31/04* (2006.01),
A61P 31/14 (2006.01), *A61P 37/04* (2006.01)

(72) Inventeurs/Inventors:
BAKER, JAMES R., JR., US;
BIELINSKA, ANNA U., US;
MANK, NICHOLAS J., US;
MAKIDON, PAUL E., US;
...

(73) Propriétaire/Owner:

(54) Titre : VACCINS A NANOEMULSION
(54) Title: NANOEMULSION VACCINES

(57) **Abrégé/Abstract:**

The present invention provides methods and compositions for the stimulation of immune responses. Specifically, the present invention provides immunogenic compositions and methods of using the same to induce immune responses (e.g., immunity (e.g., protective immunity) against Hepatitis B virus (HBV) and/or against one or a plurality of pathogens (e.g., vaccinia virus, H5N1 influenza virus, Bacillus anthracis, C. botulinum, Y. pestis, Hepatitis B, and/or HIV, etc.)) in a subject. Compositions and methods of the present invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine (e.g., vaccination)) and research applications.



(72) **Inventeurs(suite)/Inventors(continued):** KNOWLTON, JESSICA A., US; BLANCO, LUZ P., US; CAO, ZHENGYI, US;
SCOTT, ALISON J., US; NIGAVEKAR, SHRADDHA S., US

(73) **Propriétaires(suite)/Owners(continued):** THE REGENTS OF THE UNIVERSITY OF MICHIGAN, US

(74) **Agent:** SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
26 November 2009 (26.11.2009)



(10) International Publication Number
WO 2009/143524 A3

(51) International Patent Classification:
A61K 39/29 (2006.01)

(21) International Application Number:
PCT/US2009/045183

(22) International Filing Date:
26 May 2009 (26.05.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/055,818 23 May 2008 (23.05.2008) US

(71) Applicant (for all designated States except US): **THE REGENTS OF THE UNIVERSITY OF MICHIGAN** [US/US]; 1214 South University Avenue, 2nd Floor, Ann Arbor, MI 48104 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BAKER, James, R., Jr.** [US/US]; 3997 Holden Drive, Ann Arbor, MI 48103 (US). **BIELINSKA, Anna, U.** [PL/US]; 1652 Golden Lane, Ypsilanti, MI 48198 (US). **MANK, Nicholas, J.** [US/US]; 1421 West Huron St., Ann Arbor, MI 48103 (US). **MAKIDON, Paul, E.** [US/US]; 11670 Mason Road, Webberville, MI 48892 (US). **KNOWLTON, Jessica, A.** [US/US]; 1214 South University Avenue, 2nd Floor, Ann Arbor, MI 48104 (US). **BLANCO, Luz, P.** [CL/US]; 2391 Hickman Rd., Ann Arbor, MI 48105 (US). **CAO, Zhengyi** [US/US]; 1214 South University Avenue, 2nd Floor, Ann Arbor, MI 48104 (US). **SCOTT, Alison, J.** [US/US]; 1214 South University Avenue, 2nd Floor, Ann Arbor, MI 48104 (US). **NI-**

GAVEKAR, Shraddha, S. [IN/US]; 1361 Fox Pointe Circle, Ann Arbor, MI 48108 (US).

(74) Agent: **SISK, Tyler, J.**; Casimir Jones S.C., 440 Science Drive, Suite 203, Madison, WI 53711 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(88) Date of publication of the international search report:
7 January 2010

(54) Title: NANOEMULSION VACCINES

(57) Abstract: The present invention provides methods and compositions for the stimulation of immune responses. Specifically, the present invention provides immunogenic compositions and methods of using the same to induce immune responses (e.g., immunity (e.g., protective immunity) against Hepatitis B virus (HBV) and/or against one or a plurality of pathogens (e.g., vaccinia virus, H5N1 influenza virus, Bacillus anthracis, C. botulinum, Y. pestis, Hepatitis B, and/or HIV, etc.)) in a subject. Compositions and methods of the present invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine (e.g., vaccination)) and research applications.



WO 2009/143524 A3

NANOEMULSION VACCINES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to United States Provisional Patent Application Serial
5 No. 61/055,818, filed 23 May 2008.

FIELD OF THE INVENTION

The present invention provides methods and compositions for the stimulation of
immune responses. Specifically, the present invention provides immunogenic compositions
10 and methods of using the same to induce immune responses (e.g., immunity (e.g., protective
immunity) against Hepatitis B virus (HBV) and/or against one or a plurality of pathogens
(e.g., vaccinia virus, H5N1 influenza virus, *Bacillus anthracis*, *C. botulinum*, *Y. pestis*,
Hepatitis B, and/or HIV, etc.)) in a subject. Compositions and methods of the present
invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine
15 (e.g., vaccination)) and research applications.

BACKGROUND

Immunization is a principal feature for improving the health of people. Despite the
availability of a variety of successful vaccines against many common illnesses, infectious
20 diseases remain a leading cause of health problems and death. Significant problems inherent
in existing vaccines include the need for repeated immunizations, and the ineffectiveness of
the current vaccine delivery systems for a broad spectrum of diseases.

In order to develop vaccines against pathogens that have been recalcitrant to vaccine
development, and/or to overcome the failings of commercially available vaccines due to
25 expense, complexity, and underutilization, new methods of antigen presentation must be
developed which will allow for fewer immunizations, more efficient usage, and/or fewer side
effects to the vaccine.

SUMMARY OF THE INVENTION

30 The present invention provides methods and compositions for the stimulation of
immune responses. Specifically, the present invention provides immunogenic compositions
and methods of using the same to induce immune responses (e.g., immunity (e.g., protective

immunity) against Hepatitis B virus (HBV) and/or against one or a plurality of pathogens (e.g., vaccinia virus, H5N1 influenza virus, *Bacillus anthracis*, *C. botulinum*, *Y. pestis*, Hepatitis B, and/or HIV, etc.)) in a subject. Compositions and methods of the present invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine
5 (e.g., vaccination)) and research applications.

Accordingly, in some embodiments, the present invention provides a composition comprising a vaccine, the vaccine comprising an emulsion and a plurality of immunogens, the emulsion comprising an aqueous phase, an oil phase, and a solvent. In some embodiment, the plurality of immunogens comprises a plurality of pathogens (e.g., an
10 inactivated pathogens). In other embodiments, the plurality of immunogens comprises a plurality pathogen components and/or products (e.g., including, but not limited to, a protein, peptide, polypeptide, nucleic acid, polysaccharide, or a membrane component derived from the pathogens). In some embodiments, the immunogens and the emulsion are combined in a single vessel.

The present invention is not limited to any particular immunogens. For example,
15 multiple immunogens may be used in the present invention including, but not limited to, those described herein (e.g., protective antigen (PA), gp160, gp41, Tat, Nef, lethal factor, edema factor, protective antigen degradation products, etc.). Similarly, the present invention is not limited by the type of disease from which a subject is protected. Indeed, a subject can
20 be protected from a variety of diseases including, but not limited to, AIDS, the plague, disease caused by hepatitis virus, smallpox and anthrax. In some embodiments, immunity protects the subject from challenge with a subsequent exposure to live pathogen (e.g., HIV, vaccinia virus, *B. anthracis*, hepatitis virus (e.g., hepatitis B virus, hepatitis A virus, hepatitis C virus, etc.), *Yersinia pestis*, *Clostridium botulinum*, etc.).

In certain embodiments, the immunogen is selected from the group consisting of virus, bacteria, fungus and pathogen products derived from the virus, bacteria, or fungus. The present invention is not limited to a particular virus. A variety of viral immunogens are contemplated including, but not limited to, influenza A virus, avian influenza virus, H5N1 influenza virus, West Nile virus, SARS virus, Marburg virus, Arenaviruses, Nipah virus,
25 alphaviruses, filoviruses, herpes simplex virus I, herpes simplex virus II, sendai virus, sindbis virus, vaccinia virus, parvovirus, human immunodeficiency virus, hepatitis B virus, hepatitis C virus, hepatitis A virus, cytomegalovirus, human papilloma virus, picornavirus, hantavirus,

junin virus, and ebola virus. The present invention is not limited to a particular bacteria. A variety of bacterial immunogens are contemplated including, but not limited to, *Bacillus cereus*, *Bacillus circulans* and *Bacillus megaterium*, *Bacillus anthracis*, bacterial of the genus *Brucella*, *Vibrio cholera*, *Coxiella burnetii*, *Francisella tularensis*, *Chlamydia psittaci*,
5 *Ricinus communis*, *Rickettsia prowazekii*, bacteria of the genus *Salmonella*, *Cryptosporidium parvum*, *Burkholderia pseudomallei*, *Clostridium perfringens*, *Clostridium botulinum*, *Vibrio cholerae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Neisseria gonorrhea*, *Haemophilus influenzae*, *Escherichia coli*, *Salmonella typhimurium*, *Shigella dysenteriae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*,
10 *Yersinia pestis*, *Yersinia enterocolitica*, and *Yersinia pseudotuberculosis*. The present invention is also not limited to a particular fungus. A variety of fungal immunogens are contemplated including, but not limited to, *Candida* and *Aspergillus*.

In some embodiments, the present invention provides a method of inducing an immune response to a plurality of immunogens (e.g., two or more (e.g., three, four, five, six, seven, eight, or more immunogens) derived from a plurality of environmental pathogens (e.g., those described herein))) in a subject, comprising providing a nanoemulsion; and a plurality of immunogens; combining the nanoemulsion with the immunogens; and administering the combined nanoemulsion and immunogens to the subject under conditions such that the subject produces an immune response to the immunogens. In some
20 embodiments, administering comprises mucosal administration. In some embodiments, inducing an immune response induces immunity to each of the plurality of immunogens in the subject. In some embodiments, inducing an immune response to the immunogens induces immunity to the plurality of pathogens from which the immunogens are derived. In some embodiments, immunity comprises systemic immunity. In some embodiments, immunity
25 comprises mucosal immunity. In some embodiments, the immune response comprises increased expression of IFN- γ in the subject. In some embodiments, the immune response comprises increased expression of TNF- α in the subject. In some embodiments, the immune response comprises a systemic IgG response to the immunogens. In some embodiments, the immune response comprises a mucosal IgA response to the immunogens. In some
30 embodiments, the composition comprises between 15 and 75 μg of a recombinant immunogen. The present invention is not limited to this amount of immunogen. Indeed, a variety of doses of immunogen are contemplated to be useful in the present invention. In

some embodiments, a composition comprising a plurality of immunogens utilized to induce an immune response to a plurality of immunogens and/or environmental pathogens comprises two or more immunogens from virus, bacteria, fungus and pathogen products derived from one or more viruses, bacteria, or fungi. In some embodiments, the immunogens are
5 combined (e.g., in any ratio) to optimize protection against environmental pathogens from which the immunogens are derived.

The present invention further provides a kit comprising a vaccine, the vaccine comprising an emulsion and one or more immunogens, the emulsion comprising an aqueous phase, an oil phase, and a solvent. In some embodiments, the kit further comprises
10 instructions for using the kit for vaccinating a subject against the one or more immunogens.

In some embodiment, the one or more immunogens comprise a pathogen (e.g., an inactivated pathogen). In other embodiments, the immunogens comprise one or more pathogen products (e.g., including, but not limited to, a protein, peptide, polypeptide, nucleic acid, polysaccharide, or membrane component derived from the pathogen). In some
15 embodiments, the one or more immunogens and the emulsion are combined in a single vessel.

In still further embodiments, the present invention provides a method of inducing immunity to one or more immunogens, comprising providing an emulsion comprising an aqueous phase, an oil phase, and a solvent; and one or more immunogens; combining the
20 emulsion with the one or more immunogens to generate a vaccine composition; and administering the vaccine composition to a subject. In some embodiments, administering comprises contacting the vaccine composition with a mucosal surface of the subject. For example, in some embodiments, administering comprises intranasal administration. In some preferred embodiments, the administering occurs under conditions such that the subject
25 generates immunity to the one or more immunogens (e.g., via generating humoral immune responses to the one or more immunogens).

In some embodiment, the one or more immunogens comprise a pathogen (e.g., an inactivated pathogen). In other embodiments, the one or more immunogens comprise a pathogen product (e.g., including, but not limited to, a protein, peptide, polypeptide, nucleic acid, polysaccharide, or membrane component derived from the pathogen). In some
30 embodiments, the one or more immunogens and the emulsion are combined in a single vessel.

In some embodiments, the present invention provides a method of inducing an immune response to a plurality of immunogens (e.g., protective antigen (PA) of *Bacillus anthracis*, rLcrV or LcrV10 protein of *Yersinia pestis*, and rHCR/A1 of *Clostridium botulinum*) in a subject comprising providing a composition comprising a nanoemulsion and the plurality of immunogens (e.g., pathogens inactivated by a nanoemulsion of the present invention, and/or one or a plurality of protein and/or peptide antigens derived from one or a plurality of pathogens (e.g., recombinantly produced)); and administering the composition to the subject under conditions such that the subject generates an immune response to the plurality of immunogens. The present invention is not limited by the plurality of immunogens utilized. For example, in some embodiments, the plurality of immunogens are pathogens inactivated by a nanoemulsion of the present invention or are isolated, purified and/or recombinant protein or peptide antigens, or derivatives or variants thereof, derived from the plurality of pathogens (e.g., vaccinia virus inactivated by a nanoemulsion, and/or protein antigens (e.g., including, but not limited to, protective antigen (PA), lethal factor (LF), edema factor (EF), and PA degradation products from *B. anthracis* or gp160, gp120, gp41, Tat or Nef from HIV). For example, in some embodiments, the present invention provides a composition comprising a nanoemulsion (e.g., W₈₀5EC) and recombinant protective antigen of *B. anthracis*, recombinant rHCR/A1 of *C. botulinum*, and recombinant rLcrV (e.g., rLcrV10) protein of *Y. pestis*, and methods of using and/or administering (e.g., nasally administering) the composition to induce immune responses (e.g., immune responses specific to each of the immunogens) in a subject (e.g., thereby providing protective immunity to the subject from *B. anthracis*, *C. botulinum*, and/or *Y. pestis*. In some embodiments, immune responses in the subject comprise generation of antibodies to the immunogens. In some embodiments, the antibodies generated comprise IgG and/or IgA antibodies. In some embodiments, the immune responses generated in a subject via administration of a nanoemulsion composition comprising a plurality of immunogens (e.g., two, three, four, five, six, seven, eight, or more immunogens) are similar to (e.g., not detectably different than) immune responses that are generated in a subject via administration of a plurality of nanoemulsion compositions, wherein each nanoemulsion composition comprises a single immunogen (e.g., the antigen specific antibody titer levels in a subject administered a composition comprising plurality of immunogens is similar to the antigen specific antibody titer levels in a subject administered a plurality of nanoemulsion compositions wherein each

composition comprises a single immunogen).

In some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a plurality of immunogens does not comprise a preservative. For example, in some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a plurality of immunogens does not comprise a mercury based preservative (e.g., thiomersal).

In some embodiments, a composition comprising a nanoemulsion and a plurality of immunogens of the invention is utilized for the treatment and/or prophylaxis of infection or disease caused by a plurality of pathogens, especially treatment or prophylaxis, for example, of anthrax, plague and/or toxin poisoning. In some embodiments, an immunogenic

composition comprising a nanoemulsion and plurality of immunogens comprises a nanoemulsion that skews the immune response toward a Th1 type immune response. The present invention is not limited by the type of nanoemulsion utilized. Indeed, a variety of nanoemulsions can be utilized including but not limited to W₈₀5EC, although the present invention is not so limited. For example, in some embodiments, the nanoemulsion is selected

from one of the nanoemulsion formulations described herein. In some embodiments, the composition comprises between 0.5-50% nanoemulsion solution, although greater and lesser amounts also find use in the invention. For example, in some embodiments, the immunogenic composition comprises about 0.1%-0.5%, 0.5%-1.0%, 1.0% - 10%, about 10%-20%, about 20% - 30%, about 30%-40%, about 40%- 50%, about 50%-60% or more

nanoemulsion solution. In some embodiments, the immunogenic composition comprises 20% nanoemulsion solution (e.g., 20% W₈₀5EC or other emulsion described herein). In some embodiments, the immunogenic composition comprises about 10% nanoemulsion solution.

In some embodiments, the immunogenic composition comprises about 15% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 20%

nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 12% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 8% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 5% nanoemulsion solution. In some embodiments, the

immunogenic composition comprises about 2% nanoemulsion solution. In some

embodiments, the immunogenic composition comprises about 1% nanoemulsion solution. In some embodiments, an immunogenic composition (e.g., that is administered to a subject in order to generate an immune response in the subject) comprises between about 5 and 75 µg

of each of the plurality of immunogens (e.g., between 5 and 75 μ g of recombinant protective antigen (PA) of *B. anthracis*, between 5 and 75 μ g of recombinant LcrV or LcrV10 of *Y. pestis*, and between 5 and 75 μ g of recombinant receptor binding domain HCR/A1 of *C. botulinum* neurotoxin). In some embodiments, an immunogenic composition (e.g., that is administered to a subject in order to generate an immune response in the subject) comprises 20 μ g of each of the plurality of immunogens (e.g., recombinant protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and recombinant receptor binding domain HCR/A1 of *C. botulinum* neurotoxin). However, the present invention is not limited to this amount of immunogen. Indeed, a variety of doses of immunogen are contemplated to be useful in the present invention. For example, in some embodiments, it is expected that each dose (e.g., of an immunogenic composition comprising a plurality of immunogens (e.g., administered to a subject to induce an immune response (e.g., a protective immune response (e.g., protective immunity)))) comprises 0.05-5000 μ g of each immunogen (e.g., recombinant, isolated and/or purified immunogen (e.g., recombinant protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or recombinant receptor binding domain HCR/A1 of *C. botulinum* neurotoxin)). In some embodiments, each dose will comprise 1-500 μ g, in some embodiments, each dose will comprise 350-750 μ g, in some embodiments, each dose will comprise 50-200 μ g, in some embodiments, each dose will comprise 10-100 μ g of each immunogen, each dose will comprise 10-75 μ g of each immunogen, each dose will comprise 25-75 μ g of each immunogen, in some embodiments, each dose will comprise 10-25 μ g, in some embodiments, each dose will comprise 20 μ g of each immunogen (e.g., recombinant, isolated and/or purified immunogen (e.g., recombinant protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or recombinant receptor binding domain HCR/A1 of *C. botulinum* neurotoxin)). In some embodiments, each dose comprises an amount of the plurality of immunogens sufficient to generate an immune response. An effective amount of the plurality of immunogens in a dose need not be quantified, as long as the amount of immunogens generates an immune response in a subject when administered to the subject.

In some embodiments, the immunogenic composition is stable (e.g., at room temperature (e.g., for 12 hours, one day, two days, three days, four days, a week, two weeks, three weeks, a month, two months, three months, four months, five months, six months, 9

months, a year or more). In some embodiments, the immunogenic composition comprises a pharmaceutically acceptable carrier. The present invention is not limited to any particular pharmaceutically acceptable carrier. Indeed, any suitable carrier may be utilized including but not limited to those described herein. In some embodiments, the immunogenic
5 composition further comprises an adjuvant. The present invention is not limited to any particular adjuvant and any one or more adjuvants described herein find use in a composition of the invention including but not limited to adjuvants that skew toward a Th1 immune response (e.g., that induces expression and/or activity of Th1 type cytokines (e.g., IFN- γ , TNF- α , IL2 and/or IL-12). In some embodiments, the immunogenic composition comprising
10 a nanoemulsion and a plurality of immunogens comprises an adjuvant that skews the immune response toward a Th1 type immune response. In some embodiments, the immunogenic composition comprising a nanoemulsion and a plurality of immunogens does not comprise an adjuvant that skews the immune response toward a Th1 type immune response (e.g., the immunogenic composition comprising nanoemulsion and plurality of immunogens skews
15 toward a Th1 immune response due to the nanoemulsion utilized and not the presence of an adjuvant). In some embodiments, the level of Th1-type cytokines increases to a greater extent than the level of Th2-type cytokines (e.g., cytokines levels are readily assessed using standard assays, See, e.g., Mosmann and Coffman, Ann. Rev. Immunol. 7:145-173, 1989). In some embodiments, the plurality of immunogens comprise pathogen products (e.g.,
20 including, but not limited to, a protein, peptide, polypeptide, nucleic acid, polysaccharide, or a membrane component derived from the pathogen). In some embodiments, the plurality of immunogens and the nanoemulsion are combined in a single vessel.

In some embodiments, the present invention provides a method of inducing an immune response to a plurality of immunogens (e.g., protective antigen (PA) of *B. anthracis*,
25 LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) in a subject comprising: providing an immunogenic composition comprising a nanoemulsion and protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin, and administering the composition to the subject under conditions such that the subject generates an immune
30 response toward protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin. The present invention is not limited by the route chosen for administration of a composition of the present invention. In

some embodiments, administering the immunogenic composition comprises contacting a mucosal surface of the subject with the composition. In a preferred embodiment, the mucosal surface comprises nasal mucosa. In some embodiments, the immune response comprises a systemic IgG response to the immunogens. In some embodiments, the immune response
5 comprises a mucosal IgA response to the immunogens. In some embodiments, inducing an immune response induces immunity to *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin in the subject. In some embodiments, the immunity comprises systemic immunity. In some embodiments, the immunity comprises mucosal immunity. In some embodiments, a subject administered an immunogenic composition comprising a nanoemulsion and plurality of
10 immunogens generates a Th1 type immune response. In some embodiments, the Th1 type immune response comprises enhanced expression of IFN- γ and/or TNF- α . In some embodiments, the level of Th1-type cytokines increases to a greater extent than the level of Th2-type cytokines. For example, in some embodiments, a subject administered an immunogenic composition comprising a nanoemulsion and plurality of immunogens induces
15 a greater than 3 fold, greater than 5 fold, greater than 10 fold, greater than 20 fold, greater than 25 fold, greater than 30 fold or more enhanced expression of Th1 type cytokines, with lower increases (e.g., less than 3 fold, less than two fold or less) enhanced expression of Th2 type cytokines (e.g., IL-4, IL-5, and/or IL-10). In some embodiments, administration of an immunogenic composition comprising a nanoemulsion and a plurality of immunogens to a
20 subject generates immunogen specific antibodies in the subject. In some embodiments, the immunogen specific antibodies have a prevalence of IgG2b and/or IgG2a antibodies over that of IgG1 antibodies. In some embodiments, administration of an immunogenic composition comprising a nanoemulsion and a HBV immunogen to a subject generates immunogen specific IgA antibodies in the subject. The present invention is not limited to any particular
25 nanoemulsion utilized in a method of inducing an immune response to a plurality of immunogens in a subject. Indeed, a variety of nanoemulsions may be utilized including but not limited to W₈₀5EC. For example, in some embodiments, the nanoemulsion is selected from one of the nanoemulsion formulations described herein. In a preferred embodiment, the immunogenic composition comprising a nanoemulsion and plurality of immunogens does not
30 comprise an adjuvant that skews the immune response toward a Th1 type immune response (e.g., the immunogenic composition comprising nanoemulsion and plurality of immunogens skews toward a Th1 immune response due to the nanoemulsion utilized and not the presence

of an adjuvant). In some embodiments, each dose comprises an amount nanoemulsion and plurality of immunogens sufficient to generate an immune response to the plurality of immunogens in a subject. An effective amount of nanoemulsion and plurality of immunogens is a dose that need not be quantified, as long as the amount nanoemulsion and plurality of immunogens generates immunogen-specific immune responses in a subject when administered to the subject. In some embodiments, each dose will comprise 1-500 μg , in some embodiments, each dose will comprise 350-750 μg , in some embodiments, each dose will comprise 50-200 μg , in some embodiments, each dose will comprise 10-100 μg of each immunogen, each dose will comprise 10-75 μg of each immunogen, each dose will comprise 25-75 μg of each immunogen, in some embodiments, each dose will comprise 10-25 μg , in some embodiments, each dose will comprise 20 μg of each immunogen (e.g., recombinant, isolated and/or purified immunogen (e.g., recombinant protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or recombinant receptor binding domain HCR/A1 of *C. botulinum* neurotoxin)). In some embodiments, a 20% nanoemulsion solution is utilized. In some embodiments, the nanoemulsion comprises W₈₀5EC. In some embodiments, the immunity protects the subject from displaying signs or symptoms of disease caused by the immunogens. In some embodiments, the immunity protects the subject from challenge with a subsequent exposure to live pathogens from which the immunogens are derived (e.g., *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin). In some embodiments, the immunogenic composition further comprises an adjuvant. In some embodiments, the subject is a human.

In some embodiments, the present invention provides an immunogenic composition comprising a nanoemulsion and a hepatitis B virus (HBV) immunogen, the nanoemulsion comprising an aqueous phase, an oil phase, and a solvent. In some embodiments, the immunogen comprises whole HBV (e.g., inactivated HBV (e.g., inactivated using an emulsion of the invention or by other means)). In some embodiments, the immunogen is a HBV antigen. In some embodiments, the immunogen is a plurality of HBV antigens. Preferably the HBV antigen is hepatitis B surface antigen (HBsAg). In some embodiments, the antigen is hepatitis core antigen (HBcAg). In some embodiments the antigen is hepatitis B e antigen (HBeAg). The present invention is not limited by the type or source of HBV antigen (e.g., HBsAg). For example, the preparation of hepatitis B surface antigen is well documented (See for example, Harford et. al. in Develop. Biol. Standard 54, page 125 (1983),

Gregg et. al. in Biotechnology, 5, page 479 (1987), EP-A-0 226 846, EP-A-0 299 108 and references therein). In some embodiments, the HBsAg antigen is identified to be mainly free of HBsAg aggregates. In some embodiments, the HBsAg antigen is identified to be mainly composed of HBsAg aggregates. In some embodiments, the HBsAg antigen is treated (e.g.,
5 using dialysis and/or sonication (e.g., ultrasonication (e.g., to disrupt aggregates))) prior to combining with a nanoemulsion of the invention. In some embodiments, HBsAg is in particle form. In some embodiments, HBsAg comprises HBsAg S-antigen. In some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a nanoemulsion and a hepatitis B antigen (e.g., HBsAg) does not comprise a preservative. For
10 example, in some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a nanoemulsion and hepatitis B antigen (e.g., HBsAg) does not comprise a mercury based preservative (e.g., thiomersal). In some embodiments, a composition comprising a nanoemulsion and a hepatitis B immunogen of the invention is utilized for the treatment and/or prophylaxis of hepatitis B infections, especially treatment or prophylaxis,
15 for example, of chronic hepatitis B infections. In some embodiments, an immunogenic composition comprising a nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) comprises a nanoemulsion that skews the immune response toward a Th1 type immune response. The present invention is not limited by the type of nanoemulsion utilized. Indeed, a variety of nanoemulsions can be utilized including but not limited to W₈₀5EC,
20 although the present invention is not so limited. For example, in some embodiments, the nanoemulsion is selected from one of the nanoemulsion formulations described herein. In some embodiments, the composition comprises between 0.5-50% nanoemulsion solution, although greater and lesser amounts also find use in the invention. For example, in some
25 embodiments, the immunogenic composition comprises about 0.1%-0.5%, 0.5%-1.0%, 1.0% - 10%, about 10%-20%, about 20% - 30%, about 30%-40%, about 40%- 50%, about 50%- 60% or more nanoemulsion solution. In some embodiments, the immunogenic composition comprises 20% nanoemulsion solution (e.g., 20% W₈₀5EC or other emulsion described herein). In some embodiments, the immunogenic composition comprises about 10% nanoemulsion solution. In some embodiments, the immunogenic composition comprises
30 about 15% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 20% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 12% nanoemulsion solution. In some embodiments, the

immunogenic composition comprises about 8% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 5% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 2% nanoemulsion solution. In some embodiments, the immunogenic composition comprises about 1% nanoemulsion solution. In some embodiments, an immunogenic composition (e.g., that is administered to a subject in order to generate an immune response in the subject) comprises between about 5 and 75 μg of HBV immunogen (e.g., HBV antigen (e.g., HBsAg)). However, the present invention is not limited to this amount of immunogen. Indeed, a variety of doses of immunogen are contemplated to be useful in the present invention. For example, in some embodiments, it is expected that each dose (e.g., of an immunogenic composition comprising a nanoemulsion and a HBV immunogen (e.g., administered to a subject to induce an immune response (e.g., a protective immune response (e.g., protective immunity)))) comprises 0.05-5000 μg of HBV immunogen (e.g., recombinant, isolated and/or purified HBV immunogen (e.g., HBV antigen (e.g., HBsAg))). In some embodiments, each dose will comprise 1-500 μg , in some embodiments, each dose will comprise 350-750 μg , in some embodiments, each dose will comprise 50-200 μg , in some embodiments, each dose will comprise 10-100 μg of immunogen, each dose will comprise 10-75 μg of immunogen, each dose will comprise 25-75 μg of immunogen, in some embodiments, each dose will comprise 10-25 μg , in some embodiments, each dose will comprise 20 μg of HBV immunogen (e.g., recombinant, isolated and/or purified HBV immunogen (e.g., HBV antigen (e.g., HBsAg))). In some embodiments, each dose comprises an amount of the immunogen sufficient to generate an immune response. An effective amount of the immunogen in a dose need not be quantified, as long as the amount of immunogen generates an immune response in a subject when administered to the subject. In some embodiments, the immunogenic composition is stable (e.g., at room temperature (e.g., for 12 hours, one day, two days, three days, four days, a week, two weeks, three weeks, a month, two months, three months, four months, five months, six months, 9 months, a year or more). In some embodiments, the immunogenic composition comprises a pharmaceutically acceptable carrier. The present invention is not limited to any particular pharmaceutically acceptable carrier. Indeed, any suitable carrier may be utilized including but not limited to those described herein. In some embodiments, the immunogenic composition further comprises an adjuvant. The present invention is not limited to any particular adjuvant and any one or more adjuvants described herein find use in

a composition of the invention including but not limited to adjuvants that skew toward a Th1 immune response (e.g., that induces expression and/or activity of Th1 type cytokines (e.g., IFN- γ , TNF- α , IL2 and/or IL-12). In some embodiments, the immunogenic composition comprising a nanoemulsion and a HBV immunogen comprises an adjuvant that skews the immune response toward a Th1 type immune response. In some embodiments, the immunogenic composition comprising a nanoemulsion and a HBV immunogen does not comprise an adjuvant that skews the immune response toward a Th1 type immune response (e.g., the immunogenic composition comprising nanoemulsion and HBV immunogen skews toward a Th1 immune response due to the nanoemulsion utilized and not the presence of an adjuvant). In some embodiments, the level of Th1-type cytokines increases to a greater extent than the level of Th2-type cytokines (e.g., cytokines levels are readily assessed using standard assays, See, e.g., Mosmann and Coffman, Ann. Rev. Immunol. 7:145-173, 1989). In some embodiments, the immunogen comprises a pathogen product (e.g., including, but not limited to, a protein, peptide, polypeptide, nucleic acid, polysaccharide, or a membrane component derived from the pathogen). In some embodiments, the HBV immunogen and the nanoemulsion are combined in a single vessel.

In some embodiments, the present invention provides a method of inducing an immune response to hepatitis B virus (HBV) in a subject comprising: providing an immunogenic composition comprising a nanoemulsion and a HBV immunogen, and administering the composition to the subject under conditions such that the subject generates an immune response toward HBV. The present invention is not limited by the route chosen for administration of a composition of the present invention. In some embodiments, administering the immunogenic composition comprises contacting a mucosal surface of the subject with the composition. In a preferred embodiment, the mucosal surface comprises nasal mucosa. In some embodiments, the immune response comprises a systemic IgG response to HBV. In some embodiments, the immune response comprises a mucosal IgA response to the immunogen. In some embodiments, inducing an immune response induces immunity to HBV in the subject. In some embodiments, the immunity comprises systemic immunity. In some embodiments, the immunity comprises mucosal immunity. In some embodiments, a subject administered an immunogenic composition comprising a nanoemulsion and a HBV immunogen generates a Th1 type immune response. In some embodiments, the Th1 type immune response comprises enhanced expression of IFN- γ and/or

TNF- α . In some embodiments, the level of Th1-type cytokines increases to a greater extent than the level of Th2-type cytokines. For example, in some embodiments, a subject administered an immunogenic composition comprising a nanoemulsion and HBV antigen induces a greater than 3 fold, greater than 5 fold, greater than 10 fold, greater than 20 fold, greater than 25 fold, greater than 30 fold or more enhanced expression of Th1 type cytokines, with lower increases (e.g., less than 3 fold, less than two fold or less) enhanced expression of Th2 type cytokines (e.g., IL-4, IL-5, and/or IL-10). In some embodiments, administration of an immunogenic composition comprising a nanoemulsion and a HBV immunogen to a subject generates HBsAg specific antibodies in the subject. In some embodiments, the HBsAg specific antibodies have a prevalence of IgG2b and/or IgG2a antibodies over that of IgG1 antibodies. In some embodiments, administration of an immunogenic composition comprising a nanoemulsion and a HBV immunogen to a subject generates HBsAg specific IgA antibodies in the subject. The present invention is not limited to any particular nanoemulsion utilized in a method of inducing an immune response to hepatitis B virus (HBV) in a subject. Indeed, a variety of nanoemulsions may be utilized including but not limited to W₈₀5EC. For example, in some embodiments, the nanoemulsion is selected from one of the nanoemulsion formulations described herein. In a preferred embodiment, the immunogenic composition comprising a nanoemulsion and a HBV immunogen does not comprise an adjuvant that skews the immune response toward a Th1 type immune response (e.g., the immunogenic composition comprising nanoemulsion and HBV immunogen skews toward a Th1 immune response due to the nanoemulsion utilized and not the presence of an adjuvant). In some embodiments, each dose comprises an amount nanoemulsion and HBV immunogen sufficient to generate an immune response to HBV in a subject. An effective amount of nanoemulsion and HBV immunogen is a dose that need not be quantified, as long as the amount nanoemulsion and HBV immunogen generates a HBV-specific immune response in a subject when administered to the subject. In some embodiments, the immunogenic composition comprising a nanoemulsion and HBV immunogen is administered to the subject under conditions such that about 10-25 μ g of HBV immunogen (e.g., recombinant, isolated and/or purified HBV immunogen (e.g., HBV antigen (e.g., HBsAg))) is present in a dose administered to the subject, although other doses (e.g., 5-20 μ g, 20 μ g, 25-75 μ g, 50-200 μ g, 350-750 μ g or more of HBV immunogen (e.g., HBV antigen (e.g., HBsAg))) may also be utilized. In some embodiments, a 20% nanoemulsion solution is

utilized. In some embodiments, the nanoemulsion comprises W₈₀5EC. In some
embodiments, the immunity protects the subject from displaying signs or symptoms of
disease caused by HBV. In some embodiments, the immunity protects the subject from
challenge with a subsequent exposure to live HBV. In some embodiments, the immunogenic
5 composition further comprises an adjuvant. In some embodiments, the subject is a human.

The present invention is not limited to any specific nanoemulsion composition.
Indeed, a variety of nanoemulsion compositions are described herein that find use in the
present invention. Similarly, the present invention is not limited to a particular oil present in
the nanoemulsion. A variety of oils are contemplated, including, but not limited to, soybean,
10 avocado, squalene, olive, canola, corn, rapeseed, safflower, sunflower, fish, flavor, and water
insoluble vitamins. The present invention is also not limited to a particular solvent. A
variety of solvents are contemplated including, but not limited to, an alcohol (*e.g.*, including,
but not limited to, methanol, ethanol, propanol, and octanol), glycerol, polyethylene glycol,
and an organic phosphate based solvent. Nanoemulsion components including oils, solvents
15 and others are described in further detail below.

In some embodiments, the emulsion further comprises a surfactant. The present
invention is not limited to a particular surfactant. A variety of surfactants are contemplated
including, but not limited to, nonionic and ionic surfactants (*e.g.*, TRITON X-100; TWEEN
20; and TYLOXAPOL).

20 In certain embodiments, the emulsion further comprises a cationic halogen containing
compound. The present invention is not limited to a particular cationic halogen containing
compound. A variety of cationic halogen containing compounds are contemplated including,
but not limited to, cetylpyridinium halides, cetyltrimethylammonium halides,
cetyldimethylethylammonium halides, cetyldimethylbenzylammonium halides,
25 cetyltributylphosphonium halides, dodecyltrimethylammonium halides, and
tetradecyltrimethylammonium halides. The present invention is also not limited to a
particular halide. A variety of halides are contemplated including, but not limited to, halide
selected from the group consisting of chloride, fluoride, bromide, and iodide.

In still further embodiments, the emulsion further comprises a quaternary ammonium
30 containing compound. The present invention is not limited to a particular quaternary
ammonium containing compound. A variety of quaternary ammonium containing
compounds are contemplated including, but not limited to, Alkyl dimethyl benzyl ammonium

chloride,

dialkyl dimethyl ammonium chloride, n-Alkyl dimethyl benzyl ammonium chloride, n-Alkyl dimethyl ethylbenzyl ammonium chloride, Dialkyl dimethyl ammonium chloride, and n-Alkyl dimethyl benzyl ammonium chloride.

5 In some embodiments, the present invention provides a vaccine comprising an immunogenic composition comprising a nanoemulsion and plurality of immunogens. In some embodiments, the invention provides a kit comprising a vaccine, the vaccine comprising an immunogenic composition comprising a nanoemulsion and plurality of immunogens, the nanoemulsion comprising an aqueous phase, an oil phase, and a solvent. In 10 some embodiments, the kit further comprises instructions for using the kit for vaccinating a subject against a plurality of immunogens (e.g., *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin).

In still further embodiments, the present invention provides a method of inducing immunity to a plurality of immunogens (*B. anthracis*, *Y. pestis*, and/or *C. botulinum* 15 neurotoxin), comprising providing an emulsion comprising an aqueous phase, an oil phase, and a solvent; and a plurality of immunogens; combining the emulsion with the plurality of immunogens to generate a vaccine composition; and administering the vaccine composition to a subject. In some embodiments, administering comprises contacting the vaccine composition with a mucosal surface of the subject. For example, in some embodiments, 20 administering comprises intranasal administration. In some preferred embodiments, the administering occurs under conditions such that the subject generates immunity to a plurality of pathogens (e.g., *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin (e.g., via generating humoral immune responses to the one or more immunogens)).

In some embodiments, the present invention provides a vaccine comprising an 25 immunogenic composition comprising a nanoemulsion and HBV immunogen. In some embodiments, the invention provides a kit comprising a vaccine, the vaccine comprising an immunogenic composition comprising a nanoemulsion and HBV immunogen, the nanoemulsion comprising an aqueous phase, an oil phase, and a solvent. In some embodiments, the kit further comprises instructions for using the kit for vaccinating a subject 30 against HBV.

In still further embodiments, the present invention provides a method of inducing immunity to HBV, comprising providing an emulsion comprising an aqueous phase, an oil

phase, and a solvent; and one or more HBV immunogens; combining the emulsion with the one or more HBV immunogens to generate a vaccine composition; and administering the vaccine composition to a subject. In some embodiments, administering comprises contacting the vaccine composition with a mucosal surface of the subject. For example, in some
5 embodiments, administering comprises intranasal administration. In some preferred embodiments, the administering occurs under conditions such that the subject generates immunity to HBV (e.g., via generating humoral immune responses to the one or more immunogens).

The present invention is not limited by the nature of the immune response generated
10 (e.g., post administration of an immunogenic composition comprising a nanoemulsion and one or a plurality of immunogens). Indeed, a variety of immune responses may be generated and measured in a subject administered a composition comprising an immunogenic composition comprising a nanoemulsion and plurality of immunogens of the present invention including, but not limited to, activation, proliferation or differentiation of cells of
15 the immune system (e.g., B cells, T cells, dendritic cells, antigen presenting cells (APCs), macrophages, natural killer (NK) cells, etc.); up-regulated or down-regulated expression of markers and cytokines; stimulation of IgA, IgM, and/or IgG titers; splenomegaly (e.g., increased spleen cellularity); hyperplasia, mixed cellular infiltrates in various organs, and/or other responses (e.g., of cells) of the immune system that can be assessed with respect to
20 immune stimulation known in the art. In some embodiments, administering comprises contacting a mucosal surface of the subject with the composition. The present invention is not limited by the mucosal surface contacted. In some preferred embodiments, the mucosal surface comprises nasal mucosa. In some embodiments, the mucosal surface comprises vaginal mucosa. In some embodiments, administering comprises parenteral administration.
25 The present invention is not limited by the route chosen for administration of a composition of the present invention. In some embodiments, inducing an immune response induces immunity to a plurality of immunogens in the subject. In some embodiments, the immunity comprises systemic immunity. In some embodiments, the immunity comprises mucosal immunity. In some embodiments, the immune response comprises increased expression of
30 IFN- γ and/or TNF- α in the subject. In some embodiments, the immune response comprises a systemic IgG (e.g., IgG2b and/or IgG2a) response. In some embodiments, the immune response comprises a mucosal IgA response. In some embodiments, the composition

comprises a 20% nanoemulsion solution. However, the present invention is not limited to this amount (e.g., percentage) of nanoemulsion. For example, in some embodiments, an immunogenic composition comprises less than 20% nanoemulsion (e.g., 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less nanoemulsion). In some embodiments, an immunogenic composition comprises more than 20% nanoemulsion (e.g., 25%, 30%, 35%, 40%, 45%, 50%, 60% or more). In some embodiments, an immunogenic composition of the present invention comprises any of the nanoemulsions described herein. In some embodiments, the nanoemulsion comprises W₂₀5EC. In some preferred embodiments, the nanoemulsion comprises W₈₀5EC. In some embodiments, the nanoemulsion is X8P. In some embodiments, immunity protects the subject from displaying signs or symptoms of disease caused by one or a plurality of pathogens (e.g., hepatitis B virus, *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin). In some embodiments, an immunogenic composition further comprises an adjuvant. The present invention is not limited by the type of adjuvant utilized. In some embodiments, the adjuvant is a CpG oligonucleotide. In some embodiments, the adjuvant is monophosphoryl lipid A. A number of other adjuvants that find use in the present invention are described herein. In some embodiments, the subject is a human. In some embodiments, the immunity protects the subject from displaying signs or symptoms of a infection with or exposure to one or a plurality of pathogens (e.g., *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin). In some embodiments, immunity reduces the risk of infection, disease and/or morbidity upon one or more exposures to one or a plurality of pathogens (e.g., *B. anthracis*, *Y. pestis*, and/or *C. botulinum* neurotoxin).

DESCRIPTION OF THE FIGURES

The following figures form part of the present specification and are included to further demonstrate certain aspects and embodiments of the present invention. The invention may be better understood by reference to one or more of these figures in combination with the description of specific embodiments presented herein.

The following figures form part of the present specification and are included to further demonstrate certain aspects and embodiments of the present invention. The invention may be better understood by reference to one or more of these figures in combination with the description of specific embodiments presented herein.

Figure 1 shows the histopathological evaluation and metabolic analysis of animals treated intranasally with NE adjuvant or HBsAg-NE formulations.

Figure 2 shows Histopathological analysis of nasal tissue exposed to NE adjuvant or HBsAg-NE. Photomicrographs of H&E stained nasal epithelium collected from mice 14 days following the boost vaccination shown in (A-B). Nasal epithelium collected 24 hours following boost vaccination with HBsAg-NE scored as +1 shown in (C). Nasal epithelium collected 24 hours following boost vaccination with HBsAg-NE scored as a +2 grade shown in (D). Arrows indicate a single microscopic focus of accumulation of mucoid material and debris in the nasal passages in the absence of inflammatory changes (C-D). Nasal epithelium collected 14 days following final boost vaccination from rats shown in (E-F), and guinea pigs shown in (G-H) treated a total of 3 doses of HBsAg-NE administered 14 days apart. Nasal biopsies collected 24 hours following the final dose in dogs treated with a total of three doses of NE adjuvant: 200 μ l/dose shown in (I) and 400 μ l/dose shown in (J).

Figure 3 shows the effect of concentration and temperature on NE particle size.

Figure 4 shows the Stability of HBsAg by silver stained SDS-PAGE shown in (A), and by Western blot using a polyclonal anti-HBsAg antibody shown in (B).

Figure 5 shows the analysis of the interaction between HBsAg and NE droplets. Measurement of the surface charge by zeta potential shown in (A). Particle size distribution measured using a laser diffraction particle-sizer of HBsAg alone shown in (B), NE alone shown in (C), and NE with 10 μ g/ml of HBsAg shown in (D). Calorimetric titration of HBsAg with NE shown in (E). The upper panel shows differences between the sample and reference cell containing PBS. The lower panel shows enthalpy per injection of NE injected versus injection number.

Figure 6 shows the development of IgG response in serum. The effect of varying the NE adjuvant concentration from 0% to 40% is shown in (A). The effect of antigen dose escalation from 1 μ g to 40 μ g of HBsAg mixed with 20% NE is shown in (B). Arrows indicate vaccine administration.

Figure 7 shows a comparison of mucosal NE-based vaccine with conventional aluminum-based injectible HBsAg vaccine. A time course of antibody response for mice immunized with 20 μ g HBsAg antigen mixed with 20% NE for intranasal administration (HBsAg-NE), or adsorbed on aluminum hydroxide (HBsAg-Alu) for intramuscular injections is shown in (A). Avidity of anti-HBsAg IgG: Analysis of sera from mice immunized i.n.

with HBsAg-NE and with i.m. injections of HBsAg-Alu vaccines is shown in (B). An analysis of serum anti-HBsAg IgG subclass pattern for mice immunized nasally with HBsAg-NE or injected i.m. with HBsAg-Alu vaccine is shown in (C).

Figure 8 shows a characterization of immune to HBsAg-NE performed in BAL fluids obtained 23 weeks after i.n immunization with HBsAg-NE vaccines. Anti-HBsAg IgA concentrations are shown in (A). Anti-HBsAg IgG antibody concentrations are shown in (B). Pattern of Th1 (IFN- and TNF-) and Th2 (IL-4, IL-5 and IL-10) antigen-specific cytokine expression *in vitro* in splenocytes from mice intranasally immunized with HBsAg-NE is shown in (C).

Figure 9 shows immunogenicity in rats and guinea pigs immunized intranasally with either 5 g or 20 g HBsAg mixed with 20% NE.

Figure 10 shows an *in vitro* comparison of HBsAg-NE stored at test temperature conditions (1: fresh, 2: 4°C, 3: 25°C and 4: 40°C) by SDS-PAGE (S) or Western blot (W). Lanes are labeled according to sample storage conditions as follows- 1: fresh, 2: 4°C, 3: 25°C and 4: 40°C. Samples were stored for (A) 6 weeks, (B) 6 months (24 weeks), or (C) 1 year (52 weeks) at the three test temperatures. A particle size comparison of NE alone, freshly mixed HBsAg-NE, and HBsAg-NE formulation stored up to a year.

Figure 11 shows an *in vivo* analysis of HBsAg-NE stability. HBsAg specific antibody responses to freshly prepared HBsAg-NE or HBsAg-NE stored under real-time (4°C), accelerated (25°C) and stressed (40°C) temperature conditions. Comparison of serum IgG elicited by freshly prepared HBsAg-NE to formulation stored for (A) 6 weeks, (B) 3 months, (C) 6 months or (D) 1 year at indicated temperatures are shown.

Figure 12 shows particle sizing of various different lots of HBsAg (Lots D (A), F (B), and G (C)). Particle sizes were measured by quasi-elastic light scattering by using a Malvern ZETASIZER ZS. AFM images of Lots D (D), F (E), and G (F). HBsAg was imaged on silica in tapping mode using a Multimode Nanoscope IIIA AFM. The radial size distributions of the protein complexes (G) were calculated from the images.

Figure 13 shows the immunogenicity of HBsAg (Lot F)-NE in rats. Rats were immunized intranasally with 100 l of HBsAg-NE and boosted at 4 weeks. Data shown represent the serum anti-HBsAg IgG titer at 9 weeks following prime vaccination.

Figure 14 shows particle sizing of HBsAg Lot F. HBsAg Lot F was analyzed by particle sizing after a 24 hours dialysis to phosphate buffered Saline (PBS) and followed by

ultrasonication using a Fisher bath sonicator for 5 minutes.

Figure 15 shows PAGE analysis of recombinant antigens. Protein samples loaded at 0.5ug/well. Antigens appear intact and do not contain contaminants.

Figure 16 shows the immunogenicity of trivalent NE-based nasal vaccines composed of HCR/A1, PA and either LcrV10 shown in (A) or LcrV shown in (B) mixed with nanoemulsion.

Figure 17 shows a comparison of the immunogenicity if trivalent (HCR/A1-LcrV-PA)-NE vaccines versus monovalent NE-based vaccines for PA shown in (A), HCR/A1 shown in (B), and LcrV and LcrV10 shown in (C).

Figure 18 shows that immunogenicity of nasal NE-adjuvanted trivalent vaccine was compared to that of intramuscular injected alum-based vaccine.

Figure 19 shows the mucosal response following nasal immunization with NE-based vaccines for mucosal IgG shown in (A) and IgA shown in (B).

Figure 20 shows the Th-1 polarization of the immune response of subjects administered a trivalent vaccine of the invention.

Figure 21 shows the Th-1 type anti-PA IgG subclass pattern in mice immunized with the trivalent mucosal vaccine.

Figure 22 shows the Th-1 type anti-LcrV and Lcrv10 IgG subclass pattern in mice immunized with the trivalent mucosal vaccine.

Figure 23 shows Th-1 type anti-HCR/A1 subclass pattern in mice immunized with the trivalent mucosal vaccine.

Figure 24 shows protective immunity generated in a subject administered an intranasal trivalent vaccine of the invention against challenge with live *Yersinia pestis* strain C092.

Figure 25 shows protective immunity generated in a subject administered an intranasal trivalent vaccine of the invention against challenge with live *Yersinia pestis* strain C092.

Figure 26 shows protective immunity generated in a subject administered an intranasal trivalent vaccine of the invention against challenge with *B. anthracis* Ames spores.

GENERAL DESCRIPTION OF THE INVENTION

The present invention provides methods and compositions for the stimulation of

immune responses. Specifically, the present invention provides immunogenic compositions and methods of using the same to induce immune responses (e.g., immunity (e.g., protective immunity)) against Hepatitis B virus (HBV)) and/or against one or a plurality of pathogens (e.g., vaccinia virus, H5N1 influenza virus, *Bacillus anthracis*, *C. botulinum*, *Y. pestis*,
5 Hepatitis B, and/or HIV, etc.) in a subject. Compositions and methods of the present invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine (e.g., vaccination)) and research applications.

In some embodiments, one or a plurality of pathogens are mixed with a nanoemulsion prior to administration for a time period sufficient to inactivate the one or plurality of
10 pathogens. In some embodiments, one or a plurality of protein components (e.g., isolated and/or purified and/or recombinant protein) from one or a plurality of pathogens are mixed with the nanoemulsion.

Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in
15 some embodiments, NE treatment (e.g., neutralization of one or more pathogens) with a NE of the present invention) preserves important antigenic epitopes (e.g., recognizable by a subject's immune system), stabilizing their hydrophobic and hydrophilic components in the oil and water interface of the emulsion (e.g., thereby providing one or more immunogens (e.g., stabilized antigens) against which a subject can mount an immune response). In other
20 embodiments, because NE formulations penetrate the mucosa through pores, they may carry immunogens to the submucosal location of dendritic cells (e.g., thereby initiating and/or stimulating an immune response). Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, combining a NE and one or a plurality
25 of immunogenic proteins (e.g., rPA from *B. anthracis*, rHCR/A1 (fragment of *C. botulinum* neurotoxin), rLcrV (or LcrV10) protein of *Y. pestis* and/or gp120 from HIV, etc.) stabilizes the immunogens and provides a proper immunogenic material for generation of an immune response.

Furthermore, in some embodiments, a composition of the present invention (e.g., a
30 composition comprising a NE and one or a plurality of immunogens) induces (e.g., when administered to a subject) both systemic and mucosal immunity. Thus, in some preferred embodiments, administration of a composition of the present invention to a subject results in

protection against an exposure (e.g., a lethal mucosal exposure) to one or a plurality of pathogens (e.g., one or a plurality of viruses and/or bacteria). Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, mucosal administration (e.g., vaccination) provides protection against pathogen infection (e.g., that initiates at a mucosal surface).

Although it has heretofore proven difficult to stimulate secretory IgA responses and protection against pathogens that invade at mucosal surfaces (See, e.g., Mestecky et al, Mucosal Immunology. 3ed edn. (Academic Press, San Diego, 2005)), the present invention provides compositions and methods for stimulating mucosal immunity (e.g., a protective IgA response) against one or a plurality of pathogens in a subject.

In some embodiments, the present invention provides a composition (e.g., comprising a NE and one or a plurality of immunogens) to serve as a mucosal vaccine. This material can easily be produced from purified virus and/or protein or recombinant protein and induces both mucosal and systemic immunity. The ability to produce this formulation rapidly and administer it via mucosal instillation provides vaccines that can be used for general vaccination needs as well as in large-scale outbreaks or emergent situations.

DEFINITIONS

To facilitate an understanding of the present invention, a number of terms and phrases are defined below:

As used herein, the term "microorganism" refers to any species or type of microorganism, including but not limited to, bacteria, viruses, archaea, fungi, protozoans, mycoplasma, prions, and parasitic organisms. The term microorganism encompasses both those organisms that are in and of themselves pathogenic to another organism (e.g., animals, including humans, and plants) and those organisms that produce agents that are pathogenic to another organism, while the organism itself is not directly pathogenic or infective to the other organism.

As used herein the term "pathogen," and grammatical equivalents, refers to an organism (e.g., biological agent), including microorganisms, that causes a disease state (e.g., infection, pathologic condition, disease, etc.) in another organism (e.g., animals and plants) by directly infecting the other organism, or by producing agents that causes disease in another organism (e.g., bacteria that produce pathogenic toxins and the like). "Pathogens" include,

but are not limited to, viruses, bacteria, archaea, fungi, protozoans, mycoplasma, prions, and parasitic organisms.

The terms "bacteria" and "bacterium" refer to all prokaryotic organisms, including those within all of the phyla in the Kingdom Procaryotae. It is intended that the term encompass all microorganisms considered to be bacteria including *Mycoplasma*, *Chlamydia*, *Actinomyces*, *Streptomyces*, and *Rickettsia*. All forms of bacteria are included within this definition including cocci, bacilli, spirochetes, spheroplasts, protoplasts, etc.

As used herein, the term "fungi" is used in reference to eukaryotic organisms such as molds and yeasts, including dimorphic fungi.

As used herein the terms "hepatitis B surface antigen" or "HBsAg" includes any HBsAg antigen or fragment thereof displaying the antigenicity of HBV surface antigen. HBsAg may be obtained or derived from (e.g., recombinantly derived from) any of the serotypes of hepatitis B including, but not limited to, serotypes adr, adw, ayr, ayw, or from any of the various hepatitis B genotypes including, but not limited to, genotypes (A-H). It is to be further understood that in addition to the 226 amino acid sequence of the HBsAg S antigen (See, e.g., Tiollais et. al. Nature, 317, 489 (1985) and references therein) HBsAg may contain all or part of a pre-S sequence as described in U.S. Patent Application Publication No. 20090123496 and in EP-A-0 278 940. HBsAg as herein described can also refer to variants, for example the "escape mutant" described in WO 91/14703. HBsAg also refers to polypeptides described in EP 0 198 474 or EP 0 304 578

As used herein the terms "disease" and "pathologic condition" are used interchangeably, unless indicated otherwise herein, to describe a deviation from the condition regarded as normal or average for members of a species or group (e.g., humans), and which is detrimental to an affected individual under conditions that are not inimical to the majority of individuals of that species or group. Such a deviation can manifest as a state, signs, and/or symptoms (e.g., diarrhea, nausea, fever, pain, blisters, boils, rash, immune suppression, inflammation, etc.) that are associated with any impairment of the normal state of a subject or of any of its organs or tissues that interrupts or modifies the performance of normal functions. A disease or pathological condition may be caused by or result from contact with a microorganism (e.g., a pathogen or other infective agent (e.g., a virus or bacteria)), may be responsive to environmental factors (e.g., malnutrition, industrial hazards, and/or climate), may be responsive to an inherent defect of the organism (e.g., genetic anomalies) or to

combinations of these and other factors.

The terms "host" or "subject," as used herein, refer to an individual to be treated by (e.g., administered) the compositions and methods of the present invention. Subjects include, but are not limited to, mammals (*e.g.*, murines, simians, equines, bovines, porcines, canines, felines, and the like), and most preferably includes humans. In the context of the invention, the term "subject" generally refers to an individual who will be administered or who has been administered one or more compositions of the present invention (*e.g.*, a composition for inducing an immune response).

As used herein, the terms "inactivating," "inactivation" and grammatical equivalents, when used in reference to a microorganism (*e.g.*, a pathogen (*e.g.*, a bacterium or a virus)), refer to the killing, elimination, neutralization and/or reducing of the capacity of the microorganism (*e.g.*, a pathogen (*e.g.*, a bacterium or a virus)) to infect and/or cause a pathological response and/or disease in a host. For example, in some embodiments, the present invention provides a composition comprising nanoemulsion (NE)-inactivated vaccinia virus (VV). Accordingly, as referred to herein, compositions comprising "NE-inactivated VV," "NE-killed V," "NE-neutralized V" or grammatical equivalents refer to compositions that, when administered to a subject, are characterized by the absence of, or significantly reduced presence of, VV replication (*e.g.*, over a period of time (*e.g.*, over a period of days, weeks, months, or longer)) within the host.

As used herein, the term "fusigenic" is intended to refer to an emulsion that is capable of fusing with the membrane of a microbial agent (*e.g.*, a bacterium or bacterial spore). Specific examples of fusigenic emulsions are described herein.

As used herein, the term "lysogenic" refers to an emulsion (*e.g.*, a nanoemulsion) that is capable of disrupting the membrane of a microbial agent (*e.g.*, a virus (*e.g.*, viral envelope) or a bacterium or bacterial spore). In preferred embodiments of the present invention, the presence of a lysogenic and a fusigenic agent in the same composition produces an enhanced inactivating effect compared to either agent alone. Methods and compositions (*e.g.*, for inducing an immune response (*e.g.*, used as a vaccine) using this improved antimicrobial composition are described in detail herein.

The term "emulsion," as used herein, includes classic oil-in-water or water in oil dispersions or droplets, as well as other lipid structures that can form as a result of hydrophobic forces that drive apolar residues (*e.g.*, long hydrocarbon chains) away from

water and drive polar head groups toward water, when a water immiscible oily phase is mixed with an aqueous phase. These other lipid structures include, but are not limited to, unilamellar, paucilamellar, and multilamellar lipid vesicles, micelles, and lamellar phases. Similarly, the term "nanoemulsion," as used herein, refers to oil-in-water dispersions comprising small lipid structures. For example, in preferred embodiments, the nanoemulsions comprise an oil phase having droplets with a mean particle size of approximately 0.1 to 5 microns (e.g., 150 +/-25 nm in diameter), although smaller and larger particle sizes are contemplated. The terms "emulsion" and "nanoemulsion" are often used herein, interchangeably, to refer to the nanoemulsions of the present invention.

As used herein, the terms "contact," "contacted," "expose," and "exposed," when used in reference to a nanoemulsion and a live microorganism, refer to bringing one or more nanoemulsions into contact with a microorganism (e.g., a pathogen) such that the nanoemulsion inactivates the microorganism or pathogenic agent, if present. The present invention is not limited by the amount or type of nanoemulsion used for microorganism inactivation. A variety of nanoemulsion that find use in the present invention are described herein and elsewhere (e.g., nanoemulsions described in U.S. Pat. Apps. 20020045667 and 20040043041, and U.S. Pat. Nos. 6,015,832, 6,506,803, 6,635,676, and 6,559,189).

Ratios and amounts of nanoemulsion (e.g., sufficient for inactivating the microorganism (e.g., virus inactivation)) and microorganisms (e.g., sufficient to provide an antigenic composition (e.g., a composition capable of inducing an immune response)) are contemplated in the present invention including, but not limited to, those described herein.

The term "surfactant" refers to any molecule having both a polar head group, which energetically prefers solvation by water, and a hydrophobic tail that is not well solvated by water. The term "cationic surfactant" refers to a surfactant with a cationic head group. The term "anionic surfactant" refers to a surfactant with an anionic head group.

The terms "Hydrophile-Lipophile Balance Index Number" and "HLB Index Number" refer to an index for correlating the chemical structure of surfactant molecules with their surface activity. The HLB Index Number may be calculated by a variety of empirical formulas as described, for example, by Meyers, (See, e.g., Meyers, *Surfactant Science and Technology*, VCH Publishers Inc., New York, pp. 231-245 (1992)).

As used herein where appropriate, the HLB Index Number of a surfactant is the

HLB Index Number assigned to that surfactant in McCutcheon's Volume 1: Emulsifiers and Detergents North American Edition, 1996. The HLB

Index Number ranges from 0 to about 70 or more for commercial surfactants. Hydrophilic surfactants with high solubility in water and solubilizing properties are at the high end of the scale, while surfactants with low solubility in water that are good solubilizers of water in oils are at the low end of the scale.

As used herein the term "interaction enhancers" refers to compounds that act to enhance the interaction of an emulsion with a microorganism (e.g., with a cell wall of a bacteria (e.g., a Gram negative bacteria) or with a viral envelope (e.g., Vaccinia virus envelope)). Contemplated interaction enhancers include, but are not limited to, chelating agents (e.g., ethylenediaminetetraacetic acid (EDTA), ethylenebis(oxyethylenenitrilo)tetraacetic acid (EGTA), and the like) and certain biological agents (e.g., bovine serum albumin (BSA) and the like).

The terms "buffer" or "buffering agents" refer to materials, that when added to a solution, cause the solution to resist changes in pH.

The terms "reducing agent" and "electron donor" refer to a material that donates electrons to a second material to reduce the oxidation state of one or more of the second material's atoms.

The term "monovalent salt" refers to any salt in which the metal (e.g., Na, K, or Li) has a net 1+ charge in solution (i.e., one more proton than electron).

The term "divalent salt" refers to any salt in which a metal (e.g., Mg, Ca, or Sr) has a net 2+ charge in solution.

The terms "chelator" or "chelating agent" refer to any materials having more than one atom with a lone pair of electrons that are available to bond to a metal ion.

The term "solution" refers to an aqueous or non-aqueous mixture.

As used herein, the term "a composition for inducing an immune response" refers to a composition that, once administered to a subject (e.g., once, twice, three times or more (e.g., separated by weeks, months or years)), stimulates, generates and/or elicits an immune response in the subject (e.g., resulting in total or partial immunity to a microorganism (e.g., pathogen) capable of causing disease). In preferred embodiments of the invention, the composition comprises a nanoemulsion and an immunogen. In further preferred embodiments, the composition comprising a nanoemulsion and an immunogen comprises one

or more other compounds or agents including, but not limited to, therapeutic agents, physiologically tolerable liquids, gels, carriers, diluents, adjuvants, excipients, salicylates, steroids, immunosuppressants, immunostimulants, antibodies, cytokines, antibiotics, binders, fillers, preservatives, stabilizing agents, emulsifiers, and/or buffers. An immune response
5 may be an innate (e.g., a non-specific) immune response or a learned (e.g., acquired) immune response (e.g. that decreases the infectivity, morbidity, or onset of mortality in a subject (e.g., caused by exposure to a pathogenic microorganism) or that prevents infectivity, morbidity, or onset of mortality in a subject (e.g., caused by exposure to a pathogenic microorganism)).

Thus, in some preferred embodiments, a composition comprising a nanoemulsion and an
10 immunogen is administered to a subject as a vaccine (e.g., to prevent or attenuate a disease (e.g., by providing to the subject total or partial immunity against the disease or the total or partial attenuation (e.g., suppression) of a sign, symptom or condition of the disease.

As used herein, the term "adjuvant" refers to any substance that can stimulate an immune response (e.g., a mucosal immune response). Some adjuvants can cause activation
15 of a cell of the immune system (e.g., an adjuvant can cause an immune cell to produce and secrete a cytokine). Examples of adjuvants that can cause activation of a cell of the immune system include, but are not limited to, saponins purified from the bark of the *Q. saponaria* tree, such as QS21 (a glycolipid that elutes in the 21st peak with HPLC fractionation; Aquila Biopharmaceuticals, Inc., Worcester, Mass.); poly(di(carboxylatophenoxy)phosphazene
20 (PCPP polymer; Virus Research Institute, USA); derivatives of lipopolysaccharides such as monophosphoryl lipid A (MPL; Ribi ImmunoChem Research, Inc., Hamilton, Mont.), muramyl dipeptide (MDP; Ribi) and threonyl-muramyl dipeptide (t-MDP; Ribi); OM-174 (a glucosamine disaccharide related to lipid A; OM Pharma SA, Meyrin, Switzerland); and Leishmania elongation factor (a purified Leishmania protein; Corixa Corporation, Seattle,
25 Wash.). Traditional adjuvants are well known in the art and include, for example, aluminum phosphate or hydroxide salts ("alum"). In some embodiments, compositions of the present invention (e.g., comprising HIV or an immunogenic epitope thereof (e.g., gp120)) are administered with one or more adjuvants (e.g., to skew the immune response towards a Th1 or Th2 type response).

30 As used herein, the term "an amount effective to induce an immune response" (e.g., of a composition for inducing an immune response), refers to the dosage level required (e.g., when administered to a subject) to stimulate, generate and/or elicit an immune response in the

subject. An effective amount can be administered in one or more administrations (e.g., via the same or different route), applications or dosages and is not intended to be limited to a particular formulation or administration route.

As used herein, the term "under conditions such that said subject generates an
5 immune response" refers to any qualitative or quantitative induction, generation, and/or stimulation of an immune response (e.g., innate or acquired).

As used herein, the term "immune response" refers to a response by the immune system of a subject. For example, immune responses include, but are not limited to, a detectable alteration (e.g., increase) in Toll receptor activation, lymphokine (e.g., cytokine
10 (e.g., Th1 or Th2 type cytokines) or chemokine) expression and/or secretion, macrophage activation, dendritic cell activation, T cell activation (e.g., CD4+ or CD8+ T cells), NK cell activation, and/or B cell activation (e.g., antibody generation and/or secretion). Additional examples of immune responses include binding of an immunogen (e.g., antigen (e.g., immunogenic polypeptide)) to an MHC molecule and inducing a cytotoxic T lymphocyte
15 ("CTL") response, inducing a B cell response (e.g., antibody production), and/or T-helper lymphocyte response, and/or a delayed type hypersensitivity (DTH) response against the antigen from which the immunogenic polypeptide is derived, expansion (e.g., growth of a population of cells) of cells of the immune system (e.g., T cells, B cells (e.g., of any stage of development (e.g., plasma cells), and increased processing and presentation of antigen by
20 antigen presenting cells. An immune response may be to immunogens that the subject's immune system recognizes as foreign (e.g., non-self antigens from microorganisms (e.g., pathogens), or self-antigens recognized as foreign). Thus, it is to be understood that, as used herein, "immune response" refers to any type of immune response, including, but not limited to, innate immune responses (e.g., activation of Toll receptor signaling cascade) cell-
25 mediated immune responses (e.g., responses mediated by T cells (e.g., antigen-specific T cells) and non-specific cells of the immune system) and humoral immune responses (e.g., responses mediated by B cells (e.g., via generation and secretion of antibodies into the plasma, lymph, and/or tissue fluids). The term "immune response" is meant to encompass all aspects of the capability of a subject's immune system to respond to antigens and/or
30 immunogens (e.g., both the initial response to an immunogen (e.g., a pathogen) as well as acquired (e.g., memory) responses that are a result of an adaptive immune response).

As used herein, the term "immunity" refers to protection from disease (e.g.,

preventing or attenuating (e.g., suppression) of a sign, symptom or condition of the disease) upon exposure to a microorganism (e.g., pathogen) capable of causing the disease. Immunity can be innate (e.g., non-adaptive (e.g., non-acquired) immune responses that exist in the absence of a previous exposure to an antigen) and/or acquired (e.g., immune responses that are mediated by B and T cells following a previous exposure to antigen (e.g., that exhibit increased specificity and reactivity to the antigen)).

As used herein, the term "immunogen" refers to an agent (e.g., a microorganism (e.g., bacterium, virus or fungus) and/or portion or component thereof (e.g., a protein antigen (e.g., gp120 or rPA))) that is capable of eliciting an immune response in a subject. In preferred embodiments, immunogens elicit immunity against the immunogen (e.g., microorganism (e.g., pathogen or a pathogen product)) when administered in combination with a nanoemulsion of the present invention.

As used herein, the term "pathogen product" refers to any component or product derived from a pathogen including, but not limited to, polypeptides, peptides, proteins, nucleic acids, membrane fractions, and polysaccharides.

As used herein, the term "enhanced immunity" refers to an increase in the level of adaptive and/or acquired immunity in a subject to a given immunogen (e.g., microorganism (e.g., pathogen)) following administration of a composition (e.g., composition for inducing an immune response of the present invention) relative to the level of adaptive and/or acquired immunity in a subject that has not been administered the composition (e.g., composition for inducing an immune response of the present invention).

As used herein, the terms "purified" or "to purify" refer to the removal of contaminants or undesired compounds from a sample or composition. As used herein, the term "substantially purified" refers to the removal of from about 70 to 90 %, up to 100%, of the contaminants or undesired compounds from a sample or composition.

As used herein, the terms "administration" and "administering" refer to the act of giving a composition of the present invention (e.g., a composition for inducing an immune response (e.g., a composition comprising a nanoemulsion and an immunogen)) to a subject. Exemplary routes of administration to the human body include, but are not limited to, through the eyes (ophthalmic), mouth (oral), skin (transdermal), nose (nasal), lungs (inhalant), oral mucosa (buccal), ear, rectal, by injection (e.g., intravenously, subcutaneously, intraperitoneally, etc.), topically, and the like.

As used herein, the terms "co-administration" and "co-administering" refer to the administration of at least two agent(s) (e.g., a composition comprising a nanoemulsion and an immunogen and one or more other agents - e.g., an adjuvant) or therapies to a subject. In some embodiments, the co-administration of two or more agents or therapies is concurrent.

5 In other embodiments, a first agent/therapy is administered prior to a second agent/therapy. In some embodiments, co-administration can be via the same or different route of administration. Those of skill in the art understand that the formulations and/or routes of administration of the various agents or therapies used may vary. The appropriate dosage for co-administration can be readily determined by one skilled in the art. In some embodiments, 10 when agents or therapies are co-administered, the respective agents or therapies are administered at lower dosages than appropriate for their administration alone. Thus, co-administration is especially desirable in embodiments where the co-administration of the agents or therapies lowers the requisite dosage of a potentially harmful (e.g., toxic) agent(s), and/or when co-administration of two or more agents results in sensitization of a subject to 15 beneficial effects of one of the agents via co-administration of the other agent. In other embodiments, co-administration is preferable to elicit an immune response in a subject to two or more different immunogens (e.g., microorganisms (e.g., pathogens)) at or near the same time (e.g., when a subject is unlikely to be available for subsequent administration of a second, third, or more composition for inducing an immune response).

20 As used herein, the term "topically" refers to application of a compositions of the present invention (e.g., a composition comprising a nanoemulsion and an immunogen) to the surface of the skin and/or mucosal cells and tissues (e.g., alveolar, buccal, lingual, masticatory, vaginal or nasal mucosa, and other tissues and cells which line hollow organs or body cavities).

25 In some embodiments, the compositions of the present invention are administered in the form of topical emulsions, injectable compositions, ingestible solutions, and the like. When the route is topical, the form may be, for example, a spray (e.g., a nasal spray), a cream, or other viscous solution (e.g., a composition comprising a nanoemulsion and an immunogen in polyethylene glycol).

30 The terms "pharmaceutically acceptable" or "pharmacologically acceptable," as used herein, refer to compositions that do not substantially produce adverse reactions (e.g., toxic, allergic or immunological reactions) when administered to a subject.

As used herein, the term "pharmaceutically acceptable carrier" refers to any of the standard pharmaceutical carriers including, but not limited to, phosphate buffered saline solution, water, and various types of wetting agents (e.g., sodium lauryl sulfate), any and all solvents, dispersion media, coatings, sodium lauryl sulfate, isotonic and absorption delaying agents, disintegrants (e.g., potato starch or sodium starch glycolate), polyethylene glycol, and the like.. The compositions also can include stabilizers and preservatives. Examples of carriers, stabilizers and adjuvants have been described and are known in the art (See e.g., Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, Pa. (1975)).

As used herein, the term "pharmaceutically acceptable salt" refers to any salt (e.g., obtained by reaction with an acid or a base) of a composition of the present invention that is physiologically tolerated in the target subject. "Salts" of the compositions of the present invention may be derived from inorganic or organic acids and bases. Examples of acids include, but are not limited to, hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycolic, lactic, salicylic, succinic, toluene-p-sulfonic, tartaric, acetic, citric, methanesulfonic, ethanesulfonic, formic, benzoic, malonic, sulfonic, naphthalene-2-sulfonic, benzenesulfonic acid, and the like. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compositions of the invention and their pharmaceutically acceptable acid addition salts.

Examples of bases include, but are not limited to, alkali metal (e.g., sodium) hydroxides, alkaline earth metal (e.g., magnesium) hydroxides, ammonia, and compounds of formula NW_4^+ , wherein W is C_{1-4} alkyl, and the like.

Examples of salts include, but are not limited to: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, flucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, chloride, bromide, iodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, palmoate, pectinate, persulfate, phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate, undecanoate, and the like. Other examples of salts include anions of the compounds of the present invention compounded with a suitable cation such as Na^+ , NH_4^+ , and NW_4^+ (wherein W is a C_{1-4} alkyl

group), and the like. For therapeutic use, salts of the compounds of the present invention are contemplated as being pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound.

5 For therapeutic use, salts of the compositions of the present invention are contemplated as being pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable composition.

As used herein, the term "at risk for disease" refers to a subject that is predisposed to
10 experiencing a particular disease. This predisposition may be genetic (e.g., a particular genetic tendency to experience the disease, such as heritable disorders), or due to other factors (e.g., environmental conditions, exposures to detrimental compounds present in the environment, etc.). Thus, it is not intended that the present invention be limited to any particular risk (e.g., a subject may be "at risk for disease" simply by being exposed to and
15 interacting with other people), nor is it intended that the present invention be limited to any particular disease.

"Nasal application", as used herein, means applied through the nose into the nasal or sinus passages or both. The application may, for example, be done by drops, sprays, mists, coatings or mixtures thereof applied to the nasal and sinus passages.

20 "Vaginal application", as used herein, means applied into or through the vagina so as to contact vaginal mucosa. The application may contact the urethra, cervix, fornix, uterus or other area surrounding the vagina. The application may, for example, be done by drops, sprays, mists, coatings, lubricants or mixtures thereof applied to the vagina or surrounding tissue.

25 As used herein, the term "kit" refers to any delivery system for delivering materials. In the context of immunogenic agents (e.g., compositions comprising a nanoemulsion and an immunogen), such delivery systems include systems that allow for the storage, transport, or delivery of immunogenic agents and/or supporting materials (e.g., written instructions for using the materials, etc.) from one location to another. For example, kits include one or more
30 enclosures (e.g., boxes) containing the relevant immunogenic agents (e.g., nanoemulsions) and/or supporting materials. As used herein, the term "fragmented kit" refers to delivery systems comprising two or more separate containers that each contain a subportion of the

total kit components. The containers may be delivered to the intended recipient together or separately. For example, a first container may contain a composition comprising a nanoemulsion and an immunogen for a particular use, while a second container contains a second agent (e.g., an antibiotic or spray applicator). Indeed, any delivery system comprising two or more separate containers that each contains a subportion of the total kit components are included in the term "fragmented kit." In contrast, a "combined kit" refers to a delivery system containing all of the components of an immunogenic agent needed for a particular use in a single container (e.g., in a single box housing each of the desired components). The term "kit" includes both fragmented and combined kits.

DETAILED DESCRIPTION OF THE INVENTION

Infection with hepatitis B virus (HBV) remains an important global health concern, despite the availability of multiple prophylactic vaccines. The World Health Organization (WHO) estimates that more than 2 billion persons have been infected with the virus. The current prophylactic vaccines require a regimen of three intramuscular (i.m.) injections, have a 10%-15% non-responders rate, and are ineffective for limiting HBV replication in chronic carriers (See, e.g., Assad, S. and A. Francis, *Vaccine*, 1999. 18(1-2): p. 57-67; Floreani, A., et al., *Vaccine*, 2004. 22(5-6): p. 608-611; Gesemann, M. and N. Scheiermann, *Vaccine*, 1995. 13(5): p. 443-447). Large scale vaccination programs are also limited in developing populations due to compliance issues secondary to the three dose vaccination schedule, the requirement for cold storage and the availability of sterile needles (See, e.g., Weakly Epidemiological Record 1992, World Health Organization: Thirteenth meeting of the Global Advisory Group RPI. p. 1-12; CDC, Global progress toward universal childhood hepatitis B vaccination. *MMWR*, 2003. 52(36): p. 868-870). This has limited the use of hepatitis B vaccine in these populations and is partly responsible for 8%-10% of the population in areas of Africa, Asia and South America being chronically infected with HBV (See, e.g., Hepatitis B Fact sheet no. 204. 2000, World Health Organization). Chronic HBV infection increases the risk of developing liver cirrhosis, hepatocellular carcinoma and other associated complications leading to increased mortality (See, e.g., Chisari, F.V. and C. Ferrari, *Ann. Rev. Immunol*, 1995. 13(1): p. 29-60).

Hepatitis B surface antigen (HBsAg) is a major structural protein of HBV and is a protective immunogen in experimental animals and in humans (See, e.g., Peterson, D., L,

BioEssays, 1987. 6(6): p. 258-262; Schirmbeck, R., et al., J. Immunol, 1994. 152(3): p. 1110-1119; Seeger, C. and W.S. Microbiol. Mol. Biol. Rev., 2000. 64(1): p. 51-68). The hepatitis B surface (HBs) proteins are synthesized as large (L), medium (M) and small (S) envelope sub-units, which self assemble into virus-like lipid-anchored particles (about 22 nm in size)

5 (See, e.g., Gilbert, R.J.C., et al., PNAS, 2005. 102(41): p. 14783-14788; Woo, W.-P., et al., J. Virol, 2006. 80(8): p. 3975-3984). The majority of commercially available recombinant HBsAg vaccines (including RECOMBIVAX HB; MERCK, and ENGERIX-B; GSK) consist of yeast derived HBs-S antigen particles adsorbed to an aluminum salt (alum) adjuvant (See, e.g., Assad, S. and A. Francis, Vaccine, 1999. 18(1-2): p. 57-67; Lemon, S.M. and D.L.

10 Thomas, N Engl J Med, 1997. 336(3): p. 196-204). While alum is generally well tolerated and is considered safe, some adverse effects have been reported (See, e.g., Gherardi, R.K., et al., Brain, 2001. 124(9): p. 1821-1831; Pittman, P.R., Vaccine, 2002. 20(Supplement 3): p. S48-S50. Further, alum has been shown to elicit predominantly a Th2 polarization of immune response, which is associated with cellular immunity that is ineffective at producing

15 CD8 responses to virally infected cells (See, e.g., Gupta, R.K., Advanced Drug Delivery Reviews, 1998. 32(3): p. 155-172). Currently available hepatitis B vaccines have comparable thermo-stability profiles requiring unbroken cold chain storage (between 2°C and 8°C) in order to retain potency (See, e.g., Hilleman, M.R., Vaccine, 2001. 19(15-16): p. 1837-1848). The higher costs associated with guaranteed cold chain, from point of manufacture to point of

20 use, also contribute to the inaccessibility of these vaccines. Thus, an efficacious vaccine requiring fewer injections and a less stringent cold storage requirement would directly benefit underserved populations.

Development of mucosal vaccines remains limited by lack of effective mucosal adjuvants (See, e.g., Chen, H., Journal of Controlled Release, 2000. 67(2-3): p. 117-128;

25 Neutra, M.R. and P.A. Kozlowski, Nat Rev Immunol, 2006. 6(2): p. 148-158). Studies have evaluated several potential mucosal adjuvants for hepatitis B vaccines including recombinant cholera toxin (CT) (See, e.g., Isaka, M., et al., Vaccine, 2001. 19(11-12): p. 1460-1466), lipid microparticles (See, e.g., Saraf, S., et al., Vaccine, 2006. 24(1): p. 45-56), CpG oligonucleotides (See, e.g., McCluskie, M.J. and H.L. Davis, J Immunol, 1998. 161(9): p.

30 4463-4466; Payette, P., et al., Intervirology, 2006. 49(3): p. 144-151), cationic particles (See, e.g., Debin, A., et al., Vaccine, 2002. 20(21-22): p. 2752-2763), PLG microspheres (See, e.g., Jaganathan, K.S. and S.P. Vyas, Vaccine, 2006. 24(19): p. 4201-4211) or hepatitis B core

antigen (HBcAg) (See, e.g., Aguilar, J.C., et al., Biochemical and Biophysical Research Communications, 2003. 310(1): p. 59-63; Aguilar, J.C., et al., Immunol Cell Biol, 2004. 82(5): p. 539-546; Lobaina, Y., et al., Biochemical and Biophysical Research Communications, 2003. 300(3): p. 745-750). CT has been limited from use in humans due to its potential to cause CNS inflammation. Unfortunately, a CpG-adjuvanted injectable hepatitis B vaccine was recently placed on clinical hold due to inflammatory issues in a patient, further calling into question the safety of pro-inflammatory adjuvants. No other adjuvant, with the exception of using HBcAg as an adjuvant, has even been tested in clinical trials (See, e.g., Betancourt, A.A., et al., International Journal of Infectious Diseases, 2008; Zuckerman, J., N., J. Med. Virol., 2006. 78(2): p. 169-177).

Accordingly, in some embodiments, the present invention provides methods and compositions for the stimulation of immune responses. Specifically, the present invention provides immunogenic compositions and methods of using the same to induce immune responses (e.g., immunity (e.g., protective immunity)) against Hepatitis B virus (HBV)).

Compositions and methods of the present invention find use in, among other things, clinical (e.g. therapeutic and preventative medicine (e.g., vaccination)) and research applications.

In some embodiments, the present invention provides methods of inducing an immune response to HBV in a subject (e.g., a human subject) and compositions useful in such methods (e.g., immunogenic composition comprising a nanoemulsion and HBV immunogen (e.g., recombinant, isolated and/or purified HBV immunogen (e.g., HBV antigen (e.g., HBsAg))) (See, e.g., Examples 1-6). The present invention is not limited by the type or source of HBV antigen (e.g., HBsAg). Indeed, any HBV antigen (e.g., HBsAg) or fragment thereof displaying antigenicity (e.g., the antigenicity of HBV surface antigen) may be utilized. HBsAg may be obtained or derived from (e.g., recombinantly derived from) any of the serotypes of hepatitis B including, but not limited to, serotypes adr, adw, ayr, ayw, or from any of the various hepatitis B genotypes including, but not limited to, genotypes (A-H). HBsAg may contain all or part of a pre-S sequence as described in U.S. Patent Application Publication No. 20090123496 and in EP-A-0 278 940.

In some embodiments, methods of inducing an immune response provided by the present invention are used for vaccination. Thus, in some embodiments, the present invention overcomes major drawbacks to conventional HBV vaccines that require

intramuscular immunization, refrigeration of the immunogenic composition and/or at least three different administrations. For example, in some embodiments, the present invention provides an immunogenic composition comprising a nanoemulsion and HBV immunogen that is not administered via injection (e.g., can be nasally administered), that need not be
5 refrigerated for storage and/or transportation, and that produces protective immunity in a subject when administered less than three times.

In some embodiments, an immunogenic composition comprising a nanoemulsion (NE) and HBV immunogen comprises uniform lipid droplets (349 +/-17 nm) associated with HBsAg through electrostatic and hydrophobic interactions (See, e.g., physical
10 characterization using laser particle sizing, zeta potential measurement, isothermal titration calorimetry, and gel electrophoresis of Example 2). However, the present invention is not so limited. For example, in some embodiments, the droplet size is less than 350 nm (e.g., about 325 nm, about 300 nm, about 275 nm, about 250 nm, about 225 nm, about 200 nm or smaller) or larger than 350 nm (e.g., 375 nm, 400 nm, 425 nm, 450 nm, 475 nm, 500 nm, or
15 larger). HBsAg-NE formulations are stable in a broad range of both NE and antigen concentrations and in temperatures ranging from 4 to 40 °C for periods up to 6 weeks. The immunogenicity of compositions comprising nanoemulsion and HBsAg (HBsAg-NE) was evaluated in mice, rats and guinea pigs. Animals immunized intranasally developed robust and sustained systemic IgG, mucosal IgA and strong antigen-specific cellular immune
20 responses (See, e.g., Examples 3 and 4). The serum IgG concentration reached $\geq 10^6$ titers and was comparable in titer to intramuscular vaccination with an alum-adjuvanted vaccine (HBsAg-Alu). Normalization with a standardized human anti-HBsAg serum showed that intranasal NE vaccination correlated with a protective immunity equivalent or greater than 1000 IU/ml. Splenic lymphocytes from nasal HBsAg-NE vaccinated mice produced INF- γ
25 and TNF- α cytokine in response to challenge with HBsAg, and this together with elevated levels of IgG₂ subclass HBsAg-specific antibodies indicated a Th1 polarized immune response. The composition was very stable and retained immunogenicity for a year when stored at 4°C, for 6 months at 25°C and 6 weeks at 40°C. Comprehensive pre-clinical toxicology evaluation in mice, rats, guinea pigs and dogs demonstrated that HBsAg-NE
30 vaccine is safe and well tolerated in multiple animal models (See, e.g., Examples 5 and 6). In some embodiments, the present invention provides needle-free nasal immunization with an immunogenic composition comprising nanoemulsion and HBV immunogen (e.g., HBV

antigen (e.g., HBsAg)) as a safe and effective hepatitis B vaccine and/or as an alternative booster administration for parenteral hepatitis B vaccines. Immunogenic compositions comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) of the present invention induce potent Th1 cellular immunity and also provide therapeutic benefit to patients with chronic hepatitis B infection that lack cellular immune responses (e.g., in order to control viral replication in the subject).

Experiments conducted during development of embodiments of the invention documented the immunogenicity of a novel, mucosal hepatitis B vaccine comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., recombinant HBsAg)). A single nasal immunization of composition comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) produced a rapid induction of serum anti-HBsAg IgG, which is comparable to that achieved with intramuscular (i.m.) vaccination using aluminum salt-based vaccine. Serum IgG responses could be boosted and the titers persisted for 23 weeks. Normalization carried out by comparison to a standardized human anti-HBsAg serum indicated that anti-HBsAg antibody titers in mice immunized with a nasal HBsAg-NE vaccine corresponded to a greater than 1,000 mIU/ml HBsAg IgG concentration in humans which are considered to be seroprotective against HBV infection (See, e.g., Floreani, A., et al., Vaccine, 2004. 22(5-6): p. 608-611; Van Herck, K., et al., Vaccine, 1998. 16(20): p. 1933-1935). An affinity maturation in the antibody response was also observed as serum IgG from animals vaccinated with HBsAg-NE indicated that their avidity matured over time to achieve higher values at 23 weeks than at 5 weeks after vaccination. This is important since functional antibody maturation is considered a significant correlate for the protective efficacy of vaccines (See, e.g., Anttila, et al., Clinical & Experimental Immunology, 1999. 118(3): p. 402-407; Lambert, P.-H. et al., Nat Med, 2005. 11(4 suppl): p. S54-S62). The cross-reactive nature of IgG antibodies against the heterologous *ayw* serotype provides that immunization with one of the HBsAg serotypes produces IgG responses broadly reactive with HBsAg epitope variants (e.g., thereby providing protective immunity against various serotypes of HBV).

The present invention provides that nasal immunization with HBsAg-NE also induced significant mucosal immunity as documented by IgA and IgG detected in BAL fluids. Mucosal immunization with HBsAg-NE also induced antigen-specific T cell responses. *In vitro* stimulation of splenocytes harvested from vaccinated mice with HBsAg resulted in a

cytokine response characterized by significant secretion of hallmark Th1 type cytokines such as IFN- γ and TNF- α , while Th2 type cytokines IL-4, IL-5 and IL-10 showed no antigen-specific response (See, e.g., Leroux-Roels, G., et al., Vaccine, 1994. 12(9): p. 812-818; McClary, H., et al., J. Virol., 2000. 74(5): p. 2255-2264; Wieland, S.F., et al., J. Virol., 2000. 74(9): p. 4165-4173). In addition to enhancing the magnitude of antibody response, nanoemulsion adjuvant had an effect on the pattern of IgG isotypes, as indicated by prevalence of IgG2 over IgG1 subclass in contrast to vaccination with HBsAg-Alu which produced overwhelming titers of IgG1 antibodies (See, e.g., Aguilar, J.C., et al., Immunol Cell Biol, 2004. 82(5): p. 539-546; Brewer, J.M., Immunology Letters, 2006. 102(1): p. 10-15). Prevalence of IgG2b in the overall IgG response provided additional confirmation of a Th1 bias in cellular immunity produced by administration of a composition comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) to subjects. IgG1 remained at significant titers, suggesting the ability to co-activate both Th1 and Th2 immune elements (See, e.g., Khajuria, A., et al., Vaccine, 2007. 25(23): p. 4586-4594).

Thus, in some embodiments, the present invention provides composition comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) and methods of using the same (e.g., for vaccination produced immunity in a subject to HBV) that is compatible with aluminum salt-adjuvanted vaccines, but without the need for injection or an inflammatory adjuvant. The present invention also provides a straight forward approach for formulation of an immunogenic composition (e.g., for use as a Hepatitis B vaccine) that makes it suitable to be produced without special equipment. Thus, in some embodiments, compositions described herein are utilized in developing regions of the world (e.g., where refrigeration of materials is difficult to impossible). The present invention also provides that the physical association of HBsAg with the lipid phase of NE provides stability to the antigen as well as contributing to the adjuvant capability of NE. Thus, the present invention significantly decreases costs associated with conventional HBV vaccines (e.g., the need to maintain conventional vaccines at a refrigerated temperature is overcome by the present invention). Since the HBsAg-NE vaccine retained immunogenicity up to 6 months at 25°C and 3 months at 40°C, in some embodiments, the vaccine does not require refrigeration during distribution.

Adjuvants have been traditionally developed from pro-inflammatory substances, such as a toxin or microbiological component, found to trigger signaling pathways and cytokine

production (See, e.g., Graham, B.S., Plos Medicine, 2006. 3(1): p. e57). Also, enterotoxin-based adjuvants, such as cholera toxin, have been associated with inducing inflammation in the nasal mucosa and with production of the inflammatory cytokines and transport of the vaccine along olfactory neurons into the olfactory bulbs (See, e.g., van Ginkel, F.W., et al., Infect Immun., 2005. 73(10): p. 6892–6902). Some patients treated with a flu vaccine based on one of these toxins (NASALFLU, BERNA Biotech), developed Bell's palsy (See, e.g., Mutsch, M., et al., New England Journal of Medicine, 2004. 350(9): p. 896-903) presumably due to the vaccine in the olfactory bulb. This finding led to NASALFLU being withdrawn. The present invention provides a composition with no significant inflammation in HBsAg-NE treated animals and no evidence of a vaccine composition in the olfactory bulb. Thus the present invention provides, in some embodiments, compositions and methods for inducing immune responses (e.g., immunity to) to HBV utilizing needle-free mucosal administration, induction of systemic immunity comparable with conventional vaccines, as well as mucosal and cellular immune responses that are not elicited by injected, aluminum-based hepatitis vaccines.

In some embodiments, the present invention provides a composition comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) and one or more adjuvants. Preferably the adjuvant is an aluminium salt or a preferential stimulator of Th1 cell response. In some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a nanoemulsion and a hepatitis B antigen (e.g., HBsAg) does not comprise a preservative. For example, in some preferred embodiments, an immunogenic composition (e.g., vaccine) comprising a nanoemulsion and hepatitis B antigen (e.g., HBsAg) does not comprise mercury based preservative (e.g., thiomersal).

Suitable adjuvants for use in eliciting a predominantly Th1-type response include, for example a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL) together with an aluminium salt. Other known adjuvants which preferentially induce a TH1 type immune response include CpG containing oligonucleotides. The oligonucleotides are characterised in that the CpG dinucleotide is unmethylated. Such oligonucleotides are well known and are described in, for example WO 96/02555. Immunostimulatory DNA sequences are also described, for example, by Sato et al., Science 273:352, 1996. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, Mass.), which may be used alone or in

combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. A particularly potent
5 adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210. Thus, in one embodiment of the present invention there is provided a vaccine comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) which additionally comprises a Th1 inducing adjuvant. A preferred embodiment is a vaccine in which the Th1 inducing adjuvant is selected from the group of adjuvants
10 comprising: 3D-MPL, QS21, a mixture of QS21 and cholesterol, and a CpG oligonucleotide.

In some embodiments, the present invention further provides a vaccine formulation comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) of the present invention (e.g., optionally in conjunction with an adjuvant) and additionally comprising one or more antigens selected from the group comprising of: diphtheria toxoid (D),
15 tetanus toxoid (T) acellular pertussis antigens (Pa), inactivated polio virus (IPV), haemophilus influenzae antigen (Hib), hepatitis A antigen, herpes simplex virus (HSV), chlamydia, GSB, HPV, streptococcus pneumoniae and/or neisseria antigens. Antigens conferring protection for other diseases may also be combined in an immunogenic formulation comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g.,
20 HBsAg)) of the present invention.

For example, in one particular embodiment, a vaccine formulation comprises nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) formulations obtainable by a method of manufacture of the present invention in conjunction and an inactivated polio virus (e.g., inactivated using a nanoemulsion described herein or by other
25 means).

The present invention also provides a method of treatment and/or prophylaxis of hepatitis B virus infections, which comprises administering to a human or animal subject, suffering from or susceptible to hepatitis B virus infection, a safe and effective amount of composition comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g.,
30 HBsAg)) of the present invention for the prophylaxis and/or treatment of hepatitis B infection.

The invention further provides the use of a composition comprising nanoemulsion and

HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) of the present invention in the manufacture of a medicament for the treatment of patients suffering from a hepatitis B virus infection, such as chronic hepatitis B virus infection. Immunogenic compositions comprising nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) of the present invention contain an immunoprotective quantity of the antigen and may be prepared by conventional techniques.

In some embodiments, the present invention provides compositions for inducing immune responses comprising a nanoemulsion. The present invention is not limited to any particular nanoemulsion. Indeed, a variety of nanoemulsions find use in the invention including, but not limited to, those described herein and those described elsewhere (e.g., nanoemulsions described in U.S. Pat. Apps. 20020045667 and 20040043041, and U.S. Pat. Nos. 6,015,832, 6,506,803, 6,635,676, and 6,559,189).

HBV immunogens and nanoemulsions of the present invention may be combined in any suitable amount and delivered to a subject utilizing a variety of delivery methods. Any suitable pharmaceutical formulation may be utilized, including, but not limited to, those disclosed herein. Suitable formulations may be tested for immunogenicity using any suitable method. For example, in some embodiments, immunogenicity is investigated by quantitating both antibody titer and specific T-cell responses. Nanoemulsion compositions of the present invention may also be tested in animal models of infectious disease states.

An immunogenic composition comprising a nanoemulsion and HBV immunogen (e.g., HBV antigen (e.g., recombinant HBsAg)) can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce antibodies (e.g., polyclonal antibodies). If desired, a HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, keyhole limpet hemocyanin or other carrier described herein, mixed with a nanoemulsion and administered to a subject. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, nanoemulsions described herein, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially useful.

Monoclonal antibodies that specifically bind to a HBV immunogen (e.g., HBV antigen (e.g., HBsAg)) can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B cell hybridoma
5 technique, and the EBV hybridoma technique (See, e.g., Kohler et al., Nature 256, 495-497, 1985; Kozbor et al., J. Immunol. Methods 81, 3142, 1985; Cote et al., Proc. Natl. Acad. Sci. 80, 2026-2030, 1983; Cole et al., Mol. Cell. Biol. 62, 109-120, 1984).

In addition, techniques developed for the production of "chimeric antibodies," the splicing of mouse antibody genes to human antibody genes to obtain a molecule with
10 appropriate antigen specificity and biological activity, can be used (See, e.g., Morrison et al., Proc. Natl. Acad. Sci. 81, 6851-6855, 1984; Neuberger et al., Nature 312, 604-608, 1984; Takeda et al., Nature 314, 452-454, 1985). Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to
15 human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grafting of entire complementarity determining regions.

20 Alternatively, humanized antibodies can be produced using recombinant methods, as described below. Antibodies which specifically bind to a particular antigen can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Pat. No. 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can
25 be adapted using methods known in the art to produce single chain antibodies which specifically bind to a particular antigen. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (See, e.g., Burton, Proc. Natl. Acad. Sci. 88, 11120-23, 1991).

Single-chain antibodies also can be constructed using a DNA amplification method,
30 such as PCR, using hybridoma cDNA as a template (See, e.g., Thirion et al., 1996, Eur. J. Cancer Prev. 5, 507-11). Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is

taught, for example, in Coloma & Morrison, 1997, Nat. Biotechnol. 15, 159-63. Construction of bivalent, bispecific single-chain antibodies is taught, for example, in Mallender & Voss, 1994, J. Biol. Chem. 269, 199-206.

A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology (See, e.g., Verhaar et al., 1995, Int. J. Cancer 61, 497-501; Nicholls et al., 1993, J. Immunol. Meth. 165, 81-91).

Antibodies which specifically bind to a particular antigen also can be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (See, e.g., Orlandi et al., Proc. Natl. Acad. Sci. 86, 3833-3837, 1989; Winter et al., Nature 349, 293-299, 1991).

Chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, also can be prepared. Antibodies can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which the relevant antigen is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

In some embodiments, the present invention provides compositions for inducing immune responses comprising a nanoemulsion and one or more immunogens (e.g., inactivated pathogens or pathogen products). The present invention provides immunogenic compositions capable of generating an immune response against any number of pathogens (e.g., vaccines for any number of pathogens). A variety of nanoemulsions that find use in the present invention are described herein and elsewhere (e.g., nanoemulsions described in U.S. Pat. Apps. 20020045667 and 20040043041, and U.S. Pat. Nos. 6,015,832, 6,506,803, 6,635,676, and 6,559,189).

Immunogens (e.g., pathogens or pathogen products) and nanoemulsions of the present invention may be combined in any suitable amount utilizing a variety of delivery methods. Any suitable pharmaceutical formulation may be utilized, including, but not limited to, those

disclosed herein. Suitable formulations may be tested for immunogenicity using any suitable method. For example, in some embodiments, immunogenicity is investigated by quantitating both antibody titer and specific T-cell responses. Nanoemulsion compositions of the present invention may also be tested in animal models of infectious disease states. Suitable animal
5 models, pathogens, and assays for immunogenicity include, but are not limited to, those described below.

In some preferred embodiments, the present invention provides methods of inducing an immune response to a plurality of pathogenic organisms in a subject (e.g., a human subject) and compositions useful in such methods (e.g., a nanoemulsion composition
10 comprising a plurality of pathogens and/or plurality of pathogen components (e.g., a plurality of isolated and/or recombinant pathogenic proteins (e.g., protective antigen (PA) of *B. anthracis*, recombinant rHCR/A1 of *C. botulinum*, and recombinant rLcrV (e.g., rLcrV10) protein of *Y. pestis* (e.g., as described in Examples 8 and 9). In some embodiments, methods of inducing an immune response to a plurality of pathogens provided by the present invention
15 are used for vaccination. For example, in some embodiments, the present invention provides a composition comprising a nanoemulsion and one or a plurality of immunogens (e.g., derived from a plurality of pathogens (e.g., one or a plurality of pathogens inactivated by a nanoemulsion of the present invention and/or one or a plurality of protein and/or peptide antigens derived from (e.g., isolated and/or recombinantly produced from) one or a plurality
20 of pathogens); as well as methods of administering the composition (e.g., nasally administering) to a subject under conditions such that the subject generates an immune response to the one or a plurality of pathogens and/or immunogens. In some embodiments, administering comprises mucosal administration. In some embodiments, inducing an immune response induces immunity to each of the plurality of immunogens in the subject. In
25 some embodiments, inducing an immune response to the immunogens induces immunity to the plurality of pathogens from which the immunogens are derived. In some embodiments, immunity comprises systemic immunity. In some embodiments, immunity comprises mucosal immunity. In some embodiments, the immune response comprises increased expression of IFN- γ in the subject. In some embodiments, the immune response comprises a
30 systemic IgG response to the immunogens (e.g., comparable to monovalent vaccine formulations). In some embodiments, the immune response comprises a mucosal IgA response to the immunogens. In some embodiments, the immune response to a multivalent

immunogenic composition is characterized by a balanced Th1/Th2 polarization (e.g., an IgG subclass distribution and cytokine response indicative of a balanced Th1/Th2 response).

Thus, the present invention provides that a multivalent immunogenic composition comprising a nanoemulsion and a plurality of immunogens provides an environment for the plurality of

5 immunogens that maintains the conformation and/or presence of immunogenic epitopes important for protective immunity (e.g., that are recognized by and acted upon by a hosts immune system (e.g., humoral and/or cellular immune response)). In some embodiments, a plurality of immunogens are combined (e.g., in any ratio) to optimize protection against a plurality of environmental pathogens from which the immunogens are derived. For example,

10 in some embodiments, the present invention provides an immunogenic composition that is utilized as a combined anthrax, plague, and botulism vaccine. The present invention also provides that a multivalent immunogenic composition utilized as a vaccine produces immune responses in subjects that are comparable to standard alum-based vaccines. The present invention also provides that a multivalent, nanoemulsion immunogenic composition utilized

15 as a vaccine provides protective immunity to a subject against lethal infection with a pathogenic organisms from which the nanoemulsion immunogenic composition is derived (e.g., plague and/or anthrax). The present invention is not limited to any particular pathogen, combination of pathogens, and/or combination of pathogen components (e.g., peptides, proteins, etc. (e.g., utilized in an immunogenic nanoemulsion composition (e.g., utilized to

20 induce immune responses (e.g., protective immunity) to the pathogen components and/or pathogens))). In some embodiments, the present invention provides a method of inducing an immune response to a plurality of immunogens (e.g., protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) in a subject comprising: providing an immunogenic composition comprising a

25 nanoemulsion and protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin, and administering the composition to the subject under conditions such that the subject generates an immune response toward protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin.

30 Experiments conducted during development of embodiments of the invention documented the immunogenicity of a novel, mucosal multivalent vaccine that is based on a mixture of recombinant antigens (e.g., protective antigen (PA) of *B. anthracis*, LcrV or

LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) and nanoemulsion adjuvant. A single nasal immunization multivalent nanoemulsion mixture produced a rapid induction of serum anti-immunogen IgGs, which was comparable to that achieved with intramuscular (i.m.) vaccination using aluminum salt-based vaccine (See Examples 8 and 9). Mucosal NE-adjuvanted vaccines produced rapid, robust and sustainable serum IgG and mucosal responses to all three antigens. Bronchioalveolar lavages (BAL) and serum were analyzed for IgG and IgA. BAL and serum IgA levels indicated presence of mucosal immunity. Comparison of immunogenicity of trivalent vs. monovalent NE-based formulations indicated that combining PA, HCR/A1 and LcrV (or V10) did not affect the immunogenicity of individual antigen. Nasal NE-adjuvanted vaccines elicited serum IgG responses comparable with intramuscular injection of alum-based vaccines. Challenge studies with spores of Ames strain of anthrax spores and with a virulent *Y.pestis* CO92 indicate that nasal immunizations with NE-based vaccines produced protective immunity against anthrax and plague in rodent models (See Examples 8-9). Thus, in some embodiments, the present invention provides a multivalent nanoemulsion vaccine (e.g., against anthrax, plague and botulinum toxin). Thus, the present invention provides that vaccination (e.g., intranasal administration) with a multivalent vaccine obviates the need for injection or an inflammatory adjuvant. The present invention also provides a straight forward approach for formulation of an immunogenic composition (e.g., for use as a multivalent vaccine) that makes it suitable to be produced without special equipment. Thus, in some embodiments, compositions described herein are utilized in developing regions of the world. The present invention significantly decreases costs associated with conventional multivalent vaccines (e.g., the need to maintain conventional vaccines at a refrigerated temperature is overcome by the present invention). Since the multivalent vaccine retained immunogenicity after storage, in some embodiments, the vaccine does not require refrigeration during distribution.

Adjuvants have been traditionally developed from pro-inflammatory substances, such as a toxin or microbiological component, found to trigger signaling pathways and cytokine production (See, e.g., Graham, B.S., Plos Medicine, 2006. 3(1): p. e57). Also, enterotoxin-based adjuvants, such as cholera toxin, have been associated with inducing inflammation in the nasal mucosa and with production of the inflammatory cytokines and transport of the vaccine along olfactory neurons into the olfactory bulbs (See, e.g., van Ginkel, F.W., et al., Infect Immun., 2005. 73(10): p. 6892–6902). Some patients treated with a flu vaccine based

on one of these toxins (NASALFLU, BERNA Biotech), developed Bell's palsy (See, e.g., Mutsch, M., et al., New England Journal of Medicine, 2004. 350(9): p. 896-903) presumably due to the vaccine in the olfactory bulb. This finding led to NASALFLU being withdrawn. The present invention provides a composition with no significant inflammation in HBsAg-NE treated animals and no evidence of the vaccine composition in the olfactory bulb. Thus the present invention provides, in some embodiments, compositions and methods for inducing immune responses (e.g., immunity to) to a plurality of pathogens utilizing needle-free mucosal administration, induction of systemic immunity comparable with conventional vaccines, as well as mucosal and cellular immune responses that are not elicited by injected, aluminum-based vaccines.

Pathogens

The present invention is not limited to the use of any one specific type of pathogen (e.g., for providing one or more immunogens (e.g., antigens) for use in a composition comprising a nanoemulsion and a plurality of immunogens. Indeed, compositions (e.g., comprising a NE and a plurality of immunogens) useful for generating an immune response (e.g., for use as a vaccine) to a variety of pathogens are within the scope of the present invention. Accordingly, in some embodiments, the present invention provides compositions for generating an immune response to bacterial pathogens (e.g., in vegetative or spore forms) including, but not limited to, *Bacillus cereus*, *Bacillus circulans* and *Bacillus megaterium*, *Bacillus anthracis*, bacteria of the genus *Brucella*, *Vibrio cholera*, *Coxiella burnetii*, *Francisella tularensis*, *Chlamydia psittaci*, *Ricinus communis*, *Rickettsia prowazekii*, bacterial of the genus *Salmonella* (e.g., *S. typhi*), bacteria of the genus *Shigella*, *Cryptosporidium parvum*, *Burkholderia pseudomallei*, *Clostridium perfringens*, *Clostridium botulinum*, *Vibrio cholerae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus pneumonia*, *Staphylococcus aureus*, *Neisseria gonorrhea*, *Haemophilus influenzae*, *Escherichia coli*, *Salmonella typhimurium*, *Shigella dysenteriae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Yersinia pestis*, *Yersinia enterocolitica*, and *Yersinia pseudotuberculosis*). In other embodiments, the present invention provides compositions for generating an immune response to viral pathogens including, but not limited to, influenza A virus, avian influenza virus, H5N1 influenza virus, West Nile virus, SARS virus, Marburg virus, Arenaviruses, Nipah virus, alphaviruses, filoviruses, herpes simplex virus I, herpes simplex virus II, sendai, sindbis, vaccinia, parvovirus, human immunodeficiency virus,

hepatitis B virus, hepatitis C virus, hepatitis A virus, cytomegalovirus, human papilloma virus, picornavirus, hantavirus, junin virus, and ebola virus). In still further embodiments, the present invention provides compositions for generating an immune response to fungal pathogens, including, but not limited to, *Candida albicans* and *parapsilosis*, *Aspergillus*
5 *fumigatus* and *niger*, *Fusarium spp*, *Trychophyton spp*.

Bacteria for use in formulating a composition for generating an immune response of the present invention can be obtained from commercial sources, including, but not limited to, American Type Culture Collection (ATCC). In some embodiments, bacteria are passed in animals prior to being mixed with nanoemulsions in order to enhance their pathogenicity for
10 each specific animal host for 5-10 passages (Sinai *et al.*, J. Infect. Dis., 141:193 (1980)). In some embodiments, the bacteria then are then isolated from the host animals, expanded in culture and stored at -80°C. Just before use, the bacteria are thawed and grown on an appropriate solid bacterial culture medium overnight. The next day, the bacteria are collected from the agar plate and suspended in a suitable liquid solution (*e.g.*, Brain Heart Infusion
15 (BHI) broth). The concentration of bacteria is adjusted so that the bacteria count is approximately 1.5×10^8 colony forming units per ml (CFU/ml), based on the McFarland standard for bactericidal testing (Hendrichson and Krenz, 1991).

Viruses for use in formulating a composition for generating an immune response of the present invention can be obtained from commercial sources, including, but not limited to,
20 ATCC. In some embodiments, viruses are passed in the prospective animal model for 5-10 times to enhance pathogenicity for each specific animal (Ginsberg and Johnson, Infect. Immun., 13:1221 (1976)). In some embodiments, the virus is collected and propagated in tissue culture and then purified using density gradient concentration and ultracentrifugation (Garlinghouse *et al.*, Lab Anim Sci., 37:437 (1987); and Mahy, Br. Med. Bull., 41:50
25 (1985)). The Plaque Forming Units (PFU) are calculated in the appropriate tissue culture cells.

Lethal dose and/or infectious dose for each pathogen can be calculated using any suitable method, including, but not limited to, by administering different doses of the pathogens to the animals by the infective route and identifying the doses which result in the
30 expected result of either animal sickness or death based on previous publications (Fortier *et al.*, Infect Immun., 59:2922 (1991); Jacoby, Exp Gerontol., 29:89 (1994); and Salit *et al.*, Can J Microbiol., 30:1022 (1984)).

An immunogenic composition comprising a nanoemulsion and plurality of immunogens (e.g., protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce antibodies (e.g., polyclonal antibodies). If desired, one or more of a plurality of immunogens (e.g., protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, keyhole limpet hemocyanin or other carrier described herein, mixed with a nanoemulsion and administered to a subject. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, nanoemulsions described herein, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies that specifically bind one or more of a plurality of immunogens (e.g., protective antigen (PA) of *B. anthracis*, LcrV or LcrV10 of *Y. pestis*, and/or receptor binding domain HCR/A1 of *C. botulinum* neurotoxin) can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B cell hybridoma technique, and the EBV hybridoma technique (See, e.g., Kohler et al., Nature 256, 495 497, 1985; Kozbor et al., J. Immunol. Methods 81, 3142, 1985; Cote et al., Proc. Natl. Acad. Sci. 80, 2026 2030, 1983; Cole et al., Mol. Cell. Biol. 62, 109 120, 1984). Other techniques for antibody production are described herein.

Nanoemulsions

The nanoemulsion vaccine compositions of the present invention are not limited to any particular nanoemulsion. Any number of suitable nanoemulsion compositions may be utilized in the vaccine compositions of the present invention, including, but not limited to, those disclosed in Hamouda *et al.*, J. Infect Dis., 180:1939 (1999); Hamouda and Baker, J. Appl. Microbiol., 89:397 (2000); and Donovan *et al.*, Antivir. Chem. Chemother., 11:41 (2000), as well as those shown in Tables 1 and 2. Preferred nanoemulsions of the present

invention are those that are effective in killing or inactivating pathogens and that are non-toxic to animals. Accordingly, preferred emulsion formulations utilize non-toxic solvents, such as ethanol, and achieve more effective killing at lower concentrations of emulsion. In preferred embodiments, nanoemulsions utilized in the methods of the present invention are

5 stable, and do not decompose even after long storage periods (*e.g.*, one or more years).

Additionally, preferred emulsions maintain stability even after exposure to high temperature and freezing. This is especially useful if they are to be applied in extreme conditions (*e.g.*, on a battlefield). In some embodiments, one of the nanoemulsions described in Table 1 is utilized.

10 In some preferred embodiments, the emulsions comprise (i) an aqueous phase; (ii) an oil phase; and at least one additional compound. In some embodiments of the present invention, these additional compounds are admixed into either the aqueous or oil phases of the composition. In other embodiments, these additional compounds are admixed into a composition of previously emulsified oil and aqueous phases. In certain of these

15 embodiments, one or more additional compounds are admixed into an existing emulsion composition immediately prior to its use. In other embodiments, one or more additional compounds are admixed into an existing emulsion composition prior to the compositions immediate use.

Additional compounds suitable for use in the compositions of the present invention

20 include but are not limited to one or more, organic, and more particularly, organic phosphate based solvents, surfactants and detergents, quaternary ammonium containing compounds, cationic halogen containing compounds, germination enhancers, interaction enhancers, and pharmaceutically acceptable compounds. Certain exemplary embodiments of the various compounds contemplated for use in the compositions of the present invention are presented

25 below.

Table 1 Nanoemulsion Formulations		
Name	Oil Phase Formula	Water to Oil Phase Ratio (Vol/Vol)
X8P	1 vol. Tri(N-butyl)phosphate	4:1
	1 vol. TRITON X-100	

	8 vol. Soybean oil	
NN	86.5 g Glycerol monooleate	3:1
	60.1 ml Nonoxynol-9	
	24.2 g GENEROL 122	
	3.27 g Cetylpyridinium chloride	
	554 g Soybean oil	
W ₈₀ 8P	86.5 g Glycerol monooleate	3.2:1
	21.2 g Polysorbate 60	
	24.2 g GENEROL 122	
	3.27 g Cetylpyddinium chloride	
	4 ml Peppermint oil	
	554 g Soybean oil	
SS	86.5 g Glycerol monooleate	3.2:1
	21.2 g Polysorbate 60	(1% bismuth in water)
	24.2 g GENEROL 122	
	3.27 g Cetylpyridinium chloride	
	554 g Soybean oil	
Table 2 Nanoemulsion Formulations		
Nanoemulsion	Composition	
X8P	8% TRITON X-100; 8% Tributyl phosphate; 64% Soybean oil; 20% Water	
W ₂₀ 5EC	5% TWEEN 20; 8% Ethanol; 1% Cetylpyridinium Chloride; 64% Soybean oil; 22% Water	
EC	1% Cetylpyridinium Chloride; 8% Ethanol; 64% Soybean oil; 27% Water	
Y3EC	3% TYLOXAPOL; 1% Cetylpyridinium Chloride; 8% Ethanol; 64% Soybean oil; 24% Water	
X4E	4% TRITON X-100; 8% Ethanol; 64% Soybean oil; 24% Water	

Some embodiments of the present invention employ an oil phase containing ethanol.

For example, in some embodiments, the emulsions of the present invention contain (i) an aqueous phase and (ii) an oil phase containing ethanol as the organic solvent and optionally a germination enhancer, and (iii) TYLOXAPOL as the surfactant (preferably 2-5%, more preferably 3%). This formulation is highly efficacious against microbes and is also
 5 non-irritating and non-toxic to mammalian users (and can thus be contacted with mucosal membranes).

In some other embodiments, the emulsions of the present invention comprise a first emulsion emulsified within a second emulsion, wherein (a) the first emulsion comprises (i) an aqueous phase; and (ii) an oil phase comprising an oil and an organic solvent; and (iii) a
 10 surfactant; and (b) the second emulsion comprises (i) an aqueous phase; and (ii) an oil phase comprising an oil and a cationic containing compound; and (iii) a surfactant.

The following description provides a number of exemplary emulsions including formulations for compositions X8P and X₈W₆₀PC. X8P comprises a water-in oil nanoemulsion, in which the oil phase was made from soybean oil, tri-n-butyl phosphate, and
 15 TRITON X-100 in 80% water. X₈W₆₀PC comprises a mixture of equal volumes of X8P with W₈₀8P. W₈₀8P is a liposome-like compound made of glycerol monostearate, refined soya sterols (*e.g.*, GENEROL sterols), TWEEN 60, soybean oil, a cationic ion halogen-containing CPC and peppermint oil. The GENEROL family are a group of a polyethoxylated soya sterols (Henkel Corporation, Ambler, Pennsylvania). Emulsion
 20 formulations are given in Table 1 for certain embodiments of the present invention. These particular formulations may be found in U.S. Pat. Nos. 5,700,679 (NN); 5,618,840; 5,549,901 (W₈₀8P); and 5,547,677.

The X₈W₆₀PC emulsion is manufactured by first making the W₈₀8P emulsion and X8P emulsions separately. A mixture of these two emulsions is then re-emulsified to
 25 produce a fresh emulsion composition termed X₈W₆₀PC. Methods of producing such emulsions are described in U.S. Pat. Nos. 5,103,497 and 4,895,452.

These compounds have broad-spectrum antimicrobial activity, and are able to inactivate vegetative bacteria through membrane disruption.

The compositions listed above are only exemplary and those of skill in the art will be
 30 able to alter the amounts of the components to arrive at a nanoemulsion composition suitable for the purposes of the present invention. Those skilled in the art will understand that the

ratio of oil phase to water as well as the individual oil carrier, surfactant CPC and organic phosphate buffer, components of each composition may vary.

Although certain compositions comprising X8P have a water to oil ratio of 4:1, it is understood that the X8P may be formulated to have more or less of a water phase. For example, in some embodiments, there is 3, 4, 5, 6, 7, 8, 9, 10, or more parts of the water phase to each part of the oil phase. The same holds true for the W₈₀8P formulation.

Similarly, the ratio of Tri(N-butyl)phosphate:TRITON X-100:soybean oil also may be varied.

Although Table 1 lists specific amounts of glycerol monooleate, polysorbate 60, GENEROL 122, cetylpyridinium chloride, and carrier oil for W₈₀8P, these are merely exemplary. An emulsion that has the properties of W₈₀8P may be formulated that has different concentrations of each of these components or indeed different components that will fulfill the same function. For example, the emulsion may have between about 80 to about 100g of glycerol monooleate in the initial oil phase. In other embodiments, the emulsion may have between about 15 to about 30 g polysorbate 60 in the initial oil phase. In yet another embodiment the composition may comprise between about 20 to about 30 g of a GENEROL sterol, in the initial oil phase.

The nanoemulsions structure of the certain embodiments of the emulsions of the present invention may play a role in their biocidal activity as well as contributing to the non-toxicity of these emulsions. For example, the active component in X8P, TRITON-X100 shows less biocidal activity against virus at concentrations equivalent to 11% X8P. Adding the oil phase to the detergent and solvent markedly reduces the toxicity of these agents in tissue culture at the same concentrations. While not being bound to any theory (an understanding of the mechanism is not necessary to practice the present invention, and the present invention is not limited to any particular mechanism), it is suggested that the nanoemulsion enhances the interaction of its components with the pathogens thereby facilitating the inactivation of the pathogen and reducing the toxicity of the individual components. It should be noted that when all the components of X8P are combined in one composition but are not in a nanoemulsion structure, the mixture is not as effective as an antimicrobial as when the components are in a nanoemulsion structure.

Numerous additional embodiments presented in classes of formulations with like compositions are presented below. The following compositions recite various ratios and mixtures of active components. One skilled in the art will appreciate that the below recited

formulation are exemplary and that additional formulations comprising similar percent ranges of the recited components are within the scope of the present invention.

In certain embodiments of the present invention, the inventive formulation comprise from about 3 to 8 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of cetylpyridinium chloride (CPC), about 60 to 70 vol. % oil (*e.g.*, soybean oil), about 15 to 25 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS), and in some formulations less than about 1 vol. % of 1N NaOH. Some of these embodiments comprise PBS. It is contemplated that the addition of 1N NaOH and/or PBS in some of these embodiments, allows the user to advantageously control the pH of the formulations, such that pH ranges from about 4.0 to about 10.0, and more preferably from about 7.1 to 8.5 are achieved. For example, one embodiment of the present invention comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 24 vol. % of DiH₂O (designated herein as Y3EC). Another similar embodiment comprises about 3.5 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, and about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 23.5 vol. % of DiH₂O (designated herein as Y3.5EC). Yet another embodiment comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 0.067 vol. % of 1N NaOH, such that the pH of the formulation is about 7.1, about 64 vol. % of soybean oil, and about 23.93 vol. % of DiH₂O (designated herein as Y3EC pH 7.1). Still another embodiment comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 0.67 vol. % of 1N NaOH, such that the pH of the formulation is about 8.5, and about 64 vol. % of soybean oil, and about 23.33 vol. % of DiH₂O (designated herein as Y3EC pH 8.5). Another similar embodiment comprises about 4% TYLOXAPOL, about 8 vol. % ethanol, about 1% CPC, and about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as Y4EC). In still another embodiment the formulation comprises about 8% TYLOXAPOL, about 8% ethanol, about 1 vol. % of CPC, and about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as Y8EC). A further embodiment comprises about 8 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 19 vol. % of 1x PBS (designated herein as Y8EC PBS).

In some embodiments of the present invention, the inventive formulations comprise about 8 vol. % of ethanol, and about 1 vol. % of CPC, and about 64 vol. % of oil (*e.g.*,

soybean oil), and about 27 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS) (designated herein as EC).

In the present invention, some embodiments comprise from about 8 vol. % of sodium dodecyl sulfate (SDS), about 8 vol. % of tributyl phosphate (TBP), and about 64 vol. % of oil
5 (*e.g.*, soybean oil), and about 20 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS) (designated herein as S8P).

In certain embodiments of the present invention, the inventive formulation comprise from about 1 to 2 vol. % of TRITON X-100, from about 1 to 2 vol. % of TYLOXAPOL, from about 7 to 8 vol. % of ethanol, about 1 vol. % of cetylpyridinium chloride (CPC), about
10 64 to 57.6 vol. % of oil (*e.g.*, soybean oil), and about 23 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). Additionally, some of these formulations further comprise about 5 mM of L-alanine/Inosine, and about 10 mM ammonium chloride. Some of these formulations comprise PBS. It is contemplated that the addition of PBS in some of these embodiments, allows the user to advantageously control the pH of the formulations. For example, one
15 embodiment of the present invention comprises about 2 vol. % of TRITON X-100, about 2 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % CPC, about 64 vol. % of soybean oil, and about 23 vol. % of aqueous phase DiH₂O. In another embodiment the formulation comprises about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2 vol. % of ethanol, about 0.9 vol. % of CPC, about 5 mM L-
20 alanine/Inosine, and about 10 mM ammonium chloride, about 57.6 vol. % of soybean oil, and the remainder of 1x PBS (designated herein as 90% X2Y2EC/GE).

In a preferred embodiment of the present invention, the formulations comprise from about 5 vol. % of TWEEN 80, from about 8 vol. % of ethanol, from about 1 vol. % of CPC, about 64 vol. % of oil (*e.g.*, soybean oil), and about 22 vol. % of DiH₂O (designated herein
25 as W₈₀5EC).

In still other embodiments of the present invention, the formulations comprise from about 5 vol. % of TWEEN 20, from about 8 vol. % of ethanol, from about 1 vol. % of CPC, about 64 vol. % of oil (*e.g.*, soybean oil), and about 22 vol. % of DiH₂O (designated herein as W₂₀5EC).

30 In still other embodiments of the present invention, the formulations comprise from about 2 to 8 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 1 vol. % of CPC,

about 60 to 70 vol. % of oil (*e.g.*, soybean, or olive oil), and about 15 to 25 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). For example, the present invention contemplates formulations comprising about 2 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 26 vol. % of DiH₂O (designated herein as X2E). In other similar

5 embodiments, the formulations comprise about 3 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 25 vol. % of DiH₂O (designated herein as X3E). In still further embodiments, the formulations comprise about 4 vol. % TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 24 vol. % of DiH₂O (designated herein as X4E). In yet other embodiments, the formulations comprise

10 about 5 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as X5E). Another embodiment of the present invention comprises about 6 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 22 vol. % of DiH₂O (designated herein as X6E). In still further embodiments of the present invention, the formulations comprise about 8 vol.

15 % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8E). In still further embodiments of the present invention, the formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of olive oil, and about 20 vol. % of DiH₂O (designated herein as X8E O). In yet another embodiment comprises 8 vol. % of TRITON X-100, about 8 vol. %

20 ethanol, about 1 vol. % CPC, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as X8EC).

In alternative embodiments of the present invention, the formulations comprise from about 1 to 2 vol. % of TRITON X-100, from about 1 to 2 vol. % of TYLOXAPOL, from about 6 to 8 vol. % TBP, from about 0.5 to 1.0 vol. % of CPC, from about 60 to 70 vol. % of

25 oil (*e.g.*, soybean), and about 1 to 35 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS).

Additionally, certain of these formulations may comprise from about 1 to 5 vol. % of trypticase soy broth, from about 0.5 to 1.5 vol. % of yeast extract, about 5 mM L-alanine/Inosine, about 10 mM ammonium chloride, and from about 20-40 vol. % of liquid baby formula. In some of the embodiments comprising liquid baby formula, the formula

30 comprises a casein hydrolysate (*e.g.*, Neutramigen, or Progestimil, and the like). In some of these embodiments, the inventive formulations further comprise from about 0.1 to 1.0 vol. %

of sodium thiosulfate, and from about 0.1 to 1.0 vol. % of sodium citrate. Other similar embodiments comprising these basic components employ phosphate buffered saline (PBS) as the aqueous phase. For example, one embodiment comprises about 2 vol. % of TRITON X-100, about 2 vol. % TYLOXAPOL, about 8 vol. % TBP, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as X2Y2EC). In still other embodiments, the inventive formulation comprises about 2 vol. % of TRITON X-100, about 2 vol. % TYLOXAPOL, about 8 vol. % TBP, about 1 vol. % of CPC, about 0.9 vol. % of sodium thiosulfate, about 0.1 vol. % of sodium citrate, about 64 vol. % of soybean oil, and about 22 vol. % of DiH₂O (designated herein as X2Y2PC STS1). In another similar embodiment, the formulations comprise about 1.7 vol. % TRITON X-100, about 1.7 vol. % TYLOXAPOL, about 6.8 vol. % TBP, about 0.85% CPC, about 29.2% NEUTRAMIGEN, about 54.4 vol. % of soybean oil, and about 4.9 vol. % of DiH₂O (designated herein as 85% X2Y2PC/baby). In yet another embodiment of the present invention, the formulations comprise about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2 vol. % of TBP, about 0.9 vol. % of CPC, about 5mM L-alanine/Inosine, about 10mM ammonium chloride, about 57.6 vol. % of soybean oil, and the remainder vol. % of 0.1x PBS (designated herein as 90% X2Y2 PC/GE). In still another embodiment, the formulations comprise about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2 vol. % TBP, about 0.9 vol. % of CPC, and about 3 vol. % trypticase soy broth, about 57.6 vol. % of soybean oil, and about 27.7 vol. % of DiH₂O (designated herein as 90% X2Y2PC/TSB). In another embodiment of the present invention, the formulations comprise about 1.8 vol. % TRITON X-100, about 1.8 vol. % TYLOXAPOL, about 7.2 vol. % TBP, about 0.9 vol. % CPC, about 1 vol. % yeast extract, about 57.6 vol. % of soybean oil, and about 29.7 vol. % of DiH₂O (designated herein as 90% X2Y2PC/YE).

In some embodiments of the present invention, the inventive formulations comprise about 3 vol. % of TYLOXAPOL, about 8 vol. % of TBP, and about 1 vol. % of CPC, about 60 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). In a particular embodiment of the present invention, the inventive formulations comprise about 3 vol. % of TYLOXAPOL, about 8 vol. % of TBP, and about 1 vol. % of CPC, about 64 vol. % of soybean, and about 24 vol. % of DiH₂O (designated herein as Y3PC).

In some embodiments of the present invention, the inventive formulations comprise from about 4 to 8 vol. % of TRITON X-100, from about 5 to 8 vol. % of TBP, about 30 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and about 0 to 30 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). Additionally, certain of these embodiments further comprise about 1 vol. % of CPC, about 1 vol. % of benzalkonium chloride, about 1 vol. % cetylridinium bromide, about 1 vol. % cetyldimethyletylammonium bromide, 500 μ M EDTA, about 10 mM ammonium chloride, about 5 mM Inosine, and about 5 mM L-alanine. For example, in certain of these embodiments, the inventive formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8P). In another embodiment of the present invention, the inventive formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1% of CPC, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as X8PC). In still another embodiment, the formulations comprise about 8 vol. % TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as ATB-X1001). In yet another embodiment, the formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 2 vol. % of CPC, about 50 vol. % of soybean oil, and about 32 vol. % of DiH₂O (designated herein as ATB-X002). Another embodiment of the present invention comprises about 4 vol. % TRITON X-100, about 4 vol. % of TBP, about 0.5 vol. % of CPC, about 32 vol. % of soybean oil, and about 59.5 vol. % of DiH₂O (designated herein as 50% X8PC). Still another related embodiment comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 0.5 vol. % CPC, about 64 vol. % of soybean oil, and about 19.5 vol. % of DiH₂O (designated herein as X8PC_{1/2}). In some embodiments of the present invention, the inventive formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 2 vol. % of CPC, about 64 vol. % of soybean oil, and about 18 vol. % of DiH₂O (designated herein as X8PC2). In other embodiments, the inventive formulations comprise about 8 vol. % of TRITON X-100, about 8% of TBP, about 1% of benzalkonium chloride, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as X8P BC). In an alternative embodiment of the present invention, the formulation comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of cetylridinium bromide, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as

X8P CPB). In another exemplary embodiment of the present invention, the formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of cetyldimethyletylammonium bromide, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as X8P CTAB). In still further embodiments, the present
5 invention comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 500 μ M EDTA, about 64 vol. % of soybean oil, and about 15.8 vol. % DiH₂O (designated herein as X8PC EDTA). Additional similar embodiments comprise 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 10 mM ammonium chloride, about 5mM Inosine, about 5mM L-alanine, about 64 vol. % of soybean
10 oil, and about 19 vol. % of DiH₂O or PBS (designated herein as X8PC GE_{1x}). In another embodiment of the present invention, the inventive formulations further comprise about 5 vol. % of TRITON X-100, about 5% of TBP, about 1 vol. % of CPC, about 40 vol. % of soybean oil, and about 49 vol. % of DiH₂O (designated herein as X5P₅C).

In some embodiments of the present invention, the inventive formulations comprise
15 about 2 vol. % TRITON X-100, about 6 vol. % TYLOXAPOL, about 8 vol. % ethanol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X2Y6E).

In an additional embodiment of the present invention, the formulations comprise about 8 vol. % of TRITON X-100, and about 8 vol. % of glycerol, about 60 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and about 15 to 25 vol. % of aqueous phase (*e.g.*, DiH₂O or
20 PBS). Certain related embodiments further comprise about 1 vol. % L-ascorbic acid. For example, one particular embodiment comprises about 8 vol. % of TRITON X-100, about 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8G). In still another embodiment, the inventive formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of glycerol, about 1 vol. % of L-
25 ascorbic acid, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as X8GV_c).

In still further embodiments, the inventive formulations comprise about 8 vol. % of TRITON X-100, from about 0.5 to 0.8 vol. % of TWEEN 60, from about 0.5 to 2.0 vol. % of CPC, about 8 vol. % of TBP, about 60 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and
30 about 15 to 25 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). For example, in one particular embodiment the formulations comprise about 8 vol. % of TRITON X-100, about 0.70 vol. %

of TWEEN 60, about 1 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 18.3 vol. % of DiH₂O (designated herein as X8W60PC₁). Another related embodiment comprises about 8 vol. % of TRITON X-100, about 0.71 vol. % of TWEEN 60, about 1 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 18.29 vol. % of DiH₂O (designated herein as W60_{0.7}X8PC). In yet other embodiments, the inventive formulations comprise from about 8 vol. % of TRITON X-100, about 0.7 vol. % of TWEEN 60, about 0.5 vol. % of CPC, about 8 vol. % of TBP, about 64 to 70 vol. % of soybean oil, and about 18.8 vol. % of DiH₂O (designated herein as X8W60PC₂). In still other embodiments, the present invention comprises about 8 vol. % of TRITON X-100, about 0.71 vol. % of TWEEN 60, about 2 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 17.3 vol. % of DiH₂O. In another embodiment of the present invention, the formulations comprise about 0.71 vol. % of TWEEN 60, about 1 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 25.29 vol. % of DiH₂O (designated herein as W60_{0.7}PC).

In another embodiment of the present invention, the inventive formulations comprise about 2 vol. % of dioctyl sulfosuccinate, either about 8 vol. % of glycerol, or about 8 vol. % TBP, in addition to, about 60 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and about 20 to 30 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). For example, one embodiment of the present invention comprises about 2 vol. % of dioctyl sulfosuccinate, about 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 26 vol. % of DiH₂O (designated herein as D2G). In another related embodiment, the inventive formulations comprise about 2 vol. % of dioctyl sulfosuccinate, and about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 26 vol. % of DiH₂O (designated herein as D2P).

In still other embodiments of the present invention, the inventive formulations comprise about 8 to 10 vol. % of glycerol, and about 1 to 10 vol. % of CPC, about 50 to 70 vol. % of oil (*e.g.*, soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (*e.g.*, DiH₂O or PBS). Additionally, in certain of these embodiments, the compositions further comprise about 1 vol. % of L-ascorbic acid. For example, one particular embodiment comprises about 8 vol. % of glycerol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 27 vol. % of DiH₂O (designated herein as GC). An additional related embodiment comprises about 10 vol. % of glycerol, about 10 vol. % of CPC, about 60 vol. % of soybean

oil, and about 20 vol. % of DiH_2O (designated herein as GC10). In still another embodiment of the present invention, the inventive formulations comprise about 10 vol. % of glycerol, about 1 vol. % of CPC, about 1 vol. % of L-ascorbic acid, about 64 vol. % of soybean oil, and about 24 vol. % of DiH_2O (designated herein as GCV_c).

5 In some embodiments of the present invention, the inventive formulations comprise about 8 to 10 vol. % of glycerol, about 8 to 10 vol. % of SDS, about 50 to 70 vol. % of oil (e.g., soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (e.g., DiH_2O or PBS). Additionally, in certain of these embodiments, the compositions further comprise about 1 vol. % of lecithin, and about 1 vol. % of p-Hydroxybenzoic acid methyl ester.

10 Exemplary embodiments of such formulations comprise about 8 vol. % SDS, 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH_2O (designated herein as S8G). A related formulation comprises about 8 vol. % of glycerol, about 8 vol. % of SDS, about 1 vol. % of lecithin, about 1 vol. % of p-Hydroxybenzoic acid methyl ester, about 64 vol. % of soybean oil, and about 18 vol. % of DiH_2O (designated herein as S8GL1B1).

15 In yet another embodiment of the present invention, the inventive formulations comprise about 4 vol. % of TWEEN 80, about 4 vol. % of TYLOXAPOL, about 1 vol. % of CPC, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 19 vol. % of DiH_2O (designated herein as W804Y4EC).

20 In some embodiments of the present invention, the inventive formulations comprise about 0.01 vol. % of CPC, about 0.08 vol. % of TYLOXAPOL, about 10 vol. % of ethanol, about 70 vol. % of soybean oil, and about 19.91 vol. % of DiH_2O (designated herein as Y.08EC.01).

25 In yet another embodiment of the present invention, the inventive formulations comprise about 8 vol. % of sodium lauryl sulfate, and about 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH_2O (designated herein as SLS8G).

30 The specific formulations described above are simply examples to illustrate the variety of compositions that find use in the present invention. The present invention contemplates that many variations of the above formulation, as well as additional nanoemulsions, find use in the methods of the present invention. To determine if a candidate emulsion is suitable for use with the present invention, three criteria may be analyzed. Using the methods and standards described herein, candidate emulsions can be easily tested to

determine if they are suitable. First, the desired ingredients are prepared using the methods described herein, to determine if an emulsion can be formed. If an emulsion cannot be formed, the candidate is rejected. For example, a candidate composition made of 4.5% sodium thiosulfate, 0.5% sodium citrate, 10% n-butanol, 64% soybean oil, and 21% DiH₂O
5 did not form an emulsion.

Second, in preferred embodiments, the candidate emulsion should form a stable emulsion. An emulsion is stable if it remains in emulsion form for a sufficient period to allow its intended use. For example, for emulsions that are to be stored, shipped, etc., it may be desired that the composition remain in emulsion form for months to years. Typical
10 emulsions that are relatively unstable, will lose their form within a day. For example, a candidate composition made of 8% 1-butanol, 5% TWEEN 10, 1% CPC, 64% soybean oil, and 22% DiH₂O did not form a stable emulsion. The following candidate emulsions were shown to be stable using the methods described herein: 0.08% TRITON X-100, 0.08% Glycerol, 0.01% Cetylpyridinium Chloride, 99% Butter, and 0.83% diH₂O (designated
15 herein as 1% X8GC Butter); 0.8% TRITON X-100, 0.8% Glycerol, 0.1% Cetylpyridinium Chloride, 6.4% Soybean Oil, 1.9% diH₂O, and 90% Butter (designated herein as 10% X8GC Butter); 2% W₂₀5EC, 1% Natrosol 250L NF, and 97% diH₂O (designated herein as 2% W₂₀5EC L GEL); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% 70 Viscosity Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC 70 Mineral Oil); 1%
20 Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% 350 Viscosity Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC 350 Mineral Oil).

Third, the candidate emulsion should have efficacy for its intended use. For example, an anti-bacterial emulsion should kill or disable pathogens to a detectable level. As shown herein, certain emulsions of the present invention have efficacy against specific
25 microorganisms, but not against others. Using the methods described herein, one is capable of determining the suitability of a particular candidate emulsion against the desired microorganism. Generally, this involves exposing the microorganism to the emulsion for one or more time periods in a side-by-side experiment with the appropriate control samples (*e.g.*, a negative control such as water) and determining if, and to what degree, the emulsion kills or
30 disables the microorganism. For example, a candidate composition made of 1% ammonium chloride, 5% TWEEN 20, 8% ethanol, 64% soybean oil, and 22% DiH₂O was shown not to

be an effective emulsion. The following candidate emulsions were shown to be effective using the methods described herein: 5% TWEEN 20, 5% Cetylpyridinium Chloride, 10% Glycerol, 60% Soybean Oil, and 20% diH₂O (designated herein as W₂₀5GC5); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 10% Glycerol, 64% Soybean Oil, and 20% diH₂O (designated herein as W₂₀5GC); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Olive Oil, and 22% diH₂O (designated herein as W₂₀5EC Olive Oil); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Flaxseed Oil, and 22% diH₂O (designated herein as W₂₀5EC Flaxseed Oil); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Corn Oil, and 22% diH₂O (designated herein as W₂₀5EC Corn Oil); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Coconut Oil, and 22% diH₂O (designated herein as W₂₀5EC Coconut Oil); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Cottonseed Oil, and 22% diH₂O (designated herein as W₂₀5EC Cottonseed Oil); 8% Dextrose, 5% TWEEN 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C Dextrose); 8% PEG 200, 5% TWEEN 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C PEG 200); 8% Methanol, 5% TWEEN 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C Methanol); 8% PEG 1000, 5% TWEEN 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C PEG 1000); 2% W₂₀5EC, 2% Natrosol 250H NF, and 96% diH₂O (designated herein as 2% W₂₀5EC Natrosol 2, also called 2% W₂₀5EC GEL); 2% W₂₀5EC, 1% Natrosol 250H NF, and 97% diH₂O (designated herein as 2% W₂₀5EC Natrosol 1); 2% W₂₀5EC, 3% Natrosol 250H NF, and 95% diH₂O (designated herein as 2% W₂₀5EC Natrosol 3); 2% W₂₀5EC, 0.5% Natrosol 250H NF, and 97.5% diH₂O (designated herein as 2% W₂₀5EC Natrosol 0.5); 2% W₂₀5EC, 2% Methocel A, and 96% diH₂O (designated herein as 2% W₂₀5EC Methocel A); 2% W₂₀5EC, 2% Methocel K, and 96% diH₂O (designated herein as 2% W₂₀5EC Methocel K); 2% Natrosol, 0.1% X8PC, 0.1x PBS, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, and diH₂O (designated herein as 0.1% X8PC/GE+2% Natrosol); 2% Natrosol, 0.8% TRITON X-100, 0.8% Tributyl Phosphate, 6.4% Soybean Oil, 0.1% Cetylpyridinium Chloride, 0.1x PBS, 5 mM L-alanine, 5

mM Inosine, 10 mM Ammonium Chloride, and diH₂O (designated herein as 10% X8PC/GE+2% Natrosol); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Lard, and 22% diH₂O (designated herein as W₂₀5EC Lard); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Ethanol, 64% Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC Mineral Oil); 0.1% Cetylpyridinium Chloride, 2% Nerolidol, 5% TWEEN 20, 10% Ethanol, 64% Soybean Oil, and 18.9% diH₂O (designated herein as W₂₀5EC_{0.1}N); 0.1% Cetylpyridinium Chloride, 2% Farnesol, 5% TWEEN 20, 10% Ethanol, 64% Soybean Oil, and 18.9% diH₂O (designated herein as W₂₀5EC_{0.1}F); 0.1% Cetylpyridinium Chloride, 5% TWEEN 20, 10% Ethanol, 64% Soybean Oil, and 20.9% diH₂O (designated herein as W₂₀5EC_{0.1}); 10% Cetylpyridinium Chloride, 8% Tributyl Phosphate, 8% TRITON X-100, 54% Soybean Oil, and 20% diH₂O (designated herein as X8PC₁₀); 5% Cetylpyridinium Chloride, 8% TRITON X-100, 8% Tributyl Phosphate, 59% Soybean Oil, and 20% diH₂O (designated herein as X8PC₅); 0.02% Cetylpyridinium Chloride, 0.1% TWEEN 20, 10% Ethanol, 70% Soybean Oil, and 19.88% diH₂O (designated herein as W₂₀0.1EC_{0.02}); 1% Cetylpyridinium Chloride, 5% TWEEN 20, 8% Glycerol, 64% Mobil 1, and 22% diH₂O (designated herein as W₂₀5GC Mobil 1); 7.2% TRITON X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 0.1x PBS, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, and 25.87% diH₂O (designated herein as 90% X8PC/GE); 7.2% TRITON X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 1% EDTA, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, 0.1x PBS, and diH₂O (designated herein as 90% X8PC/GE EDTA); and 7.2% TRITON X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 1% Sodium Thiosulfate, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, 0.1x PBS, and diH₂O (designated herein as 90% X8PC/GE STS).

25

1. Aqueous Phase

In some embodiments, the emulsion comprises an aqueous phase. In certain preferred embodiments, the emulsion comprises about 5 to 50, preferably 10 to 40, more preferably 15 to 30, vol. % aqueous phase, based on the total volume of the emulsion (although other concentrations are also contemplated). In preferred embodiments, the aqueous phase

30

comprises water at a pH of about 4 to 10, preferably about 6 to 8. The water is preferably deionized (hereinafter "DiH₂O"). In some embodiments, the aqueous phase comprises phosphate buffered saline (PBS). In some preferred embodiments, the aqueous phase is sterile and pyrogen free.

5

2. Oil Phase

In some embodiments, the emulsion comprises an oil phase. In certain preferred embodiments, the oil phase (*e.g.*, carrier oil) of the emulsion of the present invention comprises 30-90, preferably 60-80, and more preferably 60-70, vol. % of oil, based on the total volume of the emulsion (although higher and lower concentrations also find use in emulsions described herein).

The oil in the nanoemulsion vaccine of the invention can be any cosmetically or pharmaceutically acceptable oil. The oil can be volatile or non-volatile, and may be chosen from animal oil, vegetable oil, natural oil, synthetic oil, hydrocarbon oils, silicone oils, semi-synthetic derivatives thereof, and combinations thereof.

Suitable oils include, but are not limited to, mineral oil, squalene oil, flavor oils, silicon oil, essential oils, water insoluble vitamins, Isopropyl stearate, Butyl stearate, Octyl palmitate, Cetyl palmitate, Tridecyl behenate, Diisopropyl adipate, Dioctyl sebacate, Menthyl anthranilate, Cetyl octanoate, Octyl salicylate, Isopropyl myristate, neopentyl glycol dicarpate cetols, Ceraphyls®, Decyl oleate, diisopropyl adipate, C₁₂₋₁₅ alkyl lactates, Cetyl lactate, Lauryl lactate, Isostearyl neopentanoate, Myristyl lactate, Isocetyl stearyl stearate, Octyldodecyl stearyl stearate, Hydrocarbon oils, Isoparaffin, Fluid paraffins, Isododecane, Petrolatum, Argan oil, Canola oil, Chile oil, Coconut oil, corn oil, Cottonseed oil, Flaxseed oil, Grape seed oil, Mustard oil, Olive oil, Palm oil, Palm kernel oil, Peanut oil, Pine seed oil, Poppy seed oil, Pumpkin seed oil, Rice bran oil, Safflower oil, Tea oil, Truffle oil, Vegetable oil, Apricot (kernel) oil, Jojoba oil (*simmondsia chinensis* seed oil), Grapeseed oil, Macadamia oil, Wheat germ oil, Almond oil, Rapeseed oil, Gourd oil, Soybean oil, Sesame oil, Hazelnut oil, Maize oil, Sunflower oil, Hemp oil, Bois oil, Kuki nut oil, Avocado oil, Walnut oil, Fish oil, berry oil, allspice oil, juniper oil, seed oil, almond seed oil, anise seed oil, celery seed oil, cumin seed oil, nutmeg seed oil, leaf oil, basil leaf oil, bay leaf oil, cinnamon leaf oil, common sage leaf oil, eucalyptus leaf oil, lemon grass leaf oil, melaleuca leaf oil, oregano leaf oil, patchouli leaf oil, peppermint leaf oil, pine needle oil, rosemary leaf

30

oil, spearmint leaf oil, tea tree leaf oil, thyme leaf oil, wintergreen leaf oil, flower oil, chamomile oil, clary sage oil, clove oil, geranium flower oil, hyssop flower oil, jasmine flower oil, lavender flower oil, manuka flower oil, Marhoram flower oil, orange flower oil, rose flower oil, ylang-ylang flower oil, Bark oil, cassia Bark oil, cinnamon bark oil, sassafras
5 Bark oil, Wood oil, camphor wood oil, cedar wood oil, rosewood oil, sandalwood oil), rhizome (ginger) wood oil, resin oil, frankincense oil, myrrh oil, peel oil, bergamot peel oil, grapefruit peel oil, lemon peel oil, lime peel oil, orange peel oil, tangerine peel oil, root oil, valerian oil, Oleic acid, Linoleic acid, Oleyl alcohol, Isostearyl alcohol, semi-synthetic derivatives thereof, and any combinations thereof.

10 The oil may further comprise a silicone component, such as a volatile silicone component, which can be the sole oil in the silicone component or can be combined with other silicone and non-silicone, volatile and non-volatile oils. Suitable silicone components include, but are not limited to, methylphenylpolysiloxane, simethicone, dimethicone, phenyltrimethicone (or an organomodified version thereof), alkylated derivatives of
15 polymeric silicones, cetyl dimethicone, lauryl trimethicone, hydroxylated derivatives of polymeric silicones, such as dimethiconol, volatile silicone oils, cyclic and linear silicones, cyclomethicone, derivatives of cyclomethicone, hexamethylcyclotrisiloxane, octamethylcyclotetrasiloxane, decamethylcyclopentasiloxane, volatile linear dimethylpolysiloxanes, isohexadecane, isoeicosane, isotetracosane, polyisobutene, isooctane,
20 isododecane, semi-synthetic derivatives thereof, and combinations thereof.

The volatile oil can be the organic solvent, or the volatile oil can be present in addition to an organic solvent. Suitable volatile oils include, but are not limited to, a terpene, monoterpene, sesquiterpene, carminative, azulene, menthol, camphor, thujone, thymol, nerol, linalool, limonene, geraniol, perillyl alcohol, nerolidol, farnesol, ylangene, bisabolol,
25 farnesene, ascaridole, chenopodium oil, citronellal, citral, citronellol, chamazulene, yarrow, guaiazulene, chamomile, semi-synthetic derivatives, or combinations thereof.

In one aspect of the invention, the volatile oil in the silicone component is different than the oil in the oil phase.

In some embodiments, the oil phase comprises 3-15, and preferably 5-10 vol. % of an
30 organic solvent, based on the total volume of the emulsion. While the present invention is not limited to any particular mechanism, it is contemplated that the organic phosphate-based solvents employed in the emulsions serve to remove or disrupt the lipids in the membranes of

the pathogens. Thus, any solvent that removes the sterols or phospholipids in the microbial membranes finds use in the methods of the present invention. Suitable organic solvents include, but are not limited to, organic phosphate based solvents or alcohols. In some preferred embodiments, non-toxic alcohols (*e.g.*, ethanol) are used as a solvent. The oil phase, and any additional compounds provided in the oil phase, are preferably sterile and pyrogen free.

3. Surfactants and Detergents

In some embodiments, the emulsions further comprises a surfactant or detergent. In some preferred embodiments, the emulsion comprises from about 3 to 15 %, and preferably about 10 % of one or more surfactants or detergents (although other concentrations are also contemplated). While the present invention is not limited to any particular mechanism, it is contemplated that surfactants, when present in the emulsions, help to stabilize the emulsions. Both non-ionic (non-anionic) and ionic surfactants are contemplated. Additionally, surfactants from the BRIJ family of surfactants find use in the compositions of the present invention. The surfactant can be provided in either the aqueous or the oil phase. Surfactants suitable for use with the emulsions include a variety of anionic and nonionic surfactants, as well as other emulsifying compounds that are capable of promoting the formation of oil-in-water emulsions. In general, emulsifying compounds are relatively hydrophilic, and blends of emulsifying compounds can be used to achieve the necessary qualities. In some formulations, nonionic surfactants have advantages over ionic emulsifiers in that they are substantially more compatible with a broad pH range and often form more stable emulsions than do ionic (*e.g.*, soap-type) emulsifiers.

The surfactant in the nanoemulsion vaccine of the invention can be a pharmaceutically acceptable ionic surfactant, a pharmaceutically acceptable nonionic surfactant, a pharmaceutically acceptable cationic surfactant, a pharmaceutically acceptable anionic surfactant, or a pharmaceutically acceptable zwitterionic surfactant.

Exemplary useful surfactants are described in *Applied Surfactants: Principles and Applications*. Tharwat F. Tadros, Copyright 8 2005 WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim ISBN: 3-527-30629-3).

Further, the surfactant can be a pharmaceutically acceptable ionic polymeric surfactant, a pharmaceutically acceptable nonionic polymeric surfactant, a pharmaceutically acceptable

cationic polymeric surfactant, a pharmaceutically acceptable anionic polymeric surfactant, or a pharmaceutically acceptable zwitterionic polymeric surfactant. Examples of polymeric surfactants include, but are not limited to, a graft copolymer of a poly(methyl methacrylate) backbone with multiple (at least one) polyethylene oxide (PEO) side chain,

5 polyhydroxystearic acid, an alkoxyated alkyl phenol formaldehyde condensate, a polyalkylene glycol modified polyester with fatty acid hydrophobes, a polyester, semi-synthetic derivatives thereof, or combinations thereof.

Surface active agents or surfactants, are amphipathic molecules that consist of a non-polar hydrophobic portion, usually a straight or branched hydrocarbon or fluorocarbon chain
10 containing 8-18 carbon atoms, attached to a polar or ionic hydrophilic portion. The hydrophilic portion can be nonionic, ionic or zwitterionic. The hydrocarbon chain interacts weakly with the water molecules in an aqueous environment, whereas the polar or ionic head group interacts strongly with water molecules via dipole or ion-dipole interactions. Based on the nature of the hydrophilic group, surfactants are classified into anionic, cationic,
15 zwitterionic, nonionic and polymeric surfactants.

Suitable surfactants include, but are not limited to, ethoxylated nonylphenol comprising 9 to 10 units of ethyleneglycol, ethoxylated undecanol comprising 8 units of ethyleneglycol, polyoxyethylene (20) sorbitan monolaurate, polyoxyethylene (20) sorbitan monopalmitate, polyoxyethylene (20) sorbitan monostearate, polyoxyethylene (20) sorbitan
20 monooleate, sorbitan monolaurate, sorbitan monopalmitate, sorbitan monostearate, sorbitan monooleate, ethoxylated hydrogenated ricin oils, sodium laurylsulfate, a diblock copolymer of ethyleneoxyde and propyleneoxyde, Ethylene Oxide-Propylene Oxide Block Copolymers, and tetra-functional block copolymers based on ethylene oxide and propylene oxide, Glyceryl monoesters, Glyceryl caprate, Glyceryl caprylate, Glyceryl cocate, Glyceryl erucate, Glyceryl
25 hydroxystearate, Glyceryl isostearate, Glyceryl lanolate, Glyceryl laurate, Glyceryl linolate, Glyceryl myristate, Glyceryl oleate, Glyceryl PABA, Glyceryl palmitate, Glyceryl ricinoleate, Glyceryl stearate, Glyceryl thiglycolate, Glyceryl dilaurate, Glyceryl dioleate, Glyceryl dimyristate, Glyceryl disterate, Glyceryl sesuiolate, Glyceryl stearate lactate, Polyoxyethylene cetyl/stearyl ether, Polyoxyethylene cholesterol ether, Polyoxyethylene
30 laurate or dilaurate, Polyoxyethylene stearate or distearate, polyoxyethylene fatty ethers, Polyoxyethylene lauryl ether, Polyoxyethylene stearyl ether, polyoxyethylene myristyl ether, a steroid, Cholesterol, Betasitosterol, Bisabolol, fatty acid esters of alcohols, isopropyl

myristate, Aliphatic-isopropyl n-butyrate, Isopropyl n-hexanoate, Isopropyl n-decanoate, Isopropyl palmitate, Octyldodecyl myristate, alkoxyated alcohols, alkoxyated acids, alkoxyated amides, alkoxyated sugar derivatives, alkoxyated derivatives of natural oils and waxes, polyoxyethylene polyoxypropylene block copolymers, nonoxynol-14, PEG-8 laurate, PEG-6 Cocoamide, PEG-20 methylglucose sesquistearate, PEG40 lanolin, PEG-40 castor oil, PEG-40 hydrogenated castor oil, polyoxyethylene fatty ethers, glyceryl diesters, polyoxyethylene stearyl ether, polyoxyethylene myristyl ether, and polyoxyethylene lauryl ether, glyceryl dilaurate, glyceryl dimystate, glyceryl distearate, semi-synthetic derivatives thereof, or mixtures thereof.

Additional suitable surfactants include, but are not limited to, non-ionic lipids, such as glyceryl laurate, glyceryl myristate, glyceryl dilaurate, glyceryl dimyristate, semi-synthetic derivatives thereof, and mixtures thereof.

In additional embodiments, the surfactant is a polyoxyethylene fatty ether having a polyoxyethylene head group ranging from about 2 to about 100 groups, or an alkoxyated alcohol having the structure $R_5-(OCH_2CH_2)_y-OH$, wherein R_5 is a branched or unbranched alkyl group having from about 6 to about 22 carbon atoms and y is between about 4 and about 100, and preferably, between about 10 and about 100. Preferably, the alkoxyated alcohol is the species wherein R_5 is a lauryl group and y has an average value of 23.

In a different embodiment, the surfactant is an alkoxyated alcohol which is an ethoxylated derivative of lanolin alcohol. Preferably, the ethoxylated derivative of lanolin alcohol is laneth-10, which is the polyethylene glycol ether of lanolin alcohol with an average ethoxylation value of 10.

Nonionic surfactants include, but are not limited to, an ethoxylated surfactant, an alcohol ethoxylated, an alkyl phenol ethoxylated, a fatty acid ethoxylated, a monoalkanolamide ethoxylated, a sorbitan ester ethoxylated, a fatty amino ethoxylated, an ethylene oxide-propylene oxide copolymer, Bis(polyethylene glycol bis[imidazolyl carbonyl]), nonoxynol-9, Bis(polyethylene glycol bis[imidazolyl carbonyl]), Brij[®] 35, Brij[®] 56, Brij[®] 72, Brij[®] 76, Brij[®] 92V, Brij[®] 97, Brij[®] 58P, Cremophor[®] EL, Decaethylene glycol monododecyl ether, N-Decanoyl-N-methylglucamine, n-Decyl alpha-D-glucopyranoside, Decyl beta-D-maltopyranoside, n-Dodecanoyl-N-methylglucamide, n-Dodecyl alpha-D-maltoside, n-Dodecyl beta-D-maltoside, n-Dodecyl beta-D-maltoside, Heptaethylene glycol monodecyl ether, Heptaethylene glycol monododecyl ether, Heptaethylene glycol

monotetradecyl ether, n-Hexadecyl beta-D-maltoside, Hexaethylene glycol monododecyl ether, Hexaethylene glycol monoheptadecyl ether, Hexaethylene glycol monooctadecyl ether, Hexaethylene glycol monotetradecyl ether, Igepal CA-630, Igepal CA-630, Methyl-6-O-(N-heptylcarbamoyl)-alpha-D-glucopyranoside, Nonaethylene glycol monododecyl ether, N-Nonanoyl-N-methylglucamine, N-Nonanoyl-N-methylglucamine, Octaethylene glycol monodecyl ether, Octaethylene glycol monododecyl ether, Octaethylene glycol monohexadecyl ether, Octaethylene glycol monooctadecyl ether, Octaethylene glycol monotetradecyl ether, Octyl-beta-D-glucopyranoside, Pentaethylene glycol monodecyl ether, Pentaethylene glycol monododecyl ether, Pentaethylene glycol monohexadecyl ether, Pentaethylene glycol monooctadecyl ether, Polyethylene glycol diglycidyl ether, Polyethylene glycol ether W-1, Polyoxyethylene 10 tridecyl ether, Polyoxyethylene 100 stearate, Polyoxyethylene 20 isohexadecyl ether, Polyoxyethylene 20 oleyl ether, Polyoxyethylene 40 stearate, Polyoxyethylene 50 stearate, Polyoxyethylene 8 stearate, Polyoxyethylene bis(imidazolyl carbonyl), Polyoxyethylene 25 propylene glycol stearate, Saponin from Quillaja bark, Span[®] 20, Span[®] 40, Span[®] 60, Span[®] 65, Span[®] 80, Span[®] 85, Tergitol, Type 15-S-12, Tergitol, Type 15-S-30, Tergitol, Type 15-S-5, Tergitol, Type 15-S-7, Tergitol, Type 15-S-9, Tergitol, Type NP-10, Tergitol, Type NP-4, Tergitol, Type NP-40, Tergitol, Type NP-7, Tergitol, Type NP-9, Tergitol, Tergitol, Type TMN-10, Tergitol, Type TMN-6, Tetradecyl-beta-D-maltoside, Tetraethylene glycol monodecyl ether, Tetraethylene glycol monododecyl ether, Tetraethylene glycol monotetradecyl ether, Triethylene glycol monodecyl ether, Triethylene glycol monododecyl ether, Triethylene glycol monohexadecyl ether, Triethylene glycol monooctyl ether, Triethylene glycol monotetradecyl ether, Triton CF-21, Triton CF-32, Triton DF-12, Triton DF-16, Triton GR-5M, Triton QS-15, Triton QS-44, Triton X-100, Triton X-102, Triton X-15, Triton X-151, Triton X-200, Triton X-207, Triton[®] X-100, Triton[®] X-114, Triton[®] X-165, Triton[®] X-305, Triton[®] X-405, Triton[®] X-45, Triton[®] X-705-70, TWEEN[®] 20, TWEEN[®] 21, TWEEN[®] 40, TWEEN[®] 60, TWEEN[®] 61, TWEEN[®] 65, TWEEN[®] 80, TWEEN[®] 81, TWEEN[®] 85, Tyloxapol, n-Undecyl beta-D-glucopyranoside, semi-synthetic derivatives thereof, or combinations thereof.

In addition, the nonionic surfactant can be a poloxamer. Poloxamers are polymers made of a block of polyoxyethylene, followed by a block of polyoxypropylene, followed by a block of polyoxyethylene. The average number of units of polyoxyethylene and

polyoxypropylene varies based on the number associated with the polymer. For example, the smallest polymer, Poloxamer 101, consists of a block with an average of 2 units of polyoxyethylene, a block with an average of 16 units of polyoxypropylene, followed by a block with an average of 2 units of polyoxyethylene. Poloxamers range from colorless liquids and pastes to white solids. In cosmetics and personal care products, Poloxamers are used in the formulation of skin cleansers, bath products, shampoos, hair conditioners, mouthwashes, eye makeup remover and other skin and hair products. Examples of Poloxamers include, but are not limited to, Poloxamer 101, Poloxamer 105, Poloxamer 108, Poloxamer 122, Poloxamer 123, Poloxamer 124, Poloxamer 181, Poloxamer 182, Poloxamer 183, Poloxamer 184, Poloxamer 185, Poloxamer 188, Poloxamer 212, Poloxamer 215, Poloxamer 217, Poloxamer 231, Poloxamer 234, Poloxamer 235, Poloxamer 237, Poloxamer 238, Poloxamer 282, Poloxamer 284, Poloxamer 288, Poloxamer 331, Poloxamer 333, Poloxamer 334, Poloxamer 335, Poloxamer 338, Poloxamer 401, Poloxamer 402, Poloxamer 403, Poloxamer 407, Poloxamer 105 Benzoate, and Poloxamer 182 Dibenzoate.

Suitable cationic surfactants include, but are not limited to, a quarternary ammonium compound, an alkyl trimethyl ammonium chloride compound, a dialkyl dimethyl ammonium chloride compound, a cationic halogen-containing compound, such as cetylpyridinium chloride, Benzalkonium chloride, Benzalkonium chloride, Benzyldimethylhexadecylammonium chloride, Benzyldimethyltetradecylammonium chloride, Benzyldodecyldimethylammonium bromide, Benzyltrimethylammonium tetrachloroiodate, Dimethyldioctadecylammonium bromide, Dodecylethyl dimethylammonium bromide, Dodecyltrimethylammonium bromide, Dodecyltrimethylammonium bromide, Ethylhexadecyldimethylammonium bromide, Girard's reagent T, Hexadecyltrimethylammonium bromide, Hexadecyltrimethylammonium bromide, N,N',N'-Polyoxyethylene(10)-N-tallow-1,3-diaminopropane, Thonzonium bromide, Trimethyl(tetradecyl)ammonium bromide, 1,3,5-Triazine-1,3,5(2H,4H,6H)-triethanol, 1-Decanaminium, N-decyl-N, N-dimethyl-, chloride, Didecyl dimethyl ammonium chloride, 2-(2-(p-(Diisobutyl)cresosy)ethoxy)ethyl dimethyl benzyl ammonium chloride, 2-(2-(p-(Diisobutyl)phenoxy)ethoxy)ethyl dimethyl benzyl ammonium chloride, Alkyl 1 or 3 benzyl-1-(2-hydroxyethyl)-2-imidazolinium chloride, Alkyl bis(2-hydroxyethyl) benzyl ammonium chloride, Alkyl demethyl benzyl ammonium chloride, Alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride (100% C12), Alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride

(50% C14, 40% C12, 10% C16), Alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride (55% C14, 23% C12, 20% C16), Alkyl dimethyl benzyl ammonium chloride, Alkyl dimethyl benzyl ammonium chloride (100% C14), Alkyl dimethyl benzyl ammonium chloride (100% C16), Alkyl dimethyl benzyl ammonium chloride (41% C14, 28% C12), Alkyl dimethyl benzyl ammonium chloride (47% C12, 18% C14), Alkyl dimethyl benzyl ammonium chloride (55% C16, 20% C14), Alkyl dimethyl benzyl ammonium chloride (58% C14, 28% C16), Alkyl dimethyl benzyl ammonium chloride (60% C14, 25% C12), Alkyl dimethyl benzyl ammonium chloride (61% C11, 23% C14), Alkyl dimethyl benzyl ammonium chloride (61% C12, 23% C14), Alkyl dimethyl benzyl ammonium chloride (65% C12, 25% C14), Alkyl dimethyl benzyl ammonium chloride (67% C12, 24% C14), Alkyl dimethyl benzyl ammonium chloride (67% C12, 25% C14), Alkyl dimethyl benzyl ammonium chloride (90% C14, 5% C12), Alkyl dimethyl benzyl ammonium chloride (93% C14, 4% C12), Alkyl dimethyl benzyl ammonium chloride (95% C16, 5% C18), Alkyl dimethyl benzyl ammonium chloride, Alkyl didecyl dimethyl ammonium chloride, Alkyl dimethyl benzyl ammonium chloride, Alkyl dimethyl benzyl ammonium chloride (C12-16), Alkyl dimethyl benzyl ammonium chloride (C12-18), Alkyl dimethyl benzyl ammonium chloride, dialkyl dimethyl benzyl ammonium chloride, Alkyl dimethyl dimethybenzyl ammonium chloride, Alkyl dimethyl ethyl ammonium bromide (90% C14, 5% C16, 5% C12), Alkyl dimethyl ethyl ammonium bromide (mixed alkyl and alkenyl groups as in the fatty acids of soybean oil), Alkyl dimethyl ethylbenzyl ammonium chloride, Alkyl dimethyl ethylbenzyl ammonium chloride (60% C14), Alkyl dimethyl isopropylbenzyl ammonium chloride (50% C12, 30% C14, 17% C16, 3% C18), Alkyl trimethyl ammonium chloride (58% C18, 40% C16, 1% C14, 1% C12), Alkyl trimethyl ammonium chloride (90% C18, 10% C16), Alkyldimethyl(ethylbenzyl) ammonium chloride (C12-18), Di-(C8-10)-alkyl dimethyl ammonium chlorides, Dialkyl dimethyl ammonium chloride, Dialkyl methyl benzyl ammonium chloride, Didecyl dimethyl ammonium chloride, Diisodecyl dimethyl ammonium chloride, Dioctyl dimethyl ammonium chloride, Dodecyl bis (2-hydroxyethyl) octyl hydrogen ammonium chloride, Dodecyl dimethyl benzyl ammonium chloride, Dodecylcarbamoyl methyl dinethyl benzyl ammonium chloride, Heptadecyl hydroxyethylimidazolinium chloride, Hexahydro-1,3,5 – tris(2-hydroxyethyl)-s-triazine, Hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine, Myristalkonium chloride (and) Quat RNIUM 14, N,N-Dimethyl-2-hydroxypropylammonium chloride polymer, n-Tetradecyl dimethyl benzyl ammonium

chloride monohydrate, Octyl decyl dimethyl ammonium chloride, Octyl dodecyl dimethyl ammonium chloride, Octyphenoxyethoxyethyl dimethyl benzyl ammonium chloride, Oxydiethylenebis(alkyl dimethyl ammonium chloride), Quaternary ammonium compounds, dicoco alkyldimethyl, chloride, Trimethoxysilyl propyl dimethyl octadecyl ammonium
 5 chloride, Trimethoxysilyl quats, Trimethyl dodecylbenzyl ammonium chloride, semi-synthetic derivatives thereof, and combinations thereof.

Exemplary cationic halogen-containing compounds include, but are not limited to, cetylpyridinium halides, cetyltrimethylammonium halides, cetyldimethylethylammonium halides, cetyldimethylbenzylammonium halides, cetyltributylphosphonium halides,
 10 dodecyltrimethylammonium halides, or tetradecyltrimethylammonium halides. In some particular embodiments, suitable cationic halogen containing compounds comprise, but are not limited to, cetylpyridinium chloride (CPC), cetyltrimethylammonium chloride, cetylbenzyltrimethylammonium chloride, cetylpyridinium bromide (CPB), cetyltrimethylammonium bromide (CTAB), cetyldimethylethylammonium bromide,
 15 cetyltributylphosphonium bromide, dodecyltrimethylammonium bromide, and tetradecyltrimethylammonium bromide. In particularly preferred embodiments, the cationic halogen containing compound is CPC, although the compositions of the present invention are not limited to formulation with a particular cationic containing compound.

Suitable anionic surfactants include, but are not limited to, a carboxylate, a sulphate, a
 20 sulphonate, a phosphate, chenodeoxycholic acid, chenodeoxycholic acid sodium salt, cholic acid, ox or sheep bile, Dehydrocholic acid, Deoxycholic acid, Deoxycholic acid, Deoxycholic acid methyl ester, Digitonin, Digitoxigenin, N,N-Dimethyldodecylamine N-oxide, Docusate sodium salt, Glycochenodeoxycholic acid sodium salt, Glycocholic acid hydrate, synthetic, Glycocholic acid sodium salt hydrate, synthetic, Glycodeoxycholic acid monohydrate,
 25 Glycodeoxycholic acid sodium salt, Glycodeoxycholic acid sodium salt, Glycolithocholic acid 3-sulfate disodium salt, Glycolithocholic acid ethyl ester, N-Lauroylsarcosine sodium salt, N-Lauroylsarcosine solution, N-Lauroylsarcosine solution, Lithium dodecyl sulfate, Lithium dodecyl sulfate, Lithium dodecyl sulfate, Lugol solution, Niaproof 4, Type 4, 1-Octanesulfonic acid sodium salt, Sodium 1-butanesulfonate, Sodium 1-decanesulfonate,
 30 Sodium 1-decanesulfonate, Sodium 1-dodecanesulfonate, Sodium 1-heptanesulfonate anhydrous, Sodium 1-heptanesulfonate anhydrous, Sodium 1-nonanesulfonate, Sodium 1-propanesulfonate monohydrate, Sodium 2-bromoethanesulfonate, Sodium cholate hydrate,

Sodium choleate, Sodium deoxycholate, Sodium deoxycholate monohydrate, Sodium dodecyl sulfate, Sodium hexanesulfonate anhydrous, Sodium octyl sulfate, Sodium pentanesulfonate anhydrous, Sodium taurocholate, Taurochenodeoxycholic acid sodium salt, Taurodeoxycholic acid sodium salt monohydrate, Taurohyodeoxycholic acid sodium salt hydrate, Tauroolithocholic acid 3-sulfate disodium salt, Tauroursodeoxycholic acid sodium salt, Trizma[®] dodecyl sulfate, TWEEN[®] 80, Ursodeoxycholic acid, semi-synthetic derivatives thereof, and combinations thereof.

Suitable zwitterionic surfactants include, but are not limited to, an N-alkyl betaine, lauryl amido propyl dimethyl betaine, an alkyl dimethyl glycinate, an N-alkyl amino propionate, CHAPS, minimum 98% (TLC), CHAPS, SigmaUltra, minimum 98% (TLC), CHAPS, for electrophoresis, minimum 98% (TLC), CHAPSO, minimum 98%, CHAPSO, SigmaUltra, CHAPSO, for electrophoresis, 3-(Decyldimethylammonio)propanesulfonate inner salt, 3-Dodecyldimethylammonio)propanesulfonate inner salt, SigmaUltra, 3-(Dodecyldimethylammonio)propanesulfonate inner salt, 3-(N,N-Dimethylmyristylammonio)propanesulfonate, 3-(N,N-Dimethyloctadecylammonio)propanesulfonate, 3-(N,N-Dimethyloctylammonio)propanesulfonate inner salt, 3-(N,N-Dimethylpalmitylammonio)propanesulfonate, semi-synthetic derivatives thereof, and combinations thereof.

The present invention is not limited to the surfactants disclosed herein. Additional surfactants and detergents useful in the compositions of the present invention may be ascertained from reference works (*e.g.*, including, but not limited to, McCutcheon's Volume 1: Emulsions and Detergents - North American Edition, 2000) and commercial sources.

4. Cationic Halogens Containing Compounds

In some embodiments, the emulsions further comprise a cationic halogen containing compound. In some preferred embodiments, the emulsion comprises from about 0.5 to 1.0 wt. % or more of a cationic halogen containing compound, based on the total weight of the emulsion (although other concentrations are also contemplated). In preferred embodiments, the cationic halogen-containing compound is preferably premixed with the oil phase; however, it should be understood that the cationic halogen-containing compound may be provided in combination with the emulsion composition in a distinct formulation. Suitable

halogen containing compounds may be selected from compounds comprising chloride, fluoride, bromide and iodide ions. In preferred embodiments, suitable cationic halogen containing compounds include, but are not limited to, cetylpyridinium halides, cetyltrimethylammonium halides, cetyldimethylethylammonium halides, cetyldimethylbenzylammonium halides, cetyltributylphosphonium halides, dodecyltrimethylammonium halides, or tetradecyltrimethylammonium halides. In some particular embodiments, suitable cationic halogen containing compounds comprise, but are not limited to, cetylpyridinium chloride (CPC), cetyltrimethylammonium chloride, cetylbenzyltrimethylammonium chloride, cetylpyridinium bromide (CPB), and cetyltrimethylammonium bromide (CTAB), cetyldimethylethylammonium bromide, cetyltributylphosphonium bromide, dodecyltrimethylammonium bromide, and tetradecyltrimethylammonium bromide. In particularly preferred embodiments, the cationic halogen-containing compound is CPC, although the compositions of the present invention are not limited to formulation with any particular cationic containing compound.

5. Germination Enhancers

In other embodiments of the present invention, the nanoemulsions further comprise a germination enhancer. In some preferred embodiments, the emulsions comprise from about 1 mM to 15 mM, and more preferably from about 5 mM to 10 mM of one or more germination enhancing compounds (although other concentrations are also contemplated). In preferred embodiments, the germination enhancing compound is provided in the aqueous phase prior to formation of the emulsion. The present invention contemplates that when germination enhancers are added to the nanoemulsion compositions, the sporicidal properties of the nanoemulsions are enhanced. The present invention further contemplates that such germination enhancers initiate sporicidal activity near neutral pH (between pH 6 - 8, and preferably 7). Such neutral pH emulsions can be obtained, for example, by diluting with phosphate buffer saline (PBS) or by preparations of neutral emulsions. The sporicidal activity of the nanoemulsion preferentially occurs when the spores initiate germination.

In specific embodiments, it has been demonstrated that the emulsions utilized in the vaccines of the present invention have sporicidal activity. While the present invention is not limited to any particular mechanism and an understanding of the mechanism is not required to practice the present invention, it is believed that the fusigenic component of the emulsions

acts to initiate germination and before reversion to the vegetative form is complete the lysogenic component of the emulsion acts to lyse the newly germinating spore. These components of the emulsion thus act in concert to leave the spore susceptible to disruption by the emulsions. The addition of germination enhancer further facilitates the anti-sporicidal activity of the emulsions, for example, by speeding up the rate at which the sporicidal activity occurs.

Germination of bacterial endospores and fungal spores is associated with increased metabolism and decreased resistance to heat and chemical reactants. For germination to occur, the spore must sense that the environment is adequate to support vegetation and reproduction. The amino acid L-alanine stimulates bacterial spore germination (*See e.g.*, Hills, J. Gen. Micro. 4:38 (1950); and Halvorson and Church, Bacteriol Rev. 21:112 (1957)). L-alanine and L-proline have also been reported to initiate fungal spore germination (Yanagita, Arch Mikrobiol 26:329 (1957)). Simple α -amino acids, such as glycine and L-alanine, occupy a central position in metabolism. Transamination or deamination of α -amino acids yields the glycogenic or ketogenic carbohydrates and the nitrogen needed for metabolism and growth. For example, transamination or deamination of L-alanine yields pyruvate, which is the end product of glycolytic metabolism (Embden-Meyerhof Pathway). Oxidation of pyruvate by pyruvate dehydrogenase complex yields acetyl-CoA, NADH, H^+ , and CO_2 . Acetyl-CoA is the initiator substrate for the tricarboxylic acid cycle (Kreb's Cycle), which in turns feeds the mitochondrial electron transport chain. Acetyl-CoA is also the ultimate carbon source for fatty acid synthesis as well as for sterol synthesis. Simple α -amino acids can provide the nitrogen, CO_2 , glycogenic and/or ketogenic equivalents required for germination and the metabolic activity that follows.

In certain embodiments, suitable germination enhancing agents of the invention include, but are not limited to, α -amino acids comprising glycine and the L-enantiomers of alanine, valine, leucine, isoleucine, serine, threonine, lysine, phenylalanine, tyrosine, and the alkyl esters thereof. Additional information on the effects of amino acids on germination may be found in U.S. Pat. No. 5,510,104. In some embodiments, a mixture of glucose, fructose, asparagine, sodium chloride (NaCl), ammonium chloride (NH_4Cl), calcium chloride ($CaCl_2$) and potassium chloride (KCl) also may be used. In particularly preferred embodiments of the present invention, the formulation

comprises the germination enhancers L-alanine, CaCl_2 , Inosine and NH_4Cl . In some embodiments, the compositions further comprise one or more common forms of growth media (*e.g.*, trypticase soy broth, and the like) that additionally may or may not itself comprise germination enhancers and buffers.

5 The above compounds are merely exemplary germination enhancers and it is understood that other known germination enhancers will find use in the nanoemulsions utilized in some embodiments of the present invention. A candidate germination enhancer should meet two criteria for inclusion in the compositions of the present invention: it should be capable of being associated with the emulsions disclosed herein and it should increase the
10 rate of germination of a target spore when incorporated in the emulsions disclosed herein. One skilled in the art can determine whether a particular agent has the desired function of acting as an germination enhancer by applying such an agent in combination with the nanoemulsions disclosed herein to a target and comparing the inactivation of the target when contacted by the admixture with inactivation of like targets by the composition of the present
15 invention without the agent. Any agent that increases germination, and thereby decreases or inhibits the growth of the organisms, is considered a suitable enhancer for use in the nanoemulsion compositions disclosed herein.

 In still other embodiments, addition of a germination enhancer (or growth medium) to a neutral emulsion composition produces a composition that is useful in inactivating bacterial
20 spores in addition to enveloped viruses, Gram negative bacteria, and Gram positive bacteria for use in the vaccine compositions of the present invention.

6. Interaction Enhancers

 In still other embodiments, nanoemulsions comprise one or more compounds capable
25 of increasing the interaction of the compositions (*i.e.*, "interaction enhancer") with target pathogens (*e.g.*, the cell wall of Gram negative bacteria such as *Vibrio*, *Salmonella*, *Shigella* and *Pseudomonas*). In preferred embodiments, the interaction enhancer is preferably premixed with the oil phase; however, in other embodiments the interaction enhancer is provided in combination with the compositions after emulsification. In certain preferred
30 embodiments, the interaction enhancer is a chelating agent (*e.g.*, ethylenediaminetetraacetic acid (EDTA) or ethylenebis(oxyethylenenitrilo)tetraacetic acid (EGTA) in a buffer (*e.g.*, tris buffer)). It is understood that chelating agents are merely exemplary interaction enhancing

compounds. Indeed, other agents that increase the interaction of the nanoemulsions used in some embodiments of the present invention with microbial agents and/or pathogens are contemplated. In particularly preferred embodiments, the interaction enhancer is at a concentration of about 50 to about 250 μ M. One skilled in the art will be able to determine whether a particular agent has the desired function of acting as an interaction enhancer by applying such an agent in combination with the compositions of the present invention to a target and comparing the inactivation of the target when contacted by the admixture with inactivation of like targets by the composition of the present invention without the agent. Any agent that increases the interaction of an emulsion with bacteria and thereby decreases or inhibits the growth of the bacteria, in comparison to that parameter in its absence, is considered an interaction enhancer.

In some embodiments, the addition of an interaction enhancer to nanoemulsion produces a composition that is useful in inactivating enveloped viruses, some Gram positive bacteria and some Gram negative bacteria for use in the vaccine compositions of the present invention.

7. Quaternary Ammonium Compounds

In some embodiments, nanoemulsions of the present invention include a quaternary ammonium containing compound. Exemplary quaternary ammonium compounds include, but are not limited to, Alkyl dimethyl benzyl ammonium chloride, didecyl dimethyl ammonium chloride, Alkyl dimethyl benzyl and dialkyl dimethyl ammonium chloride, N,N-Dimethyl-2-hydroxypropylammonium chloride polymer, Didecyl dimethyl ammonium chloride, n-Alkyl dimethyl benzyl ammonium chloride, n-Alkyl dimethyl ethylbenzyl ammonium chloride,

Dialkyl dimethyl ammonium chloride, n-Alkyl dimethyl benzyl ammonium chloride, n-Tetradecyl dimethyl benzyl ammonium chloride monohydrate, n-Alkyl dimethyl benzyl ammonium chloride, Dialkyl dimethyl ammonium chloride, Hexahydro-1,3,5 – tris(2-hydroxyethyl)-s-triazine, Myristalkonium chloride (and) Quat RNIUM 14, Alkyl bis(2-hydroxyethyl) benzyl ammonium chloride, Alkyl demethyl benzyl ammonium chloride, Alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride, Alkyl dimethyl benzyl ammonium chloride, Alkyl dimethyl benzyl dimethylbenzyl ammonium, Alkyl dimethyl dimethybenzyl ammonium chloride, Alkyl dimethyl ethyl ammonium bromide, Alkyl dimethyl ethyl

ammonium bromide, Alkyl dimethyl ethylbenzyl ammonium chloride, Alkyl dimethyl isopropylbenzyl ammonium chloride, Alkyl trimethyl ammonium chloride, Alkyl 1 or 3 benzyl-1-(2-hydroxyethyl)-2-imidazolinium chloride, Dialkyl methyl benzyl ammonium chloride, Dialkyl dimethyl ammonium chloride, Didecyl dimethyl ammonium chloride, 2-(2-
 5 (p-(Diisobutyl)cresosy)ethoxy)ethyl dimethyl benzyl ammonium chloride, 2-(2-(p-(Diisobutyl)phenoxy)ethoxy)ethyl dimethyl benzyl ammonium chloride, Dioctyl dimethyl ammonium chloride, Dodecyl bis (2-hydroxyethyl) octyl hydrogen ammonium chloride, Dodecyl dimethyl benzyl ammonium chloride, Dodecylcarbamoyl methyl dinethyl benzyl ammonium chloride, Heptadecyl hydroxyethylimidazolinium chloride, Hexahydro-1,3,5-
 10 tris(2-hydroxyethyl)-s-triazine, Octyl decyl dimethyl ammonium chloride, Octyl dodecyl dimethyl ammonium chloride, Octyphenoxyethoxyethyl dimethyl benzyl ammonium chloride, Oxydiethylenebis(alkyl dimethyl ammonium chloride), Quaternary ammonium compounds, dicoco alkyldimethyl, chloride, Trimethoxysilyl quats, and Trimethyl dodecylbenzyl ammonium chloride.

8. Other Components

In some embodiments, a nanoemulsion comprises one or more additional components that provide a desired property or functionality to the nanoemulsions. These components may be incorporated into the aqueous phase or the oil phase of the nanoemulsions and/or may be added prior to or following emulsification. For example, in some embodiments, the
 20 nanoemulsions further comprise phenols (e.g., triclosan, phenyl phenol), acidifying agents (e.g., citric acid (e.g., 1.5-6%), acetic acid, lemon juice), alkylating agents (e.g., sodium hydroxide (e.g., 0.3%)), buffers (e.g., citrate buffer, acetate buffer, and other buffers useful to maintain a specific pH), and halogens (e.g., polyvinylpyrrolidone, sodium hypochlorite, hydrogen peroxide).

Exemplary techniques for making a nanoemulsion (e.g., used to inactivate a pathogen and/or generation of an immunogenic composition of the present invention) are described below. Additionally, a number of specific, although exemplary, formulation recipes are also set forth below.

Formulation Techniques

Nanoemulsions of the present invention can be formed using classic emulsion forming techniques. In brief, the oil phase is mixed with the aqueous phase under relatively

high shear forces (e.g., using high hydraulic and mechanical forces) to obtain an oil-in-water nanoemulsion. The emulsion is formed by blending the oil phase with an aqueous phase on a volume-to-volume basis ranging from about 1:9 to 5:1, preferably about 5:1 to 3:1, most preferably 4:1, oil phase to aqueous phase. The oil and aqueous phases can be blended using any apparatus capable of producing shear forces sufficient to form an emulsion such as French Presses or high shear mixers (e.g., FDA approved high shear mixers are available, for example, from Admix, Inc., Manchester, NH). Methods of producing such emulsions are described in U.S. Pat. Nos. 5,103,497 and 4,895,452.

In preferred embodiments, compositions used in the methods of the present invention comprise droplets of an oily discontinuous phase dispersed in an aqueous continuous phase, such as water. In preferred embodiments, nanoemulsions of the present invention are stable, and do not decompose even after long storage periods (e.g., greater than one or more years). Furthermore, in some embodiments, nanoemulsions are stable (e.g., in some embodiments for greater than 3 months, in some embodiments for greater than 6 months, in some embodiments for greater than 12 months, in some embodiments for greater than 18 months) after combination with an immunogen (e.g., a pathogen). In preferred embodiments, nanoemulsions of the present invention are non-toxic and safe when administered (e.g., via spraying or contacting mucosal surfaces, swallowed, inhaled, etc.) to a subject.

In some embodiments, a portion of the emulsion may be in the form of lipid structures including, but not limited to, unilamellar, multilamellar, and paucilamellar lipid vesicles, micelles, and lamellar phases.

Some embodiments of the present invention employ an oil phase containing ethanol. For example, in some embodiments, the emulsions of the present invention contain (i) an aqueous phase and (ii) an oil phase containing ethanol as the organic solvent and optionally a germination enhancer, and (iii) TYLOXAPOL as the surfactant (preferably 2-5%, more preferably 3%). This formulation is highly efficacious for inactivation of pathogens and is also non-irritating and non-toxic to mammalian subjects (e.g., and thus can be used for administration to a mucosal surface).

In some other embodiments, the emulsions of the present invention comprise a first emulsion emulsified within a second emulsion, wherein (a) the first emulsion comprises (i) an aqueous phase; and (ii) an oil phase comprising an oil and an organic solvent; and (iii) a

5 The following description provides a number of exemplary emulsions including formulations for compositions BCTP and X₈W₆₀PC. BCTP comprises a water-in oil nanoemulsion, in which the oil phase was made from soybean oil, tri-n-butyl phosphate, and TRITON X-100 in 80% water. X₈W₆₀PC comprises a mixture of equal volumes of BCTP with W₈₀8P. W₈₀8P is a liposome-like compound made of glycerol monostearate, refined oya
10 sterols (e.g., GENEROL sterols), TWEEN 60, soybean oil, a cationic ion halogen-containing CPC and peppermint oil. The GENEROL family are a group of a polyethoxylated soya sterols (Henkel Corporation, Ambler, Pennsylvania). Exemplary emulsion formulations useful in the present invention are provided in Table 1B. These particular formulations may be found in U.S. Pat. Nos. 5,700,679 (NN); 5,618,840; 5,549,901 (W₈₀8P); and 5,547,677.
15 Certain other emulsion formulations are presented U.S. Pat. App. Serial No. 10/669,865.

The X₈W₆₀PC emulsion is manufactured by first making the W₈₀8P emulsion and BCTP emulsions separately. A mixture of these two emulsions is then re-emulsified to produce a fresh emulsion composition termed X₈W₆₀PC. Methods of producing such emulsions are described in U.S. Pat. Nos. 5,103,497 and 4,895,452.

	Oil Phase Formula	Water to Oil Phase Ratio (Vol/Vol)
BCTP	1 vol. Tri(N-butyl)phosphate 1 vol. TRITON X-100 8 vol. Soybean oil	4:1
NN	86.5 g Glycerol monooleate 60.1 ml Nonoxynol-9 24.2 g GENEROL 122 3.27 g Cetylpyridinium chloride	3:1

	554 g Soybean oil	
W ₈₀ 8P	86.5 g Glycerol monooleate 21.2 g Polysorbate 60 24.2 g GENEROL 122 3.27 g Cetylpyddinium chloride 4 ml Peppermint oil 554 g Soybean oil	3.2:1
SS	86.5 g Glycerol monooleate 21.2 g Polysorbate 60 24.2 g GENEROL 122 3.27 g Cetylpyridinium chloride 554 g Soybean oil	3.2:1 (1% bismuth in water)

The compositions listed above are only exemplary and those of skill in the art will be able to alter the amounts of the components to arrive at a nanoemulsion composition suitable for the purposes of the present invention. Those skilled in the art will understand that the ratio of oil phase to water as well as the individual oil carrier, surfactant CPC and organic phosphate buffer, components of each composition may vary.

Although certain compositions comprising BCTP have a water to oil ratio of 4:1, it is understood that the BCTP may be formulated to have more or less of a water phase. For example, in some embodiments, there is 3, 4, 5, 6, 7, 8, 9, 10, or more parts of the water phase to each part of the oil phase. The same holds true for the W₈₀8P formulation. Similarly, the ratio of Tri(N-butyl)phosphate:TRITON X-100:soybean oil also may be varied.

Although Table 1B lists specific amounts of glycerol monooleate, polysorbate 60, GENEROL 122, cetylpyridinium chloride, and carrier oil for W₈₀8P, these are merely exemplary. An emulsion that has the properties of W₈₀8P may be formulated that has different concentrations of each of these components or indeed different components that will fulfill the same function. For example, the emulsion may have between about 80 to about 100g of glycerol monooleate in the initial oil phase. In other embodiments, the emulsion may have between about 15 to about 30 g polysorbate 60 in the initial oil phase. In yet another embodiment the composition may comprise between about 20 to about 30 g of a GENEROL sterol, in the initial oil phase.

Individual components of nanoemulsions (e.g. in an immunogenic composition of the present invention) can function both to inactivate a pathogen as well as to contribute to the non-toxicity of the emulsions. For example, the active component in BCTP, TRITON-X100, shows less ability to inactivate a virus at concentrations equivalent to 11% BCTP. Adding
5 the oil phase to the detergent and solvent markedly reduces the toxicity of these agents in tissue culture at the same concentrations. While not being bound to any theory (an understanding of the mechanism is not necessary to practice the present invention, and the present invention is not limited to any particular mechanism), it is suggested that the nanoemulsion enhances the interaction of its components with the pathogens thereby
10 facilitating the inactivation of the pathogen and reducing the toxicity of the individual components. Furthermore, when all the components of BCTP are combined in one composition but are not in a nanoemulsion structure, the mixture is not as effective at inactivating a pathogen as when the components are in a nanoemulsion structure.

Numerous additional embodiments presented in classes of formulations with like
15 compositions are presented below. The following compositions recite various ratios and mixtures of active components. One skilled in the art will appreciate that the below recited formulation are exemplary and that additional formulations comprising similar percent ranges of the recited components are within the scope of the present invention.

In certain embodiments of the present invention, a nanoemulsion comprises from
20 about 3 to 8 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of cetylpyridinium chloride (CPC), about 60 to 70 vol. % oil (e.g., soybean oil), about 15 to 25 vol. % of aqueous phase (e.g., DiH₂O or PBS), and in some formulations less than about 1 vol. % of 1N NaOH. Some of these embodiments comprise PBS. It is contemplated that the addition of 1N NaOH and/or PBS in some of these embodiments, allows the user to
25 advantageously control the pH of the formulations, such that pH ranges from about 7.0 to about 9.0, and more preferably from about 7.1 to 8.5 are achieved. For example, one embodiment of the present invention comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 24 vol. % of DiH₂O (designated herein as Y3EC). Another similar embodiment comprises about 3.5
30 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, and about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 23.5 vol. % of DiH₂O (designated herein as Y3.5EC). Yet another embodiment comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol,

about 1 vol. % of CPC, about 0.067 vol. % of 1N NaOH, such that the pH of the formulation is about 7.1, about 64 vol. % of soybean oil, and about 23.93 vol. % of DiH₂O (designated herein as Y3EC pH 7.1). Still another embodiment comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 0.67 vol. % of 1N NaOH, such that the pH of the formulation is about 8.5, and about 64 vol. % of soybean oil, and about 23.33 vol. % of DiH₂O (designated herein as Y3EC pH 8.5). Another similar embodiment comprises about 4% TYLOXAPOL, about 8 vol. % ethanol, about 1% CPC, and about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as Y4EC). In still another embodiment the formulation comprises about 8% TYLOXAPOL, about 8% ethanol, about 1 vol. % of CPC, and about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as Y8EC). A further embodiment comprises about 8 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 19 vol. % of 1x PBS (designated herein as Y8EC PBS).

In some embodiments of the present invention, a nanoemulsion comprises about 8 vol. % of ethanol, and about 1 vol. % of CPC, and about 64 vol. % of oil (e.g., soybean oil), and about 27 vol. % of aqueous phase (e.g., DiH₂O or PBS) (designated herein as EC).

In some embodiments, a nanoemulsion comprises from about 8 vol. % of sodium dodecyl sulfate (SDS), about 8 vol. % of tributyl phosphate (TBP), and about 64 vol. % of oil (e.g., soybean oil), and about 20 vol. % of aqueous phase (e.g., DiH₂O or PBS) (designated herein as S8P).

In some embodiments, a nanoemulsion comprises from about 1 to 2 vol. % of TRITON X-100, from about 1 to 2 vol. % of TYLOXAPOL, from about 7 to 8 vol. % of ethanol, about 1 vol. % of cetylpyridinium chloride (CPC), about 64 to 57.6 vol. % of oil (e.g., soybean oil), and about 23 vol. % of aqueous phase (e.g., DiH₂O or PBS).

Additionally, some of these formulations further comprise about 5 mM of L-alanine/Inosine, and about 10 mM ammonium chloride. Some of these formulations comprise PBS. It is contemplated that the addition of PBS in some of these embodiments, allows the user to advantageously control the pH of the formulations. For example, one embodiment of the present invention comprises about 2 vol. % of TRITON X-100, about 2 vol. % of TYLOXAPOL, about 8 vol. % of ethanol, about 1 vol. % CPC, about 64 vol. % of soybean oil, and about 23 vol. % of aqueous phase DiH₂O. In another embodiment the formulation comprises about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2

vol. % of ethanol, about 0.9 vol. % of CPC, about 5 mM L-alanine/Inosine, and about 10 mM ammonium chloride, about 57.6 vol. % of soybean oil, and the remainder of 1x PBS (designated herein as 90% X2Y2EC/GE).

5 In some embodiments, a nanoemulsion comprises from about 5 vol. % of TWEEN 80, from about 8 vol. % of ethanol, from about 1 vol. % of CPC, about 64 vol. % of oil (e.g., soybean oil), and about 22 vol. % of DiH₂O (designated herein as W₈₀5EC).

In still other embodiments of the present invention, a nanoemulsion comprises from about 5 vol. % of TWEEN 20, from about 8 vol. % of ethanol, from about 1 vol. % of CPC, about 64 vol. % of oil (e.g., soybean oil), and about 22 vol. % of DiH₂O (designated herein as
10 W₂₀5EC).

In still other embodiments of the present invention, a nanoemulsion comprises from about 2 to 8 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 1 vol. % of CPC, about 60 to 70 vol. % of oil (e.g., soybean, or olive oil), and about 15 to 25 vol. % of aqueous phase (e.g., DiH₂O or PBS). For example, the present invention contemplates formulations
15 comprising about 2 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 26 vol. % of DiH₂O (designated herein as X2E). In other similar embodiments, a nanoemulsion comprises about 3 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 25 vol. % of DiH₂O (designated herein as X3E). In still further embodiments, the formulations comprise about 4 vol. % Triton of X-
20 100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 24 vol. % of DiH₂O (designated herein as X4E). In yet other embodiments, a nanoemulsion comprises about 5 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as X5E). In some embodiments, a nanoemulsion comprises about 6 vol. % of TRITON X-100, about 8 vol. % of ethanol, about
25 64 vol. % of soybean oil, and about 22 vol. % of DiH₂O (designated herein as X6E). In still further embodiments of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8E). In still further embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of ethanol, about 64 vol. % of
30 olive oil, and about 20 vol. % of DiH₂O (designated herein as X8E O). In yet another embodiment, a nanoemulsion comprises 8 vol. % of TRITON X-100, about 8 vol. % ethanol, about 1 vol. % CPC, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O

(designated herein as X8EC).

In alternative embodiments of the present invention, a nanoemulsion comprises from about 1 to 2 vol. % of TRITON X-100, from about 1 to 2 vol. % of TYLOXAPOL, from about 6 to 8 vol. % TBP, from about 0.5 to 1.0 vol. % of CPC, from about 60 to 70 vol. % of oil (e.g., soybean), and about 1 to 35 vol. % of aqueous phase (e.g., DiH₂O or PBS).

Additionally, certain of these nanoemulsions may comprise from about 1 to 5 vol. % of trypticase soy broth, from about 0.5 to 1.5 vol. % of yeast extract, about 5 mM L-alanine/Inosine, about 10 mM ammonium chloride, and from about 20-40 vol. % of liquid baby formula. In some embodiments comprising liquid baby formula, the formula comprises

a casein hydrolysate (e.g., Neutramigen, or Progestimil, and the like). In some of these embodiments, a nanoemulsion further comprises from about 0.1 to 1.0 vol. % of sodium thiosulfate, and from about 0.1 to 1.0 vol. % of sodium citrate. Other similar embodiments comprising these basic components employ phosphate buffered saline (PBS) as the aqueous phase. For example, one embodiment comprises about 2 vol. % of TRITON X-100, about 2

vol. % TYLOXAPOL, about 8 vol. % TBP, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 23 vol. % of DiH₂O (designated herein as X2Y2EC). In still other embodiments, the inventive formulation comprises about 2 vol. % of TRITON X-100, about 2 vol. % TYLOXAPOL, about 8 vol. % TBP, about 1 vol. % of CPC, about 0.9 vol. % of sodium thiosulfate, about 0.1 vol. % of sodium citrate, about 64 vol. % of soybean oil, and

about 22 vol. % of DiH₂O (designated herein as X2Y2PC STS1). In another similar embodiment, a nanoemulsion comprises about 1.7 vol. % TRITON X-100, about 1.7 vol. % TYLOXAPOL, about 6.8 vol. % TBP, about 0.85% CPC, about 29.2% NEUTRAMIGEN, about 54.4 vol. % of soybean oil, and about 4.9 vol. % of DiH₂O (designated herein as 85% X2Y2PC/baby). In yet another embodiment of the present invention, a nanoemulsion

comprises about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2 vol. % of TBP, about 0.9 vol. % of CPC, about 5mM L-alanine/Inosine, about 10mM ammonium chloride, about 57.6 vol. % of soybean oil, and the remainder vol. % of 0.1x PBS (designated herein as 90% X2Y2 PC/GE). In still another embodiment, a nanoemulsion

comprises about 1.8 vol. % of TRITON X-100, about 1.8 vol. % of TYLOXAPOL, about 7.2 vol. % TBP, about 0.9 vol. % of CPC, and about 3 vol. % trypticase soy broth, about 57.6 vol. % of soybean oil, and about 27.7 vol. % of DiH₂O (designated herein as 90%

X2Y2PC/TSB). In another embodiment of the present invention, a nanoemulsion comprises

about 1.8 vol. % TRITON X-100, about 1.8 vol. % TYLOXAPOL, about 7.2 vol. % TBP, about 0.9 vol. % CPC, about 1 vol. % yeast extract, about 57.6 vol. % of soybean oil, and about 29.7 vol. % of DiH₂O (designated herein as 90% X2Y2PC/YE).

In some embodiments of the present invention, a nanoemulsion comprises about 3
5 vol. % of TYLOXAPOL, about 8 vol. % of TBP, and about 1 vol. % of CPC, about 60 to 70 vol. % of oil (e.g., soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (e.g., DiH₂O or PBS). In a particular embodiment of the present invention, a nanoemulsion comprises about 3 vol. % of TYLOXAPOL, about 8 vol. % of TBP, and about 1 vol. % of CPC, about 64 vol. % of soybean, and about 24 vol. % of DiH₂O (designated herein as
10 Y3PC).

In some embodiments of the present invention, a nanoemulsion comprises from about 4 to 8 vol. % of TRITON X-100, from about 5 to 8 vol. % of TBP, about 30 to 70 vol. % of oil (e.g., soybean or olive oil), and about 0 to 30 vol. % of aqueous phase (e.g., DiH₂O or PBS). Additionally, certain of these embodiments further comprise about 1 vol. % of CPC,
15 about 1 vol. % of benzalkonium chloride, about 1 vol. % cetylridinium bromide, about 1 vol. % cetyldimethyletylammonium bromide, 500 μ M EDTA, about 10 mM ammonium chloride, about 5 mM Inosine, and about 5 mM L-alanine. For example, in a certain preferred embodiment, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated
20 herein as X8P). In another embodiment of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1% of CPC, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as X8PC). In still another embodiment, a nanoemulsion comprises about 8 vol. % TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O
25 (designated herein as ATB-X1001). In yet another embodiment, the formulations comprise about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 2 vol. % of CPC, about 50 vol. % of soybean oil, and about 32 vol. % of DiH₂O (designated herein as ATB-X002). In some embodiments, a nanoemulsion comprises about 4 vol. % TRITON X-100, about 4 vol. % of TBP, about 0.5 vol. % of CPC, about 32 vol. % of soybean oil, and about 59.5 vol. % of
30 DiH₂O (designated herein as 50% X8PC). In some embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 0.5 vol. % CPC, about 64 vol. % of soybean oil, and about 19.5 vol. % of DiH₂O (designated herein as X8PC_{1/2}). In

some embodiments of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 2 vol. % of CPC, about 64 vol. % of soybean oil, and about 18 vol. % of DiH₂O (designated herein as X8PC2). In other embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8% of TBP, about 1% of benzalkonium chloride, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as X8P BC). In an alternative embodiment of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of cetylyridinium bromide, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as X8P CPB). In another exemplary embodiment of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of cetyldimethyletylammonium bromide, about 50 vol. % of soybean oil, and about 33 vol. % of DiH₂O (designated herein as X8P CTAB). In still further embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 500 μ M EDTA, about 64 vol. % of soybean oil, and about 15.8 vol. % DiH₂O (designated herein as X8PC EDTA). In some embodiments, a nanoemulsion comprises 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 1 vol. % of CPC, about 10 mM ammonium chloride, about 5mM Inosine, about 5mM L-alanine, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O or PBS (designated herein as X8PC GE_{1x}). In another embodiment of the present invention, a nanoemulsion comprises about 5 vol. % of TRITON X-100, about 5% of TBP, about 1 vol. % of CPC, about 40 vol. % of soybean oil, and about 49 vol. % of DiH₂O (designated herein as X5P₅C).

In some embodiments of the present invention, a nanoemulsion comprises about 2 vol. % TRITON X-100, about 6 vol. % TYLOXAPOL, about 8 vol. % ethanol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X2Y6E).

In an additional embodiment of the present invention, a nanoemulsion comprises about 8 vol. % of TRITON X-100, and about 8 vol. % of glycerol, about 60 to 70 vol. % of oil (e.g., soybean or olive oil), and about 15 to 25 vol. % of aqueous phase (e.g., DiH₂O or PBS). Certain nanoemulsion compositions (e.g., used to generate an immune response (e.g., for use as a vaccine) comprise about 1 vol. % L-ascorbic acid. For example, one particular embodiment comprises about 8 vol. % of TRITON X-100, about 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8G). In still another embodiment, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 8

vol. % of glycerol, about 1 vol. % of L-ascorbic acid, about 64 vol. % of soybean oil, and about 19 vol. % of DiH_2O (designated herein as X8GV_c).

In still further embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, from about 0.5 to 0.8 vol. % of TWEEN 60, from about 0.5 to 2.0 vol. % of CPC, about 8 vol. % of TBP, about 60 to 70 vol. % of oil (e.g., soybean or olive oil), and about 15 to 25 vol. % of aqueous phase (e.g., DiH_2O or PBS). For example, in one particular embodiment a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 0.70 vol. % of TWEEN 60, about 1 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 18.3 vol. % of DiH_2O (designated herein as X8W60PC₁). In some
10 embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 0.71 vol. % of TWEEN 60, about 1 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 18.29 vol. % of DiH_2O (designated herein as W60_{0.7}X8PC). In yet other embodiments, a nanoemulsion comprises from about 8 vol. % of TRITON X-100, about 0.7 vol. % of TWEEN 60, about 0.5 vol. % of CPC, about 8 vol. % of TBP, about 64 to 70 vol. %
15 of soybean oil, and about 18.8 vol. % of DiH_2O (designated herein as X8W60PC₂). In still other embodiments, a nanoemulsion comprises about 8 vol. % of TRITON X-100, about 0.71 vol. % of TWEEN 60, about 2 vol. % of CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 17.3 vol. % of DiH_2O . In another embodiment of the present invention, a nanoemulsion comprises about 0.71 vol. % of TWEEN 60, about 1 vol. % of
20 CPC, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 25.29 vol. % of DiH_2O (designated herein as W60_{0.7}PC).

In another embodiment of the present invention, a nanoemulsion comprises about 2 vol. % of dioctyl sulfosuccinate, either about 8 vol. % of glycerol, or about 8 vol. % TBP, in addition to, about 60 to 70 vol. % of oil (e.g., soybean or olive oil), and about 20 to 30 vol. %
25 of aqueous phase (e.g., DiH_2O or PBS). For example, in some embodiments, a nanoemulsion comprises about 2 vol. % of dioctyl sulfosuccinate, about 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 26 vol. % of DiH_2O (designated herein as D2G). In another related embodiment, a nanoemulsion comprises about 2 vol. % of dioctyl sulfosuccinate, and about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 26 vol. % of DiH_2O
30 (designated herein as D2P).

In still other embodiments of the present invention, a nanoemulsion comprises about 8 to 10 vol. % of glycerol, and about 1 to 10 vol. % of CPC, about 50 to 70 vol. % of oil (e.g.,

soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (e.g., DiH₂O or PBS).

Additionally, in certain of these embodiments, a nanoemulsion further comprises about 1 vol. % of L-ascorbic acid. For example, in some embodiments, a nanoemulsion comprises about 8 vol. % of glycerol, about 1 vol. % of CPC, about 64 vol. % of soybean oil, and about 27

5 vol. % of DiH₂O (designated herein as GC). In some embodiments, a nanoemulsion comprises about 10 vol. % of glycerol, about 10 vol. % of CPC, about 60 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as GC10). In still another embodiment of the present invention, a nanoemulsion comprises about 10 vol. % of glycerol, about 1 vol. % of CPC, about 1 vol. % of L-ascorbic acid, about 64 vol. % of soybean or oil, and about 24
10 vol. % of DiH₂O (designated herein as GCV_c).

In some embodiments of the present invention, a nanoemulsion comprises about 8 to 10 vol. % of glycerol, about 8 to 10 vol. % of SDS, about 50 to 70 vol. % of oil (e.g., soybean or olive oil), and about 15 to 30 vol. % of aqueous phase (e.g., DiH₂O or PBS). Additionally, in certain of these embodiments, a nanoemulsion further comprise about 1 vol. % of lecithin,
15 and about 1 vol. % of p-Hydroxybenzoic acid methyl ester. Exemplary embodiments of such formulations comprise about 8 vol. % SDS, 8 vol. % of glycerol, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as S8G). A related formulation comprises about 8 vol. % of glycerol, about 8 vol. % of SDS, about 1 vol. % of lecithin, about 1 vol. % of p-Hydroxybenzoic acid methyl ester, about 64 vol. % of soybean oil, and
20 about 18 vol. % of DiH₂O (designated herein as S8GL1B1).

In yet another embodiment of the present invention, a nanoemulsion comprises about 4 vol. % of TWEEN 80, about 4 vol. % of TYLOXAPOL, about 1 vol. % of CPC, about 8 vol. % of ethanol, about 64 vol. % of soybean oil, and about 19 vol. % of DiH₂O (designated herein as W₈₀4Y4EC).

25 In some embodiments of the present invention, a nanoemulsion comprises about 0.01 vol. % of CPC, about 0.08 vol. % of TYLOXAPOL, about 10 vol. % of ethanol, about 70 vol. % of soybean oil, and about 19.91 vol. % of DiH₂O (designated herein as Y.08EC.01).

In yet another embodiment of the present invention, a nanoemulsion comprises about 8 vol. % of sodium lauryl sulfate, and about 8 vol. % of glycerol, about 64 vol. % of soybean
30 oil, and about 20 vol. % of DiH₂O (designated herein as SLS8G).

The specific formulations described above are simply examples to illustrate the variety of nanoemulsions that find use (e.g., to inactivate and/or neutralize a pathogen, and

for generating an immune response in a subject (e.g., for use as a vaccine)) in the present invention. The present invention contemplates that many variations of the above formulations, as well as additional nanoemulsions, find use in the methods of the present invention. Candidate emulsions can be easily tested to determine if they are suitable. First, 5 the desired ingredients are prepared using the methods described herein, to determine if an emulsion can be formed. If an emulsion cannot be formed, the candidate is rejected. For example, a candidate composition made of 4.5% sodium thiosulfate, 0.5% sodium citrate, 10% n-butanol, 64% soybean oil, and 21% DiH₂O does not form an emulsion.

Second, the candidate emulsion should form a stable emulsion. An emulsion is stable 10 if it remains in emulsion form for a sufficient period to allow its intended use (e.g., to generate an immune response in a subject). For example, for emulsions that are to be stored, shipped, etc., it may be desired that the composition remain in emulsion form for months to years. Typical emulsions that are relatively unstable, will lose their form within a day. For example, a candidate composition made of 8% 1-butanol, 5% Tween 10, 1% CPC, 64% 15 soybean oil, and 22% DiH₂O does not form a stable emulsion. Nanoemulsions that have been shown to be stable include, but are not limited to, 8 vol. % of TRITON X-100, about 8 vol. % of TBP, about 64 vol. % of soybean oil, and about 20 vol. % of DiH₂O (designated herein as X8P); 5 vol. % of TWEEN 20, from about 8 vol. % of ethanol, from about 1 vol. % of CPC, about 64 vol. % of oil (e.g., soybean oil), and about 22 vol. % of DiH₂O (designated 20 herein as W₂₀5EC); 0.08% Triton X-100, 0.08% Glycerol, 0.01% Cetylpyridinium Chloride, 99% Butter, and 0.83% diH₂O (designated herein as 1% X8GC Butter); 0.8% Triton X-100, 0.8% Glycerol, 0.1% Cetylpyridinium Chloride, 6.4% Soybean Oil, 1.9% diH₂O, and 90% Butter (designated herein as 10% X8GC Butter); 2% W₂₀5EC, 1% Natrosol 250L NF, and 97% diH₂O (designated herein as 2% W₂₀5EC L GEL); 1% Cetylpyridinium Chloride, 5% 25 Tween 20, 8% Ethanol, 64% 70 Viscosity Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC 70 Mineral Oil); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% 350 Viscosity Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC 350 Mineral Oil). In some embodiments, nanoemulsions of the present invention are stable for over a week, over a month, or over a year.

30 Third, the candidate emulsion should have efficacy for its intended use. For example, a nanoemulsion should inactivate (e.g., kill or inhibit growth of) a pathogen to a desired level (e.g., 1 log, 2 log, 3 log, 4 log, . . . reduction). Using the methods described herein, one is

capable of determining the suitability of a particular candidate emulsion against the desired pathogen. Generally, this involves exposing the pathogen to the emulsion for one or more time periods in a side-by-side experiment with the appropriate control samples (e.g., a negative control such as water) and determining if, and to what degree, the emulsion

5 inactivates (e.g., kills and/or neutralizes) the microorganism. For example, a candidate composition made of 1% ammonium chloride, 5% Tween 20, 8% ethanol, 64% soybean oil, and 22% diH₂O was shown not to be an effective emulsion. The following candidate emulsions were shown to be effective using the methods described herein: 5% Tween 20, 5% Cetylpyridinium Chloride, 10% Glycerol, 60% Soybean Oil, and 20% diH₂O (designated

10 herein as W₂₀5GC5); 1% Cetylpyridinium Chloride, 5% Tween 20, 10% Glycerol, 64% Soybean Oil, and 20% diH₂O (designated herein as W₂₀5GC); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Olive Oil, and 22% diH₂O (designated herein as W₂₀5EC Olive Oil); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Flaxseed Oil, and 22% diH₂O (designated herein as W₂₀5EC Flaxseed Oil); 1% Cetylpyridinium Chloride, 5%

15 Tween 20, 8% Ethanol, 64% Corn Oil, and 22% diH₂O (designated herein as W₂₀5EC Corn Oil); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Coconut Oil, and 22% diH₂O (designated herein as W₂₀5EC Coconut Oil); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Cottonseed Oil, and 22% diH₂O (designated herein as W₂₀5EC Cottonseed Oil); 8% Dextrose, 5% Tween 10, 1% Cetylpyridinium Chloride, 64% Soybean

20 Oil, and 22% diH₂O (designated herein as W₂₀5C Dextrose); 8% PEG 200, 5% Tween 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C PEG 200); 8% Methanol, 5% Tween 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as W₂₀5C Methanol); 8% PEG 1000, 5% Tween 10, 1% Cetylpyridinium Chloride, 64% Soybean Oil, and 22% diH₂O (designated herein as

25 W₂₀5C PEG 1000); 2% W₂₀5EC, 2% Natrosol 250H NF, and 96% diH₂O (designated herein as 2% W₂₀5EC Natrosol 2, also called 2% W₂₀5EC GEL); 2% W₂₀5EC, 1% Natrosol 250H NF, and 97% diH₂O (designated herein as 2% W₂₀5EC Natrosol 1); 2% W₂₀5EC, 3% Natrosol 250H NF, and 95% diH₂O (designated herein as 2% W₂₀5EC Natrosol 3); 2% W₂₀5EC, 0.5% Natrosol 250H NF, and 97.5% diH₂O (designated herein as 2% W₂₀5EC

30 Natrosol 0.5); 2% W₂₀5EC, 2% Methocel A, and 96% diH₂O (designated herein as 2% W₂₀5EC Methocel A); 2% W₂₀5EC, 2% Methocel K, and 96% diH₂O (designated herein as 2% W₂₀5EC Methocel K); 2% Natrosol, 0.1% X8PC, 0.1x PBS, 5 mM L-alanine, 5 mM

Inosine, 10 mM Ammonium Chloride, and diH₂O (designated herein as 0.1% X8PC/GE+2% Natrosol); 2% Natrosol, 0.8% Triton X-100, 0.8% Tributyl Phosphate, 6.4% Soybean Oil, 0.1% Cetylpyridinium Chloride, 0.1x PBS, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, and diH₂O (designated herein as 10% X8PC/GE+2% Natrosol); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Lard, and 22% diH₂O (designated herein as W₂₀5EC Lard); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Ethanol, 64% Mineral Oil, and 22% diH₂O (designated herein as W₂₀5EC Mineral Oil); 0.1% Cetylpyridinium Chloride, 2% Nerolidol, 5% Tween 20, 10% Ethanol, 64% Soybean Oil, and 18.9% diH₂O (designated herein as W₂₀5EC_{0.1}N); 0.1% Cetylpyridinium Chloride, 2% Farnesol, 5% Tween 20, 10% Ethanol, 64% Soybean Oil, and 18.9% diH₂O (designated herein as W₂₀5EC_{0.1}F); 0.1% Cetylpyridinium Chloride, 5% Tween 20, 10% Ethanol, 64% Soybean Oil, and 20.9% diH₂O (designated herein as W₂₀5EC_{0.1}); 10% Cetylpyridinium Chloride, 8% Tributyl Phosphate, 8% Triton X-100, 54% Soybean Oil, and 20% diH₂O (designated herein as X8PC₁₀); 5% Cetylpyridinium Chloride, 8% Triton X-100, 8% Tributyl Phosphate, 59% Soybean Oil, and 20% diH₂O (designated herein as X8PC₅); 0.02% Cetylpyridinium Chloride, 0.1% Tween 20, 10% Ethanol, 70% Soybean Oil, and 19.88% diH₂O (designated herein as W₂₀0.1EC_{0.02}); 1% Cetylpyridinium Chloride, 5% Tween 20, 8% Glycerol, 64% Mobil 1, and 22% diH₂O (designated herein as W₂₀5GC Mobil 1); 7.2% Triton X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 0.1x PBS, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, and 25.87% diH₂O (designated herein as 90% X8PC/GE); 7.2% Triton X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 1% EDTA, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, 0.1x PBS, and diH₂O (designated herein as 90% X8PC/GE EDTA); and 7.2% Triton X-100, 7.2% Tributyl Phosphate, 0.9% Cetylpyridinium Chloride, 57.6% Soybean Oil, 1% Sodium Thiosulfate, 5 mM L-alanine, 5 mM Inosine, 10 mM Ammonium Chloride, 0.1x PBS, and diH₂O (designated herein as 90% X8PC/GE STS).

In preferred embodiments of the present invention, the nanoemulsions are non-toxic (e.g., to humans, plants, or animals), non-irritant (e.g., to humans, plants, or animals), and non-corrosive (e.g., to humans, plants, or animals or the environment), while possessing potency against a broad range of microorganisms including bacteria, fungi, viruses, and spores. While a number of the above described nanoemulsions meet these qualifications, the following description provides a number of preferred non-toxic, non-irritant, non-corrosive,

anti-microbial nanoemulsions of the present invention (hereinafter in this section referred to as "non-toxic nanoemulsions").

In some embodiments the non-toxic nanoemulsions comprise surfactant lipid preparations (SLPs) for use as broad-spectrum antimicrobial agents that are effective against bacteria and their spores, enveloped viruses, and fungi. In preferred embodiments, these SLPs comprises a mixture of oils, detergents, solvents, and cationic halogen-containing compounds in addition to several ions that enhance their biocidal activities. These SLPs are characterized as stable, non-irritant, and non-toxic compounds compared to commercially available bactericidal and sporicidal agents, which are highly irritant and/or toxic.

Ingredients for use in the non-toxic nanoemulsions include, but are not limited to: detergents (e.g., TRITON X-100 (5-15%) or other members of the TRITON family, TWEEN 60 (0.5-2%) or other members of the TWEEN family, or TYLOXAPOL (1-10%)); solvents (e.g., tributyl phosphate (5-15%)); alcohols (e.g., ethanol (5-15%) or glycerol (5-15%)); oils (e.g., soybean oil (40-70%)); cationic halogen-containing compounds (e.g., cetylpyridinium chloride (0.5-2%), cetylpyridinium bromide (0.5-2%)), or cetyldimethylethyl ammonium bromide (0.5-2%)); quaternary ammonium compounds (e.g., benzalkonium chloride (0.5-2%), N-alkyldimethylbenzyl ammonium chloride (0.5-2%)); ions (calcium chloride (1mM-40mM), ammonium chloride (1mM-20mM), sodium chloride (5mM-200mM), sodium phosphate (1mM-20mM)); nucleosides (e.g., inosine (50 μ M-20mM)); and amino acids (e.g., L-alanine (50 μ M-20mM)). Emulsions are prepared, for example, by mixing in a high shear mixer for 3-10 minutes. The emulsions may or may not be heated before mixing at 82°C for 1 hour.

Quaternary ammonium compounds for use in the present include, but are not limited to, N-alkyldimethyl benzyl ammonium saccharinate; 1,3,5-Triazine-1,3,5(2H,4H,6H)-triethanol; 1-Decanaminium, N-decyl-N, N-dimethyl-, chloride (or) Didecyl dimethyl ammonium chloride; 2-(2-(p-(Diisobutyl)cresosy)ethoxy)ethyl dimethyl benzyl ammonium chloride; 2-(2-(p-(Diisobutyl)phenoxy)ethoxy)ethyl dimethyl benzyl ammonium chloride; alkyl 1 or 3 benzyl-1-(2-hydroxyethyl)-2-imidazolinium chloride; alkyl bis(2-hydroxyethyl) benzyl ammonium chloride; alkyl demethyl benzyl ammonium chloride; alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride (100% C12); alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride (50% C14, 40% C12, 10% C16); alkyl dimethyl 3,4-dichlorobenzyl ammonium chloride (55% C14, 23% C12, 20% C16); alkyl dimethyl benzyl ammonium

chloride; alkyl dimethyl benzyl ammonium chloride (100% C14); alkyl dimethyl benzyl ammonium chloride (100% C16); alkyl dimethyl benzyl ammonium chloride (41% C14, 28% C12); alkyl dimethyl benzyl ammonium chloride (47% C12, 18% C14); alkyl dimethyl benzyl ammonium chloride (55% C16, 20% C14); alkyl dimethyl benzyl ammonium chloride (58% C14, 28% C16); alkyl dimethyl benzyl ammonium chloride (60% C14, 25% C12); alkyl dimethyl benzyl ammonium chloride (61% C11, 23% C14); alkyl dimethyl benzyl ammonium chloride (61% C12, 23% C14); alkyl dimethyl benzyl ammonium chloride (65% C12, 25% C14); alkyl dimethyl benzyl ammonium chloride (67% C12, 24% C14); alkyl dimethyl benzyl ammonium chloride (67% C12, 25% C14); alkyl dimethyl benzyl ammonium chloride (90% C14, 5% C12); alkyl dimethyl benzyl ammonium chloride (93% C14, 4% C12); alkyl dimethyl benzyl ammonium chloride (95% C16, 5% C18); alkyl dimethyl benzyl ammonium chloride (and) didecyl dimethyl ammonium chloride; alkyl dimethyl benzyl ammonium chloride (as in fatty acids); alkyl dimethyl benzyl ammonium chloride (C12-C16); alkyl dimethyl benzyl ammonium chloride (C12-C18); alkyl dimethyl benzyl and dialkyl dimethyl ammonium chloride; alkyl dimethyl dimethylbenzyl ammonium chloride; alkyl dimethyl ethyl ammonium bromide (90% C14, 5% C16, 5% C12); alkyl dimethyl ethyl ammonium bromide (mixed alkyl and alkenyl groups as in the fatty acids of soybean oil); alkyl dimethyl ethylbenzyl ammonium chloride; alkyl dimethyl ethylbenzyl ammonium chloride (60% C14); alkyl dimethyl isopropylbenzyl ammonium chloride (50% C12, 30% C14, 17% C16, 3% C18); alkyl trimethyl ammonium chloride (58% C18, 40% C16, 1% C14, 1% C12); alkyl trimethyl ammonium chloride (90% C18, 10% C16); alkyl dimethyl(ethylbenzyl) ammonium chloride (C12-18); Di-(C8-10)-alkyl dimethyl ammonium chlorides; dialkyl dimethyl ammonium chloride; dialkyl dimethyl ammonium chloride; dialkyl methyl benzyl ammonium chloride; didecyl dimethyl ammonium chloride; diisodecyl dimethyl ammonium chloride; dioctyl dimethyl ammonium chloride; dodecyl bis (2-hydroxyethyl) octyl hydrogen ammonium chloride; dodecyl dimethyl benzyl ammonium chloride; dodecylcarbonyl methyl dinethyl benzyl ammonium chloride; heptadecyl hydroxyethylimidazolinium chloride; hexahydro-1,3,5-tris(2-hydroxyethyl)-s-triazine; myristalkonium chloride (and) Quat RNIUM 14; N,N-Dimethyl-2-hydroxypropylammonium chloride polymer; n-alkyl dimethyl benzyl ammonium chloride; n-alkyl dimethyl ethylbenzyl ammonium chloride; n-tetradecyl dimethyl benzyl ammonium chloride monohydrate; octyl decyl dimethyl ammonium chloride; octyl dodecyl

dimethyl ammonium chloride; octyphenoxyethoxyethyl dimethyl benzyl ammonium chloride; oxydiethylenebis (alkyl dimethyl ammonium chloride); quaternary ammonium compounds, dicoco alkyldimethyl, chloride; trimethoxysilyl propyl dimethyl octadecyl ammonium chloride; trimethoxysilyl quats, trimethyl dodecylbenzyl ammonium chloride; 5 n-dodecyl dimethyl ethylbenzyl ammonium chloride; n-hexadecyl dimethyl benzyl ammonium chloride; n-tetradecyl dimethyl benzyl ammonium chloride; n-tetradecyl dimethyl ethylbenzyl ammonium chloride; and n-octadecyl dimethyl benzyl ammonium chloride.

In general, the preferred non-toxic nanoemulsions are characterized by the following: they are approximately 200-800 nm in diameter, although both larger and smaller diameter 10 nanoemulsions are contemplated; the charge depends on the ingredients; they are stable for relatively long periods of time (e.g., up to two years), with preservation of their biocidal activity; they are non-irritant and non-toxic compared to their individual components due, at least in part, to their oil contents that markedly reduce the toxicity of the detergents and the solvents; they are effective at concentrations as low as 0.1%; they have antimicrobial activity 15 against most vegetative bacteria (including Gram-positive and Gram-negative organisms), fungi, and enveloped and nonenveloped viruses in 15 minutes (e.g., 99.99% killing); and they have sporicidal activity in 1-4 hours (e.g., 99.99% killing) when produced with germination enhancers.

20 Therapeutics and Prophylactics

Furthermore, in preferred embodiments, a composition of the present invention induces (e.g., when administered to a subject) both systemic and mucosal immunity. Thus, in some preferred embodiments, administration of a composition of the present invention to a subject results in protection against an exposure to one or a plurality of pathogens.

25 In some embodiments, the present invention provides a composition comprising a nanoemulsion and one or a plurality of immunogens to serve as a mucosal vaccine. In some embodiments, this material can easily be produced. The ability to produce this formulation rapidly and administer it via mucosal (e.g., nasal) instillation provides a vaccine that can be used in large-scale administrations (e.g., to a population of a town, village, city, state or 30 country).

In some preferred embodiments, the present invention provides a composition for generating an immune response comprising a nanoemulsion and one or a plurality of

immunogens (e.g., a purified, isolated or synthetic protein or derivative, variant, or analogue thereof from one or more serotypes of HBV).

The present invention is not limited by the plurality of immunogens utilized. For example, in some embodiments, the plurality of immunogens comprises immunogenic
5 protein antigens from HIV (e.g., gp120) to serve as a mucosal vaccine. This material can easily be produced with NE and HIV protein (e.g., viral-derived gp120, live-virus-vector-derived gp120 and gp160, recombinant mammalian gp120, recombinant denatured antigens, small peptide segments of gp120 and gp41, V3 loop peptides, and induces both mucosal and systemic immunity. The ability to produce this formulation rapidly and administer it via
10 mucosal (e.g., nasal or vaginal) instillation provides a vaccine that can be used in large-scale administrations (e.g., to a population of a town, village, city, state or country).

In some embodiments, one or a plurality of immunogens comprises one or more strains of orthopox virus (e.g., in a composition comprising a NE and immunogen (e.g., orthopox virus inactivated by the nanoemulsion). Each orthopox virus family member alone,
15 or in combination with another family member, may be used to generate a composition comprising a NE and an immunogen (e.g., used to generate an immune response) of the present invention. Orthopox virus family member include, but are not limited to, variola virus, vaccinia virus, cowpox, monkeypox, gergilpox, camelpox, and others. The present invention is not limited by the strain of vaccinia virus used. Indeed, a variety of vaccinia
20 virus strains are contemplated to be useful in the present invention including, but not limited to, classical strains of vaccinia virus (e.g., EM-63, Lister, New York City Board of Health, Elestree, and Temple of Heaven strains), attenuated strains (e.g., Ankara), non-replicating strains, modified strains (e.g., genetically or mechanically modified strains (e.g., to become more or less virulent)), Copenhagen strain, modified vaccinia Ankara, New York vaccinia
25 virus, Vaccinia Virus_{WR} and Vaccinia Virus_{WR-Luc}, or other serially diluted strain of vaccinia virus. A composition comprising a NE and one or a plurality of immunogens may comprise one or more strains of vaccinia virus and/or other type of orthopox virus. Additionally, a composition comprising a NE and one or a plurality of immunogens may comprise one or more strains of vaccinia virus, and, in addition, one or more strains of a non-vaccinia virus
30 immunogen or immunogenic epitope thereof (e.g., a bacteria (e.g., *B. anthracis*) or immunogenic epitope thereof (e.g., recombinant protective antigen) or a virus (e.g., West Nile virus, Avian Influenza virus, Ebola virus, HSV, HPV, HCV, HIV, etc.) or an

immunogenic epitope thereof (e.g., gp120)).

In some embodiments, the immunogen may comprise one or more antigens derived from a pathogen (e.g., orthopox virus). For example, in some embodiments, the immunogen is a purified, recombinant, synthetic, or otherwise isolated protein (e.g., added to the NE to
5 generate an immunogenic composition). Similarly, the immunogenic protein may be a derivative, analogue or otherwise modified (e.g., PEGylated) form of a protein from a pathogen.

In some embodiments, one or a plurality of immunogens comprises one or more strains of *Bacillus* or immunogenic proteins derived therefrom. For example, 89 different
10 strains of *B. anthracis* have been identified, ranging from virulent Ames and *Vollum* strains with biological warfare and bioterrorism applications to benign *Sterne* strain used for inoculations (See, e.g., Easterday et al., J Clin Microbiol. 2005 43(4):1995-7). The strains differ in presence and activity of various genes, determining their virulence and production of antigens and toxins. Any one of these or yet to be identified or generated strains may be used
15 in an immunogenic composition comprising a NE of the present invention.

In some embodiments, the immunogen may comprise one or more antigens derived from a pathogen (e.g., *B. anthracis*). For example, in some embodiments, the immunogen is a purified, recombinant, synthetic, or otherwise isolated protein (e.g., added to the NE to generate an immunogenic composition). Similarly, the immunogenic protein may be a
20 derivative, variant, analogue or otherwise modified form of a protein from a pathogen. The present invention is not limited by the type of protein (e.g., derived from bacteria of the genus *Bacillus*) used for generation of an immunogenic composition of the present invention. Indeed, a variety of immunogenic proteins may be used including, but not limited to, protective antigen (PA), lethal factor (LF), edema factor (EF), PA degradation products (See,
25 e.g., Farchaus, J., et al., Applied & Environmental Microbiol., 64(3):982-991 (1998)), as well as analogues, derivatives and modified forms thereof.

For example, *Bacillus* proteins of the present invention may be used in their native conformation, or more preferably, may be modified for vaccine use. These modifications may either be required for technical reasons relating to the method of purification, or they
30 may be used to biologically inactivate one or several functional properties of the *Bacillus* proteins (e.g., that would otherwise be toxic). Thus the invention encompasses derivatives of *Bacillus* proteins that may be, for example, mutated proteins (e.g., that has undergone

deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional method).

Antigens (e.g., protein antigens (e.g., recombinant protective antigen (PA))) of the present invention may be modified by chemical methods during a purification process to
5 render the proteins stable and monomeric. One method to prevent oxidative aggregation of a protein is the use of chemical modifications of the protein's thiol groups. In a first step the disulphide bridges are reduced by treatment with a reducing agent such as DTT, β -mercaptoethanol, or glutathione. In a second step the resulting thiols are blocked by reaction
10 with an alkylating agent (e.g., the protein can be carboxyamided/carbamidomethylated using iodoacetamide).

Each *Bacillus* family member alone, or in combination with another family member, may be used to generate a composition comprising a NE and plurality of immunogens (e.g., used to generate an immune response) of the present invention. A composition comprising a NE and plurality of immunogens may comprise one or more strains of *B. anthracis*.
15 Additionally, a composition comprising a NE and plurality of immunogens may comprise one or more strains of *B. anthracis*, and, in addition, one or more strains of a non-*B. anthracis* immunogen (e.g., a virus such as West Nile virus, Avian Influenza virus, Ebola virus, HSV, HPV, HCV, HIV, etc. or an immunogenic epitope thereof (e.g., gp120)).

The present invention is not limited by the type (e.g., serotype, group, or clade) of
20 HIV used or immunogenic protein derived therefrom. For example, there are currently two types of HIV: HIV-1 and HIV-2. Both types are transmitted by sexual contact, through blood, and from mother to child, and they appear to cause clinically indistinguishable AIDS. However, it seems that HIV-2 is less easily transmitted, and the period between initial infection and illness is longer in the case of HIV-2. Worldwide, the predominant virus is
25 HIV-1, and generally when people refer to HIV without specifying the type of virus they will be referring to HIV-1. The relatively uncommon HIV-2 type is concentrated in West Africa and is rarely found elsewhere.

Different levels of HIV classification exist. Each type is divided into groups, and each group is divided into subtypes and circulating recombinant forms (CRFs). The strains
30 of HIV-1 can be classified into three groups : the "major" group M, the "outlier" group O and the "new" group N.

Within group M there are known to be at least nine genetically distinct subtypes (or

clades) of HIV-1. These are subtypes A, B, C, D, F, G, H, J and K.

Any one of these or yet to be identified or generated serotypes, groups, or clades may be used in an immunogenic composition comprising a NE of the present invention.

In some embodiments, one or a plurality of immunogens may comprise one or more
5 antigens derived from a pathogen (e.g., HIV). For example, in some embodiments, the immunogen is a purified, recombinant, synthetic, or otherwise isolated protein (e.g., added to the NE to generate an immunogenic composition). Similarly, the immunogenic protein may be a derivative, analogue or otherwise modified form of a protein from a pathogen. The present invention is not limited by the type of protein (e.g., derived from HIV) used for
10 generation of an immunogenic composition of the present invention. Indeed, a variety of immunogenic proteins may be used including, but not limited to, gp160, gp120, gp41, Tat, and Nef; as well as analogues, derivatives and modified forms thereof.

For example, HIV proteins of the present invention may be used in their native conformation, or more preferably, may be modified for vaccine use. These modifications may
15 either be required for technical reasons relating to the method of purification, or they may be used to biologically inactivate one or several functional properties of HIV protein. Thus the invention encompasses derivatives of HIV proteins which may be, for example mutated proteins (e.g., that has undergone deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional
20 method.

For example, a HIV protein may be mutated so that it is biologically inactive while maintaining its immunogenic epitopes (See, e.g., Clements, Virology 235: 48-64, 1997).

Additionally, HIV proteins of the present invention may be modified by chemical methods during the purification process to render the proteins stable and monomeric. One
25 method to prevent oxidative aggregation of a HIV protein is the use of chemical modifications of the protein's thiol groups. In a first step the disulphide bridges are reduced by treatment with a reducing agent such as DTT, β -mercaptoethanol, or glutathione. In a second step the resulting thiols are blocked by reaction with an alkylating agent (e.g., the protein can be carboxyamidated/carbamidomethylated using iodoacetamide).

30 Each HIV serotype, group or clade alone, or in combination with another family member, may be used to generate a composition comprising a NE and an immunogen (e.g., used to generate an immune response) of the present invention. A composition comprising a

NE and immunogen may comprise one or more serotypes, groups or clades of HIV.

Additionally, a composition comprising a NE and immunogen may comprise one or more serotypes, groups or clades of HIV, and, in addition, one or more strains of a non-HIV immunogen (e.g., a virus such as West Nile virus, Avian Influenza virus, Ebola virus, HSV, HPV, HCV, , etc. or an immunogenic epitope thereof).

When administered to a subject, a composition of the present invention stimulates an immune response against one or a plurality of immunogens within the subject. Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, generation of an immune response (e.g., resulting from administration of a composition comprising a nanoemulsion and plurality of immunogens) provides total or partial immunity to the subject (e.g., from signs, symptoms or conditions of a disease (e.g., to a plurality of pathogens from which the immunogens are derived)). Without being bound to any specific theory, protection and/or immunity from disease (e.g., the ability of a subject's immune system to prevent or attenuate (e.g., suppress) a sign, symptom or condition of disease) after exposure to an immunogenic composition of the present invention is due to adaptive (e.g., acquired) immune responses (e.g., immune responses mediated by B and T cells following exposure to a NE comprising one or a plurality of immunogens of the present invention (e.g., immune responses that exhibit increased specificity and reactivity towards one or a plurality of pathogens from which the immunogens are derived)). Thus, in some embodiments, the compositions and methods of the present invention are used prophylactically or therapeutically to prevent or attenuate a sign, symptom or condition associated with one or a plurality of pathogens.

In some embodiments, a composition comprising a nanoemulsion and one or a plurality of immunogens is administered alone. In some embodiments, a composition comprising a nanoemulsion and one or a plurality of immunogens comprises one or more other agents (e.g., a pharmaceutically acceptable carrier, adjuvant, excipient, and the like). In some embodiments, a composition for stimulating an immune response of the present invention is administered in a manner to induce a humoral immune response. In some embodiments, a composition for stimulating an immune response of the present invention is administered in a manner to induce a cellular (e.g., cytotoxic T lymphocyte) immune response, rather than a humoral response. In some embodiments, a composition comprising a

NE and an immunogen of the present invention induces both a cellular and humoral immune response.

5 In some embodiments, the immunogen may comprise one or more antigens derived from one or a plurality of immunogens. For example, in some embodiments, the immunogen is a purified, recombinant, synthetic, or otherwise isolated protein (e.g., added to a nanoemulsion to generate an immunogenic composition). Similarly, the immunogenic protein may be a derivative, analogue or otherwise modified (e.g., PEGylated) form of a protein.

10 The present invention is not limited by the particular formulation of a composition comprising a nanoemulsion and a plurality of immunogens of the present invention. Indeed, a composition comprising a nanoemulsion and one or a plurality of immunogens of the present invention may comprise one or more different agents in addition to the nanoemulsion and plurality of immunogens. These agents or cofactors include, but are not limited to, adjuvants, surfactants, additives, buffers, solubilizers, chelators, oils, salts, therapeutic agents, 15 drugs, bioactive agents, antibacterials, and antimicrobial agents (e.g., antibiotics, antivirals, etc.). In some embodiments, a composition comprising a nanoemulsion and one or a plurality of immunogens of the present invention comprises an agent and/or co-factor that enhance the ability of the immunogen to induce an immune response (e.g., an adjuvant). In some preferred embodiments, the presence of one or more co-factors or agents reduces the amount of immunogen required for induction of an immune response (e.g., a protective immune 20 response (e.g., protective immunization)). In some embodiments, the presence of one or more co-factors or agents can be used to skew the immune response towards a cellular (e.g., T cell mediated) or humoral (e.g., antibody mediated) immune response. The present invention is not limited by the type of co-factor or agent used in a therapeutic agent of the present 25 invention.

Adjuvants are described in general in Vaccine Design--the Subunit and Adjuvant Approach, edited by Powell and Newman, Plenum Press, New York, 1995. The present invention is not limited by the type of adjuvant utilized (e.g., for use in a composition (e.g., pharmaceutical composition) comprising a NE and immunogen). For example, in some 30 embodiments, suitable adjuvants include an aluminium salt such as aluminium hydroxide gel (alum) or aluminium phosphate. In some embodiments, an adjuvant may be a salt of calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars,

cationically or anionically derivatised polysaccharides, or polyphosphazenes.

In general, an immune response is generated to an antigen through the interaction of the antigen with the cells of the immune system. Immune responses may be broadly categorized into two categories: humoral and cell mediated immune responses (e.g.,
5 traditionally characterized by antibody and cellular effector mechanisms of protection, respectively). These categories of response have been termed Th1-type responses (cell-mediated response), and Th2-type immune responses (humoral response).

Stimulation of an immune response can result from a direct or indirect response of a cell or component of the immune system to an intervention (e.g., exposure to an immunogen).

10 Immune responses can be measured in many ways including activation, proliferation or differentiation of cells of the immune system (e.g., B cells, T cells, dendritic cells, APCs, macrophages, NK cells, NKT cells etc.); up-regulated or down-regulated expression of markers and cytokines; stimulation of IgA, IgM, or IgG titer; splenomegaly (including increased spleen cellularity); hyperplasia and mixed cellular infiltrates in various organs.
15 Other responses, cells, and components of the immune system that can be assessed with respect to immune stimulation are known in the art.

Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, compositions and methods of the present invention induce expression
20 and secretion of cytokines (e.g., by macrophages, dendritic cells and CD4+ T cells).

Modulation of expression of a particular cytokine can occur locally or systemically. It is known that cytokine profiles can determine T cell regulatory and effector functions in immune responses. In some embodiments, Th1-type cytokines can be induced, and thus, the immunostimulatory compositions of the present invention can promote a Th1 type antigen-
25 specific immune response including cytotoxic T-cells (e.g., thereby avoiding unwanted Th2 type immune responses (e.g., generation of Th2 type cytokines (e.g., IL-13) involved in enhancing the severity of disease (e.g., IL-13 induction of mucus formation))).

Cytokines play a role in directing the T cell response. Helper (CD4+) T cells orchestrate the immune response of mammals through production of soluble factors that act
30 on other immune system cells, including B and other T cells. Most mature CD4+T helper cells express one of two cytokine profiles: Th1 or Th2. Th1-type CD4+ T cells secrete IL-2, IL-3, IFN- γ , GM-CSF and high levels of TNF- α . Th2 cells express IL-3, IL-4, IL-5, IL-6,

IL-9, IL-10, IL-13, GM-CSF and low levels of TNF- α . Th1 type cytokines promote both cell-mediated immunity, and humoral immunity that is characterized by immunoglobulin class switching to IgG2a in mice and IgG1 in humans. Th1 responses may also be associated with delayed-type hypersensitivity and autoimmune disease. Th2 type cytokines induce primarily humoral immunity and induce class switching to IgG1 and IgE. The antibody isotypes associated with Th1 responses generally have neutralizing and opsonizing capabilities whereas those associated with Th2 responses are associated more with allergic responses.

Several factors have been shown to influence skewing of an immune response towards either a Th1 or Th2 type response. The best characterized regulators are cytokines. IL-12 and IFN- γ are positive Th1 and negative Th2 regulators. IL-12 promotes IFN- γ production, and IFN- γ provides positive feedback for IL-12. IL-4 and IL-10 appear important for the establishment of the Th2 cytokine profile and to down-regulate Th1 cytokine production.

Thus, in preferred embodiments, the present invention provides a method of stimulating a Th1-type immune response in a subject comprising administering to a subject a composition comprising a NE and one or a plurality of immunogens. However, in other embodiments, the present invention provides a method of stimulating a Th2-type immune response in a subject (e.g., if balancing of a T cell mediated response is desired) comprising administering to a subject a composition comprising a NE and one or a plurality of immunogens. In further preferred embodiments, adjuvants can be used (e.g., can be co-administered with a composition of the present invention) to skew an immune response toward either a Th1 or Th2 type immune response. For example, adjuvants that induce Th2 or weak Th1 responses include, but are not limited to, alum, saponins, and SB-As4. Adjuvants that induce Th1 responses include but are not limited to MPL, MDP, ISCOMS, IL-12, IFN- γ , and SB-AS2.

Several other types of Th1-type immunogens can be used (e.g., as an adjuvant) in compositions and methods of the present invention. These include, but are not limited to, the following. In some embodiments, monophosphoryl lipid A (e.g., in particular 3-de-O-acylated monophosphoryl lipid A (3D-MPL)), is used. 3D-MPL is a well known adjuvant manufactured by Ribi Immunochem, Montana. Chemically it is often supplied as a mixture of 3-de-O-acylated monophosphoryl lipid A with either 4, 5, or 6 acylated chains. In some

embodiments, diphosphoryl lipid A, and 3-O-deacylated variants thereof are used. Each of these immunogens can be purified and prepared by methods described in GB 2122204B.

Other purified and synthetic

lipopolysaccharides have been described (See, e.g., U.S. Pat. No. 6,005,099 and EP 0 729 473; Hilgers et al., 1986, *Int.Arch.Allergy.Immunol.*, 79(4):392-6; Hilgers et al., 1987, *Immunology*, 60(1):141-6; and EP 0 549 074).

In some embodiments, 3D-MPL is used in the form of a particulate formulation (e.g., having a small particle size less than 0.2 μm in diameter, described in EP 0 689 454).

In some embodiments, saponins are used as an immunogen (e.g., Th1-type adjuvant) in a composition of the present invention. Saponins are well known adjuvants (See, e.g., Lacaille-Dubois and Wagner (1996) *Phytomedicine* vol 2 pp 363-386). Examples of saponins include Quil A (derived from the bark of the South American tree *Quillaja Saponaria* Molina), and fractions thereof (See, e.g., U.S. Pat. No. 5,057,540; Kensil, *Crit Rev Ther Drug Carrier Syst*, 1996, 12 (1-2):1-55; and EP 0 362 279).

Also contemplated to be useful in the present invention are the haemolytic saponins QS7, QS17, and QS21 (HPLC purified fractions of Quil A; See, e.g., Kensil et al. (1991). *J. Immunology* 146,431-437, U.S. Pat. No. 5,057,540; WO 96/33739; WO 96/11711 and EP 0 362 279).

Also contemplated to be useful are combinations of QS21 and polysorbate or cyclodextrin (See, e.g., WO 99/10008).

In some embodiments, an immunogenic oligonucleotide containing unmethylated CpG dinucleotides ("CpG") is used as an adjuvant in the present invention. CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. CpG is known in the art as being an adjuvant when administered by both systemic and mucosal routes (See, e.g., WO 96/02555, EP 468520, Davis et al., *J.Immunol*, 1998, 160(2):870-876; McCluskie and Davis, *J.Immunol.*, 1998, 161(9):4463-6; and U.S. Pat. App. No. 20050238660).

For example, in some embodiments, the immunostimulatory sequence is Purine-Purine-C-G-pyrimidine-pyrimidine; wherein the CG motif is not methylated.

Although an understanding of the mechanism is not necessary to practice the present

invention and the present invention is not limited to any particular mechanism of action, in some embodiments, the presence of one or more CpG oligonucleotides activate various immune subsets including natural killer cells (which produce IFN- γ) and macrophages. In some embodiments, CpG oligonucleotides are formulated into a composition of the present invention for inducing an immune response. In some embodiments, a free solution of CpG is co-administered together with an antigen (e.g., present within a NE solution (See, e.g., WO 96/02555)).

In some embodiments, a CpG oligonucleotide is covalently conjugated to an antigen (See, e.g., WO 98/16247),

or formulated with a carrier such as aluminium hydroxide (See, e.g., Brazolot-Millan et al., Proc.Natl.AcadSci., USA, 1998, 95(26), 15553-8).

In some embodiments, adjuvants such as Complete Freund's Adjuvant and Incomplete Freund's Adjuvant, cytokines (e.g., interleukins (e.g., IL-2, IFN- γ , IL-4, etc.), macrophage colony stimulating factor, tumor necrosis factor, etc.), detoxified mutants of a bacterial ADP-ribosylating toxin such as a cholera toxin (CT), a pertussis toxin (PT), or an E. Coli heat-labile toxin (LT), particularly LT-K63 (where lysine is substituted for the wild-type amino acid at position 63) LT-R72 (where arginine is substituted for the wild-type amino acid at position 72), CT-S109 (where serine is substituted for the wild-type amino acid at position 109), and PT-K9/G129 (where lysine is substituted for the wild-type amino acid at position 9 and glycine substituted at position 129) (See, e.g., WO93/13202 and WO92/19265),

and other immunogenic substances (e.g., that enhance the effectiveness of a composition of the present invention) are used with a composition comprising a NE and immunogen of the present invention.

Additional examples of adjuvants that find use in the present invention include poly(di(carboxylatophenoxy)phosphazene (PCPP polymer; Virus Research Institute, USA); derivatives of lipopolysaccharides such as monophosphoryl lipid A (MPL; Ribi ImmunoChem Research, Inc., Hamilton, Mont.), muramyl dipeptide (MDP; Ribi) and threonyl-muramyl dipeptide (t-MDP; Ribi); OM-174 (a glucosamine disaccharide related to lipid A; OM Pharma SA, Meyrin, Switzerland); and Leishmania elongation factor (a purified Leishmania protein; Corixa Corporation, Seattle, Wash.).

Adjuvants may be added to a composition comprising a NE and one or a plurality of immunogens, or, the adjuvant may be formulated with carriers, for example liposomes, or metallic salts (e.g., aluminium salts (e.g., aluminium hydroxide)) prior to combining with or

co-administration with a composition comprising a NE and an immunogen.

In some embodiments, a composition comprising a NE and an immunogen comprises a single adjuvant. In other embodiments, a composition comprising a NE and an immunogen comprises two or more adjuvants (See, e.g., WO 94/00153; WO 95/17210; WO 96/33739; WO 98/56414; WO 99/12565; WO 99/11241; and WO 94/00153).

In some embodiments, a composition comprising a NE and an immunogen of the present invention comprises one or more mucoadhesives (See, e.g., U.S. Pat. App. No. 20050281843).

The present invention is not limited by the type of mucoadhesive utilized. Indeed, a variety of mucoadhesives are contemplated to be useful in the present invention including, but not limited to, cross-linked derivatives of poly(acrylic acid) (e.g., carbopol and polycarbophil), polyvinyl alcohol, polyvinyl pyrrolidone, polysaccharides (e.g., alginate and chitosan), hydroxypropyl methylcellulose, lectins, fimbrial proteins, and carboxymethylcellulose. Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, use of a mucoadhesive (e.g., in a composition comprising a NE and immunogen) enhances induction of an immune response in a subject (e.g., administered a composition of the present invention) due to an increase in duration and/or amount of exposure to an immunogen that a subject experiences when a mucoadhesive is used compared to the duration and/or amount of exposure to an immunogen in the absence of using the mucoadhesive.

In some embodiments, a composition of the present invention may comprise sterile aqueous preparations. Acceptable vehicles and solvents include, but are not limited to, water, Ringer's solution, phosphate buffered saline and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed mineral or non-mineral oil may be employed including synthetic mono-ordi-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Carrier formulations suitable for mucosal, subcutaneous, intramuscular, intraperitoneal, intravenous, or administration via other routes may be found in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa.

A composition comprising a NE and an immunogen of the present invention can be used therapeutically (e.g., to enhance an immune response) or as a prophylactic (e.g., for

immunization (e.g., to prevent signs or symptoms of disease)). A composition comprising a NE and an immunogen of the present invention can be administered to a subject via a number of different delivery routes and methods.

For example, the compositions of the present invention can be administered to a
 5 subject (e.g., mucosally (e.g., nasal mucosa, vaginal mucosa, etc.)) by multiple methods, including, but not limited to: being suspended in a solution and applied to a surface; being suspended in a solution and sprayed onto a surface using a spray applicator; being mixed with a mucoadhesive and applied (e.g., sprayed or wiped) onto a surface (e.g., mucosal surface);
 10 being placed on or impregnated onto a nasal and/or vaginal applicator and applied; being applied by a controlled-release mechanism; being applied as a liposome; or being applied on a polymer.

In some preferred embodiments, compositions of the present invention are administered mucosally (e.g., using standard techniques; See, e.g., Remington: The Science and Practice of Pharmacy, Mack Publishing Company, Easton, Pa., 19th edition, 1995 (e.g.,
 15 for mucosal delivery techniques, including intranasal, pulmonary, vaginal and rectal techniques), as well as European Publication No. 517,565 and Illum et al., J. Controlled Rel., 1994, 29:133-141 (e.g., for techniques of intranasal administration)).

Alternatively, the compositions of the present invention may be administered dermally or transdermally, using standard techniques (See,
 20 e.g., Remington: The Science and Practice of Pharmacy, Mack Publishing Company, Easton, Pa., 19th edition, 1995). The present invention is not limited by the route of administration.

Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, mucosal vaccination is the preferred route of administration as it has
 25 been shown that mucosal administration of antigens has a greater efficacy of inducing protective immune responses at mucosal surfaces (e.g., mucosal immunity), the route of entry of many pathogens. In addition, mucosal vaccination, such as intranasal vaccination, may induce mucosal immunity not only in the nasal mucosa, but also in distant mucosal sites such as the genital mucosa (See, e.g., Mestecky, Journal of Clinical Immunology, 7:265-276,
 30 1987). More advantageously, in further preferred embodiments, in addition to inducing mucosal immune responses, mucosal vaccination also induces systemic immunity. In some embodiments, non-parenteral administration (e.g., mucosal administration of vaccines)

provides an efficient and convenient way to boost systemic immunity (e.g., induced by parenteral or mucosal vaccination (e.g., in cases where multiple boosts are used to sustain a vigorous systemic immunity)).

In some embodiments, a composition comprising a NE and an immunogen of the present invention may be used to protect or treat a subject susceptible to, or suffering from, disease by means of administering a composition of the present invention via a mucosal route (e.g., an oral/alimentary or nasal route). Alternative mucosal routes include intravaginal and intra-rectal routes. In preferred embodiments of the present invention, a nasal route of administration is used, termed "intranasal administration" or "intranasal vaccination" herein.

Methods of intranasal vaccination are well known in the art, including the administration of a droplet or spray form of the vaccine into the nasopharynx of a subject to be immunized. In some embodiments, a nebulized or aerosolized composition comprising a NE and immunogen is provided. Enteric formulations such as gastro resistant capsules for oral administration, suppositories for rectal or vaginal administration also form part of this invention. Compositions of the present invention may also be administered via the oral route. Under these circumstances, a composition comprising a NE and an immunogen may comprise a pharmaceutically acceptable excipient and/or include alkaline buffers, or enteric capsules. Formulations for nasal delivery may include those with dextran or cyclodextran and saponin as an adjuvant.

Compositions of the present invention may also be administered via a vaginal route. In such cases, a composition comprising a NE and an immunogen may comprise pharmaceutically acceptable excipients and/or emulsifiers, polymers (e.g., CARBOPOL), and other known stabilizers of vaginal creams and suppositories. In some embodiments, compositions of the present invention are administered via a rectal route. In such cases, a composition comprising a NE and an immunogen may comprise excipients and/or waxes and polymers known in the art for forming rectal suppositories.

In some embodiments, the same route of administration (e.g., mucosal administration) is chosen for both a priming and boosting vaccination. In some embodiments, multiple routes of administration are utilized (e.g., at the same time, or, alternatively, sequentially) in order to stimulate an immune response (e.g., using a composition comprising a NE and immunogen of the present invention).

For example, in some embodiments, a composition comprising a NE and an

immunogen is administered to a mucosal surface of a subject in either a priming or boosting vaccination regime. Alternatively, in some embodiments, a composition comprising a NE and an immunogen is administered systemically in either a priming or boosting vaccination regime. In some embodiments, a composition comprising a NE and an immunogen is administered to a subject in a priming vaccination regimen via mucosal administration and a boosting regimen via systemic administration. In some embodiments, a composition comprising a NE and an immunogen is administered to a subject in a priming vaccination regimen via systemic administration and a boosting regimen via mucosal administration. Examples of systemic routes of administration include, but are not limited to, a parenteral, intramuscular, intradermal, transdermal, subcutaneous, intraperitoneal or intravenous administration. A composition comprising a NE and an immunogen may be used for both prophylactic and therapeutic purposes.

In some embodiments, compositions of the present invention are administered by pulmonary delivery. For example, a composition of the present invention can be delivered to the lungs of a subject (e.g., a human) via inhalation (e.g., thereby traversing across the lung epithelial lining to the blood stream (See, e.g., Adjei, et al. *Pharmaceutical Research* 1990; 7:565-569; Adjei, et al. *Int. J. Pharmaceutics* 1990; 63:135-144; Braquet, et al. *J. Cardiovascular Pharmacology* 1989 143-146; Hubbard, et al. (1989) *Annals of Internal Medicine*, Vol. III, pp. 206-212; Smith, et al. *J. Clin. Invest.* 1989;84:1145-1146; Oswein, et al. "Aerosolization of Proteins", 1990; Proceedings of Symposium on Respiratory Drug Delivery II Keystone, Colorado; Debs, et al. *J. Immunol.* 1988; 140:3482-3488; and U.S. Pat. No. 5,284,656 to Platz, et al).

A method and composition for pulmonary delivery of drugs for systemic effect is described in U.S. Pat. No. 5,451,569 to Wong, et al., hereby incorporated by reference; See also U.S. Pat. No. 6,651,655 to Licalsi et al.).

Further contemplated for use in the practice of this invention are a wide range of mechanical devices designed for pulmonary and/or nasal mucosal delivery of pharmaceutical agents including, but not limited to, nebulizers, metered dose inhalers, and powder inhalers, all of which are familiar to those skilled in the art. Some specific examples of commercially available devices suitable for the practice of this invention are the Ultravent nebulizer (Mallinckrodt Inc., St. Louis, Mo.); the Acorn II nebulizer (Marquest Medical Products,

Englewood, Colo.); the Ventolin metered dose inhaler (Glaxo Inc., Research Triangle Park, N.C.); and the Spinhaler powder inhaler (Fisons Corp., Bedford, Mass.). All such devices require the use of formulations suitable for dispensing of the therapeutic agent. Typically, each formulation is specific to the type of device employed and may involve the use of an
 5 appropriate propellant material, in addition to the usual diluents, adjuvants, surfactants, carriers and/or other agents useful in therapy. Also, the use of liposomes, microcapsules or microspheres, inclusion complexes, or other types of carriers is contemplated.

Thus, in some embodiments, a composition comprising a NE and an immunogen of the present invention may be used to protect and/or treat a subject susceptible to, or suffering
 10 from, a disease by means of administering a compositions comprising a NE and an immunogen by mucosal, intramuscular, intraperitoneal, intradermal, transdermal, pulmonary, intravenous, subcutaneous or other route of administration described herein. Methods of systemic administration of the vaccine preparations may include conventional syringes and needles, or devices designed for ballistic delivery of solid vaccines (See, e.g., WO 99/27961),
 15 or needleless pressure liquid jet device (See, e.g., U.S. Pat. No. 4,596,556; U.S. Pat. No. 5,993,412), or transdermal patches (See, e.g., WO 97/48440; WO 98/28037).

The present invention may also be used to enhance the immunogenicity of antigens applied to the skin (transdermal or transcutaneous delivery, See,
 20 e.g., WO 98/20734 ; WO 98/28037).

Thus, in some embodiments, the present invention provides a delivery device for systemic administration, pre-filled with the vaccine composition of the present invention.

The present invention is not limited by the type of subject administered (e.g., in order to stimulate an immune response (e.g., in order to generate protective immunity (e.g.,
 25 mucosal and/or systemic immunity))) a composition of the present invention. Indeed, a wide variety of subjects are contemplated to be benefited from administration of a composition of the present invention. In preferred embodiments, the subject is a human. In some embodiments, human subjects are of any age (e.g., adults, children, infants, etc.) that have been or are likely to become exposed to a microorganism (e.g., HBV). In some
 30 embodiments, the human subjects are subjects that are more likely to receive a direct exposure to pathogenic microorganisms or that are more likely to display signs and symptoms of disease after exposure to a pathogen (e.g., immune suppressed subjects). In

some embodiments, the general public is administered (e.g., vaccinated with) a composition of the present invention (e.g., to prevent the occurrence or spread of disease). For example, in some embodiments, compositions and methods of the present invention are utilized to vaccinate a group of people (e.g., a population of a region, city, state and/or country) for their own health (e.g., to prevent or treat disease). In some embodiments, the subjects are non-human mammals (e.g., pigs, cattle, goats, horses, sheep, or other livestock; or mice, rats, rabbits or other animal). In some embodiments, compositions and methods of the present invention are utilized in research settings (e.g., with research animals).

A composition of the present invention may be formulated for administration by any route, such as mucosal, oral, topical, parenteral or other route described herein. The compositions may be in any one or more different forms including, but not limited to, tablets, capsules, powders, granules, lozenges, foams, creams or liquid preparations.

Topical formulations of the present invention may be presented as, for instance, ointments, creams or lotions, foams, and aerosols, and may contain appropriate conventional additives such as preservatives, solvents (e.g., to assist penetration), and emollients in ointments and creams.

Topical formulations may also include agents that enhance penetration of the active ingredients through the skin. Exemplary agents include a binary combination of N-(hydroxyethyl) pyrrolidone and a cell-envelope disordering compound, a sugar ester in combination with a sulfoxide or phosphine oxide, and sucrose monooleate, decyl methyl sulfoxide, and alcohol.

Other exemplary materials that increase skin penetration include surfactants or wetting agents including, but not limited to, polyoxyethylene sorbitan mono-oleate (Polysorbate 80); sorbitan mono-oleate (Span 80); p-isooctyl polyoxyethylene-phenol polymer (Triton WR-1330); polyoxyethylene sorbitan tri-oleate (Tween 85); dioctyl sodium sulfosuccinate; and sodium sarcosinate (Sarcosyl NL-97); and other pharmaceutically acceptable surfactants.

In certain embodiments of the invention, compositions may further comprise one or more alcohols, zinc-containing compounds, emollients, humectants, thickening and/or gelling agents, neutralizing agents, and surfactants. Water used in the formulations is preferably deionized water having a neutral pH. Additional additives in the topical formulations include, but are not limited to, silicone fluids, dyes, fragrances, pH adjusters, and vitamins.

Topical formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be present as from about 1% up to about 98% of the formulation. The ointment base can comprise one or more of petrolatum, mineral oil, ceresin, lanolin alcohol, panthenol, glycerin, bisabolol, cocoa butter and the like.

In some embodiments, pharmaceutical compositions of the present invention may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these formulations vary in the components and the consistency of the final product.

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, preferably do not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents (e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like) that do not deleteriously interact with the NE and immunogen of the formulation. In some embodiments, immunostimulatory compositions of the present invention are administered in the form of a pharmaceutically acceptable salt. When used the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically acceptable salts thereof. Such salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene sulphonic, tartaric, citric, methane sulphonic, formic, malonic, succinic, naphthalene-2-sulphonic, and benzene sulphonic. Also, such salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts of the carboxylic acid group.

Suitable buffering agents include, but are not limited to, acetic acid and a salt (1-2% w/v); citric acid and a salt (1-3% w/v); boric acid and a salt (0.5-2.5% w/v); and phosphoric

acid and a salt (0.8-2% w/v). Suitable preservatives may include benzalkonium chloride (0.003-0.03% w/v); chlorobutanol (0.3-0.9% w/v); parabens (0.01-0.25% w/v) and thimerosal (0.004-0.02% w/v).

In some embodiments, a composition comprising a NE and an immunogen is co-administered with one or more antibiotics. For example, one or more antibiotics may be administered with, before and/or after administration of a composition comprising a NE and an immunogen. The present invention is not limited by the type of antibiotic co-administered. Indeed, a variety of antibiotics may be co-administered including, but not limited to, β -lactam antibiotics, penicillins (such as natural penicillins, aminopenicillins, penicillinase-resistant penicillins, carboxy penicillins, ureido penicillins), cephalosporins (first generation, second generation, and third generation cephalosporins), and other β -lactams (such as imipenem, monobactams, β -lactamase inhibitors, vancomycin, aminoglycosides and spectinomycin, tetracyclines, chloramphenicol, erythromycin, lincomycin, clindamycin, rifampin, metronidazole, polymyxins, doxycycline, quinolones (e.g., ciprofloxacin), sulfonamides, trimethoprim, and quinolines.

There are an enormous amount of antimicrobial agents currently available for use in treating bacterial, fungal and viral infections. For a comprehensive treatise on the general classes of such drugs and their mechanisms of action, the skilled artisan is referred to Goodman & Gilman's "The Pharmacological Basis of Therapeutics" Eds. Hardman *et al.*, 9th Edition, Pub. McGraw Hill, chapters 43 through 50, 1996,

Generally, these agents include agents that inhibit cell wall synthesis (e.g., penicillins, cephalosporins, cycloserine, vancomycin, bacitracin); and the imidazole antifungal agents (e.g., miconazole, ketoconazole and clotrimazole); agents that act directly to disrupt the cell membrane of the microorganism (e.g., detergents such as polymyxin and colistimethate and the antifungals nystatin and amphotericin B); agents that affect the ribosomal subunits to inhibit protein synthesis (e.g., chloramphenicol, the tetracyclines, erythromycin and clindamycin); agents that alter protein synthesis and lead to cell death (e.g., aminoglycosides); agents that affect nucleic acid metabolism (e.g., the rifamycins and the quinolones); the antimetabolites (e.g., trimethoprim and sulfonamides); and the nucleic acid analogues such as zidovudine, gancyclovir, vidarabine, and acyclovir which act to inhibit viral enzymes essential for DNA synthesis. Various combinations of antimicrobials may be employed.

The present invention also includes methods involving co-administration of a composition comprising a NE and one or a plurality of immunogens with one or more additional active and/or immunostimulatory agents (e.g., a composition comprising a NE and a different immunogen, an antibiotic, anti-oxidant, etc.). Indeed, it is a further aspect of this invention to provide methods for enhancing prior art immunostimulatory methods (e.g., immunization methods) and/or pharmaceutical compositions by co-administering a composition of the present invention. In co-administration procedures, the agents may be administered concurrently or sequentially. In one embodiment, the compositions described herein are administered prior to the other active agent(s). The pharmaceutical formulations and modes of administration may be any of those described herein. In addition, the two or more co-administered agents may each be administered using different modes (e.g., routes) or different formulations. The additional agents to be co-administered (e.g., antibiotics, adjuvants, etc.) can be any of the well-known agents in the art, including, but not limited to, those that are currently in clinical use.

In some embodiments, a composition comprising a NE and one or a plurality of immunogens is administered to a subject via more than one route. For example, a subject that would benefit from having a protective immune response (e.g., immunity) towards a pathogenic microorganism may benefit from receiving mucosal administration (e.g., nasal administration or other mucosal routes described herein) and, additionally, receiving one or more other routes of administration (e.g., parenteral or pulmonary administration (e.g., via a nebulizer, inhaler, or other methods described herein)). In some preferred embodiments, administration via mucosal route is sufficient to induce both mucosal as well as systemic immunity towards an immunogen or organism from which the immunogen is derived. In other embodiments, administration via multiple routes serves to provide both mucosal and systemic immunity. Thus, although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments, it is contemplated that a subject administered a composition of the present invention via multiple routes of administration (e.g., immunization (e.g., mucosal as well as airway or parenteral administration of a composition comprising a NE and immunogen of the present invention) may have a stronger immune response to an immunogen than a subject administered a composition via just one route.

Other delivery systems can include time-release, delayed release or sustained release

delivery systems. Such systems can avoid repeated administrations of the compositions, increasing convenience to the subject and a physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer based systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides.

Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Pat. No. 5,075,109.

Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-di-and tri-glycerides; hydrogel release systems; sytastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which an agent of the invention is contained in a form within a matrix such as those described in U.S. Pat. Nos. 4,452,775, 4,675,189, and 5,736,152

and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Pat. Nos. 3,854,480, 5,133,974 and 5,407,686.

In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation.

In some embodiments, a composition comprising a NE and one or a plurality of immunogens of the present invention is formulated in a concentrated dose that can be diluted prior to administration to a subject. For example, dilutions of a concentrated composition may be administered to a subject such that the subject receives any one or more of the specific dosages provided herein. In some embodiments, dilution of a concentrated composition may be made such that a subject is administered (e.g., in a single dose) a composition comprising 0.5-50% of the NE and immunogen present in the concentrated composition. Concentrated compositions are contemplated to be useful in a setting in which large numbers of subjects may be administered a composition of the present invention (e.g., an immunization clinic, hospital, school, etc.). In some embodiments, a composition comprising a NE and an immunogen of the present invention (e.g., a concentrated composition) is stable at room temperature for more than 1 week, in some embodiments for more than 2 weeks, in some embodiments for more than 3 weeks, in some embodiments for more than 4 weeks, in some embodiments for more than 5 weeks, and in some embodiments

for more than 6 weeks.

In some embodiments, following an initial administration of a composition of the present invention (e.g., an initial vaccination), a subject may receive one or more boost administrations (e.g., around 2 weeks, around 3 weeks, around 4 weeks, around 5 weeks, 5 around 6 weeks, around 7 weeks, around 8 weeks, around 10 weeks, around 3 months, around 4 months, around 6 months, around 9 months, around 1 year, around 2 years, around 3 years, around 5 years, around 10 years) subsequent to a first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, and/or more than tenth administration. Although an understanding of the mechanism is not necessary to practice the present invention and the 10 present invention is not limited to any particular mechanism of action, in some embodiments, reintroduction of an immunogen in a boost dose enables vigorous systemic immunity in a subject. The boost can be with the same formulation given for the primary immune response, or can be with a different formulation that contains the immunogen. The dosage regimen will also, at least in part, be determined by the need of the subject and be dependent on the 15 judgment of a practitioner.

Dosage units may be proportionately increased or decreased based on several factors including, but not limited to, the weight, age, and health status of the subject. In addition, dosage units may be increased or decreased for subsequent administrations (e.g., boost administrations).

20 It is contemplated that the compositions and methods of the present invention will find use in various settings, including research settings. For example, compositions and methods of the present invention also find use in studies of the immune system (e.g., characterization of adaptive immune responses (e.g., protective immune responses (e.g., mucosal or systemic immunity))). Uses of the compositions and methods provided by the 25 present invention encompass human and non-human subjects and samples from those subjects, and also encompass research applications using these subjects. Compositions and methods of the present invention are also useful in studying and optimizing nanoemulsions, immunogens, and other components and for screening for new components. Thus, it is not intended that the present invention be limited to any particular subject and/or application 30 setting.

In preferred embodiments, a composition comprising a NE and an immunogen of the present invention comprises a suitable amount of the immunogen to induce an immune

response in a subject when administered to the subject. In preferred embodiments, the immune response is sufficient to provide the subject protection (e.g., immune protection) against a subsequent exposure to the immunogen or the microorganism (e.g., bacteria or virus) from which the immunogen was derived. The present invention is not limited by the amount of immunogen used. In some preferred embodiments, the amount of immunogen (e.g., virus or bacteria neutralized by the NE, or, recombinant protein) in a composition comprising a NE and immunogen (e.g., for use as an immunization dose) is selected as that amount which induces an immunoprotective response without significant, adverse side effects. The amount will vary depending upon which specific immunogen or combination thereof is/are employed, and can vary from subject to subject, depending on a number of factors including, but not limited to, the species, age and general condition (e.g., health) of the subject, and the mode of administration. Procedures for determining the appropriate amount of immunogen administered to a subject to elicit an immune response (e.g., a protective immune response (e.g., protective immunity)) in a subject are well known to those skilled in the art.

In some embodiments, it is expected that each dose (e.g., of a composition comprising a NE and one or a plurality of immunogens (e.g., administered to a subject to induce an immune response (e.g., a protective immune response (e.g., protective immunity)))) comprises 0.05-5000 μg of each immunogen (e.g., recombinant and/or purified protein), in some embodiments, each dose will comprise 1-500 μg , in some embodiments, each dose will comprise 350-750 μg , in some embodiments, each dose will comprise 50-200 μg , in some embodiments, each dose will comprise 25-75 μg of each immunogen (e.g., recombinant and/or purified protein). In some embodiments, each dose comprises an amount of the immunogen sufficient to generate an immune response. An effective amount of the immunogen in a dose need not be quantified, as long as the amount of immunogen generates an immune response in a subject when administered to the subject. An optimal amount for a particular administration (e.g., to induce an immune response (e.g., a protective immune response (e.g., protective immunity))) can be ascertained by one of skill in the art using standard studies involving observation of antibody titers and other responses in subjects.

In some embodiments, it is expected that each dose (e.g., of a composition comprising a NE and one or a plurality of antigens (e.g., administered to a subject to induce and immune response)) is from 0.001 to 15% or more (e.g., 0.001-10%, 0.5-5%, 1-3%, 2%, 6%, 10%,

15% or more) by weight immunogen (e.g., neutralized bacteria or virus, or recombinant and/or purified protein). In some embodiments, an initial or prime administration dose contains more immunogen than a subsequent boost dose

5 In some embodiments, when a NE of the present invention is utilized to inactivate a live microorganism (e.g., virus (e.g., HIV)), it is expected that each dose (e.g., administered to a subject to induce and immune response)) comprises between 10 and 10^9 pfu of the virus per dose; in some embodiments, each dose comprises between 10^5 and 10^8 pfu of the virus per dose; in some embodiments, each dose comprises between 10^3 and 10^5 pfu of the virus per dose; in some embodiments, each dose comprises between 10^2 and 10^4 pfu of the virus per
10 dose; in some embodiments, each dose comprises 10 pfu of the virus per dose; in some embodiments, each dose comprises 10^2 pfu of the virus per dose; and in some embodiments, each dose comprises 10^4 pfu of the virus per dose. In some embodiments, each dose comprises more than 10^9 pfu of the virus per dose. In some preferred embodiments, each dose comprises 10^3 pfu of the virus per dose.

15 In some embodiments, when a NE of the present invention is utilized to inactivate a live microorganism (e.g., a population of bacteria (e.g., of the genus *Bacillus* (*B. anthracis*))), it is expected that each dose (e.g., administered to a subject to induce and immune response)) comprises between 10 and 10^{10} bacteria per dose; in some embodiments, each dose comprises between 10^5 and 10^8 bacteria per dose; in some embodiments, each dose comprises between
20 10^3 and 10^5 bacteria per dose; in some embodiments, each dose comprises between 10^2 and 10^4 bacteria per dose; in some embodiments, each dose comprises 10 bacteria per dose; in some embodiments, each dose comprises 10^2 bacteria per dose; and in some embodiments, each dose comprises 10^4 bacteria per dose. In some embodiments, each dose comprises more than 10^{10} bacteria per dose. In some embodiments, each dose comprises 10^3 bacteria per dose.

25 The present invention is not limited by the amount of NE used to inactivate live microorganisms (e.g., a virus (e.g., one or more types of HIV)). In some embodiments, a 0.1% - 5% NE solution is used, in some embodiments, a 5%-20% NE solution is used, in some embodiments, a 20% NE solution is used, and in some embodiments, a NE solution greater than 20% is used order to inactivate a pathogenic microorganism. In some
30 embodiments, a 10% NE solution is used.

Similarly, the present invention is not limited by the duration of time a live microorganism is incubated in a NE of the present invention in order to become inactivated.

In some embodiments, the microorganism is incubated for 1-3 hours in NE. In some
embodiments, the microorganism is incubated for 3-6 hours in NE. In some embodiments,
the microorganism is incubated for more than 6 hours in NE. In preferred embodiments, the
microorganism is incubated for 3 hours in NE (e.g., a 10% NE solution). In some
5 embodiments, the incubation is carried out at 37°C. In some embodiments, the incubation is
carried out at a temperature greater than or less than 37°C. The present invention is also not
limited by the amount of microorganism used for inactivation. The amount of microorganism
may depend upon a number of factors including, but not limited to, the total amount of
immunogenic composition (e.g., NE and immunogen) desired, the concentration of solution
10 desired (e.g., prior to dilution for administration), the microorganism and the NE. In some
preferred embodiments, the amount of microorganism used in an inactivation procedure is
that amount that produces the desired amount of immunogen (e.g., as described herein) to be
administered in a single dose (e.g., diluted from a concentrated stock) to a subject.

In some embodiments, a composition comprising a NE and one or more immunogens
15 of the present invention is formulated in a concentrated dose that can be diluted prior to
administration to a subject. For example, dilutions of a concentrated composition may be
administered to a subject such that the subject receives any one or more of the specific
dosages provided herein. In some embodiments, dilution of a concentrated composition may
be made such that a subject is administered (e.g., in a single dose) a composition comprising
20 0.1-50% of the NE and one or more immunogens present in the concentrated composition. In
some preferred embodiments, a subject is administered in a single dose a composition
comprising 1% of the NE and one or more immunogens present in the concentrated
composition. Concentrated compositions are contemplated to be useful in a setting in which
large numbers of subjects may be administered a composition of the present invention (e.g.,
25 an immunization clinic, hospital, school, etc.). In some embodiments, a composition
comprising a NE and one or more immunogens of the present invention (e.g., a concentrated
composition) is stable at room temperature for more than 1 week, in some embodiments for
more than 2 weeks, in some embodiments for more than 3 weeks, in some embodiments for
more than 4 weeks, in some embodiments for more than 5 weeks, and in some embodiments
30 for more than 6 weeks.

Generally, the emulsion compositions of the invention will comprise at least 0.001%
to 100%, preferably 0.01 to 90%, of emulsion per ml of liquid composition. It is envisioned

that the formulations may comprise about 0.001%, about 0.0025%, about 0.005%, about 0.0075%, about 0.01%, about 0.025%, about 0.05%, about 0.075%, about 0.1 %, about 0.25%, about 0.5%, about 1.0%, about 2.5%, about 5%, about 7.5%, about 10%, about 12.5%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 50%,
5 about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95% or about 100% of emulsion per ml of liquid composition. It should be understood that a range between any two figures listed above is specifically contemplated to be encompassed within the metes and bounds of the present invention. Some variation in dosage will necessarily occur depending on the condition of the specific pathogen and the
10 subject being immunized.

In some embodiments, following an initial administration of a composition of the present invention (e.g., an initial vaccination), a subject may receive one or more boost administrations (e.g., around 2 weeks, around 3 weeks, around 4 weeks, around 5 weeks, around 6 weeks, around 7 weeks, around 8 weeks, around 10 weeks, around 3 months, around
15 4 months, around 6 months, around 9 months, around 1 year, around 2 years, around 3 years, around 5 years, around 10 years) subsequent to a first, second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, and/or more than tenth administration. Although an understanding of the mechanism is not necessary to practice the present invention and the present invention is not limited to any particular mechanism of action, in some embodiments,
20 reintroduction of an immunogen in a boost dose enables vigorous systemic immunity in a subject. The boost can be with the same formulation given for the primary immune response, or can be with a different formulation that contains the immunogen. The dosage regimen will also, at least in part, be determined by the need of the subject and be dependent on the judgment of a practitioner.

25 Dosage units may be proportionately increased or decreased based on several factors including, but not limited to, the weight, age, and health status of the subject. In addition, dosage units may be increased or decreased for subsequent administrations (e.g., boost administrations).

A composition comprising an immunogen of the present invention finds use where the
30 nature of the infectious and/or disease causing agent (e.g., for which protective immunity is sought to be elicited) is known, as well as where the nature of the infectious and/or disease causing agent is unknown (e.g., in emerging disease (e.g., of pandemic proportion (e.g.,

influenza or other outbreaks of disease))). For example, the present invention contemplates use of the compositions of the present invention in treatment of or prevention of (e.g., via immunization with an infectious and/or disease causing HIV or HIV-like agent neutralized via a NE of the present invention) infections associated with an emergent infectious and/or disease causing agent yet to be identified (e.g., isolated and/or cultured from a diseased person but without genetic, biochemical or other characterization of the infectious and/or disease causing agent).

It is contemplated that the compositions and methods of the present invention will find use in various settings, including research settings. For example, compositions and methods of the present invention also find use in studies of the immune system (e.g., characterization of adaptive immune responses (e.g., protective immune responses (e.g., mucosal or systemic immunity))). Uses of the compositions and methods provided by the present invention encompass human and non-human subjects and samples from those subjects, and also encompass research applications using these subjects. Compositions and methods of the present invention are also useful in studying and optimizing nanoemulsions, immunogens, and other components and for screening for new components. Thus, it is not intended that the present invention be limited to any particular subject and/or application setting.

The formulations can be tested *in vivo* in a number of animal models developed for the study of mucosal and other routes of delivery. As is readily apparent, the compositions of the present invention are useful for preventing and/or treating a wide variety of diseases and infections caused by viruses, bacteria, parasites, and fungi, as well as for eliciting an immune response against a variety of antigens. Not only can the compositions be used prophylactically or therapeutically, as described above, the compositions can also be used in order to prepare antibodies, both polyclonal and monoclonal (e.g., for diagnostic purposes), as well as for immunopurification of an antigen of interest. If polyclonal antibodies are desired, a selected mammal, (e.g., mouse, rabbit, goat, horse, etc.) can be immunized with the compositions of the present invention. The animal is usually boosted 2-6 weeks later with one or more--administrations of the antigen. Polyclonal antisera can then be obtained from the immunized animal and used according to known procedures (See, e.g., Jurgens et al., J. Chrom. 1985, 348:363-370).

In some embodiments, the present invention provides a kit comprising a composition

comprising a NE and one or more immunogens. In some embodiments, the kit further provides a device for administering the composition. The present invention is not limited by the type of device included in the kit. In some embodiments, the device is configured for nasal application of the composition of the present invention (e.g., a nasal applicator (e.g., a syringe) or nasal inhaler or nasal mister). In some embodiments, a kit comprises a composition comprising a NE and one or more immunogens in a concentrated form (e.g., that can be diluted prior to administration to a subject).

In some embodiments, all kit components are present within a single container (e.g., vial or tube). In some embodiments, each kit component is located in a single container (e.g., vial or tube). In some embodiments, one or more kit component are located in a single container (e.g., vial or tube) with other components of the same kit being located in a separate container (e.g., vial or tube). In some embodiments, a kit comprises a buffer. In some embodiments, the kit further comprises instructions for use.

EXAMPLES

The following examples serve to illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

In the experimental disclosure which follows, the following abbreviations apply: eq (equivalents); μ (micron); M (Molar); μ M (micromolar); mM (millimolar); N (Normal); mol (moles); mmol (millimoles); μ mol (micromoles); nmol (nanomoles); g (grams); mg (milligrams); μ g (micrograms); ng (nanograms); L (liters); ml (milliliters); μ l (microliters); cm (centimeters); mm (millimeters); μ m (micrometers); nM (nanomolar); $^{\circ}$ C (degrees Centigrade); and PBS (phosphate buffered saline).

Example 1

Nanoemulsion-based Hepatitis B (HB) vaccine and methods of using the same

Adjuvant and antigen. Nanoemulsion (NE, W₈₀5EC formulation, described herein) was supplied by NANOBIO Corporation, Ann Arbor, MI. Nanoemulsion was manufactured by emulsification of cetyl pyridinium chloride (CPC, 1%), Tween 80 (5%) and ethanol (8%) in water with soybean oil (64%) using a high speed emulsifier, with resultant mean droplet size of less than 400 nm in diameter. W₈₀5EC is formulated with surfactants and food substances that are 'Generally Recognized as Safe' (GRAS) by the FDA, and can be

economically manufactured under Good Manufacturing Practices (GMP). The nanoemulsion is stable for at least 3 years at 25°C.

Recombinant HBs antigen *adw* serotype used for immunizations (HBsAg) was supplied by Human Biologicals Institute (Indian Immunologicals, Ltd., Hyderabad, India). The antigen protein was purified from *Pichia pastoris* transfected with plasmid pPIC3K using methods according to Indian Immunologicals SOP and GMP procedures. HBsAg was dissolved in PBS (pH 7.03) and endotoxin level was determined to be < 7.5 EU/20 g of protein; below international standard of ≤ 30 EU/20 g of protein.

Reagents. Phosphate buffered saline (1 x PBS and 10 x PBS, pH 7.4) was purchased from CELLGRO (MEDTECH, Inc). Deionized water was prepared using a MILLI-Q Ultrapure Water Purification system (MILLIPORE, Billerica, MA). The bovine serum albumin (BSA) was purchased from SIGMA. Alkaline phosphatase (AP) conjugated rabbit anti-mouse IgG (H&L), IgG1, IgG2a, IgG2b, IgG3, IgA (α chain specific), goat anti-rat IgG (H&L), and goat anti-guinea pig IgG (H&L) secondary antibodies were purchased from ROCKLAND Immunochemicals, Inc.

Particle sizing. HBsAg-NE formulations were prepared by vigorously mixing concentrated NE with HBsAg and PBS. Mixtures contained a final concentration of 0.5 mg/ml or 2.5 mg/ml of antigen mixed in 1%, 20%, or 40% (v/v) NE concentrations and normalized to 1 x PBS.

The lipid-phase NE droplets were sized by quasi-elastic light scattering using an LS230 instrument (BECKMAN-COULTER, Fullerton, CA) following manufacturer's protocols. In brief, between 10 μ l and 30 μ l of NE-antigen mixtures were diluted into a flow chamber containing 1 L of deionized water. Particle size distributions were calculated using number weighting, and statistics were generated from the average of three 60 second measurement cycles. Sample concentration was optimized based on PIDS obscuration, and PIDS data was included in the instrument's Fraunhofer model calculation.

HBsAg analysis. The integrity of HBsAg protein was analyzed using SDS-PAGE and Western blotting techniques. HBsAg was mixed in 20% NE at 0.5 mg/ml and 2.5 mg/ml concentrations. Aliquots of each of the HBsAg-NE mixtures were incubated at 4°C, 25°C and 40°C for up to 72 hrs. For PAGE analysis, the HBsAg samples were resuspended in 1% SDS, reduced with β -mercaptoethanol (BME, 2.5%) and boiled for 15 minutes. The electrophoresis was performed in duplicates using 0.5ug HBsAg, 4-12% Bis-Tris PAGE gels

(INVITROGEN), and MES SDS Running Buffer. One gel of each duplicate was stained using the SILVERQUEST Silver Staining Kit (INVITROGEN). For Western blots, gels were transferred onto Immobilon-P PVDF membrane (MILLIPORE) in NuPAGE transfer buffer according to INVITROGEN's protocol. The membranes were blocked for 1 hr in 5% Milk/PBST and were probed with a polyclonal goat anti-HBsAg (ABCAM). Alkaline phosphatase-(AP) conjugated anti-goat (SIGMA) secondary antibodies were used with 1-Step NBT/BCIP AP substrate (PIERCE) for protein detection.

Zeta potential measurement. Zeta potential measurements were obtained using a NICOM 380ZLS (PSS.NICOMP, Santa Barbara, CA). Samples containing 20% NE mixed with 2.5 mg/ml HBsAg were prepared by vigorously mixing concentrated NE and HBsAg. Test mixtures were diluted in either PBS or de-ionized water. Zeta potential was measured in 200 x diluted samples at 25°C.

Isothermal titration calorimetry. The interaction of the amphiphilic HBsAg with the lipid phase of NE was studied using an isothermal titration microcalorimeter (VP-ITC MICROCALRIMITER, MICROCAL). HBsAg solutions in PBS aliquots were prepared from concentrated stock and introduced into the calorimetric reaction and reference vessels (1.3 ml). Chambers were then gently agitated until temperature equilibrium with the surroundings was reached. Concentrated NE (50% wt) was diluted in PBS to 1% (v/v). After the sample vessel had reached the equilibrium conditions, the NE solution was added in discrete injections using a syringe, into the calorimetric reaction vessel under continuous stirring (either 30°C or 40°C). The experimentally observed change of energy corresponding to a given injection of NE was measured and plotted (ORIGIN 7SR4 v. 7 ORIGIN Lab Corp., Northhampton, MA). The change in heat capacity of binding (ΔC_p) was calculated using the following equation: $\Delta C_p = (\Delta H^\circ_{T_2} - \Delta H^\circ_{T_1}) / T_2 - T_1$ where ΔH is calculated enthalpy and T is vessel temperature (VP-ITC MICROCALORIMETER User's Manual. 2007, MICROCAL, LLC.: Northhampton, MA).

Preparation of HBsAg-NE vaccine. HBsAg-NE formulations were prepared 30 to 60 minutes prior to immunization by vigorously mixing HBsAg protein solution with concentrated NE using PBS as diluent. For intranasal immunizations HBsAg-NE doses ranged from 1 µg to 40 µg HBsAg mixed with 5% to 40% NE. For intramuscular immunizations with the HBsAg/aluminum hydroxide vaccine (HBsAg-Alu), antigen was adsorped onto 0.5 mg/ml aluminium hydroxide (SIGMA) following the adsorption procedure

described in Little et al. to obtain formulation similar to that of ENERGIX (GLAXOSMITHKLINE).

Animals. Pathogen-free, outbred CD-1 mice (females 6-8 weeks old), inbred BALB/c mice (females 6-8 weeks old), and Hartley guinea pigs (females 10-11 weeks old) were purchased from CHARLES RIVER LABORATORIES. Pathogen free Sprague Dawley rats (females 7-8 weeks old) and specific pathogen free (SPF) purpose-bred American standard beagles (females, 6 month old) were obtained from HARLAN and COVANCE, respectively. Animals used in these studies were housed in SPF conditions with food and water available ad libitum in accordance to the standards of the American Association for Accreditation of Laboratory Animal Care. Mice were housed with 5 to a cage. Rats and guinea pigs were housed 3 to a cage. Dogs were housed in floor pens with soft bedding and in a rotating group setting. Daily exercise was provided as enrichment. All procedures performed on animals within this study were conducted in accordance with and by approval of the University of Michigan University Committee on Use and Care of Animals (UCUCA).

Immunization procedures. CD-1 mice were vaccinated with two administrations of HBsAg-NE vaccine six weeks apart. Both intranasal (i.n.) and intramuscular (i.m.) immunizations were performed in mice anaesthetized with isoflurane using IMPAC 6 anesthesia delivery system. For i.n. administration, animals were held in a supine position and 8 μ l (4 μ l/nare) of HBsAg-NE vaccine was administered slowly to the nares using a micropipette tip. For i.m. immunization, 50 μ l of HBsAg-Alu vaccine was injected into apaxial muscle. Rats, and guinea pigs were also manually restrained in a supine position and 100 μ l (50 μ l/nare) of HBsAg-NE vaccine was administered slowly to the nares using a micropipette tip.

Blood, bronchioalveolar lavage, and splenocyte collection. Blood samples were obtained from the saphenous vein in mice, rats, and guinea pigs and from the superficial cephalic vein in dogs at various time points during the course of the experiments. The terminal murine sample was obtained by cardiac puncture post-euthanasia. Serum was separated from whole blood by centrifugation at $1500 \times g$ for 5 minutes after allowing coagulation for 30 to 60 minutes at room temperature. Serum samples were stored at $-20^{\circ}C$ until analyzed. Bronchioalveolar lavage (BAL) fluid was obtained from mice euthanized by an overdose of isoflurane. A 22 gauge catheter (Angiocath, B-D) attached to a syringe was inserted into the distal trachea. The lungs were infused twice with 0.5 ml of PBS containing

10 μ M DTT and 0.5 mg/ml aprotinin and approximately 1 ml of aspirate was recovered. BAL samples were stored at -20° C until analyzed.

At the time of euthanasia, spleens were harvested from mice and mechanically disrupted to obtain single-cell splenocyte suspension in PBS, which was used for in vitro
5 determination of cytokine response. Red blood cells were removed by lysis with ACK buffer (150 mM NH_4Cl , 10 mM KHCO_3 , 0.1 mM Na_2EDTA), and the remaining cells were washed twice in PBS. For the cytokine expression assays, splenocytes were resuspended in RPMI 1640 medium supplemented with 2% FBS, 200 nM L-glutamine, and penicillin/streptomycin (100 U/ml and 100 μ g/ml).

10 Determination of IgG and IgA antibodies in serum and BAL fluid. Mouse, rat, and guinea pig anti-HBsAg specific IgG and mouse anti-HBsAg specific IgA levels were determined by ELISA. Microtiter plates (NUNC) were coated with 5 μ g/ml (100 μ l) of HBsAg in a coating buffer (50 mM sodium carbonate, 50 mM sodium bicarbonate, pH 9.6) and incubated overnight at 4° C. The protein solution was removed and plates were incubated
15 with blocking buffer (PBS with 1% dry milk) for 30 minutes at 37° C. After the blocking solution was aspirated, the plates were used immediately or stored sealed at 4° C until needed. For antibody detection, serum and BAL samples were serially diluted in 0.1% BSA in PBS. The 100 μ l/well aliquots were incubated in HBsAg coated plates for 1 hour at 37° C. Plates were washed three times with PBS containing 0.05% Tween 20, followed by 1 hour
20 incubation with either species specific anti-IgG or IgA alkaline phosphatase (AP)-conjugated antibodies, then washed three times and incubated with AP substrate SIGMA FAST (SIGMA). The colorimetric reaction was stopped with 1 N NaOH according to the manufacturer's protocol, and optical density (OD) measured using a SPECTRA MAX 340 ELISA reader (MOLECULAR DEVICES, Sunnyvale, CA) at 405 nm and the reference
25 wavelength of 690 nm. The antibody concentrations are presented as endpoint titers defined as the reciprocal of the highest serum dilution producing an OD above cutoff value. The cutoff value is determined as OD of the corresponding dilution of control sera + 2 (standard deviations) and plate background (Classen et al. J Clin Microbiol, 1987. 25(4): 600-604; Frey et al. J Immunol Methods, 1998. 221(35-41).). Normalization of IgG was performed at
30 UMHHC diagnostic laboratory using an ADVIA Centaur anti-HBsAg assay.

Determination of IgG avidity. The avidity index (AI) was determined by ELISA using mouse serum as described by Vermont et al. with minor modifications (Vermont et al.

Infect Immun, 2002. 70(2): 584-590). Sodium thiocyanate (NaSCN) was used for dissociation of low avidity antibody-antigen binding. Optimal assay conditions for determination of AI were established in an ELISA assay using 0 M to 3 M range of NaSCN concentrations. Incubation with 1.5 M NaSCN solution resulted in reduction of antibody binding that was discriminating between serum samples. In each assay, serial dilutions of immune serum were incubated with HBsAg as described above for standard ELISA. To differentiate antibody binding, the wells were incubated with either PBS or with 1.5 M NaSCN at room temperature for 15 minutes. Subsequently wells were washed three times and incubated with anti-mouse IgG AP-conjugate as described above. The AI was calculated as percentage of antibody titer which remained bound to antigen after incubation with NaSCN in comparison to the standard ELISA protocol.

LUMINEX analysis of cytokine expression. Freshly isolated mouse murine splenocytes were seeded at 4×10^6 cells/ml (RPMI 1640, 2% FBS) and incubated with HBsAg (5 μ g/ml) or control PHA-P mitogen (2 μ g/ml) for 72 hours. Cell culture supernatants were harvested and analyzed for the presence of cytokines. The IL-4, IL-5, IL-10, IFN- γ and TNF- α cytokine assays were performed using LUMINEX Multiplex21 multi-analyte profiling beads (LUMINEX Corporation, Austin, TX), according to the manufacturer's instructions.

Analyses of thermostability of HBsAg-NE. For vaccine thermostability studies, the formulation was made by vigorously mixing HBsAg and NE to achieve a dose of 2.5 mg/ml recombinant protein in 20% NE and a final buffered solution of 1X PBS. The vaccine was then aliquoted into sterile glass vials with TEFLON-coated caps (Wheaton) and stored at either $4 \pm 2^\circ\text{C}$, $25 \pm 2^\circ\text{C}$ or $40 \pm 2^\circ\text{C}$. Temperatures were monitored for the period of the study by Lufft OPUS10 thermographs (PalmerWahl). At time points of 6 weeks, 12 weeks (3 months), 24 weeks (6 months) and 52 weeks (1 year), an aliquot was withdrawn and used for in vitro as well as in vivo analyses. For in vitro analyses 0.5 μ g of antigen contained in vaccine product was electrophoresed per lane and detected by silver staining and Western blotting (as described above); NE particle size was also determined (as described above). In vivo immunogenicity studies were done by intranasal vaccinations (primed at 0 and boosted at 6 weeks) of about 8 week old female CD-1 mice and testing serum IgG titers at 2, 3, 5, 8, 10 and 12 weeks as described above.

Comprehensive toxicity assessments. Acute and (sub) chronic toxicity responses to either NE or HBsAg-NE were assessed in mice, rats, guinea pigs, and dogs. Numerous

species were evaluated in order to minimize the effects of animal model biasing. The end points of the study were histopathological evaluation of exposed tissues and of highly perfused organs. Metabolic changes were also measured using serum biochemical profile analysis.

5 The clinical status of each animal including the nasal cavity, body weight, body temperature, and food consumption was assessed throughout the study. Mice were non-surgically implanted with programmable temperature transponders (IPTT-3000, Bio Medic Data Systems, Inc.) for non-invasive subcutaneous temperature measurement with a handheld portable scanner (DAS-6002, Bio Medic Data Systems, Inc.). Euthanasia by isoflurane
10 asphyxiation was performed in mice whereas rats and guinea pigs were euthanized by barbiturate overdose. A complete necropsy, which included the gross pathological examination of the external surface of the body, all orifices, and the cranial thoracic and abdominal cavities and their contents, was performed on all rodent
15 species at the time of death. Vaccine exposed tissues and highly perfused organs including the sinus cavity, lungs, esophagus, trachea, brain, heart, liver, kidneys, spleen, stomach, intestines, pancreas, and adrenals were collected and immediately fixed in 10% buffered formalin (FISCHER SCIENTIFIC).

 In order to assess safety and tolerability of the adjuvant, NE was delivered to dogs using a wide angle nasal sprayer pump (Pfeiffer 62602, 415 screw enclosure). The containers
20 used were Saint Gobain Desqueres 5-mL U-SAVE Type 1 amber glass bottles with a 415 neck finish. The dose volume for the sprayer pump was 100 μ l. Dogs received either 200 μ l (100 μ /nare) or 400 μ l (200 μ /nare) administered every 14 days for a total of 3 doses as outlined (See FIG. 50). Rostral nasal sinus punch biopsy samples were collected 24 hours following the final treatment. For the biopsy procedure, dogs were anesthetized with
25 ketamine/diazepam/butorphanol (10 mg/kg, 0.5 mg/kg, 3 mg/kg) and maintained on 2.5% isoflurane after endotracheal intubation. The anterior sinus cavity and external nares were sterilely prepared. A sterile dermal punch biopsy instrument (MILTEX, 4 mm) was introduced approximately 1.5 cm into ventral portion of the anterior sinus cavity. Hemostasis was achieved using 4-0 PDS suture material. Tissues obtained for biopsy were immediately
30 fixed in 10% buffered formalin (FISCHER SCIENTIFIC). Butorphanol (3 mg/kg) administration was continued every 8 hours for three days following the biopsy procedure for analgesic management.

Histopathological analysis. Harvested tissues were fixed in 10% formalin solution for at least 24 hours. Sinus tissues including bone were decalcified for 48 hours using CAL-EX II (FISCHER SCIENTIFIC) prior to trimming and embedding in paraffin. For mice, rats, and guinea pigs, four standard cross sections of the nasal passages including the brain were taken (Herber,t RA and Leininger, JR. Pathology of the Mouse. 1999, St. Louis: Cache River Press.). Tissue blocks were processed in xylene and paraffin embedded for multi-sections and slide preparation. Routine hematoxylin and eosin (H&E) staining of each slide was carried out and blindly examined by a veterinary pathologist. Histopathological lesions were scored on a histological grading scale ranging from 0 to 10 based on severity and distribution.

The histopathology of the nasal cavity was scored using very strict criteria. Any finding other than pristine was given a positive score. A single small focus of accumulation of amorphous material and/or the presence of any cell damage no matter how slight was scored as +1 (See FIG. 2C). More than one focus of accumulation of material and/or cell damage was scored as +2 (See FIG. 2D). More than 3 foci of accumulation of material and/or cell damage or multiple locally extensive areas of pathology were scored as +3. The lesions graded as +4 to +6 were associated with increasing severity and more extensive distribution of lesions including the presence of lesions in more than one section. These lesions could be associated with morbidity. The +7 and above had increasing degrees of inflammation. Mortality would be given a score of +10.

Hematological and serum biochemical profile analysis. Whole blood samples were collected from rats and guinea pigs 2 weeks following the final vaccine dose. Dogs were phlebotomized every 14 days and at the study termination at day 43. A portion of the blood was placed in VACUTAINER tubes containing EDTA (BECTON DICKENSON) and a portion was placed in serum separator VACUTAINER tubes (BECTON DICKENSON). Anti-coagulated blood was processed to determine hematological parameters (lymphocytes, monocytes, eosinophils, basophils, red blood cells, hemoglobin , hematocrit, mean corpuscular hemoglobin, mean corpuscular volume, mean corpuscular hemoglobin concentration, and platelets) in a HEMAVET 950 hematology analyzer (Drew Scientific, Inc., Oxford, CT) in accordance to manufacturer's recommendation. Hematological data was compared to species laboratory reference values as established by the Animal Diagnostic Laboratory at the University of Michigan.

Serum samples were analyzed using a VETTEST Chemistry ANALYZER (IDEXX,

Westbrook, Maine). A complete chemistry panel including albumin, alkaline phosphatase, alanine aminotransferase, amylase, aspartate aminotransferase, total calcium, total cholesterol, creatinine, glucose, phosphorous, total bilirubin, total protein, blood urea nitrogen, sodium, potassium, chloride, globulin, and creatine kinase was performed.

5 Biochemical data was compared to species laboratory reference values as established by the Animal Diagnostic Laboratory at the University of Michigan.

Statistical Analysis. Results are expressed as mean $\pm$ standard error of the mean (SEM) or $\pm$ standard deviation (SD). Statistical significance was determined by ANOVA (analysis of variance) using the Student *t* and Fisher exact tests or a Bonferroni's Multiple
10 comparison analysis. The analyses were done with 95% confidence limits and two-tailed tests. A *p* value <0.05 was considered to be statistically significant.

Example 2

Characterization of vaccine formulation

15 A hepatitis B vaccine was formulated utilizing two components; recombinant HBsAg and NE (HBsAg-NE). The formulation was characterized by evaluating the stability of its components, as well as the physical interaction of the antigen with NE. The lipid droplet size was stable and uniform in both concentrations of antigen tested (the average size for all conditions calculated as 349 ± 17 nm), and droplet size of the mixture was not altered by
20 either temperature or NE concentration (See FIG. 3).

HBsAg integrity in the emulsion was evaluated using SDS-PAGE and Western blot (See FIG. 4). NE also did not interfere with the electrophoresis or immunoblotting procedures. After treatment with SDS, HBsAg protein migrated as a band that corresponded to HBsAg monomer ($M_w \approx 24$ kDa) with a minor fraction at twice this molecular weight
25 representing dimer, and this pattern was not altered by prior mixing in NE. In addition, antigenic recognition was retained in HBsAg mixed in NE as identified in Western Blots using a polyclonal goat antiserum raised to native HBsAg (Figure 4). No degradation products of HBsAg were detected in either analysis, and no significant aggregation was apparent during mixing or incubation with NE.

30 The surface charge of the vaccine formulation was determined by measuring the zeta potential and was compared to NE and HBsAg solutions. In either deionized water or PBS buffer as a diluent, HBsAg had negative zeta potential (See FIG. 5A)(Park, MH, et al. Tissue

Antigens, 2003. 62: p. 505-511). In contrast, there was a decrease in the positive zeta potential of the NE after mixing with the HBsAg. This suggests an electrostatic association between the negatively charged HBsAg particles and cationic CPC-containing emulsion (Talaro, K. Foundations in Microbiology. 1993, Iowa: WmC Brown Communications, Inc.).

5 The drop in charge of the emulsion was more pronounced when the HBsAg/NE formulation was made with deionized water as compared to PBS (See FIG. 5).

The interaction of HBsAg with NE was further examined using laser diffraction particle sizing and isothermal titration calorimetry (ITC). Two independent and differently sized peaks for NE and HBsAg were observed before mixing, however after formulation only
10 a single peak was detected with a dynamic diameter of ~300 nm (See FIGS. 3 and 5B-D). The absence of two separate peaks again indicated an association between the lipid phase and HBsAg protein, and suggested that no significant fraction of the antigen remained independent from the lipid in the aqueous phase of NE. Thermodynamic analysis of the interaction between the HBsAg and the NE using ITC showed a spontaneous exothermic
15 reaction with a calculated change in heat capacity of binding (ΔC_p) of -1.44 indicating an energetically favorable interaction (See FIG. 5E).

Example 3

Immunogenicity of the nasal HBsAg-NE vaccine

20 Immunogenicity of the HBsAg-NE vaccine formulation was tested by conducting *in vivo* adjuvant and antigen dose escalation studies. After a single immunization with 20 μ g of HBsAg in 5- 40% NE similar end-point, serum anti-HBsAg IgG titers averaging over 10^4 were achieved (See FIG. 6A). In contrast, lower serum titers ($<10^2$) were generated after immunization with 5% NE and low, inconsistent antibody responses were detected in mice
25 nasally vaccinated with HBsAg in PBS (See FIG. 6A). Booster immunization at six weeks caused the serum anti-HBsAg IgG titers to increase over 10 fold in all groups except in the animal immunized with HBsAg in PBS where no effect was observed. The highest anti-HBsAg antibody endpoint titers, exceeded 10^6 at 6 to 8 weeks after boost, were achieved when the animals were vaccinated with either 20% or 40% NE. The HBsAg-NE vaccine also
30 produced persistent antibody responses with serum anti-HBsAg IgG titers of 10^4 - 10^5 at 6 months after initial vaccination regardless of the concentration of NE used for vaccination. Thus, 20% was determined to be sufficient NE concentration.

To optimize the antigen concentration in the NE vaccine, mice were i.n. immunized with 1- 40 µg of HBsAg mixed with 20% NE (See FIG. 6B). After a single vaccination anti IgG HBsAg antibody responses showed a dose dependent relationship with highest titers in the 20 µg HBsAg-NE group and significantly weaker antibody responses in mice vaccinated with 1 µg of HBsAg. After a second immunization at six weeks, the anti-HBsAg IgG titers increased approximately 10 fold exceeding 10^4 , except in animals immunized with 1 µg HBsAg in NE. Intranasal immunizations with equivalent amounts of HBsAg mixed in PBS again produced only sporadic and weak antibody responses with titers less than 10^2 . This indicated that 20 µg of HBsAg was a sufficient antigen dose.

Example 4

Immunogenicity of HBsAg-NE immunization

The humoral and cell-mediated immune responses to the optimized HBsAg-NE vaccine were characterized *in vivo* in mice. Intranasal vaccination with 20 µg HBsAg-20% NE or i.m. injection of 20 µg HBsAg-Alu resulted in comparable, high levels of anti-HBsAg serum IgG antibodies reaching 10^5 to 10^6 titers within 8 weeks after primary vaccination (See FIG. 7A). Both HBsAg-NE and HBsAg-Alu vaccines produced equivalent, durable immune responses with serum anti-HBsAg IgG end point titers of 10^4 to 10^5 being maintained up to 6 months after vaccination. Nasal vaccination with HBsAg-NE elicited serum titers in mice that when normalized with standardized human anti-HBsAg serum indicated an antibody index ≥ 1000 IU/ml. This index is compatible with protective immunity in humans.

Analysis of serum IgG anti-HBsAg avidity at 23 weeks indicated significantly higher antibody avidity in HBsAg-NE immunized animals as compared to IgG from HBsAg-Alu vaccinated mice (*p value* = 0.034) (See FIG. 7B). While the overall titers were equivalent, analysis of serum IgG subclass indicated that i.n. HBsAg-NE vaccination produced anti-HBsAg IgG with a prevalence of IgG2b (and IgG2a) over IgG1 subclass antibodies, while the HBsAg-Alu vaccine produced mainly IgG1 subclass antibodies (See FIG. 7C). This demonstrates a Th1 response to the NE-based vaccine vs. the traditional Th2 response associated with alum. Immunization with HBsAg-NE composed of *adw* serotype surface antigen also produced cross-reacting IgG antibodies against the heterologous *ayw* serotype.

Mucosal immune responses were characterized in bronchioalveolar lavage (BAL) fluid of immunized animals. HBsAg specific IgA and IgG antibodies were detected in BAL

samples obtained 23 weeks after initial immunization, from mice immunized intranasally with HBsAg-NE (See FIG. 57A and B, respectively). These animals also had detectable serum levels of IgA anti-HBsAg. No anti-HBsAg antibodies were detected in BALs or serum in mice immunized with antigen administered in PBS or in intramuscularly immunized mice despite high serum titers.

HBsAg specific cellular responses were characterized in splenocytes of immunized animals obtained at 18 weeks after last immunization. The cells were stimulated with HBsAg and then evaluated for specific cytokine production (See FIG. 8C). The cytokine expression pattern included high production of the Th1-type cytokines IFN- γ and TNF- α (ranging from 5 to 40 fold) and lower increases (≤ 2 fold) in the expression of Th2-type cytokines IL-4, IL-5 and IL-10. This pattern of expression demonstrated a Th1 bias of cell-mediated response.

The serum IgG response elicited by HBsAg-NE vaccine was also studied in two alternative rodent species to ensure that the immunization effect was not species specific. Rats and guinea pigs were immunized with 5 μ g and 20 μ g doses of HBsAg mixed with 20% NE (See FIG. 9). After a single vaccination, animals showed a dose dependent response with the highest IgG antibody titers in the 20 μ g HBsAg-NE group. After a second administration at five weeks, the anti-HBsAg IgG titers increased up to 100 and 1000 fold surpassing 10^5 titers in both species. Thus, the HBsAg-NE vaccine proved to be immunogenic in all three animal species tested.

Example 5

Thermal stability of HBsAg-NE vaccine

HBsAg-NE was evaluated for thermal stability at three test temperatures. At 6 weeks, 3 months, 6 months and a year after the start of the stability study aliquots of the formulation were evaluated for physical stability *in vitro* and immunogenicity *in vivo*.

HBsAg stability in vaccine samples was analyzed by SDS-PAGE and antigenicity evaluated with Western blots (See FIG. 10) with the stored samples compared to freshly mixed vaccine at each time point. The protein stains and Western Blots of HBsAg at 6 weeks and 3 months were not different from fresh material and there were no low molecular weight degradation products appreciable at these time points (See FIG. 10A and B). After 6 months of storage (See FIG. 10B), however, the major HBsAg band was not detectable in the 40°C by silver staining or immunoblotting, whereas both 4°C and 25°C stored products were still

comparable to freshly mixed vaccine. After 1 year of storage (See FIG. 10C), the 25°C sample was also degraded, while the 4°C stored formulation was intact and comparable to freshly mixed vaccine. The stability of the NE also was evaluated by particle size characterization (See FIG. 10D). The mean diameter ($\pm$ SD) of freshly mixed HBsAg-NE samples was $0.323 \pm 0.016 \mu\text{M}$, and there were no significant differences between NE particle sizes of fresh and stored HBsAg-NE samples at any temperature or time point.

Immunogenicity of the vaccine in CD-1 mice was tested at each time point and storage temperature. Mice were immunized then boosted at six weeks post-vaccination, and anti-HBsAg serum IgG responses were determined at 2, 3, 5, 8, 10 and 12 weeks after primary vaccination. There were no significant differences in serum IgG titers elicited by HBsAg-NE vaccine stored at any temperature up to 3 months (See FIG. 11A and B). At 6 months of storage, HBsAg-NE stored at 40°C could elicit and boost HBsAg-specific antibodies, but at a significantly decreased titer when compared to freshly mixed vaccine, while 4°C and 25°C stored vaccines retained complete immunogenicity (See FIG. 11C).

After 1 year of storage, 25°C and 40°C stored HBsAg-NE elicited decreased serum IgG while the 4°C stored and freshly mixed vaccines again retained complete immunogenicity (See FIG. 60D). This demonstrated that the vaccine retained immunogenicity for 3 months at 40°C and 6 months at 25°C.

Example 6

Evaluation of the safety of NE adjuvant and HBsAg-NE vaccine

Evaluation of acute and (sub) chronic toxic effect of NE and HBsAg-NE formulations was performed in rodent models and in dogs. Multiple intranasal dose studies (See FIG. 1) for NE adjuvant or HBsAg-NE were conducted. No statistically significant changes in subcutaneous temperature or body weight were observed as compared to non-treated control groups. Likewise, no changes in activity or appetite were noted throughout the study. Hematological and serum biochemical results in rats, guinea pigs, and dogs were within normal physiological range (See FIG. 1). No lesions were reported in highly perfused organs including the olfactory bulb and frontal lobe of the brain. Cytotoxicity was not observed in nasal epithelium and other exposed tissues. The only histological lesion noted was the accumulation of amorphous material that sometimes contained cellular debris from sloughed nasal epithelial cells. None of the lesions were of clinical significance (See FIGS. 1 and 2).

Both NE and HBsAg-NE were safe and well tolerated by all animal species tested. Approximately 5% of mice developed nasal obstruction with the emulsion, but this was not observed in larger animals and determined to be related to the unique nasal anatomy of the mouse.

5

Example 7

Evaluation of HBsAg preparations

Experiments were conducted during development of embodiments of the invention in order to characterize qualitative variations in different lots of hepatitis B surface antigen (HBsAg) (e.g., supplied by Indian Immunologicals (ILL), LTD Hyderabad, India). UV spectroscopy and HPLC demonstrated fundamental differences in the protein preparation between different lots of HBsAg (e.g., supplied by ILL). It was determined that manufacturing the HBsAg stock at relatively high concentrations promoted HBsAg aggregation. A less concentrated lot of HBsAg was obtained (0.135 mg/ml, Lot G) from ILL. The lot was characterized as having a similar range of endotoxin content to other previously acquired lots. Additionally, it was shown to have *in vitro* potency comparable to commercial HBsAg produced by Aldevron (Fargo, ND), and the protein was present in a virus-like particle. Studies were conducted to examine the quality of the different HBsAg lots and the propensity of the antigen to self-aggregate.

To determine if the different lots of HBsAg self-aggregated (Lot G versus other more concentrated lots), they were sized by quasi-elastic light scattering with a Malvern ZETASIZER ZS laser diffraction particle sizer. The expected theoretical distribution for HBsAg is a single narrow peak occurring at 28 nm. As shown in Figure 12A, only a single peak maximizing at 28 nm was observed for Lot D. Two distinct peaks were observed for Lot F (28 nm and 1050 nm) (Figure 12B). However, Lot G demonstrated a significant shoulder with a broad distribution of sizes suggesting marked aggregation (Figure 12C).

To gain insight into the 3-D structure of the aggregates, the different lots of ILL HBsAg were imaged using atomic force microscopy (AFM) (Figures 12D through 12F). The protein complex radial size distributions were calculated as shown in Figure 12G. From these data, it is apparent that self-aggregation has occurred in the last two lots of HBsAg received from ILL (Lots F and G).

Although an understanding of a mechanism is not necessary to practice the present

invention and the present invention is not limited to any particular mechanism, in some embodiments, the biological relevance of using aggregated protein in an immunogenic composition (e.g., vaccine product) is important because immunity is thought to be related to the physical structure and solubility of antigen and its interaction with the host (e.g., at the mucosal inductive site). Thus, in some embodiments, aggregation is a contributing factor to changes in immune response (e.g., by limiting the epithelial uptake). An *in vivo* potency assay in guinea pigs using Lot F demonstrated immunogenicity of the vaccine product in a formulation containing 40 µg of HBsAg and 20% NE. Using 20 µg or 5 µg of HBsAg with 20% NE did not produce as robust a immune response in the guinea pigs. Studies were therefore designed to examine the relationship of concentration of NE to immunogenicity and the effects of HBsAg aggregation in the rat species.

For these studies, 10 week old female Sprague-Dawley Rats were intranasally vaccinated with either 10 or 40 µg HBsAg (Lot F) in NE ranging from 1% to 20%. The vaccination was administered on a prime and 4 week boost schedule. Serum anti-HBsAg was measured using ELISA. As shown in Figure 13, immunogenicity was observed for 1%, 10%, and 20% NE-based vaccinations.

As shown in Figure 14, it was possible to disrupt the protein aggregates via dialyzing the surface antigen to PBS and then using ultrasonication. Immunogenic effects of immunogenic compositions comprises a variety of aggregation states that can be tested by intranasally vaccinating rats using the dialyzed and sonicated HBsAg according to methods described herein. Accordingly, in some embodiments, the present invention provides immunogenic compositions comprising one or more aggregation states (e.g., marked aggregation, moderate aggregation, little to no aggregation) of one or more hepatitis B antigens (e.g., HBsAg), methods of characterizing the physical and biological characteristics of the immunogenic compositions, methods of correlating the immunogenic composition aggregation state with the ability to induce immune responses, as well as methods of using the same to induce immune responses in a host administered the same.

EXAMPLE 8

Compositions comprising a trivalent nanoemulsion-based vaccine against *Bacillus anthracis*, *Yersinia pestis*, and *Clostridium botulinum* and methods of using the same

Materials and methods

Vaccination. Anesthetized CD-1 mice were immunized with two nasal administrations of either monovalent or with a trivalent vaccine composed of the combination of recombinant proteins (20 µg of each antigen): PA (protective antigen) of *Bacillus anthracis*, rLcrV (or LcrV10) protein of *Yersinia. pestis* and rHCR/A1 (fragment of *Clostridium botulinum* neurotoxin) mixed with 20% W₈₀5EC (10ul). Control immunizations were performed with antigens suspended in PBS, or with intramuscular injection of antigens adsorbed onto aluminum hydroxide adjuvant (alum, 0.5µg/ml, 50ul).

Antibody response. To analyze humoral immunity, serum samples were collected at 2-4 weeks intervals over a course of time. Bronchoalveolar lavage fluid (BAL), spleen and cardiac blood were collected at the termination of animals. Serum and BAL antigen specific IgG and secretory IgA titers as well as IgG subclass determination were performed using ELISA.

Cellular immunity. To assess cellular responses, splenocytes were isolated from immunized mice and stimulated in vitro with specific antigen for 72 hours days. Antigen specific cytokine production was assayed in culture supernatants using specific cytokine detection kit (R&D) or Lumiex assays.

Antigens and adjuvant. HCR/A1 and LcrV provided non-commercially by Dr. Joseph Barbieri, Medical College of Wisconsin, and Dr. Olaf Schneewind, University of Chicago, respectively. PA was purchased from BEI Resources, Manassas, VA. W805EC was provided by the NanoBio Corporation, Ann Arbor, MI. Polyacrylamide gel electrophoresis (PAGE) was performed on each antigen prior to mixing with nanoemulsion (See Figure 15). Each antigen appeared to be intact and without contamination.

Immunogenic composition formulation. A trivalent, nanoemulsion-adjuvanted vaccine was designed for nasal administration. Immune responses in mice were evaluated after intranasal (i.n) immunization with either monovalent or with a trivalent vaccine composed of the combination of recombinant proteins: PA (protective antigen) of *B. anthracis*, rHCR/A1 (fragment of *C. botulinum* neurotoxin) and rLcrV (or LcrV10) protein of *Y. pestis* administered with an oil-in-water nanoemulsion (NE, W₈₀5EC formulation) adjuvant. CD-1 mice were immunized with two IN administrations of either trivalent vaccine comprised of 20 µg of each recombinant antigen or with a monovalent vaccine containing a single antigen mixed with 20% NE adjuvant. For intramuscular (IM) immunizations, mice were injected with antigens adsorbed on aluminum hydroxide (alum). For assessment of

antibody response, serum was collected at multiple time points. Spleens and cardiac blood were collected at termination of the experiment at 19 weeks. Serum antigen specific IgG and IgA subclass titers were determined by ELISA. Cellular response was characterized *in vitro* by the antigen-specific cytokine expression in splenocytes from immunized mice

5

Example 9

Characterization of immune response from trivalent nanoemulsion-based mucosal vaccine against *Bacillus anthracis*, *Yersinia pestis*, and *Clostridium botulinum*

The immunogenicity of trivalent vaccines against HCR/A1, PA and either full length LcrV protein (See FIG. 15A) or truncated LcrV10 protein (See FIG. 15B) mixed with NE
10 were tested by the analysis of antigen-specific serum IgG in immunized mice.

Development of immune response after nasal immunization with trivalent vaccine composed of PA/HCR/A1 and full length LcrV protein is shown in Figure 16. Single intranasal (IN) immunization produced rapid and significant IgG response to all antigens. The anti-HCR/A1 response was 10 to 15 fold lower than PA and LcrV antigens. After boost
15 the IgG levels reached over 10^6 titers which were sustained for 5 months without significant decrease in antibody levels. Immunizations with antigens in PBS (without combining with nanoemulsion) did not produce detectable immune responses.

Development of immune response after nasal immunization with trivalent vaccine composed of PA/HCR/A1 and truncated LcrV10 protein is shown in Figure 16B. As with
20 full length LCRV protein, single IN immunization produced rapid significant IgG response to all antigens. Again, the anti-HCR/A1 response was 10 to 15 fold lower than PA and LcrV10 antigens. After boost the IgG levels reached over 10^6 titers which were sustained for 5 months without significant decrease in antibody levels. Immunizations with antigens in PBS (without combining with nanoemulsion) did not produce detectable immune responses.

25 Thus, the present invention provides that both vaccine formulations were immunogenic and produced high anti-PA and anti-LcrV and LcrV10 antibody response surpassing 10^6 titers. Antibody response to HCR/A1 protein was consistently over 10 fold lower than response to PA, LcrV, and LcrV10 even in increased concentration of antigen .

Antigen-specific immune responses in trivalent versus monovalent vaccines is shown
30 in Figure 17. To determine if combining antigens affect their immunogenicity the efficacy of trivalent and monovalent vaccines was analyzed. Comparison of IgG titers in trivalent versus monovalent vaccines indicated that combining the antigens did not affect their immunogenicity

(e.g., either by combining the plurality of antigens with nanoemulsion or by suppressing the specific immune response generated by each antigen). Thus, the present invention provides that immune responses produced by trivalent and monovalent NE-based nasal vaccines demonstrated that combining all three proteins does not affect immunogenicity of any of the individual antigens (See Figure 17). In each case the trivalent vaccine was capable of producing nearly identical antigen specific titer level to the corresponding monovalent vaccine, demonstrating that trivalency does not reduce the immunogenicity of the individual antigens.

Immunogenicity of nasal NE-adjuvanted trivalent vaccine was compared to that of intramuscular injected alum-based vaccine by measurement of serum antigen specific titers 19 weeks after prime. The NE-adjuvanted trivalent vaccines produced immune responses comparable to intramuscular injection of conventional alum-based vaccines for each antigen tested (See Figure 18). The present invention provides that, based upon the comparison of IgG titers elicited by NE-based vaccines with conventional alum-adjuvanted vaccines, mucosal vaccination produced antibody titers comparable with intramuscular vaccines.

Figure 19 shows that nasal immunization with both trivalent vaccines produced mucosal immunity. The antigen-specific secretory IgA and IgG were detected in Bronchoalveolar lavage fluid (BAL) from mice obtained up to 5 months after vaccination. Nasal immunizations with NE-adjuvanted trivalent vaccines were capable of producing antigen-specific mucosal IgG (See Figure 19A) and IgA (See Figure 19B) in brochial secretions to the same degree as the corresponding monovalent vaccines.

In order to analyze the cellular response in mice immunized with trivalent nanoemulsion-based vaccines, cytokine levels were monitored in splenocytes of immunized mice. The pattern of antigen specific Th1- and Th-2 type cytokine expression *in vitro* shows 5 to 16 fold increase of INF- γ , and significantly lower levels of IL-4, without significant effect on other cytokines (See Figure 20). The antigen specific induction of IFN- γ , which is consider a hallmark of TH-1 response and lack of IL-4 indicated that immunization with the trivalent vaccine produced a Th-1 biased cellular response. Neither LcrV10 nor LcrV10 mutant had an effect on IL-10 synthesis. Mock immunization with nanoemulsion alone did not produce cytokine responses in splenic lymphocytes. Thus, the present invention provides that immunization with a trivalent vaccine of the invention induced a mixed Th1- Th2 immune response with a functional bias toward Th-1 cell mediated immunity.

Analysis of IgG subclass in serum after nasal immunizations with a trivalent vaccine of the invention is shown in Figures 21-23. Figure 21 shows that the analysis of IgG subclass in serum after nasal immunizations with NE adjuvanted trivalent vaccines indicated increased level of IgG2a and IgG2b anti-PA subclass antibodies in contrast to aluminum-based IM immunization that had a prevalence of IgG1 subclass of anti-PA IgG. Results presented as ratio of IgG subclass to total IgG titer. The elevated levels of IgG2 subclass antibodies indicated the induction of Th-1 polarization – in contrast to Th-2 type in aluminum based vaccine.

Figure 22 shows anti-LcrV and mutant LcrV10 IgG subclass pattern. Distribution of IgG1, IgG2a and IgG2b in mice immunized with nasal administration of trivalent [HCR/A1-LcrV-PA]-NE or [HCR/A1-LcrV10-PA]-NE, nasal monovalent LcrV-NE or LcrV10-NE and intramuscular injection of aluminum-adsorbed LcrV or LcrV10. Nasal immunizations with NE adjuvant produced increased level of IgG2a and IgG2b subclass antibodies to both full length and deletion mutant of LcrV antigen in comparison to aluminum-based IM immunization with prevalence of IgG1 subclass of anti-LcrV and V10 IgG. Results presented as ratio of IgG subclass to total IgG titer.

Figure 23 shows that similar IgG subclass patterns were detected for HCR/A1 botulism toxin in both LcrV and V10 protein mixes. Results presented as ration of IgG subclass to total IgG titer.

Mucosal immunization with NE-based vaccine produced antibody repertoire capable of recognizing LcrV epitopes similar to the natural pattern induced by surviving infection with *Y. pestis*. Loss of response to deleted regions of the full length protein resulting from immunization with the LcrV10 antigen was observed. Mucosal immunization with NE-based vaccines (trivalent or monovalent) produced antibodies capable of recognizing multiple regions of the PA protein. Frequency of the epitope recognition suggests binding to the pore formation regions of domain II and significant recognition of domains III and IV, especially enhanced in the receptor binding domain IV. LF/EF binding epitopes were recognized more frequently with serum from the monovalent rPA-NE vaccine. Thus, the present invention provides that NE based vaccine formulations maintain recognition of critical epitopes for protective immunity.

Nasal immunization with NE-based trivalent vaccine produced protective immunity against challenge with live pathogens (See Figures 24-26). For example, as shown in Figure

10, vaccination with 20 µg of either LcrV or LcrV10 protein in 20% NE produced 90% protection against 150 X LD50 challenge with a virulent strain of plague and is comparable to intramuscular vaccination based on alum. Using lower concentration of NE adjuvant produced lower protective immunity- 70%. Figure 25 shows that vaccination with lower
5 doses containing 10 µg of either protein in 20% NE produced 65 to 75%, while using 5% of NE resulted in 50 to 60% survival. Thus, the present invention provides a trivalent vaccine that provides protective immunity against challenge with the plague causing pathogen *Yersinia pestis*.

Nasal immunization with NE-based trivalent vaccine produced protective immunity
10 against challenge with anthrax spores. Guinea pigs were immunized with 10 to 100 µg doses of PA protein and intranasally challenged with 100x LD50 of anthrax Ames spores. As shown in Figure 26, survival in all groups was 70% - but higher doses of antigen produced increased time to death. Thus, the present invention provides a trivalent vaccine that provides protective immunity against inhalation challenge with anthrax spores.

15 While specific embodiments of the invention have been described and illustrated, such embodiments should be considered illustrative of the invention only and not as limiting the invention as construed in accordance with the accompanying claims.

CLAIMS:

1. An immunogenic composition comprising a nanoemulsion and hepatitis B virus (HBV) immunogen, wherein said immunogen comprises HBV surface antigen (HBsAg), and wherein the immunogenic composition induces HBV-specific humoral and cell-mediated immune responses that prevent or treat disease associated with HBV.
2. The immunogenic composition of claim 1, wherein:
 - said HBsAg is in particle form; or,
 - said HBsAg is treated with dialysis or sonication prior to combining with said nanoemulsion; or,
 - said composition does not contain a mercury based preservative; or,
 - said nanoemulsion and HBsAg skews a host immune response toward a Th1 type immune response; or,
 - said nanoemulsion comprises an aqueous phase, an oil phase, and a solvent, wherein said nanoemulsion comprises a polysorbate surfactant, ethanol, cetylpyridinium chloride (CPC), of soybean oil, and water; or,
 - a dose of said composition comprises between 10 and 100 μ g of HBsAg; or,
 - a dose of said composition comprises 20% nanoemulsion; or,
 - said HBsAg is heat stable in said nanoemulsion.
3. The immunogenic composition of claim 1 or claim 2 for use in preventing or treating HBV.
4. The immunogenic composition of claim 1 or claim 2 for use in formulating a vaccine for the prevention or treatment of HBV.
5. Use of the immunogenic composition of claim 1 or claim 2 for preventing or treating HBV.
6. Use of the immunogenic composition of claim 1 or claim 2 for formulating a vaccine for the prevention or treatment of HBV.

7. A vaccine comprising the immunogenic composition of claim 1 or claim 2 and a pharmaceutically acceptable carrier.
8. The vaccine of claim 7 for use in preventing or treating HBV infections.
9. The vaccine of claim 7 or claim 8, wherein the vaccine is formulated for nasal mucosal administration.
10. The vaccine of any one of claims 7-9, wherein the HBV-specific humoral and cell-mediated immune responses comprise a Th1 mediated immune response comprising enhanced expression of IFN-[gamma] and TNF-[alpha], or a systemic IgG response to HBV, or a mucosal IgA response to HBV.
11. The vaccine of any one of claims 7-10, wherein the HBV-specific humoral and cell-mediated immune responses decrease or prevent morbidity caused by exposure to HBV, or onset of mortality caused by exposure to HBV.
12. Use of the vaccine of any one of claims 7 and 9-11 for treating or preventing HBV.

Figure 1

Species/Strain ^a	Treatment		Number of Dose	Dose Volume (ul)	Group Average Histopathological Score ^b				Metabolic Analysis ^d
	Adjuvant SO-NE (%)	HBAg Dose (ug)			Nasal	Pulmon .	Brain	Other ^c	
Mouse/CD-1	0	20	2 ^e	10	0	0	0	0	n/a
	1	20	2 ^e	10	0	0	0	0	n/a
	5	20	2 ^e	10	1.0±0.9	0	0	0	n/a
	10	20	2 ^e	10	0.7±1.1	0	0	0	n/a
	20	20	2 ^e	10	1.4±1.3	0	0	0	n/a
	20	0	2 ^e	10	1.1±1.7	0	0	0	n/a
Mouse/BALB/c	20	0	4 ^f	6	1.2±0.4	0	0	0	n/a
	20	0	7 ^g	6	2.0±1.0	0	0	0	n/a
Rat/Wistar	20	32	3 ^h	20	0	0	0	0	Normal
Guinea Pig/Hartley	20	32	3 ^h	20	0	0	0	0	Normal
Canine/Beagle	20	0	3 ^e	200	0	n/a	n/a	n/a	Normal
	20	0	3 ^e	400	0	n/a	n/a	n/a	Normal

a- The number of animals used for analysis: CD-1 (n=10), BALB/c, Wistar rats and Hartley guinea pigs (n=5) and Beagles (n=1) per group.

b- Histological lesions were evaluated on a scale from 0 to 10 with +1 single microscopic focus, +2 at least 2 microscopic foci, +3 more than 3 foci or multiple locally extensive areas of pathology, +4 to +6 were associated increasing severity and more extensive distribution (these lesions could be associated with morbidity), +7 and above had increasing degrees of inflammation (+10 associated with mortality).

c- Other tissues evaluated include heart, liver, kidneys, spleen, esophagus, trachea, stomach, intestines, pancreas, and adrenals.

d- Metabolic analysis evaluated by standard biochemical serum profile analysis on a IDEXX Vet Test Analyzer™ and performed at the Animal Diagnostic Laboratory through the Unit for Laboratory Animal Medicine at the University of Michigan. Normal indicates all analytes fell within normal expected distributions per species.

e- Administered every 2 weeks.

f- Administered every 15 minutes.

g- Administered every 4 hours.

h- Administered every 4 weeks.

Figure 2




Figure 3

Particle Size	
Temperature Condition (°C)	
Sample description	Fresh
1% W ₈₀ 5EC	355 (+/- 130)
20% W ₈₀ 5EC	368 (+/- 255)
40% W ₈₀ 5EC	331 (+/- 154)
20% W ₈₀ 5EC + 0.5 mg/ml HBsAg	373 (+/- 229)
20% W ₈₀ 5EC + 2.5 mg/ml HBsAg	341 (+/- 143)

Average particle size 349 (+/- 17); Mean (+/- SD).

Figure 4




Figure 5




Figure 5 (Cont.)





Figure 6




Figure 7




FIGURE 7 CONTINUED

C




Figure 8




FIGURE 8 CONTINUED




Figure 9




Figure 10




Figure 11




FIGURE 11 CONTINUED




FIGURE 11 CONTINUED




FIGURE 11 CONTINUED

D.





FIGURE 12

A



B





FIGURE 12 CONTINUED

C



D





FIGURE 12 CONTINUED

E



F




FIGURE 12 CONTINUED

G




FIGURE 13




FIGURE 14




FIGURE 15

**PAGE analysis of recombinant PA, HCR/A,
LcrV and LcrV10 proteins**




FIGURE 16

A

**Time course of serum antibody response in mice
intranasally immunized with trivalent vaccine
containing LcrV10 antigen**




FIGURE 16 CONTINUED

B

**Time course of serum antibody response in mice
intranasally immunized with trivalent vaccine
containing LcrV antigen**




FIGURE 17




FIGURE 17 CONTINUED




FIGURE 18





FIGURE 19

A



B




Figure 20





FIGURE 20 CONTINUED




FIGURE 21




Trivalent-NE vaccine**Monovalent-NE vaccine****Aluminum-based injectible vaccine**

FIGURE 22




FIGURE 23




FIGURE 24

Intranasal challenge with ~150 LD₅₀ of *Yersinia pestis* C092

20 µg antigen dose




FIGURE 25




FIGURE 26

