

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2017249988 C1

(54) Title
Combination therapy with notch and PI3K/mTOR inhibitors for use in treating cancer

(51) International Patent Classification(s)
A61K 31/444 (2006.01) **A61P 35/00** (2006.01)
A61K 31/55 (2006.01) **A61P 35/02** (2006.01)

(21) Application No: **2017249988** (22) Date of Filing: **2017.04.05**

(87) WIPO No: **WO17/180385**

(30) Priority Data

(31) Number (32) Date (33) Country
62/321,307 **2016.04.12** **US**

(43) Publication Date: **2017.10.19**
(44) Accepted Journal Date: **2022.10.27**
(44) Amended Journal Date: **2023.02.16**

(71) Applicant(s)
Eli Lilly and Company

(72) Inventor(s)
Patel, Bharvin Kumar;Smith, Michele C.

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
WO 2013/016081 A1
WO 2012/097039 A1
BENDER, MH "Novel inhibitor of Notch signaling for the treatment of cancer",
CANCER RES APRIL 15, 2013 73, XP055277786, Retrieved from the Internet
[retrieved on 20211206]
CN 102085372 A

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



WIPO | PCT



(10) International Publication Number

WO 2017/180385 A1

(43) International Publication Date

19 October 2017 (19.10.2017)

(51) International Patent Classification:

A61K 31/44 (2006.01) A61P 35/00 (2006.01)
A61K 31/55 (2006.01) A61P 35/02 (2006.01)

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2017/026099

(22) International Filing Date:

5 April 2017 (05.04.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/321,307 12 April 2016 (12.04.2016) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))



WO 2017/180385 A1

(54) Title: COMBINATION THERAPY WITH NOTCH AND PI3K/MTOR INHIBITORS FOR USE IN TREATING CANCER

(57) Abstract: Medicaments for use in treating T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma, colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer in a patient comprising combination therapy with 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxoethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, and 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

COMBINATION THERAPY WITH NOTCH AND PI3K/MTOR INHIBITORS FOR USE IN TREATING CANCER

The present invention relates to cancer therapy with 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof (Compound A) and 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof (Compound B) and to methods of using combinations to treat cancer.

Cancers of the ovaries, fallopian tubes, and primary peritoneum are the fifth leading cause of cancer death in women in the U.S. and are generally combined under the phrase “ovarian cancer.” These cancers are often found at advanced stages. This is partly because they may not cause early signs or symptoms and there are no good screening tests for them. The most common type of ovarian cancer is called ovarian epithelial cancer. It begins in the tissue that covers the ovaries. Cancer sometimes begins at the end of the fallopian tube near the ovary and spreads to the ovary. Cancer can also begin in the peritoneum and spread to the ovary. The stages and treatment are generally the same for ovarian epithelial, fallopian tube, and primary peritoneal cancers.

4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, is a Notch pathway signaling inhibitor compound. Notch signaling plays an important role during development and tissue homeostasis. Dysregulation of Notch signaling due to mutation, amplification, or overexpression of ligands and/or receptors, is implicated in a number of malignancies. Inhibition of Notch signaling is a potential target for the development of cancer therapeutics. Compound A and methods of making and using this compound, including for the treatment of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, erythroleukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, colorectal cancer, head and neck cancer, cervical cancer, prostate

-2-

cancer, liver cancer, squamous cell carcinoma (oral), skin cancer and medulloblastoma are disclosed in WO 2013/016081. Compound A is being investigated in a phase 1 clinical trial and expansion cohorts having a defined molecular pathway alteration, or a tissue based malignant tumor, and in a clinical trial in patients with T-cell acute lymphoblastic leukemia 5 or T-cell lymphoblastic lymphoma (T-ALL/T-LBL).

8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof, is a dual inhibitor of phosphoinositide 3-kinases (PI3 Kinase; PI3K) and mammalian target of rapamycin (mTOR). The PI3K/mTOR pathway is stimulated by a 10 variety of growth factors and their receptors and regulates cell metabolism, cell growth, cell survival, cell proliferation, and cell motility. The PI3K/mTOR pathway is thought to be one of the most frequently mutated pathways in cancer. Compound B and methods of making and using this compound, including for the treatment of bladder cancer, colon cancer, 15 gastric cancer, head and neck cancer, non-small-cell lung cancer (NSCLC), breast cancer, melanoma, ovarian cancer, pancreatic cancer, glioblastoma, prostate cancer, lung cancer, renal cancer, sarcoma, hematopoietic and lymphoid tissue cancer, cervical cancer, endometrial cancer, liver cancer, skin cancer, stomach cancer, thyroid cancer, upper digestive tract cancer, and urinary cancer are disclosed in WO 2012/057039. Compound B is being 20 investigated in phase 1 clinical trials and tumor specific expansion cohorts for mesothelioma, breast cancer, and indolent Non-Hodgkin Lymphoma and in clinical trials in patients with squamous non-small cell lung cancer (in combination with necitumumab), endometrial cancer, prostate cancer (in combination with enzalutamide), colorectal cancer (in combination with prexasertib, LY2606368 monomesylate monohydrate; CHK1/2 inhibitor), and non-small cell lung cancer (in combination with abemaciclib; N-[5-(4-ethyl-piperazin-1-ylmethyl)-pyridin-2-yl]-[5-fluoro-4-(7-fluoro-3-isopropyl-2-methyl-3H-benzoimidazol-5-yl)-25 pyrimidin-2-yl]-amine, or a pharmaceutically acceptable salt thereof).

Combinations of a Notch pathway inhibitor and PI3K/AKT/mTOR pathway inhibition have been contemplated in the art, Sheperd et al., *Leukemia*, 2013, 27: 650-660; Gutierrez and Look, *Cancer Cell*, 2007, 12: 411-413; and Palomero and Ferrando, *Clin.*

Cancer Res., 2008, 14(17): 5314-5317. Despite existing treatment options for patients with cancer, there continues to be a need for new and different therapies affording one or both of enhanced efficacy and lower toxicity.

It is believed the present invention provides beneficial therapeutic effects from the 5 combined activity of Compound A and Compound B against T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, and 10 skin cancer as compared to the therapeutic effects provided by either agent alone.

One aspect of the present invention provides a method of treating T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma 15 (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer in a patient, comprising administering to the patient in need of treatment an effective amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, and an effective amount of 8-[5-(1- 20 hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

A further aspect of the present invention provides a method of treating ovarian cancer in a patient, comprising administering to the patient in need of treatment an effective amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, and an effective amount of 8-[5-(1- 25 hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention provides a method of treating T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma 5 (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer in a patient, comprising administering to the patient in need of such treatment, simultaneously, separately, or sequentially, an effective amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, 10 and an effective amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention provides a method of treating ovarian cancer in a patient, comprising administering to the patient in need of such treatment, 15 simultaneously, separately, or sequentially, an effective amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, and an effective amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a 20 pharmaceutically acceptable salt thereof.

A further aspect of the present invention provides a compound 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof; and a compound 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof; for simultaneous, separate, or sequential use in the treatment of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, 25

pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer.

Another aspect of the present invention provides a compound 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof; and a compound 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof; for simultaneous, separate, or sequential use in the treatment of ovarian cancer.

10 A further aspect of the present invention provides: use of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof for the manufacture of a medicament; and

15 use of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof for the manufacture of a medicament;

for the simultaneous, separate, or sequential treatment of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer.

20 A further aspect of the present invention provides: use of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof for the manufacture of a medicament; and

use of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof for the manufacture of a medicament; for the simultaneous, separate, or sequential treatment of ovarian cancer.

5 Another aspect of the present invention is a commercial package comprising a separate composition of each therapeutic agent, or a single composition comprising a combination of the therapeutic agents of the present invention, together with instructions for simultaneous, separate or sequential administration for use in treating T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, 10 chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer.

15 A still further aspect of the present invention is a commercial package comprising a separate composition of each therapeutic agent, or a single composition comprising a combination of the therapeutic agents of the present invention, together with instructions for simultaneous, separate or sequential administration for use in treating ovarian cancer.

20 The compound 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, (Compound A) has the CAS registry number 142138-81-4. Alternatively, the compound may be named: N-[(1S)-2-[(7S)-6,7-dihydro-5-(2-hydroxyethyl)-6-oxo-5H-pyrido[3,2-a][3]benzazepin-7-yl]amino]-1-methyl-2-oxoethyl]-4,4,4-trifluorobutanamide. Other names may be used to unambiguously identify 25 Compound A.

The compound 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof, (Compound B) has the CAS registry number

1386874-06-1. Alternatively, the compound may be named: 1,3-dihydro-8-[5-(1-hydroxy-1-methylethyl)-3-pyridinyl]-1-[(2S)-2-methoxypropyl]-3-methyl-2H-imidazo[4,5-c]quinolin-2-one. Other names may be used to unambiguously identify Compound B.

As used herein, the term "patient" refers to a mammal, preferably a human.

5 "Therapeutically effective amount" or "effective amount" means the dosage of Compound A, or pharmaceutically acceptable salt or hydrate thereof, or pharmaceutical composition containing Compound A, or pharmaceutically acceptable salt or hydrate thereof, and the dosage of Compound B, or pharmaceutically acceptable salt thereof, or pharmaceutical composition containing Compound B, or pharmaceutically acceptable salt thereof, necessary to inhibit tumor cell growth and eliminate or slow or arrest the progression 10 of the cancer in a patient. Anticipated dosages of Compound A, or a pharmaceutically acceptable salt or hydrate thereof, are in the range of 2.5 mg/patient to 75 mg/patient once per day every other day over a five day period followed by two days without dosing (T.I.W.). Anticipated dosages of Compound B, or a pharmaceutically acceptable salt thereof, 15 are in the range of 75 mg to 250 mg twice per day (B.I.D.) dosing. Preferred dosages of Compound A, or a pharmaceutically acceptable salt or hydrate thereof, are in the range of 5 mg to 50 mg T.I.W and Compound B, or a pharmaceutically acceptable salt thereof, are anticipated to be in the range of 100 mg to 200 mg twice daily. The exact dosage required to treat a patient and the length of treatment time will be determined by a physician in view of 20 the stage and severity of the disease as well as the specific needs and response of the individual patient. The dosing administration may be adjusted to provide a more optimal therapeutic benefit to a patient and to manage or ameliorate any drug related toxicities. Alternative dosing schedules such as once per day (QD), twice per day (B.I.D.), three times a day (T.I.D.); dosing once per day every other day (Q2D); once per day every other day over a 25 five day period followed by two days without dosing (T.I.W.); or every third day (Q3D) may be appropriate for each of Compound A and Compound B.

A combination therapy of the present invention is carried out by administering to a T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous

leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer, preferably an ovarian cancer, patient requiring treatment, an effective amount of Compound A, or a pharmaceutically acceptable salt or hydrate thereof, once per day every other day over five days and two days without dosing each week (7-days) over a 28 day cycle and Compound B, or a pharmaceutically acceptable salt thereof, once per day over a 28 day cycle.

The terms "treatment," "treat," and "treating," are meant to include the full spectrum of intervention for the cancer from which the patient is suffering, such as administration of Compounds A and B to alleviate, slow, stop, or reverse one or more of the symptoms and to delay, stop, or reverse progression of the cancer even if the cancer is not actually eliminated.

Compound A or a pharmaceutically acceptable salt or hydrate thereof, is preferably formulated as a pharmaceutical composition using a pharmaceutically acceptable carrier and administered by a variety of routes. Preferably, such compositions are for oral administration. Compound B, or a pharmaceutically acceptable salt thereof, is preferably formulated as a pharmaceutical composition using a pharmaceutically acceptable carrier and administered by a variety of routes. Preferably, such compositions are for oral administration. Such pharmaceutical compositions and processes for preparing them are well known in the art. *See, for example, HANDBOOK OF PHARMACEUTICAL EXCIPIENTS*, 5th edition, Rowe et al., Eds., Pharmaceutical Press (2006); and *REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY* (Troy, *et al.*, Eds., 21st edition, Lippincott Williams & Wilkins (2006).

Each of Compound A and Compound B are capable of reaction with a number of inorganic and organic counterions to form pharmaceutically acceptable salts. Such pharmaceutically acceptable salts and common methodology for preparing them are well known in the art. *See, for example, P. Stahl, *et al.*, HANDBOOK OF PHARMACEUTICAL SALTS: PROPERTIES, SELECTION AND USE, (VCHA/Wiley-VCH, 2002); S.M. Berge,*

et al., "Pharmaceutical Salts, " *Journal of Pharmaceutical Sciences*, Vol. 66, No. 1, January 1977.

The efficacy of the combination treatment of the invention can be measured by various endpoints commonly used in evaluating cancer treatments, including but not

5 limited to, tumor regression, tumor weight or size shrinkage, time to progression, overall survival, progression free survival, overall response rate, duration of response, inhibition of metastatic spread without tumor regression, and PET/CT imaging.

The terms "combination," "therapeutic combination" and "pharmaceutical combination" refer to either: 1) a fixed dose combination in one dosage unit form; or 2) a 10 non-fixed dose combination, optionally packaged together with instructions for combined administration where the individual therapeutic agents, Compound A, or a pharmaceutically acceptable salt or hydrate thereof, and Compound B, or a pharmaceutically acceptable salt thereof, may be administered independently at the same time or separately within time intervals that allow the therapeutic agents to exert a cooperative effect.

15 The term "simultaneous" administration means the administration of each of Compound A and Compound B to a patient in a single action such as where the two agents are incorporated into a single dosage form for administration (fixed dose combination) and where each of Compound A and Compound B are administered independently at substantially the same time or separately within time intervals that allows Compounds A and 20 B to show a cooperative therapeutic effect.

The term "separate" administration means the administration of each of Compound A and Compound B to a patient from non-fixed dose combination dosage forms simultaneously, substantially concurrently, or sequentially in any order. There may, or may not, be a specified time interval for administration of each Compound.

25 The term "sequential" administration means the administration of each of Compound A and Compound B to a patient from non-fixed (separate) dosage forms in separate actions. The two administration actions may, or may not, be linked by a specified time interval. For example, administering Compound A T.I.W. and administering Compound B once per day.

-10-

The phrase “in combination with” includes the simultaneous, separate, and sequential administration of each of Compound A and Compound B to a cancer patient in need of treatment, particularly an ovarian cancer patient.

The term “co-administration” or “combined administration” encompasses the 5 administration of the therapeutic agents to a single patient, and include treatment regimens in which the agents may be administered by different routes of administration or at different times.

The beneficial action of two therapeutic agents producing an effect in a single patient which is greater than the simple additive effects of each agent administered alone may be 10 calculated, for example, using suitable methods known in the art such as the Sigmoid-Emax equation (Holford and Scheiner, *Clin. Pharmacokinet.*, 1981, 6: 429-453), the equation of Loewe additivity (Loewe and Muischenk, *Arch. Exp. Pathol. Pharmacol.*, 1926, 114: 313-326), the median-effect equation (Chou and Talalay, *Adv. Enzyme Regul.*, 1984, 22: 27-55), and the Bliss Independence method, or known equivalents. Each equation may be applied to 15 experimental data to generate a corresponding graph to aid in assessing the effects of a drug combination as additive, within a biologically relevant range of additive, less than additive, or greater than additive.

Cancer is increasingly recognized as a heterogeneous collection of diseases whose 20 initiation and progression are induced by the aberrant function of one or more genes that regulate DNA repair, genome stability, cell proliferation, cell death, adhesion, angiogenesis, invasion, and metastasis in cell and tissue microenvironments. Variant or aberrant function of the “cancer” genes may result from naturally occurring DNA polymorphism, changes in genome copy number (through amplification, deletion, chromosome loss, or duplication), changes in gene and chromosome structure (through chromosomal translocation, inversion, 25 or other rearrangement that leads to deregulated gene expression), and point mutations.

Cancerous neoplasms may be induced by one aberrant gene function, and maintained by the same aberrant gene function, or maintenance and progression exacerbated by additional aberrant gene functions.

-11-

Beyond the genetic chromosomal aberrations mentioned above, each of the cancers may also include epigenetic modifications of the genome including DNA methylation, genomic imprinting, and histone modification by acetylation, methylation, or phosphorylation. An epigenetic modification may play a role in the induction and/or maintenance of the malignancy.

The nature of cancer, as noted is multifactorial. Under appropriate circumstances, therapeutic agents with different mechanisms of action may be combined. However, only considering a combination of therapeutic agents having different modes of action does not necessarily lead to combinations with advantageous effects. Specific therapeutic agents affording demonstrated beneficial effects (therapeutic effect such as enhanced efficacy and/or lower toxicity) compared with monotherapy of only one of the therapeutic agents is preferred.

The combination of the present invention is believed suitable for the treatment of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma (ACC), colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, or skin cancer, and particularly suitable for the treatment of ovarian cancer patients, who have failed standard therapy. This includes patients having cancer showing resistance to monotherapy or showing resistance to combinations different than the present invention.

The terms “Complete Response” (CR), “Partial Response” (PR), “Progressive Disease” (PD), “Stable Disease” (SD), “Objective Response” (OR) are used consistent with definitions according to RECIST v1.1, Eisenhauer et al., *European Journal of Cancer*, 2009, 45, 228-247.

The term “time to disease progression” (TTP) refers to the time, generally measured in weeks or months, from the time of initial treatment, until the cancer progresses (see RECIST v1.1 definition for progressive disease) which is at

-12-

least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. The appearance of one or more new lesions is also considered 5 progression. Such progression is evaluated by a clinician.

The term “extending TTP” refers to increasing the time to disease progression in a treated patient relative to i) an untreated patient, or ii) a patient treated with less than both of Compound A and Compound B.

10 The term “survival” refers to the patient remaining alive, and includes overall survival as well as progression free survival.

The term, “overall survival” refers to the patient remaining alive for a defined period of time, such as 1 year, 5 years, etc. from the time of diagnosis or treatment.

The term, “progression free survival” refers to the patient remaining alive, without the cancer progressing.

15 As used herein, the term “extending survival” is meant increasing overall or progression free survival in a treated patient relative to i) an untreated patient, ii) a patient treated with less than both of Compound A and Compound B, or iii) a control treatment protocol. Survival is monitored for a defined period of time, such as one month, six months, 1 year, 5 years, or 10 years, etc., following the initiation of treatment or following 20 the initial diagnosis of cancer.

The term “primary tumor” or “primary lesion” is meant the original cancer and not a metastatic tumor or lesion located in another tissue, organ, or location in the patient’s body.

25 In one embodiment, the dose of Compound A is escalated until the Maximum Tolerated Dosage is reached, and Compound B of the present invention is administered with a fixed dose. Alternatively, Compound A may be administered in a fixed dose and the dose of Compound B may be escalated. Each patient may receive doses of Compound A and/or Compound B either daily or intermittently. The efficacy of the treatment may be determined

-13-

in such studies, e.g., after 12, 18 or 24 weeks by evaluation of symptom scores every 6 weeks.

Compound A may be prepared by the procedures described in WO 2013/016081.

5 Compound B may be prepared by the procedures described in WO 2012/057039.

The following Example illustrates the activity of each of Compound A alone, Compound B alone and the combination of Compound A and Compound B.

10 Biological Example 1

A2780 is a human ovarian cell line (Sigma-Aldrich). The cells are grown in culture media at 37°C in 5% CO₂ with humidity in the atmosphere. Cell culture media for A2780 human ovarian carcinoma is RPMI-1640 (without phenol red) with 2.05 mM L-glutamine, 15 0.01 mg/ml insulin and 10% fetal bovine serum (FBS) added.

To evaluate *in vivo* efficacy and effect A2780 (2 x 10⁶) cells in a 1:1 Matrigel® mix (0.2 mL volume) are implanted by subcutaneous injection in the hind leg of 6-8 weeks of age athymic nude female mice (Harlan Laboratories). A total of 7 to 10 mice are used for each group. Just before implantation, animals are irradiated (450 Total Body Irradiation). Mice 20 are fed *ad libitum* on normal chow. Treatment is initiated with oral administration (gavage) of Compound A or vehicle (1% sodium carboxymethylcellulose (Na-CMC) in 0.25% Tween®-80) or Compound B in 1% hydroxyethyl cellulose (HEC)/0.25% Tween® 80/0.05% Antifoam or their vehicle in 0.2 mL volume when tumor size reached to 150 ± 50 mm³, in this case after the tumor measurements were taken on study day 9. Compound A was 25 administered at 4 or 8 mg/kg on a Monday, Wednesday and Friday schedule for 4 weeks and Compound B was administered at 15 or 30 mg/kg daily for 28 days. Tumor growth and body weight are monitored over time to evaluate efficacy and signs of toxicity. Bidimensional measurements of tumors are performed twice a week and tumor volumes are calculated based on the following formula: (Tumor Volume) = [(L) x (W2) x (H/6)] where L is mid-axis

-14-

length and W is mid-axis width. Tumor volume data are transformed to a log scale to equalize variance across time and treatment groups. The log volume data are analyzed with a two-way repeated measures analysis of variance by time and treatment using the MIXED™ procedure in SAS™ software (version 8.2). The correlation model for the repeated measures

5 is spatial power. Least squares means from the repeated measures analysis, anti-logged to the tumor volume scale, are shown in Table 1. P-values for comparing each pair of groups on study day 36 are shown in Table 2. Test Groups are:

01: Control: 1% HEC/0.25% Tween 80/0.05% Antifoam; 1% CMC/0.25% Tween 80/0.05% Antifoam;

10 02: Compound A 8 mg/kg;

03: Compound B 30 mg/kg;

04: rapamycin 4 mg/kg;

05: Compound A 8 mg/kg and Compound B 30 mg/kg;

06: Compound A 4 mg/kg and Compound B 30 mg/kg;

15 07: Compound B 15 mg/kg and Compound A 8 mg/kg;

08: Compound B 15 mg/kg and Compound A 4 mg/kg;

09: Compound A 8 mg/kg and rapamycin 4 mg/kg.

Human A2780 Xenograft

Geometric Mean

Table 1

Group	Study Days						36
	7	9	12	15	19	22	
01	137.56	92.02	134.29	217.68	453.38	928.72	1775.80
02	121.62	109.98	131.53	194.78	408.55	639.48	1140.46
03	149.36	118.70	138.06	201.95	326.33	488.21	776.88
04	130.45	100.08	130.81	215.88	428.88	786.46	1291.64
05	138.91	119.86	129.42	199.60	283.38	357.34	434.30
06	130.01	118.32	135.10	206.49	388.24	412.30	553.37
07	135.16	117.22	130.76	201.63	344.19	502.40	595.27
08	133.14	113.25	126.46	207.38	398.80	588.69	834.86
09	131.74	105.06	125.28	200.25	363.74	565.04	833.21

Table 2
Between Group P-Values (Repeated Measures ANOVA)

Group	Cmpd. A 8 mg/kg	Cmpd. B 30 mg/kg	Rapamycin 4 mg/kg	Cmpd. A 8 mg/kg and Cmpd. B 30 mg/kg	Cmpd. A 4 mg/kg and Cmpd. B 30 mg/kg	Cmpd. A 4 mg/kg and Cmpd. A 8 mg/kg	Rapamycin 4 mg/kg and Cmpd. A 8 mg/kg
Control HEK/CMV	1	< 0.001	< 0.001	0.030	< 0.001	< 0.001	< 0.001
Cmpd. A 8 mg/kg	2		0.874	0.027	< 0.001	< 0.001	0.028
Cmpd. B 30 mg/kg	3			0.038	< 0.001	< 0.001	0.018
Rapamycin 4 mg/kg	4				< 0.001	< 0.001	< 0.001
Cmpd. A 8 mg/kg and Cmpd. B 30 mg/kg	5				0.010	0.026	< 0.001
Cmpd. A 4 mg/kg and Cmpd. B 30 mg/kg	6					0.617	0.022
Cmpd. B 15 mg/kg and Cmpd. A 8 mg/kg	7						0.004
Cmpd. B 15 mg/kg and Cmpd. A 4 mg/kg	8						0.192

-17-

Table 2 shows combinations of Compound A and Compound B, in this test, demonstrated statistically significant tumor growth inhibition results over each of Compound 5 A and Compound B alone (Group 5 vs. either Group 2 or Group 3).

Combination Analysis Method

Using the repeated measures analysis previously described, a contrast statement is used to test for an interaction effect on study day 36, using the two specific treatments that 10 were combined. This test is statistically significant with $p = 0.022$, demonstrating better than additive, or synergistic, activity, since the estimated mean tumor volume in the combination group (358 mm^3) is less than the expected additive tumor volume per the Bliss Independence method ($1616 \times 1664 / 3711 = 725 \text{ mm}^3$).

15

20

25

30

Clinical Evaluation

A study of 4,4,4-Trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-

5 pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide hydrate, Compound A in combination with 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one Compound B in patients with advanced or metastatic solid tumors.

Study Design

10 This study is a multicenter, nonrandomized, open-label study consisting of a dose escalation phase in patients with advanced/metastatic cancer from a variety of solid tumors followed by a dose confirmation phase in specific tumor types. In the dose escalation phase, eligible patients will receive Compound A given orally, TIW in combination with a class I phosphatidylinositol 3-kinase (PI3K) and the mammalian target of rapamycin (mTOR) 15 (PI3K/mTOR) inhibitor Compound B given orally, every 12 hours, on a 28-day cycle. A single dose of Compound B will also be given on Day 1 during a 3-day lead-in period (dose-escalation phase only) for PK evaluation. In the dose-confirmation phase, approximately 10 patients each with advanced or metastatic colon cancer or soft tissue sarcoma will be treated. Colon cancer patients are required to have mutations, amplification, or gene expression 20 alterations related to Notch pathway signaling.

Study Objectives

The primary objective of this study is to determine the recommended Phase 2 dose of Compound A in individual combination with a PI3K/mTOR inhibitor Compound B anticancer agent.

25 The secondary objectives of the study are to characterize the safety and toxicity profile of Compound A in combination with Compound B as assessed by National Cancer Institute's (NCI) Common Terminology Criteria for Adverse Events (CTCAE) v4.0; to estimate the PK parameters of Compound B in combination with Compound A ; to

-19-

document any antitumor activity observed with Compound A in combination with Compound B; and to assess duration of response and progression free survival (PFS).

Exploratory objectives are to explore pharmacodynamic (PD) effects of Compound A on biomarkers indicative of Notch activity or Compound B; to explore the utility of 5 positron emission tomography (PET) scan to assess treatment effect with Compound A in combination with Compound B; to explore predictive biomarkers related to induction of cytochrome P450 (CYP) enzymes, such as cortisol and 6 β -hydroxycortisol; and to evaluate tumor tissue and blood for biomarkers related to the Notch signaling pathway and drug target pathways, immune functioning, mechanism of action of study drug(s) or disease state, and 10 their potential association with the objectives of the study.

Trial Drug

25 4,4,4-Trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide hydrate (Compound A), as 3 times per week, during a 28-day cycle.

15 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one (Compound B), as 100 or 200 mg capsules, or as 50, 100, 150, or 200 mg tablets for oral administration as a chronic/continuous treatment approximately every 12 hours during a 28-day cycle. For dose escalation, a 20 single dose of Compound B is administered on Day 1 during a 3-day lead-in period and administered B.I.D. on Days 1 to 28 of a 28-day cycle.

Planned Duration of Treatment/ Dosing

25 By nature of being a dose escalation study, data will be evaluated on an ongoing basis until the maximum tolerated dose (MTD) of the combination is determined. Dose escalation will be driven by the 3+3 method.

Each new dose level will have a minimum of 3 patients enrolled to it. If 1 patient, at any dose level, experiences a dose-limiting toxicity (DLT) within the first cycle of Compound A, then up to 3 additional patients will be enrolled at that dose level. If a DLT is observed in 2 or more patients at any dose level, dose escalation will cease and either the

-20-

previous dose level will be declared the MTD or, following discussions between the sponsor and investigators additional patients may be treated at intermediate doses between the previous and current dose levels.

During dose escalation, the starting dose of Compound A will be 25 mg TIW and the 5 starting dose of Compound B will be 150 mg BID. Dose escalation is scheduled to proceed according to Table 3.

10 **Table 3**
Dose-Escalation Scheme

Dose Level	Compound A Dose (mg)	Compound B Dose (mg)
1	25	150 BID
2	50	150 BID
3	50	200 BID

Abbreviation: BID = twice daily.

Criteria for Evaluation

15 Safety: NCI CTCAE, version 4.0, dose-limiting toxicities (DLT).

Efficacy: Each patient will be assessed by one or more of the following radiologic tests for tumor measurement: Computed tomography (CT) scan; Magnetic resonance imaging (MRI); and PET scan (pre- and postdose). Each patient's full extent of disease will also be assessed with:

20 Tumor measurement by RECIST 1.1 (Eisenhauer et al., *Eur. J. Cancer*, 2009; 45(2): 228-247). For tumor measurement evaluations in patients with soft tissue sarcomas, Choi et al., *J. Clin. Oncol.*, 2007; 25(13): 1753-1759 response criteria will be used in addition to RECIST 1.1. Response Assessment in Neuro-Oncology (RANO) criteria will be used for glioblastoma patients (Wen et al., *J. Clin. Oncol.*, 2010; 28(11): 1963-1972);

25 Evaluation of tumor markers, if indicated;

Evaluation of performance status (Eastern Cooperative Oncology Group (ECOG)); Oken et al., *Am. J. Clin. Oncol.*, 1982; 5: 649-655).

-21-

To confirm objective responses, all lesions should be radiologically assessed, and the same radiologic method used for the initial response determination should be repeated at least 4 weeks following the initial observation of an objective response, using the sample method that was used at baseline. If a patient is discontinued from the study, repeat radiology

5 assessments may be omitted if clear clinical signs of progressive disease are present.

Claims:

1. A method of treating a cancer, comprising administering to a patient in need of treatment an effective amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof, and an effective amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof, wherein the cancer is selected from the group consisting of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors, adenoid cystic carcinoma, colorectal cancer, head and neck cancer, cervical cancer, prostate cancer, liver cancer, and skin cancer.
2. The method of Claim 1, wherein the cancer is ovarian cancer.
3. The method of claim 1 or claim 2, wherein the administering is by simultaneous, separate, or sequential administration of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof and 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.
4. The method of any one of claims 1 to 3, wherein the amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof is between 2.5 mg and 75 mg.
5. The method of any one of claims 1 to 3, wherein the amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof is between 5 mg and 50 mg.
6. The method of any one of claims 1 to 3, wherein the amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof is 25 mg.

7. The method of any one of claims 1 to 3, wherein the amount of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof is 50 mg.

5 8. The method of any one of claims 1 to 7, wherein the 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof is for use once per day every other day over a five day period followed by two days without dosing (T.I.W.).

10 9. The method of any one of claims 1 to 8, wherein the amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof is between 75 mg and 250 mg.

15 10. The method of any one of claims 1 to 8, wherein the amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof is between 100 mg and 200 mg.

11. The method of any one of claims 1 to 8, wherein the amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof is 150 mg.

12. The method of any one of claims 1 to 8, wherein the amount of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof is 200 mg.

13. The method of any one of claims 1 to 12, wherein the 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof is for use twice per day (B.I.D.).

14. The method of any one of claims 1 to 13, wherein the 4,4,4-trifluoro-N-[(1S)-2-
[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-
2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof and the
8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-
5 dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof
are formulated for oral administration.

15. Use of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-
d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically
acceptable salt or hydrate thereof for the manufacture of a first medicament; and use of 8-
10 [5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-
dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof
for the manufacture of a second medicament; for use in combination for the treatment of a
cancer selected from the group consisting of T-cell acute lymphoblastic leukemia, acute
15 lymphoblastic leukemia, acute myelogenous leukemia, chronic myelogenous leukemia,
breast cancer, ovarian cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma,
sarcoma, desmoid tumors, adenoid cystic carcinoma, colorectal cancer, head and neck
cancer, cervical cancer, prostate cancer, liver cancer, and skin cancer.

16. The use of Claim 15, wherein the cancer is ovarian cancer.

17. The use of claim 15 or claim 16, wherein the use is for treatment of the cancer by
20 simultaneous, separate, or sequential administration of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-
(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-
ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof and 8-[5-(1-
hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-
2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

25 18. The use of any one of claims 15 to 17, wherein the first medicament comprises
between 2.5 mg and 75 mg of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-
7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a
pharmaceutically acceptable salt or hydrate thereof.

19. The use of any one of claims 15 to 17, wherein the first medicament comprises
30 between 5 mg and 50 mg of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-
7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a
pharmaceutically acceptable salt or hydrate thereof.

20. The use of any one of claims 15 to 17, wherein the first medicament comprises 25 mg of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof.

5 21. The use of any one of claims 15 to 17, wherein the first medicament comprises 50 mg of 4,4,4-trifluoro-N-[(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof.

10 22. The use of any one of claims 15 to 21, wherein the first medicament is formulated for use once per day every other day over a five day period followed by two days without dosing (T.I.W.).

15 23. The use of any one of claims 15 to 22, wherein the second medicament comprises between 75 mg and 250 mg of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

24. The use of any one of claims 15 to 22, wherein the second medicament comprises between 100 mg and 200 mg of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

20 25. The use of any one of claims 15 to 22, wherein the second medicament comprises 150 mg of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

25 26. The use of any one of claims 15 to 22, wherein the second medicament comprises 200 mg of 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a pharmaceutically acceptable salt thereof.

27. The use of any one of claims 15 to 26, wherein the second medicament is formulated for use twice per day (B.I.D.).

28. The use of any one of claims 15 to 27, wherein the first medicament and the second medicament are manufactured for oral administration.

29. A commercial package when used in treatment of a cancer, comprising:

two separate compositions, wherein one composition comprises 4,4,4-trifluoro-N-
5 [(1S)-2-[(7S)-5-(2-hydroxyethyl)-6-oxo-7H-pyrido[2,3-d][3]benzazepin-7-yl]amino]-1-
methyl-2-oxo-ethyl]butanamide, or a pharmaceutically acceptable salt or hydrate thereof,
and the other composition comprises 8-[5-(1-hydroxy-1-methylethyl)pyridin-3-yl]-1-
[(2S)-2-methoxypropyl]-3-methyl-1,3-dihydro-2H-imidazo[4,5-c]quinolin-2-one, or a
pharmaceutically acceptable salt thereof, and

10 instructions for use in treating the cancer, wherein the cancer is selected from the
group consisting of T-cell acute lymphoblastic leukemia, acute lymphoblastic leukemia,
acute myelogenous leukemia, chronic myelogenous leukemia, breast cancer, ovarian
cancer, melanoma, lung cancer, pancreatic cancer, glioblastoma, sarcoma, desmoid tumors,
adenoid cystic carcinoma, colorectal cancer, head and neck cancer, cervical cancer, prostate
15 cancer, liver cancer, and skin cancer.

Eli Lilly and Company

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON