



(86) Date de dépôt PCT/PCT Filing Date: 2015/04/14
(87) Date publication PCT/PCT Publication Date: 2015/10/29
(85) Entrée phase nationale/National Entry: 2016/10/21
(86) N° demande PCT/PCT Application No.: US 2015/025690
(87) N° publication PCT/PCT Publication No.: 2015/164121
(30) Priorité/Priority: 2014/04/24 (US61/983,732)

(51) Cl.Int./Int.Cl. *C12N 1/21* (2006.01)
(71) Demandeur/Applicant:
ADVAXIS, INC., US
(72) Inventeurs/Inventors:
WALLECHA, ANU, US;
PETIT, ROBERT, US
(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : SOUCHES DE LISTERIA UTILISEES COMME VACCIN RECOMBINANT ET PROCEDE DE PRODUCTION
(54) Title: RECOMBINANT LISTERIA VACCINE STRAINS AND METHODS OF PRODUCING THE SAME




Figure 12

(57) **Abrégé/Abstract:**

The present invention provides methods of treating, protecting against and inducing an immune response against a tumor or cancer, comprising the step of administering to a subject a recombinant *Listeria* strain. In one embodiment the present invention

(57) **Abrégé(suite)/Abstract(continued):**

relates to a recombinant *Listeria* strain, said recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising a first N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, and wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property

Organization

International Bureau

(43) International Publication Date
29 October 2015 (29.10.2015)

(10) International Publication Number

WO 2015/164121 A1

(51) International Patent Classification:

C12N 1/21 (2006.01)

(21) International Application Number:

PCT/US2015/025690

(22) International Filing Date:

14 April 2015 (14.04.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/983,732

24 April 2014 (24.04.2014)

US

(71) Applicant: ADVAXIS, INC. [US/US]; 305 College Road East, Princeton, New Jersey 08540 (US).

(72) Inventors: WALLECHA, Anu; 201 Wildflower Circle, Yardley, Pennsylvania 19067 (US). PETIT, Robert; 149 Parsons Lane, Newtown (Wrightstown), Pennsylvania 18940 (US).

(74) Agents: PEARL COHEN ZEDEK LATZER BARATZ LLP et al.; 1500 Broadway, 12th Floor, New York, New York 10036 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: RECOMBINANT LISTERIA VACCINE STRAINS AND METHODS OF PRODUCING THE SAME




Figure 12

(57) Abstract: The present invention provides methods of treating, protecting against and inducing an immune response against a tumor or cancer, comprising the step of administering to a subject a recombinant Listeria strain. In one embodiment the present invention relates to a recombinant Listeria strain, said recombinant Listeria strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising a first N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, and wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein.

**RECOMBINANT *LISTERIA* VACCINE STRAINS AND METHODS OF
PRODUCING THE SAME**

FIELD OF INVENTION

[001] The present invention provides methods of treating, protecting against, and inducing an immune response against a tumor or cancer, comprising the step of administering to a subject a recombinant *Listeria* strain comprising a nucleic acid encoding a mutant PrfA protein that partially restores PrfA function.

BACKGROUND OF THE INVENTION

[002] Persistent infection with high-oncogenic risk human papillomavirus (HR-HPV) types is recognized as a necessary, but not sufficient, cause of invasive carcinoma of the cervix (ICC) [1-3]. HPVs 16 and 18 are the most prevalent types in malignant lesions, accounting for over 70% of ICC and over 50% of high-grade precursor lesions. The HR-HPV E6 and E7 proteins are consistently expressed in dysplasias and carcinomas, disrupting the cell cycle regulatory proteins p53 and pRb, respectively. The obligatory expression of E6 and E7 by both dysplastic and invasive malignant lesions, as well as the viral origin of these proteins, make them excellent targets for HPV therapeutic vaccines.

[003] *Listeria monocytogenes* (*Lm*) is a food-borne gram-positive bacterium that can occasionally cause disease in humans, in particular elderly individuals, newborns, pregnant women and immunocompromised individuals. In addition to strongly activating innate immunity and inducing a cytokine response that enhances antigen-presenting cell (APC) function, *Lm* has the ability to replicate in the cytosol of APCs after escaping from the phagolysosome, mainly through the action of the listeriolysin O (LLO) protein. This unique intracellular life cycle allows antigens secreted by *Lm* to be processed and presented in the context of both MHC class I and II molecules, resulting in potent cytotoxic CD8⁺ and Th1 CD4⁺ T-cell-mediated immune responses. *Lm* has been extensively investigated as a vector for cancer immunotherapy in pre-clinical models. Immunization of mice with *Lm*-LLO-E7 induces regression of established tumors expressing E7 and confers long-term protection. The therapeutic efficacy of *Lm*-LLO-E7 correlates with its ability to induce E7-specific CTLs that infiltrate the tumor site, mature dendritic cells, reduce the number of intratumoral regulatory CD4⁺ CD25⁺ T cells and inhibit tumor angiogenesis.

[004] *Lm* has also a number of inherent advantages as a vaccine vector. The bacterium grows very efficiently *in vitro* without special requirements and it lacks LPS, which is a major toxicity factor in gram-negative bacteria, such as *Salmonella*. Genetically attenuated *Lm* vectors also offer additional safety as they can be readily eliminated with antibiotics, in case of serious adverse effects and unlike some viral vectors, no integration of genetic material into the host genome occurs. However, there is always great concern about the safety of a live bacterial vaccine such as *Lm*, especially regarding its mechanism of attenuation.

[005] The PrfA protein controls the expression of a regulon comprising essential virulence genes required by *Lm* to colonize its vertebrate hosts; hence the *prfA* mutation strongly impairs PrfA ability to activate expression of PrfA-dependent virulence genes. The present invention addresses this concern by providing a *prfA* mutant *Listeria* that carries a mutant *prfA* (D133V) gene in the pGG55 plasmid that restores partial PrfA function.

SUMMARY OF THE INVENTION

[006] In one embodiment, the present invention relates to a recombinant *Listeria* strain, said recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising a first an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, and wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein.

[007] In one embodiment, the present invention relates to a recombinant *Listeria* strain, said recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising a first an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein, and wherein said *Listeria* comprises a genomic mutation or deletion in the *prfA* gene. In another embodiment, the mutant PrfA protein encoded by said second open reading frame complements said genomic mutation or deletion in said *Listeria* strain's PrfA protein. In another embodiment, the mutant PrfA protein encoded by said second open reading frame restores partial PrfA function in said *Listeria* strain.

[008] In one embodiment, the present invention relates to a method for inducing an immune response against a tumor or a cancer in a subject, the method comprising the step of administering to said subject a recombinant *Listeria* strain comprising a recombinant nucleic

acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, is, wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein, thereby inducing an immune response against a tumor or a cancer. In another embodiment, the recombinant *Listeria* strain comprises a genomic mutation or deletion in the *prfA* gene.

[009] Other features and advantages of the present invention will become apparent from the following detailed description examples and figures. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] The subject matter regarded as the invention is particularly pointed out and distinctly claimed in the concluding portion of the specification. The invention, however, both as to organization and method of operation, together with objects, features, and advantages thereof, may best be understood by reference to the following detailed description when read with the accompanying drawings in which:

[0011] Figure 1. Lm-E7 and Lm-LLO-E7 use different expression systems to express and secrete E7. Lm-E7 was generated by introducing a gene cassette into the orfZ domain of the *L. monocytogenes* genome (A). The hly promoter drives expression of the hly signal sequence and the first five amino acids (AA) of LLO followed by HPV-16 E7. B), Lm-LLO-E7 was generated by transforming the prfA- strain XFL-7 with the plasmid pGG-55. pGG-55 has the hly promoter driving expression of a nonhemolytic fusion of LLO-E7. pGG-55 also contains the prfA gene to select for retention of the plasmid by XFL-7 *in vivo*.

[0012] Figure 2. Lm-E7 and Lm-LLO-E7 secrete E7. Lm-Gag (lane 1), Lm-E7 (lane 2), Lm-LLO-NP (lane 3), Lm-LLO-E7 (lane 4), XFL-7 (lane 5), and 10403S (lane 6) were grown overnight at 37°C in Luria-Bertoni broth. Equivalent numbers of bacteria, as determined by OD at 600 nm absorbance, were pelleted and 18 ml of each supernatant was TCA precipitated. E7 expression was analyzed by Western blot. The blot was probed with an anti-E7 mAb, followed by HRP-conjugated anti-mouse (Amersham), then developed using ECL detection reagents.

[0013] Figure 3. Tumor immunotherapeutic efficacy of LLO-E7 fusions. Tumor size in millimeters in mice is shown at 7, 14, 21, 28 and 56 days post tumor-inoculation. Naive mice: open-circles; Lm-LLO-E7: filled circles; Lm-E7: squares; Lm-Gag: open diamonds; and Lm-LLO-NP: filled triangles.

5 [0014] Figure 4. Splenocytes from Lm-LLO-E7-immunized mice proliferate when exposed to TC-1 cells. C57BL/6 mice were immunized and boosted with Lm-LLO-E7, Lm-E7, or control rLm strains. Splenocytes were harvested 6 days after the boost and plated with irradiated TC-1 cells at the ratios shown. The cells were pulsed with ^3H thymidine and harvested. Cpm is defined as (experimental cpm) - (no-TC-1 control).

10 [0015] Figure 5. A. Induction of E7-specific IFN-gamma-secreting CD8^+ T cells in the spleens and the numbers penetrating the tumors, in mice administered TC-1 tumor cells and subsequently administered Lm-E7, Lm-LLO-E7, Lm-ActA-E7, or no vaccine (naive). B. Induction and penetration of E7 specific CD8^+ cells in the spleens and tumors of the mice described for (A).

15 [0016] Figure 6. *Listeria* constructs containing PEST regions induce a higher percentage of E7-specific lymphocytes within the tumor. A. representative data from 1 experiment. B. average and SE of data from all 3 experiments.

[0017] Figure 7A. Effect of passaging on bacterial load (virulence) of recombinant *Listeria* vaccine vectors. Top panel. Lm-Gag. Bottom panel. Lm-LLO-E7. Figure 7B. Effect of
20 passaging on bacterial load of recombinant Lm-E7 in the spleen. Average CFU of live bacteria per milliliter of spleen homogenate from four mice is depicted.

[0018] Figure 8 shows induction of antigen-specific CD8^+ T-cells for HIV-Gag and LLO after administration of passaged Lm-Gag versus unpassaged Lm-Gag. Mice were immunized with 10^3 (A, B, E, F) or 10^5 (C, D, G, H) CFU passaged *Listeria* vaccine vectors, and antigen-
25 specific T-cells were analyzed. B, D, F, H: unpassaged *Listeria* vaccine vectors. A-D immune response to MHC class I HIV-Gag peptide. E-H: immune response to an LLO peptide. I: splenocytes from mice immunized with 10^5 CFU passaged Lm-Gag stimulated with a control peptide from HPV E7.

[0019] Figure 9A shows plasmid isolation throughout LB stability study. Figure 9B shows
30 plasmid isolation throughout TB stability study. Figure 9C shows quantitation of TB stability study.

[0020] Figure 10 shows numbers of viable bacteria chloramphenicol (CAP)-resistant and CAP-sensitive colony-forming units (CFU) from bacteria grown in LB. Dark bars: CAP⁺; white bars: CAP⁻. The two dark bars and two white bars for each time point represent duplicate samples.

5 [0021] Figure 11 shows numbers of viable bacteria CAP-resistant and CAP-sensitive CFU from bacteria grown in TB. Dark bars: CAP⁺; white bars: CAP⁻. The two dark bars and two white bars for each time point represent duplicate samples.

[0022] Figure 12. Actual chromatograms showing the region of the D133V mutation (arrows). The mixture ratio is shown in parentheses.

10 [0023] Figure 13. Representation of the location of the ADV451, 452 and 453 primers and the segment of the *prfA* gene amplified in the reaction.

[0024] Figure 14. Specificity of the PCR reaction using primers ADV451 and ADV453.

[0025] Figure 15. Specificity of the PCR reaction using primers ADV452 and ADV453.

[0026] Figure 16. Sensitivity of the PCR reaction to detect the wild-type *prfA* sequence using
15 the primer ADV452 and 1 ng as the initial amount of DNA.

[0027] Figure 17. Sensitivity of the PCR reaction to detect the wild-type *prfA* sequence using the primer ADV452 and 5 ng as the initial amount of DNA.

[0028] Figure 18. Average density of the bands from the PCR depicted in figure 16.

[0029] Figure 19. Average density of the bands from the PCR depicted in figure 17.

20 [0030] Figure 20. Validation of the PCR reaction to detect the wild-type *prfA* sequence using the primer ADV452.

[0031] Figure 21. Average density of the bands from the PCR depicted in figure 16.

[0032] Figure 22. Analysis of the D133V *prfA* mutation in the *Lm*-LLO-E7. A, Original image used for densitometry; B, Image was digitally enhanced to facilitate the visualization of
25 the low density bands.

[0033] It will be appreciated that for simplicity and clarity of illustration, elements shown in the figures have not necessarily been drawn to scale. For example, the dimensions of

some of the elements may be exaggerated relative to other elements for clarity. Further, where considered appropriate, reference numerals may be repeated among the figures to indicate corresponding or analogous elements.

DETAILED DESCRIPTION OF THE INVENTION

5 [0034] The present invention provides, in one embodiment, a recombinant *Listeria* strain, said recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising a first an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, wherein said recombinant nucleic acid further comprises a second open reading frame
10 encoding a mutant PrfA protein, and wherein said *Listeria* comprises a genomic mutation or deletion in the *prfA* gene. In another embodiment, the mutant PrfA protein encoded by said second open reading frame complements said genomic mutation or deletion in said *Listeria* strain's *prfA* gene. In another embodiment, the mutant PrfA protein encoded by said second open reading frame restores partial PrfA function in said *Listeria* strain. In one embodiment,
15 the mutant *PrfA* protein encoded by said second open reading frame comprises a point mutation in position 133. In another embodiment, the mutation on residue 133 of the PrfA amino acid sequence is from amino acid D or Asp or Aspartate (or Aspartic acid) to amino acid V or Val or Valine.

[0035] The present invention further provides immunogenic compositions comprising a
20 recombinant *Listeria* strain provided herein and methods of using the same, including methods of treating, protecting against, and inducing an immune response against a disease, where in some embodiments, the disease is a tumor or cancer.

[0036] The present invention also provides methods for inducing an anti-disease cytotoxic T-cell (CTL) response in a subject and treating disorders, and symptoms associated with said
25 disease comprising administering a recombinant *Listeria* strain provided herein, wherein in some embodiments the disease is a tumor or a cancer.

[0037] In another embodiment, a recombinant *Listeria* provided herein is an attenuated *Listeria*. "Attenuation" and "attenuated" may encompass a bacterium, virus, parasite, infectious organism, prion, tumor cell, gene in the infectious organism, and the like, that is
30 modified to reduce toxicity to a host. The host can be a human or animal host, or an organ, tissue, or cell. The bacterium, to give a non-limiting example, can be attenuated to reduce binding to a host cell, to reduce spread from one host cell to another host cell, to reduce

extracellular growth, or to reduce intracellular growth in a host cell. Attenuation can be assessed by measuring, e.g., an indicum or indicia of toxicity, the LD₅₀, the rate of clearance from an organ, or the competitive index (see, e.g., Auerbuch, et al. (2001) Infect. Immunity 69:5953-5957). Generally, an attenuation results an increase in the LD₅₀ and/or an increase in the rate of clearance by at least 25%; more generally by at least 50%; most generally by at least 100% (2-fold); normally by at least 5-fold; more normally by at least 10-fold; most normally by at least 50-fold; often by at least 100-fold; more often by at least 500-fold; and most often by at least 1000-fold; usually by at least 5000-fold; more usually by at least 10,000-fold; and most usually by at least 50,000-fold; and most often by at least 100,000-fold.

[0038] It will be well appreciated by a skilled artisan that the term “Attenuated gene” may encompass a gene that mediates toxicity, pathology, or virulence, to a host, growth within the host, or survival within the host, where the gene is mutated in a way that mitigates, reduces, or eliminates the toxicity, pathology, or virulence. The reduction or elimination can be assessed by comparing the virulence or toxicity mediated by the mutated gene with that mediated by the non-mutated (or parent) gene. “Mutated gene” encompasses deletions, point mutations, and frameshift mutations in regulatory regions of the gene, coding regions of the gene, non-coding regions of the gene, or any combination thereof.

[0039] In one embodiment, provided herein is a method for inducing an immune response against a tumor or a cancer in a subject, the method comprising the step of administering to said subject a composition comprising a recombinant *Listeria* strain provided herein, thereby inducing an immune response against a tumor or a cancer.

[0040] In one embodiment, the present invention provides a method of treating a tumor or cancer in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein. In another embodiment, the present invention provides a method of protecting a subject against a tumor or cancer, comprising the step of administering to the subject the recombinant *Listeria* strain provided herein. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a genomic mutation or deletion in the *prfA* gene.

[0041] In one embodiment, the methods provided herein further comprise the step of boosting a subject with a composition comprising a recombinant *Listeria* strain of the present

invention. In another embodiment, the method further comprises the step of boosting the subject with an immunogenic composition comprising a heterologous antigen or fragment thereof provided herein. In another embodiment, the method further comprises the step of boosting the subject with an immunogenic composition that directs a cell of the subject to
 5 express the heterologous antigen. In another embodiment, the cell is a tumor cell. In another embodiment, the cell is an antigen-presenting cell. In another embodiment, the method further comprises the step of boosting the subject with a vaccine comprising a recombinant *Listeria* strain of the present invention.

[0042] In one embodiment, the fragment thereof in the context of LLO proteins and ActA
 10 proteins provided herein refer to a peptide or polypeptide comprising an amino acid sequence of at least 5 contiguous amino acid residues of the LLO or ActA proteins. In another embodiment, the term refers to a peptide or polypeptide comprising an amino acid sequence of at least of at least 10 contiguous amino acid residues, at least 15 contiguous amino acid residues, at least 20 contiguous amino acid residues, at least 25 contiguous amino acid
 15 residues, at least 40 contiguous amino acid residues, at least 50 contiguous amino acid residues, at least 60 contiguous amino residues, at least 70 contiguous amino acid residues, at least 80 contiguous amino acid residues, at least 90 contiguous amino acid residues, at least 100 contiguous amino acid residues, at least 125 contiguous amino acid residues, at least 150 contiguous amino acid residues, at least 175 contiguous amino acid residues, at least 200
 20 contiguous amino acid residues, at least 250 contiguous amino acid residues of the amino acid sequence, at least 300 contiguous amino acid residues, at least 350 contiguous amino acid residues of, at least 400 contiguous amino acid residues, or at least 450 contiguous amino acid residues of an LLO or ActA protein or polypeptide.

[0043] In another embodiment, a “fragment” is a functional fragment that comprises a
 25 biological activity (e.g. to elicit an immune response against a heterologous antigen expressed by a tumor cell, either when administered alone or when administered in the context of a fusion protein as further described herein. In another embodiment, the fragment is functional in a non-fused form.

[0044] The present invention, in certain embodiments, provides codon optimization of a
 30 nucleic acid heterologous to *Listeria*, or of a nucleic acid endogenous to *Listeria*. The optimal codons utilized by *L. monocytogenes* for each amino acid are shown US Patent Publication 2007/0207170, which is hereby incorporated by reference herein. A nucleic acid is codon-optimized if at least one codon in the nucleic acid is replaced with a codon that is more

frequently used by *L. monocytogenes* for that amino acid than the codon in the original sequence.

[0045] An N-terminal LLO protein fragment and heterologous antigen provided herein are, in one embodiment, fused directly to one another. In another embodiment, the genes encoding the N-terminal LLO protein fragment and the heterologous antigen are fused directly to one another. In another embodiment, the N-terminal LLO protein fragment and the heterologous antigen are attached via a linker peptide. In another embodiment, the N-terminal LLO protein fragment and the heterologous antigen are attached via a heterologous peptide. In another embodiment, the N-terminal LLO protein fragment is N-terminal to the heterologous antigen. In another embodiment, the N-terminal LLO protein fragment is the N-terminal-most portion of the fusion protein. Each possibility represents a separate embodiment of the present invention.

[0046] As provided herein, recombinant *Listeria* strains expressing LLO-antigen fusions induce anti-tumor immunity (Example 1), elicit antigen-specific T cell proliferation (Example 2), generate antigen-specific, and tumor-infiltrating T cells (Example 3).

[0047] In another embodiment, the present invention provides a method of treating a tumor or cancer in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an LLO protein and an HPV E7 antigen, whereby the recombinant *Listeria* strain induces an immune response against the E7 antigen, thereby treating a tumor or cancer in a subject. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. Each possibility represents a separate embodiment of the present invention.

[0048] In one embodiment, the terms “recombinant polypeptide” and “fusion protein” are used interchangeably herein.

[0049] In another embodiment, the present invention provides a method of protecting a subject against a tumor or cancer, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an LLO protein and an HPV E7 antigen, whereby the recombinant *Listeria* strain induces an immune response against the E7 antigen, thereby protecting a subject against a tumor or cancer. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment,

the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. Each possibility represents a separate embodiment of the present invention.

[0050] In another embodiment, the present invention provides a method for inducing an immune response against a tumor or cancer in a subject, comprising the step of administering
5 to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an LLO protein and an HPV E7 antigen, thereby inducing an immune response against a tumor or cancer in a subject. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the
10 recombinant polypeptide. Each possibility represents a separate embodiment of the present invention.

[0051] In another embodiment, the present invention provides a method of treating a tumor or cancer in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide
15 comprising an N-terminal fragment of an ActA protein and heterologous antigen, whereby the recombinant *Listeria* strain induces an immune response against the heterologous antigen, thereby treating a tumor or cancer in a subject. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide.
20 Each possibility represents a separate embodiment of the present invention.

[0052] In another embodiment, the present invention provides a method of protecting a subject against a tumor or cancer, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an ActA protein and a heterologous
25 antigen, whereby the recombinant *Listeria* strain induces an immune response against the heterologous antigen, thereby protecting a subject against a tumor or cancer. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. Each possibility represents a separate embodiment of the present
30 invention.

[0053] In another embodiment, the present invention provides a method for inducing an immune response against a tumor or cancer in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an heterologous protein and a

heterologous antigen, thereby inducing an immune response against a tumor or cancer in a subject. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. Each possibility represents a separate embodiment of the present invention.

[0054] The N-terminal ActA protein fragment and the heterologous antigen are, in another embodiment, fused directly to one another. In another embodiment, the genes encoding the N-terminal ActA protein fragment and heterologous antigen are fused directly to one another. In another embodiment, the N-terminal ActA protein fragment and heterologous antigen are attached via a linker peptide. In another embodiment, the N-terminal ActA protein fragment and heterologous antigen are attached via a heterologous peptide. In another embodiment, the N-terminal ActA protein fragment is N-terminal to the heterologous antigen. In another embodiment, the N-terminal ActA protein fragment is the N-terminal-most portion of the fusion protein. Each possibility represents a separate embodiment of the present invention.

[0055] In another embodiment, the present invention provides a method of inducing an immune response against a tumor or cancer in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising a PEST amino acid sequence-containing peptide and a heterologous antigen, whereby the recombinant *Listeria* strain induces an immune response against the heterologous antigen, thereby treating a tumor or cancer in a subject. In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. In another embodiment, the method protects a subject against a tumor or cancer. In another embodiment, the method treats a tumor or cancer in said subject.

[0056] The PEST amino acid sequence-containing peptide and heterologous antigen are, in another embodiment, fused directly to one another. In another embodiment, the genes encoding the PEST amino acid sequence-containing peptide and heterologous antigen are fused directly to one another. In another embodiment, the PEST amino acid sequence-containing peptide and heterologous antigen are attached via a linker peptide. In another embodiment, the PEST amino acid sequence-containing peptide and heterologous antigen are attached via a heterologous peptide. In another embodiment, the PEST amino acid sequence-containing peptide is N-terminal to the heterologous antigen. In another embodiment, the PEST amino acid sequence-containing peptide is the N-terminal-most portion of the fusion protein. Each possibility represents a separate embodiment of the present invention.

[0057] In another embodiment, the present invention provides a method for vaccinating a subject against an HPV, comprising the step of administering to the subject a *prfA* mutant recombinant *Listeria* strain provided herein, wherein the *Listeria* expresses an HPV antigen and wherein the *Listeria* comprises a plasmid that expresses a mutant PrfA protein. In another
5 embodiment, the recombinant *Listeria* strain expresses a recombinant polypeptide comprising said HPV antigen. In another embodiment, the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. Each possibility represents a separate embodiment of the invention.

[0058] In one embodiment, provided herein is a method of increasing a ratio of T effector
10 cells to regulatory T cells (Tregs) in the spleen and tumor microenvironments of a subject, comprising administering the immunogenic composition provided herein. In another embodiment, increasing a ratio of T effector cells to regulatory T cells (Tregs) in the spleen and tumor microenvironments in a subject allows for a more profound anti-tumor response in the subject.

[0059] In one embodiment, a mutant PrfA protein provided herein comprises a D133V amino
15 acid mutation. In another embodiment, the mutant PrfA protein consists of a D133V amino acid mutation. In another embodiment, a nucleic acid comprising an open reading frame encoding a mutant PrfA protein provided herein is in a plasmid in said recombinant *Listeria*. In another embodiment, the plasmid comprising a nucleic acid encoding a mutant PrfA
20 protein provided herein is an integrative plasmid. In another embodiment, the plasmid comprising a nucleic acid encoding a mutant PrfA protein provided herein is an episomal or extrachromosomal plasmid.

[0060] In one embodiment, a *prfA* mutant recombinant *Listeria* provided herein comprises a
25 partial deletion in or a complete deletion of the chromosomal *prfA* gene. In another embodiment, the *prfA* mutant *Listeria* comprises a loss-of-function mutation in the *prfA* gene.

[0061] In one embodiment, a mutant PrfA protein provided herein complements a genomic
deletion, inactivation or mutation in the *prfA* gene in a recombinant *Listeria*. In another
embodiment, a mutant PrfA protein provided herein complements a genomic deletion,
inactivation or mutation in the *prfA* gene in the recombinant *Listeria* provided herein. In
30 another embodiment, a mutant PrfA protein provided herein restores partial *prfA* function in a recombinant *Listeria* comprising a genomic deletion, inactivation or mutation of the *prfA* gene. In another embodiment, a mutant PrfA protein provided herein restores a loss-of PrfA function mutation in a recombinant *Listeria*.

[0062] In one embodiment, a wild-type PrfA protein is encoded by the following wild-type nucleic acid sequence set forth in SEQ ID NO: 31.

```

1  atgaacgctc  aagcagaaga  attcaaaaaa  tatttagaaa  ctaacgggat
   aaaacccaaa
5      61  caatttcata  aaaaagaact  tatTTTTtaac  caatgggatc
   cacaagaata  ttgtatTTTT
      121  ctatatgatg  gtatcacaaa  gctcacgagt  attagcgaga
   acgggaccat  catgaattta
      181  caatactaca  aaggggcttt  cgTTataatg  tctggcttta
10  ttgatacaga  aacatcggtt
      241  ggctattata  atttagaagt  cattagcgag  caggctaccg
   catacgttat  caaaataaac
      301  gaactaaaag  aactactgag  caaaaatctt  acgcactttt
   tctatgtttt  ccaaacccta
15      361  caaaaacaag  tttcatacag  cctagctaaa  tttaatgatt
   tttcgattaa  cgggaagctt
      421  ggctctatTT  gcggTcaact  tttaatcctg  acctatgtgt
   atggtaaaga  aactcctgat
      481  ggcatcaaga  ttacactgga  taattttaaca  atgcaggagt
20  taggatattc  aagtggcatc
      541  gcacatagct  cagctgttag  cagaattatt  tccaaattaa
   agcaagagaa  agttatcgtg
      601  tataaaaatt  catgctttta  tgtacaaaat  cttgattatc
   tcaaaagata  tgcccctaaa
25      661  ttagatgaat  ggTTTTatTT  agcatgtcct  gctacttggg
   gaaaattaaa  ttaa (SEQ ID NO: 31)

```

[0063] In one embodiment, a wild-type PrfA protein comprises an amino acid sequence set forth in SEQ ID NO: 32.

```

MNAQAEEFKKYLETNGIKPKQFHKKELIFNQWDPQEYCIF
30  LYDGITKLTSISENGTIMNLQYYKGAFVIMSGFIDTETSVG
   YYNLEVISEQATAYVIKINELKELLSKNLTHFFYVFQTLQK
   QVSYSLAKFNDFSINGKLGSICGQLLILTYVYGKETPDGIK
   ITLDNLTMQELGYSSGIAHSSAVSRIISKLLKQEKVIVYKNS
   CFYVQNLDYLRYPKLDEWFYLA CPATW GKLN (SEQ ID
35  NO: 32).

```

[0064] In one embodiment, a nucleic acid sequence encoding a mutant prfA sequence is set forth in SEQ ID NO: 33.

```

1      atgaacgctc  aagcagaaga  attcaaaaaa  tatttagaaa
40  ctaacgggat  aaaacccaaa
   61  caatttcata  aaaaagaact  tatTTTTtaac  caatgggatc  cacaagaata
   ttgtatTTTT
   121  ctatatgatg  gtatcacaaa  gctcacgagt  attagcgaga  acgggaccat
   catgaattta
45  181  caatactaca  aaggggcttt  cgTTataatg  tctggcttta  ttgatacaga
   aacatcggtt
   241  ggctattata  atttagaagt  cattagcgag  caggctaccg  catacgttat
   caaaataaac

```

301 gaactaaaag aactactgag caaaaatctt acgcactttt tctatgtttt
 ccaaacccta
 361 caaaaacaag tttcatacag cctagctaaa tttaatgttt tttcgattaa
 cgggaagctt
 5 421 ggctctattt gcggtcaact tttaatcctg acctatgtgt atggtaaaga
 aactcctgat
 481 ggcatcaaga ttacactgga taatttaaca atgcaggagt taggatattc
 aagtggcatc
 541 gcacatagct cagctgtag cagaattatt tccaaattaa agcaagagaa
 10 agttatcgtg
 601 tataaaaatt catgctttta tgtacaaaat cgtgattatc tcaaaagata
 tgcccctaaa
 661 ttagatgaat ggttttatatt agcatgtcct gctacttggg gaaaattaaa
 ttaa (SEQ ID NO: 33)

[0065] In one embodiment, a mutant PrfA protein provided herein comprises an amino acid sequence set forth in SEQ ID NO: 34.

MNAQAEEFKKYLETNGIKPKQFHKKELIFNQWDPQEYCIFY
 DGITKLTSISENGTIMNLQYYKGAFVIMSGFIDTETSVGYYNL
 20 EVISEQATAYVIKINELKELLSKNLTHFFYVFQTLQKQVSYSL
 AKFNyFSINGKLGSICGQLLILTYVYGKETPDGIKITLDNLT
 MELGYSSGIAHSSAVSRIISKLLKQEKVIVYKNSCFYVQNRDY
 LKRYAPKLDEWFYLA CPATWGKLN (SEQ ID NO: 34). In another
 embodiment, SEQ ID NO: 34 represents a mutant PrfA protein comprising a D133V
 25 mutation. In another embodiment, a mutant PrfA protein is homologous to SEQ ID NO: 34
 and comprises a D133V mutation. In another embodiment, a mutant PrfA protein is at least
 90% homologous with SEQ ID NO: 34 and comprises a D133V mutation. In another
 embodiment, a mutant PrfA protein is at least 85% homologous with SEQ ID NO: 34, and
 comprises a D133V mutation.

30 [0066] In another embodiment, the subject is at risk for developing an HPV-mediated
 carcinogenesis (e.g. a cervical, head and neck or anal cancer). In another embodiment, the
 subject is HPV-positive.

35 [0067] In another embodiment, the subject exhibits cervical intraepithelial neoplasia. In
 another embodiment, the subject exhibits a squamous intraepithelial lesion. In another
 embodiment, the subject exhibits a dysplasia in the cervix.

[0068] The HPV that is the target of methods of the present invention is, in another
 embodiment, an HPV 16. In another embodiment, the HPV is an HPV-18. In another
 embodiment, the HPV is selected from HPV-16 and HPV-18. In another embodiment, the
 40 HPV is an HPV-31. In another embodiment, the HPV is an HPV-35. In another embodiment,
 the HPV is an HPV-39. In another embodiment, the HPV is an HPV-45. In another

embodiment, the HPV is an HPV-51. In another embodiment, the HPV is an HPV-52. In another embodiment, the HPV is an HPV-58. In another embodiment, the HPV is a high-risk HPV type. In another embodiment, the HPV is a mucosal HPV type. Each possibility represents a separate embodiment of the present invention.

5 [0069] In another embodiment, the present invention provides a method of vaccinating a subject against an antigen of interest, the method comprising the step of intravenously administering to the subject an immunogenic composition, comprising a fusion of an immunogenic peptide to the antigen of interest, wherein the immunogenic peptide is selected from (a) an N-terminal fragment of an LLO protein; (b) an ActA protein or N-terminal
10 fragment thereof; and (c) a PEST amino acid sequence-containing peptide, thereby vaccinating a subject against an antigen of interest.

[0070] In another embodiment, the present invention provides a method of vaccinating a subject against an antigen of interest, the method comprising the step of administering intravenously to the subject a recombinant *Listeria* strain comprising a recombinant
15 polypeptide, the recombinant polypeptide comprising an immunogenic peptide fused to the antigen of interest, wherein the immunogenic peptide is selected from (a) an N-terminal fragment of an LLO protein; (b) an ActA protein or N-terminal fragment thereof; and (c) a PEST amino acid sequence-containing peptide, thereby vaccinating a subject against an antigen of interest.

20 [0071] In another embodiment, the present invention provides a method of inducing a CTL response in a subject against an antigen of interest, the method comprising the step of administering to the subject a recombinant *Listeria* strain comprising or expressing the antigen of interest, thereby inducing a CTL response in a subject against an antigen of interest. In another embodiment, the step of administering is intravenous or oral
25 administration. Each possibility represents a separate embodiment of the present invention.

[0072] As provided herein, recombinant *Listeria* strains expressing LLO-antigen fusions induce anti-tumor immunity (Example 1), elicit antigen-specific T cell proliferation (Example 2), generate antigen-specific, and tumor-infiltrating T cells (Example 3). Thus, vaccines of the present invention are efficacious at inducing immune responses against HPV antigens E7 and
30 E6.

[0073] In another embodiment, the present invention provides a method for inducing a regression of a cancer in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein.

[0074] In another embodiment, the present invention provides a method for reducing an incidence of relapse of a cancer in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein.

[0075] In another embodiment, the present invention provides a method for suppressing a formation of a tumor in a subject, comprising the step of administering to the subject a composition comprising recombinant *Listeria* strain provided herein.

[0076] In another embodiment, the present invention provides a method for inducing a remission of a cancer in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein.

[0077] In another embodiment, the present invention provides a method for impeding a growth of a tumor in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein.

[0078] In another embodiment, the present invention provides a method for reducing a size of a tumor in a subject, comprising the step of administering to the subject a composition comprising a recombinant *Listeria* strain provided herein.

[0079] In one embodiment, a disease is an infectious disease, an autoimmune disease, a respiratory disease, a pre-cancerous condition or a cancer.

[0080] It will be well appreciated by the skilled artisan that the term “pre-cancerous condition” may encompass dysplasias, preneoplastic nodules; macroregenerative nodules (MRN); low-grade dysplastic nodules (LG-DN); high-grade dysplastic nodules (HG-DN); biliary epithelial dysplasia; foci of altered hepatocytes (FAH); nodules of altered hepatocytes (NAH); chromosomal imbalances; aberrant activation of telomerase; re-expression of the catalytic subunit of telomerase; expression of endothelial cell markers such as CD31, CD34, and BNH9 (see, e.g., Terracciano and Tomillo (2003) *Pathologica* 95:71-82; Su and Bannasch (2003) *Toxicol. Pathol.* 31:126-133; Rocken and Carl-McGrath (2001) *Dig. Dis.* 19:269-278; Kotoula, et al. (2002) *Liver* 22:57-69; Frachon, et al. (2001) *J. Hepatol.* 34:850-857; Shimonishi, et al. (2000) *J. Hepatobiliary Pancreat. Surg.* 7:542-550; Nakanuma, et al. (2003) *J. Hepatobiliary Pancreat. Surg.* 10:265-281). Methods for diagnosing cancer and dysplasia are disclosed (see, e.g., Riegler (1996) *Semin. Gastrointest. Dis.* 7:74-87; Benvegna, et al. (1992) *Liver* 12:80-83; Giannini, et al. (1987) *Hepatogastroenterol.* 34:95-97; Anthony (1976) *Cancer Res.* 36:2579-2583).

[0081] In one embodiment, an infectious disease is one caused by, but not limited to, any one of the following pathogens: BCG/Tuberculosis, Malaria, *Plasmodium falciparum*, *plasmodium malariae*, *plasmodium vivax*, Rotavirus, Cholera, Diphtheria-Tetanus, Pertussis,

Haemophilus influenzae, Hepatitis B, Human papilloma virus, Influenza seasonal), Influenza A (H1N1) Pandemic, Measles and Rubella, Mumps, Meningococcus A+C, Oral Polio Vaccines, mono, bi and trivalent, Pneumococcal, Rabies, Tetanus Toxoid, Yellow Fever, Bacillus anthracis (anthrax), Clostridium botulinum toxin (botulism), Yersinia pestis (plague),

5 Variola major (smallpox) and other related pox viruses, Francisella tularensis (tularemia), Viral hemorrhagic fevers, Arenaviruses (LCM, Junin virus, Machupo virus, Guanarito virus, Lassa Fever), Bunyaviruses (Hantaviruses, Rift Valley Fever), Flaviruses (Dengue), Filoviruses (Ebola , Marburg), Burkholderia pseudomallei, Coxiella burnetii (Q fever), Brucella species (brucellosis), Burkholderia mallei (glanders), Chlamydia psittaci

10 (Psittacosis), Ricin toxin (from Ricinus communis), Epsilon toxin of Clostridium perfringens, Staphylococcus enterotoxin B, Typhus fever (Rickettsia prowazekii), other Rickettsias, Food- and Waterborne Pathogens, Bacteria (Diarrheagenic *E.coli*, Pathogenic Vibrios, Shigella species, Salmonella BCG/, Campylobacter jejuni, Yersinia enterocolitica), Viruses (Caliciviruses, Hepatitis A, West Nile Virus, LaCrosse, California encephalitis, VEE, EEE,

15 WEE, Japanese Encephalitis Virus, Kyasanur Forest Virus, Nipah virus, hantaviruses, Tickborne hemorrhagic fever viruses, Chikungunya virus, Crimean-Congo Hemorrhagic fever virus, Tickborne encephalitis viruses, Hepatitis B virus, Hepatitis C virus, Herpes Simplex virus (HSV), Human immunodeficiency virus (HIV), Human papillomavirus (HPV)), Protozoa (Cryptosporidium parvum, Cyclospora cayatanensis, Giardia lamblia, Entamoeba

20 histolytica, Toxoplasma), Fungi (Microsporidia), Yellow fever, Tuberculosis, including drug-resistant TB, Rabies, Prions, Severe acute respiratory syndrome associated coronavirus (SARS-CoV), Coccidioides posadasii, Coccidioides immitis, Bacterial vaginosis, Chlamydia trachomatis, Cytomegalovirus, Granuloma inguinale, Hemophilus ducreyi, Neisseria gonorrhea, Treponema pallidum, Trichomonas vaginalis, or any other infectious disease

25 known in the art that is not listed herein.

[0082] In another embodiment, the infectious disease is a livestock infectious disease. In another embodiment, livestock diseases can be transmitted to man and are called “zoonotic diseases.” In another embodiment, these diseases include, but are not limited to, Foot and mouth disease, West Nile Virus, rabies, canine parvovirus, feline leukemia virus, equine

30 influenza virus, infectious bovine rhinotracheitis (IBR), pseudorabies, classical swine fever (CSF), IBR, caused by bovine herpesvirus type 1 (BHV-1) infection of cattle, and pseudorabies (Aujeszky's disease) in pigs, toxoplasmosis, anthrax, vesicular stomatitis virus, rhodococcus equi, Tularemia, Plague (Yersinia pestis), trichomonas.

[0083] In another embodiment, the disease provided herein is a respiratory or inflammatory disease. In another embodiment, the respiratory or inflammatory disease is chronic obstructive pulmonary disease (COPD). In another embodiment, the disease is asthma.

[0084] In one embodiment, live attenuated *Listeria* strains are capable of alleviating asthma symptoms without co-administration of other therapeutic agents, such as anti-inflammatory agents or bronchodilators. In another embodiment, the methods provided herein further comprise the step of co-administering to a subject a live attenuated *Listeria* strain and one or more therapeutic agents. In another embodiment, the therapeutic agent is an anti-asthmatic agent. In another embodiment, the agent is an anti-inflammatory agent, a non-steroidal anti-inflammatory agent, an antibiotic, an anticholinergic agent, a bronchodilator, a corticosteroid, a short-acting beta-agonist, a long-acting beta-agonist, combination inhalers, an antihistamine, or combinations thereof.

[0085] In one embodiment, a disease is a cancer or a tumor. In one embodiment, the tumor is cancerous. In another embodiment, the cancer is breast cancer. In another embodiment, the cancer is a cervical cancer. In another embodiment, the cancer is a Her2 containing cancer. In another embodiment, the cancer is a melanoma. In another embodiment, the cancer is pancreatic cancer. In another embodiment, the cancer is ovarian cancer. In another embodiment, the cancer is gastric cancer. In another embodiment, the cancer is a carcinomatous lesion of the pancreas. In another embodiment, the cancer is pulmonary adenocarcinoma. In another embodiment, it is a glioblastoma multiforme. In another embodiment, the cancer is colorectal adenocarcinoma. In another embodiment, the cancer is pulmonary squamous adenocarcinoma. In another embodiment, the cancer is gastric adenocarcinoma. In another embodiment, the cancer is an ovarian surface epithelial neoplasm (e.g. a benign, proliferative or malignant variety thereof). In another embodiment, the cancer is an oral squamous cell carcinoma. In another embodiment, the cancer is non-small-cell lung carcinoma. In another embodiment, the cancer is an endometrial carcinoma. In another embodiment, the cancer is a bladder cancer. In another embodiment, the cancer is a head and neck cancer. In another embodiment, the cancer is a prostate carcinoma. In another embodiment, the cancer is oropharyngeal cancer. In another embodiment, the cancer is lung cancer. In another embodiment, the cancer is anal cancer. In another embodiment, the cancer is colorectal cancer. In another embodiment, the cancer is esophageal cancer. The cervical tumor targeted by methods of the present invention is, in another embodiment, a squamous cell carcinoma. In another embodiment, the cervical tumor is an adenocarcinoma. In another embodiment, the cervical tumor is an adenosquamous carcinoma. In another embodiment, the

cervical tumor is a small cell carcinoma. In another embodiment, the cervical tumor is any other type of cervical tumor known in the art.

[0086] A cervical tumor targeted by methods of the present invention is, in one embodiment, a squamous cell carcinoma. In another embodiment, the cervical tumor is an adenocarcinoma.

5 In another embodiment, the cervical tumor is an adenosquamous carcinoma. In another embodiment, the cervical tumor is a small cell carcinoma. In another embodiment, the cervical tumor is any other type of cervical tumor known in the art. Each possibility represents a separate embodiment of the present invention.

[0087] In one embodiment, the terms “tumor antigen” “antigenic polypeptide,” or “foreign
10 antigen” are used interchangeably herein and include tumor antigens, tumor-associated antigens, angiogenic antigens, or infectious disease antigens. In another embodiment, an antigen provided herein is a self-antigen that is present in the host but the host does not elicit an immune response against it because of immunologic tolerance.

[0088] In one embodiment, the antigen is Human Papilloma Virus-E7 (HPV-E7) antigen,
15 which in one embodiment, is from HPV16 (in one embodiment, GenBank Accession No. AAD33253) and in another embodiment, from HPV18 (in one embodiment, GenBank Accession No. P06788). In another embodiment, the antigenic polypeptide is HPV-E6, which in one embodiment, is from HPV16 (in one embodiment, GenBank Accession No. AAD33252, AAM51854, AAM51853, or AAB67615) and in another embodiment, from
20 HPV18 (in one embodiment, GenBank Accession No. P06463). In another embodiment, the antigenic polypeptide is a Her/2-neu antigen. In another embodiment, the antigenic polypeptide is Prostate Specific Antigen (PSA) (in one embodiment, GenBank Accession No. CAD30844, CAD54617, AAA58802, or NP_001639). In another embodiment, the antigenic polypeptide is Stratum Corneum Chymotryptic Enzyme (SCCE) antigen (in one embodiment,
25 GenBank Accession No. AAK69652, AAK69624, AAG33360, AAF01139, or AAC37551). In another embodiment, the antigenic polypeptide is Wilms tumor antigen 1, which in another embodiment is WT-1 Telomerase (GenBank Accession. No. P49952, P22561, NP_659032, CAC39220.2, or EAW68222.1). In another embodiment, the antigenic polypeptide is hTERT or Telomerase (GenBank Accession. No. NM003219 (variant 1), NM198255 (variant 2), NM
30 198253 (variant 3), or NM 198254 (variant 4). In another embodiment, the antigenic polypeptide is Proteinase 3 (in one embodiment, GenBank Accession No. M29142, M75154, M96839, X55668, NM 00277, M96628 or X56606). In another embodiment, the antigenic polypeptide is Tyrosinase Related Protein 2 (TRP2) (in one embodiment, GenBank Accession

No. NP_001913, ABI73976, AAP33051, or Q95119). In another embodiment, the antigenic polypeptide is High Molecular Weight Melanoma Associated Antigen (HMW-MAA) (in one embodiment, GenBank Accession No. NP_001888, AAI28111, or AAQ62842). In another embodiment, the antigenic polypeptide is Testisin (in one embodiment, GenBank Accession
5 No. AAF79020, AAF79019, AAG02255, AAK29360, AAD41588, or NP_659206). In another embodiment, the antigenic polypeptide is NY-ESO-1 antigen (in one embodiment, GenBank Accession No. CAA05908, P78358, AAB49693, or NP_640343). In another embodiment, the antigenic polypeptide is PSCA (in one embodiment, GenBank Accession No. AAH65183, NP_005663, NP_082492, O43653, or CAB97347). In another embodiment,
10 the antigenic polypeptide is Interleukin (IL) 13 Receptor alpha (in one embodiment, GenBank Accession No. NP_000631, NP_001551, NP_032382, NP_598751, NP_001003075, or NP_999506). In another embodiment, the antigenic polypeptide is Carbonic anhydrase IX (CAIX) (in one embodiment, GenBank Accession No. CAI13455, CAI10985, EAW58359, NP_001207, NP_647466, or NP_001101426). In another embodiment, the antigenic
15 polypeptide is carcinoembryonic antigen (CEA) (in one embodiment, GenBank Accession No. AAA66186, CAA79884, CAA66955, AAA51966, AAD15250, or AAA51970.). In another embodiment, the antigenic polypeptide is MAGE-A (in one embodiment, GenBank Accession No. NP_786885, NP_786884, NP_005352, NP_004979, NP_005358, or NP_005353). In another embodiment, the antigenic polypeptide is survivin (in one embodiment,
20 GenBank Accession No. AAC51660, AAY15202, ABF60110, NP_001003019, or NP_001082350). In another embodiment, the antigenic polypeptide is GP100 (in one embodiment, GenBank Accession No. AAC60634, YP_655861, or AAB31176). In another embodiment, the antigenic polypeptide is any other antigenic polypeptide known in the art. In another embodiment, the antigenic peptide of the compositions and methods of the present invention
25 comprise an immunogenic portion of the antigenic polypeptide. Each possibility represents a separate embodiment of the present invention.

[0089] In another embodiment, the antigen is telomerase (TERT). In another embodiment, the antigen is LMP-1. In another embodiment, the antigen is p53. In another embodiment, the antigen is mesothelin. In another embodiment, the antigen is EGFRVIII. In another
30 embodiment, the antigen is carboxic anhydrase IX (CAIX). In another embodiment, the antigen is PSMA. In another embodiment, the antigen is HMW-MAA. In another embodiment, the antigen is HIV-1 Gag. In another embodiment, the antigen is Tyrosinase related protein 2. In another embodiment, the antigen is selected from Her-2, HIV-1 Gag, LMP-1, p53, PSMA, carcinoembryonic antigen (CEA), LMP-1, kallikrein-related peptidase 3

(KLK3), KLK9, Muc, Tyrosinase related protein 2, Muc1, FAP, IL-13R alpha 2, PSA (prostate-specific antigen), gp-100, heat-shock protein 70 (HSP-70), beta-HCG, EGFR-III, Granulocyte colony-stimulating factor (G-CSF), Angiogenin, Angiopoietin-1, Del-1, Fibroblast growth factors: acidic (aFGF) or basic (bFGF), Follistatin, Granulocyte colony-stimulating factor (G-CSF), Hepatocyte growth factor (HGF)/scatter factor (SF), Interleukin-8 (IL-8), Leptin, Midkine, Placental growth factor, Platelet-derived endothelial cell growth factor (PD-ECGF), Platelet-derived growth factor-BB (PDGF-BB), Pleiotrophin (PTN), Progranulin, Proliferin, Transforming growth factor-alpha (TGF-alpha), Transforming growth factor-beta (TGF-beta), Tumor necrosis factor-alpha (TNF-alpha), Vascular endothelial growth factor (VEGF)/vascular permeability factor (VPF), VEGFR, VEGFR2 (KDR/FLK-1) or a fragment thereof, FLK-1 or an epitope thereof, FLK-E1, FLK-E2, FLK-I1, endoglin or a fragment thereof, Neuropilin 1 (NRP-1), Angiopoietin 1 (Ang1), Tie2, Platelet-derived growth factor (PDGF), Platelet-derived growth factor receptor (PDGFR), Transforming growth factor-beta (TGF- β), endoglin, TGF- β receptors, monocyte chemotactic protein-1 (MCP-1), VE-cadherin, CD31, ephrin, ICAM-1, V-CAM-1, VAP-1, E-selectin, plasminogen activators, plasminogen activator inhibitor-1, Nitric oxide synthase (NOS), COX-2, AC133, or Id1/Id3, Angiopoietin 3, Angiopoietin 4, Angiopoietin 6, CD105, EDG, HHT1, ORW, ORW1 or a TGFbeta co-receptor, or a combination thereof. In another embodiment, the antigen is a chimeric Her2/neu antigen as disclosed in US Patent Application Publication No. 2011/0142791, which is incorporated by reference herein in its entirety. The use of fragments of antigens provided herein is also encompassed by the present invention.

[0090] In another embodiment, the tumor antigen provided herein is a tumor-associated antigen, which in one embodiment, is one of the following tumor antigens: a MAGE (Melanoma-Associated Antigen E) protein, e.g. MAGE 1, MAGE 2, MAGE 3, MAGE 4, a tyrosinase; a mutant ras protein; a mutant p53 protein; p97 melanoma antigen, a ras peptide or p53 peptide associated with advanced cancers; the HPV 16/18 antigens associated with cervical cancers, KLH antigen associated with breast carcinoma, CEA (carcinoembryonic antigen) associated with colorectal cancer, a MART1 antigen associated with melanoma, or the PSA antigen associated with prostate cancer. In another embodiment, the antigen for the compositions and methods provided herein are melanoma-associated antigens, which in one embodiment are TRP-2, MAGE-1, MAGE-3, gp-100, tyrosinase, HSP-70, beta-HCG, or a combination thereof. It is to be understood that a skilled artisan would be able to use any heterologous antigen not mentioned herein but known in the art for use in the methods and compositions provided herein. It is also to be understood that the present invention provides,

but is not limited by, an attenuated *Listeria* comprising a nucleic acid that encodes at least one of the antigens disclosed herein. The present invention encompasses nucleic acids encoding mutants, muteins, splice variants, fragments, truncated variants, soluble variants, extracellular domains, intracellular domains, mature sequences, and the like, of the disclosed antigens.

5 Provided are nucleic acids encoding epitopes, oligo- and polypeptides of these antigens. Also provided are codon optimized embodiments, that is, optimized for expression in *Listeria*. The cited references, GenBank Acc. Nos., and the nucleic acids, peptides, and polypeptides disclosed herein, are all incorporated herein by reference in their entirety. In another embodiment, the selected nucleic acid sequence can encode a full length or a truncated gene, a
10 fusion or tagged gene, and can be a cDNA, a genomic DNA, or a DNA fragment, preferably, a cDNA. It can be mutated or otherwise modified as desired. These modifications include codon optimizations to optimize codon usage in the selected host cell or bacteria, i.e. *Listeria*. The selected sequence can also encode a secreted, cytoplasmic, nuclear, membrane bound or cell surface polypeptide.

15 [0091] In one embodiment, vascular endothelial growth factor (VEGF) is an important signaling protein involved in both vasculogenesis (the formation of the embryonic circulatory system) and angiogenesis (the growth of blood vessels from pre-existing vasculature). In one embodiment, VEGF activity is restricted mainly to cells of the vascular endothelium, although it does have effects on a limited number of other cell types (e.g. stimulation
20 monocyte/macrophage migration). *In vitro*, VEGF has been shown to stimulate endothelial cell mitogenesis and cell migration. VEGF also enhances microvascular permeability and is sometimes referred to as vascular permeability factor.

[0092] In one embodiment, all of the members of the VEGF family stimulate cellular responses by binding to tyrosine kinase receptors (the VEGFRs) on the cell surface, causing them to
25 dimerize and become activated through transphosphorylation. The VEGF receptors have an extracellular portion consisting of 7 immunoglobulin-like domains, a single transmembrane spanning region and an intracellular portion containing a split tyrosine-kinase domain.

[0093] In one embodiment, VEGF-A is a VEGFR-2 (KDR/Flk-1) ligand as well as a VEGFR-1 (Flt-1) ligand. In one embodiment, VEGFR- mediates almost all of the known cellular responses
30 to VEGF. The function of VEGFR-1 is less well defined, although it is thought to modulate VEGFR-2 signaling, in one embodiment, via sequestration of VEGF from VEGFR-2 binding, which in one embodiment, is particularly important during vasculogenesis in the embryo. In one

embodiment, VEGF-C and VEGF-D are ligands of the VEGFR-3 receptor, which in one embodiment, mediates lymphangiogenesis.

[0094] In one embodiment, the antigen of the present invention is a VEGF receptor or a fragment thereof, which in one embodiment, is a VEGFR-2 and, in another embodiment, a VEGFR-1, and, in another embodiment, VEGFR-3.

[0095] In one embodiment, vascular Endothelial Growth Factor Receptor 2 (VEGFR2) is highly expressed on activated endothelial cells (ECs) and participates in the formation of new blood vessels. In one embodiment, VEGFR2 binds all 5 isoforms of VEGF. In one embodiment, signaling of VEGF through VEGFR2 on ECs induces proliferation, migration, and eventual differentiation. In one embodiment, the mouse homologue of VEGFR2 is the fetal liver kinase gene-1 (Flk-1), which is a strong therapeutic target, and has important roles in tumor growth, invasion, and metastasis. In one embodiment, VEGFR2 is also referred to as kinase insert domain receptor (a type III receptor tyrosine kinase) (KDR), cluster of differentiation 309 (CD309), FLK1, Ly73, Krd-1, VEGFR, VEGFR-2, or 6130401C07.

[0096] In other embodiments, the antigen is derived from a fungal pathogen, bacteria, parasite, helminth, or viruses. In other embodiments, the antigen is selected from tetanus toxoid, hemagglutinin molecules from influenza virus, diphtheria toxoid, HIV gp120, HIV gag protein, IgA protease, insulin peptide B, *Spongospora subterranea* antigen, vibriose antigens, *Salmonella* antigens, pneumococcus antigens, respiratory syncytial virus antigens, *Haemophilus influenza* outer membrane proteins, *Helicobacter pylori* urease, *Neisseria meningitidis* pilins, *N. gonorrhoeae* pilins, the melanoma-associated antigens (TRP-2, MAGE-1, MAGE-3, gp-100, tyrosinase, MART-1, HSP-70, beta-HCG), human papilloma virus antigens E1 and E2 from type HPV-16, -18, -31, -33, -35 or -45 human papilloma viruses, the tumor antigens CEA, the ras protein, mutated or otherwise, the p53 protein, mutated or otherwise, Muc1, or pSA.

[0097] In other embodiments, the antigen is associated with one of the following diseases; cholera, diphtheria, *Haemophilus*, hepatitis A, hepatitis B, influenza, measles, meningitis, mumps, pertussis, small pox, pneumococcal pneumonia, polio, rabies, rubella, tetanus, tuberculosis, typhoid, Varicella-zoster, whooping cough³ yellow fever, the immunogens and antigens from Addison's disease, allergies, anaphylaxis, Bruton's syndrome, cancer, including solid and blood borne tumors, eczema, Hashimoto's thyroiditis, polymyositis, dermatomyositis, type 1 diabetes mellitus, acquired immune deficiency syndrome, transplant rejection, such as kidney, heart, pancreas, lung, bone, and liver transplants, Graves' disease, polyendocrine autoimmune disease, hepatitis, microscopic polyarteritis, polyarteritis nodosa, pemphigus, primary biliary cirrhosis, pernicious anemia, coeliac disease, antibody-mediated

nephritis, glomerulonephritis, rheumatic diseases, systemic lupus erthematosus, rheumatoid arthritis, seronegative spondylarthritides, rhinitis, sjogren's syndrome, systemic sclerosis, sclerosing cholangitis, Wegener's granulomatosis, dermatitis herpetiformis, psoriasis, vitiligo, multiple sclerosis, encephalomyelitis, Guillain-Barre syndrome, myasthenia gravis, Lambert-
 5 Eaton syndrome, sclera, episclera, uveitis, chronic mucocutaneous candidiasis, urticaria, transient hypogammaglobulinemia of infancy, myeloma, X-linked hyper IgM syndrome, Wiskott-Aldrich syndrome, ataxia telangiectasia, autoimmune hemolytic anemia, autoimmune thrombocytopenia, autoimmune neutropenia, Waldenstrom's macroglobulinemia, amyloidosis, chronic lymphocytic leukemia, non-Hodgkin's lymphoma, malarial
 10 circumsporozite protein, microbial antigens, viral antigens, autoantigens, and lesteriosis.

[0098] In another embodiment, an HPV E6 antigen is utilized instead of or in addition to an E7 antigen in a method of the present invention for treating, protecting against, or inducing an immune response against a cervical cancer.

[0099] In another embodiment, an ActA protein fragment is utilized instead of or in addition
 15 to an LLO fragment in a method of the present invention for treating, protecting against, or inducing an immune response against a cervical cancer.

[00100] In another embodiment, a PEST amino acid sequence-containing protein fragment is utilized instead of or in addition to an LLO fragment in a method of the present invention for treating, protecting against, or inducing an immune response against a cervical cancer.

[00101] In another embodiment, the present invention provides a method for inducing an anti-
 20 E7 cytotoxic T cell (CTL) response in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an LLO protein and an HPV E7 antigen, thereby inducing an anti-E7 CTL response in a subject. In another embodiment,
 25 the recombinant *Listeria* strain comprises a plasmid that encodes the recombinant polypeptide. In another embodiment, the method further comprises the step of boosting the subject with a recombinant *Listeria* strain of the present invention. In another embodiment, the method further comprises the step of boosting the subject with an immunogenic composition comprising an E7 antigen. In another embodiment, the method further comprises
 30 the step of boosting the subject with an immunogenic composition that directs a cell of the subject to express an E7 antigen. In another embodiment, the CTL response is capable of therapeutic efficacy against an HPV-mediated disease, disorder, or symptom. In another embodiment, the CTL response is capable of prophylactic efficacy against an HPV-mediated

disease, disorder, or symptom. Each possibility represents a separate embodiment of the present invention.

[00102] In another embodiment, the present invention provides a method of treating or ameliorating an HPV-mediated disease, disorder, or symptom in a subject, comprising the step of administering to the subject a recombinant *Listeria* strain, the recombinant *Listeria* strain comprising a recombinant polypeptide comprising an N-terminal fragment of an LLO protein and an HPV E7 antigen, whereby the recombinant *Listeria* strain induces an immune response against the E7 antigen, thereby treating or ameliorating an HPV-mediated disease, disorder, or symptom in a subject. In another embodiment, the subject is a human subject. In another embodiment, the subject is a non-human mammal. In another embodiment, the subject is any other type of subject known in the art. Each possibility represents a separate embodiment of the present invention.

[00103] In another embodiment, the HPV-mediated disease, disorder, or symptom is genital warts. In another embodiment, the HPV-mediated disease, disorder, or symptom is non-genital warts. In another embodiment, the HPV-mediated disease, disorder, or symptom is a respiratory papilloma. In another embodiment, the HPV-mediated disease, disorder, or symptom is any other HPV-mediated disease, disorder, or symptom known in the art. Each possibility represents a separate embodiment of the present invention.

[00104] The antigen of methods and compositions of the present invention is, in another embodiment, an HPV E7 protein. In another embodiment, the antigen is an HPV E6 protein. In another embodiment, the antigen is any other HPV protein known in the art. Each possibility represents a separate embodiment of the present invention.

[00105] “E7 antigen” refers, in another embodiment, to an E7 protein. In another embodiment, the term refers to an E7 fragment. In another embodiment, the term refers to an E7 peptide. In another embodiment, the term refers to any other type of E7 antigen known in the art. Each possibility represents a separate embodiment of the present invention.

[00106] The E7 protein of methods and compositions of the present invention is, in another embodiment, an HPV 16 E7 protein. In another embodiment, the E7 protein is an HPV-18 E7 protein. In another embodiment, the E7 protein is an HPV-31 E7 protein. In another embodiment, the E7 protein is an HPV-35 E7 protein. In another embodiment, the E7 protein is an HPV-39 E7 protein. In another embodiment, the E7 protein is an HPV-45 E7 protein. In another embodiment, the E7 protein is an HPV-51 E7 protein. In another embodiment, the E7 protein is an HPV-52 E7 protein. In another embodiment, the E7 protein is an HPV-58 E7 protein. In another embodiment, the E7 protein is an E7 protein of a high-risk HPV type. In

another embodiment, the E7 protein is an E7 protein of a mucosal HPV type. Each possibility represents a separate embodiment of the present invention.

[00107] “E6 antigen” refers, in another embodiment, to an E6 protein. In another embodiment, the term refers to an E6 fragment. In another embodiment, the term refers to an E6 peptide. In another embodiment, the term refers to any other type of E6 antigen known in the art. Each possibility represents a separate embodiment of the present invention.

[00108] The E6 protein of methods and compositions of the present invention is, in another embodiment, an HPV 16 E6 protein. In another embodiment, the E6 protein is an HPV-18 E6 protein. In another embodiment, the E6 protein is an HPV-31 E6 protein. In another embodiment, the E6 protein is an HPV-35 E6 protein. In another embodiment, the E6 protein is an HPV-39 E6 protein. In another embodiment, the E6 protein is an HPV-45 E6 protein. In another embodiment, the E6 protein is an HPV-51 E6 protein. In another embodiment, the E6 protein is an HPV-52 E6 protein. In another embodiment, the E6 protein is an HPV-58 E6 protein. In another embodiment, the E6 protein is an E6 protein of a high-risk HPV type. In another embodiment, the E6 protein is an E6 protein of a mucosal HPV type. Each possibility represents a separate embodiment of the present invention.

[00109] In one embodiment, combinations of the E6 and E7 antigens are contemplated to fall within the scope of a “heterologous antigen” provided herein.

[00110] The immune response induced by methods and compositions of the present invention is, in another embodiment, a T cell response. In another embodiment, the immune response comprises a cytotoxic T cell response. In another embodiment, the immune response comprises a T cell response. In another embodiment, the response is a CD8⁺ T cell response. In another embodiment, the response comprises a CD8⁺ T cell response. Each possibility represents a separate embodiment of the present invention.

[00111] The N-terminal LLO protein fragment of methods and compositions of the present invention comprises, in another embodiment, SEQ ID No: 2. In another embodiment, the fragment comprises an LLO signal peptide. In another embodiment, the fragment comprises SEQ ID No: 2. In another embodiment, the fragment consists approximately of SEQ ID No: 2. In another embodiment, the fragment consists essentially of SEQ ID No: 2. In another embodiment, the fragment corresponds to SEQ ID No: 2. In another embodiment, the fragment is homologous to SEQ ID No: 2. In another embodiment, the fragment is homologous to a fragment of SEQ ID No: 2. The Δ LLO used in some of the Examples was 416 AA long (exclusive of the signal sequence), as 88 residues from the amino terminus which is inclusive of the activation domain containing cysteine 484 were truncated. It will be

clear to those skilled in the art that any Δ LLO without the activation domain, and in particular without cysteine 484, are suitable for methods and compositions of the present invention. In another embodiment, fusion of an E7 and/or E6 antigen to any Δ LLO, including a PEST amino acid AA sequence, SEQ ID NO: 1, enhances cell mediated and anti-tumor immunity of the antigen. Each possibility represents a separate embodiment of the present invention.

[00112] The LLO protein utilized to construct vaccines of the present invention has, in another embodiment, the sequence:

MKKIMLVFITLILVSLPIAQQTEAKDASAFNKENSISSMAPPASPPASPKTPIEKKHADE
IDKYIQGLDYNKNNVLVYHGDAVTNVPPrKGYKDGNEYIVVEKKKKKSINQNNADIQ
VVNAISSLTYPGALVKANSELVENQPDVLPVKRDSLTLSDLPGMTNQDNKIVVKNA
TKSNVNNAVNTLVERWNEKYAQAYPNVSAKIDYDDEMAYSESQLIKFGTAFKAV
NNSLNVNFGAISEGKMQUEEVISFKQIYYNVNVNEPTRPSRFFGKAVTKEQLQALGVN
AENPPAYISSVAYGRQVYLKLSTNSHSTKVKAADFDAVSGKSVSGDVELTNIKNSSF
KAVIYGGSAKDEVQIIDGNLGLDRDILKKGATFNRETPGVPIAYTTNFLKDNELAVIK
NNSEYIETTSKAYTDGKINIDHSGGYVAQFNISWDEVNYDPEGNEIVQHKNWSENNK
SKLAHFTSSIYLPGNARNINVYAKECTGLAWWWRTVIDDRNLPLVKNRNISIWGTT
LYPKYSNKVDNPIE (GenBank Accession No. P13128; SEQ ID NO: 3; nucleic acid
sequence is set forth in GenBank Accession No. X15127). The first 25 AA of the proprotein
corresponding to this sequence are the signal sequence and are cleaved from LLO when it is
secreted by the bacterium. Thus, in this embodiment, the full length active LLO protein is 504
residues long. In another embodiment, a full length LLO protein has an amino acid sequence
of any full length wild-type LLO protein known in the art. In another embodiment, SEQ ID
NO: 3 is used as the source of the LLO fragment incorporated in a vaccine of the present
invention. Each possibility represents a separate embodiment of the present invention.

[00113] In another embodiment, the N-terminal fragment of an LLO protein utilized in compositions and methods of the present invention has the sequence:

MKKIMLVFITLILVSLPIAQQTEAKDASAFNKENSISSVAPPASPPASPKTPIEKKHADEI
DKYIQGLDYNKNNVLVYHGDAVTNVPPrKGYKDGNEYIVVEKKKKKSINQNNADIQV
VNAISSLTYPGALVKANSELVENQPDVLPVKRDSLTLSDLPGMTNQDNKIVVKNA
KSNVNNAVNTLVERWNEKYAQAYSNVSAKIDYDDEMAYSESQLIKFGTAFKAVN
NNSLNVNFGAISEGKMQUEEVISFKQIYYNVNVNEPTRPSRFFGKAVTKEQLQALGVNA
ENPPAYISSVAYGRQVYLKLSTNSHSTKVKAADFDAVSGKSVSGDVELTNIKNSSFK

AVIYGGSAKDEVQIIDGNLGLDLRDILKKGATFNRETPGVPIAYTTNFLKDNE LAVIKN
NSEYIETTSKAYTDGKINIDHSGGYVAQFNISWDEVNYD (SEQ ID NO: 2).

[00114] In another embodiment, the LLO fragment has the sequence:

MKKIMLVFITLILVSLPIAQQTEAKDASAFNKENSISVAPPASPPASPKTPIEKKHADI
5 DKYIQGLDYNKNNVLVYHGDAVTNVPVRKGYKDGNEYIVVEKKKKKSINQNNADIQV
VNAISSLTYPGALVKANSELVENQPDVLPVKRDSLTLSDLPGMTNQDNKIVVKNAT
KSNVNNNAVNTLVERWNEKYAQAYSNVSAKIDYDDEMAYSESQLIKFGTAFKAVN
NSLNVNFGAISEGKMQUEEVISFKQIYYNVNVNEPTRPSRFFGKAVTKEQLQALGVNA
ENPPAYISSVAYGRQVYLKLSTNSHSTKVKAADFDAVSGKSVSGDVELTNIKNSSFK
10 AVIYGGSAKDEVQIIDGNLGLDLRDILKKGATFNRETPGVPIAYTTNFLKDNE LAVIKN
NSEYIETTSKAYTD (SEQ ID NO: 4).

[00115] In one embodiment, “Listeriolysin O protein,” or “LLO protein,” refer to a wild-type LLO protein unless stated to be a fragment of the same. In another embodiment, “truncated LLO” or “ΔLLO” refers to a fragment of LLO that comprises the PEST amino acid domain.

15 In another embodiment, the terms refer to an LLO fragment that comprises a PEST sequence. In another embodiment, the terms refer to an LLO fragment that comprises a putative PEST sequence.

[00116] In another embodiment, the terms refer to an LLO fragment that does not contain the activation domain at the carboxy terminus and does not include cysteine 484. In another
20 embodiment, the terms refer to an LLO fragment that is not hemolytic. In another embodiment, the LLO fragment is rendered non-hemolytic by deletion or mutation of the activation domain. In another embodiment, the LLO fragment is rendered non-hemolytic by deletion or mutation of cysteine 484. In another embodiment, the LLO fragment is rendered non-hemolytic by deletion or mutation at another location. Each possibility represents a
25 separate embodiment of the present invention.

[00117] In another embodiment, the LLO fragment consists of about the first 441 AA of a wild-type LLO protein. In another embodiment, the LLO fragment consists of about the first 420 AA of LLO. In another embodiment, the LLO fragment is a non-hemolytic form of the LLO protein.

30 [00118] In another embodiment, the LLO fragment contains residues of a homologous LLO protein that correspond to one of the above AA ranges. The residue numbers need not, in another embodiment, correspond exactly with the residue numbers enumerated above; e.g. if

the homologous LLO protein has an insertion or deletion, relative to an LLO protein utilized herein, then the residue numbers can be adjusted accordingly.

[00119] In another embodiment, the LLO fragment is any other LLO fragment known in the art. Each possibility represents a separate embodiment of the present invention.

5 [00120] In another embodiment, the recombinant polypeptide of methods of the present invention is expressed by the recombinant *Listeria* strain. In another embodiment, the expression is mediated by a nucleotide molecule carried by the recombinant *Listeria* strain.

[00121] In another embodiment, the recombinant *Listeria* strain expresses the recombinant polypeptide by means of a plasmid that encodes the recombinant polypeptide. In another
10 embodiment, the plasmid comprises a gene encoding a bacterial transcription factor. In another embodiment, the plasmid encodes a *Listeria* transcription factor. In another embodiment, the transcription factor is PrfA. In another embodiment, the PrfA is a mutant PrfA. In another embodiment, the PrfA contains a D133V amino acid mutation. In another embodiment, the transcription factor is any other transcription factor known in the art. In
15 another embodiment, the mutant PrfA encoded by said plasmid complements a genomic *prfA* mutation, deletion or inactivation in said *Listeria*. In another embodiment, the mutant PrfA encoded by said plasmid restores partial PrfA function in said *Listeria* having a genomic *prfA* mutation, deletion or inactivation. Each possibility represents a separate embodiment of the present invention.

20 [00122] In another embodiment, a plasmid comprised by a recombinant *Listeria* provided herein comprises an open reading frame encoding a metabolic enzyme. In another embodiment, the plasmid comprises a third open reading frame encoding a metabolic enzyme. In another embodiment, the metabolic enzyme is a bacterial metabolic enzyme. In another embodiment, the metabolic enzyme is a *Listerial* metabolic enzyme. In another embodiment,
25 the metabolic enzyme is an amino acid metabolism enzyme. In another embodiment, the amino acid metabolism gene is involved in a cell wall synthesis pathway. In another embodiment, the metabolic enzyme is the product of a D-amino acid aminotransferase gene (*dat*). In another embodiment, the metabolic enzyme is the product of an alanine racemase gene (*dal*). In another embodiment, the metabolic enzyme is any other metabolic enzyme
30 known in the art. In another embodiment, the plasmid carries an open reading frame encoding a *dal* protein. In another embodiment, the plasmid carries an open reading frame encoding a *dat* protein. In another embodiment, the plasmid carries an open reading frame encoding a *dal* and *dat* protein. In another embodiment, when the plasmid carries an open reading frame encoding a *dal* and/or *dat* protein, it is to complement a *dal/dat* mutation in a recombinant

Listeria strain. Hence, *dal/dat* recombinant *Listerias* are also envisioned for use in the present invention. In another embodiment, the recombinant *Listeria* provided herein comprises a *dal/dat* mutation in addition to any other mutation further described herein. Each possibility represents a separate embodiment of the present invention.

5 [00123] In another embodiment, a *Listeria* strain provided herein is deficient in an AA metabolism enzyme. In another embodiment, the *Listeria* strain is deficient in a D-glutamic acid synthase gene. In another embodiment, the *Listeria* strain is deficient in the *dat* gene. In another embodiment, the *Listeria* strain is deficient in the *dal* gene. In another embodiment, the *Listeria* strain is deficient in the *dga* gene. In another embodiment, the *Listeria* strain is
10 deficient in a gene involved in the synthesis of diaminopimelic acid (DAP). In another embodiment, the *Listeria* strain is deficient in a gene involved in the synthesis of Cysteine synthase A (*CysK*). In another embodiment, the gene is vitamin-B12 independent methionine synthase. In another embodiment, the gene is *trpA*. In another embodiment, the gene is *trpB*. In another embodiment, the gene is *trpE*. In another embodiment, the gene is *asnB*. In another
15 embodiment, the gene is *gltD*. In another embodiment, the gene is *gltB*. In another embodiment, the gene is *leuA*. In another embodiment, the gene is *argG*. In another embodiment, the gene is *thrC*. In another embodiment, the *Listeria* strain is deficient in one or more of the genes described hereinabove.

[00124] In another embodiment, a *Listeria* strain provided herein is deficient in a synthase
20 gene. In another embodiment, the gene is an AA synthesis gene. In another embodiment, the gene is *folP*. In another embodiment, the gene is dihydrouridine synthase family protein. In another embodiment, the gene is *ispD*. In another embodiment, the gene is *ispF*. In another embodiment, the gene is phosphoenolpyruvate synthase. In another embodiment, the gene is *hisF*. In another embodiment, the gene is *hisH*. In another embodiment, the gene is *fliI*. In
25 another embodiment, the gene is ribosomal large subunit pseudouridine synthase. In another embodiment, the gene is *ispD*. In another embodiment, the gene is bifunctional GMP synthase/glutamine amidotransferase protein. In another embodiment, the gene is *cobS*. In another embodiment, the gene is *cobB*. In another embodiment, the gene is *cbiD*. In another embodiment, the gene is uroporphyrin-III C-methyltransferase/ uroporphyrinogen-III
30 synthase. In another embodiment, the gene is *cobQ*. In another embodiment, the gene is *uppS*. In another embodiment, the gene is *truB*. In another embodiment, the gene is *dxs*. In another embodiment, the gene is *mvaS*. In another embodiment, the gene is *dapA*. In another embodiment, the gene is *ispG*. In another embodiment, the gene is *folC*. In another embodiment, the gene is citrate synthase. In another embodiment, the gene is *argJ*. In another

embodiment, the gene is 3-deoxy-7-phosphoheptulonate synthase. In another embodiment, the gene is indole-3-glycerol-phosphate synthase. In another embodiment, the gene is anthranilate synthase/ glutamine amidotransferase component. In another embodiment, the gene is *menB*. In another embodiment, the gene is menaquinone-specific isochorismate synthase. In another embodiment, the gene is phosphoribosylformylglycinamide synthase I or II. In another embodiment, the gene is phosphoribosylaminoimidazole-succinocarboxamide synthase. In another embodiment, the gene is *carB*. In another embodiment, the gene is *carA*. In another embodiment, the gene is *thyA*. In another embodiment, the gene is *mgsA*. In another embodiment, the gene is *aroB*. In another embodiment, the gene is *hepB*. In another embodiment, the gene is *rluB*. In another embodiment, the gene is *ilvB*. In another embodiment, the gene is *ilvN*. In another embodiment, the gene is *alsS*. In another embodiment, the gene is *fabF*. In another embodiment, the gene is *fabH*. In another embodiment, the gene is pseudouridine synthase. In another embodiment, the gene is *pyrG*. In another embodiment, the gene is *truA*. In another embodiment, the gene is *pabB*. In another embodiment, the gene is an atp synthase gene (e.g. *atpC*, *atpD-2*, *aptG*, *atpA-2*, etc).

[00125] In another embodiment, the gene is *phoP*. In another embodiment, the gene is *aroA*. In another embodiment, the gene is *aroC*. In another embodiment, the gene is *aroD*. In another embodiment, the gene is *plcB*.

[00126] In another embodiment, a *Listeria* strain provided herein is deficient in a peptide transporter. In another embodiment, the gene is ABC transporter/ ATP-binding/permease protein. In another embodiment, the gene is oligopeptide ABC transporter/ oligopeptide-binding protein. In another embodiment, the gene is oligopeptide ABC transporter/ permease protein. In another embodiment, the gene is zinc ABC transporter/ zinc-binding protein. In another embodiment, the gene is sugar ABC transporter. In another embodiment, the gene is phosphate transporter. In another embodiment, the gene is ZIP zinc transporter. In another embodiment, the gene is drug resistance transporter of the *EmrB/QacA* family. In another embodiment, the gene is sulfate transporter. In another embodiment, the gene is proton-dependent oligopeptide transporter. In another embodiment, the gene is magnesium transporter. In another embodiment, the gene is formate/nitrite transporter. In another embodiment, the gene is spermidine/putrescine ABC transporter. In another embodiment, the gene is Na/Pi-cotransporter. In another embodiment, the gene is sugar phosphate transporter. In another embodiment, the gene is glutamine ABC transporter. In another embodiment, the gene is major facilitator family transporter. In another embodiment, the gene is glycine betaine/L-proline ABC transporter. In another embodiment, the gene is molybdenum ABC

transporter. In another embodiment, the gene is teichoic acid ABC transporter. In another embodiment, the gene is cobalt ABC transporter. In another embodiment, the gene is ammonium transporter. In another embodiment, the gene is amino acid ABC transporter. In another embodiment, the gene is cell division ABC transporter. In another embodiment, the gene is manganese ABC transporter. In another embodiment, the gene is iron compound ABC transporter. In another embodiment, the gene is maltose/maltodextrin ABC transporter. In another embodiment, the gene is drug resistance transporter of the *Bcr/CflA* family. In another embodiment, the gene is a subunit of one of the above proteins.

[00127] In one embodiment, provided herein is a nucleic acid molecule that is used to transform the *Listeria* in order to arrive at a recombinant *Listeria*. In another embodiment, the nucleic acid provided herein used to transform *Listeria* lacks a virulence gene. In another embodiment, the nucleic acid molecule is integrated into the *Listeria* genome and carries a non-functional virulence gene. In another embodiment, the virulence gene is mutated in the recombinant *Listeria*. In yet another embodiment, the nucleic acid molecule is used to inactivate the endogenous gene present in the *Listeria* genome. In yet another embodiment, the virulence gene is an *actA* gene, an *inlA* gene, and *inlB* gene, an *inlC* gene, *inlJ* gene, a *plbC* gene, a *bsh* gene, or a *prfA* gene. It is to be understood by a skilled artisan, that the virulence gene can be any gene known in the art to be associated with virulence in the recombinant *Listeria*.

[00128] In one embodiment, a live attenuated *Listeria* provided herein is a recombinant *Listeria*. In another embodiment, a recombinant *Listeria* provided herein comprises a mutation of a genomic internalin C (*inlC*) gene. In another embodiment, the recombinant *Listeria* comprises a mutation or a deletion of a genomic *actA* gene and a genomic internalin C gene. In one embodiment, translocation of *Listeria* to adjacent cells is inhibited by the deletion of the *actA* gene and/or the *inlC* gene, which are involved in the process, thereby resulting in unexpectedly high levels of attenuation with increased immunogenicity and utility as a strain backbone. Each possibility represents a separate embodiment of the present invention.

[00129] It will be appreciated by a skilled artisan that the term "attenuation," may encompass a diminution in the ability of the bacterium to cause disease in an animal. In other words, for example the pathogenic characteristics of the attenuated *Listeria* strain have been lessened compared with wild-type *Listeria*, although the attenuated *Listeria* is capable of growth and maintenance in culture. Using as an example the intravenous inoculation of Balb/c mice with an attenuated *Listeria*, the lethal dose at which 50% of inoculated animals survive (LD₅₀) is

preferably increased above the LD₅₀ of wild-type *Listeria* by at least about 10-fold, more preferably by at least about 100-fold, more preferably at least about 1,000 fold, even more preferably at least about 10,000 fold, and most preferably at least about 100,000-fold. An attenuated strain of *Listeria* is thus one which does not kill an animal to which it is administered, or is one which kills the animal only when the number of bacteria administered is vastly greater than the number of wild type non-attenuated bacteria which would be required to kill the same animal. An attenuated bacterium should also be construed to mean one which is incapable of replication in the general environment because the nutrient required for its growth is not present therein. Thus, the bacterium is limited to replication in a controlled environment wherein the required nutrient is provided. The attenuated strains of the present invention are therefore environmentally safe in that they are incapable of uncontrolled replication.

[00130] In yet another embodiment, a *Listeria* strain provided herein is an *inlA* mutant, an *inlB* mutant, an *inlC* mutant, an *inlJ* mutant, *prfA* mutant, *actA* mutant, a *dal/dat* mutant, a *prfA* mutant, a *plcB* deletion mutant, or a double mutant lacking both *plcA* and *plcB*. In another embodiment, the *Listeria* comprises a deletion or mutation of these genes individually or in combination. In another embodiment, the *Listeria* provided herein lack each one of genes. In another embodiment, the *Listeria* provided herein lack at least one and up to ten of any gene provided herein, including the *actA*, *prfA*, and *dal/dat* genes. In another embodiment, the *prfA* mutant is a D133V PrfA mutant.

[00131] In one embodiment, the metabolic gene, the virulence gene, etc. is lacking in a chromosome of the *Listeria* strain. In another embodiment, the metabolic gene, virulence gene, etc. is lacking in the chromosome and in any episomal genetic element of the *Listeria* strain. In another embodiment, the metabolic gene, virulence gene, etc. is lacking in the genome of the virulence strain. In one embodiment, the virulence gene is mutated in the chromosome. In another embodiment, the virulence gene is deleted from the chromosome. In another embodiment, the metabolic gene, the virulence gene, etc. is mutated in a chromosome of the *Listeria* strain. In another embodiment, the metabolic gene, virulence gene, etc. is mutated in the chromosome and in any episomal genetic element of the *Listeria* strain. In another embodiment, the metabolic gene, virulence gene, etc. is mutated in the genome of the virulence strain. In another embodiment, the virulence gene is deleted from the chromosome. Each possibility represents a separate embodiment of the present invention.

[00132] In one embodiment, a recombinant *Listeria* strain provided herein is attenuated. In another embodiment, the recombinant *Listeria* lacks the *actA* virulence gene. In another

embodiment, the recombinant *Listeria* lacks the *prfA* virulence gene. In another embodiment, the recombinant *Listeria* lacks the *inlB* gene. In another embodiment, the recombinant *Listeria* lacks both, the *actA* and *inlB* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA* gene. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *inlB* gene. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *inlC* gene. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA* and *inlB* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA* and *inlC* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA*, *inlB*, and *inlC* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA*, *inlB*, and *inlC* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation of the endogenous *actA*, *inlB*, and *inlC* genes. In another embodiment, the recombinant *Listeria* strain provided herein comprises an inactivating mutation in any single gene or combination of the following genes: *actA*, *dal*, *dat*, *inlB*, *inlC*, *prfA*, *plcA*, *plcB*.

[00133] It will be appreciated by a skilled artisan that the term “mutation” and grammatical equivalents thereof, include any type of mutation or modification to the sequence (nucleic acid or amino acid sequence), and includes a deletion mutation, a truncation, an inactivation, a disruption, or a translocation. These types of mutations are readily known in the art.

[00134] In one embodiment, in order to select for auxotrophic bacteria comprising a plasmid encoding a metabolic enzyme or a complementing gene provided herein, transformed auxotrophic bacteria are grown on a media that will select for expression of the amino acid metabolism gene or the complementing gene. In another embodiment, a bacteria auxotrophic for D-glutamic acid synthesis is transformed with a plasmid comprising a gene for D-glutamic acid synthesis, and the auxotrophic bacteria will grow in the absence of D-glutamic acid, whereas auxotrophic bacteria that have not been transformed with the plasmid, or are not expressing the plasmid encoding a protein for D-glutamic acid synthesis, will not grow. In another embodiment, a bacterium auxotrophic for D-alanine synthesis will grow in the absence of D-alanine when transformed and expressing the plasmid of the present invention if the plasmid comprises an isolated nucleic acid encoding an amino acid metabolism enzyme for D-alanine synthesis. Such methods for making appropriate media comprising or lacking

necessary growth factors, supplements, amino acids, vitamins, antibiotics, and the like are well known in the art, and are available commercially (Becton-Dickinson, Franklin Lakes, NJ).

[00135] In another embodiment, once the auxotrophic bacteria comprising the plasmid of the present invention have been selected on appropriate media, the bacteria are propagated in the presence of a selective pressure. Such propagation comprises growing the bacteria in media without the auxotrophic factor. The presence of the plasmid expressing an amino acid metabolism enzyme in the auxotrophic bacteria ensures that the plasmid will replicate along with the bacteria, thus continually selecting for bacteria harboring the plasmid. The skilled artisan, when equipped with the present disclosure and methods herein will be readily able to scale-up the production of the *Listeria* vaccine vector by adjusting the volume of the media in which the auxotrophic bacteria comprising the plasmid are growing.

[00136] The skilled artisan will appreciate that, in another embodiment, other auxotroph strains and complementation systems are adopted for the use with this invention.

[00137] In one embodiment, a recombinant *Listeria* strain provided herein expresses a recombinant polypeptide. In another embodiment, a recombinant *Listeria* strain comprises a plasmid that encodes a recombinant polypeptide. In another embodiment, a recombinant nucleic acid provided herein is in a plasmid in the recombinant *Listeria* strain provided herein. In another embodiment, the plasmid is an episomal plasmid that does not integrate into the recombinant *Listeria* strain's chromosome. In another embodiment, the plasmid is an integrative plasmid that integrates into the *Listeria* strain's chromosome. In another embodiment, the plasmid is a multicopy plasmid. In another embodiment, the recombinant *Listeria* strain is administered to the human subject at a dose of 1×10^9 - 3.31×10^{10} CFU. In another embodiment, the dose is $5\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $7\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $10\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $20\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $30\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $50\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $70\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $100\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $150\text{-}500 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}300 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}200 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}150 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}100 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}70 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}50 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}30 \times 10^8$ CFU. In another embodiment, the dose is $5\text{-}20 \times 10^8$ CFU. In another embodiment, the dose is $1\text{-}30 \times 10^9$ CFU. In another embodiment, the

dose is $1-20 \times 10^9$ CFU. In another embodiment, the dose is $2-30 \times 10^9$ CFU. In another embodiment, the dose is $1-10 \times 10^9$ CFU. In another embodiment, the dose is $2-10 \times 10^9$ CFU. In another embodiment, the dose is $3-10 \times 10^9$ CFU. In another embodiment, the dose is $2-7 \times 10^9$ CFU. In another embodiment, the dose is $2-5 \times 10^9$ CFU. In another embodiment, the dose is $3-5 \times 10^9$ CFU.

[00138] In another embodiment, the dose is 1×10^7 organisms. In another embodiment, the dose is 1×10^8 organisms. In another embodiment, the dose is 1×10^9 organisms. In another embodiment, the dose is 1.5×10^9 organisms. In another embodiment, the dose is 2×10^9 organisms. In another embodiment, the dose is 3×10^9 organisms. In another embodiment, the dose is 4×10^9 organisms. In another embodiment, the dose is 5×10^9 organisms. In another embodiment, the dose is 6×10^9 organisms. In another embodiment, the dose is 7×10^9 organisms. In another embodiment, the dose is 8×10^9 organisms. In another embodiment, the dose is 10×10^9 organisms. In another embodiment, the dose is 1.5×10^{10} organisms. In another embodiment, the dose is 2×10^{10} organisms. In another embodiment, the dose is 2.5×10^{10} organisms. In another embodiment, the dose is 3×10^{10} organisms. In another embodiment, the dose is 3.3×10^{10} organisms. In another embodiment, the dose is 4×10^{10} organisms. In another embodiment, the dose is 5×10^{10} organisms. Each dose and range of doses represents a separate embodiment of the present invention.

[00139] In one embodiment, repeat administrations (doses) of compositions of this invention may be undertaken immediately following the first course of treatment or after an interval of days, weeks or months to achieve tumor regression. In another embodiment, repeat doses may be undertaken immediately following the first course of treatment or after an interval of days, weeks or months to achieve suppression of tumor growth. Assessment may be determined by any of the techniques known in the art, including diagnostic methods such as imaging techniques, analysis of serum tumor markers, biopsy, or the presence, absence or amelioration of tumor associated symptoms.

[00140] It will be appreciated by the skilled artisan that the term "Boosting" may encompass administering an immunogenic composition or recombinant *Listeria* strain dose to a subject. In another embodiment, of methods of the present invention, 2 boosts (or a total of 3 inoculations) are administered. In another embodiment, 3 boosts are administered. In another embodiment, 4 boosts are administered. In another embodiment, 5 boosts are administered. In another embodiment, 6 boosts are administered. In another embodiment, more than 6 boosts are administered. Each possibility represents a separate embodiment of the present invention.

[00141] In one embodiment, a method of present invention further comprises the step of boosting the human subject with a recombinant *Listeria* strain of the present invention. In another embodiment, the recombinant *Listeria* strain used in the booster inoculation is the same as the strain used in the initial “priming” inoculation. In another embodiment, the booster strain is different from the priming strain. In another embodiment, the same doses are used in the priming and boosting inoculations. In another embodiment, a larger dose is used in the booster. In another embodiment, a smaller dose is used in the booster. In another embodiment, the methods of the present invention further comprise the step of administering to the subject a booster vaccination. In one embodiment, the booster vaccination follows a single priming vaccination. In another embodiment, a single booster vaccination is administered after the priming vaccinations. In another embodiment, two booster vaccinations are administered after the priming vaccinations. In another embodiment, three booster vaccinations are administered after the priming vaccinations. In one embodiment, the period between a prime and a boost strain is experimentally determined by the skilled artisan. In another embodiment, the period between a prime and a boost strain is from 1 day and up to 1 week, in another embodiment it is up to 2 weeks, in another embodiment, it is up to 3 weeks, in another embodiment, it is up to 4 weeks, in another embodiment, it is up to 5 weeks, in another embodiment it is up to 6-8 weeks, in yet another embodiment, the boost strain is administered up to 8-12 weeks after the prime strain. Each possibility represents a separate embodiment of the present invention.

[00142] In another embodiment, a method of present invention further comprises the step of inoculating the human subject with an immunogenic composition comprising the E7 antigen. In another embodiment, the immunogenic composition comprises a recombinant E7 protein or fragment thereof. In another embodiment, the immunogenic composition comprises a nucleotide molecule expressing a recombinant E7 protein or fragment thereof. In another embodiment, the non-*Listerial* inoculation is administered after the *Listerial* inoculation. In another embodiment, the non-*Listerial* inoculation is administered before the *Listerial* inoculation. Each possibility represents a separate embodiment of the present invention.

[00143] The recombinant *Listeria* strain of methods and compositions of the present invention is, in another embodiment, a recombinant *Listeria monocytogenes* strain. In another embodiment, the *Listeria* strain is a recombinant *Listeria seeligeri* strain. In another embodiment, the *Listeria* strain is a recombinant *Listeria grayi* strain. In another embodiment, the *Listeria* strain is a recombinant *Listeria ivanovii* strain. In another embodiment, the *Listeria* strain is a recombinant *Listeria murrayi* strain. In another embodiment, the *Listeria*

strain is a recombinant *Listeria welshimeri* strain. In another embodiment, the *Listeria* strain is a recombinant strain of any other *Listeria* species known in the art. Each possibility represents a separate embodiment of the present invention.

[00144] The present invention provides a number of *Listerial* species and strains for making or engineering an attenuated *Listeria* of the present invention. In one embodiment, the *Listeria* strain is *L. monocytogenes* 10403S wild type (see Bishop and Hinrichs (1987) J. Immunol. 139: 2005–2009; Lauer, et al. (2002) J. Bact. 184: 4177–4186.) In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4056 (phage cured) (see Lauer, et al. (2002) J. Bact. 184: 4177–4186). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4027, which is phage cured and deleted in the hly gene (see Lauer, et al. (2002) J. Bact. 184: 4177–4186; Jones and Portnoy (1994) Infect. Immunity 65: 5608–5613.). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4029, which is phage cured, deleted in ActA (see Lauer, et al. (2002) J. Bact. 184: 4177–4186; Skoble, et al. (2000) J. Cell Biol. 150: 527–538). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4042 (delta PEST) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4097 (LLO-S44A) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4364 (delta lplA; lipoate protein ligase) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4405 (delta inlA) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4406 (delta inlB) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* CS-L0001 (delta ActA-delta inlB) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* CS-L0002 (delta ActA-delta lplA) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* CS-L0003 (L461T-delta lplA) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4038 (delta ActA-LLO L461T) (see Brockstedt, et al. (2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4384 (S44A-LLO L461T) (see Brockstedt, et al.

(2004) Proc. Natl. Acad. Sci. USA 101: 13832–13837; supporting information). In another embodiment, the *Listeria* strain is *L. monocytogenes*. Mutation in lipoate protein (see O'Riordan, et al. (2003) Science 302: 462–464). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4017 (10403S hly (L461T), having a point mutation in hemolysin gene (see U.S. Provisional Pat. Appl. Ser. No. 60/490,089 filed Jul. 24, 2003). In another embodiment, the *Listeria* strain is *L. monocytogenes* EGD (see GenBank Acc. No. AL591824). In another embodiment, the *Listeria* strain is *L. monocytogenes* EGD-e (see GenBank Acc. No. NC_003210. ATCC Acc. No. BAA-679). In another embodiment, the *Listeria* strain is *L. monocytogenes* DP-L4029 deleted in uvrAB (see U.S. Provisional Pat. Appl. Ser. No. 60/541,515 filed Feb. 2, 2004; US Provisional Pat. Appl. Ser. No. 60/490,080 filed Jul. 24, 2003). In another embodiment, the *Listeria* strain is *L. monocytogenes* ActA-/inlB - double mutant (see ATCC Acc. No. PTA-5562). In another embodiment, the *Listeria* strain is *L. monocytogenes* lplA mutant or hly mutant (see U.S. Pat. Applic. No. 20040013690 of Portnoy, et. al). In another embodiment, the *Listeria* strain is *L. monocytogenes* DAL/DAT double mutant. (see U.S. Pat. Applic. No. 20050048081 of Frankel and Portnoy. The present invention encompasses reagents and methods that comprise the above *Listerial* strains, as well as these strains that are modified, e.g., by a plasmid and/or by genomic integration, to contain a nucleic acid encoding one of, or any combination of, the following genes: hly (LLO; listeriolysin); iap (p60); inlA; inlB; inlC; dal (alanine racemase); dat (D-amino acid aminotransferase); plcA; plcB; actA; or any nucleic acid that mediates growth, spread, breakdown of a single walled vesicle, breakdown of a double walled vesicle, binding to a host cell, uptake by a host cell. The present invention is not to be limited by the particular strains disclosed above.

[00145] In another embodiment, a recombinant *Listeria* strain of the present invention has been passaged through an animal host. In another embodiment, the passaging maximizes efficacy of the strain as a vaccine vector. In another embodiment, the passaging stabilizes the immunogenicity of the *Listeria* strain. In another embodiment, the passaging stabilizes the virulence of the *Listeria* strain. In another embodiment, the passaging increases the immunogenicity of the *Listeria* strain. In another embodiment, the passaging increases the virulence of the *Listeria* strain. In another embodiment, the passaging removes unstable sub-strains of the *Listeria* strain. In another embodiment, the passaging reduces the prevalence of unstable sub-strains of the *Listeria* strain. In another embodiment, the *Listeria* strain contains a genomic insertion of the gene encoding the antigen-containing recombinant peptide. In another embodiment, the *Listeria* strain carries a plasmid comprising the gene encoding the

antigen-containing recombinant peptide. In another embodiment, the passaging is performed as described herein (e.g. in Example 12). In another embodiment, the passaging is performed by any other method known in the art. Each possibility represents a separate embodiment of the present invention.

5 [00146] In another embodiment, the recombinant *Listeria* strain utilized in methods of the present invention has been stored in a frozen cell bank. In another embodiment, the recombinant *Listeria* strain has been stored in a lyophilized condition. Each possibility represents a separate embodiment of the present invention.

[00147] In another embodiment, the cell bank of methods and compositions of the present
10 invention is a master cell bank. In another embodiment, the cell bank is a working cell bank. In another embodiment, the cell bank is Good Manufacturing Practice (GMP) cell bank. In another embodiment, the cell bank is intended for production of clinical-grade material. In another embodiment, the cell bank conforms to regulatory practices for human use. In another embodiment, the cell bank is any other type of cell bank known in the art. Each possibility
15 represents a separate embodiment of the present invention.

[00148] “Good Manufacturing Practices” are defined, in another embodiment, by (21 CFR 210–211) of the United States Code of Federal Regulations. In another embodiment, “Good Manufacturing Practices” are defined by other standards for production of clinical-grade material or for human consumption; e.g. standards of a country other than the United States.
20 Each possibility represents a separate embodiment of the present invention.

[00149] In another embodiment, a recombinant *Listeria* strain utilized in methods of the present invention is from a batch of vaccine doses.

[00150] In another embodiment, a recombinant *Listeria* strain utilized in methods of the present invention is from a frozen or lyophilized stock produced by methods provided in US
25 Patent Ser. No. 8,114,414, which is incorporated by reference herein.

[00151] In another embodiment, a peptide of the present invention is a fusion peptide. In another embodiment, “fusion peptide” refers to a peptide or polypeptide comprising 2 or more proteins linked together by peptide bonds or other chemical bonds. In another embodiment, the proteins are linked together directly by a peptide or other chemical bond. In another
30 embodiment, the proteins are linked together with 1 or more AA (e.g. a “spacer”) between the 2 or more proteins. Each possibility represents a separate embodiment of the present invention.

[00152] In another embodiment, a vaccine of the present invention further comprises an adjuvant. The adjuvant utilized in methods and compositions of the present invention is, in

another embodiment, a granulocyte/macrophage colony-stimulating factor (GM-CSF) protein. In another embodiment, the adjuvant comprises a GM-CSF protein. In another embodiment, the adjuvant is a nucleotide molecule encoding GM-CSF. In another embodiment, the adjuvant comprises a nucleotide molecule encoding GM-CSF. In another embodiment, the adjuvant is saponin QS21. In another embodiment, the adjuvant comprises saponin QS21. In another embodiment, the adjuvant is monophosphoryl lipid A. In another embodiment, the adjuvant comprises monophosphoryl lipid A. In another embodiment, the adjuvant is SBAS2. In another embodiment, the adjuvant comprises SBAS2. In another embodiment, the adjuvant is an unmethylated CpG-containing oligonucleotide. In another embodiment, the adjuvant comprises an unmethylated CpG-containing oligonucleotide. In another embodiment, the adjuvant is an immune-stimulating cytokine. In another embodiment, the adjuvant comprises an immune-stimulating cytokine. In another embodiment, the adjuvant is a nucleotide molecule encoding an immune-stimulating cytokine. In another embodiment, the adjuvant comprises a nucleotide molecule encoding an immune-stimulating cytokine. In another embodiment, the adjuvant is or comprises a quill glycoside. In another embodiment, the adjuvant is or comprises a bacterial mitogen. In another embodiment, the adjuvant is or comprises a bacterial toxin. In another embodiment, the adjuvant is or comprises any other adjuvant known in the art. Each possibility represents a separate embodiment of the present invention.

[00153] In another embodiment, a nucleotide of the present invention is operably linked to a promoter/regulatory sequence that drives expression of the encoded peptide in the *Listeria* strain. Promoter/regulatory sequences useful for driving constitutive expression of a gene are well known in the art and include, but are not limited to, for example, the P_{hlyA}, P_{ActA}, and p60 promoters of *Listeria*, the *Streptococcus* bac promoter, the *Streptomyces griseus* *sgiA* promoter, and the *B. thuringiensis* *phaZ* promoter. In another embodiment, inducible and tissue specific expression of the nucleic acid encoding a peptide of the present invention is accomplished by placing the nucleic acid encoding the peptide under the control of an inducible or tissue specific promoter/regulatory sequence. Examples of tissue specific or inducible promoter/regulatory sequences which are useful for his purpose include, but are not limited to the MMTV LTR inducible promoter, and the SV40 late enhancer/promoter. In another embodiment, a promoter that is induced in response to inducing agents such as metals, glucocorticoids, and the like, is utilized. Thus, it will be appreciated that the invention includes the use of any promoter/regulatory sequence, which is either known or unknown, and which is capable of driving expression of the desired protein operably linked thereto.

[00154] An N-terminal fragment of an ActA protein utilized in methods and compositions of the present invention has, in another embodiment, the sequence set forth in SEQ ID NO: 5.

MRAMMVVFITANCITINPDIIFAATDSEDSSLNTDEWEEKTEEQPSEVNTGPRYETAR
EVSSRDIKELEKSNKVRNTNKADLIAMLKEKAEEKGNINNNNSEQTENAAINEEASG
5 ADRPAIQVERRHPGLPSDSAAEIKRRRKAIASSDSELESLTYPDKPTKVNKKKVAKES
VADASESDLDSMQSADESSPQPLKANQQPFFPKVFKKIKDAGKWVRDKIDENPEVK
KAIVDKSAGLIDQLLTKKKSEEVNASDFPPPPTDEELRLALPETPMLLGFNAPATSEPS
SFEFPPPPTDEELRLALPETPMLLGFNAPATSEPSSFEFPPPPTDELEIIRETASSLDSSF
TRGDLASLRNAINRHSQNFSDFPPIPTEEELNGRGGRP. In another embodiment, the

10 ActA fragment comprises the sequence set forth in SEQ ID NO: 5. In another embodiment, the ActA fragment is any other ActA fragment known in the art. Each possibility represents a separate embodiment of the present invention.

[00155] In another embodiment, the recombinant nucleotide encoding a fragment of an ActA protein comprises the sequence set forth in SEQ ID NO: 6:

15 Atgctgctgatgatggtggttttcattactgccaattgcattacgattaaccccgacataatattgcagcgacagatagcgaagattcta
gtctaaacacagatgaatgggaagaagaaaaacagaagagcaaccaagcgaggtaaatacgggaccaagatacgaaactgcacg
tgaagtaagttcacgtgatattaaagaactagaaaaatcgaataaagtgagaaatacgaacaaagcagacctaataagcaatgttgaaag
aaaaagcagaaaaaggtccaaatatcaataataacaacagtgaacaaactgagaatgcggctataaatgaagaggcttcaggagccg
accgaccagctatacaagtggagcgtcgtcatccaggattgccatcgatagcgcagcggaattaaaaaagaaggaaagccatag
20 catcatcgatagtgagcttgaaagccttacttatccggataaaccaacaaaagtaaataagaaaaaagtggcgaaagagtcagttgcg
gatgcttctgaaagtgacttagattctagcatgcagtcagcagatgagtcttcaccacaacctttaaagcaaaccaacaaccattttccc
taaagtatttaaaaaataaaaagatgcggggaaatgggtacgtgataaaatcgacgaaaatcctgaagtaaagaaagcgattgttgata
aaagtgcagggttaattgaccaattattaacaaaaagaaaagtgaagaggtaaatagtctcggaactcccgccaccacctacggatgaa
gagttaagacttgcttgccagagacaccaatgcttcttggtttaatgctcctgctacatcagaaccgagctcattcgaattccaccacca
25 cctacggatgaagagttaagacttgcttgccagagacgccaatgcttcttggtttaatgctcctgctacatcggaaccgagctcggtcga
attccaccgcctccaacagaagatgaactagaaatcatccgggaacagcatcctcgtagattctagttttacaagaggggatttagct
agtttgagaaatgctattaatcgccatagtcaaaattctctgattcccaccaatcccaacagaagaagagttgaacgggagagggcggt
agacca. In another embodiment, the recombinant nucleotide has the sequence set forth in SEQ

ID NO: 6. In another embodiment, the recombinant nucleotide comprises any other sequence
30 that encodes a fragment of an ActA protein. Each possibility represents a separate embodiment of the present invention.

[00156] In another embodiment of the methods and compositions of the present invention, a PEST amino acid AA sequence is fused to the E7 or E6 antigen. As provided herein, recombinant *Listeria* strains expressing PEST amino acid sequence-antigen fusions induce

anti-tumor immunity (Example 3) and generate antigen-specific, tumor-infiltrating T cells (Example 4). Further, enhanced cell mediated immunity was demonstrated for fusion proteins comprising an antigen and LLO containing the PEST amino acid AA sequence KENSISSMAPPASPPASPKTPIEKKHADAIDK (SEQ ID NO: 1).

5 [00157] Thus, fusion of an antigen to other LM PEST amino acid sequences and PEST amino acid sequences derived from other prokaryotic organisms will also enhance immunogenicity of the antigen. The PEST amino acid AA sequence has, in another embodiment, a sequence selected from SEQ ID NO: 7-12. In another embodiment, the PEST amino acid sequence is a PEST amino acid sequence from the LM ActA protein. In another embodiment, the PEST amino acid sequence is KTEEQPSEVNTGPR (SEQ ID NO: 7), KASVTDTSSEGLDSSMQSADESTPQPLK (SEQ ID NO: 8), KNEEVNASDFPPPPTDEELR (SEQ ID NO: 9), or RGGIPTSEEFSSLNSGDFTDDENSETTEEEIDR (SEQ ID NO: 10). In another embodiment, the PEST amino acid sequence is from Streptolysin O protein of *Streptococcus sp.* In another embodiment, the PEST amino acid sequence is from *Streptococcus pyogenes* Streptolysin O, e.g. KQNTASTETTTTNEQPK (SEQ ID NO: 11) at AA 35-51. In another embodiment, the PEST amino acid sequence is from *Streptococcus equisimilis* Streptolysin O, e.g. KQNTANTETTTTNEQPK (SEQ ID NO: 12) at AA 38-54. In another embodiment, the PEST amino acid sequence is another PEST amino acid AA sequence derived from a prokaryotic organism. In another embodiment, the PEST amino acid sequence is any other PEST amino acid sequence known in the art. Each possibility represents a separate embodiment of the present invention.

[00158] PEST amino acid sequences of other prokaryotic organism can be identified in accordance with methods such as described by, for example Rechsteiner and Rogers (1996, Trends Biochem. Sci. 21:267-271) for LM. Alternatively, PEST amino acid AA sequences from other prokaryotic organisms can also be identified based by this method. Other prokaryotic organisms wherein PEST amino acid AA sequences would be expected to include, but are not limited to, other *Listeria* species. In another embodiment, the PEST amino acid sequence is embedded within the antigenic protein. Thus, in another embodiment, “fusion” refers to an antigenic protein comprising both the antigen and either i) an N-terminal LLO protein (tLLO), ii) an N-terminal ActA protein or iii) a PEST amino acid sequence either linked at one end of the antigen or embedded within the antigen.

[00159] In another embodiment, a PEST amino acid sequence is identified using any other method or algorithm known in the art, e.g the CaSPredictor (Garay-Malpartida HM,

Occhiucci JM, Alves J, Belizario JE. Bioinformatics. 2005 Jun;21 Suppl 1:i169-76). In another embodiment, the following method is used:

[00160] A PEST index is calculated for each 30-35 AA stretch by assigning a value of 1 to the amino acids Ser, Thr, Pro, Glu, Asp, Asn, or Gln. The coefficient value (CV) for each of the PEST residue is 1 and for each of the other AA (non-PEST) is 0.

[00161] Each method for identifying a PEST amino acid sequence represents a separate embodiment of the present invention.

[00162] In another embodiment, the LLO protein, ActA protein, or fragment thereof of the present invention need not be that which is set forth exactly in the sequences set forth herein, but rather other alterations, modifications, or changes can be made that retain the functional characteristics of an LLO or ActA protein fused to an antigen as set forth elsewhere herein. In another embodiment, the present invention utilizes an analog of an LLO protein, ActA protein, or fragment thereof. Analogs differ, in another embodiment, from naturally occurring proteins or peptides by conservative AA sequence differences or by modifications which do not affect sequence, or by both.

[00163] In another embodiment, either a whole E7 protein or a fragment thereof is fused to a LLO protein, ActA protein, or PEST amino acid sequence-containing peptide to generate a recombinant peptide of methods of the present invention. The E7 protein that is utilized (either whole or as the source of the fragments) has, in another embodiment, the sequence

[00164] MHGDTPTLHEYMLDLQPETTDLYCYEQLNDSSEEEDEIDGPAGQAEPDRAHY NIVTFCKCDSTLRRLCVQSTHVDIRTLEDLLMGTLGIVCPICSQKP (SEQ ID No: 13). In another embodiment, the E7 protein is a homologue of SEQ ID No: 13. In another embodiment, the E7 protein is a variant of SEQ ID No: 13. In another embodiment, the E7 protein is an isomer of SEQ ID No: 13. In another embodiment, the E7 protein is a fragment of SEQ ID No: 13. In another embodiment, the E7 protein is a fragment of a homologue of SEQ ID No: 13. In another embodiment, the E7 protein is a fragment of a variant of SEQ ID No: 13. In another embodiment, the E7 protein is a fragment of an isomer of SEQ ID No: 13. Each possibility represents a separate embodiment of the present invention.

[00165] In another embodiment, the sequence of the E7 protein is:

[00166] MHGPKATLQDIVLHLEPQNEIPVDLLCHEQLSDSEEENDEIDGVNHQHL PARR AEPQRHTMLCMCKCEARIELVVESSADDLRAFQQLFLNTLSFVCPWCASQQ (SEQ ID No: 14). In another embodiment, the E6 protein is a homologue of SEQ ID No: 14. In another embodiment, the E6 protein is a variant of SEQ ID No: 14. In another embodiment, the E6 protein is an isomer of SEQ ID No: 14. In another embodiment, the E6 protein is a

fragment of SEQ ID No: 14. In another embodiment, the E6 protein is a fragment of a
homologue of SEQ ID No: 14. In another embodiment, the E6 protein is a fragment of a
variant of SEQ ID No: 14. In another embodiment, the E6 protein is a fragment of an isomer
of SEQ ID No: 14. Each possibility represents a separate embodiment of the present
5 invention.

[00167] In another embodiment, the E7 protein has a sequence set forth in one of the
following GenBank entries: M24215, NC_004500, V01116, X62843, or M14119. In another
embodiment, the E7 protein is a homologue of a sequence from one of the above GenBank
entries. In another embodiment, the E7 protein is a variant of a sequence from one of the
10 above GenBank entries. In another embodiment, the E7 protein is an isomer of a sequence
from one of the above GenBank entries. In another embodiment, the E7 protein is a fragment
of a sequence from one of the above GenBank entries. In another embodiment, the E7 protein
is a fragment of a homologue of a sequence from one of the above GenBank entries. In
another embodiment, the E7 protein is a fragment of a variant of a sequence from one of the
15 above GenBank entries. In another embodiment, the E7 protein is a fragment of an isomer of
a sequence from one of the above GenBank entries. Each possibility represents a separate
embodiment of the present invention..

[00168] In another embodiment, either a whole E6 protein or a fragment thereof is fused to a
LLO protein, ActA protein, or PEST amino acid sequence-containing peptide to generate a
20 recombinant peptide of methods of the present invention. The E6 protein that is utilized
(either whole or as the source of the fragments) has, in another embodiment, the sequence

[00169] MHQKRTAMFQDPQERPRKLPQLCTELQTTIHDILECVYCKQQLLRREVDFA
FRDLCIVYRDGNPYAVCDKCLKFYISKISEYRHYCYSLYGTTLEQQYNKPLCDLLIRCI
NCQKPLCPEEKQRHLDKKQRFHNIRGRWTGRCMSCCRSSRTRRETQL (SEQ ID No:

25 15). In another embodiment, the E6 protein is a homologue of SEQ ID No: 15. In another
embodiment, the E6 protein is a variant of SEQ ID No: 15. In another embodiment, the E6
protein is an isomer of SEQ ID No: 15. In another embodiment, the E6 protein is a fragment
of SEQ ID No: 15. In another embodiment, the E6 protein is a fragment of a homologue of
SEQ ID No: 15. In another embodiment, the E6 protein is a fragment of a variant of SEQ ID
30 No: 15. In another embodiment, the E6 protein is a fragment of an isomer of SEQ ID No: 15.
Each possibility represents a separate embodiment of the present invention.

[00170] In another embodiment, the sequence of the E6 protein is:

[00171] MARFEDPTRRPYKLPDLCTELNTSLQDIEITCVYCKTVLELTEVFEFAFKDLFV
VYRDSIPHAACHKCIDFYSRIRELRHYSDSVYGDITLEKLTNTGLYNLLIRCLRCQKPL

NPAEKLRLHLNEKRRFHNIAGHYRGQCHSCCNRRARQERLQRRRETQV (SEQ ID No: 16). In another embodiment, the E6 protein is a homologue of SEQ ID No: 16. In another embodiment, the E6 protein is a variant of SEQ ID No: 16. In another embodiment, the E6 protein is an isomer of SEQ ID No: 16. In another embodiment, the E6 protein is a fragment of SEQ ID No: 16. In another embodiment, the E6 protein is a fragment of a homologue of SEQ ID No: 16. In another embodiment, the E6 protein is a fragment of a variant of SEQ ID No: 16. In another embodiment, the E6 protein is a fragment of an isomer of SEQ ID No: 16. Each possibility represents a separate embodiment of the present invention.

[00172] In another embodiment, the E6 protein has a sequence set forth in one of the following GenBank entries: M24215, M14119, NC_004500, V01116, X62843, or M14119. In another embodiment, the E6 protein is a homologue of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is a variant of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is an isomer of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is a fragment of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is a fragment of a homologue of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is a fragment of a variant of a sequence from one of the above GenBank entries. In another embodiment, the E6 protein is a fragment of an isomer of a sequence from one of the above GenBank entries. Each possibility represents a separate embodiment of the present invention.

[00173] In another embodiment, “homology” refers to identity to an LLO sequence (e.g. to one of SEQ ID No: 2-4) of greater than 60%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 64%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 68%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 72%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 75%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 78%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 80%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 82%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 83%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 85%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 87%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 88%. In another embodiment,

“homology” refers to identity to one of SEQ ID No: 2-4 of greater than 90%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 92%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 93%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 95%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 96%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 97%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 98%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of greater than 99%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 2-4 of 100%. Each possibility represents a separate embodiment of the present invention.

[00174] In another embodiment, “homology” refers to identity to an E7 sequence (e.g. to one of SEQ ID No: 13-14) of greater than 60%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 62%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 64%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 68%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 72%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 75%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 78%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 80%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 82%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 83%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 85%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 87%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 88%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 90%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 92%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 93%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 95%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 96%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 97%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14

of greater than 98%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of greater than 99%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 13-14 of 100%. Each possibility represents a separate embodiment of the present invention.

5 [00175] In another embodiment, “homology” refers to identity to an E6 sequence (e.g. to one of SEQ ID No: 15-16) of greater than 60%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 64%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 68%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 72%.
 10 In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 75%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 78%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 80%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 82%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 83%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 85%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 87%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 88%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 90%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 92%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 93%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 95%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 96%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 97%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 98%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of greater than 99%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 15-16 of 100%. Each possibility represents a separate embodiment of the present
 25 invention.
 30

[00176] In another embodiment, “homology” refers to identity to a PEST amino acid sequence (e.g. to one of SEQ ID No: 1, and 7-12) or to an ActA sequence (e.g. to one of SEQ ID No: 5-6) of greater than 60%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 60%. In another embodiment,

“homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 64%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 68%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 72%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 75%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 78%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 80%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 82%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 83%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 85%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 87%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 88%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 90%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 92%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 93%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 95%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 96%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 97%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 98%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of greater than 99%. In another embodiment, “homology” refers to identity to one of SEQ ID No: 1, and 7-12 or SEQ ID No: 5-6 of 100%. Each possibility represents a separate embodiment of the present invention.

[00177] Protein and/or peptide homology for any AA sequence listed herein is determined, in one embodiment, by methods well described in the art, including immunoblot analysis, or via computer algorithm analysis of AA sequences, utilizing any of a number of software packages available, via established methods. Some of these packages include the FASTA, BLAST, MPsrch or Scanps packages, and employ, in other embodiments, the use of the Smith and

Waterman algorithms, and/or global/local or BLOCKS alignments for analysis, for example. Each method of determining homology represents a separate embodiment of the present invention.

[00178] In another embodiment, the LLO protein, ActA protein, or fragment thereof is
5 attached to the antigen by chemical conjugation. In another embodiment, glutaraldehyde is used for the conjugation. In another embodiment, the conjugation is performed using any suitable method known in the art. Each possibility represents another embodiment of the present invention.

[00179] In another embodiment, fusion proteins of the present invention are prepared by any
10 suitable method, including, for example, cloning and restriction of appropriate sequences or direct chemical synthesis by methods discussed below. In another embodiment, subsequences are cloned and the appropriate subsequences cleaved using appropriate restriction enzymes. The fragments are then ligated, in another embodiment, to produce the desired DNA sequence. In another embodiment, DNA encoding the fusion protein is produced using DNA
15 amplification methods, for example polymerase chain reaction (PCR). First, the segments of the native DNA on either side of the new terminus are amplified separately. The 5' end of the one amplified sequence encodes the peptide linker, while the 3' end of the other amplified sequence also encodes the peptide linker. Since the 5' end of the first fragment is complementary to the 3' end of the second fragment, the two fragments (after partial
20 purification, e.g. on LMP agarose) can be used as an overlapping template in a third PCR reaction. The amplified sequence will contain codons, the segment on the carboxy side of the opening site (now forming the amino sequence), the linker, and the sequence on the amino side of the opening site (now forming the carboxyl sequence). The insert is then ligated into a plasmid.

25 [00180] In another embodiment, the LLO protein, ActA protein, or fragment thereof and the antigen, or fragment thereof are conjugated by a means known to those of skill in the art. In another embodiment, the antigen, or fragment thereof is conjugated, either directly or through a linker (spacer), to the ActA protein or LLO protein. In another embodiment, the chimeric molecule is recombinantly expressed as a single-chain fusion protein.

30 [00181] In another embodiment, a fusion peptide of the present invention is synthesized using standard chemical peptide synthesis techniques. In another embodiment, the chimeric molecule is synthesized as a single contiguous polypeptide. In another embodiment, the LLO protein, ActA protein, or fragment thereof; and the antigen, or fragment thereof are synthesized separately, then fused by condensation of the amino terminus of one molecule

with the carboxyl terminus of the other molecule, thereby forming a peptide bond. In another embodiment, the ActA protein or LLO protein and antigen are each condensed with one end of a peptide spacer molecule, thereby forming a contiguous fusion protein.

[00182] In another embodiment, the peptides and proteins of the present invention are prepared

5 by solid-phase peptide synthesis (SPPS) as described by Stewart et al. in Solid Phase Peptide Synthesis, 2nd Edition, 1984, Pierce Chemical Company, Rockford, Ill.; or as described by Bodanszky and Bodanszky (The Practice of Peptide Synthesis, 1984, Springer-Verlag, New York). In another embodiment, a suitably protected AA residue is attached through its carboxyl group to a derivatized, insoluble polymeric support, such as cross-linked polystyrene
10 or polyamide resin. "Suitably protected" refers to the presence of protecting groups on both the alpha-amino group of the amino acid, and on any side chain functional groups. Side chain protecting groups are generally stable to the solvents, reagents and reaction conditions used throughout the synthesis, and are removable under conditions which will not affect the final peptide product. Stepwise synthesis of the oligopeptide is carried out by the removal of the N-
15 protecting group from the initial AA, and couple thereto of the carboxyl end of the next AA in the sequence of the desired peptide. This AA is also suitably protected. The carboxyl of the incoming AA can be activated to react with the N-terminus of the support-bound AA by formation into a reactive group such as formation into a carbodiimide, a symmetric acid anhydride or an "active ester" group such as hydroxybenzotriazole or pentafluorophenyl
20 esters. The pharmaceutical compositions containing vaccines and compositions of the present invention are, in another embodiment, administered to a subject by any method known to a person skilled in the art, such as parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously, intra-dermally, subcutaneously, intra-peritoneally, intra-ventricularly, intra-cranially, intra-vaginally or intra-tumorally.

25 [00183] In another embodiment of the methods and compositions provided herein, the vaccines or compositions are administered orally, and are thus formulated in a form suitable for oral administration, i.e. as a solid or a liquid preparation. Suitable solid oral formulations include tablets, capsules, pills, granules, pellets and the like. Suitable liquid oral formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In another embodiment of the
30 present invention, the active ingredient is formulated in a capsule. In accordance with this embodiment, the compositions of the present invention comprise, in addition to the active compound and the inert carrier or diluent, a hard gelating capsule.

[00184] In another embodiment, the vaccines or compositions are administered by intravenous, intra-arterial, or intra-muscular injection of a liquid preparation. Suitable liquid formulations

include solutions, suspensions, dispersions, emulsions, oils and the like. In one embodiment, the pharmaceutical compositions are administered intravenously and are thus formulated in a form suitable for intravenous administration. In another embodiment, the pharmaceutical compositions are administered intra-arterially and are thus formulated in a form suitable for intra-arterial administration. In another embodiment, the pharmaceutical compositions are administered intra-muscularly and are thus formulated in a form suitable for intra-muscular administration.

[00185] It will be appreciated by a skilled artisan that the term “treating” may encompass both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or lessen the targeted pathologic condition or disorder as described herein. Thus, in one embodiment, treating may include directly affecting or curing, suppressing, inhibiting, preventing, reducing the severity of, delaying the onset of, reducing symptoms associated with the disease, disorder or condition, or a combination thereof. Thus, in one embodiment, “treating” may encompass inter alia delaying progression, expediting remission, inducing remission, augmenting remission, speeding recovery, increasing efficacy of or decreasing resistance to alternative therapeutics, or a combination thereof. In one embodiment, “preventing” or “impeding” may encompass, inter alia, delaying the onset of symptoms, preventing relapse to a disease, decreasing the number or frequency of relapse episodes, increasing latency between symptomatic episodes, or a combination thereof. In one embodiment, “suppressing” or “inhibiting”, may encompass, inter alia, reducing the severity of symptoms, reducing the severity of an acute episode, reducing the number of symptoms, reducing the incidence of disease-related symptoms, reducing the latency of symptoms, ameliorating symptoms, reducing secondary symptoms, reducing secondary infections, prolonging patient survival, or a combination thereof.

[00186] In one embodiment, symptoms are primary, while in another embodiment, symptoms are secondary. In one embodiment, “primary” refers to a symptom that is a direct result of a particular disease or disorder, while in one embodiment, “secondary” refers to a symptom that is derived from or consequent to a primary cause. In one embodiment, the compounds for use in the present invention treat primary or secondary symptoms or secondary complications. In another embodiment, “symptoms” may be any manifestation of a disease or pathological condition.

In another embodiment, the present invention provides a kit comprising vaccine of the present invention, an applicator, and instructional material that describes use of the methods of the invention. Although model kits are described below, the contents of other

useful kits will be apparent to the skilled artisan in light of the present disclosure. Each of these kits represents a separate embodiment of the present invention.

[00187] In one embodiment, the singular forms of words such as "a," "an," and "the," include their corresponding plural references unless the context clearly dictates otherwise.

5

[00188] Throughout this application, various embodiments of this invention may be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed sub ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

10

15

[00189] Whenever a numerical range is indicated herein, it is meant to include any cited numeral (fractional or integral) within the indicated range. The phrases "ranging/ranges between" a first indicate number and a second indicate number and "ranging/ranges from" a first indicate number "to" a second indicate number are used herein interchangeably and are meant to include the first and second indicated numbers and all the fractional and integral numerals there between.

20

[00190] It will be appreciated by a skilled artisan that the term "about" when used to modify a numerically defined parameter may encompass variation of the parameter in quantitative terms plus or minus 5%, or in another embodiment plus or minus 10%, or in another embodiment plus or minus 15%, or in another embodiment plus or minus 20% of stated numerical value for that parameter.

25

[00191] It is to be understood by the skilled artisan that the term "subject" can encompass a mammal including an adult human or a human child, teenager or adolescent in need of therapy for, or susceptible to, a condition or its sequelae, and also may include non-human mammals such as dogs, cats, pigs, cows, sheep, goats, horses, rats, and mice. It will also be appreciated that the term may encompass livestock. The term "subject" does not exclude an individual that is normal in all respects.

30

[00192] It will be appreciated by the skilled artisan that the term "mammal" for purposes of treatment refers to any animal classified as a mammal, including, but not limited to, humans,

domestic and farm animals, and zoo, sports, or pet animals, such as canines, including dogs, and horses, cats, cattle, pigs, sheep, etc.

[00193] In the following examples, numerous specific details are set forth in order to provide a thorough understanding of the invention. However, it will be understood by those skilled in the art that the present invention may be practiced without these specific details. In other instances, well-known methods, procedures, and components have not been described in detail so as not to obscure the present invention. Thus these examples should in no way be construed, as limiting the broad scope of the invention.

EXPERIMENTAL DETAILS SECTION

EXAMPLE 1: LLO-ANTIGEN FUSIONS INDUCE ANTI-TUMOR IMMUNITY

MATERIALS AND EXPERIMENTAL METHODS (EXAMPLES 1-2)

Cell lines

[00194] The C57BL/6 syngeneic TC-1 tumor was immortalized with HPV-16 E6 and E7 and transformed with the c-Ha-ras oncogene. TC-1, provided by T. C. Wu (Johns Hopkins University School of Medicine, Baltimore, MD) is a highly tumorigenic lung epithelial cell expressing low levels of with HPV-16 E6 and E7 and transformed with the c-Ha-ras oncogene. TC-1 was grown in RPMI 1640, 10% FCS, 2 mM L-glutamine, 100 U/ml penicillin, 100 µg/ml streptomycin, 100 µM nonessential amino acids, 1 mM sodium pyruvate, 50 micromolar (mM) 2-ME, 400 microgram (mcg)/ml G418, and 10% National Collection Type Culture-109 medium at 37° with 10% CO₂. C3 is a mouse embryo cell from C57BL/6 mice immortalized with the complete genome of HPV 16 and transformed with pEJ-ras. EL-4/E7 is the thymoma EL-4 retrovirally transduced with E7.

L. monocytogenes strains and propagation

[00195] *Listeria* strains used were Lm-LLO-E7 (hly-E7 fusion gene in an episomal expression system; Figure 1A), Lm-E7 (single-copy E7 gene cassette integrated into *Listeria* genome), Lm-LLO-NP (“DP-L2028”; hly-NP fusion gene in an episomal expression system), and Lm-Gag (“ZY-18”; single-copy HIV-1 Gag gene cassette integrated into the chromosome). E7 was amplified by PCR using the primers 5'-GGCTCGAGCATGGAGATACACC-3' (SEQ ID No: 17; XhoI site is underlined) and 5'-GGGGACTAGTTTATGGTTTCTGAGAACA-3' (SEQ ID No: 18; SpeI site is underlined) and ligated into pCR2.1 (Invitrogen, San Diego, CA). E7 was excised from pCR2.1 by XhoI/ SpeI digestion and ligated into pGG-55. The hly-E7 fusion gene and the pluripotential transcription factor prfA were cloned into pAM401, a

multicopy shuttle plasmid (Wirth R et al, J Bacteriol, 165: 831, 1986), generating pGG-55. The hly promoter drives the expression of the first 441 AA of the hly gene product, (lacking the hemolytic C-terminus, referred to below as “ΔLLO,” and having the sequence set forth in SEQ ID No: 25), which is joined by the XhoI site to the E7 gene, yielding a hly-E7 fusion gene that is transcribed and secreted as LLO-E7. Transformation of a prfA negative strain of *Listeria*, XFL-7 (provided by Dr. Hao Shen, University of Pennsylvania), with pGG-55 selected for the retention of the plasmid *in vivo* (Figures 1A-B). The hly promoter and gene fragment were generated using primers 5'-GGGGGCTAGCCCTCCTTTGATTAGTATATTC-3' (SEQ ID No: 19; NheI site is underlined) and 5'-CTCCCTCGAGATCATAATTTACTTCATC-3' (SEQ ID No: 20; XhoI site is underlined). The prfA gene was PCR amplified using primers 5'-GACTACAAGGACGATGACCGACAAGTGATAACCCGGGATCTAAATAAATCCGTTT-3' (SEQ ID No: 27; XbaI site is underlined) and 5'-CCCGTCGACCAGCTCTTCTTGGTGAAG-3' (SEQ ID No: 21; SalI site is underlined). Lm-E7 was generated by introducing an expression cassette containing the hly promoter and signal sequence driving the expression and secretion of E7 into the orfZ domain of the LM genome. E7 was amplified by PCR using the primers 5'-GCGGATCCCATGGAGATACACCTAC-3' (SEQ ID No: 22; BamHI site is underlined) and 5'-GCTCTAGATTATGGTTTCTGAG-3' (SEQ ID No: 23; XbaI site is underlined). E7 was then ligated into the pZY-21 shuttle vector. LM strain 10403S was transformed with the resulting plasmid, pZY-21-E7, which includes an expression cassette inserted in the middle of a 1.6-kb sequence that corresponds to the orfX, Y, Z domain of the LM genome. The homology domain allows for insertion of the E7 gene cassette into the orfZ domain by homologous recombination. Clones were screened for integration of the E7 gene cassette into the orfZ domain. Bacteria were grown in brain heart infusion medium with (Lm-LLO-E7 and Lm-LLO-NP) or without (Lm-E7 and ZY-18) chloramphenicol (20 µg/ml). Bacteria were frozen in aliquots at -80°C. Expression was verified by Western blotting (Figure 2).

Western blotting

[00196] *Listeria* strains were grown in Luria-Bertoni medium at 37°C and were harvested at the same optical density measured at 600 nm. The supernatants were TCA precipitated and resuspended in 1x sample buffer supplemented with 0.1 N NaOH. Identical amounts of each cell pellet or each TCA-precipitated supernatant were loaded on 4–20% Tris-glycine SDS-PAGE gels (NOVEX, San Diego, CA). The gels were transferred to polyvinylidene difluoride and probed with an anti-E7 monoclonal antibody (mAb) (Zymed Laboratories, South San

Francisco, CA), then incubated with HRP-conjugated anti-mouse secondary Ab (Amersham Pharmacia Biotech, Little Chalfont, U.K.), developed with Amersham ECL detection reagents, and exposed to Hyperfilm (Amersham Pharmacia Biotech).

Measurement of tumor growth

5 [00197] Tumors were measured every other day with calipers spanning the shortest and longest surface diameters. The mean of these two measurements was plotted as the mean tumor diameter in millimeters against various time points. Mice were sacrificed when the tumor diameter reached 20 mm. Tumor measurements for each time point are shown only for surviving mice.

10 ***Effects of Listeria recombinants on established tumor growth***

[00198] Six- to 8-wk-old C57BL/6 mice (Charles River) received 2×10^5 TC-1 cells s.c. on the left flank. One week following tumor inoculation, the tumors had reached a palpable size of 4–5 mm in diameter. Groups of eight mice were then treated with 0.1 LD₅₀ i.p. Lm-LLO-E7 (10^7 CFU), Lm-E7 (10^6 CFU), Lm-LLO-NP (10^7 CFU), or Lm-Gag (5×10^5 CFU) on days 7 and 14.

15 ***⁵¹Cr release assay***

[00199] C57BL/6 mice, 6–8 wk old, were immunized i.p. with 0.1LD₅₀ Lm-LLO-E7, Lm-E7, Lm-LLO-NP, or Lm-Gag. Ten days post-immunization, spleens were harvested. Splenocytes were established in culture with irradiated TC-1 cells (100:1, splenocytes:TC-1) as feeder cells; stimulated *in vitro* for 5 days, then used in a standard ⁵¹Cr release assay, using the following targets: EL-4, EL-4/E7, or EL-4 pulsed with E7 H-2b peptide (RAHYNIVTF). E:T cell ratios, performed in triplicate, were 80:1, 40:1, 20:1, 10:1, 5:1, and 2.5:1. Following a 4-h incubation at 37°C, cells were pelleted, and 50 µl supernatant was removed from each well. Samples were assayed with a Wallac 1450 scintillation counter (Gaithersburg, MD). The percent specific lysis was determined as [(experimental counts per minute (cpm)- spontaneous cpm)/(total cpm - spontaneous cpm)] x 100.

25 ***TC-1-specific proliferation***

[00200] C57BL/6 mice were immunized with 0.1 LD₅₀ and boosted by i.p. injection 20 days later with 1 LD₅₀ Lm-LLO-E7, Lm-E7, Lm-LLO-NP, or Lm-Gag. Six days after boosting, spleens were harvested from immunized and naive mice. Splenocytes were established in culture at 5×10^5 /well in flat-bottom 96-well plates with 2.5×10^4 , 1.25×10^4 , 6×10^3 , or 3×10^3 irradiated TC-1 cells/well as a source of E7 Ag, or without TC-1 cells or with 10 µg/ml Con A. Cells were pulsed 45 h later with 0.5 µCi [³H]thymidine/well. Plates were harvested 18 h later using a Tomtec harvester 96 (Orange, CT), and proliferation was assessed with a

Wallac 1450 scintillation counter. The change in cpm was calculated as experimental cpm - no Ag cpm.

Flow cytometric analysis

[00201] C57BL/6 mice were immunized intravenously (i.v.) with 0.1 LD₅₀ Lm-LLO-E7 or
 5 Lm-E7 and boosted 30 days later. Three-color flow cytometry for CD8 (53-6.7, PE conjugated), CD62 ligand (CD62L; MEL-14, APC conjugated), and E7 H-2Db tetramer was performed using a FACSCalibur® flow cytometer with CellQuest® software (Becton Dickinson, Mountain View, CA). Splenocytes harvested 5 days after the boost were stained at room temperature (rt) with H-2Db tetramers loaded with the E7 peptide (RAHYNIVTF) or a
 10 control (HIV-Gag) peptide. Tetramers were used at a 1/200 dilution and were provided by Dr. Larry R. Pease (Mayo Clinic, Rochester, MN) and by the NIAID Tetramer Core Facility and the NIH AIDS Research and Reference Reagent Program. Tetramer⁺, CD8⁺, CD62L^{low} cells were analyzed.

B16F0-Ova experiment

15 [00202] 24 C57BL/6 mice were inoculated with 5 x 10⁵ B16F0-Ova cells. On days 3, 10 and 17, groups of 8 mice were immunized with 0.1 LD₅₀ Lm-OVA (10⁶ cfu), Lm-LLO-OVA (10⁸ cfu) and eight animals were left untreated.

Statistics

[00203] For comparisons of tumor diameters, mean and SD of tumor size for each group were
 20 determined, and statistical significance was determined by Student's t test. $p \leq 0.05$ was considered significant.

RESULTS

[00204] Lm-E7 and Lm-LLO-E7 were compared for their abilities to impact on TC-1 growth. Subcutaneous tumors were established on the left flank of C57BL/6 mice. Seven days later
 25 tumors had reached a palpable size (4–5 mm). Mice were vaccinated on days 7 and 14 with 0.1 LD₅₀ Lm-E7, Lm-LLO-E7, or, as controls, Lm-Gag and Lm-LLO-NP. Lm-LLO-E7 induced complete regression of 75% of established TC-1 tumors, while tumor growth was controlled in the other 2 mice in the group (Figure 3). By contrast, immunization with Lm-E7 and Lm-Gag did not induce tumor regression. This experiment was repeated multiple times,
 30 always with very similar results. In addition, similar results were achieved for Lm-LLO-E7 under different immunization protocols. In another experiment, a single immunization was able to cure mice of established 5 mm TC-1 tumors.

[00205] In other experiments, similar results were obtained with 2 other E7-expressing tumor cell lines: C3 and EL-4/E7. To confirm the efficacy of vaccination with Lm-LLO-E7, animals

that had eliminated their tumors were re-challenged with TC-1 or EL-4/E7 tumor cells on day 60 or day 40, respectively. Animals immunized with Lm-LLO-E7 remained tumor free until termination of the experiment (day 124 in the case of TC-1 and day 54 for EL-4/E7).

[00206] Thus, expression of an antigen as a fusion protein with Δ LLO enhances the immunogenicity of the antigen.

EXAMPLE 2: LM-LLO-E7 TREATMENT ELICITS TC-1 SPECIFIC SPLENOCYTE PROLIFERATION

[00207] To measure induction of T cells by Lm-E7 with Lm-LLO-E7, TC-1-specific proliferative responses, a measure of antigen-specific immunocompetence, were measured in immunized mice. Splenocytes from Lm-LLO-E7-immunized mice proliferated when exposed to irradiated TC-1 cells as a source of E7, at splenocyte: TC-1 ratios of 20:1, 40:1, 80:1, and 160:1 (Figure 4). Conversely, splenocytes from Lm-E7 and rLm control-immunized mice exhibited only background levels of proliferation.

EXAMPLE 3: FUSION OF E7 TO LLO, ActA, OR A PEST AMINO ACID SEQUENCE ENHANCES E7-SPECIFIC IMMUNITY AND GENERATES TUMOR-INFILTRATING E7-SPECIFIC CD8⁺ CELLS

MATERIALS AND EXPERIMENTAL METHODS

[00208] 500 μ l (microliter) of MATRIGEL®, comprising 100 μ l of 2×10^5 TC-1 tumor cells in phosphate buffered saline (PBS) plus 400 μ l of MATRIGEL® (BD Biosciences, Franklin Lakes, N.J.) were implanted subcutaneously on the left flank of 12 C57BL/6 mice (n=3). Mice were immunized intraperitoneally on day 7, 14 and 21, and spleens and tumors were harvested on day 28. Tumor MATRIGELs were removed from the mice and incubated at 4 °C overnight in tubes containing 2 milliliters (ml) of RP 10 medium on ice. Tumors were minced with forceps, cut into 2 mm blocks, and incubated at 37 °C for 1 hour with 3 ml of enzyme mixture (0.2 mg/ml collagenase-P, 1 mg/ml DNase-1 in PBS). The tissue suspension was filtered through nylon mesh and washed with 5% fetal bovine serum + 0.05% of NaN₃ in PBS for tetramer and IFN-gamma staining.

[00209] Splenocytes and tumor cells were incubated with 1 micromole (mcm) E7 peptide for 5 hours in the presence of brefeldin A at 10^7 cells/ml. Cells were washed twice and incubated in 50 μ l of anti-mouse Fc receptor supernatant (2.4 G2) for 1 hour or overnight at 4 °C. Cells were stained for surface molecules CD8 and CD62L, permeabilized, fixed using the

permeabilization kit Golgi-stop® or Golgi-Plug® (Pharmingen, San Diego, Calif.), and stained for IFN-gamma. 500,000 events were acquired using two-laser flow cytometer FACSCalibur and analyzed using Cellquest Software (Becton Dickinson, Franklin Lakes, NJ). Percentages of IFN-gamma secreting cells within the activated (CD62L^{low}) CD8⁺ T cells were
5 calculated.

[00210] For tetramer staining, H-2D^b tetramer was loaded with phycoerythrin (PE)-conjugated E7 peptide (RAHYNIVTF, SEQ ID NO: 24), stained at rt for 1 hour, and stained with anti-allophycocyanin (APC) conjugated MEL-14 (CD62L) and FITC-conjugated CD8⁺ at 4 °C for 30 min. Cells were analyzed comparing tetramer⁺CD8⁺ CD62L^{low} cells in the spleen and in
10 the tumor.

RESULTS

[00211] To analyze the ability of Lm-ActA-E7 to enhance antigen specific immunity, mice were implanted with TC-1 tumor cells and immunized with either Lm-LLO-E7 (1 x 10⁷ CFU), Lm-E7 (1 x 10⁶ CFU), or Lm-ActA-E7 (2 x 10⁸ CFU), or were untreated (naïve). Tumors of
15 mice from the Lm-LLO-E7 and Lm-ActA-E7 groups contained a higher percentage of IFN-gamma-secreting CD8⁺ T cells (Figure 5A) and tetramer-specific CD8⁺ cells (Figure 5B) than in Lm-E7 or naïve mice.

[00212] In another experiment, tumor-bearing mice were administered Lm-LLO-E7, Lm-PEST-E7, Lm-ΔPEST-E7, or Lm-E7epi, and levels of E7-specific lymphocytes within the
20 tumor were measured. Mice were treated on days 7 and 14 with 0.1 LD₅₀ of the 4 vaccines. Tumors were harvested on day 21 and stained with antibodies to CD62L, CD8, and with the E7/Db tetramer. An increased percentage of tetramer-positive lymphocytes within the tumor were seen in mice vaccinated with Lm-LLO-E7 and Lm-PEST-E7 (Figure 6A). This result was reproducible over three experiments (Figure 6B).

[00213] Thus, Lm-LLO-E7, Lm-ActA-E7, and Lm-PEST-E7 are each efficacious at induction
25 of tumor-infiltrating CD8⁺ T cells and tumor regression.

EXAMPLE 4: PASSAGING OF *LISTERIA* VACCINE VECTORS THROUGH MICE ELICITS INCREASED IMMUNE RESPONSES TO HETEROLOGOUS AND 30 ENDOGENOUS ANTIGENS

MATERIALS AND EXPERIMENTAL METHODS

Bacterial Strains

[00214] *L. monocytogenes* strain 10403S, serotype 1 (ATCC, Manassas, Va.) was the wild type organism used in these studies and the parental strain of the constructs described below. Strain 10403S has an LD₅₀ of approximately 5×10^4 CFU when injected intraperitoneally into BALB/c mice. “Lm-Gag” is a recombinant LM strain containing a copy of the HIV-1 strain HXB (subtype B laboratory strain with a syncytia-forming phenotype) gag gene stably integrated into the *Listerial* chromosome using a modified shuttle vector pKSV7. Gag protein was expressed and secreted by the strain, as determined by Western blot. All strains were grown in brain-heart infusion (BHI) broth or agar plates (Difco Labs, Detroit, Mich).

Bacterial Culture

[00215] Bacteria from a single clone expressing the passenger antigen and/or fusion protein were selected and cultured in BHI broth overnight. Aliquots of this culture were frozen at -70°C with no additives. From this stock, cultures were grown to 0.1-0.2 O.D. at 600 nm, and aliquots were again frozen at -70°C with no additives. To prepare cloned bacterial pools, the above procedure was used, but after each passage a number of bacterial clones were selected and checked for expression of the target antigen, as described herein. Clones in which expression of the foreign antigen was confirmed were used for the next passage.

Passage of Bacteria in Mice

[00216] 6-8 week old female BALB/c (H-2d) mice were purchased from Jackson Laboratories (Bar Harbor, Me) and were maintained in a pathogen-free microisolator environment. The titer of viable bacteria in an aliquot of stock culture, stored frozen at -70 °C, was determined by plating on BHI agar plates on thawing and prior to use. In all, 5×10^5 bacteria were injected intravenously into BALB/c mice. After 3 days, spleens were harvested, homogenized, and serial dilutions of the spleen homogenate were incubated in BHI broth overnight and plated on BHI agar plates. For further passage, aliquots were again grown to 0.1-0.2 O.D., frozen at -70 °C, and bacterial titer was again determined by serial dilution. After the initial passage (passage 0), this sequence was repeated for a total of 4 times.

Intracellular Cytokine Stain for IFN-Gamma

[00217] Lymphocytes were cultured for 5 hours in complete RPMI-10 medium supplemented with 50 U/ml human recombinant IL-2 and 1 microliter/ml Brefeldin A (Golgistop™; PharMingen, San Diego, CA) in the presence or absence of either the cytotoxic T-cell (CTL) epitope for HIV-GAG (AMQMLKETI; SEQ ID No: 25), *Listeria* LLO (GYKDGNEYI; SEQ ID No: 26) or the HPV virus gene E7 (RAHYNIVTF) (SEQ ID No: 24), at a concentration of 1 micromole. Cells were first surface-stained, then washed and subjected to intracellular cytokine stain using the Cytofix/Cytoperm kit in accordance with the manufacturer's

recommendations (PharMingen, San Diego, CA). For intracellular IFN-gamma stain, FITC-conjugated rat anti-mouse IFN-gamma monoclonal antibody (clone XMG 1.2) and its isotype control Ab (rat IgG1; both from PharMingen) was used. In all, 10^6 cells were stained in PBS containing 1% Bovine Serum Albumin and 0.02% sodium azide (FACS Buffer) for 30 minutes at 4 °C. followed by 3 washes in FACS buffer. Sample data were acquired on either a FACScanTM flowcytometer or FACSCaliburTM instrument (Becton Dickinson, San Jose, CA). Three-color flow cytometry for CD8 (PERCP conjugated, rat anti-mouse, clone 53-6.7 Pharmingen, San Diego, Calif.), CD62L (APC conjugated, rat anti-mouse, clone MEL-14), and intracellular IFN-gamma was performed using a FACSCaliburTM flow cytometer, and data were further analyzed with CELLQuest software (Becton Dickinson, Mountain View, CA). Cells were gated on CD8 high and CD62L^{low} before they were analyzed for CD8⁺ and intracellular IFN-gamma staining.

RESULTS

Passaging in Mice Increases the Virulence of Recombinant *Listeria Monocytogenes*

[00218] Three different constructs were used to determine the impact of passaging on recombinant *Listeria* vaccine vectors. Two of these constructs carry a genomic insertion of the passenger antigen: the first comprises the HIV gag gene (Lm-Gag), and the second comprises the HPV E7 gene (Lm-E7). The third (Lm-LLO-E7) comprises a plasmid with the fusion gene for the passenger antigen (HPV E7) fused with a truncated version of LLO and a gene encoding prfA, the positive regulatory factor that controls *Listeria* virulence factors. This plasmid was used to complement a prfA negative mutant so that in a live host, selection pressures would favor conservation of the plasmid, because without it the bacterium is avirulent. All 3 constructs had been propagated extensively *in vitro* for many bacterial generations.

[00219] Passaging the bacteria resulted in an increase in bacterial virulence, as measured by numbers of surviving bacteria in the spleen, with each of the first 2 passages. For Lm-Gag and Lm-LLO-E7, virulence increased with each passage up to passage 2 (Figure 7A). The plasmid-containing construct, Lm-LLO-E7, demonstrated the most dramatic increase in virulence. Prior to passage, the initial immunizing dose of Lm-LLO-E7 had to be increased to 10^7 bacteria and the spleen had to be harvested on day 2 in order to recover bacteria (whereas an initial dose of 10^5 bacteria for Lm-Gag was harvested on day 3). After the initial passage, the standard dosage of Lm-LLO-E7 was sufficient to allow harvesting on day 3. For Lm-E7, virulence increased by 1.5 orders of magnitude over unpassaged bacteria (Figure 7B).

[00220] Thus, passage through mice increases the virulence of *Listeria* vaccine strains.

Passaging Increases the Ability of *L. monocytogenes* to Induce CD8⁺ T Cells

[00221] Next, the effect of passaging on induction of antigen-specific CD8⁺ T cells was determined by intracellular cytokine staining with immunodominant peptides specific for MHC-class I using HIV-Gag peptide AMQMLKETI (SEQ ID No: 25) and LLO 91-99 (GYKDGNEYI; SEQ ID No: 26). Injection of 10³ CFU passaged bacteria (Lm-Gag) into mice elicited significant numbers of HIV-Gag-specific CD8⁺ T cells, while the same dose of non-passaged Lm-Gag induced no detectable Gag-specific CD8⁺ T cells. Even increasing the dose of unpassaged bacteria 100-fold did not compensate for their relative avirulence; in fact, no detectable Gag-specific CD8⁺ T cells were elicited even at the higher dose. The same dose increase with passaged bacteria increased Gag-specific T cell induction by 50% (Figure 8). The same pattern of induction of antigen-specific CD8⁺ T cells was observed with LLO-specific CD8⁺ T cells, showing that these results were not caused by the properties of the passenger antigen, since they were observed with LLO, an endogenous *Listeria* antigen.

[00222] Thus, passage through mice increases the immunogenicity of *Listeria* vaccine strains.

EXAMPLE 5: A PrfA-CONTAINING PLASMID IS STABLE IN AN LM STRAIN WITH A PrfA DELETION IN THE ABSENCE OF ANTIBIOTICS

MATERIALS AND EXPERIMENTAL METHODS***Bacteria***

[00223] *L. monocytogenes* strain XFL7 contains a 300 base pair deletion in the prfA gene. XFL7 carries pGG55 which partially restores virulence and confers CAP resistance, and is described in United States Patent Application Publication No. 200500118184.

Development of protocol for plasmid extraction from *Listeria*

[00224] 1 mL of *Listeria monocytogenes* Lm-LLO-E7 research working cell bank vial was inoculated into 27 mL BH1 medium containing 34 µg/mL CAP and grown for 24 hours at 37°C and 200 rpm.

[00225] Seven 2.5 mL samples of the culture were pelleted (15000 rpm for 5 minutes), and pellets were incubated at 37°C with 50 µl lysozyme solution for varying amounts of time, from 0-60 minutes.

[00226] Lysozyme solution:

- 29 µl 1 M dibasic Potassium Phosphate
- 21 µl 1 M monobasic Potassium Phosphate
- 500 µl 40% Sucrose (filter sterilized through 0.45 µm filter)
- 450 µl water

- 60 µl lysozyme (50 mg/mL)

[00227] After incubation with the lysozyme, the suspensions were centrifuged as before and the supernatants discarded. Each pellet was then subjected to plasmid extraction by a modified version of the QIAprep Spin Miniprep Kit® (Qiagen, Germantown, Maryland) protocol. The changes to the protocol were as follows:

1. The volumes of buffers P1, P2 and N3 were all increased threefold to allow complete lysis of the increased biomass.
2. 2 mg/mL of lysozyme was added to the resuspended cells before the addition of P2. The lysis solution was then incubated at 37°C for 15 minutes before neutralization.
3. The plasmid DNA was resuspended in 30 µL rather than 50 µL to increase the concentration.

[00228] In other experiments, the cells were incubated for 15min in P1 buffer + Lysozyme, then incubated with P2 (lysis buffer) and P3 (neutralization buffer) at room temperature.

[00229] Equal volumes of the isolated plasmid DNA from each subculture were run on an 0.8% agarose gel stained with ethidium bromide and visualized for any signs of structural or segregation instability.

[00230] The results showed that plasmid extraction from *L. monocytogenes* Lm-LLO-E7 increases in efficiency with increasing incubation time with lysozyme, up to an optimum level at approximately 50 minutes incubation.

[00231] These results provide an effective method for plasmid extraction from *Listeria* vaccine strains.

Replica plating

[00232] Dilutions of the original culture were plated onto plates containing LB or TB agar in the absence or presence of 34 µg/mL CAP. The differences between the counts on selective and non-selective agar were used to determine whether there was any gross segregational instability of the plasmid.

RESULTS

[00233] The genetic stability (i.e. the extent to which the plasmid is retained by or remains stably associated with the bacteria in the absence of selection pressure; e.g. antibiotic selection pressure) of the pGG55 plasmid in *L. monocytogenes* strain XFL7 in the absence of antibiotic was assessed by serial sub-culture in both Luria-Bertani media (LB: 5 g/L NaCl, 10 g/ml soy peptone, 5 g/L yeast extract) and Terrific Broth media (TB: 10 g/L glucose, 11.8 g/L soy peptone, 23.6 g/L yeast extract, 2.2 g/L KH₂PO₄, 9.4 g/L K₂HPO₄), in duplicate cultures. 50 mL of fresh media in a 250 mL baffled shake flask was inoculated with a fixed number of

cells (1 ODmL), which was then subcultured at 24 hour intervals. Cultures were incubated in an orbital shaker at 37°C and 200 rpm. At each subculture the OD₆₀₀ was measured and used to calculate the cell doubling time (or generation) elapsed, until 30 generations were reached in LB and 42 in TB. A known number of cells (15 OD mL) at each subculture stage
 5 (approximately every 4 generations) were pelleted by centrifugation, and the plasmid DNA was extracted using the Qiagen QIAprep Spin Miniprep® protocol described above. After purification, plasmid DNA was subjected to agarose gel electrophoresis, followed by ethidium bromide staining. While the amount of plasmid in the preps varied slightly between samples, the overall trend was a constant amount of plasmid with respect to the generational number of
 10 the bacteria (Figures 9A-B). Thus, pGG55 exhibited stability in strain XFL7, even in the absence of antibiotic.

[00234] Plasmid stability was also monitored during the stability study by replica plating on agar plates at each stage of the subculture. Consistent with the results from the agarose gel electrophoresis, there was no overall change in the number of plasmid-containing cells
 15 throughout the study in either LB or TB liquid culture (Figures 10 and 11, respectively).

[00235] These findings demonstrate that prfA-encoding plasmids exhibit stability in the absence of antibiotic in *Listeria* strains containing mutations in prfA.

MATERIALS AND METHODS (examples 6-10)

20 [00236] PCR reagents:

[00237] The primers used for amplification of the prfA gene and discrimination of the D133V mutation are shown in Table 1. Stock solutions of the primers ADV451, 452 and 453 were prepared by diluting the primers in TE buffer to 400 µM. An aliquot of the stock solution was further diluted to 20µM in water (PCR grade) to prepare a working solution. Primers were
 25 stored at -20°C. The reagents used in the PCR are shown in Table 2.

[00238] **Table 1.** Primers ADV451, 452 and 453.

Primer	Orientation	Sequence (5'→3')	Specificity
ADV451	Forward	CCTAGCTAAATTTAATGT (SEQ ID NO: 28)	D133V mutation
ADV452	Forward	CCTAGCTAAATTTAATGA (SEQ ID NO: 29)	Wild-type sequence
ADV453	Reverse	TAATTTTCCCCAAGTAGCAGG (SEQ ID NO: 30)	Shared sequence

[00239] **Table 2.** PCR reagents.

	Description	Provider	Catalog number
1	0.2 ml thin-walled PCR tubes: GeneAmp autoclaved reaction tube with cap	Applied Biosystems	N801-0612
2	Water (PCR reagent)	Sigma	W1754

3	Taq DNA Polymerase with 10x reaction buffer containing 15 mM MgCl ₂	Sigma	D1806
4	Set of deoxynucleotides (dNTPs), 10 mM each	Sigma	D7295
5	Primers ADV451, ADV452 and ADV453	Invitrogen	
6	Template DNA, midipreparations of pGG55 plasmids		
7	Thermal cycler PTC200 (48 wells block)	MJ Research	

Plasmid DNA preparation

[00240] pGG55 plasmids with (pGG55 D133V) and without (pGG55 WT) the *prfA* mutation were extracted and purified by midipreparations either from *E. coli* or *Listeria monocytogenes* using the PureLink™ HiPure Plasmid Midiprep Kit (Invitrogen, K2100-05), according to the manufacturer's instructions. For plasmid purification from *Listeria*, bacterial strains carrying the pGG55 D133V or WT plasmids were streak plated from frozen stocks in BHI agar plates supplemented with chloramphenicol (25 µg/ml). A single colony from each strain was grown in 5 ml of selective medium (BHI broth with 25 µg/ml of chloramphenicol) for 6 hours with vigorous shaking at 37°C and subinoculated 1:500 in 100 ml of selective medium for overnight growth under similar conditions. Bacteria from the overnight culture were harvested by centrifugation at 4,000 x g for 10 minutes and resuspended buffer R3 (resuspension buffer) containing 2 mg/ml of lysozyme (Sigma, L7001). The bacteria suspension was incubated for at least 1 hour at 37°C before proceeding to the regular protocol. Concentration and purity of the eluted plasmids were measured in a spectrophotometer at 260nm and 280nm. To prepare the template DNAs, the pGG55 D133V and WT plasmids were resuspended in water to a final concentration of 1 ng/µl from the midiprep stock solution. For the pGG55 WT plasmid, serial 10-fold dilutions from the 1 ng/µl solution were prepared, corresponding to dilutions from 10⁻¹ to 10⁻⁷.

prfA specific PCR protocol to test clinical grade material

[00241] The reaction mixture contained 1x PCR buffer, 1.5 mM MgCl₂, 0.8 mM dNTPs, 0.4 µM of each primer, 0.05 U/µl of Taq DNA polymerase and 0.04 ng/µl of the pGG55 D133V template plasmid. For each test, 10 tubes were required and the key components in each tube in a 25 µl reaction are shown in the Table 3. For the PCR reaction, a master mix was prepared with enough reagents for 11 reactions as shown in Table 4, and 24 µl of this PCR mix was

added to each tube. Subsequently, a total of 1 µl of the serially diluted pGG55 WT plasmid was added to the corresponding tubes: 1 ng in tube 3; 100 pg in tube 4; 10 pg in tube 5; 1 pg in tube 6; 100 fg in tube 7; 10 fg in tube 8; 1 fg in tube 9; 0.1 fg in tube 10. This serial dilution was used to calibrate a standard curve to determine the method sensitivity. Additionally, 0.5 µl of water and 0.5 µl of primer ADV451 (20 µM stock) were added in tube 1, and 1 µl of water added in tube 2, completing 25 µl of final volume. The quantities of each reagent per tube for a 25 µl reaction are shown in Table 5. The PCR cycling conditions used in the reaction are shown in Table 6.

[00242] After conclusion of the PCR reaction, 5 µl of gel-loading buffer (6x, with bromophenol blue) was added to each sample and 10 µl were analyzed by electrophoresis in 1.2% agarose gel in TBE buffer. The gel dimensions were 7 cm x 7 cm x 1 cm with a 15 sample wells (1 mm x 2 mm) comb. The gel was run at 100 V for ~30 minutes, until the bromophenol blue dye reached the middle of the gel. The gel was stained in ethidium bromide (0.5 µg/ml) for 20 minutes, destaining in water for 10 minutes. The gel is visualized by illumination with UV light and photographed. The image was analyzed using a band densitometry software (Quantity One version 4.5.1, BioRad).

[00243] **Table 3.** Set of individual PCR reactions to validate the method to detect the presence of wild-type prfA sequence in *Lm*-LLO-E7 samples.

Tube	Primer A	Primer B	Template DNA	Function	Expected result
1	ADV451	ADV453	1 ng of pGG55 (D133V)	Positive control for the ADV451 reaction	Positive
2	ADV452	ADV453	1 ng of pGG55 (D133V)	Negative control for the ADV452 reaction (specificity)	Negative
3	ADV452	ADV453	1 ng of pGG55 (wild-type) + 1 ng of pGG55 (D133V)	Positive control for the ADV452 reaction	Positive
4	ADV452	ADV453	100 pg of pGG55 (wild-type) + 1 ng of pGG55 (D133V)	Test the sensitivity of the reaction	Positive
5	ADV452	ADV453	10 pg of pGG55 (wild-type) + 1 ng of pGG55 (D133V)	Test the sensitivity of the reaction	Positive
6	ADV452	ADV453	1 pg of pGG55 (wild-type) + 1 ng of pGG55 (D133V)	Test the sensitivity of the reaction	Positive
7	ADV452	ADV453	100 fg of pGG55 (wild-type) + 1 ng pGG55 (D133V)	Test the sensitivity of the reaction	Positive
8	ADV452	ADV453	10 fg of pGG55 (wild-type) + pGG55 (D133V)	Test the sensitivity of the reaction	Positive
9	ADV452	ADV453	1 fg of pGG55 (wild-type) + pGG55 (D133V)	Test the sensitivity of the reaction	Weakly positive
10	ADV452	ADV453	0.1 fg of pGG55	Test the sensitivity of	To be

(wild-type) pGG55 (D133V)	+	the reaction	determined
------------------------------	---	--------------	------------

[00244] **Table 4.** Master PCR mix preparation.

Reagent	Quantity (μl)
Water	206.25
Taq DNA Polymerase 10x reaction buffer containing 15 mM MgCl ₂	27.5
Deoxynucleotides (dNTPs) 10 mM each	5.5
Primers ADV452 (20 μM in water)	5.5
Primers ADV453 (20 μM in water)	5.5
pGG55 D133V (<i>Lm</i> -LLO-E7) plasmid (1 ng/μl)	11
Taq DNA Polymerase (5 U/ μl)	2.75
Total	264

5 [00245] **Table 5.** PCR protocol for validation of the method to detect the presence of wild-type prfA sequence using primers ADV451, 452 and 453.

Reagent	PCR
Water	18.75 μl
PCR Buffer 10x + MgCl ₂ 15mM	2.5 μl
Deoxynucleotides mix (dATP, dCTP, dGTP and dTTP) 10mM each	0.5 μl
Primer ADV452 (20 μM)	0.5 μl
Primer ADV453 (20 μM)	0.5 μl
Taq DNA polymerase (5 U/μl)	0.25 μl
Template DNA (1 ng/μl) pGG55 D133V	1 μl
Template DNA pGG55 WT (tubes 3 to 10) ^a	1 μl
Final volume per tube ^b	25 μl

^a pGG55 WT (1 ng in tube 3; 100 pg in tube 4; 10 pg in tube 5; 1 pg in tube 6; 100 fg in tube 7; 10 fg in tube 8; 1 fg in tube 9; 0.1 fg in tube 10).

^b In tube 1, add 0.5 μl of water and 0.5 μl of primer ADV451 (20 μM stock); in tube 2 add 1 μl of water.

10

[00246] **Table 6.** PCR cycling conditions to detect the presence of wild-type prfA sequence using primers ADV451, 452 and 453.

Step	Temperature	Time	Number of cycles
1.	94°C	2 minutes and 30 seconds	1
2.	94°C	30 seconds	1
3.	53°C	30 seconds	1
4.	72°C	30 seconds	1
5.		Repeat steps 2 to 4	12
6.	94°C	30 seconds	1

7.	50°C	30 seconds	1
8.	72°C	30 seconds	1
9.	Repeat steps 6 to 8		23
10.	72°C	10 minutes	1

Sequencing:

[00247] Sequencing of the plasmids was done using the dideoxy sequencing method. The plasmids pGG55 D133V and pGG55 WT were mixed at different ratios (1:1, 1:10, 1:100, 1:1,000 and 1:10,000). The total amount of plasmid in the mixture was kept constant (500 µg) and the plasmid containing the wild-type sequence was 10-fold serially diluted in relation to the D133V plasmid to determine the sensitivity of the method.

RESULTS

EXAMPLE 6: SEQUENCING IS NOT A SENSITIVE METHOD TO DETECT THE REVERSION OF THE D133V MUTATION.

[00248] To estimate the sensitivity of sequencing in detecting the wild-type *prfA* sequence, the pGG55 D133V and WT plasmids were mixed at the different ratios and sequenced. The results are shown in Figure 12 and reveal that sequencing has a high specificity in discriminating the *prfA* D133V mutation (Figure 12). On the other hand, the sensitivity is low and the maximum dilution of wild-type *prfA* pGG55 plasmid with a detectable peak in the sequence was 1 in 10 (Figure 12). In conclusion, although sequencing is very specific, the sensitivity of the method is low and not appropriate to screen for the presence of rare events such as revertants of the *prfA* D133V mutation in *Lm*-LLO-E7 samples.

EXAMPLE 7: DEVELOPMENT OF A HIGHLY SPECIFIC AND SENSITIVE PCR METHOD TO DETECT REVERSION OF THE D133V MUTATION.

[00249] Given the low sensitivity of sequencing to detect rare events, it became imperative to develop a more sensitive method with similar specificity to detect reversion of the D133V mutation to wild-type. To achieve this goal, we designed a PCR-based method that specifically amplifies the wild-type sequence and is sensitive enough to detect at least 1 wild-type copy of *prfA* in 10,000,000 copies of the D133V mutated sequence. We designed 3 primers for this method: ADV451, ADV452 and ADV453 (Table 1). Both ADV451 and ADV452 are forward primers and differ in the last nucleotide at the 3' position to discriminate the A→T (D133V) mutation at position 398 of the *prfA* gene. The ADV453 primer is the

reverse primer located approximately 300 bp downstream the annealing site of the ADV451 and ADV452 primers (Figure 13). The expected PCR band obtained with the primers ADV451 or ADV452 and ADV453 is 326 bp. Under stringent conditions, the ADV451 primer should only amplify the pGG55 D133V plasmid, whereas the ADV452 would be
 5 specific to the wild-type *prfA* sequence.

EXAMPLE 8: SPECIFICITY OF THE PCR METHOD.

[00250] The reaction using the primer ADV451 was very specific and amplified the mutated D133V *prfA* sequence (lanes 1 to 3), but not the wild-type sequence (lanes 4 to 6). However, a
 10 very faint band can be detected in lane 4, when 5 ng of template DNA was used, but not with 1 ng (Figure 14).

[00251] As shown in Figure 15, the reaction with the ADV452 primer only amplified the wild-type *prfA* sequence (lanes 4, 5 and 6), and no bands were detected when the pGG55 carrying the D133V *prfA* mutation was used as a template (lanes 1, 2 and 3), even when using 5 ng of
 15 plasmid in the reaction (Figure 16). In conclusion, the PCR reactions with primers ADV451 and ADV452 are very specific and able to discriminate the A↔T (D133V) mutation at position 398 of the *prfA* gene in the pGG55 plasmid. Based on these results, we selected the amount of 1 ng as the standard amount of template DNA to be used in the reaction.

EXAMPLE 9: SENSITIVITY OF THE PCR METHOD.

[00252] The sensitivity of the reaction was tested using 1 ng of template DNA. For the plasmid carrying the wild-type *prfA* sequence, decreasing amounts of DNA (corresponding to 10-fold dilutions from 10^{-1} to 10^{-7}), were also included in the reaction to estimate the sensitivity. In these reactions only the primers ADV452 and ADV453 were used. In a PCR reaction with 30
 25 cycles (10 cycles with annealing temperature of 53°C and an additional 20 cycles with annealing temperature of 50°C), the sensitivity of the method was 1 in 100,000 (data not shown). As shown in figure 5, increasing the number of PCR cycles to 37 improved the visual sensitivity of the method to 10^{-6} for the detection of D133V revertants, without significantly compromising the specificity. A clear band was visible at the 10^{-6} dilution, corresponding to a
 30 detection level of 1 copy of the wild-type sequence in a million of the D133V mutant, when 1 ng of plasmid was used as the initial amount of DNA. Only a very weak band can be visualized in lanes 1 and 9 after longer exposure, reassuring the robust specificity of the method. On the other hand, when starting with 5 ng of DNA, a band could be easily detected at the 10^{-7} dilution, increasing the sensitivity of the PCR. However, a similar band in intensity

could also be detected with the pGG55 D133V plasmid, indicating the specificity limit of the method (Figure 17). This band observed with the pGG55 D133V plasmid is likely due to non-specific amplification of the D133V mutation with primer ADV452 that can significantly accumulate with the increased number of cycles. These results indicate that the sensitivity limit for this method, without significantly compromising the specificity, is situated between 1 to 1,000,000 and 1 to 10,000,000.

EXAMPLE 10: RECOMBINANT *LISTERIA* EXPRESSING A FUSION PROTEIN OF LLO TO E7 (LM-LLO-E7)

[00253] This strain is approx. 4 -5 logs more attenuated than the wild-type parent strain 10403S and secretes the fusion protein tLLO-E7. This immunotherapy is based on the backbone XFL7, which is derived from 10403S by the irreversible deletion in the virulence gene transcription activator *prfA*. PrfA regulates the transcription of several virulence genes such as Listeriolysin O (LLO), ActA, PlcA (phospholipase A), PlcB (phospholipase B) etc that are required for *in vivo* intracellular growth and survival of *L. monocytogenes*. The plasmid pGG55 is retained by the Lm-LLO-E7 *in vitro* by means of selection with ‘chloramphenicol’. However for *in vivo* retention of the plasmid by Lm-LLO-E7, it carries a copy of mutated *prfA* (D133V), which has been demonstrated to be less active than wild-type PrfA in DNA binding and activating the transcription of virulence genes. We have observed that complementation with mutated *prfA* resulted in approx. 40 fold reduction in the amount of secreted LLO from Lm-LLO-E7 when compared to wild-type strain 10403S. This implicates that the strain Lm-LLO-E7 likely exhibits a reduced expression of the virulence genes that are regulated by PrfA such as *actA*, *inlA*, *inlB*, *inlC*, *plcB* etc. In Lm-LLO-E7, the complementation with mutated copy of *prfA* likely causes a reduction in the expression of different virulence genes that are regulated by PrfA resulting in overall attenuation of approx. 4-5 logs.

[00254] While certain features of the invention have been illustrated and described herein, many modifications, substitutions, changes, and equivalents will now occur to those of ordinary skill in the art. It is, therefore, to be understood that the appended claims are intended to cover all such modifications and changes as fall within the true spirit of the invention.

What is claimed:

1. A recombinant *Listeria* strain, said recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, and wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein.
2. The recombinant *Listeria* of claim 1, wherein said *Listeria* comprises a deletion, inactivation or mutation in the *prfA* gene.
3. The recombinant *Listeria* of any one of claims 1-2, wherein said mutant PrfA protein comprises a D133V mutation.
4. The recombinant *Listeria* of any one of claims 2-3, wherein said mutant PrfA protein encoded by said second open reading frame complements said *prfA* genomic mutation, deletion or inactivation in said *Listeria* strain or restores partial PrfA function in said *Listeria* strain.
5. The recombinant *Listeria* of any one of claims 1-4, wherein said heterologous antigen is Human Papilloma Virus-E7 (HPV-E7) or HPV-E6.
6. The recombinant *Listeria* strain of any one of claims 1-5, wherein said N-terminal fragment of an LLO protein is selected from a sequence comprising SEQ ID NO: 2 or SEQ ID NO: 4.
7. The recombinant *Listeria* strain of any one of claims 1-6, wherein said recombinant *Listeria* strain is a recombinant *Listeria monocytogenes* strain.

8. The recombinant *Listeria* of any one of claims 1-7, wherein said recombinant *Listeria* strain has been passaged through an animal host, prior to the step of administering.
9. The recombinant *Listeria* of any one of claims 1-8, wherein said recombinant polypeptide is expressed by said recombinant *Listeria* strain.
10. The recombinant *Listeria* of any one of claims 1-9, wherein said recombinant *Listeria* strain comprises a plasmid that comprises said recombinant nucleic acid.
11. The recombinant *Listeria* of any one of claims 1-10, wherein said plasmid comprises a gene encoding a metabolic enzyme.
12. A pharmaceutical composition comprising the recombinant *Listeria* of any one of claims 1-11.
13. A method of inducing an immune response against a tumor or a cancer in a human subject, the method comprising the step of administering to said subject a recombinant *Listeria* strain comprising a recombinant nucleic acid, said nucleic acid comprising a first open reading frame encoding a recombinant polypeptide comprising an N-terminal fragment of an LLO protein fused to a heterologous antigen or fragment thereof, wherein said recombinant nucleic acid further comprises a second open reading frame encoding a mutant PrfA protein, thereby inducing an immune response against a tumor or a cancer.
14. The recombinant *Listeria* of claim 13, wherein said *Listeria* comprises a deletion, inactivation or mutation in the *prfA* gene.
15. The method of any one of claims 12-14, wherein said mutant PrfA protein comprises a D133V mutation.

16. The method *Listeria* of any one of claims 14-15, wherein said mutant PrfA protein encoded by said second open reading frame complements a *prfA* genomic mutation, deletion or inactivation in said *Listeria* strain or restores partial PrfA function in said *Listeria* strain.
17. The method of any one of claims 14-16, wherein said administering is intravenous or oral administering.
18. The method of any one of claims 14-17, wherein said heterologous antigen is Human Papilloma Virus-E7 (HPV-E7) or HPV-E6.
19. The method of any one of claims 14-18, wherein said N-terminal fragment of an LLO protein is selected from a sequence comprising SEQ ID NO: 2 or SEQ ID NO: 4.
20. The method of any one of claims 14-18, wherein said recombinant *Listeria* strain is administered to said human subject at a dose of 1×10^9 - 3.31×10^{10} organisms.
21. The method of claim 20, wherein said recombinant *Listeria* strain is stored in a frozen or lyophilized condition prior to administering.
22. The method of any one of claims 14-22, wherein said recombinant *Listeria* strain is a recombinant *Listeria monocytogenes* strain
23. The method of any one of claims 14-21, wherein said recombinant *Listeria* strain has been passaged through an animal host, prior to the step of administering.
24. The method of any one of claims 14-23, wherein said recombinant polypeptide is expressed by said recombinant *Listeria* strain.
25. The method of any one of claims 14-24, wherein said recombinant *Listeria* strain comprises a plasmid that encodes said recombinant polypeptide.

26. The method of any one of claims 14-25, wherein said plasmid comprises a gene encoding a metabolic enzyme.
27. The method of any one of claims 14-26, further comprising the step of boosting said human subject with said recombinant *Listeria* strain.
28. The method of any one of claims 14-27, further comprising the step of inoculating said human subject with an immunogenic composition that comprises or directs expression of said E7 antigen.
29. The method of any one of claims 14-28, wherein said immune response is a cytotoxic T cell anti-tumor immune response.
30. The method of any one of claims 14-29, wherein said method allows protecting a subject against a tumor or cancer.
31. The method of any one of claims 14-29, wherein said method allows treating a subject against a tumor or cancer.
32. The method of claims 30-31, wherein said cancer is cervical cancer, head and neck cancer (HNC) or anal cancer.




Figure 1




Figure 2

Figure 3




Figure 4

4/17








Figure 5

**Figure 6**


6/17

**Figure 7A****Figure 7B**

7/17

**Figure 8**


8/17



Lane	Sample	Lane	Sample
1	1Kb ladder	9	LB B, generation 5
2	100ng reference pGG55	10	LB A, generation 9
3	LB A, generation 5	11	LB B, generation 14
4	LB A, generation 9	12	LB B, generation 19*
5	LB A, generation 14	13	LB B, generation 24*
6	LB A, generation 19	14	LB B, generation 29*
7	LB A, generation 24	15	1Kb ladder
8	LB A, generation 29		

* Residual ethanol remaining in sample, therefore the majority of the sample did not load into the well, resulting in a less intense plasmid band

Figure 9A



Lane	Sample	Lane	Sample
1	1Kb ladder	5	TB, generation 21
2	100ng reference pGG55	6	TB, generation 28
3	TB, generation 7	7	TB, generation 35
4	TB, generation 14	8	TB, generation 42

Figure 9B




Figure 9C

10/17




Figure 10






Figure 11

11/17


pGG55 D133V only (1:0)



pGG55 WT only (0:1)




pGG55 D133V + pGG55 WT (1:1)




Two peaks are clearly visible and indicated by the arrow when plasmids are mixed in equimolar amounts.

pGG55 D133V + pGG55 WT (10:1)



1 in 10 is the maximum limit of detection and the peak from the wild-type sequence is still clearly detectable.

pGG55 D133V + pGG55 WT (100:1)



The peak from the wild-type sequence has disappeared completely when the wild-type prfA plasmid is diluted 1 to 100 or more.

Figure 12






Figure 13

13/17




1. 5 ng of pGG55 D133V;
2. 1 ng of pGG55 D133V;
3. 0.2 ng of pGG55 D133V;
4. 5 ng of pGG55 wild-type;
5. 1 ng of pGG55 wild-type;
6. 0.2 ng of pGG55 wild-type;

Figure 14



- 1) 5 ng of pGG55 D133V;
- 2) 1 ng of pGG55 D133V;
- 3) 0.2 ng of pGG55 D133V;
- 4) 5 ng of pGG55 wild-type;
- 5) 1 ng of pGG55 wild-type;
- 6) 0.2 ng of pGG55 wild-type;


Figure 15



1. 1 ng of pGG55 D133V
2. 1 ng of pGG55 wild-type
3. 100 pg of pGG55 wild-type (10^{-1})
4. 10 pg of pGG55 wild-type (10^{-2})
5. 1 pg of pGG55 wild-type (10^{-3})
6. 100 fg of pGG55 wild-type (10^{-4})
7. 10 fg of pGG55 wild-type (10^{-5})
8. 1 fg of pGG55 wild-type (10^{-6})

Figure 16

14/17



1. 5 ng of pGG55 D133V
2. 5 ng of pGG55 wild-type
3. 500 pg of pGG55 wild-type (10^{-1})
4. 50 pg of pGG55 wild-type (10^{-2})
5. 5 pg of pGG55 wild-type (10^{-3})
6. 500 fg of pGG55 wild-type (10^{-4})
7. 50 fg of pGG55 wild-type (10^{-5})
8. 5 fg of pGG55 wild-type (10^{-6})

Figure 17




Figure 18

15/17




Figure 19




Figure 20

16/17




Figure 21






Figure 22


pGG55 D133V only (1:0)



pGG55 WT only (0:1)




pGG55 D133V + pGG55 WT (1:1)




Two peaks are clearly visible and indicated by the arrow when plasmids are mixed in equimolar amounts.

pGG55 D133V + pGG55 WT (10:1)



1 in 10 is the maximum limit of detection and the peak from the wild-type sequence is still clearly detectable.

pGG55 D133V + pGG55 WT (100:1)



The peak from the wild-type sequence has disappeared completely when the wild-type prfA plasmid is diluted 1 to 100 or more.

Figure 12