



(12) **DEMANDE DE BREVET CANADIEN**
CANADIAN PATENT APPLICATION
(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2018/03/13
(87) Date publication PCT/PCT Publication Date: 2018/10/25
(85) Entrée phase nationale/National Entry: 2019/10/16
(86) N° demande PCT/PCT Application No.: JP 2018/009679
(87) N° publication PCT/PCT Publication No.: 2018/193752
(30) Priorités/Priorities: 2017/04/18 (JP2017-082423);
2017/04/24 (JP2017-085741);
2017/05/25 (JP2017-103935);
2017/09/12 (JP PCT/JP2017/032931)

(51) Cl.Int./Int.Cl. *A61K 45/00* (2006.01),
A23L 33/10 (2016.01), *A61K 31/194* (2006.01),
A61K 9/20 (2006.01), *A61P 13/12* (2006.01),
A61P 19/06 (2006.01), *A61P 9/00* (2006.01),
A61P 9/10 (2006.01)
(71) Demandeurs/Applicants:
TOHOKU UNIVERSITY, JP;
NIPPON CHEMIPHAR CO., LTD., JP
(72) Inventeurs/Inventors:
ABE, MICHIAKI, JP;
KOSHIBA, SEIZO, JP;
NISHIOKA, KOICHIRO, JP;
KAWAGUCHI, KAZUHIKO, JP; ...

(54) Titre : PROCEDE DE PURIFICATION DU SANG A L'AIDE D'UN AGENT D'ALCALINISATION
(54) Title: BLOOD PURIFICATION BY ALKALINIZING AGENT

(57) **Abrégé/Abstract:**

The present invention relates to a food composition containing an alkalifying agent, in particular an alkali metal salt of citric acid. Renal function is maintained through ingestion of this food composition.

(72) **Inventeurs(suite)/Inventors(continued):** YAMASAKI, SATOMI, JP; TERANAKA, YASUYUKI, JP

(74) **Agent:** BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

174

ABSTRACT

The present invention relates to a food composition comprising an alkalinizing agent, particularly an alkali metal salt of citric acid. Ingestion of the food composition
5 maintains kidney function.

DESCRIPTION

BLOOD PURIFICATION BY ALKALINIZING AGENT

5

TECHNICAL FIELD

[0001]

The present invention relates to blood purification by an alkalinizing agent.

Priority is claimed on Japanese Patent Application No. 2017-82423, filed on April 18, 2017; Japanese Patent Application No. 2017-85741, filed on April 24, 2017; Japanese Patent Application No. 2017-103935, filed on May 25, 2017; and International Application No. PCT/JP2017/032931, filed on September 12, 2017, the contents of which are incorporated herein by reference.

BACKGROUND ART

15

[0002]

The number of patients with end-stage kidney disease (ESKD) who require dialysis and transplantation is increasing worldwide. The number has tended to increase also in Japan, and the number of dialysis patients at the end of 2014 was 320,000.

Chronic kidney disease (CKD) is recognized to be likely to get ESKD. CKD is a concept encompassing kidney disease that chronically progresses regardless of the underlying disease, and is a concept encompassing all clinical conditions in which kidney function indicated by glomerular filtration rate (GFR) deteriorates, or findings that suggest damage in kidneys is maintained chronically (3 months or longer). Because CKD has not only a risk of progression to ESKD, but also a strong risk of developing into cardiovascular disease (CVD) etc., it is very important to detect CKD in early stages

and perform appropriate treatment. Many CKD treatment methods have been established so far, but they are still insufficient, and further development of nephroprotective agents is required.

[0003]

5 In CKD, various uremic substances accumulate in the body as renal clearance decreases. Among them, a concentration of indoxyl sulfate, which is a metabolic end-product of tryptophan, in blood increases with progression of CKD, and a high concentration (100 μ M to 1 mM) of indoxyl sulfate accumulates in the blood. Indoxyl sulfate is known to be deeply involved in progression of kidney damage due to kidney
10 fibrosis, and CKD complications such as CVD due to vascular calcification. It is reported that a concentration of indoxyl sulfate in serum correlates with a mortality rate and incidence of cardiovascular events in dialysis patients (Non-Patent Literature 1). In addition, it is considered that, by reducing a concentration of indoxyl sulfate in blood in CKD patients, progression to ESKD can be suppressed, and onset of CVD related to
15 kidney failure can be suppressed. In fact, a spherical carbonaceous adsorbent (KREMEZIN (registered trademark)) which adsorbs indole, which is a precursor of indoxyl sulfate, in the intestinal tract to decrease a concentration of indoxyl sulfate in blood, delays introduction of dialysis in CKD patients, thereby ameliorating arteriosclerosis (Non-Patent Literature 2).

20 [0004]

Meanwhile, because a concentration of bicarbonate ions (HCO_3^-) in blood decreases and metabolic acidosis develops in patients with advanced CKD, an alkalinizing agent such as sodium bicarbonate or a citric acid preparation is administered. In addition, it is reported that progression of CKD is suppressed by administration of
25 sodium bicarbonate that is an alkalinizing agent (Non-Patent Literature 3).

Furthermore, it is reported that oral administration of sodium bicarbonate suppresses kidney tubular cell damage due to acidic urine in an animal model with nephrosis caused by protein overload (Non-Patent Literature 4).

However, there is no report regarding suppression of progression of kidney damage by administering an alkalinizing agent to early-stage CKD patients, and there is also no report regarding a decrease in concentration of uremic substance in blood.

CITATION LIST

NON-PATENT LITERATURE

10 [0005]

[Non-Patent Literature 1] Barreto, F. C., et al.: Serum indoxyl sulfate is associated with vascular disease and mortality in chronic kidney disease patients. Clin. J. Am. Soc. Nephrol., 4: 1551-1558, 2009.

[Non-Patent Literature 2] Nakamura T., et al.: Oral ADSORBENT AST-120 decreases carotid intima-media thickness and arterial stiffness in patients with chronic renal failure. Kidney Blood Press Res, 27: 121-6, 2004.

[Non-Patent Literature 3] Brito-Ashurst, I. D., et al.: Bicarbonate supplementation slows progression of CKD and improves nutritional status. J. Am. Soc. Nephrol., 20: 2075-2084, 2009.

20 [Non-Patent Literature 4] Souma T., et al.: Luminal alkalinization attenuates proteinuria-induced oxidative damage in proximal tubular cells. J. Am. Soc. Nephrol., 22: 635-648, 2011.

SUMMARY OF THE INVENTION

25 PROBLEMS TO BE SOLVED BY THE INVENTION

[0006]

An object of the present invention is to provide a medicine useful for blood purification in a patient with kidney disease. Another object of the present invention is to provide a medicine useful for suppressing progression of chronic kidney disease (advancing in severity of chronic kidney disease), treating and preventing uremic symptoms, and delaying introduction of dialysis. Still another object of the present invention is to provide a medicine useful for suppressing progression from acute kidney failure to chronic kidney disease. Still another object of the present invention is to provide a food for promoting excretion of uremic substance outside the body. Still another object of the present invention is to provide a food for maintenance of kidney function (for example, for suppression of kidney tubular damage, for protection of kidney tubular cells, or for maintenance of kidney tubular function). Still another object of the present invention is to provide a method for determining suppression of progression of chronic kidney disease, or a method for determining a decrease in concentration of uremic substance in blood, and/or a promotion of excretion of uremic substance into urine.

MEANS TO SOLVE THE PROBLEMS

[0007]

The inventors of the present invention have performed extensive studies to achieve the above-mentioned objects. As a result, they have found that an agent that alkalinizes a body fluid is useful for promoting excretion of uremic substance from the body of a patient with kidney disease (for example, promoting excretion of uremic substance into urine), and therefore have completed the present invention.

25 [0008]

In one aspect, the present invention provides a pharmaceutical composition for promoting excretion of uremic substance outside the body, comprising an alkalinizing agent.

[0009]

5 In one aspect, the present invention provides a pharmaceutical composition for decreasing a concentration of uremic substance in blood, comprising an alkalinizing agent.

[0010]

10 In one aspect, the present invention provides a pharmaceutical composition for promoting excretion into urine in chronic kidney disease, comprising an alkalinizing agent.

[0011]

15 In one aspect, the present invention provides a pharmaceutical composition for ameliorating uremic symptoms in chronic kidney disease, comprising an alkalinizing agent.

[0012]

In one aspect, the present invention provides a pharmaceutical composition for delaying introduction of dialysis in chronic kidney disease, comprising an alkalinizing agent.

20 [0013]

In one aspect, the present invention provides a pharmaceutical composition for treating or preventing cardiovascular disease associated with chronic kidney disease, comprising an alkalinizing agent.

[0014]

25 In one aspect, the present invention provides a pharmaceutical composition for

suppressing progression from acute kidney failure to chronic kidney disease, comprising an alkalinizing agent.

[0015]

In one aspect, the present invention provides a food composition for promoting
5 excretion of uremic substance outside the body, comprising an alkalinizing agent.

[0016]

In one aspect, the present invention provides a method for determining suppression of progression of chronic kidney disease.

[0017]

10 In one aspect, the present invention provides a method for determining a decrease in concentration of uremic toxins in blood of a human, and/or a promotion of excretion of uremic toxins into urine.

[0018]

That is, the present invention has the following aspects.

15 (1) A pharmaceutical composition for decreasing a concentration of uremic substance in blood, comprising an alkalinizing agent.

(2) A pharmaceutical composition for promoting excretion of uremic substance into urine, comprising an alkalinizing agent.

(3) A pharmaceutical composition for promoting excretion of uremic substance
20 outside the body, comprising an alkalinizing agent.

(4) The pharmaceutical composition according to (2) or (3), wherein the excretion into urine or the excretion outside the body depends on a concentration of uremic substance in blood.

(5) The pharmaceutical composition according to any one of (1) to (4), which is
25 administered to a patient with chronic kidney disease or acute kidney failure.

(6) The pharmaceutical composition according to any one of (1) to (5), wherein the uremic substance is at least one selected from the group consisting of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.

5 (7) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance are indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.

(8) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance are indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine.

10 (9) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance is indoxyl sulfate and phenylacetyl-L-glutamine.

(10) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance is indoxyl sulfate and hippuric acid.

(11) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance is phenylacetyl-L-glutamine and p-cresyl sulfate.

15 (12) The pharmaceutical composition according to any one of (1) to (6), wherein the uremic substance is indoxyl sulfate.

(13) A pharmaceutical composition for ameliorating uremic symptoms in chronic kidney disease, comprising an alkalinizing agent.

20 (14) A pharmaceutical composition for delaying introduction of dialysis in chronic kidney disease, comprising an alkalinizing agent.

(15) A pharmaceutical composition for treating or preventing cardiovascular disease associated with chronic kidney disease, comprising an alkalinizing agent.

(16) The pharmaceutical composition according to (15), which ameliorates arteriosclerosis.

25 (17) A pharmaceutical composition for suppressing progression of chronic

kidney disease, comprising an alkalinizing agent.

(18) A pharmaceutical composition for treating or preventing kidney tubular damage, comprising an alkalinizing agent.

(19) The pharmaceutical composition according to any one of (1) to (18), which
5 is administered to a patient with early-stage chronic kidney disease.

(20) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient with stage G3b or lower chronic kidney disease.

(21) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient with stage G2 to stage G3b chronic kidney disease.

10 (22) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient with stage G2 and stage G3a chronic kidney disease.

(23) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient with stage G2 chronic kidney disease.

(24) The pharmaceutical composition according to any one of (1) to (18), which
15 is administered to a patient in which a urinary β 2-microglobulin concentration is 290 μ g/L or less.

(25) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient in which a urinary β 2-microglobulin concentration is 50 to 150 μ g/L.

20 (26) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient in which a blood cystatin C concentration is 0.5 to 2.2 mg/L.

(27) The pharmaceutical composition according to any one of (1) to (18), which is administered to a patient in which a blood cystatin C concentration is 1.0 to 1.3 mg/L.

(28) A pharmaceutical composition for suppressing progression from acute
25 kidney failure to chronic kidney disease, comprising an alkalinizing agent.

(29) The pharmaceutical composition according to any one of (1) to (28), wherein the alkalinizing agent is a pharmaceutically acceptable salt of citric acid or a hydrate thereof, or a mixture thereof.

(30) The pharmaceutical composition according to any one of (1) to (29),
5 wherein the alkalinizing agent is a sodium citrate, potassium citrate, or a hydrate thereof, or a mixture thereof.

(31) The pharmaceutical composition according to any one of (1) to (30), wherein the alkalinizing agent comprises a mixture of sodium citrate or a hydrate thereof, and potassium citrate or a hydrate thereof.

10 (32) The pharmaceutical composition according to any one of (1) to (31), wherein the alkalinizing agent is sodium citrate or a hydrate thereof.

(33) The pharmaceutical composition according to any one of (1) to (32), which is a tablet.

(34) The pharmaceutical composition according to any one of (1) to (33),
15 wherein a decrease in concentration of the uremic substance in blood compared to before the start of administration of the alkalinizing agent is detected 12 weeks after the administration of the alkalinizing agent.

(35) The pharmaceutical composition according to any one of (1) to (34), wherein an increase in concentration of the uremic substance in urine compared to before
20 the start of administration of the alkalinizing agent is detected 12 weeks after the administration of the alkalinizing agent.

(36) The pharmaceutical composition according to any one of (1) to (35), wherein an increase in amount of the uremic substance excreted outside the body compared to before the start of administration of the alkalinizing agent is detected 12
25 weeks after the administration of the alkalinizing agent.

(37) The pharmaceutical composition according to any one of (1) to (36), wherein administration of the alkalinizing agent suppresses an increase of a urinary β 2-microglobulin concentration.

(38) The pharmaceutical composition according to any one of (1) to (37),
5 wherein, by administration of the alkalinizing agent, after 12 weeks from the administration, an increase of a urinary β 2-microglobulin concentration is suppressed.

(39) The pharmaceutical composition according to any one of (1) to (38), wherein, by administration of the alkalinizing agent, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the start of the
10 administration.

(40) The pharmaceutical composition according to any one of (1) to (39), wherein, by administration of the alkalinizing agent, after 12 weeks from the administration, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the start of the administration.

(41) The pharmaceutical composition according to any one of (1) to (40),
15 wherein administration of the alkalinizing agent does not substantially increase blood cystatin C compared to before the start of the administration.

(42) The pharmaceutical composition according to any one of (1) to (41), wherein, by administration of the alkalinizing agent, after 12 weeks from the
20 administration, blood cystatin C is not substantially increased compared to before the start of the administration.

(43) The pharmaceutical composition according to any one of (1) to (42), wherein, by administration of the alkalinizing agent, amelioration of kidney proximal tubular damage and/or amelioration of glomerular damage is not recognized compared to
25 before the start of the administration; and a decrease in a concentration of the uremic

substance in blood, a promotion of excretion of the uremic substance into urine, and/or a promotion of excretion of the uremic substance outside the body is recognized compared to before the start of the administration.

(44) The pharmaceutical composition according to any one of (1) to (43),
5 wherein, by administration of the alkalinizing agent, after 12 weeks from the administration of the alkalinizing agent, amelioration of kidney proximal tubular damage and/or amelioration of glomerular damage is not recognized compared to before the start of the administration; and a decrease in a concentration of the uremic substance in blood, a promotion of excretion of the uremic substance into urine, and/or a promotion of
10 excretion of the uremic substance outside the body is recognized, compared to before the start of the administration.

(45) The pharmaceutical composition according to any one of (1) to (44), wherein the alkalinizing agent is administered at 1 to 3 g/day.

(46) The pharmaceutical composition according to any one of (1) to (45),
15 wherein the alkalinizing agent is administered at 1 to 1.5 g/day.

(47) The pharmaceutical composition according to any one of (1) to (46), wherein the alkalinizing agent is administered such that a pH of early morning urine becomes pH 5.2 to pH 6.8.

(48) The pharmaceutical composition according to any one of (1) to (47),
20 wherein the alkalinizing agent is administered such that a pH of early morning urine becomes pH 6.0 or more and less than pH 6.2.

(49) The pharmaceutical composition according to any one of (1) to (48), wherein the alkalinizing agent is administered for 12 weeks or longer.

(50) The pharmaceutical composition according to any one of (1) to (49),
25 wherein the alkalinizing agent is administered for 12 weeks.

(50-1) The pharmaceutical composition according to any one of (1) to (50), wherein administration of the alkalinizing agent increases an amount of uremic substance in urine.

(50-2) The pharmaceutical composition according to any one of (1) to (50) and (50-1), wherein administration of the alkalinizing agent increases a concentration of uremic substance in urine.

(50-3) The pharmaceutical composition according to any one of (1) to (50), (50-1), and (50-2), wherein administration of the alkalinizing agent exerts an effect of decreasing a concentration of uremic substance in blood, an effect of promoting excretion of uremic substance into urine, and/or an effect of promoting excretion of uremic substance outside the body, and the effects are recognized with respect to placebo administration.

(50-4) The pharmaceutical composition according to any one of (1) to (50) and (50-1) to (50-3), wherein administration of the alkalinizing agent exerts an effect of decreasing a concentration of uremic substance in blood, an effect of promoting excretion of uremic substance into urine, and/or an effect of promoting excretion of uremic substance outside the body, and the effects are recognized with respect to before administration of the alkalinizing agent.

(50-5) The pharmaceutical composition according to any one of (1) to (50) and (50-1) to (50-4), which is a tablet.

(50-6) The pharmaceutical composition according to any one of (1) to (50) and (50-1) to (50-5), wherein the urine is early morning urine.

(51) A food composition for decreasing a concentration of uremic substance in blood, promoting excretion of uremic substance into urine, and/or promoting excretion of uremic substance outside the body, wherein the food composition comprises an

alkalinizing agent.

(52) The food composition according to (51), wherein the alkalinizing agent is an alkali metal salt of citric acid or a hydrate thereof, or a mixture thereof.

5 (52-1) The food composition according to (51) or (52), which maintains kidney function (suppresses kidney tubular damage, protects kidney tubular cells, or maintains kidney tubular function).

(52-2) The food composition according to any one of (51), (52), and (52-1), wherein the alkalinizing agent is ingested at 1 to 3 g/day.

10 (52-3) The food composition according to any one of (51), (52), (52-1), and (52-2), wherein the alkalinizing agent is ingested at 1 to 1.5 g/day.

(52-4) The food composition according to any one of (51), (52), and (52-1) to (52-3), wherein a concentration of uremic substance in blood is decreased compared to before the start of ingesting the alkalinizing agent, or compared to placebo ingestion.

15 (52-5) The food composition according to any one of (51), (52), and (52-1) to (52-4), wherein a concentration of uremic substance in urine increases compared to before the start of ingesting the alkalinizing agent, or compared to placebo ingestion.

(52-6) The food composition according to any one of (51), (52), and (52-1) to (52-5), wherein an amount of uremic substance in urine increases compared to before the start of ingesting the alkalinizing agent, or compared to placebo ingestion.

20 (52-7) The food composition according to any one of (51), (52), and (52-1) to (52-6), wherein an amount of uremic substance excreted outside the body increases compared to before the start of ingesting the alkalinizing agent, or compared to placebo ingestion.

25 (52-8) The food composition according to any one of (51), (52), and (52-1) to (52-7), wherein ingestion of the alkalinizing agent suppresses an increase of a urinary β 2-

microglobulin concentration.

(52-9) The food composition according to any one of (51), (52), and (52-1) to (52-8), wherein, by ingestion of the alkalinizing agent, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the ingestion of the
5 alkalinizing agent or ingestion of placebo.

(52-10) The food composition according to any one of (51), (52), and (52-1) to (52-9), wherein ingestion of the alkalinizing agent does not substantially increase blood cystatin C.

(52-11) The food composition according to any one of (51), (52), and (52-1) to
10 (52-10), wherein the alkalinizing agent is ingested for 12 weeks or longer.

(52-12) The food composition according to any one of (51), (52), and (52-1) to (52-11), wherein the alkalinizing agent is ingested for 12 weeks.

(53) A method for determining suppression of progression of chronic kidney disease, comprising measuring a pH of urine.

15 (54) A method for determining a decrease in concentration of uremic substance in blood of a patient with chronic kidney disease, comprising measuring a pH of urine.

(55) A method for determining promotion of excretion of uremic substance into urine of a patient with chronic kidney disease, comprising measuring a pH of urine.

[0019]

20 Furthermore, the present invention has the following aspects.

(56) A pharmaceutical composition for decreasing a concentration of uremic substance in blood, wherein the pharmaceutical composition comprises an alkalinizing agent, and the pharmaceutical composition is a tablet.

(57) A pharmaceutical composition for promoting excretion of uremic substance
25 into urine, wherein the pharmaceutical composition comprises an alkalinizing agent, and

the pharmaceutical composition is a tablet.

(58) The pharmaceutical composition according to (56) or (57), which is administered to a patient with chronic kidney disease or acute kidney failure.

(59) The pharmaceutical composition according to any one of (56) to (58),
5 wherein the uremic substance is at least one selected from the group consisting of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.

(60) The pharmaceutical composition according to any one of (56) to (59), wherein the uremic substance is indoxyl sulfate.

10 (61) A pharmaceutical composition for suppressing progression of chronic kidney disease, wherein the pharmaceutical composition comprises an alkalinizing agent, and the pharmaceutical composition is a tablet.

(62) A pharmaceutical composition for treating or preventing kidney tubular damage, wherein the pharmaceutical composition comprises an alkalinizing agent, and
15 the pharmaceutical composition is a tablet.

(63) The pharmaceutical composition according to any one of (56) to (62), wherein the alkalinizing agent is a pharmaceutically acceptable salt of citric acid or a hydrate thereof, or a mixture thereof.

(64) The pharmaceutical composition according to any one of (56) to (63),
20 wherein the alkalinizing agent comprises a mixture of sodium citrate or a hydrate thereof, and potassium citrate or a hydrate thereof.

[0020]

Furthermore, the present invention has the following aspects.

(65) A food composition which is for maintenance of kidney function,
25 comprising an alkalinizing agent.

(66) The food composition according to (65), wherein the maintenance of kidney function is suppression of kidney tubular damage, protection of kidney tubular cells, or maintenance of kidney tubular function.

(67) The food composition according to (66), wherein the kidney tubule is a
5 kidney proximal tubule.

(68) The food composition according to any one of (65) to (67), wherein the alkalinizing agent is an acceptable salt of citric acid as food or a hydrate thereof, or a mixture thereof.

(69) The food composition according to any one of (65) to (68), wherein the
10 alkalinizing agent comprises a mixture of sodium citrate or a hydrate thereof, and potassium citrate or a hydrate thereof.

(70) The food composition according to any one of (65) to (69), wherein the alkalinizing agent is sodium citrate or a hydrate thereof.

(71) The food composition according to any one of (65) to (70), which is a
15 tablet.

(72) The food composition according to any one of (65) to (71), wherein an effect of the maintenance of kidney function is indicated on packaging, a container, or an instruction leaflet for the food composition.

(73) The food composition according to any one of (65) to (71), wherein an
20 effect of suppression of kidney tubular damage, protection of kidney tubular cells, or maintenance of kidney tubular function is indicated on packaging, a container, or an instruction leaflet for the food composition.

(74) The food composition according to any one of (65) to (72), which is ingested by a healthy subject concerned about kidney health.

25 (75) The food composition according to any one of (65) to (71) and (73), which

is ingested by a healthy subject concerned about kidney tubular health.

(76) The food composition according to any one of (65) to (75), which is ingested by a subject in which a urinary β 2-microglobulin concentration is 290 μ g/L or less.

5 (77) The food composition according to any one of (65) to (76), which is ingested by a subject in which a urinary β 2-microglobulin concentration is 50 to 150 μ g/L.

(78) The food composition according to any one of (65) to (77), which is ingested by a subject in which a blood cystatin C concentration is 0.5 to 2.2 mg/L.

10 (79) The food composition according to any one of (65) to (78), which is ingested by a subject in which a blood cystatin C concentration is 1.0 to 1.3 mg/L.

(80) The food composition according to any one of (65) to (79), wherein ingestion of the food composition suppresses an increase of a urinary β 2-microglobulin concentration.

15 (81) The food composition according to any one of (65) to (80), wherein, by ingestion of the food composition, after 12 weeks from the ingestion, an increase of a urinary β 2-microglobulin concentration is suppressed.

(82) The food composition according to any one of (65) to (81), wherein, by ingestion of the food composition, a urinary β 2-microglobulin concentration is not
20 substantially decreased compared to before the start of the ingestion or placebo.

(83) The food composition according to any one of (65) to (82), wherein, by ingestion of the food composition, after 12 weeks from the ingestion, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the start of administration or placebo.

25 (84) The food composition according to any one of (65) to (83), wherein

ingestion of the food composition does not substantially increase blood cystatin C compared to before the start of the ingestion.

(85) The food composition according to any one of (65) to (84), wherein ingestion of the food composition does not substantially increase blood cystatin C compared to placebo.

(86) The food composition according to any one of (65) to (85), wherein ingestion of the food composition suppresses, in early morning urine, an increase of a β 2-microglobulin amount which is associated with progression of a stage of chronic kidney disease.

(87) The food composition according to any one of (65) to (86), wherein ingestion of the food composition does not affect glomerular function in a patient with chronic kidney disease, and suppresses kidney proximal tubular cell damage associated with progression of a stage of chronic kidney disease to protect kidney proximal tubular cells.

EFFECTS OF THE INVENTION

[0021]

By the pharmaceutical composition and the like provided by the present invention, uremic substance is excreted outside the body in mammals. By the method provided by the present invention, it is possible to preliminarily determine as to whether or not uremic substance is excreted outside the body and/or whether or not a progression of chronic kidney disease can be suppressed. By the food composition and the like provided by the present invention, it is possible to maintain kidney function in mammals, more specifically, to suppress kidney tubular damage, protect kidney tubular cells, or maintain kidney tubular function.

BRIEF DESCRIPTION OF DRAWINGS

[0022]

FIG. 1 is a graph showing a correlation between a concentration of indoxyl sulfate in urine and a concentration of indoxyl sulfate in plasma of patients of a control group after 6, 12, and 24 weeks from the start of the test.

FIG. 2 is a graph showing a correlation between a concentration of indoxyl sulfate in urine and a concentration of indoxyl sulfate in plasma of patients of a group to which a combination preparation of hydrates of potassium citrate and sodium citrate has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 3 is a graph showing a correlation between a concentration of indoxyl sulfate in urine and a concentration of indoxyl sulfate in plasma of patients of a group to which a sodium bicarbonate preparation has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 4 is a graph showing a correlation between a concentration of indoxyl sulfate in urine and a concentration of indoxyl sulfate in plasma of all the patients after 6, 12, and 24 weeks from the start of the test.

FIG. 5 is a graph showing a correlation between a concentration of p-cresyl sulfate in urine and a concentration of p-cresyl sulfate in plasma of patients of a control group after 6, 12, and 24 weeks from the start of the test.

FIG. 6 is a graph showing a correlation between a concentration of p-cresyl sulfate in urine and a concentration of p-cresyl sulfate in plasma of patients of a group to which a combination preparation of hydrates of potassium citrate and sodium citrate has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 7 is a graph showing a correlation between a concentration of p-cresyl

sulfate in urine and a concentration of p-cresyl sulfate in plasma of patients of a group to which a sodium bicarbonate preparation has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 8 is a graph showing a correlation between a concentration of p-cresyl sulfate in urine and a concentration of p-cresyl sulfate in plasma of all the patients after 6, 12, and 24 weeks from the start of the test.

FIG. 9 is a graph showing a correlation between a concentration of hippuric acid in urine and a concentration of hippuric acid in plasma of patients of a control group after 6, 12, and 24 weeks from the start of the test.

FIG. 10 is a graph showing a correlation between a concentration of hippuric acid in urine and a concentration of hippuric acid in plasma of patients of a group to which a combination preparation of hydrates of potassium citrate and sodium citrate has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 11 is a graph showing a correlation between a concentration of hippuric acid in urine and a concentration of hippuric acid in plasma of patients of a group to which a sodium bicarbonate preparation has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 12 is a graph showing a correlation between a concentration of hippuric acid in urine and a concentration of hippuric acid in plasma of all the patients after 6, 12, and 24 weeks from the start of the test.

FIG. 13 is a graph showing a correlation between a concentration of argininosuccinic acid in urine and a concentration of argininosuccinic acid in plasma of patients of a control group after 6, 12, and 24 weeks from the start of the test.

FIG. 14 is a graph showing a correlation between a concentration of argininosuccinic acid in urine and a concentration of argininosuccinic acid in plasma of

patients of a group to which a combination preparation of hydrates of potassium citrate and sodium citrate has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 15 is a graph showing a correlation between a concentration of argininosuccinic acid in urine and a concentration of argininosuccinic acid in plasma of patients of a group to which a sodium bicarbonate preparation has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 16 is a graph showing a correlation between a concentration of argininosuccinic acid in urine and a concentration of argininosuccinic acid in plasma of all the patients after 6, 12, and 24 weeks from the start of the test.

FIG. 17 is a graph showing a correlation between a concentration of phenylacetyl-L--L-glutamine in urine and a concentration of phenylacetyl-L--L-glutamine in plasma of patients of a control group after 6, 12, and 24 weeks from the start of the test.

FIG. 18 is a graph showing a correlation between a concentration of phenylacetyl-L--L-glutamine in urine and a concentration of phenylacetyl-L--L-glutamine in plasma of patients of a group to which a combination preparation of hydrates of potassium citrate and sodium citrate has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 19 is a graph showing a correlation between a concentration of phenylacetyl-L--L-glutamine in urine and a concentration of phenylacetyl-L-glutamine in plasma of patients of a group to which a sodium bicarbonate preparation has been administered, after 6, 12, and 24 weeks from the start of the test.

FIG. 20 is a graph showing a correlation between a concentration of phenylacetyl-L-glutamine in urine and a concentration of phenylacetyl-L-glutamine in

plasma of all the patients after 6, 12, and 24 weeks from the start of the test.

EMBODIMENTS FOR CARRYING OUT THE INVENTION

[0023]

5 1. Pharmaceutical composition

A pharmaceutical composition provided by the present invention can comprise an alkalinizing agent as an active ingredient.

The alkalinizing agent is an agent having the ability to increase the HCO_3^- concentration and pH of body fluids of mammals (particularly a human), such as blood or
10 urine. Examples of alkalinizing agents include a pharmaceutically acceptable salt of citric acid or a hydrate thereof, or a mixture thereof, and sodium bicarbonate (baking soda). Examples of a pharmaceutically acceptable salt of citric acid include an alkali metal salt of citric acid. Examples of an alkali metal salt of citric acid include potassium citrate and sodium citrate. Potassium citrate may be a hydrate such as a
15 stable potassium citrate monohydrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$), and sodium citrate may be a hydrate such as sodium citrate dihydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$).

Examples of preferable alkalinizing agents include sodium citrate, potassium citrate, or a hydrate thereof, or a mixture thereof. For example, an alkalinizing agent may be a mixture of a potassium citrate monohydrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and sodium
20 citrate dihydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$). A mixing ratio of a potassium citrate monohydrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and a sodium citrate dihydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$) can be appropriately set by those skilled in the art. For example, a molar ratio of a potassium citrate monohydrate to a sodium citrate dihydrate can be set to 1 (for a potassium citrate monohydrate) to 0.01 to 100 (for a sodium citrate dihydrate). The

mixing ratio may be about 1:1 as a molar ratio.

In addition, additional examples of preferable alkalinizing agents include sodium citrate or a hydrate thereof, for example, a sodium citrate dihydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$).

5 Furthermore, additional examples of preferable alkalinizing agents include potassium citrate or a hydrate thereof, for example, a potassium citrate monohydrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$).

In one embodiment, the alkalinizing agent contained in the pharmaceutical composition of the present invention may comprise a mixture of sodium citrate or a
10 hydrate thereof and potassium citrate or a hydrate thereof.

In another embodiment, the alkalinizing agent contained in the pharmaceutical composition of the present invention may be composed of only a mixture of sodium citrate or a hydrate thereof and potassium citrate or a hydrate thereof.

In the present specification, when referring to the weight of an alkalinizing agent
15 (for example, a potassium citrate monohydrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and a sodium citrate dihydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$)), this weight may be a dry weight.

[0024]

In the present specification, a uremic substance is a substance (waste products, toxins, and the like) excreted by normal kidneys, and it means a substance that increases
20 (accumulates) in the blood and causes uremic symptoms or diseases when the excretory function deteriorates due to some cause such as deteriorated kidney function. Examples of uremic substance include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.

Among them, indoxyl sulfate is produced by oxidation and sulfate conjugation

of indole produced by enteric bacteria from tryptophan derived from dietary protein. Most of the indoxyl sulfate is present by being bound to albumin in the blood and is not metabolized. In healthy subjects, it is excreted from the kidneys into the urine, but in a case of a patient with kidney disease, it remains accumulated at high concentrations in the blood.

Indoxyl sulfate, which is a uremic substance, not only causes uremia in a patient with kidney disease, but also causes a patient with chronic kidney disease to be subjected to introduction of dialysis.

Accordingly, by reducing a concentration of indoxyl sulfate in blood, the uremic symptoms of a patient with kidney disease are ameliorated, and thereby treatment and/or prevention of uremia becomes possible. In addition, it is possible to delay the introduction of dialysis in a patient with chronic kidney disease by decreasing a concentration of indoxyl sulfate in blood. In one embodiment, a patient with chronic kidney disease has progressive chronic kidney disease.

In the present specification, the expression [A, B, and/or C] represents “at least one selected from the group consisting of A, B, and C.” Accordingly, for example, “indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid” means at least one selected from the group consisting of “indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.”

[0025]

In addition, indoxyl sulfate, which is a uremic substance, causes myocardial fibrosis, arteriosclerosis, vascular smooth muscle cell proliferation, vascular endothelial cell injury, thickening of the arterial wall, calcification of the aorta, and the like, and allows cardiovascular disease (for example, heart failure, myocardial infarction), which is

one of the complications in a patient with chronic kidney disease, and/or a stroke, which is a cerebrovascular disease, to develop.

Accordingly, decreasing a concentration of indoxyl sulfate in blood suppresses myocardial fibrosis, arteriosclerosis, vascular smooth muscle cell proliferation, vascular
5 endothelial cell injury, thickening of the arterial wall, calcification of the aorta, and the like, and thereby it becomes possible to treat and/or prevent cardiovascular disease, which is one of the complications in a patient with chronic kidney disease, and/or cerebrovascular disease.

[0026]

10 In one aspect, in the pharmaceutical composition provided by the present invention, it is possible to decrease a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof) in the blood. Examples of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid,
15 and a combination thereof include indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and phenylacetyl-L-glutamine; hippuric acid and
20 phenylacetyl-L-glutamine; indoxyl sulfate; p-cresyl sulfate; hippuric acid; and phenylacetyl-L-glutamine.

In the present specification, “decreasing a concentration of uremic substance in the blood” means that the concentration of uremic substance in the blood after administration of the pharmaceutical composition provided by the present invention
25 decreases compared to the concentration of uremic substance in the blood before the

administration, or means that the concentration of uremic substance in the blood is decreased by administration of the pharmaceutical composition provided by the present invention, compared to placebo administration.

In one embodiment, administration of the pharmaceutical composition provided by the present invention decreases a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate; p-cresyl sulfate; hippuric acid; or phenylacetyl-L-glutamine)) in the blood compared to before administration, but an amount of decrease thereof is 1% to 5%, 3% to 5%, 1% to 10%, 3% to 10%, 5% to 10%, 1% to 15%, 3% to 15%, 5% to 15%, 1% to 30%, 1% to 40%, 3% to 40%, 5% to 40%, 1% to 50%, 3% to 50%, 5% to 50%, 30% to 50%, 1% to 60%, 5% or more, 10% or more, or 30% or more of a concentration of uremic substance in the blood before administration.

In one embodiment, an amount of decrease in concentration of uremic substance in the blood is calculated by Calculation Equation (1).

Amount of decrease in concentration of uremic substance in blood (%) =

$$\left[\frac{(\text{concentration (ng/mL) of uremic substance in blood before administration of pharmaceutical composition} - \text{concentration (ng/mL) of uremic substance in blood after administration of pharmaceutical composition})}{\text{concentration (ng/mL) of uremic substance in blood before administration of pharmaceutical composition}} \right] \times 100$$

(1)

In one embodiment, continuous administration of the pharmaceutical composition provided by the present invention for 6, 12, or 24 weeks decreases a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate; p-cresyl sulfate; hippuric acid; or phenylacetyl-L-glutamine)) in the blood compared to before administration, but an amount of decrease thereof is 1% to 5%, 3% to 5%, 1% to 10%, 3% to 10%, 5% to 10%, 1% to 15%, 3% to 15%, 5% to 15%, 1% to 30%, 1% to 40%, 3% to 40%, 5% to 40%, 1% to 50%, 3% to 50%, 5% to 50%, 30% to 50%, 1% to 60%, 5% or more, 10% or more, or 30% or more of a concentration of uremic substance in the blood before administration.

In one aspect, in the pharmaceutical composition provided by the present invention, it is possible to promote excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof) into urine. Examples of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-

glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid; p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid; phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate; hippuric acid; argininosuccinic acid; and phenylacetyl-L-glutamine.

In the present specification, “promoting excretion of uremic substance into the urine” means that a concentration of uremic substance in the urine after administration increases compared to the concentration of uremic substance in the urine before administration of the pharmaceutical composition provided by the present invention; means that a concentration of uremic substance in the urine increases by administration of the pharmaceutical composition provided by the present invention compared to placebo administration; means that an amount of uremic substance in the urine after administration increases compared to the amount of uremic substance in urine before administration of the pharmaceutical composition provided by the present invention; or means that the amount of uremic substance in the urine increases by administration of the pharmaceutical composition provided by the present invention compared to placebo administration.

In the present specification, “in urine” means, for example, “in early morning urine.”

In one embodiment, by administration of the pharmaceutical composition provided by the present invention, a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid,

argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid; p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid; phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate; hippuric acid; argininosuccinic acid; or phenylacetyl-L-glutamine)) in urine increases by 1% to 100%, 1% to 50%, 3% to 50%, 5% to 50%, 10% to 50%, 15% to 50%, 1% to 40%, 5% to 40%, 10% to 40%, 1% to 30%, 5% to 30%, 10% to 30%, 15% to 30%, 10% or more, 20% or more, 30% or more, or 40% or more, compared to before administration.

In one embodiment, the amount of increase in concentration of uremic substance in urine is calculated by Calculation Equation (2).

$$\begin{aligned} &\text{Amount of increase in concentration of uremic substance in urine (\%)} = \\ &[(\text{concentration (ng/mL) of uremic substance in urine after administration of} \\ &\text{pharmaceutical composition} - \text{concentration (ng/mL) of uremic substance in urine before} \\ &\text{administration of pharmaceutical composition}) / \text{concentration (ng/mL) of uremic} \\ &\text{substance in urine before administration of pharmaceutical composition}] \times 100 \\ &(2) \end{aligned}$$

In one embodiment, by continuous administration of the pharmaceutical

composition provided by the present invention for 6, 12, or 24 weeks, a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and

5 argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid; indoxyl sulfate

10 and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid; p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid; phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate; hippuric acid;

15 argininosuccinic acid; or phenylacetyl-L-glutamine)) in urine increases by 1% to 100%, 1% to 50%, 3% to 50%, 5% to 50%, 10% to 50%, 15% to 50%, 1% to 40%, 5% to 40%, 10% to 40%, 1% to 30%, 5% to 30%, 10% to 30%, 15% to 30%, 10% or more, 20% or more, 30% or more, or 40% or more, compared to before administration.

In one aspect, in the pharmaceutical composition provided by the present

20 invention, it is possible to promote excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof) from blood into urine, thereby promoting excretion of the uremic substance outside the body. Examples of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination

25 thereof include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid,

and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine,
 and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and
 phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-
 glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl
 5 sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid;
 indoxyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-
 glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-
 glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid;
 p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid;
 10 phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate;
 hippuric acid; argininosuccinic acid; and phenylacetyl-L-glutamine.

In one embodiment, by administration of the pharmaceutical composition
 provided by the present invention, a ratio of a concentration of uremic substance in urine
 to a concentration of uremic substance in blood after administration increases compared
 15 to a ratio of a concentration of uremic substance in urine to a concentration of uremic
 substance in blood before administration of the pharmaceutical composition provided by
 the present invention. In one embodiment, administration of the pharmaceutical
 composition provided by the present invention increases the ratio of a concentration of
 uremic substance in urine to a concentration of uremic substance in blood, compared to
 20 placebo administration.

In one embodiment, by administration of the pharmaceutical composition
 provided by the present invention, a ratio of a concentration of uremic substance (for
 example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid,
 argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl
 25 sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl

sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid; p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid; phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate; hippuric acid; argininosuccinic acid; or phenylacetyl-L-glutamine)) in urine with respect to a concentration of uremic substance in blood (a concentration (ng/mL) in urine/a concentration (ng/mL) in blood) increases by 1% to 100%, 1% to 50%, 3% to 50%, 5% to 50%, 10% to 50%, 15% to 50%, 1% to 40%, 5% to 40%, 10% to 40%, 1% to 30%, 5% to 30%, 10% to 30%, 15% to 30%, 10% or more, 20% or more, 30% or more, or 40% or more, compared to before administration.

In one embodiment, an increase in ratio of a concentration of uremic substance in urine to a concentration of uremic substance in blood (a concentration (ng/mL) in urine/a concentration (ng/mL) in blood) is calculated by Calculation Equation (3).

Amount of increase in ratio of a concentration of uremic substance in urine to a concentration of uremic substance in blood (%) = [(ratio of concentration of uremic substance in urine to concentration of uremic substance in blood after administration of pharmaceutical composition – ratio of concentration of uremic substance in urine to concentration of uremic substance in blood before administration of pharmaceutical composition)/ratio of concentration of uremic substance in urine to concentration of uremic substance in blood before administration of pharmaceutical composition] × 100

(3)

In one embodiment, by continuous administration of the pharmaceutical composition provided by the present invention for 6, 12, or 24 weeks, a ratio of a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate, phenylacetyl-L-glutamine, and argininosuccinic acid; indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine; indoxyl sulfate and hippuric acid; indoxyl sulfate and phenylacetyl-L-glutamine; hippuric acid and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; p-cresyl sulfate and hippuric acid; indoxyl sulfate and argininosuccinic acid; p-cresyl sulfate and argininosuccinic acid; hippuric acid and argininosuccinic acid; phenylacetyl-L-glutamine and argininosuccinic acid; indoxyl sulfate; p-cresyl sulfate; hippuric acid; argininosuccinic acid; or phenylacetyl-L-glutamine)) in urine with respect to a concentration of uremic substance in blood (a concentration (ng/mL) in urine/a concentration (ng/mL) in blood) increases by 1% to 100%, 1% to 50%, 3% to 50%, 5% to 50%, 10% to 50%, 15% to 50%, 1% to 40%, 5% to 40%, 10% to 40%, 1% to 30%, 5% to 30%, 10% to 30%, 15% to 30%, 10% or more, 20% or more, 30% or more, or 40% or more, compared to before administration.

In one embodiment, the administration of the pharmaceutical composition provided by the present invention promotes excretion of the uremic substance outside the body depending on a concentration of uremic substance in blood.

In one embodiment, the administration of the pharmaceutical composition provided by the present invention promotes excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof) into urine depending on a concentration of uremic substance in blood. For example, in a case where a concentration of uremic substance in blood is high, the amount of uremic substance excreted into the urine becomes a high value in accordance with the high concentration of uremic substance in blood. In a case where the concentration of uremic substance in blood is low, the amount of uremic substance excreted into the urine becomes a low value. Such excretion of uremic substance into urine depending on a concentration of uremic substance in blood suggests that the pharmaceutical composition provided by the present invention has a low risk of side effects and excellent safety. Examples of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; indoxyl sulfate; p-cresyl sulfate; and phenylacetyl-L-glutamine.

In one embodiment, the administration of the pharmaceutical composition provided by the present invention promotes excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination thereof) into urine depending on a concentration of uremic substance in blood, and thereby a concentration of uremic substance in blood is decreased. Examples of indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, argininosuccinic acid, and a combination

thereof include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid; indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine; indoxyl sulfate and phenylacetyl-L-glutamine; indoxyl sulfate and p-cresyl sulfate; p-cresyl sulfate and phenylacetyl-L-glutamine; indoxyl sulfate; p-cresyl sulfate; and phenylacetyl-L-glutamine.

In one embodiment, administration of the pharmaceutical composition provided by the present invention excretes indoxyl sulfate into the urine depending on a concentration of indoxyl sulfate in blood, and as a result, a ratio of a concentration of indoxyl sulfate in urine to a concentration of indoxyl sulfate in blood (a concentration in urine/a concentration in blood) can become 1 to 1000, preferably 1 to 200, more preferably 1 to 100, and even more preferably 10 to 100. In this embodiment, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of indoxyl sulfate in blood is 0.01 to 100 $\mu\text{g/mL}$ (for example, 0.1 to 30 $\mu\text{g/mL}$). Furthermore, as a result of the administration, the concentration of indoxyl sulfate in blood may become 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.03 to 10 $\mu\text{g/mL}$).

In one embodiment, administration of the pharmaceutical composition provided by the present invention excretes p-cresyl sulfate into the urine depending on a concentration of p-cresyl sulfate in blood, and as a result, a ratio of a concentration of p-cresyl sulfate in urine to a concentration of p-cresyl sulfate in blood (a concentration in urine/a concentration in blood) can become 0.1 to 1000, preferably 1 to 300, more preferably 1 to 150, and even more preferably 1 to 100. In this embodiment, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of p-cresyl sulfate in blood is 0.003 to 300 $\mu\text{g/mL}$ (for example, 0.01 to 30 $\mu\text{g/mL}$).

Furthermore, as a result of the administration, the concentration of p-cresyl sulfate in blood may become 0.001 to 100 $\mu\text{g/mL}$ (for example, 0.001 to 30 $\mu\text{g/mL}$).

In one embodiment, administration of the pharmaceutical composition provided by the present invention excretes phenylacetyl-L-glutamine into the urine depending on a concentration of phenylacetyl-L-glutamine in blood, and as a result, a ratio of a concentration of phenylacetyl-L-glutamine in urine to a concentration of phenylacetyl-L-glutamine in blood (a concentration in urine/a concentration in blood) can become 1 to 1500, preferably 1 to 1000, more preferably 1 to 800, and even more preferably 10 to 600. In this embodiment, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of phenylacetyl-L-glutamine in blood is 0.03 to 30 $\mu\text{g/mL}$ (for example, 0.1 to 10 $\mu\text{g/mL}$). Furthermore, as a result of the administration, the concentration of phenylacetyl-L-glutamine in blood may become 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.03 to 10 $\mu\text{g/mL}$).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a plurality of humans (for example, patients with chronic kidney disease), and a ratio of a concentration of indoxyl sulfate in urine to a concentration of indoxyl sulfate in blood in each individual (a concentration in urine/a concentration in blood) shows high correlation. In this embodiment, high correlation may be indicated by a Pearson test in which an r value is 0.4 to 1, 0.5 to 1, 0.6 to 1, or 0.7 to 1 (preferably 0.7 to 1). In addition, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of indoxyl sulfate in blood is 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.1 to 10 $\mu\text{g/mL}$), and as a result of the administration, a concentration of indoxyl sulfate in blood may become 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.1 to 10 $\mu\text{g/mL}$).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a plurality of humans (for example, patients with chronic kidney disease), and a ratio of a concentration of p-cresyl sulfate in urine to a concentration of p-cresyl sulfate in blood in each individual (a concentration in urine/a concentration in blood) shows high correlation. In this embodiment, high correlation may be indicated by a Pearson test in which an r value is 0.4 to 1, 0.5 to 1, 0.6 to 1, or 0.7 to 1 (preferably 0.7 to 1). In addition, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of p-cresyl sulfate in blood is 0.001 to 100 $\mu\text{g/mL}$ (for example, 0.01 to 50 $\mu\text{g/mL}$), and as a result of the administration, a concentration of p-cresyl sulfate in blood may become 0.001 to 100 $\mu\text{g/mL}$ (for example, 0.01 to 50 $\mu\text{g/mL}$).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a plurality of humans (for example, patients with chronic kidney disease), and a ratio of a concentration of phenylacetyl-L-glutamine in urine to a concentration of phenylacetyl-L-glutamine in blood in each individual (a concentration in urine/a concentration in blood) shows high correlation. In this embodiment, high correlation may be indicated by a Pearson test in which an r value is 0.4 to 1, 0.5 to 1, 0.6 to 1, or 0.7 to 1 (preferably 0.7 to 1). In addition, the pharmaceutical composition provided by the present invention may be administered to a human (for example, a patient with chronic kidney disease) in which a concentration of phenylacetyl-L-glutamine in blood is 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.05 to 10 $\mu\text{g/mL}$), and as a result of the administration, a concentration of phenylacetyl-L-glutamine in blood may become 0.01 to 10 $\mu\text{g/mL}$ (for example, 0.05 to 10 $\mu\text{g/mL}$).

Because of the characteristics of the pharmaceutical composition provided by

the present invention, the pharmaceutical composition provided by the present invention, in one aspect, not only can be used as a pharmaceutical composition for decreasing a concentration of uremic substance in blood and/or a pharmaceutical composition for promoting excretion of uremic substance into urine, but also can be used as any of a
5 pharmaceutical composition for ameliorating uremic symptoms in a patient with kidney disease, a pharmaceutical composition for treating and/or preventing uremic symptoms in a patient with kidney disease, a pharmaceutical composition for suppressing progression of chronic kidney disease, and a pharmaceutical composition for delaying introduction of dialysis in a patient with chronic kidney disease.

10 In the present specification, the term “amelioration” is a concept encompassing bringing “pathological” or “abnormal” symptoms, conditions, or diseases closer to “healthy” or “normal” conditions, or action therefor; and making “pathological” or “abnormal” symptoms, conditions, or diseases “healthy” or “normal” conditions, or action therefor. Accordingly, in one embodiment, the term “amelioration” encompasses
15 a concept in which a numerical value indicative of “pathological” or “abnormal” symptoms or conditions decreases or increases in accordance with the “amelioration,” and becomes closer to a normal value or becomes a normal value. In another embodiment, the term “amelioration” encompasses a concept in which, in accordance with the “amelioration,” a concentration of uremic substance in blood is decreased, a
20 concentration of uremic substance in urine increases, and a concentration of uremic substance in urine may turn to a decrease when a concentration of uremic substance in blood becomes sufficiently small.

In the present specification, the term “healthy” represents a state in which there is no acute or chronic disease or disorder, and the term “normal” represents in a state that
25 a healthy subject usually expresses.

In the present specification, the term “treatment” encompasses a concept in which “pathological” or “abnormal” symptoms, conditions, or diseases are eliminated, completely cured, healed, or in remission, and action therefor; a deterioration in “pathological” or “abnormal” symptoms, conditions, or diseases is suppressed, and action therefor; and the “amelioration.” The term suppression has the meaning to be described later. In one embodiment, the term “treatment” means that “pathological” or “abnormal” symptoms, conditions, or diseases are eliminated, completely cured, healed, or in remission, and means action therefor. In another embodiment, the term “treatment” means that “pathological” or “abnormal” symptoms, conditions, or diseases are eliminated, completely cured, healed, or in remission.

In the present specification, the term “prevention” encompasses a concept in which onset of “pathological” or “abnormal” symptoms, conditions, or diseases is prevented before they occur, and action therefor.

In the present specification, the term “delay” encompasses a concept in which a time for an object phenomenon to occur is extended, and action therefor; and a time is extended so that the object phenomenon does not occur.

In the present specification, the term “suppression” encompasses a concept in which a deterioration or progression of symptoms, conditions, or diseases is stopped or slowed down, or action therefor; and the symptoms, conditions, or the diseases are ameliorated, or action therefor. The term amelioration has the meaning as described above. The above-mentioned “deterioration or progression of symptoms, conditions, or diseases” includes a deterioration or progression of “pathological” or “abnormal” symptoms, conditions, or diseases; and a deterioration or progression of “pathological” or “abnormal” symptoms, conditions, or diseases from “healthy” or “normal” conditions. In one embodiment, the term “suppression” means that a deterioration or progression of

symptoms, conditions, or diseases is stopped or slowed down, or means action therefor.

In another embodiment, the term “suppression” means that a deterioration or progression of symptoms, conditions, or diseases is stopped or slowed down.

The symptoms, conditions, or diseases are compared before and after
5 administration of the pharmaceutical composition provided by the present invention.

In addition, because of the characteristics of the pharmaceutical composition provided by the present invention as described above, in one aspect, the pharmaceutical composition provided by the present invention can be used as any of a pharmaceutical composition for suppressing myocardial fibrosis in a patient with kidney disease, a
10 pharmaceutical composition for suppressing arteriosclerosis in a patient with kidney disease, a pharmaceutical composition for suppressing vascular smooth muscle cell proliferation in a patient with kidney disease, a pharmaceutical composition for suppressing vascular endothelial cell injury in a patient with kidney disease, a pharmaceutical composition for suppressing thickening of the arterial wall of a patient
15 with kidney disease, a pharmaceutical composition for suppressing calcification of the aorta of a patient with kidney disease, and a pharmaceutical composition for treating and/or preventing cardiovascular disease associated with chronic kidney disease.

[0027]

In addition, it has been reported that, when a drug that decreases a concentration
20 of indoxyl sulfate in blood was administered to a patient with non-diabetic chronic kidney disease, the pulse wave velocity and carotid intima-media complex thickness, which are indicators of arteriosclerosis, were significantly improved compared to those before administration (Nakamura T., et al.: Oral ADSORBENT AST-120 decreases carotid intima-media thickness and arterial stiffness in patients with chronic renal failure.
25 Kidney Blood Press Res, 27:121-6, 2004.). Accordingly, in one aspect, the

pharmaceutical composition provided by the present invention which decreases the concentration of indoxyl sulfate in blood can be used as a pharmaceutical composition for ameliorating arteriosclerosis or a pharmaceutical composition for ameliorating thickening of the arterial wall (for example, carotid artery) of a patient with kidney
5 disease (preferably a patient with chronic kidney disease, more preferably a patient with non-diabetic chronic kidney disease).

[0028]

In addition, drugs that decrease the concentration of indoxyl sulfate in blood have been reported to suppress cisplatin-induced acute kidney injury (Morisaki T., et.
10 Al.,: Regulation of renal organic ion transporters in cisplatin-induced acute kidney injury and uremia in rats. Pharm. Res., 25 (11): 2526-33, 2008). Accordingly, in one aspect, the pharmaceutical composition provided by the present invention which decreases the concentration of indoxyl sulfate in blood can be used as a pharmaceutical composition for treating acute kidney failure, or a pharmaceutical composition for suppressing
15 progression from acute kidney failure to chronic kidney disease.

[0029]

In addition, p-cresyl sulfate, which is a uremic substance, has been reported to be a causative substance of vascular endothelial damage (Meijers B.K., et. Al.,: The uremic retention solute p-cresyl sulfate and markers of endothelial damage., Am. J.
20 Kidney Dis., 54: 891-901, 2009).

In one aspect, the pharmaceutical composition provided by the present invention which promotes excretion of p-cresyl sulfate into urine can be used as a pharmaceutical composition for suppressing vascular endothelial damage in a patient with kidney disease (preferably a patient with chronic kidney disease).

25 In addition, phenylacetyl-L-glutamine, which is a uremic substance, has been

reported to increase the risk of developing cardiovascular disease in a patient with chronic kidney disease.

In one aspect, the pharmaceutical composition provided by the present invention which promotes excretion of phenylacetyl-L-glutamine into urine can be used as a pharmaceutical composition for treating and/or preventing cardiovascular disease in a patient with chronic kidney disease.

In one embodiment, since the pharmaceutical composition provided by the present invention promotes excretion of uremic substance into urine, such as indoxyl sulfate, p-cresyl sulfate, hippuric acid, argininosuccinic acid, and phenylacetyl-L-glutamine, the pharmaceutical composition provided by the present invention can be used as a pharmaceutical composition for promoting excretion of indoxyl sulfate, p-cresyl sulfate, hippuric acid, argininosuccinic acid, and/or phenylacetyl-L-glutamine into urine of a patient with kidney disease (preferably a patient with chronic kidney disease).

In one embodiment, in order to decrease indoxyl sulfate concentration in the blood, p-cresyl sulfate concentration in blood, hippuric acid concentration in blood, and/or phenylacetyl-L-glutamine concentration in blood, a pharmaceutically acceptable salt of citric acid, or a hydrate thereof, or a mixture thereof (for example, a mixture of potassium citrate monohydrate and sodium citrate dihydrate) is administered to a patient with kidney disease (preferably a patient with chronic kidney disease).

In one embodiment, in order to increase indoxyl sulfate concentration in urine, p-cresyl sulfate concentration in urine, urinary hippuric acid concentration, urinary argininosuccinic acid concentration, and/or urinary phenylacetyl-L-glutamine concentration (preferably to increase indoxyl sulfate concentration in urine, p-cresyl sulfate concentration in urine, and phenylacetyl-L-glutamine concentration in urine), a pharmaceutically acceptable salt of citric acid, or a hydrate thereof, or a mixture thereof

(for example, a mixture of potassium citrate monohydrate and sodium citrate dihydrate) is administered to a patient with kidney disease (preferably a patient with chronic kidney disease).

In one embodiment, sodium bicarbonate is administered to a patient with kidney
5 disease (preferably a patient with chronic kidney diseases) to decrease blood p-cresyl sulfate concentration and/or blood phenylacetyl-L-glutamine concentration.

In one embodiment, sodium bicarbonate is administered to a patient with kidney disease (preferably a patient with chronic kidney disease) to increase the urinary argininosuccinic acid concentration.

10 [0030]

In one aspect, the pharmaceutical composition provided by the present invention can be used as a pharmaceutical composition for treating kidney tubular damage, a pharmaceutical composition for preventing kidney tubular damage, or a pharmaceutical composition for suppressing kidney tubular damage. A kidney tubule may be, for
15 example, a kidney proximal tubule.

In another aspect, the pharmaceutical composition provided by the present invention can be used as a pharmaceutical composition for maintaining kidney function.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition for suppressing kidney tubular cell damage, a
20 pharmaceutical composition for protecting kidney tubular cells, or a pharmaceutical composition for maintaining kidney tubular cell function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like).

In one embodiment, the pharmaceutical composition provided by the present
25 invention is a pharmaceutical composition for suppressing kidney proximal tubular cell

damage, a pharmaceutical composition for protecting kidney proximal tubular cells, or a pharmaceutical composition for maintaining kidney proximal tubular cell function (for example, reabsorption of glucose, amino acid, vitamin, and the like).

In one embodiment, the pharmaceutical composition provided by the present invention suppresses an increase in the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) associated with progression of a stage of chronic kidney disease.

In one embodiment, the pharmaceutical composition provided by the present invention does not affect the glomerular function of a patient with chronic kidney disease, whereas it suppresses kidney proximal tubular cell damage associated with progression of a stage of chronic kidney disease to protect kidney proximal tubular cells.

In the present specification, “maintenance of kidney function” means, for example, suppression of kidney tubular damage, protection of kidney tubular cells, or maintenance of kidney tubular function. A kidney tubule may be, for example, a kidney proximal tubule, and one aspect of maintenance of kidney tubular function or kidney proximal tubular function is to maintain kidney tubular cell function or to maintain kidney proximal tubular cell function.

In the present specification, “protection of cells” means maintenance or preservation of cell states or suppression of cell damage. The term suppression has the above-mentioned meaning.

In the present specification, “maintenance of cell function” means preservation of cell function or suppression of deterioration of cell function. The term suppression has the above-mentioned meaning.

The state or function of cells is compared before and after administration of the pharmaceutical composition provided by the present invention.

In the present specification, “early morning urine” represents the first urine after getting up.

In one embodiment, the pharmaceutical composition provided by the present invention suppresses an increase in the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) 6 weeks, 12 weeks, and/or 24 weeks after administration, as compared with that before the start of administration.

In one embodiment, the pharmaceutical composition provided by the present invention suppresses an increase in the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) associated with progression of a stage of chronic kidney disease, but an amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) is not substantially decreased compared to that before the start of administration of the pharmaceutical composition provided by the present invention. In this embodiment, for example, in a case where the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) before the start of administration of the pharmaceutical composition provided by the present invention is 1, the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) after administration may be 0.7 to 1.0 or 1.0 or more, 0.8 to 1.0 or 1.0 or more, 0.85 to 1.0 or 1.0 or more, 0.9 to 1.0 or 1.0 or more 0.7 to 2.0, 0.8 to 2.0, 0.85 to 2.0, 0.9 to 2.0, 0.7 to 1.6, 0.8 to 1.6, 0.85 to 1.6, or 0.9 to 1.6.

In one embodiment, the pharmaceutical composition provided by the present invention suppresses an increase in the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) associated with progression of a stage of chronic kidney disease, but an amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) is not substantially decreased 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration,

compared to that before the start of administration of the pharmaceutical composition provided by the present invention. In this embodiment, for example, in a case where the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) before the start of administration of the pharmaceutical composition provided by the present invention is 1, the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine) 6 weeks after administration, 12 weeks after administration, or 24 weeks after administration may be 0.7 to 1.0 or 1.0 or more, 0.8 to 1.0 or 1.0 or more, 0.85 to 1.0 or 1.0 or more, 0.9 to 1.0 or 1.0 or more 0.7 to 2.0, 0.8 to 2.0, 0.85 to 2.0, 0.9 to 2.0, 0.7 to 1.6, 0.8 to 1.6, 0.85 to 1.6, or 0.9 to 1.6.

In one embodiment, the pharmaceutical composition provided by the present invention does not substantially increase the amount (concentration) of cystatin C in the blood (for example, plasma) compared to before the start of administration. In this embodiment, for example, in a case where the amount (concentration) of cystatin C in blood (for example, in plasma) before the start of administration of the pharmaceutical composition provided by the present invention is 1, the amount (concentration) of cystatin C in blood (for example, in plasma) after administration may be 1.0 or less or 1.0 to 1.2, 1.0 or less or 1.0 to 1.15, 1.0 or less or 1.0 to 1.1, 1.0 or less or 1.0 to 1.05, 0.9 to 1.2, 0.9 to 1.15, 0.9 to 1.1, 0.9 to 1.05, 0.95 to 1.2, 0.95 to 1.15, 0.95 to 1.1, or 0.95 to 1.05.

In one embodiment, the pharmaceutical composition provided by the present invention does not substantially increase the amount (concentration) of cystatin C in the blood (for example, plasma) 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration, compared to before the start of administration. In this embodiment, for example, in a case where the amount (concentration) of cystatin C in blood (for example, in plasma) before the start of administration of the pharmaceutical

composition provided by the present invention is 1, the amount (concentration) of cystatin C in blood (for example, in plasma) 6 weeks after administration, 12 weeks after administration, or 24 weeks after administration may be 1.0 or less or 1.0 to 1.2, 1.0 or less or 1.0 to 1.15, 1.0 or less or 1.0 to 1.1, 1.0 or less or 1.0 to 1.05, 0.9 to 1.2, 0.9 to 1.15, 0.9 to 1.1, 0.9 to 1.05, 0.95 to 1.2, 0.95 to 1.15, 0.95 to 1.1, or 0.95 to 1.05.

In one embodiment, by administration of the pharmaceutical composition provided by the present invention, despite no amelioration in kidney proximal tubular damage and/or glomerular damage is recognized 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration as compared to before the start of administration, a decrease in the blood concentration of the uremic substance, promotion of urinary excretion of the uremic substance, and/or promotion of excretion of the uremic substance outside of the body is observed 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration, as compared to before the start of administration.

In one embodiment, by administration of the pharmaceutical composition provided by the present invention, despite no amelioration in kidney proximal tubular damage and/or glomerular damage is recognized 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration as compared to before the start of administration, a decrease in the blood concentration of the uremic substance, promotion of urinary excretion of the uremic substance, and/or promotion of excretion of the uremic substance outside the body is observed 6 weeks after administration, 12 weeks after administration, and/or 24 weeks after administration, as compared to before the start of administration.

[0031]

The pharmaceutical composition provided by the present invention is orally or

parenterally administered to a human or other mammals. Examples of parenteral administration include intravenous administration, subcutaneous administration, intramuscular administration, intra-articular administration, transmucosal administration, transdermal administration, nasal administration, rectal administration, intrathecal administration, intraperitoneal administration, and local administration

The pharmaceutical composition provided by the present invention may be prepared as an alkalinizing agent as it is or may be prepared by mixing an alkalinizing agent with a pharmaceutically acceptable carrier such as excipients (for example, lactose, D-mannitol, crystalline cellulose, and glucose), binders (for example, hydroxypropylcellulose (HPC), gelatin, and polyvinylpyrrolidone (PVP)), lubricants (for example, magnesium stearate, and talc), disintegrants (for example, starch and carboxymethylcellulose calcium (CMC-Ca)), diluents (for example, water for injection and saline), and other additives if necessary (for example, pH adjusters, surfactants, solubilizers, preservatives, emulsifiers, tonicity agents, and stabilizers), or may be preparation such as a tablet, capsule, suspension, injection, or suppository. For example, in the case of a tablet, the alkalinizing agent may be mixed with excipients (for example, lactose, D-mannitol, crystalline cellulose, and glucose), disintegrants (for example, starch and carboxymethylcellulose calcium (CMC-Ca)), binders (for example, hydroxypropylcellulose (HPC), gelatin, and polyvinylpyrrolidone (PVP)), lubricants (for example, magnesium stearate and talc), and the like to be formulated.

The tablet according to the present invention will be described in more detail below.

[0032]

In one embodiment, the pharmaceutical composition provided by the present invention is a tablet. The tablet provided by the present invention may comprise

pharmaceutically acceptable additives customary in the pharmaceutical field, in addition to an alkalinizing agent (for example, potassium citrate or a hydrate thereof; sodium citrate or a hydrate thereof; a mixture of potassium citrate monohydrate and sodium citrate dihydrate; or sodium bicarbonate). Examples of such additives include

5 excipients, binders, disintegrants, fluidizers, flavoring agents, lubricants, pH adjusters, surfactants, stabilizers, and fragrances.

The content of the alkalinizing agent in the tablet provided by the present invention may be 10% to 95% by weight, preferably 30% to 90% by weight, and more preferably 60% to 85% by weight with respect of the tablet.

10 [0033]

Examples of excipients that can be used in the tablets provided by the present invention include lactose (for example, lactose hydrate, and anhydrous lactose), sugars such as glucose, sucrose, fructose, and maltose, sugar alcohols such as erythritol, sorbitol, maltitol, xylitol, and D-mannitol, starch (for example, corn starch, potato starch,

15 rice starch, and wheat starch), crystalline cellulose, magnesium aluminate metasilicate, anhydrous calcium phosphate, precipitated calcium carbonate, calcium silicate, calcium lactate, and ethylcellulose, among which crystalline cellulose is particularly preferable.

The content of the excipient in the tablet provided by the present invention may be 1% to 95% by weight, preferably 1% to 80% by weight, more preferably 3% to 80%

20 by weight, and even more preferably 3% to 20% by weight with respect to the tablet.

[0034]

Examples of binders that can be used in the tablets provided by the present invention include hydroxypropylcellulose, hydroxypropyl methylcellulose, polyvinylpyrrolidone, dextrin, methylcellulose, polyvinyl alcohol, sodium alginate,

25 aminoalkyl methacrylate copolymer, polyethylene glycol, pregelatinized starch (for

example, partially pregelatinized starch), agar, and gelatin, among which hydroxypropylcellulose is particularly preferable.

The content of the binder in the tablet provided by the present invention may be 0.1% to 30% by weight, preferably 0.1% to 10% by weight, and more preferably 0.3% to 3% by weight with respect to the tablet.

[0035]

Examples of disintegrants that may be used in tablets provided by the present invention include croscarmellose sodium, carmellose calcium, carboxymethyl starch sodium, low substituted hydroxypropylcellulose, crospovidone, starch (for example, wheat starch, corn starch, and partially pregelatinized starch), and carmellose, among which partially pregelatinized starch is particularly preferable.

The content of the disintegrant in the tablet provided by the present invention may be 0.3% to 20% by weight, preferably 1% to 10% by weight, and more preferably 3% to 10% by weight with respect to the tablet.

[0036]

Examples of fluidizers that can be used in the tablets provided by the present invention include light anhydrous silicic acid, talc, and magnesium aluminate metasilicate.

The content of the fluidizers in the tablet provided by the present invention may be 0.03% to 3% by weight, preferably 0.1% to 3% by weight, and more preferably 0.3% to 3% by weight with respect to the tablet.

[0037]

Examples of flavoring agents that can be used in the tablets provided by the present invention include citric acid (for example, anhydrous citric acid), acidulants such as malic acid, acetic acid, tartaric acid, fumaric acid, and ascorbic acid (where, the said

flavoring agent does not comprise the alkalizing agent according to the present invention), saccharin sodium, dipotassium glycyrrhizinate, aspartame (registered trademark), and sweeteners such as stevia, thaumatin, and sucralose.

The content of the flavoring agent in the tablet provided by the present invention may be 0.03% to 3% by weight, preferably 0.1% to 3% by weight, and more preferably 0.3% to 3% by weight with respect to the tablet.

[0038]

Examples of lubricants that can be used in the tablets provided by the present invention include magnesium stearate, calcium stearate, talc, light anhydrous silicic acid, sucrose fatty acid esters, carnauba wax, macrogol, and sodium stearyl fumarate, among which magnesium stearate is particularly preferable.

The lubricant content in the tablet provided by the present invention may be 0.1% to 30% by weight, preferably 0.3% to 10% by weight, and more preferably 1% to 3% by weight with respect to the tablet.

[0039]

Examples of pH adjusters that can be used in the tablets provided by the present invention include citric acid, phosphates (for example, sodium dihydrogen phosphate and potassium dihydrogen phosphate), carbonates (for example, magnesium carbonate and sodium carbonate), tartrate, fumarate, acetate, and amino acid salts (where, the pH adjuster does not include the alkalizing agent according to the present invention).

The content of the pH adjuster in the tablet provided by the present invention may be 0.1% to 30% by weight, preferably 0.3% to 10% by weight, and more preferably 1% to 5% by weight with respect to the tablet.

[0040]

Examples of surfactants that can be used in the tablets provided by the present

invention include sodium lauryl sulfate, polysorbate, sucrose fatty acid ester, polyoxyethylene hydrogenated castor oil, polyoxyl stearate, macrogol, and poloxamer. The content of the surfactant in the tablet provided by the present invention may be 0.01% to 3% by weight, preferably 0.03% to 1% by weight, and more preferably 0.03% to 0.5% by weight with respect to the tablet.

Examples of stabilizers that can be used in tablets provided by the present invention include citric acid (for example, anhydrous citric acid), malic acid, acetic acid, tartaric acid, maleic acid, ascorbic acid, edetate sodium, and tocopherol (where, the stabilizer does not include the alkalinizing agent according to the present invention), among which anhydrous citric acid is particularly preferable.

The content of the stabilizer in the tablet provided by the present invention may be 0.01% to 30% by weight, preferably 0.1% to 30% by weight, and more preferably 1% to 20% by weight with respect to the tablet.

[0041]

Examples of fragrances that can be used in the tablets provided by the present invention include citrus flavors such as lemon, orange, and grapefruit, peppermint, spearmint, and menthol. An appropriate amount thereof can be contained in the tablet (for example, 0.01% to 1% by weight, and more preferably 0.01% to 0.1% by weight with respect to the tablet).

The total content of the alkalinizing agent and the pharmaceutically acceptable additives in the tablet provided by the present invention does not exceed 100% by weight with respect to the tablet.

[0042]

The tablet provided by the present invention can be an uncoated tablet comprising the above components and not having a coating layer, or a film-coated tablet

having a coating layer. The content of the coating layer can be appropriately set by those skilled in the art. For example, it may be 0.1% to 10% by weight with respect to the uncoated tablet. In the coating layer, in addition to the coating base, a plasticizer, a coloring agent, a brightening agent, and the like can be appropriately incorporated.

5 Examples of coating bases that can be used for tablets provided by the present invention include hydroxypropylcellulose, hydroxypropyl methylcellulose, ethylcellulose, cellulose acetate phthalate, methacrylic acid copolymer, and polyvinylpyrrolidone, among which hydroxypropyl methylcellulose is particularly preferable. The content of the coating base in the tablet provided by the present
10 invention may be 0.01% to 10% by weight, and preferably 0.3% to 3% by weight with respect to the tablet.

 Examples of coating plasticizers that can be used in tablets provided by the present invention include triethyl citrate, medium chain fatty acid triglyceride, triacetin, glycerin, and propylene glycol and polyethylene glycol (for example, Macrogol 6000),
15 among which Macrogol 6000 is particularly preferable. The content of the coating plasticizer in the tablet provided by the present invention may be 0.01% to 1% by weight, and preferably 0.03% to 3% by weight with respect to the tablet.

 Examples of coating colorants that can be used in the tablets provided by the present invention include titanium oxide, yellow iron sesquioxide, iron sesquioxide,
20 black iron oxide, food blue No. 2, and food blue No. 2 aluminum lake. The content of the coating colorant in the tablet provided by the present invention may be 0.01% to 1% by weight, and preferably 0.03% to 3% by weight with respect to the tablet.

 Examples of coating brighteners that can be used in tablets provided by the present invention include carnauba wax. The content of the coating brightener in the
25 tablet provided by the present invention may be 0.0001% to 0.1% by weight, and

preferably 0.001% to 0.01% by weight with respect to the tablet.

[0043]

The pharmaceutical composition provided by the present invention can be produced by a method known in the pharmaceutical field. For example, in the case of
5 tablets, the production method may comprise a mixing step of mixing an additive with an alkalinizing agent (for example, potassium citrate or a hydrate thereof; sodium citrate or a hydrate thereof; a mixture of potassium citrate monohydrate and sodium citrate dihydrate; or sodium bicarbonate), a granulation step, a tableting step, and/or a coating step.

10 The mixing step may comprise a step of mixing an alkalinizing agent and an additive such as an excipient, a stabilizer, a disintegrant and/or a binder. In addition, before the tableting step, a step in which a mixture comprising the alkalinizing agent and an additive is mixed with a lubricant, flavoring agent, and/or fragrance may further be provided. Mixing can be performed using a V-type mixer, a W-type mixer, a container
15 mixer, a tumbler mixer, a stirring mixer, or the like.

The granulation step can be performed by a known granulation method in the pharmaceutical field. Examples of granulation methods include a dry granulation method, a wet granulation method, and a fluidized bed granulation method.

As one embodiment, the mixture obtained in the mixing step and the granulated
20 product obtained in the granulation step are appropriately pulverized and/or sieved to obtain a mixture or granulated product having a desired particle size. The pulverization can be performed by a pulverizer known in the pharmaceutical field, such as a ball mill, a jet mill, or a hammer mill. The sieving can be performed using a 16 mesh sieve (aperture of 1000 μm) to 32 mesh sieve (aperture of 500 μm), and the like.

25 The tableting step can be performed by a tableting method known in the

pharmaceutical field. Examples of tableting methods include a direct tableting method, a dry tableting method, a wet tableting method, and an external lubricant tableting method. For example, the mixture or granulated product obtained in the above steps can be tableted using a tableting machine known in the pharmaceutical field, such as a
5 single-shot tableting machine or a rotary tableting machine. When using a single tableting machine, a rotary tableting machine, and the like, a tableting pressure of 1 kN to 30 kN can be employed.

The coating step can be performed by a method known in the pharmaceutical field. For example, the coating can be performed by spray coating the outside of the
10 uncoated tablet with a coating liquid appropriately comprising a coating base, and a plasticizer, colorant, brightener, or the like.

In one embodiment, the tablet provided by the present invention can be produced by mixing an alkalizing agent with excipients (for example, lactose, D-mannitol, crystalline cellulose, and/or glucose), binders (for example,
15 hydroxypropylcellulose (HPC), gelatin, and/or polyvinylpyrrolidone (PVP)), stabilizers (for example, anhydrous citric acid), disintegrants (for example, starch (for example, partially pregelatinized starch) and/or carboxymethylcellulose calcium (CMC-Ca)), and lubricants (for example, magnesium stearate) to obtain uncoated tablets by tableting; and forming, on the outside of the uncoated tablet, a coating layer comprising a coating base
20 (for example, hydroxypropylcellulose, hydroxypropyl methylcellulose, and/or PVP), plasticizers (for example, triethyl citrate and/or Macrogol 6000), colorants (for example, iron sesquioxide and/or titanium oxide), and brighteners (for example, carnauba wax).

In one embodiment, the tablets obtained can have a hardness of 10 to 200 N, preferably 30 to 150 N.

The amount of the alkalinizing agent in the pharmaceutical composition provided by the present invention can be appropriately set.

In one embodiment, the amount of the alkalinizing agent in the pharmaceutical composition provided by the present invention may be set to an amount that enables
5 amelioration in acid urine in gout or hyperuricemia by administering the alkalinizing agent to a human, or an amount smaller than the above amount. For example, an amount may be set to be 1% to 50% or 10% to 20% of the daily dose approved in Japan for amelioration in acid urine in gout or hyperuricemia (for example, in a case where the
alkalinizing agent is a citric acid preparation: a tablet in which one tablet comprises 231.5
10 mg of potassium citrate ($C_6H_5K_3O_7 \cdot H_2O$) and 195.0 mg of sodium citrate hydrate ($C_6H_5Na_3O_7 \cdot 2H_2O$) is orally administered 3 times a day, 2 tablets at one time; and in a case where the alkalinizing agent is sodium bicarbonate: oral administration of 3 to 5 g a day).

In one embodiment, the pharmaceutical composition provided by the present
15 invention is a tablet, and it may comprise 10 mg to 1 g, preferably, 100 mg to 500 mg, and more preferably 400 mg to 500 mg of potassium citrate monohydrate or sodium citrate dihydrate as an alkalinizing agent in one tablet.

In one embodiment, the pharmaceutical composition provided by the present invention is a tablet, and it may comprise, in one tablet, 10 mg to 300 mg for each, a total
20 of 20 mg to 600 mg, preferably, 150 to 250 mg for each, a total of 400 to 500 mg, and more preferably 190 to 240 mg for each, a total of 400 to 450 mg of potassium citrate monohydrate and sodium citrate dihydrate.

In one embodiment, the pharmaceutical composition provided by the present invention is a tablet, and may comprise, in one tablet, 10 mg to 1 g, preferably 100 mg to

500 mg of sodium bicarbonate as an alkalinizing agent.

In one embodiment, the pharmaceutical composition provided by the present invention is a tablet, and it comprises, as an alkalinizing agent, potassium citrate monohydrate 231.5 mg and sodium citrate dihydrate 195.0 mg; and may comprise, as an additive, anhydrous citric acid, crystalline cellulose, partially pregelatinized starch, hydroxypropylcellulose, magnesium stearate, hypromellose, macrogol 6000, titanium oxide, and carnauba wax.

As one embodiment, a tablet comprising potassium citrate monohydrate 231.5 mg and sodium citrate dihydrate 195.0 mg may be a single dosage unit.

In the present specification, “dosage unit” represents a unit of the preparation, and “one dosage unit” represents the minimum unit of the preparation. Accordingly, for example, in the case of tablets, the dosage unit is each tablet, and one dosage unit represents one tablet. In the case of an injection, the dosage unit is an injection contained in a sealed container such as an ampoule or vial, and one dosage unit represents an injection contained in a sealed container such as one ampoule or vial.

In a case where the pharmaceutical composition provided by the present invention is administered to a human or other mammals, one or more of the above dosage units may be administered at one time, or the one dosage unit may be divided and administered.

[0045]

The dose of the alkalinizing agent is appropriately determined according to the type of the alkalinizing agent, the method of administration, the age, weight, sex, symptom, sensitivity to the drug, and the like of a subject of administration, but it may be adjusted depending on the condition of symptom amelioration.

In one embodiment, in a case of an oral administration of a mixture of potassium

citrate monohydrate and sodium citrate dihydrate or sodium bicarbonate as an
 alkalinizing agent to a human, half of the daily dose approved in Japan for amelioration
 in acid urine in gout or hyperuricemia (for example, in a case where the alkalinizing
 agent is a citric acid preparation: a tablet in which one tablet comprises 231.5 mg of
 5 potassium citrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and 195.0 mg of sodium citrate hydrate
 ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$) is orally administered 3 times a day, 2 tablets at one time; and in a
 case where the alkalinizing agent is sodium bicarbonate: oral administration of 3 to 5 g a
 day) may be set to a daily dose.

In one embodiment, in a case of an oral administration of a mixture of potassium
 10 citrate monohydrate and sodium citrate dihydrate or sodium bicarbonate as an
 alkalinizing agent to a human, the daily dose approved in Japan for amelioration in acid
 urine in gout or hyperuricemia (for example, in a case where the alkalinizing agent is a
 citric acid preparation: a tablet in which one tablet comprises 231.5 mg of potassium
 citrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and 195.0 mg of sodium citrate hydrate ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$) is
 15 orally administered 3 times a day, 2 tablets at one time; and in a case where the
 alkalinizing agent is sodium bicarbonate: oral administration of 3 to 5 g a day) may be set
 to a daily dose.

In one embodiment, in a case of an oral administration of a mixture of potassium
 citrate monohydrate and sodium citrate dihydrate or sodium bicarbonate as an
 20 alkalinizing agent to a human, half of the daily dose approved in Japan for amelioration
 in acid urine in gout or hyperuricemia (for example, in a case where the alkalinizing
 agent is a citric acid preparation: a tablet in which one tablet comprises 231.5 mg of
 potassium citrate ($\text{C}_6\text{H}_5\text{K}_3\text{O}_7 \cdot \text{H}_2\text{O}$) and 195.0 mg of sodium citrate hydrate
 ($\text{C}_6\text{H}_5\text{Na}_3\text{O}_7 \cdot 2\text{H}_2\text{O}$) is orally administered 3 times a day, 2 tablets at one time; and in a

case where the alkalinizing agent is sodium bicarbonate: oral administration of 3 to 5 g a day) may be set to a daily dose to start administration, and thereafter, the dosage may be increased to a daily dosage approved in Japan for the amelioration of acidic urine in gout and hyperuricemia.

5 In one embodiment, the dosage amount of the alkalinizing agent may be a dosage amount such that, when the alkalinizing agent is orally administered, the pH of human urine (for example, early morning urine) becomes pH 5.2 to pH 6.8, pH 5.5 to pH 6.8, pH 5.8 to pH 6.8, pH 5.8 to pH 6.5, pH 5.8 to pH 6.2, pH 5.8 or more and less than pH 6.2, pH 6.0 to pH 6.5, pH 6.0 to pH 6.4, pH 6.0 to pH 6.3, pH 6.0 to pH 6.2, pH 6.0
10 or more and less than pH 6.2, pH 6.1 to pH 6.3, pH 6.2 to 6.8, pH 6.2 to pH 6.5, or pH 6.5 to 6.8.

 In one embodiment, the dosage amount of the alkalinizing agent may be a dosage amount such that, when the alkalinizing agent is orally administered, the pH of human urine (for example, early morning urine) becomes pH 5.2 to pH 6.8, pH 5.5 to pH
15 6.8, pH 5.8 to pH 6.8, pH 5.8 to pH 6.5, pH 5.8 to pH 6.2, pH 5.8 or more and less than pH 6.2, pH 6.0 to pH 6.5, pH 6.0 to pH 6.4, pH 6.0 to pH 6.3, pH 6.0 to pH 6.2, pH 6.0 or more and less than pH 6.2, pH 6.1 to pH 6.3, pH 6.2 to 6.8, pH 6.2 to pH 6.5, or pH 6.5 to 6.8, 6 weeks, 12 weeks, or 24 weeks after the oral administration of the alkalinizing agent.

20 In one embodiment, when a mixture of potassium citrate monohydrate and sodium citrate dihydrate as an alkalinizing agent is orally administered to a human, potassium citrate monohydrate and sodium citrate dihydrate may be administered at 0.1 to 5g/day for each, a total of 0.2 to 10g/day, 0.1 to 3g/day for each, a total of 0.2 to 6g/day, 0.5 to 3g/day for each, a total of 1 to 6g/day, preferably, 0.5 to 1.5 g/day for each,
25 a total of 1 to 3 g/day, 1 to 1.5g/day for each, a total of 2 to 3g/day, or 0.5 to 1 g/day for

each, a total of 1 to 2 g/day; or may be divided and administered 1 to 5 times a day, preferably 3 times a day.

In one embodiment, when potassium citrate monohydrate or sodium citrate dihydrate is orally administered to a human as an alkalinizing agent, it may be
5 administered at 1 to 10 g/day, 1 to 6 g/day, 2 to 5.5 g/day, 1 to 3 g/day, 2 to 3 g/day, or 1 to 1.5 g/day; or may be divided and administered 1 to 5 times a day, preferably 3 times a day.

In one embodiment, when sodium bicarbonate is orally administered to a human as an alkalinizing agent, it may be administered at 1 to 6 g/day, preferably 1 to 3 g/day, or
10 3 to 5 g/day; or may be divided and administered 1 to 5 times a day, preferably 3 times a day.

[0046]

In one embodiment, the alkalinizing agent may be administered for a long period of time, for example, for 1 week, 2 weeks, 3 weeks, 6 weeks, 8 weeks, 10 weeks, 12
15 weeks, 24 weeks, 40 weeks, 60 weeks, 80 weeks, 100 weeks, 120 weeks, 1 week or more, 2 weeks or more, 3 weeks or more, 6 weeks or more, 8 weeks or more, 10 weeks or more, 12 weeks or more, 24 weeks or more, 40 weeks or more, 60 weeks or more, 80 weeks or more, 100 weeks or more, 120 weeks or more, 6 to 24 weeks, 12 to 24 weeks, 6
20 weeks, 12 to 30 weeks, 6 to 40 weeks, 12 to 40 weeks, 6 to 60 weeks, 12 to 60 weeks, 6 to 80 weeks, 12 to 80 weeks, 6 to 100 weeks, 12 to 100 weeks, 6 to 120 weeks, or 12 to 120 weeks.

In one embodiment, by continuous administration for 6 weeks, continuous administration for 12 weeks, and/or continuous administration for 24 weeks of the pharmaceutical composition provided by the present invention, beneficial effects for a
25 patient with kidney disease (for example, chronic kidney disease) (for example, an effect

of decreasing a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid) in blood, an effect of increasing a concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid) in urine (an effect of promoting excretion into urine), and/or an effect of increasing a urinary β 2-microglobulin concentration) can be detected.

[0047]

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a human suffering from kidney disease. Kidney disease includes acute kidney failure and chronic kidney disease unless otherwise specified.

Examples of acute kidney failure include acute kidney failure due to drugs (for example, platinum preparations such as non-steroidal anti-inflammatory drugs, angiotensin converting enzyme inhibitors, angiotensin II receptor antagonists, aminoglycoside antibiotics, new quinolone antibacterials, iodinated contrast agents, platinum preparations such as cisplatin, and the like), and acute kidney failure caused by kidney ischemia.

Chronic kidney disease (CKD) is a concept encompassing kidney disease that chronically progresses regardless of the underlying disease, and is a concept encompassing all clinical conditions in which kidney function expressed by glomerular filtration rate (GFR) deteriorates, or findings that suggest damage in kidneys is maintained chronically (3 months or longer).

[0048]

According to the CKD Medical Guide 2012 (Nichijinkaishi 2012), the severity of chronic kidney disease is evaluated by classification according to cause (Cause: C), kidney function (GFR: G), and proteinuria (albuminuria: A).

The classification of GFR is as follows.

G1: GFR is normal or high (≥ 90 mL/min/1.73 m²)

G2: Normal or slight decrease in GFR (60 to 89 mL/min/1.73 m²)

G3a: Mild to moderate decrease in GFR (45 to 59 mL/min/1.73 m²)

5 G3b: Moderate to high decrease in GFR (30 to 44 mL/min/1.73 m²)

G4: GFR is highly reduced (15 to 29 mL/min/1.73 m²)

G5: End-stage kidney disease (ESKD) (< 15 mL/min/1.73 m²)

The classification by proteinuria (albuminuria: A) is classified as follows using the urine albumin/creatinine (Cr) ratio when the primary disease is diabetes.

10 A1: Normal (less than 30 mg/gCr)

A2: Microalbuminuria (30 to 299 mg/gCr)

A3: Overt albuminuria (300 mg/gCr or more)

In addition, in a case where the primary disease is hypertension other than diabetes, nephritis, polycystic kidneys, transplanted kidneys, and the like, proteinuria (albuminuria: A) is classified using the urine protein/creatinine (Cr) ratio as follows.

A1: Normal (less than 0.15 g/gCr)

A2: Mild proteinuria (0.15 to 0.49 g/gCr)

A3: High proteinuria (0.50 g/gCr or more)

According to CKD medical care guide 2012 (Nichijinkaishi 2012), the severity classification of chronic kidney disease (CKD) is expressed as, for example, diabetes G2A3, chronic nephritis G3bA1, and the like using the above C, G, and A.

However, it is considered that the severity of chronic kidney disease has conventionally been described only in the stages classified by GFR, and as in the related art, the severity of chronic kidney disease can be expressed in stages of G1, G2, G3a, G3b, G4, and G5.

25 G3b, G4, and G5.

[0049]

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with low-severity, early-stage chronic kidney diseases.

5 In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease of stage G3b or lower, preferably stage G2 or lower.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease in stage G2 or more
10 and stage G3b or less (for example, stage G2 and stage G3a; or stage G2, stage G3a and stage G3b).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease in stage G3b or less and microalbuminuria, preferably administered to a patient with kidney disease in stage
15 G2 and chronic albuminuria.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease in stage G2 or more and stage G3b or less (for example, stage G2 and stage G3a; or stage G2, stage G3a and stage G3b) and which is microalbuminuria.

20 In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease in stage G3b or lower and whose urinary protein excretion is less than 3.5 g/gCr, and is preferably administered to a patient with chronic kidney disease in stage G2 and whose urinary protein excretion is less than 3.5 g/gCr.

25 In one embodiment, the pharmaceutical composition provided by the present

invention is administered to a patient with chronic kidney disease in stage G2 or more and stage G3b or less (for example, stage G2 and stage G3a; or stage G2, stage G3a and stage G3b) and whose urinary protein excretion is less than 3.5 g/gCr.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with progressive chronic kidney disease.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose concentration of β 2-microglobulin in urine (for example, early morning urine) is 2000 μ g/L or less, 1000 μ g/L or less, 800 μ g/L or less, 290 μ g/L or less, 200 μ g/L or less, 1 to 2000 μ g/L, 1 to 1000 μ g/L, 1 to 800 μ g/L, 1 to 290 μ g/L, 1 to 200 μ g/L, 10 to 2000 μ g/L, 10 to 1000 μ g/L, 10 to 800 μ g/L, 10 to 290 μ g/L, 10 to 200 μ g/L, or 80 to 200 μ g/L.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose cystatin C in blood (for example, in plasma or in serum) is 0.1 to 3.0 mg/L, 0.1 to 2.0 mg/L, 0.1 to 1.6 mg/L, 0.1 to 1.3 mg/L, 0.5 to 3.0 mg/L, 0.5 to 2.0 mg/L, 0.5 to 1.6 mg/L, 0.5 to 1.3 mg/L, or 0.9 to 1.3 mg/mL.

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose indoxyl sulfate concentration in blood is 0.001 to 100 μ g/mL (for example, 0.1 to 30 μ g/mL).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose concentration of p-cresyl sulfate in blood is 0.003 to 300 μ g/mL (for example, 0.01 to 30 μ g/mL).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose hippuric acid concentration in blood is 0.01 to 100 μ g/mL (for example, 0.01 to 10 μ g/mL).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose argininosuccinic acid concentration in blood is 0.01 to 100 $\mu\text{g/mL}$ (for example, 0.1 to 10 $\mu\text{g/mL}$).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease whose concentration of phenylacetyl-L-glutamine in blood is 0.03 to 30 $\mu\text{g/mL}$ (for example, 0.1 to 10 $\mu\text{g/mL}$).

[0050]

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient receiving treatment according to a CKD medical guide. For example, it is administered to a patient subjected to, according to CKD medical guide, blood pressure management (administration of RA inhibitors such as ARB and ACE inhibitors, diuretics, Ca antagonists), anti-proteinuria measures (administration of RA inhibitors, and the like), blood glucose level control (administration of α -glucosidase inhibitor, and the like), lipid management (administration of statins, fibrates, and the like), anemia management (for example, erythropoietin administration), and/or bone/mineral measures (bisphosphonate administration and the like).

In one embodiment, the pharmaceutical composition provided by the present invention is used in combination with an antihypertensive agent (for example, ARB, ACE inhibitor, diuretic, Ca antagonist).

In one embodiment, the pharmaceutical composition provided by the present invention is used in combination with a spherical carbonaceous adsorbent (sold in Japan as KREMEZIN (registered trademark)) obtained by oxidation and reduction treatment of spherical fine porous carbon derived from petroleum hydrocarbons at high temperature.

[0051]

In one embodiment, the pharmaceutical composition provided by the present

invention is administered to a patient with low-severity, early-stage chronic kidney disease (for example, a patient with chronic kidney disease in stage G3b or lower, preferably stage G2 to stage G3b, more preferably stage G2 and stage G3a, and even more preferably stage G2), and thereby a concentration of uremic substance (for example, indoxyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine) in blood of a patient is decreased, and excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, hippuric acid, argininosuccinic acid, and/or phenylacetyl-L-glutamine) outside the body (for example, into urine) is promoted. In this embodiment, the pharmaceutical composition provided by the present invention may be a pharmaceutical composition for ameliorating uremic symptoms, a pharmaceutical composition for treating or preventing uremia, a pharmaceutical composition for suppressing progression of chronic kidney disease, a pharmaceutical composition for delaying introduction of dialysis, a pharmaceutical composition for suppressing myocardial fibrosis, a pharmaceutical composition for suppressing arteriosclerosis, a pharmaceutical composition for ameliorating arteriosclerosis, a pharmaceutical composition for suppressing vascular smooth muscle cell proliferation, a pharmaceutical composition for suppressing vascular endothelial cell injury, a pharmaceutical composition for suppressing arterial wall thickening, a pharmaceutical composition for ameliorating arterial wall thickening, a pharmaceutical composition for suppressing calcification of the aorta, or a pharmaceutical composition for treating or preventing cardiovascular disease, which is a complication (for example, heart failure, myocardial infarction, stroke, and the like).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with low-severity, early-stage chronic kidney disease (for example, a patient with chronic kidney disease in stage G3b or lower,

preferably stage G2 to stage G3b, more preferably stage G2 and stage G3a, and even more preferably stage G2), and thereby an increase in urinary β 2-microglobulin in the patient is suppressed. In this embodiment, the pharmaceutical composition provided by the present invention may be a composition for suppressing damage of kidney tubule (for example, kidney proximal tubule), a composition for suppressing damage of kidney tubular cells (for example, kidney proximal tubular cells), a composition for protecting kidney tubular cells (for example, kidney proximal tubular cells), or a pharmaceutical composition for maintaining kidney tubular cell function (for example, reabsorption of glucose, amino acids, and the like) (for example, kidney proximal tubular cell function).

10 In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease having a moderate severity or higher (for example, a patient with chronic kidney disease in stage G3b or higher), and thereby a concentration of uremic substance (for example, indoxyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine) in blood of the patient is decreased, and
15 excretion of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, hippuric acid, argininosuccinic acid, and/or phenylacetyl-L-glutamine) outside the body (for example, into urine) is promoted. In this embodiment, the pharmaceutical composition provided by the present invention may be a pharmaceutical composition for ameliorating uremic symptoms, a pharmaceutical composition for treating or preventing uremia, a
20 pharmaceutical composition for suppressing progression of chronic kidney disease, a pharmaceutical composition for delaying introduction of dialysis, a pharmaceutical composition for suppressing myocardial fibrosis, a pharmaceutical composition for suppressing arteriosclerosis, a pharmaceutical composition for ameliorating arteriosclerosis, a pharmaceutical composition for suppressing vascular smooth muscle
25 cell proliferation, a pharmaceutical composition for suppressing vascular endothelial cell

injury, a pharmaceutical composition for suppressing arterial wall thickening, a pharmaceutical composition for ameliorating arterial wall thickening, a pharmaceutical composition for suppressing calcification of the aorta, a pharmaceutical composition for suppressing a decrease in energy production in muscle cells, a pharmaceutical composition for suppressing a decrease in muscle mass and/or muscle strength, or a pharmaceutical composition for treating or preventing cardiovascular disease, which is a complication (for example, heart failure, myocardial infarction, stroke, and the like).

In one embodiment, the pharmaceutical composition provided by the present invention is administered to a patient with chronic kidney disease having a moderate severity or higher (for example, a patient with chronic kidney disease in stage G3b or higher), and thereby an increase in urinary β 2-microglobulin in the patient is suppressed. In this embodiment, the pharmaceutical composition provided by the present invention may be a composition for suppressing damage of kidney tubule (for example, kidney proximal tubule), a composition for suppressing damage of kidney tubular cells (for example, kidney proximal tubular cells), a composition for protecting kidney tubular cells (for example, kidney proximal tubular cells), or a pharmaceutical composition for maintaining kidney tubular cell function (for example, reabsorption of glucose, amino acids, and the like) (for example, kidney proximal tubular cell function).

[0052]

Examples of other embodiments of the present invention include the following:

a) A method for decreasing a concentration of uremic substance in blood in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which a concentration of uremic substance in blood is required to be decreased;

b) A method for promoting excretion of uremic substance into urine in a

mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which excretion of uremic substance into urine is required to be promoted;

c) A method for ameliorating uremic symptoms in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which amelioration of uremic symptoms is required;

d) A method for treating or preventing uremia in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which uremic symptoms are required to be treated or prevented;

e) A method for suppressing progression of chronic kidney disease in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which progression of chronic kidney disease is required to be suppressed;

f) A method for delaying introduction of dialysis in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from chronic kidney disease and in which delaying introduction of dialysis is required;

g) A method for suppressing myocardial fibrosis in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which myocardial fibrosis is required to be suppressed;

h) A method for suppressing arteriosclerosis in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of

an alkalinizing agent to a subject who suffers from kidney disease and in which arteriosclerosis is required to be suppressed;

i) A method for suppressing vascular smooth muscle cell proliferation in a mammalian subject (for example, a human), wherein the method comprises
5 administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which vascular smooth muscle cell proliferation is required to be suppressed;

j) A method for suppressing vascular endothelial cell injury in a mammalian subject (for example, a human), wherein the method comprises administering an effective
10 amount of an alkalinizing agent to a subject who suffers from kidney disease and in which vascular endothelial cell injury is required to be suppressed;

k) A method for suppressing thickening of the arterial wall in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in
15 which thickening of the arterial wall is required to be suppressed;

l) A method for suppressing calcification of the aorta in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which calcification of the aorta is required to be suppressed;

20 m) A method for treating or preventing cardiovascular disease in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject who suffers from kidney disease and in which cardiovascular disease is required to be treated or prevented;

n) A method for ameliorating arteriosclerosis in a mammalian subject (for
25 example, a human), wherein the method comprises administering an effective amount of

an alkalinizing agent to a subject who suffers from kidney disease and in which arteriosclerosis is required to be ameliorated;

o) A method for ameliorating thickening of the arterial wall in a mammalian subject (for example, a human), wherein the method comprises administering an effective
5 amount of an alkalinizing agent to a subject who suffers from kidney disease and in which thickening of the arterial wall is required to be ameliorated;

p) A method for treating acute kidney failure in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which acute kidney failure is required to be treated;

10 q) A method for suppressing progression from acute kidney failure to chronic kidney disease in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which progression from acute kidney failure to chronic kidney disease is required to be suppressed;

15 r) A method for treating or preventing kidney tubular damage in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which kidney tubular damage is required to be treated or prevented;

s) A method for suppressing kidney tubular damage in a mammalian subject (for
20 example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which kidney tubular damage is required to be suppressed;

t) A method for suppressing kidney proximal tubular cell damage in a mammalian subject (for example, a human), wherein the method comprises
25 administering an effective amount of an alkalinizing agent to a subject in which kidney

proximal tubular cell damage is required to be suppressed;

u) A method for protecting kidney proximal tubular cells in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which kidney proximal tubular cells are
5 required to be protected;

v) A method for maintaining kidney proximal tubular cell function in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which kidney proximal tubular cell function is required to be maintained;

10 w) A method for promoting excretion of uremic substance outside the body in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which excretion of uremic substance outside the body is required to be promoted;

x) A method for excreting uremic substance into urine depending on a
15 concentration of uremic substance in blood in a mammalian subject (for example, a human), wherein the method comprises administering an effective amount of an alkalinizing agent to a subject in which uremic substance is required to be excreted into urine;

[0053]

20 aa) An alkalinizing agent for use in decreasing a concentration of uremic substance in blood;

bb) An alkalinizing agent for use in promoting excretion of uremic substance into urine;

cc) An alkalinizing agent for use in ameliorating uremic symptoms in a patient
25 with kidney disease;

dd) An alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease;

ee) An alkalinizing agent for use in suppressing progression of chronic kidney disease;

5 ff) An alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease;

gg) An alkalinizing agent for use in suppressing myocardial fibrosis in a patient with kidney disease;

10 hh) An alkalinizing agent for use in suppressing arteriosclerosis in a patient with kidney disease;

ii) An alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease;

jj) An alkalinizing agent for use in suppressing vascular endothelial cell injury in a patient with kidney disease;

15 kk) An alkalinizing agent for use in suppressing thickening of the arterial wall in a patient with kidney disease;

ll) An alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease;

20 mm) An alkalinizing agent for use in treating or preventing cardiovascular disease in a patient with kidney disease;

nn) An alkalinizing agent for use in ameliorating arteriosclerosis in a patient with kidney disease;

oo) An alkalinizing agent for use in ameliorating thickening of the arterial wall of a patient with kidney disease;

25 pp) An alkalinizing agent for use in treating acute kidney failure;

qq) An alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease;

rr) An alkalinizing agent for use in treating or preventing kidney tubular damage;

5 ss) An alkalinizing agent for use in suppressing kidney tubular damage;

tt) An alkalinizing agent for use in suppressing kidney proximal tubular cell damage;

uu) An alkalinizing agent for use in protecting kidney proximal tubular cells;

vv) An alkalinizing agent for use in maintaining kidney proximal tubular cell
10 function;

ww) An alkalinizing agent for use in promoting excretion of uremic substance outside the body;

xx) An alkalinizing agent for use in excreting of uremic substance into urine depending on a concentration of uremic substance in blood;

15 [0054]

aaa) A pharmaceutical composition comprising an alkalinizing agent for use in decreasing a concentration of uremic substance in blood;

bbb) A pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance into urine;

20 ccc) A pharmaceutical composition comprising an alkalinizing agent for use in ameliorating uremic symptoms in a patient with kidney disease;

ddd) A pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease;

eeee) A pharmaceutical composition comprising an alkalinizing agent for use in
25 suppressing progression of chronic kidney disease;

fff) A pharmaceutical composition comprising an alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease;

ggg) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing myocardial fibrosis in a patient with kidney disease;

5 hhh) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing arteriosclerosis in a patient with kidney disease;

iii) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease;

10 jjj) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular endothelial cell injury in a patient with kidney disease;

kkk) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing thickening of the arterial wall of a patient with kidney disease;

lll) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease;

15 mmm) A pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing cardiovascular disease in a patient with kidney disease;

nnn) A pharmaceutical composition comprising an alkalinizing agent for use in ameliorating arteriosclerosis in a patient with kidney disease;

20 ooo) A pharmaceutical composition comprising an alkalinizing agent for use in ameliorating thickening of the arterial wall of a patient with kidney disease;

ppp) A pharmaceutical composition comprising an alkalinizing agent for use in treating acute kidney failure;

qqq) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease;

25 rrr) A pharmaceutical composition comprising an alkalinizing agent for use in

treating or preventing kidney tubular damage;

sss) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney tubular damage;

5 ttt) A pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney proximal tubular cell damage;

uuu) A pharmaceutical composition comprising an alkalinizing agent for use in protecting kidney proximal tubular cells;

vvv) A pharmaceutical composition comprising an alkalinizing agent for use in maintaining kidney proximal tubular cell function;

10 www) A pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance outside the body;

xxx) A pharmaceutical composition comprising an alkalinizing agent for use in excretion of uremic substance into urine depending on a concentration of uremic substance in blood;

15 [0055]

aaaa) Use of an alkalinizing agent to produce a pharmaceutical composition for decreasing a concentration of uremic substance in blood;

bbbb) Use of an alkalinizing agent to produce a pharmaceutical composition for promoting excretion of uremic substance into urine;

20 cccc) Use of an alkalinizing agent to produce a pharmaceutical composition for ameliorating uremic symptoms in a patient with kidney disease;

dddd) Use of an alkalinizing agent to produce a pharmaceutical composition for treating or preventing uremia in a patient with kidney disease;

25 eeee) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing progression of chronic kidney disease;

ffff) Use of an alkalinizing agent to produce a pharmaceutical composition for delaying introduction of dialysis to a patient with chronic kidney disease;

gggg) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing myocardial fibrosis in a patient with kidney disease;

5 hhhh) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing arteriosclerosis in a patient with kidney disease;

 iiii) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing vascular smooth muscle cell proliferation in a patient with kidney disease;

 jjjj) Use of an alkalinizing agent to produce a pharmaceutical composition for
10 suppressing vascular endothelial cell injury in a patient with kidney disease;

 kkkk) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing thickening of the arterial wall of a patient with kidney disease;

 llll) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing calcification of the aorta of a patient with kidney disease;

15 mmmm) Use of an alkalinizing agent to produce a pharmaceutical composition for treating preventing cardiovascular disease in a patient with kidney disease;

 nnnn) Use of an alkalinizing agent to produce a pharmaceutical composition for ameliorating arteriosclerosis in a patient with kidney disease;

 oooo) Use of an alkalinizing agent to produce a pharmaceutical composition for
20 ameliorating thickening of the arterial wall of a patient with kidney disease;

 pppp) Use of an alkalinizing agent to produce a pharmaceutical composition for treating acute kidney failure;

 qqqq) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing progression from acute kidney failure to chronic kidney disease;

25 rrrr) Use of an alkalinizing agent to produce a pharmaceutical composition for

treating or preventing kidney tubular damage;

ssss) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing kidney tubular damage;

5 tttt) Use of an alkalinizing agent to produce a pharmaceutical composition for suppressing kidney proximal tubular cell damage;

uuuu) Use of an alkalinizing agent to produce a pharmaceutical composition for protecting kidney proximal tubular cells;

vvvv) Use of an alkalinizing agent to produce a pharmaceutical composition for maintaining kidney proximal tubular cell function;

10 wwww) Use of an alkalinizing agent to produce a pharmaceutical composition for promoting excretion of uremic substance outside the body; and

xxxx) Use of an alkalinizing agent to produce a pharmaceutical composition for excretion of uremic substance into urine depending on a concentration of uremic substance in blood.

15 [0056]

2. Food composition

In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent and exhibits an effect of promoting excretion, outside the body, of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, p-cresyl sulfate, and phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate).

20

In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent and exhibits an effect of decreasing the concentration of uremic substance in blood.

25

In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent and exhibits an effect of promoting excretion of uremic substance into urine.

In one embodiment, the food composition provided by the present invention
5 comprises an alkalinizing agent and exhibits a kidney function maintenance effect.

In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent and exhibits an effect of suppressing kidney tubular damage. Examples of kidney tubules include a kidney proximal tubule.

In one embodiment, the food composition provided by the present invention
10 comprises an alkalinizing agent and exhibits an effect of suppressing kidney proximal tubular cell damage.

In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent and exhibits an effect of protecting kidney proximal tubular cells.

15 In one embodiment, the food composition provided by the present invention comprises an alkalinizing agent, and exhibits an effect of maintaining kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like). Examples of kidney tubules include a kidney proximal tubule, and examples of kidney
20 proximal tubular function include reabsorption of glucose, amino acids, vitamins, and the like.

In the above-described embodiment, the food composition provided by the present invention exhibits an effect of suppressing an increase in the amount (concentration) of β 2-microglobulin in urine (for example, early morning urine)
25 associated with progression of a stage of chronic kidney disease.

In the above-described embodiment, the food composition provided by the present invention does not affect the glomerular function of a patient with chronic kidney disease, whereas it exhibits an effect of suppressing kidney proximal tubular cell damage associated with progression of a stage of chronic kidney disease to protect kidney proximal tubular cells.

For the alkalinizing agent, the alkalinizing agent described in “1. Pharmaceutical composition” above can be applied. Examples of alkalinizing agents include a pharmaceutically acceptable salt of citric acid as acceptable salts of citric acid as food (for example, an alkali metal salt of citric acid or a hydrate thereof, or a mixture thereof), and sodium bicarbonate. A mixture of a potassium citrate monohydrate ($C_6H_5K_3O_7 \cdot H_2O$) and a sodium citrate dihydrate ($C_6H_5Na_3O_7 \cdot 2H_2O$), or sodium citrate dihydrate is preferable.

Uremic substance is also as described in “1. Pharmaceutical composition” above. Examples of uremic substances include indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid.

The content of the alkalinizing agent in the food composition provided by the present invention can be appropriately determined depending on the type of food. Examples of food compositions include foods for specified health use, dietary supplements, functional foods, hospital patient foods, and supplements. The form of these food compositions is not particularly limited as long as it comprises an effective amount of an alkalinizing agent for exerting the above-mentioned effects and can be taken orally. The form may be a form of a normal food or drink, or the food composition may be provided as a preparation suitable for oral administration, for example, a preparation such as a tablet, a capsule, or a suspension, among preparations that can be applied to the pharmaceutical composition. Regarding the constitution and

production method of these preparations, in the present specification, the constitution and production method of the pharmaceutical preparation described in the above “1.

Pharmaceutical composition” can be applied as it is, and known formulation techniques in the field of pharmaceutical preparation technology itself can be also be applied.

5 For example, in the case of foods for specified health use, dietary supplements, functional foods, or foods for hospital patients, per serving of food, an amount of 1/3 of a total of 1 to 3 g of a potassium citrate monohydrate and a sodium citrate dihydrate may be contained as an alkalinizing agent, or an amount of 1/3 of 1 to 6 g of a sodium
10 hydrogen carbonate may be contained as an alkalinizing agent. When specified health foods, dietary supplements, functional foods, hospital patient foods, or supplements are provided as tablets, for example, per tablet, 70% to 80% by weight of the alkalinizing agent may be contained in 300 mg to 600 mg of a tablet.

 When the food composition provided by the present invention is not formulated and is provided in the form of a normal food or drink, it can be appropriately produced
15 by those skilled in the art depending on the type of the food. For example, it can be produced by blending an alkalinizing agent (for example, potassium citrate and/or sodium citrate) with the food material.

 The form of the food and drink is a liquid or milky or pasty food such as a beverage, soy sauce, milk, yogurt, or miso; a semi-solid food such as jelly or gummy;
20 solid foods such as rice cakes, gums, tofu, and supplements; powdered foods; and the like.

 Examples of beverages include fruit juice, fruit drinks, coffee drinks, oolong tea drinks, green tea drinks, tea drinks, barley tea drinks, vegetable drinks, carbonated soft
25 drinks, fruit extract drinks, vegetable extract juices, near water, sports drinks, diet drinks, and the like.

In beverages, additives such as antioxidants, fragrances, various esters, organic acids, organic acid salts, inorganic acids, inorganic acid salts, inorganic salts, pigments, emulsifiers, preservatives, seasonings, sweeteners, acidulants, fruit juice extracts, vegetable extracts, nectar extract, pH adjusters, and quality stabilizers can be blended in
5 individually or in combination.

The food composition provided by the present invention can be used in the same manner as the method of using the pharmaceutical composition described in the above “1. Pharmaceutical composition,” and can also be used within a range not intended to treat or prevent diseases. That is, when the alkalinizing agent contained in the food
10 composition according to the present invention is used as a reference, it can be applied to an application target of the pharmaceutical composition such that the amount of the alkalinizing agent used in the food composition becomes the same as the alkalinizing agent contained in the pharmaceutical composition. In addition, in one embodiment, the “food composition” according to the present invention can be applied to a subject (for
15 example, a human or other mammals) who does not have “morbid” or “abnormal” symptoms, conditions, or diseases, that is, a subject (for example, a human or other mammals) in a “healthy” or “normal” state in order to maintain or enhance a “healthy” or “normal” state. Furthermore, it can be applied to a “healthy subject concerned about kidney health” or “healthy subject concerned about kidney tubule health” in order to
20 maintain or improve a “healthy” or “normal” state. In this case, even if the alkalinizing agent is a component of the pharmaceutical composition or a component of the food composition, the pharmacological effect of the alkalinizing agent itself is basically the same. Accordingly, the application amount and application method of the food composition can be appropriately adjusted based on the alkalinizing agent according to
25 an expected effect.

A food composition applied to a subject (for example, a human or other mammals) who does not have “morbid” or “abnormal” symptoms, conditions, or diseases, that is, a subject (for example, a human or other mammals) in a “healthy” or “normal” state in order to maintain or enhance a “healthy” or “normal” state, may be particularly referred to as “functional food” in some cases.

The term “administration” described in the above “1. Pharmaceutical composition” can also be applied to the “food composition” according to the present invention. Furthermore, regarding the “food composition” according to the present invention, the term “administration” can be read as “ingestion.” Accordingly, for example, the terms “administer,” “administered,” and the like can be read as “is ingested,” “ingest,” “ingesting,” and the like, with different word forms depending on the context.

Accordingly, the embodiments of the food composition according to the present invention include the following:

<1> A food composition for decreasing a concentration of uremic substance in blood, wherein the food composition comprises an alkalinizing agent;

<2> A food composition for promoting excretion of uremic substance into urine, wherein the food composition comprises an alkalinizing agent;

<3> A food composition for maintenance of kidney function, wherein the food composition comprises an alkalinizing agent;

<4> A food composition for suppression of kidney tubular damage, wherein the food composition comprises an alkalinizing agent;

<5> A food composition for suppression of kidney tubular cell damage, preferably for suppression of kidney proximal tubular cell damage, wherein the food composition comprises an alkalinizing agent;

<6> A food composition for protection of kidney tubular cells, preferably for protection of kidney proximal tubular cells, wherein the food composition comprises an alkalinizing agent;

<7> A food composition for maintenance of kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like), preferably for maintenance of kidney proximal tubular function (for example, reabsorption of glucose, amino acids, vitamins, and the like), wherein the food composition comprises an alkalinizing agent;

<11> A method for decreasing a concentration of uremic substance in blood, wherein the method comprises ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which a concentration of uremic substance in blood is required to be decreased;

<22> A method for promoting excretion of uremic substance into urine, wherein the method comprises ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which promotion of uremic substance to be excreted into urine is required;

<33> A method for maintaining kidney function, wherein the method comprises ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which maintenance of kidney function is required;

<44> A method for suppressing kidney tubular damage, wherein the method comprises ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which suppression of kidney tubular damage is required;

<55> A method for suppressing damage of kidney tubular cells, preferably kidney proximal tubular cells, wherein the method comprises ingesting a food

composition comprising an effective amount of an alkalinizing agent to a subject in which suppression of damage of kidney tubular cells, preferably kidney proximal tubular cells is required;

5 <66> A method for protecting kidney tubular cells, preferably kidney proximal tubular cells, wherein the method comprises ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which protection of kidney tubular cells, preferably kidney proximal tubular cells is required;

10 <77> A method for maintaining kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like), preferably kidney proximal tubular function (for example, reabsorption of glucose, amino acids, vitamins, and the like), the method including ingesting a food composition comprising an effective amount of an alkalinizing agent to a subject in which maintenance of kidney tubular function, preferably kidney proximal tubular function is required;

15 <111> A food composition comprising an alkalinizing agent for decreasing a concentration of uremic substance in blood;

 <222> A food composition comprising an alkalinizing agent for promoting excretion of uremic substance into urine;

20 <333> A food composition comprising an alkalinizing agent for maintenance of kidney function;

 <444> A food composition comprising an alkalinizing agent for suppression of kidney tubular damage;

 <555> A food composition comprising an alkalinizing agent for suppression of damage of kidney tubular cells, preferably kidney proximal tubular cells;

25 <666> A food composition comprising an alkalinizing agent for protection of

kidney tubular cells, preferably kidney proximal tubular cells;

<777> A food composition for maintenance of kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like), preferably kidney proximal tubular function (for example, reabsorption of glucose, amino acids, vitamins, and the like), wherein the food composition comprises an alkalinizing agent;

<1111> Use of an alkalinizing agent to produce a food composition for decreasing a concentration of uremic substance in blood;

<2222> Use of an alkalinizing agent to produce a food composition for promoting excretion of uremic substance into urine;

<3333> Use of an alkalinizing agent to produce a food composition for maintenance of kidney function;

<4444> Use of an alkalinizing agent to produce a food composition for suppression of kidney tubular damage;

<5555> Use of an alkalinizing agent to produce a food composition for suppression of damage of kidney tubular cells, preferably kidney proximal tubular cell damage;

<6666> Use of an alkalinizing agent to produce a food composition for protection of kidney tubular cells, preferably kidney proximal tubular cells; and

<7777> Use of an alkalinizing agent to produce a food composition for maintenance of kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like), preferably kidney proximal tubular function (for example, reabsorption of glucose, amino acids, vitamins, and the like)

On packaging, a container, or an instruction leaflet for the food composition

according to the present invention, effects of decreasing a concentration of uremic substance in blood; effects of promotion of excretion of uremic substance into urine; effects of maintenance of kidney function; effects of suppression of kidney tubular damage; effects of suppression of kidney tubular cell damage; effects of suppression of kidney proximal tubular cell damage; effects of protection of kidney tubular cells; effects of protection of kidney proximal tubular cells; effects of maintenance of kidney tubular function (for example, reabsorption of water, sodium ion, potassium ion, calcium ion, phosphate ion, bicarbonate ion, chloride ion, glucose, amino acid, vitamin, and the like), preferably maintenance of kidney proximal tubular function (for example, reabsorption of glucose, amino acids, vitamins, and the like); and the like; are preferably indicated.

In one embodiment, the “food composition” according to the present invention is ingested by a subject (for example, a human or other mammals) in which a urinary β 2-microglobulin concentration is 290 μ g/L or less, preferably 50 to 150 μ g/L.

In one embodiment, the “food composition” according to the present invention is ingested by a subject (for example, a human or other mammals) in which a blood cystatin C concentration is 0.5 to 2.2 mg/L, preferably 1.0 to 1.3 mg/L.

In one embodiment, the ingestion of the “food composition” according to the present invention suppresses an increase in urinary β 2-microglobulin concentration.

In one embodiment, the ingestion of the “food composition” according to the present invention suppresses an increase in urinary β 2-microglobulin concentration 12 weeks after the administration.

In one embodiment, the ingestion of the “food composition” according to the present invention does not substantially decrease a urinary β 2-microglobulin concentration compared to before the start of administration.

In one embodiment, the ingestion of the “food composition” according to the

present invention does not substantially decrease a urinary β 2-microglobulin concentration 12 weeks after administration, compared to before the start of administration.

In one embodiment, the ingestion of the “food composition” according to the present invention does not substantially increase blood cystatin C compared to before the start of administration.

In one embodiment, the ingestion of the “food composition” according to the present invention does not substantially increase blood cystatin C compared to before the start of administration.

In one embodiment, the ingestion of the “food composition” according to the present invention suppresses, in early morning urine, an increase of a β 2-microglobulin amount which is associated with progression of a stage of chronic kidney disease.

In one embodiment, the ingestion of the “food composition” according to the present invention does not affect the glomerular function of a patient with chronic kidney disease, whereas it suppresses kidney proximal tubular cell damage associated with progression of a stage of chronic kidney disease progression to protect kidney proximal tubular cells.

[0057]

3. Method for determining decrease in concentration of uremic substance in blood

In one embodiment, the present invention provides a method for determining a decrease in concentration of uremic substance (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, hippuric acid, and phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate) in blood

of a patient with chronic kidney disease, comprising measuring a pH of urine.

In addition, in one embodiment, the present invention provides a method for determining a promotion of excretion of uremic substance (indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably
5 indoxyl sulfate, p-cresyl sulfate, hippuric acid, and phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate) into urine of a patient with chronic kidney disease, comprising measuring a pH of urine.

Measurement of the content of uremic substance (for example, indoxyl sulfate)
10 in body fluid can be performed by a HPLC method or enzyme method. However, these measurement methods require specialized and expensive reagents.

As described in the present specification, a concentration of uremic substance in blood (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid) is decreased by administering the alkalinizing agent,
15 and thereby excretion of these uremic substance into the urine is promoted.

Accordingly, by measuring a pH of urine of a patient with chronic kidney disease, very simply and inexpensively, it is possible to determine a decrease in concentration of uremic substance in blood (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid), and/or a promotion of excretion
20 of uremic substance into urine (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid). For the measurement of pH, a well-known technique may be used. For example, a pH test paper, a pH test solution, or a simple pH measuring device can be used.

In one embodiment, a patient with chronic kidney disease measures a pH of
25 early morning urine (wake up first urine) over time from the start of taking an

alkalinizing agent (for example, a mixture of a potassium citrate monohydrate and a sodium citrate dihydrate, or sodium citrate dihydrate), and if the urine pH is high, it can be easily determined that a decrease in concentration of uremic substance in blood (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate), and/or a promotion of excretion of uremic substance into urine (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, p-cresyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate) is achieved.

In addition, in one embodiment, a patient with chronic kidney disease measures a pH of early morning urine (wake up first urine) over time after taking an alkalinizing agent (for example, a mixture of a potassium citrate monohydrate and a sodium citrate dihydrate, or sodium citrate dihydrate). If the urine pH is within a range of 5.2 to 6.8 (for example, the pH of urine is within a range of pH 5.5 to pH 6.8, pH 5.8 to pH 6.8, pH 5.8 to pH 6.5, pH 5.8 to pH 6.2, pH 5.8 or more and less than pH 6.2, pH 6.0 to pH 6.5, pH 6.0 to pH 6.4, pH 6.0 to pH 6.3, pH 6.0 to pH 6.2, pH 6.0 or more and less than pH 6.2, pH 6.1 to pH 6.3, pH 6.2 to 6.8, pH 6.2 to pH 6.5, or pH 6.5 to 6.8), it can be easily determined that a decrease in concentration of uremic substance in blood (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate), and/or a promotion of excretion of uremic substance into

urine (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid, preferably indoxyl sulfate, p-cresyl sulfate, hippuric acid, and/or phenylacetyl-L-glutamine, more preferably indoxyl sulfate and phenylacetyl-L-glutamine, and even more preferably indoxyl sulfate) is achieved.

5 The determination, which is thus obtained as above, as to whether or not a decrease in concentration of uremic substance in blood (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic acid), and/or a promotion of excretion of uremic substance into urine (for example, indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and/or argininosuccinic
10 acid) is achieved can aid the diagnosis of whether or not progression of chronic kidney disease is suppressed.

Accordingly, in one embodiment, the present invention provides a method for determining suppression of progression of chronic kidney disease, comprising measuring the pH of urine (for example, early morning urine) of a patient to which an alkalinizing
15 agent (for example, a mixture of a potassium citrate monohydrate and a sodium citrate dihydrate, or sodium citrate dihydrate) has been administered. If it is recognized that the urine pH increases over time or the urine pH is within the range of 5.8 to 6.8 (for example, the urine pH is within the range of 6.0 to 6.2), this can then aid the diagnosis that progression of a stage of chronic kidney disease progression are suppressed.

20 [0058]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in decreasing a concentration of uremic substance in blood, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the
25 alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day,

preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in decreasing a concentration of uremic substance in blood, in which one dosage unit
5 (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0059]

In one embodiment, the pharmaceutical composition provided by the present
10 invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance into urine, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

15 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance into urine, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided
20 doses per day.

[0060]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating uremic symptoms in a patient with kidney disease, in which 0.5 to 1.5 g/day
25 of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the

alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating uremic symptoms in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0061]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0062]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression of chronic kidney disease, in which 0.5 to 1.5 g/day of each of a

potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression of chronic kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

10 [0063]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0064]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in

suppressing myocardial fibrosis in a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

5 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing myocardial fibrosis in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in
10 three divided doses per day.

[0065]

 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing arteriosclerosis in a patient with kidney disease, in which 0.5 to 1.5 g/day of
15 each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
20 suppressing arteriosclerosis in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0066]

25 In one embodiment, the pharmaceutical composition provided by the present

invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5
5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate
10 monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0067]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
15 suppressing vascular endothelial cell injury in a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present
20 invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular endothelial cell injury in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

25 [0068]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing thickening of the arterial wall of a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing thickening of the arterial wall of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0069]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0070]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing cardiovascular disease in a patient with kidney disease, in which
5 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
10 treating or preventing cardiovascular disease in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0071]

15 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating arteriosclerosis in a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day,
20 preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating arteriosclerosis in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg
25 sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three

divided doses per day.

[0072]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating thickening of the arterial wall of a patient with kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating thickening of the arterial wall of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0073]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating of acute kidney failure, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating of acute kidney failure, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0074]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0075]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing kidney tubular damage, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing kidney tubular damage, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses

per day.

[0076]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
5 suppressing kidney tubular damage, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
10 suppressing kidney tubular damage, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0077]

In one embodiment, the pharmaceutical composition provided by the present
15 invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney proximal tubular cell damage, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

20 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney proximal tubular cell damage, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided
25 doses per day.

[0078]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in protecting kidney proximal tubular cells, in which 0.5 to 1.5 g/day of each of a potassium
5 citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in protecting kidney proximal tubular cells, in which one dosage unit (preferably one tablet)
10 contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0079]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
15 maintaining kidney proximal tubular cell function, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present
20 invention is a pharmaceutical composition comprising an alkalinizing agent for use in maintaining kidney proximal tubular cell function, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

25 [0080]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance outside the body, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the
5 alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance outside the body, in which one dosage unit
10 (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0081]

In one embodiment, the pharmaceutical composition provided by the present
15 invention is a pharmaceutical composition comprising an alkalinizing agent for use in excreting uremic substance into urine depending on a concentration of uremic substance in blood, in which 0.5 to 1.5 g/day of each of a potassium citrate monohydrate and a sodium citrate dihydrate as the alkalinizing agent are orally administered at a total of 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

20 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in excreting uremic substance into urine depending on a concentration of uremic substance in blood, in which one dosage unit (preferably one tablet) contains 231.5 mg potassium citrate monohydrate and 195.0 mg sodium citrate dihydrate, and 3 to 6 dosage units per
25 day are orally administered in three divided doses per day.

[0082]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in decreasing a concentration of uremic substance in blood, in which sodium bicarbonate as
5 the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in decreasing a concentration of uremic substance in blood, in which one dosage unit
10 (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0083]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
15 promoting excretion of uremic substance into urine, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
20 promoting excretion of uremic substance into urine, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0084]

In one embodiment, the pharmaceutical composition provided by the present
25 invention is a pharmaceutical composition comprising an alkalinizing agent for use in

ameliorating uremic symptoms in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating uremic symptoms in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0085]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing uremia in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0086]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression of chronic kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression of chronic kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0087]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in delaying introduction of dialysis to a patient with chronic kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0088]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing myocardial fibrosis in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing myocardial fibrosis in a patient with kidney disease, in which one dosage

unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0089]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing arteriosclerosis in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing arteriosclerosis in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0090]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular smooth muscle cell proliferation in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0091]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular endothelial cell injury in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing vascular endothelial cell injury in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0092]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing thickening of the arterial wall of a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing thickening of the arterial wall of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0093]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease, in which sodium

bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing calcification of the aorta of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0094]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing cardiovascular disease in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating or preventing cardiovascular disease in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0095]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use ameliorating arteriosclerosis in a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present

invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating arteriosclerosis in a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

5 [0096]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use ameliorating thickening of the arterial wall of a patient with kidney disease, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5
10 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in ameliorating thickening of the arterial wall of a patient with kidney disease, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6
15 dosage units per day are orally administered in three divided doses per day.

[0097]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating acute kidney failure, in which sodium bicarbonate as the alkalinizing agent is
20 orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in treating acute kidney failure, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in
25 three divided doses per day.

[0098]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease, in which
5 sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing progression from acute kidney failure to chronic kidney disease, in which
10 one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0099]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
15 treating or preventing kidney tubular damage, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
20 treating or preventing kidney tubular damage, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0100]

In one embodiment, the pharmaceutical composition provided by the present
25 invention is a pharmaceutical composition comprising an alkalinizing agent for use in

suppressing kidney tubular damage, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in
5 suppressing kidney tubular damage, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0101]

In one embodiment, the pharmaceutical composition provided by the present
10 invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney proximal tubular cell damage, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present
15 invention is a pharmaceutical composition comprising an alkalinizing agent for use in suppressing kidney proximal tubular cell damage, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0102]

20 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in protecting kidney proximal tubular cells, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present
25 invention is a pharmaceutical composition comprising an alkalinizing agent for use in

protecting kidney proximal tubular cells, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0103]

5 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in maintaining kidney proximal tubular cell function, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

10 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in maintaining kidney proximal tubular cell function, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

15 [0104]

 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance outside the body, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day,
20 preferably 3 times a day.

 In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in promoting excretion of uremic substance outside the body, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per
25 day are orally administered in three divided doses per day.

[0105]

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in excreting uremic substance into urine depending on a concentration of uremic substance in blood, in which sodium bicarbonate as the alkalinizing agent is orally administered at 1 to 3 g/day, 1 to 5 times a day, preferably 3 times a day.

In one embodiment, the pharmaceutical composition provided by the present invention is a pharmaceutical composition comprising an alkalinizing agent for use in excreting uremic substance into urine depending on a concentration of uremic substance in blood, in which one dosage unit (preferably one tablet) contains 500 mg sodium bicarbonate, and 3 to 6 dosage units per day are orally administered in three divided doses per day.

[0106]

Hereinafter, the present invention will be further described by examples, but the present invention is not limited thereto.

EXAMPLES

[0107]

A human clinical trial was conducted to examine whether oral administration of a combination preparation of hydrates of potassium citrate and sodium citrate, and a sodium bicarbonate preparation, which are oral alkalinizing agents, promotes excretion of uremic substance into urine.

[0108]

1. Method

47 patients with chronic kidney diseases in stage G2 to G3b (eGFR: 30 to 89 ml/min/1.73m²) were randomly divided into a group to which a combination preparation

of hydrates of potassium citrate and sodium citrate had been administered (a group A: 16 patients), a group to which a sodium bicarbonate preparation (sodium hydrogen carbonate) had been administered (a group B: 16 patients), and a control group (a group C: 15 patients). Patients were assigned to each group so that age, sex, presence of diabetes, and eGFR were not biased. Each group was treated according to the “CKD medical care guide-summary of treatment” (hereinafter referred to as standard treatment).

An alkalinizing agent was not administered to the control group. To the group A, 3 tablets a day in which each tablet contained potassium citrate ($C_6H_5K_3O_7 \cdot H_2O$) 231.5 mg and a sodium citrate hydrate ($C_6H_5Na_3O_7 \cdot 2H_2O$) 195.0 mg were orally administered 3 times a day (morning, noon, evening) for 24 weeks. A pH of early morning urine was controlled over time so that the dosage could be increased up to 6 tablets, 3 times a day (morning, noon, evening) as needed according to the discretion of the physician in cases where a pH of early morning urine was less than pH 6.5. To the group B, 3 tablets a day in which each tablet contained 500 mg of sodium bicarbonate were orally administered 3 times a day (morning, noon, evening) for 24 weeks. A pH of early morning urine was controlled over time so that the dosage could be increased up to 6 tablets, 3 times a day (morning, noon, evening) as needed according to the discretion of the physician in cases where a pH of early morning urine was less than pH 6.5.

Early morning urine and blood were collected before the start of administration, and 6 weeks, 12 weeks, and 24 weeks after the start of administration, and each specimen was stored at $-80^{\circ}C$. Methods used in the present field for indoxyl sulfate, p-cresyl sulfate, phenylacetyl-L-glutamine, hippuric acid, and argininosuccinic acid in urine and blood (Sato, E., et. al., Metabolic alteration by indoxyl sulfate in skeletal muscle induce uremic sarcopenia in chronic kidney disease., Sci Rep. 2016 Nov 10;6:36618. doi:

10.1038/srep36618. and the like) were referred to, and quantitative analysis was performed using the following liquid chromatography triple quadrupole mass spectrometer (LC-MS/MS).

For LC, NANOSPACE SI-2 (manufactured by Shiseido) was used, and
5 CAPCELLPAK MGIII was selected as the analytical column.

For MS, TSQ Quantiva (manufactured by Thermo Fisher Scientific Co., Ltd.) was used, and five compounds were ionized in the negative mode and detected using the Selected Reaction Monitoring method. The quantitative value was calculated using a calibration curve prepared with a standard solution of each compound.

10 In addition, the amount of urinary β 2-microglobulin was measured by latex agglutination method using the LZ test “Eiken” β 2-M and LZ- β 2-M standard U “Eiken” (Eiken Chemical Co., Tokyo, Japan). Furthermore, the amount of cystatin C in the serum was measured by a colloidal gold agglutination method using Nescoat GC Cystatin C (Nm) (Alfresa Pharma, Osaka, Japan).

15 For statistical analysis, the Mann-Whitney test was used for comparison between groups, and the Wilcoxon test was used for comparison of changes over time. The Pearson test was used for correlation.

[0109]

2. Result

20 Based on the measurement results using LC-MS/MS, for respective patients in the group A (the group to which the combination preparation of hydrates of potassium citrate and sodium citrate was administered), the group B (the group to which the sodium bicarbonate preparation was administered), and the group C (the control group), the following was calculated:

25 (i) Concentration of each uremic substance in plasma before administration

(ii) Concentration of each uremic substance in early morning urine before administration

(iii) Ratio of uremia substance concentration in early morning urine and plasma uremic substance concentration before the start of administration (amount of uremic substance in urine/amount of uremic substance in plasma)

(iv) Concentration of each uremic substance in plasma 6 weeks, 12 weeks, and 24 weeks after the start of administration

(v) Concentration of each uremic substance in early morning urine 6 weeks, 12 weeks, and 24 weeks after the start of administration

(vi) Ratio of uremic substance concentration in early morning urine and uremic substance concentration in plasma at 6 weeks, 12 weeks, and 24 weeks after the start of administration (uremic substance amount in urine/uremic substance amount in plasma)

(vii) Amount of change in the concentration of each uremic substance in the plasma 6 weeks, 12 weeks, and 24 weeks after the start of administration from before the start of administration

(viii) Amount of change in the concentration of each uremic substance in the early morning urine 6 weeks, 12 weeks, and 24 weeks, from before the start of administration

(ix) Amount of change in ratio of uremic substance concentration in early morning urine and uremic substance concentration in plasma 6 weeks, 12 weeks, and 24 weeks after the start of administration (amount of uremic substance in urine/amount of uremic substance in plasma), from before the start of administration

Then, the average value and SD of each group were calculated for (i) to (ix) above. For each of (iv) to (ix) above, the average value and SD of each group were calculated for all data of 6 weeks, 12 weeks, and 24 weeks after the start of

administration in each group.

The results are shown in tables below. In the tables and drawings, the group A: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered was described as “Citrate,” and the group B: the
5 group to which the sodium bicarbonate preparation had been administered was described as “Bicarbonate.” In addition, the numerical value in the parenthesis in the tables indicates the number of cases.

Table 1-1-1: Amount (ng/mL) of indoxyl sulfate in plasma

Table 1-1-2: Change in amount (ng/mL) of indoxyl sulfate in plasma from
10 before the start of administration

Table 1-2-1: Amount (ng/mL) of indoxyl sulfate in early morning urine

Table 1-2-2: Change in amount (ng/mL) of indoxyl sulfate in early morning
urine from before the start of administration

Table 1-3-1: Ratio of amount of indoxyl sulfate in urine to an amount of indoxyl
15 sulfate in plasma

Table 1-3-2: Amount of change in ratio of amount of indoxyl sulfate in urine to
amount of indoxyl sulfate in plasma from before the start of administration

Table 2-1-1: Amount (ng/mL) of p-cresyl sulfate in plasma

Table 2-1-2: Change in amount (ng/mL) of p-cresyl sulfate in plasma from
20 before the start of administration

Table 2-2-1: Amount (ng/mL) of p-cresyl sulfate in early morning urine

Table 2-2-2: Change in amount (ng/mL) of p-cresyl sulfate in early morning
urine from before the start of administration

Table 2-3-1: Ratio of amount of p-cresyl sulfate in urine to an amount of p-
25 cresyl sulfate in plasma

Table 2-3-2: Amount of change in ratio of amount of p-cresyl sulfate in urine to amount of p-cresyl sulfate in plasma from before the start of administration

Table 3-1-1: Amount (ng/mL) of hippuric acid in plasma

Table 3-1-2: Change in amount (ng/mL) of hippuric acid in plasma from before
5 the start of administration

Table 3-2-1: Amount (ng/mL) of hippuric acid in early morning urine

Table 3-2-2: Change in amount (ng/mL) of hippuric acid in early morning urine
from before the start of administration

Table 3-3-1: Ratio of amount of hippuric acid in urine to amount of hippuric
10 acid in plasma

Table 3-3-2: Amount of change in ratio of amount of hippuric acid in urine to amount of hippuric acid in plasma from before the start of administration

Table 4-1-1: Amount (ng/mL) of argininosuccinic acid in plasma

Table 4-1-2: Change in amount (ng/mL) of argininosuccinic acid in plasma from
15 before the start of administration

Table 4-2-1: Amount (ng/mL) of argininosuccinic acid in early morning urine

Table 4-2-2: Change in amount (ng/mL) of argininosuccinic acid in early
morning urine from before the start of administration

Table 4-3-1: Ratio of amount of argininosuccinic acid in urine to amount of
20 argininosuccinic acid in plasma

Table 4-3-2: Amount of change in ratio of amount of argininosuccinic acid in urine to amount of argininosuccinic acid in plasma from before the start of administration

Table 5-1-1: Amount (ng/mL) of phenylacetyl-L-glutamine (PAG) in plasma

Table 5-1-2: Change in amount (ng/mL) of phenylacetyl-L-glutamine (PAG) in
25 plasma from before the start of administration

Table 5-2-1: Amount (ng/mL) of phenylacetyl-L-glutamine (PAG) in early morning urine

Table 5-2-2: Change in amount (ng/mL) of phenylacetyl-L-glutamine (PAG) in early morning urine from before the start of administration

5 Table 5-3-1: Ratio of amount of phenylacetyl-L-glutamine (PAG) in urine to amount of phenylacetyl-L-glutamine (PAG) in plasma

Table 5-3-2: Amount of change in ratio of amount of phenylacetyl-L-glutamine (PAG) in urine to amount of phenylacetyl-L-glutamine (PAG) in plasma from before the start of administration

10 [0110]

In the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), an indoxyl sulfate concentration in plasma 6, 12, and 24 weeks after the administration was a lower value, compared to those of the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered) and the group C (Control: the control group) (refer to Table 1-1-1). In addition, in the group A, an indoxyl sulfate (IS) concentration in early morning urine 12 and 24 weeks after administration was a higher value than that in the group C (refer to Table 1-2-1). The indoxyl sulfate concentration in plasma at 6 to 24 weeks was a significantly lower value in the group A than those in the groups B and C (refer to Table 1-1-1), and the indoxyl sulfate concentration in early morning urine at 6 to 24 weeks was significantly greater in the group A than those in the groups B and C (refer to Table 1-2-2).

In addition, by administering the combination preparation of hydrates of potassium citrate and sodium citrate to patients with chronic kidney disease, the concentration of indoxyl sulfate, which is a uremic substance, in urine increased

compared to before administration, and the concentration of indoxyl sulfate in blood decreased compared to before administration. Even with the same alkalinizing agent, such an effect was not recognized in the case of the sodium bicarbonate preparation. Compared with the sodium bicarbonate preparation, the combination preparation of
5 hydrates of potassium citrate and sodium citrate exhibited an excellent effect of decreasing an indoxyl sulfate concentration in blood, and an excellent effect of increasing an indoxyl sulfate concentration in urine. The effect of decreasing the indoxyl sulfate concentration in blood and the effect of increasing the indoxyl sulfate concentration in urine by the combination preparation of hydrates of potassium citrate and sodium citrate
10 were recognized from 12 weeks after the administration.

Based on the value of the ratio of indoxyl sulfate concentration in urine to indoxyl sulfate concentration in plasma, it was shown that administration of the combination preparation of hydrates of potassium citrate and sodium citrate led to the excretion of indoxyl sulfate from blood into urine, and thereby excretion thereof outside
15 the body was promoted. The effect of excretion of indoxyl sulfate from blood into urine was recognized by administration of the combination preparation of hydrates of potassium citrate and sodium citrate, but not by administration of the sodium bicarbonate preparation (refer to Table 1-3-1 and Table 1-3-2).

[0111]

20 [Table 1-1-1]

Plasma Indoxyl Sulfate

Group	N	0W	6W	12W	24W	6-24W (45-47)
Control	15	1189 ± 634	1229 ± 710	1326 ± 793	1134 ± 371	1230 ± 641
Citrate	16	1130 ± 1157	1170 ± 1056	1105 ± 1213 ^b	1050 ± 762	1108 ± 770 ^{a,c}
Bicarbonate	16	1338 ± 770	1365 ± 821	1432 ± 1031	1481 ± 988	1426 ± 930
Mean ± SD						

^ap=0.0288, ^bp=0.0451 vs Control and ^cp=0.0350 vs Bicarbonate (Mann-Whitney)

[0112]

[Table 1-1-2]

Plasma Indoxyl Sulfate (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-47)
Control	15	39.5 ± 446.4	136.5 ± 468.1	-55.8 ± 564.2	40.1 ± 490.6
Citrate	16	40.0 ± 460.7	-36.8 ± 380.8	-80.3 ± 583.7	-25.5 ± 487.6
Bicarbonate	16	27.5 ± 523.0	153.1 ± 887.4	143.4 ± 682.5	107.0 ± 695.8
Mean ± SD					

5

[0113]

[Table 1-2-1]

Urine Indoxyl Sulfate

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	41026 ± 26159	34608 ± 23011	30959 ± 15288	29139 ± 16714 ^a	31568 ± 18358
Citrate	16	41081 ± 45508 ^a	32728 ± 19842	58648 ± 96652	46808 ± 43377	46061 ± 64823
Bicarbonate	16	54379 ± 40259	52510 ± 39229	37860 ± 26401	56482 ± 39593	48954 ± 35769 ^a
Mean ± SD						

^ap=0.0073 vs Control and ^ap=0.0385 vs Bicarbonate (Mann-Whitney)

^ap=0.0103 vs 0 week (Wilcoxon)

[0114]

[Table 1-2-2]

Urine Indoxyl Sulfate (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-6419 $\pm$ 26152	-10067 $\pm$ 26839	-11887 $\pm$ 21994	-9458 $\pm$ 24617
Citrate	16	-8352 $\pm$ 41966	17566 $\pm$ 64586 ^a	5727 $\pm$ 36794	4980 $\pm$ 49397 ^{a,d}
Bicarbonate	16	-1869 $\pm$ 19390	-16519 $\pm$ 40380	2113 $\pm$ 49356	-5425 $\pm$ 38514
Mean $\pm$ SD					

^ap=0.0141 vs Control and ^bp=0.0233, ^cp=0.0438 vs Bicarbonate (Mann-Whitney)

[0115]

[Table 1-3-1]

Indoxyl Sulfate - Urine/Plasma ratio

Group	N	0W	6W	12W	24W	6-24W (45-47)
Control	15	39.2 ± 21.1	31.8 ± 21.2 ^a	27.7 ± 16.1 ^c	26.4 ± 13.8 ^e	28.6 ± 17.1
Citrate	16	37.5 ± 24.7	38.6 ± 31.8	51.5 ± 50.6	52.4 ± 64.4 ^b	47.4 ± 50.0
Bicarbonate	16	55.5 ± 43.6	51.7 ± 36.4	35.3 ± 26.4	50.7 ± 36.5 ^c	46.2 ± 33.7 ^d
Mean ± SD						

^ap=0.0482, ^bp=0.0322 and ^cp=0.0141 vs Control (Mann-Whitney)^dp=0.0478, ^ep=0.0413 and ^fp=0.0151 vs 0 week (Wilcoxon)

5

[0116]

[Table 1-3-2]

Indoxyl Sulfate - Urine/Plasma ratio (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-47)
Control	15	-7.41 $\pm$ 24.69	-11.52 $\pm$ 9.87	-12.79 $\pm$ 16.00	-10.57 $\pm$ 20.16
Citrate	16	1.10 $\pm$ 22.99	10.70 $\pm$ 39.08 ^d	14.91 $\pm$ 56.11 ^b	8.90 $\pm$ 40.99 ^{c,e}
Bicarbonate	16	-3.81 $\pm$ 37.48	-22.41 $\pm$ 35.49	-4.78 $\pm$ 35.33	-10.33 $\pm$ 36.35
Mean $\pm$ SD					

^ap=0.0239, ^bp=0.0006 vs Control and ^cp=0.0457, ^dp=0.0292 vs Bicarbonate (Mann-Whitney)

[0117]

Regarding p-cresyl sulfate (PCS), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the concentration of p-cresyl sulfate in the early morning urine 12 and 24

weeks after administration was a higher value than that of the group C (Control: the control group). In the group A (the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), even when compared to the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), the p-cresyl sulfate concentration in the early morning urine at 6, 12, and 24 weeks after administration was a higher value (refer to Table 2-2-1). An increase in p-cresyl sulfate concentration in early morning urine at 6 to 24 weeks was recognized only in the group A (refer to Table 2-2-2).

In addition, administration of the combination preparation of hydrates of potassium citrate and sodium citrate to patients with chronic kidney diseases increased the concentration of p-cresyl sulfate, which is a uremic substance, in urine compared to before administration (refer to Table 2-2-1 and Table 2-2-2). Even with the same alkalinizing agent, such an effect was not recognized in the case of the sodium bicarbonate preparation, and the combination preparation of hydrates of potassium citrate and sodium citrate exhibited a stronger effect of increasing a concentration of p-cresyl sulfate in urine as compared to the sodium bicarbonate preparation. The effect of increasing the p-cresyl sulfate concentration in urine by the combination preparation of hydrates of potassium citrate and sodium citrate was recognized from 12 weeks after administration.

On the other hand, administration of the sodium bicarbonate preparation decreased the concentration of p-cresyl sulfate, which is a uremic substance, in plasma as compared to before administration (refer to Table 2-1-2). Such an effect of decreasing a concentration of p-cresyl sulfate in plasma was more strongly recognized in the group B as compared to the group A and group C (refer to Table 2-1-2).

Based on the value of the ratio of p-cresyl sulfate concentration in urine to p-

cresyl sulfate concentration in plasma, it was shown that administration of the combination preparation of hydrates of potassium citrate and sodium citrate led to the excretion of p-cresyl sulfate from blood into urine, and thereby excretion thereof outside the body was promoted. It was shown that the effect of excretion of p-cresyl sulfate from blood into urine was more strongly recognized by administration of the combination preparation of hydrates of potassium citrate and sodium citrate than by administration of the sodium bicarbonate preparation (refer to Table 2-3-1 and Table 2-3-2).

[0118]

[Table 2-1-1]

Plasma *p*-Cresyl Sulfate

Group	N	0W	6W	12W	24W	6-24W (44-48)
Control	15	4467 ± 3531	5289 ± 4389	4482 ± 4570	3852 ± 3454	4524 ± 4106
Citrate	16	3491 ± 5159	3414 ± 3432	3306 ± 4140	3673 ± 4853	3468 ± 4094
Bicarbonate	16	5402 ± 5209	4506 ± 5300	3771 ± 4139	4284 ± 3682	4187 ± 4354
Mean ± SD						

[0119]

[Table 2-1-2]

Plasma *p*-Cresyl Sulfate (Conversion to *Δ* actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-47)
Control	15	536.1 ± 2801	14.3 ± 3291	-615.2 ± 2280	-34.3 ± 2796
Citrate	16	-77.2 ± 2341	-6.8 ± 2982	181.8 ± 2283	33.4 ± 2491 ^c
Bicarbonate	16	-895.9 ± 2153	-1103 ± 2478	-1119 ± 3631	-1038 ± 2773
Mean ± SD					

^cp=0.0377 vs Bicarbonate (Mann-Whitney)

[0120]

[Table 2-2-1]

126

Urine *p*-Cresyl Sulfate

Group	N	0W	6W	12W	24W	6-24W (44-48)
Control	15	58782 ± 59346	60188 ± 72317	39257 ± 57116	34669 ± 51832	44353 ± 60343
Citrate	16	39252 ± 53214	38343 ± 26834	42970 ± 43893	55979 ± 68542	45764 ± 49000
Bicarbonate	16	66164 ± 97704	67744 ± 105548	37730 ± 48411	52029 ± 57779	52501 ± 73583
Mean ± SD						

[0121]

[Table 2-2-2]

Urine *p*-Cresyl Sulfate (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	-2403 ± 56332	-19525 ± 50834	-24113 ± 47778	-15641 ± 51296
Citrate	16	-909 ± 48618	3718 ± 38763	16727 ± 42639 ^a	6512 ± 43256 ^{b,c}
Bicarbonate	16	1580 ± 51468	-28434 ± 72253	-14163 ± 95244	-13663 ± 74566
Mean ± SD					

^ap=0.0170, ^bp=0.0109 vs Control and ^cp=0.0242 vs Bicarbonate (Mann-Whitney)

5

[0122]

[Table 2-3-1]

p-Cresyl Sulfate - Urine/Plasma ratio

Group	N	0W	6W	12W	24W	6-24W (44-48)
Control	15	11.9 ± 9.4	10.9 ± 4.9	9.8 ± 5.9	9.3 ± 8.2	10.0 ± 5.6
Citrate	16	14.0 ± 9.2	16.1 ± 13.7	16.3 ± 12.9	20.5 ± 21.5 ^a	17.7 ± 16.3 ^b
Bicarbonate	16	15.6 ± 11.1	17.0 ± 12.0	16.5 ± 17.9	17.0 ± 11.8 ^a	16.8 ± 13.9 ^d
Mean ± SD						

^ap=0.0221, ^bp=0.0094, ^cp=0.0364 and ^dp=0.0055 vs Control (Mann-Whitney)

[0123]

[Table 2-3-2]

***p*-Cresyl Sulfate - Urine/Plasma ratio** (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	-1.08 $\pm$ 10.03	-2.14 $\pm$ 9.15	-2.59 $\pm$ 9.68	-1.96 $\pm$ 9.41
Citrate	16	2.08 $\pm$ 11.30	1.22 $\pm$ 11.94	6.51 $\pm$ 17.93	3.21 $\pm$ 13.94
Bicarbonate	16	1.39 $\pm$ 8.25	0.91 $\pm$ 15.68	1.31 $\pm$ 9.81	1.20 $\pm$ 11.45
Mean $\pm$ SD					

[0124]

Regarding hippuric acid (HA), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), a hippuric acid concentration in plasma 24 weeks after the administration was a lower value, compared to those of the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered) and the group C (Control: the control group) (refer to Table 3-1-1). Such an effect was not seen in the administration of the sodium bicarbonate preparation. In addition, in the group A, the hippuric acid concentration in the early morning urine 12 and 24 weeks after administration was a higher value than that in the group C (refer to Table 3-2-1).

In addition, by administering the combination preparation of hydrates of potassium citrate and sodium citrate to patients with chronic kidney diseases, the concentration of hippuric acid, which is a uremic substance, in plasma 24 weeks after administration was decreased as compared to that before administration (refer to Tables 3-1-1 and 3-2-2), and the concentration of hippuric acid in urine increased compared to that before administration (refer to Tables 3-2-1 and 3-2-2). An increase in hippuric acid concentration in the early morning urine at 6 to 24 weeks was recognized only in the group A (refer to Table 2-2-2). Even with the same alkalinizing agent, compared with the sodium bicarbonate preparation, the combination preparation of hydrates of

potassium citrate and sodium citrate exhibited an excellent effect of increasing a hippuric acid concentration in urine. The effect of increasing the hippuric acid concentration in urine by the combination preparation of hydrates of potassium citrate and sodium citrate was recognized from 6 weeks after administration.

- 5 Based on the value of the ratio of hippuric acid concentration in urine to hippuric acid concentration in plasma, it was shown that administration of the combination preparation of hydrates of potassium citrate and sodium citrate led to the excretion of hippuric acid from blood into urine, and thereby excretion thereof outside the body was promoted. In addition, it was shown that the effect of excretion of
- 10 hippuric acid from blood into urine at 6 to 24 weeks was more strongly recognized by administration of the combination preparation of hydrates of potassium citrate and sodium citrate than by administration of the sodium bicarbonate preparation (refer to Table 3-3-1 and Table 3-3-2).

[0125]

- 15 [Table 3-1-1]

Plasma Hippuric Acid									
Group	N	0W		6W		12W		24W	6-24W (45-47)
Control	15	1096 ±	828	752 ±	192 ^b	975 ±	615	818 ± 615	848 ± 572
Citrate	16	868 ±	1015	1464 ±	2213	985 ±	1376	520 ± 437	990 ± 1546
Bicarbonate	16	742 ±	706	809 ±	496	994 ±	806	1465 ± 1387 ^c	1091 ± 991
Mean ± SD									
^b p=0.0181 and ^c p=0.0215 vs 0 week (Wilcoxon)									

[0126]

[Table 3-1-2]

Plasma Hippuric Acid (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-47)
Control	15	-343.6 $\pm$ 620.1	-120.6 $\pm$ 700.9	-227.9 $\pm$ 893.8	-247.3 $\pm$ 736.1
Citrate	16	596.5 $\pm$ 2370 ^a	153.7 $\pm$ 1671	-347.1 $\pm$ 1080 ^a	133.7 $\pm$ 1794
Bicarbonate	16	66.6 $\pm$ 681.8	230.0 $\pm$ 655.6	722.3 $\pm$ 1023 ^a	342.1 $\pm$ 838.5 ^a
Mean $\pm$ SD					

^ap=0.0430, ^ap=0.0164, ^ap=0.0008 vs Control
and ^ap=0.0093 vs Bicarbonate (Mann-Whitney)

[0127]

[Table 3-2-1]

Urine Hippuric Acid

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	128966 ± 110600	163650 ± 81018	86202 ± 43032	88804 ± 43032	94885 ± 64566
Citrate	16	94469 ± 57885	112946 ± 80846	112486 ± 86932	99287 ± 70822	108240 ± 78356
Bicarbonate	16	109249 ± 96133	98283 ± 80730	96346 ± 70315	114453 ± 81562	103027 ± 76475
Mean ± SD						

5

[0128]

[Table 3-2-2]

Urine Hippuric Acid (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	-19316 $\pm$ 96398	-42764 $\pm$ 105143	-40162 $\pm$ 85472	-34080 $\pm$ 94400
Citrate	16	18478 $\pm$ 89920	18018 $\pm$ 62771	4818 $\pm$ 81234	13771 $\pm$ 77363 ^a
Bicarbonate	16	-10966 $\pm$ 75133	-12903 $\pm$ 61236	5204 $\pm$ 85546	-6221 $\pm$ 73493
Mean $\pm$ SD					

^ap=0.0431 vs Control (Mann-Whitney)

[0129]

[Table 3-3-1]

130

Hippuric Acid - Urine/Plasma ratio

Group	N	0W	6W	12W	24W	6-24W (41-47)
Control	15	139.0 ± 89.6	146.2 ± 70.2	119.3 ± 75.1	134.4 ± 100.5	133.3 ± 81.9
Citrate	16	316.8 ± 374.8	160.8 ± 179.9	275.1 ± 315.5	430.1 ± 619.6 ^{b,c}	288.9 ± 61.7 ^a
Bicarbonate	16	304.5 ± 361.8	148.7 ± 119.2 ^f	151.8 ± 115.8	523.0 ± 1438	265.8 ± 812.9
Mean ± SD						

^bp=0.0239 vs Control and ^cp=0.0355, ^fp=0.0478 vs Bicarbonate (Mann-Whitney)

^fp=0.0479 vs 0 week (Wilcoxon)

[0130]

[Table 3-3-2]

Hippuric Acid - Urine/Plasma ratio (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	7.2 ± 73.2	-19.1 ± 83.1	-4.6 ± 82.5	-5.7 ± 78.7
Citrate	16	-156.0 ± 362.1	-60.2 ± 475.5	113.3 ± 588.8	-33.7 ± 487.4
Bicarbonate	15	-155.8 ± 312.2 ^a	-161.2 ± 302.1	191.1 ± 1147	-50.2 ± 688.3 ^b
Mean ± SD					

^ap=0.0451, ^bp=0.0170 vs Control (Mann-Whitney)

5 [0131]

Regarding argininosuccinic acid (ASA), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), a concentration of argininosuccinic acid in the early morning urine was a higher value than that of the group C (Control: the control group), whereas the value was lower than that of the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered) (refer to Table 4-2-1).

In addition, administration of the combination preparation of hydrates of potassium citrate and sodium citrate to patients with chronic kidney diseases increased the concentration of argininosuccinic acid, which is a uremic substance, in urine compared to before administration (refer to Table 4-2-1 and Table 4-2-2). The effect of

increasing the argininosuccinic acid concentration in urine by the combination preparation of hydrates of potassium citrate and sodium citrate was recognized from 12 weeks after administration. The increase in argininosuccinic acid concentration in early morning urine at 6 to 24 weeks was more increased in the group B than in the group A (refer to Table 4-2-2).

Based on the value of the ratio of an argininosuccinic acid concentration in urine to an argininosuccinic acid concentration in plasma, it was shown that administration of the combination preparation of hydrates of potassium citrate and sodium citrate led to the excretion of argininosuccinic acid from blood into urine, and thereby excretion thereof outside the body was promoted. In addition, it was shown that the effect of excretion of argininosuccinic acid from blood into urine at 6 to 24 weeks was more strongly recognized by administration of the sodium bicarbonate preparation than that by administration of the combination preparation of hydrates of potassium citrate and sodium citrate (refer to Table 4-3-1 and Table 4-3-2).

[0132]

[Table 4-1-1]

Plasma Arginino Succinic Acid						
Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	1973 ± 3343	1493 ± 1193	1464 ± 1211	1142 ± 692	1366 ± 1047
Citrate	16	1286 ± 789	1241 ± 847	1472 ± 915	1404 ± 887	1370 ± 869
Bicarbonate	16	1178 ± 655	1245 ± 748	1576 ± 2160	1160 ± 977	1327 ± 1416
Mean ± SD						

[0133]

[Table 4-1-2]

Plasma Arginino Succinic Acid (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-479.8 $\pm$ 3374	-508.4 $\pm$ 3352	-830.6 $\pm$ 3349	-606.3 $\pm$ 3285
Citrate	16	-47.0 $\pm$ 723.7	135.8 $\pm$ 710.2	116.4 $\pm$ 666.6	66.9 $\pm$ 690.0
Bicarbonate	16	66.5 $\pm$ 565.5	397.7 $\pm$ 1733	-17.6 $\pm$ 776.9	148.9 $\pm$ 1134
Mean $\pm$ SD					

[0134]

[Table 4-2-1]

Urine Arginino Succinic Acid

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	2861 $\pm$ 1563	2726 $\pm$ 1757	2289 $\pm$ 1381	1977 $\pm$ 1076 ^b	2331 $\pm$ 1433
Citrate	16	2483 $\pm$ 1649	2399 $\pm$ 979	2625 $\pm$ 1485	2796 $\pm$ 1526	2607 $\pm$ 1333
Bicarbonate	16	2911 $\pm$ 1691	3346 $\pm$ 2636	2647 $\pm$ 2123	3221 $\pm$ 2466	3071 $\pm$ 2386
Mean $\pm$ SD						

^bp=0.0215 vs 0 week (Wilcoxon)

5

[0135]

[Table 4-2-2]

Urine Arginino Succinic Acid (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	-110.7 $\pm$ 1578	-571.8 $\pm$ 1159	-884.0 $\pm$ 1363	-531.5 $\pm$ 1377
Citrate	16	-84.7 $\pm$ 1617	142.0 $\pm$ 1785	312.1 $\pm$ 1591 ^a	123.1 $\pm$ 1639 ^b
Bicarbonate	16	434.7 $\pm$ 1299	-264.1 $\pm$ 1491	309.9 $\pm$ 2335	160.2 $\pm$ 1756
Mean $\pm$ SD					

^ap=0.0105 and ^bp=0.0175 vs Control (Mann-Whitney)

[0136]

[Table 4-3-1]

Arginino Succinic Acid ~ Urine/Plasma ratio

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	3.17 ± 2.57	3.67 ± 3.69	3.28 ± 3.47	2.16 ± 1.35	3.04 ± 3.02
Citrate	16	2.39 ± 1.60	2.92 ± 2.31	2.87 ± 3.27	3.24 ± 3.17	3.01 ± 2.88
Bicarbonate	16	3.82 ± 4.93	5.75 ± 9.80	2.81 ± 1.86	6.36 ± 9.60	4.97 ± 7.98
Mean ± SD						

[0137]

[Table 4-3-2]

ASA - Urine/Plasma ratio (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (44-48)
Control	15	0.49 ± 3.25	0.11 ± 3.74	-1.01 ± 2.43	-0.14 ± 3.18
Citrate	16	0.53 ± 2.61	0.49 ± 3.55	0.84 ± 2.90	0.62 ± 2.97
Bicarbonate	16	1.93 ± 7.90	-1.01 ± 4.47	2.54 ± 10.20*	1.15 ± 7.87
Mean ± SD					

*p=0.0403 vs Control (Mann-Whitney)

5 [0138]

Regarding phenylacetyl-L-glutamine (PAG), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), a phenylacetyl-L-glutamine concentration in plasma was a lower value compared to that of the group C (Control: the control group) (refer to Table 5-1-1).

10 In the group A, a phenylacetyl-L-glutamine concentration in early morning urine 12 and 24 weeks after administration was a higher value than that in the group C (refer to Table 5-2-1).

15 In addition, administration of the combination preparation of hydrates of potassium citrate and sodium citrate to patients with chronic kidney diseases increased the concentration of phenylacetyl-L-glutamine, which is a uremic substance, in urine 12 and 24 weeks after administration, compared to before administration (refer to Table 5-2-

1 and Table 5-2-2). Even with the same alkalinizing agent, such an effect was not
recognized in the case of the sodium bicarbonate preparation. The effect of increasing
the phenylacetyl-L-glutamine concentration in urine by the combination preparation of
hydrates of potassium citrate and sodium citrate was recognized from 12 weeks after
5 administration. Administration of the combination preparation of hydrates of potassium
citrate and sodium citrate to patients with chronic kidney diseases decreased the
concentration of phenylacetyl-L-glutamine, which is a uremic substance, in plasma
compared to before administration (refer to Table 5-1-1 and Table 5-2-2). The effect of
decreasing a phenylacetyl-L-glutamine concentration in plasma was strongly recognized
10 by the sodium bicarbonate preparation than that by the combination preparation of
hydrates of potassium citrate and sodium citrate (refer to Table 5-1-2).

Based on the value of the ratio of phenylacetyl-L-glutamine concentration in
urine to phenylacetyl-L-glutamine concentration in plasma, it was shown that
administration of the combination preparation of hydrates of potassium citrate and
15 sodium citrate led to the excretion of phenylacetyl-L-glutamine from blood into urine,
and thereby excretion thereof outside the body was promoted. In addition, it was shown
that the effect of excretion of phenylacetyl-L-glutamine from blood into urine was more
strongly recognized by administration of the combination preparation of hydrates of
potassium citrate and sodium citrate than that by administration of the sodium
20 bicarbonate preparation (refer to Table 5-3-1 and Table 5-3-2).

[0139]

[Table 5-1-1]

135

Plasma Phenyl Acetyl L-Glutamine (PAG)

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	1121 ± 534	1144 ± 644	1138 ± 623	1088 ± 495	1123 ± 575
Citrate	16	919 ± 962 ^a	818 ± 523	934 ± 843	907 ± 772	885 ± 709 ^c
Bicarbonate	16	1324 ± 949	1240 ± 1062	1153 ± 1000	1284 ± 945	1226 ± 983
Mean ± SD						

^ap=0.0326 and ^cp=0.0065 vs Control (Mann-Whitney)

[0140]

[Table 5-1-2]

Plasma PAG (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	22.9 ± 521.1	17.1 ± 445.3	-32.7 ± 468.1	2.4 ± 468.9
Citrate	16	-101.4 ± 666.0	45.6 ± 521.2	-12.8 ± 578.6	-24.4 ± 583.2
Bicarbonate	16	-84.0 ± 636.1	-171.6 ± 896.9	-40.3 ± 514.7	-98.6 ± 688.1
Mean ± SD					

5

[0141]

[Table 5-2-1]

Urine Phenyl Acetyl L-Glutamine (PAG)

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	94801 ± 62107	86419 ± 91066	76897 ± 68238	60028 ± 90932 ^a	74115 ± 73634
Citrate	16	73688 ± 76268	73505 ± 49199	85591 ± 104730	91951 ± 106696	93682 ± 89253
Bicarbonate	16	124676 ± 149757	144753 ± 246773	82478 ± 112943 ^b	100902 ± 91424	109378 ± 163923
Mean ± SD						

^ap=0.0151, ^bp=0.0335 vs 0 week (Wilcoxon)

[0142]

[Table 5-2-2]

136

Urine PAG (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-8382 $\pm$ 71407	-18904 $\pm$ 61477	-34773 $\pm$ 49511	-20687 $\pm$ 61034
Citrate	16	-183 $\pm$ 55548	11902 $\pm$ 57092 ^a	18263 $\pm$ 60993 ^b	9994 $\pm$ 57168 ^{a,b}
Bicarbonate	16	20077 $\pm$ 118811	-42198 $\pm$ 85069	-23774 $\pm$ 117957	-13663 $\pm$ 74566
Mean $\pm$ SD					

^ap=0.0105, ^bp=0.0062 vs Control and ^ap=0.0287, ^bp=0.0282 vs Bicarbonate (Mann-Whitney)

[0143]

[Table 5-3-1]

Phenyl Acetyl L-Glutamine (PAG) - Urine/Plasma ratio

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	97.4 ± 88.9	71.1 ± 43.6	62.3 ± 30.4	53.0 ± 33.5 ^a	62.2 ± 36.2
Citrate	16	95.0 ± 52.9	105.4 ± 74.6	99.1 ± 88.6	108.5 ± 94.5 ^a	104.4 ± 84.4 ^b
Bicarbonate	16	102.2 ± 96.5	93.7 ± 84.0	70.4 ± 46.8	97.9 ± 116.4	87.3 ± 86.2
Mean ± SD						

^ap=0.0358 and ^bp=0.0107 vs Control (Mann-Whitney)^ap=0.0103 vs 0 week (Wilcoxon)

5

[0144]

[Table 5-3-2]

PAG - Urine/Plasma ratio (Conversion to Δ actual measurement value from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-26.3 $\pm$ 82.0	-35.1 $\pm$ 76.6	-44.4 $\pm$ 74.7	-35.2 $\pm$ 76.4
Citrate	16	10.4 $\pm$ 58.2	1.4 $\pm$ 81.4	13.6 $\pm$ 77.5	8.6 $\pm$ 71.5 ^b
Bicarbonate	16	-8.5 $\pm$ 57.8	-31.8 $\pm$ 81.8	-4.3 $\pm$ 133.9	-14.9 $\pm$ 95.3
Mean $\pm$ SD					

^bp=0.0194 vs Control (Mann-Whitney)

[0145]

Effects of potassium citrate/sodium citrate hydrate (Citrate) and sodium

10 bicarbonate (Bicarbonate) on a concentration of each uremic substance in plasma, a

concentration of each uremic substance in early morning urine, and a ratio of a concentration of uremic substance in early morning urine to a concentration of uremic substance in plasma are summarized in the following tables. In the tables below, the sodium bicarbonate preparation (Bicarbonate) is a reference drug for the case of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate), and the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) is a reference drug for the case of the sodium bicarbonate preparation (Bicarbonate). When the effect is significantly superior to the control group or the reference drug group, O is described, when the effect is significantly inferior to the control group or the reference drug group, X is described, and when there is no significant difference, - is described. The effect of the group to which the sodium bicarbonate preparation (Bicarbonate) had been administered with respect to the control group for the indoxyl sulfate concentration in the early morning urine is indicated as (O) in the tables below. This is because in the group to which the sodium bicarbonate preparation (Bicarbonate) had been administered, the indoxyl sulfate concentration in the early morning urine significantly increased compared to that of the control group, but the indoxyl sulfate concentration in the early morning urine after administration of the sodium bicarbonate preparation (Bicarbonate) decreased compared to before the start of the administration, and thus whether or not there is an effect of promoting excretion of indoxyl sulfate into the urine cannot be determined.

Based on the tables below, the effect of decreasing the concentration of uremic substance in blood (plasma) by the alkalinizing agent was clearly recognized in the cases of indoxyl sulfate (IS) and phenylacetyl-L-glutamine (PAG). Among them, for the case of indoxyl sulfate (IS), administration of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) significantly decreased the concentration of

uremic substance in blood (plasma) compared to administration of the sodium bicarbonate preparation (Bicarbonate).

In addition, an effect of increasing a uremic substance concentration in urine by the alkalinizing agent (an effect of excreting uremic toxin substances into urine) is clearly
5 recognized by indoxyl sulfate (IS), p-cresyl sulfate (PCS), hippuric acid (HA), argininosuccinic acid (ASA), and phenylacetyl-L-glutamine (PAG). Among them, in the cases of indoxyl sulfate (IS), p-cresyl sulfate (PCS), and phenylacetyl-L-glutamine (PAG), administration of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrates) significantly increased the concentration of uremic substance
10 in urine (excretion into urine) than that by administration of the sodium bicarbonate preparation (Bicarbonate).

The excretion of uremic substance from the blood into the urine by the alkalinizing agent (an effect of excretion outside of the body) is clearly recognized in the cases of indoxyl sulfate (IS), p-cresyl sulfate (PCS), and phenylacetyl-L-glutamine
15 (PAG). Among them, for the case of indoxyl sulfate (IS), administration of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrates) significantly increased excretion of uremic substance from the blood into the urine (excretion outside of the body) compared to administration of the sodium bicarbonate preparation (Bicarbonate).

[0146]

[Table 6]

Influence of alkalinizing agent on concentration of uremic substance in plasma and in urine
 – Comparison between Cont and reference drug –

Drug	Sample	IS		PCS		HA		ASA		PAG	
		Cont	Reference drug	Cont	Reference drug	Cont	Reference drug	Cont	Reference drug	Cont	Reference drug
Citrate	Plasma	O	O	-	X	-	-	-	-	O	-
	Urine	O	O	O	O	O	-	O	-	O	O
	U/P ratio	O	O	O	-	-	-	-	-	O	-
Bicarbonate	Plasma	-	X	-	O	X	-	-	-	-	-
	Urine	(O)	X	-	X	-	-	-	-	-	X
	U/P ratio	O	X	O	-	X	-	-	-	-	-

Preferable results: Plasma ↓, Urine ↑, U/P ratio ↑

Excellent effects: O, Poor effects: X, No significant difference: -

5

[0147]

Based on the above table, it can be generally understood that the combination preparation of hydrates of potassium citrate and sodium citrate exhibits a greater effect of excreting the uremic substance outside the body compared to the sodium bicarbonate preparation. In addition, administration of the alkalinizing agent to patients with stage G2 chronic kidney disease as well as stage G3b can suppress progression of chronic kidney disease, and it was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate further suppresses the progression of chronic kidney disease than the sodium bicarbonate preparation.

10

[0148]

15

The results of measuring the amount of urinary β 2-microglobulin and the amount of cystatin C in serum are shown below.

[0149]

[Table 7-1]

140

Urine β_2 -microglobulin ($\mu\text{g/L}$)						
Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	16	100.8 $\pm$ 83.2	145.8 $\pm$ 163.2	159.8 $\pm$ 167.3	137.2 $\pm$ 366.8	164.3 $\pm$ 245.9
Citrate	16	100.7 $\pm$ 81.2	93.1 $\pm$ 46.9	88.7 $\pm$ 65.3 ^a	154.4 $\pm$ 181.8	112.6 $\pm$ 117.1 ^b
Bicarbonate	16	99.0 $\pm$ 62.8	216.3 $\pm$ 223.2 ^a	264.5 $\pm$ 301.9	261.0 $\pm$ 301.6 ^{a,f}	208.2 $\pm$ 255.8 ^a
Mean $\pm$ SD						
^a p=0.0338, ^c p=0.0220 vs Control and ^b p=0.0077, ^d p=0.0186 vs Bicarbonate (Mann-Whitney)						
^e p=0.0110 and ^f p=0.0063 vs 0 week (Wilcoxon)						
(No significant difference between groups at week 0)						

[0150]

[Table 7-2]

Urine β_2 -microglobulin (Conversion to $\Delta \mu\text{g/L}$ from week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	45.0 $\pm$ 153.3	59.1 $\pm$ 161.7	86.4 $\pm$ 361.8	63.5 $\pm$ 240.3
Citrate	16	-14.6 $\pm$ 59.3 ^a	-22.2 $\pm$ 98.7 ^a	46.7 $\pm$ 189.9	3.9 $\pm$ 129.8 ^a
Bicarbonate	16	117.3 $\pm$ 220.7	165.5 $\pm$ 282.0	162.0 $\pm$ 281.8 ^c	148.3 $\pm$ 258.4 ^b
Mean $\pm$ SD					
^a p=0.0095, ^b p=0.0415, ^c p=0.0295 vs Bicarbonate and ^d p=0.0002, ^e p=0.0437 vs Control (Mann-Whitney)					
(No significant difference between groups at week 0)					

5

[0151]

[Table 8]

Plasma Cystatine C (mg/L)						
Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	1.209 $\pm$ 0.397	1.191 $\pm$ 0.423	1.155 $\pm$ 0.372 ^a	1.173 $\pm$ 0.339	1.173 $\pm$ 0.371
Citrate	16	1.040 $\pm$ 0.262	1.069 $\pm$ 0.223	1.065 $\pm$ 0.254	1.089 $\pm$ 0.290	1.075 $\pm$ 0.263
Bicarbonate	16	1.165 $\pm$ 0.296	1.171 $\pm$ 0.281	1.161 $\pm$ 0.296	1.141 $\pm$ 0.275	1.154 $\pm$ 0.278
Mean $\pm$ SD						
Not Significant between Groups (Mann-Whitney)						
^a p=0.0303 vs 0 week (Wilcoxon)						
(No significant difference between groups at week 0)						

[0152]

The concentration of cystatin C in the plasma was not different between the

group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and the group C (Control: the control group), and there was no influence on the function of glomeruli by the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered, and the group to which the sodium bicarbonate preparation had been administered (Table 8).

On the other hand, regarding a urinary β 2-microglobulin concentration, in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), a urinary β 2-microglobulin concentration was a lower value compared to that of the group C (Control: the control group), and even in comparison with the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), a urinary β 2-microglobulin concentration of the group A (the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered) was a lower value. The group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered) had a higher urinary β 2-microglobulin concentration than that of the group C (Control: the control group). Administration of the combination preparation of hydrates of potassium citrate and sodium citrate suppresses an increase in urinary β 2-microglobulin concentration associated with progression of the stage, and it was recognized that there is no change in the urinary β 2-microglobulin concentration compared to before administration.

Based on these results, it was shown that administration of the combination preparation of hydrates of potassium citrate and sodium citrate suppresses kidney tubular damage (kidney proximal tubular damage) associated with progression of the stage. In

addition, it was shown that administration of the sodium bicarbonate preparation did not suppress kidney tubular damage (kidney proximal tubular damage) associated with progression of the stage, and it rather worsened the progression. These effects were recognized from 6 weeks after administration.

5 In addition, an effect of increasing a concentration of uremic substance in urine and an effect of decreasing a concentration of uremic substance in blood by administration of the combination preparation of hydrates of potassium citrate and sodium citrate was not recognized to have a correlation with an effect of suppressing an increase in urinary β 2-microglobulin concentration by administration of the combination
10 preparation of hydrates of potassium citrate and sodium citrate. It was suggested that an effect of promoting excretion of uremic substance into urine by administration of the combination preparation of hydrates of potassium citrate and sodium citrate was not caused only by suppression of glomerular and kidney proximal tubular damage.

[0153]

15 Relevance of each of a concentrations of indoxyl sulfate (IS) in early morning urine and a concentration of indoxyl sulfate (IS) in plasma at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been
20 administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and all the patients (all the patients in the groups A, B, and C). The results are shown in FIGS. 1 to 4.

For indoxyl sulfate, a higher correlation between plasma concentration and urine concentration was recognized in the group to which the combination preparation of
25 hydrates of potassium citrate and sodium citrate had been administered as compared to

the control group, and a high correlation was recognized even when compared to the group to which the sodium bicarbonate preparation had been administered (refer to r values in FIGS. 1 to 3). Based on FIGS. 1 to 4, it was suggested that administration of the combination preparation of hydrates of potassium citrate and sodium citrate causes excretion of indoxyl sulfate into urine depending on the indoxyl sulfate concentration in blood. It was suggested that excretion of indoxyl sulfate into the urine depending on the indoxyl sulfate concentration in blood suppresses an increase in the concentration of indoxyl sulfate in blood, and a ratio of the indoxyl sulfate concentration in blood to the indoxyl sulfate concentration in urine was within a certain range.

10 [0154]

Relevance of each of a concentrations of p-cresyl sulfate (PCS) in early morning urine and a concentration of p-cresyl sulfate (PCS) in plasma at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and all the patients (all the patients in the groups A, B, and C). The results are shown in FIGS. 5 to 8.

Regarding p-cresyl sulfate, a correlation between plasma concentration and urine concentration was recognized between the control group, the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered, and the group to which the sodium bicarbonate preparation had been administered, and a higher correlation was recognized in the group to which the sodium bicarbonate preparation had been administered as compared to the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been

administered (refer to r values in FIGS. 5 to 7). Based on FIGS. 5 to 8, it was suggested that administration of the sodium bicarbonate preparation or the combination preparation of hydrates of potassium citrate and sodium citrate causes excretion of p-cresyl sulfate into urine depending on the p-cresyl sulfate concentration in blood. It was suggested
5 that excretion of p-cresyl sulfate into the urine depending on the p-cresyl sulfate concentration in blood suppresses an increase in the concentration of p-cresyl sulfate in blood, and a ratio of the p-cresyl sulfate concentration in blood to the p-cresyl sulfate concentration in urine was within a certain range.

[0155]

10 Relevance of each of a concentrations of hippuric acid (HA) in early morning urine and a concentration of hippuric acid (HA) in plasma at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been
15 administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and all the patients (all the patients in the groups A, B, and C). The results are shown in FIGS. 9 to 12.

 Regarding hippuric acid, no high correlation between plasma concentration and urinary concentration in the control group, the group to which the sodium bicarbonate
20 preparation had been administered, and the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered was recognized.

[0156]

 Relevance of each of a concentrations of argininosuccinic acid (ASA) in early
25 morning urine and a concentration of argininosuccinic acid (ASA) in plasma at 6, 12, and

24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and all the patients (all the patients in the groups A, B, and C). The results are shown in FIGS. 13 to 16.

Regarding argininosuccinic acid, no high correlation between plasma concentration and urine concentration in the control group, the group to which the sodium bicarbonate preparation had been administered, and the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered was recognized.

[0157]

Relevance of each of a concentrations of phenylacetyl-L-glutamine (PAG) in early morning urine and a concentration of phenylacetyl-L-glutamine (PAG) in plasma at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), and all the patients (all the patients in the groups A, B, and C). The results are shown in FIGS. 17 to 20.

Regarding phenylacetyl-L-glutamine, a correlation between plasma concentration and urine concentration was recognized between the control group, the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered, and the group to which the sodium bicarbonate preparation had been administered, and a higher correlation was recognized in the group to which the

sodium bicarbonate preparation had been administered as compared to the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered (refer to r values in FIGS. 17 to 19). Based on FIGS. 17 to 20, it was suggested that administration of the sodium bicarbonate preparation or the combination preparation of hydrates of potassium citrate and sodium citrate causes excretion of phenylacetyl-L-glutamine into the urine depending on the concentration of phenylacetyl-L-glutamine in the blood. It was suggested that excretion of phenylacetyl-L-glutamine into the urine depending on the phenylacetyl-L-glutamine concentration in blood suppresses an increase in the concentration of phenylacetyl-L-glutamine in blood, and a ratio of the phenylacetyl-L-glutamine concentration in blood to the phenylacetyl-L-glutamine concentration in urine falls within a certain range.

[0158]

Relevance of respective concentrations of indoxyl sulfate (IS), p-cresyl sulfate (PCS), hippuric acid (HA), argininosuccinic acid (ASA), and phenylacetyl-L-glutamine (PAG) in early morning urine at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W) was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). The results are shown in Table 9. In the table, "Contro" indicates the control group, "Citrate" indicates the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered, and "Bicarb" indicates the group to which the sodium bicarbonate preparation had been administered.

As a result, in the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered high correlation was

observed between indoxyl sulfate and phenylacetyl-L-glutamine, p-cresyl sulfate and phenylacetyl-L-glutamine, and argininosuccinic acid and phenylacetyl-L-glutamine. In the group to which the sodium bicarbonate preparation had been administered, high correlation was observed between indoxyl sulfate and argininosuccinic acid, indoxyl sulfate and phenylacetyl-L-glutamine, p-cresyl sulfate and argininosuccinic acid, p-cresyl sulfate and phenylacetyl-L-glutamine, and argininosuccinic acid and phenylacetyl-L-glutamine. It was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate may increase concentrations of indoxyl sulfate, phenylacetyl-L-glutamine, p-cresyl sulfate, and argininosuccinic acid in urine by the same mechanism.

In addition, it was suggested that the sodium bicarbonate preparation may increase concentrations of indoxyl sulfate, phenylacetyl-L-glutamine, p-cresyl sulfate, and argininosuccinic acid in urine by the same mechanism.

[0159]

[Table 9]

Relevance and influence of five uremic substances on concentration in urine

	IS			PCS			HA			ASA		
	Control	Citrate	Bicarb	Control	Citrate	Bicarb	Control	Citrate	Bicarb	Control	Citrate	Bicarb
PCS	0.5785	0.5789	0.6506									
	0.0001	<0.0001	<0.0001									
HA	0.5439	0.4184	0.6281	0.4105	0.5114	0.4178						
	0.0001	0.0031	<0.0001	0.0051	0.0002	0.0031						
ASA	0.6935	0.8207	0.7537	0.7531	0.6964	0.8598	0.5781	0.2946	0.4949			
	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.0421	0.0064			
PAG	0.6454	0.8379	0.7636	0.5811	0.8582	0.8185	0.5582	0.3843	0.5934	0.6519	0.7108	0.7285
	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.0157	0.0079	<0.0001	<0.0001	<0.0001	<0.0001

Upper: correlation coefficient (r), lower: significance level (p), Pearson. $r > 0.7$ is underlined.
3 group comparison (6 to 24 weeks, n = 45 - 48)

[0160]

Relevance of respective concentrations of indoxyl sulfate (IS), p-cresyl sulfate (PCS), hippuric acid (HA), argininosuccinic acid (ASA), and phenylacetyl-L-glutamine (PAG) in plasma at 6, 12, and 24 weeks after the start of the test (6W, 12W, and 24W)

was analyzed using a Pearson test for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). The results are shown in Table 10. In the table, “Contro” indicates the control group, “Citrate” indicates the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered, and “Bicarb” indicates the group to which the sodium bicarbonate preparation had been administered.

As a result, in the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered high correlation was observed between indoxyl sulfate and p-cresyl sulfate, indoxyl sulfate and phenylacetyl-L-glutamine, and p-cresyl sulfate and phenylacetyl-L-glutamine. In the group to which the sodium bicarbonate preparation had been administered, a high correlation was recognized between indoxyl sulfate and p-cresyl sulfate. It was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate may reduce concentrations of indoxyl sulfate, phenylacetyl-L-glutamine, and p-cresyl sulfate in blood by the same mechanism. In addition, it was suggested that the sodium bicarbonate preparation may reduce concentrations of p-cresyl sulfate and phenylacetyl-L-glutamine in blood by the same mechanism. It was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate and the sodium bicarbonate preparation have different mechanisms for reducing the concentration of uremic substance in blood.

[0161]

[Table 10]

Relevance and influence of five uremic substances on concentration in plasma

	IS			PCS			HA			ASA		
	Control	Citrate	Bicarb	Control	Citrate	Bicarb	Control	Citrate	Bicarb	Control	Citrate	Bicarb
PCS	0.1488	0.7128	0.5973	--	--	--	--	--	--	--	--	--
	0.2351	<0.0001	<0.0001									
HA	-0.6056	0.2504	0.0056	0.0424	0.2841	0.2949	--	--	--	--	--	--
	0.8686	0.6895	0.0062	0.7810	0.6503	0.6586						
ASA	0.4901	0.3069	0.3695	0.3629	0.1660	-0.0486	-0.0538	0.0068	0.2653	--	--	--
	0.0066	0.0359	0.0107	0.0155	0.2649	0.7428	0.7257	0.9640	0.0896			
PAG	0.9353	0.7664	0.5513	0.7552	0.9637	0.8486	0.1058	0.2463	0.3978	0.6792	0.1886	0.1431
	0.5280	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	0.4890	0.0959	0.0091	0.6050	0.2057	0.3597

Upper: correlation coefficient (r), lower: significance level (p), Pearson. $r > 0.7$ is underlined.
3 group comparison (6 to 24 weeks, n = 45 - 48)

[0162]

A urinary specific gravity of early morning urine before the start of the test (0 W) and at 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) was analyzed for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). The results are shown in Table 11-0-1. In addition, amounts of change in each urinary specific gravity after 6, 12, and 24 weeks after the start of the test are respectively shown as a % relative value with respect to each urinary specific gravity before the start of the test, and a difference from urinary specific gravity before the start of the test in Tables 11-0-2 and 11-0-3. The urinary specific gravity was measured using a urine hydrometer (PAL-09S, Atago Co., Ltd., Tokyo, Japan).

[0163]

[Table 11-0-1]

Urine Specific Gravity

Group	N	0W	6W	12W	24W	6-24W (45-48)
Control	15	1.014 ± 0.005	1.013 ± 0.005	1.011 ± 0.004 ^a	1.012 ± 0.005	1.012 ± 0.005
Citrate	16	1.012 ± 0.006	1.013 ± 0.005	1.014 ± 0.006	1.014 ± 0.006	1.014 ± 0.006
Bicarbonate	16	1.014 ± 0.007	1.014 ± 0.006	1.013 ± 0.005	1.016 ± 0.007	1.014 ± 0.006

Mean ± SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0464 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

[0164]

[Table 11-0-2]

Urine Specific Gravity (% Relative value vs week 0)

Group	N	6W	12W ^b	24W	6-24W ^a (45-48)
Control	15	99.96 ± 0.52	99.77 ± 0.40 ^{a,d}	99.80 ± 0.51 ^c	99.84 ± 0.48 ^{a,b,c}
Citrate	16	100.11 ± 0.65	100.22 ± 0.54	100.23 ± 0.60	100.20 ± 0.59
Bicarbonate	16	100.02 ± 0.53	99.89 ± 0.58 ^e	100.17 ± 0.61	100.00 ± 0.57

Mean ± SD

^ap=0.0023, ^bp=0.0577, ^cp=0.0734, ^dp=0.0289 and ^ep=0.0715 vs Citrate (Mann-Whitney)^ap=0.0100, ^bp=0.0652 (Kruskal-Wallis) and ^cp=0.0084, ^dp=0.0941 vs Citrate (Dunn)

5

[0165]

[Table 11-0-3]

Urine Specific Gravity (Δ Actual measurement value vs week 0)

Group	N	6W	12W ^b	24W	6-24W ^a (45-48)
Control	15	-0.0004 ± 0.0053	-0.0023 ± 0.0041 ^{a,d}	-0.0017 ± 0.0043 ^c	-0.0015 ± 0.0046 ^{a,c}
Citrate	16	0.0011 ± 0.0066	0.0022 ± 0.0055	0.0023 ± 0.0061	0.0019 ± 0.0060
Bicarbonate	16	0.0002 ± 0.0054	-0.0012 ± 0.0058 ^e	0.0017 ± 0.0062	0.0002 ± 0.0058 ^b

Mean ± SD

^ap=0.0023, ^bp=0.0592, ^cp=0.0704, ^dp=0.0289 and ^ep=0.0746 vs Citrate (Mann-Whitney)^ap=0.0102, ^bp=0.0668 (Kruskal-Wallis) and ^cp=0.0085, ^dp=0.0956 vs Citrate (Dunn)

[0166]

As a result of the above measurement, compared with the group C (Control: the control group), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered) and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), the urinary specific gravity value was maintained or increased over 6, 12, and 24 weeks after the start of the test. In addition, compared to the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), the urinary specific gravity showed tendency to be more favorably maintained or to increase.

It could be understood that maintaining or increasing the urinary specific gravity is based on maintenance or improvement of kidney function. Accordingly, based on the above results, administration of the alkalinizing agent to patients with stage G2 chronic kidney disease as well as stage G3b can suppress progression of chronic kidney disease, and it was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate further suppresses the progression of chronic kidney disease than the sodium bicarbonate preparation.

[0167]

Values obtained by correcting respective concentrations of uremic substance in early morning urine, that is, indoxyl sulfate (IS), p-cresyl sulfate (PCS), phenylacetyl-L-glutamine (PAG), hippuric acid (HA), and argininosuccinic acid (ASA) before the start of the test (0 W), and 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) by the following equation using the urinary specific gravity before the start of the test (0 W), and at 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) were

analyzed for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). Medical Examination, Vol. 44, No. 1, 1995, pages 79-83 by Tetsuo Aoki et al. was referred to. A urinary specific gravity 1.022 was converted into a reference value.

Specific gravity correction value (unit/1.022·UG) = actual measurement value × (1.022 - 1.000)/(specific gravity value - 1.000)

The results are shown in Table 11-1-1, Table 11-2-1, Table 11-3-1, Table 11-4-1, and Table 11-5-1. In addition, amounts of change in the values obtained by correcting, with the urinary specific gravity, the concentration of the uremic substance in the early morning urine 6, 12, and 24 weeks after the start of the test are respectively shown as a % relative value with respect to the correction value before the start of the test, and a difference from the correction value before the start of the test in Table 11-1-2, Table 11-1-3, Table 11-2-2, Table 11-2-3, Table 11-3-2, Table 11-3-3, Table 11-4-2, Table 11-4-3, Table 11-5-2, and Table 11-5-3.

[0168]

[Table 11-1-1]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$)						
Group	N	0W ^a	6W	12W	24W	6-24W ^b (45-48)
Control	15	65.0 ± 27.5 ^c	58.7 ± 32.0 ^d	60.9 ± 24.2	58.5 ± 19.6 ^{e,f}	57.7 ± 25.8 ^{e,f}
Citrate	16	66.1 ± 51.1 ^b	58.6 ± 34.6	74.9 ± 81.5	71.1 ± 54.6	68.2 ± 59.2 ^d
Bicarbonate	16	92.6 ± 43.5 ^a	83.3 ± 41.9	65.5 ± 29.0 ^e	81.1 ± 41.6	76.7 ± 38.0

Mean ± SD

^bp=0.0849, ^cp=0.0559, ^dp=0.0065, ^ep=0.0729, ^fp=0.0266 and ^gp=0.0849 vs Bicarbonate (Mann-Whitney)
^ap=0.0052 and ^bp=0.0730 vs 0 week (Wilcoxon)

^bp=0.0844, ^cp=0.0361 (Kruskal-Wallis) and ^dp=0.0202, ^ep=0.0385 vs Citrate (Dunn)
 (There is a difference between groups at week 0)

153

[0169]

[Table 11-1-2]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W ^a (45-48)
Control	15	88 ± 26	118 ± 101	96 ± 55	101 ± 67
Citrate	16	103 ± 67	110 ± 62	117 ± 70	110 ± 65
Bicarbonate	16	95 ± 30	77 ± 42	95 ± 59	89 ± 45

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0170]

5

[Table 11-1-3]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, Δ Actual measurement value vs week 0)

Group	N	6W	12W ^b	24W	6-24W ^a (45-48)
Control	15	-6.3 ± 17.4	-4.2 ± 27.0 ^c	-13.8 ± 20.9	-8.1 ± 22.0
Citrate	16	-7.5 ± 36.7	8.8 ± 44.6 ^{b,d}	5.0 ± 44.4	2.1 ± 41.8 ^{a,c}
Bicarbonate	16	-9.3 ± 27.7	-27.0 ± 32.9	-11.4 ± 40.8	-15.9 ± 34.4

Mean ± SD

^ap=0.0251, ^bp=0.0135 and ^cp=0.0651 vs Bicarbonate (Mann-Whitney)^ap=0.0648, ^bp=0.0368 (Kruskal-Wallis) and ^cp=0.0581, ^dp=0.0429 vs Citrate (Dunn)

[0171]

[Table 11-2-1]

154

Urine PCS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$)

Group	N	0W	6W	12W	24W	6-24W(47-48)
Control	14-15	56.7 $\pm$ 80.6	91.8 $\pm$ 72.3	73.8 $\pm$ 69.6	59.8 $\pm$ 69.8	74.5 $\pm$ 77.2
Citrate	16	61.1 $\pm$ 64.3	74.6 $\pm$ 53.6	68.7 $\pm$ 60.2	91.2 $\pm$ 101.0	78.2 $\pm$ 75.1
Bicarbonate	16	113.0 $\pm$ 119.5	98.1 $\pm$ 118.3	66.7 $\pm$ 73.0	84.8 $\pm$ 86.0	88.2 $\pm$ 93.2
Mean $\pm$ SD						
Not Significant between Groups (Mann-Whitney)						
^a p=0.0507 vs 0 week (Wilcoxon)						
Not Significant between Groups (Kruskal-Wallis & Dunn)						
(No significant difference between groups at week 0)						

[0172]

[Table 11-2-2]

Urine PCS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, % Relative value vs week 0)

Group	N	6W ^a	12W	24W ^b	6-24W ^c (45-48)
Control	14-15	152 $\pm$ 170	117 $\pm$ 169	85 $\pm$ 97 ^{cd}	121 $\pm$ 149 ^{a,d}
Citrate	16	193 $\pm$ 160	343 $\pm$ 829	209 $\pm$ 216	248 $\pm$ 497
Bicarbonate	16	83 $\pm$ 34 ^{de}	105 $\pm$ 112	265 $\pm$ 540	151 $\pm$ 323 ^{b,e}
Mean $\pm$ SD					
^a p=0.0095, ^b p=0.0125, ^c p=0.0149, ^d p=0.0240 vs Citrate (Mann-Whitney)					
^a p=0.0129, ^b p=0.0449, ^c p=0.0569 (Kruskal-Wallis) and ^d p=0.0250, ^e p=0.0439, ^f p=0.0393,					
^g p=0.0849 vs Citrate (Dunn)					

5

[0173]

[Table 11-2-3]

Urine PCS ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, Δ Actual measurement value vs week 0)

Group	N	6W	12W ^a	24W ^b	6-24W ^c (45-48)
Control	14-15	-1.0 $\pm$ 41.7	-12.9 $\pm$ 58.7	-28.7 $\pm$ 50.2 ^{cd}	-14.3 $\pm$ 51.4 ^{a,e}
Citrate	16	13.5 $\pm$ 39.7	7.6 $\pm$ 53.3	30.1 $\pm$ 64.2	17.1 $\pm$ 53.1
Bicarbonate	16	-14.9 $\pm$ 36.6 ^e	-46.3 $\pm$ 68.4 ^{de}	-28.3 $\pm$ 84.4	-29.8 $\pm$ 66.1 ^{b,d}
Mean $\pm$ SD					
^a p=0.0226, ^b p=0.0007, ^c p=0.0214, ^d p=0.0287 and ^e p=0.0465 vs Citrate (Mann-Whitney)					
^a p=0.0035, ^b p=0.0661, ^c p=0.0795 (Kruskal-Wallis) and ^d p=0.0020, ^e p=0.0736, ^f p=0.0863,					
^g p=0.0849 vs Citrate (Dunn)					

155

[0174]

[Table 11-3-1]

Urine PAG ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	146.4 $\pm$ 83.2	143.8 $\pm$ 110.4	144.8 $\pm$ 105.4	103.9 $\pm$ 82.4 ^a	132.5 $\pm$ 99.3
Citrate	16	124.2 $\pm$ 94.3	143.8 $\pm$ 145.6	126.4 $\pm$ 108.7	145.3 $\pm$ 147.4	133.5 $\pm$ 132.5
Bicarbonate	16	203.6 $\pm$ 169.1	202.9 $\pm$ 250.9	136.0 $\pm$ 141.4 ^b	147.1 $\pm$ 116.4	162.0 $\pm$ 182.5

Mean $\pm$ SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0302 and ^bp=0.0110 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

[0175]

5

[Table 11-3-2]

Urine PAG ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W*(45-48)
Control	15	108 $\pm$ 70	113 $\pm$ 67 ^b	85 $\pm$ 48	102 $\pm$ 62
Citrate	16	132 $\pm$ 104	113 $\pm$ 93	122 $\pm$ 108	123 $\pm$ 100 ^a
Bicarbonate	16	97 $\pm$ 71	76 $\pm$ 60	88 $\pm$ 49	87 $\pm$ 60

Mean $\pm$ SD^ap=0.0328 and ^bp=0.0799 vs Bicarbonate (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0176]

[Table 11-3-3]

156

Urine PAG ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, Δ Actual measurement value vs week 0)

Group	N	0W	12W ^b	24W	6-24W ^a (45-48)
Control	15	-2.6 ± 86.0	-1.6 ± 84.1 ^d	-36.0 ± 70.7	-13.4 ± 80.4
Citrate	16	19.7 ± 90.3	2.3 ± 61.7 ^c	18.0 ± 59.8 ^b	14.4 ± 79.9 ^{a,b}
Bicarbonate	16	-0.7 ± 162.0	-67.6 ± 91.9	-56.5 ± 113.7	-41.6 ± 126.8

Mean ± SD

^ap=0.0078, ^bp=0.0385, ^cp=0.0491 vs Bicarbonate and ^bp=0.0712 vs Control (Mann-Whitney)^ap=0.0248, ^bp=0.0613 (Kruskal-Wallis) and ^bp=0.0197 vs Bicarbonate (Dunn)

[0177]

[Table 11-4-1]

Urine HA ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	219.0 ± 157.4	189.8 ± 124.5	169.2 ± 64.3	164.3 ± 90.9	174.1 ± 94.9
Citrate	16	201.2 ± 109.8	209.3 ± 160.3	176.9 ± 97.7	176.1 ± 129.8	187.7 ± 120.5
Bicarbonate	16	202.3 ± 185.6	161.8 ± 111.1	171.1 ± 87.0	177.7 ± 111.0	170.2 ± 101.6

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

5

[0178]

[Table 11-4-2]

Urine HA ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W(45-48)
Control	15	97 ± 52	110 ± 87	119 ± 113	109 ± 86
Citrate	16	146 ± 157	119 ± 104	125 ± 121	130 ± 127
Bicarbonate	16	104 ± 89	133 ± 134	118 ± 96	118 ± 106

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

157

[0179]

[Table 11-4-3]

Urine HA ($\mu\text{g} \cdot \text{mL}^{-1}$ / 1.022·UG, Δ Actual measurement value vs week 0)

Group	N	0W	12W ^b	24W	6-24W(45-48)
Control	15	-30.2 ± 113.2	-49.8 ± 125.1	-52.6 ± 121.9	-44.2 ± 117.8
Citrate	16	8.0 ± 139.4	-24.3 ± 91.6	-25.1 ± 162.6	-13.8 ± 132.5
Bicarbonate	16	-40.5 ± 171.7	-31.2 ± 163.7	-24.6 ± 155.3	-22.1 ± 160.3

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0180]

5

[Table 11-5-1]

Urine ASA ($\mu\text{g} \cdot \text{mL}^{-1}$ / 1.022·UG)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	4.60 ± 1.77	4.43 ± 1.78	4.31 ± 2.07	3.66 ± 1.15 ^a	4.13 ± 1.71
Citrate	16	4.77 ± 1.75	4.96 ± 1.90	4.37 ± 1.80	4.64 ± 2.19	4.46 ± 1.93
Bicarbonate	16	5.87 ± 5.12	5.47 ± 3.87	4.66 ± 2.78	4.93 ± 3.53	5.02 ± 3.27

Mean ± SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0412 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

[0181]

[Table 11-5-2]

Urine ASA ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W(45-48)
Control	15	101 ± 39	97 ± 40	88 ± 44	96 ± 41
Citrate	16	101 ± 60	103 ± 66	106 ± 52	103 ± 59
Bicarbonate	16	101 ± 32	93 ± 29	97 ± 33	97 ± 31

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0182]

[Table 11-5-3]

Urine ASA ($\mu\text{g} \cdot \text{mL}^{-1} / 1.022 \cdot \text{UG}$, Δ Actual measurement value vs week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-0.17 ± 1.65	-0.29 ± 1.54	-1.14 ± 1.60	-0.53 ± 1.62
Citrate	16	-0.41 ± 1.95	-0.14 ± 1.74	-0.13 ± 1.55	-0.32 ± 2.05
Bicarbonate	16	-0.40 ± 2.25	-0.21 ± 3.04	-0.94 ± 3.02	-0.85 ± 2.76

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

5 [0183]

As a result, in the cases of indoxyl sulfate (IS), p-cresyl sulfate (PCS), and phenylacetyl-L-glutamine (PAG), the concentration of uremic substance in urine (excretion into urine) significantly increased in the group to which the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) had been administered than that group to which the sodium bicarbonate preparation (Bicarbonate) had been administered and the control group. Also regarding the cases of hippuric acid (HA) and argininosuccinic acid (ASA), the concentration of uremic substance in urine (excretion into urine) increased in the group to which the combination preparation of

hydrates of potassium citrate and sodium citrate (Citrate) had been administered than that group to which the sodium bicarbonate preparation (Bicarbonate) had been administered and the control group. In addition, administration of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) increased concentrations of indoxyl sulfate (IS), p-cresyl sulfate (PCS), and phenylacetyl-L-glutamine (PAG) in the urine (excretion into urine) compared to before the start of the test (0 W).

[0184]

The osmotic pressure of early morning urine before the start of the test (0 W) and at 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) was analyzed for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). The results are shown in Table 12-0-1. In addition, amounts of change in the osmotic pressure of each early morning urine after 6, 12, and 24 weeks after the start of the test are respectively shown as a % relative value with respect to the osmotic pressure of the early morning urine before the start of the test, and a difference from osmotic pressure of early morning urine before the start of the test in Tables 12-0-2 and 12-0-3. The osmotic pressure was measured using the freezing point depression method.

[0185]

[Table 12-0-1]

160

Urine Osmotic Pressure (mOsm/kg)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	489 ± 162	487 ± 167	413 ± 157 ^a	428 ± 169	442 ± 158
Citrate	16	418 ± 197	440 ± 168	477 ± 209	482 ± 177	467 ± 182
Bicarbonate	16	492 ± 256	488 ± 154	456 ± 143	515 ± 185	486 ± 164

Mean ± SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0616 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

[0186]

[Table 12-0-2]

Urine Osmotic Pressure (% Relative value vs week 0)

Group	N	6W	12W	24W	6-24W ^a (45-48)
Control	15	108 ± 45	89 ± 26 ^c	95 ± 38	97 ± 37 ^{a,b}
Citrate	16	121 ± 52	123 ± 48	142 ± 101	129 ± 70
Bicarbonate	16	108 ± 34	105 ± 57	114 ± 45	109 ± 45 ^b

Mean ± SD

^ap=0.0123, ^bp=0.0997 and ^cp=0.0436 vs Citrate (Mann-Whitney)^ap=0.0382 (Kruskal-Wallis) and ^bp=0.0358 vs Citrate (Dunn)

5

[0187]

[Table 12-0-3]

Urine Osmotic Pressure (Δ mOsm/kg vs week 0)

Group	N	6W	12W	24W	6-24W ^a (45-48)
Control	15	-2 $\pm$ 170	-76 $\pm$ 132 ^b	-47 $\pm$ 147	-41 $\pm$ 150 ^{a,b}
Citrate	16	22 $\pm$ 206	59 $\pm$ 184	64 $\pm$ 230	48 $\pm$ 204
Bicarbonate	16	-4 $\pm$ 165	-36 $\pm$ 176	23 $\pm$ 169	-6 $\pm$ 168

Mean $\pm$ SD^ap=0.0179 and ^bp=0.0375 vs Citrate (Mann-Whitney)^ap=0.0597 (Kruskal-Wallis) and ^bp=0.0618 vs Citrate (Dunn)

[0188]

As a result of the above measurement, compared with the group C (Control: the control group), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered) and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), the osmotic pressure value of early morning urine was maintained or increased over 6, 12, and 24 weeks after the start of the test. In addition, compared to the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered), in the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), an osmotic pressure in the early morning urine showed tendency to be more favorably maintained or to increase.

It could be understood that maintaining or increasing the osmotic pressure of early morning urine is based on maintenance or improvement of kidney function.

Accordingly, based on the above results, administration of the alkalinizing agent to patients with stage G2 chronic kidney disease as well as stage G3b can suppress progression of chronic kidney disease, and it was suggested that the combination preparation of hydrates of potassium citrate and sodium citrate further suppresses the

progression of chronic kidney disease than the sodium bicarbonate preparation.

[0189]

Values obtained by correcting respective concentrations of uremic substance in early morning urine, that is, indoxyl sulfate (IS), p-cresyl sulfate (PCS), phenylacetyl-L-glutamine (PAG), hippuric acid (HA), and argininosuccinic acid (ASA) before the start of the test (0 W), and 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) by the following equation using the osmotic pressure of early morning urine before the start of the test (0 W), and at 6, 12, and 24 weeks after the start of the test (6 W, 12 W, and 24 W) were analyzed for each of the group C (Control: the control group), the group A (Citrate: the group to which the combination preparation of hydrates of potassium citrate and sodium citrate had been administered), and the group B (Bicarbonate: the group to which the sodium bicarbonate preparation had been administered). Medical Examination, Vol. 44, No. 1, 1995, pages 79-83 by Tetsuo Aoki et al. was referred to. In addition, a urine osmotic pressure 770 mOsm/kg was converted into a reference value.

Osmotic pressure correction value (unit/500 mOsm·P) = actual measurement value x 500/osmotic pressure value

The results are shown in Table 12-1-1, Table 12-2-1, Table 12-3-1, Table 12-4-1, and Table 12-5-1. In addition, amounts of change in the values obtained by correcting, with the osmotic pressure of the early morning urine, the concentration of the uremic substance in the early morning urine 6, 12, and 24 weeks after the start of the test are respectively shown as a % relative value with respect to the correction value before the start of the test, and a difference from the correction value before the start of the test in Table 12-1-2, Table 12-1-3, Table 12-2-2, Table 12-2-3, Table 12-3-2, Table 12-3-3, Table 12-4-2, Table 12-4-3, Table 12-5-2, and Table 12-5-3.

[0190]

[Table 12-1-1]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-2} / 770 \text{ mOsm} \cdot \text{P}$)

Group	N	0W ^a	6W	12W	24W	6-24W ^b (45-48)
Control	15	65.4 ± 32.0	56.5 ± 32.1 ^c	59.7 ± 26.0	51.0 ± 19.7 ^{d1}	55.7 ± 26.1 ^{d2}
Citrate	16	67.1 ± 59.5 ^{ac}	59.4 ± 36.8	77.7 ± 91.9	70.8 ± 58.1	69.3 ± 65.3 ^c
Bicarbonate	16	87.9 ± 40.0	82.9 ± 44.6	69.9 ± 32.4 ^c	84.4 ± 48.4	77.0 ± 42.5

Mean ± SD

^ap=0.0424, ^bp=0.0090, ^cp=0.0839, ^dp=0.0214 and ^ep=0.0650 vs Bicarbonate (Mann-Whitney)^ap=0.0092 and ^bp=0.0413 vs 0 week (Wilcoxon)^bp=0.0839, ^cp=0.0266 (Kruskal-Wallis) and ^dp=0.0880, ^ep=0.0416 vs Bicarbonate (Dunn)

(There is a significant difference between groups at week 0)

[0191]

[Table 12-1-2]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-2} / 770 \text{ mOsm} \cdot \text{P}$, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W ^a (45-48)
Control	15	87 ± 29	119 ± 107	93 ± 54	100 ± 71
Citrate	16	107 ± 68	114 ± 64	119 ± 68	114 ± 66
Bicarbonate	16	98 ± 30	77 ± 39	104 ± 68	93 ± 49

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0192]

[Table 12-1-3]

Urine IS ($\mu\text{g} \cdot \text{mL}^{-1} / 770 \text{ mOsm} \cdot \text{P}$, Δ Actual measurement value vs week 0)

Group	N	6W	12W ^a	24W	6-24W(45-48)
Control	15	- 8.9 ± 20.4	- 5.7 ± 29.5	- 16.7 ± 25.0	- 10.4 ± 25.1
Citrate	16	- 7.6 ± 41.0	10.6 ± 45.6 ^{ab}	2.7 ± 46.5	2.2 ± 44.1
Bicarbonate	16	- 5.1 ± 28.3	- 24.1 ± 33.2	- 3.6 ± 49.0	- 10.9 ± 38.2

Mean ± SD

^ap=0.0169 vs Bicarbonate (Mann-Whitney)^ap=0.0492 (Kruskal-Wallis) and ^bp=0.0445 vs Bicarbonate (Dunn)

164

[0193]

[Table 12-2-1]

Urine PCS ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P)

Group	N	0W ^a	6W	12W	24W	6-24W(45-48)
Control	15	88.1 ± 82.8	90.4 ± 75.6	72.0 ± 87.1	56.4 ± 64.6 ^c	72.5 ± 75.9
Citrate	16	62.4 ± 72.2	75.3 ± 61.9	70.6 ± 62.1 ^b	90.2 ± 105.0	78.7 ± 77.8
Bicarbonate	16	108.8 ± 118.2	100.7 ± 126.6	66.9 ± 74.4	88.5 ± 93.1	85.4 ± 99.2

Mean ± SD
^aNot Significant between Groups (Mann-Whitney)
^bp=0.0637 and ^cp=0.0507 vs 0 week (Wilcoxon)
^cNot Significant between Groups (Kruskal-Wallis & Dunn)
 (No significant difference between groups at week 0)

[0194]

5

[Table 12-2-2]

Urine PCS ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, % Relative value vs week 0)

Group	N	6W	12W	24W ^b	6-24W ^c (44-48)
Control	14-15	159 ± 162	121 ± 186	85 ± 103 ^{c,e}	121 ± 153 ^{a,c}
Citrate	16	206 ± 173	375 ± 917	203 ± 200	262 ± 545
Bicarbonate	16	86 ± 35 ^d	102 ± 100	294 ± 590 ^e	161 ± 352 ^{b,d}

Mean ± SD
^ap=0.0054, ^bp=0.0127, ^cp=0.0085, ^dp=0.0259 vs Citrate and ^ep=0.0704 vs Control (Mann-Whitney)
^ap=0.0085, ^bp=0.0266 (Kruskal-Wallis) and ^cp=0.0120, ^dp=0.0524, ^ep=0.0241 vs Citrate (Dunn)

[0195]

[Table 12-2-3]

Urine PCS ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, Δ Actual measurement value vs week 0)

Group	N	6W	12W	24W ^b	6-24W ^a (45-48)
Control	15	- 3.5 ± 49.0	- 16.1 ± 63.1	- 33.5 ± 56.5 ^{a,c}	- 18.0 ± 56.7 ^{a,c}
Citrate	16	12.9 ± 43.9	8.2 ± 62.3	27.8 ± 62.0	16.3 ± 56.2
Bicarbonate	16	- 8.1 ± 32.0	- 41.9 ± 67.7 ^d	- 20.4 ± 95.1	- 29.5 ± 69.8 ^{b,d}

Mean ± SD

^ap=0.0117, ^bp=0.0047, ^cp=0.0093, ^dp=0.0465 vs Citrate (Mann-Whitney)

^ap=0.0080, ^bp=0.0436 (Kruskal-Wallis) and ^cp=0.0407, ^dp=0.0130, ^ep=0.0370 vs Citrate (Dunn)

[0196]

[Table 12-3-1]

Urine PAG ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	148.2 ± 87.4	139.0 ± 109.5	139.9 ± 100.3	103.3 ± 76.5 ^a	127.4 ± 95.8
Citrate	16	125.0 ± 107.6	147.3 ± 157.9	129.5 ± 115.2	145.9 ± 158.5	140.9 ± 142.4
Bicarbonate	16	197.7 ± 170.9	209.0 ± 279.2	135.4 ± 145.4 ^b	153.8 ± 128.0	166.1 ± 194.6

Mean ± SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0256 and ^bp=0.0092 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)
(No significant difference between groups at week 0)

5

[0197]

[Table 12-3-2]

Urine PAG ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W ^a (45-48)
Control	15	105 ± 68	114 ± 72 ^b	82 ± 46	100 ± 63
Citrate	16	137 ± 107	119 ± 99	123 ± 107	127 ± 103 ^a
Bicarbonate	16	99 ± 68	75 ± 55	97 ± 58	90 ± 60

Mean ± SD

^ap=0.0764 and ^bp=0.0912 vs Bicarbonate (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0198]

[Table 12-3-3]

Urine PAG ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, Δ Actual measurement value vs week 0)

Group	N	6W	12W ^b	24W	6-24W ^a (45-48)
Control	15	-7.2 ± 84.9	-6.3 ± 82.9	-42.3 ± 70.5 ^c	-18.6 ± 79.7 ^c
Citrate	16	22.3 ± 87.1	4.6 ± 65.8	20.9 ± 85.4	15.9 ± 78.7
Bicarbonate	16	11.3 ± 167.3	-62.4 ± 89.7 ^{d,e}	-44.0 ± 132.2	-31.7 ± 134.5 ^b

Mean ± SD

^ap=0.0912, ^bp=0.0386, ^cp=0.0712, ^dp=0.0310 vs Bicarbonate and ^ep=0.0491 vs Control (Mann-Whitney)^ap=0.0794, ^bp=0.0727 (Kruskal-Wallis) and ^cp=0.0903 vs Bicarbonate (Dunn)

[0199]

[Table 12-4-1]

Urine HA ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P)

Group	N	0W	6W	12W	24W	6-24W(45-48)
Control	15	225.6 ± 180.2	182.5 ± 123.7	166.0 ± 68.6	157.1 ± 90.4	168.5 ± 95.3
Citrate	16	198.4 ± 109.5	210.0 ± 156.9	181.2 ± 103.3	171.7 ± 131.4	187.6 ± 130.5
Bicarbonate	16	188.9 ± 161.7	162.2 ± 114.0	166.3 ± 89.9	182.0 ± 114.5	170.1 ± 104.8

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

5

[0200]

[Table 12-4-2]

167

Urine HA ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	97 ± 56	112 ± 97	116 ± 109	108 ± 89
Citrate	16	150 ± 161	124 ± 109	129 ± 128	134 ± 132
Bicarbonate	16	106 ± 91	130 ± 122	127 ± 106	121 ± 105

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

[0201]

[Table 12-4-3]

Urine HA ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P, Δ Actual measurement value vs week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	-43.1 ± 131.0	-59.6 ± 146.6	-67.4 ± 137.5	-56.7 ± 135.7
Citrate	16	11.6 ± 144.1	-17.1 ± 96.6	-26.7 ± 168.7	-10.8 ± 137.7
Bicarbonate	16	-26.8 ± 151.8	-22.7 ± 135.9	-7.0 ± 136.0	-18.8 ± 138.7

Mean ± SD

Not Significant between Groups (Mann-Whitney)

Not Significant between Groups (Kruskal-Wallis & Dunn)

5

[0202]

[Table 12-5-1]

Urine ASA ($\mu\text{g}\cdot\text{mL}^{-1}$ / 770 mOsm·P)

Group	N	6W	6W	12W	24W	6-24W(45-48)
Control	15	4.62 ± 2.07	4.27 ± 1.81	4.27 ± 2.28	3.48 ± 1.13 ^a	4.01 ± 1.80
Citrate	16	4.72 ± 1.96	4.49 ± 2.06	4.47 ± 1.96	4.70 ± 2.36	4.56 ± 2.09
Bicarbonate	16	5.48 ± 4.23	5.50 ± 4.08	4.63 ± 3.04	5.14 ± 3.95	5.09 ± 3.66

Mean ± SD

Not Significant between Groups (Mann-Whitney)

^ap=0.0302 vs 0 week (Wilcoxon)

Not Significant between Groups (Kruskal-Wallis & Dunn)

(No significant difference between groups at week 0)

[0203]

[Table 12-5-2]

Urine ASA (μg·mL⁻¹ / 770 mOsm·P, % Relative value vs week 0)

Group	N	6W	12W	24W	6-24W (45-48)
Control	15	99 ± 40	98 ± 45	85 ± 44	94 ± 42
Citrate	16	124 ± 61	109 ± 70	111 ± 57	115 ± 62 ^a
Bicarbonate	16	104 ± 33	93 ± 30	107 ± 42	101 ± 35

Mean ± SD
^ap=0.0716 vs Control (Mann-Whitney)
Not Significant between Groups (Kruskal-Wallis & Dunn)

[0204]

[Table 12-5-3]

Urine ASA (μg·mL⁻¹ / 770 mOsm·P, Δ Actual measurement value vs week 0)

Group	N	6W	12W	24W ^a	6-24W (45-48)
Control	15	-0.36 ± 1.78	-0.35 ± 1.71	-1.35 ± 1.84	-0.69 ± 1.80
Citrate	16	-0.24 ± 2.02	-0.25 ± 2.31	-0.03 ± 2.53 ^{a,b}	-0.17 ± 2.25
Bicarbonate	16	0.02 ± 1.67	-0.85 ± 2.02	-0.34 ± 2.72	-0.39 ± 2.17

Mean ± SD
^ap=0.0364 vs Control (Mann-Whitney)
^bp=0.0935 (Kruskal-Wallis) and ^cp=0.0962 vs Control (Dunn)

5

[0205]

As a result, in the cases of p-cresyl sulfate (PCS) and phenylacetyl-L-glutamine (PAG), the concentration of uremic substance in urine (excretion into urine) significantly increased in the group to which the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) had been administered than that group to which the sodium bicarbonate preparation (Bicarbonate) had been administered and the control group. Also regarding the cases of indoxyl sulfate (IS), hippuric acid (HA), and argininosuccinic acid (ASA), the concentration of uremic substance in urine (excretion

10

into urine) increased in the group to which the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) had been administered than that group to which the sodium bicarbonate preparation (Bicarbonate) had been administered and the control group. In addition, administration of the combination preparation of hydrates of potassium citrate and sodium citrate (Citrate) increased concentrations of indoxyl sulfate (IS), p-cresyl sulfate (PCS), and phenylacetyl-L-glutamine (PAG) in the urine (excretion into urine) compared to before the start of the test (0 W).

INDUSTRIAL APPLICABILITY

10 [0206]

By the pharmaceutical composition and the like provided by the present invention, uremic substance are excreted outside the body in mammals. By the method provided by the present invention, it is possible to preliminarily determine as to whether or not uremic substances are excreted outside the body and/or whether or not progression of chronic kidney disease can be suppressed.

CLAIMS

1. A food composition for maintenance of kidney function, comprising an alkalinizing agent.

5

2. The food composition according to Claim 1, wherein the maintenance of kidney function is suppression of kidney tubular damage, protection of kidney tubular cells, or maintenance of kidney tubular function.

10

3. The food composition according to Claim 2, wherein the kidney tubule is a kidney proximal tubule.

15

4. The food composition according to any one of Claims 1 to 3, wherein the alkalinizing agent is an acceptable salt of citric acid as food or a hydrate thereof, or a mixture thereof.

20

5. The food composition according to any one of Claims 1 to 4, wherein the alkalinizing agent comprises a mixture of sodium citrate or a hydrate thereof, and potassium citrate or a hydrate thereof.

6. The food composition according to any one of Claims 1 to 5, wherein the alkalinizing agent is sodium citrate or a hydrate thereof.

25

7. The food composition according to any one of Claims 1 to 6, which is a tablet.

8. The food composition according to any one of Claims 1 to 7, wherein an effect of the maintenance of kidney function is indicated on packaging, a container, or an instruction leaflet for the food composition.

5

9. The food composition according to any one of Claims 1 to 7, wherein an effect of suppression of kidney tubular damage, protection of kidney tubular cells, or maintenance of kidney tubular function is indicated on packaging, a container, or an instruction leaflet for the food composition.

10

10. The food composition according to any one of Claims 1 to 8, which is ingested by a healthy subject concerned about kidney health.

11. The food composition according to any one of Claims 1 to 7 and 9, which
15 is ingested by a healthy subject concerned about kidney tubular health.

12. The food composition according to any one of Claims 1 to 11, which is ingested by a subject in which a urinary β 2-microglobulin concentration is 290 μ g/L or less.

20

13. The food composition according to any one of Claims 1 to 12, which is ingested by a subject in which a urinary β 2-microglobulin concentration is 50 to 150 μ g/L.

25 14. The food composition according to any one of Claims 1 to 13, which is

ingested by a subject in which a blood cystatin C concentration is 0.5 to 2.2 mg/L.

15. The food composition according to any one of Claims 1 to 14, which is ingested by a subject in which a blood cystatin C concentration is 1.0 to 1.3 mg/L.

5

16. The food composition according to any one of Claims 1 to 15, wherein ingestion of the food composition suppresses an increase of a urinary β 2-microglobulin concentration.

10 17. The food composition according to any one of Claims 1 to 16, wherein, by ingestion of the food composition, after 12 weeks from the ingestion, an increase of a urinary β 2-microglobulin concentration is suppressed.

15 18. The food composition according to any one of Claims 1 to 17, wherein, by ingestion of the food composition, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the start of the ingestion.

19. The food composition according to any one of Claims 1 to 18, wherein, by ingestion of the food composition, after 12 weeks from the ingestion, a urinary β 2-microglobulin concentration is not substantially decreased compared to before the start of administration.

20 20. The food composition according to any one of Claims 1 to 19, wherein ingestion of the food composition does not substantially increase blood cystatin C
25 compared to before the start of the ingestion.

21. The food composition according to any one of Claims 1 to 20, wherein ingestion of the food composition does not substantially increase blood cystatin C compared to before the start of the ingestion.

5

22. The food composition according to any one of Claims 1 to 21, wherein ingestion of the food composition suppresses, in early morning urine, an increase of a β 2-microglobulin amount which is associated with progression of a stage of chronic kidney disease.

10

23. The food composition according to any one of Claims 1 to 22, wherein ingestion of the food composition does not affect glomerular function in a patient with chronic kidney disease, and suppresses kidney proximal tubular cell damage associated with progression of a stage of chronic kidney disease to protect kidney proximal tubular cells.

15

1/10

FIG. 1



Urine IS vs Plasma IS : Control

FIG. 2

Urine IS vs Plasma IS : Citrate

2/10

FIG. 3



Urine IS vs Plasma IS : Bicarbonate

FIG. 4

Urine IS vs Plasma IS : All

3/10

FIG. 5

Urine PCS vs Plasma PCS : Control





FIG. 6

Urine PCS vs Plasma PCS : Citrate



4/10

FIG. 7



Urine PCS vs Plasma PCS : Bicarbonate

FIG. 8

Urine PCS vs Plasma PCS : All

5/10

FIG. 9



Urine HA vs Plasma HA : Control

FIG. 10

Urine HA vs Plasma HA : Citrate

6/10

FIG. 11



Urine HA vs Plasma HA : Bicarbonate

FIG. 12

Urine HA vs Plasma HA : All

7/10

FIG. 13



Urine ASA vs Plasma ASA : Control

FIG. 14

Urine ASA vs Plasma ASA : Citrate

8/10

FIG. 15





Urine ASA vs Plasma ASA : Bicarbonate

FIG. 16

Urine ASA vs Plasma ASA : All

9/10

*FIG. 17*Urine PAG vs Plasma PAG : Control*FIG. 18*Urine PAG vs Plasma PAG : Citrate

10/10

FIG. 19


Urine PAG vs Plasma PAG : Bicarbonate

FIG. 20

Urine PAG vs Plasma PAG : All