

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2018255352 B2**

(54) Title
P-ethoxy nucleic acids for STAT3 inhibition

(51) International Patent Classification(s)
A61K 9/127 (2006.01) **A61K 47/69** (2017.01)
A61K 31/7125 (2006.01) **A61P 35/00** (2006.01)
A61K 47/54 (2017.01) **C07H 21/02** (2006.01)

(21) Application No: **2018255352** (22) Date of Filing: **2018.04.19**

(87) WIPO No: **WO18/195249**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/487,292	2017.04.19	US

(43) Publication Date: **2018.10.25**

(44) Accepted Journal Date: **2023.11.16**

(71) Applicant(s)
Bio-Path Holdings, Inc.

(72) Inventor(s)
Ashizawa, Ana Tari

(74) Agent / Attorney
Spruson & Ferguson, Level 24, Tower 2 Darling Park, 201 Sussex Street, Sydney, NSW, 2000, AU

(56) Related Art
US 2014/0336237 A1



(51) International Patent Classification:

A61K 9/127 (2006.01) A61K 47/69 (2017.01)
A61K 31/7125 (2006.01) A61P 35/00 (2006.01)
A61K 47/54 (2017.01) C07H 21/02 (2006.01)

(21) International Application Number:

PCT/US2018/028259

(22) International Filing Date:

19 April 2018 (19.04.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/487,292 19 April 2017 (19.04.2017) US

(71) Applicant: BIO-PATH HOLDINGS, INC. [US/US];
4710 Bellaire Blvd., Suite 210, Bellaire, TX 77401 (US).

(72) Inventor: ASHIZAWA, Ana, Tari; c/o Bio-Path Holdings, Inc., 4710 Bellaire Blvd., Suite 210, Bellaire, TX 77401 (US).

(74) Agent: SCHNEPP, Amanda, S.J.; Parker Highlander PLLC, 1120 S. Capital of Texas Highway, Bldg. One, Suite 200, Austin, TX 78746 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: P-ETHOXY NUCLEIC ACIDS FOR STAT3 INHIBITION




FIG. 1

(57) **Abstract:** Provided herein are improved delivery systems for oligonucleotides, said delivery system comprising a liposome that comprises neutral phospholipids and a P-ethoxy oligonucleotide, which targets a STAT3-encoding polynucleotide. Methods of treating patients with said delivery systems are also provided.



Published:

- *with international search report (Art. 21(3))*
- *with sequence listing part of description (Rule 5.2(a))*

DESCRIPTION

P-ETHOXY NUCLEIC ACIDS FOR STAT3 INHIBITION

[0001] The present application claims the priority benefit of United States provisional application number 62/487,292, filed April 19, 2017, the entire contents of which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0002] The present invention relates generally to the field of medicine. More particularly, it concerns liposomal formulations of P-ethoxy oligonucleotides that hybridize to a *STAT3* polynucleotide gene product and methods of making and using such formulations in medicine, even more particularly in the treatment of solid and hematological cancers that have high expression or increased activity of the *STAT3* gene.

2. Description of Related Art

[0003] Many genes are implicated in the sustained proliferation and survival of cancer cells in pre-malignant and advanced refractory tumors. Members of the Signal Transducer and Activator of Transcription (STAT) signaling pathways have been known to play an essential role in this capacity. STAT3 is a transcription factor that is over-activated in about 70% of all solid tumors and blood cancers. When activated STAT3 translocates from the cytosol into the nucleus to drive the expression of a large number of genes including Survivin, Bcl-X_L, Mcl-1, c-Myc, Cyclin D1, p21, Cyclin E, Matrix metalloproteinase-9 and -2, and VEGF to regulate a broad range of cellular processes such as cell survival, growth, migration, invasion, metastasis, and angiogenesis (Wu *et al.*, 2011; Hardee *et al.*, 2013). Additionally STAT3 regulates several important signaling pathways implicated in cancer progression including the IL-3, IL-6, and NF- κ B cascades, while also activating feedback pathways reinforcing its own activation (Hardee *et al.*, 2013).

[0004] Whereas STAT3 is usually expressed at low levels in the cytosol of healthy cells, it is frequently overexpressed and activated (phosphorylated and localized to the nucleus) in cancer, including diffuse large B-cell lymphoma (DLBCL). In DLBCL, it is most frequently activated in ABC subtype of the disease, and less often in GC subtype DLBCL (Wu *et al.*, 2011; Hardee *et al.*, 2013; Scuto *et al.*, 2011). Importantly, STAT3

overexpression, hyperphosphorylation, and nuclear localization correlate with a worse overall prognosis in these tumors compared with tumors expressing lower levels of STAT3. Additionally tumors bearing over-activated STAT3 also exhibit increased NF- κ B signaling in ABC sub-type DLBCL suggesting that STAT3 signaling plays a role in the progression of DLBCL.

[0005] Due to the nature of the STAT3 protein, designing small molecules to directly target its signaling activity is difficult, making it an ideal target for antisense and other genetic knockdown approaches. Given the central role that STAT3 plays in so many forms of cancer and other diseases there have been a number attempts however, to design peptide, decoy oligonucleotide, and small-molecule based compounds to block STAT3 activation, dimerization, DNA binding, or to accelerate inactivation of the enzyme (Miklossy *et al.*, 2013). As an alternative approach, multiple groups are working to develop Janus Kinase (JAK) inhibitors to block STAT3 phosphorylation and activation in LCL and other diseases with varied success (Amin *et al.*, 2003; Fagard *et al.*, 2013).

[0006] Approaches that target STAT3 specifically involve the use of oligonucleotides to block the translation of the *STAT3* mRNA into protein or to induce RNA degradation through RNaseH mechanisms *via* antisense oligonucleotides (ASO) and siRNA based technology. These strategies offer a way to reduce toxicity and are less adaptable to resistance mutations. Several attempts at this approach for STAT3 have not been successful. Thus, there is a need for improved antisense compositions targeting STAT3 for use in treatment of disease.

[0006a] The term “comprise” and variants of the term such as “comprises” or “comprising” are used herein to denote the inclusion of a stated integer or stated integers but not to exclude any other integer or any other integers, unless in the context or usage an exclusive interpretation of the term is required.

[0006b] Any reference to any prior art in this specification is not, and should not be taken as an acknowledgement or any form of suggestion that the prior art forms part of the common general knowledge.

SUMMARY OF THE INVENTION

[0006c] According to a first aspect, the present invention provides a composition comprising a population of oligonucleotides, phospholipids, and a surfactant, wherein the oligonucleotides hybridize to a *STAT3* polynucleotide gene product, wherein oligonucleotides of the population are composed of nucleoside molecules linked together through phosphate backbone linkages, wherein 60% to 75% of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage, and wherein 25% to 40% of the phosphate backbone linkages in each oligonucleotide are phosphodiester backbone linkages.

[0006d] According to a second aspect, the present invention provides a method for reducing the expression level of STAT3 protein in a cell comprising contacting the cell with a composition of the first aspect.

[0006e] According to a third aspect, the present invention provides a pharmaceutical composition comprising a composition according to the first aspect and a pharmaceutically acceptable carrier.

[0006f] According to a fourth aspect, the present invention provides a method of treating a subject with a cancer comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition of the third aspect.

[0006g] According to a fifth aspect, the present invention provides use of the composition of the first aspect in the manufacture of a medicament for treating cancer.

[0007] Provided herein are compositions and methods that induce growth inhibition and/or apoptosis in a wide range of cancer cells controlled by STAT3. STAT3 protein expression is prevented by a non-toxic nuclease resistant oligonucleotide that targets STAT3-encoding polynucleotides in combination with a neutral liposome, thus eliminating the pool of available STAT3 protein.

[0008] In one embodiment, compositions are provided comprising a population of oligonucleotides that hybridize to a *STAT3* polynucleotide gene product. In some aspects, the oligonucleotides of the population are composed of nucleoside molecules linked together through phosphate backbone linkages, wherein at least one of the phosphate backbone

linkages in each oligonucleotide is a P-ethoxy backbone linkage, and wherein no more than 80% of the phosphate backbone linkages in each oligonucleotide are P-ethoxy backbone linkages. In some aspects, at least one of the phosphate backbone linkages in each oligonucleotide is a phosphodiester backbone linkage. In some aspects, the oligonucleotides of the population comprise a sequence according to any one of SEQ ID NOs: 1-4. In some aspects, the oligonucleotides of the population comprise a sequence according to SEQ ID NO: 1. In some aspects, the oligonucleotides of the population comprise a sequence according to SEQ ID NO: 2. In some aspects, the oligonucleotides of the population comprise a sequence according to SEQ ID NO: 3. In some aspects, the oligonucleotides of the population comprise a sequence according to SEQ ID NO: 4. In various aspects, the oligonucleotides of the population inhibit the expression of STAT3. In some aspects, the composition is lyophilized.

[0009] In some aspects, 10% to 80% of the phosphate backbone linkages are P-ethoxy backbone linkages; 20% to 80% of the phosphate backbone linkages are P-ethoxy backbone linkages; 30% to 80% of the phosphate backbone linkages are P-ethoxy backbone linkages; 40% to 80% of the phosphate backbone linkages are P-ethoxy backbone linkages; 50% to 80% of the phosphate backbone linkages are P-ethoxy backbone linkages; or 60% to 70% of the phosphate backbone linkages are P-ethoxy backbone linkages, or any range derivable therein. In some aspects, 20% to 90% of the phosphate backbone linkages are phosphodiester backbone linkages; 20% to 80% of the phosphate backbone linkages are phosphodiester backbone linkages; 20% to 70% of the phosphate backbone linkages are phosphodiester backbone linkages; 20% to 60% of the phosphate backbone linkages are phosphodiester backbone linkages; 20% to 50% of the phosphate backbone linkages are phosphodiester backbone linkages; or 30% to 40% of the phosphate backbone linkages are phosphodiester backbone linkages, or any range derivable therein. In various aspects, at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, or any value therein, of the phosphate backbone linkages are P-ethoxy backbone linkages. In various aspects, at most 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, or any value therein, of the phosphate backbone linkages are phosphodiester backbone linkages. In certain aspects, the phosphodiester backbone linkages are distributed throughout the oligonucleotides. As such, the oligonucleotides are not chimeric molecules. In some aspects, the oligonucleotides do not comprise a phosphorothioate backbone linkage.

[0010] In some aspects, the oligonucleotides of the population have a size ranging from 7 to 30 nucleotides. In certain aspects, the oligonucleotides of the population have a size ranging from 12 to 25 nucleotides. In various aspects, the oligonucleotides of the population have a size of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides. The size range may be an average size of the oligonucleotides in the population.

[0011] In some aspects, the oligonucleotides of the population have an average size of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides, wherein no more than 5, 6, 7, 8, 8, 9, 10, 11, 11, 12, 13, 14, 15, 15, 16, 17, 18, 19, 20, 20, 21, 22, 23, or 24, respectively, of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage. In some aspects, the oligonucleotides of the population have an average size of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides and at least 2, 2, 2, 2, 3, 3, 3, 3, 4, 4, 4, 4, 4, 5, 5, 5, 5, 5, 6, 6, 6, 6, or 6, respectively, of the phosphate backbone linkages in each oligonucleotide is a phosphodiester backbone linkage. By way of example, the oligonucleotides of the population may have an average size of 18 nucleotides, wherein no more than 14 of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage; the oligonucleotides of the population may have an average size of 20 nucleotides, wherein no more than 16 of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage; the oligonucleotides of the population may have an average size of 25 nucleotides, wherein no more than 20 of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage; or the oligonucleotides of the population may have an average size of 30 nucleotides, wherein no more than 24 of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage.

[0012] In some aspects, the population of oligonucleotides comprises a single species of oligonucleotides. In other aspects, the population of oligonucleotides comprises at least two species of oligonucleotides. A single species of oligonucleotide may have the same nucleotide sequence but either have or lack P-ethoxy linkages in different positions within the molecule. As such, the population may be homogeneous as to the nucleotide sequence and heterogeneous as to the distribution of phosphodiester backbone linkages among the oligonucleotides of the population. In addition, the population may be heterogeneous as to the number of P-ethoxy backbone linkages and phosphodiester backbone linkages among the

oligonucleotides of the population. As a non-limiting example, a first portion of the oligonucleotides of the population may have 70% P-ethoxy linkages and 30% phosphodiester linkages while a second portion of the oligonucleotides of the population may have 60% P-ethoxy linkages and 40% phosphodiester linkages. In some aspects, the population of oligonucleotides comprises antisense oligonucleotides, short interfering RNAs (siRNAs), microRNAs (miRNAs), or piwiRNAs (piRNAs).

[0013] In various aspects, the composition further comprises phospholipids. In some aspects, the phospholipids and oligonucleotides are present at a molar ratio of from about 5:1 to about 100:1. In some aspects, the oligonucleotides and phospholipids form an oligonucleotide–lipid complex, such as, for example, a liposome complex. In some aspects, at least 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the liposomes are less than 5 microns in diameter. In some aspects, at least 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the liposomes are less than 4 microns in diameter. In some aspects, the population of oligonucleotides are incorporated in the population of liposomes.

[0014] In some aspects, the phospholipids are uncharged or have a neutral charge at physiologic pH. In some aspects, the phospholipids are neutral phospholipids. In certain aspects, the neutral phospholipids are phosphatidylcholines. In certain aspects, the neutral phospholipids are dioleoylphosphatidyl choline. In some aspects, the phospholipids are essentially free of cholesterol.

[0015] In one embodiment, pharmaceutical compositions are provided comprising a composition of oligonucleotides and phospholipids of the present embodiments and a pharmaceutically acceptable carrier. In some aspects, the composition further comprises a chemotherapeutic agent.

[0016] In one embodiment, methods are provided for delivering a therapeutically effective amount of an oligonucleotide to a cell comprising contacting the cell with a pharmaceutical composition of the present embodiments. In some aspects, the method is a method of treating hyperplasia, cancer, an autoimmune disease, or an infectious disease.

[0017] In one embodiment, methods are provided for treating a subject with cancer, an autoimmune disease, or an infectious disease comprising administering to the subject a

therapeutically effective amount of a pharmaceutical composition of the present embodiments. In some aspects, the subject is a human. In some aspects, the cancer is a bladder, blood, lymphoma, pancreas, bone, bone marrow, brain, breast, colon, esophagus, stomach, head and neck, kidney, liver, lung, prostate, skin, testis, tongue, ovary, or uterine cancer. In various aspects, the cancer is a lung adenocarcinoma, a leukemia, a lymphoma, such as, for example, diffuse large B cell lymphoma, or a pancreatic ductal adenocarcinoma. In some aspects, the autoimmune disease is systemic Lupus erythematosus, Spondyloarthropathy, Sjogren's disease, Crohn's disease, diabetes mellitus, multiple sclerosis, or rheumatoid arthritis. In some aspects, the infectious disease is a bacterial infection, fungal infection, viral infection, or parasitic infection. In some aspects, the composition is administering subcutaneously, intravenously, or intraperitoneally. In some aspects, the method further comprises administering at least a second anticancer therapy to the subject. In some aspects, the second anticancer therapy is a surgical therapy, chemotherapy, radiation therapy, cryotherapy, hormone therapy, immunotherapy, or cytokine therapy. In some aspects, the immunotherapy is a checkpoint blockade therapy. In some aspects, administration of the composition reduces expression of STAT3 protein in the patient.

[0018] An oligonucleotide includes an antisense nucleic acid molecule that specifically hybridizes to a nucleic acid molecule encoding a target protein or regulating the expression of the target protein. "Specific hybridization" means that the antisense nucleic acid molecule hybridizes to the targeted nucleic acid molecule and regulates its expression. Preferably, "specific hybridization" also means that no other genes or transcripts are affected. An oligonucleotide can be a single-stranded nucleic acid and may comprise 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more nucleobases. In particular aspects the oligonucleotide can comprise 15 to 30, 19 to 25, 20 to 23, or 21 contiguous nucleobases. In certain embodiments, the oligonucleotide inhibits the translation of a gene that promotes growth of a cancerous or pre-cancerous or hyperplastic mammalian cell (*e.g.*, a human cell). An oligonucleotide may induce apoptosis in the cell, and/or inhibit the translation of an oncogene or other target gene. In certain embodiments, the oligonucleotide component comprises a single species of oligonucleotide. In other embodiments, the oligonucleotide component comprises a 2, 3, 4 or more species of oligonucleotide that target 1, 2, 3, 4, or more genes. The composition may further comprise a chemotherapeutic or other anti-cancer agent, which may or may not be incorporated in a lipid

component or liposome of the invention. In further embodiments, the oligonucleotide component is incorporated within the liposome or lipid component.

[0019] “Entrap,” “encapsulate,” and “incorporate” refer to the lipid or liposome forming an impediment to free diffusion into solution by an association with or around an agent of interest, e.g., a liposome may encapsulate an agent within a lipid layer or within an aqueous compartment inside or between lipid layers. In certain embodiments, the composition is comprised in a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier may be formulated for administration to a human subject or patient.

[0020] In certain embodiments, the lipid component has an essentially neutral charge because it comprises a neutral phospholipid or a net neutral charge. In certain aspects a neutral phospholipid may be a phosphatidylcholine, such as DOPC, egg phosphatidylcholine (“EPC”), dilauroylphosphatidylcholine (“DLPC”), dimyristoylphosphatidylcholine (“DMPC”), dipalmitoylphosphatidylcholine (“DPPC”), distearoylphosphatidylcholine (“DSPC”), dilinoleoylphosphatidylcholine, 1,2-diarachidoyl-sn-glycero-3-phosphocholine (“DAPC”), 1,2-dieicosenoyl-sn-glycero-3-phosphocholine (“DEPC”), 1-myristoyl-2-palmitoyl phosphatidylcholine (“MPPC”), 1-palmitoyl-2-myristoyl phosphatidylcholine (“PMPC”), 1-palmitoyl-2-stearoyl phosphatidylcholine (“PSPC”), 1-stearoyl-2-palmitoyl phosphatidylcholine (“SPPC”), 1-palmitoyl-2-oleoyl phosphatidylcholine (“POPC”), 1-oleoyl-2-palmitoyl phosphatidylcholine (“OPPC”), or lysophosphatidylcholine. In other aspects the neutral phospholipid can be a phosphatidylethanolamine, such as dioleoylphosphatidylethanolamine (“DOPE”), distearoylphosphatidylethanolamine (“DSPE”), dimyristoyl phosphatidylethanolamine (“DMPE”), dipalmitoyl phosphatidylethanolamine (“DPPE”), palmitoyl-oleoyl phosphatidylethanolamine (“POPE”), or lysophosphatidylethanolamine. In certain embodiments, the phospholipid component can comprise 1, 2, 3, 4, 5, 6, 7, 8, or more kinds or types of neutral phospholipid. In other embodiments, a phospholipid component can comprise 2, 3, 4, 5, 6 or more kinds or type of neutral phospholipids.

[0021] In certain embodiments, a lipid component can have an essentially neutral charge because it comprises a positively charged lipid and a negatively charged lipid. The lipid component may further comprise a neutrally charged lipid(s) or phospholipid(s). The positively charged lipid may be a positively charged phospholipid. The negatively charged lipid may be a negatively charged phospholipid. The negatively charged phospholipid may be

a phosphatidylserine, such as dimyristoyl phosphatidylserine (“DMPS”), dipalmitoyl phosphatidylserine (“DPPS”), or brain phosphatidylserine (“BPS”). The negatively charged phospholipid may be a phosphatidylglycerol, such as dilauroylphosphatidylglycerol (“DLPG”), dimyristoylphosphatidylglycerol (“DMPG”), dipalmitoylphosphatidylglycerol (“DPPG”), distearoylphosphatidylglycerol (“DSPG”), or dioleoylphosphatidylglycerol (“DOPG”). In certain embodiments, the composition further comprises cholesterol or polyethyleneglycol (PEG). In other embodiments, the composition is essentially free of cholesterol. In certain embodiments, a phospholipid is a naturally-occurring phospholipid. In other embodiments, a phospholipid is a synthetic phospholipid.

[0022] Liposomes can be made of one or more phospholipids, as long as the lipid material is substantially uncharged. It is important that the composition be substantially free of anionic and cationic phospholipids and cholesterol. Suitable phospholipids include phosphatidylcholines and others that are well known to persons that are skilled in this field.

[0023] Another aspect of the present invention involves methods for delivering oligonucleotide to a cell comprising contacting the cell with a neutral lipid composition of the invention. The methods will provide an inventive composition in an effective amount. An effective amount is an amount of therapeutic component that attenuates, slows, reduces or eliminates a cell, condition, or disease state in a subject. The cell may be comprised in a subject or patient, such as a human. The method may further comprise a method of treating cancer or other hyperplastic condition. The cancer may have originated in the bladder, blood, bone, bone marrow, brain, breast, colon, esophagus, gastrointestinal, gum, head, kidney, liver, lymph node, lung, nasopharynx, neck, prostate, skin, stomach, testis, tongue, ovary, or uterus. In certain embodiments, the method further comprises a method of treating a non-cancerous disease or hyperplastic condition. The cell may be a pre-cancerous or a cancerous cell. In certain embodiments, the compositions and methods inhibit the growth of the cell, induce apoptosis in the cell, and/or inhibit the translation of an oncogene. The oligonucleotide may inhibit the translation of a gene that is overexpressed in the cancerous cell.

[0024] In certain embodiments, the methods of the invention further comprise administering an additional therapy to the subject. The additional therapy may comprise administering a chemotherapeutic (*e.g.*, paclitaxel or docetaxel), a surgery, a radiation therapy, and/or a gene therapy. In certain aspects the chemotherapy is docetaxel, paclitaxel, cisplatin (CDDP), carboplatin, procarbazine, mechlorethamine, cyclophosphamide,

camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatin, 5-fluorouracil, vincristin, vinblastin, methotrexate, or combinations thereof. In certain embodiments the chemotherapy is a taxane such as docetaxel or paclitaxel. The chemotherapy can be delivered before, during, after, or combinations thereof relative to a neutral lipid composition of the invention. A chemotherapy can be delivered within 0, 1, 5, 10, 12, 20, 24, 30, 48, or 72 hours or more of the neutral lipid composition. The neutral lipid composition, the second anti-cancer therapy, or both the neutral lipid composition and the anti-cancer therapy can be administered intratumorally, intravenously, intraperitoneally, subcutaneously, orally or by various combinations thereof.

[0025] It is contemplated that any embodiment discussed in this specification can be implemented with respect to any method or composition of the invention, and vice versa. Furthermore, compositions of the invention can be used to achieve the methods of the invention.

[0026] As used herein, “essentially free,” in terms of a specified component, is used herein to mean that none of the specified component has been purposefully formulated into a composition and/or is present only as a contaminant or in trace amounts. The total amount of the specified component resulting from any unintended contamination of a composition is therefore well below 0.05%, preferably below 0.01%. Most preferred is a composition in which no amount of the specified component can be detected with standard analytical methods.

[0027] As used herein the specification, “a” or “an” may mean one or more. As used herein in the claim(s), when used in conjunction with the word “comprising,” the words “a” or “an” may mean one or more than one.

[0028] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” As used herein “another” may mean at least a second or more.

[0029] Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

[0030] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0031] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0032] **FIG. 1** – Inhibition of STAT3 expression by liposomal *STAT3* antisense. The ability of liposomal *STAT3* antisense to inhibit the expression of STAT3 was tested in two lung adenocarcinoma cell lines, H1975 and HCC 827. Liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 (180 µg/mL) or an equivalent amount of empty liposomes were incubated with each cell line for four days.

[0033] **FIGs. 2A-C** – Inhibition of non-small cell lung cancer cell viability by liposomal *STAT3* antisense. The ability of liposomal *STAT3* antisense to inhibit the viability of non-small cell lung cancer cells was tested in three lung adenocarcinoma cell lines: H1975 (FIG. 2A), HCC 827 (FIG. 2B), and H358 (FIG. 2C). Liposomal *STAT3* antisense corresponding to one of SEQ ID NOs: 1-4 was incubated with each cell line for four days.

[0034] **FIGs. 3A-B** – Inhibition of leukemia and lymphoma cell viability by liposomal *STAT3* antisense. The ability of liposomal *STAT3* antisense to inhibit the growth of leukemia cells was tested in three human leukemia cell lines (K-562, MV4-11, and Kasumi-1) (FIG. 3A) and five human lymphoma cell lines (DOHH-2, SU-DHL-4, SU-DHL-6, SU-DHL-10, and RL) (FIG. 3B). Liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 was incubated with each cell line for four days.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0035] The present invention provides compositions and methods for delivery of an anti-*STAT3* oligonucleotide (*e.g.*, an inhibitor of gene expression) to a cell *via* a lipid composition, in certain aspects a lipid composition with a net charge of about zero, *i.e.*, a neutral lipid composition. In certain embodiments the lipid composition is a non-charged liposome. These methods may be effectively used to treat a cancer or an auto-immune disease.

I. Lipids and Liposomes

[0036] “Liposomes” is used herein to mean lipid-containing vesicles having a lipid bilayer, as well as other lipid carrier particles that can entrap or incorporate antisense oligonucleotides. As such, liposome is a generic term encompassing a variety of unilamellar, multilamellar, and multivesicular lipid vehicles formed by the generation of enclosed lipid bilayers or aggregates. In addition, liposomes may have an undefined lamellar structure. Liposomes may be characterized as having vesicular structures with a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). However, the present invention also encompasses compositions that have different structures in solution than the normal vesicular structure. For example, the lipids may assume a micellar structure or merely exist as non-uniform aggregates of lipid molecules.

[0037] Liposomes are a form of nanoparticles that are carriers for delivering a variety of drugs into a diseased tissue. Optimal liposome size depends on the target tissue. In tumor tissue, the vasculature is discontinuous, and pore sizes vary from 100 to 780 nm (Siwak *et al.*, 2002). By comparison, pore size in normal vascular endothelium is <2 nm in most tissues, and 6 nm in post-capillary venules. Negatively charged liposomes are thought to be more rapidly removed from circulation than neutral or positively charged liposomes; however, recent studies have indicated that the type of negatively charged lipid affects the rate of liposome uptake by the reticulo-endothelial system (RES). For example, liposomes containing negatively charged lipids that are not sterically shielded (phosphatidylserine,

phosphatidic acid, and phosphatidylglycerol) are cleared more rapidly than neutral liposomes. Interestingly, cationic liposomes (1,2-dioleoyl-3-trimethylammonium-propane [DOTAP]) and cationic-liposome-DNA complexes are more avidly bound and internalized by endothelial cells of angiogenic blood vessels via endocytosis than anionic, neutral, or sterically stabilized neutral liposomes (Thurston *et al.*, 1998; Krasnici *et al.*, 2003). Cationic liposomes may not be ideal delivery vehicles for tumor cells because surface interactions with the tumor cells create an electrostatically derived binding-site barrier effect, inhibiting further association of the delivery systems with tumor spheroids (Kostarelos *et al.*, 2004). However, neutral liposomes appear to have better intratumoral penetration. Toxicity with specific liposomal preparations has also been a concern. Cationic liposomes elicit dose-dependent toxicity and pulmonary inflammation by promoting release of reactive oxygen intermediates, and this effect is more pronounced with multivalent cationic liposomes than monovalent cationic liposomes, such as DOTAP (Dokka *et al.*, 2000). Neutral and negative liposomes do not appear to exhibit lung toxicity (Guitierrez-Puente *et al.*, 1999). Cationic liposomes, while efficiently taking up nucleic acids, have had limited success for *in vivo* gene down-regulation, perhaps because of their stable intracellular nature and resultant failure to release nucleic acid contents. Lipids with neutral charge or lipid compositions with a neutralized charge, *e.g.*, 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), are used herein because of the neutral properties and success in delivering antisense oligonucleotides *in vivo*.

[0038] The present invention provides methods and compositions for associating an oligonucleotide, such as an antisense oligonucleotide, with a lipid and/or liposome. The oligonucleotide may be incorporated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome *via* a linking molecule that is associated with both the liposome and the oligonucleotide, entrapped in a liposome, complexed with a liposome, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid. The liposome or liposome/oligonucleotide-associated compositions provided herein are not limited to any particular structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a “collapsed” structure. They may also simply be interspersed in a solution, possibly forming aggregates that are not uniform in either size or shape.

A. Lipids

[0039] Lipids are fatty substances that may be naturally occurring or synthetic. For example, lipids include the fatty droplets that naturally occur in the cytoplasm as well as the class of compounds that are well known to those of skill in the art that contain long-chain aliphatic hydrocarbons and their derivatives, such as fatty acids, alcohols, amines, amino alcohols, and aldehydes. An example is the lipid 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC).

[0040] Lipid compositions of the present invention may comprise phospholipids. In certain embodiments, a single kind or type of phospholipid may be used in the creation of lipid compositions, such as liposomes. In other embodiments, more than one kind or type of phospholipid may be used.

[0041] Phospholipids include glycerophospholipids and certain sphingolipids. Phospholipids include, but are not limited to, dioleoylphosphatidylcholine (“DOPC”), egg phosphatidylcholine (“EPC”), dilauryloylphosphatidylcholine (“DLPC”), dimyristoylphosphatidylcholine (“DMPC”), dipalmitoylphosphatidylcholine (“DPPC”), distearoylphosphatidylcholine (“DSPC”), dilinoleoylphosphatidylcholine, 1,2-diarachidoyl-sn-glycero-3-phosphocholine (“DAPC”), 1,2-dieicosenoyl-sn-glycero-3-phosphocholine (“DEPC”), 1-myristoyl-2-palmitoyl phosphatidylcholine (“MPPC”), 1-palmitoyl-2-myristoyl phosphatidylcholine (“PMPC”), 1-palmitoyl-2-stearoyl phosphatidylcholine (“PSPC”), 1-stearoyl-2-palmitoyl phosphatidylcholine (“SPPC”), palmitoyloleoyl phosphatidylcholine (“POPC”), 1-oleoyl-2-palmitoyl phosphatidylcholine (“OPPC”), dilauryloylphosphatidylglycerol (“DLPG”), dimyristoylphosphatidylglycerol (“DMPG”), dipalmitoylphosphatidylglycerol (“DPPG”), distearoylphosphatidylglycerol (“DSPG”), dioleoylphosphatidylglycerol (“DOPG”), dimyristoyl phosphatidic acid (“DMPA”), dipalmitoyl phosphatidic acid (“DPPA”), distearoyl phosphatidic acid (“DSPA”), dioleoyl phosphatidic acid (“DOPA”), dimyristoyl phosphatidylethanolamine (“DMPE”), dipalmitoyl phosphatidylethanolamine (“DPPE”), distearoylphosphatidylethanolamine (“DSPE”), dioleoylphosphatidylethanolamine (“DOPE”), palmitoyloleoyl phosphatidylethanolamine (“POPE”), dimyristoyl phosphatidylserine (“DMPS”), dipalmitoyl phosphatidylserine (“DPPS”), brain phosphatidylserine (“BPS”), distearoyl sphingomyelin (“DSSP”), brain sphingomyelin (“BSP”), dipalmitoyl sphingomyelin (“DPSP”), lysophosphatidylcholine, and lysophosphatidylethanolamine.

[0042] Phospholipids include, for example, phosphatidylcholines, phosphatidylglycerols, and phosphatidylethanolamines; because phosphatidylethanolamines and phosphatidylcholines are non-charged under physiological conditions (*i.e.*, at about pH 7), these compounds may be particularly useful for generating neutral liposomes. In certain embodiments, the phospholipid DOPC is used to produce non-charged liposomes or lipid compositions. In certain embodiments, a lipid that is not a phospholipid (*e.g.*, a cholesterol) can also be used

[0043] Phospholipids may be from natural or synthetic sources. However, phospholipids from natural sources, such as egg or soybean phosphatidylcholine, brain phosphatidic acid, brain or plant phosphatidylinositol, heart cardiolipin, and plant or bacterial phosphatidylethanolamine, are not used in certain embodiments as the primary phosphatide (*i.e.*, constituting 50% or more of the total phosphatide composition) because this may result in instability and leakiness of the resulting liposomes.

B. Neutral Liposomes

[0044] “Neutral liposomes or lipid composition” or “non-charged liposomes or lipid composition,” as used herein, are defined as liposomes or lipid compositions having one or more lipids that yield an essentially-neutral net charge (substantially non-charged). In certain embodiments, neutral liposomes or lipid compositions may include mostly lipids and/or phospholipids that are themselves neutral. In certain embodiments, amphipathic lipids may be incorporated into or used to generate neutral liposomes or lipid compositions. For example, a neutral liposome may be generated by combining positively and negatively charged lipids so that those charges substantially cancel one another, thereby yielding an essentially-neutral net charge. By “essentially neutral” or “essentially non-charged,” it is meant that few, if any, lipids within a given population (*e.g.*, a population of liposomes) include a charge that is not canceled by an opposite charge of another component (*e.g.*, fewer than 10% of components include a non-canceled charge, more preferably fewer than 5%, and most preferably fewer than 1%). In certain embodiments of the present invention, a composition may be prepared wherein the lipid component of the composition is essentially neutral but is not in the form of liposomes.

[0045] The size of the liposomes varies depending on the method of synthesis. A liposome suspended in an aqueous solution is generally in the shape of a spherical vesicle, and may have one or more concentric layers of lipid bilayer molecules. Each layer consists of

a parallel array of molecules represented by the formula XY, wherein X is a hydrophilic moiety and Y is a hydrophobic moiety. In aqueous suspension, the concentric layers are arranged such that the hydrophilic moieties tend to remain in contact with an aqueous phase and the hydrophobic regions tend to self-associate. For example, when aqueous phases are present within the liposome, the lipid molecules may form a bilayer, known as a lamella, of the arrangement XY-YX. Aggregates of lipids may form when the hydrophilic and hydrophobic parts of more than one lipid molecule become associated with each other. The size and shape of these aggregates will depend upon many different variables, such as the nature of the solvent and the presence of other compounds in the solution.

[0046] Liposomes within the scope of the present invention can be prepared in accordance with known laboratory techniques, such as, for example, the method of Bangham *et al.* (1965), the contents of which are incorporated herein by reference; the method of Gregoriadis (1979), the contents of which are incorporated herein by reference; the method of Deamer and Uster (1983), the contents of which are incorporated by reference; and the reverse-phase evaporation method as described by Szoka and Papahadjopoulos (1978). The aforementioned methods differ in their respective abilities to entrap aqueous material and their respective aqueous space-to-lipid ratios.

[0047] In certain embodiments, a neutral liposome may be used to deliver an oligonucleotide, such as an antisense oligonucleotide. The neutral liposome may contain a single species of oligonucleotide directed to the suppression of translation of a single gene, or the neutral liposome may contain multiple species of oligonucleotides that are directed to the suppression of translation of multiple genes. Further, the neutral liposome may also contain a chemotherapeutic in addition to the oligonucleotide; thus, in certain embodiments, a chemotherapeutic and an oligonucleotide may be delivered to a cell (*e.g.*, a cancerous cell in a human subject) in the same or separate compositions.

[0048] Dried lipids or lyophilized liposomes may be dehydrated and reconstituted at an appropriate concentration with a suitable solvent (*e.g.*, DPBS or Hepes buffer). The mixture may then be vigorously shaken in a vortex mixer. The liposomes may be resuspended at an appropriate total phospholipid concentration (*e.g.*, about 10-200 mM). Unencapsulated oligonucleotide may be removed by centrifugation at 29,000 g and the liposomal pellets washed. Alternatively, the unencapsulated oligonucleotides may be

removed by dialyzing against an excess of solvent. The amount of oligonucleotide encapsulated can be determined in accordance with standard methods.

II. Inhibition of Gene Expression

[0049] An inhibitory oligonucleotide can inhibit the transcription or translation of a gene in a cell. An oligonucleotide may be from 5 to 50 or more nucleotides long, and in certain embodiments from 7 to 30 nucleotides long. In certain embodiments, the oligonucleotide may be 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides long. The oligonucleotide may comprise a nucleic acid and/or a nucleic acid analog. Typically, an inhibitory oligonucleotide will inhibit the translation of a single gene within a cell; however, in certain embodiments, an inhibitory oligonucleotide may inhibit the translation of more than one gene within a cell.

[0050] Within an oligonucleotide, the components of the oligonucleotide need not be of the same type or homogenous throughout (*e.g.*, an oligonucleotide may comprise a nucleotide and a nucleic acid or nucleotide analog). In certain embodiments of the present invention, the oligonucleotide may comprise only a single nucleic acid or nucleic acid analog. The inhibitory oligonucleotide may comprise 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30 or more contiguous nucleobases, including all ranges therebetween, that hybridize with a complementary nucleic acid to form a double-stranded structure.

III. Nucleic Acids

[0051] The present invention provides methods and compositions for the delivery of an oligonucleotide *via* neutral liposomes. Because an oligonucleotide is composed of a nucleic acid, methods relating to nucleic acids (*e.g.*, production of a nucleic acid, modification of a nucleic acid, *etc.*) may also be used with regard to an oligonucleotide.

[0052] The term “nucleic acid” is well known in the art. A “nucleic acid” as used herein generally refers to a molecule (*i.e.*, a strand) of DNA, RNA, or a derivative or analog thereof, comprising a nucleobase. These definitions refer to a single-stranded or double-stranded nucleic acid. Double-stranded nucleic acids may be formed by fully complementary binding; however, in some embodiments, a double-stranded nucleic acid may be formed by partial or substantial complementary binding. As used herein, a single-stranded nucleic acid may be denoted by the prefix “ss” and a double-stranded nucleic acid by the prefix “ds.”

A. Nucleobases

[0053] As used herein a “nucleobase” refers to a heterocyclic base, such as, for example, a naturally occurring nucleobase (*i.e.*, an A, T, G, C or U) found in at least one naturally occurring nucleic acid (*i.e.*, DNA and RNA), and naturally or non-naturally occurring derivative(s) and analogs of such a nucleobase. A nucleobase generally can form one or more hydrogen bonds (*i.e.*, “anneal” or “hybridize”) with at least one naturally occurring nucleobase in a manner that may substitute for naturally occurring nucleobase pairing (*e.g.*, the hydrogen bonding between A and T, G and C, and A and U). A nucleobase may be comprised in a nucleoside or nucleotide, using any chemical or natural synthesis method described herein or known to one of ordinary skill in the art.

[0054] “Purine” and/or “pyrimidine” nucleobase(s) encompass naturally occurring purine and/or pyrimidine nucleobases and also derivative(s) and analog(s) thereof, including but not limited to, a purine or pyrimidine substituted by one or more of an alkyl, carboxyalkyl, amino, hydroxyl, halogen (*i.e.*, fluoro, chloro, bromo, or iodo), thiol, or alkylthiol moiety. Preferred alkyl (*e.g.*, alkyl, caboxyalkyl, *etc.*) moieties comprise of from about 1, about 2, about 3, about 4, about 5, to about 6 carbon atoms. Other non-limiting examples of a purine or pyrimidine include a deazapurine, a 2,6-diaminopurine, a 5-fluorouracil, a xanthine, a hypoxanthine, a 8-bromoguanine, a 8-chloroguanine, a bromothylane, a 8-aminoguanine, a 8-hydroxyguanine, a 8-methylguanine, a 8-thioguanine, an azaguanine, a 2-aminopurine, a 5-ethylcytosine, a 5-methylcytosine, a 5-bromouracil, a 5-ethyluracil, a 5-iodouracil, a 5-chlorouracil, a 5-propyluracil, a thiouracil, a 2-methyladenine, a methylthioadenine, a N,N-dimethyladenine, an azaadenines, a 8-bromoadenine, a 8-hydroxyadenine, a 6-hydroxyaminopurine, a 6-thiopurine, a 4-(6-aminoethyl/cytosine), and the like. Purine and pyrimidine derivatives or analogs include, but are not limited to (abbreviation/modified base description): ac4c/4-acetylcytidine, Mam5s2u/5-methoxyaminomethyl-2-thiouridine, Chm5u/5-(carboxyhydroxymethyl) uridine, Man q/Beta, D-mannosylqueosine, Cm/2'-O-methylcytidine, Mcm5s2u/5-methoxycarbonylmethyl-2-thiouridine, Cmm5s2u/5-carboxymethylamino-methyl-2-thioridine, Mcm5u/5-methoxycarbonylmethyluridine, Cmm5u/5-carboxymethylaminomethyluridine, Mo5u/5-methoxyuridine, D/Dihydrouridine, Ms2i6a, 2-methylthio-N6-isopentenyladenosine, Fm/2'-O-methylpseudouridine, Ms2t6a/N-((9-beta-D-ribofuranosyl-2-methylthiopurine-6-yl)carbamoyl)threonine, Gal q/Beta, D-galactosylqueosine, Mt6a/N-((9-beta-D-ribofuranosylpurine-6-yl)N-methyl-

carbamoyl)threonine, Gm/2'-O-methylguanosine, Mv/Uridine-5-oxyacetic acid methylester, I/Inosine, o5u/Uridine-5-oxyacetic acid (v), I6a/N6-isopentenyladenosine, Osyw/Wybutoxosine, m1a/1-methyladenosine, P/Pseudouridine, m1f/1-methylpseudouridine, Q/Queosine, m1g/1-methylguanosine, s2c/2-thiocytidine, m1I/1-methylinosine, s2t/5-methyl-2-thiouridine, m22g/2,2-dimethylguanosine, s2u/2-thiouridine, m2a/2-methyladenosine, s4u/4-thiouridine, m2g/2-methylguanosine, T/5-methyluridine, m3c/3-methylcytidine, t6a/N-((9-beta-D-ribofuranosylpurine-6-yl)carbamoyl)threonine, m5c/5-methylcytidine, Tm/2'-O-methyl-5-methyluridine, m6a/N6-methyladenosine, Um/2'-O-methyluridine, m7g/7-methylguanosine, Yw/Wybutosine, Mam5u/5-methylaminomethyluridine, or X/3-(3-amino-3-carboxypropyl)uridine, (acp3)u.

B. Nucleosides

[0055] As used herein, a “nucleoside” refers to an individual chemical unit comprising a nucleobase covalently attached to a nucleobase linker moiety. A non-limiting example of a “nucleobase linker moiety” is a sugar comprising 5-carbon atoms (*i.e.*, a “5-carbon sugar”), including but not limited to a deoxyribose, a ribose, an arabinose, or a derivative or an analog of a 5-carbon sugar. Non-limiting examples of a derivative or an analog of a 5-carbon sugar include a 2'-fluoro-2'-deoxyribose or a carbocyclic sugar where a carbon is substituted for an oxygen atom, in the sugar ring. As used herein, a “moiety” generally refers to a smaller chemical or molecular component of a larger chemical or molecular structure.

[0056] Different types of covalent attachment(s) of a nucleobase to a nucleobase linker moiety are known in the art. By way of non-limiting example, a nucleoside comprising a purine (*i.e.*, A or G) or a 7-deazapurine nucleobase typically comprises a covalent attachment of the 9 position of the purine or 7-deazapurine to a 1'-position of a 5-carbon sugar. In another non-limiting example, a nucleoside comprising a pyrimidine nucleobase (*i.e.*, C, T, or U) typically comprises a covalent attachment of the 1 position of the pyrimidine to a 1'-position of a 5-carbon sugar (Kornberg and Baker, 1992).

C. Nucleotides

[0057] As used herein, a “nucleotide” refers to a nucleoside further comprising a “backbone linkage.” A backbone linkage generally covalently attaches a nucleotide to another molecule comprising a nucleotide, or to another nucleotide to form a nucleic acid.

The “backbone linkage” in naturally occurring nucleotides typically comprises a phosphate moiety (*e.g.*, a phosphodiester backbone linkage), which is covalently attached to a 5-carbon sugar. The attachment of the backbone moiety typically occurs at either the 3'- or 5'-position of the 5-carbon sugar. However, other types of attachments are known in the art, particularly when a nucleotide comprises derivatives or analogs of a naturally occurring 5-carbon sugar or phosphate moiety.

D. Nucleic Acid Analogs

[0058] A nucleic acid may comprise, or be composed entirely of, a derivative or analog of a nucleobase, a nucleobase linker moiety, and/or backbone linkage that may be present in a naturally occurring nucleic acid. As used herein a “derivative” refers to a chemically modified or altered form of a naturally occurring molecule, while the terms “mimic” or “analog” refer to a molecule that may or may not structurally resemble a naturally occurring molecule or moiety, but possesses similar functions. Nucleobase, nucleoside, and nucleotide analogs or derivatives are well known in the art.

[0059] Non-limiting examples of nucleosides, nucleotides, or nucleic acids comprising 5-carbon sugar and/or backbone linkage derivatives or analogs, include those in U.S. Pat. No. 5,681,947 which describes oligonucleotides comprising purine derivatives that form triple helices with and/or prevent expression of dsDNA; U.S. Pat. Nos. 5,652,099 and 5,763,167 which describe nucleic acids incorporating fluorescent analogs of nucleosides found in DNA or RNA, particularly for use as fluorescent nucleic acids probes; U.S. Pat. No. 5,614,617 which describes oligonucleotide analogs with substitutions on pyrimidine rings that possess enhanced nuclease stability; U.S. Pat. Nos. 5,670,663, 5,872,232 and 5,859,221 which describe oligonucleotide analogs with modified 5-carbon sugars (*i.e.*, modified 2'-deoxyfuranosyl moieties) used in nucleic acid detection; U.S. Pat. No. 5,446,137 which describes oligonucleotides comprising at least one 5-carbon sugar moiety substituted at the 4' position with a substituent other than hydrogen that can be used in hybridization assays; U.S. Pat. No. 5,886,165 which describes oligonucleotides with both deoxyribonucleotides with 3'-5' backbone linkages and ribonucleotides with 2'-5' backbone linkages; U.S. Pat. No. 5,714,606 which describes a modified backbone linkage wherein a 3'-position oxygen of the backbone linkage is replaced by a carbon to enhance the nuclease resistance of nucleic acids; U.S. Pat. No. 5,672,697 which describes oligonucleotides containing one or more 5' methylene phosphonate backbone linkages that enhance nuclease resistance; U.S. Pat. Nos.

5,466,786 and 5,792,847 which describe the linkage of a substituent moiety that may comprise a drug or label to the 2' carbon of an oligonucleotide to provide enhanced nuclease stability and ability to deliver drugs or detection moieties; U.S. Pat. No. 5,223,618 which describes oligonucleotide analogs with a 2 or 3 carbon backbone linkage attaching the 4' position and 3' position of adjacent 5-carbon sugar moiety to enhanced cellular uptake, resistance to nucleases, and hybridization to target RNA; U.S. Pat. No. 5,470,967 which describes oligonucleotides comprising at least one sulfamate or sulfamide backbone linkage that are useful as nucleic acid hybridization probes; U.S. Pat. Nos. 5,378,825, 5,777,092, 5,623,070, 5,610,289 and 5,602,240 which describe oligonucleotides with a three or four atom backbone linkage moiety replacing the phosphodiester backbone linkage used for improved nuclease resistance, cellular uptake, and regulating RNA expression; U.S. Pat. No. 5,858,988 which describes hydrophobic carrier agent attached to the 2'-O position of oligonucleotides to enhance their membrane permeability and stability; U.S. Pat. No. 5,214,136 which describes oligonucleotides conjugated to anthraquinone at the 5' terminus that possess enhanced hybridization to DNA or RNA; enhanced stability to nucleases; U.S. Pat. No. 5,700,922 which describes PNA-DNA-PNA chimeras wherein the DNA comprises 2'-deoxy-erythro-pentofaranosyl nucleotides for enhanced nuclease resistance, binding affinity, and ability to activate RNase H; U.S. Pat. No. 5,708,154 which describes RNA linked to a DNA to form a DNA-RNA hybrid; U.S. Pat. No. 5,908,845 which describes polyether nucleic acids wherein one or more nucleobases are linked to chiral carbon atoms in a polyether backbone; U.S. Pat. Nos. 5,786,461, 5,891,625, 5,786,461, 5,773,571, 5,766,855, 5,736,336, 5,719,262, 5,714,331, 5,539,082, and WO 92/20702 which describe peptide nucleic acids (PNA or peptide-based nucleic acid analog; or PENAM) that generally comprise one or more nucleotides or nucleosides that comprise a nucleobase moiety, a nucleobase linker moiety that is not a 5-carbon sugar (*e.g.*, aza nitrogen atoms, amido and/or ureido tethers), and/or a backbone linkage that is not a phosphate backbone linkage (*e.g.*, aminoethylglycine, polyamide, polyethyl, polythioamide, polysulfonamide, or polysulfonamide backbone linkage); and U.S. Pat. No. 5,855,911 which describes the hydrophobic, nuclease resistant P-ethoxy backbone linkage.

[0060] Other modifications and uses of nucleic acid analogs are known in the art, and it is anticipated that these techniques and types of nucleic acid analogs may be used with the present invention.

E. Preparation of Nucleic Acids

[0061] A nucleic acid may be made by any technique known to one of ordinary skill in the art, such as chemical synthesis, enzymatic production or biological production. Non-limiting examples of a synthetic nucleic acid (*e.g.*, a synthetic oligonucleotide) include a nucleic acid made by *in vitro* chemical synthesis using phosphotriester, phosphite, or phosphoramidite chemistry and solid phase techniques, such as described in EP 266,032, incorporated herein by reference, or by deoxynucleoside H-phosphonate intermediates as described by Froehler *et al.* (1986) and U.S. Pat. No. 5,705,629, each incorporated herein by reference. In the methods of the present invention, one or more species of oligonucleotide may be used. Various mechanisms of oligonucleotide synthesis have been disclosed in, for example, U.S. Pat. Nos. 4,659,774, 4,816,571, 5,141,813, 5,264,566, 4,959,463, 5,428,148, 5,554,744, 5,574,146, 5,602,244, each of which is incorporated herein by reference.

F. Purification of Nucleic Acids

[0062] A nucleic acid may be purified on polyacrylamide gels, cesium chloride centrifugation gradients, or by any other means known to one of ordinary skill in the art (see for example, Sambrook *et al.* (2001), incorporated herein by reference).

[0063] In certain embodiments, the present invention concerns a nucleic acid that is an isolated nucleic acid. As used herein, the term “isolated nucleic acid” refers to a nucleic acid molecule (*e.g.*, an RNA or DNA molecule) that has been isolated free of, or is otherwise free of, the bulk of the total genomic and transcribed nucleic acids of one or more cells. In certain embodiments, “isolated nucleic acid” refers to a nucleic acid that has been isolated free of, or is otherwise free of, the bulk of cellular components or *in vitro* reaction components, such as, for example, macromolecules, such as lipids or proteins, small biological molecules, and the like.

G. Hybridization

[0064] As used herein, “hybridization,” “hybridize(s),” or “capable of hybridizing” is understood to mean the forming of a double or triple stranded molecule or a molecule with partial double or triple stranded nature. The term “anneal” as used herein is synonymous with “hybridize.”

[0065] As used herein “stringent condition(s)” or “high stringency” are those conditions that allow hybridization between or within one or more nucleic acid strand(s) containing complementary sequence(s), but precludes hybridization of random sequences. Stringent conditions tolerate little, if any, mismatch between a nucleic acid and a target strand. Such conditions are well known to those of ordinary skill in the art, and are preferred for applications requiring high selectivity.

[0066] Stringent conditions may comprise low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.15 M NaCl at temperatures of about 50°C to about 70°C. It is understood that the temperature and ionic strength of a desired stringency are determined in part by the length of the particular nucleic acid(s), the length and nucleobase content of the target sequence(s), the charge composition of the nucleic acid(s), and to the presence or concentration of formamide, tetramethylammonium chloride, or other solvent(s) in a hybridization mixture.

[0067] It is also understood that these ranges, compositions and conditions for hybridization are mentioned by way of non-limiting examples only, and that the desired stringency for a particular hybridization reaction is often determined empirically by comparison to one or more positive or negative controls. Depending on the application envisioned it is preferred to employ varying conditions of hybridization to achieve varying degrees of selectivity of a nucleic acid towards a target sequence. In a non-limiting example, identification or isolation of a related target nucleic acid that does not hybridize to a nucleic acid under stringent conditions may be achieved by hybridization at low temperature and/or high ionic strength. Such conditions are termed “low stringency” or “low stringency conditions,” and non-limiting examples of low stringency include hybridization performed at about 0.15 M to about 0.9 M NaCl at a temperature range of about 20°C to about 50°C. Of course, it is within the skill of one in the art to further modify the low or high stringency conditions to suit a particular application.

IV. Method of Manufacturing Liposomal P-ethoxy Antisense Drug Product

[0068] Antisense oligonucleotides (oligos) complementary to specific regions of a target mRNA have been used to inhibit the expression of endogenous genes. When the antisense oligonucleotides bind to a target mRNA, a DNA-RNA hybrid is formed. This hybrid formation inhibits the translation of the mRNA and, thus, the expression of the

encoded protein. If the protein is essential for the survival of the cell, the inhibition of its expression may lead to cell death. Therefore, antisense oligonucleotides can be useful tools in anticancer and antiviral therapies.

[0069] The main obstacles in using antisense oligonucleotides to inhibit gene expression are cellular instability, low cellular uptake, and poor intercellular delivery. Natural phosphodiester are not resistant to nuclease hydrolysis; thus high concentrations of antisense oligonucleotides are needed before any inhibitory effect is observed. Modified phosphodiester analogs, such as P-ethoxy, have been made to overcome this nuclease hydrolysis problem, but they have not provided a satisfactory solution to the problem.

[0070] The cellular uptake of antisense oligonucleotides is low. To solve this problem, physical techniques, such as calcium-phosphate precipitation, DEAE-dextran mediation, or electroporation, have been used to increase the cellular uptake of oligonucleotides. These techniques are difficult to reproduce and are inapplicable *in vivo*. Cationic lipids, such as Lipofectin, have also been used to deliver oligonucleotides. An electrostatic interaction is formed between the cationic lipids and the negatively charged oligonucleotides, which results in a complex that is then taken up by the target cells. Since these cationic lipids do not protect the oligonucleotides from nuclease digestion and are harmful to the cell membrane, they are only useful in delivering the nuclease-resistant phosphorothioates, but not the nuclease-cleavable phosphodiesters.

[0071] Another modified phosphodiester analog that has been prepared is P-ethoxy. The P-ethoxy antisense backbone does not have an adverse effect on bleeding and complement activation, which are some of the toxicities that have been reported for other antisense analogs. The modifications of P-ethoxy oligonucleotides are made in the phosphate backbone so that the modification will not interfere with the binding of these oligonucleotides to a target mRNA. P-ethoxy oligonucleotides are made by adding an ethyl group to the non-bridging oxygen atom of the phosphate backbone, thus rendering these oligonucleotides uncharged compounds. In spite of their resistance to nucleases, the cellular uptake and intracellular delivery of P-ethoxy oligonucleotides is poor because upon internalization, these oligonucleotides remain sequestered inside the endosomal/lysosomal vacuoles, impeding their access to target mRNA.

A. P-ethoxy antisense drug product

[0072] The liposomal P-ethoxy antisense drug product is composed of two cGMP products, both of which have a FDA-required Certificate of Analysis with FDA-approved release criteria. The raw materials, solvents, and final drug product are described herein. When manufactured, the drug product is a lyophilized crystal or powder of amber or white color that comprises the following materials: oligonucleotide (*e.g.*, P-ethoxy antisense drug substance), neutral lipids (*e.g.*, DOPC), and surfactant (*e.g.*, polysorbate 20). In preparation for administration to a patient, normal saline is added to the vial, at which time liposomes are formed with the P-ethoxy antisense incorporated into the interior.

B. P-ethoxy antisense drug substance

[0073] Specific physical properties (*e.g.*, solubility and hydrophobicity, which then affect drug product solubility in saline, incorporation of oligo into liposomes, and liposome particle size) of the finished product can be defined using a pre-determined P-ethoxy and phosphodiester amidite raw material mix during production of the P-ethoxy antisense drug substance. While loss of the P-ethoxy backbone group randomly occurs during oligonucleotide manufacturing resulting in phosphodiester bonds at those linkages, that loss may not generate the preferred ratio of P-ethoxy : phosphodiester backbone linkage within the oligonucleotide. In this case, the mix of P-ethoxy and phosphodiester amidite raw material supplements the expected value of P-ethoxy backbone deletions, thus generating an oligonucleotide with the desired ratio. Increasing the number of P-ethoxy molecules in the backbone of the oligonucleotide causes the molecule to be more hydrophobic (which results in larger liposome particles; Table 1), less polar, and less soluble (Table 2). Methods of testing the charge-neutral, hydrophobic P-ethoxy drug substance include mass spectrometry to determine the distribution of oligonucleotide lengths and assays to determine the solubility of drug substance, which for practical purposes for solubility is a visual inspection of the drug product reconstituted in saline. As the oligonucleotide becomes less soluble due to a greater number of P-ethoxy backbone linkages the reconstituted solution becomes whiter until particulates form as hydrophobicity becomes too high.

[0074] Formulation must use a particle size, wherein the 90% value is less than 5000 nm in size and is soluble, which is a function of the nucleotide composition. By way of example, if an oligonucleotide is 18-20 nucleotides in length, then at least five of the phosphate backbone linkages should be phosphodiester backbone linkages. This is supported

by the Experiments 7-10 below in Table 1, which provides data from 18mer oligonucleotides.. Wherein if an oligonucleotide is 25 nucleotides in length, then at least six of the phosphate backbone linkages should be phosphodiester backbone linkages.

Table 1. Liposome Particle Size Variability with Antisense Backbone Composition

Experiment	Engineered Antisense Backbone	Post-Manufacturing Backbone Ethyl Deletion		Particle Size Characteristics: Cumulative Distribution Function		
		Principal Peak ^d	Composite Deletion ^e	90% Value (nm) **	50% Value (nm)	300 nm Value (%)
1	3 amidite substitution	-6	-5.67	2130	911	15.30
2	3 amidite substitution	-6	-5.67	2420	1004	15.50
3	3 amidite substitution	-6	-6.12	3682	943	15.50
4	3 amidite substitution	-7	-6.66	3805	978	14.60
5	100% P-ethoxy	-5	-5.66	3924	976	16.00
6	2 amidite substitution	-5	-5.32	4387	1888	11.60
7 ^a	100% P-ethoxy	-4	-4.22	5057	1131	17.70
8	100% P-ethoxy	-4	-4.52	5659	1359	10.00
9 ^b	100% P-ethoxy	-4	-4.38	7571	1909	2.60
10 ^c	100% P-ethoxy	-4	-4.38	7994	1653	14.40

** Drug product release criteria is for 90% of the liposome particles to be less than or equal to 5000 nm.

a. This lot was discarded due to poor solubility; specifically, antisense particles in the reconstituted solution.

b. This lot had lower DMSO and tBA volume with 2 mg antisense in a 20 mL vial, which added an additional component to liposome enlargement.

c. This lot was not released because it failed the particle size release spec.

d. The principal peak represents the most common number of p-ethoxy deletions in the oligonucleotides of the population.

e. The composite deletion represents the average number of p-ethoxy deletions in the population of oligonucleotides.

Table 2. Liposome Particle Solubility with Antisense Backbone Composition

Experiment	Engineered	Post-Manufacturing Backbone Ethyl Deletion		Drug Solubility	
		Principal	Composite	Visual	Solubility

	Antisense Backbone	Peak	Deletion	Observation **	Assessment
1	3 amidite substitution	-6	-5.67	skim milk solution	good
2	3 amidite substitution	-6	-5.67	skim milk solution	good
3	3 amidite substitution	-6	-6.12	skim milk solution	good
4	3 amidite substitution	-7	-6.66	skim milk solution	good
5	100% P-ethoxy	-5	-5.66	skim milk solution	good
6	2 amidite substitution	-5	-5.32	skim milk solution	good
7	100% P-ethoxy	-4	-4.52	white solution	pass
8 ^b	100% P-ethoxy	-4	-4.38	white solution	pass
9 ^c	100% P-ethoxy	-4	-4.38	white solution	pass
10 ^a	100% P-ethoxy	-4	-4.22	white solution particles	fail

** If the drug product sample has particles the lot will be rejected

a. This lot was discarded due to poor solubility; specifically, antisense particles in the reconstituted solution.

b. This lot had lower DMSO and tBA volume with 2 mg antisense in a 20 mL vial, which added an additional component to liposome enlargement.

c. This lot was not released because it failed the particle size release spec.

C. Formulation, filtration, and lyophilization of liposomal P-ethoxy antisense drug product

[0075] One gram (1 g) of pE oligos is dissolved in DMSO at a ratio of 10 mg oligonucleotide per 1 mL DMSO. Next, DOPC is added to tert-butyl alcohol at a ratio of 1 g DOPC per 1719 mL of tert-butyl alcohol. The oligo and DOPC are combined and mixed at a ratio of 1 g oligonucleotide per 2.67 g DOPC. Then, 20 mL of a 0.835% (v/v) solution of polysorbate 20 is added to the mixture resulting in a final concentration of 0.039 mg/mL. The solution is passed through a sterile filter prior to dispensing into glass vials for lyophilization.

[0076] The effect of the surfactant on liposome particle size was determined by titrating the amount of surfactant (Table 3). In the absence of polysorbate 20, only 2.8% of the particles had a diameter of 300 nm or less. In the presence of 1x polysorbate 20, 12.5%

of the particles had a diameter of 300 nm or less. With the addition of 3x-10x polysorbate 20, around 20% of the particles had a diameter of 300 nm or less. Thus an increase in surfactant from 1x to 3x results in a decrease in particle size.

Table 3. Liposome Particle Size Variability with Surfactant

Experiment	Amount of Surfactant	Particle Size Characteristics: Cumulative Distribution Function		
		50% Value	90% Value **	300 nm Value
1	0x	5301 nm	10719 nm	2.8%
2	1x	1053 nm	4054 nm	12.5%
3	3x	785 nm	2926 nm	19.1%
4	5x	721 nm	2691 nm	21.9%
5	10x	734 nm	2937 nm	21.4%

** Drug product release criteria is for 90% of the liposome particles to be less than or equal to 5000 nm.

D. Preparation of liposomal P-ethoxy antisense drug product for administration

[0077] The lyophilized preparation was hydrated with normal saline (0.9%/10 mM NaCl) at a final oligo concentration of 10-5000 μ M. The liposomal-P-ethoxy oligos were mixed by hand shaking.

E. Methods of Testing Liposomal P-ethoxy Antisense Drug Product

[0078] *Visual Inspection of Manufactured Drug Product:* After manufacturing, a sample vial containing drug product is selected and visually inspected. The absence of liquid is mandatory, and then amber crystals at the bottom of the vial are acceptable, and increasing in acceptance to a white, flocculated powder or appearance, the best result. The white appearance indicates a better drying process, with a high surface area to mass ratio, which is very conducive to reconstitution for use.

[0079] *Visual Inspection of Reconstituted Drug Ready for Patient IV:* Normal saline is added to a vial containing the manufactured Liposomal P-ethoxy Antisense Drug Product and shaken to reconstitute into a solution with the drug crystal or powder completely dissolved. Three main observations are made: 1) that the crystal or powder is completely dissolved, 2) there are no white clumps of undissolved material, and 3) the appearance is a milky white or skim milk appearance. The bluer the appearance of the reconstituted liquid, the better, as this signals a smaller liposome particle size that reflects light in the blue spectrum.

[0080] *Mass Spectrometry:* Mass spectrometry (mass spec) is used to display the profile of the various masses in a sample. When P-ethoxy antisense material is produced, a mass spec is run on the sample. The result shows peaks of material present on a grid that has increasing mass on the “x” axis to the right, and relative mass abundance on the “y” axis increasing upward. The profile from a sample is analyzed to determine the relative quantity of P-ethoxy backbones in the P-ethoxy sample, recognizing that the profile of peaks represents (starting farthest to the right), full length material with all backbones comprised of the P-ethoxy linkage, the next peak moving left a full length with one backbone with a P-ethoxy deletion (and therefore, the ethyl being knocked off and the result being a normal phosphodiester backbone linkage), and continuing. The mass spec pattern shifted to the right represents a P-ethoxy sample having more P-ethoxy backbones, and therefore having the properties of being more hydrophobic and less soluble; and likewise, shifted to the left having the opposite effects. Inspection of the mass spec chart of a sample also can be used to determine if filtration during manufacturing produces any adverse effects on oligonucleotide composition present in the filtered drug product.

[0081] *UV Testing:* Ultraviolet light testing is used to determine the mass of oligonucleotide present in a sample. Oligonucleotides absorb light in the 260 nanometer range. As a result, UV testing of the finished reconstituted drug product has come to be used as a method in determining the quantity of oligonucleotide drug substance in a vial of drug product. In terms of manufacturing development and innovations, UV testing was used to determine if there were problems experienced during filtration in manufacturing or poor solubility of the P-ethoxy antisense drug substance, resulting in less oligonucleotide in solution and therefore a lower UV reading. The method will be validated and likely become part of the final product release testing.

[0082] *Liposome Particle Size:* A vial of finished drug product is reconstituted and tested for liposome particle size. The result is often a roughly normal distribution, having a central point, tails and average values or a roughly normal distribution of the majority of the particles and smaller, secondary peaks of the smaller liposomes particles resulting from second-order particle formation effects. It is important that liposome particles not be too large, as they may create adverse effects in patients (for example, create blood flow problems in smaller blood vessels in the lungs). As a result, the drug product release criteria include that particle size testing show that 90% of liposomes be 5 microns or less in size. In addition,

smaller liposomes are preferred because they will have better uptake into cells, and secondly, smaller liposomes can penetrate vascular pores, thereby allowing the liposomes to penetrate inside tumors, increasing treatment effectiveness of a Liposomal P-ethoxy Antisense Drug Product.

V. Methods of Treatment

[0083] Certain aspects of the present invention provide an oligonucleotide–lipid complex (*e.g.*, an oligonucleotide incorporated into a non-charged liposome) for treating diseases, such as cancer, autoimmune disease, or infectious disease. Particularly, the oligonucleotide may have a sequence that allows for base pairing with a human nucleotide sequence and thus may inhibit the expression of a protein encoded by the human nucleotide sequence.

[0084] “Treatment” and “treating” refer to administration or application of a therapeutic agent to a subject or performance of a procedure or modality on a subject for the purpose of obtaining a therapeutic benefit of a disease or health-related condition. For example, a treatment may include administration of a pharmaceutically effective amount of an oligonucleotide–lipid complex.

[0085] “Subject” and “patient” refer to either a human or non-human, such as primates, mammals, and vertebrates. In particular embodiments, the subject is a human.

[0086] The term “therapeutic benefit” or “therapeutically effective” as used throughout this application refers to anything that promotes or enhances the well-being of the subject with respect to the medical treatment of this condition. This includes, but is not limited to, a reduction in the frequency or severity of the signs or symptoms of a disease. For example, treatment of cancer may involve, for example, a reduction in the size of a tumor, a reduction in the invasiveness of a tumor, reduction in the growth rate of the cancer, or prevention of metastasis. Treatment of cancer may also refer to prolonging survival of a subject with cancer. Treatment of an autoimmune disease may involve, for example, reducing the expression of a self-antigen against which there is an undesired immune response, inducing tolerance of a self-antigen against which there is an undesired immune response, or inhibiting the immune response towards the self-antigen. Treatment of an infectious disease may involve, for example, eliminate the infectious agent, reduce the level of the infectious agent, or maintain the level of the infectious agent at a certain level.

[0087] Tumors for which the present treatment methods are useful include any malignant cell type, such as those found in a solid tumor, a hematological tumor, metastatic cancer, or non-metastatic cancer. Exemplary solid tumors can include, but are not limited to, a tumor of an organ selected from the group consisting of pancreas, colon, cecum, esophagus, gastrointestinal, gum, liver, skin, stomach, testis, tongue, uterus, stomach, brain, head, neck, ovary, kidney, larynx, sarcoma, bone, lung, bladder, melanoma, prostate, and breast. Exemplary hematological tumors include tumors of the bone marrow, T or B cell malignancies, leukemias, lymphomas, such as, for example, diffuse large B cell lymphoma, blastomas, myelomas, and the like. Further examples of cancers that may be treated using the methods provided herein include, but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, leukemia, squamous cell cancer, lung cancer (including small-cell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, and squamous carcinoma of the lung), cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer (including gastrointestinal cancer and gastrointestinal stromal cancer), pancreatic cancer (including pancreatic ductal adenocarcinoma), glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, vulval cancer, thyroid cancer, various types of head and neck cancer, melanoma, superficial spreading melanoma, lentigo malignant melanoma, acral lentiginous melanomas, nodular melanomas, as well as B-cell lymphoma (including low grade/follicular non-Hodgkin's lymphoma (NHL); small lymphocytic (SL) NHL; intermediate grade/follicular NHL; intermediate grade diffuse NHL; high grade immunoblastic NHL; high grade lymphoblastic NHL; high grade small non-cleaved cell NHL; bulky disease NHL; diffuse large B-cell lymphoma; mantle cell lymphoma; AIDS-related lymphoma; and Waldenstrom's macroglobulinemia), chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), Hairy cell leukemia, multiple myeloma, acute myeloid leukemia (AML) and chronic myeloblastic leukemia.

[0088] The cancer may specifically be of the following histological type, though it is not limited to these: neoplasm, malignant; carcinoma; carcinoma, undifferentiated; giant and spindle cell carcinoma; small cell carcinoma; papillary carcinoma; squamous cell carcinoma; lymphoepithelial carcinoma; basal cell carcinoma; pilomatrix carcinoma; transitional cell carcinoma; papillary transitional cell carcinoma; adenocarcinoma; gastrinoma, malignant; cholangiocarcinoma; hepatocellular carcinoma; combined hepatocellular carcinoma and cholangiocarcinoma; trabecular adenocarcinoma; adenoid cystic carcinoma; adenocarcinoma

in adenomatous polyp; adenocarcinoma, familial polyposis coli; solid carcinoma; carcinoid tumor, malignant; branchiolo-alveolar adenocarcinoma; papillary adenocarcinoma; chromophobe carcinoma; acidophil carcinoma; oxyphilic adenocarcinoma; basophil carcinoma; clear cell adenocarcinoma; granular cell carcinoma; follicular adenocarcinoma; papillary and follicular adenocarcinoma; nonencapsulating sclerosing carcinoma; adrenal cortical carcinoma; endometroid carcinoma; skin appendage carcinoma; apocrine adenocarcinoma; sebaceous adenocarcinoma; ceruminous adenocarcinoma; mucoepidermoid carcinoma; cystadenocarcinoma; papillary cystadenocarcinoma; papillary serous cystadenocarcinoma; mucinous cystadenocarcinoma; mucinous adenocarcinoma; signet ring cell carcinoma; infiltrating duct carcinoma; medullary carcinoma; lobular carcinoma; inflammatory carcinoma; paget's disease, mammary; acinar cell carcinoma; adenosquamous carcinoma; adenocarcinoma w/squamous metaplasia; thymoma, malignant; ovarian stromal tumor, malignant; thecoma, malignant; granulosa cell tumor, malignant; androblastoma, malignant; sertoli cell carcinoma; leydig cell tumor, malignant; lipid cell tumor, malignant; paraganglioma, malignant; extra-mammary paraganglioma, malignant; pheochromocytoma; glomangiosarcoma; malignant melanoma; amelanotic melanoma; superficial spreading melanoma; malignant melanoma in giant pigmented nevus; epithelioid cell melanoma; blue nevus, malignant; sarcoma; fibrosarcoma; fibrous histiocytoma, malignant; myxosarcoma; liposarcoma; leiomyosarcoma; rhabdomyosarcoma; embryonal rhabdomyosarcoma; alveolar rhabdomyosarcoma; stromal sarcoma; mixed tumor, malignant; mullerian mixed tumor; nephroblastoma; hepatoblastoma; carcinosarcoma; mesenchymoma, malignant; brenner tumor, malignant; phyllodes tumor, malignant; synovial sarcoma; mesothelioma, malignant; dysgerminoma; embryonal carcinoma; teratoma, malignant; struma ovarii, malignant; choriocarcinoma; mesonephroma, malignant; hemangiosarcoma; hemangioendothelioma, malignant; kaposi's sarcoma; hemangiopericytoma, malignant; lymphangiosarcoma; osteosarcoma; juxtacortical osteosarcoma; chondrosarcoma; chondroblastoma, malignant; mesenchymal chondrosarcoma; giant cell tumor of bone; ewing's sarcoma; odontogenic tumor, malignant; ameloblastic odontosarcoma; ameloblastoma, malignant; ameloblastic fibrosarcoma; pinealoma, malignant; chordoma; glioma, malignant; ependymoma; astrocytoma; protoplasmic astrocytoma; fibrillary astrocytoma; astroblastoma; glioblastoma; oligodendroglioma; oligodendroblastoma; primitive neuroectodermal; cerebellar sarcoma; ganglioneuroblastoma; neuroblastoma; retinoblastoma; olfactory neurogenic tumor; meningioma, malignant; neurofibrosarcoma; neurilemmoma, malignant; granular cell tumor, malignant; malignant lymphoma; hodgkin's disease; hodgkin's; paraganuloma; malignant

lymphoma, small lymphocytic; malignant lymphoma, large cell, diffuse; malignant lymphoma, follicular; mycosis fungoides; other specified non-hodgkin's lymphomas; malignant histiocytosis; multiple myeloma; mast cell sarcoma; immunoproliferative small intestinal disease; leukemia; lymphoid leukemia; plasma cell leukemia; erythroleukemia; lymphosarcoma cell leukemia; myeloid leukemia; basophilic leukemia; eosinophilic leukemia; monocytic leukemia; mast cell leukemia; megakaryoblastic leukemia; myeloid sarcoma; and hairy cell leukemia.

[0089] Autoimmune diseases for which the present treatment methods are useful include, without limitation, systemic lupus erythematosus, spondyloarthropathy, ankylosing spondylitis, psoriatic arthritis, reactive arthritis, enteropathic arthritis, diabetes mellitus, celiac disease, autoimmune thyroid disease, autoimmune liver disease, Addison's disease, transplant rejection, graft vs. host disease, host vs. graft disease, ulcerative colitis, Crohn's disease, irritable bowel disease, inflammatory bowel disease, rheumatoid arthritis, juvenile rheumatoid arthritis, familial Mediterranean fever, amyotrophic lateral sclerosis, Sjogren's syndrome, early arthritis, viral arthritis, multiple sclerosis, or psoriasis. The diagnosis and treatment of these diseases are well documented in the literature.

[0090] Infectious diseases for which the present treatment methods are useful include, without limitation, bacterial infections, viral infections, fungal infections, and parasitic infections. Exemplary viral infections include hepatitis B virus, hepatitis C virus, human immunodeficiency virus 1, human immunodeficiency virus 2, human papilloma virus, herpes simplex virus 1, herpes simplex virus 2, herpes zoster, varicella zoster, coxsackievirus A16, cytomegalovirus, ebola virus, enterovirus, Epstein-Barr virus, hanta virus, hendra virus, viral meningitis, respiratory syncytial virus, rotavirus, west nile virus, adenovirus, and influenza virus infections. Exemplary bacterial infections include *Chlamydia trachomatis*, *Listeria monocytogenes*, *Helicobacter pylori*, *Escherichia coli*, *Borrelia burgdorferi*, *Legionella pneumophila*, *Mycobacteria* sps (e.g., *M. tuberculosis*, *M. avium*, *M. intracellulare*, *M. kansasii*, *M. goodii*), *Staphylococcus aureus*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Streptococcus pyogenes* (Group A Streptococcus), *Streptococcus agalactiae* (Group B Streptococcus), *Streptococcus* (viridans group), *Streptococcus faecalis*, *Streptococcus bovis*, *Streptococcus* (anaerobic sps.), *Streptococcus pneumoniae*, pathogenic *Campylobacter* sp., *Enterococcus* sp., *Haemophilus influenzae*, *Bacillus anthracis*, *Corynebacterium diphtheriae*, *Corynebacterium* sp., *Erysipelothrix rhusiopathiae*,

Clostridium perfringens, *Clostridium tetani*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Pasturella multocida*, *Bacteroides* sp., *Fusobacterium nucleatum*, *Streptobacillus moniliformis*, *Treponema pallidum*, *Treponema pertemue*, *Leptospira*, *Rickettsia*, *Actinomyces israeli*, *Shigella* sps (e.g., *S. flexneri*, *S. sonnei*, *S. dysenteriae*), and *Salmonella* spp infections. Exemplary fungal infections include *Candida albicans*, *Candida glabrata*, *Aspergillus fumigatus*, *Aspergillus terreus*, *Cryptococcus neoformans*, *Histoplasma capsulatum*, *Coccidioides immitis*, *Blastomyces dermatitidis*, and *Chlamydia irachomatis* infections.

[0091] The oligonucleotide–lipid complex may be used herein as an antitumor, antiviral, antibacterial, antifungal, antiparasite, or anti-autoimmune agent in a variety of modalities. In a particular embodiment, the invention contemplates methods of using an oligonucleotide–lipid complex comprises contacting a population of diseased cells with a therapeutically effective amount of an oligonucleotide–lipid complex for a time period sufficient to inhibit or reverse disease.

[0092] In one embodiment, the contacting *in vivo* is accomplished by administering, by intravenous, intraperitoneal, subcutaneous, or intratumoral injection, a therapeutically effective amount of a physiologically tolerable composition comprising an oligonucleotide–lipid complex of this invention to a patient. The oligonucleotide–lipid complex can be administered parenterally by injection or by gradual infusion over time.

[0093] Therapeutic compositions comprising oligonucleotide–lipid complex are conventionally administered intravenously or subcutaneously, such as by injection of a unit dose, for example. The term “unit dose” when used in reference to a therapeutic composition refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent, *i.e.*, carrier, or vehicle.

[0094] The compositions are administered in a manner compatible with the dosage formulation, and in a therapeutically effective amount. The quantity to be administered depends on the subject to be treated, capacity of the subject's system to utilize the active ingredient, and degree of therapeutic effect desired. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual. However, suitable dosage ranges for systemic application are

disclosed herein and depend on the route of administration. Suitable regimes for initial and booster administration are also contemplated and are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration. Exemplary multiple administrations are described herein and are particularly preferred to maintain continuously high serum and tissue levels of polypeptide. Alternatively, continuous intravenous infusion sufficient to maintain concentrations in the blood in the ranges specified for *in vivo* therapies are contemplated.

[0095] It is contemplated that an oligonucleotide of the invention can be administered systemically or locally to treat disease, such as to inhibit tumor cell growth or to kill cancer cells in cancer patients with locally advanced or metastatic cancers. They can be administered intravenously, intrathecally, subcutaneously, and/or intraperitoneally. They can be administered alone or in combination with anti-proliferative drugs. In one embodiment, they are administered to reduce the cancer load in the patient prior to surgery or other procedures. Alternatively, they can be administered after surgery to ensure that any remaining cancer (*e.g.*, cancer that the surgery failed to eliminate) does not survive.

[0096] A therapeutically effective amount of an oligonucleotide is a predetermined amount calculated to achieve the desired effect, *i.e.*, to inhibit the expression of a target protein. Thus, the dosage ranges for the administration of oligonucleotides of the invention are those large enough to produce the desired effect. The dosage should not be so large as to cause adverse side effects, such as hyperviscosity syndromes, pulmonary edema, congestive heart failure, neurological effects, and the like. Generally, the dosage will vary with age of, condition of, sex of, and extent of the disease in the patient and can be determined by one of skill in the art. The dosage can be adjusted by the individual physician in the event of any complication.

[0097] A composition of the present invention is preferably administered to a patient parenterally, for example by intravenous, intraarterial, intramuscular, intralymphatic, intraperitoneal, subcutaneous, intrapleural, or intrathecal injection, or may be used *ex vivo*. Preferred dosages are between 5-25 mg/kg. The administration is preferably repeated on a timed schedule until the cancer disappears or regresses, and may be in conjunction with other forms of therapy.

VI. Pharmaceutical Preparations

[0098] A pharmaceutical composition comprising the liposomes will usually include a sterile, pharmaceutically acceptable carrier or diluent, such as dextrose or saline solution.

[0099] Where clinical application of non-charged lipid component (*e.g.*, in the form of a liposome) containing an oligonucleotide is undertaken, it will generally be beneficial to prepare the lipid complex as a pharmaceutical composition appropriate for the intended application. This will typically entail preparing a pharmaceutical composition that is essentially free of pyrogens, as well as any other impurities that could be harmful to humans or animals. One may also employ appropriate buffers to render the complex stable and allow for uptake by target cells.

[00100] The phrases “pharmaceutical or pharmacologically acceptable” refers to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, such as a human, as appropriate. The preparation of a pharmaceutical composition that contains at least one non-charged lipid component comprising an oligonucleotide or additional active ingredient will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington: The Science and Practice of Pharmacy, 21st, 2005, incorporated herein by reference. Moreover, for animal (*e.g.*, human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards.

[00101] As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (*e.g.*, antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art. A pharmaceutically acceptable carrier is preferably formulated for administration to a human, although in certain embodiments it may be desirable to use a pharmaceutically acceptable carrier that is formulated for administration to a non-human animal but which would not be acceptable (*e.g.*, due to governmental regulations) for administration to a human. Except insofar as any

conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated.

[00102] The actual dosage amount of a composition of the present invention administered to a patient or subject can be determined by physical and physiological factors such as body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the patient and on the route of administration. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject.

[00103] In certain embodiments, pharmaceutical compositions may comprise, for example, at least about 0.1% of an active compound. In other embodiments, the an active compound may comprise between about 2% to about 75% of the weight of the unit, or between about 25% to about 60%, for example, and any range derivable therein. In other non-limiting examples, a dose may also comprise from about 1 microgram/kg/body weight, about 5 microgram/kg/body weight, about 10 microgram/kg/body weight, about 50 microgram/kg/body weight, about 100 microgram/kg/body weight, about 200 microgram/kg/body weight, about 350 microgram/kg/body weight, about 500 microgram/kg/body weight, about 1 milligram/kg/body weight, about 5 milligram/kg/body weight, about 10 milligram/kg/body weight, about 50 milligram/kg/body weight, about 100 milligram/kg/body weight, about 200 milligram/kg/body weight, about 350 milligram/kg/body weight, about 500 milligram/kg/body weight, to about 1000 mg/kg/body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 5 µg/kg/body weight to about 1000 mg/kg/body weight, about 5 microgram/kg/body weight to about 500 milligram/kg/body weight, etc., can be administered.

[00104] An oligonucleotide of the present embodiments may be administered in a dose of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100 or more µg of nucleic acid per dose. Each dose may be in a volume of 1, 10, 50, 100, 200, 500, 1000 or more µl or ml.

[00105] Solutions of therapeutic compositions can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions also can be

prepared in glycerol, liquid polyethylene glycols, mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[00106] The therapeutic compositions of the present invention are advantageously administered in the form of injectable compositions either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. These preparations also may be emulsified. A typical composition for such purpose comprises a pharmaceutically acceptable carrier. For instance, the composition may contain 10 mg, 25 mg, 50 mg or up to about 100 mg of human serum albumin per milliliter of phosphate buffered saline. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like.

[00107] Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, saline solutions, parenteral vehicles such as sodium chloride, Ringer's dextrose, etc. Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial agents, anti-oxidants, chelating agents and inert gases. The pH and exact concentration of the various components the pharmaceutical composition are adjusted according to well known parameters.

[00108] The therapeutic compositions of the present invention may include classic pharmaceutical preparations. Administration of therapeutic compositions according to the present invention will be via any common route so long as the target tissue is available via that route. This includes oral, nasal, buccal, rectal, vaginal or topical. Topical administration may be particularly advantageous for the treatment of skin cancers, to prevent chemotherapy-induced alopecia or other dermal hyperproliferative disorder. Alternatively, administration may be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients. For treatment of conditions of the lungs, aerosol delivery can be used. Volume of the aerosol is between about 0.01 ml and 0.5 ml.

[00109] An effective amount of the therapeutic composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete

units suitable for use in a subject, each unit containing a predetermined-quantity of the therapeutic composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and treatment regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the protection or effect desired.

[00110] Precise amounts of the therapeutic composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting the dose include the physical and clinical state of the patient, the route of administration, the intended goal of treatment (*e.g.*, alleviation of symptoms versus cure) and the potency, stability and toxicity of the particular therapeutic substance.

VII. Combination Treatments

[00111] In certain embodiments, the compositions and methods of the present invention involve an inhibitory oligonucleotide, or oligonucleotide capable of expressing an inhibitor of gene expression, in combination with a second or additional therapy. The methods and compositions including combination therapies enhance the therapeutic or protective effect, and/or increase the therapeutic effect of another anti-cancer or anti-hyperproliferative therapy. Therapeutic and prophylactic methods and compositions can be provided in a combined amount effective to achieve the desired effect, such as the killing of a cancer cell and/or the inhibition of cellular hyperproliferation. This process may involve contacting the cells with both an inhibitor of gene expression and a second therapy. A tissue, tumor, or cell can be contacted with one or more compositions or pharmacological formulation(s) including one or more of the agents (i.e., inhibitor of gene expression or an anti-cancer agent), or by contacting the tissue, tumor, and/or cell with two or more distinct compositions or formulations, wherein one composition provides 1) an inhibitory oligonucleotide; 2) an anti-cancer agent, or 3) both an inhibitory oligonucleotide and an anti-cancer agent. Also, it is contemplated that such a combination therapy can be used in conjunction with a chemotherapy, radiotherapy, surgical therapy, or immunotherapy.

[00112] An inhibitory oligonucleotide may be administered before, during, after or in various combinations relative to an anti-cancer treatment. The administrations may be in intervals ranging from concurrently to minutes to days to weeks. In embodiments where the inhibitory oligonucleotide is provided to a patient separately from an anti-cancer agent,

one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the two compounds would still be able to exert an advantageously combined effect on the patient. In such instances, it is contemplated that one may provide a patient with the inhibitory oligonucleotide therapy and the anti-cancer therapy within about 12 to 24 or 72 h of each other and, more preferably, within about 6-12 h of each other. In some situations it may be desirable to extend the time period for treatment significantly where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between respective administrations.

[00113] In certain embodiments, a course of treatment will last 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90 days or more. It is contemplated that one agent may be given on day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, and/or 90, any combination thereof, and another agent is given on day 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, and/or 90, or any combination thereof. Within a single day (24-hour period), the patient may be given one or multiple administrations of the agent(s). Moreover, after a course of treatment, it is contemplated that there is a period of time at which no anti-cancer treatment is administered. This time period may last 1, 2, 3, 4, 5, 6, 7 days, and/or 1, 2, 3, 4, 5 weeks, and/or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 months or more, depending on the condition of the patient, such as their prognosis, strength, health, etc.

[00114] Various combinations may be employed. For the example below an inhibitory oligonucleotide therapy is “A” and an anti-cancer therapy or an autoimmune therapy is “B”:

A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B/B

B/A/B/B B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A

B/B/A/A B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A

A/A/B/A

[00115] Administration of any compound or therapy of the present invention to a patient will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the agents. Therefore, in some embodiments there is a step of monitoring toxicity that is attributable to combination therapy. It is expected that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies, as well as surgical intervention, may be applied in combination with the described therapy.

[00116] In specific aspects, it is contemplated that a standard therapy will include chemotherapy, radiotherapy, immunotherapy, surgical therapy or gene therapy and may be employed in combination with the inhibitor of gene expression therapy, anticancer therapy, or both the inhibitor of gene expression therapy and the anti-cancer therapy, as described herein.

A. Chemotherapy

[00117] A wide variety of chemotherapeutic agents may be used in accordance with the present embodiments. The term “chemotherapy” refers to the use of drugs to treat cancer. A “chemotherapeutic agent” is used to connote a compound or composition that is administered in the treatment of cancer. These agents or drugs are categorized by their mode of activity within a cell, for example, whether and at what stage they affect the cell cycle. Alternatively, an agent may be characterized based on its ability to directly cross-link DNA, to intercalate into DNA, or to induce chromosomal and mitotic aberrations by affecting nucleic acid synthesis.

[00118] Examples of chemotherapeutic agents include alkylating agents, such as thiotepa and cyclophosphamide; alkyl sulfonates, such as busulfan, improsulfan, and piposulfan; aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines, including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide, and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and

cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards, such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, and uracil mustard; nitrosureas, such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics, such as the enediyne antibiotics (*e.g.*, calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin, anthracycline, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, such as mitomycin C, mycophenolic acid, nogalarnycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, and zorubicin; anti-metabolites, such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues, such as denopterin, pteropterin, and trimetrexate; purine analogs, such as fludarabine, 6-mercaptopurine, thiamiprine, and thioguanine; pyrimidine analogs, such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, and floxuridine; androgens, such as calusterone, dromostanolone propionate, epitostanol, mepitostane, and testolactone; anti-adrenals, such as mitotane and trilostane; folic acid replenisher, such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids, such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSKpolysaccharide complex; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; taxoids, *e.g.*, paclitaxel and docetaxel gemcitabine; 6-thioguanine; mercaptopurine; platinum coordination complexes, such as

cisplatin, oxaliplatin, and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (*e.g.*, CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids, such as retinoic acid; capecitabine; carboplatin, procarbazine, plicomycin, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatin, and pharmaceutically acceptable salts, acids, or derivatives of any of the above.

B. Radiotherapy

[00119] Other factors that cause DNA damage and have been used extensively include what are commonly known as γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also contemplated such as microwaves, proton beam irradiation (U.S. Pat. Nos. 5,760,395 and 4,870,287) and UV-irradiation. It is most likely that all of these factors affect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

[00120] The terms “contacted” and “exposed,” when applied to a cell, are used herein to describe the process by which a therapeutic construct and a chemotherapeutic or radiotherapeutic agent are delivered to a target cell or are placed in direct juxtaposition with the target cell. To achieve cell killing, for example, both agents are delivered to a cell in a combined amount effective to kill the cell or prevent it from dividing.

C. Immunotherapy

[00121] In the context of cancer treatment, immunotherapeutics, generally, rely on the use of immune effector cells and molecules to target and destroy cancer cells. Trastuzumab (Herceptin™) is such an example. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually affect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve merely as a targeting agent. Alternatively,

the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells. The combination of therapeutic modalities, i.e., direct cytotoxic activity and inhibition or reduction of ErbB2 would provide therapeutic benefit in the treatment of ErbB2 overexpressing cancers.

[00122] Another immunotherapy could also be used as part of a combined therapy with gene silencing therapy discussed above. In one aspect of immunotherapy, the tumor cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumor markers exist and any of these may be suitable for targeting in the context of the present invention. Common tumor markers include carcinoembryonic antigen, prostate specific antigen, urinary tumor associated antigen, fetal antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, estrogen receptor, laminin receptor, erb B and p155. An alternative aspect of immunotherapy is to combine anticancer effects with immune stimulatory effects. Immune stimulating molecules also exist including: cytokines such as IL-2, IL-4, IL-12, GM-CSF, gamma-IFN, chemokines such as MIP-1, MCP-1, IL-8 and growth factors such as FLT3 ligand. Combining immune stimulating molecules, either as proteins or using gene delivery in combination with a tumor suppressor has been shown to enhance anti-tumor effects. Moreover, antibodies against any of these compounds can be used to target the anti-cancer agents discussed herein.

[00123] Examples of immunotherapies currently under investigation or in use are immune adjuvants e.g., Mycobacterium bovis, Plasmodium falciparum, dinitrochlorobenzene and aromatic compounds (U.S. Pat. Nos. 5,801,005 and 5,739,169; Hui and Hashimoto, 1998; Christodoulides et al., 1998), cytokine therapy, e.g., interferons α , β and γ ; IL-1, GM-CSF and TNF (Bukowski et al., 1998; Davidson et al., 1998; Hellstrand et al., 1998) gene therapy, e.g., TNF, IL-1, IL-2, p53 (Qin et al., 1998; Austin-Ward and Villaseca, 1998; U.S. Pat. Nos. 5,830,880 and 5,846,945) and monoclonal antibodies, e.g., anti-ganglioside GM2, anti-HER-2, anti-p185 (Pietras et al., 1998; Hanibuchi et al., 1998; U.S. Pat. No. 5,824,311). It is contemplated that one or more anti-cancer therapies may be employed with the gene silencing therapies described herein.

[00124] In active immunotherapy, an antigenic peptide, polypeptide or protein, or an autologous or allogenic tumor cell composition or “vaccine” is administered, generally

with a distinct bacterial adjuvant (Ravindranath and Morton, 1991; Morton et al., 1992; Mitchell et al., 1990; Mitchell et al., 1993).

[00125] In adoptive immunotherapy, the patient's circulating lymphocytes, or tumor infiltrated lymphocytes, are isolated in vitro, activated by lymphokines such as IL-2 or transduced with genes for tumor necrosis, and readministered (Rosenberg et al., 1988; 1989).

[00126] In some embodiments, the immunotherapy may be an immune checkpoint inhibitor. Immune checkpoints either turn up a signal (*e.g.*, co-stimulatory molecules) or turn down a signal. Inhibitory immune checkpoints that may be targeted by immune checkpoint blockade include adenosine A2A receptor (A2AR), B7-H3 (also known as CD276), B and T lymphocyte attenuator (BTLA), cytotoxic T-lymphocyte-associated protein 4 (CTLA-4, also known as CD152), indoleamine 2,3-dioxygenase (IDO), killer-cell immunoglobulin (KIR), lymphocyte activation gene-3 (LAG3), programmed death 1 (PD-1), T-cell immunoglobulin domain and mucin domain 3 (TIM-3) and V-domain Ig suppressor of T cell activation (VISTA). In particular, the immune checkpoint inhibitors target the PD-1 axis and/or CTLA-4.

[00127] The immune checkpoint inhibitors may be drugs such as small molecules, recombinant forms of ligand or receptors, or, in particular, are antibodies, such as human antibodies (*e.g.*, International Patent Publication WO2015016718; Pardoll, *Nat Rev Cancer*, 12(4): 252-64, 2012; both incorporated herein by reference). Known inhibitors of the immune checkpoint proteins or analogs thereof may be used, in particular chimerized, humanized or human forms of antibodies may be used. As the skilled person will know, alternative and/or equivalent names may be in use for certain antibodies mentioned in the present disclosure. Such alternative and/or equivalent names are interchangeable in the context of the present disclosure. For example, it is known that lambrolizumab is also known under the alternative and equivalent names MK-3475 and pembrolizumab.

[00128] In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its ligand binding partners. In a specific aspect, the PD-1 ligand binding partners are PDL1 and/or PDL2. In another embodiment, a PDL1 binding antagonist is a molecule that inhibits the binding of PDL1 to its binding partners. In a specific aspect, PDL1 binding partners are PD-1 and/or B7-1. In another embodiment, the PDL2 binding antagonist is a molecule that inhibits the binding of PDL2 to its binding partners. In a

specific aspect, a PDL2 binding partner is PD-1. The antagonist may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide. Exemplary antibodies are described in U.S. Patent Nos. 8,735,553, 8,354,509, and 8,008,449, all incorporated herein by reference. Other PD-1 axis antagonists for use in the methods provided herein are known in the art such as described in U.S. Patent Publication Nos. 20140294898, 2014022021, and 20110008369, all incorporated herein by reference.

[00129] In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody (*e.g.*, a human antibody, a humanized antibody, or a chimeric antibody). In some embodiments, the anti-PD-1 antibody is selected from the group consisting of nivolumab, pembrolizumab, and CT-011. In some embodiments, the PD-1 binding antagonist is an immunoadhesin (*e.g.*, an immunoadhesin comprising an extracellular or PD-1 binding portion of PDL1 or PDL2 fused to a constant region (*e.g.*, an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 binding antagonist is AMP- 224. Nivolumab, also known as MDX-1106-04, MDX-1106, ONO-4538, BMS-936558, and OPDIVO[®], is an anti-PD-1 antibody described in WO2006/121168. Pembrolizumab, also known as MK-3475, Merck 3475, lambrolizumab, KEYTRUDA[®], and SCH-900475, is an anti-PD-1 antibody described in WO2009/114335. CT-011, also known as hBAT or hBAT-1, is an anti-PD-1 antibody described in WO2009/101611. AMP-224, also known as B7-DCIg, is a PDL2-Fc fusion soluble receptor described in WO2010/027827 and WO2011/066342.

[00130] Another immune checkpoint that can be targeted in the methods provided herein is the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), also known as CD152. The complete cDNA sequence of human CTLA-4 has the Genbank accession number L15006. CTLA-4 is found on the surface of T cells and acts as an “off” switch when bound to CD80 or CD86 on the surface of antigen-presenting cells. CTLA4 is a member of the immunoglobulin superfamily that is expressed on the surface of Helper T cells and transmits an inhibitory signal to T cells. CTLA4 is similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to CD80 and CD86, also called B7-1 and B7-2 respectively, on antigen-presenting cells. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T cells and may be important to their function. T cell activation through the T cell receptor and CD28 leads to increased expression of CTLA-4, an inhibitory receptor for B7 molecules.

[00131] In some embodiments, the immune checkpoint inhibitor is an anti-CTLA-4 antibody (*e.g.*, a human antibody, a humanized antibody, or a chimeric antibody), an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide.

[00132] Anti-human-CTLA-4 antibodies (or VH and/or VL domains derived therefrom) suitable for use in the present methods can be generated using methods well known in the art. Alternatively, art recognized anti-CTLA-4 antibodies can be used. For example, the anti-CTLA-4 antibodies disclosed in: US Patent No. 8,119,129, WO 01/14424, WO 98/42752; WO 00/37504 (CP675,206, also known as tremelimumab; formerly ticilimumab), U.S. Patent No. 6,207,156; Hurwitz *et al.* (1998) *Proc Natl Acad Sci USA* 95(17): 10067-10071; Camacho *et al.* (2004) *J Clin Oncology* 22(145): Abstract No. 2505 (antibody CP-675206); and Mokyr *et al.* (1998) *Cancer Res* 58:5301-5304 can be used in the methods disclosed herein. The teachings of each of the aforementioned publications are hereby incorporated by reference. Antibodies that compete with any of these art-recognized antibodies for binding to CTLA-4 also can be used. For example, a humanized CTLA-4 antibody is described in International Patent Application No. WO2001014424, WO2000037504, and U.S. Patent No. 8,017,114; all incorporated herein by reference.

[00133] An exemplary anti-CTLA-4 antibody is ipilimumab (also known as 10D1, MDX- 010, MDX- 101, and Yervoy®) or antigen binding fragments and variants thereof (see, *e.g.*, WO 01/14424). In other embodiments, the antibody comprises the heavy and light chain CDRs or VRs of ipilimumab. Accordingly, in one embodiment, the antibody comprises the CDR1, CDR2, and CDR3 domains of the VH region of ipilimumab, and the CDR1, CDR2 and CDR3 domains of the VL region of ipilimumab. In another embodiment, the antibody competes for binding with and/or binds to the same epitope on CTLA-4 as the above-mentioned antibodies. In another embodiment, the antibody has at least about 90% variable region amino acid sequence identity with the above-mentioned antibodies (*e.g.*, at least about 90%, 95%, or 99% variable region identity with ipilimumab).

[00134] Other molecules for modulating CTLA-4 include CTLA-4 ligands and receptors such as described in U.S. Patent Nos. 5844905, 5885796 and International Patent Application Nos. WO1995001994 and WO1998042752; all incorporated herein by reference, and immunoadhesins such as described in U.S. Patent No. 8329867, incorporated herein by reference.

[00135] In some embodiment, the immune therapy could be adoptive immunotherapy, which involves the transfer of autologous antigen-specific T cells generated *ex vivo*. The T cells used for adoptive immunotherapy can be generated either by expansion of antigen-specific T cells or redirection of T cells through genetic engineering (Park, Rosenberg et al. 2011). Isolation and transfer of tumor specific T cells has been shown to be successful in treating melanoma. Novel specificities in T cells have been successfully generated through the genetic transfer of transgenic T cell receptors or chimeric antigen receptors (CARs) (Jena, Dotti et al. 2010). CARs are synthetic receptors consisting of a targeting moiety that is associated with one or more signaling domains in a single fusion molecule. In general, the binding moiety of a CAR consists of an antigen-binding domain of a single-chain antibody (scFv), comprising the light and variable fragments of a monoclonal antibody joined by a flexible linker. Binding moieties based on receptor or ligand domains have also been used successfully. The signaling domains for first generation CARs are derived from the cytoplasmic region of the CD3zeta or the Fc receptor gamma chains. CARs have successfully allowed T cells to be redirected against antigens expressed at the surface of tumor cells from various malignancies including lymphomas and solid tumors (Jena, Dotti et al. 2010).

[00136] In one embodiment, the present application provides for a combination therapy for the treatment of cancer wherein the combination therapy comprises adoptive T-cell therapy and a checkpoint inhibitor. In one aspect, the adoptive T-cell therapy comprises autologous and/or allogenic T cells. In another aspect, the autologous and/or allogenic T cells are targeted against tumor antigens.

D. Surgery

[00137] Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative, and palliative surgery. Curative surgery is a cancer treatment that may be used in conjunction with other therapies, such as the treatment of the present invention, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy and/or alternative therapies.

[00138] Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and microscopically controlled surgery

(Mohs' surgery). It is further contemplated that the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

[00139] Upon excision of part or all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

E. Other Agents

[00140] It is contemplated that other agents may be used in combination with certain aspects of the present embodiments to improve the therapeutic efficacy of treatment. These additional agents include agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhesion, agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers, or other biological agents. Increases in intercellular signaling by elevating the number of GAP junctions would increase the anti-hyperproliferative effects on the neighboring hyperproliferative cell population. In other embodiments, cytostatic or differentiation agents can be used in combination with certain aspects of the present embodiments to improve the anti-hyperproliferative efficacy of the treatments. Inhibitors of cell adhesion are contemplated to improve the efficacy of the present embodiments. Examples of cell adhesion inhibitors are focal adhesion kinase (FAKs) inhibitors and Lovastatin. It is further contemplated that other agents that increase the sensitivity of a hyperproliferative cell to apoptosis, such as the antibody c225, could be used in combination with certain aspects of the present embodiments to improve the treatment efficacy.

VIII. Kits and Diagnostics

[00141] In various aspects of the invention, a kit is envisioned containing therapeutic agents and/or other therapeutic and delivery agents. In some embodiments, the present invention contemplates a kit for preparing and/or administering a therapy of the invention. The kit may comprise reagents capable of use in administering an active or effective agent(s) of the invention. Reagents of the kit may include at least one inhibitor of

gene expression (*e.g.*, a STAT3 oligonucleotides), one or more lipid component, one or more anti-cancer component of a combination therapy, as well as reagents to prepare, formulate, and/or administer the components of the invention or perform one or more steps of the inventive methods.

[00142] In some embodiments, the kit may also comprise a suitable container means, which is a container that will not react with components of the kit, such as an eppendorf tube, an assay plate, a syringe, a bottle, or a tube. The container may be made from sterilizable materials such as plastic or glass.

[00143] The kit may further include an instruction sheet that outlines the procedural steps of the methods, and will follow substantially the same procedures as described herein or are known to those of ordinary skill.

IX. Examples

[00144] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1 – STAT3-targeted P-ethoxy oligonucleotides

[00145] Oligonucleotides targeting *STAT3* were designed for use in a liposomal *STAT3* antisense drug product to inhibit the expression of STAT3. The contiguous cDNA sequence of *STAT3* is provided in SEQ ID NO: 5 and the protein sequence of STAT3 is provided in SEQ ID NO: 6. The sequence of each of the oligonucleotides is provided in Table 4.

Table 4. *STAT3* antisense sequences

SEQ ID NO:	Sequence
------------	----------

1	5' - CAA AGT GGC ATG TGA TTC -3'
2	5' - GCT ATA CTG CTG GTC AAT -3'
3	5' - CTT CGT AGA TTG TGC TGA -3'
4	5' - CTG ATA ATT CAA CTC AGG -3'

[00146] The liposomal *STAT3* antisense drug product was manufactured according to the methods described herein. Mass spectrometry testing showed that about 80% of the oligonucleotide drug substance had between three and seven phosphodiester backbone linkages. Particle testing showed that 90% of the liposomes had a particle size diameter of 2319 nm or less, 50% of the liposomes had a particle size diameter of 368 nm or less, and about 18% of the liposomes had a particle size diameter of 300 nm or less.

Example 2 – Inhibition of *STAT3* expression with P-ethoxy oligonucleotides

[00147] The ability of liposomal *STAT3* antisense to inhibit the expression of *STAT3* was tested in two lung adenocarcinoma cell lines, H1975 and HCC 827. Liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 (180 µg/mL) or an equivalent amount of empty liposomes was incubated with each cell line for four days. As the data in FIG. 1 show, incubation with liposomal *STAT3* antisense, but not empty liposomes, reduced the level of *STAT3* expression relative to β-actin.

Example 3 – Inhibition of cancer cell viability by liposomal *STAT3* antisense

[00148] The ability of liposomal *STAT3* antisense to inhibit the viability of non-small cell lung cancer cells was tested in three lung adenocarcinoma cell lines: H1975, HCC 827, and H358. Liposomal *STAT3* antisense corresponding to one of SEQ ID NOs: 1-4 was incubated with each cell line for four days. As the data in FIGs. 2A-C show, incubation with liposomal *STAT3* antisense reduced the cell viability of each cell line, with liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 being the most potent.

[00149] The ability of liposomal *STAT3* antisense to inhibit the growth of leukemia cells was tested in three human leukemia cell lines: K-562, MV4-11, and Kasumi-1. Liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 was incubated with each cell line for four days. As the data in FIG. 3A show, incubation with liposomal *STAT3* antisense reduced the cell viability of each cell line.

[00150] The ability of liposomal *STAT3* antisense to inhibit the growth of lymphoma cells was tested in five human lymphoma cell lines: DOHH-2, SU-DHL-4, SU-DHL-6, SU-DHL-10, and RL. Liposomal *STAT3* antisense corresponding to SEQ ID NO: 4 was incubated with each cell line for four days. As the data in FIG. 3B show, incubation with liposomal *STAT3* antisense reduced the cell viability of each cell line.

* * *

[00151] All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Pat. No. 4,659,774
U.S. Pat. No. 4,816,571
U.S. Pat. No. 4,870,287
U.S. Pat. No. 4,959,463
U.S. Pat. No. 5,141,813
U.S. Pat. No. 5,214,136
U.S. Pat. No. 5,223,618
U.S. Pat. No. 5,264,566
U.S. Pat. No. 5,378,825
U.S. Pat. No. 5,428,148
U.S. Pat. No. 5,446,137
U.S. Pat. No. 5,466,786
U.S. Pat. No. 5,470,967
U.S. Pat. No. 5,539,082
U.S. Pat. No. 5,554,744
U.S. Pat. No. 5,574,146
U.S. Pat. No. 5,602,240
U.S. Pat. No. 5,602,244
U.S. Pat. No. 5,610,289
U.S. Pat. No. 5,614,617
U.S. Pat. No. 5,623,070
U.S. Pat. No. 5,652,099
U.S. Pat. No. 5,670,663
U.S. Pat. No. 5,672,697
U.S. Pat. No. 5,681,947
U.S. Pat. No. 5,700,922
U.S. Pat. No. 5,705,629
U.S. Pat. No. 5,708,154

U.S. Pat. No. 5,714,331
U.S. Pat. No. 5,714,606
U.S. Pat. No. 5,719,262
U.S. Pat. No. 5,736,336
U.S. Pat. No. 5,739,169
U.S. Pat. No. 5,760,395
U.S. Pat. No. 5,763,167
U.S. Pat. No. 5,766,855
U.S. Pat. No. 5,773,571
U.S. Pat. No. 5,777,092
U.S. Pat. No. 5,786,461
U.S. Pat. No. 5,792,847
U.S. Pat. No. 5,801,005
U.S. Pat. No. 5,824,311
U.S. Pat. No. 5,830,880
U.S. Pat. No. 5,846,945
U.S. Pat. No. 5,855,911
U.S. Pat. No. 5,858,988
U.S. Pat. No. 5,859,221
U.S. Pat. No. 5,872,232
U.S. Pat. No. 5,886,165
U.S. Pat. No. 5,891,625
U.S. Pat. No. 5,908,845
U.S. Pat. No. 9,744,187

Amin et al., *Oncogene*, 22:5399-5407, 2013.

Austin-Ward and Villaseca, *Revista Medica de Chile*, 126(7):838-845, 1998.

Bailey and Sullivan, *Biochimica. Biophys. Acts.*, 239-252, 2000.

Bangham et al., *J. Mol. Biol.*, 13(1):253-259, 1965.

Bukowski et al., *Clinical Cancer Res.*, 4(10):2337-2347, 1998.

Christodoulides et al., *Microbiology*, 144(Pt 11):3027-3037, 1998.

Davidson et al., *J. Immunother.*, 21(5):389-398, 1998.

Deamer and Uster, In: *Liposome Preparation: Methods and Mechanisms*, Ostro (Ed.),
Liposomes, 1983.

- Dokka et al., *Pharm Res*, 17: 521-25, 2000.
- duBois et al., *J Clin Oncol*, 17: 46-51, 1999.
- Dubey et al., *J. Drug Target*, 12:257-264, 2004.
- Duxbury et al., *Biochem. Biophys. Res. Commun.*, 311:786-792, 2003.
- Duxbury et al., *Oncogene*, 23:1448-1456, 2004.
- Egholm et al., *Nature*, 365(6446):566-568, 1993.
- Elbashir et al., *Nature*, 411 (6836):494-498, 2001.
- European Appln. 01219
- European Appln. 266,032
- Fagard et al., *JAKSTAT*, 2:e22882, 2013.
- Farhood et al., *Biochim. Biophys. Act*, 289-295, 1995.
- Fire et al., *Nature*, 391(6669):806-811, 1998.
- Flenniken et al., *Dev. Biol.*, 179:382-401, 1996.
- Froehler et al., *Nucleic Acids Res.*, 14(13):5399-5407, 1986.
- Gabizon, *Cancer Invest.*, 19:424-436, 2001.
- Ghosh and Bachhawat, In: *Liver Diseases, Targeted Diagnosis and Therapy Using Specific Receptors and Ligands*, Wu et al. (Eds.), Marcel Dekker, NY, 87-104, 1991.
- Gregoriadis, In: *Drug Carriers in Biology and Medicine*, Gregoriadis (Ed.), 287-341, 1979.
- Gutierrez-Puente et al., *J. Pharmacol. Exp. Ther.*, 291:865-869, 1999.
- Halder et al., *Clinical Cancer Research*, 11: 8829-36, 2005.
- Han et al., *Ann Surg Oncol*, 4:264-268, 1997.
- Hanibuchi et al., *Int. J. Cancer*, 78(4):480-485, 1998.
- Hannon and Rossi, *Nature*, 431:371-378, 2004.
- Hardee et al., *G3 (Bethesda)* 3:2173-2185, 2013.
- Hassani et al., *J. Gene Med.*, 7(2):198-207, 2005.
- Hecker et al., *Cancer Research*, 62:2699-2707, 2002.
- Hellstrand et al., *Acta Oncologica*, 37(4):347-353, 1998.
- Hortobagyi et al., *J. Clin. Oncol.*, 19:3422-3433, 2001.
- Hsia et al., *J Cell Biol*, 160:753-67, 2003.
- Hui and Hashimoto, *Infection Immun.*, 66(11):5329-5336, 1998.
- Jackson et al., *Nat. Biotechnol.*, 21:635-637, 2003.
- Jemal et al., *CA Cancer J. Clin.*, 55(1):10-30, 2005.
- Judson et al., *Cancer*, 86: 1551-56, 1999.
- Kaneda et al., *Science*, 243:375-378, 1989.

- Kato et al., J. Biol. Chem., 266:3361-3364, 1991.
- Kim et al., Nat. Biotechnol., 22:321-325, 2004.
- Kinch et al., Clin. Exp. Metastasis, 20:59-68, 2003.
- Klein et al., Gastroenterology, 125:9-18, 2003.
- Kohnno et al., Int J Cancer, 97:336-43, 2002.
- Kornberg and Baker, DNA Replication, 2nd Ed., Freeman, San Francisco, 1992.
- Kornberg et al., Invest Ophthalmol Vis Sci, 45:4463-69, 2004.
- Kornberg, Head Neck, 20: 634-639, 1998.
- Kostarelos et al., Int J Cancer, 112: 713-21, 2004.
- Krasnici et al., Int. J. Cancer, 105(4):561-567, 2003.
- Landen, Cancer Res, 65: 6910-18, 2005.
- Langley et al., Cancer Research, 63: 2971-76, 2003.
- Lewis et al., Cell, 115:787-798, 2003.
- Lewis et al., Nat. Genet., 32:107-108, 2002.
- Lori et al., Am. J. Pharmacogenomics, 2:245-252, 2002.
- Matsuda et al., Proc. Natl. Acad. Sci. USA, 101:16-22, 2004.
- McCaffrey et al., Nature, 418:38-39, 2002.
- McGuire et al., New England Journal of Medicine, 334:1-6, 1996.
- McLean et al., Expert Opin Pharmacother, 4: 227-34, 2003.
- Miklossy et al., Nat. Rev. Drug Discov., 12:611-629, 2013.
- Miller et al., Biochemistry, 37(37):12875-83, 1998.
- Mitchell et al., Ann. NY Acad. Sci., 690:153-166, 1993.
- Mitchell et al., J. Clin. Oncol., 8(5):856-869, 1990.
- Mitra et al., Nature Reviews Molecular Cell Biology, 6: 56-68, 2005.
- Mitra et al., Proc Am Assoc Cancer Res, 2005.
- Morton et al., Arch. Surg., 127:392-399, 1992.
- Nemoto et al., Pathobiology, 65:195-203, 1997.
- Nicolau et al., Methods Enzymol., 149:157-176, 1987.
- Noblitt et al., Cancer Gene Ther., 11:757-766, 2004.
- Ogawa et al, Oncogene, 19:6043-6052, 2000.
- Owens et al., Cancer Res, 55:2752-2755, 1995.
- Park et al., Cancer Lett., 118:153-160, 1997.
- PCT Appln. WO 92/20702
- PCT Appln. WO02/100435A1

PCT Appln. WO03/015757A1

PCT Appln. WO04/002453A1

PCT Appln. WO04029213A2

Pietras et al., *Oncogene*, 17(17):2235-2249, 1998.

Qin et al., *Proc. Natl. Acad. Sci. USA*, 95(24):14411-14416, 1998.

Ravindranath and Morton, *Intern. Rev. Immunol.*, 7: 303-329, 1991.

Reich et al., *Mol. Vis.*, 9:210-216, 2003.

Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1289-1329, 1990.

Rosenberg et al., *Ann. Surg.* 210(4):474-548, 1989.

Rosenberg et al., *N. Engl. J. Med.*, 319:1676, 1988.

Ryther et al., *Gene Ther.*, 12(1):5-11, 2004.

Sambrook et al., In: *Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001.

Schaller and Parsons, *Trends in Cell Biology*, 3:258-62, 1993.

Schaller et al., *Mol Biol Cell*, 10:3489-3505, 1999.

Schaller, *Biochim Biophys Acta*, 1540:1-21, 2001.

Schaller, *J Endocrinol*, 150:1-7, 1996.

Schaller, *Trends Cell Biol*, 3:258-262, 1993.

Scheit, In: *Synthesis and Biological Function*, Wiley-Interscience, NY, 171-172, 1980.

Schlaepfer and Hunter, *Trends in Cell Biology*, 8: 151-57, 1998.

Schlaepfer et al., *Prog Biophys Mol Biol*, 71: 435-78, 1999.

Scuto et al., *Cancer Res.*, 71:3182-3188, 2011.

Sein et al., *Oncogene*, 19: 5539-42, 2000.

Sheta et al., *J Natl Cancer inst*, 92: 1065-73, 2000.

Shibata et al., *Cancer Res*, 58: 900-903, 1998.

Sieg et al., *Nat Cell Biol*, 2:249-56, 2000.

Sioud and Sorensen, *Biochem. Biophys. Res. Comm.*, 312:1220-1225, 2003.

Siwak et al., *Clin Cancer Res*, 8: 955-56, 2002.

Sledz et al., *Nat. Cell Biol.*, 5:834-839, 2003.

Song et al., *Nature Med.* 9:347-351, 2003.

Sonoda et al., *Journal of Biological Chemistry*, 275:16309-15, 2000.

Sood et al., *Am J Pathol*, 165:1087-1095, 2004.

Sood et al., *Cancer Biology & Therapy*, 1: 511-17, 2002.

Sorensen et al., *J. Mol. Biol.*, 327:761-66, 2003.

- Soutschek et al., *Nature*, 432:173-178, 2004.
- Spagnou et al., *Biochemistry*, 43:13348-13356, 2004.
- Sulman et al., *Genomics*, 40:371-374, 1997.
- Szoka and Papahadjopoulos, *Proc. Natl. Acad. Sci. USA*, 75:4194-4198, 1978.
- Thaker et al., 36th Annual Meeting of the Society of Gynecologic Oncologists, Miami, Fla., 2005.
- Thaker et al., *Clin. Cancer Res.*, 10:5145-5150, 2004.
- Thurston et al., *J. Clin. Invest.*, 101(7):1401-1413, 1998.
- Uchida et al., *Mol. Ther.*, 10:162-171, 2004.
- Voskoglou-Nomikos et al., *Clin. Cancer Res.*, 9:4227-4239, 2003.
- Walker-Daniels et al., *Prostate*, 41:275-80, 1999.
- Wianny et al., *Nat. Cell Biol.*, 2:70-75, 2000.
- Wong et al., *Gene*, 10:87-94, 1980.
- Wu et al., *J. Hematol. Oncol.*, 4:31, 2011.
- Xia et al., *Nat. Biotechnol.*, 20:1006-10, 2002.
- Yang et al., *Oncogene*, 22:5694-701, 2003.
- Zelinski et al., *Cancer Res.*, 61:2301, 2001.
- Zhang et al., *J. Biol. Chem.*, 279:10677-684, 2004.

WHAT IS CLAIMED IS:

1. A composition comprising a population of oligonucleotides, phospholipids, and a surfactant, wherein the oligonucleotides hybridize to a *STAT3* polynucleotide gene product, wherein oligonucleotides of the population are composed of nucleoside molecules linked together through phosphate backbone linkages, wherein 60% to 75% of the phosphate backbone linkages in each oligonucleotide is a P-ethoxy backbone linkage, and wherein 25% to 40% of the phosphate backbone linkages in each oligonucleotide are phosphodiester backbone linkages.
2. The composition of claim 1, wherein oligonucleotides of the population comprise a sequence according to any one of SEQ ID NOs: 1-4.
3. The composition of claim 2, wherein oligonucleotides of the population comprise a sequence according to SEQ ID NO: 1.
4. The composition of claim 2, wherein oligonucleotides of the population comprise a sequence according to SEQ ID NO: 2.
5. The composition of claim 2, wherein oligonucleotides of the population comprise a sequence according to SEQ ID NO: 3.
6. The composition of claim 2, wherein oligonucleotides of the population comprise a sequence according to SEQ ID NO: 4.
7. The composition of claim 1, wherein the phosphodiester backbone linkages are distributed throughout each oligonucleotide.
8. The composition of claim 1, wherein the phosphodiester backbone linkages are not clustered within a portion of each oligonucleotide.
9. The composition of claim 1, wherein the population of oligonucleotides is heterogeneous as to the number of P-ethoxy backbone linkages and phosphodiester backbone linkages present in the oligonucleotides of the population.
10. The composition of claim 1, wherein the oligonucleotides of the population have a size ranging from 18 to 30 nucleotides.

11. The composition of claim 1, wherein the population of oligonucleotides comprises a single species of oligonucleotides.

12. The composition of claim 1, wherein the population of oligonucleotides comprises at least two species of oligonucleotides.

13. The composition of claim 1, wherein the population of oligonucleotides is heterogeneous as to the distribution of phosphodiester backbone linkages among the oligonucleotides of the population.

14. The composition of claim 1, wherein the oligonucleotides of the population inhibit the expression of STAT3 protein.

15. The composition of claim 1, wherein the oligonucleotides of the population are composed of deoxyribose nucleoside molecules.

16. The composition of claim 1, wherein the surfactant is polysorbate 20.

17. The composition of claim 1, wherein the oligonucleotides and phospholipids form an oligonucleotide–lipid complex.

18. The composition of claim 1, wherein the phospholipids are uncharged or have a neutral charge at physiologic pH.

19. The composition of claim 18, wherein the phospholipids are neutral phospholipids.

20. The composition of claim 19, wherein the neutral phospholipids are phosphatidylcholines.

21. The composition of claim 19, wherein the neutral phospholipids are dioleoylphosphatidyl choline.

22. The composition of claim 1, wherein the phospholipids and oligonucleotides are present at a molar ratio of from about 5:1 to about 100:1.

23. The composition of claim 1, wherein the oligonucleotide–lipid complex is further defined as a population of liposomes.

24. The composition of claim 23, wherein at least 90% of the liposomes are less than 5 microns in diameter.

25. The composition of claim 23, wherein the population of oligonucleotides is incorporated in the population of liposomes.

26. The composition of claim 1, wherein the composition is lyophilized.

27. A method for reducing the expression level of STAT3 protein in a cell comprising contacting the cell with a composition of claim 1.

28. A pharmaceutical composition comprising a composition according to claim 17 and a pharmaceutically acceptable carrier.

29. A method of treating a subject with a cancer comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition of claim 28.

30. Use of the composition of claim 1 in the manufacture of a medicament for treating cancer.

31. The method of claim 29, wherein the subject is a human.

32. The method or use of claim 29 or 30, wherein the cancer is a non-small cell lung cancer, pancreatic adenocarcinoma, breast cancer, prostate cancer, melanoma, colon cancer, leukemia, lymphoma, glioblastoma, osteosarcoma, oral cavity cancer, ovarian cancer, uterine cancer, bone cancer, brain cancer, prostate cancer, kidney cancer, stomach cancer, esophageal cancer, rectal cancer, bladder cancer, testicular cancer, or liver cancer.

33. The method of claim 29, wherein the composition is administered subcutaneously, intravenously, or intraperitoneally.

34. The method of claim 29, further comprising administering at least a second anticancer therapy to the subject.

35. The method of claim 34, wherein the second anticancer therapy is a surgical therapy, chemotherapy, radiation therapy, cryotherapy, hormone therapy, immunotherapy, anti-viral therapy, immune suppression therapy, anti-bacterial therapy, anti-parasite therapy, anti-fungal therapy, or cytokine therapy.

36. The method of claim 29, wherein administration of the composition reduces expression of STAT3 protein in the patient.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B

SEQUENCE LISTING

<110> Bio-Path Holdings, Inc.
 <120> P-ETHOXY NUCLEIC ACIDS FOR STAT3 INHIBITION
 <130> BPHI.P0004WO
 <140> Unknown
 <141> 2018-04-19
 <150> 62/487,292
 <151> 2017-04-19
 <160> 6
 <170> PatentIn version 3.5
 <210> 1
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic polynucleotide
 <400> 1
 caaagtggca tgtgattc 18
 <210> 2
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic polynucleotide
 <400> 2
 gctataactgc tgggtcaat 18
 <210> 3
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic polynucleotide
 <400> 3
 cttcgtagat tgtgctga 18
 <210> 4
 <211> 18
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic polynucleotide
 <400> 4

ctgataattc aactcagg 18

<210> 5

<211> 4978

<212> DNA

<213> Homo sapiens

<400> 5

ggttttccgga gctgcggcgg cgcagactgg gagggggagc cggggggttcc gacgtcgcag	60
ccgaggggaac aagccccaac cggatcctgg acaggcaccc cggcttggcg ctgtctctcc	120
ccctcggctc ggagaggccc ttccggcctga gggagcctcg cggcccgtcc ccggcacacg	180
cgcagccccg gcctctcggc ctctgccgga gaaacagttg ggaccctga ttttagcagg	240
atggcccaat ggaatcagct acagcagctt gacacacggt acctggagca gctccatcag	300
ctctacagtg acagcttccc aatggagctg cggcagtttc tggccccttg gattgagagt	360
caagattggg catatgcggc cagcaaagaa tcacatgcca ctttggtgtt tcataatctc	420
ctgggagaga ttgaccagca gtatagccgc ttctgcaag agtcgaatgt tctctatcag	480
cacaatctac gaagaatcaa gcagtttctt cagagcaggt atcttgagaa gccaatggag	540
attgcccgga ttgtggcccg gtgcctgtgg gaagaatcac gccttctaca gactgcagcc	600
actgcggccc agcaaggggg ccaggccaac cccccacag cagccgtggt gacggagaag	660
cagcagatgc tggagcagca ccttcaggat gtccggaaga gagtgcagga tctagaacag	720
aaaatgaaag tggtagagaa tctccaggat gactttgatt tcaactataa aaccctcaag	780
agtcaaggag acatgcaaga tctgaatgga aacaaccagt cagtgaccag gcagaagatg	840
cagcagctgg aacagatgct cactgcgctg gaccagatgc ggagaagcat cgtgagttag	900
ctggcggggc ttttgtcagc gatggagtac gtgcagaaaa ctctcacgga cgaggagctg	960
gctgactgga agaggcggca acagattgcc tgcattggag gcccgcccaa catctgccta	1020
gatcggctag aaaactggat aacgtcatta gcagaatctc aacttcagac ccgtcaacaa	1080
attaagaaac tggaggagtt gcagcaaaaa gtttcctaca aaggggaccc cattgtacag	1140
caccggccga tgctggagga gagaatcgtg gagctgttta gaaacttaat gaaaagtgcc	1200
tttgtggtgg agcggcagcc ctgcatgccc atgcatcctg accggcccct cgtcatcaag	1260
accggcgtcc agttcactac taaagtcagg ttgctggtca aattccctga gttgaattat	1320
cagcttaaaa ttaaagtgtg cattgacaaa gactctgggg acgttgcagc tctcagagga	1380
tcccggaaat ttaacattct gggcacaaac acaaaagtga tgaacatgga agaatccaac	1440
aacggcagcc tctctgcaga attcaaacac ttgaccctga gggagcagag atgtgggaat	1500
gggggcccag ccaattgtga tgcttcctg attgtgactg aggagctgca cctgatcacc	1560
tttgagaccg aggtgtatca ccaaggcctc aagattgacc tagagacca ctcccttgcca	1620
gttgtggtga tctccaacat ctgtcagatg ccaaatgcct gggcgtccat cctgtggtac	1680

aacatgctga ccaacaatcc caagaatgta aacttttttta ccaagcccc aattggaacc	1740
tgggatcaag tggccgaggt cctgagctgg cagttctcct ccaccaccaa gcgaggactg	1800
agcatcgagc agctgactac actggcagag aaactcttgg gacctggtgt gaattattca	1860
gggtgtcaga tcacatgggc taaattttgc aaagaaaaca tggctggcaa gggcttctcc	1920
ttctgggtct ggctggacaa tatcattgac cttgtgaaaa agtacatcct ggccttttgg	1980
aacgaagggt acatcatggg ctttatcagt aaggagcggg agcgggccat cttgagcact	2040
aagcctccag gcaccttct gctaagattc agtgaaagca gcaaagaagg aggcgtcact	2100
ttcacttggg tggagaagga catcagcggg aagaccaga tccagtccgt ggaaccatac	2160
acaaagcagc agctgaacaa catgtcattt gctgaaatca tcatgggcta taagatcatg	2220
gatgctacca atatcctggg gtctccactg gtctatctct atcctgacat tccaaggag	2280
gaggcattcg gaaagtattg tcggccagag agccaggagc atcctgaagc tgaccaggt	2340
agcgtgccc catacctgaa gaccaagttt atctgtgtga caccaacgac ctgcagcaat	2400
accattgacc tgccgatgtc ccccgact ttagattcat tgatgcagtt tggaaataat	2460
ggtgaagggtg ctgaaccctc agcaggaggg cagtttgagt ccctcacctt tgacatggag	2520
ttgacctcg agtgcgctac ctcccccatg tgaggagctg agaacggaag ctgcagaaaag	2580
atacgactga ggcgcctacc tgcattctgc caccctcac acagccaaac ccagatcat	2640
ctgaaactac taactttgtg gttccagatt ttttttaatc tcctacttct gctatctttg	2700
agcaatctgg gcacttttaa aaatagagaa atgagtgaat gtgggtgatc tgcttttatc	2760
taaatgcaaa taaggatgtg ttctctgaga cccatgatca ggggatgtgg cggggggtgg	2820
ctagagggag aaaaaggaaa tgtcttgtgt tgtttgttc ccctgccctc ctttctcagc	2880
agctttttgt tattgttgtt gttgttctta gacaagtgcc tcctggtgcc tgcggcatcc	2940
ttctgcctgt ttctgtaagc aaatgccaca ggccacctat agctacatac tcctggcatt	3000
gcacttttta accttgctga catccaaata gaagatagga ctatctaagc cctaggtttc	3060
tttttaaatt aagaaataat aacaattaaa gggcaaaaaa cactgtatca gcatagcctt	3120
tctgtattta agaaacttaa gcagccgggc atgggtggctc acgcctgtaa tcccagcact	3180
ttgggaggcc gaggcggatc ataaggtcag gagatcaaga ccatcctggc taacacggtg	3240
aaaccccgtc tctactaaaa gtacaaaaaa ttagctgggt gtggtgggtg gcgcctgtag	3300
tcccagctac tcgggaggct gaggcaggag aatcgcttga acctgagagg cggaggttgc	3360
agtgagccaa aattgcacca ctgcacactg cactccatcc tgggcgacag tctgagactc	3420
tgtctcaaaa aaaaaaaaaa aaaaaagaaa cttcagttaa cagcctcctt ggtgctttaa	3480
gcattcagct tccttcaggc tggtaattta tataatccct gaaacgggct tcaggtaaaa	3540
cccttaagac atctgaagct gcaacctggc ctttggtgtt gaaataggaa ggtttaagga	3600

```

gaatctaagc atttttagact tttttttata aatagactta ttttcctttg taatgtattg 3660
gccttttagt gagtaaggct gggcagaggg tgcttacaac cttgactccc tttctccctg 3720
gacttgatct gctgtttcag aggctagggt gtttctgtgg gtgccttatc agggctggga 3780
tactttgat tctggcttcc ttctgcccc accctcccga cccagtccc cctgatcctg 3840
ctagaggcat gtctccttgc gtgtctaaag gtccctcatc ctgtttgttt taggaatcct 3900
ggctctcagga cctcatggaa gaagaggggg agagagttac aggttggaaca tgatgcacac 3960
tatggggccc cagcgacgtg tctgggtgag ctcagggaat atggttctta gccagtttct 4020
tggtgatata cagtggcact tgtaatggcg tcttcattca gttcatgcag ggcaaaggct 4080
tactgataaa cttgagtctg ccctcgatat aggggtgtata cctggcctcc ctctgaggct 4140
ggtgactcct ccctgctggg gccccacagg tgaggcagaa cagctagagg gcctccccgc 4200
ctgcccgcct tggctggcta gctcgctct cctgtgcgta tgggaacacc tagcacgtgc 4260
tggatgggct gcctctgact cagaggcatg gccggatttg gcaactcaaa accaccttgc 4320
ctcagctgat cagagtttct gtggaattct gtttgttaaa tcaaattagc tggctctctga 4380
attaaggggg agacgacctt ctctaagatg aacaggggtc gcccagtc tctgctgg 4440
agacagttga tgtgtcatgc agagctctta cttctccagc aacactcttc agtacataat 4500
aagcttaact gataaacaga atatttagaa aggtgagact tgggcttacc attgggttta 4560
aatcataggg acctagggcg agggttcagg gcttctctgg agcagatatt gtcaagttca 4620
tggccttagg tagcatgtat ctggtcttaa ctctgattgt agcaaaagtt ctgagaggag 4680
ctgagccctg ttgtggccca ttaaagaaca gggtcctcag gccctgcccg cttcctgtcc 4740
actgccccct ccccatcccc agcccagccg aggggaatccc gtgggttgct tacctaccta 4800
taaggtggtt tataagctgc tgtcctggcc actgcattca aattccaatg tgtacttcat 4860
agtgtaaaaa tttatattat tgtgaggttt tttgtctttt tttttttttt ttttttttgg 4920
tatattgctg tatctacttt aacttccaga aataaacgtt atataggaac cgtaaaaa 4978

```

```

<210> 6
<211> 770
<212> PRT
<213> Homo sapiens

```

```
<400> 6
```

```

Met Ala Gln Trp Asn Gln Leu Gln Gln Leu Asp Thr Arg Tyr Leu Glu
1              5              10              15

```

```

Gln Leu His Gln Leu Tyr Ser Asp Ser Phe Pro Met Glu Leu Arg Gln
20              25              30

```

```
Phe Leu Ala Pro Trp Ile Glu Ser Gln Asp Trp Ala Tyr Ala Ala Ser
```

35					40					45					
Lys	Glu	Ser	His	Ala	Thr	Leu	Val	Phe	His	Asn	Leu	Leu	Gly	Glu	Ile
50						55				60					
Asp	Gln	Gln	Tyr	Ser	Arg	Phe	Leu	Gln	Glu	Ser	Asn	Val	Leu	Tyr	Gln
65					70					75					80
His	Asn	Leu	Arg	Arg	Ile	Lys	Gln	Phe	Leu	Gln	Ser	Arg	Tyr	Leu	Glu
			85						90					95	
Lys	Pro	Met	Glu	Ile	Ala	Arg	Ile	Val	Ala	Arg	Cys	Leu	Trp	Glu	Glu
			100					105					110		
Ser	Arg	Leu	Leu	Gln	Thr	Ala	Ala	Thr	Ala	Ala	Gln	Gln	Gly	Gly	Gln
		115					120					125			
Ala	Asn	His	Pro	Thr	Ala	Ala	Val	Val	Thr	Glu	Lys	Gln	Gln	Met	Leu
	130					135					140				
Glu	Gln	His	Leu	Gln	Asp	Val	Arg	Lys	Arg	Val	Gln	Asp	Leu	Glu	Gln
145					150					155					160
Lys	Met	Lys	Val	Val	Glu	Asn	Leu	Gln	Asp	Asp	Phe	Asp	Phe	Asn	Tyr
				165					170					175	
Lys	Thr	Leu	Lys	Ser	Gln	Gly	Asp	Met	Gln	Asp	Leu	Asn	Gly	Asn	Asn
		180						185					190		
Gln	Ser	Val	Thr	Arg	Gln	Lys	Met	Gln	Gln	Leu	Glu	Gln	Met	Leu	Thr
		195					200						205		
Ala	Leu	Asp	Gln	Met	Arg	Arg	Ser	Ile	Val	Ser	Glu	Leu	Ala	Gly	Leu
	210					215					220				
Leu	Ser	Ala	Met	Glu	Tyr	Val	Gln	Lys	Thr	Leu	Thr	Asp	Glu	Glu	Leu
225					230					235					240
Ala	Asp	Trp	Lys	Arg	Arg	Gln	Gln	Ile	Ala	Cys	Ile	Gly	Gly	Pro	Pro
				245					250					255	
Asn	Ile	Cys	Leu	Asp	Arg	Leu	Glu	Asn	Trp	Ile	Thr	Ser	Leu	Ala	Glu
			260					265					270		
Ser	Gln	Leu	Gln	Thr	Arg	Gln	Gln	Ile	Lys	Lys	Leu	Glu	Glu	Leu	Gln
		275					280					285			
Gln	Lys	Val	Ser	Tyr	Lys	Gly	Asp	Pro	Ile	Val	Gln	His	Arg	Pro	Met

290																	
Leu	Glu	Glu	Arg	Ile	Val	Glu	Leu	Phe	Arg	Asn	Leu	Met	Lys	Ser	Ala		
305					310					315					320		
Phe	Val	Val	Glu	Arg	Gln	Pro	Cys	Met	Pro	Met	His	Pro	Asp	Arg	Pro		
				325					330					335			
Leu	Val	Ile	Lys	Thr	Gly	Val	Gln	Phe	Thr	Thr	Lys	Val	Arg	Leu	Leu		
			340					345					350				
Val	Lys	Phe	Pro	Glu	Leu	Asn	Tyr	Gln	Leu	Lys	Ile	Lys	Val	Cys	Ile		
		355					360					365					
Asp	Lys	Asp	Ser	Gly	Asp	Val	Ala	Ala	Leu	Arg	Gly	Ser	Arg	Lys	Phe		
	370					375					380						
Asn	Ile	Leu	Gly	Thr	Asn	Thr	Lys	Val	Met	Asn	Met	Glu	Glu	Ser	Asn		
385					390					395					400		
Asn	Gly	Ser	Leu	Ser	Ala	Glu	Phe	Lys	His	Leu	Thr	Leu	Arg	Glu	Gln		
				405					410					415			
Arg	Cys	Gly	Asn	Gly	Gly	Arg	Ala	Asn	Cys	Asp	Ala	Ser	Leu	Ile	Val		
			420					425					430				
Thr	Glu	Glu	Leu	His	Leu	Ile	Thr	Phe	Glu	Thr	Glu	Val	Tyr	His	Gln		
		435					440					445					
Gly	Leu	Lys	Ile	Asp	Leu	Glu	Thr	His	Ser	Leu	Pro	Val	Val	Val	Ile		
	450					455					460						
Ser	Asn	Ile	Cys	Gln	Met	Pro	Asn	Ala	Trp	Ala	Ser	Ile	Leu	Trp	Tyr		
465					470				475						480		
Asn	Met	Leu	Thr	Asn	Asn	Pro	Lys	Asn	Val	Asn	Phe	Phe	Thr	Lys	Pro		
				485					490					495			
Pro	Ile	Gly	Thr	Trp	Asp	Gln	Val	Ala	Glu	Val	Leu	Ser	Trp	Gln	Phe		
			500					505					510				
Ser	Ser	Thr	Thr	Lys	Arg	Gly	Leu	Ser	Ile	Glu	Gln	Leu	Thr	Thr	Leu		
		515					520					525					
Ala	Glu	Lys	Leu	Leu	Gly	Pro	Gly	Val	Asn	Tyr	Ser	Gly	Cys	Gln	Ile		
	530					535					540						
Thr	Trp	Ala	Lys	Phe	Cys	Lys	Glu	Asn	Met	Ala	Gly	Lys	Gly	Phe	Ser		

545		550		555		560
Phe Trp Val Trp	Leu Asp Asn Ile Ile	Asp Leu Val Lys Lys	Tyr Ile			
	565	570	575			
Leu Ala Leu Trp	Asn Glu Gly Tyr Ile	Met Gly Phe Ile	Ser Lys Glu			
	580	585	590			
Arg Glu Arg Ala	Ile Leu Ser Thr	Lys Pro Pro Gly	Thr Phe Leu Leu			
	595	600	605			
Arg Phe Ser Glu	Ser Ser Lys Glu	Gly Gly Val Thr	Phe Thr Trp Val			
	610	615	620			
Glu Lys Asp Ile	Ser Gly Lys Thr	Gln Ile Gln Ser	Val Glu Pro Tyr			
	625	630	635			640
Thr Lys Gln Gln	Leu Asn Asn Met	Ser Phe Ala Glu	Ile Ile Met Gly			
	645	650	655			
Tyr Lys Ile Met	Asp Ala Thr Asn	Ile Leu Val Ser	Pro Leu Val Tyr			
	660	665	670			
Leu Tyr Pro Asp	Ile Pro Lys Glu	Glu Ala Phe Gly	Lys Tyr Cys Arg			
	675	680	685			
Pro Glu Ser Gln	Glu His Pro Glu	Ala Asp Pro Gly	Ser Ala Ala Pro			
	690	695	700			
Tyr Leu Lys Thr	Lys Phe Ile Cys	Val Thr Pro Thr	Thr Cys Ser Asn			
	705	710	715			720
Thr Ile Asp Leu	Pro Met Ser Pro	Arg Thr Leu Asp	Ser Leu Met Gln			
	725	730	735			
Phe Gly Asn Asn	Gly Glu Gly Ala	Glu Pro Ser Ala	Gly Gly Gln Phe			
	740	745	750			
Glu Ser Leu Thr	Phe Asp Met Glu	Leu Thr Ser Glu	Cys Ala Thr Ser			
	755	760	765			
Pro Met						
770						