




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/47, 16/18, G01N 33/50, 33/53, C12Q 1/68		A1	(11) International Publication Number: WO 98/46755 (43) International Publication Date: 22 October 1998 (22.10.98)
(21) International Application Number: PCT/US98/07894 (22) International Filing Date: 16 April 1998 (16.04.98)		(72) Inventor; and (75) Inventor/Applicant (for US only): MCCARTHY, Sean, A. [GB/US]; 62 Commonwealth Avenue #4, Boston, MA 02116 (US).	
(30) Priority Data: 08/843,704 16 April 1997 (16.04.97) US 08/842,898 17 April 1997 (17.04.97) US 60/071,589 15 January 1998 (15.01.98) US 09/009,802 20 January 1998 (20.01.98) US		(74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US) et al.	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 08/843,704 (CIP) Filed on 16 April 1997 (16.04.97) US 08/842,898 (CIP) Filed on 17 April 1997 (17.04.97) US 60/071,589 (CIP) Filed on 15 January 1998 (15.01.98) US 09/009,802 (CIP) Filed on 20 January 1998 (20.01.98)		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
<p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>			
(71) Applicant (for all designated States except US): MILLENIUM BIOTHERAPEUTICS, INC. [US/US]; 620 Memorial Drive, Cambridge, MA 02319 (US).			

(54) Title: CRSP PROTEIN (CYSTEINE-RICH SECRETED PROTEINS), NUCLEIC ACID MOLECULES ENCODING THEM AND USES THEREFOR

The Human CRSP Family


h-CRSP-1 (T59)



h-CRSP-2




h-CRSP-3



h-CRSP-4



h-CRSP-n



(57) Abstract

CRSP (Cysteine-Rich Secreted Proteins) polypeptides, proteins, and nucleic acid molecules are disclosed. In addition to isolated, full-length CRSP proteins, the invention further provides isolated CRSP fusion proteins, antigenic peptides and anti-CRSP antibodies. The invention also provides CRSP nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a CRSP gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CRSP PROTEIN (CYSTEINE-RICH SECRETED PROTEINS), NUCLEIC ACID MOLECULES ENCODING THEM AND USES THEREFOR

Background of the Invention

5 Secreted proteins play an integral role in the formation, differentiation, and maintenance of cells in multicellular organisms. For instance, secretory proteins are known in the art to be involved in signalling between cells which are not in direct contact. Such secreted signaling molecules are particularly important in the development of vertebrate tissue during embryogenesis as well as in the maintenance of 10 the differentiated state of adult tissues. For example, inductive interactions that occur between neighboring cell layers and tissues in the developing embryo are largely dependent on the existence and regulation of secreted signaling molecules. In inductive interactions, biochemical signals secreted by one cell population influence the developmental fate of a second cell population, typically by altering the fate of the 15 second cell population. For example, the Wnt proteins are now recognized as one of the major families of developmentally important signaling molecules in organisms ranging from *Drosophila* to mice. For review see Cadigan, K.M. and R. Nusse (1997) *Genes & Development*, 11:3286-3305.

20 It is also now well recognized that major families of signaling molecules have a dual role to play in both the development of an organism as well as in promoting or maintaining the differentiated state of tissues in the adult animal. Furthermore, major families of signaling molecules have been implicated in controlling proliferation of cells in mature adult tissue, for example, during normal cell turnover in the adult organism as well as in tissue regeneration activated as a result of damage to the adult tissue.

25

Summary of the Invention

The present invention is based, at least in part, on the discovery of nucleic acid molecules which encode a novel family of secreted proteins, referred to herein as the Cysteine-Rich Secreted Proteins ("CRSPs" or "CRSP proteins"). The CRSP molecules 30 of the present invention are useful as modulating agents in regulating a variety of cellular processes. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding CRSP proteins or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of CRSP-encoding nucleic acids.

35

In one embodiment, a CRSP nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, or complement thereof. In yet another embodiment, a CRSP nucleic acid molecule is 80% homologous to the

- 2 -

nucleotide sequence shown in SEQ ID NO:4, SEQ ID NO:6, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, or a complement thereof. In yet another embodiment, a CRSP nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, the
5 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof. In yet another embodiment, a CRSP nucleic acid molecule is 85% homologous to the nucleotide sequence shown in SEQ ID NO:7, SEQ ID NO:9, the nucleotide sequence of
10 the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof. In yet another embodiment, a CRSP nucleic acid molecule is 70% homologous to the nucleotide sequence shown in SEQ ID NO:10, SEQ ID NO:12, or a complement thereof. In a preferred embodiment,
15 an isolated CRSP nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:3, or a complement thereof. In another embodiment, a CRSP nucleic acid molecule further comprises nucleotides 1-37 of SEQ ID NO:1. In yet another preferred embodiment, a CRSP nucleic acid molecule further comprises nucleotides 1088-2479 of SEQ ID NO:1. In another preferred embodiment, an isolated CRSP nucleic acid
20 molecule has the nucleotide sequence shown in SEQ ID NO:1.

In another preferred embodiment, an isolated CRSP nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:6, or a complement thereof. In another embodiment, a CRSP nucleic acid molecule further comprises nucleotides 1-125 of SEQ ID NO:4. In yet another preferred embodiment, a CRSP nucleic acid molecule further
25 comprises nucleotides 797-848 of SEQ ID NO:4. In another preferred embodiment, an isolated CRSP nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:4.

In another preferred embodiment, an isolated CRSP nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:9, the nucleotide sequence of the DNA
30 insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof. In another embodiment, a CRSP nucleic acid molecule further comprises nucleotides 1-92 of SEQ ID NO:7. In yet another preferred embodiment, a CRSP nucleic acid molecule further comprises nucleotides 891-1529 of
35 SEQ ID NO:7. In another preferred embodiment, an isolated CRSP nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:7.

In another preferred embodiment, an isolated CRSP nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:12, or a complement thereof. In another embodiment, a CRSP nucleic acid molecule further comprises nucleotides 538-702 of SEQ ID NO:10. In yet another preferred embodiment, an isolated CRSP nucleic acid 5 molecule has the nucleotide sequence shown in SEQ ID NO:10.

In another embodiment, a CRSP nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID 5, SEQ ID NO:8, or SEQ ID NO:11. In another preferred embodiment, a CRSP nucleic acid molecule includes a nucleotide 10 sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:2. In yet another preferred embodiment, a CRSP nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:5. In yet another preferred embodiment, a CRSP nucleic acid molecule includes a 15 nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a CRSP nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 75% homologous to the amino acid sequence of SEQ ID NO:8. In yet another preferred embodiment, a CRSP nucleic acid 20 molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 65% homologous to the amino acid sequence of SEQ ID NO:11.

In another embodiment, an isolated nucleic acid molecule of the present invention encodes a CRSP protein which includes a signal sequence and at least one cysteine-rich domain, and is secreted. In another embodiment, an isolated nucleic acid 25 molecule of the present invention encodes a CRSP protein which includes a signal sequence and at least one cysteine-rich domain, preferably a cysteine-rich region, and is secreted. In yet another embodiment, a CRSP nucleic acid molecule encodes a CRSP protein and is a naturally occurring nucleotide sequence.

Another embodiment of the invention features CRSP nucleic acid molecules 30 which specifically detect CRSP nucleic acid molecules relative to nucleic acid molecules encoding non-CRSP proteins. For example, in one embodiment, a CRSP nucleic acid molecule hybridizes under stringent conditions to a nucleic acid molecule comprising the nucleotides 470-2479 of nucleotide sequence shown in SEQ ID NO:1, to nucleotides 1-475 of nucleotide sequence shown in SEQ ID NO:4, or to nucleotides 1-600 of 35 nucleotide sequence shown in SEQ ID NO:7, or hybridizes under stringent conditions to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, to the nucleotide sequence of the DNA insert of the plasmid

deposited with ATCC as Accession Number 98633, or to the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452. In another embodiment, the CRSP nucleic acid molecule is at least 500 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising 5 the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10 or a complement thereof.

Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of a CRSP nucleic acid.

Another aspect of the invention provides a vector comprising a CRSP nucleic 10 acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. The invention also provides a method for producing a CRSP protein by culturing in a suitable medium, a host cell of the invention containing a recombinant expression vector such that a CRSP protein is produced.

15 Another aspect of this invention features isolated or recombinant CRSP proteins and polypeptides. In one embodiment, an isolated CRSP protein has a signal sequence and at least one cysteine-rich domain, preferably a cysteine-rich region, and is secreted. In another embodiment, an isolated CRSP protein has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, 20 SEQ ID NO:8, or SEQ ID NO:11. In a preferred embodiment, a CRSP protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2. In another preferred embodiment, a CRSP protein has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:5. In another preferred embodiment, a CRSP protein has an amino acid sequence at least about 60% 25 homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a CRSP protein has an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:8. In another preferred embodiment, a CRSP protein has an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:11. In another embodiment, a 30 CRSP protein has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11.

Another embodiment of the invention features an isolated CRSP protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:1, or a complement thereof. 35 Another embodiment of the invention features an isolated CRSP protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 80% homologous to a nucleotide sequence of SEQ ID NO:4, or a complement thereof.

Another embodiment of the invention features an isolated CRSP protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof.

Another embodiment of the invention features an isolated CRSP protein which is 5 encoded by a nucleic acid molecule having a nucleotide sequence at least about 85% homologous to a nucleotide sequence of SEQ ID NO:7, or a complement thereof.

Another embodiment of the invention features an isolated CRSP protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 70% homologous to a nucleotide sequence of SEQ ID NO:10, or a complement thereof. This 10 invention further features an isolated CRSP protein which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, or a complement thereof.

The CRSP proteins of the present invention, or biologically active portions 15 thereof, can be operatively linked to a non-CRSP polypeptide to form CRSP fusion proteins. The invention further features antibodies that specifically bind CRSP proteins, such as monoclonal or polyclonal antibodies. In addition, the CRSP proteins or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers.

20 In another aspect, the present invention provides a method for detecting CRSP expression in a biological sample by contacting the biological sample with an agent capable of detecting a CRSP nucleic acid molecule, protein or polypeptide such that the presence of a CRSP nucleic acid molecule, protein or polypeptide is detected in the biological sample.

25 In another aspect, the present invention provides a method for detecting the presence of CRSP activity in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of CRSP activity such that the presence of CRSP activity is detected in the biological sample.

30 In another aspect, the invention provides a method for modulating CRSP activity comprising contacting the cell with an agent that modulates CRSP activity such that CRSP activity in the cell is modulated. In one embodiment, the agent inhibits CRSP activity. In another embodiment, the agent stimulates CRSP activity. In one embodiment, the agent is an antibody that specifically binds to a CRSP protein. In another embodiment, the agent modulates expression of CRSP by modulating transcription of a CRSP gene or translation of a CRSP mRNA. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of a CRSP mRNA or a CRSP gene.

In one embodiment, the methods of the present invention are used to treat a subject having a disorder characterized by aberrant CRSP protein or nucleic acid expression or activity by administering an agent which is a CRSP modulator to the subject. In one embodiment, the CRSP modulator is a CRSP protein. In another 5 embodiment the CRSP modulator is a CRSP nucleic acid molecule. In yet another embodiment, the CRSP modulator is a peptide, peptidomimetic, or other small molecule. In a preferred embodiment, the disorder characterized by aberrant CRSP protein or nucleic acid expression is a developmental, differentiative, or proliferative disorder.

The present invention also provides a diagnostic assay for identifying the 10 presence or absence of a genetic alteration characterized by at least one of (i) aberrant modification or mutation of a gene encoding a CRSP protein; (ii) mis-regulation of said gene; and (iii) aberrant post-translational modification of a CRSP protein, wherein a wild-type form of said gene encodes a protein with a CRSP activity.

In another aspect the invention provides a method for identifying a compound 15 that binds to or modulates the activity of a CRSP protein, by providing a indicator composition comprising a CRSP protein having CRSP activity, contacting the indicator composition with a test compound, and determining the effect of the test compound on CRSP activity in the indicator composition to identify a compound that modulates the activity of a CRSP protein.

20 Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 depicts the cDNA sequence and predicted amino acid sequence of 25 human CRSP-1. The nucleotide sequence corresponds to nucleic acids 1 to 2479 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 350 of SEQ ID NO:2.

Figure 2 depicts the cDNA sequence and predicted amino acid sequence of 30 human CRSP-2. The nucleotide sequence corresponds to nucleic acids 1 to 848 of SEQ ID NO:4. The amino acid sequence corresponds to amino acids 1 to 224 of SEQ ID NO:5.

Figure 3 depicts the cDNA sequence and predicted amino acid sequence of 35 human CRSP-3. The nucleotide sequence corresponds to nucleic acids 1 to 1529 of SEQ ID NO:7. The amino acid sequence corresponds to amino acids 1 to 266 of SEQ ID NO:8.

Figure 4 depicts the cDNA sequence and predicted amino acid sequence of human CRSP-4. The nucleotide sequence corresponds to nucleic acids 1 to 702 of SEQ

ID NO:10. The amino acid sequence corresponds to amino acids 1 to 179 of SEQ ID NO:11.

5 *Figure 5* depicts the cDNA sequence and predicted amino acid sequence of human CRSP-like-N. The nucleotide sequence corresponds to nucleic acids 1 to 928 of SEQ ID NO:13. The amino acid sequence corresponds to amino acids 1 to 242 of SEQ ID NO:14.

10 *Figure 6* is depicts the amino acid sequences of human CRSP-1, human CRSP-2, human CRSP-3, and human CRSP-4, as well as the consensus sequence generated from the alignment of these sequences. The figure details the relationship between the CRSP proteins of the instant invention.

15 *Figure 7* is a schematic diagram depicting the the relationship between the CRSP proteins of the instant invention. The figure depicts the biological and functional domains of human CRSP. Signal peptides are indicated by filled boxes. The cysteine-rich domains of a CRSP cysteine-rich region are depicted as CRD-1 and CRD-2.

15 *Figure 8* depicts the cDNA sequence and predicted amino acid sequence of murine CRSP-1. The nucleotide sequence corresponds to nucleic acids 1 to 928 of SEQ ID NO:16. The amino acid sequence corresponds to amino acids 1 to 349 of SEQ ID NO:17.

20 *Figure 9* is a schematic diagram depicting the relationship between the CRSP-1 nucleotide sequence and those of RIG and RIG-like 7-1 (Accession Nos. U32331 and AF034208, respectively).

Detailed Description of the Invention

25 The present invention is based on the discovery of novel molecules, referred to herein as CRSP protein and nucleic acid molecules, which comprise a family of molecules having certain conserved structural and functional features. The term "family" when referring to the protein and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having a common structural domain and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members of a family may also have common functional characteristics.

35 In one embodiment, a CRSP family member is identified based on the presence of at least one "cysteine-rich domain" in the protein or corresponding nucleic acid

molecule. As defined herein, a "cysteine-rich domain" refers to a portion of a CRSP protein (e.g., CRSP-1) which is rich in cysteine residues, i.e., at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues. Preferably, a "cysteine-rich domain" is a protein domain having an amino acid sequence of about 30-

5 100 amino acids, preferably about 35-95 amino acids, more preferably about 40-90 amino acids, more preferably about 45-85 amino acids, even more preferably about 50-80 amino acids, and even more preferably about 55-75, 60-70, or 65 amino acids, of which at least about 3-20, preferably about 5-15, or more preferably about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids are cysteine residues.

10 A preferred cysteine-rich domain has at least about 10, at least about 15, 16, 17, 18, 19, or preferably 20 cysteine residues located at the same relative amino acid position as the cysteine residues in human CRSP-1 having SEQ ID NO:2. For example, as shown in Figure 6, CRSP-2 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human CRSP-1 having SEQ ID NO:2 (e.g., cys151 in CRSP-2, SEQ ID NO:5, is located at the same relative amino acid position as cys214 in CRSP-1, SEQ ID NO:2; cys156 in CRSP-2, SEQ ID NO:5, is located at the same relative amino acid position as cys219 in CRSP-1, SEQ ID NO:2; and cys157 in CRSP-2, SEQ ID NO:5, is located at the same relative amino acid position as cys220 in CRSP-1, SEQ ID NO:2). Similarly, as shown in Figure 6, CRSP-3 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human CRSP-1 having SEQ ID NO:2. As also shown in Figure 6, CRSP-4 has at least about 10 cysteine residues located at the same relative amino acid position as the cysteine residues in human CRSP-1 having SEQ ID NO:2.

15 A preferred CRSP protein of the present invention is a human protein (e.g., encoded by a nucleotide sequence corresponding to a naturally-occurring human gene.) For example, in one embodiment, a CRSP-1 protein (human CRSP-1) contains a first cysteine-rich domain containing about amino acids 147-195 of SEQ ID NO:2, having 10 cysteine residues, and a second cysteine-rich domain containing about amino acids 201-284 of SEQ ID NO:2, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a CRSP-2 (human CRSP-2) protein contains a first cysteine-rich domain containing about amino acids 41-90 of SEQ ID NO:5, having 10 cysteine residues, and a second cysteine-rich domain containing about amino acids 41-218 of SEQ ID NO:5, having 10 cysteine residues (the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a CRSP-3 protein (human CRSP-3) contains a first cysteine-rich domain containing about amino acids 85-138 of SEQ ID NO:8, having 10 cysteine residues, and a second cysteine-rich domain containing about amino acids 182-263 of SEQ ID NO:8, having 10 cysteine residues

(the positions of the cysteine residues are depicted in Figure 6). In another embodiment, a CRSP-4 protein (human CRSP-4) contains a first cysteine-rich domain containing about amino acids 1-47 of SEQ ID NO:11, having 8 cysteine residues, and a second cysteine-rich domain containing about amino acids 96-176 of SEQ ID NO:11, having 10 5 cysteine residues (the positions of the cysteine residues are depicted in Figure 6).

Preferred CRSP proteins have more than one cysteine-rich domains, more preferably have at least two cysteine-rich domains and, thus, have a cysteine-rich region. As used herein, the term "cysteine-rich region" refers to a protein domain which includes at least two cysteine-rich domains and has an amino acid sequence of about 120-200, 10 preferably about 130-190, more preferably about 140-180 amino acid residues, and even more preferably at least about 135-175 amino acids of which at least about 10-30, preferably about 15-20, and more preferably about 16, 17, 18, or 19 of the amino acids are cysteine residues. In a preferred embodiment, a cysteine-rich region is located in the C-terminal region of a CRSP protein. For example, in one embodiment, a CRSP-1 15 protein contains a cysteine rich region containing about amino acids 147-284 of SEQ ID NO:2, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a CRSP-2 protein contains a cysteine rich region containing about amino acids 41-218 of SEQ ID NO:5, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a CRSP-3 protein contains a cysteine rich region 20 containing about amino acids 85-263 of SEQ ID NO:8, having 20 cysteine residues at the positions indicated in Figure 6. In another embodiment, a CRSP-4 protein contains a cysteine rich region containing about amino acids 1-176 of SEQ ID NO:2, having 18 cysteine residues at the positions indicated in Figure 6.

In another embodiment, in addition to cysteine-rich domains, the cysteine-rich 25 region contains a spacer region which separates the first and second cysteine-rich domains. As used herein, the "spacer region" refers to amino acid residues which are located between the first and second cysteine-rich domains of a cysteine-rich region and includes amino acid residues located C-terminal to the first cysteine-rich domain and N-terminal to the second cysteine-rich domain. As defined herein, a "spacer region" refers 30 to a protein domain of about 5-70 amino acids, preferably about 10-65 amino acids, more preferably about 15-60 amino acids, even more preferably about 20-55 amino acids, and even more preferably about 25-50, 30-45 or 35-40 amino acids. For example, CRSP-1 protein contains a spacer region of about amino acids 196-200 of SEQ ID NO:2; CRSP-2 protein contains a spacer region of about amino acids 91-137 of SEQ ID NO:5; CRSP-3 protein contains a spacer region of about amino acids 139-181 of SEQ ID NO:8; and CRSP-4 protein contains a spacer region of about amino acids 48-95 of SEQ ID NO:11.

In another embodiment of the invention, the CRSP protein has at least one cysteine-rich domain, preferably a cysteine-rich region, and a signal sequence. As used herein, a "signal sequence" refers to a peptide containing about 20 amino acids which occurs at the N-terminus of secretory and integral membrane proteins and which

5 contains a large number of hydrophobic amino acid residues. For example, a signal sequence contains at least about 14-28 amino acid residues, preferably about 16-26 amino acid residues, more preferably about 18-24 amino acid residues, and more preferably about 20-22 amino acid residues, and has at least about 40-70%, preferably about 50-65%, and more preferably about 55-60% hydrophobic amino acid residues

10 (e.g., Alanine, Valine, Leucine, Isoleucine, Phenylalanine, Tyrosine, Tryptophan, or Proline). Such a "signal sequence", also referred to in the art as a "signal peptide", serves to direct a protein containing such a sequence to a lipid bilayer. For example, in one embodiment, a CRSP-1 protein contains a signal sequence of about amino acids 1-23 of SEQ ID NO:2. In another embodiment, a CRSP-2 protein contains a signal

15 sequence of about amino acids 1-19 of SEQ ID NO:5. In another embodiment, a CRSP-3 protein contains a signal sequence of about amino acids 1-20 of SEQ ID NO:8.

CRSP proteins of the present invention can be used to identify additional CRSP family members. For example, a protein having homology to CRSP-1 was identified using the nucleotide sequence encoding the N-terminal unique region of CRSP-1 to

20 search a protein sequence database. Such a protein, referred to herein as "CRSP-like-N" is depicted in Figure 5. The nucleotide sequence of CRSP-like-N (SEQ ID NO:13) encodes a protein having 242 amino acids (SEQ ID NO:14). The nucleotide sequence of CRSP-like-N includes a 5' untranslated region containing nucleotides 1-74 of SEQ ID NO:13, a coding region containing nucleotides 75-800 of SEQ ID NO:13 (corresponding to nucleotides 1-726 of SEQ ID NO:15), and a 3' untranslated region containing nucleotides 801-928 of SEQ ID NO:13.

Accordingly, one embodiment of the invention features a CRSP protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region. Another embodiment features a CRSP protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least one cysteine-rich domains. Another embodiment features a CRSP protein having at least one cysteine-rich region, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment features a protein having 20, 30, 40, 50, 60, 70, 80, 90, 95, or 99% homology to a cysteine-rich domain of a CRSP protein of the invention (e.g., CRSP-1, CRSP-2, CRSP-3, or CRSP-4).

Yet another embodiment of the invention features a CRSP protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal

peptide. Another embodiment features a CRSP protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains. Another embodiment features a CRSP protein having at least one cysteine-rich domain, preferably at least one cysteine-rich region and a signal peptide, wherein the cysteine-rich region includes at least two cysteine-rich domains and a spacer.

Preferred CRSP molecules of the present invention have an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11. As used herein, the term "sufficiently homologous" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (e.g., an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains and/or a common functional activity. For example, amino acid or nucleotide sequences which share common structural domains have at least about 40% homology, preferably 50% homology, more preferably 60%-70% homology across the amino acid sequences of the domains and contain at least one, preferably two, more preferably three, and even more preferably four, five or six structural domains, are defined herein as sufficiently homologous. Furthermore, amino acid or nucleotide sequences which share at least 40%, preferably 50%, more preferably 60, 70, or 80% homology and share a common functional activity are defined herein as sufficiently homologous.

As used interchangeably herein, a "CRSP activity", "biological activity of CRSP" or "functional activity of CRSP", refers to an activity exerted by a CRSP protein, polypeptide or nucleic acid molecule on a CRSP responsive cell as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, a CRSP activity is a direct activity, such as an association with a CRSP-target molecule. As used herein, a "target molecule" is a molecule with which a CRSP protein binds or interacts in nature, such that CRSP-mediated function is achieved. A CRSP target molecule can be a non-CRSP molecule or a CRSP protein or polypeptide of the present invention. In an exemplary embodiment, a CRSP target molecule is a membrane-bound protein (e.g., a cell-surface receptor or "CRSP receptor") or a modified form of such a protein which has been altered such that the protein is soluble (e.g., recombinantly produced such that the protein does not express a membrane-binding domain). In another embodiment, a CRSP target is a second soluble protein molecule (e.g., a "CRSP binding partner" or "CRSP substrate"). In such an exemplary embodiment, a CRSP binding partner can be a second soluble non-CRSP protein or a second CRSP protein molecule of the present

invention. Alternatively, a CRSP activity is an indirect activity, such as a cellular signaling activity mediated by interaction of the CRSP protein with a second protein (e.g., a CRSP receptor). As used herein, the term "CRSP receptor" refers to a protein or protein complex, to which a CRSP protein, e.g., human CRSP, can bind. A receptor can 5 be a cell surface receptor, e.g., a nuclear hormone receptor. CRSP receptors can be isolated by methods known in the art and further described herein. Interaction of a CRSP protein with a CRSP receptor can result in transduction of a signal from the cell surface to the nucleus. The signal transduced can be, an increase in intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, e.g., in 10 phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

In a preferred embodiment, a CRSP activity is at least one or more of the following activities: (i) interaction of a CRSP protein with and/or binding to a second molecule, (e.g., a protein, such as a CRSP receptor, a soluble form of a CRSP receptor, a 15 receptor for a member of the *wnt* family of signaling proteins, or a non-CRSP signaling molecule); (ii) interaction of a CRSP protein with an intracellular protein via a membrane-bound CRSP receptor; (iii) complex formation between a soluble CRSP protein and a second soluble CRSP binding partner (e.g., a non-CRSP protein molecule or a second CRSP protein molecule); (iv) interaction with other extracellular proteins 20 (e.g., regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (e.g., a detoxifying activity or defense function); and/or (vi) an enzymatic activity. In yet another preferred embodiment, a CRSP activity is at least one or more of the following activities: (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (e.g., antagonism of 25 the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction); (2) regulation of communication between cells (e.g., regulation of *wnt*-dependent cell-cell interactions); (3) regulation of expression of genes whose expression is modulated by binding of CRSP (e.g., CRSP-1) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, 30 either *in vitro* or *in vivo* (e.g., induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) regulation of gene transcription in a cell involved in development or differentaition, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in 35 development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of cellular proliferation, either *in vitro* or *in vivo* (e.g., induction of cellular proliferation or inhibition of proliferation as in the case of suppression of

tumorigenesis (e.g., suppression of glioblastoma formation)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (e.g., induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as 5 stimulation of cell survival; (11) regulating cell migration; and/or (12) immune modulation.

As referred to herein, "differentiation-specific proteins" include proteins involved in the transition of a cell from the undifferentiated to the differentiated phenotype. For example, such proteins can be differentiation specific structural proteins 10 or differentiation-specific transcription factors. Such differentiation-specific proteins are generally expressed at higher levels in cells which are making the transition from the undifferentiated to the differentiated phenotype (e.g., during embryonic development or during regeneration of mature tissue in the adult animal), or are expressed at higher levels in fully-differentiated or terminally-differentiated cells as compared to their 15 undifferentiated counterparts. Also, as referred to herein, "differentiation-specific genes" include nucleic acid molecules which encode differentiation-specific proteins.

Accordingly, another embodiment of the invention features isolated CRSP proteins and polypeptides having a CRSP activity. Preferred CRSP proteins have at least one cysteine-rich region and a CRSP activity. In another preferred embodiment, 20 the CRSP protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least one cysteine-rich domain, and a CRSP activity. In another preferred embodiment, the CRSP protein has at least one cysteine-rich region, wherein the cysteine-rich region comprises at least two cysteine-rich domains, and a CRSP activity. In yet another preferred embodiment, a CRSP protein further comprises a signal 25 sequence. In still another preferred embodiment, a CRSP protein has a cysteine-rich region, a CRSP activity, and an amino acid sequence sufficiently homologous to an amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11.

The human CRSP-1 cDNA, which is approximately 2479 nucleotides in length, 30 encodes a protein which is approximately 350 amino acid residues in length. The human CRSP protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A CRSP cysteine-rich region can be found at least, for example, from about amino acids 147-284 of SEQ ID NO:2. The CRSP-1 cysteine-rich region comprises a first cysteine-rich domain from about amino acids 147- 35 195 of SEQ ID NO:2 and a second cysteine-rich domain from about amino acids 201-284 of SEQ ID NO:2. The human CRSP-1 protein is a secreted protein which further contains a signal sequence at about amino acids 1-23 of SEQ ID NO:2. The prediction

of such a signal peptide can be made, for example, utilizing the computer algorithm SIGNALP (Henrik, et al. (1997) *Protein Engineering* 10:1-6).

The human CRSP-2 cDNA, which is approximately 848 nucleotides in length, encodes a protein which is approximately 224 amino acid residues in length. The 5 human CRSP protein contains an N-terminal signal sequence and a cysteine-rich region comprising two cysteine-rich domains. A CRSP cysteine-rich region can be found at least, for example, from about amino acids 41-218 of SEQ ID NO:5. The CRSP-2 cysteine-rich region comprises a first cysteine-rich domain from about amino acids 41-90 of SEQ ID NO:5 and a second cysteine-rich domain from about amino acids 138-218 10 of SEQ ID NO:5. The human CRSP-2 protein is a secreted protein which further contains a signal sequence at about amino acids 1-19 of SEQ ID NO:5.

The human CRSP-3 cDNA, which is approximately 1529 nucleotides in length, encodes a protein which is approximately 266 amino acid residues in length. The human CRSP protein contains an N-terminal signal sequence and a cysteine-rich region 15 comprising two cysteine-rich domains. A CRSP cysteine-rich region can be found at least, for example, from about amino acids 85-263 of SEQ ID NO:8. The CRSP-3 cysteine-rich region comprises a first cysteine-rich domain from about amino acids 85-138 of SEQ ID NO:8 and a second cysteine-rich domain from about amino acids 182-263 of SEQ ID NO:2. The human CRSP-3 protein is a secreted protein which further 20 contains a signal sequence at about amino acids 1-20 of SEQ ID NO:8.

The human CRSP-4 cDNA, which is approximately 702 nucleotides in length, encodes a protein which is approximately 179 amino acid residues in length. The human CRSP protein contains a cysteine-rich region comprising two cysteine-rich domains. A CRSP cysteine-rich region can be found at least, for example, from about 25 amino acids 1-176 of SEQ ID NO:11. The CRSP-4 cysteine-rich region comprises a first cysteine-rich domain from about amino acids 1-47 of SEQ ID NO:11 and a second cysteine-rich domain from about amino acids 96-176 of SEQ ID NO:11.

Various aspects of the invention are described in further detail in the following 30 subsections:

I. Isolated Nucleic Acid Molecules

One aspect of the invention pertains to isolated nucleic acid molecules that encode CRSP proteins or biologically active portions thereof, as well as nucleic acid 35 fragments sufficient for use as hybridization probes to identify CRSP-encoding nucleic acids (e.g., CRSP mRNA) and fragments for use as PCR primers for the amplification or mutation of CRSP nucleic acid molecules. As used herein, the term "nucleic acid

molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA.

5 An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, 10 the isolated CRSP nucleic acid molecule can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, 15 or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with 20 ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or portion of the nucleic acid sequence 25 of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 as a hybridization probe, 30 CRSP nucleic acid molecules can be isolated using standard hybridization and cloning techniques (e.g., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID 35 NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the 5 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452.

A nucleic acid of the invention can be amplified using cDNA, mRNA or 10 alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to CRSP nucleotide sequences can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

15 In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:1. The sequence of SEQ ID NO:1 corresponds to the human CRSP-1 cDNA. This cDNA comprises sequences encoding the human CRSP-1 protein (i.e., "the coding region", from nucleotides 38-1087), as well as 5' untranslated sequences (nucleotides 1 to 37) and 3' untranslated 20 sequences (nucleotides 1088 to 2479). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:1 (e.g., nucleotides 38 to 1087, corresponding to SEQ ID NO:3). A plasmid containing the full-length nucleotide sequence encoding CRSP-1 was deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on June 11, 1998 and assigned Accession Number 25 98452. A plasmid containing the full-length nucleotide sequence encoding CRSP-1 was deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on January 16, 1998 and assigned Accession Number 98634.

In another preferred embodiment, an isolated nucleic acid molecule of the 30 invention comprises the nucleotide sequence shown in SEQ ID NO:4. The sequence of SEQ ID NO:4 corresponds to the human CRSP-2 cDNA. This cDNA comprises sequences encoding the human CRSP-2 protein (i.e., "the coding region", from nucleotides 126-796), as well as 5' untranslated sequences (nucleotides 1 to 125) and 3' untranslated sequences (nucleotides 797 to 848). Alternatively, the nucleic acid 35 molecule can comprise only the coding region of SEQ ID NO:4 (e.g., nucleotides 126 to 796, corresponding to SEQ ID NO:6).

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:7. The sequence of

SEQ ID NO:7 corresponds to the human CRSP-3 cDNA. This cDNA comprises sequences encoding the human CRSP-3 protein (i.e., "the coding region", from nucleotides 93-890), as well as 5' untranslated sequences (nucleotides 1 to 92) and 3' untranslated sequences (nucleotides 891 to 1529). Alternatively, the nucleic acid
5 molecule can comprise only the coding region of SEQ ID NO:7 (e.g., nucleotides 93 to 890, corresponding to SEQ ID NO:9). A plasmid containing the full-length nucleotide sequence encoding CRSP-3 was deposited with the American Type Culture Collection (ATCC), Rockville, Maryland, on January 16, 1998 and assigned Accession Number 98633.

10 In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:10. The sequence of SEQ ID NO:10 corresponds to the human CRSP-4 cDNA. This cDNA comprises sequences encoding the human CRSP-4 protein (i.e., "the coding region", from nucleotides 1-537), as well as 3' untranslated sequences (nucleotides 538 to 702).

15 Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:10 (e.g., nucleotides 1 to 537, corresponding to SEQ ID NO:12).

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, the
20 nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a portion of any of these nucleotide sequences. A nucleic acid molecule which is complementary to
25 the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, is
30 one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, such that it can hybridize to the
35 nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, or the nucleotide sequence of the DNA insert of the plasmid deposited with

ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, thereby forming a stable duplex.

5 In still another preferred embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about 60-65%, and even more preferably at least about 70-75%, 80-85%, 90-95% or more homologous to the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:4, SEQ
10 ID NO:7, or SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a portion of any of these nucleotide sequences.

15 Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the
20 DNA insert of the plasmid deposited with ATCC as Accession Number 98452, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a CRSP protein. The nucleotide sequence determined from the cloning of the human CRSP genes allows for the generation of probes and primers designed for use in identifying and/or cloning CRSP homologues in other cell
25 types, e.g., from other tissues, as well as CRSP homologues from other mammals. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense sequence of SEQ ID NO:1, SEQ
30 ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, of an anti-sense sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or

the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or of a naturally occurring mutant of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of 5 the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising nucleotides 470-2479 of 10 SEQ ID NO:1 or to a nucleic acid molecule comprising nucleotides 1-475 of SEQ ID NO:4.

Probes based on the human CRSP nucleotide sequence can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. For instance, primers based on the nucleic acid represented in SEQ ID NOS:1 or 3 can be 15 used in PCR reactions to clone CRSP homologs (e.g., CRSP-1 homologues). In a preferred embodiment of the invention, CRSP homologs are cloned by PCR amplification (e.g., RT-PCR) using primers hybridizing to a portion of the nucleotide sequence encoding the CRSP cysteine rich domain. Likewise, probes based on the subject CRSP sequences can be used to detect transcripts or genomic sequences 20 encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, e.g., the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a CRSP protein, such as by measuring a level of a CRSP-encoding nucleic acid in a sample of 25 cells from a subject e.g., detecting CRSP mRNA levels or determining whether a genomic CRSP gene has been mutated or deleted.

A nucleic acid fragment encoding a "biologically active portion of a CRSP protein" can be prepared by isolating a portion of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid 30 deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, which encodes a polypeptide having a CRSP biological activity (the biological activities of the CRSP proteins have previously been described), 35 expressing the encoded portion of the CRSP protein (e.g., by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the CRSP protein.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid 5 deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, due to degeneracy of the genetic code and thus encode the same CRSP proteins as those encoded by the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid 10 deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid 15 sequence shown in SEQ ID NO:2, SEQ ID NO: 5, SEQ ID NO:8, or SEQ ID NO:11.

In addition to the human CRSP nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 20 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the CRSP proteins may exist within a population (e.g., the human population). Such genetic polymorphism in the CRSP genes may exist among individuals within a population due 25 to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a CRSP protein, preferably a mammalia CRSP protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a CRSP gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in CRSP genes 30 that are the result of natural allelic variation and that do not alter the functional activity of a CRSP protein are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding CRSP proteins from other species, and thus which have a nucleotide sequence which differs from the human sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence 35 of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid

deposited with ATCC as Accession Number 98452, are intended to be within the scope of the invention. For example, a murine CRSP-1 cDNA has been identified based of the nucleotide sequence of human CRSP-1. The nucleotide sequence of murine CRSP-1 (SEQ ID NO:16) encodes a CRSP-1 protein having 349 amino acids. The nucleotide 5 and amino acid sequences of murine CRSP-1 are depicted in Figure 8. The coding region of murine CRSP-1 is represented by SEQ ID NO:18.

Nucleic acid molecules corresponding to natural allelic variants and homologues of the CRSP cDNAs of the invention can be isolated based on their homology to the human CRSP nucleic acids disclosed herein using the human cDNA, or a portion 10 thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Examples of tissues and/or libraries suitable for isolation of the subject nucleic acids include thymus, lymph nodes and inflammatory tissue. cDNA encoding a CRSP protein (e.g., a CRSP-1 protein) can be obtained by isolating total mRNA from a cell, e.g., a vertebrate cell, a mammalian cell, or a human 15 cell, including embryonic cells. Double stranded cDNAs can then be prepared from the total mRNA, and subsequently inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. The gene encoding a CRSP-1 protein can also be cloned using established polymerase chain reaction techniques in accordance 20 with the nucleotide sequence information provided by the invention. The nucleic acid of the invention can be DNA or RNA or analogs thereof.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the 25 plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452. In other embodiment, the nucleic acid is at least 30, 50, 100, 250 or 500 nucleotides in length. As used herein, the term "hybridizes under stringent 30 conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% homologous to each other typically remain hybridized to each other. Such 35 stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization

in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 corresponds to a naturally-occurring nucleic acid molecule. As used 5 herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural protein).

In addition to naturally-occurring allelic variants of the CRSP sequences that may exist in the population, the skilled artisan will further appreciate that changes can be 10 introduced by mutation into the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as 15 Accession Number 98452, thereby leading to changes in the amino acid sequence of the encoded CRSP proteins, without altering the functional ability of the CRSP proteins. For example, nucleotide substitutions leading to amino acid substitutions (particularly conservative amino acid substitutions) at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, 20 the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence 25 of CRSP (e.g., the sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the CRSP proteins of the present invention (e.g., cysteine residues within cysteine-rich domains), are predicted to be particularly unamenable to alteration. 30 Furthermore, amino acid residues that are conserved between CRSP protein and other proteins having cysteine-rich domains are not likely to be amenable to alteration.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding CRSP proteins that contain changes in amino acid residues that are not essential for activity. Such CRSP proteins differ in amino acid sequence from SEQ ID 35 NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about

60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11. Preferably, the protein encoded by the nucleic acid molecule is at least about 65-70% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, more preferably at least about 75-80% homologous to SEQ ID NO:2, 5 SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, even more preferably at least about 85-90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, and most preferably at least about 95% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11.

An isolated nucleic acid molecule encoding a CRSP protein homologous to the 10 protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid 15 deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid 20 deposited with ATCC as Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, by standard techniques, such as site-directed mutagenesis and 25 PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, 30 threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue 35 in a CRSP protein (e.g., one not located in a cysteine-rich domain) is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a CRSP

coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for CRSP biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

5 Accession Number 98634, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

10 In a preferred embodiment, a mutant CRSP protein can be assayed for intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, e.g., in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

15 In a preferred embodiment, a mutant CRSP protein can also be assayed for the ability to (1) modulate cellular signal transduction, either *in vitro* or *in vivo*; (2) regulate communication between cells; (3) regulate expression of genes whose expression is modulated by binding of CRSP (e.g., CRSP-1) to a receptor; (4) regulate gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo*; (5) regulate cellular proliferation, either *in vitro* or *in vivo*; (6) form and/or 20 maintain ordered spatial arrangements of differentiated tissues in vertebrates; (7) modulate cell death (e.g. cell survival); (8) regulate cell migration; and/or (9) modulate immune system function.

25 In addition to the nucleic acid molecules encoding CRSP proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be 30 complementary to an entire CRSP coding strand, or to only a portion thereof. In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding CRSP. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (e.g., the coding region of human CRSP-1 corresponds to SEQ 35 ID NO:3, the coding region of human CRSP-2 corresponds to SEQ ID NO:6, the coding region of human CRSP-3 corresponds to SEQ ID NO:9, the coding region of human CRSP-4 corresponds to SEQ ID NO:12, and the coding region of human CRSP-like-N

corresponds to SEQ ID NO:15). In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding CRSP. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (i.e., also referred 5 to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding CRSP disclosed herein (e.g., SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, or SEQ ID NO:12), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of 10 CRSP mRNA, but more preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of CRSP mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of CRSP mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention 15 can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the 20 antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5- 25 carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 30 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced 35 biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of

an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a CRSP protein to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier et al. (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue et al. (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al. (1987) *FEBS Lett.* 215:327-330).

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave CRSP mRNA transcripts to thereby inhibit translation of CRSP mRNA. A ribozyme having specificity for a CRSP-encoding nucleic acid can be designed based upon the nucleotide sequence of a CRSP cDNA disclosed herein (i.e., SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as

Accession Number 98633). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a CRSP-encoding mRNA. See, e.g., Cech et al. U.S. Patent No. 4,987,071; and Cech et al. U.S. Patent No. 5,116,742. Alternatively, 5 CRSP mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

Alternatively, CRSP gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the CRSP (e.g., the CRSP 10 promoter and/or enhancers) to form triple helical structures that prevent transcription of the CRSP gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. et al. (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

In yet another embodiment, the CRSP nucleic acid molecules of the present 15 invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, e.g., the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup B. et al. (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" 20 refer to nucleic acid mimics, e.g., DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis 25 protocols as described in Hyrup B. et al. (1996) *supra*; Perry-O'Keefe et al. PNAS 93: 14670-675.

PNAs of CRSP nucleic acid molecules can be used therapeutic and diagnostic 30 applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of CRSP nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (e.g., by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (e.g., S1 nucleases (Hyrup B. (1996) *supra*)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. et al. (1996) *supra*; Perry- 35 O'Keefe *supra*).

In another embodiment, PNAs of CRSP can be modified, (e.g., to enhance their stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by

the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of CRSP nucleic acid molecules can be generated which may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, (e.g., RNase H and DNA 5 polymerases), to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup B. (1996) *supra* and Finn P.J. *et al.* 10 (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs, e.g., 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. *et al.* (1989) *Nucleic Acid Res.* 17: 5973-88). PNA monomers are then coupled in a 15 stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn P.J. *et al.* (1996) *supra*). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser, K.H. *et al.* (1975) *Bioorganic Med. Chem. Lett.* 5: 1119-1124).

In other embodiments, the oligonucleotide may include other appended groups 20 such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. US* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. W089/10134, published April 25, 1988). In 25 addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, e.g., Krol *et al.* (1988) *BioTechniques* 6:958-976) or intercalating agents. (See, e.g., Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

30

II. Isolated CRSP Proteins and Anti-CRSP Antibodies

One aspect of the invention pertains to isolated CRSP proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise anti-CRSP antibodies. In one embodiment, native CRSP proteins can be 35 isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, CRSP proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, a

CRSP protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or 5 tissue source from which the CRSP protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of CRSP protein in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language "substantially free of 10 cellular material" includes preparations of CRSP protein having less than about 30% (by dry weight) of non-CRSP protein (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-CRSP protein, still more preferably less than about 10% of non-CRSP protein, and most preferably less than about 5% non-CRSP protein. When the CRSP protein or biologically active portion thereof is 15 recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" 20 includes preparations of CRSP protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" 25 includes preparations of CRSP protein having less than about 30% (by dry weight) of chemical precursors or non-CRSP chemicals, more preferably less than about 20% chemical precursors or non-CRSP chemicals, still more preferably less than about 10% chemical precursors or non-CRSP chemicals, and most preferably less than about 5% chemical precursors or non-CRSP chemicals.

Biologically active portions of a CRSP protein include peptides comprising 30 amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the CRSP protein, e.g., the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, which include less amino acids than the full length CRSP proteins, and exhibit at least one activity of a CRSP protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the CRSP protein. A biologically active portion of a CRSP protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

35 In one embodiment, a biologically active portion of a CRSP protein comprises at least a cysteine-rich region. In another embodiment, a biologically active portion of a CRSP protein comprises at least a cysteine-rich region, wherein the cysteine-rich region

includes at least one cysteine-rich domain. In yet another embodiment, a biologically active portion of a CRSP protein comprises at least a signal sequence.

In an alternative embodiment, a biologically active portion of a CRSP protein comprises a CRSP amino acid sequence lacking a signal sequence.

5 It is to be understood that a preferred biologically active portion of a CRSP protein of the present invention may contain at least one of the above-identified structural domains. A more preferred biologically active portion of a CRSP protein may contain at least two of the above-identified structural domains. An even more preferred biologically active portion of a CRSP protein may contain at least three of the
10 above-identified structural domains. A particularly preferred biologically active portion of a CRSP protein of the present invention may contain at least four of the above-identified structural domains.

15 Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native CRSP protein.

In a preferred embodiment, the CRSP protein has an amino acid sequence shown in SEQ ID NO:2 or an amino acid sequence at least about 55% homologous to SEQ ID NO:2. In another preferred embodiment, the CRSP protein has an amino acid sequence shown in SEQ ID NO:5 or an amino acid sequence at least about 35% homologous to
20 SEQ ID NO:5. In another preferred embodiment, the CRSP protein has an amino acid sequence shown in SEQ ID NO:8 or an amino acid sequence at least about 85% homologous to SEQ ID NO:8. In another preferred embodiment, the CRSP protein has an amino acid sequence shown in SEQ ID NO:11 or an amino acid sequence at least about 35% homologous to SEQ ID NO:11. In still another preferred embodiment, a
25 CRSP protein of the present invention comprises an amino acid sequence which is at least about 30-35%, preferably about 40-45%, more preferably about 50-55%, even more preferably about 60-65%, and even more preferably at least about 70-75%, 80-85%, 90-95% or more homologous to the amino acid sequences shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11.

30 In other embodiments, the CRSP protein is substantially homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, and, preferably, retains the functional activity of the protein of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another
35 embodiment, the CRSP protein is a protein which comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11 and, preferably, retains the functional activity

of the CRSP proteins of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, respectively. Preferably, the protein is at least about 70% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, more preferably at least about 80% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, even more preferably at least about 90% homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, and most preferably at least about 95% or more homologous to SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11.

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (i.e., as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = # of identical positions/total # of positions x 100). The determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to CRSP nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to CRSP protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for

comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The invention also provides CRSP chimeric or fusion proteins. As used herein, a CRSP "chimeric protein" or "fusion protein" comprises a CRSP polypeptide operatively linked to a non-CRSP polypeptide. A "CRSP polypeptide" refers to a polypeptide having an amino acid sequence corresponding to CRSP, whereas a "non-CRSP polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the CRSP protein, e.g., a protein which is different from the CRSP protein and which is derived from the same or a different 5 organism. Within a CRSP fusion protein the CRSP polypeptide can correspond to all or a portion of a CRSP protein. In a preferred embodiment, a CRSP fusion protein comprises at least one biologically active portion of a CRSP protein. In another preferred embodiment, a CRSP fusion protein comprises at least two biologically active 10 portions of a CRSP protein. In another preferred embodiment, a CRSP fusion protein comprises at least three biologically active portions of a CRSP protein. Within the fusion protein, the term "operatively linked" is intended to indicate that the CRSP polypeptide and the non-CRSP polypeptide are fused in-frame to each other. The non-CRSP polypeptide can be fused to the N-terminus or C-terminus of the CRSP 15 polypeptide.

20 For example, in one embodiment, the fusion protein is a GST-CRSP fusion protein in which the CRSP sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant CRSP.

25 In another embodiment, the fusion protein is a CRSP protein containing a heterologous signal sequence at its N-terminus. For example, the native CRSP signal sequence (i.e., about amino acids 1 to 23 of SEQ ID NO:2) can be removed and replaced with a signal sequence from another protein. In certain host cells (e.g., mammalian host cells), expression and/or secretion of CRSP can be increased through use of a heterologous signal sequence.

30 In yet another embodiment, the fusion protein is a CRSP-immunoglobulin fusion protein in which the CRSP sequences comprising primarily the CRSP cysteine-rich regions are fused to sequences derived from a member of the immunoglobulin protein family. Soluble derivatives have also been made of cell surface glycoproteins in the immunoglobulin gene superfamily consisting of an extracellular domain of the cell 35 surface glycoprotein fused to an immunoglobulin constant (Fc) region (see e.g., Capon, D.J. *et al.* (1989) *Nature* 337:525-531 and Capon U.S. Patents 5,116,964 and 5,428,130 [CD4-IgG1 constructs]; Linsley, P.S. *et al.* (1991) *J. Exp. Med.* 173:721-730 [a CD28-IgG1 construct and a B7-1-IgG1 construct]; and Linsley, P.S. *et al.* (1991) *J. Exp. Med.*

174:561-569 and U.S. Patent 5,434,131[a CTLA4-IgG1]). Such fusion proteins have proven useful for modulating receptor-ligand interactions. Soluble derivatives of cell surface proteins of the tumor necrosis factor receptor (TNFR) superfamily proteins have been made consisting of an extracellular domain of the cell surface receptor fused to an 5 immunoglobulin constant (Fc) region (See for example Moreland *et al.* (1997) *N. Engl. J. Med.* 337(3):141-147; van der Poll *et al.* (1997) *Blood* 89(10):3727-3734; and Ammann *et al.* (1997) *J. Clin. Invest.* 99(7):1699-1703.)

The CRSP-immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction 10 between a CRSP ligand and a CRSP receptor on the surface of a cell, to thereby suppress CRSP-mediated signal transduction *in vivo*. The CRSP-immunoglobulin fusion proteins can be used to affect the bioavailability of a CRSP cognate receptor. Inhibition of the CRSP ligand/CRSP interaction may be useful therapeutically for both the treatment of differentiative or proliferative disorders, as well as modulating (e.g., 15 promoting or inhibiting) developmental responses, cell adhesion, and/or cell fate. Moreover, the CRSP-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-CRSP antibodies in a subject, to purify CRSP ligands and in screening assays to identify molecules which inhibit the interaction of CRSP with a CRSP ligand.

20 Preferably, a CRSP chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, 25 filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene 30 fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). A CRSP-encoding nucleic acid can be cloned into such an expression vector such that the fusion 35 moiety is linked in-frame to the CRSP protein.

The present invention also pertains to variants of the CRSP proteins which function as either CRSP agonists (mimetics) or as CRSP antagonists. Variants of the

CRSP proteins can be generated by mutagenesis, e.g., discrete point mutation or truncation of a CRSP protein. An agonist of the CRSP proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a CRSP protein. An antagonist of a CRSP protein can inhibit one or more of the activities 5 of the naturally occurring form of the CRSP protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the CRSP protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the 10 protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the CRSP protein.

In one embodiment, variants of a CRSP protein which function as either CRSP agonists (mimetics) or as CRSP antagonists can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, of a CRSP protein for CRSP protein 15 agonist or antagonist activity. In one embodiment, a variegated library of CRSP variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of CRSP variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential CRSP sequences is expressible as 20 individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g., for phage display) containing the set of CRSP sequences therein. There are a variety of methods which can be used to produce libraries of potential CRSP variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an 25 appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential CRSP sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477.

In addition, libraries of fragments of a CRSP protein coding sequence can be used to generate a variegated population of CRSP fragments for screening and subsequent selection of variants of a CRSP protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR 35 fragment of a CRSP coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from

different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the CRSP protein.

5 Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of CRSP proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which 10 enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify CRSP variants (Arkin and Yourvan 15 (1992) *PNAS* 89:7811-7815; Delgrave et al. (1993) *Protein Engineering* 6(3):327-331).

In one embodiment, cell based assays can be exploited to analyze a variegated CRSP library. For example, a library of expression vectors can be transfected into a cell 20 line which ordinarily responds to a particular ligand in a CRSP-dependent manner. The transfected cells are then contacted with the ligand and the effect of expression of the mutant on signaling by the ligand can be detected, e.g., by measuring any of a number of immune cell responses. Plasmid DNA can then be recovered from the cells which score 25 for inhibition, or alternatively, potentiation of ligand induction, and the individual clones further characterized.

An isolated CRSP protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind CRSP using standard techniques for polyclonal and monoclonal antibody preparation. A full-length CRSP protein can be used or, alternatively, the invention provides antigenic peptide fragments of CRSP for 30 use as immunogens. The antigenic peptide of CRSP comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11 and encompasses an epitope of CRSP such that an antibody raised against the peptide forms a specific immune complex with CRSP. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 35 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at least 30 amino acid residues.

Preferred epitopes encompassed by the antigenic peptide are regions of CRSP that are located on the surface of the protein, e.g., hydrophilic regions.

A CRSP immunogen typically is used to prepare antibodies by immunizing a suitable subject, (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed CRSP protein or a chemically synthesized CRSP polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic CRSP preparation induces a polyclonal anti-CRSP antibody response.

Accordingly, another aspect of the invention pertains to anti-CRSP antibodies. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as CRSP. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind CRSP. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of CRSP. A monoclonal antibody composition thus typically displays a single binding affinity for a particular CRSP protein with which it immunoreacts.

Polyclonal anti-CRSP antibodies can be prepared as described above by immunizing a suitable subject with a CRSP immunogen. The anti-CRSP antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized CRSP. If desired, the antibody molecules directed against CRSP can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the anti-CRSP antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also, Brown et al. (1981) *J. Immunol.* 127:539-46; Brown et al. (1980) *J. Biol. Chem.* 255:4980-83; Yeh et al. (1976) *PNAS* 76:2927-31; and Yeh et al. (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor et al. (1983) *Immunol Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and Cancer Therapy*,

Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*,

5 54:387-402; M. L. Gefter et al. (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a CRSP immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds CRSP.

10 Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating an anti-CRSP monoclonal antibody (see, e.g., G. Galfre et al. (1977) *Nature* 266:55052; Gefter et al. *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell

15 lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma

20 cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for

25 antibodies that bind CRSP, e.g., using a standard ELISA assay.

30 Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal anti-CRSP antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with CRSP to thereby isolate immunoglobulin library members that bind CRSP. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally,

examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner et al. U.S. Patent No. 5,223,409; Kang et al. PCT International Publication No. WO 92/18619; Dower et al. PCT International Publication No. WO 91/17271; Winter et al. PCT International Publication WO 92/20791; Markland et al. PCT International Publication No. WO 92/15679; Breitling et al. PCT International Publication WO 93/01288; McCafferty et al. PCT International Publication No. WO 92/01047; Garrard et al. PCT International Publication No. WO 92/09690; Ladner et al. PCT International Publication No. WO 90/02809; Fuchs et al. (1991) *Bio/Technology* 9:1370-1372; Hay et al. (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J* 12:725-734; Hawkins et al. (1992) *J. Mol. Biol.* 226:889-896; Clarkson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *PNAS* 89:3576-3580; Garrad et al. (1991) *Bio/Technology* 9:1373-1377; Hoogenboom et al. (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas et al. (1991) *PNAS* 88:7978-7982; and McCafferty et al. *Nature* (1990) 348:552-554.

Additionally, recombinant anti-CRSP antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson et al. International Application No. PCT/US86/02269; Akira, et al. European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT International Publication No. WO 86/01533; Cabilly et al. U.S. Patent No. 4,816,567; Cabilly et al. European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *PNAS* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *PNAS* 84:214-218; Nishimura et al. (1987) *Canc. Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; and Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559); Morrison, S. L. (1985) *Science* 229:1202-1207; Oi et al. (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones et al. (1986) *Nature* 321:552-525; Verhoeyen et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060.

An anti-CRSP antibody (e.g., monoclonal antibody) can be used to isolate CRSP by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-CRSP antibody can facilitate the purification of natural CRSP from cells and of recombinantly produced CRSP expressed in host cells. Moreover, an anti-CRSP antibody can be used to detect CRSP protein (e.g., in a cellular lysate or cell

supernatant) in order to evaluate the abundance and pattern of expression of the CRSP protein. Anti-CRSP antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, e.g., to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include 5 streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include 10 luciferase, luciferin, and aequorin, and examples of suitable radioactive material include 15 ^{125}I , ^{131}I , ^{35}S or ^3H .

III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding CRSP (or a portion thereof). As used herein, 20 the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication 25 in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they 30 are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral 35 vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is

5 operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (e.g., in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory

10 sequence" is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and

15 those which direct expression of the nucleotide sequence only in certain host cells (e.g., tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce

20 proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (e.g., CRSP proteins, mutant forms of CRSP, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of CRSP in prokaryotic or eukaryotic cells. For example, CRSP can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

30 Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein;

35 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion

moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B.

5 and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in CRSP activity assays, in CRSP ligand

binding (e.g., direct assays or competitive assays described in detail below), to generate

10 antibodies specific for CRSP proteins, as examples. In a preferred embodiment, a CRSP fusion expressed in a retroviral expression vector of the present invention can be utilized to infect bone marrow cells which are subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g six (6) weeks).

15 Examples of suitable inducible non-fusion *E. coli* expression vectors include

pTrc (Amann et al., (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *Gene*

Expression Technology: Methods in Enzymology 185, Academic Press, San Diego,

California (1990) 60-89). Target gene expression from the pTrc vector relies on host

RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene

20 expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion

promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral

polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident λ

prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

25 One strategy to maximize recombinant protein expression in *E. coli* is to express

the protein in a host bacteria with an impaired capacity to proteolytically cleave the

recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in*

Enzymology 185, Academic Press, San Diego, California (1990) 119-128). Another

30 strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an

expression vector so that the individual codons for each amino acid are those

preferentially utilized in *E. coli* (Wada et al., (1992) *Nucleic Acids Res.* 20:2111-2118).

Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the CRSP expression vector is a yeast expression vector.

35 Examples of vectors for expression in yeast *S. cerevisiae* include pYEPSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-

943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (InVitrogen Corp, San Diego, CA).

Alternatively, CRSP can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured 5 insect cells (e.g., Sf 9 cells) include the pAc series (Smith et al. (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC 10 (Kaufman et al. (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, 15 cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type 20 (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert et al. (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and 25 Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji et al. (1983) *Cell* 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (e.g., the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund et al. (1985) *Science* 230:912-916), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and 30 European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a 35 DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an

RNA molecule which is antisense to CRSP mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which 5 direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene 10 expression using antisense genes see Weintraub, H. et al., Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such 15 terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

20 A host cell can be any prokaryotic or eukaryotic cell. For example, CRSP protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

25 Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or 30 transfecting host cells can be found in Sambrook, et al. (*Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

35 For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred

selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding CRSP or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) CRSP protein. Accordingly, the invention further provides methods for producing CRSP protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding CRSP has been introduced) in a suitable medium such that CRSP protein is produced. In another embodiment, the method further comprises isolating CRSP from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which CRSP-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous CRSP sequences have been introduced into their genome or homologous recombinant animals in which endogenous CRSP sequences have been altered. Such animals are useful for studying the function and/or activity of CRSP and for identifying and/or evaluating modulators of CRSP activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous CRSP gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, e.g., an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing CRSP-encoding nucleic acid into the male pronuclei of a fertilized oocyte, e.g., by microinjection, retroviral infection, and allowing the oocyte to develop in a

pseudopregnant female foster animal. The human CRSP cDNA sequence of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a nonhuman homologue of a human CRSP gene, such as a mouse CRSP gene, can be isolated based on hybridization 5 to the human CRSP cDNA (described further in subsection I above) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably linked to the CRSP transgene to direct expression of CRSP protein to particular cells. Methods for generating transgenic 10 animals via embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder et al., U.S. Patent No. 4,873,191 by Wagner et al. and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of 15 other transgenic animals. A transgenic founder animal can be identified based upon the presence of the CRSP transgene in its genome and/or expression of CRSP mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene encoding CRSP can further be bred to other transgenic animals carrying other 20 transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a CRSP gene into which a deletion, addition or substitution has been introduced to thereby alter, e.g., functionally disrupt, the CRSP gene. The CRSP gene can be a human gene (e.g., the cDNA of SEQ ID NO:3, SEQ ID NO: 6, SEQ ID NO:9 25 or SEQ ID NO:12), but more preferably, is a non-human homologue of a human CRSP gene. For example, a mouse CRSP gene of SEQ ID NO:16 can be used to construct a homologous recombination vector suitable for altering an endogenous CRSP gene in the mouse genome. In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous CRSP gene is functionally disrupted (i.e., 30 no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous CRSP gene is mutated or otherwise altered but still encodes functional protein (e.g., the upstream regulatory region can be altered to thereby alter the expression of the endogenous CRSP protein). In the homologous recombination vector, 35 the altered portion of the CRSP gene is flanked at its 5' and 3' ends by additional nucleic acid of the CRSP gene to allow for homologous recombination to occur between the exogenous CRSP gene carried by the vector and an endogenous CRSP gene in an

embryonic stem cell. The additional flanking CRSP nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see e.g., Thomas, K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of 5 homologous recombination vectors). The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced CRSP gene has homologously recombined with the endogenous CRSP gene are selected (see e.g., Li, E. et al. (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (e.g., a mouse) to form aggregation chimeras (see e.g., Bradley, A. in 10 *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed 15 animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, A. (1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International Publication Nos.: WO 90/11354 by Le Mouellec et al.; WO 91/01140 by Smithies et al.; WO 92/0968 by Zijlstra et al.; and WO 93/04169 by Berns et al.

20 In another embodiment, transgenic non-humans animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, e.g., Lakso et al. (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase 25 system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, e.g., by mating two transgenic animals, one containing a transgene encoding a 30 selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. et al. (1997) *Nature* 385:810-813. In brief, a cell, e.g., a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be 35 fused, e.g., through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyst and then transferred to

pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell, e.g., the somatic cell, is isolated.

IV. Pharmaceutical Compositions

5 The CRSP nucleic acid molecules, CRSP proteins, and anti-CRSP antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is 10 intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is 15 contemplated. Supplementary active compounds can also be incorporated into the compositions.

 A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), 20 transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as 25 ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be 30 enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

 Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic 35 water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage

and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity
5 can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include
10 isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active
15 compound (e.g., a CRSP protein or anti-CRSP antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In
20 the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They
25 can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically
30 compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For 5 transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active 10 compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

15 In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. 20 Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled 25 in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound 30 calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

35 Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose

therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a 5 delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little 10 or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of 15 the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The nucleic acid molecules of the invention can be inserted into vectors and used 20 as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see e.g., Chen et al. (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy 25 vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene 30 delivery system.

The pharmaceutical compositions can be included in a container, pack, or 35 dispenser together with instructions for administration.

V. Uses and Methods of the Invention

The molecules of the present invention (e.g., nucleic acid molecules, proteins, protein homologues, and antibodies described herein) can be used in one or more of the 35 following methods: a) screening assays; b) predictive medicine (e.g., diagnostic assays, prognostic assays, monitoring clinical trials, and pharmacogenetics); and c) methods of treatment (e.g., therapeutic and prophylactic). As described herein, a CRSP protein of

the invention has one or more of the following activities: intracellular calcium, an increase in phosphatidylinositol or other molecule, and can result, e.g., in phosphorylation of specific proteins, a modulation of gene transcription and any of the other biological activities set forth herein.

5 In a preferred embodiment, a CRSP activity is at least one or more of the following activities: (i) interaction of a CRSP protein with and/or binding to a second molecule, (e.g., a protein, such as a CRSP (e.g., CRSP-1) receptor, a soluble form of a CRSP receptor, a receptor for a member of the *wnt* family of signaling proteins, or a non-CRSP signaling molecule); (ii) interaction of a CRSP protein with an intracellular 10 protein via a membrane-bound CRSP receptor; (iii) complex formation between a soluble CRSP protein and a second soluble CRSP binding partner (e.g., a non-CRSP protein molecule or a second CRSP protein molecule); (iv) interaction with other extracellular proteins (e.g., regulation of *wnt*-dependent cellular adhesion to extracellular matrix components); (v) binding to and eliminating an undesirable molecule (e.g., a 15 detoxifying activity or defense function); and/or (vi) an enzymatic activity, and can thus be used in, for example, (1) modulation of cellular signal transduction, either *in vitro* or *in vivo* (e.g., antagonism of the activity of members of the *wnt* family of secreted proteins or suppression of *wnt*-dependent signal transduction); (2) regulation of communication between cells (e.g., regulation of *wnt*-dependent cell-cell interactions); 20 (3) regulation of expression of genes whose expression is modulated by binding of CRSP (e.g., CRSP-1) to a receptor; (4) regulation of gene transcription in a cell involved in development or differentiation, either *in vitro* or *in vivo* (e.g., induction of cellular differentiation); (5) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a differentiation-specific protein; (6) 25 regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a second secreted protein; (7) regulation of gene transcription in a cell involved in development or differentiation, wherein at least one gene encodes a signal transduction molecule; (8) regulation of cellular proliferation, either *in vitro* or *in vivo* (e.g., induction of cellular proliferation or inhibition of 30 proliferation as in the case of suppression of tumorogenesis (e.g., suppression of glioblastoma formation)); (9) formation and maintenance of ordered spatial arrangements of differentiated tissues in vertebrates, both adult and embryonic (e.g., induction of head formation during vertebrate development or maintenance of hematopoietic progenitor cells); (10) modulation of cell death, such as stimulation of 35 cell survival; (11) regulating cell migration; and/or (12) immune modulation.

Accordingly one embodiment of the present invention involves a method of use (e.g., a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of

treatment) wherein a molecule of the present invention (e.g., a CRSP protein, CRSP nucleic acid, or a CRSP modulator) is used, for example, to diagnose, prognosis and/or treat a disease and/or condition in which any of the aforementioned activities (i.e., activities (i) - (vi) and (1) - (12) in the above paragraph) is indicated. In another 5 embodiment, the present invention involves a method of use (e.g., a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (e.g., a CRSP protein, CRSP nucleic acid, or a CRSP modulator) is used, for example, for the diagnosis, prognosis, and/or treatment of subjects, preferably a human subject, in which any of the aforementioned activities is 10 pathologically perturbed. In a preferred embodiment, the methods of use (e.g., diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a subject, preferably a human subject, a molecule of the present invention (e.g., a CRSP protein, CRSP nucleic acid, or a CRSP modulator) for the diagnosis, prognosis, and/or therapeutic treatment. In another embodiment, the methods 15 of use (e.g., diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a human subject a molecule of the present invention (e.g., a CRSP protein, CRSP nucleic acid, or a CRSP modulator).

Other embodiments of the invention pertain to the use of isolated nucleic acid molecules of the invention can be used, for example, to express CRSP protein (e.g., via a 20 recombinant expression vector in a host cell in gene therapy applications), to detect CRSP mRNA (e.g., in a biological sample) or a genetic alteration in a CRSP gene, and to modulate CRSP activity, as described further below. In addition, the CRSP proteins can be used to screen drugs or compounds which modulate the CRSP activity as well as to treat disorders characterized by insufficient or excessive production of CRSP protein 25 or production of CRSP protein forms which have decreased or aberrant activity compared to CRSP wild type protein (e.g., developmental disorders or proliferative diseases such as cancer as well as diseases, conditions or disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in a defense mechanism). Moreover, the anti-CRSP antibodies of the 30 invention can be used to detect and isolate CRSP proteins, regulate the bioavailability of CRSP proteins, and modulate CRSP activity. The term "an aberrant activity", as applied to an activity of a protein such as CRSP (e.g., CRSP-1), refers to an activity which differs from the activity of the wild-type or native protein or which differs from the activity of the protein in a healthy subject. An activity of a protein can be aberrant 35 because it is stronger than the activity of its native counterpart. Alternatively, an activity can be aberrant because it is weaker or absent related to the activity of its native counterpart. An aberrant activity can also be a change in an activity. For example an

aberrant protein can interact with a different protein relative to its native counterpart. A cell can have an aberrant CRSP (e.g., CRSP-1) activity due to overexpression or underexpression of the gene encoding CRSP.

5

A. Screening Assays:

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, i.e., candidate or test compounds or agents (e.g., peptides, peptidomimetics, small molecules or other drugs) which bind to CRSP proteins or have 10 a stimulatory or inhibitory effect on, for example, CRSP expression or CRSP activity.

Modulators can include, for example, CRSP agonist and/or CRSP antagonists. The term "agonist", as used herein, is meant to refer to an agent that mimics or upregulates (e.g. potentiates or supplements) a CRSP (e.g., CRSP-1) bioactivity. A CRSP agonist can be a compound which mimics a bioactivity of a CRSP protein, such as transduction of a 15 signal from a CRSP receptor, by, e.g., interacting with a CRSP-1 receptor. A CRSP agonist can also be a compound that upregulates expression of a CRSP gene. A CRSP agonist can also be a compound which modulates the expression or activity of a protein which is located downstream of a CRSP receptor, thereby mimicking or enhancing the effect of binding of CRSP to a CRSP receptor.

20 "Antagonist" as used herein is meant to refer to an agent that inhibits, decreases or suppresses a CRSP (e.g., CRSP-1) bioactivity. An antagonist can be a compound which decreases signalling from a CRSP protein, e.g., a compound that is capable of binding to CRSP-1 or to a CRSP-1 receptor. A preferred CRSP antagonist inhibits the interaction between a CRSP protein and another molecule, such as a CRSP receptor.

25 Alternatively, a CRSP antagonist can be a compound that downregulates expression of a CRSP gene. A CRSP antagonist can also be a compound which modulates the expression or activity of a protein which is located downstream of a CRSP receptor, thereby antagonizing the effect of binding of CRSP to a CRSP receptor.

In one embodiment, the invention provides assays for screening candidate or test 30 compounds which bind to or modulate the activity of a CRSP protein or polypeptide or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of a CRSP receptor. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, 35 including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-

compound' library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

5 Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in
10 Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds may be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage
15 (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra*).

In one embodiment, an assay is a cell-based assay in which a cell which
20 expresses a CRSP receptor on the cell surface is contacted with a test compound and the
ability of the test compound to bind to a CRSP receptor determined. The cell, for
example, can be of mammalian origin or a yeast cell. Determining the ability of the test
compound to bind to a CRSP receptor can be accomplished, for example, by coupling
the test compound with a radioisotope or enzymatic label such that binding of the test
compound to the CRSP receptor can be determined by detecting the labeled compound
25 in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H ,
either directly or indirectly, and the radioisotope detected by direct counting of
radioemmission or by scintillation counting. Alternatively, test compounds can be
enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase,
or luciferase, and the enzymatic label detected by determination of conversion of an
30 appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a test
compound to interact with a CRSP receptor without the labeling of any of the
interactants. For example, a microphysiometer can be used to detect the interaction of a
test compound with a CRSP receptor without the labeling of either the test compound or
35 the receptor. McConnell, H. M. *et al.* (1992) *Science* 257:1906-1912. As used herein, a
"microphysiometer" (e.g., CytosensorTM) is an analytical instrument that measures the

rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between ligand and receptor.

In a preferred embodiment, the assay comprises contacting a cell which
5 expresses a CRSP receptor on the cell surface with a CRSP protein or biologically-
active portion thereof, to form an assay mixture, contacting the assay mixture with a test
compound, and determining the ability of the test compound to interact with a CRSP
receptor, wherein determining the ability of the test compound to interact with a CRSP
receptor comprises determining the ability of the test compound to preferentially bind to
10 the CRSP receptor as compared to the ability of CRSP, or a biologically active portion
thereof, to bind to the receptor.

In another embodiment, an assay is a cell-based assay comprising contacting a
cell expressing a CRSP target molecule with a test compound and determining the
ability of the test compound to modulate (e.g. stimulate or inhibit) the activity of the
15 CRSP target molecule. Determining the ability of the test compound to modulate the
activity of a CRSP target molecule can be accomplished, for example, by determining
the ability of the CRSP protein to bind to or interact with the CRSP target molecule.

Determining the ability of the CRSP protein to bind to or interact with a CRSP
target molecule can be accomplished by one of the methods described above for
20 determining direct binding. In a preferred embodiment, determining the ability of the
CRSP protein to bind to or interact with a CRSP target molecule can be accomplished
by determining the activity of the target molecule. For example, the activity of the
target molecule can be determined by detecting induction of a cellular second messenger
of the target (i.e. intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting
25 catalytic/enzymatic activity of the target an appropriate substrate, detecting the induction
of a reporter gene (comprising a CRSP-responsive regulatory element operatively linked
to a nucleic acid encoding a detectable marker, e.g., luciferase), or detecting a cellular
response, for example, development, differentiation or rate of proliferation.

In yet another embodiment, an assay of the present invention is a cell-free assay
30 in which a CRSP protein or biologically active portion thereof is contacted with a test
compound and the ability of the test compound to bind to the CRSP protein or
biologically active portion thereof is determined. Binding of the test compound to the
CRSP protein can be determined either directly or indirectly as described above. In a
preferred embodiment, the assay includes contacting the CRSP protein or biologically
35 active portion thereof with a known compound which binds CRSP to form an assay
mixture, contacting the assay mixture with a test compound, and determining the ability
of the test compound to interact with a CRSP protein, wherein determining the ability of

the test compound to interact with a CRSP protein comprises determining the ability of the test compound to preferentially bind to CRSP or biologically active portion thereof as compared to the known compound.

In another embodiment, the assay is a cell-free assay in which a CRSP protein or 5 biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate (e.g., stimulate or inhibit) the activity of the CRSP protein or biologically active portion thereof is determined. Determining the ability of the test compound to modulate the activity of a CRSP protein can be accomplished, for example, by determining the ability of the CRSP protein to bind to a CRSP target 10 molecule by one of the methods described above for determining direct binding.

Determining the ability of the CRSP protein to bind to a CRSP target molecule can also be accomplished using a technology such as real-time Biomolecular Interaction Analysis (BIA). Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo 15 *et al.* (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIACoreTM). Changes in the optical phenomenon surface plasmon resonance (SPR) 20 can be used as an indication of real-time reactions between biological molecules.

In an alternative embodiment, determining the ability of the test compound to modulate the activity of a CRSP protein can be accomplished by determining the ability 25 of the CRSP protein to further modulate the activity of a CRSP target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay involves contacting a CRSP protein or biologically active portion thereof with a known compound which binds the 25 CRSP protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the CRSP protein, wherein determining the ability of the test compound to interact with the CRSP protein comprises determining the ability of the CRSP protein to preferentially bind to or modulate the activity of a CRSP target molecule.

30 In many drug screening programs which test libraries of compounds and natural extracts, high throughput assays are desirable in order to maximize the number of compounds surveyed in a given period of time. Assays which are performed in cell-free systems, such as may be derived with purified or semi-purified proteins, are often preferred as "primary" screens in that they can be generated to permit rapid development 35 and relatively easy detection of an alteration in a molecular target which is mediated by a test compound. Moreover, the effects of cellular toxicity and/or bioavailability of the test compound can be generally ignored in the *in vitro* system, the assay instead being

focused primarily on the effect of the drug on the molecular target as may be manifest in an alteration of binding affinity with upstream or downstream elements. Accordingly, in an exemplary screening assay of the present invention, the compound of interest is contacted with a CRSP (e.g., CRSP-1) protein or a CRSP (e.g., CRSP-1) binding

5 partner, e.g., a receptor. The receptor can be soluble or the receptor can be present on a cell surface. To the mixture of the compound and the CRSP protein or CRSP binding partner is then added a composition containing a CRSP binding partner or a CRSP protein, respectively. Detection and quantification of complexes of CRSP proteins and CRSP binding partners provide a means for determining a compound's efficacy at
10 inhibiting (or potentiating) complex formation between CRSP and a binding partner. The efficacy of the compound can be assessed by generating dose response curves from data obtained using various concentrations of the test compound. Moreover, a control assay can also be performed to provide a baseline for comparison. In the control assay, isolated and purified CRSP polypeptide or binding partner is added to a composition
15 containing the CRSP binding partner or CRSP polypeptide, and the formation of a complex is quantitated in the absence of the test compound.

The cell-free assays of the present invention are amenable to use of both soluble and/or membrane-bound forms of isolated proteins (e.g. CRSP proteins or biologically active portions thereof or CRSP target molecules). In the case of cell-free assays in

20 which a membrane-bound form an isolated protein is used (e.g., a CRSP target molecule or receptor) it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-
25 methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecygly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylammonio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylammonio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate.

In more than one embodiment of the above assay methods of the present
30 invention, it may be desirable to immobilize either CRSP or its target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to a CRSP protein, or interaction of a CRSP protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel
35 suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a

matrix. For example, glutathione-S-transferase/ CRSP fusion proteins or glutathione-S-transferase/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed 5 target protein or CRSP protein, and the mixture incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can 10 be dissociated from the matrix, and the level of CRSP binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either a CRSP protein or a CRSP target molecule can be immobilized utilizing conjugation of biotin and streptavidin.

15 Biotinylated CRSP protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with CRSP protein or target molecules but which do not interfere with binding of the CRSP protein to its 20 target molecule can be derivatized to the wells of the plate, and unbound target or CRSP protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the CRSP protein or target molecule, as well as enzyme-linked assays which rely on detecting an 25 enzymatic activity associated with the CRSP protein or target molecule.

In another embodiment, modulators of CRSP expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of CRSP mRNA or protein in the cell is determined. The level of expression of CRSP mRNA or protein in the presence of the candidate compound is compared to the level of 30 expression of CRSP mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of CRSP expression based on this comparison. For example, when expression of CRSP mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of CRSP mRNA or protein 35 expression. Alternatively, when expression of CRSP mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of CRSP mRNA or protein

expression. The level of CRSP mRNA or protein expression in the cells can be determined by methods described herein for detecting CRSP mRNA or protein.

In yet another aspect of the invention, the CRSP proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Patent No.

5 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.* 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with CRSP ("CRSP-binding proteins" or "CRSP-bp") and modulate CRSP activity. Such CRSP-binding proteins are also likely to be involved in 10 the propagation of signals by the CRSP proteins as, for example, downstream elements of a CRSP-mediated signaling pathway. Alternatively, such CRSP-binding proteins are likely to be cell-surface molecules associated with non-CRSP expressing cells, wherein such CRSP-binding proteins are involved in signal transduction.

The two-hybrid system is based on the modular nature of most transcription 15 factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for a CRSP protein is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is 20 fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a CRSP-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the 25 transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the CRSP protein.

This invention further pertains to novel agents identified by the above-described screening assays and to processes for producing such agents by use of these assays.

30 Accordingly, in one embodiment, the present invention includes a compound or agent obtainable by a method comprising the steps of any one of the aforementioned screening assays (e.g., cell-based assays or cell-free assays). For example, in one embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a CRSP target molecule with a test compound and the determining 35 the ability of the test compound to bind to, or modulate the activity of, the CRSP target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a CRSP target

molecule with a CRSP protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of, the CRSP target molecule. In another embodiment, the invention includes a compound or agent

- 5 obtainable by a method comprising contacting a CRSP protein or biologically active portion thereof with a test compound and determining the ability of the test compound to bind to, or modulate (e.g., stimulate or inhibit) the activity of, the CRSP protein or biologically active portion thereof. In yet another embodiment, the present invention included a compound or agent obtainable by a method comprising contacting a CRSP
- 10 protein or biologically active portion thereof with a known compound which binds the CRSP protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of the CRSP protein.

Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a CRSP modulating agent, an antisense CRSP nucleic acid molecule, a CRSP-specific antibody, or a CRSP-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent.

The present invention also pertains to uses of novel agents identified by the above-described screening assays for diagnoses, prognoses, and treatments as described herein. Accordingly, it is within the scope of the present invention to use such agents in the design, formulation, synthesis, manufacture, and/or production of a drug or pharmaceutical composition for use in diagnosis, prognosis, or treatment, as described herein. For example, in one embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition by reference to the structure and/or properties of a compound obtainable by one of the above-described screening assays. For example, a drug or pharmaceutical composition can be synthesized based on the structure and/or properties of a compound obtained by a method in which a cell which expresses a CRSP target molecule is contacted with a test compound and the ability of the test compound to bind to, or modulate the activity of, the CRSP target molecule is determined. In another exemplary embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition based on the structure and/or properties of a compound obtainable by a method in which a CRSP protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to, or modulate (e.g.,

stimulate or inhibit) the activity of, the CRSP protein or biologically active portion thereof is determined.

B. Detection Assays

5 Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue 10 typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

1. Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this 15 sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the CRSP nucleotide sequences, described herein, can be used to map the location of the CRSP genes on a chromosome. The mapping of the CRSP sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

20 Briefly, CRSP genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the CRSP nucleotide sequences. Computer analysis of the CRSP sequences can be used to predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing 25 individual human chromosomes. Only those hybrids containing the human gene corresponding to the CRSP sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (e.g., human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the 30 mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse 35 chromosomes, allowing easy mapping of individual genes to specific human chromosomes. (D'Eustachio P. et al. (1983) *Science* 220:919-924). Somatic cell hybrids

containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be 5 assigned per day using a single thermal cycler. Using the CRSP nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a 9o, 1p, or 1v sequence to its chromosome include *in situ* hybridization (described in Fan, Y. et al. (1990) *PNAS*, 87:6223-27), pre-screening with labeled flow- 10 sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been 15 blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher 20 likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will suffice to get good results at a reasonable amount of time. For a review of this technique, see Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques* (Pergamon Press, New York 1988).

25 Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the 30 chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. (Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man*, available on-line through Johns Hopkins University Welch Medical Library). The 35 relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and unaffected with a disease associated with the CRSP gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease.

5 Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

10

2. Tissue Typing

The CRSP sequences of the present invention can also be used to identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for

15 identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The sequences of the present invention are useful as additional DNA markers

20 for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the CRSP nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3' ends of the

25 sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

30 Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the present invention can be used to obtain such identification sequences from individuals and from tissue. The CRSP nucleotide sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of

35 about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the

noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID NO:10, can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100 5 bases. If predicted coding sequences, such as those in SEQ ID NO:3, SEQ ID NO:6, SEQ ID NO:9, or SEQ ID NO:12 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

If a panel of reagents from CRSP nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can 10 later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

3. Use of Partial CRSP Sequences in Forensic Biology

15 DNA-based identification techniques can also be used in forensic biology. Forensic biology is a scientific field employing genetic typing of biological evidence found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify DNA sequences taken from very small biological samples such as tissues, e.g., hair or 20 skin, or body fluids, e.g., blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide 25 reagents, e.g., PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (i.e. another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments. Sequences targeted to noncoding regions of SEQ ID NOs:1, SEQ 30 ID NO:4, SEQ ID NO:7, or SEQ ID NO:10 are particularly appropriate for this use as greater numbers of polymorphisms occur in the noncoding regions, making it easier to differentiate individuals using this technique. Examples of polynucleotide reagents include the CRSP nucleotide sequences or portions thereof, e.g., fragments derived from the noncoding regions of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, or SEQ ID 35 NO:10, having a length of at least 20 bases, preferably at least 30 bases.

The CRSP nucleotide sequences described herein can further be used to provide polynucleotide reagents, e.g., labeled or labelable probes which can be used in, for

- 65 -

example, an *in situ* hybridization technique, to identify a specific tissue, e.g., brain tissue. This can be very useful in cases where a forensic pathologist is presented with a tissue of unknown origin. Panels of such CRSP probes can be used to identify tissue by species and/or by organ type.

5 In a similar fashion, these reagents, e.g., CRSP primers or probes can be used to screen tissue culture for contamination (i.e. screen for the presence of a mixture of different types of cells in a culture).

C. Predictive Medicine:

10 The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the present invention relates to diagnostic assays for determining CRSP protein and/or nucleic acid expression as well as CRSP activity, in 15 the context of a biological sample (e.g., blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant CRSP expression or activity, such as aberrant cell proliferation, differentiation, and/or survival resulting for example in a neurodegenerative disease or cancer. The invention also provides for prognostic (or 20 predictive) assays for determining whether an individual is at risk of developing a disorder associated with CRSP protein, nucleic acid expression or activity. For example, mutations in a CRSP gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with CRSP 25 protein, nucleic acid expression or activity.

Another aspect of the invention pertains to monitoring the influence of agents (e.g., drugs, compounds) on the expression or activity of CRSP in clinical trials.

These and other agents are described in further detail in the following sections.

30 1. Diagnostic Assays

An exemplary method for detecting the presence or absence of CRSP protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting CRSP protein or nucleic acid (e.g., mRNA, genomic DNA) that encodes CRSP 35 protein such that the presence of CRSP protein or nucleic acid is detected in the biological sample. A preferred agent for detecting CRSP mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to CRSP mRNA or genomic DNA.

The nucleic acid probe can be, for example, a full-length CRSP nucleic acid, such as the nucleic acid of SEQ ID NO: 1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the DNA insert of

5 the plasmid deposited with ATCC as Accession Number 98452, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to CRSP mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

10 A preferred agent for detecting CRSP protein is an antibody capable of binding to CRSP protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling

15 (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin. The term

20 "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect CRSP mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of CRSP mRNA include Northern hybridizations and *in situ*

25 hybridizations. *In vitro* techniques for detection of CRSP protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of CRSP genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of CRSP protein include introducing into a subject a labeled anti-CRSP antibody. For example, the

30 antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a

compound or agent capable of detecting CRSP protein, mRNA, or genomic DNA, such that the presence of CRSP protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of CRSP protein, mRNA or genomic DNA in the control sample with the presence of CRSP protein, mRNA or genomic DNA in the test 5 sample.

The invention also encompasses kits for detecting the presence of CRSP in a biological sample. For example, the kit can comprise a labeled compound or agent capable of detecting CRSP protein or mRNA in a biological sample; means for determining the amount of CRSP in the sample; and means for comparing the amount of 10 CRSP in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect CRSP protein or nucleic acid.

2. Prognostic Assays

15 The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant CRSP expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with CRSP protein, nucleic acid 20 expression or activity such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder as well as diseases, conditions or disorders characterized by abnormal cell survival, abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a differentiative or 25 proliferative disease (e.g., cancer). Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant CRSP expression or activity in which a test sample is obtained from a subject and CRSP protein or nucleic acid (e.g., mRNA, genomic DNA) is detected, wherein the presence of CRSP protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder 30 associated with aberrant CRSP expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (e.g., serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (e.g., an agonist, antagonist, 35 peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant CRSP expression or activity. For example, such methods can be used to determine whether a subject can be effectively

treated with an agent for a disorder, such as a proliferative disorder, a differentiative or developmental disorder, a hematopoietic disorder, as well disorders characterized by abnormal cell survival, an abnormal extracellular structure, or an abnormality in a defense mechanism. Alternatively, such methods can be used to determine whether a subject can be effectively treated with an agent for a differentiative or proliferative disease (e.g., cancer). Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant CRSP expression or activity in which a test sample is obtained and CRSP protein or nucleic acid expression or activity is detected (e.g., wherein the abundance of CRSP protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant CRSP expression or activity.)

The methods of the invention can also be used to detect genetic alterations in a CRSP gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by aberrant development, aberrant cellular differentiation, aberrant cellular proliferation or an aberrant hematopoietic response. In preferred embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a CRSP-protein, or the mis-expression of the CRSP gene. For example, such genetic alterations can be detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a CRSP gene; 2) an addition of one or more nucleotides to a CRSP gene; 3) a substitution of one or more nucleotides of a CRSP gene, 4) a chromosomal rearrangement of a CRSP gene; 5) an alteration in the level of a messenger RNA transcript of a CRSP gene, 6) aberrant modification of a CRSP gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a CRSP gene, 8) a non-wild type level of a CRSP-protein, 9) allelic loss of a CRSP gene, and 10) inappropriate post-translational modification of a CRSP-protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting alterations in a CRSP gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject.

In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, e.g., U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the CRSP-gene (see Abravaya *et al.* (1995)

Nucleic Acids Res. 23:675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (e.g., genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a CRSP gene under conditions such that

5 hybridization and amplification of the CRSP-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

10 Alternative amplification methods include: self sustained sequence replication (Guatelli, J.C. *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh, D.Y. *et al.*, 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. *et al.*, 1988, *Bio/Technology* 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified 15 molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

20 In an alternative embodiment, mutations in a CRSP gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel 25 electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

30 In other embodiments, genetic mutations in CRSP can be identified by hybridizing a sample and control nucleic acids, e.g., DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in CRSP can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by 35 making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays

complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the CRSP gene and detect mutations by comparing the sequence of the sample CRSP with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert ((1977) *PNAS* 74:560) or Sanger ((1977) *PNAS* 74:5463). It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques* 19:448), including sequencing by mass spectrometry (see, e.g., PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the CRSP gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type CRSP sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in CRSP cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a CRSP

sequence, e.g., a wild-type CRSP sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.

5 In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in CRSP genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids (Orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; and Hayashi (1992) 10 *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control CRSP nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay 15 may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment the movement of mutant or wild-type fragments in 20 polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of 25 high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective 30 primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different 35 mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention.

Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or 5 reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for 10 amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it 15 possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent 15 described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a CRSP gene.

Furthermore, any cell type or tissue in which CRSP is expressed may be utilized 20 in the prognostic assays described herein.

3. Monitoring of Effects During Clinical Trials

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or 25 activity of CRSP (*e.g.*, modulation of cellular signal transduction, regulation of gene transcription in a cell involved in development or differentiation, regulation of cellular proliferation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase CRSP gene expression, protein levels, or upregulate CRSP activity, 30 can be monitored in clinical trials of subjects exhibiting decreased CRSP gene expression, protein levels, or downregulated CRSP activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease CRSP gene expression, protein levels, or downregulate CRSP activity, can be monitored in clinical trials of subjects exhibiting increased CRSP gene expression, protein levels, or upregulated CRSP activity. In such clinical trials, the expression or activity of CRSP and, preferably, other genes that have been implicated in, for example, a proliferative 35 disorder can be used as a "read out" or markers of the phenotype of a particular cell.

For example, and not by way of limitation, genes, including CRSP, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule)

which modulates CRSP activity (e.g., identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on proliferative disorders, developmental or differentiative disorder, hematopoietic disorder as well disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular 5 structure, or an abnormality in a defense mechanism, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of CRSP and other genes implicated in the proliferative disorder, developmental or differentiative disorder, hematopoietic disorder as well as disorders characterized by abnormal cell differentiation and/or survival, an abnormal extracellular structure, or an abnormality in 10 a defense mechanism, respectively. The levels of gene expression (i.e., a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of CRSP or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the 15 physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (e.g., an agonist, 20 antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a CRSP protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples 25 from the subject; (iv) detecting the level of expression or activity of the CRSP protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the CRSP protein, mRNA, or genomic DNA in the pre-administration sample with the CRSP protein, mRNA, or genomic DNA in the post-administration sample or samples; and (vi) altering the administration of the agent to the 30 subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of CRSP to higher levels than detected, i.e., to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of CRSP to lower levels than detected, i.e. to decrease the effectiveness of the agent. According to such an 35 embodiment, CRSP expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

C. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant CRSP expression or activity. With regards to both prophylactic and 5 therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics.

"Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More specifically, the term refers the study of

10 how a patient's genes determine his or her response to a drug (e.g., a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the CRSP molecules of the present invention or CRSP modulators according to that individual's drug response genotype. Pharmacogenomics allows a 15 clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

20 1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant CRSP expression or activity, by administering to the subject an agent which modulates CRSP expression or at least one CRSP activity. Subjects at risk for a disease which is caused or contributed to by 25 aberrant CRSP expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the CRSP aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of CRSP aberrancy, for example, a CRSP agonist or CRSP antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the present invention are further discussed in the following subsections.

30 2. Therapeutic Methods

35 Another aspect of the invention pertains to methods of modulating CRSP expression or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of

CRSP protein activity associated with the cell. An agent that modulates CRSP protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring target molecule of a CRSP protein, a peptide, a CRSP peptidomimetic, or other small molecule. In one embodiment, the agent stimulates one or more CRSP protein activity. Examples of such stimulatory agents include active CRSP protein and a nucleic acid molecule encoding CRSP that has been introduced into the cell. In another embodiment, the agent inhibits one or more CRSP protein activity. Examples of such inhibitory agents include antisense CRSP nucleic acid molecules and anti-CRSP antibodies. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a CRSP protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) CRSP expression or activity. In another embodiment, the method involves administering a CRSP protein or nucleic acid molecule as therapy to compensate for reduced or aberrant CRSP expression or activity.

Stimulation of CRSP activity is desirable in situations in which CRSP is abnormally downregulated and/or in which increased CRSP activity is likely to have a beneficial effect. Likewise, inhibition of CRSP activity is desirable in situations in which CRSP is abnormally upregulated and/or in which decreased CRSP activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant development or cellular differentiation. Another example of such a situation is where the subject has a proliferative disease (e.g., cancer) or a disorder characterized by an aberrant hematopoietic response. Yet another example of such a situation is where it is desireable to achieve tissue regeneration in a subject (e.g., where a subject has undergone brain or spinal cord injury and it is desirable to regenerate neuronal tissue in a regulated manner.)

Accordingly, in one embodiment, the disease is a disease characterized by an abnormal cell proliferation, differentiation, and/or survival. For example, the disease can be a hyper- or hypoproliferative disease. The invention also provides methods for treating diseases characterized by an abnormal cell proliferation, differentiation, and/or survival in a subject, which are not characterized by an abnormal CRSP activity (e.g., CRSP-1 activity). In fact, since CRSP is likely to be capable of modulating the proliferative state of a cell (i.e., state of proliferation, differentiation, and/or survival of a

cell), CRSP can regulate disease wherein the abnormal proliferative state of a cell results from a defect other than an abnormal CRSP activity.

Hyperproliferative diseases can be treated with CRSP (e.g., CRSP-1) therapeutics include neoplastic and hyperplastic diseases, such as various forms of cancers and leukemias, and fibroproliferative disorders. Other hyperproliferative diseases that can be treated or prevented with the subject CRSP therapeutics (e.g. CRSP-1 therapeutics) include malignant conditions, premalignant conditions, and benign conditions. The condition to be treated or prevented can be a solid tumor, such as a tumor arising in an epithelial tissue. Accordingly, treatment of such a cancer could comprise administration to the subject of a CRSP therapeutic decreasing the interaction of CRSP with a CRSP receptor. Other cancers that can be treated or prevented with a CRSP protein include sarcomas and carcinomas, e.g., lung cancer, cancer of the colon, prostate, breast, ovary, esophagus, lung cancer, melanoma, seminoma, and squamous adenocarcinoma. Additional solid tumors within the scope of the invention include those that can be found in a medical textbook.

The condition to be treated or prevented can also be a soluble tumor, such as leukemia, either chronic or acute, including chronic or acute myelogenous leukemia, chronic or acute lymphocytic leukemia, promyelocytic leukemia, monocytic leukemia, myelomonocytic leukemia, and erythroleukemia. Yet other proliferative disorders that can be treated with a CRSP therapeutic of the invention include heavy chain disease, multiple myeloma, lymphoma, e.g., Hodgkin's lymphoma and non-Hodgkin's lymphoma, and Waldenstroem's macroglobulemia.

Diseases or conditions characterized by a solid or soluble tumor can be treated by administrating a CRSP therapeutic either locally or systemically, such that aberrant cell proliferation is inhibited or decreased. Methods for administering the compounds of the invention are further described below.

The invention also provides methods for preventing the formation and/or development of tumors. For example, the development of a tumor can be preceded by the presence of a specific lesion, such as a pre-neoplastic lesion, e.g., hyperplasia, metaplasia, and dysplasia, which can be detected, e.g., by cytologic methods. Such lesions can be found, e.g., in epithelial tissue. Thus, the invention provides a method for inhibiting progression of such a lesion into a neoplastic lesion, comprising administering to the subject having a preneoplastic lesion an amount of a CRSP-1 therapeutic sufficient to inhibit progression of the preneoplastic lesion into a neoplastic lesion.

The invention also provides for methods for treating or preventing diseases or conditions in which proliferation of cells is desired. For example, CRSP therapeutics can be used to stimulate tissue repair or wound healing, such as after surgery or to

stimulate tissue healing from burns. Other diseases in which proliferation of cells is desired are hypoproliferative diseases, i.e., diseases characterized by an abnormally low proliferation of certain cells.

In yet another embodiment, the invention provides a method for treating or preventing diseases or conditions characterized by aberrant cell differentiation.

Accordingly, the invention provides methods for stimulating cellular differentiation in conditions characterized by an inhibition of normal cell differentiation which may or may not be accompanied by excessive proliferation. Alternatively, CRSP therapeutics can be used to inhibit differentiation of specific cells.

10 In a preferred method, the aberrantly proliferating and/or differentiating cell is a cell present in the nervous system. A role for CRSP in the nervous system is suggested at least in part from the fact that human CRSP-1 is expressed in human fetal brain. Accordingly, the invention provides methods for treating diseases or conditions associated with a central or peripheral nervous system. For example, the invention

15 provides methods for treating lesions of the nervous system associated with an aberrant proliferation, differentiation or survival of any of the following cells: neurons, Schwann cells, glial cells, and other types of neural cells. Disorders of the nervous system include, but are not limited to: spinal cord injuries, brain injuries, lesions associated with surgery, ischemic lesions, malignant lesions, infectious lesions, degenerative

20 lesions (Parkinson's disease, Alzheimer's disease, Huntington's chorea, amyotrophic lateral sclerosis), demyelinating diseases (multiple sclerosis, human immunodeficiency associated myelopathy, transverse myelopathy, progressive multifocal leukoencephalopathy, pontine myelinolysis), motor neuron injuries, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and

25 juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis, and hereditary mototsensory neuropathy (Charcot-Marie-Tooth disease).

In another embodiment, the invention provides a method for enhancing the survival and/or stimulating proliferation and/or differentiation of cells and tissues *in vitro*. In a preferred embodiment, CRSP therapeutics are used to promote tissue regeneration and/or repair (e.g., to treat nerve injury). For example, tissues from a subject can be obtained and grown *in vitro* in the presence of a CRSP therapeutic, such that the tissue cells are stimulated to proliferate and/or differentiate. The tissue can then be readministered to the subject.

30 Among the approaches which may be used to ameliorate disease symptoms involving an aberrant CRSP activity and/or an abnormal cell proliferation, differentiation, and/or survival, are, for example, antisense, ribozyme, and triple helix

molecules described above. Examples of suitable compounds include the antagonists, agonists or homologues described in detail above.

Yet other CRSP therapeutics consist of a first peptide comprising a CRSP peptide capable of binding to a CRSP receptor, and a second peptide which is cytotoxic.

5 Such therapeutics can be used to specifically target and lyse cells expressing or overexpressing a receptor for CRSP.

3. Pharmacogenomics

The CRSP molecules of the present invention, as well as agents, or modulators 10 which have a stimulatory or inhibitory effect on CRSP activity (e.g., CRSP gene expression) as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (e.g, proliferative or developmental disorders) associated with aberrant CRSP activity. In conjunction with such treatment, pharmacogenomics (i.e., the study of the relationship between an 15 individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to 20 administer a CRSP molecule or CRSP modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a CRSP molecule or CRSP modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See e.g., Eichelbaum, M., *Clin Exp Pharmacol Physiol*, 1996, 23(10-11) :983-25 985 and Linder, M.W., *Clin Chem*, 1997, 43(2):254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic 30 defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug 35 response, known as "a genome-wide association", relies primarily on a high-resolution map of the human genome consisting of already known gene-related markers (e.g., a "biallelic" gene marker map which consists of 60,000-100,000 polymorphic or variable

sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a

5 high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-
10 associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

15 Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (e.g., a CRSP protein or CRSP receptor of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is
20 associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (e.g., N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation
25 as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic
30 and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed
35 metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (e.g., a CRSP molecule or CRSP modulator of the present invention) can give an indication whether gene pathways related to toxicity have been 5 turned on.

Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus 10 enhance therapeutic or prophylactic efficiency when treating a subject with a CRSP molecule or CRSP modulator, such as a modulator identified by one of the exemplary screening assays described herein.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent 15 applications cited throughout this application are hereby incorporated by reference.

EXAMPLES

Example 1: Isolation And Characterization of Human CRSP-1 cDNA

20 In this example, the isolation and characterization of the gene encoding human CRSP-1 (also referred to as "CRISPY-1" or "TANGO 59") is described.

Isolation of the human CRSP-1 cDNA

25 The invention is based at least in part on the discovery of a human gene encoding a secreted protein, referred to herein as Cysteine Rich Secreted Protein-1 (CRSP-1). A partial cDNA was isolated using a Signal Sequence Trap method. This methodology takes advantage of the fact that molecules such as CRSP have an amino terminal signal sequence which directs certain secreted and membrane-bound proteins through the cellular secretory apparatus.

30 Briefly, a randomly primed cDNA library using mRNA prepared from human fetal brain tissue (Clontech, Palo Alto CA) was made by using the Stratagene-ZAP-cDNA Synthesis™ kit, (catalog #20041). The cDNA was ligated into the mammalian expression vector pTrap adjacent to a cDNA encoding placental alkaline phosphatase lacking a secretory signal. The plasmids were transformed into *E. coli* and DNA was 35 prepared using the Wizard™ DNA purification kit (Promega). DNA was transfected into COS-7 cells with lipofectamine™ (Gibco-BRL). After 48 hours incubation the COS cell supernatants were assayed for alkaline phosphatase on a Wallac Micro-Beta scintillation

counter using the Phospha-Light™ kit (Tropix Inc. Catalog #BP300). The individual plasmid DNAs scoring positive in the COS cell Alkaline Phosphatase secretion assay were further analyzed by DNA sequencing using standard procedures.

Using a partial cDNA isolated by the above-described method, a full length 5 cDNA encoding human CRSP-1 was cloned. The nucleotide sequence encoding the full length human CRSP-1 protein is shown in Figure 1 and is set forth as SEQ ID NO: 1. The full length protein encoded by this nucleic acid is comprised of about 350 amino acids and has the amino acid sequence shown in Figure 1 and set forth as SEQ ID NO:2. The coding portion (open reading frame) of SEQ ID NO:1 is set forth as SEQ ID NO:3. 10 DNA for the clone jthKb075a10 was deposited with the ATCC as Accession No. 98634.

Analysis of Human CRSP-1

Determination of the hydrophobicity profile of human CRSP-1 having the amino acid sequence set forth in SEQ ID NO:2 indicated the presence of a hydrophobic region 15 from about amino acid 1 to about amino acid 23 of SEQ ID NO:2. Further analysis of the amino acid sequence SEQ ID NO:2 using signal peptide prediction programs predicted the presence of a signal peptide from about amino acid 1 to about amino acid 19, 21, or 23 of SEQ ID NO:2. Accordingly, the mature CRSP-1 protein includes about 327, 329, or 331 amino acids spanning from about amino acid 20, 22, or 24 to about 20 amino acid 350 of SEQ ID NO:2. The presence of the signal sequence, in addition to the fact that CRSP-1 has been identified using a Signal Sequence Trap system, indicates that CRSP-1 is a secreted protein. Furthermore, the prediction of such a signal peptide and signal peptide cleavage site can be made, for example, utilizing the computer algorithm SIGNALP (Henrik, et al. (1997) *Protein Engineering* 10:1-6).

25 Furthermore, when the cDNA encoding human CRSP-1 was expressed in bacterial cells with a C-terminal FLAG tag, the FLAG-tagged CRSP-1 protein product was found to be secreted into the cellular medium.

Examination of the cDNA sequence depicted in Figure 1 shows that human CRSP-1 is particularly rich in cysteine residues. As shown in Figure 1, CRSP-1 30 contains 20 cysteine residues located between amino acid 147 and amino acid 284 of SEQ ID NO: 2. These cysteine residues can possibly form 10 disulfide bridges.

A BLAST search (Altschul et al. (1990) *J. Mol. Biol.* 215:403) of the nucleotide and the amino acid sequences of CRSP-1 has revealed that CRSP-1 is significantly similar to a chicken cDNA encoding a protein of unknown function having GenBank 35 Accession No. D26311. This cDNA was isolated from a chicken lens cDNA library and was shown to be expressed in lens fibers and lens epithelium, but not in neural retina nor in liver cells. (Sawada et al. (1996) *Int. J. Dev. Biol.* 40:531). CRSP-1 and the chicken

protein have 56% amino acid sequence identity and 72% amino acid sequence similarity. The amino acid sequence similarity between the chicken protein and human CRSP-1 is particularly high in the cysteine-rich domain of CRSP-1 which is located between amino acids 147 and 284 of SEQ ID NO: 2. In particular, the 20 cysteine residues of CRSP-1 located in this region are present in the chicken protein (see Figure 3).

Two genes recently identified in a screen for suppressors of glioblastoma formation (Ligon et al., 1997 *Oncogene* 14 1075-1081) also show significant homology to hCRSP-1. These genes, RIG ("Regulated In Glioblastoma") and RIG-like 7-1 (GenBank Accession Nos. U32331 and AF034208, respectively), were identified in a differential screen for mRNAs regulated by the introduction of a normal copy of chromosome 10 into a glioblastoma cell line harboring a deletion in chromosome 10 that promotes tumorigenesis. A schematic diagram summarizing the relationship between the sequences of the CRSP-1 and the RIG genes presented as Figure 9. The indicated region of identity between CRSP-1, RIG and RIG-like 7-1 comprises a region of the 3' UTR of the human CRSP-1 mRNA. This cDNA was initially isolated in the described differential screen and subsequently used to isolate full length RIG and RIG-like 7-1 messages as depicted. RIG-like 7-1 is highly homologous (96.8% similar as determined using the Lipman-Pearson protein alignment program, Ktuple: 2; Gap Penalty: 4, Gap Length Penalty: 12; and 64.7% homologous using the Wilbur Lipman DNA alignment program, Ktuple: 3; Gap Penalty: 3; Window: 20) to CRSP-1 although the encoded protein lacks the N-terminal signal sequence and is not therefore predicted to be a secreted protein. The full length RIG mRNA only displays homology to CRSP-1 in the region of the initial cDNA. These data associate CRSP-1 with human glioblastoma and suggest that CRSP-1 may be important in the suppression of the tumorigenic phenotype. A role in glioblastoma is also consistent with the high level of CRSP-1 mRNA expression observed in human brain tissue. In addition, the co-localization of the CRSP-1, RIG and RIG-like genes to a region of chromosome 11 (11p15.1) implicated in the development of human malignant astrocytoma (Ligon et al. 1997 *Oncogene* 14 1075-1081) further indicates a role for these genes in tumorigenesis.

Human CRSP-1 protein has also some amino acid sequence similarity to metallothionein, particularly in the cysteine-rich domain.

Tissue Distribution of CRSP-1 mRNA

This Example describes the tissue distribution of CRSP-1 mRNA, as determined by Northern blot hybridization.

Northern blot hybridizations with the various RNA samples were performed under standard conditions and washed under stringent conditions, i.e., 0.2xSSC at 65°C.

In each sample, the probe hybridized to a single RNA of about 2.5 kb. The results of hybridization of the probe to various mRNA samples are described below.

Hybridization of a Clontech Fetal Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from fetal brain, lung, liver, and kidney 5 indicated the presence of high levels of CRSP-1 mRNA in fetal brain, lung, and slightly lower levels of CRSP-1 mRNA in fetal kidney. However, no significant level of CRSP-1 mRNA was found in fetal liver.

Hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) containing RNA from adult heart, brain, placenta, lung, liver, 10 skeletal muscle, kidney, and pancreas with a human CRSP-1 probe indicated the presence of high levels of CRSP-1 mRNA in heart, slightly lower levels in brain, and much lower levels in placenta and lung. Some CRSP-1 mRNA was also found in adult skeletal muscle. However, no significant levels of CRSP-1 mRNA was observed in adult liver, kidney, or pancreas. Interestingly, the chicken gene which is homologous to 15 CRSP-1 was not expressed at detectable levels in liver either (Sawada et al. (1996) Int. J. Dev. Biol. 40:531).

Further hybridization of a Clontech human Multiple Tissue Northern (MTN) blot (Clontech, LaJolla, CA) including RNA from adult spinal cord revealed high levels of expression of CRSP-1 in mRNA isolated from adult spinal cord.

20 In situ analysis further revealed the following expression patterns when tissue sections were hybridized with CRSP-1 probes. CRSP-1 mRNA was expressed in heart, lung, aorta, eye (retina and lens), and brain (cortex) in 14.5 day mouse embryos. CRSP-1 mRNA was expressed in heart, lung, eye (retina and lens), brain (cortical), and cartilage (foot, trachea, larynx, head, sternum, and spinal column) in 1.5 day post natal 25 mouse tissues. CRSP-1 mRNA was expressed in brain (cortex, hippocampus, brainstem), heart (atria only), eye (retina and lens outer layer) in adult mouse tissues.

Thus, CRSP-1 is expressed in a tissue specific manner, with the strongest expression observed in brain, heart, and spinal cord.

30 CRSP-1 protein expression data

Constructs

Expression constructs for two forms of CRSP-1 were prepared using the mammalian expression vector pMET-stop. Form-1 comprised a cDNA incorporating the complete 350aa CRSP-1 protein coding sequence (CRSP-1flag.long) and form-2 35 comprised the entire CRSP-1 protein coding sequence except for the final 18 amino acids (CRSP-1flag.short). A C-terminal sequence encoding the FLAG epitope (DYKDDDDK) was added to both CRSP-1 forms for ease of detection and purification.

CRSP-1flag cDNAs were generated by PCR from a full length CRSP-1 cDNA template and ligated into pMET-stop using EcoR1 and Sal1 restriction sites.

Trial transfection - small scale expression

5 Expression constructs for CRSP-1flag.long and CRSP-1flag.short were transfected into 293T cells using 10 µl of lipofectamine (GIBCO/BRL) and 2 µg of DNA per well of a 6-well plate of cells which were 70-80% confluent. After 5 hours at 37°C, cells were fed with 1ml of 20%FCS/DMEM. After incubation overnight at 37°C, cells were conditioned in 1ml OptiMEM for 48 hours at 37°C. Samples of supernatant
10 and cell pellets were solubilized in boiling SDS-PAGE gel buffer, run out on a 4-20% SDS-PAGE gel, transferred to a nylon membrane and probed with the anti-FLAG monoclonal antibody M2. Samples from both supernatant and pellet samples showed significant immunoreactivity within a molecular weight range of 40-65 kDa on autoradiographic film using a HRP conjugated secondary antibody and ECL detection
15 reagents. Thus, both forms of CRSP-1 tested are secreted from 293T cells thereby confirming experimentally that CRSP-1 is a secreted protein. It should be noted that the molecular weights of both forms of CRSP-1 tested are greater than predicted from the amino acid sequence, suggesting that the CRSP-1 proteins secreted by 293T cells may be glycosylated. This is consistent with the presence of four potential sites for N-linked
20 glycosylation in the CRSP-1 protein.

Large scale CRSP-1 protein production

For scale-up of CRSP-1flag.long protein expression, 30 x 150mM plates of 293T cells at 70-80% confluence were transfected with 27 µg DNA, 100 µl lipofectamine in
25 18ml OptiMEM for 5 hours at 37°C. 18 ml of 10%FCS/DMEM was added to each plate and incubated overnight at 37°C. 24 hours after the start of transfection, transfection supernatant was aspirated and 35 mls OptiMEM was added to each plate and the plates incubated at 37°C for 72 hours. Conditioned medium was harvested, spun at 4000 rpm for 30 min. at 4°C, and filtered through a 0.45 micron filter unit. 1100 ml was passed
30 over a 1.6 x 10 cm anti-FLAG M2 affinity column pre-equilibrated in PBS pH7.4 buffer at a flow rate of 2.0 ml per minute. After washing with 200 ml of PBS pH 7.4 buffer, bound material was eluted by a step of 200 mM Glycine pH 3.0 buffer and 0.5 ml fractions collected. Upon elution, a significant protein peak was detected by absorbance at 280nm. Samples corresponding to conditioned medium, flow through and eluted
35 fractions were analyzed by Coomassie blue and silver stained SDS-PAGE and by western blot analysis as described above. Significant immunoreactivity within a molecular weight range of 40-65 kDa was detected in conditioned medium and eluted

fractions but not in the flow through sample, indicating that the secreted CRSP-1flag.long protein bound to the affinity column specifically and was eluted efficiently by the described conditions. Coomassie blue staining of SDS-PAGE gels suggested that the predominant immunoreactive protein constituted >90% of the protein present in the 5 bound and eluted protein peak. Peak fractions of eluted protein were pooled and dialysed against Phosphate Buffered Saline resulting in a 4 ml volume of recombinant CRSP-1flag.long protein at a concentration of approximately 1mg/ml.

Example 2: Isolation And Characterization of Human CRSP-2, CRSP-3, and
10 **CRSP-4 cDNA**

To identify novel proteins related to CRSP-1, the human CRSP-1 amino acid sequence was used to search the private and dbEST databases using TBLASTN (WashUversion, 2.0, BLOSUM62 search matrix). From this search, four distinct partial 15 protein sequences were identified which displayed significant homology to human CRSP-1. These sequences were designated as partial sequences for the novel hCRSPs; hCRSP-2, h-CRSP-3, h-CRSP-4 and h-CRSP-like-n.

hCRSP-2 partial protein sequence was derived from a single dbEST clone with accession number AA565546. This clone was subsequently obtained from the IMAGE 20 collection and sequenced fully to define the entire hCRSP-2 sequence depicted in Figure 2.

hCRSP-3 partial protein sequence was derived from a jthKb075a10 clone with sequence identification code jthKb075a10. This clone was sequenced further and this sequence information combined with additional sequences from dbEST using the 25 assembly program Phrap (P. Green, U. Washington) to define the entire hCRSP-3 sequence depicted in Figure 3. DNA for the clone jthKb075a10 was deposited with the ATCC as Accession No. 98633. Northern blot analysis of various tissues using a probe specific for hCRSP-3 revealed that CRSP-3 mRNA expression was restricted to 30 placenta. CRSP-3 mRNA expression has not yet been demonstrated in other normal adult human tissues.

A murine and *Xenopus* homologue of CRSP-3 has recently been described called dkk-1 ("dickkopf", German, thickhead) which is proposed to encode a secreted signaling molecule (Nature, 391, 357-368). dkk-1 is described as a powerful inducer of head formation during early *Xenopus* development, its mechanism of action apparently 35 involving inhibition of the *wnt* family of secreted factors.

Given the homology between CRSP-3 and dkk-1, it is suggested that hCRSP-3 may display profound head-inducing activity during early human embryonic

development. Furthermore, the mechanisms of action of dkk-1 is thought to involve antagonism of members of the *wnt* family of secreted proteins. Overexpression of *wnt* proteins has been associated with certain malignancies. For example, activation of *wnt-1* expression by proviral insertion of MMTV causes breast cancer in the mouse (see 5 Cadigan and Nusse, Genes and Dev., 1997 11 3286-3305). Therefore, at least CRSP-3, and potentially other CRSP family members, may also play a role in the suppression of *wnt* signaling in cancer.

h-CRSP-4 partial protein sequence was derived from a single dbEST clone with accession number W55979. This clone was subsequently obtained from the IMAGE 10 collection and sequenced fully to define a hCRSP-4 sequence depicted in Figure 4. Northern blot analysis of various tissues using a probe specific for hCRSP-4 revealed that CRSP-4 mRNA was expressed in several adult human tissues. CRSP-4 mRNA expression was highest in heart, brain, placenta, lung, and skeletal muscle.

h-CRSP-like-n partial protein sequence was derived from a single dbEST clone 15 with accession number AA397836. This clone was subsequently obtained from the IMAGE collection and sequenced fully to define the entire hCRSP-like-n sequence depicted in Figure 5.

Structure of the CRSP Family proteins

20 An alignment of the amino acid sequences of human CRSP-1, human CRSP-2, human CRSP-3, and human CRSP-4 is shown in Figure 6. Amino acid residues which are conserved between human CRSP family members are boxed. The 20 conserved cysteine residues are indicated by an asterix. The cysteine-rich domains are indicated as CRD-1 and CRD-2. The domain structure of the full length CRSP proteins of the 25 present invention are depicted in Figure 7. The amino acid and nucleotide homology between CRSP family members is as follows in Tables 1 and 2.

Table 1:

	CRSP-1	CRSP-2	CRSP-3	CRSP-4	mdkk-1	xdkk-1	CLFEST
CRSP-1	100	16.0	18.6	15.1	18.5	16.5	53.0
CRSP-2		100	33.7	35.2	32.6	33.7	16.2
CRSP-3			100	33.1	80.2	53.5	17.4
CRSP-4				100	30.5	33.7	12.5

Amino acid percent homologies were determined using the ALIGN program, (version 2.0) (GCG software package) using a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4.

5 Table 2:

	CRSP-1	CRSP-2	CRSP-3	CRSP-4	mdkk-1	xdkk-1	CLFEST
CRSP-1	100	30.0	37.2	34.7	31.5	45.4	58.8
CRSP-2		100	43.0	35.9	38.8	38.4	36.7
CRSP-3			100	59.3	66.4	53.7	32.1
CRSP-4				100	38.8	38.4	36.7

Nucleotide percent homologies were determined using the using the Wilbur Lipman DNA alignment program, Ktuple: 3; Gap Penalty: 3; Window: 20.

10

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

15

- 88 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT:

- (A) NAME: MILLENNIUM BIOTHERAPEUTICS, INC.
- (B) STREET: 620 MEMORIAL DRIVE
- (C) CITY: CAMBRIDGE
- (D) STATE: MASSACHUSETTS
- (E) COUNTRY: US
- (F) POSTAL CODE: 02319
- (G) TELEPHONE:
- (H) TELEFAX:

15

(ii) TITLE OF INVENTION: NOVEL CRSP PROTEIN AND NUCLEIC ACID
MOLECULES AND USES THEREFOR

20

(iii) NUMBER OF SEQUENCES: 18

25

(iv) CORRESPONDENCE ADDRESS:

- (A) ADDRESSEE: LAHIVE & COCKFIELD, LLP
- (B) STREET: 28 STATE STREET
- (C) CITY: BOSTON
- (D) STATE: MASSACHUSETTS
- (E) COUNTRY: US
- (F) ZIP: 02109

30

(v) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

35

(vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: PCT/US98/
- (B) FILING DATE: 16 APRIL 1998
- (C) CLASSIFICATION:

40

(vii) PRIOR APPLICATION DATA:

- (A) APPLICATION NUMBER: US 08/843,704
- (B) FILING DATE: 16 APRIL 1997
- (C) APPLICATION NUMBER: US 08/842,898
- (D) FILING DATE: 17 APRIL 1997
- (E) APPLICATION NUMBER: 60/071,589
- (F) FILING DATE: 15 JANUARY 1998
- (G) APPLICATION NUMBER: 09/009,802
- (H) FILING DATE: 20 JANUARY 1998

45

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: MANDRAGOURAS, AMY E.
- (B) REGISTRATION NUMBER: 36,207
- (C) REFERENCE/DOCKET NUMBER: MEI-008CPPC

- 89 -

(ix) TELECOMMUNICATION INFORMATION:
 (A) TELEPHONE: (617) 227-7400
 (B) TELEFAX: (617) 742-4214

5

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2479 base pairs
 10 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 38..1087

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGCACGAGGG	GGCGGCGGCT	GCGGGCGCAG	AGCGGAG	ATG	CAG	CGG	CTT	GGG	GCC	55							
Met Gln Arg Leu Gly Ala																	
				1					5								
25	ACC	CTG	CTG	TGC	CTG	CTG	GCG	GCG	GTC	CCC	ACG	GCC	CCC	GCG	103		
	Thr	Leu	Leu	Cys	Leu	Leu	Leu	Ala	Ala	Ala	Val	Pro	Thr	Ala	Pro	Ala	
				10			15				20						
30	CCC	GCT	CCG	ACG	GCG	ACC	TCG	GCT	CCA	GTC	AAG	CCC	GGC	CCG	GCT	CTC	151
	Pro	Ala	Pro	Thr	Ala	Thr	Ser	Ala	Pro	Val	Lys	Pro	Gly	Pro	Ala	Leu	
				25			30			35							
35	AGC	TAC	CCG	CAG	GAG	GAG	GCC	ACC	CTC	AAT	GAG	ATG	TTC	CGC	GAG	GTT	199
	Ser	Tyr	Pro	Gln	Glu	Ala	Thr	Leu	Asn	Glu	Met	Phe	Arg	Glu	Val		
				40			45			50							
40	GAG	GAA	CTG	ATG	GAG	GAC	ACG	CAG	CAC	AAA	TTG	CGC	AGC	GCG	GTG	GAA	247
	Glu	Glu	Leu	Met	Glu	Asp	Thr	Gln	His	Lys	Leu	Arg	Ser	Ala	Val	Glu	
				55			60			65		70					
45	GAG	ATG	GAG	GCA	GAA	GAA	GCT	GCT	GCT	AAA	GCA	TCA	TCA	GAA	GTG	AAC	295
	Glu	Met	Glu	Ala	Glu	Ala	Ala	Ala	Ala	Lys	Ala	Ser	Ser	Glu	Val	Asn	
				75			80			85							
50	CTG	GCA	AAC	TTA	CCT	CCC	AGC	TAT	CAC	AAT	GAG	ACC	AAC	ACA	GAC	ACG	343
	Leu	Ala	Asn	Leu	Pro	Pro	Ser	Tyr	His	Asn	Glu	Thr	Asn	Thr	Asp	Thr	
				90			95			100							
	AAC	GTT	GGA	AAT	AAT	ACC	ATC	CAT	GTG	CAC	CGA	GAA	ATT	CAC	AAG	ATA	391
	Asn	Val	Gly	Asn	Asn	Thr	Ile	His	Val	His	Arg	Glu	Ile	His	Lys	Ile	
				105			110			115							

- 90 -

	ACC AAC AAC CAG ACT GGA CAA ATG GTC TTT TCA GAG ACA GTT ATC ACA	439
	Thr Asn Asn Gln Thr Gly Gln Met Val Phe Ser Glu Thr Val Ile Thr	
	120 125 130	
5	TCT GTG GGA GAC GAA GAA GGC AGA AGG AGC CAC GAG TGC ATC ATC GAC	487
	Ser Val Gly Asp Glu Glu Gly Arg Arg Ser His Glu Cys Ile Ile Asp	
	135 140 145 150	
10	GAG GAC TGT GGG CCC AGC ATG TAC TGC CAG TTT GCC AGC TTC CAG TAC	535
	Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln Phe Ala Ser Phe Gln Tyr	
	155 160 165	
15	ACC TGC CAG CCA TGC CGG GGC CAG AGG ATG CTC TGC ACC CGG GAC AGT	583
	Thr Cys Gln Pro Cys Arg Gly Gln Arg Met Leu Cys Thr Arg Asp Ser	
	170 175 180	
	GAG TGC TGT GGA GAC CAG CTG TGT GTC TGG GGT CAC TGC ACC AAA ATG	631
	Glu Cys Cys Gly Asp Gln Leu Cys Val Trp Gly His Cys Thr Lys Met	
20	185 190 195	
	GCC ACC AGG GGC AGC AAT GGG ACC ATC TGT GAC AAC CAG AGG GAC TGC	679
	Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys Asp Asn Gln Arg Asp Cys	
	200 205 210	
25	CAG CCG GGG CTG TGC TGT GCC TTC CAG AGA GGC CTG CTG TTC CCT GTG	727
	Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg Gly Leu Leu Phe Pro Val	
	215 220 225 230	
30	TGC ACA CCC CTG CCC GTG GAG GGC GAG CTT TGC CAT GAC CCC GCC AGC	775
	Cys Thr Pro Leu Pro Val Glu Gly Glu Leu Cys His Asp Pro Ala Ser	
	235 240 245	
35	CGG CTT CTG GAC CTC ATC ACC TGG GAG CTA GAG CCT GAT GGA GCC TTG	823
	Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu Glu Pro Asp Gly Ala Leu	
	250 255 260	
	GAC CGA TGC CCT TGT GCC AGT GGC CTC CTC TGC CAG CCC CAC AGC CAC	871
	Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu Cys Gln Pro His Ser His	
	265 270 275	
40	AGC CTG GTG TAT GTG TGC AAG CCG ACC TTC GTG GGG AGC CGT GAC CAA	919
	Ser Leu Val Tyr Val Cys Lys Pro Thr Phe Val Gly Ser Arg Asp Gln	
	280 285 290	
45	GAT GGG GAG ATC CTG CTG CCC AGA GAG GTC CCC GAT GAG TAT GAA GTT	967
	Asp Gly Glu Ile Leu Leu Pro Arg Glu Val Pro Asp Glu Tyr Glu Val	
	295 300 305 310	
50	GGC AGC TTC ATG GAG GAG GTG CGC CAG GAG CTG GAG GAC CTG GAG AGG	1015
	Gly Ser Phe Met Glu Glu Val Arg Gln Glu Leu Glu Asp Leu Glu Arg	
	315 320 325	
55	AGC CTG ACT GAA GAG ATG GCG CTG AGG GAG CCT GCG GCT GCC GCC GCT	1063
	Ser Leu Thr Glu Glu Met Ala Leu Arg Glu Pro Ala Ala Ala Ala	
	330 335 340	

	GCA CTG CTG GGA AGG GAA GAG ATT TAGATCTGGA CCAGGCTGTG GGTAGATGTG	1117
	Ala Leu Leu Gly Arg Glu Glu Ile	
	345 350	
5	CAATAGAAAT AGCTAATTAA TTTCCCCANG TGTGTGCTTT AAGCGTGGGC TGACCAGGCT	1177
	TCTTCCTTACA TCTTCTTCCC AGTAAGTTTC CCCTCTGGCT TGACAGCATG AGGTGTTGTG	1237
10	CATTTGTTCA GCTCCCCAG GCTGTTCTCC AGGCTTCACA GTCTGGTGT TGGGAGAGTC	1297
	AGGCAGGGTT AAACTGCAGG AGCAGTTGC CACCCCTGTC CAGATTATTG GCTGCTTGC	1357
15	CTCTACCAGT TGGCAGACAG CCGTTGTTTC TACATGGCTT TGATAATTGT TTGAGGGAG	1417
	GAGATGGAAA CAATGTGGAG TCTCCCTCTG ATTGGTTTG GGGAAATGTG GAGAAGAGTG	1477
	CCCTGCTTTG CAAACATCAA CCTGGCAAAA ATGCAACAAA TGAATTTCC ACGCAGTTCT	1537
20	TTCCATGGGC ATAGGTAAGC TGTGCCTTCA GCTGTTGCAG ATGAAATGTT CTGTTCACCC	1597
	TGCATTACAT GTGTTTATTTC ATCCAGCAGT GTTGCTCAGC TCCTACCTCT GTGCCAGGGC	1657
25	AGCATTTCAT TATCCAAGAT CAATTCCCTC TCTCAGCACA GCCTGGGGAG GGGGTCATTG	1717
	TTCTCCTCGT CCATCAGGGA TTTCAGAGGC TCAGAGACTG CAAGCTGCTT GCCCAAGTCA	1777
	CACAGCTAGT GAAGACCAGA GCAGTTTCAT CTGGTTGTGA CTCTAAGCTC AGTGCTCTCT	1837
30	CCACTACCCC ACACCAGCCT TGGTGCCACC AAAAGTGCTC CCCAAAAGGA AGGAGAATGG	1897
	GATTTTCCTT TTGAGGCATG CACATCTGGA ATTAAGGTCA AACTAATTCT CACATCCCTC	1957
35	TAAAAGTAAA CTACTGTTAG GAACAGCAGT GTTCTCACAG TGTGGGGCAG CCGTCCTTCT	2017
	AATGAAGACA ATGATATTGA CACTGTCCCT CTTTGGCAGT TGCATTAGTA ACTTTGAAAG	2077
	GTATATGACT GAGCGTAGCA TACAGGTTAA CCTGCAGAAA CAGTACTTAG GTAATTGTAG	2137
40	GGCGAGGATT ATAAATGAAA TTTGCAAAAT CACTTAGCAG CAACTGAAGA CAATTATCAA	2197
	CCACGTGGAG AAAATCAAAC CGAGCAGGGC TGTGTGAAAC ATGGTTGTAA TATGCGACTG	2257
	CGAACACTGA ACTCTACGCC ACTCCACAAA TGATGTTTTC AGGTGTCATG GACTGTTGCC	2317
45	ACCATGTATT CATCCAGAGT TCTTAAAGTT TAAAGTTGCA CATGATTGTA TAAGCATGCT	2377
	TTCTTGTAGT TTTAAATTAT GTATAAACAT AAGTTGCATT TAGAAATCAA GCATAAATCA	2437
50	CTTCAACTGC TAAAAAAAAA AAAAAAAA AAAAAAAA AA	2479

- 92 -

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

5 (A) LENGTH: 350 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Leu Ala Ala Ala
1 5 10 15

15 Val Pro Thr Ala Pro Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val
20 25 30

Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn
35 40 45

20 Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
50 55 60

25 Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys
65 70 75 80

Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn
85 90 95

30 Glu Thr Asn Thr Asp Thr Asn Val Gly Asn Asn Thr Ile His Val His
100 105 110

Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe
115 120 125

35 Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Arg Arg Ser
130 135 140

40 His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln
145 150 155 160

Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met
165 170 175

45 Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp
180 185 190

Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys
195 200 205

50 Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
210 215 220

Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
225 230 235 240

- 93 -

Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu
 245 250 255

5 Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
 260 265 270

Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe
 275 280 285

10 Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val
 290 295 300

15 Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu
 305 310 315 320

Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu
 325 330 335

20 Pro Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile
 340 345 350

(2) INFORMATION FOR SEQ ID NO:3:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1050 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1050

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

40 ATG CAG CGG CTT GGG GCC ACC CTG CTG TGC CTG CTG CTG GCG GCG GCG
 48 Met Gln Arg Leu Gly Ala Thr Leu Leu Cys Leu Leu Ala Ala Ala
 1 5 10 15

45 GTC CCC ACG GCC CCC GCG CCC GCT CCG ACG GCG ACC TCG GCT CCA GTC
 96 Val Pro Thr Ala Pro Ala Pro Thr Ala Thr Ser Ala Pro Val
 20 25 30

50 AAG CCC GGC CCG GCT CTC AGC TAC CCG CAG GAG GAG GCC ACC CTC AAT
 144 Lys Pro Gly Pro Ala Leu Ser Tyr Pro Gln Glu Glu Ala Thr Leu Asn
 35 40 45

55 GAG ATG TTC CGC GAG GTT GAG GAA CTG ATG GAG GAC ACG CAG CAC AAA
 192 Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys
 50 55 60

- 94 -

TTG CGC AGC GCG GTG GAA GAG ATG GAG GCA GAA GAA GCT GCT GCT AAA	240
Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys	
65 70 75 80	
5 GCA TCA TCA GAA GTG AAC CTG GCA AAC TTA CCT CCC AGC TAT CAC AAT	288
Ala Ser Ser Glu Val Asn Leu Ala Asn Leu Pro Pro Ser Tyr His Asn	
85 90 95	
10 GAG ACC AAC ACA GAC ACG AAC GTT GGA AAT AAT ACC ATC CAT GTG CAC	336
Glu Thr Asn Thr Asp Thr Asn Val Gly Asn Asn Thr Ile His Val His	
100 105 110	
15 CGA GAA ATT CAC AAG ATA ACC AAC CAG ACT GGA CAA ATG GTC TTT	384
Arg Glu Ile His Lys Ile Thr Asn Asn Gln Thr Gly Gln Met Val Phe	
115 120 125	
20 TCA GAG ACA GTT ATC ACA TCT GTG GGA GAC GAA GAA GGC AGA AGG AGC	432
Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Gly Arg Arg Ser	
130 135 140	
25 CAC GAG TGC ATC ATC GAC GAG GAC TGT GGG CCC AGC ATG TAC TGC CAG	480
His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Ser Met Tyr Cys Gln	
145 150 155 160	
30 TTT GCC AGC TTC CAG TAC ACC TGC CAG CCA TGC CGG GGC CAG AGG ATG	528
Phe Ala Ser Phe Gln Tyr Thr Cys Gln Pro Cys Arg Gly Gln Arg Met	
165 170 175	
35 CTC TGC ACC CGG GAC AGT GAG TGC TGT GGA GAC CAG CTG TGT GTC TGG	576
Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Val Trp	
180 185 190	
40 GGT CAC TGC ACC AAA ATG GCC ACC AGG GGC AGC AAT GGG ACC ATC TGT	624
Gly His Cys Thr Lys Met Ala Thr Arg Gly Ser Asn Gly Thr Ile Cys	
195 200 205	
45 GAC AAC CAG AGG GAC TGC CAG CCG GGG CTG TGC TGT GCC TTC CAG AGA	672
Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg	
210 215 220	
50 GGC CTG CTG TTC CCT GTG TGC ACA CCC CTG CCC GTG GAG GGC GAG CTT	720
Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu	
225 230 235 240	
55 TGC CAT GAC CCC GCC AGC CGG CTT CTG GAC CTC ATC ACC TGG GAG CTA	768
Cys His Asp Pro Ala Ser Arg Leu Leu Asp Leu Ile Thr Trp Glu Leu	
245 250 255	
50 GAG CCT GAT GGA GCC TTG GAC CGA TGC CCT TGT GCC AGT GGC CTC CTC	816
Glu Pro Asp Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu	
260 265 270	
55 TGC CAG CCC CAC AGC CAC AGC CTG GTG TAT GTG TGC AAG CCG ACC TTC	864
Cys Gln Pro His Ser His Ser Leu Val Tyr Val Cys Lys Pro Thr Phe	
275 280 285	

- 95 -

GTG GGG AGC CGT GAC CAA GAT GGG GAG ATC CTG CTG CCC AGA GAG GTC	912																												
Val Gly Ser Arg Asp Gln Asp Gly Glu Ile Leu Leu Pro Arg Glu Val																													
290	295		300	5 CCC GAT GAG TAT GAA GTT GGC AGC TTC ATG GAG GAG GTG CGC CAG GAG	960	Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu		305	310		315		320	10 CTG GAG GAC CTG GAG AGG AGC CTG ACT GAA GAG ATG GCG CTG AGG GAG	1008	Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu		325	330		335	15 CCT GCG GCT GCC GCC GCT GCA CTG CTG GGA AGG GAA GAG ATT	1050	Pro Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile		340	345		350
	300																												
5 CCC GAT GAG TAT GAA GTT GGC AGC TTC ATG GAG GAG GTG CGC CAG GAG	960																												
Pro Asp Glu Tyr Glu Val Gly Ser Phe Met Glu Glu Val Arg Gln Glu																													
305	310		315		320	10 CTG GAG GAC CTG GAG AGG AGC CTG ACT GAA GAG ATG GCG CTG AGG GAG	1008	Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu		325	330		335	15 CCT GCG GCT GCC GCC GCT GCA CTG CTG GGA AGG GAA GAG ATT	1050	Pro Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile		340	345		350								
	315																												
	320																												
10 CTG GAG GAC CTG GAG AGG AGC CTG ACT GAA GAG ATG GCG CTG AGG GAG	1008																												
Leu Glu Asp Leu Glu Arg Ser Leu Thr Glu Glu Met Ala Leu Arg Glu																													
325	330		335	15 CCT GCG GCT GCC GCC GCT GCA CTG CTG GGA AGG GAA GAG ATT	1050	Pro Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile		340	345		350																		
	335																												
15 CCT GCG GCT GCC GCC GCT GCA CTG CTG GGA AGG GAA GAG ATT	1050																												
Pro Ala Ala Ala Ala Ala Leu Leu Gly Arg Glu Glu Ile																													
340	345		350																										
	350																												

20 (2) INFORMATION FOR SEQ ID NO:4:

25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 848 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: cDNA
 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 125..796

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

35 GAATTCTGGCA CGAGAGACCA CGTGCTGAGC TGCCAGCTTA GTGGAAGCTC TGCTCTGGT	60
GGAGAGCAGC CTCGCTTGG TGACGCACAG TGCTGGGACC CTCCAGGAGC CCCGGGATTG	120
40 AAGG ATG GTG GCG GCC GTC CTG CTG GGG CTG AGC TGG CTC TGC TCT CCC	169
Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro	
1 5 10 15	

45 CTG GGA GCT CTG GTC CTG GAC TTC AAC AAC ATC AGG AGC TCT GCT GAC	217
Leu Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp	
20 25 30	

50 CTG CAT GGG GCC CGG AAG GGC TCA CAG TGC CTG TCT GAC ACG GAC TGC	265
Leu His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys	
35 40 45	

50 AAT ACC AGA AAG TTC TGC CTC CAG CCC CGC GAT GAG AAG CCG TTC TGT	313
Asn Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys	
55 60	

- 96 -

	GCT ACA TGT CGT GGG TTG CGG AGG AGG TGC CAG CGA GAT GCC ATG TGC	361
	Ala Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys	
	65 70 75	
5	TGC CCT GGG ACA CTC TGT GTG AAC GAT GTT TGT ACT ACG ATG GAA GAT	409
	Cys Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp	
	80 85 90 95	
10	GCA ACC CCA ATA TTA GAA AGG CAG CTT GAT GAG CAA GAT GGC ACA CAT	457
	Ala Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His	
	100 105 110	
15	GCA GAA GGA ACA ACT GGG CAC CCA GTC CAG GAA AAC CAA CCC AAA AGG	505
	Ala Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg	
	115 120 125	
20	AAG CCA AGT ATT AAG AAA TCA CAA GGC AGG AAG GGA CAA GAG GGA GAA	553
	Lys Pro Ser Ile Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu	
	130 135 140	
25	AGT TGT CTG AGA ACT TTT GAC TGT GGC CCT GGA CTT TGC TGT GCT CGT	601
	Ser Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg	
	145 150 155	
30	CAT TTT TGG ACG AAA ATT TGT AAG CCA GTC CTT TTG GAG GGA CAG GTC	649
	His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val	
	160 165 170 175	
35	TGC TCC AGA AGA GGG CAT AAA GAC ACT GCT CAA GCT CCA GAA ATC TTC	697
	Cys Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe	
	180 185 190	
40	CAG CGT TGC GAC TGT GGC CCT GGA CTA CTG TGT CGA AGC CAA TTG ACC	745
	Gln Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr	
	195 200 205	
45	AGC AAT CGG CAG CAT GCT CGA TTA AGA GTA TGC CAA AAA ATA GAA AAG	793
	Ser Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys	
	210 215 220	
50	CTA TAAATATTTC AAAATAAAGA AGAATCCACA TTGCAAAAAA AAAAAAAA AA	848
	Leu	

45 (2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 224 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

55

- 97 -

Met Val Ala Ala Val Leu Leu Gly Leu Ser Trp Leu Cys Ser Pro Leu			
1	5	10	15
Gly Ala Leu Val Leu Asp Phe Asn Asn Ile Arg Ser Ser Ala Asp Leu			
5	20	25	30
His Gly Ala Arg Lys Gly Ser Gln Cys Leu Ser Asp Thr Asp Cys Asn			
	35	40	45
10	Thr Arg Lys Phe Cys Leu Gln Pro Arg Asp Glu Lys Pro Phe Cys Ala		
	50	55	60
Thr Cys Arg Gly Leu Arg Arg Arg Cys Gln Arg Asp Ala Met Cys Cys			
	65	70	75
15	Pro Gly Thr Leu Cys Val Asn Asp Val Cys Thr Thr Met Glu Asp Ala		
	85	90	95
20	Thr Pro Ile Leu Glu Arg Gln Leu Asp Glu Gln Asp Gly Thr His Ala		
	100	105	110
Glu Gly Thr Thr Gly His Pro Val Gln Glu Asn Gln Pro Lys Arg Lys			
	115	120	125
25	Pro Ser Ile Lys Lys Ser Gln Gly Arg Lys Gly Gln Glu Gly Glu Ser		
	130	135	140
Cys Leu Arg Thr Phe Asp Cys Gly Pro Gly Leu Cys Cys Ala Arg His			
	145	150	155
30	Phe Trp Thr Lys Ile Cys Lys Pro Val Leu Leu Glu Gly Gln Val Cys		
	165	170	175
Ser Arg Arg Gly His Lys Asp Thr Ala Gln Ala Pro Glu Ile Phe Gln			
35	180	185	190
Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser			
	195	200	205
40	Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu		
	210	215	220

(2) INFORMATION FOR SEQ ID NO:6:

45 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 672 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

50 (ii) MOLECULE TYPE: cDNA

- 98 -

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..672

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATG	GTG	GCG	GCC	GTC	CTG	CTG	GGG	CTG	AGC	TGG	CTC	TGC	TCT	CCC	CTG	48	
Met	Val	Ala	Ala	Val	Leu	Leu	Gly	Leu	Ser	Trp	Leu	Cys	Ser	Pro	Leu		
1	5				10					15							
10	GGA	GCT	CTG	GTC	CTG	GAC	TTC	AAC	AAC	ATC	AGG	AGC	TCT	GCT	GAC	CTG	96
	Gly	Ala	Leu	Val	Leu	Asp	Phe	Asn	Asn	Ile	Arg	Ser	Ser	Ala	Asp	Leu	
	20				25						30						
15	CAT	GGG	GCC	CGG	AAG	GGC	TCA	CAG	TGC	CTG	TCT	GAC	ACG	GAC	TGC	AAT	144
	His	Gly	Ala	Arg	Lys	Gly	Ser	Gln	Cys	Leu	Ser	Asp	Thr	Asp	Cys	Asn	
	35				40						45						
20	ACC	AGA	AAG	TTC	TGC	CTC	CAG	CCC	CGC	GAT	GAG	AAG	CCG	TTC	TGT	GCT	192
	Thr	Arg	Lys	Phe	Cys	Leu	Gln	Pro	Arg	Asp	Glu	Lys	Pro	Phe	Cys	Ala	
	50				55						60						
25	ACA	TGT	CGT	GGG	TTG	CGG	AGG	AGG	TGC	CAG	CGA	GAT	GCC	ATG	TGC	TGC	240
	Thr	Cys	Arg	Gly	Leu	Arg	Arg	Arg	Cys	Gln	Arg	Asp	Ala	Met	Cys	Cys	
	65				70					75				80			
30	CCT	GGG	ACA	CTC	TGT	GTG	AAC	GAT	GTT	TGT	ACT	ACG	ATG	GAA	GAT	GCA	288
	Pro	Gly	Thr	Leu	Cys	Val	Asn	Asp	Val	Cys	Thr	Thr	Met	Glu	Asp	Ala	
	85				90					95							
35	ACC	CCA	ATA	TTA	GAA	AGG	CAG	CTT	GAT	GAG	CAA	GAT	GGC	ACA	CAT	GCA	336
	Thr	Pro	Ile	Leu	Glu	Arg	Gln	Leu	Asp	Glu	Gln	Asp	Gly	Thr	His	Ala	
	100				105						110						
40	GAA	GGA	ACA	ACT	GGG	CAC	CCA	GTC	CAG	GAA	AAC	CAA	CCC	AAA	AGG	AAG	384
	Glu	Gly	Thr	Thr	Gly	His	Pro	Val	Gln	Glu	Asn	Gln	Pro	Lys	Arg	Lys	
	115				120						125						
45	CCA	AGT	ATT	AAG	AAA	TCA	CAA	GGC	AGG	AAG	GGA	CAA	GAG	GGA	GAA	AGT	432
	Pro	Ser	Ile	Lys	Lys	Ser	Gln	Gly	Arg	Lys	Gly	Gln	Glu	Gly	Glu	Ser	
	130				135						140						
50	TGT	CTG	AGA	ACT	TTT	GAC	TGT	GGC	CCT	GGA	CTT	TGC	TGT	GCT	CGT	CAT	480
	Cys	Leu	Arg	Thr	Phe	Cys	Gly	Pro	Gly	Leu	Cys	Cys	Ala	Arg	His		
	145				150					155			160				
	165				170					175							
	TTT	TGG	ACG	AAA	ATT	TGT	AAG	CCA	GTC	CTT	TTG	GAG	GGA	CAG	GTC	TGC	528
	Phe	Trp	Thr	Lys	Ile	Cys	Lys	Pro	Val	Leu	Leu	Glu	Gly	Gln	Val	Cys	
	165				170					175							
	TCC	AGA	AGA	GGG	CAT	AAA	GAC	ACT	GCT	CAA	GCT	CCA	GAA	ATC	TTC	CAG	576
	Ser	Arg	Arg	Gly	His	Lys	Asp	Thr	Ala	Gln	Ala	Pro	Glu	Ile	Phe	Gln	
	180				185					185			190				

- 99 -

CGT TGC GAC TGT GGC CCT GGA CTA CTG TGT CGA AGC CAA TTG ACC AGC 624
 Arg Cys Asp Cys Gly Pro Gly Leu Leu Cys Arg Ser Gln Leu Thr Ser
 195 200 205

5 AAT CGG CAG CAT GCT CGA TTA AGA GTA TGC CAA AAA ATA GAA AAG CTA 672
 Asn Arg Gln His Ala Arg Leu Arg Val Cys Gln Lys Ile Glu Lys Leu
 210 215 220

10 (2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1529 base pairs
 (B) TYPE: nucleic acid
 15 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 93..890

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

25 CCGGACGCGT GGGCGGCACG GTTTCGTGGG GACCCAGGCT TGCAAAGTGA CGGTCATT 60

CTCTTTCTTT CTCCCTCTTG AGTCCTTCTG AG ATG ATG GCT CTG GGC GCA GCG 113
 Met Met Ala Leu Gly Ala Ala
 30 1 5

GGA GCT ACC CGG GTC TTT GTC GCG ATG GTA GCG GCG GCT CTC GGC GGC 161
 Gly Ala Thr Arg Val Phe Val Ala Met Val Ala Ala Ala Leu Gly Gly
 10 15 20

35 CAC CCT CTG CTG GGA GTG AGC GCC ACC TTG AAC TCG GTT CTC AAT TCC 209
 His Pro Leu Leu Gly Val Ser Ala Thr Leu Asn Ser Val Leu Asn Ser
 25 30 35

40 AAC GCT ATC AAG AAC CTG CCC CCA CCG CTG GGC GGC GCT GCG GGG CAC 257
 Asn Ala Ile Lys Asn Leu Pro Pro Leu Gly Gly Ala Ala Gly His
 40 45 50 55

45 CCA GGC TCT GCA GTC AGC GCC GCG CCG GGA ATC CTG TAC CCG GGC GGG 305
 Pro Gly Ser Ala Val Ser Ala Ala Pro Gly Ile Leu Tyr Pro Gly Gly
 60 65 70

50 AAT AAG TAC CAG ACC ATT GAC AAC TAC CAG CCG TAC CCG TGC GCA GAG 353
 Asn Lys Tyr Gln Thr Ile Asp Asn Tyr Gln Pro Tyr Pro Cys Ala Glu
 75 80 85

GAC GAG GAG TGC GGC ACT GAT GAG TAC TGC GCT AGT CCC ACC CGC GGA 401
 Asp Glu Glu Cys Gly Thr Asp Glu Tyr Cys Ala Ser Pro Thr Arg Gly
 90 95 100

- 100 -

- 101 -

ACCCCATTTN ATTCTAGAGT CNAGAACGCA AGGATCTCTT GGAATGACAA ATGATAGGTA 1350
 CCTAAAATGT AACATGAAAA TACTAGCTTA TTTTCTGAAA TGTACTATCT TAATGCTTAA 1410
 5 ATTATATTTC CCTTTAGGCT GTGATAGTTT TTGAAATAAA ATTTAACATT TAATATCATG 1470
 AAATGKTATA AGTAGACATA AAAAAAAA AAAAAAAA AGGGCGGCCG CTAGACTAG 1529

10

(2) INFORMATION FOR SEQ ID NO:8:

15

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 266 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
 1 5 10 15

30

Val Ala Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr
 20 25 30

Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro
 35 40 45

35

Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro
 50 55 60

40

Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr
 65 70 75 80

Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr
 85 90 95

45

Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu
 100 105 110

Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys
 115 120 125

50

Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn
 130 135 140

55

His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn
 145 150 155 160

Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser
 165 170 175

- 102 -

Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser
 180 185 190

5 Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys
 195 200 205

Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg
 210 215 220

10 Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly
 225 230 235 240

Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn
 245 250 255

15 Ser Ser Arg Leu His Thr Cys Gln Arg His
 260 265

(2) INFORMATION FOR SEQ ID NO:9:

20 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 798 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 25 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

30 (A) NAME/KEY: CDS
 (B) LOCATION: 1..798

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

35 ATG ATG GCT CTG GGC GCA GCG GGA GCT ACC CGG GTC TTT GTC GCG ATG 48
 Met Met Ala Leu Gly Ala Ala Gly Ala Thr Arg Val Phe Val Ala Met
 1 5 10 15

40 GTA GCG GCG GCT CTC GGC GGC CAC CCT CTG CTG GGA GTG AGC GCC ACC 96
 Val Ala Ala Leu Gly Gly His Pro Leu Leu Gly Val Ser Ala Thr
 20 25 30

45 TTG AAC TCG GTT CTC AAT TCC AAC GCT ATC AAG AAC CTG CCC CCA CCG 144
 Leu Asn Ser Val Leu Asn Ser Asn Ala Ile Lys Asn Leu Pro Pro Pro
 35 40 45

50 CTG GGC GGC GCT GCG GGG CAC CCA GGC TCT GCA GTC AGC GCC GCG CCG 192
 Leu Gly Gly Ala Ala Gly His Pro Gly Ser Ala Val Ser Ala Ala Pro
 50 55 60

55 GGA ATC CTG TAC CCG GGC GGG AAT AAG TAC CAG ACC ATT GAC AAC TAC 240
 Gly Ile Leu Tyr Pro Gly Gly Asn Lys Tyr Gln Thr Ile Asp Asn Tyr
 65 70 75 80

- 103 -

	CAG CCG TAC CCG TGC GCA GAG GAC GAG GAG TGC GGC ACT GAT GAG TAC	288
	Gln Pro Tyr Pro Cys Ala Glu Asp Glu Glu Cys Gly Thr Asp Glu Tyr	
	85 90 95	
5	TGC GCT AGT CCC ACC CGC GGA GGG GAC GCA GGC GTG CAA ATC TGT CTC	336
	Cys Ala Ser Pro Thr Arg Gly Gly Asp Ala Gly Val Gln Ile Cys Leu	
	100 105 110	
10	GCC TGC AGG AAG CGC CGA AAA CGC TGC ATG CGT CAC GCT ATG TGC TGC	384
	Ala Cys Arg Lys Arg Arg Lys Arg Cys Met Arg His Ala Met Cys Cys	
	115 120 125	
15	CCC GGG AAT TAC TGC AAA AAT GGA ATA TGC GTG TCT TCT GAT CAA AAT	432
	Pro Gly Asn Tyr Cys Lys Asn Gly Ile Cys Val Ser Ser Asp Gln Asn	
	130 135 140	
20	CAT TTC CGA GGA GAA ATT GAG GAA ACC ATC ACT GAA AGC TTT GGT AAT	480
	His Phe Arg Gly Glu Ile Glu Glu Thr Ile Thr Glu Ser Phe Gly Asn	
	145 150 155 160	
	CAT TGT GAT GGG TAT TCC AGA AGA ACC ACC TTG TCT TCA	528
	Asp His Ser Thr Leu Asp Gly Tyr Ser Arg Arg Thr Thr Leu Ser Ser	
	165 170 175	
25	AAA ATG TAT CAC ACC AAA GGA CAA GAA GGT TCT GTT TGT CTC CGG TCA	576
	Lys Met Tyr His Thr Lys Gly Gln Glu Gly Ser Val Cys Leu Arg Ser	
	180 185 190	
30	TCA GAC TGT GCC TCA GGA TTG TGT TGT GCT AGA CAC TTC TGG TCC AAG	624
	Ser Asp Cys Ala Ser Gly Leu Cys Cys Ala Arg His Phe Trp Ser Lys	
	195 200 205	
35	ATC TGT AAA CCT GTC CTG AAA GAA GGT CAA GTG TGT ACC AAG CAT AGG	672
	Ile Cys Lys Pro Val Leu Lys Glu Gly Gln Val Cys Thr Lys His Arg	
	210 215 220	
40	AGA AAA GGC TCT CAT GGA CTA GAA ATA TTC CAG CGT TGT TAC TGT GGA	720
	Arg Lys Gly Ser His Gly Leu Glu Ile Phe Gln Arg Cys Tyr Cys Gly	
	225 230 235 240	
	GAA GGT CTG TCT TGC CGG ATA CAG AAA GAT CAC CAT CAA GCC AGT AAT	768
	Glu Gly Leu Ser Cys Arg Ile Gln Lys Asp His His Gln Ala Ser Asn	
	245 250 255	
45	TCT TCT AGG CTT CAC ACT TGT CAG AGA CAC	798
	Ser Ser Arg Leu His Thr Cys Gln Arg His	
	260 265	
50	(2) INFORMATION FOR SEQ ID NO:10:	

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 702 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

55

- 104 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..537

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

10	GAA TTC GGC ACG AGG GTT GGG AGG TAT TGC CAC AGT CCC CAC CAA GGA Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly 1 5 10 15	48
15	TCA TCG GCC TGC ATG GTG TGT CGG AGA AAA AAG AAG CGC TGC CAC CGA Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg 20 25 30	96
20	GAT GGC ATG TGC TGC CCC AGT ACC CGC TGC AAT AAT GGC ATC TGT ATC Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile 35 40 45	144
25	CCA GTT ACT GAA AGC ATC TTA ACC CCT CAC ATC CCG GCT CTG GAT GGT Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly 50 55 60	192
30	ACT CGG CAC AGA GAT CGA AAC CAC GGT CAT TAC TCA AAC CAT GAC TTG Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu 65 70 75 80	240
35	GGA TGG CAG AAT CTA GGA AGA CCA CAC ACT AAG ATG TCA CAT ATA AAA Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys 85 90 95	288
40	GGG CAT GAA GGA GAC CCC TGC CTA CGA TCA TCA GAC TGC ATT GAA GGG Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly 100 105 110	336
45	TTT TGC TGT GCT CGT CAT TTC TGG ACC AAA ATC TGC AAA CCA GTG CTC Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu 115 120 125	384
50	CAT CAG GGG GAA GTC TGT ACC AAA CAA CGC AAG AAG GGT TCT CAT GGG His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly 130 135 140	432
	CTG GAA ATT TTC CAG CGT TGC GAC TGT GCG AAG GGC CTG TCT TGC AAA Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys 145 150 155 160	480
	GTA TGG AAA GAT GCC ACC TAC TCC TCC AAA GCC AGA CTC CAT GTG TGT Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys 165 170 175	528

- 105 -

CAG AAA ATT TGATCACCAT TGAGGAACAT CATCAATTGC AGACTGTGAA 577
Gln Lys Ile

GTTGTGTATT TAATGCATTA TAGCATGGTG GAAAATAAGG TTCAGATGCA GAAGAATGGC 637
5

TAAAATAAGA AACGTGATAA GAATATAGAT GATCACAAAA AAAAAAAA AAAAGATGCG 697

GCCGC 702

10

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 179 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Glu Phe Gly Thr Arg Val Gly Arg Tyr Cys His Ser Pro His Gln Gly
1 5 10 15

25

Ser Ser Ala Cys Met Val Cys Arg Arg Lys Lys Lys Arg Cys His Arg
20 25 30

Asp Gly Met Cys Cys Pro Ser Thr Arg Cys Asn Asn Gly Ile Cys Ile
35 40 45

30

Pro Val Thr Glu Ser Ile Leu Thr Pro His Ile Pro Ala Leu Asp Gly
50 55 60

35

Thr Arg His Arg Asp Arg Asn His Gly His Tyr Ser Asn His Asp Leu
65 70 75 80

Gly Trp Gln Asn Leu Gly Arg Pro His Thr Lys Met Ser His Ile Lys
85 90 95

40

Gly His Glu Gly Asp Pro Cys Leu Arg Ser Ser Asp Cys Ile Glu Gly
100 105 110

Phe Cys Cys Ala Arg His Phe Trp Thr Lys Ile Cys Lys Pro Val Leu
115 120 125

45

His Gln Gly Glu Val Cys Thr Lys Gln Arg Lys Lys Gly Ser His Gly
130 135 140

50

Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys
145 150 155 160

Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys
165 170 175

55

Gln Lys Ile

- 106 -

(2) INFORMATION FOR SEO ID NO:12:

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 537 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: cDNA

15 (ix) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 1..537

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GAA	TTC	GGC	ACG	AGG	GTT	GGG	AGG	TAT	TGC	CAC	AGT	CCC	CAC	CAA	GGA	48
Glu	Phe	Gly	Thr	Arg	Val	Gly	Arg	Tyr	Cys	His	Ser	Pro	His	Gln	Gly	
1	5				10				15							

TCA	TCG	GCC	TGC	ATG	GTG	TGT	CGG	AGA	AAA	AAG	AAG	CGC	TGC	CAC	CGA	96
Ser	Ser	Ala	Cys	Met	Val	Cys	Arg	Arg	Lys	Lys	Lys	Lys	Arg	Cys	His	Arg
20	25					25								30		

GAT	GGC	ATG	TGC	TGC	CCC	AGT	ACC	CGC	TGC	AAT	AAT	GGC	ATC	TGT	ATC	144
Asp	Gly	Met	Cys	Cys	Pro	Ser	Thr	Arg	Cys	Asn	Asn	Gly	Ile	Cys	Ile	
35	40					40			45							

CCA	GTT	ACT	GAA	AGC	ATC	TTA	ACC	CCT	CAC	ATC	CCG	GCT	CTG	GAT	GGT	192
Pro	Val	Thr	Glu	Ser	Ile	Leu	Thr	Pro	His	Ile	Pro	Ala	Leu	Asp	Gly	
50	55					55			60							

ACT	CGG	CAC	AGA	GAT	CGA	AAC	CAC	GGT	CAT	TAC	TCA	AAC	CAT	GAC	TTG	240
Thr	Arg	His	Arg	Asp	Arg	Asn	His	Gly	His	Tyr	Ser	Asn	His	Asp	Leu	
65	70					70			75					80		

GGA	TGG	CAG	AAT	CTA	GGA	AGA	CCA	CAC	ACT	AAG	ATG	TCA	CAT	ATA	AAA	288
Gly	Trp	Gln	Asn	Leu	Gly	Arg	Pro	His	Thr	Lys	Met	Ser	His	Ile	Lys	
85	90				90				95							

GGG	CAT	GAA	GGA	GAC	CCC	TGC	CTA	CGA	TCA	TCA	GAC	TGC	ATT	GAA	GGG	336
Gly	His	Glu	Gly	Asp	Pro	Cys	Leu	Arg	Ser	Ser	Asp	Cys	Ile	Glu	Gly	
45	100					105			110							

TTT	TGC	TGT	GCT	CGT	CAT	TTC	TGG	ACC	AAA	ATC	TGC	AAA	CCA	GTG	CTC	384
Phe	Cys	Cys	Ala	Arg	His	Phe	Trp	Thr	Lys	Ile	Cys	Lys	Pro	Val	Leu	
115	120					120			125							

CAT	CAG	GGG	GAA	GTC	TGT	ACC	AAA	CAA	CGC	AAG	AAG	GGT	TCT	CAT	GGG	432
His	Gln	Gly	Glu	Val	Cys	Thr	Lys	Gln	Arg	Lys	Lys	Gly	Ser	His	Gly	
130	135				135				140							

- 107 -

CTG GAA ATT TTC CAG CGT TGC GAC TGT GCG AAG GGC CTG TCT TGC AAA	480
Leu Glu Ile Phe Gln Arg Cys Asp Cys Ala Lys Gly Leu Ser Cys Lys	
145 150 155 160	
5 GTA TGG AAA GAT GCC ACC TAC TCC TCC AAA GCC AGA CTC CAT GTG TGT	528
Val Trp Lys Asp Ala Thr Tyr Ser Ser Lys Ala Arg Leu His Val Cys	
165 170 175	
10 CAG AAA ATT	537
Gln Lys Ile	

(2) INFORMATION FOR SEQ ID NO:13:

15 (i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 928 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: single	
(D) TOPOLOGY: linear	
20 (ii) MOLECULE TYPE: cDNA	
(ix) FEATURE:	
(A) NAME/KEY: CDS	
25 (B) LOCATION: 75..800	

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

30 CTCGAGGCCA AAATTCGGCA CGAGGCCGGG CTGTGGTCTA GCATAAAGGC GGAGCCCAGA	60
AGAAGGGGCG GGGT ATG GGA GAA GCC TCC CCA CCT GCC CCC GCA AGG CGG	110
Met Gly Ala Ser Pro Pro Ala Pro Ala Arg Arg	
1 5 10	
35 CAT CTG CTG GTC CTG CTG CTC CTC TCT ACC CTG GTG ATC CCC TCC	158
His Leu Leu Val Leu Leu Leu Ser Thr Leu Val Ile Pro Ser	
15 20 25	
40 GCT GCA GCT CCT ATC CAT GAT GCT GAC GCC CAA GAG AGC TCC TTG GGT	206
Ala Ala Ala Pro Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly	
30 35 40	
45 CTC ACA GGC CTC CAG AGC CTA CTC CAA GGC TTC AGC CGA CTT TTC CTG	254
Leu Thr Gly Leu Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu	
45 50 55 60	
50 AAA GGT AAC CTG CTT CGG GGC ATA GAC AGC TTA TTC TCT GCC CCC ATG	302
Lys Gly Asn Leu Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met	
65 70 75	
50 GAC TTC CGG GGC CTC CCT GGG AAC TAC CAC AAA GAG GAG AAC CAG GAG	350
Asp Phe Arg Gly Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu	
80 85 90	

- 108 -

CAC CAG CTG GGG AAC AAC ACC CTC TCC AGC CAC CTC CAG ATC GAC AAG His Gln Leu Gly Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys 95 100 105	398
5 ATG ACC GAC AAC AAG ACA GGA GAG GTG CTG ATC TCC GAG AAT GTG GTG Met Thr Asp Asn Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val 110 115 120	446
10 GCA TCC ATT CAA CCA GCG GAG GGG AGC TTC GAG GGT GAT TTG AAG GTA Ala Ser Ile Gln Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val 125 130 135 140	494
15 CCC AGG ATG GAG GAG AAG GAG GCC CTG GTA CCC ATC CAG AAG GCC ACG Pro Arg Met Glu Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr 145 150 155	542
20 GAC AGC TTC CAC ACA GAA CTC CAT CCC CGG GTG GCC TTC TGG ATC ATT Asp Ser Phe His Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile 160 165 170	590
25 AAG CTG CCA CGG CGG AGG TCC CAC CAG GAT GCC CTG GAG GGC GGC CAC Lys Leu Pro Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His 175 180 185	638
30 TGG CTC AGC GAG AAG CGA CAC CGC CTG CAG GCC ATC CGG GAT GGA CTC Trp Leu Ser Glu Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu 190 195 200	686
35 CGC AAG GGG ACC CAC AAG GAC GTC CTA GAA GAG GGG ACC GAG AGC TCC Arg Lys Gly Thr His Lys Asp Val Leu Glu Gly Thr Glu Ser Ser 205 210 215 220	734
40 TCC CAC TCC AGG CTG TCC CCC CGA AAG ACC CAC TTA CTG TAC ATC CTC Ser His Ser Arg Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu 225 230 235	782
45 AGG CCC TCT CGG CAG CTG TAGGGTGGG GACCGGGGAG CACCTGCCTG Arg Pro Ser Arg Gln Leu 240	830
50 TAGCCCCAT CAGACCCTGC CCCAAGCACC ATATGGAAAT AAAGTTCTTT CTTACATCTA AAAAAAAAAAA AAAAAAAAAA AAAAAAATTG GCGGCCGC	890
55 (2) INFORMATION FOR SEQ ID NO:14: (i) SEQUENCE CHARACTERISTICS: (A) LENGTH: 242 amino acids (B) TYPE: amino acid (D) TOPOLOGY: linear (ii) MOLECULE TYPE: protein (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:	928

- 109 -

Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val
 1 5 10 15

5 Leu Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro
 20 25 30

Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu
 35 40 45

10 Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu
 50 55 60

15 Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly
 65 70 75 80

Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly
 85 90 95

20 Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn
 100 105 110

Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln
 115 120 125

25 Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu
 130 135 140

30 Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr Asp Ser Phe His
 145 150 155 160

Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile Lys Leu Pro Arg
 165 170 175

35 Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His Trp Leu Ser Glu
 180 185 190

Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu Arg Lys Gly Thr
 195 200 205

40 His Lys Asp Val Leu Glu Glu Gly Thr Glu Ser Ser Ser His Ser Arg
 210 215 220

45 Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu Arg Pro Ser Arg
 225 230 235 240

Gln Leu

50 (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 726 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single

- 110 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

5 (ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..726

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

10	ATG GGA GAA GCC TCC CCA CCT GCC CCC GCA AGG CGG CAT CTG CTG GTC Met Gly Glu Ala Ser Pro Pro Ala Pro Ala Arg Arg His Leu Leu Val 1 5 10 15	48
15	CTG CTG CTG CTC CTC TCT ACC CTG GTG ATC CCC TCC GCT GCA GCT CCT Leu Leu Leu Leu Ser Thr Leu Val Ile Pro Ser Ala Ala Ala Pro 20 25 30	96
20	ATC CAT GAT GCT GAC GCC CAA GAG AGC TCC TTG GGT CTC ACA GGC CTC Ile His Asp Ala Asp Ala Gln Glu Ser Ser Leu Gly Leu Thr Gly Leu 35 40 45	144
25	CAG AGC CTA CTC CAA GGC TTC AGC CGA CTT TTC CTG AAA GGT AAC CTG Gln Ser Leu Leu Gln Gly Phe Ser Arg Leu Phe Leu Lys Gly Asn Leu 50 55 60	192
30	CTT CGG GGC ATA GAC AGC TTA TTC TCT GCC CCC ATG GAC TTC CGG GGC Leu Arg Gly Ile Asp Ser Leu Phe Ser Ala Pro Met Asp Phe Arg Gly 65 70 75 80	240
35	CTC CCT GGG AAC TAC CAC AAA GAG GAG AAC CAG GAG CAC CAG CTG GGG Leu Pro Gly Asn Tyr His Lys Glu Glu Asn Gln Glu His Gln Leu Gly 85 90 95	288
40	AAC AAC ACC CTC TCC AGC CAC CTC CAG ATC GAC AAG ATG ACC GAC AAC Asn Asn Thr Leu Ser Ser His Leu Gln Ile Asp Lys Met Thr Asp Asn 100 105 110	336
45	AAG ACA GGA GAG GTG CTG ATC TCC GAG AAT GTG GTG GCA TCC ATT CAA Lys Thr Gly Glu Val Leu Ile Ser Glu Asn Val Val Ala Ser Ile Gln 115 120 125	384
50	CCA GCG GAG GGG AGC TTC GAG GGT GAT TTG AAG GTA CCC AGG ATG GAG Pro Ala Glu Gly Ser Phe Glu Gly Asp Leu Lys Val Pro Arg Met Glu 130 135 140	432
55	GAG AAG GAG GCC CTG GTA CCC ATC CAG AAG GCC ACG GAC AGC TTC CAC Glu Lys Glu Ala Leu Val Pro Ile Gln Lys Ala Thr Asp Ser Phe His 145 150 155 160	480
60	ACA GAA CTC CAT CCC CGG GTG GCC TTC TGG ATC ATT AAG CTG CCA CGG Thr Glu Leu His Pro Arg Val Ala Phe Trp Ile Ile Lys Leu Pro Arg 165 170 175	528

- 111 -

	CGG AGG TCC CAC CAG GAT GCC CTG GAG GGC GGC CAC TGG CTC AGC GAG	576
	Arg Arg Ser His Gln Asp Ala Leu Glu Gly Gly His Trp Leu Ser Glu	
	180 185 190	
5	AAG CGA CAC CGC CTG CAG GCC ATC CGG GAT GGA CTC CGC AAG GGG ACC	624
	Lys Arg His Arg Leu Gln Ala Ile Arg Asp Gly Leu Arg Lys Gly Thr	
	195 200 205	
10	CAC AAG GAC GTC CTA GAA GAG GGG ACC GAG AGC TCC TCC CAC TCC AGG	672
	His Lys Asp Val Leu Glu Gly Thr Glu Ser Ser Ser His Ser Arg	
	210 215 220	
15	CTG TCC CCC CGA AAG ACC CAC TTA CTG TAC ATC CTC AGG CCC TCT CGG	720
	Leu Ser Pro Arg Lys Thr His Leu Leu Tyr Ile Leu Arg Pro Ser Arg	
	225 230 235 240	
	CAG CTG	726
	Gln Leu	

20

(2) INFORMATION FOR SEQ ID NO:16:

	(i) SEQUENCE CHARACTERISTICS:	
	(A) LENGTH: 2381 base pairs	
25	(B) TYPE: nucleic acid	
	(C) STRANDEDNESS: single	
	(D) TOPOLOGY: linear	
	(ii) MOLECULE TYPE: cDNA	
30	(ix) FEATURE:	
	(A) NAME/KEY: CDS	
	(B) LOCATION: 110..1156	
35	(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:	
	FGTCGACCCA CGCGTCCGCT GTGGCAGCCC AGCTACCGGT CGTGACCAGA TCCAGCTTGC	60
	AGCTCAGCTT TGTTCATTCTG AATTGGGCGG CGGCCAGCGC GGAACAAAC ATG CAG	115
40	Met Gln	
	1	
	CGG CTC GGG GGT ATT TTG CTG TGT ACA CTG CTG GCG GCG GCG GTC CCC	163
	Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Val Pro	
45	5 10 15	
	ACT GCT CCT GCT CCT TCC CCG ACG GTC ACT TGG ACT CCG GCG GAG CCG	211
	Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala Glu Pro	
	20 25 30	
50	GGC CCA GCT CTC AAC TAC CCT CAG GAG GAA GCT ACG CTC AAT GAG ATG	259
	Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn Glu Met	
	35 40 45 50	

- 112 -

	TTT CGA GAG GTG GAG GAG CTG ATG GAA GAC ACT CAG CAC AAA CTG CGC	307
	Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys Leu Arg	
	55 60 65	
5	AGT GCC GTG GAG GAG ATG GAG GCG GAA GAA GCA GCT GCT AAA ACG TCC	355
	Ser Ala Val Glu Glu Met Glu Ala Glu Ala Ala Lys Thr Ser	
	70 75 80	
10	TCT GAG GTG AAC CTG GCA AGC TTA CCT CCC AAC TAT CAC AAT GAG ACC	403
	Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn Glu Thr	
	85 90 95	
15	AGC ACG GAG ACC AGG GTG GGA AAT AAC ACA GTC CAT GTG CAC CAG GAA	451
	Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His Gln Glu	
	100 105 110	
20	GTT CAC AAG ATA ACC AAC AAC CAG AGT GGA CAG GTG GTC TTT TCT GAG	499
	Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe Ser Glu	
	115 120 125 130	
	ACA GTC ATT ACA TCT GTA GGG GAT GAA GAA GGC AAG AGG AGC CAT GAA	547
	Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser His Glu	
	135 140 145	
25	TGT ATC ATT GAT GAA GAC TGT GGG CCC ACC AGG TAC TGC CAG TTC TCC	595
	Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln Phe Ser	
	150 155 160	
30	AGC TTC AAG TAC ACC TGC CAG CCA TGC CGG GAC CAG CAG ATG CTA TGC	643
	Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met Leu Cys	
	165 170 175	
35	ACC CGA GAC AGT GAG TGC TGT GGA GAC CAG CTG TGT GCC TGG GGT CAC	691
	Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp Gly His	
	180 185 190	
40	TGC ACC CAA AAG GCC ACC AAA GGT GGC AAT GGG ACC ATC TGT GAC AAC	739
	Cys Thr Gln Lys Ala Thr Lys Gly Asn Gly Thr Ile Cys Asp Asn	
	195 200 205 210	
	CAG AGG GAT TGC CAG CCT GGC CTG TGT TGT GCC TTC CAA AGA GGC CTG	787
	Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg Gly Leu	
	215 220 225	
45	CTG TTC CCC GTG TGC ACA CCC CTG CCC GTG GAG GGA GAG CTC TGC CAT	835
	Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu Cys His	
	230 235 240	
50	GAC CCC ACC AGC CAG CTG CTG GAT CTC ATC ACC TGG GAA CTG GAG CCT	883
	Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu Glu Pro	
	245 250 255	
55	GAA GGA GCT TTG GAC CGA TGC CCC TGC GCC AGT GGC CTC CTA TGC CAG	931
	Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu Cys Gln	
	260 265 270	

CCA CAC AGC CAC AGT CTG GTG TAC ATG TGC AAG CCA GCC TTC GTG GGC	979
Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe Val Gly	
275 280 285 290	295
5 AGC CAT GAC CAC AGT GAG GAG AGC CAG CTG CCC AGG GAG GCC CCG GAT	1027
Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala Pro Asp	
295 300 305	
10 GAG TAC GAA GAT GTT GGC TTC ATA GGG GAA GTG CGC CAG GAG CTG GAA	1075
Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu Leu Glu	
310 315 320	
15 GAC CTG GAG CGG AGC CTA GCC CAG GAG ATG GCA TTT GAG GGG CCT GCC	1123
Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly Pro Ala	
325 330 335	
20 CCT GTG GAG TCA CTA GGC GGA GAG GAG GAG ATT TAGGCCAGA CCCAGCTGAG	1176
Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile	
340 345	
25 TCACTGGTAG ATGTGCAATA GAAATGGCTA ATTTATTTTC CCAGGAGTGT CCCCAAGTGT	1236
GGAATGGCCG CAGCTCCTTC CCAGTAGCTT TTCTCTGGC TTGACAAGGT ACAGTGCAGT	1296
25 ACATTTCTTC CAGCCGCCCT GCTTCTCTGA CTTGGAAAG ACAGGCATGG CGGGTAAGGG	1356
CAGCGGTGAG TCGTCCCTCG CTGTTGCTAG AAACGCTGTC TTGTTCTCA TGGATGGAAG	1416
30 ATTTGTTGA AGGGAGAGGA TGGGAAGGGG TGAAGTCTGC TCATGATGGA TTTGGGGGAT	1476
ACAGGGAGGA GGATGCCTGC CTTGCAGACG TGGACTTGGC AAAATGTAAC CTTTGCTTT	1536
35 GTCTTGCAGCC GCTCCCATGG GCTGAGGCAG TGGCTACACA AGAGCTATGC TGCTCTGTGG	1596
CCTCCACAT ATTCACTCCCT GTGTTTCAGC TCCTACCTCA CTGTCAGCAC AGCCCTTCAT	1656
AGCCACGCC CCTCTTGCTC ACCACAGCCT AGGAGGGGAC CAGAGGGGAC TTCTCTCAGA	1716
40 GCCCCATGCT CTCTCTCTCA ACCCCATACC AGCCTCTGTG CCAGCGACAG TCCTTCCAAA	1776
TGGAGGGAGT GAAATCCTT GGTTAATTAA TTTCTCCTT CAAGGCACGC CTGCCACTAA	1836
45 GGTCAGGCTG ACTTGCATGT CCCTCTAACG TTCTAGCAG TGTGGTGGAC ACTGTCTTCC	1896
ACCGACTGCT TCAATACCTC TGAAAGCCAG TGCTCGGAGT GCAGTTCGTG TAAATTAAATT	1956
TGCAGGAAGT ATACTTGGCT AATTGTAGGG CTAGGATTGT GAATGAAATT TGCAAAGTCG	2016
50 CTTAGCAACA ATGGAAAGCC TTTCTCAGTC ACACCGAGAA GTCACAACCA AGCCAGGTTG	2076
TGTAGAGTAC AGCTGTGACA TACAGACAGA AGAAGGCTGG GCTGGATGTC AGGCCTCAGA	2136
55 TGACGGTTTC AGGTGCCAGG AACTATTACC ATTCTGTATC TATCCAGAGT TATTAAAATT	2196

- 114 -

GAAAGTTGCA	CACATTTGTA	TAAGCATGCC	TTTCTCCTGA	GTTTTAAATT	ATATGTATAC	2256
ACAAACATGT	GGCCCTCAAA	GATCATGCAC	AAACCACTAC	TCTTGCTAA	TTCTTGGACT	2316
5	TTTCTCTTG	ATTTCAATA	AATACAAATC	CCCTTCATGC	AAAAAAAAAA	AAAAAGGGCG
						2376
	GCCGC					2381

10 (2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 349 amino acids
 (B) TYPE: amino acid
 15 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

20	Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala	1	5	10	15
	Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala	20	25	30	
25	Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Glu Ala Thr Leu Asn	35	40	45	
30	Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys	50	55	60	
	Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Glu Ala Ala Ala Lys	65	70	75	80
35	Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn	85	90	95	
40	Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His	100	105	110	
	Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe	115	120	125	
45	Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser	130	135	140	
	His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln	145	150	155	160
50	Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met	165	170	175	
55	Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp	180	185	190	

- 115 -

Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys
 195 200 205

5 Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg
 210 215 220

Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu
 225 230 235 240

10 Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu
 245 250 255

Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
 15 260 265 270

Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe
 275 280 285

20 Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala
 290 295 300

Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu
 305 310 315 320

25 Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly
 325 330 335

30 Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
 340 345

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 1047 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40 (ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS
 (B) LOCATION: 1..1047

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ATG CAG CGG CTC GGG GGT ATT TTG CTG TGT ACA CTG CTG GCG GCG GCG
 Met Gln Arg Leu Gly Gly Ile Leu Leu Cys Thr Leu Leu Ala Ala Ala
 50 1 5 10 15

48

GTC CCC ACT GCT CCT GCT TCC CCG ACG GTC ACT TGG ACT CCG GCG
 Val Pro Thr Ala Pro Ala Pro Ser Pro Thr Val Thr Trp Thr Pro Ala
 20 25 30

96

55

- 116 -

	GAG CCG GGC CCA GCT CTC AAC TAC CCT CAG GAG GAA GCT ACG CTC AAT		144
	Glu Pro Gly Pro Ala Leu Asn Tyr Pro Gln Glu Ala Thr Leu Asn		
	35 40 45		
5	GAG ATG TTT CGA GAG GTG GAG GAG CTG ATG GAA GAC ACT CAG CAC AAA		192
	Glu Met Phe Arg Glu Val Glu Glu Leu Met Glu Asp Thr Gln His Lys		
	50 55 60		
10	CTG CGC AGT GCC GTG GAG GAG ATG GAG GCG GAA GAA GCA GCT GCT AAA		240
	Leu Arg Ser Ala Val Glu Glu Met Glu Ala Glu Ala Ala Ala Lys		
	65 70 75 80		
15	ACG TCC TCT GAG GTG AAC CTG GCA AGC TTA CCT CCC AAC TAT CAC AAT		288
	Thr Ser Ser Glu Val Asn Leu Ala Ser Leu Pro Pro Asn Tyr His Asn		
	85 90 95		
20	GAG ACC AGC ACG GAG ACC AGG GTG GGA AAT AAC ACA GTC CAT GTG CAC		336
	Glu Thr Ser Thr Glu Thr Arg Val Gly Asn Asn Thr Val His Val His		
	100 105 110		
	CAG GAA GTT CAC AAG ATA ACC AAC AAC CAG AGT GGA CAG GTG GTC TTT		384
	Gln Glu Val His Lys Ile Thr Asn Asn Gln Ser Gly Gln Val Val Phe		
	115 120 125		
25	TCT GAG ACA GTC ATT ACA TCT GTA GGG GAT GAA GAA GGC AAG AGG AGC		432
	Ser Glu Thr Val Ile Thr Ser Val Gly Asp Glu Glu Gly Lys Arg Ser		
	130 135 140		
30	CAT GAA TGT ATC ATT GAT GAA GAC TGT GGG CCC ACC AGG TAC TGC CAG		480
	His Glu Cys Ile Ile Asp Glu Asp Cys Gly Pro Thr Arg Tyr Cys Gln		
	145 150 155 160		
35	TTC TCC AGC TTC AAG TAC ACC TGC CAG CCA TGC CGG GAC CAG CAG ATG		528
	Phe Ser Ser Phe Lys Tyr Thr Cys Gln Pro Cys Arg Asp Gln Gln Met		
	165 170 175		
40	CTA TGC ACC CGA GAC AGT GAG TGC TGT GGA GAC CAG CTG TGT GCC TGG		576
	Leu Cys Thr Arg Asp Ser Glu Cys Cys Gly Asp Gln Leu Cys Ala Trp		
	180 185 190		
	GGT CAC TGC ACC CAA AAG GCC ACC AAA GGT GGC AAT GGG ACC ATC TGT		624
	Gly His Cys Thr Gln Lys Ala Thr Lys Gly Gly Asn Gly Thr Ile Cys		
	195 200 205		
45	GAC AAC CAG AGG GAT TGC CAG CCT GGC CTG TGT TGT GCC TTC CAA AGA		672
	Asp Asn Gln Arg Asp Cys Gln Pro Gly Leu Cys Cys Ala Phe Gln Arg		
	210 215 220		
50	GGC CTG CTG TTC CCC GTG TGC ACA CCC CTG CCC GTG GAG GGA GAG CTC		720
	Gly Leu Leu Phe Pro Val Cys Thr Pro Leu Pro Val Glu Gly Glu Leu		
	225 230 235 240		
55	TGC CAT GAC CCC ACC AGC CAG CTG CTG GAT CTC ATC ACC TGG GAA CTG		768
	Cys His Asp Pro Thr Ser Gln Leu Leu Asp Leu Ile Thr Trp Glu Leu		
	245 250 255		

- 117 -

GAG CCT GAA GGA GCT TTG GAC CGA TGC CCC TGC GCC AGT GGC CTC CTA 816
Glu Pro Glu Gly Ala Leu Asp Arg Cys Pro Cys Ala Ser Gly Leu Leu
260 265 270

5 TGC CAG CCA CAC AGC CAC AGT CTG GTG TAC ATG TGC AAG CCA GCC TTC 864
Cys Gln Pro His Ser His Ser Leu Val Tyr Met Cys Lys Pro Ala Phe
275 280 285

10 GTG GGC AGC CAT GAC CAC AGT GAG GAG AGC CAG CTG CCC AGG GAG GCC 912
Val Gly Ser His Asp His Ser Glu Glu Ser Gln Leu Pro Arg Glu Ala
290 295 300

15 CCG GAT GAG TAC GAA GAT GTT GGC TTC ATA GGG GAA GTG CGC CAG GAG 960
Pro Asp Glu Tyr Glu Asp Val Gly Phe Ile Gly Glu Val Arg Gln Glu
305 310 315 320

20 CTG GAA GAC CTG GAG CGG AGC CTA GCC CAG GAG ATG GCA TTT GAG GGG 1008
Leu Glu Asp Leu Glu Arg Ser Leu Ala Gln Glu Met Ala Phe Glu Gly
325 330 335

25 CCT GCC CCT GTG GAG TCA CTA GGC GGA GAG GAG GAG ATT 1047
Pro Ala Pro Val Glu Ser Leu Gly Gly Glu Glu Glu Ile
340 345

What is claimed:

1. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof;
 - b) a nucleic acid molecule comprising a fragment of at least 1000 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof;
 - c) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, or SEQ ID NO:11, the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452;
 - d) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the

plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452; and

5 e) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452,

10 wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA

15 insert of the plasmid deposited with ATCC as Accession Number 98452 under stringent conditions.

2. The isolated nucleic acid molecule of claim 1 which is selected from the

20 group consisting of:

25 a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, or a complement thereof; and

30 b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452.

35

3. The nucleic acid molecule of claim 1 further comprising vector nucleic acid sequences.

4. The nucleic acid molecule of claim 1 further comprising nucleic acid sequences encoding a heterologous polypeptide.

5. A host cell which contains the nucleic acid molecule of claim 1.

6. The host cell of claim 5 which is a mammalian host cell.

7. A non-human mammalian host cell containing the nucleic acid molecule 10 of claim 1.

8. An isolated polypeptide selected from the group consisting of:

15 a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 20 98452, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 25 98452;

30 b) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited 35

with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 under stringent conditions;

5 c) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the DNA insert of the plasmid deposited with

10 ATCC as Accession Number 98452; and

15 d) a polypeptide comprising an amino acid sequence which is at least 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452.

9. The isolated polypeptide of claim 8 comprising the amino acid sequence
20 of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide
encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number
98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC
as Accession Number 98633, or the polypeptide encoded by the DNA insert of the
plasmid deposited with ATCC as Accession Number 98452.

25

10. The polypeptide of claim 8 further comprising heterologous amino acid
sequences.

30

11. An antibody which selectively binds to a polypeptide of claim 8.

35 12. A method for producing a polypeptide selected from the group consisting
of:
 a) a polypeptide comprising the amino acid sequence of SEQ ID
NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded
by the DNA insert of the plasmid deposited with ATCC as Accession Number
98634, the polypeptide encoded by the DNA insert of the plasmid deposited with

ATCC as Accession Number 98633, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452;

5 b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452; and

10 15 20 25 c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98452, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, the DNA insert of the plasmid deposited with ATCC as Accession Number 98634, the DNA insert of the plasmid deposited with ATCC as Accession Number 98633, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98452 under stringent conditions;

30 comprising culturing the host cell of claim 5 under conditions in which the nucleic acid molecule is expressed.

13. A method for detecting the presence of a polypeptide of claim 8 in a 35 sample comprising:

a) contacting the sample with a compound which selectively binds to the polypeptide; and

b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 8 in the sample.

5 14. The method of claim 13, wherein the compound which binds to the polypeptide is an antibody.

10 15. A kit comprising a compound which selectively binds to a polypeptide of claim 8 and instructions for use.

15 16. A method for detecting the presence of a nucleic acid molecule in claim 1 in a sample comprising:

a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and

b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of claim 1 in the sample.

20 17. The method of claim 16, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

25 18. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

19. A method for identifying a compound which binds to a polypeptide of claim 8 comprising:

a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and

b) determining whether the polypeptide binds to the test compound.

30 20. The method of claim 19, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:

a) detection of binding by direct detection of test compound/polypeptide binding;

b) detection of binding using a competition binding assay; and

c) detection of binding using an assay for CRSP activity.

- 124 -

21. A method of modulating the activity of a polypeptide of claim 8 comprising contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.

5

22. A method for identifying a compound which modulates the activity of a polypeptide of claim 8 comprising:

10

- a) contacting a polypeptide of claim 8 with a test compound; and
- b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

Figure 1A

Figure 1B

ATCTGGACCGGGCTGGCTAGATGTGCAATAGAAATAGCTAATTATTTCCTCCANGTGTGCTTAAGCGTGGGCTG	1169
ACCAAGGCTTCTCTACATCTCTTCCCTAGTAAGTTCCCTCTGGCTGACAGCATGAGGTGTTGCAATTGTTCAAG	1248
CTCCCCCAGGGCTTCTCAGGCTTCAAGTCTGGCTGGAGAGTCAGGCAGGGTTAAACTGCGAGGAGCTGGCTTGGC	1327
CACCCCTGTCAGATTATGGCTGTTGGCTCTACCAAGTGGCAGACAGCCGTTCTACATGGCTTGTATAATTG	1406
TTTGAGGGAGGAGATGGAAACAAATGTGGAGTCTCCCTCTGATTGGTTTGGGAAATGTGGAGAAGAGTCCCCTG	1485
TGCAAAACATCAACCTGGAAAATGCAAAATGAATTTCACGGCAGTTCTCCATGGGCATAGGTAAAGCTGTG	1564
TCAGCTGTTGAGATGAAATGTTCTGTTCACCCCTGCATTACATGTGTTATTACATCCAGCAGTGTGCTCAGT	1643
CTCTGTGCCAGGGCAGGATTTCAATCCAAGATCAATTCCCTCTCAGCACAGCCTGGGGGGTCATTGTTCTC	1722
CTCGTCCATCAGGGATTTCAGAGGACTGCAAGCTGCCAAGTCAAGCTGCAACACAGCTAGTGAAGACAGAGC	1801
TTTCATCTGGTTCTGACTCTAAAGCTCAAGTGCCTCCACTACCCACACCCAGCCTGGCACACAAAGTGTCT	1880
AAAAGGAAGGAAATGGGATTTTTCTTTGAGGCATGCAACTAACTAAAGGTCACATCTGGAATTAAAGTCAACT	1959
AAAGTAACACTGTTAGAACAGCAGTGTCTCACAGTGTGGGGCACGGCTCTTAATGAAAGACAATGATATTG	2038
ACTGTCCCTCTTGGCAGGGCATTAGTAACCTGAAAGGTATATGACTGAGCTGAGCTGACGGTTAACCTGCAG	2117
CAGGACTTAGGTAATTGTAGGGCAGGATTATAATGAAATTGCAAAATCACTTAGCAGCAACTGAAAGACAATT	2196
ACCAAGCTGGAGAAAATCAACCGGAGCAGGGCTCTGTGAAACATGGTTGTAATATGCAACTGCAACTCTA	2275
CCACTCCACAAATGATGTGTTCAAGGTGTCACATGGACTGTCACCATGATTACATCAGAGTCTAAAGTTAA	2354
GCACATGATTGATAAGCATGCTTCTTGAGTTAAATTATGTATAAACATAAGGTGCATTAGAAATCAAGCATA	2433
ATCACTCAACTGCTAAAAA.....	2479

2/12

3/12

Figure 2

GAATTGGCACGGAGAGAACGACGGTGGCTGAGCTAGTGGAAAGCTCTGGGGAGAGGCCCTGGCTTGTG 79

GTGACGCACAGTGTGGACCCCTCCAGGAGCCCCGATTGAGG ATG GTG GCG GCG CTG CTG GGG 8
CTG AGC TGG CTC TGC TCT CCC CTG GGA GCT CTG GTC CTG GAC TTC AAC ATC AGG AGC 148

S A D L H G A R K G S Q C L S D T D C N I R S 28
TCT GCT GAC CTG CAT GGG GCC CGG AAG GGC TCA CAG TGC CTG TCT GAC ACG GAC TGC AAT 268

T R K F C L Q P R D E K P F C A T C R G 68
ACC AGA AAG TTC TGC CTC CAG CCC CGC GAT GAG AAG CCG TTC TGT GCT ACA TGT CGT GGG 328

L R R R C Q R D A M C C P G T L C V N D 88
TTG CGG AGG AGG TGC CAG CGA GAT GCC ATG TGC CCT GGG ACA CTC TGT GTG AAC GAT 388

V C T T M E D A T P I L E R Q L D E Q D 108
GTT TGT ACT ACG ATG GAA GAT GCA ACC CCA ATA TTA GAA AGG CAG CTT GAT GAG CAA GAT 448

G T H A E G T T G H P V Q E N Q P K R K 128
GGC ACA CAT GCA GAA GGA ACA ACT GGG CAC CCA GTC CAG GAA AAC CAA CCC AAA AGG AAG 508

P S I K S Q G R K G Q E G E S C L R T 148
CCA AGT ATT AAG AAA TCA CAA CGC AGG AAG GGA CAA GAG GGA GAA AGT TGT CTG AGA ACT 568

F D C G P G L C C A R H F W T K I C K P 168
TTT GAC TGT GGC CCT GGA CTT TGC TGT GCT CGT CAT TTT TGG ACG AAA ATT TGT AAG CCA 628

V L L E G Q V C S R R G H K D T A Q A P 188
GTC CTT TTG GAG GGA CAG GTC TCC AGA AGA GGG CAT AAA GAC ACT GCT CAA GCT CCA 688

E I F Q R C D C G P G L L C R S Q L T S 208
GAA ATC TTC CAG CGT TGC GAC TGT GGC CCT GGA CTA CTG TGT CGA AGC CAA TTG ACC AGC 748

N R Q H A R L R V C Q K I E K L * 225
AAT CGG CAG CAT GCT CGA TTA AGA GTA TGC CAA AAA ATA GAA AAG CTA TAA 799

ATATTTCAAAATAAGAGAATCCACATTGCAAAAAAAAAAAAAAA 848

Figure 3A

CGGGACCGCTGGCGGCACGGTTCTGGGGACCCAGGCTTGC
 AAGTGACGGTCATTTCTCTTCTCCCTCTT 79
 M M A L G A A G A T R V F V A M 16
 GAGTCCTCTGAG ATG ATG GCT CTG GGC GCA GCG GGA GCT ACC CGG GTC TTT GTC GCG ATG 140
 V A A A L G G H P L L G V S A T L N S V 36
 GTA GCG GCG GCT CTC GGC GGC CAC CCT CTG CTG GGA GTG AGC GCC ACC TTG AAC TCG GTT 200
 L N S N A I K N L P P P L G G A A G H P 56
 CTC AAT TCC AAC GCT ATC AAG AAC CTG CCC CCA CCG CTG GGC GGC GCT GCG GGG CAC CCA 260
 G S A V S A A P G I L Y P G G N K Y Q T 76
 GGC TCT GCA GTC AGC GCC GCG CCG GGA ATC CTG TAC CCG GGC GGG AAT AAG TAC CAG ACC 320
 I D N Y Q P Y P C A E D E E C G T D E Y 96
 ATT GAC AAC TAC CAG CCG TAC CCG TGC GCA GAG GAC GAG GAG TGC GGC ACT GAT GAG TAC 380
 C A S P T R G G D A G V Q I C L A C R K 116
 TGC GCT AGT CCC ACC CGC GGA GGG GAC GCA GGC GTG CAA ATC TGT CTC GCC TGC AGG AAG 440
 R R K R C M R H A M C C P G N Y C K N G 136
 CGC CGA AAA CGC TGC ATG CGT CAC GCT ATG TGC TGC CCC GGG AAT TAC TGC AAA AAT GGA 500
 I C V S S D Q N H F R G E I E E T I T E 156
 ATA TGC GTG TCT TCT GAT CAA AAT CAT TTC CGA GGA GAA ATT GAG GAA ACC ATC ACT GAA 560
 S F G N D H S T L D G Y S R R T T L S S 176
 AGC TTT GGT AAT GAT CAT AGC ACC TTG GAT GGG TAT TCC AGA AGA ACC ACC TTG TCT TCA 620
 K M Y H T K G Q E G S V C L R S S D C A 196
 AAA ATG TAT CAC ACC AAA GGA CAA GAA GGT TCT GTT TGT CTC CGG TCA TCA GAC TGT GCC 680
 S G L C C A R H F W S K I C K P V L K E 216
 TCA GGA TTG TGT TGT GCT AGA CAC TTC TGG TCC AAG ATC TGT AAA CCT GTC CTG AAA GAA 740
 G Q V C T K H R R K G S H G L E I F Q R 236
 GGT CAA GTG TGT ACC AAG CAT AGG AGA AAA GGC TCT CAT GGA CTA GAA ATA TTC CAG CGT 800
 C Y C G E G L S C R I Q K D H H Q A S N 256
 TGT TAC TGT GGA GAA GGT CTG TCT TGC CGG ATA CAG AAA GAT CAC CAT CAA GCC AGT AAT 860
 S S R L H T C Q R H * 267
 TCT TCT AGG CTT CAC ACT TGT CAG AGA CAC TAA 893
 ACCAGCTATCCAAATGCAGTGAACCTCTTTATATAATAGATGCTATGAAACCTTTATGACCTTCATCAACTCAAT 972
 CCTAAGGATATACAAGTTCTGTGGTTTCAGTTAACGCATTCCAATAACACCTCCAAAAACCTGGAGTGTAAAGAGCTTG 1051
 TTTCTTATGAAACTCCCCTGTGATTGCACTGCCTATTTCTCTTGTAAATTCTCAGTGTGGCACTTACCTGTAAATGCA 1130
 ATGAAACTTTAATTATTTCTAAAGGTGCTGCACTGCCTATTTCTCTTGTAAATTCTCAGTGTGGCACTTACCTGTAAATGCA 1209
 TTGTTATCTGACTGACAAATATTCTATATTGAACGTGAAGTAAATCATTCACTTACCTTAAAGCATAACCCT 1288
 TTACCCCATTNATTCTAGAGTCNAGAACGCAAGGATCTCTTGAATGACAAATGATAGGTACCTAAAATGTAACATGA 1367

Figure 3B

AAATACTAGCTTATTCTCTGAAATGTAATGCTTAATCTTAATGCTTAATTATATTCCCTTACGGCTGTGATAGTTTGAAA 1446
TAAAATTAAACATTAAATATCATGAAATGKATAAGTAGACATAAAAAAAAGGGGGGGCCGCTAGA 1525
CTAG 1529

6/12

Figure 4

E	F	G	T	R	V	G	R	Y	C	H	S	P	H	Q	G	S	S	A	C	20
GAA	TTC	GGC	ACG	AGG	GTT	GGG	AGG	TAT	TGC	CAC	AGT	CCC	CAC	CAA	GGG	TCA	TCG	GCC	TGC	60
M	V	C	R	R	K	K	K	R	C	H	R	D	G	M	C	C	P	S	T	40
ATG	GTG	TGT	CGG	AGA	AAA	AAG	AAG	CAC	TGC	CGC	GAT	GGC	ATG	TGC	CCC	AGT	ACC	120		
R	C	N	N	G	I	C	I	P	V	T	E	S	I	L	T	P	H	I	P	60
CGC	TGC	AAT	AAT	GGC	ATC	TGT	ATC	CCA	GTT	ACT	GAA	AGC	ATC	TTA	ACC	CCT	CAC	ATC	CCG	180
A	L	D	G	T	R	H	R	D	R	N	H	G	H	Y	S	N	H	D	L	80
GCT	CTG	GAT	GGT	ACT	CGG	CAC	AGA	GAT	CGA	AAC	CAC	GGT	CAT	TAC	TCA	AAC	CAT	GAC	TG	240
G	W	Q	N	L	G	R	P	H	T	K	M	S	H	I	K	G	H	E	G	100
GGA	TGG	CAG	AAT	CTA	GGA	AGA	CCA	CAC	ACT	AAG	ATG	TCA	CAT	ATA	AAA	GGG	CAT	GAA	GGA	300
D	P	C	L	R	S	S	D	C	I	E	G	F	C	C	A	R	H	F	W	120
GAC	CCC	TGC	CTA	CGA	TCA	TCA	GAC	TGC	ATT	GAA	GGG	TTT	TGC	TGT	GCT	CGT	CAT	TTC	TGG	360
T	K	I	C	K	P	V	L	H	Q	G	E	V	C	T	K	Q	R	K	K	140
ACC	AAA	ATC	TGC	AAA	CCA	GTG	CTC	CAT	CAG	GGG	GAA	GTC	TGT	ACC	AAA	CAA	CGC	AAG	AAG	420
G	S	H	G	L	E	I	F	Q	R	C	D	C	A	K	G	L	S	C	K	160
GGT	TCT	CAT	GGG	CTG	GAA	ATT	TTC	CAG	CGT	TGC	GAC	TGT	GCG	AAG	GGC	CTG	TCT	TGC	AAA	480
V	W	K	D	A	T	Y	S	S	K	A	R	L	H	V	C	Q	K	I	*	180
GTA	TGG	AAA	GAT	GCC	ACC	TAC	TCC	TCC	AAA	GCC	AGA	CTC	CAT	GTG	TGT	CAG	AAA	ATT	TGA	540
TCACCATTTGAGGAACATCATCAATTGAGACTGTGAAGTTGTTATTIAATGCATTATAGCATGGTGGAAATAAGGTT	CAGATGGCAGAAGAATGGCTAAAATAAGAACGTTGATAAGAATAATAGATGATCACAAAAAAAGATGCGG	619																		
CCGC		698																		

7/12

Figure 5

CTCGAGGCCAAAATCGGCACGAGGCCGGCTGTTCTAGCATAAAGGCAGGCCAGAAGAAGGGCGGGGT	M 1
ATG	77
G E A S P P A P A R R H L L L V L L L L L L	21
GGA GAA GCC TCC CCA CCT GCC CCC GCA AGG CGG CAT CTG CTG GTC CTG CTG CTG CTC CTC	137
S T L V I P S A A A P I H D A D A Q E S	41
TCT ACC CTG GTG ATC CCC TCC GCT GCA GCT CCT ATC CAT GAT GCT GAC GCC CAA GAG ACC	197
S L G L T G L Q S L L Q G F S R L F L K	61
TCC TTG GGT CTC ACA GGC CTC CAG AGC CTA CTC CAA GGC TTC AGC CGA CTT TTC CTG AAA	257
G N L L R G I D S L F S A P M D F R G L	81
GGT AAC CTG CTT CGG GGC ATA GAC AGC TTA TTC TCT GCC CCC ATG GAC TTC CGG GGC CTC	317
P G N Y H K E E N Q E H Q L G N N T L S	101
CCT GGG AAC TAC CAC AAA GAG GAG AAC CAG GAG CAC CAG CTG GGG AAC AAC ACC CTC TCC	377
S H L Q I D K M T D N K T G E V L I S E	121
AGC CAC CTC CAG ATC GAC AAG ACC GAC AAC AAG ACA GGA GAG GTG CTG ATC TCC GAG	437
N V V A S I Q P A E G S F E G D L K V P	141
AAT GTG GTG GCA TCC ATT CAA CCA GCG GAG GGG AGC TTC GAG GGT GAT TTG AAG GTA CCC	497
R M E E K E A L V P I Q K A T D S F H T	161
AGG ATG GAG GAG AAG GAG GCC CTG GTA CCC ATC CAG AAG GCC ACG GAC AGC TTC CAC ACA	557
E L H P R V A F W I I K L P R R R S H Q	181
GAA CTC CAT CCC CGG GTG GCC TTC TGG ATC ATT AAG CTG CCA CGG CGG AGG TCC CAC CAG	617
D A L E G G H W L S E K R H R L Q A I R	201
GAT GCC CTG GAG GGC GGC CAC TGG CTC AGC GAG AAG CGA CAC CGC CTG CAG GCC ATC CGG	677
D G L R K G T H K D V L E E G T E S S S	221
GAT GGA CTC CGC AAG GGG ACC CAC AAG GAC GTC CTA GAA GAG GGG ACC GAG AGC TCC TCC	737
H S R L S P R K T H L L Y I L R P S R Q	241
CAC TCC AGG CTG TCC CCC CGA AAG ACC CAC TTA CTG TAC ATC CTC AGG CCC TCT CGG CAG	797
L *	243
CTG TAG	803
GGGTGGGGACGGGGAGCACCTGCCTGTAGCCCCATCAGACCTGCCAAGCACCATATGGAAATAAGTTCTTCT	882
TACATCTAAAAAAAAAAAAAAAAAAAAAAATTGGCGCCGC	928


8/12

FIGURE 6

9/12

The Human CRSP Family

FIGURE 7



10/12

Figure 8A


FGTCGACCCACCGCGTCCGCTGTGGCAGCCCAGCTACCGGTCGTGACCAGATCCAGCTTGAGCTCAGCTTGTTCATTC 79

M	Q	R	L	G	G	I	L	L	C	T	L	12								
GAATTGGCGGCCAGCGCGAACAAAC	ATG	CAG	CGG	CTC	GGG	GGT	ATT	TTG	CTG	TGT	ACA	CTG	145							
L	A	A	A	V	P	T	A	P	P	S	P	T	32							
CTG	GCG	GCG	GCG	GTC	CCC	ACT	GCT	CCT	GCT	CCT	TCC	CCG	ACG	GTC	ACT	TGG	ACT	CCG	GCG	205
E	P	G	P	A	L	N	Y	P	Q	E	E	A	T	L	N	E	M	F	R	52
GAG	CCG	GGC	CCA	GCT	CTC	AAC	TAC	CCT	CAG	GAG	GAA	GCT	ACG	CTC	AAT	GAG	ATG	TTT	CGA	265
E	V	E	E	L	M	E	D	T	Q	H	K	L	R	S	A	V	E	E	M	72
GAG	GTG	GAG	GAG	CTG	ATG	GAA	GAC	ACT	CAG	CAC	AAA	CTG	CGC	AGT	GCC	GTG	GAG	GAG	ATG	325
E	A	E	E	A	A	A	K	T	S	S	E	V	N	L	A	S	L	P	P	92
GAG	GCG	GAA	GAA	GCA	GCT	GCT	AAA	ACG	TCC	TCT	GAG	GTG	AAC	CTG	GCA	AGC	TTA	CCT	CCC	385
N	Y	H	N	E	T	S	T	E	T	R	V	G	N	N	T	V	H	V	H	112
AAC	TAT	CAC	AAT	GAG	ACC	AGC	ACG	GAG	ACC	AGG	GTG	GGA	AAT	AAC	ACA	GTC	CAT	GTG	CAC	445
Q	E	V	H	K	I	T	N	N	Q	S	G	Q	V	V	F	S	E	T	V	132
CAG	GAA	GTT	CAC	AAG	ATA	ACC	AAC	AAAC	CAG	AGT	GGGA	CAG	GTG	GTC	TTT	TCT	GAG	ACA	GTC	505
I	T	S	V	G	D	E	E	G	K	R	S	H	E	C	I	I	D	E	D	152
ATT	ACA	TCT	GTA	GGG	GAT	GAA	GAA	GGC	AAG	AGG	AGC	CAT	GAA	TGT	ATC	ATT	GAT	GAA	GAC	565
C	G	P	T	R	Y	C	Q	F	S	S	F	K	Y	T	C	Q	P	C	R	172
TGT	GGG	CCC	ACC	AGG	TAC	TGC	CAG	TTC	TCC	AGC	TTC	AAG	TAC	ACC	TGC	CAG	CCA	TGC	CGG	625
D	Q	Q	M	L	C	T	R	D	S	E	C	C	G	D	Q	L	C	A	W	192
GAC	CAG	CAG	ATG	CTA	TGC	ACC	CGA	GAC	AGT	GAG	TGC	TGT	GGA	GAC	CAG	CTG	TGT	GCC	TGG	685
G	H	C	T	Q	K	A	T	K	G	G	N	G	T	I	C	D	N	Q	R	212
GGT	CAC	TGC	ACC	CAA	AAG	GCC	ACC	AAA	GGT	GGC	AAT	GGG	ACC	ATC	TGT	GAC	AAC	CAG	AGG	745
D	C	Q	P	G	L	C	C	A	F	Q	R	G	L	L	F	P	V	C	T	232
GAT	TGC	CAG	CCT	GGC	CTG	TGT	TGT	GCC	TTC	CAA	AGA	GGC	CTG	CTG	TTC	CCC	GTG	TGC	ACA	805
P	L	P	V	E	G	E	L	C	H	D	P	T	S	Q	L	L	D	L	I	252
CCC	CTG	CCC	GTG	GAG	GGG	GAG	CTC	TGC	CAT	GAC	CCC	ACC	AGC	CAG	CTG	CTG	GAT	CTC	ATC	865
T	W	E	L	E	P	E	G	A	L	D	R	C	P	C	A	S	G	L	L	272
ACC	TGG	GAA	CTG	GAG	CCT	GAA	GGA	GCT	TTG	GAC	CGA	TGC	CCC	TGC	GCC	AGT	GGC	CTC	CTA	925
C	Q	P	H	S	H	S	L	V	Y	M	C	K	P	A	F	V	G	S	H	292
TGC	CAG	CCA	CAC	AGC	CAC	AGT	CTG	GTG	TAC	ATG	TGC	AAG	CCA	GCC	TTC	GTG	GGC	AGC	CAT	985
D	H	S	E	E	S	Q	P	R	E	A	P	D	E	Y	E	D	V	G	312	
GAC	CAC	AGT	GAG	GAG	AGC	CAG	CTG	CCC	AGG	GAG	GCC	CCG	GAT	GAG	TAC	GAA	GAT	GTT	GGC	1045
F	I	G	E	V	R	Q	E	L	E	D	L	E	R	S	L	A	Q	E	M	332
TTC	ATA	GGG	GAA	GTG	CGC	CAG	GAG	CTG	GAA	GAC	CTG	GAG	CGG	AGC	CTA	GCC	CAG	GAG	ATG	1105

Figure 8B

12/12

FIGURE 9



INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/07894

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C07K16/18 G01N33/50 G01N33/53
C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^o	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL database entry HS266245; accession number H99266; 16. Dec. 1995; Hillier et al.: 'The WashU-Merck EST Project.' XP002077278 see abstract ---	1,16-18
X	EMBL database entry HSAA53464; accession number AA253464; 15. March 1997; Hillier et al.: The WashU-Merck EST Project.' XP002077279 see abstract --- -/-	1,16-18

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

^o Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

16 September 1998

25/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mandl, B

INTERNATIONAL SEARCH REPORT

Int'l. Application No

PCT/US 98/07894

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EMBL database entry HSA37322; accession number AA037322; 28. Aug. 1996; Hillier et al.: 'The WashU-Merck EST project.' XP002077280 see abstract ---	1,16-18
X	EMBL database entry HS979350; accession number W55979; 6. June 1996; Hillier et al.: 'The WashU-Merck EST Project.' XP002077281 see abstract ---	1,16-18
A	US 5 525 486 A (HONJO TASUKU ET AL) 11 June 1996 see figure 1 ---	1-22
A	KLEIN R. D. ET AL.: "Selection for genes encoding secreted proteins and receptors." PROC. NATL. ACAD. SCI. USA, vol. 93, July 1996, pages 7108-7113, XP002077277 see the whole document ---	1-22
P, X	EMBL database entry AF034208; accession number AF034208; 6. Jan. 1998; Ligon A.H. et al.: 'Homo sapiens RIG-like 7-1 mRNA.' XP002077282 cited in the application see abstract ---	1,3,5,8, 12
P, X	EMBL database entry AA565546; accession number AA565546; 11. Sept. 1997; Robert Strausberg: 'NCI, Cancer Genome Anatomy Project.' XP002077283 see abstract -----	1,16-18

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/07894

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

REMARK: The search for claims 13, 15 and 21 was limited to an antibody as being the 'compound which selectively binds to a polypeptide of claim 8' because no other compounds are sufficiently specified in the present application.
3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking(Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT**Information on patent family members**International Application No
PCT/US 98/07894

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
US 5525486	A 11-06-1996	CA	2113363 A	15-07-1994
		EP	0607054 A	20-07-1994
		JP	6315380 A	15-11-1994