



(22) Date de dépôt/Filing Date: 1992/02/11
 (41) Mise à la disp. pub./Open to Public Insp.: 1992/08/14
 (45) Date de délivrance/Issue Date: 2004/01/27
 (30) Priorité/Priority: 1991/02/13 (654,480) US

(51) Cl.Int.⁵/Int.Cl.⁵ C07F 7/00, A61K 7/38
 (72) Inventeurs/Inventors:
 SHIN, CHUNG T., US;
 ROSENBERG, ALLAN H., US;
 MARKARIAN, HERAND M., US
 (73) Propriétaire/Owner:
 SIGNAL INVESTMENT & MANAGEMENT CO., US
 (74) Agent: GOWLING LAFLEUR HENDERSON LLP

(54) Titre : COMPOSE ANTISUDORIFIQUE ZIRCONIUM/ALUMINIUM, ET COMPOSITIONS ANTISUDORIFIQUES EN
 CONTENANT
 (54) Title: ZIRCONIUM/ALUMINUM ANTIPERSPIRANT MATERIAL AND ANTIPERSPIRANT COMPOSITIONS
 CONTAINING SAME

(57) **Abrégé/Abstract:**

A novel ZAG complex and a process for making same, the process comprising mixing zirconium hydroxychloride, an aluminum chloro species of the formula $Al_2(OH)_{6-x}Cl_x$ wherein $x = 1.5$ to 6 and an amino acid in aqueous solution and, optionally, drying said aqueous solution to obtain a dry ZAG salt. The invention further includes antiperspirant compositions and products containing the ZAG complex and an activated aluminum chlorhydroxide, and processes for the manufacture of said compositions.



ABSTRACT OF THE DISCLOSURE

A novel ZAG complex and a process for making same, the process comprising mixing zirconium hydroxychloride, an aluminum chloro species of the formula $Al_2(OH)_{6-x}Cl_x$ wherein $x = 1.5$ to 6 and an amino acid in aqueous solution and, optionally, drying said aqueous solution to obtain a dry ZAG salt. The invention further includes antiperspirant compositions and products containing the ZAG complex and an activated aluminum chlorhydroxide, and processes for the manufacture of said compositions.

[BP-8906A]

Background of the Invention

The preparation of aluminum- and zirconium-based antiperspirant materials generally involves combining compounds of these metals with an amino acid under suitable conditions. Such materials are generally referred to as ZAG complexes and have often been shown to be chemically distinct species having antiperspirant efficacy, i.e., the inhibition of perspiration when applied to human skin, especially the axilla.

In U.S. Patent 4,775,528 to Callaghan et al, an antiperspirant composition is produced by heating a 2 to 18% aqueous solution of aluminum chlorhydroxide to at least 50°C, mixing zirconyl hydroxychloride with the solution before, during or after heating, continuing heating of the aqueous mixture until prescribed characterizations are achieved, and thereafter rapidly drying the heated aqueous mixture to obtain a solid material having an Al:Zr atomic ratio of from 6:1 to 1:1 and also having peak 4 and peak 3 Al chromatograph peaks, the ratio of peak 4 to peak 3 being at least 2:1. Patentees specify in Claim 1 that the zirconyl hydroxychloride be mixed with the aluminum chlorhydroxide solution before the solution loses all of its water, i.e., "before completion of . . . drying".

U.S. 4,818,512 to Markarian et al -- which is assigned to applicants' assignee -- describes the production of an improved activated aluminum chlorhydroxide material from a conventional activated aluminum chlorhydroxide material, the improved material being characterized by containing a high concentration (77 to 85% by weight) of the K_d 0.4 fraction. The improved aluminum chlorhydride of the '512 patent is prepared by heating a 5% aqueous solution of the conventional aluminum chlorhydroxide material under certain conditions and spray-drying the product. The material made in the '512 patent may be combined with zirconium-containing species to yield zirconium-aluminum-glycine (ZAG) complexes or compounds, which are highly efficacious as antiperspirants.

The Invention

It has been discovered that, under certain conditions, solutions of certain aluminum chloro species (as hereinafter described) and zirconium hydroxychloride, i.e., $ZrO(OH)Cl$, can be admixed in the presence of an amino acid, as to yield a ZAG material, the solid form of the material being obtained by conventional drying means, such as by steam bath evaporation, spray-drying and freeze-drying. The ZAG material is believed to be a new chemical entity having a unique chemical formula. Antiperspirant compositions having superior antiperspirant properties are obtained when the ZAG material is formulated with activated aluminum chlorhydroxide, $Al_2(OH)_5Cl$ (hereinafter also referred to as "ACH"), as hereinafter described.

This new material is a zirconium-aluminum-glycine (ZAG) or Al/Zr polychlorohydroxide-glycine material, which is referred to herein variously as a "ZAG compound", "ZAG material", "ZAG complex" or "ZAG salt". For simplicity, applicants will generally refer to the dried ZAG material as a ZAG salt. The term "ZAG complex" is often used by practitioners of this art, but in a general sense, since the mechanism by which the ZAG material is formed is not well understood. The ZAG material is a complex in the sense that the various constituents interact in a chemical or physical manner to provide the subject ZAG species.


By "aluminum chloro species" applicants mean aluminum chloride and aluminum hydroxy chloro species defined by the formula $Al_2(OH)_{6-x}Cl_x$ wherein $x = 1.5$ to 6 , in particular, $2/3$ basic aluminum chloride, $Al_2(OH)_4Cl_2$, and $3/4$ basic aluminum chloride, $Al_2(OH)_{4.5}Cl_{1.5}$. Thus, when $x = 6$ the salt is aluminum chloride, $AlCl_3$, which salt may be hydrated with up to 6 moles water.

As used herein, "antiperspirant composition" means a composition obtained by dry-blending the ZAG salt of the present invention and an activated aluminum chlorhydroxide or, alternatively, by mixing a solution of the ZAG material and a solution of an activated aluminum chlorhydroxide, and then rapidly drying

same. As used herein, "antiperspirant product" means the antiperspirant composition incorporated into a cosmetically acceptable carrier or vehicle, as hereinafter described.

In this specification and in the claims, unless otherwise indicated, all percents are percent by weight and all parts are parts by weight.

The ZAG complex of the present invention is formed by preparing an aqueous solution of $ZrO(OH)Cl$, the aluminum chloro species and the amino acid, and, preferably, heating said solution. The ZAG salt is obtained thereafter by drying the solution. In a preferred embodiment of this process, 66.8 parts of a nominal 50% $ZrO(OH)Cl$ solution, 19.4 parts of a 50% $AlCl_3$ solution and 13.8 parts of glycine were heated together on a steam bath and dried, to yield a dry powder (ZAG salt) containing on assay 26.6% zirconium, 2.0% aluminum, 24.4% glycine (percent glycine calculated by difference) and 18.1% chloride. A ZAG salt containing 25.5% Zr, 1.9% Al, 23.6% glycine and 17.5% chloride -- which values are only slightly less than the assayed ZAG salt made by the described procedure -- has an empirical formula of



wherein $n = 0-9$, preferably 2-7.

An antiperspirant composition containing the ZAG salt and an ACH is obtained by dry-blending of these two antiperspirant active materials. An antiperspirant composition was obtained by dry-blending 48 parts of the above-described ZAG salt with 52 parts ACH powder. The resulting antiperspirant composition contained 13.3% Al, 12.4% Zr and 16.8% Cl and had an Al/Zr atomic ratio of 3.6:1 and a metal/chloride ratio of 1.3:1.

Similarly, a solution containing 66.8 parts of a nominal 50% $ZrO(OH)Cl$ solution, 19.4 parts of a 50% $AlCl_3$ solution and 13.8 parts glycine was spray-dried to yield a ZAG salt containing on assay 28.7% Zr, 2.1% Al, 26.6% glycine and 19.4% chloride. The empirical formula was similar to that given above. This ZAG salt was blended or mixed in a dry state with dry ACH to yield a highly useful antiperspirant composition.

In another embodiment of the present invention, the antiperspirant composition is made by first preparing an aqueous solution of $ZrO(OH)Cl$, the aluminum chloro species and the amino acid, preferably by mixing the solution at about room temperature to obtain the ZAG material, combining said solution containing the ZAG material with a solution of ACH preferably at about room temperature, and thereafter rapidly drying the solution.

Advantages

The compounds and compositions of the invention have several advantages over other antiperspirants. The ZAG/ACH combinations made in accordance with the invention are significantly more efficacious than are commercially available tetra ZAG salts.

In addition, the ZAG salts made in accordance with the process of the invention are characterized by their

- (1) Ease of manufacture,
- (2) More uniform chemical properties, and
- (3) Good stability during the mixing process.

These and other advantages and aspects of the invention will be apparent after a consideration of the following description and claims.

Detailed Description of the Invention

The description of the invention will focus on discussions of the new ZAG salt, the ACH with which it is combined, and the final antiperspirant compositions.

ZAG Ingredient (Component I)

The new ZAG compounds of the present invention are aluminum-zirconium polychlorohydrate-amino acid materials. Glycine is the preferred amino acid. The description herein shall refer, for simplicity, to glycine when generally describing the amino acid, except when the context indicates otherwise, as in the illustrative examples and specific embodiments.

In the past Al-Zr-glycine complexes have been made by heating an aqueous solution of aluminum chlorohydrate with zirconium and glycine for appropriate periods of time. Previously cited U.S. Patent 4,775,528 to Callaghan et al disclosed such a process.

The ZAG complexes of the present invention are polymeric species which have proportions of zirconium, aluminum, glycine and chloride as set forth in Table I. The atomic ratio of Zr:Al is in the range of about 6:1 to about 2.75:1.

The ZAG complexes of the present invention are made by preparing an aqueous solution of zirconium hydroxychloride

(ZrO(OH)Cl); an aluminum chloro species; an amino acid, for example, glycine, lysine and alanine and heating the solution. The aqueous solution contains from about 25 to 40%, preferably from 30 to 37% ZrO(OH)Cl, from about 8 to 12%, preferably from about 9 to 11% AlCl₃, and from 11 to 17%, preferably from about 12 to 15% amino acid.

The individual components may be added as dry powders to water to form the aqueous solution. Preferably, premixes of ZrO(OH)Cl and AlCl₃ are employed, with amino acid being added to either premix or to the combined premix solutions.

Commercially available ZrO(OH)Cl solutions (i.e., 25 to 50% aqueous ZrO(OH)Cl are especially useful. Basic zirconyl chloride or zirconyl hydroxychloride available from Westwood Chemical (Middletown, New York) and Magnesium Electron, Inc. can be used. Mixtures are operable. As previously described, the aluminum chloro species component of the ZAG material is a salt of the general formula Al₂(OH)_{6-x}Cl_x wherein x = 1.5-6, for example, AlCl₃, 2/3 and 3/4 basic aluminum chlorides. The aluminum chloro species are typically available commercially and used as aqueous solutions. Aluminum chloride is the preferred aluminum chloro species, as this salt is the most acidic of those included in the general formula. Aqueous aluminum chloride solutions having a 50% active concentration are available from Reheis Chemical Co. These

commercially available zirconium and aluminum salt solutions are provided as "nominal" solutions, e.g., a 50% solution contains approximately 50% of the active ingredient.

The zirconium-containing solution is combined with the aluminum chloride solution via simple mixing at a temperature in the range of from about 70 up to about 240°F, preferably from about 70 to about 140°F. Generally, mixing time is not critical, and is completed in under 15 minutes, preferably within 10 minutes, the time required for mixing depending on the size, temperature and degree of agitation of the batch. Longer mixing times are possible, but are not economically practical. Heating of the solution when elevated temperature is used may be conducted by conventional means such as a steam bath, steam jacket and closed loop circulation through a heat exchanger. At higher temperatures system pressures would be such as to avoid boiling of the aqueous solution and to ensure suitable control of the process. The pressure can be determined easily by using steam tables. It is believed that the ZAG complex is formed during this mixing procedure. However, applicants do not wish to be bound to any particular theory regarding complexation or salt formation.

The third component is a commercial amino acid typically of 99 to 100 percent purity, preferably glycine. Suitable glycines are glycine crystals made by Robeco Company of New York and glycine

powders made by Chattem Company of Tennessee. Other amino acids, e.g., alanine, glycine and the like, are also suitable.

It is convenient to form aqueous premixes as of the $ZrO(OH)Cl$ and the aluminum chloro constituents, and thereafter to combine these solutions, along with the glycine component. The glycine component may be added to either premix or to the combined premix solutions. An aqueous premix of the glycine component is also acceptable. Alternatively, an aqueous solution of $ZrO(OH)Cl$ and the aluminum chloro species can be prepared based on the use of the dry materials, although this is not preferred. Also, any one of the dry components can be added to an aqueous premix of another component.

As previously mentioned, aqueous solutions of $ZrO(OH)Cl$ and the aluminum chloro species are commercially available and may be used as the premix solutions with or without dilution. The use of premix solutions is preferred for ease of handling.

After completion of mixing, the ZAG salt, i.e., the dry ZAG complex, can be obtained by drying the solution to substantial dryness. The time to effect drying is not critical, and will depend on the size of the batch, the temperature and pressure of the mixing vessel, and the drying equipment used. It is preferable to effect drying rapidly, which can be accomplished best by

spray-drying techniques well known in the art. A portion of the water in the aqueous solution may evaporate during the mixing step when elevated temperatures are used. Drying methods include those which employ a steam bath, spray-drying, freeze-drying, etc. Drying is carried out until an essentially powdery state is reached.

The process described above may be conducted by batch, semi-batch or continuous techniques. Typically, a semi-batch process is employed with preparation of a given quantity of the solution mixture containing the components, with direct feed to a spray drying chamber. In this case drying is rapid, especially when elevated temperatures are used. The solution mixture may be prepared by simultaneously charging the mixture vessel with premix solutions.

Once the powder has been obtained, it may be ground (e.g., via jet-milling or other commercial process) to yield an impalpable powder for storage, shipment, or further processing. Typically, 97% of this powder passes through a 325 mesh screen, and provides good organoleptic properties when provided in an antiperspirant product.

Table I sets forth a typical assay of the ZAG salt (Component I) obtained by the process of the present invention.

TABLE I
COMPOSITION OF ZAG INGREDIENT
(COMPONENT I)

	<u>Broad Range</u>	<u>Preferred Range</u>	<u>Highly Preferred</u>
% Zr	23.0-34.7	25.8-31.6	28.7
% Al	1.7- 2.5	1.9- 2.3	2.1
% Amino Acid	21.3-31.9	23.9-29.2	26.6
% Chloride	15.5-23.3	17.5-21.3	19.4

While the exact nature of the ZAG salt is not readily ascertained, the ZAG salt, based on a theoretical correlation of the assays made on the material, generally conforms to the empirical formula



wherein n = 0 to 9.

Activated ACH (Component II)

The ZAG compound may be employed to form an improved antiperspirant composition also containing an activated aluminum chlorhydroxide (ACH) component. ACH components are well known in the art, as described, for example, in Antiperspirants and Deodorants, K. Laden (Editor), pp. 214-8 (1989).

See also U.S. 3,904,741 to Jones et al.

One highly useful ACH is described in the previously cited U.S. Patent 4,818,512 to Markarian et al.

The preferred ACH component of the antiperspirant composition typically has the following approximate distribution for the Al species:

TABLE II
PERCENT ALUMINUM PEAK HEIGHTS FOR ACH

	$K_d = 0$	$K_d = 0.25$	$K_d = 0.4$	$K_d = 0.6$
Preferred Range	0-3	25-35	50-60	5-20

Commercially available ACH powders are available as the products sold under the trade names DM200 or DM250 by Westwood Chemical (Middletown, New York), RE101 and RE103 by Reheis (Berkely, New Jersey) and Q5-7171 (AAH) by Dow Corning (Midland, Michigan).

Antiperspirant (ZAG/ACH) Combination Compositions

The ZAG salt and the ACH components, discussed above, may be combined, undiluted with carrier(s), via simple mixing in a substantially dry state to obtain a homogeneous blend, typically

for between 30 to 120 minutes, preferably for between 60 to 90 minutes, to yield a highly efficacious antiperspirant composition. The weight ratio of ZAG salt to ACH present in the antiperspirant composition is from about 2:1 to about 1:2, preferably from about 1.5:1 to about 1:1.5, most preferably about 1:1. A typical antiperspirant composition contains 13.3% Al, 12.4% Zr, 16.8% Cl and 11.5% glycine, and has an Al/Zr atomic ratio of about 3.6:1 and a metal/chloride atomic ratio of 1.3:1.

In another embodiment of the present invention, the antiperspirant composition is prepared by first forming the aqueous solution containing the $ZrO(OH)Cl$, the aluminum chloro species and the glycine constituents, combining said solution with the activated aluminum chlorhydroxide, and rapidly thereafter drying said solution to form the solid antiperspirant composition. In this embodiment it is important that the mixing of the constituents be conducted at lower temperatures, typically less than about 125°F, and preferably at ambient. The weight ratio of the ZAG salt to the ACH present in the antiperspirant composition is as previously described. The assay and Al/Zr and metal/chloride atomic ratios for the antiperspirant composition in accordance with this process embodiment are also as previously described.

In the process described in the paragraph above, the time in which the ACH and the ZAG material are present together in the

in the aqueous phase should not be excessive, in order to obtain a final antiperspirant composition having a high proportion of K_d 0.4 for the aluminum species. Accordingly, the preparation of the combined ZAG-ACH solution and the drying process should both preferably be effected rapidly, within the practical time limits imposed by conventional processing equipment. The combined ZAG complex-ACH solution is typically ready for processing in the drying equipment as soon as an essentially homogeneous mixture is obtained. Generally, this is within about 5 to 20 minutes, preferably within 5 minutes. Drying should also be completed rapidly. In general, using a shorter mixing residence time provides greater leeway for the time in which to carry out the drying procedure. Spray-drying provides essentially instantaneous drying, and accordingly is highly preferred. The mixing and drying steps, in total, should not exceed about 30 minutes, and preferably are effected in less than about 15 minutes.

The final dry antiperspirant composition obtained in accordance with the embodiments of the present invention was analyzed using the gel chromatographic technique set forth below. The ZAG/ACH antiperspirant composition of the invention has the following Al species K_d distribution for the Al species:

TABLE III

% Al K_d PEAK HEIGHTS FOR ZAG/ACH COMBINATIONS
 $K_d = 0$ $K_d = 0.25$ $K_d = 0.4$ $K_d = 0.6$

	$K_d = 0$	$K_d = 0.25$	$K_d = 0.4$	$K_d = 0.6$
Broad Range	0-5	30-40	40-60	5-20
Preferred Range	0-3	25-35	50-60	5-20

In general, combinations in which K_d 0.4 for 50% or more of the Al peaks are preferred.

Polymer Characterization Techniques

The character of the ACH and of the ZAG/ACH compositions was determined using the following procedure.

Aluminum species distribution for ACH and ZAG/ACH mixtures are obtained from gel filtration chromatography. The principles of gel filtration are discussed in Gel Filtration Theory and Practice published by Pharmacia Fine Chemicals, Uppsala, Sweden (November 1981-82).

As explained in Gel Filtration Theory and Practice, K_d is the distribution coefficient, which is defined by the equation

$$K_d = (V_e - V_o) / V_s$$

where

V_e = elution volume

V_o = void volume

V_s = volume of the stationary phase

Generally, the gel chromatograph measures K_{av} , defined by the equation

$$K_{av} = (V_e - V_o) / (V_t - V_o)$$

where

V_t = total volume

K_{av} represents the fraction of the stationery gel volume which is available for diffusion of a given solute species. For a given gel, there is a constant ratio $K_{av}:K_d$ which is independent of the nature of the solute or its concentration. In the characterizations of the active antiperspirant components and compositions, the gel chromatography technique measures K_{av} . Measurements using the gel chromatography techniques herein described and used in the characterization of the antiperspirant actives, in fact, measure the parameter K_{av} . However, because the ratio $K_{av}:K_d$ is close to unity, it is common practice in the antiperspirant art to use these parameters interchangeably.

Metal analysis of the separated species eluting off the column are obtained by ICP spectroscopy. The following conditions are employed.

Column : Sephadex* G-25 resin (superfine) 50 cm
manufactured by Pharmacia Fine Chemicals
Eluent : 0.2M KCl/0.0035 M HCl
Sample Size : 100 μ l of a freshly prepared 13% aqueous
solution of activated ACH or ZAG/ACH
antiperspirant composition
Flow Rate : 0.4 ml/min.
RI Detector : Shimatzu RID 6A
Metal Analyzer: Perkin Elmer ICP 6500

Other Ingredients

The ZAG salt and the antiperspirant compositions of the present invention may be incorporated into a cosmetically acceptable carrier or vehicle to provide commercially useful antiperspirant products.

The antiperspirant products of the invention may contain a wide variety of materials conventionally employed in the manufacture of such products.

*Trade mark

Excipients such as fillers, diluents, coupling agents, pH modifiers, buffers, colorants (e.g., opacifiers, whiteners), perfumes, flow-control agents (e.g., anticaking agents), thickeners and the like and mixtures thereof can be used in suitable quantities in the compositions of the invention, as is well known in the art. Suitable excipients and their concentrations in an antiperspirant product are described in U.S. 4,781,917 to Leubbe et al, U.S. 4,774,079 to Shin et al and U.S. 4,083,956 to Shelton.

The antiperspirant products of the invention may be solid, semisolid or liquid in form. When desired, they may be made into gels, pastes, creams, foams, or aerosol liquids using suitable amounts of diluents or other extenders. The formulations are contemplated for use in deodorant/antiperspirant products which are applied as sticks, sprays, roll-ons, creams, lotions, and the like.

The following examples illustrate the invention.

Example 1: Preparation of ZAG salt (Component I)

This example illustrates the preparation of ZAG salts. Solution A below was prepared by admixing the recited ingredients.

Solution A

<u>Ingredient</u>	<u>(w/w%)</u>
ZrO(OH)Cl (50% aqueous solution)	66.8
AlCl ₃ (50% aqueous solution)	19.4
Glycine	<u>13.8</u>
	100.00

Solution A was heated at about 180°F with gentle stirring over a steam bath until a highly thickened, aqueous residue remained, after which the residue was placed in an oven until a powdery solid was obtained.

All dilute solutions used contained water as the diluent. The resultant dried ZAG ingredient (Component I) powder was analyzed as follows.

<u>Ingredient</u>	<u>Theoretical Amount</u>	<u>Assay Amount</u>	<u>% of Theoretical</u>
Zirconium	25.5	26.6	104
Aluminum	1.9	2.0	105
Chloride	17.5	18.1	103

Example 2

The ZAG salt of Example 1 was dry-mixed with an ACH powder obtained from Reheis (RE101) to provide an antiperspirant composition containing 48% ZAG salt and 52% ACH. The ACH was characterized as having a K_d 0.4 concentration of 59%. The dry-mixing was conducted in a twin shell blender for about one hour. This resulted in the final ZAG/ACH antiperspirant composition.

Example 3

Solution A (from Example 1) was spray-dried to obtain the ZAG salt (Component I) in lieu of oven-drying. The resulting ZAG was dry-blended with ACH using the procedure of Example 2 to obtain a final ZAG/ACH antiperspirant composition. Analytical results on these systems were similar to those obtained in Example 1. Spray-drying is the preferred method used to prepare ZAG ingredient (I) powders, rather than evaporation using a steam bath or oven-drying.

The antiperspirant composition was incorporated into a vehicle, the antiperspirant product containing the following ingredients:

<u>Antiperspirant Product</u>	<u>Wt. %</u>
Antiperspirant composition	24.00
Cyclomethicone 7158	53.05
Stearyl alcohol	13.00
Hydrogenated castor oil	4.00
PEG-25 Propylene glycol stearate	1.00
Glyceryl monostearate	0.35
Butylated hydroxytholuene	0.05
Talc	4.00
Fumed silica	0.50
Pentadecalactone	<u>0.05</u>
	100.00

Three clinical studies comparing this ZAG/ACH antiperspirant product against Dow Corning 369 (normal tetra ZAG salt) in BAN^R Solid were conducted. The tests showed that the combination of this invention was 9.7% drier than Dow Corning 369 in the first study, 6.9% drier in the second study, and 8.6% drier in the third study.

Example 4 (Comparative)

100 g of Solution A (Example 1) was mixed with 61.66 g of RE101 ACH powder, heated for 90 minutes on a steam bath and dried in an oven as in Example 1. The resultant powder had a K_d 0.4 value of 17%, which is unacceptably low.

Example 5

The antiperspirant composition was prepared by the alternate process of the present invention, as follows:

66.8 g of a 50% aqueous $ZrO(OH)Cl$ solution was mixed with 19.4 g of a 50% aqueous aluminum chloride solution and 13.8 g glycine at room temperature. This solution, which had a ZAG material concentration of 13.8%, was combined with 100 g of a 15.0% aqueous solution of activated ACH in a common vessel. Immediately, the combined solutions were spray-dried to produce a ZAG/ACH antiperspirant composition in powder form.

The concentrations of ingredients in the ZAG and ACH solutions can be adjusted to produce the final ZAG/ACH ratios discussed above under "Antiperspirant (ZAG/ACH) Combination Compositions."

Reasonable variations, such as those which would occur to a skilled artisan, can be made herein without departing from the scope of the invention.

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE AS FOLLOWS:

1. A process for making a zirconium-aluminum-amino acid (ZAG) material comprising admixing zirconium hydroxychloride, an aluminum chloro species of the formula $\text{Al}_2(\text{OH})_{6-x}\text{Cl}_x$ wherein $x = 1.5$ to 6 and an amino acid selected from the group consisting of glycine, lysine and alanine in an aqueous solution at from about 70 to 240°F , whereby the ZAG material is obtained in solution, said ZAG material following drying essentially to dryness having the following analysis on a weight basis: 23.3 - 34.7% Zr, 1.7 - 2.5% Al, 21.3 - 32.9% amino acid and 15.5 - 23.3% chloride, and being bound with zero to nine moles water per mole of the ZAG complex.
2. The process of Claim 1 wherein the aqueous solution initially contains from about 25 to 40% zirconium hydroxychloride, from about 8 to 12% aluminum chloro species and from about 11 to 17% amino acid.
3. The process of Claim 1 wherein the aluminum chloro species is selected from the group consisting of AlCl_3 , $\text{Al}_2(\text{OH})_4\text{Cl}_2$ and $\text{Al}_2(\text{OH})_5\text{Cl}_{1.5}$; the amino acid is glycine, and the dry ZAG material contains on assay 25.8 - 31.6% Zr, 1.9 - 2.3% Al, 23.9 - 29.2% glycine, and 17.5 - 21.3% chloride.
4. The process of Claim 1 wherein the aluminum chloro species is selected from the group consisting of AlCl_3 , $\text{Al}_2(\text{OH})_4\text{Cl}_2$ and $\text{Al}_2(\text{OH})_5\text{Cl}_{1.5}$; the amino acid is glycine, and the dry ZAG material contains about 28.7% Zr, about 2.1% Al, about 26.6% glycine and about 19.4% chloride.

5. The process of Claim 1 wherein the aluminum chloro species is selected from the group consisting of AlCl_3 , $\text{Al}_2(\text{OH})_4\text{Cl}_2$ and $\text{Al}_2(\text{OH})_{4.5}\text{Cl}_{1.5}$ and the amino acid is glycine.

6. The process of Claim 1 wherein amino acid is glycine and the dry ZAG material has the empirical formula



wherein $n = 0$ to 9.

7. The process of Claim 1, 3 or 6 further comprising the step of drying the aqueous ZAG solution until a dry ZAG salt is obtained.

8. The process of Claim 7 wherein the aluminum chloro species is AlCl_3 .

9. The process of Claim 3, 4 or 6 wherein the aluminum chloro species is AlCl_3 .

10. The process of Claim 9 further comprising the step of drying the aqueous ZAG solution until a dry ZAG salt is obtained.

11. The process of Claim 7 wherein the aqueous ZAG solution is spray-dried.

12. The process of Claim 10 wherein the aqueous ZAG solution is spray-dried.

13. A zirconium-aluminum-amino acid ZAG salt containing on a weight basis: 23.0-34.7% Zr, 1.7-2.5% Al, 21.3-31.9% amino acid selected from the group consisting of glycine, lysine and alanine, and 15.5-23.3% chloride, said salt being bound with zero to nine moles water per mole of the zirconium-aluminum-amino acid salt.

14. A zirconium-aluminum-amino acid ZAG salt containing on a weight basis: 25.8-31.6% Zr, 1.9-2.3% Al, 23.9-29.2% amino acid selected from the group consisting of glycine, lysine and alanine, and 17.5-21.3% chloride, the salt being bound with zero to nine moles water per mole of the zirconium-aluminum-amino acid salt.

15. The ZAG salt of Claim 13 or 14 wherein the amino acid is glycine.

16. A zirconium-aluminum-glycine ZAG salt having the empirical formula



wherein $n = 0$ to 9.

17. A zirconium-aluminum-amino acid salt made by the process of Claim 11.

18. A zirconium-aluminum-amino acid salt made by the process of Claim 12.

19. A solid antiperspirant composition comprising

a zirconium-aluminum-amino acid (ZAG) salt containing on a weight basis 23.0-34.7% Zr, 1.7-2.5% Al, 21.3-31.9% amino acid selected from the group consisting of glycine, lysine and alanine, and 15.5-23.3% chloride, the ZAG salt being bound with zero to nine moles water per mole of the ZAG salt, and

an activated aluminum chlorhydroxide

wherein the weight ratio of ZAG salt to activated aluminum chlorhydroxide is from about 2:1 to about 1:2,

more than 40% of the Al peaks as determined by gel chromatography of the antiperspirant composition having a K_d value of 0.4.

20. The composition of Claim 19 wherein the ZAG salt contains 25.8-31.6% Zr, 1.9-2.3% Al, 23.9-29.2% amino acid, and 17.5-21.3% chloride.

21. The composition of Claim 19 wherein the amino acid is glycine and the ZAG salt has the empirical formula:



wherein $n = 0$ to 9.

22. The composition of Claim 19, 20 or 21 wherein the weight ratio of the ZAG salt to the activated aluminum chlorhydroxide is from 2:1 to 1:2.

23. The composition of Claim 22 wherein the weight ratio of the ZAG salt to the activated aluminum chlorhydroxide is from 1.5:1 to 1:1.5.

24. A process for making an antiperspirant composition, the process comprising the step of mixing, in a dry state, the ZAG salt of Claim 13 and an activated aluminum chlorhydroxide,

wherein the weight ratio of ZAG salt to activated aluminum chlorhydroxide is from about 2:1 to about 1:2,

more than 40% of the Al peaks as determined by gel chromatography of the antiperspirant composition having a K_d value of 0.4.

25. A process for making an antiperspirant composition, the process comprising the steps of preparing at less than about 125°F the aqueous solution containing the ZAG material, as set forth in Claim 1;

admixing an activated aluminum chlorhydroxide and said ZAG solution, wherein the weight ratio of ZAG salt to activated aluminum chlorhydroxide is from about 2:1 to about 1:2, and

rapidly drying said admixture,

more than 40% of the aluminum peaks as determined by gel chromatography of the antiperspirant composition having a K_d value of 0.4.

26. The process of Claim 24 or 25 wherein the weight ratio of the ZAG material to the activated aluminum chlorhydroxide is from 2:1 to 1:2.

27. The process of Claim 26 wherein the weight ratio of the ZAG material to the activated aluminum chlorhydroxide is from 1.5:1 to 1:1.5.

28. The process of Claim 25 wherein the solution of the ZAG material is prepared at about room temperature and the admixture is dried by spray-drying.

29. The process of Claim 24 or 25 wherein the ZAG salt is defined by Claim 14.
30. The process of Claim 29 wherein the ZAG salt is defined by Claim 16.
31. An antiperspirant composition made by the process of Claim 25.
32. An antiperspirant product comprising the antiperspirant composition of Claim 19, in a cosmetically acceptable carrier or vehicle.
33. The use of a compound or composition according to any of Claims 13 to 23, 31 or 32, as an antiperspirant.