



(86) Date de dépôt PCT/PCT Filing Date: 2005/09/02

(87) Date publication PCT/PCT Publication Date: 2006/03/16

(45) Date de délivrance/Issue Date: 2013/01/15

(85) Entrée phase nationale/National Entry: 2007/03/01

(86) N° demande PCT/PCT Application No.: US 2005/031327

(87) N° publication PCT/PCT Publication No.: 2006/028967

(30) Priorité/Priority: 2004/09/02 (US60/606,855)

(51) Cl.Int./Int.Cl. *C12N 15/11* (2006.01),
A61K 31/713 (2006.01), *A61K 48/00* (2006.01),
A61P 35/00 (2006.01)

(72) Inventeurs/Inventors:
SLACK, FRANK J., US;
JOHNSON, STEVEN M., US;
GROSSHANS, HELGE, US

(73) Propriétaire/Owner:
YALE UNIVERSITY, US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : REGULATION D'ONCOGENES PAR DES MICRO-ARN

(54) Title: REGULATION OF ONCOGENES BY MICRORNAS

(57) **Abrégé/Abstract:**

Naturally occurring miRNAs that regulate human oncogenes and methods of use thereof are described. Suitable nucleic acids for use in the methods and compositions described herein include, but are not limited to, pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of the mature miRNA and DNA encoding a pri-miRNA, pre-miRNA, mature miRNA, fragments or variants thereof, or regulatory elements of the miRNA. The compositions containing nucleic acids are administered to a patient in need of treatment or prophylaxis of at least one symptom or manifestation of cancer. In one embodiment, the compositions are administered in an effective amount to inhibit gene expression of one or more oncogenes. Methods for treatment or prevention of at least one symptom or manifestation of cancer are also described.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 March 2006 (16.03.2006)

PCT

(10) International Publication Number
WO 2006/028967 A3

(51) International Patent Classification:

C12N 15/11 (2006.01) *A61K 48/00* (2006.01)
A61K 31/713 (2006.01) *A61P 35/00* (2006.01)

(21) International Application Number:

PCT/US2005/031327

(22) International Filing Date:

2 September 2005 (02.09.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/606,855 2 September 2004 (02.09.2004) US

(71) Applicant (for all designated States except US): **YALE UNIVERSITY** [US/US]; 451 College Street, New Haven, CT 06511 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SLACK, Frank, J.** [US/US]; 111 grist Mill Circle, Guilford, CT 06437 (US). **JOHNSON, Steven, M.** [US/US]; 1103 King Street, Redwood City, CA 94061 (US). **GROSSHANS, Helge** [DE/US]; 137 Cottage Street, Apartment E1, New Haven, CT 06511 (US).

(74) Agents: **PABST, Patrea, L.** et al.; Pabst Patent Group LLP, 1201 Peachtree Street, Suite 1200, Atlanta, GA 30361 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report

(88) Date of publication of the international search report:

22 February 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: REGULATION OF ONCOGENES BY MICRORNAS

(57) Abstract: Naturally occurring miRNAs that regulate human oncogenes and methods of use thereof are described. Suitable nucleic acids for use in the methods and compositions described herein include, but are not limited to, pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of the mature miRNA and DNA encoding a pri-miRNA, pre-miRNA, mature miRNA, fragments or variants thereof, or regulatory elements of the miRNA. The compositions containing nucleic acids are administered to a patient in need of treatment or prophylaxis of at least one symptom or manifestation of cancer. In one embodiment, the compositions are administered in an effective amount to inhibit gene expression of one or more oncogenes. Methods for treatment or prevention of at least one symptom or manifestation of cancer are also described.



WO 2006/028967 A3

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

REGULATION OF ONCOGENES BY MICRORNAS
CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to U.S. Serial No. 60/606,855 filed September 2, 2004 entitled "*Regulation of Oncogenes by MicroRNAs*" by Frank J. Slack, Steven M. Johnson, Kristy L. Reinert, and Helge Grosshans.

STATEMENT REGARDING FEDERALLY FUNDED RESEARCH

The United States Federal Government has certain rights in this invention by virtue of Grant No. 1R01GM062594-01A1 and Grant No. 1R01GM064701-01 from the National Institute of Health to Frank J. Slack.

BACKGROUND OF THE INVENTION

Cancer is a group of diseases characterized by uncontrolled growth and spread of abnormal cells. Cancer is caused by both external factors (tobacco, chemicals, radiation, and infections organisms) and internal factors (inherited mutations, hormones, immune conditions, and DNA damage).

These factors may act together or sequentially to initiate and/or promote carcinogenesis. Cancer causes 1 of every 4 deaths and is the leading cause of death in people under age 85 in the United States. Nearly half of all men and a little over one third of all women in the U.S. will develop cancer during their lifetimes. Today, millions of people are living with cancer or have had cancer. The sooner a cancer is found and treatment begins, the better are the chances for living for many years.

Lung cancer is the leading cause of cancer deaths for both men and women in the United States. According to the American Cancer Society, about 160,000 people die annually of this disease, with about 170,000 newly diagnosed cases each year. Despite the use of surgery, chemotherapy, and radiation, the survival rate for patients remains extremely poor (>15% over 5 years). As estimated by the American Cancer Society, the 5-year survival rate for all cancers is about 64% for cancers diagnosed between 1995-2000. However, the survival rate varies depending on cancer type and the stage of cancer at time of detection. For example, the survival rate for brain, breast, and colon cancer is 33, 88, and 63%, respectively for cancers diagnosed between 1995-2000. Therefore, treatments in addition to the standard

methods of treatment that include surgery, radiation, chemotherapy, immunotherapy, and hormone therapy are needed.

Misregulation of genes that control cell fate determination often contributes to cancer. Such altered genes are known as oncogenes.

- 5 Oncogenes are called proto-oncogenes when they are normal (i.e., not mutated). Proto-oncogenes encode components of the cell's normal growth-control pathway. Some of these components are growth factors, receptors, signaling enzymes, and transcription factors.

- 10 Ras is one such oncogene. Mammalian ras genes code for closely related, small proteins (H-ras, K-ras and N-ras). Ras is found in normal cells, where it helps to relay signals by acting as a switch. When receptors on the cell surface are stimulated (by a hormone, for example), Ras is switched on and transduces signals that tell the cell to grow. If the cell-surface receptor is not stimulated, Ras is not activated and so the pathway
15 that results in cell growth is not initiated. In about 30% of human cancers, Ras is mutated so that it is permanently switched on, telling the cell to grow regardless of whether receptors on the cell surface are activated or not. A high incidence of ras gene mutations is found in all lung cancers and adenocarcinomas (10% and 25%, respectively, K-ras), in malignant tumors
20 of the pancreas (80-90%, K-ras), in colorectal carcinomas (30-60%, K-ras), in non-melanoma skin cancer (30-50%, H-ras), in hematopoietic neoplasia of myeloid origin (18-30%, K-and N-ras), and in seminoma (25-40%, K-ras). In other tumors, a mutant ras gene is found at a lower frequency: for example, in breast carcinoma (0-12%, K-ras), glioblastoma and
25 neuroblastoma (0-10%, K- and N-ras).

- Other oncogenes include members of the MYC family (c-MYC, N-MYC, and L-MYC), which have been widely studied, and amplification of myc genes has been found in a variety of tumor types including lung (c-MYC, N-MYC, L-MYC), colon (c-MYC), breast (c-MYC), and
30 neuroblastoma (N-MYC). Genes that inhibit apoptosis have also been identified as oncogenes. The prototype of these genes is BCL-2. Originally identified at the chromosomal breakpoint in follicular lymphoma, this protein was found to inhibit cell death rather than promote cell growth. BCL-2

belongs to a family of intracellular proteins whose role is to regulate caspase activation that leads to DNA fragmentation and cell death. In melanoma, BCL-2 has been reported to be overexpressed in primary and metastatic lesions and this phenotype is associated with tumor progression.

5 Micro RNAs (referred to as “miRNAs”) are small non-coding RNAs, belonging to a class of regulatory molecules found in plants and animals that control gene expression by binding to complementary sites on target messenger RNA (mRNA) transcripts. miRNAs are generated from large RNA precursors (termed pri-miRNAs) that are processed in the nucleus into
10 approximately 70 nucleotide pre-miRNAs, which fold into imperfect stem-loop structures (Lee, Y., et al., *Nature* (2003) 425(6956):415-9) (Figure 1) SEQ ID Nos. 1, 2, and 3). The pre-miRNAs undergo an additional processing step within the cytoplasm where mature miRNAs of 18-25 nucleotides in length are excised from one side of the pre-miRNA hairpin by
15 an RNase III enzyme, Dicer (Hutvagner, G., et al., *Science* (2001) 12:12 and Grishok, A., et al., *Cell* (2001) 106(1):23-34). MiRNAs have been shown to regulate gene expression in two ways. First, miRNAs that bind to protein-coding mRNA sequences that are exactly complementary to the miRNA induce the RNA-mediated interference (RNAi) pathway. Messenger RNA
20 targets are cleaved by ribonucleases in the RISC complex. This mechanism of miRNA-mediated gene silencing has been observed mainly in plants (Hamilton, A.J. and D.C. Baulcombe, *Science* (1999) 286(5441):950-2 and Reinhart, B.J., et al., *MicroRNAs in plants*. *Genes and Dev.* (2002) 16:1616-1626), but an example is known from animals (Yekta, S., I.H. Shih, and D.P. Bartel, *Science* (2004) 304(5670):594-6). In the second mechanism,
25 miRNAs that bind to imperfect complementary sites on messenger RNA transcripts direct gene regulation at the posttranscriptional level but do not cleave their mRNA targets. MiRNAs identified in both plants and animals use this mechanism to exert translational control of their gene targets (Bartel,
30 D.P., *Cell* (2004) 116(2):281-97).

Hundreds of miRNAs have been identified in the fly, worm, plant and mammalian genomes. The biological role for the majority of the miRNAs remains unknown because almost all of these were found through

5

cloning and bioinformatic approaches (Lagos-Quintana, M., et al., Curr Biol (2002) 12(9):735-9; Lagos-Quintana, M., et al., RNA (2003) 9(2): 175-179; Lagos-Quintana, M., et al., Science (2001) 294(5543): 853-8; Lee, R.C. and V. Ambros, Science (2001) 294(5543):862-4; Lau, N.C., et al., Science (2001) 294(5543):858-62; Lim, L.P., et al., Genes Dev (2003) 17(8):991-1008; Johnston, R.J. and O. Hobert, Nature (2003) 426(6968):845-9; and Chang, S., et al. Nature (2004) 430(7001):785-9).

It is likely that these uncharacterized miRNAs act as important gene regulators during development to coordinate proper organ formation, embryonic patterning, and body growth, but this remains to be established. In zebrafish, most miRNAs are expressed from organogenesis onward (Chen, P.Y., et al., Genes Dev (2005) 19(11):1288-93 and Wienholds, E., et al., Science, (2005)).

The biological roles for several miRNAs have been elucidated. These studies highlight the importance of these regulatory molecules in a variety of developmental and metabolic processes. For example, the *Drosophila* miRNA, *bantam*, was identified in a gain-of-function genetic screen for factors that caused abnormal tissue growth (Brennecke, J., et al., Cell (2003) 113(1):25-36). *Bantam* was found to induce tissue growth in the fly by both stimulating cell proliferation and inhibiting apoptosis (Brennecke, J., et al., Cell (2003). 113(1):25-36). Although the proliferation targets for *bantam* have not been identified, a pro-apoptotic gene, *hid*, was shown to have multiple *bantam* complementary sites in its 3'UTR. Since *hid* gene expression was repressed by the *bantam* miRNA, this implicates a role for *bantam* in controlling apoptosis by blocking *hid* function. Another *Drosophila* miRNA, *mir-14*, was identified in a genetic screen for factors that modified Reaper-induced apoptosis in the fly eye (Xu, P., et al., Curr Biol (2003) 13(9):790-5). *mir-14* was shown to be a strong suppressor of apoptosis. In addition, *mir-14* also appears to play a role in the *Drosophila* stress response as well as in regulating fat metabolism. MiRNAs also regulate *Notch* pathway genes in *Drosophila* (Lai, et al. Genes Dev (2005) 19(9):1067-80). A mammalian miRNA, *mir-181*, was shown to direct the

differentiation of human B cells (Chen, C.Z., et al., Science (2004) 303(5654): 83-6), *mir-373* regulates insulin secretion (Poy, M.N., et al., Nature (2004) 432(7014):226-30), while other miRNAs regulate viral infections (Lecellier, C.H., et al., Science (2005) 308(5721):557-60 and Sullivan, C.S., et al., Nature (2005) 435(7042):682-6).

Studies to understand the mechanism of RNAi in *C. elegans*, *Drosophila* and human cells have shown that the miRNA and RNAi pathways may intersect (Grishok, A., et al. Cell (2001) 106(1):23-34 and Hutvagner, G., et al., Science (2001) 293(5531):834-8). MiRNAs copurify with components of the RNAi effector complex, RISC, suggesting a link between miRNAs and siRNAs involved in RNAi (Mourelatos, Z., et al., Genes Dev (2002) 16(6):720-8; Hutvagner, G. and P.D. Zamore, Science (2002) 297(5589):2056-60; and Caudy, A.A., et al., Nature (2003) 425(6956): 411-4). There is also an indication that some protein factors may play a role in both the miRNA ribonucleoprotein (miRNP) and RISC and others might be unique to the miRNP (Grishok, A., et al., Cell (2001) 106(1): 23-34 and Carmell, M.A., et al., Genes Dev (2002) 16(21): 2733-42). For example, proteins of the argonaute/PAZ/PIWI family are components of both RISC and miRNPs. There is also mounting evidence that genes encoding these proteins are linked to cancer. *hAgo3*, *hAgo1*, and *hAgo4* reside in region 1p34-35, often lost in Wilms' tumors, and *Hiwi*, is located on chromosome 12q24.33, which has been linked to the development of testicular germ cell tumors (Carmell, M.A., et al., Genes Dev (2002) 16(21):2733-42). In addition, DICER, the enzyme which processes miRNAs and siRNAs, is poorly expressed in lung cancers (Karube, Y., et al., Cancer Sci (2005) 96(2):111-5).

It is therefore an object of the present invention to provide naturally occurring miRNAs for inhibition of expression of one or more oncogenes.

It is further an object of the present invention to provide naturally occurring nucleic acids for treatment or prophylaxis of one or more symptoms of cancer.

BRIEF SUMMARY OF THE INVENTION

Genes that control cell differentiation and development are frequently mutated in human cancers. These include, but are not limited to, oncogenes such as *RAS*, *c-myc* and *bcl-2*. Naturally occurring microRNAs, in particular *let-7*, have been found that down regulate these oncogenes in humans. Some of the *let-7* genes are located in chromosomal regions that are deleted in certain cancers. Therefore, up-regulating these specific microRNAs or providing analogous pharmaceutical compounds exogenously, should be effective cancer therapies for tumors resulting from activation or over-expression of these oncogenes. MiRNAs nucleic acids including pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of the mature miRNA and DNA encoding a pri-miRNA, pre-miRNA, mature miRNA, fragments or variants thereof, or regulatory elements of the miRNA, referred to jointly as "miRNAs" unless otherwise stated, are described. In one embodiment, the size range of the miRNA can be from 21 nucleotides to 170 nucleotides, although miRNAs of up to 2000 nucleotides can be utilized. In a preferred embodiment the size range of the miRNA is from 70 to 170 nucleotides in length. In another preferred embodiment, mature miRNAs of from 21 to 25 nucleotides in length can be used.

These miRNAs are useful as diagnostics and as therapeutics. The compositions are administered to a patient in need of treatment or prophylaxis of at least one symptom or manifestation (since disease can occur/progress in the absence of symptoms) of cancer. Aberrant expression of oncogenes is a hallmark of cancer, for example, lung cancer. In one embodiment, the compositions are administered in an effective amount to inhibit expression of one or more oncogenes. In a more preferred embodiment, the compositions are administered in an effective amount to inhibit expression of *RAS*, *MYC*, and/or *BCL-2*. Effective, safe dosages can be experimentally determined in model organisms and in human trials by methods well known to one of ordinary skill in the art. The compositions can be administered alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy,

immunotherapy, hormone therapy and laser therapy, to provide a beneficial effect, e.g. reduce tumor size, reduce cell proliferation of the tumor, inhibit angiogenesis, inhibit metastasis, or otherwise improve at least one symptom or manifestation of the disease.

5 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the predicted secondary structure of the *let-7* pre-miRNA from various organisms (SEQ ID Nos. 1, 2, and 3). The shaded residues indicate the mature miRNA transcript excised by Dicer.

Figure 2 is a schematic of potential RNA/RNA duplexes between a
10 *C. elegans* miRNA, *let-7* (SEQ ID No. 6), and the target mRNA, *lin-41* (SEQ ID Nos. 4 and 5). The position of a loss-of-function mutation for the *let-7* miRNA, *let-7(n2853)*, is shown by an arrow below the duplexes.

Figure 3 depicts *let-7* homologues from various species (SEQ ID Nos. 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, and 25),
15 including humans and mouse. There are 2 human homologues of *lin-4* (SEQ ID No. 22), *mir-125a* (SEQ ID No. 25) and *mir-125b* (SEQ ID No. 23). *mir-237* (SEQ ID No. 24) is a *C. elegans lin-4* homologue.

Figures 4A-D depict potential LCSs in *C. elegans let-60/RAS* and in mammalian *RAS* genes. Figure 4A *C. elegans let-60/RAS* mRNA 3'UTR,
20 black arrows indicate sites with similarity between *C. elegans* and *C. briggsae* and white arrows indicate non-similar sites. Shown below are predicted duplexes formed by LCSs (SEQ ID Nos. 26, 27, 28, 29, 32, 33, 34 and 35) (top) and miR-84 (SEQ ID Nos. 30 and 31) (bottom). *let-7* and miR-84 are so similar that most *let-7* sites are also potential miR-84 sites. Figures
25 4B, 4C, and 4D depict that *H.s. NRAS*, *KRAS* and *HRAS* mRNA 3'UTRs have 9 (SEQ ID Nos. 36, 37, 38, 40, 41, 42, 43, 44 and 45), 8 (SEQ ID Nos. 36, 37, 38, 40, 41, 42, 43, 44 and 45), and 3 (SEQ ID Nos. 54, 55 and 56) potential LCSs, respectively. Black arrows indicate sites conserved among mammalian species (in most cases human, rat, mouse, hamster and guinea
30 pig). Shown below are hypothesized duplexes formed by LCSs (top) and *let-7a* miRNA (SEQ ID No. 39) (bottom).

Figure 5A is a graph of the quantitative analysis of the expression pattern from five independent wild-type transgenic lines grown at 20°C. At least 25% repression was observed in all lines. A non-regulated *lin-4l* 3'UTR missing its LCSs (pFS1031), tested in duplicate is shown as a control. Figure 5B is a graph depicting the down-regulation of the reporter gene expression is lost in *let-7(n2853)* mutant worms grown at the

permissive temperature, 15°C. The parental (N2) line was tested in triplicate; four isogenic *let-7(n2853)* mutant lines were tested. Error bars represent standard deviations.

Figures 6A-C are graphs of the quantification of expression data.
 5 *gfp54* is a fusion of *gfp* to the *unc-54* 3'UTR, driven by the *lin-31* promoter. Error bars represent standard deviations.

Figures 7A and B are graphs showing that the presence of *let-7* influences the expression of *RAS* in human cells. HEPG2 cells were transfected with 10 and 30 nM of a *let-7* or negative control precursor
 10 miRNA. Immunofluorescence using an antibody specific to NRAS, VRAS, and KRAS revealed that the *let-7* transfected cells have much lower levels of the RAS proteins. Figure 7A is a graph of the quantification of the RAS antibody fluorescence from replicates of the transfections. Alternatively, HeLa cells were transfected with 100 nM *let-7* inhibitor or negative control
 15 inhibitor. RAS immunofluorescence revealed that cells transfected with the *let-7* inhibitor has increased levels of the RAS proteins relative to the negative control transfected cells. Figure 7B is a graph of the quantification of the RAS antibody fluorescence from replicates of the transfections.

Figures 8A-C depict that the 3'UTRs of NRAS and KRAS enable *let-7*
 20 regulation. Figure 8A is a schematic showing the *NRAS* short (NRAS S) *NRAS* long (NRAS L) and *KRAS* 3'UTRs. Arrows indicate LCSs. The blackened areas indicate the sequence cloned behind the reporter. Figure 8B is a graph of the relative repression of firefly luciferase expression standardized to a transfection control, renilla luciferase. pGL3-Cont is the
 25 empty vector. Figure 8C is a graph of the induction of firefly luciferase expression when reporter plasmids with 3'UTR domains corresponding to KRAS and NRAS are co-transfected with an inhibitor of *let-7*, relative to a control inhibitor.

Figure 9A is a graph of the expression of *let-7* in 21 breast, colon,
 30 and lung tumors relative to associated normal adjacent tissue (NAT). Fluorescently labeled miRNA was hybridized to microarrays that included probes specific to *let-7a* and *let-7c*. Fluorescence intensities for the tumor

and NAT were normalized by total fluorescence signal for all elements and the relative average signal from the let-7 probes in the tumor and normal adjacent samples are expressed as log ratios.

Figure 9B is a graph of the correlation between RAS protein and *let-7c* expression in tumor and normal adjacent tissue samples from three lung squamous cell carcinomas.

10 *NRAS* mRNA in
the tumor and normal adjacent tissues samples was measured by real-time PCR. The real-time data were normalized based on the real-time PCR detection of 18S rRNA in the various samples. The relative expression of *NRAS* in the normal adjacent tissues was taken to be 100% and the Ct value
15 of *NRAS* in the tumor samples was used to assign the relative expression of *NRAS* in the tumor samples.

Figure 10A is a sequence comparison of the let-7 family of miRNAs (SEQ ID Nos. 19, 21, 57, 58 and 59) in *C. elegans*. Figure 10B is a dendrogram of let-7 family members.

20 Figure 11 depicts potential LCSs (SEQ ID Nos. 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 and 80) in let-60/RAS in *C. elegans* mRNA.

Figure 12 is a sequence alignment of *C. elegans* (SEQ ID No. 81) (top) and *C. briggsae* (SEQ ID No. 82) (bottom) let-60/RAS mRNAs
25 (consensus sequence shown in middle) (SEQ ID No. 83). LCSs are shown as black boxes.

Figures 13A-C show that Let-60 is the ortholog of human HRAS, KRAS, and NRAS proteins. Figure 13A is a sequence alignment of *C. elegans* LET-60 (SEQ ID No. 87) with other RAS and RAS related proteins
30 from humans (SEQ ID Nos. 84, 85 and 86) and *C. elegans* (SEQ ID Nos. 88, 89, 90 and 91). Figure 13B is a dendrogram showing the relationship of LET-60 to other RAS related proteins. Figure 13C is a graph showing

quantification of the partial suppression of let-60(gf) alleles. o84-X are lines overexpressing mir-84, and TOPO-X are lines with the empty vector control. The average of each of the three experimental and control lines is indicated.

Figure 14 is a sequence alignment of partial sequences from rodent (SEQ ID Nos. 94, 95 and 96) and human (SEQ ID Nos. 92 and 93) *NRAS* 3'UTRs . The LCSs shown in this alignment are boxed in black.

Figure 15 (SEQ ID Nos. 97, 98, 99, 100, 101, 102 and 103) depicts
5 potential *let-7*::LCS duplexes formed with *Xenopus laevis* and *Danio rerio* *NRAS* 3'UTRs.

Figures 16A and B show that the presence of *let-7* does not influence the expression of control proteins in human cells. Figure 16A is a graph of the quantification of the antibody fluorescence from replicates of HepG2
10 cells transfected with *let-7* or negative control siRNAs using antibodies specific to GAPDH or p21. Figure 16B is a graph of the quantification of fluorescent signal from a single field of 50-100 cells for both the *NRAS*- and Negative Control siRNA transfections.

Figure 17 is a graph of the expression of *let-7c* and *let-7g* in lung
15 tumors relative to associated normal adjacent tissue (NAT).

Figure 18A is a graph of the inhibition of *let-7* that results in a 100% increase of A549 cell numbers, compared to a control transfection and a control anti-miRNA (*mir-19a*). Figure 18B is a graph of Extra *let-7* that causes a decrease in A549 cell numbers compared to a control miRNA (NC).

Figure 19 depicts potential *let-7* complementary sites (boxed) in the
20 3'UTR of human MYC (SEQ ID No. 104) and other vertebrates (SEQ ID Nos. 105, 106 and 107) and potential duplexes between *let-7* (SEQ ID No. 99 and 110) and human MYC (SEQ ID Nos. 108 and 109).

Figure 20A is a graph of the reduced expression of MYC and BCL-2
25 protein in cells treated with exogenous *let-7* miRNA compared to a control miRNA with a scrambled *let-7* sequence. Figure 20B is a graph of the increased expression of MYC and BCL-2 in HeLa cells transfected with an anti-*let-7* molecule.

DETAILED DESCRIPTION OF THE INVENTION

30 Definitions

As used herein the term "nucleic acid" refers to multiple nucleotides (i.e. molecules comprising a sugar (e.g. ribose or deoxyribose) linked to a

phosphate group and to an exchangeable organic base, which is either a substituted pyrimidine (e.g. cytosine (C), thymidine (T) or uracil (U)) or a

substituted purine (e.g. adenine (A) or guanine (G)). The term shall also include polynucleosides (i.e. a polynucleotide minus the phosphate) and any other organic base containing polymer. Purines and pyrimidines include but are not limited to adenine, cytosine, guanine, thymidine, inosine, 5-methylcytosine, 2-aminopurine, 2-amino-6-chloropurine, 2,6-diaminopurine, hypoxanthine, and other naturally and non-naturally occurring nucleobases, substituted and unsubstituted aromatic moieties. Other such modifications are well known to those of skill in the art. Thus, the term nucleic acid also encompasses nucleic acids with substitutions or modifications, such as in the bases and/or sugars.

As used herein, the term "microRNA" refers to any type of interfering RNA, including but not limited to, endogenous microRNA and artificial microRNA. Endogenous microRNA are small RNAs naturally present in the genome which are capable of modulating the productive utilization of mRNA. The term artificial microRNA includes any type of RNA sequence, other than endogenous microRNA, which is capable of modulating the productive utilization of mRNA.

"MicroRNA flanking sequence" as used herein refers to nucleotide sequences including microRNA processing elements. MicroRNA processing elements are the minimal nucleic acid sequences which contribute to the production of mature microRNA from precursor microRNA. Precursor miRNA termed pri-miRNAs are processed in the nucleus into about 70 nucleotide pre-miRNAs, which fold into imperfect stem-loop structures.

The microRNA flanking sequences may be native microRNA flanking sequences or artificial microRNA flanking sequences. A native microRNA flanking sequence is a nucleotide sequence that is ordinarily associated in naturally existing systems with microRNA sequences, i.e., these sequences are found within the genomic sequences surrounding the minimal microRNA hairpin *in vivo*. Artificial microRNA flanking sequences are nucleotide sequences that are not found to be flanking to microRNA sequences in naturally existing systems. The artificial microRNA flanking sequences may be flanking sequences found naturally in

the context of other microRNA sequences. Alternatively they may be composed of minimal microRNA processing elements which are found within naturally occurring flanking sequences and inserted into other random nucleic acid sequences that do not naturally occur as flanking sequences or
5 only partially occur as natural flanking sequences.

The microRNA flanking sequences within the precursor microRNA molecule may flank one or both sides of the stem-loop structure encompassing the microRNA sequence. Preferred structures have flanking sequences on both ends of the stem-loop structure. The flanking sequences
10 may be directly adjacent to one or both ends of the stem-loop structure or may be connected to the stem-loop structure through a linker, additional nucleotides or other molecules.

As used herein a "stem-loop structure" refers to a nucleic acid having a secondary structure that includes a region of nucleotides which are known
15 or predicted to form a double strand (stem portion) that is linked on one side by a region of predominantly single-stranded nucleotides (loop portion). The terms "hairpin" and "fold-back" structures are also used herein to refer to stem-loop structures. Such structures and terms are well known in the art. The actual primary sequence of nucleotides within the stem-loop structure is
20 not critical as long as the secondary structure is present. As is known in the art, the secondary structure does not require exact base-pairing. Thus, the stem may include one or more base mismatches. Alternatively, the base-pairing may not include any mismatches.

As used herein, the term "let-7" refers to the nucleic acid encoding
25 the let-7 miRNA and homologues and variants thereof including conservative substitutions, additions, and deletions therein not adversely affecting the structure or function. Preferably, let-7 refers to the nucleic acid encoding let-7 from *C. elegans* (NCBI Accession No. AY390762), most preferably, let-7 refers to the nucleic acid encoding a let-7 family member
30 from humans, including but not limited to, NCBI Accession Nos. AJ421724, AJ421725, AJ421726, AJ421727, AJ421728, AJ421729, AJ421730, AJ421731, AJ421732, and biologically active sequence variants of let-7,

including alleles, and *in vitro* generated derivatives of let-7 that demonstrate let-7 activity.

Sequence variants of let-7 fall into one or more of three classes: substitutional, insertional or deletional variants. Insertions include 5' and/or
5 3' terminal fusions as well as intrasequence insertions of single or multiple residues. Insertions can also be introduced within the mature sequence of let-7. These, however, ordinarily will be smaller insertions than those at the 5' or 3' terminus, on the order of 1 to 4 residues.

Insertional sequence variants of let-7 are those in which one or more
10 residues are introduced into a predetermined site in the target let-7. Most commonly insertional variants are fusions of nucleic acids at the 5' or 3' terminus of let-7.

Deletion variants are characterized by the removal of one or more residues from the let-7 RNA sequence. These variants ordinarily are
15 prepared by site specific mutagenesis of nucleotides in the DNA encoding let-7, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. However, variant let-7 fragments may be conveniently prepared by *in vitro* synthesis. The variants typically exhibit the same qualitative biological activity as the naturally-occurring analogue,
20 although variants also are selected in order to modify the characteristics of let-7.

Substitutional variants are those in which at least one residue sequence has been removed and a different residue inserted in its place. While the site for introducing a sequence variation is predetermined, the
25 mutation *per se* need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target region and the expressed let-7 variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a
30 known sequence are well known.

Nucleotide substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 residues; and deletions will

WO 2006/028967

PCT/US2005/031327

range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs; i.e. a deletion of 2 residues or insertion of 2 residues. Substitutions, deletion, insertions or any combination thereof may be combined to arrive at a final construct. Changes may be made to increase the activity of the miRNA, to increase its biological stability or half-life, and the like. All such modifications to the nucleotide sequences encoding such miRNA are encompassed.

A DNA isolate is understood to mean chemically synthesized DNA, cDNA or genomic DNA with or without the 3' and/or 5' flanking regions. DNA encoding let-7 can be obtained from other sources by a) obtaining a cDNA library from cells containing mRNA, b) conducting hybridization analysis with labeled DNA encoding let-7 or fragments thereof (usually, greater than 100bp) in order to detect clones in the cDNA library containing homologous sequences, and c) analyzing the clones by restriction enzyme analysis and nucleic acid sequencing to identify full-length clones.

As used herein nucleic acids and/or nucleic acid sequences are homologous when they are derived, naturally or artificially, from a common ancestral nucleic acid or nucleic acid sequence. Homology is generally inferred from sequence similarity between two or more nucleic acids or proteins (or sequences thereof). The precise percentage of similarity between sequences that is useful in establishing homology varies with the nucleic acid and protein at issue, but as little as 25% sequence similarity is routinely used to establish homology. Higher levels of sequence similarity, e.g., 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 99% or more can also be used to establish homology. Methods for determining sequence similarity percentages (e.g., BLASTN using default parameters) are generally available. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information.

30 I. Compositions

Genes that control cell differentiation and development are frequently mutated in human cancers. These include, but are not limited to, oncogenes such as *RAS*, *c-myc* and *bcl-2*. Naturally occurring microRNAs, in particular

let-7, have been found that down regulate these oncogenes in humans. Some of the *let-7* genes are located in chromosomal regions that are deleted in certain cancers. Therefore, up-regulating these specific microRNAs or providing analogous pharmaceutical compounds exogenously, should be effective cancer therapies for tumors resulting from activation or over-expression of these oncogenes.

In preferred embodiments, the miRNA formulations are administered to individuals with a cancer that expresses one or more targets of *let-7* or *lin-4*. More preferably, the formulations are administered to individuals with a cancer that over expresses RAS, MYC and/or BCL-2 or other target having one or more binding sites for the *let-7*. Multiple different pathways, along with the RAS/MAPK pathway, are implicated in cancer. The 3'UTRs of many known cancer genes have been examined and potential *let-7* complementary sites have been identified in the 3'UTR of many of them (Figure 19 and Table 1). These sites possess features of established *let-7* complementary sites in known *C. elegans let-7* targets (Reinhart, et al., Nature (2000) 403:901-906; Johnson, et al., Cell (2005) 120(5):635-47; Grosshans, et al., Dev Cell (2005) 8(3):321-30; Lin, et al., Dev Cell (2003) 4(5):639-50; Slack, et al., Molec. Cell (2000) 5:659-669; Vella, et al., Genes Dev (2004) 18(2):132-7). *let-7* complementary sites (LCS) in the 3'UTRs of human c-MYC and BCL-2 have been identified (see Figure 19 for c-MYC). Many of the potential target genes shown in Table 1 are also up-regulated in lung cancer as well as other cancers, leading to the conclusion that *let-7* is responsible for repressing their expression in normal tissues. Some genes, like GRB2, have a similar number of LCSs as known *let-7* target genes like KRAS (Table 1). VEGF was also identified as having *let-7* complementary sites. These results indicate that *let-7* can repress expression of multiple oncogenes. In addition to inhibiting cell proliferation, *let-7* may also inhibit angiogenesis. Therefore, administration of *let-7* may inhibit multiple pathways that promote survival of cancer or tumor cells (i.e., angiogenesis, decreased apoptosis and increased cell proliferation).

Table 1. Genes implicated in cancer that contain *let-7* binding sites.

<i>Homo sapiens</i> Gene	Number of LCS sites
EGF	1
EGFR	1
ERBB3	3
GRB2	10
NRAS	9
KRAS2	8
HRAS	3
RAF1	1
ARAF	3
MAP2K2	2
MAPK1	1
MAPK3	4
MET	3
KIT	3
TP73L(AIX)	5
MYC	2
MYCL1	6
MYCN	4
BCL2	5
BCL2L1	5
BCL2L2	6
CCND1/cyclinD	3
CDK4	1
MDM2/HDM2	4
FES	2
FURIN	2
INSL3	2

CSF1R/FMS	1
MYBL2	1
MYB	1
PIK3CD	6
PIK3C2B	4
PIK3CG	1
PIK3R5	1
TERT	3
AKT1	3
AKT3	2
VEGF	3
HLIN-41	4
VDR	7
PXR	3
FOXA1	2
FOXA2	A
ASH1L	2
ARID1B	5
GR	2
GLI2	1
14-3-3zeta	6
MO25	1
SMG1	2
FRAP1	3
PER2	4

miRNAs have known roles in human cancer

Recent studies from different laboratories show roles for miRNAs in human cancer (McManus, Seminars in Cancer Biology (2003) 13:253-258).

- 5 The human miRNAs, mir-15 and mir-16, are preferentially deleted or down-regulated in patients with a common form of adult leukemia, B cell chronic lymphocytic leukemia (Calin, et al., Proc Natl Acad Sci U S A (2002))

99(24):15524-9). This study suggests that miRNAs may function as tumor suppressor genes. The bic locus, which encodes the mir-155 miRNA works cooperatively with c-myc and induces B-cell lymphomas, presumably acting as a proto-oncogene (Haasch, D., et al., Cell Immunol (2002) 217(1-2):78-86). miR142 acts as a tumor suppressor in chronic lymphocytic leukemia (Calin, G.A., et al., Proc Natl Acad Sci U S A (2002) 99(24):15524-9; Lagos-Quintana, M., et al., Curr Biol (2002) 12(9):735-9). His-1 acts as an oncogene in B cell lymphoma (Haasch, D., et al., (2002); Lagos-Quintana, M., et al., (2002); Lagos-Quintana, M., et al., Science (2001) 294(5543):853-8; Li, et al. Am J Pathol (1997) 150:1297-305). Translocation of myc to the mir-142 locus causes B cell lymphoma (Lagos-Quintana, M., et al., (2002); Gauwerky, C.E., et al., Proc Natl Acad Sci U S A, 1989. 86(22):8867-71). Mir-143 and mir-145 are poorly expressed in colorectal cancer (Michael, M.Z., et al., Mol Cancer Res (2003) 1(12):882-91). Over-expression of the mir-17, 18, 19, 20 locus is able to cause lymphomas in a mouse model and is up-regulated by MYC (He, L., et al., Nature (2005) 435(7043):828-33; O'Donnell, K.A., et al., Nature (2005) 435(7043):839-43).

Misregulation of genes that control cell fate determination often contributes to cancer. Genes that control cell differentiation and development are frequently mutated in human cancer. The model organism *Caenorhabditis elegans* has been used to identify genes required for cell differentiation in a stem-cell like lineage in the epidermis. *C. elegans* growth and development is divided into three major stages called embryo, larva and adult. Larval growth is subdivided into four larval stages (L1, L2, L3 and L4). Each larval stage ends in a molt and ultimately the animal matures into an adult. The genes that regulate timing of stage-appropriate cell division and differentiation are known as heterochronic genes (Slack, F. and G. Ruvkun, Annu Rev Genet (1997) 31:611-34; Banerjee, D. and F. Slack, Bioessays (2002) 24(2):119-29). In *C. elegans*, heterochronic genes control the timing of cell fate determination and differentiation. In heterochronic mutants, cells frequently fail to terminally differentiate, and instead divide again, a hallmark of cancer.

The founding members of the miRNA family, *lineage defective-4* (*lin-4*) and *lethal-7* (*let-7*), were identified through genetic analysis to control the timing of stage-appropriate cell division and differentiation in *C. elegans* (Lee, et al. *Cell* (1993) 75(5):843-854; Reinhart, B., et al., *Nature* (2000) 403: 901-906; Slack, F. and G. Ruvkun, *Annu Rev Genet* (1997) 31: 611-34; Banerjee, D. and F. Slack, *Bioessays* (2002) 24(2):119-29). *let-7* and *lin-4* control the timing of proliferation versus differentiation decisions. Some of these genes, like *lin-4* and *let-7*, encode microRNAs (miRNAs) that are conserved in humans. Mutations in the *lin-4* and *let-7* miRNAs result in inappropriate reiterations of the first larval stage (L1) and the fourth larval stage (L4) fates, respectively, and these defects lead to disruptions in cell cycle exit (Lee, et al. *Cell* (1993) 75(5):843-854; Reinhart, B., et al., *Nature* (2000) 403:901-906). For example, in wild-type animals, specialized skin cells, known as seam cells, divide with a stem cell pattern and terminally differentiate at the beginning of the adult stage. The seam cells fail to terminally differentiate in *lin-4* and *let-7* mutant animals, and instead reiterate the larval fate and divide again. Lack of cell cycle control and failure to terminally differentiate are hallmarks of cancer.

The expression patterns for *lin-4* and *let-7* correlate with their role in directing developmental timing. *lin-4* RNA accumulates during the L1 stage and is responsible for the L1/L2 transition in nematodes by inhibiting the expression of *lin-14* and *lin-28*, repressors of post-L1 fates (Lee, et al. *Cell* (1993) 75(5):843-854; Ambros, V. and H.R. Horvitz, *Science* (1984) 226:409-416; Wightman, et al. *Cell* (1993) 75(5):855-862; Moss, et al. *Cell* (1997) 88(5): 37-46; and Feinbaum, R. and V. Ambros, *Dev Biol* (1999) 210(1):87-95). *let-7* RNA accumulates during the L4 stage and is responsible for the L4/Adult transition by down-regulating the expression of *lin-41*, *hbl-1* and *RAS* (Johnson, et al., *Cell* (2005) 120(5):635-47; Grosshans, et al., *Dev Cell* (2005) 8(3):321-30; Lin, et al., *Dev Cell* (2003) 4(5):639-50; Slack, F.J., *Molec. Cell* (2000) 5:659-669).

These 21-22 nucleotide miRNAs exert their effect by binding to imperfect complementary sites within the 3'-untranslated regions (3'UTRs) of their target protein-coding mRNAs and repress the expression of these

genes at the level of translation (Lee, et al. Cell (1993) 75(5):843-854; Reinhart, B., et al., Nature (2000) 403:901-906; Moss, et al. Cell (1997) 88(5):637-46; Lin, S.Y., et al., Dev Cell (2003) 4(5):639-50; Slack, F.J., et al., Molec. Cell (2000) 5:659-669; Abrahante, J.E., et al., Dev Cell (2003) 4(5):625-37; and Olsen, P.H. and V. Ambros, Dev Biol (1999) 216(2):671-80). Deletion of *let-7* miRNA complementary sites (LCS) (SEQ ID No. 4 and SEQ ID No. 5) from the *lin-41* 3'UTR (Figure 2) showed abrogation of the normal down-regulation of *lin-41* during the L4 and adult stages and recent work has shown that these complementary sites alone are sufficient for regulation on *lin-41* (Reinhart, B., et al., Nature (2000) 403:901-906; Slack, F.J., et al., Molec. Cell (2000) 5:659-669; Vella, M.C., et al., Genes Dev (2004) 18(2):132-7).

The *let-7* target gene, *lin-41* is similar to known oncogenes. *C. elegans lin-41* loss-of-function (*lf*) mutations cause cells to terminally differentiate precociously, opposite to of the effect seen with *let-7(lf)*, while over-expression of *lin-41* causes *let-7(lf)*-like seam cell proliferation (Reinhart, B., et al., Nature (2000) 403: 901-906; Slack, F.J., et al., Molec. Cell (2000) 5:659-669). Like *let-7*, *lin-41* is an important cell proliferation and differentiation gene. *let-7* and *lin-41* work together to cause cells to proliferate or differentiate at the right time.

lin-41 encodes a member of the RBCC (RING finger, B box, Coiled Coil (Freemont, Ann. New York. Acad. Sci. (1993) 684:174-192) family of proteins. Members of this family have diverse proposed functions, such as transcription and RNA binding, and include the PML (Kakizuka, A., et al., Cell (1991) 66:663-674), TIF1 (Le Dourarin, B., et al., EMBO J. (1995) 14(9):2020-2033) and Rfp proto-oncogenes. The most common form of promyelocytic leukemia involves a translocation that fuses PML to the RAR α gene. The N-terminal part of TIF1, is fused to B-raf in the oncogenic protein T18 (Le Dourarin, B., et al., EMBO J. (1995) 14(9):2020-2033). Emu-ret mice, carrying an RFP/RET fusion gene under the transcriptional control of the immunoglobulin heavy chain enhancer, develop B lineage

leukemias and lymphomas (Wasserman, et al. Blood (1998) 92(1):273-82).
In transformed NIH 3T3 cells, the amino-terminal half of Rfp with a RING

finger motif is fused to a truncated Ret receptor tyrosine kinase, Rfp/Ret (Hasegawa, N., et al., Biochem Biophys Res Commun (1996) 225(2):627-31). Members of this family are associated with cancer progression. It is expected that mammalian *lin-41* is also a proto-oncogene.

5 In addition to *lin-41*, *let-7* regulates other target genes in a 3' UTR dependent manner, including *hunchback-like1* (*hbl-1*) (Lin Shin-Yi, J., et al. Dev. Cell, 2003(4):1-20) and *let-60*, the *C. elegans* *RAS* oncogene homologue (see Example 1). As shown in Table 2, *let-60/RAS* contains multiple *let-7* complimentary sites in its 3'UTR and *let-60(lf)* suppresses *let-*
10 7 mutants. *let-60/RAS* is best understood for its role in *C. elegans* vulval development and *let-60/RAS* 3'UTR is sufficient to restrict *let-60/RAS* expression only to the vulval precursor cell (VPC) that absolutely requires *let-60/ras* activity (the primary induced cell, 1° or P6.p cell). In a normal animal, a *let-7* family member, *mir-84*, is expressed in all the VPCs except
15 the primary induced cell, and represses *let-60/ras* expression in these cells.

In animals carrying *let-60* activating mutations, more than one VPC is induced to differentiate into the 1° cell fate, leading to excess vulvae. Over-expression of *mir-84* suppresses activating mutations in *let-60/RAS*. Many activating mutations in the human *NRAS*, *KRAS* and *HRAS* genes alter
20 the same amino acid affected by the *C. elegans* *let-60* activating mutation. Since *RAS* is mutated in multiple human cancers (Malumbres, et al. Nat Rev Cancer (2003) 3(6):459-65), the hypothesis that human *RAS* is a target of human *let-7* was tested and determined to be (see example 1 below).

lin-4 and *let-7* miRNAs are evolutionarily conserved in higher
25 animals, including humans (Figure 3) (SEQ ID Nos. 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 and 25) and temporally expressed (see examples below) which implies a universal role for these miRNAs during animal development (Lagos-Quintana, M., et al., Mouse. Curr Biol (2002) 12(9):735-9 and Pasquinelli, A.E., et al., Nature (2000)
30 408(6808):86-9). *let-7* orthologues have been identified in mammals, including humans, and *let-7* is expressed in human lung tissues as judged by northern blot (Pasquinelli, A.E., et al.). There are 3 exact copies of the

3
mature *let-7* sequence in the sequenced human genome (referred to as *let-7a1*, *let-7a2*, *let-7a3* under

control of separate promoters) and a variety of close homologues that differ from *let-7* at certain nucleotide positions (e.g. *let-7c* (SEQ ID No. 16), see Figure 3). The nematode, fly and human *let-7* genes are processed from a precursor form (*pre-let-7*) that is predicted to form a stem loop structure,
 5 which is also conserved (Figure 1). Similarly, there are 2 human and mouse homologues of *lin-4* (SEQ ID No. 22), (SEQ ID no. 25) and *mir-125b* (SEQ ID No. 23) (Figure 3).

Recent work has demonstrated that microRNA expression profiles can accurately diagnose particular cancers better than standard messenger
 10 RNA expression profiles (Lu, et al., *Nature* 435:834-838 (2005)).

miRNAs useful to regulate Human Oncogenes

Naturally occurring microRNAs that regulate human oncogenes, pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of the mature miRNA and DNA encoding a pri-
 15 miRNA, pre-miRNA, mature miRNA, fragments or variants thereof, or regulatory elements of the miRNA, have been identified. The size of the miRNA is typically from 21 nucleotides to 170 nucleotides, although nucleotides of up to 2000 nucleotides can be utilized. In a preferred embodiment the size range of the pre-miRNA is between 70 to 170
 20 nucleotides in length and the mature miRNA is between 21 and 25 nucleotides in length.

Nucleic acids

General Techniques

General texts which describe molecular biological techniques include
 25 Sambrook, *Molecular Cloning: a Laboratory Manual* (2nd ed.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989); *Current Protocols in Molecular Biology*, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); *Laboratory Techniques in Biochemistry and Molecular Biology: Hybridization With Nucleic Acid Probes, Part I. Theory and Nucleic Acid*
 30 *Preparation*, P. Tijssen, ed. Elsevier, N.Y. (1993); Berger and Kimmel, *Guide to Molecular Cloning Techniques Methods in Enzymology* volume

152 Academic Press, Inc., San Diego, Calif. These texts describe mutagenesis, the use of vectors, promoters and many other relevant topics

related to, e.g., the generation and expression of genes that encode let-7 or any other miRNA activity. Techniques for isolation, purification and manipulation of nucleic acids, genes, such as generating libraries, subcloning into expression vectors, labeling probes, and DNA hybridization are also described in the texts above and are well known to one of ordinary skill in the art.

The nucleic acids, whether miRNA, DNA, cDNA, or genomic DNA, or a variant thereof, may be isolated from a variety of sources or may be synthesized *in vitro*. Nucleic acids as described herein can be administered to or expressed in humans, transgenic animals, transformed cells, in a transformed cell lysate, or in a partially purified or a substantially pure form.

Nucleic acids are detected and quantified in accordance with any of a number of general means well known to those of skill in the art. These include, for example, analytical biochemical methods such as spectrophotometry, radiography, electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), and hyperdiffusion chromatography, various immunological methods, such as fluid or gel precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immuno-fluorescent assays, and the like, Southern analysis, Northern analysis, Dot-blot analysis, gel electrophoresis, RT-PCR, quantitative PCR, other nucleic acid or target or signal amplification methods, radiolabeling, scintillation counting, and affinity chromatography.

Various types of mutagenesis can be used, e.g., to modify a nucleic acid encoding a gene with let-7 or other miRNA activity. They include but are not limited to site-directed, random point mutagenesis, homologous recombination (DNA shuffling), mutagenesis using uracil containing templates, oligonucleotide-directed mutagenesis, phosphorothioate-modified DNA mutagenesis, and mutagenesis using gapped duplex DNA or the like. Additional suitable methods include point mismatch repair, mutagenesis using repair-deficient host strains, restriction-selection and restriction-

purification, deletion mutagenesis, mutagenesis by total gene synthesis, double-strand break repair, and the like. Mutagenesis, e.g., involving chimeric constructs, are also included in the present invention. In one embodiment, mutagenesis can be guided by known information of the naturally occurring molecule or altered or mutated naturally occurring molecule, e.g., sequence, sequence comparisons, physical properties, crystal structure or the like. Changes may be made to increase the activity of the miRNA, to increase its biological stability or half-life, and the like.

Comparative hybridization can be used to identify nucleic acids encoding genes with let-7 or other miRNA activity, including conservative variations of nucleic acids.

Nucleic acids "hybridize" when they associate, typically in solution. Nucleic acids hybridize due to a variety of well characterized physico-chemical forces, such as hydrogen bonding, solvent exclusion, base stacking and the like. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology-Hybridization with Nucleic Acid Probes* part I chapter 2, "Overview of principles of hybridization and the strategy of nucleic acid probe assays," (Elsevier, N.Y.), as well as in Ausubel, supra. Hames and Higgins (1995) *Gene Probes* 1 IRL Press at Oxford University Press, Oxford, England, (Hames and Higgins 1) and Hames and Higgins (1995) *Gene Probes* 2 IRL Press at Oxford University Press, Oxford, England (Hames and Higgins 2) provide details on the synthesis, labeling, detection and quantification of DNA and RNA, including oligonucleotides.

Nucleic acids which do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides which they encode are substantially identical. This occurs, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code.

Suitable nucleic acids for use in the methods described herein include, but are not limited to, pri-miRNA, pre-miRNA, mature miRNA or fragments of variants thereof that retain the biological activity of the miRNA

and DNA encoding a pri-miRNA, pre-miRNA, mature miRNA, fragments or variants thereof, or DNA encoding regulatory elements of the miRNA.

Viral Vectors

In one embodiment the nucleic acid encoding a miRNA molecule is
5 on a vector. These vectors include a sequence encoding a mature microRNA and *in vivo* expression elements. In a preferred embodiment, these vectors include a sequence encoding a pre-miRNA and *in vivo* expression elements such that the pre-miRNA is expressed and processed *in vivo* into a mature miRNA. In another embodiment, these vectors include a sequence encoding
10 the pri-miRNA gene and *in vivo* expression elements. In this embodiment, the primary transcript is first processed to produce the stem-loop precursor miRNA molecule. The stem-loop precursor is then processed to produce the mature microRNA.

Vectors include, but are not limited to, plasmids, cosmids,
15 phagemids, viruses, other vehicles derived from viral or bacterial sources that have been manipulated by the insertion or incorporation of the nucleic acid sequences for producing the microRNA, and free nucleic acid fragments which can be attached to these nucleic acid sequences. Viral and retroviral vectors are a preferred type of vector and include, but are not limited to,
20 nucleic acid sequences from the following viruses: retroviruses, such as: Moloney murine leukemia virus; Murine stem cell virus, Harvey murine sarcoma virus; murine mammary tumor virus; Rous sarcoma virus; adenovirus; adeno-associated virus; SV40-type viruses; polyoma viruses; Epstein-Barr viruses; papilloma viruses; herpes viruses; vaccinia viruses;
25 polio viruses; and RNA viruses such as any retrovirus. One of skill in the art can readily employ other vectors known in the art.

Viral vectors are generally based on non-cytopathic eukaryotic viruses in which non-essential genes have been replaced with the nucleic acid sequence of interest. Non-cytopathic viruses include retroviruses, the
30 life cycle of which involves reverse transcription of genomic viral RNA into DNA with subsequent proviral integration into host cellular DNA. Retroviruses have been approved for human gene therapy trials. Genetically

altered retroviral expression vectors have general utility for the high-efficiency transduction of nucleic acids in vivo. Standard protocols for producing replication-deficient retroviruses (including the steps of incorporation of exogenous genetic material into a plasmid, transfection of a packaging cell lined with plasmid, production of recombinant retroviruses by the packaging cell line, collection of viral particles from tissue culture media, and infection of the target cells with viral particles) are provided in Kriegler, M., "Gene Transfer and Expression, A Laboratory Manual," W.H. Freeman Co., New York (1990) and Murry, E. J. Ed. "Methods in Molecular Biology," vol. 7, Humana Press, Inc., Clifton, N.J. (1991).

Promoters

The "in vivo expression elements" are any regulatory nucleotide sequence, such as a promoter sequence or promoter-enhancer combination, which facilitates the efficient expression of the nucleic acid to produce the microRNA. The *in vivo* expression element may, for example, be a mammalian or viral promoter, such as a constitutive or inducible promoter or a tissue specific promoter. Examples of which are well known to one of ordinary skill in the art. Constitutive mammalian promoters include, but are not limited to, polymerase promoters as well as the promoters for the following genes: hypoxanthine phosphoribosyl transferase (HPTR), adenosine deaminase, pyruvate kinase, and beta.-actin. Exemplary viral promoters which function constitutively in eukaryotic cells include, but are not limited to, promoters from the simian virus, papilloma virus, adenovirus, human immunodeficiency virus (HIV), Rous sarcoma virus, cytomegalovirus, the long terminal repeats (LTR) of moloney leukemia virus and other retroviruses, and the thymidine kinase promoter of herpes simplex virus. Other constitutive promoters are known to those of ordinary skill in the art. Inducible promoters are expressed in the presence of an inducing agent and include, but are not limited to, metal-inducible promoters and steroid-regulated promoters. For example, the metallothionein promoter is induced to promote transcription in the presence of certain metal ions. Other inducible promoters are known to those of ordinary skill in the art.

Examples of tissue-specific promoters include, but are not limited to, the promoter for creatine kinase, which has been used to direct expression in muscle and cardiac tissue and immunoglobulin heavy or light chain promoters for expression in B cells. Other tissue specific promoters include
5 the human smooth muscle alpha-actin promoter.

Exemplary tissue-specific expression elements for the liver include but are not limited to HMG-COA reductase promoter, sterol regulatory element 1, phosphoenol pyruvate carboxy kinase (PEPCK) promoter, human C-reactive protein (CRP) promoter, human glucokinase promoter, cholesterol
10 7-alpha hydroxylase (CYP-7) promoter, beta-galactosidase alpha-2,6 sialyltransferase promoter, insulin-like growth factor binding protein (IGFBP-1) promoter, aldolase B promoter, human transferrin promoter, and collagen type I promoter.

Exemplary tissue-specific expression elements for the prostate
15 include but are not limited to the prostatic acid phosphatase (PAP) promoter, prostatic secretory protein of 94 (PSP 94) promoter, prostate specific antigen complex promoter, and human glandular kallikrein gene promoter (hgt-1).

Exemplary tissue-specific expression elements for gastric tissue include but are not limited to the human H⁺/K⁺-ATPase alpha subunit
20 promoter.

Exemplary tissue-specific expression elements for the pancreas include but are not limited to pancreatitis associated protein promoter (PAP), elastase 1 transcriptional enhancer, pancreas specific amylase and elastase enhancer promoter, and pancreatic cholesterol esterase gene promoter.

25 Exemplary tissue-specific expression elements for the endometrium include, but are not limited to, the uteroglobin promoter.

Exemplary tissue-specific expression elements for adrenal cells include, but are not limited to, cholesterol side-chain cleavage (SCC) promoter.

30 Exemplary tissue-specific expression elements for the general nervous system include, but are not limited to, gamma-gamma enolase (neuron-specific enolase, NSE) promoter.

Exemplary tissue-specific expression elements for the brain include, but are not limited to, the neurofilament heavy chain (NF-H) promoter.

Exemplary tissue-specific expression elements for lymphocytes include, but are not limited to, the human CGL-1/granzyme B promoter, the
5 terminal deoxy transferase (TdT), lambda 5, VpreB, and lck (lymphocyte specific tyrosine protein kinase p56lck) promoter, the humans CD2 promoter and its 3' transcriptional enhancer, and the human NK and T cell specific activation (NKG5) promoter.

Exemplary tissue-specific expression elements for the colon include,
10 but are not limited to, pp60c-src tyrosine kinase promoter, organ-specific neoantigens (OSNs) promoter, and colon specific antigen-P promoter.

Exemplary tissue-specific expression elements for breast cells include, but are not limited to, the human alpha-lactalbumin promoter.

Exemplary tissue-specific expression elements for the lung include,
15 but are not limited to, the cystic fibrosis transmembrane conductance regulator (CFTR) gene promoter.

Other elements aiding specificity of expression in a tissue of interest can include secretion leader sequences, enhancers, nuclear localization signals, endosmolytic peptides, etc. Preferably, these elements are derived
20 from the tissue of interest to aid specificity.

In general, the *in vivo* expression element shall include, as necessary, 5' non-transcribing and 5' non-translating sequences involved with the initiation of transcription. They optionally include enhancer sequences or upstream activator sequences.

25 **Methods and Materials for Production of miRNA**

The miRNA can be isolated from cells or tissues, recombinantly produced, or synthesized *in vitro* by a variety of techniques well known to one of ordinary skill in the art.

In one embodiment, miRNA is isolated from cells or tissues.
30 Techniques for isolating miRNA from cells or tissues are well known to one of ordinary skill in the art. For example, miRNA can be isolated from total RNA using the mirVana miRNA isolation kit from Ambion, Inc. Another

techniques utilizes the flashPAGE™ Fractionator System (Ambion, Inc.) for PAGE purification of small nucleic acids.

The miRNA can be obtained by preparing a recombinant version thereof (i.e., by using the techniques of genetic engineering to produce a recombinant nucleic acid which can then be isolated or purified by techniques well known to one of ordinary skill in the art). This embodiment involves growing a culture of host cells in a suitable culture medium, and purifying the miRNA from the cells or the culture in which the cells are grown. For example, the methods include a process for producing a miRNA in which a host cell containing a suitable expression vector that includes a nucleic acid encoding an miRNA is cultured under conditions that allow expression of the encoded miRNA. In a preferred embodiment the nucleic acid encodes let-7. The miRNA can be recovered from the culture, from the culture medium or from a lysate prepared from the host cells, and further purified. The host cell can be a higher eukaryotic host cell such as a mammalian cell, a lower eukaryotic host cell such as a yeast cell, or the host cell can be a prokaryotic cell such as a bacterial cell. Introduction of a vector containing the nucleic acid encoding the miRNA into the host cell can be effected by calcium phosphate transfection, DEAE, dextran mediated transfection, or electroporation (Davis, L. et al., *Basic Methods in Molecular Biology* (1986)).

Any host/vector system can be used to express one or more of the miRNAs. These include, but are not limited to, eukaryotic hosts such as HeLa cells and yeast, as well as prokaryotic host such as *E. coli* and *B. subtilis*. miRNA can be expressed in mammalian cells, yeast, bacteria, or other cells where the miRNA gene is under the control of an appropriate promoter. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor, N.Y. (1989). In the preferred embodiment, the miRNA is expressed in mammalian cells. Examples of mammalian expression systems include C127, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human

kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells.

5 Mammalian expression vectors will comprise an origin of replication, a suitable promoter, polyadenylation site, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and polyadenylation sites may be used to provide

10 the required nontranscribed genetic elements. Potentially suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing miRNA. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of

15 expressing miRNA.

In a preferred embodiment, genomic DNA encoding let-7 is isolated, the genomic DNA is expressed in a mammalian expression system, RNA is purified and modified as necessary for administration to a patient. In a preferred embodiment the let-7 is in the form of a pre-miRNA, which can be

20 modified as desired (i.e. for increased stability or cellular uptake).

Knowledge of DNA sequences of miRNA allows for modification of cells to permit or increase expression of an endogenous miRNA. Cells can be modified (e.g., by homologous recombination) to provide increased miRNA expression by replacing, in whole or in part, the naturally occurring

25 promoter with all or part of a heterologous promoter so that the cells express the miRNA at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the desired miRNA encoding sequences. See, for example, PCT International Publication No. WO 94/12650 by Transkaryotic Therapies, Inc., PCT International Publication

30 No. WO 92/20808 by Cell Genesys, Inc., and PCT International Publication No. WO 91/09955 by Applied Research Systems. Cells also may be engineered to express an endogenous gene comprising the miRNA under the

control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. Gene activation techniques are described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.;
5 PCT/US92/09627 (WO93/09222) by Selden et al.; and PCT/US90/06436 (WO91/06667) by Skoultchi et al.

The miRNA may be prepared by culturing transformed host cells under culture conditions suitable to express the miRNA. The resulting expressed miRNA may then be purified from such culture (i.e., from culture
10 medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the miRNA may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl™ or Cibacrom blue 3GA Sepharose™; one or
15 more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; immunoaffinity chromatography, or complementary cDNA affinity chromatography.

The miRNA may also be expressed as a product of transgenic animals, which are characterized by somatic or germ cells containing a
20 nucleotide sequence encoding the miRNA. A vector containing DNA encoding miRNA and appropriate regulatory elements can be inserted in the germ line of animals using homologous recombination (Capecchi, Science 244:1288-1292 (1989)), such that the express the miRNA. Transgenic animals, preferably non-human mammals, are produced using methods as
25 described in U.S. Patent No 5,489,743 to Robinson, et al., and PCT Publication No. WO 94/28122 by Ontario Cancer Institute. miRNA can be isolated from cells or tissue isolated from transgenic animals as discussed above.

In a preferred embodiment, the miRNA can be obtained synthetically,
30 for example, by chemically synthesizing a nucleic acid by any method of synthesis known to the skilled artisan. The synthesized miRNA can then be purified by any method known in the art. Methods for chemical synthesis of

nucleic acids include, but are not limited to, *in vitro* chemical synthesis using phosphotriester, phosphate or phosphoramidite chemistry and solid phase techniques, or via deosynucleoside H-phosphonate intermediates (see U.S. Patent No. 5,705,629 to Bhongle).

5 In some circumstances, for example, where increased nuclease stability is desired, nucleic acids having nucleic acid analogs and/or modified internucleoside linkages may be preferred. Nucleic acids containing modified internucleoside linkages may also be synthesized using reagents and methods that are well known in the art. For example, methods of
10 synthesizing nucleic acids containing phosphonate phosphorothioate, phosphorodithioate, phosphoramidate methoxyethyl phosphoramidate, formacetal, thioformacetal, diisopropylsilyl, acetamidate, carbamate, dimethylene-sulfide (-CH₂-S-CH₂), diethylene-sulfoxide (-CH₂-SO-CH₂), dimethylene-sulfone (-CH₂-SO₂-CH₂), 2'-O-alkyl, and 2'-deoxy-2'-fluoro
15 phosphorothioate internucleoside linkages are well known in the art (see Uhlmann et al., 1990, *Chem. Rev.* 90:543-584; Schneider et al., 1990, *Tetrahedron Lett.* 31:335 and references cited therein). U.S. Patent Nos. 5,614,617 and 5,223,618 to Cook, et al., 5,714,606 to Acevedo, et al., 5,378,825 to Cook, et al., 5,672,697 and 5,466,786 to Buhr, et al., 5,777,092
20 to Cook, et al., 5,602,240 to De Mesmaeker, et al., 5,610,289 to Cook, et al. and 5,858,988 to Wang, also describe nucleic acid analogs for enhanced nuclease stability and cellular uptake.

Formulations

The compositions are administered to a patient in need of treatment
25 or prophylaxis of at least one symptom or manifestation (since disease can occur/progress in the absence of symptoms) of cancer. Aberrant expression of oncogenes is a hallmark of cancer. In a preferred embodiment, the cancer is lung cancer. In one embodiment, the compositions are administered in an effective amount to inhibit gene expression of one or more oncogenes. In
30 preferred embodiments, the compositions are administered in an effective amount to inhibit gene expression of RAS, MYC, and/or BCL-2.

Methods for treatment or prevention of at least one symptom or manifestation of cancer are also described consisting of administration of an effective amount of a composition containing a nucleic acid molecule to alleviate at least one symptom or decrease at least one manifestation. In a preferred embodiment, the cancer is lung cancer. The compositions described herein can be administered in effective dosages alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy, immunotherapy, hormone therapy and laser therapy, to provide a beneficial effect, e.g. reduce tumor size, reduce cell proliferation of the tumor, inhibit angiogenesis, inhibit metastasis, or otherwise improve at least one symptom or manifestation of the disease.

The nucleic acids described above are preferably employed for therapeutic uses in combination with a suitable pharmaceutical carrier. Such compositions comprise an effective amount of the compound, and a pharmaceutically acceptable carrier or excipient. The formulation is made to suit the mode of administration. Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions containing the nucleic acids some of which are described herein.

It is understood by one of ordinary skill in the art that nucleic acids administered *in vivo* are taken up and distributed to cells and tissues (Huang, et al., *FEBS Lett.* 558(1-3):69-73 (2004)). For example, Nyce *et al.* have shown that antisense oligodeoxynucleotides (ODNs) when inhaled bind to endogenous surfactant (a lipid produced by lung cells) and are taken up by lung cells without a need for additional carrier lipids (Nyce and Metzger, *Nature*, 385:721-725 (1997)). Small nucleic acids are readily taken up into T24 bladder carcinoma tissue culture cells (Ma, et al., *Antisense Nucleic Acid Drug Dev.* 8:415-426 (1998)). siRNAs have been used for therapeutic silencing of an endogenous genes by systemic administration (Soutschek, et al., *Nature* 432, 173-178 (2004)).

The nucleic acids described above may be in a formulation for administration topically, locally or systemically in a suitable pharmaceutical carrier. Remington's Pharmaceutical Sciences, 15th Edition by E. W. Martin (Mark Publishing Company, 1975), discloses typical carriers and methods of preparation. The nucleic acids may also be encapsulated in suitable biocompatible microcapsules, microparticles or microspheres formed of biodegradable or non-biodegradable polymers or proteins or liposomes for targeting to cells. Such systems are well known to those skilled in the art and may be optimized for use with the appropriate nucleic acid.

Various methods for nucleic acid delivery are described, for example in Sambrook et al., 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York; and Ausubel et al., 1994, *Current Protocols in Molecular Biology*, John Wiley & Sons, New York. Such nucleic acid delivery systems comprise the desired nucleic acid, by way of example and not by limitation, in either "naked" form as a "naked" nucleic acid, or formulated in a vehicle suitable for delivery, such as in a complex with a cationic molecule or a liposome forming lipid, or as a component of a vector, or a component of a pharmaceutical composition. The nucleic acid delivery system can be provided to the cell either directly, such as by contacting it with the cell, or indirectly, such as through the action of any biological process. By way of example, and not by limitation, the nucleic acid delivery system can be provided to the cell by endocytosis, receptor targeting, coupling with native or synthetic cell membrane fragments, physical means such as electroporation, combining the nucleic acid delivery system with a polymeric carrier such as a controlled release film or nanoparticle or microparticle, using a vector, injecting the nucleic acid delivery system into a tissue or fluid surrounding the cell, simple diffusion of the nucleic acid delivery system across the cell membrane, or by any active or passive transport mechanism across the cell membrane. Additionally, the nucleic acid delivery system can be provided to the cell using techniques such as antibody-related targeting and antibody-mediated immobilization of a viral vector.

Formulations for topical administration may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like can be used as desired.

5 Formulations suitable for parenteral administration, such as, for example, by intraarticular (in the joints), intravenous, intramuscular, intradermal, intraperitoneal, and subcutaneous routes, include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain
10 antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions, solutions or emulsions that can include suspending agents, solubilizers, thickening agents, dispersing agents, stabilizers, and preservatives. Formulations for injection may be presented in unit dosage form, e.g., in ampules or in multi-dose containers, with an
15 added preservative. The compositions may take such forms as.

 Preparations include sterile aqueous or nonaqueous solutions, suspensions and emulsions, which can be isotonic with the blood of the subject in certain embodiments. Examples of nonaqueous solvents are polypropylene glycol, polyethylene glycol, vegetable oil such as olive oil,
20 sesame oil, coconut oil, arachis oil, peanut oil, mineral oil, injectable organic esters such as ethyl oleate, or fixed oils including synthetic mono or di-glycerides. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, 1,3-butandiol, Ringer's dextrose,
25 dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, antioxidants, chelating agents and inert gases and the like. In addition,
30 sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid

may be used in the preparation of injectables. Carrier formulation can be found in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa. Those of skill in the art can readily determine the various parameters for preparing and formulating the compositions without resort to
5 undue experimentation.

The nucleic acids alone or in combination with other suitable components, can also be made into aerosol formulations (i.e., they can be "nebulized") to be administered via inhalation. Aerosol formulations can be placed into pressurized acceptable propellants, such as
10 dichlorodifluoromethane, propane, nitrogen, and the like. For administration by inhalation, the nucleic acids are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant.

In some embodiments, the nucleic acids described above may include
15 pharmaceutically acceptable carriers with formulation ingredients such as salts, carriers, buffering agents, emulsifiers, diluents, excipients, chelating agents, fillers, drying agents, antioxidants, antimicrobials, preservatives, binding agents, bulking agents, silicas, solubilizers, or stabilizers. In one embodiment, the nucleic acids are conjugated to lipophilic groups like
20 cholesterol and lauric and lithocholic acid derivatives with C32 functionality to improve cellular uptake. For example, cholesterol has been demonstrated to enhance uptake and serum stability of siRNA *in vitro* (Lorenz, et al., *Bioorg. Med. Chem. Lett.* 14(19):4975-4977 (2004)) and *in vivo* (Soutschek, et al., *Nature* 432(7014):173-178 (2004)). In addition, it has been shown that
25 binding of steroid conjugated oligonucleotides to different lipoproteins in the bloodstream, such as LDL, protect integrity and facilitate biodistribution (Rump, et al., *Biochem. Pharmacol.* 59(11):1407-1416 (2000)). Other groups that can be attached or conjugated to the nucleic acids described above to increase cellular uptake, include, but are not limited to,
30 acridinederivatives; cross-linkers such as psoralen derivatives, azidophenacyl, proflavin, and azidoproflavin; artificial endonucleases; metal complexes such as EDTA-Fe(II) and porphyrin-Fe(II); alkylating moieties,;

nucleases such as alkaline phosphatase; terminal transferases; abzymes; cholesteryl moieties; lipophilic carriers; peptide conjugates; long chain alcohols; phosphate esters; radioactive markers; non-radioactive markers; carbohydrates; and polylysine or other polyamines. U.S. Patent No. 6,919,208 to Levy, et al., also described methods for enhanced delivery of nucleic acids molecules.

These pharmaceutical formulations may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

The formulations described herein of the nucleic acids embrace fusions of the nucleic acids or modifications of the nucleic acids, wherein the nucleic acid is fused to another moiety or moieties, e.g., targeting moiety or another therapeutic agent. Such analogs may exhibit improved properties such as activity and/or stability. Examples of moieties which may be linked or unlinked to the nucleic acid include, for example, targeting moieties which provide for the delivery of nucleic acid to specific cells, e.g., antibodies to pancreatic cells, immune cells, lung cells or any other preferred cell type, as well as receptor and ligands expressed on the preferred cell type. Preferably, the moieties target cancer or tumor cells. For example, since cancer cells have increased consumption of glucose, the nucleic acids can be linked to glucose molecules. Monoclonal humanized antibodies that target cancer or tumor cells are preferred moieties and can be linked or unlinked to the nucleic acids. In the case of cancer therapeutics, the target antigen is typically a protein that is unique and/or essential to the tumor cells (e.g., the receptor protein HER-2).

II. Methods of Treatment

Method of Administration

In general, methods of administering nucleic acids are well known in the art. In particular, the routes of administration already in use for nucleic acid therapeutics, along with formulations in current use, provide preferred

routes of administration and formulation for the nucleic acids described above.

Nucleic acid compositions can be administered by a number of routes including, but not limited to: oral, intravenous, intraperitoneal, intramuscular, 5 transdermal, subcutaneous, topical, sublingual, or rectal means. Nucleic acids can also be administered via liposomes. Such administration routes and appropriate formulations are generally known to those of skill in the art.

Administration of the formulations described herein may be accomplished by any acceptable method which allows the miRNA or nucleic 10 acid encoding the miRNA to reach its target. The particular mode selected will depend of course, upon factors such as the particular formulation, the severity of the state of the subject being treated, and the dosage required for therapeutic efficacy. As generally used herein, an "effective amount" of a nucleic acids is that amount which is able to treat one or more symptoms of 15 cancer or related disease, reverse the progression of one or more symptoms of cancer or related disease, halt the progression of one or more symptoms of cancer or related disease, or prevent the occurrence of one or more symptoms of cancer or related disease in a subject to whom the formulation is administered, as compared to a matched subject not receiving the compound 20 or therapeutic agent. The actual effective amounts of drug can vary according to the specific drug or combination thereof being utilized, the particular composition formulated, the mode of administration, and the age, weight, condition of the patient, and severity of the symptoms or condition being treated.

25 Any acceptable method known to one of ordinary skill in the art may be used to administer a formulation to the subject. The administration may be localized (i.e., to a particular region, physiological system, tissue, organ, or cell type) or systemic, depending on the condition being treated.

30 Injections can be e.g., intravenous, intradermal, subcutaneous, intramuscular, or intraperitoneal. The composition can be injected intradermally for treatment or prevention of cancer, for example. In some embodiments, the injections can be given at multiple locations. Implantation

includes inserting implantable drug delivery systems, e.g., microspheres, hydrogels, polymeric reservoirs, cholesterol matrixes, polymeric systems, e.g., matrix erosion and/or diffusion systems and non-polymeric systems, e.g., compressed, fused, or partially-fused pellets. Inhalation includes
5 administering the composition with an aerosol in an inhaler, either alone or attached to a carrier that can be absorbed. For systemic administration, it may be preferred that the composition is encapsulated in liposomes.

Preferably, the agent and/or nucleic acid delivery system are provided in a manner which enables tissue-specific uptake of the agent and/or nucleic
10 acid delivery system. Techniques include using tissue or organ localizing devices, such as wound dressings or transdermal delivery systems, using invasive devices such as vascular or urinary catheters, and using interventional devices such as stents having drug delivery capability and configured as expansive devices or stent grafts.

15 The formulations may be delivered using a bioerodible implant by way of diffusion or by degradation of the polymeric matrix. In certain embodiments, the administration of the formulation may be designed so as to result in sequential exposures to the miRNA over a certain time period, for example, hours, days, weeks, months or years. This may be accomplished,
20 for example, by repeated administrations of a formulation or by a sustained or controlled release delivery system in which the miRNA is delivered over a prolonged period without repeated administrations. Administration of the formulations using such a delivery system may be, for example, by oral dosage forms, bolus injections, transdermal patches or subcutaneous
25 implants. Maintaining a substantially constant concentration of the composition may be preferred in some cases.

Other delivery systems suitable include, but are not limited to, time-release, delayed release, sustained release, or controlled release delivery systems. Such systems may avoid repeated administrations in many cases,
30 increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include, for example, polymer-based systems such as polylactic

and/or polyglycolic acids, polyanhydrides, polycaprolactones, copolyoxalates, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and/or combinations of these. Microcapsules of the foregoing polymers containing nucleic acids are described in, for example, U.S. Patent No. 5,075,109. Other examples include nonpolymer systems that are lipid-based including sterols such as cholesterol, cholesterol esters, and fatty acids or neutral fats such as mono-, di- and triglycerides; hydrogel release systems; liposome-based systems; phospholipid based-systems; silastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; or partially fused implants. Specific examples include, but are not limited to, erosional systems in which the miRNA is contained in a formulation within a matrix (for example, as described in U.S. Patent Nos. 4,452,775, 4,675,189, 5,736,152, 4,667,013, 4,748,034 and 5,239,660), or diffusional systems in which an active component controls the release rate (for example, as described in U.S. Patent Nos. 3,832,253, 3,854,480, 5,133,974 and 5,407,686). The formulation may be as, for example, microspheres, hydrogels, polymeric reservoirs, cholesterol matrices, or polymeric systems. In some embodiments, the system may allow sustained or controlled release of the composition to occur, for example, through control of the diffusion or erosion/degradation rate of the formulation containing the miRNA. In addition, a pump-based hardware delivery system may be used to deliver one or more embodiments.

Examples of systems in which release occurs in bursts includes, e.g., systems in which the composition is entrapped in liposomes which are encapsulated in a polymer matrix, the liposomes being sensitive to specific stimuli, e.g., temperature, pH, light or a degrading enzyme and systems in which the composition is encapsulated by an ionically-coated microcapsule with a microcapsule core degrading enzyme. Examples of systems in which release of the inhibitor is gradual and continuous include, e.g., erosional systems in which the composition is contained in a form within a matrix and diffusional systems in which the composition permeates at a controlled rate,

e.g., through a polymer. Such sustained release systems can be e.g., in the form of pellets, or capsules.

Use of a long-term release implant may be particularly suitable in some embodiments. "Long-term release," as used herein, means that the
5 implant containing the composition is constructed and arranged to deliver therapeutically effective levels of the composition for at least 30 or 45 days, and preferably at least 60 or 90 days, or even longer in some cases. Long-term release implants are well known to those of ordinary skill in the art, and include some of the release systems described above.

10 Dosages for a particular patient can be determined by one of ordinary skill in the art using conventional considerations, (e.g. by means of an appropriate, conventional pharmacological protocol). A physician may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. The dose administered to a
15 patient is sufficient to effect a beneficial therapeutic response in the patient over time, or, e.g., to reduce symptoms, or other appropriate activity, depending on the application. The dose is determined by the efficacy of the particular formulation, and the activity, stability or serum half-life of the miRNA employed and the condition of the patient, as well as the body
20 weight or surface area of the patient to be treated. The size of the dose is also determined by the existence, nature, and extent of any adverse side-effects that accompany the administration of a particular vector, formulation, or the like in a particular patient.

Therapeutic compositions comprising one or more nucleic acids are
25 optionally tested in one or more appropriate *in vitro* and/or *in vivo* animal models of disease, to confirm efficacy, tissue metabolism, and to estimate dosages, according to methods well known in the art. In particular, dosages can be initially determined by activity, stability or other suitable measures of treatment vs. non-treatment (e.g., comparison of treated vs. untreated cells or
30 animal models), in a relevant assay. Formulations are administered at a rate determined by the LD50 of the relevant formulation, and/or observation of any side-effects of the nucleic acids at various concentrations, e.g., as applied

WO 2006/028967

PCT/US2005/031327

to the mass and overall health of the patient. Administration can be accomplished via single or divided doses.

In vitro models can be used to determine the effective doses of the nucleic acids as a potential cancer treatment. Suitable *in vitro* models include, but are not limited to, proliferation assays of cultured tumor cells, growth of cultured tumor cells in soft agar (see Freshney, (1987) Culture of Animal Cells: A Manual of Basic Technique, Wiley-Liss, New York, N.Y. Ch 18 and Ch 21), tumor systems in nude mice as described in Giovanella et al., J. Natl. Can. hist., 52: 921-30 (1974), mobility and invasive potential of tumor cells in Boyden Chamber assays as described in Pilkington et al., Anticancer Res., 17: 4107-9 (1997), and angiogenesis assays such as induction of vascularization of the chick chorioallantoic membrane or induction of vascular endothelial cell migration as described in Ribatta et al., Intl. J. Dev. Biol., 40: 1189-97 (1999) and Li et al., Clin. Exp. Metastasis, 17:423-9 (1999), respectively. Suitable tumor cells lines are available, e.g. from American Type Tissue Culture Collection catalogs.

In vivo models are the preferred models to determine the effective doses of nucleic acids described above as potential cancer treatments. Suitable *in vivo* models include, but are not limited to, mice that carry a mutation in the *KRAS* oncogene (Lox-Stop-Lox K-Ras^{G12D} mutants, *Kras2^{tm4Tyj}*) available from the National Cancer Institute (NCI) Frederick Mouse Repository. Other mouse models known in the art and that are available include but are not limited to models for gastrointestinal cancer, hematopoietic cancer, lung cancer, mammary gland cancer, nervous system cancer, ovarian cancer, prostate cancer, skin cancer, cervical cancer, oral cancer, and sarcoma cancer.

In determining the effective amount of the miRNA to be administered in the treatment or prophylaxis of disease the physician evaluates circulating plasma levels, formulation toxicities, and progression of the disease.

The dose administered to a 70 kilogram patient is typically in the range equivalent to dosages of currently-used therapeutic antisense oligonucleotides such as Vitravene® (fomivirsen sodium injection) which is

approved by the FDA for treatment of cytomegaloviral RNA, adjusted for the altered activity or serum half-life of the relevant composition.

The formulations described herein can supplement treatment conditions by any known conventional therapy, including, but not limited to, antibody administration, vaccine administration, administration of cytotoxic agents, natural amino acid polypeptides, nucleic acids, nucleotide analogues, and biologic response modifiers. Two or more combined compounds may be used together or sequentially. For example, the nucleic acids can also be administered in therapeutically effective amounts as a portion of an anti-cancer cocktail. An anti-cancer cocktail is a mixture of the oligonucleotide or modulator with one or more anti-cancer drugs in addition to a pharmaceutically acceptable carrier for delivery. The use of anti-cancer cocktails as a cancer treatment is routine. Anti-cancer drugs that are well known in the art and can be used as a treatment in combination with the nucleic acids described herein include, but are not limited to: Actinomycin D, Aminoglutethimide, Asparaginase, Bleomycin, Busulfan, Carboplatin, Carmustine, Chlorambucil, Cisplatin (cis-DDP), Cyclophosphamide, Cytarabine HCl (Cytosine arabinoside), Dacarbazine, Dactinomycin, Daunorubicin HCl, Doxorubicin HCl, Estramustine phosphate sodium, Etoposide (V16-213), Floxuridine, 5-Fluorouracil (5-Fu), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alpha-2a, Interferon Alpha-2b, Leuprolide acetate (LHRH-releasing factor analog), Lomustine, Mechlorethamine HCl (nitrogen mustard), Melphalan, Mercaptopurine, Mesna, Methotrexate (MTX), Mitomycin, Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Vincristine sulfate, Amsacrine, Azacitidine, Hexamethylmelamine, Interleukin-2, Mitoguazone, Pentostatin, Semustine, Teniposide, and Vindesine sulfate.

Diseases to be Treated

Cancer treatments promote tumor regression by inhibiting tumor cell proliferation, inhibiting angiogenesis (growth of new blood vessels that is necessary to support tumor growth) and/or prohibiting metastasis by

reducing tumor cell motility or invasiveness. Therapeutic formulations described herein may be effective in adult and pediatric oncology including in solid phase tumors/malignancies, locally advanced tumors, human soft tissue sarcomas, metastatic cancer, including lymphatic metastases, blood cell malignancies including multiple myeloma, acute and chronic leukemias, and lymphomas, head and neck cancers including mouth cancer, larynx cancer and thyroid cancer, lung cancers including small cell carcinoma and non-small cell cancers, breast cancers including small cell carcinoma and ductal carcinoma, gastrointestinal cancers including esophageal cancer, stomach cancer, colon cancer, colorectal cancer and polyps associated with colorectal neoplasia, pancreatic cancers, liver cancer, urologic cancers including bladder cancer and prostate cancer, malignancies of the female genital tract including ovarian carcinoma, uterine (including endometrial) cancers, and solid tumor in the ovarian follicle, kidney cancers including renal cell carcinoma, brain cancers including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, bone cancers including osteomas, skin cancers including malignant melanoma, tumor progression of human skin keratinocytes, squamous cell carcinoma, basal cell carcinoma, hemangiopericytoma and Kaposi's sarcoma. Therapeutic formulations can be administered in therapeutically effective dosages alone or in combination with adjuvant cancer therapy such as surgery, chemotherapy, radiotherapy, thermotherapy, immunotherapy, hormone therapy and laser therapy, to provide a beneficial effect, e.g. reducing tumor size, slowing rate of tumor growth, reducing cell proliferation of the tumor, promoting cancer cell death, inhibiting angiogenesis, inhibiting metastasis, or otherwise improving overall clinical condition, without necessarily eradicating the cancer.

Cancers include, but are not limited to, biliary tract cancer; bladder cancer; breast cancer; brain cancer including glioblastomas and medulloblastomas; cervical cancer; choriocarcinoma; colon cancer including colorectal carcinomas; endometrial cancer; esophageal cancer; gastric cancer; head and neck cancer; hematological neoplasms including acute

lymphocytic and myelogenous leukemia, multiple myeloma, AIDS-associated leukemias and adult T-cell leukemia lymphoma; intraepithelial neoplasms including Bowen's disease and Paget's disease; liver cancer; lung cancer including small cell lung cancer and non-small cell lung cancer; lymphomas including Hodgkin's disease and lymphocytic lymphomas; neuroblastomas; oral cancer including squamous cell carcinoma; osteosarcomas; ovarian cancer including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells; pancreatic cancer; prostate cancer; rectal cancer; sarcomas including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma, synovial sarcoma and osteosarcoma; skin cancer including melanomas, Kaposi's sarcoma, basocellular cancer, and squamous cell cancer; testicular cancer including germinal tumors such as seminoma, non-seminoma (teratomas, choriocarcinomas), stromal tumors, and germ cell tumors; thyroid cancer including thyroid adenocarcinoma and medullar carcinoma; transitional cancer and renal cancer including adenocarcinoma and Wilms tumor. In a preferred embodiment, the formulations are administered for treatment or prevention of lung cancer.

In addition, therapeutic nucleic acids may be used for prophylactic treatment of cancer. There are hereditary conditions and/or environmental situations (e.g. exposure to carcinogens) known in the art that predispose an individual to developing cancers. Under these circumstances, it may be beneficial to treat these individuals with therapeutically effective doses of the nucleic acids to reduce the risk of developing cancers. In one embodiment, a nucleic acid in a suitable formulation may be administered to a subject who has a family history of cancer, or to a subject who has a genetic predisposition for cancer. In other embodiments, the nucleic acid in a suitable formulation is administered to a subject who has reached a particular age, or to a subject more likely to get cancer. In yet other embodiments, the nucleic acid in a suitable formulation is administered to subjects who exhibit symptoms of cancer (e.g., early or advanced). In still other embodiments, the nucleic acid in a suitable formulation may be administered to a subject as a

preventive measure. In some embodiments, the nucleic acid in a suitable formulation may be administered to a subject based on demographics or epidemiological studies, or to a subject in a particular field or career.

Examples

5 The present invention will be further understood by reference to the following non-limiting examples.

Example 1. *RAS* is regulated by the *let-7* microRNA family

10 *C. elegans let-7, mir-48, mir-84* and *mir-241* encode four developmentally regulated miRNAs that comprise the *let-7* family (Lau et al., (2001) Science 294, 858-862; Lim et al., (2003) Genes Dev 17, 991-1008; Reinhart et al., (2000) Nature 403, 901-906). This family displays high sequence identity, with particular conservation at the 5' end of the mature miRNAs (Figure 10A, 10B). The *C. elegans let-7* family miRNAs, *mir-84*, plays a role in vulval development, a model for dissecting

15 *RAS/MAP Kinase* signaling (Wang and Sternberg, (2001) Curr Top Dev Biol 51, 189-220). *C. elegans let-60/RAS* is regulated by members of the *let-7* family. *let-7* and *mir-84* are complementary to multiple sites in the 3' UTR of *let-60/RAS*. *let-7* and *mir-84* are expressed in a reciprocal manner to *let-60/RAS* in the hypodermis and the vulva respectively.

20 *let-7* and *mir-84* genetically interact with *let-60/RAS*, consistent with negative regulation of *RAS* expression by *let-7* and *mir-84*. The results demonstrate for the first time that miRNAs can regulate *RAS*, a critical human oncogene. Moreover, all three human *RAS* genes have *let-7* complementary sites in their 3'UTRs that subject the oncogenes to *let-7*

25 miRNA-mediated regulation in cell culture. Lung tumor tissues display significantly reduced levels of *let-7* and significantly increased levels of *RAS* protein relative to normal lung tissue, suggesting *let-7* regulation of *RAS* as a mechanism for *let-7* in lung oncogenesis.

Experimental Procedures

30 **Plasmid Constructs.** PSJ840 (*mir-84::gfp*) was made by amplifying 2.2 kb of genomic sequence (base pairs -2201 to -9) upstream of the mature *mir-84* sequence from N2 genomic DNA and adding a *SmaI* site and an *AgeI* site to the 5' and 3' ends, respectively, using the polymerase

chain reaction (PCR) with primers MIR84UP and MIR84DN (all primer sequences available upon request). This product was digested with SmaI and AgeI and then cloned into the pPD95.70 vector digested with SmaI and AgeI. The *mir-84* upstream DNA contained various sequence elements that are also found in the *let-7* upstream DNA. PSJo84 (*mir-84(+++)*) was made by amplifying 3.0 kb of genomic sequence (base pairs -2201 to +792) upstream and downstream of the mature *mir-84* sequence including the *mir-84* sequence itself from N2 genomic DNA and adding SmaI sites to both the 5' and 3' ends using PCR with primers MIR84UP and MIR84DN2. This product was ligated into the pCR4-TOPO vector using the TOPO TA Cloning Kit as described by the manufacturer (Invitrogen). The empty TOPO control vector was made by digesting PSJo84 with EcoRI, extracting the 4 kb vector band and self ligating it. The miR-84 deletion plasmid, o84Δ84 (Δ*mir-84(+++)*), was made by overlap extension PCR, starting with two separate PCR reactions using primers MIR84UP with DEL84DN and MIR84DN2 with DEL84UP and PSJo84 plasmid as template. The two PCR products were purified (QIAGEN) and then used together as template for the final PCR using primers MIR84UP and MIR84DN2. This final product, identical to PSJo84 except for the deletion of the 75 nt pre-miR84 encoding sequence, was ligated into the pCR2.1-TOPO vector using the TOPO TA Cloning Kit (Invitrogen) and called o84Δ84. GFP60 was made by amplifying the 819 bp of genomic sequence encoding the *let-60* 3'UTR from N2 genomic DNA and adding EcoRI and SpeI sites to the 5' and 3' ends, respectively, using PCR with primers 3LET60UP and 3LET60DN. This product was digested with EcoRI and SpeI and then cloned into the pPD95.70 vector (Fire Lab) digested with EcoRI and SpeI, thus replacing the *unc-54* 3'UTR found in pPD95.70 with the *let-60* 3'UTR. The green fluorescent protein gene (GFP) followed by the *let-60* 3'UTR was then amplified out of this plasmid and BglII and NotI sites were added to the 5' and 3' ends respectively using PCR with primers BGLGFP and 3UTRNOT. This PCR product was digested with BglII and NotI and ligated into PB255 digested with BglII and NotI, resulting in the plasmid GFP60 containing the *lin-31* promoter and enhancer driving GFP with the *let-60* 3'UTR. GFP54

was made by amplifying *gfp* followed by the *unc-54* 3'UTR out of pPD95.70 using PCR with primers BGLGFP and 3UTRNOT. This PCR product was digested and ligated into PB255 digested with BglII and NotI, resulting in the plasmid GFP54 containing the *lin-31* promoter and enhancer driving GFP
 5 with the *unc-54* 3'UTR. pGL3-NRAS S and pGL3-NRAS L were made by amplifying the entire 1140 bp shorter form of the *H.s. NRAS* 3'UTR out of a IMAGE cDNA clone (accession # AI625442), or the entire 3616 bp longer form of the *H.s. NRAS* 3'UTR (excluding the first 43 bp) from *H.s.* genomic DNA, and adding NheI sites to the ends by PCR using primers SMJ100 and
 10 SMJ101 or SMJ102 and SMJ103 respectively. These products were digested with NheI and ligated into pGL3-Control (Promega) digested with XbaI and treated with CIP, resulting in *Rrluc*-expressing plasmids containing either the short or long form of the *H.s. NRAS* 3'UTR. To generate pFS1047, containing the *col-10-lacZ-let-60* 3'UTR reporter gene, the entire *let-60*
 15 3'UTR was subcloned into the *SacII* and *NcoI* sites of plasmid B29 (Wightman, B., Ha, I., and Ruvkun, G. (1993) *Cell* 75, 855–862).

***let-60; let-7* Double Mutants.** *let-60(n2021); let-7(mn112)* double mutant animals were generated by crossing *let-60(n2021)* heterozygotes with *let-7(mn112) unc-3/0; mnDp1* hemizygotes. From this cross, 180 individual
 20 homozygous *let-7(mn112)* F2 animals were female, as determined by their Unc-phenotype, due to the tight linkage of the *let-7(mn112)* to *unc-3*. Of these, 10 survived into adulthood and produced eggs. The resulting progeny from these animals died as larvae with the rod-like phenotype that is characteristic of *let-60* mutant animals, thus showing that these animals all
 25 contained the *let-60(n2021)* mutation, and that adult survival was likely due to the *let-60(n2021)* mutation, confirming our RNAi data. From 180 F2s, 25% (or about 45 animals) would have been predicted to be *let-60(n2021)* homozygous. Since one saw survival of only 10 of these animals, this indicates a suppression of about 22%, similar to what was observed with
 30 RNAi. However, because the brood sizes were low (usually about 10 hatched larvae and several unhatched eggs); and because of a combination of larval lethality as well as limited parental survival, double mutant lines could not be established for further analysis; all progeny died as larvae.

***C. elegans* and Transgenic reporter analysis.** All animal experiments were performed at room temperature or 20°C unless stated otherwise. All experimental plasmids were injected in animals at 50-100 ng/μl. Two different markers, *rol-6* (100 ng/μl) and *myo-3::gfp* (50 ng/μl), were separately co-injected with PSJo84: *myo-3::gfp* (50 ng/μl) was co-injected with o84Δ84; and *myo-2::gfp* (5 ng/μl) was co-injected with GFP60 and GFP54. These animals are mosaic for the transgenes. To compare expression between individual lines, the percent expression of GFP in each of the Pn.p cells was normalized relative to the expression of the highest expressing Pn.p cell and represented as a fraction of the highest expresser for each individual line of animals. For each construct the average of the lines was calculated along with the standard deviation for each construct represented as error bars (*mir-84::gfp* n=239, *gfp60* n=42 and *gfp54* n=40). For the *mir-84*(+++) analysis, animals were examined using DIC optics to score seam cell and vulval anatomy. LacZ reporter analysis was as described (Vella et al., (2004) *Genes Dev* 18, 132-137). The *lin-41* 3'UTR missing its LCSs (pFS1031) was used as a control (Reinhart et al., (2000)). RNAi methods were standard feeding procedures using synchronized L1s (Timmons et al., (2001) *Gene* 263, 103-112). All RNAi experiments were done in parallel to an empty vector (L4440) feeding control. See Supplemental Experimental procedures for details on the *let-60*; *let-7* double mutant cross.

***let-7/RAS* association in mammalian cells.** HeLa S3 cells grown in D-MEM (GIBCO) supplemented with 10% fetal bovine serum (GIBCO) were cotransfected in 12-well plates using Lipofectamine 2000 (Invitrogen) according to the manufacturer's protocol using 1.0 μg/well of *Pp*-luc-expressing plasmid (pGL3-Control from Promega, pGL3-NRAS S pGL3-NRAS L and pGL3-KRAS) and 0.1 μg/well of *Rr*-luc-expressing plasmid (pRL-TK from Promega). 24 hrs post transfection, the cells were harvested and assayed using the Dual Luciferase assay as described by the manufacturer (Promega). HeLa cells grown as above were transfected in 24-well plates with 30 pmoles of Anti-miR *let-7* or negative control #1 inhibitors (Ambion) using Lipofectamine 2000. Three days post-

transfection, RAS expression was monitored by immunofluorescence using an FITC conjugated primary antibody against RAS protein (US Biological). The resulting fluorescent signal was analyzed using the appropriate filter set and was quantified using MetaMorph software. The fluorescence intensity
5 of 150-300 cells was typically measured in one or a few viewing areas. The experiments with both the precursors and the inhibitors were performed three times. The photos represent single viewing fields from one of the experiments and are representative of the triplicate experiment. Identically grown HeLa cells were cotransfected in 24-well plates using Lipofectamine
10 2000 (Invitrogen) according to the manufacturer's protocol using 200 ng /well of *Pp*-luc-expressing plasmid (pGL3-Control from Promega, pGL3-NRAS S pGL3-NRAS L and pGL3-KRAS). 48 hrs post transfection, the cells were harvested and assayed using the Luciferase assay as described by the manufacturer (Promega).

15 HepG2 cells grown in D-MEM (GIBCO) supplemented with 10% fetal bovine serum (GIBCO) were transfected with 15 or 5 pmole of Pre-miR *Let-7c* or negative control #1 Precursor miRNAs (Ambion) in 24-well plates using siPort Neo-FX (Ambion) according to the manufacturer's protocol. Three days post-transfection, RAS expression was monitored by
20 immunofluorescence as described above.

MiRNA microarray analysis procedures used by Ambion, Inc.
Total RNA from tumor and normal adjacent tissue (NAT) samples from 3 breast cancer, 6 colon cancer, and 12 lung cancer patients was isolated using the mirVana RNA Isolation Kit (Ambion). Twenty µg of each total RNA
25 sample was fractionated by polyacrylamide gel electrophoresis (PAGE) using a 15% denaturing polyacrylamide gel and the miRNA fractions for each sample were recovered. The miRNAs from all of the samples were subjected to a poly(A) polymerase reaction wherein amine modified uridines were incorporated as part of ~40 nt long tails (Ambion). The tailed tumor
30 samples were fluorescently labeled using an amine-reactive Cy3 (Amersham) and the normal adjacent tissue samples were labeled with Cy5 (Amersham). The fluorescently labeled miRNAs were purified by glass-

fiber filter binding and elution (Ambion) and the tumor and normal adjacent tissue samples from the same patient were mixed. Each sample mixture was hybridized for 14 hours with slides upon which 167 miRNA probes were arrayed. The microarrays were washed 3 X 2 minutes (min) in 2X SSC and scanned using a GenePix 4000B (Axon). Fluorescence intensities for the Cy3- and Cy5-labeled samples for each element were normalized by total Cy3 and Cy5 signal on the arrays. The normalized signal intensity for each element was compared between the tumor and NAT samples from each pair of patient samples and expressed as a log ratio of the tumor to normal adjacent sample.

Northern Analysis. *mir-84* northern blots were performed as described (Johnson et al., (2003) Dev Biol 259, 364-379). For human tissues, 1 ug of total RNA from the tumor and normal adjacent tissues of patients 1 and 5 were fractionated by PAGE using a 15% denaturing polyacrylamide gel. The RNA was transferred to a positively charged nylon membrane by electroblotting at 200 mA in 0.5X TBE for 2 hours. The Northern blot was dried and then incubated overnight in 10 ml of ULTRAhyb-Oligo (Ambion) with 10^7 cpm of a radio-labeled transcript complementary to *let-7c*. The blot was washed 3 X 10 min at room temperature in 2X SSC, 0.5% SDS and then 1X15 min at 42°C in 2X SSC, 0.5% SDS. Overnight phosphorimaging using the Storm system (Amersham), revealed *let-7c*. The process was repeated using a radio-labeled probe for 5S rRNA.

Lung tumor protein/northern/mRNA analysis. Total RNA and protein were isolated from tumor and normal adjacent tissue samples from three lung cancer patients using the mirVana PARIS Kit (Ambion). *let-7* miRNA and U6 snRNA were measured using the Northern procedure described above. NRAS and B-actin mRNA as well as 18S rRNA were quantified by real-time RT-PCR using primers specific to each of the target RNAs. RAS and GAPDH protein were measured by Western analysis using the RAS antibody described above and an antibody for GAPDH (Ambion).

Accession numbers. The following sequences were searched for 3'UTR LCSs: *Hs KRAS* (Genbank M54968), *Hs HRAS* (NM176795), *Hs NRAS* (BC005219). NRAS is known to exist as a 2 Kb and a 4.3 Kb form.

BC005219 represents the short form with a 1151 nt polyadenylated 3'UTR. Two human EST clones (Genbank BU177671 and BG388501) were sequenced to obtain additional *NRAS* 3'UTR sequence. This revealed that the *NRAS* 3'UTR exists in a 3642 nt polyadenylated 3'UTR version, utilizing an alternative polyadenylation and cleavage site, 2491 nt downstream of the first. This presumably corresponds to the long *NRAS* form. The sequence was deposited with accession numbers AY941100 and AY941101.

Results

Additional targets including *let-60/RAS* of the *let-7* miRNA in *C.*

elegans. The *let-7* miRNA is temporally expressed in *C. elegans* (Johnson et al., (2003) Nature 426, 845-849; Pasquinelli et al., (2000) Nature 408, 86-89; Reinhart et al., (2000) Nature 403, 901-906) where it down-regulates at least two target genes, *lin-41* (Slack et al., (2000) Molec Cell 5, 659-669) and *hbl-1* (Abrahante et al., (2003) Dev Cell 4, 625-637; Lin et al., (2003) Dev Cell 4, 639-650), mutations which lead to precocious terminal differentiation of seam cells. To better understand the role of *let-7* in *C. elegans* seam cell differentiation and its potential role in humans, additional targets of *let-7* were identified. A computational screen for *C. elegans* genes was performed with *let-7* family complementary sites (LCS) in their 3'UTR (Grosshans, H., et al., Dev Cell (2005) 8(3):321-30). One of the top scoring genes was *let-60*, encoding the *C. elegans* orthologue of the human oncogene *RAS*. 8 LCSs were identified in the 3'UTR of *let-60* with features resembling validated LCSs (Lin et al., (2003) Dev Cell 4, 639-650; Reinhart et al., (2000) Nature 403, 901-906; Slack et al., (2000) Molec Cell 5, 659-669; Vella et al., (2004) Genes Dev 18, 132-137) (Figure 4A) (SEQ ID Nos. 26, 27, 28, 29, 32, 33, 34 and 35). Many of the identified sites were found in the 3'UTR of *let-60* from the closely related nematode *C. briggsae* (Stein et al., (2003) PLoS Biol 1, E45) (Figure 4A, Figure 11 and Figure 12), suggesting that they are likely to be biologically significant. An additional three sites were found in the *let-60/RAS* coding sequence as well as 10 other non-conforming 3'UTR sites that may also bind to *let-7* family miRNAs (Figure 11) (SEQ ID Nos. 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 and 80).

let-7(n2853ts) loss of function (lf) mutants express reduced *let-7* miRNA and die by bursting at the vulva at the non-permissive temperature

(Reinhart et al., (2000); Slack et al., (2000) Molec Cell 5, 659-669). Loss of function mutations in two previously identified targets of *let-7*, *lin-41* and *hbl-1*, have the property of partially suppressing the *let-7* lethal phenotype (Abrahante et al., (2003) Dev Cell 4, 625-637; Lin et al., (2003) Dev Cell 4, 639-650; Reinhart et al., (2000) Nature 403, 901-906; Slack et al., (2000) Molec Cell 5, 659-669). It was found that post-embryonic reduction of function of *let-60* by feeding RNA interference (RNAi), also partially suppressed *let-7(n2853)* in a reproducible manner. While 5% of *let-7* mutants grown on control RNAi survived at the non-permissive temperature of 25°C (n=302), 27% of *let-7(n2853); let-60(RNAi)* animals survived (n=345). Thus, similar to other known *let-7* targets, *let-60* partially suppresses the *let-7(n2853)* lethal phenotype, suggesting that *let-7* lethality may at least partially be caused by over-expression of *let-60*. However *let-60(RNAi)* did not appear to suppress the *let-7* seam cell terminal differentiation defect and did not cause precocious seam cell terminal differentiation. In addition, wild-type animals subjected to *let-60(RNAi)* did not display typical lethal and vulvaless phenotypes associated with *let-60* alleles (Beitel et al., (1990) Nature 348, 503-509; Han et al., (1990) Genetics 126, 899-913; Han and Sternberg, (1990) Cell 63, 921-931). *let-60(RNAi)* resulted in approximately 80% knock down of *let-60* mRNA (Grosshans, H., et al., (2005) Dev Cell 8(3):321-30), suggesting that the remaining *let-60* is still sufficient for seam cell differentiation and vulval development. To verify the specificity of the *let-60(RNAi)*, it was shown that while *let-7(mn112)* adults all die, *let-60(n2021); let-7(mn112)* adults can live (see Experimental Procedures). Interestingly, *let-7(n2853); let-60(RNAi)* animals delivered a brood and could lay some eggs, suggesting that the vulval bursting phenotype of *let-7* was not suppressed merely because of the lack of a vulva.

let-60 and *let-7* are both expressed in hypodermal seam cells (Dent and Han, 1998 Post-embryonic expression pattern of *C. elegans let-60 ras* reporter constructs. Mech Dev 72, 179-182; Johnson et al., 2003 A microRNA controlling left/right neuronal asymmetry in *Caenorhabditis elegans*. Nature 426, 845-849). The *let-60* 3'UTR was fused behind the

Escherichia coli lacZ gene driven by the hypodermally-expressing *col-10* promoter. It was found that reporter gene activity is down-regulated around the L4 stage (Figure 5A), around the same time that *let-7* is expressed in the seam cells (Johnson et al., (2003) Nature 426, 845-849). In contrast, the same reporter gene fused to an unregulated control 3'UTR was expressed at all stages (Figure 5A) (Reinhart et al., (2000) Nature 403, 901-906; Slack et al., (2000) Molec Cell 5, 659-669; Vella et al., (2004) Genes Dev 18, 132-137; Wightman et al., (1993) Cell 75, 855-862). It was found that reporter down-regulation directed by the *let-60* 3'UTR depended on a wild-type *let-7* gene, since down-regulation failed in *let-7(n2853)* mutants (Figure 5B). Thus, multiple lines of evidence strongly suggest that *let-60* is negatively regulated by *let-7*. First, the *let-60* 3'UTR contains multiple elements complementary to *let-7*; second, the *let-60* 3'UTR directs down-regulation of a reporter gene in a *let-7* dependent manner; third, this down-regulation is reciprocal to *let-7* up-regulation in the hypodermis; and finally, *let-60* loss of function partially suppresses the *let-7* lethal phenotype.

The *let-7* family member *mir-84* is dynamically expressed in the vulval precursor cells. *let-60/RAS* is best understood for its role in vulval development (Wang and Sternberg, (2001) Curr Top Dev Biol 51, 189-220), however *let-7* has not been reported to be expressed in the vulva. In *C. elegans*, *let-7*, *mir-48*, *mir-84* and *mir-241* comprise the *let-7* family (Lau et al., (2001) Science 294, 858-862; Lim et al., (2003) Genes Dev 17, 991-1008; Reinhart et al., (2000) Nature 403, 901-906) (Figures 10A, 10B). Previous work demonstrated that a *let-7::gfp* fusion faithfully recapitulates the temporal expression of *let-7* and is temporally expressed in seam cell tissues affected in the *let-7* mutant (Johnson et al., (2003) Dev Biol 259, 364-379). The expression pattern of *mir-84*, the closest *let-7* relative, was examined by fusing 2.2 kilobases (Kb) of genomic sequence immediately upstream of the miR-84 encoding sequence to the green fluorescent protein (*gfp*) gene. *mir-84::gfp* was first observed in the somatic gonad in larval stage 1 (L1). In L3 animals, strong expression was observed in uterine cells including the anchor cell (AC), and weak dynamic expression was observed in the vulval precursor cells (VPCs) (Figures 6A-C). VPCs are multipotent

ventral hypodermal cells that generate the vulva during L3 and later stages (Sulston and Horvitz, (1977) Dev Biol 56, 110-156). VPCs adopt one of three fates depending on EGF signaling from the AC (Wang and Sternberg, (2001) Curr Top Dev Biol 51, 189-220). The cell closest to the AC, P6.p, receives the most LIN-3/EGF signal (Katz et al., (1995) Cell 82, 297-307) and adopts the primary (1^0) fate through activation of a *RAS*/MAPK signal transduction pathway (Beitel et al., (1990) Nature 348, 503-509; Han et al., (1990) Genetics 126, 899-913; Han and Sternberg, (1990) Cell 63, 921-931): P5.p and P7.p receive less LIN-3 as well as receiving a secondary lateral signal (Sternberg, (1988) Nature 335, 551-554) from P6.p, and adopt the secondary (2^0) fate: P3.p, P4.p and P8.p adopt the uninduced tertiary (3^0) fate. *mir-84::gfp* expression was observed during the early to mid L3 stage in all the VPCs except for P6.p, in which expression was rarely observed (Figure 6A). Subsequent VPC expression in the mid to late L3 stage was restricted to the daughters (Pn.px) of P5.p and P7.p with weaker GFP first appearing in the P6.p daughters just before their division into P6.pxx. Thereafter, equivalent expression was observed in the granddaughters (Pn.pxx) of P5.p, P6.p and P7.p. *mir-84::gfp* expression was observed in all the VPCs except for P6.p at the stage when their fate in vulval development is determined by signaling from the AC (Ambros, (1999) Development 126, 1947-1956) suggesting that *mir-84* could play a role in vulval cell fate determination. In the L4 stage, GFP expression was maintained in the AC and other uterine cells, appeared weakly in hypodermal seam cells, and was up-regulated to higher levels in many P5.p-P7.p descendants. A second *let-7* family member, *mir-48* was also expressed in non-P6.p VPCs, suggesting the potential for redundancy between *mir-48* and *mir-84* in the VPCs.

***mir-84* overexpression causes vulval and seam defects.** miR-84 was overexpressed by generating transgenic animals harboring a multi-copy array of a 3.0 Kb genomic DNA fragment that spans from 2.2 Kb upstream to 0.8 Kb downstream of the miR-84 encoding sequence (called *mir-84(+++)*). These animals expressed elevated levels of miR-84 and displayed abnormal vulval development phenotypes, including protrusion and bursting of the vulva (40% of animals, n=40). Consistent with *mir-*

84::*gfp* expression in seam cells, it was found that *mir-84(+++)* animals also exhibited precocious seam cell terminal differentiation and alae formation in the L4 stage, a characteristic seen in precocious developmental timing mutants. In fact, *let-7* over-expressing strains also exhibit precocious seam cell terminal differentiation in the L4 stage (Reinhart et al., (2000) Nature 403, 901-906). In contrast, animals carrying an array containing a construct identical to *mir-84(+++)* except for a 75 nucleotide (nt) deletion of sequences encoding the predicted pre-*mir-84* (Δ *mir-84* (+++)) did not display any vulval or seam defects, demonstrating that the phenotypes observed in *mir-84(+++)* are dependent on the miR-84 sequence.

A search was conducted for *let-7* family miRNA complementary sequences (LCS) in the 3'UTRs of all genes known to play a role in vulval development (Table 2). LCSs have the potential to bind all members of the *let-7* family, including *mir-84*. Approximately 11 vulval genes contained at least one LCS (Table 2), raising the possibility that the *let-7* family may regulate multiple genes in the vulva. In this analysis though, *let-60/RAS* stood out due to the high number of LCS sites.

Table 2. LCSs in the 3'UTRs of Known Vulval Genes

Gene	Chromosome	LCS in 3'UTR
<i>eor-1</i>	IV	Yes
<i>had-1</i>	V	Yes
<i>let-60</i>	IV	Yes
<i>lin-3</i>	IV	Yes
<i>lin-9</i>	III	Yes
<i>lin-11</i>	I	Yes
<i>lin-36</i>	III	Yes
<i>lin-39</i>	III	Yes
<i>lin-45</i>	IV	Yes
<i>mpk-1</i>	III	Yes
<i>sem-5</i>	X	Yes

***mir-84* overexpression partially suppresses *let-60/RAS* gain of function phenotypes.** *let-60/RAS* is active in P6.p following a *lin-3* EGF signal from the anchor cell that activates a MAPK signal transduction cascade transforming P6.p to the 1⁰ vulval fate (Han and Sternberg, (1990)

Cell 63, 921-931). Since *mir-84* is expressed in all VPCs except P6.p, the possibility that *mir-84* negatively regulates expression of *let-60/RAS* in cells not destined to adopt the 1⁰ fate was examined. Activating mutations in *let-60/RAS* cause multiple VPCs (including the non-P6.p VPCs) to adopt 1⁰ or 2⁰ fates leading to a multivulva (Muv) phenotype (Han et al., 1990). It was found that over-expression of *mir-84* partially suppressed the Muv phenotype of *let-60(gf)* mutations. In the study, 41% (n=51) of *let-60(ga89)* (Eisenmann and Kim, (1997) Genetics 146, 553-565) animals displayed a Muv phenotype, while only 13% (n=168) did so when also over-expressing *mir-84* from a multi-copy array (p<<0.0001 Chi square test). The same suppression was observed with a second *let-60(gf)* allele, *let-60(n1046)* (Han et al., (1990) Genetics 126, 899-913): 77% (n=39) of *let-60(n1046)* animals displayed a Muv phenotype, while only 50% (n=113) did so when also over-expressing *mir-84* (p<<0.0001 Chi square test). *let-60(n1046)* animals displayed an average of 1.54 pseudovulvae per animal compared to an average of 0.66 pseudovulvae per *let-60(n1046)* animal over-expressing *mir-84*. For both *let-60(gf)* alleles, animals exhibiting low mosaicism for the *myo-3::gfp* co-injection marker, were completely suppressed, suggesting that the partial suppression was likely due to mosaicism of the transgeneic array. Neither an empty vector control (TOPO) (n=111) (p=0.1435 Chi square test), nor the Δ *mir-84* (+++) array (n=129), suppressed the Muv phenotype of *let-60(n1046)*. For all *let-60(gf)* experiments, three independent lines behaved similarly (Figure 13C).

The *let-60/RAS* 3'UTR confines expression to P6.p. The promoter of *let-60/RAS* drives reporter expression in all VPCs (Dent and Han, (1998) Mech Dev 72, 179-182). However, the transgenic reporters used in this earlier work did not include the *let-60* 3'UTR. GFP was fused to the *let-60* 3'UTR and drove GFP expression in all the VPCs using the VPC-specific *lin-31* (Tan et al., (1998) Cell 93, 569-580) promoter (*gfp60*). In the late L2 and early L3 stages, GFP was expressed in all the Pn.p cells, but by mid to late L3 stages, GFP was largely restricted to the P6.p cell (Figure 6B), with some expression in the P5.p and P7.p cell descendants. A similar fusion

construct in which the *let-60* 3'UTR was replaced by the unregulated *unc-54* 3'UTR showed GFP expression in all Pn.p cells (Figure 6C). Since the *lin-31* promoter is active in all Pn.p cells (Tan et al., (1998) Cell 93, 569-580), this result demonstrates that the *let-60/RAS* 3'UTR is sufficient to down-regulate a reporter gene in the non-P6.p cells.

The *let-60* 3'UTR was replaced with the unregulated *unc-54* 3'UTR in a *let-60* genomic DNA fragment. While one could generate viable lines using a *let-60::let-60(+):let-60* 3'UTR construct at 10 ng/μl, it was not possible to generate viable transformants using this *let-60::let-60(+):unc-54* 3'UTR construct, even at 0.1 ng/μl. The results suggest that the removal of the *let-60* 3'UTR may severely over-express *let-60* and cause lethality.

***let-60/RAS* is a likely target of *mir-84* in the vulva.** Previous work has demonstrated that VPCs are sensitive to the levels of *let-60/RAS* (Beitel et al., (1990) Nature 348, 503-509; Han et al.(1990)). Animals carrying extra copies of the wild-type *let-60/RAS* gene display a Muv phenotype, where non-P6.p VPCs can adopt the 1° fate. The data strongly suggest that *mir-84* negatively regulates *let-60* in non-P6.p VPCs. First, *mir-84* is complementary to multiple sites in the *let-60* 3'UTR. Second, *mir-84* is expressed in a reciprocal manner to *let-60* in the VPCs. *mir-84* is largely absent from P6.p, at the same time as the *let-60* 3'UTR confines GFP expression mainly to the P6.p cell lineage. Finally, *mir-84* over-expression partially suppresses the effects of activating mutations in the *let-60* gene. *mir-84* modulates the expression of *let-60/RAS* in non-P6.p VPCs to reduce flux through the RAS/MAPK signaling pathway and hence decrease the likelihood that these cells will also adopt the 1° fate. However, *mir-84* is clearly not the only regulator of *let-60/RAS* in non-P6.p cells: *daf-12(rh61)* mutants do not express *mir-84* in any VPC (n=60 animals), and yet *daf-12(rh61)* animals do not display a Muv phenotype. Other known factors, e.g. *synmuv* genes, (Berset et al., (2001) Science 291, 1055-1058; Ceol and Horvitz, (2004) Dev Cell 6, 563-576; Hopper et al., (2000) Mol Cell 6, 65-75; Lee et al., (1994) Genes Dev 8, 60-73; Wang and Sternberg, (2001) Curr Top Dev Biol 51, 189-220; Yoo et al., (2004) Science 303, 663-666; Yoon et al.,

(1995) Science 269, 1102-1105) or unknown factors may also regulate *let-60/RAS* signaling in these cells.

The combined results provide strong evidence that *let-7* and *mir-84* regulate *let-60/RAS* expression through its 3'UTR in seam and vulval cells, cells in which they are all naturally expressed. Given that the 3'UTR of *let-60/RAS* contains multiple *let-7/mir-84* complementary sites, it is expected that this regulation is direct.

***let-7* complementary sites in human *RAS* 3'UTRs.** Numerous miRNAs are altered in human cancers (Calin et al., (2002) Proc Natl Acad Sci U S A 99, 15524-15529; Calin et al., (2004) Proc Natl Acad Sci U S A 101, 2999-3004; Michael et al., (2003) Mol Cancer Res 1, 882-891; Tam et al., (2002) J Virol 76, 4275-4286) and three of the best understood miRNAs, *lin-4* (Lee et al., (1993) Cell 75, 843-854), *let-7* (Reinhart et al., (2000) Nature 403, 901-906) and *bantam* (Brennecke et al., (2003) Cell 113, 25-36), all regulate cell proliferation and differentiation. The closest human homologues of *let-7* and *mir-84* are the *H.s. let-7* family miRNAs (Lagos-Quintana et al., (2002) Curr Biol 12, 735-739; Pasquinelli et al., (2000) Nature 408, 86-89). *let-60/RAS* (SEQ ID No. 87) is the *C. elegans* orthologue of human *HRAS* (SEQ ID No. 84), *KRAS* (SEQ ID No. 86), and *NRAS* (SEQ ID No. 85) (Figures 13A-B), which are commonly mutated in human cancer (Malumbres and Barbacid, (2003) Nat Rev Cancer 3, 459-465), including lung cancer. It was found that all three human *RAS* 3'UTRs contain multiple putative *let-7* complementary sites with features of validated *C. elegans* LCSs (SEQ ID Nos. 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 55) (Figures 4B-D). Many of these are conserved in rodents, amphibians and fish (Figure 14 and Figure 15 (SEQ ID Nos. 97, 98, 99, 100, 101, 102, 103)), suggesting functional relevance. The presence of putative LCSs in human *RAS* 3'UTRs indicates that mammalian *let-7* family members may regulate human *RAS* in a manner similar to the way *let-7* and *mir-84* regulate *let-60/RAS* in *C. elegans*.

Human *RAS* expression is regulated by *let-7* in cell culture.

Microarray analysis performed by Ambion, Inc. on six different cell

lines revealed that HepG2 cells express *let-7* at levels too low to detect by microarray analysis. Therefore, by request, Ambion, Inc., transfected HepG2 cells with a double-stranded (ds) RNA that mimics the *let-7a* precursor.

Consistent with the prediction that RAS expression is negatively regulated by *let-7*, immunofluorescence with a RAS-specific antibody revealed that the protein is reduced by approximately 70% in HepG2 cells transfected with exogenous *let-7a* miRNA relative to the same cells transfected with a negative control miRNA (Figure 7A). The protein expression levels of GAPDH and p21^{CIP1} were largely unaffected by the transfected *let-7a* and negative control pre-miRNAs (Figure 16A), indicating that *let-7a* regulation is specific to RAS. To confirm that the RAS antibody is specific to RAS protein in the transfected cells, HepG2 cells were also independently transfected with two exactly complementary siRNAs targeting independent regions of *NRAS*. Both siRNAs reduced cell fluorescence by more than 60% as compared to negative control siRNA-transfected cells (Figure 16B).

It was predicted that cells expressing native *let-7* may express less RAS protein and that inhibition of *let-7* may lead to derepression of RAS expression. To test this, by request, Ambion, Inc. transfected HeLa cells, which express endogenous *let-7* (Lagos-Quintana et al., (2001) Science 294, 853-858; Lim et al., (2003) Genes Dev 17, 991-1008), with anti-sense molecules designed to inhibit the activity of *let-7* (Hutvagner et al., (2004) PLoS Biol 2, E98; Meister et al., (2004) RNA 10, 544-550). Reducing the activity of *let-7* in HeLa cells resulted in an about 70% increase in RAS protein levels (Figure 7B). These results, combined with the reciprocal experiment using pre-*let-7* miRNAs discussed above, indicates that *let-7* negatively regulates the expression of RAS in human cells.

The 3'UTR of human *NRAS* and *KRAS* was fused to a luciferase reporter gene and these constructs transfected along with transfection controls into HeLa cells. *NRAS* contains two naturally occurring 3'UTRs that utilize alternative polyadenylation and cleavage sites, such that one of the 3'UTRs is 2.5 kb longer than the other. It was found that while the long *NRAS* 3'UTR strongly repressed reporter expression compared to an unregulated control 3'UTR (Figures 8A-B), the short *NRAS* 3'UTR led to only slight, but reproducible, repression of the reporter. The short 3'UTR contains 4 LCSs, while the long form contains 9 LCSs. The *KRAS* 3'UTR

also repressed the luciferase reporter (Figure 8A-B), while *HRAS* was not tested. The results demonstrate that the 3'UTRs of *NRAS* and *KRAS* contain regulatory information, sufficient to down-regulate the reporter.

As with the endogenous *RAS* experiments described above, the
5 reciprocal experiment was performed wherein HeLa cells were transfected with the *RAS* 3' UTR reporter constructs and the *let-7a* anti-sense inhibitor molecule (or a control scrambled molecule). Cells transfected with the *let-7a* inhibitor relieved repression exerted on the reporter relative to the control transfections (Figure 8C). Since, a loss in the extent of down-regulation is
10 observed when *let-7* is inhibited, these results strongly indicate that *let-7* regulates *NRAS* and *KRAS* in human cells through their 3'UTRs.

let-7, RAS and lung cancer. Like *let-60/ras*, human *RAS* is dose-sensitive, since over-expression of *RAS* results in oncogenic transformation of human cells (McKay et al., (1986) *Embo J* 5, 2617-2621; Pulciani et al.,
15 (1985) *Mol Cell Biol* 5, 2836-2841). It is plausible that loss of miRNA control of *RAS* could also lead to over-expression of *RAS* and contribute to human cancer. Recent work has mapped *let-7* family members to human chromosomal sites implicated in a variety of cancers (Calin et al., (2004) *Proc Natl Acad Sci U S A* 101, 2999-3004). In particular *let-7a-2*, *let-7c* and
20 *let-7g* have been linked to small chromosomal intervals that are deleted in lung cancers (Calin et al., (2004) *Proc Natl Acad Sci U S A* 101, 2999-3004), a cancer type in which *RAS* mis-regulation is known to be a key oncogenic event (Ahrendt et al., (2001) *Cancer* 92, 1525-1530; Johnson et al., (2001) *Nature* 410, 1111-1116).

25 miRNA microarray analysis was performed by Ambion, Inc. to examine expression levels of members of the *let-7* gene family in tissue from twenty-one different cancer patients, including twelve lung cancer patients with squamous cell carcinomas (stage IB or IIA). *let-7* is poorly expressed in lung tumors, as shown by expression of *let-7* in 21 breast, colon, and lung
30 tumors relative to associated normal adjacent tissue ("NAT"). Fluorescently labeled miRNA was hybridized to microarrays that included probes specific to *let-7a* and *let-7c*. Fluorescence intensities for the tumor and NAT were

normalized by total fluorescence signal for all elements and the relative average signal from the *let-7* probes in the tumor and normal adjacent samples are expressed as log ratios. *let-7a* and *let-7c* had similar profiles suggesting cross-hybridization between the two closely related miRNAs. *let-*
5 *7* miRNAs were reduced in expression in a number of the tumors relative to the normal adjacent tissue samples from the same patients. *let-7* was expressed at lower levels in all of the lung tumor tissues (Figure 9A), but only sporadically in other tumor types. A similar finding was independently discovered (Takamizawa et al., (2004) Cancer Res 64, 3753-3756). On
10 average, *let-7* was expressed in lung tumors at less than 50% of expression in the associated normal lung samples. Northern analysis was used to measure *let-7c* in the tumor and NAT samples for the two patients from which RNA was purified (samples represented by the first and fifth lung cancer bars in Figure 9B). Consistent with the microarray results by Ambion, Inc.,
15 northern analysis verified that the expression of *let-7c* was 65% lower in the tumor of patient #1 and 25% lower in the tumor of patient #5. Seven of eight examined samples also had on average 30% less *let-7g* expression in the tumor tissue (Figure 17). The miRNA arrays were used to compare the lung tumors and NAT included probes for 167 total miRNAs. The expression of
20 the vast majority of these were unchanged in the lung tumors indicating that *let-7* might be important in lung cancer. In theory, down regulation of *let-7* could result in up-regulation of RAS and thus induce or accentuate oncogenesis.

To test this hypothesis, Ambion, Inc. isolated total RNA and total
25 protein from the tumor and normal adjacent tissues of three new lung cancer patients with squamous cell carcinoma. The RNA samples were split and half was used for northern analysis to measure *let-7c* and U6 snRNA. The other halves of the RNA samples were used for real-time PCR to measure the *NRAS* mRNA, 18S rRNA, and B-actin mRNA. The protein samples were
30 used for western analysis to assess RAS and GAPDH protein levels. RAS protein was present in the tumors at levels at least ten-fold higher than in the normal adjacent samples from the same patients. Consistent with the

miRNA array results of Ambion, Inc. for other lung cancer samples, all three lung tumor samples had 4- to 8-fold lower levels of *let-7* than did the corresponding NAT samples. The first and third lung cancer samples had similar levels of *NRAS* mRNA in both the tumor and NAT while the second sample pair had significantly higher levels of *NRAS* mRNA in the tumor sample. RAS protein levels correlate poorly with *NRAS* mRNA levels but very well with *let-7* levels, suggesting that the expression of the oncogene is significantly influenced at the level of translation, consistent with the known mechanism of *let-7* in invertebrates.

10 The reciprocal expression pattern between *let-7* and RAS in cancer cells closely resembles what was seen with *let-7* and RAS in *C. elegans* and in the human tissue culture experiments. The correlation between reduced *let-7* expression and increased RAS protein expression in the lung tumor samples indicates that one or more members of the *let-7* gene family
15 regulates *RAS* expression *in vivo* and that the level of expression of the miRNA is an important factor in limiting or contributing to oncogenesis.

 These results demonstrate that the *let-7* miRNA family negatively regulates *RAS* in two different *C. elegans* tissues and in two different human cell lines. Strikingly, *let-7* is expressed in normal adult lung tissue
20 (Pasquinelli et al., (2000) *Nature* 408, 86-89), but is poorly expressed in lung cancer cell lines and lung cancer tissue (Takamizawa et al., (2004) *Cancer Res* 64, 3753-3756). The expression of *let-7* inversely correlates with expression of RAS protein in lung cancer tissues, suggesting a possible causal relationship. In addition, over-expression of *let-7* inhibited growth of
25 a lung cancer cell line *in vitro* (Takamizawa et al., (2004) *Cancer Res* 64, 3753-3756), suggesting a causal relationship between *let-7* and cell growth in these cells.

 These results demonstrate that the expression of the *RAS* oncogene is regulated by *let-7* and that over-expression of *let-7* can inhibit tumor cell line
30 growth. The combined observations that *let-7* expression is reduced in lung tumors, that several *let-7* genes map to genomic regions that are often deleted in lung cancer patients, that over-expression of *let-7* can inhibit lung tumor

cell line growth, that the expression of the *RAS* oncogene is regulated by *let-7*, and that *RAS* is significantly over-expressed in lung tumor samples strongly implicates *let-7* as a tumor suppressor in lung tissue and provides evidence of a mechanism forming the basis for treatment.

5 **Example 2. Expression patterns of *let-7* and *mir-125*, the *lin-4* homologue.**

It was found that both *let-7* and *mir-125*, the *lin-4* homologue, are expressed in a variety of adult tissues, with prominent expression in the lung and brain. Interestingly, both are expressed at low levels in the pancreas and
10 testis. Past work has shown that the *C. elegans lin-4* and *let-7* miRNAs are temporally expressed (Reinhart, B., et al., (2000) 403:901-906 and
Feinbaum, R. and V. Ambros, (1999) Dev Biol 210(1):87-95). Similarly, it has been shown that the mammalian homologues for these miRNAs are also temporally expressed during mouse development. Northern blots reveal that
15 *let-7* and *mir-125* have very similar expression profiles. Both become expressed at around day E9.5. Interestingly, this coincides approximately with the time of lung organogenesis, and when other major organs begin to develop.

An *in situ* protocol using an oligonucleotide based on the mouse
20 sequence of the *let-7c* miRNA, which has been digoxigenin-labeled, has been developed. Since it has been confirmed by northern analysis that *let-7c* is expressed at E12.5, frozen sections taken from similarly aged embryos have been analyzed. Preliminary results show *let-7* is expressed in the lung epithelium by *in situ*. In addition to other cancers, these results indicate that
25 *let-7* is useful as a therapeutic for lung cancer therapy because *let-7* is a natural compound in lung cells.

Example 3. Effect of Inhibition and Overexpression of *let-7* in Lung Cancer Cells.

Inhibition of *let-7* function in A549 lung cancer cells via transfection
30 (with anti-*let-7* molecules) causes increased cell division of A549 lung cancer cells (Figure 18A), while *let-7* over-expression (with transfected pre-*let-7*) caused a reduction in A549 cell number (Figure 18B). These results

are consistent with the tumor suppressing activity of *let-7*. Moreover, the *let-7* over-expression phenotype resembled that caused by MYC down-regulation (Figure 18A), suggesting that the effects of *let-7* on cell proliferation may also be through repression of MYC. Preliminary evidence indicates that MYC is also a direct target of *let-7* in human cells. Therefore, these results indicate that *let-7* may be a potential therapeutic in cancers with aberrant expression of MYC.

Example 4. *let-7* affects expression of MYC and BCL-2.

let-7 regulates RAS, MYC and BCL-2 protein levels, all three of which are major cancer oncogenes. Addition of *let-7* to HepG2 cells (that do not make *let-7* endogenously, reduces the expression of all three of these important oncoproteins (Figures 20, 7A, and 8). HeLa cells that make endogenous *let-7* express increased levels of all three oncoproteins when transfected with an anti-*let-7* inhibitor (Figures 20A, B and 7B). These results demonstrate that *let-7* represses expression of these genes. Multiple *mir-125* and *mir-143* complementary sites have been identified in the human *KRAS* and *BCL2* 3'UTRs and it is expected these oncomirs may also regulate these oncogenes in a manner similar to that seen with *let-7*. The results indicate that *let-7* is a master regulator of cancer pathways, regulating proliferation (RAS and MYC) and survival pathways (BCL2). It is possible that *let-7* also regulates telomerase (TERT) and angiogenesis (VEGF) pathways (Table 1). Since cancer is the result of multiple genetic mutations, these results indicate that introduction of *let-7* to cancer patients could repress the expression of multiple oncogenes, and provide an effective therapy (effectively a one drug cocktail).

It is understood that the disclosed invention is not limited to the particular methodology, protocols, and reagents described as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific

embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**

THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

Claims:

1. A use of a let-7 miRNA that binds to and inhibits expression of an oncogene containing a let-7 complementary site (LCS) selected from the group consisting of
5 NRAS, KRAS, and HRAS, for inhibiting the expression of the oncogene in a cell of a patient, wherein the let-7 miRNA is selected from the group consisting of a pri-miRNA, a pre-miRNA, and a mature miRNA.
2. The use of the let-7 miRNA of claim 1, wherein the let-7 miRNA is encoded by a
10 nucleic acid.
3. The use of the let-7 miRNA of claim 2, wherein the nucleic acid is located on a vector.
- 15 4. The use of the let-7 miRNA of claim 3, wherein the vector is selected from the group consisting of a plasmid, cosmid, phagemid, and virus.
5. The use of the let-7 miRNA of claim 4, wherein the vector further comprises one or more *in vivo* expression elements.
- 20 6. The use of the let-7 miRNA of claim 5, wherein the *in vivo* expression element is selected from the group consisting of a promoter, enhancer, and combinations thereof.
7. The use of the let-7 miRNA of any one of claims 1 to 6, wherein the miRNA is
25 from 21 nucleotides to 170 nucleotides in length.
8. The use of the let-7 miRNA of claim 7, wherein the miRNA is from 21 to 25 nucleotides in length.
- 30 9. The use of the let-7 miRNA of any one of claims 1 to 8, for treatment or prevention of cancer in the patient.
10. The use of the let-7 miRNA of claim 9, wherein the cancer is selected from the group consisting of lung cancer, pancreatic cancer, skin cancer, hematological

neoplasms, breast cancer, brain cancer, colon cancer, follicular lymphoma, bladder cancer, cervical cancer, endometrial cancer, esophageal cancer, gastric cancer, head and neck cancer, multiple myeloma, liver cancer, lymphomas, oral cancer, osteosarcomas, ovarian cancer, prostate cancer, testicular cancer, and thyroid cancer.

5

11. The use of the let-7 miRNA of claim 9, wherein the patient is undergoing one or more cancer therapies selected from the group consisting of surgery, chemotherapy, radiotherapy, thermotherapy, immunotherapy, hormone therapy and laser therapy.

10 12. The use of the let-7 miRNA of any one of claims 1-11, wherein the patient is a human.

13. An *in vitro* method for determining the sensitivity of a cancer to a let-7 miRNA comprising providing a let-7 miRNA, or increasing expression of the let-7 miRNA,
15 that binds to and inhibits the expression of a mRNA encoded by an oncogene containing a let-7 complementary site (LCS) selected from the group consisting of NRAS, KRAS, and HRAS in a cancerous or transformed cell and determining if expression of the oncogene is inhibited, wherein the let-7 miRNA is selected from the group consisting of a pri-miRNA, a pre-miRNA, and a mature miRNA.



20

14. A therapeutic composition for inhibiting the expression of an oncogene containing a let-7 complementary site (LCS) selected from the group consisting of NRAS, KRAS, and HRAS, comprising an effective amount of a let-7 miRNA and a pharmaceutically acceptable carrier that binds to and inhibits the expression of a
25 mRNA encoded by the oncogene in a transformed or cancerous cell in a patient, wherein the let-7 miRNA is selected from the group consisting of a pri-miRNA, a pre-miRNA, and a mature miRNA.

15. The composition of claim 14, wherein the let-7 miRNA is encoded by a nucleic
30 acid located on a vector selected from the group consisting of a plasmid, cosmid, phagemid, and virus, comprising one or more *in vivo* expression elements selected from the group consisting of a promoter, enhancer, and combinations thereof.

16. The composition of claim 14 or claim 15, wherein the let-7 miRNA is from 21 nucleotides to 170 nucleotides in length.
17. The composition of claim 16, wherein the let-7 miRNA is from 21 to 25
5 nucleotides in length.
18. The composition of any one of claims 14 to 17, wherein the cancer is selected from the group consisting of lung cancer, pancreatic cancer, skin cancer, hematological neoplasms, breast cancer, brain cancer, colon cancer, follicular
10 lymphoma, bladder cancer, cervical cancer, endometrial cancer, esophageal cancer, gastric cancer, head and neck cancer, multiple myeloma, liver cancer, lymphomas, oral cancer, osteosarcomas, ovarian cancer, prostate cancer, testicular cancer, and thyroid cancer.

1 / 28

**FIG. 1****FIG. 2**

2/28






FIG. 3

3/28



FIG. 4B

FIG. 4D


SUBSTITUTE SHEET (RULE 26)





6 / 28

**FIG. 6A****FIG. 6B**




7/28

**FIG. 6C**

8/28

**FIG. 7A****FIG. 7B**

9/28

FIG. 8A**FIG. 8B****FIG. 8C**

10 / 28




FIG. 9A




FIG. 9B






FIG. 10A





13 / 28



FIG. 12-1

14 / 28

FIG. 12-2

15 / 28

FIG. 13A



1	CAAGATACCTTT-TAAAGTTTGTCA---GAAAGAGCCACCTTTCAAGCTGCACCTGACCCCTGGTCTGACTTCCCTGGAGGAGAAAGTATTCCTGTTG	Hs NRAS short 3'UTR
1	CAAGATACCTTT-TAAAGTTTGTCA---GAAAGAGCCACCTTTCAAGCTGCACCTGACCCCTGGTCTGACTTCCCTGGAGGAGAAAGTATTCCTGTTG	Hs NRAS long 3'UTR
1	TAAGACCCCTTT-TAAAGTTTCTGTCTATCAAGAAAGAGCCACCTTTCAAGCTGCACCTGACCTGAGTCCCTGGTCTGACTTCCCTGGAGGAGAAAGTATTCCTGTTG	Rn nras 3'UTR
1	-----TAAAGTTTCTGTCTATCAAGAAAGAGCCACCTTTCAAGCTGCACCTGACCTGAGTCCCTGGTCTGACTTCCCTGGAGGAGAAAGTATTCCTGTTG	Mn nras 3'UTR
1	CAAGATATTTAAACAAAGTTCTGTCA---GAAAGAGCCACCTTTCAAGCTGCACCTGACCTGCTGACTTCCCTGGAGGAGAAAGTATTCCTGTTG	Cp nras 3'UTR
95	CTGCTTCAGTCTACAGAGAAGCTCCCTGCTACTTCCCAAGCT---CTCAGTATGTTTATGTAACAAATATCTCTATTTTGAGAGTCTTCAGAGAATAACTACC	Hs NRAS short 3'UTR
95	CTGCTTCAGTCTACAGAGAAGCTCCCTGCTACTTCCCAAGCT---CTCAGTATGTTTATGTAACAAATATCTCTATTTTGAGAGTCTTCAGAGAATAACTACC	Hs NRAS long 3'UTR
98	CTCTCTGCA---TCTCAGAGAAGCTCCCTGCTTCTGCTTCCCAAGCT---CTCAGTATGTTTATGTAACAAATATCTCTATTTTGAGAGTCTTCAGAGAATAACTACC	Rn nras 3'UTR
85	CTCTCTGCA---TCTCAGAGAAGCTCCCTGCTTCTGCTTCCCAAGCT---CTCAGTATGTTTATGTAACAAATATCTCTATTTTGAGAGTCTTCAGAGAATAACTACC	Mn nras 3'UTR
96	CTCTCTTCA---TCTCAGAGAAGCTCCCTGCTGTTTCTGCTTCCCAAGCT---CTCAGTATGTTTATGTAACAAATATCTCTATTTTGAGAGTCTTCAGAGAATAACTACC	Cp nras 3'UTR
191	TCCTCACTTGGCTGTCTGA-CCAGAGAA-TGCACCTCTTGTGTA-CTCCCTGTTATTTTCTGCCCTGGGTCTCTCC---ACAGCACAAACACACACCTCTG	Hs NRAS short 3'UTR
191	TCCTCACTTGGCTGTCTGA-CCAGAGAA-TGCACCTCTTGTGTA-CTCCCTGTTATTTTCTGCCCTGGGTCTCTCC---ACAGCACAAACACACACCTCTG	Hs NRAS long 3'UTR
192	TCCTCACTTGGCTGTCTGA-CCAGAGAAATGACCTT---GTTT-CTCCCTCAGTATGTTTCTGCTGCCCTGGGTCTCCCTAGAAACAAACACACACCTGCTGAG	Rn nras 3'UTR
179	TCCTCACTTGGCTGTCTGA-CCAGAGAAATGACCTT---GTTT-CTCCCTCAGTATGTTTCTGCTGCCCTGGGTCTCCCTAGAAACAAACACACACCTGCTGAG	Mn nras 3'UTR
186	TCCTCACTTGGCTGTCTGA-CCAGAGAAATGACCTTCTGTTATTTCTGCCCTGGGTCTCTCC---ACAGCACAAACACACACCTCTG	Cp nras 3'UTR
284	CCACCCACAGGTTTTCATCTGAAAGCAATTTCTATGCTCTGAACA-GAGAACCAAAACGCGTAACGTGAATTTCTATTGAAACACAGTGTCTTGAGCTCTA	Hs NRAS short 3'UTR
284	CCACCCACAGGTTTTCATCTGAAAGCAATTTCTATGCTCTGAACA-GAGAACCAAAACGCGTAACGTGAATTTCTATTGAAACACAGTGTCTTGAGCTCTA	Hs NRAS long 3'UTR
286	CTGGCT---TTTCTCTCCTGAAAGCAGTTTACAT-TGATGCAAGAGAACCAAACTATAGACAAACCAATCTGTTTCTAACAGTCTCTTACGCTCTA	Rn nras 3'UTR
270	CTGGCT---TTTCTCTCCTGAAAGCAGTTTACAT-TGATGCAAGAGAACCAAACTATAGACAAACCAATCTGTTTCTAACAGTCTCTTACGCTCTA	Mn nras 3'UTR
283	CCATCTC---TTTCTCTCCTGAAAGCAGTTTACAT-TGATGCAAGAGAACCAAACTATAGACAAACCAATCTGTTTCTAACAGTCTCTTACGCTCTA	Cp nras 3'UTR
381	AAGT-AGCAACTGCTGGTGATTTT-TTT-TTCTTTTACTGTGTGAACCTTGAACCT---ATGCTAAATTTTGGAGAAA-TGTCATAAAATTACTGTT	Hs NRAS short 3'UTR
381	AAGT-AGCAACTGCTGGTGATTTT-TTT-TTCTTTTACTGTGTGAACCTTGAACCT---ATGCTAAATTTTGGAGAAA-TGTCATAAAATTACTGTT	Hs NRAS long 3'UTR
378	AGGT-ACCAATGCTGGTGATTTTCCCTTTGGCCCCCACTGTGTGAACCTTGGCCCTT---GTTAGTTTGGGGGAAA-TGTAAATTTTATTCCTC	Rn nras 3'UTR
359	AGGT-ACCAATGCTGGTGATTTTCCCTTTGGCCCCCACTGTGTGAACCTTGGCCCTT---GTTAGTTTGGGGGAAA-TGTAAATTTTATTCCTC	Mn nras 3'UTR
358	AAGTTAGCAACGCTGGTGATTTTATTT-TCCTTTATTTTGAACCTTGGAACTGACCTATGTTAGTTTGGAGAAA-TGTCATAAAAGTACTGTT	Cp nras 3'UTR
470	TTGCC-AAGAATATAGTTATTTTGTCTGTTTGGTTTGTATTAATGTTATCGGCTCTATTCTCTAAACTGGCATCTGCTCTAGATTTCATAAAATACAAA	Hs NRAS short 3'UTR
470	TTGCC-AAGAATATAGTTATTTTGTCTGTTTGGTTTGTATTAATGTTATCGGCTCTATTCTCTAAACTGGCATCTGCTCTAGATTTCATAAAATACAAA	Hs NRAS long 3'UTR
467	TTCCCGAGAAATAGATTATGTTGTTGCTGATTTGCGGATTTTCAATGTCATGCTATATTCTCTGCTCTGTTATTCATAAAATGCAAA	Rn nras 3'UTR
447	TTCCCGAGAAATAGATTATGTTGTTGCTGATTTTCAATGTCATGCTATATTCTCTGCTCTGTTATTCATAAAATGCAAA	Mn nras 3'UTR
454	GTGCC-AAGAAAG	Cp nras 3'UTR

FIG. 14-1

18 / 28

FIG. 14-2

19 / 28

FIG. 14-3

171		Hs NRAS short 3'UTR
156	AGATAGATGTTACTCAATCTTGGGCTTTGTGCCAATAGCCCTTGTAATGATGTAACCTTAGAGTTTCCAGTATGTTCTTAAGCACACAGAGTTCTCAAA	Hs NRAS long 3'UTR
167	CCTTAGTAAACTG--CTGC-----CTTTTACAAACATAGGCTCAGTCAATCC--TCCCTGAGCTGCTTGACACACACTTGGCTCTGTTCTTTTAA	Rn nras 3'UTR
099	CCCTGGCCAACTTAAATGCT--GCCCTTTTACAAACA--AGGCTCAGTCACTCT--TCCCTGAGCTGCTT--GAGACCAGCACTTGGCTCTTCTTTGTTTAA	Mn nras 3'UTR
64		Cp nras 3'UTR

21 / 28

Hs NRAS short 3'UTR
Hs NRAS long 3'UTR
Rn nras 3'UTR
Mn nras 3'UTR
Cp nras 3'UTR

1171 TGGGCGCCAAATTCAGACTTGGACTATGTCCTTTGGATTACCTTAAGAAG
2254 TGTGTCGTAT----GACTGGTGGGATCTCTAAATAAGTTTATTAA
2256 TATGTCATAT----GACTGGTGGGATCTCTAAATAAGTTTATTAA
2190
464

FIG. 14-5

22/28

X.I. NRAS 3'UTR



CUUU UA	CU G	A U AG
AACAU CAA CCACUUACCUC	AAUU UGCU UUAUGCCUCA	AAC AAU GUUG CACCUC
UUGUA GUU GGUGGAUGGAGU	UGA AUGU GAUAUGGAGU	UUG UUG UGAU GUGGAGU
A U A	U UG G	AA UG A
LCS1	LCS2	LCS3

D.r. NRAS 3'UTR


U C	G U U
AAUUAUACAUUUAC CCUCA	AAU AAUAAUUAUU CCUCA
UGAUUAUGUGGAUG GGAGU	UUG UUGUUGAUGA GGAGU
U AU	AA U
LCS1	LCS2

FIG. 15


23 / 28

**FIG. 16A****FIG. 16B**

24 / 28

**FIG. 17**

25/28

**FIG. 18A**

26 / 28







FIG. 18B



28 / 28

**FIG. 20A****FIG. 20B**