

(19) **DANMARK**

(10) **DK/EP 2141982 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 01 N 63/04 (2006.01)** **A 01 H 17/00 (2006.01)** **A 01 N 65/00 (2009.01)**
A 01 P 17/00 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2019-01-07**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-09-12**
- (86) Europæisk ansøgning nr.: **08753855.9**
- (86) Europæisk indleveringsdag: **2008-04-28**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-01-13**
- (86) International ansøgning nr.: **NZ2008000094**
- (87) Internationalt publikationsnr.: **WO2008133533**
- (30) Prioritet: **2007-04-27 US 914656 P**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Grasslanz Technology Limited, Tennent Drive RD 4, Palmerston North 4474, New Zealand**
- (72) Opfinder: **PENNELL, Christopher, Gerald, Lee, 18 Bridget Lane, Rangiora 7400, Canterbury, New Zealand**
ROLSTON, Maurice, Phillip, 303 Tai Tapu Road, RD 2, Christchurch 7672, New Zealand
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **GRÆSBASERET FUGLEAFSKRÆKNINGSMIDDEL**
- (56) Fremdragne publikationer:
WO-A1-00/62600
WO-A1-2007/021200
WO-A2-02/13616
WO-A2-2004/106487
US-A- 6 072 107
US-A- 6 111 170
PENNELL C. ET AL: "The effect of grass-endophyte associations on feeding of Canada geese (Branta canadensis)", PROCEEDINGS OF THE NEW ZEALAND GRASSLAND ASSOCIATION, vol. 65, 1 January 2003 (2003-01-01), pages 239-244, XP055090896,
PENNELL C G L ET AL: "Development of a bird -deterrent fungal endophyte in turf tall fescue", NEW ZEALAND JOURNAL OF AGRICULTURAL RESEARCH,, vol. 53, no. 2, 1 June 2010 (2010-06-01), pages 145-150, XP009174691, ISSN: 0028-8233, DOI: 10.1080/00288231003777681
CONOVER M.R.: 'Impact of consuming tall fescue seeds infected with the endophytic fungus, Neotyphodium coenophialum, on reproduction of chickens' THERIOGENOLOGY vol. 59, no. 5-6, 2003, pages 1313 - 1323, XP008125305
CONOVER M.R. AND MESSMER T.A.: 'Feeding preferences and changes in mass of Canada geese grazing

Fortsættes ...

endophyte-infected tall fescue' THE CONDOR vol. 98, no. 4, 1996, pages 859 - 862, XP008131903

MADEJ C.W. AND CLAY K.: 'Avian seed preference and weight loss experiments: the effect of fungal

endophyte-infected tall fescue seeds' OECOLOGIA vol. 88, no. 2, 1991, pages 296 - 302, XP008125367

DESCRIPTION

FIELD OF THE INVENTION

[0001] This invention relates to a grass based avian deterrent. More specifically, the invention relates to a grass cultivar and endophyte combination that produces alkaloid toxins sufficient to deter avian species from the grass cultivar and endophyte combination.

DESCRIPTION OF THE RELATED ART

[0002] Avian populations can be undesirable in certain environments. Bird populations can have a devastating effect on horticultural and agricultural crops, cause severe damage to an aircraft in the event of a bird strike and can simply be an irritant, for example in recreational areas.

[0003] More specifically, bird populations can quickly destroy fruit crop(s) such as apricots, peaches, apples, kiwifruit and the like through feeding on the ripening fruit. Similarly, birds can also significantly reduce the harvest of seed or grains from various cereal and grain crops (e.g., wheat production).

[0004] For the aviation industry, the problem is highly significant as a bird strike incident can severely disrupt normal aircraft handling and can even be a significant causal factor in aircraft crashes.

[0005] In the above situations as well as in general, bird populations around waterways or in recreation areas and even around domestic housing can also be a nuisance. Other problems such as water and ground contamination from bird excrement can also be significant.

[0006] Various attempts have been made to deter birds from an area. These act on primary initial response mechanisms e.g., a loud noise or act using a secondary response, for example, making the bird ill by feeding the bird bait which includes a chemical irritant.

[0007] In more detail, primary deterrents include acoustical distress calls, gas guns, lights, lasers, dogs, falconry, kites and balloons which all work to frighten the birds from an area. Other methods such as sticky pastes, spikes, wires and netting may also be used.

[0008] Secondary deterrents include a (UV) light product known as ultra violet anthraquinone, methyl anthranilate (MA), taste aversion sprays and poisons such as methiocarb carbamate (Mesurol 50 HBT).

[0009] However, each of the above methods has disadvantages. Primary deterrents generally do not work over a longer term as birds 'learn' that the danger is not real. In addition, these

methods usually involve significant financial and labour costs to install and run the devices and arrangements used to deter the birds.

[0010] Secondary methods have the obvious problem of allowing at least a first strike by the birds. In the case of baits, the birds may not all eat the bait and hence it may take some time before the full deterrent effect is obtained. In addition the deterrents need to be re-applied on a regular basis and in some environments, simply cannot be used for risk of the chemical agents contaminating food or water.

[0011] A further problem is that the above methods need to be carried out numerous times during a given period of time. For example, baits may need to be used on numerous occasions to achieve the deterrent effect and, later another cycle may need to be completed to deal with new populations of birds arriving.

[0012] Pennell and Rolston (2003, Proceedings of the New Zealand Grassland Association 65:229-238) investigated the effect of feeding Canada geese with ryegrass herbage and seed infected with the endophyte *Neotyphodium lolii*. There was an 80% rejection of the seed containing the endophyte on day 2, compared to endophyte free seed. The rejection was still observed after a 3 month endophyte-free feeding period.

[0013] Therefore, the subject matter of the present invention is drawn to providing the public with useful compositions and their uses as well as methods for addressing such problems.

SUMMARY OF THE INVENTION

[0014] The present invention is defined as set out in the claims.

[0015] For the purposes of this specification, the term 'endophyte' or grammatical variations thereof refer to fungi living within cultivated grasses or axenic culture medium.

[0016] The term 'cultivated grasses' or 'cultivar' or grammatical variations thereof refer to varieties of grasses that have been created or selected intentionally and maintained through cultivation.

[0017] The grass cultivar may also be a synthetic grass cultivar. The term 'synthetic grass cultivar' refers to the grass cultivar being produced through selective breeding techniques including selection and development from an uncultivated population. For example, a synthetic grass cultivar refers to where -

1. (a) reproducible units are from a cross-pollinated crop which can encompass clones or inbred grasses;
2. (b) materials used are selected from their performance in combining ability or progeny tests;

3. (c) the cultivar is constituted by random inter-mating of the units;
4. (d) the units are maintained so that the synthetic can be reconstituted.

[0018] The term 'herbage' refers to the plant generally including the stems, pseudostems, leaves, flowers and seeds.

[0019] The term 'combination' or grammatical variations thereof refer to the combination of an endophyte and a grass cultivar infected with endophyte fungi.

[0020] The term 'pests' for the purpose of this invention refers to insects, birds or other animals that destroy or harm plants, crops, food, or even endanger or threaten human safety.

[0021] The term 'low temperature' for the purpose of this invention include temperatures up to and including 7°C. While the term 'high temperature' for the purpose of this invention, include temperatures from 27°C and above.

[0022] The term 'sufficient levels' used in the context of alkaloid compounds in the grass refers to the levels being sufficient to exhibit a deterrent effect on pests as defined above.

[0023] The term 'AR601, AR602, AR603, AR604, AR4, AR5, AR8 or AR94 endophyte' refers to an endophyte which can be distinguished from other endophytes through their particular microsatellite pattern. In particular, Example 14 below summarizes these endophytes and the simple sequence repeat sizes at various alleles. Also described is the methodology for achieving the microsatellite pattern. The endophyte strains have also been deposited at The National Measurement Institute (NMI) (1 Suakin Street, Pymble, NSW, Australia,) on 23 July 2007 and accorded with accession numbers: V07/029058, V07/029059, V07/029060, V07/029061, V07/029054, V07/029055, V07/029056, V07/029057, respectively.

[0024] The present invention provides the use of an endophyte and grass cultivar combination to repel avian species from the grass combination. The grass combination is characterised as a combination that includes a level of at least one ergot-peptide alkaloid sufficient to deter the avian species from the grass combination.

[0025] The inventors have found that by manipulating the properties of grass and endophyte combinations, endophyte and grass combinations can be produced and selected to repel avian species from the grass. This has significant advantages as the grass is a relatively permanent fixture and easy to maintain once it is sown. Therefore the deterrent can be used for minimal cost and labour.

[0026] The present disclosure describes an endophyte and grass cultivar combination that repels avian species from the combination, characterised in that the endophyte in the combination produces ergovaline alkaloid compound sufficient to repel the avian species from the cultivar herbage and seeds produced therefrom.

[0027] The present disclosure describes an endophyte and grass cultivar combination including an endophyte selected from the group consisting of: AR601, AR602, AR603, AR604, AR4, AR5, AR8 and AR94 (Deposit Nos. V07/029058, V07/029059, V07/029060, V07/029061, V07/029054, V07/029055, V07/029056 V07/029057, and combinations thereof; characterised in that avian species are repelled from the cultivar herbage and seeds produced therefrom.

[0028] Preferably, the endophyte may be from the genera *Neotyphodium*. More preferably, the endophyte may be from the *Neotyphodium coenophialum* species. More preferably, the endophyte may be selected from the group consisting of: AR601, AR602, AR603, AR604, AR4, AR5, AR8, AR94 (Deposit Nos. V07/029058, V07/029059, V07/029060, V07/029061, V07/029054, V07/029055, V07/029056, V07/029057, and combinations thereof. For example, as noted above, a characteristic in determining repellent effect is the amount of ergovaline alkaloid. Other endophytes can be found that also produce levels of ergovaline alkaloid compounds sufficient to result in a repellent effect. Preferably, the ergovaline alkaloid compound or compounds are present in at least a portion of the cultivar herbage. The ergovaline alkaloid may be present in at least the pseudostem and/or seeds of the cultivar. Preferably, the combinations as described above include repellent levels of ergovaline throughout the herbage of the plant including the pseudostem as well as the leaf, stems and seeds. Levels are provided with respect to the pseudostem to illustrate alkaloid spread through the herbage. It should be apparent to one of skill in the art that the ergovaline alkaloid (and other alkaloids) can be present in other parts of the grass cultivar herbage and that reference to the pseudostem should not be seen as being limiting.

[0029] Although reference is made in this specification with respect to ergovaline, it should be appreciated by those skilled in the art that the disclosure may be applied to other ergot-peptide alkaloid compounds.

[0030] The invention also provides an endophyte and a *Festuca* grass cultivar combination, or grass cultivars exemplified by *Festuca* characteristics, that repels avian species from the combination, characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0031] Preferably, the *Festuca* grass cultivar may be tall fescue (*Festuca arundinacea* , or its synonyms) meadow fescue (*F. pratensis*), chewing fescue (*F. rubra*), and the like or other *Festuca* grass cultivars with similar characteristics, or other grass species such as brown tops (*Agrostis tenuis*) and other *Agrostis* species.

[0032] The endophytes for *Festuca* grass cultivar of the invention are endophytes selected from the group consisting of: AR601, AR602, AR603, AR604, (Deposit Nos. V07/029058, V07/029059, V07/029060, V07/029061) and combinations thereof. However, it should be appreciated that other endophytes may be used together with other grass species which produce similar levels of ergovaline.

[0033] The invention also provides an endophyte and a *Lolium* grass cultivar combination, or grass cultivars exemplified by *Lolium* characteristics, that repels avian species from the combination, characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0034] In preferred embodiments, the grass may be perennial ryegrass or *Lolium perenne*. In some other embodiments other *Lolium* cultivars may be used including *L. multiflorum*, *L. hybridum* and hybrids thereof.

[0035] The endophytes for *Lolium* grass cultivars of the invention are endophytes selected from the group consisting of: of AR4, AR5, AR8, AR94 (Deposit Nos. V07/029054, V07/029055, V07/029056, V07/029057 and combinations thereof. However, it should be appreciated that other endophytes and *Lolium* cultivars may be used which produce similar levels of ergovaline.

[0036] Preferably, the grass and endophyte combination repel avian species via a secondary deterrent effect. More preferably, the repellent effect is a secondary effect resulting from a post digestive feedback (PDF) mechanism.

[0037] The avian species may be birds including waterfowl such as geese, seabirds such as seagulls and city birds (*Passeriformes* bird types) such as sparrows, and finches. The avian species can include Lapwings which can pose a problem around airports.

[0038] The inventors have found that a level of at least 3 ppm of the ergovaline alkaloid compound is sufficient to cause a deterrent effect. Preferably, the ergovaline may be present in a range from 3ppm to 100ppm.

[0039] Preferably, the endophyte may also confer resistance to the combination against biotic and abiotic stresses.

[0040] Biotic stresses may include resistance to attack from insects and pests, including resistance to Grass Grub, (*Costelytra zealandica*) Argentine Stem Weevil (*Listronotus bonariensis*) and sucking insects (for example aphids of the *Rhopalosiphum padi* species).

[0041] Abiotic stresses may include resistance to drought or dry periods as well as resistance to high or low temperature climates.

[0042] The inventors have found that by selecting an endophyte that also produces peramine and loline alkaloids, the grass cultivar may not only deter avian species, but also have a high tolerance to biotic and abiotic stresses. This is particularly advantageous in locations where the grass is subject to little care and attention and where the grass is not likely to be grazed by animals, such as grass adjacent an airport or other areas where birds are attracted to, for example areas where plant predatory insects are formed.

[0043] Preferably, the endophyte may also produce peramine alkaloids compounds within at least a portion of the cultivar herbage and seeds produced therefrom. Preferably, the peramine alkaloid may be present in a range from 1ppm to 100ppm or more. More preferably, the peramine alkaloid may be present at level of at least 1 ppm.

[0044] Preferably, the endophyte may also produce loline alkaloids within at least a portion of the cultivar herbage and seeds produced therefrom. Preferably, the loline alkaloid may be present in a range from 1ppm to 9500ppm or more. More preferably, the loline alkaloid may be present at level of at least 1500 ppm.

[0045] Preferably, the endophyte may also produce lolitrem alkaloids compounds within at least a portion of the cultivar herbage and seeds produced therefrom. Preferably, the lolitrem alkaloid may be present in a range from 1ppm to 50 ppm or more. More preferably, the lolitrem alkaloid may be present at level of at least 1 ppm.

[0046] The present disclosure describes the use of an endophyte and grass cultivar combination that repels avian species from the combination, characterised in that the endophyte in the combination produces ergovaline alkaloid compound sufficient to repel the avian species from the cultivar herbage and seeds produced therefrom.

[0047] The present disclosure describes the use of an endophyte and grass cultivar combination including an endophyte selected from the group consisting of: AR601, AR602, AR603, AR604, AR4, AR5, AR8 and AR94 (Deposit Nos.V07/029058, V07/029059, V07/029060, V07/029061, V07/029054, V07/029055, V07/029056, V07/029057, and combinations thereof; characterised in that avian species are repelled from the cultivar herbage and seeds produced therefrom.

[0048] The present disclosure describes the use of an endophyte and a *Festuca* grass cultivar combination, or grass cultivars exemplified by *Festuca* characteristics, that repels avian species from the combination, characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0049] The present disclosure describes the use of an endophyte and a *Lolium* grass cultivar combination, or grass cultivars exemplified by *Lolium* characteristics, that repels avian species from the combination, characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0050] The present invention provides a method of repelling avian species from an area of land by planting an endophyte and grass cultivar of the invention on or adjacent the land from which the avian species are to be repelled from characterised in that the endophyte in the combination produces ergovaline alkaloid compound sufficient to repel the avian species from the cultivar herbage and seeds produced therefrom.

[0051] The present disclosure describes a method of repelling avian species from an area of land by planting an endophyte and grass cultivar on or adjacent the land from which the avian species are to be repelled from characterised in that the endophyte and grass cultivar combination including an endophyte selected from the group consisting of: AR601, AR602, AR603, AR604, AR4, AR5, AR8, AR94 (Deposit Nos.V07/029058, V07/029059, V07/029060, V07/029061, V07/029054, V07/029055, V07/029056, V07/029057, and combinations thereof; characterised in that avian species are repelled from the cultivar herbage and seeds produced therefrom.

[0052] The present disclosure describes a method of repelling avian species from an area of land by planting an endophyte and *Festuca* grass cultivar or grass cultivars exemplified by *Festuca* characteristics, on or adjacent to the land from which the avian species are to be repelled from characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0053] The present disclosure describes a method of repelling avian species from an area of land by planting an endophyte and *Lolium* grass cultivar or grass cultivars exemplified by *Lolium* characteristics, on or adjacent the land from which the avian species are to be repelled from characterised in that the endophyte in the combination produces a level of at least 3ppm of ergovaline alkaloid compound within a portion of the cultivar herbage and seeds produced therefrom.

[0054] The present invention provides a method of screening an endophyte and grass cultivar combination of the invention, for the combinations ability to repel avian species by the steps of:

1. (a) cultivating the combination;
2. (b) measuring the level of an ergovaline alkaloid in the cultivated combination; and;
3. (c) selecting the combination if the level of ergovaline is above at least 3ppm.

[0055] The present disclosure describes a method of screening an endophyte and grass cultivar combination, for the combinations ability to repel avian species by the steps of:

1. (a) cultivating the combination;
2. (b) measuring the level of an ergovaline alkaloid in the cultivated combination; and;
3. (c) selecting the combination if the level of ergovaline is above at least 3ppm;

characterised in that the endophyte in the combination endophyte selected from the group consisting of: AR601, AR602, AR603, AR604, AR94 and AR4, AR5, AR8 (Deposit Nos.V07/029058, V07/029059, V07/029060, V07/029061, V07/029057, V07/029054, V07/029055, V07/029056), and combinations thereof.

[0056] The present disclosure describes a method of screening an endophyte and *Festuca*

grass cultivar combination, or grass cultivars exemplified by *Festuca* characteristics, for the combinations ability to repel avian species by the steps of:

1. (a) cultivating the combination;
2. (b) measuring the level of an ergovaline alkaloid in the cultivated combination; and;
3. (c) selecting the combination if the level of ergovaline is above at least 3ppm.

[0057] The present disclosure describes a method of screening an endophyte and *Lolium* grass cultivar combination, or grass cultivars exemplified by *Lolium* characteristics, for the combinations ability to repel avian species by the steps of:

1. (a) cultivating the combination;
2. (b) measuring the level of an ergovaline alkaloid in the cultivated combination; and;
3. (c) selecting the combination if the level of ergovaline is above at least 3ppm.

[0058] It should be appreciated from the above description that there is provided a method of selecting and using an endophyte to repel birds from the grass and endophyte combination or area on which the combination grows. The invention not only provides identification of specific endophytes that confer repellent properties but also describes methods to select new endophytes for use as a repellent. The invention provides methods to overcome difficulties in previous studies which allude to an endophyte and grass combination having a repellent effect but which use wild type endophyte and do not provide any guidance on methods of selection and key characteristics needed in a commercial product.

BRIEF DESCRIPTION OF THE DRAWINGS

[0059]

Figure 1 illustrates the percentage of seeds of different endophyte treatment consumed by 'naïve geese' in a feeding experiment;

Figure 2 illustrates the percentage of seeds of different endophyte treatment consumed by 'learned geese' in a feeding experiment;

Figure 3 illustrates the percentage of seeds of different endophyte treatment consumed by 'learned geese' in a feeding experiment conducted after 3 months from the initial learning feeding;

Figure 4 illustrates the amount in grams of herbage consumed in a further feeding experiment of 'learned geese';

Figure 5 illustrates the average amount of plant material (dry weight) consumed by the geese

per day in a no-choice feed experiment of nil endophyte infected plant material and endophyte infected plant material.

Figure 6 illustrates the amount of weight loss experienced by geese feeding on endophyte infected plant material.

Figure 7 illustrates the percentage of seed from five different grass and endophyte cultivars consumed in three different feeding sessions, by the same group of geese;

Figure 8 illustrates the alkaloid levels of three different alkaloids, ergovaline, lolitrem and peramine from four grass and endophyte cultivars against against a nil endophyte seed;

Figure 9 illustrates ergovaline concentration in different cultivars infected with the same wild type endophyte strain;

Figure 10 illustrates the differences of leaf colonisation of four different tall fescue cultivars infected with three different endophyte strains;

Figure 11 illustrates peramine expression in six different ryegrass endophyte strains in the same ryegrass cultivar;

Figure 12 illustrates ergovaline expression in six different ryegrass endophyte strains (same to that in Figure 11) in the same ryegrass cultivar;

Figure 13 illustrates peramine expression in five different endophyte strains, four in tall fescue grass cultivars and one in a ryegrass cultivar;

Figure 14 illustrates loline expression in five different endophyte strains, four in tall fescue grass cultivars and one in a ryegrass cultivar;

Figure 15 illustrates ergovaline expression in five different endophyte strains, four in tall fescue grass cultivars and one in a ryegrass cultivar;

Figure 16 illustrates the concentration of ergovaline in various plant anatomy of "Jackal" grass cultivar inoculated with either AR601 or AR604;

Figure 17 illustrates the concentration of lolines in various plant anatomy of "Jackal" grass cultivar inoculated with either AR601 or AR604;

Figure 18 illustrates the percentage of pellets consumed by gulls for different treatments;

Figure 19 illustrates the level of nil endophyte feed eaten by finches compared to endophyte containing feed

Figure 20 is a graph showing the average amount of seed consumed for finches between nil endophyte feed consumed versus endophyte containing feed consumed;

Figure 21 is a graph showing the total bird numbers present in the endophyte-enhanced plots with the control plots per month for the Christchurch International Airport (CIAL) trial;

Figure 22 illustrates the relative grazing score 1st assessment (0=not grazed). LSD 5% shown on right for the Rangiora Sewage treatment plant trial;

Figure 23 illustrates the relative grazing score 5th assessment for the Rangiora Sewage treatment plant trial; and

Figure 24 illustrates the average grazing scores (mean of 8 grazings) of each endophyte and grass cultivar combination for the Rangiora Sewage treatment plant trial.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0060] The present invention relates to a grass based avian deterrent. More specifically, the invention relates to a grass cultivar and endophyte combination that produces alkaloid toxins sufficient to deter avian species from the grass cultivar and endophyte combination. As mentioned above, various primary and secondary deterrence methods have been attempted. However, a need exists for improved deterrence methods.

[0061] One secondary deterrent method is a repellence created via the food source of birds inducing a "post digestion feed back" (PDF) response. In feeling ill, the bird, through education and memory, avoids the earlier feeding area. PDF, as described by Conover et al., 1996 is the deterrent effect resulting from an animal eating something that is not initially associated with tastes but causes malaise over time and can affect the animals' long term food preference.

[0062] Studies have documented the PDF responses in geese and other pests. For example, a report published by The Berryman Institute (Gosser et al, 1997,) describes methods for deterring Canadian geese by planting grass that is less palatable to the geese, therefore discouraging the geese from feeding. The report describes that geese dislike tall fescue grasses, specifically the Kentucky-31 (K-31) cultivar, as the grass cultivar contains endophytes which produce toxic alkaloids that cause the PDF repellent effect. The report illustrates the PDF effect in relation to the K-31 variety but does not consider what particular endophytes cause this effect or the way the effect is achieved.

[0063] Durham et al., 1998 describes the use of endophyte infected grass seeds to cause a repellent effect in rodents. Durham et al., 1998 showed that over a number of 'learning experiments', vole rodents avoid consuming the endophyte infected tissue.

[0064] Conover et al., 1996, describes a study where Canadian geese fed K-31 tall fescue grass in a choice situation showed over time a preference to K-31 without endophyte present.

[0065] The above studies show a link between endophytes and animal deterrent, but no further details on this effect have been found. For example, there is no teaching as to what alkaloids cause the deterrent effect and how the deterrent effect can be maximised. Further, none of the

studies discuss or mention other grass cultivars and/or endophyte combinations other than the use of K-31, which had been naturally infected with the endophyte *Acremonium coenophialum* (later reclassified as *Neotyphodium coenophialum*). Given the varying habitats and climates where bird problems are prevalent and the K-31 characteristics, K-31 infected with *N. coenophialum* is unlikely to be the most suitable grass cultivar or endophyte combination to use in all habitats and climates.

[0066] Fungal endophytes of the genus *Neotyphodium* infect a number of temperate climate Pooideae grasses. The *Neotyphodium* endophytes can produce alkaloids such as ergovaline, peramine and lolines which are considered to confer degrees of pest and disease protection upon the plants in which they naturally occur. U.S. Patent 6,111,170 and U.S. Patent 6,072,107 provide further background on grass and endophyte combinations.

[0067] The instant invention relates to controlling and selecting an endophyte and grass combination to maximise the deterrent effect, where the combination is also suitable for varying habitats and climates. The invention also relates to methods that can be used to develop a robust grass and endophyte combination that is well understood, and a selection method that can be used to select combinations that are useful in a variety of habitats and climates.

[0068] The present invention is now explained in more detail using examples below. In these experiments, the inventors show that a bird post digestive feedback (PDF) reaction occurs as measured by the learned behaviour observed.

1. Target Avian Species: Canadian Geese (*Branta canadensis*)

[0069] Approximately 50 geese were obtained from the wild. The geese were wing clipped and contained in a fenced off area to become familiar with their surroundings with little human contact. No endophytic material, either in seed or forage, was fed to these geese for a period of three weeks. Body weights were maintained with feed of crushed barley and Timothy grass forage. The geese were then split into various groups, for the two trials assessing the PDF response to inoculated endophyte seed (Example 1) and inoculated endophyte herbage (Example 2).

EXAMPLE 1 - SEED, CAFETERIA CHOICE TRIAL

[0070] The inventors ran a series of cafeteria choice trials' where the gaggle of approximately 20 geese chose between (i) seed without endophyte infection (Nil-endophyte); or (ii) seed that had been inoculated with endophyte A; or (iii) seed that had been inoculated with endophyte B. This tested the PDF and learnt behaviour of post digestion feed back in geese in seed feeding trials.

Seed and Endophyte Combinations Tested

[0071] Three different seed and endophyte combinations were tested during this experiment. The seed from the perennial ryegrass cultivar, Kingston, was inoculated with either endophyte A (Wild Type) or with endophyte B (Endosafe AR5).

[0072] Therefore, the three different seed and endophyte combinations that were tested consisted of:

1. (i) Kingston seed without endophyte (Nil-Endophyte);
2. (ii) Kingston seed inoculated with Endophyte A (Wild type endophyte); and
3. (iii) Greenstone seed inoculated with Endophyte B (AR 5 endophyte).

Run 1 - Day 1;

[0073] The seed was distributed in trays that were designed so only the head and necks of the geese could reach the food source. During the trials, the geese were feed a selection of 500g from each of the seed selections, as notes above in points (i) to (iii) above.

[0074] The amount of test seed consumed during the trials was determined by weighting the trays and determining the amount of remaining seed after a two hour feeding period. At the end of this first run, the geese consumed the majority of the seed from each selection. No trends between each of the three selections were shown towards one type or another. As shown in Figure 1, this group of naive geese consumed the majority of the seed from each feeding station.

Run 2 - Five Days later;

[0075] The trial in Run 1 above was then repeated each day over a period of five days. It was observed that between the beginning of the trials to the end of the five days in Run 2, the geese in the trials 'learned' from their feeding experiences and consumed all of the 'nil' endophyte seed and approximately 10% of the seed from the Endophyte A and B feeding stations. The total percentage of seed consumed from each feeding station is shown in the graph in Figure 2. This showed that a PDF effect had occurred and that the endophyte and grass combination exhibited a repellent effect.

Run 3 - Learned Geese, Three Months Later;

[0076] The inventors then conducted a further trial subsequent to Runs 1 and 2. The learned

geese from the Runs 1 and 2, who had been exposed to endophyte and had displayed PDF behaviour were removed from the trial and fed on a high maintenance nil endophyte diet of timothy forage grass and crushed barley for three months.

[0077] Following this three month exposure to the nil endophyte grass, these learned geese were then exposed to feeding stations containing endophyte infected seed or nil-endophyte infected seed. At the same time, in a separate enclosure, a trial was conducted with naïve geese also being exposed to feeding stations containing endophyte infected seed or nil-endophyte infected seed.

[0078] The graph in Figure 3 shows the consumption rate of the learned geese from this run. As shown, the percentage of nil-endophyte infected seed consumed was approximately 95% whereas the percentage of endophyte infected seed consumed was only approximately 5%. Again, this Run showed that the grass and endophyte combination exhibited a repellent effect and that the geese retained the PDF learned behaviour over the 3 month time period. In practice, this illustrates that birds are not only repelled in the short term by the endophyte and grass combination but that the endophyte and grass combination also provide a longer term repellence solution.

EXAMPLE 2 - HERBAGE, CHOICE TRIAL

[0079] In this trial, the methodology used was of similar manner to that of Example 1. The main difference in this trial was that the geese were fed on herbage rather than seeds.

[0080] During this trial, 20 naive geese that had no exposure to any endophytes were confined to an area of an approximate size of 15m². Within this area, perennial ryegrass herbage known as 'Grasslands Nui' (Nui) was grown and random sections (of an approximate size of 0.5m²) of the grass were either infected with AR1 endophyte or not infected at all (nil-endophyte). Each of these plots was randomly distributed within the larger area and cut to ground level.

[0081] The geese were exposed to a number of areas of the same approximate size as detailed above and were only moved when the grass levels were significantly reduced or fouled. After 10 days of geese exposure to the grass areas, the amount of herbage consumed was assessed. As shown in the graph of Figure 4, the geese consumed approximately 40.0g of non-infected 'nil' endophyte grass herbage and approximately 10.0g of the endophyte infected grass. This shows that the geese had developed a grazing preference within this time period and developed PDF behaviour, even when exposed to a different portion of the plant. This also illustrates that the repellent effect results from alkaloid toxins throughout the herbage and not just from the seed.

EXAMPLE 3 - HERBAGE, NO CHOICE TRIAL

[0082] A similar trial as described in Example 2 was conducted to assess the amount of herbage a gaggle of geese would consume when the geese were put onto a no-choice diet.

[0083] In this trial, two groups of four naive geese were confined to cages of 1m by 3.5m. The geese were then put onto a diet of nil endophyte Nui ryegrass for a period of eight days. Following this, the geese were then moved to a section of Nui ryegrass that had been infected with AR1 endophyte, also for an eight day period. Post grazing samples of the amount of herbage consumed were taken and assessed.

[0084] As shown in Figure 5 the average amount of dry weight consumed per day of plant material containing nil endophyte was approximately 20g. This was in comparison to the 5g of plant material consumed that was infected with endophyte. Also shown in Figure 6, during the eight-day consumption period on the endophyte infected plant material, the geese experienced a weight loss of approximately 2.2g, while only approximately 1g of weight was lost during the consumption period on nil-endophyte infected grass.

EXAMPLE 4

[0085] A further trial was then completed to assess geese feeding habits using seed collected from four different grass cultivars infected with endophyte to determine which grass cultivar(s) the geese preferred and hence which cultivars had the highest repellent effect.

[0086] The four different perennial ryegrass cultivars and hybrids (now referred herein as 'ryegrass') (and control) that were tested include the following:

1. (i) Grass seed with no endophyte infection (control);
2. (ii) Nui cultivar infected with AR1 endophyte;
3. (iii) Kingston cultivar infected with wild type endophyte;
4. (iv) 'Grasslands Greenstone' (Greenstone) cultivar infected with Endosafe®; and,
5. (v) Aries HD cultivar infected with wild type endophyte.

[0087] The trials were conducted in a similar manner as previously outlined above in Example 1. Here the feeding preferences of 20 naive geese of seed from 5 different grass cultivars and endophyte infection were assessed. The amount of seed consumed by the geese was assessed over a time period of three hours after distributing the seed. This trial ran for three days, with the seed being distributed once a day.

[0088] The results of this trial are shown in Figure 7. As shown, during the first feeding session, a high percentage (from 60% to 100%) of the seed from all five different feeding stations was consumed. During the second and third feeding sessions, there was a sharp decrease in the percentage of seed consumed from three out of the five combinations offered to the geese. For

example, the percentage of Kingston cultivar seed consumed dropped from a 60% consumption rate in the first feed to an approximately 5% consumption in the second and third feeding sessions. Greenstone cultivar consumption dropped from 80% consumption in the first feeding session to 10% consumption in the second and third feeding sessions. Airies cultivar went from a 100% consumption rate to an approximately 2% consumption rate in the second and third feeding sessions.

2. Alkaloid Concentration Analysis

[0089] From the above results the inventors concluded that the degree of repellence was based on the level of one or more alkaloids in the combination causing the PDF response behaviour.

EXAMPLE 5

[0090] To verify the above findings, the inventors conducted a further trial to assess the alkaloid concentrations of each grass cultivar to assess if there was a common alkaloid between the grass cultivars.

[0091] Figure 8 shows the content of three alkaloids, ergovaline, lolitrem and peramine from the five different grass cultivars tested in Example 3. From these results, the combinations that exhibited the strongest PDF responses had high levels of the alkaloid ergovaline. As shown, of the cultivars that displayed the PDF responses, Airies had a 10 ppm concentration of ergovaline while Greenstone and Kingstone had a 20 ppm concentration. In comparison, the combinations that did not display the PDF response effect, specifically the 'nil' infected seed and Nui, had no ergovaline present.

[0092] There was no clear pattern from the results observed for other alkaloids. For example, Kingstone and Airies both had small concentration levels of lolitrem, while Greenstone had no detectable levels of lolitrem. Levels of peramine alkaloid were observed in all grass cultivars, including the 'nil' infected seed and the Nui cultivar, hence it is concluded that peramine or lolitrem was not a cause of the exhibited PDF response.

EXAMPLE 6

[0093] Given the above results, further trials were completed to select preferred candidates of endophyte and grass cultivar that maximised ergovaline levels. Selections were also made based on alkaloid levels being exhibited throughout the herbage of the plant, rather than just those levels concentrated within a particular area. A further selection criteria was to screen for levels of peramine and loline alkaloids which are known to confer resistance to biotic and abiotic stresses.

Alkaloid concentration - tall fescue***Ergovaline concentration***

[0094] Firstly, the inventors assessed the alkaloid, ergovaline, in different tall fescue grass cultivars infected with the same New Zealand wild endophyte strain. The inventors assessed the concentration of ergovaline throughout the total herbage of the plant, as well as throughout the leaf and pseudostem. The results of this assessment are shown in Figure 9. As shown in Figure 9, ergovaline expression in New Zealand wild fescue was the highest of the grass cultivars tested.

Leaf Colonisation

[0095] The inventors then assessed the percentage of leaf colonisation of three New Zealand wild tall fescue endophyte strains in four tall fescue cultivars. The results of this assessment are shown in Figure 10. Here, the grass cultivar Resolute showed a high percentage of leaf colonisation overall between all three endophyte strains and all the grass cultivars assessed.

Alkaloid concentration - Lolium species***Peramine Alkaloid***

[0096] The alkaloid, peramine has been shown to have an increased effect in pest resistance, such as insects. Therefore, this is also a further preferred characteristic of a grass and endophyte combination.

[0097] To assess the peramine expression within particular ryegrass specific endophyte strains in one ryegrass cultivar, using High Performance Liquid Chromatography (HPLC), the inventors assessed the peramine expression in both the leaf and pseudostem from six different endophyte strains (AR41, AR5, AR8, AR4 and AR9001 in combination with the same ryegrass cultivar.

[0098] As shown in Figure 11, the majority of the endophytes displayed a peramine concentration level of approximately 20 ppm in both the leaf and the pseudostem. In particular, the combination with the Waiau endophyte displayed the highest peramine concentration of 35 ppm and 40 ppm in the leaf and pseudostem respectively.

Ergovaline Alkaloid

[0099] As previously discussed, trial results illustrated that ergovaline is an important alkaloid in producing the PDF response displayed by the geese (Example 5). Therefore, it is preferable to have a high concentration of this alkaloid within the herbage of grass and endophyte combination. The inventors assessed the same endophyte strains as above (AR41, AR5, AR8, AR4 and AR9001 for the level of ergovaline concentration, in the same ryegrass cultivar.

[0100] Figure 12 shows the ergovaline concentrations expressed within these ryegrass and endophyte combinations. Here, endophytes AR4, AR5, and AR8 displayed the highest concentration of ergovaline expression. The expression was greatest throughout the pseudostem; however, ergovaline expression was also displayed throughout the leaf.

[0101] The tests in this Example showed that endophytes AR5 and AR8 have high concentration levels of both the alkaloids ergovaline and peramine. Therefore, combinations with these endophytes can have high bird repellent properties as well as good resistance to biotic (insect) stresses. Endophyte AR4 can further provide a bird repelling cultivar, but as it expresses lower levels of peramine, is better suited to environments with fewer insect stresses.

Alkaloid Concentration - Lolium and Festuca species

[0102] The inventors also assessed the degree of alkaloid expression (peramine, loline and ergovaline) throughout the leaf and pseudostem from four different tall fescue endophytes (AR601, AR602, AR603 and unknown endophyte) infected in the same tall fescue cultivar and from one endophyte strain, AR94 infected in a ryegrass cultivar.

[0103] As shown in Figure 13, peramine expression in the unknown Nui endophyte and tall fescue cultivars was high (>10 ppm concentration) throughout the pseudostem of the cultivars.

[0104] The tall fescue and AR602 combination also produced high peramine expression, as did the ryegrass and AR94 combination.

[0105] Figure 14 shows the expression of loline alkaloid of the various endophyte and tall fescue/ryegrass cultivar combinations. As illustrated, tall fescue and AR601 combination showed a high expression of loline concentration. Other tall fescue and endophyte combinations also displayed moderate levels of loline expression. By comparison, ryegrass and AR94 combination did not express any detectable loline alkaloids.

[0106] Figure 15 shows the concentration of ergovaline expression of grass and endophyte combinations as discussed above. As shown, the tall fescue and AR601 combination exhibited a high level of ergovaline concentration throughout the plant. There were comparatively high levels of ergovaline throughout the leaf and pseudostem. In comparison, the ryegrass and AR94 combination exhibited high levels of ergovaline expression in the pseudostem but with

only a small amount within the leaf.

EXAMPLE 7

[0107] Further trials are conducted to select preferred candidates of endophyte and grass cultivar that maximise ergovaline levels. Selections are also made based on alkaloid levels being exhibited throughout the herbage of the plant, rather than just those levels concentrated within a particular area. A further selection criteria is to screen for levels of peramine and loline alkaloids which are known to confer resistance to biotic and abiotic stresses.

Ergovaline concentration

[0108] Firstly, levels of the alkaloid ergovaline are assessed in different tall fescue grass cultivars infected with the same New Zealand wild endophyte strain. The concentration of ergovaline is also assessed throughout the total herbage of the plant, as well as throughout the leaf and pseudostem.

Leaf Colonisation

[0109] The percentage of leaf colonisation of three New Zealand wild tall fescue endophyte strains in four tall fescue cultivars is subsequently evaluated.

Alkaloid concentration - Ryegrass

Peramine Alkaloid

[0110] The alkaloid, peramine has been shown to have an increased effect in pest resistance, such as insects. Therefore, this is also a further preferred characteristic of a grass and endophyte combination.

[0111] To assess the peramine expression within particular ryegrass specific endophyte strains in one perennial ryegrass cultivar, the level of peramine expression in both the leaf and pseudostem from different endophyte strains (AR4, AR5 and AR94), in combination with the same ryegrass cultivar, is evaluated. Expression levels of at least 15 ppm in the leaf and, or alternatively, in the pseudostem is detected.

Ergovaline Alkaloid

[0112] As previously discussed, trial results illustrated that ergovaline is an important alkaloid in producing the PDF response displayed by the geese (Example 5). Therefore, it is preferable to have a high concentration of this alkaloid within the herbage of grass and endophyte combination.

[0113] The same endophyte strains (AR4, AR5 and AR94) in combination with the ryegrass cultivar as described above are evaluated for the level of ergovaline concentration. Levels of ergovaline of at least 5 ppm, 10 ppm or 15 ppm within the endophyte and ryegrass combination are detected. The expression is greatest throughout the pseudostem; however, ergovaline expression is also displayed throughout the leaf.

[0114] Combinations with these endophytes can have high bird repellent properties as well as good resistance to biotic (insect) stresses.

Alkaloid Concentration - Perennial Ryegrass and Tall Fescue

[0115] The degree of alkaloid expression (peramine, lolitrem and ergovaline) throughout the leaf and pseudostem of different ryegrass endophytes (AR94, AR4, AR5, AR8) infected in the same ryegrass cultivar is assessed.

[0116] Peramine expression of at least 15 ppm concentration is detected throughout the pseudostem of the cultivars.

[0117] The expression of lolitrem alkaloid of the various endophyte and ryegrass cultivar combinations is evaluated. High expression of lolitrem concentration within the range of 1-50ppm is detected in the combinations.

[0118] The concentration of ergovaline expression of grass and endophyte combinations as discussed above is evaluated. High levels of ergovaline concentration of at least 5 ppm, 10 ppm, or 15 ppm are detected throughout the plant in the various ryegrass and endophyte combinations. Comparatively high levels of ergovaline are found throughout the leaves and pseudostems of the combinations

[0119] The combinations displaying the highest levels of ergovaline, peramine and/or loline levels demonstrate efficacy in repelling avian species using protocols described in Examples 1-4.

[0120] It is found that by selecting an endophyte that produces peramine and loline alkaloids in addition to levels of ergovaline effective for repelling avian species, the grass not only deters avian species but also produces a high tolerance to biotic and abiotic stresses. This finding is also the case with ryegrass cultivars. Biotic stresses include resistance to attack from insects and pests. Specific examples include resistance to Grass Grub, (*Costelytra zealandica*)

Argentine Stem Weevil (*Listronotus bonariensis*) and sucking insects such as, for example *Rhopalosiphum padi*. Abiotic stresses include resistance to drought or dry periods as well as resistance to high or low temperature climates. High temperatures include temperatures from 27°C and above. For example, temperatures may include 27°C, 30°C, 32 °C, 35°C or 40°C. While low temperatures include any temperatures below, and including, 7°C, 5°C, 3 °C or 0°C for example.

EXAMPLE 8

Tall fescue turf cultivar "Jackal" inoculated with AR601 and AR604

[0121] Following the above trials, the inventors assessed the further concentration of alkaloids in a tall fescue turf cultivar "Jackal" inoculated with AR601 & AR604 endophytes. The alkaloid concentrations over a time period of approximately 1 year were assessed using the High Performance Liquid Chromatography (HPLC), methods described above.

[0122] Figures 16 and 17 show the results of the concentration analysis of the alkaloids lolines and ergovaline respectively from Jackal inoculated with either AR601 or AR604. Alkaloid measurements were taken from both the leaf and stem of the plants. As shown in the comparison, the stems of the plants had a higher alkaloid concentration. Additional seasonal variations can also be shown in the Figures.

Table 1: Shows the average alkaloid concentration in pseudostems of Jackal tall fescue infected with AR601 or AR604.

	Ergovaline (ppm)	Lolines (ppm)
AR601	8.3	1816
AR604	7.7	1778

3. Target Avian Species: Gulls (*Larus dominicus*)

[0123] The inventors conducted a further trial to determine the effect of introducing the repellent into an animal feed (pig feed) and determines the repellence effect on gulls (omnivorous birds).

EXAMPLE 9

Treatments

[0124] To assess the PDF effect in gulls from endophytes the following treatments were assessed: (i) Commercially available pig pellets made from crushed grain or (ii) commercially available pig pellets externally hand-coated with ground powder produced from a ground endophytic seed.

[0125] To produce the ground endophytic seed for coating the pellets, 40 kg of endophyte-containing seed was commercially ground, with the temperature never exceeding 30 deg Celsius. The pellets were then manufactured by treating 20 kg lots of standard pig pellets with a 2kg coating of ground seed powder in a concrete mixer, wherein the pellets were sprayed with warm water to produce a sticky surface that was then dusted with powder and allowed to air dry. Most of the pellets being coated ranged in size from 25 to 30 mm long by 15 mm diameter. Broken pellets smaller than 5mm by 5 mm were also treated in the mix. A trial to confirm that the birds would not reject pellets containing the endophyte powder was conducted eight weeks prior to the trial and no rejection was noted.

Run 1

Observation of gulls behaviour prior to testing

[0126] Observation of the birds flocking to the pig farm at feeding times showed two distinct groups of birds that, when finished with opportunistic feeding for the day, loitered at specific locations post-feeding. These post-feeding locations were on bare, raised stony paddocks that offer security and good vision.

Pre-trial manipulating of the gulls feeding habits

[0127] Each day for three weeks, the gulls were fed 50 kg of untreated pig pellets per day in the mornings at one of these loitering areas. The gulls appeared to consume all that was offered to them. The area was fenced off from the pigs.

[0128] During this pre-trial behaviour manipulation, the birds learned that there was regular feeding at this area and consumed the pig feed on most fine days through the three week pre-feeding period. On inspection in the mornings the birds were not in these rest areas but actively feeding from troughs around the whole farm by following the feed carts for opportunistic feeding. Once these troughs were emptied by the communal feeding, the gulls spent the rest of the day in these rest areas and were observed there until disturbed in the evenings. Gulls were observed to swallow the pellets whole.

Trial

[0129] At the end of the three week pre-feeding time period, the trial testing above treatments commenced. Each morning, pellets treated as described were placed into separate troughs that were located in this fenced off area. The trough and pellets were weighed before feeding in order to measure the amount of pellets consumed the following morning.

[0130] For 15 days, the troughs were measured for daily consumption. During this period, it was noticed that there was some rejection of the smaller treated pellets. Visual observations of the treated pellets consumed showed a marked preference by the gulls in consuming whole large pellets rather than the broken smaller pieces.

Results

[0131] On day 1, the birds consumed less endophyte treated pellets. This demonstrates that the birds were able to detect differences by indicators such as, smell, taste or visual differences between the troughs containing untreated and treated pellets. However, there was no total rejection from the onset.

[0132] As shown in Figure 18, on days 1 to 3, up to 70% of the endophyte-treated pellets offered were consumed. For the following trial days 4 to 15, of the total amount of pellets consumed, only an average of 30% of the treated pellets per day were consumed. This developed aversion to the endophyte treated pellets shows a learned response or Post Digestion Feed Back (PDF).

[0133] Although there was a significant difference between the consumption of treated pellets compared with the untreated control at the onset, this difference rapidly grew larger up to day 15 when the trial was stopped.

[0134] Even though total rejection of the control pellets was not achieved, over time, the divergence of the slopes in Figure 19 between the treatments is positive. The results with the gulls show promising results for the development of a pellet that can deliver selected endophyte compounds and induce PDF in birds, so that they will subsequently associate normal untreated pig pellets as a source of their malaise. This learned behaviour is dependent on treated pellets looking, smelling and tasting exactly the same as the untreated pellets. The pellets manufactured in this experiment were visually different, and this can contribute to the birds' ability to differentiate between the treated and control pellets, as illustrated at the onset of the trial.

4. Target Avian Species: Finches (*Carduelis choris*)

[0135] The inventors in this trial assessed the PDF response of Finches and whether learned

behaviour can be shown with a choice between nil-endophyte seed and endophyte containing seed.

EXAMPLE 10***Capturing and conditioning the birds***

[0136] A number of green finches that were attracted to a granary outside Christchurch were humanely trapped and contained in a facility built for this purpose. The birds were conditioned for six weeks to their new environment and fed bird seed and water on demand.

[0137] In the seventh week, nil endophyte ryegrass seed was introduced into the bird seed mix at increasing percentages to condition the birds to this food source. This was conducted for a further six week period, and consumption of the nil endophyte ryegrass was recorded to ascertain how much of this grass seed the birds normally eat.

[0138] After the above time period, the birds were fed only water for a period of four hours. Four randomly chosen birds were then captured and separated into individual enclosures. Each bird was then fed on endophyte containing ryegrass seed and later on nil endophyte ryegrass seed.

Results

[0139] The results of the birds' eating behaviour of the nil endophyte seed and endophyte seed is shown in Figures 20 and 21. These Figures show a marked decrease in consumption of feed that contained endophyte (10 to 20% of the normal feed consumed for endophyte seed versus 60 to 65% of the normal feed consumed for nil endophyte seed. The results show a genuine repellency by the finches towards endophyte containing seed. Therefore, these results also illustrate a Post Digestive Feed back response to the endophyte containing seed.

5. Christchurch International Airport Trial:

Target Avian Species: Wild birds, various species

[0140] During the autumn of 2007, a trial assessing PDF response on wild birds present at Christchurch International Airport (CIAL) was conducted.

EXAMPLE 11

Methodology

[0141] A number of plots, as outlined in Table 2 below, containing either existing vegetation (control) or planted "Jackal" Tall Fescue (*Festuca arundinacea*) was set up. Each plot was the square size of 50m by 50m.

TABLE2: PLOT OUTLINE

CONTROL EXISTING SPECIES REP 1	CONTROL EXISTING SPECIES REP 1	CONTROL EXISTING SPECIES REP 2
TALL FESCUE ENDOPHYTE REP 1	TALL FESCUE ENDOPHYTE REP 2	CONTROL EXISTING SPECIES REP 2

[0142] The existing vegetation on the control plots consists of plant species, including Hair grass (*Vulpia* sp) and browntop (*Agrostis tenuis*), with small amounts of Yorkshire fog (*Holcus lanatus*), Bromus species, Cocksfoot (*Dactylis glomerata*), Barley grass (*Hordeum* sp), Couch (*Agropyron* sp) and many broadleaf weeds including Wire weed (*Polygonum* sp), Storksbill (*Erodium* sp), and. It was assumed that this vegetation did not contain endophytes.

[0143] In comparison, the Tall fescue contained either AR601 or AR604 endophyte strains from the *Neotyphodium coenophialum* species.

[0144] During the trial, each plot was monitored daily, at the morning and evening for two minutes. During this time the bird species and numbers was assessed and recorded.

Results

[0145] The total bird numbers present in the endophyte-enhanced plots with the control plots per month for the trial are shown in Figure 22. As shown, there was a significant reduction in the number of birds visiting the endophyte plots in comparison to the control plots over the six month period.

[0146] The statistical analysis of variance of this trial (shown in Tables 3 to 5 below) showed that even though the data are very variable, the endophyte effects on bird numbers were significant.

TABLE 3: Shows the logarithms of numbers of Birds - 6 month period

	SS	df	MS	F	P-value		
Endophyte effect	222.2033	1	222.2033	7.714026	0.04995		
residual	115.2204	4	28.8051				

	SS	df	MS	F	P-value	
Total	337.4237	5				
Bird numbers were converted to logarithms to smooth variation before statistical analysis						

TABLE 4: Shows the means for Endophyte

	Endophyte	601/604	Control (existing grasses)			
	mean	11	30.83			
	s.e.	5.999	4.242			

Two bird counts were made each day. Counts were added for each month.

TABLE 5: Shows the mean accumulated totals per plot for Month

Date	Oct	Nov	Dec	Jan	Feb	Mar	
	49.83	16.83	40.17	8	8.17	22.33	
Standard error 8.48							

6. Rangiora Sewage Treatment Centre:

Target Avian Species: Wild birds, various species

[0147] Following the alkaloid assessment of plants as described under Example 8 above, a field trial to assess the PDF from a variety of bird species was conducted.

EXAMPLE 12

Method

[0148] From the Inoculated Jackal tall fescue plants that were chemo-typed for their alkaloid profile, a number of plants with AR601 and AR604, covering a range of ergovaline (EV) and loline profiles representing three groups, moderate, high and very high EV's. Table 6 below summarises the grass species and ergovaline levels assessed in this trial.

Table 6: Selected range of the average ergovaline alkaloid treatments:

Grass species	average level of ergovaline (ppm)
Tall Fescue	15.1
Tall Fescue	8.7
Tall Fescue	2.8
Tall Fescue-nil Endo	0

[0149] The plants were each divided into four ramets of the same approximate size to get four replicates. All plants were grown over the winter period and rimmed to the same relative size prior to transplanting the plants to the test area - a field at a sewage treatment plant in Rangiora, New Zealand, where wild herbivorous birds are known to frequent. The plants were placed in groups and identified by location using a grid mapping system.

[0150] Treatments (nil endophyte and endophyte-infected with 3 levels of ergovaline) were placed in random blocks in the grazing area. Each treatment plant was placed so that the birds had the ability to select in a choice situation.

Assessment

[0151] After a grazing period of approximately 5 days depending on bird numbers in the area, each plant was inspected for grazing and assessed on a scale of 0 (no grazing) to 5 (severely grazed). Following scoring, plants were protected from further bird predation to allow sufficient re-growth. The plants were then trimmed off to the same level with a motor mower, watered and fertilised as necessary then fenced to allow for growth recovery. This recovery period was approximately one week depending on the seasonal rainfall.

[0152] The area was then re-opened to birds and all plants were inspected every second day and scored when evidence of flock grazing had occurred.

[0153] The dominant bird species in October were Canada geese. Later paradise shelducks (in fact another goose species) dominated. Initial assessment cycles were frequent when the local geese, about 20 in number, found the plots. Larger numbers of paradise shelduck (600 to 1500 in the 10 ha pond site) were present from December and due to moulting birds remaining in the area.

[0154] For statistical analysis of grazing scores, a mean over all scoring occasions was calculated for each consolidated block of plants of the same group (based on ergovaline measurements).

Results

Feeding scores

[0155] Grazing preference of the birds over the eight observation periods between October 2007 and April 2008.

[0156] At the first sampling period with low bird numbers, most of the endophyte grasses had some repellent properties (See Figure 23). At the second and third scoring there were no significant differences between any of the treatments. On the next two scorings between the end of October and early December there was a significant result between endophyte plants and nil endophyte (Figure 24).

[0157] For the remaining scorings from mid December to early April all plants were under stress from very large bird numbers. As the birds have to learn that the endophyte grasses can make them ill over time it is our opinion that there were insufficient endophyte plants (or too many geese) in this trial to teach these 600 + birds to avoid grazing.

[0158] Analysis of all scorings for October 2007 to April 2008 looking at the grazing pressure over time has shown that the fescue plants infected with AR601 & AR604 endophyte strains had significant repellence, despite the large bird numbers (Table 7 and 8 and Figure 24).

Table 7: Analysis of variance results of all assessments. Accumulated feeding scores for tall fescue plots

Source	d.f.	s.s.	m.s.	v.r.	F pr.
endo	2	2.023	1.011	14.28	<.001
rep	3	2.434	0.811	11.45	<.001
Residual	17	1.204	0.071		
Total	22	5.661	0.257		

Table 8: Mean accumulated feeding scores for endophyte treatments for tall fescue plots.

Endophyte strain	Accumulated feeding score
AR601	1.40
AR604	1.18
nil	2.01
LSD 5%*	0.34
*mean for all comparisons	

[0159] An additional analysis of the endophyte-infected plots alone showed that the feeding scores on AR604 plots were significantly lower (i.e. less grazed) ($P < 0.001$ probability level) than those on AR601 (1.1 v 1.5 respectively).

Survival

[0160] At the end of the trial, the number of plants that had survived in this environment under both bird grazing pressure and compacted low fertility stony soils were assessed. The results are shown in Table 9 below.

Table 9: Percentage surviving plants in April 2008.

Cultivar-endophyte	surviving (%)
Fescue-601	53
Fescue-604	52
Fescue-nil	14
LSD 5%	31

[0161] The data shows that endophyte infection has enhanced the survival of tall fescue. Drought, severe grazing and perhaps pest pressure have contributed to mortality, but endophyte has influenced the overall outcome.

Minimum effective ergovaline levels:

[0162] In analysing the levels of ergovaline necessary to obtain some significant degree of repellence in plants infected with endophyte, there is no difference between levels of material that had previously tested with ergovaline in the range levels between 2.8 and 15 ppm. As shown in Table 10, there is no evidence at this stage that high levels have further reduced grazing scores compared to moderate levels of ergovaline.

Table 10: Mean ergovaline levels determined pre-trial (March 2007) and grazing scores for plants.

Description	Endophyte	Mean ergovaline level (ppm)	Mean grazing score over the season
Jackal tall fescue	AR601	2.8 †	1.5 *
Jackal tall fescue	AR604	2.8	1.2 *
Jackal tall fescue	AR601	8.7	1.5 *
Jackal tall fescue	AR604	8.7	1.6
Jackal tall fescue	AR601	15.1	1.3 *
Jackal tall fescue	AR604	15.1	1.1 *

Description	Endophyte	Mean ergovaline level (ppm)	Mean grazing score over the season
Jackal tall fescue	NIL	0	2.0
LSD 5% = . 0.4 * = significantly different from nil endophyte control @ 5% † ppm = parts per million (mg/kg)			

Conclusion

[0163] The trial showed that endophyte infection of tall fescue and ryegrass effectively reduces feeding by herbivorous birds and increases plant survival and has shown that increasing the ergovaline concentrations above those in the lower range of the distribution does not affect the result.

7. Endophyte Genotype Analysis

[0164] This example provides microsatellite data to enable a skilled person to identify and confirm whether they are in possession of an endophyte strain or a variation thereof.

EXAMPLE 13

[0165] This example enables a person skilled in the art to distinguish endophytes AR601, AR602, AR603, AR604, AR4, AR5, AR8 and AR94 and variations thereof, as described above from other endophytes. In Table 11 below, the endophyte strains and allele sizes at various Simple Sequence Repeats (SSR) are listed, along with the methodology used to arrive at the SSR sizes.

TABLE 11: Summary of Microsatellite data for distinguishing the endophytes referred to in the present application.

SSR	Allele ¹	Allele size (bp) ²	AR5	AR8	AR94	AR60 1	AR60 2	AR60 3	AR60 4
B10	b10_01	140				140			
	b10_06	162					162	162	162
	b10_07	165							165
	b10_09	171				171	171	171	171
	b10_11	176	176	176	176				
	b10_14	185				185	185	185	

SSR	Allele ¹	Allele size (bp) ²	AR5	AR8	AR94	AR60 1	AR60 2	AR60 3	AR60 4
B11	b11_16	165				165	165	165	165
	b11_18	177							
	b11_20	184			184				
	b11_22	192		192		192	192	192	192
	b11_28	237	237						
ans014	ans014_02	310				310	310	310	310
	ans014_05	314	314	314	314				
	ans014_07	316				316	316	316	316
ans016	ans016_04	290				290	290	290	290
	ans016_05	293	293	293	293	293	293	293	293
	ans016_09	305				305	305	305	305
ans017	ans017_12	300				300			
	ans017_13	305		305	305				
	ans017_18	315							315
	ans017_19	318					318	318	
	ans017_28	362	362						
	ans017_31	388							
ans019	ans019_01	193				193	193	193	193
	ans019_03	200	200	200	200				
ans025	ans025_02	286				286	286	286	286
	ans025_07	305		305	305				
	ans025_11	311	311						
ans030	ans030_06	325	325	325	325				
	ans030_08	331				331	331	331	331
ans031	ans031_10	317	317	no data		317	317	317	317
	ans031_17	341		no data	341				

SSR	Allele ¹	Allele size (bp) ²	AR5	AR8	AR94	AR60 1	AR60 2	AR60 3	AR60 4
	ans031_19	353		no data		353	353	353	353
ans036	ans036_04	262				262	262	262	262
	ans036_05	265	265	265	265	265	265	265	265
ans042	ans042_01	123				123	123	123	123
	ans042_03	130				130	130	130	130
	ans042_06	140		140	140				
	ans042_09	150							
	ans042_14	184	184						
¹ Allele name is a temporary designation only, based on alleles identified in strain surveys to date ² Allele size bin +/- 0.5 bp									

Methodology used to locate Simple Sequence Repeats above in Table 1

DNA isolation

[0166] Genomic DNA was isolated from segments of fresh pseudostem material from endophyte-infected grass using the FastDNA® kit (Q-BIOgene, Irvine, California, USA) in conjunction with a FastPrep Cell Disruptor (Q-BIOgene). Genomic DNA is diluted 10x in sterile MilliQ water prior to preparation of polymerase chain reactions (PCR).

Simple sequence repeat (SSR) marker analysis

[0167] The endophyte strains were characterized using primer pairs for SSR markers loci B10, B11, ans014, ans016, ans017, ans019, ans025, ans030, ans031, ans036 and ans042. Primer sequences for these loci are given in Table 12 below. Strain characterizations are conducted using primers synthesized commercially by Integrated DNA Technologies (Coralville, Iowa, USA).

TABLE 12: Primer sequences for simple sequence repeat (SSR) loci used to characterize endophyte strains *in planta*.

SSR locus	Forward primer sequence (5' - 3')	Reverse primer sequence (5' - 3')
ans014	CAATCATTCTGAACCAAACCA	CCTTATACGCAGGACCAAAGAC
ans016	CACAAAGACAAACGCCAAAAG	GCAAAGCTCACAGACAAAGGTC
ans017	GCCATCCGATAACCTGTCTCTA	TCGTTTCGGTTCGATTAAGAGT
ans019	TACCTCTGCACGGTGTATTCC	TGCATAACACTCACCTTATAGTCG
ans025	CAGCCCTATTTTATGTTTGAAGG	GCTCCTGCTTTTATTCCTGCTA
ans030	GCATAGGAAGTGCTGTTAATTTGA	AGAGTAGAACCTGCTTGCGTTA
ans031	ACTCGGAGACA TGAAAACCA TC	GACGTGCTCTGTGATGTTGAAT
ans036	ATTTGCAGCAGAGATGATGTGT	CCTGCACCGGACTGTTAGTAAT
ans042	GATGACTACCCGAGTGAGAACC	AACCCAACAACGTCTTTTCATT
B10	CGCTCAGGGCTACATACACCATGG	CTCATCGAGTAACGCAGGCGACG
B11	CATGGATGGACAAGAGATTGCACG	TTCAGTGCTACAATTCTGTGGAGC

PCR amplifications using SSR markers B10 and B11

[0168] For both markers B10 and B11 the 5' end of the forward primer is labeled with 6-carboxyfluorescein (6-FAM™) for detection purposes. PCR amplifications were conducted in a 10 µL reaction volume containing 2 µL of diluted genomic DNA, 1.5mM magnesium chloride, 1x PCR buffer (Invitrogen, Carlsbad, California, USA), 0.1 mM of each deoxynucleotide triphosphate (dATP, dGTP, dTTP and dCTP), 0.2 µM each of the forward and reverse primers, and 1 U of Platinum Taq DNA polymerase (Invitrogen). PCR was carried out with the following profile: (1) 94°C for 4:00 minutes, (2) 30 cycles of: 94°C for 30 seconds, 60°C for 30 seconds and 72°C for 30 seconds, (3) 72°C for 7 minutes with an iCycler thermocycler (BioRad, Hercules, California, USA).

PCR amplifications using SSR markers with the 'ans' prefix

[0169] PCR amplifications use a three-primer system for fluorophore addition (Schuelke 2000). To facilitate this, forward primers are synthesized with a 21 nucleotide M13 tail sequence at the 5'-terminus (sequence 5'-TGTAACGACGGCCAGT-3'). The fluorophore used in the PCR reactions is 6-carboxyfluorescein (6-FAM™), but alternative dyes may be used. Reverse primers are synthesized with the sequence 5'-GTTTCTT-3' at the 5'-terminus end, to promote non-templated adenylation at the 3'-terminus end of PCR product (Brownstein et al. 1996). The result of adding the M13 tail sequence and 'pig tail' sequence to the primers is that the expected and observed lengths of PCR products using the 'ans' SSR primer pairs are increased by 25 basepairs (bp). PCR amplifications are conducted in a 10 µL reaction volume containing 2 µL of diluted genomic DNA, 2.5 mM magnesium chloride, 1x PCR buffer, 0.05 mM of each

deoxynucleotide triphosphate (dATP, dGTP, dTTP and dCTP), 0.0375 µM forward primer, 0.15 µM reverse primer, 0.15 µM of fluorescent-labelled M13 primer and 0.5 U of Platinum Taq DNA polymerase. PCR is carried out with the following profile: (1) 94°C for 4:00 minutes, (2) 30 cycles of: 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 30 seconds, (3) 8 cycles of: 94°C for 30 seconds, 53°C for 30 seconds and 72°C for 30 seconds, (4) 72°C for 30 minutes (Schuelke 2000) with an iCycler thermocycler.

Genotypic analysis of SSR markers

[0170] Genotypic analysis was conducted by capillary electrophoresis on an ABI 3100 Genetic Analyser (Applied Biosystems, Foster City, California, USA) used POP-7™ polymer (Applied Biosystems) and GeneScan™ 500 LIZ™ (Applied Biosystems) as an internal size standard. A run module based on the default Genescan22_POP4DefaultModule is used, with the following changes: run current 200 µA, pre-run voltage 0 kV, pre-run time 1 second, injection voltage 3 kV, injection time 10 seconds, number of steps 1 nk, voltage step interval 1 second, run time 560 seconds. Electropherograms are analysed and PCR fragment size determined using ABI Prism GeneScan (v 3.7, Applied Biosystems). The Genotype data was scored using ABI Prism Genotyper (v 3.7, Applied Biosystems).

CONCLUSIONS

[0171] It can therefore be concluded from the above results that:

- Selected endophyte infected grasses exhibit a repellent effect on birds.
- This repellent effect is a secondary repellent or PDF effect.
- Birds are repelled from seeds or herbage of the endophyte and grass cultivar combination.
- Ryegrass or tall fescue grasses show the repellent effect depending on the endophyte chosen.
- Ergovaline alkaloid is an important toxin that produces the avian repellent effect.
- Peramine and loline alkaloids are also useful to confer resistance to biotic and abiotic stresses on the grass.
- A combination of AR4, AR5, AR8 or AR94 endophyte with a ryegrass cultivar are good candidates for avian repellent applications, owing to their repellent levels of the alkaloid ergovaline and insect resistance conferred by peramine and/or lolitrem alkaloids.
- A tall fescue and AR601 or AR604 endophyte combination can be used as avian repellent and additionally provide good insect resistance.
- Other suitable candidates include, AR602 or AR603 with tall fescue as these combinations also display higher levels of ergovaline and peramine and/or loline alkaloids.

[0172] The present invention has been described by way of example only, and it is appreciated that modifications and additions can be made thereto without departing from the scope thereof as defined in the appended claims.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US6111170A **[0066]**
- US6072107A **[0066]**

Non-patent literature cited in the description

- **PENNELLROLSTON** Proceedings of the New Zealand Grassland Association, 2003, vol. 65, 229-238 **[0012]**

Patentkrav

1. Endophyt- og græskultivarkombination, der frastøder fuglearter fra kombinationen, kendetegnet ved, at endophytten
5 er valgt fra den gruppe, der består af: AR601, AR602, AR603, AR604, (opbevaret ved The National Measurement Institute (NMI) Australia, Accession Nos.V07/029058, V07/029059, V07/029060, and V07/029061) og kombinationer deraf.
- 10 2. Endophyt- og græskultivarkombination ifølge krav 1, kendetegnet ved, at græskultivaren er en Festuca-græskultivar eller græskultivar, der udviser Festuca-egenskaber, som frastøder fuglearter fra kombinationen, kendetegnet ved, at endophytten i kombinationen producerer et niveau på mindst 3
15 ppm ergoalinalkaloidforbindelse i en del af kultivargræsset og de frø, der produceres ud fra det.
3. Endophyt- og græskultivarkombination ifølge krav 1 eller krav 2, kendetegnet ved, at græskultivaren er en
20 strandsvingelkultivar (Festuca).
4. Endophyt- og græskultivarkombination ifølge et af kravene 1 til 3, kendetegnet ved, at endophytten i kombinationen producerer lolinalkaloider i mindst en del af kultivargræsset
25 og de frø, der produceres ud fra det.
5. Endophyt- og græskultivarkombination ifølge krav 4, kendetegnet ved, at endophytten i kombinationen producerer lolinalkaloider i et omfang fra 1 ppm til 9500 ppm.
30
6. Endophyt- og græskultivarkombination, der frastøder fuglearter fra kombinationen, kendetegnet ved, at endophytten er valgt fra den gruppe, der består af: AR4, AR5, AR8 and AR94 (registreret ved The National Measurement Institute (NMI),
35 Australia, Accession Nos. V07/029054, V07/029055, V07/029056, V07/029057) og kombinationer deraf.
7. Endophyt- og græskultivarkombination ifølge krav 6,

kendetegnet ved, at græskultivaren er en Lulium-græskultivar eller græskultivar, der udviser Lulium-egenskaber, kendetegnet ved, at endophytten i kombinationen producerer et niveau på mindst 3 ppm ergovalinalkaloidforbindelse i en del af kultivargræsset og de frø, der produceres ud fra det.

10

8. Endophyt- og græskultivarkombination ifølge krav 6 eller krav 7, kendetegnet ved, at græskultivaren er en flerårig rajgræskultivar (*Lolium perenne*).

9. Endophyt- og græskultivarkombination ifølge et af kravene 1 til 8, kendetegnet ved, at fuglearterne frastødes fra kultivargræsset og de frø, der produceres ud fra det.

15

10. Endophyt- og græskultivarkombination ifølge et af kravene 1 til 9, kendetegnet ved, at endophyt- og græskultivarkombinationen frastøder fuglearter via en sekundær afskrækkende virkning, der omfatter en post-digestiv feedback- (PDF) -mekanisme.

20

11. Endophyt- og græskultivarkombination ifølge et af kravene 1 til 10, kendetegnet ved, at endophytten i kombinationen producerer et niveau på mindst 3 ppm ergovalinalkaloidforbindelse i en del af kultivargræsset og de frø, der produceres ud fra det.

25

12. Endophyt- og græskultivarkombination ifølge krav 11 kendetegnet ved, at:

a) kombinationen omfatter en syntetisk græskultivar, eller

30

b) fuglearterne er fuglearter, som er valgt fra den gruppe, der består af gæs, måger, spurve, finker, viber og kombinationer deraf, eller

c) endophytten bibringer kombinationen resistens over for biotisk og abiotisk stress, hvor den biotiske stress omfatter resistens over for angreb fra insekter og skadedyr, og hvor den abiotiske stress omfatter resistens over for tørke eller tørre perioder såvel som resistens over for høje eller lave temperaturer, eller

35

d) endophytten i kombinationen producerer peraminalkaloidforbindelser i mindst en del af kultivargræsset og de frø, der produceres ud fra det, eller

5 e) endophytten i kombinationen producerer peramin i mindst en del af kultivargræsset og de frø, der produceres ud fra det, i et omfang fra 1 ppm til 100 ppm.

13. Fremgangsmåde til frastødning af fuglearter fra et jordareal ved plantning af en endophyt- og græskultivar ifølge
10 et hvilket som helst af kravene 1 til 12 på eller ved siden af det land, hvorfra fuglearterne skal frastødes.

14. Fremgangsmåde til screening af en endophyt- og græskultivarkombination ifølge et hvilket som helst af kravene
15 1 til 12 for kombinationens evne til at frastøde fuglearter ved hjælp af følgende trin:

a) dyrkning af kombinationen

b) måling af niveauet af et ergovalinalkaloid i den dyrkede kombination og

20 c) udvælgelse af kombinationen, hvis niveauet af ergovalin er mindst 3 ppm større.

15. Endophyt, som er valgt fra den gruppe, der består af: AR601, AR602, AR603, AR604, (opbevaret ved The National
25 Measurement Institute (NMI) Australia, Accession Nos.V07/029058, V07/029059, V07/029060, and V07/029061) og kombinationer deraf.

16. Endophyt, som er valgt fra den gruppe, der består af AR4,
30 AR5, AR8 and AR94 (opbevaret ved The National Measurement Institute (NMI), Australia, Accession Nos. V07/029054, V07/029055, V07/029056, V07/029057) og kombinationer deraf.

DRAWINGS

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16

LOLINES FROM JACKAL CIAL 2006/07




Figure 17




Figure 18




Figure 19





Figure 20



3489773
030207

Figure 21




Figure 22




Figure 23




Figure 24

