

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2017211120 C1**

(54) Title
Maytansinoid derivatives, conjugates thereof, and methods of use

(51) International Patent Classification(s)
C07D 498/18 (2006.01) **A61P 35/00** (2006.01)
A61K 31/535 (2006.01)

(21) Application No: **2017211120** (22) Date of Filing: **2017.01.24**

(87) WIPO No: **WO17/132173**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/286,858	2016.01.25	US

(43) Publication Date: **2017.08.03**

(44) Accepted Journal Date: **2021.04.01**

(44) Amended Journal Date: **2021.10.07**

(71) Applicant(s)
Regeneron Pharmaceuticals, Inc.

(72) Inventor(s)
Nittoli, Thomas;Markotan, Thomas P.

(74) Agent / Attorney
Phillips Ormonde Fitzpatrick, PO Box 323, COLLINS STREET WEST, VIC, 8007, AU

(56) Related Art
WO 2012177837 A2
WO 2012061590 A1



- (51) International Patent Classification:
C07D 498/18 (2006.01) *A61P 35/00* (2006.01)
A61K 31/535 (2006.01)
- (21) International Application Number:
PCT/US2017/014782
- (22) International Filing Date:
24 January 2017 (24.01.2017)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
62/286,858 25 January 2016 (25.01.2016) US
- (71) Applicant: **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US).
- (72) Inventors: **NITTOLI, Thomas**; c/o Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, New York 10591 (US). **MARKOTAN, Thomas P.**; 518 Washington Crossing Road, Newtown, Pennsylvania 19840 (US).
- (74) Agents: **OSTOMEL, Todd** et al.; Squire Patton Boggs (US) LLP, 620 Hansen way, Palo Alto, California 94304 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

[Continued on next page]

- (54) Title: MAYTANSINOID DERIVATIVES, CONJUGATES THEREOF, AND METHODS OF USE




FIG. 1

- (57) Abstract: Provided herein are maytansinoid compounds, derivatives thereof, conjugates thereof, and methods of treating or preventing proliferative diseases with the same.

Published:

— *with international search report (Art. 21(3))*

**MAYTANSINOID DERIVATIVES,
CONJUGATES THEREOF, AND METHODS OF USE**

[0001] This application claims priority to, and the benefit of, US Provisional Patent Application No. 62/286,858, entitled MAYTANSINOID DERIVATIVES, CONJUGATES THEREOF AND METHODS OF USE, which was filed January 25, 2016. The content of this provisional patent application is herein incorporated by reference in its entirety for all purposes.

FIELD

[0002] The present disclosure concerns maytansinoid compounds, derivatives thereof, conjugates thereof, and methods of treating or preventing proliferative diseases with the same.

BACKGROUND


[0003] Proliferative diseases, for example cancer, are characterized by the uncontrolled growth of abnormal cells. Current treatments of proliferative diseases include surgery, radiation, chemotherapy, hormone-based therapy and/or immunotherapy. A number of these treatments, particularly chemotherapy, utilize anti-proliferative drugs that limit the spread of the abnormal cells. However, these drugs are typically indiscriminate in their ability to kill cells, affecting both normal and abnormal cells. To address this problem, various approaches to targeted drug delivery have been explored, including the use of conjugates of tumor-targeted probes (such as antibodies or growth factors) with toxins, to selectively target abnormal cells. Antibody drug conjugates (ADCs) are compounds composed of an antibody that is linked, via a chemical linker, to a cytotoxic agent. Such compounds leverage the antibody's binding specificity for its target to deliver a cytotoxic agent to an abnormal cell. Thus, there is a need for anti-proliferative compounds and their conjugates.

[0003a] The discussion of documents, acts, materials, devices, articles and the like is included in this specification solely for the purpose of providing a context for the present invention. It is not suggested or represented that any or all of these matters formed part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

[0003b] Where the terms "comprise", "comprises", "comprised" or "comprising" are used in this specification (including the claims) they are to be interpreted as specifying the presence of the stated features, integers, steps or components, but not precluding the presence of one or more other features, integers, steps or components, or group thereof.

SUMMARY

[0004] Provided herein are compounds of Formula (I):



(I)

or a pharmaceutically acceptable salt thereof,

wherein:


A is a arylene or heteroarylene, optionally substituted as described herein;

L is a linker, optionally substituted as described herein;

BA is a binding agent, wherein the binding agent is an antibody or an antigen binding fragment thereof; and

k is an integer from 1 to 30. Also provided herein are stereoisomers of compounds of Formula (I).


[0005] Provided herein are also compounds of Formula (II):



(II)

or a pharmaceutically acceptable salt thereof, wherein A is arylene or heteroarylene, optionally substituted as described herein. Also provided herein are stereoisomers of compounds of Formula (II)

[0006] Provided herein are also compounds of Formula PP5:




PP5

or a salt thereof, wherein A is arylene or heteroarylene, optionally substituted as described herein. Also provided herein are stereoisomers of compounds of Formula PP5.

[0007] Furthermore, provided herein are methods of treating proliferative diseases comprising administering the compounds described herein.

[0008] Furthermore, provided herein are methods of treating proliferative diseases comprising administering the conjugates described herein.

[0009] Furthermore, provided herein are methods of making a compound of Formula (I) comprising contacting a compound of Formula P1 with a binding agent,



[0010] Furthermore, provided herein are methods of making a compound of Formula P1, comprising contacting a compound of Formula (II) with a reactive linker (RL) described herein.

[0011] Furthermore, provided herein are methods of making a compound of Formula (II), comprising contacting a compound of Formula PP5 with a suitable reducing agent.

[0012] Furthermore, provided herein are methods of making a compound of Formula PP5 comprising contacting a maytansinol with a nitro-phenyl-isocyanate.

BRIEF DESCRIPTIONS OF THE DRAWINGS

- [0013] FIG. 1 depicts the plot of % Cell Viability OVCAR3 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0014] FIG 2 depicts the plot of % Cell Viability SKBr3 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0015] FIG. 3 depicts the plot of % Cell Viability SKBr3 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0016] FIG 4 depicts the plot of % Cell Viability SKBr3 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0017] FIG. 5 depicts the plot of % Cell Viability SKBr3 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0018] FIG 6 depicts the plot of % Cell Viability C4-2 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0019] FIG. 7 depicts the plot of % Cell Viability C4-2 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0020] FIG 8 depicts the plot of % Cell Viability C4-2 vs. Log_{10} [M] of certain compounds tested in Example 28.
- [0021] FIG. 9 depicts a reaction scheme referenced in Example 1.
- [0022] FIG. 10 depicts a reaction scheme referenced in Example 2.
- [0023] FIG. 11 depicts a reaction scheme referenced in Example 3.
- [0024] FIG. 12 depicts a reaction scheme referenced in Example 4.
- [0025] FIG. 13 depicts a reaction scheme referenced in Example 5.
- [0026] FIG. 14 depicts a reaction scheme referenced in Example 6.
- [0027] FIG. 15 depicts a reaction scheme referenced in Example 7.
- [0028] FIG. 16 depicts a reaction scheme referenced in Example 8.
- [0029] FIG. 17 depicts a reaction scheme referenced in Example 9.
- [0030] FIG. 18 shows the Deconvoluted Mass Spectrum of MCC-DM1 Conjugated mAb from Example 26.

DETAILED DESCRIPTION**A. DEFINITIONS**

[0031] As used herein, “alkyl” refers to a monovalent and saturated hydrocarbon radical moiety. Alkyl is optionally substituted and can be linear, branched, or cyclic, *i.e.*, cycloalkyl. Alkyl includes, but is not limited to, those having 1-20 carbon atoms, *i.e.*, C₁₋₂₀ alkyl; 1-12 carbon atoms, *i.e.*, C₁₋₁₂ alkyl; 1-8 carbon atoms, *i.e.*, C₁₋₈ alkyl; 1-6 carbon atoms, *i.e.*, C₁₋₆ alkyl; and 1-3 carbon atoms, *i.e.*, C₁₋₃ alkyl. Examples of alkyl moieties include, but are not limited to methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *s*-butyl, *t*-butyl, *i*-butyl, a pentyl moiety, a hexyl moiety, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

[0032] As used herein, “haloalkyl” refers to alkyl, as defined above, wherein the alkyl includes at least one substituent selected from a halogen, *e.g.*, F, Cl, Br, or I.

[0033] As used herein, “alkenyl” refers to a monovalent hydrocarbon radical moiety containing at least two carbon atoms and one or more non-aromatic carbon-carbon double bonds. Alkenyl is optionally substituted and can be linear, branched, or cyclic. Alkenyl includes, but is not limited to, those having 2-20 carbon atoms, *i.e.*, C₂₋₂₀ alkenyl; 2-12 carbon atoms, *i.e.*, C₂₋₁₂ alkenyl; 2-8 carbon atoms, *i.e.*, C₂₋₈ alkenyl; 2-6 carbon atoms, *i.e.*, C₂₋₆ alkenyl; and 2-4 carbon atoms, *i.e.*, C₂₋₄ alkenyl. Examples of alkenyl moieties include, but are not limited to vinyl, propenyl, butenyl, and cyclohexenyl.

[0034] As used herein, “alkynyl” refers to a monovalent hydrocarbon radical moiety containing at least two carbon atoms and one or more carbon-carbon triple bonds. Alkynyl is optionally substituted and can be linear, branched, or cyclic. Alkynyl includes, but is not limited to, those having 2-20 carbon atoms, *i.e.*, C₂₋₂₀ alkynyl; 2-12 carbon atoms, *i.e.*, C₂₋₁₂ alkynyl; 2-8 carbon atoms, *i.e.*, C₂₋₈ alkynyl; 2-6 carbon atoms, *i.e.*, C₂₋₆ alkynyl; and 2-4 carbon atoms, *i.e.*, C₂₋₄ alkynyl. Examples of alkynyl moieties include, but are not limited to ethynyl, propynyl, and butynyl.

[0035] As used herein, “alkoxy” refers to a monovalent and saturated hydrocarbon radical moiety wherein the hydrocarbon includes a single bond to an oxygen atom and wherein the radical is localized on the oxygen atom, *e.g.*, CH₃CH₂-O· for ethoxy. Alkoxy substituents bond to the compound which they substitute through this oxygen atom of the alkoxy substituent. Alkoxy is optionally substituted and can be linear, branched, or cyclic, *i.e.*, cycloalkoxy. Alkoxy includes, but is not limited to, those having 1-20 carbon atoms, *i.e.*, C₁₋₂₀ alkoxy; 1-12 carbon atoms, *i.e.*, C₁₋₁₂ alkoxy; 1-8 carbon atoms, *i.e.*, C₁₋₈ alkoxy; 1-6 carbon atoms, *i.e.*, C₁₋₆ alkoxy; and 1-3 carbon atoms, *i.e.*, C₁₋₃ alkoxy. Examples of alkoxy moieties include, but are not limited

to methoxy, ethoxy, *n*-propoxy, *i*-propoxy, *n*-butoxy, *s*-butoxy, *t*-butoxy, *i*-butoxy, a pentoxy moiety, a hexoxy moiety, cyclopropoxy, cyclobutoxy, cyclopentoxy, and cyclohexoxy.

[0036] As used herein, “haloalkoxy” refers to alkoxy, as defined above, wherein the alkoxy includes at least one substituent selected from a halogen, *e.g.*, F, Cl, Br, or I.

[0037] As used herein, “aryl” refers to a monovalent moiety that is a radical of an aromatic compound wherein the ring atoms are carbon atoms. Aryl is optionally substituted and can be monocyclic or polycyclic, *e.g.*, bicyclic or tricyclic. Examples of aryl moieties include, but are not limited to those having 6 to 20 ring carbon atoms, *i.e.*, C₆₋₂₀ aryl; 6 to 15 ring carbon atoms, *i.e.*, C₆₋₁₅ aryl, and 6 to 10 ring carbon atoms, *i.e.*, C₆₋₁₀ aryl. Examples of aryl moieties include, but are limited to phenyl, naphthyl, fluorenyl, azulenyl, anthryl, phenanthryl, and pyrenyl.

[0038] As used herein, “arylene” refers to a divalent moiety of an aromatic compound wherein the ring atoms are only carbon atoms. Arylene is optionally substituted and can be monocyclic or polycyclic, *e.g.*, bicyclic or tricyclic. Examples of aryl moieties include, but are not limited to those having 6 to 20 ring carbon atoms, *i.e.*, C₆₋₂₀ arylene; 6 to 15 ring carbon atoms, *i.e.*, C₆₋₁₅ arylene, and 6 to 10 ring carbon atoms, *i.e.*, C₆₋₁₀ arylene.

[0039] As used herein, “alkaryl” refers to an aryl that is substituted with at least one alkyl. Alkaryl is optionally substituted.

[0040] As used herein, “heteroalkyl” refers to an alkyl in which one or more carbon atoms are replaced by heteroatoms. As used herein, “heteroalkenyl” refers to an alkenyl in which one or more carbon atoms are replaced by heteroatoms. As used herein, “heteroalkynyl” refers to an alkenyl in which one or more carbon atoms are replaced by heteroatoms. Suitable heteroatoms include, but are not limited to, nitrogen, oxygen, and sulfur atoms. Heteroalkyl is optionally substituted.

[0041] As used herein, “heteroaryl” refers to a monovalent moiety that is a radical of an aromatic compound wherein the ring atoms contain carbon atoms and at least one oxygen, sulfur, nitrogen, or phosphorus atom. Examples of heteroaryl moieties include, but are not limited to those having 5 to 20 ring atoms; 5 to 15 ring atoms; and 5 to 10 ring atoms. Heteroaryl is optionally substituted.

[0042] As used herein, “heteroarylene” refers to an arylene in which one or more carbon ring atoms of the aromatic ring are replaced with an oxygen, sulfur, nitrogen, or phosphorus

atom. Heteroarylene is optionally substituted and can be monocyclic or polycyclic, e.g., bicyclic or tricyclic.

[0043] As used herein, “optionally substituted,” when used to describe a radical moiety, e.g., optionally substituted alkyl, means that such moiety is optionally bonded to one or more substituents. Examples of such substituents include, but are not limited to halo, cyano, nitro, haloalkyl, azido, epoxy, optionally substituted heteroaryl, optionally substituted

heterocycloalkyl, $-\xi-OR^A$, $-\xi-SR^A$, $-\xi-NR^A R^B$, $-\xi-\overset{O}{\parallel}-R^A$, $-\xi-\overset{O}{\parallel}-OR^A$, $-\xi-\overset{O}{\parallel}-NR^A R^B$,
 $-\xi-NR^C-\overset{O}{\parallel}-R^A$, $-\xi-\overset{N}{\parallel}-NR^A R^B$, $-\xi-NR^C-\overset{N}{\parallel}-NR^A R^B$, $-\xi-S(O)-R^A$, $-\xi-S(O)_2-R^A$, $\overset{R^A}{\underset{|}{\underset{|}{N}}}$, $\overset{O}{\underset{|}{\underset{|}{N}}}$, or
 $\overset{S}{\underset{|}{\underset{|}{N}}}$, wherein R^A , R^B , and R^C are, independently at each occurrence, a hydrogen atom, alkyl, alkenyl, alkynyl, aryl, alkaryl, arylalkyl, heteroaryl, or heterocycloalkyl, or R^A and R^B , together with the atoms to which they are bonded, form a saturated or unsaturated carbocyclic ring, wherein the ring is optionally substituted and wherein one or more ring atoms is optionally replaced with a heteroatom. In some embodiments, R^A , R^B , and R^C are not hydrogen atoms.

[0044] As used herein, “binding agent” refers to any molecule capable of binding with specificity to a given binding partner.

[0045] As used herein, “linker” refers to a divalent moiety that covalently links the binding agent to the maytansinoid compounds and derivatives described herein.

[0046] As used herein, “amide synthesis conditions” refers to reaction conditions suitable to effect the formation of an amide, e.g., by the reaction of a carboxylic acid, activated carboxylic acid, or acyl halide with an amine. In some examples, amide synthesis conditions refer to reaction conditions suitable to effect the formation of an amide bond between a carboxylic acid and an amine. In some of these examples, the carboxylic acid is first converted to an activated carboxylic acid before the activated carboxylic acid reacts with an amine to form an amide. Suitable conditions to effect the formation of an amide include, but are not limited to, those utilizing reagents to effect the reaction between a carboxylic acid an amine, including, but not limited to, dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), (7-azabenzotriazol-1-


xyloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP), bromotripyrrolidinophosphonium hexafluorophosphate (PyBrOP), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate (HATU), 2-Ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), 2-Chloro-1,3-dimethylimidazolidinium hexafluorophosphate (CIP), 2-chloro-4,6-dimethoxy-1,3,5-triazine (CDMT), and carbonyldiimidazole (CDI). In some examples, a carboxylic acid is first converted to an activated carboxylic ester before reacting with an amine to form an amide bond. In certain embodiments, the carboxylic acid is reacted with a reagent. The reagent activates the carboxylic acid by deprotonating the carboxylic acid and then forming a product complex with the deprotonated carboxylic acid as a result of nucleophilic attack by the deprotonated carboxylic acid onto the protonated reagent. For certain carboxylic acids, this activated ester is more susceptible subsequently to nucleophilic attack by an amine than the carboxylic acid is before it is converted into an activated ester. This results in amide bond formation. As such, the carboxylic acid is described as activated. Exemplary reagents include DCC and DIC.

[0047] As used herein, “therapeutically effective amount” refers to an amount (of a compound) that is sufficient to provide a therapeutic benefit to a patient in the treatment or management of a disease or disorder, or to delay or minimize one or more symptoms associated with the disease or disorder.


[0048] As used herein, “Lewis acid” refers to a molecule or ion that accepts an electron lone pair. The Lewis acids used in the methods described herein are those other than protons. Lewis acids include, but are not limited to, non-metal acids, metal acids, hard Lewis acids, and soft Lewis acids. Lewis acids include, but are not limited to, Lewis acids of aluminum, boron, iron, tin, titanium, magnesium, copper, antimony, phosphorus, silver, ytterbium, scandium, nickel, and zinc. Illustrative Lewis acids include, but are not limited to, AlBr₃, AlCl₃, BCl₃, boron trichloride methyl sulfide, BF₃, boron trifluoride methyl etherate, boron trifluoride methyl sulfide, boron trifluoride tetrahydrofuran, dicyclohexylboron trifluoromethanesulfonate, iron (III) bromide, iron (III) chloride, tin (IV) chloride, titanium (IV) chloride, titanium (IV) isopropoxide, Cu(OTf)₂, CuCl₂, CuBr₂, zinc chloride, alkylaluminum

halides (R_nAlX_{3-n} , wherein R is hydrocarbonyl), $Zn(OTf)_2$, $ZnCl_2$, $Yb(OTf)_3$, $Sc(OTf)_3$, $MgBr_2$, $NiCl_2$, $Sn(OTf)_2$, $Ni(OTf)_2$, and $Mg(OTf)_2$.


[0049] Certain groups, moieties, substituents, and atoms are depicted with a wiggly line that intersects a bond or bonds to indicate the atom through which the groups, moieties, substituents, atoms are bonded. For example, a phenyl group that is substituted with a propyl group depicted as:




has the following structure:





As used herein, illustrations showing substituents bonded to a cyclic group (*e.g.*, aromatic, heteroaromatic, fused ring, and saturated or unsaturated cycloalkyl or heterocycloalkyl) through a bond between ring atoms are meant to indicate, unless specified otherwise, that the cyclic group may be substituted with that substituent at any ring position in the cyclic group or on any ring in the fused ring group, according to techniques set forth herein or which are known in the field to which the instant disclosure pertains. For example, the group,





, wherein subscript q is an integer from 0 to 4 and in which the positions of substituent R^1 are described generically, includes the following groups in which the positions of substituent







example, the group, in which the positions of substituents other than R¹ which are bonded to the cyclic group through a bond between ring atoms are described generically, includes the following groups in which the positions of these substituents other than R¹ are described specifically:



Also, for example, the group, in which the positions of substituents other than R¹ which are bonded to the cyclic group through a bond between ring atoms are described generically, includes the following groups in which the positions of these substituents other than R¹ are described specifically:




each of these structures in which the positions of the substituents other than R^1 are described specifically, the substituent R^1 may be bonded to any ring position in the cyclic group or on any ring in the fused ring group which is not occupied by one of these substituents other than R^1 . The following non-limiting representative illustrations indicate that the cyclic group can be substituted with the indicated substituent at any ring position or on either ring in the fused ring




[0050] When a group described herein is optionally substituted, the substituent bonded to the group is unsubstituted unless otherwise specified.

[0051] As used herein, the phrase “reactive linker,” or the abbreviation “RL” refers to a monovalent group that includes a reactive group and spacer group, depicted for example as $RG-SP-\xi$, wherein RG is the reactive group and SP is the spacer group. As described herein, a reactive linker may include more than one reactive group and more than one spacer group. The spacer group is any divalent moiety that bridges the reactive group to another group, such as a payload. The reactive linkers (RL), together with the payloads to which they are bonded, provide intermediates (“linker-payloads”) useful as synthetic precursors for the preparation of the antibody conjugates described herein. The reactive linker contains a reactive group (“RG”), which is a functional group or moiety that is capable of reacting with a reactive portion of another group, for instance, an antibody, modified antibody, or antigen binding fragment thereof, or an enhancement group. The moiety resulting from the reaction of the reactive group with the


antibody, modified antibody, or antigen binding fragment thereof, together with the linking group, include the “binding agent linker” (“BL”) portion of the conjugate, described herein. In certain embodiments, the “reactive group” is a functional group or moiety (*e.g.*, maleimide or *N*-hydroxysuccinimide (NHS) ester) that reacts with a cysteine or lysine residue of an antibody or antigen-binding fragment thereof. In certain embodiments, the “reactive group” is a functional group or moiety that is capable of undergoing a click chemistry reaction (*see, e.g.*, click chemistry, Huisgen *Proc. Chem. Soc.* **1961**, Wang *et al. J. Am. Chem. Soc.* **2003**, and Agard *et al. J. Am. Chem. Soc.* **2004**). In some embodiments of said click chemistry reaction, the reactive group is an alkyne that is capable of undergoing a 1,3-cycloaddition reaction with an azide. Such suitable reactive groups include, but are not limited to, strained alkynes, *e.g.*, those suitable for strain-promoted alkyne-azide cycloadditions (SPAAC), cycloalkynes, *e.g.*, cyclooctynes, benzannulated alkynes, and alkynes capable of undergoing 1,3-cycloaddition reactions with alkynes in the absence of copper catalysts. Suitable alkynes also include, but are not limited to,



derivatives thereof. Particularly useful alkynes include




and




. Linker-payloads including such reactive groups are useful for conjugating antibodies that have been functionalized with azido groups. Such functionalized antibodies include antibodies functionalized with azido-polyethylene glycol groups. In certain embodiments, such functionalized antibody is derived by treating an antibody having at least one glutamine residue, *e.g.*, heavy chain Gln195, with a compound bearing an azide group, in the presence of the enzyme transglutaminase.

[0052] In some examples, the reactive group is an alkyne, *e.g.*,



, which can react via click chemistry with an azide, *e.g.*, $\text{N}=\text{N}=\text{N}$, to form a click


chemistry product, *e.g.*,



or


. In some examples, the group reacts with an azide on a modified antibody or antigen binding fragment thereof. In some examples, the

reactive group is an alkyne, *e.g.*,




, which can react via click chemistry with an

azide, *e.g.*,




to form a click chemistry product, *e.g.*,






. In some


examples, the reactive group is an alkyne, *e.g.*,




, which can react via click chemistry


with an azide, e.g., $\text{N}=\text{N}=\text{N}$, to form a click chemistry product, e.g.,  or .

In some examples, the reactive group is a functional group, e.g., , which reacts with a cysteine residue on an antibody or antigen-binding fragment thereof, to form a bond thereto, e.g.,



, wherein Ab refers to an antibody or antigen-binding fragment thereof and S refers to the S atom on a cysteine residue through which the functional group bonds to the Ab. In some


examples, the reactive group is a functional group, e.g., , which reacts with a lysine residue on an antibody or antigen-binding fragment thereof, to form a bond thereto, e.g.,



, wherein Ab refers to an antibody or antigen-binding fragment thereof and NH refers to the NH atom on a lysine side chain residue through which the functional group bonds to the Ab.

B. CONJUGATES

[0053] Provided herein are compounds of Formula (I):



(I)

or a pharmaceutically acceptable salt thereof,
wherein:

A is arylene or heteroarylene;

L is a linker;

BA is a binding agent, wherein the binding agent is an antibody or an antigen binding fragment thereof; and

k is an integer from 1 to 30.

1. "A" moieties

[0054] In some embodiments, A is arylene. In some embodiments, A is heteroarylene. In some embodiments, the arylene or heteroarylene is substituted with one or more electron withdrawing groups and/or one or more electron donating groups.


[0055] In some embodiments, A is a divalent radical of benzene, of pyridine, of naphthalene, or of quinolone, which are optionally substituted.

[0056] In some embodiments, A is a divalent radical of benzene which is optionally substituted with a member selected from the group consisting of amino, amido, alkyl, halo, haloalkyl, alkoxy, and haloalkoxy.

[0057] In some embodiments, A is a divalent radical of benzene which is optionally substituted with a member selected from the group consisting of alkyl, alkoxy, haloalkyl, and halo. In some embodiments, A is a divalent radical of benzene which is optionally substituted with a member selected from the group consisting of methyl, methoxy, trifluoromethyl, fluoro, chloro, and bromo. In some embodiments, A is a divalent radical of benzene which is optionally substituted with a member selected from the group consisting of halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl,

cyano, nitro, amino, $-\text{OR}^a$, $-\text{SO}_2\text{R}^a$, $-\text{C}(=\text{O})\text{R}^a$, or azido, wherein R^a is alkyl.

[0058] In some embodiments, A is:



wherein:

R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido,


wherein R^A is alkyl;

n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5.



[0059] In some embodiments, A is:


wherein:

R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido,

wherein R^A is alkyl; and

n is an integer from 0 to 4. In some embodiments, n is 0, 1, 2, or 3. In some embodiments, n is 0, 1, or 2. In some embodiments, n is 0 or 1. In some embodiments, n is 1.




[0060] In some embodiments, A is:

wherein:

R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; and

n is an integer from 0 to 4. In some embodiments, n is 0, 1, 2, or 3. In some embodiments, n is 0, 1, or 2. In some embodiments, n is 0 or 1. In some embodiments, n is 1.

[0061] In some embodiments, A is:



wherein:

R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido,

wherein R^A is alkyl;

n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5.

In some embodiments, R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl,

cyano, nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido,

wherein R^A is alkyl;


[0062] In some embodiments, R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.

[0063] In some embodiments, R^1 is C_{1-6} alkyl or C_{1-6} alkoxy. In some embodiments, R^1 is C_1 alkyl. In some embodiments, R^1 is C_1 alkoxy. In some embodiments, R^1 is halo. In some embodiments, R^1 is haloalkyl. In some embodiments, R^1 is heterocycloalkyl. In some of these embodiments, R^1 is methyl, ethyl, methoxy, or ethoxy. In some of these embodiments, R^1 is methyl. In some of these embodiments, R^1 is methoxy. In some of these embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro, chloro, or bromo. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some of these embodiments, R^1 is pyrrolidinyl. In some of these embodiments, R^1 is morpholinyl.


[0064] In some embodiments, R^1 is, independently, alkyl or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} haloalkyl, or halo. In some embodiments, R^1 is, independently, halo. In some embodiments, R^1 is, independently, fluoro, chloro, bromo, iodo, or trifluoromethyl. In some embodiments, n , m , p , or q is 0, 1 or 2. In some embodiments, n , m , p , or q is 0 or 1. In some embodiments, n , m , p , or q is 0.

[0065] In some embodiments, R^1 is, independently, alkyl, alkoxy, or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or halo. In some embodiments, R^1 is, independently, halo. In some embodiments, R^1 is, independently, fluoro, chloro, bromo, iodo, or trifluoromethyl. In some embodiments, R^1 is C_1 alkoxy. In some embodiments, R^1 is halo. In some of these embodiments, R^1 is methyl. In some of these embodiments, R^1 is methoxy. In some of these embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro, chloro, or bromo. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some embodiments, n, m, p, or q is 0, 1 or 2. In some embodiments, n, m, p, or q is 0 or 1. In some embodiments, n, m, p, or q is 0.


[0066] In some embodiments, A is:



[0067] In some embodiments, A is:




[0068] In some embodiments, A is:



wherein n is 0, 1, 2, 3, or 4.

[0069] In some embodiments, A is:




wherein n is 0, 1, or 2. In some examples, n is 0 or 1; and R^1 is alkyl, alkoxy, halo, haloalkyl, or haloalkoxy. In some examples, R^1 is alkyl, alkoxy, halo, haloalkyl, or heterocycloalkyl. In some embodiments, R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl,

alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,


nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido, wherein R^A is alkyl.

[0070] In some embodiments, A is:




wherein n is 0, 1, or 2. In some examples, n is 0 or 1; and R^1 is C_{1-6} alkyl, C_{1-6} alkoxy, halo, C_{1-6} haloalkyl, C_{1-6} haloalkoxy. In some examples, R^1 is heterocycloalkyl. In some embodiments, R^1 is methyl, methoxy, trifluoromethyl, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is bromo. In some embodiments, R^1 is chloro. In some examples, R^1 is pyrrolidinyl. In some examples, R^1 is morpholinyl.

[0071] In some embodiments, A is:




wherein n is 0, 1, or 2. In some examples, n is 0 or 1; R^1 is C_{1-6} alkyl, C_{1-6} alkoxy, halo, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or heterocycloalkyl; and L is



, wherein b is an integer from 2 to 8 and ξ^A is a bond


to the binding agent.

[0072] In some embodiments, A is:




wherein n is 0, 1, 2, 3, or 4.

[0073] In some embodiments, A is:



wherein R^1 is C_{1-6} alkyl, C_{1-6} alkoxy, halo, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or heterocycloalkyl. In certain of these embodiments, R^1 is methoxy or methyl. In some specific embodiments, R^1 is methoxy. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methyl, methoxy, trifluoromethyl, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is bromo. In some embodiments, R^1 is chloro. In some examples, R^1 is pyrrolidinyl. In some examples, R^1 is morpholinyl. In some examples, R^1 is alkyl, alkoxy, halo, haloalkyl, or heterocycloalkyl. In some embodiments, R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano, nitro, amino, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido, wherein R^A is alkyl.


[0074] In some embodiments, A is:




wherein:

R^1 is halo, methyl, methoxy, or trifluoromethyl; and
n is 0, 1 or 2.

[0075] In some embodiments, A is:




[0076] In some embodiments, A is:



wherein:

X is a hydrogen atom, halo, methyl, methoxy, or trifluoromethyl.

[0077] In some embodiments, A is:




wherein:

X is a hydrogen atom, halo, methyl, methoxy, or trifluoromethyl;

$\text{---}\xi^1\text{---}$ is the bond to the nitrogen atom of the amino-ester which is directly bonded to the drug molecule; and $\text{---}\xi^2\text{---}$ is the bond to the nitrogen atom which is bonded to the linker.

[0078] In some embodiments, A is:




wherein:

R^1 is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, heterocycloalkyl, halo, haloalkyl, or haloalkoxy;

$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above. In some embodiments, R^1 is 1-methylethylthiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some embodiments, R^1 is hydrogen.

[0079] In some embodiments, A is:




wherein:

R^1 is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, heterocycloalkyl, halo, haloalkyl, haloalkoxy;

$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ as a defined above. In some embodiments, R^1 is 1-methylethylthiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is methyl. In some

embodiments, R¹ is methoxy. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo. In some embodiments, R¹ is hydrogen.


[0080] In some embodiments, A is:



wherein:


R¹ is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, heterocycloalkyl, halo, haloalkyl, or haloalkoxy; ξ^1 and ξ^2 are as defined above. In some embodiments, R¹ is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, heterocycloalkyl, halo, haloalkyl, or haloalkoxy. In some embodiments, R¹ is alkyl or alkoxy. In some specific embodiments, R¹ is propylamino, difluoro-methoxy, phenyl, 2-fluorophenyl. In some embodiments, R¹ is trifluoromethyl. In some embodiments, R¹ is methoxy. In some embodiments, R¹ is methyl. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo. In some embodiments, R¹ is hydrogen. In some examples, R¹ is pyrrolidinyl. In some examples, R¹ is morpholinyl.

[0081] In some embodiments, A is:




ξ^1 and ξ^2 are as defined above.

[0082] In some embodiments, A is:




ξ^1 and ξ^2 are as defined above.

[0083] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0084] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0085] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0086] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0087] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0088] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0089] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0090] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0091] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0092] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0093] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0094] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0095] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0096] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0097] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0098] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0099] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0100] In some embodiments, A is:




$\text{---}\overset{s1}{\text{~}}\text{---}$ and $\text{---}\overset{s2}{\text{~}}\text{---}$ are as defined above.

[0101] In some embodiments, A is:




$\text{---}\overset{s1}{\text{~}}\text{---}$ and $\text{---}\overset{s2}{\text{~}}\text{---}$ are as defined above.

[0102] In some embodiments, A is:




$\text{---}\overset{s1}{\text{~}}\text{---}$ and $\text{---}\overset{s2}{\text{~}}\text{---}$ are as defined above.

[0103] In some embodiments, A is:




$\text{---}\overset{s1}{\text{~}}\text{---}$ and $\text{---}\overset{s2}{\text{~}}\text{---}$ are as defined above.

[0104] In some embodiments, A is:




$\text{---}\frac{s^1}{\text{z}}\text{---}$ and $\text{---}\frac{s^2}{\text{z}}\text{---}$ are as defined above.

[0105] In some embodiments, A is:




$\text{---}\frac{s^1}{\text{z}}\text{---}$ and $\text{---}\frac{s^2}{\text{z}}\text{---}$ are as defined above.

[0106] In some embodiments, A is:




$\text{---}\frac{s^1}{\text{z}}\text{---}$ and $\text{---}\frac{s^2}{\text{z}}\text{---}$ are as defined above.

[0107] In some embodiments, A is:




$\text{---}\frac{s^1}{\text{z}}\text{---}$ and $\text{---}\frac{s^2}{\text{z}}\text{---}$ are as defined above.

[0108] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0109] In some embodiments, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above. In some embodiments, A is:



$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above .


[0110] In some embodiments, A is:



wherein each R¹ is independently, at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkyl, or haloalkoxy;


$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above . In some embodiments, R¹ is hydrogen, fluoro, methyl, trifluoromethyl, or methoxy. In some embodiments, R¹ is fluoro, chloro, bromo, or iodo.

[0111] In some embodiments, A is:




wherein $\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above .

[0112] In some embodiments, A is:




wherein each R¹ is independently, at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkyl, or haloalkoxy; wherein $\text{---}\xi^1$ and $\text{---}\xi^2$ are as defined above .

[0113] In some embodiments, A is:




wherein $\text{---}\xi^1$ and $\text{---}\xi^2$ are as defined above .


[0114] In some embodiments, A is:




[0115] In some embodiments, A is:



[0116] In some embodiments, A is:




[0117] In some embodiments, A is:



wherein:

wherein $\text{---}\xi^1$ and $\text{---}\xi^2$ are as defined above .


[0118] In some embodiments, A is:



wherein:

wherein $\text{-}\frac{S^1}{\text{Z}}$ and $\text{-}\frac{S^2}{\text{Z}}$ are as defined above .


[0119] In some embodiments, A is:



wherein:

wherein $\text{-}\frac{S^1}{\text{Z}}$ and $\text{-}\frac{S^2}{\text{Z}}$ are as defined above .


[0120] In some embodiments, A is:




wherein:

wherein $\text{-}\frac{S^1}{\text{Z}}$ and $\text{-}\frac{S^2}{\text{Z}}$ are as defined above .


[0121] In some embodiments, A is:



[0122] In some embodiments, A is:




[0123] In some embodiments, A is:




wherein n is 0, 1 2, or 3.

[0124] In some embodiments, A is:



wherein n is 0, 1 2, or 3.


[0125] In some embodiments, A is:




wherein:

R¹ is C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ heteroalkyl, or heterocycloalkyl; and n is 0, 1 2, 3 or 4. In some embodiments, R¹ is C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ heteroalkyl, or heterocycloalkyl.


[0126] In some embodiments, A is:



[0127] In some embodiments, A is:




[0128] In some embodiments, A is:



wherein:

X is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X is independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X is fluoro, chloro, bromo, iodo, dimethylamino, methylamino, methyl, methoxy, ethoxy, or trifluoromethyl. In some embodiments, X is methyl. In some embodiments, X is methoxy. In some embodiments, X is trifluoromethyl. In some embodiments, X is fluoro. In some embodiments, X is chloro. In some embodiments, X is bromo. In some embodiments, X is morpholinyl. In some embodiments, X is pyrrolidinyl.

[0129] In some embodiments, A is:




wherein:

X is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X is independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X, independently at each occurrence, fluoro, chloro, bromo, iodo, dimethylamino, methyl, methylamino, methoxy, ethoxy, trifluoromethyl methoxy, pyrrolidinyl, or morpholinyl;

wherein $\text{-}\overset{1}{\underset{2}{\text{C}}}$ and $\text{-}\overset{2}{\underset{1}{\text{C}}}$ are as defined above. In some embodiments, X is methyl. In some embodiments, X is methoxy. In some embodiments, X is trifluoromethyl. In some embodiments, X is fluoro. In some embodiments, X is chloro. In some embodiments, X is bromo. In some embodiments, X is morpholinyl. In some embodiments, X is pyrrolidinyl.


[0130] In some embodiments, A is:



wherein:

X is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X is independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X is fluoro, chloro, bromo, iodo, dimethylamino, methylamino, methyl, methoxy, ethoxy, or trifluoromethyl. In some embodiments, X is methyl. In some embodiments, X is methoxy. In some embodiments, X is trifluoromethyl. In some embodiments, X is fluoro. In some embodiments, X is chloro. In some embodiments, X is bromo. In some embodiments, X is morpholinyl. In some embodiments, X is pyrrolidinyl.

[0131] In some embodiments, A is:



wherein:

X is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, heteroalkyl, or heterocycloalkyl. In some embodiments, X, independently at each occurrence, fluoro, chloro, bromo, iodo, dimethylamino, methyl, methylamino, methoxy, ethoxy, trifluoromethyl methoxy, pyrrolidinyl, or morpholinyl;


wherein $\frac{\xi^1}{\xi}$ and $\frac{\xi^2}{\xi}$ are as defined above. In some embodiments, X is methyl. In some embodiments, X is methoxy. In some embodiments, X is trifluoromethyl. In some embodiments, X is fluoro. In some embodiments, X is chloro. In some embodiments, X is bromo. In some embodiments, X is morpholinyl. In some embodiments, X is pyrrolidinyl.


2. Linkers

[0132] The linker portion of the conjugates described herein is a divalent moiety that covalently links the binding agent to the maytansinoid compounds and derivatives described herein. Suitable linkers include those that release at least the maytansinoid portion in the presence of an enzyme or at a particular pH range or value.

[0133] In some embodiments, the linker comprises an enzyme-cleavable moiety. Illustrative enzyme-cleavable moieties include, but are not limited to, peptide bonds, ester linkages, hydrazones, and disulfide linkages. In some embodiments, the linker comprises a cathepsin-cleavable linker.

[0134] In some embodiments, the reactive linker comprises a non-cleavable moiety. In

some embodiments, the non-cleavable reactive linker is  or a residue


thereof. In some embodiments, the non-cleavable reactive linker is  or a residue thereof.

[0135] Suitable linkers also include, but are not limited to, those that are chemically bonded to two cysteine residues of a single binding agent, *e.g.*, antibody. Such linkers can serve to mimic the antibody's disulfide bonds that are disrupted as a result of the conjugation process.

[0136] In some embodiments, the linker comprises one or more amino acids. Suitable amino acids include natural, non-natural, standard, non-standard, proteinogenic, non-proteinogenic, and L-, or D- α - amino acids. In some embodiments, the linker comprises alanine, valine, leucine, isoleucine, methionine, tryptophan, phenylalanine, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartic acid, glutamic acid, lysine, arginine, histidine, or citrulline, or derivative thereof.

[0137] In some embodiments, the linker comprises valine and citrulline.

[0138] In some embodiments, the linker is:



wherein:

SP is a spacer;

$-\overset{A}{\xi}-$ is one or more bonds to the binding agent;

AA¹ is an amino acid; and

AA² is an amino acid.

[0139] In some embodiments, the linker is:




wherein:

SP is a spacer; and

$-\overset{A}{\xi}-$ is one or more bonds to the binding agent

[0140] In some examples, the spacer is a divalent moiety that connects the AA¹-AA² moiety to the binding agent (BA). Suitable spacers include, but are not limited to, those comprising alkylene or polyethylene glycol. The ends of the spacers, *i.e.*, the portion of the spacer directly bonded to the binding agent or AA¹, can be moieties derived from reactive moieties that are used for purposes of coupling the naked antibody or AA¹ to the spacer during the chemical synthesis of the conjugate. In some examples, the ends of the spacers, *i.e.*, the portion of the spacer directly bonded to the binding agent or AA¹, can be residues of reactive moieties that are used for purposes of coupling the naked antibody or AA¹ to the spacer during the chemical synthesis of the conjugate.

[0141] In some embodiments, the spacer comprises an alkylene. In some embodiments, the spacer comprises a C₅₋₇ alkylene. In some embodiments, the spacer is:




wherein:

$\overset{A}{\sim}$ is a bond to the binding agent; and

b is an integer from 2 to 8. In some examples, b is selected from 2, 3, 4, 5, 6, 7, or 8. In some examples, b is 2. In some examples, b is 3. In some examples, b is 4. In some examples, b is 5. In some examples, b is 6. In some examples, b is 7. In some examples, b is 8.


[0142] In some embodiments, the spacer is:



wherein:

$\overset{A}{\sim}$ is a bond to the binding agent.


[0143] In some embodiments, the spacer is:



wherein:

$\overset{A}{\sim}$ is a bond to the binding agent.

[0144] In some embodiments, the spacer is:




wherein:

R^N is a hydrogen atom or alkyl;

R^M is alkyl;

the two bonds represented by $\overset{BA-S}{\text{---}}$ are bonds to cysteines of a binding agent; and
 b is an integer from 2 to 8.


[0145] In some embodiments, the spacer is:



wherein:

$\overset{A}{\text{---}}$ is a bond to the binding agent; and
 b is an integer from 2 to 8.


[0146] In some embodiments, the spacer is:




wherein:

$\overset{A}{\text{---}}$ is a bond to the binding agent; and
 g is an integer from 2 to 20. In some embodiments, g is 2-8. In some
 embodiments, g is 2, 4, 6, or 8.


[0147] In some embodiments, the spacer is:




[0148] In some embodiments, the spacer is:




[0149] In some embodiments, the spacer is:




[0150] In some embodiments, the spacer is:




[0151] In some embodiments, the spacer is:




[0152] In some embodiments, the spacer is:




[0153] In some embodiments, the spacer is:




[0154] In some embodiments, the spacer is:



[0155] In some embodiments, the spacer is:



[0156] In some embodiments, the spacer is:




[0157] wherein ---^A is a bond to the binding agent;

X is N or O; R^N and R^M are each, independently, hydrogen or alkyl; and b is an integer from 1 to 8.

[0158] In some embodiments, AA^1-AA^2 is: valine-citrulline, citrulline-valine, lysine-phenylalanine, phenylalanine-lysine, valine-asparagine, asparagine-valine, asparagine-threonine, threonine-asparagine, serine-asparagine, asparagine-serine, phenylalanine-asparagine, asparagine-phenylalanine, leucine-asparagine, asparagine-leucine, isoleucine-asparagine, asparagine-isoleucine, glycine-asparagine, asparagine-glycine, glutamic acid-asparagine, asparagine-glutamic acid, citrulline-asparagine, asparagine-citrulline, alanine-asparagine, asparagine-alanine, valine-alanine, alanine-valine, valine-glycine, or glycine-valine.

[0159] In some embodiments, AA^1-AA^2 is: valine-citrulline or citrulline-valine. In some embodiments, AA^1-AA^2 is: valine-citrulline.

[0160] In some embodiments, the linker is:



wherein:

SP is a spacer;


---^A is one or more bonds to the binding agent;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.

[0161] As used herein, “amino acid side chain” refers the monovalent non-hydrogen substituent bonded to the α -carbon of an α -amino acid, including natural and non-natural amino acids. Exemplary amino acid side chains include, but are not limited to, the α -carbon substituent of alanine, valine, leucine, isoleucine, methionine, tryptophan, phenylalanine, proline, serine, threonine, cysteine, tyrosine, asparagine, glutamine, aspartic acid, glutamic acid, lysine, arginine, histidine, and citrulline.

[0162] In some embodiments, the linker is:




wherein:

SP is a spacer; and

$\overset{A}{\text{---}}$ is one or more bonds to the binding agent.

[0163] In some embodiments, the linker is:




wherein:

$\overset{A}{\text{---}}$ is a bond to the binding agent; and

b is an integer from 2 to 8.

[0164] In some embodiments, the linker is:




wherein:

$\overset{A}{\text{---}}$ is a bond to the binding agent; and

b is an integer from 2 to 8.


[0165] In some embodiments, BA is an antibody and the linker is:



wherein:

the two bonds represented by $\overset{\text{BA-S}}{\text{---}}$ are bonds to cysteines of the antibody; and
b is an integer from 2 to 8.


[0166] In some embodiments, BA is an antibody and the linker is:



wherein:

the two bonds represented by $\overset{\text{BA-S}}{\text{---}}$ are bonds to cysteines of the antibody; and
b is an integer from 2 to 8.

[0167] In some embodiments, BA is an antibody and the linker is:



wherein:


R^N is a hydrogen atom or alkyl;

R^M is alkyl;


the two bonds represented by $\overset{\text{BA-S}}{\text{---}}$ are bonds to cysteines of the antibody; and

b is an integer from 2 to 8.

[0168] In some embodiments, the linker is:




wherein:


 is a bond to the binding agent; and

b is an integer from 2 to 8.

[0169] In some embodiments, the linker is:




wherein:


 is a bond to the binding agent; and

g is an integer from 2 to 20. In some embodiments, g is 2 to 8. In some embodiments, g is 2, 4, 6, or 8.


[0170] In some embodiments, the linker is:




[0171] In some embodiments, the linker is:




[0172] In some embodiments, the linker is:




[0173] In some embodiments, the linker is:




[0174] In some embodiments, the linker is:




[0175] In some embodiments, the linker is:



[0176] In some embodiments, the linker is:



[0177] In some embodiments, the linker is:



3. Binding agents

[0178] Suitable binding agents include, but are not limited to, antibodies, lymphokines, hormones, growth factors, viral receptors, interleukins, or any other cell binding or peptide binding molecules or substances.

[0179] In some embodiments, the binding agent is an antibody. In some embodiments, the antibody is a monoclonal antibody, polyclonal antibody, antibody fragment (Fab, Fab', and F(ab)2, minibody, diabody, tribody, and the like), or bispecific antibody. Antibodies herein can be humanized using methods described in US Patent No. 6,596,541 and US Publication No. 2012/0096572, each incorporated by reference in their entirety.

[0180] Where the binding agent is an antibody, it binds to an antigen binding partner that is a polypeptide and may be a transmembrane molecule (*e.g.*, receptor) or a growth factor that

might be glycosylated or phosphorylated. Exemplary antigens include, but are not limited to, molecules such as renin; a growth hormone, including human growth hormone and bovine growth hormone; growth hormone releasing factor; parathyroid hormone; thyroid stimulating hormone; lipoproteins; alpha1-antitrypsin; insulin A-chain; insulin B-chain; proinsulin; follicle stimulating hormone; calcitonin; luteinizing hormone; glucagon; clotting factors such as factor vmc, factor IX, tissue factor (TF), and von Willebrands factor; anti-clotting factors such as Protein C; atrial natriuretic factor; lung surfactant; a plasminogen activator, such as urokinase or human urine or tissue-type plasminogen activator (t-PA); bombesin; thrombin; hemopoietic growth factor; tumor necrosis factor-alpha and -beta; enkephalinase; RANTES (regulated on activation normally T-cell expressed and secreted); human macrophage inflammatory protein (MIP-I-alpha); a serum albumin, such as human serum albumin; Muellerian-inhibiting substance; relaxin A-chain; relaxin B-chain; prorelaxin; mouse gonadotropin-associated peptide; a microbial protein, such as betalactamase; DNase; 19E; a cytotoxic T-lymphocyte associated antigen (CTLA), such as CTLA-4; inhibin; activin; vascular endothelial growth factor (VEGF); receptors for hormones or growth factors; protein A or D; rheumatoid factors; a neurotrophic factor such as bone-derived neurotrophic factor (BDNF), neurotrophin-3, -4, -5, or -6 (NT-3, NT4, NT-5, or NT-6), or a nerve growth factor such as NGF- β ; platelet-derived growth factor (PDGF); fibroblast growth factor such as aFGF and bFGF; fibroblast growth factor receptor 2 (FGFR2), epidermal growth factor (EGF); transforming growth factor (TGF) such as TGF-alpha and TGF-beta, including TGF- β 1, TGF- β 2, TGF- β 3, TGF- β 4, or TGF- β 5; insulin-like growth factor-I and -II (IGF-I and IGF-II); des(I-3)-IGF-I (brain IGF-I), insulin-like growth factor binding proteins, EpCAM, GD3, FLT3, PSMA, PSCA, MUC1, MUC16, STEAP, STEAP2, CEA, TENB2, EphA receptors, EphB receptors, folate receptor, FOLRI, mesothelin, cripto, alphavbeta6, integrins, VEGF, VEGFR, EGFR, transferrin receptor, IRTA1, IRTA2, IRTA3, IRTA4, IRTA5; CD proteins such as CD2, CD3, CD4, CD5, CD6, CD8, CD11, CD14, CD19, CD20, CD21, CD22, CD25, CD26, CD28, CD30, CD33, CD36, CD37, CD38, CD40, CD44, CD52, CD55, CD56, CD59, CD70, CD79, CD80, CD81, CD103, CD105, CD134, CD137, CD138, CD152, or an antibody which binds to one or more tumor-associated antigens or cell-surface receptors disclosed in US Publication No. 2008/0171040 or US Publication No. 2008/0305044 and incorporated in their entirety by reference; erythropoietin; osteoinductive factors; immunotoxins; a bone morphogenetic protein (BMP); an interferon, such as interferon-

alpha, -beta, and -gamma; colony stimulating factors (CSFs), *e.g.*, M-CSF, GM-CSF, and G-CSF; interleukins (ILs), *e.g.*, IL-1 to IL-10; superoxide dismutase; T-cell receptors; surface membrane proteins; decay accelerating factor; viral antigen such as, for example, a portion of the HIV envelope; transport proteins; homing receptors; addressins; regulatory proteins; integrins, such as CD11a, CD11b, CD11c, CD18, an ICAM, VLA-4 and VCAM; a tumor associated antigen such as AFP, ALK, B7H4, BAGE proteins, β -catenin, bcr-abl, BRCA1, BORIS, CA9 (carbonic anhydrase IX), caspase-8, CD20, CD40, CD123, CDK4, CEA, CLEC12A, c-kit, cMET, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, endoglin, Epcam, EphA2, ErbB2/Her2, ErbB3/Her3, ErbB4/Her4, ETV6-AML, Fra-1, FOLR1, GAGE proteins, GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/EBNA1, HLA/k-ras, HLA/MAGE-A3, hTERT, IGF1R, LGR5, LMP2, MAGE proteins, MART-1, mesothelin, ML-IAP, Muc1, Muc16, CA-125, MUM1, NA17, NGEP, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PDGFR- α , PDGFR- β , PDGF-A, PDGF-B, PDGF-C, PDGF-D, PLAC1, PRLR, PRAME, PSCA, PSGR, PSMA (FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, Steap-1, Steap-2, STn, survivin, TAG-72, TGF- β , TMPRSS2, Tn, TNFRSF17, TRP-1, TRP-2, tyrosinase, and uroplakin-3, and fragments of any of the above-listed polypeptides. In some examples, the GAGE proteins are selected from GAGE-1 and GAGE-2. In some examples, the MAGE proteins are selected from MAGE-1, -2, -3, -4, -6, and -12

[0181] Exemplary antigens also include, but are not limited to, BCMA, SLAMF7, B7H4, GPNMB, UPK3A, and LGR5. Exemplary antigens also include, but are not limited to, MUC16, PSMA, STEAP2, and HER2.

[0182] In some embodiments, the antigens include prolactin receptor (PRLR) or prostate-specific membrane antigen (PSMA). In some embodiments, the antigens include MUC16. In some embodiments, the antigens include STEAP2. In some embodiments, the antigens include PSMA. In some embodiments, the antigens include HER2. In some embodiments, the antigen is prolactin receptor (PRLR) or prostate-specific membrane antigen (PSMA). In some embodiments, the antigen is MUC16. In some embodiments, the antigens include PSMA. In some embodiments, the antigen is HER2. In some embodiments, the antigen is STEAP2.

[0183] Binding agents also include, but are not limited to, ankyrin repeat proteins, interferons, lymphokines such as IL-2 or IL-3, hormones like insulin and glucocorticoids, growth factors such as EGF, transferrin and fibronectin type III.

[0184] In some embodiments, the binding agents interact with or bind to tumor antigens, including antigens specific for a type of tumor or antigens that are shared, overexpressed or modified on a particular type of tumor. Examples include, but are not limited to: alpha-actinin-4 with lung cancer, ARTC1 with melanoma, BCR-ABL fusion protein with chronic myeloid leukemia, B-RAF, CLPP or Cdc27 with melanoma, CASP-8 with squamous cell carcinoma, and hsp70-2 with renal cell carcinoma as well as the following shared tumor-specific antigens, for example: BAGE-1, GAGE, GnTV, KK-LC-1, MAGE-A2, NA88-A, TRP2-INT2.

[0185] In some embodiments, the binding agent is an antibody. In some embodiments, the binding agent is a monoclonal antibody. In some embodiments, the binding agent is a polyclonal antibody. In some embodiments, the antibody is an anti-PSMA, anti-MUC16, anti-HER2, or anti-EGFRvIII, or anti-STEAP-2 antibody.

[0186] The linkers can be bonded to the binding agent, *e.g.*, antibody or antigen-binding molecule, through an attachment at a particular amino acid within the antibody or antigen-binding molecule. Exemplary amino acid attachments that can be used in the context of this aspect of the disclosure include, *e.g.*, lysine (*see, e.g.*, US 5,208,020; US 2010/0129314; Hollander *et al.*, *Bioconjugate Chem.*, 2008, 19:358-361; WO 2005/089808; US 5,714,586; US 2013/0101546; and US 2012/0585592), cysteine (*see, e.g.*, US 2007/0258987; WO 2013/055993; WO 2013/055990; WO 2013/053873; WO 2013/053872; WO 2011/130598; US 2013/0101546; and US 7,750,116), selenocysteine (*see, e.g.*, WO 2008/122039; and Hofer *et al.*, *Proc. Natl. Acad. Sci., USA*, 2008, 105:12451-12456), formyl glycine (*see, e.g.*, Carrico *et al.*, *Nat. Chem. Biol.*, 2007, 3:321-322; Agarwal *et al.*, *Proc. Natl. Acad. Sci., USA*, 2013, 110:46-51, and Rabuka *et al.*, *Nat. Protocols*, 2012, 10:1052-1067), non-natural amino acids (*see, e.g.*, WO 2013/068874, and WO 2012/166559), and acidic amino acids (*see, e.g.*, WO 2012/05982). Linkers can be conjugated via glutamine via transglutaminase-based chemo-enzymatic conjugation (*see, e.g.*, Denmler *et al.*, *Bioconjugate Chem.* 2014, 25, 569-578). Linkers can also be conjugated to an antigen-binding protein via attachment to carbohydrates (*see, e.g.*, US 2008/0305497, WO 2014/065661, and Ryan *et al.*, *Food & Agriculture Immunol.*, 2001, 13:127-


130) and disulfide linkers (*see, e.g.*, WO 2013/085925, WO 2010/010324, WO 2011/018611, WO 2014/197854, and Shaunak *et al.*, *Nat. Chem. Biol.*, 2006, 2:312-313).

[0187] In some embodiments, the binding agent is an antibody, and the antibody is bonded to the linker through a lysine residue. In some embodiments, the antibody is bonded to the linker through a cysteine residue.

4. Illustrative embodiments

[0188] In some embodiments,

A is:



wherein:

R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano,

nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5; and

L is:



wherein:

SP is a spacer;


$-\xi^A-$ is one or more bonds to the binding agent;

AA^1 is an amino acid; and

AA^2 is an amino acid.

[0189] In some embodiments,

A is:



wherein:

R¹, independently at each occurrence, is C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, or halo;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5; and

L is:



wherein:

SP is a spacer;

$-\overset{\text{A}}{\underset{\text{Z}}{\text{Z}}}-$ is one or more bonds to the binding agent;


AA¹ is an amino acid; and

AA² is an amino acid.

[0190]

In some embodiments,

A is:



wherein:


R¹, independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano,

nitro, $-\overset{\text{Z}}{\text{Z}}-\text{OR}^{\text{A}}$, $-\overset{\text{Z}}{\text{Z}}-\text{SO}_2\text{R}^{\text{A}}$, $-\overset{\text{Z}}{\text{Z}}-\overset{\text{O}}{\parallel}-\text{R}^{\text{A}}$, or azido,

wherein R^A is alkyl;


m is an integer from 0 to 3;
 p is an integer from 0 to 6; and
 q is an integer from 0 to 5; and

L is:



wherein:

SP is:



wherein:

$-\overset{A}{\xi}-$ is a bond to the binding agent; and

b is an integer from 2 to 8; and


AA¹ is an amino acid; and

AA² is an amino acid.

[0191] In some embodiments,

wherein:


A is:



R¹, independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl;

n is an integer from 0 to 4; and

L is:



wherein:

SP is a spacer;


$\overset{\text{A}}{\text{---}}\text{---}$ is one or more bonds to the binding agent;

AA¹ is an amino acid; and

AA² is an amino acid.

[0192] In some embodiments,
wherein:

A is:



R¹, independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl;

n is an integer from 0 to 4; and

L is:




wherein:

SP is a spacer; and

$\overset{\text{A}}{\text{---}}\text{---}$ is one or more bonds to the binding agent.

[0193] In some embodiments,

A is:




wherein

R¹ is, independently at each occurrence, is C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, halo, or heterocycloalkyl; and

n is 0, 1, or 2; and

L is:



wherein:


SP is a spacer;

$\overset{\text{A}}{\text{---}}$ is the one or more bonds to the binding agent;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.

[0194] In some embodiments, A is:



wherein:


R^1 , independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, ---OR^{A} , $\text{---SO}_2\text{R}^{\text{A}}$, $\text{---C(=O)R}^{\text{A}}$, or azido,

wherein R^{A} is alkyl;

wherein n is an integer from 0 to 4;

L is:



wherein:

SP is a spacer;


$\overset{\text{A}}{\text{---}}$ is the one or more bonds to the binding agent;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.

[0195] In some embodiments,

A is:




wherein

R^1 is, independently at each occurrence, is halo; and

n is 0, 1, or 2; and

L is:




wherein:

$\text{---}\overset{\text{A}}{\text{---}}$ is a bond to the binding agent; and

b is an integer from 2 to 8.

[0196] In some embodiments, A is:




wherein:

R^1 , independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, $\text{---}\xi\text{---OR}^A$, $\text{---}\xi\text{---SO}_2\text{R}^A$, $\text{---}\xi\text{---C(=O)---R}^A$, or azido,

wherein R^A is alkyl;

L is:



wherein:


$\text{---}\overset{\text{A}}{\text{~}}\text{---}$ is a bond to the binding agent;

wherein n is an integer from 0 to 4; and

b is an integer from 2 to 8.

[0197] In some embodiments,

A is:




wherein

R^1 is, independently at each occurrence, is halo; and


n is 0, 1, or 2; and

L is:



wherein $\text{---}\overset{\text{A}}{\text{~}}\text{---}$ is a bond to the binding agent.

[0198] In some embodiments, A is:



wherein:


R¹, independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}-R^A$, or azido,

wherein R^A is alkyl;

wherein n is an integer from 0 to 4;


L is:




wherein $-\xi^A$ is a bond to the binding agent.

[0199] In some embodiments,


A is:



L is




[0200] In some embodiments, A is:



wherein:

R^1 is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, halo, haloalkoxy, or haloalkyl; ξ^1 is the bond to the nitrogen atom of the amino-ester which is directly bonded to the drug molecule; and ξ^2 is the bond to the nitrogen atom which is bonded to the linker. In some embodiments, R^1 is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some embodiments, R^1 is hydrogen.

[0201] In some embodiments, A is:




wherein:

R^1 is, independently at each occurrence, a hydrogen atom, alkyl, alkoxy, aryl, heteroalkyl, halo, haloalkyl, haloalkoxy;


ξ^1 is the bond to the nitrogen atom of the amino-ester which is directly bonded to the drug molecule; and ξ^2 is the bond to the nitrogen atom which is bonded to the linker. In some embodiments, R^1 is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some embodiments, R^1 is hydrogen.

[0202] In some embodiments,

A is:




L is




[0203] In some embodiments,

A is:




L is



[0204] In some embodiments,

BA is an antibody,

A is:




wherein


R^1 is, independently at each occurrence, is halo; and

n is 0, 1, or 2; and

L is:



wherein $\overset{A}{\sim}$ is a bond to the binding agent.




[0205] In some embodiments, including any of the foregoing, A is:

wherein:

R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; and


n is an integer from 0 to 4.

[0206] In some embodiments, including any of the foregoing, A is:




$\overset{1}{\sim}$ and $\overset{2}{\sim}$ are as defined above.

[0207] In some embodiments, including any of the foregoing, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0208] In some embodiments, including any of the foregoing, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0209] In some embodiments, including any of the foregoing, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0210] In some embodiments, including any of the foregoing, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0211] In some embodiments, including any of the foregoing, A is:




$\text{---}\xi^1\text{---}$ and $\text{---}\xi^2\text{---}$ are as defined above.

[0212] In some embodiments, A is:




$\text{---}\overset{1}{\text{wavy}}\text{---}$ and $\text{---}\overset{2}{\text{wavy}}\text{---}$ are as defined above.

[0213] In some embodiments, A is:




$\text{---}\overset{1}{\text{wavy}}\text{---}$ and $\text{---}\overset{2}{\text{wavy}}\text{---}$ are as defined above.

[0214] In some embodiments, including any of the foregoing, A is:




$\text{---}\overset{1}{\text{wavy}}\text{---}$ and $\text{---}\overset{2}{\text{wavy}}\text{---}$ are as defined above.

[0215] In some embodiments, including any of the foregoing, A is:




$\text{---}\overset{1}{\text{wavy}}\text{---}$ and $\text{---}\overset{2}{\text{wavy}}\text{---}$ are as defined above.

[0216] In some embodiments, including any of the foregoing, A is:




$\text{---}\overset{1}{\underset{\sim}{\text{C}}}\text{---}$ and $\text{---}\overset{2}{\underset{\sim}{\text{C}}}\text{---}$ are as defined above.

[0217] In some embodiments, including any of the foregoing, A is:




$\text{---}\overset{1}{\underset{\sim}{\text{C}}}\text{---}$ and $\text{---}\overset{2}{\underset{\sim}{\text{C}}}\text{---}$ are as defined above.

[0218] In some embodiments, including any of the foregoing, A is:




[0219] In some embodiments, including any of the foregoing, A is:



wherein n is 0, 1, 2, or 3.

[0220] In some embodiments,

A is:




wherein

R¹, independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl,

heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido; and

q is an integer from 0 to 5; and

L is:



wherein:

SP is a spacer;


$-\xi^A$ is the one or more bonds to the binding agent;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.

[0221]

In some embodiments, A is:




wherein:

R¹, independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido;

wherein q is an integer from 0 to 5;

L is:



wherein:

SP is a spacer;


$\text{---}\overset{\text{A}}{\underset{\text{A}}{\text{C}}}\text{---}$ is the one or more bonds to the binding agent;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.

[0222] In some embodiments,

A is:




wherein:

R^1 , independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, $\text{---}\text{OR}^{\text{A}}$, $\text{---}\text{SO}_2\text{R}^{\text{A}}$, $\text{---}\overset{\text{O}}{\parallel}\text{---}\text{R}^{\text{A}}$, or azido;

wherein q is an integer from 0 to 5; and

L is:




wherein:

$\text{---}\overset{\text{A}}{\underset{\text{A}}{\text{C}}}\text{---}$ is a bond to the binding agent; and

b is an integer from 2 to 8.

[0223] In some embodiments,

A is:




wherein:

R^1 , independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido; and


q is an integer from 0 to 5; and

L is:



wherein $-\xi-$ is a bond to the binding agent.

[0224] In some embodiments, A is:




wherein:

R^1 , independently at each occurrence, is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano, nitro, $-\xi-OR^A$,

$-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido;

wherein q is an integer from 0 to 5;


L is:




wherein $\overset{A}{\sim}$ is a bond to the binding agent.

[0225] In some embodiments,


A is:



L is




some embodiments, the compound of Formula I is:



wherein A is arylene or heteroarylene, L^1 and L^2 are linkers, BA is a binding agent, k is an integer from 0 to 30, and t is an integer from 0 to 8. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, the subscript, k,

represents the number of linkers, L^1 , bonded to the BA through lysine residues on the BA. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue. In some of these embodiments, the subscript, t , represents the number of linkers, L^2 , bonded to the BA through cysteine residues on the BA. In some embodiments, when the linker, L^2 , is a monodentate linker, t is an integer from 0 to 8. In some embodiments, when the linker, L^2 , is a bidentate linker, t is an integer from 0 to 4. In some of these examples, the sum of $k + t$ is equal to 1-8.


[0226] In some embodiments, the compound of Formula I is:



wherein A is arylene or heteroarylene, L^1 and L^2 are linkers, and BA is a binding agent. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue.

[0227] In some embodiments,

A is:



wherein:

R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido,


wherein R^A is alkyl;

m is an integer from 0 to 3;


p is an integer from 0 to 6; and

q is an integer from 0 to 5;

L is




some embodiments, the compound of Formula I is:




wherein A is arylene or heteroarylene, L^1 and L^2 are linkers, BA is a binding agent, k is an integer from 0 to 30, and t is an integer from 0 to 8. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, the subscript, k , represents the number of linkers, L^1 , bonded to the BA through lysine residues on the BA . In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue. In some of these embodiments, the subscript, t , represents the number of linkers, L^2 , bonded to the BA through cysteine residues on the BA . In some embodiments, when the linker, L^2 , is a monodentate linker, t is an integer from 0 to 8. In some embodiments, when the linker, L^2 , is a bidentate linker, t is an integer from 0 to 4. In some of these examples, the sum of $k + t$ is equal to 1-8.

[0228] In some embodiments, the compound of Formula I is:



wherein A is arylene or heteroarylene, L^1 and L^2 are linkers, and BA is a binding agent. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue.


[0229] In some embodiments, the compound of Formula I is:



wherein A is arylene or heteroarylene, L^1 and L^2 are linkers, and BA is a binding agent. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue.

[0230] In some embodiments,

A is:



wherein:

R¹, independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano, nitro, $\text{-}\overset{\xi}{\text{C}}\text{-OR}^{\text{A}}$,

$\text{-}\overset{\xi}{\text{C}}\text{-SO}_2\text{R}^{\text{A}}$, $\text{-}\overset{\xi}{\text{C}}\overset{\text{O}}{\parallel}\text{-R}^{\text{A}}$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;


p is an integer from 0 to 6; and

q is an integer from 0 to 5;

L is




the compound of Formula I is:




wherein A is arylene or heteroarylene, L^1 , L^2 , and L^3 are linkers, BA is a binding agent, k is an integer from 0 to 30, t is an integer from 0 to 8, and g is an integer from 0 to 4. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, the subscript, k, represents the number of linkers, L^1 , bonded to the BA through lysine residues on the BA. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue. In some of these embodiments, the subscript, t, represents the number of linkers, L^2 , bonded to the BA through cysteine residues on the BA. In some embodiments, when the linker, L^2 , is a monodentate linker, t is an integer from 0 to 8. In some embodiments, when the linker, L^2 , is a bidentate linker, t is an integer from 0 to 4. In some of these examples, the sum of $k + t$ is equal to 1-8. In some of these embodiments, L^3 is a linker which binds to the BA through a glutamine residue. In some of these embodiments, the subscript, g, represents the number of linkers, L^3 , bonded to the BA through glutamine residues on the BA.

[0231] In some embodiments, the compound of Formula I is:



wherein A is arylene or heteroarylene, L^1 , L^2 , and L^3 are linkers, and BA is a binding agent. In some of these embodiments, L^1 is a linker which binds to the BA through a lysine residue. In some of these embodiments, L^2 is a linker which binds to the BA through a cysteine residue. In some of these embodiments, L^3 is a linker which binds to the BA through a glutamine residue.

[0232] In some embodiments, A is:



wherein:

R^1 is, independently at each occurrence, halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, haloalkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl,

heterocycloalkyl, cyano, nitro, $-\overset{\xi}{\underset{\xi}{\text{C}}}-OR^A$, $-\overset{\xi}{\underset{\xi}{\text{C}}}-SO_2R^A$, $-\overset{\xi}{\underset{\xi}{\text{C}}}-\overset{\text{O}}{\parallel}R^A$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5.

[0233] In some embodiments, the linker is:



wherein:

SP is a spacer;


$-\overset{\xi}{\underset{\xi}{\text{C}}}-$ is one or more bonds to the binding agent;

AA^1 is an amino acid; and


AA^2 is an amino acid.

[0234] The spacer is a divalent moiety that connects the AA^1 - AA^2 moiety to the binding agent (BA). Suitable spacers include, but are not limited to, those comprising alkylene or polyethylene glycol. The ends of the spacers, *i.e.*, the portion of the spacer directly bonded to the binding agent or AA^1 , can be moieties derived from reactive moieties that are used for purposes of coupling the antibody or AA^1 to the spacer during the chemical synthesis of the conjugate.


[0235] In some embodiments, the spacer comprises an alkylene. In some embodiments, the spacer comprises a C_{5-7} alkylene. In some embodiments, the spacer is:




wherein:

 is a bond to the binding agent; and
 b is an integer from 2 to 8.


[0236] In some embodiments, the spacer is:



wherein:

 is a bond to the binding agent.


[0237] In some embodiments, the spacer is:




wherein:

R^N is a hydrogen atom or alkyl;


R^M is alkyl;

the two bonds represented by  are bonds to cysteines of a binding agent; and
 b is an integer from 2 to 8.

[0238] In some embodiments, the spacer is:




wherein:

 is a bond to the binding agent; and

b is an integer from 2 to 8.


[0239] In some embodiments, the spacer is:



wherein:

is a bond to the binding agent.

[0240] In some embodiments, the spacer is:




wherein:


is a bond to the binding agent; and

g is an integer from 2 to 20. In some embodiments, g is 2-8. In some embodiments, g is 2, 4, 6, or 8.


[0241] In some embodiments, the spacer is:




[0242] In some embodiments, the spacer is:




[0243] In some embodiments, the spacer is:




[0244] In some embodiments, the spacer is:




[0245] In some embodiments, the spacer is:




[0246] In some embodiments, the spacer is:




[0247] In some embodiments, the spacer is:



[0248] In some embodiments, the spacer is:



[0249] In some embodiments, the spacer is:



[0250] wherein $\overset{A}{\sim}$ is a bond to the binding agent;


X is N or O; R^N and R^M are each, independently, hydrogen or alkyl; and b is an integer from 1 to 8.

[0251] In some embodiments, AA¹-AA² is: valine-citrulline, citrulline-valine, lysine-phenylalanine, phenylalanine-lysine, valine-asparagine, asparagine-valine, asparagine-threonine, threonine-asparagine, serine-asparagine, asparagine-serine, phenylalanine-asparagine, asparagine-phenylalanine, leucine-asparagine, asparagine-leucine, isoleucine-asparagine, asparagine-isoleucine, glycine-asparagine, asparagine-glycine, glutamic acid- asparagine,


asparagine-glutamic acid, citrulline-asparagine, asparagine-citrulline, alanine-asparagine, asparagine-alanine, valine-alanine, alanine-valine, valine-glycine, or glycine-valine.

[0252] In some embodiments, AA¹-AA² is: valine-citrulline or citrulline-valine. In some embodiments, AA¹-AA² is: valine-citrulline.

[0253] In some embodiments, the compound of Formula I is:



or



wherein X is N or O,





R^N and R^M are each, independently, hydrogen or aryl,





b is an integer from 1 to 8,


A is aryl or heteroaryl, and




t is an integer from 1-8.




[0254] In some embodiments, the compound of Formula I is:



































































































































































































wherein:

Ab is an antibody or an antigen binding fragment thereof;

—S— is a bond to a cysteine of the antibody or the antigen binding fragment thereof;

—N— is a bond to a lysine of the antibody or the antigen binding fragment thereof;


k is an integer from 1 to 30; and

t is an integer from 1 to 8; —Gln— is a bond to a glutamine of the antibody or the antigen binding fragment thereof. In some examples, k is an integer from 1 to 8. In some examples, t is an integer from 1 to 4. In some examples, when —S— is a bond to a cysteine of the antibody, up to 8 conjugates set forth herein may be bonded to the antibody. In some examples, when —N— is a bond to a lysine of the antibody, up to 30 conjugates set forth herein may be bonded to the antibody.

[0255] In some embodiment, k is an integer from 1 to 30. In some embodiment, k is an integer from 1 to 8. In some embodiment, k is an integer from 1 to 6. In some embodiments, k is an integer from 1 to 4. In some embodiments, k is an integer from 1 to 3. In some embodiments, the drug-antibody ratio (DAR) of the conjugate is from 1 to 30.

C. COMPOUNDS

[0256] Provided herein are compounds of Formula (II):



(II)

or a pharmaceutically acceptable salt thereof,
wherein A is arylene or heteroarylene.


[0257] In certain embodiments, these compounds represent the payload portion of the conjugates described herein and are released, *e.g.*, by enzyme proteolysis, following internalization of the conjugate into a cell. The methods provided herein include methods of treating a proliferative disease, *e.g.*, cancer, comprising administering to a patient a therapeutically effective amount of a conjugate, *e.g.*, antibody-drug conjugate that releases a compound of Formula (II) following internalization of said conjugate into a cell in said patient.

[0258] In some embodiments, these compounds represent the metabolic product of the conjugates described herein, *e.g.*, enzyme proteolysis product. In some embodiments, these compounds represent the catabolic product of the conjugates described herein. In some embodiments, these compounds represent the cellular product of the conjugates described herein.

[0259] In some embodiments, A is a divalent radical of benzene, of pyridine, of naphthalene, or of quinoline, which are optionally substituted.

[0260] In some embodiments, A is arylene.

[0261] In some embodiments, A is:



wherein:

R¹, independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroaryl, heteroalkyl, heterocycloalkyl, cyano,

nitro, amino, $\text{-}\xi\text{-OR}^{\text{A}}$, $\text{-}\xi\text{-SO}_2\text{R}^{\text{A}}$, $\text{-}\xi\text{-C(=O)R}^{\text{A}}$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and


q is an integer from 0 to 5.

[0262] In some embodiments, the compound of Formula (II) is a compound having the Formula:



wherein R^1 and n are as defined herein.


[0263] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIA):



(IIA)

wherein R^1 and n are as defined herein. In some embodiments, R^1 is alkyl, alkoxy, haloalkyl, or halo. In some embodiments, R^1 is methyl, trifluoromethyl, methoxy, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo.


[0264] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB):



(IIB)

wherein R¹ and q are as defined herein.


[0265] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB2):



(IIB2)

wherein R¹ and q are as defined herein.


[0266] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB3):



(IIB3)


wherein R¹ and q are as defined herein.

[0267] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIC):




wherein R¹ and q are as defined herein.

[0268] In some embodiments, the compound of Formula (II) is a compound of the Formula (IID):




wherein R¹ and q are as defined herein.

[0269] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIE):




wherein R¹ and q are as defined herein.

[0270] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIF):



wherein R^1 and q are as defined herein.

[0271] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIG):




wherein R^1 and q are as defined herein.

[0272] In some embodiments, R^1 is, independently, alkyl or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} haloalkyl, or halo. In some embodiments, R^1 is, independently, C_{1-6} haloalkyl or halo. In some embodiments, R^1 is, independently, halo. In some embodiments, R^1 is, independently, fluoro, chloro, bromo, iodo, or trifluoromethyl. In some embodiments, n , m , p , or q is 0, 1 or 2. In some embodiments, n , m , p , or q is 0 or 1. In some embodiments, n , m , p , or q is 0.

[0273] In some embodiments, R^1 is, independently, alkyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl or C_{1-6} alkoxy. In some embodiments, R^1 is, independently, alkoxy. In some embodiments, R^1 is,

independently, methoxy, ethoxy, propoxy. In some embodiments, n, m, p, or q is 0, 1 or 2. In some embodiments, n, m, p, or q is 0 or 1. In some embodiments, n, m, p, or q is 0.


[0274] In some embodiments, the compound of Formula (II) is a compound of Formula (IIA):



wherein:

R^1 is, independently at each occurrence, halo, methyl, methoxy, or trifluoromethyl; and n is 0, 1, or 2. In some embodiments, R^1 is alkyl, alkoxy, haloalkyl, or halo. In some embodiments, R^1 is methyl, trifluoromethyl, methoxy, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some examples, R^1 is heterocycloalkyl. In some examples, R^1 is pyrrolidinyl. In some examples, R^1 is morpholinyl.


[0275] A compound of Formula IIA1-3:



or a pharmaceutically acceptable salt thereof,


wherein:

A is:




R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; n is an integer from 0 to 4.


[0276] In some embodiments, the compound is selected from a compound of the Formula (IIA1), a compound of the Formula (IIA2), and a compound of the Formula (IIA3):



(IIA1)



(IIA2)



(IIA3).

[0277] In some embodiments, the compound of any of claims 31-32, wherein n is 0, 1 or 2.

[0278] In some embodiments, the compound of any of claims 31-32, wherein R^1 is, independently at each occurrence, selected from C_{1-6} alkyl, C_{1-6} alkoxy, halo, C_{1-6} haloalkyl, and heterocycloalkyl.

[0279] In some embodiments, R^1 is, independently at each occurrence, selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, pyrrolidinyl, and morpholinyl.

[0280] In some embodiments, R^1 is methyl.

[0281] In some embodiments, R^1 , independently at each occurrence, is selected from fluoro, chloro, and bromo.


[0282] In some embodiments, R^1 is chloro.

[0283] In some embodiments, R^1 is trifluoromethyl.

[0284] In some embodiments, R^1 is methoxy.

[0285] In some embodiments, R^1 is, independently at each occurrence, selected from methyl, morpholinyl, and pyrrolidinyl.

[0286] In some embodiments, the compound of Formula (II) is a compound of Formula (IIB):












(IIB)









wherein:









R^1 is, independently at each occurrence, halo or trifluoromethyl; and
 q is 0, 1, or 2.

[0287] In some embodiments, the compound of Formula (II) is:



















and




[0288] In some embodiments, the compound of Formula (II) is:







or




[0289] In certain embodiments, these compounds represent the payload portion of the conjugates described herein and are released, *e.g.*, by enzyme proteolysis, following internalization of the conjugate into a cell. The methods provided herein include methods of treating a proliferative disease, *e.g.*, cancer, comprising administering to a patient a therapeutically effective amount of a conjugate, *e.g.*, antibody-drug conjugate that releases a compound of Formula (II) following internalization of said conjugate into a cell in said patient.

[0290] In some embodiments, these compounds represent the metabolic product of the conjugates described herein, *e.g.*, enzyme proteolysis product.

[0291] In some embodiments, A is a divalent radical of benzene, of pyridine, of naphthalene, or of quinolone, which are optionally substituted.

[0292] In some embodiments, A is arylene.

[0293] In some embodiments, A is:



wherein:

R^1 is, independently at each occurrence, alkyl, alkenyl, alkynyl, aryl, alkaryl, arylalkyl,

halo, haloalkoxy, heteroaryl, heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5.

[0294] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIA):




(IIA)

wherein R^1 is, independently at each occurrence, methyl, methoxy, halo or trifluoromethyl; and n is 0, 1, or 2. In some embodiments, R^1 is methyl, trifluoromethyl, methoxy, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some


embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo.

[0295] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB):




wherein R^1 is, independently at each occurrence, methoxy, methyl, halo or trifluoromethyl; and q is 0, 1, or 2.

[0296] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB2):



wherein R^1 is, independently at each occurrence, methoxy, halo or trifluoromethyl; and q is 0, 1, or 2.

[0297] In some embodiments, the compound of Formula (II) is a compound of the Formula (IIB3):



(IIB3)

[0298] wherein R¹ is, independently at each occurrence, methoxy, halo or trifluoromethyl; and q is 0, 1, or 2. In some embodiments, R¹ is, independently, alkyl or halo. In some embodiments, R¹ is, independently, C₁₋₆ alkyl, C₁₋₆ haloalkyl, or halo. In some embodiments, R¹ is, independently, C₁₋₆ haloalkyl or halo. In some embodiments, R¹ is, independently, halo. In some embodiments, R¹ is, independently, fluoro, chloro, bromo, iodo, or trifluoromethyl. In some embodiments, n, m, p, or q is 0, 1 or 2. In some embodiments, n, m, p, or q is 0 or 1. In some embodiments, n, m, p, or q is 0.


[0299] In some embodiments, R¹ is, independently, alkyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy. In some embodiments, R¹ is, independently, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, or halo. In some embodiments, R¹ is, independently, C₁₋₆ alkyl or C₁₋₆ alkoxy. In some embodiments, R¹ is, independently, alkoxy. In some embodiments, R¹ is, independently, methoxy, ethoxy, propoxy. In some embodiments, n, m, p, or q is 0, 1 or 2. In some embodiments, n, m, p, or q is 0 or 1. In some embodiments, n, m, p, or q is 0.

[0300] In some embodiments, provided herein are compounds which include any of the compounds described herein covalently bonded to a linker and/or binding agent as described herein. In some examples, provided herein are any of the maytansinoid compounds described herein covalently bonded to a linker and/or binding agent as described herein.

D. PREPARATION OF COMPOUNDS

[0301] Compounds of Formula I can be synthesized by coupling compounds of Formula P1 with a binding agent, *e.g.*, antibody under standard conjugation conditions (*see, e.g.*, Doronina *et al.*, *Nature Biotechnology* 2003, 21, 7, 778, which is incorporated herein by reference). When the binding agent is an antibody, the antibody can be coupled to a compound


of Formula P1 via one or more cysteine or lysine residues of the antibody. Compounds of Formula P1 can be coupled to cysteine residues, for example, by subjecting the antibody to a reducing agent, *e.g.*, dithioerithol, to cleave the disulfide bonds of the antibody, purifying the reduced antibody, *e.g.*, by gel filtration, and subsequently reacting the antibody with a compound of formula P1 containing a reactive moiety, *e.g.*, a maleimido group. Suitable solvents include, but are not limited to water, DMA, DMF, and DMSO. Compounds of formula P1 containing a reactive moiety, *e.g.*, activated ester or acid halide group, can be coupled to lysine residues. Suitable solvents include, but are not limited to water, DMA, DMF, and DMSO. The compounds of Formula I can be purified using known protein techniques, including, for example, size exclusion chromatography, dialysis, and ultrafiltration/diafiltration.



wherein RL is a reactive linker, A is arylene or heteroarylene, L is a linker, and BA is a binding agent.


[0302] In some examples, set forth herein are methods of making a compound of Formula (I) wherein the method includes contacting a compound of Formula P1 with a binding agent. In some of these methods, the contacting is under standard antibody conjugation conditions. In some of these examples, the binding agent is an antibody. In some of these examples, the binding agent is a fragment of an antibody. In some examples herein, the methods include contacting a compound of Formula P1 with one or more cysteine or lysine residues of an antibody. In some examples herein, the methods include contacting a compound of Formula P1 with one or more cysteine or lysine residues of an antibody in a solvent. In some embodiments, the solvent is a single compound. In some embodiments, the solvent is a mixture of two or more compounds. In some examples, the methods include contacting a compound of Formula P1 with one or more cysteine or lysine residues of an antibody in a solvent selected from water, DMA, DMF, or DMSO.

[0303] In some examples, set forth herein are methods of making a compound of Formula P1, wherein the methods include reacting a compound of Formula (II) with a reactive linker (RL), wherein RL is a reactive linker as set forth and described herein.



[0304] In some examples, set forth herein are methods of making a compound of Formula P1, wherein the methods include reacting a compound of Formula (II) with a reactive linker set forth and described herein. In some examples, the methods include reacting a compound of Formula P1 with a RL in the presence of a solvent. In some examples, the solvent is DCM. In some other examples, the solvent is methanol. In some other examples, the solvent is anhydrous methanol. In certain other examples, the solvent is a combination of DCM and methanol. In some other examples, the solvent is a combination of DCM and anhydrous methanol.

[0305] In some examples, the RL is selected from a compound of Formula PP7



, wherein SP^R is a reactive spacer; R^{AA1} is an amino acid side chain; R^{AA2} is an amino acid side chain. In some examples, R^{AA1} and R^{AA2} are independently natural or non-natural amino acid side chains. In some examples of the methods of making a compound of Formula P1, the methods include reacting a compound of Formula (II) with a compound of Formula PP7 under amide synthesis conditions. In some of these examples, the compound of Formula PP7 is converted into an activated ester before reacting with a compound of Formula (II).

[0306] In some examples of the methods of making a compound of Formula P1, the methods include contacting a compound of Formula (II) with a RL. In some these examples, the methods include contacting a compound of Formula (II) with a RL under amide synthesis conditions. In some examples, the RL includes a carboxylic acid group at one terminal end of


the RL compound. As shown in Formula (II), a compound of Formula (II) includes at least one amino group at one terminal end of the compound. In some examples, the methods include converting a carboxylic acid group of the RL compound into an activated ester. In some examples, the methods further include reacting this activated ester with an amino group on a compound of Formula (II). In some examples herein, the methods of converting the carboxylic acid into an activated ester include contacting the carboxylic acid with a reagent. In some examples herein, the methods of converting the carboxylic acid of the RL into an activated ester include contacting the RL with a reagent selected from dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIC), (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), (7-azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP), bromotripyrrolidinophosphonium hexafluorophosphate (PyBrOP), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU), 1-[Bis(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate (HATU), 2-Ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC), 2-Chloro-1,3-dimethylimidazolidinium hexafluorophosphate (CIP), 2-chloro-4,6-dimethoxy-1,3,5-triazine (CDMT), or carbonyldiimidazole (CDI). In some examples, the methods include, converting a carboxylic acid group of the RL compound into an acyl chloride. In some examples, the methods include converting a carboxylic acid group of the RL compound into an acyl chloride, for example by contacting the carboxylic acid of the RL with EEDQ. In some examples herein, the carboxylic acid of the RL is converted into a mixed anhydride, for example by contacting the carboxylic acid of the RL with EEDQ. In some examples herein, the methods of converting the carboxylic acid of the RL into a mixed anhydride include contacting a carboxylic acid of the RL with EEDQ. In some examples herein, the methods of converting a carboxylic acid of the RL into a mixed anhydride include contacting a carboxylic acid of the RL with EEDQ in a solvent selected from DCM.

[0307] In some examples, set forth herein are methods of making a compound of Formula P1, wherein the methods include reacting a compound of Formula (II) with a RL wherein the RL is 6-maleimidyl-caproamidyl-L-valine-L-citrulline. In some examples, the 6-

maleimidyl-caproamidyl-L-valine-L-citrulline is activated to form an activated ester before reacting with a compound of Formula (II). In some examples, the methods herein include providing 6-maleimidyl-caproamidyl-L-valine-L-citrulline by reacting BOC-protected L-valine-L-citrulline with 2-(2-amino-3-methyl-butyrylamino)-5-ureido-pentanoic acid and 6-(2,5-Dioxo-2,5-dihydro-pyrrol-1-yl)-hexanoic acid 2,5-dioxo-pyrrolidin-1-yl ester. In some methods herein, the methods include providing 6-maleimidyl-caproamidyl-L-valine-L-citrulline by reacting BOC-protected L-valine-L-citrulline with 2-(2-amino-3-methyl-butyrylamino)-5-ureido-pentanoic acid and 6-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-hexanoic acid 2,5-dioxo-pyrrolidin-1-yl ester in DMF. In some methods herein, the methods include providing 6-maleimidyl-caproamidyl-L-valine-L-citrulline by reacting BOC-protected L-valine-L-citrulline with 2-(2-amino-3-methyl-butyrylamino)-5-ureido-pentanoic acid and 6-(2,5-dioxo-2,5-dihydro-pyrrol-1-yl)-hexanoic acid 2,5-dioxo-pyrrolidin-1-yl ester in DIEA.


[0308] In some examples of the methods of making a compound of Formula P1, the methods include reacting a compound of Formula (II) with adipic anhydride. In these methods, the adipic anhydride is an activated ester. In some of these examples, the methods include reacting a compound of Formula (II) with adipic anhydride and triethylamine. In some of examples, the methods include reacting a compound of Formula (II) with adipic anhydride and triethylamine in tetrahydrofuran. In some of these examples, the methods include reacting a compound of Formula (II) with adipic anhydride and triethylamine in dimethylformamide. In some of these examples, the methods include reacting a compound of Formula (II) with adipic anhydride and triethylamine in tetrahydrofuran and dimethylformamide (DMF). In some examples, these methods further include reacting N-hydroxysuccinimide with the products of the aforementioned methods. In some other of these examples, these methods further include reacting N-hydroxysuccinimide and EDC hydrochloride with the products of the aforementioned methods.

[0309] In some examples, set forth herein are methods of making a compound of Formula (II). In some examples, these methods include reacting a compound of Formula PP5 with a suitable reducing agent.



[0310] In some of these methods of making a compound of Formula (II), the suitable reducing agent includes a metal, a metal foil, a metal powder, dust of a metal, a metal amalgam, or metal filings. In certain embodiments, the metal is selected from zinc, iron, aluminum, palladium, or Raney nickel. In some of these methods, the suitable reducing agent is Zn foil, Zn powder, Zn dust, Zn amalgam, or Zn filings. In some of these methods, the suitable reducing agent is Zn foil. In some of these methods, the suitable reducing agent is Zn powder. In some of these methods, the suitable reducing agent is Zn dust. In some of these methods, the suitable reducing agent is Zn amalgam. In some of these methods, the suitable reducing agent is Zn filings. In some methods, the methods further include reducing the compound of formula PP5 in the presence of a solvent. In some examples, the solvent is acetic acid. In some examples, the solvent is tetrahydrofuran. In some examples, the solvent is a combination of acetic acid and tetrahydrofuran. In some examples, the solvent is acetonitrile. In some examples, the solvent is acetonitrile and water.

[0311] In some examples, set forth herein are methods of making a compound of Formula PP5. In some examples, these methods include reacting a compound of Formula P2 with a compound of Formula PP6.



In some of these examples, A is arylene or heteroarylene, optionally substituted as set forth herein.


[0312] In some examples of the methods of making a compound of Formula PP5, the methods include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions. In some examples, the compound of PP6 is a para-nitro-phenyl-isocyanate. In some examples, the compound of PP6 is a meta-nitro-phenyl-isocyanate. In some examples, the Lewis acid conditions include using a Lewis acid during the reaction of a compound of Formula P2 with a compound of Formula PP6. In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of a Lewis Acid. Suitable Lewis acids include, but are not limited to AlBr_3 , AlCl_3 , BCl_3 , boron trichloride methyl sulfide, BF_3 , boron trifluoride methyl etherate, boron trifluoride methyl sulfide, boron trifluoride tetrahydrofuran, dicyclohexylboron trifluoromethanesulfonate, iron (III) bromide, iron (III) chloride, tin (IV) chloride, titanium (IV) chloride, titanium (IV) isopropoxide, $\text{Cu}(\text{OTf})_2$, CuCl_2 , CuBr_2 , zinc chloride, alkylaluminum halides ($\text{R}_n\text{AlX}_{3-n}$, wherein R is hydrocarbyl and n is from 0 to 3), $\text{Zn}(\text{OTf})_2$, $\text{Yb}(\text{OTf})_3$, $\text{Sc}(\text{OTf})_3$, MgBr_2 , NiCl_2 , $\text{Sn}(\text{OTf})_2$, $\text{Ni}(\text{OTf})_2$, or $\text{Mg}(\text{OTf})_2$. In some examples, the Lewis acid is ZnCl_2 . In some examples, the Lewis acid conditions include using a Lewis acid and an organic solvent during the reaction of a compound of Formula P2 with a compound of Formula PP6. In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of a Lewis Acid and an organic solvent. In some examples, the organic solvent is dichloromethane (DCM). In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of ZnCl_2 and DCM. In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of ZnCl_2 , DCM, and diethyl ether. In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of DCM and also a diethyl ether solution of ZnCl_2 . In some examples, a compound of Formula P2 is reacted with a compound of Formula PP6 in the presence of DCM wherein ZnCl_2 is added during this reaction as a diethyl ether solution of ZnCl_2 . In some examples, the methods herein include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions for at least 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, or 24 hours. In some examples, the methods herein include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions for 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, or 24 hours. In some examples, the methods herein include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions for 16 hours. In some examples, the methods herein include reacting a compound of

Formula P2 with a compound of Formula PP6 under Lewis acid conditions for 18 hours. In some examples, the methods herein include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions for 20 hours. In some examples, the methods herein include reacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions for 24 hours.

[0313] In some examples, after reacting a compound of Formula P2 with a compound of Formula PP6, the product is diluted with water and extracted once, twice, or three times with an organic solvent. In some examples the organic solvent used for this extraction is selected from ethyl acetate (EtOAc).


[0314] In some examples, after reacting a compound of Formula P2 with a compound of Formula PP6, the product is concentrated to dryness. In some examples, after reacting a compound of Formula P2 with a compound of Formula PP6, the product is concentrated to dryness and purified by chromatography

[0315] In some embodiments, the compound of formula P1 includes A, wherein A is:




wherein n is 0 or 1; and R¹ is alkyl, alkoxy, halo, haloalkoxy, or haloalkyl.

[0316] In some embodiments, the compound of formula P1 includes A, wherein A is:




wherein n is 0 or 1; and R¹ is C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkoxy, or C₁₋₆ haloalkyl.

[0317] In some embodiments, the compound of formula P1 includes A, wherein A is:




wherein n is 0 or 1; R¹ is C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkoxy, or C₁₋₆ haloalkyl; and RL



to the binding agent. In some embodiments, R¹ is alkyl, alkoxy, haloalkyl, or halo. In some embodiments, R¹ is methyl, trifluoromethyl, methoxy, fluoro, chloro, or bromo. In some embodiments, R¹ is methyl. In some embodiments, R¹ is methoxy. In some embodiments, R¹ is trifluoromethyl. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo.


[0318] The reactive linker is a moiety comprising a portion in its structure that is capable of reacting with the binding agent (*e.g.*, reacting with an antibody at its cysteine or lysine residues) to form the compound of Formula I. Following conjugation to the binding agent, the reactive linker becomes the linker (L) moiety of the compound of Formula I. Illustrative reactive linkers include, but are not limited to, those that comprise haloacetyl, isothiocyanate, terminal primary amine or maleimide portions that are capable of reacting with the binding agent.

Reactive portions also include moieties having the following structure:



wherein X is –O– or –NH– and LG is a leaving group, *e.g.*, Br.

[0319] In some embodiments, the reactive linker is:



wherein:

SP^R is a reactive spacer;


AA¹ is an amino acid; and

AA² is an amino acid.

[0320] The reactive spacer is a moiety that contains the above-described reactive linker portion that is capable of reacting with the binding agent and connects this portion to AA¹.


Suitable spacers include, but are not limited to, those comprising alkylene or polyethylene glycol connecting the AA¹ to the portion capable of reacting with binding agent (*e.g.*, haloacetyl, isothiocyanate, or maleimide).

[0321] In some embodiments, the reactive spacer is:




wherein b is an integer from 2 to 8.


[0322] In some embodiments, the reactive spacer is:



[0323] In some embodiments, the spacer is




[0324] In some embodiments, the spacer is




, wherein g is an integer from 1 to 24.

[0325] In some embodiments, the reactive spacer is:



wherein b is an integer from 2 to 8 and g is an integer from 2 to 20.


[0326] In some embodiments, the reactive spacer is:



[0327] In some embodiments, AA¹-AA² is: valine-citrulline, citrulline-valine, lysine-phenylalanine, phenylalanine-lysine, valine-asparagine, asparagine-valine, threonine-asparagine, serine-asparagine, asparagine-serine, phenylalanine-asparagine, asparagine-phenylalanine, leucine-asparagine, asparagine-leucine, isoleucine-asparagine, asparagine-isoleucine, glycine-asparagine, asparagine-glycine, glutamic acid-asparagine, asparagine-glutamic acid, citrulline-asparagine, asparagine-citrulline, alanine-asparagine, asparagine-alanine, valine-alanine, alanine-valine, valine-glycine, or glycine-valine.

[0328] In some embodiments, AA¹-AA² is: valine-citrulline or citrulline-valine. In some embodiments, AA¹-AA² is: valine-citrulline.

[0329] In some embodiments, the reactive linker is:




wherein:

SP^R is a reactive spacer;

R^{AA1} is an amino acid side chain; and

R^{AA2} is an amino acid side chain.


[0330] In some embodiments, the reactive linker is:



wherein:


SP is a reactive spacer.

[0331] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0332] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0333] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0334] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0335] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8, R^N is a hydrogen atom or alkyl, and R^M is alkyl.

[0336] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0337] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0338] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8; R^N is a hydrogen atom or alkyl; and R^M is alkyl.

[0339] In some embodiments, the reactive linker is:




wherein b is an integer from 2 to 8.

[0340] In some embodiments, the reactive linker is:




wherein g is an integer from 2 to 8.


[0341] In some embodiments, the reactive linker is:




[0342] In some embodiments, the reactive linker is:




[0343] In some embodiments, the reactive linker is:




[0344] In some embodiments, the reactive linker is:




[0345] In some embodiments, the reactive linker is:




[0346] In some embodiments, the reactive linker is:




[0347] In some embodiments, the reactive linker is:




[0348] In some embodiments, the reactive linker is:




[0349] In some embodiments, the reactive linker is:




[0350] In some embodiments, the reactive linker is:




[0351] In some embodiments, the reactive linker is:



[0352] In some embodiments, the reactive linker is:




[0353] In some embodiments, the compound of Formula P1 is a compound of Formula P1A:



wherein:

A is:



wherein:

R¹ is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl,

cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, or azido, wherein R^A is alkyl;

n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and


q is an integer from 0 to 5;

SP^R is a reactive spacer;

AA^1 is an amino acid; and


AA^2 is an amino acid.

[0354] In some embodiments, the compound of Formula P1A is a compound which includes A wherein A is:




wherein n is 0 or 1; and R^1 is alkoxy, alkyl, halo, haloalkoxy, or haloalkyl.

[0355] In some embodiments, the compound of Formula P1A is a compound which includes A wherein A is:




wherein n is 0 or 1; and R^1 is C_{1-6} alkoxy, C_{1-6} alkyl, halo, C_{1-6} haloalkoxy, or C_{1-6} haloalkyl. In some embodiments, R^1 is alkyl, alkoxy, haloalkyl, or halo. In some embodiments, R^1 is methyl, trifluoromethyl, methoxy, fluoro, chloro, or bromo. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo.

[0356] In some embodiments, the compound of Formula P1A is a compound which includes A wherein A is:




wherein q is an integer from 0 to 5; and R^1 is C_{1-6} alkoxy, C_{1-6} alkyl, halo, C_{1-6} haloalkoxy, or C_{1-6} haloalkyl.

[0357] In some embodiments, the compound of Formula P1A is a compound which includes A wherein A is:




wherein q is an integer from 0 to 5; and R^1 is C_{1-6} alkoxy, C_{1-6} alkyl, halo, C_{1-6} haloalkoxy, or C_{1-6} haloalkyl.

[0358] In some embodiments, the compound of Formula P1 is a compound of Formula P1A1-3:



wherein:


A is:



R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl;


n is an integer from 0 to 4; and RL is a reactive linker.

[0359] In some embodiments, the compound of Formula P1A1-3 is a compound of Formula P2A1-3:




wherein:


A is selected from:




SP^R is a reactive spacer.

[0360] In some embodiments, the compound of Formula P1A1-3 is selected from a compound of Formula P1H1, a compound of Formula P1I1, a compound of Formula P1V1, a compound of Formula P1W1, a compound of Formula P1K1, a compound of Formula P1TG1, a compound of Formula P1ZZ1, and a compound of Formula P1ZZ2, wherein b is an integer from 2 to 8:







(PII1)





(PIV1)




(P1W1)



(PIK1)




, and



[0361]
P1B:


In some embodiments, the compound of Formula P1 is a compound of Formula



P1B

wherein

A is:



wherein:

R^1 is halo, haloalkoxy, haloalkyl, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, or azido, wherein R^A is alkyl;

n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and


q is an integer from 0 to 5; and

SP^R is a reactive spacer.

[0362]

In some embodiments, the compound of Formula P1 is a compound of Formula

P1C:



P1C

wherein:

SP^R is a reactive spacer;


AA^1 is an amino acid;

AA^2 is an amino acid;

R^1 is, independently at each occurrence, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and

n is 0, 1, or 2.

[0363] In some embodiments, the compound of Formula P1 is a compound of Formula P1D:



P1D

wherein:

SP^R is a reactive spacer;


AA^1 is an amino acid;

AA^2 is an amino acid;

R^1 is, independently at each occurrence, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and n is 0, 1, or 2. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.


[0364] In some embodiments, the compound of Formula P1 is a compound of Formula P1D, wherein R^1 is alkoxy, alkyl, halo, haloalkoxy, or haloalkyl. In some embodiments, and R^1 is C_{1-6} alkoxy, C_{1-6} alkyl, halo, C_{1-6} haloalkoxy, or C_{1-6} haloalkyl. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 2.

[0365] In some embodiments, the compound of Formula P1 is a compound of Formula P1D, wherein and R^1 is C_{1-6} alkoxy, C_{1-6} alkyl, halo, C_{1-6} haloalkoxy, or C_{1-6} haloalkyl; and SP^R -



embodiments b is 4. In some embodiments b is 5. In some embodiments b is 6. In some embodiments b is 7. In some embodiments b is 8.

[0366] In some embodiments, the compound of Formula P1 is a compound of Formula P1E:



wherein:

SP^R is a reactive spacer;


R^{AA1} is an amino acid side chain;

R^{AA2} is an amino acid side chain;

R¹ is, independently at each occurrence, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and

n is 0, 1, or 2.

[0367] In some embodiments, the compound of Formula P1 is a compound of Formula P1F:



P1F

wherein:

SP^R is a reactive spacer;


R^1 is, independently at each occurrence, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and

n is 0, 1, or 2.

[0368]

In some embodiments, the compound of Formula P1 is a compound of Formula

P1G:



P1G

wherein:


SP^R is a reactive spacer; and

R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.

[0369]

In some embodiments, the compound of Formula P1 is a compound of Formula

P1H:




P1H

wherein:

R^1 is hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl; and

b is an integer from 2 to 8.

[0370] In some embodiments, the compound of Formula P1 is a compound of Formula P1I:




P1I

wherein:

R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl; and

b is an integer from 2 to 8.

[0371] In some embodiments, the compound of Formula P1 is a compound of Formula P1J:




P1J

wherein:

R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl; and

b is an integer from 2 to 8.


[0372] In some embodiments, the compound of Formula P1 is a compound of Formula P1K:



P1K

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0373] In some embodiments, the compound of Formula P1 is a compound of Formula P1L:



P1L

wherein R¹ is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0374] In some embodiments, the compound of Formula P1 is a compound of Formula P1M:



P1M

wherein R¹ is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0375] In some embodiments, the compound of Formula P1 is a compound of Formula P1N:



P1N

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0376] In some embodiments, the compound of Formula P1 is a compound of Formula P1O:



P1O

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0377] In some embodiments, the compound of Formula P1 is a compound of Formula P1P:



P1P

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0378] In some embodiments, the compound of Formula P1 is a compound of Formula P1Q:



P1Q

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0379] In some embodiments, the compound of Formula P1 is a compound of Formula P1R:



P1R

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0380] In some embodiments, the compound of Formula P1 is a compound of Formula P1S:



P1S

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl.


[0381] In some embodiments, the compound of Formula P1 is a compound of Formula P1T:



P1T

wherein R^N is a hydrogen atom or alkyl, R^M is alkyl, R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0382] In some embodiments, the compound of Formula P1 is a compound of Formula P1U:



P1U

wherein R^N is a hydrogen atom or alkyl, R^M is alkyl, R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0383] In some embodiments, the compound of Formula P1 is a compound of Formula P1V:



P1V

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, and b is an integer from 2 to 8.


[0384] In some embodiments, the compound of Formula P1 is a compound of Formula P1W:



P1W

wherein R^1 is a hydrogen atom, alkyl, alkoxy, halo, haloalkoxy, haloalkyl, or trifluoromethyl, g is an integer from 2 to 20; and b is an integer from 2 to 8.

[0385] In some embodiments, the compound of Formula P1 is a compound of Formula P1X:




P1X

wherein:

R^1 is alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy; and b is an integer from 2 to 8. In some embodiments, R^1 is methyl, ethyl, methoxy, or ethoxy. In some of these embodiments, R^1 is methoxy. In some embodiments, R^1 is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is hydrogen.


[0386] In some embodiments, the compound of Formula P1 is a compound of Formula P1Y:



P1Y




wherein R^1 is alkyl, alkenyl, alkynyl, alkoxy, heteroalkyl, halo, haloalkyl, haloalkoxy. In some embodiments, R^1 is methyl, ethyl, methoxy, or ethoxy. In some of these embodiments, R^1 is methoxy. In some embodiments, R^1 is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is hydrogen.





[0387] In some embodiments, the compound of Formula P1 is a compound of Formula P1Z:











P1Z,





wherein R^c is selected from alkyl or haloalkyl and wherein the alkyl or haloalkyl is linear, branched, or cyclic. In some embodiments, the compound of Formula P1 is a compound having one of the following structures:












































































































































































































































































or










[0388] In some embodiments, the compound of Formula P1 is a compound having one of the following structures:















[0389] Compounds of Formula (II) can be synthesized by contacting compounds of Formula PP5 with a suitable reducing agent:



PP5,


wherein A is arylene or heteroarylene.

[0390] In some embodiments, the suitable reducing agent includes a metal, a metal foil, a metal powder, a dust of a metal, a metal amalgam, or metal filings. In certain embodiments, the metal is selected from zinc, iron, aluminum, palladium, or Raney nickel.

[0391] For example, in some embodiments, the following reducing agent conditions are employed. With respect to the amount of compound PP5, for example, in some of the methods herein about twenty (20) equivalents of zinc dust and forty (40) equivalents of acetic acid were combined. In some examples, the reducing reaction was conducted at room temperature for about from 1 to 24 hours. In some of these examples, the aforementioned acetic acid is substituted with another suitable mild acid or proton donor. Examples of suitable mild acids or proton donors include, but are not limited to formic acid, pTsOH, and NH_4Cl . In some of these examples, the reducing metal is substituted with a suitable reducing agent selected from iron, aluminum, palladium, or Raney nickel. In some of these examples, suitable solvents includes those solvents having 10-50 % water (by volume) in a miscible organic solvent. Example miscible organic solvents include, but are not limited to THF, Dioxane, and diethyl ether. In some examples, the reducing reactions set forth herein are conducted at reaction temperatures which range from 0 to 50°C . In some examples, the reducing reactions set forth herein are conducted at reaction times which range from 1 to 40 hours.

[0392] Suitable acids include, but are not limited to, acetic acid.

[0393] In some embodiments, A is:



wherein:

R^1 is, independently at each occurrence, alkyl, alkenyl, alkynyl, aryl, alkaryl, arylalkyl, halo, haloalkoxy, haloalkyl, heteroalkyl, heteroaryl, heterocycloalkyl, cyano, nitro,

$-\xi-\text{OR}^A$, $-\xi-\text{SO}_2\text{R}^A$, $-\xi-\overset{\text{O}}{\parallel}-\text{R}^A$, or azido,

wherein R^A is alkyl;


n is an integer from 0 to 4;

m is an integer from 0 to 3;

p is an integer from 0 to 6; and

q is an integer from 0 to 5.


[0394] In some embodiments, the compound of Formula PP5 is a compound of the Formula PP5A:



PP5A,

wherein R^1 and n are as defined herein.

In some embodiments, the compound of PP5 is the a compound of Formula PP5A1:



(PP5A1)

wherein:


R^1 , independently at each occurrence, selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; and

n is an integer from 0 to 4.

[0395] In some embodiments, R^1 is, independently, alkyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl or C_{1-6} alkoxy. In some embodiments, R^1 is halo. In some embodiments, R^1 is, independently, alkoxy. In some embodiments, R^1 is, independently, methoxy, ethoxy, or propoxy. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In

some embodiments, R^1 is bromo. In some embodiments, n , m , p , or q is 0, 1 or 2. In some embodiments, n , m , p , or q is 0 or 1. In some embodiments, n , m , p , or q is 0.

[0396] In some embodiments, the compound of Formula PP5 is a compound of the Formula PP5A:




PP5A,

wherein:

R^1 is, independently at each occurrence, halo or trifluoromethyl; and
 n is 0, 1, or 2.

[0397] In some embodiments, the compound of Formula PP5 is a compound of the Formula PP5A2:




PP5A2,

wherein:

R^1 is, independently at each occurrence, halo or trifluoromethyl; and
 q is an integer from 0 to 5

[0398] In some embodiments, the compound of Formula PP5 is a compound of the Formula PP5A3:




PP5A3,


wherein:

R^1 is, independently at each occurrence, halo or trifluoromethyl; and q is an integer from 0 to 5. In some embodiments, R^1 is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is hydrogen.

[0399] Compounds of Formula PP5 can be synthesized by contacting compounds of Formula P2 with compounds of Formula PP6 under Lewis acid conditions:



P2,




PP6.

[0400] Suitable compounds of Formula PP6 include, but are not limited to, 4-isocyanato-2-methyl-1-nitro-benzene, 1-isocyanato-2-methyl-4-nitro-benzene, 2-bromo-4-isocyanato-1-nitro-benzene, 2-chloro-1-isocyanato-4-nitro-benzene, 3-chloro-1-isocyanato-4-nitro-benzene, 2-fluoro-1-isocyanato-4-nitro-benzene, 2-bromo-1-isocyanato-4-nitro-benzene, 4-isocyanato-2-methoxy-1-nitro-benzene,

1-isocyanato-2-methoxy-4-nitro-benzene, 4-isocyanato-1-nitro-2-trifluoromethyl-benzene, or 1-isocyanato-4-nitro-2-trifluoromethyl-benzene.


[0401] Suitable compounds of Formula PP6 include compounds having any one of the following formula:



wherein R¹ is, independently at each occurrence, C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkyl, or C₁₋₆ haloalkoxy, wherein n is 0, 1, 2, 3, or 4. In certain of these embodiments, R¹ is methoxy or methyl. In some specific embodiments, R¹ is methoxy, methyl, fluoro, chloro, bromo or trifluoromethyl. In certain embodiments, n is 1 or 2. In some of these embodiments, n is 1. In some embodiments, R¹ is fluoro, chloro, bromo, or iodo. In some embodiments, R¹ is methyl. In some embodiments, R¹ is methoxy. In some embodiments, R¹ is trifluoromethyl. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo.

[0402] In some embodiments, R¹ is 1-methylethyl-thiol, phenyl, 2-fluorophenyl, pyridinyl, 4-pyridinyl, pyrrolidinyl, or 1-pyrrolidinyl. In some embodiments, R¹ is trifluoromethyl. In some embodiments, R¹ is methoxy. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is hydrogen.





[0403] In some embodiments, provided herein are compounds of Formula PP5:



















PP5,









wherein A is arylene or heteroarylene.


[0404] In some embodiments, the compound of Formula PP5 is a compound selected from



















[0406] In some examples, set forth herein are methods of making a compound of Formula PP5. In some examples, these methods include contacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid conditions:




P2,



PP6.

[0407] In some of the methods herein, the contacting of a compound of Formula P2 with a compound of Formula PP6 is with a Lewis acid and an aprotic solvent in this contacting step. In some examples, the Lewis acid is selected from AlBr_3 , AlCl_3 , BCl_3 , boron trichloride methyl sulfide, BF_3 , boron trifluoride methyl etherate, boron trifluoride methyl sulfide, boron trifluoride tetrahydrofuran, dicyclohexylboron trifluoromethanesulfonate, iron (III) bromide, iron (III) chloride, tin (IV) chloride, titanium (IV) chloride, titanium (IV) isopropoxide, $\text{Cu}(\text{OTf})_2$, CuCl_2 , CuBr_2 , zinc chloride, alkylaluminum halides ($\text{R}_n\text{AlX}_{3-n}$, wherein R is hydrocarbyl), $\text{Zn}(\text{OTf})_2$, $\text{Yb}(\text{OTf})_3$, $\text{Sc}(\text{OTf})_3$, MgBr_2 , NiCl_2 , $\text{Sn}(\text{OTf})_2$, $\text{Ni}(\text{OTf})_2$, or $\text{Mg}(\text{OTf})_2$. In some examples, the Lewis acid is zinc chloride. In some examples, the aprotic solvent is selected from perfluorohexane ($\text{CF}_3(\text{CF}_2)_4\text{CF}_3$), a,a,a-trifluorotoluene, pentane, hexane, cyclohexane, methylcyclohexane, dioxane, carbon tetrachloride, freon-11 (CFCl_3), benzene, toluene, triethyl amine, carbon disulfide, diisopropyl ether, diethyl ether, t-butyl methyl ether, chloroform, ethyl acetate, 1,2-dimethoxyethane (glyme), 2-methoxyethyl ether (diglyme), tetrahydrofuran, dichloromethane (methylene chloride), pyridine, 2-butanone (MEK), acetone, hexamethylphosphoramide, N-methylpyrrolidinone, nitromethane, dimethylformamide, acetonitrile, sulfolane, dimethyl sulfoxide, or propylene carbonate. In some examples, the aprotic solvent is dichloromethane. In some of the methods herein, the contacting of a compound of Formula P2 with a compound of Formula PP6 includes contacting zinc chloride and dichloromethane with a compound of Formula P2 and a compound of Formula PP6.

[0408] In some of these methods, the compound of Formula PP5 has the Formula PP5A:




PP5A,

wherein R^1 and n are as defined herein.

[0409] In some of these methods, R^1 is, independently, alkyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy. In some embodiments, R^1 is, independently, C_{1-6} alkyl, C_{1-6} alkoxy, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, or halo. In some embodiments, R^1 is, independently, C_{1-6} alkyl or C_{1-6} alkoxy. In some embodiments, R^1 is halo. In some embodiments, R^1 is, independently, alkoxy. In some embodiments, R^1 is, independently, methoxy, ethoxy, or propoxy. In some embodiments, R^1 is methyl. In some embodiments, R^1 is methoxy. In some embodiments, R^1 is trifluoromethyl. In some embodiments, R^1 is fluoro. In some embodiments, R^1 is chloro. In some embodiments, R^1 is bromo. In some embodiments, n, m, p, or q is 0, 1 or 2. In some embodiments, n, m, p, or q is 0 or 1. In some embodiments, n, m, p, or q is 0.

[0410] In some of these methods, the compound of Formula PP5 has the Formula PP5A:




PP5A,

wherein:

R^1 is, independently, alkyl, alkoxy, heteroalkyl, halo, haloalkyl, or haloalkoxy; and n is 0, 1, or 2.


[0411] In some embodiments, the methods disclosed herein include a compound of PP6 selected from 4-isocyanato-2-methyl-1-nitro-benzene, 1-isocyanato-2-methyl-4-nitro-benzene, 2-bromo-4-isocyanato-1-nitro-benzene, 2-chloro-1-isocyanato-4-nitro-benzene, 3-chloro-1-isocyanato-4-nitro-benzene, 2-fluoro-1-isocyanato-4-nitro-benzene, 2-bromo-1-isocyanato-4-nitro-benzene, 4-isocyanato-2-methoxy-1-nitro-benzene, 1-isocyanato-2-methoxy-4-nitro-benzene, 4-isocyanato-1-nitro-2-trifluoromethyl-benzene, or 1-isocyanato-4-nitro-2-trifluoromethyl-benzene.

[0412] In some embodiments, the methods disclosed herein include a compound of PP6 having any one of the following formula:



wherein R¹ is, independently at each occurrence, C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, C₁₋₆ haloalkyl, or C₁₋₆ haloalkoxy, wherein n is 0, 1, 2, 3, or 4. In certain of these embodiments, R¹ is methoxy or methyl. In some specific embodiments, R¹ is methoxy, methyl, fluoro, chloro, bromo or trifluoromethyl. In certain embodiments, n is 1 or 2. In some of these embodiments, n is 1. In some embodiments, R¹ is fluoro, chloro, bromo, or iodo. In some embodiments, R¹ is methyl. In some embodiments, R¹ is methoxy. In some embodiments, R¹ is trifluoromethyl. In some embodiments, R¹ is fluoro. In some embodiments, R¹ is chloro. In some embodiments, R¹ is bromo.

[0413] In some embodiments, the methods disclosed herein further including reducing a compound of Formula PP5 by contacting a compound of Formula PP5 with a suitable reducing agent:



PP5,

wherein A is arylene or heteroarylene.


[0414] In some of these methods, the suitable reducing agent includes a metal, a metal foil, a metal powder, a dust of a metal, a metal amalgam, or metal filings. In certain

embodiments, the metal is selected from zinc, iron, aluminum, palladium, or Raney nickel. In some of these methods, the suitable reducing agent is zinc. In some of these methods, the suitable reducing agent is zinc in acetic acid. In some of these methods, the suitable reducing agent is zinc dust in acetic acid.

[0415] In some embodiments, the methods disclosed herein further including reducing a compound of Formula PP5 by contacting a compound of Formula PP5 with a suitable reducing agent under particular reducing agent conditions. In some examples, the methods include reducing a compound of Formula PP5 with a reducing agent selected from zinc dust, iron, aluminum, palladium, or Raney nickel. In some examples, the methods include reducing a compound of Formula PP5 with zinc dust. In some examples, the methods include reducing a compound of Formula PP5 with zinc dust and a solvent selected from acetic acid, formic acid, pTsOH, or NH₄Cl. In some examples, the methods include reducing a compound of Formula PP5 with zinc dust and acetic acid. In some examples, the methods include reducing a compound of Formula PP5 with about twenty (20) equivalents of zinc dust and forty (40) equivalents of acetic acid. In some examples, the reducing is conducted at a temperature between 0 and 70 °C. In some examples, the reducing is conducted at a temperature of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, or 70 °C. In some examples, the reducing is conducted for 1 to 40 hours. In some examples, the reducing is conducted for 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 hours. In some examples, the reducing is conducted for 24 hours. In some examples, the reducing is conducted at room temperature for about from 1 to 24 hours.


[0416] In some examples, the above method steps are performed or conducted in a solvent having 10-50 % water (by volume) in a miscible organic solvent. In some examples, the miscible organic solvents is THF, Dioxane, or diethyl ether.

[0417] Compounds of Formula III:



III

can be synthesized by contacting compounds of Formula PP1:



PP1

with a binding agent under conjugation conditions,
wherein:

BA is a binding agent;

SP is a spacer;

SP^R is a reactive spacer;


R^{AA1} is an amino acid side chain;

R^{AA2} is an amino acid side chain;

A is arylene or heteroarylene; and

k is an integer from 1 to 30.

[0418] Compounds of Formula PP1 can be prepared by contacting a compound of Formula (II) with the compound of Formula PP7:




wherein:


SP^R is a reactive spacer;

R^{AA1} is an amino acid side chain;
 R^{AA2} is an amino acid side chain; and
 A is arylene or heteroarylene.


[0419] Compounds of Formula PP7 can be prepared by contacting a compound of Formula PP9 with a bifunctional spacer:



[0420] Illustrative bifunctional spacers include, but are not limited to:




[0421] Compounds of Formula PP1 can be prepared by contacting an activated form of a compound of Formula PP10 with a compound of Formula (II):



wherein Y is a moiety that renders the carboxylic group to which it is attached electrophilic. Compound of Formula PP10 can be prepared by first coupling its corresponding amino acids using standard amino acid coupling techniques, including, for example, active ester formation using HATU, BOP/HOBt, or EDC/N-hydroxysuccinamide in the presence of DIEA, DBU, EEDG or tributylamine. Compound of Formula PP10 can be prepared by then subsequently attaching the RL using standard coupling techniques, including, for example, active ester

formation using HATU, BOP/HOBt, or EDC/N-hydroxysuccinamide in the presence of DIEA, DBU, EEDQ or tributylamine.

[0422] Bifunctional spacers are compounds that react with the compound of Formula PP9 to append the SP^R moiety present in the compounds of Formula PP7. Illustrative bifunctional spacers include, but are not limited to:



E. METHODS OF USE AND PHARMACEUTICAL COMPOSITIONS

[0423] The present disclosure includes methods of treating or preventing diseases, conditions, or disorders *e.g.*, proliferative diseases such as cancer, comprising administering a therapeutically effective amount or one or more of the compounds disclosed herein, *e.g.*, one or more of the compounds of Formula (I) or (II). Diseases, disorders, and/or conditions include, but are not limited to, those associated with the antigens listed herein. In some embodiments, the antigen is PSMA, MUC16, STEAP2, or EGFRvIII.

[0424] The compounds disclosed herein can be used for treating primary and/or metastatic tumors arising in the brain and meninges, oropharynx, lung and bronchial tree, gastrointestinal tract, male and female reproductive tract, muscle, bone, skin and appendages, connective tissue, spleen, immune system, blood forming cells and bone marrow, liver and urinary tract, and special sensory organs such as the eye. In certain embodiments, the compounds provided herein are used to treat one or more of the following cancers: renal cell carcinoma, pancreatic carcinoma, head and neck cancer, prostate cancer, malignant gliomas, osteosarcoma, colorectal cancer, gastric cancer (*e.g.*, gastric cancer with MET amplification), malignant mesothelioma, multiple myeloma, ovarian cancer, small cell lung cancer, non-small cell lung cancer, synovial sarcoma, thyroid cancer, breast cancer, or melanoma. In some embodiments, the cancer is breast cancer.

[0425] The compounds described herein can be administered alone or together with one or more additional therapeutic agents. The one or more additional therapeutic agents can be administered just prior to, concurrent with, or shortly after the administration of the compounds described herein. The present disclosure also includes pharmaceutical compositions comprising any of the compounds described herein in combination with one or more additional therapeutic agents, and methods of treatment comprising administering such combinations to subjects in need thereof.

[0426] Suitable additional therapeutic agents include, but are not limited to: an EGFR antagonist (*e.g.*, an anti-EGFR antibody [*e.g.*, cetuximab or panitumumab] or small molecule inhibitor of EGFR [*e.g.*, gefitinib or erlotinib]), an antagonist of another EGFR family member such as Her2/ErbB2, ErbB3 or ErbB4 (*e.g.*, anti-ErbB2 [*e.g.*, trastuzumab or T-DM1 {KADCYLA®}], anti-ErbB3 or anti-ErbB4 antibody or small molecule inhibitor of ErbB2, ErbB3 or ErbB4 activity), an antagonist of EGFRvIII (*e.g.*, an antibody that specifically binds EGFRvIII), a cMET antagonist (*e.g.*, an anti-cMET antibody), an IGF1R antagonist (*e.g.*, an anti-IGF1R antibody), a B-raf inhibitor (*e.g.*, vemurafenib, sorafenib, GDC-0879, PLX-4720), a PDGFR- α inhibitor (*e.g.*, an anti-PDGFR- α antibody), a PDGFR- β inhibitor (*e.g.*, an anti-PDGFR- β antibody or small molecule kinase inhibitor such as, *e.g.*, imatinib mesylate or sunitinib malate), a PDGF ligand inhibitor (*e.g.*, anti-PDGF-A, -B, -C, or -D antibody, aptamer, siRNA, etc.), a VEGF antagonist (*e.g.*, a VEGF-Trap such as aflibercept, *see, e.g.*, US 7,087,411 (also referred to herein as a "VEGF-inhibiting fusion protein"), anti-VEGF antibody (*e.g.*,

bevacizumab), a small molecule kinase inhibitor of VEGF receptor (*e.g.*, sunitinib, sorafenib or pazopanib)), a DLL4 antagonist (*e.g.*, an anti-DLL4 antibody disclosed in US 2009/0142354 such as REGN421), an Ang2 antagonist (*e.g.*, an anti-Ang2 antibody disclosed in US 2011/0027286 such as H1H685P), a FOLH1 antagonist (*e.g.*, an anti-FOLH1 antibody), a STEAP1 or STEAP2 antagonist (*e.g.*, an anti-STEAP1 antibody or an anti-STEAP2 antibody), a TMPRSS2 antagonist (*e.g.*, an anti-TMPRSS2 antibody), a MSLN antagonist (*e.g.*, an anti-MSLN antibody), a CA9 antagonist (*e.g.*, an anti-CA9 antibody), a uroplakin antagonist (*e.g.*, an anti-uroplakin [*e.g.*, anti-UPK3A] antibody), a MUC16 antagonist (*e.g.*, an anti-MUC16 antibody), a Tn antigen antagonist (*e.g.*, an anti-Tn antibody), a CLEC12A antagonist (*e.g.*, an anti-CLEC12A antibody), a TNFRSF17 antagonist (*e.g.*, an anti-TNFRSF17 antibody), a LGR5 antagonist (*e.g.*, an anti-LGR5 antibody), a monovalent CD20 antagonist (*e.g.*, a monovalent anti-CD20 antibody such as rituximab), etc. Other agents that may be beneficially administered in combination with compounds of the disclosure include, *e.g.*, tamoxifen, aromatase inhibitors, and cytokine inhibitors, including small-molecule cytokine inhibitors and antibodies that bind to cytokines such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-8, IL-9, IL-11, IL-12, IL-13, IL-17, IL-18, or to their respective receptors.

[0427] Suitable therapeutic agents also include, but are not limited to chemotherapeutic agents, including alkylating agents such as thiotepa and cyclophosphamide (CytosanTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as

methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as froinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfornithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSKTM; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxanes, e.g. paclitaxel (TaxolTM, Bristol-Myers Squibb Oncology, Princeton, N.J.) and docetaxel (TaxotereTM; Aventis Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY 117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0428] The compounds described herein can also be administered and/or co-formulated in combination with antivirals, antibiotics, analgesics, corticosteroids, steroids, oxygen, antioxidants, COX inhibitors, cardioprotectants, metal chelators, IFN-gamma, and/or NSAIDs.

[0429] In some embodiments of the methods described herein, multiple doses of a compound described herein (or a pharmaceutical composition comprising a combination of an

compound described herein and any of the additional therapeutic agents mentioned herein) may be administered to a subject over a defined time course. The methods according to this aspect of the disclosure comprise sequentially administering to a subject multiple doses of a compound described herein. As used herein, "sequentially administering" means that each dose of the compound is administered to the subject at a different point in time, *e.g.*, on different days separated by a predetermined interval (*e.g.*, hours, days, weeks or months). The present disclosure includes methods which comprise sequentially administering to the patient a single initial dose of a compound described herein, followed by one or more secondary doses of the compound, and optionally followed by one or more tertiary doses of the compound.

[0430] The terms "initial dose," "secondary doses," and "tertiary doses," refer to the temporal sequence of administration of the compounds described herein. Thus, the "initial dose" is the dose which is administered at the beginning of the treatment regimen (also referred to as the "baseline dose"); the "secondary doses" are the doses which are administered after the initial dose; and the "tertiary doses" are the doses which are administered after the secondary doses. The initial, secondary, and tertiary doses can all contain the same amount the compound described herein, but generally can differ from one another in terms of frequency of administration. In certain embodiments, the amount of the compound contained in the initial, secondary and/or tertiary doses varies from one another (*e.g.*, adjusted up or down as appropriate) during the course of treatment. In certain embodiments, two or more (*e.g.*, 2, 3, 4, or 5) doses are administered at the beginning of the treatment regimen as "loading doses" followed by subsequent doses that are administered on a less frequent basis (*e.g.*, "maintenance doses").

[0431] In certain exemplary embodiments of the present disclosure, each secondary and/or tertiary dose is administered 1 to 26 (*e.g.*, 1, 1½, 2, 2½, 3, 3½, 4, 4½, 5, 5½, 6, 6½, 7, 7½, 8, 8½, 9, 9½, 10, 10½, 11, 11½, 12, 12½, 13, 13½, 14, 14½, 15, 15½, 16, 16½, 17, 17½, 18, 18½, 19, 19½, 20, 20½, 21, 21½, 22, 22½, 23, 23½, 24, 24½, 25, 25½, 26, 26½, or more) weeks after the immediately preceding dose. The phrase "the immediately preceding dose," as used herein, means, in a sequence of multiple administrations, the dose the compound which is administered to a patient prior to the administration of the very next dose in the sequence with no intervening doses.

[0432] The methods according to this aspect of the disclosure may comprise administering to a patient any number of secondary and/or tertiary doses of the compound. For example, in certain embodiments, only a single secondary dose is administered to the patient. In other embodiments, two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, or more) secondary doses are administered to the patient. Likewise, in certain embodiments, only a single tertiary dose is administered to the patient. In other embodiments, two or more (*e.g.*, 2, 3, 4, 5, 6, 7, 8, or more) tertiary doses are administered to the patient. The administration regimen may be carried out indefinitely over the lifetime of a particular subject, or until such treatment is no longer therapeutically needed or advantageous.

[0433] In embodiments involving multiple secondary doses, each secondary dose may be administered at the same frequency as the other secondary doses. For example, each secondary dose may be administered to the patient 1 to 2 weeks or 1 to 2 months after the immediately preceding dose. Similarly, in embodiments involving multiple tertiary doses, each tertiary dose may be administered at the same frequency as the other tertiary doses. For example, each tertiary dose may be administered to the patient 2 to 12 weeks after the immediately preceding dose. In certain embodiments of the disclosure, the frequency at which the secondary and/or tertiary doses are administered to a patient can vary over the course of the treatment regimen. The frequency of administration may also be adjusted during the course of treatment by a physician depending on the needs of the individual patient following clinical examination.

[0434] The present disclosure includes administration regimens in which 2 to 6 loading doses are administered to a patient at a first frequency (*e.g.*, once a week, once every two weeks, once every three weeks, once a month, once every two months, etc.), followed by administration of two or more maintenance doses to the patient on a less frequent basis. For example, according to this aspect of the disclosure, if the loading doses are administered at a frequency of once a month, then the maintenance doses may be administered to the patient once every six weeks, once every two months, once every three months, *etc.*

[0435] The present disclosure includes pharmaceutical compositions of the compounds and/or conjugates described herein, *e.g.*, the compounds of Formula (I) and (II), *e.g.*, compositions comprising a compound described herein, a salt, stereoisomer, polymorph thereof, and a pharmaceutically acceptable carrier, diluent, and/or excipient. Examples of suitable carriers, diluents and excipients include, but are not limited to: buffers for maintenance of proper

composition pH (*e.g.*, citrate buffers, succinate buffers, acetate buffers, phosphate buffers, lactate buffers, oxalate buffers and the like), carrier proteins (*e.g.*, human serum albumin), saline, polyols (*e.g.*, trehalose, sucrose, xylitol, sorbitol, and the like), surfactants (*e.g.*, polysorbate 20, polysorbate 80, polyoxolate, and the like), antimicrobials, and antioxidants.

F. EXAMPLES

[0436] Proton NMR spectra of linkers were acquired on a Varian 300 MHz instrument, while spectra of payloads and payload-linkers were acquired on a Varian 500 MHz instrument. Mass spectra were collected on an Agilent LCMS instruments using electrospray ionization and either an ion trap mass analyzer or a single quadrupole analyzer. For all Examples, only the most abundant isotopes acquired by mass spectrometry are reported. All starting materials were purchased commercially and used without purification, unless otherwise noted, while solvents were purchased commercially and dried where necessary via methods well known in the art. The following is a list of the abbreviations used in the Examples, with their full chemical names in parentheses: Ahx (6-aminohexanoic acid), Boc (*N-tert*-butoxycarbonyl), Cap (caproyl), CDCl₃ (chloroform-*d*), CD₃OD (methanol-*d*₄), Cit (L-citrulline), DCM (dichloromethane), DIEA (*N,N*-diisopropylethylamine), DMF (*N,N*-dimethylformamide), EDC (*N*-(3-Dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride), EEDQ (2-Ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline), EtOAc (ethyl acetate), Fmoc (9-fluorenylmethoxycarbonyl), HATU (1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate), HCl (aqueous hydrochloric acid), HOAc (glacial acetic acid), HOAT (1-hydroxy-7-azabenzotriazole), HPLC (high-performance liquid chromatography), LCMS (tandem HPLC and mass spectrometry), Mal (maleimide), MeCN (acetonitrile), MeOH (methanol), MS (low resolution mass spectrometry), NaHCO₃ (sodium bicarbonate), Na₂SO₄ (anhydrous sodium sulfate), NH₄Cl (ammonium chloride), NHS (*N*-hydroxysuccinimide), NMR (nuclear magnetic resonance spectroscopy), OSu (succinate ester), TEA (triethylamine), TFA (trifluoroacetic acid), THF (tetrahydrofuran), TLC (thin-layer chromatography), Val (L-valine), ZnCl₂ (zinc chloride).

EXAMPLE 1

[0437] Example 1 refers to compounds illustrated in FIG. 9.

Maytan-3-O-carbamoyl-N-phenyl-p-(amino-Cit-Val-Cap-Mal) (9)

Step A: L-valine-L-citrulline (2):

[0438] To a 50 mL round bottom flask equipped with a magnetic stirrer and nitrogen inlet was charged Boc-L-valine-L-citrulline (**1**, 0.5 g; 1.3 mmol), acetonitrile (10 mL) and water (10 mL). An aqueous solution of 1 M HCl (13 mL; 13 mmol) was added and this solution was stirred at ambient temperature for 18 hours (h). The solution was frozen and lyophilized to a white solid. The solid was extracted with 10% MeOH in DCM, the organic layer was dried over Na₂SO₄, filtered and concentrated to afford the title compound (0.4 g, 98% yield) as a clear colorless oil. MS (ESI, pos. & neg.): calc'd for C₁₁H₂₂N₄O₄, 274.2; found 275.2 (M+H), 273.2 (M-H).

Step B: 6-Maleimidyl-caproamidyl-L-valine-L-citrulline (4):

[0439] To a 25 mL round bottom flask equipped with a magnetic stirrer and nitrogen inlet was charged L-valine-L-citrulline (**2**, 0.4 g; 1.3 mmol), 6-maleimido-caproic acid succinate ester (**3**, 0.6 g; 2.0 mmol) in DMF (3 mL) followed by DIEA (0.4 mL; 2.0 mmol). The solution was stirred at ambient temperature for 18 hours, diluted with water (2 mL) and loaded onto a C18 column eluting with 5-95% MeCN in water (containing 0.05% HOAc). The fractions containing product were combined, frozen and lyophilized to afford the title compound (0.3 g, 47% yield) as a white solid. MS (ESI, pos. & neg.): calc'd for C₂₁H₃₃N₅O₇, 467.2; found 468.4 (M+H), 466.4 (M-H).

Step C: Maytan-3-O-carbamoyl-N-(4-nitrobenzene) (7):

[0440] To a dry round bottom flask was weighed maytansinol (**6**) (prepared as previously described in the art, 114 mg, 0.202 mmol) and p-nitrophenyl isocyanate (**5**, 73 mg, 0.445 mmol). The mixture was dissolved in dry DCM (10 mL) and treated dropwise with a 1.0 M solution of zinc chloride in diethyl ether (0.50 mL, 0.50 mmol). The flask was sealed with a rubber septum, purged with argon via needle, and the reaction stirred at ambient temperature for 18 h. LCMS showed complete conversion of the maytansinol. The reaction was diluted with water and

extracted three times with EtOAc. The combined organic layers were washed with saturated aqueous (aq.) NaHCO₃ and brine, dried over Na₂SO₄, and filtered. The evaporated product was not very soluble in MeCN/water, so it was concentrated to dryness again *in vacuo*, dissolved in EtOAc/DCM, and filtered over Na₂SO₄. The filtrate was then purified by flash chromatography on silica gel (gradient elution: 50 – 100% EtOAc in hexanes) and the fractions containing product were combined, concentrated, and dried *in vacuo* giving the title compound as a yellow solid (50 mg, 34%). MS (ESI, pos.): calc'd for C₃₅H₄₁N₄O₁₁Cl, 728.3; found 729.2 (M+H).

Step D: Maytan-3-O-carbamoyl-N-(4-aminobenzene) (8):

[0441] The product of the preceding step (7, 49 mg, 0.067 mmol) and zinc dust (85 mg, 1.30 mmol) were weighed into a round bottom flask and treated with THF (4 mL), water (1 mL), and HOAc (0.100 mL, 1.75 mmol). The flask was sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature for 18 h. The reaction was filtered over Celite, solids rinsed with acetonitrile (MeCN, 3 x 5 mL), and the filtrate concentrated *in vacuo* to an amber oil. This was purified directly on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both, over 12 min). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 3 days (d) giving the title compound as a cream-colored solid (32 mg, 63%). MS (ESI, pos.): calc'd for C₃₅H₄₃N₄O₉Cl, 698.3; found 699.2 (M+H).

Step E: Maytan-3-O-carbamoyl-N-phenyl-p-(amino-Cit-Val-Cap-Mal) (9):

[0442] The product of the preceding step (8, 16 mg, 0.021 mmol), 6-Maleimidyl-caproamidyl-L-valine-L-citrulline (4, 16 mg, 0.034 mmol), and EEDQ (15 mg, 0.061 mmol) were weighed into a round bottom flask and dissolved in dry DCM (2 mL) and anhydrous methanol (1 mL). The flask was sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature for 18 h. The reaction was concentrated *in vacuo*, dissolved in MeCN/water, and purified directly on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both, over 12 min). The product did not separate well, so the impure product fractions were lyophilized and repurified

twice by preparative HPLC on a Phenomenex Gemini 5u, 30x150mm C18 column (first 40 – 80% then 20 – 80% MeCN in water, 0.1% HOAc both, over 20 min, 30 mL/min). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized overnight giving the title compound as a white solid (5 mg, 21%). MS (ESI, pos.): calc'd for C₅₆H₇₄N₉O₁₅Cl, 1147.5; found 1148.5 (M+H), 1130.4 (M-H₂O+H), 1170.5 (M+Na). ¹H NMR (500 MHz, Methanol-*d*₄) δ: 7.48 – 7.42 (m, 2H), 7.40 (t, J = 8.3 Hz, 2H), 7.07 (s, 1H), 7.02 (d, J = 1.8 Hz, 1H), 6.69 (d, J = 4.7 Hz, 1H), 6.49 (dd, J = 15.4, 11.1 Hz, 1H), 6.14 (d, J = 8.9 Hz, 1H), 5.33 (dd, J = 15.4, 8.7 Hz, 1H), 4.47 (dd, J = 11.9, 2.9 Hz, 1H), 4.40 (dd, J = 8.9, 5.2 Hz, 1H), 4.25 – 4.14 (m, 1H), 4.07 (dd, J = 12.0, 7.6 Hz, 1H), 3.88 (s, 3H), 3.50 (d, J = 12.7 Hz, 1H), 3.43 (d, J = 8.9 Hz, 1H), 3.38 (t, J = 7.1 Hz, 2H), 3.18 – 3.13 (m, 2H), 3.11 (s, 4H), 3.09 – 2.97 (m, 2H), 2.90 (dd, J = 9.8, 3.7 Hz, 1H), 2.58 – 2.47 (m, 1H), 2.18 (t, J = 7.5 Hz, 2H), 2.14 – 2.04 (m, 2H), 2.00 – 1.92 (m, 1H), 1.87 (d, J = 4.9 Hz, 1H), 1.80 (dd, J = 10.0, 4.8 Hz, 1H), 1.67 (dd, J = 13.7, 2.1 Hz, 2H), 1.63 (s, 4H), 1.59 – 1.41 (m, 8H), 1.41 – 1.33 (m, 2H), 1.25 – 1.17 (m, 4H), 1.17 – 1.09 (m, 4H), 1.04 (d, J = 6.5 Hz, 1H), 0.94 (dd, J = 11.7, 6.7 Hz, 1H), 0.87 (dt, J = 7.9, 4.0 Hz, 6H), 0.77 (s, 3H).

EXAMPLE 2

[0443] Example 2 refers to compounds illustrated in FIG. 10.

Maytan-3-O-carbamoyl-N-phenyl-p-amino-adipic Acid Succinate Ester (10)

Step A: Maytan-3-O-carbamoyl-N-phenyl-p-amino-adipic acid:

[0444] The product of Example 1, step D (**8**, 20 mg, 0.0263 mmol) and adipic anhydride (a.k.a. oxepane-2,7-dione, 20 mg, 0.156 mmol) were weighed into a round bottom flask, dissolved in THF (2.0 mL), and treated with triethylamine (0.010 mL, 0.0717 mmol). The solids would not dissolve, so DMF (1.0 mL) was added, the flask sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature. After 18 h, LCMS showed complete conversion of **2**, so the reaction was concentrated to an oil *in vacuo*, dissolved in MeCN/water, and purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in

water, 0.05% acetic acid in both). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 18 h giving the title compound as a white solid (19 mg, 86%). MS (ESI, pos.): calc'd for C₄₁H₅₁ClN₄O₁₂, 826.3; found 827.3 (M+H), 849.3 (M+Na).

Step B: Maytan-3-O-carbamoyl-N-phenyl-p-amino-adipic Acid Succinate Ester (10):

[0445] The product of the preceding step (18 mg, 0.0218 mmol), N-hydroxysuccinimide (19 mg, 0.165 mmol), and EDC hydrochloride (34 mg, 0.177 mmol) were weighed into a round bottom flask, dissolved in DCM (3 mL), the flask sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature. After 20 h the reaction was concentrated *in vacuo*, dissolved in MeCN, treated with a couple drops of 10% aqueous acetic acid, and purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 30 – 90% MeCN in water, 0.05% acetic acid in both). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 18 h giving the title compound as a white solid (10 mg, 50%). MS (ESI, pos.): calc'd for C₄₅H₅₄N₅O₁₄Cl, 923.3; found 924.3 (M+H), 946.3 (M+Na). ¹H NMR (500 MHz, CDCl₃) δ: 7.62 (s, 1H), 7.54 (d, 2H, *J* = 9 Hz), 7.39 (d, 2H, *J* = 9 Hz), 6.84 (d, 1H, *J* = 2 Hz), 6.69 (d, 1H, *J* = 2 Hz), 6.58 (s, 1H), 6.47 (dd, 1H, *J* = 16 Hz, 11 Hz), 6.18 (s, 1H), 6.10 (d, 1H, *J* = 11 Hz), 5.37 (dd, 1H, *J* = 16 Hz, 9 Hz), 4.81 (dd, 1H, *J* = 12 Hz, 2 Hz), 4.32 (t, 1H, *J* = 10 Hz), 4.00 (s, 3H), 3.50 (d, 1H, *J* = 14 Hz), 3.47 (d, 1H, *J* = 9 Hz), 3.31 (s, 3H), 3.27 (d, 1H, *J* = 13 Hz), 3.20 (s, 3H), 2.89 (m, 6H), 2.68 (m, 2H), 2.51 (dd, 1H, *J* = 14 Hz, 12 Hz), 2.43 (m, 2H), 2.26 (dd, 1H, *J* = 14 Hz, 2 Hz), 1.89 (m, 5H), 1.71 (s, 3H), 1.29 (m, 5H), 0.91 (s, 3H).

EXAMPLE 3

Example 3 refers to compounds illustrated in FIG. 11.

Maytan-3-O-carbamoyl-N-phenyl-p-amino-Cit-Val-adipic Acid Succinate Ester (14):

Step A: Maytan-3-O-carbamoyl-N-phenyl-p-amino-Cit-Val-Fmoc (12):

[0446] The product of Example 1, step D (**8**, 54 mg, 0.071 mmol) and Fmoc-L-valine-L-citrulline (**11**, 151 mg, 0.304 mmol) were coupled via the method of Example 1, step E giving the title compound as a white solid (51 mg, 61%). MS (ESI, pos.): calc'd for C₆₁H₇₃ClN₈O₁₄, 1176.5; found 1178.1 (M+H), 1200.1 (M+Na).

Step B: Maytan-3-O-carbamoyl-N-phenyl-p-amino-Cit-Val (**13**):

[0447] The product of the preceding step (**12**, 49 mg, 0.042 mmol) was dissolved in a 5% v/v solution of piperidine in DMF (2.0 mL), and the mixture stirred at ambient temperature. After 20 h the reaction was acidified with a few drops of 10% aqueous acetic acid, diluted with MeCN and water (ca. 1 mL each), and purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 18 h giving the title compound as a cream-colored solid (34 mg, 81%). MS (ESI, pos.): calc'd for C₄₆H₆₃ClN₈O₁₂, 954.5; found 955.9 (M+H), 977.9 (M+Na).

Step C: Maytan-3-O-carbamoyl-N-phenyl-p-amino-Cit-Val-adipic Acid:

[0448] The product of the preceding step (**13**, 49 mg, 0.042 mmol) and adipic anhydride (a.k.a. oxepane-2,7-dione, 20 mg, 0.156 mmol) were weighed into a round bottom flask, dissolved in pyridine (2 mL), the flask sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature. After 18 h the reaction was concentrated to dryness under high vacuum, acidified with a few drops of 10% aqueous acetic acid, dissolved in MeCN, and purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 3 d giving the title compound as a white solid (14 mg, 40%). MS (ESI, pos.): calc'd for C₅₂H₇₁ClN₈O₁₅, 1082.5; found 1084.0 (M+H), 1106.1 (M+Na).

Step D: Maytan-3-O-carbamoyl-N-phenyl-p-amino-Cit-Val-adipic Acid Succinate Ester

(14):

[0449] Using the method of Example 2, step B, the title compound was prepared from the product of the preceding step (13 mg, 0.012 mmol) as a white solid (7 mg, 50%). MS (ESI, pos.): calc'd for C₅₆H₇₄ClN₉O₁₇, 1179.5; found 1182.1 (M+H), 1203.2 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.91 (s, 1H), 8.48 (s, 1H), 8.08 (d, *J* = 7.6 Hz, 1H), 7.84 (d, *J* = 8.6 Hz, 1H), 7.54 (d, *J* = 8.8 Hz, 2H), 7.44 (d, *J* = 8.5 Hz, 2H), 7.26 (s, 1H), 7.19 (s, 1H), 6.77 (s, 1H), 6.59 – 6.46 (m, 1H), 6.36 (s, 1H), 5.98 (s, 1H), 5.53 (s, 1H), 5.40 (s, 2H), 5.30 (dd, *J* = 15.1, 9.1 Hz, 1H), 4.44 – 4.31 (m, 2H), 4.27 – 4.18 (m, 1H), 4.14 (t, *J* = 11.2 Hz, 1H), 3.94 (s, 3H), 3.63 (d, *J* = 12.0 Hz, 1H), 3.42 (d, *J* = 9.1 Hz, 1H), 3.18 (s, 4H), 3.10 (s, 3H), 3.06 – 2.98 (m, 1H), 2.98 – 2.91 (m, 1H), 2.89 (d, *J* = 9.7 Hz, 1H), 2.81 (s, 4H), 2.73 – 2.65 (m, 2H), 2.30 – 2.22 (m, 1H), 2.21 – 2.14 (m, 1H), 2.06 (d, *J* = 13.2 Hz, 1H), 1.97 (dd, *J* = 12.5, 5.6 Hz, 1H), 1.66 (s, 3H), 1.60 (d, *J* = 5.5 Hz, 5H), 1.53 – 1.30 (m, 6H), 1.23 (s, 1H), 1.11 (d, *J* = 6.3 Hz, 3H), 0.85 (dd, *J* = 15.2, 6.7 Hz, 6H), 0.75 (s, 3H).

EXAMPLE 4

[0450] Example 4 refers to compounds illustrated in FIG. 12.

Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-Cap-Mal) (19)

Step A: Maytan-3-O-carbamoyl-N-(2-chloro-4-nitro)benzene (16):

[0451] To a dry round bottom flask was weighed 2-chloro-4-nitrophenyl isocyanate (**15**, 525 mg, 2.64 mmol) and maytansinol (**6**, 503 mg, 0.890 mmol). The mixture was dissolved in dry DCM (10 mL) and treated dropwise with a 1.0 M solution of zinc chloride in diethyl ether (1.20 mL, 1.20 mmol). The flask was sealed with a rubber septum, purged with argon via needle, and the reaction stirred at ambient temperature. After 18 h the reaction was incomplete by LCMS, so the flask was topped with a water condenser and the reaction heated to 50 °C. After 6 h, most of the maytansinol was consumed by LCMS so the reaction was cooled to ambient temperature and stirred overnight. After another 18 h the reaction was concentrated *in vacuo*, dissolved in MeCN/DCM, and purified by flash chromatography on silica gel (gradient

elution: 10 – 50% MeCN in DCM). The slow-running product fractions were combined, concentrated, and dried *in vacuo* for 3 d giving the title compound as an orange solid (396 mg, 58%). MS (ESI, pos.): calc'd for C₃₅H₄₀Cl₂N₄O₁₁, 762.2; found 763.1 (M+H).

Step B: Maytan-3-O-carbamoyl-N-(2-chloro-4-amino)benzene (17):

[0452] The title compound was prepared from the product of the preceding step (**16**, 388 mg, 0.509 mmol) as a white solid (301 mg, 75%) using the method of Example 1, step D, except that MeCN was used instead of THF and the reaction was stirred for only 2 h. MS (ESI, pos.): calc'd for C₃₅H₄₂Cl₂N₄O₉, 732.2; found 733.1 (M+H). ¹H NMR (500 MHz, Chloroform-*d*) δ: 7.83 (d, *J* = 8.8 Hz, 1H), 6.85 (s, 2H), 6.79 (s, 1H), 6.74 (d, *J* = 2.6 Hz, 1H), 6.60 (dd, *J* = 8.8, 2.6 Hz, 1H), 6.42 (dd, *J* = 15.4, 11.0 Hz, 1H), 6.30 (d, *J* = 10.8 Hz, 1H), 6.20 – 6.13 (m, 1H), 5.23 (dd, *J* = 15.3, 9.0 Hz, 1H), 4.80 (dd, *J* = 11.8, 2.5 Hz, 1H), 4.37 (ddd, *J* = 12.4, 10.5, 2.0 Hz, 1H), 4.00 (s, 3H), 3.56 (d, *J* = 13.0 Hz, 1H), 3.43 (d, *J* = 9.0 Hz, 1H), 3.26 (s, 4H), 3.20 (s, 3H), 3.04 (d, *J* = 9.8 Hz, 1H), 2.85 – 2.73 (m, 1H), 2.49 (dd, *J* = 13.7, 11.8 Hz, 1H), 2.23 (dd, *J* = 13.7, 2.5 Hz, 1H), 1.78 (d, *J* = 13.5 Hz, 1H), 1.69 (s, 3H), 1.54 – 1.43 (m, 2H), 1.31 (d, *J* = 6.4 Hz, 3H), 1.23 (t, *J* = 12.6 Hz, 1H), 0.88 (s, 3H).

Step C: Maytan-3-O-carbamoyl-N-2-chloro-4-(aminophenyl-Cit-Val-Fmoc):

[0453] Using the method of Example 3, step A, the title compound was prepared from the product of the preceding step (**17**, 150 mg, 0.189 mmol) as a white solid (174 mg, 76%). MS (ESI, pos.): calc'd for C₆₁H₇₂Cl₂N₈O₁₄, 1210.5; found 1213.2 (M+H), 1235.3 (M+Na).

Step D: Maytan-3-O-carbamoyl-N-2-chloro-4-(aminophenyl-Cit-Val) (18):

[0454] Using the method of Example 3, step B, the title compound was prepared from the product of the preceding step (172 mg, 0.142 mmol) as a white solid (51 mg, 34%). MS (ESI, pos.): calc'd for C₄₆H₆₂Cl₂N₈O₁₂, 988.4; found 989.3 (M+H), 1011.3 (M+Na).

Step E: Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-Cap-Mal) (19):

[0455] The product of the preceding step (**18**, 14 mg, 0.013 mmol) and 6-maleimidyl-caproic acid succinate ester (**3**, 20 mg, 0.065 mmol) were dissolved in MeCN (2.0 mL) and water (0.5 mL), treated with one drop of saturated aqueous NaHCO₃, the flask sealed with a rubber septum, purged with argon, and the reaction stirred at ambient temperature. After 20 h the reaction was acidified with a few drops of 10% aqueous acetic acid, dissolved in MeCN (ca. 2 mL), and purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 30 – 90% MeCN in water, 0.05% acetic acid in both). The fractions containing product were combined, partially concentrated *in vacuo*, frozen on dry ice, and lyophilized for 4 d giving the title compound as a white solid (10 mg, 63%). MS (ESI, pos.): calc'd for C₅₆H₇₃Cl₂N₉O₁₅, 1181.5; found 1183.1 (M+H), 1204.1 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 10.16 (s, 1H), 8.14 (d, *J* = 7.3 Hz, 1H), 7.93 (dd, *J* = 15.2, 2.3 Hz, 1H), 7.81 (d, *J* = 9.1 Hz, 2H), 7.71 – 7.59 (m, 1H), 7.46 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 1.7 Hz, 1H), 7.07 (s, 1H), 6.99 (d, *J* = 11.8 Hz, 2H), 6.90 (d, *J* = 8.2 Hz, 1H), 6.57 (dd, *J* = 15.1, 11.2 Hz, 1H), 6.45 (d, *J* = 10.6 Hz, 1H), 5.98 (d, *J* = 6.2 Hz, 1H), 5.91 – 5.84 (m, 1H), 5.47 – 5.34 (m, 3H), 4.45 (dd, *J* = 12.0, 2.6 Hz, 1H), 4.40 – 4.29 (m, 1H), 4.19 (dd, *J* = 8.6, 6.8 Hz, 1H), 4.13 (t, *J* = 11.1 Hz, 1H), 3.94 (s, 3H), 3.57 (d, *J* = 12.4 Hz, 1H), 3.48 (d, *J* = 9.1 Hz, 1H), 3.21 (d, *J* = 3.7 Hz, 4H), 3.07 (s, 3H), 3.05 – 2.98 (m, 1H), 2.98 – 2.88 (m, 1H), 2.78 (d, *J* = 9.8 Hz, 1H), 2.43 (t, *J* = 13.1 Hz, 1H), 2.24 – 2.05 (m, 2H), 2.01 (d, *J* = 13.8 Hz, 1H), 1.98 – 1.91 (m, 1H), 1.69 (d, *J* = 7.9 Hz, 1H), 1.65 (s, 3H), 1.61 (dd, *J* = 9.2, 4.9 Hz, 1H), 1.54 – 1.40 (m, 8H), 1.40 – 1.30 (m, 1H), 1.19 (td, *J* = 15.8, 14.8, 8.2 Hz, 3H), 1.11 (d, *J* = 6.4 Hz, 3H), 0.88 (d, *J* = 6.7 Hz, 1H), 0.84 (dd, *J* = 17.0, 6.8 Hz, 6H), 0.81 (s, 2H).

EXAMPLE 5

[0456] Example 5 refers to compounds illustrated in FIG. 13.

Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-adipic Acid Succinate Ester) (20):

Step A: Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-adipic Acid):

[0457] Using the method of Example 3, step C, the title compound was prepared from the product of Example 4, step D (**18**, 19 mg, 0.018 mmol) as a white solid (11 mg, 55%). MS (ESI, pos.): calc'd for C₅₂H₇₀Cl₂N₈O₁₅, 1116.4; found 1119.2 (M+H), 1139.2 (M+Na).

Step B: *Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-adipic Acid Succinate Ester)* (**20**):

[0458] Using the method of Example 2, step B, the title compound was prepared from the product of the preceding step (11 mg, 0.010 mmol) as a white solid (4 mg, 33%). MS (ESI, pos.): calc'd for C₅₆H₇₃Cl₂N₉O₁₇, 1213.5; found 1216.2 (M+H), 1236.3 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 10.16 (s, 1H), 8.16 (d, *J* = 7.3 Hz, 1H), 7.91 (d, *J* = 2.4 Hz, 1H), 7.84 (d, *J* = 8.7 Hz, 1H), 7.78 (s, 1H), 7.65 (d, *J* = 8.7 Hz, 1H), 7.45 (dd, *J* = 8.7, 2.4 Hz, 1H), 7.21 (d, *J* = 1.8 Hz, 1H), 7.06 (s, 1H), 6.88 (s, 1H), 6.64 – 6.50 (m, 1H), 6.44 (d, *J* = 11.0 Hz, 1H), 5.97 (d, *J* = 6.2 Hz, 1H), 5.87 (d, *J* = 1.3 Hz, 1H), 5.41 (q, *J* = 7.6, 6.4 Hz, 3H), 4.50 – 4.39 (m, 1H), 4.34 (d, *J* = 7.0 Hz, 1H), 4.28 – 4.17 (m, 1H), 4.13 (t, *J* = 11.6 Hz, 1H), 3.94 (s, 3H), 3.56 (d, *J* = 13.9 Hz, 1H), 3.52 – 3.42 (m, 1H), 3.21 (s, 3H), 3.07 (s, 3H), 3.03 (dt, *J* = 13.1, 6.7 Hz, 2H), 2.98 – 2.90 (m, 2H), 2.82 – 2.74 (m, 5H), 2.69 – 2.64 (m, 2H), 2.24 (d, *J* = 9.9 Hz, 1H), 2.21 – 2.13 (m, 1H), 2.00 (d, *J* = 13.3 Hz, 1H), 1.98 – 1.93 (m, 1H), 1.64 (s, 3H), 1.60 (d, *J* = 6.5 Hz, 5H), 1.51 (d, *J* = 13.1 Hz, 1H), 1.47 – 1.40 (m, 3H), 1.39 (d, *J* = 6.5 Hz, 1H), 1.11 (d, *J* = 6.4 Hz, 3H), 0.85 (dd, *J* = 16.2, 6.8 Hz, 6H), 0.80 (s, 3H).

EXAMPLE 6

[0459] Example 6 refers to compounds illustrated in FIG. 14.

Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-6-Ahx) (**22**):

Step A: *Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cit-Val-6-Ahx-Fmoc)*:

[0460] The product of Example 4, step D (**18**, 40 mg, 0.040 mmol) and Fmoc-6-aminohexanoic acid succinate ester (**21**, 92 mg, 0.204 mmol) were coupled via the method of

Example 4, step E, giving the title compound as a white solid (31 mg, 57%). MS (ESI, pos.): calc'd for C₆₇H₈₃Cl₂N₉O₁₅, 1323.5; found 1326.5 (M+H), 1346.6 (M+Na).

Step B: *Maytan-3-O-carbamoyl-N-2-chloro-4-(aminophenyl-Cit-Val-6-Ahx)* (22):

[0461] Using the method of Example 3, step B, the title compound was prepared from the product of the preceding step (30 mg, 0.023 mmol) as a white solid (9 mg, 35%) after repurification by preparative HPLC on a Phenomenex Gemini 5u, 30x150mm C18 column (10 – 90% MeCN in water, 0.05% HOAc both, over 22 min, 40 mL/min). MS (ESI, pos.): calc'd for C₅₂H₇₃Cl₂N₉O₁₃, 1101.5; found 1102.8 (M+H). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 7.93 – 7.85 (m, 1H), 7.65 (d, *J* = 8.7 Hz, 1H), 7.49 (d, *J* = 10.2 Hz, 1H), 7.21 (d, *J* = 1.8 Hz, 1H), 7.06 (s, 1H), 6.57 (dd, *J* = 15.1, 11.1 Hz, 1H), 6.44 (d, *J* = 11.2 Hz, 1H), 6.05 (s, 1H), 5.50 – 5.32 (m, 2H), 4.49 – 4.39 (m, 1H), 4.34 (s, 1H), 4.23 – 4.06 (m, 1H), 3.94 (s, 2H), 3.21 (s, 3H), 3.07 (s, 3H), 3.03 – 2.92 (m, 3H), 2.78 (d, *J* = 9.8 Hz, 1H), 2.41 (d, *J* = 12.7 Hz, 1H), 2.30 – 2.23 (m, 1H), 2.18 (q, *J* = 6.7 Hz, 2H), 2.05 – 1.93 (m, 3H), 1.81 (s, 2H), 1.75 – 1.67 (m, 1H), 1.65 (s, 3H), 1.54 – 1.41 (m, 7H), 1.41 – 1.32 (m, 4H), 1.30 – 1.17 (m, 5H), 1.11 (d, *J* = 6.3 Hz, 3H), 0.85 (dd, *J* = 11.5, 6.7 Hz, 6H), 0.80 (s, 2H).

EXAMPLE 7

[0462] Example 7 refers to compounds illustrated in FIG. 15.

Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-Cap-Mal) (24):

[0463] The product of Example 4, step B (**17**, 20 mg, 0.025 mmol) and 6-maleimidyl-caproic acid (**23**, 28 mg, 0.133 mmol) were coupled via the method of Example 1, step E giving the title compound as a white solid (21 mg, 91%) after a second ISCO purification. MS (ESI, pos.): calc'd for C₄₅H₅₃Cl₂N₅O₁₂, 925.3; found 926.2 (M+H), 948.2 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 9.96 (s, 1H), 7.90 (d, *J* = 2.4 Hz, 1H), 7.65 (d, *J* = 8.8 Hz, 1H), 7.60 (s, 1H), 7.43 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 1.8 Hz, 1H), 7.04 (d, *J* = 1.8 Hz, 1H), 6.98 (s, 2H), 6.72 (s, 1H), 6.58 (dd, *J* = 15.2, 11.1 Hz, 1H), 6.42 (d, *J* = 11.0 Hz, 1H), 5.75 (s, 1H), 5.43 (dd, *J* = 15.3, 9.1 Hz, 1H), 4.51 (dd, *J* = 11.9, 2.6 Hz, 1H), 4.18 (ddd, *J* = 12.4, 10.4, 2.3 Hz, 1H), 3.96

(s, 3H), 3.57 (d, $J = 12.4$ Hz, 1H), 3.50 (d, $J = 9.1$ Hz, 1H), 3.42 (t, $J = 7.0$ Hz, 2H), 3.09 (s, 4H), 2.79 (d, $J = 9.8$ Hz, 1H), 2.45 (dd, $J = 13.8, 11.9$ Hz, 1H), 2.31 (t, $J = 7.4$ Hz, 2H), 2.09 – 1.98 (m, 1H), 1.88 (s, 1H), 1.67 (s, 3H), 1.65 – 1.57 (m, 2H), 1.55 (q, $J = 7.2$ Hz, 3H), 1.46 (d, $J = 12.7$ Hz, 2H), 1.34 – 1.21 (m, 2H), 1.14 (d, $J = 6.3$ Hz, 3H), 0.83 (s, 3H).

EXAMPLE 8

[0464] Example 8 refers to compounds illustrated in FIG. 16.

Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-adipic Acid Succinate Ester) (26):

Step A: Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-adipic Acid):

[0465] The product of Example 4, step B (17, 17 mg, 0.021 mmol) and adipic acid (25, 18 mg, 0.123 mmol) were coupled via the method of Example 1, step E giving the title compound as a white solid (11 mg, 59%). MS (ESI, pos.): calc'd for $C_{41}H_{50}Cl_2N_4O_{12}$, 860.3; found 861 (M+H), 883 (M+Na).

Step B: Maytan-3-O-carbamoyl-N-phenyl-2-chloro-4-(amino-adipic Acid Succinate Ester) (26):

[0466] Using the method of Example 2, step B, the title compound was prepared from the product of the preceding step (10 mg, 0.012 mmol) as a white solid (11 mg, 100%). MS (ESI, pos.): calc'd for $C_{45}H_{53}Cl_2N_5O_{14}$, 957.3; found 958.0 (M+H), 980.0 (M+Na). 1H NMR (500 MHz, Chloroform- d) δ : 8.18 (d, $J = 2.4$ Hz, 1H), 8.06 (d, $J = 8.9$ Hz, 1H), 7.78 (s, 1H), 7.08 – 6.93 (m, 2H), 6.85 (d, $J = 1.8$ Hz, 1H), 6.81 (d, $J = 1.8$ Hz, 1H), 6.40 (dd, $J = 15.0, 10.9$ Hz, 1H), 6.33 (d, $J = 11.0$ Hz, 1H), 6.14 (d, $J = 1.4$ Hz, 1H), 6.04 (s, 1H), 5.19 (dd, $J = 15.1, 9.0$ Hz, 1H), 4.81 (dd, $J = 11.8, 2.6$ Hz, 1H), 4.39 – 4.28 (m, 1H), 4.00 (s, 3H), 3.74 – 3.65 (m, 3H), 3.55 (d, $J = 13.0$ Hz, 1H), 3.41 (d, $J = 8.8$ Hz, 1H), 3.28 – 3.21 (m, 4H), 3.19 (s, 3H), 3.05 (d, $J = 9.8$ Hz, 1H), 2.93 – 2.87 (m, 6H), 2.86 (s, 6H), 2.82 (d, $J = 5.4$ Hz, 8H), 2.68 (d, $J = 6.9$ Hz, 2H), 2.51 (dd, $J = 13.9, 11.9$ Hz, 1H), 2.42 (t, $J = 7.2$ Hz, 2H), 2.23 (dd, $J = 13.8, 2.6$ Hz, 1H), 1.88

(d, $J = 5.4$ Hz, 5H), 1.74 (d, $J = 13.7$ Hz, 1H), 1.69 (s, 3H), 1.52 (s, 1H), 1.30 (d, $J = 6.4$ Hz, 3H), 1.25 – 1.18 (m, 1H), 0.87 (s, 3H).

EXAMPLE 9

[0467] Examples 9-25 refer to compounds illustrated in FIG. 17, which is a general scheme for making payloads **30a-q**.

Maytan-3-O-carbamoyl-N-(3-chloro-4-amino)benzene (30a)

Step A: 1-isocyanato-3-chloro-4-nitrobenzene (28a):

[0468] Following the procedure of Cai et al. (*Org. Lett.* **2012**, *14*, 3332-35), triphosgene (301 mg, 1.01 mmol) was dissolved in 2 mL of dry EtOAc, and the reaction cooled to 0 °C under Ar atmosphere. A solution of Et₃N (15 μL, 0.11 mmol) and 3-chloro-4-nitroaniline (**27a**, 173 mg, 1.00 mmol) in 3 mL dry EtOAc was added dropwise over 20 min. The reaction flask was warmed to ambient temperature and stirred for 1h, then topped with a water condenser and heated to 80 °C for 18h. The solvent was evaporated under vacuum and the residue was washed with Et₂O/hexane 1:1 (3 x 10 mL) on a fritted filter. The filtrate was evaporated and dried *in vacuo* to give the title compound as a yellow solid (177 mg, 89%). ¹H NMR (300 MHz, CDCl₃) δ: 7.29 (d, $J = 2.3$ Hz, 1H), 7.14 (d, $J = 2.3$ Hz, 1H), 7.11 (d, $J = 2.3$ Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(3-chloro-4-nitro)benzene (29a):

[0469] Using the method of Example 1, step C, the title compound was prepared from the product of the preceding step (**28a**, 87 mg, 0.438 mmol) and maytansinol (**6**, 103 mg, 0.182 mmol) as a light yellow solid (78 mg, 56%). MS (ESI, pos.): calc'd for C₃₅H₄₀Cl₂N₄O₁₁, 762.2; found 763.1 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(3-chloro-4-amino)benzene (30a):

[0470] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29a**, 77 mg, 0.101 mmol) as a white solid (59 mg, 74%). MS (ESI, pos.): calc'd for $C_{35}H_{42}Cl_2N_4O_9$, 732.2; found 733.1 (M+H). 1H NMR (500 MHz, Chloroform-*d*) δ : 7.43 (s, 1H), 7.13 (d, $J = 8.4$ Hz, 1H), 6.83 (d, $J = 1.8$ Hz, 1H), 6.76 (s, 1H), 6.70 (s, 1H), 6.46 (dd, $J = 15.4, 11.1$ Hz, 2H), 6.22 (s, 1H), 6.07 (d, $J = 10.9$ Hz, 1H), 5.38 (dd, $J = 15.4, 8.7$ Hz, 1H), 4.78 (d, $J = 11.5$ Hz, 1H), 4.34 (t, $J = 11.5$ Hz, 1H), 3.99 (s, 3H), 3.98 (s, 1H), 3.53 – 3.44 (m, 2H), 3.32 (s, 3H), 3.26 (d, $J = 14.5$ Hz, 1H), 3.21 (s, 3H), 2.98 (s, 1H), 2.90 (d, $J = 9.1$ Hz, 1H), 2.50 (t, $J = 12.8$ Hz, 1H), 2.26 (d, $J = 13.9$ Hz, 1H), 1.87 (d, $J = 13.6$ Hz, 1H), 1.70 (s, 3H), 1.50 (dd, $J = 10.8, 4.4$ Hz, 2H), 1.30 (d, $J = 6.3$ Hz, 3H), 1.24 (t, $J = 12.7$ Hz, 1H), 0.90 (s, 3H).

EXAMPLE 10

Maytan-3-O-carbamoyl-N-(4-chloro-2-amino)benzene (30b)

Step A: 1-isocyanato-4-chloro-2-nitrobenzene (28b):

[0471] Using the method of Example 9, step A, the title compound was prepared from 4-chloro-2-nitroaniline (**27b**, 172 mg, 0.997 mmol) as a yellow solid (198 mg, 100%). 1H NMR (300 MHz, Chloroform-*d*) δ : 8.14 (d, $J = 2.5$ Hz, 1H), 7.57 (dd, $J = 8.6, 2.5$ Hz, 1H), 7.21 (d, $J = 8.6$ Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(4-chloro-2-nitro)benzene (29b):

[0472] Using the method of Example 4, step A, the title compound was prepared from the product of the preceding step (**28b**, 98 mg, 0.493 mmol) and maytansinol (**6**, 54 mg, 0.096 mmol) as an orange solid (46 mg, 63%). MS (ESI, pos.): calc'd for $C_{35}H_{40}Cl_2N_4O_{11}$, 762.2; found 763.5 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-chloro-2-amino)benzene (30b):

[0473] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29b**, 44 mg, 0.058 mmol) as a white solid (34 mg, 74%). MS

(ESI, pos.): calc'd for $C_{35}H_{42}Cl_2N_4O_9$, 732.2; found 733.5 (M+H). 1H NMR (500 MHz, Chloroform-*d*) δ : 7.43 (s, 1H), 7.13 (d, $J = 8.4$ Hz, 1H), 6.83 (d, $J = 1.8$ Hz, 1H), 6.76 (s, 1H), 6.70 (s, 1H), 6.46 (dd, $J = 15.4, 11.1$ Hz, 2H), 6.22 (s, 1H), 6.07 (d, $J = 10.9$ Hz, 1H), 5.38 (dd, $J = 15.4, 8.7$ Hz, 1H), 4.78 (d, $J = 11.5$ Hz, 1H), 4.34 (t, $J = 11.5$ Hz, 1H), 3.99 (s, 3H), 3.98 (s, 1H), 3.54 – 3.44 (m, 2H), 3.32 (s, 3H), 3.26 (d, $J = 14.5$ Hz, 1H), 3.21 (s, 3H), 2.98 (s, 1H), 2.90 (d, $J = 9.1$ Hz, 1H), 2.50 (t, $J = 12.8$ Hz, 1H), 2.26 (d, $J = 13.9$ Hz, 1H), 1.87 (d, $J = 13.6$ Hz, 1H), 1.70 (s, 3H), 1.50 (dd, $J = 10.8, 4.4$ Hz, 2H), 1.30 (d, $J = 6.3$ Hz, 3H), 1.24 (t, $J = 12.7$ Hz, 1H), 0.90 (s, 3H).

EXAMPLE 11

Maytan-3-O-carbamoyl-N-(2-methyl-4-amino)benzene (30c)

Step A: Maytan-3-O-carbamoyl-N-(2-methyl-4-nitro)benzene (29c):

[0474] The title compound was prepared from 2-methyl-4-nitrophenyl isocyanate (**28c**, 41 mg, 0.230 mmol) and maytansinol (**6**, 53 mg, 0.094 mmol) as a yellow solid (10 mg, 14%), using the method of Example 4, step A, except that it was purified on a C18 Aq RediSep Gold column via Teledyne ISCO purification system (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both). MS (ESI, pos.): calc'd for $C_{36}H_{43}ClN_4O_{11}$, 742.3; found 743.2 (M+H).

Step B: Maytan-3-O-carbamoyl-N-(2-methyl-4-amino)benzene (30c):

[0475] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29c**, 10 mg, 0.014 mmol) as a cream-colored solid (6 mg, 60%). MS (ESI, pos.): calc'd for $C_{35}H_{42}Cl_2N_4O_9$, 712.3; found 713.2 (M+H). 1H NMR (500 MHz, Chloroform-*d*) δ : 7.33 – 7.30 (m, 1H), 6.84 (d, $J = 1.8$ Hz, 1H), 6.79 (s, 1H), 6.55 (d, $J = 2.6$ Hz, 1H), 6.52 (dd, $J = 8.4, 2.7$ Hz, 1H), 6.47 (dd, $J = 15.5, 10.9$ Hz, 1H), 6.24 (s, 1H), 6.22 – 6.15 (m, 1H), 6.10 (d, $J = 10.9$ Hz, 1H), 5.46 (dd, $J = 15.5, 8.9$ Hz, 1H), 4.80 (dd, $J = 11.7, 2.4$ Hz, 1H), 4.42 – 4.29 (m, 1H), 4.00 (s, 3H), 3.53 – 3.41 (m, 2H), 3.32 (s, 3H), 3.24 (d, $J = 13.0$ Hz, 1H), 3.19 (s, 3H), 3.14 (s, 1H), 2.89 (d, $J = 9.8$ Hz, 1H), 2.49 (dd, $J = 13.7, 11.7$ Hz, 1H), 2.25

(s, 3H), 2.22 (d, $J = 2.4$ Hz, 1H), 1.94 (d, $J = 13.6$ Hz, 1H), 1.70 (s, 3H), 1.51 – 1.45 (m, 1H), 1.34 – 1.21 (m, 5H), 0.91 (s, 3H).

EXAMPLE 12

Maytan-3-O-carbamoyl-N-(2-methyl-3-amino)benzene (30d)

Step A: 1-isocyanato-2-methyl-3-nitrobenzene (28d):

[0476] Using the method of Example 9, step A, the title compound was prepared from 2-methyl-3-nitroaniline (**27d**, 153 mg, 1.01 mmol) as a pale yellow solid (178 mg, 99%). ^1H NMR (300 MHz, Chloroform- d) δ : 7.68 (dd, $J = 7.5, 1.8$ Hz, 1H), 7.43 – 7.26 (m, 2H), 2.49 (s, 3H).

Step B: Maytan-3-O-carbamoyl-N-(2-methyl-3-nitro)benzene (29d):

[0477] Using the method of Example 4, step A, the title compound was prepared from the product of the preceding step (**28d**, 70 mg, 0.393 mmol) and maytansinol (**6**, 73 mg, 0.129 mmol) as a yellow solid (41 mg, 43%). MS (ESI, pos.): calc'd for $\text{C}_{36}\text{H}_{43}\text{ClN}_4\text{O}_{11}$, 742.3; found 743.5 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(2-methyl-3-amino)benzene (30d):

[0478] The title compound was prepared from the product of the preceding step (**29d**, 44 mg, 0.058 mmol) as a white solid (12 mg, 29%) using the method of Example 4, step B, except that it had to be repurified by preparative HPLC on a Phenomenex Gemini 5u, 30x150mm C18 column (20 – 80% MeCN in water, 0.05% HOAc both, over 15 min, 50 mL/min). MS (ESI, pos.): calc'd for $\text{C}_{36}\text{H}_{45}\text{ClN}_4\text{O}_9$, 712.3; found 713.5 (M+H). ^1H NMR (500 MHz, DMSO- d_6) δ : 7.92 (s, 1H), 7.19 (d, $J = 1.9$ Hz, 1H), 7.15 (s, 1H), 7.00 (s, 1H), 6.90 (s, 2H), 6.63 – 6.53 (m, 1H), 6.49 (d, $J = 11.1$ Hz, 1H), 5.54 (dd, $J = 15.0, 9.1$ Hz, 1H), 4.48 (dd, $J = 12.1, 2.7$ Hz, 1H), 4.13 (t, $J = 11.3$ Hz, 1H), 3.94 (s, 3H), 3.58 (d, $J = 12.2$ Hz, 1H), 3.25 (s, 3H), 3.16 (d, $J = 12.1$ Hz, 2H), 3.09 (s, 3H), 2.73 (d, $J = 9.7$ Hz, 1H), 2.45 – 2.38 (m, 1H), 2.06 – 1.94 (m, 5H), 1.66 (s, 3H), 1.56 (d, $J = 13.3$ Hz, 1H), 1.44 (d, $J = 12.2$ Hz, 2H), 1.10 (d, $J = 6.3$ Hz, 3H), 0.79 (s, 3H).

EXAMPLE 13Maytan-3-O-carbamoyl-N-(6-methyl-3-amino)benzene (30e)Step A: 1-isocyanato-6-methyl-3-nitrobenzene (28e):

[0479] Using the method of Example 9, step A, the title compound was prepared from 6-methyl-3-nitroaniline (**27e**, 1.00 mmol) as a yellow solid (100 mg, 56% yield). ¹H NMR (500 MHz, Chloroform-*d*) δ: 7.98 (d, 1H, *J* = 2.5 Hz), 7.95 (dd, 1H, *J* = 2.5, 8.5 Hz), 7.36 (d, 1H, *J* = 8.5 Hz).

Step B: Maytan-3-O-carbamoyl-N-(6-methyl-3-nitro)benzene (29e):

[0480] The title compound was prepared from the product of the preceding step (**28e**, 0.422 mmol) and maytansinol (**6**, 100 mg, 0.176 mmol) as a yellow solid (21 mg, 16% yield) using the method of Example 1, step C, except that it was purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both). MS (ESI, pos.): calc'd for C₃₆H₄₃ClN₄O₁₁, 742.3; found 725.2 (M+H-H₂O), 765.2 (M+Na).

Step C: Maytan-3-O-carbamoyl-N-(6-methyl-3-amino)benzene (30e):

[0481] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29e**, 21 mg, 0.028 mmol) as a white solid (6.0 mg, 29% yield). MS (ESI, pos.): calc'd for C₃₆H₄₅ClN₄O₉, 712.3, found 713.3 (M+H), 695.3 (M+H-H₂O). ¹H NMR (500 MHz, CDCl₃): δ 7.25-7.22 (m, 1H), 6.96 (d, 1H, *J* = 7.5 Hz), 6.84 (s, 1H), 6.77 (s, 1H), 6.45-6.40 (m, 1H), 6.26 (s, 1H), 6.10 (d, 1H, *J* = 11.0 Hz), 5.35 (dd, 1H, *J* = 8.5, 11.7 Hz), 4.77 (dd, 1H, *J* = 2.5, 11.7 Hz), 4.37 (dd, 1H, *J* = 10.5, 10.5 Hz), 3.99 (s, 3H), 3.48 (d, 1H, *J* = 13.5 Hz), 3.45 (d, 1H, *J* = 9.0 Hz), 3.28 (s, 3H), 3.24 (d, 1H, *J* = 13.5 Hz), 3.19 (s, 3H), 2.97 (d, 1H, *J* = 10.0 Hz), 2.50 (dd, 1H, *J* = 11.5, 11.5 Hz), 2.28-2.22 (m, 4H), 1.86 (d, 1H, *J* = 13.5 Hz), 1.70 (s, 3H), 1.53-1.49 (m, 1H), 1.31 (s, 3H), 1.27-1.22 (m, 1H), 0.89 (s, 3H).

EXAMPLE 14Maytan-3-O-carbamoyl-N-(6-methyl-2-amino)benzene (30f)Step A: 1-isocyanato-6-methyl-2-nitrobenzene (28f):

[0482] Using the method of Example 9, step A, the title compound was prepared from 6-methyl-2-nitroaniline (**27f**, 1.00 mmol) as a yellow solid (135 mg, 76% yield). ¹H NMR (500 MHz, CDCl₃): δ 7.92 (d, 1H, *J* = 8.5 Hz), 7.49 (d, 1H, *J* = 8.5 Hz), 7.22 (dd, 1H, *J* = 8.5, 8.5 Hz), 2.43 (s, 3H).

Step B: Maytan-3-O-carbamoyl-N-(6-methyl-2-nitro)benzene (29f):

[0483] Using the method of Example 13, step B, the title compound was prepared from the product of the preceding step (**28f**, 0.422 mmol) and maytansinol (**6**, 100 mg, 0.176 mmol) as a yellow solid (33 mg, 25% yield). MS (ESI, pos.): calc'd for C₃₆H₄₃ClN₄O₁₁, 742.3; found 743.2 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(6-methyl-2-amino)benzene (30f):

[0484] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29f**, 33 mg, 0.044 mmol) as a yellow solid (15.0 mg, 53% yield). MS (ESI, pos.): calc'd for C₃₆H₄₅ClN₄O₉, 712.3, found 713.30 (M+H). ¹H NMR (500 MHz, CDCl₃): δ 7.01- 6.98 (m, 1H), 6.83 (s, 1H), 6.78 (s, 1H), 6.67-6.62 (m, 2H), 6.50-6.45 (m, 1H), 6.35 (bs, 1H), 6.16 (d, 1H, *J* = 11.0 Hz), 5.98 (s, 1H), 5.63 (dd, 1H, *J* = 8.5, 14.7 Hz), 4.96 (d, 1H, *J* = 11.5 Hz), 4.28 (dd, 1H, *J* = 10.5, 10.5 Hz), 3.98 (s, 3H), 3.50-3.45 (m, 2H), 3.32 (s, 3H), 3.21 (d, 1H, *J* = 13.5 Hz), 3.18 (s, 3H), 2.81 (d, 1H, *J* = 10.0 Hz), 2.54 (dd, 1H, *J* = 13.5, 13.5 Hz), 2.25-2.23 (m, 4H), 1.90 (d, 1H, *J* = 12.5 Hz), 1.70 (s, 3H), 1.52-1.48 (m, 1H), 1.35-1.27 (m, 4H), 0.92 (s, 3H).

EXAMPLE 15*Maytan-3-O-carbamoyl-N-(4-fluoro-2-amino)benzene (30g)**Step A: 1-isocyanato-4-fluoro-2-nitrobenzene (28g):*

[0485] Using the method of Example 9, step A, the title compound was prepared from 4-fluoro-2-nitroaniline (**27g**, 1.00 mmol) as an orange solid (223 mg, 81% yield). ¹H NMR (300 MHz, CDCl₃): δ 7.88 (dd, 1H, *J* = 3.0, 7.5 Hz), 7.33 (dd, 1H, *J* = 3.0, 7.5 Hz), 7.25 (dd, 1H, *J* = 4.0, 9.0 Hz).

Step B: Maytan-3-O-carbamoyl-N-(4-fluoro-2-nitro)benzene (29g):

[0486] Using the method of Example 13, step B, the title compound was prepared from the product of the preceding step (**28g**, 0.422 mmol) and maytansinol (**6**, 10 mg, 0.176 mmol) as a yellow solid (16 mg, 12% yield). MS (ESI, pos.): calc'd for C₃₅H₄₀ClFN₄O₁₁, 746.2; found 747.2 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-fluoro-2-amino)benzene (30g):

[0487] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29g**, 16 mg, 0.021 mmol) as a yellow solid (4.0 mg, 26% yield). MS (ESI, pos.): calc'd for C₃₅H₄₂ClFN₄O₉, 716.3; found 717.2 (M+H). ¹H NMR (500 MHz, CDCl₃): δ 7.25-7.21 (m, 1H), 6.83 (s, 1H), 6.73 (bs, 1H), 6.52 (dd, 1H, *J* = 2.5, 8.5 Hz), 6.50-6.48 (2H, m), 6.39 (s, 1H), 6.27 (s, 1H), 6.11 (1H, bs), 5.52 (dd, 1H, *J* = 8.5, 15.7 Hz), 4.87 (dd, 1H, *J* = 2.5, 11.5 Hz), 4.27 (dd, 1H, *J* = 11.5, 11.5 Hz), 3.99 (s, 3H), 3.49 (d, 1H, *J* = 8.5 Hz), 3.45 (bs, 1H), 3.34 (s, 3H), 3.21 (s, 3H), 3.18 (bs, 1H), 2.84-2.83 (m, 1H), 2.52 (dd, 1H, *J* = 11.5, 13.5 Hz), 2.28 (d, 1H, *J* = 13.5 Hz), 1.86 (d, 1H, *J* = 13.5 Hz), 1.68 (s, 3H), 1.51-1.45 (m, 1H), 1.27 (d, 3H, *J* = 6.5 Hz), 1.27-1.23 (m, 1H), 0.89 (s, 3H).

EXAMPLE 16*Maytan-3-O-carbamoyl-N-(2-fluoro-5-amino)benzene (30h)**Step A: 1-isocyanato-2-fluoro-5-nitrobenzene (28h):*

[0488] Using the method of Example 9, step A, the title compound was prepared from 2-fluoro-5-nitroaniline (**27h**, 155 mg, 0.99 mmol) as an orange solid (168 mg, 93% yield). ¹H NMR (300 MHz, Chloroform-*d*) δ: 8.10 (dd, *J* = 8.7, 4.1 Hz, 1H), 8.01 (dd, *J* = 6.7, 2.8 Hz, 1H), 7.31 (t, *J* = 8.8 Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(2-fluoro-5-nitro)benzene (29h):

[0489] Using the method of Example 13, step B, the title compound was prepared from the product of the preceding step (**28h**, 61 mg, 0.335 mmol) and maytansinol (**6**, 67 mg, 0.119 mmol) as an orange solid (21 mg, 24% yield). MS (ESI, pos.): calc'd for C₃₅H₄₀ClFN₄O₁₁, 746.2; found 747.5 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(2-fluoro-5-amino)benzene (30h):

[0490] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29h**, 20 mg, 0.027 mmol) as a white solid (6 mg, 29% yield) after a second reverse-phase ISCO purification (gradient elution: 30 – 90% MeCN in water, 0.05% acetic acid in both). MS (ESI, pos.): calc'd for C₃₅H₄₂ClFN₄O₉, 716.3; found 717.5 (M+H). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 7.70 (s, 1H), 7.19 (d, *J* = 1.9 Hz, 1H), 7.07 (s, 1H), 7.03 (dd, *J* = 7.0, 2.8 Hz, 1H), 6.90 (dd, *J* = 10.6, 8.7 Hz, 1H), 6.81 (s, 1H), 6.56 (dd, *J* = 15.3, 11.1 Hz, 1H), 6.38 (d, *J* = 11.1 Hz, 1H), 6.32 (dt, *J* = 8.7, 3.3 Hz, 1H), 5.59 (s, 1H), 5.49 (dd, *J* = 15.2, 9.1 Hz, 1H), 4.90 (s, 2H), 4.50 (dd, *J* = 11.7, 2.5 Hz, 1H), 4.14 (t, *J* = 11.2 Hz, 1H), 3.94 (s, 3H), 3.56 (d, *J* = 12.4 Hz, 1H), 3.48 (d, *J* = 9.1 Hz, 1H), 3.23 (s, 3H), 3.07 (s, 3H), 2.79 (d, *J* = 9.7 Hz, 1H), 2.44 (dd, *J* = 14.0, 11.9 Hz, 2H), 2.03 (d, *J* = 13.7 Hz, 1H), 1.66 (s, 3H), 1.52 (d, *J* = 13.5 Hz, 1H), 1.44 (dd, *J* = 18.3, 10.2 Hz, 2H), 1.12 (d, *J* = 6.3 Hz, 3H), 0.80 (s, 3H).

EXAMPLE 17Maytan-3-O-carbamoyl-N-(4-methoxy-2-amino)benzene (30i)Step A: 1-isocyanato-4-methoxy-2-nitrobenzene (28i):

[0491] Using the method of Example 9, step A, the title compound was prepared from 4-methoxy-2-nitroaniline (**27i**, 1.00 mmol) as a yellow solid (100 mg, 51% yield). ¹H NMR (300 MHz, CDCl₃): δ 8.10 (dd, 1H, *J* = 3.0, 8.7 Hz), 7.90 (d, 1H, *J* = 3.0 Hz), 6.96 (d, 1H, *J* = 8.7 Hz), 4.06 (s, 3H).

Step B: Maytan-3-O-carbamoyl-N-(4-methoxy-2-nitro)benzene (29i):

[0492] Using the method of Example 13, step B, the title compound was prepared from the product of the preceding step (**28i**, 0.422 mmol) and maytansinol (**6**, 100 mg, 0.176 mmol) as a yellow solid (25 mg, 19% yield). MS (ESI, pos.): calc'd for C₃₆H₄₃ClN₄O₁₂, 758.3; found 759.2 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-methoxy-2-amino)benzene (30i):

[0493] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29i**, 25 mg, 0.033 mmol) as a yellow solid (2.0 mg, 9% yield). MS (ESI, pos.): calc'd for C₃₆H₄₅ClN₄O₁₀, 728.3; found 729.2 (M+H). ¹H NMR (500 MHz, CDCl₃): δ 7.14 (d, 1H, *J* = 8.5 Hz), 6.83 (s, 1H), 6.49-6.38 (m, 2H), 6.30 (s, 1H), 6.20 (bs, 1H), 5.55-5.51 (m, 1H), 4.89 (d, 1H, *J* = 11 Hz), 4.30 (dd, 1H, *J* = 11.0, 11.0 Hz), 3.99 (s, 3H), 3.76 (s, 3H), 3.49-3.48 (m, 1H), 3.33 (s, 3H), 3.20-3.17 (m, 1H), 3.21 (s, 3H), 2.84 (m, 1H), 2.54-2.49 (m, 1H), 2.25 (d, 1H, *J* = 13.5 Hz), 1.90 (d, 1H, *J* = 13.5 Hz), 1.69 (s, 3H), 1.69-1.57 (m, 2H), 1.29 (s, 3H), 0.91 (s, 3H).

EXAMPLE 18Maytan-3-O-carbamoyl-N-(3-methoxy-4-amino)benzene (30j)

Step A: 1-isocyanato-3-methoxy-4-nitrobenzene (28j):

[0494] Using the method of Example 9, step A, the title compound was prepared from 3-methoxy-4-nitroaniline (**27j**, 169 mg, 1.00 mmol) as a yellow solid (102 mg, 52% yield). ¹H NMR (300 MHz, Chloroform-*d*) δ: 8.01 – 7.81 (m, 1H), 6.85 – 6.65 (m, 2H), 3.96 (s, 4H).

Step B: Maytan-3-O-carbamoyl-N-(3-methoxy-4-nitro)benzene (29j):

[0495] The title compound was prepared from the product of the preceding step (**28j**, 100 mg, 0.515 mmol) and maytansinol (**6**, 71 mg, 0.126 mmol) as a yellow solid (64 mg, 67%) using the method of Example 4, step A, except that the reaction was run at ambient temperature for 18 h. MS (ESI, pos.): calc'd for C₃₆H₄₃ClN₄O₁₂, 758.3; found 759.5 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(3-methoxy-4-amino)benzene (30j):

[0496] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29j**, 63 mg, 0.083 mmol) as a white solid (37 mg, 57%). MS (ESI, pos.): calc'd for C₃₆H₄₅ClN₄O₁₀, 728.3; found 729.5 (M+H). ¹H NMR (500 MHz, Chloroform-*d*) δ: 6.84 (d, *J* = 1.8 Hz, 1H), 6.71 (s, 1H), 6.66 (d, *J* = 8.2 Hz, 1H), 6.56 (dd, *J* = 8.2, 2.3 Hz, 1H), 6.52 – 6.38 (m, 2H), 6.22 (s, 1H), 6.13 (d, *J* = 10.9 Hz, 1H), 5.41 (dd, *J* = 15.5, 8.6 Hz, 1H), 4.78 (d, *J* = 11.3 Hz, 1H), 4.33 (t, *J* = 11.2 Hz, 1H), 3.99 (s, 3H), 3.87 (s, 3H), 3.55 – 3.39 (m, 2H), 3.31 (s, 3H), 3.27 (d, *J* = 13.3 Hz, 1H), 3.19 (s, 3H), 2.90 (d, *J* = 9.8 Hz, 1H), 2.49 (dd, *J* = 13.9, 11.6 Hz, 1H), 2.25 (d, *J* = 14.0 Hz, 1H), 2.11 (s, 2H), 1.87 (d, *J* = 13.4 Hz, 1H), 1.70 (s, 3H), 1.30 (d, *J* = 6.2 Hz, 3H), 1.25 (dd, *J* = 15.1, 10.5 Hz, 1H), 0.90 (s, 3H).

EXAMPLE 19

Maytan-3-O-carbamoyl-N-(2-methoxy-4-amino)benzene (30k)

Step A: Maytan-3-O-carbamoyl-N-(2-methoxy-4-nitro)benzene (29k):

[0497] The title compound was prepared from 1-isocyanato-2-methoxy-4-nitrobenzene (**28k**, 62 mg, 0.319 mmol) and maytansinol (**6**, 53 mg, 0.094 mmol) as a pale yellow solid (25 mg, 35%) using the method of Example 4, step A, except that the reaction was purified on a C18 Aq RediSep Gold column via ISCO (gradient elution: 20 – 80% MeCN in water, 0.05% acetic acid in both). MS (ESI, pos.): calc'd for C₃₆H₄₃ClN₄O₁₂, 758.3; found 759.5 (M+H).

Step B: Maytan-3-O-carbamoyl-N-(2-methoxy-4-amino)benzene (30k):

[0498] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29k**, 24 mg, 0.032 mmol) as a white solid (9 mg, 36%). MS (ESI, pos.): calc'd for C₃₆H₄₅ClN₄O₁₀, 728.3; found 729.5 (M+H), 751.5 (M+Na). ¹H NMR (500 MHz, Chloroform-*d*) δ: 7.83 (d, *J* = 8.3 Hz, 1H), 6.90 (s, 1H), 6.86 (d, *J* = 1.9 Hz, 1H), 6.82 (d, *J* = 1.8 Hz, 1H), 6.42 (dd, *J* = 15.1, 10.9 Hz, 1H), 6.36 – 6.24 (m, 2H), 6.15 (s, 1H), 5.31 (dd, *J* = 15.1, 9.1 Hz, 1H), 4.79 (dd, *J* = 11.8, 2.4 Hz, 1H), 4.41 – 4.29 (m, 1H), 4.00 (s, 3H), 3.94 (s, 3H), 3.52 (d, *J* = 12.9 Hz, 1H), 3.41 (d, *J* = 9.1 Hz, 1H), 3.29 (d, *J* = 12.7 Hz, 1H), 3.25 (s, 3H), 3.19 (s, 3H), 3.02 (d, *J* = 9.9 Hz, 1H), 2.73 (s, 1H), 2.47 (dd, *J* = 13.8, 11.8 Hz, 1H), 2.22 (dd, *J* = 13.7, 2.4 Hz, 1H), 2.13 (s, 2H), 1.78 (d, *J* = 13.6 Hz, 1H), 1.70 (s, 3H), 1.50 (dd, *J* = 10.2, 6.4 Hz, 1H), 1.30 (d, *J* = 6.4 Hz, 3H), 1.20 (t, *J* = 12.9 Hz, 1H), 0.86 (s, 3H).

EXAMPLE 20

Maytan-3-O-carbamoyl-N-(3-trifluoromethyl-4-amino)benzene (30l)

Step A: 1-isocyanato-3-trifluoromethyl-4-nitrobenzene (28l):

[0499] Using the method of Example 9, step A, the title compound was prepared from 3-trifluoromethyl-4-nitroaniline (**27l**, 103 mg, 0.500 mmol) as a yellow oily solid (63 mg, 54% yield). ¹H NMR (300 MHz, Chloroform-*d*) δ: 7.95 (d, *J* = 8.7 Hz, 1H), 7.53 (dt, *J* = 2.4, 0.5 Hz, 1H), 7.41 (ddd, *J* = 8.7, 2.3, 0.5 Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(3-trifluoromethyl-4-nitro)benzene (29l):

[0500] The title compound was prepared from the product of the preceding step (**28l**, 59 mg, 0.254 mmol) and maytansinol (**6**, 57 mg, 0.101 mmol) as a yellow solid (27 mg, 34%) using the method of Example 19, step A. MS (ESI, pos.): calc'd for $C_{36}H_{40}ClF_3N_4O_{11}$, 796.2; found 797.4 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(3-trifluoromethyl-4-amino)benzene (**30l**):

[0501] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29l**, 25 mg, 0.031 mmol) as a white solid (16 mg, 62%) after a second reverse-phase ISCO purification. MS (ESI, pos.): calc'd for $C_{36}H_{42}ClF_3N_4O_9$, 766.3; found 767.5 (M+H). 1H NMR (500 MHz, Chloroform-*d*) δ : 7.47 (dd, $J = 8.7, 2.6$ Hz, 1H), 7.40 (d, $J = 2.5$ Hz, 1H), 6.84 (d, $J = 1.9$ Hz, 1H), 6.73 (d, $J = 8.7$ Hz, 1H), 6.71 – 6.61 (m, 1H), 6.50 (d, $J = 2.3$ Hz, 1H), 6.49 – 6.42 (m, 1H), 6.22 (s, 1H), 6.06 (d, $J = 10.9$ Hz, 1H), 5.37 (dd, $J = 15.5, 8.5$ Hz, 1H), 4.79 (dd, $J = 11.5, 2.2$ Hz, 1H), 4.32 (ddd, $J = 12.2, 10.5, 1.9$ Hz, 1H), 4.11 (s, 2H), 3.99 (s, 3H), 3.54 – 3.44 (m, 2H), 3.32 (s, 3H), 3.26 (d, $J = 13.5$ Hz, 1H), 3.20 (s, 3H), 2.97 (s, 1H), 2.88 (d, $J = 9.8$ Hz, 1H), 2.50 (dd, $J = 14.0, 11.6$ Hz, 1H), 2.27 (dd, $J = 14.0, 2.3$ Hz, 1H), 1.86 (d, $J = 13.5$ Hz, 1H), 1.71 (s, 3H), 1.50 (td, $J = 10.2, 6.3$ Hz, 1H), 1.29 (d, $J = 6.4$ Hz, 3H), 1.25 (t, $J = 12.6$ Hz, 1H), 0.90 (s, 3H).

EXAMPLE 21

Maytan-3-O-carbamoyl-N-(2-bromo-4-amino)benzene (**30m**)

Step A: 1-isocyanato-2-bromo-4-nitrobenzene (**28m**):

[0502] Using the method of Example 9, step A, the title compound was prepared from 2-bromo-4-nitroaniline (**27m**, 108 mg, 0.498 mmol) as a yellow solid (109 mg, 90% yield). 1H NMR (300 MHz, Chloroform-*d*) δ : 8.49 (d, $J = 2.5$ Hz, 1H), 8.16 (dd, $J = 8.8, 2.6$ Hz, 1H), 7.28 (dd, $J = 8.8, 0.3$ Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(2-bromo-4-nitro)benzene (29m):

[0503] The title compound was prepared from the product of the preceding step (28m, 55 mg, 0.226 mmol) and maytansinol (6, 46 mg, 0.081 mmol) as a yellow solid (30 mg, 45%) using the method of Example 19, step A. MS (ESI, pos.): calc'd for C₃₅H₄₀BrClN₄O₁₁, 806.2; found 809.0 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(2-bromo-4-amino)benzene (30m):

[0504] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (29m, 28 mg, 0.035 mmol) as a white solid (10 mg, 34%). MS (ESI, pos.): calc'd for C₃₅H₄₂BrClN₄O₉, 776.2; found 779.0 (M+H). ¹H NMR (300 MHz, Chloroform-*d*) δ: 7.79 (d, *J* = 8.7 Hz, 1H), 6.94 (d, *J* = 1.8 Hz, 1H), 6.90 (d, *J* = 2.5 Hz, 1H), 6.85 (d, *J* = 1.8 Hz, 1H), 6.77 (s, 1H), 6.64 (dd, *J* = 8.8, 2.6 Hz, 1H), 6.50 – 6.29 (m, 2H), 6.19 (s, 1H), 5.22 (dd, *J* = 14.6, 9.0 Hz, 1H), 4.79 (dd, *J* = 11.8, 2.6 Hz, 1H), 4.38 (t, *J* = 11.1 Hz, 1H), 4.00 (s, 3H), 3.69 (s, 1H), 3.60 (d, *J* = 12.9 Hz, 1H), 3.43 (d, *J* = 9.0 Hz, 1H), 3.26 (s, 4H), 3.20 (s, 4H), 3.05 (d, *J* = 9.8 Hz, 1H), 2.89 (s, 1H), 2.49 (dd, *J* = 13.7, 11.9 Hz, 1H), 2.22 (dd, *J* = 13.7, 2.6 Hz, 1H), 1.79 (d, *J* = 13.5 Hz, 1H), 1.69 (s, 4H), 1.49 (dd, *J* = 10.3, 6.2 Hz, 1H), 1.31 (d, *J* = 6.3 Hz, 3H), 1.28 – 1.16 (m, 2H), 0.87 (s, 3H).

EXAMPLE 22

Maytan-3-O-carbamoyl-N-(4-trifluoromethoxy-2-amino)benzene (30n)

Step A: 1-isocyanato-4-trifluoromethoxy-2-nitrobenzene (28n):

[0505] Using the method of Example 9, step A, the title compound was prepared from 4-trifluoromethoxy-2-nitroaniline (27n, 223 mg, 1.00 mmol) as an orange oily solid (228 mg, 92% yield). ¹H NMR (300 MHz, Chloroform-*d*) δ: 8.03 (d, *J* = 2.8 Hz, 1H), 7.52 – 7.44 (m, 1H), 7.32 (dt, *J* = 9.0, 0.6 Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(4-trifluoromethoxy-2-nitro)benzene (29n):

[0506] The title compound was prepared from the product of the preceding step (**28n**, 215 mg, 0.867 mmol) and maytansinol (**6**, 84 mg, 0.149 mmol) as a yellow solid (87 mg, 72%) using the method of Example 4, step A. MS (ESI, pos.): calc'd for $C_{36}H_{40}ClF_3N_4O_{12}$, 812.2; found 813.5 (M+H), 835.4 (M+Na).

Step C: Maytan-3-O-carbamoyl-N-(4-trifluoromethoxy-2-amino)benzene (30n):

[0507] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29n**, 85 mg, 0.105 mmol) as a white solid (59 mg, 67%) after a second reverse-phase ISCO purification. MS (ESI, pos.): calc'd for $C_{36}H_{42}ClF_3N_4O_{10}$, 782.3; found 783.5 (M+H), 805.5 (M+Na). 1H NMR (500 MHz, Chloroform-*d*) δ : 7.31 (d, $J = 8.8$ Hz, 1H), 6.81 (s, 1H), 6.70 (s, 1H), 6.63 (m, 2H), 6.43 (dd, $J = 15.4, 10.9$ Hz, 1H), 6.09 (s, 1H), 5.58 (dd, $J = 15.4, 8.8$ Hz, 1H), 4.88 – 4.77 (m, 1H), 4.28 (t, $J = 11.2$ Hz, 1H), 3.96 (s, 3H), 3.44 (d, $J = 8.7$ Hz, 2H), 3.39 (p, $J = 1.7$ Hz, 2H), 3.33 (s, 3H), 3.19 (s, 3H), 3.15 (d, $J = 13.1$ Hz, 1H), 2.84 (d, $J = 9.6$ Hz, 1H), 2.57 – 2.47 (m, 1H), 2.28 – 2.22 (m, 1H), 1.84 (d, $J = 13.7$ Hz, 1H), 1.65 (s, 3H), 1.54 – 1.43 (m, 1H), 1.30 (d, $J = 12.6$ Hz, 1H), 1.26 (d, $J = 6.2$ Hz, 3H), 0.87 (s, 3H).

EXAMPLE 23

Maytan-3-O-carbamoyl-N-(4-fluoro-5-chloro-2-amino)benzene (30o)

Step A: 1-isocyanato-4-fluoro-5-chloro-2-nitrobenzene (28o):

[0508] Using the method of Example 9, step A, the title compound was prepared from 4-fluoro-5-chloro-2-nitroaniline (**27o**, 70 mg, 0.367 mmol) as a yellow oil (72 mg, 91% yield). 1H NMR (300 MHz, Chloroform-*d*) δ : 8.01 (d, $J = 8.3$ Hz, 1H), 7.35 (d, $J = 6.5$ Hz, 1H).

Step B: Maytan-3-O-carbamoyl-N-(4-fluoro-5-chloro-2-nitro)benzene (29o):

[0509] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**28o**, 70 mg, 0.323 mmol) and maytansinol (**6**, 48 mg, 0.085

mmol) as a yellow solid (35 mg, 53% yield). MS (ESI, pos.): calc'd for $C_{35}H_{39}Cl_2FN_4O_{11}$, 780.2; found 781.4 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-fluoro-5-chloro-2-amino)benzene (30o):

[0510] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29o**, 34 mg, 0.044 mmol) as a white solid (21 mg, 60% yield). MS (ESI, pos.): calc'd for $C_{35}H_{41}Cl_2FN_4O_9$, 750.2; found 751.5 (M+H); 1H NMR (500 MHz, DMSO- d_6) δ : 7.97 (s, 1H), 7.19 (s, 1H), 6.91 (s, 1H), 6.73 (d, $J = 11.4$ Hz, 1H), 6.66 – 6.30 (m, 2H), 5.84 (s, 1H), 5.52 (dd, $J = 14.9, 9.3$ Hz, 1H), 5.17 (s, 2H), 4.52 (d, $J = 11.8$ Hz, 1H), 4.15 (t, $J = 11.1$ Hz, 1H), 3.94 (s, 3H), 3.47 (d, $J = 9.2$ Hz, 1H), 3.24 (s, 3H), 3.17 (d, $J = 12.2$ Hz, 1H), 3.10 (s, 3H), 2.75 (d, $J = 9.9$ Hz, 1H), 2.43 (d, $J = 13.4$ Hz, 1H), 2.08 (d, $J = 13.7$ Hz, 1H), 1.81 (s, 1H), 1.66 (s, 3H), 1.52 (d, $J = 13.5$ Hz, 1H), 1.45 (t, $J = 12.9$ Hz, 2H), 1.11 (d, $J = 6.3$ Hz, 3H), 0.77 (s, 3H).

EXAMPLE 24

Maytan-3-O-carbamoyl-N-(4-amino-2-methyl-5-morpholin-4-yl)benzene (30p)

Step A: (4-nitro-2-methyl-5-morpholin-4-yl)phenyl isocyanate (28p):

[0511] Using the procedure of Example 9, step A, the title compound was prepared from 2-methyl-5-morpholin-4-yl-4-nitrophenylamine (237 mg, 1.00 mmol) as a bright orange solid (230 mg, 87% yield). 1H NMR (500 MHz, $CDCl_3$): δ 7.72 (s, 1H), 6.79 (s, 1H), 3.84 (t, 4H, $J = 5$ Hz), 3.01 (t, 4H, $J = 5$ Hz), 2.31 (s, 3H).

Step B: Maytan-3-O-carbamoyl-N-(4-nitro-2-methyl-5-morpholin-4-yl)benzene (29p):

[0512] Using the method of Example 19, step A, the title compound was prepared from the product of the preceding step (**28p**, 56 mg, 0.21 mmol) and maytansinol (**6**, 50 mg, 0.09 mmol) as an off-white solid (8 mg, 11%). MS (ESI, pos.): calc'd for $C_{40}H_{50}ClN_5O_{12}$, 827.3; found 828.2 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-amino-2-methyl-5-morpholin-4-yl)benzene (30p):

[0513] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29p**, 8 mg, 0.01 mmol) as a white solid (3.2 mg, 42%). MS (ESI, pos.): calc'd for C₄₀H₅₂ClN₅O₁₀, 797.3; found 798.2 (M+H), 820.2 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 7.54 (s, 1H), 7.17 (s, 2H), 6.88 (s, 1H), 6.84 (s, 1H), 6.62 – 6.52 (m, 2H), 6.51 (s, 1H), 5.69 – 5.51 (m, 2H), 4.69 (s, 2H), 4.45 (dd, *J* = 12.0, 2.7 Hz, 1H), 4.16 (t, *J* = 10.9 Hz, 1H), 3.93 (s, 3H), 3.73 (t, *J* = 4.8 Hz, 4H), 3.64 (d, *J* = 12.3 Hz, 1H), 3.53 – 3.44 (m, 1H), 3.25 (s, 3H), 3.14 (d, *J* = 12.2 Hz, 1H), 3.05 (s, 3H), 2.77 (q, *J* = 4.2, 3.2 Hz, 4H), 2.69 (d, *J* = 9.6 Hz, 1H), 2.40 (d, *J* = 12.7 Hz, 1H), 2.01 (s, 3H), 1.97 (d, *J* = 13.4 Hz, 1H), 1.65 (s, 3H), 1.56 (d, *J* = 13.6 Hz, 1H), 1.42 (t, *J* = 12.9 Hz, 2H), 1.09 (d, *J* = 6.4 Hz, 3H), 0.78 (s, 3H).

EXAMPLE 25

Maytan-3-O-carbamoyl-N-(4-amino-2-methyl-5-pyrrolidin-1-yl)benzene (30q)

Step A: (4-nitro-2-methyl-5-pyrrolidin-1-yl)phenyl isocyanate (28q):

[0514] Using the procedure of Example 9, step A, the title compound was prepared from 2-methyl-4-nitro-5-pyrrolidin-1-yl-phenylamine (221 mg, 1.00 mmol) as a bright orange solid (210 mg, 85% yield). ¹H-NMR (300 MHz, CDCl₃): δ 7.65 (s, 1H), 6.59 (s, 1H), 3.25-3.14 (m, 4H), 2.00-1.94 (m, 4H).

Step B: Maytan-3-O-carbamoyl-N-(4-nitro-2-methyl-5-pyrrolidin-1-yl)phenyl (29q):

[0515] Using the method of Example 19, step A, the title compound was prepared from the product of the preceding step (**28q**, 53 mg, 0.21 mmol) and maytansinol (**6**, 50 mg, 0.09 mmol) as an off-white solid (28 mg, 39%). MS (ESI, pos.): calc'd for C₄₀H₅₀ClN₅O₁₁, 811.3; found 812.2 (M+H).

Step C: Maytan-3-O-carbamoyl-N-(4-amino-2-methyl-5-pyrrolidin-1-yl)benzene (30q):

[0516] Using the method of Example 4, step B, the title compound was prepared from the product of the preceding step (**29q**, 28 mg, 0.03 mmol) as a white solid (8.6 mg, 32%). MS (ESI, pos.): calc'd for C₄₀H₅₂ClN₅O₉, 781.3; found 782.2 (M+H), 804.2 (M+Na). ¹H NMR (500 MHz, DMSO-*d*₆) δ: 7.53 (s, 1H), 7.19 (s, 2H), 6.93 – 6.86 (m, 1H), 6.83 (s, 1H), 6.66 – 6.50 (m, 2H), 6.49 (s, 1H), 5.62 (dd, *J* = 14.6, 9.2 Hz, 1H), 5.56 (s, 1H), 4.54 (s, 2H), 4.47 (dd, *J* = 12.0, 2.7 Hz, 1H), 4.17 (t, *J* = 11.1 Hz, 1H), 3.94 (s, 3H), 3.65 (d, *J* = 12.2 Hz, 1H), 3.55 – 3.45 (m, 1H), 3.25 (s, 3H), 3.15 (d, *J* = 12.2 Hz, 1H), 3.07 (s, 3H), 2.93 (d, *J* = 6.7 Hz, 4H), 2.70 (d, *J* = 9.7 Hz, 1H), 2.41 (d, *J* = 12.9 Hz, 1H), 2.02 (s, 3H), 1.98 (d, *J* = 13.2 Hz, 1H), 1.89 – 1.76 (m, 4H), 1.66 (s, 3H), 1.58 (d, *J* = 13.5 Hz, 1H), 1.43 (t, *J* = 12.9 Hz, 2H), 1.10 (d, *J* = 6.4 Hz, 3H), 0.79 (s, 3H).

EXAMPLE 26

Conjugate Preparation and Characterization

[0517] Five antibodies were conjugated to the linker-payload compounds of the disclosure using the procedure below. The four targeting antibodies used in these experiments were: (1) a PSMA antibody having the heavy and light chain variable domains of clone AB-PG1-XG1-006 as set forth in International Patent Application Publication No. WO2007/002222 A2, entitled PSMA ANTIBODY-DRUG CONJUGATES, (2) anti-MUC16 antibody having variable regions derived from clone 3A5 from International Patent Application Publication No. WO2007001851, entitled COMPOSITIONS AND METHODS FOR THE DIAGNOSIS AND TREATMENT OF TUMOR, (3) anti-HER2 antibody having variable regions derived from humAb4D5-8 from Carter *et al*, PNAS 1992 89 4285, and (4) an anti-STEAP2 antibody. These patent application publications and journal citations are herein incorporated by reference in their entirety for all purposes. All the monoclonal antibodies were expressed in CHO cells and purified by Protein A. A non-binding isotype control (5) derived from an immunological antigen having no relation to oncology was also used.

[0518] *Method A:* The antibody (10 mg/ml) in 50 mM HEPES, 150 mM NaCl, pH 7.5, was treated with 1 mM dithiothreitol at 37 °C for 30 min. After gel filtration (G-25, pH 4.5 sodium acetate), the maleimido linker payload derivative (for linker-payloads **9**, **19**, or **24**; 1.2 equivalents/S_H group) in DMSO (10 mg/ml) was added to the reduced antibody and the mixture adjusted to pH 7.0 with 1 M HEPES (pH 7.4). The conjugation reaction was optionally quenched

with N-ethyl maleimide (NEM). The conjugates were purified by size exclusion chromatography and sterile filtered. Protein concentrations and payload to antibody ratios were determined by UV spectral analysis according to Hamblett *et al* (American Association for Cancer Research, 2004 Oct 15;10(20):7063-70) or mass difference, native versus conjugated. Size-exclusion HPLC established that all conjugates used were >95% monomeric, and RP-HPLC established that there was <0.5% unconjugated linker payload. Yields and payload to antibody ratios are reported in Table 1.

[0519] *Method B:* The antibodies (10-20 mg/ml) in 50 mM HEPES, 150 mM NaCl, pH 8.0, and 10-15% (v/v) DMA were conjugated with a 5-6 fold excess of **10**, **14**, **20**, **22**, or **26** for 2 hours (hrs) at ambient temperature. The conjugate was purified by size exclusion chromatography or extensive ultrafiltration and sterile filtered. Protein concentrations were determined by UV spectral analysis. Size-exclusion HPLC established that all conjugates used were >90% monomeric, and RP-HPLC established that there was <1% unconjugated linker payload. All conjugated antibodies were analyzed by mass difference, native versus conjugated. Payload to antibody ratios are reported in Table 1.

[0520] *Method C - Bacterial Transglutaminase Conjugation:* Aglycosylated, deglycosylated, or glutamine tagged antibodies can be conjugated at 1-10 mg/mL in PBS pH 7.4. Linker payload **22** can be added in a 10 to 25-fold molar excess over antibody and the enzymatic reaction can be initiated by addition of 1-5 units of bacterial transglutaminase (Zedira, T1001) per mg antibody and incubated with shaking at 37°C for 4-16 hours. The conjugates can be purified by size exclusion chromatography and sterile filtered. Protein and linker payload concentrations can be determined by UV spectral analysis. Size-exclusion HPLC can establish that conjugates can be >95% monomeric. All conjugated antibodies can be analyzed by UV for linker payload loading values according to Hamblett et al, Cancer Res 2004 10 7063. In addition, the conjugates can be analyzed by ESI-MS for linker payload loadings using a Waters Synapt G2-Si QTOF mass spectrometry coupled with Acquity UPLC. The chromatographic separation can be achieved on a C4 column (Waters protein BEH C4, 50 mm X 1 mm, 1.7 µm) in a 25 minute gradient (minute:percentage of mobile phase B; 0:20%, 1:20%, 18:40%, 18.1:90, 20:95%, 20.8:95%, 20.9:20% 25:20%). The mobile phase A can be 0.1% formic acid in water and mobile phase B can be 0.1% formic acid in acetonitrile. The flow rate can be set at 100

μl/min. The detector TOF scan can be set from m/z 700-5000 for 25 minutes with major parameters as listed (Capillary voltage 3.2 kV; Sampling Cone 150; Source Offset at 80; Source temperatures 120 °C; Desolvation temperature 500 °C; Trap collision Energy 30; Transfer Collision Energy Off; Gas controls OFF; Resolving Quadrupole: LM resolution at 4.7). The combined spectra can be deconvoluted with MaxEnt function within MassLynx software.

Conjugates of Method A DAR Determination

[0521] To determine the loading of the linker-payloads on the antibody (cysteine conjugates), the conjugates were deglycosylated, reduced, and analyzed by LC-MS.

[0522] For the assay, 50 μg of the conjugate was diluted with mili-Q water to a final concentration of 1 mg/mL. Ten μL of PNGase F solution [PNGase F solution was prepared by adding 150 μL of PNGase F stock (New England Biolabs, Cat#P0704L) and 850 μL of mili-Q water and mixed well] was added to the diluted conjugate solution and then incubated at 37°C overnight. 2.4 μL of 0.5 M TCEP was added to the sample such that the resulting material had a final TCEP concentration of 20 mM and this was then incubated at 50°C for 30 minutes.

Injections of 10 μL of each sample were made onto LC-MS (Waters Synat G2-Si) and eluted with 0.1 mL/minute of a gradient mobile phase 20-40% of mobile phase B over 25 minutes (Mobile Phase A: 0.1% v/v FA in H₂O; Mobile Phase B: 0.1% v/v FA in Acetonitrile). The LC separation was achieved on Waters Acquity BEH C18 column (1.0 X 50 mM, 1.7 μM).

[0523] The mass spectrometry spectra were deconvoluted and the identified light and heavy chain peaks represent the light chain (L) with linker-payload values = 0 and 1, heavy chain (H) with linker-payload values = 0, 1, 2, and 3. From the intensity values of each species, the drug to antibody ratio (DAR) was calculated using equation 1 below for a homo-dimer antibody conjugate.

Equation 1:

$$\text{DAR} = 2 * \left[\frac{\text{L1}}{\text{L0} + \text{L1}} + \frac{\text{H1} + 2 * \text{H2} + 3 * \text{H3}}{\text{H0} + \text{H1} + \text{H2} + \text{H3}} \right]$$

Conjugates of Method B DAR Determination

[0524] To determine the loading of the linker-payloads on the antibody (lysine conjugates), the conjugates were deglycosylated, and analyzed by LC-MS.

[0525] For the assay, 50 μg of the conjugate was diluted with milli-Q water to a final concentration of 1 mg/mL. Ten μL of PNGase F solution [PNGase F solution was prepared by adding 150 μL of PNGase F stock (New England Biolabs, Cat#P0704L) and 850 μL of milli-Q water and mixed well] was added to the diluted conjugate solution and then incubated at 37°C overnight. Injections of 5 μL of each sample were made onto LC-MS (Waters Synat G2-Si) and eluted with 0.1 mL/minute of a gradient mobile phase 20-50% of mobile phase B over 25 minutes (Mobile Phase A: 0.1%v/v FA in H₂O; Mobile Phase B: 0.1% v/v FA in Acetonitrile). The LC separation was achieved on a Waters Acquity BEH C4 column (1.0 X 50 mM, 1.7 μM) at 80 °C.

[0526] The mass spectrometry spectra were deconvoluted using Masslynx software and the drug to antibody ratio (DAR) was calculated using the following equations. *See* FIG. 18.

1. Relative percentage (%) of drug (D_n) by distribution peak intensity (PI):

a. $D_n\% = \text{PI}_n / \Sigma(\text{PI}_0 + \text{PI}_1 + \text{PI}_2 + \dots + \text{PI}_i) \times 100$

b. (n= 0,1,2,3,...,i)

2. Average DAR calculation:

a. $\text{DAR} = \Sigma(1 \times D_1\% + 2 \times D_2\% + 3 \times D_3\% + \dots + i \times D_i\%)$

Table 1

Antibody	$\epsilon_{252 \text{ nm}} (\text{cm}^{-1} \text{ M}^{-1})$	$\epsilon_{280 \text{ nm}} (\text{cm}^{-1} \text{ M}^{-1})$
PSMA	77652	224320
MUC16	85888	247360
HER2	81847	215388
STEAP2	-	-
Isotype Control	75113	218360

Antibody Conjugate	Payload:Antibody (UV)	Yield %
PSMA-9	1.3	80
MUC16-9	1.4	70
HER2	1.5	75
Isotype Control-9	1.5	80

Antibody Conjugate	Payload:Antibody (ESI-MS)	Yield %
STEAP2-14	2.7	50
STEAP2-20	1.2	45
STEAP2-24	3.0	40
HER2-10	2.3	60
HER2-20	0.7	50
HER2-24	3.3	40
Isotype Control-10	2.4	75
Isotype Control-14	1.3	70
Isotype Control-20	1.0	50
Isotype Control-24	3.6	40

EXAMPLE 27*In Vitro Linker-Payload Cell-free Enzymatic Assays*Cathepsin B incubation

[0527] *In vitro* cell-free enzymatic assay procedure was adopted from Dubowchik, *et al.* Bioconjugate Chem. **2002** 13 855, the entire contents of which are herein incorporated by

reference in its entirety for all purposes. The linker payload **9** was set at 100 µg/mL final in 25 mM sodium acetate buffer, 1 mM EDTA, pH 5.0 and pre-incubated at 37°C. Cathepsin B (Sigma # C8571) was activated at room temperature for 15 minutes with 1 equivalent of 30 mM DTT, 15 mM EDTA to 2 equivalents of cathepsin B stock. The activated cathepsin B solution was added to the substrate solutions at a 1:20 molar ratio (purified H₂O, instead of activated cathepsin B was added for the control sample.) Samples were incubated at 37°C overnight and the resulting samples are detected by LC-MS through Q1 Scan.

LC-MS detection

[0528] Samples are centrifuged at 12,000 g for 5 min. Supernatant was recovered and analyzed by liquid chromatography-mass spectrometry (Waters Xevo TQ-S) by combined infusion of 0.3 ml/min of 30:70 mobile phase B:A (Mobile Phase A: 0.1%FA in H₂O; Mobile Phase B: 0.1% FA in Acetonitrile) at 20 µl/min from supernatant. MS1 is set at an appropriate range for detection of molecular ion of either linker payload or payload. The supernatant contained the predicted payload, Maytan-3-O-carbamoyl-N-(4-aminobenzene) (**8**), with a mass of 720.9 (M+Na) (calc'd monoisotopic mass for C₃₅H₄₃N₄O₉Cl, 698.3) and the control samples without cathepsin B contained **9** with a mass of 1170.4 (M+Na) (calc'd monoisotopic mass for C₆₀H₈₀ClN₉O₁₆, 1147.50). No predicted payload molecular ion was detected in the control samples.

[0529] The results of this Example are significant in part because cathepsin B proteolysis of **9** should only occur after internalization of the ADC in the cell where the enzyme exists. Off target effects should be reduced since the antibody delivers the cytotoxic payload directly to targeted cells.

EXAMPLE 28

In Vitro Cytotoxicity Assays

[0530] In this Example, the ability of various antibody-drug conjugates and naked payloads to kill antigen-expressing tumor cells *in vitro* was assessed.

[0531] Ovc3 (Muc16+) cells were seeded in 96 well plates at 3000 cells per well in complete growth media and grown overnight. For cell viability curves, serially diluted

conjugates or naked payloads were added to the cells at final concentrations ranging from 100 nM to 5 pM and incubated for 3 days. To measure viability, cells were incubated with CCK8 (Dojindo) for the final 2 hours and the absorbance at 450nm (OD₄₅₀) was determined on a Victor (Perkin Elmer). Background OD₄₅₀ levels determined from digitonin (40 nM) treated cells were subtracted from all wells and viability is expressed as a percentage of the untreated controls. IC₅₀ values were determined from a four-parameter logistic equation over a 10-point response curve (GraphPad Prism). The IC₅₀ for the MUC16-9 ADC was +++ which indicates an IC₅₀ (nM) greater than, or equal to, 0.1 and less than 1.0. All curves and IC₅₀ values are corrected for payload equivalents. The plot of % Cell Viability vs. Log₁₀ [M] Drug is shown in FIG 1. Naked payload IC₅₀ values and percent cell kill are listed in Table 2.

[0532] SKBR3 (Her2+) cells were seeded in 96 well plates at 8000 cells per well in complete growth media and grown overnight. For cell viability curves, serially diluted conjugates or naked payloads were added to the cells at final concentrations ranging from 100 nM to 5 pM and incubated for 3 days. To measure viability, cells were incubated with CCK8 (Dojindo) for the final 2 hours and the absorbance at 450nm (OD₄₅₀) was determined on a Victor (Perkin Elmer). Background OD₄₅₀ levels determined from digitonin (40 nM) treated cells were subtracted from all wells and viability is expressed as a percentage of the untreated controls. IC₅₀ values were determined from a four-parameter logistic equation over a 10-point response curve (GraphPad Prism). The IC₅₀ for the HER2-9, 10, 20, and 24 ADCs was +++, +++, +++, and +++, respectively, wherein +++ indicates IC₅₀ (nM) greater than, or equal to, 0.1 and less than 1.0. All curves and IC₅₀ values are corrected for payload equivalents. The plot of % Cell Viability vs. Log₁₀ [M] Drug is shown in FIGS 2-5. Naked payload IC₅₀ values and percent cell kill are listed in Table 2.

[0533] C4-2 (STEAP2+) cells were seeded in 96 well plates at 4000 cells per well in complete growth media and grown overnight. For cell viability curves, serially diluted conjugates or naked payloads were added to the cells at final concentrations ranging from 100 nM to 5 pM and incubated for 5 days. To measure viability, cells were incubated with CellTiter-Glo reagents for 5 minutes and luminescence was determined on a Victor plate reader (PerkinElmer). IC₅₀ values were determined from a four-parameter logistic equation over a 10-point response curve (GraphPad Prism). The IC₅₀ for the STEAP2-14, 20, and 24 ADCs were ++, +++, + nM, respectively. +++ indicates IC₅₀ (nM) greater than, or equal to, 0.1 and less than

1.0; ++ indicates IC₅₀ (nM) greater than, or equal to, 1.0 and less than 10; and + indicates IC₅₀ (nM) greater than, or equal to, 10.

[0534] All IC₅₀s are expressed in nM concentration and % kill is reported for the highest dose. All curves and IC₅₀ values are corrected for payload equivalents. The plot of % Cell Viability vs. Log₁₀ [M] Drug is shown in FIGS 6-8. Naked payload IC₅₀ values and percent cell kill are listed in Table 2.

[0535] The embodiments and examples described above are intended to be merely illustrative and non-limiting. Those skilled in the art will recognize or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials and procedures. All such equivalents are considered to be within the scope and are encompassed by the appended claims.

Table 2

Compound #	Ovcar3		C4-2		SKBr-3	
	IC50 (nM)	% kill	IC50 (nM)	% kill	IC50 (nM)	% kill
8	++	**	+	****	++	****
17	+++++	***	+++	****	++++	*****
30a	+++	***	++	*****	+++	****
30b			+	*	+++	****
30c	++	**	+	**	++	***
30d			++	****	++	****
30e			++	****	++	***
30f			++	****	+++	*****
30g			++	*****	+++	*****
30h			++	*	+++	****
30i			+	*****	++	*****
30j	++	**	+	***	+++	****
30k	+++	***	+		++	*****
30l	+++	***	+		+++	*****
30m	+++++	**	++++	*****	++++	*****
30n			+	*****	++	*****
30o			+	*	+++	*****
30p			+		+	*
30q			+	***	++	*****

+++++ indicates IC₅₀ (nM) greater than, or equal to, 0.001 and less than 0.01;

++++ indicates IC₅₀ (nM) greater than, or equal to, 0.01 and less than 0.1;

+++ indicates IC₅₀ (nM) greater than, or equal to, 0.1 and less than 1.0;

++ indicates IC₅₀ (nM) greater than, or equal to, 1.0 and less than 10; and

+ indicates IC₅₀ (nM) greater than, or equal to, 10.

***** indicates % kill greater than, or equal to, 99;

**** indicates % kill greater than, or equal to, 95 and less than 99;

*** indicates % kill greater than, or equal to, 90 and less than 95;


** indicates % kill greater than, or equal to, 85 and less than 90; and

* indicates % kill less than 85.

[0536] The embodiments and examples described above are intended to be merely illustrative and non-limiting. Those skilled in the art will recognize or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials and procedures. All such equivalents are considered to be within the scope and are encompassed by the appended claims.

The claims defining the invention are as follows:

1. A compound of Formula (I):



(I)

or a pharmaceutically acceptable salt or stereoisomer thereof,

wherein:

A is arylene or heteroarylene;


L is a linker;

BA is a binding agent, wherein the binding agent is an antibody or antigen-binding fragment thereof; and

k is an integer from 1 to 30.

2. The compound of claim 1, wherein

A is:



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,

heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}-R^A$, and azido;

R^A is alkyl;

n is an integer from 0 to 4; and


ξ and ξ each represents a bond from A to -NH-.

3. The compound of claim 2, wherein R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.

4. The compound of any of claim 2 or 3,

wherein:


A is selected from:




n is 0, 1, 2, or 3.

5. The compound of any of claims 2-4, wherein R^1 , independently at each occurrence, is selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, pyrrolidinyl, and morpholinyl.

6. The compound of any of claims 2-5, wherein R^1 is methyl or chloro.

7. The compound of any of claims 2-6, wherein A is .

8. The compound of any of claims 1-7, wherein L is:



wherein:

SP is a spacer;

$\overset{\text{A}}{\underset{\sim}{\sim}}$ is one or more bonds to the binding agent;


$\overset{\sim}{\sim}$ is a bond from L to -NH-A-;

AA¹ is an amino acid; and

AA² is an amino acid.

9. The compound of claim 8, wherein AA¹-AA² is: valine-citrulline, citrulline-valine, lysine-phenylalanine, phenylalanine-lysine, valine-asparagine, asparagine-valine, threonine-asparagine, asparagine-threonine, serine-asparagine, asparagine-serine, phenylalanine-asparagine, asparagine-phenylalanine, leucine-asparagine, asparagine-leucine, isoleucine-asparagine, asparagine-isoleucine, glycine-asparagine, asparagine-glycine, glutamic acid-asparagine, asparagine-glutamic acid, citrulline-asparagine, asparagine-citrulline, alanine-asparagine, asparagine-alanine, valine-alanine, alanine-valine, valine-glycine, or glycine-valine.

10. The compound of any of claims 1-8, wherein L is:




wherein:

SP is a spacer;

$\overset{\text{A}}{\underset{\sim}{\sim}}$ is one or more bonds to the binding agent;

$\text{---}\overset{\text{A}}{\text{---}}$ is a bond from L to -NH-A-;
 R^{AA1} is an amino acid side chain; and
 R^{AA2} is an amino acid side chain.

11. The compound of claim 10, wherein L is:




wherein:

SP is a spacer; and

$\text{---}\overset{\text{A}}{\text{---}}$ is one or more bonds to the binding agent; and

$\text{---}\overset{\text{A}}{\text{---}}$ is a bond from L to -NH-A-.

12. The compound of any of claims 8-11, wherein SP comprises a C_{5-7} alkylene.
13. The compound of any of claims 8-12, wherein SP is selected from:




wherein:

b, independently in each instance, is an integer from 2 to 8;


$\text{---}\overset{\text{A}}{\text{---}}$ is one or more bonds to the binding agent; and





$\text{---}\overset{\text{A}}{\text{---}}$ is a bond from L to -NH-A-.







wherein $\text{---}\overset{\text{A}}{\text{S}}\text{---}$ is a bond to the binding agent and $\text{---}\overset{\text{L}}{\text{N}}\text{---}$ is a bond from L to -NH-A-.


16. The compound of claim 1, wherein the compound is selected from:





































k ,




k ,




k ,






k ,









k ,








k ,




























wherein:

Ab is an antibody or an antigen binding fragment thereof;

S is a bond to a cysteine of the antibody or the antigen binding fragment thereof;


N is a bond to a lysine of the antibody or the antigen binding fragment thereof;

k is an integer from 1 to 30; and

t is an integer from 1 to 8; and

Gln is a bond to a glutamine of the antibody or the antigen binding fragment thereof.

17. The compound of claim 16, wherein the compound is:




wherein Ab is an antibody or antigen-binding fragment thereof.

18. The compound of any of claims 1-15, wherein the BA is an antibody.

19. The compound of claim 18, wherein the antibody is an anti-MUC16 antibody, an anti-PSMA antibody, an anti-STEAP2 antibody or an anti-HER2 antibody.

20. A compound of Formula II:




(II)

or a pharmaceutically acceptable salt or stereoisomer thereof,
wherein A is arylene or heteroarylene.

21. The compound of claim 20,

wherein:

A is:



R^1 , independently at each occurrence, is halo, haloalkyl, haloalkoxy, alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, aralkyl, heteroaryl, heterocycloalkyl, cyano, nitro, $-\text{OR}^A$,


$-\text{SO}_2\text{R}^A$, $-\text{C}(=\text{O})\text{R}^A$, or azido;

R^A is alkyl; and

n is an integer from 0 to 4.

22. The compound of claim 21, wherein R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl;

23. The compound of any of claims 20-22, wherein the compound of Formula (II) is selected from a compound of the Formula (IIA1), a compound of the Formula (IIA2), and a compound of the Formula (IIA3):














24. The compound of any of claims 21-23, wherein n is 0, 1, 2, or 3; and wherein R^1 , independently at each occurrence, is selected from C_{1-6} alkyl, C_{1-6} alkoxy, halo, C_{1-6} haloalkyl, and heterocycloalkyl.









25. The compound of any of claims 21-24, wherein R^1 is, independently at each occurrence, selected from methyl, methoxy, fluoro, chloro, bromo, trifluoromethyl, pyrrolidinyl, and morpholinyl.









26. The compound of any of claims 21-24, wherein R^1 is methyl or chloro.


27. The compound of claim 20, wherein the compound is selected from:













28. A pharmaceutical composition comprising the compound of any of claims 1-27 and a pharmaceutically acceptable excipient.
29. A method of treating a proliferative disorder comprising administering to a patient having said disorder a therapeutically effective amount of a compound of any of claims 1-27 or a composition of claim 28.
30. A compound of Formula P1:



wherein A is arylene or heteroarylene; and RL is a reactive linker.

31. The compound of claim 30, wherein:

A is:



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,


heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{\text{O}}{\parallel}-R^A$, and azido;

R^A is alkyl; and

n is an integer from 0 to 4.


32. The compound of claim 31, wherein R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.

33. The compound of any of claims 30-32, wherein the compound of Formula P1 is a compound of Formula P2A1-3:




wherein:

A is selected from:




SP^R is a spacer-reactive group.


34. The compound of any of claims 30-32, wherein the compound of Formula P1 is selected from a compound of Formula P1H1, a compound of Formula P1I1, a compound of




(P1W1)



(P1K1)




(P1TG1)





(P1ZZ1)





, and










(P1ZZ2)





35. The compound of claim 30, wherein the compound is selected from:



































































36. A compound of Formula PPII:



wherein A is arylene or heteroarylene.

37. The compound of claim 36, wherein:

A is:



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,


heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, and azido;

R^A is alkyl;

n is an integer from 0 to 4.

38. The compound of claim ~~52~~ 37, wherein R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.


39. The compound of any of claims ~~51-53~~ 36-38, wherein the compound of Formula PPII is a compound of Formula PP5A1:


















wherein:









R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; and
 n is an integer from 0 to 4.





40. The compound of any of claims 36-38, wherein the compound is selected from:













41. A method of preparing a compound of Formula (I):



comprising contacting a compound of P1:



with a binding agent under conjugation conditions,

wherein:

A is arylene or heteroarylene;

L is a linker;


BA is a binding agent, wherein the binding agent is an antibody or antigen-binding fragment thereof;

k is an integer from 1 to 10; and

RL is a reactive linker.

42. The method of claim 41, wherein:

A is:



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,

heterocycloalkyl, cyano, nitro, $-\text{OR}^A$, $-\text{SO}_2\text{R}^A$, $-\text{C}(=\text{O})\text{R}^A$, and azido;

R^A is alkyl; and


n is an integer from 0 to 4.

43. The method of claim 42, wherein:

R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl; and

n is an integer from 0 to 4.

44. The method of claim 41, wherein
 the compound of Formula P1 is prepared by contacting a compound of Formula II
 with a RL,
 RL is a reactive linker; and
 the compound of Formula II is prepared by contacting a compound of Formula
 PP5A with a suitable reducing agent:



wherein:


R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,

heterocycloalkyl, cyano, nitro, $-\text{OR}^A$, $-\text{SO}_2\text{R}^A$, $-\text{C}(=\text{O})\text{R}^A$, and azido;
 R^A is alkyl;

n is an integer from 0 to 4; and

the suitable reducing agent is selected from a metal, a metal foil, a metal powder, dust of a metal, a metal amalgam, or metal filings.

45. A method of making a compound of Formula II:




(II),

wherein:

A is arylene or heteroarylene


comprising:

(a) contacting a compound of formula P2




P2

with a compound of formula PP6:



PP6

under Lewis acid synthesis conditions to form a compound of PPII:



; and


(PPII)

(b) reducing the compound of formula PPII by contacting the compound of formula PPII with a reducing agent to form a compound of Formula II.

46. The method of claim 45:

wherein:

A is:



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,


heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-C(=O)R^A$, and azido;


R^A is alkyl;

n is an integer from 0 to 4.

47. The method of claim 46, wherein R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.

48. A method of making a compound of Formula PPII, comprising contacting a compound of Formula P2 with a compound of Formula PP6 under Lewis acid synthesis conditions:






wherein:

A is arylene or heteroarylene.

49. The method of claim 48, wherein:

A is



R^1 , independently at each occurrence, is selected from alkyl, alkenyl, alkynyl, alkoxy, aryl, alkaryl, arylalkyl, halo, haloalkyl, haloalkoxy, heteroalkyl, heteroaryl,

heterocycloalkyl, cyano, nitro, $-\xi-OR^A$, $-\xi-SO_2R^A$, $-\xi-\overset{O}{\parallel}R^A$, and azido;

R^A is alkyl;

n is an integer from 0 to 4.

50. The method of claim 49, wherein:




R^1 , independently at each occurrence, is selected from alkyl, alkoxy, halo, haloalkyl, and heterocycloalkyl.




51. The method of claim 29, wherein the proliferative disorder is cancer.


52. The method of claim 51, wherein the cancer is renal cell carcinoma, pancreatic carcinoma, head and neck cancer, prostate cancer, malignant gliomas, osteosarcoma, colorectal cancer, gastric cancer, malignant mesothelioma, multiple myeloma, ovarian cancer, small cell lung cancer, non-small cell lung cancer, synovial sarcoma, thyroid cancer, breast cancer, or melanoma.
53. The method of claim 29, wherein the binding agent in the compound of any one of claims 1-27 or the composition of claim 28 interacts with or binds to a tumor antigen.
54. The method of claim 53, wherein the tumor antigen is selected from the group consisting of prolactin receptor (PRLR), prostate-specific membrane antigen (PSMA), MUC16, STEAP2, and EGFRvIII.
55. The method of claim 53, wherein the tumor antigen is selected from the group consisting of AFP, ALK, B7H4, BAGE proteins, β -catenin, bcr-abl, BRCA1, BORIS, CA9 (carbonic anhydrase IX), caspase-8, CD20, CD40, CD123, CDK4, CEA, CLEC12A, c-kit, cMET, CTLA4, cyclin-B1, CYP1B1, EGFR, EGFRvIII, endoglin, Epcam, EphA2, ErbB2/Her2, ErbB3/Her3, ErbB4/Her4, ETV6-AML, Fra-1, FOLR1, GAGE proteins, GD2, GD3, GloboH, glypican-3, GM3, gp100, Her2, HLA/B-raf, HLA/EBNA1, HLA/k-ras, HLA/MAGE-A3, hTERT, IGF1R, LGR5, LMP2, MAGE proteins, MART-1, mesothelin, ML-IAP, Muc1, Muc16, CA-125, MUM1, NA17, NGEF, NY-BR1, NY-BR62, NY-BR85, NY-ESO1, OX40, p15, p53, PAP, PAX3, PAX5, PCTA-1, PDGFR- α , PDGFR- β , PDGF-A, PDGF-B, PDGF-C, PDGF-D, PLAC1, PRLR, PRAME, PSCA, PSGR, PSMA (FOLH1), RAGE proteins, Ras, RGS5, Rho, SART-1, SART-3, Steap-1, Steap-2, STn, survivin, TAG-72, TGF- β , TMPRSS2, Tn, TNFRSF17, TRP-1, TRP-2, tyrosinase, and uroplakin-3, alpha-actinin-4 with lung cancer, ARTC1 with melanoma, BCR-ABL fusion protein with chronic myeloid leukemia, B-RAF, CLPP or Cdc27 with melanoma, CASP-8 with squamous cell carcinoma, and hsp70-2 with renal cell carcinoma, BAGE-1, GAGE, GnTV, KK-LC-1, MAGE-A2, NA88-A, and TRP2-INT2, BCMA, SLAMF7, B7H4, GPNMB, UPK3A, and LGR5, and fragments thereof.
56. The method of claim 29, wherein the binding agent in the compound of any one of claims 1-27 or the composition of claim 28 is an antibody selected from the group consisting

of an anti-PSMA antibody, anti-MUC16 antibody, anti-HER2 antibody, anti-EGFRvIII antibody, and anti-STEAP-2 antibody.

57. The compound of claim 1, wherein the compound is selected from:







wherein:

Ab is an antibody, wherein the binding agent is an antibody or antigen-binding fragment thereof;

S is a bond to a cysteine of the antibody or the antigen binding fragment thereof;


N is a bond to a lysine of the antibody or the antigen binding fragment thereof;

k is an integer from 1 to 30; and




t is an integer from 1 to 8; and





Gln is a bond to a glutamine of the antibody or the antigen binding fragment thereof.

58. The compound of claim 27 that is selected from:




59. The compound of claim 35, wherein the compound is selected from:





, and



60. Use of a compound of any of claims 1-27, 57, 58, and 59 in the manufacture of a medicament for treatment of a proliferative disorder.
61. The use of claim 60 wherein the proliferative disorder is cancer.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18