

(19) **DANMARK**

(10) **DK/EP 2257263 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 8/02 (2006.01)** **A 61 K 9/00 (2006.01)** **A 61 K 33/30 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2014-12-01**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2014-09-03**
- (86) Europæisk ansøgning nr.: **08744462.6**
- (86) Europæisk indleveringsdag: **2008-03-27**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-12-08**
- (86) International ansøgning nr.: **US2008058428**
- (87) Internationalt publikationsnr.: **WO2009108210**
- (30) Prioritet: **2008-02-27 US 38283**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Clji Corporation, 550 Northlake Boulevard, North Palm Beach, FL 33408, USA**
- (72) Opfinder: **LICHTBLAU, Craig, 18826 Loxahatchee River Road, Jupiter, FL 33458, USA**
IPARRAGUIRRE, Jose, I., 8950 N. Kendall Drive, Suite 302, Miami, FL 33176, USA
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Højbro Plads 10, 1200 København K, Danmark**
- (54) Benævnelse: **TOPIKE FORMULERINGER TIL FOREBYGGELSE AF SEKSUELT OVERFØRTE SYGDOMME OG FREMGANGSMÅDER TIL FREMSTILLING HERAF**
- (56) Fremdragne publikationer:
EP-A1- 1 506 781
US-A1- 2003 031 727
US-A1- 2005 048 139
US-A1- 2006 018 937
US-B1- 6 183 762
US-B2- 7 097 828
TENNICAN P O ET AL: "The diverse effects of topical and systemic administration of zinc on the virulence of Herpes simplex genitalis", LIFE SCIENCES, PERGAMON PRESS, OXFORD, GB, vol. 24, no. 20, 14 May 1979 (1979-05-14), pages 1877-1883, XP023768417, ISSN: 0024-3205, DOI: 10.1016/0024-3205(79)90239-X [retrieved on 1979-05-14]
EBY G A ET AL: "Use of topical zinc to prevent recurrent herpes simplex infection: Review of literature and suggested protocols", MEDICAL HYPOTHESES, EDEN PRESS, PENRITH, US, vol. 17, no. 2, 1 June 1985 (1985-06-01), pages 157-165, XP023026809, ISSN: 0306-9877, DOI: 10.1016/0306-9877(85)90142-2 [retrieved on 1985-06-01]
DATABASE GNPD [Online] Mintel; April 2003 (2003-04), "Sensitive Facial Skin Extra Protective Lotion", XP002675579, Database accession no. 10134434
SKOLER-KARPOFF S ET AL: "Efficacy of Carraguard for prevention of HIV infection in women in South Africa:

Fortsættes ...

a randomised, double-blind, placebo-controlled trial", THE LANCET, LANCET LIMITED. LONDON, GB, vol. 372, no. 9654, 6 December 2008 (2008-12-06), pages 1977-1987, XP025745749, ISSN: 0140-6736, DOI: 10.1016/S0140-6736(08)61842-5 [retrieved on 2008-12-04]

Description

FIELD OF THE INVENTION

[0001] This invention relates to topical protective formulations for use in the prevention of the spread of sexually transmitted diseases (STDs); and particularly to unique topical formulations including a plurality of film forming excipients which act in concert to provide a barrier to help inhibit the transmission of STDs.

BACKGROUND OF THE INVENTION

[0002] Sexually transmitted diseases (STDs) are a universal concern, effecting millions of individuals and straining health care systems. More than 20 different sexually transmitted diseases have been identified by the medical community and generally fall into two groups, including bacterial types (e.g., gonorrhea, chlamydia and syphilis) and viral types (human immunodeficiency virus (HIV), human papilloma virus (HPV) and hepatitis). The numerous diseases affect men, women and children of all backgrounds, races and economic classifications. Despite years of research and educational programs, the transmission of sexually transmitted diseases remains a global health threat. Although specific methods of transmission may vary depending on the disease-causing organism, STDs are usually transmitted to the uninfected person through injured or exposed skin or mucous membranes during sexual contact.

[0003] Treatments may be available for some types of STDs (e.g. antibiotic treatment for gonorrhea or chlamydia). However, most people who suffer from these types of STDs are unaware that they have the disease and therefore do not get the necessary treatment. Moreover, because of the sociological impact and generally negative stigma associated with these diseases, people are reluctant to seek such treatments. The continued emphasis on educating the population as to the use of "mechanical barriers" such as condoms has helped to decrease the morbidity of most STDs but more preventative methods need to be developed to prevent STD transmission.

[0004] Chemical actives such as microbicides, antimicrobials, and spermicides, most notably, nonylphenoxypoly(ethyleneoxy)-ethanol (also referred to as nonoxynol-9) have been used in topical formulations to effectively reduce the rate of STD transmission, especially when used in conjunction with prophylactics. However, many of these chemical actives are harsh and have been shown to induce local irritation, inflammation, and ulcerations which might actually favor the transmission of STDs. Thus, a need exists for topical formulations that do not cause irritation and will help inhibit the spread of STDs, especially when used in combination with condoms, and thereby provide an additional degree of protection from contamination, should the condom become damaged.

PRIOR ART

[0005] Although there are numerous patents and publications directed to formulations containing chemical actives, such as, microbicides, antimicrobials, spermicides, and drug delivery carriers (liposomes, micelles), none of the known prior art teach formulations comprising non-irritating agents that provide a physical barrier to the permeation of pathogens.

5 [0006] U.S. Patent Application 2003/0143189 A1 to Askill et al., is directed to a method of treating skin lesions by forming a polymeric film over the lesions to inhibit proliferation of infectious agents in the lesions. These compositions also include one or more chemical agents in combination therewith.

[0007] U.S. Patent 6,835,717 to Hildreth discloses a method of reducing the risk of a sexually transmitted pathogen by contacting a pathogen within the composition which consists of β -cyclodextrin.

10 [0008] U.S. Patent No. 6,821,958 to Hershline discloses a method of preventing viral transmission using an alkylsulfate derivative of sulfated dextrin as a topical formulation.

[0009] U.S. Patent 6,582,711 to Asmus et al., discloses an antimicrobial hydroalcoholic composition having a cationic polymeric thickener.

15 [0010] U.S. Patent No. 6,355,235 to Cone et al., consists of an antibody capable of trapping sperm and a pharmaceutical carrier.

[0011] U.S. Patent No. 6,328,991 to Myhling discloses a chemical composition to prevent the transmission of sexually transmitted diseases comprising nonylphenoxypoly-(ethyleneoxy)-ethanol, benzalkonium chloride and povidone iodine.

20 [0012] U.S. Patent 5,439,685 to Augros is a composition of an agent active against microorganisms responsible for sexually transmitted diseases together with a film-forming agent such as dimethylpolysiloxane, and benzalkonium chloride as a spermicidal.

[0013] U.S. Patent 4,952,411 to Fox, Jr. et al., is a composition of silver sulfadiazine, alone or in combination with chlorhexidine or sodium deoxycholate (antimicrobial and detergent).

25 [0014] Tennican et al., 1979 discusses the diverse effects of topical and systemic administration of zinc on the virulence of Herpes simplex genitals.

[0015] Eby, 1985 discusses the use of topical zinc to prevent recurrent herpes simplex infection; review of literature and suggested protocols.

Summary of the Invention

30 [0016] In a first aspect, the invention provides an oil-in-water formulation for the inhibition of the transmission of sexually transmitted pathogens, characterized by enhanced skin protective properties consisting of a combination of:

(1) Dimethicone 20, in the range of 10.0 to 20.0% wt/wt;

(2) Dimethicone 12500, in the range of 3.0-10.0% wt/wt;

(3) Zinc oxide, in the range of 1.0 to 8.0% wt/wt;

(4) a mixture of Cetearyl alcohol and Ceteareth-20, in the range of 6.0 to 10.0% wt/wt;

(5) Glycerin, in the range of 2.0 to 10.0% wt/wt;

(6) Ceteareth-12, in the range of 0.5 to 3.0% wt/wt;

5 (7) disodium edetate, in the range of 0.01 to 0.1% wt/wt;

(8) Benzalkonium Chloride, in the range of 0.01 to 0.25% wt/wt;

(9) Anhydrous lanolin, in the range of 0.5 to 3.0% wt/wt;

(10) Cetostearyl alcohol, in the range of 1.0 to 5.0% wt/wt;

(11) Cetyl alcohol, in the range of 1.0 to 5.0% wt/wt; and

10 (12) a sufficient quantity of deionized water to form the formulation;

wherein contact with the skin results in the in situ formation of a skin protective barrier layer and wherein said sexually transmitted pathogens are viral pathogens.

[0017] In a further aspect, the invention provides a process for forming an oil-in-water formulation consisting of:

15 a water phase including (in % w/w)

(1) Deionized Water (USP) in quantities sufficient to incorporate therein ingredients 2-5;

(2) Disodium Edetate 0.01-0.1

(3) Benzalkonium Chloride 0.01-0.25

(4) Zinc Oxide 1.0-8.0

20 (5) Glycerin (USP) 2.0-10.0

and an oil phase including (in % w/w)

(6) Dimethicone 20 10.0-20.0

(7) Dimethicone 12500 3.0-10.0

(8) Cetearyl Alcohol And/Or Ceteareth-20 6.0-10.0

(9) Cetareth-12 0.5-3.0

(10) Anhydrous Lanolin 0.5-3.0

(11) Cetostearyl Alcohol 1.0-5.0

(12) Cetyl Alcohol 1.0-5.0

5 wherein said oil-in-water formulation is produced in accordance with the following steps:

1. heating to 75°C. Dimethicone 20, Dimethicone 12500, Cetearyl Alcohol, Cetareth-20, Cetareth-12, Anhydrous Lanolin, Cetostearyl Alcohol and Cetyl Alcohol to allow the materials to melt;

2. dissolving Disodium Edetate in the deionized water;

10 3. Adding Benzalkonium Chloride to Step #2;

4. Heating Step #3 to 75-78°C.;

5. Dispersing the Zinc Oxide in the Glycerin to form a homogeneous dispersion;

6. With high speed mixing, slowly adding the materials from Step #4 (Water phase) to the materials from Step #1 (Oil phase) and initiating cooling;

15 7. When at 60°C.; adding the dispersion from Step#5 to the materials from Step #6;

8. When at 50°C., reducing mixing speed;

9. When at 40°C., further reducing the mixing speed;

10. When at 25°C. stopping mixing.

20 **[0018]** In a further aspect, the invention provides an oil-in-water formulation according to the invention for use in the inhibition of the transmission of sexually transmitted pathogens, wherein said sexually transmitted pathogens are viral pathogens.

[0019] The purpose of this invention is to provide non-irritating protective formulations to be used as an adjunct in preventing the spread of a broad range of sexually transmitted diseases. The products of the present invention can be formulated

25 **[0020]** as a topical lotion, cream, solution, emulsion, or the like, and will be hereinafter referred to as creams. The use of antimicrobial/antiviral active agents and film-forming excipients in the following

formulations have demonstrated unique skin protective barrier properties with enhanced persistence that inhibit "skin to skin" and "sore to sore" transmission of pathogens (e.g., viruses, bacteria, fungi, parasites, ectoparasites and mycoplasmas) linked to communicable diseases.

[0021] Skin protectant products are regulated under CFR 21 Part 347 "Skin Protectant Drug Products for Over the Counter Human Use". The official description of a skin protectant is "a drug product that temporarily protects injured or exposed skin or mucous membrane surfaces from harmful or annoying stimuli, and may help provide relief to such surfaces." These regulations cover applicable ingredients, as well as labeling requirements for over the counter skin protectants. Active ingredients officially classified as skin protectants compositions as well as certain combinations of these compositions are listed in the CFR 21 Sec. 347.10, reproduced in Table 1 below. In accordance with the instant disclosure, the phrase "a skin protectant composition provided in a skin protecting effective concentration" will be understood to mean any of the following ingredients within their designated range of concentrations:

TABLE 1

These include any of the following within the concentration range specified:	
Ingredient	Concentration
Allantoin	0.5 to 2.0 %
Aluminum hydroxide gel	0.15 to 5.0%
Calamine	1.0 to 25.0 %
Cocoa Butter	50.0 to 100.0%
Cod liver oil	5.0 to 13.56 %(in accordance with Sec. 347.20 (a) (1) or (a) (2), provided the product is labeled so that the quantity used in a 24-hr period does not exceed 10,000 USP Units vitamin A and 400 USP Units cholecalciferol.
Colloidal oatmeal	0.007 minimum; 0.003 minimum in combination with mineral oil in accordance with Sec. 347.20 (a) (4).
Dimethicone	1.0 to 30.0%
Glycerin	20.0 to 45.0%
Hard fat	50.0 to 100.0%
Kaolin	4.0 to 20.0%
Lanolin	12.5 to 50.0%

These include any of the following within the concentration range specified:	
Ingredient	Concentration
Mineral oil	50.0 to 100.0%; 30.0 to 35.0% in combination with colloidal oatmeal in accordance with Sec. 347.20 (a) (4).
Petrolatum	30.0 to 100.0%
Topical starch	10.0 to 98.0%
White petrolatum	30.0 to 100.0%
Zinc acetate	0.1 to 2.0%
Zinc carbonate	0.2 to 2.0%
Zinc oxide	1.0 to 25.0%

[0022] It has been discovered that incorporation of at least one of the aforementioned skin protectants compositions listed in Table 1 in combination with other film forming excipients, in particular, film-forming emollients (e.g., Cetostearyl alcohol, Cetyl alcohol), silicone-containing excipients/skin protectants (e.g., polydimethylsiloxane derivatives of varying viscosities, utilized either singly or in combination, and marketed under names such as Dimethicone 20 and Dimethicone 12500), emulsifying agents (e.g. selected from Cetearyl alcohol (a mixture of fatty alcohols, predominantly stearyl and cetyl alcohols, polyoxyethylene ethers of a mixture of high molecular mass saturated fatty acids (mainly cetyl alcohol and stearyl alcohol, having a number of ethylene oxide residues in the polyoxyethylene chain, e.g. 20, 12 or the like, and referred to as Ceteareth-20, Ceteareth-12, or the like, and/or mixtures thereof), humectants (e.g. glycerin and anhydrous lanolin) and chelating compounds (e.g. disodium edetate) in a formulation for topical application results in a product that temporarily protects injured or exposed skin or mucous membrane surfaces from harmful or annoying stimuli, thereby preventing cross-contamination of pathogenic microorganisms responsible for sexually transmitted diseases (e.g., Herpes simplex virus type 1 and type 2 (HSV-1, HSV-2), human immunodeficiency virus (HIV), or the like) through skin or mucous membrane surfaces. In addition to the hydrophobic, barrier-forming formulation, active agents that prevent bacterial or viral transmission are also included. The novel formulation products of this invention are persistent in that they form a film or barrier on healthy or even damaged skin.

[0023] The hydrophobic portions of the instant formulations utilize a combination of film-forming excipients, including skin protectant(s) listed in Table 1, silicones and silicone derivatives or like equivalents, and film-forming emollients. These ingredients are dispersed by the emulsifiers throughout the continuous aqueous phase. They liquify and spread over the skin as a result of exposure to body heat. This forms a physical hydrophobic layer which resides on the skin surface (including mucosal

membranes) and provides a barrier which would inhibit penetration of liquids and pathogens which are primarily hydrophilic in nature. When used in combination with consistent and careful use of condoms, the products of the present invention help inhibit the spread of STDs and protect the skin from contamination should the condom become damaged.

5 [0024] Accordingly, it is an objective of the instant invention to teach various topical protective formulations to be used in preventing the spread of a broad range of sexually transmitted diseases.

[0025] Another objective of the present invention is to teach formulations which are condom compatible, meaning that they are latex friendly and provide effective lubricity.

10 [0026] It is yet another objective of the instant invention to teach skin protective formulations wherein contact with the skin results in the formation of a hydrophobic skin protective surface layer.

[0027] It is yet another objective of the instant invention to teach skin protective formulations wherein contact with the skin results in the formation of a mechanical barrier skin protective surface layer.

[0028] It is yet another objective of the instant invention to teach skin protective formulations wherein contact with the skin results in the formation of a hydrophobic skin protective surface layer containing
15 antiviral/antimicrobial agents.

[0029] Yet another objective of the invention is to teach products formulated as a lotion, cream, solution, emulsion, or other topically applied product.

[0030] Other objects and advantages of this invention will become apparent from the following description, wherein are set forth, by way of illustration and example, certain embodiments of this
20 invention.

DETAILED DESCRIPTION OF THE INVENTION

[0031] Zinc oxide is an inert, non-toxic chemical compound with the chemical formula of ZnO. Zinc Oxide can be used as a skin barrier. When applied to skin, Zinc Oxide acts as a mechanical barrier that physically excludes isolates and prevents the skin from any contact with harmful stimuli. Zinc Oxide is
25 often used in creams or lotions and provides a continuous barrier, which also helps prevent the loss of active ingredients due to friction and rubbing. Because of its inert, non-toxic characteristics and general non-solubility in water, it can be applied to skin as many times as may be needed. While zinc oxide is the preferred skin protectant composition of the present invention, other suitable skin protectant compositions and their amounts effective to provide skin protection as listed in Table 1 above could be
30 used herein (e.g., Dimethicone).

[0032] Glycerin (INCI name: Glycerin) is a non-irritating humectant and film former. It is also water-soluble. Moreover, glycerin is compatible with latex products and provides extended lubricity, which makes it a common base for many products designed for genital use.

[0033] Lanolin is a humectant isolated from wool-bearing animals such as sheep. It is a product of the
35 sebaceous gland and consists mostly of a mixture of cholesterol and the esters

of several fatty acids. It has many commercial uses, including within the medical and cosmetic industries.

[0034] Benzalkonium Chloride is a member of the quaternary ammonium compounds. These compounds are a group of ammonium salts in which organic radicals have been substituted for all four hydrogens of the original ammonium cation. Benzalkonium Chloride has been reported to be an effective anti-viral wetting agent in the prevention of transmission of viral diseases such as HIV and herpes simplex virus. Benzalkonium Chloride is also listed in the FDA Topical Antimicrobial Drug Products Monograph as a First Aid Antiseptic. Although the use of Benzalkonium Chloride is preferred, it is contemplated that any other quaternary ammonium salt compound could be used without departing from the scope of the present invention, such as CETRIMIDE (alkyltrimethylammonium bromide).

[0035] The phrase "film-forming emollient" refers to any excipient suitable for cosmetic and pharmaceutical applications, which forms a water-repelling film. According to the invention, the film-forming emollients are cetyl alcohol and cetostearyl alcohol. Cetyl alcohol (IUPAC name of 1-hexadecanol) is a member of the alcohol class of compounds. It is a solid organic compound and belongs to a group of fatty alcohols. In addition to its use as an emollient, it is often used in the cosmetic industry as a surfactant in shampoos, emulsifiers or thickening agents in the manufacture of skin creams and lotions. Cetostearyl alcohol is a blend of cetyl alcohol and stearyl alcohol, two fatty alcohols derived from vegetable sources.

[0036] The phrase "silicone-containing excipient" refers to any silicone or silicone derivatives, including silicone-based skin protectants, suitable for cosmetic and pharmaceutical applications, which acts as an emollient and forms a water-repelling film, including silicone oils. In addition to skin barriers, silicone and silicone oils are highly substantive on the skin. As a result of this property, silicon and silicone oils are often used in topical formulations, such as creams and lotions, to improve the substantivity of active ingredients on the skin.

[0037] After applying the topical formulations to the skin and removing volatile substances, the film formed as a result of using a silicone-containing excipient helps to maintain the active ingredient in close contact with the skin and prevents the loss of the active ingredient by abrasion. The ability to form hydrophobic films which can easily be applied and spread over skin accounts for high resistance to wash-off and rub-off. Use of silicones in topical formulations over petroleum-based products is more desirable because silicones do not exhibit the negative aesthetics of petroleum and, unlike petroleum, are known to be compatible with the materials used to manufacture condoms. According to one non-limiting embodiment, Dimethicone (preferably Dimethicone 20 cSt and Dimethicone 12500 cSt) is preferred. Dimethicone is a highly pure, non-volatile silicone fluid which is colorless and odorless. It can be used as a lubricant and emollient skin protectant.

[0038] Ceteareth-12 is part of a family with the INCI name of Ceteareth-n and refers to polyoxyethylene ethers of a mixture of high molecular mass saturated fatty alcohols. The "n" indicates the average number of ethylene oxide residues in the polyoxyethylene chain. Ceteareth-12 is a non-

toxic surfactant which is frequently used as an emulsifier in the cosmetic industry. The mixture of Cetearyl alcohol and Ceteareth-20 (EMULGADE 1000NI, Cognis Corporation) is a non-ionic, self-emulsifying base commonly used in the production of oil/water creams and lotions.

[0039] Disodium Edetate, having the chemical name of disodium (ethylene-dinitrilo) tetra-acetate

5 dihydrate, is also commonly known as the disodium salt of EDTA. The chemical acts as a chelating compound by preventing ingredients from binding to trace elements that may be present.

[0040] Excipients and antimicrobial/antiviral ingredients useful in forming the skin protective creams, according to the present invention are described in the following non-limiting example.

EXAMPLE 1

10 **[0041]** In formulating a batch of a skin protective cream according to the invention, active ingredients and excipients useful in the manufacture of this product, a particularly effective product resulted when ingredients were utilized within the following approximate ranges:

<u>ACTIVE INGREDIENTS/EXCIPIENT</u>		<u>WT/WT %</u>
		<u>RANGES</u>
Water phase		
(1)	DEIONIZED WATER	Q.S.
(2)	EDETATE DISODIUM	0.01-0.1
(3)	BENZALKONIUM CHLORIDE	0.01-0.25
(4)	ZINC OXIDE	1.0-8.0
(5)	GLYCERIN (USP)	2.0-10.0
Oil Phase		
(6)	DIMETHICONE 20	10.0-20.0
(7)	DIMETHICONE 12500	3.0-10.0
(8)	CETEARYL ALCOHOL and/or CETEARETH-20	6.0-10.0
(9)	CETEARETH-12	0.5-3.0
(10)	ANYHDROUS LANOLIN	0.5-3.0
(11)	CETOSTEARYL ALCHOHOL	1.0-5.0
(12)	CETYL ALCOHOL	1.0-5.0

[0042] A preferred, albeit non-limiting procedure for manufacture of the formulation comprises the steps of:

Procedure:

[0043]

- 5 1. Heat to 75°C -Dimethicone 20⁺, Dimethicone 12500, Cetearyl Alcohol and Cetearth-20 (available as EMULGADE 1000 NI from Cognis), Cetearth-12 (available as (EUMULGIN B1 from Cognis), Anhydrous Lanolin, Cetostearyl alcohol and Cetyl alcohol. Allow the materials to melt..
2. Dissolve Disodium Edetate in the deionized water (USP).
- 10 3. Add Benzalkonium Chloride to Step #2
4. Heat Step #3 to 75-78 °C.
5. Disperse the Zinc Oxide in the Glycerin to form a homogeneous dispersion.
6. With high speed mixing, slowly add the materials from Step #4 (Water phase) to the materials from Step #1 (Oil phase). Initiate cooling.
- 15 7. When at 60 °C; add the dispersion from Step#5 to the materials from Step #6 (Emulsion).
8. When at 50 °C, reduce mixing speed.
9. When at 40 °C, further reduce the mixing speed.
10. When at 25°C stop mixing.

[0044] While not wishing to be bound to any particular theory, it is believed that these film forming excipients and active ingredient form a neutral and hydrophobic layer barrier which is also believed to interact with the skin thereby having a stabilizing effect upon the hydrophobic layer, which results in the enhanced persistence of the product. The use of a silicone-containing skin protectant, e.g. Dimethicone 20 and Dimethicone 12500, or a like equivalent, acts in coordination with at least one of the skin protectants compositions as defined in CFR 21 Sec. 347.10, e.g. Zinc Oxide, or others from Table 1, excipients and the film-forming emollients, and melts when contacted with the heat of the body. This in turn forms a physical hydrophobic layer that provides a barrier, which appears to inhibit penetration of liquids, and sexually transmitted pathogens (viral, bacterial) which are primarily

hydrophilic in nature. This property helps protect the user from contamination due to sexual contact with infected individuals.

5

10

15

20

25

Patentkrav

1. Olie-i-vand-formulering til inhibering af overførsel af seksuelt overførte patogener, **kendetegnet ved** forbedrede hudbeskyttelsesegenskaber, hvilken formulering består af en kombination af:

- 5 (1) dimethicone 20, i området fra 10,0 til 20,0 vægt-%;
(2) dimethicone 12500, i området fra 3,0 til 10,0 vægt-%;
(3) zinkoxid, i området fra 1,0 til 8,0 vægt-%;
(4) en blanding af cetearylalkohol og ceteareth-20, i området fra 6,0 til 10,0 vægt-%;
(5) glycerin, i området fra 2,0 til 10,0 vægt-%;
10 (6) ceteareth-12, i området fra 0,5 til 3,0 vægt-%;
(7) dinatriumedetat, i området fra 0,01 til 0,1 vægt-%;
(8) benzalkoniumchlorid, i området fra 0,01 til 0,25 vægt-%;
(9) vandfri lanolin, i området fra 0,5 til 3,0 vægt-%;
(10) cetostearylalkohol, i området fra 1,0 til 5,0 vægt-%;
15 (11) cetylalkohol, i området fra 1,0 til 5,0 vægt-%; og
(12) en tilstrækkelig mængde deioniseret vand til at danne formuleringen;
hvor kontakt med huden fører til in situ-dannelse af et hudbeskyttende barrierelag, og hvor de seksuelt overførte patogener er virale patogener.

- 20 2. Olie-i-vand-formulering ifølge krav 1, der består af en kombination af:

- (1) dimethicone 20 i en koncentration på ca. 15,0 vægt-%;
(2) dimethicone 12500 i en koncentration på ca. 5,0 vægt-%;
(3) zinkoxid i en koncentration på ca. 2,0 vægt-%;
25 (4) en blanding af cetearylalkohol og ceteareth-20 i en koncentration på ca. 8,0 vægt-%;
(5) glycerin i en koncentration på ca. 5,0 vægt-%;
(6) ceteareth-12 i en koncentration på ca. 1,0 vægt-%;
(7) dinatriumedetat i en koncentration på ca. 0,05 vægt-%;
(8) benzalkoniumchlorid i en koncentration på ca. 0,2 vægt-%;
30 (9) vandfri lanolin i en koncentration på ca. 1,0 vægt-%;
(10) cetostearylalkohol i en koncentration på ca. 3,0 vægt-%;
(11) cetylalkohol i en koncentration på ca. 2,0 vægt-%; og
(12) en tilstrækkelig mængde deioniseret vand til at danne formuleringen.

3. Olie-i-vand-formulering ifølge krav 1, der består af en kombination af:

- 35 (1) dimethicone 20 i en koncentration på ca. 15,0 vægt-%;
(2) dimethicone 12500 i en koncentration på ca. 5,0 vægt-%;
(3) zinkoxid i en koncentration på ca. 5,0 vægt-%;

- (4) en blanding af cetearylalkohol og ceteareth-20 i en koncentration på ca. 7,0 vægt-%;
- (5) glycerin i en koncentration på ca. 8,0 vægt-%;
- (6) ceteareth-12 i en koncentration på ca. 1,0 vægt-%;
- (7) dinatriumedetat i en koncentration på ca. 0,5 vægt-%;
- 5 (8) benzalkoniumchlorid i en koncentration på ca. 0,2 vægt-%;
- (9) vandfri lanolin i en koncentration på ca. 1,0 vægt-%;
- (10) cetostearylalkohol i en koncentration på ca. 3,0 vægt-%;
- (11) cetylalkohol i en koncentration på ca. 2,0 vægt-%; og
- (12) en tilstrækkelig mængde deioniseret vand til at danne formuleringen.

10

4. Olie-i-vand-formulering ifølge krav 1, hvor formuleringen omfatter en vandfase, der omfatter

- (1) DEIONISERET VAND (USP) i tilstrækkelige mængder til at opløse bestanddelene 2-5;
- (2) DINATRIUMEDETAT 0,01 – 0,1
- 15 (3) BENZALKONIUMCHLORID 0,01 – 0,25
- (4) ZINKOXID 1,0 – 8,0
- (5) GLYCERIN (USP) 2,0 – 10,0
- og en oliefase, der omfatter
- (6) DIMETHICONE 20 10,0 – 20,0
- 20 (7) DIMETHICONE 12500 3,0 – 10,0
- (8) CETEARYLALKOHOL og/eller CETEARETH-20 6,0 – 10,0
- (9) CETEARETH-12 0,5 – 3,0
- (10) VANDFRI LANOLIN 0,5 – 3,0
- (11) CETOSTEARYLALKOHOL 1,0 – 5,0
- 25 (12) CETYLALKOHOL 1,0 – 5,0.

5. Olie-i-vand-formulering ifølge et hvilket som helst af kravene 1 til 4, hvor formuleringen er en creme.

30 6. Olie-i-vand-formulering ifølge et hvilket som helst af kravene 1 til 5, hvor det virale patogen er valgt fra herpes simplex virus type 1 (HSV-1), herpes simplex virus type 2 (HSV-2) og human immundeficiensvirus (HIV).

7. **Fremgangsmåde** til dannelsen af en olie-i-vand-formulering bestående af:
35 en vandfase, der omfatter (i vægt-%)

- (1) deioniseret vand (USP) i tilstrækkelige mængder til at indarbejde bestanddelene 2-5 deri;

- (2) dinatriumedetat 0,01 – 0,1
 (3) benzalkoniumchlorid 0,01 – 0,25
 (4) zinkoxid 1,0 – 8,0
 (5) glycerin (USP) 2,0 – 10,0
 5 og en oliefase, der omfatter (i vægt-%)
 (6) dimethicone 20 10,0 – 20,0
 (7) dimethicone 12500 3,0 – 10,0
 (8) cetearylalkohol og/eller cetareth-20 6,0 – 10,0
 (9) cetareth-12 0,5 – 3,0
 10 (10) vandfri lanolin 0,5 – 3,0
 (11) cetostearylalkohol 1,0 – 5,0
 (12) cetylalkohol 1,0 – 5,0.

hvor olie-i-vand-formuleringen er fremstillet i henhold til følgende trin:

- 15
1. Opvarmning til 75 °C af dimethicone 20, dimethicone 12500, cetearylalkohol, cetareth-20, cetareth-12, vandfri lanolin, cetostearylalkohol og cetylalkohol for at tillade materialerne at smelte;
 2. Opløsning af dinatriumedetat i det deioniserede vand;
 - 20 3. Tilsætning af benzalkoniumchlorid til trin 2;
 4. Opvarmningstrin 3 til 75-78 °C;
 5. Dispergering af zinkoxid i glycerin for at danne en homogen dispersion;
 6. Med højhastighedsblanding, langsom tilsætning af materialerne fra trin 4 (vandfase) til materialerne fra trin 1 (oliefase) og begyndende afkøling;
 - 25 7. Når der nås 60 °C, tilføjelse af dispersionen fra trin 5 til materialerne fra trin 6.
 8. Når der nås 50 °C, reducere af blandingshastigheden;
 9. Når der nås 40 °C, yderligere reducere af blandingshastigheden;
 10. Når der nås 25 °C, afbrydelse af blandingen.
- 30 **8.** Olie-i-vand-formulering ifølge et hvilket som helst af kravene 1 til 5 til anvendelse i inhiberingen af overførsel af seksuelt overførte patogener, hvor de seksuelt overførte patogener er virale patogener.
- 9.** Olie-i-vand-formulering ifølge krav 8, hvor det virale patogen er valgt fra herpes simplex virus
- 35 type 1 (HSV-1), herpes simplex virus type 2 (HSV-2) og human immundeficiensvirus (HIV).