

AUSTRALIA
Patents Act 1990

NOTICE OF ENTITLEMENT

I/we, ANGELINI RICERCHE S.p.A. SOCIETA' CONSORTILE
of PIAZZALE DELA STAZIONE S.N.C.
00040 S. PALOMBA - POMEZIA (ROMA)
ITALY

being the applicant in respect of Application No.47034/93, state the following:-

The nominated person acquired has entitlement from the actual inventors by way of assignment as far as actual inventor Rosario NICOLETTI is concerned and by way or employment as far as actual inventor Leandro BAIOCCHI is concerned.

The nominated person has entitlement from the applicant of the basic application listed in the declaration under Article 8 of the PCT by virtue of a merger and change of name of the basic applicant to that of the nominated person.

DATED this 27th day of February 1997

ANGELINI RICERCHE S.p.A. SOCIETA' CONSORTILE

GRIFFITH HACK & CO

GRIFFITH HACK
MELBOURNE OFFICE

Julie M. Wilkie
JULIE M. WILKIE

.....
Patent Attorneys for and
on behalf of the applicants



AU9347034

(12) PATENT ABRIDGMENT (11) Document No. **AU-B-47034/93**
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. **680200**

- (54) Title
GLYCOGEN POLYSACCHARIDES
- International Patent Classification(s)
(51)⁵ **C08B 037/18**
- (21) Application No. : **47034/93** (22) Application Date : **28.07.93**
- (87) PCT Publication Number : **WO94/03502**
- (30) Priority Data
- | | | |
|------------------|-----------------|-----------------|
| (31) Number | (32) Date | (33) Country |
| MI92A1927 | 04.08.92 | IT ITALY |
- (43) Publication Date : **03.03.94**
- (44) Publication Date of Accepted Application : **24.07.97**
- (71) Applicant(s)
ANGELINI RICERCHE S.P.A. SOCIETA CONSORTILE
- (72) Inventor(s)
ROSARIO NICOLETTI; LEANDRO BAIOCCHI
- (74) Attorney or Agent
GRIFFITH HACK , GPO Box 1285K, MELBOURNE VIC 3001
- (57) Claim


1. A glycogen polysaccharide wherein the nitrogen content is less than 60 ppm when measured by the Kjeldahl method and reducing sugars content is less than 0.25% when measured with the method of F.D. Snell and Snell.
5. A process for preparing a glycogen polysaccharide wherein the nitrogen content is less than 60 ppm when measured by the Kjeldahl method and reducing sugars content is less than 0.25% when measured with the method of F.D. Snell and Snell comprising boiling an animal tissue rich in glycogen in an aqueous solution of a strong base, cooling the broth thus obtained, adding a non-acidic, volatile solvent miscible in water, separating the precipitate formed by filtration and dissolving the said precipitate in water, characterized in that the pH of the aqueous solution of the precipitate is adjusted to neutral and the solution is treated with a cationic resin, filtered to separate the

(11) AU-B-47034/93
(10) 680200

-2-

cationic resin, treated with a non-acidic, volatile solvent miscible in water to precipitate the polysaccharide free of nitrogenous compounds and reducing sugars and filtered to recover the precipitate formed.

OPI DATE 03/03/94 APPLN. ID 47034/93
AOJP DATE 26/05/94 PCT NUMBER PCT/EP93/02006



AU9347034

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C08B 37/18	A1	(11) International Publication Number: WO 94/03502 (43) International Publication Date: 17 February 1994 (17.02.94)
(21) International Application Number: PCT/EP93/02006 (22) International Filing Date: 28 July 1993 (28.07.93) (30) Priority data: MI92A001927 4 August 1992 (04.08.92) IT (71) Applicant (for all designated States except US): ISTITUTO RICERCA FRANCESCO ANGELINI S.P.A. [IT/IT]; Piazzale della Stazione, I-00040 S. Palomba (IT). (72) Inventors; and (75) Inventors/Applicants (for US only) : NICOLETTI, Rosario [IT/IT]; Viale Parioli, 96, I-00197 Roma (IT). BAIOC- CHI, Leandro [IT/IT]; Via Belluno, 3, I-00161 Roma (IT). (71) ANGELINI RICERCHE S.p.A. SOCIETA' CONSORTILE PIAZZALE DELLA STAZIONE S.N.C 00040 S. PALOMBA - POMEZIA (ROMA) ITALY		(74) Agents: MARCHI, Massimo et al.; Marchi & Mittler s.r.l., Viale Lombardia, 20, I-20131 Milano (IT). (81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> 68 0200
(54) Title: GLYCOGEN POLYSACCHARIDES (57) Abstract Glycogen polysaccharides substantially free of nitrogenous compounds and reducing sugars.		

- 1 -

"Glycogen polysaccharides"

* * * * *

DESCRIPTION

5 This invention relates to glycogen polysaccharides, the process for their preparation and their use. More particularly, this invention relates to glycogen polysaccharides substantially free of nitrogenous compounds and reducing sugars.

10 The term "glycogen" is commonly used to mean a group of similar, but not identical, proteoglycans widespread in the animal kingdom.

15 The most investigated glycogen is the one extracted from rabbit liver and it is believed to be a protein (glycogenin) having a molecular weight of about 37,000 daltons, linked by a glycosidic link of a tyrosine to a highly branched polysaccharide of glucose having a molecular weight of 10,000,000 daltons (beta particle). Many beta particles, up to 50, can aggregate together to yield a compound (alpha particles) having a molecular weight of 500,000,000 daltons, which is the unit of native glycogen (D.J. Manners, Carbohydrate Polymers, 16, pp 37-82 (1991)).

20 The various glycogens of the different animal species show, as far as ^{it} is known, differences in the degree of the polysaccharide branching. Thus, for example, Stuart A.S. Craig et al. (Carbohydrate Research 179, pp 327-340 (1988)) reported significant differences in the polysaccharide branching of glycogen samples extracted from mammals and invertebrates.

25 Indeed, although several different processes for the extraction of glycogen from animal tissues are described in the literature, they mainly have two diff. ~~forms~~ ^{forms}:



- 2 -

- (i) a quantitative extraction of glycogen with an analytical scope for biochemistry; that is, an extraction aimed at the subsequent quantitative determination of the level of glycogen in a certain tissue;
- 5 (ii) the extraction of glycogen samples, minimizing the denaturation of the original polymer, for subsequent biochemical and conformational studies.

As a consequence, irrespective of their origin, in all the types of glycogen commercially available there is always some amount of nitrogen (500-600 ppm), which corresponds to at least that calculated for proteoglycan (D.J. Manners et al., as cited above).

From an examination of the literature, a notable difference is found between the points of view of the various authors, some of whom consider the said amount of nitrogen as a trace impurity, and others who consider it a minor constituent of glycogen. On the other hand, the various commercial sources do not specify any difference between proteoglycan glycogen and its polysaccharide.

20 While the extraction of glycogen has been thoroughly investigated, poor attention has been devoted to the extraction of glycogen polysaccharide so far. Only an old paper describes a method for the preparation of glycogen samples "free of nitrogen" from rat livers (M. Somogyi, J.Biol.Chem., 104, 245 (1934)). However, it should be noted that in this preparation glycogen undergoes an acid treatment for one night, that is under hydrolysing conditions for the glycogen polysaccharide. Moreover, the sensitivity of the analytical method used by the authors to test the nitrogen content is not known.

30 When we attempted to practise the said method it was found



- 3 -

to be poorly reproducible and the product obtained had a small, but variable, quantity of nitrogen and/or a noticeable quantity of reducing sugars (more than 0.15%) as a consequence of hydrolytic degradation.

5 Various pharmaceutical uses have been proposed for glycogen, especially as an emollient (JP-A-87-178 505) and as an excipient for its hydrating properties (JP-A-88-290 809) and in dermatological products against the skin ageing (US-5 093 109).

10 Moreover, it has been proposed to use it as a nutrient medium for bacilli which produce lactic acid, in a pharmaceutical form to regulate vaginal pH (EP-A-0 257 007). However, it is not enough stable and it is likely that traces of proteins, nucleic acids and fragments thereof give rise to sensitivity phenomena. Living contaminants could be even
15 present.

Therefore, we aimed at the preparation of a compound which maintains for the most part the structure of glycogen polysaccharide and also ensures the maximum innocuity and safety. That is to say, a compound which is free of nitrogenous
20 compounds and reducing sugars.

After many fruitless attempts, it was unexpectedly found that an aqueous solution of crude glycogen treated for a sufficiently long time with a cationic resin leaves in solution the desired polysaccharide. The polysaccharide can then be
25 easily precipitated by adding a solvent miscible in water.

Thus, the first object of this invention is to provide a glycogen polysaccharide substantially free of nitrogenous compounds and reducing sugars.

As used in this specification and in the claims appended thereto, the expression "substantially free of nitrogenous
30

compounds and reducing sugars" is intended to mean that the of
nitrogen content is less than 60 ppm as measured by the
Kjeldahl method and the reducing sugar content is less than
0.25% as measured by the method of F.D. Snell and ^{C.T.} Snell,
("Colorimetric Methods of Analysis, including some
Turbidimetric and Nephelometric methods. 3rd Edition, New York,
D. Van Nostrand Co 1954, vol.III, p 204).

27 FEB 1997

Mytilus Edulis and Mytilus Gallus Provincialis are a
particularly interesting source of glycogen; in fact, these
molluscs are found in large quantities at a moderately low cost
and have a fairly high glycogen content. Thus, the preferred
glycogen polysaccharide according to this invention is that
obtained from Mytilus Edulis and Mytilus Gallus Provincialis.

However, this invention is not limited to glycogen
polysaccharide from Mytilus Edulis and Mytilus Gallus
Provincialis. Other suitable sources of glycogen for preparing
the corresponding polysaccharide according to this invention
include other molluscs, such as oysters and Crepidula
fornicata, or the organs of animal vertebrates rich in
glycogen, such as the liver and muscle.

The glycogen polysaccharide of this invention is further
characterized by a carbon content of from about 44 to about
45%, a molecular weight of about $(2.5 \pm 0.1) \times 10^6$ daltons and a
rotatory power $(\alpha)_D^{20}$ of 197 ± 2.0 (c = 1 in water).

The first step in the method to prepare the glycogen
polysaccharide of this invention is performed according to
conventional techniques which comprise boiling the selected
tissues in an aqueous solution of a strong base, cooling the
broth thus obtained and then precipitating the glycogen by
adding a non-acidic, volatile solvent, miscible in water.

As is already well known, the quantity of glycogen contained



- 5 -

in the various tissues ranges widely, not only with regards to the tissue and the animal species but also, for the same animal tissue of the same species, with regards to various other factors such as the nutritional state and the time of the year.

5 Thus, the quantity of glycogen extracted according to the abovementioned treatment depends largely on the quantity contained in the treated animal tissues.

10 The novel step of the process of this invention lies in adjusting to neutral the pH of the solution obtained by dissolving the said precipitate in water and then treating the solution with a cationic resin.

15 Thus, a further object of this invention is to provide a process for preparing a glycogen polysaccharide substantially free of nitrogenous compounds and reducing sugars comprising boiling an animal tissue rich in glycogen in an aqueous solution of a strong base, cooling the broth thus obtained, adding a non-acidic, volatile solvent miscible in water, separating the precipitate formed by filtration and dissolving the said precipitate in water, characterized in that the pH of the aqueous solution of the precipitate is adjusted to neutral and treated with a cationic resin, filtered to separate the cationic resin, treated with a non-acidic, volatile solvent miscible in water to precipitate the polysaccharide free of nitrogenous compounds and reducing sugars and filtered to recover the precipitate formed.

25 The neutralization step is preferably performed with a weak organic acid, soluble in water such as acetic acid.

30 The treatment with a cationic resin is preferably carried out for from 8 to 48 hours under stirring and at room temperature.

- 6 -

An example of a suitable cationic resin is AmberliteTM IR-120 in acid form; other cationic resins having similar properties may also be used.

5 The cationic resin is then separated by filtration and a solvent miscible in water is added.

Examples of preferred solvents are the lower alcohols and ketons such as ethyl alcohol and acetone.

10 A precipitate is thus formed, which is the glycogen polysaccharide free of nitrogenous compounds and reducing sugars and is separated by filtration. No further purification is needed apart from the removal of the solvent.

The yield, with respect to the glycogen precipitated from the broth, is approximately quantitative.

15 Thereby, one of the further advantages of the process of this invention is that the production cost for the polysaccharide of the invention is substantially the same as the extraction cost for glycogen. In other words, for substantially the same cost, this invention yields a product which can substitute glycogen in all its known uses without
20 having the drawbacks of containing proteins, nucleic acids or their fragments and/or reducing sugars.

The following examples are intended to illustrate this invention without, however, limiting it in any way.

Example 1

25 A) Extraction of glycogen

1000 g of Mytilus Gallus Provincialis pulp were placed in a steel vessel with 1.0 l of 30% KOH and heated to 100°C for 1 hour.

30 The solution thus obtained was cooled to room temperature and 1.5 l of 95% ethyl alcohol added. The solid precipitate

- 7 -

formed (62 g) was separated by filtration.

After drying, this product had the following characteristics

C: 44.44%

N: 0.18 - 0.34%

5 B) Preparation of the glycogen polysaccharide

The solid separated in Step A was dissolved in 1 l of water, the pH of the resulting solution was adjusted to neutral with glacial acetic acid and then filtered to complete limpidity.

10 To the solution thus obtained 60 g of AmberliteTM IR-120, in acid form, were added and the mixture was maintained under stirring at room temperature for 24 hours.

The resin was separated from the solution by filtration and the glycogen polysaccharide was precipitated by adding an equal volume of 95% ethyl alcohol and then separated by filtration.

15 After drying, the glycogen polysaccharide thus prepared (61 g) exhibited the following physico-chemical characteristics;

C: 44.44%

N: absent^{*}

reducing sugars: absent^{**}

20 molecular weight: $(2.5 \pm 0.1) \times 10^{6***}$

$(\alpha)_D^{20}$: 198 ± 1.0 (c=1 in water)

Hydrolysis with 1N H₂SO₄ (3 hours at 100°C) yields only glucose (tested by gas chromatography according to M. Ochiai, J. Crom. 194, 224 (1980)).

25 * (sensitivity of method 60 ppm)

** (tested according to F.D. Snell and Snell, Colorimetric Methods of Analysis New York, 1954, vol.III, p 204, sensitivity of method 0.25%)

30 *** (calculated from the value of (eta) by applying the Flory equation with the following values $k = 1.80 \times 10^{-4}$,

- 8 -

$\alpha = 0.70$, obtained by L.P. Yu and J.E. Rolling for glycogen (J. Applied Pol. Sci. 33, 1909 (1987)).

Example 2

The glycogen polysaccharide from Mytilus Gallus Provincialis was prepared as described in part B of Example 1 except that the precipitation step was performed with acetone instead of ethyl alcohol.

Yield, 60.5 g;

(α)_D²⁰: 196 ± 1.0 (c=1 in water);

Reducing sugars: absent.

Example 3

5 g of glycogen from pig liver extracted according to Bell et al. "Biochem. J. 28, 882 (1934)" were dissolved in water (85 ml), treated with AmberliteTM IR-120 (5 g) and then precipitated with 95% ethyl alcohol (85 ml) as described in part B of Example 1.

Nitrogen and reducing sugars were absent; the rotatory power was similar to that of the glycogen polysaccharides of the Examples 1 and 2.

CLAIMS

1. A glycogen polysaccharide wherein the nitrogen content is less than 60 ppm when measured by the Kjeldahl method and reducing sugars content is less than 0.25% when measured with the method of F.D. Snell and ^{C.T.} ~~Snell~~ as identified herein.
2. A polysaccharide according to claim 1, characterized in that it has a carbon content of from about 44 to about 45% ^{by weight.}
3. A polysaccharide according to any of claims from 1 to 2, characterized in that it has a molecular weight of about $(2.5 \pm 0.1) \times 10^6$ daltons.
4. A polysaccharide according to any of claims from 1 to 3, characterized in that it has a rotatory power $(\alpha)_D^{20}$ of 197 ± 2.0 (c = 1 in water).
5. A process for preparing a glycogen polysaccharide wherein the nitrogen content is less than 60 ppm when measured by the Kjeldahl method and reducing sugars content is less than 0.25% when measured with the method of F.D. Snell and ^{C.T.} ~~as identified herein,~~ comprising boiling an animal tissue rich in glycogen in an aqueous solution of a strong base, cooling the broth thus obtained, adding a non-acidic, volatile solvent miscible in water, separating the precipitate formed by filtration and dissolving the said precipitate in water, characterized in that the pH of the aqueous solution of the precipitate is adjusted to neutral and the solution is treated with a cationic resin, filtered to separate the cationic resin, treated with a non-acidic, volatile solvent miscible in water to precipitate the polysaccharide free of nitrogenous compounds and reducing sugars and filtered to recover the precipitate formed.
6. A process according to claim 5, characterized in that the



neutralization step is performed with a weak organic acid.

7. A process according to any of claims 5 and 6, characterized in that the treatment with a cationic resin is carried out at room temperature.
- 5 8. A process according to any of claims from 5 to 7, characterized in that the treatment with a cationic resin is carried out for from 8 to 48 hours.
9. A process according to claim 8, characterized in that the treatment with a cationic resin is carried out for 24
- 10 hours.
10. A process according to any of claims from 5 to 9, characterized in that the cationic resin is AmberliteTM IR-120 in acid form.
11. A process according to any of claims from 5 to 10,
- 15 characterized in that the solvent is ethyl alcohol or acetone.

20

25



AMENDED SHEET

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 93/02006

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C08B37/18		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C08B	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	THE JOURNAL OF BIOLOGICAL CHEMISTRY vol. 25, no. 671, 1931, BALTIMORE M. SOMOGYI 'The solubility and preparation of phosphorus- and nitrogen-free glycogen' cited in the application *summary*	1-5
A	PATENT ABSTRACTS OF JAPAN vol. 12, no. 28 (C-471)(2875) 1988 & JP,A,62 178 505 (MARUZEN KASEI K.K.) 5 August 1987 cited in the application see abstract	1,6
¹⁰ Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
25 OCTOBER 1993		- 2 11. 93
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		MAZET Jean-François