

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2016206059 B2**

(54) Title
CGRP antagonist peptides

(51) International Patent Classification(s)
C07D 513/04 (2006.01) **A61P 25/06** (2006.01)
A61K 38/12 (2006.01) **C07K 7/64** (2006.01)

(21) Application No: **2016206059** (22) Date of Filing: **2016.01.06**

(87) WIPO No: **WO16/110499**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/100,371	2015.01.06	US

(43) Publication Date: **2016.07.14**

(44) Accepted Journal Date: **2019.07.25**

(71) Applicant(s)
Ferring B.V.

(72) Inventor(s)
Wisniewski, Kazimierz;Jiang, Guangcheng;Rabinovich, Aleksandr K.;Sueiras-Diaz, Javier J.

(74) Agent / Attorney
Davies Collison Cave Pty Ltd, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000, AU

(56) Related Art
LIANG Z. Y. et al., 'Discovery of potent, cyclic calcitonin gene-related peptide receptor antagonists', Journal of Peptide Science, 2011, Vol. 17, No. 5, pages 383-386



(51) International Patent Classification:

C07D 513/04 (2006.01) A61K 38/12 (2006.01)
C07K 7/64 (2006.01) A61P 25/06 (2006.01)

(21) International Application Number:

PCT/EP2016/050110

(22) International Filing Date:

6 January 2016 (06.01.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/100,371 6 January 2015 (06.01.2015) US

(71) Applicant: FERRING B.V. [NL/NL]; Polaris Avenue
144, 2132 JX Hoofddorp (NL).(72) Inventors: WISNIEWSKI, Kazimierz; 8274 Chandler
Hill Ct., San Diego, California 92127 (US). JIANG,
Guangcheng; 13330 Kibbings Road, San Diego, Califor-
nia 92130 (US). RABINOVICH, Aleksandr, K; 8234
Caminito Maritimo, La Jolla, California 92037 (US).
SUEIRAS-DIAZ, Javier, J; Apart #137, 8526 Villa La
Jolla Drive, La Jolla, California 92037-2315 (US).(74) Agent: BATES, Philip Ian; Reddie & Grose LLP, 16
Theobalds Road, London Greater London WC1X 8PL
(GB).


(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: CGRP ANTAGONIST PEPTIDES



(I)

(57) Abstract: This disclosure relates to compounds of formula (I) or a pharmaceutically acceptable salt thereof: in which m, p, A, Ar¹, Ar², Ar³, R¹, R², and R³ are defined in the specification. The compounds of formula (I) can be used as CGRP antagonists and can be used to treat

CGRP Antagonist Peptides

TECHNICAL FIELD

This disclosure relates to CGRP antagonist peptides, as well as related compositions and methods.

BACKGROUND

5 Migraine is a debilitating condition characterized by recurrent attacks of often severe throbbing headache, typically together with nausea and sensitivity to light and sound. Migraines are frequently preceded by a focal neurological symptom called an aura. Current standard of care for treating migraine is the use of the triptan class of drugs. However, approximately 30% of patients do not find relief from triptans. In
10 addition, triptans are contraindicated in migraineurs with high risk for cardiovascular diseases (e.g., diabetes, obesity, and hypercholesterolemia). Thus, there remains a need for new therapeutic paradigms for the treatment of migraine.

 Calcitonin gene related peptide (CGRP) is a 37 amino acid peptide, resulting from alternative splicing of the calcitonin gene. CGRP is implicated in many physiological
15 and pathophysiological conditions. It was discovered that truncated peptides (e.g., CGRP(8-37) or CGRP(27-37)) could act as antagonists at the CGRP receptor. These peptides were useful as research tools, but such peptides were not pursued in clinical trials. Drug discovery efforts focusing on non-peptidic small molecules yielded several compounds that advanced to clinical development, such as olcegepant and telcagepant.
20 Despite apparent effectiveness in treating migraine, these programs were all stopped, mostly due to concerns of liver toxicity. More recently, drug development efforts targeting the CGRP pathway for migraine have refocused on monoclonal antibodies against CGRP or its receptor.


 The CGRP receptor is the seven transmembrane CLR (calcitonin like receptor) in
25 complex with RAMP1 (Receptor Activity Modulating Peptide 1). In addition to CGRP receptor, CGRP also activates the adrenomedullin (AM) receptors AM1 and AM2 (CLR+RAMP2 and CLR+RAMP3, respectively) at higher concentrations. The AM

receptors are thought to have an effect on reproduction; cardiovascular and kidney function; inflammation and other conditions. A selective CGRP-R antagonist with reduced activity at the AM receptors would reduce risk of adverse events due to disruption of AM signaling.

5

SUMMARY

In one aspect, this disclosure features a compound of formula (I) or a salt thereof:



in which m is 0, 1, 2, 3, 4, or 5; p is 0, 1, 2, or 3; A is single or double carbon-carbon bond; Ar¹ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen, nitro, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, OR_a, or N(R_aR_a'), in which each R_a, independently, is H or C₁-C₄ alkyl and each R_a', independently, is H or C₁-C₄ alkyl; Ar² is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen, cyano, nitro, C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, OR_b, N(R_bR_b'), C(O)-N(R_bR_b'), or NH-C(O)-N(R_bR_b'), in which each R_b, independently, is H or C₁-C₄ alkyl and each R_b', independently, is H or C₁-C₄ alkyl; Ar³ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, OR_c, or N(R_cR_c'), in which each R_c, independently, is H or C₁-C₄ alkyl and each R_c', independently, is H or C₁-C₄ alkyl; each R¹, independently, is C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, OR_d, or C(O)-N(R_dR_d'), in which each R_d,

2016206059 03 Jul 2019

independently, is H or C₁-C₄ alkyl and each R_d¹, independently, is H or C₁-C₄ alkyl; R² is -(CH₂)_n-R, in which n is 0, 1, 2, or 3 and R is guanidino optionally substituted by CN or CH₃, aminoacyl, C₁-C₄ alkylaminoacyl, OR_e, N(R_eR_e[']), NH-C(O)-CH(NH₂)-(CH₂)₄-N(R_eR_e[']), NH-C(O)-CH₂-(OCH₂CH₂)₂-N(R_eR_e[']), or 5-membered heterocycloalkyl
5 optionally substituted with C₁-C₄ alkyl or N(R_eR_e[']), in which each R_e, independently, is H or C₁-C₄ alkyl and each R_e['], independently, is H or C₁-C₄ alkyl; and each R³, independently, is halogen, C₁-C₄ alkyl, or OR_f, in which each R_f, independently, is H or C₁-C₄ alkyl; with the provisos that, when n is 0, R is not amino or guanidino and that, when the amino acid residue bonded to Ar^lC(O) is L-Val, Ar^l is not unsubstituted phenyl.


10 In another aspect, this disclosure features a pharmaceutical composition that includes a compound of formula (I) described herein and a pharmaceutically acceptable carrier.

In still another aspect, this disclosure features a method of treating migraine that includes administering to a patient in need thereof an effective amount of the
15 pharmaceutical composition described herein.

DETAILED DESCRIPTION

This disclosure generally relates to CGRP antagonist peptides and their use for treating migraine. In particular, this disclosure is based on the unexpected discovery that certain peptides are CGRP antagonists that exhibit improved potency for CGRP receptor,
20 and can be used effectively for treating migraine. In certain embodiments, the CGRP antagonist peptides are more selective for CGRP receptor vs. AM2 receptor. In certain embodiments, the CGRP antagonist peptides have improved solubility. In certain embodiments, the CGRP antagonist peptides have improved bioavailability.

In some embodiments, the CGRP antagonist peptides described herein are those of
25 formula (I) or a pharmaceutically acceptable salt thereof:



In formula (I), m is 0, 1, 2, 3, 4, or 5; p is 0, 1, 2, or 3; A is single or double carbon-carbon bond; Ar¹ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen (e.g., F, Cl, Br, or I), nitro, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl (e.g., CH₂OH), OR_a, or N(R_aR_{a'}), in which each R_a, independently, is H or C₁-C₄ alkyl and each R_{a'}, independently, is H or C₁-C₄ alkyl; Ar² is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen, cyano, nitro, C₁-C₄ alkyl, C₁-C₄ aminoalkyl (e.g., CH₂NH₂), C₁-C₄ hydroxyalkyl, OR_b, N(R_bR_{b'}), C(O)-N(R_bR_{b'}), or NH-C(O)-N(R_bR_{b'}), in which each R_b, independently, is H or C₁-C₄ alkyl and each R_{b'}, independently, is H or C₁-C₄ alkyl; Ar³ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more substituents, each substituent independently being halogen, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, OR_c, or N(R_cR_{c'}), in which each R_c, independently, is H or C₁-C₄ alkyl and each R_{c'}, independently, is H or C₁-C₄ alkyl; each R¹, independently, is C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, OR_d, or C(O)-N(R_dR_{d'}), in which each R_d, independently, is H or C₁-C₄ alkyl and each R_{d'}, independently, is H or C₁-C₄ alkyl; R² is -(CH₂)_n-R, in which n is 0, 1, 2, or 3 and R is substituted or unsubstituted guanidino, aminoacyl (i.e., C(O)NH₂), C₁-C₄ alkylaminoacyl (e.g., C(O)NHCH₃), OR_e, N(R_eR_{e'}), NH-C(O)-CH(NH₂)-(CH₂)₄-N(R_eR_{e'}), NH-C(O)-CH₂-(OCH₂CH₂)₂-N(R_eR_{e'}), or 5-membered heterocycloalkyl optionally substituted with C₁-C₄ alkyl or N(R_eR_{e'}), in

which each R_e , independently, is H or C₁-C₄ alkyl and each $R_{e'}$, independently, is H or C₁-C₄ alkyl; and each R^3 , independently, is halogen, C₁-C₄ alkyl, or OR_f , in which each R_f , independently, is H or C₁-C₄ alkyl; with the provisos that, when n is 0, R is not amino or guanidino and that, when the amino acid residue bonded to $Ar^1C(O)$ is L-Val, Ar^1 is not unsubstituted phenyl.

The term “alkyl” refers to a saturated, linear or branched hydrocarbon moiety, such as -CH₃ or -CH(CH₃)₂. The term “cycloalkyl” refers to a saturated, cyclic hydrocarbon moiety, such as cyclohexyl. The term “heterocycloalkyl” refers to a saturated, cyclic moiety having at least one ring heteroatom (e.g., N, O, or S), such as 4-tetrahydropyranyl. The term “aryl” refers to a hydrocarbon moiety having one or more aromatic rings. Examples of aryl moieties include phenyl (Ph), phenylene, naphthyl, naphthylene, pyrenyl, anthryl, and phenanthryl. The term “heteroaryl” refers to a moiety having one or more aromatic rings that contain at least one heteroatom (e.g., N, O, or S). Examples of heteroaryl moieties include furyl, furylene, fluorenyl, pyrrolyl, thienyl, oxazolyl, imidazolyl, thiazolyl, pyridinyl, pyrimidinyl, quinazolinyl, quinolyl, isoquinolyl and indolyl.

In some embodiments, the amino acid residue bonded to $Ar^1C(O)$ can be D-Val.

In some embodiments, Ar^1 can be phenyl, pyridinyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, pyrrolyl, or triazolyl, each of which is optionally substituted with one or more substituents, such as F, Cl, NO₂, CH₃, CH₂OH, or NH₂.

In some embodiments, Ar^2 can be phenyl or pyridinyl, each of which is optionally substituted with one or more substituents, such as CH₂NH₂, C(O)NH₂, OH, CN, CH₂OH, NH₂, or NH-C(O)-NH₂.

In some embodiments, Ar^3 can be pyridinyl.

In some embodiments, R^1 can be OH, C(O)NH₂, or CH₂NH₂. In such embodiments, m in formula (I) can be 1.

In some embodiments, n in R^2 in formula (I) can be 0, 1, or 2.

In some embodiments, R can be $N(R_eR_{e'})$, $NH-C(O)-CH(NH_2)-(CH_2)_4-N(R_eR_{e'})$, $NH-C(O)-CH_2-(OCH_2CH_2)_2-N(R_eR_{e'})$, triazolyl optionally substituted with NH₂, or guanidino optionally substituted with CN or CH₃, in which each R_e , independently, is H

or C₁-C₃ alkyl and each R_e, independently, is H or C₁-C₃ alkyl. For example, R can be NH₂, N(CH₃)₂, N(CH₂CH₃)₂, NH(CH(CH₃)₂), NH-C(O)-CH(NH₂)-(CH₂)₄-N(CH₃)₂, NH-C(O)-CH₂-(OCH₂CH₂)₂-NH₂, NH-C(O)-CH₂-(OCH₂CH₂)₂-NH(CH(CH₃)₂), 3-amino-1,2,4-triazol-5-yl, or guanidino optionally substituted with CN or CH₃.

5 In some embodiments, p in formula (I) is 0.

Exemplary compounds of formula (I) (i.e., Compounds 1-70) include those listed in Table 1 below.

Table 1

Cpd #	Sequence ID
1	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
2	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
3	Picolinoyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
4	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Dhp-Phe-Cys)-3Pal-NH ₂
5	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe(3-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
6	Bz(4-F)-D-Val-Tyr-c(Cys-Dpr-Asp-Val-Gly -Pro-Phe(3-Cbm)-Cys)-3Pal-NH ₂
7	Bz(4-F)-D-Val-Tyr-c(Cys-Dpr-Asp-Val-Gly -Pro-Tyr-Cys)-3Pal-NH ₂
8	Picolinoyl-D-Val-Tyr-c(Cys-Dab(Et ₂)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
9	Picolinoyl-D-Val-Tyr-c(Cys-Dab(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
10	Picolinoyl-D-Val-Tyr-c(Cys-Dpr(CO-CH ₂ -(O-(CH ₂) ₂) ₂ -NH ₂)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
11	Oxazole-2-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
12	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
13	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(CO-CH ₂ -(O-(CH ₂) ₂) ₂ -NH-iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
14	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
15	Bz(4-F)-D-Val-Phe(2-Cbm)-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
16	Pyrimidine-4-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
17	Picolinoyl(3-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
18	Picolinoyl(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
19	Bz(4-F)-D-Val-Tyr-c(Cys-norArg(CN)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
20	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe(4-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
21	Bz(4-F)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe(3-CN)-Cys)-3Pal-NH ₂
22	Bz(4-F)-D-Val-Phe(3-CH ₂ NH ₂)-c(Cys-Arg-Asp-Val-Gly-Dhp-Tyr-Cys)-3Pal-NH ₂
23	Oxazole-2-carbonyl-D-Val-Phe(3-CH ₂ NH ₂)-c(Cys-Arg-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-NH ₂
24	Picolinoyl-D-Val-Tyr-c(Cys-Orn(Et ₂)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
25	Picolinoyl-D-Val-Tyr-c(Cys-Orn-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂

26	Picolinoyl-D-Val-Tyr-c(Cys-Arg(CN)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
27	Picolinoyl-D-Val-Tyr-c(Cys-Orn(Atz)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
28	Picolinoyl-D-Val-Tyr-c(Cys-Arg(Me)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
29	Picolinoyl(3-F)-D-Val-Tyr-c(Cys-Orn-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
30	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
31	Pyrimidine-4-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Arg-Asp-Val-Gly-Dhp-Phe(3-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
32	Picolinoyl(3-F)-D-Val-Phe(2-Cbm)-c(Cys-Arg-Asp-Val-Gly-Dhp-Phe(3-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
33	Thiazole-2-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
34	Picolinoyl(3-Me)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
35	Picolinoyl(3,5-F ₂)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
36	Picolinoyl(3-NH ₂)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
37	1H-imidazole-5-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
38	Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-3Pal-Cys)-3Pal-NH ₂
39	Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
40	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
41	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Lys(iPr)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
42	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Dab(Atz)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
43	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
44	Oxazole-2-carbonyl-D-Val-Tyr-c(Cys-Orn(Lys(Me ₂))-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
45	Pyrimidine-4-carbonyl-D-Val-Tyr-c(Cys-Orn(Lys(Me ₂))-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
46	Picolinoyl(3-F)-D-Val-Tyr-c(Cys-Orn(Me ₂))-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
47	1H-imidazole-4-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
48	1H-1,2,4-triazole-5-carbonyl(3-Me)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
49	1H-pyrrole-2-carbonyl-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
50	Picolinoyl(3-NO ₂)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
51	Picolinoyl(3-Cl)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂

52	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(Atz)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-NH ₂
53	1H-1,2,4-triazole-5-carbonyl(3-Me)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe(2-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
54	Picolinoyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-CH ₂ NH ₂)-Cys)-3Pal-NH ₂
55	Picolinoyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
56	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
57	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Tyr-Cys)-3Pal-NH ₂
58	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-3Pal-Cys)-3Pal-NH ₂
59	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-4Aph-Cys)-3Pal-NH ₂
60	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-4Uph-Cys)-3Pal-NH ₂
61	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
62	Picolinoyl(3,5-F ₂)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH ₂
63	Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
64	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
65	Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-NH ₂
66	Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(2-Cbm)-Cys)-3Pal-NH ₂
67	Oxazole-2-carbonyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-NH ₂
68	Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-NH ₂
69	Picolinoyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂
70	Oxazole-2-carbonyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(4-CH ₂ OH)-Cys)-3Pal-NH ₂

Unless specified otherwise, the amino acid code in Table 1 refers to its L-isomer. For example, Orn refers to L-ornithine, 3Pal refers to 3-(3-Pyridyl)-L-alanine, Dhp refers to 3,4-dehydro-L-proline, and Phe(2-Cbm) refers to 3-(2-carbamoyl)phenyl-L-alanine.

Exemplary Compounds 1-70 are those of formula (I), in which m is 1, p is 0, Ar³ is 3-pyridinyl, and A, Ar¹, Ar², R¹, n, and R are those shown in Table 2 below.

Table 2

Cpd #	A	Ar ¹	Ar ²	R ¹	n	R
1	single	4-fluorophenyl	phenyl	4-OH	2	guanidino
2	single	4-fluorophenyl	3-pyridinyl	4-OH	2	guanidino
3	single	2-pyridinyl	3-pyridinyl	4-OH	2	guanidino
4	double	4-fluorophenyl	phenyl	4-OH	2	guanidino
5	single	4-fluorophenyl	3-aminomethylphenyl	4-OH	2	guanidino
6	single	4-fluorophenyl	3-carbamoylphenyl	4-OH	0	NH ₂
7	single	4-fluorophenyl	4-hydroxyphenyl	4-OH	0	NH ₂
8	single	2-pyridinyl	phenyl	4-OH	1	N(Et ₂)
9	single	2-pyridinyl	phenyl	4-OH	1	NH-iPr
10	single	2-pyridinyl	phenyl	4-OH	0	NH-CO-CH ₂ -(O-(CH ₂) ₂) ₂ -NH ₂
11	single	1,3-oxazol-2-yl	3-pyridinyl	4-OH	2	guanidino
12	single	5-fluoropyridin-2-yl	phenyl	4-OH	2	NH ₂
13	single	5-fluoropyridin-2-yl	phenyl	4-OH	2	NH-CO-CH ₂ -(O-(CH ₂) ₂) ₂ -NH-iPr
14	single	5-fluoropyridin-2-yl	phenyl	4-OH	2	NH-iPr
15	single	4-fluorophenyl	phenyl	2-C(O)-NH ₂	2	guanidino
16	single	pyrimidinyl	phenyl	4-OH	2	guanidino
17	single	3-fluoropyridin-2-yl	phenyl	4-OH	2	guanidino
18	single	4-fluoropyridin-2-yl	phenyl	4-OH	2	guanidino
19	single	4-fluorophenyl	phenyl	4-OH	1	cyanoguanidino
20	single	4-fluorophenyl	4-aminomethylphenyl	4-OH	2	guanidino
21	single	4-fluorophenyl	3-cyanophenyl	4-OH	2	guanidino
22	double	4-fluorophenyl	4-hydroxyphenyl	3-CH ₂ -NH ₂	2	guanidino
23	double	1,3-oxazol-2-yl	3-carbamoylphenyl	3-CH ₂ -NH ₂	2	guanidino
24	single	2-pyridinyl	phenyl	4-OH	2	N(Et ₂)
25	single	2-pyridinyl	4-hydroxymethylphenyl	4-OH	2	NH ₂
26	single	2-pyridinyl	phenyl	4-OH	2	cyanoguanidino
27	single	2-pyridinyl	phenyl	4-OH	2	3-amino-1,2,4-triazol-5-yl
28	single	2-pyridinyl	phenyl	4-OH	2	methylguanidino
29	single	3-fluoropyridin-2-yl	4-hydroxymethylphenyl	4-OH	2	NH ₂
30	single	1,3-oxazol-2-yl	phenyl	2-C(O)-NH ₂	2	NH-iPr
31	double	pyrimidinyl	3-aminomethylphenyl	2-C(O)-NH ₂	2	guanidino
32	double	3-fluoropyridin-2-yl	3-aminomethylphenyl	2-C(O)-NH ₂	2	guanidino
33	single	2-thiazolyl	phenyl	4-OH	2	guanidino
34	single	3-methylpyridin-2-yl	phenyl	4-OH	2	guanidino
35	single	3,5-difluoropyridin-2-yl	phenyl	4-OH	2	guanidino

36	single	3-aminopyridin-2-yl	phenyl	4-OH	2	guanidino
37	single	5-imidazolyl	phenyl	4-OH	2	guanidino
38	double	5-fluoropyridin-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	NH-iPr
39	single	5-fluoropyridin-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	NH-iPr
40	single	1,3-oxazol-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	NH-iPr
41	single	1,3-oxazol-2-yl	3-pyridinyl	2-C(O)-NH ₂	3	NH-iPr
42	single	1,3-oxazol-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	3-amino-1,2,4-triazol-5-yl
43	double	1,3-oxazol-2-yl	3-aminomethylphenyl	2-C(O)-NH ₂	2	NH-iPr
44	single	1,3-oxazol-2-yl	phenyl	4-OH	2	NH-C(O)-CH(NH ₂)-(CH ₂) ₄ -N(CH ₃) ₂
45	single	pyrimidinyl	phenyl	4-OH	2	NH-C(O)-CH(NH ₂)-(CH ₂) ₄ -N(CH ₃) ₂
46	single	3-fluoropyridin-2-yl	phenyl	4-OH	2	NMe ₂
47	single	5-imidazolyl	phenyl	4-OH	2	guanidino
48	single	1,2,4-triazol-5-yl	phenyl	4-OH	2	guanidino
49	single	2-pyryl	phenyl	4-OH	2	guanidino
50	single	3-nitropyridin-2-yl	phenyl	4-OH	2	guanidino
51	single	3-chloropyridin-2-yl	phenyl	4-OH	2	guanidino
52	single	1,3-oxazol-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	3-amino-1,2,4-triazol-5-yl
53	single	1,2,4-triazol-5-yl	2-aminomethylphenyl	4-OH	2	guanidino
54	double	2-pyridinyl	3-aminomethylphenyl	2-C(O)-NH ₂	2	NH-iPr
55	single	2-pyridinyl	4-hydroxymethylphenyl	4-OH	2	NH-iPr
56	single	5-fluoropyridin-2-yl	4-hydroxymethylphenyl	4-OH	2	NH ₂
57	double	1,3-oxazol-2-yl	4-hydroxyphenyl	2-C(O)-NH ₂	2	NH-iPr
58	double	1,3-oxazol-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	NH-iPr
59	double	1,3-oxazol-2-yl	4-aminophenyl	2-C(O)-NH ₂	2	NH-iPr
60	double	1,3-oxazol-2-yl	4-ureidophenyl	2-C(O)-NH ₂	2	NH-iPr
61	single	5-fluoropyridin-2-yl	4-hydroxymethylphenyl	4-OH	2	NH-iPr
62	single	3,5-difluoropyridin-2-yl	3-pyridinyl	4-OH	2	NH-iPr
63	double	5-fluoropyridin-2-yl	3-pyridinyl	2-C(O)-NH ₂	2	NH-iPr
64	single	1,3-oxazol-2-yl	4-hydroxymethylphenyl	2-C(O)-NH ₂	2	NH-iPr
65	double	1,3-oxazol-2-yl	3-carbamoylphenyl	2-C(O)-NH ₂	2	NH-iPr
66	double	5-fluoropyridin-2-yl	2-carbamoylphenyl	2-C(O)-NH ₂	2	NH-iPr
67	double	1,3-oxazol-2-yl	3-carbamoylphenyl	4-OH	2	NH-iPr
68	double	5-fluoropyridin-2-yl	3-carbamoylphenyl	4-OH	2	NH-iPr
69	double	2-pyridinyl	4-hydroxymethylphenyl	4-OH	2	NH-iPr
70	double	1,3-oxazol-2-yl	4-hydroxymethylphenyl	4-OH	2	NH-iPr

The compounds of formula (I) can be made by methods known in the art or methods described herein. Examples 1-5 below provide detailed descriptions of how compounds 1-70 were actually prepared.

This disclosure also features pharmaceutical compositions containing a therapeutically effective amount of at least one (e.g., two or more) of the CGRP antagonist peptides described herein (i.e., the compounds of formula (I)) or a pharmaceutically acceptable salt thereof as an active ingredient, as well as at least one pharmaceutically acceptable carrier (e.g., adjuvant or diluent). Examples of pharmaceutically acceptable salts include acid addition salts, e.g., salts formed by reaction with hydrohalogen acids (such as hydrochloric acid or hydrobromic acid), mineral acids (such as sulfuric acid, phosphoric acid and nitric acid), and aliphatic, alicyclic, aromatic or heterocyclic sulfonic or carboxylic acids (such as formic acid, acetic acid, propionic acid, succinic acid, glycolic acid, lactic acid, malic acid, tartaric acid, citric acid, benzoic acid, ascorbic acid, maleic acid, hydroxymaleic acid, pyruvic acid, p-hydroxybenzoic acid, embonic acid, methanesulphonic acid, ethanesulphonic acid, hydroxyethanesulphonic acid, halobenzenesulphonic acid, trifluoroacetic acid, trifluoromethanesulphonic acid, toluenesulphonic acid, and naphthalenesulphonic acid).

The carrier in the pharmaceutical composition must be “acceptable” in the sense that it is compatible with the active ingredient of the composition (and preferably, capable of stabilizing the active ingredient) and not deleterious to the subject to be treated. One or more solubilizing agents can be utilized as pharmaceutical carriers for delivery of an active CGRP antagonist peptide. Examples of other carriers include colloidal silicon oxide, magnesium stearate, cellulose, sodium lauryl sulfate, and D&C Yellow # 10.

The pharmaceutical composition described herein can optionally include at least one further additive selected from a disintegrating agent, binder, lubricant, flavoring agent, preservative, colorant and any mixture thereof. Examples of such and other additives can be found in “Handbook of Pharmaceutical Excipients”; Ed. A.H. Kibbe, 3rd Ed., American Pharmaceutical Association, USA and Pharmaceutical Press UK, 2000.

The pharmaceutical composition described herein can be adapted for parenteral, oral, topical, nasal, rectal, buccal, or sublingual administration or for administration via

the respiratory tract, e.g., in the form of an aerosol or an air-suspended fine powder. The term "parenteral" as used herein refers to subcutaneous, intracutaneous, intravenous, intramuscular, intraarticular, intraarterial, intrasynovial, intrasternal, intrathecal, intralesional, intraperitoneal, intraocular, intra-aural, or intracranial injection, as well as any suitable infusion technique. In some embodiments, the composition can be in the form of tablets, capsules, powders, microparticles, granules, syrups, suspensions, solutions, nasal spray, transdermal patches or suppositories.

In some embodiments, the pharmaceutical composition described herein can contain a CGRP antagonist peptide described herein that is dissolved in an aqueous solution. For example, the composition can include a sodium chloride aqueous solution (e.g., containing 0.9 wt% of sodium chloride) to serve as a diluent.

In addition, this disclosure features a method of using a CGRP antagonist peptide as outlined above for treating migraine or for the manufacture of a medicament for such a treatment. The method can include administering to a patient in need thereof an effective amount of the pharmaceutical composition described herein. "An effective amount" refers to the amount of the pharmaceutical composition that is required to confer a therapeutic effect on the treated subject. Effective doses will vary, as recognized by those skilled in the art, depending on the types of diseases treated, route of administration, excipient usage, and the possibility of co-usage with other therapeutic treatment.

As used herein, the terms "treatment," "treat," and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of, a migraine or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

The typical dosage of the CGRP antagonist peptide described herein can vary within a wide range and will depend on various factors such as the individual needs of

each patient and the route of administration. Exemplary daily dosages (e.g., for subcutaneous administration) can be at least about 0.5 mg (e.g., at least about 1 mg, at least about 5 mg, at least about 10 mg, or at least about 15 mg) and/or at most about 100 mg (e.g., at most about 75 mg, at most about 50 mg, at most about 20 mg, or at most about 15 mg) of a CGRP antagonist peptide. The skilled person or physician may consider relevant variations to this dosage range and practical implementations to accommodate the situation at hand.

In some embodiments, the pharmaceutical composition described herein can be administered once daily. In some embodiments, the pharmaceutical composition can be administered more than once daily (e.g., twice daily, three times daily, or four times daily).

The contents of all publications cited herein (e.g., patents, patent application publications, and articles) are hereby incorporated by reference in their entirety.

The following examples are illustrative and not intended to be limiting.

Examples

General Synthetic Methods

1. Amino acid derivatives

Amino acid derivatives were purchased from commercial providers (such as Aapptec, Chem Impex International, EMD Millipore, PPL, PepTech and Peptides International), except for Fmoc-Orn(iPr,Boc)-OH. Fmoc-Orn(iPr,Boc)-OH was prepared as follows:

50.0 g (105.8 mmol) of Fmoc-Orn(Boc)-OH was dissolved in 100 mL of dichloromethane (DCM). 100 mL of trifluoroacetic acid (TFA) was subsequently added. The reaction mixture was magnetically stirred for 1 hour and the solvents were evaporated. To remove excess TFA, the residue was reconstituted in DCM and evaporated several times. The oily residue was dissolved in 400 mL of MeOH and 100 mL of acetone, followed by 30 mL of acetic acid. The reaction mixture was vigorously stirred and 120.0 g (0.57 mol, 5.4 eq) of solid NaBH(OAc)₃ was added in 10 g portions until Fmoc-Orn-OH was consumed (about 2 hours, monitored by analytical HPLC). The

solvents were then evaporated and the resulting solid residue was used in the next step without purification.

The residue obtained in the previous step was dissolved in 100 mL of water and the pH of the solution was adjusted to about 9.5 with solid Na₂CO₃. 100 mL of t-BuOH was subsequently added to the magnetically stirred reaction mixture. Boc₂O (60.0 g, 275 mmol, 2.6 eq) in 100 mL of t-BuOH was then added portionwise over 10 hours. The pH of the reaction mixture was maintained at about 9.5 with the addition of saturated Na₂CO₃ (aq). After the last portion of Boc₂O was added, the reaction mixture was stirred for 9 more hours. The reaction mixture was diluted with 1 L of water and extracted with 2 x 200 mL of hexane. The water phase was acidified with 2 M HCl and the product was extracted with diethyl ether (3 x 300 mL). The combined organic extracts were thoroughly washed with 2 M HCl (3 x 200 mL) and water, and were subsequently dried over anhydrous MgSO₄. The drying agent was filtered off and the solvent was evaporated. The resulting solid residue was treated with petroleum ether, decanted and dried *in vacuo*. The crystalline product was dissolved in 200 mL of t-BuOH and lyophilized. 41.8 g (84 mmol, 79.5% yield) of the lyophilized derivative was obtained.

2. Peptide Synthesis

Resins were purchased from commercial suppliers (e.g., PCAS BioMatrix Inc. and EMD Millipore). Carboxylic acids for the N-terminal acyl group introduction were obtained from AstaTech, ChemBridge Corp. Frontier Scientific, J&W Pharmalab, Oakwood Products and TCI America. All additional reagents, chemicals and solvents were purchased from Sigma-Aldrich and VWR.

The compounds described herein were synthesized by standard methods in solid phase peptide chemistry utilizing Fmoc methodology. The peptides were assembled either manually or automatically using a Tribute peptide synthesizer (Protein Technologies Inc., Tucson, Arizona) or an Applied Biosystems 433A peptide synthesizer, or by combination of manual and automatic syntheses.

Preparative HPLC was performed on a Waters Prep LC System using a PrepPack cartridge Delta-Pack C18, 300Å, 15 µm, 47 x 300 mm at a flow rate of 100 mL/min

and/or on a Phenomenex Luna C18 column, 100Å, 5 µm, 30 x 100 mm at a flow rate of 40 mL/min. Analytical reverse phase HPLC was performed on an Agilent Technologies 1200rr Series liquid chromatograph using an Agilent Zorbax C18 column, 1.8 µm, 4.6 x 110 mm at a flow rate of 1.5 mL/min. Final compound analyses were performed on an
5 Agilent Technologies 1200 Series chromatograph by reverse phase HPLC on a Phenomenex Gemini 110Å C18 column, 3 µm, 2 x 150 mm at a flow rate of 0.3 mL/min. Mass spectra were recorded on a MAT Finnigan LCQ electrospray mass spectrometer. Unless stated otherwise, all reactions were performed at room temperature. The following references provides further guidance on general experimental set up, as well as
10 on the availability of required starting material and reagents: Kates, S.A., Albericio, F., Eds., Solid Phase Synthesis: A Practical Guide, Marcel Dekker, New York, Basel, 2000; Greene, T.W., Wuts, P.G.M., Protective Groups in Organic Synthesis, John Wiley Sons Inc., 2nd Edition, 1991; Stewart, J.M., Young, J.D., Solid Phase Synthesis, Pierce Chemical Company, 1984; Bisello, et al., J. Biol. Chem. 1998, 273, 22498-22505;
15 Merrifield, J. Am. Chem. Soc. 1963, 85, 2149-2154; and Chang and White P.D., 'Fmoc Solid Phase Peptide Synthesis: a Practical Approach', Oxford University Press, Oxford, 2000.

The following protecting groups were utilized to protect the given amino acid side chain functional groups: Pbf (2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl) for
20 Arg; tBu (t-butyl) for Tyr and Asp; Boc (t-butoxycarbonyl) for Dab, Orn, Orn(iPr) and Lys; and Trt (trityl) for Cys.

Couplings of Fmoc-protected amino acids on the Tribute synthesizer were mediated with HBTU/NMM in DMF except for cysteine derivatives that were coupled with DIC/HOBt in DMF. Single cycles of 30-60 minutes with a 5-fold excess of
25 activated Fmoc-protected amino acids were used during the synthesis. Removal of the Fmoc protecting group was monitored by UV. Multiple (up to 10 times, as needed) two-minute washes of the peptide resin with 20% piperidine in DMF were performed.

Cycle protocols specified by Applied Biosystems were used on the 433A synthesizer. Couplings were mediated with HATU/DIPEA or DIC/HOBt in DMF/NMP.
30 Single couplings of 35-50 minutes with a 4-fold excess of activated Fmoc-protected

amino acids were employed. Removal of the Fmoc protecting group was monitored by UV and was achieved by a single 20-minute wash with 20% piperidine/NMP.

DIC/HOBt mediated couplings in DMF were employed for all amino acids in manual mode. Single cycles of at least 2 hours with up to 3-fold excess of activated Fmoc-protected amino acids were used during the synthesis. The completeness of couplings was assessed with ninhydrine (Kaiser) test. Removal of the Fmoc protecting group was achieved with a single 30-minute wash of the peptide resin with 20% piperidine in DMF.

Upon completion of the peptide synthesis, the peptide resins were washed with DCM and dried *in vacuo*. The resins were treated with TFA containing variable amounts of H₂O (up to 10%) and diisopropylsilane (TIS; up to 4%) for 2 hours to remove the side-chain protecting groups with concomitant cleavage of the peptide from the resin. The peptides were filtered, precipitated with diethyl ether and decanted. To obtain peptides with disulfide bridges, the precipitate was dissolved in neat TFA or AcOH and the solution was subsequently poured into 10 % acetonitrile in water. In some cases an additional amount of acetonitrile was added to solubilize the substrate. The linear peptide was oxidized with 0.1M I₂ in MeOH or AcOH. The oxidizer solution was added dropwise until yellow color persisted. The excess of iodine was reduced with ascorbic acid. The pH was then adjusted to about 4 with concentrated ammonia. The obtained solution was loaded directly onto an HPLC prep column and eluted with a gradient of Component B shown in Table 3 below.

Each crude peptide was purified with Buffer T shown in Table 3. The fractions with a purity exceeding 90%, determined by reverse-phase analytical HPLC, were pooled and reloaded onto the column and eluted with Buffer T to provide trifluoroacetate salts. In some cases, an additional purification with Buffer C shown in Table 3 was performed. To obtain hydrochloride salts, the fractions from runs with Buffer T or C were reloaded onto the column and the column was washed with 3-5 volumes of 0.1 M sodium chloride in 1 mM HCl. The final product was eluted with Buffer H shown in Table 3. The fractions were pooled and lyophilized. The compounds thus prepared were typically found to be at least about 90% pure.

Table 3. Prep HPLC Buffer Compositions

Buffer	Component A	Component B
C	0.25 M Triethylammonium Perchlorate, pH 2.3	60% acetonitrile, 40% Component A
T	0.1% Trifluoroacetic acid (TFA)	60% acetonitrile, 0.1% TFA
H	1 mM HCl	60% acetonitrile, 1 mM HCl

Syntheses of certain exemplary compounds of formula (I) described herein are provided below.

5 **Example 1: Synthesis of Compound 30**

The peptide was assembled manually starting from 3.0 g (1.95 mmol) of Fmoc-Rink amide MBHA resin (EMD Millipore, catalog number 855003, 0.65 mmol/g). DIC/HOBt mediated couplings in DMF were employed. Single cycles of at least 2 hours with up to 3-fold excess of activated Fmoc-protected amino acids were used during the

10 synthesis. The completeness of couplings was assessed with ninhydrine test. Removal of the Fmoc protecting group was achieved with a single 30-minute wash of the peptide resin with 20% piperidine in DMF. The following amino acid derivatives were used to assemble the resin-bound peptide: Fmoc-3Pal-OH, Fmoc-Cys(Trt)-OH, Fmoc-Phe-OH, Fmoc-Pro-OH, Fmoc-Gly-OH, Fmoc-Val-OH, Fmoc-Asp(tBu)-OH, Fmoc-Orn(iPr,Boc)-

15 OH, Fmoc-Cys(Trt)-OH, Fmoc-Phe(2-Cbm)-OH and Fmoc-D-Val-OH. After the 1-11 peptide fragment was assembled, the resin was capped with oxazole-2-carboxylic acid/DIC/HOBt (4 eq), washed thoroughly with DCM and dried *in vacuo*. The crude linear peptide was cleaved from the resin with 50 mL of TFA/H₂O/TIS 96:2:2 (v/v/v) for 2 hours. After the solvent was evaporated, the crude peptide was precipitated with

20 diethyl ether and decanted. The precipitate was dissolved in 1 L of 1% aqueous TFA and oxidized with 0.1M I₂/MeOH. The oxidizer solution was added dropwise until yellow color persisted. The excess iodine was reduced with solid ascorbic acid. The pH was then adjusted to about 4 with concentrated ammonia. The obtained solution was loaded directly onto an HPLC prep column and purified with Buffer T. The fractions with purity

>90%, determined by reverse-phase analytical HPLC, were pooled and reloaded onto the column. The column was equilibrated with 1 mM HCl, washed with 3 volumes of 0.1 mM NaCl in 1 mM HCl and the compound was eluted with Buffer H to provide hydrochloride salt. The fractions were pooled and lyophilized. 1009.8 mg (0.63 mmol, 32.3% overall based on 89.6% peptide content) of white peptide powder (Compound 30) was obtained.

The product purity was determined by analytical HPLC as 90.7%. The observed and calculated MS data (i.e., M+H) are provided in Table 4 below.

Example 2: Synthesis of Compound 40

The solid phase synthesis of this peptide was performed on the Tribute Peptide Synthesizer using Fmoc-strategy. The starting resin was 0.23 g (0.15 mmol) of Rink Amide MBHA resin (EMD Millipore, catalog number 855003, 100-200 mesh, 0.65 mmol/g). DIC/HOBt mediated couplings in DMF were employed for all amino acids except for the N-terminal oxazole-2-carboxylic acid that required HBTU/NMM coupling method. Single cycles of 2 hours with 3-fold excess of activated Fmoc-protected amino acids were used during the synthesis. Fmoc protecting group was removed by treatment with 20% piperidine in DMF, 1 × 5 min and 1 × 25 min. The following amino acid derivatives were used consecutively to assemble the resin-bound peptide: Fmoc-3Pal-OH, Fmoc-Cys(Trt)-OH, Fmoc-3Pal-OH, Fmoc-Pro-OH, Fmoc-Gly-OH, Fmoc-Val-OH, Fmoc-Asp(tBu)-OH, Fmoc-Orn(iPr,Boc)-OH, Fmoc-Cys(Trt)-OH, Fmoc-Phe(2-Cbm)-OH and Fmoc-D-Val-OH. The N-terminal acyl group was introduced by treating the resin-bound (1-11) peptide fragment with pre-activated mixture of oxazole-2-carboxylic acid (0.5 mmol), HBTU (0.5 mmol), and DIEA (1.0 mmol) in DMF for 4 hours. The final assembled peptide resin was washed with DCM and dried *in vacuo*. The crude linear peptide was cleaved from the resin with 25 mL of TFA/H₂O/TIS (94:3:3, v/v/v) for 2.5 hours. The solvent was evaporated under vacuum and the crude peptide was precipitated with diethyl ether. The precipitate was collected by filtration and then dissolved in 400 mL of 0.1% TFA in 5% ACN and oxidized with 0.1M I₂/AcOH. The iodine solution was added dropwise until yellow color persisted. The excess of iodine

was reduced with a saturated solution of ascorbic acid in water. The resulting solution was loaded directly onto a preparative HPLC column and purified with Buffer T. The fractions with purity >90%, determined by reverse-phase analytical HPLC, were pooled and freeze dried on a lyophilizer. 80.2 mg (45.0 μ mol, 30% overall yield based on 80%
5 estimated peptide content) of white peptide powder (Compound 40) was obtained.

The product purity was determined by analytical HPLC as 96.8%. The observed and calculated MS data (i.e., M+H) are provided in Table 4 below.

Example 3: Synthesis of Compound 62

The peptide was assembled manually starting from 3.0 g (1.77 mmol) of Rink
10 amide MBHA resin (Novabiochem, catalog number 8.55003, 0.59 mmol/g), using high-temperature SPPS (75°C, Lauda E100 water bath, jacketed 50 mL SPPS reaction vessel). Single cycles of at least 15 minutes with up to 4-fold excess of preactivated Fmoc-protected amino acids (HOBt, DIC, no preactivation for Fmoc-Orn(iPr,Boc)-OH) were used during the synthesis. Fmoc protecting group was removed with a 2x 5-minute wash
15 of the peptide resin with 25% piperidine in DMF. The following amino acid derivatives were used to assemble the resin-bound peptide: Fmoc-3Pal-OH, Fmoc-Cys(Trt)-OH, Fmoc-Phe-OH, Fmoc-Pro-OH, Fmoc-Gly-OH, Fmoc-Val-OH, Fmoc-Asp(tBu)-OH, Fmoc-Orn(iPr,Boc)-OH, Fmoc-Cys(Trt)-OH, and Fmoc-D-Val-OH. After the 1-11 peptide fragment was assembled, the resin was capped with 3,5-difluoropicolinic
20 acid/HATU/DIPEA (4 eq), washed thoroughly with MeOH, and dried *in vacuo*. The crude linear peptide was cleaved from the resin with 75 mL of TFA/H₂O/TIS 96:2:2 (v/v/v) for 2 hours. After the solvent was evaporated, the crude peptide was precipitated with diethyl ether and decanted. The precipitate was dissolved in 1 L of 10% MeCN in 0.5% aqueous TFA and oxidized with 0.05M I₂/AcOH. The oxidizer solution was added
25 dropwise until yellow color persisted. The excess of iodine was reduced with 1 M ascorbic acid. The obtained solution was loaded directly onto an HPLC prep column and purified with modified Buffer T (Component A: 0.01% TFA, Component B: 95% acetonitrile in 0.01% TFA). The fractions with purity >95%, determined by reverse-phase analytical HPLC, were pooled and reloaded onto the column. The column was

equilibrated with 1 mM HCl, washed with 3 volumes of 0.1 mM NaCl in 1 mM HCl and the compound was subsequently eluted with Buffer H to provide hydrochloride salt. The fractions were pooled and lyophilized. 583 mg (0.63 mmol, 20% overall based on 87.3% peptide content and 98.8% purity) of white peptide powder (Compound 62) was obtained.

5 The product purity was determined by analytical HPLC as 98.8%. The observed and calculated MS data (i.e., M+H) are provided in Table 4 below.

Example 4: Synthesis of Compound 65

10 The compound was assembled on solid phase by a combination of manual and automatic syntheses. First, the C-terminal tripeptide was synthesized manually starting from 7.3 g (3.5 mmol) of Fmoc-Rink amide Chem Matrix resin (Biotage, catalog number 7-600-1310-25, 0.48 mmol/g). HATU/DIPEA mediated couplings in DMF were employed for 3Pal and Phe(3-Cbm), and DIC/HOBt mediated coupling in DMF was used for Cys. Single cycles of at least 2 hours with up to 3-fold excess of activated Fmoc-protected amino acids were used during the synthesis. The completeness of couplings was assessed with ninhydrine test. Removal of the Fmoc protecting group was achieved with 30% piperidine in DMF using two washes of 5 and 25 minutes, respectively. The following amino acid derivatives were used to assemble the resin-bound peptide: Fmoc-3Pal-OH, Fmoc-Cys(Trt)-OH and Fmoc-Phe(3-Cbm)-OH. The synthesis was continued on the 433A Synthesizer with one eighth (0.44 mmol) of the resin. HATU/DIPEA or 15 DIC/HOBt (for Cys) mediated couplings in NMP/DMF were employed. Single cycles of at least 30 minutes with up to 5-fold excess of activated Fmoc-protected amino acids were used during the synthesis. Removal of the Fmoc protecting group was achieved with a single 30-minute wash of the peptide resin with 20% piperidine in NMP. The following amino acid derivatives were used to finish the assembly of the resin-bound peptide: Fmoc-Dhp-OH, Fmoc-Gly-OH, Fmoc-Val-OH, Fmoc-Asp(tBu)-OH, Fmoc-Orn(iPr,Boc)-OH, Fmoc-Cys(Trt)-OH, Fmoc-Phe(2-Cbm)-OH and Fmoc-D-Val-OH. 20 After the 1-11 peptide fragment was assembled, the resin was capped manually with oxazole-2-carboxylic acid/HATU/DIPEA (4 eq), washed thoroughly with DCM, and dried *in vacuo*. The crude linear peptide was cleaved from the resin with 50 mL of

TFA/H₂O/TIS 90:8:2 (v/v/v) for 2 hour. The solvent was evaporated and the crude peptide was precipitated with MTBA, centrifuged and decanted. The precipitate was dissolved in 15 mL of AcOH and poured into 250 mL of 10% (v/v) aqueous acetonitrile and oxidized with 0.1M I₂/MeOH. The oxidizer solution was added dropwise until yellow color persisted. The excess of iodine was reduced with solid ascorbic acid. The pH was then adjusted to about 4 with concentrated ammonia. The obtained solution was loaded directly onto an HPLC prep column and purified with Buffer T (see table above). The fractions with purity >90%, determined by reverse-phase analytical HPLC, were pooled and lyophilized. 116.2 mg (0.06 mmol, 14.1% overall based on 78.5% peptide content) of white peptide powder (Compound 65) was obtained.

The product purity was determined by analytical HPLC as 98.3%. The observed and calculated MS data (i.e., M+H) are provided in Table 4 below.

Example 5: Synthesis of Compounds 1-29, 31-39, 41-61, 63, 64, and 66-70

Compounds 1-29, 31-39, 41-61, 63, 64, and 66-70 were synthesized by using the methods described in Examples 1-4.

The observed and calculated MS data (i.e., M+H) of Compounds 1-70 are summarized in Table 4 below.

Table 4

Compound No.	Calculated M+H	Observed M+H
1	1425.6	1425.6
2	1426.6	1426.6
3	1408.6	1408.6
4	1423.6	1423.5
5	1454.6	1454.6
6	1398.5	1398.5
7	1371.5	1371.5
8	1408.6	1408.8
9	1394.6	1394.7
10	1628.7	1628.9
11	1398.6	1398.8
12	1384.6	1384.7
13	1571.7	1571.8

14	1426.6	1426.7
15	1452.6	1452.7
16	1409.6	1409.7
17	1426.6	1426.7
18	1426.6	1426.7
19	1436.6	1436.7
20	1454.6	1454.6
21	1450.6	1450.6
22	1452.6	1452.7
23	1452.6	1452.7
24	1422.6	1422.7
25	1396.6	1396.7
26	1433.6	1433.7
27	1448.6	1488.7
28	1422.6	1422.8
29	1414.6	1414.7
30	1425.6	1425.7
31	1463.6	1463.8
32	1480.6	1480.7
33	1414.5	1414.7
34	1422.6	1422.7
35	1444.6	1444.7
36	1423.6	1423.6
37	1397.6	1397.7
38	1452.6	1452.6
39	1454.6	1454.6
40	1426.6	1426.5
41	1440.6	1440.6
42	1452.6	1452.5
43	1452.6	1452.6
44	1512.7	1512.7
45	1523.7	1523.8
46	1412.6	1412.7
47	1397.6	1397.6
48	1412.6	1412.7
49	1396.6	1396.6
50	1453.6	1453.6
51	1442.6	1442.6
52	1466.6	1466.6
53	1441.6	1441.8

54	1462.6	1462.7
55	1438.6	1438.8
56	1414.6	1414.9
57	1439.6	1439.8
58	1424.6	1424.7
59	1438.6	1438.9
60	1481.6	1481.7
61	1456.6	1456.7
62	1444.6	1444.7
63	1481.6	1481.6
64	1455.6	1455.6
65	1466.6	1466.7
66	1494.6	1494.7
67	1439.6	1439.6
68	1467.6	1467.7
69	1436.6	1436.7
70	1426.6	1426.5

Example 6: CGRP receptor antagonist activity measured by cAMP assay

CGRP receptor agonists increase intracellular cyclic adenosine mono-phosphate
 5 (cAMP). CGRP receptor antagonists can reduce the agonist effect. Antagonist activity
 was assessed by measurement of cyclic adenosine mono-phosphate (cAMP) using cell
 line stably expressing the hCGRP receptor (GeneBLAzer® CALCRL:RAMP1-CRE-bla
 Freestyle™ 293F, Invitrogen). hCGRP receptor expressing cells were maintained in
 DMEM high-glucose with GlutaMAX™ containing 10% (v/v) FBS, 0.1 mM NEAA,
 10 25 mM HEPES, 5 ug/ml Blasticidin, 100 ug/ml Hygromycin, and 400 ug/ml Geneticin at
 37°C under 5% CO₂ in a humidified atmosphere. For cAMP measurement, cells were
 washed once with 5 ml 1X PBS, cell maintenance media was replaced with compound
 buffer ((CB): DMEM containing 0.1% BSA and 0.5 mM IBMX), and flasks were
 incubated for 1 hour at 37°C under 5% CO₂ in a humidified atmosphere. Cells were
 15 removed from culture flasks using non-enzymatic cell dissociation buffer and harvested
 in CB. The reaction was performed in 384 well white small volume plates (Greiner) at a
 density of 10,000 cells/well. Cells were exposed to varying concentrations of antagonist

compounds for 30 minutes in the presence of a fixed concentration of agonist (human α -CGRP). cAMP levels were measured using an HTRF (homogeneous time resolved fluorescence)-based competitive cAMP immunoassay (cAMP Dynamic 2 Kit, Cisbio), according to the manufacturer's instructions. The ratio of 665 nm and 615 nm time-resolved fluorescence readings (RFU) was calculated and a single-binding site, four parameter concentration response model: $(MIN + ((MAX - MIN) / (1 + ((EC50/x)^{Hill}))))$, was used to perform non-linear regression analysis, generating the concentration response curve. Reported parameters include antagonist potency IC₅₀ (the concentration causing half-maximal inhibition of the agonist response for antagonist compounds) and efficacy (%MPE: percent of the maximal possible effect).

Compounds 1-70 and three reference peptide compounds were tested in the above assay. The three references peptide compounds are: (1) Bz(4-F)-D-Val-Tyr-c(Cys-Agp-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂ ("Reference Compound 1", i.e., Compound 36 in Yan et al., *J. Pept. Sci.* 2011, 17, 383-386), (2) Bz-D-Val-Tyr-c(Cys-Dpr-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂ ("Reference Compound 2", Compound 33 in Yan et al., *J. Pept. Sci.* 2011, 17, 383-386), and (3) H-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂ ("Reference Compound 3", human α -CGRP(8-37)-NH₂ antagonist). The results are summarized in Table 5 below.

As shown in Table 5, Compounds 1-70 generally exhibited improved potency compared to Reference Compounds 1-3.

Example 7: AM2 receptor antagonist activity measured by cAMP assay

Antagonist activity for the adrenomedullin receptor AM2 was determined using the method described in Example 6 above, with the following modifications. Instead of GeneBLAzer® CALCRL:RAMP1-CRE-bla Freestyle™ 293F cells, GeneBLAzer® CALCRL:RAMP3-CRE-bla Freestyle™ 293F cells were used to test activity at hAM2 receptors. The agonist was human adrenomedullin, instead of α -CGRP.

Compounds 1-70 and the three reference peptide compounds described in Example 6 were tested in this assay. The results are summarized in Table 5 below. In

Table 5, the selectivity ratio for hCGRP over hAM2 is calculated as hCGRP-R IC_{50} /hAM2-R IC_{50} .

As shown in Table 5, a majority of Compounds 1-70 exhibited improved selectivity to hCGRP receptor over hAM2 receptor compared to Reference Compounds 1-3.

Table 5.

Patent Cpd No.	hCGRP-R		hAM2-R		
	IC50 Ave (nM)	Efficacy Antag (%) Ave	IC50 Ave (nM)	Efficacy Antag (%) Ave	selectivity ratio CGRP/AM2
1	0.14	100	52	101	362
2	0.08	100	45	100	555
3	0.12	100	35	101	285
4	0.17	100	29	113	172
5	0.13	100	26	99	192
6	0.17	100	101	113	612
7	0.13	100	43	100	334
8	0.09	100	30	100	347
9	0.15	100	51	106	351
10	0.19	100	114	107	589
11	0.13	100	25	108	198
12	0.11	100	18	100	172
13	0.10	100	26	100	249
14	0.12	100	20	100	167
15	0.10	100	58	99	610
16	0.07	100	6.8	101	93
17	0.05	100	7.1	100	148
18	0.16	100	8.1	98	52
19	0.19	100	48	100	248
20	0.18	100	54	100	309
21	0.13	100	33	99	258
22	0.19	100	33	101	175
23	0.05	100	21	100	400
24	0.11	100	128	98	1213
25	0.07	100	78	100	1071
26	0.06	100	38	100	636
27	0.15	100	119	98	765
28	0.07	100	68	99	1031
29	0.09	100	148	98	1699
30	0.11	100	700	98	6628
31	0.12	100	46	100	390
32	0.12	100	61	99	498

33	0.10	100	11	100	117
34	0.14	100	7.9	100	58
35	0.06	100	33	101	580
36	0.17	100	16	101	93
37	0.17	100	379	100	2295
38	0.06	100	87	101	1478
39	0.13	100	275	101	2125
40	0.09	100	980	98	10884
41	0.13	100	1020	97	8075
42	0.14	100	1225	97	9068
43	0.04	100	180	101	4693
44	0.17	100	114	100	668
45	0.18	100	198	98	1082
46	0.15	100	184	99	1189
47	0.16	100	232	99	1460
48	0.07	100	402	96	5610
49	0.19	100	89	99	461
50	0.18	100	8.1	99	46
51	0.14	100	13	100	92
52	0.15	100	770	97	5203
53	0.11	100	901	98	8130
54	0.17	100	150	99	898
55	0.18	100	137	100	746
56	0.17	100	55	100	328
57	0.12	100	624	99	5225
58	0.18	100	476	99	2588
59	0.17	100	467	99	2770
60	0.12	100	444	100	3759
61	0.10	100	89	100	934
62	0.14	100	85	100	592
63	0.10	101	85	98	848
64	0.12	101	1246	99	10594
65	0.06	101	318	100	5492
66	0.08	101	84	100	1027
67	0.09	101	103	99	1094
68	0.06	101	18	100	300
69	0.13	100	64	100	497
70	0.15	100	107	100	727
Ref. Cpd. 1	0.20	100	42	98	210
Ref. Cpd. 2	0.52	100	200	100	387
Ref. Cpd. 3	4.1	100	35	101	9

Other embodiments are within the scope of the following claims.

2016206059 03 Jul 2019


Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

2016206059 03 Jul 2019

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound of formula (I) or a pharmaceutically acceptable salt thereof:



5 wherein

m is 0, 1, 2, 3, 4, or 5;

p is 0, 1, 2, or 3;

A is single or double carbon-carbon bond;

Ar¹ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted
 10 with one or more substituents, each substituent independently being halogen, nitro, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, OR_a, or N(R_aR_a'), in which each R_a, independently, is H or C₁-C₄ alkyl and each R_a', independently, is H or C₁-C₄ alkyl;

Ar² is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted
 with one or more substituents, each substituent independently being halogen, cyano, nitro,
 15 C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, OR_b, N(R_bR_b'), C(O)-N(R_bR_b'), or
 NH-C(O)-N(R_bR_b'), in which each R_b, independently, is H or C₁-C₄ alkyl and each R_b',
 independently, is H or C₁-C₄ alkyl;

Ar³ is aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted
 with one or more substituents, each substituent independently being halogen, C₁-C₄ alkyl,
 20 C₁-C₄ hydroxyalkyl, OR_c, or N(R_cR_c'), in which each R_c, independently, is H or C₁-C₄
 alkyl and each R_c', independently, is H or C₁-C₄ alkyl;

each R¹, independently, is C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl,
 OR_d, or C(O)-N(R_dR_d'), in which each R_d, independently, is H or C₁-C₄ alkyl and each R_d',

2016206059 03 Jul 2019

independently, is H or C₁-C₄ alkyl;

R² is -(CH₂)_n-R, in which n is 0, 1, 2, or 3 and R is guanidino optionally substituted by CN or CH₃, aminoacyl, C₁-C₄ alkylaminoacyl, OR_e, N(R_eR_{e'}), NH-C(O)-CH(NH₂)-(CH₂)₄-N(R_eR_{e'}), NH-C(O)-CH₂-(OCH₂CH₂)₂-N(R_eR_{e'}), or 5-membered heterocycloalkyl optionally substituted with C₁-C₄ alkyl or N(R_eR_{e'}), in which each R_e, independently, is H or C₁-C₄ alkyl and each R_{e'}, independently, is H or C₁-C₄ alkyl; and

each R³, independently, is halogen, C₁-C₄ alkyl, or OR_f, in which each R_f, independently, is H or C₁-C₄ alkyl;

with the provisos that, when n is 0, R is not amino or guanidino and that, when the amino acid residue bonded to Ar¹C(O) is L-Val, Ar¹ is not unsubstituted phenyl.

2. The compound of claim 1, wherein Ar¹ is phenyl, pyridinyl, oxazolyl, thiazolyl, imidazolyl, pyrimidinyl, pyrrolyl, or triazolyl, each of which is optionally substituted with one or more substituents, each substituent independently being F, Cl, NO₂, CH₃, CH₂OH, or NH₂.

3. The compound of claim 1 or claim 2, wherein Ar² is phenyl or pyridinyl, each of which is optionally substituted with one or more substituents, each substituent independently being CH₂NH₂, C(O)NH₂, OH, CN, CH₂OH, NH₂, or NH-C(O)-NH₂.

4. The compound of any one of claims 1-3, wherein Ar³ is pyridinyl.

5. The compound of any one of claims 1-4, wherein m is 1.

6. The compound of any one of claims 1-5, wherein R¹ is OH, C(O)NH₂, or CH₂NH₂.

7. The compound of any one of claims 1-6, wherein n is 0, 1, or 2.

8. The compound of any one of claims 1-7, wherein R is N(R_eR_{e'}), NH-C(O)-CH(NH₂)-(CH₂)₄-N(R_eR_{e'}), NH-C(O)-CH₂-(OCH₂CH₂)₂-N(R_eR_{e'}), triazolyl optionally

2016206059 03 Jul 2019

substituted with NH₂, or guanidino optionally substituted with CN or CH₃, in which each R_e, independently, is H or C₁-C₃ alkyl and each R_e', independently, is H or C₁-C₃ alkyl.

9. The compound of any one of claims 1-7, wherein R is NH₂, N(CH₃)₂,
 5 N(CH₂CH₃)₂, NH(CH(CH₃)₂), NH-C(O)-CH(NH₂)-(CH₂)₄-N(CH₃)₂, NH-C(O)-CH₂-
 (OCH₂CH₂)₂-NH₂, NH-C(O)-CH₂-(OCH₂CH₂)₂-NH(CH(CH₃)₂), 3-amino-1,2,4-triazol-5-
 yl, or guanidino optionally substituted with CN or CH₃.

10. The compound of any one of claims 1-9, wherein p is 0.

11. The compound of claim 1, wherein the compound is Oxazole-2-carbonyl-D-
 Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂.

12. The compound of claim 1, wherein the compound is
 15 Picolinoyl-D-Val-Tyr-c(Cys-Dab(Et₂)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl-D-Val-Tyr-c(Cys-Dab(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl-D-Val-Tyr-c(Cys-Orn(Et₂)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl-D-Val-Tyr-c(Cys-Orn-Asp-Val-Gly-Pro-Phe(4-CH₂OH)-Cys)-3Pal-NH₂;
 20 Picolinoyl(3,5-F₂)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-3Pal-Cys)-3Pal-
 NH₂;
 Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-
 NH₂;
 25 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-3Pal-Cys)-3Pal-
 NH₂;
 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-
 CH₂NH₂)-Cys)-3Pal-NH₂;
 1H-1,2,4-triazole-5-carbonyl(3-Me)-D-Val-Tyr-c(Cys-Arg-Asp-Val-Gly-Pro-Phe(2-
 30 CH₂NH₂)-Cys)-3Pal-NH₂;
 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-4Aph-Cys)-

2016206059 03 Jul 2019

- 3Pal-NH₂;
 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-4Uph-Cys)-
 3Pal-NH₂;
 Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe(4-CH₂OH)-Cys)-3Pal-
 5 NH₂;
 Picolinoyl(3,5-F₂)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe-Cys)-3Pal-NH₂;
 Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(4-CH₂OH)-
 Cys)-3Pal-NH₂;
 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Pro-Phe(4-
 10 CH₂OH)-Cys)-3Pal-NH₂;
 Oxazole-2-carbonyl-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-
 Cys)-3Pal-NH₂;
 Picolinoyl(5-F)-D-Val-Phe(2-Cbm)-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(2-Cbm)-Cys)-
 3Pal-NH₂;
 15 Oxazole-2-carbonyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-
 NH₂;
 Picolinoyl(5-F)-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(3-Cbm)-Cys)-3Pal-
 NH₂; or
 Picolinoyl-D-Val-Tyr-c(Cys-Orn(iPr)-Asp-Val-Gly-Dhp-Phe(4-CH₂OH)-Cys)-3Pal-NH₂.

20

13. A pharmaceutical composition, comprising the compound of any of claims 1-12 and a pharmaceutically acceptable carrier.

14. The pharmaceutical composition of claim 13, wherein the composition
 25 comprises an aqueous solution.

15. The pharmaceutical composition of claim 14, wherein the composition comprises a sodium chloride aqueous solution.

16. The pharmaceutical composition of claim 15, wherein the aqueous solution
 30 comprises about 0.9 wt% of sodium chloride.

2016206059 03 Jul 2019

17. A method of treating migraine, comprising administering to a patient in need thereof an effective amount of the pharmaceutical composition of claim 13.

5 18. A compound of any of claims 1 to 12 for, or for use in, the treatment of migraine.

19. Use of a compound of any of claims 1 to 12 in the manufacture of a medicament for the treatment of migraine.
10