



(51) International Patent Classification:

C07D 498/04 (2006.01) A61P 35/00 (2006.01)
A61K 31/553 (2006.01)

(21) International Application Number:

PCT/GB2020/052588

(22) International Filing Date:

14 October 2020 (14.10.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1914860.0 14 October 2019 (14.10.2019) GB

(71) Applicants: **CANCER RESEARCH TECHNOLOGY LIMITED** [GB/GB]; 2 Redman Place, London E20 1JQ (GB). **THE INSTITUTE OF CANCER RESEARCH: ROYAL CANCER HOSPITAL** [GB/GB]; 123 Old Brompton Road, London SW7 3RP (GB).

(72) Inventors: **BELLENIE, Benjamin Richard**; The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **BRENNAN, Alfie**; c/o The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **CHEUNG, Kwai Ming Jack**; The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **DAVIS, Owen Alexander**; c/o The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **HARNDEN, Alice Claire**; The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **HOELDER, Swen**; The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB). **HUCKVALE, Rosemary**; The Institute of Cancer Research: Royal Cancer Hospital, 123 Old Brompton Road, London SW7 3RP (GB).

(74) Agent: **HGF LIMITED**; Document Handling - HGF (Manchester), 1 City Walk, LEEDS Yorkshire LS11 9DX (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW,

SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: [1,4]OXAZEPINO[2,3-C]QUI NOLI NONE DERIVATIVES AS BCL6 INHIBITORS

(57) Abstract: The present invention relates to compounds that function as inhibitors of BCL6 (B-cell lymphoma 6) activity. The present invention also relates to processes for the preparation of these compounds, to pharmaceutical compositions comprising them, and to their use in the treatment of proliferative disorders, such as cancer, as well as other diseases or conditions in which BCL6 activity is implicated.

[1 ,4]OXAZEPINO[2,3-C]QUINOLINONE DERIVATIVES AS BCL6 INHIBITORS**INTRODUCTION**

[0001] The present invention relates to certain compounds that function as inhibitors of BCL6 (B-cell lymphoma 6) activity. The present invention also relates to processes for the preparation of these compounds, to pharmaceutical compositions comprising them, and to their use in the treatment of proliferative disorders, such as cancer, as well as other diseases or conditions in which BCL6 activity is implicated.

BACKGROUND OF THE INVENTION

[0002] BCL6 is a zinc finger transcription repressor that plays a key role in the formation and development of germinal centres, in which B cells undergo somatic hypermutation and recombination of the immunoglobulin genes, in order to generate diversity in antibodies against a variety of foreign antigens (Dent et al., Science, 1997, 276, 589-592). BCL6 allows the proliferation of antibody producing B cells by repressing genes involved in DNA damage response, cell cycle arrest and apoptosis. BCL6 mediates this repression by recruiting the corepressor proteins SMRT, NCoR and BCoR to an extended groove motif that forms along the dimer interface of the BCL6 BTB (BR-C, Ttk and Bab) domain (Ahmad et al., Mol Cell, 2003, 12, 1551-1564; Ghetu et al., Mol Cell, 2008, 29, 384-391). Genetic upregulation of the BCL6 gene, as seen in many lymphomas, leads to malignant B cell proliferation (Hatzi & Melnick, Trends Mol Med, 2014, 20, 343-352). Therefore, there exists a need to develop agents that inhibit the tumourigenic effects of BCL6, either by selectively binding to the BTB domain and preventing corepressor recruitment, or by binding to the BTB domain and inducing protein degradation (Kerres et al. Cell Rep., 2017, 20, 2860-2875; Bellenie et al., J. Med. Chem., 2020, 63, 4047-4068).

SUMMARY OF THE INVENTION

[0003] According to a first aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein.

[0004] According to a further aspect of the present invention, there is provided a pharmaceutical composition comprising a compound as defined herein, or a pharmaceutically acceptable salt, hydrate or solvate thereof, in admixture with a pharmaceutically acceptable diluent or carrier.

[0005] According to a further aspect of the present invention, there is provided a method of inhibiting BCL6 activity, *in vitro* or *in vivo*, said method comprising contacting a cell with an effective amount of a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein.

[0006] According to a further aspect of the present invention, there is provided a method of inhibiting cell proliferation, *in vitro* or *in vivo*, said method comprising contacting a cell with an

effective amount of a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein, or a pharmaceutical composition as defined herein.

[0007] According to a further aspect of the present invention, there is provided a method of treating a disease or disorder in which BCL6 activity is implicated in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein, or a pharmaceutical composition as defined herein.

[0008] According to a further aspect of the present invention, there is provided a method of treating a proliferative disorder in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein, or a pharmaceutical composition as defined herein.

[0009] According to a further aspect of the present invention, there is provided a method of treating cancer in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein, or a pharmaceutical composition as defined herein.

[0010] According to a further aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in therapy.

[0011] According to a further aspect of the present invention, there is provided a compound or a pharmaceutically acceptable salt, hydrate or solvate thereof as defined herein, or a pharmaceutical composition as defined herein, for use in the treatment of a proliferative condition.

[0012] According to a further aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in the treatment of cancer, HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease, non-alcoholic steatohepatitis or auto-immune diseases. In a particular embodiment, the cancer is human cancer.

[0013] According to a further aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein for use in the inhibition of BCL6 activity.

[0014] According to a further aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein for use in the treatment of a disease or disorder in which BCL6 activity is implicated (such as cancer, HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease, non-alcoholic steatohepatitis

or auto-immune diseases).

[0015] According to a further aspect of the present invention, there is provided the use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of a proliferative condition.

- 5 [0016] Suitably, the proliferative disorder is cancer, suitably a human cancer (for example haematological cancers such as lymphomas (including diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL) and angioimmunoblastic T-cell lymphoma (AITL)), leukaemias (including acute lymphoblastic leukaemia (ALL), acute myeloid leukaemia (AML) and chronic myeloid leukaemia (CML)) and multiple myeloma, and solid tumours (including
10 glioma, breast cancer, non-small cell lung cancer (NSCLC) and squamous cell carcinomas (SCC) (including SCC of the head and neck, oesophagus, lung and ovary)).

[0017] According to a further aspect of the present invention, there is provided the use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of cancer.

- 15 [0018] According to a further aspect of the present invention, there is provided a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the inhibition of BCL6 activity.

[0019] According to a further aspect of the present invention, there is provided a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in
20 the manufacture of a medicament for the treatment of a disease or disorder in which BCL6 activity is implicated.

[0020] According to a further aspect of the present invention, there is provided a process for preparing a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein.

- 25 [0021] According to a further aspect of the present invention, there is provided a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, obtainable by, or obtained by, or directly obtained by a process of preparing a compound as defined herein.

[0022] According to a further aspect of the present invention, there are provided novel intermediates as defined herein which are suitable for use in any one of the synthetic methods
30 set out herein.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0023] Unless otherwise stated, the following terms used in the specification and claims have the following meanings set out below.

- 35 [0024] It is to be appreciated that references to “treating” or “treatment” include prophylaxis as

well as the alleviation of established symptoms of a condition. "Treating" or "treatment" of a state, disorder or condition therefore includes: (1) preventing or delaying the appearance of clinical symptoms of the state, disorder or condition developing in a human that may be afflicted with or predisposed to the state, disorder or condition but does not yet experience or display clinical or subclinical symptoms of the state, disorder or condition, (2) inhibiting the state, disorder or condition, i.e., arresting, reducing or delaying the development of the disease or a relapse thereof (in case of maintenance treatment) or at least one clinical or subclinical symptom thereof, or (3) relieving or attenuating the disease, i.e., causing regression of the state, disorder or condition or at least one of its clinical or subclinical symptoms.

- 10 **[0025]** A "therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity and the age, weight, etc., of the mammal to be treated.

Compounds of the Invention

- 15 **[0026]** In one aspect, the present invention relates to compounds, or pharmaceutically acceptable salts, hydrates or solvates thereof, selected from the following compounds:

(S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1a);

- (S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1b);

(S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1c);

(S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1d);

- 25 (S)-10-((2-(((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1e);

N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide

- 30 (Example 1f);

N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1g);

- (S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1h);

(S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1i);

(S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1j);

5 (S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1k);

(S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1l);

10 (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1m);

(S)-10-((5-chloro-2-(3-hydroxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1n);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
15 (Example 2a);

(S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2b);

(S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one
20 (Example 3a);

(S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4a);

(S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4b);

25 (S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4c);

(S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4d);

(S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4e);
30

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 5a);

(S)-10-((5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 6a);
35

(2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7a);

(2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7b);

(S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 8a);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9b);

(3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);

- (S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);
- 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- 5 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9h);
- (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9i);
- 10 (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- 15 N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide (Example 9k);
- N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide (Example 9k);
- 20 (S)-10-((5-chloro-2-((1R,4R)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9l);
- 25 (S)-10-((5-chloro-2-((1S,4S)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9l);
- N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide (Example 9m);
- 30 N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide (Example 9n);

(S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9o);

(S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9p);

(S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9q);

(S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((5-chloro-2-((3S,5R)-3,5-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9s);

(S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9t);

(S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9u);

(S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9v);

(S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9w);

(2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9x);

(2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9y);

(2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9za);

(S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9zb);

(S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

(S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

(S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);

(S)-10-((5-chloro-2-((3R,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);

(S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);

(S)-10-((5-chloro-2-((S)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);

(S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);

(S)-10-((5-fluoro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);

(2S)-10-[[5-chloro-2-(4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl]amino]-2-cyclopropyl-3,3-difluoro-7-methyl-2,4-dihydro-1H-[1,4]oxazepino[2,3-c]quinolin-6-one (Example 11a);

(S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);

(S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);

(S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);

(S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);

(S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);

(S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);

(S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11e);

(S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

5 (S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

(S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

10 (S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

(S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11h);

15 (S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11h);

(S)-10-((5-chloro-2-(methyl(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11i);

20 (S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11j);

(S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11j);

25 (S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11k);

(S)-10-((5-chloro-2-(5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11l);

30 (S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);

(S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);

35 (S)-10-((5-chloro-2-(3,3-dimethylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11n);

- (S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11o);
- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile (Example 11p);
- 5 (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11q);
- (S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);
- (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);
- 10 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile (Example 11s);
- (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile (Example 11t);
- 15 (S)-10-((5-chloro-2-(3-(difluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11u);
- (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11v);
- 20 (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11w);
- (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11x);
- (S)-10-((5-chloro-2-((R)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11y);
- 25 (S)-10-((5-chloro-2-((S)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11y);
- (S)-10-((5-chloro-2-((R)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11z);
- 30 (S)-10-((5-chloro-2-((S)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11z);
- (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- (S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- 35

- (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12b);
- (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12c);
- 5 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12d);
- (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- (S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- 10 (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- 15 (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-
- 20 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);
- (S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);
- 25 (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);
- (S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
- 30 (Example 12i);
- (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);
- (S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-
- 35 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);

- (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13a);
((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13b);
- 5 (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);
(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);
(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);
- 10 (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);
(S)-10-((5-chloro-2-((S)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x01);
- 15 (S)-10-((5-chloro-2-((R)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x02);
(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x03);
- 20 (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x04);
(S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x05);
- 25 (2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x06);
(2S)-10-((5-chloro-2-(3,4-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x07);
- 30 (2S)-10-((5-chloro-2-(2,3-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x08);
(2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x09);

(S)-10-((5-chloro-2-((2R,5S)-2-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x10);

5 (2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x11);

(2S)-10-((5-chloro-2-(2,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x12);

10 (S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x13);

(2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x14);

15 (2S)-10-((5-chloro-2-(6-hydroxy-4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x15);

(2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x15-*d*₃);

20 (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16);

(2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

25 (2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

30 (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17);

(2S)-10-((5-chloro-2-((3R,5S)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17a/b);

(2S)-10-((5-chloro-2-((3S,5R)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17a/b);

5 (2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x18);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x19);

10 (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x20);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x21);

15 (2S)-10-((5-chloro-2-(3-ethynylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x22);
1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile (Example 12x23);

(S)-10-((5-chloro-2-((4,4-difluorocyclohexyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x24);

(2S)-10-((2-(2-azabicyclo[4.1.0]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x25);

25 (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x26);

(2S)-10-((5-chloro-2-(3-methyl-5-oxopiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x27);

30 (S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x28);

(S)-10-((5-chloro-2-((3R,5S)-3-(3-(dimethylamino)propoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x29);

(2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x31);

(S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x32);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33);

(S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33a);

(S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33b);

(2S)-10-((5-chloro-2-(2-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x34);

(S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x35);

(2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x36);

(2S)-10-((5-chloro-2-(4,4-difluoro-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x37);

(S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile (Example 12x38);

1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide (Example 12x39);

(2S)-10-((5-chloro-2-(3-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x40);

(2S)-10-((5-chloro-2-(4,4-difluoro-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x41);

(S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x42);

(S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x44);

(2S)-10-((5-chloro-2-(2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45);

(S)-10-((5-chloro-2-((S)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45a);

(S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45b);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x46);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47);

(2S)-10-((5-chloro-2-(4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48);

(S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48a);

(S)-10-((5-chloro-2-((2S,4R)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48b);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x49);

(2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x50);

(2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p01);

- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p02);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p06);
- 5 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p07);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p08);
- (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p09);
- 10 (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p11);
- 15 (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p12);
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p13);
- 20 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c01);
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c02);
- 25 (2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c03);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c06);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c07);
- 30 (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c08);
- (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c09);
- 35

(S)-10-((2-(((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c10);

5 (S)-10-((2-(((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c11);

(S)-2-cyclopropyl-10-((2-(dimethylamino)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c12);

10 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c13);

(2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f01);

15 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f02);

(2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f03);

20 (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f04);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f05);

25 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f06);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f07);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f08);

30 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f09);

(S)-10-((2-(((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f10);

(S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f11);

(S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f12); and

(2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f13).

[0027] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

(S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1a);

(S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1b);

(S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1c);

(S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1d);

(S)-10-((2-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1e);

N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1f);

N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1g);

(S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1h);

(S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1i);

(S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1j);

- (S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1k);
- (S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1l);
- 5 (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1m);
- (S)-10-((5-chloro-2-(3-hydroxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1n);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2a);
- 10 (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2b);
- (S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one (Example 3a);
- 15 (S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4a);
- (S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4b);
- 20 (S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4c);
- (S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4d);
- 25 (S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4e);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 5a);
- 30 (S)-10-((5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 6a);
- (2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7a);

(2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7b);

(S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 8a);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9b);

(3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);

(S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);

2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

- 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9h);
- 5 (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9i);
- (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- 10 (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-
- 15 ethylacetamide (Example 9k);
- N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide (Example 9k);
- (S)-10-((5-chloro-2-((1R,4R)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-
- 20 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9l);
- (S)-10-((5-chloro-2-((1S,4S)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9l);
- 25 N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide (Example 9m);
- N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-
- 30 methylacetamide (Example 9n);
- (S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9o);
- (S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-
- 35 difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9p);

(S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9q);

(S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((5-chloro-2-((3S,5R)-3,5-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9s);

(S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9t);

(S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9u);

(S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9v);

(S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9w);

(2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9x);

(2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9y);

(2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9za);

(S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9zb);

(S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

- (S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);
- (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);
- 5 (S)-10-((5-chloro-2-((3R,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);
- (S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);
- (S)-10-((5-chloro-2-((S)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);
- 10 (S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);
- (S)-10-((5-fluoro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);
- 15 (2S)-10-[[5-chloro-2-(4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl]amino]-2-cyclopropyl-3,3-difluoro-7-methyl-2,4-dihydro-1H-[1,4]oxazepino[2,3-c]quinolin-6-one (Example 11a);
- (S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);
- (S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);
- 20 (S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);
- (S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);
- 25 (S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);
- (S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);
- (S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11e);
- 30 (S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

(S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

5 (S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

(S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

10 (S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11h);

(S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
15 (Example 11h);

(S)-10-((5-chloro-2-(methyl(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11i);

(S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
20 (Example 11j);

(S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11j);

(S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11k);
25

(S)-10-((5-chloro-2-(5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11l);

(S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);

30 (S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);

(S)-10-((5-chloro-2-(3,3-dimethylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11n);

(S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11o);
35

- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile (Example 11p);
- (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11q);
- 5 (S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);
- (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);
- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile
- 10 (Example 11s);
- (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile (Example 11t);
- (S)-10-((5-chloro-2-(3-(difluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11u);
- 15 (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11v);
- (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11w);
- 20 (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11x);
- (S)-10-((5-chloro-2-((R)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11y);
- (S)-10-((5-chloro-2-((S)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11y);
- 25 (S)-10-((5-chloro-2-((R)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11z);
- (S)-10-((5-chloro-2-((S)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11z);
- 30 (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- (S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-
- 35 3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12b);

- (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12c);
- (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12d);
- 5 (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- (S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- 10 (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- 15 (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);
- 20 (S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);
- (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);
- 25 (S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);
- (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);
- 30 (S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);
- (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13a);
- 35

((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13b);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);

5 (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);

10 (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);

(S)-10-((5-chloro-2-((S)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x01);

(S)-10-((5-chloro-2-((R)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x02);

15 (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x03);

(S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

20 (Example 12x05);

(2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x06);

(2S)-10-((5-chloro-2-(3,4-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x07);

25 (2S)-10-((5-chloro-2-(2,3-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x08);

(2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x09);

30 (S)-10-((5-chloro-2-((2R,5S)-2-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x10);

(2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

35 (Example 12x11);

(2S)-10-((5-chloro-2-(2,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x12);

(S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

5 (Example 12x13);

(2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x14);

(2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

10 (Example 12x15-*d*₃);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16);

(2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

15

(2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

(2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17);

20

(2S)-10-((5-chloro-2-((3R,5S)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-

25 [1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17a/b);

(2S)-10-((5-chloro-2-((3S,5R)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x17a/b);

(2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x18);

30

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x19);

(2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x20);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x21);

(2S)-10-((5-chloro-2-(3-ethynylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x22);

1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile (Example 12x23);

(S)-10-((5-chloro-2-((4,4-difluorocyclohexyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x24);

(2S)-10-((2-(2-azabicyclo[4.1.0]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x25);

(2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x26);

(S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x28);

(2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x31);

(S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x32);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33);

(S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33a);

(S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33b);

(S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x35);

(2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x36);

(S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile (Example 12x38);

1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide (Example 12x39);

(2S)-10-((5-chloro-2-(3-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x40);

(S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x42);

(S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x44);

(2S)-10-((5-chloro-2-(2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45);

(S)-10-((5-chloro-2-((S)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45a);

(S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45b);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x46);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47);

(2S)-10-((5-chloro-2-(4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48);

(S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48a);

- (S)-10-((5-chloro-2-((2S,4R)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48b);
- 5 (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x49);
- (2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x50);
- 10 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p01);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p06);
- 15 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p08);
- (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p11);
- 20 (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p12);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c06);
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c08);
- 25 (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13c12);
- (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f04); and
- 30 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f05).

[0028] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

- (S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1b);
- (S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1c);
- 5 (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1d);
- (S)-10-((2-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1e);
- 10 N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1f);
- N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide
- 15 (Example 1g);
- (S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1h);
- (S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-
- 20 7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1i);
- (S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1j);
- (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1m);
- 25 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2a);
- (S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one
- 30 (Example 3a);
- (S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4a);
- (S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4b);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 5a);

(2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-

5 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7a);

(2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-

cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7b);

10 (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-

cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-

15 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-

cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9b);

(3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-

20 [1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-

[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

25 (S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-

cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-

30 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

35 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-

[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

- 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9h);
- 5 (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9i);
- (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- 10 (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-
- 15 ethylacetamide (Example 9k);
- N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide (Example 9k);
- N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-
- 20 methylacetamide (Example 9m);
- N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide (Example 9n);
- 25 (S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9o);
- (S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9p);
- 30 (S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9q);
- (S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9r);

(S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9t);

(S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9u);

(S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9v);

(S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9w);

(2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9x);

(2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9y);

(2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9za);

(S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9zb);

(S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);

(S)-10-((5-chloro-2-((S)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10c);

(S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11e);

(S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-

[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

(S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11f);

5 (S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

(S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11g);

10 (S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11h);

(S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
15 (Example 11h);

(S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11j);

(S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
20 (Example 11j);

(S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11k);

(S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);
25

(S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11m);

(S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11o);

30 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile (Example 11p);

(S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11q);

(S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile
35 (Example 11s);

- (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile (Example 11t);
- (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11v);
- 5 (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11w);
- (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11x);
- (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- 10 (S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12a);
- (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12b);
- 15 (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12c);
- (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12d);
- (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- 20 (S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12e);
- (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- 25 (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);
- (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);
- 30 (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);

(S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);

5 (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);

(S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);

10 (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);

(S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
15 (Example 12j);

(S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13a);

(S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
20 (Example 12x05);

(2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x09);

(2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
25 (Example 12x11);

(S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x13);

30 (2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x14);

(2S)-10-((5-chloro-2-(6-hydroxy-4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x15);

(2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x15-d3);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x16);

(2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x16a/b);

(2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x16a/b);

(2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x17);

(2S)-10-((5-chloro-2-((3R,5S)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x17a/b);

(2S)-10-((5-chloro-2-((3S,5R)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x17a/b);

(2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x18);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x19);

(2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x20);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Example 12x21);

1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-*c*]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile (Example 12x23);

- (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x26);
- (S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x28);
- (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);
- (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x31);
- (S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x32);
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33);
- (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33a);
- (S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33b);
- (S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x35);
- (2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x36);
- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile (Example 12x38);
- 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide (Example 12x39);
- (S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x42);
- (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x44);

(S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x45b);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47);

(S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x48a);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x49);

(2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x50);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f05); and

(S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p11).

[0029] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

(S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1b);

(S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1c);

N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1f);

N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1g);

(S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1i);

(S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1j);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

5 (Example 2a);

(S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one (Example 3a);

10 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 5a);

(2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7a);

15 (2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 7b);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

20 (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

25 (S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9b);

2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

30 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

(S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9i);

N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-

35 methylacetamide (Example 9m);

(S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9p);

(S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9w);

(2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9x);

(2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9y);

(S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11e);

(S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12b);

(S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);

(S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12g);

(S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);

(S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12h);

(S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);

(S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12i);

(S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);

(S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12j);

5 (2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x09);

(2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x11);

10 (2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x15-d3);

(2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

(2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16a/b);

20 (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x26);

(S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x28);

25 (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);

(S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33a);
and

30 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide (Example 12x39).

[0030] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1g);

(S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1k);

(S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1l);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2a);

(S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2b);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);

- (S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9f);
- 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- 5 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);
- (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- 10 (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide (Example 9j);
- (S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);
- 15 (S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);
- (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);
- (S)-10-((5-chloro-2-((3R,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10b);
- 20 (S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);
- (S)-10-((5-fluoro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10d);
- 25 (S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);
- (S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11b);
- (S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);
- 30 (S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11c);
- (S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);

(S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11d);

(S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);

5 (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11r);

(S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12c);

10 (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);

(S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);

(S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13a);

15 ((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13b);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);

20 (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14a);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 14b);

25 (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x03);

(S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x05);

30 (2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x06);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16);

35

(S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x28);

5 (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x31);

10 (S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x42);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47);

15 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13p01);

(S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f04); and

20 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f05).

[0031] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

25 N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide (Example 1g);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
30 (Example 2a);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (Example 9c);

(S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9d);

(S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9e);

2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (Example 9g);

(S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

(S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

(S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12c);

(S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12f);

(S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13a);

(S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x03);

(S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x05);

(2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x16);

(2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x30);

(2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x31);

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47); and

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 13f05).

[0032] Particular compounds of the present invention include any of the following compounds, or a pharmaceutically acceptable salt or solvate thereof:

(S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 1a);

(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2a);

(S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 2b);

(S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one (Example 3a);

(S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4b);

(S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4c);

(S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4d);

(S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 4e);

5 (S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 8a);

(S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

10 (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 9a);

(S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 10a);

15 (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11q);

(S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11v);

(S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 11w);

20 (S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x05);

(2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x33);
and

(S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Example 12x47).

30 **[0033]** A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulfuric, phosphoric, trifluoroacetic, formic, citric, methane sulfonate or maleic acid. In addition, a suitable pharmaceutically acceptable salt of a compound of the
35 invention which is sufficiently acidic is an alkali metal salt, for example a sodium or potassium

salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a pharmaceutically acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

5 **[0034]** Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers". Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers" and those that are non-superimposable mirror images of each other are termed "enantiomers". When a
10 compound has an asymmetric center, for example, it is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center and is described by the R- and S-sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist
15 as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".


[0035] The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the
20 specification and claims is intended to include both individual enantiomers and mixtures, racemic or otherwise, thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art (see discussion in Chapter 4 of "Advanced Organic Chemistry", 4th edition J. March, John Wiley and Sons, New York, 2001), for example by synthesis from optically active starting materials or by resolution of a racemic form. Some of the
25 compounds of the invention may have geometric isomeric centres (E- and Z- isomers). It is to be understood that the present invention encompasses all optical, diastereoisomers and geometric isomers and mixtures thereof that possess antiproliferative activity.

[0036] The present invention also encompasses compounds of the invention as defined herein which comprise one or more isotopic substitutions. For example, H may be in any isotopic form,
30 including 1H , 2H(D) , and 3H (T) ; C may be in any isotopic form, including 12C , 13C , and 14C ; and O may be in any isotopic form, including 16O and 18O ; and the like.

[0037] It is also to be understood that certain compounds of the invention may exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms that possess antiproliferative activity.

[0038] It is also to be understood that certain compounds of the invention may exhibit polymorphism, and that the invention encompasses all such forms that possess antiproliferative activity.

[0039] Compounds of the invention may exist in a number of different tautomeric forms and references to compounds of the invention include all such forms. For the avoidance of doubt, where a compound can exist in one of several tautomeric forms, and only one is specifically described or shown, all others are nevertheless embraced by the invention. Examples of tautomeric forms include keto-, enol-, and enolate-forms, as in, for example, the following tautomeric pairs: keto/enol (illustrated below), imine/enamine, amide/imino alcohol, amidine/amidine, nitroso/oxime, thioketone/enethiol, and nitro/aci-nitro.



[0040] The compounds of the invention may be administered in the form of a pro-drug which is broken down in the human or animal body to release a compound of the invention. A pro-drug may be used to alter the physical properties and/or the pharmacokinetic properties of a compound of the invention. A pro-drug can be formed when the compound of the invention contains a suitable group or substituent to which a property-modifying group can be attached. Examples of pro-drugs include *in vivo* cleavable ester derivatives that may be formed at a carboxy group or a hydroxy group in a compound of invention, and *in-vivo* cleavable amide derivatives that may be formed at a carboxy group or an amino group in a compound of the invention.

[0041] Accordingly, the present invention includes those compounds of the invention, as defined hereinbefore, when made available by organic synthesis and when made available within the human or animal body by way of cleavage of a pro-drug thereof. Accordingly, the present invention includes those compounds of the invention that are produced by organic synthetic means and also such compounds that are produced in the human or animal body by way of metabolism of a precursor compound, that is a compound of the invention may be a synthetically-produced compound or a metabolically-produced compound.

[0042] A suitable pharmaceutically acceptable pro-drug of a compound of the invention is one that is based on reasonable medical judgement as being suitable for administration to the human or animal body without undesirable pharmacological activities and without undue toxicity.

[0043] Various forms of pro-drug have been described, for example in the following documents :-

- a) Methods in Enzymology, Vol. 42, p. 309-396, edited by K. Widder, et al. (Academic Press, 1985);
- b) Design of Pro-drugs, edited by H. Bundgaard, (Elsevier, 1985);

- c) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Pro-drugs", by H. Bundgaard p. 113-191 (1991);
- d) H. Bundgaard, Advanced Drug Delivery Reviews, 8, 1-38 (1992);
- 5 e) H. Bundgaard, et al., Journal of Pharmaceutical Sciences, 77, 285 (1988);
- f) N. Kakeya, et al., Chem. Pharm. Bull., 32, 692 (1984);
- g) T. Higuchi and V. Stella, "Pro-Drugs as Novel Delivery Systems", A.C.S. Symposium Series, Volume 14;
- h) E. Roche (editor), "Bioreversible Carriers in Drug Design", Pergamon Press, 1987; and
- 10 i) J. Rautio, et al., Nature Reviews Drug Discov., 17, 559-587 (2018).

[0044] A suitable pharmaceutically acceptable pro-drug of a compound of the invention that possesses a carboxy group is, for example, an *in vivo* cleavable ester thereof. An *in vivo* cleavable ester of a compound of the invention containing a carboxy group is, for example, a pharmaceutically acceptable ester which is cleaved in the human or animal body to produce the

15 parent acid. Suitable pharmaceutically acceptable esters for carboxy include (1-6C)alkyl esters such as methyl, ethyl and *tert*-butyl, (1-6C)alkoxymethyl esters such as methoxymethyl esters, (1-6C)alkanoyloxymethyl esters such as pivaloyloxymethyl esters, 3-phthalidyl esters, (3-8C)cycloalkylcarbonyloxy-(1-6C)alkyl esters such as cyclopentylcarbonyloxymethyl and 1-cyclohexylcarbonyloxyethyl esters, 2-oxo-1,3-dioxolenylmethyl esters such as 5-methyl-2-oxo-

20 1,3-dioxolen-4-ylmethyl esters and (1-6C)alkoxycarbonyloxy-(1-6C)alkyl esters such as methoxycarbonyloxymethyl and 1-methoxycarbonyloxyethyl esters.

[0045] A suitable pharmaceutically acceptable pro-drug of a compound of the invention that possesses a hydroxy group is, for example, an *in vivo* cleavable ester or ether thereof. An *in vivo* cleavable ester or ether of a compound of the invention containing a hydroxy group is, for

25 example, a pharmaceutically acceptable ester or ether which is cleaved in the human or animal body to produce the parent hydroxy compound. Suitable pharmaceutically acceptable ester forming groups for a hydroxy group include inorganic esters such as phosphate esters (including phosphoramidic cyclic esters). Further suitable pharmaceutically acceptable ester forming groups for a hydroxy group include (1-10C)alkanoyl groups such as acetyl, benzoyl, phenylacetyl and

30 substituted benzoyl and phenylacetyl groups, (1-10C)alkoxycarbonyl groups such as ethoxycarbonyl, N,N-(1-6C)₂carbamoyl, 2-dialkylaminoacetyl and 2-carboxyacetyl groups. Examples of ring substituents on the phenylacetyl and benzoyl groups include aminomethyl, N-alkylaminomethyl, N,N-dialkylaminomethyl, morpholinomethyl, piperazin-1-ylmethyl and 4-(1-4C)alkylpiperazin-1-ylmethyl. Suitable pharmaceutically acceptable ether forming groups for a

35 hydroxy group include α -acyloxyalkyl groups such as acetoxymethyl and pivaloyloxymethyl groups.

[0046] A suitable pharmaceutically acceptable pro-drug of a compound of the invention that possesses a carboxy group is, for example, an *in vivo* cleavable amide thereof, for example an amide formed with an amine such as ammonia, a (1-4C)alkylamine such as methylamine, a [(1-4C)alkyl]₂amine such as dimethylamine, N-ethyl-N-methylamine or diethylamine, a (1-4C)alkoxy-(2-4C)alkylamine such as 2-methoxyethylamine, a phenyl-(1-4C)alkylamine such as benzylamine and amino acids such as glycine or an ester thereof.

[0047] A suitable pharmaceutically acceptable pro-drug of a compound of the invention that possesses an amino group is, for example, an *in vivo* cleavable amide derivative thereof. Suitable pharmaceutically acceptable amides from an amino group include, for example an amide formed with (1-10C)alkanoyl groups such as an acetyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl groups. Examples of ring substituents on the phenylacetyl and benzoyl groups include aminomethyl, N-alkylaminomethyl, N,N-dialkylaminomethyl, morpholinomethyl, piperazin-1-ylmethyl and 4-(1-4C)alkyl)piperazin-1-ylmethyl.

[0048] The *in vivo* effects of a compound of the invention may be exerted in part by one or more metabolites that are formed within the human or animal body after administration of a compound of the invention. As stated hereinbefore, the *in vivo* effects of a compound of the invention may also be exerted by way of metabolism of a precursor compound (a pro-drug).

Synthesis

[0049] The compounds of the present invention can be prepared by any suitable technique known in the art. Particular processes for the preparation of these compounds are described further in the accompanying examples.

[0050] In the description of the synthetic methods described herein and in any referenced synthetic methods that are used to prepare the starting materials, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, can be selected by a person skilled in the art.

[0051] It is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule must be compatible with the reagents and reaction conditions utilised.

[0052] It will be appreciated that during the synthesis of the compounds of the invention in the processes defined herein, or during the synthesis of certain starting materials, it may be desirable to protect certain substituent groups to prevent their undesired reaction. The skilled chemist will appreciate when such protection is required, and how such protecting groups may be put in place, and later removed.

[0053] For examples of protecting groups see one of the many general texts on the subject, for example, 'Protective Groups in Organic Synthesis' by Theodora Green (publisher: John Wiley & Sons). Protecting groups may be removed by any convenient method described in the literature or known to the skilled chemist as appropriate for the removal of the protecting group in question, such methods being chosen so as to effect removal of the protecting group with the minimum disturbance of groups elsewhere in the molecule.

[0054] Thus, if reactants include, for example, groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

[0055] By way of example, a suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or t-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed by, for example, hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *tert*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulfuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

[0056] A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium, sodium hydroxide or ammonia. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

[0057] A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a t-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon. Resins may also be used as a protecting group.

Biological Activity

[0058] The biological assays described in the Examples section herein may be used to measure the pharmacological effects of the compounds of the present invention.

[0059] Although the pharmacological properties of the compounds of the invention vary with structural change, as expected, the compounds of the invention were found to be active in the HTRF *in vitro* assay, and in some cases also in the OCI-LY1 degradation assay described in the Examples section.

[0060] In general, as illustrated by the Example compound data in Table 1, the compounds of the invention demonstrate a pIC₅₀ of 7.0 or more, in the HTRF assay described in the Examples section. Preferred compounds of the invention demonstrate a pIC₅₀ of 7.5 or more. The more preferred compounds of the invention demonstrate a pIC₅₀ of 8.0 or more. The most preferred compounds of the invention demonstrate a pIC₅₀ of 8.5 or more.

[0061] In the OCI-LY1 degradation assay described herein in the Examples section, as illustrated by the Example compound data in Table 2, the compounds of Formula I may also demonstrate a pDC₅₀ of 7.0 or more (preferably 8.0 or more).

[0062] The following data were generated for the Examples:

Table 1

Example	HTRF avg pIC ₅₀	Example	HTRF avg pIC ₅₀	Example	HTRF avg pIC ₅₀	Example	HTRF avg pIC ₅₀
1a	7.99	9l	7.93	11u	7.94	12x23	8.29
1b	8.71	9m	8.60	11v	8.27	12x24	7.84
1c	8.63	9n	8.31	11w	8.43	12x25	7.8
1d	8.44	9o	8.03	11x	8.45	12x26	8.72
1e	8.49	9p	8.54	11y	7.33	12x28	8.5
1f	8.60	9q	8.37	11z	7.88	12x30	8.7
1g	8.50	9r	8.44	12a	8.20	12x31	8.42
1h	8.44	9s	7.71	12b	8.60	12x32	8.43
1i	8.51	9t	8.37	12c	8.53	12x33a	8.67
1j	8.53	9u	8.38	12d	8.31	12x33b	8.11
1k	7.42	9v	8.33	12e	8.32	12x35	8.27
1l	7.59	9w	8.56	12f	8.02	12x36	8.11
1m	8.03	9x	8.46	12g	8.58	12x38	8.43
1n	7.86	9y	8.56	12h	8.69	12x39	8.6
2a	8.62	9za	8.12	12i	8.89	12x40	7.97
2b	7.65	9zb	8.01	12j	8.58	12x42	8.21
3a	8.79	10a	7.91	12x01	7.57	12x44	8.3
4a	8.46	10b	7.80	12x02	7.67	12x45a	7.95
4b	8.20	10c	8.30	12x03	7.81	12x45b	8.32
4c	7.86	10d	7.68	12x05	8.43	12x46	7.95
4d	7.78	11a	8.56	12x06	7.52	12x47	8.08

4e	7.73	11b	7.48	12x07	7.33	12x48a	8.37
5a	8.82	11c	7.26	12x08	7.23	12x48b	7.73
6a	7.85	11d	7.19	12x09	8.77	12x49	8.2
7a	9.11	11e	9.06	12x10	7.93	12x50	8.15
7b	8.79	11f	8.26	12x11	8.79	13a	8.17
8a	7.97	11g	8.28	12x12	7.25	13b	7.51
9a	8.60	11h	8.25	12x13	8.25	13c12	7.71
9a-1	8.74	11i	7.51	12x14	8.46	13p01	7.07
9a-2	8.53	11j	8.22	12x15-d3	8.75	13p06	7.93
9b	8.51	11k	8.01	12x16a	8.69	13p08	7.98
9c	8.46	11l	7.35	12x16b	8.27	13p11	8.14
9d	8.38	11m	8.41	12x17a	8.4	13p12	7.72
9e	8.43	11n	7.83	12x17b	7.88	13f04	7.23
9f	7.99	11o	8.12	12x18	8.26	13f05	8.18
9g	8.64	11p	8.39	12x19	8.23	14a	7.81
9h	8.47	11q	8.32	12x20	8.43	14b	7.33
9i	8.57	11r	8.00	12x21	8.45		
9j	8.38	11s	8.41	12x22	7.64		
9k	8.37	11t	8.46				

Table 2

Example	OCI-LY1 avg	Example	OCI-LY1 avg	Example	OCI-LY1 avg	Example	OCI-LY1 avg
1g	8.76	9f	7.72	11r	7.57	12x16a	8.4
1k	7.8	9g	8.16	12c	8.44	12x28	6.43
1l	7.56	9j	7.99	12f	8.1	12x30	8.2
2a	8.86	9p	8.33	13a	8.49	12x31	8.27
2b	7.87	10a	8.46	13b	7.39	12x42	7.02
9a	8.42	10b	7.23	14a	7.8	12x47	8.08
9a-1	8.61	10d	7.54	14b	7.41	13p01	7.95
9c	8.79	11b	7.22	12x03	8.21	13f04	7.5
9d	8.13	11c	7.33	12x05	8.47	13f05	8.53
9e	8.3	11d	7.04	12x06	7.51		

Pharmaceutical Compositions

[0063] According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, in association with a pharmaceutically acceptable diluent or carrier.

[0064] The compositions of the invention may be in a form suitable for oral use (for example as tablets, lozenges, hard or soft capsules, aqueous or oily suspensions, emulsions, dispersible powders or granules, syrups or elixirs), for topical use (for example as creams, ointments, gels, or aqueous or oily solutions or suspensions), for administration by inhalation (for example as a finely divided powder or a liquid aerosol), for administration by insufflation (for example as a finely divided powder) or for parenteral administration (for example as a sterile aqueous or oily solution

for intravenous, subcutaneous, intramuscular, intraperitoneal or intramuscular dosing or as a suppository for rectal dosing).

[0065] The compositions of the invention may be obtained by conventional procedures using conventional pharmaceutical excipients, well known in the art. Thus, compositions intended for oral use may contain, for example, one or more colouring, sweetening, flavouring and/or preservative agents.

[0066] An effective amount of a compound of the present invention for use in therapy is an amount sufficient to treat or prevent a proliferative condition referred to herein, slow its progression and/or reduce the symptoms associated with the condition.

[0067] The amount of active ingredient that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the individual treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 0.5 g of active agent (more suitably from 0.5 to 100 mg, for example from 1 to 30 mg) compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition.

[0068] The size of the dose for therapeutic or prophylactic purposes of a compound of the formula I will naturally vary according to the nature and severity of the conditions, the age and sex of the animal or patient and the route of administration, according to well-known principles of medicine.

[0069] In using a compound of the invention for therapeutic or prophylactic purposes it will generally be administered so that a daily dose in the range, for example, 0.1 mg/kg to 75 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed. Thus, for example, for intravenous or intraperitoneal administration, a dose in the range, for example, 0.1 mg/kg to 30 mg/kg body weight will generally be used. Similarly, for administration by inhalation, a dose in the range, for example, 0.05 mg/kg to 25 mg/kg body weight will be used. Oral administration may also be suitable, particularly in tablet form. Typically, unit dosage forms will contain about 0.5 mg to 0.5 g of a compound of this invention.

Therapeutic Uses and Applications

[0070] The present invention provides compounds that function as inhibitors of BCL6 activity.

[0071] The present invention therefore provides a method of inhibiting BCL6 activity *in vitro* or *in vivo*, said method comprising contacting a cell with an effective amount of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein.

[0072] The present invention also provides a method of treating a disease or disorder in which BCL6 activity is implicated in a patient in need of such treatment, said method comprising

administering to said patient a therapeutically effective amount of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein.

5 [0073] The present invention provides a method of inhibiting cell proliferation, *in vitro* or *in vivo*, said method comprising contacting a cell with an effective amount of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein.

[0074] The present invention provides a method of treating a proliferative disorder in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof,
10 or a pharmaceutical composition as defined herein.

[0075] The present invention provides a method of treating cancer in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein.

15 [0076] The present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in therapy.

[0077] The present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in the treatment of a proliferative condition.

20 [0078] The present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in the treatment of cancer. In a particular embodiment, the cancer is human cancer.

[0079] The present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein for use in the inhibition of BCL6 activity (i.e. in the
25 inhibition of BCL6 transcriptional repression and/or co-repressor binding).

[0080] Certain compounds of the present invention have been found to bind to BCL6 and initiated the degradation of BCL6. Thus, the present invention also provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein for use in the degradation of BCL6.

30 [0081] The present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein for use in the treatment of a disease or disorder in which BCL6 activity is implicated.

[0082] The present invention provides a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the
35 treatment of a proliferative condition.

[0083] The present invention provides a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of cancer. Suitably, the medicament is for use in the treatment of human cancers.

[0084] The present invention provides a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the inhibition of BCL6 activity (i.e. in the inhibition of BCL6 transcriptional repression and/or co-repressor binding).

[0085] The present invention provides a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the degradation of BCL6.

[0086] The present invention provides a use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, as defined herein in the manufacture of a medicament for the treatment of a disease or disorder in which BCL6 activity is implicated.

[0087] The term "proliferative disorder" and "proliferative condition" are used interchangeably herein and pertain to an unwanted or uncontrolled cellular proliferation of excessive or abnormal cells which is undesired, such as, neoplastic or hyperplastic growth, whether *in vitro* or *in vivo*. Examples of proliferative conditions include, but are not limited to, pre-malignant and malignant cellular proliferation, including but not limited to, malignant neoplasms and tumours, cancers (including breast cancer, non-small cell lung cancer (NSCLC) and squamous cell carcinomas (SCC) (including SCC of the head and neck, oesophagus, lung and ovary), leukemias (including acute lymphoblastic leukaemia (ALL) and chronic myeloid leukaemia (CML)), lymphomas (including acute lymphoblastic leukaemia (ALL) and chronic myeloid leukaemia (CML)), psoriasis, bone diseases, fibroproliferative disorders (e.g., of connective tissues), and atherosclerosis. Any type of cell may be treated, including but not limited to, lymphatic, blood, lung, colon, breast, ovarian, prostate, liver, pancreas, brain, and skin.

[0088] The anti-cancer effect may arise through one or more mechanisms, including but not limited to, the regulation of cell proliferation, the inhibition of angiogenesis (the formation of new blood vessels), the inhibition of metastasis (the spread of a tumour from its origin), the inhibition of invasion (the spread of tumour cells into neighbouring normal structures), or the promotion of apoptosis (programmed cell death).

[0089] The compounds disclosed herein, or pharmaceutically acceptable salts thereof, being inhibitors of BCL6, have potential therapeutic uses in a variety of BCL6-mediated disease states. BCL6 expression has been linked to a variety of lymphomas (Wagner et al., *British J Haematology*, 2010, 152, 3-12). BCL6 is involved in chromosomal translocations in diffuse large B-cell lymphoma (DLBCL) and inhibitors of BCL6 have been reported to kill DLBCL cells (Cerchietti et al., *Cancer Cell*, 2010, 17, 400-411), primary low grade follicular lymphoma cells

(Cardenas et al., *Clin Cancer Res*, 2017, 23(4), 885-893) and Burkitt lymphoma cells (Polo et al., *Nat. Med.*, 2004, 10, 1329-1335). BCL6 is required for the formation of follicular helper T cells (Hatzl et al., *J Exp. Med.*, 2015, 212(4), 539-553), which raises the possibility that BCL6 inhibitors may be used to treat angioimmunoblastic T-cell lymphoma (AITL), in which BCL6 is strongly expressed (Cortes & Palomero, *Curr Opin Hematol*, 2016, 23, 434-443).

[0090] BCL6 has also been implicated in leukaemia cells which have acquired resistance to tyrosine kinase inhibitors (TKIs). TKIs typically fail to eradicate leukaemia-initiating cells, which may often cause recurrence of leukaemia after initial treatment. BCL6 has been identified as an important component of the TKI drug-resistance pathway in both Ph⁺ acute lymphoblastic leukaemia (ALL) (Duy et al., *Nature*, 2011, 473, 384-388) and Ph⁺ chronic myeloid leukaemia (CML) (Hurtz et al., *J Exp Med*, 2011, 208(11), 2163-2174). Inhibitors of BCL6 may therefore be used to treat ALL and CML in combination with a TKI.

[0091] Further non-haematological, solid tumours may be treated with an inhibitor of BCL6. BCL6 is amplified in approximately 50% of breast tumours and is expressed in many breast cancer cell lines, including triple negative breast cancer cell lines (Walker et al., *Oncogene*, 2015, 34, 1073-1082). BCL6 is also important for the survival and proliferation of non-small cell lung cancer (NSCLC) cells, primarily due to repression of genes involved in DNA damage repair (Marullo et al., *Proc 107th Annual Meeting AACR*, 2016, Abstract nr 1271 and Deb et al., *Cancer Res.*, 2017, Apr. 4, doi: 10.1158/0008-5472.CAN-15-3052). BCL6 amplification may also be prevalent in squamous cell carcinomas (SCC) (including SCC of the head & neck, oesophagus, lung and ovary). Furthermore, inhibition of BCL6 has recently been reported to be a suitable therapeutic target for glioma and glioblastoma (Xu et al., *Proc. Natl. Acad. Sci. U.S.A.*, 2017, 114(15), 3981-3986; Fabre et al., *PLoS One*, 2020, 15(4): e0231470).

[0092] According to a further aspect of the specification there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined herein for use in the treatment of haematological cancers such as lymphomas (including diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL) and angioimmunoblastic T-cell lymphoma (AITL)), leukaemias (including acute lymphoblastic leukaemia (ALL), acute myeloid leukaemia (AML), and chronic myeloid leukaemia (CML)) and multiple myeloma, and of solid tumours (including glioma, breast cancer, non-small cell lung cancer (NSCLC) and squamous cell carcinomas (SCC) (including SCC of the head and neck, oesophagus, lung and ovary)).

[0093] According to a further feature of this aspect of the specification there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined herein for use in the treatment of lymphomas, including DLBCL, FL, BL and AITL.

[0094] According to a further feature of this aspect of the specification there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined herein for use in the treatment of DLBCL and FL.

[0095] According to a further feature of this aspect of the specification there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined herein for use in the treatment of leukaemias, including ALL, AML and CML.

[0096] According to a further feature of this aspect of the specification there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined herein for use in the treatment of solid tumours, including glioma, breast cancer, NSCLC and SCC.

[0097] According to a further feature of this aspect of the specification there is provided a method for treating haematological cancers such as lymphomas (including DLBCL, FL, BL and AITL), leukaemias (including ALL, AML and CML) and multiple myeloma, and of solid tumours (including glioma, breast cancer, NSCLC and SCC (including SCC of the head and neck, oesophagus, lung and ovary)) in a warm-blooded animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[0098] According to a further feature of this aspect of the specification there is provided a method for treating lymphomas, including DLBCL, FL, BL and AITL, in a warm-blooded animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[0099] According to a further feature of this aspect of the specification there is provided a method for treating DLBCL and FL, in a warm-blooded animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[00100] According to a further feature of this aspect of the specification there is provided a method for treating leukaemias, including ALL, AML and CML, in a warm-blooded animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[00101] According to a further feature of this aspect of the specification there is provided a method for treating solid tumours (including glioma, breast cancer, NSCLC and SCC (including SCC of the head and neck, oesophagus, lung and ovary)), in a warm-blooded animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

[00102] According to a further feature of this aspect of the specification there is provided the use of a compound, or a pharmaceutically acceptable salt thereof, as defined herein in the

manufacture of a medicament for use in the treatment of haematological cancers such as lymphomas (including DLBCL, FL, BL and AITL), leukaemias (including ALL, AML and CML) and multiple myeloma, and of solid tumours (including glioma, breast cancer, NSCLC and SCC (including SCC of the head and neck, oesophagus, lung and ovary)).

5 **[00103]** According to a further feature of this aspect of the specification there is provided the use of a compound, or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for use in the treatment of lymphomas, including DLBCL, FL, BL and AITL.

[00104] According to a further feature of this aspect of the specification there is provided the use
10 of a compound, or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for use in the treatment of DLBCL and FL.

[00105] According to a further feature of this aspect of the specification there is provided the use of a compound, or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for use in the treatment of leukaemias, including ALL, AML and
15 CML.

[00106] According to a further feature of this aspect of the specification there is provided the use of a compound, or a pharmaceutically acceptable salt thereof, as defined herein in the manufacture of a medicament for use in the treatment of solid tumours (including glioma, breast cancer, NSCLC and SCC (including SCC of the head and neck, oesophagus, lung and ovary)).

20 **[00107]** BCL6 has also been linked to other disease states outside the cancer field. These include HIV (Cai et al, *J. Virol.*, 2019, 93, e01073/1-e01073/15), sepsis (Zhang et al., *Int. Immunopharmacol.*, 2019, 75, 105789), graft-versus-host disease (Paz et al. *Blood*, 2019, 133, 94–99), non-alcoholic fatty liver disease including non-alcoholic steatohepatitis (Sommars et al., *eLife*, 2019, 8, e43922/1-e43922/25; Chikada et al., *Sci. Rep.*, 2020, 10, 9704) and auto-immune
25 diseases (Crotty, *Immunity*, 2014, 41, 529-542; Lintermann et al., *J. Exp. Med. Vol.*, 2008, 206(3), 561-576). Examples of auto-immune diseases or conditions include rheumatoid arthritis, systemic lupus erythematosus, Graves' disease, autoimmune haemolytic anaemia, multiple sclerosis, Type 1 diabetes, Goodpasture's syndrome, Hashimoto's thyroiditis, Guillain-Barre syndrome, immune thrombocytopenic purpura, atherosclerosis, Crohn's disease, ulcerative colitis, inflammatory bowel
30 disease, ankylosing spondylitis, seronegative spondylarthropathies, autoimmune thyroiditis, Sjogren's syndrome, Hughes' syndrome, psoriasis, psoriatic arthritis, myasthenia gravis, thrombocytopenic purpura, Addison's disease, primary biliary cirrhosis, diffuse scleroderma, polymyositis, dermatomyositis, autoimmune hepatitis, autoimmune sclerosing cholangitis, limited scleroderma, autoimmune uveitis, acquired haemophilia, pernicious anaemia, pemphigus,
35 pemphigoid and vitiligo.

[00108] In an embodiment, the present invention provides a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein for use in the treatment of auto-immune diseases (such as rheumatoid arthritis, systemic lupus erythematosus, Graves' disease, autoimmune haemolytic anaemia, multiple sclerosis, Type 1 diabetes, Goodpasture's syndrome, Hashimoto's thyroiditis, Guillain-Barre syndrome, immune thrombocytic purpura, atherosclerosis, Crohn's disease, ulcerative colitis, inflammatory bowel disease, ankylosing spondylitis, seronegative spondylarthropathies, autoimmune thyroiditis, Sjogren's syndrome, Hughes' syndrome, psoriasis, psoriatic arthritis, myasthenia gravis, thrombocytopenic purpura, Addison's disease, primary biliary cirrhosis, diffuse scleroderma, polymyositis, dermatomyositis, autoimmune hepatitis, autoimmune sclerosing cholangitis, limited scleroderma, autoimmune uveitis, acquired haemophilia, pernicious anaemia, pemphigus, pemphigoid or vitiligo), HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease or non-alcoholic steatohepatitis.

[00109] In an embodiment, there is provided the use of a compound, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition as defined herein in the manufacture of a medicament for use in the treatment of auto-immune diseases (such as rheumatoid arthritis, systemic lupus erythematosus, Graves' disease, autoimmune haemolytic anaemia, multiple sclerosis, Type 1 diabetes, Goodpasture's syndrome, Hashimoto's thyroiditis, Guillain-Barre syndrome, immune thrombocytic purpura, atherosclerosis, Crohn's disease, ulcerative colitis, inflammatory bowel disease, ankylosing spondylitis, seronegative spondylarthropathies, autoimmune thyroiditis, Sjogren's syndrome, Hughes' syndrome, psoriasis, psoriatic arthritis, myasthenia gravis, thrombocytopenic purpura, Addison's disease, primary biliary cirrhosis, diffuse scleroderma, polymyositis, dermatomyositis, autoimmune hepatitis, autoimmune sclerosing cholangitis, limited scleroderma, autoimmune uveitis, acquired haemophilia, pernicious anaemia, pemphigus, pemphigoid or vitiligo), HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease or non-alcoholic steatohepatitis.

[00110] In an embodiment, there is provided a method of treating auto-immune diseases (such as rheumatoid arthritis, systemic lupus erythematosus, Graves' disease, autoimmune haemolytic anaemia, multiple sclerosis, Type 1 diabetes, Goodpasture's syndrome, Hashimoto's thyroiditis, Guillain-Barre syndrome, immune thrombocytic purpura, atherosclerosis, Crohn's disease, ulcerative colitis, inflammatory bowel disease, ankylosing spondylitis, seronegative spondylarthropathies, autoimmune thyroiditis, Sjogren's syndrome, Hughes' syndrome, psoriasis, psoriatic arthritis, myasthenia gravis, thrombocytopenic purpura, Addison's disease, primary biliary cirrhosis, diffuse scleroderma, polymyositis, dermatomyositis, autoimmune hepatitis, autoimmune sclerosing cholangitis, limited scleroderma, autoimmune uveitis, acquired haemophilia, pernicious anaemia, pemphigus, pemphigoid or vitiligo), HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease or non-alcoholic steatohepatitis, in a warm-blooded

animal such as man that is in need of such treatment, which comprises administering an effective amount of a compound, or a pharmaceutically acceptable salt thereof, as defined herein.

Routes of Administration

[00111] The compounds of the invention or pharmaceutical compositions comprising these compounds may be administered to a subject by any convenient route of administration, whether systemically, peripherally or topically (i.e., at the site of desired action).

[00112] Routes of administration include, but are not limited to, oral (e.g., by ingestion); buccal; sublingual; transdermal (including, e.g., by a patch, plaster, etc.); transmucosal (including, e.g., by a patch, plaster, etc.); intranasal (e.g., by nasal spray); ocular (e.g., by eye drops); pulmonary (e.g., by inhalation or insufflation therapy using, e.g., via an aerosol, e.g., through the mouth or nose); rectal (e.g., by suppository or enema); vaginal (e.g., by pessary); parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intra-arterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal; by implant of a depot or reservoir, for example, subcutaneously or intramuscularly.

Combination Therapies

[00113] The antiproliferative treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents:-

(i) other antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, oxaliplatin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan, temozolamide and nitrosoureas); antimetabolites (for example cytarabine, gemcitabine and antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, and hydroxyurea); antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and taxotere and polokinas inhibitors); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);

(ii) cytostatic agents such as antioestrogens (for example tamoxifen, fulvestrant, toremifene, raloxifene, droloxifene and idoxifene), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), steroid hormones, including progestogens (for example megestrol acetate) and corticosteroids (for example dexamethasone, prednisone and prednisolone),

aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;

(iii) anti-invasion agents [for example c-Src kinase family inhibitors like 4-(6-chloro-2,3-methylenedioxyanilino)-7-[2-(4-methylpiperazin-1-yl)ethoxy]-5-tetrahydropyran-4-yloxyquin-
5 azoline (AZD0530; International Patent Application WO 01/94341), *N*-(2-chloro-6-methylphenyl)-2-[6-[4-(2-hydroxyethyl)piperazin-1-yl]-2-methylpyrimidin-4-ylamino]thiazole-5-carboxamide (dasatinib, BMS-354825; *J. Med. Chem.*, 2004, 47, 6658-6661) and bosutinib (SKI-606), and metalloproteinase inhibitors like marimastat, inhibitors of urokinase plasminogen activator receptor function or antibodies to Heparanase];

(iv) inhibitors of growth factor function: for example such inhibitors include growth factor antibodies and growth factor receptor antibodies (for example the anti-erbB2 antibody trastuzumab [HerceptinTM], the anti-EGFR antibody panitumumab, the anti-erbB1 antibody cetuximab [Erbix, C225] and any growth factor or growth factor receptor antibodies disclosed by Stern *et al.* (Critical reviews in oncology/haematology, 2005, Vol. 54, pp11-29); such inhibitors
15 also include tyrosine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as *N*-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, ZD1839), *N*-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-*N*-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)-quinazolin-4-amine (CI 1033), erbB2 tyrosine kinase
20 inhibitors such as lapatinib); inhibitors of the hepatocyte growth factor family; inhibitors of the insulin growth factor family; inhibitors of the platelet-derived growth factor family such as imatinib and/or nilotinib (AMN107); inhibitors of serine/threonine kinases (for example Ras/Raf signalling inhibitors such as farnesyl transferase inhibitors, for example sorafenib (BAY 43-9006), tipifarnib (R115777) and lonafarnib (SCH66336)), inhibitors of cell signalling through MEK and/or AKT
25 kinases, c-kit inhibitors, abl kinase inhibitors, PI3 kinase inhibitors, Plt3 kinase inhibitors, CSF-1R kinase inhibitors, IGF receptor (insulin-like growth factor) kinase inhibitors; aurora kinase inhibitors (for example AZD1152, PH739358, VX-680, MLN8054, R763, MP235, MP529, VX-528 AND AX39459) and cyclin dependent kinase inhibitors such as CDK2 and/or CDK4 inhibitors;

(v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth
30 factor, [for example the anti-vascular endothelial cell growth factor antibody bevacizumab (AvastinTM) and for example, a VEGF receptor tyrosine kinase inhibitor such as vandetanib (ZD6474), vatalanib (PTK787), sunitinib (SU11248), axitinib (AG-013736), pazopanib (GW 786034) and 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (AZD2171; Example 240 within WO 00/47212), compounds such as those
35 disclosed in International Patent Applications WO97/22596, WO 97/30035, WO 97/32856 and WO 98/13354 and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin)];

- (vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 and WO 02/08213;
- (vii) an endothelin receptor antagonist, for example zibotentan (ZD4054) or atrasentan;
- 5 (viii) inhibitors of DNA damage response (DDR) pathways, for example inhibitors of ATM (such as KU-60019, M3541, AZD0156 or AZD1390) and ATR (such as M6620, AZD6738 or BAY1895344);
- (ix) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- 10 (x) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy; and
- 15 (xi) immunotherapy approaches, including for example *ex-vivo* and *in-vivo* approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine-transfected dendritic cells, approaches using cytokine-transfected tumour cell lines, approaches using
- 20 anti-idiotypic antibodies and approaches using checkpoint inhibitors (such as inhibitors targeting CTLA-4 (e.g. ipilimumab), PD-1 (e.g. nivolumab, pembrolizumab or cemiplimab) or PD-L1 (e.g. atezolizumab, avelumab or durvalumab)).

[00114] In a particular embodiment, the antiproliferative treatment defined hereinbefore may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or

25 chemotherapy, wherein the chemotherapy may include one or more anti-tumour agents selected from procarbazine, carmustine, cytarabine, lomustine, irinotecan, temozolomide, cisplatin, carboplatin, doxorubicin, methotrexate, etoposide, cyclophosphamide, ifosfamide, and vincristine.

[00115] In another particular embodiment, the antiproliferative treatment defined hereinbefore

30 may involve, in addition to the compound of the invention, conventional surgery or radiotherapy or chemotherapy, wherein the chemotherapy may include one or more chemotherapeutic agents selected from a BCL-2 family inhibitor (e.g. Venetoclax and/or navitoclax), a BTK inhibitor (e.g. Ibrutinib, Acalabrutinib, Tirabrutinib (ONO/GS-4059), BGB-3111 or Spebrutinib (CC-292), a TNF inhibitor (e.g. Lenalidomide), an EZH2 inhibitor (e.g. Tazemetostat, CPI-0209, CPI-1205, DS-

35 3201, HH2853 PF-06821497, GSK126, GSK343, SHR2554 or EPZ011989), a corticosteroid (e.g. dexamethasone, prednisone or prednisolone), a HDAC inhibitor (e.g. panobinostat, entinostat,

romidepsin, belinostat or vorinostat), an ATR or ATM kinase inhibitor and an EGFR tyrosine kinase inhibitor.

[00116] Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate dosing of the individual components of the treatment. Such combination products employ the compounds of this invention within the dosage range described hereinbefore and the other pharmaceutically-active agent within its approved dosage range.

[00117] According to this aspect of the invention there is provided a combination for use in the treatment of a cancer (for example a cancer involving a solid tumour) comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and another anti-tumour agent.

[00118] According to this aspect of the invention there is provided a combination for use in the treatment of a proliferative condition, such as cancer (for example a cancer involving a solid tumour), comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and any one of the anti-tumour agents listed herein above.

[00119] According to this aspect of the invention there is provided a combination for use in the treatment of a cancer comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and a tyrosine kinase inhibitor.

[00120] According to this aspect of the invention there is provided a combination for use in the treatment of leukaemia (such as ALL or CML) comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and a tyrosine kinase inhibitor.

[00121] According to this aspect of the invention there is provided a combination for use in the treatment of lymphomas comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and an EZH2 inhibitor.

[00122] According to this aspect of the invention there is further provided a combination for use in the treatment of cancer comprising a compound of the invention as defined hereinbefore, or a pharmaceutically acceptable salt, hydrate or solvate thereof, and a corticosteroid.

[00123] In a further aspect of the invention there is provided a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate thereof, for use in the treatment of cancer in combination with another anti-tumour agent, optionally selected from one listed herein above.

[00124] In a further aspect of the invention there is provided a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate thereof, for use in the treatment of cancer in combination with a tyrosine kinase inhibitor, optionally selected from one listed herein above.

[00125] In a further aspect of the invention there is provided a compound of the invention or a

pharmaceutically acceptable salt, hydrate or solvate thereof, for use in the treatment of leukaemia (such as ALL or CML) in combination with a tyrosine kinase inhibitor, optionally selected from one listed herein above.

[00126] In a further aspect of the invention there is provided a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate thereof, for use in the treatment of cancer in combination with an EZH2 inhibitor, optionally selected from one listed herein above.

[00127] In a further aspect of the invention there is provided a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate thereof, for use in the treatment of lymphomas in combination with an EZH2 inhibitor, optionally selected from one listed herein above.

[00128] Herein, where the term “combination” is used it is to be understood that this refers to simultaneous, separate or sequential administration. In one aspect of the invention “combination” refers to simultaneous administration. In another aspect of the invention “combination” refers to separate administration. In a further aspect of the invention “combination” refers to sequential administration. Where the administration is sequential or separate, the delay in administering the second component should not be such as to lose the beneficial effect of the combination.

[00129] According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of the invention, or a pharmaceutically acceptable salt, hydrate or solvate thereof, in combination with an anti-tumour agent (optionally selected from one listed herein above), in association with a pharmaceutically acceptable diluent or carrier.

EXAMPLES

[00130] Particular embodiments of the invention are further described hereinafter, with reference to the accompanying drawings, in which:

Figure 1 shows the flow cytometry splenocyte hierarchical gating strategy: GC B cells were gated according to the following hierarchical gating strategy: viable->single cell (doublet exclusion)-> lymphocytes-> CD19+CD20+ ->GL7+CD95+ (light squares); non-GC B cells were gated according to the following hierarchical gating strategy: viable->single cell (doublet exclusion)-> lymphocytes-> CD19+CD20+ ->GL7-CD95- (dark squares).

Figure 2 shows the flow cytometry expression levels of BCL6 (geometric mean fluorescence intensity - GeoMFI) in non-GC B cells, without (vehicle) or with Example 4b at various concentrations. Stimulated and unstimulated B cell BCL6 expression is indicated with the upper and middle dashed lines respectively, while the lower dashed line represents the biological background expression for CD4 stimulated T cells.

Figure 3 shows the flow cytometry expression levels of Blimp (geometric mean fluorescence intensity - GeoMFI) in GC B cells, without (vehicle) or with Example 4b at various concentrations. Unstimulated and stimulated non-GC B cell Blimp expression is indicated with the dashed line.

Figure 4 shows the number of GC B cells as defined by the expression of the lineage markers GL7 and CD95 on CD19+CD21+ spleen cells without (vehicle) or with Example 4b (at 10 μ M).

Abbreviations

	APCI	Atmospheric pressure chemical ionization
5	aq.	Aqueous
	Ar	Argon
	Boc	t-butyloxycarbonyl protecting group
	br	broad (in NMR spectrum)
	conc.	concentrated
10	d	doublet (in NMR spectrum)
	DCM	dichloromethane
	DIPEA	N,N-diisopropylethylamine
	DMF	N,N-dimethylformamide
	DMSO	dimethylsulfoxide
15	ESI	electrospray ionisation
	EtOAc	ethyl acetate
	EtOH	ethanol
	FID	free induction decay
	h	hour(s)
20	HATU	N-[(Dimethylamino)-1H-1,2,3-triazolo-[4,5-b]pyridin-1-ylmethylene]-N-methylmethanaminium hexafluorophosphate N-oxide
	HPLC	High Performance Liquid Chromatography
	HRMS	high resolution mass spectrometry
	Josiphos	ligand for Pd catalysed reactions: (R)-1-[(S)-2-
25		(Dicyclohexylphosphino)ferrocenyl]ethyl- <i>tert</i> -butylphosphine
	KP-Sil	Biotage KP-Sil (50uM irregular silica)
	LCMS	liquid chromatography and mass spectrometry
	MeOH	methanol
	MeCN	acetonitrile
30	MS	mass spectrometry
	Ms	mesyl (methanesulfonyl)
	m	multiplet (in NMR spectrum)
	MHz	megahertz
	min	minute(s)
35	mins	minute(s)
	mL	milliliter(s)
	m/z	mass to charge ratio
	NMP	N-methylpyrrolidinone

	NMR	nuclear magnetic resonance
	Pd/C	palladium on activated charcoal
	ppm	parts per million
	q	quartet (in NMR spectrum)
5	QToF	Quadrupole Time-of-flight
	quin.	quintet (in NMR spectrum)
	Rt, RT	retention time (in LCMS)
	rt	room temperature
	s	singlet (in NMR spectrum)
10	SCX-2	strong cation exchange (e.g. Isolute® SCX-2 columns)
	t	triplet (in NMR spectrum)
	Tf	triflate (trifluoromethane sulfonate)
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
15	T3P	propylphosphonic anhydride
	uL	microliters
	UPLC	Ultra-Performance Liquid Chromatography
	Xantphos	4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene

Analytical methods: LCMS

20 Method T2

[00131] LC/MS and HRMS analysis was performed on an Agilent 1200 series HPLC and diode array detector coupled to a 6210 time of flight mass spectrometer with dual multimode APCI/ESI source. Analytical separation was carried out at 40°C on a Merck Chromolith Flash column (RP-18e, 25 x 2 mm) using a flow rate of 1.5 mL/min in a 2 minute gradient elution with detection at 254 nm. The mobile phase was a mixture of methanol (solvent A) and water (solvent B), both containing formic acid at 0.1%. Gradient elution was as follows: 5:95 (A/B) to 100:0 (A/B) over 1.25 min, 100:0 (A/B) for 0.5 min, and then reversion back to 5:95 (A/B) over 0.05 min, finally 5:95 (A/B) for 0.2 min.

Method T4

30 [00132] As for method T2 except at 30°C, using a flow rate of 0.75 mL/min in a 4 minute gradient elution as follows: 5:95 (A/B) to 100:0 (A/B) over 2.5 min, 100:0 (A/B) for 1 min, and then reversion back to 5:95 (A/B) over 0.1 min, finally 5:95 (A/B) for 0.4 min.

Method X2

35 [00133] LC/MS and HRMS analysis was performed on a Waters Acquity UPLC and diode array detector coupled to a Waters G2 QToF mass spectrometer fitted with a multimode ESI/APCI source. Analytical separation was carried out at 30°C on a Phenomenex Kinetex C18 column (30

x 2.1 mm, 2.6u, 100A) using a flow rate of 0.5 mL/min in a 2 minute gradient elution with detection at 254 nm. The mobile phase was a mixture of methanol (solvent A) and water (solvent B), both containing formic acid at 0.1%. Gradient elution was as follows: 10:90 (A/B) to 90:10 (A/B) over 1.25 min, 90:10 (A/B) for 0.5 min, and then reversion back to 10:90 (A/B) over 0.15 min, finally 10:90 (A/B) for 0.1 min.

Method X4

[00134] As for method X2, except using a flow rate of 0.3 mL/min in a 4 minute gradient elution as follows: 10:90 (A/B) to 90:10 (A/B) over 3 min, 90:10 (A/B) for 0.5 min, and then reversion back to 10:90 (A/B) over 0.3 min, finally 10:90 (A/B) for 0.2 min.

Analytical methods: NMR

[00135] NMR data was collected on a Bruker Avance 500 spectrometer equipped with a 5 mm BBO/QNP probe, or on a Bruker Avance Neo 600 spectrometer equipped with a 5 mm TCI Cryo-Probe. The ^1H and ^{13}C spectra were referenced to the internal deuterated solvent. All NMR data were acquired at the temperature of 298 K. All data were acquired and processed using Bruker Topspin 2.1 or Bruker Topspin 4.

[00136] The ^1H NMR spectra were acquired using a Bruker standard 1D zg30 pulse sequence with 16 scans. The sweep width was 20.5 ppm, and the FID contained 64k time-domain data points.

Purification methods


[00137] Unless otherwise described in the text, preparative HPLC purification was carried out on an Agilent 6120 MS-Prep LC using an ACE 5 C18-PFP 250 x 21.2 mm (or 30mm) column using a 15 min gradient of water:methanol (both modified with 0.1% formic acid) – for example 90:10 to 0:100 or 60:40 to 0:100 – at a flow rate of 20 mLmin⁻¹ (or 40 mLmin⁻¹ for the 30mm column).

[00138] Flash column chromatography was carried out using prepacked Biotage SNAP KP-Sil columns. Reverse phase chromatography was carried out using a Biotage SNAP Ultra C-18 12g and 30g columns as required.

Example Compounds

Example 1a: (S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




76










[00139] A mixture of (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A1, 11.5 mg, 0.025 mmol), 3,3-difluoroazetidine hydrochloride (8 mg, 0.062 mmol) and DIPEA (80 μ L, 0.46 mmol) in acetonitrile (0.6 mL) under argon was heated to 80 $^{\circ}$ C for 1 hour, then further heated under microwave irradiation to 120 $^{\circ}$ C for 19h. The crude product was concentrated under reduced pressure, dissolved in DMSO (1 mL) and purified by reverse-phase chromatography (12g Biotage Ultra C18, 10-60-90-100% methanol in water, 0.1% formic acid modifier). Fractions containing product were combined and passed through an SCX-2 (2g) column, washing with methanol (10 mL) and eluting with 2M methanolic ammonia (25 mL). The ammonia fractions were combined and evaporated under reduced pressure to give the title compound as an off-white solid (4.3 mg, 0.008 mmol). ^1H NMR (600 MHz, Methanol- d_4) δ 8.14 (d, J = 2.1 Hz, 1 H), 8.02 (s, 1 H), 7.95 (dd, J = 9.1, 2.1 Hz, 1 H), 7.55 (d, J = 9.1 Hz, 1 H), 4.52–4.34 (m, 6 H), 3.72 (s, 3 H), 3.36–3.29 (m, 1 H), 1.44–1.37 (m, 1 H), 0.83–0.76 (m, 1 H), 0.71–0.65 (m, 1 H), 0.65–0.59 (m, 1 H), 0.39–0.33 (m, 1 H); HRMS (Method X4) RT = 3.19 min; m/z calcd for $\text{C}_{23}\text{H}_{22}\text{ClF}_4\text{N}_6\text{O}_2^+$ $[\text{M}+\text{H}]^+$: 525.1429, Found: 525.1412.



[00140] The following tabulated examples were prepared by a method analogous to that used for the preparation of **Example 1a**, using the appropriate amine (as free base or as hydrochloride salt). For examples 1b – 1e, the reaction mixtures were heated to 140 $^{\circ}$ C for 1 – 2h using microwave irradiation. For examples 1f – 1l, the reaction mixtures were heated to 80 $^{\circ}$ C for 1.5 – 3h. Purification was carried out by reverse-phase chromatography, and in some cases products were desalted using an SCX-2 column.

Example	Data and comments	Amine used
Example 1b: (S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	^1H NMR (600 MHz, Methanol- d_4) δ 8.04 (d, J = 2.2 Hz, 1 H), 8.00 (s, 1 H), 7.92 (dd, J = 9.1, 2.2 Hz, 1 H), 7.56 (d, J = 9.1 Hz, 1 H), 4.53–4.38 (m, 2 H), 4.22 (d, J = 18.3 Hz, 1 H), 4.16 (d, J = 18.3 Hz, 1 H), 3.92–3.87 (m, 1 H), 3.86–3.81 (m, 1 H), 3.72 (s, 3 H), 3.40 (t, J = 5.4 Hz, 2 H), 3.35–3.28 (m, 1 H), 2.78–2.73 (m, 1 H), 1.45–1.36 (m, 1 H), 0.83–0.76 (m, 3 H), 0.71–0.67 (m, 2 H), 0.66–0.58 (m, 2 H), 0.37–0.31 (m, 1 H); HRMS m/z calcd for $\text{C}_{27}\text{H}_{29}\text{ClF}_2\text{N}_7\text{O}_3^+$	1-cyclopropyl-piperazin-2-one


	[M+H] ⁺ : 572.1989, Found: 572.1987.	
Example 1c: (S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.02-7.99 (m, 2 H), 7.96 (dd, <i>J</i> = 9.1, 2.3 Hz, 1 H), 7.57 (d, <i>J</i> = 9.1 Hz, 1 H), 4.53-4.38 (m, 3 H), 4.26 (dd, <i>J</i> = 13.4, 2.7 Hz, 1 H), 3.93 (d, <i>J</i> = 18.5 Hz, 1 H), 3.72 (s, 3 H), 3.62-3.57 (m, 1 H), 3.51 (dd, <i>J</i> = 13.4, 3.6 Hz, 1 H), 3.35-3.28 (m, 1 H), 2.97 (s, 3 H), 1.43-1.36 (m, 1 H), 1.21 (d, <i>J</i> = 6.4 Hz, 3 H), 0.81-0.75 (m, 1 H), 0.68-0.57 (m, 2 H), 0.38-0.33 (m, 1 H); HRMS <i>m/z</i> calcd for C₂₆H₂₉ClF₂N₇O₃⁺ [M+H]⁺: 560.1989, Found: 560.1982.	(S)-1,6-dimethylpiperazin-2-one hydrochloride
Example 1d: (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:formic acid (1:1) 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.24 (s, 1H), 8.11 (s, 1H), 7.94 (dd, <i>J</i> = 9.0, 2.4 Hz, 1H), 7.92 (s, 1H), 7.57 (d, <i>J</i> = 9.1 Hz, 1H), 4.51 (ddd, <i>J</i> = 15.8, 13.2, 2.4 Hz, 1H), 4.43 (ddd, <i>J</i> = 23.9, 13.3, 7.2 Hz, 1H), 3.74 (s, 3H), 3.37 (s, 2H), 3.32 - 3.30 (m, 1H), 1.47 - 1.40 (m, 1H), 1.15 (s, 6H), 0.84 - 0.76 (m, 1H), 0.72 - 0.65 (m, 1H), 0.65 - 0.59 (m, 1H), 0.40 - 0.34 (m, 1H). HRMS <i>m/z</i> calcd for C₂₄H₂₈ClF₂N₆O₃⁺ [M+H]⁺: 521.1874, found 521.1841.	1-amino-2-methylpropan-2-ol
Example 1e: (S)-10-((2-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.11 (s, 1H), 8.02 - 7.95 (m, 2H), 7.57 (d, <i>J</i> = 9.1 Hz, 1H), 4.85 - 4.79 (m, 1H), 4.66 (s, 1H), 4.55 - 4.37 (m, 2H), 3.86 - 3.79 (m, 2H), 3.74 (s, 3H), 3.52 - 3.44 (m, 2H), 3.37 (s, 1H), 2.00 - 1.89 (m, 2H), 1.46 - 1.37 (m, 1H), 0.85 - 0.78 (m, 1H), 0.74 - 0.67 (m, 1H), 0.65 - 0.60 (m, 1H), 0.41 - 0.34 (m, 1H). HRMS <i>m/z</i> calcd for C₂₅H₂₆ClF₂N₆O₃⁺ [M+H]⁺: 531.1717; found 531.1725	(1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptane
Example 1f: N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.08 (d, <i>J</i> = 9.1 Hz, 1H), 7.97 - 7.92 (m, 2H), 7.54 (d, <i>J</i> = 9.1 Hz, 1H), 4.78 - 4.67 (m, 1H), 4.57 - 4.34 (m, 3H), 3.80 - 3.73 (m, 1H), 3.73 (s, 3H), 3.40 - 3.26 (m, 1H), 2.43 (dd, <i>J</i> = 12.6, 11.2 Hz, 1H), 2.31 (dd, <i>J</i> = 13.1, 11.4 Hz, 1H), 1.97 (m, 4H), 1.66 (m, 1H), 1.46 - 1.36 (m, 1H), 1.14 - 1.05 (m, 1H), 0.94 (d, <i>J</i> = 6.5	N-((3S,5R)-5-methylpiperidin-3-yl)acetamide [Prepared from tert-butyl (3S,5R)-3-amino-5-methylpiperidine-1-

methylpiperidin-3-yl)acetamide 	Hz, 3H), 0.83 – 0.77 (m, 1H), 0.72 – 0.65 (m, 1H), 0.65 – 0.59 (m, 1H), 0.41 – 0.35 (m, 1H). HRMS: m/z calcd for C₂₈H₃₃ClF₂N₇O₃ [M+H]⁺: 588.2302, Found: 588.2307.	carboxylate by acylation with acetic anhydride followed by Boc deprotection.]
Example 1g: N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.08 (d, J = 9.1 Hz, 1H), 7.96 (s, 1H), 7.94 (s, 1H), 7.54 (d, J = 9.1 Hz, 1H), 4.76 – 4.71 (m, 1H), 4.56 – 4.37 (m, 3H), 3.79 – 3.74 (m, 1H), 3.73 (s, 3H), 3.38 – 3.27 (m, 1H), 2.44 (dd, J = 12.6, 11.1 Hz, 1H), 2.32 (dd, J = 13.1, 11.4 Hz, 1H), 2.03 – 1.98 (m, 1H), 1.96 (s, 3H), 1.71 – 1.61 (m, 1H), 1.45 – 1.38 (m, 1H), 1.10 (q, J = 12.1 Hz, 1H), 0.94 (d, J = 6.6 Hz, 3H), 0.84 – 0.78 (m, 1H), 0.72 – 0.65 (m, 1H), 0.65 – 0.58 (m, 1H), 0.41 – 0.33 (m, 1H). HRMS: m/z calcd for C₂₈H₃₃ClF₂N₇O₃ [M+H]⁺: 588.2302, Found: 588.2302.	N-((3R,5S)-5-methylpiperidin-3-yl)acetamide [Prepared from tert-butyl (3R,5S)-3-amino-5-methylpiperidine-1-carboxylate by acylation with acetic anhydride followed by Boc deprotection.]
Example 1h: (S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, DMSO-<i>d</i>₆) δ 8.85 (s, 1H), 8.09 (s, 1H), 8.02 (s, 1H), 7.72 (dd, J = 9.1, 2.2 Hz, 1H), 7.40 (d, J = 9.0 Hz, 1H), 6.25 (s, 1H), 4.65 (s, 1H), 4.55 – 4.25 (m, 3H), 3.55 (s, 3H), 3.26 – 3.17 (m, 1H), 2.36 – 2.11 (m, 9H), 1.97 – 1.86 (m, 1H), 1.51 – 1.42 (m, 1H), 1.35 – 1.27 (m, 1H), 1.10 – 0.98 (m, 1H), 0.84 (d, J = 5.5 Hz, 3H), 0.74 – 0.69 (m, 1H), 0.55 – 0.46 (m, 2H), 0.39 – 0.25 (m, 1H). HRMS: m/z calcd for C₂₈H₃₅ClF₂N₇O₂ [M+H]⁺: 574.2509, Found: 574.2503.	(3S,5R)-N,N,5-trimethylpiperidin-3-amine [prepared from tert-butyl (3S,5R)-3-amino-5-methylpiperidine-1-carboxylate by reductive amination with formaldehyde, followed by boc deprotection]
Example 1i: (S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.05 (d, J = 2.3 Hz, 1H), 7.97 (dd, J = 9.1, 2.3 Hz, 1H), 7.94 (s, 1H), 7.56 (d, J = 9.1 Hz, 1H), 4.54 – 4.35 (m, 2H), 4.28 – 4.20 (m, 1H), 4.11 – 4.02 (m, 1H), 3.72 (s, 3H), 3.67 – 3.59 (m, 1H), 3.39 – 3.26 (m, 1H), 3.20 – 3.12 (m, 1H), 3.08 (dd, 1H), 2.03 – 1.93 (m, 1H), 1.83 – 1.74 (m, 1H), 1.57 – 1.37 (m, 3H), 0.86 – 0.75 (m, 1H), 0.72 – 0.65 (m, 1H), 0.65 – 0.58 (m,	(S)-piperidin-3-ol hydrochloride

[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	1H), 0.42 – 0.32 (m, 1H). HRMS: m/z calcd for C₂₅H₂₈ClF₂N₆O₃ [M+H]⁺: 533.1874, Found: 533.1880	
Example 1j: (S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.06 (d, <i>J</i> = 2.4 Hz, 1H), 7.98 (dd, <i>J</i> = 9.1, 2.3 Hz, 1H), 7.95 (s, 1H), 7.56 (d, <i>J</i> = 9.1 Hz, 1H), 4.58 – 4.36 (m, 2H), 4.31 – 4.20 (m, 1H), 4.11 – 4.00 (m, 1H), 3.73 (s, 3H), 3.66 – 3.55 (m, 1H), 3.38 – 3.26 (m, 1H), 3.21 – 3.14 (m, 1H), 3.09 (dd, <i>J</i> = 12.9, 8.3 Hz, 1H), 2.02 – 1.90 (m, 1H), 1.86 – 1.73 (m, 1H), 1.59 – 1.34 (m, 3H), 0.87 – 0.76 (m, 1H), 0.73 – 0.64 (m, 1H), 0.64 – 0.56 (m, 1H), 0.44 – 0.33 (m, 1H). HRMS: m/z calcd for C₂₅H₂₈ClF₂N₆O₃ [M+H]⁺: 533.1874, Found: 533.1882	(R)-piperidin-3-ol hydrochloride
Example 1k: (S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.07 (d, <i>J</i> = 2.3 Hz, 1H), 7.96 (dd, <i>J</i> = 9.1, 2.3 Hz, 1H), 7.93 (s, 1H), 7.54 (d, <i>J</i> = 9.2 Hz, 1H), 4.59 – 4.29 (m, 4H), 3.73 (s, 3H), 3.39 – 3.27 (m, 1H), 2.83 (td, <i>J</i> = 12.7, 3.0 Hz, 1H), 2.51 (dd, <i>J</i> = 12.8, 10.6 Hz, 1H), 1.88 – 1.79 (m, 1H), 1.73 – 1.62 (m, 1H), 1.60 – 1.52 (m, 1H), 1.51 – 1.37 (m, 2H), 1.22 – 1.13 (m, 1H), 0.90 (d, <i>J</i> = 6.6 Hz, 3H), 0.83 – 0.76 (m, 1H), 0.71 – 0.64 (m, 1H), 0.65 – 0.58 (m, 1H), 0.41 – 0.31 (m, 1H). HRMS: m/z calcd for C₂₆H₃₀ClF₂N₆O₂ [M+H]⁺: 531.2081, Found: 531.2076	(S)-3-methylpiperidine hydrochloride
Example 1l: (S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	¹H NMR (600 MHz, Methanol-<i>d</i>₄) δ 8.05 (d, <i>J</i> = 2.3 Hz, 1H), 7.98 (dd, <i>J</i> = 9.1, 2.3 Hz, 1H), 7.93 (s, 1H), 7.54 (d, <i>J</i> = 9.1 Hz, 1H), 4.53 – 4.35 (m, 5H), 3.72 (s, 3H), 2.84 (ddd, <i>J</i> = 13.1, 11.9, 3.0 Hz, 1H), 2.51 (dd, <i>J</i> = 13.0, 10.6 Hz, 1H), 1.87 – 1.78 (m, 1H), 1.72 – 1.65 (m, 1H), 1.62 – 1.37 (m, 3H), 1.22 – 1.12 (m, 1H), 0.90 (d, <i>J</i> = 6.7 Hz, 3H), 0.85 – 0.77 (m, 1H), 0.70 – 0.65 (m, 1H), 0.64 – 0.59 (m, 1H), 0.41 – 0.33 (m, 1H). HRMS: m/z calcd for	(R)-3-methylpiperidine hydrochloride

	$C_{26}H_{30}ClF_2N_6O_2$ $[M+H]^+$: 531.2081, Found: 531.2084	
Example 1m: (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	^1H NMR (500 MHz, DMSO- d_6) δ 8.89 (s, 1H), 8.26 (d, J = 2.3 Hz, 1H), 8.07 (s, 1H), 7.77 (dd, J = 9.1, 2.2 Hz, 1H), 7.43 (d, J = 9.1 Hz, 1H), 6.21 (s, 1H), 4.55 – 4.27 (m, 2H), 4.16 – 3.93 (m, 4H), 3.56 (s, 3H), 3.29 – 3.18 (m, 1H), 1.56 (d, J = 22.1 Hz, 3H), 1.37 – 1.27 (m, 1H), 0.75 – 0.67 (m, 1H), 0.60 – 0.48 (m, 2H), 0.43 – 0.26 (m, 1H). HRMS: m/z calcd for $C_{24}H_{25}ClF_3N_6O_2$ $[M+H]^+$: 521.1680, Found: 521.1675	3-fluoro-3-methylazetidine hydrochloride
Example 1n: (S)-10-((5-chloro-2-(3-hydroxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	^1H NMR (600 MHz, Methanol- d_4) δ 8.21 (d, J = 2.3 Hz, 1H), 8.04 (dd, J = 9.1, 2.3 Hz, 1H), 7.94 (s, 1H), 7.54 (d, J = 9.2 Hz, 1H), 4.56 – 4.35 (m, 2H), 4.04 – 3.89 (m, 4H), 3.72 (s, 3H), 3.41 – 3.25 (m, 1H), 1.51 (s, 3H), 1.47 – 1.36 (m, 1H), 0.83 – 0.77 (m, 1H), 0.77 – 0.69 (m, 1H), 0.68 – 0.61 (m, 1H), 0.49 – 0.31 (m, 1H). HRMS: m/z calcd for $C_{24}H_{26}ClF_2N_6O_3$ $[M+H]^+$: 519.1723, Found: 519.1720	3-methylazetidin-3-ol hydrochloride

Example 2a: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




5

10

[00141] A mixture of (S)-10-((2-chloro-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A2, 12.6 mg, 0.028 mmol), (3R,5S)-5-methylpiperidin-3-ol (Intermediate M2a, 6.4 mg, 0.0556 mmol) and DIPEA (20.00 μ L, 0.1159 mmol) in NMP (0.6 mL) was heated at 140°C for 12 h, then diluted with DMSO (0.8 mL) and purified by reverse-phase chromatography (Biotage 12 g Ultra C-18 column; 10-30-70-100% methanol in water (0.1% formic acid modifier)). Fractions containing product were combined and passed through an SCX-2 (2g) column, washing with methanol (20 mL) and eluting


with 2M methanolic ammonia (25 mL). The ammonia fractions were combined and evaporated under reduced pressure to give the title compound as an off-white solid (7.5 mg, 0.014 mmol). ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.03 (dd, *J* = 9.1, 2.1 Hz, 1 H), 7.98 (d, *J* = 2.1 Hz, 1 H), 7.84 (d, *J*_{HF} = 3.7 Hz, 1 H), 7.54 (d, *J* = 9.1 Hz, 1 H), 4.72–4.66 (m, 1 H), 4.53–4.35 (m, 3 H), 3.70 (s, 3 H), 3.54–3.48 (m, 1 H), 3.34–3.29 (m, 1 H), 2.46 (dd, *J* = 12.2, 10.7 Hz, 1 H), 2.29 (app. t, *J* = 12.2 Hz, 1H), 2.09–2.02 (m, 1 H), 1.66–1.56 (m, 1 H), 1.45–1.37 (m, 1 H), 1.06 (q, *J* = 11.7 Hz, 1 H), 0.94 (d, *J* = 6.6 Hz, 3 H), 0.82–0.76 (m, 1 H), 0.69–0.57 (m, 2 H), 0.43–0.34 (m, 1 H); HRMS (Method X4) RT = 2.56 min; *m/z* calcd for C₂₆H₃₀F₃N₆O₃⁺ [M+H]⁺: 531.2332, Found: 531.2316.

Example 2b: (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one




[00142] 4,4-Difluoropiperidine hydrochloride (19 mg, 0.12 mmol), DIPEA (35 μ L, 0.20 mmol) and (S)-10-((2-chloro-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Intermediate A2, 23 mg, 0.05 mmol) were combined in NMP (0.40 mL) in a microwave vial. The vessel was sealed and heated at 140 °C for 24 h. The resulting mixture was purified by reverse-phase flash chromatography (40% - 85% methanol in water (0.1% formic acid)). Fractions containing product were loaded onto an Isolute SCX-2 column (1 g), and the column was washed with methanol and then eluted with 20% 1.4 M methanolic ammonia. The product was further purified by flash column chromatography (3-10% methanol in DCM), and the fractions containing product were combined, the solvent removed under reduced pressure to yield the title compound (18.5 mg). ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.06 (br s, 1H), 7.93 (dd, *J* = 9.1, 2.3 Hz, 1H), 7.89 (d, *J* = 2.3 Hz, 1H), 7.54 (br d, *J* = 9.1 Hz, 1H), 4.55 – 4.35 (m, 2H), 3.82 (m, 4H), 3.70 (s, 3H), 3.37 – 3.23 (m, 1H), 2.00 – 1.86 (m, 4H), 1.48 – 1.33 (m, 1H), 0.83 – 0.76 (m, 1H), 0.68 – 0.58 (m, 2H), 0.39 – 0.33 (m, 1H). LCMS (Method T4): Rt 2.91min, *m/z* calcd for C₂₅H₂₆F₅N₆O₂⁺ [M+H]⁺: 537.2032, Found: 537.2036.

Example 3a: (S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-*d*3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*][1,8]naphthyridin-6(7H)-one



[00143] A mixture of 3-oxa-8-azabicyclo[3.2.1]octane hydrochloride (13.4 mg, 0.09 mmol), DIPEA (23 μ L, 0.13 mmol) and (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthayridin-6(7H)-one (Intermediate A3, 9 mg, 0.019 mmol) in NMP (0.8 mL) under argon was heated at 140 °C in a heating block for 10 h. The crude product was diluted with DMSO (1 mL) and purified by reverse-phase chromatography (12g Biotage Ultra C18, 10-60-90-100% methanol in water, 0.1% formic acid modifier). Fractions containing product were combined and passed through an SCX-2 (2g) column, washing with methanol (15 mL) and eluting with 2M methanolic ammonia (30 mL). The ammonia fractions were combined and evaporated under reduced pressure to give the title compound as an off-white solid (8 mg, 0.015 mmol). ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.85 (d, *J* = 2.3 Hz, 1 H), 8.46 (d, *J* = 2.3 Hz, 1 H), 8.00 (s, 1 H), 4.54–4.38 (m, 4 H), 3.71 (d, *J* = 10.8 Hz, 2 H), 3.55 (d, *J* = 10.8 Hz, 2 H), 3.35–3.28 (m, 1 H), 2.04–1.91 (m, 4 H), 1.41–1.34 (m, 1 H), 0.82–0.75 (m, 1 H), 0.68–0.56 (m, 2 H), 0.40–0.32 (m, 1 H); HRMS (Method T4) RT = 2.96 min; *m/z* calcd for C₂₅H₂₄D₃ClF₂N₇O₃⁺ [M+H]⁺: 549.2015, Found: 549.2015.


Example 4a: (S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



[00144] A mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 11 mg, 0.034 mmol), 5-(4,5-dichloropyridin-2-yl)-3-methyl-1,2,4-oxadiazole (Intermediate C1, 9.8 mg, 0.043 mmol, cesium carbonate (90.8 mg, 0.28 mmol), Tris(dibenzylideneacetone)dipalladium(0) (3 mg, 0.003 mmol), palladium (II) acetate (3.2 mg), and Xantphos (11.9 mg, 0.021 mmol) in DMF (0.35 mL) and toluene (0.35 mL) under argon was heated to 100 °C for 90 min. The reaction mixture was cooled to rt. Water (10 mL) was added and the aqueous mixture was extracted with EtOAc (5 x 10 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na₂SO₄) and concentrated under reduced pressure. The crude product was dissolved in DMSO (1.0 mL) and

directly purified by reverse-phase chromatography (Biotage 12 g Ultra C-18 column; 10-60-80-100% methanol in water (0.1% formic acid modifier)) affording the title compound (1.7 mg, 0.003 mmol) as a pale yellow solid. ¹H NMR (600 MHz, CDCl₃) δ 8.57 (s, 1H), 7.61 (s, 1H), 7.50 - 7.46 (m, 2H), 7.45 - 7.43 (m, 1H), 6.75 (s, 1H), 4.65 (dd, *J* = 16.2, 13.3 Hz, 1H), 4.44 (ddd, *J* = 26.8, 13.3, 6.1 Hz, 1H), 4.21 (br s, 1H), 3.77 (s, 3H), 3.38 - 3.31 (m, 1H), 2.46 (s, 3H), 1.34 - 1.28 (m, 1H), 0.87 - 0.82 (m, 1H), 0.72 - 0.67 (m, 1H), 0.66-0.60 (m, 1H), 0.33 - 0.28 (m, 1H); HRMS (Method X4) RT = 2.93 min; *m/z* calcd for C₂₄H₂₂ClF₂N₆O₃⁺ [M+H]⁺: 515.1404, Found: 515.1397.

Example 4b: (S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




[00145] A mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 10 mg, 0.031 mmol), 4-bromo-3-chloro-2-fluoropyridine (8 mg, 0.038 mmol), cesium carbonate (81 mg, 0.25 mmol), tris(dibenzylideneacetone)dipalladium(0) (3.9 mg, 0.004 mmol) and Xantphos (11.6 mg, 0.020 mmol) in DMF (0.3 mL) and toluene (0.3 mL) under argon was heated at 80°C for 90 min. Water (15 mL) was added and the aqueous mixture was extracted with CH₂Cl₂ (4 x 10 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na₂SO₄), and stirred overnight in the presence of MP-TMT to remove residual Pd. The beads were filtered off, washed with CH₂Cl₂ and the filtrate was concentrated under reduced pressure. The crude product was dissolved in DMSO (1.0 mL) and purified by reverse-phase chromatography (Biotage 12 g Ultra C-18 column; 10%-60%-80%-100% methanol in water (0.1% formic acid modifier)) affording the title compound (5.8 mg, 0.013 mmol) as an off-white solid. ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.00 (d, *J* = 2.3 Hz, 1 H), 7.68 (d, *J* = 5.9 Hz, 1 H), 7.62 (d, *J* = 9.0 Hz, 1 H), 7.54 (dd, *J* = 9.0, 2.3 Hz, 1 H), 6.69 (d, *J* = 5.9 Hz, 1 H), 4.53–4.37 (m, 2 H), 3.73 (s, 3 H), 3.28 (ddd, *J* = 18.8, 10.1, 5.1 Hz, 1 H), 1.42–1.35 (m, 1 H), 0.80–0.74 (m, 1 H), 0.66–0.55 (m, 2 H), 0.35–0.29 (m, 1 H); HRMS (Method X4) RT = 2.91 min; *m/z* calcd for C₂₁H₁₉ClF₃N₄O₂⁺ [M+H]⁺: 451.1143, Found: 451.1139.

[00146] Example 4b was also prepared by the following procedure; (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 1.7 g, 5.3 mmol), 4-bromo-3-chloro-2-fluoropyridine (1.41 g, 6.7 mmol), palladium (II) acetate (0.143 g, 0.64 mmol), 4,5-bis(diphenylphosphino)-9,9-dimethylxanthene (0.456 g, 0.79 mmol) and cesium carbonate (2.7 g, 8.29 mmol) were combined under N₂ in 1,4-dioxane (50mL) and the resulting mixture was heated at reflux for 2 h. The mixture was then cooled, and the volatiles were removed under reduced pressure. The residue was purified by flash column chromatography


(DCM/Celite dry-loading, 0-100% EtOAc/n-heptane), giving the title compound (1.6 g) as a yellow/brown solid. This was further purified by trituration in diethyl ether:propan-2-ol (9:1), followed by lyophilisation from a MeCN/H₂O suspension to give the title compound as a white solid. ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.84 (s, 1H), 8.08 (d, *J* = 2.0 Hz, 1H), 7.71 (d, *J* = 5.8 Hz, 1H), 7.50 - 7.57 (m, 1H), 7.43 - 7.50 (m, 1H), 6.53 (d, *J* = 5.8 Hz, 1H), 6.31 - 6.45 (m, 1H), 4.32 - 4.52 (m, 2H), 3.58 (s, 3 H), 3.15 - 3.28 (m, 1H), 1.23 - 1.34 (m, 1H), 0.64 - 0.76 (m, 1H), 0.44 - 0.56 (m, 2H), 0.25 - 0.37 (m, 1H). ¹⁹F NMR (376 MHz, DMSO-*d*₆): δ -74.11 (s, 1F), -101.45 to -100.49 (m, 1F), -116.69 to -115.21 (m, 1F).

Example 4c: (S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one




[00147] To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one (Intermediate B1, 10.2 mg, 0.032 mmol), 2,3-dichloro-4-iodopyridine (10.3 mg, 0.038 mmol), cesium carbonate (80 mg, 0.246 mmol), tris(dibenzylideneacetone)dipalladium(0) (3.6 mg, 0.004 mmol) and Xantphos (11.4 mg, 0.020 mmol) under argon was added DMF (0.3 mL) and toluene (0.3 mL). The resulting mixture was heated to 80°C for 1 h, then cooled to rt. Water (15 mL) was added and the aqueous mixture was extracted with DCM (4 x 10 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na₂SO₄), and stirred overnight with MP-TMT to remove residual Pd. The beads were removed by filtration, washing with DCM, and the combined filtrate was concentrated under reduced pressure. The crude product was dissolved in DMSO (1.0 mL) and directly purified by reverse-phase chromatography (12 g Ultra C-18; 10%-60%-80%-100% methanol in water (containing 0.1% formic acid)) affording the title compound (7.4 mg, 50%, 0.0158 mmol) as an off-white solid. ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.00 (d, *J* = 2.3 Hz, 1 H), 7.83 (d, *J* = 5.8 Hz, 1 H), 7.63 (d, *J* = 9.0 Hz, 1 H), 7.54 (dd, *J* = 9.0, 2.3 Hz, 1 H), 6.74 (d, *J* = 5.8 Hz, 1 H), 4.53–4.38 (m, 2 H), 3.73 (s, 3 H), 3.32–3.24 (m, 1 H), 1.42–1.35 (m, 1 H), 0.80–0.74 (m, 1 H) 0.66–0.55 (m, 2 H), 0.35–0.28 (m, 1 H); HRMS (Method X4) RT = 2.99 min; *m/z* calcd for C₂₁H₁₉Cl₂F₂N₄O₂⁺ [M+H]⁺: 467.0853, Found: 467.0853.

Example 4d: (S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one




[00148] Example 4d was prepared by a method analogous to that used for the synthesis of example 4b, using 2-chloro-3-fluoro-4-iodo-pyridine. ¹H NMR (600 MHz, methanol-*d*₄) 7.95 (d, *J* = 2.3 Hz, 1H), 7.78 (d, *J* = 5.7 Hz, 1H), 7.60 (d, *J* = 9.0 Hz, 1H), 7.52 (dd, *J* = 9.0, 2.3 Hz, 1H), 6.94 (app. t, *J* = 6.0 Hz, 1H), 4.53 - 4.37 (m, 2H), 3.72 (s, 3H), 3.28 (ddd, *J* = 18.7, 10.1, 5.2 Hz, 1H), 1.43 - 1.35 (m, 1H), 0.81 - 0.74 (m, 1H), 0.66 - 0.55 (m, 2H), 0.36 - 0.30 (m, 1H); HRMS (Method X4) RT = 2.90 min; *m/z* calcd for C₂₁H₁₉ClF₃N₄O₂⁺ [M+H]⁺: 451.1148, Found: 451.1144.

Example 4e: (S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



[00149] To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 35 mg, 0.11 mmol), 3,4-dichloro-2-methylpyridine (26.5 mg, 0.16 mmol), tris(dibenzylideneacetone)dipalladium(0) (10 mg, 0.011 mmol), Xantphos (38 mg, 0.065 mmol) and cesium carbonate (284 mg, 0.87 mmol) under argon was added DMF (0.5 mL) and toluene (0.4 mL). The resulting mixture was heated at 140 °C under microwave irradiation for 90 min, then further heated at 100 °C for 6 h, then again at 140 °C for 3 hours. The reaction mixture was cooled to rt, concentrated under reduced pressure, redissolved in DCM (~20 mL) and washed with water (2 x 10 mL). The solvent was removed under reduced pressure and the reaction was purified by column chromatography (0% to 10% methanol in DCM) affording a dark orange oil. The product was loaded onto an Isolute SCX-2 column (2 g), and the column was washed with methanol and then eluted with 20% 1.4 M methanolic ammonia. The product was further purified by reverse phase column chromatography (C18 Sfar; 40 - 90% methanol in water (0.1% formic acid)) to afford the title compound as a white solid (25.2 mg). ¹H NMR (600 MHz, Methanol-*d*₄) δ 7.98 (d, *J* = 2.3 Hz, 1 H), 7.91 (d, *J* = 5.8 Hz, 1 H), 7.62 (d, *J* = 9.0 Hz, 1 H), 7.55 (dd, *J* = 9.0, 2.3 Hz, 1 H), 6.72 (d, *J* = 5.8 Hz, 1 H), 4.54 - 4.37 (m, 2 H), 3.74 (s, 3 H), 3.33 - 3.15 (m, 1 H), 2.56 (s, 3H), 1.43 - 1.35 (m, 1 H), 0.81 - 0.73 (m, 1 H), 0.67 - 0.55 (m, 2 H), 0.36 - 0.29 (m, 1 H); HRMS (Method X4) RT = 1.93 min; *m/z* calcd for C₂₂H₂₂ClF₂N₄O₂⁺ [M+H]⁺: 447.1399, Found: 447.1384.

Example 5a: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



5 Step 1: 2-chloro-5-fluoro-4-(methylthio)pyrimidine

[00150] Sodium thiomethoxide (127 mg, 1.81 mmol) was added to a stirred solution of 2,4-dichloro-5-fluoropyrimidine (0.3 g, 1.8 mmol) in THF (2 mL) and water (2 mL) at 0°C. The resulting mixture was stirred at rt for 24 h, then water (20 mL) was added and the aqueous mixture was extracted with EtOAc (2 x 20 mL). The organic extracts were combined, washed with brine (20 mL), dried (Na₂SO₄) and concentrated *in vacuo* affording the title compound (228 mg, 71%) as a white solid which was used without further purification. . ¹H NMR (500 MHz, CDCl₃) δ 8.10 (d, J = 1.4 Hz, 1 H), 2.62 (s, 3 H); LCMS (Method T2); RT 1.23 min; m/z 178.9987 [M+H]⁺.

10

Step 2: (1R,5S,7s)-9-(5-fluoro-4-(methylthio)pyrimidin-2-yl)-3-oxa-9-azabicyclo[3.3.1]nonan-7-ol [endo isomer]

15 [00151] A sealed vial containing 2-chloro-5-fluoro-4-(methylthio)pyrimidine (0.23 g, 1.27 mmol), (1R,5S,7s)-3-oxa-9-azabicyclo[3.3.1]nonan-7-ol hydrochloride (0.25 g, 1.39 mmol) and DIPEA (0.89 mL, 5.11 mmol) in IPA (2.5 mL) was heated at 120 °C for 22 h, further heated at 140 °C for 5 days, then concentrated *in vacuo*. Purification by flash chromatography (10 g KP-sil; 0% to 70% EtOAc in cyclohexane) afforded the title compound (0.22 g, 60%) as a white solid which was used without further purification in the subsequent step. ¹H NMR (500 MHz, CDCl₃) δ 7.89 (d, J = 1.9 Hz, 1 H), 5.64 (d, J = 12.6 Hz, 1 H), 4.69 (br s, 2 H), 4.00-3.92 (m, 3 H), 3.88-3.82 (m, 2 H), 2.50 (s, 3 H), 2.24 - 2.16 (m, 2 H), 1.87 (dd, J = 15.0, 1.1 Hz, 2 H); LCMS (Method T2); RT 1.44 min; m/z 286.2016 [M+H]⁺.

20

Step 3: (1R,5S,7s)-9-(5-fluoro-4-(methylsulfonyl)pyrimidin-2-yl)-3-oxa-9-azabicyclo[3.3.1]-nonan-7-ol [endo isomer]

25 [00152] 3-Chloroperoxybenzoic acid (0.42 g, 1.88 mmol) was added portionwise to a stirred solution of (1R,5S,7s)-9-(5-fluoro-4-(methylthio)pyrimidin-2-yl)-3-oxa-9-azabicyclo[3.3.1]-nonan-7-ol (0.22 g, 0.76 mmol) in DCM (4 mL) and MeCN (4 mL) at 0°C under Ar. The reaction mixture was stirred at 0°C for 5 min, then warmed to rt and stirred for 2 h. Reaction mixture was diluted with DCM (30 mL) and the organic mixture was washed with 1 M sodium sulfite (30 mL) and sat. aq. NaHCO₃ (20 mL). The aq. washings were combined and extracted with DCM (20 mL). The organic extracts were combined, dried (Na₂SO₄) and concentrated *in vacuo*. Purification by flash


30

chromatography (10 g KP-sil; 0% to 10% MeOH in DCM) afforded the title compound (0.25 g, 106%) as a sticky yellow solid, which was used without further purification in the subsequent steps. ¹H NMR (500 MHz, MeOD-*d*₄) δ 8.63 (d, *J* = 2.1 Hz, 1 H), 4.81–4.64 (m, 2 H), 4.00–3.96 (m, 2 H), 3.93–3.88 (m, 1 H), 3.86–3.80 (m, 2 H), 3.32 (s, 3 H), 2.29–2.15 (m, 2 H), 1.88 (dd, *J* = 14.9, 1.3 Hz, 2 H). LCMS (Method X2) RT 1.06; *m/z* 300.0919 [M-H₂O+H]⁺.

Step 4: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

[00153] To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 29.1 mg, 0.091 mmol), (1R,5S,7s)-9-(5-fluoro-4-(methylsulfonyl)pyrimidin-2-yl)-3-oxa-9-azabicyclo[3.3.1]nonan-7-ol (from step 3, 32 mg, 0.10 mmol) in 2,2,2-trifluoroethanol (0.9 mL) was added trifluoroacetic acid (7.7 μL, 0.10 mmol). The resulting mixture was heated under argon to 70°C for 16 h, then concentrated under reduced pressure. The residue was re-dissolved in DMSO (1 mL) and purified by reverse-phase chromatography (Biotage 12 g C-18 column; 10-75-85-100% methanol in water (0.1% formic acid modifier)). Fractions containing product were combined and passed through an SCX-2 (2g) column, washing with methanol (30 mL) and eluting with 2M methanolic ammonia (30 mL). The ammonia fractions were combined and evaporated under reduced pressure to give a beige solid, which was triturated with methanol / diethyl ether to give the title compound (10 mg, 0.018 mmol) as an off-white solid. ¹H NMR (600 MHz, DMSO-*d*₆) δ 9.39 (s, 1 H), 8.17 (br s, 1 H), 8.05 (d, *J* = 3.6 Hz, 1 H), 7.71 (app. d, *J* = 8.5 Hz, 1 H), 7.43 (d, *J* = 9.1 Hz, 1 H), 6.61 (br s, 1 H), 5.16 (d, *J* = 11.7 Hz, 1 H), 4.64–4.21 (m, 4 H), 3.80 (br s, 2 H), 3.70–3.62 (m, 3 H), 3.55 (s, 3 H), 3.23–3.17 (m, 1 H), 2.13–1.99 (m, 2 H), 1.61 (app. d, *J* = 14.3 Hz, 2 H), 1.35–1.27 (m, 1 H), 0.74–0.67 (m, 1 H), 0.53–0.46 (m, 2 H), 0.38–0.32 (m, 1 H); HRMS (Method X4) RT 2.72 min; *m/z* calcd for C₂₇H₃₀F₃N₆O₄⁺ [M+H]⁺: 559.2280, Found: 559.2290.


Example 6a: (S)-10-((5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



[00154] A mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 9.2 mg, 0.029 mmol) and 4,5-Dichloropyrimidine (12 mg, 0.08 mmol) and DIPEA (30 μL, 0.17 mmol) in acetonitrile (0.6 mL) in a sealed tube under argon was heated to 140°C for 18 h, then concentrated under reduced pressure. The residue was dissolved in DMSO (0.8 mL) and purified by reverse-phase


chromatography (Biotage 12 g C-18 column; 10-55-80-100% methanol in water (0.1% formic acid modifier)). The resulting material was further purified using an SCX-2 (1 g) column, washing with methanol (10 mL) and eluting with 2M methanolic ammonia (10 mL). Fractions containing product were concentrated under reduced pressure to give the title compound (2.4 mg, 0.0055 mmol) as a yellow solid. ¹H NMR (600 MHz, Methanol-*d*₄) δ 8.43 (s, 1 H), 8.37 (s, 1 H), 8.19 (d, *J* = 2.3 Hz, 1 H), 7.86 (dd, *J* = 9.1, 2.3 Hz, 1 H), 7.56 (d, *J* = 9.1 Hz, 1 H), 4.53–4.37 (m, 2 H), 3.72 (s, 3 H), 3.35–3.28 (m, 1 H), 1.44–1.37 (m, 1 H), 0.82–0.76 (m, 1 H), 0.70–0.64 (m, 1 H), 0.63–0.57 (m, 1 H), 0.40–0.34 (m, 1 H); HRMS (Method X4) RT = 2.68 min; *m/z* calcd for C₂₀H₁₉ClF₂N₅O₂⁺ [M+H]⁺: 434.1195, Found: 434.1188.

Example 7a: (2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



and;

Example 7b: (2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



Step 1: tert-butyl 3-acetyl-3,8-diazabicyclo[3.2.1]octane-8-carboxylate

[00155] Acetyl chloride (0.11 mL, 1.55 mmol) was added dropwise to a stirred solution of tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (0.3 g, 1.41 mmol) and triethylamine (0.43 mL, 3.09 mmol) in anhydrous dichloromethane (4 mL) at 0 °C under argon. The reaction mixture was stirred at rt for 6 h, then concentrated *in vacuo*. The residue was re-dissolved in EtOAc (30 mL) and washed with 1 M HCl (2 x 15 mL), saturated aq. NaHCO₃ (30 mL) and brine (30 mL). The organic layer was dried (Na₂SO₄) and concentrated *in vacuo* affording the title compound (266 mg, 74%) as a yellow oil that was used without further purification. ¹H NMR (500 MHz, CDCl₃) δ 4.36 - 4.16 (m, 3 H), 3.50 - 3.45 (m, 1 H), 3.40 (br s, 1 H), 2.85 (br s, 1 H), 2.09 (s, 3 H), 2.00 - 1.86 (m, 2 H), 1.73 - 1.60 (m, 2 H), 1.48 (s, 9H).

Step 2: 1-(3,8-diazabicyclo[3.2.1]octan-3-yl)ethan-1-one hydrochloride

[00156] 4 M HCl in 1,4-dioxane (3.50 mL, 14 mmol) was added dropwise to a solution of tert-butyl 3-acetyl-3,8-diazabicyclo[3.2.1]octane-8-carboxylate (from Step 1, 266 mg, 1.05 mmol) in DCM (5 mL) at 0 °C. The reaction mixture was allowed to warm to rt and stirred for 15 h, then concentrated *in vacuo* and dried under vacuum affording the title compound (216 mg, 108%, 1.1328 mmol) as an off-white hygroscopic solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.57 - 9.22 (m, 2 H), 4.17 (app d, *J* = 14.1 Hz, 1 H), 4.00 (br s, 2 H), 3.75-3.69 (m, 1 H), 3.56-3.50 (m, 1 H), 3.00 (app d, *J* = 14.1 Hz, 1 H), 2.03 (s, 3 H), 1.95-1.79 (m, 3 H), 1.65-1.53 (m, 1 H)

Step 3: (2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one and (2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one


[00157] A mixture of (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A1, 14.80 mg, 0.0316 mmol), 1-(3,8-diazabicyclo[3.2.1]octan-3-yl)ethan-1-one hydrochloride (from Step 2, 12.8 mg, 0.067 mmol) and DIPEA (44 μL, 0.26 mmol) in NMP under argon was heated at 140°C for 16 h. The reaction mixture was dissolved in DMSO (0.8 mL) and purified by reverse-phase chromatography (Biotage 12 g Ultra C-18 column; 10-60-80-100% methanol in water (0.1% formic acid modifier)). Fractions containing product were combined and passed through an SCX-2 (2g) column, washing with methanol (10 mL) and eluting with 2M methanolic ammonia (30 mL). The ammonia fractions were combined and evaporated under reduced pressure to give a mixture of regioisomers. This was dissolved in a 1:1 mixture of DMSO/MeCN (1 mL) and purified by HPLC (2 injections; Phenomenex Gemini C18 110A column (5 μM, 250 x 10 mm); 15min gradient of 45:55 to 30:70 H₂O:MeOH (both modified with 0.1 % formic acid); flow rate 5 mLmin⁻¹; 1260 Infinity IIMS-Prep LC). The earlier eluting major product, example 7a, was obtained as a white solid (6.2 mg, 0.011 mmol). The later eluting minor product, example 7b, presumably resulting from the acetyl group moving to the less hindered position in the starting material, was obtained as an off-white solid (2.5 mg, 0.004 mmol).

Example 7a: ¹H NMR (600 MHz, methanol-*d*₄) δ 8.06 - 8.02 (m, 1H), 8.02 - 7.99 (m, 1H), 7.95 (ddd, *J* = 9.1, 4.6, 2.3 Hz, 1H), 7.57 (d, *J* = 9.1 Hz, 1H), 4.63 - 4.56 (m, 2H), 4.53 - 4.38 (m, 2H), 4.23 - 4.16 (m, 1H), 3.72 (s, 3H), 3.67 - 3.61 (m, 1H), 3.43 (d, *J* = 12.5 Hz, 1H), 3.36 - 3.27 (m, 1H), 2.92 (d, *J* = 13.0 Hz, 1H), 2.07 (s, 3H), 2.03 - 1.92 (m, 2H), 1.83 - 1.76 (m, 1H), 1.71 - 1.65 (m, 1H), 1.46 - 1.35 (m, 1H), 0.82 - 0.75 (m, 1H), 0.69 - 0.63 (m, 1H), 0.63 - 0.57 (m, 1H), 0.38 - 0.31 (m, 1H); HRMS (Method X4) RT = 2.93 min; *m/z* calcd for C₂₈H₃₁ClF₂N₇O₃⁺ [M+H]⁺: 586.2139, Found: 586.2120.

Example 7b: ¹H NMR (600 MHz, methanol-*d*₄) δ 8.07 (dd, *J* = 7.4, 2.3 Hz, 1 H), 7.98 (d, *J* = 1.3 Hz, 1 H), 7.92 (dd, *J* = 9.1, 2.3 Hz, 1 H), 7.57 (d, *J* = 9.1 Hz, 1 H), 4.71-4.63 (m, 1 H), 4.54-4.38

(m, 2 H), 4.36–4.24 (m, 3 H), 3.73 (s, 3 H), 3.36–3.27 (m, 1 H), 3.08–2.98 (m, 2 H), 2.11 (d, J = 1.4 Hz, 3 H), 2.03–1.95 (m, 1 H), 1.89–1.81 (m, 1 H), 1.81–1.75 (m, 1 H), 1.74–1.68 (m, 1 H), 1.47–1.35 (m, 1 H), 0.83–0.76 (m, 1 H), 0.69–0.58 (m, 2 H), 0.38–0.32 (m, 1 H); HRMS (Method X4) RT = 2.97 min; m/z calcd for $C_{28}H_{31}ClF_2N_7O_3^+$ $[M+H]^+$: 586.2145, Found: 586.2132.

5 **Example 8a: (S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one**




10 **[00158]** To a sealed tube containing (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A1, 12.7 mg, 0.027 mmol) and DIPEA (20 μ L, 0.12 mmol) in DMF (0.55 mL) under argon was added dimethylamine (2 M in THF, 0.11 mL, 0.2200 mmol). The resulting mixture was heated at 140°C under microwave irradiation for 1 h, then diluted with DMSO (0.8 mL) and purified by reverse-phase chromatography (Biotage 12 g Ultra C-18 column; 10-50-70-100% methanol in water (0.1% formic acid modifier)). Fractions containing product were combined and passed

15 through an SCX-2 (2g) column, washing with methanol (10 mL) and eluting with 2M methanolic ammonia (25 mL). The ammonia fractions were combined and evaporated under reduced pressure to give the title compound (6.8 mg, 0.014 mmol) as an off-white solid. ^1H NMR (600 MHz, Methanol- d_4) δ 8.12 (d, J = 2.3 Hz, 1 H), 8.02 (dd, J = 9.1, 2.3 Hz, 1 H), 7.93 (s, 1 H), 7.53 (d, J = 9.1 Hz, 1 H), 4.53–4.34 (m, 2 H), 3.71 (s, 3 H), 3.36–3.28 (m, 1 H), 3.09 (s, 6 H), 1.43–

20 1.36 (m, 1 H), 0.83–0.76 (m, 1 H), 0.70–0.64 (m, 1 H), 0.64–0.57 (m, 1 H), 0.39–0.32 (m, 1 H); HRMS (Method X4) RT = 2.47 min; m/z calcd for $C_{22}H_{24}ClF_2N_6O_2^+$ $[M+H]^+$: 477.1617, Found: 477.1596.

25 **Example 9a: (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)**



[00159] A mixture of (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A1, 8 mg, 0.017 mmol), rac-(3R,5S)-5-fluoropiperidin-3-ol hydrochloride (8 mg, 0.051 mmol) and DIPEA (0.02 mL, 0.10 mmol) in NMP (0.68 mL) was heated under microwave irradiation at 140°C for 1h. The resulting mixture was purified by HPLC (1260 Infinity II Series Preparative HPLC (Agilent, Santa Clara, USA) using an Phenomenex Gemini C18 110A column (250 x 21.2 mm) using a 15 min gradient elution of water:methanol from 60:40 to 0:100 (modified with 0.1% formic acid) at a flow rate of 20 mLmin⁻¹), and further by SCX-2 to give the title compound (mixture of diastereoisomers) as a beige solid (5 mg, 0.009 mmol). HRMS (Method X4) RT = 2.81 min; m/z calcd for C₂₅H₂₇ClF₃N₆O₃⁺ [M+H]⁺: 551.1785, Found: 551.1785.



Example 9a-1: (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one, and Example 9a-2: (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




[00160] The preparation of example 9a was repeated on larger scale to provide a further 21 mg of this mixture of diastereoisomers, which was separated by preparative HPLC (column and conditions as above, over a 30 minute isocratic elution with 55% methanol in water (modified with 0.1% formic acid). Fractions containing only the earlier-running diastereoisomer were combined and evaporated to give example 9a-1 as a beige solid (6.5 mg). Fractions containing only the later-running diastereoisomer were combined and evaporated to give example 9a-2 as a beige solid (4.5 mg). Both compounds are *cis*-piperidine diastereoisomers, but absolute stereochemistry has not been unambiguously determined and is provisionally assigned as shown above.




[00161] Example 9a-1: ¹H NMR (600 MHz, Methanol-d₄) δ 8.05 (d, J = 2.3 Hz, 1H), 7.98 (s, 1H), 7.95 (dd, J = 9.1, 2.3 Hz, 1H), 7.56 (d, J = 9.1 Hz, 1H), 4.61 – 4.39 (m, 4H), 4.37 – 4.30 (m, 1H), 3.72 (s, 3H), 3.70 – 3.62 (m, 1H), 3.37 – 3.28 (m, 1H), 3.23 – 3.16 (m, 1H), 3.02 (dd, J = 12.9, 8.8 Hz, 1H), 2.46 – 2.34 (m, 1H), 1.70 (tt, J = 11.8, 9.2 Hz, 1H), 1.49 – 1.39 (m, 1H), 0.86 – 0.75 (m, 1H), 0.72 – 0.64 (m, 1H), 0.64 – 0.57 (m, 1H), 0.44 – 0.32 (m, 1H). LCMS (method X4) Rt 2.97 min, m/z calculated for C₂₅H₂₇ClF₃N₆O₃ [M+H]⁺: 551.1785, found 551.1779.




[00162] Example 9a-2: ¹H NMR (600 MHz, Methanol-d₄) δ 8.04 (d, J = 2.3 Hz, 1H), 7.98 (s, 1H), 7.95 (dd, J = 9.1, 2.3 Hz, 1H), 7.57 (d, J = 9.1 Hz, 1H), 4.62 – 4.39 (m, 4H), 4.38 – 4.34 (m, 1H), 3.73 (s, 3H), 3.70 – 3.63 (m, 1H), 3.38 – 3.28 (m, 1H), 3.22 – 3.11 (m, 1H), 2.97 (dd, J = 12.9, 9.0 Hz, 1H), 2.46 – 2.36 (m, 1H), 1.69 (tt, J = 11.5, 9.3 Hz, 1H), 1.49 – 1.39 (m, 1H), 0.86 – 0.75 (m, 1H), 0.73 – 0.65 (m, 1H), 0.65 – 0.57 (m, 1H), 0.41 – 0.31 (m, 1H). LCMS (method X4) Rt 2.99 min, m/z calculated for C₂₅H₂₇ClF₃N₆O₃ [M+H]⁺: 551.1785, found 551.1785.






[00163] The following tabulated examples were prepared by a method analogous to that used for the preparation of Example 9a, using the appropriate amine (as free base or as hydrochloride salt). Amines were obtained from commercial vendors, or prepared by common transformations from commercially available intermediates, as described in the table. For examples 9j – 9n, example 9s, and examples 9y, 9za and 9zb, extended heating times of 3 – 8 h were required. Where racemic amines were used, two diastereoisomers were formed, as shown.





Example	Data and comments	Amine used
Example 9b: (S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.14, m/z = 574.3 HRMS (Method X4): m/z calcd for C ₂₈ H ₃₅ ClF ₂ N ₇ O ₂ [M+H] ⁺ : 574.2509, Found: 574.2510 (Rt = 2.34 min)	N,N-dimethyl-1-[(3R)-3-piperidyl]methanamine dihydrochloride
Example 9c: (3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide: (3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide (1:1) 	LCMS (Method X2): Rt = 1.41, m/z = 602.2 HRMS (Method X4): m/z calcd for C ₂₉ H ₃₅ ClF ₂ N ₇ O ₃ [M+H] ⁺ : 602.2458, Found: 602.2458 (Rt = 2.97 min)	<i>rac</i> -(3R,5S)-N-ethyl-5-methylpiperidine-3-carboxamide [prepared from <i>rac</i> -(3R,5S)-1-(tert-butoxycarbonyl)-5-methylpiperidine-3-carboxylic acid by HATU coupling followed by boc-deprotection]
Example 9d: (S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-	LCMS (Method X2): Rt = 1.41, m/z = 561.2 HRMS (Method X4): m/z calcd for C ₂₇ H ₃₂ ClF ₂ N ₆ O ₃ [M+H] ⁺ : 561.2192,	Rac-cis-[(3R,5S)-5-methyl-3-piperidyl]-methanol [prepared from reduction of <i>rac</i> -cis-(3R,5S)-5-methylpiperidine-3-carboxylic acid by borane






cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1) 	Found: 561.2177 (Rt = 2.88 min)	tetrahydrofuran complex solution followed by boc-deprotection]
Example 9e: (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one and (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1). 	LCMS (Method X2): Rt = 1.42, m/z = 569.2 HRMS (Method X4): m/z calcd for C₂₅H₂₆ClF₄N₆O₃ [M+H]⁺: 569.1691, Found: 569.1684 (Rt = 2.99 min)	4,4-difluoropiperidin-3-ol hydrochloride
Example 9f: (S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one and (S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1) 	LCMS (Method X2): Rt = 1.48, m/z = 549.2 HRMS (Method X4): m/z calcd for C₂₆H₂₉ClF₃N₆O₂ [M+H]⁺: 549.1992, Found: 549.1992 (Rt = 3.14 min)	3-(fluoromethyl)piperidine hydrochloride
Example 9g: 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-	LCMS (Method X2): Rt = 1.43, m/z = 610.2	2-(4,4-difluoropiperidin-3-yl)acetamide




1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide and 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide (1:1) 	HRMS (Method X4): m/z calcd for $C_{27}H_{29}ClF_4N_7O_3$ [M+H]⁺: 610.1956, Found: 610.1957 (Rt = 3.06 min)	[prepared from 2-(1-tert-butoxycarbonyl-4,4-difluoro-3-piperidyl)acetic acid by amide coupling and boc deprotection]
Example 9h: (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.15, m/z = 560.2 HRMS (Method X4): m/z calcd for $C_{27}H_{33}ClF_2N_7O_2$ [M+H]⁺: 560.2352, Found: 560.2356 (Rt = 2.36 min)	(3S)-N,N-dimethylpiperidin-3-amine hydrochloride
Example 9i: (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.16, m/z = 560.2 HRMS (Method X4): m/z calcd for $C_{27}H_{33}ClF_2N_7O_2$ [M+H]⁺: 560.2352, Found: 560.2338 (Rt = 2.42 min)	(3R)-N,N-dimethylpiperidin-3-amine hydrochloride
Example 9j: (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide and (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-	LCMS (Method X2): Rt = 1.41, m/z = 624.2 HRMS (Method X4): m/z calcd for $C_{28}H_{31}ClF_4N_7O_3$ [M+H]⁺: 624.2113, Found: 624.2101 (Rt = 2.97 min)	N-ethyl-4,4-difluoropiperidine-3-carboxamide [prepared by HATU coupling followed by benzyl deprotection of 1-benzyl-4,4-difluoro-piperidine-3-carboxylic acid]

difluoropiperidine-3-carboxamide (1:1) 		
Example 9k: N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide and N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide (1:1) 	LCMS (Method X2): Rt = 1.50, m/z = 652.2 HRMS (Method X4): m/z calcd for C₃₀H₃₅ClF₄N₇O₃ [M+H]⁺: 652.2426, Found: 652.2422 (Rt = 3.23 min)	N-((4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide [prepared by mesylation of (1-benzyl-4,4-difluoropiperidin-3-yl)methanol, followed by displacement of mesylate with ethylamine, acylation and benzyl deprotection]
Example 9l: (S)-10-((5-chloro-2-((1R,4R)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((1S,4S)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:formic acid (1:1:0.5) 	LCMS (Method X2): Rt = 1.47, m/z = 565.2 HRMS (Method X4): m/z calcd for C₂₆H₂₆ClF₄N₆O₂ [M+H]⁺: 565.1742, Found: 565.1745 (Rt = 3.12 min)	5,5-difluoro-2-azabicyclo[2.2.1]heptane hydrochloride
Example 9m: N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-	LCMS (Method X2): Rt = 1.45, m/z = 624.2 HRMS (Method X4): m/z calcd for C₂₈H₃₁ClF₄N₇O₃ [M+H]⁺: 624.2113, Found: 624.2104 (Rt = 3.08 min)	(R)-N-(4,4-difluoropiperidin-3-yl)-N-methylacetamide [prepared by sequential acylation, alkylation and boc-deprotection of tert-butyl (R)-3-amino-4,4-difluoropiperidine-1-carboxylate]

difluoropiperidin-3-yl)-N-methylacetamide 		
Example 9n: N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide 	LCMS (Method X2): Rt = 1.46, m/z = 624.2 HRMS (Method X4): m/z calcd for $C_{28}H_{31}ClF_4N_7O_3$ [M+H]⁺: 624.2113, Found: 624.2112 (Rt = 3.11 min)	(S)-N-(4,4-difluoropiperidin-3-yl)-N-methylacetamide [prepared by sequential acylation, alkylation and boc-deprotection of tert-butyl (S)-3-amino-4,4-difluoropiperidine-1-carboxylate]
Example 9o: (S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.33, m/z = 565.2 HRMS (Method X4): m/z calcd for $C_{26}H_{29}ClF_3N_6O_3$ [M+H]⁺: 565.1942, Found: 565.1929 (Rt = 2.78 min)	(4-fluoropiperidin-4-yl)methanol hydrochloride
Example 9p: (S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method T2): Rt = 1.38, m/z = 547.2 HRMS (Method X4): m/z calcd for $C_{26}H_{30}ClF_2N_6O_3$ [M+H]⁺: 547.2036, Found: 547.2032 (Rt = 2.81 min)	(R)-piperidin-3-ylmethanol
Example 9q: (S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method T2): Rt = 1.38, m/z = 547.2 HRMS (Method X4): m/z calcd for	(S)-piperidin-3-ylmethanol








	$C_{26}H_{30}ClF_2N_6O_3$ $[M+H]^+$: 547.2036, Found: 547.2027 (R_t = 2.80 min)	
Example 9r: (S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1) 	LCMS (Method X2): R_t = 1.57, m/z = 586.2 HRMS (Method X4): m/z calcd for $C_{28}H_{31}ClF_2N_7O_3$ $[M+H]^+$: 586.2145, Found: 586.2133 (R_t = 3.37 min)	Rac-3-ethyl-3,8-diazabicyclo[3.2.1]octan-4-one [prepared from tert-butyl 2-oxo-3,8-diazabicyclo[3.2.1]octane-8-carboxylate by ethylation and boc-deprotection]
Example 9s: (S)-10-((5-chloro-2-((3S,5R)-3,5-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): R_t = 1.45, m/z = 553.2 HRMS (Method X4): m/z calcd for $C_{25}H_{26}ClF_4N_6O_2$ $[M+H]^+$: 553.1742, Found: 553.1741 (R_t = 3.05 min)	(3S,5R)-3,5-difluoropiperidine hydrochloride
Example 9t: (S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): R_t = 1.34, m/z = 598.2 HRMS (Method X4): m/z calcd for $C_{29}H_{31}ClF_2N_7O_3$ $[M+H]^+$: 598.2145, Found: 598.2135 (R_t = 2.77 min)	cyclopropyl(2,6-diazaspiro[3.3]heptan-2-yl)methanone [prepared from tert-butyl 2,6-diazaspiro[3.3]heptane-2-carboxylate by HATU coupling with cyclopropanecarboxylic acid followed by Boc deprotection]
Example 9u: (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-	LCMS (Method X2): R_t = 1.29, m/z = 535.2	2-methyl-1-(methylamino)propan-2-ol

1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	HRMS (Method X4): m/z calcd for $C_{25}H_{30}ClF_2N_6O_3$ [M+H]⁺: 535.2036, Found: 535.2054 (Rt = 2.65 min)	
Example 9v: (S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.28, m/z = 549.2 HRMS (Method X4): m/z calcd for $C_{26}H_{32}ClF_2N_6O_3$ [M+H]⁺: 549.2192, Found: 549.2175 (Rt = 2.63 min)	2-methyl-4-(methylamino)butan-2-ol
Example 9w: (S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.16, m/z = 584.2 HRMS (Method T4): m/z calcd for $C_{29}H_{33}ClF_2N_7O_2$ [M+H]⁺: 584.2347, Found: 584.2343 (Rt = 2.32 min)	2-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptane (made from tert-butyl 2,6-diazaspiro[3.3]heptane-2-carboxylate by alkylation and boc-deprotection)
Example 9x: (2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method X2): Rt = 1.18, m/z = 620.2 HRMS (Method T4): m/z calcd for $C_{29}H_{31}ClF_4N_7O_2$ [M+H]⁺: 620.2158, Found: 620.2153 (Rt = 2.39 min)	2-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptane [made from tert-butyl 2,6-diazaspiro[3.3]heptane-2-carboxylate by alkylation and boc-deprotection]
Example 9y: (2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c] 	LCMS (Method T2): Rt = 1.32, m/z = 598.3 HRMS (Method T4): m/z calcd for	8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octane [made from tert-butyl 3,8-


c]quinolin-6(7H)-one 	$C_{30}H_{35}ClF_2N_7O_2$ $[M+H]^+$: 598.2503, Found: 598.2499 (R_t = 2.50 min)	diazabicyclo[3.2.1]octane-3-carboxylate by alkylation and boc-deprotection]
Example 9za: (2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method T2): R_t = 1.34, m/z = 634.2 HRMS (Method X4): m/z calcd for $C_{30}H_{33}ClF_4N_7O_2$ $[M+H]^+$: 634.2321, Found: 634.2312 (R_t = 2.59 min)	8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octane [made from tert-butyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylate by alkylation and boc-deprotection]
Example 9zb: (S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	LCMS (Method T4): R_t = 2.64 min; m/z calcd for $C_{26}H_{30}ClF_2N_6O_3$ $[M+H]^+$ 547.2030; found 547.2024	(3R,5R)-5-methylpiperidin-3-ol [prepared by analogy to the procedure described in Pan <i>et al</i> , Bioorganic & Medicinal Chemistry Letters, 24(8), 1983-1986; 2014.]

[00164] The following tabulated examples were prepared by a method analogous to that used for the preparation of **Example 9a**. 1-1.5 eq of the appropriate amine (as free base or as hydrochloride salt) was used. Amines used were racemic, leading to a mixture of isomers as shown. Potassium carbonate (3 - 4 eq) was used in place of DIPEA, and acetonitrile was used as solvent. Heating times were 4 - 5h at 140 - 150 °C in the microwave. Example 10d used (S)-10-((2-chloro-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A2), and was heated to 140 °C for 16 hours.

Example 10a: (S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-	HRMS (Method X4) m/z calc for $C_{25}H_{26}ClF_4N_6O_2$ $[M+H]^+$ = 553.1742, found
---	--

Example 10a: (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:  	553.1734. RT 3.28 mins
Example 10b: (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:  	HRMS (Method T4) m/z calc for C₂₅H₂₆ClF₄N₆O₂ [M+H]⁺ = 553.1736, found 553.1751. RT 2.96 mins
Example 10c: (S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: 	HRMS (Method X4) m/z calc for C₂₆H₃₀ClF₂N₆O₃ [M+H]⁺ = 547.2036, found 547.2032. RT 3.11 mins
Example 10d: (S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:  	HRMS (Method X4) m/z calc for C₂₅H₂₆F₅N₆O₂ [M+H]⁺ = 537.2037, found 537.2032. RT = 3.18 mins





Example 11a: (2S)-10-[(5-chloro-2-(4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino]-2-cyclopropyl-3,3-difluoro-7-methyl-2,4-dihydro-1H-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one















[00165] 1-methyl-1,4-diazepane (5 mg, 0.044 mmol), (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate A1, 12 mg, 0.026 mmol) and DIPEA (22 μ L, 0.13 mmol) were combined in acetonitrile (0.5 mL) and heated in a sealed vessel at 70°C for 2 days. The resulting mixture was evaporated, dissolved in DMSO and purified using reverse phase flash chromatography (Biotage 12g SNAP Ultra C18, 20-35-75-100% methanol in water, 0.1% formic acid modifier). Fractions containing product were loaded onto an SCX-2 (1g) column which was washed with methanol and eluted with 20% (1.6M methanolic ammonia) in DCM to give the title compound (10.3 mg, 0.017 mmol) as a yellow solid after evaporation and drying. HRMS (Method X4) RT = 2.38 min; m/z calcd for $C_{26}H_{31}ClF_2N_7O_2^+$ [M+H]⁺: 546.2190, found 546.2186







[00166] The following tabulated examples were prepared by a method analogous to that used for the preparation of **Example 11a**. Amines were obtained from commercial vendors and used as free base or hydrochloride salt. Some amines were used as racemates, leading to a mixture of isomers as shown. For example 12a and 12d, the reactions were heated to 140 °C in the microwave. For reactions forming examples 12b and 12c, (3S,5R)-3-(Boc-amino)-5-methylpiperidine or (3R,5S)-3-(Boc-amino)-5-methylpiperidine were used. Boc groups were removed by treatment with TFA in DCM at room temperature to give the products shown. Example 12e was further heated to 160 °C in the microwave for 4 hours.







Example 11b: (S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 3.05 mins m/z calc for $C_{27}H_{32}ClF_2N_8O_2$ [M+H]⁺ = 545.2238, found 545.2233.
Example 11c: (S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-		LCMS (Method T4) RT 2.91 mins m/z calc for $C_{27}H_{32}ClF_2N_8O_2$ [M+H]⁺ = 545.2238, found 545.2240.

yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		
Example 11d: (S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1))		LCMS (Method T4) RT 2.94 mins m/z calc for $C_{27}H_{32}ClF_2N_6O_2$ [M+H] ⁺ = 545.2238, found 545.2248.
Example 11e: (S)-10-((5-chloro-2-((3R,5S)-3,5-dihydropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		LCMS (Method T4) RT 3.05 mins m/z calc for $C_{25}H_{28}ClF_2N_6O_4$ [M+H] ⁺ = 549.1823, found 549.1827.
Example 11f: (S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.66 mins m/z calc for $C_{27}H_{30}ClF_2N_6O_3$ [M+H] ⁺ = 559.2030, found 559.2026.
Example 11g: (S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.72 mins m/z calc for $C_{25}H_{27}ClF_3N_6O_3$ [M+H] ⁺ = 551.1780, found 551.1785.


Example 11h: (S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.73 mins m/z calc for $C_{27}H_{30}ClF_2N_6O_3$ [M+H]⁺ = 559.2030, found 559.2023.
Example 11i: (S)-10-((5-chloro-2-(methyl(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		LCMS (Method T4) RT 2.68 mins m/z calc for $C_{24}H_{28}ClF_2N_6O_2$ [M+H]⁺ = 505.1925, found 505.1929.
Example 11j: (S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one : (S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.55 mins m/z calc for $C_{26}H_{30}ClF_2N_6O_3$ [M+H]⁺ = 547.2030, found 547.2033.
Example 11k: (S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		LCMS (Method T4) RT 2.60 mins m/z calc for $C_{25}H_{30}ClF_2N_6O_3$ [M+H]⁺ = 535.2030, found 535.2035.
Example 11l: (S)-10-((5-chloro-2-(5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		LCMS (Method T4) RT 2.95 mins m/z calc for $C_{27}H_{30}ClF_2N_6O_2$ [M+H]⁺ = 543.2081, found 543.2086.
Example 11m: (S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one : (S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.53 mins m/z calc for $C_{25}H_{30}ClF_2N_6O_3$ [M+H]⁺ = 535.2030, found 535.2033.

Example 11n: (S)-10-((5-chloro-2-(3,3-dimethylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method T4) m/z calc for $C_{25}H_{28}ClF_2N_6O_2$ [M+H]⁺ =: 517.1925, Found: 517.1923
Example 11o: (S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method T4) m/z calc for $C_{24}H_{26}ClF_2N_6O_2$ [M+H]⁺ = 503.1769, Found: 503.1768
Example 11p: (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile		HRMS (Method X4) m/z calcd for $C_{24}H_{23}ClF_2N_7O_2$ [M+H]⁺: 514.1570, Found: 514.1570
Example 11q: (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method X4) m/z calcd for $C_{23}H_{23}ClF_3N_6O_2$ [M+H]⁺: 507.1518, Found: 507.1519
¹H NMR (600 MHz, Methanol-d₄) δ 8.16 (d, J = 2.3 Hz, 1H), 8.00 (dd, J = 9.1, 2.3 Hz, 1H), 7.97 (s, 1H), 7.54 (d, J = 9.2 Hz, 1H), 5.39 (dtt, J = 57.2, 6.1, 3.2 Hz, 1H), 4.54 – 4.29 (m, 4H), 4.16 – 4.05 (m, 2H), 3.72 (s, 3H), 3.42 – 3.25 (m, 1H), 1.48 – 1.36 (m, 1H), 0.85 – 0.77 (m, 1H), 0.74 – 0.66 (m, 1H), 0.66 – 0.58 (m, 1H), 0.45 – 0.32 (m, 1H).		
Example 11r: (S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		HRMS (Method T4) RT 3.05 mins m/z calc for $C_{26}H_{28}ClF_4N_6O_2$ [M+H]⁺ = 567.1893, found 567.1900.
Example 11s: (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile		HRMS (Method T4) m/z calcd for $C_{25}H_{25}ClF_2N_7O_2$ [M+H]⁺: 528.1721, Found: 528.1721



Example 11t: (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile		HRMS (Method T4) m/z calcd for C₂₅H₂₅ClF₂N₇O₂ [M+H]⁺: 528.1721, Found: 528.1726
Example 11u: (S)-10-((5-chloro-2-(3-(difluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method T4) m/z calcd for C₂₄H₂₄ClF₄N₆O₂ [M+H]⁺: 539.1580, Found: 539.1585
Example 11v: (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	 ¹H NMR (600 MHz, MeOD-<i>d</i>₄) δ 8.17 (d, <i>J</i> = 2.3 Hz, 1H), 8.02 (dd, <i>J</i> = 9.1, 2.3 Hz, 1H), 7.90 (s, 1H), 7.50 (d, <i>J</i> = 9.1 Hz, 1H), 4.54 - 4.34 (m, 2H), 4.10-4.03 (m, 4H), 3.69 (s, 3H), 3.37-3.28 (m, 1H), 2.32 (quin, <i>J</i> = 7.5 Hz, 2H), 1.42-1.34 (m, 1H), 0.83 - 0.77 (m, 1H), 0.71 - 0.64 (m, 1H), 0.61 - 0.58 (m, 1H), 0.40 - 0.33 (m, 1H).	HRMS (Method T4) m/z calcd for C₂₃H₂₄ClF₂N₆O₂⁺ [M+H]⁺: 489.1617, Found: 489.1602.
Example 11w: (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	 ¹H NMR (600 MHz, MeOD-<i>d</i>₄) δ 8.18 (d, <i>J</i> = 2.3 Hz, 1H), 8.00 (dd, <i>J</i> = 9.1, 2.3 Hz, 1H), 7.93 (s, 1H), 7.52 (d, <i>J</i> = 9.1 Hz, 1H), 4.52 - 4.37 (m, 2H), 4.31 - 4.26 (m, 1H), 4.25 - 4.19 (m, 2H), 3.90 - 3.86 (m, 2H), 3.71 (s, 3H), 3.36 - 3.28 (m, 4H), 1.43 - 1.35 (m, 1H), 0.83 - 0.76 (m, 1H), 0.72 - 0.66 (m, 1H), 0.64 - 0.59 (m, 1H), 0.40 - 0.34 (m, 1H)	HRMS (Method T4) m/z calcd for C₂₄H₂₆ClF₂N₆O₃⁺ [M+H]⁺: 519.1723, Found: 519.1708.
Example 11x: (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method T4) m/z calcd for C₂₅H₂₈ClF₂N₆O₃ [M+H]⁺: 533.1874, Found: 533.1884
Example 11y: (S)-10-((5-chloro-2-((R)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((S)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-		LCMS (Method X2) RT 1.63min; m/z calcd for C₂₈H₃₀ClF₂N₆O₂⁺ 531.2081 found 531.2085






[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		
Example 11z: (S)-10-((5-chloro-2-((R)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method X4) RT 3.29min; m/z calcd for $C_{26}H_{29}ClF_3N_6O_2+$ [M+H]⁺ 549.1992 found 549.1983
Example 12a: (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.49 mins m/z calc for $C_{25}H_{28}ClF_2N_6O_3$ [M+H]⁺ = 533.1874, found 533.1873
Example 12b: (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method X4) m/z calcd for $C_{26}H_{31}ClF_2N_7O_2$ [M+H]⁺: 546.2195, Found: 546.2202.
Example 12c: (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		HRMS (Method X4) m/z calcd for $C_{26}H_{31}ClF_2N_7O_2$ [M+H]⁺: 546.2195, Found: 546.2182.
Example 12d: (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one		LCMS (Method T4) RT 2.62 mins m/z calcd for $C_{25}H_{26}ClF_2N_6O_3$ ([M+H]⁺) 531.1717; found 531.1679
Example 12e: (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one:(S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-		LCMS (Method X4) RT 3.20min; m/z calcd for $C_{25}H_{26}ClF_4N_6O_3+$ [M+H]⁺ 569.1691 found 569.1688





yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		
Example 12f: (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method X4) RT 3.63min m/z calcd for C ₂₅ H ₂₈ ClF ₂ N ₈ O ₂ S + [M+H] ⁺ 549.1646 found 549.1635
Example 12g: (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.50 min m/z calcd for C ₂₇ H ₃₃ ClF ₂ N ₇ O ₂ [M+H] ⁺ 560.2347 found 560.2351
Example 12h: (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one: (S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) RT 2.49min; m/z calcd for C ₂₇ H ₃₁ ClF ₂ N ₇ O ₂ [M+H] ⁺ 558.2190 found 558.2188
Example 12i: (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one : (S)-10-((5-chloro-2-((s)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method T4) Rt 2.81min; m/z calcd for C ₂₇ H ₂₉ ClF ₂ N ₇ O ₃ [M+H] ⁺ 572.1983 found 572.1994





Example 12j: (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one : (S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (1:1)		LCMS (Method X4) Rt 3.20min; m/z calcd for C ₂₈ H ₃₂ ClF ₂ N ₆ O ₃ [M+H] ⁺ 573.2192 found 573.2183
--	--	---





[00167] The following examples 12x01-12x50 may be prepared by methods analogous to those previously described for examples 9a, 11a or 12a. Amines may be obtained from commercial vendors or prepared by known methods, and used as free base or hydrochloride salt. Some amines were used as racemates, leading to a mixture of isomers as shown. Protecting groups may be used to mask reactive functionality; for example in the preparation of example 12x14 tert-butyl 6-hydroxy-1,4-diazepane-1-carboxylate was used, and the Boc group was removed by treatment with TFA in DCM at room temperature. In the preparation of example 12x26, tert-butyl 2-(difluoromethyl)piperazine-1-carboxylate was used, and Boc deprotection occurred during the reaction.





Example 12x01: (S)-10-((5-chloro-2-((S)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (Method X4) rt 2.75min; m/z calculated for C ₂₅ H ₂₈ ClF ₂ N ₆ O ₂ [M+H] ⁺ 517.1925 found 517.1929
Example 12x02: (S)-10-((5-chloro-2-((R)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (Method T4) rt 2.70min; m/z calculated for C ₂₅ H ₂₈ ClF ₂ N ₆ O ₂ [M+H] ⁺ 517.1925 found 517.1894
Example 12x03: (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared using a method analogous to 11a, heated to 120 °C for 18h. LCMS (Method X4) rt 2.84 min; m/z calculated





	for $C_{26}H_{29}ClF_3N_6O_2$ $[M+H]^+$ 549.1987; found 549.1973
Example 12x04: (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x05: (S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared [(as 1:1 mixture with (3S,5R) isomer)] from rac-(3S,5R)-3-methoxy-5-methylpiperidine:TFA using a method analogous to 11a LCMS (Method T4) Rt 3.03min; m/z calculated for $C_{27}H_{32}ClF_2N_6O_3$ $[M+H]^+$ 561.2187 found 561.2188
Example 12x06: (2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as mixture of diastereoisomers on thiomorpholine ring using a method analogous to 11a. LCMS (method X4) 1:1 ratio of peaks with rt 3.54min, m/z 563.1790 and rt 3.62min, m/z 563.1786. Calculated m/z 563.1807 for $C_{26}H_{30}ClF_2N_6O_2S$ $[M+H]^+$
Example 12x07: (2S)-10-((5-chloro-2-(3,4-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as mixture of diastereoisomers on piperidine ring using a method analogous to 11a. LCMS (method X4) 2:1 ratio of rt 3.34 min, m/z 545.2227 and rt 3.42min, m/z 545.2229. Calculated m/z 545.2243 for $C_{27}H_{32}ClF_2N_6O_2$ $[M+H]^+$





Example 12x08: (2S)-10-((5-chloro-2-(2,3-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as mixture of diastereoisomers on piperidine ring using a method analogous to 11a, heated additionally to 160 °C for 2h in the microwave. LCMS (method X4) rt 3.27min, 93% purity; m/z calculated 545.2243 for C₂₇H₃₂ClF₂N₆O₂ [M+H]⁺, found 545.2237
Example 12x09: (2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiophosphorin-4-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as mixture of diastereoisomers on thiomorpholine ring using a method analogous to 11a, heated additionally to 160 °C for 5h in the microwave. LCMS (method T4) rt 2.85min; m/z calculated 579.1751 for C₂₈H₃₀ClF₂N₆O₃S⁺ [M+H]⁺, found 579.1752
Example 12x10: (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of (2R,5S) and (2S,5R) piperidine diastereoisomers using a method analogous to 11a, additionally heated in a 1:1 mixture of acetonitrile and NMP at 160 °C for 16h in the microwave. LCMS (method X4) rt 2.86min; m/z calculated 561.2192 for C₂₇H₃₂ClF₂N₆O₃ [M+H]⁺, found 561.2199
Example 12x11: (2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxothiophosphorin-4-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of diastereoisomers on thiomorpholine ring, using a method analogous to 11a, with additional heating in a 1:1 mixture of acetonitrile and NMP at 160 °C for 4h in the microwave. Major diastereoisomer (~80%) is <i>cis</i>. LCMS (Method T4) Rt 2.93min; m/z calculated 595.1700 for C₂₈H₃₀ClF₂N₆O₄S⁺ [M+H]⁺, found 595.1700
Example 12x12: (2S)-10-((5-chloro-2-(2,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared as a 1:1 mixture of (2R,5R) and (2S,5S) piperidine diastereoisomers using a method analogous to 9a, heated in a 1:1 mixture of





	acetonitrile and NMP at 160 °C for 8h in the microwave. LCMS (method X4) rt 3.20min; m/z calculated 545.2243 for C₂₇H₃₂ClF₂N₆O₂ [M+H]⁺, found 545.2235
Example 12x13: (S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as formic acid salt using a method analogous to 11a, heated to 140 °C for 22h. LCMS (Method X4) RT = 2.66 min; m/z calc 547.2036 for C₂₆H₃₀ClF₂N₆O₃ [M+H]⁺; found 547.2020
Example 12x14: (2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as 1:1 mixture of R and S hydroxy epimers using a method analogous to 11a. LCMS (method T4) rt 2.45min, m/z calculated 548.1983 for C₂₅H₂₉ClF₂N₇O₃ [M+H]⁺, found 548.1987
Example 12x15: (2S)-10-((5-chloro-2-(6-hydroxy-4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Trideuteromethyl version was prepared – see below
Example 12x15-d3: (2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-d₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a 1:1 mixture of R and S hydroxy epimers using a method analogous to 11a. LCMS (method T4) Rt 2.22min; m/z calculated 565.2328 for C₂₆H₂₈D₃ClF₂N₇O₃ [M+H]⁺ found 565.2330
Example 12x16: (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared using a method analogous to 9a. Obtained two peaks following HPLC purification containing different piperidine diastereoisomers.






	Example 12x16a: LCMS (method X4) rt 2.96min; m/z calculated 628.2614 for $C_{31}H_{37}ClF_2N_7O_3$ $[M+H]^+$, found 628.2601. Example 12x16b: LCMS (method X4) rt 3.06 and 3.09, approx. 1:1 ratio, m/z found 628.2614.
Example 12x17: (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 9. Obtained two peaks following HPLC purification containing different piperidine diastereoisomers Example 12x17a: LCMS (method X4) rt 3.07min; m/z calculated 664.2426 for $C_{31}H_{35}ClF_4N_7O_3$ $[M+H]^+$, found 664.2413. Example 12x17b: LCMS (method X4) rt 3.18 and 3.20, approx. 1:1 ratio, m/z found 664.2410.
Example 12x18: (2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of epimers at the piperidine 3-position using a method analogous to 9a, with additional heating in the microwave at 160 °C for 8h LCMS (method X4) rt 2.99min; m/z calculated 636.2113 for $C_{29}H_{31}ClF_4N_7O_3$ $[M+H]^+$, found 636.2112
Example 12x19: (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of epimers at the piperidine 3-position using a method analogous to 9a, heating in the microwave at 140 °C for 12h LCMS (method T4) rt 2.76min; m/z calculated 672.1919 for $C_{29}H_{29}ClF_6N_7O_3$ $[M+H]^+$, found 672.1930
Example 12x20: (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-	Prepared as a mixture of epimers at the piperidine 3-






yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	position using a method analogous to 9a, heating in the microwave at 140 °C for 4h LCMS (method X4) rt 3.42min; m/z calculated 686.2081 for C₃₀H₃₁ClF₆N₇O₃, found 686.2072
Example 12x21: (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of epimers at the piperidine 3-position using a method analogous to 9a, heating in the microwave at 140 °C for 4h LCMS (method X4) rt 3.39min; m/z calculated 650.2269 for C₃₀H₃₃ClF₄N₇O₃, found 650.2280
Example 12x22: (2S)-10-((5-chloro-2-(3-ethynylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a 1:1 mixture of (3R) and (3S)-ethynyl epimers using a method analogous to 11a. LCMS (method T4) rt 2.93min; m/z calculated 541.1925 for C₂₇H₂₈ClF₂N₆O₂, found 541.1919
Example 12x23: 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile 	Prepared using a method analogous to 9a. LCMS (method T4) rt 2.67min; m/z calculated 542.1877 for C₂₆H₂₆ClF₂N₇O₂ [M+H]⁺, found 542.1875
Example 12x24: (S)-10-((5-chloro-2-((4,4-difluorocyclohexyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared using a method analogous to 11a, heated additionally to 160 °C for 4h in the microwave, then for a





	further 4h at 140 °C in presence of >10eq amine. LCMS (method X4) rt 2.72min; m/z calculated 567.1898 for C₂₆H₂₈ClF₄N₆O₂ [M+H]⁺, found 567.1898
Example 12x25: (2S)-10-((2-(2-azabicyclo[4.1.0]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of diastereoisomers on fused piperidine ring, using a method analogous to 11a. LCMS (method T4) rt 2.64min; m/z calculated 529.1925 for C₂₆H₂₈ClF₂N₆O₂ [M+H]⁺, found 529.1926
Example 12x26: (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of (3R) and (3S)-difluoromethyl epimers using a method analogous to 11a, additionally heated in a 1:1 mixture of acetonitrile and NMP at 120 °C for 1h then 160 °C for 4h in the microwave. LCMS (method T4) rt 2.49min; m/z calculated 568.1845 for C₂₅H₂₇ClF₄N₇O₂ [M+H]⁺, found 568.1849
Example 12x27: (2S)-10-((5-chloro-2-(3-methyl-5-oxopiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x28: (S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared as a mixture of piperidine diastereoisomers using a method analogous to 9a, LCMS (method X4) rt 2.29min; m/z expected 618.2771 for C₃₀H₃₉ClF₂N₇O₃ [M+H]⁺, found 618.2766





	
Example 12x29: (S)-10-((5-chloro-2-((3R,5S)-3-(3-(dimethylamino)propoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x30: (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of (3S,5R) and (3R,5S)-piperidine diastereoisomers using a method analogous to 9a LCMS (method X4) rt 2.91min; m/z calculated 614.2458 for C₃₀H₃₅ClF₂N₇O₃ [M+H]⁺, found 614.2452
Example 12x31: (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of (3S,5R) and (3R,5S)-piperidine diastereoisomers using a method analogous to 9a, heating in the microwave at 160 °C for 1h LCMS (method X4) rt 3.05min; m/z expected 650.2269 for C₃₀H₃₃ClF₄N₇O₃ [M+H]⁺, found 650.2259
Example 12x32: (S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared using a method analogous to 11a, heated to 80 °C for 90 mins. LCMS (method X4) rt 2.31 min; m/z calculated 519.1723 for

	$C_{24}H_{26}ClF_2N_6O_3$ $[M+H]^+$, found 519.1715
Example 12x33: (2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared as single diastereoisomers, examples 12x33a and 12x33b, see below.
Example 12x33a: (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a, heated to 80 °C for 90 mins. LCMS (method X4) rt 2.50min; m/z calculated 519.1723 for $C_{24}H_{26}ClF_2N_6O_3$ $[M+H]^+$, found 519.1723
Example 12x33b: (S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a, heated to 80 °C for 90 mins. LCMS (method T4) rt 2.17min; m/z calculated 519.1723 for $C_{24}H_{26}ClF_2N_6O_3$ $[M+H]^+$, found 519.1727
Example 12x34: (2S)-10-((5-chloro-2-(2-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x35: (S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared using a method analogous to 11a, heated to 80 °C for 90 mins. LCMS (method T4) rt 2.68min; m/z calculated 521.1680 for


	$C_{24}H_{25}ClF_3N_6O_2$ $[M+H]^+$, found 521.1668
Example 12x36: (2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as mixture of diastereoisomers using a method analogous to 11a, heated to 80 °C for 90 mins. LCMS (method T4) rt 2.63min; m/z calculated 503.1774 for $C_{24}H_{26}ClF_2N_6O_2$ $[M+H]^+$, found 503.1772
Example 12x37: (2S)-10-((5-chloro-2-(4,4-difluoro-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x38: (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile 	Prepared using a method analogous to 11a LCMS (method T4) rt 2.79min; m/z calculated 542.1877 for $C_{26}H_{27}ClF_2N_7O_2$ $[M+H]^+$, found 542.1875
Example 12x39: 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide 	Prepared as a mixture of piperidine diastereoisomers using a method analogous to 9a, heating in the microwave at 160 °C for 1h LCMS (method X4) rt 2.94min; m/z calculated 606.2207 for $C_{28}H_{32}ClF_3N_7O_3$ $[M+H]^+$, found 606.2200

	
Example 12x40: (2S)-10-((5-chloro-2-(3-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of (3S)- and (3R)-fluoropiperidine epimers using a method analogous to 9a, heating in the microwave at 140 °C for 4h LCMS (method X4) rt 3.06 and 3.11 min; m/z calculated 535.1836 for $C_{25}H_{27}ClF_3N_6O_2$ [M+H]⁺, found 535.1836 and 535.1837
Example 12x41: (2S)-10-((5-chloro-2-(4,4-difluoro-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	
Example 12x42: (S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (method X4) rt 3.12min; m/z calculated 535.1836 for $C_{25}H_{27}ClF_3N_6O_2$ [M+H]⁺, found 535.1836
Example 12x44: (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (method X4) rt 2.46min; m/z calculated 533.1879 for $C_{25}H_{28}ClF_2N_6O_3$ [M+H]⁺, found 533.1873

Example 12x45: (2S)-10-((5-chloro-2-(2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one.	Prepared as single diastereoisomers example 12x45a and 12x45b, see below
Example 12x45a: (S)-10-((5-chloro-2-((S)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (method X4) rt 2.75min; m/z calculated 547.2036 for C ₂₆ H ₃₀ ClF ₂ N ₆ O ₃ [M+H] ⁺ , found 547.2061
Example 12x45b: (S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (method T4) rt 2.63min; m/z calculated 547.2036 for C ₂₆ H ₃₀ ClF ₂ N ₆ O ₃ [M+H] ⁺ , found 547.2022
Example 12x46: (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of piperidine diastereoisomers using a method analogous to 9a, heating in the microwave at 160 °C for 12h. LCMS (method X4) rt 2.83 and 2.87min, ratio 2:3, m/z calculated 561.2192 for C ₂₇ H ₃₂ ClF ₂ N ₆ O ₃ [M+H] ⁺ , found 561.2200
Example 12x47: (S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 9a. Additionally heated to 160 °C for 8h in the microwave. LCMS (method X4) rt 3.07min, m/z calculated 569.1691 for C ₂₅ H ₂₆ ClF ₄ N ₆ O ₃ [M+H] ⁺ , found 569.1688

Example 12x48: (2S)-10-((5-chloro-2-(4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Prepared as single diastereoisomers, examples 12x48a and 12x48b, see below.
Example 12x48a: (S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 11a. LCMS (method X4) rt 2.64min; m/z calculated 551.1785 for C₂₅H₂₇ClF₃N₆O₃ [M+H]⁺, found 551.1783
Example 12x48b: (S)-10-((5-chloro-2-((2S,4R)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared using a method analogous to 12a, heated in the microwave to 140 °C for 3h then 160 °C for 90mins LCMS (method T4) rt 2.44min; m/z calculated 551.1785 for C₂₅H₂₇ClF₃N₆O₃ [M+H]⁺, found 551.1782
Example 12x49: (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of pyrrolidine diastereoisomers using a method analogous to 11a. LCMS (method X4) rt 2.58 min; m/z calculated 547.2036 for C₂₆H₃₀ClF₂N₆O₃ [M+H]⁺, found 547.2042
Example 12x50: (2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one 	Prepared as a mixture of 2-piperidine epimers using a method analogous to 9a, heating in the microwave at 160 °C for 12h then 180 °C for 12h LCMS (method X4) rt 3.07 min; m/z calculated 583.1848 for C₂₆H₂₈ClF₄N₆O₃ [M+H]⁺, found 583.1846

Example 13a: (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




5 Step 1: (3R,5S)-1-(5-chloro-4-iodopyridin-2-yl)-5-methylpiperidin-3-ol

[00168] (3R,5S)-5-methylpiperidin-3-ol (Intermediate M2a, 8 mg, 0.097 mmol), 5-chloro-2-fluoro-4-iodopyridine (25 mg, 0.097 mmol), and potassium carbonate (26.8 mg, 0.19 mmol) were combined in MeCN (0.5 mL) and the reaction mixture was heated to 120 °C for 12 h. The reaction mixture was purified by reverse phase chromatography (50 - 100% MeOH in water, 0.1% formic acid), and fractions containing product were loaded onto an SCX-2 cartridge, washed with methanol, and eluted with 2M NH₃ in MeOH to afford the title compound as a white solid (8.3 mg). LCMS (Method X2) RT 1.61 mins m/z 352.9928 [M+H]⁺

Step 2: (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

[00169] (3R,5S)-1-(5-chloro-4-iodopyridin-2-yl)-5-methylpiperidin-3-ol (Step 1, 8.3 mg, 0.023 mmol), (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino-[2,3-c]quinolin-6(7H)-one (Intermediate B1, 6 mg, 0.019 mmol), and trifluoroacetic acid (3.6 µL, 0.047 mmol) were combined in TFE (0.5 mL) and the reaction mixture was heated to 170 °C for 4 h. Reaction mixture loaded directly onto a reverse phase column and purified (30 - 60% MeOH in water, 0.1% formic acid), then further purified by flash column chromatography (1 - 10% MeOH in DCM, 10 g KP Sil) to give the title compound (3 mg). ¹H NMR (600 MHz, Methanol-*d*₄) δ 7.97 (d, *J* = 2.3 Hz, 1H), 7.87 (d, *J* = 1.4 Hz, 1H), 7.63 (d, *J* = 9.0 Hz, 1H), 7.58 (dd, *J* = 9.0, 2.3 Hz, 1H), 6.18 (s, 1H), 4.55 – 4.39 (m, 2H), 4.06 – 3.94 (m, 2H), 3.75 (s, 3H), 3.59 – 3.51 (m, 1H), 3.33 – 3.26 (m, 1H), 2.39 (dd, *J* = 12.2, 10.4 Hz, 1H), 2.21 (dd, *J* = 12.8, 11.3 Hz, 1H), 2.08 – 2.02 (m, 1H), 1.69 – 1.60 (m, 1H), 1.46 – 1.37 (m, 1H), 1.01 (q, *J* = 11.8 Hz, 1H), 0.93 (d, *J* = 6.6 Hz, 3H), 0.83 – 0.76 (m, 1H), 0.69 – 0.57 (m, 2H), 0.38 – 0.31 (m, 1H). HRMS (method X4) RT = 2.38 mins m/z calc for C₂₇H₃₁ClF₂N₅O₃ [M+H]⁺ = 546.2083, found 546.2070








Example 13b: ((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




[00170] Example 13b was prepared by a two-step procedure analogous to that used to prepare example 12a. ¹H NMR (600 MHz, Methanol-d₄) δ 7.97 (d, J = 2.3 Hz, 1H), 7.91 (s, 1H), 7.62 (d, J = 9.0 Hz, 1H), 7.57 (dd, J = 8.9, 2.3 Hz, 1H), 6.24 (s, 1H), 4.54 – 4.38 (m, 2H), 3.74 (s, 3H), 3.56 – 3.46 (m, 4H), 3.33 – 3.26 (m, 1H), 1.95 (tt, J = 13.6, 5.7 Hz, 4H), 1.45 – 1.36 (m, 1H), 0.83 – 0.76 (m, 1H), 0.68 – 0.57 (m, 2H), 0.36 – 0.29 (m, 1H). HRMS (method X4) RT = 2.75 mins m/z calc for C₂₆H₂₇ClF₄N₅O₂ [M+H]⁺ = 552.1790, found 552.1780.

[00171] The following tabulated examples may be prepared by methods analogous to those previously described for examples 2a, 4a, 9a, 10a, 12a and 13a.

	Example 13p01: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c01: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f01: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	
	Example 13p02: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c02: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f02: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	
	[Example 10d, described above]	Example 13c03: (2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f03: (2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	
	[Example 2b, described above]	[Example 13b, described above]	Example 13f04: (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-	

			[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	[Example 2a, described above]	[Example 13a, described above]	Example 13f05: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p06: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c06: (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f06: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p07: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c07: (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f07: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p08: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c08: (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f08: (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p09: (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c09: (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f09: (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
		Example 13c10: (S)-10-((2-((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f10: (S)-10-((2-((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p11: (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c11: (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f11: (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

	Example 13p12: (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c12: (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f12: (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one
	Example 13p13: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13c13: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one	Example 13f13: (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

[00172] For example:

Example 13p01 was prepared by analogy to example 2a, with heating to 120 °C for 4 days. LCMS (method X4) rt 3.46min, m/z calculated 565.2350 for C₂₇H₃₀F₅N₆O₂ [M+H]⁺, found 565.2383.

Example 13p06 was prepared by analogy to example 2a, with heating to 120 °C for 4 days. LCMS (method T4) rt 2.37min, m/z calculated 503.2013 for C₂₄H₂₆F₃N₆O₃ [M+H]⁺, found 503.2014.

Example 13p08 was prepared by analogy to example 2a, with heating to 120 °C for 4 days. LCMS (method X4) rt 2.81min, m/z calculated 517.2175 for C₂₅H₂₈F₃N₆O₃ [M+H]⁺, found 517.2200.

Example 13p11 was prepared by analogy to example 9a, with additional heating to 160 °C for 8h, then to 180 °C for 3h. LCMS (method X4) rt 2.71 min; m/z calculated 529.2175 for C₂₆H₂₈ClF₃N₆O₃⁺ [M+H]⁺, found 529.2155.

Example 13p12 was prepared by analogy to example 9a. LCMS (method T4) rt 2.00 min; m/z calculated 461.1907 for C₂₂H₂₄F₃N₆O₂ [M+H]⁺, found 461.1909.

Example 13c06 was prepared by analogy to example 4a: To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 16.5 mg, 0.051 mmol), 4-(5-fluoro-4-iodopyridin-2-yl)morpholine (Intermediate C4, 24.5 mg, 0.076 mmol), dicesium carbonate (126 mg, 0.39 mmol), tris(dibenzylideneacetone)dipalladium(0) (4.7 mg, 0.005 mmol), and xantphos (18.7 mg, 0.032 mmol) under argon was added DMF (0.3 mL) and toluene (0.3 mL). The resulting mixture was heated to 80 °C for 18h. The solvent was removed under reduced pressure and the residue was purified by column chromatography (0-10% MeOH in DCM). Fractions containing product were combined and the solvent removed under reduced pressure. The crude product was dissolved

in DMSO (1.0 mL) and directly purified by reverse-phase chromatography (C-18; 40%-90% methanol in water (containing 0.1% formic acid)). The fractions containing product were loaded onto an Isolute SCX-2 column (2 g), and the column was washed with methanol and then flushed with 20% 1.4 M methanolic ammonia, the solvent was removed under reduced pressure to yield the title compound (16 mg) as a white solid. LCMS (method T4): Rt 1.95 min; m/z calculated 518.1765 for C₂₅H₂₇ClF₂N₅O₂ [M+H]⁺: found: 518.1751.

Example 13c08 was prepared by analogy to example 4a: To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 16.5 mg, 0.051 mmol), (S)-4-(5-fluoro-4-iodopyridin-2-yl)-2-methylmorpholine (Intermediate C5, 25.7 mg, 0.076 mmol), dicesium carbonate (126 mg, 0.39 mmol), tris(dibenzylideneacetone)dipalladium(0) (4.7 mg, 0.005 mmol), and xantphos (18.7 mg, 0.032 mmol) under argon was added DMF (0.3 mL) and toluene (0.3 mL). The resulting mixture was heated to 80 °C for 18h. The solvent was removed under reduced pressure and the residue was purified by column chromatography (0-10% MeOH in DCM). Fractions containing product were combined and the solvent removed under reduced pressure. The crude product was dissolved in DMSO (1.0 mL) and directly purified by reverse-phase chromatography (C-18; 40%-90% methanol in water (containing 0.1% formic acid)). The fractions containing product were loaded onto an Isolute SCX-2 column (2 g), and the column was washed with methanol and then flushed with 20% 1.4 M methanolic ammonia, the solvent was removed under reduced pressure. The crude product was purified by column chromatography (20-100% EtOAc in hexane). Fractions containing product were combined and the solvent removed under reduced pressure to yield the title compound (4.8 mg) as a white solid. LCMS (method T4): Rt 2.08 min; m/z calculated 532.1921 for C₂₆H₂₉ClF₂N₅O₃ [M+H]⁺: found: 532.1917.

Example 13c12 was prepared by analogy to example 4a: To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 16.5 mg, 0.051 mmol), 5-chloro-4-iodo-N,N-dimethylpyridin-2-amine (Intermediate C3, 21.4 mg, 0.076 mmol), dicesium carbonate (126 mg, 0.39 mmol), tris(dibenzylideneacetone)dipalladium(0) (4.7 mg, 0.005 mmol), and xantphos (18.7 mg, 0.032 mmol) under argon was added DMF (0.3 mL) and toluene (0.3 mL). The resulting mixture was heated to 80 °C for 18h, then further heated in the microwave to 140 °C for 20h. The solvent was removed under reduced pressure and the residue was purified by column chromatography (0-10% MeOH in DCM). Fractions containing product were combined and the solvent removed under reduced pressure. The crude product was dissolved in DMSO (1.0 mL) and directly purified by reverse-phase chromatography (C-18; 40%-90% methanol in water (containing 0.1% formic acid)). The fractions containing product were loaded onto an Isolute SCX-2 column (2 g), and the column was washed with methanol and then flushed with 20% 1.4 M methanolic ammonia, the solvent was removed under reduced pressure to yield the title compound (13 mg, 53%, 0.0273


mmol) as a white solid. LCMS (method T4): Rt 1.90 min; m/z calculated 476.1659 for $C_{23}H_{25}ClF_2N_5O_2$ $[M+H]^+$: found: 476.1673.

Example 13f04 was prepared as mono-formic acid salt by analogy to example 4a: To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1, 10 mg, 0.031 mmol), 2-(4,4-difluoropiperidin-1-yl)-5-fluoro-4-iodopyridine (Intermediate C2, 10.7 mg, 0.031 mmol), dicesium carbonate (41 mg, 0.125 mmol), tris(dibenzylideneacetone)dipalladium(0) (3.7 mg, 0.004 mmol), and xantphos (4.5 mg, 0.008 mmol) under argon was added DMF (0.3 mL) and Toluene (0.3 mL). The resulting mixture was heated to 90 °C for 1 h, then loaded onto silica gel and purified by flash column chromatography (25g KP-Sil, 0 - 10% MeOH in DCM) to afford 27 mg of an orange oil. This was dissolved in methanol, 5 drops of formic acid were added, and the mixture loaded onto an SCX2 cartridge, washed with MeOH, and eluted with 2M NH_3 in MeOH to afford 14 mg of a yellow oil which solidified on scratching. This was further purified by reverse phase flash chromatography (Biotage 12g SNAP Ultra C18, 30-100% methanol in water, 0.1% formic acid modifier) to afford 8 mg of a white solid. ^1H NMR (600 MHz, Methanol- d_4) δ 7.91 (d, J = 2.3 Hz, 1H), 7.83 (d, J = 3.4 Hz, 1H), 7.59 (d, J = 9.0 Hz, 1H), 7.55 (dd, J = 9.0, 2.3 Hz, 1H), 6.40 (d, J = 6.1 Hz, 1H), 4.54 – 4.38 (m, 2H), 3.72 (s, 3H), 3.55 – 3.44 (m, 4H), 3.35 – 3.26 (m, 1H), 1.97 (tt, J = 13.8, 5.7 Hz, 4H), 1.46 – 1.37 (m, 1H), 0.83 – 0.76 (m, 1H), 0.68 – 0.57 (m, 2H), 0.38 – 0.31 (m, 1H). LCMS (method T4) rt 2.30min, m/z calculated 536.2079 for $C_{26}H_{27}F_5N_5O_2$ $[M+H]^+$ found 536.2058.

Example 13f05 was prepared by analogy to example 13f04 from (3R,5S)-1-(5-fluoro-4-iodopyridin-2-yl)-5-methylpiperidin-3-ol (prepared by analogy to intermediate C2). ^1H NMR (600 MHz, Methanol- d_4) δ 7.94 (d, J = 2.3 Hz, 1H), 7.81 (d, J = 3.7 Hz, 1H), 7.62 (d, J = 9.0 Hz, 1H), 7.56 (dd, J = 9.0, 2.3 Hz, 1H), 6.31 (d, J = 6.2 Hz, 1H), 4.54 – 4.34 (m, 2H), 3.99 (ddt, J = 12.2, 4.1, 1.7 Hz, 1H), 3.89 (ddt, J = 12.7, 3.9, 1.6 Hz, 1H), 3.72 (s, 3H), 3.59 (tt, J = 10.7, 4.5 Hz, 1H), 3.34 – 3.26 (m, 1H), 2.45 (dd, J = 12.2, 10.4 Hz, 1H), 2.27 (dd, J = 12.7, 11.2 Hz, 1H), 2.09 – 2.03 (m, 1H), 1.77 – 1.64 (m, 1H), 1.46 – 1.37 (m, 1H), 1.03 (q, J = 11.9 Hz, 1H), 0.95 (d, J = 6.6 Hz, 3H), 0.84 – 0.76 (m, 1H), 0.68 – 0.57 (m, 2H), 0.41 – 0.33 (m, 1H). LCMS (method T4) rt 1.99 min, m/z calculated 530.2374 for $C_{27}H_{30}F_3N_5O_3$, found 530.2375.

[00173] Examples **14a** and **14b** represent epimers at the 3-methylpiperidine centre. Each is a single compound, resulting from chiral separation of an intermediate at an earlier stage of the synthesis, but the stereochemistry of each has not been determined.


Example 14a: (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one or (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



[00174] A vial containing (S)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine or (R)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine (**Intermediate L2a**; 7 mg, 0.025 mmol), DIPEA (30 μ L, 0.17 mmol), and (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (**Intermediate B1**, 8 mg, 0.025 mmol) in NMP (2 mL) was heated at 140°C under microwave irradiation for 5.5 h. The reaction mixture was directly purified by reverse-phase chromatography (Biotage reverse-phase 12 g Ultra C-18 column; 35-80% MeOH in H₂O (containing 0.1% formic acid)) to give the title compound (0.3 mg). LCMS (Method T4) RT = 3.17 min, m/z calculated 567.1893 for C₂₆H₂₈ClF₄N₆O₂ [M+H]⁺; found 567.1898. ¹H NMR (600 MHz, Methanol-d₄) δ 8.08 (d, J = 2.3 Hz, 1H), 8.00 (s, 1H), 7.90 (dd, J = 9.1, 2.3 Hz, 1H), 7.58 (d, J = 9.1 Hz, 1H), 4.55 – 4.36 (m, 3H), 4.33 (d, J = 13.7 Hz, 1H), 3.75 (s, 3H), 3.38 – 3.24 (m, 1H), 3.03 – 2.95 (m, 1H), 2.11 – 1.95 (m, 1H), 1.92 – 1.79 (m, 1H), 1.46 – 1.37 (m, 1H), 1.31 (s, 2H), 1.01 (d, J = 6.7 Hz, 3H), 0.84 – 0.77 (m, 1H), 0.71 – 0.59 (m, 2H), 0.40 – 0.32 (m, 1H).


[00175] The following tabulated examples were prepared by a method analogous to that used for the preparation of **Example 14a**, starting from the intermediate shown in the table.

Example	Data and comments	Intermediate
Example 14b: (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one or (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-	LCMS (Method T4) RT = 3.17 min m/z calcd for C ₂₆ H ₂₈ ClF ₄ N ₆ O ₂ [M+H] ⁺ = 567.1893; found 567.1892. ¹ H NMR (600 MHz, Methanol-d ₄) δ 8.06 (d, J = 2.3 Hz, 1H), 8.00 (s, 1H), 7.92 (dd, J = 9.1, 2.3 Hz, 1H), 7.58 (d, J = 9.1 Hz, 1H), 4.55 – 4.36 (m, 3H), 4.30 (d, J = 13.8 Hz, 1H), 3.75 (s, 3H), 3.33 – 3.21 (m, 1H), 3.00 (dd, J = 13.5, 10.2 Hz, 1H), 2.12 – 1.95 (m, 1H), 1.92 – 1.77 (m, 1H), 1.46 – 1.39 (m, 1H), 1.35 – 1.30 (m, 2H), 1.00 (d, J = 6.8 Hz, 3H), 0.85 – 0.77 (m, 1H), 0.71 – 0.59 (m, 2H), 0.40 – 0.33 (m, 1H).	Intermediate L2b: (R)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine or (S)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine



Intermediate compounds

Intermediate A1: (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one




5

[00176] To a mixture of (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate B1; 1.7 g, 5.3 mmol) and 2,4,5-trichloropyrimidine (0.67 mL, 5.82 mmol) in NMP (5 mL) under argon was added DIPEA (3.7 mL, 21 mmol). The reaction mixture was heated at 140°C under microwave irradiation for 1 h then cooled to rt. The reaction mixture was added to water (5 mL) and a beige precipitate formed. Additional water (10 mL) was added and the aq. mixture was stirred for 5 min. The precipitate was filtered, washed with water (100 mL) and dried, then purified by flash column chromatography (10 g KP-Sil, 0-10% methanol in DCM) affording the title compound (2.16 g, 88%) as a beige solid. ¹H NMR (500 MHz, Chloroform-*d*) δ 8.39 (d, *J* = 2.4 Hz, 1H), 8.26 (s, 1H), 7.48 (dd, *J* = 9.1, 2.3 Hz, 1H), 7.41 (s, 1H), 7.36 (d, *J* = 9.0 Hz, 1H), 4.70 – 4.58 (m, 1H), 4.49 – 4.29 (m, 2H), 3.71 (s, 3H), 3.42 – 3.32 (m, 1H), 1.41 – 1.31 (m, 1H), 0.90 – 0.82 (m, 1H), 0.81 – 0.74 (m, 1H), 0.73 – 0.66 (m, 1H), 0.42 – 0.27 (m, 1H); LCMS (Method X4); RT 3.19 min; *m/z* 468.0801 [M+H]⁺.

15


Intermediate A2: (S)-10-((2-chloro-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one

129




[00177] Prepared by a method analogous to that used in the preparation of intermediate A1, using 2,4-dichloro-5-fluoropyrimidine. ¹H NMR (500 MHz, DMSO-*d*₆) δ 10.12 (s, 1H), 8.32 (d, *J* = 3.3 Hz, 1H), 8.20 (d, *J* = 2.3 Hz, 1H), 7.66 (dd, *J* = 9.0, 2.2 Hz, 1H), 7.50 (d, *J* = 9.0 Hz, 1H), 6.12 (d, *J* = 3.2 Hz, 1H), 4.53 – 4.43 (m, 1H), 4.43 – 4.32 (m, 1H), 3.57 (s, 3H), 3.32 – 3.22 (m, 1H), 1.38 – 1.25 (m, 1H), 0.75 – 0.67 (m, 1H), 0.58 – 0.48 (m, 2H), 0.38 – 0.32 (m, 1H); LCMS (Method X2); RT 1.52 min; *m/z* 452.1098 [M+H]⁺.

Intermediate A3: (S)-2-cyclopropyl-10-((2,5-dichloropyrimidin-4-yl)amino)-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one



[00178] Prepared by a method analogous to that used in the preparation of intermediate A1, using (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-(methyl-d₃)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one (Intermediate B2). LCMS (Method T2); RT 1.54 min; *m/z* 472.09 [M+H]⁺.


Intermediate B1: (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one



[00179] Three reactions were set up in parallel as follows: to (S)-2-cyclopropyl-3,3-difluoro-7-methyl-10-nitro-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (Intermediate E1; 2.5 g, 7.1 mmol) and 10 wt% Pd/C (100 mg) under an argon atmosphere was added ethanol (50 mL). The reaction mixture was stirred at 60°C under an atmosphere of hydrogen for 90 minutes then allowed to cool to rt. The three reaction mixtures were combined and filtered through a pad of celite, washing with EtOH (200 mL). The filtrate was concentrated in vacuo affording the title

compound (6.78 g, 99%) as an orange solid which was used without further purification. LCMS (Method X4); RT 1.93 min; m/z 322.14 $[M+H]^+$.

Intermediate B2: (S)-10-amino-2-cyclopropyl-3,3-difluoro-7-(methyl-d3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one




5

[00180] A sealed vial containing (S)-10-chloro-2-cyclopropyl-3,3-difluoro-7-(methyl-d3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one (Intermediate E2, 20.3 mg, 0.059 mmol), sodium tert-butoxide (8.8 mg, 0.092 mmol), diphenylmethanimine (16 mg, 0.088 mmol), palladium(II) acetate (1.3 mg, 0.006 mmol) and Josiphos (3.2 mg, 0.006 mmol) in 1,2-dimethoxyethane (0.24 mL) under Ar was heated at 70°C for 2 h. The reaction mixture was cooled to rt and 3 M HCl (0.5 mL) was added. The mixture was stirred at rt for 2 h, then filtered through a pad of Celite onto an SCX-2 (2 g) column, which was eluted with water (10 mL), MeOH (20 mL) and 2 N methanolic ammonia (20 mL). The basic fraction was concentrated under reduced pressure affording (2S)-10-amino-2-cyclopropyl-3,3-difluoro-7-(trideuteriomethyl)-2,4-dihydro-1H-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6-one (18.5 mg) as a dark yellow solid. LCMS (Method T2); RT 1.20 min; m/z 326.1471 $[M+H]^+$.

15

Intermediate C1: 5-(4,5-dichloropyridin-2-yl)-3-methyl-1,2,4-oxadiazole




20

[00181] To 4,5-dichloropicolinic acid (80 mg, 0.42 mmol) in THF (2.7 mL) under Ar was added T3P (50 wt% in EtOAc, 0.34 g, 0.54 mmol), DIPEA (0.15 mL, 0.86 mmol) and acetamide oxime (31 mg, 0.42 mmol). The reaction mixture was sealed and stirred at rt for 3 h, then heated at 80°C in a heating block for 15 h. Water (5 mL) was added and the aqueous mixture was extracted with EtOAc (2 x 10 mL). The organic extracts were combined, washed with saturated aq. NaHCO_3 (5 mL), brine (10 mL), dried (Na_2SO_4) and concentrated *in vacuo* affording the title compound (44 mg, 45%) as an off-white solid. ^1H NMR (500 MHz, CDCl_3) δ 8.80 (s, 1H), 8.29 (s, 1H), 2.52 (s, 3H); LCMS (2 min); RT 1.32 min; m/z 229.9880 $[M+H]^+$


25

Intermediate C2: 2-(4,4-difluoropiperidin-1-yl)-5-fluoro-4-iodopyridine




[00182] 4,4-Difluoropiperidine hydrochloride (63 mg, 0.40 mmol), 2,5-difluoro-4-iodo-pyridine (80 mg, 0.33 mmol), and potassium carbonate (153 mg, 1.11 mmol) were combined in MeCN (1.0 mL) and heated to 170 °C for 12 h in the microwave. The resulting mixture was loaded onto silica gel and purified by flash column chromatography (25g KP-Sil, 0 - 5% MeOH in DCM) to give the title compound (10 mg) as a clear oil. LCMS (Method X2) rt 1.51 min; m/z 342.9942 [M+H]⁺.

Intermediate C3: 5-chloro-4-iodo-N,N-dimethylpyridin-2-amine




[00183] A mixture of dimethylamine hydrochloride (21 mg, 0.26 mmol), 5-chloro-2-fluoro-4-iodopyridine (60 mg, 0.23 mmol), and *N,N*-diisopropylethylamine (80 uL, 0.47 mmol) in NMP (0.5 mL) was heated to 140 °C for 1 h, then concentrated under reduced pressure and the residue purified by column chromatography (0-5% MeOH in DCM) to yield title compound as an off white solid (56 mg). ¹H NMR (500 MHz, Chloroform-d) δ 8.07 (s, 1H), 6.99 (s, 1H), 3.05 (s, 6H).

Intermediate C4: 4-(5-fluoro-4-iodopyridin-2-yl)morpholine



[00184] A mixture of morpholine (18 uL, 0.21 mmol), 5-chloro-2-fluoro-4-iodopyridine (50 mg, 0.19 mmol), and *N,N*-diisopropylethylamine (67 uL, 0.38 mmol) in MeCN (0.5 mL) was heated to 80 °C for 18 h, then concentrated under reduced pressure and the residue purified by column chromatography (0-5% MeOH in DCM) to yield title compound as an off white solid (29 mg). ¹H NMR (600 MHz, Chloroform-d) δ 8.12 (s, 1H), 7.13 (s, 1H), 3.84 – 3.79 (m, 4H), 3.50 – 3.45 (m, 4H).


Intermediate C5: (S)-4-(5-fluoro-4-iodopyridin-2-yl)-2-methylmorpholine



[00185] A mixture of (S)-2-Methylmorpholine hydrochloride (29 mg, 0.21 mmol), 5-chloro-2-fluoro-4-iodopyridine (50 mg, 0.19 mmol), and *N,N*-diisopropylethylamine (67 uL, 0.38 mmol) in MeCN (0.5 mL) was heated to 80 °C for 18 h, then concentrated under reduced pressure and the

residue purified by column chromatography (0-5% MeOH in DCM) to yield title compound as an off white solid (27 mg). ¹H NMR (600 MHz, Chloroform-d) δ 8.11 (s, 1H), 7.12 (s, 1H), 4.00 (tdd, J = 12.6, 3.1, 1.5 Hz, 2H), 3.90 (ddt, J = 12.7, 3.1, 1.6 Hz, 1H), 3.73 – 3.62 (m, 2H), 3.01 – 2.91 (m, 1H), 2.60 (dd, J = 12.7, 10.4 Hz, 1H), 1.27 (d, J = 6.3 Hz, 3H).

5 **Intermediate E1: (S)-2-cyclopropyl-3,3-difluoro-7-methyl-10-nitro-1,2,3,4-tetrahydro-[1,4]oxazepino-[2,3-c]quinolin-6(7H)-one**



Step 1: (S)-4-((1-cyclopropyl-2,2-difluoro-3-hydroxypropyl)amino)-1-methyl-6-nitroquinolin-2(1H)-one

10 **[00186]** An oven-dried microwave vial (10–20 mL volume) was charged with (S)-3-amino-3-cyclopropyl-2,2-difluoropropan-1-ol hydrochloride (3 g, 13.3 mmol) and ethyl 4-chloro-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate (Intermediate F1; 3 g, 9.7 mmol). The reaction vial was flushed with Ar and sealed with a cap. Anhydrous acetonitrile (15 mL) was added followed by DIPEA (4.2 mL). The reaction mixture was heated at 160 °C under microwave irradiation for 15

15 h. Four further batches were prepared by the same method. Batches were combined, 2M sodium hydroxide (150 mL) was added, and the reaction mixture was heated at 85 °C for 2 h. The reaction mixture was cooled to rt. Water (80 mL) was added and the reaction mixture was acidified to pH 5 - 6 with 3 M HCl. The resulting precipitate was filtered, washed with H₂O (500 mL) and dried affording the title compound (15.83 g, 93%) as a beige solid which was used without further

20 purification. ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.31 (d, J = 2.5 Hz, 1 H), 8.40 (dd, J = 9.4, 2.5 Hz, 1 H), 7.62 (d, J = 9.4 Hz, 1 H), 7.47 (d, J = 8.7 Hz, 1 H), 5.72 (s, 1 H), 5.60 (t, J = 6.1 Hz, 1 H), 3.90–3.71 (m, 2 H), 3.57–3.45 (m, 1 H) overlapping with 3.55 (s, 3H), 1.38–1.29 (m, 1 H), 0.71–0.64 (m, 1 H), 0.63–0.56 (m, 1 H), 0.53–0.46 (m, 1 H), 0.27–0.21 (m, 1 H); LCMS (Method X2) RT 1.17 min; m/z 354.1264 [M+H]⁺.

25 Step 2: (S)-3-bromo-4-((1-cyclopropyl-2,2-difluoro-3-hydroxypropyl)amino)-1-methyl-6-nitroquinolin-2(1H)-one

[00187] Trifluoroacetic acid (8.83 mL) was added to a stirred mixture of (S)-4-((1-cyclopropyl-2,2-difluoro-3-hydroxypropyl)amino)-1-methyl-6-nitroquinolin-2(1H)-one (from step 1; 8.15 g, 23 mmol) and freshly recrystallised *N*-bromosuccinimide (4.1 g, 23 mmol) in anhydrous CH₂Cl₂ (150

30 mL) at 0 °C under Ar. The reaction mixture was stirred at 0 °C for 15 min. The reaction mixture was diluted with CH₂Cl₂ (100 mL) and washed with saturated aq. NaHCO₃ (3 x 100 mL). The aqueous washings were further extracted with CH₂Cl₂ (100 mL). The organic extracts were combined, washed with brine (100 mL), dried (Na₂SO₄) and concentrated. A second batch was


prepared by an analogous method, starting from 7.69g of the quinolinone from step 1. Batches of products were combined to give the title compound as a yellow solid (18.6 g, 96%). ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.95 (d, *J* = 2.5 Hz, 1 H), 8.43 (dd, *J* = 9.4, 2.5 Hz, 1 H), 7.75 (d, *J* = 9.4 Hz, 1 H), 5.85 (d, *J* = 11.1 Hz, 1 H), 5.63 (t, *J* = 5.8 Hz, 1 H), 4.05–3.95 (m, 1 H), 3.88 - 3.73 (m, 2 H), 3.71 (s, 3 H), 1.29 – 1.21 (m, 1 H), 0.68 – 0.62 (m, 1 H), 0.62 – 0.51 (m, 2 H), 0.50–0.44 (m, 1 H); LCMS (Method T2) RT 1.45 min; *m/z* 432.0373 [M+H]⁺.

Step 3: (S)-2-cyclopropyl-3,3-difluoro-7-methyl-10-nitro-1,2,3,4-tetrahydro-[1,4]oxazepino-[2,3-*c*]quinolin-6(7H)-one

[00188] Lithium *tert*-butoxide (1 M in THF; 35 mL) was added to a stirred suspension of (S)-3-bromo-4-((1-cyclopropyl-2,2-difluoro-3-hydroxypropyl)amino)-1-methyl-6-nitro-quinolin-2(1H)-one (from step 2; 9.48 g, 22 mmol) in THF (220 mL) under Ar. A reflux condenser and Ar balloon were fitted and the reaction mixture was heated at 60 °C for 15 min. The reaction mixture was cooled to rt. Water (100 mL) was added and the aqueous mixture was extracted with CH₂Cl₂ (3 x 100 mL). The organic extracts were combined, washed with brine (2 x 100 mL), dried (Na₂SO₄) and concentrated under reduced pressure. A second batch was prepared by the same method using a further 9.17g of compound from step 2. Batches were combined to give the title compound as a yellow solid (15.4 g, 100%). ¹H NMR (500 MHz, DMSO-*d*₆) δ 9.12 (d, *J* = 2.6 Hz, 1 H), 8.35 (dd, *J* = 9.3, 2.5 Hz, 1 H), 7.66 (d, *J* = 9.3 Hz, 1 H), 7.01 (br d, *J* = 5.0 Hz, 1 H), 4.54–4.37 (m, 2 H), 3.62 (s, 3 H, NCH₃), 3.29–3.22 (m, 1 H), 1.39–1.31 (m, 1 H), 0.76–0.69 (m, 1 H), 0.58–0.49 (m, 2 H), 0.37–0.30 (m, 1 H); LCMS (Method X2) RT 1.34 min; *m/z* 352.1105 [M+H]⁺. The product was found to be ~88% ee by chiral SFC [YMC Chiral ART Amylose-C, (4.6mm x 250mm, 5μm); 30:70 methanol:CO₂ (isocratic, 0.2% v/v ammonia modifier); flowrate 4 mL/min; 40 °C; detection at 210-400nm; 1 μL injection]. Rt of desired (S) enantiomer 2.06min; (R) enantiomer 1.75min. 10 g of material was therefore dissolved in DCM:methanol (1:1 v/v, concentration 15 mg/mL) and purified by chiral SFC [Chiralpak SA (21.2mm x 250mm, 5μm); 30:70 methanol:CO₂ (isocratic, 0.2% v/v ammonia modifier); flowrate 21 mL/min; ambient temperature; detection at 229nm; 2.5 mL per injection]. Combined fractions of the desired (S) enantiomer were then evaporated to near dryness using a rotary evaporator, transferred into final vessels with DCM, which was removed under a stream of compressed air at 35 °C before being stored in a vacuum oven at 35 °C and 5mbar until constant weight to afford a beige solid (7.51g). Chiral SFC [Amylose-C, as above] rt 2.09 min; no minor isomer observed.


Intermediate E2: (S)-10-chloro-2-cyclopropyl-3,3-difluoro-7-(methyl-*d*3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*][1,8]naphthyridin-6(7H)-one

134



[00189] Prepared by a three step method analogous to that used for the preparation of intermediate E1, starting from ethyl 4,6-dichloro-1-(methyl-d3)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxylate (Intermediate F2). Chiral SFC analysis / purification was not carried out. ¹H NMR (500 MHz, DMSO-d₆) δ 8.76 (d, J = 2.4 Hz, 1 H), 8.60 (d, J = 2.4 Hz, 1 H), 6.60 - 6.50 (m, 1 H), 4.56-4.35 (m, 2 H), 3.30 - 3.23 (m, 1 H), 1.31 - 1.24 (m, 1 H), 0.76 - 0.69 (m, 1 H), 0.60-0.48 (m, 2 H), 0.37-0.31 (m, 1 H); LCMS (Method T2); RT 1.49 min; m/z 345.1034 [M+H]⁺.

Intermediate F1: ethyl 4-chloro-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate



Step 1: 1-methyl-6-nitro-2H-benzo[d][1,3]oxazine-2,4(1H)-dione

[00190] To a solution of 5-nitro-isatoic anhydride (25.1 g, 120.6 mmol) in DMF (241 mL) at rt was added sodium hydride (60% in mineral oil; 7.24 g, 180.9 mmol). The solution was allowed to stir for 15 min with warming to RT. Iodomethane (18.8 mL, 301.5 mmol) was added and the mixture was stirred at rt for 4 h. The reaction mixture was poured onto ice, the resulting precipitate filtered and washed with water (5 litres). The solid was collected and dried under vacuum overnight affording 1-methyl-6-nitro-2H-benzo[d][1,3]oxazine-2,4(1H)-dione (19.7 g, 73%) as an orange powder. LCMS (Method T2) RT 1.35 min, m/z 211.069 [M+MeOH-CO₂]⁺.

Step 2: ethyl 4-hydroxy-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate


[00191] To a solution of 1-methyl-6-nitro-2H-benzo[d][1,3]oxazine-2,4(1H)-dione (from step 1; 19.6 g, 88.2 mmol) in DMF (177 mL) was added diethyl malonate (40.4 mL, 264.7 mmol). The solution was cooled to 0°C, then Sodium hydride (60% in mineral oil) (7.06 g, 176.46 mmol) was added in 4 portions over 30 min. The solution was allowed to warm to rt and stirred at that temperature for 3 h. Water was added with care to the reaction mixture followed by 10% aq. HCl until the pH of the mixture was ~pH5. The resulting precipitate was filtered through a sinter funnel and washed with water (5 litres). The resulting solid was transferred to a round bottom flask and dried under vacuum affording ethyl 4-hydroxy-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate (24.2 g, 94%) as a pale yellow solid. LCMS (Method T2) RT 1.45 min; m/z 293.074 [M+H]⁺.

Step 3: ethyl 4-chloro-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate

[00192] Phosphorus oxychloride (250 mL, 2700 mmol) was added to a flask containing ethyl 4-hydroxy-1-methyl-6-nitro-2-oxo-1,2-dihydroquinoline-3-carboxylate (24.1 g, 82.5 mmol). The flask was fitted with a suba-seal and an argon balloon then heated to 80°C with stirring for 2.5 h. The mixture was concentrated *in vacuo*. The residue was diluted with water and extracted with EtOAc.

5 The combined organic extracts were washed with brine, dried (MgSO₄) and concentrated *in vacuo*. Purification by flash chromatography (340 g KP-sil; 0% to 10% MeOH in CH₂Cl₂) afforded the title compound (14.5 g, 57%) as a dark orange solid. ¹H NMR (500 MHz, CDCl₃) δ 8.95 (d, *J* = 2.5 Hz, 1H), 8.50 (dd, *J* = 9.3, 2.5 Hz, 1H), 7.53 (d, *J* = 9.3 Hz, 1H), 4.48 (q, *J* = 7.1 Hz, 2H), 3.78 (s, 3H), 1.42 (t, *J* = 7.1 Hz, 3H); LCMS (Method T2) RT 1.42 min; *m/z* 311.043 [M+H]⁺.

10 **Intermediate F2: ethyl 4,6-dichloro-1-(methyl-*d*₃)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxylate**



Step 1: methyl 5-chloro-2-((methyl-*d*₃)amino)nicotinate

[00193] A microwave vial (10–20 mL volume) was charged with methyl 5-chloro-2-fluoronicotinate (261 mg, 1.38 mmol) and methan-*d*₃-amine hydrochloride (192 mg, 2.73 mmol). The reaction vial was flushed with Ar, sealed with a cap and then further flushed with Ar. THF (4 mL) was added followed by DIPEA (0.71 mL, 4.11 mmol). The reaction mixture was stirred at 40°C for 22 h then concentrated *in vacuo*. The residue was diluted with water (10 mL) and extracted with EtOAc (2 x 15 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by flash chromatography (10 g KP-sil; 0% to 30% EtOAc in cyclohexane) afforded the title compound (239 mg, 85%) as a white solid. LCMS (Method T2) RT 1.43 min, *m/z* 204.06 [M+H]⁺

Step 2: ethyl 6-chloro-4-hydroxy-1-(methyl-*d*₃)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxylate

25 [00194] A microwave vial (2.0–5.0 mL volume) was charged with methyl 5-chloro-2-((methyl-*d*₃)amino)nicotinate (from step 1; 239 mg, 1.17 mmol). The reaction vial was flushed with Ar, sealed with a cap and then further flushed with Ar. DCM (4.5 mL) was added followed by triethylamine (0.98 mL, 7.0 mmol) and ethyl 3-chloro-3-oxopropanoate (0.6 mL, 4.7 mmol). The reaction mixture was heated at 60°C in a heating block for 2 h. Additional ethyl 3-chloro-3-oxopropanoate (0.6 mL, 4.7 mmol) and triethylamine (0.98 mL, 7.0 mmol) were added and the reaction mixture was stirred at 60°C for a further 6 h. The reaction mixture was concentrated *in vacuo*. Water (20 mL) was added, followed by 10% aq. HCl (10 mL). The aqueous mixture was extracted with EtOAc (3 x 20 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na₂SO₄) and concentrated *in vacuo*. Purification by flash chromatography (10 g KP-


sil; 0% to 10% MeOH in DCM) afforded the title compound (335 mg, 100%) as an orange solid. LCMS (Method T2) RT 1.55 min; m/z 286.066 $[M+H]^+$

Step 3: ethyl 4,6-dichloro-1-(methyl- d_3)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxylate

[00195] Phosphorus oxychloride (3.5 mL, 37.4 mmol) was added to a flask containing ethyl 6-chloro-4-hydroxy-1-(methyl- d_3)-2-oxo-1,2-dihydro-1,8-naphthyridine-3-carboxylate (from step 2; 335 mg, 1.2 mmol). The flask was fitted with a reflux condenser with a suba-seal and an argon balloon then heated to 80°C with stirring for 2.5 h. The reaction mixture was cooled to rt and concentrated *in vacuo*. The crude reaction mixture was dissolved in EtOAc (20 mL) and washed with water (2 x 10 mL) and saturated aq. NaHCO_3 (10 mL). The aqueous washings were combined and extracted with EtOAc (20 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na_2SO_4) and concentrated *in vacuo*. Purification by flash chromatography (10 g KP-sil; 0-60% EtOAc in cyclohexane) afforded the title compound (52 mg, 15%) as a yellow solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.87 (d, $J = 2.4$ Hz, 1 H), 8.46 (d, $J = 2.4$ Hz, 1 H), 4.38 (q, $J = 7.1$ Hz, 2 H), 1.31 (t, $J = 7.1$ Hz, 3 H); LCMS (Method T2) RT 1.51 min; m/z 304.0287 $[M+H]^+$

[00196] Intermediate L1a and Intermediate L1b represent a pair of enantiomers where one is the (*R*)- and the other is the (*S*)-methyl-piperidine. It has not been determined which is the (*R*)- and which is the (*S*)-enantiomer. The compounds were separated by preparative chiral SFC during Step 3 using the method described below.

Intermediate L1a and Intermediate L1b: (*S*)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol and (*R*)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol



Step 1: 2,5-dichloropyrimidin-4-ol

[00197] 2 M sodium hydroxide (6 mL, 12 mmol) was added to a stirred solution of 2,4,5-trichloropyrimidine (1.29 g, 7.0 mmol) in THF (4 mL). The reaction mixture was stirred at rt for 24 h. The reaction mixture was concentrated *in vacuo* and the aqueous mixture was neutralised with 3 M HCl. The aqueous mixture was extracted with Et_2O (2 x 10 mL) followed by EtOAc (2 x 10 mL). The organic extracts were combined, washed with brine (10 mL), dried (Na_2SO_4) and concentrated *in vacuo* affording 2,5-dichloropyrimidin-4-ol (923 mg, 80%) as a yellow solid which was used without further purification. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 8.26 (s, 1 H); LCMS (Method T2) RT 0.19 min; m/z 164.9602 $[M+H]^+$

Step 2: rac-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol

[00198] A microwave vial (10–20 mL volume) was charged with 2,5-dichloropyrimidin-4-ol (from step 1; 502 mg, 3.0 mmol) and rac-4,4-difluoro-3-methylpiperidine hydrochloride (522 mg, 3.0


mmol). The reaction vial was flushed with Ar, sealed with a cap and then further flushed with Ar. Ethanol (5 mL) was added followed by DIPEA (1.4 mL, 8.0 mmol). The reaction mixture was heated at 80°C in a heating block for 10 h. The reaction mixture was concentrated *in vacuo*. The residue was dissolved in DMSO (1 mL) and directly purified by reverse-phase chromatography (Biotage reverse-phase 12 g C-18 column; 25-80% MeOH in H₂O (containing 0.1% formic acid)) affording *rac*-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol (264 mg, 33%) as an off-white solid. ¹H NMR (600 MHz, CDCl₃) δ 12.18 (br s, 1 H), 7.89 (s, 1 H), 4.49–4.41 (m, 1 H), 4.34–4.27 (m, 1 H), 3.37–3.30 (m, 1 H), 3.08 (dd, *J* = 13.8, 10.9 Hz, 1 H), 2.26–2.17 (m, 1 H), 2.17–2.08 (m, 1 H), 2.02–1.89 (m, 1 H), 1.16 (d, *J* = 6.8 Hz, 3 H); LCMS (Method T2) RT 1.21 min; *m/z* 264.0675 [M+H]⁺

Step 3: (S)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol and (R)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol

[00199] *rac*-5-Chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol (250 mg) was dissolved to 25 mg/mL in MeOH:CH₂Cl₂ (4:1) (1% v/v NH₃) and was then purified by SFC (Amy-C (20 mm x 250 mm, 5 μm), 10:90 MeOH:CO₂ (0.2% v/v NH₃); flow rate 50 mLmin⁻¹). The earlier eluting enantiomer was identified as **Intermediate L1a** and the later eluting enantiomer was identified as **Intermediate L1b**. Combined fractions of **Intermediate L1a** were concentrated *in vacuo* before being stored in a vacuum oven at 35°C and 5mbar affording **Intermediate L1a** (85 mg) as a white solid. Combined fractions of **Intermediate L1b** were concentrated *in vacuo* and re-purified and isolated as above to afford **Intermediate L1b** (68 mg) as a white solid.

[00200] Chiral purity analysis was determined by SFC (Amy-C (4.6 mm x 250 mm, 5 μm), 10:90 MeOH:CO₂ (0.2% v/v NH₃); flow rate 4 mLmin⁻¹). **Intermediate L1a**: ee = 99.2 %; RT 3.58 min. **Intermediate L1b**: ee = 99.0 %; RT 3.83 min.


Intermediate L2a: (S)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine or (R)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine



[00201] To a vial containing (S)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol or (R)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol (**Intermediate L1a**; 43 mg, 0.16 mmol) was added POCl₃ (0.6 mL, 6.4 mmol) and the vial was sealed and heated to 90°C for 3 h. The excess POCl₃ was removed *in vacuo*, and the residue was partitioned between water and EtOAc. The layers were separated and the organic layer was dried (MgSO₄), and concentrated *in vacuo* to give the title compound (41 mg, 89%) as a colourless oil which was used without further purification. ¹H NMR (500 MHz, CDCl₃) δ 8.22 (s, 1 H), 4.56–4.50 (m, 1 H), 4.47–4.41 (m,

1 H), 3.40–3.33 (m, 1 H), 3.11–3.04 (m, 1 H), 2.21–2.11 (m, 1 H), 2.11–1.98 (m, 1 H), 1.96–1.82 (m, 1 H), 1.09 (d, $J = 6.8$ Hz, 3 H); LCMS (Method T2); RT 1.71 min; m/z 282 $[M+H]^+$.

[00202] The following tabulated examples were prepared by a method analogous to that used for the preparation of **Intermediate L2a**, starting from the aryl sulfide shown in the table.

Intermediate	Data and comments	Pyrimidin-4-ol
Intermediate L2b: (<i>R</i>)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine or (<i>S</i>)-4,5-dichloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidine 	LCMS (Method T2); RT 1.71 min; m/z 282 $[M+H]^+$.	Intermediate L1b: (<i>R</i>)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol or (<i>S</i>)-5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-ol

5

Intermediate M1a: (*3R,5S*)-1-benzyl-5-methylpiperidin-3-ol [and **Intermediate M1b:** (*3S,5R*)-1-benzyl-5-methylpiperidin-3-ol]



[00203] The commercially available *rac*-1-benzyl-5-methylpiperidin-3-ol.HCl (10 g) was dissolved to 66 mg/mL in 2:1 MeOH:DCM and was then purified by SFC (Phenomenex Lux Amylose-1 (30 mm x 250 mm, 5 μ m), 10:90 MeOH:CO₂ (0.2% v/v DEA; flow rate 50 ml min⁻¹; 40 °C; 1.7 mL per injection). The earlier eluting enantiomer was assigned as Intermediate M1a and the later eluting enantiomer was assigned as Intermediate M1b. Combined fractions of Intermediate M1a were evaporated to dryness using a rotary evaporator, redissolved in DCM, precipitated with heptane and solvents removed using a rotary evaporator, before being stored in a vacuum oven at 35 °C and 5 mbar until constant weight. This gave Intermediate M1a (3.61 g) as a white solid. Combined fractions of Intermediate M1b were evaporated to dryness using a rotary evaporator, redissolved in DCM, washed with water, dried over magnesium sulfate, precipitated with heptane and solvents removed using a rotary evaporator, before being stored in a vacuum oven at 35 °C and 5 mbar until constant weight. This gave Intermediate M1b (3.26 g) as a white solid. Small molecule X-ray of intermediate M1b, recrystallized from ether/DCM gave an assignment of this isomer as (*3S,5R*)-1-benzyl-5-methylpiperidin-3-ol. Intermediate M1a was therefore assigned as the (*3R,5S*)-1-benzyl-5-methylpiperidin-3-ol isomer. Analysis of chiral purity was carried out using SFC [YMC Chiral ART Amylose-C, (4.6 mm x 250 mm, 5 μ m), 10:90 MeOH:CO₂ (0.2% v/v DEA);


10

15

20

flow rate 4 mLmin⁻¹). Intermediate M1a: rt 2.19 min, other isomer not detected. Intermediate M1b: rt 3.00 min, ee = 99 %.

Intermediate M2a: (3R,5S)-5-methylpiperidin-3-ol



- 5 **[00204]** To a solution of (3R,5S)-1-benzyl-5-methylpiperidin-3-ol (Intermediate M1a, 1.5 g, 0.73 mmol) in ethanol (120 mL) under argon was added Pd/C (10 wt%; 778 mg). The flask was evacuated and back-filled with hydrogen twice before being stirred at room temperature under a hydrogen balloon for 1 h. The reaction was filtered through celite (eluent methanol) and the filtrate was concentrated *in vacuo* to give the title compound (814 mg, 97%) as a white solid which was
- 10 used without further purification. ¹H NMR (500 MHz, methanol-*d*₄) δ 3.62 – 3.53 (m, 1H), 3.12 – 3.05 (m, 1H), 2.91 – 2.83 (m, 1H), 2.22 (dd, J = 11.8, 10.5 Hz, 1H), 2.08 – 2.00 (m, 2H), 1.69 – 1.56 (m, 1H), 0.96 (td, J = 12.1, 11.0 Hz, 1H), 0.91 (d, J = 6.7 Hz, 3H); LCMS (Method T4) RT 0.14 min; m/z 116.1216 [M+H]⁺.

Biological assays

15 ***HTRF assay***

- [00205]** Assays were performed in a 384-well black Proxiplate (Perkin Elmer) containing 1 nM Trx-6xHis-BCL6 (in house-produced, human BCL6 BTB domain covering amino-acid sequence 5-129), 300 nM BCOR-AF633 peptide (RSEIISTAPSSWVPGP-Cys-AlexaFluor 633-amide, Cambridge Research Biochemical) and 0.5nM anti-6xHis-Terbium cryptate (CisBio Bioassays,
- 20 France), in assay buffer (25 mM Hepes pH8, 100 mM NaCl, 0.05% Tween20, 0.5 mM TCEP, 0.05% bovine serum albumin). Test compounds in DMSO or DMSO alone were added to the wells using an ECHO550 acoustic dispenser (Labcyte Inc) to give the appropriate test concentration in 0.7% v/v DMSO final. After 2 hours incubation at room temperature the plate was read on a Pherastar FSX (BMG Labtech) plate reader equipped with 337 nm laser excitation filter,
- 25 a first emission filter at 620 nm and a second emission filter at 665 nm. The % inhibition at each concentration was calculated by normalising FRET ratio to the appropriate high (DMSO with all reagents) and low (DMSO without BCL6) controls. The compound IC50s were determined using GraphPad Prism 6.0 or Dotmatics (Bishops Stortford, UK) software by fitting the normalised data to a sigmoidal four-parameter logistic fit equation.

- 30 **[00206]** The results of this assay are shown in Table 1 above.

Mesoscale Discovery (MSD) Degradation Assay

[00207] An MSD assay was developed for screening of compounds to determine the degradation of endogenous BCL6 in OCI-Ly1 cells. Briefly, test compounds in DMSO or DMSO alone to a total of 1332.5 nL/well were dispensed to a 96-well Nunc™ Edge 2 plate (ThermoFisher

Scientific, 267544) using an Echo® 550 acoustic dispenser (Labcyte Inc) to give the appropriate test concentration in 0.67% v/v DMSO final. OCI-Ly1 passaged in Iscove's Modified Dulbecco's Medium (IMDM, ThermoFisher Scientific, 12440053) supplemented with 10% heat-inactivated fetal bovine serum (Sera Plus, PAN Biotech, P30-3702) were added to the compound/DMSO at a density of 2.5×10^5 cells/mL in 200 μ L media/ well. After 2 hours incubation at 37 °C + 5% CO₂ the plate was centrifuged at 300 x g for 5 minutes. Using a BioTek 405TS plate washer for all wash and aspiration steps, media was removed and cells were rinsed with PBS, centrifuged as before and PBS removed. Cells were lysed in 50 μ L ice-cold lysis buffer at pH 7.4 containing 50 mM Tris-HCl, 150 mM NaCl, 1 % Triton X-100 (v/v), 1 mM PMSF, 1 mM activated sodium vanadate, 1 mM EDTA, phosphatase inhibitors 2 and 3 (1:50 dilution) and protease inhibitor cocktail (1:100 dilution). The plate was shaken briefly and incubated on ice for 5-10 minutes before being frozen overnight at -20 °C or continuing the assay. A 96-well MSD standard bind plate (MSD, L15XA-3) which was coated overnight with 1 μ g/mL anti-human BCL6 goat antibody in PBS (R&D Systems, AF5046) was blocked in 3% BSA in TBS + 0.1% Tween-20 for 1 hour with shaking. Wells were emptied thoroughly before 40 μ L/well of lysate was transferred from the cell plate. The MSD plate was incubated at RT for 1 hour with shaking then washed three times with TBS + 0.1 % Tween-20. Antibodies were prepared in TBS + 0.1% Tween-20 + 1% BSA. Anti-human BCL6 rabbit antibody (Cell Signalling Technology, 14895S) was added at 1:100 dilution at 25 μ L/well. The plate was incubated and washed as before. MSD Sulfo-tag anti-rabbit detection antibody (MSD, R32AB-1) at 1:1000 dilution at 25 μ L/well was added. The plate was incubated and washed as before and MSD Gold read buffer (MSD R92TG) at 1:1 dilution with water at 150 μ L/well was added. The plate was read within 10 minutes on the MSD Quickplex reader. The % inhibition at each concentration was calculated by normalising the electrochemiluminescent signal to the appropriate high (DMSO) and low ((5-((5-chloro-2-((3R,5S)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one) at 2 μ M) controls. DC50 values (compound concentration at which 50% of endogenous BCL6 protein is degraded) were determined using GraphPad Prism 6.0 or Dotmatics (Bishops Stortford, UK) software by fitting the normalised data to a sigmoidal four-parameter logistic fit equation.

[00208] The results obtained using this assay are shown in Table 2 above.

Mouse Splenocyte Flow Cytometry Immunology Experiments

[00209] Male C57BL/6 mice were immunised in both flanks subcutaneously with CFA (100 μ L of 4mg/ml CFA). On day 8 the animals were terminated, spleens harvested and processed to single cell suspensions.

[00210] Splenocytes were resuspended in complete media (RPMI 1640 + 10% foetal calf serum, supplemented with 4mM L-glutamine, 100U/ml-100 μ g/ml) penicillin/streptomycin and 50 μ M B-mercaptoethanol (Gibco)). The cells were activated for 48 hours with 2 μ g/ml α CD40

(ThermoFisher) + 0.002 μ g/ml IL-4 (Bio-technique) at 37°C in a humidified CO₂ incubator. Example 4b was added at the same time as activation in an 8-point dose titration. Following the culture period, cells were collected, washed with PBS (Gibco) and selected wells were counted using a LUNATMCounter (Logos Biosystems).

- 5 **[00211]** Harvested cells were washed in PBS (Gibco) and stained with InvitrogenTM eBioscienceTM Fixable Viability Dye eFluor 780 (Fisher scientific), then FcR blocked (Miltenyi Biotec) and stained with surface receptor antibodies (see Table 3 below). Cells were then washed in flow cytometry buffer (PBS supplemented with 1%BSA and 0.05% azide) and the secondary streptavidin was added. Cells were then washed, fixed and permeabilized in Transcription factor
- 10 staining buffer (Fisher scientific); according to the manufacturer's instructions. Intracellular antibodies were then added to the samples. Following final washes, cells were fixed in 1% PFA and acquired on a BD LSR FortessaTM (BD Bioscience). All analysis was carried out using FlowJo software v10.

Table 3 – surface receptor antibody stains

Antibody	Fluorochrome	Clone	Supplier
CD19	BV785	1D3	BD
CD4	BUV395	GK1.5	BD
Bcl6	PEDazzle	K112-91	BD
Blimp-1	AF647	5 e 7	BD
CD20	Percpcy5.5	SA275A11	Biolegend
CD95	BV605	SA367H8	Biolegend
GL7	Biotin	GL7	Biolegend

15

- [00212]** In order to test the hypothesis that degradation or inhibition of Bcl6 may be able to modulate primary cells of the immune system, resulting in a decrease in their potential to generate autoantibodies, the above experiment was performed. Mice were immunised in order to drive the generation of germinal centre (GC) B cells. Spleens were then cultured in the presence or
- 20 absence of a Bcl6 inhibitor (Example 4b) for 48 hours and non-GC B cells or GC B cells were identified using flow cytometry (Figure 1).

- [00213]** As expected the addition of the Bcl6 inhibitor did not result in a reduction of BCL6 expression in non-GC B cells (Figure 2), whereas it did result in an increase in Blimp expression by GC B cells (Figure 3). Blimp is involved in the repression of B cell metabolism and GC B cell
- 25 formation. It blocks antibody class switching and antigen presentation by B cells.

- [00214]** The addition of the Bcl6 inhibitor to spleen cell cultures decreased the numbers of GC B cells as determined by expression of the lineage markers GL7 and CD95 (Figure 4).

CLAIMS

1. A compound, or a pharmaceutically acceptable salt or solvate thereof, selected from:
- (S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((2-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide;
- 15 N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide;
- (S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-hydroxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one ;
- (S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-
- 5 [1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-
- 15 cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide;
- (3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-
- 25 carboxamide;
- (S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-
- 35 difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide;
- 40 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide;
- (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-
- 45 difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide;
- 5 (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide;
- N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide;
- 10 N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide;
- (S)-10-((5-chloro-2-((1R,4R)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((1S,4S)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide;
- 20 N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide;
- (S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((3S,5R)-3,5-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((S)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-fluoro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-[[5-chloro-2-(4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl]amino]-2-cyclopropyl-3,3-difluoro-7-methyl-2,4-dihydro-1H-[1,4]oxazepino[2,3-c]quinolin-6-one;
- 30 (S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-(methyl(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (;
- (S)-10-((5-chloro-2-(5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3,3-dimethylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile;
- 35 (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile;
- (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile;

- (S)-10-((5-chloro-2-(3-(difluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((S)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((S)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 ((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((R)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3,4-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (2S)-10-((5-chloro-2-(2,3-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((2R,5S)-2-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-((3R,5S)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-((3S,5R)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-ethynylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-*c*]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile;
- (S)-10-((5-chloro-2-((4,4-difluorocyclohexyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(2-azabicyclo[4.1.0]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile;
- 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide;
- 20 (2S)-10-((5-chloro-2-(3-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((2S,4R)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (2S)-10-((5-chloro-2-(6-hydroxy-4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-methyl-5-oxopiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((3R,5S)-3-(3-(dimethylamino)propoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (2S)-10-((5-chloro-2-(4,4-difluoro-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(4,4-difluoro-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-chloropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-methylpiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-2-cyclopropyl-10-((2-(3,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((R)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((1R,5S)-3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((2-((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one; and
- (2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one.
- 40

2. A compound, or a pharmaceutically acceptable salt or solvate thereof, according to claim 1 selected from:

- (S)-10-((5-chloro-2-(3,3-difluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(4-cyclopropyl-3-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-((S)-3,4-dimethyl-5-oxopiperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((1R,4R)-2-oxa-5-azabicyclo[2.2.1]heptan-5-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 N-((3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide;
- N-((3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-methylpiperidin-3-yl)acetamide;
- 15 (S)-10-((5-chloro-2-((3S,5R)-3-(dimethylamino)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((S)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-(3-fluoro-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-hydroxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-(3-oxa-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-(methyl-d3)-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c][1,8]naphthyridin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-(3-methyl-1,2,4-oxadiazol-5-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((3-chloro-2-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-2-cyclopropyl-10-((2,3-dichloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one ;
- 40 (S)-10-((2-chloro-3-fluoropyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((3-chloro-2-methylpyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((1R,5S,7R)-7-hydroxy-3-oxa-9-azabicyclo[3.3.1]nonan-9-yl)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (2S)-10-((2-(3-acetyl-3,8-diazabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((2-(8-acetyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((S)-3-((dimethylamino)methyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (3R,5S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide;
- 20 (3S,5R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-5-methylpiperidine-3-carboxamide;
- (S)-10-((5-chloro-2-((3R,5S)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((3S,5R)-3-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(fluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 2-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide;
- 2-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidin-3-yl)acetamide;
- 40 (S)-10-((5-chloro-2-((S)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(dimethylamino)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide;
- 5 (S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-N-ethyl-4,4-difluoropiperidine-3-carboxamide;
- N-(((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide;
- 10 N-(((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)methyl)-N-ethylacetamide;
- (S)-10-((5-chloro-2-((1R,4R)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((1S,4S)-5,5-difluoro-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- N-((R)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide;
- 20 N-((S)-1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-4,4-difluoropiperidin-3-yl)-N-methylacetamide;
- (S)-10-((5-chloro-2-(4-fluoro-4-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((R)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((1S,5R)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((1R,5S)-3-ethyl-2-oxo-3,8-diazabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((3S,5R)-3,5-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-(6-acetyl-2,6-diazaspiro[3.3]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2-hydroxy-2-methylpropyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((3-hydroxy-3-methylbutyl)(methyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(6-(cyclopropylmethyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(6-((2,2-difluorocyclopropyl)methyl)-2,6-diazaspiro[3.3]heptan-2-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(8-(cyclopropylmethyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(8-((2,2-difluorocyclopropyl)methyl)-3,8-diazabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((3R,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((3S,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((S)-3-methoxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-fluoro-2-((3S,4R)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-fluoro-2-((3R,4S)-3,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-[[5-chloro-2-(4-methyl-1,4-diazepan-1-yl)pyrimidin-4-yl]amino]-2-cyclopropyl-3,3-difluoro-7-methyl-2,4-dihydro-1H-[1,4]oxazepino[2,3-c]quinolin-6-one;
- 30 (S)-10-((5-chloro-2-((R)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-ethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((S)-3-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((S)-4-methylazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3,5-dihydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-10-((5-chloro-2-((1R,5R)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-((1S,5S)-1-(hydroxymethyl)-5-methyl-3-azabicyclo[3.1.0]hexan-3-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5R)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((3S,5S)-3-fluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-7-hydroxy-5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-(methyl(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,4S)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3S,4R)-3-(hydroxymethyl)-4-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((2-hydroxyethyl)(propyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one (;
- (S)-10-((5-chloro-2-(5-azaspiro[2.5]octan-5-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-(((R)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(((S)-3-methoxy-2-methylpropyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3,3-dimethylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-(3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidine-3-carbonitrile;
- 35 (S)-10-((5-chloro-2-(3-fluoroazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-(difluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-3-methylazetidine-3-carbonitrile;
- (S)-2-(1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)azetidin-3-yl)acetonitrile;

- (S)-10-((5-chloro-2-(3-(difluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-(azetidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-(3-methoxyazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-methoxy-3-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((S)-3-ethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((S)-3-fluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-hydroxy-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((2-((3S,5R)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((2-((3R,5S)-3-amino-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (2S)-10-((2-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3,3-difluoro-5-hydroxypiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((R)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-methylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((R)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,6-dimethyl-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((R)-hexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (S)-10-((5-chloro-2-((S)-6-oxohexahydropyrrolo[1,2-a]pyrazin-2(1H)-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-((S)-8-hydroxy-6-azaspiro[3.5]nonan-6-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 ((S)-10-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-10-((5-chloro-2-((R)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 20 (S)-10-((5-chloro-2-((R)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-fluoro-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((3R,5S)-3-methoxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethylthiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3,4-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (2S)-10-((5-chloro-2-(2,3-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethyl-1-oxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((2R,5S)-2-(hydroxymethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,6-dimethyl-1,1-dioxidothiomorpholino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((3S,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(6-hydroxy-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(6-hydroxy-4-(methyl-*d*₃)-1,4-diazepan-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-((3S,5R)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-((3R,5S)-3-(2-(azetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-((3R,5S)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-((3S,5R)-3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(azetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(2-(3,3-difluoroazetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(2-(azetidin-1-yl)-2-oxoethyl)-4,4-difluoropiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-ethynylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-*c*]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile;
- (S)-10-((5-chloro-2-((4,4-difluorocyclohexyl)amino)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(2-azabicyclo[4.1.0]heptan-2-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(3-(difluoromethyl)piperazin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((3R,5S)-3-(2-(dimethylamino)ethoxy)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;
- (2S)-10-((2-(3-(azetidine-1-carbonyl)-5-methylpiperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-*c*]quinolin-6(7H)-one;

- (2S)-10-((5-chloro-2-(3-(3,3-difluoroazetidine-1-carbonyl)-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 5 (S)-10-((5-chloro-2-(3-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 10 (S)-10-((5-chloro-2-((R)-2-(hydroxymethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(3-(fluoromethyl)azetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-methylazetidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 15 (S)-1-(5-chloro-4-((2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile;
- 1-(5-chloro-4-(((S)-2-cyclopropyl-3,3-difluoro-7-methyl-6-oxo-1,2,3,4,6,7-hexahydro-[1,4]oxazepino[2,3-c]quinolin-10-yl)amino)pyrimidin-2-yl)-5-fluoro-N,N-dimethylpiperidine-3-carboxamide;
- 20 (2S)-10-((5-chloro-2-(3-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-(4-fluoropiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 25 (S)-10-((5-chloro-2-((S)-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((S)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 30 (S)-10-((5-chloro-2-((R)-2-(methoxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 35 (S)-10-((5-chloro-2-((S)-4,4-difluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (S)-10-((5-chloro-2-((2S,4S)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- 40 (S)-10-((5-chloro-2-((2S,4R)-4-fluoro-2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
- (2S)-10-((5-chloro-2-(2-(hydroxymethyl)-3-methylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;

(2S)-10-((5-chloro-2-(4,4-difluoro-2-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(2S)-2-cyclopropyl-10-((2-(4,4-difluoro-3,5-dimethylpiperidin-1-yl)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
5 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-morpholinopyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((S)-2-methylmorpholino)pyrimidin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(S)-10-(((1R,5S)-8-oxa-3-azabicyclo[3.2.1]octan-3-yl)-5-fluoropyrimidin-4-yl)amino)-2-cyclopropyl-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
10 (S)-2-cyclopropyl-10-((2-(dimethylamino)-5-fluoropyrimidin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(S)-2-cyclopropyl-10-((2-(dimethylamino)-5-chloropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
15 (S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-morpholinopyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(S)-2-cyclopropyl-3,3-difluoro-10-((5-chloro-2-((S)-2-methylmorpholino)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one;
(S)-2-cyclopropyl-10-((2-(4,4-difluoropiperidin-1-yl)-5-fluoropyridin-4-yl)amino)-3,3-difluoro-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one; and
20 (S)-2-cyclopropyl-3,3-difluoro-10-((5-fluoro-2-((3R,5S)-3-hydroxy-5-methylpiperidin-1-yl)pyridin-4-yl)amino)-7-methyl-1,2,3,4-tetrahydro-[1,4]oxazepino[2,3-c]quinolin-6(7H)-one.

3. A pharmaceutical composition comprising a compound according to claim 1 or claim 2, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier or excipient.

4. A compound according to claim 1 or claim 2, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition according to claim 3, for use in therapy.

5. A compound according to claim 1 or claim 2, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition according to claim 3, for use in the treatment of cancer, HIV, sepsis, graft-versus-host disease, non-alcoholic fatty liver disease, non-alcoholic steatohepatitis or auto-immune diseases.

6. A compound according to claim 1 or claim 2, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition according to claim 3, for use in the treatment of cancer in combination with an additional therapeutic agent.

7. A compound or a pharmaceutical composition for the use according to claim 6, wherein the additional therapeutic agent is selected from a BCL-2 family inhibitor, a BTK inhibitor, a TNF inhibitor, an EZH2 inhibitor, a HDAC inhibitor, an ATR kinase inhibitor, an ATM kinase inhibitor, an EGFR tyrosine kinase inhibitor and a corticosteroid.

8. A compound or a pharmaceutical composition for the use according to claims 5 to 7, wherein said cancer is diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL), angioimmunoblastic T-cell lymphoma (AITL), acute lymphoblastic leukaemia (ALL), acute myeloid leukaemia (AML), chronic myeloid leukaemia (CML), multiple myeloma, breast cancer, non-small cell lung cancer (NSCLC) or squamous cell carcinomas (SCC) of the head and neck, oesophagus, lung or ovary.

9. A method for the treatment of cancer in a subject in need of such treatment, said method comprising administering a therapeutically effective amount of a compound according to claim 1 or claim 2, or a pharmaceutically acceptable salt or solvate thereof, or a pharmaceutical composition according to claim 3.

10. A method according to claim 9, wherein said cancer is diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL), angioimmunoblastic T-cell lymphoma (AITL), acute lymphoblastic leukaemia (ALL), acute myeloid leukaemia (AML), chronic myeloid leukaemia (CML), multiple myeloma, breast cancer, non-small cell lung cancer (NSCLC) or squamous cell carcinomas (SCC) of the head and neck, oesophagus, lung or ovary.




FIG. 1




FIG. 2




FIG. 3




FIG. 4

INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2020/052588

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D498/04 A61K31/553 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2018/215798 A1 (CANCER RESEARCH TECH LTD [GB]) 29 November 2018 (2018-11-29) see structure of the BCL6 inhibitors according to claim 1 and examples e.g. 1k, 3g and 3o -----	1-10
A	WO 2018/108704 A1 (BOEHRINGER INGELHEIM INT [DE]) 21 June 2018 (2018-06-21) see structure of the BCL6 inhibitors according to claim 1 and e.g. examples 1-161 to 164 and 166 to 169 -----	1-10
X,P	WO 2019/197842 A1 (CANCER RESEARCH TECH LTD [GB] ET AL.) 17 October 2019 (2019-10-17) see the structure of the compounds of claim 1 as BCL6 inhibitors -----	1-10

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

11 December 2020

Date of mailing of the international search report

22/12/2020

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Traegler-Goeldel, M

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2020/052588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2018215798 A1	29-11-2018	EP 3630749 A1 WO 2018215798 A1	08-04-2020 29-11-2018

WO 2018108704 A1	21-06-2018	EP 3555063 A1 JP 2020503374 A US 2020071297 A1 WO 2018108704 A1	23-10-2019 30-01-2020 05-03-2020 21-06-2018

WO 2019197842 A1	17-10-2019	AU 2019253510 A1 CA 3095371 A1 SG 11202009735Q A WO 2019197842 A1	22-10-2020 17-10-2019 29-10-2020 17-10-2019
