



HU000025619T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 025 619**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 09 722278**(22) A bejelentés napja: **2009. 03. 13.**

(96) Az európai bejelentés bejelentési száma:

EP 20090722278

(97) Az európai bejelentés közzétételi adatai:

EP 2265251 A2 **2009. 09. 24.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2265251 B1 **2015. 05. 27.**(51) Int. Cl.: **A61K 47/44** (2006.01)**A61K 312/15** (2006.01)**A61K 312/16** (2006.01)**A61K 315/35** (2006.01)**A61K 315/42** (2006.01)**A61K 315/75** (2006.01)**A61K 45/06** (2006.01)

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 09/037077

(87) A nemzetközi közzétételi szám:

WO 09117316

(30) Elsőbbségi adatok:

37117 P **2008. 03. 17.** **US****111920 P** **2008. 11. 06.** **US**

(72) Feltalálók(k):

KABRA, Bhagwati, P., EulestTX 76039 (US)**CARRERAS, Nuria, 08140 Caldes De Montbui
(barcelona) (ES)****CUCHI, Mónica, Barcelona (ES)****GALÁN, Francisco, Javier, 08329 Teia (Barcelona) (ES)****MRIDVIKA, 08348 Cabriels (Barcelona) (ES)****PUIG, Elena, González, 08013 Barcelona (ES)****JIMÉNEZ, Nuria, 08005 Barcelona (ES)****MARTÍNEZ, Carmen, 08185 Lliçá De Vall (Barcelona)
(ES)**

(73) Jogosult(ak):

Alcon Research, Ltd., Fort Worth, TX**76134-2099 (US)**

(74) Képviselő:

**ifj. Szentpéteri Ádám, SBGK Szabadalmi
Ügyvivői Iroda, Budapest**

(54)

Előnyös biológiai hozzáférhetőségű gyógyászati készítmények

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 265 251 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
27.05.2015 Bulletin 2015/22

(21) Application number: **09722278.0**

(22) Date of filing: **13.03.2009**

(51) Int Cl.:
A61K 47/44 ^(2006.01) **A61K 31/535** ^(2006.01)
A61K 31/5575 ^(2006.01) **A61K 31/542** ^(2006.01)
A61K 45/06 ^(2006.01) **A61K 31/215** ^(2006.01)
A61K 31/216 ^(2006.01)

(86) International application number:
PCT/US2009/037077

(87) International publication number:
WO 2009/117316 (24.09.2009 Gazette 2009/39)

(54) **PHARMACEUTICAL COMPOSITIONS HAVING DESIRABLE BIOAVAILABILITY**

PHARMAZEUTISCHE ZUSAMMENSETZUNGEN MIT ERWÜNSCHTER BIOVERFÜGBARKEIT
COMPOSITIONS PHARMACEUTIQUES AYANT LA BIODISPONIBILITÉ SOUHAITABLE

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK TR**

(30) Priority: **17.03.2008 US 37117 P**
06.11.2008 US 111920 P

(43) Date of publication of application:
29.12.2010 Bulletin 2010/52

(60) Divisional application:
12193597.7 / 2 599 475
13184332.8 / 2 705 836

(73) Proprietor: **Alcon Research, Ltd.**
Fort Worth, TX 76134-2099 (US)

(72) Inventors:
• **KABRA, Bhagwati, P.**
Euless
TX 76039 (US)
• **CARRERAS, Nuria**
08140 Caldes De Montbui (barcelona) (ES)
• **CUCHI, Mónica**
Barcelona (ES)
• **GALÁN, Francisco, Javier**
08329 Teia (Barcelona) (ES)
• **MRIDVIKA**
08348 Cabriels (Barcelona) (ES)
• **PUIG, Elena, González**
08013 Barcelona (ES)

• **JIMÉNEZ, Nuria**
08005 Barcelona (ES)
• **MARTÍNEZ, Carmen**
08185 Lliçà De Vall (Barcelona) (ES)

(74) Representative: **Teipel, Stephan et al**
Lederer & Keller
Patentanwälte Partnerschaft mbB
Unsöldstrasse 2
80538 München (DE)

(56) References cited:
EP-A1- 1 547 599 **WO-A1-00/03736**
WO-A1-97/29752 **WO-A1-98/19680**
WO-A1-98/25620 **WO-A1-98/41208**
WO-A1-98/53809 **US-A- 5 932 572**
US-A1- 2004 082 660 **US-B1- 6 743 439**

• **FELDMAN ET AL: "Comparison of the Ocular
Hypotensive Efficacy of Adjunctive Brimonidine
0.15% or Brinzolamide 1% in Combination with
Travoprost 0.004%" OPTHALMOLOGY, J. B.
LIPPINCOTT CO., PHILADELPHIA, PA, US, vol.
114, no. 7, 1 July 2007 (2007-07-01), pages
1248-1254.E2, XP022137571 ISSN: 0161-6420**

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 265 251 B1

Description

Cross-Reference to Related Applications

5 [0001] The present application claims priority based on U.S. Provisional Patent Application Serial No. 61/037,117 filed March 17, 2008, and U.S. Provisional Patent Application Serial No. 61/111,920, filed November 6, 2008.

Technical Field of the Invention

10 [0002] The present invention is related to pharmaceutical compositions that contain surfactant concentrations that promote bioavailability of a therapeutic agent in the composition. More specifically the present invention relates to topical pharmaceutical compositions (e.g., multi-dose ophthalmic compositions) having relatively low concentrations of surfactant that promote the bioavailability of a therapeutic agent (a prostaglandin such as travoprost).

15 **Background of the Invention**

[0003] The present invention is directed to pharmaceutical compositions formulated to exhibit enhanced bioavailability of a therapeutic agent of the composition. The composition may also exhibit other additional or alternative desired characteristics. For example, the composition may also be sterile, may exhibit desired antimicrobial or preservation efficacy, may exhibit a desired degree of stability, combinations thereof or the like.

20 [0004] Therapeutic agents (e.g., ophthalmic drugs) of many pharmaceutical compositions are often required to be stable within those compositions. It is typically undesirable for the therapeutic agents or overall compositions to decompose or chemically or physically change to a significant degree prior to application of the agents to an individual or otherwise. For maintaining stability, pharmaceutical compositions are typically formulated with ingredients that can enhance such stability or ingredients that can minimize destabilizing effects of other entities (e.g., chemicals, ambient conditions or the like).

25 [0005] Surfactants are one preferred family of ingredients that have exhibited significant efficacy in stabilizing pharmaceutical compositions and/or therapeutic agents thereof (particularly aqueous ophthalmic solutions including relatively lipophilic and/or relatively insoluble components). Moreover, it has generally been believed that stability of an agent or composition can often be achieved by using larger concentrations of surfactant within that pharmaceutical composition.

30 [0006] In addition to stability of the agents or compositions, it is also typically desirable for the therapeutic agents of the pharmaceutical compositions to exhibit relatively high degrees of bioavailability. Such bioavailability often becomes particularly important where the method or manner of application or dosing of a particular pharmaceutical composition provides only a limited amount of time for the therapeutic agent of that composition to be absorbed or otherwise taken in by a biological target such as an eye, ear, throat or nose of an individual. As an example, topically applied ophthalmic pharmaceutical compositions may only dwell in or on an individual's eye for a limited period of time (e.g., before tears transport the composition elsewhere). Thus, it is often desirable to limit the concentration of any ingredient in a pharmaceutical composition where that ingredient tends to inhibit bioavailability of a therapeutic agent of that composition.

35 [0007] Recently, it has been discovered that surfactants, when used at certain higher concentrations, may act as an ingredient that can limit the bioavailability of a therapeutic agent, particularly an ophthalmic therapeutic agent. As a consequence, the addition of surfactant to pharmaceutical compositions and particularly ophthalmic compositions can limit the bioavailability and, in turn, the efficacy of the therapeutic agent in the composition. However, as suggested above, surfactants can also be quite desirable in a pharmaceutical composition since they can significantly enhance the stability of a pharmaceutical composition or therapeutic agent.

40 [0008] US 6,743,439 discloses aqueous pharmaceutical solution compositions comprising a cationic drug, a cationic preservative, and a sulfonated styrene/mealeic anhydride copolymer.

[0009] WO 00/03736 discloses a pharmaceutical product comprising an aqueous prostaglandin formulation and a polypropylene container.

[0010] EP-A-1 547 599 discloses an ophthalmic solution comprising latanoprost.

50 [0011] US 2004/0082660 A1 discloses an ophthalmic composition which contains substantially no benzalkonium chloride.

[0012] WO 98/41208 discloses an ophthalmic composition including prostaglandin, non-ionic surfactants, preservatives, and non-ionic toxicity adjusting agents.

[0013] WO 98/25620 discloses ophthalmic suspensions containing brinzolamide.

55 [0014] US 5,932,572 discloses ophthalmic compositions comprising a combination of a beta-blocker and a carbonic anhydrate inhibitor.

[0015] WO 98/19680 discloses a combination of carbonic anhydrase inhibitors and prostaglandins for treating glaucoma.

[0016] WO 98/23809 discloses combinations of a prostaglandin and a topical carbonic anhydrase inhibitor for the treatment of ocular hypertension.

[0017] In view of the above, it would be desirable to provide a pharmaceutical composition with a lower surfactant concentration and/or a higher bioavailability of a therapeutic agent. Moreover, it would also be desirable, although not required unless otherwise specifically stated, for such composition to exhibit a desirable level of stability.

Summary of the Invention

[0018] Accordingly, the present invention is directed to a pharmaceutical composition, namely an ophthalmic composition, that combines a therapeutic agent with a relatively low surfactant concentration. Typically, the composition will exhibit a higher bioavailability of the therapeutic agent particularly when used for topical applications. In one embodiment, an effectively low amount of surfactant is provided such that an area under a concentration/time curve when determined for the pharmaceutical composition of the present invention as applied to a biological target is at least 130%, more typically at least 200% and even possibly at least 250% relative to an area under a similar concentration/time curve when determined for a control composition as applied to the biological target. For such embodiment, the control composition will typically have at least double the amount of surfactant relative to the pharmaceutical composition.

[0019] The present invention has been found particularly suitable for use in multi-dose ophthalmic solutions. The combination of therapeutic agent and surfactant for such compositions is the combination of a prostaglandin (e.g., travoprost) with an ethoxylated and/or hydrogenated vegetable oil (e.g., Polyoxyl 40 Hydrogenated castor oil).

Brief Description of the Drawing

[0020]

Fig. 1 is an exemplary graph of free therapeutic agent relative to surfactant concentration for an exemplary ophthalmic composition in accordance with an aspect of the present invention.

Fig. 2 is an exemplary graph of concentration of prostaglandin therapeutic agent at a biological target versus time.

Fig. 3 is an exemplary graph of concentration of prostaglandin therapeutic agent at a biological target versus time in accordance with an aspect of the present invention.

Fig. 4. is an exemplary graph of concentration of prostaglandin therapeutic agent at a biological target versus time in accordance with an aspect of the present invention.

Fig. 5. is an exemplary graph of concentration of therapeutic agent (i.e., carbonic anhydrase inhibitor) at a biological target versus time in accordance with an aspect of the present invention.

Fig. 6 is an exemplary graph of concentration of therapeutic agent (i.e., carbonic anhydrase inhibitor) at a biological target versus time in accordance with an aspect of the present invention.

Detailed Description of the Invention

[0021] The present invention is predicated upon the provision of a pharmaceutical composition having a relatively low amount of surfactant and enhanced bioavailability of a therapeutic agent of the composition. The pharmaceutical composition is particularly desirable as a solution suitable for topical application to a biological target of the human body, namely the eye. In a highly preferred embodiment, the pharmaceutical composition is an aqueous ophthalmic composition that is provided as a solution. Moreover, it is preferred that the ophthalmic solution have a therapeutic agent suitable for treatment of one or more eye or ophthalmic maladies such as allergies, glaucoma, dry eye, macular degeneration, cataracts, combinations thereof or the like. As one highly preferred example, a therapeutic agent such as travoprost might be combined in an ophthalmic composition with a relatively low amount of surfactant for the treatment of glaucoma.

[0022] Unless otherwise indicated, percentages provided for the ingredients of the pharmaceutical composition of the present invention are weight/volume percentages (w/v %).

[0023] A combination of surfactant and therapeutic agent can be used according to the present invention to enhance the bioavailability of the therapeutic agent in the pharmaceutical composition. Examples of compositions include: ophthalmic pharmaceutical compositions, such as topical compositions used in the treatment of glaucoma, infections, allergies or inflammation; compositions for treating contact lenses, such as cleaning products and products for enhancing the ocular comfort of patients wearing contact lenses; and various other types of ophthalmic compositions, such as

ocular lubricating products, artificial tears, astringents, and so on. As suggested, the compositions can be completely aqueous solutions, suspensions or otherwise.

[0024] The compositions of the present invention may contain various types of therapeutic agent. The invention can include therapeutic agents that are nonionic, anionic, cationic or combinations thereof. The therapeutic agent that exhibits higher bioavailability according to the present invention will typically be substantially or entirely non-ionic. The compositions of the present invention can also include one or more therapeutic agents where the bioavailability of those agents are not significantly affected by surfactant concentrations while the bioavailability of one or more other therapeutic agents are affected. For example, one of the former therapeutic agents could be part of a suspension while one of the latter therapeutic agents may be in the solution (e.g., dissolved in aqueous solution) of the suspension.

[0025] Examples of therapeutic agents that may be contained in the ophthalmic or other compositions of the present invention include timolol (e.g., timolol maleate), olopatadine (e.g., olopatadine hydrochloride), brinzolamide, tandospirone, roscovitine, nepafenac, combinations thereof or the like. Examples of therapeutic agents that may exhibit increased bioavailability in accordance with the present invention include, without limitation, hypotensive lipids (e.g., bimatoprost), and glucocorticoids (e.g., prednisolone, dexamethasone and lotoporednol). Therapeutic agents that typically exhibit significant increased bioavailability in accordance with the present invention are prostaglandins (e.g., latanoprost, travoprost and unoprostone).

[0026] As used herein, it is contemplated that the term "prostaglandin" includes, without limitation, natural prostaglandins, prostaglandin analogs, prostaglandin derivatives or any combination thereof.

[0027] The amount of therapeutic agent in the pharmaceutical composition will depend upon factors such as the efficacy of therapeutic agent at different concentrations, the compatibility of the therapeutic agent with other ingredients in the composition, the ability of biological target to accept various amounts of therapeutic agent, combinations thereof or the like. Generally speaking, the pharmaceutical composition can include at least 0.0001% by weight or w/v %, at least 0.001% by weight or w/v% or even at least 0.01% or 0.1% by weight or w/v % or more of the therapeutic agent.

[0028] Therapeutic agent that exhibits a desired degree of improved bioavailability according to the present invention is typically composed of agent that exhibits relatively low solubility in water. Thus, it is contemplated that the therapeutic agent of the pharmaceutical composition, particularly when it is employed in an aqueous ophthalmic composition, can exhibit solubility in water that is less than 0.1%, more typically less than 0.05%. It is also typically desirable for the therapeutic agent to be non-ionic, particularly in aqueous solution. It is also typically desirable for the therapeutic agent to be dissolved in solution of the ophthalmic or pharmaceutical composition which is typically accomplished with the assistance of the surfactant.

[0029] Further, therapeutic agent that exhibits a desired degree of improved bioavailability according to the present invention is typically lipophilic (i.e., it prefers an organic phase as compared to water or an aqueous phase). Such agent typically has a relatively high octanol/water partition coefficient. Thus, it is contemplated that the therapeutic agent of the pharmaceutical composition, particularly when it is employed in an aqueous ophthalmic composition, can exhibit an octanol/water partition coefficient that is typically at least 5 and more typically at least 10.

[0030] It is contemplated that the therapeutic agent can be partially or substantially entirely of one or more therapeutic agents that have the aforementioned solubility and/or partition component. As used herein, the term "substantially entirely", when used to describe what ingredient[s] are part of a component of the ophthalmic composition, means that it is contemplated that the component is formed entirely of one or more particular ingredient[s] or is formed substantially entirely of those one or more particular ingredient[s] with only a nominal amount (e.g., less than 5% or 1% by weight) of the component being formed of other than those one or more particular ingredients.

[0031] The surfactant included in the pharmaceutical composition of the present invention will often depend upon the therapeutic agent in the composition or other ingredients of the composition. Preferably, the surfactant can increase the solubility of the therapeutic agent and/or at least assist in assuring that the agent is distributed evenly in the composition. The surfactant may also promote the ability of the therapeutic agent to penetrate human tissue (e.g., corneal tissue of the eye) thereby further increasing the bioavailability of the agent.

[0032] The amount of surfactant will typically depend upon the therapeutic agent employed in the composition. The amount of surfactant employed is typically chosen so as to increase the bioavailability of the therapeutic agent. Generally speaking, the pharmaceutical composition can include at least 0.01% by weight or w/v% or at least 0.05% by weight or w/v% of the surfactant.

[0033] The surfactant can include non-ionic, an anionic, a cationic, or an amphoteric or zwitterionic surfactant or a combination of such surfactants. It is highly preferred that at least a portion or substantially the entirety of the surfactant be non-ionic for assisting in providing enhanced bioavailability of the therapeutic agent. As used herein, the phrase "substantially the entirety of the surfactant" is used to suggest either the entirety of the surfactant or the entirety of the surfactant with the exception of a nominal amount of surfactant or both.

[0034] Examples of potentially suitable surfactant include, without limitation, ethers of fatty alcohols and/or polyoxyethylene alkyl ethers, e.g., macrogol ethers such as cetomacrogol 1000, polyoxyethylene castor oil derivatives, polyoxyethylene sorbitan fatty acid esters, e.g., the commercially available Tweens.TM., polyoxyethylene stearates, combi-

nations thereof or the like.

[0035] Surfactant that assists in providing for a desired degree of bioavailability according to the present invention is typically composed of agent that exhibits a relatively high hydrophile/lipophile/balance (HLB). Thus, it is contemplated that the surfactant of the pharmaceutical composition, particularly the ophthalmic composition, has an HLB value greater than 8, preferably greater than 10 and even possibly greater than 12.

[0036] The surfactant may include polysorbate 20 (TWEEN 20) (polyoxyethylene 20 sorbitan monolaurate), TWEEN 40, TWEEN 60, polysorbate 80 (TWEEN 80), Zwittergent 312, TEEPOL HB7, SPAN 85, pluronic or poloxamers, especially, PLURONIC L62LF, L101, and L64, F68, L44, L121, F-84 and P-103, PEG1000, and/or TETRONIC 1501, 150R1, 701, 901, 1301, and 130R1, poloxamer 333, poloxamer 334, and poloxamer 335, sorbitan oleate, polysorbate 81, polysorbate 85, polysorbate 120, sodium taurocholates, sodium deoxytaurocholates, chenodeoxycholic acid, ursodeoxycholic acid, or combinations thereof.

[0037] The surfactant includes vegetable oils that have been hydrogenated, ethoxylated or both. Such surfactants include, but are not limited, to babassu oil, almond oil, maize oil, palm kernel oil, castor oil, coconut oil, cotton seed oil, jojoba oil, linseed oil, mustard oil, olive oil, peanut oil, safflower oil sesame oil, soybean oil, sunflower-seed oil and wheat germ oil, their hydrogenated or ethoxylated derivatives or combinations thereof. Preferred oils are castor oil, babassu oil, almond oil, maize oil and palm kernel oil, most preferably castor oil and cababassu oil.

[0038] Particularly preferred surfactants include Polyoxyethylene (POE) (40) Hydrogenated castor oil (or PEG (40 Hydrogenated castor oil) (HCO-40), POE (60) Hydrogenated castor oil (HCO-60), and POE (200) Hydrogenated castor oil (HCO-200).

[0039] Without being bound by theory, it is believed that use of higher amounts or concentrations of surfactant relative to the therapeutic agent can results in higher amounts of the therapeutic agent being absorbed into the micelle of the surfactants as the therapeutic agent is solubilized. In turn, it is believed that such absorption can limit the amount of therapeutic agent readily available to a biological target (e.g., the cornea of a human eye) during topical application of the pharmaceutical composition (e.g., topical application of an ophthalmic solution). It should be understood that this theory is not binding upon the scope of the present invention unless otherwise specifically recited.

[0040] It is contemplated that the pharmaceutical composition of the present invention can include an effectively low amount of surfactant such that the concentration of the therapeutic agent located at a biological target is substantially greater than the concentration of therapeutic agent located at the same biological target after a separate application of a control composition. As used herein, a "separate application of a control composition" is an application of the control composition to a same biologic target of a separate animal. For example, testing can be performed on two sets of ten rabbits apiece wherein the composition of the present invention is applied to an eye of each rabbit of the first set while the control composition is applied to an eye of each rabbit of the second set. In such embodiment, the control composition is substantially identical to the pharmaceutical composition with the exception that the concentration of surfactant is at least doubled, more preferably tripled and even more typically quadrupled in the control composition relative to the pharmaceutical composition of the present invention. Moreover, the pharmaceutical composition of the present invention is applied in an amount that is equivalent to the amount of control composition applied.

[0041] For quantifying such concentrations, a graph is developed, an example of which is shown in Fig. 2, plotting concentration at the biological target relative to time after application of the pharmaceutical composition. The concentration is determined at three separate times at the biological target. In particular, the concentration is determined at 1 hour, 2 hours and 4 hours after application. These points are then plotted on the graph. Examples of such points 10, 12, 14 are shown in Fig. 2. Those points are then connected by line segments and the area 16 under those line segments (referred to herein as area under the concentration/time curve) is determined as a quantification of concentration. Using this area under the curve measurement, it has been determined that the area under the concentration/time curve when determined for the pharmaceutical composition of the present invention as applied to the biological target is at least 130%, more typically at least 150% and even possibly at least 200% relative to the area under the concentration/time curve when determined for the control composition as applied to the biological target. One preferred protocol for application of control composition and pharmaceutical composition as well as measurement of concentration of therapeutic agent is provided in the example section below.

[0042] As an example of area under the curve measurements, an exemplary situation is provided wherein the concentrations at 1, 2 and 4 hours for the composition of the present invention is 20 nanograms per milliliter (ng/ml) and the concentrations at 1, 2 and 4 hours for the control composition are 10 ng/ml. In such a situation the area under the curve for the composition of the present invention is 20 ng/ml x 3 hr, which is 60 while the area under curve for the control composition is 10 ng/ml x 3 hr, which is 30. In this scenario, the area under the curve for the composition of the present invention is 200% that of the area under the curve for the composition of the control composition.

[0043] It has been found that the bioavailability of certain therapeutic agents is sensitive to the amount of surfactant employed in conjunction with those therapeutic agents. This is particularly true when the pharmaceutical composition of the present invention is a single-dose or multi-dose aqueous ophthalmic composition. As a class, it is believed that the therapeutic agent classified herein as prostaglandins exhibit a higher degree of bioavailability when employed in

conjunction with relatively low concentrations of surfactants classified herein as vegetable oil surfactants. It has been discovered that travoprost in particular exhibits a higher degree of bioavailability in an aqueous ophthalmic composition of the present invention when used in conjunction with a relatively low concentration of an ethoxylated and/or hydrogenated vegetable oil surfactant such as Polyoxyethylene (POE) (40) Hydrogenated castor oil (or PEG (40) Hydrogenated castor oil) (HCO-40), POE (60) Hydrogenated castor oil (HCO-60), and POE (200) Hydrogenated castor oil (HCO-200), combinations thereof or the like. As such, it is contemplated that the surfactant of the present invention can be entirely or substantially entirely one or more ethoxylated and/or hydrogenated vegetable oils such as Polyoxyethylene (POE) (40) Hydrogenated castor oil (or PEG (40) Hydrogenated castor oil) (HCO-40), POE (60) Hydrogenated castor oil (HCO-60), and POE (200) Hydrogenated castor oil (HCO-200), combinations thereof or the like and the therapeutic agent can be entirely or substantially entirely one or more prostaglandins such as latanoprost, travoprost, unoprostone or combinations thereof.

[0044] In such an aqueous ophthalmic composition having prostaglandin therapeutic agent (e.g., travoprost) and hydrogenated and/or ethoxylated vegetable oil surfactant (e.g., HCO-40), the amount of such therapeutic agent is at least 0.00001 w/v %, at least 0.0001% w/v % or even at least 0.001 w/v % of the composition. Moreover, such composition includes less than 5 w/v %, more typically less than 0.05% w/v % and still more typically less than 0.01 w/v % such therapeutic agent. Further, the composition includes at least 0.005 w/v %, at least 0.01 w/v % or even at least 0.03 w/v % such surfactant. The composition also includes less than 0.3 w/v % and even possibly less than 0.15 w/v % such surfactant.

[0045] It is contemplated that area under the concentration/time curve measurements as described above can specifically be performed for such prostaglandin/surfactant combinations. The ophthalmic composition having prostaglandin therapeutic agent and hydrogenated and/or ethoxylated vegetable oil surfactant preferably includes an effectively low amount of surfactant such that the area under the concentration/time curve when determined for the ophthalmic composition of the present invention as for the aqueous humor of an eye is at least 130%, more typically at least 150% and even possibly at least 200% relative to the area under the concentration/time curve when determined for the control composition as for the aqueous humor of an eye. In such embodiment, the control composition is substantially identical to the pharmaceutical composition with the exception that the concentration of surfactant is at least doubled, more preferably tripled and even more typically quadrupled in the control composition relative to the ophthalmic composition of the present invention. Moreover, the ophthalmic composition of the present invention is applied in an amount that is equivalent to the amount of control composition applied. It is further contemplated that such concentrations can be similarly taken for the iris-ciliary body.

[0046] As suggested previously, concentrations for the composition of the present invention and the control composition can be carried out, for at least one embodiment of the present invention, according to the testing protocol provided below.

[0047] The pharmaceutical compositions of the present invention will generally be formulated as sterile aqueous solutions. These compositions are also formulated so as to be compatible with the eye to be treated with the compositions. The ophthalmic compositions intended for direct application to the eye will typically be formulated so as to have a pH and tonicity that are compatible with the eye. It is also contemplated that the compositions can be suspensions or other types of solutions. The ophthalmic compositions will typically have a pH in the range of 4 to 9, preferably 5.5 to 8.5, and most preferably 5.5 to 8.0. Particularly desired pH ranges are 6.0 to 7.8 and more specifically 6.4 to 7.2 or 7.5.

[0048] The pharmaceutical compositions of the present invention can include a polymer or a viscosity agent that further enhances bioavailability by extending the retention time of the compositions in or on the tear film of the eye, the cull-de-sac of the eye or elsewhere on the eye or another biological target. The preferred polymers include, without limitation, hydroxyethyl cellulose, hydroxypropylmethyl cellulose, carbomer, carbopol, xanthan gum, any combination thereof or the like.

[0049] In one embodiment, the pharmaceutical composition of the present invention is an ophthalmic aqueous solution. Such a solution will typically include a preservative system. As used herein, a "preservative system" is one or a group of ingredients included within the ophthalmic solution for maintaining antimicrobial activity in the solution. A solution may be a self-preserving solution if its ingredients naturally provide for anti-microbial activity and such solution still includes a preservative system. Examples of ophthalmic solutions or ingredients suitable for such solutions where those solutions may benefit from the teaching of the present invention are disclosed in U.S. Patent Nos. 3,931,319; 4,027,020; 4,407,791; 4,525,346; 4,836,986; 5,037,647; 5,300,287; 5,817,277; 6,503,497; 5,741,817; 6,319,464; 6,348,190; 6,348,190; 6,482,799; 5,320,843; 5,221,664; 6,034,043; 4,522,806; 6,017,861 and U.S. Patent Publications: 2002/0122831; and PCT application WO 91/09523 (Dziabo et al.); and JP 2003-104870, all of which are incorporated herein by reference for all purposes. One particular Ophthalmic Solution which may benefit from the teachings of the present invention is disclosed in U.S. Patent Application attorney docket no. 3464, titled "Aqueous Pharmaceutical Compositions Containing Borate-Polyol Complexes", filed on the same date as the present invention and incorporated herein by reference for all purposes.

[0050] Accordingly, the preservative system of the pharmaceutical composition of the present invention can include a borate. As used herein, the term "borate" shall refer to boric acid, salts of boric acid and other pharmaceutically

acceptable borates, or combinations thereof. Most suitable are: boric acid, sodium borate, potassium borate, calcium borate, magnesium borate, manganese borate, and other such borate salts. Borate interacts with polyols, such as glycerol, propylene glycol, sorbitol and mannitol, to form borate polyol complexes. The type and ratio of such complexes depends on the number of OH groups of a polyol on adjacent carbon atoms that are not in *trans* configuration relative to each other. It shall be understood that weight/volume percentages of the ingredients polyol and borate include those amounts whether as part of a complex or not.

[0051] When used, borate is generally used in the composition of the present invention in an amount that is greater than about 0.001 w/v %, more typically greater than about 0.01 w/v % and even more typically greater than about 0.07 w/v % of the pharmaceutical composition. Moreover, when used, the borate is generally used in the compositions of the present invention in an amount that is less than about 5 w/v %, more typically less than about 1.2 w/v % and even more typically less than about 0.8 w/v % of the pharmaceutical composition.

[0052] The preservative system of the pharmaceutical composition may also include one or more polyols. As used herein, the term "polyol" includes any compound having at least one hydroxyl group on each of two adjacent carbon atoms that are not in *trans* configuration relative to each other. The polyols can be linear or cyclic, substituted or unsubstituted, or mixtures thereof, so long as the resultant complex is water soluble and pharmaceutically acceptable. Examples of such compounds include: sugars, sugar alcohols, sugar acids and uronic acids. Preferred polyols are sugars, sugar alcohols and sugar acids, including, but not limited to: mannitol, glycerin, xylitol, sorbitol and propylene glycol.

[0053] When used, polyol generally used in the composition of the present invention in an amount that is greater than about 0.001 w/v %, more typically greater than about 0.01 w/v % and even more typically greater than about 0.07 w/v % of the pharmaceutical composition. Moreover, when used, polyol is generally used in the compositions of the present invention in an amount that is less than about 5 w/v %, more typically less than about 1.2 w/v % and even more typically less than about 0.8 w/v % of the pharmaceutical composition.

[0054] The compositions of the present invention can include a preservative. Potential preservatives include, without limitation, hydrogen peroxide, chlorine containing preservatives, or others. According to a preferred aspect, however, the ophthalmic composition of the present invention is substantially free of any chloride containing preservatives. Particularly, the composition is free of benzalkonium chloride. Highly preferred preservatives included for ophthalmic uses are polymeric quaternary ammonium compounds. It is noted that use of the amounts of surfactant specified herein can increase bioavailability in a manner that can at least partially or substantially entirely offset losses in bioavailability that may occur when benzalkonium chloride or other such ingredient are not present.

[0055] As used herein, the phrase "substantially free of" as it refers to an ingredient of the ophthalmic composition means that the ophthalmic solution is either entirely devoid of that particular ingredient or includes only a nominal amount of that particular ingredient.

[0056] The polymeric quaternary ammonium compounds useful in the compositions of the present invention are those which have an antimicrobial effect and which are ophthalmically acceptable. Preferred compounds of this type are described in U.S. Pat. Nos. 3,931,319; 4,027,020; 4,407,791; 4,525,346; 4,836,986; 5,037,647 and 5,300,287; and PCT application WO 91/09523 (Dziabo et al.). The most preferred polymeric ammonium compound is polyquaternium 1, otherwise known as POLYQUAD.RTM. or ONAMERM.RTM. with a number average molecular weight between 2,000 to 30,000. Preferably, the number average molecular weight is between 3,000 to 14,000.

[0057] When used, the polymeric quaternary ammonium compounds or other preservatives are generally used in the compositions of the present invention in an amount that is greater than about 0.00001 w/v %, more typically greater than about 0.0003 w/v % and even more typically greater than about 0.0007 w/v % of the pharmaceutical composition. Moreover, when used, the polymeric quaternary ammonium compounds or other preservatives are generally used in the compositions of the present invention in an amount that is less than about 3 w/v %, more typically less than about 0.003 w/v % and even more typically less than about 0.0015 w/v % of the pharmaceutical composition.

[0058] The pharmaceutical compositions of the present invention can be a multi-dose ophthalmic compositions having sufficient antimicrobial activity to allow the composition to satisfy the USP preservative efficacy requirements, as well as other preservative efficacy standards for aqueous pharmaceutical compositions.

[0059] The preservative efficacy standards for multi-dose ophthalmic solutions in the U.S. and other countries/regions are set forth in the following table:

Preservative Efficacy Test ("PET") Criteria (

Log Order Reduction of Microbial Inoculum Over Time

	Bacteria	Fungi
USP 27	A reduction of 1 log (90%), by day 7; 3 logs (99.9%) by day 14; and no increase after day 14	The compositions must demonstrate over the entire test period, which means no increases of 0.5 logs or greater, relative to the initial inoculum

(continued)

<u>Log Order Reduction of Microbial Inoculum Over Time</u>		
	Bacteria	Fungi
Japan	3 logs by 14 days; and no increase from day 14 through day 28	No increase from initial count at 14 and 28 days
Ph. Eur. A ¹	A reduction of 2 logs (99%) by 6 hours; 3 logs by 24 hours; and no recovery after 28 days	A reduction of 2 logs (99%) by 7 days, and no increase thereafter
Ph. Eur. B	A reduction of 1 log at 24 hours; 3 logs by day 7; and no increase thereafter	A reduction of 1 log (90%) by day 14, and no increase thereafter
FDA/ISO 14730	A reduction of 3 logs from initial challenge at day 14; and a reduction of 3 logs from rechallenge	No increase higher than the initial value at day 14, and no increase higher than the day 14 rechallenge count through day 28
¹ There are two preservative efficacy standards in the European Pharmacopoeia ' "A" and "B".		

[0060] The standards identified above for the USP 27 are substantially identical to the requirements set forth in prior editions of the USP, particularly USP 24, USP 25 and USP 26.

[0061] Table A below provides a listing of exemplary ingredients suitable for an exemplary preferred formulation of the ophthalmic composition of the present invention and a desired percentage for those ingredients.

TABLE A

<u>Ingredient</u>	<u>w/v Percent</u>
Travoprost	0.004
Polyoxyl 40 Hydrogenated Castor Oil (HCO-40)	≤ 0.2 or ≥ 0.05
Boric Acid	0.3
Zinc Chloride	0.0025
Sorbitol	0.25
Propylene Glycol	1.6
Sodium Chloride	0.35
NaOH	sufficient to achieve pH = 6.8
purified water	Q.S. 100

[0062] It is understood that the weight/volume percents in table A can be varied by $\pm 10\%$, $\pm 20\%$, $\pm 30\%$, $\pm 90\%$ of those weight/volume percents or more and that those variances can be specifically used to create ranges for the ingredients of the present invention. For example, an ingredient weight/volume percent of 10% with a variance of $\pm 20\%$ means that the ingredient can have a weight/volume percentage range of 8% to 12 w/v %.

[0063] With reference to Fig. 1, experiments were performed for several pharmaceutical compositions varying the amount of surfactant in those compositions. In particular, the amount or percentage of free or available travoprost was determined for compositions having the same ingredients as the compositions of table E with the exception that surfactant level were varied. As can be seen, the concentration of surfactant can have a significant effect upon the availability of the travoprost.

[0064] Notably, these preservative systems can at least assist in providing stability to the compositions of the present invention. Such systems can, in many instances, offset stability that may be lost through the use of less surfactant.

[0065] Applicants specifically incorporate the entire contents of all cited references in this disclosure. Further, when an amount, concentration, or other value or parameter is given as either a range, preferred range, or a list of upper preferable values and lower preferable values, this is to be understood as specifically disclosing all ranges formed from any pair of any upper range limit or preferred value and any lower range limit or preferred value, regardless of whether ranges are separately disclosed. Where a range of numerical values is recited herein, unless otherwise stated, the range is intended to include the endpoints thereof, and all integers and fractions within the range. It is not intended that the

scope of the invention be limited to the specific values recited when defining a range.

[0066] As suggested above, the relatively low concentrations of surfactant can be employed in formulations having multiple different therapeutic agents for producing desired bioavailability of those agents. Additionally, however, it has been found, that composition with particular combinations of therapeutic (combination compositions) agents gain significant benefits from the surfactant of the present invention, particularly when it is used at lower concentrations, but also potentially when it is used at higher concentrations.

[0067] In particular, it has been found that the surfactant that, when used at low concentrations, assists in solubilizing and/or increases bioavailability of the particular therapeutic agents discussed herein (prostaglandins such as travoprost) and can also act as a wetting agent for other therapeutic agents such as carbonic anhydrase inhibitors such as brinzolamide. Thus, the surfactant can provide a dual function within the composition. This dual function can then allow the composition to include only a single surfactant. Of course, multiple surfactants may be employed unless otherwise specifically stated.

[0068] It is contemplated that any of the preferred surfactants discussed herein as being capable of enhancing the bioavailability of a therapeutic agent and/or having the properties that provide such capability may be used to perform this dual function. Thus, a preferred surfactant for combination composition, like the preferred surfactant for the rest of the compositions herein, is a non-ionic vegetable oil that has been hydrogenated, ethoxylated or both. Such surfactants include, but are not limited to, babassu oil, almond oil, maize oil, palm kernel oil, castor oil, coconut oil, cotton seed oil, jojoba oil, linseed oil, mustard oil, olive oil, peanut oil, safflower oil, sesame oil, soybean oil, sunflower-seed oil and wheat germ oil, their hydrogenated or ethoxylated derivatives or combinations thereof. Preferred oils are castor oil, babassu oil, almond oil, maize oil and palm kernel oil, most preferably castor oil and cababassu oil.

[0069] Particularly preferred surfactants include Polyoxyethylene (POE) (40) Hydrogenated castor oil (or PEG (40 Hydrogenated castor oil) (HCO-40), POE (60) Hydrogenated castor oil (HCO-60), and POE (200) Hydrogenated castor oil (HCO-200).

[0070] Where higher bioavailability is desired, the surfactant for the combination composition can be used at the low concentrations that have already been disclosed herein. However, it is also contemplated that higher concentrations of such surfactant may be employed in circumstances where bioavailability is of less concern.

[0071] The combination composition benefits from particular advantages when it is formulated as a suspension. The use of the surfactant as discussed for the combination composition can allow for substantially complete (i.e., at least 80 or 90%) or complete solubilization of one therapeutic agent (a prostaglandin such as travoprost). At the same time, the surfactant can provide wetting and/or stabilization to the other therapeutic agent (e.g., a carbonic anhydrase inhibitor such as brinzolamide) which is suspended within the suspension. When the surfactant is used at the low concentrations discussed herein, it can also significantly improve the bioavailability of the non-suspended agent. Further, the preferred surfactant can often prevent the solubilized therapeutic agent from binding to container walls as would be seen if other surfactants were employed.

[0072] It is to be understood that any of the excipients discussed herein can be used in these combination compositions whether the compositions are suspensions, other aqueous solutions or other compositions. Moreover, the skilled artisan will understand that the discussions of the therapeutic agents and surfactants, when discussing any of the compositions herein, also applies to the combination compositions.

[0073] Other embodiments of the present invention will be apparent to those skilled in the art from consideration of the present specification and practice of the present invention disclosed herein.

COMPARATIVE EXAMPLES

[0074] Table B below provides two compositions, which include a preferred ophthalmic composition of the present invention as well as a control composition. The control composition has a surfactant level higher than the preferred composition according to the parameters discussed above.

TABLE B

INGREDIENTS	CONTROL	PREFERRED COMPOSITION
Travoprost	0.004	0.004
Polyoxyethylene 40 Hydrogenated Castor Oil (HCO-40)	0.5	0.1
Boric Acid	0.3	0.3
Zinc Chloride	0.0025	0.0025
Sorbitol	0.25	0.25
Propylene Glycol	1.6	1.6

EP 2 265 251 B1

(continued)

INGREDIENTS	CONTROL	PREFERRED COMPOSITION
Sodium Hydroxide and/or Hydrochloric Acid	Adjust pH 6.0	Adjust pH 6.0
Purified Water	QS 100 w/v%	QS 100 w/v%

[0075] The amount of surfactant (i.e., HCO-40) in the control composition is 5 times the amount of surfactant in the preferred composition. The preferred composition was applied to the eyes of a first set of rabbits and then a separate application of the control composition was applied to the eyes of a second set of rabbits. After application of the preferred composition and after application of the control composition, concentrations in nanograms per milliliter (ng/mL) of therapeutic agent (i.e., travoprost free acid) were determined for the Aqueous Humor and the Iris Ciliary of the eyes at various times. The results are respectively shown below in TABLE C and TABLE D.

TABLE C

Time	Control	Standard Deviation	Preferred Composition	Standard Deviation
0.25	1.13	0.37	2.01	0.87
0.5	3.82	1.31	9.17	2.72
1	6.87	2.40	20.78	4.35
2	5.90	1.17	18.40	6.08
4	5.05	2.78	10.24	3.69
6	0.57	0.15	1.20	0.77

TABLE D

Time	Control	Standard Deviation	Preferred Composition	Standard Deviation
0.25	1.40	0.29	1.71	0.42
0.5	2.43	0.94	5.86	1.88
1	3.33	0.76	9.32	2.68
2	2.79	0.39	7.59	3.25
4	1.98	1.14	4.69	1.59
6	BLQ	--	0.75	0.26

[0076] As can be seen from Tables B through D, the amount of therapeutic agent in the tested portion of the eyes was substantially higher in the preferred composition relative to the control composition.

[0077] Table E below shows further preferred compositions A, B and C of the present invention. After application of these preferred compositions in the eyes of rabbits, concentrations (in nanograms per milliliter (ng/mL) of therapeutic agent (i.e., travoprost free acid) were determined for the Aqueous Humor of the eyes at various times.

TABLE E

Ingredients	Composition A	Composition B	Composition C
Travoprost	0.004	0.004	0.002
Polyoxyethylene 40 Hydrogenated Castor Oil (HCO-40)	0.1	0.03	0.1
Boric Acid	0.3	0.3	0.3
Mannitol	0.3	0.3	0.3
Propylene Glycol	0.75	0.75	0.75
Sodium chloride	0.35	0.35	0.35
Polyquaternium-1	0.001	0.001	0.001
Hydrochloric Acid and/or Sodium HCl	Adj pH to 6.8	Adj pH to 6.8	Adj pH to 6.8
Purified Water	QS to 100 %	QS to 100 %	QS to 100 %

EP 2 265 251 B1

[0078] Table F below shows the concentrations in nanograms per milliliter (ng/mL) of the therapeutic agent of compositions A, B and C in the Aqueous humor of eyes of rabbits at various times after the application of the compositions.

TABLE F

Sampling Time (n)	Composition A	Composition B	Composition C
60 minutes (9-10)	22.79 ± 7.62	25.64 ± 9.34	13.21 ± 7.79
120 minutes (10)	15.46 ± 6.73	23.86 ± 4.67	9.23 ± 4.02
240 minutes (10)	4.73 ± 1.23	12.68 ± 9.81	2.59 ± 1.46

[0079] As can be seen, concentrations of the therapeutic agent from Compositions A through C in the Aqueous humor of the rabbits are desirably high.

[0080] Table G below shows a further preferred composition D of the present invention.

TABLE G

INGREDIENT	COMPOSITION D
Travoprost	0.004
Timolol Maleate	0.68
Polyoxyl 40 Hydrogenated Castor Oil (HCO-40)	0.1
Boric Acid	0.3
Mannitol	0.3
Propylene Glycol	0.75
Sodium Chloride	0.25
Polyquaternium-1	0.001
Sodium Hydroxide	Adj. pH 6.8
Hydrochloric Acid	Adj. pH 6.8
Purified Water	QS 100 %

[0081] As discussed above, the amount of surfactant can affect the stability of the compositions. Table H shows results of stability testing of composition B through E relative to Composition A, which has a higher surfactant concentration. All formulations A through E were formulated like the compositions of table E and contained 0.3% boric acid, 0.3% mannitol, 0.001% Polyquaternium-1 and were adjusted to pH 6.8 with sodium hydroxide and or hydrochloric acid. Composition A and composition each contained 0.66% sodium chloride while compositions C through D included 0.75% propylene glycol and 0.35% sodium chloride instead.

[0082] Each of the compositions A-E was packaged utilizing a syndiotactic polypropylene container, polypropylene plug and a polypropylene cap. They were stored at a stress condition of 55°C for 8 weeks. Control samples were stored in a refrigerator at 4 °C. These samples were analyzed for travoprost assay and the travoprost degradation product (travoprost free acid). The results are provided in Table. The formulations loose about 6 to 8% moisture upon 8 weeks storage at 55°C. As a result they show increase in travoprost assay at 8 weeks 55°C.

EP 2 265 251 B1

TABLE H

	Travoprost Concentration w/v%	HCO-40 Concentration w/v%	Travoprost Assay at 4C (% Label)	Travoprost Assay at 8 Wks/55C (% Label)	Travoprost free acid Assay at 4C (% of Travoprost)	Travoprost free acid (% of Travoprost)
A	0.004	0.5	96 96	102 101	0.0 0.0	0.7 0.7
B	0.004	0.1	99 99	104 103	0.0 0.0	2.4 2.4
C	0.004	0.1	97 97	101 102	NT	2.7 2.7
D	0.004	0.05	93 95	100 100	0.3 0.3	3.1 3.1
E	0.004	0.03	91 92	93 96	0.0 0.0	3.4 3.4

[0083] As can be seen, the amount of the degradation product increases with decreasing HCO-40 concentration. However, such increase is relatively insignificant for the overall compositions since the amount of degradation product is still quite small compare to the total amount of travoprost that remains in solution.

Bioavailability Examples

[0084] Polyoxyl hydrogenated castor oil 40 (HCO-40) is a surfactant that solubilizes travoprost and incorporates travoprost into its micelles. HCO-40 can be obtained by reacting 40 to 45 moles of ethylene oxide with 1 mole of hydrogenated castor oil. Thus, it typically has a high enough molecular weight such that it will not be filtered through a 3000 molecular weight cut off filter. An estimate of free fraction of travoprost, (i.e., fraction travoprost outside the HCO-40 micelle), can be obtained by filtering the solution through a 3000 molecular weight cut off filter.

[0085] The following procedure was used to determine the free fraction of travoprost. About 2 ml of formulation was placed in a centrifuge tube fitted with a 3000 Molecular weight cut off regenerated cellulose filter. The sample was centrifuged for approximately 90 minutes at 2000 r.p.m. to pass approximately 1 to 1.2 mL of formulation through the filter. The filtrate and retentate were then collected and assayed for travoprost using a HPLC procedure. The results are provided in Table I below. Note that the compositions 1 through 15 are all formulated like those in Table E and each composition contains 0.3% boric acid, 0.3% mannitol, 0.001% Polyquaternium-1 and were adjusted to pH 6.8 with sodium hydroxide and/or hydrochloric acid.

[0086] The assay values of filtrate represent the free fraction of travoprost. The results show that free fraction of travoprost increases with decreasing HCO-40 concentration. It is desired to have free fraction of travoprost greater than 1%, preferably greater than 2% and most preferably greater than 4%. The topical ocular bioavailability is expected to increase with increasing travoprost concentration and free fraction of travoprost.

TABLE I

#	Other Excipients	Travoprost Concentration	HCO-40 Concentration	Travoprost Assay of Filtrate, % Label	Travoprost Assay of Retentate, % Label	Weight of Filtrate	Weight of Retentate
1	0.66% NaCl	0.004	0.5	0.8	176	1.00	1.00
2	0.66% NaCl	0.004	0.25	1.7	188	1.01	0.99
3	0.66% NaCl	0.004	0.25	1.6	210	1.12	0.86
4	0.66% NaCl	0.002	0.1	3.6	222	1.19	0.80

EP 2 265 251 B1

(continued)

#	Other Excipients	Travoprost Concentration	HCO-40 Concentration	Travoprost Assay of Filtrate, % Label	Travoprost Assay of Retentate, % Label	Weight of Filtrate	Weight of Retentate
5	0.75% propylene glycol,, 0.35% NaCl	0.002	0.1	3.7	199	1.11	0.89
6	0.75% propylene glycol,, 0.35% NaCl	0.004	0.1	5.2	226	1.18	0.81
7	0.75% propylene glycol,, 0.35% NaCl	0.004	0.1	4.8	197	1.00	0.99
8	0.75% propylene glycol,, 0.35% NaCl	0.004	0.1	5.0	205	1.13	0.88
9	0.66% NaCl	0.004	0.1	4.4	216	1.13	0.87
10	0.75% propylene glycol,, 0.35% NaCl	0.001	0.05	8.7	187	1.03	0.99
11	0.75% propylene glycol,, 0.35% NaCl	0.002	0.05	9.0	214	1.18	0.81
12	0.75% propylene glycol,, 0.35% NaCl	0.004	0.05	9.8	196	1.14	0.85
13	0.75% propylene glycol,, 0.35% NaCl	0.004	0.05	12.5	217	1.25	0.76
14	0.75% propylene glycol,, 0.35% NaCl	0.001	0.03	13.1	219	1.28	0.73
15	0.75% propylene glycol,, 0.35% NaCl	0.004	0.03	15.6	166	1.15	0.86
16	0.75% propylene glycol,, 0.35% NaCl	0.004	0.03	15.4	201	1.26	0.75

[0087] Thus, in general for the present invention, half (1 ml) of a volumetric amount (2 ml) of composition of the present invention is forced through a molecular weight cut-off filter to form half of the composition into a filtrate and half into a retentate. For such composition, it is typically desirable for the concentration of surfactant in the composition to be such that the concentration of therapeutic agent is relatively high in the filtrate. This can be quantified as a filtrate/retentate ratio, which is equal to the weight/volume concentration of therapeutic agent in the filtrate divided by the weight/volume concentration of therapeutic agent in the retentate. This ratio for the present invention is determined when half (1 ml) of an amount (2 ml) of composition of the present invention is forced through a molecular weight cut-off filter to form a filtrate and a retentate where the molecular weight cut-off filter does not allow any or any substantial portion of the surfactant to flow through the weight cut-off filter. For the present invention, this ratio is typically greater than 0.0060, more typically greater than 0.014 and even possibly greater than 0.035.

Concentration Testing Protocol

[0088] The following protocol is at least one method of determining concentration of therapeutic agent at a biologic target. New Zealand white rabbits each receive a single 30 μ l topical ocular dose to each eye. Aqueous humor samples are collected immediately after euthanasia at the time points provided in results Table C. The concentration of travoprost free acid is then determined using tandem mass spectrometry (LC-MS/MS) analysis. Travoprost free acid is extracted from the aqueous humor samples and reconstituted in a water ethanol mixture. LC-MS-MS analysis for the determination of travoprost free acid concentrations is carried out using a Perkin Elmer Sciex AP 3, atmospheric pressure ionization mass spectrometer with an electrospray inlet and turbo ion spray, in the negative ion mode. Phenomenex ODS C18 (2) HPLC column is used. 5 mM ammonium formate buffer pH 6.3: methanol (30:70) is used as mobile phase.

COMBINATION COMPOSITION EXAMPLES

[0089] Table J below provides two exemplary preferred formulations (formulations R and S) of combination compositions, which are aqueous suspensions. The suspensions include two therapeutic agents, the carbonic anhydrase inhibitor brinzolamide and the prostaglandin travoprost. The compositions also include low concentrations of an exemplary preferred surfactant, HCO-40.

TABLE J

Composition W/V	Composition R	Composition S
Brinzolamide	1.0 g	1.0 g
Travoprost	0.0015 g	0.004 g
HCO-40	0.2 g	0.2 g
Carbomer 974 P	0.4 g	0.4 g
Mannitol	3.5 g	2.6 g
Edetate disodium	0.01 g	0.01 g
Benzalkonium Chloride	0.01 g	0.015 g
Sodium chloride	0.18 g	0.35 g
NaOH/HCl	q.s. to 6.5	q.s. to 6.5
Purified water	q.s. 100 ml	q.s. 100 ml

[0090] For purposes of comparison, table K below provides two formulations (formulations T and U) of combination compositions, which are also aqueous suspensions. The suspensions include two therapeutic agents, the carbonic anhydrase inhibitor brinzolamide and the prostaglandin travoprost. The compositions also include relatively higher concentrations of the surfactant HCO-40 and include tyloxapol as an additional surfactant.

TABLE K

Composition W/V	Composition T	Composition U
Brinzolamide	1.0 g + 3% excess	1.0 g + 3% excess
Tyloxapol	0.025 g	0.025 g

(continued)

Composition W/V	Composition T	Composition U
Travoprost	0.0015 g	0.004 g
HCO-40	0.5 g	0.5 g
Carbomer 974 P	0.4 g	0.4 g
Povidone K29-32	0.2 g	0.2 g
Mannitol	3.5 g	2.6 g
Edetate disodium	0.01 g	0.01 g
Benzalkonium Chloride	0.01 g	0.015 g
Sodium chloride	0.18 g	0.35 g
NaOH/HCl	q.s. to 6.5	q.s. to 6.5
Purified water	q.s. 100 ml	q.s. 100 ml

[0091] With reference to Fig. 3, it can be seen that the concentration of travoprost in the aqueous humor of rabbits was significantly greater for composition R than it was for composition T or for a composition that includes a same or similar level of a single therapeutic agent travoprost in combination with HCO-40 at a concentration of 0.5 w/v%. With reference to Fig. 4, it can be seen that the concentration of travoprost in the aqueous humor of rabbits was significantly greater for composition S than it was for composition U or the composition that includes a same or similar level of a single therapeutic agent (travoprost) in combination with HCO-40 at a concentration 0.5 w/v%. With reference to Fig. 5, it can be seen that the concentrations of brinzolamide in the aqueous humor of rabbits showed a difference that is considered small for composition R than it showed for composition T or a composition that includes a same or similar level of a single therapeutic agent brinzolamide in combination with HCO-40 at a concentration of 0.5 w/v%. With reference to Fig. 6, it can be seen that the concentrations of brinzolamide in the aqueous humor of rabbits showed a difference that is considered small for composition S than it showed for composition U or a composition that includes a same or similar level of a single therapeutic agent brinzolamide in combination with HCO-40 at a concentration of 0.5 w/v%.

Claims

1. An aqueous ophthalmic pharmaceutical composition, comprising:

a pharmaceutical vehicle suitable for topical application to an eye;
 an amount of therapeutic agent;
 an effectively low amount of surfactant; and
 polyquaternium-1 as a preservative;
 wherein the therapeutic agent includes prostaglandin therapeutic agent and the surfactant includes hydrogenated and/or ethoxylated vegetable oil surfactant;
 wherein the amount of prostaglandin therapeutic agent is at least 0.00001 w/v % but is less than 5 w/v % of the composition;
 wherein the effectively low amount of hydrogenated and/or ethoxylated vegetable oil surfactant is at least 0.005 w/v % but is less than 0.3 w/v % of the composition; and
 wherein the composition is free of benzalkonium chloride.

2. A composition as in claim 1 wherein the therapeutic agent is entirely or substantially entirely prostaglandin.

3. A composition as in any of claims 1-2 wherein the therapeutic agent, the surfactant or both are non-ionic.

4. A composition as in any of claims 1-3 wherein the therapeutic agent is travoprost.

5. A composition as in any of claims 1-4 wherein the surfactant is entirely or substantially entirely ethoxylated and/or hydrogenated vegetable oil.

6. A composition as in any of claims 1-5 wherein the surfactant is entirely or substantially entirely castor oil.
7. A composition as in any of claims 1-6 wherein the surfactant includes or is entirely or substantially entirely Polyoxyl 40 Hydrogenated castor Oil.
8. A composition as in any of claims 1-7 wherein the surfactant is less than 0.2 % or 0.15% by weight of the composition.
9. A composition as in any of claims 1-8 wherein the composition satisfies Ph. Eur. A, Ph. Eur. B or both.
10. A composition as in any of claims 1-9 wherein the therapeutic agent is less than 0.01 w/v % of the composition.
11. A composition as in any of claims 1-9 wherein the therapeutic agent is less than 0.006 w/v % of the composition.
12. A composition of any of claims 1-11 for use in a method comprising:
providing the composition to an eye of a human being.
13. A composition as in claim 12 wherein the composition is provided as drops from an eye dropper.

Patentansprüche

1. Wässrige ophthalmische pharmazeutische Zusammensetzung, umfassend:

ein pharmazeutisches Vehikel, das für die topische Verabreichung an ein Auge geeignet ist;
eine Menge eines therapeutischen Mittels;
eine wirksam geringe Menge eines oberflächenaktiven Mittels und
Polyquaternium-1 als ein Konservierungsmittel;
wobei das therapeutische Mittel das therapeutische Mittel Prostaglandin umfasst und das oberflächenaktive
Mittel das oberflächenaktive Mittel hydriertes und/oder ethoxyliertes pflanzliches Öl umfasst;
wobei die Menge des therapeutischen Mittels Prostaglandin mindestens 0,00001 % Gew./Vol., aber weniger
als 5 % Gew./Vol. der Zusammensetzung beträgt;
wobei die wirksam geringe Menge des oberflächenaktiven Mittels hydriertes und/oder ethoxyliertes pflanzliches
Öl mindestens 0,005 % Gew./Vol., aber weniger als 0,3 % Gew./Vol. der Zusammensetzung beträgt und
wobei die Zusammensetzung frei von Benzalkoniumchlorid ist.

2. Zusammensetzung nach Anspruch 1, wobei das therapeutische Mittel gänzlich oder im Wesentlichen gänzlich Prostaglandin ist.
3. Zusammensetzung nach einem der Ansprüche 1 - 2, wobei das therapeutische Mittel, das oberflächenaktive Mittel oder beide nicht ionisch sind.
4. Zusammensetzung nach einem der Ansprüche 1 - 3, wobei das therapeutische Mittel Travoprost ist.
5. Zusammensetzung nach einem der Ansprüche 1 - 4, wobei das oberflächenaktive Mittel gänzlich oder im Wesentlichen gänzlich ethoxyliertes und/oder hydriertes pflanzliches Öl ist.
6. Zusammensetzung nach einem der Ansprüche 1 - 5, wobei das oberflächenaktive Mittel gänzlich oder im Wesentlichen gänzlich Rizinusöl ist.
7. Zusammensetzung nach einem der Ansprüche 1 - 6, wobei das oberflächenaktive Mittel Polyoxyl-40-hydriertes Rizinusöl umfasst oder dieses gänzlich oder im Wesentlichen gänzlich ist.
8. Zusammensetzung nach einem der Ansprüche 1 - 7, wobei das oberflächenaktive Mittel weniger als 0,2 oder 0,15 Gew.-% der Zusammensetzung beträgt.
9. Zusammensetzung nach einem der Ansprüche 1 - 8, wobei die Zusammensetzung Ph. Eur. A, Ph. Eur. B oder beiden genügt.

10. Zusammensetzung nach einem der Ansprüche 1 - 9, wobei das therapeutische Mittel weniger als 0,01 % Gew./Vol. der Zusammensetzung beträgt.

11. Zusammensetzung nach einem der Ansprüche 1 - 9, wobei das therapeutische Mittel weniger als 0,006 % Gew./Vol. der Zusammensetzung beträgt.

12. Zusammensetzung nach einem der Ansprüche 1 - 11 zur Verwendung in einem Verfahren, umfassend:

das Abgeben der Zusammensetzung an ein Auge eines menschlichen Wesens.

13. Zusammensetzung nach Anspruch 12, wobei die Zusammensetzung als Tropfen aus einem Augentropfer abgegeben wird.

Revendications

1. Composition pharmaceutique ophtalmique aqueuse comprenant :

un véhicule pharmaceutique approprié pour une application topique au niveau d'un oeil ;
une quantité d'agent thérapeutique ;
une quantité efficacement faible de tensioactif ; et
du polyquaternium-1 en tant qu'agent de conservation ;
dans laquelle l'agent thérapeutique comprend un agent thérapeutique de prostaglandine et le tensioactif comprend un tensioactif d'huile végétale hydrogénée et/ou éthoxylée ;
dans laquelle la quantité d'agent thérapeutique de prostaglandine est d'au moins 0,00001 % m/v mais est inférieure à 5 % m/v de la composition ;
dans laquelle la quantité efficacement faible du tensioactif d'huile végétale hydrogénée et/ou éthoxylée est d'au moins 0,005 % m/v mais est inférieure à 0,3 % m/v de la composition ; et
dans laquelle la composition est exempte de chlorure de benzalconium.

2. Composition selon la revendication 1, dans laquelle l'agent thérapeutique est totalement ou presque totalement la prostaglandine.

3. Composition selon la revendication 1 ou 2, dans laquelle l'agent thérapeutique, le tensioactif ou les deux sont non ioniques.

4. Composition selon l'une quelconque des revendications 1 à 3, dans laquelle l'agent thérapeutique est le travoprost.

5. Composition selon l'une quelconque des revendications 1 à 4, dans laquelle le tensioactif est totalement ou presque totalement une huile végétale hydrogénée et/ou éthoxylée.

6. Composition selon l'une quelconque des revendications 1 à 5, dans laquelle le tensioactif est totalement ou presque totalement une huile de ricin.

7. Composition selon l'une quelconque des revendications 1 à 6, dans laquelle le tensioactif comprend ou est totalement ou presque totalement une huile de ricin polyoxyl 40 hydrogénée.

8. Composition selon l'une quelconque des revendications 1 à 7, dans laquelle le tensioactif est inférieur à 0,2 % ou 0,15 % en poids de la composition.

9. Composition selon l'une quelconque des revendications 1 à 8, dans laquelle la composition satisfait la Ph. Eur. A, la Ph. Eur. B ou les deux.

10. Composition selon l'une quelconque des revendications 1 à 9, dans laquelle l'agent thérapeutique est inférieur à 0,01 % m/v de la composition.

11. Composition selon l'une quelconque des revendications 1 à 9, dans laquelle l'agent thérapeutique est inférieur à 0,006 % m/v de la composition.

EP 2 265 251 B1

12. Composition selon l'une quelconque des revendications 1 à 11, pour une utilisation dans un procédé comprenant à :

administrer la composition au niveau d'un oeil d'un être humain.

5 **13.** Composition selon la revendication 12, dans laquelle la composition est administrée sous forme de gouttes à partir d'un compte-goutte oculaire.

10

15

20

25

30

35

40

45

50

55




FIG. 1




Fig. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61037117 A [0001]
- US 61111920 A [0001]
- US 6743439 B [0008]
- WO 0003736 A [0009]
- EP 1547599 A [0010]
- US 20040082660 A1 [0011]
- WO 9841208 A [0012]
- WO 9825620 A [0013]
- US 5932572 A [0014]
- WO 9819680 A [0015]
- WO 9823809 A [0016]
- US 3931319 A [0049] [0056]
- US 4027020 A [0049] [0056]
- US 4407791 A [0049] [0056]
- US 4525346 A [0049] [0056]
- US 4836986 A [0049] [0056]
- US 5037647 A [0049] [0056]
- US 5300287 A [0049] [0056]
- US 5817277 A [0049]
- US 6503497 B [0049]
- US 5741817 A [0049]
- US 6319464 B [0049]
- US 6348190 B [0049]
- US 6482799 B [0049]
- US 5320843 A [0049]
- US 5221664 A [0049]
- US 6034043 A [0049]
- US 4522806 A [0049]
- US 6017861 A [0049]
- US 20020122831 A [0049]
- WO 9109523 A, Dziabo [0049] [0056]
- JP 2003104870 A [0049]

Előnyös biológiai hozzáférhetőségű gyógyászati készítmények

Szabadalmi igénypontok

1. Vizes szemgyógyászati készítmény, mely:

egy szemhez topikálisan adagolható gyógyszerészetileg elfogadható hordozóanyagot;

meghatározott mennyiségű terápiás szert;

hatékonyan alacsony mennyiségű felületaktív szert; és

konzerválószerként polikvaternium-1-et tartalmaz,

ahol a terápiás szer prosztaglandint tartalmaz és a felületaktív szer hidrogénezett és/vagy etoxilezett növényi olaj felületaktív szert tartalmaz;

ahol a prosztaglandin terápiás szer mennyisége a készítményre számítva legalább 0,00001 tömeg/térfogat%, de kevesebb, mint 5 tömeg/térfogat%;

ahol a hatékonyan alacsony mennyiségű hidrogénezett és/vagy etoxilezett növényi olaj felületaktív szer mennyisége a készítményre számítva legalább 0,005 tömeg/térfogat%, de kevesebb, mint 0,3 tömeg/térfogat%; és

ahol a készítmény benzalkonium-kloridtól mentes.

2. Az 1. igénypont szerinti készítmény, ahol a terápiás szer kizárólag vagy lényegében kizárólag prosztaglandin.

3. Az 1. vagy 2. igénypont szerinti készítmény, ahol a terápiás szer, a felületaktív szer vagy mindkettő nem-ionos.

4. Az 1-3. igénypontok bármelyike szerinti készítmény, ahol a terápiás szer travoprost.

5. Az 1-4. igénypontok bármelyike szerinti készítmény, ahol a felületaktív szer kizárólag vagy lényegében kizárólag etoxilezett és/vagy hidrogénezett növényi olaj.

6. Az 1-5. igénypontok bármelyike szerinti készítmény, ahol a felületaktív szer kizárólag vagy lényegében kizárólag ricinusolaj.

7. Az 1-6. igénypontok bármelyike szerinti készítmény, ahol a felületaktív szer polioxil 40 hidrogénezett ricinusolajat tartalmaz, vagy kizárólag vagy lényegében kizárólag tartalmaz.

8. Az 1-7. igénypontok bármelyike szerinti készítmény, ahol a felületaktív szer mennyisége a készítményre számított 0,2 tömeg%-nál vagy 0,15 tömeg%-nál kevesebb.

9. Az 1-8. igénypontok bármelyike szerinti készítmény, ahol a készítmény megfelel a Ph. Eur. A., Ph. Eur. B. vagy mindkettő előírásainak.

10. Az 1-9. igénypontok bármelyike szerinti készítmény, ahol a terápiás szer a készítményben 0,01 tömeg/térfogat%-nál kisebb mennyiségben van jelen.

11. Az 1-9. igénypontok bármelyike szerinti készítmény, ahol a terápiás szer a készítményben 0,006 tömeg/térfogat%-nál kisebb mennyiségben van jelen.

12. Az 1-12. igénypontok bármelyike szerinti készítményegy eljárásban történő alkalmazására, melynek során a készítményt ember szemébe visszük be.

13. A 12. igénypont szerinti készítmény, ahol a készítményt cseppentve, szemcseppentőből adagoljuk.

Dr. Szentpéteri Ádám
szabadalmi ügyvivő
SOGK Szabadalmi Ügyvivői Iroda
H-1067 Budapest, Andrassy út 113.
Telefon: +361-1067223, +361-1067
Email: szentpeteri@sogk.hu