



(86) **Date de dépôt PCT/PCT Filing Date:** 2009/08/03
(87) **Date publication PCT/PCT Publication Date:** 2010/02/11
(45) **Date de délivrance/Issue Date:** 2019/02/26
(85) **Entrée phase nationale/National Entry:** 2011/02/01
(86) **N° demande PCT/PCT Application No.:** EP 2009/060052
(87) **N° publication PCT/PCT Publication No.:** 2010/015608
(30) **Priorité/Priority:** 2008/08/05 (US61/086,355)

(51) **Cl.Int./Int.Cl. C12N 15/13** (2006.01),
A61K 39/395 (2006.01), **C07K 16/18** (2006.01),
C07K 16/46 (2006.01)

(72) **Inventeurs/Inventors:**
DIEFENBACH-STREIBER, BEATE, DE;
EBERTH, ADINA, DE;
GUILD, BRAYDON CHARLES, US;
KIM, YONG-IN, US;
ROGUSKA, MICHAEL, US;
SPLAWSKI, IGOR, US

(73) **Propriétaire/Owner:**
NOVARTIS AG, CH

(74) **Agent:** FETHERSTONHAUGH & CO.

(54) **Titre : COMPOSITIONS ET PROCEDES POUR DES ANTICORPS CIBLANT UNE PROTEINE DU COMPLEMENT C5**
(54) **Title: COMPOSITIONS AND METHODS FOR ANTIBODIES TARGETING COMPLEMENT PROTEIN C5**

(57) **Abrégé/Abstract:**

The present invention relates to antibodies targeting complement protein C5 and compositions and methods of use thereof.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 February 2010 (11.02.2010)

PCT

(10) International Publication Number
WO 2010/015608 A1

(51) International Patent Classification:
A61K 39/395 (2006.01) *C07K 16/18* (2006.01)

(21) International Application Number:
PCT/EP2009/060052

(22) International Filing Date:
3 August 2009 (03.08.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/086,355 5 August 2008 (05.08.2008) US

(71) Applicant (for all designated States except US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DIEFENBACH-STREIBER, Beate** [DE/DE]; Reiherweg 6, 86949 Windach (DE). **EBERTH, Adina** [DE/DE]; Friedrich-List Strasse 114, 81377 München (DE). **GUILD, Braydon Charles** [US/US]; 109 Riverdale Road, Concord, Massachusetts 01742 (US). **KIM, Yong-In** [KR/US]; Novartis Institutes for BioMedical Research, Inc., 500 Technology Square, Cambridge, Massachusetts 02139 (US). **ROGUSKA, Michael** [US/US]; Novartis Institutes for BioMedical Research, Inc., 500 Technology Square, Cambridge, Massachusetts 02139 (US). **SPLAWSKI, Igor** [PL/US]; Novartis Institutes for BioMedical Research, Inc., 500 Technology Square, Cambridge, Massachusetts 02139 (US).

(74) Agent: **VOEGELI-LANGE, Regina**; Novartis Pharma Ag, Patent Department, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))

(54) Title: COMPOSITIONS AND METHODS FOR ANTIBODIES TARGETING COMPLEMENT PROTEIN C5

(57) Abstract: The present invention relates to antibodies targeting complement protein C5 and compositions and methods of use thereof.



WO 2010/015608 A1

COMPOSITIONS AND METHODS FOR ANTIBODIES TARGETING COMPLEMENT PROTEIN C5

1. INTRODUCTION

The present invention relates to antibodies targeting complement protein C5 and compositions and methods of use thereof.

2. BACKGROUND OF THE INVENTION

The normal role of complement, which is part of the innate immune system, is in host defense. Complement defends against bacterial infection, links adaptive and innate immunity, and disposes immune complexes and the products of inflammatory injury.

The defensive functions are accomplished by biologically active products generated in the course of complement activation, which opsonise infectious agents, promote inflammation or lyse susceptible targets (Marzari *et al.*, Eur J Immunol 32:2773-2782 (2002)). The complement system consists of about 25-30 plasma proteins which play a role in the immune system. The complement cascade is activated by at least three major pathways. The classical pathway is typically activated by immune-complexes, the alternative pathway can be activated by unprotected cell surfaces, and the mannose binding lectin (MBL) pathway is initiated by binding of MBL to cell surface carbohydrates (Trendelenburg, Swiss Med Wkly 137:413-417 (2007)).

All three pathways lead to the cleavage of C5 by the C5 convertase. The result of this cleavage is release of C5a fragment, a potent inflammatory molecule, and C5b which initiates the membrane attack complex (MAC). The complement products, once released, do not differentiate between foreign and self targets and, if not tightly regulated, often cause extensive damage of bystander cells and tissues in clinical conditions associated with unrestricted complement activation (Marzari *et al.*, 2002).

C5 is expressed intracellularly as a single pro-C5 peptide of 1676 amino acids that consist of an 18 residue signal sequence and an Arg-rich linker sequence (RPRR) situated between the mature N-terminal β chain and the C-terminal α chain. The mature C5 has a molecular weight of about 190 kDa, and consists of two polypeptide chains (α , 115 kDa and β , 75 kDa) which are connected by disulfide bonds. The C5 convertase cleaves C5 between residues 74 and 75 of the alpha chain to release the 74 amino acid C5a peptide and the C5b fragment which is subsequently incorporated into the membrane-attack complex (MAC).

Macular degeneration is a medical condition predominantly found in the elderly in which the center of the inner lining of the eye, known as the macula area of the retina, suffers thinning, atrophy, and in some cases, bleeding. This can result in loss of central vision, which entails inability to see fine details, to read, or to recognize faces. Pathogenesis of new

choroidal vessel formation is poorly understood, but factors such as inflammation, ischemia, and local production of angiogenic factors are thought to be important.

Despite current treatment options for treating diseases and disorders associated with the classical or alternative component pathways, particularly AMD, there remains a need for finding specific targets that lead to treatments which are effective and well-tolerated.

3. SUMMARY OF THE INVENTION

The present invention provides isolated complement C5-binding molecules (*e.g.*, C5-binding antibodies or antigen binding fragments thereof), pharmaceutical compositions comprising such molecules, methods of making such molecules and compositions, and methods of use thereof.

In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibody has an affinity constant (K_A) of at least $1 \times 10^7 \text{ M}^{-1}$, 10^8 M^{-1} , 10^9 M^{-1} , 10^{10} M^{-1} , or 10^{11} M^{-1} .

In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, and inhibit the alternative complete pathway as measured by in vitro hemolytic assay with an IC_{50} range from about 20pM to about 200 pM.

In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, and cross compete with an antibody described in Table 1 below. In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof that bind to the same epitope of C5 protein as an antibody described in Table 1 below.

In some embodiments, the antibodies of the invention are isolated monoclonal antibodies that specifically bind to a C5 protein. In some embodiments, the antibodies of the invention are isolated human or humanized monoclonal antibodies that specifically bind to a C5 protein. In some embodiments, the antibodies of the invention are isolated chimeric antibodies that specifically bind to a C5 protein. In some embodiments, the antibodies of the invention comprise a human heavy chain constant region and a human light chain constant region.

In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies are single chain antibodies. In some embodiments, the antibodies of the invention are Fab fragments. In some embodiments, the antibodies of the invention are scFv.

In some embodiments, the present invention provides isolated antibodies or antigen-binding fragments thereof that specifically bind to both human C5 and cynomolgus C5. In some embodiments, the antibodies of the invention are an IgG isotype.

In some embodiments, the present invention provides isolated antibodies or antigen binding fragments thereof comprising a framework in which amino acids have been substituted into the antibody framework from the respective human VH or VL germline sequences.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise at least one complementarity determining (CDR) sequence having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 1, 2, 3, 4, 5, 6, 17, 18, 19, 20, 21, 22, 33, 34, 35, 36, 37, 38, 49, 50, 61, 62, 63, 64, 65, 66, 77, 78, 89, 95, 101, 107, 113, 119, 120, 131, 132, 133, 134, 135, 136, 145, 146, 147, 148, 149, 150, 159, 160, 161, 162, 163, 164, 173, 174, 175, 176, 177, 178, 195, 196, 197, 198, 199, 200, 209, 226, 235, 236, 237, 238, 239, or 240.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise at least one heavy chain CDR sequence that is identical to SEQ ID NO: 1, 2, 3, 17, 18, 19, 33, 34, 35, 49, 61, 62, 63, 77, 77, 95, 107, 113, 119, 132, 131, 133, 145, 146, 147, 159, 160, 161, 173, 174, 175, 195, 196, 197, 226, 235, 236, or 237.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise at least one light chain CDR sequence that is identical to SEQ ID NO: 4, 5, 6, 20, 21, 22, 36, 37, 38, 50, 64, 65, 66, 78, 89, 101, 120, 134, 135, 136, 148, 149, 150, 162, 163, 164, 176, 177, 178, 198, 199, 200, 209, 238, 239, or 240.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a heavy chain CDR 1 selected from the group consisting SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235; a heavy chain CDR2 selected from the group consisting SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236; and a heavy chain CDR3 selected from the group consisting SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237. In some embodiments, such antibodies or antigen binding fragments thereof further comprise a light chain CDR1 selected from the group consisting SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238; a light chain CDR2 selected from the group consisting SEQ ID NOs: 5, 21, 37, 65, 135, 149,

163, 177, 199, and 239; and a light chain CDR3 selected from the group consisting SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a light chain CDR 1 selected from the group consisting SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238; a light chain CDR2 selected from the group consisting SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239; and a light chain CDR3 selected from the group consisting SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a heavy chain variable region having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281. In some embodiments, such antibodies or antigen binding fragments thereof further comprise a light chain variable region having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, and 289.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a light chain variable region having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, and 289.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a heavy chain having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 9, 25, 41, 53, 69, 81, 97, 109, 115, 123, 139, 153, 167, 181, 189, 203, 212, 220, 229, 243, 249, 254, 258, 274, 278, or 282. In some embodiments, such antibodies further comprise a light chain having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154, 168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, or 290.

In some embodiments, the present invention provides isolated monoclonal antibodies or antigen binding fragments thereof that specifically bind to a C5 protein, wherein said antibodies comprise a light chain having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154, 168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, or 290.

The present invention also comprises pharmaceutical compositions comprising one or more C5-binding molecules of the invention (e.g., C5 binding antibodies or antigen binding fragments thereof) and a pharmaceutically acceptable carrier.

In some embodiments, the present invention provides nucleic acids comprising a nucleotide sequence encoding a polypeptide comprising a heavy chain variable region having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281.

In some embodiments, the present invention provides nucleic acids comprising a nucleotide sequence encoding a polypeptide comprising a light chain variable region having at least 90%, 95%, 97%, 98% or at least 99% sequence identity to SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, and 289.

The present invention also provides vectors and host cells comprising such nucleic acids. In one embodiment, the present invention provides isolated host cells comprising (1) a recombinant DNA segment encoding a heavy chain of the antibodies of the invention, and (2) a second recombinant DNA segment encoding a light chain of the antibodies of the invention; wherein said DNA segments are respectively operably linked to a first and a second promoter, and are capable of being expressed in said host cell. In another embodiment, the present invention provides isolated host cells comprising a recombinant DNA segment encoding a heavy chain, and a light chain of the antibodies of the invention, respectively, wherein said DNA segment is operably linked to a promoter, and is capable of being expressed in said host cells. In some embodiments, the host cells are non-human mammalian cell line. In some embodiments, the antibodies or antigen binding fragments thereof are a human monoclonal antibody, or an antigen binding fragment thereof.

The present invention further provides treatment of diagnostic methods using the C5 binding molecules (e.g., C5 binding antibodies or antigen binding fragments thereof) of the invention. In one embodiment, the present invention provides methods of treating age related macular degeneration comprising administering to a subject in need thereof an effective amount of a composition comprising an antibody or an antigen binding fragment thereof of the invention.

In another embodiment, the present invention provides methods of treating a disease comprising administering to a subject in need thereof an effective amount of a composition comprising an antibody or an antigen binding fragment thereof of the invention, wherein said disease is asthma, arthritis, autoimmune heart disease, multiple sclerosis, inflammatory bowel disease, ischemia-reperfusion injuries, Barraquer-Simons Syndrome, hemodialysis,

systemic lupus, lupus erythematosus, psoriasis, multiple sclerosis, transplantation, Alzheimer's disease, glomerulonephritis, or MPGN II.

The present invention also provides methods of treating paroxysmal nocturnal hemoglobinuria (PNH) comprising administering to a subject in need thereof an effective
5 amount of a composition comprising an antibody or antigen binding fragment thereof of the invention.

The present invention further provides methods of ameliorating a symptom associated with extracorporeal circulation comprising administering to a subject in need thereof an effective amount of a composition comprising an antibody or antigen binding
10 fragment thereof of the invention.

In another embodiment, there is provided an isolated monoclonal human C5 protein antibody, or an antigen binding fragment thereof comprising: (i) a heavy chain CDR1 comprising the sequence of SEQ ID NO:1; a heavy chain CDR2 comprising the sequence of SEQ ID NO: 2; and a heavy chain CDR3 sequence comprising the sequence of SEQ ID
15 NO:3; and (ii) a light chain CDR1 comprising the sequence of SEQ ID NO:4; a light chain CDR2 comprising the sequence of SEQ ID NO:5; and a light chain CDR3 sequence comprising the sequence of SEQ ID NO: 6, wherein said antibody or antigen binding fragment thereof binds to human C5 protein with a K_D of 10^{-10} M or less as measured by surface plasmon resonance.

20 In another embodiment, there is provided a pharmaceutical composition comprising the antibody or antigen binding fragment as described herein and a pharmaceutically acceptable carrier.

In another embodiment, there is provided an isolated nucleic acid comprising a nucleotide sequence encoding the antibody or antigen binding fragment as described herein.

25 In another embodiment, there is provided use of an antibody as described herein for treating age related macular degeneration in a subject.

In another embodiment, there is provided use of an antibody as described herein for treating a disease, wherein said disease is hyperacute allograft rejection, xenograft rejection, asthma, arthritis, autoimmune heart disease, multiple sclerosis, inflammatory bowel disease,
30 ischemia-reperfusion injuries, Barraquer-Simons Syndrome, hemodialysis, systemic lupus, lupus erythematosus, psoriasis, multiple sclerosis, transplantation, Alzheimer's disease,

81538694

glomerulonephritis, MPGN II, wet age related macular degeneration, dry age related macular degeneration or geographic atrophy.

In another embodiment, there is provided use of an antibody as described herein for treating paroxysmal nocturnal hemoglobinuria (PNH).

5 3.1. Definitions

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which this invention pertains.

The term "antibody" as used herein includes whole antibodies and any antigen
 10 binding fragment (*i.e.*, "antigen-binding portion") or single chains thereof. A naturally occurring "antibody" is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3.
 15 Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each VH and VL is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1,
 20 CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (*e.g.*, effector cells) and the first component
 25 (C1q) of the classical complement system.

The term "antigen binding portion" of an antibody, as used herein, refers to one or more fragments of an intact antibody that retain the ability to specifically bind to a given antigen (*e.g.*, C5). Antigen binding functions of an antibody can be performed by fragments of an intact antibody. Examples of binding fragments encompassed within the term "antigen
 30 binding portion" of an antibody include a Fab fragment, a monovalent fragment consisting of

the VL, VH, CL and CH1 domains; a F(ab)₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; an Fd fragment consisting of the VH and CH1 domains; an Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a single domain antibody (dAb) fragment (Ward *et al.*, 1989 Nature 341:544-546), which consists of a VH domain; and an isolated complementarity determining region (CDR).

Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by an artificial peptide linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see, *e.g.*, Bird *et al.*, 1988 Science 242:423-426; and Huston *et al.*, 1988 Proc. Natl. Acad. Sci. 85:5879-5883). Such single chain antibodies include one or more "antigen binding portions" of an antibody. These antibody fragments are obtained using conventional techniques known to those of skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies.

Antigen binding portions can also be incorporated into single domain antibodies, maxibodies, minibodies, intrabodies, diabodies, triabodies, tetrabodies, v-NAR and bis-scFv (see, *e.g.*, Hollinger and Hudson, 2005, Nature Biotechnology, 23, 9, 1126-1136). Antigen binding portions of antibodies can be grafted into scaffolds based on polypeptides such as Fibronectin type III (Fn3) (see U.S. Pat. No. 6,703,199, which describes fibronectin polypeptide monobodies).

Antigen binding portions can be incorporated into single chain molecules comprising a pair of tandem Fv segments (VH-CH1-VH-CH1) which, together with complementary light chain polypeptides, form a pair of antigen binding regions (Zapata *et al.*, 1995 Protein Eng. 8(10):1057-1062; and U.S. Pat. No. 5,641,870).

As used herein, the term "Affinity" refers to the strength of interaction between antibody and antigen at single antigenic sites. Within each antigenic site, the variable region of the antibody "arm" interacts through weak non-covalent forces with antigen at numerous sites; the more interactions, the stronger the affinity.

As used herein, the term "Avidity" refers to an informative measure of the overall stability or strength of the antibody-antigen complex. It is controlled by three major factors: antibody epitope affinity; the valency of both the antigen and antibody; and the structural arrangement of the interacting parts. Ultimately these factors define the specificity of the antibody, that is, the likelihood that the particular antibody is binding to a precise antigen epitope.

The term "amino acid" refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, *e.g.*, hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refer to compounds that have the same basic chemical structure as a naturally occurring amino acid, *i.e.*, an alpha carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, *e.g.*, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (*e.g.*, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

The term "binding specificity" as used herein refers to the ability of an individual antibody combining site to react with only one antigenic determinant. The combining site of the antibody is located in the Fab portion of the molecule and is constructed from the hypervariable regions of the heavy and light chains. Binding affinity of an antibody is the strength of the reaction between a single antigenic determinant and a single combining site on the antibody. It is the sum of the attractive and repulsive forces operating between the antigenic determinant and the combining site of the antibody.

Specific binding between two entities means a binding with an equilibrium constant (K_A) of at least $1 \times 10^7 \text{ M}^{-1}$, 10^8 M^{-1} , 10^9 M^{-1} , 10^{10} M^{-1} , or 10^{11} M^{-1} . The phrase "specifically (or selectively) binds" to an antibody (*e.g.*, a C5-binding antibody) refers to a binding reaction that is determinative of the presence of a cognate antigen (*e.g.*, a human C5 or cynomolgus C5) in a heterogeneous population of proteins and other biologics. In addition to the equilibrium constant (K_A) noted above, an C5-binding antibody of the invention typically also has a dissociation rate constant (K_d) of about $1 \times 10^{-2} \text{ s}^{-1}$, $1 \times 10^{-3} \text{ s}^{-1}$, $1 \times 10^{-4} \text{ s}^{-1}$, $1 \times 10^{-4} \text{ s}^{-1}$, or lower, and binds to C5 with an affinity that is at least two-fold greater than its affinity for binding to a non-specific antigen (*e.g.*, C3, C4, BSA). The phrases "an antibody recognizing an antigen" and "an antibody specific for an antigen" are used interchangeably herein with the term "an antibody which binds specifically to an antigen".

The term "chimeric antibody" is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, *e.g.*, an enzyme, toxin, hormone, growth factor, drug, etc.; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a

different or altered antigen specificity. For example, a mouse antibody can be modified by replacing its constant region with the constant region from a human immunoglobulin. Due to the replacement with a human constant region, the chimric antibody can retain its specificity in recognizing the antigen while having reduced antigenicity in human as compared to the original mouse antibody.

The term "complement C5 protein" or "C5" are used interchangeably, and refers to the C5 protein in different species. For example, human C5 has the sequence as set in SEQ ID NO: 296, cynomolgus C5 has the sequence as set in SEQ ID NO: 297 (*Macaca fascicularis*) (see Table 1). Human C5 can be obtained from Quidel (Cat. Number A403). Cynomolgus C5 can be produced as illustrated in the Example section below.

The term "conservatively modified variant" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid that encodes a polypeptide is implicit in each described sequence.

For polypeptide sequences, "conservatively modified variants" include individual substitutions, deletions or additions to a polypeptide sequence which result in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention. The following eight groups contain amino acids that are conservative substitutions for one another: 1) Alanine (A), Glycine (G); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (K); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); 7) Serine (S), Threonine (T); and 8) Cysteine (C), Methionine (M) (see, e.g., Creighton, Proteins (1984)). In some embodiments, the term "conservative sequence

modifications" are used to refer to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody containing the amino acid sequence.

The terms "cross-block", "cross-blocked" and "cross-blocking" are used interchangeably herein to mean the ability of an antibody or other binding agent to interfere with the binding of other antibodies or binding agents to C5 in a standard competitive binding assay.

The ability or extent to which an antibody or other binding agent is able to interfere with the binding of another antibody or binding molecule to C5, and therefore whether it can be said to cross-block according to the invention, can be determined using standard competition binding assays. One suitable assay involves the use of the Biacore technology (e.g. by using the BIAcore 3000 instrument (Biacore, Uppsala, Sweden)), which can measure the extent of interactions using surface plasmon resonance technology. Another assay for measuring cross-blocking uses an ELISA-based approach.

The term "epitope" means a protein determinant capable of specific binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

As used herein, the term "high affinity" for an IgG antibody refers to an antibody having a K_D of 10^{-8} M or less, 10^{-9} M or less, or 10^{-10} M, or 10^{-11} M or less for a target antigen. However, "high affinity" binding can vary for other antibody isotypes. For example, "high affinity" binding for an IgM isotype refers to an antibody having a K_D of 10^{-7} M or less, or 10^{-8} M or less.

The term "human antibody", as used herein, is intended to include antibodies having variable regions in which both the framework and CDR regions are derived from sequences of human origin. Furthermore, if the antibody contains a constant region, the constant region also is derived from such human sequences, e.g., human germline sequences, or mutated versions of human germline sequences. The human antibodies of the invention may include amino acid residues not encoded by human sequences (e.g., mutations introduced by random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*).

The term "human monoclonal antibody" refers to antibodies displaying a single binding specificity which have variable regions in which both the framework and CDR regions are derived from human sequences. In one embodiment, the human monoclonal antibodies are produced by a hybridoma which includes a B cell obtained from a transgenic nonhuman

animal, *e.g.*, a transgenic mouse, having a genome comprising a human heavy chain transgene and a light chain transgene fused to an immortalized cell.

A "humanized" antibody is an antibody that retains the reactivity of a non-human antibody while being less immunogenic in humans. This can be achieved, for instance, by retaining the non-human CDR regions and replacing the remaining parts of the antibody with their human counterparts (*i.e.*, the constant region as well as the framework portions of the variable region). See, *e.g.*, Morrison *et al.*, Proc. Natl. Acad. Sci. USA, 81:6851-6855, 1984; Morrison and Oi, Adv. Immunol., 44:65-92, 1988; Verhoeven *et al.*, Science, 239:1534-1536, 1988; Padlan, Molec. Immun., 28:489-498, 1991; and Padlan, Molec. Immun., 31:169-217, 1994. Other examples of human engineering technology include, but is not limited to Xoma technology disclosed in US 5,766,886.

The terms "identical" or percent "identity," in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same. Two sequences are "substantially identical" if two sequences have a specified percentage of amino acid residues or nucleotides that are the same (*i.e.*, 60% identity, optionally 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identity over a specified region, or, when not specified, over the entire sequence), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Optionally, the identity exists over a region that is at least about 50 nucleotides (or 10 amino acids) in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides (or 20, 50, 200 or more amino acids) in length.

For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A "comparison window", as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith and Waterman (1970) Adv.

Appl. Math. 2:482c, by the homology alignment algorithm of Needleman and Wunsch, J. Mol. Biol. 48:443, 1970, by the search for similarity method of Pearson and Lipman, Proc. Nat'l. Acad. Sci. USA 85:2444, 1988, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., Brent *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons, Inc. (ringbou ed., 2003)).

Two examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, Nuc. Acids Res. 25:3389-3402, 1977; and Altschul *et al.*, J. Mol. Biol. 215:403-410, 1990, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) or 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1989) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul, Proc. Natl. Acad. Sci. USA 90:5873-5787, 1993). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is

21489-11410

considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

The percent identity between two amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (Comput. Appl. Biosci., 4:11-17, 1988) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch (J. Mol. Biol. 48:444-453, 1970) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

Other than percentage of sequence identity noted above, another indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequence.

The term "isolated antibody" refers to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds C5 is substantially free of antibodies that specifically bind antigens other than C5). An isolated antibody that specifically binds C5 may, however, have cross-reactivity to other antigens. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

The term "isotype" refers to the antibody class (e.g., IgM, IgE, IgG such as IgG1 or IgG4) that is provided by the heavy chain constant region genes. Isotype also includes modified versions of one of these classes, where modifications have been made to alter the Fc function, for example, to enhance or reduce effector functions or binding to Fc receptors.

The term "Kassoc" or "Ka", as used herein, is intended to refer to the association rate of a particular antibody-antigen interaction, whereas the term "Kdis" or "Kd," as used herein, is intended to refer to the dissociation rate of a particular antibody-antigen interaction. The term "K_D", as used herein, is intended to refer to the dissociation constant, which is obtained

from the ratio of K_D to K_A (*i.e.* K_D/K_A) and is expressed as a molar concentration (M). K_D values for antibodies can be determined using methods well established in the art. A method for determining the K_D of an antibody is by using surface plasmon resonance, or using a biosensor system such as a Biacore[®] system.

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of single molecular composition. A monoclonal antibody composition displays a single binding specificity and affinity for a particular epitope.

The term "nucleic acid" is used herein interchangeably with the term "polynucleotide" and refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, as detailed below, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.*, Nucleic Acid Res. 19:5081, 1991; Ohtsuka *et al.*, J. Biol. Chem. 260:2605-2608, 1985; and Rossolini *et al.*, Mol. Cell. Probes 8:91-98, 1994).

The term "operably linked" refers to a functional relationship between two or more polynucleotide (*e.g.*, DNA) segments. Typically, it refers to the functional relationship of a transcriptional regulatory sequence to a transcribed sequence. For example, a promoter or enhancer sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, *i.e.*, they are *cis*-acting. However, some transcriptional regulatory sequences, such as enhancers, need not be physically contiguous or located in close proximity to the coding sequences whose transcription they enhance.

As used herein, the term, "optimized" means that a nucleotide sequence has been altered to encode an amino acid sequence using codons that are preferred in the production

cell or organism, generally a eukaryotic cell, for example, a cell of Pichia, a Chinese Hamster Ovary cell (CHO) or a human cell. The optimized nucleotide sequence is engineered to retain completely or as much as possible the amino acid sequence originally encoded by the starting nucleotide sequence, which is also known as the "parental" sequence. The optimized sequences herein have been engineered to have codons that are preferred in mammalian cells. However, optimized expression of these sequences in other eukaryotic cells or prokaryotic cells is also envisioned herein. The amino acid sequences encoded by optimized nucleotide sequences are also referred to as optimized.

The terms "polypeptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer. Unless otherwise indicated, a particular polypeptide sequence also implicitly encompasses conservatively modified variants thereof.

The term "recombinant human antibody", as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom, antibodies isolated from a host cell transformed to express the human antibody, e.g., from a transfectoma, antibodies isolated from a recombinant, combinatorial human antibody library, and antibodies prepared, expressed, created or isolated by any other means that involve splicing of all or a portion of a human immunoglobulin gene, sequences to other DNA sequences. Such recombinant human antibodies have variable regions in which the framework and CDR regions are derived from human germline immunoglobulin sequences. In certain embodiments, however, such recombinant human antibodies can be subjected to *in vitro* mutagenesis (or, when an animal transgenic for human Ig sequences is used, *in vivo* somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human germline VH and VL sequences, may not naturally exist within the human antibody germline repertoire *in vivo*.

The term "recombinant host cell" (or simply "host cell") refers to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term "host cell" as used herein.

The term “subject” includes human and non-human animals. Non-human animals include all vertebrates, e.g., mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, and reptiles. Except when noted, the terms “patient” or “subject” are used herein interchangeably.

The term “treating” includes the administration of compositions or antibodies to prevent or delay the onset of the symptoms, complications, or biochemical indicia of a disease (e.g., AMD), alleviating the symptoms or arresting or inhibiting further development of the disease, condition, or disorder. Treatment may be prophylactic (to prevent or delay the onset of the disease, or to prevent the manifestation of clinical or subclinical symptoms thereof) or therapeutic suppression or alleviation of symptoms after the manifestation of the disease.

The term “vector” is intended to refer to a polynucleotide molecule capable of transporting another polynucleotide to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

4. BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows variable-region alignments of selected antibodies with their most closely related human germline sequences.

FIG. 2 shows a hemolytic assay in which human C5 is titrated into human C5-depleted serum to determine C5 activity. FIG. 3 shows titration of cynomolgus serum into human C5-depleted serum to determine optimal cynomolgus C5 concentration for alternative pathway hemolytic assay.

FIG. 4 shows examples of classical pathway hemolytic assays with 20% human serum.

FIG. 5 shows example of alternative pathway hemolytic assays with 100pM purified human C5 added to human C5-depleted serum.

FIG. 6 shows examples of alternative pathway hemolytic assays with 0.025% cynomolgus serum added to human C5-depleted serum.

FIG. 7 shows examples of classical pathway hemolytic assays (20% human serum) with matured Fabs in comparison to their respective parentals.

FIG. 8 shows examples of classical pathway hemolytic assays (5% cynomolgus serum) with matured Fabs.

FIG. 9 shows affinity matured Fab characterization in alternative pathway hemolytic assay using 100 pM human C5 added to 20% human C5-depleted serum.

FIG. 10 shows affinity matured Fab characterization in alternative pathway hemolytic assay using 20% human serum.

FIG. 11 shows affinity matured Fab characterization in alternative pathway hemolytic assay using 100 pM cynomolgus C5 added to 20% human C5-depleted serum.

FIG. 12 shows characterization of germlined IgGs in classical pathway hemolytic assay using 20% human serum.

FIG. 13 shows characterization of germlined IgGs in classical pathway hemolytic assay using 5% cynomolgus serum.

FIG. 14 shows characterization of germlined IgGs in alternative pathway hemolytic assay, 100 pM human C5.

FIG. 15 shows characterization of final germlined IgGs in alternative pathway hemolytic assay and C5a generation ELISA using 20% human serum.

FIG. 16 shows affinity matured Fab characterization in the C5a ELISA using supernatant from 20% human serum hemolytic assays.

FIG. 17 shows specificity solution ELISA on human C3, C4, C5 and cynomolgus C5 testing antibody 7091 and its derivatives.

FIG. 18 shows serum stability assays (binding to human C5 in the presence of 50% serum) with the Fabs.

FIG. 19 shows epitope binning of some affinity improved Fabs.

FIG. 20 shows an ELISA for antibody binding to mouse-human chimeric C5 or human C5 to determine alpha chain versus beta chain binders. C5 was presented by 5G1.1 to determine competition with 5G1.1.

FIG. 21 shows ELISA for testing alpha chain versus beta chain binders with 5G1.1 capture.

FIG. 22 shows results of hemolytic assay for testing alpha chain versus beta chain binders.

FIG. 23 show thermolysin proteolysis of parental Fabs at 37°C (0, 30, 60 and 90 minutes).

FIG. 24 show thermolysin proteolysis of parental Fabs at 55°C (0, 30, 60 and 90 minutes).

FIG. 25 shows thermolysin sensitivity of matured Fabs at 37°C.

FIG. 26 shows thermolysin sensitivity of matured Fabs at 55°C.

FIG. 27 shows examples of Fab inhibition of alternative pathway in MAC deposition assay.

5. DETAILED DESCRIPTION OF THE INVENTION

The present invention provides antibodies that specifically bind to complement C5 protein (e.g., human C5, cynomologus C5), pharmaceutical compositions, production methods, and methods of use of such antibodies and compositions.

5.1. C5 Antibodies

The present invention provides antibodies that specifically bind to C5 (e.g., human C5, cynomologus C5). In some embodiments, the present invention provides antibodies that specifically bind to both human and cynomologus C5. Antibodies of the invention include, but are not limited to, the human monoclonal antibodies, isolated as described, in the Examples (see Section 6 below).

The present invention provides antibodies that specifically bind a C5 protein (e.g., human and/or cynomologus C5), said antibodies comprising a VH domain having an amino acid sequence of SEQ ID NO: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281. The present invention also provides antibodies that specifically bind to a C5 protein (e.g., human and/or cynomologus C5), said antibodies comprising a VH CDR having an amino acid sequence of any one of the VH CDRs listed in Table 1, *infra*. In particular, the invention provides antibodies that specifically bind to a C5 protein (e.g., human and/or cynomologus C5), said antibodies comprising (or alternatively, consisting of) one, two, three, four, five or more VH CDRs having an amino acid sequence of any of the VH CDRs listed in Table 1, *infra*.

The present invention provides antibodies that specifically bind to a C5 protein (e.g., human and/or cynomologus C5), said antibodies comprising a VL domain having an amino acid sequence of SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188,

202, 211, 219, 228, 242, 261, 265, 269, 285, or 289. The present invention also provides antibodies that specifically bind to a C5 protein (*e.g.*, human and/or cynomologus C5), said antibodies comprising a VL CDR having an amino acid sequence of any one of the VL CDRs listed in Table 1, *infra*. In particular, the invention provides antibodies that specifically bind to a C5 protein (*e.g.*, human and/or cynomologus C5), said antibodies comprising (or alternatively, consisting of) one, two, three or more VL CDRs having an amino acid sequence of any of the VL CDRs listed in Table 1, *infra*.

Other antibodies of the invention include amino acids that have been mutated, yet have at least 60, 70, 80, 90 or 95 percent identity in the CDR regions with the CDR regions depicted in the sequences described in Table 1. In some embodiments, it includes mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated in the CDR regions when compared with the CDR regions depicted in the sequence described in Table 1.

The present invention also provides nucleic acid sequences that encode VH, VL, the full length heavy chain, and the full length light chain of the antibodies that specifically bind to a C5 protein (*e.g.*, human and/or cynomologus C5). Such nucleic acid sequences can be optimized for expression in mammalian cells (for example, Table 1 shows the optimized nucleic acid sequences for the heavy chain and light chain of antibodies 8109, 8110, 8111, 8113, 8114, 8112, 8125, 8126, 8127, 8128, 8129, 8130, 8131, 8132, and 8091).

Table 1. Examples of C5 Antibodies of the Present Invention and C5 Proteins

Antibody 8109	Sequence Identifier (SEQ ID NO:) or comments/details
CDRH1	1: SYAIS
CDRH2	2: GIGPFFGTANYAQKFQG
CDRH3	3: DTPYFDY
CDRL1	4: SGDSIPNYYVY
CDRL2	5: DDSNRPS
CDRL3	6: QSFDSLNAEV
VH	7: EVQLVQSGAEVKKPGSSVKVCKASGGTFSSYAISWVRQAPGQGLEWMGGIGPFFGTANYAQKFQGRVTITADE STSTAYMELSSLRSEDVAVYYCARDTPYFDYWGGQTLTVSS
VL	8: SYELTQPLSVSVALGQTARITCSGDSIPNYYVYVYQQKPGQAPVLIYDDSNRPSGIPERFSGSNSGNTATLTISR AQAGDEADYYCQSFDSLNAEVFGGGTKLTVL
Heavy chain	9: EVQLVQSGAEVKKPGSSVKVCKASGGTFSSYAISWVRQAPGQGLEWMGGIGPFFGTANYAQKFQGRVTITADE STSTAYMELSSLRSEDVAVYYCARDTPYFDYWGGQTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVKPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMHREALHNHYTQKSLSLSPGK
Light chain	10: SYELTQPLSVSVALGQTARITCSGDSIPNYYVYVYQQKPGQAPVLIYDDSNRPSGIPERFSGSNSGNTATLTISR AQAGDEADYYCQSFDSLNAEVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKAD SSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding	11: GAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG

SEQ ID NO:7	CCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGAT GGGCGGTATCGGTCCGTTTTTTGGCACTGCGAATTACGCGCAGAAAGTTTCAGGGCCGGGTGACCATTACCGCG GATGAAAGCACCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCG CGCGTGATACTCCTTATTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:8	12:TCCTATGAACTCACACAGCCCCTGAGCGTGAGCGTGGCCCTGGGCCAGACCGCCCGGATCACCTGCTCCG GCGACAGCATCCCCAACTACTACGTGTACTGGTACCAGCAGAAGCCCGGCCAGGCCCCCGTGTGGTGTATCTA CGACGACAGCAACCGGCCAGCGGCATCCCCGAGCGGTTGAGCGGCAGCAACAGCGGCAACACCGCCACCC TGACCATTTCAGAGCACAGGCAGGCGACGAGGCCGACTACTACTGCCAGAGCTTCGACAGCAGCCTGAACGC CGAGGTGTTTCGGCGGAGGGACCAAGTTAACCGTCCTA
PN encoding SEQ ID NO:9	13:GAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGAT GGGCGGTATCGGTCCGTTTTTTGGCACTGCGAATTACGCGCAGAAAGTTTCAGGGCCGGGTGACCATTACCGCG GATGAAAGCACCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCG CGCGTGATACTCCTTATTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGT CCATCGGTCTTCCCCCTGGCACCCCTCCTCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTC AAGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAAGTCAAGCGCCCTGACCAGCGGCGTGACACCTTC CCGGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCA CCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATC TTGTGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCCTCTTC CCCCAAAACCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCC ACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGCG GGAGGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTACCGTCCTGCACCAGGACTGGCTGAATGG CAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAGCCAAA GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGC CTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGG ACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACAC GCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:10	14:TCCTATGAACTCACACAGCCCCTGAGCGTGAGCGTGGCCCTGGGCCAGACCGCCCGGATCACCTGCTCCG GCGACAGCATCCCCAACTACTACGTGTACTGGTACCAGCAGAAGCCCGGCCAGGCCCCCGTGTGGTGTATCTA CGACGACAGCAACCGGCCAGCGGCATCCCCGAGCGGTTGAGCGGCAGCAACAGCGGCAACACCGCCACCC TGACCATTTCAGAGCACAGGCAGGCGACGAGGCCGACTACTACTGCCAGAGCTTCGACAGCAGCCTGAACGC CGAGGTGTTTCGGCGGAGGGACCAAGTTAACCGTCCTAGGTGAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTT CCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAG CCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAAC AAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCACAGAAGCTA CAGCTGCCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:9	15:GAGGTGCAGCTGGTGACAGAGCGGAGCCGAGGTGAAGAAGCCCGGTAGCAGCGTCAAGGTGTCCTGCAAG GCCAGCGGCGGCACCTTCAGCAGCTACGCCATCAGCTGGGTGCGGCAGGCCCCAGGCCAGGGCCTGGAGTG GATGGGCGGCATCGGCCATTCTTCGGCACCGCCAACTACGCCCAGAAGTTCCAGGGCAGGGTCACCATCAC CGCCGACGAGAGCACACCGCCTACATGGAGCTGTCCAGCCTGAGAAGCGAGGACACCGCCGTGTACTA CTGCGCCAGAGACACCCCTACTTCGACTACTGGGGCCAGGGCACCCCTGGTGACCGTGAGCAGCGCTAGCAC CAAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCT GCCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTG ACACCTTCCCCGCCGTGCTGCAGAGCAGCGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCAGCAGCA GCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGA GCCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGT GTTCTGTTCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGT GGACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCACAACGCCAA GACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGA CTGGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCATCGAAAAGACCATC

	AGCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAG AACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAAC GGCCAGCCCGAGAACAATAACAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCCTGTACAGCA AGCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTACAGCTGCAGCGTGATGCACGAGGCCCTGC ACAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:10	16:AGCTACGAGCTGACCCAGCCCCTGAGCGTGAGCGTGGCCCTGGGCCAGACCGCCAGGATCACCTGCAGCG GCGACAGCATCCCCAACTACTACGTGTACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATCTA CGACGACAGCAACAGGCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACAGCGGCAACACCGCCACCCT GACCATCAGCAGAGCCCAGGCCGGCGACGAGGCCGACTACTACTGCCAGAGCTTCGACAGCTCACTGAACGC CGAGGTGTTTCGGCGGAGGGACCAAGCTGACCGTGCTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCTGTT CCCCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAGG CGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGAGACCACCACCCCCAGCA AGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAGAGCCACAGGT CCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAGAGACCGTGCCCCCAACCGAGTGACAGC
Antibody 8110	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	17: NYIS
CDRH2	18: IIDPDDSYTEYSPSFQG
CDRH3	19: YEYGGFDI
CDRL1	20: SGDNIGNSYVH
CDRL2	21: KDNDRPS
CDRL3	22: GTYDIESYV
VH	23:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTEYSPSFQGGVVISADKSI STAYLQWSSLKASDTAMYYCAREYEGGFDIWGQGLTVTVSS
VL	24:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCGTYDIESYVFGGGTKLTVL
Heavy chain	25:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTEYSPSFQGGVVISADKSI STAYLQWSSLKASDTAMYYCAREYEGGFDIWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMHENHHTQKSLSLSPGK
Light chain	26:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCGTYDIESYVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADS SPVKAGVETTTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:23	27:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGCA TTATTGATCCTGATGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATAGCGCGGATAAAAG CATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTATG AGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:24	28:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGG CGATAATATTGGTAATTCTTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGGTACTTATGATATTGAGTCTTATGTGTTTGG CGGCGGCACGAAGTTAACCGTCCTA
PN encoding SEQ ID NO:25	29:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGCA TTATTGATCCTGATGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATAGCGCGGATAAAAG CATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTATG AGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGTCCATC GGTCTTCCCCCTGGCACCCCTCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGA

	CTACTTCCCCGAACCGGTGACGGTGTCTGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGC TGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAG ACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGTGA CAAACTCACACATGCCACCGTGCCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCTCTTCCCCCA AAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAG ACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGG AGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGG AGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCA GCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGAC CTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA CTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAG AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGA AGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:26	30:AGTTACGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGACACCGCGCGTATCTCGTGTAGCGG CGATAATATTGGTAATTCTTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGGTACTTATGATATTGAGTCTTATGTGTTTGG CGGCGGCACGAAGTTAACCGTCCTAGGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTCT GAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGG CCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAGCAACAACA AGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCACAGAAGCTACAGCTGCCAGG TCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:25	31:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAGG GCAGCGGCTACAGCTTACCAACTACATCAGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGATGG GCATCATCGACCCCGACGACAGCTACACCGAGTACAGCCCCAGCTTCCAGGGCCAGGTGACCATCAGCGCCG ACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGCG CCAGATACGAGTACGGCGGCTTCGACATCTGGGGCCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACCA AGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTGC CTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGAC ACCTTCCCCGCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCCAGCAGCAGC CTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGAGC CCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGTGT TCCTGTTCCCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGTGG ACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAGA CCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGACT GGCTGAACGGCAAAGAATAAAGTGAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCAG CAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGAA CCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAACGG CCAGCCCGAGAACAATAAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAAG CTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTGAGCTGCAGCGTGATGCACGAGGCCCTGCAC AACCACTACACCCAGAAGAGCCTGAGCCTGTACCCGGCAAG
Optimized PN encoding SEQ ID NO:26	32:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGGCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAACATCGGCAACAGCTACGTGCACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATC TACAAGGACAACGACAGGCCCAGCGGCATCCCCGAGAGGTTTACGCGGCAGCAACTCCGGCAACACCGCCACC CTGACCATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCGGCACCTACGACATCGAGTCATAC GTGTTTGGCGGAGGGACCAAGCTGACCGTGCTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCTGTTCCCC CCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAGGCGCC GTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACCACCCCCAGCAAGCA GAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAGAGCCACAGGTCTTA CAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 8111	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS

23

	AGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTATGATTCTCAGTCTAT TGTGTTTGGCGGCGGCACGAAGTTAACCGTCCTAGGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCG CCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCG TGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAA GCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAG CTGCCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:41	47:GAGGTGACCCTGAAGGAGAGCGGCCAGCCCTGGTGAAGCCCACCCAGACCCTGACCCTGACTTGACCT TCAGCGGCTTCAGCCTGAGCACCAGCGGAGGGGGCGTGAGCTGGATCAGGCAGCCCCCAGGTAAGGCCCTG GAGTGGCTGGCCAATATCGACGACGCCGATATCAAGGACTACAGCCCCAGCCTGAAGAGCAGGCTGACCATCA GCAAGGACACCAGCAAGAACCAGGTGGTGTGACCATGACCAATATGGACCCCGTGGACACCGCCACCTACTA CTGCGCCAGAGGCCCCCTACGGCTTCGACAGCTGGGGCCAGGGCACCCTGGTGACCGTCAGCTCAGCTAGCAC CAAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCT GCCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCAGCGGCGTG ACACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCCAGCAGCA GCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGA GCCCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGT GTTCTGTTCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGT GGACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCACAACGCCAA GACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGA CTGGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATC AGCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAG AACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAAC GGCCAGCCCGAGAACAATAACAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCA AGCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTACGTGCAGCGTGATGCACGAGGCCCTGC ACAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:42	48:GAGAGCGCCCTGACCCAGCCCGCCAGCGTGAGCGGCAGCCCAGGCCAGTCTATCACAATCAGCTGCACCG GCACCTCCAGCGATATCGGCACCTACAATACTAGTGAGCTGGTATCAGCAGCACCCCGGCAAGGCCCCCAAGCT GATGATCTACGACGACAGCAACAGGCCAGCGGCGTGAGCAACAGGTTTCAGCGGCAGCAAGAGCGGCAACAC CGCCAGCCTGACAATCAGCGGCCTGCAGGCCGAGGACGAGGCCGACTACTACTGCCAGAGCTACGACAGCCA GTCAATCGTGTTGCGCGGAGGGACCAAGCTGACCGTGCTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCT GTTCCCCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCC AGGCGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGAGACCACCACCCCCA GCAAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAGAGCCACA GGTCCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCCAACCGAGTGCAGC
Antibody 8113	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	49: IIDPDDSYTRYSPSFQG
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	50:ATWGSEDQV
VH	51:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTRYSPSFQGGQVTISADKSI STAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGLVTVSS
VL	52:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCATWGSEDQVFGGGTKLTVL
Heavy chain	53:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTRYSPSFQGGQVTISADKSI STAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE

	SNGQPENNYKTTTPVLDSDFSFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
Light chain	54:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCATWGSSEQVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKAD SSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:51	55:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGCA TTATCGATCCGGATGATAGCTATACCCGTTATTCTCCGAGCTTTCAGGGACAGGTGACCATTAGCGCGGATAAA AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:52	56:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGAGACCGCGCGTATCTCGTGTAGCGG CGATAATATTGGTAATTCTTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGCTACTTGGGGTTCTGAGGATCAGGTGTTTG GCGGCGGCACGAAGTTAACCGTCCTA
PN encoding SEQ ID NO:53	57:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGCA TTATCGATCCGGATGATAGCTATACCCGTTATTCTCCGAGCTTTCAGGGACAGGTGACCATTAGCGCGGATAAA AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGTCCA TCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAG GACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGCGGCGTGACACCTTCCCG GCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCC AGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGT GACAAAACCTCACACATGCCACCGTGCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCTCTTCCCCC CAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGA AGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA GGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCTCACCCTGCTGACCAGGACTGGCTGAATGGCAA GGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATCTCAAAGCCAAAGGG CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTG ACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC AACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACA AGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCA GAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:54	58:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGAGACCGCGCGTATCTCGTGTAGCGG CGATAATATTGGTAATTCTTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGCTACTTGGGGTTCTGAGGATCAGGTGTTTG GCGGCGGCACGAAGTTAACCGTCCTAGGTGAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTC TGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGAAGTCTACCCGGGAGCCGTGACAGTG GCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAAC AAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCTGCCAG GTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:53	59:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAGG GCAGCGGCTACAGCTTACCAACTACATCAGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGATGG GCATCATCGACCCCCGACGACAGCTACACCAGGTACAGCCCCAGCTTCCAGGGCCAGGTGACCATCAGCGCCG ACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGCG CCAGATACGAGTACGGCGGCTTCGACATCTGGGGCCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACC AGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTGC CTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGAC ACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCAGCAGCAGC CTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGAGC

	CCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGTGT TCCTGTTCCCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGTTGGTGG ACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAGA CCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGACT GGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCAG CAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGAA CCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAACGG CCAGCCCCGAGAACAACTACAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAAG CTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTTCAGCTGCAGCGTGATGCACGAGGCCCTGCAC AACCCTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:54	60:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGGCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAATATCGGCAACAGCTACGTGCACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATC TACAAGGACAACGACAGGCCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACTCCGGCAACACCGCCACC CTGACAATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCGCCACCTGGGGCTCAGAGGACCAG GTGTTTCGGCGGAGGGACCAAGCTGACCGTGCTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCGTGTTCCCC CCCAGCAGCGAGGAGCTGCAGGCCAACAAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAGGCGCC GTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACCACCCCCAGCAAGCA GAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAGAGCCACAGGTCCTA CAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAGACCGTGCCCCCAACCGAGTGCAGC
Antibody 8114	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	61: SYYIG
CDRH2	62: IIDPTDSQTAYSPSFQG
CDRH3	63: YMMRGFDH
CDRL1	64: SGDSLGDYYAY
CDRL2	65: KDNNRPS
CDRL3	66: QTWDTGESGV
VH	67: EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSQTAYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVTVSS
VL	68: SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTWDTGESGVFGGGTKLTVL
Heavy chain	69: EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSQTAYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKHTCPP CPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
Light chain	70: SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTWDTGESGVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTTPSKQSNKNYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:67	71: GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGATGG GCATTATTGATCCTACTGATTCTCAGACTGCTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATA AAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGT TATATGATGCGTGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:68	72: AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGACACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATACTGGTGAGTCTGGTGTGTT TGGCGGCGGCACGAAGTTAACCGTCCTA
PN encoding	73: GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG

SEQ ID NO:69	TTCCGGATATTCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGG GCATTATTGATCCTACTGATTCTCAGACTGCTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATA AAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGT TATATGATGCGTGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGTC CATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGACAGCGGCCCTGGGCTGCCTGGTCA AGGACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCC CGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCAC CCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTT GTGACAAAACCTCACACATGCCCCACCGTGCCCGAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCTCTTCCC CCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGCGG GAGGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGC AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAG GGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGCC TGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGA ACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGA CAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACG CAGAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:70	74:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGAGACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATACTGGTGAGTCTGGTGTGTT TGGCGGCGGCACGAAGTTAACCGTCCTAGGTGAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCC TCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGAATTCTACCCGGGAGCCGTGACAG TGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACACACCCTCCAAACAAAGCAACA ACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCA GGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:69	75:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAGG GCAGCGGCTACAGCTTACCAGCTACTACATCGGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGA TGGGCATCATCGACCCACCGACAGCCAGACCGCCTACAGCCCCAGCTTCCAGGGCCAGGTGACCATCAGCG CCGACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACT GCGCCCCGTACATGATGAGGGGCTTCGACCACTGGGGTCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCA CCAAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGC TGCCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTG CACACCTTCCCCGCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCCAGCAGC AGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGG AGCCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCG TGTTCTGTTCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGT GGACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCACAACGCCAA GACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGA CTGGCTGAACGGCAAAGAATAACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATC AGCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAG AACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAAC GGCCAGCCCCGAGAACAATAACAAGACCACCCCCCAGTGTGGACAGCGACGGCAGCTTCTTCTGTACAGCA AGCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTGAGCTGCAGCGTGATGCACGAGGCCCTGC ACAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:70	76:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGGCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAGCCTGGGCGACTACTACGCCTACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCCGTGCTGGTGATC TACAAGGACAACAACAGGCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACAGCGGCAACACCGCCACC CTGACAATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCCAGACCTGGGACACCGGCCGAGTCA GGCGTGTTCCGGCGAGGGACCAAGCTGACCGTGCTGGGTGAGCCTAAGGCTGCCCCAGCGTGACCCCTGTTT

	CCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAGGC GCCGTGACCGTGCCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGAGACCACCACCCCCAGCAA GCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAGAGCCACAGGTC CTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 8112	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	77: IIDPSDSHTTYSPSFQG
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	78:QTDILPHGLV
VH	79:EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPSDSHTTYSPSFQGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVT VSS
VL	80:SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPV LVIYKDN NRP SGIPERFSGSNSGNTATLTIS GTQAEDEADYYC QTDILPHGLVFGGGTKLTVL
Heavy chain	81:EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPSDSHTTYSPSFQGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVT VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV PSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPP CPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS VLT V L H Q D W L N G K E Y K C K V S N K A L P A P I E K T I S K A K G Q P R E P Q V Y T L P P S R E E M T K N Q V S L T C L V K G F Y P S D I A V E W ESNGQPENNYKTTTPVLDS DGSFFLYSKLTVDKSRWQQGNV FSCSV MHEALHNHYTQKSLSLSPGK
Light chain	82:SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPV LVIYKDN NRP SGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTDILPHGLVFGGGTKLTVLGQPKAAPS VTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:79	83:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGG GCATTATCGATCCGTCTGATAGCCATACCACTTATTCTCCGAGCTTTACAGGGCCAGGTGACCATTAGCGCGGAT AAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:80	84:AGTTACGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATATTCTTCCTCATGGTCTTGT GTTTGGCGGCGGCACGAAGTTAACCGTCCTA
PN encoding SEQ ID NO:81	85:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGG GCATTATCGATCCGTCTGATAGCCATACCACTTATTCTCCGAGCTTTACAGGGCCAGGTGACCATTAGCGCGGAT AAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGG TCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGT CAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTT CCCGGCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGC ACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAAT CTTGTGACAAAACCTCACACATGCCACCGTGCCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCCTCTT CCCCCAAACCCAAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGACGTGAGC CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGC GGGAGGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTACCGTCCTGCACCAGGACTGGCTGAATG GCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATCTCCAAGCCAA AGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAG CCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGA

	GAACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTG GACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:82	86:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATATTCTTCTCATGGTCTTGT GTTTGGCGGCGGCACGAAGTTAACCGTCCTAGGTACGCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCC TCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTATAAGTGAAGTCTACCCGGGAGCCGTGA CAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACACCCCTCCAAACAAAGCA ACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTG CCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:81	87:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAGG GCAGCGGCTACAGCTTCACCAGCTACTACATCGGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGA TGGGCATTATCGATCCGTCTGATAGCCATACCACTTATTCTCCGAGCTTTAGGGGCCAGGTGACCATCAGCGCC GACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGC GCCCCGTACATGATGAGGGGCTTCGACCACTGGGGTCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACC AAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTG CCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGCA CACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGGCCAGCAGCAG CCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCAGCAACACCAAGGTGGACAAGAGAGTGGAG CCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGTG TTCCTGTTCCCCCCTAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGTG GACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAG ACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGAC TGGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCA GCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGA ACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCAGCGACATCGCCGTGGAGTGGGAGAGCAACG GCCAGCCCCGAGAACAACACTACAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAA GCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTAGCTGCAGCGTGATGCACGAGGCCCTGCA CAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:82	88:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGGCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAGCCTGGGCGACTACTACGCCTACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATC TACAAGGACAACAACAGGCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACAGCGGCAACACCGCCACC CTGACAATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCCAGACTTGGGATATTCTTCTCATG GTCTTGTGTTGCGCGGAGGGACCAAGCTGACCGTGCTGGGTGAGCCTAAGGCTGCCCCCAGCGTGACCCCTGT TCCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCCTGGTGTGCCTGATCAGCGACTTCTACCCAG GCGCCGTGACCGTGCCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACACCCCCAGC AAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCGAGCAGTGGAAGAGCCACAGG TCCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 8125	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:77
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	89:QAWTDSPTGLV
VH	SEQ ID NO:79
VL	90:SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQKPGQAPVLVIYKDNRRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQAWTDSPTGLVFGGGTKLTVL
Heavy chain	SEQ ID NO:81

Light chain	91:SYELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQAWTDSPTGLVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWK ADSSPVKAGVETTTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:79	SEQ ID NO:83
PN encoding SEQ ID NO:90	92:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGGCTTGGACTGATTCTCCTACTGGTCTTGT GTTTGGCGGCGGCACGAAGTTAACCGTCCTA
PN encoding SEQ ID NO:81	SEQ ID NO:85
PN encoding SEQ ID NO:91	93:AGTTACGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGG CGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATAA GGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGACC ATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGGCTTGGACTGATTCTCCTACTGGTCTTGT GTTTGGCGGCGGCACGAAGTTAACCGTCCTAGGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCC TCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGAAGTCTACCCGGGAGCCGTGA CAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACACACCCCTCCAAACAAAGCA ACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCTG CCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:81	SEQ ID NO:87
Optimized PN encoding SEQ ID NO:91	94:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGCCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAGCCTGGGCGACTACTACGCCTACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATC TACAAGGACAACAACAGGCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACAGCGGCAACACCGCCACC CTGACAATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCCAGGCTTGGACTGATTCTCCTACTG GTCTTGTGTTTCGGCGGAGGGACCAAGCTGACCGTGCTGGGTCAGCCTAAGGCTGCCCCCAGCGTGACCCCTGT TCCCCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAG GCGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACACCCCCAGC AAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGAAGAGCCACAGG TCCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 8126	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:62
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:89
VH	SEQ ID NO:67
VL	SEQ ID NO:90
Heavy chain	SEQ ID NO:69
Light chain	SEQ ID NO:91
PN encoding SEQ ID NO:79	SEQ ID NO:71
PN encoding SEQ ID NO:90	SEQ ID NO:92
PN encoding SEQ ID NO:81	SEQ ID NO:73

PN encoding SEQ ID NO:91	SEQ ID NO:93
Optimized PN encoding SEQ ID NO:81	SEQ ID NO:75
Optimized PN encoding SEQ ID NO:91	SEQ ID NO:94
Antibody 8127	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	95: IIDPTDSYTVYSPSFQG
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:89
VH	96:EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSYTVYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGTLVTVSS
VL	SEQ ID NO:90
Heavy chain	97:EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSYTVYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPP CPAPEAAGGPSVFLFPPKPKDITLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVS VLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK
Light chain	SEQ ID NO:91
PN encoding SEQ ID NO:96	98:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGG GCATTATTGATCCTACTGATTCTTATACTGTTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAA AAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTT ATATGATGCGTGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGC
PN encoding SEQ ID NO:90	SEQ ID NO:92
PN encoding SEQ ID NO:97	99:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGG TTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGG GCATTATTGATCCTACTGATTCTTATACTGTTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAA AAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTT ATATGATGCGTGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGCCTCCACCAAGGGT CCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTC AAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCCCTGACCAGCGGCGTGACACCTTC CCGGCTGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCA CCCAGACCTACATCTGCAACGTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATC TTGTGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCTCTTC CCCCAAAACCAAGGACACCCCTCATGATCTCCCGAACCCCTGAGGTACATGCGTGGTGGTGACGTGAGCC ACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGTGCATAATGCCAAGACAAAGCCGCG GGAGGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTACCGTCCTGCACCAGGACTGGCTGAATGG CAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAAGCCAAA GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGC CTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGG ACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACAC

	GCAGAAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:91	SEQ ID NO:93
Optimized PN encoding SEQ ID NO:97	100:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAG GGCAGCGGCTACAGCTTCACCAGCTACTACATCGGCTGGGTGCGGCAGATGCCCCGCAAGGGCCTGGAGTGG ATGGGCATTATTGATCCTACTGATTCTTATACTGTTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATCAGCGCC GACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGC GCCCCGTACATGATGAGGGGCTTCGACCACTGGGGTCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACCC AAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTG CCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGCA CACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCAGCAGCAG CCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGAG CCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCCTTCCGTG TTCCTGTTCCCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGTG GACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAG ACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGAC TGGCTGAACGGCAAAGAATACAAGTGAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCA GCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGA ACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAACG GCCAGCCCCGAGAACAATAACAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAA GCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTAGCTGCAGCGTGATGCACGAGGCCCTGCA CAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:91	SEQ ID NO:94
Antibody 8128	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:49
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	101:STWDIEPTYV
VH	SEQ ID NO:51
VL	102:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCSTWDIEPTYVFGGGTKLTVL
Heavy chain	SEQ ID NO:53
Light chain	103:SYELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCSTWDIEPTYVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKAD SSPVKAGVETTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:51	SEQ ID NO:55
PN encoding SEQ ID NO:102	104:AGTTACGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCTCTACTTGGGATATTGAGCCTACTTATGTGT TTGGCGGCGGCACGAAGTTAACCGTCCTA
PN encoding SEQ ID NO:53	SEQ ID NO:57
PN encoding SEQ ID	105:AGTTACGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA

NO:103	AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCTCTACTTGGGATATTGAGCCTACTTATGTGT TTGGCGGCGGCACGAAGTTAACCGTCCTAGGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTC CTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACA GTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAAC AACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCCACAGAAGCTACAGCTGCC AGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:53	SEQ ID NO:59
Optimized PN encoding SEQ ID NO:103	106:AGCTACGAGCTGACCCAGCCCCCAGCGTGAGCGTGGCCCCAGGCCAGACCGCCAGGATCAGCTGCAGC GGCGACAATATCGGCAACAGCTACGTGCACTGGTATCAGCAGAAGCCCGGCCAGGCCCCCGTGCTGGTGATC TACAAGGACAACGACAGGCCCAGCGGCATCCCCGAGAGGTTTCAGCGGCAGCAACTCCGGCAACACCGCCACC CTGACAATCAGCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCTCTACTTGGGATATTGAGCCTACTT ATGTGTTTCGGCGGAGGGACCAAGCTGACCGTGCTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCTGTTCC CCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAGGCG CCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGAGACCACCACCCCCAGCAAG CAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGGAAAGAGCCACAGGTCC TACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 8129	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	107: IIDPQDSYTEYSPSFQG
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	108:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPQDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSS
VL	SEQ ID NO:24
Heavy chain	109:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPQDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVMEALHNHYTQKSLSLSPGK
Light chain	SEQ ID NO:26
PN encoding SEQ ID NO:108	110:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTCAGGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:24	SEQ ID NO:28
PN encoding SEQ ID NO:109	111:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTCAGGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGTCCAT CGGTCTTCCCCCTGGCACCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGG ACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGG

	CTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCA GACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGTG ACAAAACCTCACACATGCCCACCGTGCCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCCTCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGTTGGTGGACGTGAGCCACGAA GACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAG GAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAG GAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGC AGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTGAGCCTGA CCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACA ACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAA GAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAG AAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:26	SEQ ID NO:30
Optimized PN encoding SEQ ID NO:109	112:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAG GGCAGCGGCTACAGCTTCACCAACTACATCAGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGATG GGCATCATCGACCCCCAGGACAGCTACACCGAGTACAGCCCCAGCTTCAGGGCCAGGTGACCATCAGCGCC GACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGC GCCAGATACGAGTACGGCGGCTTCGACATCTGGGGCCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACC AAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTG CCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGCA CACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCAGCAGCAG CCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGAG CCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGTG TTCCTGTTCCCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGTG GACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAG ACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGAC TGGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCA GCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGA ACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAACG GCCAGCCCCGAGAACAATAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAA GCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTGAGCTGCAGCGTGATGCACGAGGCCCTGCA CAACCACTACACCCAGAAGAGCCTGAGCCTGTCACCCGGCAAG
Optimized PN encoding SEQ ID NO:26	SEQ ID NO:32
Antibody 8130	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:107
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:101
VH	SEQ ID NO:108
VL	SEQ ID NO:102
Heavy chain	SEQ ID NO:109
Light chain	SEQ ID NO:103
PN encoding SEQ ID NO:108	SEQ ID NO:110
PN encoding	SEQ ID NO:104

SEQ ID NO:102	
PN encoding SEQ ID NO:109	SEQ ID NO:111
PN encoding SEQ ID NO:103	SEQ ID NO:105
Optimized PN encoding SEQ ID NO:109	SEQ ID NO:112
Optimized PN encoding SEQ ID NO:103	SEQ ID NO:106
Antibody 8131	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	113: IIDPEDSHTEYSPSFQG
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	114:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPEDSHTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGTLVTVSS
VL	SEQ ID NO:24
Heavy chain	115:EVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPEDSHTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPC PAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
Light chain	SEQ ID NO:26
PN encoding SEQ ID NO:114	116:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGAGGATTCTCATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAAA AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:24	SEQ ID NO:28
PN encoding SEQ ID NO:115	117:GAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGAGGATTCTCATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAAA AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCCTCCACCAAGGGTCCA TCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAG GACTACTTCCCCGAACCGGTGACGGTGTCGTGGAACCTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCG GCTGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCC AGACCTACATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTTGAGCCCAAATCTTGT GACAAAACCTCACACATGCCCACCGTGCCCAGCACCTGAAGCAGCGGGGGGACCGTCAGTCTTCTCTTCCCCC CAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCACGA AGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA

	GGAGCAGTACAACAGCACGTACCGGGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAA GGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGG CAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTG ACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAAC AACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACA AGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCA GAAGAGCCTCTCCCTGTCTCCGGGTAAA
PN encoding SEQ ID NO:26	SEQ ID NO:30
Optimized PN encoding SEQ ID NO:115	118:GAGGTGCAGCTGGTGCAGAGCGGAGCCGAGGTGAAAAAGCCCGGTGAGAGCCTGAAGATCAGCTGCAAG GGCAGCGGCTACAGCTTCACCAACTACATCAGCTGGGTGCGGCAGATGCCCGGCAAGGGCCTGGAGTGGATG GGCATCATCGACCCCCGAGGACAGCCATACCGAGTACAGCCCCAGCTTCCAGGGCCAGGTGACCATCAGCGCC GACAAGAGCATCAGCACCGCCTACCTGCAGTGGAGCAGCCTGAAGGCCAGCGACACCGCCATGTACTACTGC GCCAGATACGAGTACGGCGGCTTCGACATCTGGGGCCAGGGCACCCCTGGTGACCGTCAGCTCAGCTAGCACC AAGGGCCCCAGCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCTCCGGCGGCACAGCCGCCCTGGGCTG CCTGGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACAGCGGAGCCCTGACCAGCGGCGTGCA CACCTTCCCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGTCCAGCGTGGTGACAGTGCCCAGCAGCAG CCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAGAGTGGAG CCCAAGAGCTGCGACAAGACCCACACCTGCCCCCCTGCCAGCCCCCGAAGCTGCAGGCGGCCCTTCCGTG TTCCTGTTCCCCCCCCAAGCCCAAGGACACCCTGATGATCAGCAGGACCCCCGAGGTGACCTGCGTGGTGGTG GACGTGAGCCACGAGGACCCAGAGGTGAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAG ACCAAGCCCAGAGAGGAGCAGTACAACAGCACCTACAGGGTGGTGTCCGTGCTGACCGTGCTGCACCAGGAC TGGCTGAACGGCAAAGAATACAAGTGCAAGGTCTCCAACAAGGCCCTGCCTGCCCCCATCGAAAAGACCATCA GCAAGGCCAAGGGCCAGCCACGGGAGCCCCAGGTGTACACCCTGCCCCCTTCTCGGGAGGAGATGACCAAGA ACCAGGTGTCCCTGACCTGTCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAACG GCCAGCCCCGAGAACAATAAGACCACCCCCCAGTGCTGGACAGCGACGGCAGCTTCTTCTGTACAGCAA GCTGACCGTGGACAAGAGCAGGTGGCAGCAGGGCAACGTGTTTCAGCTGCAGCGTGATGCACGAGGCCCTGCA CAACCACTACACCCAGAAGAGCCTGAGCCTGTACCCGGCAAG
Optimized PN encoding SEQ ID NO:26	SEQ ID NO:32
Antibody 8132	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:113
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:101
VH	SEQ ID NO:114
VL	SEQ ID NO:102
Heavy chain	SEQ ID NO:115
Light chain	SEQ ID NO:103
PN encoding SEQ ID NO:114	SEQ ID NO:116
PN encoding SEQ ID NO:102	SEQ ID NO:104
PN encoding	SEQ ID NO:117

SEQ ID NO:115	
PN encoding SEQ ID NO:103	SEQ ID NO:105
Optimized PN encoding SEQ ID NO:115	SEQ ID NO:118
Optimized PN encoding SEQ ID NO:103	SEQ ID NO:106
Antibody 8091	Sequence Identifier (SEQ ID NO:) or comments/details
CDRH1	SEQ ID NO:1
CDRH2	119: NIGPFFGIANYAQKFQG
CDRH3	SEQ ID NO:3
CDRL1	SEQ ID NO:4
CDRL2	SEQ ID NO:5
CDRL3	120: QTYDDGSTAEV
VH	121: QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAIWVRQAPGQGLEWMGNIGPFFGIANYAQKFQGRVTITAD ESTSTAYMELSSLRSEDTAVYYCARDTPYFDYWGGQGLTVTVSS
VL	122: DIELTQPPSVSVAPGQTARISCSGDSIPNYYVYWYQQKPGQAPVLVIYDDSNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTYDDGSTAEVFGGGTKLTVL
Heavy chain	123: QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAIWVRQAPGQGLEWMGNIGPFFGIANYAQKFQGRVTITAD ESTSTAYMELSSLRSEDTAVYYCARDTPYFDYWGGQGLTVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCVECP PCP APPVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTV VHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK
Light chain	124: DIELTQPPSVSVAPGQTARISCSGDSIPNYYVYWYQQKPGQAPVLVIYDDSNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTYDDGSTAEVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTTPSKQSNKYAASSYLSLTPEQWKSRSYSCQVTHEGSTVEKTVAPTECS
PN encoding SEQ ID NO:121	125: CAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGAT GGGCAATATCGGTCCGTTTTTTGGCATTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCG GATGAAAGCACCAGCACCGCGTATATGGAAGTACGAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCG CGCGTGATACTCCTTATTTTATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:122	126: GATATCGAAGTACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTACAGCCGCGGTATCTCGTGTAGCG GCGATTCTATTCTAATTATTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATG ATGATTCTAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTATGATGATGGTTCTACTGCTGAGG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:123	127: CAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGAT GGGCAATATCGGTCCGTTTTTTGGCATTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCG GATGAAAGCACCAGCACCGCGTATATGGAAGTACGAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCG CGCGTGATACTCCTTATTTTATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCTTCCACCAAGGGC CCCAGCGTGTTCCCCCTGGCCCCCTGCAGCAGAAGCACCAGCGAGAGCACAGCCGCCCTGGGCTGCCTGGTG AAGGACTACTTCCCCGAGCCCGTGACCGTGAGCTGGAACAGCGGAGCCCTGACCAGCGGCGTGACACCTTC CCCGCCGTGCTGCAGAGCAGCGGCCTGTACAGCCTGAGCAGCGTGGTGACCGTGCCCAGCAGCAACTTCGGC ACCCAGACCTACACCTGCAACGTGGACCACAAGCCCAGCAACACCAAGGTGGACAAGACCGTGGAGCGGAAG TGCTGCGTGGAGTGCCCCCCTGCCCTGCCCTCCTGTGGCCGGACCCTCCGTGTTCTTCCCCCAAG

	CCCAAGGACACCCTGATGATCAGCCGGACCCCCGAGGTGACCTGCGTGGTGGTGGACGTGAGCCACGAGGAC CCCGAGGTGCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAGACCAAGCCCCGGGAGGAA CAGTTCAACAGCACCTTCCGGGTGGTGTCCGTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAAGAAT ACAAGTGCAAGGTGTCCAACAAGGGCCTGCCTGCCCCCATCGAGAAAACCATCAGCAAGACAAAGGGGCCAGCC CAGGGAACCCAGGTGTACACCCTGCCCCCAGCCGGGAGGAAATGACCAAGAACCAGGTGTCCCTGACCTG TCTGGTGAAGGGCTTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAACGGCCAGCCCGAGAACAATA CAAGACCACCCCCCATGCTGGACAGCGACGGCAGCTTCTTCCTGTACAGCAAGCTGACAGTGGACAAGAGC CGGTGGCAGCAGGGCAACGTGTTGAGCTGCAGCGTGATGCACGAGGCCCTGCACAACCACTACACCCAGAAG AGCCTGAGCCTGTCCCCCGGCAAA
PN encoding SEQ ID NO:124	128:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGACACCGCGCGTATCTCGTGTAGCG GCGATTCTATTCTAATTATTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATG ATGATTCTAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTATGATGATGGTTCTACTGCTGAGG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCC CTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTG ACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGC AACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCT GCCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGACAGTGGCCCCTACAGAATGTTCA
Optimized PN encoding SEQ ID NO:123	129:CAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAGCCCCGGCTCCTCCGTGAAGGTGTCCTGCAAG GCCTCCGGCGGCACCTTCTCCTCCTACGCCATCTCCTGGGTGCGGCAGGCCCCCGGCCAGGGCCTGGAGTGG ATGGGCAACATCGGCCCCCTTCTTCGGCATCGCCAACTACGCCCAGAAGTTCCAGGGCCGGGTGACCATCACCG CCGACGAGTCCACCTCCACCGCCTACATGGAGCTGTCTCCTGCGGTCCGAGGACACCGCCGTGTACTACTG CGCCCGGGACACCCCCTACTTCGACTACTGGGGCCAGGGCACCCCTGGTGACCGTGTCTCCTCCGCCTCCACCAA GGGCCCCCTCCGTGTTCCCCCTGGCCCCCTGCTCCCGGTCCACCTCCGAGTCCACCGCCGCCCTGGGCTGCCT GGTGAAGGACTACTTCCCCGAGCCCGTGACCGTGTCTGGAACCTCCGGCGCCCTGACCTCCGGCGTGACAC CTTCCCCGCCGTGCTGCAGTCTCCTCGGCCTGTACTCCTGTCTCCTCCGTGGTGACCGTGCCCTCCTCCAACCTC GGCACCCAGACCTACACCTGCAACGTGGACCACAAGCCCTCCAACACCAAGGTGGACAAGACCGTGGAGCGG AAGTGCTGCGTGGAGTGCCCCCCTGCCCGCCCCCCCCCGTGGCCGGCCCCCTCCGTGTTCTCTGTTCCCCCCC AAGCCCAAGGACACCCTGATGATCTCCCGGACCCCCGAGGTGACCTGCGTGGTGGTGGACGTGTCCACGAG GACCCCGAGGTGCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCACAACGCCAAGACCAAGCCCCGGGAG GAGCAGTTCAACTCCACCTTCCGGGTGGTGTCCGTGCTGACCGTGGTGCACCAGGACTGGCTGAACGGCAAG GAGTACAAGTGCAAGGTGTCCAACAAGGGCCTGCCCGCCCCCATCGAGAAGACCATCTCCAAGACCAAGGGC CAGCCCCGGGAGCCCCAGGTGTACACCCTGCCCCCCTCCCGGGAGGAGATGACCAAGAACCAGGTGTCCCTG ACCTGCCTGGTGAAGGGCTTCTACCCCTCCGACATCGCCGTGGAGTGGGAGTCCAACGGCCAGCCCGAGAAC AACTACAAGACCACCCCCCCCCATGCTGGACTCCGACGGCTCCTTCTTCCTGTACTCCAAGCTGACCGTGGACAA GTCCCGGTGGCAGCAGGGCAACGTGTTCTCCTGCTCCGTGATGCACGAGGCCCTGCACAACCACTACACCCA GAAGTCCCTGTCCCTGTCCCCCGGCAAG
Optimized PN encoding SEQ ID NO:124	130:GACATCGAGCTGACCCAGCCCCCTCCGTGTCCGTGGCCCCCGGCCAGACCGCCCGGATCTCCTGCTCC GGCGACTCCATCCCCAACTACTACGTGTACTGGTACCAGCAGAAGCCCGGCCAGGCCCCCGTGTGGTGATCT ACGACGACTCCAACCGGCCCTCCGGCATCCCGAGCGGTTCTCCGGCTCCAACCTCCGGCAACACCGCCACCC TGACCATCTCCGGCACCCAGGCCGAGGACGAGGCCGACTACTACTGCCAGACCTACGACGACGGCTCCACCG CCGAGGTGTTCCGGCGGCGGCACCAAGCTGACCGTGTGGGCCAGCCTAAGGCTGCCCCCAGCGTGACCCTGT TCCCCCCCAGCAGCGAGGAGCTGCAGGCCAACAAGGCCACCCTGGTGTGCCTGATCAGCGACTTCTACCCAG GCGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACCACCCCCAGC AAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCCGAGCAGTGAAGAGCCACAGG TCCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAAAAGACCGTGGCCCCAACCGAGTGCAGC
Antibody 6525	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	131: SYWIS
CDRH2	132: IIDPDDSKTNYSFQG
CDRH3	133:RSYYPMDY
CDRL1	134:TGTSSDVGVYNFVS

CDRL2	135: YVDNRPS
CDRL3	136: QSFDGFGIDMV
VH	137: QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKGLEWMGIIDPDDSKTNYSPSFQGGVTTISAD KSISTAYLQWSSLKASDTAMYYCARRSYYPMDYWGGGLTVTVSS
VL	138: DIALTQPASVSGSPGQSITISCTGTSSDVGVYNFVSWYQQHPGKAPKLMIIYYVDNRPSGVSNRFSGSKSGNTA SLTISGLQAEDEADYYCQSFDGFGIDMVFGGGTKLTVL
Heavy chain	139: QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWISWVRQMPGKGLEWMGIIDPDDSKTNYSPSFQGGVTTISAD KSISTAYLQWSSLKASDTAMYYCARRSYYPMDYWGGGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDY FPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	140: DIALTQPASVSGSPGQSITISCTGTSSDVGVYNFVSWYQQHPGKAPKLMIIYYVDNRPSGVSNRFSGSKSGNTA SLTISGLQAEDEADYYCQSFDGFGIDMVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTV AWKADSSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKSRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:137	141: CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTGGATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATCGATCCGGATGATAGCAAGACCAATTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGA TAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGGATACGGCCATGTATTATTGCGCGC GTCGTTCTTATTATCCTATGGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:138	142: GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTGAGAGCATTACCATCTCGTGACGG GTACTAGCAGCGATGTTGTTGGTGTTTATAATTTTGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAAA CTTATGATTTATTATGTTGATAATCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACC GCGAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTTTGATGGTTTTGG TATTGATATGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:139	143: CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTGGATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATCGATCCGGATGATAGCAAGACCAATTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGA TAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGGATACGGCCATGTATTATTGCGCGC GTCGTTCTTATTATCCTATGGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGT CCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTT AAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGATACCTTTC CGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCA CTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAGC X (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:140	144: GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTGAGAGCATTACCATCTCGTGACGG GTACTAGCAGCGATGTTGTTGGTGTTTATAATTTTGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAAA CTTATGATTTATTATGTTGATAATCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACC GCGAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTTTGATGGTTTTGG TATTGATATGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACG CTGTTTCCGCCGAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATC CGGGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGAGACCACCACACCC TCCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCACAC GAAGCTACAGCTGCCAGGTACGCATGAGGGGAGCACCGTGGAAGAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 6756	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	145: SYWIA
CDRH2	146: ILYPGDSDTNYSPSFQG
CDRH3	147: SKYGSFDY
CDRL1	148: TGTSSDVGGYNYVS
CDRL2	149: NVNSRPS
CDRL3	150: QSYDDGQDNEV

VH	151:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIAWVRQMPGKGLEWMGIIYPGDSDTNYSFQGGQVTISADKSISTAYLQWSSLKASDTAMYYCARSKYGSFDYWGQGLTVTVSS
VL	152:DIALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIYNVNSRPSGVSNRFGSGSKSGNTASLTISGLQAEDEADYYCQSYDDGQDNEVFGGGTLTVL
Heavy chain	153:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIAWVRQMPGKGLEWMGIIYPGDSDTNYSFQGGQVTISADKSISTAYLQWSSLKASDTAMYYCARSKYGSFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	154:DIALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMIYNVNSRPSGVSNRFGSGSKSGNTASLTISGLQAEDEADYYCQSYDDGQDNEVFGGGTLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:151	155:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGGTTCCGGATATTCTTTACTTCTTATTGGATTGCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGCATTATCTATCCGGGTGATAGCGATACCAATTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGATAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTCTAAGTATGGTTCTTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:152	156:GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTCAGAGCATTACCATCTCGTGACGGGTACTAGCAGCGATGTTGGTGGTTATAATTATGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAAACTTATGATTTATAATGTTAATTCTCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACCGCGAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTATGATGATGGTCAGGATAATGAGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:153	157:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAGGTTCCGGATATTCTTTACTTCTTATTGGATTGCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGCATTATCTATCCGGGTGATAGCGATACCAATTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGATAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTCTAAGTATGGTTCTTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGATACCTTTCCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCCATCAGACCTATATTTGCAACGTGAACCATAAACCAGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:154	158:GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTCAGAGCATTACCATCTCGTGACGGGTACTAGCAGCGATGTTGGTGGTTATAATTATGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAAACTTATGATTTATAATGTTAATTCTCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACCGCGAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTATGATGATGGTCAGGATAATGAGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACCGTGTTTCCGCCGAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCATGAGGGGAGCACCGTGGA AAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 6757	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	159: SYAMH
CDRH2	160: AISSSGSSTYYADSVKG
CDRH3	161:ESWFLDL
CDRL1	162:RASQISINWLA
CDRL2	163: LASSLQS
CDRL3	164:QYYDFSDT
VH	165:QVQLVESGGGLVQPGGSLRLSCAASGFTFTSYAMHWVRQAPGKGLEWVSAISSSGSSTYYADSVKGRFTISR

	NSKNTLYLQMNSLRAEDTAVYYCARESWFLDLWGQGLTVTVSS
VL	166:DIQMTQSPSSLSASVGDRVTITCRASQISNWLAWYQQKPGKAPKLLIYLASSLQSGVPSRFSGSGSGTDFTLTIS SSLQPEDFAVYYCQQYYDFSDTFGQGTKVEIK
Heavy chain	167:QVQLVESGGGLVQPGGSLRLSCAASGFTFTSYAMHWVRQAPGKGLEWVSAISSSGSSTYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCARESWFLDLWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	168:DIQMTQSPSSLSASVGDRVTITCRASQISNWLAWYQQKPGKAPKLLIYLASSLQSGVPSRFSGSGSGTDFTLTIS SSLQPEDFAVYYCQQYYDFSDTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNNFYPREAKVQWKVDN ALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEX (X can be C or A)
PN encoding SEQ ID NO:165	169:CAGGTGCAATTGGTGAAAGCGGCGGCGCCTGGTGCAACCGGGCGGCAGCCTGCGTCTGAGCTGCGC GGCCTCCGGATTTACCTTTACTTCTTATGCTATGCATTGGGTGCGCCAAGCCCCTGGGAAGGGTCTCGAGTGG GTGAGCGCTATCTCTTCTTCTGGTAGCTCTACCTATTATGCGGATAGCGTGAAAGGCCGTTTTACCATTTCACGT GATAATTCGAAAAACACCCTGTATCTGCAAATGAACAGCCTGCGTGCGGAAGATACGGCCGTGTATTATTGCGC GCGTGAGTCTTGGTTTCTTGATCTTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:166	170:GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCGAGCGTGGGTGATCGTGTGACCATTACCTGCA GAGCGAGCCAGTCTATTTCTAATTGGCTGGCTTGGTACCAGCAGAAACCAGGTAAAGCACCGAAACTATTAATT TATCTTGCTTCTTCTTTGCAAAGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCT GACCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTATTGCCAGCAGTATTATGATTTTTCTGATACCTTT GGCCAGGGTACGAAAGTTGAAATTA
PN encoding SEQ ID NO:167	171:CAGGTGCAATTGGTGAAAGCGGCGGCGCCTGGTGCAACCGGGCGGCAGCCTGCGTCTGAGCTGCGC GGCCTCCGGATTTACCTTTACTTCTTATGCTATGCATTGGGTGCGCCAAGCCCCTGGGAAGGGTCTCGAGTGG GTGAGCGCTATCTCTTCTTCTGGTAGCTCTACCTATTATGCGGATAGCGTGAAAGGCCGTTTTACCATTTCACGT GATAATTCGAAAAACACCCTGTATCTGCAAATGAACAGCCTGCGTGCGGAAGATACGGCCGTGTATTATTGCGC GCGTGAGTCTTGGTTTCTTGATCTTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGT CCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTT AAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGATACCTTTC CGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCA CTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAGC X (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:168	172:GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCGAGCGTGGGTGATCGTGTGACCATTACCTGCA GAGCGAGCCAGTCTATTTCTAATTGGCTGGCTTGGTACCAGCAGAAACCAGGTAAAGCACCGAAACTATTAATT TATCTTGCTTCTTCTTTGCAAAGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCT GACCATTAGCAGCCTGCAACCTGAAGACTTTGCGGTTTATTATTGCCAGCAGTATTATGATTTTTCTGATACCTTT GGCCAGGGTACGAAAGTTGAAATTAACGTACGGTGGCTGCTCCGAGCGTGTTTATTTTTCCGCCGAGCGATGA ACAACTGAAAAGCGGCACGGCGAGCGTGGTGTGCCTGCTGAACAACTTTATCCGCGTGAAGCGAAAGTTTCAG TGGAAGTAGACAACGCGCTGCAAAGCGGCAACAGCCAGGAAAGCGTGACCGAACAGGATAGCAAAGATAGCA CCTATTCTCTGAGCAGCACCTGACCCTGAGCAAAGCGGATTATGAAAAACATAAAGTGTATGCGTGCGAAGTG ACCCATCAAGGTCTGAGCAGCCCGGTGACTAAATCTTTAATCGTGGCGAGX (X can be TGC or GCC)
Antibody 6763	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	173: NYGMH
CDRH2	174: VSYAGSFTNYADSVKG
CDRH3	175:SWLFGYPDIFDY
CDRL1	176:TGTSSDVGGYNYVS
CDRL2	177: DVNNRPS
CDRL3	178:SSYDKFQTV
VH	179:QVQLVESGGGLVQPGGSLRLSCAASGFTFSNYGMHWVRQAPGKGLEWVSVSYAGSFTNYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCARSWLFGYPDIFDYWGQGLTVTVSS
VL	180:DIALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVNNRPSGVSNRFSGSKSGNTA SLTISGLQAEDEADYYCSSYDKFQTVFGGGTKLTVL

Heavy chain	181:QVQLVESGGGLVQPGGSLRLSCAASGFTFSNYGMHWVRQAPGKGLEWVSYSYAGSFTNYADSVKGRFTISRDN NSKNTLYLQMNSLRAEDTAVYYCARSWLFGYPDIFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	182:DIALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLMYDVNNRPSGVSNRFSGSKSGNTA SLTISGLQAEDEADYYCSSYDKFQTVFGGGTLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAW KADSSPVKAGVETTTTPSKQSNKNYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:179	183:CAGGTGCAATTGGTGAAAGCGGCGGCGGCCTGGTGCAACCGGGCGGCAGCCTGCGTCTGAGCTGCGC GGCCTCCGGATTTACCTTTTCTAATTATGGTATGCATTGGGTGCGCCAAGCCCCTGGGAAGGGTCTCGAGTGG GTGAGCGTTTCTTATGCTGGTAGCTTTACCAATTATGCGGATAGCGTGAAAGGCCGTTTTACCATTTCACGTGAT AATTCGAAAAACACCCTGTATCTGCAATGAACAGCCTGCGTGCGGAAGATACGGCCGTGTATTATTGCGCGCG TTCTTGGCTTTTTGGTTATCCTGATATTTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:180	184:GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTCAGAGCATTACCATCTCGTGACGG GTACTAGCAGCGATGTTGGTGGTTATAATTATGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAACTT ATGATTTATGATGTTAATAATCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACCGC GAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCTCTTCTTATGATAAGTTTCAGA CTGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:181	185:CAGGTGCAATTGGTGAAAGCGGCGGCGGCCTGGTGCAACCGGGCGGCAGCCTGCGTCTGAGCTGCGC GGCCTCCGGATTTACCTTTTCTAATTATGGTATGCATTGGGTGCGCCAAGCCCCTGGGAAGGGTCTCGAGTGG GTGAGCGTTTCTTATGCTGGTAGCTTTACCAATTATGCGGATAGCGTGAAAGGCCGTTTTACCATTTCACGTGAT AATTCGAAAAACACCCTGTATCTGCAATGAACAGCCTGCGTGCGGAAGATACGGCCGTGTATTATTGCGCGCG TTCTTGGCTTTTTGGTTATCCTGATATTTTTGATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTC GACCAAAGGTCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGG CTGCCTGGTTAAAGATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTG CATACCTTTCCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCA GCTTAGGCACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGGAA CCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:182	186:GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTCAGAGCATTACCATCTCGTGACGG GTACTAGCAGCGATGTTGGTGGTTATAATTATGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAACTT ATGATTTATGATGTTAATAATCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACCGC GAGCCTGACCATTAGCGGCCTGCAAGCGGAAGACGAAGCGGATTATTATTGCTCTTCTTATGATAAGTTTCAGA CTGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCC GCCGAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCC GTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGAGACCACCACACCCTCCAAACAA AGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACA GCTGCCAGGTCACGCATGAGGGGAGCACCGTGGA AAAAACC GTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7086	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:1
CDRH2	SEQ ID NO:2
CDRH3	SEQ ID NO:3
CDRL1	SEQ ID NO:4
CDRL2	SEQ ID NO:5
CDRL3	SEQ ID NO:6
VH	187:QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIGPFFGTANYAQKFQGRVTITA DESTSTAYMELSSLRSEDVAVYYCARDTPYFDYWGQGLTVTVSS
VL	188:DIELTQPPSVSVAPGQTARISCSGDSIPNYYVYVYQQKPGQAPVLVIYDDSNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQSFDSLNAEVFGGGTKLTVL
Heavy chain	189:QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIGPFFGTANYAQKFQGRVTITA DESTSTAYMELSSLRSEDVAVYYCARDTPYFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDY

	FPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	190:DIELTQPPSVSVAPGQTARISCSGDSIPNYYVYWYQQKPGQAPVLVIYDDSNRPSGIPERFSGSNSGNTATLTISGTQAEDEADYYCQSFDSLSNAEVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:187	191:CAGGTGCAATTGGTTTCAGTCTGGCGCGGAAGTGAAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAGCCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGATGGGCGGTATCGGTCCGTTTTTTGGCACTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCGGATGAAAGCACCCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCGCGCGTGATACTCCTTATTTTATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:188	192:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGGCGATTCTATTCTAATTATTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATGATGATTCTAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACCATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTTTGATTCTTCTCTTAATGCTGAGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:189	193:CAGGTGCAATTGGTTTCAGTCTGGCGCGGAAGTGAAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAGCCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGATGGGCGGTATCGGTCCGTTTTTTGGCACTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCGGATGAAAGCACCCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCGCGCGTGATACTCCTTATTTTATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGATACCTTTCGGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCCACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:190	194:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCGGCGATTCTATTCTAATTATTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATGATGATTCTAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGACCATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTTTGATTCTTCTCTTAATGCTGAGGTGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGACGCTGTTTCCGCCGAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGAGAGACCACACACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCATGAGGGGAGCACCGTGGAATAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7087	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	195: SYYIS
CDRH2	196: GIPIFGTANYAQKFQG
CDRH3	197: GEIWHVHQPYKSGVYGAAY
CDRL1	198: RASQGISNWLN
CDRL2	199: GTSSLQS
CDRL3	200: QQLDSFPAT
VH	201:QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYYISWVRQAPGQGLEWMGGIPIFGTANYAQKFQGRVTITADESTSTAYMELSSLRSEDNAVYYCARGEIWHVHQPYKSGVYGAAYWGQGTLVTVSS
VL	202:DIQMTQSPSSLSASVGDRVTITCRASQGISNWLNWYQQKPGKAPKLLIYGTSSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQLDSFPATFGQGTKVEIK
Heavy chain	203:QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYYISWVRQAPGQGLEWMGGIPIFGTANYAQKFQGRVTITADESTSTAYMELSSLRSEDNAVYYCARGEIWHVHQPYKSGVYGAAYWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	204:DIQMTQSPSSLSASVGDRVTITCRASQGISNWLNWYQQKPGKAPKLLIYGTSSLQSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQLDSFPATFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN

	ALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEX (X can be C or A)
PN encoding SEQ ID NO:201	205:CAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATTATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGATG GGCGGTATCATTCCGATTTTTGGCACTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCGG ATGAAAGCACCAGCACCGCGTATATGGAAGTACGAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCGC GCGTGGTGAGATTTGGCATGTTTCATCAGCCTTATAAGTCTGGTGTTTATGGTGCTGCTTATTGGGGCCAAGGCA CCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:202	206:GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCGAGCGTGGGTGATCGTGTGACCATTACCTGCA GAGCGAGCCAGGGTATTTCTAATTGGCTGAATTGGTACCAGCAGAAACCAGGTAAAGCACCGAACTATTAATT TATGGTACTTCTTCTTTGCAAAGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCT GACCATTAGCAGCCTGCAACCTGAAGACTTTGCGACTTATTATTGCCAGCAGCTTGATTCTTTTCCTGCTACCTT TGCCAGGGTACGAAAGTTGAAATTAA
PN encoding SEQ ID NO:203	207:CAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATTATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGATG GGCGGTATCATTCCGATTTTTGGCACTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCGG ATGAAAGCACCAGCACCGCGTATATGGAAGTACGAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCGC GCGTGGTGAGATTTGGCATGTTTCATCAGCCTTATAAGTCTGGTGTTTATGGTGCTGCTTATTGGGGCCAAGGCA CCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCAC CAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAAGATTATTTCCCGGAACCAGTCACCGTGAGCTGGAAC AGCGGGGCGCTGACCAGCGGCGTGACATACCTTTCCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGC AGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAA CACCAAAGTGGATAAAAAAGTGGAAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:204	208:GATATCCAGATGACCCAGAGCCCGTCTAGCCTGAGCGCGAGCGTGGGTGATCGTGTGACCATTACCTGCA GAGCGAGCCAGGGTATTTCTAATTGGCTGAATTGGTACCAGCAGAAACCAGGTAAAGCACCGAACTATTAATT TATGGTACTTCTTCTTTGCAAAGCGGGGTCCCGTCCCGTTTTAGCGGCTCTGGATCCGGCACTGATTTTACCCT GACCATTAGCAGCCTGCAACCTGAAGACTTTGCGACTTATTATTGCCAGCAGCTTGATTCTTTTCCTGCTACCTT TGCCAGGGTACGAAAGTTGAAATTAAACGTACGGTGGCTGCTCCGAGCGTGTTTATTTTCCGCCGAGCGATG AACAACTGAAAAGCGGCACGGCGAGCGTGGTGTGCCTGCTGAACAACTTTTATCCGCGTGAAGCGAAAGTTCA GTGGAAGTAGACAACGCGCTGCAAAGCGGCAACAGCCAGGAAAGCGTGACCGAACAGGATAGCAAAGATAG CACCTATTCTCTGAGCAGCACCTGACCCTGAGCAAAGCGGATTATGAAAAACATAAAGTGTATGCGTGCGAAG TGACCCATCAAGGTCTGAGCAGCCCGGTGACTAAATCTTTAATCGTGGCGAGX (X can be TGC or GCC)
Antibody 7091	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:77
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	209:QSWTDSPTLV
VH	210:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPSDSHTTYSFQGGQVTISADK SISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGTLVTVSS
VL	211:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQSWTDSPTLVFGGGTKLTVL
Heavy chain	212:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPSDSHTTYSFQGGQVTISADK SISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	213:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQSWTDSPTLVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID	214:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG

NO:210	GGCATTATCGATCCGTCTGATAGCCATACCACTTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGA TAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:211	215:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTGGACTGATTCTCCTAATACTCTTG TGTTTGGCGGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:212	216:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATCGATCCGTCTGATAGCCATACCACTTATTCTCCGAGCTTTCAGGGCCAGGTGACCATTAGCGCGGA TAAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGGTTTTGATCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGG TCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGT TAAAGATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGACATACCTTT CCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGC ACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAG CX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:213	217:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTGAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGTCTTGGACTGATTCTCCTAATACTCTTG TGTTTGGCGGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCC GAGCAGCGAAGAATTGCAGGCGAACAAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTG ACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGC AACAAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCACAGAAGCTACAGCT GCCAGGTCACGCATGAGGGGAGCACCGTGGA AAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7092	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:49
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	218:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTRYSPSFQGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGTLVTVSS
VL	219:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCGTYDIESYVFGGGTKLTVL
Heavy chain	220:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTRYSPSFQGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIWGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	221:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCGTYDIESYVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADS SPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:218	222:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATCGATCCGGATGATAGCTATACCCGTTATTCTCCGAGCTTTCAGGGACAGGTGACCATTAGCGCGGATAA AAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTT ATGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA

PN encoding SEQ ID NO:219	223:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGGTACTTATGATATTGAGTCTTATGTGTTTG GCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:220	224:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATCGATCCGGATGATAGCTATAACCGTTATTCTCCGAGCTTTCAGGGACAGGTGACCATTAGCGCGGATAA AAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTT ATGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCC AAGCGTGTTCGCTGGCTCCGAGCAGCAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAA AGATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGCATACCTTTCCG GCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCACT CAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:221	225:GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGGTACTTATGATATTGAGTCTTATGTGTTTG GCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCCGAGCA GCGAAGAATTGCAGGCGAACAAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGACAGT GGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGCAACAA CAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCACAGAAGCTACAGTGCCA GGTCACGCATGAGGGGAGCACCGTGGAAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7093	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:33
CDRH2	226: HIFSDDDKYYSTSLKT
CDRH3	SEQ ID NO:35
CDRL1	SEQ ID NO:36
CDRL2	SEQ ID NO:37
CDRL3	SEQ ID NO:38
VH	227:QVQLKESGPALVKPTQTLTLCTFSGFSLSSTSGGGVSWIRQPPGKALEWLAHIFSDDDKYYSTSLKTRLTISKDT SKNQVVLMTNMDPVDATYYCARGPYGFDSWGQGTLLTVSS
VL	228:DIALTQPASVSGSPGQSITISCTGTSSDIGTYNYVSWYQQHPGKAPKLMYDDSNRPSGVSNRFSKSGNTAS LTISGLQAEDEADYYCQSYDSQSIVFGGGTKLTVL
Heavy chain	229:QVQLKESGPALVKPTQTLTLCTFSGFSLSSTSGGGVSWIRQPPGKALEWLAHIFSDDDKYYSTSLKTRLTISKDT SKNQVVLMTNMDPVDATYYCARGPYGFDSWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	230:DIALTQPASVSGSPGQSITISCTGTSSDIGTYNYVSWYQQHPGKAPKLMYDDSNRPSGVSNRFSKSGNTAS LTISGLQAEDEADYYCQSYDSQSIVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWK ADSSPVKAGVETTTPSKQSNNKYAASSYLSLTPEQWWSHRYSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:227	231:CAGGTGCAATTGAAAGAAAGCGGCCCGGCCCTGGTGAAACCGACCCAAACCCTGACCCTGACCTGTACCT TTTCCGGATTTAGCCTGTCTACTTCTGGTGGTGGTGTGCTTGGATTGCGCCAGCCGCTGGGAAAGCCCTCGAG TGGCTGGCTCATATCTTTCTGATGATGATAAGTATTATAGCACCAGCCTGAAAACGCGTCTGACCATTAGCAAA GATACTTCGAAAAATCAGGTGGTGCTGACTATGACCAACATGGACCCGGTGGATACGGCCACCTATTATTGCGC GCGTGGTCCTTATGGTTTTGATTCTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:228	232:GATATCGCACTGACCCAGCCAGCTTCAGTGAGCGGCTCACCAGGTCAGAGCATTACCATCTCGTGTACGG GTACTAGCAGCGATATTGGTACTTATAATTATGTGTCTTGGTACCAGCAGCATCCCGGGAAGGCGCCGAACTT ATGATTTATGATGATTCTAATCGTCCCTCAGGCGTGAGCAACCGTTTTAGCGGATCCAAAAGCGGCAACACCGC

47

PN encoding SEQ ID NO:243	247:CAGGTGCAATTGAAAGAAAGCGGCCCGGCCCTGGTGAAACCGACCCAAACCCTGACCCTGACCTGTACCT TTTCCGGATTTAGCCTGTCTACTTCTGGTATGTCTGTGGGTGGATTGCGCCAGCCGCCTGGGAAAGCCCTCGAG TGGCTGGCTCTTATCGATTGGGATGAGGATAAGTCTTATAGCACCAGCCTGAAAACGCGTCTGACCATTAGCAA AGATACTTCGAAAAATCAGGTGGTGTCTGACTATGACCAACATGGACCCGGTGGATACGGCCACCTATTATTGCG CGCGTTATAATTGGTATAATCCTCCTGGTTTTGATAATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCG TCGACCAAAGGTCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTG GGCTGCCTGGTTAAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGC GTGCATACCTTTCCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCA GCAGCTTAGGCACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTG GAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:244	248:GATATCGTGCTGACCCAGCCGCCTTCAGTGAGTGGCGCACCAGGTCAGCGTGTGACCATCTCGTGTAGCG GCAGCAGCAGCAACATTGGTTCTAATTATGTGTCTTGGTACCAGCAGTTGCCCGGGACGGCGCCGAAACTTCT GATTTATCGTAATGATAAGCGTCCCTCAGGCGTGCCGGATCGTTTTAGCGGATCCAAAAGCGGCACCAGCGCG AGCCTTGCGATTACGGGCCTGCAAAGCGAAGACGAAGCGGATTATTATTGCCAGTCTGCTGATTCTTCTTCTAT GGTGTGTTGGCGGCGGCACGAAGTTAACC GTTCTTGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCG CCGAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCG TGACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAA GCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAG CTGCCAGGTCACGCATGAGGGGAGCACCGTGGAAAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7821	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:1
CDRH2	SEQ ID NO:119
CDRH3	SEQ ID NO:3
CDRL1	SEQ ID NO:4
CDRL2	SEQ ID NO:5
CDRL3	SEQ ID NO:6
VH	SEQ ID NO:121
VL	SEQ ID NO:188
Heavy chain	249:QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAIWVRQAPGQGLEWMGNIGPFFGIANYAQKFQGRVTITAD ESTSTAYMELSSLRSED TAVYYCARDTPYFDYWGGQTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFP EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:190
PN encoding SEQ ID NO:121	SEQ ID NO:125
PN encoding SEQ ID NO:188	SEQ ID NO:192
PN encoding SEQ ID NO:249	250:CAGGTGCAATTGGTTCAGTCTGGCGCGGAAGTGAAAAAACCGGGCAGCAGCGTGAAAGTGAGCTGCAAAG CCTCCGGAGGCACTTTTTCTTCTTATGCCATTTCTTGGGTGCGCCAAGCCCCTGGGCAGGGTCTCGAGTGGAT GGGCAATATCGGTCCGTTTTTTGGCATTGCGAATTACGCGCAGAAGTTTCAGGGCCGGGTGACCATTACCGCG GATGAAAGCACCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGTAGCGAAGATACGGCCGTGTATTATTGCG CGCGTGATACTCCTTATTTTATTATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGT CCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTT AAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGATACCTTTC CGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCA CTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGGAACCGAAAAGC X (X can be TGC, GAATTC or TGCGAATTC)

PN encoding SEQ ID NO:190	SEQ ID NO:194
Antibody 7865	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:1
CDRH2	SEQ ID NO:2
CDRH3	SEQ ID NO:3
CDRL1	SEQ ID NO:4
CDRL2	SEQ ID NO:5
CDRL3	SEQ ID NO:120
VH	SEQ ID NO:187
VL	SEQ ID NO:122
Heavy chain	SEQ ID NO:189
Light chain	251:DIELTQPPSVSVAPGQTARISCSGDSIPNYYVYWYQQKPGQAPVLVIYDDSNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTYDDGSTAEVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:187	SEQ ID NO:191
PN encoding SEQ ID NO:122	SEQ ID NO:126
PN encoding SEQ ID NO:189	SEQ ID NO:193
PN encoding SEQ ID NO:251	252:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTATTCTAATTATTATGTTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATG ATGATTCTAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTATGATGATGGTTCTACTGCTGAGG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCC GAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTG ACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCCTCCAAACAAAGC AACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCT GCCAGGTCACGCATGAGGGGAGCACCGTGAAAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7829	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:62
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:209
VH	253:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSQTAYSPSFQGGQVTISADK SISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVT VSS
VL	SEQ ID NO:211
Heavy chain	254:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSQTAYSPSFQGGQVTISADK SISTAYLQWSSLKASDTAMYYCARYMMRGFDHWGQGT LVT VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:213
PN encoding	255:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG

SEQ ID NO:253	GTTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATTGATCCTACTGATTCTCAGACTGCTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGAT AAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGTTTGGTTCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:211	SEQ ID NO:215
PN encoding SEQ ID NO:254	256:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATTGATCCTACTGATTCTCAGACTGCTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGAT AAAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGC GTTATATGATGCGTGTTTGGTTCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGG TCCAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGT TAAAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGCATACCTTT CCGGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGC ACTCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGAACCGAAAAG CX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:213	SEQ ID NO:217
Antibody 7830	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:95
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:209
VH	257:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSYTVYSPSFQGGQVTISADK SISTAYLQWSSLKASDTAMYCYCARYMMRGFDHWGQGTLLTVSS
VL	SEQ ID NO:211
Heavy chain	258:QVQLVQSGAEVKKPGESLKISCKGSGYSFTSYIIGWVRQMPGKGLEWMGIIDPTDSYTVYSPSFQGGQVTISADK SISTAYLQWSSLKASDTAMYCYCARYMMRGFDHWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:213
PN encoding SEQ ID NO:257	259:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATTGATCCTACTGATTCTTATACTGTTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATA AAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGT TATATGATGCGTGTTTGGTTCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:211	SEQ ID NO:215
PN encoding SEQ ID NO:258	260:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCCTTTACTTCTTATTATATTGGTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATG GGCATTATTGATCCTACTGATTCTTATACTGTTTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATA AAAGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGT TATATGATGCGTGTTTGGTTCATTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTC CAAGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTA AAGATTATTTCCCGGAACCAAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGCATACCTTTCC

	GGCGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCAC TCAGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGGAACCGAAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
PN encoding SEQ ID NO:213	SEQ ID NO:217
Antibody 7871	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:77
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:66
VH	SEQ ID NO:210
VL	261:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTDWDTGESGVFGGGTKLTVL
Heavy chain	SEQ ID NO:212
Light chain	262:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTDWDTGESGVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTPSKQSNKYYAASSYLSLTPEQWKSRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:210	SEQ ID NO:214
PN encoding SEQ ID NO:261	263:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATACTGGTGAGTCTGGTGTG TTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:212	SEQ ID NO:216
PN encoding SEQ ID NO:262	264:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATACTGGTGAGTCTGGTGTG TTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCCGA GCAGCGAAGAATTGCAGGCGAACAAGCGACCCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGAC AGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGAGACCACCACACCCTCCAAACAAAGCAA CAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCACAGAAGCTACAGCTGC CAGGTCACGCATGAGGGGAGCACCGTGGAAAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7872	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:77
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:78
VH	SEQ ID NO:210
VL	265:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTDWDLPHGLVFGGGTKLTVL

Heavy chain	SEQ ID NO:212
Light chain	266:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQTDILPHGLVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKA DSSPVKAGVETTTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:210	SEQ ID NO:214
PN encoding SEQ ID NO:265	267:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATATTCTTCCTCATGGTCTTG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:212	SEQ ID NO:216
PN encoding SEQ ID NO:266	268:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGACTTGGGATATTCTTCCTCATGGTCTTG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCC GAGCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTG ACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGC AACAAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGAAGTCCCACAGAAGCTACAGCT GCCAGGTCACGCATGAGGGGAGCACCGTGGA AAAAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7873	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:61
CDRH2	SEQ ID NO:77
CDRH3	SEQ ID NO:63
CDRL1	SEQ ID NO:64
CDRL2	SEQ ID NO:65
CDRL3	SEQ ID NO:89
VH	SEQ ID NO:210
VL	269:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQAWTDSPTGLVFGGGTKLTVL
Heavy chain	SEQ ID NO:212
Light chain	270:DIELTQPPSVSVAPGQTARISCSGDSLGDYYAYWYQQKPGQAPVLVIYKDNNRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCQAWTDSPTGLVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWK ADSSPVKAGVETTTTPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:210	SEQ ID NO:214
PN encoding SEQ ID NO:269	271:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGGCTTGGACTGATTCTCCTACTGGTCTTG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:212	SEQ ID NO:216
PN encoding	272:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG

SEQ ID NO:270	GCGATTCTCTTGGTGATTATTATGCTTATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATAATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCCAGGCTTGGACTGATTCTCCTACTGGTCTTG TGTTTGGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCC GAGCAGCGAAGAATTGCAGGCGAACAAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTG ACAGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCAAACAAAGC AACAAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCACAGAAGCTACAGCT GCCAGGTCACGCATGAGGGGAGCACCGTGGAACCAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7832	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:18
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	273:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSS
VL	SEQ ID NO:219
Heavy chain	274:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPDDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:221
PN encoding SEQ ID NO:273	275:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGATGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:219	SEQ ID NO:223
PN encoding SEQ ID NO:274	276:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGATGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCAA GCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAAG ATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGCATACCTTTCCGGC GGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCACTCA GACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
Antibody 7909	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:107
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	277:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPQDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSS

VL	SEQ ID NO:219
Heavy chain	278:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPQDSYTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:221
PN encoding SEQ ID NO:277	279:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTCAGGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:219	SEQ ID NO:223
PN encoding SEQ ID NO:278	280:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTCAGGATTCTTATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTCACCATTAGCGCGGATAAAA GCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTAT GAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCAA GCGTGTTTCCGCTGGCTCCGAGCAGCAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAAG ATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGACATACCTTTCCGGC GGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCACTCA GACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGATAAAAAAGTGGAACCGAAAAGCX (X can be TGC, GAATTC or TGCGAATTC)
Antibody 7910	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:113
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:22
VH	281:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPEDSHTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSS
VL	SEQ ID NO:219
Heavy chain	282:QVQLVQSGAEVKKPGESLKISCKGSGYSFTNYISWVRQMPGKGLEWMGIIDPEDSHTEYSPSFQGGQVTISADKS ISTAYLQWSSLKASDTAMYYCARYEYGGFDIW/GQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPE PVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (X can be C, EF or CEF)
Light chain	SEQ ID NO:221
PN encoding SEQ ID NO:281	283:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGAGGATTCTCATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAAA AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCA
PN encoding SEQ ID NO:219	SEQ ID NO:223
PN encoding SEQ ID NO:282	284:CAGGTGCAATTGGTTCAGAGCGGCGCGGAAGTGAAAAAACCGGGCGAAAGCCTGAAAATTAGCTGCAAAG GTTCCGGATATTCTTTACTAATTATATTTCTTGGGTGCGCCAGATGCCTGGGAAGGGTCTCGAGTGGATGGGC ATTATTGATCCTGAGGATTCTCATACTGAGTATTCTCCTTCTTTTCAGGGTCAGGTGACCATTAGCGCGGATAAA

	AGCATTAGCACCGCGTATCTTCAATGGAGCAGCCTGAAAGCGAGCGATACGGCCATGTATTATTGCGCGCGTTA TGAGTATGGTGGTTTTGATATTTGGGGCCAAGGCACCCTGGTGACGGTTAGCTCAGCGTCGACCAAAGGTCCA AGCGTGTTTCCGCTGGCTCCGAGCAGCAAAAGCACCAGCGGCGGCACGGCTGCCCTGGGCTGCCTGGTTAAA GATTATTTCCCGGAACCAGTCACCGTGAGCTGGAACAGCGGGGCGCTGACCAGCGGCGTGCATACCTTTCCGG CGGTGCTGCAAAGCAGCGGCCTGTATAGCCTGAGCAGCGTTGTGACCGTGCCGAGCAGCAGCTTAGGCACTC AGACCTATATTTGCAACGTGAACCATAAACCGAGCAACACCAAAGTGGATAAAAAAGTGGAACCGAAAAGCX (X can be TGC, GAATTC or TCGAATTC)
Antibody 7876	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:49
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:50
VH	SEQ ID NO:218
VL	285:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCATWGSEDQVFGGGTKLTVL
Heavy chain	SEQ ID NO:220
Light chain	286:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCATWGSEDQVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKAD SSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:218	SEQ ID NO:222
PN encoding SEQ ID NO:285	287:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGCTACTTGGGGTTCTGAGGATCAGGTGTTT GGCGGCGGCACGAAGTTAACCGTTCTT
PN encoding SEQ ID NO:220	SEQ ID NO:224
PN encoding SEQ ID NO:286	288:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCGCTACTTGGGGTTCTGAGGATCAGGTGTTT GGCGGCGGCACGAAGTTAACCGTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCCGAGC AGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGACAG TGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCAACACACCCTCCAAACAAAGCAACA ACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAAGTCCCACAGAAGCTACAGCTGCCA GGTCACGCATGAGGGGAGCACCGTGGAACCAACCGTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Antibody 7878	SEQUENCE IDENTIFIER (SEQ ID NO:) OR COMMENTS/DETAILS
CDRH1	SEQ ID NO:17
CDRH2	SEQ ID NO:49
CDRH3	SEQ ID NO:19
CDRL1	SEQ ID NO:20
CDRL2	SEQ ID NO:21
CDRL3	SEQ ID NO:101
VH	SEQ ID NO:218
VL	289:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS

	GTQAEDEADYYCSTWDIEPTYVFGGGTKLTVL
Heavy chain	SEQ ID NO:220
Light chain	290:DIELTQPPSVSVAPGQTARISCSGDNIGNSYVHWYQQKPGQAPVLVIYKDNDRPSGIPERFSGSNSGNTATLTIS GTQAEDEADYYCSTWDIEPTYVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKAD SSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTEX (X can be CS or A)
PN encoding SEQ ID NO:218	SEQ ID NO:222
PN encoding SEQ ID NO:289	291:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCTCTACTTGGGATATTGAGCCTACTTATGTGT TTGGCGGCGGCACGAAGTTAACC GTTCTT
PN encoding SEQ ID NO:220	SEQ ID NO:224
PN encoding SEQ ID NO:290	292:GATATCGAACTGACCCAGCCGCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGCGTATCTCGTGTAGCG GCGATAATATTGGTAATTCTTATGTTTCATTGGTACCAGCAGAAACCCGGGCAGGCGCCAGTTCTTGTGATTTATA AGGATAATGATCGTCCCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACACCGCGACCCTGAC CATTAGCGGCACTCAGGCGGAAGACGAAGCGGATTATTATTGCTCTACTTGGGATATTGAGCCTACTTATGTGT TTGGCGGCGGCACGAAGTTAACC GTTCTTGGCCAGCCGAAAGCCGCACCGAGTGTGACGCTGTTTCCGCCGA GCAGCGAAGAATTGCAGGCGAACAAGCGACCCTGGTGTGCCTGATTAGCGACTTTTATCCGGGAGCCGTGAC AGTGGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGAGACCACCACACCCTCCAAACAAAGCAA CAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGC CAGGTCACGCATGAGGGGAGCACCGTGGA AAAAACC GTTGCGCCGACTGAGX (X can be TGCAGC or GCC)
Human (<i>Homo sapiens</i>) C5	296:MGLLGILCFLIFLGKTWGEQTYVISAPKIFRVGASENIVIQVYGYTEAFDATISIKSYPDKKFSYSSGHVHLSSSENK FQNSAILTIQPKQLPGGQNPVSYVYLEVVSKHFSKSKRMPITYDNGFLFIHTDKPVYTPDQSVKVRVYSLNDDLKPAK RETVLTFIDPEGSEVDMVEEIDHIGISFPDFKIPSNPRYGMWTIKAKYKEDFSTTGTAYFEVKEYVLPHFVSVSIEPEYNF IGYKNFKNFEITIKARYFYNKVVTEADVYITFGIREDLKDDQKEMMQTAMQNTMLINGIAQVTFDSETAVKELSYYSLED LNNKYLYIAVTVIESTGGFSEEAIEPIGIKYLSPYKLNLVATPLFLKPGIPYPIKVQKDSLQDLVGGVPVTLNAQTIDVN QETSDLDPSKSVTRVDDGVASFVNLNPSGVTVLEFNVKTDAPDLPEENQAREGYRAIAYSSLSQSYLYIDWTDNHKA LLVGEHLNIIVTPKSPYIDKITHYNYLILSKGKIIHFGTREKFSDASYQSINIPVTQNMVPSSRLLVYIVTGEQTAELVSD SVWLNIEEKCGNQLQVHLSPDADAYSPGQTVSLNMGMDSWVALAAVDSAVYGVQRGAKKPLERVVFQFLEKSDL GCGAGGGLNNANVFHLAGLTFLTNANADDSQENDEPCKEILRPRRTLQKKIEEIAAKYKHSVVKCCYDGACVNNDE TCEQRAARISLGPRCIKAFTECCVVASQLRANISHKDMQLGRLHMKTLTPVSKPEIRSYFPESWLWEVHLVPRRKQL QFALPDSLTTWEIQGVGISNTGICVADTVKAKVFKDVFLEMNIPYSVVRGEQIQLKGTVYNYRTSGMQFCVKMSAVE GICTSESPVIDHQGTKSSKCVRQKVEGSSSHLVFTFTVLPLEIGHLNINFSLETWFGKEILVKTLRVVPEGVKRESYSGV TLDPRGIYGTISRKEFPYRIPDLVPKTEIKRILSVKLLVGEILSAVLSQEGINILTHLPKGSAAELMSVVPVYFVHY LETGNHWNIFHSDPLIEKQKLKKKLKEGMLSIMSYRNADYSYSVWKGGSASTWLTAFAALRVLGQVNKYVEQNQNSIC NSLLWLVENYQLDNGSFKENSQYQPIKLQGTLPVEARENSLYLTAFTVIGIRKAFDICPLVKIDTALIKADNFFLENTLPA QSTFTLAISAYALSLGDKTHPQFRSIVSALKREALVKGNPPIYRFWKDNLQHKDSSVPNTGTARMVETTAYALLTSLNL KDINYVNPVIKWLSEEQRYGGGFYSTQDTINAIEGLTEYSLLVKQLRLSMDIDVSYKHKGALHNYKMTDKNFLGRPVE VLLNDDLIVSTGFGSGLATVHVTTVVHKTSTSEEVCSFYLKIDTQDIEASHYRGYGNSDYKRIVACASYKPSREESSSG SSHAVMDISLPTGISANEEDLKALVEGVDQLFTDYQIKDGHVILQLNSIPSSDFLCVRFRI FELFEVGFLSPATFTVY EY HRPDKQCTMFYSTSNIKIQKVCEGAACKCEADCGQMQEELDTISAETRKTACKPEIAYAYKVSITSITVENVFKY KATLLDIYKTGEAVAEKDSEITFIKKVTCTNAELVKGRQYLIMGKEALQIKYNFSFRYIYPLDSL TWIEYWPRD TTCSSC QAFLANLDEFAEDIFLNGC
Cynomolgus Macaque (<i>Macaca fascicularis</i>)	297:MGLLGILCFLIFLGKTWGEQTYVISAPKIFRVGASENIVIQVYGYTEAFDATISIKSYPDKKFSYSSGHVHLSSSENK FQNSAVLTIQPKQLPGGQNPVSYVYLEVVSKHFSKSKIPITYDNGFLFIHTDKPVYTPDQSVKVRVYSLNDDLKPAK RETVLTFIDPEGSEIDMVEEIDHIGISFPDFKIPSNPRYGMWTIQAKYKEDFSTTGTAF FEVKEYVLPHFVS VSEPE SNF IGYKNFKNFEITIKARYFYNKVVTEADVYITFGIREDLKDDQKEMMQTAMQNTMLINGIAQVTFDSETAVKELSYYSLED

C5	LNNKYLYIAVTVIESTGGFSEEAIEPGIKYVLSPLYKLNLVATPLFLKPGIPYSIKVQVKDALDQLVGGVPVTLNAQTIDVN QETSDLEPRKSVTRVDDGVASFVNLPSPGVTVLEFNVKTDAPDLPDENQAREGYRAIAYSSLSQSYLYIDWTDNHKA LLVGEYLNIIIVTPKSPYIDKITHYNYLILSKGKIIHFGTREKLSDASYQSINIPVTQNMVPSSRLLVYYIVTGEQTAELVSDS VWLNIEEKCGNQLQVHLSPADTYSFGQTVSLNMVGTGMDSWVALTAVDSAVYGVQRRAKKPLERVVFQFLEKSDLG CGAGGGLNNANVFHLAGLTFLTANADDSQENDEPCKEIRPRRMLQEKIEEIAAKYKHLVVKCCYDGVRIHDETC EQRAARISVGPRCVKAFTECCVVASQLRANNSHKDLQLGRLHMKTLTPVSKPEIRSYFPESWLWEVHLVPRRKQLQ FALPDSVTTWEIQGVGISNSGICVADTIKAKVFKDVFLEMNIPYSVVRGEQVQLKGTVYNYRTSGMQFCVKMSAVEGI CTSESPVIDHQGTKSSKCVRQKVEGSSNHLVTFTVLPLEIGLQNINFSLETSFGKEILVKSLRVVPEGVKRESYSGITLD PRGIYGTISRRKEFPYRIPLDLVPKTEIKRILSVKGLLVGEILSAVLSREGINILTHLPKGSAAELMSVVPVFYVFHYLET GNHWNIFHSDPLIEKRNLEKKLKEGMVSIMSRYNADYSYSVWKGGSASTWLTAFALRVLGQVHKYVEQNQNSICNS LLWLVENYQLDNGSFKENSQYQPIKLQGTLPVEARENSLYLTAFTVIGIRKAFDICPLVKINTALIKADTFLENTLPAQS TFTLAISAYALSLGDKTHPQFRSIVSALKREALVKGNPPIYRFWKDSLQHKDSSVPNTGTARMVETTAYALLTSLNLKD INYVNPPIKWLLSEEQRYGGGFYSTQDTINAIEGLTEYSLLVKQLRLNMDIDVAYKHKGPLHNYKMTDKNFLGRPVEVLL NDDL VVSTGFGSGLATVHVTTVVHKTSTSEEVCSFYLKIDTQDIEASHYRGYGNSDYKRIVACASYKPSKEESSSGSS HAVMDISLPTGINANEEDLKALVEGVDQLFTDYQIKDGHVILQLNSIPSSDFLCVRFRI FELFEVGFLSPATFTVYEHYR PDKQCTMFYSTSNIKIQKVCEGATCKCIEADCGMQKELDLTISAETRKQTACNPEIAYAYKVIITSITTENVFVKYKAT LLDIYKTGEAVAEKDSEITFIKKVTCTNAELVKGRQYLIMGKEALQIKYNFTFRYIYPLDSL TWIEYWPRD T T C S S C Q A F LANLDEFAEDIFLNGC
----	--

Other antibodies of the invention include those where the amino acids or nucleic acids encoding the amino acids have been mutated, yet have at least 60, 70, 80, 90 or 95 percent identity to the sequences described in Table 1. In some embodiments, it include mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated in the variable regions when compared with the variable regions depicted in the sequence described in Table 1, while retaining substantially the same therapeutic activity.

Since each of these antibodies can bind to C5, the VH, VL, full length light chain, and full length heavy chain sequences (amino acid sequences and the nucleotide sequences encoding the amino acid sequences) can be "mixed and matched" to create other C5-binding antibodies of the invention. Such "mixed and matched" C5-binding antibodies can be tested using the binding assays known in the art (e.g., ELISAs, and other assays described in the Example section). When these chains are mixed and matched, a VH sequence from a particular VH/VL pairing should be replaced with a structurally similar VH sequence. Likewise a full length heavy chain sequence from a particular full length heavy chain / full length light chain pairing should be replaced with a structurally similar full length heavy chain sequence. Likewise, a VL sequence from a particular VH/VL pairing should be replaced with a structurally similar VL sequence. Likewise a full length light chain sequence from a particular full length heavy chain / full length light chain pairing should be replaced with a structurally similar full length light chain sequence. Accordingly, in one aspect, the invention provides an isolated monoclonal antibody or antigen binding region thereof having: a heavy chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, and 281; and a light chain variable region comprising an

amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, and 289; wherein the antibody specifically binds to C5 (e.g., human and/or cynomolgus C5).

In another aspect, the invention provides (i) an isolated monoclonal antibody having: a full length heavy chain comprising an amino acid sequence that has been optimized for expression in the cell of a mammalian selected from the group consisting of SEQ ID NOs: 9, 25, 41, 53, 69, 81, 97, 109, 115, 123, 139, 153, 167, 181, 189, 203, 212, 220, 229, 243, 249, 254, 258, 274, 278, and 282; and a full length light chain comprising an amino acid sequence that has been optimized for expression in the cell of a mammalian selected from the group consisting of SEQ ID NOs: 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154, 168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, and 290; or (ii) a functional protein comprising an antigen binding portion thereof.

In another aspect, the present invention provides C5-binding antibodies that comprise the heavy chain and light chain CDR1s, CDR2s and CDR3s as described in Table 1, or combinations thereof. The amino acid sequences of the VH CDR1s of the antibodies are shown in SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235. The amino acid sequences of the VH CDR2s of the antibodies are shown in SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236. The amino acid sequences of the VH CDR3s of the antibodies are shown in SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237. The amino acid sequences of the VL CDR1s of the antibodies are shown in SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238. The amino acid sequences of the VL CDR2s of the antibodies are shown in SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239. The amino acid sequences of the VL CDR3s of the antibodies are shown in SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240. The CDR regions are delineated using the Kabat system (Kabat, E. A., *et al.*, 1991 Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242).

Given that each of these antibodies can bind to C5 and that antigen-binding specificity is provided primarily by the CDR1, 2 and 3 regions, the VH CDR1, 2 and 3 sequences and VL CDR1, 2 and 3 sequences can be "mixed and matched" (*i.e.*, CDRs from different antibodies can be mixed and match, although each antibody must contain a VH CDR1, 2 and 3 and a VL CDR1, 2 and 3 to create other C5-binding binding molecules of the invention. Such "mixed and matched" C5-binding antibodies can be tested using the binding assays known in the art and those described in the Examples (e.g., ELISAs). When VH CDR sequences are mixed and matched, the CDR1, CDR2 and/or CDR3 sequence from a particular VH sequence should be replaced with a structurally similar CDR sequence(s). Likewise, when VL CDR sequences are mixed and matched, the CDR1, CDR2 and/or CDR3

sequence from a particular VL sequence should be replaced with a structurally similar CDR sequence(s). It will be readily apparent to the ordinarily skilled artisan that novel VH and VL sequences can be created by substituting one or more VH and/or VL CDR region sequences with structurally similar sequences from the CDR sequences shown herein for monoclonal antibodies of the present invention.

Accordingly, the present invention provides an isolated monoclonal antibody or antigen binding region thereof comprising a heavy chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235; a heavy chain variable region CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236; a heavy chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237; a light chain variable region CDR1 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238; a light chain variable region CDR2 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239; and a light chain variable region CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240; wherein the antibody specifically binds C5.

In a specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO:1; a heavy chain variable region CDR2 of SEQ ID NO: 2; a heavy chain variable region CDR3 of SEQ ID NO: 3; a light chain variable region CDR1 of SEQ ID NO: 4; a light chain variable region CDR2 of SEQ ID NO: 5; and a light chain variable region CDR3 of SEQ ID NO: 6. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 18; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 22.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 33; a heavy chain variable region CDR2 of SEQ ID NO: 34; a heavy chain variable region CDR3 of SEQ ID NO: 35; a light chain variable region CDR1 of SEQ ID NO: 36; a light chain variable region CDR2 of SEQ ID NO: 37; and a light chain variable region CDR3 of SEQ ID NO: 38. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 49; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of

SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 50.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 62; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 66. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 78.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 89. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 62; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 89.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 95; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 89. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 49; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 101.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 107; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO:

21; and a light chain variable region CDR3 of SEQ ID NO: 22. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 107; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 101.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 113; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 22. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 113; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 101.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 1; a heavy chain variable region CDR2 of SEQ ID NO: 119; a heavy chain variable region CDR3 of SEQ ID NO: 3; a light chain variable region CDR1 of SEQ ID NO: 4; a light chain variable region CDR2 of SEQ ID NO: 5; and a light chain variable region CDR3 of SEQ ID NO: 120. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 131; a heavy chain variable region CDR2 of SEQ ID NO: 132; a heavy chain variable region CDR3 of SEQ ID NO: 133; a light chain variable region CDR1 of SEQ ID NO: 134; a light chain variable region CDR2 of SEQ ID NO: 135; and a light chain variable region CDR3 of SEQ ID NO: 136.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 145; a heavy chain variable region CDR2 of SEQ ID NO: 146; a heavy chain variable region CDR3 of SEQ ID NO: 147; a light chain variable region CDR1 of SEQ ID NO: 148; a light chain variable region CDR2 of SEQ ID NO: 149; and a light chain variable region CDR3 of SEQ ID NO: 150. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 159; a heavy chain variable region CDR2 of SEQ ID NO: 160; a heavy chain variable region CDR3 of SEQ ID NO: 161; a light chain variable region CDR1 of SEQ ID NO: 162; a light chain variable region CDR2 of SEQ ID NO: 163; and a light chain variable region CDR3 of SEQ ID NO: 164.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 173; a heavy chain variable region CDR2 of SEQ ID NO: 174; a heavy chain variable region CDR3 of SEQ ID NO: 175; a light chain variable region CDR1 of SEQ ID NO: 176; a light chain variable region CDR2 of SEQ ID NO: 177; and a light chain variable region CDR3 of SEQ ID NO: 178. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 195; a heavy chain variable region CDR2 of SEQ ID NO: 196; a heavy chain variable region CDR3 of SEQ ID NO: 197; a light chain variable region CDR1 of SEQ ID NO: 198; a light chain variable region CDR2 of SEQ ID NO: 199; and a light chain variable region CDR3 of SEQ ID NO: 200.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 209. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 49; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 22.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 33; a heavy chain variable region CDR2 of SEQ ID NO: 226; a heavy chain variable region CDR3 of SEQ ID NO: 35; a light chain variable region CDR1 of SEQ ID NO: 36; a light chain variable region CDR2 of SEQ ID NO: 37; and a light chain variable region CDR3 of SEQ ID NO: 38. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 235; a heavy chain variable region CDR2 of SEQ ID NO: 236; a heavy chain variable region CDR3 of SEQ ID NO: 237; a light chain variable region CDR1 of SEQ ID NO: 238; a light chain variable region CDR2 of SEQ ID NO: 239; and a light chain variable region CDR3 of SEQ ID NO: 240.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 1; a heavy chain variable region CDR2 of SEQ ID NO: 119; a heavy chain variable region CDR3 of SEQ ID NO: 3; a light chain variable region CDR1 of SEQ ID NO: 4; a light chain variable region CDR2 of SEQ ID NO: 5; and a light chain variable region CDR3 of SEQ ID NO: 6. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 1; a heavy chain variable region CDR2 of SEQ ID NO: 2; a heavy chain variable

region CDR3 of SEQ ID NO: 3; a light chain variable region CDR1 of SEQ ID NO: 4; a light chain variable region CDR2 of SEQ ID NO: 5; and a light chain variable region CDR3 of SEQ ID NO: 120.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 62; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 209. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 95; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 209.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 66. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 78.

In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 61; a heavy chain variable region CDR2 of SEQ ID NO: 77; a heavy chain variable region CDR3 of SEQ ID NO: 63; a light chain variable region CDR1 of SEQ ID NO: 64; a light chain variable region CDR2 of SEQ ID NO: 65; and a light chain variable region CDR3 of SEQ ID NO: 89. In another specific embodiment, an antibody that specifically binds to C5 comprising a heavy chain variable region CDR1 of SEQ ID NO: 17; a heavy chain variable region CDR2 of SEQ ID NO: 107; a heavy chain variable region CDR3 of SEQ ID NO: 19; a light chain variable region CDR1 of SEQ ID NO: 20; a light chain variable region CDR2 of SEQ ID NO: 21; and a light chain variable region CDR3 of SEQ ID NO: 22.

In certain embodiments, an antibody that specifically binds to C5 is an antibody that is described in Table 1.

As used herein, a human antibody comprises heavy or light chain variable regions or full length heavy or light chains that are "the product of" or "derived from" a particular germline sequence if the variable regions or full length chains of the antibody are obtained from a system that uses human germline immunoglobulin genes. Such systems include immunizing a transgenic mouse carrying human immunoglobulin genes with the antigen of interest or screening a human immunoglobulin gene library displayed on phage with the antigen of interest. A human antibody that is "the product of" or "derived from" a human germline immunoglobulin sequence can be identified as such by comparing the amino acid sequence of the human antibody to the amino acid sequences of human germline immunoglobulins and selecting the human germline immunoglobulin sequence that is closest in sequence (*i.e.*, greatest % identity) to the sequence of the human antibody. A human antibody that is "the product of" or "derived from" a particular human germline immunoglobulin sequence may contain amino acid differences as compared to the germline sequence, due to, for example, naturally occurring somatic mutations or intentional introduction of site-directed mutations. However, in the VH or VL framework regions, a selected human antibody typically is at least 90% identical in amino acids sequence to an amino acid sequence encoded by a human germline immunoglobulin gene and contains amino acid residues that identify the human antibody as being human when compared to the germline immunoglobulin amino acid sequences of other species (*e.g.*, murine germline sequences). In certain cases, a human antibody may be at least 60%, 70%, 80%, 90%, or at least 95%, or even at least 96%, 97%, 98%, or 99% identical in amino acid sequence to the amino acid sequence encoded by the germline immunoglobulin gene. Typically, a recombinant human antibody will display no more than 10 amino acid differences from the amino acid sequence encoded by the human germline immunoglobulin gene in the VH or VL framework regions. In certain cases, the human antibody may display no more than 5, or even no more than 4, 3, 2, or 1 amino acid difference from the amino acid sequence encoded by the germline immunoglobulin gene.

Homologous antibodies

In yet another embodiment, the present invention provides an antibody or an antigen-binding fragment thereof comprising amino acid sequences that are homologous to the sequences described in Table 1, and said antibody binds to a C5 protein (*e.g.*, human and/or cynomolgus C5), and retains the desired functional properties of those antibodies described in Table 1.

For example, the invention provides an isolated monoclonal antibody (or a functional antigen binding fragment thereof) comprising a heavy chain variable region and a light chain variable region, wherein the heavy chain variable region comprises an amino acid sequence that is at least 80%, at least 90%, or at least 95% identical to an amino acid sequence

selected from the group consisting of SEQ ID NOs: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281; the light chain variable region comprises an amino acid sequence that is at least 80%, at least 90%, or at least 95% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, or 289; the antibody specifically binds to C5 (e.g., human and/or cynomolgus C5), and the antibody can inhibit red blood cell lysis in a hemolytic assay. In a specific example, such antibodies have an IC_{50} value in a hemolytic assay of 20-200 pM when using human C5-depleted serum that is reconstituted with 100 pM human C5.

In other embodiments, the VH and/or VL amino acid sequences may be 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth in Table 1. In other embodiments, the VH and/or VL amino acid sequences may be identical except an amino acid substitution in no more than 1,2,3,4 or 5 amino acid position. An antibody having VH and VL regions having high (*i. e.*, 80% or greater) identity to the VH and VL regions of those described in Table 1 can be obtained by mutagenesis (e.g., site-directed or PCR-mediated mutagenesis) of nucleic acid molecules encoding SEQ ID NOs: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281; and 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, or 289 respectively, followed by testing of the encoded altered antibody for retained function using the functional assays described herein.

In other embodiments, the full length heavy chain and/or full length light chain amino acid sequences may be 50% 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth in Table 1. An antibody having a full length heavy chain and full length light chain having high (*i.e.*, 80% or greater) identity to the full length heavy chains of any of SEQ ID NOs : 9, 25, 41, 53, 69, 81, 97, 109, 115, 123, 139, 153, 167, 181, 189, 203, 212, 220, 229, 243, 249, 254, 258, 274, 278, and 282 and full length light chains of any of SEQ ID NOs 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154, 168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, and 290 respectively, can be obtained by mutagenesis (e.g., site-directed or PCR-mediated mutagenesis) of nucleic acid molecules encoding such polypeptides respectively, followed by testing of the encoded altered antibody for retained function using the functional assays described herein.

In other embodiments, the full length heavy chain and/or full length light chain nucleotide sequences may be 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth above.

In other embodiments, the variable regions of heavy chain and/or light chain nucleotide sequences may be 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98% or 99% identical to the sequences set forth above

As used herein, the percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % identity equals number of identical positions/total number of positions x 100), taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

Additionally or alternatively, the protein sequences of the present invention can further be used as a "query sequence" to perform a search against public databases to, for example, identify related sequences. For example, such searches can be performed using the BLAST program (version 2.0) of Altschul, *et al.*, 1990 J.Mol. Biol. 215:403-10.

Antibodies with Conservative Modifications

In certain embodiments, an antibody of the invention has a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein one or more of these CDR sequences have specified amino acid sequences based on the antibodies described herein or conservative modifications thereof, and wherein the antibodies retain the desired functional properties of the C5-binding antibodies of the invention. Accordingly, the invention provides an isolated monoclonal antibody, or a functional antigen binding fragment thereof, consisting of a heavy chain variable region comprising CDR1, CDR2, and CDR3 sequences and a light chain variable region comprising CDR1, CDR2, and CDR3 sequences, wherein: the heavy chain variable region CDR1 amino acid sequences are selected from the group consisting of SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235, and conservative modifications thereof; the heavy chain variable region CDR2 amino acid sequences are selected from the group consisting of SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236, and conservative modifications thereof; the heavy chain variable region CDR3 amino acid sequences are selected from the group consisting of SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237, and conservative modifications thereof; the light chain variable regions CDR1 amino acid sequences are selected from the group consisting of SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238, and conservative modifications thereof; the light chain variable regions CDR2 amino acid sequences are selected from the group consisting of SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239, and conservative modifications thereof; the light chain

variable regions of CDR3 amino acid sequences are selected from the group consisting of SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240, and conservative modifications thereof; the antibody or the antigen-binding fragment thereof specifically binds to C5, and inhibits red blood cell lysis in a hemolytic assay as described herein.

In other embodiments, an antibody of the invention optimized for expression in a mammalian cell has a full length heavy chain sequence and a full length light chain sequence, wherein one or more of these sequences have specified amino acid sequences based on the antibodies described herein or conservative modifications thereof, and wherein the antibodies retain the desired functional properties of the C5-binding antibodies of the invention. Accordingly, the invention provides an isolated monoclonal antibody optimized for expression in a mammalian cell consisting of a full length heavy chain and a full length light chain wherein: the full length heavy chain has amino acid sequences selected from the group of SEQ ID NOs: : 9, 25, 41, 53, 69, 81, 97, 109, 115, 123, 139, 153, 167, 181, 189, 203, 212, 220, 229, 243, 249, 254, 258, 274, 278, and 282, and conservative modifications thereof; and the full length light chain has amino acid sequences selected from the group of SEQ ID NOs: 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154, 168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, and 290, and conservative modifications thereof; the antibody specifically binds to C5 (*e.g.*, human and/or cynomolgus C5); and the antibody inhibits red blood cell lysis in a hemolytic assay as described herein. In a specific embodiment, such antibodies have an IC_{50} value in a hemolytic assay of 20-200 pM when using human C5-depleted serum that is reconstituted with 100 pM human C5.

Antibodies That Bind to the Same Epitope

The present invention provides antibodies that bind to the same epitope as do the C5-binding antibodies described in Table 1. Additional antibodies can therefore be identified based on their ability to cross-compete (*e.g.*, to competitively inhibit the binding of, in a statistically significant manner) with other antibodies of the invention in C5 binding assays. The ability of a test antibody to inhibit the binding of antibodies of the present invention to a C5 protein (*e.g.*, human and/or cynomolgus C5) demonstrates that the test antibody can compete with that antibody for binding to C5; such an antibody may, according to non-limiting theory, bind to the same or a related (*e.g.*, a structurally similar or spatially proximal) epitope on the C5 protein as the antibody with which it competes. In a certain embodiment, the antibody that binds to the same epitope on C5 as the antibodies of the present invention is a human monoclonal antibody. Such human monoclonal antibodies can be prepared and isolated as described herein.

Engineered and Modified Antibodies

An antibody of the invention further can be prepared using an antibody having one or more of the VH and/or VL sequences shown herein as starting material to engineer a modified antibody, which modified antibody may have altered properties from the starting antibody. An antibody can be engineered by modifying one or more residues within one or both variable regions (*i. e.*, VH and/or VL), for example within one or more CDR regions and/or within one or more framework regions. Additionally or alternatively, an antibody can be engineered by modifying residues within the constant region(s), for example to alter the effector function(s) of the antibody.

One type of variable region engineering that can be performed is CDR grafting. Antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific naturally occurring antibodies by constructing expression vectors that include CDR sequences from the specific naturally occurring antibody grafted onto framework sequences from a different antibody with different properties (see, *e.g.*, Riechmann, L. *et al.*, 1998 *Nature* 332:323-327; Jones, P. *et al.*, 1986 *Nature* 321:522-525; Queen, C. *et al.*, 1989 *Proc. Natl. Acad., U.S.A.* 86:10029-10033; U.S. Patent No. 5,225,539 to Winter, and U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen *et al.*)

Accordingly, another embodiment of the invention pertains to an isolated monoclonal antibody, or an antigen binding fragment thereof, comprising a heavy chain variable region comprising CDR1 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235; CDR2 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236; CDR3 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237, respectively; and a light chain variable region having CDR1 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238; CDR2 sequences having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239; and CDR3 sequences consisting of an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240, respectively. Thus, such antibodies contain the VH and VL CDR sequences of monoclonal antibodies, yet may contain different framework sequences from these antibodies.

21489-11410

Such framework sequences can be obtained from public DNA databases or published references that include germline antibody gene sequences. For example, germline DNA sequences for human heavy and light chain variable region genes can be found in the "VBase" human germline sequence database (MRC Centre for Protein Engineering), as well as in Kabat, E. A., *et al.*, 1991 Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson, I. M., *et al.*, 1992 J. Mol. Biol. 227:776-798; and Cox, J. P. L. *et al.*, 1994 Eur. J. Immunol. 24:827-836.

An example of framework sequences for use in the antibodies of the invention are those that are structurally similar to the framework sequences used by selected antibodies of the invention, *e.g.*, consensus sequences and/or framework sequences used by monoclonal antibodies of the invention. The VH CDR1, 2 and 3 sequences, and the VL CDR1, 2 and 3 sequences, can be grafted onto framework regions that have the identical sequence as that found in the germline immunoglobulin gene from which the framework sequence derive, or the CDR sequences can be grafted onto framework regions that contain one or more mutations as compared to the germline sequences. For example, it has been found that in certain instances it is beneficial to mutate residues within the framework regions to maintain or enhance the antigen binding ability of the antibody (see *e.g.*, U.S. Patent Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen *et al.*).

Another type of variable region modification is to mutate amino acid residues within the VH and/or VL CDR1, CDR2 and/or CDR3 regions to thereby improve one or more binding properties (*e.g.*, affinity) of the antibody of interest, known as "affinity maturation." Site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the mutation(s) and the effect on antibody binding, or other functional property of interest, can be evaluated in *in vitro* or *in vivo* assays as described herein and provided in the Examples. Conservative modifications (as discussed above) can be introduced. The mutations may be amino acid substitutions, additions or deletions. Moreover, typically no more than one, two, three, four or five residues within a CDR region are altered.

Accordingly, in another embodiment, the invention provides isolated C5-binding monoclonal antibodies, or an antigen binding fragment thereof, consisting of a heavy chain variable region having: a VH CDR1 region consisting of an amino acid sequence selected from the group having SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235 or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235; a VH CDR2 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236, or

an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236; a VH CDR3 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237; a VL CDR1 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238; a VL CDR2 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239; and a VL CDR3 region having an amino acid sequence selected from the group consisting of SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240, or an amino acid sequence having one, two, three, four or five amino acid substitutions, deletions or additions as compared to SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240.

Grafting Antigen-binding Domains Into Alternative Frameworks or Scaffolds

A wide variety of antibody/ immunoglobulin frameworks or scaffolds can be employed so long as the resulting polypeptide includes at least one binding region which specifically binds to C5. Such frameworks or scaffolds include the 5 main idiotypes of human immunoglobulins, or fragments thereof, and include immunoglobulins of other animal species, preferably having humanized aspects. Single heavy-chain antibodies such as those identified in camelids are of particular interest in this regard. Novel frameworks, scaffolds and fragments continue to be discovered and developed by those skilled in the art.

In one aspect, the invention pertains to generating non-immunoglobulin based antibodies using non- immunoglobulin scaffolds onto which CDRs of the invention can be grafted. Known or future non-immunoglobulin frameworks and scaffolds may be employed, as long as they comprise a binding region specific for the target C5 protein (e.g., human and/or cynomolgus C5). Known non-immunoglobulin frameworks or scaffolds include, but are not limited to, fibronectin (Compound Therapeutics, Inc., Waltham, MA), ankyrin (Molecular Partners AG, Zurich, Switzerland), domain antibodies (Domantis, Ltd., Cambridge, MA, and Ablynx nv, Zwijnaarde, Belgium), lipocalin (Pieris Proteolab AG, Freising, Germany), small modular immuno-pharmaceuticals (Trubion Pharmaceuticals Inc.,

Seattle, WA), maxybodies (Avidia, Inc., Mountain View, CA), Protein A (Affibody AG, Sweden), and affilin (gamma-crystallin or ubiquitin) (Scil Proteins GmbH, Halle, Germany).

The fibronectin scaffolds are based on fibronectin type III domain (*e.g.*, the tenth module of the fibronectin type III (10 Fn3 domain)). The fibronectin type III domain has 7 or 8 beta strands which are distributed between two beta sheets, which themselves pack against each other to form the core of the protein, and further containing loops (analogous to CDRs) which connect the beta strands to each other and are solvent exposed. There are at least three such loops at each edge of the beta sheet sandwich, where the edge is the boundary of the protein perpendicular to the direction of the beta strands (see US 6,818,418). These fibronectin-based scaffolds are not an immunoglobulin, although the overall fold is closely related to that of the smallest functional antibody fragment, the variable region of the heavy chain, which comprises the entire antigen recognition unit in camel and llama IgG. Because of this structure, the non-immunoglobulin antibody mimics antigen binding properties that are similar in nature and affinity to those of antibodies. These scaffolds can be used in a loop randomization and shuffling strategy *in vitro* that is similar to the process of affinity maturation of antibodies *in vivo*. These fibronectin-based molecules can be used as scaffolds where the loop regions of the molecule can be replaced with CDRs of the invention using standard cloning techniques.

The ankyrin technology is based on using proteins with ankyrin derived repeat modules as scaffolds for bearing variable regions which can be used for binding to different targets. The ankyrin repeat module is a 33 amino acid polypeptide consisting of two anti-parallel α -helices and a β -turn. Binding of the variable regions is mostly optimized by using ribosome display.

Avimers are derived from natural A-domain containing protein such as LRP-1. These domains are used by nature for protein-protein interactions and in human over 250 proteins are structurally based on A-domains. Avimers consist of a number of different "A-domain" monomers (2-10) linked via amino acid linkers. Avimers can be created that can bind to the target antigen using the methodology described in, for example, U.S. Patent Application Publication Nos. 20040175756; 20050053973; 20050048512; and 20060008844.

Affibody affinity ligands are small, simple proteins composed of a three-helix bundle based on the scaffold of one of the IgG-binding domains of Protein A. Protein A is a surface protein from the bacterium *Staphylococcus aureus*. This scaffold domain consists of 58 amino acids, 13 of which are randomized to generate affibody libraries with a large number of ligand variants (See *e.g.*, US 5,831,012). Affibody molecules mimic antibodies, they have a molecular weight of 6 kDa, compared to the molecular weight of antibodies, which is 150

kDa. In spite of its small size, the binding site of affibody molecules is similar to that of an antibody.

Anticalins are products developed by the company Pieris ProteoLab AG. They are derived from lipocalins, a widespread group of small and robust proteins that are usually involved in the physiological transport or storage of chemically sensitive or insoluble compounds. Several natural lipocalins occur in human tissues or body liquids. The protein architecture is reminiscent of immunoglobulins, with hypervariable loops on top of a rigid framework. However, in contrast with antibodies or their recombinant fragments, lipocalins are composed of a single polypeptide chain with 160 to 180 amino acid residues, being just marginally bigger than a single immunoglobulin domain. The set of four loops, which makes up the binding pocket, shows pronounced structural plasticity and tolerates a variety of side chains. The binding site can thus be reshaped in a proprietary process in order to recognize prescribed target molecules of different shape with high affinity and specificity. One protein of lipocalin family, the bilin-binding protein (BBP) of *Pieris Brassicae* has been used to develop anticalins by mutagenizing the set of four loops. One example of a patent application describing anticalins is in PCT Publication No. WO 199916873.

Affilin molecules are small non-immunoglobulin proteins which are designed for specific affinities towards proteins and small molecules. New affilin molecules can be very quickly selected from two libraries, each of which is based on a different human derived scaffold protein. Affilin molecules do not show any structural homology to immunoglobulin proteins. Currently, two affilin scaffolds are employed, one of which is gamma crystalline, a human structural eye lens protein and the other is "ubiquitin" superfamily proteins. Both human scaffolds are very small, show high temperature stability and are almost resistant to pH changes and denaturing agents. This high stability is mainly due to the expanded beta sheet structure of the proteins. Examples of gamma crystalline derived proteins are described in WO200104144 and examples of "ubiquitin-like" proteins are described in WO2004106368.

Protein epitope mimetics (PEM) are medium-sized, cyclic, peptide-like molecules (MW 1-2kDa) mimicking beta-hairpin secondary structures of proteins, the major secondary structure involved in protein-protein interactions.

In some embodiments, the Fabs are converted to silent IgG1 format by changing the Fc region. For example, antibodies 6525-7910 in Table 1 can be converted to silent IgG1 formate by substituting the "X" in the amino acid sequences for the heavy chain with:

CDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVD
GVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG

QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS CSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 293)

and substituting the “X” in the amino acid sequence for the light chain with: CS if the light chain is lambda, or C if the light chain is kappa. As used herein, a “silent IgG1” is an IgG1 Fc sequence in which the amino acid sequence has been altered to reduce Fc-mediated effector functions (for example ADCC and/or CDC). Such an antibody will typically have reduced binding to Fc receptors and/or C1q.

In some other embodiments, the Fabs are converted to IgG2 format. For example, antibodies 6525-7910 in Table 1 can be converted to IgG2 format by substituting the constant sequence

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSX (SEQ ID NO: 294)

with the constant sequence for the heavy chain of IgG2:

ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSNFGTQTYTCNV DHKPSNTKVDKTV ERKCCVECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFS CSVMHEALHNHYTQKSLSLSPGK (SEQ ID NO: 295)

and substituting the “X” in the amino acid sequence for the light chain with CS if the light chain is lambda, or C if the light chain is kappa.

Human or humanized antibodies

The present invention provides fully human antibodies that specifically bind to a C5 protein (e.g., human and/or cynomolgus C5). Compared to the chimeric or humanized antibodies, the human C5-binding antibodies of the invention have further reduced antigenicity when administered to human subjects.

The human C5-binding antibodies can be generated using methods that are known in the art. For example, the humaneering technology used to converting non-human antibodies into engineered human antibodies. U.S. Patent Publication No. 20050008625 describes an *in vivo* method for replacing a nonhuman antibody variable region with a human variable region in an antibody while maintaining the same or providing better binding characteristics relative to that of the nonhuman antibody. The method relies on epitope guided replacement of variable regions of a non-human reference antibody with a fully human antibody. The resulting human antibody is generally unrelated structurally to the reference nonhuman antibody, but binds to the same epitope on the same antigen as the reference antibody.

Briefly, the serial epitope-guided complementarity replacement approach is enabled by setting up a competition in cells between a "competitor" and a library of diverse hybrids of the reference antibody ("test antibodies") for binding to limiting amounts of antigen in the presence of a reporter system which responds to the binding of test antibody to antigen. The competitor can be the reference antibody or derivative thereof such as a single-chain Fv fragment. The competitor can also be a natural or artificial ligand of the antigen which binds to the same epitope as the reference antibody. The only requirements of the competitor are that it binds to the same epitope as the reference antibody, and that it competes with the reference antibody for antigen binding. The test antibodies have one antigen-binding V-region in common from the nonhuman reference antibody, and the other V-region selected at random from a diverse source such as a repertoire library of human antibodies. The common V-region from the reference antibody serves as a guide, positioning the test antibodies on the same epitope on the antigen, and in the same orientation, so that selection is biased toward the highest antigen-binding fidelity to the reference antibody.

Many types of reporter system can be used to detect desired interactions between test antibodies and antigen. For example, complementing reporter fragments may be linked to antigen and test antibody, respectively, so that reporter activation by fragment complementation only occurs when the test antibody binds to the antigen. When the test antibody- and antigen-reporter fragment fusions are co-expressed with a competitor, reporter activation becomes dependent on the ability of the test antibody to compete with the competitor, which is proportional to the affinity of the test antibody for the antigen. Other reporter systems that can be used include the reactivator of an auto-inhibited reporter reactivation system (RAIR) as disclosed in U.S. Patent Application Ser. No. 10/208,730 (Publication No. 20030198971), or competitive activation system disclosed in U.S. Patent Application Ser. No. 10/076,845 (Publication No. 20030157579).

With the serial epitope-guided complementarity replacement system, selection is made to identify cells expresses a single test antibody along with the competitor, antigen, and reporter components. In these cells, each test antibody competes one-on-one with the competitor for binding to a limiting amount of antigen. Activity of the reporter is proportional to the amount of antigen bound to the test antibody, which in turn is proportional to the affinity of the test antibody for the antigen and the stability of the test antibody. Test antibodies are initially selected on the basis of their activity relative to that of the reference antibody when expressed as the test antibody. The result of the first round of selection is a set of "hybrid" antibodies, each of which is comprised of the same non-human V-region from the reference antibody and a human V-region from the library, and each of which binds to the same epitope on the antigen as the reference antibody. One or more of the hybrid antibodies

selected in the first round will have an affinity for the antigen comparable to or higher than that of the reference antibody.

In the second V-region replacement step, the human V-regions selected in the first step are used as guide for the selection of human replacements for the remaining non-human reference antibody V-region with a diverse library of cognate human V-regions. The hybrid antibodies selected in the first round may also be used as competitors for the second round of selection. The result of the second round of selection is a set of fully human antibodies which differ structurally from the reference antibody, but which compete with the reference antibody for binding to the same antigen. Some of the selected human antibodies bind to the same epitope on the same antigen as the reference antibody. Among these selected human antibodies, one or more binds to the same epitope with an affinity which is comparable to or higher than that of the reference antibody.

Using one of the mouse or chimeric C5-binding antibodies described above as the reference antibody, this method can be readily employed to generate human antibodies that bind to human C5 with the same binding specificity and the same or better binding affinity. In addition, such human C5-binding antibodies can also be commercially obtained from companies which customarily produce human antibodies, *e.g.*, KaloBios, Inc. (Mountain View, CA).

Camelid antibodies

Antibody proteins obtained from members of the camel and dromedary (*Camelus bactrianus* and *Camelus dromedarius*) family including new world members such as llama species (*Lama pacos*, *Lama glama* and *Lama vicugna*) have been characterized with respect to size, structural complexity and antigenicity for human subjects. Certain IgG antibodies from this family of mammals as found in nature lack light chains, and are thus structurally distinct from the typical four chain quaternary structure having two heavy and two light chains, for antibodies from other animals. See PCT/EP93/02214 (WO 94/04678 published 3 March 1994).

A region of the camelid antibody which is the small single variable domain identified as VHH can be obtained by genetic engineering to yield a small protein having high affinity for a target, resulting in a low molecular weight antibody-derived protein known as a "camelid nanobody". See U.S. patent number 5,759,808 issued June 2, 1998; see also Stijlemans, B. *et al.*, 2004 J Biol Chem 279: 1256-1261; Dumoulin, M. *et al.*, 2003 Nature 424: 783-788; Pleschberger, M. *et al.* 2003 Bioconjugate Chem 14: 440-448; Cortez-Retamozo, V. *et al.* 2002 Int J Cancer 89: 456-62; and Lauwereys, M. *et al.* 1998 EMBO J 17: 3512-3520. Engineered libraries of camelid antibodies and antibody fragments are commercially available, for example, from Ablynx, Ghent, Belgium. As with other antibodies of non-human

origin, an amino acid sequence of a camelid antibody can be altered recombinantly to obtain a sequence that more closely resembles a human sequence, *i.e.*, the nanobody can be "humanized". Thus the natural low antigenicity of camelid antibodies to humans can be further reduced.

The camelid nanobody has a molecular weight approximately one-tenth that of a human IgG molecule, and the protein has a physical diameter of only a few nanometers. One consequence of the small size is the ability of camelid nanobodies to bind to antigenic sites that are functionally invisible to larger antibody proteins, *i.e.*, camelid nanobodies are useful as reagents detect antigens that are otherwise cryptic using classical immunological techniques, and as possible therapeutic agents. Thus yet another consequence of small size is that a camelid nanobody can inhibit as a result of binding to a specific site in a groove or narrow cleft of a target protein, and hence can serve in a capacity that more closely resembles the function of a classical low molecular weight drug than that of a classical antibody.

The low molecular weight and compact size further result in camelid nanobodies being extremely thermostable, stable to extreme pH and to proteolytic digestion, and poorly antigenic. Another consequence is that camelid nanobodies readily move from the circulatory system into tissues, and even cross the blood-brain barrier and can treat disorders that affect nervous tissue. Nanobodies can further facilitated drug transport across the blood brain barrier. See U.S. patent application 20040161738 published August 19, 2004. These features combined with the low antigenicity to humans indicate great therapeutic potential. Further, these molecules can be fully expressed in prokaryotic cells such as *E. coli* and are expressed as fusion proteins with bacteriophage and are functional.

Accordingly, a feature of the present invention is a camelid antibody or nanobody having high affinity for C5. In certain embodiments herein, the camelid antibody or nanobody is naturally produced in the camelid animal, *i.e.*, is produced by the camelid following immunization with C5 or a peptide fragment thereof, using techniques described herein for other antibodies. Alternatively, the C5-binding camelid nanobody is engineered, *i.e.*, produced by selection for example from a library of phage displaying appropriately mutagenized camelid nanobody proteins using panning procedures with C5 as a target as described in the examples herein. Engineered nanobodies can further be customized by genetic engineering to have a half life in a recipient subject of from 45 minutes to two weeks. In a specific embodiment, the camelid antibody or nanobody is obtained by grafting the CDRs sequences of the heavy or light chain of the human antibodies of the invention into nanobody or single domain antibody framework sequences, as described for example in PCT/EP93/02214.

Bispecific Molecules and Multivalent Antibodies

In another aspect, the present invention features bispecific or multispecific molecules comprising an C5-binding antibody, or a fragment thereof, of the invention. An antibody of the invention, or antigen-binding regions thereof, can be derivatized or linked to another functional molecule, e.g., another peptide or protein (e.g., another antibody or ligand for a receptor) to generate a bispecific molecule that binds to at least two different binding sites or target molecules. The antibody of the invention may in fact be derivatized or linked to more than one other functional molecule to generate multi-specific molecules that bind to more than two different binding sites and/or target molecules; such multi-specific molecules are also intended to be encompassed by the term "bispecific molecule" as used herein. To create a bispecific molecule of the invention, an antibody of the invention can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other binding molecules, such as another antibody, antibody fragment, peptide or binding mimetic, such that a bispecific molecule results.

Accordingly, the present invention includes bispecific molecules comprising at least one first binding specificity for C5 and a second binding specificity for a second target epitope. For example, the second target epitope is another epitope of C5 different from the first target epitope.

Additionally, for the invention in which the bispecific molecule is multi-specific, the molecule can further include a third binding specificity, in addition to the first and second target epitope.

In one embodiment, the bispecific molecules of the invention comprise as a binding specificity at least one antibody, or an antibody fragment thereof, including, e.g., an Fab, Fab', F(ab')₂, Fv, or a single chain Fv. The antibody may also be a light chain or heavy chain dimer, or any minimal fragment thereof such as a Fv or a single chain construct as described in Ladner *et al.* U.S. Patent No. 4,946,778.

Diabodies are bivalent, bispecific molecules in which VH and VL domains are expressed on a single polypeptide chain, connected by a linker that is too short to allow for pairing between the two domains on the same chain. The VH and VL domains pair with complementary domains of another chain, thereby creating two antigen binding sites (see e.g., Holliger *et al.*, 1993 Proc. Natl. Acad. Sci. USA 90:6444-6448; Poljak *et al.*, 1994 Structure 2:1121-1123). Diabodies can be produced by expressing two polypeptide chains with either the structure VHA-VLB and VHB-VLA (VH-VL configuration), or VLA-VHB and VLB-VHA (VL-VH configuration) within the same cell. Most of them can be expressed in soluble form in bacteria. Single chain diabodies (scDb) are produced by connecting the two diabody-forming polypeptide chains with linker of approximately 15 amino acid residues (see

Holliger and Winter, 1997 *Cancer Immunol. Immunother.*, 45(3-4):128-30; Wu *et al.*, 1996 *Immunotechnology*, 2(1):21-36). scDb can be expressed in bacteria in soluble, active monomeric form (see Holliger and Winter, 1997 *Cancer Immunol. Immunother.*, 45(34): 128-30; Wu *et al.*, 1996 *Immunotechnology*, 2(1):21-36; Pluckthun and Pack, 1997 *Immunotechnology*, 3(2): 83-105; Ridgway *et al.*, 1996 *Protein Eng.*, 9(7):617-21). A diabody can be fused to Fc to generate a "di-diabody" (see Lu *et al.*, 2004 *J. Biol. Chem.*, 279(4):2856-65).

Other antibodies which can be employed in the bispecific molecules of the invention are murine, chimeric and humanized monoclonal antibodies.

The bispecific molecules of the present invention can be prepared by conjugating the constituent binding specificities, using methods known in the art. For example, each binding specificity of the bispecific molecule can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-S-acetyl-thioacetate (SATA), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfosuccinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC) (see *e.g.*, Karpovsky *et al.*, 1984 *J. Exp. Med.* 160:1686; Liu, MA *et al.*, 1985 *Proc. Natl. Acad. Sci. USA* 82:8648). Other methods include those described in Paulus, 1985 *Behring Ins. Mitt.* No. 78,118-132; Brennan *et al.*, 1985 *Science* 229:81-83), and Glennie *et al.*, 1987 *J. Immunol.* 139: 2367-2375). Conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, IL).

When the binding specificities are antibodies, they can be conjugated by sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In a particularly embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, for example one, prior to conjugation.

Alternatively, both binding specificities can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific molecule is a mAb x mAb, mAb x Fab, Fab x F(ab')₂ or ligand x Fab fusion protein. A bispecific molecule of the invention can be a single chain molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific molecules may comprise at least two single chain molecules. Methods for preparing bispecific molecules are described for example in U.S. Patent Number 5,260,203; U.S. Patent Number 5,455,030; U.S. Patent Number 4,881,175; U.S. Patent Number 5,132,405; U.S. Patent Number 5,091,513; U.S. Patent Number

5,476,786; U.S. Patent Number 5,013,653; U.S. Patent Number 5,258,498; and U.S. Patent Number 5,482,858.

Binding of the bispecific molecules to their specific targets can be confirmed by, for example, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (REA), FACS analysis, bioassay (e.g., growth inhibition), or Western Blot assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody) specific for the complex of interest.

In another aspect, the present invention provides multivalent compounds comprising at least two identical or different antigen-binding portions of the antibodies of the invention binding to C5. The antigen-binding portions can be linked together via protein fusion or covalent or non covalent linkage. Alternatively, methods of linkage has been described for the bispecific molecules. Tetravalent compounds can be obtained for example by cross-linking antibodies of the invention with an antibody that binds to the constant regions of the antibodies of the invention, for example the Fc or hinge region.

Trimerizing domain are described for example in Borean patent EP 1 012 280B1. Pentamerizing modules are described for example in PCT/EP97/05897.

Antibodies with Extended Half Life

The present invention provides for antibodies that specifically bind to C5 protein which have an extended half-life *in vivo*.

Many factors may affect a protein's half life *in vivo*. For examples, kidney filtration, metabolism in the liver, degradation by proteolytic enzymes (proteases), and immunogenic responses (e.g., protein neutralization by antibodies and uptake by macrophages and dendritic cells). A variety of strategies can be used to extend the half life of the antibodies of the present invention. For example, by chemical linkage to polyethyleneglycol (PEG), reCODE PEG, antibody scaffold, polysialic acid (PSA), hydroxyethyl starch (HES), albumin-binding ligands, and carbohydrate shields; by genetic fusion to proteins binding to serum proteins, such as albumin, IgG, FcRn, and transferrin; by coupling (genetically or chemically) to other binding moieties that bind to serum proteins, such as nanobodies, Fabs, DARPins, avimers, affibodies, and anticalins; by genetic fusion to rPEG, albumin, domain of albumin, albumin-binding proteins, and Fc; or by incorporation into nanocarriers, slow release formulations, or medical devices.

To prolong the serum circulation of antibodies *in vivo*, inert polymer molecules such as high molecular weight PEG can be attached to the antibodies or a fragment thereof with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C-terminus of the antibodies or via epsilon-amino groups present on lysine residues. To

pegylate an antibody, the antibody, or fragment thereof, typically is reacted with polyethylene glycol (PEG), such as a reactive ester or aldehyde derivative of PEG, under conditions in which one or more PEG groups become attached to the antibody or antibody fragment. The pegylation can be carried out by an acylation reaction or an alkylation reaction with a reactive PEG molecule (or an analogous reactive water-soluble polymer). As used herein, the term "polyethylene glycol" is intended to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol or polyethylene glycol-maleimide. In certain embodiments, the antibody to be pegylated is an aglycosylated antibody. Linear or branched polymer derivatization that results in minimal loss of biological activity will be used. The degree of conjugation can be closely monitored by SDS-PAGE and mass spectrometry to ensure proper conjugation of PEG molecules to the antibodies. Unreacted PEG can be separated from antibody-PEG conjugates by size-exclusion or by ion-exchange chromatography. PEG-derivatized antibodies can be tested for binding activity as well as for *in vivo* efficacy using methods well-known to those of skill in the art, for example, by immunoassays described herein. Methods for pegylating proteins are known in the art and can be applied to the antibodies of the invention. See for example, EP 0 154 316 by Nishimura *et al.* and EP 0 401 384 by Ishikawa *et al.*

Other modified pegylation technologies include reconstituting chemically orthogonal directed engineering technology (ReCODE PEG), which incorporates chemically specified side chains into biosynthetic proteins via a reconstituted system that includes tRNA synthetase and tRNA. This technology enables incorporation of more than 30 new amino acids into biosynthetic proteins in *E. coli*, yeast, and mammalian cells. The tRNA incorporates a nonnative amino acid any place an amber codon is positioned, converting the amber from a stop codon to one that signals incorporation of the chemically specified amino acid.

Recombinant pegylation technology (rPEG) can also be used for serum half-life extension. This technology involves genetically fusing a 300-600 amino acid unstructured protein tail to an existing pharmaceutical protein. Because the apparent molecular weight of such an unstructured protein chain is about 15-fold larger than its actual molecular weight, the serum half-life of the protein is greatly increased. In contrast to traditional PEGylation, which requires chemical conjugation and repurification, the manufacturing process is greatly simplified and the product is homogeneous.

Polysialylation is another technology, which uses the natural polymer polysialic acid (PSA) to prolong the active life and improve the stability of therapeutic peptides and proteins. PSA is a polymer of sialic acid (a sugar). When used for protein and therapeutic peptide drug delivery, polysialic acid provides a protective microenvironment on conjugation. This increases the active life of the therapeutic protein in the circulation and prevents it from being

recognized by the immune system. The PSA polymer is naturally found in the human body. It was adopted by certain bacteria which evolved over millions of years to coat their walls with it. These naturally polysialylated bacteria were then able, by virtue of molecular mimicry, to foil the body's defence system. PSA, nature's ultimate stealth technology, can be easily produced from such bacteria in large quantities and with predetermined physical characteristics. Bacterial PSA is completely non-immunogenic, even when coupled to proteins, as it is chemically identical to PSA in the human body.

Another technology include the use of hydroxyethyl starch ("HES") derivatives linked to antibodies. HES is a modified natural polymer derived from waxy maize starch and can be metabolized by the body's enzymes. HES solutions are usually administered to substitute deficient blood volume and to improve the rheological properties of the blood. Hesylation of an antibody enables the prolongation of the circulation half-life by increasing the stability of the molecule, as well as by reducing renal clearance, resulting in an increased biological activity. By varying different parameters, such as the molecular weight of HES, a wide range of HES antibody conjugates can be customized.

Antibodies having an increased half-life *in vivo* can also be generated introducing one or more amino acid modifications (*i.e.*, substitutions, insertions or deletions) into an IgG constant domain, or FcRn binding fragment thereof (preferably a Fc or hinge Fc domain fragment). See, *e.g.*, International Publication No. WO 98/23289; International Publication No. WO 97/34631; and U.S. Patent No. 6,277,375.

Further, antibodies can be conjugated to albumin in order to make the antibody or antibody fragment more stable *in vivo* or have a longer half life *in vivo*. The techniques are well-known in the art, see, *e.g.*, International Publication Nos. WO 93/15199, WO 93/15200, and WO 01/77137; and European Patent No. EP 413,622.

The strategies for increasing half life is especially useful in nanobodies, fibronectin-based binders, and other antibodies or proteins for which increased *in vivo* half life is desired.

Antibody Conjugates

The present invention provides antibodies or fragments thereof that specifically bind to a C5 protein recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or polypeptide (or fragment thereof, preferably to a polypeptide of at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids) to generate fusion proteins. In particular, the invention provides fusion proteins comprising an antigen-binding fragment of an antibody described herein (*e.g.*, a Fab fragment, Fd fragment, Fv fragment, F(ab)₂ fragment, a VH domain, a VH CDR, a VL domain or a VL CDR) and a heterologous protein, polypeptide, or peptide. Methods for fusing or conjugating proteins, polypeptides, or

21489-11410

peptides to an antibody or an antibody fragment are known in the art. See, e.g., U.S. Patent Nos. 5,336,603, 5,622,929, 5,359,046, 5,349,053, 5,447,851, and 5,112,946; European Patent Nos. EP 307,434 and EP 367,166; International Publication Nos. WO 96/04388 and WO 91/06570; Ashkenazi *et al.*, 1991, Proc. Natl. Acad. Sci. USA 88: 10535-10539; Zheng *et al.*, 1995, J. Immunol. 154:5590-5600; and Vil *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:11337- 11341.

Additional fusion proteins may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to alter the activities of antibodies of the invention or fragments thereof (e.g., antibodies or fragments thereof with higher affinities and lower dissociation rates). See, generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458; Patten *et al.*, 1997, Curr. Opinion Biotechnol. 8:724-33; Harayama, 1998, Trends Biotechnol. 16(2):76-82; Hansson, *et al.*, 1999, J. Mol. Biol. 287:265-76; and Lorenzo and Blasco, 1998, Biotechniques 24(2):308- 313. Antibodies or fragments thereof, or the encoded antibodies or fragments thereof, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. A polynucleotide encoding an antibody or fragment thereof that specifically binds to a C5 protein may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Moreover, the antibodies or fragments thereof can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, 1989, Proc. Natl. Acad. Sci. USA 86:821-824, for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin ("HA") tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, 1984, Cell 37:767), and the "flag" tag.

In other embodiments, antibodies of the present invention or fragments thereof conjugated to a diagnostic or detectable agent. Such antibodies can be useful for monitoring or prognosing the onset, development, progression and/or severity of a disease or disorder as part of a clinical testing procedure, such as determining the efficacy of a particular therapy. Such diagnosis and detection can accomplished by coupling the antibody to detectable substances including, but not limited to, various enzymes, such as, but not limited to, horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as, but not limited to, streptavidin/biotin and

21489-11410

avidin/biotin; fluorescent materials, such as, but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent materials, such as, but not limited to, luminol; bioluminescent materials, such as but not limited to, luciferase, luciferin, and aequorin; radioactive materials, such as, but not limited to, iodine (^{131}I , ^{125}I , ^{123}I , and ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{115}In , ^{113}In , ^{112}In , and ^{111}In), technetium (^{99}Tc), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn , ^{75}Se , ^{113}Sn , and ^{117}Tm ; and positron emitting metals using various positron emission tomographies, and noradioactive paramagnetic metal ions.

The present invention further encompasses uses of antibodies or fragments thereof conjugated to a therapeutic moiety. An antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells.

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety or drug moiety that modifies a given biological response. Therapeutic moieties or drug moieties are not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein, peptide, or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, cholera toxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, an anti-angiogenic agent; or, a biological response modifier such as, for example, a lymphokine.

Moreover, an antibody can be conjugated to therapeutic moieties such as a radioactive metal ion, such as alpha-emitters such as ^{213}Bi or macrocyclic chelators useful for conjugating radiometal ions, including but not limited to, ^{131}In , ^{131}Lu , ^{131}Y , ^{131}Ho , ^{131}Sm , to polypeptides. In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane- $\text{N},\text{N}',\text{N}'',\text{N}'''$ -tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo *et al.*, 1998, Clin Cancer Res. 4(10):2483-90; Peterson *et al.*, 1999, Bioconjug. Chem. 10(4):553-7; and Zimmerman *et al.*, 1999, Nucl. Med. Biol. 26(8):943-50.

Techniques for conjugating therapeutic moieties to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer

Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies 84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16 (Academic Press 1985), and Thorpe *et al.*, 1982, *Immunol. Rev.* 62:119-58.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

5.2. Methods of Producing Antibodies of the Invention

5.2.1. Nucleic Acids Encoding the Antibodies

The invention provides substantially purified nucleic acid molecules which encode polypeptides comprising segments or domains of the C5-binding antibody chains described above. Some of the nucleic acids of the invention comprise the nucleotide sequence encoding the heavy chain variable region shown in SEQ ID NO: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281, and/or the nucleotide sequence encoding the light chain variable region shown in SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, or 289. In a specific embodiment, the nucleic acid molecules are those identified in Table 1. Some other nucleic acid molecules of the invention comprise nucleotide sequences that are substantially identical (*e.g.*, at least 65, 80%, 95%, or 99%) to the nucleotide sequences of those identified in Table 1. When expressed from appropriate expression vectors, polypeptides encoded by these polynucleotides are capable of exhibiting C5 antigen binding capacity.

Also provided in the invention are polynucleotides which encode at least one CDR region and usually all three CDR regions from the heavy or light chain of the C5-binding antibody set forth above. Some other polynucleotides encode all or substantially all of the variable region sequence of the heavy chain and/or the light chain of the C5-binding antibody set forth above. Because of the degeneracy of the code, a variety of nucleic acid sequences will encode each of the immunoglobulin amino acid sequences.

The nucleic acid molecules of the invention can encode both a variable region and a constant region of the antibody. Some of nucleic acid sequences of the invention comprise

nucleotides encoding a mature heavy chain variable region sequence that is substantially identical (e.g., at least 80%, 90%, or 99%) to the mature heavy chain variable region sequence set forth in SEQ ID NO: 7, 23, 39, 51, 67, 79, 96, 108, 114, 121, 137, 151, 165, 179, 187, 201, 210, 218, 227, 241, 253, 257, 273, 277, or 281. Some other nucleic acid sequences comprising nucleotide encoding a mature light chain variable region sequence that is substantially identical (e.g., at least 80%, 90%, or 99%) to the mature light chain variable region sequence set forth in SEQ ID NO: 8, 24, 40, 52, 68, 80, 90, 102, 122, 138, 152, 166, 180, 188, 202, 211, 219, 228, 242, 261, 265, 269, 285, or 289.

The polynucleotide sequences can be produced by de novo solid-phase DNA synthesis or by PCR mutagenesis of an existing sequence (e.g., sequences as described in the Examples below) encoding an C5-binding antibody or its binding fragment. Direct chemical synthesis of nucleic acids can be accomplished by methods known in the art, such as the phosphotriester method of Narang *et al.*, 1979, Meth. Enzymol. 68:90; the phosphodiester method of Brown *et al.*, Meth. Enzymol. 68:109, 1979; the diethylphosphoramidite method of Beaucage *et al.*, Tetra. Lett., 22:1859, 1981; and the solid support method of U.S. Patent No. 4,458,066. Introducing mutations to a polynucleotide sequence by PCR can be performed as described in, e.g., PCR Technology: Principles and Applications for DNA Amplification, H.A. Erlich (Ed.), Freeman Press, NY, NY, 1992; PCR Protocols: A Guide to Methods and Applications, Innis *et al.* (Ed.), Academic Press, San Diego, CA, 1990; Mattila *et al.*, Nucleic Acids Res. 19:967, 1991; and Eckert *et al.*, PCR Methods and Applications 1:17, 1991.

Also provided in the invention are expression vectors and host cells for producing the C5-binding antibodies described above. Various expression vectors can be employed to express the polynucleotides encoding the C5-binding antibody chains or binding fragments. Both viral-based and nonviral expression vectors can be used to produce the antibodies in a mammalian host cell. Nonviral vectors and systems include plasmids, episomal vectors, typically with an expression cassette for expressing a protein or RNA, and human artificial chromosomes (see, e.g., Harrington *et al.*, Nat Genet 15:345, 1997). For example, nonviral vectors useful for expression of the C5-binding polynucleotides and polypeptides in mammalian (e.g., human) cells include pThioHis A, B & C, pcDNA3.1/His, pEBVHis A, B & C, (Invitrogen, San Diego, CA), MPSV vectors, and numerous other vectors known in the art for expressing other proteins. Useful viral vectors include vectors based on retroviruses, adenoviruses, adenoassociated viruses, herpes viruses, vectors based on SV40, papilloma virus, HBP Epstein Barr virus, vaccinia virus vectors and Semliki Forest virus (SFV). See, Brent *et al.*, supra; Smith, Annu. Rev. Microbiol. 49:807, 1995; and Rosenfeld *et al.*, Cell 68:143, 1992.

The choice of expression vector depends on the intended host cells in which the vector is to be expressed. Typically, the expression vectors contain a promoter and other regulatory sequences (e.g., enhancers) that are operably linked to the polynucleotides encoding an C5-binding antibody chain or fragment. In some embodiments, an inducible promoter is employed to prevent expression of inserted sequences except under inducing conditions. Inducible promoters include, e.g., arabinose, lacZ, metallothionein promoter or a heat shock promoter. Cultures of transformed organisms can be expanded under noninducing conditions without biasing the population for coding sequences whose expression products are better tolerated by the host cells. In addition to promoters, other regulatory elements may also be required or desired for efficient expression of an C5-binding antibody chain or fragment. These elements typically include an ATG initiation codon and adjacent ribosome binding site or other sequences. In addition, the efficiency of expression may be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, e.g., Scharf *et al.*, Results Probl. Cell Differ. 20:125, 1994; and Bittner *et al.*, Meth. Enzymol., 153:516, 1987). For example, the SV40 enhancer or CMV enhancer may be used to increase expression in mammalian host cells.

The expression vectors may also provide a secretion signal sequence position to form a fusion protein with polypeptides encoded by inserted C5-binding antibody sequences. More often, the inserted C5-binding antibody sequences are linked to a signal sequences before inclusion in the vector. Vectors to be used to receive sequences encoding C5-binding antibody light and heavy chain variable domains sometimes also encode constant regions or parts thereof. Such vectors allow expression of the variable regions as fusion proteins with the constant regions thereby leading to production of intact antibodies or fragments thereof. Typically, such constant regions are human.

The host cells for harboring and expressing the C5-binding antibody chains can be either prokaryotic or eukaryotic. *E. coli* is one prokaryotic host useful for cloning and expressing the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which typically contain expression control sequences compatible with the host cell (e.g., an origin of replication). In addition, any number of a variety of well-known promoters will be present, such as the lactose promoter system, a tryptophan (*trp*) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters typically control expression, optionally with an operator sequence, and have ribosome binding site sequences and the like, for initiating and completing transcription and translation. Other microbes, such as yeast, can also be

employed to express C5-binding polypeptides of the invention. Insect cells in combination with baculovirus vectors can also be used.

In some preferred embodiments, mammalian host cells are used to express and produce the C5-binding polypeptides of the present invention. For example, they can be either a hybridoma cell line expressing endogenous immunoglobulin genes (*e.g.*, the 1D6.C9 myeloma hybridoma clone as described in the Examples) or a mammalian cell line harboring an exogenous expression vector (*e.g.*, the SP2/0 myeloma cells exemplified below). These include any normal mortal or normal or abnormal immortal animal or human cell. For example, a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed including the CHO cell lines, various Cos cell lines, HeLa cells, myeloma cell lines, transformed B-cells and hybridomas. The use of mammalian tissue cell culture to express polypeptides is discussed generally in, *e.g.*, Winnacker, FROM GENES TO CLONES, VCH Publishers, N.Y., N.Y., 1987. Expression vectors for mammalian host cells can include expression control sequences, such as an origin of replication, a promoter, and an enhancer (see, *e.g.*, Queen, *et al.*, Immunol. Rev. 89:49-68, 1986), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. These expression vectors usually contain promoters derived from mammalian genes or from mammalian viruses. Suitable promoters may be constitutive, cell type-specific, stage-specific, and/or modulatable or regulatable. Useful promoters include, but are not limited to, the metallothionein promoter, the constitutive adenovirus major late promoter, the dexamethasone-inducible MMTV promoter, the SV40 promoter, the MRP polIII promoter, the constitutive MPSV promoter, the tetracycline-inducible CMV promoter (such as the human immediate-early CMV promoter), the constitutive CMV promoter, and promoter-enhancer combinations known in the art.

Methods for introducing expression vectors containing the polynucleotide sequences of interest vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts. (See generally Sambrook, *et al.*, *supra*). Other methods include, *e.g.*, electroporation, calcium phosphate treatment, liposome-mediated transformation, injection and microinjection, ballistic methods, virosomes, immunoliposomes, polycation:nucleic acid conjugates, naked DNA, artificial virions, fusion to the herpes virus structural protein VP22 (Elliot and O'Hare, Cell 88:223, 1997), agent-enhanced uptake of DNA, and *ex vivo* transduction. For long-term, high-yield production of recombinant proteins, stable expression will often be desired. For example, cell lines which stably express C5-binding antibody chains or binding fragments can be prepared using expression vectors of the invention which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector,

cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth of cells which successfully express the introduced sequences in selective media. Resistant, stably transfected cells can be proliferated using tissue culture techniques appropriate to the cell type.

5.2.2. Generation of monoclonal antibodies of the invention

Monoclonal antibodies (mAbs) can be produced by a variety of techniques, including conventional monoclonal antibody methodology e.g., the standard somatic cell hybridization technique of Kohler and Milstein, 1975 Nature 256: 495. Many techniques for producing monoclonal antibody can be employed e.g., viral or oncogenic transformation of B lymphocytes.

An animal system for preparing hybridomas is the murine system. Hybridoma production in the mouse is a well established procedure. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known.

Chimeric or humanized antibodies of the present invention can be prepared based on the sequence of a murine monoclonal antibody prepared as described above. DNA encoding the heavy and light chain immunoglobulins can be obtained from the murine hybridoma of interest and engineered to contain non-murine (e.g., human) immunoglobulin sequences using standard molecular biology techniques. For example, to create a chimeric antibody, the murine variable regions can be linked to human constant regions using methods known in the art (see e.g., U.S. Patent No. 4,816,567 to Cabilly *et al.*). To create a humanized antibody, the murine CDR regions can be inserted into a human framework using methods known in the art. See e.g., U.S. Patent No. 5225539 to Winter, and U.S. Patent Nos. 5530101; 5585089; 5693762 and 6180370 to Queen *et al.*

In a certain embodiment, the antibodies of the invention are human monoclonal antibodies. Such human monoclonal antibodies directed against C5 can be generated using transgenic or transchromosomic mice carrying parts of the human immune system rather than the mouse system. These transgenic and transchromosomic mice include mice referred to herein as HuMAb mice and KM mice, respectively, and are collectively referred to herein as "human Ig mice."

The HuMAb mouse[®] (Medarex, Inc.) contains human immunoglobulin gene miniloci that encode un-rearranged human heavy (μ and γ) and κ light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous μ and κ chain loci (see e.g., Lonberg, *et al.*, 1994 Nature 368(6474): 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or κ , and in response to immunization, the introduced human heavy and light chain transgenes undergo class switching and somatic

21489-11410

mutation to generate high affinity human IgGκ monoclonal (Lonberg, N. *et al.*, 1994 *supra*; reviewed in Lonberg, N., 1994 *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N. and Huszar, D., 1995 *Intern. Rev. Immunol.* 13: 65-93, and Harding, F. and Lonberg, N., 1995 *Ann. N. Y. Acad. Sci.* 764:536-546). The preparation and use of HuMAb mice, and the genomic modifications carried by such mice, is further described in Taylor, L. *et al.*, 1992 *Nucleic Acids Research* 20:6287-6295; Chen, J. *et al.*, 1993 *International Immunology* 5: 647-656; Tuailon *et al.*, 1993 *Proc. Natl. Acad. Sci. USA* 94:3720-3724; Choi *et al.*, 1993 *Nature Genetics* 4:117-123; Chen, J. *et al.*, 1993 *EMBO J.* 12: 821-830; Tuailon *et al.*, 1994 *J. Immunol.* 152:2912-2920; Taylor, L. *et al.*, 1994 *International Immunology* 579-591; and Fishwild, D. *et al.*, 1996 *Nature Biotechnology* 14: 845-851. See further, U.S. Patent Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay; U.S. Patent No. 5,545,807 to Surani *et al.*; PCT Publication Nos. WO 92103918, WO 93/12227, WO 94/25585, WO 97113852, WO 98/24884 and WO 99/45962, all to Lonberg and Kay; and PCT Publication No. WO 01/14424 to Korman *et al.*

In another embodiment, human antibodies of the invention can be raised using a mouse that carries human immunoglobulin sequences on transgenes and transchromosomes such as a mouse that carries a human heavy chain transgene and a human light chain transchromosome. Such mice, referred to herein as "KM mice", are described in detail in PCT Publication WO 02/43478 to Ishida *et al.*

Still further, alternative transgenic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise C5-binding antibodies of the invention. For example, an alternative transgenic system referred to as the Xenomouse (Abgenix, Inc.) can be used. Such mice are described in, *e.g.*, U.S. Patent Nos. 5,939,598; 6,075,181; 6,114,598; 6,150,584 and 6,162,963 to Kucherlapati *et al.*

Moreover, alternative transchromosomic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise C5-binding antibodies of the invention. For example, mice carrying both a human heavy chain transchromosome and a human light chain transchromosome, referred to as "TC mice" can be used; such mice are described in Tomizuka *et al.*, 2000 *Proc. Natl. Acad. Sci. USA* 97:722-727. Furthermore, cows carrying human heavy and light chain transchromosomes have been described in the art (Kuroiwa *et al.*, 2002 *Nature Biotechnology* 20:889-894) and can be used to raise C5-binding antibodies of the invention.

Human monoclonal antibodies of the invention can also be prepared using phage display methods for screening libraries of human immunoglobulin genes. Such phage display methods for isolating human antibodies are established in the art or described in the

examples below. See for example: U.S. Patent Nos. 5,223,409; 5,403,484; and 5,571,698 to Ladner *et al.*; U.S. Patent Nos. 5,427,908 and 5,580,717 to Dower *et al.*; U.S. Patent Nos. 5,969,108 and 6,172,197 to McCafferty *et al.*; and U.S. Patent Nos. 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915 and 6,593,081 to Griffiths *et al.*

Human monoclonal antibodies of the invention can also be prepared using SCID mice into which human immune cells have been reconstituted such that a human antibody response can be generated upon immunization. Such mice are described in, for example, U.S. Patent Nos. 5,476,996 and 5,698,767 to Wilson *et al.*

5.2.3. Framework or Fc engineering

Engineered antibodies of the invention include those in which modifications have been made to framework residues within VH and/or VL, *e.g.* to improve the properties of the antibody. Typically such framework modifications are made to decrease the immunogenicity of the antibody. For example, one approach is to "backmutate" one or more framework residues to the corresponding germline sequence. More specifically, an antibody that has undergone somatic mutation may contain framework residues that differ from the germline sequence from which the antibody is derived. Such residues can be identified by comparing the antibody framework sequences to the germline sequences from which the antibody is derived. To return the framework region sequences to their germline configuration, the somatic mutations can be "backmutated" to the germline sequence by, for example, site-directed mutagenesis. Such "backmutated" antibodies are also intended to be encompassed by the invention.

Another type of framework modification involves mutating one or more residues within the framework region, or even within one or more CDR regions, to remove T cell -epitopes to thereby reduce the potential immunogenicity of the antibody. This approach is also referred to as "deimmunization" and is described in further detail in U.S. Patent Publication No. 20030153043 by Carr *et al.*

In addition or alternative to modifications made within the framework or CDR regions, antibodies of the invention may be engineered to include modifications within the Fc region, typically to alter one or more functional properties of the antibody, such as serum half-life, complement fixation, Fc receptor binding, and/or antigen-dependent cellular cytotoxicity. Furthermore, an antibody of the invention may be chemically modified (*e.g.*, one or more chemical moieties can be attached to the antibody) or be modified to alter its glycosylation, again to alter one or more functional properties of the antibody. Each of these embodiments is described in further detail below. The numbering of residues in the Fc region is that of the EU index of Kabat.

In one embodiment, the hinge region of CH1 is modified such that the number of cysteine residues in the hinge region is altered, e.g., increased or decreased. This approach is described further in U.S. Patent No. 5,677,425 by Bodmer *et al.* The number of cysteine residues in the hinge region of CH1 is altered to, for example, facilitate assembly of the light and heavy chains or to increase or decrease the stability of the antibody.

In another embodiment, the Fc hinge region of an antibody is mutated to decrease the biological half-life of the antibody. More specifically, one or more amino acid mutations are introduced into the CH2-CH3 domain interface region of the Fc-hinge fragment such that the antibody has impaired Staphylococcal protein A (SpA) binding relative to native Fc-hinge domain SpA binding. This approach is described in further detail in U.S. Patent No. 6,165,745 by Ward *et al.*

In another embodiment, the antibody is modified to increase its biological half-life. Various approaches are possible. For example, one or more of the following mutations can be introduced: T252L, T254S, T256F, as described in U.S. Patent No. 6,277,375 to Ward. Alternatively, to increase the biological half life, the antibody can be altered within the CH1 or CL region to contain a salvage receptor binding epitope taken from two loops of a CH2 domain of an Fc region of an IgG, as described in U.S. Patent Nos. 5,869,046 and 6,121,022 by Presta *et al.*

In yet other embodiments, the Fc region is altered by replacing at least one amino acid residue with a different amino acid residue to alter the effector functions of the antibody. For example, one or more amino acids can be replaced with a different amino acid residue such that the antibody has an altered affinity for an effector ligand but retains the antigen-binding ability of the parent antibody. The effector ligand to which affinity is altered can be, for example, an Fc receptor or the C1 component of complement. This approach is described in further detail in U.S. Patent Nos. 5,624,821 and 5,648,260, both by Winter *et al.*

In another embodiment, one or more amino acids selected from amino acid residues can be replaced with a different amino acid residue such that the antibody has altered C1q binding and/or reduced or abolished complement dependent cytotoxicity (CDC). This approach is described in further detail in U.S. Patent Nos. 6,194,551 by Idusogie *et al.*

In another embodiment, one or more amino acid residues are altered to thereby alter the ability of the antibody to fix complement. This approach is described further in PCT Publication WO 94/29351 by Bodmer *et al.*

In yet another embodiment, the Fc region is modified to increase the ability of the antibody to mediate antibody dependent cellular cytotoxicity (ADCC) and/or to increase the affinity of the antibody for an Fcγ receptor by modifying one or more amino acids. This approach is described further in PCT Publication WO 00/42072 by Presta. Moreover, the

binding sites on human IgG1 for FcγRI, FcγRII, FcγRIII and FcRn have been mapped and variants with improved binding have been described (see Shields, R.L. *et al.*, 2001 J. Biol. Chem. 276:6591-6604).

In still another embodiment, the glycosylation of an antibody is modified. For example, an aglycosylated antibody can be made (*i.e.*, the antibody lacks glycosylation). Glycosylation can be altered to, for example, increase the affinity of the antibody for "antigen". Such carbohydrate modifications can be accomplished by, for example, altering one or more sites of glycosylation within the antibody sequence. For example, one or more amino acid substitutions can be made that result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site. Such aglycosylation may increase the affinity of the antibody for antigen. Such an approach is described in further detail in U.S. Patent Nos. 5,714,350 and 6,350,861 by Co *et al.*

Additionally or alternatively, an antibody can be made that has an altered type of glycosylation, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNac structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Such carbohydrate modifications can be accomplished by, for example, expressing the antibody in a host cell with altered glycosylation machinery. Cells with altered glycosylation machinery have been described in the art and can be used as host cells in which to express recombinant antibodies of the invention to thereby produce an antibody with altered glycosylation. For example, EP 1,176,195 by Hang *et al.* describes a cell line with a functionally disrupted FUT8 gene, which encodes a fucosyl transferase, such that antibodies expressed in such a cell line exhibit hypofucosylation. PCT Publication WO 03/035835 by Presta describes a variant CHO cell line, Lecl3 cells, with reduced ability to attach fucose to Asn(297)-linked carbohydrates, also resulting in hypofucosylation of antibodies expressed in that host cell (see also Shields, R.L. *et al.*, 2002 J. Biol. Chem. 277:26733-26740). PCT Publication WO 99/54342 by Umana *et al.* describes cell lines engineered to express glycoprotein-modifying glycosyl transferases (*e.g.*, beta(1,4)-N acetylglucosaminyltransferase III (GnTIII)) such that antibodies expressed in the engineered cell lines exhibit increased bisecting GlcNac structures which results in increased ADCC activity of the antibodies (see also Umana *et al.*, 1999 Nat. Biotech. 17:176-180).

5.2.4. Methods of Engineering Altered Antibodies

As discussed above, the C5-binding antibodies having VH and VL sequences or full length heavy and light chain sequences shown herein can be used to create new C5-binding antibodies by modifying full length heavy chain and/or light chain sequences, VH and/or VL sequences, or the constant region(s) attached thereto. Thus, in another aspect of the

invention, the structural features of a C5-binding antibody of the invention are used to create structurally related C5-binding antibodies that retain at least one functional property of the antibodies of the invention, such as binding to human C5 and also inhibiting one or more functional properties of C5 (e.g., inhibit red blood cell lysis in a hemolytic assay).

For example, one or more CDR regions of the antibodies of the present invention, or mutations thereof, can be combined recombinantly with known framework regions and/or other CDRs to create additional, recombinantly-engineered, C5-binding antibodies of the invention, as discussed above. Other types of modifications include those described in the previous section. The starting material for the engineering method is one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. To create the engineered antibody, it is not necessary to actually prepare (*i.e.*, express as a protein) an antibody having one or more of the VH and/or VL sequences provided herein, or one or more CDR regions thereof. Rather, the information contained in the sequence(s) is used as the starting material to create a "second generation" sequence(s) derived from the original sequence(s) and then the "second generation" sequence(s) is prepared and expressed as a protein.

Accordingly, in another embodiment, the invention provides a method for preparing an C5-binding antibody consisting of: a heavy chain variable region antibody sequence having a CDR1 sequence selected from the group consisting of SEQ ID NOs: 1, 17, 33, 61, 131, 145, 159, 173, 195, and 235, a CDR2 sequence selected from the group consisting of SEQ ID NOs: 2, 18, 34, 49, 62, 77, 95, 107, 113, 119, 132, 146, 160, 174, 196, 226, and 236, and/or a CDR3 sequence selected from the group consisting of SEQ ID NOs: 3, 19, 35, 63, 133, 147, 161, 175, 197, and 237; and a light chain variable region antibody sequence having a CDR1 sequence selected from the group consisting of SEQ ID NOs: 4, 20, 36, 64, 134, 148, 162, 176, 198, and 238, a CDR2 sequence selected from the group consisting of SEQ ID NOs: 5, 21, 37, 65, 135, 149, 163, 177, 199, and 239, and/or a CDR3 sequence selected from the group consisting of SEQ ID NOs: 6, 22, 38, 50, 66, 78, 89, 101, 120, 136, 150, 164, 178, 200, 209, and 240; altering at least one amino acid residue within the heavy chain variable region antibody sequence and/or the light chain variable region antibody sequence to create at least one altered antibody sequence; and expressing the altered antibody sequence as a protein.

Accordingly, in another embodiment, the invention provides a method for preparing an C5-binding antibody optimized for expression in a mammalian cell consisting of: a full length heavy chain antibody sequence having a sequence selected from the group of SEQ ID NOs: 9, 25, 41, 53, 69, 81, 97, 109, 115, 123, 139, 153, 167, 181, 189, 203, 212, 220, 229, 243, 249, 254, 258, 274, 278, and 282; and a full length light chain antibody sequence having a sequence selected from the group of 10, 26, 42, 54, 70, 82, 91, 103, 124, 140, 154,

168, 182, 190, 204, 213, 221, 230, 244, 251, 262, 266, 270, 286, and 290; altering at least one amino acid residue within the full length heavy chain antibody sequence and/or the full length light chain antibody sequence to create at least one altered antibody sequence; and expressing the altered antibody sequence as a protein.

The altered antibody sequence can also be prepared by screening antibody libraries having fixed CDR3 sequences or minimal essential binding determinants as described in US2005025552 and diversity on CDR1 and CDR2 sequences. The screening can be performed according to any screening technology appropriate for screening antibodies from antibody libraries, such as phage display technology.

Standard molecular biology techniques can be used to prepare and express the altered antibody sequence. The antibody encoded by the altered antibody sequence(s) is one that retains one, some or all of the functional properties of the C5-binding antibodies described herein, which functional properties include, but are not limited to, specifically binding to human and/or cynomolgus C5; and the antibody inhibit red blood cell lysis in a hemolytic assay.

The functional properties of the altered antibodies can be assessed using standard assays available in the art and/or described herein, such as those set forth in the Examples (*e.g.*, ELISAs).

In certain embodiments of the methods of engineering antibodies of the invention, mutations can be introduced randomly or selectively along all or part of an C5-binding antibody coding sequence and the resulting modified C5-binding antibodies can be screened for binding activity and/or other functional properties as described herein. Mutational methods have been described in the art. For example, PCT Publication WO 02/092780 by Short describes methods for creating and screening antibody mutations using saturation mutagenesis, synthetic ligation assembly, or a combination thereof. Alternatively, PCT Publication WO 03/074679 by Lazar *et al.* describes methods of using computational screening methods to optimize physiochemical properties of antibodies.

5.3. Characterization of the Antibodies of the Invention

The antibodies of the invention can be characterized by various functional assays. For example, they can be characterized by their ability to inhibit red blood cell lysis in hemolytic assays, their affinity to a C5 protein (*e.g.*, human and/or cynomolgus C5), the epitope binning, their resistance to proteolysis, and their ability to block the complement cascade, for example, their ability to inhibit MAC formation.

Various methods can be used to measure presence of complement pathway molecules and activation of the complement system (see, *e.g.*, U.S. Pat. No. 6,087,120; and

Newell *et al.*, J Lab Clin Med, 100:437-44, 1982). For example, the complement activity can be monitored by (i) measurement of inhibition of complement-mediated lysis of red blood cells (hemolysis); (ii) measurement of ability to inhibit cleavage of C3 or C5; and (iii) inhibition of alternative pathway mediated hemolysis.

The two most commonly used techniques are hemolytic assays (see, *e.g.*, Baatrup *et al.*, Ann Rheum Dis, 51:892-7, 1992) and immunological assays (see, *e.g.*, Auda *et al.*, Rheumatol Int, 10:185-9, 1990). The hemolytic techniques measure the functional capacity of the entire sequence-either the classical or alternative pathway. Immunological techniques measure the protein concentration of a specific complement component or split product. Other assays that can be employed to detect complement activation or measure activities of complement components in the methods of the present invention include, *e.g.*, T cell proliferation assay (Chain *et al.*, J Immunol Methods, 99:221-8, 1987), and delayed type hypersensitivity (DTH) assay (Forstrom *et al.*, 1983, Nature 303:627-629; Halliday *et al.*, 1982, in Assessment of Immune Status by the Leukocyte Adherence Inhibition Test, Academic, New York pp. 1-26; Koppi *et al.*, 1982, Cell. Immunol. 66:394-406; and U.S. Pat. No. 5,843,449).

In hemolytic techniques, all of the complement components must be present and functional. Therefore hemolytic techniques can screen both functional integrity and deficiencies of the complement system (see, *e.g.*, Dijk *et al.*, J Immunol Methods 36: 29-39, 1980; Minh *et al.*, Clin Lab Haematol. 5:23-34 1983; and Tanaka *et al.*, J Immunol 86: 161-170, 1986). To measure the functional capacity of the classical pathway, sheep red blood cells coated with hemolysin (rabbit IgG to sheep red blood cells) or chicken red blood cells that are sensitized with rabbit anti-chicken antibodies are used as target cells (sensitized cells). These Ag-Ab complexes activate the classical pathway and result in lysis of the target cells when the components are functional and present in adequate concentration. To determine the functional capacity of the alternative pathway, rabbit red blood cells are used as the target cell (see, *e.g.*, U.S. Pat. No. 6,087,120).

To test the ability of an antibody to inhibit MAC (membrane attack complex) formation, a MAC deposition assay can be performed. Briefly, zymosan can be used to activate the alternative pathway and IgM can be used to active the classic pathway. Fabs are pre-inclubated with human serum and added to plates coated with zymosan or IgM. Percentage inhibition of MAC deposition can be calculated for each sample relative to baseline (EDTA treated human serum) and positive control (human serum).

The ability of an antibody to bind to C5 can be detected by labelling the antibody of interest directly, or the antibody may be unlabelled and binding detected indirectly using various sandwich assay formats known in the art.

In some embodiments, the C5-binding antibodies of the invention block or compete with binding of a reference C5-binding antibody to a C5 polypeptide. These can be fully human C5-binding antibodies described above. They can also be other mouse, chimeric or humanized C5-binding antibodies which bind to the same epitope as the reference antibody. The capacity to block or compete with the reference antibody binding indicates that a C5-binding antibody under test binds to the same or similar epitope as that defined by the reference antibody, or to an epitope which is sufficiently proximal to the epitope bound by the reference C5-binding antibody. Such antibodies are especially likely to share the advantageous properties identified for the reference antibody. The capacity to block or compete with the reference antibody may be determined by, e.g., a competition binding assay. With a competition binding assay, the antibody under test is examined for ability to inhibit specific binding of the reference antibody to a common antigen, such as a C5 polypeptide. A test antibody competes with the reference antibody for specific binding to the antigen if an excess of the test antibody substantially inhibits binding of the reference antibody. Substantial inhibition means that the test antibody reduces specific binding of the reference antibody usually by at least 10%, 25%, 50%, 75%, or 90%.

There are a number of known competition binding assays that can be used to assess competition of a C5-binding antibody with the reference C5-binding antibody for binding to a C5 protein. These include, e.g., solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahl *et al.*, Methods in Enzymology 9:242-253, 1983); solid phase direct biotin-avidin EIA (see Kirkland *et al.*, J. Immunol. 137:3614-3619, 1986); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow & Lane, *supra*); solid phase direct label RIA using I-125 label (see Morel *et al.*, Molec. Immunol. 25:7-15, 1988); solid phase direct biotin-avidin EIA (Cheung *et al.*, Virology 176:546-552, 1990); and direct labeled RIA (Moldenhauer *et al.*, Scand. J. Immunol. 32:77-82, 1990). Typically, such an assay involves the use of purified antigen bound to a solid surface or cells bearing either of these, an unlabelled test C5-binding antibody and a labelled reference antibody. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test antibody. Usually the test antibody is present in excess. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur.

To determine if the selected C5-binding monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (e.g., reagents from Pierce, Rockford, IL). Competition studies using unlabeled monoclonal

antibodies and biotinylated monoclonal antibodies can be performed using a C5 polypeptide coated-ELISA plates. Biotinylated MAb binding can be detected with a strep-avidin-alkaline phosphatase probe. To determine the isotype of a purified C5-binding antibody, isotype ELISAs can be performed. For example, wells of microtiter plates can be coated with 1 µg/ml of anti-human IgG overnight at 4°C. After blocking with 1% BSA, the plates are reacted with 1 µg/ml or less of the monoclonal C5-binding antibody or purified isotype controls, at ambient temperature for one to two hours. The wells can then be reacted with either human IgG1 or human IgM-specific alkaline phosphatase-conjugated probes. Plates are then developed and analyzed so that the isotype of the purified antibody can be determined.

To demonstrate binding of monoclonal C5-binding antibodies to live cells expressing a C5 polypeptide, flow cytometry can be used. Briefly, cell lines expressing C5 (grown under standard growth conditions) can be mixed with various concentrations of a C5-binding antibody in PBS containing 0.1% BSA and 10% fetal calf serum, and incubated at 37°C for 1 hour. After washing, the cells are reacted with Fluorescein-labeled anti-human IgG antibody under the same conditions as the primary antibody staining. The samples can be analyzed by FACScan instrument using light and side scatter properties to gate on single cells. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells can be stained exactly as described above and examined by fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

C5-binding antibodies of the invention can be further tested for reactivity with a C5 polypeptide or antigenic fragment by Western blotting. Briefly, purified C5 polypeptides or fusion proteins, or cell extracts from cells expressing C5 can be prepared and subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens are transferred to nitrocellulose membranes, blocked with 10% fetal calf serum, and probed with the monoclonal antibodies to be tested. Human IgG binding can be detected using anti-human IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (Sigma Chem. Co., St. Louis, MO).

Examples of functional assays are also described in the Example section below.

5.4. Prophylactic and Therapeutic Uses

The present invention provides methods of treating a disease or disorder associated with increased complement activity by administering to a subject in need thereof an effective amount of the antibodies of the invention. In a specific embodiment, the present invention provides a method of treating age-related macular degeneration (AMD) by administering to a subject in need thereof an effective amount of the antibodies of the invention.

The antibodies of the invention can be used, *inter alia*, to prevent progression of dry AMD to wet AMD, to slow and/or prevent progression of geographic atrophy, and to improve vision lost due to dry AMD progression. It can also be used in combination with anti-VEGF therapies for the treatment of wet AMD patients.

In some embodiments, the present invention provides methods of treating a complement related disease or disorder by administering to a subject in need thereof an effective amount of the antibodies of the invention. Examples of known complement related diseases or disorders include: neurological disorders, multiple sclerosis, stroke, Guillain Barre Syndrome, traumatic brain injury, Parkinson's disease, disorders of inappropriate or undesirable complement activation, hemodialysis complications, hyperacute allograft rejection, xenograft rejection, interleukin-2 induced toxicity during IL-2 therapy, inflammatory disorders, inflammation of autoimmune diseases, Crohn's disease, adult respiratory distress syndrome, thermal injury including burns or frostbite, post-ischemic reperfusion conditions, myocardial infarction, balloon angioplasty, post-pump syndrome in cardiopulmonary bypass or renal bypass, hemodialysis, renal ischemia, mesenteric artery reperfusion after aortic reconstruction, infectious disease or sepsis, immune complex disorders and autoimmune diseases, rheumatoid arthritis, systemic lupus erythematosus (SLE), SLE nephritis, proliferative nephritis, hemolytic anemia, and myasthenia gravis. In addition, other known complement related disease are lung disease and disorders such as dyspnea, hemoptysis, ARDS, asthma, chronic obstructive pulmonary disease (COPD), emphysema, pulmonary embolisms and infarcts, pneumonia, fibrogenic dust diseases, inert dusts and minerals (e.g., silicon, coal dust, beryllium, and asbestos), pulmonary fibrosis, organic dust diseases, chemical injury (due to irritant gasses and chemicals, e.g., chlorine, phosgene, sulfur dioxide, hydrogen sulfide, nitrogen dioxide, ammonia, and hydrochloric acid), smoke injury, thermal injury (e.g., burn, freeze), asthma, allergy, bronchoconstriction, hypersensitivity pneumonitis, parasitic diseases, Goodpasture's Syndrome, pulmonary vasculitis, and immune complex-associated inflammation.

In a specific embodiment, the present invention provides methods of treating a complement related disease or disorder by administering to a subject in need thereof an effective amount of the antibodies of the invention, wherein said disease or disorder is asthma, arthritis (e.g., rheumatoid arthritis), autoimmune heart disease, multiple sclerosis, inflammatory bowel disease, ischemia-reperfusion injuries, Barraquer-Simons Syndrome, hemodialysis, systemic lupus, lupus erythematosus, psoriasis, multiple sclerosis, transplantation, diseases of the central nervous system such as Alzheimer's disease and other neurodegenerative conditions, aHUS, glomerulonephritis, bullous pemphigoid or MPGN II.

In a specific embodiment, the present invention provides methods of treating glomerulonephritis by administering to a subject in need thereof an effective amount of a composition comprising an antibody of the present invention. Symptoms of glomerulonephritis include, but not limited to, proteinuria; reduced glomerular filtration rate (GFR); serum electrolyte changes including azotemia (uremia, excessive blood urea nitrogen--BUN) and salt retention, leading to water retention resulting in hypertension and edema; hematuria and abnormal urinary sediments including red cell casts; hypoalbuminemia; hyperlipidemia; and lipiduria. In a specific embodiment, the present invention provides methods of treating paroxysmal nocturnal hemoglobinuria (PNH) by administering to a subject in need thereof an effective amount of a composition comprising an antibody of the present invention.

In a specific embodiment, the present invention provides methods of reducing the dysfunction of the immune and hemostatic systems associated with extracorporeal circulation by administering to a subject in need thereof an effective amount of a composition comprising an antibody of the present invention. The antibodies of the present invention can be used in any procedure which involves circulating the patient's blood from a blood vessel of the patient, through a conduit, and back to a blood vessel of the patient, the conduit having a luminal surface comprising a material capable of causing at least one of complement activation, platelet activation, leukocyte activation, or platelet-leukocyte adhesion. Such procedures include, but are not limited to, all forms of ECC, as well as procedures involving the introduction of an artificial or foreign organ, tissue, or vessel into the blood circuit of a patient.

Subjects to be treated with therapeutic agents of the present invention can also be administered other therapeutic agents with known methods of treating conditions associated with macular degeneration, such as antibiotic treatments as described in U.S. Pat. No. 6,218,368. In other treatments, immunosuppressive agents such as cyclosporine, are agents capable of suppressing immune responses. These agents include cytotoxic drugs, corticosteroids, nonsteroidal anti-inflammatory drugs (NSAIDs), specific T-lymphocyte immunosuppressants, and antibodies or fragments thereof (see Physicians' Desk Reference, 53rd edition, Medical Economics Company Inc., Montvale, N.J. (1999)). Immunosuppressive treatment is typically continued at intervals for a period of a week, a month, three months, six months or a year. In some patients, treatment is administered for up to the rest of a patient's life.

When the therapeutic agents of the present invention are administered together with another agent, the two can be administered sequentially in either order or simultaneously. In some aspects, an antibody of the present invention is administered to a subject who is also

receiving therapy with a second agent (e.g., verteporfin). In other aspects, the binding molecule is administered in conjunction with surgical treatments.

Suitable agents for combination treatment with C5-binding antibodies include agents known in the art that are able to modulate the activities of complement components (see, e.g., U.S. Pat. No. 5,808,109). Other agents have been reported to diminish complement-mediated activity. Such agents include: amino acids (Takada, Y. *et al.* Immunology 1978, 34, 509); phosphonate esters (Becker, L. Biochem. Biophys. Acta 1967, 147, 289); polyanionic substances (Conrow, R. B. *et al.* J. Med. Chem. 1980, 23, 242); sulfonyl fluorides (Hansch, C.; Yoshimoto, M. J. Med. Chem. 1974, 17, 1160, and references cited therein); polynucleotides (DeClercq, P. F. *et al.* Biochem. Biophys. Res. Commun. 1975, 67, 255); pimaric acids (Glovsky, M. M. *et al.* J. Immunol. 1969, 102, 1); porphines (Lapidus, M. and Tomasco, J. Immunopharmacol. 1981, 3, 137); several antiinflammatories (Burge, J. J. *et al.* J. Immunol. 1978, 120, 1625); phenols (Muller-Eberhard, H. J. 1978, in Molecular Basis of Biological Degradative Processes, Berlin, R. D. *et al.*, eds. Academic Press, New York, p. 65); and benzamidines (Vogt, W. *et al.* Immunology 1979, 36, 138). Some of these agents function by general inhibition of proteases and esterases. Others are not specific to any particular intermediate step in the complement pathway, but, rather, inhibit more than one step of complement activation. Examples of the latter compounds include the benzamidines, which block C1, C4 and C5 utilization (see, e.g., Vogt *et al.* Immunol. 1979, 36, 138).

Additional agents known in the art that can inhibit activity of complement components include K-76, a fungal metabolite from *Stachybotrys* (Corey *et al.*, J. Amer. Chem. Soc. 104: 5551, 1982). Both K-76 and K-76 COOH have been shown to inhibit complement mainly at the C5 step (Hong *et al.*, J. Immunol. 122: 2418, 1979; Miyazaki *et al.*, Microbiol. Immunol. 24: 1091, 1980), and to prevent the generation of a chemotactic factor from normal human complement (Bumpers *et al.*, Lab. Clin. Med. 102: 421, 1983). At high concentrations of K-76 or K-76 COOH, some inhibition of the reactions of C2, C3, C6, C7, and C9 with their respective preceding intermediaries is exhibited. K-76 or K-76 COOH has also been reported to inhibit the C3b inactivator system of complement (Hong *et al.*, J. Immunol. 127: 104-108, 1981). Other suitable agents for practicing methods of the present invention include griseofulvin (Weinberg, in Principles of Medicinal Chemistry, 2d Ed., Foye, W. O., ed., Lea & Febiger, Philadelphia, Pa., p. 813, 1981), isopannarin (Djura *et al.*, Aust. J. Chem. 36: 1057, 1983), and metabolites of *Siphonodictyon coralli-phagum* (Sullivan *et al.*, Tetrahedron 37: 979, 1981).

A combination therapy regimen may be additive, or it may produce synergistic results (e.g., reductions in complement pathway activity more than expected for the combined use of the two agents). In some embodiments, the present invention provide a combination therapy for preventing and/or treating AMD or another complement related disease as described

above with a C5-binding antibody of the invention and an anti-angiogenic, such as anti-VEGF agent.

5.5. Diagnostic Uses

In one aspect, the invention encompasses diagnostic assays for determining C5 protein and/or nucleic acid expression as well as C5 protein function, in the context of a biological sample (e.g., blood, serum, cells, tissue) or from individual is afflicted with a disease or disorder, or is at risk of developing a disorder associated with AMD.

Diagnostic assays, such as competitive assays rely on the ability of a labelled analogue (the "tracer") to compete with the test sample analyte for a limited number of binding sites on a common binding partner. The binding partner generally is insolubilized before or after the competition and then the tracer and analyte bound to the binding partner are separated from the unbound tracer and analyte. This separation is accomplished by decanting (where the binding partner was preinsolubilized) or by centrifuging (where the binding partner was precipitated after the competitive reaction). The amount of test sample analyte is inversely proportional to the amount of bound tracer as measured by the amount of marker substance. Dose-response curves with known amounts of analyte are prepared and compared with the test results in order to quantitatively determine the amount of analyte present in the test sample. These assays are called ELISA systems when enzymes are used as the detectable markers. In an assay of this form, competitive binding between antibodies and C5-binding antibodies results in the bound C5 protein, preferably the C5 epitopes of the invention, being a measure of antibodies in the serum sample, most particularly, neutralising antibodies in the serum sample.

A significant advantage of the assay is that measurement is made of neutralising antibodies directly (*i.e.*, those which interfere with binding of C5 protein, specifically, epitopes). Such an assay, particularly in the form of an ELISA test has considerable applications in the clinical environment and in routine blood screening.

Immunologic techniques employ polyclonal or monoclonal antibodies against the different epitopes of the various complement components (e.g., C3, C4, C5) to detect, e.g., the split products of complement components (see, e.g., Hugli *et al.*, Immunoassays Clinical Laboratory Techniques 443-460, 1980; Gorski *et al.*, J Immunol Meth 47: 61-73, 1981; Linder *et al.*, J Immunol Meth 47: 49-59, 1981; and Burger *et al.*, J Immunol 141: 553-558, 1988). Binding of the antibody with the split product in competition with a known concentration of labeled split product could then be measured. Various assays such as radio-immunoassays, ELISA's, and radial diffusion assays are available to detect complement split products.

The immunologic techniques provide high sensitivity to detect complement activation, since they allow measurement of split-product formation in blood from a test subject and

control subjects with or without macular degeneration-related disorders. Accordingly, in some methods of the present invention, diagnosis of a disorder associated with ocular disorders is obtained by measurement of abnormal complement activation through quantification of the soluble split products of complement components in blood plasma from a test subject. The measurements can be performed as described, *e.g.*, in Chenoweth *et al.*, N Engl J Med 304: 497-502, 1981; and Bhakdi *et al.*, Biochim Biophys Acta 737: 343-372, 1983. Preferably, only the complement activation formed *in vivo* is measured. This can be accomplished by collecting a biological sample from the subject (*e.g.*, serum) in medium containing inhibitors of the complement system, and subsequently measuring complement activation (*e.g.*, quantification of the split products) in the sample.

In the clinical diagnosis or monitoring of patients with disorders associated with ocular diseases or disorders, the detection of complement proteins in comparison to the levels in a corresponding biological sample from a normal subject is indicative of a patient with disorders associated with macular degeneration.

In vivo diagnostic or imaging is described in US2006/0067935. Briefly, these methods generally comprise administering or introducing to a patient a diagnostically effective amount of a C5 binding molecule that is operatively attached to a marker or label that is detectable by non-invasive methods. The antibody-marker conjugate is allowed sufficient time to localize and bind to complement proteins within the eye. The patient is then exposed to a detection device to identify the detectable marker, thus forming an image of the location of the C5 binding molecules in the eye of a patient. The presence of C5 binding antibody or an antigen-binding fragment thereof is detected by determining whether an antibody-marker binds to a component of the eye. Detection of an increased level in selected complement proteins or a combination of protein in comparison to a normal individual without AMD disease is indicative of a predisposition for and/or on set of disorders associated with macular degeneration. These aspects of the invention are also preferred for use in eye imaging methods and combined angiogenic diagnostic and treatment methods.

The invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically.

The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with dysregulation of complement pathway activity. For example, mutations in a C5 gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with C5 protein, nucleic acid expression or activity.

Another aspect of the invention provides methods for determining C5 nucleic acid expression or C5 protein activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (e.g., drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (e.g., the genotype of the individual examined to determine the ability of the individual to respond to a particular agent.)

Yet another aspect of the invention pertains to monitoring the influence of agents (e.g., drugs) on the expression or activity of C5 protein in clinical trials.

5.6. Pharmaceutical Compositions

The invention provides pharmaceutical compositions comprising the C5-binding antibodies (intact or binding fragments) formulated together with a pharmaceutically acceptable carrier. The compositions can additionally contain one or more other therapeutic agents that are suitable for treating or preventing a complement-associated disease (e.g., AMD). Pharmaceutically carriers enhance or stabilize the composition, or to facilitate preparation of the composition. Pharmaceutically acceptable carriers include solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible.

A pharmaceutical composition of the present invention can be administered by a variety of methods known in the art. The route and/or mode of administration vary depending upon the desired results. It is preferred that administration be intravenous, intramuscular, intraperitoneal, or subcutaneous, or administered proximal to the site of the target. In a specific embodiment, the antibodies of the invention are formulated so that they can be administered intravitreally into the eye. The pharmaceutically acceptable carrier should be suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound, *i.e.*, antibody, bispecific and multispecific molecule, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

The composition should be sterile and fluid. Proper fluidity can be maintained, for example, by use of coating such as lecithin, by maintenance of required particle size in the case of dispersion and by use of surfactants. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol or sorbitol, and sodium chloride in the composition. Long-term absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate or gelatin.

Pharmaceutical compositions of the invention can be prepared in accordance with methods well known and routinely practiced in the art. See, e.g., Remington: The Science and Practice of Pharmacy, Mack Publishing Co., 20th ed., 2000; and Sustained and Controlled Release Drug Delivery Systems, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978. Pharmaceutical compositions are preferably manufactured under GMP conditions. Typically, a therapeutically effective dose or efficacious dose of the C5-binding antibody is employed in the pharmaceutical compositions of the invention. The C5-binding antibodies are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention can be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level depends upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors.

A physician or veterinarian can start doses of the antibodies of the invention employed in the pharmaceutical composition at levels lower than that required to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved. In general, effective doses of the compositions of the present invention, for the treatment of an allergic inflammatory disorder described herein vary depending upon many different factors, including means of administration, target site, physiological state of the patient, whether the patient is human or an animal, other medications administered, and whether treatment is prophylactic or therapeutic. Treatment dosages need to be titrated to optimize safety and efficacy. For systemic administration with an antibody, the dosage

ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 15 mg/kg, of the host body weight. An exemplary treatment regime entails systemic administration once per every two weeks or once a month or once every 3 to 6 months. For intravitreal administration with an antibody, the dosage ranges from about 0.0001 to about 10 mg. An exemplary treatment regime entails systemic administration once per every two weeks or once a month or once every 3 to 6 months.

Antibody is usually administered on multiple occasions. Intervals between single dosages can be weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of C5-binding antibody in the patient. In some methods of systemic administration, dosage is adjusted to achieve a plasma antibody concentration of 1–1000 µg/ml and in some methods 25–500 µg/ml. Alternatively, antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, humanized antibodies show longer half life than that of chimeric antibodies and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

6. EXAMPLES

The following examples are provided to further illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims.

Example 1 : Generation of Cynomolgus C5 and Human C5

1. Generation of Cynomolgus C5

Cynomolgus C5 was purified successfully from cynomolgus serum by affinity chromatography using MOR07086 hu IgG1. Cynomolgus C5 was quality tested by SDS-PAGE, Western blot, mass spectrometry and hemolytic assays. Quality of purified cynomolgus C5 was shown to be high by SDS-PAGE and Western blot. Lack of C3 contamination was confirmed by SDS and Western blot. In addition, the identity of cynomolgus C5 sequence was determined by mass spectrometric analysis and the activity of purified cynomolgus C5 was tested in hemolytic assays. In hemolytic assays the new preparation was equipotent to human C5 (*e.g.*, Sample 6, which was used in affinity

21489-11410

maturation panning, reconstituted complement activity of 20% human C5-depleted serum with similar activity to purified human C5).

2. Quality Control of Human and Cynomolgus Biotinylated and Non-Biotinylated C5 Proteins

Bioactivity of purified human C5 was characterized and confirmed by the alternative pathway hemolytic activity. C5 was spiked into C5-depleted human serum at varying concentrations to obtain an EC50. EC50 values ranging between 0.02-0.1nM were considered acceptable.

Before using, the bioactivity of every purified human C5 protein lot was tested in the hemolytic assay. The same quality control was done for cynomolgus C5 after purification from cynomolgus serum. After biotinylation of human and cynomolgus C5, the bioactivity of the material was also tested in hemolytic assays, in order to analyze if there was a loss of activity caused by biotinylation.

Example 2: Generation of C5-specific antibodies from the HuCAL GOLD® Library

C5 antibodies were generated by selection of clones having high binding affinities, using as the source of antibody variant proteins a commercially available phage display library, the MorphoSys HuCAL GOLD® library.

HuCAL GOLD® library is a Fab library (Knappik *et al.*, 2000) in which all six CDRs are diversified by appropriate mutation, and which employs the CysDisplay™ technology for linking the Fab to the phage surface (WO01/05950, Löhning *et al.*, 2001).

1. Selection by panning of C5 specific antibodies from the library

For the selection of antibodies against C5, two different panning strategies were applied. The six different pools were individually subjected to three rounds of: (a) a solid phase panning where the antigens (human and cynomolgus C5) were directly coated on Maxisorp 96 well microtiter plates (Nunc, Wiesbaden, Germany); or (b) a solution panning with biotinylated human and cyno C5 where the phage-antigen complex was captured by Streptavidin magnetic beads (Dynabeads M-280; Dynal) for each panning pool.

The HuCAL GOLD® library was amplified in 2xYT medium containing 34µg/ml chloramphenicol and 1% glucose (2xYT-CG). After infection with VCSM13 helper phage at an OD_{600nm} of 0.5 (30 min at 37°C without shaking; 30 min at 37°C shaking at 250 rpm), cells were spun down (4120 g; 5 min; 4°C), resuspended in 2xYT/ 34µg/ml chloramphenicol/ 50 µg/ml kanamycin/ 0.25mM IPTG and grown overnight at 22°C. Phage were PEG-precipitated from the supernatant, resuspended in PBS/ 20% glycerol and stored at -80°C. Phage amplification between two panning rounds was conducted as follows: mid-log phase *E. coli* TG1 cells were infected with eluted phage and plated onto LB-agar supplemented with

21489-11410

1% of glucose and 34 µg/ml of chloramphenicol (LB-CG plates). After overnight incubation at 30°C, the TG1 colonies were scraped off the agar plates and used to inoculate 2xYT-CG until an OD_{600nm} of 0.5 was reached. VCSM13 helper phage were added for infection as described above.

Taken together 354 clones derived from all panning strategies were sequenced, resulting in 64 unique clones with the desired profile: binding to human and cynomolgus C5 and no binding to the counter targets C3 and C4.

45 clones derived from solid phase pannings and 19 clones from solution pannings were selected for protein expression and purification. Four Fabs from solid phase pannings (MOR06525, MOR06756, MOR06757 and MOR06763) and 6 Fabs from solution pannings (MOR07086, MOR07087, MOR07091, MOR07092, MOR07093 and MOR07094) entered affinity maturation.

Solid Phase Panning Against C5 on Directly Coated Protein

The first panning variant was solid phase panning alternating human C5 (first and third round of selection) and cynomolgus C5 (second round of selection).

Three wells of a Maxisorp plate (F96 Nunc-Immunoplate) were coated with 200 µl of 50nM C5 each o/n at 4°C. The coated wells were washed 2x with 400 µl PBS and blocked with 350 µl 5% MPBS for 2h at RT on a microtiter plate shaker. For each panning about 10¹³ HuCAL GOLD[®] phage-antibodies were blocked with equal volume of PBST/5% milk powder for 2h at room temperature. The coated wells were washed 2x with 400 µl PBS after the blocking procedure. 200 µl of pre-blocked HuCAL GOLD[®] phage-antibodies were added to each coated well and incubated for 2h at RT on a shaker. Washing was performed by adding five times 350 µl PBS/0.05% Tween[™], followed by washing another five times with PBS. For some panning conditions a more stringent wash procedure was applied.

Elution of phage from the plate was performed with 200 µl 20mM DTT in 10mM Tris/HCl pH8 per well for 10 min. The DTT phage eluate was added to 15ml of E.coli TG1, which were grown to an OD₆₀₀ of 0.6-0.8 at 37°C in 2YT medium and incubated in 50ml plastic tubes for 45min at 37°C without shaking for phage infection. After centrifugation for 5 min at 4120 x g, the bacterial pellets were each resuspended in 600 µl 2xYT medium, plated on 3xYT-CG agar plates and incubated overnight at 37°C. Colonies were scraped off the plates and phages were rescued and amplified as described above.

The second and third rounds of solid phase panning were performed according to the protocol of the first round. In the second selection round for some panning conditions the output of the first round was used for selections on cynomolgus C5 in order to enrich for cynomolgus cross-reactive antibodies.

For some panning conditions washing stringency was increased and antigen concentration was decreased within the three round of selection in order to generate high affinity antibodies.

The HuCAL GOLD[®] phagemid library was used to select specific Fab antibody fragments against human C5. First strategy was a solid phase panning on directly coated human C5 protein (panning procedure described above).

After the the 3rd panning round, the enriched phage pools were subcloned from the pMORPH[®]23 library vector (allowing efficient antibody display on the phage surface) into the pMORPH[®]x9_Fab_MH expression vector which mediates periplasmic expression of soluble Fabs. Single clones were picked and soluble Fabs were expressed from these single clones.

In total, 6624 clones were analyzed in primary screening which was performed by binding of the Fabs directly from the bacterial lysates to human C5 immobilized on Maxisorp microtiter plates. 1660 hits were obtained from the primary screening on human C5 with signals >5-fold over background. 384 hits were further analyzed in a secondary screening to confirm binding on human C5 and to screen for binding to the counter targets human C3 and C4.

Many primary hits could be confirmed on human C5 and showed no cross-reactivity to human C3 and C4, but only 6 Fabs had weak cross-reactivity to cynomolgus C5.

As a first consequence new solid phase pannings were performed alternating on human and cynomolgus C5. In parallel, quality controls of the purified cynomolgus C5 batch revealed a high amount of cynomolgus C3 within the cynomolgus C5 batch. Considering this results, a new method to screen for cynomolgus cross-reactive antibodies was applied. Cynomolgus C5 was captured from cynomolgus serum using an C5-binding polyclonal antibody (see Example 3, section 3). Using this method the initial primary hits were screened again on cynomolgus C5 and 56 clones were confirmed for binding to cynomolgus C5.

For the alternating solid phase pannings, the 1st round output of the most successful 12 human solid phase pannings was used for selections on cynomolgus C5 (protein batch contaminated with cynomolgus C3; not known during pannings). 376 clones were confirmed in a secondary screening for binding to human C5 and 361 clones for binding to cynomolgus C5 captured from cynomolgus serum.

Solution Panning on Biotinylated C5 Protein

The second panning variant was solution panning against biologically active (in hemolytic assays) biotinylated human C5 and biotinylated cynomolgus C5.

For this panning 200µl of Streptavidin magnetic beads (Dynabeads M-280; Dynal) were washed once with PBS and blocked with Chemiblocker for 2h at RT. 300µl of the PBS diluted phage were blocked also with Chemiblocker for 1-2h at RT on a rotator. The blocked phages were twice pre-adsorbed against 50µl blocked Streptavidin magnetic beads for 30min. The phage supernatant was transferred to a new blocked 2ml reaction tube and human biotinylated C5 was added and incubated for 1h at RT on a rotator. 100µl of the blocked Streptavidin magnetic beads were added to each panning pool and incubated for 10min on a rotator. The beads were collected with a particle separator (Dynal MPC-E) for approx. 2.5min and the solution was removed carefully.

Beads were then washed 7x in PBS/0.05% Tween using a rotator, followed by washing another three times with PBS. Elution of phage from the Dynabeads was performed by adding 200µl 20mM DTT in 10mM Tris/HCl pH 8 to each tube and incubation for 10min. Dynabeads were removed by the magnetic particle separator and the supernatant was added to 15ml of an E.coli TG-1 culture grown to OD_{600nm} of 0.6-0.8. Beads were then washed once with 200µl PBS and together with additionally removed phages the PBS was added to the 15ml E.coli TG-1 culture. For phage infection, the culture was incubated in 50ml plastic tubes for 45min at 37°C without shaking. After centrifugation for 5min at 4120 x g, the bacterial pellets were resuspended each in 600µl 2xYT medium, plated on 3xYT-CG agar plates and incubated overnight at 37 °C. Colonies were scraped off the plates and phages were rescued and amplified as described above. The second and third rounds of selection were performed in an identical way to the first round of selection.

A further panning strategy was solution panning using human C5 and alternating human and cynomolgus C5 (protein batch contaminated with cynomolgus C3, not known during pannings). Therefore the proteins were biotinylated and the retained bio-functionality after the biotinylation procedure was confirmed in hemolytic bioassays.

The phage-antigen complex was captured on Streptavidin magnetic beads via the biotin moiety of the antigen. After washing only specific bound phage were eluted (panning procedure described above).

First screening was done on directly coated proteins (see Example 3, section 1) and only 80 clones could be confirmed on human C5. Due to the fact that during the pannings the antigen was kept in solution, a new screening method was developed. In a solution ELISA the Fabs were incubated with biotinylated antigen on a NeutrAvidin plate. Using this solution screening procedure, a significantly higher amount of human and cynomolgus C5 specific clones could be selected. These results confirmed that many Fabs derived from solution pannings recognize C5 only in solution or when captured (e.g. via a polyclonal C5-binding antibody).

2. Subcloning and Expression of Selected Fab Fragments

To facilitate rapid expression of soluble Fabs, the Fab-encoding inserts of the selected HuCAL GOLD[®] phages were subcloned via XbaI and EcoRI into the E. coli expression vector pMORPH[®]x9_MH. Fab fragments carry a C-terminal Myc tag and as a second C-terminal tag the 6x His-tag (Chen *et al.*, Gene 139:73-75 (1994)). After transformation of the expression plasmids into E. coli TG1 F- cells chloramphenicol-resistant single clones were picked into the wells of a sterile 384-well microtiter plate pre-filled with 60µl 2xYT-CG medium and grown o/n at 30°C. 5µl of each E.coli TG-1 o/n culture were transferred to a fresh, sterile 96-well microtiter plate pre-filled with 40µl 2xYT medium supplemented with 34µg/ml chloramphenicol per well. The microtiter plates were incubated at 30°C shaking at 400rpm on a microplate shaker until the cultures were slightly turbid (~2-4 h) with an OD600nm of ~0.5. To these expression plates, 10µl 2xYT medium supplemented with 34µg/ml chloramphenicol and 3mM IPTG (isopropyl-β-D-thiogalactopyranoside) was added per well (end concentration 0.5mM IPTG). The microtiter plates were sealed with a gas-permeable tape, and incubated o/n at 30°C shaking at 400rpm. To each well of the expression plates, 15µl BEL buffer was added containing 2.5mg/ml lysozyme, 4mM EDTA and 10U/µl Benzonase and incubated for 1h at 22°C on a microtiter plate shaker (400rpm) followed by an optional freezing step for at least 2h at -80°C. The BEL extracts were used for binding analysis by ELISA or Fab SET screening after affinity maturation.

Expression of Fab fragments encoded by pMORPH[®]x9_Fab_MH in TG-1 cells was carried out in shaker flask cultures using 750ml of 2xYT medium supplemented with 34µg/ml chloramphenicol. Cultures were shaken at 30°C until the OD600nm reached 0.5. Expression was induced by addition of 0.75mM IPTG for 20h at 30°C. Cells were disrupted using lysozyme and Fab fragments isolated by Ni-NTA chromatography (Qiagen, Hilden, Germany). Buffer exchange to 1 x Dulbecco's PBS (pH 7.2) was performed using PD10 columns. Samples were filtered sterile (0.2µm, Millipore). Purity of samples was determined in denatured, reduced state by SDS-PAGE (15% Criterion Gels, BioRad) and in native state by size exclusion chromatography (HP-SEC). Protein concentrations were determined by UV-spectrophotometry (Krebs *et al.*, J. Immunol. Methods 254:67-84 (2001)).

On Fab level, the overall expression rates and the percentage of monomeric fraction in SEC (Size Exclusion Chromatography) ranged from acceptable to good for most of the identified antibody fragments. 64 parental Fabs were expressed and 61 Fabs could be purified. 60 affinity matured Fabs were purified in the mg scale. Most of the Fabs were good expressors and had no aggregation tendency.

Example 3: Identification of C5-specific antibodies from the HuCAL GOLD[®] Library

Below four different Enzyme Linked Immunosorbent Assay (ELISA) methods describe the screening of C5-binding antibodies (as bacterial BEL lysates or purified Fabs) on specific and counter antigens.

1. Screening on Directly Coated protein

Maxisorp (Nunc, Rochester, NY, USA) 384 well plates were coated with 20µl per well of 2.5µg/ml antigen (human C5 and the counter proteins human C3 and C4) in PBS, pH 7.4 o/n at 4°C. In parallel, plates were coated with 20µl per well of 5µg/ml sheep anti-human IgG, Fd fragment specific (The Binding Site, Birmingham, UK), diluted in PBS, pH 7.4 to check for Fab expression level.

The plates were blocked with PBS/ 0.05% Tween 20 (PBST) containing 5% milk powder for 1-2h at RT. After washing the wells with PBST, BEL-extracts, purified HuCAL GOLD® Fabs or control Fabs diluted in PBS were added and incubated for 1 h at RT. To detect Fab binding, anti-HIS6 antibody coupled to peroxidase was applied (Roche).

For detection of POD-conjugates fluorogenic substrate QuantaBlu (Pierce) was used according to manufacturer's instructions. Between all incubation steps, the wells of the microtiter plates were washed three times and five times with PBST after the final incubation with the secondary antibody. Fluorescence was measured in a Tecan GENios Pro plate reader.

2. Solution Screening with Biotinylated Proteins

The ELISA method described below was used for screening of HuCAL GOLD® Fabs after solution panning using biotinylated complement proteins.

NeutrAvidin plates were blocked with 1x Chemiblocker (Chemicon) diluted in PBS o/n at 4°C. These plates were used to screen for binding to human C5 and to the counter targets C3 and C4. In parallel, Maxisorp 384 well plates (Nunc, Rochester, NY, USA) were coated with 20µl per well of 5 µg/ml sheep anti-human IgG, Fd fragment specific (The Binding Site, Birmingham, UK), diluted in PBS, pH 7.4. These plates were used to check for Fab expression levels and for non-specific biotin binding. On the next day, coated Maxisorp plates were washed 2x with PBST and blocked with 3% BSA in TBS for 1-2h at RT. Periplasmic BEL extracts containing Fabs or purified HuCAL GOLD® Fabs were added to both blocked NeutrAvidin and Maxisorp plates.

Subsequently, 20µl per well of biotinylated human C5 (to detect specific binding) and in parallel, biotinylated human C3 and C4 (to detect unwanted binding) were added to wells of the NeutrAvidin plates. The biotinylated antigens were incubated with the HuCAL GOLD® Fabs for 1-2h at RT. Biotinylated unrelated antigen Transferrin was then added to the

21489-11410

Maxisorp plates to check for biotin binding Fabs (in this case the HuCAL®-Fab fragments were previously captured via anti-Fd antibody).

Following secondary antibodies were applied for detection: Alkaline phosphatase (AP)-conjugated Streptavidin-AP AffiniPure F(ab')₂ fragment, goat anti-human, was added to the Maxisorp expression plates; anti-HIS6 Peroxidase conjugated mouse antibody, Roche, was added to the NeutrAvidin plates and Streptavidin-Alkaline Phosphatase, ZYMED, was added to the Maxisorp plates with the biotinylated Transferrin.

For detection of AP-conjugates, fluorogenic substrate AttoPhos (Roche Diagnostics, Mannheim, Germany) and for detection of POD-conjugates, fluorogenic substrate QuantaBlu (Pierce) were used according to manufacturer's instructions. Fluorescence was measured in a Tecan GENios Pro plate reader.

Using this method it was possible to screen for anti-human C5 Fabs which recognize human C5 in solution and to exclude antibodies binding to the biotin moiety of the target antigens.

3. Determination of Cross-reactivity to Cynomolgus C5

A polyclonal C5-binding antibody (US Biological Cat#C7850-24) was used to capture cynomolgus C5 from cynomolgus serum.

384 well Maxisorp plates were coated with 20µl/well of 5µg/ml polyclonal C5-binding in PBS and incubated o/n at 4°C. On the next day the plates were washed 3x with PBST and blocked with 100µl/well of diluent (4% BSA/ 0.1% Tween20/ 0.1% Triton-X 100TM/ PBS) for 2 hours at RT. Cynomolgus serum was diluted 1:20 in diluent (4% BSA/ 0.1% Tween20/ 0.1% Triton-X 100/ PBS) (~approx. concentration of cynomolgus C5 4µg/ml) and 20µl/well was added to the 2x PBST washed Maxisorp plates. After 1h incubation at RT the plates were washed 3x PBST and BEL lysates containing Fab fragments or purified Fabs were added and incubated for 1h at RT. The plates were washed again and detection antibody anti-HIS6-POD (Roche #1965085), was added. POD substrate, BM Blue, soluble, (Roche Applied Science) was added and the reaction was stopped with 1M H₂SO₄. Absorbance was read at 450 nm using the BMG Reader device.

Example 4: Affinity Maturation

1. Construction of Affinity Maturation Libraries of Selected C5-Binding Fabs

To increase affinity and biological activity of selected antibody fragments, L-CDR3 and H-CDR2 regions were optimized in parallel by cassette mutagenesis using trinucleotide directed mutagenesis (see e.g., Virnekas *et al.*, Nucleic Acids Res. 22:5600-5607 (1994)), while the framework regions were kept constant. Prior to cloning for affinity maturation, all parental Fab fragments were transferred from the corresponding expression vector

(pMORPH[®]x9_MH) into the CysDisplayTM vector pMORPH[®]25 via XbaI/EcoRI. pMORPH[®]25 was created from the HuCAL GOLD[®] display vector pMORPH[®]23 by removal of one BssHII site interfering with library cloning for H-CDR2 optimization. For optimizing L-CDR3 of parental Fabs, the L-CDR3, framework 4 and the constant region of the light chains (405 bp) of the binders were removed by BpiI/SphI and replaced by a repertoire of diversified L-CDR3s together with framework 4 and the constant domain.

10 parental C5-binding Fabs were divided in 7 pools according to different selection criteria and only Fabs with same framework were put together: (1) MOR07086; (2) MOR06525 + 6756 (same framework); (3) MOR06757; (4) MOR06763; (5) MOR07087; (6) MOR07091 + 7092 (same framework); (7) MOR07093 + 7094 (same framework).

Approximately 1.5µg of the single Fab vector fragment and of the Fab pool were ligated with a 3 to 5-fold molar excess of the insert fragment carrying the diversified L-CDR3s. In a second library set, the H-CDR2 (XhoI/BssHII) was diversified while the connecting framework regions were kept constant. In order to monitor the cloning efficiency, the parental H-CDR2 was replaced by a dummy before the diversified H-CDR2 cassette was cloned in.

Ligation mixtures of the different libraries were electroporated into E.coli TOP10 F' cells (Invitrogen) yielding from 2×10^7 to 2×10^8 independent colonies. The libraries were amplified. For quality control, several single clones per library were randomly picked and sequenced using primers CFR84 (VL) and OCAL_Seq_Hp (VH).

As described above, seven maturation sub pools were generated and kept separate during the subsequent selection process.

14 different affinity maturation libraries (one LCDR3 and one HCDR3 library for each lead or pool) were generated by standard cloning procedures and transformation of the diversified clones into electro-competent E. coli TOP10F' cells (Invitrogen). Library sizes were good, being in the range of 2×10^7 – 5×10^8 . Sequencing of randomly picked clones showed a diversity of 100%. No parental binders but derivatives of all respective parental input binders were found. Finally phages of all 14 libraries were prepared separately.

Table 2. Overview of maturation libraries

MOR0	Maturation	VH/VL Type	Library Size
6757	HCDR2	VH3	3.70x10E7
6763	HCDR2	VH3	4.95x10E7
7086	HCDR2	VH1A	1.58x10E8
7087	HCDR2	VH1A	7.85x10E7
6525 + 6756	HCDR2	VH5	5.22x10E7
7091+ 7092	HCDR2	VH5	3.51x10E7
7093 + 7094	HCDR2	VH2	2.01x10E7
6757	LCDR3	Vkappa1	1.89x10E7
6763	LCDR3	Vlambda2	7.35x10E7
7086	LCDR3	Vlambda3	7.54x10E7
7087	LCDR3	Vkappa1	5.46x10E7
6525 + 6756	LCDR3	Vlambda2	8.50x10E7
7091+ 7092	LCDR3	Vlambda3	4.93x10E8
7093 + 7094	LCDR3	Vlambda2	1.33x10E8

2. Preparation of Antibody-Phages for Affinity Maturation

The HuCAL® maturation libraries were amplified in 2xYT medium containing 34µg/ml chloramphenicol and 1% glucose (2xYT-CG). After infection with VCSM13 helper phage at an OD_{600nm} of 0.5 (30min at 37°C without shaking; 30min at 37°C shaking at 250rpm), cells were spun down (4120 x g; 5min; 4°C), resuspended in 2xYT/ 34µg/ml chloramphenicol/ 50µg/ml kanamycin/ 0.25mM IPTG and grown o/n at 22°C. Phages were PEG-precipitated twice from the supernatant, resuspended in PBS and used for the maturation pannings described below.

3. Standard Solution Maturation Panning on Biotinylated C5 Protein

About 10¹² phages rescued from the generated affinity maturation libraries, as described above, were subjected to pannings performed under very stringent conditions to select for affinity improved C5 specific Fabs.

Solution pannings using the respective phage pools were either performed using biotinylated human C5 or alternating biotinylated human and cynomolgus C5 proteins. In order to increase panning stringency and to select for improved off-rates, antigen concentration was decreased and prolonged washing periods were applied (washing conditions are listed in Table 3).

Table 3. Increased washing conditions within the selection rounds of solution maturation pannings

Selection Rd.	Washing conditions (modified: stringent)
1st round	4x PBS/ 0.05% Tween 5min on rotator 3x PBS/ 0.05% Tween 15min on rotator-> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube 4x PBS quick 3x PBS 5min on rotator-> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube

2nd round	• 3x PBS/ 0.05% Tween quick • 7x PBS/ 0.05% Tween 15min on rotator-> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube • 3x PBS quick • 7x PBS 15min on rotator-> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube
3rd round	• 5x PBS/ 0.05% Tween quick • 8x PBS/ 0.05% Tween 15min on rotator • 1x PBS/ 0.05% Tween on on rotator • 3x PBS/ 0.05% Tween quick • 6x PBS/ 0.05% Tween 15min on rotator -> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube • 5x PBS quick • 8x PBS 15min on rotator -> transfer magnetic beads with the captured antigen and phages to a fresh blocked tube

Pre-blocked phage (1:2 mixture with 2x Chemiblocker incubated for 1h at RT) were incubated with low concentration of biotinylated C5 protein for 1-2h at RT. The panning strategy is similar to a standard solution panning described above. The phage antigen complex was captured via the biotin moiety of C5 to pre-blocked Streptavidin magnetic beads 30min at RT. Beads were then washed more stringently compared to a normal panning. Elution and amplification of phage was performed as described above.

The second and third rounds of selection were performed in an identical way to the first round, but at higher stringency washing conditions and lower antigen concentrations. For each antibody lead or pool several different pannings were performed. For each panning strategy different stringency conditions were applied. Panning strategies are summarized in Table 4.

Table 4. Overview of solution maturation pannings 1783 and 1784 on biotinylated human C5 and biotinylated cynomolgus C5

Panning #	Library	Panning mode	Antigen 1st round	Antigen 2nd round	Antigen 3rd round	Antigen Conc.	Washing
1783.1	MOR06525 + 6756 HCDR2	solution Streptavidin beads	human C5	human C5	human C5	50nM human / 5nM human / 0.25nM human	modified (more stringent)
1783.2	MOR07086 HCDR2						
1783.3	MOR06763 HCDR2						
1783.4	MOR07087 HCDR2						
1783.5	MOR06525 + 6756 LCDR3						
1783.6	MOR07086 LCDR3						
1783.7	MOR06763 LCDR3						
1783.8	MOR07087 LCDR3						
1783.9	MOR06525 + 6756 HCDR2	solution Streptavidin beads	human C5	cyno C5	human C5	25nM human / 5nM cyno / 0.25nM human	modified (more stringent)
1783.10	MOR07086 HCDR2						
1783.11	MOR06763 HCDR2						
1783.12	MOR06525 + 6756 LCDR3						
1783.13	MOR07086 LCDR3						
1783.14	MOR06763 LCDR3						
1784.1	MOR06757 HCDR2	solution Streptavidin beads	human C5	human C5	human C5	50nM human / 5nM human / 0.25nM human	modified (more stringent)
1784.2	MOR07091 + 7092 HCDR2						
1784.3	MOR07093 + 7094 HCDR2						

1784.4	MOR06757 LCDR3						
1784.5	MOR07091 + 7092 LCDR3						
1784.6	MOR07093 + 7094 LCDR3						
1784.7	MOR06757 HCDR2	solution Streptavidin beads	human C5	cyno C5	human C5	25nM human / 5nM cyno / 0.25nM human	modified (more stringent)
1784.8	MOR07091 + 7092 HCDR2						
1784.9	MOR07093 + 7094 HCDR2						
1784.10	MOR07087 HCDR2						
1784.11	MOR06757 LCDR3						
1784.12	MOR07091 + 7092 LCDR3						
1784.13	MOR07093 + 7094 LCDR3						
1784.14	MOR07087 LCDR3						

After maturation pannings, the enriched phagemid pools were sub-cloned into pMORPH[®]x9_MH expression vector.

4. Cross-Combination of Optimized VL (L-CDR3) with Optimized VH (H-CDR2)

For further improvement of affinity and potency, the independently optimized heavy and light chains from matured antibodies, derived from the same parental clone, were combined (see *e.g.*, Rauchenberger *et al.*, J. Biol. Chem. 278:38194-38205 (2003); Chen *et al.*, J. Mol. Biol. 293:865-881 (1999); and Schier *et al.*, J. Mol. Biol. 263:551-567 (1996)). This procedure, called cross-cloning, was applied for binders deriving from the same parental clones.

5. Affinity Screening and Maturation Panning Outcome

A total of 2640 clones derived from all pannings were screened as bacterial lysates for improved affinities on human C5. Preliminary affinities were estimated by solution equilibrium titration (SET). Based on their estimated affinities, clones derived from each parental Fab or Fab pools were sequenced. Table 5 shows number of sequenced clones and number of obtained unique sequences for each panning condition.

Table 5. Overview of affinity improved clones selected for sequence analysis

Parental/Maturation	Antigen	Sequenced clones	Unique Sequences	Parental of unique
MOR06525 + 6756 HCDR2	hu/hu/hu	10	9	6525
MOR07086 HCDR2	hu/hu/hu	10	4	7086
MOR06763 HCDR2	hu/hu/hu	22	10	6763(8x), 7086(2x)
MOR07087 HCDR2	hu/hu/hu	10	4	7087
MOR06757 HCDR2	hu/hu/hu	10	0	
MOR07091 + 7092 HCDR2	hu/hu/hu	24	7	7092
MOR07093 + 7094 HCDR2	hu/hu/hu	10	10	7093
MOR06525 + 6756 LCDR3	hu/hu/hu	20	5	6756
MOR07086 LCDR3	hu/hu/hu	10	5	7086
MOR06763 LCDR3	hu/hu/hu	10	8	7086
MOR07087 LCDR3	hu/hu/hu	6	1	7086
MOR06757 LCDR3	hu/hu/hu	16	0	
MOR07091 + 7092 LCDR3	hu/hu/hu	6	6	7091(1x), 7092(5x)
MOR07093 + 7094 LCDR3	hu/hu/hu	10	9	7094
MOR06525 + 6756 HCDR2	hu/cyno/hu	10	8	6525
MOR07086 HCDR2	hu/cyno/hu	10	6	7086
MOR06763 HCDR2	hu/cyno/hu	22	5	6763
MOR06757 HCDR2	hu/cyno/hu	15	2	6757
MOR07091 + 7092 HCDR2	hu/cyno/hu	15	6	7091(3x), 7092(3x)
MOR07093 + 7094 HCDR2	hu/cyno/hu	10	10	7093
MOR07087 HCDR2	hu/cyno/hu	10	6	7087(5x), 7086(1x)
MOR06525 + 6756 LCDR3	hu/cyno/hu	12	0	
MOR07086 LCDR3	hu/cyno/hu	10	1	7086
MOR06763 LCDR3	hu/cyno/hu	10	0	
MOR06757 LCDR3	hu/cyno/hu	9	1	7094
MOR07091 + 7092 LCDR3	hu/cyno/hu	11	9	7091(6x), 7092(3x)
MOR07093 + 7094 LCDR3	hu/cyno/hu	10	7	7094
MOR07087 LCDR3	hu/cyno/hu	10	0	
Sum		338	139	

6. Sequence Analysis and Selection of Affinity Optimized Fabs for Protein Production

A very good diversity was maintained by recovering derivatives of all 10 parental Fabs. The nucleotide sequences of the heavy chain (VH) for 188 HCDR2 improved clones and the light chain (VL) variable regions for 150 improved LCDR3 clones were determined. 87 unique HCDR2 and 52 unique LCDR3 sequences were selected for a detailed analysis of sequence diversity within the matured CDRs. Fabs containing possible glycosylations sites in the CDRs were omitted from further characterizations.

The VH and VL sequence analysis and affinity data showed that all 10 parental Fabs yielded affinity-improved successors. Parental Fabs MOR06525, MOR06757, MOR06763, MOR07087 and MOR07094 yielded only HCDR2 improved clones and parentals MOR06756 and MOR07093 yielded only LCDR3 improved clones. MOR07086, MOR07091 and MOR07092 had matured clones for both VH and VL. This later allowed cross-cloning of VH and VL matured chains. From all data, 60 clones with best affinity and highest diversity in the matured CDRs were selected for Fab expression. Selected VH and VL amino acid, as well as nucleotide sequences, are listed in Table 1.

Example 5: IgG Conversion

1. Conversion into Human IgG2 Format

In order to express full length immunoglobulin (Ig), variable domain fragments of heavy (VH) and light chains (VL) were subcloned from the pMORPH[®]x9_MH Fab expression vectors into pMORPH[®]2_h_Ig vector series for human IgG2. Restriction enzymes MfeI, and BlnI were used for subcloning of the VH domain fragment into pMORPH[®]2_h_IgG2. Subcloning of the VL domain fragment into pMORPH[®]2_h_Igk was performed via the EcoRV and BsiWI sites, whereas subcloning into pMORPH[®]2_h_Igλ2 was done using EcoRV and HpaI.

All ten parental Fabs (MOR06525, 6756, 6757, 6763, 7086, 7087, MOR07091, 7092, 7093 and 7094) were converted into human IgG2. The IgGs were also expressed.

2. Conversion into Human IgG1AA Format

In order to express full length immunoglobulin, variable domain fragments of Fab heavy (VH) and light chains (VL) were subcloned from the Fab expression vectors into IgG1 expression vectors. Restriction enzymes MfeI, and BlnI were used for subcloning of the VH domain fragment into pMORPH[®]2_h_IgG1AA, in which leucines at positions 234 and 235 were mutated to alanines to abrogate FcRγ binding and attenuate effector functions. The restriction enzymes EcoRV and HpaI were used to subclone of the VL domain fragment into pMORPH[®]2_h_Igλ2.

Following matured Fabs with desired profile were subcloned into human IgG1AA format: MOR07832, 7834, 7872, 7876, 7829, 7871, 7865, 7873, 7830, 7878, 7910. Cross-cloning on IgG level was achieved by transfecting cells with combinations of light and heavy chain constructs. For example, MOR08114 was the product of the germlined heavy chain from MOR07829 and the germlined light chain from MOR07871. Table 6 summarizes the most relevant cross-cloned germlined IgGs.

Table 6. Overview of most relevant cross-cloned germlined IgGs

MOR0 Nr.	VH/VL germlined	VH/VL		matured CDRs		format
		VH	VL	matured VH	matured VL	
8114	yes	7829	7871	7091/HCDR2	7091/LCDR3	hulg1AA
8125	yes	7091	7873	-	7091/LCDR3	hulg1AA
8126	yes	7829	7873	7091/HCDR2	7091/LCDR3	hulg1AA
8127	yes	7830	7873	7091/HCDR2	7091/LCDR3	hulg1AA
8128	yes	7092	7878	-	7092/LCDR3	hulg1AA
8129	yes	7909	7092	7092/HCDR2	-	hulg1AA
8130	yes	7909	7878	7092/HCDR2	7092/LCDR3	hulg1AA
8131	yes	7910	7092	7092/HCDR2	-	hulg1AA
8132	yes	7910	7878	7092/HCDR2	7092/LCDR3	hulg1AA

3. Transient Expression and Purification of Human IgG

Eukaryotic HKB11 and HEK293 cells were transfected with an equimolar ratio of IgG heavy and light chain expression vector DNA. Cell culture supernatant was harvested at 3 or

7 days post transfection and subjected to standard protein A affinity chromatography (rProteinA FF or MabSelect SURE, GE Healthcare). As not otherwise stated, buffer exchange was performed to 1 x Dulbecco's PBS (pH 7.2, Invitrogen) and samples were sterile filtered (0.2µm). Purity of IgG was analyzed under denaturing, reducing and non-reducing conditions in SDS-PAGE or by using Agilent BioAnalyzer and in native state by HP-SEC.

Example 6: Germlining

IgG constructs were germlined via site-directed mutagenesis using QuickChange® Site-Directed Mutagenesis Kit (Stratagene). The N-terminal DI of MOR08111 Vλ2 were changed to ES to match human germline sequence as well as to avoid a terminal Q (N-terminal Q can form pyroglutamine). N-terminal DI of MOR08110 Vλ3, MOR08113 Vλ3, and MOR08114 Vλ3 were germlined to SY, the most commonly found sequence in human λ3 genes. N-terminal QVQ of MOR08111 VH2 was germlined to EVT to match a λ2 gene and avoid terminal Q. N-terminal Q in MOR08109 VH5, MOR08110 VH5, MOR08113 VH5 and MOR08114 VH5 was also mutated to E.

Framework sequences for MOR08109 Vλ3 were synthesized to match the human λ3j gene and cloned into the expression vector using NheI and HpaI restriction sites. Sequence alignments of the antibodies variable domains with their respective closest related human germline sequences are shown in FIG. 1.

Example 7: Affinity Determination

1. Kon/Koff and K_D Determination of anti-human C5 antibodies using Surface Plasmon Resonance (Biacore)

It was determined that anti-Fab antibodies used to immobilize Fabs to the Biacore chip were influencing differently the binding affinity of each Fab for human C5, thus making the comparison of the Fabs to each other difficult. Biacore analysis was performed on IgG antibodies.

A CM4 chip was coated with 50µg/ml goat anti-human Fc antibody (500-2000 RU) in 10mM acetate buffer, pH 4.5, using standard EDC-NHS amine coupling chemistry. Each anti-human C5 IgG was captured on the chip in HBS-EP buffer at constant flow rate of 10µl/min for a contact time leading to a ligand density around 20 RU. After capturing the anti-hu C5 IgG, different concentrations of human or cynomolgus C5, in the range between 0.156nM to 2.5nM, were injected. Each cycle was completed with two regeneration steps with phosphoric acid. All running conditions were carried out at 25°C in 1x HBS-EP buffer. The resulting signals were adjusted by double referencing, subtracting the refraction index values from the reference flow cell and the binding step with no analyte. Data were collected

at 10Hz and analyzed using the Biacore T100 Evaluation Software Version 1.1 (GE). This program uses a global fitting analysis method for the determination of rate and affinity constants for each interaction.

The specificity of the antibodies were measured. Preferably, the K_{on} and K_{off} values for binding to human and cynomolgus C5 are as follows: $K_{on} > 1 \times 10^5$, $K_{off} < 1 \times 10^{-4}$). These measurements were performed in Biacore for the germlined IgGs and resulting data are listed in Table 7.

Table 7. K_D , K_{on} and K_{off} values of the germlined IgGs determined in Biacore

antiC5 final IgG	C5 sample	k_a [1/Ms]	k_d [1/s]	K_D [pM]
MOR08109	huC5	2.13E+06	2.56E-05	12
	cynoC5	1.23E+06	4.49E-05	37
MOR08110	huC5	4.15E+06	4.69E-05	12
	cynoC5	1.81E+06	9.24E-05	60
MOR08111	huC5	1.00E+06	3.07E-05	31
	cynoC5	8.91E+05	1.28E-04	144
MOR08113	huC5	2.51E+06	6.77E-05	28
	cynoC5	1.53E+06	1.27E-04	83
MOR08114	huC5	2.09E+06	3.12E-05	15
	cynoC5	1.06E+06	3.13E-05	31
5G1.1	huC5	1.29E+06	7.22E-05	56

2. Determination of Picomolar Affinities Using Solution Equilibrium Titration (SET) for Purified Fabs or Fabs Bacterial Lysates (Meso Scale Discovery (MSD))

For K_D determination by solution equilibrium titration (SET), monomer fractions (at least 90% monomer content, analyzed by analytical SEC; Superdex75, Amersham Pharmacia) of Fab protein were used. Affinity determination in solution was basically performed as described in the literature (Friguet *et al.*, J. Immunol Methods 77:305-319 (1985)). In order to improve the sensitivity and accuracy of the SET method, it was transferred from classical ELISA to ECL based technology (Haenel *et al.*, Anal Biochem 339:182-184 (2005)).

1mg/ml goat-anti-human (Fab)₂ fragment specific antibodies (Dianova) were labelled with ECL Sulfo-TAGTM NHS-Ester (Meso Scale Discovery, Gaithersburg, MD, USA) according to manufacturer's instructions. Experiments were carried out in polypropylene microtiter plates and PBS pH 7.4 with 0.5% BSA and 0.02% Tween 20 as assay buffer. Unlabelled antigen was diluted in 2ⁿ series, starting with a concentration at least 10 times

higher than the K_D . Wells without antigen were used to determine B_{max} values; wells with neither antigen nor Fab were used to determine background. After addition of e.g. 10pM Fab (final concentration in 60µl final volume), the mixture was incubated over night at RT. The applied Fab concentration was similar to or below the expected K_D .

Streptavidin MSD plates were coated with 0.2µg/ml biotinylated human C5 (30µl/well) and blocked with 5% BSA in PBS. Subsequently the equilibrated samples were transferred to those plates (30µl per well) and incubated for 20min. After washing, 30µl/well of the ECL Sulfo-tag labeled detection antibody (goat anti-human (Fab)₂) in a final dilution of 1:1500 was added to the MSD plate and incubated for 30min on an Eppendorf shaker (700rpm).

After washing and adding 30µl/well MSD Read Buffer T with surfactant Electrochemiluminescence signals were detected using a Sector Imager 6000 (Meso Scale Discovery, Gaithersburg, MD, USA).

Data were evaluated with XLfit (IDBS) software applying customized fitting models. For data evaluation *i.e.* K_D determination of Fab molecules the following fit model was used (model of Abraham et al¹⁶, modified according to *et al.*, 2005¹⁵): $y = B_{max} - (B_{max} / (2 * c_{Fab}) * (x + c_{Fab} + K_D - \sqrt{(x + c_{Fab} + K_D)^2 - 4 * x * c_{Fab}}))$; c_{Fab} : applied Fab concentration; x : applied total soluble antigen concentration (binding sites); sqrt: square root. Using the assay conditions described above (monomeric) affinities for the affinity-optimized C5-binding Fabs were determined in solution.

Parental Fabs

In order to further characterize the C5-binding antibodies, affinity of the parental Fabs to human C5 was determined. Because characterization focus was on efficacy in hemolytic assays, affinity measurements were done only for the most relevant Fabs. For a reliable determination of monovalent affinities only Fab batches were used for measurements which showed $\geq 90\%$ monomeric fraction in a qualitative size exclusion chromatography.

Affinities of the 10 parental Fabs which entered affinity maturation are summarized in Table 8. Affinities ranged from 72 pM to 3.7 nM.

Table 8. Affinities of the 10 parental Fabs determined in SET

	SET	
MOR0 Number	KD [pM]	(n = 1)
6525	72	
6756	1521	
6757	1186	
6763	820	
7086	108	
7087	3793	
7091	324	
7092	229	
7093	576	
7094	1364	
3207 (negative control)	no binding	

Matured Fabs

Monovalent affinities of the purified Fabs to human C5 were measured in SET. Affinities were in the low pM range and best affinities were obtained for derivatives of MOR07086, 7091, 7092 and 7093. Subsequently affinity measurements of these derivatives to cynomolgus C5 showed affinities in the mid to low pM range.

The affinity maturation process was very successful resulting in a repertoire of binders with markedly improved affinity. Table 9 summarizes affinities to human and cynomolgus C5 of the best improved binders. Certain Fabs have K_D to human C5 ≤ 30 pM and to cynomolgus C5 ≤ 150 pM.

Table 9. Overview of affinities to human and cynomolgus C5 for the best affinity improved Fabs

MOR	Matured	SET hu C5 (n=1-2)	SET cyno C5 (n=1)	MOR	Matured	SET hu C5 (n=1-2)	SET cyno C5 (n=1)
		KD [pM]	KD [pM]			KD [pM]	KD [pM]
6525		273 / 29		7871	LCDR3	3	4
7813	HCDR2	437		7872	LCDR3	2	3
7814	HCDR2	137		7873	LCDR3	13 / 13	6
7816	HCDR2	116		7874	LCDR3	35	8
6757		3650 / 1245		7092		96	481
7818	HCDR2	491	70	7831	HCDR2	10	36
7907	HCDR2	179		7832	HCDR2	4	13
6763		673 / 962		7909	HCDR2	7	18
7820	HCDR2	62		7910	HCDR2	27	31
7086		12 / 65	10	7876	LCDR3	78	60
7821	HCDR2	7	39	7877	LCDR3	29	144
7822	HCDR2	5	14	7878	LCDR3	33	70
7823	HCDR2	5	15	7879	LCDR3	25	122
7824	HCDR2	55 / 130		7093		431 / 992	3146
7864	LCDR3	22	974	7833	HCDR2	47	107
7865	LCDR3	10	88	7834	HCDR2	4	15
7866	LCDR3	10	191	7835	HCDR2	29	28
7867	LCDR3	19	154	7836	HCDR2	11	
7868	LCDR3		384	7890	HCDR2	46	
7869	LCDR3	2	83	7094			
7870	LCDR3	12	500	7880	LCDR3	13	13
7087		120		7881	LCDR3	88	
7827	HCDR2	361		7882	LCDR3	70	
7828	HCDR2	2477 / 1730		7883	LCDR3	49	
7091		135 / 138	704	7884	LCDR3	83	
7829	HCDR2	429	116	7885	LCDR3	35	
7830	HCDR2	399	75	criterion: KD hu C5 <30pM, cy C5 <150pM			
7908	HCDR2	15	39	*scattering (no reliable measurement)			

3. K_D determination of IgG molecules using Solution Equilibrium Titration (SET)

Affinities of the germlined IgGs (human IgG1AA format) to human and cynomolgus C5 were determined in SET as described below. Similar data sets between two independent measurements showed higher affinities of the lead IgGs to human C5 than reference IgG 5G1.1 (see U.S. Pat. NO. 6,355,245). Final IgGs had affinities for human C5 ranging from 1 to 14pM and affinities to cynomolgus C5 ranging from 3 to 29pM.

Table 10. K_D values determination for the final lead IgGs (human IgG1AA format) in SET

		1 st measurement		2 nd measurement	
		human C5	cyno C5	human C5	cyno C5
		KD [pM]	KD [pM]	KD [pM]	KD [pM]
hu IgG1AA germlined	MOR08109	4	13	2	6
	MOR08110	7	18	3	8
	MOR08111	5	14	3	17
	MOR08113	14	29	8	16
	MOR08114	1	5	2	4
hu IgG2/4 (reference IgG)	5G1.1	24	no binding	19	no binding

For K_D determination by solution equilibrium titration (SET), monomer fractions of IgG protein were used (at least 90% monomer content, analyzed by analytical SEC MALS; Tosoh TSKgel G3000SWXL, Wyatt Treos miniDAWN). Affinity determination in solution was basically performed as described in the literature (Friguet *et al.*, J. Immunol Methods 77:305-

319 (1985)). In order to improve the sensitivity and accuracy of the SET method, it was transferred from classical ELISA to ECL based technology (Haenel *et al.*, Anal Biochem 339:182-184 (2005)).

1mg/ml goat-anti-human (Fab)₂ fragment specific antibodies (Dianova) were labelled with ECL Sulfo-TAGTM NHS-Ester (Meso Scale Discovery, Gaithersburg, MD, USA) according to the manufacturer's instructions. The experiments were carried out in polypropylene microtiter plates and PBS pH 7.4 with 0.5% BSA and 0.02% Tween 20 as assay buffer. Unlabeled antigen was diluted in 2n or 1.75n series, respectively, starting with a concentration at least 10 timer higher than the K_D. Wells without antigen were used to determine Bmax values; wells containing neither antigen nor IgG were used to determine background. After addition of e.g. 10pM IgG (final concentration in 60µl final volume), the mixture was incubated over night at RT. The applied IgG concentration was similar to or below the expected K_D.

Streptavidin MSD plates were coated with 0.2µg/ml biotinylated human C5 (30µl/well) and blocked with 5% BSA in PBS. Subsequently the equilibrated samples were transferred to those plates (30µl per well) and incubated for 20min. After washing, 30µl/well of the ECL Sulfo-tag labeled detection antibody (goat anti-human (Fab)₂) in a final dilution of 1:1500 was added to the MSD plate and incubated for 30min on an Eppendorf shaker (700 rpm).

Electrochemiluminescence signals were detected after washing and adding 30µl/well MSD Read Buffer T with surfactant using a Sector Imager 6000 (Meso Scale Discovery, Gaithersburg, MD, USA).

Data were evaluated with XLfit (IDBS) software applying customized fitting models. For data evaluation *i.e.* K_D determination of IgG molecules the following fit model for IgG was used (modified according to Piehler *et al.*, 199717): $y = B_{max} / ((x + c_{lgG} + K_D) / 2 - ((x + c_{lgG} + K_D)^2 / 4 - x * c_{lgG}^{0.5})^{0.5} / (2 * c_{lgG}^{0.5}))$; c_{lgG} = applied IgG concentration, complete molecule (not binding sites); x = applied total soluble antigen concentration (binding sites); sqrt : square root.

Example 8: Characterization by Hemolytic Assays

The hemolytic assay is a basic functional assay that tests for complement activation and has been used to evaluate the ability of anti-human C5 mAbs and Fab molecules to block lysis of red blood cells (RBCs) by complement pathways (see e.g., Evans *et al.*, Mol. Immunol 32: 1183-1195 (1995); Thomas *et al.*, Mol Immunol 33:1389-1401 (1996); Rinder *et al.*, J Clin Invest 96:1564-1572 (1995)). Briefly, for classical pathway assays, sensitized red blood cells are used as targets for lysis by complement proteins present in serum. This assay is of interest for the characterization and screening of high-affinity anti-human C5 mAbs.

1. Classical Pathway

The desired number of chicken red blood cells was washed four times with cold gelatin veronal buffer (GVB++) and resuspended to 5×10^7 cells/ml. To sensitize the cells rabbit anti-chRBC IgG was added to RBC cell suspension to a final concentration of 1 µg/ml IgG. After 15 minutes incubation on ice, the sensitized chRBCs were centrifuged, washed twice with GVB++ and diluted to 8.33×10^7 cells/ml.

Round-bottom 96 well plates were used for hemolytic assay. Antibodies were diluted in GVB++ buffer and added to the wells (when calculating the required concentration of C5-binding Abs, it was considered that the sample will be diluted two-fold when serum is added). 50 µl of 40% human serum (diluted in GVB++) was added to 50 µl antibody dilutions, resulting in a final serum assay concentration of 20%.

The control and blank wells were prepared as described here: control wells: i) 0% lysis control → 100 µl GVB++, ii) 100% lysis control → 100 µl 0.1% NP-40, iii) 20% serum control → 100 µl of 20% serum (0% Ab control). blank wells: i) 20% serum blank → 100 µl 20% serum, ii) GVB++ blank → 100 µl GVB++, iii) NP-40 blank → 100 µl 0.1% NP-40.

2.5×10^6 (30 µl) sensitized chRBCs/well were added to all sample and control wells. To the blank wells PBS was added instead of cells. Assay plate was incubated 30 min at 37°C, centrifuged (2,000 rpm, 5 min) and 85 µl supernatant was transferred to a new, flat-bottomed 96-well plate. The new plate was centrifuged (2,000 rpm, 3 min) to get rid of any bubbles. Hemoglobin release was measured by reading absorbance at 415 nm. Percentage of hemolysis was calculated with respect to the control and blank wells using the following calculation algorithms:

$$\%Hemolysis = 100 \times \frac{OD_{sample} - OD_{negativecontrol}}{OD_{positivecontrol} - OD_{negativecontrol}}$$

where

$$OD_{sample} = [AverageOD_{sample}] - [AverageOD_{20\%SerumBlank}]$$

$$OD_{negativecontrol} = [AverageOD_{0\%Lysis}] - [AverageOD_{GVB++Blank}]$$

$$OD_{positivecontrol} = [AverageOD_{100\%Lysis}] - [AverageOD_{NP-40Blank}]$$

Using this procedure, anti human-C5 antibodies which were able to inhibit red blood cell lysis could be identified. To screen for cross-reactivity to cynomolgus C5, the classical pathway was performed using 5% cynomolgus serum.

2. Alternative Pathway

Hemolytic assays undergoing the alternative pathway were done in a similar way to the classical pathway hemolytic assays. In the alternative pathway RBCs cells from rabbit were used and there was no need to sensitize the cells. The rabbit RBCs are different from chicken RBCs in that they are sensitive to lysis caused by the complement alternative pathway.

The working buffer was GVB++ supplemented with 10mM EGTA and 5mM Mg++, since the C5 convertase of the alternative pathway is Mg++ dependent and the C5 convertase of the classical pathway is Ca++ dependent.

Hemolytic assays of the alternative pathway were run with: i) 20% human serum, ii) 100pM human C5 added to 20% human C5-depleted serum, iii) 0.025% cynomolgus serum added to 20% human C5-depleted serum, iv) 100pM cynomolgus C5 added to 20% human C5-depleted serum, v) 10% cynomolgus serum. These settings were used to screen for antibodies with high affinity to the human and cynomolgus C5 proteins which were able to inhibit very effectively the red blood cell lysis induced by the alternative complement pathway.

3. Hemolytic Assays with Parental Fabs

Hemolytic assays were used as a basic bio-functional assay to evaluate the ability of anti-human C5 mAbs to block complement mediated lysis of red blood cells. C5 convertase cleaves C5 into C5a peptide and C5b fragment, that is subsequently incorporated into the membrane-attack complex (MAC), which leads to cell lysis. C5 convertase of the classical pathway, formed by a C3bC4bC2a complex has a different structure than the C5 convertase of the alternative pathway which is formed by a C3bC3bBb complex. HuCAL GOLD® generated antibodies should be inhibitory in both classical and alternative pathway, but with focus on the alternative pathway because mainly the alternative pathway (factor H, factor B and factor H-related genes) is implicated in AMD.

The classical and alternative pathway assays were performed with 20% human serum (~80nM C5). To increase sensitivity of alternative pathway assays, new assay formats were developed. 10-100pM purified human C5 or 0.025% cynomolgus serum (~100pM cynomolgus C5) were added to human C5-depleted serum (but containing all other serum and complement components).

FIG. 2 shows that considerable hemolysis could be observed between 10 and 100 pM purified human C5 added to human C5-depleted serum. Cynomolgus serum was added to human C5-depleted serum to test for cross-reactivity. FIG. 3 shows that 0.025% of cynomolgus serum (~100pM C5) added to human C5 depleted serum restores hemolytic activity.

Classical Pathway

First Fab selection was done in the classical pathway (20% human serum). Approximately half of the 61 purified parental Fabs were weak to strong inhibitors of the classical pathway. IC₅₀ values of the best inhibitory Fabs were between 35 and 900nM.

Assays were done showing congruent results (as shown in FIG. 4). % hemolysis was calculated with respect to the control and blank wells. Fab inhibition of cell lysis was compared to a maximum lysis caused by 20% human serum (=100%). An irrelevant human Fab (hen egg white lysozyme binder MOR03207) was used as negative control and anti-human C5 IgG monoclonal antibody (Quidel) as positive control. FIG. 4 show an example with the best inhibitory Fabs.

Alternative Pathway

Fabs which showed inhibitory activity in the classical pathway were further evaluated in the alternative pathway. Hemolytic assays were run with 100pM purified human C5 or 0.025% cynomolgus serum added to human C5-depleted serum. IC₅₀ values for the human alternative assays were between 0.1 and 90nM (examples of assays with the most relevant Fabs are shown in FIG. 5).

The positive control of the classical pathway (anti-human C5 antibody, Quidel) was not inhibitory in the alternative pathway. Therefore an anti-complement factor P antibody (Quidel) was used as positive control. As shown in FIG. 5, MOR07086 had best inhibitory activity and NVS data revealed a better potency than for the reference antibody 5G1.1.

To test for cynomolgus cross-reactivity, hemolytic assays of the alternative pathway were performed with 0.025% cynomolgus serum added to human C5-depleted serum. A comparison to 5G1.1 was not possible, since 5G1.1 does not recognize cynomolgus C5. The anti-Factor P antibody was used as positive control. Results of assays revealed IC₅₀ values between 0.1 and 400nM for the best inhibitory Fabs. Again, MOR07086 showed best potency (shown in FIG. 6).

A consistent inhibitory activity of the Fabs was noticed in both classical and alternative pathway. Table 11 below summarizes the results of hemolytic assays for the most relevant 22 Fabs. To have a reliable comparison between different experiments, lysis caused by 20% human serum was normalized to 100%.

Table 11. Summary of hemolytic assays with the most relevant Fabs

MOR-Nr	MOR IC50 [nM]			NVS IC50 [nM]		
	CP [human] normalized	AP (0.1nM C5) [human] normalized	AP (0.025% cyno serum) [cyno] normalized	CP [human] normalized	AP (0.1nM C5) [human] normalized	AP (0.025% cyno serum) [cyno] normalized
6525	190	15	11	185	7	5
6756	320	80	400	225	70	2500
6757	500	90	30	305	130	25
6763	250	45	110	195	20	360
6764	n.t.	50	n.t.	n.t.	25	30% inh
6776	>4000	40		n.t.	20*	50% inh
6952	90	20	>1000	110	15	200
6961	100	25	600	85	15	30
7081	180	5	40% inh	170	3	10
7082	70	2.5	1	90	1	1
7083	100	30	300	140	10	5
7084	120	10	1.2	160	5	1.5
7086	35	0.2 / 0.2	0.2 / 0.4	85	0.1	0.1
7087	>4000	50	100	775	10	1
7088	110	15	230	130	5	15
7089	150	75	900	250	20	50
7090	105	20	10	120	10	1
7091	82	7	40	110	3	4
7092	100	1	1.5	90	0.5	1.5
7093	>4000	7	190	230	5	15
7094				770	40	190
7095*	120*	0.5**	1.3**	n.t.		

*not pure as MH

**as pMx9_FS

4. Hemolytic Assays with Matured Fabs

Classical Pathway

(1) Classical Pathway Using 20% Human Serum

Matured Fabs were tested in the classical pathway with 20% human serum. Derivatives of MOR07086, 7091, 7092 and 7093 showed highest potency (IC50 values in the low nM range). Descendants of MOR07091, 7092 and 7093 showed strongly improved potency. FIG. 7 shows examples of hemolytic assays with derivatives of MOR07086, 7091, 7092 and 7093.

(2) Classical Pathway Using 5% Cynomolgus Serum

Assays of the complement pathway were also run in the presence of 5% cynomolgus serum in order to test for cross-reactivity. Derivatives of MOR07086, 7091, 7092 and 7093 could very effectively inhibit red blood cell lysis. The negative control, MOR03207 (anti-lysozyme Fab), had no impact on the complement pathway. Results of these assays are shown in FIG. 8.

Alternative Pathway

(1) Alternative Pathway Using 100pM Human C5

Matured Fabs were tested in the alternative pathway hemolytic assay with 100pM human C5. Some derivatives of MOR06525, 6757, 6763, and 7087 showed potency improvement compared to their parentals. MOR07086-, 7091-, 7092-, 7093-, and 7094-derived Fabs showed highest potency (IC50 values in the low nM range). Descendants of MOR07091, 7092, 7093, and 7094 showed highly improved potency, many of which are more potent than reference antibody 5G1.1. FIG. 9 shows examples of hemolytic assay results for the affinity matured Fabs and 5G1.1.

(2) Alternative Pathway Using 20% Human Serum

Matured Fabs were tested in the alternative pathway hemolytic assay with 20% human serum. MOR07086-, 7091-, 7092- and 7093-derived Fabs showed best inhibitory activity. Many of these Fabs had better inhibitory activity than 5G1.1. FIG. 10 shows examples of hemolytic assay results for the affinity matured Fabs and reference antibody 5G1.1.

(3) Alternative Pathway Using 100pM Cynomolgus C5

Matured Fabs were tested in the alternative pathway hemolytic assay using 100pM cynomolgus C5 added to 20% human C5-depleted serum. MOR07091-, 7092- and 7093-derived Fabs showed best inhibitory activity; 5G1.1 does not crossreact with cynomolgus C5. FIG. 11 shows examples of hemolytic assay results for the affinity matured Fabs.

5. Hemolytic Assays with Germlined IgGs (Human IgG1AA Format)

Classical Pathway

(1) Classical Pathway Using 20% Human Serum

Classical pathway assays using 20% human serum were run at MOR. IC50 values of the final germlined hu IgGAA - MOR08109, 8110, 8113, 8114 - were better or similar to reference IgG 5G1.1 (see FIG. 12).

(2) Classical Pathway Using 5% Cynomolgus Serum

A comparison to 5G1.1 in the classical pathway using 5% cynomolgus serum was not applicable, since this reference antibody does not recognize cynomolgus C5. The final germlined IgGs could completely inhibit lysis of the red blood cells induced by cynomolgus serum except MOR08111. Data are shown in FIG. 13.

Alternative Pathway

(1) Alternative Pathway Using 100pM Human C5

The germlined IgGs were tested in the alternative pathway hemolytic assay using 100pM human C5. All antibodies showed potent inhibitory activity with IC50 values between

28 and 128pM (with the exception of MOR08111, see FIG. 14), all were equal to or better than 5G1.1. FIG. 14 shows examples of hemolytic assay results for the IgGs.

(2) Alternative pathway Using 20% Human Serum and C5a Generation ELISA

The germlined IgGs were also tested in the alternative pathway hemolytic assay with 20% human serum. The majority of the antibodies tested achieve complete inhibition with IC50 values lower than 80nM. Reference antibody 5G1.1 does not fully inhibit hemolysis in this assay. FIG. 15 shows examples of hemolytic assay results for the IgGs. Inhibition of C5a generation by the final IgGs was similar to 5G1.1 (IC50 values in the low nM range).

(3) Alternative Pathway Using 100pM Cynomolgus C5

Hemolytic assays of the alternative pathway in 20% human C5-depleted serum were reconstituted with 100 pM cynomolgus C5. Potency of the germlined final candidates against cynomolgus C5 was within 5-fold of that for human C5 (IC50 values in the low pM range).

(4) Alternative Pathway Using 10% Cynomolgus Serum

In hemolytic assays of the alternative pathway using 10% cynomolgus serum ([C5] ~ 40 nM) the potency of the germlined candidates was similar to the potency in human serum (success criterion was to have a potency not more than 5-fold weaker than for the functional assay using human C5).

Example 9: C5a Generation ELISA

C5a-des-Arg ELISA was developed to measure C5a generation during hemolysis to confirm that antibodies that were inhibitory in the hemolytic assay also inhibited cleavage of C5 into C5a and C5b.

A Maxisorp plate was coated with 100µl/well mouse anti-human C5a-des-Arg (US Biologics) at 1µg/ml in coating buffer (bicarbonate pH 9.5-9.8) and was incubated overnight at 4°C. After washing 3x with PBST, the plate was blocked with 300µl/well diluent (Synblock, AbD Serotec) for 2 hours at room temperature. After aspirating the blocking solution, 100µl samples or standards diluted with diluent were incubated for 1 hour at room temperature. Standards were prepared as follows: start was at 20ng/ml standard (rC5a-des-Arg) and 1:4 serial dilutions were prepared for a 7-point curve. Samples of hemolytic assays were diluted 1:5 in diluent (hemolytic assay supernatants should be stored at -80°C until used in C5a ELISA). In between the plate was washed 3x with PBST. 100µl/well of 0.4µg/ml detection antibody (biotin-goat anti-human c5a, R&D Systems) diluted in diluent was added and after 1 hour incubation at room temperature, 100µl/well Strep-HRP (poly-HRP streptavidin) diluted 1:5000 in HRP diluent (poly-HRP diluent) was added for 30 minutes. After washing 4x with PBST, 100µl/well TMB Substrate (Ultra TMB substrate solution) was added for 5-10 minutes.

Reaction was stopped with 50µl/well stop solution (2N H₂SO₄). Absorbance was read (A450-A570) and data were analyzed using SoftMax Pro.

Matured Fabs were tested for C5a generation during hemolysis to confirm that inhibitory activity was due to blocking C5 cleavage into C5a and C5b. The supernatants from hemolytic assays in 20% human serum were used for quantifying the C5a formation.

All Fabs tested brought C5a levels down to baseline. FIG.16 shows examples of C5a ELISA results.

Example 10: Specificity ELISA on Human C3, C4, C5 and Cynomolgus C5

All purified Fabs were analyzed in a solution ELISA (method described above) for binding to human C3, C4 and C5. Fabs were incubated with biotinylated antigen on a Neutravidin plate and detected via the histidin tag.

Improved binding was seen for almost all matured Fabs compared to their respective parental. No binding to the counter targets human C4 and C3 was detected up to 100nM Fab. These results hit the success criteria for specificity: binding to human and cynomolgus C5 and no binding to human complement proteins C3 and C4. Examples for derivatives of parental Fab MOR07091 are shown in FIG. 17.

Example 11: Serum Stability Assays

Retained binding activity to human C5 in a binding assay at 50% human serum of C5-binding antibodies was determined as described below.

Antibodies (Fab format) were incubated up to 8h at 37°C with 100% human C5-depleted serum or with PBST/0.5% BSA (positive control). Wells of a blocked polypropylene plate were used for incubation to ensure no binding of the antibodies to the surface over the long incubation time. Samples were collected at different time points and stored at -20°C.

Samples were tested in a solution ELISA on NeutrAvidin plates to check binding ability to human C5. To the NeutrAvidin plates, which were blocked o/n with 1x ChemiBlocker-PBST. 20µl of serial dilutions of the different collected samples were added. First dilution of the samples was 1:2 (final serum concentration 50%), followed by 1:3 dilutions steps. After 1h incubation the plate was washed 3x with PBST and 20µl biotinylated human C5 was applied to a concentration of 2.5 µg/ml. After 1h plate was washed again 5x with PBST (0.05% Tween) and anti-HIS6-POD detection antibody for Fabs was added.

Fluorescence of the substrate (Quanta Blue or AttoPhos) was measured after 5-10 min and retained binding activity was calculated compared with the respective maximum signal (antibody incubated with PBST/0.5% BSA).

One of the “must” criteria for the C5-binding antibodies is to retain 75-80% of binding activity in human serum i) in a functional assay at 10% serum and ii) in a binding assay at 50% serum. Because hemolytic assays were run in the presence of 20% serum it was only necessary to show retained binding in a binding assay at 50% serum.

Therefore matured final Fabs were incubated with 100% human C5-depleted serum at 37°C for 8h. Samples were collected at different time points and tested for binding to human C5 in a solution ELISA. Fab + serum samples used for ELISA were diluted to a concentration of 50% serum + 10nM Fab.

FIG. 18 illustrates the results of the final C5-binding final antibodies in the Fab format. 70-93% of the binding activity was retained after an 8 hour incubation time at 37°C in 50% serum compared to incubation in PBS.

Example 12: Characterization by Epitope Binning

This procedure was used to group anti-human C5 Fabs into different epitope bins binding to the same or an overlapping epitope of the C5 protein.

Competition of each biotinylated anti-human C5 antibody with each unlabelled anti-human C5 antibody in 100-fold excess was tested in an ELISA (capture mode). It was compared with the highest signal of each antibody (biotinylated Fab without competition).

Human C5 was captured via a polyclonal anti-human C5 IgG (US Biological), which was coated previously o/n at 4°C on a 384 well black Maxisorp plates. Next day the plate was washed twice with PBST and blocked for 2h with 3% BSA-PBST. After washing 3x with PBST, 20µl human C5 was added and incubated 2h at RT. The plate was washed 3x with PBST before adding the Fabs.

20µl unlabelled Fab (200µg/ml or 400µg/ml) (100-fold excess) was added to the wells of a Maxisorp plate and subsequently 20ng/ml or 40ng/ml of biotinylated Fab. The biotinylated and unlabelled Fabs were incubated for 1h at RT. The plate was washed 3x with PBST and Strep-AP Zymax Streptavidin-Alkaline Phosphatase, ZYMED, Code: 43-8322, Lot: 50799648 was added for detection of the biotinylated Fab binding via C5 to the plates. AttoPhos substrate (Roche) was added to the plates and Fluorescence was read after 5-10min.

Parental Fabs

C5 was captured (via a polyclonal antibody) and unlabelled FabY was applied in excess to biotinylated FabX. Binding of biotinylated FabX to human C5 was detected. Six groups of Fabs could be defined: Group 1: MOR06952, 6961; Group 2: MOR06525, 6756, 6757, 6763; Group 3: MOR07087; Group 4: MOR06764, 6776, 7081; Group 5: MOR07089; Group 6: MOR07082, 7083, 7084, 7086, 7088, 7090, 7091, 7092, 7093, 7095.

The Fabs were also divided into different epitope binding groups using a different method: FabX was immobilized, then FabY pre-incubated with biotinylated C5 was added. Following groups of Fabs could be defined: Group 1: MOR06952, 6961; Group 2: MOR06525, 6757, 7083; Group 3: MOR07087; Group 4: MOR06763; Group 5: MOR07081; Group 6: MOR07082, 7083, 7084, 7086, 7088, 7091, 7092, 7093 (7089 competes with 7084). The conclusion was drawn that using two different methods, similar results could be obtained.

Matured Fabs

In order to complete Fab characterization competition of biotinylated Fab with unlabelled Fab (applied in 100-fold excess) was measured in solution ELISA. Results were compared with the highest signal (biotinylated Fab without competition).

As shown in FIG. 19, biotinylated Fabs compete with identical unlabelled Fabs and all Fabs compete for binding to the same or overlapping epitope. These results correlate with epitope binning data for the parental Fabs.

Example 13: Screening of C5 Alpha versus Beta Chain Binders and Competition Assays

Two ELISA experiments and hemolytic assays were performed to test if a Fab was an alpha or beta chain binder as described below.

In the first experiment, Fab was coated on a plate and purified C5 or supernatant from chimeric C5 preparation (human alpha, mouse beta chain) was added. As a next step 5G1.1 was applied and detection was done via an anti-human IgG.

In a second experiment, 5G1.1 was coated on a plate, purified C5 or supernatant from chimeric C5 preparation (human alpha, mouse beta chain) was added, then Fab, which was detected with an anti-Myc antibody.

Reference IgG 5G1.1 recognizes the alpha chain and was used to determine if the MorhpSys generated Fabs compete with 5G1.1 for binding. In the hemolytic assays supernatant from chimeric C5 preparation was added to human C5-depleted serum and Fabs were tested for inhibition of hemolysis.

Parental Fabs

FIG. 20 shows the results of an ELISA experiment where the Fabs were coated on a plate, C5 or supernatant of a chimeric C5 preparation (human alpha chain and mouse beta chain) was added, then 5G1.1. FIG. 21 shows the results of an ELISA experiment where purified C5 and supernatant from chimeric C5 were captured via 5G1.1.

MOR06525, 6756, 6763 were beta chain binders (bind to C5 but not chimeric C5). Most MOR070XX Fabs (derived from solution pannings) are alpha chain binders (bind to C5 and chimeric C5). MOR06952 and 6961 compete with 5G1.1 so they are negative for both C5 and chimeric C5 and, thus, are most likely alpha chain binders as 5G1.1. MOR06757 behaves like MOR06952 and 6961, *i.e.* it likely is an alpha chain binder. However, MOR06757 does not inhibit hemolysis of chimeric C5 supernatant spiked into C5-depleted serum, while all the other alpha chain binders do (see FIG. 22).

In the hemolytic assay supernatant from chimeric C5 prep was added to human C5-depleted serum and Fabs were tested for inhibition of hemolysis. MOR06525, 6756, 6757 and 6763 did not inhibit hemolysis with chimeric C5 and thus, could be beta chain binders. MOR06952, 6961, 7081, 7082, 7083, 7084, 7086, 7087, 7088, 7089, 7090, 7091, 7092, 7093, 7094, 7095 inhibited hemolysis and thus could be alpha chain binders.

Example 14: Resistance to Proteolysis

To investigate the structural rigidity of Fabs, resistance of Fabs to proteolysis by thermolysin was performed (thermolysin bacterial protease, Calbiochem). Fab was incubated with thermolysin (Fab : thermolysin = 3 : 1 (w/w), reaction volume of 8µL) either at 37°C or at 55°C (thermolysin activity is optimal at 55°C). The reaction was stopped by adding 4µL of 0.5 M EDTA and 4µL of 4x LDS sample buffer (Invitrogen) and the stopped samples were run on 4-12% SDS-PAGE at non-reducing condition. Proteolysis of Fabs was analyzed by monitoring the disappearance of Fab bands that were visualized by Coomassie staining.

Parental Fabs

Parental Fabs were tested for resistance to thermolysin proteolysis at 37°C and 55°C. Fab from a humanized IL-1β antibody was used as control. Most tested Fabs were resistant to degradation by thermolysin at 37°C up to 90 min. To further differentiate the structural rigidity of Fabs, proteolysis was performed at higher temperature of 55°C. Many of the Fabs tested were quickly degraded at 55°C (>90% Fab was degraded within 30 min), while some Fabs were still resistant to proteolysis after 90 min (*e.g.*, 7094). The resistant Fabs were suggested to have a more rigid structure such that they might show better *in vivo* pharmacokinetic properties. Results of these experiments are shown in the FIG. 23 and FIG. 24.

Matured Fabs

Fabs with the highest potency in hemolytic assays were tested for sensitivity to thermolysin at 37°C and 55°C. In FIG. 25 and FIG. 26, experiments with derivatives of MOR07086, 7091, 7092 and 7093 are shown.

Results of these tests revealed that derivatives of parentals MOR07091, 7092 and 7093 were less sensitive to proteolysis, while MOR07086 derivatives were more sensitive to proteolysis.

Example 15: MAC Deposition Assay

As the terminal complement cascade ends up with formation of the MAC, inhibition of MAC formation was a further hint for the antibody ability to block the complement cascade. The rational was to have an additional set-up independent of cells and cell behaviour

Zymosan (Sigma), which is an insoluble carbohydrate from the cell wall of yeast, used especially in the immunoassay of the alternative pathway, was coated to activate the Alternative Pathway and IgM (Sigma) was coated to activate the Classical Pathway for determination of MAC (membrane attack complex) deposition. Fabs were pre-incubated with human serum (6% for AP, 2% for CP) and added to plate. Percentage (%) inhibition of MAC deposition was calculated for each sample relative to baseline (EDTA treated human serum) and positive control (human serum), and used to generate the IC₅₀ curve with XLFit.

Parental Fabs

Parental Fabs were used in different concentrations and the maximal inhibition (if applicable also IC₅₀ values) were determined (example shown in FIG. 27). Most Fabs completely inhibited MAC deposition indicating blocking of C5 cleavage. Potency and ranking of Fabs were similar to data from hemolytic assays.

SEQUENCE LISTING IN ELECTRONIC FORM

In accordance with Section 111(1) of the Patent Rules, this description contains a sequence listing in electronic form in ASCII text format (file: 21489-11410 Seq 10-MAY-13 v2.txt).

A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office.

CLAIMS:

1. An isolated monoclonal human C5 protein antibody, or an antigen binding fragment thereof comprising:
 - (i) a heavy chain CDR1 comprising the sequence of SEQ ID NO:1; a heavy chain CDR2 comprising the sequence of SEQ ID NO: 2; and a heavy chain CDR3 sequence comprising the sequence of SEQ ID NO:3; and
 - (ii) a light chain CDR1 comprising the sequence of SEQ ID NO:4; a light chain CDR2 comprising the sequence of SEQ ID NO:5; and a light chain CDR3 sequence comprising the sequence of SEQ ID NO: 6, wherein said antibody or antigen binding fragment thereof binds to human C5 protein with a K_D of 10^{-10} M or less as measured by surface plasmon resonance.
2. An isolated monoclonal antibody, or antigen binding fragment as claimed in claim 1 comprising a heavy chain variable region having at least 80%, 90%, 95%, 97%, 98% or 99% sequence identity to an amino acid sequence of SEQ ID NO: 7; and a light chain variable region having at least 80%, 90%, 95%, 97%, 98% or 99% sequence identity to an amino acid sequence of SEQ ID NO: 8.
3. An isolated monoclonal antibody, or antigen binding fragment as claimed in claim 1 or claim 2 comprising a heavy chain variable region consisting of the amino acid sequence of SEQ ID NO: 7; and a light chain variable region consisting of the amino acid sequence of SEQ ID NO: 8.
4. An isolated monoclonal antibody, or antigen binding fragment as claimed in any one of claims 1 to 3 comprising a heavy chain having at least 95%, 96%, 97%, 98% or 99% sequence identity to SEQ ID NO: 9; and a light chain having at least 95%, 97%, 98% or 99% sequence identity to SEQ ID NO: 10.
5. An isolated monoclonal antibody, or antigen binding fragment as claimed in claim 4 comprising a heavy chain consisting of the amino acid sequence of

SEQ ID NO: 9; and a light chain consisting of the amino acid sequence of SEQ ID NO: 10.

6. The antibody as claimed in any one of claims 1 to 5 which inhibits the alternative complement pathway with an IC_{50} of 20-200 pM as measured by *in vitro* hemolytic assay when using human C5 depleted serum that is reconstituted with 100 pM human C5.
7. The antibody or antigen binding fragment as claimed in any one of claims 1 to 6, wherein said antibody is a human or chimeric antibody.
8. The antibody as claimed in any one of claims 1 to 7, wherein said antibody is an IgG isotype.
9. A pharmaceutical composition comprising the antibody or antigen binding fragment as claimed in any one of claims 1-8 and a pharmaceutically acceptable carrier.
10. An isolated nucleic acid comprising a nucleotide sequence encoding the antibody or antigen binding fragment as claimed in any one of claims 1 to 8.
11. The nucleic acid as claimed in claim 10, comprising a nucleotide sequence encoding a polypeptide comprising a heavy chain variable region having at least 95% sequence identity to SEQ ID NO: 7, or a light chain variable region having at least 95% sequence identity to SEQ ID NO: 8.
12. The nucleic acid as claimed in claim 11 wherein the nucleotide sequence encoding the polypeptide comprising a heavy chain variable region consists of the sequence of SEQ ID NO: 11, and the nucleotide sequence encoding the polypeptide comprising a light chain variable region consists of the sequence of SEQ ID NO: 12.
13. A vector comprising the nucleic acid of any one of claims 10 to 12.

14. An isolated host cell comprising recombinant DNA segments encoding a heavy chain and a light chain of the antibody as claimed in any one of claims 1 to 8; wherein said DNA segments are operably linked to a promoter, and are capable of being expressed in said host cell.
- 5 15. Use of an antibody of any one of claims 1-8 for treating age related macular degeneration in a subject.
16. The use according to claim 15, wherein the subject is a human.
17. Use of an antibody of any one of claims 1-8 for treating a disease,
wherein said disease is hyperacute allograft rejection, xenograft rejection, asthma,
10 arthritis, autoimmune heart disease, multiple sclerosis, inflammatory bowel disease, ischemia-reperfusion injuries, Barraquer-Simons Syndrome, hemodialysis, systemic lupus, lupus erythematosus, psoriasis, multiple sclerosis, transplantation, Alzheimer's disease, glomerulonephritis, MPGN II, wet age related macular degeneration, dry age related macular degeneration or geographic atrophy.
- 15 18. The use according to claim 17, wherein said disease is glomerulonephritis.
19. Use of an antibody of any one of claims 1-8 for treating paroxysmal nocturnal hemoglobinuria (PNH).
20. The pharmaceutical composition according to claim 9 for use in the
20 treatment of age related macular degeneration in a subject.
21. The pharmaceutical composition according to claim 20, wherein the subject is a human.
22. The pharmaceutical composition according to claim 9 for use in the treatment of a disease, wherein said disease is hyperacute allograft rejection,
25 xenograft rejection, asthma, arthritis, autoimmune heart disease, multiple sclerosis, inflammatory bowel disease, ischemia-reperfusion injuries, Barraquer-Simons

Syndrome, hemodialysis, systemic lupus, lupus erythematosus, psoriasis, multiple sclerosis, transplantation, Alzheimer's disease, glomerulonephritis, MPGN II, wet age related macular degeneration, dry age related macular degeneration or geographic atrophy.

5 23. The pharmaceutical composition according to claim 22, wherein said disease is glomerulonephritis.

24. The pharmaceutical composition according to claim 9 for use in the treatment of paroxysmal nocturnal hemoglobinuria (PNH).

MOR08109	VH	EVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIGPFFGTANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCARDTPYFDYWGQGT	LTVTVSS
1-69	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISWVRQAPGQGLEWMGGIIPITFGTANYAQKFQGRVTITADESTSTAYMELSSLRSEDTAVYYCAR		
MOR08109	VL	SYELTQPLSVSVALGQTARITCSGDSIPNYVYVWYQQKPGQAPVLIYDDSNRPSGIPERFSGSNSGNTATLTISRQAQGEADYYCQSFDSLNAEVFGGGTKL	TVL
3j	SYELTQPLSVSVALGQTARITCSGNNIGSKNVHWYQQKPGQAPVLIYRDSNRPSGIPERFSGSNSGNTATLTISRQAQGEADYYCQVWDSSTA		
MOR08110	VH	EVQLVQSGAEVKKPGESLKISCKGSGYSFTN-YISWVRQMPGKGLEWMGIIIDPDDSYTEYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCARYEYGGFDI
5-51	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIIPGDSITRYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCAR	
MOR08110	VL	SYELTQPPSVSVAPGQTARISCGDNIIGNSYVHWYQQKPGQAPVLIYKDNDRPSGIPERFSGSNSGNTATLTISGTQAE	DEADYYCGTYDIESYVFGGGTKL
3r	SYELTQPPSVSVSPGQTASITCSGDKLGDKYACWYQQKPGQSPVLVIYQDSKRPSGIPERFSGSNSGNTATLTISGTQAM	DEADYYCQAWDESTA	
MOR08111	VH	EVTLKESGPALVKPTQTLTLTCTFSGFSLSSTSGGGVSWIRQPPGKALEWLANIIDDAIDKDYSPSLKSRRLTISKDTSKNQVVL	TMTNMDPVD
2-70	QVTLKESGPALVKPTQTLTLTCTFSGFSLSSTSGMRVSWIRQPPGKALEWLANIIDDDDKFYSTSLKTRLTISKDTSKNQVVL	TMTNMDPVD	TATYYCARPYGFD
MOR08111	VL	HSALTQPASVSGSPGQSITISQGTSSDITGTYNYVSWYQQHPGKAPKLMYDDSNRPSGVSNRFSGSKSGNTASLTISGLQAE	DEADYYCQSYDSQSYIVFGGGTKL
2a2	QSALTQPASVSGSPGQSITISQGTSSDVGGYNYVSWYQQHPGKAPKLMYEVSNRPSGVSNRFSGSKSGNTASLTISGLQAE	DEADYYCQSSYTSSTL	
MOR08113	VH	EVQLVQSGAEVKKPGESLKISCKGSGYSFTN-YISWVRQMPGKGLEWMGIIIDPDDSYTRYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCARYEYGGFDI
5-51	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIIPGDSITRYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCAR	
MOR08H3	VL	SYELTQPPSVSVAPGQTARISCGDNIIGNSYVHWYQQKPGQAPVLIYKDNDRPSGIPERFSGSNSGNTATLTISGTQAD	DEADYYCATI
3r	SYELTQPPSVSVSPGQTASITCSGDKLGDKYACWYQQKPGQSPVLVIYQDSKRPSGIPERFSGSNSGNTATLTISGTQAM	DEADYYCQAWDSSTA	
MOR08114	VH	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIIDPDDSYTRYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCARYMMRGFDH
5-51	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGWVRQMPGKGLEWMGIIIPGDSITRYSPSFQGGQVTISADKSI	STAYLQWSSLKASDTAMYYCAR	
MOR08114	VL	SYELTQPPSVSVAPGQTARISCGDSLGDYVYAWYQQKPGQAPVLIYKDNDRPSGIPERFSGSNSGNTATLTISGTQAE	DEADYYCQTD
3r	SYELTQPPSVSVSPGQTASITCSGDKLGDKYACWYQQKPGQSPVLVIYQDSKRPSGIPERFSGSNSGNTATLTISGTQAM	DEADYYCQAWDSSTA	

FIG.1

2/73




FIG.2

3/73




FIG.3




FIG.4

5/73






FIG.5



7/73

6GHA23_hemolytic assay CP normalized*




*100% lysis = 20% serum lysis

Max. inhibition (%):

MOR07091: 80
 MOR07829: 95
 MOR07830: 90
 MOR07871: 90
 MOR07872: 95
 MOR07873: 97
 MOR07874: 100
 MOR07908: 97

FIG.7-1

8/73



*100% lysis = 20% serum lysis


Max. inhibition (%):

MOR07092: 100
MOR07831: 100
MOR07832: 100
MOR07875: 30
MOR07876: 100
MOR07877: 100
MOR07878: 100
MOR07879: 95
MOR07909: 100
MOR07910: 100

FIG.7-2

9/73

6GHA23_hemolytic assay CP normalized*



*100% lysis = 20% serum lysis


Max. inhibition (%):

MOR07093: 30
 MOR07833: 75
 MOR07834: 70
 MOR07835: 60
 MOR07836: 75
 MOR07890: 65

FIG.7-3

10/73

6GHA23_hemolytic assay CP normalized*

Max. inhibition (%):

MOR07086: 95
MOR07864: 100
MOR07865: 100
MOR07866: 100
MOR07867: 100
MOR07868: 95
MOR07869: 100
MOR07870: 97

FIG.7-4

11/73




FIG.8-1

12/73




FIG.8-2

13/73




FIG.8-3

14/73




FIG.8-4




FIG. 9a




FIG.9b




FIG.9c

18/73




FIG.9d




FIG.9e




FIG.9f




FIG.9g

22/73




FIG.9h

23/73




FIG.9i

24/73




FIG.9j




FIG. 10a




FIG. 10b

27/73




FIG. 10C




FIG. 10d




FIG.10e




FIG. 10f

31/73




FIG.11a




FIG.11b




FIG.11C




FIG.11d




FIG.11e

36/73




FIG.11f

37/73

3SGI35_Hemolytic assay: CP, 20% human serum




FIG.12




FIG.13

39/73




FIG. 14a

40/73




FIG.14b

41/73




FIG.14c

42/73




FIG.15a

43/73




FIG.15b

44/73




FIG.15C




FIG.15d




FIG.15e

47/73




FIG.15f




FIG. 16a




FIG. 16b

50/73




FIG.16C




FIG.16d




FIG.17-1

53/73




FIG.17-2

54/73




FIG.17-3

55/73

6GHA18_QC ELISA:
MOR07091 + HCDR2- and LCDR3-derivates on huC4




FIG.17-4

MOR0	retained binding activity
MOR07086	88.02%
MOR07832	69.06%
MOR07834	66.95%
MOR07872	82.72%
MOR07876	78.55%
MOR07086	92.94%
MOR07832	74.84%
MOR07834	92.00%
MOR07872	90.70%
MOR07876	75.60%
MOR07086	89.57%
MOR07832	91.56%
MOR07834	86.05%
MOR07872	72.10%
MOR07876	70.34%




FIG.18

Rows: unlabelled Fab (100-fold excess)						Columns: biotinylated Fab					
MOR0	7086	7832	7834	7871	7872	7873	7876	7878			
7086	691	2597	853	1191	825	694	1002	495			
7832	668	1186	404	586	3601	499	595	251			
7834	618	1727	391	625	609	455	756	560			
7871	426	1384	422	758	619	1009	392	451			
7872	1553	3199	773	1486	944	882	842	781			
7873	800	3342	965	1414	1491	922	1188	691			
7876	892	2535	652	978	1366	1394	621	701			
7878	850	2643	784	1228	2218	891	1180	643			
100%	36816	37112	37340	43250	41651	33768	38078	37405			
competition with identical Fab						biotinylated Fab without competition = 100% signal					

FIG.19

58/73





FIG. 20






FIG.22-1




FIG. 22-2




FIG.22-3

63/73




Fig. 23

64/73




Fig. 24

65/73




FIG. 25-1

66/73




FIG. 25-2

67/73




FIG. 25-3

68/73




FIG. 25-4

69/73




FIG. 26-1




FIG.26-2

71/73




FIG. 26-3

72/73




FIG.26-4




FIG.27