

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 2247577 T4**

(12) Oversættelse af ændret
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 D 231/12 (2006.01)** **C 07 C 251/80 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2022-07-18**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om opretholdelse af patentet i ændret form: **2022-04-27**
- (86) Europæisk ansøgning nr.: **09715350.6**
- (86) Europæisk indleveringsdag: **2009-02-12**
- (87) Den europæiske ansøgnings publiceringsdag: **2010-11-10**
- (86) International ansøgning nr.: **EP2009000958**
- (87) Internationalt publikationsnr.: **WO2009106230**
- (30) Prioritet: **2008-02-25 EP 08151891**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK TR**
- (73) Patenthaver: **Bayer Intellectual Property GmbH, Alfred-Nobel-Strasse 10, 40789 Monheim am Rhein, Tyskland**
- (72) Opfinder: **PAZENOK, Sergii, Wipperauer Strasse 57, 42699 Solingen, Tyskland**
LUI, Norbert, Am Geus Garten 2, 51519 Odenthal, Tyskland
HEINRICH, Jens-Dietmar, Nagelsbaum 52a, 51399 Burscheid, Tyskland
WOLLNER, Thomas, Lessingstrasse 62, 50825 Köln, Tyskland
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **FREMANGSMÅDE TIL REGIOSELEKTIV SYNTSE AF 1-ALKYL-3-HALOALKYL-PYRAZOL-4-CARBOXYLSYREDERIVATER**
- (56) Fremdragne publikationer:
EP-A1- 1 854 788
WO-A1-2005/042468
R.J. ALTENBACH ET AL: JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, Bd. 50, Nr. 22, 1. November 2007 (2007-11-01), Seiten 5439-5448, XP002488452, ISSN: 0022-2623, DOI: DOI:10.1021/JM0705051 [gefundet am 2007-10-06]
E. OKADA ET AL: HETEROCYCLES, ELSEVIER SCIENCE PUBLISHERS B.V. AMSTERDAM, NL, Bd. 34, Nr. 4, 1. Januar 1992 (1992-01-01), Seiten 791-798, XP009103121, ISSN: 0385-5414
C. WAKSELMAN ET AL: JOURNAL OF THE CHEMICAL SOCIETY, CHEMICAL COMMUNICATIONS, Nr. 23, 1975, XP008131157, ISSN: 0022-4936


Description

The present invention relates to a process for the regioselective synthesis of 1-alkyl-3-haloalkylpyrazole-4-carboxylic acid derivatives by cyclization of 2,3-disubstituted acrylic acid derivatives, and to the hydrazones formed as intermediates in the process.

2-Dihaloacyl-3-dialkylaminoacrylic esters of the formula II (Y=COOAlk, Z=O) are useful intermediates for the preparation of dihalomethyl-substituted pyrazolylcarboxylic acid derivatives which for their part are precursors of fungicidally active compounds (cf. WO 03/070705).


Pyrazolecarboxylic acid derivatives are usually prepared by reacting acrylic acid derivatives having two leaving groups (Z and A) with hydrazines.

The reaction with the monoalkylhydrazines gives mainly 1-alkylpyrazoles. However, the cyclization is frequently not regioselective. As a consequence, depending on the substrate and the reaction conditions, the unwanted 5-alkylpyrazoles are formed in amounts between 10 and 80% (see Scheme 1).



Scheme 1

The synthesis of 1-alkylpyrazolecarboxylic acid derivatives by alkylation of pyrazole derivatives unsubstituted in the 1-position frequently also proceeds with formation of both regioisomers (see Scheme 2).



Scheme 2

- 5 An alternative route of preparing fluorohaloalkylpyrazolecarboxylic acids is the cyclization of, for example, 4,4-dichloro-2-[(dimethylamino)methylidene]-3-oxobutanoate with alkylhydrazines followed by a halogen exchange.
- 10 WO 2005/042468 discloses a process for preparing 2-dihaloacyl-3-aminoacrylic esters by reacting acid halides with dialkylaminoacrylic esters, followed by their cyclization with alkylhydrazines.
- 15 The European patent application No. 07117232.4, hitherto unpublished, describes a process for preparing HCl-free 2-dihaloacyl-3-aminoacrylic esters by reacting acid fluorides with dialkylaminoacrylic acid derivatives. The process can be carried out in the absence of a base, as a result of which the removal
- 20 of halide salts is not required.
- WO 2008/092583 describes a process for preparing 3-dihalomethylpyrazole-4-carboxylic acid derivatives by reacting α,α -fluoroamines in the presence of Lewis acids with acrylic
- 25 acid derivatives, followed by their reaction with alkylhydrazines.
- WO 2006/090778 discloses a process for preparing 1-methyl-3-difluoromethylpyrazolecarboxylic esters by cyclization of 2-alkoxymethylenefluoroacylacetate with methylhydrazine in the
- 30 presence of water and a base.
- EP 1854788 describes a regioselective process for preparing alkyl 1-alkyl-3-fluoroalkyl-1H-pyrazole-4-carboxylates by

cyclization of 2-alkoxymethylenefluoroacyl acetates with alkyl hydrazines in the presence of water and a base.

5 R.J. ALTENBACH ET AL.: JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN
CHEMICAL SOCIETY, WASHINGTON, US, Vol. 50, No. 22,
1 November 2007 (2007-11-01), pages 5439-5448 describes a
process for preparing 1-(1,3-dimethyl-1H-pyrazol-4-yl)ethanones
where, in a first step, 3-ethoxymethylenepentane-2,4-dione is
reacted with 1-methyl-2-(propan-2-ylidene)hydrazone to give
10 3-((1-methyl-2-(propan-2-ylidene)hydrazinyl)methylene)pentane-
2,4-dione, which is cyclized in a second step.

E. OKADA ET AL.: HETEROCYCLES, ELSEVIER SCIENCE PUBLISHERS B.V.
AMSTERDAM, NL, Vol. 34, No. 4, 1 January 1992 (1992-01-01),
15 pages 791-798 describes a process for preparing 4-
trifluoroacetyl-3-trifluoromethyl-1-methyl-1H-pyrazole and 4-
trifluoroacetyl-3-trifluoromethyl-1-phenyl-1H-pyrazole by
reacting β,β -bis(trifluoroacetyl)vinyl isobutyl ether with
benzaldehyde methylhydrazone or benzaldehyde phenylhydrazone to
20 give benzaldehyde N- $[\beta,\beta$ -bis(trifluoroacetyl)vinyl]-
methylhydrazone and benzaldehyde N- $[\beta,\beta$ -
bis(trifluoroacetyl)vinyl]phenylhydrazone, respectively, and
their subsequent cyclization.

25 WO 2005/042468 describes processes for preparing 2-dihaloacyl-
3-aminoacrylates by reacting acid halides with
dialkylaminoacrylates, and their subsequent cyclization with
alkylhydrazines to give 1-alkyl-3-dihalomethyl-1H-pyrazole
carboxylates.


30 C. WAKSELMAN ET AL.: JOURNAL OF THE CHEMICAL SOCIETY, CHEMICAL
COMMUNICATIONS, No. 23, 1975 discloses a process for the
iminoalkylation/acylation of electron-rich aromatic compounds
using fluorinated immonium salts which can be obtained by
35 reacting α,α -difluoroamines with BF_3 .

However, the processes above-described all have the disadvantage
that the cyclization, even at low temperatures, proceeds only

with unsatisfactory regioselectivity.

In view of the prior art described above, it is an object of the present invention to provide a process which does not have the disadvantages mentioned above, thus providing regioselective access to 1-alkyl-3-haloalkylpyrazole-4-carboxylic acid derivatives.

The object described above was achieved by a process for preparing 1-alkyl-3-haloalkylpyrazole-4-carboxylic acid derivatives of the formula (I)



in which

R^1 is selected from the group consisting of methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl and tert-butyl;

R^2 is selected from C_1 - C_4 -alkyl groups which may be substituted by one, two or three halogen atoms selected from the group consisting of F, Cl and Br or by a CF_3 group;


Y is selected from the group consisting of $(C=O)OR^3$, CN and $(C=O)NR^4R^5$, where R^3 , R^4 and R^5 independently of one another are selected from the group consisting of C_{1-12} -alkyl, C_{3-8} -cycloalkyl, C_{2-12} -alkenyl, C_{2-12} -alkynyl, C_{6-8} -aryl, C_{7-19} -arylalkyl and C_{7-19} -alkylaryl groups, each of which may be substituted by one or more groups selected from the group consisting of - R' , -X, - OR' , - SR' , - NR'_2 , - SiR'_3 , - $COOR'$, - $(C=O)R'$, -CN and - $CONR'_2$, where R' is hydrogen or a C_{1-12} -alkyl group, and R^4 and R^5 together with the nitrogen atom to which they are attached may form a 5- or 6-membered ring which may optionally contain one or more further heteroatoms selected from the group

consisting of O, S and an SO₂ group and which may be substituted by one or more groups selected from the group consisting of -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN and -CONR'₂, where R' is hydrogen or a C₁₋₁₂-alkyl group;

5

comprising at least one of the reaction sequences consisting of steps (A) and (D), (B) and (D) or (C) and (D) below:

(A) reaction of a 2-acylated or 2-iminoalkylated acrylic acid
10 derivative of the formula (II),



in which

15


Z is selected from the group consisting of O, S and N⁺R¹⁰R¹¹, where R¹⁰ and R¹¹ independently of one another are selected from the group consisting of C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₆₋₈-aryl, C₇₋₁₉-arylalkyl and C₇₋₁₉-alkylaryl groups, each of which may be substituted by one or more groups selected from the group consisting of -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN and -CONR'₂, where R' is hydrogen or a C₁₋₁₂-alkyl group, and R¹⁰ and R¹¹ together with the nitrogen atom to which they are attached
20 may form a 5- or 6-membered ring which may optionally contain one or two further heteroatoms selected from the group consisting of O, S and an SO₂ group and which may be substituted by one or more groups selected from the group consisting of -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN and -CONR'₂, where R' is hydrogen or a C₁₋₁₂-alkyl group;

30

A is a leaving group NR¹²R¹³, where R¹² and R¹³ independently of one another are selected from the group consisting of C₁₋₁₂-alkyl, C₃₋₈-cycloalkyl, C₂₋₁₂-alkenyl, C₂₋₁₂-alkynyl, C₆₋₈-aryl, C₇₋₁₉-arylalkyl and C₇₋₁₉-alkylaryl groups, each of which may be
35

substituted by one or more groups selected from the group consisting of -
 R' , -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN
 and -CONR'₂, where R' is hydrogen or a C₁₋₁₂-alkyl group, and R¹²
 5 and R¹³ together with the nitrogen atom to which they are attached
 may form a 5- or 6-membered ring which may optionally contain
 one or two further heteroatoms selected from the group
 consisting of O, S and an SO₂ group and which may be substituted
 by one or more groups selected from the group consisting of -
 10 R' , -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN
 and -CONR'₂, where R' is hydrogen or a C₁₋₁₂-alkyl group;

with an N-alkylhydrazone of the formula (III)




in which

R⁸ is selected from the group consisting of H, methyl, ethyl,
 20 n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, phenyl,
 tolyl, cyclohexyl;

R⁹ is selected from the group consisting of H, methyl, ethyl,
 n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, phenyl,
 25 tolyl, cyclohexyl;


where R⁸ and R⁹ together with the carbon atom to which they are
 attached may form a 5- or 6-membered ring;

30 (B) acylation of an acrylic acid hydrazone of the formula (VI);




with an acyl halide of the formula (X), where X is selected from the group consisting of F, Cl, Br and I,

5



(C) iminoalkylation of an acrylic acid hydrazone of the formula (VI) above with an α,α -dihaloamine of the formula (XI), where X is selected from the group consisting of F, Cl, Br and I,

10



(D) cyclization of the intermediates obtained in step (A), (B) or (C) to give the 3-haloalkylpyrazole-4-carboxylic acid derivative of the formula (I).

15

Surprisingly, under the conditions according to the invention, the 1-alkyl-3-haloalkylpyrazole-4-carboxylic acid derivatives of the formula (I) can be prepared with good yields, regioselectivities and in high purity, the process according to the invention thus overcoming the disadvantages mentioned above of the preparation processes described in the prior art.

20

25 General definitions

In the context of the present invention, the term halogens (X) comprises, unless defined otherwise, elements selected from the group consisting of fluorine, chlorine, bromine and iodine,

where fluorine, chlorine and bromine are used with preference and fluorine and chlorine with particular preference.

Optionally substituted groups may be mono- or polysubstituted, where in the case of polysubstitution the substituents may be identical or different.

Alkyl groups substituted by one or more halogen atoms (-X) are, for example, selected from the group consisting of trifluoromethyl (CF₃), difluoromethyl (CHF₂), CF₃CH₂, ClCH₂, CF₃CCl₂.

In the context of the present invention, unless defined otherwise, alkyl groups are straight-chain or branched hydrocarbon groups which may optionally have one, two or more heteroatoms selected from the group consisting of O, N, P and S. Moreover, the alkyl groups according to the invention may optionally be substituted by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-C(=O)R') and amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may have one or more heteroatoms selected from the group consisting of N, O, P and S.

In the context of the present invention, cycloalkyl groups are, unless defined otherwise, monocyclic saturated hydrocarbon groups having 3 to 8 carbon ring members, which groups may optionally contain one, two or more heteroatoms selected from the group consisting of O, N, P and S. Moreover, the cycloalkyl groups according to the invention may optionally be substituted by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino- (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-C(=O)R') and amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may contain one

or more heteroatoms selected from the group consisting of N, O, P and S.

Specifically, this definition comprises, for example, the meanings cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The definition C₁-C₁₂-alkyl comprises the largest range defined herein for an alkyl group. Specifically, this definition comprises, for example, the meanings methyl, ethyl, n-, isopropyl, n-, iso-, sec- and t-butyl, n-pentyl, n-hexyl, 1,3-dimethylbutyl, 3,3-dimethylbutyl, n-heptyl, n-nonyl, n-decyl, n-undecyl, n-dodecyl.

In the context of the present invention, alkenyl groups are, unless defined otherwise, straight-chain or branched hydrocarbon groups which contain at least one single unsaturation (double bond) and may optionally have one, two or more single or double unsaturations or one, two or more heteroatoms selected from the group consisting of O, N, P and S. Moreover, the alkenyl groups according to the invention may optionally be substituted by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-C(=O)R') and amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may contain one or more heteroatoms selected from the group consisting of N, O, P and S.

The definition C₁-C₁₂-alkenyl comprises the largest range defined herein for an alkenyl group. Specifically, this definition comprises, for example, the meanings vinyl; allyl (2-propenyl), isopropenyl (1-methylethenyl); but-1-enyl (crotyl), but-2-enyl, but-3-enyl; hex-1-enyl, hex-2-enyl, hex-3-enyl, hex-4-enyl, hex-5-enyl; hept-1-enyl, hept-2-enyl, hept-3-enyl, hept-4-enyl, hept-5-enyl, hept-6-enyl; oct-1-enyl, oct-2-enyl, oct-3-enyl, oct-4-enyl, oct-5-enyl, oct-6-enyl, oct-7-enyl; non-1-enyl, non-2-enyl, non-3-enyl, non-4-enyl, non-5-enyl, non-6-enyl,

non-7-enyl, non-8-enyl; dec-1-enyl, dec-2-enyl, dec-3-enyl, dec-4-enyl, dec-5-enyl, dec-6-enyl, dec-7-enyl, dec-8-enyl, dec-9-enyl; undec-1-enyl, undec-2-enyl, undec-3-enyl, undec-4-enyl, undec-5-enyl, undec-6-enyl, undec-7-enyl, undec-8-enyl, undec-9-enyl, undec-10-enyl; dodec-1-enyl, dodec-2-enyl, dodec-3-enyl, dodec-4-enyl, dodec-5-enyl, dodec-6-enyl, dodec-7-enyl, dodec-8-enyl, dodec-9-enyl, dodec-10-enyl, dodec-11-enyl; buta-1,3-dienyl, penta-1,3-dienyl.

- 10 In the context of the present invention, cycloalkenyl groups are, unless defined otherwise, monocyclic nonaromatic hydrocarbon groups having 3 to 8 carbon ring members and at least one double bond, which groups may optionally contain one, two or more heteroatoms selected from the group consisting of
- 15 O, N, P and S. Moreover, the cycloalkenyl groups according to the invention may optionally be substituted by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-C(=O)R') and
- 20 amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may contain one or more heteroatoms selected from the group consisting of N, O, P and S.
- 25 Specifically, this definition comprises, for example, the meanings cyclopenten-1-yl, cyclohexen-1-yl, cyclohepta-1,3-dien-1-yl.

- In the context of the present invention, alkynyl groups are,
- 30 unless defined otherwise, straight-chain, branched or cyclic hydrocarbon groups which contain at least one double unsaturation (triple bond) and may optionally have one, two or more single or double unsaturations or one, two or more heteroatoms selected from the group consisting of O, N, P and
- 35 S. Moreover, the alkynyl groups according to the invention may optionally be substituted by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl

(-COOR'), cyano (-CN), acyl $-(C=O)R'$ and amide $(-CONR'_2)$ groups, where R' is hydrogen or a straight-chain, branched or cyclic C_{1-12} -alkyl group which may contain one or more heteroatoms selected from the group consisting of N, O, P and S.

5

The definition C_2-C_{12} -alkynyl comprises the largest range defined herein for an alkynyl group. Specifically, this definition comprises, for example, the meanings ethynyl (acetylenyl); prop-1-ynyl and prop-2-ynyl.

10

In the context of the present invention, aryl groups are, unless defined otherwise, aromatic hydrocarbon groups which may contain one, two or more heteroatoms selected from the group consisting of O, N, P and S and which may optionally be substituted by further groups selected from the group consisting of $-R'$, halogen $(-X)$, alkoxy $(-OR')$, thioether or mercapto $(-SR')$, amino $(-NR'_2)$, silyl $(-SiR'_3)$, carboxyl $(-COOR')$, cyano $(-CN)$, acyl $-(C=O)R'$ and amide $(-CONR'_2)$ groups, where R' is hydrogen or a C_{1-12} -alkyl group, preferably a C_{2-10} -alkyl group, particularly preferably a C_{3-8} -alkyl group, which may contain one or more heteroatoms, selected from the group consisting of N, O, P and S.

20

The definition C_{5-18} -aryl comprises the largest range defined herein for an aryl groups having 5 to 18 skeleton atoms, where the carbon atoms may be replaced by heteroatoms. Specifically, this definition comprises, for example, the meanings cyclopentadienyl, phenyl, cycloheptatrienyl, cyclooctatetraenyl, naphthyl and anthracenyl; 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyrrolyl, 3-pyrrolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,2,4-triazol-3-yl, 1,3,4-oxadiazol-2-yl, 1,3,4-thiadiazol-2-yl and 1,3,4-triazol-2-yl; 1-pyrrolyl, 1-pyrazolyl, 1,2,4-triazol-1-yl, 1-imidazolyl, 1,2,3-triazol-1-yl, 1,3,4-triazol-1-yl; 3-pyridazinyl,

35

4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 1,3,5-triazin-2-yl and 1,2,4-triazin-3-yl.

In the context of the present invention, arylalkyl groups (aralkyl groups) are, unless defined otherwise, alkyl groups which are substituted by aryl groups, which may have a C₁₋₈-alkylene chain and which may be substituted in the aryl skeleton or the alkylene chain by one or more heteroatoms selected from the group consisting of O, N, P and S and optionally by further groups selected from the group consisting of -R', halogen- (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-(C=O)R') and amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may contain one or more heteroatoms selected from the group consisting of N, O, P and S.

The definition C₇₋₁₉-aralkyl group comprises the largest range defined herein for an arylalkyl group having a total of 7 to 19 atoms in the skeleton and the alkylene chain. Specifically, this definition comprises, for example, the meanings benzyl and phenylethyl.


In the context of the present invention, alkylaryl groups (alkaryl groups) are, unless defined otherwise, aryl groups which are substituted by alkyl groups, which may have a C₁₋₈-alkylene chain and which may be substituted in the aryl skeleton or the alkylene chain by one or more heteroatoms selected from the groups consisting of O, N, P and S and optionally by further groups selected from the group consisting of -R', halogen (-X), alkoxy (-OR'), thioether or mercapto (-SR'), amino (-NR'₂), silyl (-SiR'₃), carboxyl (-COOR'), cyano (-CN), acyl (-(C=O)R') and amide (-CONR'₂) groups, where R' is hydrogen or a C₁₋₁₂-alkyl group, preferably a C₂₋₁₀-alkyl group, particularly preferably a C₃₋₈-alkyl group, which may contain one or more heteroatoms selected from the group consisting of N, O, P and S.

The definition C₇₋₁₉-alkylaryl group comprises the largest range defined herein for an alkylaryl group having a total of 7 to 19 atoms in the skeleton and the alkylene chain. Specifically, this definition comprises, for example, the meanings tolyl, 2,3-,
5 2,4-, 2,5-, 2,6-, 3,4- or 3,5-dimethylphenyl.

The alkyl, alkenyl, alkynyl, aryl, alkaryl and aralkyl groups may furthermore contain one or more heteroatoms which - unless defined otherwise - are selected from the group consisting of
10 N, O, P and S. Here, the heteroatoms replace the carbon atoms specified.


If appropriate, the compounds according to the invention can be present as mixtures of various possible isomeric forms, in
15 particular of stereoisomers, such as, for example, E and Z, threo and erythro and also optical isomers, and, if appropriate, also of tautomers. What is disclosed and claimed are both the E and the Z isomers, and also the threo and erythro isomers and the optical isomers, any mixtures of these isomers, and also the
20 possible tautomeric forms.

Scheme 3 below summarizes the individual variants of the process according to the invention:



Scheme 3

- 5 The 1-alkyl-3-haloalkylpyrazole-4-carboxylic acid derivatives obtainable by the process according to the invention are compounds of the formula (I)



(I).

According to the invention, the radicals in formula (I) have the abovementioned meanings.

5 In a *preferred* embodiment of the present invention, the radicals in formula (I) have the meanings below:

R^1 is selected from the group consisting of methyl, ethyl, n-propyl and isopropyl,

10 R^2 is selected from the group consisting of chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl,
15 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, pentafluoroethyl and 1,1,1-trifluoroprop-2-yl; 1,2,2,2-tetrafluoroethyl,

20 Y is selected from the group consisting of $(C=O)OR^3$, CN and $(C=O)NR^4R^5$, where R^3 , R^4 and R^5 independently of one another are selected from the group consisting of methyl, ethyl, n-propyl and isopropyl.

25 In a particularly preferred embodiment of the present invention, the radicals in formula (I) have the meanings below:

R^1 is methyl,

30 R^2 is selected from the group consisting of trifluoromethyl and difluoromethyl,


Y is selected from the group consisting of $(C=O)OR^3$, where R^3 is methyl or ethyl.

35

Step (A)

In a first embodiment of the present process, which is the result


of combining reaction steps (A) and (D), initially 2-acylated or 2-iminoalkylated acrylic acid derivatives of the formula (II) are reacted with N-alkylhydrazones of the formula (III) (Scheme 4). Subsequently, the intermediates formed in step (A) are cyclized to give the 3-haloalkylpyrazole-4-carboxylic acid derivatives of the formula (I) (Step D).



10 Scheme 4



Possible intermediates which may, in accordance with the present invention, result from reaction step (A) are the acrylhydrazones of the formula (VII)

15



This formula includes in particular the 2-acylhydrazinoprop-2-enoates of the formula (VIIa) and the hydrazinoaminium salts of the formula (VIIb)

20



5 in which the radicals R^1 , R^2 , R^8 to R^{11} and Y have the meanings described above and which optionally have a counterion (anion⁻) selected from the group consisting of Cl^- , BF_4^- , PF_6^- , SbF_6^- and $AlCl_4^-$.

10 The process step (A) according to the invention is preferably carried out in a temperature range of from $-20^\circ C$ to $+150^\circ C$, particularly preferably at temperatures of from $-10^\circ C$ to $+70^\circ C$

The process step (A) according to the invention is generally
15 carried out under atmospheric pressure. However, alternatively, it is also possible to operate under reduced pressure to remove the highly volatile dialkylamines.

The reaction time is not critical and may, depending on the
20 batch size and the temperature, be chosen from a range of between a few minutes and a number of hours.

When carrying out the process step (A) according to the invention, 1 mol of the acrylic acid derivative of the formula (II) is

reacted with from 0.5 mol to 3 mol, preferably from 0.5 mol to 1.5 mol, particularly preferably with an equimolar amount, of the hydrazone of the formula (III).

5 Preferably, the hydrazone of the formula (III), dissolved in the solvent, is initially charged, and the acrylic acid derivative of the formula (II) is added. However, it is also possible to reverse the order.

10 The reaction can be promoted by adding catalysts, such as, for example, CH_3COOH , H_2SO_4 , KHSO_4 , NaH_2PO_4 , HCl , CF_3COOH , CH_3COONa .


Work-up and isolation of the intermediates is generally not required, and the reaction mixture obtainable from step (A) can
15 be used immediately or after storage for the cyclization step (D).

The acrylic acid derivatives of the formula (II) can be obtained by the processes described above, in connection with the prior
20 art.

In the context of the present invention, preference is given to using 2-acylated or 2-iminoalkylated acrylic acid derivatives of the formula (II) selected from the group consisting of N-1-
25 (trifluoromethyl)-3-(dimethylamino)-2-(ethoxycarbonyl)prop-2-en-1-ylidene]-N-methylmethanaminium chloride, 2-(, N-1-(difluoromethyl)-3-(dimethylamino)-2-(ethoxycarbonyl)prop-2-en-1-ylidene]-N-methylmethanaminium tetrafluoroborate and N-[3-(dimethylamino)-2-(ethoxycarbonyl)-1-(1,1,2,2-tetra-
30 fluoroethyl)prop-2-en-1-ylidene]-N-methylmethanaminium chloride.

In the context of the present invention, the N-alkylhydrazones of the formula (III) are preferably selected from the group
35 consisting of 1-methyl-2-(1-methylethylidene)hydrazine, 1-methyl-2-(1,2,2-trimethylpropylidene)hydrazine, 1-methyl-2-(1-methylpropylidene)hydrazine, 1-cyclohexylidene-2-methylhydrazine, 1-methyl-2-(phenylmethylidene)hydrazine, 1-

methyl-2-(1-phenylethylidene)hydrazine, 1-(diphenylmethyl-
 idene)-2-methylhydrazine, ethyl 2-[(dimethylamino)methylidene]-
 4,4,4-trifluoro-3-oxobutanoate. The N-alkylhydrazones of the
 formula (III) have been described in the literature (Zhurnal
 5 Organicheskoi Khimii (1968), 4(6), 986-92.) and can be
 obtained by reacting commercially available hydrazines of the
 formula (VIII) with carbonyl compounds of the formula (IX) (see
 Scheme 5).



Scheme 5

According to the invention, the radicals in formulae (III),
 15 (VIII) and (IX) have the following meanings:

R¹ represents methyl, ethyl, n-propyl, isopropyl, n-butyl,
 sec-butyl, tert-butyl;

20 R⁸ is selected from the group consisting of H, methyl, ethyl,
 n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, phenyl,
 tolyl, cyclohexyl;

R⁹ is selected from the group consisting of H, methyl, ethyl,
 25 n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, phenyl,
 tolyl, cyclohexyl.

Alternatively, R⁸ and R⁹ together with the carbon atom to which
 they are attached may form a 5- or 6-membered ring.

30 According to the invention, the radicals in formulae (III),
 (VIII) and (IX) have the following particularly preferred
 meanings:

35 R¹ is methyl;

R⁸ is selected from the group consisting of H, methyl, tert-butyl, phenyl;

5 R⁹ is selected from the group consisting of H, methyl, tert-butyl, phenyl.

According to the invention, preference is given to using ketones of the formula (IX) selected from the group consisting of
10 acetone, benzophenone, pinacolone, cyclohexanone, benzaldehyde, with acetone and benzaldehyde being particularly preferably used.

A further advantage of the process according to the invention is the fact that, to prepare the N-methylhydrazones of the
15 formula (III), it is possible to use aqueous methylhydrazine solutions and the explosive concentrated methylhydrazine, which is also used as rocket fuel, is not necessarily required.

In two further embodiments of the process according to the invention which result when the reactions steps (B) and (D) or
20 (C) and (D) are combined, acrylic acid derivatives of the formula (VI) are acylated (step (B)) or iminoalkylated (step (C)) and finally cyclized to give the 3-haloalkylpyrazole-4-carboxylic acid derivatives of the formula (I) (step D).

25


Step (B): Acylation

In one embodiment of the process according to the invention, the acrylic acid hydrazones of the formula (VI) are acylated with
30 haloalkylcarbonyl halides of the formula (X) according to Scheme 6 below to give the 2-acylacrylic acid hydrazones of the formula (VIIa). In formula (X), R² has the above meanings and X is a halogen atom selected from the group consisting of F, Cl or Br, preferably from the group consisting of F and Cl, particularly
35 preferably F.

In the context of the present invention, the acylating agent used is preferably a haloalkylcarbonyl halide of the formula (X)

selected from the group consisting of difluoroacetyl chloride, difluoroacetic anhydride, difluoroacetyl fluoride, trifluoroacetyl chloride and trifluoroacetyl fluoride.

- 5 The acrylic acid hydrazones of the formula (VI) used in step (B) are preferably selected from the group consisting of ethyl (2E)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, ethyl (2Z)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, propyl 3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, ethyl (2E)-3-[1-methyl-2-(3,3-dimethylbutan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(3,3-dimethylbutan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(phenylmethylidene)hydrazinyl]prop-2-enoate, (2E)-3-[1-methyl-2-(phenylmethylidene)hydrazinyl]prop-2-enenitrile and methyl 3-[1-ethyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate.



Scheme 6

According to the invention, the acylation is carried out at temperatures of from -20°C to $+150^{\circ}\text{C}$, preferably at temperatures of from -20°C to $+100^{\circ}\text{C}$, particularly preferably at from -10 to 50°C , and under atmospheric pressure.

The reaction time is not critical and may, depending on the batch size and the temperature, be chosen from a range between a few minutes and a number of hours.

According to the invention, 1 mol of the acrylic acid hydrazone

of the formula (VI) is reacted with from 0.5 mol to 3 mol, preferably from 1.2 mol to 1.5 mol, particularly preferably with equimolar amounts, of the haloalkylcarbonyl halide of the formula (X).

5

Preferably, the hydrazinoacrylic acid derivative of the formula (VI) dissolved in the solvent is initially charged, and the haloalkylcarbonyl halide of the formula (X) is added. However, a reverse order is also possible.

10

The acylation is preferably carried out in the presence of a base. Preference is given here to organic bases, such as trialkylamines, alkylpyridines, phosphazenes and 1,8-diazabicyclo[5.4.0]undecene (DBU); alkali metal hydroxides, such as, for example, lithium hydroxide, sodium hydroxide or potassium hydroxide, alkali metal carbonates (Na_2CO_3 , K_2CO_3) and alkoxides, such as, for example, NaOMe, NaOEt, NaOt-Bu, KOt-Bu.

15

Suitable solvents are, for example, aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, n-hexane, n-heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decaline, and halogenated hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane, ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; nitriles, such as acetonitrile, propionitrile, n- or isobutyronitrile or benzonitrile; ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; sulphoxides, such as dimethyl sulphoxide, or sulphones, such as sulfolane. Particular preference is given to using toluene, xylene, chlorobenzene, n-hexane, cyclohexane or methylcyclohexane, very particularly preferably toluene or xylene.

20

25

30


35

Possible intermediates which, according to the present invention, may result from reaction step (B) are, usually, the 2-acylhydrazinoacrylic acid derivatives of the formula (VIIa).


- 5 The intermediates formed can, without prior work-up, be used for the cyclization step (step D).

Alternatively, the intermediates can be isolated by suitable work-up steps and, if appropriate, further purification, and
10 they can be used at a later point in time for the cyclization (step D).

The acrylic acid hydrazones of the formula (VI) have been described in the literature (see Chem. Ber. 1975, 108 (7), 2161-
15 2170; Bulletin de la Societe Chimique de France (1976), (1-2, Pt. 2), 260-4), and they can be prepared according to Scheme 7 or 8 below.




20 Scheme 7



25 Scheme 8

Step (C) Iminoalkylation

In a further embodiment of the process according to the invention, the acrylic acid hydrazones of the formula (VI) are iminoalkylated with at least one α,α -dihaloamine of the formula (XI), according to Scheme 9 below to give the hydrazinoaminium salts of the formula (VIIb). In formula (XI), X and R² have the above meanings.



Scheme 9

In Scheme 9 [L] is a Lewis acid.

In the context of the present invention, the iminoalkylating agent of the formula (XI) used may be 1,1,2,2-tetrafluoroethyl-N,N-dimethylamine, 1,1,2,2-tetrafluoroethyl-N,N-diethylamine, 1,1,2-trifluoro-2-(trifluoromethyl)ethyl-N,N-dimethylamine, 1,1,2-trifluoro-2-(trifluoromethyl)ethyl-N,N-diethylamine (Ishikawa reagent), 1,1,2-trifluoro-2-chloroethyl-N,N-dimethylamine or 1,1,2-trifluoro-2-chloroethyl-N,N-diethylamine (Yarovenko reagent). Here, preference is given to 1,1,2,2-tetrafluoroethyl-N,N-dimethylamine and 1,1,2,2-tetrafluoroethyl-N,N-diethylamine, and particular preference is given to 1,1,2,2-tetrafluoroethyl-N,N-dimethylamine.

The acrylic acid hydrazones of the formula (VI) used in step (C) are preferably selected from the group consisting of ethyl (2E)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate,

ethyl (2Z)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, propyl 3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate, ethyl (2E)-3-[1-methyl-2-(3,3-dimethylbutan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(3,3-dimethylbutan-2-ylidene)hydrazinyl]prop-2-enoate, methyl 3-[1-methyl-2-(phenylmethylidene)hydrazinyl]prop-2-enoate, (2E)-3-[1-methyl-2-(phenylmethylidene)hydrazinyl]prop-2-enenitrile and methyl 3-[1-ethyl-2-(propan-2-ylidene)hydrazinyl]prop-2-enoate.

In a preferred embodiment of the process according to the invention, the α,α -dihaloamine is initially reacted with a Lewis acid [L], such as, for example, BF_3 , AlCl_3 , SbCl_5 , SbF_5 , ZnCl_2 , PF_5 , and the mixture of the hydrazinoacrylic ester of the formula (VI) and a base, neat or dissolved in a suitable solvent, is then added.

Preference is given to organic bases such as trialkylamines, alkyipyridines, phosphazenes and 1,8-diazabicyclo[5.4.0]undecene (DBU), alkali metal hydroxides, such as, for example, lithium hydroxide, sodium hydroxide or potassium hydroxide, alkali metal carbonates (Na_2CO_3 , K_2CO_3) and alkoxides, such as, for example, NaOMe , NaOEt , NaOt-Bu , KOt-Bu .

The reaction of the α,α -dihaloamines of the formula (XI) with the acrylic acid hydrazones of the formula (VI) is usually carried out at temperatures of from -50 to 70°C , preferably from -20 to 60°C , particularly preferably from -10 to 50°C .

The reaction can be carried out under reduced pressure, under atmospheric pressure or under elevated pressure and is preferably carried out under atmospheric pressure.

The reaction can be carried out in the absence of a solvent or in a solvent. The reaction is preferably carried out in a solvent. Suitable solvents are, for example, selected from the group consisting of aliphatic and aromatic hydrocarbons, such as, for

example, n-hexane, benzene or toluene, which may be substituted by fluorine and chlorine atoms, such as methylene chloride, dichloroethane, fluorobenzene, chlorobenzene or dichlorobenzene; ethers, such as, for example, diethyl ether, diphenyl ether, methyl tert-butyl ether, isopropyl ethyl ether, dioxane, diglym, dimethylglycol, dimethoxyethane (DME) or THF; nitriles, such as methylnitrile, butylnitrile or phenylnitrile; amides, such as dimethylformamide (DMF) or N-methylpyrrolidone (NMP), or mixtures of such solvents, with acetonitrile, dichloromethane, THF, DME and ethyl acetate being particularly preferred.

The base and the acrylic acid hydrazone of the formula (VI) are preferably employed in equimolar amounts, based on the α,α -dihaloamines of the formula (XI). The base can also be used in excess. According to the invention, the ratio of base:hydrazinoacrylic acid derivative is from 1.5:1 to 0.9:1, preferably from 1.4:1 to 0.95:1, particularly preferably from 1.3:1 to 1.05:1.

Owing to the susceptibility of the α,α -dihaloamines to hydrolysis, the reaction is to be carried out under an atmosphere of inert gas in apparatus free from water.

Possible intermediates which may result in accordance with the present invention from reaction step (C) are usually the hydrazinoaminium salts of the formula (VIIb).

The intermediates formed can be used without prior work-up for the cyclization step (step D).

Alternatively, the intermediates can be isolated by suitable work-up steps and, if appropriate, further purification and can be used at a later point in time for the cyclization (step D).

Step (D): Cyclization

According to the invention, the cyclization of the intermediates obtainable from steps (A), (B) or (C) is carried out at

temperatures of from -20°C to $+150^{\circ}\text{C}$, preferably at temperatures of from -10°C to $+100^{\circ}\text{C}$, particularly preferably at from -10 to 50°C , and under atmospheric pressure.

- 5 The reaction time is not critical and can be chosen from a relatively wide range, depending on the batch size.

According to the invention, the ring closure to the pyrazole is carried out in the presence of an acid. Here, the carbonyl
10 compound $\text{R}^8\text{R}^9\text{C}=\text{O}$ is cleaved off. The carbonyl compounds can be separated off and recycled into the preparation of hydrazones of the formula (III).

Usually, step (D) is carried out after step (A), (B) or (C)
15 without a change of solvent.

Ideally, all reaction steps of the process according to the invention are carried out in the same solvent.

20 Suitable solvents are, for example, aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, n-hexane, n-heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decaline, and halogenated hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloro-
25 methane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane, ethers, such as diethyl ether, diisopropyl ether, methyl tert-butyl ether, methyl tert-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; nitriles, such as acetonitrile, propionitrile, n- or
30 isobutyronitrile or benzonitrile; ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; sulfoxides, such as dimethyl sulfoxide, or sul-
35 phones, such as sulpholane; alcohols, such as methanol, ethanol, n- or isopropanol, butanol. Particular preference is given to using toluene, xylene, chlorobenzene, n-hexane, cyclohexane or methylcyclohexane, ethanol, very particularly preferably

ethanol, toluene, xylene and water.


Suitable acids are selected from the group consisting of HCl, H₂SO₄, CF₃COOH, CF₃SO₃H, CH₃COOH; particular preference is given to HCl and H₂SO₄.

After the reaction has ended, for example, the solvents are removed and the product is isolated by filtration or initially extracted with water, the organic phase is separated off and the solvent is removed by distillation.

The process according to the invention is illustrated in more detail by the examples below. However, the examples are not to be interpreted as limiting.

Preparation Examples

Example 1: Ethyl 4,4-difluoro-2-[[1-methyl-2-(phenylmethylidene)hydrazino]methylidene]-3-oxobutanoate




6.2 g of 2-methyl-1-phenylhydrazone and 9.3 g of ethyl 2-(difluoroacetyl)-3-(dimethylamino)acrylate were initially charged in 100 ml of toluene, and 5.7 g of potassium bisulphate were added. The reaction mixture was then heated at 35-40°C and stirred at the same temperature for 8 h. The salts were separated off, and the solvent was then removed under reduced pressure. This gave ethyl 4,4-difluoro-2-[[1-methyl-2-(phenylmethylidene)hydrazino]methylidene]-3-oxobutanoate as a

solid (91% yield). The isomer ratio according to NMR was 53/39%.

¹H-NMR (278 K, CDCN₃): δ = 1.08 (t, 3H); 1.23 (t, 3H); 3.44 (s, 3H); 3.52 (s, 3H); 4.14 (m, 2H); 6.0 (t, 1H); 6.37 (t, 1H); 7.4
5 - 7.98 (m, 7H) ppm.

Patentkrav

1. Fremgangsmåde til fremstilling af 1-alkyl-3-haloalkyl-pyrazol-4-carboxylsyrederivater med formel (I)




hvor

R¹ er udvalgt af methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl eller tert-butyl;

R² er udvalgt af C₁-C₄-alkylgrupper, som kan være substitueret med et, to eller tre halogenatomer udvalgt af F, Cl og Br eller af en CF₃-gruppe;

Y er udvalgt af gruppen bestående af (C=O)OR³, CN og (C=O)NR⁴R⁵, idet R³, R⁴ og R⁵ uafhængigt af hinanden er udvalgt af C₁-₁₂-alkyl-, C₃-₈-cycloalkyl-, C₂-₁₂-alkenyl-, C₂-₁₂-alkynyl-, C₆-₈-aryl-, C₇-₁₉-arylalkyl- eller C₇-₁₉-alkylaryl-grupper, som hver især kan være substitueret med en eller flere grupper, som er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁-₁₂-alkylgruppe, og R⁴ og R⁵ sammen med det nitrogenatom, som de er bundet til, kan danne 5- til 6-leddet ring, som i givet fald kan indeholde et eller to yderligere heteroatomer udvalgt af O, S og en SO₂-gruppe, og som kan være substitueret med en eller flere grupper, der er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁-₁₂-alkylgruppe; omfattende i det mindste en af reaktionssekvenserne, som består af de følgende trin (A) og (D), (B) og (D) eller (C) og (D):

(A) omsætning af et 2-acyleret eller 2-iminoalkyleret acrylsyrederivat med formel (II),




hvor

Z er udvalgt af O, S og N⁺R¹⁰R¹¹, idet R¹⁰ og R¹¹ uafhængigt af hinanden er udvalgt af C₁-₁₂-alkyl-, C₃-₈-cycloalkyl-, C₂-₁₂-

alkenyl-, C₂₋₁₂-alkynyl-, C₆₋₈-aryl-, C₇₋₁₉-arylalkyl- eller C₇₋₁₉-alkylarylgrupper, som hver især kan være substitueret med en eller flere grupper, der er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁₋₁₂-alkylgruppe, og R¹⁰ og R¹¹ sammen med det nitrogenatom, som de er bundet til, kan danne en 5- til 6-leddet ring, som i givet fald kan indeholde et eller to yderligere heteroatomer, udvalgt af O, S og en SO₂-gruppe, og som kan være substitueret med en eller flere grupper, som er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁₋₁₂-alkylgruppe;

A er en udgangsgruppe NR¹²R¹³, idet R¹² og R¹³ uafhængigt af hinanden er udvalgt af C₁₋₁₂-alkyl-, C₃₋₈-cycloalkyl-, C₂₋₁₂-alkenyl-, C₂₋₁₂-alkynyl-, C₆₋₈-aryl-, C₇₋₁₉-arylalkyl- eller C₇₋₁₉-alkylaryl-grupper, som hver især kan være substitueret med en eller flere grupper, som er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁₋₁₂-alkylgruppe, og R¹² og R¹³ sammen med det nitrogenatom, som de er bundet til, kan danne en 5- til 6-leddet ring, som i givet fald kan indeholde et eller to yderligere heteroatomer udvalgt af O, S og en SO₂-gruppe, og som kan være substitueret med en eller flere grupper, som er udvalgt af gruppen bestående af -R', -X, -OR', -SR', -NR'₂, -SiR'₃, -COOR', -(C=O)R', -CN og -CONR'₂, idet R' er hydrogen eller en C₁₋₁₂-alkylgruppe;

med en N-alkyl-hydrazon med formel (III)



(III);


hvor

R⁸ er udvalgt af H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, phenyl, tolyl, cyclohexyl;


R⁹ er udvalgt af H, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, tert-butyl, phenyl, tolyl, cyclohexyl;

idet R⁸ og R⁹ i fællesskab med det C-atom, som de er knyttet til, kan danne en 5- eller 6-leddet ring;


(B) acylering af en acrylsyrehydrazon med formel (VI);



med et acylhalogenid med formel (X), idet X er udvalgt af F, Cl, Br eller I;




- 5 (C) iminoalkylering af et acrylsyrehydrazon med den ovenstående formel (VI) med en α,α -dihalogenamin med formel (XI), idet X er udvalgt af F, Cl, Br eller I,



- 10 (D) cyklisering af de i trin (A), (B) eller (C) opnåede mellemprodukter til 3-haloalkyl-pyrazol-4-carboxylylsyre-derivatet med formel (I).

2. Fremgangsmåde ifølge krav 1, kendetegnet ved, at der som mellemprodukter i trinnene (A), (B) eller (C) dannes et 2-acyleret eller et 2-iminoalkyleret hydrazinoacrylsyrederivat med formel (VII),



hvor grupperne R¹, R², R⁸, R⁹, Y og Z har de ovenfor beskrevne betegnelser.

- 20 3. Fremgangsmåde ifølge et af kravene 1 eller 2, kendetegnet ved, at reaktionssekvensen består af trinnene (A) og (D).

- 25 4. Fremgangsmåde ifølge et af kravene 1 eller 2, kendetegnet ved, at reaktionssekvensen består af trinnene (B) og (D).

5. Fremgangsmåde ifølge et af kravene 1 eller 2, kendetegnet ved, at reaktionssekvensen består af trinnene (C) og (D).

5 6. Fremgangsmåde ifølge et af kravene 1 til 2 eller 4, kendetegnet ved, at acylhalogenidet med formel (X) i trin (B) er udvalgt af gruppen bestående af difluoreddikesyrefluorid eller difluoreddikesyrechlorid.

10 7. Fremgangsmåde ifølge et af kravene 1 til 2 eller 5, kendetegnet ved, at α,α -dihalogenaminen med formel (XI) i trin (C) er 1,1,2,2-tetrafluorethyl-N,N-dimethylamin.


8. Fremgangsmåde ifølge et af kravene 1 til 3, kendetegnet ved,
 15 at det 2-acylerede eller 2-iminoalkylerede acrylsyre-derivat med formel (II) er udvalgt af gruppen bestående af N-1-(trifluormethyl)-3-(dimethylamino)-2-(ethoxycarbonyl)prop-2-en-1-ylidene]-N-methylmethanaminiumchlorid, N-1-(difluormethyl)-3-(dimethyl-amino)-2-(ethoxycarbonyl)prop-2-
 20 en-1-ylidene]-N-methylmethanaminium tetrafluorborat og N-[3-(dimethylamino)-2-(ethoxycarbonyl)-1-(1,1,2,2-tetrafluoroethyl)prop-2-en-1-ylidene]-N-methylmethanaminiumchlorid.

25 9. Fremgangsmåde ifølge et af kravene 1 til 3 eller 8, kendetegnet ved, at N-alkyl-hydrazonen med formel (III) er udvalgt af gruppen bestående af 1-methyl-2-(1-methylethylidene)hydrazin, 1-methyl-2-(1-phenylethylidene)hydrazin, 1-methyl-2-(1,2,2-trimethylpropylidene)hydrazin, 1-methyl-2-(1-methylpropylidene)hydrazin, 1-cyclohexylidene-2-methylhydrazin, 1-methyl-2-(phenyl-methylidene)hydrazin, 1-(diphenylmethylidene)-2-methylhydrazin, ethyl-2-[(dimethyl-amino)methylidene]-4,4,4-trifluoro-3-oxobutanoat.

35 10. Fremgangsmåde ifølge et af kravene 1 til 2 eller 4 til 7, kendetegnet ved, at acrylsyrehydrazonen med formel (VI) i trinnene (B) og (C) er udvalgt af gruppen bestående af ethyl


(2E)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]-prop-2-enoate,
 ethyl (2Z)-3-[1-methyl-2-(propan-2-ylidene)hydrazinyl]prop-2-
 enoate, methyl -3-[1-methyl-2-(propan-2-
 ylidene)hydrazinyl]prop-2-enoate, propyl-3-[1-methyl-2-
 5 (propan-2-ylidene)hydrazinyl]prop-2-enoate, ethyl (2E)-3-[1-
 methyl-2-(3,3-dimethylbutan-2-ylidene)hydrazinyl]prop-2-enoate,
 methyl 3-[1-methyl-2-(3,3-dimethylbutan-2-
 ylidene)hydrazinyl]prop-2-enoate, methyl-3-[1-methyl-2-
 (phenylmethylidene)-hydrazinyl]prop-2-enoate, (2E)-3-[1-
 10 methyl-2-(phenylmethylidene)hydrazinyl]prop-2-enenitrile og
 methyl -3-[1-ethyl-2-(propan-2-ylidene)hydrazinyl]prop-2-
 enoate.

11. 2-acyleret eller 2-iminoalkyleret acrylsyrehydrazon med
 15 formel (VII)




hvor grupperne R^1 , R^2 , R^8 , R^9 , Y og Z har de ovenfor beskrevne
 betegnelser.

20 12. 2-acylacrylhydrazon med formel (VIIa)



hvor grupperne R^1 , R^2 , R^8 , R^9 og Y har de ovenfor beskrevne
 betegnelser.

25 13. Salte med formel (VIIb)

Anion⁻

(VIIb),

hvor grupperne R^1 til R^8 og Y har de ovenfor beskrevne betegnelser, og anionen er udvalgt af gruppen bestående af Cl^- , BF_4^- , PF_6^- , SbF_6^- og $AlCl_4^-$.